

CS 207: Discrete Structures

Graph theory

Applications of Hall's theorem, matchings and vertex covers

Lecture 31

Oct 6 2015

Topic 3: Graph theory

Basic definitions and concepts

Characterizations

1. **Eulerian graphs:** Using degrees of vertices.
2. **Bipartite graphs:** Using odd length cycles.
3. **Connected components:** Using cycles.
4. **Maximum matchings:** Using augmenting paths.

Topic 3: Graph theory

Basic definitions and concepts

Characterizations

1. **Eulerian graphs:** Using degrees of vertices.
2. **Bipartite graphs:** Using odd length cycles.
3. **Connected components:** Using cycles.
4. **Maximum matchings:** Using augmenting paths.
5. **Perfect matchings in bipartite graphs:** Using neighbour sets. – **Hall's theorem**

Recap: Matchings


Definitions

- ▶ **Matching:** set of edges with no shared end-points.
- ▶ The vertices incident to edges in a matching are called **saturated**. Others are **unsaturated**.
- ▶ **Perfect matching:** saturates every vertex in graph.
- ▶ **Maximum matching:** matching of maximum size (# edges).
- ▶ **Maximal matching:** cannot be enlarged by adding an edge.

Recap: Matchings

Definitions

- ▶ **Matching**: set of edges with no shared end-points.
- ▶ The vertices incident to edges in a matching are called **saturated**. Others are **unsaturated**.
- ▶ **Perfect matching**: saturates every vertex in graph.
- ▶ **Maximum matching**: matching of maximum size ($\#$ edges).
- ▶ **Maximal matching**: cannot be enlarged by adding an edge.
- ▶ **M -alternating path**: alternates between edges in/out of M .
- ▶ **M -augmenting path**: An M -alternating path whose endpoints are unsaturated by M .



Recap: Matchings

Definitions

- ▶ **Matching:** set of edges with no shared end-points.
- ▶ The vertices incident to edges in a matching are called **saturated**. Others are **unsaturated**.
- ▶ **Perfect matching:** saturates every vertex in graph.
- ▶ **Maximum matching:** matching of maximum size (# edges).
- ▶ **Maximal matching:** cannot be enlarged by adding an edge.
- ▶ **M -alternating path:** alternates between edges in/out of M .
- ▶ **M -augmenting path:** An M -alternating path whose endpoints are unsaturated by M .

Theorem

A matching M in G is a maximum matching iff G has no M -augmenting path.

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

- ▶ For $v \in V$, its neighbour-set $N(v) = \{u \in V \mid (u, v) \in E\}$.
- ▶ For $S \subseteq V$, $N(S) = \{u \in V \mid (u, v) \in E \text{ for some } v \in S\}$.

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\implies) is straightforward:

- ▶ Let M be a matching.
- ▶ Then for any $S \subseteq X$, each vertex of S is matched to a distinct vertex in $N(S)$
- ▶ So $|N(S)| \geq |S|$.

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow)

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow)

- ▶ **Converse:** If for all $S \subseteq X$, $|N(S)| \geq |S|$, then G has a matching that saturates X .
- ▶ **Contrapositive:**

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow)

- ▶ **Converse:** If for all $S \subseteq X$, $|N(S)| \geq |S|$, then G has a matching that saturates X .
- ▶ **Contrapositive:** If G does not have any matching that saturates X , then there must exist $S \subseteq X$, $|N(S)| < |S|$.

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow)

- ▶ **Converse:** If for all $S \subseteq X$, $|N(S)| \geq |S|$, then G has a matching that saturates X .
- ▶ **Contrapositive:** If G does not have any matching that saturates X , then there must exist $S \subseteq X$, $|N(S)| < |S|$.
- ▶ If G does not have any matching that saturates X , then surely any maximum matching of G does not saturate X .

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow)

- ▶ **Converse:** If for all $S \subseteq X$, $|N(S)| \geq |S|$, then G has a matching that saturates X .
- ▶ **Contrapositive:** If G does not have any matching that saturates X , then there must exist $S \subseteq X$, $|N(S)| < |S|$.
- ▶ If G does not have any matching that saturates X , then surely any maximum matching of G does not saturate X .
- ▶ Let M be such a maximum matching. Then, we will construct $S \subseteq X$ s.t. $|N(S)| < |S|$.

Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow) Thus, starting from a maximum matching M which does not saturate X , we construct $S \subseteq X$, $|N(S)| < |S|$.

- Let $u \in X$ be any unsaturated vertex of M .


Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow) Thus, starting from a maximum matching M which does not saturate X , we construct $S \subseteq X$, $|N(S)| < |S|$.

- Consider vertices V_u from u by M -alternating paths in G and let $S = V_u \cap X$ and $T = V_u \cap Y$.




Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow) Thus, starting from a maximum matching M which does not saturate X , we construct $S \subseteq X$, $|N(S)| < |S|$.

- Consider vertices V_u from u by M -alternating paths in G and let $S = V_u \cap X$ and $T = V_u \cap Y$.



Claim: M matches T with $S \setminus \{u\}$ and $|N(S)| = |T|$.


Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow) Thus, starting from a maximum matching M which does not saturate X , we construct $S \subseteq X$, $|N(S)| < |S|$.

- Consider vertices V_u from u by M -alternating paths in G and let $S = V_u \cap X$ and $T = V_u \cap Y$.



Claim: M matches T with $S \setminus \{u\}$ and $|N(S)| = |T|$.

- Every vertex of $S \setminus \{u\}$ has an edge in M to a vertex in T .


Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow) Thus, starting from a maximum matching M which does not saturate X , we construct $S \subseteq X$, $|N(S)| < |S|$.

- ▶ Consider vertices V_u from u by M -alternating paths in G and let $S = V_u \cap X$ and $T = V_u \cap Y$.



Claim: M matches T with $S \setminus \{u\}$ and $|N(S)| = |T|$.

- ▶ Every vertex of $S \setminus \{u\}$ has an edge in M to a vertex in T .
- ▶ Every vertex of T extends via M to a unique vertex of S .


Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow) Thus, starting from a maximum matching M which does not saturate X , we construct $S \subseteq X$, $|N(S)| < |S|$.

- ▶ Consider vertices V_u from u by M -alternating paths in G and let $S = V_u \cap X$ and $T = V_u \cap Y$.



Claim: M matches T with $S \setminus \{u\}$ and $|N(S)| = |T|$.

- ▶ Every vertex of $S \setminus \{u\}$ has an edge in M to a vertex in T .
- ▶ Every vertex of T extends via M to a unique vertex of S .
- ▶ Thus, there is a bijection between T and $S \setminus \{u\}$.


Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow) Thus, starting from a maximum matching M which does not saturate X , we construct $S \subseteq X$, $|N(S)| < |S|$.

- Consider vertices V_u from u by M -alternating paths in G and let $S = V_u \cap X$ and $T = V_u \cap Y$.



Claim: M matches T with $S \setminus \{u\}$ and $|N(S)| = |T|$.

- $T \subseteq N(S)$ (from T any M -alternating path will reach S).


Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow) Thus, starting from a maximum matching M which does not saturate X , we construct $S \subseteq X$, $|N(S)| < |S|$.

- ▶ Consider vertices V_u from u by M -alternating paths in G and let $S = V_u \cap X$ and $T = V_u \cap Y$.



Claim: M matches T with $S \setminus \{u\}$ and $|N(S)| = |T|$.

- ▶ $T \subseteq N(S)$ (from T any M -alternating path will reach S).
- ▶ Conversely, if $v \in S$ has edge to $y \in Y \setminus T$, then path from u to v via M to y is an M -alternating path, implies $y \in T$.


Characterizing perfect matchings in bipartite graphs

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Proof: (\Leftarrow) Thus, starting from a maximum matching M which does not saturate X , we construct $S \subseteq X$, $|N(S)| < |S|$.

- Consider vertices V_u from u by M -alternating paths in G and let $S = V_u \cap X$ and $T = V_u \cap Y$.



Claim: M matches T with $S \setminus \{u\}$ and $|N(S)| = |T|$.

Thus, $|N(S)| = |T| = |S| - 1 < |S|$

□.

Applications of Hall's condition

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Corollary, Marriage theorem

- ▶ Consider n women and n men. If every woman is compatible with k men and every man compatible with k women, then a perfect matching must exist.
- ▶ What is the formal statement?

Applications of Hall's condition

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Corollary, Marriage theorem

- ▶ Consider n women and n men. If every woman is compatible with k men and every man compatible with k women, then a perfect matching must exist.
- ▶ For $k > 0$, every **k -regular** bipartite graph (i.e, **every vertex has degree exactly k**) has a perfect matching. **Prove this!**

Applications of Hall's condition

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Corollary, Marriage theorem

- ▶ Consider n women and n men. If every woman is compatible with k men and every man compatible with k women, then a perfect matching must exist.
- ▶ For $k > 0$, every **k -regular** bipartite graph (i.e, **every vertex has degree exactly k**) has a perfect matching. **Prove this!**
- ▶ If G is a k -regular X, Y bipartite graph, then $|X| = |Y|$. (why?)

Applications of Hall's condition

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Corollary, Marriage theorem

- ▶ Consider n women and n men. If every woman is compatible with k men and every man compatible with k women, then a perfect matching must exist.
- ▶ For $k > 0$, every **k -regular** bipartite graph (i.e, **every vertex has degree exactly k**) has a perfect matching. **Prove this!**
- ▶ If G is a k -regular X, Y bipartite graph, then $|X| = |Y|$.
- ▶ If a matching saturates X then it saturates Y .

Applications of Hall's condition

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Corollary, Marriage theorem

- ▶ Consider n women and n men. If every woman is compatible with k men and every man compatible with k women, then a perfect matching must exist.
- ▶ For $k > 0$, every **k -regular** bipartite graph (i.e, **every vertex has degree exactly k**) has a perfect matching. **Prove this!**
- ▶ If G is a k -regular X, Y bipartite graph, then $|X| = |Y|$.
- ▶ If a matching saturates X then it saturates Y .
- ▶ Can you now verify Hall's condition?

Applications of Hall's condition

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Corollary, Marriage theorem

- ▶ Consider n women and n men. If every woman is compatible with k men and every man compatible with k women, then a perfect matching must exist.
- ▶ For $k > 0$, every **k -regular** bipartite graph (i.e, **every vertex has degree exactly k**) has a perfect matching. **Prove this!**
- ▶ If G is a k -regular X, Y bipartite graph, then $|X| = |Y|$.
- ▶ If a matching saturates X then it saturates Y .
 - ▶ Let $S \subseteq X$. Let $m = \#$ edges from S to $N(S)$.

Applications of Hall's condition

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Corollary, Marriage theorem

- ▶ Consider n women and n men. If every woman is compatible with k men and every man compatible with k women, then a perfect matching must exist.
- ▶ For $k > 0$, every **k -regular** bipartite graph (i.e, **every vertex has degree exactly k**) has a perfect matching. **Prove this!**
- ▶ If G is a k -regular X, Y bipartite graph, then $|X| = |Y|$.
- ▶ If a matching saturates X then it saturates Y .
 - ▶ Let $S \subseteq X$. Let $m = \#$ edges from S to $N(S)$.
 - ▶ Since G is k -regular, $m = k|S|$. And since they touch $N(S)$, $m \leq k|N(S)|$.

Applications of Hall's condition

Theorem (Hall'35)

A bipartite graph G with bipartitions X, Y has a matching that saturates X iff for all $S \subseteq X$, $|N(S)| \geq |S|$.

Corollary, Marriage theorem

- ▶ Consider n women and n men. If every woman is compatible with k men and every man compatible with k women, then a perfect matching must exist.
- ▶ For $k > 0$, every **k -regular** bipartite graph (i.e, **every vertex has degree exactly k**) has a perfect matching. **Prove this!**
- ▶ If G is a k -regular X, Y bipartite graph, then $|X| = |Y|$.
- ▶ If a matching saturates X then it saturates Y .
 - ▶ Let $S \subseteq X$. Let $m = \#$ edges from S to $N(S)$.
 - ▶ Since G is k -regular, $m = k|S|$. And since they touch $N(S)$, $m \leq k|N(S)|$. Hence $k|S| \leq k|N(S)|$, $k > 0$ completes proof. □

Let's play a game

A two player game on a graph


1. Given a graph G , two players will alternatively choose distinct vertices.
2. One player starts by choosing any vertex.
3. Subsequent move must be adjacent to preceding choice (of other player).
4. Last player who can move wins.

Let's play a game

A two player game on a graph


1. Given a graph G , two players will alternatively choose distinct vertices.
2. One player starts by choosing any vertex.
3. Subsequent move must be adjacent to preceding choice (of other player).
4. Last player who can move wins.

Two volunteers please. Who wants to start?





Another game

Consider the graph of a road network in a city. When a minister is visiting, the Chief of Police wants to place a policeman to watch every road. What is the minimum number of policemen required?




Another game

Consider the graph of a road network in a city. When a minister is visiting, the Chief of Police wants to place a policeman to watch every road. What is the minimum number of policemen required?



Another game

Consider the graph of a road network in a city. When a minister is visiting, the Chief of Police wants to place a policeman to watch every road. What is the minimum number of policemen required?



Definition

A **vertex cover** of a graph G is a set $Q \subseteq V$ that contains at least one endpoint of every edge. The vertices in Q are said to **cover** E .

Matchings and vertex covers

So, what is the link between matchings and vertex covers?

Matchings and vertex covers

So, what is the link between matchings and vertex covers?

Examples and properties of vertex covers

Matchings and vertex covers

So, what is the link between matchings and vertex covers?

Examples and properties of vertex covers

- ▶ The set of all vertices is always a vertex cover.

Matchings and vertex covers

So, what is the link between matchings and vertex covers?

Examples and properties of vertex covers

- ▶ The set of all vertices is always a vertex cover.
- ▶ The end-points of a maximal matching

Matchings and vertex covers

So, what is the link between matchings and vertex covers?

Examples and properties of vertex covers

- ▶ The set of all vertices is always a vertex cover.
- ▶ The end-points of a maximal matching form a vertex cover.

Matchings and vertex covers

So, what is the link between matchings and vertex covers?

Examples and properties of vertex covers

- ▶ The set of all vertices is always a vertex cover.
- ▶ The end-points of a maximal matching form a vertex cover.
- ▶ What about the complement of a vertex cover?

Matchings and vertex covers

So, what is the link between matchings and vertex covers?

Examples and properties of vertex covers

- ▶ The set of all vertices is always a vertex cover.
- ▶ The end-points of a maximal matching form a vertex cover.
- ▶ What about the complement of a vertex cover?
- ▶ What is the size of the minimum vertex cover in $K_{m,n}$?

Matchings and vertex covers

So, what is the link between matchings and vertex covers?

Examples and properties of vertex covers

- ▶ The set of all vertices is always a vertex cover.
- ▶ The end-points of a maximal matching form a vertex cover.
- ▶ What about the complement of a vertex cover?
- ▶ What is the size of the minimum vertex cover in $K_{m,n}$?

So, how do you compute min no. of policemen required, i.e., the size of the minimum vertex covers?

Matchings and vertex covers

So, what is the link between matchings and vertex covers?

Examples and properties of vertex covers

- ▶ The set of all vertices is always a vertex cover.
- ▶ The end-points of a maximal matching form a vertex cover.
- ▶ What about the complement of a vertex cover?
- ▶ What is the size of the minimum vertex cover in $K_{m,n}$?

So, how do you compute min no. of policemen required, i.e., the size of the minimum vertex covers?

Let's consider bipartite graphs...

A min-max theorem

Theorem (Konig '31)

If G is a bipartite graph, then the size of the maximum matching of G equals the size of the minimum vertex cover of G .