

CS 207: Discrete Structures

Lecture 14 – Counting and Combinatorics

Aug 18 2015

Last two classes

Basic counting techniques and applications

1. Sum and product, bijection, double counting principles
2. Counting no. of (ordered) subsets, partitions, relations...
3. Handshake lemma

Last two classes

Basic counting techniques and applications

1. Sum and product, bijection, double counting principles
2. Counting no. of (ordered) subsets, partitions, relations...
3. Handshake lemma
4. Binomial coefficients and binomial theorem
5. Pascal's triangle and its applications
6. Permutations and combinations with/without repetitions

Estimating $n!$

How big is $n!$?

- ▶ It is clearly bigger than n and n^2 .

Estimating $n!$

How big is $n!$?

- ▶ It is clearly bigger than n and n^2 .
- ▶ Is it bigger than 2^n , n^n ?

Estimating $n!$

How big is $n!$?

- ▶ It is clearly bigger than n and n^2 .
- ▶ Is it bigger than 2^n , n^n ?
- ▶ Easy to see: for all $n \geq 4$,

$$2^n \leq n! \leq n^n$$

Estimating $n!$

How big is $n!$?

- ▶ It is clearly bigger than n and n^2 .
- ▶ Is it bigger than 2^n , n^n ?
- ▶ Easy to see: for all $n \geq 4$,

$$2^n \leq n! \leq n^n$$

- ▶ Can we do better?
Can we quantify how much more n^n is compared to $n!$?

Estimating $n!$

How big is $n!$?

- ▶ It is clearly bigger than n and n^2 .
- ▶ Is it bigger than 2^n , n^n ?
- ▶ Easy to see: for all $n \geq 4$,

$$2^n \leq n! \leq n^n$$

- ▶ Can we do better?
Can we quantify how much more n^n is compared to $n!$?

Theorem (Stirling's Approximation)

$$e \left(\frac{n}{e}\right)^n \leq n! \leq ne \left(\frac{n}{e}\right)^n$$

where e is the base of natural logarithms, $\log(e) = e^{\log(e)} = 1$.

Approximating the factorial

Theorem (Stirling's Approximation)

$$e \left(\frac{n}{e}\right)^n \leq n! \leq ne \left(\frac{n}{e}\right)^n$$

where e is the base of natural logarithms, $\log(e) = e^{\log(e)} = 1$.

Approximating the factorial

Theorem (Stirling's Approximation)

$$e \left(\frac{n}{e}\right)^n \leq n! \leq ne \left(\frac{n}{e}\right)^n$$

where e is the base of natural logarithms, $\log(e) = e^{\log(e)} = 1$.

Proof: Let $S = \log(n!) = \sum_{i=1}^n \log(i)$. Thus, $e^S = n!$

Approximating the factorial


Theorem (Stirling's Approximation)

$$e \left(\frac{n}{e}\right)^n \leq n! \leq ne \left(\frac{n}{e}\right)^n$$

where e is the base of natural logarithms, $\log(e) = e^{\log(e)} = 1$.

Proof: Let $S = \log(n!) = \sum_{i=1}^n \log(i)$. Thus, $e^S = n!$

Now, we relate it to natural log function as shown in the figure.



$$\sum_{i=1}^n \log(i) \geq \int_1^n \log(x) dx \geq \sum_{i=1}^{n-1} \log(i)$$

Approximating the factorial

Theorem (Stirling's Approximation)

$$e \left(\frac{n}{e}\right)^n \leq n! \leq ne \left(\frac{n}{e}\right)^n$$

where e is the base of natural logarithms, $\log(e) = e^{\log(e)} = 1$.

Proof: Let $S = \log(n!) = \sum_{i=1}^n \log(i)$. Thus, $e^S = n!$

$$\sum_{i=1}^n \log(i) \geq \int_1^n \log(x) dx \geq \sum_{i=1}^{n-1} \log(i)$$

Approximating the factorial

Theorem (Stirling's Approximation)

$$e \left(\frac{n}{e}\right)^n \leq n! \leq ne \left(\frac{n}{e}\right)^n$$

where e is the base of natural logarithms, $\log(e) = e^{\log(e)} = 1$.

Proof: Let $S = \log(n!) = \sum_{i=1}^n \log(i)$. Thus, $e^S = n!$

$$\sum_{i=1}^n \log(i) \geq \int_1^n \log(x) dx \geq \sum_{i=1}^{n-1} \log(i)$$

$$S \geq x \log(x) - x \Big|_1^n \geq S - \log(n)$$

Approximating the factorial

Theorem (Stirling's Approximation)

$$e \left(\frac{n}{e}\right)^n \leq n! \leq ne \left(\frac{n}{e}\right)^n$$

where e is the base of natural logarithms, $\log(e) = e^{\log(e)} = 1$.

Proof: Let $S = \log(n!) = \sum_{i=1}^n \log(i)$. Thus, $e^S = n!$

$$\sum_{i=1}^n \log(i) \geq \int_1^n \log(x) dx \geq \sum_{i=1}^{n-1} \log(i)$$

$$S \geq x \log(x) - x \Big|_1^n \geq S - \log(n)$$

$$S \geq n \log(n) - n + 1 \geq S - \log(n).$$

Approximating the factorial

Theorem (Stirling's Approximation)

$$e \left(\frac{n}{e}\right)^n \leq n! \leq ne \left(\frac{n}{e}\right)^n$$

where e is the base of natural logarithms, $\log(e) = e^{\log(e)} = 1$.

Proof: Let $S = \log(n!) = \sum_{i=1}^n \log(i)$. Thus, $e^S = n!$

$$\sum_{i=1}^n \log(i) \geq \int_1^n \log(x) dx \geq \sum_{i=1}^{n-1} \log(i)$$

$$S \geq x \log(x) - x \Big|_1^n \geq S - \log(n)$$

$$S \geq n \log(n) - n + 1 \geq S - \log(n).$$

Raising both sides to power of e we get

► l.h.s. $n! \geq e^{n \log(n) - n + 1} = (n/e)^n e$ and

Approximating the factorial

Theorem (Stirling's Approximation)

$$e \left(\frac{n}{e}\right)^n \leq n! \leq ne \left(\frac{n}{e}\right)^n$$

where e is the base of natural logarithms, $\log(e) = e^{\log(e)} = 1$.

Proof: Let $S = \log(n!) = \sum_{i=1}^n \log(i)$. Thus, $e^S = n!$

$$\sum_{i=1}^n \log(i) \geq \int_1^n \log(x) dx \geq \sum_{i=1}^{n-1} \log(i)$$

$$S \geq x \log(x) - x \Big|_1^n \geq S - \log(n)$$

$$S \geq n \log(n) - n + 1 \geq S - \log(n).$$

Raising both sides to power of e we get

► l.h.s. $n! \geq e^{n \log(n) - n + 1} = (n/e)^n e$ and

► r.h.s. $n! \leq e^{(n+1) \log(n) - n + 1} = n^{n+1} / e^{n-1} = ne(n/e)^n$. □

Next: Recurrence relations and generating functions

Recall: No. of subsets of a set of n elements

How many subsets does a set A of n elements have?

- ▶ **Induction**
- ▶ **Product principle**: two choices for each element, hence $2 \cdot 2 \cdots 2 \cdot 2$ (n -times).
- ▶ **Bijection**: between $\mathcal{P}(X)$ and n -length $\{0, 1\}$ -sequences.
- ▶ **Sum principle**: Subsets of size 0 + subsets of size 1 + \dots + subsets of size n = Total number of subsets.

Next: Recurrence relations and generating functions

Recall: No. of subsets of a set of n elements

How many subsets does a set A of n elements have?

- ▶ **Induction**
- ▶ **Product principle**: two choices for each element, hence $2 \cdot 2 \cdots 2 \cdot 2$ (n -times).
- ▶ **Bijection**: between $\mathcal{P}(X)$ and n -length $\{0, 1\}$ -sequences.
- ▶ **Sum principle**: Subsets of size 0 + subsets of size 1 + \dots + subsets of size n = Total number of subsets.
- ▶ **Recurrence**: $F(n) = 2 \cdot F(n - 1), F(0) = 1$.

Next: Recurrence relations and generating functions

Recall: No. of subsets of a set of n elements

How many subsets does a set A of n elements have?

- ▶ **Induction**
- ▶ **Product principle**: two choices for each element, hence $2 \cdot 2 \cdots 2 \cdot 2$ (n -times).
- ▶ **Bijection**: between $\mathcal{P}(X)$ and n -length $\{0, 1\}$ -sequences.
- ▶ **Sum principle**: Subsets of size 0 + subsets of size 1 + \dots + subsets of size n = Total number of subsets.
- ▶ **Recurrence**: $F(n) = 2 \cdot F(n - 1), F(0) = 1$.

Next: Recurrence relations and generating functions


Recall: No. of subsets of a set of n elements

How many subsets does a set A of n elements have?

- ▶ **Induction**
- ▶ **Product principle**: two choices for each element, hence $2 \cdot 2 \cdots 2 \cdot 2$ (n -times).
- ▶ **Bijection**: between $\mathcal{P}(X)$ and n -length $\{0, 1\}$ -sequences.
- ▶ **Sum principle**: Subsets of size 0 + subsets of size 1 + ... + subsets of size n = Total number of subsets.
- ▶ **Recurrence**: $F(n) = 2 \cdot F(n - 1), F(0) = 1$.


But how do you solve it?

Another example of recurrence: The Fibonacci Sequence




- Fibonacci sequence: 1, 1, 2, 3, 5, 8, 13, 21, ...

Another example of recurrence: The Fibonacci Sequence




- ▶ Fibonacci sequence: $1, 1, 2, 3, 5, 8, 13, 21, \dots$
- ▶ Recurrence relation: $u_n = u_{n-1} + u_{n-2}$ where $u_1 = u_0 = 1$

Another example of recurrence: The Fibonacci Sequence



- ▶ Fibonacci sequence: 1, 1, 2, 3, 5, 8, 13, 21, ...
- ▶ Recurrence relation: $u_n = u_{n-1} + u_{n-2}$ where $u_1 = u_0 = 1$
- ▶ But rabbits die!

Another example of recurrence: The Fibonacci Sequence



- ▶ Fibonacci sequence: $1, 1, 2, 3, 5, 8, 13, 21, \dots$
- ▶ Recurrence relation: $u_n = u_{n-1} + u_{n-2}$ where $u_1 = u_0 = 1$
- ▶ But rabbits die!
- ▶ Consider $u_n = u_{n-1} + u_{n-2} - u_{n-3}$ where $u_2 = 2, u_1 = u_0 = 1$

Recurrence and linear recurrence relations

Definition

- ▶ A **recurrence relation** for a sequence is an equation that expresses its n^{th} term using one or more of the previous terms of the sequence.
- ▶ A **linear recurrence relation** is of the form

$$u_n = a_{k-1}u_{n-1} + \dots + a_1u_{n-k+1} + a_0u_{n-k}$$

where $a_0, \dots, a_{k-1} \in \mathbb{R}$ are constants.

- ▶ k is called the **degree/depth** of the sequence.
- ▶ The first few (e.g., k elements u_0, \dots, u_{k-1}) are **initial conditions** and they determine the whole sequence.

Some more examples of recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 7?

Some more examples of recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 7?

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

Some more examples of recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 7?

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Example: $n = 3 : ((a + b) + c), (a + (b + c))$

Some more examples of recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 7?

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Example: $n = 3 : ((a + b) + c), (a + (b + c))$
- ▶ $n = 4 : (((a + b) + c) + d), ((a + b) + (c + d)), ((a + (b + c)) + d), \dots$

Some more examples of recurrences

How many bit strings of length n are there that do not have two consecutive 0's?

- ▶ Find a recurrence relation for this
- ▶ Give the initial conditions
- ▶ How many such bit strings are there of length 7?

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Example: $n = 3 : ((a + b) + c), (a + (b + c))$
- ▶ $n = 4 : (((a + b) + c) + d), ((a + b) + (c + d)), ((a + (b + c)) + d), \dots$
In general, let $C(n)$ be the number of ways of doing this.

Some more examples of recurrences

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Let $C(n)$ be the number of ways of doing this.
- ▶ If outermost bracketing $(A + B)$ appears between x_k and x_{k+1} , then there are $C(k) \cdot C(n - k)$ ways of bracketing it.

Some more examples of recurrences

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Let $C(n)$ be the number of ways of doing this.
- ▶ If outermost bracketing $(A + B)$ appears between x_k and x_{k+1} , then there are $C(k) \cdot C(n - k)$ ways of bracketing it.
- ▶ k can be anything from 1 till $n - 1$

Some more examples of recurrences

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Let $C(n)$ be the number of ways of doing this.
- ▶ If outermost bracketing $(A + B)$ appears between x_k and x_{k+1} , then there are $C(k) \cdot C(n - k)$ ways of bracketing it.
- ▶ k can be anything from 1 till $n - 1$
- ▶ Thus,
$$C(n) = \sum_{i=1}^{n-1} C(i)C(n - i) \text{ for } n > 1$$


Some more examples of recurrences

How many ways are there to bracket a sum of n terms so that it can be computed by adding two numbers at a time?

- ▶ Let $C(n)$ be the number of ways of doing this.
 - ▶ If outermost bracketing $(A + B)$ appears between x_k and x_{k+1} , then there are $C(k) \cdot C(n - k)$ ways of bracketing it.
 - ▶ k can be anything from 1 till $n - 1$
 - ▶ Thus, $C(n) = \sum_{i=1}^{n-1} C(i)C(n - i)$ for $n > 1$
-
- ▶ Initial conditions are $C(0) = C(1) = 1$.
 - ▶ This sequence are called Catalan numbers...

How do we **solve** such recurrences? We start with the Fibonacci sequence.

An aside: find the Fibonacci sequence!



- ▶ $F(n) = F(n-1) + F(n-2)$.
- ▶ 1, 1, 2, 3, 5, 8, 13, ...
- ▶ Can you observe the sum of which terms in the Pascal's triangle gives rise to the terms of the Fibonacci sequence?