CS 207: Discrete Structures

Abstract algebra and Number theory — Modular arithmetic and RSA

Lecture 41 Nov 3 2015

Definition

For integers a, b and positive integer m, if m|(a-b), then we say that a is congruent to b modulo m, denoted $a \equiv b \mod m$.

).

Definition

For integers a, b and positive integer m, if m|(a-b), then we say that a is congruent to b modulo m, denoted $a \equiv b \mod m$.

- $(\mathbb{Z}_p \setminus \{0\}, \times_p)$ is a group.
 - ▶ What are its elements?

Definition

For integers a, b and positive integer m, if m|(a-b), then we say that a is congruent to b modulo m, denoted $a \equiv b \mod m$.

- $(\mathbb{Z}_p \setminus \{0\}, \times_p)$ is a group.
 - ▶ What are its elements? $\mathbb{Z}_p \setminus \{0\} = \{1, 2, \dots, p-1\}$
 - ▶ What is the order of this group?

).

Definition

For integers a, b and positive integer m, if m|(a-b), then we say that a is congruent to b modulo m, denoted $a \equiv b \mod m$.

- $(\mathbb{Z}_p \setminus \{0\}, \times_p)$ is a group.
 - ▶ What are its elements? $\mathbb{Z}_p \setminus \{0\} = \{1, 2, \dots, p-1\}$
 - ▶ What is the order of this group? p-1.

).

Definition

For integers a, b and positive integer m, if m|(a-b), then we say that a is congruent to b modulo m, denoted $a \equiv b \mod m$.

- $(\mathbb{Z}_p \setminus \{0\}, \times_p)$ is a group.
 - ▶ What are its elements? $\mathbb{Z}_p \setminus \{0\} = \{1, 2, \dots, p-1\}$
 - ▶ What is the order of this group? p-1.
- ▶ What about $(\mathbb{Z}_n \setminus \{0\}, \times_n)$? Which elements have inverses?

Definition

For integers a, b and positive integer m, if m | (a - b), then we say that a is congruent to b modulo m, denoted $a \equiv b \mod m$.

- $(\mathbb{Z}_p \setminus \{0\}, \times_p)$ is a group.
 - ▶ What are its elements? $\mathbb{Z}_p \setminus \{0\} = \{1, 2, \dots, p-1\}$
 - ▶ What is the order of this group? p-1.
- ▶ What about $(\mathbb{Z}_n \setminus \{0\}, \times_n)$? Which elements have inverses?
- ightharpoonup Only those co-prime to n

Definition

For integers a, b and positive integer m, if m | (a - b), then we say that a is congruent to b modulo m, denoted $a \equiv b \mod m$.

- $(\mathbb{Z}_p \setminus \{0\}, \times_p)$ is a group.
 - ▶ What are its elements? $\mathbb{Z}_p \setminus \{0\} = \{1, 2, \dots, p-1\}$
 - ▶ What is the order of this group? p-1.
- ▶ What about $(\mathbb{Z}_n \setminus \{0\}, \times_n)$? Which elements have inverses?
- \triangleright Only those co-prime to n because,
 - ▶ if gcd(a, n) = 1 then $\exists r, s \text{ s.t.}$, ar + ns = gcd(a, n) = 1. Again implies $ar \equiv 1 \mod n$, i.e., r is inverse of $a \mod n$.

).

Definition

For integers a, b and positive integer m, if m|(a-b), then we say that a is congruent to b modulo m, denoted $a \equiv b \mod m$.

- $(\mathbb{Z}_p \setminus \{0\}, \times_p)$ is a group.
 - ▶ What are its elements? $\mathbb{Z}_p \setminus \{0\} = \{1, 2, \dots, p-1\}$
 - ▶ What is the order of this group? p-1.
- ▶ What about $(\mathbb{Z}_n \setminus \{0\}, \times_n)$? Which elements have inverses?
- \triangleright Only those co-prime to n because,
 - if gcd(a, n) = 1 then $\exists r, s \text{ s.t.}$, ar + ns = gcd(a, n) = 1. Again implies $ar \equiv 1 \mod n$, i.e., r is inverse of $a \mod n$.
- ▶ Thus, numbers co-prime to n, denoted $Coprime(\mathbb{Z}_n)$ form a group under \times_n . Check!

).

Definition

For integers a, b and positive integer m, if m|(a-b), then we say that a is congruent to b modulo m, denoted $a \equiv b \mod m$.

- $(\mathbb{Z}_p \setminus \{0\}, \times_p)$ is a group.
 - ▶ What are its elements? $\mathbb{Z}_p \setminus \{0\} = \{1, 2, \dots, p-1\}$
 - ▶ What is the order of this group? p-1.
- ▶ What about $(\mathbb{Z}_n \setminus \{0\}, \times_n)$? Which elements have inverses?
- \triangleright Only those co-prime to n because,
 - ▶ if gcd(a, n) = 1 then $\exists r, s \text{ s.t.}, ar + ns = gcd(a, n) = 1$. Again implies $ar \equiv 1 \mod n$, i.e., r is inverse of $a \mod n$.
- ▶ Thus, numbers co-prime to n, denoted $Coprime(\mathbb{Z}_n)$ form a group under \times_n . Check!
- ▶ The number of elements co-prime to n, i.e., $|Coprime(\mathbb{Z}_n)|$ is denoted $\phi(n)$ called the Euler totient no./function.

- ▶ $\forall n \in \mathbb{Z}^+$, $Coprime(\mathbb{Z}_n)$ denotes numbers coprime to n.
- ▶ $(Coprime(\mathbb{Z}_n), \times_n)$ is a group, where $a \times_n b = ab \mod n$.
- ▶ Order of this group is

- $\blacktriangleright \forall n \in \mathbb{Z}^+, Coprime(\mathbb{Z}_n)$ denotes numbers coprime to n.
- $(Coprime(\mathbb{Z}_n), \times_n)$ is a group, where $a \times_n b = ab \mod n$.
- ▶ Order of this group is $\phi(n)$, no. of elements co-prime to n.

- ▶ $\forall n \in \mathbb{Z}^+$, $Coprime(\mathbb{Z}_n)$ denotes numbers coprime to n.
- $(Coprime(\mathbb{Z}_n), \times_n)$ is a group, where $a \times_n b = ab \mod n$.
- ▶ Order of this group is $\phi(n)$, no. of elements co-prime to n.

- 1. Suppose p is a prime, what is $(Coprime(\mathbb{Z}_p), \times_p)$?
- 2. What is $\phi(p)$?
- 3. Suppose n = pq is a product of primes, what is $\phi(pq)$, i.e., how many numbers are there co-prime to pq in $\{1, \ldots, pq\}$?

- ▶ $\forall n \in \mathbb{Z}^+$, $Coprime(\mathbb{Z}_n)$ denotes numbers coprime to n.
- $(Coprime(\mathbb{Z}_n), \times_n)$ is a group, where $a \times_n b = ab \mod n$.
- ▶ Order of this group is $\phi(n)$, no. of elements co-prime to n.

- 1. Suppose p is a prime, what is $(Coprime(\mathbb{Z}_p), \times_p)$?
- 2. What is $\phi(p)$?
- 3. Suppose n = pq is a product of primes, what is $\phi(pq)$, i.e., how many numbers are there co-prime to pq in $\{1, \ldots, pq\}$?
- 4. (Euler's theorem) If gcd(a, n) = 1, then $a^{\phi(n)} \equiv 1 \mod n$.

- $\blacktriangleright \forall n \in \mathbb{Z}^+, Coprime(\mathbb{Z}_n)$ denotes numbers coprime to n.
- $(Coprime(\mathbb{Z}_n), \times_n)$ is a group, where $a \times_n b = ab \mod n$.
- ▶ Order of this group is $\phi(n)$, no. of elements co-prime to n.

- 1. Suppose p is a prime, what is $(Coprime(\mathbb{Z}_p), \times_p)$?

 the usual (\mathbb{Z}_p, \times_p)
- 2. What is $\phi(p)$? p-1
- 3. Suppose n=pq is a product of primes, what is $\phi(pq)$, i.e., how many numbers are there co-prime to pq in $\{1,\ldots,pq\}$? $\phi(pq)=(p-1)(q-1)$
- 4. (Euler's theorem) If gcd(a, n) = 1, then $a^{\phi(n)} \equiv 1 \mod n$.

- $\blacktriangleright \forall n \in \mathbb{Z}^+, Coprime(\mathbb{Z}_n)$ denotes numbers coprime to n.
- $(Coprime(\mathbb{Z}_n), \times_n)$ is a group, where $a \times_n b = ab \mod n$.
- ▶ Order of this group is $\phi(n)$, no. of elements co-prime to n.

- 1. Suppose p is a prime, what is $(Coprime(\mathbb{Z}_p), \times_p)$?
- 2. What is $\phi(p)$?
- 3. Suppose n=pq is a product of primes, what is $\phi(pq)$, i.e., how many numbers are there co-prime to pq in $\{1,\ldots,pq\}$? $\phi(pq)=(p-1)(q-1)$
- 4. (Euler's theorem) If gcd(a, n) = 1, then $a^{\phi(n)} \equiv 1 \mod n$.
 - ▶ Let gcd(a, n) = 1, $\phi(n) = |(Coprime(\mathbb{Z}_n), \times_n)|$

- ▶ $\forall n \in \mathbb{Z}^+$, $Coprime(\mathbb{Z}_n)$ denotes numbers coprime to n.
- $(Coprime(\mathbb{Z}_n), \times_n)$ is a group, where $a \times_n b = ab \mod n$.
- ▶ Order of this group is $\phi(n)$, no. of elements co-prime to n.

- 1. Suppose p is a prime, what is $(Coprime(\mathbb{Z}_p), \times_p)$?
- 2. What is $\phi(p)$?
- 3. Suppose n=pq is a product of primes, what is $\phi(pq)$, i.e., how many numbers are there co-prime to pq in $\{1,\ldots,pq\}$? $\phi(pq)=(p-1)(q-1)$
- 4. (Euler's theorem) If gcd(a, n) = 1, then $a^{\phi(n)} \equiv 1 \mod n$.
 - ▶ Let gcd(a, n) = 1, $\phi(n) = |(Coprime(\mathbb{Z}_n), \times_n)|$
 - ▶ By Lagrange's theorem, order of element a divides order of this group, i.e., $a^{(ord.ofgrp)} = (id \text{ of this grp})$.

- ▶ $\forall n \in \mathbb{Z}^+$, $Coprime(\mathbb{Z}_n)$ denotes numbers coprime to n.
- $(Coprime(\mathbb{Z}_n), \times_n)$ is a group, where $a \times_n b = ab \mod n$.
- ▶ Order of this group is $\phi(n)$, no. of elements co-prime to n.

- 1. Suppose p is a prime, what is $(Coprime(\mathbb{Z}_p), \times_p)$?
- 2. What is $\phi(p)$?
- 3. Suppose n=pq is a product of primes, what is $\phi(pq)$, i.e., how many numbers are there co-prime to pq in $\{1,\ldots,pq\}$? $\phi(pq)=(p-1)(q-1)$
- 4. (Euler's theorem) If gcd(a, n) = 1, then $a^{\phi(n)} \equiv 1 \mod n$.
 - Let gcd(a, n) = 1, $\phi(n) = |(Coprime(\mathbb{Z}_n), \times_n)|$
 - ▶ By Lagrange's theorem, order of element a divides order of this group, i.e., $a^{(ord.ofgrp)} = (id \text{ of this grp})$.
 - That is, $a^{\phi(n)} = 1 + kn$ for some k, i.e., $a^{\phi(n)} \equiv 1 \mod n$

- ▶ Alice wants to send a message M (a number, lets say between 1 and 20000) to Bob that only he can figure out.
- ▶ But any message can be intercepted (hacker: Carol!)
- ► How can Bob ensure privacy of messages?

Bob starts by choosing two large primes p, q

- ▶ Alice wants to send a message M (a number, lets say between 1 and 20000) to Bob that only he can figure out.
- ▶ But any message can be intercepted (hacker: Carol!)
- ▶ How can Bob ensure privacy of messages?

Bob starts by choosing two large primes p, q

Also chooses numbers D and E such that $ED \equiv 1 \mod \phi(pq)$ (i.e., $\phi(pq) \mid (ED - 1)$).

- ▶ Alice wants to send a message M (a number, lets say between 1 and 20000) to Bob that only he can figure out.
- ▶ But any message can be intercepted (hacker: Carol!)
- ▶ How can Bob ensure privacy of messages?

Bob starts by choosing two large primes p, q

Also chooses numbers D and E such that $ED \equiv 1 \mod \phi(pq)$ (i.e., $\phi(pq) \mid (ED-1)$).

- 1. Bob shouts out N = pq and E (encryptor!)...
- 2. But he keeps p, q and the decryptor D secret.

- ▶ Alice wants to send a message M (a number, lets say between 1 and 20000) to Bob that only he can figure out.
- ▶ But any message can be intercepted (hacker: Carol!)
- ► How can Bob ensure privacy of messages?

Bob starts by choosing two large primes p, q

Also chooses numbers D and E such that $ED \equiv 1 \mod \phi(pq)$ (i.e., $\phi(pq) \mid (ED-1)$).

- 1. Bob shouts out N = pq and E (encryptor!)...
- 2. But he keeps p, q and the decryptor D secret.
- 3. Alice wants to send message M < p, q that only Bob understands. What does she send?

- ▶ Alice wants to send a message M (a number, lets say between 1 and 20000) to Bob that only he can figure out.
- ▶ But any message can be intercepted (hacker: Carol!)
- ▶ How can Bob ensure privacy of messages?

Bob starts by choosing two large primes p, q

Also chooses numbers D and E such that $ED \equiv 1 \mod \phi(pq)$ (i.e., $\phi(pq) \mid (ED-1)$).

- 1. Bob shouts out N = pq and E (encryptor!)...
- 2. But he keeps p, q and the decryptor D secret.
- 3. Alice wants to send message M < p, q that only Bob understands. What does she send?
- 4. Alice sends $X = M^E \mod N$

- ▶ Alice wants to send a message M (a number, lets say between 1 and 20000) to Bob that only he can figure out.
- ▶ But any message can be intercepted (hacker: Carol!)
- ▶ How can Bob ensure privacy of messages?

Bob starts by choosing two large primes p, q

Also chooses numbers D and E such that $ED\equiv 1\mod \phi(pq)$ (i.e., $\phi(pq)\mid (ED-1)).$

- 1. Bob shouts out N = pq and E (encryptor!)...
- 2. But he keeps p, q and the decryptor D secret.
- 3. Alice wants to send message M < p, q that only Bob understands. What does she send?
- 4. Alice sends $X = M^E \mod N$
- 5. Bob can decrypt it by:

- ▶ Alice wants to send a message M (a number, lets say between 1 and 20000) to Bob that only he can figure out.
- ▶ But any message can be intercepted (hacker: Carol!)
- ▶ How can Bob ensure privacy of messages?

Bob starts by choosing two large primes p, q

Also chooses numbers D and E such that $ED \equiv 1 \mod \phi(pq)$ (i.e., $\phi(pq) \mid (ED-1)$).

- 1. Bob shouts out N = pq and E (encryptor!)...
- 2. But he keeps p, q and the decryptor D secret.
- 3. Alice wants to send message M < p, q that only Bob understands. What does she send?
- 4. Alice sends $X = M^E \mod N$
- 5. Bob can decrypt it by: $X^D = M^{ED} \mod N = M^{1+m\phi(pq)} \mod pq = M$ (by Euler's theorem)

RSA cryptography

Why does this work?

▶ Because Carol knows only X, N, E and has to solve $ED \equiv 1 \mod \phi(N)$ to obtain D after computing $\phi(N)$.

RSA cryptography

Why does this work?

- ▶ Because Carol knows only X, N, E and has to solve $ED \equiv 1 \mod \phi(N)$ to obtain D after computing $\phi(N)$.
- ▶ But there is no known fast way to get $\phi(N)$ from N.

RSA cryptography

Why does this work?

- ▶ Because Carol knows only X, N, E and has to solve $ED \equiv 1 \mod \phi(N)$ to obtain D after computing $\phi(N)$.
- ▶ But there is no known fast way to get $\phi(N)$ from N.
- ▶ Only known way is to factorize N and finding a poly-time algo for this is open!



Figure: Rivest, Shamir, Addleman, 1977 - Turing Award in 2002

Compare this to Diffie-hellman

Start with any finite cyclic group G and generator $g \in G$

- 1. Alice picks a random $a \in \mathbb{N}$ and sends g^a to Bob.
- 2. Bob picks a random $b \in \mathbb{N}$ and sends g^b to Alice.
- 3. Alice computes $(g^b)^a$ and Bob computes $(g^a)^b$.
- 4. Shared key is g^{ab} .
- ▶ Of course, we know modular logarithm we could do it!
- ▶ i.e., if $g^a = g'$ and g and g' are given, what is a?
- ► Called the discrete logarithm problem and it is also open!

Summary

What we covered in this course

- 1. Mathematical proofs and structures
- 2. Counting and combinatorics
- 3. Introduction to graph theory
- 4. Elements of group theory and applications to number theory

Summary: Till half time

► Mathematical proofs and structures

► Counting and Combinatorics

Summary: Till half time

- ► Mathematical proofs and structures
 - ▶ Propositions, proof techniques: contradiction, contrapositive
 - ▶ Induction: strong induction, well-ordering principle
 - ▶ Sets: finite and infinite sets, countable and uncountable sets
 - ▶ Functions: bijections (from e.g., $\mathbb{N} \times \mathbb{N} \to \mathbb{N}$), injections and surjections, Cantor's diagonalization technique
 - ▶ Relations: equivalence relations and partitions; partial orders, chains, anti-chains, lattices
- ► Counting and Combinatorics

Summary: Till half time

- ► Mathematical proofs and structures
 - ▶ Propositions, proof techniques: contradiction, contrapositive
 - ▶ Induction: strong induction, well-ordering principle
 - ▶ Sets: finite and infinite sets, countable and uncountable sets
 - ▶ Functions: bijections (from e.g., $\mathbb{N} \times \mathbb{N} \to \mathbb{N}$), injections and surjections, Cantor's diagonalization technique
 - Relations: equivalence relations and partitions; partial orders, chains, anti-chains, lattices
- ► Counting and Combinatorics
 - ▶ Basic counting principles, double counting
 - ightharpoonup Binomial theorem, permutations and combinations, Estimating n!
 - ▶ Recurrence relations and generating functions
 - ▶ Principle of Inclusion-Exclusion (PIE) and its applications.
 - ▶ Pigeon-Hole Principle (PHP) and its applications.
 - Some special numbers: Fibonacci, Catalan, Stirling of the second kind . . .
 - ▶ Introduction to Ramsey theory

Summary: Graph theory

Topics in Graph theory

- ▶ Basics: graphs, paths, cycles, walks, trails, . . .
- ▶ Cliques and independent sets.
- ▶ Graph representations, isomorphisms and automorphisms.
- ▶ Matchings: perfect, maximal and maximum.

Summary: Graph theory

Graph theory: Characterizations

- 1. Basics concepts and definitions.
- 2. Eulerian graphs: Using degrees of vertices.
- 3. Bipartite graphs: Using odd length cycles.
- 4. Connected components: Using cycles.
- 5. Maximum matchings: Using augmenting paths.
- 6. Perfect matchings in bipartite graphs: Using neighbour sets. Hall's theorem
- 7. Maximum matchings in bipartite graphs: Minimum vertex covers. Konig-Egervary's theorem
- 8. Stable matchings... and the Gale Shapley Algo

Summary: Abstract Algebra and Number theory

- ▶ Definition of an abstract group; basic properties
- ► Examples:
 - ▶ Invertible matrices, Symmetries of a regular polygon
 - ▶ Permutation groups, Graph automorphisms
 - \triangleright $(\mathbb{Z},+), (\mathbb{Z}_n,+n), (\mathbb{Z}_p,\times_p), \ldots$
- ► Abelian groups, Cyclic groups
- ▶ Group Isomorphisms and subgroups of a group.
- ▶ Order of a group and order of an element.
- ▶ Lagrange's theorem; corollaries and some applications

Summary: Abstract Algebra and Number theory

- ▶ Definition of an abstract group; basic properties
- ► Examples:
 - ▶ Invertible matrices, Symmetries of a regular polygon
 - Permutation groups, Graph automorphisms
 - \triangleright $(\mathbb{Z},+), (\mathbb{Z}_n,+n), (\mathbb{Z}_p,\times_p), \ldots$
- ► Abelian groups, Cyclic groups
- ▶ Group Isomorphisms and subgroups of a group.
- ▶ Order of a group and order of an element.
- ▶ Lagrange's theorem; corollaries and some applications

Applications to number theory and cryptography.

- ▶ Modular arithmetic
- ▶ Diffie-Hellman key exchange
- ► RSA cryptosystems.

Summary: Abstract Algebra and Number theory

- ▶ Definition of an abstract group; basic properties
- ► Examples:
 - ▶ Invertible matrices, Symmetries of a regular polygon
 - ▶ Permutation groups, Graph automorphisms
 - \triangleright $(\mathbb{Z},+), (\mathbb{Z}_n,+n), (\mathbb{Z}_p,\times_p), \ldots$
- ▶ Abelian groups, Cyclic groups
- ▶ Group Isomorphisms and subgroups of a group.
- Order of a group and order of an element.
- ▶ Lagrange's theorem; corollaries and some applications

Applications to number theory and cryptography.

- ► Modular arithmetic
- ▶ Diffie-Hellman key exchange
- ► RSA cryptosystems.

... the end.