

# CS 207: Discrete Structures

Instructor : S. Akshay

July 28, 2015

Lecture 05 – Comparing infinite sets via functions

## Chapter 2: Basic mathematical structures

### Sets and Functions

- ▶ Definition of a set, Russel's paradox, axioms of ZFC.
- ▶ Infinite sets and using functions to compare them.

# Hilbert's hotel



- ▶ Suppose there is a hotel with infinitely many rooms.
  - ▶ And suppose they are all full (like in IIT guest house).
1. Can you accomodate 1 or finitely many more guests, by shifting around the existing guests?
  2. What if infinitely many more guests arrive?
  3. What if infinitely many more trains with infinitely many more guests arrive and no room should be empty? (H.W)


# Functions

## Definition

Let  $A, B$  be two sets. A **function**  $f$  from  $A$  to  $B$  is an assignment of exactly one element of  $B$  to each element of  $A$ . i.e.,  $f : A \rightarrow B$  is a subset  $R$  of  $A \times B$  such that

- (i)  $\forall a \in A, \exists b \in B$  such that  $(a, b) \in R$ , and
- (ii) if  $(a, b) \in R$  and  $(a, c) \in R$ , then  $b = c$ .

# Comparing (finite and infinite) sets




- **Surjective or onto:**  $f : A \rightarrow B$  is surjective if  $\forall y \in B$ ,  $\exists x \in A$  such that  $f(x) = y$ .
- **Injective or 1-1:**  $f : A \rightarrow B$  is injective if  $\forall x, y \in A$ , if  $f(x) = f(y)$ , then  $x = y$ .
- **Bijective or 1-1 correspondence:** A function is bijective if it is surjective and injective.

If  $f$  is a bijection, then its inverse function exists and


$$f \circ f^{-1} = f^{-1} \circ f = \text{id}$$

# Comparing (finite and infinite) sets



- **Surjective or onto:**  $f: A \rightarrow B$  is surjective if  $\forall y \in B$ ,  $\exists x \in A$  such that  $f(x) = y$ .
  - **Injective or 1-1:**  $f: A \rightarrow B$  is injective if  $\forall x, y \in A$ , if  $f(x) = f(y)$ , then  $x = y$ .
  - **Bijective or 1-1 correspondence:** A function is bijective if it is surjective and injective.
1.  $f: \mathbb{Z} \rightarrow \mathbb{Z}$  such that  $f(x) = x^2$ .
  2.  $f: \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$  such that  $f(x) = x^2$ .

# Comparing (finite and infinite) sets



- **Surjective or onto:**  $f : A \rightarrow B$  is surjective if  $\forall y \in B$ ,  $\exists x \in A$  such that  $f(x) = y$ .
    - If  $A, B$  finite,  $|A| \geq |B|$
  - **Injective or 1-1:**  $f : A \rightarrow B$  is injective if  $\forall x, y \in A$ , if  $f(x) = f(y)$ , then  $x = y$ .
    - If  $A, B$  finite,  $|A| \leq |B|$
  - **Bijective or 1-1 correspondence:** A function is bijective if it is surjective and injective.
    - If  $A, B$  finite,  $|A| = |B|$
1.  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  such that  $f(x) = x^2$ .
  2.  $f : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$  such that  $f(x) = x^2$ .

# Properties of finite and infinite sets

## Relative notion of “size”

Thus, two finite/infinite sets have the same “size” iff there is a bijection between them.



# Properties of finite and infinite sets

## Similarities between finite and infinite sets

- ▶  $\exists$  **bij** from  $A$  to  $B$  and  $B$  to  $C$ , implies  $\exists$  **bij** from  $A$  to  $C$ .
- ▶  $\exists$  **bij** from  $A$  to  $B$ , then  $\exists$  **bij** from  $B$  to  $A$ .
- ▶  $\exists$  **surj** from  $A$  to  $B$  and  $\exists$  **surj**  $B$  to  $A$ , implies  $\exists$  **bij** from  $A$  to  $B$ .

# Properties of finite and infinite sets

## Similarities between finite and infinite sets

- ▶  $\exists$  **bij** from  $A$  to  $B$  and  $B$  to  $C$ , implies  $\exists$  **bij** from  $A$  to  $C$ .
- ▶  $\exists$  **bij** from  $A$  to  $B$ , then  $\exists$  **bij** from  $B$  to  $A$ .
- ▶ (**Schröder-Bernstein Theorem**):  $\exists$  **surj** from  $A$  to  $B$  and  $\exists$  **surj**  $B$  to  $A$ , implies  $\exists$  **bij** from  $A$  to  $B$ . (H.W: Read this!)

# Properties of finite and infinite sets

## Similarities between finite and infinite sets

- ▶  $\exists$  **bij** from  $A$  to  $B$  and  $B$  to  $C$ , implies  $\exists$  **bij** from  $A$  to  $C$ .
- ▶  $\exists$  **bij** from  $A$  to  $B$ , then  $\exists$  **bij** from  $B$  to  $A$ .
- ▶ (**Schröder-Bernstein Theorem**;)  $\exists$  **surj** from  $A$  to  $B$  and  $\exists$  **surj**  $B$  to  $A$ , implies  $\exists$  **bij** from  $A$  to  $B$ . (H.W: Read this!)

## Differences between finite and infinite sets

- ▶ For finite sets, if  $A$  is a set and  $b \notin A$ , then  $|A| \neq |A \cup \{b\}|$ .

# Properties of finite and infinite sets

## Similarities between finite and infinite sets

- ▶  $\exists$  **bij** from  $A$  to  $B$  and  $B$  to  $C$ , implies  $\exists$  **bij** from  $A$  to  $C$ .
- ▶  $\exists$  **bij** from  $A$  to  $B$ , then  $\exists$  **bij** from  $B$  to  $A$ .
- ▶ (**Schröder-Bernstein Theorem**;)  $\exists$  **surj** from  $A$  to  $B$  and  $\exists$  **surj**  $B$  to  $A$ , implies  $\exists$  **bij** from  $A$  to  $B$ . (H.W: Read this!)

## Differences between finite and infinite sets

- ▶ For finite sets, if  $A$  is a set and  $b \notin A$ , then  $|A| \neq |A \cup \{b\}|$ .
- ▶ What about infinite sets?

# Properties of finite and infinite sets

## Similarities between finite and infinite sets

- ▶  $\exists$  **bij** from  $A$  to  $B$  and  $B$  to  $C$ , implies  $\exists$  **bij** from  $A$  to  $C$ .
- ▶  $\exists$  **bij** from  $A$  to  $B$ , then  $\exists$  **bij** from  $B$  to  $A$ .
- ▶ (**Schröder-Bernstein Theorem**;)  $\exists$  **surj** from  $A$  to  $B$  and  $\exists$  **surj**  $B$  to  $A$ , implies  $\exists$  **bij** from  $A$  to  $B$ . (H.W: Read this!)

## Differences between finite and infinite sets

- ▶ For finite sets, if  $A$  is a set and  $b \notin A$ , then  $|A| \neq |A \cup \{b\}|$ .
- ▶ What about infinite sets?

## Theorem

Let  $A$  be a set and  $b \notin A$ . Then  $A$  is infinite iff there is a bijection from  $A$  to  $A \cup \{b\}$ .

# Properties of finite and infinite sets

## Similarities between finite and infinite sets

- ▶  $\exists$  **bij** from  $A$  to  $B$  and  $B$  to  $C$ , implies  $\exists$  **bij** from  $A$  to  $C$ .
- ▶  $\exists$  **bij** from  $A$  to  $B$ , then  $\exists$  **bij** from  $B$  to  $A$ .
- ▶ (**Schröder-Bernstein Theorem**):  $\exists$  **surj** from  $A$  to  $B$  and  $\exists$  **surj**  $B$  to  $A$ , implies  $\exists$  **bij** from  $A$  to  $B$ . (H.W: Read this!)

## Differences between finite and infinite sets

- ▶ For finite sets, if  $A$  is a set and  $b \notin A$ , then  $|A| \neq |A \cup \{b\}|$ .
- ▶ What about infinite sets?

## Theorem

Let  $A$  be a set and  $b \notin A$ . Then  $A$  is infinite iff there is a bijection from  $A$  to  $A \cup \{b\}$ .

Proof: essentially Hilbert's hotel but be careful...

# Comparing infinite sets using functions

## Theorem

There is a bijection from  $\mathbb{Z}$  to  $\mathbb{N}$ .

Proof: Hilbert's hotel argument. But how do you formalize it?

# Comparing infinite sets using functions

## Theorem

There is a bijection from  $\mathbb{Z}$  to  $\mathbb{N}$ .

Proof: Hilbert's hotel argument. But how do you formalize it?

$$f(x) = \begin{cases} -2x & \text{if } x \leq 0 \\ 2x - 1 & \text{else} \end{cases}$$



# Comparing infinite sets using functions

## Theorem

There is a bijection from  $\mathbb{Z}$  to  $\mathbb{N}$ .

Proof: Hilbert's hotel argument. But how do you formalize it?

$$f(x) = \begin{cases} -2x & \text{if } x \leq 0 \\ 2x - 1 & \text{else} \end{cases}$$

## Some questions...

- ▶ Is there a bijection between  $\mathbb{N} \times \mathbb{N}$  to  $\mathbb{N}$ ?
- ▶ Is there a bijection between  $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$  to  $\mathbb{N}$ ?
- ▶ Is there a bijection from  $\mathbb{Q}$  to  $\mathbb{N}$ ?
- ▶ Is there a bijection from the set of all subsets of  $\mathbb{N}$  to  $\mathbb{N}$ ?
- ▶ Is there a bijection from  $\mathbb{R}$  to  $\mathbb{N}$ ?