

# CS 207: Discrete Structures

## Lecture 12 – Basic counting techniques

Aug 13, 2015

# Course Outline

## Topics to be covered

1. Proofs and structures
2. Counting and combinatorics
  - ▶ Basics of counting
  - ▶ Subsets, partitions, Permutations and combinations
  - ▶ Recurrence relations and generating functions
  - ▶ Principle of Inclusion and Exclusion and its applications
  - ▶ Pigeonhole Principle and its extensions
3. Elements of graph theory
4. Introduction to abstract algebra and number theory

## Two examples

### Letters and envelopes

Given  $n$  letters and  $n$  addressed envelopes, in how many ways can letters be placed in envelopes so that none of them is in its correctly addressed envelope.

- ▶ Total no. of ways of putting  $n$  letters in  $n$  envelopes is

## Two examples

### Letters and envelopes

Given  $n$  letters and  $n$  addressed envelopes, in how many ways can letters be placed in envelopes so that none of them is in its correctly addressed envelope.

- ▶ Total no. of ways of putting  $n$  letters in  $n$  envelopes is  $n!$ .
- ▶ We will see that the fraction which are incorrectly addressed is close to  $1/e$ , where  $e = 2.71828\dots$  is the base of the natural logarithm!

# Two examples

## Letters and envelopes

Given  $n$  letters and  $n$  addressed envelopes, in how many ways can letters be placed in envelopes so that none of them is in its correctly addressed envelope.

- ▶ Total no. of ways of putting  $n$  letters in  $n$  envelopes is  $n!$ .
- ▶ We will see that the fraction which are incorrectly addressed is close to  $1/e$ , where  $e = 2.71828\dots$  is the base of the natural logarithm!

## A Ramsey game

- ▶ There are six points on board and two colored chalks.

# Two examples

## Letters and envelopes

Given  $n$  letters and  $n$  addressed envelopes, in how many ways can letters be placed in envelopes so that none of them is in its correctly addressed envelope.

- ▶ Total no. of ways of putting  $n$  letters in  $n$  envelopes is  $n!$ .
- ▶ We will see that the fraction which are incorrectly addressed is close to  $1/e$ , where  $e = 2.71828\dots$  is the base of the natural logarithm!

## A Ramsey game

- ▶ There are six points on board and two colored chalks.
- ▶ Divide class into 2 groups. Group 1 draws white lines and Group 2 draws blue lines between points.

# Two examples

## Letters and envelopes

Given  $n$  letters and  $n$  addressed envelopes, in how many ways can letters be placed in envelopes so that none of them is in its correctly addressed envelope.

- ▶ Total no. of ways of putting  $n$  letters in  $n$  envelopes is  $n!$ .
- ▶ We will see that the fraction which are incorrectly addressed is close to  $1/e$ , where  $e = 2.71828\dots$  is the base of the natural logarithm!

## A Ramsey game

- ▶ There are six points on board and two colored chalks.
- ▶ Divide class into 2 groups. Group 1 draws white lines and Group 2 draws blue lines between points.
- ▶ You lose if you are first to draw a triangle of your color.

# Two examples

## Letters and envelopes

Given  $n$  letters and  $n$  addressed envelopes, in how many ways can letters be placed in envelopes so that none of them is in its correctly addressed envelope.

- ▶ Total no. of ways of putting  $n$  letters in  $n$  envelopes is  $n!$ .
- ▶ We will see that the fraction which are incorrectly addressed is close to  $1/e$ , where  $e = 2.71828\dots$  is the base of the natural logarithm!

## A Ramsey game

- ▶ There are six points on board and two colored chalks.
- ▶ Divide class into 2 groups. Group 1 draws white lines and Group 2 draws blue lines between points.
- ▶ You lose if you are first to draw a triangle of your color.
- ▶ Can you ever have a draw?



## Basic counting principles illustrated...

- ▶ How many reflexive relations are there on a set  $A$  of size  $n$ ?
  - ▶ Reflexive relations are ordered pairs of which there are  $n^2$ .
  - ▶ Of these, all  $n$  pairs of  $(a, a)$  have to be present.
  - ▶ Of the remaining, we can choose any of them to be in or out.
  - ▶ there are  $n^2 - n$  of them, so  $2^{n^2 - n}$  of them.
  - ▶ We used the so-called “product principle”...

## Basic counting principles illustrated...

- ▶ How many reflexive relations are there on a set  $A$  of size  $n$ ?
  - ▶ Reflexive relations are ordered pairs of which there are  $n^2$ .
  - ▶ Of these, all  $n$  pairs of  $(a, a)$  have to be present.
  - ▶ Of the remaining, we can choose any of them to be in or out.
  - ▶ there are  $n^2 - n$  of them, so  $2^{n^2 - n}$  of them.
  - ▶ We used the so-called “product principle”...

### The product principle

If there are  $n_1$  ways of doing something and  $n_2$  ways of doing another thing, then there are  $n_1 \cdot n_2$  ways of performing both actions.

## Basic counting principles illustrated...

- ▶ How many reflexive relations are there on a set  $A$  of size  $n$ ?
- ▶ How many functions are there from a set of size  $n$  to itself?
- ▶ If 20 teams play in the IITB-premier league and every game has a winner/loser and loser is always eliminated. How many games are played before a champion is chosen?
- ▶ How many subsets does a set  $A$  of  $n$  elements have?

## Basic counting principles illustrated...

- ▶ How many reflexive relations are there on a set  $A$  of size  $n$ ?
- ▶ How many functions are there from a set of size  $n$  to itself?
- ▶ How many subsets does a set  $A$  of  $n$  elements have?
  - ▶ **Product principle**: two choices for each element, hence  $2 \cdot 2 \cdots 2 \cdot 2$  ( $n$ -times).
  - ▶ **Bijection**: between  $\mathcal{P}(X)$  and  $n$ -length sequences over  $\{0, 1\}$  (characteristic vector).
  - ▶ **Induction**: Since we already know the answer!
  - ▶ **Recurrence**:  $F(n) = 2 \cdot F(n-1), F(0) = 1$ . solve it?
  - ▶ **Sum principle**: Subsets of size 0 + subsets of size 1 + ... + subsets of size  $n$  = Total number of subsets.

## Basic counting principles illustrated...

- ▶ How many reflexive relations are there on a set  $A$  of size  $n$ ?
- ▶ How many functions are there from a set of size  $n$  to itself?
- ▶ How many subsets does a set  $A$  of  $n$  elements have?
  - ▶ **Product principle**: two choices for each element, hence  $2 \cdot 2 \cdots 2 \cdot 2$  ( $n$ -times).
  - ▶ **Bijection**: between  $\mathcal{P}(X)$  and  $n$ -length sequences over  $\{0, 1\}$  (characteristic vector).
  - ▶ **Induction**: Since we already know the answer!
  - ▶ **Recurrence**:  $F(n) = 2 \cdot F(n-1)$ ,  $F(0) = 1$ . solve it?
  - ▶ **Sum principle**: Subsets of size 0 + subsets of size 1 + ... + subsets of size  $n$  = Total number of subsets.

### Sum Principle

If something can be done in  $n_1$  **or**  $n_2$  ways such that none of the  $n_1$  ways is the same as any of the  $n_2$  ways, then the total number of ways to do this is  $n_1 + n_2$ .

## Basic counting principles illustrated...

- ▶ How many reflexive relations are there on a set  $A$  of size  $n$ ?
- ▶ How many functions are there from a set of size  $n$  to itself?
- ▶ How many subsets does a set  $A$  of  $n$  elements have?
  - ▶ **Product principle**: two choices for each element, hence  $2 \cdot 2 \cdots 2 \cdot 2$  ( $n$ -times).
  - ▶ **Bijection**: between  $\mathcal{P}(X)$  and  $n$ -length sequences over  $\{0, 1\}$  (characteristic vector).
  - ▶ **Induction**: Since we already know the answer!
  - ▶ **Recurrence**:  $F(n) = 2 \cdot F(n-1)$ ,  $F(0) = 1$ . solve it?
  - ▶ **Sum principle**: Subsets of size 0 + subsets of size 1 + ... + subsets of size  $n$  = Total number of subsets.
- ▶ But, how many subsets of size  $k$  does a set of  $n$  elements have?

## Basic counting principles illustrated...

- ▶ How many reflexive relations are there on a set  $A$  of size  $n$ ?
- ▶ How many functions are there from a set of size  $n$  to itself?
- ▶ How many subsets does a set  $A$  of  $n$  elements have?
  - ▶ **Product principle**: two choices for each element, hence  $2 \cdot 2 \cdots 2 \cdot 2$  ( $n$ -times).
  - ▶ **Bijection**: between  $\mathcal{P}(X)$  and  $n$ -length sequences over  $\{0, 1\}$  (characteristic vector).
  - ▶ **Induction**: Since we already know the answer!
  - ▶ **Recurrence**:  $F(n) = 2 \cdot F(n-1)$ ,  $F(0) = 1$ . solve it?
  - ▶ **Sum principle**: Subsets of size 0 + subsets of size 1 + ... + subsets of size  $n$  = Total number of subsets.
- ▶ But, how many subsets of size  $k$  does a set of  $n$  elements have? This number, denoted  $\binom{n}{k}$ , is called a **binomial coefficient**.

## Basic counting principles illustrated...

- ▶ How many reflexive relations are there on a set  $A$  of size  $n$ ?
- ▶ How many functions are there from a set of size  $n$  to itself?
- ▶ How many subsets does a set  $A$  of  $n$  elements have?
  - ▶ **Product principle**: two choices for each element, hence  $2 \cdot 2 \cdots 2 \cdot 2$  ( $n$ -times).
  - ▶ **Bijection**: between  $\mathcal{P}(X)$  and  $n$ -length sequences over  $\{0, 1\}$  (characteristic vector).
  - ▶ **Induction**: Since we already know the answer!
  - ▶ **Recurrence**:  $F(n) = 2 \cdot F(n-1)$ ,  $F(0) = 1$ . solve it?
  - ▶ **Sum principle**: Subsets of size 0 + subsets of size 1 + ... + subsets of size  $n$  = Total number of subsets.
- ▶ But, how many subsets of size  $k$  does a set of  $n$  elements have? This number, denoted  $\binom{n}{k}$ , is called a **binomial coefficient**.
- ▶ We all know(?) that  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$ . Prove it!



# Permutations and combinations

Binomial Coefficients. Let  $n, k$  be integers s.t.,  $n \geq k \geq 0$ .

How many subsets of size  $k$  does a set of  $n$  elements have? This number, denoted  $\binom{n}{k}$ , is called a **binomial coefficient**.

One proof of  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$  is as follows:

# Permutations and combinations

Binomial Coefficients. Let  $n, k$  be integers s.t.,  $n \geq k \geq 0$ .

How many subsets of size  $k$  does a set of  $n$  elements have? This number, denoted  $\binom{n}{k}$ , is called a **binomial coefficient**.

One proof of  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$  is as follows: Let us count the number of “**ordered**” subsets of size  $k$ .

# Permutations and combinations

Binomial Coefficients. Let  $n, k$  be integers s.t.,  $n \geq k \geq 0$ .

How many subsets of size  $k$  does a set of  $n$  elements have? This number, denoted  $\binom{n}{k}$ , is called a **binomial coefficient**.

One proof of  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$  is as follows: Let us count the number of “**ordered**” subsets of size  $k$ .

- ▶ No. of **ordered** subsets of size  $k = n \cdot (n-1) \cdots (n-k+1)$ .

# Permutations and combinations

**Binomial Coefficients.** Let  $n, k$  be integers s.t.,  $n \geq k \geq 0$ .

How many subsets of size  $k$  does a set of  $n$  elements have? This number, denoted  $\binom{n}{k}$ , is called a **binomial coefficient**.

One proof of  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$  is as follows: Let us count the number of “**ordered**” subsets of size  $k$ .

- ▶ No. of **ordered** subsets of size  $k = n \cdot (n-1) \cdots (n-k+1)$ .
- ▶ No. of **ordered** subsets of size  $k = (\text{no. of unordered subsets}) \times (\text{no. of ways to order them}) = \binom{n}{k} \times k!$ .

# Permutations and combinations

**Binomial Coefficients.** Let  $n, k$  be integers s.t.,  $n \geq k \geq 0$ .

How many subsets of size  $k$  does a set of  $n$  elements have? This number, denoted  $\binom{n}{k}$ , is called a **binomial coefficient**.

One proof of  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$  is as follows: Let us count the number of “**ordered**” subsets of size  $k$ .

- ▶ No. of **ordered** subsets of size  $k = n \cdot (n-1) \cdots (n-k+1)$ .
- ▶ No. of **ordered** subsets of size  $k = (\text{no. of unordered subsets}) \times (\text{no. of ways to order them}) = \binom{n}{k} \times k!$ .
- ▶ Equate them! Principle of **double counting**.
  - ▶ if you can't count something, count something else and count it twice over!

# Permutations and combinations

**Binomial Coefficients.** Let  $n, k$  be integers s.t.,  $n \geq k \geq 0$ .

How many subsets of size  $k$  does a set of  $n$  elements have? This number, denoted  $\binom{n}{k}$ , is called a **binomial coefficient**.

One proof of  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$  is as follows: Let us count the number of “**ordered**” subsets of size  $k$ .

- ▶ No. of **ordered** subsets of size  $k = n \cdot (n-1) \cdots (n-k+1)$ .
- ▶ No. of **ordered** subsets of size  $k = (\text{no. of unordered subsets}) \times (\text{no. of ways to order them}) = \binom{n}{k} \times k!$ .
- ▶ Equate them! Principle of **double counting**.

## Permutations and combinations

- ▶ No. of  $k$ -size subsets of set of size  $n = \text{No. of } k\text{-combinations of a set of } n \text{ (distinct) elements} = \binom{n}{k}$ .
- ▶ No. of  $k$ -size ordered subsets of set of size  $n = \text{No. of } k\text{-permutations of a set of } n \text{ (distinct) elements}$ .

## Simple examples to illustrate “double counting”

Prove the following identities (by only using double counting!)

$$1. \sum_{k=0}^n \binom{n}{k} = 2^n.$$

$$2. \binom{n}{k} = \binom{n}{n-k}.$$

$$3. k \binom{n}{k} = n \binom{n-1}{k-1}$$

$$4. \binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$$

## Simple examples to illustrate “double counting”

Prove the following identities (by only using double counting!)

$$1. \sum_{k=0}^n \binom{n}{k} = 2^n.$$

$$2. \binom{n}{k} = \binom{n}{n-k}.$$

$$3. k \binom{n}{k} = n \binom{n-1}{k-1}$$

$$4. \binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$$

The latter two are in fact recursive definitions for  $\binom{n}{k}$ . What are the boundary conditions?



## A more interesting example with double counting

### Handshake Lemma

At a meeting with  $n$  people, the number of people who shake hands an odd number of times is even.

What will you count here?

## A more interesting example with double counting

### Handshake Lemma

At a meeting with  $n$  people, the number of people who shake hands an odd number of times is even.

Proof in six steps:

1. Define a relation  $R$ :  $iRj$  if  $i$  and  $j$  shook hands.
2. Is this relation symmetric (trans/refl.)? Draw its graph.

## A more interesting example with double counting

### Handshake Lemma

At a meeting with  $n$  people, the number of people who shake hands an odd number of times is even.

Proof in six steps:

1. Define a relation  $R$ :  $iRj$  if  $i$  and  $j$  shook hands.
2. Is this relation symmetric (trans/refl.)? Draw its graph.
3. Let  $m_i$  be the number of times  $i$  shakes hands.

## A more interesting example with double counting

### Handshake Lemma

At a meeting with  $n$  people, the number of people who shake hands an odd number of times is even.

Proof in six steps:

1. Define a relation  $R$ :  $iRj$  if  $i$  and  $j$  shook hands.
2. Is this relation symmetric (trans/refl.)? Draw its graph.
3. Let  $m_i$  be the number of times  $i$  shakes hands. i.e.,  $m_i$  is the number of directed edges going out from  $i$ .

## A more interesting example with double counting

### Handshake Lemma

At a meeting with  $n$  people, the number of people who shake hands an odd number of times is even.

Proof in six steps:

1. Define a relation  $R$ :  $iRj$  if  $i$  and  $j$  shook hands.
2. Is this relation symmetric (trans/refl.)? Draw its graph.
3. Let  $m_i$  be the number of times  $i$  shakes hands. i.e.,  $m_i$  is the number of directed edges going out from  $i$ .
4. Therefore, **Total no. of directed edges**  $= \sum_i m_i$ .

## A more interesting example with double counting

### Handshake Lemma

At a meeting with  $n$  people, the number of people who shake hands an odd number of times is even.

Proof in six steps:

1. Define a relation  $R$ :  $iRj$  if  $i$  and  $j$  shook hands.
2. Is this relation symmetric (trans/refl.)? Draw its graph.
3. Let  $m_i$  be the number of times  $i$  shakes hands. i.e.,  $m_i$  is the number of directed edges going out from  $i$ .
4. Therefore, **Total no. of directed edges**  $= \sum_i m_i$ .
5. But now, let  $X$  be the total number of handshakes. Clearly this is an integer.

## A more interesting example with double counting

### Handshake Lemma

At a meeting with  $n$  people, the number of people who shake hands an odd number of times is even.

Proof in six steps:

1. Define a relation  $R$ :  $iRj$  if  $i$  and  $j$  shook hands.
2. Is this relation symmetric (trans/refl.)? Draw its graph.
3. Let  $m_i$  be the number of times  $i$  shakes hands. i.e.,  $m_i$  is the number of directed edges going out from  $i$ .
4. Therefore, **Total no. of directed edges**  $= \sum_i m_i$ .
5. But now, let  $X$  be the total number of handshakes. Clearly this is an integer. **Total no. of directed edges**  $= 2 \cdot X$ .

## A more interesting example with double counting

### Handshake Lemma

At a meeting with  $n$  people, the number of people who shake hands an odd number of times is even.

Proof in six steps:

1. Define a relation  $R$ :  $iRj$  if  $i$  and  $j$  shook hands.
2. Is this relation symmetric (trans/refl.)? Draw its graph.
3. Let  $m_i$  be the number of times  $i$  shakes hands. i.e.,  $m_i$  is the number of directed edges going out from  $i$ .
4. Therefore, **Total no. of directed edges**  $= \sum_i m_i$ .
5. But now, let  $X$  be the total number of handshakes. Clearly this is an integer. **Total no. of directed edges**  $= 2 \cdot X$ .
6. This implies,  **$\sum_i m_i = 2 \cdot X$** . Which means that number of  $i$  such that  $m_i$  is odd is even!



# What about partitions?

We have looked at:

- ▶ number of subsets of a set (combinations)
- ▶ number of ordered subsets of a set (permutations)

What about number of partitions of a set of  $n$  distinct elements, denoted  $B_n$ ?

# What about partitions?

We have looked at:

- ▶ number of subsets of a set (combinations)
- ▶ number of ordered subsets of a set (permutations)

What about number of partitions of a set of  $n$  distinct elements, denoted  $B_n$ ?

- ▶ Same as number of equivalence relations!
- ▶ What are  $B_3$ ,  $B_2$ ,  $B_1$ ? What about  $B_0$ ?
- ▶ What about  $B_n$  in general?