

CS 207: Discrete Structures

Instructor : S. Akshay

Aug 3, 2015

Lecture 07 – Uncountable sets, relations, equivalence relation

Recap: Countable and countably infinite sets

Definition

- ▶ For a given set C , if there is a bijection from C to \mathbb{N} , then C is called **countably infinite**. A set is **countable** if it is finite or countably infinite.
- ▶ Examples: \mathbb{Z} , $\mathbb{N} \times \mathbb{N}$, \mathbb{Q} .
- ▶ Properties: Closure under (countable) unions and (finite) products.

Recap: Countable and countably infinite sets

Definition

- ▶ For a given set C , if there is a bijection from C to \mathbb{N} , then C is called **countably infinite**. A set is **countable** if it is finite or countably infinite.
- ▶ Examples: \mathbb{Z} , $\mathbb{N} \times \mathbb{N}$, \mathbb{Q} .
- ▶ Properties: Closure under (countable) unions and (finite) products.

Are all sets countable?

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	✓	×	×	×	...
$f(1)$	✓	×	✓	✓	...
$f(2)$	×	×	×	×	...
$f(3)$	×	✓	×	✓	...

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	✓ ×	×	×	×	...
$f(1)$	✓	× ✓	✓	✓	...
$f(2)$	×	×	× ✓	×	...
$f(3)$	×	✓	×	✓ ×	...

- Consider the set $S \subseteq \mathbb{N}$ obtained by switching the diagonal elements, i.e., $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$.

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	✓ ×	×	×	×	...
$f(1)$	✓	× ✓	✓	✓	...
$f(2)$	×	×	× ✓	×	...
$f(3)$	×	✓	×	✓ ×	...

- ▶ Consider the set $S \subseteq \mathbb{N}$ obtained by switching the diagonal elements, i.e., $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$.
- ▶ As f is bij, $\exists j \in \mathbb{N}, f(j) = S$.

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	$\checkmark \times$	\times	\times	\times	...
$f(1)$	\checkmark	$\not\checkmark \checkmark$	\checkmark	\checkmark	...
$f(2)$	\times	\times	$\not\checkmark \checkmark$	\times	...
$f(3)$	\times	\checkmark	\times	$\not\checkmark \times$...

- ▶ Consider the set $S \subseteq \mathbb{N}$ obtained by switching the diagonal elements, i.e., $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$.
- ▶ As f is bij, $\exists j \in \mathbb{N}, f(j) = S$.
- ▶ But S and $f(j)$ differ at position j .

Comparing \mathbb{N} and set of all subsets of \mathbb{N}

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

Proof by contradiction: Suppose there is such a bijection, say f . This would imply that each $i \in \mathbb{N}$ maps to some set $f(i) \subseteq \mathbb{N}$.

	0	1	2	3	...
$f(0)$	✓ ×	×	×	×	...
$f(1)$	✓	× ✓	✓	✓	...
$f(2)$	×	×	× ✓	×	...
$f(3)$	×	✓	×	✓ ×	...

- ▶ Consider the set $S \subseteq \mathbb{N}$ obtained by switching the diagonal elements, i.e., $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$.
- ▶ As f is bij, $\exists j \in \mathbb{N}, f(j) = S$.
- ▶ But S and $f(j)$ differ at position j .
- ▶ Thus, $S \neq f(j)$ which is a contradiction! □

Cantor's diagonalization




Figure : Cantor and Russell

- $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$ is like the one from Russell's paradox.

Cantor's diagonalization




Figure : Cantor and Russell

- ▶ $S = \{i \in \mathbb{N} \mid i \notin f(i)\}$ is like the one from Russell's paradox.
- ▶ If $\exists j \in \mathbb{N}$ such that $f(j) = S$, then we have a contradiction.
 - ▶ If $j \in S$, then $j \notin f(j) = S$.
 - ▶ If $j \notin S$, then $j \in f(j) = S$.

Cantor's diagonalization




Figure : Cantor and Russell

In fact, using diagonalization Cantor showed that...

- ▶ There cannot be a bijection between **any** set and its power set (i.e., its set of subsets). (H.W)
- ▶ So there is an infinite hierarchy of “larger” infinities...

Cantor's diagonalization




Figure : Cantor and Russell

In fact, using diagonalization Cantor showed that...

- ▶ There cannot be a bijection between **any** set and its power set (i.e., its set of subsets). **(H.W)**
- ▶ So there is an infinite hierarchy of “larger” infinities...
- ▶ There is no bijection from \mathbb{R} to \mathbb{N} **(H.W)**. Moreover, there is a bijection from \mathbb{R} to set of subsets of \mathbb{N} .

Cantor's diagonalization




Figure : Cantor and Russell

In fact, using diagonalization Cantor showed that...

- ▶ There cannot be a bijection between **any** set and its power set (i.e., its set of subsets). **(H.W)**
- ▶ So there is an infinite hierarchy of “larger” infinities...
- ▶ There is no bijection from \mathbb{R} to \mathbb{N} **(H.W)**. Moreover, there is a bijection from \mathbb{R} to set of subsets of \mathbb{N} .
- ▶ What about **the set of all sets**??

One infinity is “strictly” bigger than another!

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

One infinity is “strictly” bigger than another!

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

- ▶ But, there is a surjection from set of all subsets of \mathbb{N} to \mathbb{N} .
- ▶ Thus, the “size” of $\mathcal{P}(\mathbb{N})$ is strictly greater than \mathbb{N} !

One infinity is “strictly” bigger than another!

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

- ▶ But, there is a surjection from set of all subsets of \mathbb{N} to \mathbb{N} .
- ▶ Thus, the “size” of $\mathcal{P}(\mathbb{N})$ is strictly greater than \mathbb{N} !

Can there be some set whose “size” is in between the two?

One infinity is “strictly” bigger than another!

Theorem (Cantor, 1891)

There is no bijection between \mathbb{N} and the set of all subsets of \mathbb{N} .

- ▶ But, there is a surjection from set of all subsets of \mathbb{N} to \mathbb{N} .
- ▶ Thus, the “size” of $\mathcal{P}(\mathbb{N})$ is strictly greater than \mathbb{N} !

Can there be some set whose “size” is in between the two?

Cantor’s Continuum hypothesis

There is no set whose “cardinality” is strictly between \mathbb{N} and $\mathcal{P}(\mathbb{N})$ (i.e., between naturals and reals).




Figure : 1st of Hilbert’s 23 problems for the 20th century in 1900.

What did the world think about these proofs (in 1890s?)



(a) Kronecker



(b) Poincaré



(c) Theologians

- ▶ **Kronecker:** Only constructive proofs are proofs! “Scientific Charlatan”, “Corruptor of youth”!
- ▶ **Poincaré:** Set theory is a “disease” from which mathematics will be cured.
- ▶ **Christian Theologians:** God=Uniqueness of an absolute infinity. So, what is all this different infinities...?!

What did the world think about these proofs (in 1890s?)



(a) Kronecker



(b) Poincaré



(c) Theologians

- ▶ **Kronecker:** Only constructive proofs are proofs! “Scientific Charlatan”, “Corruptor of youth”!
- ▶ **Poincaré:** Set theory is a “disease” from which mathematics will be cured.
- ▶ **Christian Theologians:** God=Uniqueness of an absolute infinity. So, what is all this different infinities...?!
- ▶ **Hilbert:** No one can expel us from the paradise that Cantor has created for us.

Summary and moving on...

- ▶ Finite and infinite sets.
- ▶ Using functions to compare sets: focus on bijections.
- ▶ Countable, countably infinite and uncountable sets.
- ▶ Cantor's diagonalization argument (A new powerful proof technique!).

Summary and moving on...

- ▶ Finite and infinite sets.
- ▶ Using functions to compare sets: focus on bijections.
- ▶ Countable, countably infinite and uncountable sets.
- ▶ Cantor's diagonalization argument (A new powerful proof technique!).

Next: Basic Mathematical Structures – Relations

Relations

Definition: Function

Let A, B be two sets. A **function** f from A to B is a subset R of $A \times B$ such that

- (i) $\forall a \in A, \exists b \in B$ such that $(a, b) \in R$, and
 - (ii) if $(a, b) \in R$ and $(a, c) \in R$, then $b = c$.
- Now, suppose A is the set of all Btech students and B is the set of all courses. Clearly, we can assign to each student the set of courses he/she is taking. Is this a function?

Relations

Definition: Function

Let A, B be two sets. A **function** f from A to B is a subset R of $A \times B$ such that

- (i) $\forall a \in A, \exists b \in B$ such that $(a, b) \in R$, and
- (ii) if $(a, b) \in R$ and $(a, c) \in R$, then $b = c$.

- ▶ Now, suppose A is the set of all Btech students and B is the set of all courses. Clearly, we can assign to each student the set of courses he/she is taking. Is this a function?
- ▶ By removing the two extra assumptions in the defn, we get:

Definition: Relation

- ▶ A **relation** R from A to B is a subset of $A \times B$. If $(a, b) \in R$, we also write this as $a R b$.
- ▶ Thus, a relation is a way to relate the elements of two (not necessarily different) sets.

Examples and representations of relations

We write $R(A, B)$ for a relation from A to B and just $R(A)$ if $A = B$. Also if A is clear from context, we just write R .

Examples of relations

- ▶ All functions are relations.
- ▶ $R_1(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a - b \text{ is even} \}$.
- ▶ $R_2(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$.
- ▶ Let S be a set, $R_3(\mathcal{P}(S)) = \{(A, B) \mid A, B \subseteq S, A \subseteq B\}$.
- ▶ Relational databases are practical examples.

Examples and representations of relations

We write $R(A, B)$ for a relation from A to B and just $R(A)$ if $A = B$. Also if A is clear from context, we just write R .

Examples of relations

- ▶ All functions are relations.
- ▶ $R_1(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a - b \text{ is even} \}$.
- ▶ $R_2(\mathbb{Z}) = \{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$.
- ▶ Let S be a set, $R_3(\mathcal{P}(S)) = \{(A, B) \mid A, B \subseteq S, A \subseteq B\}$.
- ▶ Relational databases are practical examples.

Representations of a relation from A to B .

- ▶ As a set of **ordered pairs of elements**, i.e., subset of $A \times B$.
- ▶ As a **directed graph**.
- ▶ As a **(database) table**.

Use of relations

Practical application in relational databases: IMDB, university records, etc.

Use of relations

Practical application in relational databases: IMDB, university records, etc.

But, why study relations in this course?

Use of relations

Practical application in relational databases: IMDB, university records, etc.

But, why study relations in this course?

- ▶ Functions were special kinds of relations that were useful to compare sets.
- ▶ Are there other **special** relations? What are they useful for?

Use of relations

Practical application in relational databases: IMDB, university records, etc.

But, why study relations in this course?

- ▶ Functions were special kinds of relations that were useful to compare sets.
- ▶ Are there other **special** relations? What are they useful for?
 - ▶ Equivalence relations
 - ▶ Partial orders

Partitions of a set – grouping “like” elements

Examples

- ▶ Natural numbers are partitioned into even and odd.
- ▶ This class is partitioned into set of all those who are taking the **same** set of courses.

How do you define a partition?

Partitions of a set – grouping “like” elements

Examples

- ▶ Natural numbers are partitioned into even and odd.
- ▶ This class is partitioned into set of all those who are taking the **same** set of courses.

How do you define a partition?

Definition

A partition of a set S is a set P of its subsets such that

- ▶ if $S' \in P$, then $S' \neq \emptyset$.
- ▶ $\bigcup_{S' \in P} S' = S$: its union covers entire set S .
- ▶ If $S_1, S_2 \in P$, then $S_1 \cap S_2 = \emptyset$: sets are disjoint.

Partitions of a set – grouping “like” elements

Examples

- ▶ Natural numbers are partitioned into even and odd.
- ▶ This class is partitioned into set of all those who are taking the **same** set of courses.

How do you define a partition?

Definition

A partition of a set S is a set P of its subsets such that

- ▶ if $S' \in P$, then $S' \neq \emptyset$.
- ▶ $\bigcup_{S' \in P} S' = S$: its union covers entire set S .
- ▶ If $S_1, S_2 \in P$, then $S_1 \cap S_2 = \emptyset$: sets are disjoint.

Can you think of two trivial partitions that any set must have?

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same set of courses...

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same set of courses...
- ▶ But, this sounds like relation, right? Which one?

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same set of courses...
- ▶ But, this sounds like relation, right? Which one?

Relation generated by a partition

- ▶ Clearly all elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ So can we define this as a relation?

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same set of courses...
- ▶ But, this sounds like relation, right? Which one?

Relation generated by a partition

- ▶ Clearly all elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ So can we define this as a relation? aRb if a is “like” b .

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same set of courses...
- ▶ But, this sounds like relation, right? Which one?

Relation generated by a partition

- ▶ Clearly all elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ So can we define this as a relation? aRb if a is “like” b .
- ▶ Formally, we define $R(S)$ by aRb if a and b belong to the same set in the partition of S .

Interpreting partitions as relations

Thus, a partition divides S into subsets that each contain (only) elements that share some common property. e.g.,

- ▶ Evenness or oddness (formally, remainder modulo 2!).
- ▶ Same set of courses...
- ▶ But, this sounds like relation, right? Which one?

Relation generated by a partition

- ▶ Clearly all elements in a set of the partition are related by the “sameness” or “likeness” property.
- ▶ So can we define this as a relation? aRb if a is “like” b .
- ▶ Formally, we define $R(S)$ by aRb if a and b belong to the same set in the partition of S .

What properties does this relation have?

Properties of a relation generated by a partition

1. All elements must be related to themselves (why?)

Properties of a relation generated by a partition

1. All elements must be related to themselves (why?)
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .

Properties of a relation generated by a partition

1. All elements must be related to themselves (why?)
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .

Properties of a relation generated by a partition

1. All elements must be related to themselves (why?)
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .

Properties of a relation generated by a partition

1. All elements must be related to themselves (why?)
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .

Properties of a relation generated by a partition

1. All elements must be related to themselves (why?)
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .
 - ▶ A relation $R(S)$ called **transitive** if for all $a, b, c \in S$, we have aRb and bRc implies aRc .

Properties of a relation generated by a partition

1. All elements must be related to themselves (why?)
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .
 - ▶ A relation $R(S)$ called **transitive** if for all $a, b, c \in S$, we have aRb and bRc implies aRc .

Any other defining properties?

Properties of a relation generated by a partition

1. All elements must be related to themselves (why?)
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .
 - ▶ A relation $R(S)$ called **transitive** if for all $a, b, c \in S$, we have aRb and bRc implies aRc .

Any other defining properties?

We will see that any relation that satisfies these three properties defines a partition!

Properties of a relation generated by a partition

1. All elements must be related to themselves (why?)
 - ▶ A relation $R(S)$ is called **reflexive** if for all $a \in S$, aRa .
2. If a is “like” b , then b must be “like” a .
 - ▶ A relation $R(S)$ is called **symmetric** if for all $a, b \in S$, we have aRb implies bRa .
3. If a is “like” b and b is “like” c , then a must be “like” c .
 - ▶ A relation $R(S)$ called **transitive** if for all $a, b, c \in S$, we have aRb and bRc implies aRc .

Any other defining properties?

We will see that any relation that satisfies these three properties defines a partition!

Definition

A relation which satisfies all these three properties is called an **equivalence relation**.

Examples

- ▶ **Reflexive:** $\forall a \in S, aRa$.
- ▶ **Symmetric:** $\forall a, b \in S, aRb$ implies bRa .
- ▶ **Transitive:** $\forall a, b, c \in S, aRb, bRc$ implies aRc .
- ▶ **Equivalence:** Reflexive, Symmetric and Transitive.

Relation	Refl.	Symm.	Trans.	Equiv.
aRb if students a and b take same set of courses	✓	✓	✓	✓
$\{(a, b) \mid a, b \in \mathbb{Z}, (a - b) \bmod 2 = 0\}$	✓	✓	✓	✓
$\{(a, b) \mid a, b \in \mathbb{Z}, a \leq b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}, a < b\}$				
$\{(a, b) \mid a, b \in \mathbb{Z}, a \mid b\}$				
$\{(a, b) \mid a, b \in \mathbb{R}, a - b < 1\}$				
$\{((a, b), (c, d)) \mid (a, b), (c, d) \in \mathbb{Z} \times (\mathbb{Z} \setminus \{0\}), (ad = bc)\}$				