

CS 207: Discrete Structures

Lecture 19 – Counting and Combinatorics Pigeon-Hole Principle

Aug 27 2015

Topics in Combinatorics

Basic counting techniques and applications

1. Basic counting techniques, double counting
2. Binomial theorem, permutations and combinations, Estimating $n!$
3. Recurrence relations and generating functions
4. Principle of Inclusion-Exclusion (PIE) and its applications.

Topics in Combinatorics

Basic counting techniques and applications

1. Basic counting techniques, double counting
2. Binomial theorem, permutations and combinations, Estimating $n!$
3. Recurrence relations and generating functions
4. Principle of Inclusion-Exclusion (PIE) and its applications.
 - ▶ Counting the number of surjections on $[n]$.
 - ▶ Combinatorial proof of PIE.
 - ▶ Number of derangements – $> \frac{1}{e}$.
 - ▶ Stirling numbers of the first kind.

Today's lecture

Pigeon-Hole Principle (PHP) and its applications

Today's lecture

Pigeon-Hole Principle (PHP) and its applications

A simple formulation

Let $k \in \mathbb{N}$. If $k + 1$ (or more) objects are to be placed in k boxes, then at least one box will have 2 (or more) objects.

Today's lecture

Pigeon-Hole Principle (PHP) and its applications

A simple formulation

Let $k \in \mathbb{N}$. If $k + 1$ (or more) objects are to be placed in k boxes, then at least one box will have 2 (or more) objects.

How do you prove it?

Today's lecture

Pigeon-Hole Principle (PHP) and its applications

A simple formulation

Let $k \in \mathbb{N}$. If $k + 1$ (or more) objects are to be placed in k boxes, then at least one box will have 2 (or more) objects.

How do you prove it?

A simple corollary

- Can a function from a set of $k + 1$ or more elements to a set with k elements be injective?

Pigeon-Hole Principle (PHP)

Another simple application

For every $n \in \mathbb{Z}^+$, there exists a multiple of n whose decimal expansion only has 0's and 1's.

Pigeon-Hole Principle (PHP)

Another simple application

For every $n \in \mathbb{Z}^+$, there exists a multiple of n whose decimal expansion only has 0's and 1's.

- ▶ Consider $n + 1$ integers $k_1 = 1, k_2 = 11, k_3 = 111, \dots, k_{n+1} = 1 \dots 1$ (with $n + 1$ 1's).

Pigeon-Hole Principle (PHP)

Another simple application

For every $n \in \mathbb{Z}^+$, there exists a multiple of n whose decimal expansion only has 0's and 1's.

- ▶ Consider $n + 1$ integers $k_1 = 1, k_2 = 11, k_3 = 111, \dots, k_{n+1} = 1 \dots 1$ (with $n + 1$ 1's).
- ▶ When any integer is divided by n , the remainder can be either $0, 1, \dots, n - 1$, i.e., n choices.

Pigeon-Hole Principle (PHP)

Another simple application

For every $n \in \mathbb{Z}^+$, there exists a multiple of n whose decimal expansion only has 0's and 1's.

- ▶ Consider $n + 1$ integers $k_1 = 1, k_2 = 11, k_3 = 111, \dots, k_{n+1} = 1 \dots 1$ (with $n + 1$ 1's).
- ▶ When any integer is divided by n , the remainder can be either $0, 1, \dots, n - 1$, i.e., n choices.
- ▶ So among the $n + 1$ integers, by PHP, at least 2 must have the same remainder.

Pigeon-Hole Principle (PHP)

Another simple application

For every $n \in \mathbb{Z}^+$, there exists a multiple of n whose decimal expansion only has 0's and 1's.

- ▶ Consider $n + 1$ integers $k_1 = 1, k_2 = 11, k_3 = 111, \dots, k_{n+1} = 1 \dots 1$ (with $n + 1$ 1's).
- ▶ When any integer is divided by n , the remainder can be either $0, 1, \dots, n - 1$, i.e., n choices.
- ▶ So among the $n + 1$ integers, by PHP, at least 2 must have the same remainder.
- ▶ That is, $\exists i, j, k_i = pn + d, k_j = qn + d$.
- ▶ But then $|k_i - k_j|$ is a multiple of n and its decimal expansion only has 0's and 1's. □

A (slightly) more general PHP

Pigeon-Hole Principle (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

A (slightly) more general PHP

Pigeon-Hole Principle (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

Suppose not. Then each box has strictly less than $\lfloor N/k \rfloor$ objects. Therefore, totally there can be strictly less than N objects, which is a contradiction.

A (slightly) more general PHP

Pigeon-Hole Principle (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

Suppose not. Then each box has strictly less than $\lfloor N/k \rfloor$ objects. Therefore, totally there can be strictly less than N objects, which is a contradiction.

A simple example

How many cards **must** be selected from a pack of 52 cards so that at least three cards of the same suit are chosen?

A (slightly) more general PHP

Pigeon-Hole Principle (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

Suppose not. Then each box has strictly less than $\lfloor N/k \rfloor$ objects. Therefore, totally there can be strictly less than N objects, which is a contradiction.

A simple example

How many cards **must** be selected from a pack of 52 cards so that at least three cards of the same suit are chosen?

- ▶ 4 boxes, one for each suit; each selected card is put in one.

A (slightly) more general PHP

Pigeon-Hole Principle (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

Suppose not. Then each box has strictly less than $\lceil N/k \rceil$ objects. Therefore, totally there can be strictly less than N objects, which is a contradiction.

A simple example

How many cards **must** be selected from a pack of 52 cards so that at least three cards of the same suit are chosen?

- ▶ 4 boxes, one for each suit; each selected card is put in one.
- ▶ If N cards are selected then at least 1 box has $\lceil N/4 \rceil$ cards.

A (slightly) more general PHP

Pigeon-Hole Principle (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

Suppose not. Then each box has strictly less than $\lceil N/k \rceil$ objects. Therefore, totally there can be strictly less than N objects, which is a contradiction.

A simple example

How many cards **must** be selected from a pack of 52 cards so that at least three cards of the same suit are chosen?

- ▶ 4 boxes, one for each suit; each selected card is put in one.
- ▶ If N cards are selected then at least 1 box has $\lceil N/4 \rceil$ cards.
- ▶ To have ≥ 3 cards from same suit, suffices $\lceil N/4 \rceil \geq 3$.

A (slightly) more general PHP

Pigeon-Hole Principle (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

Suppose not. Then each box has strictly less than $\lceil N/k \rceil$ objects. Therefore, totally there can be strictly less than N objects, which is a contradiction.

A simple example

How many cards **must** be selected from a pack of 52 cards so that at least three cards of the same suit are chosen?

- ▶ 4 boxes, one for each suit; each selected card is put in one.
- ▶ If N cards are selected then at least 1 box has $\lceil N/4 \rceil$ cards.
- ▶ To have ≥ 3 cards from same suit, suffices $\lceil N/4 \rceil \geq 3$.
- ▶ Thus, $N = 9$.

A (slightly) more general PHP

Pigeon-Hole Principle (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

Suppose not. Then each box has strictly less than $\lceil N/k \rceil$ objects. Therefore, totally there can be strictly less than N objects, which is a contradiction.

A simple example

How many cards **must** be selected from a pack of 52 cards so that at least three cards of the same suit are chosen?

- ▶ 4 boxes, one for each suit; each selected card is put in one.
- ▶ If N cards are selected then at least 1 box has $\lceil N/4 \rceil$ cards.
- ▶ To have ≥ 3 cards from same suit, suffices $\lceil N/4 \rceil \geq 3$.
- ▶ Thus, $N = 9$. But can we do better?

A (slightly) more general PHP

Pigeon-Hole Principle (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

Suppose not. Then each box has strictly less than $\lceil N/k \rceil$ objects. Therefore, totally there can be strictly less than N objects, which is a contradiction.

A simple example

How many cards **must** be selected from a pack of 52 cards so that at least three cards of the same suit are chosen?

- ▶ 4 boxes, one for each suit; each selected card is put in one.
- ▶ If N cards are selected then at least 1 box has $\lceil N/4 \rceil$ cards.
- ▶ To have ≥ 3 cards from same suit, suffices $\lceil N/4 \rceil \geq 3$.
- ▶ Thus, $N = 9$. But can we do better? **No.** □

Another application of PHP

Question: Give a sequence of 10 real numbers with no subsequence of length 4 which is increasing or decreasing.

Another application of PHP

Question: Give a sequence of 10 real numbers with no subsequence of length 4 which is increasing or decreasing.

Theorem

Every sequence of $n^2 + 1$ distinct real numbers contains a subsequence of length $n + 1$ which is either increasing or decreasing.

Another application of PHP

Theorem

Every sequence of $n^2 + 1$ distinct real numbers contains a subsequence of length $n + 1$ which is either increasing or decreasing.

1. Let a_1, \dots, a_{n^2+1} be a sequence of distinct real numbers.

Another application of PHP

Theorem

Every sequence of $n^2 + 1$ distinct real numbers contains a subsequence of length $n + 1$ which is either increasing or decreasing.

1. Let a_1, \dots, a_{n^2+1} be a sequence of distinct real numbers.
2. For each $k \in \{1 \dots n^2 + 1\}$, let (i_k, d_k) denote a pair:
 - i_k = length of longest increasing subsequence starting from a_k
 - d_k = length of longest decreasing subsequence starting from a_k

Another application of PHP

Theorem

Every sequence of $n^2 + 1$ distinct real numbers contains a subsequence of length $n + 1$ which is either increasing or decreasing.

1. Let a_1, \dots, a_{n^2+1} be a sequence of distinct real numbers.
2. For each $k \in \{1 \dots n^2 + 1\}$, let (i_k, d_k) denote a pair:
 i_k = length of longest increasing subsequence starting from a_k
 d_k = length of longest decreasing subsequence starting from a_k
3. Suppose, there are no increasing/decreasing subsequences of length $n + 1$. Then $\forall k, i_k \leq n$ and $d_k \leq n$.
4. \therefore by PHP, $\exists \ell, m, 1 \leq \ell < m \leq n^2 + 1$ s.t. $(i_\ell, d_\ell) = (i_m, d_m)$
5. We will show that this is not possible:
 - ▶ Case 1: $a_\ell < a_m$. Then $i_m \geq i_\ell + 1$, a contradiction.
 - ▶ Case 2: $a_\ell > a_m$. Then $d_\ell \geq d_m + 1$, a contradiction.
6. All a_i 's are distinct so this completes the proof. □

Another variant of PHP

Theorem (PHP Variant 2)

Suppose there are $n \geq 1 + r(\ell - 1)$ objects which are colored with r different colors. Then there exist ℓ objects all with the same color.

Another variant of PHP

Theorem (PHP Variant 2)

Suppose there are $n \geq 1 + r(\ell - 1)$ objects which are colored with r different colors. Then there exist ℓ objects all with the same color.

Proof: (H.W)

Another variant of PHP

Theorem (PHP Variant 2)

Suppose there are $n \geq 1 + r(\ell - 1)$ objects which are colored with r different colors. Then there exist ℓ objects all with the same color.

Proof: (H.W)

- Is this coloring optimal?

Another variant of PHP

Theorem (PHP Variant 2)

Suppose there are $n \geq 1 + r(\ell - 1)$ objects which are colored with r different colors. Then there exist ℓ objects all with the same color.

Proof: (H.W)

- ▶ Is this coloring optimal?
- ▶ That is, if fewer than $1 + r(\ell - 1)$ objects are given, is there a way of coloring them such that no ℓ have the same color?

Back to the coloring game

The coloring game

- ▶ There are six points on board and two colored chalks.

Back to the coloring game

The coloring game

- ▶ There are six points on board and two colored chalks.
- ▶ Divide class into 2 groups. Group 1 draws white lines and Group 2 draws blue lines between points.

Back to the coloring game

The coloring game

- ▶ There are six points on board and two colored chalks.
- ▶ Divide class into 2 groups. Group 1 draws white lines and Group 2 draws blue lines between points.
- ▶ You lose if you are first to draw a triangle of your color.

Back to the coloring game

The coloring game

- ▶ There are six points on board and two colored chalks.
- ▶ Divide class into 2 groups. Group 1 draws white lines and Group 2 draws blue lines between points.
- ▶ You lose if you are first to draw a triangle of your color.
- ▶ Can you ever have a draw?

Back to the coloring game

The coloring game

- ▶ There are six points on board and two colored chalks.
- ▶ Divide class into 2 groups. Group 1 draws white lines and Group 2 draws blue lines between points.
- ▶ You lose if you are first to draw a triangle of your color.
- ▶ Can you ever have a draw?

We will now show that this is impossible. That is,

Back to the coloring game

The coloring game

- ▶ There are six points on board and two colored chalks.
- ▶ Divide class into 2 groups. Group 1 draws white lines and Group 2 draws blue lines between points.
- ▶ You lose if you are first to draw a triangle of your color.
- ▶ Can you ever have a draw?

We will now show that this is impossible. That is,

Lemma

Any 2-coloring of edges of a graph on 6 nodes has a monochromatic triangle.

- ▶ **2-coloring of edges**: coloring all edges of the graph using atmost 2 colors.
- ▶ **monochromatic (triangle)**: all edges have the same color.

Back to the coloring game

Lemma

Any 2-coloring of edges of a graph on 6 nodes has a monochromatic triangle.

Proof:

Back to the coloring game

Lemma

Any 2-coloring of edges of a graph on 6 nodes has a monochromatic triangle.

Proof:

- ▶ Let $1, \dots, 6$ be the points, and red/blue the colors.
- ▶ Consider the edges $16, 26, 36, 46, 56$.
- ▶ By PHP at least 3 of them must be same color, say $16, 26, 36$ are red.
- ▶ Now there are two possibilities:
 - ▶ Either one of $12, 23, 31$ is red (then we have a red triangle).
 - ▶ Else none of them are red, implies 123 is a blue triangle. \square

Back to the coloring game

Lemma

Any 2-coloring of edges of a graph on 6 nodes has a monochromatic triangle.

Proof:

- ▶ Let $1, \dots, 6$ be the points, and red/blue the colors.
- ▶ Consider the edges $16, 26, 36, 46, 56$.
- ▶ By PHP at least 3 of them must be same color, say $16, 26, 36$ are red.
- ▶ Now there are two possibilities:
 - ▶ Either one of $12, 23, 31$ is red (then we have a red triangle).
 - ▶ Else none of them are red, implies 123 is a blue triangle. \square
- ▶ What if there were 5 or lesser nodes?