

CS 207: Discrete Structures

Lecture 20 – Counting and Combinatorics

PHP and its extensions

Ramsey Theory - A search for order in disorder

Aug 31 2015

Recap: Topics in Combinatorics

Counting techniques and applications

1. Basic counting techniques, double counting
2. Binomial theorem, permutations and combinations, Estimating $n!$
3. Recurrence relations and generating functions
4. Principle of Inclusion-Exclusion (PIE) and its applications.
 - ▶ Hand-shake Lemma
 - ▶ Counting the number of surjections on $[n]$.
 - ▶ Number of derangements $- > \frac{1}{e}$.
 - ▶ Number of partitions of size k .

Recap: Topics in Combinatorics

Counting techniques and applications

1. Basic counting techniques, double counting
2. Binomial theorem, permutations and combinations, Estimating $n!$
3. Recurrence relations and generating functions
4. Principle of Inclusion-Exclusion (PIE) and its applications.
 - ▶ Hand-shake Lemma
 - ▶ Counting the number of surjections on $[n]$.
 - ▶ Number of derangements $- > \frac{1}{e}$.
 - ▶ Number of partitions of size k .
5. Pigeon-Hole Principle (PHP) and its applications.
 - ▶ Every sequence of $n^2 + 1$ distinct real numbers contains a subsequence of length $n + 1$ which is either increasing or decreasing.
 - ▶ The coloring game.

Different variants of PHP

Simplest formulation (Variant 0)

Let $k \in \mathbb{N}$. If $k + 1$ (or more) objects are to be placed in k boxes, then at least one box will have 2 (or more) objects.

Different variants of PHP

Simplest formulation (Variant 0)

Let $k \in \mathbb{N}$. If $k + 1$ (or more) objects are to be placed in k boxes, then at least one box will have 2 (or more) objects.

PHP (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

Different variants of PHP

Simplest formulation (Variant 0)

Let $k \in \mathbb{N}$. If $k + 1$ (or more) objects are to be placed in k boxes, then at least one box will have 2 (or more) objects.

PHP (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

PHP (Variant 2)

If there are $n \geq 1 + r(\ell - 1)$ objects colored with r different colors, then there exist ℓ objects all with the same color.

Different variants of PHP

Simplest formulation (Variant 0)

Let $k \in \mathbb{N}$. If $k + 1$ (or more) objects are to be placed in k boxes, then at least one box will have 2 (or more) objects.

PHP (Variant 1)

If N objects are placed into k boxes then there is at least one box with at least $\lceil N/k \rceil$ objects.

PHP (Variant 2)

If there are $n \geq 1 + r(\ell - 1)$ objects colored with r different colors, then there exist ℓ objects all with the same color.

PHP (Variant 3)

If $n \geq k_1 + k_2 + \dots + k_r - r + 1$ objects are colored with r colors, then for some $i \in \{1 \dots r\}$, there exist k_i objects all of color i .

Back to the coloring game

Lemma

Any 2-coloring of edges of a graph on 6 nodes has a monochromatic triangle.

Back to the coloring game

Lemma

Any 2-coloring of edges of a graph on 6 nodes has a monochromatic triangle.

Proof:

- ▶ Let $1, \dots, 6$ be the points, and red/blue the colors.
- ▶ Consider the edges $16, 26, 36, 46, 56$.
- ▶ By PHP at least 3 of them must be same color, say $16, 26, 36$ are red.
- ▶ Now there are two possibilities:
 - ▶ Either one of $12, 23, 31$ is red (then we have a red triangle).
 - ▶ Else none of them are red, implies 123 is a blue triangle. \square

Back to the coloring game

Lemma

Any 2-coloring of edges of a graph on 6 nodes has a monochromatic triangle.

Proof:

- ▶ Let $1, \dots, 6$ be the points, and red/blue the colors.
- ▶ Consider the edges $16, 26, 36, 46, 56$.
- ▶ By PHP at least 3 of them must be same color, say $16, 26, 36$ are red.
- ▶ Now there are two possibilities:
 - ▶ Either one of $12, 23, 31$ is red (then we have a red triangle).
 - ▶ Else none of them are red, implies 123 is a blue triangle. \square
- ▶ What if there were 5 or lesser nodes?

Another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.


- ▶ **complete**: all pairs of edges are present.
- ▶ How do you prove this? Any ideas?
- ▶ How is this different from the previous problem?

Another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

Proof:




- Consider all edges from some node x .
- By PHP, either 4 edges have red color or 6 have blue.

Another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

Proof:




- ▶ Case 1: 4 red edges
 - ▶ Either one of edges between a, b, c, d is red or all are blue. So, we are done.

Another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

Proof:




- ▶ Case 2: 6 blue edges
 - ▶ But this means that there are 6 nodes a, \dots, f .

Another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

Proof:




- ▶ Case 2: 6 blue edges
 - ▶ But this means that there are 6 nodes a, \dots, f .
 - ▶ Any 2-coloring on 6 vertices has a red or blue triangle.
 - ▶ Thus we are done again.

Another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

Proof:



- ▶ Case 2: 6 blue edges
 - ▶ But this means that there are 6 nodes a, \dots, f .
 - ▶ Any 2-coloring on 6 vertices has a red or blue triangle.
 - ▶ Thus we are done again.
- ▶ And this completes the proof. □

Another coloring problem...

Thus, we have showed...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

Another coloring problem...

Thus, we have showed...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

- But, is this optimal?

Another coloring problem...

Thus, we have showed...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

- ▶ But, is this optimal?
- ▶ That is, does this fail for a graph on 9 nodes?
- ▶ Can you find 2-coloring on a graph of 9 nodes such that the statement above does NOT hold?

Another coloring problem...

Thus, we have showed...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

- ▶ But, is this optimal?
- ▶ That is, does this fail for a graph on 9 nodes?
- ▶ Can you find 2-coloring on a graph of 9 nodes such that the statement above does NOT hold?

Answer: No! In fact, it does hold on 9 nodes!

Another coloring problem...

Thus, we have showed...

Theorem

Any 2-coloring (say red and blue) of a graph on 10 nodes has either a **red triangle** or a **blue complete graph on 4 nodes**.

- ▶ But, is this optimal?
- ▶ That is, does this fail for a graph on 9 nodes?
- ▶ Can you find 2-coloring on a graph of 9 nodes such that the statement above does NOT hold?

Answer: No! In fact, it does hold on 9 nodes!

Theorem

Any 2-coloring (say red and blue) of a graph on **9 nodes** has either a **red triangle** or a **blue complete graph on 4 nodes**.

Yet another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on **9 nodes** has either a **red triangle** or a **blue complete graph on 4 nodes**.

Proof:

- ▶ Try the same proof as above?

Yet another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on **9 nodes** has either a **red triangle** or a **blue complete graph on 4 nodes**.

Proof:

- ▶ Try the same proof as above?
- ▶ The only new case, where the previous proof does not work, is if all nodes have 3 red edges and 5 blue edges. (Why?)

Yet another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on **9 nodes** has either a **red triangle** or a **blue complete graph on 4 nodes**.

Proof:

- ▶ Try the same proof as above?
- ▶ The only new case, where the previous proof does not work, is if all nodes have 3 red edges and 5 blue edges. (Why?)
- ▶ But is this case possible?

Yet another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on 9 nodes has either a red triangle or a blue complete graph on 4 nodes.

Proof:

- ▶ Try the same proof as above?
- ▶ The only new case, where the previous proof does not work, is if all nodes have 3 red edges and 5 blue edges. (Why?)
- ▶ But is this case possible?
- ▶ Recall the Handshake lemma!
 - ▶ In any graph, the number of nodes having odd degree is even.

Yet another coloring problem...

Theorem

Any 2-coloring (say red and blue) of a graph on 9 nodes has either a red triangle or a blue complete graph on 4 nodes.

Proof:

- ▶ Try the same proof as above?
- ▶ The only new case, where the previous proof does not work, is if all nodes have 3 red edges and 5 blue edges. (Why?)
- ▶ But is this case possible?
- ▶ Recall the Handshake lemma!
 - ▶ In any graph, the number of nodes having odd degree is even.
- ▶ Thus, this case is impossible and we are done. □

Can we generalize the above?

In general,

How many nodes should a (complete) graph have so that any 2 coloring of its edges has

- ▶ either, a k -sized complete graph with all red edges
- ▶ or, a ℓ -sized complete graph with all blue edges

Can we generalize the above?

In general,

How many nodes should a (complete) graph have so that any 2 coloring of its edges has

- ▶ either, a k -sized complete graph with all red edges
 - ▶ or, a ℓ -sized complete graph with all blue edges
- ▶ The minimal such number is denoted $R(k, \ell)$ and is called **Ramsey number**.

Can we generalize the above?

In general,

How many nodes should a (complete) graph have so that any 2 coloring of its edges has

- ▶ either, a k -sized complete graph with all red edges
- ▶ or, a ℓ -sized complete graph with all blue edges
- ▶ The minimal such number is denoted $R(k, \ell)$ and is called **Ramsey number**.
- ▶ We have seen that $R(3, 3) = 6$.
- ▶ Also $R(3, 4) =$

Can we generalize the above?

In general,

How many nodes should a (complete) graph have so that any 2 coloring of its edges has

- ▶ either, a k -sized complete graph with all red edges
 - ▶ or, a ℓ -sized complete graph with all blue edges
-
- ▶ The minimal such number is denoted $R(k, \ell)$ and is called **Ramsey number**.
 - ▶ We have seen that $R(3, 3) = 6$.
 - ▶ Also $R(3, 4) = 9$ (not quite...).

Can we generalize the above?

In general,

How many nodes should a (complete) graph have so that any 2 coloring of its edges has

- ▶ either, a k -sized complete graph with all red edges
- ▶ or, a ℓ -sized complete graph with all blue edges
- ▶ The minimal such number is denoted $R(k, \ell)$ and is called **Ramsey number**.
- ▶ We have seen that $R(3, 3) = 6$.
- ▶ Also $R(3, 4) = 9$ (not quite...).

What about $R(k, \ell)$ in general?

Ramsey's theorem



Figure : Frank Plumpton Ramsey (1903-1930)

Ramsey's theorem (simplified version)

For any $k, \ell \in \mathbb{N}$, there exists $R(k, \ell) \in \mathbb{N}$ such that any 2-coloring of a (complete) graph on $R(k, \ell)$ nodes has

- ▶ either, a k -sized complete graph with all red edges
- ▶ or, a ℓ -sized complete graph with all blue edges

Moreover, we have

$$R(k, \ell) \leq \binom{k + \ell - 2}{k - 1}$$

Edge coloring problems

Summary of results till now

1. Any 2-coloring of a graph on **6 nodes** has either a **red triangle** or a **blue triangle**.
 - ▶ 6 is the optimal such number. Thus, $R(3, 3) = 6$.

Edge coloring problems

Summary of results till now

1. Any 2-coloring of a graph on **6 nodes** has either a **red triangle** or a **blue triangle**.
 - ▶ 6 is the optimal such number. Thus, $R(3, 3) = 6$.
2. Any 2-coloring of a graph on **10 nodes** has either a **red triangle** or a **blue complete graph on 4 nodes**.

Edge coloring problems

Summary of results till now

1. Any 2-coloring of a graph on **6 nodes** has either a **red triangle** or a **blue triangle**.
 - ▶ 6 is the optimal such number. Thus, $R(3, 3) = 6$.
2. Any 2-coloring of a graph on **10 nodes** has either a **red triangle** or a **blue complete graph on 4 nodes**.
3. Any 2-coloring of a graph on **9 nodes** has either a **red triangle** or a **blue complete graph on 4 nodes**.
 - ▶ Is 9 the optimal such number? $R(3, 4) \leq 9$.
 - ▶ (H.W?) Prove that $R(3, 4) = 9$!

Edge coloring problems

Summary of results till now

1. Any 2-coloring of a graph on **6 nodes** has either a **red triangle** or a **blue triangle**.
 - ▶ 6 is the optimal such number. Thus, $R(3, 3) = 6$.
 2. Any 2-coloring of a graph on **10 nodes** has either a **red triangle** or a **blue complete graph on 4 nodes**.
 3. Any 2-coloring of a graph on **9 nodes** has either a **red triangle** or a **blue complete graph on 4 nodes**.
 - ▶ Is 9 the optimal such number? $R(3, 4) \leq 9$.
 - ▶ (H.W?) Prove that $R(3, 4) = 9$!
- ▶ (H.W) Prove that any 2-coloring of a graph on **18 nodes** has a **monochromatic complete graph on 4 nodes**.
(hint: you may use any of the above results)