CS 207: Discrete Structures


Graph theory

Matchings, maximum matchings, augmenting paths

Lecture 29 Oct 1 2015


- ▶ Suppose *m* people are applying for *n* different jobs. But not all applicants are qualified for all jobs, and each can hold at most one job.
- ► Then can you find a unique way to match jobs to applicants?

- ▶ Suppose *m* people are applying for *n* different jobs. But not all applicants are qualified for all jobs, and each can hold at most one job.
- ► Then can you find a unique way to match jobs to applicants?



).

- ▶ Suppose *m* people are applying for *n* different jobs. But not all applicants are qualified for all jobs, and each can hold at most one job.
- ► Then can you find a unique way to match jobs to applicants?



- ▶ What are the properties of such an assignment?
- ▶ Another practical example: the dating scene!

Definitions

- ightharpoonup A matching in a graph G is a set of (non-loop) edges with no shared end-points.
- ► The vertices incident to edges in a matching are called matched or saturated. Others are unsaturated.
- ▶ A perfect matching in a graph is a matching that saturates every vertex.

Definitions

- ▶ A matching in a graph G is a set of (non-loop) edges with no shared end-points.
- ► The vertices incident to edges in a matching are called matched or saturated. Others are unsaturated.
- ▶ A perfect matching in a graph is a matching that saturates every vertex.
- ▶ Is there a perfect matching if everyone is fully qualified/likes everyone?

Definitions

- ▶ A matching in a graph G is a set of (non-loop) edges with no shared end-points.
- ► The vertices incident to edges in a matching are called matched or saturated. Others are unsaturated.
- ► A perfect matching in a graph is a matching that saturates every vertex.
- ▶ Is there a perfect matching if everyone is fully qualified/likes everyone?
- ▶ How many perfect matchings are possibly in $K_{n,n}$?

- ightharpoonup A matching in a graph G is a set of (non-loop) edges with no shared end-points.
- ► The vertices incident to edges in a matching are called matched or saturated. Others are unsaturated.
- ► A perfect matching in a graph is a matching that saturates every vertex.

- ightharpoonup A matching in a graph G is a set of (non-loop) edges with no shared end-points.
- ► The vertices incident to edges in a matching are called matched or saturated. Others are unsaturated.
- ► A perfect matching in a graph is a matching that saturates every vertex.
- \triangleright Consider the problem of choosing roommates from a bunch of n people who may like or not like everyone.
- ▶ Or study-pairs...

How many perfect matchings are possible in K_n ?

- ightharpoonup A matching in a graph G is a set of (non-loop) edges with no shared end-points.
- ► The vertices incident to edges in a matching are called matched or saturated. Others are unsaturated.
- ► A perfect matching in a graph is a matching that saturates every vertex.
- \triangleright Consider the problem of choosing roommates from a bunch of n people who may like or not like everyone.
- ▶ Or study-pairs...

How many perfect matchings are possible in K_n ?

▶ If even, then

- ▶ A matching in a graph G is a set of (non-loop) edges with no shared end-points.
- ► The vertices incident to edges in a matching are called matched or saturated. Others are unsaturated.
- ► A perfect matching in a graph is a matching that saturates every vertex.
- \triangleright Consider the problem of choosing roommates from a bunch of n people who may like or not like everyone.
- ▶ Or study-pairs...

How many perfect matchings are possible in K_n ?

▶ If even, then f(n) = (2n-1)f(n-1)

- ▶ A matching in a graph G is a set of (non-loop) edges with no shared end-points.
- ► The vertices incident to edges in a matching are called matched or saturated. Others are unsaturated.
- ► A perfect matching in a graph is a matching that saturates every vertex.
- \triangleright Consider the problem of choosing roommates from a bunch of n people who may like or not like everyone.
- ▶ Or study-pairs...

How many perfect matchings are possible in K_n ?

▶ If even, then $f(n) = (2n-1)f(n-1) = \frac{(2n)!}{2^n n!}$.

- ▶ A matching in a graph G is a set of (non-loop) edges with no shared end-points.
- ► The vertices incident to edges in a matching are called matched or saturated. Others are unsaturated.
- ► A perfect matching in a graph is a matching that saturates every vertex.
- \triangleright Consider the problem of choosing roommates from a bunch of n people who may like or not like everyone.
- ▶ Or study-pairs...

How many perfect matchings are possible in K_n ?

▶ If even, then $f(n) = (2n-1)f(n-1) = \frac{(2n)!}{2^n n!}$.

Our goal is to characterize perfect matchings

But first...

Since we have seen that perfect matchings are not always possible, we can ask what is the best we can do.

Since we have seen that perfect matchings are not always possible, we can ask what is the best we can do.

Definition

A maximal matching in a graph is a matching that cannot be enlarged by adding an edge.

Since we have seen that perfect matchings are not always possible, we can ask what is the best we can do.

Definition

A maximal matching in a graph is a matching that cannot be enlarged by adding an edge. A maximum matching is

Since we have seen that perfect matchings are not always possible, we can ask what is the best we can do.

Definition

A maximal matching in a graph is a matching that cannot be enlarged by adding an edge. A maximum matching is a matching of maximum size among all matchings in a graph.

Since we have seen that perfect matchings are not always possible, we can ask what is the best we can do.

Definition

A maximal matching in a graph is a matching that cannot be enlarged by adding an edge. A maximum matching is a matching of maximum size among all matchings in a graph.

▶ A matching is maximal if every edge not in it is incident to some edge in it.

Since we have seen that perfect matchings are not always possible, we can ask what is the best we can do.

Definition

A maximal matching in a graph is a matching that cannot be enlarged by adding an edge. A maximum matching is a matching of maximum size among all matchings in a graph.

- ▶ A matching is maximal if every edge not in it is incident to some edge in it.
- ▶ Does maximum matching imply maximal?

Since we have seen that perfect matchings are not always possible, we can ask what is the best we can do.

Definition

A maximal matching in a graph is a matching that cannot be enlarged by adding an edge. A maximum matching is a matching of maximum size among all matchings in a graph.

- ▶ A matching is maximal if every edge not in it is incident to some edge in it.
- ▶ Does maximum matching imply maximal? Does maximal imply maximum?

Since we have seen that perfect matchings are not always possible, we can ask what is the best we can do.

Definition

A maximal matching in a graph is a matching that cannot be enlarged by adding an edge. A maximum matching is a matching of maximum size among all matchings in a graph.

- ▶ A matching is maximal if every edge not in it is incident to some edge in it.
- ▶ Does maximum matching imply maximal? Does maximal imply maximum?
- ► Find the smallest graph that has a maximal matching that is not maximum.

Since we have seen that perfect matchings are not always possible, we can ask what is the best we can do.

Definition

A maximal matching in a graph is a matching that cannot be enlarged by adding an edge. A maximum matching is a matching of maximum size among all matchings in a graph.

- ▶ A matching is maximal if every edge not in it is incident to some edge in it.
- ▶ Does maximum matching imply maximal? Does maximal imply maximum?
- ▶ Find the smallest graph that has a maximal matching that is not maximum. P_4 .

Definition

Given a matching M, an M-alternating path is a path that alternates between edges in M and edges not in M. An M-alternating path whose endpoints are unmatched by M is an M-augmenting path.


Definition

Given a matching M, an M-alternating path is a path that alternates between edges in M and edges not in M. An M-alternating path whose endpoints are unmatched by M is an M-augmenting path.




Definition

Given a matching M, an M-alternating path is a path that alternates between edges in M and edges not in M. An M-alternating path whose endpoints are unmatched by M is an M-augmenting path.



Definition

Given a matching M, an M-alternating path is a path that alternates between edges in M and edges not in M. An M-alternating path whose endpoints are unmatched by M is an M-augmenting path.



► Clearly, a maximum matching cannot have an *M*-augmenting path.

- Clearly, a maximum matching cannot have an M-augmenting path.
- ▶ In fact, this is the characterization!

- ► Clearly, a maximum matching cannot have an *M*-augmenting path.
- ▶ In fact, this is the characterization!


Theorem

A matching M in G is a maximum matching iff G has no M-augmenting path.

We need a definition and a lemma.

Definition


If M, M' are matchings in a graph H, the symmetric difference $M \triangle M'$ is the set of edges which are either in M or in M' but not both, i.e., $M \triangle M' = (M \setminus M') \cup (M' \setminus M)$.



We need a definition and a lemma.

Definition


If M, M' are matchings in a graph H, the symmetric difference $M \triangle M'$ is the set of edges which are either in M or in M' but not both, i.e., $M \triangle M' = (M \setminus M') \cup (M' \setminus M)$.




We need a definition and a lemma.

Definition


If M, M' are matchings in a graph H, the symmetric difference $M \triangle M'$ is the set of edges which are either in M or in M' but not both, i.e., $M \triangle M' = (M \setminus M') \cup (M' \setminus M)$.





Lemma


Every component of the symmetric difference of two matchings is either a path or an even cycle.



Lemma

Every component of the symmetric difference of two matchings is either a path or an even cycle.

▶ Let $F = M \triangle M'$. F has at most 2 edges at each vertex, hence every component is a path or a cycle.



Lemma

Every component of the symmetric difference of two matchings is either a path or an even cycle.

- Let $F = M \triangle M'$. F has at most 2 edges at each vertex, hence every component is a path or a cycle.
- ▶ Further every path/cycle alternates between edges of $M \setminus M'$ and $M' \setminus M$.



Lemma

Every component of the symmetric difference of two matchings is either a path or an even cycle.

- ▶ Let $F = M \triangle M'$. F has at most 2 edges at each vertex, hence every component is a path or a cycle.
- ▶ Further every path/cycle alternates between edges of $M \setminus M'$ and $M' \setminus M$.
- ▶ Thus, each cycle has even length with equal edges from M and M'.

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M-augmenting path.

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M-augmenting path.

Proof:

▶ One direction is trivial (which one?!).

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M-augmenting path.

- ▶ One direction is trivial (which one?!).
- \blacktriangleright (\Leftarrow) For the other, we will show the contrapositive.

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M-augmenting path.

- ▶ One direction is trivial (which one?!).
- \blacktriangleright (\Leftarrow) For the other, we will show the contrapositive.
- ▶ i.e., if \exists matching M' larger than M, we will construct an M-augmenting path.

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M-augmenting path.

- ▶ One direction is trivial (which one?!).
- \blacktriangleright (\Leftarrow) For the other, we will show the contrapositive.
- ▶ i.e., if \exists matching M' larger than M, we will construct an M-augmenting path.
- ▶ Let $F = M \triangle M'$. By Lemma, F has only paths and even cycles with equal no. of edges from M and M'.

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M-augmenting path.

- ▶ One direction is trivial (which one?!).
- ► (⇐=) For the other, we will show the contrapositive.
- ▶ i.e., if \exists matching M' larger than M, we will construct an M-augmenting path.
- ▶ Let $F = M \triangle M'$. By Lemma, F has only paths and even cycles with equal no. of edges from M and M'.
- ▶ But then since |M'| > |M| it must have a component with more edges in M' than M.

Theorem (Berge'57)

A matching M in G is a maximum matching iff G has no M-augmenting path.

- ▶ One direction is trivial (which one?!).
- ► (⇐=) For the other, we will show the contrapositive.
- ▶ i.e., if \exists matching M' larger than M, we will construct an M-augmenting path.
- ▶ Let $F = M \triangle M'$. By Lemma, F has only paths and even cycles with equal no. of edges from M and M'.
- ▶ But then since |M'| > |M| it must have a component with more edges in M' than M.
- ▶ This component can only be a path that starts and ends with an edge of M''; i.e., it is an M-augmenting path in G.