

CS 207: Discrete Structures

Graph theory

Basic terminology, Eulerian graphs and a characterization

Lecture 23

Sept 14 2014

Topic 3: Graph theory

Textbook Reference

- ▶ Introduction to Graph Theory, 2nd Ed., by Douglas West.
- ▶ Low cost Indian edition available, published by PHI Learning Private Ltd.

Topic 3: Graph theory

Textbook Reference

- ▶ **Introduction to Graph Theory, 2nd Ed., by Douglas West.**
- ▶ Low cost Indian edition available, published by PHI Learning Private Ltd.

Topics covered in the first two lectures:

- ▶ What is a Graph?
- ▶ Paths, cycles, walks and trails; connected graphs.
- ▶ Eulerian graphs and a characterization in terms of degrees of vertices.
- ▶ Bipartite graphs and a characterization in terms of odd length cycles.

Reference: Section 1.1, 1.2 of Chapter 1 from **Douglas West**.

What are graphs

Recall:

Definition

A simple graph G is a pair (V, E) of a set of vertices/nodes V and edges E between unordered pairs of vertices called end-points: $e = vu$ means that e is an edge between v and u ($u \neq v$).

What are graphs

Recall:

Definition

A simple graph G is a pair (V, E) of a set of vertices/nodes V and edges E between unordered pairs of vertices called end-points: $e = vu$ means that e is an edge between v and u ($u \neq v$).

- ▶ What about loops?
- ▶ What about directed edges?
- ▶ What about multiple edges?

What are graphs

Recall:

Definition

A simple graph G is a pair (V, E) of a set of vertices/nodes V and edges E between unordered pairs of vertices called end-points: $e = vu$ means that e is an edge between v and u ($u \neq v$).


- ▶ What about loops?
- ▶ What about directed edges?
- ▶ What about multiple edges?

General Definition

A graph G is a triple V, E, R where V is a set of vertices, E is a set of edges and $R \subseteq E \times V \times V$ is a relation that associates each edge with two vertices called its end-points.


We will consider **only** finite graphs (i.e., $|V|, |E|$ are finite) and **often** deal with simple graphs.

Königsberg Bridge problem





- ▶ In 18th century Prussia, the city on river Pregel...
- ▶ Question was to find a walk from home, crossing every bridge exactly once and returning home.

Königsberg Bridge problem





- ▶ In 18th century Prussia, the city on river Pregel...
- ▶ Question was to find a walk from home, crossing every bridge exactly once and returning home.
- ▶ Leonard Euler showed in 1735 that this is impossible.
- ▶ Thus, he “gave birth” to the area of graph theory.

Königsberg Bridge problem





- ▶ In 18th century Prussia, the city on river Pregel...
- ▶ Question was to find a walk from home, crossing every bridge exactly once and returning home.
- ▶ Leonard Euler showed in 1735 that this is impossible.
- ▶ Thus, he “gave birth” to the area of graph theory.

Königsberg Bridge problem





- ▶ Qn: Find a walk from home, which crosses every bridge exactly once and returns home.
- ▶ Leonard Euler showed in 1735 that this is impossible. The argument is as follows:

Königsberg Bridge problem





- ▶ Qn: Find a walk from home, which crosses every bridge exactly once and returns home.
- ▶ Leonard Euler showed in 1735 that this is impossible. The argument is as follows:
 - ▶ Each time you enter or leave a vertex, you use an edge.

Königsberg Bridge problem





- ▶ Qn: Find a walk from home, which crosses every bridge exactly once and returns home.
- ▶ Leonard Euler showed in 1735 that this is impossible. The argument is as follows:
 - ▶ Each time you enter or leave a vertex, you use an edge.
 - ▶ So each vertex must be connected to an even no. of vertices.

Königsberg Bridge problem





- ▶ Qn: Find a walk from home, which crosses every bridge exactly once and returns home.
- ▶ Leonard Euler showed in 1735 that this is impossible. The argument is as follows:
 - ▶ Each time you enter or leave a vertex, you use an edge.
 - ▶ So each vertex must be connected to an even no. of vertices.
 - ▶ Which is not the case here, hence it is impossible.

Königsberg Bridge problem





- ▶ Qn: Find a walk from home, which crosses every bridge exactly once and returns home.
- ▶ Leonard Euler showed in 1735 that this is impossible. The argument is as follows:
 - ▶ Each time you enter or leave a vertex, you use an edge.
 - ▶ So each vertex must be connected to an even no. of vertices.
 - ▶ Which is not the case here, hence it is impossible.
- ▶ Clearly, this is a sufficient condition, but is it necessary?

Königsberg Bridge problem



- ▶ Qn: Find a walk from home, which crosses every bridge exactly once and returns home.
- ▶ Leonard Euler showed in 1735 that this is impossible. The argument is as follows:
 - ▶ Each time you enter or leave a vertex, you use an edge.
 - ▶ So each vertex must be connected to an even no. of vertices.
 - ▶ Which is not the case here, hence it is impossible.
- ▶ Clearly, this is a sufficient condition, but is it necessary?
- ▶ If every vertex is connected to an even no. of vertices in a graph, is there such a walk?

Königsberg Bridge problem



- ▶ Qn: Find a walk from home, which crosses every bridge exactly once and returns home.
- ▶ Leonard Euler showed in 1735 that this is impossible. The argument is as follows:
 - ▶ Each time you enter or leave a vertex, you use an edge.
 - ▶ So each vertex must be connected to an even no. of vertices.
 - ▶ Which is not the case here, hence it is impossible.
- ▶ Clearly, this is a sufficient condition, but is it necessary?
- ▶ If every vertex is connected to an even no. of vertices in a graph, is there such a walk? This is called **Eulerian walk**.

Basic terminology

The **degree** $d(v)$ of a **vertex** v (in an undirected loopless graph) is the number of edges incident to it, i.e., $|\{vw \in E \mid w \in V\}|$. A vertex of degree 0 is called an **isolated vertex**.

Basic terminology

The **degree** $d(v)$ of a **vertex** v (in an undirected loopless graph) is the number of edges incident to it, i.e., $|\{vw \in E \mid w \in V\}|$. A vertex of degree 0 is called an **isolated vertex**.

- ▶ A **walk** is a sequence of vertices v_1, \dots, v_k such that $\forall i \in \{1, \dots, k-1\}, (v_i, v_{i+1}) \in E$. The vertices v_1 and v_k are called the **end-points** and others are called **internal vertices**.

Basic terminology

The **degree** $d(v)$ of a vertex v (in an undirected loopless graph) is the number of edges incident to it, i.e., $|\{vw \in E \mid w \in V\}|$. A vertex of degree 0 is called an **isolated vertex**.

- ▶ A **walk** is a sequence of vertices v_1, \dots, v_k such that $\forall i \in \{1, \dots, k-1\}, (v_i, v_{i+1}) \in E$. The vertices v_1 and v_k are called the **end-points** and others are called **internal vertices**.
- ▶ A **path** is a walk in which no vertex is repeated. Its **length** is the number of **edges** in it.

Basic terminology

The **degree** $d(v)$ of a vertex v (in an undirected loopless graph) is the number of edges incident to it, i.e., $|\{vw \in E \mid w \in V\}|$. A vertex of degree 0 is called an **isolated vertex**.

- ▶ A **walk** is a sequence of vertices v_1, \dots, v_k such that $\forall i \in \{1, \dots, k-1\}, (v_i, v_{i+1}) \in E$. The vertices v_1 and v_k are called the **end-points** and others are called **internal vertices**.
- ▶ A **path** is a walk in which no vertex is repeated. Its **length** is the number of **edges** in it.
- ▶ A walk is called **closed** if it starts and ends with the same vertex, i.e., its endpoints are the same.

Basic terminology

The **degree** $d(v)$ of a vertex v (in an undirected loopless graph) is the number of edges incident to it, i.e., $|\{vw \in E \mid w \in V\}|$. A vertex of degree 0 is called an **isolated vertex**.

- ▶ A **walk** is a sequence of vertices v_1, \dots, v_k such that $\forall i \in \{1, \dots, k-1\}, (v_i, v_{i+1}) \in E$. The vertices v_1 and v_k are called the **end-points** and others are called **internal vertices**.
- ▶ A **path** is a walk in which no vertex is repeated. Its **length** is the number of **edges** in it.
- ▶ A walk is called **closed** if it starts and ends with the same vertex, i.e., its endpoints are the same.
- ▶ A closed walk is called a **cycle** if its internal vertices are all distinct from each other and from the end-point.

Basic terminology

The **degree** $d(v)$ of a vertex v (in an undirected loopless graph) is the number of edges incident to it, i.e., $|\{vw \in E \mid w \in V\}|$. A vertex of degree 0 is called an **isolated vertex**.

- ▶ A **walk** is a sequence of vertices v_1, \dots, v_k such that $\forall i \in \{1, \dots, k-1\}, (v_i, v_{i+1}) \in E$. The vertices v_1 and v_k are called the **end-points** and others are called **internal vertices**.
- ▶ A **path** is a walk in which no vertex is repeated. Its **length** is the number of **edges** in it.
- ▶ A walk is called **closed** if it starts and ends with the same vertex, i.e., its endpoints are the same.
- ▶ A closed walk is called a **cycle** if its internal vertices are all distinct from each other and from the end-point.

A graph is called **connected** if there is a path (or walk) between any two of its vertices.

Eulerian graphs

Definition

A graph is called **Eulerian** if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an **Eulerian walk**.

Eulerian graphs

Definition

A graph is called **Eulerian** if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an **Eulerian walk**.

- ▶ If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

Eulerian graphs

Definition

A graph is called **Eulerian** if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an **Eulerian walk**.

- ▶ If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

Eulerian graphs

Definition

A graph is called **Eulerian** if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an **Eulerian walk**.

- ▶ If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

Proof:

Eulerian graphs

Definition

A graph is called **Eulerian** if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an **Eulerian walk**.

- ▶ If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

Proof: (\implies)

- ▶ Suppose G is Eulerian: every vertex has even degree.
 - ▶ each passage through a vertex uses two edges (in and out).
 - ▶ at the first vertex first edge is paired with last.

Eulerian graphs

Definition

A graph is called **Eulerian** if it has a closed walk that contains all edges, and each edge occurs exactly once. Such a walk is called an **Eulerian walk**.

- ▶ If G is Eulerian, each vertex must have an even degree.
- ▶ This is a necessary condition, but is it sufficient?

Theorem

A graph G with no isolated vertices is Eulerian iff it is connected and all its vertices have even degree.

Proof: (\implies)

- ▶ Suppose G is Eulerian: every vertex has even degree.
 - ▶ each passage through a vertex uses two edges (in and out).
 - ▶ at the first vertex first edge is paired with last.
- ▶ Any two edges are in the same walk implies graph is connected (unless it has isolated vertices).

Eulerian graphs

Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

Eulerian graphs

A path P is said to be **maximal** in G if it is not contained in any longer path. In a finite graph maximal paths exist.

Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.


Eulerian graphs

A path P is said to be **maximal** in G if it is not contained in any longer path. In a finite graph maximal paths exist.

Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

Proof:



- Let P be a maximal path in G . It starts from some u .


Eulerian graphs

A path P is said to be **maximal** in G if it is not contained in any longer path. In a finite graph maximal paths exist.

Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

Proof:



- ▶ Let P be a maximal path in G . It starts from some u .
- ▶ Since P cannot be extended, every nbr of u is already in P .


Eulerian graphs

A path P is said to be **maximal** in G if it is not contained in any longer path. In a finite graph maximal paths exist.

Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

Proof:



- ▶ Let P be a maximal path in G . It starts from some u .
- ▶ Since P cannot be extended, every nbr of u is already in P .
- ▶ As $d(u) \geq 2$, $\exists v$ in P such that $uv \notin P$.
- ▶ Thus we have the cycle $u \dots vu$. □


Eulerian graphs

A path P is said to be **maximal** in G if it is not contained in any longer path. In a finite graph maximal paths exist.

Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

Proof:



- ▶ Let P be a maximal path in G . It starts from some u .
- ▶ Since P cannot be extended, every nbr of u is already in P .
- ▶ As $d(u) \geq 2$, $\exists v$ in P such that $uv \notin P$.
- ▶ Thus we have the cycle $u \dots vu$. □


Is this true if G were infinite?

Eulerian graphs

Lemma

If every vertex of a graph G has degree at least 2, then G contains a cycle.

Proof:



- ▶ Let P be a maximal path in G . It starts from some u .
- ▶ Since P cannot be extended, every nbr of u is already in P .
- ▶ As $d(u) \geq 2$, $\exists v$ in P such that $uv \notin P$.
- ▶ Thus we have the cycle $u \dots vu$. □

Is this true if G were infinite?

No! Consider $V = \mathbb{Z}$, $E = \{ij : |i - j| = 1\}$.

