

CS 207: Discrete Structures

Lecture 13 – Counting and Combinatorics

Aug 17 2015

Last class

Basic counting techniques

- ▶ Sum and product principles
- ▶ Bijection principle
- ▶ Binomial coefficients, permutations and combinations
- ▶ Double counting

Last class

Basic counting techniques

- ▶ Sum and product principles
- ▶ Bijection principle
- ▶ Binomial coefficients, permutations and combinations
- ▶ Double counting

Applications

- ▶ No. of subsets of a set, no. of subsets of fixed size, no. of ordered subsets of a fixed size.
- ▶ No. of reflexive relations, no. of symmetric relations.
- ▶ Proving identities on binomial coefficients.
- ▶ Handshake lemma: Number of people who shake hands an odd number of times is even.
- ▶ No. of equivalence relations or partitions of a set.

What about partitions?

We have looked at:

- ▶ number of subsets of a set (combinations)
- ▶ number of ordered subsets of a set (permutations)

What about number of partitions of a set of n distinct elements, denoted B_n ?

What about partitions?

We have looked at:

- ▶ number of subsets of a set (combinations)
- ▶ number of ordered subsets of a set (permutations)

What about number of partitions of a set of n distinct elements, denoted B_n ?

- ▶ Same as number of equivalence relations!
- ▶ What are B_3 , B_2 , B_1 ? What about B_0 ?
- ▶ What about B_n in general?

What about partitions?

We have looked at:

- ▶ number of subsets of a set (combinations)
- ▶ number of ordered subsets of a set (permutations)

What about number of partitions of a set of n distinct elements, denoted B_n ?

- ▶ Same as number of equivalence relations!
- ▶ What are B_3 , B_2 , B_1 ? What about B_0 ?
- ▶ What about B_n in general?

Turns out

- ▶ there is no closed-form answer to this question!
- ▶ can we get a recurrence?

What about partitions?

We have looked at:

- ▶ number of subsets of a set (combinations)
- ▶ number of ordered subsets of a set (permutations)

What about number of partitions of a set of n distinct elements, denoted B_n ?

- ▶ Same as number of equivalence relations!
- ▶ What are B_3 , B_2 , B_1 ? What about B_0 ?
- ▶ What about B_n in general?

Turns out

- ▶ there is no closed-form answer to this question!
- ▶ can we get a recurrence?

- ▶ Prove by counting:
$$B_{n+1} = \sum_{k=0}^n \binom{n}{k} B_k$$

Next, we will see

Basic counting techniques... (contd.)

1. Binomial coefficients and Binomial theorem
2. Pascal's triangle
3. Permutations and combinations with repetitions
4. Estimating $n!$

Binomial theorem

Recall: $\sum_{k=0}^n \binom{n}{k} = 2^n.$

We generalize this...

Binomial theorem

Recall: $\sum_{k=0}^n \binom{n}{k} = 2^n$.

We generalize this...

Binomial Theorem

Let x, y be variables and $n \in \mathbb{Z}^{\geq 0}$. Then,

$$(x + y)^n = \sum_{j=0}^n \binom{n}{j} x^{n-j} y^j$$

$$(x + y)^1 = x + y$$

$$(x + y)^2 = (x + y)(x + y) = x^2 + 2xy + y^2$$

$$(x + y)^3 = (x + y)(x + y)^2 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$(x + y)^4 = (x + y)(x + y)^3 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$$

...

Binomial theorem

Binomial Theorem

Let x, y be variables and $n \in \mathbb{Z}^{\geq 0}$. Then,

$$(x + y)^n = \sum_{j=0}^n \binom{n}{j} x^{n-j} y^j$$

(H.W-1) Prove this by induction.

Binomial theorem

Binomial Theorem

Let x, y be variables and $n \in \mathbb{Z}^{\geq 0}$. Then,

$$(x + y)^n = \sum_{j=0}^n \binom{n}{j} x^{n-j} y^j$$

Proof (combinatorial):

1. Consider any term $x^i y^j$, where $i + j = n$.
2. To get $x^i y^j$ term in

$$(x + y)(x + y) \cdots (x + y) \quad (n \text{ times})$$

we need to pick j y 's from n sums and remaining x 's.

Binomial theorem

Binomial Theorem

Let x, y be variables and $n \in \mathbb{Z}^{\geq 0}$. Then,

$$(x + y)^n = \sum_{j=0}^n \binom{n}{j} x^{n-j} y^j$$

Proof (combinatorial):

1. Consider any term $x^i y^j$, where $i + j = n$.
2. To get $x^i y^j$ term in

$$(x + y)(x + y) \cdots (x + y) \quad (n \text{ times})$$

we need to pick j y 's from n sums and remaining x 's.

3. Thus, the coefficient of this term = number of ways to get this term = number of ways to pick j y 's from n elts = $\binom{n}{j}$.

Binomial theorem

Binomial Theorem

Let x, y be variables and $n \in \mathbb{Z}^{\geq 0}$. Then,

$$(x + y)^n = \sum_{j=0}^n \binom{n}{j} x^{n-j} y^j$$

Binomial theorem

Binomial Theorem

Let x, y be variables and $n \in \mathbb{Z}^{\geq 0}$. Then,

$$(x + y)^n = \sum_{j=0}^n \binom{n}{j} x^{n-j} y^j$$

Corollaries:

1. $\binom{n}{j} = \binom{n}{n-j}$,
2. $\sum_{j=0}^n \binom{n}{j} 2^j = 3^n$.
3. No. of subsets of n -element set having even cardinality = ?
(H.W-2)


Pascal's Triangle

A recursive way to compute binomial coefficients

$$\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$$

$$\begin{array}{ccccccc}
& & \binom{0}{0} & & & & \\
& & & & 1 & & \\
& \binom{1}{0} & \binom{1}{1} & & & & \\
& & & & 1 & 1 & \\
& \binom{2}{0} & \binom{2}{1} & \binom{2}{2} & & & \\
& & & & 1 & 2 & 1 \\
& \binom{3}{0} & \binom{3}{1} & \binom{3}{2} & \binom{3}{3} & & \\
& & & & 1 & 3 & 3 & 1 \\
& \binom{4}{0} & \binom{4}{1} & \binom{4}{2} & \binom{4}{3} & \binom{4}{4} & & \\
& & & & 1 & 4 & 6 & 4 & 1 \\
& \binom{5}{0} & \binom{5}{1} & \binom{5}{2} & \binom{5}{3} & \binom{5}{4} & \binom{5}{5} & & \\
& & & & 1 & 5 & 10 & 10 & 5 & 1
\end{array}$$


Fun with Pascal's triangle



Some simple observations. Recall: $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

1. Row i adds up to 2^i , Row $i + 1$ adds up to twice of row i .
2. Sequence of numbers, squares, cubes?


Fun with Pascal's triangle



Some simple observations. Recall: $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

1. Row i adds up to 2^i , Row $i + 1$ adds up to twice of row i .
2. Sequence of numbers, squares, cubes?
3. Hockey stick patterns

Fun with Pascal's triangle



Some simple observations. Recall: $\binom{n+1}{k} = \binom{n}{k-1} + \binom{n}{k}$

1. Row i adds up to 2^i , Row $i + 1$ adds up to twice of row i .
2. Sequence of numbers, squares, cubes?
3. Hockey stick patterns: (H.W-3)

$$\binom{n+1}{m} = \binom{n}{m} + \binom{n-1}{m-1} + \dots + \binom{n-m}{0}$$

Fun with Pascal's triangle

Pascal's Triangle:

					1						
				1		1					
			1		2		1				
		1		3		3		1			
	1		4		6		4		1		
	1	5		10		10		5		1	
	1	6	15		20		15	6		1	
	1	7	21	35		35	21	7		1	
	1	8	28	56	70		56	28	8		1
1		9	36	84	126	126		84	36	9	
1	10		45	120	200	252	200	120	45	10	1

Some **not so** simple observations

- For some rows, all values in the row (except first and last) are divisible by the second!

Fun with Pascal's triangle

					1																				
					1		1																		
					1		2		1																
					1		3		3		1														
					1		4		6		4		1												
					1		5		10		10		5		1										
					1		6		15		20		15		6		1								
					1		7		21		35		35		21		7		1						
					1		8		28		56		70		56		28		8		1				
					1		9		36		84		126		126		84		36		9		1		
					1		10		45		120		200		252		200		120		45		10		1

Some **not so** simple observations

- ▶ For some rows, all values in the row (except first and last) are divisible by the second!
- ▶ In fact, for all prime rows? why should p divide $\binom{p}{r}$, $r < p$?


Fun with Pascal's triangle

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Some **not so** simple observations

- ▶ For some rows, all values in the row (except first and last) are divisible by the second!
- ▶ In fact, for all prime rows? why should p divide $\binom{p}{r}$, $r < p$?
- ▶ **Corollary:** $2^p - 2$ is a multiple of p , for any prime p .

Fun with Pascal's triangle




Some **not so** simple observations

- For some rows, all values in the row (except first and last) are divisible by the second!
- In fact, for all prime rows? why should p divide $\binom{p}{r}$, $r < p$?
- **Corollary:** $2^p - 2$ is a multiple of p , for any prime p .
- **Interesting Ex.:** Count no. of odd numbers in each row...

An application to path counting

Map problems


From the top corner, how many shortest routes lead to a particular junction?



An application to path counting

Map problems

From the top corner, how many shortest routes lead to a particular junction?




- Denote path sequences of $\{L, R\}$, e.g., $RRRLL$ reaches $*$.

An application to path counting

Map problems

From the top corner, how many shortest routes lead to a particular junction?




- ▶ Denote path sequences of $\{L, R\}$, e.g., $RRRLL$ reaches $*$.
- ▶ Our path has 5 segments of which there must be 3 R's.

An application to path counting

Map problems

From the top corner, how many shortest routes lead to a particular junction?




- ▶ Denote path sequences of $\{L, R\}$, e.g., $RRRLL$ reaches $*$.
- ▶ Our path has 5 segments of which there must be 3 R's.
- ▶ Bijection from set of paths to subsets of fixed size.

An application to path counting

Map problems

From the top corner, how many shortest routes lead to a particular junction?




- ▶ Denote path sequences of $\{L, R\}$, e.g., $RRRLL$ reaches $*$.
- ▶ Our path has 5 segments of which there must be 3 R's.
- ▶ Bijection from set of paths to subsets of fixed size.
- ▶ No. of paths = ways of choosing 3 R's out of 5 elts = $\binom{5}{3}$.

An application to path counting

Map problems

From the top corner, how many shortest routes lead to a particular junction?



- ▶ Denote path sequences of $\{L, R\}$, e.g., $RRRLL$ reaches $*$.
- ▶ Our path has 5 segments of which there must be 3 R's.
- ▶ Bijection from set of paths to subsets of fixed size.
- ▶ No. of paths = ways of choosing 3 R's out of 5 elts = $\binom{5}{3}$.

H.W-4: Prove/verify this formally.

Permutations and Combinations with repetitions

How many ways can you select k objects from a set of n elements?

- ▶ Depends on whether order is significant: If yes permutations, else combinations.
- ▶ What if repetitions are allowed?

	Order significant	Order not significant
Repetitions not allowed	$\frac{n!}{(n-k)!}$	$\binom{n}{k}$
Repetitions are allowed	n^k	??

Combinations with repetitions

Theorem

The no. of ways k elements can be chosen from n -elements, when repetition is allowed is $\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$.

Combinations with repetitions

Theorem

The no. of ways k elements can be chosen from n -elements, when repetition is allowed is $\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$.

1. Represent them as a list of $n - 1$ separators of k objects.

Combinations with repetitions

Theorem

The no. of ways k elements can be chosen from n -elements, when repetition is allowed is $\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$.

1. Represent them as a list of $n - 1$ separators of k objects.
2. e.g., suppose we want to select 5 elts from a set of 4 with repetitions. Then, $** | * || **$ represents: 2 of the first element, 1 of the second, none of third and 2 of fourth.

Combinations with repetitions

Theorem

The no. of ways k elements can be chosen from n -elements, when repetition is allowed is $\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$.

1. Represent them as a list of $n - 1$ separators of k objects.
2. e.g., suppose we want to select 5 elts from a set of 4 with repetitions. Then, $** | * || **$ represents: 2 of the first element, 1 of the second, none of third and 2 of fourth.
3. There is a bijection between such lists and k -sets of n -elements with repetitions allowed (why?).

Combinations with repetitions

Theorem

The no. of ways k elements can be chosen from n -elements, when repetition is allowed is $\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$.

1. Represent them as a list of $n - 1$ separators of k objects.
2. e.g., suppose we want to select 5 elts from a set of 4 with repetitions. Then, $** | * || **$ represents: 2 of the first element, 1 of the second, none of third and 2 of fourth.
3. There is a bijection between such lists and k -sets of n -elements with repetitions allowed (why?).
4. Thus, question reduces to no. of ways to choose k stars or $n - 1$ bars from a set of $n - k + 1$ positions $= \binom{n+k-1}{k}$. \square

Combinations with repetitions

Theorem

The no. of ways k elements can be chosen from n -elements, when repetition is allowed is $\binom{n+k-1}{k} = \binom{n+k-1}{n-1}$.

1. Represent them as a list of $n - 1$ separators of k objects.
 2. e.g., suppose we want to select 5 elts from a set of 4 with repetitions. Then, $** | * || **$ represents: 2 of the first element, 1 of the second, none of third and 2 of fourth.
 3. There is a bijection between such lists and k -sets of n -elements with repetitions allowed (why?).
 4. Thus, question reduces to no. of ways to choose k stars or $n - 1$ bars from a set of $n - k + 1$ positions $= \binom{n+k-1}{k}$. \square
- **H.W-5:** How many solutions does the equation $x_1 + x_2 + x_3 + x_4 = 17$ have such that $x_1, x_2, x_3, x_4 \in \mathbb{Z}^{\geq 0}$?