

CS 207: Discrete Structures

Abstract algebra and Number theory

— subgroups, cyclic groups, group isomorphisms

Lecture 37

Oct 26 2015

Recap

Definition

A **group** is a set S along with an operator $*$ such that:

- ▶ **Closure**: $\forall a, b \in S, a * b \in S$.
- ▶ **Associativity**: $\forall a, b, c \in S, a * (b * c) = (a * b) * c$.
- ▶ **Identity**: $\exists e \in S$ s.t., $\forall a \in S, a * e = e * a = a$.
- ▶ **Inverse**: $\forall a \in S, \exists a' \in S$ s.t., $a * a' = a' * a = e$.

Recap

Definition

A **group** is a set S along with an operator $*$ such that:

- ▶ **Closure**: $\forall a, b \in S, a * b \in S$.
- ▶ **Associativity**: $\forall a, b, c \in S, a * (b * c) = (a * b) * c$.
- ▶ **Identity**: $\exists e \in S$ s.t., $\forall a \in S, a * e = e * a = a$.
- ▶ **Inverse**: $\forall a \in S, \exists a' \in S$ s.t., $a * a' = a' * a = e$.

Egs:

1. permutations of a set,
2. automorphisms of a graph,
3. symmetries of geometrical figures,
4. Numbers $(\mathbb{Z}, +)$, $(\mathbb{Q} \setminus 0, \times)$, etc
5. Modular counting $(\mathbb{Z}_p \setminus 0, \times)$,
6. Invertible matrices $GL_n(\mathbb{R})$, $SL_n(\mathbb{R})$.

Some types of groups

- ▶ If any two elements in a group commute, then the group is called a **commutative** or an **Abelian** group.

Some types of groups

- ▶ If any two elements in a group commute, then the group is called a **commutative** or an **Abelian** group.
- ▶ Let G be a group under operation $*$. A subset H of G is called a **subgroup** if H is also a group under $*$.

Some types of groups

- ▶ If any two elements in a group commute, then the group is called a **commutative** or an **Abelian** group.
- ▶ Let G be a group under operation $*$. A subset H of G is called a **subgroup** if H is also a group under $*$.

Proposition

$H \subseteq G$ is a subgroup of G iff $H \neq \emptyset$ and $\forall x, y \in H, xy^{-1} \in H$.

Some types of groups

- ▶ If any two elements in a group commute, then the group is called a **commutative** or an **Abelian** group.
- ▶ Let G be a group under operation $*$. A subset H of G is called a **subgroup** if H is also a group under $*$.

Proposition

$H \subseteq G$ is a subgroup of G iff $H \neq \emptyset$ and $\forall x, y \in H, xy^{-1} \in H$.

- ▶ The order of a finite group is the number of elements in it.

Some types of groups

- ▶ If any two elements in a group commute, then the group is called a **commutative** or an **Abelian** group.
- ▶ Let G be a group under operation $*$. A subset H of G is called a **subgroup** if H is also a group under $*$.

Proposition

$H \subseteq G$ is a subgroup of G iff $H \neq \emptyset$ and $\forall x, y \in H, xy^{-1} \in H$.

- ▶ The order of a finite group is the number of elements in it.
- ▶ If x is an element of a finite group, then the **order of x in G** is the least positive integer m such that $x^m = e$.

Some types of groups

- ▶ If any two elements in a group commute, then the group is called a **commutative** or an **Abelian** group.
- ▶ Let G be a group under operation $*$. A subset H of G is called a **subgroup** if H is also a group under $*$.

Proposition

$H \subseteq G$ is a subgroup of G iff $H \neq \emptyset$ and $\forall x, y \in H, xy^{-1} \in H$.

- ▶ The order of a finite group is the number of elements in it.
- ▶ If x is an element of a finite group, then the **order of x in G** is the least positive integer m such that $x^m = e$.

Proposition

Let x be an element of order m in a finite group G . $x^s = e$ iff

Some types of groups

- ▶ If any two elements in a group commute, then the group is called a **commutative** or an **Abelian** group.
- ▶ Let G be a group under operation $*$. A subset H of G is called a **subgroup** if H is also a group under $*$.

Proposition

$H \subseteq G$ is a subgroup of G iff $H \neq \emptyset$ and $\forall x, y \in H, xy^{-1} \in H$.

- ▶ The order of a finite group is the number of elements in it.
- ▶ If x is an element of a finite group, then the **order of x in G** is the least positive integer m such that $x^m = e$.

Proposition

Let x be an element of order m in a finite group G . $x^s = e$ iff s is a multiple of m .

Some types of groups

- ▶ If any two elements in a group commute, then the group is called a **commutative** or an **Abelian** group.
- ▶ Let G be a group under operation $*$. A subset H of G is called a **subgroup** if H is also a group under $*$.

Proposition

$H \subseteq G$ is a subgroup of G iff $H \neq \emptyset$ and $\forall x, y \in H, xy^{-1} \in H$.

- ▶ The order of a finite group is the number of elements in it.
- ▶ If x is an element of a finite group, then the **order of x in G** is the least positive integer m such that $x^m = e$.

Proposition

Let x be an element of order m in a finite group G . $x^s = e$ iff s is a multiple of m .

- ▶ What is the order of $(\mathbb{Z}_n, +_n)$; $(\mathbb{Z}_p \setminus \{0\}, \times_p)$?

Another special type of group: cyclic group

Definition

A group G is said to be **cyclic** if it contains an element x and every member of G is a power of x , i.e., obtained by repeatedly (possible zero times) applying x to itself!

Another special type of group: cyclic group

Definition

A group G is said to be **cyclic** if it contains an element x and every member of G is a power of x , i.e., obtained by repeatedly (possible zero times) applying x to itself!

- ▶ x is said to **generate** G and we write $G = \langle x \rangle$.
- ▶ If $G = \langle x \rangle$ and all powers of x , i.e., $\dots x^{-2}, x^{-1}, e, x^1, x^2, \dots$ are distinct, then G is called an **infinite cyclic group**.

Another special type of group: cyclic group

Definition

A group G is said to be **cyclic** if it contains an element x and every member of G is a power of x , i.e., obtained by repeatedly (possible zero times) applying x to itself!

- ▶ x is said to **generate** G and we write $G = \langle x \rangle$.
- ▶ If $G = \langle x \rangle$ and all powers of x , i.e., $\dots x^{-2}, x^{-1}, e, x^1, x^2, \dots$ are distinct, then G is called an **infinite cyclic group**.
- ▶ Give an example of an infinite and a finite cyclic group.

Another special type of group: cyclic group

Definition

A group G is said to be **cyclic** if it contains an element x and every member of G is a power of x , i.e., obtained by repeatedly (possible zero times) applying x to itself!

- ▶ x is said to **generate** G and we write $G = \langle x \rangle$.
- ▶ If $G = \langle x \rangle$ and all powers of x , i.e., $\dots x^{-2}, x^{-1}, e, x^1, x^2, \dots$ are distinct, then G is called an **infinite cyclic group**.
- ▶ Give an example of an infinite and a finite cyclic group.
 $(\mathbb{Z}, +)$, $(\mathbb{Z}_n, +_n)$, symmetries of a polygon?

Another special type of group: cyclic group

Definition

A group G is said to be **cyclic** if it contains an element x and every member of G is a power of x , i.e., obtained by repeatedly (possible zero times) applying x to itself!

- ▶ x is said to **generate** G and we write $G = \langle x \rangle$.
- ▶ If $G = \langle x \rangle$ and all powers of x , i.e., $\dots x^{-2}, x^{-1}, e, x^1, x^2, \dots$ are distinct, then G is called an **infinite cyclic group**.
- ▶ Give an example of an infinite and a finite cyclic group.
 $(\mathbb{Z}, +)$, $(\mathbb{Z}_n, +_n)$, rotational symmetries of a polygon

Another special type of group: cyclic group

Definition

A group G is said to be **cyclic** if it contains an element x and every member of G is a power of x , i.e., obtained by repeatedly (possible zero times) applying x to itself!

- ▶ x is said to **generate** G and we write $G = \langle x \rangle$.
- ▶ If $G = \langle x \rangle$ and all powers of x , i.e., $\dots x^{-2}, x^{-1}, e, x^1, x^2, \dots$ are distinct, then G is called an **infinite cyclic group**.
- ▶ Give an example of an infinite and a finite cyclic group. $(\mathbb{Z}, +)$, $(\mathbb{Z}_n, +_n)$, rotational symmetries of a polygon
- ▶ Are any two cyclic groups of same order the “same”?

Another special type of group: cyclic group

Definition

A group G is said to be **cyclic** if it contains an element x and every member of G is a power of x , i.e., obtained by repeatedly (possible zero times) applying x to itself!

- ▶ x is said to **generate** G and we write $G = \langle x \rangle$.
- ▶ If $G = \langle x \rangle$ and all powers of x , i.e.,
 $\dots x^{-2}, x^{-1}, e, x^1, x^2, \dots$ are distinct, then G is called an **infinite cyclic group**.
- ▶ Give an example of an infinite and a finite cyclic group.
 $(\mathbb{Z}, +)$, $(\mathbb{Z}_n, +_n)$, rotational symmetries of a polygon
- ▶ Are any two cyclic groups of same order the “same”?
- ▶ In general, when are two groups the “same”?

Another special type of group: cyclic group

Definition

A group G is said to be **cyclic** if it contains an element x and every member of G is a power of x , i.e., obtained by repeatedly (possible zero times) applying x to itself!

- ▶ x is said to **generate** G and we write $G = \langle x \rangle$.
- ▶ If $G = \langle x \rangle$ and all powers of x , i.e., $\dots x^{-2}, x^{-1}, e, x^1, x^2, \dots$ are distinct, then G is called an **infinite cyclic group**.
- ▶ Give an example of an infinite and a finite cyclic group. $(\mathbb{Z}, +)$, $(\mathbb{Z}_n, +_n)$, rotational symmetries of a polygon
- ▶ Are any two cyclic groups of same order the “same”?
- ▶ In general, when are two groups the “same”?
- ▶ What about $(\mathbb{Q} \setminus \{0\}, \times)$ and $(\mathbb{Z}, +)$?

“Sameness” in groups: group isomorphisms

Definition

- ▶ If G_1 and G_2 are groups, then a bijection $f : G_1 \rightarrow G_2$ is called an **isomorphism** if for all $g, g' \in G$,
 $f(gg') = f(g)f(g')$.
- ▶ G_1 and G_2 are said to be **isomorphic**, denoted $G_1 \equiv G_2$, if there exists an isomorphism f between them.

“Sameness” in groups: group isomorphisms

Definition

- ▶ If G_1 and G_2 are groups, then a bijection $f : G_1 \rightarrow G_2$ is called an **isomorphism** if for all $g, g' \in G$,
 $f(gg') = f(g)f(g')$.
- ▶ G_1 and G_2 are said to be **isomorphic**, denoted $G_1 \equiv G_2$, if there exists an isomorphism f between them.

Questions:

- ▶ Do $g, f(g)$ have the same order?

“Sameness” in groups: group isomorphisms

Definition

- ▶ If G_1 and G_2 are groups, then a bijection $f : G_1 \rightarrow G_2$ is called an **isomorphism** if for all $g, g' \in G$,
 $f(gg') = f(g)f(g')$.
- ▶ G_1 and G_2 are said to be **isomorphic**, denoted $G_1 \equiv G_2$, if there exists an isomorphism f between them.

Questions:

- ▶ Do $g, f(g)$ have the same order?
- ▶ Is the isomorphism relation, \equiv , an equivalence relation?

“Sameness” in groups: group isomorphisms

Definition

- ▶ If G_1 and G_2 are groups, then a bijection $f : G_1 \rightarrow G_2$ is called an **isomorphism** if for all $g, g' \in G$,
 $f(gg') = f(g)f(g')$.
- ▶ G_1 and G_2 are said to be **isomorphic**, denoted $G_1 \equiv G_2$, if there exists an isomorphism f between them.

Questions:

- ▶ Do $g, f(g)$ have the same order?
- ▶ Is the isomorphism relation, \equiv , an equivalence relation?
- ▶ Show that every infinite cyclic group is isomorphic to $(\mathbb{Z}, +)$.

“Sameness” in groups: group isomorphisms

Definition

- ▶ If G_1 and G_2 are groups, then a bijection $f : G_1 \rightarrow G_2$ is called an **isomorphism** if for all $g, g' \in G$,
 $f(gg') = f(g)f(g')$.
- ▶ G_1 and G_2 are said to be **isomorphic**, denoted $G_1 \equiv G_2$, if there exists an isomorphism f between them.

Questions:

- ▶ Do $g, f(g)$ have the same order?
- ▶ Is the isomorphism relation, \equiv , an equivalence relation?
- ▶ Show that every infinite cyclic group is isomorphic to $(\mathbb{Z}, +)$.
- ▶ Which of the following are isomorphic:
 $(\mathbb{R}, +), (\mathbb{Z}, +), (\mathbb{R}^+, \times), (\mathbb{C}, +)$?