


CS 207: Discrete Structures

Lecture 15 – Counting and Combinatorics Solving Recurrence relations and generating functions


Aug 20 2015

Find the Fibonacci sequence!



- ▶ $F(n) = F(n-1) + F(n-2)$.
- ▶ 1, 1, 2, 3, 5, 8, 13,

Find the Fibonacci sequence!



- ▶ $F(n) = F(n - 1) + F(n - 2)$.
- ▶ 1, 1, 2, 3, 5, 8, 13,

Last few classes

Basic counting techniques and applications

1. Sum and product, bijection, double counting principles
2. Binomial coefficients and binomial theorem, Pascal's triangle
3. Permutations and combinations with/without repetitions
4. Counting subsets, relations, partitions, Handshake lemma
5. Stirling's approximation: Estimating $n!$
6. Recurrence relations

Last few classes

Basic counting techniques and applications

1. Sum and product, bijection, double counting principles
2. Binomial coefficients and binomial theorem, Pascal's triangle
3. Permutations and combinations with/without repetitions
4. Counting subsets, relations, partitions, Handshake lemma
5. Stirling's approximation: Estimating $n!$
6. Recurrence relations

Today

Solving recurrence relations and generating functions.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation...

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation... \therefore if we find any solutions, we can take linear combinations.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation... \therefore if we find any solutions, we can take linear combinations.
2. i.e., if $F_n = F_{n-1} - F_{n-2}$, $G_n = G_{n-1} - G_{n-2}$ and $H_n = aF_n + bG_n$, then what about H_n ?

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation... \therefore if we find any solutions, we can take linear combinations.
2. i.e., if $F_n = F_{n-1} - F_{n-2}$, $G_n = G_{n-1} - G_{n-2}$ and $H_n = aF_n + bG_n$, then $H_n = H_{n-1} - H_{n-2}$.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

1. This is a linear recurrence relation... \therefore if we find any solutions, we can take linear combinations.
2. i.e., if $F_n = F_{n-1} - F_{n-2}$, $G_n = G_{n-1} - G_{n-2}$ and $H_n = aF_n + bG_n$, then $H_n = H_{n-1} - H_{n-2}$.

Proof method 1 (for linear recurrences: try α^n !)

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it!

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
5. Thus, general solution is $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$.

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
5. Thus, general solution is $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$.
6. How do we get a and b ?

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
5. Thus, general solution is $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$.
6. Use F_0 and F_1 – initial conditions: $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

How to “solve” the Fibonacci recurrence?

By solving, we mean give a closed-form expression for n^{th} term.

Fibonacci recurrence relation

For $n \geq 2$, $F_n = F_{n-1} + F_{n-2}$, $F_0 = F_1 = 1$.

Proof method 1 (for linear recurrences: try α^n !)

1. try $F_n = \alpha^n \dots$
2. $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$ implies $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$.
3. So if $\alpha^2 - \alpha - 1 = 0$, the recurrence holds for all n .
4. Solve it! $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
5. Thus, general solution is $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$.
6. Use F_0 and F_1 – initial conditions: $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

$$\text{Thus, } F_n = \frac{\sqrt{5}+1}{2\sqrt{5}} \left(\frac{1+\sqrt{5}}{2} \right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left(\frac{1-\sqrt{5}}{2} \right)^n.$$

Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?

Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?
- ▶ Find a solution for $a_n = 6a_{n-1} - 9a_{n-2}$, with $a_0 = 1, a_1 = 6$? Can you apply the same method for this?

Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?
- ▶ Find a solution for $a_n = 6a_{n-1} - 9a_{n-2}$, with $a_0 = 1, a_1 = 6$? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of n terms s.t. it can be computed by adding two numbers at a time?

Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?
- ▶ Find a solution for $a_n = 6a_{n-1} - 9a_{n-2}$, with $a_0 = 1, a_1 = 6$? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of n terms s.t. it can be computed by adding two numbers at a time?

This method does not work if we have repeated roots (but this can be fixed) and non-linear recurrences.

Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?
- ▶ Find a solution for $a_n = 6a_{n-1} - 9a_{n-2}$, with $a_0 = 1, a_1 = 6$? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of n terms s.t. it can be computed by adding two numbers at a time?

This method does not work if we have repeated roots (but this can be fixed) and non-linear recurrences.

Reading assignment

Read examples/generalizations from Sections 6.1 and 6.2 from Rosen's book (6th Edition).

Solving general linear recurrence sequences

- ▶ (C.W) Find a solution for $a_n = a_{n-1} + 2a_{n-2}$ with $a_0 = 2$, $a_1 = 7$?
- ▶ Find a solution for $a_n = 6a_{n-1} - 9a_{n-2}$, with $a_0 = 1, a_1 = 6$? Can you apply the same method for this? What went wrong?
- ▶ Recall the recurrence for Catalan Numbers:

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = C(1) = 1.$$

No. of ways to bracket a sum of n terms s.t. it can be computed by adding two numbers at a time?

This method does not work if we have repeated roots (but this can be fixed) and non-linear recurrences.

Reading assignment

Read examples/generalizations from Sections 6.1 and 6.2 from Rosen's book (6th Edition).

We next consider a method of much wider applicability...

Proof Method 2: Using generating functions

Yet another way to count number of subsets of a set.

Recurrence: $F(n) = 2 \cdot F(n - 1), F(0) = 1.$

Proof Method 2: Using generating functions

Yet another way to count number of subsets of a set.

Recurrence: $F(n) = 2 \cdot F(n-1), F(0) = 1.$

- Set $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$. (does it converge?!)

Proof Method 2: Using generating functions

Yet another way to count number of subsets of a set.

Recurrence: $F(n) = 2 \cdot F(n-1), F(0) = 1$.

- Set $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$. (does it converge?!)

$$\begin{aligned}\text{Now, } 2t\phi(t) &= \sum_{n=0}^{\infty} 2 \cdot F(n)t^{n+1} \\ &= \sum_{n=0}^{\infty} F(n+1)t^{n+1} \\ &= \phi(t) - 1\end{aligned}$$

Proof Method 2: Using generating functions

Yet another way to count number of subsets of a set.

Recurrence: $F(n) = 2 \cdot F(n-1), F(0) = 1$.

- Set $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$. (does it converge?!)

$$\begin{aligned}\text{Now, } 2t\phi(t) &= \sum_{n=0}^{\infty} 2 \cdot F(n)t^{n+1} \\ &= \sum_{n=0}^{\infty} F(n+1)t^{n+1} \\ &= \phi(t) - 1\end{aligned}$$

- Thus, $\phi(t) = \frac{1}{1-2t} = \sum_{n=0}^{\infty} (2t)^n$.

Proof Method 2: Using generating functions

Yet another way to count number of subsets of a set.

Recurrence: $F(n) = 2 \cdot F(n-1), F(0) = 1$.

- ▶ Set $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$. (does it converge?!)

$$\begin{aligned}\text{Now, } 2t\phi(t) &= \sum_{n=0}^{\infty} 2 \cdot F(n)t^{n+1} \\ &= \sum_{n=0}^{\infty} F(n+1)t^{n+1} \\ &= \phi(t) - 1\end{aligned}$$

- ▶ Thus, $\phi(t) = \frac{1}{1-2t} = \sum_{n=0}^{\infty} (2t)^n$.
- ▶ Comparing to original series, we conclude $F(n) = 2^n$.
- ▶ (If two power series are equal, then all coeffs coincide).

Proof Method 2: Using generating functions

Yet another way to count number of subsets of a set.

Recurrence: $F(n) = 2 \cdot F(n-1), F(0) = 1$.

- ▶ Set $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$. (does it converge?!)

$$\begin{aligned}\text{Now, } 2t\phi(t) &= \sum_{n=0}^{\infty} 2 \cdot F(n)t^{n+1} \\ &= \sum_{n=0}^{\infty} F(n+1)t^{n+1} \\ &= \phi(t) - 1\end{aligned}$$

- ▶ Thus, $\phi(t) = \frac{1}{1-2t} = \sum_{n=0}^{\infty} (2t)^n$.
- ▶ Comparing to original series, we conclude $F(n) = 2^n$.
- ▶ (If two power series are equal, then all coeffs coincide).
- ▶ $\forall t$ with $|t| < 1/2$, power series does converge by analysis.