

# CS 207: Discrete Structures

## Lecture 16 – Counting and Combinatorics Solving Recurrence relations and generating functions

Aug 24 2015

## Solving recurrence relations

By solving, we mean give a closed-form expression for  $n^{th}$  term.

## Solving recurrence relations

By solving, we mean give a closed-form expression for  $n^{th}$  term.

Fibonacci recurrence:  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2},$   
 $F_0 = F_1 = 1.$

# Solving recurrence relations

By solving, we mean give a closed-form expression for  $n^{th}$  term.

Fibonacci recurrence:  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2}$ ,  
 $F_0 = F_1 = 1$ .

Proof method 1 (for linear recurrences: try  $F_n = \alpha^n$ !)

1.  $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$  implies  $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$ .
2. So if  $\alpha^2 - \alpha - 1 = 0$ , the recurrence holds for all  $n$ .
3. Solving,  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
4. Thus, general solution is  $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$ .
5. Use  $F_0$  and  $F_1$  – initial conditions:  $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

# Solving recurrence relations

By solving, we mean give a closed-form expression for  $n^{th}$  term.

Fibonacci recurrence:  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2},$   
 $F_0 = F_1 = 1.$

Proof method 1 (for linear recurrences: try  $F_n = \alpha^n$ )

1.  $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$  implies  $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0.$
2. So if  $\alpha^2 - \alpha - 1 = 0$ , the recurrence holds for all  $n.$
3. Solving,  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
4. Thus, general solution is  $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n.$
5. Use  $F_0$  and  $F_1$  – initial conditions:  $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

$$\text{Thus, } F_n = \frac{\sqrt{5}+1}{2\sqrt{5}} \left( \frac{1+\sqrt{5}}{2} \right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left( \frac{1-\sqrt{5}}{2} \right)^n.$$

# Solving recurrence relations

By solving, we mean give a closed-form expression for  $n^{\text{th}}$  term.

Fibonacci recurrence:  $\forall n \geq 2, F_n = F_{n-1} + F_{n-2}$ ,  
 $F_0 = F_1 = 1$ .

Proof method 1 (for linear recurrences: try  $F_n = \alpha^n$ )

1.  $\alpha^n = \alpha^{n-1} + \alpha^{n-2}$  implies  $\alpha^{n-2}(\alpha^2 - \alpha - 1) = 0$ .
2. So if  $\alpha^2 - \alpha - 1 = 0$ , the recurrence holds for all  $n$ .
3. Solving,  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$
4. Thus, general solution is  $F_n = a(\frac{1+\sqrt{5}}{2})^n + b(\frac{1-\sqrt{5}}{2})^n$ .
5. Use  $F_0$  and  $F_1$  – initial conditions:  $a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

$$\text{Thus, } F_n = \frac{\sqrt{5}+1}{2\sqrt{5}} \left(\frac{1+\sqrt{5}}{2}\right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left(\frac{1-\sqrt{5}}{2}\right)^n.$$

But this method may not work if we have repeated roots (but this can be fixed) and non-linear recurrences.

## Proof Method 2: Using generating functions

### Fibonacci recurrence relation

For  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = F_1 = 1$ .

Compute  $F_n$  in terms of  $n$ .

- Consider the power series...  $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$ .

## Proof Method 2: Using generating functions

### Fibonacci recurrence relation

For  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = F_1 = 1$ .

Compute  $F_n$  in terms of  $n$ .

- Consider the power series...  $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$ .

$$t\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+1} = \sum_{n=1}^{\infty} F(n-1)t^n$$



## Proof Method 2: Using generating functions

### Fibonacci recurrence relation

For  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = F_1 = 1$ .

Compute  $F_n$  in terms of  $n$ .

- Consider the power series...  $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$ .

$$t\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+1} = \sum_{n=1}^{\infty} F(n-1)t^n$$

$$t^2\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+2} = \sum_{n=2}^{\infty} F(n-2)t^n$$

# Proof Method 2: Using generating functions

## Fibonacci recurrence relation

For  $n \geq 2$ ,  $F_n = F_{n-1} + F_{n-2}$ ,  $F_0 = F_1 = 1$ .

Compute  $F_n$  in terms of  $n$ .

► Consider the power series...  $\phi(t) = \sum_{n=0}^{\infty} F(n)t^n$ .

$$t\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+1} = \sum_{n=1}^{\infty} F(n-1)t^n$$

$$t^2\phi(t) = \sum_{n=0}^{\infty} F(n)t^{n+2} = \sum_{n=2}^{\infty} F(n-2)t^n$$

$$(t + t^2)\phi(t) = \sum_{n=0}^{\infty} F(n)t^n - 1$$

$$(t + t^2)\phi(t) = \phi(t) - 1$$

## Proof Method 2: Using generating functions

- ▶  $\phi(t) = \frac{1}{1-t-t^2}$

## Proof Method 2: Using generating functions

$$\blacktriangleright \phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)}$$

## Proof Method 2: Using generating functions

- ▶  $\phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)} = \frac{a}{1-\alpha t} + \frac{b}{1-\beta t}$
- ▶ Solving we get  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}, a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$

## Proof Method 2: Using generating functions

- ▶  $\phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)} = \frac{a}{1-\alpha t} + \frac{b}{1-\beta t}$
- ▶ Solving we get  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}, a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$
- ▶ Now  $\phi(t) = a(1 + \alpha t + \alpha^2 t^2 + \dots) + b(1 + \beta t + \beta^2 t^2 + \dots)$

## Proof Method 2: Using generating functions

- ▶  $\phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)} = \frac{a}{1-\alpha t} + \frac{b}{1-\beta t}$
- ▶ Solving we get  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}, a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$
- ▶ Now  $\phi(t) = a(1 + \alpha t + \alpha^2 t^2 + \dots) + b(1 + \beta t + \beta^2 t^2 + \dots)$
- ▶ Equating coefficients of  $t^n$  we get  $F_n = a\alpha^n + b\beta^n$ .

## Proof Method 2: Using generating functions

- ▶  $\phi(t) = \frac{1}{1-t-t^2} = \frac{1}{(1-\alpha t)(1-\beta t)} = \frac{a}{1-\alpha t} + \frac{b}{1-\beta t}$
- ▶ Solving we get  $\alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}, a = \frac{\sqrt{5}+1}{2\sqrt{5}}, b = \frac{\sqrt{5}-1}{2\sqrt{5}}$
- ▶ Now  $\phi(t) = a(1 + \alpha t + \alpha^2 t^2 + \dots) + b(1 + \beta t + \beta^2 t^2 + \dots)$
- ▶ Equating coefficients of  $t^n$  we get  $F_n = a\alpha^n + b\beta^n$ .

thus, as before

$$F(n) = \frac{\sqrt{5}+1}{2\sqrt{5}} \left( \frac{1+\sqrt{5}}{2} \right)^n + \frac{\sqrt{5}-1}{2\sqrt{5}} \left( \frac{1-\sqrt{5}}{2} \right)^n$$



# Properties of generating functions

## Definition

The **(ordinary) generating function** for a sequence  $a_0, a_1, \dots \in \mathbb{R}$  is the infinite series  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .

- Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ ,  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ . Then
1. If  $f(x) = g(x)$ , then  $a_k = b_k$  for all  $k$ .
  2.  $f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$ ,
  3.  $f(x)g(x) = \sum_{k=0}^{\infty} (\sum_{j=0}^k a_j b_{k-j}) x^k$ ,
  4.  $\frac{d}{dx} (\sum_{k=0}^{\infty} a_k x^k) = \sum_{k=1}^{\infty} (k a_k) x^{k-1}$

# Properties of generating functions

## Definition

The (ordinary) generating function for a sequence  $a_0, a_1, \dots \in \mathbb{R}$  is the infinite series  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .

- ▶ Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ ,  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ . Then
  1. If  $f(x) = g(x)$ , then  $a_k = b_k$  for all  $k$ .
  2.  $f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$ ,
  3.  $f(x)g(x) = \sum_{k=0}^{\infty} (\sum_{j=0}^k a_j b_{k-j}) x^k$ ,
  4.  $\frac{d}{dx} (\sum_{k=0}^{\infty} a_k x^k) = \sum_{k=1}^{\infty} (k a_k) x^{k-1}$
- ▶ Let  $u \in \mathbb{R}$ ,  $k \in \mathbb{Z}^{\geq 0}$ , Then extended binomial coefficient  $\binom{u}{k}$  is defined as  $\binom{u}{k} = \frac{u(u-1)\dots(u-k+1)}{k!}$  if  $k > 0$  and  $= 1$  if  $k = 0$ .
- ▶ What if  $u = -n$  for  $n \in \mathbb{N}$ ?

# Properties of generating functions

## Definition

The **(ordinary) generating function** for a sequence  $a_0, a_1, \dots \in \mathbb{R}$  is the infinite series  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .

- ▶ Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ ,  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ . Then
  1. If  $f(x) = g(x)$ , then  $a_k = b_k$  for all  $k$ .
  2.  $f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$ ,
  3.  $f(x)g(x) = \sum_{k=0}^{\infty} (\sum_{j=0}^k a_j b_{k-j}) x^k$ ,
  4.  $\frac{d}{dx} (\sum_{k=0}^{\infty} a_k x^k) = \sum_{k=1}^{\infty} (k a_k) x^{k-1}$
- ▶ Let  $u \in \mathbb{R}$ ,  $k \in \mathbb{Z}^{\geq 0}$ , Then **extended binomial coefficient**  $\binom{u}{k}$  is defined as  $\binom{u}{k} = \frac{u(u-1)\dots(u-k+1)}{k!}$  if  $k > 0$  and  $= 1$  if  $k = 0$ .
- ▶ What if  $u = -n$  for  $n \in \mathbb{N}$ ?

## The extended binomial theorem

Let  $u \in \mathbb{R}$ ,  $(1+x)^u = \sum_{k=0}^{\infty} \binom{u}{k} x^k$ .

# Properties of generating functions

## Definition

The (ordinary) generating function for a sequence  $a_0, a_1, \dots \in \mathbb{R}$  is the infinite series  $\phi(x) = \sum_{k=0}^{\infty} a_k x^k$ .

- ▶ Let  $f(x) = \sum_{k=0}^{\infty} a_k x^k$ ,  $g(x) = \sum_{k=0}^{\infty} b_k x^k$ . Then
  1. If  $f(x) = g(x)$ , then  $a_k = b_k$  for all  $k$ .
  2.  $f(x) + g(x) = \sum_{k=0}^{\infty} (a_k + b_k) x^k$ ,
  3.  $f(x)g(x) = \sum_{k=0}^{\infty} (\sum_{j=0}^k a_j b_{k-j}) x^k$ ,
  4.  $\frac{d}{dx} (\sum_{k=0}^{\infty} a_k x^k) = \sum_{k=1}^{\infty} (k a_k) x^{k-1}$
- ▶ Let  $u \in \mathbb{R}$ ,  $k \in \mathbb{Z}^{\geq 0}$ , Then extended binomial coefficient  $\binom{u}{k}$  is defined as  $\binom{u}{k} = \frac{u(u-1)\dots(u-k+1)}{k!}$  if  $k > 0$  and  $= 1$  if  $k = 0$ .
- ▶ What if  $u = -n$  for  $n \in \mathbb{N}$ ?

## The extended binomial theorem

Let  $u \in \mathbb{R}$ ,  $(1+x)^u = \sum_{k=0}^{\infty} \binom{u}{k} x^k$ .

If you don't like this, take  $x \in \mathbb{R}$ ,  $|x| < 1$ .

# Simple examples using generating functions

Standard identities:

- ▶  $\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k$
- ▶  $\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk}$
- ▶  $e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$

# Simple examples using generating functions

## Standard identities:

- ▶  $\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k$
- ▶  $\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk}$
- ▶  $e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$

## Class work:

1. Solve the recurrence  $a_k = 4a_{k-1}$  with  $a_0 = 3$ .

# Simple examples using generating functions

## Standard identities:

- ▶  $\frac{1}{1-ax} = \sum_{k=0}^{\infty} a^k x^k$
- ▶  $\frac{1}{1-x^r} = \sum_{k=0}^{\infty} x^{rk}$
- ▶  $e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$

## Class work:

1. Solve the recurrence  $a_k = 4a_{k-1}$  with  $a_0 = 3$ .
2. Solve the recurrence  $a_k = 8a_{k-1} + 10^{k-1}$  with  $a_0 = 1, a_1 = 9$ .

# Solving Catalan numbers using generating functions

## Catalan Numbers

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = 0, C(1) = 1.$$



# Solving Catalan numbers using generating functions

## Catalan Numbers

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = 0, C(1) = 1.$$

- Let  $\phi(x) = \sum_{k=1}^{\infty} C(k)x^k$ .

# Solving Catalan numbers using generating functions

## Catalan Numbers

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = 0, C(1) = 1.$$

- ▶ Let  $\phi(x) = \sum_{k=1}^{\infty} C(k)x^k$ .
- ▶ Now consider  $\phi(x)^2$ .
- ▶ 
$$\begin{aligned}\phi(x)^2 &= \left(\sum_{k=1}^{\infty} C(k)x^k\right)\left(\sum_{k=1}^{\infty} C(k)x^k\right) \\ &= \left(\sum_{k=2}^{\infty} \sum_{i=1}^{k-1} C(i)C(k-i)x^k\right) \\ &= \left(\sum_{k=2}^{\infty} C(k)x^k\right) = \phi(x) - x\end{aligned}$$

# Solving Catalan numbers using generating functions

## Catalan Numbers

$$C(n) = \sum_{i=1}^{n-1} C(i)C(n-i) \text{ for } n > 1, C(0) = 0, C(1) = 1.$$

- ▶ Let  $\phi(x) = \sum_{k=1}^{\infty} C(k)x^k$ .
- ▶ Now consider  $\phi(x)^2$ .
- ▶ 
$$\begin{aligned}\phi(x)^2 &= \left(\sum_{k=1}^{\infty} C(k)x^k\right)\left(\sum_{k=1}^{\infty} C(k)x^k\right) \\ &= \left(\sum_{k=2}^{\infty} \sum_{i=1}^{k-1} C(i)C(k-i)x^k\right) \\ &= \left(\sum_{k=2}^{\infty} C(k)x^k\right) = \phi(x) - x\end{aligned}$$
- ▶ Solving for  $\phi(x)$  we get,  $\phi(x) = \frac{1}{2}(1 \pm (1 - 4x)^{1/2})$
- ▶ But since  $\phi(0) = 0$ , we have
$$\phi(x) = \frac{1}{2}(1 - (1 - 4x)^{1/2}) = \frac{1}{2} + (-\frac{1}{2}(1 - 4x)^{1/2}).$$

# Catalan numbers

Recall: Extended binomial theorem

Let  $\alpha \in \mathbb{R}$ ,  $(1+x)^\alpha = \sum_{n=0}^{\infty} \binom{\alpha}{n} x^n$ , where  $\binom{\alpha}{n} = \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}$ .

# Catalan numbers

Recall: Extended binomial theorem

Let  $\alpha \in \mathbb{R}$ ,  $(1+x)^\alpha = \sum_{n=0}^{\infty} \binom{\alpha}{n} x^n$ , where  $\binom{\alpha}{n} = \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}$ .

$$\blacktriangleright \sum_{k=1}^{\infty} C(k)x^k = \phi(x) = \frac{1}{2} + (-\frac{1}{2}(1-4x)^{1/2}) = \frac{1}{2} + (-\frac{1}{2} \sum_{k=0}^{\infty} \binom{1/2}{k} (-4x)^k).$$

# Catalan numbers

## Recall: Extended binomial theorem

Let  $\alpha \in \mathbb{R}$ ,  $(1+x)^\alpha = \sum_{n=0}^{\infty} \binom{\alpha}{n} x^n$ , where  $\binom{\alpha}{n} = \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}$ .

- ▶  $\sum_{k=1}^{\infty} C(k)x^k = \phi(x) = \frac{1}{2} + (-\frac{1}{2}(1-4x)^{1/2}) = \frac{1}{2} + (-\frac{1}{2} \sum_{k=0}^{\infty} \binom{1/2}{k} (-4x)^k).$
- ▶ The coefficient of  $x^k$  is  $C(k) = -\frac{1}{2} \binom{1/2}{k} (-4)^k$   
 $= -\frac{1}{2} (\frac{1}{2}(\frac{1}{2}-1)(\frac{1}{2}-2)\dots(\frac{1}{2}-k+1)) \frac{(-4)^k}{k!}$   
 $= -\frac{1}{2} (\frac{1}{2})(-\frac{3}{2})(-\frac{5}{2})\dots(-\frac{2k-3}{2}) \frac{(-4)^k}{k!}$

# Catalan numbers

## Recall: Extended binomial theorem

Let  $\alpha \in \mathbb{R}$ ,  $(1+x)^\alpha = \sum_{n=0}^{\infty} \binom{\alpha}{n} x^n$ , where  $\binom{\alpha}{n} = \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}$ .

- ▶  $\sum_{k=1}^{\infty} C(k)x^k = \phi(x) = \frac{1}{2} + (-\frac{1}{2}(1-4x)^{1/2}) = \frac{1}{2} + (-\frac{1}{2} \sum_{k=0}^{\infty} \binom{1/2}{k} (-4x)^k).$
- ▶ The coefficient of  $x^k$  is  $C(k) = -\frac{1}{2} \binom{1/2}{k} (-4)^k$   
 $= -\frac{1}{2} (\frac{1}{2}(\frac{1}{2}-1)(\frac{1}{2}-2)\dots(\frac{1}{2}-k+1)) \frac{(-4)^k}{k!}$   
 $= -\frac{1}{2} (\frac{1}{2})(-\frac{3}{2})(-\frac{5}{2})\dots(-\frac{2k-3}{2}) \frac{(-4)^k}{k!}$
- ▶  $C(k) = \frac{(-1)^k (-4)^k}{2^{k+1} k!} \cdot 1 \cdot 3 \dots (2k-3)$

# Catalan numbers

## Recall: Extended binomial theorem

Let  $\alpha \in \mathbb{R}$ ,  $(1+x)^\alpha = \sum_{n=0}^{\infty} \binom{\alpha}{n} x^n$ , where  $\binom{\alpha}{n} = \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}$ .

- ▶  $\sum_{k=1}^{\infty} C(k)x^k = \phi(x) = \frac{1}{2} + (-\frac{1}{2}(1-4x)^{1/2}) = \frac{1}{2} + (-\frac{1}{2} \sum_{k=0}^{\infty} \binom{1/2}{k} (-4x)^k)$ .
- ▶ The coefficient of  $x^k$  is  $C(k) = -\frac{1}{2} \binom{1/2}{k} (-4)^k$   
 $= -\frac{1}{2} (\frac{1}{2}(\frac{1}{2}-1)(\frac{1}{2}-2)\dots(\frac{1}{2}-k+1)) \frac{(-4)^k}{k!}$   
 $= -\frac{1}{2} (\frac{1}{2})(-\frac{3}{2})(-\frac{5}{2})\dots(-\frac{2k-3}{2}) \frac{(-4)^k}{k!}$
- ▶  $C(k) = \frac{(-1)^k (-4)^k}{2^{k+1} k!} \cdot 1 \cdot 3 \dots (2k-3)$
- ▶  $C(k) = \frac{1 \cdot 4^k}{2^{k+1} \cdot k!} \cdot \frac{1 \cdot 2 \dots (2k-3)(2k-2)}{2^{k-1} (k-1)!} = \frac{(2k-2)!}{k!(k-1)!}$ .



# Catalan numbers

## Recall: Extended binomial theorem

Let  $\alpha \in \mathbb{R}$ ,  $(1+x)^\alpha = \sum_{n=0}^{\infty} \binom{\alpha}{n} x^n$ , where  $\binom{\alpha}{n} = \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!}$ .

- ▶  $\sum_{k=1}^{\infty} C(k)x^k = \phi(x) = \frac{1}{2} + (-\frac{1}{2}(1-4x)^{1/2}) = \frac{1}{2} + (-\frac{1}{2} \sum_{k=0}^{\infty} \binom{1/2}{k} (-4x)^k)$ .
- ▶ The coefficient of  $x^k$  is  $C(k) = -\frac{1}{2} \binom{1/2}{k} (-4)^k$   
 $= -\frac{1}{2} (\frac{1}{2}(\frac{1}{2}-1)(\frac{1}{2}-2)\dots(\frac{1}{2}-k+1)) \frac{(-4)^k}{k!}$   
 $= -\frac{1}{2} (\frac{1}{2})(-\frac{3}{2})(-\frac{5}{2})\dots(-\frac{2k-3}{2}) \frac{(-4)^k}{k!}$
- ▶  $C(k) = \frac{(-1)^k (-4)^k}{2^{k+1} k!} \cdot 1 \cdot 3 \dots (2k-3)$
- ▶  $C(k) = \frac{1 \cdot 4^k}{2^{k+1} \cdot k!} \cdot \frac{1 \cdot 2 \dots (2k-3)(2k-2)}{2^{k-1} (k-1)!} = \frac{(2k-2)!}{k!(k-1)!}$ .

Thus, the  $n^{\text{th}}$  Catalan number is given by

$$C(n) = \frac{(2n-2)!}{n!(n-1)!} = \frac{1}{n} \binom{2n-2}{n-1}$$

## Other examples

- ▶ (H.W) Write a recurrence for the number of derangements. That is, no. of ways to arrange  $n$  letters into  $n$  addressed envelopes such that no letter goes to the correct envelope.
- ▶ (H.W) How many ways can a convex  $n$ -sided polygon be cut into triangles by adding non-intersecting diagonals (i.e., connecting vertices with non-crossing lines)? Write a recurrence and solve it!