

# **CS 228 : Logic in Computer Science**

Krishna. S

# Summary

---

- ▶ Started looking at FO nondefinability
- ▶ Defined quantifier depth or quantifier rank of a formula
- ▶ Showed that there are finitely many FO formulae of quantifier rank  $r$
- ▶ Introduced some new notations for words, mimicking assignments of values to free variables

# FO Definability

---

Let  $\varphi$  be a FO formula. Define the **quantifier rank** of  $\varphi$  denoted  $c(\varphi)$

- ▶ If  $\varphi$  is atomic ( $x = y, x < y, S(x, y), Q_a(x)$ ) then  $c(\varphi) = 0$

# FO Definability

---

Let  $\varphi$  be a FO formula. Define the **quantifier rank** of  $\varphi$  denoted  $c(\varphi)$

- ▶ If  $\varphi$  is atomic ( $x = y, x < y, S(x, y), Q_a(x)$ ) then  $c(\varphi) = 0$
- ▶  $c(\neg\varphi) = c(\varphi)$

# FO Definability

---

Let  $\varphi$  be a FO formula. Define the **quantifier rank** of  $\varphi$  denoted  $c(\varphi)$

- ▶ If  $\varphi$  is atomic ( $x = y$ ,  $x < y$ ,  $S(x, y)$ ,  $Q_a(x)$ ) then  $c(\varphi) = 0$
- ▶  $c(\neg\varphi) = c(\varphi)$
- ▶  $c(\varphi \wedge \psi) = \max(c(\varphi), c(\psi))$

# FO Definability

---

Let  $\varphi$  be a FO formula. Define the **quantifier rank** of  $\varphi$  denoted  $c(\varphi)$

- ▶ If  $\varphi$  is atomic ( $x = y$ ,  $x < y$ ,  $S(x, y)$ ,  $Q_a(x)$ ) then  $c(\varphi) = 0$
- ▶  $c(\neg\varphi) = c(\varphi)$
- ▶  $c(\varphi \wedge \psi) = \max(c(\varphi), c(\psi))$
- ▶  $c(\exists\varphi) = c(\varphi) + 1$

# FO Definability

---

Let  $\varphi$  be a FO formula. Define the **quantifier rank** of  $\varphi$  denoted  $c(\varphi)$

- ▶ If  $\varphi$  is atomic ( $x = y, x < y, S(x, y), Q_a(x)$ ) then  $c(\varphi) = 0$
- ▶  $c(\neg\varphi) = c(\varphi)$
- ▶  $c(\varphi \wedge \psi) = \max(c(\varphi), c(\psi))$
- ▶  $c(\exists\varphi) = c(\varphi) + 1$
- ▶ Quantifier free formulae written in DNF :  $C_1 \vee C_2 \vee \dots \vee C_n$

# FO Definability

---

Let  $\varphi$  be a FO formula. Define the **quantifier rank** of  $\varphi$  denoted  $c(\varphi)$

- ▶ If  $\varphi$  is atomic ( $x = y, x < y, S(x, y), Q_a(x)$ ) then  $c(\varphi) = 0$
- ▶  $c(\neg\varphi) = c(\varphi)$
- ▶  $c(\varphi \wedge \psi) = \max(c(\varphi), c(\psi))$
- ▶  $c(\exists\varphi) = c(\varphi) + 1$
- ▶ Quantifier free formulae written in DNF :  $C_1 \vee C_2 \vee \dots \vee C_n$
- ▶ Formulae of quantifier rank  $c + 1$  written as a disjunction of the conjunction of formulae, each formula of the form  $\exists x\varphi, \neg\exists x\varphi$  or  $\varphi$ , with  $c(\varphi) \leq c$ . Eliminate repeated disjuncts/conjuncts



# Number of FO formulae of rank $\leq c$

---

Let  $\mathcal{V}$  be a finite set of first order variables. Fix a finite signature  $\tau$ . Let there be  $m$  atomic formulae over  $\tau$  having variables from  $\mathcal{V}$ .

# Number of FO formulae of rank $\leq c$

---

Let  $\mathcal{V}$  be a finite set of first order variables. Fix a finite signature  $\tau$ . Let there be  $m$  atomic formulae over  $\tau$  having variables from  $\mathcal{V}$ .

- ▶ If  $\mathcal{V}$  has 2 variables  $x, y$ , and  $\tau$  has  $Q_a, S, <$ .
- ▶ Atomic formulae :  $\{Q_a(x), Q_a(y), S(x, y), x < y\}$
- ▶  $G = \{Q_a(x), \neg Q_a(x), Q_a(y), \neg Q_a(y), S(x, y), \neg S(x, y), x < y, \neg(x < y)\}$
- ▶ Each subset of  $G$  is a possible conjunct  $C_i$ .
- ▶ All possible disjuncts using each  $C_i$  : formulae in DNF of rank 0

# Number of FO formulae of rank $\leq c$

---

Let  $\mathcal{V}$  be a finite set of first order variables. Fix a finite signature  $\tau$ . Let there be  $m$  atomic formulae over  $\tau$  having variables from  $\mathcal{V}$ .

# Number of FO formulae of rank $\leq c$

---

Let  $\mathcal{V}$  be a finite set of first order variables. Fix a finite signature  $\tau$ . Let there be  $m$  atomic formulae over  $\tau$  having variables from  $\mathcal{V}$ .

- ▶  $2m$  atomic/negated atomic formulae

# Number of FO formulae of rank $\leq c$

---

Let  $\mathcal{V}$  be a finite set of first order variables. Fix a finite signature  $\tau$ . Let there be  $m$  atomic formulae over  $\tau$  having variables from  $\mathcal{V}$ .

- ▶  $2m$  atomic/negated atomic formulae
- ▶ Number of conjunctions  $C_i$  possible  $\leq 2^{2m}$

# Number of FO formulae of rank $\leq c$

---

Let  $\mathcal{V}$  be a finite set of first order variables. Fix a finite signature  $\tau$ . Let there be  $m$  atomic formulae over  $\tau$  having variables from  $\mathcal{V}$ .

- ▶  $2m$  atomic/negated atomic formulae
- ▶ Number of conjunctions  $C_i$  possible  $\leq 2^{2m}$
- ▶ Number of formulae in DNF  $\leq 2^{2^{2m}}$  ( $c = 0$ )

# Rank 1

---

Let there be  $p$  formulae  $\varphi$  of rank 0.

- ▶  $2p$  formulae of the form  $\exists x\varphi, \neg\exists x\varphi$
- ▶  $2^{2p}$  conjunctions of rank 1
- ▶ Conjoining any one of the  $p$  formulae of rank 0 gives all conjuncts of rank  $\leq 1$  :  $p2^{2p}$  more
- ▶ Possible conjuncts of rank  $\leq 1$  is  $q = (p + 1)2^{2p}$
- ▶ Possible disjuncts of these :  $2^q$

# Number of FO formulae of rank $c$

---

Let  $\mathcal{V}$  be a finite set of first order variables, and let  $c \geq 0$ . There are finitely many FO formulae in DNF with rank  $c$  over  $\mathcal{V}$ .



# Some Notation

---

Given a word  $w = a_1 \dots a_n$ , and a finite set of variables  $\mathcal{V}$ , define a  $\mathcal{V}$ -structure with respect to  $w$  as

- ▶  $(a_1, U_1)(a_2, U_2) \dots (a_n, U_n)$  where
- ▶  $\bigcup_i U_i = \mathcal{V}$
- ▶  $U_i \cap U_j = \emptyset$

- ▶ Think of a  $\mathcal{V}$ -structure as a word over the alphabet  $\Sigma \times 2^{\mathcal{V}}$
- ▶  $(a, \{x\})(b, \{y, z\})(c, \emptyset)(d, \{u, v\})$  is a  $\{x, y, z, u, v\}$ -structure with respect to the word  $abcd$ .

# Notational Semantics

---

Given a  $\mathcal{V}$ -structure  $w = (a_1, S_1) \dots (a_n, S_n)$ ,

# Notational Semantics

---

Given a  $\mathcal{V}$ -structure  $w = (a_1, S_1) \dots (a_n, S_n)$ ,

- ▶  $w \models Q_a(x)$  iff there exists  $j$  such that  $a_j = a$  and  $x \in S_j$ 
  - ▶  $(a, \{y\})(b, \{u, v\})(a, \{x\}) \models Q_a(x)$

# Notational Semantics

---

Given a  $\mathcal{V}$ -structure  $w = (a_1, S_1) \dots (a_n, S_n)$ ,

- ▶  $w \models Q_a(x)$  iff there exists  $j$  such that  $a_j = a$  and  $x \in S_j$ 
  - ▶  $(a, \{y\})(b, \{u, v\})(a, \{x\}) \models Q_a(x)$
- ▶  $w \models (x = y)$  iff there exists  $j$  such that  $x, y \in S_j$ 
  - ▶  $(a, \{x\})(b, \{y, z\})(c, \emptyset) \not\models (x = y)$

# Notational Semantics

---

Given a  $\mathcal{V}$ -structure  $w = (a_1, S_1) \dots (a_n, S_n)$ ,

- ▶  $w \models Q_a(x)$  iff there exists  $j$  such that  $a_j = a$  and  $x \in S_j$ 
  - ▶  $(a, \{y\})(b, \{u, v\})(a, \{x\}) \models Q_a(x)$
- ▶  $w \models (x = y)$  iff there exists  $j$  such that  $x, y \in S_j$ 
  - ▶  $(a, \{x\})(b, \{y, z\})(c, \emptyset) \not\models (x = y)$
- ▶  $w \models x < y$  iff there exists  $i < j$  such that  $x \in S_i, y \in S_j$ 
  - ▶  $(a, \{x\})(b, \{y, z\})(c, \emptyset) \models x < y$

# Notational Semantics

---

Given a  $\mathcal{V}$ -structure  $w = (a_1, S_1) \dots (a_n, S_n)$ ,

- ▶  $w \models Q_a(x)$  iff there exists  $j$  such that  $a_j = a$  and  $x \in S_j$ 
  - ▶  $(a, \{y\})(b, \{u, v\})(a, \{x\}) \models Q_a(x)$
- ▶  $w \models (x = y)$  iff there exists  $j$  such that  $x, y \in S_j$ 
  - ▶  $(a, \{x\})(b, \{y, z\})(c, \emptyset) \not\models (x = y)$
- ▶  $w \models x < y$  iff there exists  $i < j$  such that  $x \in S_i, y \in S_j$ 
  - ▶  $(a, \{x\})(b, \{y, z\})(c, \emptyset) \models x < y$
- ▶  $w \models \exists x Q_a(x)$  iff there exists  $i$  such that  $(a_1, S_1) \dots (a_i, S_i \cup \{x\}) \dots (a_n, S_n) \models Q_a(x)$ 
  - ▶  $(b, \{y, z\})(a, \{u\})(c, \emptyset) \models \exists x Q_a(x)$  since  
 $(b, \{y, z\})(a, \{x, u\})(c, \emptyset) \models Q_a(x)$

# Notational Semantics

---

- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \forall x \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff

# Notational Semantics

---

- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \forall x \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \neg \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff



# Notational Semantics

---

- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \forall x \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \neg \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \not\models \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff

# Notational Semantics

---

- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \forall x \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \neg \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \not\models \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
  - ▶  $(a, \{x\})(a, \emptyset)(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and

# Notational Semantics

---

- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \forall x \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \neg \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \not\models \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
  - ▶  $(a, \{x\})(a, \emptyset)(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and
  - ▶  $(a, \emptyset)(a, \{x\})(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and

# Notational Semantics

---

- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \forall x \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \neg \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \not\models \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
  - ▶  $(a, \{x\})(a, \emptyset)(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and
  - ▶  $(a, \emptyset)(a, \{x\})(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and
  - ▶  $(a, \emptyset)(a, \emptyset)(b, \{x\}) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$

# Notational Semantics

---

- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \forall x \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \neg \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \not\models \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
  - ▶  $(a, \{x\})(a, \emptyset)(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and
  - ▶  $(a, \emptyset)(a, \{x\})(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and
  - ▶  $(a, \emptyset)(a, \emptyset)(b, \{x\}) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$
- ▶  $(a, \{x\})(a, \emptyset)(b, \emptyset) \models \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff

# Notational Semantics

---

- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \forall x \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \neg \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \not\models \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
  - ▶  $(a, \{x\})(a, \emptyset)(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and
  - ▶  $(a, \emptyset)(a, \{x\})(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and
  - ▶  $(a, \emptyset)(a, \emptyset)(b, \{x\}) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$
- ▶  $(a, \{x\})(a, \emptyset)(b, \emptyset) \models \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \{x\})(a, \emptyset)(b, \{y\}) \models (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$

# Notational Semantics

- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \forall x \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \models \neg \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
- ▶  $(a, \emptyset)(a, \emptyset)(b, \emptyset) \not\models \exists x \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  iff
  - ▶  $(a, \{x\})(a, \emptyset)(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and
  - ▶  $(a, \emptyset)(a, \{x\})(b, \emptyset) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$  and
  - ▶  $(a, \emptyset)(a, \emptyset)(b, \{x\}) \not\models \neg [\exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])]$
- ▶  $(a, \{x\})(a, \emptyset)(b, \emptyset) \models \exists y (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  iff
- ▶  $(a, \{x\})(a, \emptyset)(b, \{y\}) \models (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$

Similarly,  $(a, \emptyset)(a, \{x\})(b, \{y\}) \models (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$  and  
 $(a, \emptyset)(a, \emptyset)(b, \{x, y\}) \models (Q_a(x) \rightarrow [(x < y) \wedge Q_b(y)])$

# Notational Semantics

---

- ▶  $(a_1, \emptyset) \dots (a_n, \emptyset) \models \exists x \varphi$  iff
- ▶ There is some position  $i$  such that  $(a_1, \emptyset) \dots (a_i, \{x\}) \dots (a_n, \emptyset) \models \varphi$



# Notational Semantics

---

- ▶  $(a_1, \emptyset) \dots (a_n, \emptyset) \models \exists x \varphi$  iff
- ▶ There is some position  $i$  such that  $(a_1, \emptyset) \dots (a_i, \{x\}) \dots (a_n, \emptyset) \models \varphi$
- ▶ For a formula  $\varphi(x_1, \dots, x_m)$ ,  $L(\varphi)$  is the set of all  $\{x_1, \dots, x_m\}$  structures satisfying  $\varphi$

# Notational Semantics

---

- ▶  $(a_1, \emptyset) \dots (a_n, \emptyset) \models \exists x \varphi$  iff
- ▶ There is some position  $i$  such that  $(a_1, \emptyset) \dots (a_i, \{x\}) \dots (a_n, \emptyset) \models \varphi$
- ▶ For a formula  $\varphi(x_1, \dots, x_m)$ ,  $L(\varphi)$  is the set of all  $\{x_1, \dots, x_m\}$  structures satisfying  $\varphi$
- ▶ For a sentence  $\varphi$ ,  $L(\varphi)$  is the set of all  $\emptyset$  structures satisfying  $\varphi$

# Logical Equivalence

---

- ▶ Let  $w_1, w_2$  be two  $\mathcal{V}$ -structures and let  $r \geq 0$ .

# Logical Equivalence

---

- ▶ Let  $w_1, w_2$  be two  $\mathcal{V}$ -structures and let  $r \geq 0$ .
- ▶ Write  $w_1 \sim_r w_2$  iff  $w_1, w_2$  satisfy the same set of FO formulae of rank  $\leq r$ .

# Logical Equivalence

---

- ▶ Let  $w_1, w_2$  be two  $\mathcal{V}$ -structures and let  $r \geq 0$ .
- ▶ Write  $w_1 \sim_r w_2$  iff  $w_1, w_2$  satisfy the same set of FO formulae of rank  $\leq r$ .
- ▶  $(a, \emptyset)(b, \emptyset) \sim_0 (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶  $(a, \emptyset)(b, \emptyset) \approx_2 (a, \emptyset)(b, \emptyset)(a, \emptyset)$

# Logical Equivalence

---

- ▶ Let  $w_1, w_2$  be two  $\mathcal{V}$ -structures and let  $r \geq 0$ .
- ▶ Write  $w_1 \sim_r w_2$  iff  $w_1, w_2$  satisfy the same set of FO formulae of rank  $\leq r$ .
- ▶  $(a, \emptyset)(b, \emptyset) \sim_0 (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶  $(a, \emptyset)(b, \emptyset) \approx_2 (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶  $\sim_r$  is an equivalence relation

# Logical Equivalence

---

- ▶ Let  $w_1, w_2$  be two  $\mathcal{V}$ -structures and let  $r \geq 0$ .
- ▶ Write  $w_1 \sim_r w_2$  iff  $w_1, w_2$  satisfy the same set of FO formulae of rank  $\leq r$ .
- ▶  $(a, \emptyset)(b, \emptyset) \sim_0 (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶  $(a, \emptyset)(b, \emptyset) \approx_2 (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶  $\sim_r$  is an equivalence relation
- ▶ **Finitely** many equivalence classes : each class consists of words that behave the same way on formulae of rank  $\leq r$

## Non-Expressibility in FO : The Game Begins



# Come, Lets Play

---

- ▶ Given two  $\mathcal{V}$ -structures  $w_1, w_2$ , lets play a game on the pair of words  $w_1, w_2$

# Come, Lets Play

---

- ▶ Given two  $\mathcal{V}$ -structures  $w_1, w_2$ , lets play a game on the pair of words  $w_1, w_2$
- ▶ There are 2 players : Spoiler and Duplicator

# Come, Lets Play

---

- ▶ Given two  $\mathcal{V}$ -structures  $w_1, w_2$ , lets play a game on the pair of words  $w_1, w_2$
- ▶ There are 2 players : Spoiler and Duplicator
- ▶ Play for  $r$ -rounds,  $r \geq 0$

# Come, Lets Play

---

- ▶ Given two  $\mathcal{V}$ -structures  $w_1, w_2$ , lets play a game on the pair of words  $w_1, w_2$
- ▶ There are 2 players : Spoiler and Duplicator
- ▶ Play for  $r$ -rounds,  $r \geq 0$
- ▶ Spoiler wants to show that  $w_1, w_2$  are different ( $w_1 \not\sim_r w_2$ )

# Come, Lets Play

---

- ▶ Given two  $\mathcal{V}$ -structures  $w_1, w_2$ , lets play a game on the pair of words  $w_1, w_2$
- ▶ There are 2 players : Spoiler and Duplicator
- ▶ Play for  $r$ -rounds,  $r \geq 0$
- ▶ Spoiler wants to show that  $w_1, w_2$  are different ( $w_1 \not\sim_r w_2$ )
- ▶ Duplicator wants to show that they are same ( $w_1 \sim_r w_2$ )

# Come, Lets Play

---

- ▶ Given two  $\mathcal{V}$ -structures  $w_1, w_2$ , lets play a game on the pair of words  $w_1, w_2$
- ▶ There are 2 players : Spoiler and Duplicator
- ▶ Play for  $r$ -rounds,  $r \geq 0$
- ▶ Spoiler wants to show that  $w_1, w_2$  are different ( $w_1 \not\sim_r w_2$ )
- ▶ Duplicator wants to show that they are same ( $w_1 \sim_r w_2$ )
- ▶ Each player has  $r$  pebbles  $z_1, \dots, z_r$

# Moves of the Game

---

- ▶ At the start of each round, spoiler chooses a structure.

# Moves of the Game

---

- ▶ At the start of each round, spoiler chooses a structure.
- ▶ Duplicator gets the other structure



# Moves of the Game

---

- ▶ At the start of each round, spoiler chooses a structure.
- ▶ Duplicator gets the other structure
- ▶ Spoiler places his pebble say  $z_i$  on one of the positions of his chosen word

# Moves of the Game

---

- ▶ At the start of each round, spoiler chooses a structure.
- ▶ Duplicator gets the other structure
- ▶ Spoiler places his pebble say  $z_i$  on one of the positions of his chosen word
- ▶ Duplicator must keep the pebble  $z_i$  on one of the positions of her word

# Moves of the Game

---

- ▶ At the start of each round, spoiler chooses a structure.
- ▶ Duplicator gets the other structure
- ▶ Spoiler places his pebble say  $z_i$  on one of the positions of his chosen word
- ▶ Duplicator must keep the pebble  $z_i$  on one of the positions of her word
- ▶ A pebble once placed, cannot be removed

# Moves of the Game

---

- ▶ At the start of each round, spoiler chooses a structure.
- ▶ Duplicator gets the other structure
- ▶ Spoiler places his pebble say  $z_i$  on one of the positions of his chosen word
- ▶ Duplicator must keep the pebble  $z_i$  on one of the positions of her word
- ▶ A pebble once placed, cannot be removed
- ▶ The game ends after  $r$  rounds, when both players have used all their pebbles

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$
- ▶ Spoiler picks  $w_2$ , duplicator picks  $w_1$

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$
- ▶ Spoiler picks  $w_2$ , duplicator picks  $w_1$
- ▶ Round 1:
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \emptyset)$



# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$
- ▶ Spoiler picks  $w_2$ , duplicator picks  $w_1$
- ▶ Round 1:
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \emptyset)$
  - ▶ Duplicator :  $(a, \{z_1\})(b, \emptyset)$

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$
- ▶ Spoiler picks  $w_2$ , duplicator picks  $w_1$
- ▶ Round 1:
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \emptyset)$
  - ▶ Duplicator :  $(a, \{z_1\})(b, \emptyset)$
  - ▶ After round 1, we have two  $\{z_1\}$  structures  $(w'_1, w'_2)$

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$
- ▶ Spoiler picks  $w_2$ , duplicator picks  $w_1$
- ▶ Round 1:
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \emptyset)$
  - ▶ Duplicator :  $(a, \{z_1\})(b, \emptyset)$
  - ▶ After round 1, we have two  $\{z_1\}$  structures  $(w'_1, w'_2)$
- ▶ Round 2:

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$
- ▶ Spoiler picks  $w_2$ , duplicator picks  $w_1$
- ▶ Round 1:
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \emptyset)$
  - ▶ Duplicator :  $(a, \{z_1\})(b, \emptyset)$
  - ▶ After round 1, we have two  $\{z_1\}$  structures  $(w'_1, w'_2)$
- ▶ Round 2:
  - ▶ Spoiler continues on the structure  $w'_2$

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$
- ▶ Spoiler picks  $w_2$ , duplicator picks  $w_1$
- ▶ Round 1:
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \emptyset)$
  - ▶ Duplicator :  $(a, \{z_1\})(b, \emptyset)$
  - ▶ After round 1, we have two  $\{z_1\}$  structures  $(w'_1, w'_2)$
- ▶ Round 2:
  - ▶ Spoiler continues on the structure  $w'_2$
  - ▶ Duplicator gets  $w'_1$  to play

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$
- ▶ Spoiler picks  $w_2$ , duplicator picks  $w_1$
- ▶ Round 1:
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \emptyset)$
  - ▶ Duplicator :  $(a, \{z_1\})(b, \emptyset)$
  - ▶ After round 1, we have two  $\{z_1\}$  structures  $(w'_1, w'_2)$
- ▶ Round 2:
  - ▶ Spoiler continues on the structure  $w'_2$
  - ▶ Duplicator gets  $w'_1$  to play
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\})$

# A Play

---

- ▶  $w_1 = (a, \emptyset)(b, \emptyset)$  and  $w_2 = (a, \emptyset)(b, \emptyset)(a, \emptyset)$
- ▶ 2 rounds, so 2 pebbles :  $z_1, z_2$
- ▶ Spoiler picks  $w_2$ , duplicator picks  $w_1$
- ▶ Round 1:
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \emptyset)$
  - ▶ Duplicator :  $(a, \{z_1\})(b, \emptyset)$
  - ▶ After round 1, we have two  $\{z_1\}$  structures  $(w'_1, w'_2)$
- ▶ Round 2:
  - ▶ Spoiler continues on the structure  $w'_2$
  - ▶ Duplicator gets  $w'_1$  to play
  - ▶ Spoiler :  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\})$
  - ▶ Duplicator :  $(a, \{z_1, z_2\})(b, \emptyset)$  or  $(a, \{z_1\})(b, \{z_2\})$

# Winner

---

- ▶ Start with two  $\emptyset$  structures  $(w_1, w_2)$



# Winner

---

- ▶ Start with two  $\emptyset$  structures  $(w_1, w_2)$
- ▶  $r$ -round game, pebble set  $\mathcal{V} = \{z_1, \dots, z_r\}$

# Winner

---

- ▶ Start with two  $\emptyset$  structures  $(w_1, w_2)$
- ▶  $r$ -round game, pebble set  $\mathcal{V} = \{z_1, \dots, z_r\}$
- ▶ Each round changes the structures

# Winner

---

- ▶ Start with two  $\emptyset$  structures  $(w_1, w_2)$
- ▶  $r$ -round game, pebble set  $\mathcal{V} = \{z_1, \dots, z_r\}$
- ▶ Each round changes the structures
- ▶ At the end of  $r$ -rounds, we have two  $\mathcal{V}$ -structures  $(w'_1, w'_2)$

# Winner

---

- ▶ Start with two  $\emptyset$  structures  $(w_1, w_2)$
- ▶  $r$ -round game, pebble set  $\mathcal{V} = \{z_1, \dots, z_r\}$
- ▶ Each round changes the structures
- ▶ At the end of  $r$ -rounds, we have two  $\mathcal{V}$ -structures  $(w'_1, w'_2)$
- ▶ Duplicator wins iff for every atomic formula  $\alpha$ ,  
 $w'_1 \models \alpha$  iff  $w'_2 \models \alpha$

# Winner

---

- ▶ Start with two  $\emptyset$  structures  $(w_1, w_2)$
- ▶  $r$ -round game, pebble set  $\mathcal{V} = \{z_1, \dots, z_r\}$
- ▶ Each round changes the structures
- ▶ At the end of  $r$ -rounds, we have two  $\mathcal{V}$ -structures  $(w'_1, w'_2)$
- ▶ Duplicator wins iff for every atomic formula  $\alpha$ ,  
 $w'_1 \models \alpha$  iff  $w'_2 \models \alpha$
- ▶ That is,  $w'_1 \sim_0 w'_2$

# Winner

---

- ▶ Start with two  $\emptyset$  structures  $(w_1, w_2)$
- ▶  $r$ -round game, pebble set  $\mathcal{V} = \{z_1, \dots, z_r\}$
- ▶ Each round changes the structures
- ▶ At the end of  $r$ -rounds, we have two  $\mathcal{V}$ -structures  $(w'_1, w'_2)$
- ▶ Duplicator wins iff for every atomic formula  $\alpha$ ,  
 $w'_1 \models \alpha$  iff  $w'_2 \models \alpha$
- ▶ That is,  $w'_1 \sim_0 w'_2$
- ▶ Spoiler wins otherwise.

# Winner

---

Given two word structures  $(w_1, w_2)$ , duplicator wins on  $(w_1, w_2)$  if for every atomic formula  $\alpha$ ,  $w_1 \models \alpha$  iff  $w_2 \models \alpha$

# Play continues

---

- ▶ Who won in the earlier play?
- ▶ We had
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\})$  and  $(a, \{z_1, z_2\})(b, \emptyset)$
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\}) \models (z_1 < z_2)$
  - ▶  $(a, \{z_1, z_2\})(b, \emptyset) \not\models (z_1 < z_2)$  or



# Play continues

---

- ▶ Who won in the earlier play?
- ▶ We had
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\})$  and  $(a, \{z_1, z_2\})(b, \emptyset)$
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\}) \models (z_1 < z_2)$
  - ▶  $(a, \{z_1, z_2\})(b, \emptyset) \not\models (z_1 < z_2)$  or
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\})$  and  $(a, \{z_1\})(b, \{z_2\})$
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\}) \models Q_a(z_2)$
  - ▶  $(a, \{z_1\})(b, \{z_2\}) \not\models Q_a(z_2)$
- ▶ Spoiler wins in two rounds

# Play continues

---

- ▶ Who won in the earlier play?
- ▶ We had
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\})$  and  $(a, \{z_1, z_2\})(b, \emptyset)$
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\}) \models (z_1 < z_2)$
  - ▶  $(a, \{z_1, z_2\})(b, \emptyset) \not\models (z_1 < z_2)$  or
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\})$  and  $(a, \{z_1\})(b, \{z_2\})$
  - ▶  $(a, \{z_1\})(b, \emptyset)(a, \{z_2\}) \models Q_a(z_2)$
  - ▶  $(a, \{z_1\})(b, \{z_2\}) \not\models Q_a(z_2)$
- ▶ Spoiler wins in two rounds
- ▶ If the game was played only for one round, who will win?

# Unique Winner

---

Given structures  $w_1$ ,  $w_2$ , and a number of rounds  $r$ , exactly one of the players win.

# Logical Equivalence and Winning

---

Let  $w_1, w_2$  be  $\mathcal{V}$ -structures and let  $r \geq 0$ . Then  $w_1 \sim_r w_2$  iff Duplicator has a winning strategy in the  $r$ -round game on  $(w_1, w_2)$ .

# Logical Equivalence and Winning

---

Assume  $w_1 \sim_r w_2$ , and induct on  $r$

- ▶ Base :  $r = 0$  and  $w_1 \sim_0 w_2$ . Duplicator wins, since by assumption,  $w_1, w_2$  agree on all atomic formulae.

# Logical Equivalence and Winning

---

Assume  $w_1 \sim_r w_2$ , and induct on  $r$

- ▶ Base :  $r = 0$  and  $w_1 \sim_0 w_2$ . Duplicator wins, since by assumption,  $w_1, w_2$  agree on all atomic formulae.
- ▶ Assume for  $r - 1$  :  $w_1 \sim_{r-1} w_2 \Rightarrow$  Duplicator has a winning strategy in a  $r - 1$  round game

# Logical Equivalence and Winning

---

- ▶ Now, let  $w_1 \sim_r w_2$ , and assume spoiler wins the  $r$ -round game on  $(w_1, w_2)$ .
  - ▶ Assume spoiler starts on  $w_1$ , places a pebble  $z_1$  somewhere on  $w_1$

# Logical Equivalence and Winning

---

- ▶ Now, let  $w_1 \sim_r w_2$ , and assume spoiler wins the  $r$ -round game on  $(w_1, w_2)$ .
  - ▶ Assume spoiler starts on  $w_1$ , places a pebble  $z_1$  somewhere on  $w_1$
  - ▶ The resultant structure is  $w'_1$



# Logical Equivalence and Winning

---

- ▶ Now, let  $w_1 \sim_r w_2$ , and assume spoiler wins the  $r$ -round game on  $(w_1, w_2)$ .
  - ▶ Assume spoiler starts on  $w_1$ , places a pebble  $z_1$  somewhere on  $w_1$
  - ▶ The resultant structure is  $w'_1$
  - ▶ In response, duplicator places her pebble somewhere on  $w_2$

# Logical Equivalence and Winning

---

- ▶ Now, let  $w_1 \sim_r w_2$ , and assume spoiler wins the  $r$ -round game on  $(w_1, w_2)$ .
  - ▶ Assume spoiler starts on  $w_1$ , places a pebble  $z_1$  somewhere on  $w_1$
  - ▶ The resultant structure is  $w'_1$
  - ▶ In response, duplicator places her pebble somewhere on  $w_2$
  - ▶ The resultant structure is  $w'_2$

# Logical Equivalence and Winning

---

- ▶ Now, let  $w_1 \sim_r w_2$ , and assume spoiler wins the  $r$ -round game on  $(w_1, w_2)$ .
  - ▶ Assume spoiler starts on  $w_1$ , places a pebble  $z_1$  somewhere on  $w_1$
  - ▶ The resultant structure is  $w'_1$
  - ▶ In response, duplicator places her pebble somewhere on  $w_2$
  - ▶ The resultant structure is  $w'_2$
  - ▶ By assumption, spoiler wins the  $r - 1$  round game on  $(w'_1, w'_2)$

# Logical Equivalence and Winning

---

- ▶ Now, let  $w_1 \sim_r w_2$ , and assume spoiler wins the  $r$ -round game on  $(w_1, w_2)$ .
  - ▶ Assume spoiler starts on  $w_1$ , places a pebble  $z_1$  somewhere on  $w_1$
  - ▶ The resultant structure is  $w'_1$
  - ▶ In response, duplicator places her pebble somewhere on  $w_2$
  - ▶ The resultant structure is  $w'_2$
  - ▶ By assumption, spoiler wins the  $r - 1$  round game on  $(w'_1, w'_2)$
  - ▶ By inductive hypothesis,  $w'_1 \sim_{r-1} w'_2$

# Logical Equivalence and Winning

---

- ▶ Now, let  $w_1 \sim_r w_2$ , and assume spoiler wins the  $r$ -round game on  $(w_1, w_2)$ .
  - ▶ Assume spoiler starts on  $w_1$ , places a pebble  $z_1$  somewhere on  $w_1$
  - ▶ The resultant structure is  $w'_1$
  - ▶ In response, duplicator places her pebble somewhere on  $w_2$
  - ▶ The resultant structure is  $w'_2$
  - ▶ By assumption, spoiler wins the  $r - 1$  round game on  $(w'_1, w'_2)$
  - ▶ By inductive hypothesis,  $w'_1 \sim_{r-1} w'_2$
  - ▶ Let  $\psi$  be the conjunction of all formulae of rank  $\leq r - 1$  in normal form that are satisfied by  $w'_1$

# Logical Equivalence and Winning

---

- ▶ Now, let  $w_1 \sim_r w_2$ , and assume spoiler wins the  $r$ -round game on  $(w_1, w_2)$ .
  - ▶ Assume spoiler starts on  $w_1$ , places a pebble  $z_1$  somewhere on  $w_1$
  - ▶ The resultant structure is  $w'_1$
  - ▶ In response, duplicator places her pebble somewhere on  $w_2$
  - ▶ The resultant structure is  $w'_2$
  - ▶ By assumption, spoiler wins the  $r - 1$  round game on  $(w'_1, w'_2)$
  - ▶ By inductive hypothesis,  $w'_1 \sim_{r-1} w'_2$
  - ▶ Let  $\psi$  be the conjunction of all formulae of rank  $\leq r - 1$  in normal form that are satisfied by  $w'_1$
  - ▶ Then  $w'_1 \models \psi$ ,  $w'_2 \not\models \psi$

# Logical Equivalence and Winning

---

- ▶ Now, let  $w_1 \sim_r w_2$ , and assume spoiler wins the  $r$ -round game on  $(w_1, w_2)$ .
  - ▶ Assume spoiler starts on  $w_1$ , places a pebble  $z_1$  somewhere on  $w_1$
  - ▶ The resultant structure is  $w'_1$
  - ▶ In response, duplicator places her pebble somewhere on  $w_2$
  - ▶ The resultant structure is  $w'_2$
  - ▶ By assumption, spoiler wins the  $r - 1$  round game on  $(w'_1, w'_2)$
  - ▶ By inductive hypothesis,  $w'_1 \sim_{r-1} w'_2$
  - ▶ Let  $\psi$  be the conjunction of all formulae of rank  $\leq r - 1$  in normal form that are satisfied by  $w'_1$
  - ▶ Then  $w'_1 \models \psi$ ,  $w'_2 \not\models \psi$
  - ▶ We thus have

$$w_1 \models \exists z_1 \psi, w_2 \not\models \exists z_1 \psi$$

contradicting  $w_1 \sim_r w_2$