

# **CS 228 : Logic in Computer Science**

Krishna. S

# So Far

---

- ▶ Proof Rules
- ▶ Soundness : If  $\varphi \vdash \psi$ , then  $\varphi \models \psi$ .
- ▶ Completeness : If  $\varphi \models \psi$ , then  $\varphi \vdash \psi$ .
- ▶ Normal Forms
- ▶ Horn Formulae
- ▶ Resolution
- ▶ Quiz 1 on Jan 27

# Resolution

---

- ▶ Resolution is a technique used to check if a formula in CNF is satisfiable. Call each conjunct in the CNF a clause.

# Resolution

---

- ▶ Resolution is a technique used to check if a formula in CNF is satisfiable. Call each conjunct in the CNF a clause.
- ▶ If  $\psi = (a \vee b \vee \neg d) \wedge (d \vee \neg a \vee b)$ , then  $C_1 = \{a, b, \neg d\}$  and  $C_2 = \{d, \neg a, b\}$  are the two clauses.

# Resolution

---

- ▶ Resolution is a technique used to check if a formula in CNF is satisfiable. Call each conjunct in the CNF a clause.
- ▶ If  $\psi = (a \vee b \vee \neg d) \wedge (d \vee \neg a \vee b)$ , then  $C_1 = \{a, b, \neg d\}$  and  $C_2 = \{d, \neg a, b\}$  are the two clauses.
- ▶ We also write  $\psi$  as  $\{\{a, b, \neg d\}, \{d, \neg a, b\}\}$ .

# Resolution

---

- ▶ Resolution is a technique used to check if a formula in CNF is satisfiable. Call each conjunct in the CNF a clause.
- ▶ If  $\psi = (a \vee b \vee \neg d) \wedge (d \vee \neg a \vee b)$ , then  $C_1 = \{a, b, \neg d\}$  and  $C_2 = \{d, \neg a, b\}$  are the two clauses.
- ▶ We also write  $\psi$  as  $\{\{a, b, \neg d\}, \{d, \neg a, b\}\}$ .
- ▶ Let  $C_1, C_2$  be two clauses. Assume  $a \in C_1$  and  $\neg a \in C_2$  for some atomic formula  $a$ . Then the clause  $R = (C_1 - \{a\}) \cup (C_2 - \{\neg a\})$  is a **resolvent** of  $C_1$  and  $C_2$ .

# Resolution

---

- ▶ Resolution is a technique used to check if a formula in CNF is satisfiable. Call each conjunct in the CNF a clause.
- ▶ If  $\psi = (a \vee b \vee \neg d) \wedge (d \vee \neg a \vee b)$ , then  $C_1 = \{a, b, \neg d\}$  and  $C_2 = \{d, \neg a, b\}$  are the two clauses.
- ▶ We also write  $\psi$  as  $\{\{a, b, \neg d\}, \{d, \neg a, b\}\}$ .
- ▶ Let  $C_1, C_2$  be two clauses. Assume  $a \in C_1$  and  $\neg a \in C_2$  for some atomic formula  $a$ . Then the clause  $R = (C_1 - \{a\}) \cup (C_2 - \{\neg a\})$  is a **resolvent** of  $C_1$  and  $C_2$ .
- ▶ Let  $C_1 = \{a_1, \neg a_2, a_3\}$  and  $C_2 = \{a_2, \neg a_3, a_4\}$ . As  $a_3 \in C_1$  and  $\neg a_3 \in C_2$ , we can find the resolvent. A resolvent is  $\{a_1, a_2, \neg a_2, a_4\}$ .

# Resolution

---

- ▶ Resolution is a technique used to check if a formula in CNF is satisfiable. Call each conjunct in the CNF a clause.
- ▶ If  $\psi = (a \vee b \vee \neg d) \wedge (d \vee \neg a \vee b)$ , then  $C_1 = \{a, b, \neg d\}$  and  $C_2 = \{d, \neg a, b\}$  are the two clauses.
- ▶ We also write  $\psi$  as  $\{\{a, b, \neg d\}, \{d, \neg a, b\}\}$ .
- ▶ Let  $C_1, C_2$  be two clauses. Assume  $a \in C_1$  and  $\neg a \in C_2$  for some atomic formula  $a$ . Then the clause  $R = (C_1 - \{a\}) \cup (C_2 - \{\neg a\})$  is a **resolvent** of  $C_1$  and  $C_2$ .
- ▶ Let  $C_1 = \{a_1, \neg a_2, a_3\}$  and  $C_2 = \{a_2, \neg a_3, a_4\}$ . As  $a_3 \in C_1$  and  $\neg a_3 \in C_2$ , we can find the resolvent. A resolvent is  $\{a_1, a_2, \neg a_2, a_4\}$ .
- ▶ Resolvent **not** unique :  $\{a_1, a_3, \neg a_3, a_4\}$  is also a resolvent.



# Resolution

---

- ▶ Let  $F$  be a formula in CNF, and let  $C$  be a clause in  $F$ . Then  $F \vdash C$  (Prove!)

# Resolution

---

- ▶ Let  $F$  be a formula in CNF, and let  $C$  be a clause in  $F$ . Then  $F \vdash C$  (Prove!)
- ▶ Let  $F$  be a formula in CNF. Let  $R$  be a resolvent of two clauses of  $F$ . Then  $F \vdash R$  (Prove!)

# Completeness of Resolution

---

Show that resolution can be used to determine whether any given formula is satisfiable.

- ▶ Given  $F$  in CNF, let  $Res^0(F) = \{C \mid C \text{ is a clause in } F\}$ .

# Completeness of Resolution

---

Show that resolution can be used to determine whether any given formula is satisfiable.

- ▶ Given  $F$  in CNF, let  $Res^0(F) = \{C \mid C \text{ is a clause in } F\}$ .
- ▶  $Res(F) = Res^0(F) \cup \{R \mid R \text{ is a resolvent of two clauses in } Res^0(F)\}$

# Completeness of Resolution

---

Show that resolution can be used to determine whether any given formula is satisfiable.

- ▶ Given  $F$  in CNF, let  $Res^0(F) = \{C \mid C \text{ is a clause in } F\}$ .
- ▶  $Res(F) = Res^0(F) \cup \{R \mid R \text{ is a resolvent of two clauses in } Res^0(F)\}$
- ▶  $Res^n(F) = Res^{n-1}(F) \cup \{R \mid R \text{ is a resolvent of two clauses in } Res^{n-1}(F)\}$

# Completeness of Resolution

---

Show that resolution can be used to determine whether any given formula is satisfiable.

- ▶ Given  $F$  in CNF, let  $Res^0(F) = \{C \mid C \text{ is a clause in } F\}$ .
- ▶  $Res(F) = Res^0(F) \cup \{R \mid R \text{ is a resolvent of two clauses in } Res^0(F)\}$
- ▶  $Res^n(F) = Res^{n-1}(F) \cup \{R \mid R \text{ is a resolvent of two clauses in } Res^{n-1}(F)\}$
- ▶  $Res^0(F)$  is finite : there are finitely many clauses that can be derived from  $F$ .

# Completeness of Resolution

---

Show that resolution can be used to determine whether any given formula is satisfiable.

- ▶ Given  $F$  in CNF, let  $Res^0(F) = \{C \mid C \text{ is a clause in } F\}$ .
- ▶  $Res(F) = Res^0(F) \cup \{R \mid R \text{ is a resolvent of two clauses in } Res^0(F)\}$
- ▶  $Res^n(F) = Res^{n-1}(F) \cup \{R \mid R \text{ is a resolvent of two clauses in } Res^{n-1}(F)\}$
- ▶  $Res^0(F)$  is finite : there are finitely many clauses that can be derived from  $F$ .
- ▶  $Res^1(F)$  is finite : finitely many resolvents can be obtained from clauses of  $F$

# Completeness of Resolution

---

Show that resolution can be used to determine whether any given formula is satisfiable.

- ▶ Given  $F$  in CNF, let  $Res^0(F) = \{C \mid C \text{ is a clause in } F\}$ .
- ▶  $Res(F) = Res^0(F) \cup \{R \mid R \text{ is a resolvent of two clauses in } Res^0(F)\}$
- ▶  $Res^n(F) = Res^{n-1}(F) \cup \{R \mid R \text{ is a resolvent of two clauses in } Res^{n-1}(F)\}$
- ▶  $Res^0(F)$  is finite : there are finitely many clauses that can be derived from  $F$ .
- ▶  $Res^1(F)$  is finite : finitely many resolvents can be obtained from clauses of  $F$
- ▶ There is some  $m$  such that  $Res^m(F) = Res^{m+1}(F)$ . Denote it by  $Res^*(F)$ .



# Example

---

Let  $F = \{\{a_1, a_2, \neg a_3\}, \{\neg a_2, a_3\}\}$ .

►  $Res^0(F) = F$

# Example

---

Let  $F = \{\{a_1, a_2, \neg a_3\}, \{\neg a_2, a_3\}\}$ .

- ▶  $Res^0(F) = F$
- ▶  $Res^1(F) = F \cup \{a_1, a_2, \neg a_2\} \cup \{a_1, \neg a_3, a_3\}$ .

# Example

---

Let  $F = \{\{a_1, a_2, \neg a_3\}, \{\neg a_2, a_3\}\}$ .

- ▶  $Res^0(F) = F$
- ▶  $Res^1(F) = F \cup \{a_1, a_2, \neg a_2\} \cup \{a_1, \neg a_3, a_3\}$ .
- ▶  $Res^2(F) = Res^1(F) \cup \{a_1, a_2, \neg a_3\} \cup \{a_1, a_3, \neg a_2\}$

# Resolution

---

Let  $F$  be a formula in CNF. If  $\emptyset \in \text{Res}^*(F)$ , then  $F$  is unsatisfiable.

- If  $\emptyset \in \text{Res}^*(F)$ . Then  $\emptyset \in \text{Res}^n(F)$  for some  $n$ .

# Resolution

---

Let  $F$  be a formula in CNF. If  $\emptyset \in \text{Res}^*(F)$ , then  $F$  is unsatisfiable.

- ▶ If  $\emptyset \in \text{Res}^*(F)$ . Then  $\emptyset \in \text{Res}^n(F)$  for some  $n$ .
- ▶ Since  $\emptyset \notin \text{Res}^0(F)$  ( $\emptyset$  is not a clause), there is an  $m > 0$  such that  $\emptyset \notin \text{Res}^m(F)$  and  $\emptyset \in \text{Res}^{m+1}(F)$ .

# Resolution

---

Let  $F$  be a formula in CNF. If  $\emptyset \in \text{Res}^*(F)$ , then  $F$  is unsatisfiable.

- ▶ If  $\emptyset \in \text{Res}^*(F)$ . Then  $\emptyset \in \text{Res}^n(F)$  for some  $n$ .
- ▶ Since  $\emptyset \notin \text{Res}^0(F)$  ( $\emptyset$  is not a clause), there is an  $m > 0$  such that  $\emptyset \notin \text{Res}^m(F)$  and  $\emptyset \in \text{Res}^{m+1}(F)$ .
- ▶ Then  $\{A\}, \{\neg A\} \in \text{Res}^m(F)$ . By the rules of resolution, we have  $F \vdash A, \neg A$ , and thus  $F \vdash \perp$ . Hence,  $F$  is unsatisfiable.

# Resolution

---

If  $F$  in CNF is unsatisfiable, then  $\emptyset \in \text{Res}^*(F)$ .

- ▶ Let  $F$  have  $k$  clauses  $C_1, \dots, C_k$ .

# Resolution

---

If  $F$  in CNF is unsatisfiable, then  $\emptyset \in \text{Res}^*(F)$ .

- ▶ Let  $F$  have  $k$  clauses  $C_1, \dots, C_k$ .
- ▶ wlg, assume that no  $C_i$  has both  $p$  and  $\neg p$



# Resolution

---

If  $F$  in CNF is unsatisfiable, then  $\emptyset \in \text{Res}^*(F)$ .

- ▶ Let  $F$  have  $k$  clauses  $C_1, \dots, C_k$ .
- ▶ wlg, assume that no  $C_i$  has both  $p$  and  $\neg p$
- ▶ Induct on the number  $n$  of propositional variables that occur in  $F$ .

# Resolution

---

If  $F$  in CNF is unsatisfiable, then  $\emptyset \in \text{Res}^*(F)$ .

- ▶ Let  $F$  have  $k$  clauses  $C_1, \dots, C_k$ .
- ▶ wlg, assume that no  $C_i$  has both  $p$  and  $\neg p$
- ▶ Induct on the number  $n$  of propositional variables that occur in  $F$ .
- ▶ If  $n = 1$ , then the possible clauses are  $p$ ,  $\neg p$  and  $p \vee \neg p$ . The third one is ruled out, by assumption.

# Resolution

---

If  $F$  in CNF is unsatisfiable, then  $\emptyset \in \text{Res}^*(F)$ .

- ▶ Let  $F$  have  $k$  clauses  $C_1, \dots, C_k$ .
- ▶ wlg, assume that no  $C_i$  has both  $p$  and  $\neg p$
- ▶ Induct on the number  $n$  of propositional variables that occur in  $F$ .
- ▶ If  $n = 1$ , then the possible clauses are  $p$ ,  $\neg p$  and  $p \vee \neg p$ . The third one is ruled out, by assumption.
- ▶ If  $F = \{\{p\}\}$  or  $F = \{\{\neg p\}\}$ ,  $F$  is satisfiable.

# Resolution

---

If  $F$  in CNF is unsatisfiable, then  $\emptyset \in \text{Res}^*(F)$ .

- ▶ Let  $F$  have  $k$  clauses  $C_1, \dots, C_k$ .
- ▶ wlg, assume that no  $C_i$  has both  $p$  and  $\neg p$
- ▶ Induct on the number  $n$  of propositional variables that occur in  $F$ .
- ▶ If  $n = 1$ , then the possible clauses are  $p$ ,  $\neg p$  and  $p \vee \neg p$ . The third one is ruled out, by assumption.
- ▶ If  $F = \{\{p\}\}$  or  $F = \{\{\neg p\}\}$ ,  $F$  is satisfiable.
- ▶ Hence,  $F = \{\{p\}, \{\neg p\}\}$ . Clearly,  $\emptyset \in \text{Res}(F)$ .

# Resolution

---

- ▶ Inductive hypothesis : If  $F$  has  $\leq n$  variables and is unsat, then  $\emptyset \in Res^*(F)$ .

# Resolution

---

- ▶ Inductive hypothesis : If  $F$  has  $\leq n$  variables and is unsat, then  $\emptyset \in Res^*(F)$ .
- ▶ Let  $F$  have  $n + 1$  variables  $p_1, \dots, p_{n+1}$ .

# Resolution

---

- ▶ Inductive hypothesis : If  $F$  has  $\leq n$  variables and is unsat, then  $\emptyset \in Res^*(F)$ .
- ▶ Let  $F$  have  $n + 1$  variables  $p_1, \dots, p_{n+1}$ .
  - ▶ Let  $G_0$  be the conjunction of all  $C_i$  in  $F$  such that  $\neg p_{n+1} \notin C_i$ .
  - ▶ Let  $G_1$  be the conjunction of all  $C_i$  in  $F$  such that  $p_{n+1} \notin C_i$ .

# Resolution

---

- ▶ Inductive hypothesis : If  $F$  has  $\leq n$  variables and is unsat, then  $\emptyset \in Res^*(F)$ .
- ▶ Let  $F$  have  $n + 1$  variables  $p_1, \dots, p_{n+1}$ .
  - ▶ Let  $G_0$  be the conjunction of all  $C_i$  in  $F$  such that  $\neg p_{n+1} \notin C_i$ .
  - ▶ Let  $G_1$  be the conjunction of all  $C_i$  in  $F$  such that  $p_{n+1} \notin C_i$ .
- ▶ Clauses in  $F =$  Clauses in  $G_0 \cup$  Clauses in  $G_1$



# Resolution

- ▶ Inductive hypothesis : If  $F$  has  $\leq n$  variables and is unsat, then  $\emptyset \in Res^*(F)$ .
- ▶ Let  $F$  have  $n + 1$  variables  $p_1, \dots, p_{n+1}$ .

- ▶ Let  $G_0$  be the conjunction of all  $C_i$  in  $F$  such that  $\neg p_{n+1} \notin C_i$ .
- ▶ Let  $G_1$  be the conjunction of all  $C_i$  in  $F$  such that  $p_{n+1} \notin C_i$ .

- ▶ Clauses in  $F =$  Clauses in  $G_0 \cup$  Clauses in  $G_1$

- ▶ Let  $F_0 = \{C_i - \{p_{n+1}\} \mid C_i \in G_0\}$
- ▶ Let  $F_1 = \{C_i - \{\neg p_{n+1}\} \mid C_i \in G_1\}$

# Resolution

---

Let  $F = \{\{p_1, p_3\}, \{p_2\}, \{\neg p_1, \neg p_2, p_3\}, \{\neg p_2, \neg p_3\}\}$  and  $n = 2$ .

- ▶  $G_0 = \{\{p_1, p_3\}, \{p_2\}, \{\neg p_1, \neg p_2, p_3\}\}$ ,  $G_1 = \{\{p_2\}, \{\neg p_2, \neg p_3\}\}$ .
- ▶  $F_0 = \{\{p_1\}, \{p_2\}, \{\neg p_1, \neg p_2\}\}$  and  $F_1 = \{\{p_2\}, \{\neg p_2\}\}$
- ▶ If  $p_{n+1} = \text{false}$  in  $F$ , then  $F$  is equivalent to  $F_0$

# Resolution

---

Let  $F = \{\{p_1, p_3\}, \{p_2\}, \{\neg p_1, \neg p_2, p_3\}, \{\neg p_2, \neg p_3\}\}$  and  $n = 2$ .

- ▶  $G_0 = \{\{p_1, p_3\}, \{p_2\}, \{\neg p_1, \neg p_2, p_3\}\}$ ,  $G_1 = \{\{p_2\}, \{\neg p_2, \neg p_3\}\}$ .
- ▶  $F_0 = \{\{p_1\}, \{p_2\}, \{\neg p_1, \neg p_2\}\}$  and  $F_1 = \{\{p_2\}, \{\neg p_2\}\}$
- ▶ If  $p_{n+1} = \text{false}$  in  $F$ , then  $F$  is equivalent to  $F_0$
- ▶ If  $p_{n+1} = \text{true}$  in  $F$ , then  $F$  is equivalent to  $F_1$

# Resolution

---

Let  $F = \{\{p_1, p_3\}, \{p_2\}, \{\neg p_1, \neg p_2, p_3\}, \{\neg p_2, \neg p_3\}\}$  and  $n = 2$ .

- ▶  $G_0 = \{\{p_1, p_3\}, \{p_2\}, \{\neg p_1, \neg p_2, p_3\}\}$ ,  $G_1 = \{\{p_2\}, \{\neg p_2, \neg p_3\}\}$ .
- ▶  $F_0 = \{\{p_1\}, \{p_2\}, \{\neg p_1, \neg p_2\}\}$  and  $F_1 = \{\{p_2\}, \{\neg p_2\}\}$
- ▶ If  $p_{n+1} = \text{false}$  in  $F$ , then  $F$  is equivalent to  $F_0$
- ▶ If  $p_{n+1} = \text{true}$  in  $F$ , then  $F$  is equivalent to  $F_1$
- ▶ Hence  $F \equiv F_0 \vee F_1$ .

# Resolution

---

Let  $F = \{\{p_1, p_3\}, \{p_2\}, \{\neg p_1, \neg p_2, p_3\}, \{\neg p_2, \neg p_3\}\}$  and  $n = 2$ .

- ▶  $G_0 = \{\{p_1, p_3\}, \{p_2\}, \{\neg p_1, \neg p_2, p_3\}\}$ ,  $G_1 = \{\{p_2\}, \{\neg p_2, \neg p_3\}\}$ .
- ▶  $F_0 = \{\{p_1\}, \{p_2\}, \{\neg p_1, \neg p_2\}\}$  and  $F_1 = \{\{p_2\}, \{\neg p_2\}\}$
- ▶ If  $p_{n+1} = \text{false}$  in  $F$ , then  $F$  is equivalent to  $F_0$
- ▶ If  $p_{n+1} = \text{true}$  in  $F$ , then  $F$  is equivalent to  $F_1$
- ▶ Hence  $F \equiv F_0 \vee F_1$ .
- ▶ As  $F$  is unsatisfiable,  $F_0$  and  $F_1$  are both unsatisfiable.

# Resolution

---

- ▶ By induction hypothesis,  $\emptyset \in Res^*(F_0)$  and  $\emptyset \in Res^*(F_1)$ .

# Resolution

---

- ▶ By induction hypothesis,  $\emptyset \in Res^*(F_0)$  and  $\emptyset \in Res^*(F_1)$ .
- ▶ Hence,  $\emptyset \in Res^*(G_0)$  or  $\{p_{n+1}\} \in Res^*(G_0)$ , and  $\emptyset \in Res^*(G_1)$  or  $\{\neg p_{n+1}\} \in Res^*(G_1)$ .

# Resolution

---

- ▶ By induction hypothesis,  $\emptyset \in Res^*(F_0)$  and  $\emptyset \in Res^*(F_1)$ .
- ▶ Hence,  $\emptyset \in Res^*(G_0)$  or  $\{p_{n+1}\} \in Res^*(G_0)$ , and  $\emptyset \in Res^*(G_1)$  or  $\{\neg p_{n+1}\} \in Res^*(G_1)$ .
- ▶ If  $\emptyset \in Res^*(G_0)$  or  $\emptyset \in Res^*(G_1)$ , then  $\emptyset \in Res^*(F)$ .



# Resolution

---

- ▶ By induction hypothesis,  $\emptyset \in Res^*(F_0)$  and  $\emptyset \in Res^*(F_1)$ .
- ▶ Hence,  $\emptyset \in Res^*(G_0)$  or  $\{p_{n+1}\} \in Res^*(G_0)$ , and  $\emptyset \in Res^*(G_1)$  or  $\{\neg p_{n+1}\} \in Res^*(G_1)$ .
- ▶ If  $\emptyset \in Res^*(G_0)$  or  $\emptyset \in Res^*(G_1)$ , then  $\emptyset \in Res^*(F)$ .
- ▶ Else,  $\{p_{n+1}\} \in Res^*(G_0)$  and  $\{\neg p_{n+1}\} \in Res^*(G_1)$ .

# Resolution

---

- ▶ By induction hypothesis,  $\emptyset \in Res^*(F_0)$  and  $\emptyset \in Res^*(F_1)$ .
- ▶ Hence,  $\emptyset \in Res^*(G_0)$  or  $\{p_{n+1}\} \in Res^*(G_0)$ , and  $\emptyset \in Res^*(G_1)$  or  $\{\neg p_{n+1}\} \in Res^*(G_1)$ .
- ▶ If  $\emptyset \in Res^*(G_0)$  or  $\emptyset \in Res^*(G_1)$ , then  $\emptyset \in Res^*(F)$ .
- ▶ Else,  $\{p_{n+1}\} \in Res^*(G_0)$  and  $\{\neg p_{n+1}\} \in Res^*(G_1)$ .
- ▶ Hence  $\emptyset \in Res^*(F)$ .

# Resolution Summary

---

Given a formula  $\psi$ , convert it into CNF, say  $\zeta$ .  $\psi$  is satisfiable iff  $\emptyset \notin \text{Res}^*(\zeta)$ .

- If  $\psi$  is unsat, we might get  $\emptyset$  before reaching  $\text{Res}^*(\zeta)$ .

# Resolution Summary

---

Given a formula  $\psi$ , convert it into CNF, say  $\zeta$ .  $\psi$  is satisfiable iff  $\emptyset \notin Res^*(\zeta)$ .

- ▶ If  $\psi$  is unsat, we might get  $\emptyset$  before reaching  $Res^*(\zeta)$ .
- ▶ If  $\psi$  is sat, then truth tables are faster : stop when some row evaluates to 1.

# Resolution Summary

---

Given a formula  $\psi$ , convert it into CNF, say  $\zeta$ .  $\psi$  is satisfiable iff  $\emptyset \notin \text{Res}^*(\zeta)$ .

- ▶ If  $\psi$  is unsat, we might get  $\emptyset$  before reaching  $\text{Res}^*(\zeta)$ .
- ▶ If  $\psi$  is sat, then truth tables are faster : stop when some row evaluates to 1.
- ▶ Time to move on!