

# **CS 228 : Logic in Computer Science**

Krishna. S

# So Far


---

- ▶  $\omega$ -automata with Büchi acceptance, also called Büchi automata
- ▶ Non-determinism versus determinism

## Büchi Acceptance

For Büchi Acceptance,  $Acc$  is specified as a set of states,  $G \subseteq Q$ . The  $\omega$ -word  $\alpha$  is accepted if there is a run  $\rho$  of  $\alpha$  such that  $Inf(\rho) \cap G \neq \emptyset$ .

# $\omega$ -Automata with Büchi Acceptance



- ▶ Left (T-B): Inf many  $b$ 's, Inf many  $a$ 's
- ▶ Right (T-B): Finitely many  $b$ 's,  $(a + b)^{\omega}$

# Büchi Acceptance


---

A language  $L \subseteq \Sigma^\omega$  is called  $\omega$ -regular if there exists a NBA  $\mathcal{A}$  such that  $L = L(\mathcal{A})$ . Recall definition of regular languages and NFA/DFA acceptance.

# NBA and DBA

---



- ▶ Is every DBA as expressible as a NBA, like in the case of DFA and NFA?
- ▶ Can we do subset construction on NBA and obtain DBA?



# NBA and DBA

---

- ▶ Is every DBA as expressible as a NBA, like in the case of DFA and NFA?
- ▶ Can we do subset construction on NBA and obtain DBA?



# NBA and DBA


---

There does not exist a deterministic Büchi automata capturing the language finitely many  $a$ 's.

# NBA and DBA

---

There does not exist a deterministic Büchi automata capturing the language finitely many  $a$ 's.






# NBA and DBA


---

There does not exist a deterministic Büchi automata capturing the language finitely many  $a$ 's.




# NBA and DBA

There does not exist a deterministic Büchi automata capturing the language finitely many  $a$ 's.




# NBA and DBA

There does not exist a deterministic Büchi automata capturing the language finitely many  $a$ 's.




# NBA and DBA

There does not exist a deterministic Büchi automata capturing the language finitely many  $a$ 's.




# Union and Intersection of NBA

---




# Union and Intersection of NBA




- States as  $Q_1 \times Q_2 \times \{1, 2\}$ , start state  $(q_0, q'_0, 1)$

# Union and Intersection of NBA



- ▶ States as  $Q_1 \times Q_2 \times \{1, 2\}$ , start state  $(q_0, q'_0, 1)$
- ▶  $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 1)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_1 \notin G_1$
- ▶  $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 2)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_1 \in G_1$


# Union and Intersection of NBA



- ▶ States as  $Q_1 \times Q_2 \times \{1, 2\}$ , start state  $(q_0, q'_0, 1)$
- ▶  $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 1)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_1 \notin G_1$
- ▶  $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 2)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_1 \in G_1$
- ▶  $(q_1, q_2, 2) \xrightarrow{a} (q'_1, q'_2, 2)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_2 \notin G_2$
- ▶  $(q_1, q_2, 2) \xrightarrow{a} (q'_1, q'_2, 1)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_2 \in G_2$



# Union and Intersection of NBA



- ▶ States as  $Q_1 \times Q_2 \times \{1, 2\}$ , start state  $(q_0, q'_0, 1)$
- ▶  $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 1)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_1 \notin G_1$
- ▶  $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 2)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_1 \in G_1$
- ▶  $(q_1, q_2, 2) \xrightarrow{a} (q'_1, q'_2, 2)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_2 \notin G_2$
- ▶  $(q_1, q_2, 2) \xrightarrow{a} (q'_1, q'_2, 1)$  if  $q_1 \xrightarrow{a} q'_1$  and  $q_2 \xrightarrow{a} q'_2$  and  $q_2 \in G_2$
- ▶ Good states =  $Q_1 \times G_2 \times \{2\}$  or  $G_1 \times Q_2 \times \{1\}$