

# **CS 228 : Logic in Computer Science**

Krishna. S

# Free and Bound Variables

---

- ▶ For a wff  $\varphi = \forall x\psi$ ,  $\psi$  is said to be the scope of the quantifier  $x$

# Free and Bound Variables

---

- ▶ For a wff  $\varphi = \forall x\psi$ ,  $\psi$  is said to be the scope of the quantifier  $x$
- ▶ Every occurrence of  $x$  in  $\forall x\psi$  is **bound**
- ▶ Any occurrence of  $x$  which is not bound is called **free**

# Free and Bound Variables

---

- ▶ For a wff  $\varphi = \forall x\psi$ ,  $\psi$  is said to be the scope of the quantifier  $x$
- ▶ Every occurrence of  $x$  in  $\forall x\psi$  is **bound**
- ▶ Any occurrence of  $x$  which is not bound is called **free**
- ▶  $\varphi = P(x, y) \rightarrow \forall x((\forall yR(x, y)) \rightarrow Q(x, y))$ 
  - ▶  $y$  is free in  $Q(x, y)$  and bound in  $R(x, y)$ ,
  - ▶  $x$  is free in  $P(x, y)$ , and bound in  $Q(x, y), R(x, y)$


# Free and Bound Variables

---

- ▶ For a wff  $\varphi = \forall x\psi$ ,  $\psi$  is said to be the scope of the quantifier  $x$
- ▶ Every occurrence of  $x$  in  $\forall x\psi$  is **bound**
- ▶ Any occurrence of  $x$  which is not bound is called **free**
- ▶  $\varphi = P(x, y) \rightarrow \forall x((\forall yR(x, y)) \rightarrow Q(x, y))$ 
  - ▶  $y$  is free in  $Q(x, y)$  and bound in  $R(x, y)$ ,
  - ▶  $x$  is free in  $P(x, y)$ , and bound in  $Q(x, y), R(x, y)$
- ▶ Given  $\varphi$ , denote by  $\varphi(x_1, \dots, x_n)$ , that  $x_1, \dots, x_n$  are the free variables of  $\varphi$ , also  $\text{free}(\varphi)$
- ▶ A **sentence** is a formula  $\varphi$  **none** of whose variables are **free**


$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$

---




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$

---



$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$


---






$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$

---




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$

---




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$

---




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$

---




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$

---



$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$


---



$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$



---

$$\varphi(s, t) = P(s, t) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, t))$$



$$\forall x(R(x, y) \rightarrow \forall yP(y))$$


---






$$\forall x(R(x, y) \rightarrow \forall yP(y))$$

---




$$\forall x(R(x, y) \rightarrow \forall yP(y))$$

---




$$\forall x(R(x, y) \rightarrow \forall yP(y))$$

---




$$\forall x(R(x, y) \rightarrow \forall yP(y))$$

---




$$\forall x(R(x, y) \rightarrow \forall yP(y))$$

---



$$\forall x(R(x, y) \rightarrow \forall yP(y))$$

---



$$\varphi(t) = \forall x(R(x, t) \rightarrow \forall yP(y))$$

# Assignments on $\tau$ -structures

---

## Assignments

For a  $\tau$ -structure  $\mathcal{A}$ , an assignment over  $\mathcal{A}$  is a function  $\alpha : \mathcal{V} \rightarrow u(\mathcal{A})$  that assigns every variable  $x \in \mathcal{V}$  a value  $\alpha(x) \in u(\mathcal{A})$ . If  $t$  is a constant symbol  $c$ , then  $\alpha(t)$  is  $c^{\mathcal{A}}$

# Assignments on $\tau$ -structures

---

## Assignments

For a  $\tau$ -structure  $\mathcal{A}$ , an assignment over  $\mathcal{A}$  is a function  $\alpha : \mathcal{V} \rightarrow u(\mathcal{A})$  that assigns every variable  $x \in \mathcal{V}$  a value  $\alpha(x) \in u(\mathcal{A})$ . If  $t$  is a constant symbol  $c$ , then  $\alpha(t)$  is  $c^{\mathcal{A}}$

## Binding on a Variable

For an assignment  $\alpha$  over  $\mathcal{A}$ ,  $\alpha[x \mapsto a]$  is the assignment

$$\alpha[x \mapsto a](y) = \begin{cases} \alpha(y), & y \neq x, \\ a, & y = x \end{cases}$$



# Satisfaction

---

We define the relation  $\mathcal{A} \models_{\alpha} \varphi$  (read as  $\varphi$  is true in  $\mathcal{A}$  under the assignment  $\alpha$ ) inductively:

# Satisfaction

---

We define the relation  $\mathcal{A} \models_{\alpha} \varphi$  (read as  $\varphi$  is true in  $\mathcal{A}$  under the assignment  $\alpha$ ) inductively:

- ▶  $\mathcal{A} \not\models_{\alpha} \perp$

# Satisfaction

---

We define the relation  $\mathcal{A} \models_{\alpha} \varphi$  (read as  $\varphi$  is true in  $\mathcal{A}$  under the assignment  $\alpha$ ) inductively:

- ▶  $\mathcal{A} \not\models_{\alpha} \perp$
- ▶  $\mathcal{A} \models_{\alpha} t_1 = t_2$  iff  $\alpha(t_1) = \alpha(t_2)$

# Satisfaction

---

We define the relation  $\mathcal{A} \models_{\alpha} \varphi$  (read as  $\varphi$  is true in  $\mathcal{A}$  under the assignment  $\alpha$ ) inductively:

- ▶  $\mathcal{A} \not\models_{\alpha} \perp$
- ▶  $\mathcal{A} \models_{\alpha} t_1 = t_2$  iff  $\alpha(t_1) = \alpha(t_2)$
- ▶  $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$  iff  $(\alpha(t_1), \dots, \alpha(t_k)) \in R^{\mathcal{A}}$

# Satisfaction

---

We define the relation  $\mathcal{A} \models_{\alpha} \varphi$  (read as  $\varphi$  is true in  $\mathcal{A}$  under the assignment  $\alpha$ ) inductively:

- ▶  $\mathcal{A} \not\models_{\alpha} \perp$
- ▶  $\mathcal{A} \models_{\alpha} t_1 = t_2$  iff  $\alpha(t_1) = \alpha(t_2)$
- ▶  $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$  iff  $(\alpha(t_1), \dots, \alpha(t_k)) \in R^{\mathcal{A}}$
- ▶  $\mathcal{A} \models_{\alpha} (\varphi \rightarrow \psi)$  iff  $\mathcal{A} \not\models_{\alpha} \varphi$  or  $\mathcal{A} \models_{\alpha} \psi$

# Satisfaction

---

We define the relation  $\mathcal{A} \models_{\alpha} \varphi$  (read as  $\varphi$  is true in  $\mathcal{A}$  under the assignment  $\alpha$ ) inductively:

- ▶  $\mathcal{A} \not\models_{\alpha} \perp$
- ▶  $\mathcal{A} \models_{\alpha} t_1 = t_2$  iff  $\alpha(t_1) = \alpha(t_2)$
- ▶  $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$  iff  $(\alpha(t_1), \dots, \alpha(t_k)) \in R^{\mathcal{A}}$
- ▶  $\mathcal{A} \models_{\alpha} (\varphi \rightarrow \psi)$  iff  $\mathcal{A} \not\models_{\alpha} \varphi$  or  $\mathcal{A} \models_{\alpha} \psi$
- ▶  $\mathcal{A} \models_{\alpha} (\forall x)\varphi$  iff for every  $a \in u(\mathcal{A})$ ,  $\mathcal{A} \models_{\alpha[x \mapsto a]} \varphi$

# Satisfaction

---

We define the relation  $\mathcal{A} \models_{\alpha} \varphi$  (read as  $\varphi$  is true in  $\mathcal{A}$  under the assignment  $\alpha$ ) inductively:

- ▶  $\mathcal{A} \not\models_{\alpha} \perp$
- ▶  $\mathcal{A} \models_{\alpha} t_1 = t_2$  iff  $\alpha(t_1) = \alpha(t_2)$
- ▶  $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$  iff  $(\alpha(t_1), \dots, \alpha(t_k)) \in R^{\mathcal{A}}$
- ▶  $\mathcal{A} \models_{\alpha} (\varphi \rightarrow \psi)$  iff  $\mathcal{A} \not\models_{\alpha} \varphi$  or  $\mathcal{A} \models_{\alpha} \psi$
- ▶  $\mathcal{A} \models_{\alpha} (\forall x)\varphi$  iff for every  $a \in u(\mathcal{A})$ ,  $\mathcal{A} \models_{\alpha[x \mapsto a]} \varphi$
- ▶  $\mathcal{A} \models_{\alpha} (\exists x)\varphi$  iff there is some  $a \in u(\mathcal{A})$ ,  $\mathcal{A} \models_{\alpha[x \mapsto a]} \varphi$

Last two cases,  $\alpha$  has no effect on the value of  $x$ . Thus, assignments matter to free variables.

# Example of Satisfaction

---

- ▶  $\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$ 
  - ▶ For any assignment  $\alpha$ ,  $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$  iff for every  $a, b \in u(\mathcal{A})$ ,  $\mathcal{A} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$



# Example of Satisfaction

---

- ▶  $\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$ 
  - ▶ For any assignment  $\alpha$ ,  $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$  iff for every  $a, b \in u(\mathcal{A})$ ,  $\mathcal{A} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
  - ▶ There is an assignment  $\alpha$  which satisfies  $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$

# Example of Satisfaction

---

- ▶  $\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$ 
  - ▶ For any assignment  $\alpha$ ,  $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$  iff for every  $a, b \in u(\mathcal{A})$ ,  $\mathcal{A} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
  - ▶ There is an assignment  $\alpha$  which satisfies  $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$
  - ▶ There is no assignment  $\alpha$  which satisfies  $\exists x \forall y (E(x, y))$
- ▶  $\mathcal{W} = (abaaa, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}}, Q_b^{\mathcal{W}})$ .
  - ▶ There is an assignment  $\alpha$  for which  $\mathcal{W} \models_{\alpha} (Q_a(x) \wedge Q_a(y) \wedge S(x, y))$

# Example of Satisfaction

---

- ▶  $\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$ 
  - ▶ For any assignment  $\alpha$ ,  $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$  iff for every  $a, b \in u(\mathcal{A})$ ,  $\mathcal{A} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
  - ▶ There is an assignment  $\alpha$  which satisfies  $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$
  - ▶ There is no assignment  $\alpha$  which satisfies  $\exists x \forall y (E(x, y))$
- ▶  $\mathcal{W} = (abaaa, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}}, Q_b^{\mathcal{W}})$ .
  - ▶ There is an assignment  $\alpha$  for which  $\mathcal{W} \models_{\alpha} (Q_a(x) \wedge Q_a(y) \wedge S(x, y))$
  - ▶ There is no assignment  $\alpha$  which satisfies  $\exists x \exists y (Q_b(x) \wedge Q_b(y) \wedge x \neq y)$

# Satisfiability, Validity and Equivalence

---

- ▶ A formula  $\varphi$  over a signature  $\tau$  is said to be **satisfiable** iff for some  $\tau$ -structure  $\mathcal{A}$  and assignment  $\alpha$ ,  $\mathcal{A} \models_{\alpha} \varphi$

# Satisfiability, Validity and Equivalence

---

- ▶ A formula  $\varphi$  over a signature  $\tau$  is said to be **satisfiable** iff for some  $\tau$ -structure  $\mathcal{A}$  and assignment  $\alpha$ ,  $\mathcal{A} \models_{\alpha} \varphi$
- ▶ A formula  $\varphi$  over a signature  $\tau$  is said to be **valid** iff for every  $\tau$ -structure  $\mathcal{A}$  and assignment  $\alpha$ ,  $\mathcal{A} \models_{\alpha} \varphi$

# Satisfiability, Validity and Equivalence

---

- ▶ A formula  $\varphi$  over a signature  $\tau$  is said to be **satisfiable** iff for some  $\tau$ -structure  $\mathcal{A}$  and assignment  $\alpha$ ,  $\mathcal{A} \models_{\alpha} \varphi$
- ▶ A formula  $\varphi$  over a signature  $\tau$  is said to be **valid** iff for every  $\tau$ -structure  $\mathcal{A}$  and assignment  $\alpha$ ,  $\mathcal{A} \models_{\alpha} \varphi$
- ▶ Formulae  $\varphi$  and  $\psi$  are **equivalent** denoted  $\varphi \equiv \psi$  iff for every  $\mathcal{A}$  and  $\alpha$ ,  $\mathcal{A} \models_{\alpha} \varphi$  iff  $\mathcal{A} \models_{\alpha} \psi$

# True or False?

---

For a formula  $\varphi$  and assignments  $\alpha_1$  and  $\alpha_2$  such that for every  $x \in \text{free}(\varphi)$ ,  $\alpha_1(x) = \alpha_2(x)$ ,  $\mathcal{A} \models_{\alpha_1} \varphi$  iff  $\mathcal{A} \models_{\alpha_2} \varphi$

# True or False?

---

For a formula  $\varphi$  and assignments  $\alpha_1$  and  $\alpha_2$  such that for every  $x \in \text{free}(\varphi)$ ,  $\alpha_1(x) = \alpha_2(x)$ ,  $\mathcal{A} \models_{\alpha_1} \varphi$  iff  $\mathcal{A} \models_{\alpha_2} \varphi$

- For example,

$$\varphi = \forall x(R(x, y) \rightarrow \forall yP(y)) \equiv \forall x(R(x, y) \rightarrow \forall zP(z))$$
$$\text{free}(\varphi) = \{y\}$$



# True or False?

---

For a formula  $\varphi$  and assignments  $\alpha_1$  and  $\alpha_2$  such that for every  $x \in \text{free}(\varphi)$ ,  $\alpha_1(x) = \alpha_2(x)$ ,  $\mathcal{A} \models_{\alpha_1} \varphi$  iff  $\mathcal{A} \models_{\alpha_2} \varphi$

- ▶ For example,  
 $\varphi = \forall x(R(x, y) \rightarrow \forall yP(y)) \equiv \forall x(R(x, y) \rightarrow \forall zP(z))$   
 $\text{free}(\varphi) = \{y\}$
- ▶ Consider two assignments  $\alpha_1, \alpha_2$  such that  
 $\alpha_1(y) = \alpha_2(y) = \alpha(\text{say})$

# True or False?

---

For a formula  $\varphi$  and assignments  $\alpha_1$  and  $\alpha_2$  such that for every  $x \in \text{free}(\varphi)$ ,  $\alpha_1(x) = \alpha_2(x)$ ,  $\mathcal{A} \models_{\alpha_1} \varphi$  iff  $\mathcal{A} \models_{\alpha_2} \varphi$

- ▶ For example,  
 $\varphi = \forall x(R(x, y) \rightarrow \forall yP(y)) \equiv \forall x(R(x, y) \rightarrow \forall zP(z))$   
 $\text{free}(\varphi) = \{y\}$
- ▶ Consider two assignments  $\alpha_1, \alpha_2$  such that  
 $\alpha_1(y) = \alpha_2(y) = \alpha(\text{say})$
- ▶ Evaluate for all  $a, b \in u(\mathcal{A})$ ,  $R(a, \alpha) \rightarrow P(b)$

# True or False?

---

For a formula  $\varphi$  and assignments  $\alpha_1$  and  $\alpha_2$  such that for every  $x \in \text{free}(\varphi)$ ,  $\alpha_1(x) = \alpha_2(x)$ ,  $\mathcal{A} \models_{\alpha_1} \varphi$  iff  $\mathcal{A} \models_{\alpha_2} \varphi$

- ▶ For example,  
 $\varphi = \forall x(R(x, y) \rightarrow \forall yP(y)) \equiv \forall x(R(x, y) \rightarrow \forall zP(z))$   
 $\text{free}(\varphi) = \{y\}$
- ▶ Consider two assignments  $\alpha_1, \alpha_2$  such that  
 $\alpha_1(y) = \alpha_2(y) = \alpha(\text{say})$
- ▶ Evaluate for all  $a, b \in u(\mathcal{A})$ ,  $R(a, \alpha) \rightarrow P(b)$
- ▶  $\mathcal{A} \models_{\alpha_1} \varphi$  iff  $\mathcal{A} \models_{\alpha_2} \varphi$

# True or False?

---

For a sentence  $\varphi$ , and any two assignments  $\alpha_1$  and  $\alpha_2$ ,  $\mathcal{A} \models_{\alpha_1} \varphi$  iff  $\mathcal{A} \models_{\alpha_2} \varphi$

# True or False?

---

For a sentence  $\varphi$ , and any two assignments  $\alpha_1$  and  $\alpha_2$ ,  $\mathcal{A} \models_{\alpha_1} \varphi$  iff  $\mathcal{A} \models_{\alpha_2} \varphi$

No free variables!