

# The Discrete Fourier Transform

Abhiram Ranade

January 13, 2016

# Tutorial

1. Which of the following statements convey information? Which can be simplified while conveying the same information?
  - 1.1 Algorithm  $A$  takes time at least  $O(n^2)$ .
  - 1.2 Algorithm  $A$  times time at most  $O(n^2)$
2. Suppose I have 27 coins, one of which is heavier than the rest, which have the same weight. You are to find the heavier coin using a two pan balance. With the balance, you can compare whether a given set of coins is heavier than another set, or is lighter, or both have equal weight. Show that you can determine the heavy coin using 3 comparisons.
3. Show that three comparisons are necessary. Hint: Argue that every algorithm must be a decision tree..
4. Harder: Suppose one coin is known to have a *different* weight than the rest – you do not know whether it is heavier or lighter. Determine the number of comparisons needed as a function of  $n$ . Give upper and lower bounds.

# Convolution using Divide and Conquer

# Convolution using Divide and Conquer

Convolution:

# Convolution using Divide and Conquer

Convolution:

Input: Numbers  $a_0, \dots, a_{n-1}$ ,  $b_0, \dots, b_{m-1}$

# Convolution using Divide and Conquer

Convolution:

Input: Numbers  $a_0, \dots, a_{n-1}, b_0, \dots, b_{m-1}$

Output:  $c_0, \dots, c_{m+n-2}$  where  $c_k = \sum_{i,j|i+j=k} a_i b_j$

# Convolution using Divide and Conquer

Convolution:

Input: Numbers  $a_0, \dots, a_{n-1}, b_0, \dots, b_{m-1}$

Output:  $c_0, \dots, c_{m+n-2}$  where  $c_k = \sum_{i,j|i+j=k} a_i b_j$

Alternative definition:

# Convolution using Divide and Conquer

Convolution:

Input: Numbers  $a_0, \dots, a_{n-1}, b_0, \dots, b_{m-1}$

Output:  $c_0, \dots, c_{m+n-2}$  where  $c_k = \sum_{i,j|i+j=k} a_i b_j$

Alternative definition:

Let  $A(x) = \sum_{i=0}^{n-1} a_i x^i, B(x) = \sum_{j=0}^{m-1} b_j x^j$



# Convolution using Divide and Conquer

Convolution:

Input: Numbers  $a_0, \dots, a_{n-1}, b_0, \dots, b_{m-1}$

Output:  $c_0, \dots, c_{m+n-2}$  where  $c_k = \sum_{i,j|i+j=k} a_i b_j$

Alternative definition:

Let  $A(x) = \sum_{i=0}^{n-1} a_i x^i, B(x) = \sum_{j=0}^{m-1} b_j x^j$

Then  $c_k$  are coefficients of  $C(x) = A(x)B(x)$

# Convolution using Divide and Conquer

Convolution:

Input: Numbers  $a_0, \dots, a_{n-1}, b_0, \dots, b_{m-1}$

Output:  $c_0, \dots, c_{m+n-2}$  where  $c_k = \sum_{i,j|i+j=k} a_i b_j$

Alternative definition:

Let  $A(x) = \sum_{i=0}^{n-1} a_i x^i, B(x) = \sum_{j=0}^{m-1} b_j x^j$

Then  $c_k$  are coefficients of  $C(x) = A(x)B(x)$

Time to convolve using "direct" method:  $O(mn)$

# Convolution using Divide and Conquer

Convolution:

Input: Numbers  $a_0, \dots, a_{n-1}, b_0, \dots, b_{m-1}$

Output:  $c_0, \dots, c_{m+n-2}$  where  $c_k = \sum_{i,j|i+j=k} a_i b_j$

Alternative definition:

Let  $A(x) = \sum_{i=0}^{n-1} a_i x^i, B(x) = \sum_{j=0}^{m-1} b_j x^j$

Then  $c_k$  are coefficients of  $C(x) = A(x)B(x)$

Time to convolve using "direct" method:  $O(mn)$

Today: Time  $O(n \log n)$

Assume  $n \geq m$

# Convolution using Divide and Conquer

Convolution:

Input: Numbers  $a_0, \dots, a_{n-1}, b_0, \dots, b_{m-1}$

Output:  $c_0, \dots, c_{m+n-2}$  where  $c_k = \sum_{i,j|i+j=k} a_i b_j$

Alternative definition:

Let  $A(x) = \sum_{i=0}^{n-1} a_i x^i, B(x) = \sum_{j=0}^{m-1} b_j x^j$

Then  $c_k$  are coefficients of  $C(x) = A(x)B(x)$

Time to convolve using "direct" method:  $O(mn)$

Today: Time  $O(n \log n)$

Assume  $n \geq m$

Polynomial multiplication is related to integer multiplication.  
Fastest integer multiplication algorithm is based on polynomial multiplication algorithms.

# Convolution using Divide and Conquer

Convolution:

Input: Numbers  $a_0, \dots, a_{n-1}, b_0, \dots, b_{m-1}$

Output:  $c_0, \dots, c_{m+n-2}$  where  $c_k = \sum_{i,j|i+j=k} a_i b_j$

Alternative definition:

Let  $A(x) = \sum_{i=0}^{n-1} a_i x^i, B(x) = \sum_{j=0}^{m-1} b_j x^j$

Then  $c_k$  are coefficients of  $C(x) = A(x)B(x)$

Time to convolve using "direct" method:  $O(mn)$

Today: Time  $O(n \log n)$

Assume  $n \geq m$

Polynomial multiplication is related to integer multiplication.  
Fastest integer multiplication algorithm is based on polynomial multiplication algorithms.

Convolution is very important in Signal Processing.

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0



# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0

Output: Pulses of heights  $b_j$  at time  $j$ , for  $j = 0, \dots, m - 1$ .

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0

Output: Pulses of heights  $b_j$  at time  $j$ , for  $j = 0, \dots, m - 1$ .

$b_0, \dots, b_{m-1}$  = characteristic of the filter. "Impulse response"

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0

Output: Pulses of heights  $b_j$  at time  $j$ , for  $j = 0, \dots, m - 1$ .

$b_0, \dots, b_{m-1}$  = characteristic of the filter. "Impulse response"

Input: Unit height pulse at time  $t$ .

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0

Output: Pulses of heights  $b_j$  at time  $j$ , for  $j = 0, \dots, m - 1$ .

$b_0, \dots, b_{m-1}$  = characteristic of the filter. "Impulse response"

Input: Unit height pulse at time  $t$ .

Output: Pulses of height  $b_j$  at time  $t + j$ , for  $j = 0, \dots, m - 1$ .

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0

Output: Pulses of heights  $b_j$  at time  $j$ , for  $j = 0, \dots, m - 1$ .

$b_0, \dots, b_{m-1}$  = characteristic of the filter. "Impulse response"

Input: Unit height pulse at time  $t$ .

Output: Pulses of height  $b_j$  at time  $t + j$ , for  $j = 0, \dots, m - 1$ .

Time invariance

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0

Output: Pulses of heights  $b_j$  at time  $j$ , for  $j = 0, \dots, m - 1$ .

$b_0, \dots, b_{m-1}$  = characteristic of the filter. "Impulse response"

Input: Unit height pulse at time  $t$ .

Output: Pulses of height  $b_j$  at time  $t + j$ , for  $j = 0, \dots, m - 1$ .

Time invariance

Input: Pulses of height  $a_0, \dots, a_{n-1}$  at times  $0, \dots, n - 1$

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0

Output: Pulses of heights  $b_j$  at time  $j$ , for  $j = 0, \dots, m-1$ .

$b_0, \dots, b_{m-1}$  = characteristic of the filter. "Impulse response"

Input: Unit height pulse at time  $t$ .

Output: Pulses of height  $b_j$  at time  $t+j$ , for  $j = 0, \dots, m-1$ .

Time invariance

Input: Pulses of height  $a_0, \dots, a_{n-1}$  at times  $0, \dots, n-1$

Output: Pulses of height  $c_k$ , at times  $k = 0, \dots, m+n-2$ , where

$$c_k = \sum_{i,j|k=i+j} a_i b_j$$

# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0

Output: Pulses of heights  $b_j$  at time  $j$ , for  $j = 0, \dots, m-1$ .

$b_0, \dots, b_{m-1}$  = characteristic of the filter. "Impulse response"

Input: Unit height pulse at time  $t$ .

Output: Pulses of height  $b_j$  at time  $t+j$ , for  $j = 0, \dots, m-1$ .

Time invariance

Input: Pulses of height  $a_0, \dots, a_{n-1}$  at times  $0, \dots, n-1$

Output: Pulses of height  $c_k$ , at times  $k = 0, \dots, m+n-2$ , where

$$c_k = \sum_{i,j|k=i+j} a_i b_j$$

Linearity



# The basic signal processing problem

Input Signal  $\rightarrow$  "Filter"  $\rightarrow$  Output signal

Action of a "linear time invariant" filter:

Input: Unit height pulse at time 0

Output: Pulses of heights  $b_j$  at time  $j$ , for  $j = 0, \dots, m-1$ .

$b_0, \dots, b_{m-1}$  = characteristic of the filter. "Impulse response"

Input: Unit height pulse at time  $t$ .

Output: Pulses of height  $b_j$  at time  $t+j$ , for  $j = 0, \dots, m-1$ .

Time invariance

Input: Pulses of height  $a_0, \dots, a_{n-1}$  at times  $0, \dots, n-1$

Output: Pulses of height  $c_k$ , at times  $k = 0, \dots, m+n-2$ , where

$$c_k = \sum_{i,j|k=i+j} a_i b_j$$

Linearity

Output is a convolution of input and impulse response

# Alternate representation of polynomials

# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

**Theorem:** There is a unique degree  $d$  polynomial through  $(x_0, y_0), (x_1, y_1), \dots, (x_d, y_d)$  if  $x_i$  are distinct.

# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

**Theorem:** There is a unique degree  $d$  polynomial through  $(x_0, y_0), (x_1, y_1), \dots, (x_d, y_d)$  if  $x_i$  are distinct.

$(x_0, A(x_0)), (x_1, A(x_1)), \dots, (x_{n-1}, A(x_{n-1}))$  is "point-value representation".

# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

**Theorem:** There is a unique degree  $d$  polynomial through  $(x_0, y_0), (x_1, y_1), \dots, (x_d, y_d)$  if  $x_i$  are distinct.

$(x_0, A(x_0)), (x_1, A(x_1)), \dots, (x_{n-1}, A(x_{n-1}))$  is "point-value representation".

Coefficient  $\rightarrow$  point-value : Evaluate!

# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

**Theorem:** There is a unique degree  $d$  polynomial through  $(x_0, y_0), (x_1, y_1), \dots, (x_d, y_d)$  if  $x_i$  are distinct.

$(x_0, A(x_0)), (x_1, A(x_1)), \dots, (x_{n-1}, A(x_{n-1}))$  is "point-value representation".

Coefficient  $\rightarrow$  point-value : Evaluate!

point-value  $\rightarrow$  Coefficient : Lagrange Interpolation

# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

**Theorem:** There is a unique degree  $d$  polynomial through  $(x_0, y_0), (x_1, y_1), \dots, (x_d, y_d)$  if  $x_i$  are distinct.

$(x_0, A(x_0)), (x_1, A(x_1)), \dots, (x_{n-1}, A(x_{n-1}))$  is "point-value representation".

Coefficient  $\rightarrow$  point-value : Evaluate!

point-value  $\rightarrow$  Coefficient : Lagrange Interpolation

Suppose  $C(x) = A(x)B(x)$ .



# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

**Theorem:** There is a unique degree  $d$  polynomial through  $(x_0, y_0), (x_1, y_1), \dots, (x_d, y_d)$  if  $x_i$  are distinct.

$(x_0, A(x_0)), (x_1, A(x_1)), \dots, (x_{n-1}, A(x_{n-1}))$  is "point-value representation".

Coefficient  $\rightarrow$  point-value : Evaluate!

point-value  $\rightarrow$  Coefficient : Lagrange Interpolation

Suppose  $C(x) = A(x)B(x)$ .

If we have the  $A, B$  in point-value form, at same  $2n$  points, then we get  $C$  in point-value form using  $2n$  multiplications!

# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

**Theorem:** There is a unique degree  $d$  polynomial through  $(x_0, y_0), (x_1, y_1), \dots, (x_d, y_d)$  if  $x_i$  are distinct.

$(x_0, A(x_0)), (x_1, A(x_1)), \dots, (x_{n-1}, A(x_{n-1}))$  is "point-value representation".

Coefficient  $\rightarrow$  point-value : Evaluate!

point-value  $\rightarrow$  Coefficient : Lagrange Interpolation

Suppose  $C(x) = A(x)B(x)$ .

If we have the  $A, B$  in point-value form, at same  $2n$  points, then we get  $C$  in point-value form using  $2n$  multiplications!

Representation of  $C$  is  $(\dots, (x_i, A(x_i)B(x_i)), \dots)$

# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

**Theorem:** There is a unique degree  $d$  polynomial through  $(x_0, y_0), (x_1, y_1), \dots, (x_d, y_d)$  if  $x_i$  are distinct.

$(x_0, A(x_0)), (x_1, A(x_1)), \dots, (x_{n-1}, A(x_{n-1}))$  is "point-value representation".

Coefficient  $\rightarrow$  point-value : Evaluate!

point-value  $\rightarrow$  Coefficient : Lagrange Interpolation

Suppose  $C(x) = A(x)B(x)$ .

If we have the  $A, B$  in point-value form, at same  $2n$  points, then we get  $C$  in point-value form using  $2n$  multiplications!

Representation of  $C$  is  $(\dots, (x_i, A(x_i)B(x_i)), \dots)$

**Alternate convolution algorithm:** Convert  $A, B$  to point-value form ( $2n$  points), pointwise multiply, Interpolate result.

# Alternate representation of polynomials

$a_0, \dots, a_{n-1}$  is "coefficient representation"

**Theorem:** There is a unique degree  $d$  polynomial through  $(x_0, y_0), (x_1, y_1), \dots, (x_d, y_d)$  if  $x_i$  are distinct.

$(x_0, A(x_0)), (x_1, A(x_1)), \dots, (x_{n-1}, A(x_{n-1}))$  is "point-value representation".

Coefficient  $\rightarrow$  point-value : Evaluate!

point-value  $\rightarrow$  Coefficient : Lagrange Interpolation

Suppose  $C(x) = A(x)B(x)$ .

If we have the  $A, B$  in point-value form, at same  $2n$  points, then we get  $C$  in point-value form using  $2n$  multiplications!

Representation of  $C$  is  $(\dots, (x_i, A(x_i)B(x_i)), \dots)$

**Alternate convolution algorithm:** Convert  $A, B$  to point-value form ( $2n$  points), pointwise multiply, Interpolate result.

Need fast evaluation/interpolation.

Can we evaluate/interpolate quickly?

# Can we evaluate/interpolate quickly?

**Naive evaluation method:** Evaluating degree  $n - 1$  polynomial at 1 point takes  $O(n)$  time. At  $2n$  points:  $O(n^2)$  time.

# Can we evaluate/interpolate quickly?

**Naive evaluation method:** Evaluating degree  $n - 1$  polynomial at 1 point takes  $O(n)$  time. At  $2n$  points:  $O(n^2)$  time.

**Main Result 1:** It is possible to evaluate a degree  $n - 1$  polynomial at  $n$  carefully chosen points in time  $O(n \log n)$ .

# Can we evaluate/interpolate quickly?

**Naive evaluation method:** Evaluating degree  $n - 1$  polynomial at 1 point takes  $O(n)$  time. At  $2n$  points:  $O(n^2)$  time.

**Main Result 1:** It is possible to evaluate a degree  $n - 1$  polynomial at  $n$  carefully chosen points in time  $O(n \log n)$ .

**Main Result 2:** It is possible to construct the coefficients of a degree  $n - 1$  polynomial given the values at  $n$  carefully chosen points in time  $O(n \log n)$ .



# Can we evaluate/interpolate quickly?

**Naive evaluation method:** Evaluating degree  $n - 1$  polynomial at 1 point takes  $O(n)$  time. At  $2n$  points:  $O(n^2)$  time.

**Main Result 1:** It is possible to evaluate a degree  $n - 1$  polynomial at  $n$  carefully chosen points in time  $O(n \log n)$ .

**Main Result 2:** It is possible to construct the coefficients of a degree  $n - 1$  polynomial given the values at  $n$  carefully chosen points in time  $O(n \log n)$ .

For convolving, we need to evaluate the polynomial at  $2n - 1$  points. To fit with the above results, we simply consider the polynomials to be of degree  $n' = 2n - 2$ , using 0s for higher coefficients. Then we use the above results substituting  $n'$  for  $n$ .

# Can we evaluate/interpolate quickly?

**Naive evaluation method:** Evaluating degree  $n - 1$  polynomial at 1 point takes  $O(n)$  time. At  $2n$  points:  $O(n^2)$  time.

**Main Result 1:** It is possible to evaluate a degree  $n - 1$  polynomial at  $n$  carefully chosen points in time  $O(n \log n)$ .

**Main Result 2:** It is possible to construct the coefficients of a degree  $n - 1$  polynomial given the values at  $n$  carefully chosen points in time  $O(n \log n)$ .

For convolving, we need to evaluate the polynomial at  $2n - 1$  points. To fit with the above results, we simply consider the polynomials to be of degree  $n' = 2n - 2$ , using 0s for higher coefficients. Then we use the above results substituting  $n'$  for  $n$ .

**Convolution time:** Evaluate + multiply + Interpolate :  
 $O(n \log n) + O(n) + O(n \log n) = O(n \log n)$

# Choosing the points at which to evaluate

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2)$$



## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2)$$

$$A(-x) = E(x^2) - xO(x^2)$$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2)$$

$$A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$

in  $O(1)$  time

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2)$$

$$A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$

in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2)$$

$$A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$

in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$

$\rightarrow$  value of  $A$  at  $\pm x_0, \pm x_1, \pm x_{n/2-1}$

in  $O(n)$  time

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2)$$

$$A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$

in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$

$\rightarrow$  value of  $A$  at  $\pm x_0, \pm x_1, \pm x_{n/2-1}$

in  $O(n)$  time

**Alternatively:** Value of  $E, O$  at  $v_0, v_1, v_{n/2-1}$

degree  $n/2 - 1$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2)$$

$$A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$

in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$

$\rightarrow$  value of  $A$  at  $\pm x_0, \pm x_1, \dots, \pm x_{n/2-1}$

in  $O(n)$  time

**Alternatively:** Value of  $E, O$  at  $v_0, v_1, v_{n/2-1}$

degree  $n/2 - 1$

$\rightarrow$  Value of  $A$  at  $\pm\sqrt{v_0}, \pm\sqrt{v_1}, \dots, \pm\sqrt{v_{n/2-1}}$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2)$$

$$A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$

in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$

$\rightarrow$  value of  $A$  at  $\pm x_0, \pm x_1, \dots, \pm x_{n/2-1}$

in  $O(n)$  time

**Alternatively:** Value of  $E, O$  at  $v_0, v_1, \dots, v_{n/2-1}$

degree  $n/2 - 1$

$\rightarrow$  Value of  $A$  at  $\pm\sqrt{v_0}, \pm\sqrt{v_1}, \dots, \pm\sqrt{v_{n/2-1}}$

Value of  $EE, OE, EO, OO$  at  $w_0, w_1, \dots, w_{n/4-1}$

degree  $n/4 - 1$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i$$

$$A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i$$

$$\text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2)$$

$$A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$

in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$

$\rightarrow$  value of  $A$  at  $\pm x_0, \pm x_1, \dots, \pm x_{n/2-1}$

in  $O(n)$  time

**Alternatively:** Value of  $E, O$  at  $v_0, v_1, \dots, v_{n/2-1}$

degree  $n/2 - 1$

$\rightarrow$  Value of  $A$  at  $\pm\sqrt{v_0}, \pm\sqrt{v_1}, \dots, \pm\sqrt{v_{n/2-1}}$

Value of  $EE, OE, EO, OO$  at  $w_0, w_1, \dots, w_{n/4-1}$

degree  $n/4 - 1$

$\rightarrow$  Value of  $E, O$  at  $\pm\sqrt{w_0}, \pm\sqrt{w_1}, \dots, \pm\sqrt{w_{n/4-1}}$



## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i \qquad A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i \qquad \text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2) \qquad A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$  in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$   
 $\rightarrow$  value of  $A$  at  $\pm x_0, \pm x_1, \pm x_{n/2-1}$  in  $O(n)$  time

**Alternatively:** Value of  $E, O$  at  $v_0, v_1, v_{n/2-1}$  degree  $n/2 - 1$   
 $\rightarrow$  Value of  $A$  at  $\pm\sqrt{v_0}, \pm\sqrt{v_1}, \dots, \pm\sqrt{v_{n/2-1}}$

Value of  $EE, OE, EO, OO$  at  $w_0, w_1, \dots, w_{n/4-1}$  degree  $n/4 - 1$   
 $\rightarrow$  Value of  $E, O$  at  $\pm\sqrt{w_0}, \pm\sqrt{w_1}, \dots, \pm\sqrt{w_{n/4-1}}$   
 $\rightarrow$  Value of  $A$  at 4th roots of each of  $w_0, w_1, \dots, w_{n/4-1}$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i \qquad A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i \qquad \text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2) \qquad A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$  in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$   
 $\rightarrow$  value of  $A$  at  $\pm x_0, \pm x_1, \pm x_{n/2-1}$  in  $O(n)$  time

**Alternatively:** Value of  $E, O$  at  $v_0, v_1, v_{n/2-1}$  degree  $n/2 - 1$   
 $\rightarrow$  Value of  $A$  at  $\pm\sqrt{v_0}, \pm\sqrt{v_1}, \dots, \pm\sqrt{v_{n/2-1}}$

Value of  $EE, OE, EO, OO$  at  $w_0, w_1, \dots, w_{n/4-1}$  degree  $n/4 - 1$   
 $\rightarrow$  Value of  $E, O$  at  $\pm\sqrt{w_0}, \pm\sqrt{w_1}, \dots, \pm\sqrt{w_{n/4-1}}$   
 $\rightarrow$  Value of  $A$  at 4th roots of each of  $w_0, w_1, \dots, w_{n/4-1}$

Value of  $n$  degree-0-polynomials at single point  $u_0$

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i \qquad A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i \qquad \text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$

$$A(x) = E(x^2) + xO(x^2) \qquad A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$  in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$   
 $\rightarrow$  value of  $A$  at  $\pm x_0, \pm x_1, \pm x_{n/2-1}$  in  $O(n)$  time

**Alternatively:** Value of  $E, O$  at  $v_0, v_1, v_{n/2-1}$  degree  $n/2 - 1$   
 $\rightarrow$  Value of  $A$  at  $\pm\sqrt{v_0}, \pm\sqrt{v_1}, \dots, \pm\sqrt{v_{n/2-1}}$

Value of  $EE, OE, EO, OO$  at  $w_0, w_1, \dots, w_{n/4-1}$  degree  $n/4 - 1$   
 $\rightarrow$  Value of  $E, O$  at  $\pm\sqrt{w_0}, \pm\sqrt{w_1}, \dots, \pm\sqrt{w_{n/4-1}}$   
 $\rightarrow$  Value of  $A$  at 4th roots of each of  $w_0, w_1, \dots, w_{n/4-1}$

Value of  $n$  degree-0-polynomials at single point  $u_0$   
 $\rightarrow \dots \rightarrow$  Value of  $A$  at  $n$   $n$ th roots of  $u_0$ .

## Choosing the points at which to evaluate

$$A(x) = \sum_i a_i x^i \qquad A(-x) = \sum_{i \text{ even}} a_i x^i - \sum_{i \text{ odd}} a_i x^i$$

$$\text{"even(A)": } E(y) = \sum_i a_{2i} y^i \qquad \text{"odd(A)": } O(y) = \sum_i a_{2i+1} y^i$$


$$A(x) = E(x^2) + xO(x^2) \qquad A(-x) = E(x^2) - xO(x^2)$$

Value of  $E, O$  at  $x^2 \rightarrow$  value of  $A$  at  $\pm x$  in  $O(1)$  time

Value of  $E, O$  at  $x_0^2, x_1^2, \dots, x_{n/2-1}^2$   
 $\rightarrow$  value of  $A$  at  $\pm x_0, \pm x_1, \pm x_{n/2-1}$  in  $O(n)$  time

**Alternatively:** Value of  $E, O$  at  $v_0, v_1, v_{n/2-1}$  degree  $n/2 - 1$   
 $\rightarrow$  Value of  $A$  at  $\pm\sqrt{v_0}, \pm\sqrt{v_1}, \dots, \pm\sqrt{v_{n/2-1}}$

Value of  $EE, OE, EO, OO$  at  $w_0, w_1, \dots, w_{n/4-1}$  degree  $n/4 - 1$   
 $\rightarrow$  Value of  $E, O$  at  $\pm\sqrt{w_0}, \pm\sqrt{w_1}, \dots, \pm\sqrt{w_{n/4-1}}$   
 $\rightarrow$  Value of  $A$  at 4th roots of each of  $w_0, w_1, \dots, w_{n/4-1}$

Value of  $n$  degree-0-polynomials at single point  $u_0$   
 $\rightarrow \dots \rightarrow$  Value of  $A$  at  $n$   $n$ th roots of  $u_0$ . Choose  $u_0 = 1$  

# On complex roots of 1

## On complex roots of 1

$\omega_n = e^{2\pi\sqrt{-1}/n}$  is a principal  $n$ th root of 1.

## On complex roots of 1

$\omega_n = e^{2\pi\sqrt{-1}/n}$  is a principal  $n$ th root of 1.

Let  $\omega = \omega_n$

## On complex roots of 1

$\omega_n = e^{2\pi\sqrt{-1}/n}$  is a principal  $n$ th root of 1.

Let  $\omega = \omega_n$

$\omega^0, \omega^1, \omega^2, \dots, \omega^{n-1}$  are all  $n$ th roots of 1.



# On complex roots of 1

$\omega_n = e^{2\pi\sqrt{-1}/n}$  is a principal  $n$ th root of 1.

Let  $\omega = \omega_n$

$\omega^0, \omega^1, \omega^2, \dots, \omega^{n-1}$  are all  $n$ th roots of 1.

$A$  is evaluated at  $\omega^0, \dots, \omega^{n-1}$

## On complex roots of 1

$\omega_n = e^{2\pi\sqrt{-1}/n}$  is a principal  $n$ th root of 1.

Let  $\omega = \omega_n$

$\omega^0, \omega^1, \omega^2, \dots, \omega^{n-1}$  are all  $n$ th roots of 1.

$A$  is evaluated at  $\omega^0, \dots, \omega^{n-1}$

Note that second half of sequence = negative of first half

# On complex roots of 1

$\omega_n = e^{2\pi\sqrt{-1}/n}$  is a principal  $n$ th root of 1.

Let  $\omega = \omega_n$

$\omega^0, \omega^1, \omega^2, \dots, \omega^{n-1}$  are all  $n$ th roots of 1.

$A$  is evaluated at  $\omega^0, \dots, \omega^{n-1}$

Note that second half of sequence = negative of first half

$E, O$  are evaluated at  $(\omega^0)^2, \dots, (\omega^i)^2, \dots, (\omega^{n/2-1})^2$

# On complex roots of 1

$\omega_n = e^{2\pi\sqrt{-1}/n}$  is a principal  $n$ th root of 1.

Let  $\omega = \omega_n$

$\omega^0, \omega^1, \omega^2, \dots, \omega^{n-1}$  are all  $n$ th roots of 1.

$A$  is evaluated at  $\omega^0, \dots, \omega^{n-1}$

Note that second half of sequence = negative of first half

$E, O$  are evaluated at  $(\omega^0)^2, \dots, (\omega^i)^2, \dots, (\omega^{n/2-1})^2$

$$(\omega^i)^2 = (e^{2\pi\sqrt{-1}/n})^{2i} = (e^{2\pi\sqrt{-1}/(n/2)})^i = \omega_{n/2}^i$$

# On complex roots of 1

$\omega_n = e^{2\pi\sqrt{-1}/n}$  is a principal  $n$ th root of 1.

Let  $\omega = \omega_n$

$\omega^0, \omega^1, \omega^2, \dots, \omega^{n-1}$  are all  $n$ th roots of 1.

$A$  is evaluated at  $\omega^0, \dots, \omega^{n-1}$

Note that second half of sequence = negative of first half

$E, O$  are evaluated at  $(\omega^0)^2, \dots, (\omega^i)^2, \dots, (\omega^{n/2-1})^2$

$$(\omega^i)^2 = (e^{2\pi\sqrt{-1}/n})^{2i} = (e^{2\pi\sqrt{-1}/(n/2)})^i = \omega_{n/2}^i$$

$E, O$  are evaluated at powers of  $\omega_{n/2} = \omega^2$ .

# On complex roots of 1

$\omega_n = e^{2\pi\sqrt{-1}/n}$  is a principal  $n$ th root of 1.

Let  $\omega = \omega_n$

$\omega^0, \omega^1, \omega^2, \dots, \omega^{n-1}$  are all  $n$ th roots of 1.

$A$  is evaluated at  $\omega^0, \dots, \omega^{n-1}$

Note that second half of sequence = negative of first half

$E, O$  are evaluated at  $(\omega^0)^2, \dots, (\omega^i)^2, \dots, (\omega^{n/2-1})^2$

$$(\omega^i)^2 = (e^{2\pi\sqrt{-1}/n})^{2i} = (e^{2\pi\sqrt{-1}/(n/2)})^i = \omega_{n/2}^i$$

$E, O$  are evaluated at powers of  $\omega_{n/2} = \omega^2$ .

$E, O$  can be evaluated recursively:  $n/2 - 1$  degree polynomial to be evaluated at  $n/2$   $n/2$ th roots of 1.

Evaluate( $a_0, a_1, \dots, a_{n-1}, \omega$ ) {

Evaluate( $a_0, a_1, \dots, a_{n-1}, \omega$ ) {

// Evaluate  $A$  at  $\omega^0, \dots, \omega^{n-1}$



Evaluate( $a_0, a_1, \dots, a_{n-1}, \omega$ ) {

// Evaluate  $A$  at  $\omega^0, \dots, \omega^{n-1}$

//  $\omega^{n/2+i} = -\omega^i$  for  $i = 0, \dots, n/2 - 1$ .

Evaluate( $a_0, a_1, \dots, a_{n-1}, \omega$ ) {

// Evaluate  $A$  at  $\omega^0, \dots, \omega^{n-1}$

//  $\omega^{n/2+i} = -\omega^i$  for  $i = 0, \dots, n/2 - 1$ .

1. Base case: if  $n = 1$  return  $a_0$
2.  $E = \text{Evaluate}(a_0, a_2, \dots, a_{n-2}, \omega^2)$
3.  $O = \text{Evaluate}(a_1, a_3, \dots, a_{n-1}, \omega^2)$
4. for  $i = 0$  to  $n/2 - 1$
5.      $A[i] = E[i] + \omega^i O[i]$       $A[n/2 + i] = E[i] - \omega^i O[i]$
6. end for
7. Return  $A$  // array

Evaluate( $a_0, a_1, \dots, a_{n-1}, \omega$ ) {

// Evaluate  $A$  at  $\omega^0, \dots, \omega^{n-1}$

//  $\omega^{n/2+i} = -\omega^i$  for  $i = 0, \dots, n/2 - 1$ .

1. Base case: if  $n = 1$  return  $a_0$
2.  $E = \text{Evaluate}(a_0, a_2, \dots, a_{n-2}, \omega^2)$
3.  $O = \text{Evaluate}(a_1, a_3, \dots, a_{n-1}, \omega^2)$
4. for  $i = 0$  to  $n/2 - 1$
5.      $A[i] = E[i] + \omega^i O[i]$       $A[n/2 + i] = E[i] - \omega^i O[i]$
6. end for
7. Return  $A$  // array

$$T(n) = 2T(n/2) + O(n)$$

Evaluate( $a_0, a_1, \dots, a_{n-1}, \omega$ ) {

// Evaluate  $A$  at  $\omega^0, \dots, \omega^{n-1}$

//  $\omega^{n/2+i} = -\omega^i$  for  $i = 0, \dots, n/2 - 1$ .

1. Base case: if  $n = 1$  return  $a_0$
2.  $E = \text{Evaluate}(a_0, a_2, \dots, a_{n-2}, \omega^2)$
3.  $O = \text{Evaluate}(a_1, a_3, \dots, a_{n-1}, \omega^2)$
4. for  $i = 0$  to  $n/2 - 1$
5.      $A[i] = E[i] + \omega^i O[i]$       $A[n/2 + i] = E[i] - \omega^i O[i]$
6. end for
7. Return  $A$  // array

$$T(n) = 2T(n/2) + O(n)$$

So  $T(n) = O(n \log n)$

# Polynomial Evaluation as matrix vector multiplication

$$A(\omega^i) = \sum_{j=0}^{n-1} a_j (\omega^i)^j$$

# Polynomial Evaluation as matrix vector multiplication

$$A(\omega^i) = \sum_{j=0}^{n-1} a_j (\omega^i)^j = \sum_{j=0}^{n-1} a_j \omega^{ij}$$

# Polynomial Evaluation as matrix vector multiplication

$$A(\omega^i) = \sum_{j=0}^{n-1} a_j (\omega^i)^j = \sum_{j=0}^{n-1} a_j \omega^{ij}$$

$$\begin{bmatrix} \vdots \\ A(\omega^i) \\ \vdots \end{bmatrix} = \begin{bmatrix} \omega^0 & \omega^i & \dots & \omega^{ij} & \dots & \omega^{i(n-1)} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_j \\ \vdots \\ a_{n-1} \end{bmatrix}$$

# Polynomial Evaluation as matrix vector multiplication

$$A(\omega^i) = \sum_{j=0}^{n-1} a_j (\omega^i)^j = \sum_{j=0}^{n-1} a_j \omega^{ij}$$

$$\begin{bmatrix} \vdots \\ A(\omega^i) \\ \vdots \end{bmatrix} = \begin{bmatrix} \vdots & & & \vdots & & \\ \omega^0 & \omega^i & \dots & \omega^{ij} & \dots & \omega^{i(n-1)} \\ & & & \vdots & & \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_j \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Interpolation = multiplying lhs by inverse of matrix,  $M(\omega)$ .



# Polynomial Evaluation as matrix vector multiplication

$$A(\omega^i) = \sum_{j=0}^{n-1} a_j (\omega^i)^j = \sum_{j=0}^{n-1} a_j \omega^{ij}$$

$$\begin{bmatrix} \vdots \\ A(\omega^i) \\ \vdots \end{bmatrix} = \begin{bmatrix} \vdots & & & \vdots & & \\ \omega^0 & \omega^i & \dots & \omega^{ij} & \dots & \omega^{i(n-1)} \\ & & & \vdots & & \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_j \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Interpolation = multiplying lhs by inverse of matrix,  $M(\omega)$ .

Matrix  $M(\omega)$  has orthogonal columns!

Next

# Polynomial Evaluation as matrix vector multiplication

$$A(\omega^i) = \sum_{j=0}^{n-1} a_j (\omega^i)^j = \sum_{j=0}^{n-1} a_j \omega^{ij}$$

$$\begin{bmatrix} \vdots \\ A(\omega^i) \\ \vdots \end{bmatrix} = \begin{bmatrix} \vdots & & & \vdots & & \\ \omega^0 & \omega^i & \dots & \omega^{ij} & \dots & \omega^{i(n-1)} \\ & & & \vdots & & \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_j \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Interpolation = multiplying lhs by inverse of matrix,  $M(\omega)$ .

Matrix  $M(\omega)$  has orthogonal columns!

Each column has norm  $\sqrt{n}$

Next

Next

# Polynomial Evaluation as matrix vector multiplication

$$A(\omega^i) = \sum_{j=0}^{n-1} a_j (\omega^i)^j = \sum_{j=0}^{n-1} a_j \omega^{ij}$$

$$\begin{bmatrix} \vdots \\ A(\omega^i) \\ \vdots \end{bmatrix} = \begin{bmatrix} \vdots & & & \vdots & & \\ \omega^0 & \omega^i & \dots & \omega^{ij} & \dots & \omega^{i(n-1)} \\ & & & \vdots & & \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_j \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Interpolation = multiplying lhs by inverse of matrix,  $M(\omega)$ .

Matrix  $M(\omega)$  has orthogonal columns!

Next

Each column has norm  $\sqrt{n}$

Next

Matrix inverse =  $\frac{1}{n} M(\omega)^* = \frac{1}{n} M(\omega^{-1})$

Next

# Polynomial Evaluation as matrix vector multiplication

$$A(\omega^i) = \sum_{j=0}^{n-1} a_j (\omega^i)^j = \sum_{j=0}^{n-1} a_j \omega^{ij}$$

$$\begin{bmatrix} \vdots \\ A(\omega^i) \\ \vdots \end{bmatrix} = \begin{bmatrix} \omega^0 & \omega^i & \dots & \omega^{ij} & \dots & \omega^{i(n-1)} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_j \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Interpolation = multiplying lhs by inverse of matrix,  $M(\omega)$ .

Matrix  $M(\omega)$  has orthogonal columns!

Next

Each column has norm  $\sqrt{n}$

Next

Matrix inverse =  $\frac{1}{n} M(\omega)^* = \frac{1}{n} M(\omega^{-1})$

Next

Multiplication by inverse is similar to multiplication by original.

# Polynomial Evaluation as matrix vector multiplication

$$A(\omega^i) = \sum_{j=0}^{n-1} a_j (\omega^i)^j = \sum_{j=0}^{n-1} a_j \omega^{ij}$$

$$\begin{bmatrix} \vdots \\ A(\omega^i) \\ \vdots \end{bmatrix} = \begin{bmatrix} \omega^0 & \omega^i & \dots & \omega^{ij} & \dots & \omega^{i(n-1)} \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ \vdots \\ a_j \\ \vdots \\ a_{n-1} \end{bmatrix}$$

Interpolation = multiplying lhs by inverse of matrix,  $M(\omega)$ .

Matrix  $M(\omega)$  has orthogonal columns!

Next

Each column has norm  $\sqrt{n}$

Next

Matrix inverse =  $\frac{1}{n} M(\omega)^* = \frac{1}{n} M(\omega^{-1})$

Next

Multiplication by inverse is similar to multiplication by original.

Interpolation time is also  $O(n \log n)$ .

# Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

## Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

To find its norm, we have to multiply by conjugate transpose

# Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

To find its norm, we have to multiply by conjugate transpose

Conjugate of  $\omega$  is  $\omega^{-1}$

Argand diagram



# Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

To find its norm, we have to multiply by conjugate transpose

Conjugate of  $\omega$  is  $\omega^{-1}$

Argand diagram

Conjugate of  $\omega^{ik}$  is  $\omega^{-ik}$ .

# Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

To find its norm, we have to multiply by conjugate transpose

Conjugate of  $\omega$  is  $\omega^{-1}$

Argand diagram

Conjugate of  $\omega^{ik}$  is  $\omega^{-ik}$ .

$$R_i R_i^* = \sum_k \omega^{ik} (\omega^{-ik}) = n$$

# Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

To find its norm, we have to multiply by conjugate transpose

Conjugate of  $\omega$  is  $\omega^{-1}$

Argand diagram

Conjugate of  $\omega^{ik}$  is  $\omega^{-ik}$ .

$$R_i R_i^* = \sum_k \omega^{ik} (\omega^{-ik}) = n$$

Hence norm =  $\sqrt{n}$ .

# Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

To find its norm, we have to multiply by conjugate transpose

Conjugate of  $\omega$  is  $\omega^{-1}$

Argand diagram

Conjugate of  $\omega^{ik}$  is  $\omega^{-ik}$ .

$$R_i R_i^* = \sum_k \omega^{ik} (\omega^{-ik}) = n$$

Hence norm =  $\sqrt{n}$ .

$$\text{For } i \neq j : R_i R_j^* = \sum_k \omega^{ik} (\omega^{-jk}) = \sum_{k=0}^{n-1} \omega^{(i-j)k}$$

# Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

To find its norm, we have to multiply by conjugate transpose

Conjugate of  $\omega$  is  $\omega^{-1}$

Argand diagram

Conjugate of  $\omega^{ik}$  is  $\omega^{-ik}$ .

$$R_i R_i^* = \sum_k \omega^{ik} (\omega^{-ik}) = n$$

Hence norm =  $\sqrt{n}$ .

$$\text{For } i \neq j : R_i R_j^* = \sum_k \omega^{ik} (\omega^{-jk}) = \sum_{k=0}^{n-1} \omega^{(i-j)k}$$

$$= \frac{(\omega^{i-j})^n - 1}{\omega^{i-j} - 1}$$

Sum of geometric series

# Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

To find its norm, we have to multiply by conjugate transpose

Conjugate of  $\omega$  is  $\omega^{-1}$

Argand diagram

Conjugate of  $\omega^{ik}$  is  $\omega^{-ik}$ .

$$R_i R_i^* = \sum_k \omega^{ik} (\omega^{-ik}) = n$$

Hence norm =  $\sqrt{n}$ .

$$\text{For } i \neq j : R_i R_j^* = \sum_k \omega^{ik} (\omega^{-jk}) = \sum_{k=0}^{n-1} \omega^{(i-j)k}$$

$$= \frac{(\omega^{i-j})^n - 1}{\omega^{i-j} - 1}$$

$$= 0$$

Sum of geometric series

$$\omega^{in} = \omega^{jn} = 1$$

# Conjugation, orthogonality and norm

$i$ th row of  $M(\omega)$  is:  $R_i = [\dots \omega^{ik} \dots]$

To find its norm, we have to multiply by conjugate transpose

Conjugate of  $\omega$  is  $\omega^{-1}$

Argand diagram

Conjugate of  $\omega^{ik}$  is  $\omega^{-ik}$ .

$$R_i R_i^* = \sum_k \omega^{ik} (\omega^{-ik}) = n$$

Hence norm =  $\sqrt{n}$ .

$$\text{For } i \neq j : R_i R_j^* = \sum_k \omega^{ik} (\omega^{-jk}) = \sum_{k=0}^{n-1} \omega^{(i-j)k}$$

$$= \frac{(\omega^{i-j})^n - 1}{\omega^{i-j} - 1}$$

$$= 0$$

Sum of geometric series

$$\omega^{in} = \omega^{jn} = 1$$

Hence orthogonal.

# Remarks



# Remarks

- ▶ Values at roots of unity = Discrete Fourier Transform of coefficients.

# Remarks

- ▶ Values at roots of unity = Discrete Fourier Transform of coefficients.
- ▶ Algorithm is called FFT : Fast Fourier Transform.  
Cooley-Tukey 1965.

# Remarks

- ▶ Values at roots of unity = Discrete Fourier Transform of coefficients.
- ▶ Algorithm is called FFT : Fast Fourier Transform.  
Cooley-Tukey 1965.  
Like many great ideas, it was apparently known to Gauss!

# Remarks

- ▶ Values at roots of unity = Discrete Fourier Transform of coefficients.
- ▶ Algorithm is called FFT : Fast Fourier Transform.  
Cooley-Tukey 1965.  
Like many great ideas, it was apparently known to Gauss!
- ▶ Many ideas. Multiple representations. Exercising your freedom carefully (where to evaluate). Divide and conquer.

# Remarks

- ▶ Values at roots of unity = Discrete Fourier Transform of coefficients.
- ▶ Algorithm is called FFT : Fast Fourier Transform.  
Cooley-Tukey 1965.  
Like many great ideas, it was apparently known to Gauss!
- ▶ Many ideas. Multiple representations. Exercising your freedom carefully (where to evaluate). Divide and conquer.
- ▶ Version exists for ring of integers modulo a number. Useful for doing arithmetic.

# Remarks

- ▶ Values at roots of unity = Discrete Fourier Transform of coefficients.
- ▶ Algorithm is called FFT : Fast Fourier Transform.  
Cooley-Tukey 1965.  
Like many great ideas, it was apparently known to Gauss!
- ▶ Many ideas. Multiple representations. Exercising your freedom carefully (where to evaluate). Divide and conquer.
- ▶ Version exists for ring of integers modulo a number. Useful for doing arithmetic.
- ▶ Reading: [Dasgupta, Papadimitriou, Vazirani].