



CS 228 : Logic in Computer Science

Krishna. S

Monadic Second Order Logic (MSO)

Symbols in MSO

Formulae of MSO, over signature τ , are sequences of symbols, where each symbol is one of the following:

Symbols in MSO

Formulae of MSO, over signature τ , are sequences of symbols, where each symbol is one of the following:

- ▶ The symbol \perp called **false**

Symbols in MSO

Formulae of MSO, over signature τ , are sequences of symbols, where each symbol is one of the following:

- ▶ The symbol \perp called **false**
- ▶ An element of the infinite set $\mathcal{V}_1 = \{x_1, x_2, \dots\}$ of **first order variables**

Symbols in MSO

Formulae of MSO, over signature τ , are sequences of symbols, where each symbol is one of the following:

- ▶ The symbol \perp called **false**
- ▶ An element of the infinite set $\mathcal{V}_1 = \{x_1, x_2, \dots\}$ of **first order variables**
- ▶ An element of the infinite set $\mathcal{V}_2 = \{X_1, X_2, \dots\}$ of **second order variables**, where each variable has arity 1

Symbols in MSO

Formulae of MSO, over signature τ , are sequences of symbols, where each symbol is one of the following:

- ▶ The symbol \perp called **false**
- ▶ An element of the infinite set $\mathcal{V}_1 = \{x_1, x_2, \dots\}$ of **first order variables**
- ▶ An element of the infinite set $\mathcal{V}_2 = \{X_1, X_2, \dots\}$ of **second order variables**, where each variable has arity 1
- ▶ Constants and relations from τ

Symbols in MSO

Formulae of MSO, over signature τ , are sequences of symbols, where each symbol is one of the following:

- ▶ The symbol \perp called **false**
- ▶ An element of the infinite set $\mathcal{V}_1 = \{x_1, x_2, \dots\}$ of **first order variables**
- ▶ An element of the infinite set $\mathcal{V}_2 = \{X_1, X_2, \dots\}$ of **second order variables**, where each variable has arity 1
- ▶ Constants and relations from τ
- ▶ The symbol \rightarrow called **implication**

Symbols in MSO

Formulae of MSO, over signature τ , are sequences of symbols, where each symbol is one of the following:

- ▶ The symbol \perp called **false**
- ▶ An element of the infinite set $\mathcal{V}_1 = \{x_1, x_2, \dots\}$ of **first order variables**
- ▶ An element of the infinite set $\mathcal{V}_2 = \{X_1, X_2, \dots\}$ of **second order variables**, where each variable has arity 1
- ▶ Constants and relations from τ
- ▶ The symbol \rightarrow called **implication**
- ▶ The symbol \forall called the **universal quantifier**

Symbols in MSO

Formulae of MSO, over signature τ , are sequences of symbols, where each symbol is one of the following:

- ▶ The symbol \perp called **false**
- ▶ An element of the infinite set $\mathcal{V}_1 = \{x_1, x_2, \dots\}$ of **first order variables**
- ▶ An element of the infinite set $\mathcal{V}_2 = \{X_1, X_2, \dots\}$ of **second order variables**, where each variable has arity 1
- ▶ Constants and relations from τ
- ▶ The symbol \rightarrow called **implication**
- ▶ The symbol \forall called the **universal quantifier**
- ▶ The symbols (and) called **paranthesis**

Well formed Formulae

A well-formed formula (wff) over a signature τ is inductively defined as follows:

Well formed Formulae

A well-formed formula (wff) over a signature τ is inductively defined as follows:

- ▶ \perp is a wff

Well formed Formulae

A well-formed formula (wff) over a signature τ is inductively defined as follows:

- ▶ \perp is a wff
- ▶ If t_1, t_2 are either variables or constants in τ , then $t_1 = t_2$ is a wff

Well formed Formulae

A well-formed formula (wff) over a signature τ is inductively defined as follows:

- ▶ \perp is a wff
- ▶ If t_1, t_2 are either variables or constants in τ , then $t_1 = t_2$ is a wff
- ▶ If t_i is either a first order variable or a constant, for $1 \leq i \leq k$ and R is a k -ary relation symbol in τ , then $R(t_1, \dots, t_k)$ is a wff

Well formed Formulae

A well-formed formula (wff) over a signature τ is inductively defined as follows:

- ▶ \perp is a wff
- ▶ If t_1, t_2 are either variables or constants in τ , then $t_1 = t_2$ is a wff
- ▶ If t_i is either a first order variable or a constant, for $1 \leq i \leq k$ and R is a k -ary relation symbol in τ , then $R(t_1, \dots, t_k)$ is a wff
- ▶ If t is either a first order variable or a constant, X is a second order variable, then $X(t)$ is a wff

Well formed Formulae

A well-formed formula (wff) over a signature τ is inductively defined as follows:

- ▶ \perp is a wff
- ▶ If t_1, t_2 are either variables or constants in τ , then $t_1 = t_2$ is a wff
- ▶ If t_i is either a first order variable or a constant, for $1 \leq i \leq k$ and R is a k -ary relation symbol in τ , then $R(t_1, \dots, t_k)$ is a wff
- ▶ If t is either a first order variable or a constant, X is a second order variable, then $X(t)$ is a wff
- ▶ If φ and ψ are wff, then $\varphi \rightarrow \psi$ is a wff

Well formed Formulae

A well-formed formula (wff) over a signature τ is inductively defined as follows:

- ▶ \perp is a wff
- ▶ If t_1, t_2 are either variables or constants in τ , then $t_1 = t_2$ is a wff
- ▶ If t_i is either a first order variable or a constant, for $1 \leq i \leq k$ and R is a k -ary relation symbol in τ , then $R(t_1, \dots, t_k)$ is a wff
- ▶ If t is either a first order variable or a constant, X is a second order variable, then $X(t)$ is a wff
- ▶ If φ and ψ are wff, then $\varphi \rightarrow \psi$ is a wff
- ▶ If φ is a wff and x is a first order variable, then $(\forall x)\varphi$ is a wff

Well formed Formulae

A well-formed formula (wff) over a signature τ is inductively defined as follows:

- ▶ \perp is a wff
- ▶ If t_1, t_2 are either variables or constants in τ , then $t_1 = t_2$ is a wff
- ▶ If t_i is either a first order variable or a constant, for $1 \leq i \leq k$ and R is a k -ary relation symbol in τ , then $R(t_1, \dots, t_k)$ is a wff
- ▶ If t is either a first order variable or a constant, X is a second order variable, then $X(t)$ is a wff
- ▶ If φ and ψ are wff, then $\varphi \rightarrow \psi$ is a wff
- ▶ If φ is a wff and x is a first order variable, then $(\forall x)\varphi$ is a wff
- ▶ If φ is a wff and X is a second order variable, then $(\forall X)\varphi$ is a wff

Free and Bound Variables

- ▶ Free, Bound Variables and Scope same as in FO

Free and Bound Variables

- ▶ Free, Bound Variables and Scope same as in FO
- ▶ In a wff $\varphi = \forall X\psi$, every occurrence of X in ψ is bound

Free and Bound Variables

- ▶ Free, Bound Variables and Scope same as in FO
- ▶ In a wff $\varphi = \forall X\psi$, every occurrence of X in ψ is bound
- ▶ A sentence is a formula with no free first order and second order variables

Assignments on τ -structures

Assignments

For a τ -structure \mathcal{A} , an assignment over \mathcal{A} is a pair of functions (α_1, α_2) , where

- ▶ $\alpha_1 : \mathcal{V}_1 \rightarrow u(\mathcal{A})$ assigns every first order variable $x \in \mathcal{V}_1$ a value $\alpha_1(x) \in u(\mathcal{A})$. If t is a constant symbol c , then $\alpha_1(t)$ is $c^{\mathcal{A}}$.

Assignments on τ -structures

Assignments

For a τ -structure \mathcal{A} , an assignment over \mathcal{A} is a pair of functions (α_1, α_2) , where

- ▶ $\alpha_1 : \mathcal{V}_1 \rightarrow u(\mathcal{A})$ assigns every first order variable $x \in \mathcal{V}_1$ a value $\alpha_1(x) \in u(\mathcal{A})$. If t is a constant symbol c , then $\alpha_1(t)$ is $c^{\mathcal{A}}$.
- ▶ $\alpha_2 : \mathcal{V}_2 \rightarrow 2^{u(\mathcal{A})}$ assigns to every second order variable $X \in \mathcal{V}_2$, $\alpha_2(X) \subseteq u(\mathcal{A})$.

Assignments on τ -structures

Assignments

For a τ -structure \mathcal{A} , an assignment over \mathcal{A} is a pair of functions (α_1, α_2) , where

- ▶ $\alpha_1 : \mathcal{V}_1 \rightarrow u(\mathcal{A})$ assigns every first order variable $x \in \mathcal{V}_1$ a value $\alpha_1(x) \in u(\mathcal{A})$. If t is a constant symbol c , then $\alpha_1(t)$ is $c^{\mathcal{A}}$.
- ▶ $\alpha_2 : \mathcal{V}_2 \rightarrow 2^{u(\mathcal{A})}$ assigns to every second order variable $X \in \mathcal{V}_2$, $\alpha_2(X) \subseteq u(\mathcal{A})$.

Binding on a Variable

For an assignment $\alpha = (\alpha_1, \alpha_2)$ over \mathcal{A} , and $x \in \mathcal{V}_i$, $i = 1, 2$, $\alpha_i[x \mapsto a]$ is the assignment $\alpha_i[x \mapsto a](y) = \begin{cases} \alpha_i(y), & y \neq x, \\ a, & y = x \end{cases}$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha_1(t_1) = \alpha_1(t_2)$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha_1(t_1) = \alpha_1(t_2)$
- ▶ $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$ iff $(\alpha_1(t_1), \dots, \alpha_1(t_k)) \in R^{\mathcal{A}}$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha_1(t_1) = \alpha_1(t_2)$
- ▶ $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$ iff $(\alpha_1(t_1), \dots, \alpha_1(t_k)) \in R^{\mathcal{A}}$
- ▶ $\mathcal{A} \models_{\alpha} X(t)$ iff $\alpha_1(t) \in \alpha_2(X)$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha_1(t_1) = \alpha_1(t_2)$
- ▶ $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$ iff $(\alpha_1(t_1), \dots, \alpha_1(t_k)) \in R^{\mathcal{A}}$
- ▶ $\mathcal{A} \models_{\alpha} X(t)$ iff $\alpha_1(t) \in \alpha_2(X)$
- ▶ $\mathcal{A} \models_{\alpha} (\varphi \rightarrow \psi)$ iff $\mathcal{A} \not\models_{\alpha} \varphi$ or $\mathcal{A} \models_{\alpha} \psi$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha_1(t_1) = \alpha_1(t_2)$
- ▶ $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$ iff $(\alpha_1(t_1), \dots, \alpha_1(t_k)) \in R^{\mathcal{A}}$
- ▶ $\mathcal{A} \models_{\alpha} X(t)$ iff $\alpha_1(t) \in \alpha_2(X)$
- ▶ $\mathcal{A} \models_{\alpha} (\varphi \rightarrow \psi)$ iff $\mathcal{A} \not\models_{\alpha} \varphi$ or $\mathcal{A} \models_{\alpha} \psi$
- ▶ $\mathcal{A} \models_{\alpha} (\forall x)\varphi$ iff for every $a \in u(\mathcal{A})$, $\mathcal{A} \models_{\alpha[x \mapsto a]} \varphi$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha_1(t_1) = \alpha_1(t_2)$
- ▶ $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$ iff $(\alpha_1(t_1), \dots, \alpha_1(t_k)) \in R^{\mathcal{A}}$
- ▶ $\mathcal{A} \models_{\alpha} X(t)$ iff $\alpha_1(t) \in \alpha_2(X)$
- ▶ $\mathcal{A} \models_{\alpha} (\varphi \rightarrow \psi)$ iff $\mathcal{A} \not\models_{\alpha} \varphi$ or $\mathcal{A} \models_{\alpha} \psi$
- ▶ $\mathcal{A} \models_{\alpha} (\forall x)\varphi$ iff for every $a \in u(\mathcal{A})$, $\mathcal{A} \models_{\alpha[x \mapsto a]} \varphi$
- ▶ $\mathcal{A} \models_{\alpha} (\forall X)\varphi$ iff for every $S \subseteq u(\mathcal{A})$, $\mathcal{A} \models_{\alpha[X \mapsto S]} \varphi$

Examples

Recall the signature for the graph structure, $\tau = \{E\}$

- ▶ The graph is 3-colorable

Examples

Recall the signature for the graph structure, $\tau = \{E\}$

- ▶ The graph is 3-colorable

$$\exists X \exists Y \exists Z (\forall x [X(x) \vee Y(x) \vee Z(x)] \wedge$$

$$\forall x \forall y [E(x, y) \rightarrow \{\neg(X(x) \wedge X(y)) \wedge \neg(Y(x) \wedge Y(y)) \wedge \neg(Z(x) \wedge Z(y))\}])$$

Examples

Recall the signature for the graph structure, $\tau = \{E\}$

- ▶ The graph has an independent set of size $\geq k$

Examples

Recall the signature for the graph structure, $\tau = \{E\}$

- ▶ The graph has an independent set of size $\geq k$

$$\exists I \{ \forall x \forall y [(\neg(x = y) \wedge I(x) \wedge I(y)) \rightarrow \neg E(x, y)] \wedge$$

$$\exists x_1 \dots x_k [\bigwedge_{i \neq j} \neg(x_i = x_j) \wedge \bigwedge_i I(x_i)] \}$$

Examples

Recall the signature τ for the word structure, $\tau = \{Q_a, Q_b, <, S\}$ for $\Sigma = \{a, b\}$

- ▶ Words of even length

Examples

Recall the signature τ for the word structure, $\tau = \{Q_a, Q_b, <, S\}$ for $\Sigma = \{a, b\}$

- Words of even length

$$\exists E \exists O \{ \forall x [(first(x) \rightarrow E(x)) \wedge (last(x) \rightarrow O(x))] \}$$

Examples

Recall the signature τ for the word structure, $\tau = \{Q_a, Q_b, <, S\}$ for $\Sigma = \{a, b\}$

- Words of even length

$$\exists E \exists O \{ \forall x [(first(x) \rightarrow E(x)) \wedge (last(x) \rightarrow O(x))] \}$$

$$\wedge \forall x [(E(x) \vee O(x)) \wedge \neg(E(x) \wedge O(x))]$$

Examples

Recall the signature τ for the word structure, $\tau = \{Q_a, Q_b, <, S\}$ for $\Sigma = \{a, b\}$

- Words of even length

$$\exists E \exists O \{ \forall x [(first(x) \rightarrow E(x)) \wedge (last(x) \rightarrow O(x))] \}$$

$$\wedge \forall x [(E(x) \vee O(x)) \wedge \neg(E(x) \wedge O(x))]$$

$$\wedge \forall x \forall y [S(x, y) \wedge O(x) \rightarrow E(y)]$$

$$\wedge \forall x \forall y [S(x, y) \wedge E(x) \rightarrow O(y)] \}$$