

# **CS 228 : Logic in Computer Science**

Krishna. S

# More Rules

---

- ▶ Thanks to MT, we have  $p \rightarrow q, \neg q \vdash \neg p$ .

# More Rules

---

- ▶ Thanks to MT, we have  $p \rightarrow q, \neg q \vdash \neg p$ .
- ▶ Can we prove  $p \rightarrow q \vdash \neg q \rightarrow \neg p$ ?

# More Rules

---

- ▶ Thanks to MT, we have  $p \rightarrow q, \neg q \vdash \neg p$ .
- ▶ Can we prove  $p \rightarrow q \vdash \neg q \rightarrow \neg p$ ?
- ▶ So far, no proof rule that can do this.

# More Rules

---

- ▶ Thanks to MT, we have  $p \rightarrow q, \neg q \vdash \neg p$ .
- ▶ Can we prove  $p \rightarrow q \vdash \neg q \rightarrow \neg p$ ?
- ▶ So far, no proof rule that can do this.
- ▶ Given  $p \rightarrow q$ , let us assume  $\neg q$ . Can we then prove  $\neg p$ ?

# More Rules

---

- ▶ Thanks to MT, we have  $p \rightarrow q, \neg q \vdash \neg p$ .
- ▶ Can we prove  $p \rightarrow q \vdash \neg q \rightarrow \neg p$ ?
- ▶ So far, no proof rule that can do this.
- ▶ Given  $p \rightarrow q$ , let us assume  $\neg q$ . Can we then prove  $\neg p$ ?
- ▶ Yes, using MT.

# The implies introduction rule $\rightarrow i$

---

►  $p \rightarrow q \vdash \neg q \rightarrow \neg p$

1.  $p \rightarrow q$  premise

2.  $\neg q$  assumption

3.  $\neg p$  MT 1,2

4.  $\neg q \rightarrow \neg p$   $\rightarrow i$  2-3

## More on $\rightarrow$ *i*

---

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.        *true*

premise

2.



## More on $\rightarrow i$

---

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.		

## More on $\rightarrow i$

---

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.	$\neg q \rightarrow \neg p$	assumption
4.		

# More on $\rightarrow i$

---

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.	$\neg q \rightarrow \neg p$	assumption
4.	$p$	assumption
5.		

# More on $\rightarrow i$

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.	$\neg q \rightarrow \neg p$	assumption
4.	$p$	assumption
5.	$\neg\neg p$	$\neg\neg i$ 4
6.		

# More on $\rightarrow i$

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.	$\neg q \rightarrow \neg p$	assumption
4.	$p$	assumption
5.	$\neg\neg p$	$\neg\neg i$ 4
6.	$\neg\neg q$	MT 3,5
7.		

# More on $\rightarrow i$

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.	$\neg q \rightarrow \neg p$	assumption
4.	$p$	assumption
5.	$\neg\neg p$	$\neg\neg i$ 4
6.	$\neg\neg q$	MT 3,5
7.	$q$	$\neg\neg e$ 6
8.		

# More on $\rightarrow i$

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.	$\neg q \rightarrow \neg p$	assumption
4.	$p$	assumption
5.	$\neg\neg p$	$\neg\neg i$ 4
6.	$\neg\neg q$	MT 3,5
7.	$q$	$\neg\neg e$ 6
8.	$r$	MP 2,7

# More on $\rightarrow i$

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.	$\neg q \rightarrow \neg p$	assumption
4.	$p$	assumption
5.	$\neg\neg p$	$\neg\neg i$ 4
6.	$\neg\neg q$	MT 3,5
7.	$q$	$\neg\neg e$ 6
8.	$r$	MP 2,7
9.	$p \rightarrow r$	$\rightarrow i$ 4-8



# More on $\rightarrow i$

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.	$\neg q \rightarrow \neg p$	assumption
4.	$p$	assumption
5.	$\neg\neg p$	$\neg\neg i$ 4
6.	$\neg\neg q$	MT 3,5
7.	$q$	$\neg\neg e$ 6
8.	$r$	MP 2,7
9.	$p \rightarrow r$	$\rightarrow i$ 4-8
10.	$(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)$	$\rightarrow i$ 3-9
11.		

# More on $\rightarrow i$

►  $\vdash (q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$

1.	<i>true</i>	premise
2.	$q \rightarrow r$	assumption
3.	$\neg q \rightarrow \neg p$	assumption
4.	$p$	assumption
5.	$\neg\neg p$	$\neg\neg i$ 4
6.	$\neg\neg q$	MT 3,5
7.	$q$	$\neg\neg e$ 6
8.	$r$	MP 2,7
9.	$p \rightarrow r$	$\rightarrow i$ 4-8
10.	$(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)$	$\rightarrow i$ 3-9
11.	$(q \rightarrow r) \rightarrow [(\neg q \rightarrow \neg p) \rightarrow (p \rightarrow r)]$	$\rightarrow i$ 2-10

# Transforming Proofs

---

- ▶  $(q \rightarrow r), (\neg q \rightarrow \neg p), p \vdash r$
- ▶ Transform any proof  $\varphi_1, \dots, \varphi_n \vdash \psi$  to  $\vdash \varphi_1 \rightarrow (\varphi_2 \rightarrow \dots (\varphi_n \rightarrow \psi) \dots)$  by adding  $n$  lines of the rule  $\rightarrow i$

# More Examples

---

►  $p \rightarrow (q \rightarrow r) \vdash (p \wedge q) \rightarrow r$

1.  $p \rightarrow (q \rightarrow r)$  premise

2.

# More Examples

---

►  $p \rightarrow (q \rightarrow r) \vdash (p \wedge q) \rightarrow r$

1.  $p \rightarrow (q \rightarrow r)$  premise

2.  $p \wedge q$  assumption

3.

# More Examples

---

►  $p \rightarrow (q \rightarrow r) \vdash (p \wedge q) \rightarrow r$

1.  $p \rightarrow (q \rightarrow r)$  premise

2.  $p \wedge q$  assumption

3.  $p$   $\wedge e_1$  2

4.

# More Examples

---

►  $p \rightarrow (q \rightarrow r) \vdash (p \wedge q) \rightarrow r$

1.  $p \rightarrow (q \rightarrow r)$  premise

2.  $p \wedge q$  assumption

3.  $p$   $\wedge e_1$  2

4.  $q$   $\wedge e_2$  2

5.

# More Examples

---

►  $p \rightarrow (q \rightarrow r) \vdash (p \wedge q) \rightarrow r$

1.	$p \rightarrow (q \rightarrow r)$	premise
2.	$p \wedge q$	assumption
3.	$p$	$\wedge e_1$ 2
4.	$q$	$\wedge e_2$ 2
5.	$q \rightarrow r$	MP 1,3
6.		



# More Examples

---

►  $p \rightarrow (q \rightarrow r) \vdash (p \wedge q) \rightarrow r$

- |    |                                   |                |
|----|-----------------------------------|----------------|
| 1. | $p \rightarrow (q \rightarrow r)$ | premise        |
| 2. | $p \wedge q$                      | assumption     |
| 3. | $p$                               | $\wedge e_1$ 2 |
| 4. | $q$                               | $\wedge e_2$ 2 |
| 5. | $q \rightarrow r$                 | MP 1,3         |
| 6. | $r$                               | MP 4,5         |
| 7. |                                   |                |

# More Examples

---

►  $p \rightarrow (q \rightarrow r) \vdash (p \wedge q) \rightarrow r$

1.	$p \rightarrow (q \rightarrow r)$	premise
2.	$p \wedge q$	assumption
3.	$p$	$\wedge e_1$ 2
4.	$q$	$\wedge e_2$ 2
5.	$q \rightarrow r$	MP 1,3
6.	$r$	MP 4,5
7.	$p \wedge q \rightarrow r$	$\rightarrow i$ 2-6

# More Rules

---

The or introduction rule  $\vee i_1$

$$\frac{\varphi}{\varphi \vee \psi}$$

The or introduction rule  $\vee i_2$

$$\frac{\psi}{\varphi \vee \psi}$$

# More Rules

---

The or elimination rule  $\vee e$

$$\frac{\varphi \vee \psi \quad \varphi \vdash \chi \quad \psi \vdash \chi}{\chi}$$

# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.  $q \rightarrow r$  premise

2.

# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.	$q \rightarrow r$	premise
2.	$p \vee q$	assumption
3.		

# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.	$q \rightarrow r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.		

# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.	$q \rightarrow r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee r$	$\vee i_1$ 3
5.		



# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.	$q \rightarrow r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee r$	$\vee i_1$ 3
5.	$q$	assumption
6.		

# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.	$q \rightarrow r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee r$	$\vee i_1$ 3
5.	$q$	assumption
6.	$r$	MP 1,5
7.		

# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.	$q \rightarrow r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee r$	$\vee i_1$ 3
5.	$q$	assumption
6.	$r$	MP 1,5
7.	$p \vee r$	$\vee i_2$ 6

# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.	$q \rightarrow r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee r$	$\vee i_1$ 3
5.	$q$	assumption
6.	$r$	MP 1,5
7.	$p \vee r$	$\vee i_2$ 6
8.	$p \vee r$	$\vee e$ 2, 3-4, 5-7

# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.	$q \rightarrow r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee r$	$\vee i_1$ 3
5.	$q$	assumption
6.	$r$	MP 1,5
7.	$p \vee r$	$\vee i_2$ 6
8.	$p \vee r$	$\vee e$ 2, 3-4, 5-7
9.		

# Or Elimination Example

---

►  $q \rightarrow r \vdash (p \vee q) \rightarrow (p \vee r)$

1.	$q \rightarrow r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee r$	$\vee i_1$ 3
5.	$q$	assumption
6.	$r$	MP 1,5
7.	$p \vee r$	$\vee i_2$ 6
8.	$p \vee r$	$\vee e$ 2, 3-4, 5-7
9.	$(p \vee q) \rightarrow (p \vee r)$	$\rightarrow i$ 2-8

# Associativity Using Or Elimination

---

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.  $(p \vee q) \vee r$  premise

2.

# Associativity Using Or Elimination

---

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

- |    |                     |            |
|----|---------------------|------------|
| 1. | $(p \vee q) \vee r$ | premise    |
| 2. | $p \vee q$          | assumption |
| 3. |                     |            |



# Associativity Using Or Elimination

---

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.	$(p \vee q) \vee r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.		

# Associativity Using Or Elimination

---

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.  $(p \vee q) \vee r$  premise

2.  $p \vee q$  assumption

3.  $p$  assumption

4.  $p \vee (q \vee r)$   $\vee i_1$  3

5.

# Associativity Using Or Elimination

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.	$(p \vee q) \vee r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee (q \vee r)$	$\vee i_1$ 3
5.	$q$	assumption
6.		

# Associativity Using Or Elimination

---

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.	$(p \vee q) \vee r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee (q \vee r)$	$\vee i_1$ 3
5.	$q$	assumption
6.	$q \vee r$	$\vee i_1$ 5
7.		

# Associativity Using Or Elimination

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.  $(p \vee q) \vee r$  premise

2.  $p \vee q$  assumption

3.  $p$  assumption

4.  $p \vee (q \vee r)$   $\vee i_1$  3

5.  $q$  assumption

6.  $q \vee r$   $\vee i_1$  5

7.  $p \vee (q \vee r)$   $\vee i_2$  6

8.

# Associativity Using Or Elimination

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.  $(p \vee q) \vee r$  premise

2.  $p \vee q$  assumption

3.  $p$  assumption

4.  $p \vee (q \vee r)$   $\vee i_1$  3

5.  $q$  assumption

6.  $q \vee r$   $\vee i_1$  5

7.  $p \vee (q \vee r)$   $\vee i_2$  6

8.  $p \vee (q \vee r)$   $\vee e$  2, 3-4, 5-7

9.

# Associativity Using Or Elimination

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.	$(p \vee q) \vee r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee (q \vee r)$	$\vee i_1$ 3
5.	$q$	assumption
6.	$q \vee r$	$\vee i_1$ 5
7.	$p \vee (q \vee r)$	$\vee i_2$ 6
8.	$p \vee (q \vee r)$	$\vee e$ 2, 3-4, 5-7
9.	$r$	assumption
10.		

# Associativity Using Or Elimination

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.	$(p \vee q) \vee r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee (q \vee r)$	$\vee i_1$ 3
5.	$q$	assumption
6.	$q \vee r$	$\vee i_1$ 5
7.	$p \vee (q \vee r)$	$\vee i_2$ 6
8.	$p \vee (q \vee r)$	$\vee e$ 2, 3-4, 5-7
9.	$r$	assumption
10.	$q \vee r$	$\vee i_2$ 9
11.		



# Associativity Using Or Elimination

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.	$(p \vee q) \vee r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee (q \vee r)$	$\vee i_1$ 3
5.	$q$	assumption
6.	$q \vee r$	$\vee i_1$ 5
7.	$p \vee (q \vee r)$	$\vee i_2$ 6
8.	$p \vee (q \vee r)$	$\vee e$ 2, 3-4, 5-7
9.	$r$	assumption
10.	$q \vee r$	$\vee i_2$ 9
11.	$p \vee (q \vee r)$	$\vee i_2$ 10

# Associativity Using Or Elimination

►  $(p \vee q) \vee r \vdash p \vee (q \vee r)$

1.	$(p \vee q) \vee r$	premise
2.	$p \vee q$	assumption
3.	$p$	assumption
4.	$p \vee (q \vee r)$	$\vee i_1$ 3
5.	$q$	assumption
6.	$q \vee r$	$\vee i_1$ 5
7.	$p \vee (q \vee r)$	$\vee i_2$ 6
8.	$p \vee (q \vee r)$	$\vee e$ 2, 3-4, 5-7
9.	$r$	assumption
10.	$q \vee r$	$\vee i_2$ 9
11.	$p \vee (q \vee r)$	$\vee i_2$ 10
12.	$p \vee (q \vee r)$	$\vee e$ 1, 2-8, 9-11

# The Copy Rule

---

►  $\vdash p \rightarrow (q \rightarrow p)$

1.     *true*             premise

2.

# The Copy Rule

---

►  $\vdash p \rightarrow (q \rightarrow p)$

1.	<i>true</i>	premise
2.	<i>p</i>	assumption
3.		

# The Copy Rule

---

►  $\vdash p \rightarrow (q \rightarrow p)$

1.	<i>true</i>	premise
2.	$p$	assumption
3.	$q$	assumption
4.		

# The Copy Rule

---

►  $\vdash p \rightarrow (q \rightarrow p)$

1.	<i>true</i>	premise
2.	<i>p</i>	assumption
3.	<i>q</i>	assumption
4.	<i>p</i>	copy 2
5.		

# The Copy Rule

---

►  $\vdash p \rightarrow (q \rightarrow p)$

1.	<i>true</i>	premise
2.	<i>p</i>	assumption
3.	<i>q</i>	assumption
4.	<i>p</i>	copy 2
5.	$q \rightarrow p$	$\rightarrow i$ 3-4
6.		

# The Copy Rule

---

►  $\vdash p \rightarrow (q \rightarrow p)$

1.	<i>true</i>	premise
2.	$p$	assumption
3.	$q$	assumption
4.	$p$	copy 2
5.	$q \rightarrow p$	$\rightarrow i$ 3-4
6.	$p \rightarrow (q \rightarrow p)$	$\rightarrow i$ 2-5



# The Rules of Single Negtion

---

- ▶ We have seen  $\neg\neg e$  and  $\neg\neg i$ , the elimination and introduction of double negation.

# The Rules of Single Negtion

---

- ▶ We have seen  $\neg\neg e$  and  $\neg\neg i$ , the elimination and introduction of double negation.
- ▶ How about introducing and eliminating single negations?

# The Rules of Single Negtion

---

- ▶ We have seen  $\neg\neg e$  and  $\neg\neg i$ , the elimination and introduction of double negation.
- ▶ How about introducing and eliminating single negations?
- ▶ We use the notion of contradictions, an expression of the form  $\varphi \wedge \neg\varphi$ , where  $\varphi$  is any propositional logic formula.

# The Rules of Single Negtion

---

- ▶ We have seen  $\neg\neg e$  and  $\neg\neg i$ , the elimination and introduction of double negation.
- ▶ How about introducing and eliminating single negations?
- ▶ We use the notion of contradictions, an expression of the form  $\varphi \wedge \neg\varphi$ , where  $\varphi$  is any propositional logic formula.
- ▶ Any two contradictions are equivalent :  $p \wedge \neg p$  is equivalent to  $\neg r \wedge r$ .

# The Rules of Single Negation

---

- ▶ We have seen  $\neg\neg e$  and  $\neg\neg i$ , the elimination and introduction of double negation.
- ▶ How about introducing and eliminating single negations?
- ▶ We use the notion of contradictions, an expression of the form  $\varphi \wedge \neg\varphi$ , where  $\varphi$  is any propositional logic formula.
- ▶ Any two contradictions are equivalent :  $p \wedge \neg p$  is equivalent to  $\neg r \wedge r$ .
- ▶  $\perp \rightarrow \varphi$  for any formula  $\varphi$ .

# Rules with $\perp$

---

The  $\perp$  elimination rule  $\perp e$

$$\frac{\perp}{\psi}$$

The  $\perp$  introduction rule  $\perp i$

$$\frac{\varphi \quad \neg\varphi}{\perp}$$

# An Example

---

►  $\neg p \vee q \vdash p \rightarrow q$

1.  $\neg p \vee q$  premise

2.

# An Example

---

►  $\neg p \vee q \vdash p \rightarrow q$

1.  $\neg p \vee q$  premise

2.  $\neg p$  premise

3.



# An Example

---

►  $\neg p \vee q \vdash p \rightarrow q$

1.  $\neg p \vee q$  premise

2.  $\neg p$  premise

3.  $p$  assumption

4.

# An Example

---

►  $\neg p \vee q \vdash p \rightarrow q$

1.  $\neg p \vee q$  premise

2.  $\neg p$  premise

3.  $p$  assumption

4.  $\perp$   $\perp i$  2,3

5.

# An Example

---

►  $\neg p \vee q \vdash p \rightarrow q$

1.  $\neg p \vee q$  premise

2.  $\neg p$  premise

3.  $p$  assumption

4.  $\perp$   $\perp i$  2,3

5.  $q$   $\perp e$  4

6.

# An Example

---

►  $\neg p \vee q \vdash p \rightarrow q$

1.	$\neg p \vee q$	premise
2.	$\neg p$	premise
3.	$p$	assumption
4.	$\perp$	$\perp i$ 2,3
5.	$q$	$\perp e$ 4
6.	$p \rightarrow q$	$\rightarrow i$ 3-5
7.	$q$	premise
8.		

# An Example

►  $\neg p \vee q \vdash p \rightarrow q$

1.	$\neg p \vee q$	premise
2.	$\neg p$	premise
3.	$p$	assumption
4.	$\perp$	$\perp i$ 2,3
5.	$q$	$\perp e$ 4
6.	$p \rightarrow q$	$\rightarrow i$ 3-5
7.	$q$	premise
8.	$p$	assumption
9.		

# An Example

►  $\neg p \vee q \vdash p \rightarrow q$

1.	$\neg p \vee q$	premise
2.	$\neg p$	premise
3.	$p$	assumption
4.	$\perp$	$\perp i$ 2,3
5.	$q$	$\perp e$ 4
6.	$p \rightarrow q$	$\rightarrow i$ 3-5
7.	$q$	premise
8.	$p$	assumption
9.	$q$	copy 7
10.	$p \rightarrow q$	$\rightarrow i$ 8-9
11.	$p \rightarrow q$	$\vee e$ 1, 2-6, 7-10

# Introducing Negations

---

- ▶ In the course of a proof, if you assume  $\varphi$  (by opening a box) and obtain  $\perp$  in the box, then we conclude  $\neg\varphi$
- ▶ This rule is denoted  $\neg i$  and is read as  $\neg$  introduction.

# An Example

---

►  $p \rightarrow \neg p \vdash \neg p$

1.  $p \rightarrow \neg p$  premise

2.



# An Example

---

►  $p \rightarrow \neg p \vdash \neg p$

- |    |                        |            |
|----|------------------------|------------|
| 1. | $p \rightarrow \neg p$ | premise    |
| 2. | $p$                    | assumption |
| 3. |                        |            |

# An Example

---

►  $p \rightarrow \neg p \vdash \neg p$

1.	$p \rightarrow \neg p$	premise
2.	$p$	assumption
3.	$\neg p$	MP 1,2
4.		

# An Example

---

►  $p \rightarrow \neg p \vdash \neg p$

1.  $p \rightarrow \neg p$  premise

2.  $p$  assumption

3.  $\neg p$  MP 1,2

4.  $\perp$   $\perp i$  2,3

5.  $\neg p$   $\neg i$  2-4

# Natural Deduction : Summary

---

- ▶ Given a sequent  $\varphi \vdash \psi$ , start writing all the premises on the top (think of  $\varphi$  as a conjunction of several formulae)

# Natural Deduction : Summary

---

- ▶ Given a sequent  $\varphi \vdash \psi$ , start writing all the premises on the top (think of  $\varphi$  as a conjunction of several formulae)
- ▶ If  $\psi$  has no  $\rightarrow$  in it, then use the premises, and the rules of natural deduction and arrive at  $\psi$ , or

# Natural Deduction : Summary

---

- ▶ Given a sequent  $\varphi \vdash \psi$ , start writing all the premises on the top (think of  $\varphi$  as a conjunction of several formulae)
- ▶ If  $\psi$  has no  $\rightarrow$  in it, then use the premises, and the rules of natural deduction and arrive at  $\psi$ , or
- ▶ Assume (opening a box)  $\neg\psi$ , then obtain using the premises  $\perp$ . Infer henceforth that  $\psi$  holds.

# Natural Deduction : Summary

---

- ▶ Given a sequent  $\varphi \vdash \psi$ , start writing all the premises on the top (think of  $\varphi$  as a conjunction of several formulae)
- ▶ If  $\psi$  has no  $\rightarrow$  in it, then use the premises, and the rules of natural deduction and arrive at  $\psi$ , or
- ▶ Assume (opening a box)  $\neg\psi$ , then obtain using the premises  $\perp$ . Infer henceforth that  $\psi$  holds.
- ▶ If  $\psi$  is of the form  $\eta \rightarrow \kappa$ , then open a box assuming  $\eta$ , use the proof rules, as well as the premises and obtain  $\kappa$ .

# Natural Deduction : Summary

---

- ▶ Given a sequent  $\varphi \vdash \psi$ , start writing all the premises on the top (think of  $\varphi$  as a conjunction of several formulae)
- ▶ If  $\psi$  has no  $\rightarrow$  in it, then use the premises, and the rules of natural deduction and arrive at  $\psi$ , or
- ▶ Assume (opening a box)  $\neg\psi$ , then obtain using the premises  $\perp$ . Infer henceforth that  $\psi$  holds.
- ▶ If  $\psi$  is of the form  $\eta \rightarrow \kappa$ , then open a box assuming  $\eta$ , use the proof rules, as well as the premises and obtain  $\kappa$ .
- ▶ If  $\varphi$  is a disjunction of the form  $\psi_1 \vee \psi_2$ , then show that  $\psi_1 \vdash \psi$  and  $\psi_2 \vdash \psi$



# The Proofs So Far

---

- ▶ So far, the “proof” we have seen is purely syntactic, no true/false meanings were attached

# The Proofs So Far

---

- ▶ So far, the “proof” we have seen is purely syntactic, no true/false meanings were attached
- ▶ Intuitively,  $p \rightarrow q \vdash \neg p \vee q$  makes sense because you think semantically. However, we never used any semantics so far.

# The Proofs So Far

---

- ▶ So far, the “proof” we have seen is purely syntactic, no true/false meanings were attached
- ▶ Intuitively,  $p \rightarrow q \vdash \neg p \vee q$  makes sense because you think semantically. However, we never used any semantics so far.
- ▶ Now we show that whatever can be proved makes sense semantically too.

# Semantics

---

- ▶ Each propositional variable is assigned values true/false. Truth tables for each of the operators  $\vee, \wedge, \neg, \rightarrow$  to determine truth values of complex formulae.

# Semantics

---

- ▶ Each propositional variable is assigned values true/false. Truth tables for each of the operators  $\vee, \wedge, \neg, \rightarrow$  to determine truth values of complex formulae.
- ▶  $\varphi_1, \dots, \varphi_n \models \psi$  iff whenever  $\varphi_1, \dots, \varphi_n$  evaluate to true, so does  $\psi$ .  $\models$  is read as **semantically entails**

# Semantics

---

- ▶ Each propositional variable is assigned values true/false. Truth tables for each of the operators  $\vee, \wedge, \neg, \rightarrow$  to determine truth values of complex formulae.
- ▶  $\varphi_1, \dots, \varphi_n \models \psi$  iff whenever  $\varphi_1, \dots, \varphi_n$  evaluate to true, so does  $\psi$ .  $\models$  is read as **semantically entails**
- ▶ Two formulae  $\varphi$  and  $\psi$  are **provably equivalent** iff  $\varphi \vdash \psi$  and  $\psi \vdash \varphi$

# Semantics

---

- ▶ Each propositional variable is assigned values true/false. Truth tables for each of the operators  $\vee, \wedge, \neg, \rightarrow$  to determine truth values of complex formulae.
- ▶  $\varphi_1, \dots, \varphi_n \models \psi$  iff whenever  $\varphi_1, \dots, \varphi_n$  evaluate to true, so does  $\psi$ .  $\models$  is read as **semantically entails**
- ▶ Two formulae  $\varphi$  and  $\psi$  are **provably equivalent** iff  $\varphi \vdash \psi$  and  $\psi \vdash \varphi$
- ▶ Two formulae  $\varphi$  and  $\psi$  are **semantically equivalent** iff  $\varphi \models \psi$  and  $\psi \models \varphi$

# Soundness of Propositional Logic

---

$$\varphi_1, \dots, \varphi_n \vdash \psi \Rightarrow \varphi_1, \dots, \varphi_n \models \psi$$

Whenever  $\psi$  can be proved from  $\varphi_1, \dots, \varphi_n$ , then  $\psi$  evaluates to true whenever  $\varphi_1, \dots, \varphi_n$  evaluate to true



# Soundness

---

- ▶ Assume  $\varphi_1, \dots, \varphi_n \vdash \psi$ .

# Soundness

---

- ▶ Assume  $\varphi_1, \dots, \varphi_n \vdash \psi$ .
- ▶ There is some proof (of length  $k$  lines) that yields  $\psi$ . Induct on  $k$ .

# Soundness

---

- ▶ Assume  $\varphi_1, \dots, \varphi_n \vdash \psi$ .
- ▶ There is some proof (of length  $k$  lines) that yields  $\psi$ . Induct on  $k$ .
- ▶ When  $k = 1$ , there is only one line in the proof, say  $\varphi$ , which is the premise. Then we have  $\varphi \vdash \varphi$ , since  $\varphi$  is given. But then we also have  $\varphi \models \varphi$ .

# Soundness

---

- ▶ Assume  $\varphi_1, \dots, \varphi_n \vdash \psi$ .
- ▶ There is some proof (of length  $k$  lines) that yields  $\psi$ . Induct on  $k$ .
- ▶ When  $k = 1$ , there is only one line in the proof, say  $\varphi$ , which is the premise. Then we have  $\varphi \vdash \varphi$ , since  $\varphi$  is given. But then we also have  $\varphi \models \varphi$ .
- ▶ Assume that whenever  $\varphi_1, \dots, \varphi_n \vdash \psi$  using a proof of length  $\leq k - 1$ , we have  $\varphi_1, \dots, \varphi_n \models \psi$ .

# Soundness

---

- ▶ Assume  $\varphi_1, \dots, \varphi_n \vdash \psi$ .
- ▶ There is some proof (of length  $k$  lines) that yields  $\psi$ . Induct on  $k$ .
- ▶ When  $k = 1$ , there is only one line in the proof, say  $\varphi$ , which is the premise. Then we have  $\varphi \vdash \varphi$ , since  $\varphi$  is given. But then we also have  $\varphi \models \varphi$ .
- ▶ Assume that whenever  $\varphi_1, \dots, \varphi_n \vdash \psi$  using a proof of length  $\leq k - 1$ , we have  $\varphi_1, \dots, \varphi_n \models \psi$ .
- ▶ Consider now a proof with  $k$  lines.

# Soundness : Case $\wedge i$

---

- ▶ How did we arrive at  $\psi$ ? Which proof rule gave  $\psi$  as the last line?

# Soundness : Case $\wedge i$

---

- ▶ How did we arrive at  $\psi$ ? Which proof rule gave  $\psi$  as the last line?
- ▶ Assume  $\psi$  was obtained using  $\wedge i$ . Then  $\psi$  is of the form  $\psi_1 \wedge \psi_2$ .

# Soundness : Case $\wedge i$

---

- ▶ How did we arrive at  $\psi$ ? Which proof rule gave  $\psi$  as the last line?
- ▶ Assume  $\psi$  was obtained using  $\wedge i$ . Then  $\psi$  is of the form  $\psi_1 \wedge \psi_2$ .
- ▶  $\psi_1$  and  $\psi_2$  were proved earlier, say in lines  $k_1, k_2 < k$ .



# Soundness : Case $\wedge i$

---

- ▶ How did we arrive at  $\psi$ ? Which proof rule gave  $\psi$  as the last line?
- ▶ Assume  $\psi$  was obtained using  $\wedge i$ . Then  $\psi$  is of the form  $\psi_1 \wedge \psi_2$ .
- ▶  $\psi_1$  and  $\psi_2$  were proved earlier, say in lines  $k_1, k_2 < k$ .
- ▶ We have the shorter proofs  $\varphi_1, \dots, \varphi_n \vdash \psi_1$  and  $\varphi_1, \dots, \varphi_n \vdash \psi_2$

# Soundness : Case $\wedge i$

---

- ▶ How did we arrive at  $\psi$ ? Which proof rule gave  $\psi$  as the last line?
- ▶ Assume  $\psi$  was obtained using  $\wedge i$ . Then  $\psi$  is of the form  $\psi_1 \wedge \psi_2$ .
- ▶  $\psi_1$  and  $\psi_2$  were proved earlier, say in lines  $k_1, k_2 < k$ .
- ▶ We have the shorter proofs  $\varphi_1, \dots, \varphi_n \vdash \psi_1$  and  $\varphi_1, \dots, \varphi_n \vdash \psi_2$
- ▶ By inductive hypothesis, we have  $\varphi_1, \dots, \varphi_n \models \psi_1$  and  $\varphi_1, \dots, \varphi_n \models \psi_2$ . By semantics, we have  $\varphi_1, \dots, \varphi_n \models \psi_1 \wedge \psi_2$ .

# Soundness : Case $\rightarrow i$

---

- ▶ Assume  $\psi$  was obtained using  $\rightarrow i$ . Then  $\psi$  is of the form  $\psi_1 \rightarrow \psi_2$ .

# Soundness : Case $\rightarrow i$

---

- ▶ Assume  $\psi$  was obtained using  $\rightarrow i$ . Then  $\psi$  is of the form  $\psi_1 \rightarrow \psi_2$ .
- ▶ A box starting with  $\psi_1$  was opened at some line  $k_1 < k$ .

# Soundness : Case $\rightarrow i$

---

- ▶ Assume  $\psi$  was obtained using  $\rightarrow i$ . Then  $\psi$  is of the form  $\psi_1 \rightarrow \psi_2$ .
- ▶ A box starting with  $\psi_1$  was opened at some line  $k_1 < k$ .
- ▶ The last line in the box was  $\psi_2$ .

# Soundness : Case $\rightarrow i$

---

- ▶ Assume  $\psi$  was obtained using  $\rightarrow i$ . Then  $\psi$  is of the form  $\psi_1 \rightarrow \psi_2$ .
- ▶ A box starting with  $\psi_1$  was opened at some line  $k_1 < k$ .
- ▶ The last line in the box was  $\psi_2$ .
- ▶ The line just after the box was  $\psi$ .

# Soundness : Case $\rightarrow i$

---

- ▶ Assume  $\psi$  was obtained using  $\rightarrow i$ . Then  $\psi$  is of the form  $\psi_1 \rightarrow \psi_2$ .
- ▶ A box starting with  $\psi_1$  was opened at some line  $k_1 < k$ .
- ▶ The last line in the box was  $\psi_2$ .
- ▶ The line just after the box was  $\psi$ .
- ▶ Consider adding  $\psi_1$  in the premises along with  $\varphi_1, \dots, \varphi_n$ . Then we will get a proof  $\varphi_1, \dots, \varphi_n, \psi_1 \vdash \psi_2$ , of length  $k - 1$ . By inductive hypothesis,  $\varphi_1, \dots, \varphi_n, \psi_1 \models \psi_2$ . By semantics, this is same as  $\varphi_1, \dots, \varphi_n \models \psi_1 \rightarrow \psi_2$

# Soundness : Case $\rightarrow i$

---

- ▶ Assume  $\psi$  was obtained using  $\rightarrow i$ . Then  $\psi$  is of the form  $\psi_1 \rightarrow \psi_2$ .
- ▶ A box starting with  $\psi_1$  was opened at some line  $k_1 < k$ .
- ▶ The last line in the box was  $\psi_2$ .
- ▶ The line just after the box was  $\psi$ .
- ▶ Consider adding  $\psi_1$  in the premises along with  $\varphi_1, \dots, \varphi_n$ . Then we will get a proof  $\varphi_1, \dots, \varphi_n, \psi_1 \vdash \psi_2$ , of length  $k - 1$ . By inductive hypothesis,  $\varphi_1, \dots, \varphi_n, \psi_1 \models \psi_2$ . By semantics, this is same as  $\varphi_1, \dots, \varphi_n \models \psi_1 \rightarrow \psi_2$
- ▶ The equivalence of  $\varphi_1, \dots, \varphi_n \vdash \psi_1 \rightarrow \psi_2$  and  $\varphi_1, \dots, \varphi_n, \psi_1 \vdash \psi_2$  gives the proof.