

A decorative blue crosshair consisting of a vertical line and a horizontal line intersecting in the upper-left quadrant of the slide.

CS 228 : Logic in Computer Science

Krishna. S



So Far

- ▶ ω -automata with Büchi acceptance, also called Büchi automata
- ▶ Non-determinism versus determinism
- ▶ Closure under union, intersection




Büchi Acceptance

For Büchi Acceptance, Acc is specified as a set of states, $G \subseteq Q$. The ω -word α is accepted if there is a run ρ of α such that $Inf(\rho) \cap G \neq \emptyset$.




Union and Intersection of NBA






Union and Intersection of NBA






Union and Intersection of NBA






Union and Intersection of NBA






Union and Intersection of NBA






Union and Intersection of NBA




Union and Intersection of NBA



Union and Intersection of NBA




Union and Intersection of NBA




- States as $Q_1 \times Q_2 \times \{1, 2\}$, start state $(q_0, q'_0, 1)$

Union and Intersection of NBA




- ▶ States as $Q_1 \times Q_2 \times \{1, 2\}$, start state $(q_0, q'_0, 1)$
- ▶ $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 1)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_1 \notin G_1$
- ▶ $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 2)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_1 \in G_1$

Union and Intersection of NBA



- ▶ States as $Q_1 \times Q_2 \times \{1, 2\}$, start state $(q_0, q'_0, 1)$
- ▶ $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 1)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_1 \notin G_1$
- ▶ $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 2)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_1 \in G_1$
- ▶ $(q_1, q_2, 2) \xrightarrow{a} (q'_1, q'_2, 2)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_2 \notin G_2$
- ▶ $(q_1, q_2, 2) \xrightarrow{a} (q'_1, q'_2, 1)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_2 \in G_2$

Union and Intersection of NBA




- ▶ States as $Q_1 \times Q_2 \times \{1, 2\}$, start state $(q_0, q'_0, 1)$
- ▶ $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 1)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_1 \notin G_1$
- ▶ $(q_1, q_2, 1) \xrightarrow{a} (q'_1, q'_2, 2)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_1 \in G_1$
- ▶ $(q_1, q_2, 2) \xrightarrow{a} (q'_1, q'_2, 2)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_2 \notin G_2$
- ▶ $(q_1, q_2, 2) \xrightarrow{a} (q'_1, q'_2, 1)$ if $q_1 \xrightarrow{a} q'_1$ and $q_2 \xrightarrow{a} q'_2$ and $q_2 \in G_2$
- ▶ Good states = $Q_1 \times G_2 \times \{2\}$ or $G_1 \times Q_2 \times \{1\}$

Emptiness



Given an NBA/DBA \mathcal{A} , how do you check if $L(\mathcal{A}) = \emptyset$?

- ▶ Enumerate SCCs
- ▶ Check if there is an SCC containing a good state



Complementation of DBA





Complementation of DBA



Complementation of DBA



Complementation of DBA



Complementation of DBA

- ▶ Given \mathcal{A} is a DBA, and $w \notin L(\mathcal{A})$, then after some finite prefix, the unique run of w settles in bad states.
- ▶ Idea for complement: “copy” states of $Q - G$, once you enter this block, you stay there.
- ▶ View this as the set of good states, any word w that was rejected by \mathcal{A} has two possible runs in this automaton: the original run, and one another, that will settle in the $Q - G$ copy, and will be accepted.
- ▶ What we get now is an NBA for $\overline{L(\mathcal{A})}$, not a DBA.

Complementing NBA non-trivial, can be done.

Normal Form for ω -regular languages

An ω -regular language $L \subseteq \Sigma^\omega$ can be written as $L = \bigcup_{i=1}^n U_i V_i^\omega$, where U_i, V_i are regular languages.

One direction : Assume L is accepted by an NBA/DBA.



- ▶ Define $U_g = \{w \in \Sigma^* \mid q_0 \xrightarrow{w} g\}$
- ▶ Define $V_g = \{w \in \Sigma^* \mid g \xrightarrow{w} g\}$
- ▶ Then $L = \bigcup_{g \in G} U_g V_g^\omega$, where U_g, V_g are regular
- ▶ Show that U_g, V_g are regular.

Normal Form for ω -regular languages

An ω -regular language $L \subseteq \Sigma^\omega$ can be written as $L = \bigcup_{i=1}^n U_i V_i^\omega$, where U_i, V_i are regular languages.

Other direction : Assume $L = \bigcup_{i=1}^n U_i V_i^\omega$. Show that L is accepted by an NBA/DBA.

1. If V is regular, V^ω is ω -regular





Normal Form for ω -regular languages

An ω -regular language $L \subseteq \Sigma^\omega$ can be written as $L = \bigcup_{i=1}^n U_i V_i^\omega$, where U_i, V_i are regular languages.

Other direction : Assume $L = \bigcup_{i=1}^n U_i V_i^\omega$. Show that L is accepted by an NBA/DBA.

1. If V is regular, V^ω is ω -regular





Normal Form for ω -regular languages

An ω -regular language $L \subseteq \Sigma^\omega$ can be written as $L = \bigcup_{i=1}^n U_i V_i^\omega$, where U_i, V_i are regular languages.

Other direction : Assume $L = \bigcup_{i=1}^n U_i V_i^\omega$. Show that L is accepted by an NBA/DBA.

1. If V is regular, V^ω is ω -regular





Normal Form for ω -regular languages

An ω -regular language $L \subseteq \Sigma^\omega$ can be written as $L = \bigcup_{i=1}^n U_i V_i^\omega$, where U_i, V_i are regular languages.

Other direction : Assume $L = \bigcup_{i=1}^n U_i V_i^\omega$. Show that L is accepted by an NBA/DBA.

1. If V is regular, V^ω is ω -regular



Normal Form for ω -regular languages

1. If V is regular, V^ω is ω -regular
 - ▶ Let $D = (Q, \Sigma, q_0, \delta, F)$ be a DFA accepting V
 - ▶ Construct NBA $E = (Q \cup \{p_0\}, \Sigma, p_0, \Delta, G)$ such that $G = \{p_0\}$,
 - ▶ $\Delta = \delta \cup \{p_0 \in \Delta(q, a) \mid \delta(q, a) \in F\} \cup \{\Delta(p_0, a) = s \mid \delta(q_0, a) = s\}$
2. Show that if U is regular and V^ω is ω -regular, then UV^ω is ω -regular
 - ▶ $D = (Q_1, \Sigma, q_0, \delta_1, F)$ be a DFA, $L(D) = U$ and $E = (Q_2, \Sigma, q'_0, \delta_2, G)$ be an NBA, $L(E) = V^\omega$.
 - ▶ $A = (Q_1 \cup Q_2, \Sigma, q_0, \delta', G)$ NBA such that $\delta' = \delta_1 \cup \delta_2 \cup \{(q, a, q'_0) \mid \delta_1(q, a) \in F\}$