

# CS310 Automata Theory – 2016-2017

Nutan Limaye

Indian Institute of Technology, Bombay

[nutan@cse.iitb.ac.in](mailto:nutan@cse.iitb.ac.in)

Lecture 25: Turing machines, computability

March 20, 2017

# Module III so far...

## Introduction to Turing machines

What are Turing machines? Informal and formal definitions and examples.

# Module III so far...

## Introduction to Turing machines

What are Turing machines? Informal and formal definitions and examples.

Configurations of a Turing machine.

# Module III so far...

## Introduction to Turing machines

What are Turing machines? Informal and formal definitions and examples.

Configurations of a Turing machine.

Turing recognizable and Turing decidable languages.

# Module III so far...

## Introduction to Turing machines

What are Turing machines? Informal and formal definitions and examples.

Configurations of a Turing machine.

Turing recognizable and Turing decidable languages.

$k$ -tape TMs equivalent to 1-tape TMs.

Existence of unrecognizable languages.

$A_{TM}$  is recognizable but not decidable.

# A decision problem about TMs

$$A_{TM} = \{(M, w) \mid M \text{ accepts } w\}$$

# A decision problem about TMs

$$A_{TM} = \{(M, w) \mid M \text{ accepts } w\}$$

## Lemma

*$A_{TM}$  is Turing recognizable.*

## Proof sketch

Design a TM, say  $N$  such that,

$N$  behaves like  $M$  on  $w$  at each step,

if  $M$  reaches  $q_{acc}$  then  $N$  also accepts.



Is  $A_{TM}$  decidable?

# A decision problem about TMs

## Lemma

$A_{TM}$  is not Turing decidable.

Assume that there exists  $M$  such that  $M$  decides  $A_{TM}$ .






# A decision problem about TMs


## Lemma

$A_{TM}$  is not Turing decidable.

Assume that there exists  $M$  such that  $M$  decides  $A_{TM}$ .




What happens if we give  $D$  as input to itself?



# A decision problem about TMs

## Lemma

$A_{TM}$  is not Turing decidable.



If  $D$  accepts  $\langle D \rangle$  then  $D$  rejects  $\langle D \rangle$ .

If  $D$  rejects  $\langle D \rangle$  then  $D$  accepts  $\langle D \rangle$ . ☹️

# Diagonalization inside the proof

Behaviour of the machines.

	$\langle M_1 \rangle$	$\langle M_2 \rangle$	$\langle M_3 \rangle$	.....	.....
$M_1$	✓		✓	✓...	.....

# Diagonalization inside the proof

Behaviour of the machines.

	$\langle M_1 \rangle$	$\langle M_2 \rangle$	$\langle M_3 \rangle$	.....	.....
$M_1$	✓		✓	✓ ...	.....
$M_2$	✓	×		×	... ✓ ... × ✓ ...

# Diagonalization inside the proof

Behaviour of the machines.

	$\langle M_1 \rangle$	$\langle M_2 \rangle$	$\langle M_3 \rangle$	.....	.....
$M_1$	✓		✓	✓ ...	.....
$M_2$	✓	×		×	... ✓ ... × ✓ ...
$M_3$	×	×	✓	... ×	✓ .....
⋮					
⋮					

# Diagonalization inside the proof

Behaviour of  $H$ .

	$\langle M_1 \rangle$	$\langle M_2 \rangle$	$\langle M_3 \rangle$	.....	.....
$M_1$	✓	×	✓	✓ ...	.....
$M_2$	✓	×	×	×	... ✓ ... × ✓ ...
$M_3$	×	×	✓	... ×	✓ .....
⋮					
⋮					

# Diagonalization inside the proof

Behaviour of  $H$ .

	$\langle M_1 \rangle$	$\langle M_2 \rangle$	$\langle M_3 \rangle$	.....	.....
$M_1$	✓	×	✓	✓ ...	.....
$M_2$	✓	×	×	×	... ✓ ... × ✓ ...
$M_3$	×	×	✓	... ×	✓ .....
⋮					
⋮					

# Diagonalization inside the proof

Behaviour of  $D$ .

	$\langle M_1 \rangle$	$\langle M_2 \rangle$	$\langle M_3 \rangle$	.....	.....
$M_1$	<del>✓</del> / <del>✓</del> / <del>✓</del> ×	×	✓	✓ ...	.....
$M_2$	✓	<del>×</del> ✓	×	×	... ✓ ... × ✓ ...
$M_3$	×	×	<del>✓</del> / <del>✓</del> / <del>✓</del> ×	... ×	✓ .....
⋮					
⋮					



# Diagonalization inside the proof

Behaviour of  $D$  on itself.

	$\langle M_1 \rangle$	$\langle M_2 \rangle$	$\langle M_3 \rangle$	$\dots \langle D \rangle \dots$	$\dots$
$M_1$	<del>✓</del> ×	×	✓	✓ ...	$\dots$
$M_2$	✓	<del>×</del> ✓	×	×	✓ ... × ✓ ...
$M_3$	×	×	<del>✓</del> ×	... ×	✓ $\dots$
$\vdots$					
$\vdots$					
$D$				... ? ...	$\dots$

# Other undecidable problems and reducibility

## Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

## Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

# Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

Assume that Halt is decidable.

## Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

Assume that Halt is decidable. Let  $\mathcal{H}$  be the TM deciding Halt.

## Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

Assume that Halt is decidable. Let  $\mathcal{H}$  be the TM deciding Halt.

$\mathcal{A}$ : Run  $\mathcal{H}$  on  $(M, w)$ .

## Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

Assume that Halt is decidable. Let  $\mathcal{H}$  be the TM deciding Halt.

$\mathcal{A}$ : Run  $\mathcal{H}$  on  $(M, w)$ . If it rejects then reject,



## Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

Assume that Halt is decidable. Let  $\mathcal{H}$  be the TM deciding Halt.

$\mathcal{A}$ : Run  $\mathcal{H}$  on  $(M, w)$ . If it rejects then reject, else do as per  $M$  on  $w$ .

## Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

Assume that Halt is decidable. Let  $\mathcal{H}$  be the TM deciding Halt.

$\mathcal{A}$ : Run  $\mathcal{H}$  on  $(M, w)$ . If it rejects then reject, else do as per  $M$  on  $w$ .

$\mathcal{A}$  accepts  $(M, w)$  if  $M$  accepts  $w$

## Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

Assume that Halt is decidable. Let  $\mathcal{H}$  be the TM deciding Halt.

$\mathcal{A}$ : Run  $\mathcal{H}$  on  $(M, w)$ . If it rejects then reject, else do as per  $M$  on  $w$ .

$\mathcal{A}$  accepts  $(M, w)$  if  $M$  accepts  $w$  and rejects it if either  $M$  rejects  $w$

# Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

Assume that Halt is decidable. Let  $\mathcal{H}$  be the TM deciding Halt.

$\mathcal{A}$ : Run  $\mathcal{H}$  on  $(M, w)$ . If it rejects then reject, else do as per  $M$  on  $w$ .

$\mathcal{A}$  accepts  $(M, w)$  if  $M$  accepts  $w$  and rejects it if either  $M$  rejects  $w$  or  $M$  loops forever on  $w$ .

# Other undecidable problems and reducibility

Reducing  $A_{TM}$  to another problem to prove undecidability.

$$\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$$

We would like to show that Halt is undecidable.

Assume that Halt is decidable. Let  $\mathcal{H}$  be the TM deciding Halt.

$\mathcal{A}$ : Run  $\mathcal{H}$  on  $(M, w)$ . If it rejects then reject, else do as per  $M$  on  $w$ .

$\mathcal{A}$  accepts  $(M, w)$  if  $M$  accepts  $w$  and rejects it if either  $M$  rejects  $w$  or  $M$  loops forever on  $w$ .

$\mathcal{H}$  decides Halt if and only if  $\mathcal{A}$  decides  $A_{TM}$ .

# The halting problem

## Lemma

*The halting problem,  $\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$ , is undecidable.*

# The halting problem

## Lemma

*The halting problem,  $\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$ , is undecidable.*


Another way to describe the same proof.

# The halting problem

## Lemma

*The halting problem,  $\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$ , is undecidable.*

Another way to describe the same proof.






# The halting problem

## Lemma

*The halting problem,  $\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$ , is undecidable.*

Another way to describe the same proof.




# The halting problem

## Lemma

*The halting problem,  $\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$ , is undecidable.*

Another way to describe the same proof.




If Halt is decidable then  $\mathcal{A}$  decides  $A_{TM}$

# The halting problem

## Lemma

*The halting problem,  $\text{Halt} = \{(M, w) \mid M \text{ halts on } w\}$ , is undecidable.*

Another way to describe the same proof.



If Halt is decidable then  $\mathcal{A}$  decides  $A_{TM}$ , which is a contradiction.

# Emptiness problem for TM

## Lemma

*The emptiness problem for TMs,  $E_{TM} = \{\langle M \rangle \mid L(M) = \emptyset\}$ , is undecidable.*

Assume for the sake of contradiction that it is decidable.

# Emptiness problem for TM

## Lemma

*The emptiness problem for TMs,  $E_{TM} = \{\langle M \rangle \mid L(M) = \emptyset\}$ , is undecidable.*

Assume for the sake of contradiction that it is decidable. Let  $T$  be a machine that decides  $E_{TM}$ .

# Emptiness problem for TM

## Lemma

*The emptiness problem for TMs,  $E_{TM} = \{\langle M \rangle \mid L(M) = \emptyset\}$ , is undecidable.*

Assume for the sake of contradiction that it is decidable. Let  $T$  be a machine that decides  $E_{TM}$ .

Let  $T'_{M,w}$  be as follows:

# Emptiness problem for TM

## Lemma

*The emptiness problem for TMs,  $E_{TM} = \{\langle M \rangle \mid L(M) = \emptyset\}$ , is undecidable.*

Assume for the sake of contradiction that it is decidable. Let  $T$  be a machine that decides  $E_{TM}$ .

Let  $T'_{M,w}$  be as follows:

On input  $x$

{  
if  $w \neq x$  then reject  
else do as per  $M$   
}

# Emptiness problem for TM

## Lemma

*The emptiness problem for TMs,  $E_{TM} = \{\langle M \rangle \mid L(M) = \emptyset\}$ , is undecidable.*

Assume for the sake of contradiction that it is decidable. Let  $T$  be a machine that decides  $E_{TM}$ .

Let  $T'_{M,w}$  be as follows:

Let  $A$  be as follows:

On input  $x$

{  
if  $w \neq x$  then reject  
else do as per  $M$   
}



# Emptiness problem for TM

## Lemma

*The emptiness problem for TMs,  $E_{TM} = \{\langle M \rangle \mid L(M) = \emptyset\}$ , is undecidable.*

Assume for the sake of contradiction that it is decidable. Let  $T$  be a machine that decides  $E_{TM}$ .

Let  $T'_{M,w}$  be as follows:

On input  $x$

{  
if  $w \neq x$  then reject  
else do as per  $M$   
}

Let  $A$  be as follows:

On input  $M, w$

{  
Create machine  $T'_{M,w}$ .  
If  $T$  on  $\langle T'_{M,w} \rangle$  rejects  
then accept  
else reject  
}

# Equality for TM

## Lemma

*The equality problem for TMs,  $EQ_{TM} = \{(M_1, M_2) \mid L(M_1) = L(M_2)\}$ , is undecidable.*

# Equality for TM

## Lemma

*The equality problem for TMs,  $EQ_{TM} = \{(M_1, M_2) \mid L(M_1) = L(M_2)\}$ , is undecidable.*

Assume for the sake of contradiction that  $EQ_{TM}$  is decidable. Let  $M$  be the TM for it.

# Equality for TM

## Lemma

*The equality problem for TMs,  $EQ_{TM} = \{(M_1, M_2) \mid L(M_1) = L(M_2)\}$ , is undecidable.*

Assume for the sake of contradiction that  $EQ_{TM}$  is decidable. Let  $M$  be the TM for it.

Let  $M_1$  be a machine that rejects all strings. That is,  $L(M_1) = \emptyset$ .

# Equality for TM

## Lemma

*The equality problem for TMs,  $EQ_{TM} = \{(M_1, M_2) \mid L(M_1) = L(M_2)\}$ , is undecidable.*

Assume for the sake of contradiction that  $EQ_{TM}$  is decidable. Let  $M$  be the TM for it.

Let  $M_1$  be a machine that rejects all strings. That is,  $L(M_1) = \emptyset$ .

Given a machine  $M_2$  as an input, use  $M$  to check whether  $L(M_2) = L(M_1)$

# Equality for TM

## Lemma

*The equality problem for TMs,  $EQ_{TM} = \{(M_1, M_2) \mid L(M_1) = L(M_2)\}$ , is undecidable.*

Assume for the sake of contradiction that  $EQ_{TM}$  is decidable. Let  $M$  be the TM for it.

Let  $M_1$  be a machine that rejects all strings. That is,  $L(M_1) = \emptyset$ .

Given a machine  $M_2$  as an input, use  $M$  to check whether  $L(M_2) = L(M_1)$ , i.e. to check whether  $L(M_2) = \emptyset$  or not.

# Equality for TM

## Lemma

*The equality problem for TMs,  $EQ_{TM} = \{(M_1, M_2) \mid L(M_1) = L(M_2)\}$ , is undecidable.*

Assume for the sake of contradiction that  $EQ_{TM}$  is decidable. Let  $M$  be the TM for it.

Let  $M_1$  be a machine that rejects all strings. That is,  $L(M_1) = \emptyset$ .

Given a machine  $M_2$  as an input, use  $M$  to check whether  $L(M_2) = L(M_1)$ , i.e. to check whether  $L(M_2) = \emptyset$  or not.

This implies that if  $EQ_{TM}$  is decidable then  $E_{TM}$  is decidable.

# Equality for TM

## Lemma

*The equality problem for TMs,  $EQ_{TM} = \{(M_1, M_2) \mid L(M_1) = L(M_2)\}$ , is undecidable.*

Assume for the sake of contradiction that  $EQ_{TM}$  is decidable. Let  $M$  be the TM for it.

Let  $M_1$  be a machine that rejects all strings. That is,  $L(M_1) = \emptyset$ .

Given a machine  $M_2$  as an input, use  $M$  to check whether  $L(M_2) = L(M_1)$ , i.e. to check whether  $L(M_2) = \emptyset$  or not.

This implies that if  $EQ_{TM}$  is decidable then  $E_{TM}$  is decidable.

But from the previous result we know that  $E_{TM}$  is undecidable.



# Regularity checking

## Lemma

$REG_{TM} = \{\langle M \rangle \mid L(M) \text{ is regular}\}$  is undecidable.

# Regularity checking

## Lemma

$REG_{TM} = \{\langle M \rangle \mid L(M) \text{ is regular}\}$  is undecidable.