

# Strong Normalization for Intuitionistic Simple Type Theory

Wolfram Pohlers

March, 2008

## Intuitionistic simple type theory

- The system ITT

- Conversion and Reduction

- The Main Theorem

## Type free $\lambda$ -calculus $\lambda K$

## Foundation-predicates and the system IFT

- Foundation-predicates

- The Theory IFT

- Evaluating IFT-terms

- Properties of the evaluation

## Embedding IFT into $\lambda K$

## Results

# The system ITT

The basic symbols of the language of ITT

# The system ITT

The basic symbols of the language of ITT

Basic types are defined inductively by

- ▶ 0 and 1 are basic types. The type 0 stands for ground objects, the type 1 for formulas.

# The system ITT

## The basic symbols of the language of ITT

Basic types are defined inductively by

- ▶ 0 and 1 are basic types. The type 0 stands for ground objects, the type 1 for formulas.
- ▶ If  $\tau_1, \dots, \tau_n$  are basic types then  $(\tau_1, \dots, \tau_n)$  is a basic type. The type  $(\tau_1, \dots, \tau_n)$  stands for predicates taking objects of types  $\tau_1, \dots, \tau_n$  as arguments.

# The system ITT

## The basic symbols of the language of ITT

Basic types are defined inductively by

- ▶ 0 and 1 are basic types. The type 0 stands for ground objects, the type 1 for formulas.
- ▶ If  $\tau_1, \dots, \tau_n$  are basic types then  $(\tau_1, \dots, \tau_n)$  is a basic type. The type  $(\tau_1, \dots, \tau_n)$  stands for predicates taking objects of types  $\tau_1, \dots, \tau_n$  as arguments.

The non logical basic symbols of ITT comprise

- ▶ A constant 0 of type 0.

# The system ITT

## The basic symbols of the language of ITT

Basic types are defined inductively by

- ▶ 0 and 1 are basic types. The type 0 stands for ground objects, the type 1 for formulas.
- ▶ If  $\tau_1, \dots, \tau_n$  are basic types then  $(\tau_1, \dots, \tau_n)$  is a basic type. The type  $(\tau_1, \dots, \tau_n)$  stands for predicates taking objects of types  $\tau_1, \dots, \tau_n$  as arguments.

The non logical basic symbols of ITT comprise

- ▶ A constant 0 of type 0.
- ▶ Symbols for  $n$ -ary functions mapping ground object to ground objects

# The system ITT

The basic symbols of the language of ITT

The logical symbols of ITT comprise

- ▶ Basic variables



# The system ITT

The basic symbols of the language of ITT

The logical symbols of ITT comprise

- ▶ Basic variables
- ▶ The connective  $\rightarrow$  and the quantifier  $\forall$ .

# The system ITT

The basic symbols of the language of ITT

The logical symbols of ITT comprise

- ▶ Basic variables
- ▶ The connective  $\rightarrow$  and the quantifier  $\forall$ .
- ▶ The abstractor  $\{ | \}$

# The system ITT

The basic symbols of the language of ITT

The logical symbols of ITT comprise

- ▶ Basic variables
- ▶ The connective  $\rightarrow$  and the quantifier  $\forall$ .
- ▶ The abstractor  $\{ | \}$
- ▶ The abstractor  $\lambda$

# The system ITT

## Terms of ITT and their types

Inductive definition of the relation  $a \vdash \sigma$ . ( $a$  is an ITT-term of type  $\sigma$ ).

# The system ITT

## Terms of ITT and their types

Inductive definition of the relation  $a \vdash \sigma$ . ( $a$  is an ITT-term of type  $\sigma$ ).

- ▶ If  $\tau$  is a basic type and  $x$  a basic variable then  $x^\tau \vdash \tau$ .

# The system ITT

## Terms of ITT and their types

Inductive definition of the relation  $a \vdash \sigma$ . ( $a$  is an ITT-term of type  $\sigma$ ).

- ▶ If  $\tau$  is a basic type and  $x$  a basic variable then  $x^\tau \vdash \tau$ .
- ▶ If  $a \vdash (\tau_1, \dots, \tau_n)$  and  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  then  $(at_1 \dots t_n) \vdash 1$ .

# The system ITT

## Terms of ITT and their types

Inductive definition of the relation  $a \vdash \sigma$ . ( $a$  is an ITT-term of type  $\sigma$ ).

- ▶ If  $\tau$  is a basic type and  $x$  a basic variable then  $x^\tau \vdash \tau$ .
- ▶ If  $a \vdash (\tau_1, \dots, \tau_n)$  and  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  then  $(at_1 \dots t_n) \vdash 1$ .
- ▶ If  $A \vdash 1$  and  $B \vdash 1$  then  $A \rightarrow B \vdash 1$

# The system ITT

## Terms of ITT and their types

Inductive definition of the relation  $a \vdash \sigma$ . ( $a$  is an ITT-term of type  $\sigma$ ).

- ▶ If  $\tau$  is a basic type and  $x$  a basic variable then  $x^\tau \vdash \tau$ .
- ▶ If  $a \vdash (\tau_1, \dots, \tau_n)$  and  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  then  $(at_1 \dots t_n) \vdash 1$ .
- ▶ If  $A \vdash 1$  and  $B \vdash 1$  then  $A \rightarrow B \vdash 1$
- ▶ If  $A \vdash 1$  and  $\tau$  is a basic type,  $x$  basic variable then  $(\forall x^\tau)A \vdash 1$ .



# The system ITT

## Terms of ITT and their types

Inductive definition of the relation  $a \vdash \sigma$ . ( $a$  is an ITT-term of type  $\sigma$ ).

- ▶ If  $\tau$  is a basic type and  $x$  a basic variable then  $x^\tau \vdash \tau$ .
- ▶ If  $a \vdash (\tau_1, \dots, \tau_n)$  and  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  then  $(at_1 \dots t_n) \vdash 1$ .
- ▶ If  $A \vdash 1$  and  $B \vdash 1$  then  $A \rightarrow B \vdash 1$
- ▶ If  $A \vdash 1$  and  $\tau$  is a basic type,  $x$  basic variable then  $(\forall x^\tau)A \vdash 1$ .
- ▶ If  $A \vdash 1$ ,  $\tau_1, \dots, \tau_n$  are basic types and  $x$  is a basic variable then  $\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\} \vdash (\tau_1, \dots, \tau_n)$

# The system ITT

Terms of ITT and their types (continued)

- ▶ If  $A \vdash 1$  and  $x$  is a basic variable then  $x^A \vdash A$ .

# The system ITT

## Terms of ITT and their types (continued)

- ▶ If  $A \vdash 1$  and  $x$  is a basic variable then  $x^A \vdash A$ .
- ▶ If  $A \vdash 1$ ,  $B \vdash 1$  and  $a \vdash B$  then  $\lambda x^A. a \vdash A \rightarrow B$ .

# The system ITT

## Terms of ITT and their types (continued)

- ▶ If  $A \vdash 1$  and  $x$  is a basic variable then  $x^A \vdash A$ .
- ▶ If  $A \vdash 1$ ,  $B \vdash 1$  and  $a \vdash B$  then  $\lambda x^A. a \vdash A \rightarrow B$ .
- ▶ Assume  $A \vdash 1$ ,  $\tau$  is a basic type,  $x$  a basic variable and  $a \vdash A$  such that  $x^\tau$  does not occur in any type  $\sigma$  of a variable  $y^\sigma$  occurring freely in  $a$ . Then  $\lambda x^\tau. a \vdash (\forall x^\tau)A$

# The system ITT

## Terms of ITT and their types (continued)

- ▶ If  $A \vdash 1$  and  $x$  is a basic variable then  $x^A \vdash A$ .
- ▶ If  $A \vdash 1$ ,  $B \vdash 1$  and  $a \vdash B$  then  $\lambda x^A. a \vdash A \rightarrow B$ .
- ▶ Assume  $A \vdash 1$ ,  $\tau$  is a basic type,  $x$  a basic variable and  $a \vdash A$  such that  $x^\tau$  does not occur in any type  $\sigma$  of a variable  $y^\sigma$  occurring freely in  $a$ . Then  $\lambda x^\tau. a \vdash (\forall x^\tau) A$
- ▶ Let  $A \vdash 1$ ,  $\tau_1, \dots, \tau_n$  be basic types,  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$ . Then  $a \vdash A \left( \begin{smallmatrix} t_1, \dots, t_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)$  iff  $a \vdash (\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\} t_1 \dots t_n)$ .

# The system ITT

## Terms of ITT and their types (continued)

- ▶ If  $A \vdash 1$  and  $x$  is a basic variable then  $x^A \vdash A$ .
- ▶ If  $A \vdash 1$ ,  $B \vdash 1$  and  $a \vdash B$  then  $\lambda x^A. a \vdash A \rightarrow B$ .
- ▶ Assume  $A \vdash 1$ ,  $\tau$  is a basic type,  $x$  a basic variable and  $a \vdash A$  such that  $x^\tau$  does not occur in any type  $\sigma$  of a variable  $y^\sigma$  occurring freely in  $a$ . Then  $\lambda x^\tau. a \vdash (\forall x^\tau) A$
- ▶ Let  $A \vdash 1$ ,  $\tau_1, \dots, \tau_n$  be basic types,  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$ . Then  $a \vdash A \left( \begin{smallmatrix} t_1, \dots, t_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)$  iff  $a \vdash (\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\} t_1 \dots t_n)$ .
- ▶ If  $a \vdash A \rightarrow B$  and  $b \vdash A$  then  $(ab) \vdash B$

# The system ITT

## Terms of ITT and their types (continued)

- ▶ If  $A \vdash 1$  and  $x$  is a basic variable then  $x^A \vdash A$ .
- ▶ If  $A \vdash 1$ ,  $B \vdash 1$  and  $a \vdash B$  then  $\lambda x^A. a \vdash A \rightarrow B$ .
- ▶ Assume  $A \vdash 1$ ,  $\tau$  is a basic type,  $x$  a basic variable and  $a \vdash A$  such that  $x^\tau$  does not occur in any type  $\sigma$  of a variable  $y^\sigma$  occurring freely in  $a$ . Then  $\lambda x^\tau. a \vdash (\forall x^\tau)A$
- ▶ Let  $A \vdash 1$ ,  $\tau_1, \dots, \tau_n$  be basic types,  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$ . Then  $a \vdash A \left( \begin{smallmatrix} t_1, \dots, t_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)$  iff  $a \vdash (\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\} t_1 \dots t_n)$ .
- ▶ If  $a \vdash A \rightarrow B$  and  $b \vdash A$  then  $(ab) \vdash B$
- ▶ If  $\tau$  is a basic type,  $a \vdash (\forall x^\tau)A$  and  $t \vdash \tau$  then  $(at) \vdash A \left( \begin{smallmatrix} t \\ x^\tau \end{smallmatrix} \right)$

# The system ITT

## Notions

The type of an ITT-term is either a basic type (denoted by  $\tau, \tau_1, \dots$ ) or a formula denoted by  $A, B, \dots F, G \dots$

We use  $\sigma, \rho \dots$  as syntactical variables for arbitrary types.



# The system ITT

## Notions

The type of an ITT-term is either a basic type (denoted by  $\tau, \tau_1, \dots$ ) or a formula denoted by  $A, B, \dots F, G \dots$

We use  $\sigma, \rho \dots$  as syntactical variables for arbitrary types.

A variable is a basic variable equipped with a type.

# The system ITT

## Notions

The type of an ITT-term is either a basic type (denoted by  $\tau, \tau_1, \dots$ ) or a formula denoted by  $A, B, \dots F, G \dots$

We use  $\sigma, \rho \dots$  as syntactical variables for arbitrary types.

A variable is a basic variable equipped with a type.

If  $a \vdash A$  and  $A \vdash 1$  we call  $a$  a derivation for  $A$ .

# The system ITT

## Notions

The type of an ITT-term is either a basic type (denoted by  $\tau, \tau_1, \dots$ ) or a formula denoted by  $A, B, \dots F, G \dots$

We use  $\sigma, \rho \dots$  as syntactical variables for arbitrary types.

A variable is a basic variable equipped with a type.

If  $a \vdash A$  and  $A \vdash 1$  we call  $a$  a derivation for  $A$ .

By  $\overline{FV(a)}$  we denote the set of all free variables  $x^\sigma$  occurring in  $a$  such that  $\sigma \vdash 1$ .

# The system ITT

## Notions

The type of an ITT-term is either a basic type (denoted by  $\tau, \tau_1, \dots$ ) or a formula denoted by  $A, B, \dots F, G \dots$

We use  $\sigma, \rho \dots$  as syntactical variables for arbitrary types.

A variable is a basic variable equipped with a type.

If  $a \vdash A$  and  $A \vdash 1$  we call  $a$  a derivation for  $A$ .

By  $\overline{FV(a)}$  we denote the set of all free variables  $x^\sigma$  occurring in  $a$  such that  $\sigma \vdash 1$ .

A formula  $A$  is derivable in simple type theory if there is an  $a \vdash A$  such that  $\overline{FV(a)} = \emptyset$ .

# The system ITT

## Remarks

The system ITT corresponds to a natural deduction system of intuitionistic simple type theory. The set  $\overline{FV(a)}$  is the set of open assumptions in the derivation  $a \vdash F$ .

# The system ITT

## Remarks

The system ITT corresponds to a natural deduction system of intuitionistic simple type theory. The set  $\overline{FV}(a)$  is the set of open assumptions in the derivation  $a \vdash F$ . Observe that falsum and thus negation and the other connectives and quantifiers are definable

# The system ITT

## Remarks

The system ITT corresponds to a natural deduction system of intuitionistic simple type theory. The set  $\overline{FV(a)}$  is the set of open assumptions in the derivation  $a \vdash F$ . Observe that falsum and thus negation and the other connectives and quantifiers are definable

- ▶  $\perp$  is the formula  $(\forall x^1)x^1$  and  $\neg A$  the formula  $A \rightarrow \perp$ .

# The system ITT

## Remarks

The system ITT corresponds to a natural deduction system of intuitionistic simple type theory. The set  $\overline{FV(a)}$  is the set of open assumptions in the derivation  $a \vdash F$ . Observe that falsum and thus negation and the other connectives and quantifiers are definable

- ▶  $\perp$  is the formula  $(\forall x^1)x^1$  and  $\neg A$  the formula  $A \rightarrow \perp$ .
- ▶  $A \wedge B$  is defined by  $(\forall x^1)[(A \rightarrow (B \rightarrow x^1)) \rightarrow x^1]$



# The system ITT

## Remarks

The system ITT corresponds to a natural deduction system of intuitionistic simple type theory. The set  $\overline{FV}(a)$  is the set of open assumptions in the derivation  $a \vdash F$ . Observe that falsum and thus negation and the other connectives and quantifiers are definable

- ▶  $\perp$  is the formula  $(\forall x^1)x^1$  and  $\neg A$  the formula  $A \rightarrow \perp$ .
- ▶  $A \wedge B$  is defined by  $(\forall x^1)[(A \rightarrow (B \rightarrow x^1)) \rightarrow x^1]$
- ▶  $A \vee B$  is defined by  $(\forall x^1)[(A \rightarrow x^1) \rightarrow ((B \rightarrow x^1) \rightarrow x^1)]$

# The system ITT

## Remarks

The system ITT corresponds to a natural deduction system of intuitionistic simple type theory. The set  $\overline{FV(a)}$  is the set of open assumptions in the derivation  $a \vdash F$ . Observe that falsum and thus negation and the other connectives and quantifiers are definable

- ▶  $\perp$  is the formula  $(\forall x^1)x^1$  and  $\neg A$  the formula  $A \rightarrow \perp$ .
- ▶  $A \wedge B$  is defined by  $(\forall x^1)[(A \rightarrow (B \rightarrow x^1)) \rightarrow x^1]$
- ▶  $A \vee B$  is defined by  $(\forall x^1)[(A \rightarrow x^1) \rightarrow ((B \rightarrow x^1) \rightarrow x^1)]$
- ▶  $(\exists x^\tau)F(x^\tau)$  is defined by  $(\forall x^1)[(\forall x^\tau)(F(x^\tau) \rightarrow x^1) \rightarrow x^1]$ .

# Conversion and Reduction

## Simple reductions

Conversion  $\succ_0$  is defined as usual by:

if  $b \vdash \sigma$  then  $(\lambda x^\sigma. a)b \succ_0 a \left( \begin{smallmatrix} b \\ x^\sigma \end{smallmatrix} \right)$ .

# Conversion and Reduction

## Simple reductions

Conversion  $\succ_0$  is defined as usual by:

if  $b \vdash \sigma$  then  $(\lambda x^\sigma. a)b \succ_0 a \left( \begin{smallmatrix} b \\ x^\sigma \end{smallmatrix} \right)$ .

We define the simple reduction step  $a \succ_1 b$  by the following clauses

$$(R_0) \ a \succ_0 b \Rightarrow a \succ_1 b$$

$$(R_1) \ a \succ_1 b \Rightarrow \lambda x^\sigma. a \succ_1 \lambda x^\sigma. b$$

$$(R_2) \ a \succ_1 b \Rightarrow ac \succ_1 bc \text{ and } ca \succ_1 cb$$

# Conversion and Reduction

## Simple reductions

Conversion  $\succ_0$  is defined as usual by:

if  $b \vdash \sigma$  then  $(\lambda x^\sigma. a)b \succ_0 a \left( \begin{smallmatrix} b \\ x^\sigma \end{smallmatrix} \right)$ .

We define the simple reduction step  $a \succ_1 b$  by the following clauses

$$(R_0) \ a \succ_0 b \Rightarrow a \succ_1 b$$

$$(R_1) \ a \succ_1 b \Rightarrow \lambda x^\sigma. a \succ_1 \lambda x^\sigma. b$$

$$(R_2) \ a \succ_1 b \Rightarrow ac \succ_1 bc \text{ and } ca \succ_1 cb$$

A term  $a$  is normal iff there is no term  $b$  such that  $a \succ_1 b$ .

# Conversion and Reduction

## Reduction Chains

A *reduction chain* for an ITT-term  $a$  is a sequence  $\mathfrak{R} := \langle a_i \mid i \in I \rangle$  for  $I \subseteq \omega$  such that

# Conversion and Reduction

## Reduction Chains

A *reduction chain* for an ITT-term  $a$  is a sequence

$\mathfrak{R} := \langle a_i \mid i \in I \rangle$  for  $I \subseteq \omega$  such that

►  $a_0 = a$

# Conversion and Reduction

## Reduction Chains

A *reduction chain* for an ITT-term  $a$  is a sequence

$\mathfrak{R} := \langle a_i \mid i \in I \rangle$  for  $I \subseteq \omega$  such that

- ▶  $a_0 = a$
- ▶ If  $a_i \in \mathfrak{R}$  is normal, then  $a_i$  is the last element of the chain.



# Conversion and Reduction

## Reduction Chains

A *reduction chain* for an ITT-term  $a$  is a sequence

$\mathfrak{R} := \langle a_i \mid i \in I \rangle$  for  $I \subseteq \omega$  such that

- ▶  $a_0 = a$
- ▶ If  $a_i \in \mathfrak{R}$  is normal, then  $a_i$  is the last element of the chain.
- ▶ If  $a_i \in \mathfrak{R}$  is not normal, then  $a_i \succ_1 a_{i+1}$ .

# Conversion and Reduction

## Reduction Chains

A *reduction chain* for an ITT-term  $a$  is a sequence

$\mathfrak{R} := \langle a_i \mid i \in I \rangle$  for  $I \subseteq \omega$  such that

- ▶  $a_0 = a$
- ▶ If  $a_i \in \mathfrak{R}$  is normal, then  $a_i$  is the last element of the chain.
- ▶ If  $a_i \in \mathfrak{R}$  is not normal, then  $a_i \succ_1 a_{i+1}$ .

We introduce the following notions

# Conversion and Reduction

## Reduction Chains

A *reduction chain* for an ITT-term  $a$  is a sequence

$\mathfrak{R} := \langle a_i \mid i \in I \rangle$  for  $I \subseteq \omega$  such that

- ▶  $a_0 = a$
- ▶ If  $a_i \in \mathfrak{R}$  is normal, then  $a_i$  is the last element of the chain.
- ▶ If  $a_i \in \mathfrak{R}$  is not normal, then  $a_i \succ_1 a_{i+1}$ .

We introduce the following notions

- ▶ By  $a \succ b$  we denote that there is a reduction chain  $\mathfrak{R}$  for  $a$  such that  $b \in \mathfrak{R}$ .

# Conversion and Reduction

## Reduction Chains

A *reduction chain* for an ITT-term  $a$  is a sequence  $\mathfrak{R} := \langle a_i \mid i \in I \rangle$  for  $I \subseteq \omega$  such that

- ▶  $a_0 = a$
- ▶ If  $a_i \in \mathfrak{R}$  is normal, then  $a_i$  is the last element of the chain.
- ▶ If  $a_i \in \mathfrak{R}$  is not normal, then  $a_i \succ_1 a_{i+1}$ .

We introduce the following notions

- ▶ By  $a \succ b$  we denote that there is a reduction chain  $\mathfrak{R}$  for  $a$  such that  $b \in \mathfrak{R}$ .
- ▶ An ITT-term  $a$  is well-founded, if every reduction chain for  $a$  is finite.

# The Main Theorem

# The Main Theorem

Our aim is to prove the following strong normalization theorem for intuitionistic type theory.

# The Main Theorem

Our aim is to prove the following strong normalization theorem for intuitionistic type theory.

## Theorem

*Every closed ITT-term is well-founded.*

# Type free $\lambda$ -calculus

## Basic facts

We assume familiarity with the type free lambda calculus  $\lambda K$ . As a peculiarity we require that there is a constant  $u$  in the  $\lambda$ -calculus. Conversion  $\succ_0$ , simple reduction  $\succ_1$  and reduction chains and the reduction relation  $a \succ b$  are defined as for [ITT](#).



# Type free $\lambda$ -calculus

## Basic facts

We assume familiarity with the type free lambda calculus  $\lambda K$ . As a peculiarity we require that there is a constant  $u$  in the  $\lambda$ -calculus. Conversion  $\succ_0$ , simple reduction  $\succ_1$  and reduction chains and the reduction relation  $a \succ b$  are defined as for [ITT](#).

- Terms of the form  $\lambda x. a$  are called  $L$ -terms.

# Type free $\lambda$ -calculus

## Basic facts

We assume familiarity with the type free lambda calculus  $\lambda K$ . As a peculiarity we require that there is a constant  $u$  in the  $\lambda$ -calculus. Conversion  $\succ_0$ , simple reduction  $\succ_1$  and reduction chains and the reduction relation  $a \succ b$  are defined as for [ITT](#).

- ▶ Terms of the form  $\lambda x. a$  are called  $L$ -terms.
- ▶ A  $\lambda$ -term  $t$  is strongly normal iff it is normal and not an  $L$ -term.

# Type free $\lambda$ -calculus

## Basic facts

We assume familiarity with the type free lambda calculus  $\lambda K$ . As a peculiarity we require that there is a constant  $u$  in the  $\lambda$ -calculus. Conversion  $\succ_0$ , simple reduction  $\succ_1$  and reduction chains and the reduction relation  $a \succ b$  are defined as for [ITT](#).

- ▶ Terms of the form  $\lambda x. a$  are called  $L$ -terms.
- ▶ A  $\lambda$ -term  $t$  is strongly normal iff it is normal and not an  $L$ -term.
- ▶ A reduction chain  $\Re$  for an  $\lambda$ -term is normal, if it is finite.

# Type free $\lambda$ -calculus

## Basic facts

We assume familiarity with the type free lambda calculus  $\lambda K$ . As a peculiarity we require that there is a constant  $u$  in the  $\lambda$ -calculus. Conversion  $\succ_0$ , simple reduction  $\succ_1$  and reduction chains and the reduction relation  $a \succ b$  are defined as for [ITT](#).

- ▶ Terms of the form  $\lambda x. a$  are called  $L$ -terms.
- ▶ A  $\lambda$ -term  $t$  is strongly normal iff it is normal and not an  $L$ -term.
- ▶ A reduction chain  $\Re$  for an  $\lambda$ -term is normal, if it is finite.
- ▶ We call a reduction chain strongly normal if it is normal and its last term is strongly normal.

# Type free $\lambda$ -calculus

## Basic facts

### Lemma

*If  $\mathfrak{R} = \langle c_i \mid i \in I \rangle$  is a reduction chain for an  $\lambda$ -term  $(ab)$  and  $i$  is its least index such that  $c_i \succ_1 c_{i+1}$  is a reduction according to  $R_0$ , i.e., a conversion, then for all  $j \leq i$  the term  $c_j$  has the form  $(a'_j b'_j)$  such that  $a \succ a'_j$  and  $b \succ b'_j$*

# Type free $\lambda$ -calculus

## Basic facts

### Lemma

*If  $\mathfrak{R} = \langle c_i \mid i \in I \rangle$  is a reduction chain for an  $\lambda$ -term  $(ab)$  and  $i$  is its least index such that  $c_i \succ_1 c_{i+1}$  is a reduction according to  $R_0$ , i.e., a conversion, then for all  $j \leq i$  the term  $c_j$  has the form  $(a'_j b'_j)$  such that  $a \succ a'_j$  and  $b \succ b'_j$*

### Lemma (Lemma I)

*If  $a$  and  $b$  are well-founded  $\lambda$ -terms then every reduction chain  $\mathfrak{R}$  for  $(ab)$  which contains no conversion according to  $R_0$  is strongly normal.*

► Back to Main Lemma (3.1)

► Back to Main Lemma (3.2)

# Foundation—predicates

# Foundation–predicates

## Definition

$(F_0)$  Every closed object term of ITT is a foundation–predicate of type 0



# Foundation–predicates

## Definition

- $(F_0)$  Every closed object term of ITT is a foundation–predicate of type 0
- $(F_1)$  A set  $\mathfrak{F}^1$  of closed  $\lambda$ –terms is a foundation–predicate of type 1 iff the following conditions are satisfied

# Foundation–predicates

## Definition

- $(F_0)$  Every closed object term of ITT is a foundation–predicate of type 0
- $(F_1)$  A set  $\mathfrak{F}^1$  of closed  $\lambda$ –terms is a foundation–predicate of type 1 iff the following conditions are satisfied
  - $(F_{11})$  All  $a \in \mathfrak{F}^1$  are well–founded

# Foundation–predicates

## Definition

- $(F_0)$  Every closed object term of ITT is a foundation–predicate of type 0
- $(F_1)$  A set  $\mathfrak{F}^1$  of closed  $\lambda$ –terms is a foundation–predicate of type 1 iff the following conditions are satisfied
  - $(F_{11})$  All  $a \in \mathfrak{F}^1$  are well–founded
  - $(F_{12})$   $a \in \mathfrak{F}^1$  and  $a \succ b \Rightarrow b \in \mathfrak{F}^1$

# Foundation–predicates

## Definition

- $(F_0)$  Every closed object term of ITT is a foundation–predicate of type 0
- $(F_1)$  A set  $\mathfrak{F}^1$  of closed  $\lambda$ –terms is a foundation–predicate of type 1 iff the following conditions are satisfied
  - $(F_{11})$  All  $a \in \mathfrak{F}^1$  are well–founded
  - $(F_{12})$   $a \in \mathfrak{F}^1$  and  $a \succ b \Rightarrow b \in \mathfrak{F}^1$
- $(F_2)$  A foundation–predicate of type  $(\tau_1, \dots, \tau_n)$  is a mapping which assigns a foundation–predicate of type 1 to any tuple  $\mathfrak{F}_1^{\tau_1}, \dots, \mathfrak{F}_n^{\tau_n}$  of foundation–predicates of type  $\tau_i (i = 1, \dots, n)$ .

# The theory IFT

The language

# The theory IFT

## The language

We extend the basic symbols of the language ITT by names for every foundation–predicate of type  $\tau$  and extend the definition of the ITT–terms by the following clauses:

# The theory IFT

## The language

We extend the basic symbols of the language ITT by names for every foundation–predicate of type  $\tau$  and extend the definition of the ITT–terms by the following clauses:

- ▶ If  $\mathfrak{F}^\tau$  is a name for a foundation–predicate of type  $\tau$ , then  $\mathfrak{F}^\tau \vdash \tau$ .

# The theory IFT

## The language

We extend the basic symbols of the language ITT by names for every foundation–predicate of type  $\tau$  and extend the definition of the ITT–terms by the following clauses:

- ▶ If  $\mathfrak{F}^\tau$  is a name for a foundation–predicate of type  $\tau$ , then  $\mathfrak{F}^\tau \vdash \tau$ .
- ▶ Let  $\tau_1, \dots, \tau_n$  be a tuple of basic types,  $t \vdash \tau_i$  for  $i = 1, \dots, n$  and  $\mathfrak{F}^\tau$  a foundation–predicate of type  $\tau = (\tau_1, \dots, \tau_n)$ . Then  $(\mathfrak{F}^\tau t_1 \dots t_n) \vdash 1$ .



# Evaluating IFT-terms

Defining  $|a|$

# Evaluating IFT-terms

Defining  $|a|$

## Definition

We define the evaluation  $|a|$  for every closed IFT-term  $a \vdash \sigma$  inductively by the following clauses:

# Evaluating IFT-terms

Defining  $|a|$

## Definition

We define the evaluation  $|a|$  for every closed IFT-term  $a \vdash \sigma$  inductively by the following clauses:

$(B_0)$  If  $a \vdash 0$  then  $|a| := a$ .

# Evaluating IFT-terms

Defining  $|a|$

## Definition

We define the evaluation  $|a|$  for every closed IFT-term  $a \vdash \sigma$  inductively by the following clauses:

$(B_0)$  If  $a \vdash 0$  then  $|a| := a$ .

$(B_1)$  For a closed  $\lambda$ -term  $t$  we put  $t \in |\mathfrak{F}^1|$  if every reduction chain  $\mathfrak{R}$  of  $t$  which is not strongly normal contains an element  $t' \in \mathfrak{F}^1$  which is not preceded by an  $L$ -term in  $\mathfrak{R}$ .

# Evaluating IFT-terms

Defining  $|a|$

## Definition

We define the evaluation  $|a|$  for every closed IFT-term  $a \vdash \sigma$  inductively by the following clauses:

- $(B_0)$  If  $a \vdash 0$  then  $|a| := a$ .
- $(B_1)$  For a closed  $\lambda$ -term  $t$  we put  $t \in |\mathfrak{F}^1|$  if every reduction chain  $\mathfrak{R}$  of  $t$  which is not strongly normal contains an element  $t' \in \mathfrak{F}^1$  which is not preceded by an  $L$ -term in  $\mathfrak{R}$ .
- $(B_2)$  If  $\tau = (\tau_1, \dots, \tau_n)$  is a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  are closed IFT-terms with  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$ , then  $t \in |at_1 \dots t_n|$  holds true iff  $t$  is a closed  $\lambda$ -term such that every reduction chain  $\mathfrak{R}$  of  $t$  which is not strongly normal contains an element  $s \in |a|(|t_1|, \dots, |t_n|)$  which is not preceded by an  $L$ -term in  $\mathfrak{R}$ .

# Evaluating IFT-terms

Defining  $|a|$  continued

## Definition

(continued)

$(B_3)$   $t \in |A \rightarrow B|$  holds true iff  $t$  is a closed  $\lambda$ -term such that every reduction chain  $\mathfrak{R}$  of  $t$ , which is not strongly normal, contains a first  $L$ -term  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} b \\ x \end{smallmatrix} \right) \in |B|$  for all  $b \in |A|$

# Evaluating IFT-terms

Defining  $|a|$  continued

## Definition

(continued)

- $(B_3)$   $t \in |A \rightarrow B|$  holds true iff  $t$  is a closed  $\lambda$ -term such that every reduction chain  $\mathfrak{R}$  of  $t$ , which is not strongly normal, contains a first  $L$ -term  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} b \\ x \end{smallmatrix} \right) \in |B|$  for all  $b \in |A|$
- $(B_4)$   $t \in |(\forall x^\tau A)|$  holds true iff  $t$  is a closed  $\lambda$ -term such that every reduction chain  $\mathfrak{R}$  of  $t$ , which is not strongly normal, contains a first  $L$ -term  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} u \\ x \end{smallmatrix} \right) \in |A \left( \begin{smallmatrix} \mathfrak{F}^\tau \\ x^\tau \end{smallmatrix} \right)|$  for all foundation-predicates  $\mathfrak{F}^\tau$  of type  $\tau$ .

# Evaluating IFT-terms

Defining  $|a|$

## Definition

(continued)

( $B_5$ ) If  $\tau := (t_1, \dots, t_n)$  is a basis type and  $\mathfrak{F}^\tau$  is a name for a foundation-predicate of type  $\tau$ , then then  $|\mathfrak{F}^\tau|$  is the mapping defined by  $|\mathfrak{F}^\tau|(\mathfrak{F}_1, \dots, \mathfrak{F}_n) := |\mathfrak{F}^\tau(\mathfrak{F}_1, \dots, \mathfrak{F}_n)|$  for any tuple  $\mathfrak{F}_1, \dots, \mathfrak{F}_n$  of foundation-predicates of types  $\tau_1, \dots, \tau_n$ .



# Evaluating IFT-terms

Defining  $|a|$

## Definition

(continued)

- ( $B_5$ ) If  $\tau := (t_1, \dots, t_n)$  is a basis type and  $\mathfrak{F}^\tau$  is a name for a foundation-predicate of type  $\tau$ , then then  $|\mathfrak{F}^\tau|$  is the mapping defined by  $|\mathfrak{F}^\tau|(\mathfrak{F}_1, \dots, \mathfrak{F}_n) := |\mathfrak{F}^\tau(\mathfrak{F}_1, \dots, \mathfrak{F}_n)|$  for any tuple  $\mathfrak{F}_1, \dots, \mathfrak{F}_n$  of foundation-predicates of types  $\tau_1, \dots, \tau_n$ .
- ( $B_6$ ) If  $\tau = (\tau_1, \dots, \tau_n)$  is a basic type then
- $$|\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\}|(\mathfrak{F}_1, \dots, \mathfrak{F}_n) := |A \left( \begin{smallmatrix} \mathfrak{F}_1, \dots, \mathfrak{F}_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)| \text{ for any}$$
- tuple  $\mathfrak{F}_1, \dots, \mathfrak{F}_n$  of foundation-predicates of types  $\tau_1, \dots, \tau_n$ .

# Evaluating IFT-terms

Properties of  $|a|$

## Lemma

*If  $\tau$  is a basic type and  $a \vdash \tau$  is a closed IFT-term then  $|a|$  is a foundation-predicate of type  $\tau$ .*

# Evaluating IFT-terms

Properties of  $|a|$

## Lemma

*If  $\tau$  is a basic type and  $a \vdash \tau$  is a closed IFT-term then  $|a|$  is a foundation-predicate of type  $\tau$ .*

► Proof ( $B_1$ )

► Proof ( $B_2$ )

► Proof ( $B_3$ )

► Proof ( $B_4$ )

► Proof ( $B_5$ )

► Proof ( $B_6$ )

# Evaluating IFT-terms

Properties of  $|a|$

## Lemma

*If  $\tau$  is a basic type and  $a \vdash \tau$  is a closed IFT-term then  $|a|$  is a foundation-predicate of type  $\tau$ .*

► Proof ( $B_1$ )

► Proof ( $B_2$ )

► Proof ( $B_3$ )

► Proof ( $B_4$ )

► Proof ( $B_5$ )

► Proof ( $B_6$ )

## Lemma

*Let  $A$  be an IFT-sentence and  $t$  a  $\lambda$ -functional. If every reduction chain  $\Re$  of  $t$ , which is not strongly normal, contains an element  $t' \in |A|$  which is not preceeded by  $L$ -terms in  $\Re$  then  $t \in |A|$ .*

# Evaluating IFT-terms

Properties of  $|a|$

## Lemma

*If  $\tau$  is a basic type and  $a \vdash \tau$  is a closed IFT-term then  $|a|$  is a foundation-predicate of type  $\tau$ .*

► Proof ( $B_1$ )

► Proof ( $B_2$ )

► Proof ( $B_3$ )

► Proof ( $B_4$ )

► Proof ( $B_5$ )

► Proof ( $B_6$ )

## Lemma

*Let  $A$  be an IFT-sentence and  $t$  a  $\lambda$ -functional. If every reduction chain  $\Re$  of  $t$ , which is not strongly normal, contains an element  $t' \in |A|$  which is not preceeded by  $L$ -terms in  $\Re$  then  $t \in |A|$ .*

► Proof (atomic case)

► Proof ( $\rightarrow$  case)

► Proof ( $\forall$  case)

# Evaluating IFT-terms

Properties of  $|a|$

## Lemma

*Let  $\tau$  be a basic type and  $a \vdash \tau$  for a closed IFT-term  $a$ . Then  $|a| = \|a\|$ .*

► Proof

# Evaluating IFT-terms

Properties of  $|a|$

## Lemma

*Let  $\tau$  be a basic type and  $a \vdash \tau$  for a closed IFT-term  $a$ . Then  $|a| = \|a\|$ .*

► Proof

## Corollary

*Let  $FV(a) = \{x^\tau\}$ . For any closed IFT-term  $b \vdash \tau$  we obtain  $|a \left( \begin{smallmatrix} b \\ x^\tau \end{smallmatrix} \right)| = |a \left( \begin{smallmatrix} |b| \\ x^\tau \end{smallmatrix} \right)|$ .*

# Embedding IFT into $\lambda K$

Defining the embedding



# Embedding IFT into $\lambda K$

Defining the embedding

## Definition

For  $a \vdash \sigma$  we define  $\bar{a} \in \lambda K$  inductively by the following clauses:

# Embedding IFT into $\lambda K$

## Defining the embedding

### Definition

For  $a \vdash \sigma$  we define  $\bar{a} \in \lambda K$  inductively by the following clauses:

- ▶ If  $a \vdash \tau$  and  $\tau$  is a basic type then  $\bar{a} := u$ .

# Embedding IFT into $\lambda K$

## Defining the embedding

### Definition

For  $a \vdash \sigma$  we define  $\bar{a} \in \lambda K$  inductively by the following clauses:

- ▶ If  $a \vdash \tau$  and  $\tau$  is a basic type then  $\bar{a} := u$ .
- ▶ If  $A \vdash 1$  and  $x$  is a basic variable then  $\overline{x^A} := x$ .

# Embedding IFT into $\lambda K$

## Defining the embedding

### Definition

For  $a \vdash \sigma$  we define  $\bar{a} \in \lambda K$  inductively by the following clauses:

- ▶ If  $a \vdash \tau$  and  $\tau$  is a basic type then  $\bar{a} := u$ .
- ▶ If  $A \vdash 1$  and  $x$  is a basic variable then  $\overline{x^A} := x$ .
- ▶  $\overline{\lambda x^\sigma. a} = \lambda x. \bar{a}$ .

# Embedding IFT into $\lambda K$

## Defining the embedding

### Definition

For  $a \vdash \sigma$  we define  $\bar{a} \in \lambda K$  inductively by the following clauses:

- ▶ If  $a \vdash \tau$  and  $\tau$  is a basic type then  $\bar{a} := u$ .
- ▶ If  $A \vdash 1$  and  $x$  is a basic variable then  $\overline{x^A} := x$ .
- ▶  $\overline{\lambda x^\sigma. a} = \lambda x. \bar{a}$ .
- ▶  $\overline{(ab)} := (\bar{a} \bar{b})$ .

# Embedding IFT into $\lambda K$

## Properties of the embedding

### Lemma

$$\blacktriangleright \overline{a \left( \begin{smallmatrix} b \\ x^\sigma \end{smallmatrix} \right)} = \bar{a} \left( \begin{smallmatrix} \bar{b} \\ x \end{smallmatrix} \right).$$

# Embedding IFT into $\lambda K$

## Properties of the embedding

### Lemma

- ▶  $\overline{a \left( \overline{b} \right)_{x^\sigma}} = \bar{a} \left( \overline{\bar{b}} \right)_x$ .
- ▶  $a \succ_0 b$  implies  $\bar{a} \succ_0 \bar{b}$ .

# Embedding IFT into $\lambda K$

## Properties of the embedding

### Lemma

- ▶  $\overline{a \left( \overline{b} \right)_{x^\sigma}} = \bar{a} \left( \overline{\bar{b}} \right)_x$ .
- ▶  $a \succ_0 b$  implies  $\bar{a} \succ_0 \bar{b}$ .
- ▶  $a \succ_1 b$  implies  $\bar{a} \succ_1 \bar{b}$ .



# Embedding IFT into $\lambda K$

## Properties of the embedding

### Lemma

- ▶  $\overline{a \left( \overline{b} \right)_{x^\sigma}} = \overline{a} \left( \overline{\overline{b}} \right)_x$ .
- ▶  $a \succ_0 b$  implies  $\overline{a} \succ_0 \overline{b}$ .
- ▶  $a \succ_1 b$  implies  $\overline{a} \succ_1 \overline{b}$ .
- ▶ If  $\mathfrak{R} = \langle c_i \mid i \in I \rangle$  is a reduction chain for  $a$  then  $\overline{\mathfrak{R}} := \langle \overline{c_i} \mid i \in I \rangle$  is a reduction chain for  $\overline{a}$ .

# Embedding IFT into $\lambda K$

## Properties of the embedding

### Lemma

- ▶  $\overline{a \left( \overline{b} \right)_{x^\sigma}} = \bar{a} \left( \bar{b} \right)_x$ .
- ▶  $a \succ_0 b$  implies  $\bar{a} \succ_0 \bar{b}$ .
- ▶  $a \succ_1 b$  implies  $\bar{a} \succ_1 \bar{b}$ .
- ▶ If  $\mathfrak{R} = \langle c_i \mid i \in I \rangle$  is a reduction chain for  $a$  then  $\bar{\mathfrak{R}} := \langle \bar{c}_i \mid i \in I \rangle$  is a reduction chain for  $\bar{a}$ .

### Theorem

*Well-foundedness of  $\bar{a}$  entails the well-foundedness of  $a$ .*

# Results

## The Main Lemma

# Results

## The Main Lemma

### Lemma

Let  $A \vdash 1$  and  $a \vdash A$  be IFT-terms. Let  $FV(a) \setminus \overline{FV(a)} = \{x^{\tau_1}, \dots, x^{\tau_n}\}$  and  $c_i$  closed terms such that  $c_i \vdash \tau_i$  for  $i = 1, \dots, n$ . Then  $FV(a \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)) = \{y^{B_1}, \dots, y^{B_m}\}$  and  $b_i \in |B_i|$  imply

$$\overline{a \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)} \left( \begin{smallmatrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{smallmatrix} \right) \in |A \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)|.$$

► Proof (case 1)

► Proof (case 2.1)

► Proof (case 2.2)

► Proof (case 3.1)

► Proof (case 3.2)

## Theorem

*Let  $A$  be a sentence and  $a \vdash A$  for a closed IFT-term  $a$ . Then  $\bar{a} \in |A|$ .*

## Theorem

*Let  $A$  be a sentence and  $a \vdash A$  for a closed IFT-term  $a$ . Then  $\bar{a} \in |A|$ .*

## Corollary

*Every closed ITT-term is strongly normalizable.*

Thank you for your attention

# Proof of the Evaluation Lemma

Case ( $B_1$ )

We prove the lemma by induction on the definition of  $|a|$ .



# Proof of the Evaluation Lemma

Case ( $B_1$ )

We prove the lemma by induction on the definition of  $|a|$ .  
The claim holds trivially for ( $B_0$ ).

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ .

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal.

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in \mathfrak{F}^1$ .

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in \mathfrak{F}^1$ . Since  $\mathfrak{F}^1$  is a foundation-predicate, every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite.

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in \mathfrak{F}^1$ . Since  $\mathfrak{F}^1$  is a foundation-predicate, every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite. But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in \mathfrak{F}^1$ . Since  $\mathfrak{F}^1$  is a foundation-predicate, every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite.

But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ .

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in \mathfrak{F}^1$ . Since  $\mathfrak{F}^1$  is a foundation-predicate, every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite.

But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is strongly normal then so is  $\mathfrak{R}'$ .



# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in \mathfrak{F}^1$ . Since  $\mathfrak{F}^1$  is a foundation-predicate, every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite.

But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is strongly normal then so is  $\mathfrak{R}'$ . Assume that  $\mathfrak{R}$  is not strongly normal. Then  $\mathfrak{R}'$  is also not strongly normal and therefore contains a  $\lambda$ -functional  $s' \in \mathfrak{F}^1$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$ .

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in \mathfrak{F}^1$ . Since  $\mathfrak{F}^1$  is a foundation-predicate, every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite.

But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is strongly normal then so is  $\mathfrak{R}'$ . Assume that  $\mathfrak{R}$  is not strongly normal. Then  $\mathfrak{R}'$  is also not strongly normal and therefore contains a  $\lambda$ -functional  $s' \in \mathfrak{F}^1$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$ .

If  $s' \notin \mathfrak{R}$  we get  $s' \succ s$  and thus also  $s \in \mathfrak{F}^1$ .

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in \mathfrak{F}^1$ . Since  $\mathfrak{F}^1$  is a foundation-predicate, every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite.

But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is strongly normal then so is  $\mathfrak{R}'$ . Assume that  $\mathfrak{R}$  is not strongly normal. Then  $\mathfrak{R}'$  is also not strongly normal and therefore contains a  $\lambda$ -functional  $s' \in \mathfrak{F}^1$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$ .

If  $s' \notin \mathfrak{R}$  we get  $s' \succ s$  and thus also  $s \in \mathfrak{F}^1$ . Hence  $s \in |\mathfrak{F}^1|$  by  $(B_1)$ .

# Proof of the Evaluation Lemma

## Case $(B_1)$

We prove the lemma by induction on the definition of  $|a|$ .

The claim holds trivially for  $(B_0)$ .

To handle the case of  $(B_1)$  let  $t \in |\mathfrak{F}^1|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in \mathfrak{F}^1$ . Since  $\mathfrak{F}^1$  is a foundation-predicate, every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite.

But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is strongly normal then so is  $\mathfrak{R}'$ . Assume that  $\mathfrak{R}$  is not strongly normal. Then  $\mathfrak{R}'$  is also not strongly normal and therefore contains a  $\lambda$ -functional  $s' \in \mathfrak{F}^1$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$ .

If  $s' \notin \mathfrak{R}$  we get  $s' \succ s$  and thus also  $s \in \mathfrak{F}^1$ . Hence  $s \in |\mathfrak{F}^1|$  by  $(B_1)$ .

If  $s' \in \mathfrak{R}$  then there are no  $L$ -terms in  $\mathfrak{R}$  which precede  $s'$ . Hence  $s \in |\mathfrak{F}^1|$  by  $(B_1)$ .

► Back

# Proof of the Evaluation Lemma

## Case $(B_2)$

To handle the case of  $(B_2)$  let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ .

# Proof of the Evaluation Lemma

## Case $(B_2)$

To handle the case of  $(B_2)$  let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ . Let  $\Re$  be a reduction chain for  $t$  which is not strongly normal.

# Proof of the Evaluation Lemma

## Case $(B_2)$

To handle the case of  $(B_2)$  let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |a|(|t_1|, \dots, |t_n|)$ .

# Proof of the Evaluation Lemma

## Case $(B_2)$

To handle the case of  $(B_2)$  let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |a|(|t_1|, \dots, |t_n|)$ . Since by induction hypothesis  $|a|$  is a foundation-predicate of type  $\tau$  and  $|t_i|$  a foundation-predicate of type  $\tau_i$  we obtain  $|a|(|t_1|, \dots, |t_n|)$  as a foundation-predicate of type 1.



# Proof of the Evaluation Lemma

## Case $(B_2)$

To handle the case of  $(B_2)$  let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |a|(|t_1|, \dots, |t_n|)$ . Since by induction hypothesis  $|a|$  is a foundation-predicate of type  $\tau$  and  $|t_i|$  a foundation-predicate of type  $\tau_i$  we obtain  $|a|(|t_1|, \dots, |t_n|)$  as a foundation-predicate of type 1. Thus every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite. But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

# Proof of the Evaluation Lemma

## Case ( $B_2$ )

To handle the case of ( $B_2$ ) let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |a|(|t_1|, \dots, |t_n|)$ . Since by induction hypothesis  $|a|$  is a foundation-predicate of type  $\tau$  and  $|t_i|$  a foundation-predicate of type  $\tau_i$  we obtain  $|a|(|t_1|, \dots, |t_n|)$  as a foundation-predicate of type 1. Thus every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite. But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ .

# Proof of the Evaluation Lemma

## Case $(B_2)$

To handle the case of  $(B_2)$  let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |a|(|t_1|, \dots, |t_n|)$ . Since by induction hypothesis  $|a|$  is a foundation-predicate of type  $\tau$  and  $|t_i|$  a foundation-predicate of type  $\tau_i$  we obtain  $|a|(|t_1|, \dots, |t_n|)$  as a foundation-predicate of type 1. Thus every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite. But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is not strongly normal then  $\mathfrak{R}'$  is not strongly normal, too, and thus contains a  $\lambda$ -functional  $s' \in |a|(|t_1|, \dots, |t_n|)$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$ .

# Proof of the Evaluation Lemma

## Case ( $B_2$ )

To handle the case of ( $B_2$ ) let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |a|(|t_1|, \dots, |t_n|)$ . Since by induction hypothesis  $|a|$  is a foundation-predicate of type  $\tau$  and  $|t_i|$  a foundation-predicate of type  $\tau_i$  we obtain  $|a|(|t_1|, \dots, |t_n|)$  as a foundation-predicate of type 1. Thus every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite. But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is not strongly normal then  $\mathfrak{R}'$  is not strongly normal, too, and thus contains a  $\lambda$ -functional  $s' \in |a|(|t_1|, \dots, |t_n|)$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$ . If  $s' \notin \mathfrak{R}$  we get  $s' \succ s$  and thus also  $s \in |a|(|t_1|, \dots, |t_n|)$ .

# Proof of the Evaluation Lemma

## Case ( $B_2$ )

To handle the case of ( $B_2$ ) let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |a|(|t_1|, \dots, |t_n|)$ . Since by induction hypothesis  $|a|$  is a foundation-predicate of type  $\tau$  and  $|t_i|$  a foundation-predicate of type  $\tau_i$  we obtain  $|a|(|t_1|, \dots, |t_n|)$  as a foundation-predicate of type 1. Thus every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite. But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is not strongly normal then  $\mathfrak{R}'$  is not strongly normal, too, and thus contains a  $\lambda$ -functional  $s' \in |a|(|t_1|, \dots, |t_n|)$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$ . If  $s' \notin \mathfrak{R}$  we get  $s' \succ s$  and thus also  $s \in |a|(|t_1|, \dots, |t_n|)$ . Since  $s$  is not preceded by any term in  $\mathfrak{R}$  we get  $s \in |at_1 \dots t_n|$  by ( $B_3$ ).

# Proof of the Evaluation Lemma

## Case $(B_2)$

To handle the case of  $(B_2)$  let  $\tau = (\tau_1, \dots, \tau_n)$  be a basic type,  $a$  a closed IFT-term such that  $a \vdash \tau$  and  $t_1, \dots, t_n$  be closed IFT-terms such that  $t_i \vdash \tau_i$  for  $i = 1, \dots, n$  and  $t \in |at_1 \dots t_n|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |a|(|t_1|, \dots, |t_n|)$ . Since by induction hypothesis  $|a|$  is a foundation-predicate of type  $\tau$  and  $|t_i|$  a foundation-predicate of type  $\tau_i$  we obtain  $|a|(|t_1|, \dots, |t_n|)$  as a foundation-predicate of type 1. Thus every reduction chain  $\mathfrak{R}'$  for  $t'$  is finite. But then  $\mathfrak{R}$  is finite, too. Therefore  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is not strongly normal then  $\mathfrak{R}'$  is not strongly normal, too, and thus contains a  $\lambda$ -functional  $s' \in |a|(|t_1|, \dots, |t_n|)$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$ . If  $s' \notin \mathfrak{R}$  we get  $s' \succ s$  and thus also  $s \in |a|(|t_1|, \dots, |t_n|)$ . Since  $s$  is not preceded by any term in  $\mathfrak{R}$  we get  $s \in |at_1 \dots t_n|$  by  $(B_3)$ .

If  $s' \in \mathfrak{R}$  then there are no  $L$ -terms in  $\mathfrak{R}$  which precede  $s'$ . Hence  $s \in |at_1 \dots t_n|$  by  $(B_3)$ .

# Proof of the Evaluation Lemma

Case  $(B_3)$

To handle the case of  $(B_3)$  let  $t \in |A \rightarrow B|$ .

# Proof of the Evaluation Lemma

## Case $(B_3)$

To handle the case of  $(B_3)$  let  $t \in |A \rightarrow B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal.



# Proof of the Evaluation Lemma

## Case $(B_3)$

To handle the case of  $(B_3)$  let  $t \in |A \rightarrow B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right) \in |B|$  for all  $s \in |A|$ .

# Proof of the Evaluation Lemma

## Case $(B_3)$

To handle the case of  $(B_3)$  let  $t \in |A \rightarrow B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right) \in |B|$  for all  $s \in |A|$ . By induction hypothesis  $|B|$  is a foundation-predicate of type 1.

# Proof of the Evaluation Lemma

## Case $(B_3)$

To handle the case of  $(B_3)$  let  $t \in |A \rightarrow B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \binom{s}{x} \in |B|$  for all  $s \in |A|$ . By induction hypothesis  $|B|$  is a foundation-predicate of type 1. Therefore  $t' \binom{s}{x}$  is well-founded which implies that  $\lambda x. t'$  and thus also  $t$  is well-founded.

# Proof of the Evaluation Lemma

## Case $(B_3)$

To handle the case of  $(B_3)$  let  $t \in |A \rightarrow B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right) \in |B|$  for all  $s \in |A|$ . By induction hypothesis  $|B|$  is a foundation-predicate of type 1. Therefore  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right)$  is well-founded which implies that  $\lambda x. t'$  and thus also  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ .

# Proof of the Evaluation Lemma

## Case $(B_3)$

To handle the case of  $(B_3)$  let  $t \in |A \rightarrow B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right) \in |B|$  for all  $s \in |A|$ . By induction hypothesis  $|B|$  is a foundation-predicate of type 1. Therefore  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right)$  is well-founded which implies that  $\lambda x. t'$  and thus also  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is not strongly normal then  $\mathfrak{R}'$  is also not strongly normal and thus contains a  $\lambda$ -functional  $\lambda x. t'$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$  such that  $t' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right) \in |B|$  for all  $r \in |A|$ .

# Proof of the Evaluation Lemma

## Case ( $B_3$ )

To handle the case of ( $B_3$ ) let  $t \in |A \rightarrow B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right) \in |B|$  for all  $s \in |A|$ . By induction hypothesis  $|B|$  is a foundation-predicate of type 1. Therefore  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right)$  is well-founded which implies that  $\lambda x. t'$  and thus also  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is not strongly normal then  $\mathfrak{R}'$  is also not strongly normal and thus contains a  $\lambda$ -functional  $\lambda x. t'$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$  such that  $t' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right) \in |B|$  for all  $r \in |A|$ . If  $\lambda x. t' \notin \mathfrak{R}$  we get  $\lambda x. t' \succ s$  which implies that  $s$  is a functional  $\lambda x. s'$  such that  $t' \succ s'$ .

# Proof of the Evaluation Lemma

## Case ( $B_3$ )

To handle the case of ( $B_3$ ) let  $t \in |A \rightarrow B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x.t'$  such that  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right) \in |B|$  for all  $s \in |A|$ . By induction hypothesis  $|B|$  is a foundation-predicate of type 1. Therefore  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right)$  is well-founded which implies that  $\lambda x.t'$  and thus also  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is not strongly normal then  $\mathfrak{R}'$  is also not strongly normal and thus contains a  $\lambda$ -functional  $\lambda x.t'$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$  such that  $t' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right) \in |B|$  for all  $r \in |A|$ . If  $\lambda x.t' \notin \mathfrak{R}$  we get  $\lambda x.t' \succ s$  which implies that  $s$  is a functional  $\lambda x.s'$  such that  $t' \succ s'$ . This entails  $t' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right) \succ s' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right)$  for all  $r \in |A|$ .

# Proof of the Evaluation Lemma

## Case ( $B_3$ )

To handle the case of ( $B_3$ ) let  $t \in |A \rightarrow B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x.t'$  such that  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right) \in |B|$  for all  $s \in |A|$ . By induction hypothesis  $|B|$  is a foundation-predicate of type 1. Therefore  $t' \left( \begin{smallmatrix} s \\ x \end{smallmatrix} \right)$  is well-founded which implies that  $\lambda x.t'$  and thus also  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is not strongly normal then  $\mathfrak{R}'$  is also not strongly normal and thus contains a  $\lambda$ -functional  $\lambda x.t'$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$  such that  $t' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right) \in |B|$  for all  $r \in |A|$ . If  $\lambda x.t' \notin \mathfrak{R}$  we get  $\lambda x.t' \succ s$  which implies that  $s$  is a functional  $\lambda x.s'$  such that  $t' \succ s'$ . This entails  $t' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right) \succ s' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right)$  for all  $r \in |A|$ . Hence  $s' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right) \in |B|$  which implies  $\lambda x.s' \in |A \rightarrow B|$ .



### Case ( $B_3$ )

If  $\lambda x.t' \in \mathfrak{R}$  then there are no  $L$ -terms in  $\mathfrak{R}$  which precede  $\lambda x.t'$ . Hence  $s \in |A \rightarrow B|$ .

# Proof of the Evaluation Lemma

Case  $(B_4)$

To handle the case of  $(B_4)$  let  $t \in |(\forall x^\tau)B|$ .

# Proof of the Evaluation Lemma

## Case $(B_4)$

To handle the case of  $(B_4)$  let  $t \in |(\forall x^\tau)B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal.

# Proof of the Evaluation Lemma

## Case $(B_4)$

To handle the case of  $(B_4)$  let  $t \in |(\forall x^\tau)B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} u \\ x \end{smallmatrix} \right) \in |B \left( \begin{smallmatrix} \mathfrak{F} \\ x^\tau \end{smallmatrix} \right)|$  for all foundation-predicates  $\mathfrak{F}$  of type  $\tau$ .

# Proof of the Evaluation Lemma

## Case $(B_4)$

To handle the case of  $(B_4)$  let  $t \in |(\forall x^\tau)B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \left( \begin{smallmatrix} u \\ x \end{smallmatrix} \right) \in |B \left( \begin{smallmatrix} \mathfrak{F} \\ x^\tau \end{smallmatrix} \right)|$  for all foundation-predicates  $\mathfrak{F}$  of type  $\tau$ .

By induction hypothesis  $|B \left( \begin{smallmatrix} \mathfrak{F} \\ x^\tau \end{smallmatrix} \right)|$  is a foundation-predicate of type

1.

# Proof of the Evaluation Lemma

## Case $(B_4)$

To handle the case of  $(B_4)$  let  $t \in |(\forall x^\tau)B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \left( \frac{u}{x} \right) \in |B \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  for all foundation-predicates  $\mathfrak{F}$  of type  $\tau$ .

By induction hypothesis  $|B \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  is a foundation-predicate of type

1. This implies that  $t' \left( \frac{u}{x^\tau} \right)$  is well-founded which in turn implies that  $\lambda x. t'$  and therefore also  $t$  is well-founded.

# Proof of the Evaluation Lemma

## Case $(B_4)$

To handle the case of  $(B_4)$  let  $t \in |(\forall x^\tau)B|$ . Let  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $\lambda x. t'$  such that  $t' \left( \frac{u}{x} \right) \in |B \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  for all foundation-predicates  $\mathfrak{F}$  of type  $\tau$ .

By induction hypothesis  $|B \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  is a foundation-predicate of type

1. This implies that  $t' \left( \frac{u}{x^\tau} \right)$  is well-founded which in turn implies that  $\lambda x. t'$  and therefore also  $t$  is well-founded.

Now assume  $t \succ s$  and let  $\mathfrak{R}$  be a reduction chain for  $s$ . Then  $\mathfrak{R}$  is a subchain of a reduction chain  $\mathfrak{R}'$  for  $t$ . If  $\mathfrak{R}$  is not strongly normal then so is  $\mathfrak{R}'$  and thus contains a  $\lambda$ -functional  $\lambda x. t'$  which is not preceded by  $L$ -terms in  $\mathfrak{R}'$  such that  $t' \left( \frac{u}{x} \right) \in |B \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  for all foundation-predicates  $\mathfrak{F}$  of type  $\tau$ .

# Proof of the Evaluation Lemma

Case ( $B_4$ ) (continued)

If  $\lambda x. t' \notin \mathfrak{R}$  we get  $\lambda x. t' \succ s$  which implies that  $s$  is a functional  $\lambda x. s'$  such that  $t' \succ s'$ .



# Proof of the Evaluation Lemma

Case ( $B_4$ ) (continued)

If  $\lambda x.t' \notin \mathfrak{R}$  we get  $\lambda x.t' \succ s$  which implies that  $s$  is a functional  $\lambda x.s'$  such that  $t' \succ s'$ . This entails  $t' \left(\frac{u}{x}\right) \succ s' \left(\frac{u}{x}\right)$ .

# Proof of the Evaluation Lemma

## Case $(B_4)$ (continued)

If  $\lambda x. t' \notin \mathfrak{R}$  we get  $\lambda x. t' \succ s$  which implies that  $s$  is a functional  $\lambda x. s'$  such that  $t' \succ s'$ . This entails  $t' \left( \frac{u}{x} \right) \succ s' \left( \frac{u}{x} \right)$ . Hence  $s' \left( \frac{u}{x} \right) \in |B \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  which implies  $\lambda x. s' \in |(\forall x^\tau) B|$ .

# Proof of the Evaluation Lemma

## Case ( $B_4$ ) (continued)

If  $\lambda x. t' \notin \mathfrak{R}$  we get  $\lambda x. t' \succ s$  which implies that  $s$  is a functional  $\lambda x. s'$  such that  $t' \succ s'$ . This entails  $t' \left( \frac{u}{x} \right) \succ s' \left( \frac{u}{x} \right)$ . Hence

$s' \left( \frac{u}{x} \right) \in |B \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  which implies  $\lambda x. s' \in |(\forall x^\tau) B|$ .

If  $\lambda x. t' \in \mathfrak{R}$  then there are no  $L$ -terms in  $\mathfrak{R}$  which precede  $\lambda x. t'$ . Hence  $s \in |(\forall x^\tau) B|$ .

► Back

# Proof of the Evaluation Lemma

## Case ( $B_5$ )

In the case of ( $B_5$ ) we get  $|\mathfrak{F}^\tau \mathfrak{F}_1 \dots \mathfrak{F}_n|$  as a foundation-predicate of type 1. Therefore  $|\mathfrak{F}^\tau|$  maps tuples of foundation-predicates of adequate types to foundation-predicates of type 1. So  $|\mathfrak{F}^\tau|$  is a foundation-predicates of type  $\tau$ .

► Back

# Proof of the Evaluation Lemma

## Case $(B_6)$

In the case of  $(B_6)$  we get

$|\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\}|(\mathfrak{F}_1, \dots, \mathfrak{F}_n) := |A\left(\frac{\mathfrak{F}_1}{x^{\tau_1}}, \dots, \frac{\mathfrak{F}_n}{x^{\tau_n}}\right)|$  which is a foundation-predicate of type 1 by induction hypothesis.

# Proof of the Evaluation Lemma

## Case ( $B_6$ )

In the case of ( $B_6$ ) we get

$|\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\}|(\mathfrak{F}_1, \dots, \mathfrak{F}_n) := |A\left(\frac{\mathfrak{F}_1}{x^{\tau_1}}, \dots, \frac{\mathfrak{F}_n}{x^{\tau_n}}\right)|$  which is a foundation-predicate of type 1 by induction hypothesis. Therefore  $|\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\}|$  maps tuples of foundation-predicates of adequate types to foundation-predicates of type 1. So  $|\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\}|$  is a foundation-predicates of type  $\tau$ .

► Back

# Proof of the Lemma

We distinguish the following cases:

(Atomic case) Let  $A$  be the formula  $(at_1 \dots t_n)$  for  $n \geq 0$  (If  $n = 0$  we assume that  $a$  is of type 1).

# Proof of the Lemma

We distinguish the following cases:

(Atomic case) Let  $A$  be the formula  $(at_1 \dots t_n)$  for  $n \geq 0$  (If  $n = 0$  we assume that  $a$  is of type 1). Let  $t$  be a  $\lambda$ -functional and  $\mathfrak{R}$  be a reduction chain for  $t$  which is not strongly normal but contains an element  $t' \in |a|(|t_1|, \dots, |t_n|)$ . Let  $\mathfrak{R}'$  be the subchain which starts with  $t'$ . Then  $\mathfrak{R}'$  is not strongly normal. Therefore it contains an element  $t'' \in \mathfrak{R}'$  which is not preceded by an  $L$ -functional in  $\mathfrak{R}'$ . But then  $t''$  is also not preceded by an  $L$ -functional in  $\mathfrak{R}$ . Hence  $t \in |at_1 \dots t_n|$ .

► Back



# Proof of the Lemma

(continued)

( $\rightarrow$  case) Now assume that  $A$  is a formula  $B \rightarrow C$  and let  $\mathfrak{R}$  be a reduction chain of a  $\lambda$ -functional  $t$  which is not strongly normal.

# Proof of the Lemma

(continued)

( $\rightarrow$  case) Now assume that  $A$  is a formula  $B \rightarrow C$  and let  $\mathfrak{R}$  be a reduction chain of a  $\lambda$ -functional  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |B \rightarrow C|$  which is not preceded by an  $L$ -functional in  $\mathfrak{R}$ .

# Proof of the Lemma

(continued)

( $\rightarrow$  case) Now assume that  $A$  is a formula  $B \rightarrow C$  and let  $\mathfrak{R}$  be a reduction chain of a  $\lambda$ -functional  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |B \rightarrow C|$  which is not preceded by an  $L$ -functional in  $\mathfrak{R}$ . Let  $\mathfrak{R}'$  be the subchain of  $\mathfrak{R}$  which starts with  $t'$ .

# Proof of the Lemma

(continued)

( $\rightarrow$  case) Now assume that  $A$  is a formula  $B \rightarrow C$  and let  $\mathfrak{R}$  be a reduction chain of a  $\lambda$ -functional  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |B \rightarrow C|$  which is not preceded by an  $L$ -functional in  $\mathfrak{R}$ . Let  $\mathfrak{R}'$  be the subchain of  $\mathfrak{R}$  which starts with  $t'$ . Then  $\mathfrak{R}'$  contains a first functional  $\lambda x. t''$  such that  $t'' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right) \in |C|$  for all  $r \in |B|$ .

# Proof of the Lemma

(continued)

( $\rightarrow$  case) Now assume that  $A$  is a formula  $B \rightarrow C$  and let  $\mathfrak{R}$  be a reduction chain of a  $\lambda$ -functional  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |B \rightarrow C|$  which is not preceded by an  $L$ -functional in  $\mathfrak{R}$ . Let  $\mathfrak{R}'$  be the subchain of  $\mathfrak{R}$  which starts with  $t'$ . Then  $\mathfrak{R}'$  contains a first functional  $\lambda x.t''$  such that  $t'' \left( \begin{smallmatrix} r \\ x \end{smallmatrix} \right) \in |C|$  for all  $r \in |B|$ . But then  $\lambda x.t''$  is also the first functional with this property in  $\mathfrak{R}$ . Hence  $t \in |B \rightarrow C|$ .

► Back

# Proof of the Lemma

Continued

( $\forall$ )–case In the last case assume that  $A$  is a sentence  $(\forall x^\tau)B$  and let  $\mathfrak{R}$  be a reduction chain of  $t$  which is not strongly normal.

# Proof of the Lemma

Continued

( $\forall$ )–case In the last case assume that  $A$  is a sentence  $(\forall x^\tau)B$  and let  $\mathfrak{R}$  be a reduction chain of  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |(\forall x^\tau)B|$ .

# Proof of the Lemma

Continued

( $\forall$ )–case In the last case assume that  $A$  is a sentence  $(\forall x^\tau)B$  and let  $\mathfrak{R}$  be a reduction chain of  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |(\forall x^\tau)B|$ . Let  $\mathfrak{R}'$  be the subchain of  $\mathfrak{R}$  starting with  $t'$ . Then  $\mathfrak{R}'$  is not strongly normal and therefore contains a functional  $\lambda x. t''$  which is not preceded in  $\mathfrak{R}'$  by  $L$ –functional such that  $t'' \left( \begin{smallmatrix} u \\ x \end{smallmatrix} \right) \in |B \left( \begin{smallmatrix} \mathfrak{F} \\ x^\tau \end{smallmatrix} \right)|$  for all foundation–predicates  $\mathfrak{F}$  of type  $\tau$ .



# Proof of the Lemma

Continued

( $\forall$ )–case In the last case assume that  $A$  is a sentence  $(\forall x^\tau)B$  and let  $\mathfrak{R}$  be a reduction chain of  $t$  which is not strongly normal. Then  $\mathfrak{R}$  contains an element  $t' \in |(\forall x^\tau)B|$ . Let  $\mathfrak{R}'$  be the subchain of  $\mathfrak{R}$  starting with  $t'$ . Then  $\mathfrak{R}'$  is not strongly normal and therefore contains a functional  $\lambda x. t''$  which is not preceded in  $\mathfrak{R}'$  by  $L$ –functional such that  $t'' \left( \begin{smallmatrix} u \\ x \end{smallmatrix} \right) \in |B \left( \begin{smallmatrix} \mathfrak{F} \\ x^\tau \end{smallmatrix} \right)|$  for all foundation–predicates  $\mathfrak{F}$  of type  $\tau$ . But then  $\lambda x. t''$  is also the first functional with this property in  $\mathfrak{R}$  which implies  $t \in |(\forall x^\tau)B|$ . □

► Back

# Proof of the Lemma

Let  $a \vdash \tau$ . Firstly we observe that  $|a|$  as well as  $\|a\|$  are foundation-predicates predicates of type  $\tau$ .

# Proof of the Lemma

Let  $a \vdash \tau$ . Firstly we observe that  $|a|$  as well as  $\|a\|$  are foundation–predicates predicates of type  $\tau$ . We prove the lemma by induction on the complexity of  $\tau$  and distinguish the following cases:

# Proof of the Lemma

Let  $a \vdash \tau$ . Firstly we observe that  $|a|$  as well as  $\|a\|$  are foundation–predicates predicates of type  $\tau$ . We prove the lemma by induction on the complexity of  $\tau$  and distinguish the following cases:  $\tau = 0$ . Then  $|a| = a$  which trivially entails  $\|a\| = |a| = a$ .

# Proof of the Lemma

Let  $a \vdash \tau$ . Firstly we observe that  $|a|$  as well as  $\|a\|$  are foundation-predicates predicates of type  $\tau$ . We prove the lemma by induction on the complexity of  $\tau$  and distinguish the following cases:  
 $\tau = 0$ . Then  $|a| = a$  which trivially entails  $\|a\| = |a| = a$ .

$\tau = 1$ . If  $t \in |a|$  then every reduction chain of  $t$  contains an element in  $|a|$ . Hence  $t \in \|a\|$ .

# Proof of the Lemma

Let  $a \vdash \tau$ . Firstly we observe that  $|a|$  as well as  $\|a\|$  are foundation-predicates predicates of type  $\tau$ . We prove the lemma by induction on the complexity of  $\tau$  and distinguish the following cases:  
 $\tau = 0$ . Then  $|a| = a$  which trivially entails  $\|a\| = |a| = a$ .

$\tau = 1$ . If  $t \in |a|$  then every reduction chain of  $t$  contains an element in  $|a|$ . Hence  $t \in \|a\|$ .

If vice versa  $t \in \|a\|$  then every reduction chain  $\mathfrak{R}$  of  $t$  which is not strongly normal contains an element  $t'$  which is not preceded in  $\mathfrak{R}$  by an  $L$ -term. By the previous lemma we then get  $t \in |a|$ .

# Proof of the Lemma

Let  $a \vdash \tau$ . Firstly we observe that  $|a|$  as well as  $\|a\|$  are foundation–predicates predicates of type  $\tau$ . We prove the lemma by induction on the complexity of  $\tau$  and distinguish the following cases:  
 $\tau = 0$ . Then  $|a| = a$  which trivially entails  $\|a\| = |a| = a$ .

$\tau = 1$ . If  $t \in |a|$  then every reduction chain of  $t$  contains an element in  $|a|$ . Hence  $t \in \|a\|$ .

If vice versa  $t \in \|a\|$  then every reduction chain  $\mathfrak{R}$  of  $t$  which is not strongly normal contains an element  $t'$  which is not preceeded in  $\mathfrak{R}$  by an  $L$ –term. By the previous lemma we then get  $t \in |a|$ .

$\tau = (\tau_1, \dots, \tau_n)$ . Then  $a$  is a name  $\mathfrak{F}^\tau$  for a foundation–predicate or  $a$  is a term  $\{((x^{\tau_1} \dots x^{\tau_n})) \mid A\}$ . In both cases  $|a|$  is a foundation–predicate of type  $\tau$ . In the first case we get for any tuple  $\mathfrak{F}_1, \dots, \mathfrak{F}_n$  of foundation–predicates of adequate types

$$\begin{aligned} \|\mathfrak{F}^\tau\|(\mathfrak{F}_1, \dots, \mathfrak{F}_n) &= \|\mathfrak{F}^\tau\|(\mathfrak{F}_1, \dots, \mathfrak{F}_n) = \|\mathfrak{F}^\tau(\mathfrak{F}_1, \dots, \mathfrak{F}_n)\| = \\ &|\mathfrak{F}^\tau(\mathfrak{F}_1, \dots, \mathfrak{F}_n)| = |\mathfrak{F}^\tau|(\mathfrak{F}_1, \dots, \mathfrak{F}_n). \end{aligned}$$



# Proof of the Lemma

(continued)

In the second case we get  $\|\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\}(\mathfrak{F}_1, \dots, \mathfrak{F}_n) =$   
 $\|\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\}(\mathfrak{F}_1, \dots, \mathfrak{F}_n)\| = \|A\left(\begin{smallmatrix} \mathfrak{F}_1, \dots, \mathfrak{F}_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix}\right)\| =$   
 $|A\left(\begin{smallmatrix} \mathfrak{F}_1, \dots, \mathfrak{F}_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix}\right)| = \|\{(x^{\tau_1} \dots x^{\tau_n}) \mid A\}(\mathfrak{F}_1, \dots, \mathfrak{F}_n).$

► Back



# Proof of the Main Lemma

We prove the lemma by induction on the length of  $a$ .

# Proof of the Main Lemma

We prove the lemma by induction on the length of  $a$ .

Firstly assume  $a = y^A$ . Then  $a \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right) = y^{A \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)} = y^{B_i}$

for some  $i \in \{1, \dots, m\}$ .

# Proof of the Main Lemma

We prove the lemma by induction on the length of  $a$ .

Firstly assume  $a = y^A$ . Then  $a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) = y^A \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) = y^{B_i}$  for some  $i \in \{1, \dots, m\}$ . Hence  $a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) = y$ , which for  $b \in |B_i|$  implies  $a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) = b \in |B_i| = |A \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|$ .

▶ Back

# Proof of the Main Lemma

continued

Assume that  $a = \lambda x^\sigma . d$ .

# Proof of the Main Lemma

continued

Assume that  $a = \lambda x^\sigma. d$ .

If  $\sigma$  is a basic type  $\tau$  then  $A$  is a formula  $(\forall x^\tau)C$  and  $d \vdash C$  with  $FV(d) \setminus \overline{FV(d)} = \{x^{\tau_1}, \dots, x^{\tau_n}, x^\tau\}$ .

# Proof of the Main Lemma

continued

Assume that  $a = \lambda x^\sigma. d$ .

If  $\sigma$  is a basic type  $\tau$  then  $A$  is a formula  $(\forall x^\tau)C$  and  $d \vdash C$  with

$FV(d) \setminus \overline{FV(d)} = \{x^{\tau_1}, \dots, x^{\tau_n}, x^\tau\}$ . We have to show that

$$\frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) \left( \frac{u}{x} \right) \in |C \left( \frac{c_1, \dots, c_m}{x^{\tau_1}, \dots, x^{\tau_m}} \right) \left( \frac{\mathfrak{F}}{x^\tau} \right)|}{\text{holds true}}$$

for every foundation-predicate  $\mathfrak{F}$  of type  $\tau$ .

# Proof of the Main Lemma

continued

Assume that  $a = \lambda x^\sigma. d$ .

If  $\sigma$  is a basic type  $\tau$  then  $A$  is a formula  $(\forall x^\tau)C$  and  $d \vdash C$  with

$FV(d) \setminus \overline{FV(d)} = \{x^{\tau_1}, \dots, x^{\tau_n}, x^\tau\}$ . We have to show that

$\frac{}{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) \left( \frac{u}{x} \right) \in |C \left( \frac{c_1, \dots, c_m}{x^{\tau_1}, \dots, x^{\tau_m}} \right) \left( \frac{\mathfrak{F}}{x^\tau} \right)|}$  holds true

for every foundation–predicate  $\mathfrak{F}$  of type  $\tau$ .

By the induction hypothesis we have

$\frac{}{d \left( \frac{c_1, \dots, c_m}{x^{\tau_1}, \dots, x^{\tau_m}} \right) \left( \frac{\mathfrak{F}}{x^\tau} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) \in |C \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{\mathfrak{F}}{x^\tau} \right)|}$

# Proof of the Main Lemma

continued

Assume that  $a = \lambda x^\sigma . d$ .

If  $\sigma$  is a basic type  $\tau$  then  $A$  is a formula  $(\forall x^\tau)C$  and  $d \vdash C$  with

$FV(d) \setminus \overline{FV(d)} = \{x^{\tau_1}, \dots, x^{\tau_n}, x^\tau\}$ . We have to show that

$\overline{d \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right) \left( \begin{smallmatrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{smallmatrix} \right) \left( \begin{smallmatrix} u \\ x \end{smallmatrix} \right)} \in |C \left( \begin{smallmatrix} c_1, \dots, c_m \\ x^{\tau_1}, \dots, x^{\tau_m} \end{smallmatrix} \right) (\mathfrak{F})_{x^\tau}|$  holds true

for every foundation–predicate  $\mathfrak{F}$  of type  $\tau$ .

By the induction hypothesis we have

$\overline{d \left( \begin{smallmatrix} c_1, \dots, c_m \\ x^{\tau_1}, \dots, x^{\tau_m} \end{smallmatrix} \right) (\mathfrak{F})_{x^\tau} \left( \begin{smallmatrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{smallmatrix} \right)} \in |C \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right) (\mathfrak{F})_{x^\tau}|$

and compute easily  $\overline{d \left( \begin{smallmatrix} c_1, \dots, c_m \\ x^{\tau_1}, \dots, x^{\tau_m} \end{smallmatrix} \right) (\mathfrak{F})_{x^\tau} \left( \begin{smallmatrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{smallmatrix} \right)} =$

$\overline{d \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right) \left( \begin{smallmatrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{smallmatrix} \right) \left( \begin{smallmatrix} u \\ x \end{smallmatrix} \right)}$ .

► Back



# Proof of the Main Lemma

continued

Assume that  $a = \lambda x^\sigma . d$ .

# Proof of the Main Lemma

continued

Assume that  $a = \lambda x^\sigma . d$ .

If  $\sigma$  is a formula  $B$  then  $A$  is a formula  $B \rightarrow C$  and  $d \vdash C$ . Then  
 $FV(d \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)) = \{y^{B_1}, \dots, y^{B_n}, x^B\}$ .

# Proof of the Main Lemma

continued

Assume that  $a = \lambda x^\sigma . d$ .

If  $\sigma$  is a formula  $B$  then  $A$  is a formula  $B \rightarrow C$  and  $d \vdash C$ . Then  $FV(d \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)) = \{y^{B_1}, \dots, y^{B_n}, x^B\}$ . We have to show that

$\overline{d \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)} \left( \begin{smallmatrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{smallmatrix} \right) \left( \begin{smallmatrix} b \\ x \end{smallmatrix} \right) \in |C \left( \begin{smallmatrix} c_1, \dots, c_m \\ x^{\tau_1}, \dots, x^{\tau_m} \end{smallmatrix} \right)|$  holds true for all  $b \in |B|$ .

# Proof of the Main Lemma

continued

Assume that  $a = \lambda x^\sigma . d$ .

If  $\sigma$  is a formula  $B$  then  $A$  is a formula  $B \rightarrow C$  and  $d \vdash C$ . Then  $FV(d \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)) = \{y^{B_1}, \dots, y^{B_n}, x^B\}$ . We have to show that

$\overline{d \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right) \left( \begin{smallmatrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{smallmatrix} \right) \left( \begin{smallmatrix} b \\ x \end{smallmatrix} \right)} \in |C \left( \begin{smallmatrix} c_1, \dots, c_m \\ x^{\tau_1}, \dots, x^{\tau_m} \end{smallmatrix} \right)|$  holds true for all  $b \in |B|$ .

But this is in fact the induction hypothesis. [▶ Back](#)

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash (\forall x^\tau)D$  and  $e \vdash \tau$  for a basic type  $\tau$ .

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash (\forall x^\tau)D$  and  $e \vdash \tau$  for a basic type  $\tau$ .

Then  $A$  is the formula  $D \left( \frac{e}{x^\tau} \right)$ ,  $\bar{e} = u$  and

$$\frac{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{\left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) u}{\left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \right)} =$$

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash (\forall x^\tau)D$  and  $e \vdash \tau$  for a basic type  $\tau$ .

Then  $A$  is the formula  $D \left( \frac{e}{x^\tau} \right)$ ,  $\bar{e} = u$  and

$$\begin{aligned} \frac{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{\left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{u} \right)} &= \left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) u}{\left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \right) = \\ &= \left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{u} \right). \quad \text{By induction hypothesis we have} \\ \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{\left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} &\in |(\forall x^\tau)D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|. \end{aligned}$$

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash (\forall x^\tau)D$  and  $e \vdash \tau$  for a basic type  $\tau$ .

Then  $A$  is the formula  $D \left( \frac{e}{x^\tau} \right)$ ,  $\bar{e} = u$  and

$$\frac{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{\left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{u} \right)} = \left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) u}{\left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \right) =$$

$$\left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{u} \right) u. \quad \text{By induction hypothesis we have}$$

$$\frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{u} \in |(\forall x^\tau)D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|. \quad \text{Hence}$$

$$\frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{u} \text{ is well-founded.}$$



# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash (\forall x^\tau)D$  and  $e \vdash \tau$  for a basic type  $\tau$ .

Then  $A$  is the formula  $D \left( \frac{e}{x^\tau} \right)$ ,  $\bar{e} = u$  and

$$\frac{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{\left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{u} \right)} = \left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) u}{\left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \right) =$$

$$\left( \frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{u} \right) u. \quad \text{By induction hypothesis we have}$$

$$\frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{\left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \in |(\forall x^\tau)D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|. \quad \text{Hence}$$

$$\frac{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)}{\left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \text{ is well-founded. Let } \mathfrak{R} = \langle r_i \mid i \in I \rangle \text{ be a}$$

$$\text{reduction chain of } a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right).$$

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash (\forall x^\tau)D$  and  $e \vdash \tau$  for a basic type  $\tau$ .

Then  $A$  is the formula  $D \left( \frac{e}{x^\tau} \right)$ ,  $\bar{e} = u$  and

$$\overline{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} = \overline{\left( d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) u \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} = \left( \overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \right) u. \quad \text{By induction hypothesis we have}$$

$$\overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \in |(\forall x^\tau)D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|. \quad \text{Hence}$$

$$\overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \text{ is well-founded. Let } \mathfrak{R} = \langle r_i \mid i \in I \rangle \text{ be a}$$

reduction chain of  $a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)$ . If  $\mathfrak{R}$  does not contain

a conversion according to  $R_0$  then  $\mathfrak{R}$  is strongly normal by [▶ Lemma I](#).

# Proof of the Main Lemma

continued

Otherwise let  $k$  be the least index such that  $r_k \succ r_{k+1}$  is a reduction according to  $R_0$ . Then  $r_k = (\lambda x. r')u$  and

$$d \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}} \left( b_1, \dots, b_m \right)_{y_1, \dots, y_m} \succ \lambda x. r'.$$

# Proof of the Main Lemma

continued

Otherwise let  $k$  be the least index such that  $r_k \succ r_{k+1}$  is a reduction according to  $R_0$ . Then  $r_k = (\lambda x.r')u$  and

$\overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \succ \lambda x.r'$ . Hence

$\lambda x.r' \in |(\forall x^\tau) D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) |$

# Proof of the Main Lemma

continued

Otherwise let  $k$  be the least index such that  $r_k \succ r_{k+1}$  is a reduction according to  $R_0$ . Then  $r_k = (\lambda x. r')u$  and

$\overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \succ \lambda x. r'$ . Hence

$\lambda x. r' \in |(\forall x^\tau) D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|$  which implies that

$r' \left( \frac{u}{x} \right) \in |D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  for all foundation-predicates  $\mathfrak{F}$  of type  $\tau$ .

# Proof of the Main Lemma

continued

Otherwise let  $k$  be the least index such that  $r_k \succ r_{k+1}$  is a reduction according to  $R_0$ . Then  $r_k = (\lambda x. r')u$  and

$\overline{d \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right) \left( \begin{smallmatrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{smallmatrix} \right)} \succ \lambda x. r'$ . Hence

$\lambda x. r' \in |(\forall x^\tau) D \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)|$  which implies that

$r' \left( \begin{smallmatrix} u \\ x \end{smallmatrix} \right) \in |D \left( \begin{smallmatrix} c_1, \dots, c_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right) (\mathfrak{F})|$  for all foundation-predicates  $\mathfrak{F}$  of type  $\tau$ . But  $|e|$  is a foundation-predicate of type  $\tau$ .

# Proof of the Main Lemma

continued

Otherwise let  $k$  be the least index such that  $r_k \succ r_{k+1}$  is a reduction according to  $R_0$ . Then  $r_k = (\lambda x. r')u$  and

$\overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \succ \lambda x. r'$ . Hence

$\lambda x. r' \in |(\forall x^\tau) D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|$  which implies that

$r' \left( \frac{u}{x} \right) \in |D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  for all foundation-predicates  $\mathfrak{F}$  of type  $\tau$ . But  $|e|$  is a foundation-predicate of type  $\tau$ . Therefore

we get  $r_{k+1} = r' \left( \frac{u}{x} \right) \in |D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{|e|}{x^\tau} \right)| =$

$|D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{e}{x^\tau} \right)|$ .

# Proof of the Main Lemma

continued

Otherwise let  $k$  be the least index such that  $r_k \succ r_{k+1}$  is a reduction according to  $R_0$ . Then  $r_k = (\lambda x. r')u$  and

$\overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \succ \lambda x. r'$ . Hence

$\lambda x. r' \in |(\forall x^\tau) D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|$  which implies that

$r' \left( \frac{u}{x} \right) \in |D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{\mathfrak{F}}{x^\tau} \right)|$  for all foundation-predicates  $\mathfrak{F}$  of type  $\tau$ . But  $|e|$  is a foundation-predicate of type  $\tau$ . Therefore

we get  $r_{k+1} = r' \left( \frac{u}{x} \right) \in |D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{|e|}{x^\tau} \right)| =$

$|D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{e}{x^\tau} \right)|$ . Since  $r_{k+1}$  is not preceded by

$L$ -functionals in  $\mathfrak{R}$  we obtain

$\overline{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right)} \in |D \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right) \left( \frac{e}{x^\tau} \right)|$ .

► Back



# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash B \rightarrow A$  and  $e \vdash B$  for a formula  $B$ .

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash B \rightarrow A$  and  $e \vdash B$  for a formula  $B$ .

By the induction hypothesis we get

$$d' := \overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) \in |(B \rightarrow A) \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)| \text{ and}$$

$$e' := \overline{e \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_n}{y_1, \dots, y_n} \right) \in |B \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|$$

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash B \rightarrow A$  and  $e \vdash B$  for a formula  $B$ .

By the induction hypothesis we get

$$d' := \overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) \in |(B \rightarrow A) \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)| \text{ and}$$

$$e' := \overline{e \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_n}{y_1, \dots, y_n} \right) \in |B \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)| \text{ and}$$

$$a' := \overline{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) = (d'e').$$

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash B \rightarrow A$  and  $e \vdash B$  for a formula  $B$ .

By the induction hypothesis we get

$$d' := \overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) \in |(B \rightarrow A) \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)| \text{ and}$$

$$e' := \overline{e \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_n}{y_1, \dots, y_n} \right) \in |B \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)| \text{ and}$$

$$a' := \overline{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) = (d'e').$$

Let  $\Re$  be a reduction chain of  $a'$ .

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash B \rightarrow A$  and  $e \vdash B$  for a formula  $B$ .

By the induction hypothesis we get

$$d' := \overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) \in |(B \rightarrow A) \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)| \text{ and}$$

$$e' := \overline{e \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_n}{y_1, \dots, y_n} \right) \in |B \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)| \text{ and}$$

$$a' := \overline{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) = (d' e').$$

Let  $\mathfrak{R}$  be a reduction chain of  $a'$ . If  $\mathfrak{R}$  does not contain conversions according to  $R_0$  then  $\mathfrak{R}$  is strongly normal by [▶ Lemma 1](#) since  $d'$  and  $e'$  are well-founded.

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash B \rightarrow A$  and  $e \vdash B$  for a formula  $B$ .

By the induction hypothesis we get

$$d' := d \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}} \left( b_1, \dots, b_m \right)_{y_1, \dots, y_m} \in |(B \rightarrow A) \left( c_1, \dots, c_n \right)_{x^{\tau_1}, \dots, x^{\tau_n}}| \text{ and}$$

$$e' := e \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}} \left( b_1, \dots, b_n \right)_{y_1, \dots, y_n} \in |B \left( c_1, \dots, c_n \right)_{x^{\tau_1}, \dots, x^{\tau_n}}| \text{ and}$$

$$a' := a \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}} \left( b_1, \dots, b_m \right)_{y_1, \dots, y_m} = (d' e').$$

Let  $\mathfrak{R}$  be a reduction chain of  $a'$ . If  $\mathfrak{R}$  does not contain conversions according to  $R_0$  then  $\mathfrak{R}$  is strongly normal by [▶ Lemma I](#) since  $d'$  and  $e'$  are well-founded. If  $\mathfrak{R}$  is not strongly normal then it contains an element with minimal index  $k$  such that  $r_k \succ_1 r_{k+1}$  is a conversion according to  $R_0$ . Then  $r_k$  is of the form  $(\lambda x. r')s$  such that  $d' \succ \lambda x. r'$  and  $e' \succ s$ .

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash B \rightarrow A$  and  $e \vdash B$  for a formula  $B$ .

By the induction hypothesis we get

$$d' := \overline{d \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) \in |(B \rightarrow A) \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)| \text{ and}$$

$$e' := \overline{e \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_n}{y_1, \dots, y_n} \right) \in |B \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)| \text{ and}$$

$$a' := \overline{a \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)} \left( \frac{b_1, \dots, b_m}{y_1, \dots, y_m} \right) = (d' e').$$

Let  $\mathfrak{R}$  be a reduction chain of  $a'$ . If  $\mathfrak{R}$  does not contain conversions according to  $R_0$  then  $\mathfrak{R}$  is strongly normal by [Lemma I](#) since  $d'$  and  $e'$  are well-founded. If  $\mathfrak{R}$  is not strongly normal then it contains an element with minimal index  $k$  such that  $r_k \succ_1 r_{k+1}$  is a conversion according to  $R_0$ . Then  $r_k$  is of the form  $(\lambda x. r')s$  such that  $d' \succ \lambda x. r'$  and  $e' \succ s$ . Hence  $\lambda x. r' \in |(B \rightarrow A) \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|$  and  $s \in |B \left( \frac{c_1, \dots, c_n}{x^{\tau_1}, \dots, x^{\tau_n}} \right)|$

# Proof of the Main Lemma

continued

Assume that  $a = (de)$  such that  $d \vdash B \rightarrow A$  and  $e \vdash B$  for a formula  $B$ .

By the induction hypothesis we get

$$d' := d \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}} \left( \begin{matrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{matrix} \right) \in |(B \rightarrow A) \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}}| \text{ and}$$

$$e' := e \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}} \left( \begin{matrix} b_1, \dots, b_n \\ y_1, \dots, y_n \end{matrix} \right) \in |B \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}}| \text{ and}$$

$$a' := a \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}} \left( \begin{matrix} b_1, \dots, b_m \\ y_1, \dots, y_m \end{matrix} \right) = (d' e').$$

Let  $\Re$  be a reduction chain of  $a'$ . If  $\Re$  does not contain conversions

according to  $R_0$  then  $\Re$  is strongly normal by [Lemma 1](#) since  $d'$  and  $e'$  are well-founded. If  $\Re$  is not strongly normal then it contains an element with minimal index  $k$  such that  $r_k \succ_1 r_{k+1}$  is a conversion according to  $R_0$ . Then  $r_k$  is of the form  $(\lambda x. r')s$  such that  $d' \succ \lambda x. r'$  and  $e' \succ s$ .

Hence  $\lambda x. r' \in |(B \rightarrow A) \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}}|$  and  $s \in |B \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}}|$  and therefore  $r' \left( \begin{matrix} s \\ x \end{matrix} \right) \in |A \left( \overline{c_1, \dots, c_n} \right)_{x^{\tau_1}, \dots, x^{\tau_n}}|$ .



# Proof of the Main Lemma

continued

But  $r_{k+1}$  is the term  $r' \binom{s}{x}$ . Since  $r_{k+1}$  is not preceded by  $L$ -functionals in  $\mathfrak{R}$  we have  $a' \in |A \left( \begin{smallmatrix} b_1, \dots, b_n \\ x^{\tau_1}, \dots, x^{\tau_n} \end{smallmatrix} \right)|$ . □

► Back



J.-Y. GIRARD, *Une extension de l'interpretation de Gödel a l'analyse et son application a l'elimination des coupures dans l'analyse et la theorie des types*, **Proceedings of the 2nd Scandinavian logic symposium** (J. E. Fenstad, editor), Studies in Logic and the Foundations of Mathematics, vol. 63, North-Holland Publishing Company, Amsterdam, 1971, pp. 63–92.



W. POHLERS, *Ein starker Normalisationssatz für die intuitionistische Typentheorie*, **Manuscripta Mathematica**, vol. 8 (1973), pp. 371–387.



D. PRAWITZ, **Natural deduction. A proof-theoretical study**, Almqvist & Wiksell Foerlag, Stockholm, 1965.



K. SCHÜTTE, *Syntactical and semantical properties of simple type theory*, **Journal of Symbolic Logic**, vol. 25 (1960), pp. 305–326.



G. TAKEUTI, *On a generalized logic calculus*, **Japanese Journal of Mathematics**, vol. 24 (1953), pp. 149–156.