

Magnetic Transport Along Translationally Invariant Obstacles

A Bachelor's Thesis Defense

Michal Grňo

September 10, 2021

Magnetic Transport Along Translationally Invariant Obstacles

Magnetic Transport Along Translationally Invariant Obstacles

A magnetic Hamiltonian

$$H_0 = (-i\vec{\nabla} + \vec{A})^2,$$

Magnetic Transport Along Translationally Invariant Obstacles

A magnetic Hamiltonian

$$H_0 = (-i\vec{\nabla} + \vec{A})^2,$$

$$\vec{\nabla} \times \vec{A} = \vec{B}_0 = \text{const.}$$

Magnetic Transport Along Translationally Invariant Obstacles

A magnetic Hamiltonian

$$H_0 = (-i\vec{\nabla} + \vec{A})^2,$$

$$\vec{\nabla} \times \vec{A} = \vec{B}_0 = \text{const.}$$

restricted to a 2D plane orthogonal to \vec{B}_0 .

Magnetic Transport Along Translationally Invariant Obstacles

A magnetic Hamiltonian

$$H_0 = (-i\vec{\nabla} + \vec{A})^2,$$

$$\vec{\nabla} \times \vec{A} = \vec{B}_0 = \text{const.}$$

restricted to a 2D plane orthogonal to \vec{B}_0 .

This is the *Landau Hamiltonian*.

Magnetic Transport Along Translationally Invariant Obstacles

Magnetic Transport Along Translationally Invariant Obstacles

- potential obstacle: $H = H_0 + V(x, y)$

Magnetic Transport Along Translationally Invariant Obstacles

- potential obstacle: $H = H_0 + V(x, y)$
- magnetic obstacle: $\vec{B} = \vec{B}_0 + \vec{b}(x, y)$

Magnetic Transport Along Translationally Invariant Obstacles

- potential obstacle: $H = H_0 + V(x, y)$
- magnetic obstacle: $\vec{B} = \vec{B}_0 + \vec{b}(x, y)$, $\vec{b} \parallel \vec{B}_0$

Magnetic Transport Along Translationally Invariant Obstacles

- potential obstacle: $H = H_0 + V(x, y)$
- magnetic obstacle: $\vec{B} = \vec{B}_0 + \vec{b}(x, y)$, $\vec{b} \parallel \vec{B}_0$
- geometric obstacle:
 - ▶ the system is not restricted to a plane, but to a **thin layer**

Magnetic Transport Along Translationally Invariant Obstacles

- potential obstacle: $H = H_0 + V(x, y)$
- magnetic obstacle: $\vec{B} = \vec{B}_0 + \vec{b}(x, y)$, $\vec{b} \parallel \vec{B}_0$
- geometric obstacle:
 - ▶ the system is not restricted to a plane, but to a **thin layer**
 - ▶ the layer is smoothly **bent**

Magnetic Transport Along Translationally Invariant Obstacles

- potential obstacle: $H = H_0 + V(x, y)$
- magnetic obstacle: $\vec{B} = \vec{B}_0 + \vec{b}(x, y)$, $\vec{b} \parallel \vec{B}_0$
- geometric obstacle:
 - ▶ the system is not restricted to a plane, but to a **thin layer**
 - ▶ the layer is smoothly **bent**
 - ▶ Dirichlet boundary is assumed


Magnetic Transport Along Translationally Invariant Obstacles

- potential obstacle: $H = H_0 + V(x)$
- magnetic obstacle: $\vec{B} = \vec{B}_0 + \vec{b}(x)$, $\vec{b} \parallel \vec{B}_0$
- geometric obstacle:
 - ▶ the system is not restricted to a plane, but to a thin layer
 - ▶ the layer is smoothly bent and invariant under translation $y \mapsto y + c$
 - ▶ Dirichlet boundary is assumed

Magnetic Transport Along Translationally Invariant Obstacles


Magnetic Transport Along Translationally Invariant Obstacles

Classically:




Magnetic Transport Along Translationally Invariant Obstacles

Classically:



Magnetic Transport Along Translationally Invariant Obstacles

Classically:



Magnetic Transport Along Translationally Invariant Obstacles

Quantum Mechanics:

Magnetic Transport Along Translationally Invariant Obstacles

Quantum Mechanics:

- Spectrum of H is pure point

Magnetic Transport Along Translationally Invariant Obstacles

Quantum Mechanics:

- Spectrum of H is pure point
 ↔ there is a basis consisting of stationary states

Magnetic Transport Along Translationally Invariant Obstacles

Quantum Mechanics:

- Spectrum of H is pure point
 - ↔ there is a basis consisting of stationary states
 - ↔ time evolution is trivial

Magnetic Transport Along Translationally Invariant Obstacles

Quantum Mechanics:

- Spectrum of H is pure point
 - ↔ there is a basis consisting of stationary states
 - ↔ time evolution is trivial
- Spectrum of H is continuous

Magnetic Transport Along Translationally Invariant Obstacles

Quantum Mechanics:

- Spectrum of H is pure point
 - ↔ there is a basis consisting of stationary states
 - ↔ time evolution is trivial
- Spectrum of H is continuous
 - ↔ there are no stationary states

Magnetic Transport Along Translationally Invariant Obstacles

Quantum Mechanics:

- Spectrum of H is pure point
 - ↔ there is a basis consisting of stationary states
 - ↔ time evolution is trivial
- Spectrum of H is continuous
 - ↔ there are no stationary states
 - ↔ **Magnetic Transport!**

Magnetic Transport Along Translationally Invariant Obstacles

The Hamiltonian is either of these:

- (a) $H = (-i\vec{\nabla} + \vec{A})^2 + V(x)$ on $L^2(\Omega \subset \mathbb{R}^2)$
- (b) $H = (-i\vec{\nabla} + \vec{A}(x))^2$ on $L^2(\Omega \subset \mathbb{R}^2)$
- (c) $H = (-i\vec{\nabla} + \vec{A})$ on $L^2(\Omega)$, Ω being a thin layer in \mathbb{R}^3

And we are interested in its pure point / continuous spectrum.

The two parts

① Summary of known results

- ▶ Steep potential wall (Macris et al., 1999) and (Fröhlich et al., 2000)
- ▶ Half-plane with Dirichlet boundary (Fröhlich et al., 2000)
- ▶ Bounded magnetic perturbation (Iwatsuka, 1983 and 1985)
- ▶ Layer with one-sided fold, asymptotically flat layer, very thin layer (Exner et al., 2018)

The two parts

① Summary of known results

- ▶ Steep potential wall (Macris et al., 1999) and (Fröhlich et al., 2000)
- ▶ Half-plane with Dirichlet boundary (Fröhlich et al., 2000)
- ▶ Bounded magnetic perturbation (Iwatsuka, 1983 and 1985)
- ▶ Layer with one-sided fold, asymptotically flat layer, very thin layer (Exner et al., 2018)

② Original work

- ▶ Half-plane with Robin boundary
- ▶ Dirac δ -interaction on a line

Frame Title

Frame Title