

OMENTUM, RIGHT HEMICOLECTOMY
 DIFFERENTIATED ADENOCARCINOMA OF THE CECUM (6.5 CM)
 INVADING THROUGH THE MUSCULARIS PROPRIA INTO PERICECAL FAT WITH MULTIPLE ISOLATED
 TUMOR NODULES. NO DEFINITIVE SEROSAL INVASION IS IDENTIFIED.
 TUMORS PRESENT AT THE RADIAL PERICECAL SOFT TISSUE MARGIN AND INTRAVASCULAR TUMOR IS
 PRESENT AT THE PROXIMAL MESENTERIC MARGIN (see comment).
 EXTENSIVE ANGIOLYMPHATIC, VASCULAR, AND PERINEURAL INVASION.
 ISOLATED FOCI OF METASTATIC ADENOCARCINOMA ARE PRESENT ON THE SURFACE OF THE OMENTUM
 AND WITHIN THE MUSCULARIS PROPRIA OF THE TERMINAL ILEUM.
 PROXIMAL ILEUM AND DISTAL COLON MARGINS OF RESECTION ARE FREE OF CARCINOMA.
 METASTATIC ADENOCARCINOMA IN THIRTEEN OF TWENTY-THREE LYMPH NODES (13/23).
 SEPARATE TUBULAR ADENOMA.
 PATHOLOGIC STAGE: pT3c/d N2 M1.
 APPENDIX NOT IDENTIFIED.

CASE SYNOPSIS:

SYNOPTIC DATA - PRIMARY COLON, RECTAL, APPENDIX TUMORS

SPECIMEN TYPE: Right hemicolectomy

SPECIMEN LENGTH: 40 cm

TUMOR SITE: Cecum

TUMOR CONFIGURATION: Infiltrative

TUMOR SIZE: Greatest dimension: 6.5 cm

Additional dimensions: 6.0 x 5.5 cm

INTACTNESS OF MESORECTUM: Not applicable

HISTOLOGIC TYPE: Adenocarcinoma

HISTOLOGIC GRADE: High-grade (poorly differentiated to undifferentiated)

PATHOLOGIC STAGING (pTNM): pT3c/d

pN2

Number of nodes examined: 23

Number of nodes involved: 13

pM1

Site(s): omentum, muscularis propria of ileum

Treatment: Untreated

Proximal margin uninvolved by invasive carcinoma

Distal margin uninvolved by invasive carcinoma

Circumferential (radial) margin involved by invasive carcinoma (tumor present 0-1 mm

from CRM)

Mesenteric margin involved by invasive carcinoma

Present

Present

TUMOR BORDER CONFIGURATION: Infiltrating

ADDITIONAL TUMOR CHARACTERISTICS: None

ADDITIONAL PATH FINDINGS (check all that apply):

Adenoma(s)

TCGA-AZ-4616