	Lesson 14
	Concentration
-	Trequalities
	concentration inequality
	ives us a probability
	, and on certain random
Vieta de la companya della companya della companya della companya de la companya della companya	

Nariable		
Nalues	· · · · · · · · · · · · · · · · · · ·	 

1. Markov's Ine	quality
Assume X is a non-neg	ative r.v.
with E[X] <00,	then for
any a>o	1/21-0/2
$P(X7d) \leq$	#LXJ

Remark: This inequality is
non-trivial when d> ECXJ.

-		
	VIV. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.	
<del>,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,</del>		
-		
-		

Question: The 'decay rate" is
Vd in Markov Tnequality.
Can we have a facay rate
that is faster than 1/2?
The answer is YES:

2. Chebysher's Inequality

Assume that X is a

r.v. with expectation u


and Variance or 200,


then

$P( X-\mu >k\sigma)\leq 1/k^2$ $\forall k>0$
4K>0
OR
$\mathbb{P}( X-\mu >c)\leq \frac{\sigma^2}{c^2}$

Proof:	

·
Classical Tadualia
Cheby sher's Inequality
Chebysher's Inequality
Chebysher's Inequality for Various k's
for Various k's
for Various k's
for Various ks  K=1 P(   X-\mu   > \sigma) \leq
for Various ks  K=1 P(   X-\mu   > \sigma) \leq
for Various k's
for Various ks  K=1 P(   X-\mu   > \sigma) \leq
for Various ks  K=1 P(   X-\mu   > \sigma) \leq





Chebysher In equality yields
a bound that decays as
fast as 1/2, an improvement
over the basic Murkor
Inequality.

Question: Can we find bounds

that decay faster than

polynomias, i.e., exponentially?

YES!