

On connections between k-coloring and Euclidean k-means

Karthik C. S.
(Rutgers University)


Joint Work with

Enver Aman
Undergrad at Rutgers
Graduated May 2024




Sharath Punna
Masters at Rutgers
Graduated May 2023

Graph Coloring




k-means Clustering




Graph Coloring

- **Input:** Graph $G = (V, E)$ and integer k




k-means Clustering

- **Input:** Points $P \subset \mathbb{R}^d$ and integer k



Graph Coloring

- **Input:** Graph $G = (V, E)$ and integer k
- **Output:** $V := V_1 \dot{\cup} \dots \dot{\cup} V_{k'}$, such that V_i is an **independent** set




k-means Clustering

- **Input:** Points $P \subset \mathbb{R}^d$ and integer k
- **Output:** $P := P_1 \dot{\cup} \dots \dot{\cup} P_{k'}$, such that


$\sum_{i \in [k]} \sum_{p \in P_i} \|p - c_i\|_2^2$ is **minimized**,

$$\text{where } c_i = \sum_{p \in P_i} \frac{p}{|P_i|}$$




Graph Coloring

- **Input:** Graph $G = (V, E)$ and integer k
- **Output:** $V := V_1 \dot{\cup} \dots \dot{\cup} V_{k'}$, such that V_i is an **independent** set
- Classic NP-hard problem



k-means Clustering

- **Input:** Points $P \subset \mathbb{R}^d$ and integer k
- **Output:** $P := P_1 \dot{\cup} \dots \dot{\cup} P_{k'}$, such that $\sum_{i \in [k]} \sum_{p \in P_i} \|p - c_i\|_2^2$ is **minimized**, where $c_i = \frac{p}{|P_i|}$
- Important NP-hard problem



k -coloring to k -means

A **simple** reduction

Graph k -Coloring

- Input: Graph $G = (V, E)$, G is d -regular

k -means Clustering

- Input: Points $P \subset \mathbb{R}^d$

k-coloring to k-means

A **simple** reduction

Graph k-Coloring

- Input: Graph $G = (V, E)$, G is d -regular

k-means Clustering

- Input: Points $P \subset \mathbb{R}^d$

Orient E arbitrarily

$$v \in V \longrightarrow p_v \in \mathbb{R}^{|E|}$$

$$p_v(e) = \begin{cases} +1 & \text{if } e \text{ is outgoing edge of } v \\ -1 & \text{if } e \text{ is incoming edge of } v \\ 0 & \text{otherwise} \end{cases}$$

k-coloring to k-means


A simple reduction

Graph k-Coloring

- Input: Graph $G = (V, E)$, G is d -regular

k-means Clustering

- Input: Points $P \subset \mathbb{R}^d$



Orient E arbitrarily

$$v \in V \longrightarrow p_v \in \mathbb{R}^{|E|}$$


$$p_v(e) = \begin{cases} +1 & \text{if } e \text{ is outgoing edge of } v \\ -1 & \text{if } e \text{ is incoming edge of } v \\ 0 & \text{otherwise} \end{cases}$$

k-coloring to k-means

A simple reduction


Graph k-Coloring

- Input: Graph $G = (V, E)$, G is d -regular



k-means Clustering

- Input: Points $P \subset \mathbb{R}^d$



Orient E arbitrarily

$$v \in V \longrightarrow p_v \in \mathbb{R}^{|E|}$$

$$p_v(e) = \begin{cases} +1 & \text{if } e \text{ is outgoing edge of } v \\ -1 & \text{if } e \text{ is incoming edge of } v \\ 0 & \text{otherwise} \end{cases}$$

k-coloring to k-means

A **simple** reduction

Graph k-Coloring

- Input: Graph $G = (V, E)$, G is d -regular

k-means Clustering

- Input: Points $P \subset \mathbb{R}^{|E|}$

$$\|p_u - p_v\|_2^2 = \begin{cases} 2d & \text{if } \{u, v\} \notin E \\ 2d + 2 & \text{if } \{u, v\} \in E \end{cases}$$

k-coloring to k-means

A **simple** reduction

Graph k-Coloring

- Input: Graph $G = (V, E)$, G is d -regular

k-means Clustering

- Input: Points $P \subset \mathbb{R}^{|E|}$

$$\|p_u - p_v\|_2^2 = \begin{cases} 2d & \text{if } \{u, v\} \notin E \\ 2d + 2 & \text{if } \{u, v\} \in E \end{cases}$$

$\sum_{i \in [k]} \sum_{p \in P_i} \|p - c_i\|_2^2$ is minimized,

$$\text{where } c_i = \sum_{p \in P_i} \frac{p}{|P_i|}$$



$\sum_{i \in [k]} \frac{1}{2|P_i|} \sum_{p,q \in P_i} \|p - q\|_2^2$ is minimized,

k-coloring to k-means

A **simple** reduction

$$\|p_u - p_v\|_2^2 = \begin{cases} 2d & \text{if } \{u, v\} \notin E \\ 2d + 2 & \text{if } \{u, v\} \in E \end{cases}$$

Completeness

Soundness

k-coloring to k-means

A **simple** reduction

$$\|p_u - p_v\|_2^2 = \begin{cases} 2d & \text{if } \{u, v\} \notin E \\ 2d + 2 & \text{if } \{u, v\} \in E \end{cases}$$

Completeness

- $V := V_1 \dot{\cup} \dots \dot{\cup} V_{k'}$ such that
 V_i is an **independent** set

Soundness

$$\begin{aligned} \sum_{i \in [k]} \frac{1}{2|P_i|} \sum_{p,q \in P_i} \|p - q\|_2^2 &= \sum_{i \in [k]} \frac{1}{2|P_i|} \sum_{p,q \in P_i} 2d \\ &= \sum_{i \in [k]} d \cdot (|P_i| - 1) = d \cdot (|V| - k) \end{aligned}$$

k-coloring to k-means

A **simple** reduction

$$\|p_u - p_v\|_2^2 = \begin{cases} 2d & \text{if } \{u, v\} \notin E \\ 2d + 2 & \text{if } \{u, v\} \in E \end{cases}$$

Completeness

- $V := V_1 \dot{\cup} \dots \dot{\cup} V_{k'}$ such that V_i is an **independent** set

$$\begin{aligned} \sum_{i \in [k]} \frac{1}{2|P_i|} \sum_{p,q \in P_i} \|p - q\|_2^2 &= \sum_{i \in [k]} \frac{1}{2|P_i|} \sum_{p,q \in P_i} 2d \\ &= \sum_{i \in [k]} d \cdot (|P_i| - 1) = d \cdot (|V| - k) \end{aligned}$$

Soundness

- $P := P_1 \dot{\cup} \dots \dot{\cup} P_k$ is some **clustering**
- $\forall i \in [k], \sum_{p,q \in P_i} \|p - q\|_2^2 \geq 2d \cdot |P_i| \cdot (|P_i| - 1)$
- $\exists i \in [k], \sum_{p,q \in P_i} \|p - q\|_2^2 > 2d \cdot |P_i| \cdot (|P_i| - 1)$

k -coloring to k -means

A simple reduction

$$\|p_u - p_v\|_2^2 = \begin{cases} 2d & \text{if } \{u, v\} \notin E \\ 2d + 2 & \in E \end{cases}$$

Completeness

Soundness

- $V := V_1 \dot{\cup} \dots \dot{\cup} V_{k'}$ such that V_i is an independent set

$$\begin{aligned} & \sum_{i \in [k]} \frac{1}{2|P_i|} \sum_{p,q \in P_i} \|p - q\|_2^2 = \sum_{i \in [k]} \frac{1}{2|P_i|} \sum_{p,q \in P_i} 2d \\ &= \sum_{i \in [k]} d \cdot (|P_i| - 1) = d \cdot (|V| - k) \end{aligned}$$

3-coloring is NP-hard \iff **3-means is NP-hard**

- $\forall i \in [k], \sum_{p,q \in P_i} \|p - q\|_2^2 \geq 2d \cdot |P_i| \cdot (|P_i| - 1)$
- $\exists i \in [k], \sum_{p,q \in P_i} \|p - q\|_2^2 > 2d \cdot |P_i| \cdot (|P_i| - 1)$

What about 1-means?



What about 2-means?

- 2-coloring reduces to 2-means
- 2-coloring is in P so 2-means is in P

What about 2-means?

- 2-coloring reduces to 2-means
- 2-coloring is in P so 2-means is in P



What about 2-means?


- 2-means is **NP-hard** [Dasgupta-Freund'09]

What about 2-means?

- 2-means is NP-hard [Dasgupta-Freund'09]
- **Proof Strategy:**
 - (i) Structured 3-NAE-SAT is NP-hard
 - (ii) NAE-SAT is reduced to distance matrix of points of 2-means instance
 - (iii) Distance matrix can be realized in Euclidean space (PSD check)

What about 2-means?

- 2-means is NP-hard [Dasgupta-Freund'09]
- **Proof Strategy:**
 - (i) Structured 3-NAE-SAT is NP-hard
 - (ii) NAE-SAT is reduced to distance matrix of points of 2-means instance
 - (iii) Distance matrix can be realized in Euclidean space (PSD check)



Almost 2-coloring to 2-means

New Reduction

Almost 2-coloring to 2-means

New Reduction

Balanced Max-Cut

- Input: Graph $G = (V, E)$, G is d -regular
- Output: $V := V_1 \dot{\cup} V_2$, to minimize:

$$|E(V_1, V_2)| \cdot \left(1 + \frac{||V_1| - |V_2||}{|V|} \right)$$

2-means Clustering

- Input: Points $P \subset \mathbb{R}^d$
- Output: $P := P_1 \dot{\cup} P_2$, to minimize:

$$\frac{\sum_{p,q \in P_1} \|p - q\|_2^2}{2|P_1|} + \frac{\sum_{p,q \in P_2} \|p - q\|_2^2}{2|P_2|}$$

Almost 2-coloring to 2-means

New Reduction

Balanced Max-Cut

- Input: Graph $G = (V, E)$, G is d -regular
- Output: $V := V_1 \dot{\cup} V_2$, to minimize:

$$|E(V_1, V_2)| \cdot \left(1 + \frac{||V_1| - |V_2||}{|V|} \right)$$

- We give NP-hardness reduction from
Linear 4-regular 3-NAE-SAT

2-means Clustering

- Input: Points $P \subset \mathbb{R}^d$
- Output: $P := P_1 \dot{\cup} P_2$, to minimize:

$$\frac{\sum_{p,q \in P_1} \|p - q\|_2^2}{2|P_1|} + \frac{\sum_{p,q \in P_2} \|p - q\|_2^2}{2|P_2|}$$

Almost 2-coloring to 2-means

New Reduction

Balanced Max-Cut

Orient E arbitrarily

- Input: Graph $G = (V, E)$, G is d -regular

- Output: $V := V_1 \dot{\cup} V_2$, to minimize:

$$|E(V_1, V_2)| \cdot \left(1 + \frac{||V_1| - |V_2||}{|V|}\right)$$

- We give NP-hardness reduction from
Linear 4-regular 3-NAE-SAT

$$v \in V \longrightarrow p_v \in \mathbb{R}^{|E|}$$

$$p_v(e) = \begin{cases} +1 & \text{if } e \text{ is outgoing edge of } v \\ -1 & \text{if } e \text{ is incoming edge of } v \\ 0 & \text{otherwise} \end{cases}$$

$$\|p_u - p_v\|_2^2 = \begin{cases} 2d & \text{if } \{u, v\} \notin E \\ 2d + 2 & \text{if } \{u, v\} \in E \end{cases}$$

Reverse the Reduction


Max-Cut on n vertices
can be solved in
 $2^{\omega n/3}$ time

[Williams'05]

Reverse the Reduction




Reverse the Reduction



Reverse the Reduction

Max-Cut on n vertices
can be solved in
 $2^{\omega n/3}$ time


[Williams'05]



Reverse the Reduction

Max-Cut on n vertices
can be solved in
 $2^{\omega n/3}$ time

[Williams'05]






Reverse the Reduction

Max-Cut on n vertices
can be solved in
 $2^{\omega n/3}$ time

[Williams'05]

- Enumerate all cuts (A_i, B_i) of G_i




Reverse the Reduction

Max-Cut on n vertices
can be solved in
 $2^{\omega n/3}$ time

[Williams'05]

- Enumerate all cuts (A_i, B_i) of G_i
- Construct graph H on $3 \cdot 2^{n/3}$ nodes
- Edge Weight of H is sum of cut edges




Reverse the Reduction

Max-Cut on n vertices
can be solved in
 $2^{\omega n/3}$ time

[Williams'05]

- Enumerate all cuts (A_i, B_i) of G_i
- Construct graph H on $3 \cdot 2^{n/3}$ nodes
- Edge Weight of H is sum of cut edges
- Solve weighted triangle detection on $3 \cdot 2^{n/3}$ node graph H




Reverse the Reduction

2-means on n points
can be solved in
 $2^{\omega n/3}$ time

Reverse the Reduction


2-means on n points
can be solved in
 $2^{\omega n/3}$ time



Reverse the Reduction

2-means on n points
can be solved in
 $2^{\omega n/3}$ time


- Enumerate all clustering (S_i, T_i) of P_i



Reverse the Reduction

2-means on n points
can be solved in
 $2^{\omega n/3}$ time


- Enumerate all clustering (S_i, T_i) of P_i
- Construct graph H on $3 \cdot 2^{n/3}$ nodes
- Edge Weight of H is sum of pairwise intracluster distances
- Solve weighted triangle detection on $3 \cdot 2^{n/3}$ node graph H



Open Directions

3-coloring \Rightarrow **3**-means

Open Directions



Can we use ideas from
3-coloring algorithms
to obtain $(2 - \varepsilon)^n$ time algorithm
for 3-means?

Open Directions

Can we use ideas from
2-means algorithm
to obtain $(2 - \varepsilon)^n$ time algorithm
for 2-median or 2-center?

Thank you for engaging!