

Review

Glucocorticoids and the regulation of memory in health and disease

Dominique J.-F. de Quervain^{a,*}, Amanda Aerni^a, Gustav Schelling^b, Benno Roozendaal^c^a Division of Psychiatry Research, University of Zürich, Lenggstr. 31, 8032 Zürich, Switzerland^b Department of Anesthesiology, Ludwig Maximilians University, 81377 Munich, Germany^c Department of Neuroscience, Section Anatomy, University Medical Center Groningen, University of Groningen, A. Deusinglaan 1, 9713 AV Groningen, The Netherlands

ARTICLE INFO

Article history:

Available online 31 March 2009

Keywords:

Glucocorticoids

Cortisol

Cortisone

Emotion

Memory

Retrieval

Extinction

PTSD

Phobia

Amygdala

ABSTRACT

Over the last decades considerable evidence has accumulated indicating that glucocorticoids – stress hormones released from the adrenal cortex – are crucially involved in the regulation of memory. Specifically, glucocorticoids have been shown to enhance memory consolidation of emotionally arousing experiences, but impair memory retrieval and working memory during emotionally arousing test situations. Furthermore, growing evidence indicates that these different glucocorticoid effects all depend on emotional arousal-induced activation of noradrenergic transmission within the basolateral complex of the amygdala (BLA) and on interactions of the BLA with other brain regions, such as the hippocampus and neocortical regions. Here we review findings from both animal and human experiments and present an integrated perspective of how these opposite glucocorticoid effects might act together to serve adaptive processing of emotionally significant information. Furthermore, as intense emotional memories also play a crucial role in the pathogenesis and symptomatology of anxiety disorders, such as posttraumatic stress disorder (PTSD) or phobias, we discuss to what extent the basic findings on glucocorticoid effects on emotional memory might have implications for the understanding and treatment of these clinical conditions. In this context, we review data suggesting that the administration of glucocorticoids might ameliorate chronic anxiety by reducing retrieval of aversive memories and enhancing fear extinction.

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

Stress activates the hypothalamus–pituitary–adrenal axis, which results in the release of glucocorticoid hormones (cortisol in humans, corticosterone in rodents) from the adrenal cortex. It has long been recognized that glucocorticoids readily enter the brain and affect cognition. Early reports on both enhancing and impairing properties of glucocorticoids on memory [9,16,21,63,91] have indicated that these hormones have complex effects on cognitive functions. More recent studies investigating glucocorticoid effects on distinct memory phases and studies discerning acute from chronic effects helped to disentangle the multifaceted actions of these stress hormones. For example, acute elevations of glucocorticoids are known to enhance the consolidation of memory of new information, but to impair the retrieval of already stored information [51,146,156]. Conditions with chronically elevated glucocorticoid levels are usually associated with impaired cognitive performance and these deficits are thought to result from a cumulative and long-lasting burden on hippocampal function and morphology [102,172]. Recently, however, it became clear that memory deficits observed under such chronic conditions can also

result, at least in part, from acute and reversible glucocorticoid actions on memory retrieval processes [39].

In the present review, we will summarize and discuss how glucocorticoids affect memory consolidation, retrieval and working memory and why these stress hormones specifically modulate memory of emotionally arousing experiences. Furthermore, because emotional memory plays a crucial role in the pathogenesis and symptomatology of anxiety disorders, such as posttraumatic stress disorder (PTSD) or phobias, we will discuss to what extent the basic findings on glucocorticoid effects on emotional memory might have clinical implications.

2. Glucocorticoid effects on memory consolidation

Memory consolidation is the process by which a fragile short-term memory trace is transferred into stable long-term memory. However, not all information is equally well transferred into long-term storage. In fact, it is well recognized that especially emotionally arousing experiences are well remembered, even after decades [107]. Successful memory consolidation depends on de-novo protein synthesis and on long-term changes in synaptic plasticity [76]. There is extensive evidence that glucocorticoids, along with other components of the stress response, are critically involved in regulating memory consolidation of emotionally arousing experiences [106]. Blockade of glucocorticoid production with

* Corresponding author. Fax: +41 44 384 2686.

E-mail address: quervain@bli.unizh.ch (D.J.-F. de Quervain).

the synthesis inhibitor metyrapone impairs memory consolidation in both animals and humans [95,148] and prevents stress- or epinephrine-induced memory enhancement [90,149]. Acute systemic administration of glucocorticoids enhances long-term memory consolidation when given either before or immediately after a training experience [2,27,41,63,155,166,171]. Such glucocorticoid effects on memory consolidation follow an inverted U-shape dose–response relationship: Moderate doses enhance memory, whereas higher doses are typically less effective or may even impair memory consolidation [166]. In rodents, enhancing effects of glucocorticoids on memory consolidation have been observed in many different kinds of learning tasks, including inhibitory avoidance, contextual and cued fear conditioning, water-maze spatial and cued training, object recognition and conditioned taste aversion [161]. These findings indicate that, in animals, glucocorticoids not only enhance memory of training on hippocampus-dependent tasks that have a strong spatial/contextual component, but also memory of recognition- and procedural-training that are known to depend on other brain systems. In humans, glucocorticoid effects on memory consolidation have mostly been investigated with respect to declarative tasks [72,197].

Glucocorticoids also play a role in the consolidation of memory of extinction training. Extinction occurs when conditioned responding to a stimulus decreases when the reinforcer is omitted [139]. Like other forms of learning, extinction acquisition is followed by a consolidation phase and it has been found that the administration of glucocorticoids facilitates the consolidation of extinction memory, whereas a suppression of glucocorticoid function impairs such extinction processes [14,21,35,200]. Glucocorticoids have been shown to enhance memory consolidation and synaptic plasticity by influencing a wide variety of cellular functions, including cell signaling, ion channel properties as well as cell structure [20,78,145].

2.1. Role of emotional arousal in enabling glucocorticoid effects on memory consolidation

Recent findings suggest that glucocorticoids modulate memory consolidation of emotionally arousing experiences but do not affect memory consolidation of neutral information. Learning tasks in animal experiments are usually emotionally arousing because of the motivational stimulation necessary to elicit changes in behavior. We investigated the importance of emotional arousal in mediating glucocorticoid effects on memory consolidation in rats trained on an object recognition task [124]. Although no rewarding or aversive stimulation is used during object recognition training, such training induces modest novelty-induced stress or arousal [46]. However, extensive habituation of rats to the training apparatus (in the absence of any objects) prior to the training reduces the arousal level induced by object recognition training. We found that corticosterone administered systemically immediately after training enhanced 24-h retention performance of rats that were not previously habituated to the experimental context (i.e., emotionally aroused rats). In contrast, post-training corticosterone did not affect 24-h retention of rats that had received extensive prior habituation to the experimental context and, thus, had decreased novelty-induced emotional arousal during training [124] (Fig. 1).

A link between the level of emotional arousal at encoding and the efficacy of adrenocortical stress hormones in influencing memory consolidation has also been demonstrated in humans. Cortisol administered shortly before or after learning selectively enhances long-term memory of emotionally arousing, but not of emotionally neutral, items [27,86], but see [2]. Moreover, a cold pressor stress in humans (i.e., placing the arm in ice water for up to 3 min), a procedure that significantly elevates endogenous cortisol levels, enhances memory of emotionally arousing slides, but does not




Fig. 1. The enhancing effect of post-training administration of corticosterone on 24-h object recognition performance depends on emotional arousal. Rats received a single injection of corticosterone or vehicle immediately after the 3-min training trial. Corticosterone administered in a dose of 1.0 mg/kg significantly enhanced 24-h object recognition memory of aroused rats in the WITHOUT-habituation condition (A) but failed to affect memory of non-aroused rats in the WITH-habituation condition (B). ** $P < 0.0001$ compared with the corresponding vehicle control group. Data are presented as means \pm SEM. Reprinted from Okuda et al. [124].

affect memory of relatively neutral slides [32]. Consistent with these findings, it has been reported that levels of endogenous cortisol at the time of learning correlated with enhanced memory consolidation only in individuals who were emotionally aroused [3]. Furthermore, it has been found that a stress-induced elevation of cortisol levels was associated with enhanced consolidation of emotionally arousing memory of conditioned fear [210]. Thus, these findings from animal and human studies indicate that training-associated endogenous emotional arousal is essential for enabling glucocorticoid effects on memory consolidation.

2.2. Role of arousal-induced noradrenergic activation in the amygdala

Why do stress hormones selectively enhance memory consolidation of emotionally arousing experiences? Our findings suggest that interactions between stress hormones and amygdala activity may be key in determining this selectivity. It is well established that emotional experiences that induce the release of adrenal stress hormones also activate the amygdala [36,129]. Evidence from a large number of studies in both animals and humans has indicated that the amygdala plays a critical role in a variety of processes involving emotionally arousing information, including emotional influences on attention, perception, and learning and memory [89,130]. Extensive evidence from our as well as other laboratories indicates that the enhancing effects of stress hormone administration on the consolidation of memory of emotionally arousing experiences involve the amygdala. Lesions of the basolateral complex of the amygdala (BLA) block the memory-modulatory effects induced by post-training systemic injections of glucocorticoids [105,155]. As lesions of the adjacent central nucleus of the amygdala (CEA) are ineffective, the BLA appears to be the critical region of the amygdala in mediating stress and arousal effects on memory consolidation. Moreover, and in support of this view, post-training infusions of a specific glucocorticoid receptor (GR) agonist administered into the BLA, but not the CEA, enhance memory consolidation, whereas intra-BLA infusions of a GR antagonist impair memory consolidation or block the facilitatory effects of stress exposure [40,59,158]. These findings thus indicate that the BLA may be a critical gateway in mediating stress hormone effects on memory consolidation.

In addition, there is evidence that glucocorticoids require noradrenergic activation within the BLA to influence memory of emotionally arousing training. In vivo microdialysis experiments in rats have shown that electric footshock, or training on an inhibitory avoidance task, induces the release of norepinephrine in the amygdala [67,108,136]. A blockade of the β -adrenoceptor or of

the downstream cyclic adenosine monophosphate (cAMP) signaling pathway in the BLA blocks the memory-enhancing effect of systemically administered glucocorticoids [137,154,162,164]. Glucocorticoid hormones readily enter the brain and bind directly to adrenal steroid receptors in the BLA and other brain regions [144]. Extensive evidence indicates that glucocorticoid effects on memory consolidation involve a selective activation of the GR [122,163]. Glucocorticoids are known to act through intracellular and intranuclear receptors and can affect gene transcription by direct binding of receptor homodimers to DNA or via protein–protein interactions with other transcription factors [15,44,123]. However, glucocorticoids may also act more rapidly by interacting with membrane-associated steroid receptors [43,77,125,189]. In line with this evidence, we reported that GR activation in the BLA may facilitate memory consolidation via a rapid potentiation of the norepinephrine signal cascade through an interaction with G-protein-mediated events [164]. Activation of β -adrenoceptors in the BLA enhances memory consolidation via stimulation of the cAMP-dependent protein kinase pathway [62]. Intra-BLA infusions of the GR antagonist RU 38486 (mifepristone) attenuated the dose-response effects of a β -adrenoceptor agonist on retention enhancement for inhibitory avoidance training [164]. As the GR antagonist had no effect on memory enhancement induced by post-training intra-BLA infusions of the synthetic cAMP analog 8-Br-cAMP, these findings indicate that cAMP activation is downstream from the interaction of glucocorticoids with the noradrenergic system. In addition to such postsynaptic actions, glucocorticoid administration may increase the availability of norepinephrine in the BLA via an activation of GRs located in brainstem noradrenergic cell groups [166]. Fig. 2 summarizes the interactions of glucocorticoids and other major neuromodulatory systems with the noradrenergic system of the BLA in regulating memory consolidation. Recent findings indicate that the stress-responsive neuropeptide corticotropin-releasing factor (CRF) also enhances memory consolidation via an interaction with the β -adrenoceptor–cAMP cascade in the BLA [165]. Moreover, it was found that the CRF and glucocorticoid systems interact in influencing β -adrenoceptor–cAMP effects on memory consolidation.

Based on the evidence summarized above, it may be hypothesized that emotional arousal-induced increases in noradrenergic activity within the BLA are essential in enabling glucocorticoid hormone effects on memory consolidation. Such a mechanism may then provide a direct explanation of the findings that glucocorticoid hormones selectively enhance memory consolidation of emotionally arousing experiences. We recently investigated this issue in rats trained on an object recognition task. As is discussed above, corticosterone enhances memory of object recognition training when administered to naive rats, but is ineffective in rats that have reduced training-associated emotional arousal because of prior habituation to the experimental context [124]. We found that, in non-habituated (i.e., emotionally aroused) rats, the β -adrenoceptor antagonist propranolol blocked the corticosterone-induced memory enhancement [162]. Propranolol infused directly into the BLA also blocked the enhancing effects of corticosterone on object recognition memory. To determine whether the failure of corticosterone to enhance memory consolidation under low-arousing conditions is due to insufficient training-induced noradrenergic activation, low doses of the α_2 -adrenoceptor antagonist yohimbine, which increases norepinephrine levels in the brain, were co-administered with the corticosterone to well-habituated rats immediately after object recognition training. The critical finding of this study was that such an augmented noradrenergic tone was sufficient to mimic the effects of emotional arousal in that simultaneously administered corticosterone enhanced memory consolidation [162]. Further, in habituated rats, corticosterone activated BLA neurons, as assessed by phosphorylated cAMP response-element binding (pCREB) protein immunoreactivity levels, only in animals also given yohimbine. Such observations strongly suggest that because glucocorticoid effects on memory consolidation require noradrenergic activation within the BLA, they only modulate memory under emotionally arousing conditions that induce the release of norepinephrine.

Human studies have also provided evidence that stress hormone effects on memory enhancement for emotionally arousing experiences require amygdala activity. Memory for emotionally arousing material is not enhanced in human subjects with selective




Fig. 2. Interactions of glucocorticoids and other major neuromodulatory systems with the amygdala noradrenergic system in regulating memory consolidation. Adrenal stress hormones are released during training in emotionally arousing tasks and are known to enhance memory consolidation. Epinephrine, which does not cross the blood–brain barrier, induces the release of norepinephrine (NE) in the BLA by activating vagal afferents to the nucleus of the solitary tract (NTS). Noradrenergic neurons in the NTS project directly to the BLA, and indirectly via the locus coeruleus (LC). Norepinephrine binds to both β -adrenoceptors and α_1 -adrenoceptors at postsynaptic sites and activates cAMP formation. Glucocorticoids freely enter the brain and bind to glucocorticoid receptors (GRs) in brainstem noradrenergic neurons to potentiate norepinephrine release in the BLA, as well as postsynaptically in BLA neurons to facilitate the norepinephrine signaling cascade. Glucocorticoids may influence the β -adrenoceptor–cAMP system via a coupling with α_1 -adrenoceptors. These stress hormone effects on noradrenergic activation in the BLA are required for regulating memory consolidation in other brain regions. Furthermore, CRF may facilitate the β -adrenoceptor–cAMP response, but independently from the α_1 -adrenoceptor-induced modulation. Glucocorticoids enhance memory consolidation via a synergistic interaction with both the CRF and α_1 -adrenoceptor systems in potentiation training-induced β -adrenoceptor–cAMP activation. $\alpha_1 = \alpha_1$ -adrenoceptor; $\beta = \beta$ -adrenoceptor; cAMP = adenosine 3',5'-cyclic monophosphate; CREB, cAMP response-element binding; PGV, nucleus paraventricularis; CRF-R, corticotropin-releasing factor receptor. Adapted from Roozendaal [146].

bilateral damage to the amygdala, as it is in normal subjects [5,31]. Studies using brain positron emission tomography (PET) and functional magnetic resonance imaging (fMRI) techniques have provided evidence that amygdala activity during viewing of emotionally arousing stimuli correlates highly with the subjects' recall of the material assessed weeks later [33,37,70]. Furthermore, the relationship between amygdala activity during encoding and subsequent memory was greatest for the material rated as being the most emotionally intense. In a recent study it has been shown that amygdala activity during viewing of emotionally arousing pictures was greatest for those subjects who responded with a large increase in endogenous cortisol [193] and the combination of pharmacologically induced elevation of central norepinephrine and cortisol in healthy subjects produced a negative response bias in the amygdala [87]. Furthermore, a β -adrenoceptor antagonist blocked the increase in amygdala activity and the enhanced retention induced by either emotional arousal or endogenous cortisol [34,188,194].

2.3. Interaction of the amygdala with other brain regions

As indicated, glucocorticoids are known to enhance memory consolidation of training on many different kinds of emotionally arousing tasks, including inhibitory avoidance, contextual and cued fear conditioning, water-maze spatial and cued training, object recognition and conditioned taste aversion. As these different training experiences are known to engage different brain systems during both training and the consolidation occurring after training [69,73,128], the stress hormone-induced modulation involves receptor activation in a variety of brain regions. For example, glucocorticoids administered into the hippocampus influence memory of water-maze spatial training [157], which is consistent with a role of the hippocampus in spatial/contextual learning and memory [96,118,169]. Glucocorticoids infused into the hippocampus also enhance memory of inhibitory avoidance training [157]. In fear conditioning tasks, including inhibitory avoidance, the rats learn that footshock occurs in a specific context. Importantly, lesions of the BLA or the administration of a β -adrenoceptor antagonist into the BLA block the enhancing effect of post-training intra-hippocampal infusions of a GR agonist on inhibitory avoidance memory [157,160]. BLA lesions also block the impairing effect of a GR antagonist infused into the hippocampus on spatial memory in water-maze [157]. Thus, these findings indicate that an intact and functional BLA is required for enabling memory modulation of spatial/contextual information induced by manipulation of GR activity in the hippocampus. Importantly, corticosterone enhances memory consolidation of conditioned taste aversion when infused post-training into the insular cortex or the BLA, but is ineffective when administered into the hippocampus, a region that does not play a significant role in taste learning [116]. Other findings suggest that corticosterone infusions into the caudate nucleus may selectively modulate the consolidation of memory of implicit (i.e., stimulus response) forms of learning [109] and there is evidence that BLA activity also regulates insular cortex- and caudate nucleus-dependent memory consolidation [115,127]. Taken together, these findings indicate that glucocorticoids act in different brain regions to enhance the consolidation of different aspects of information acquired during training. Although the BLA plays a critical role in regulating stress hormone effects on memory consolidation, extensive evidence indicates that it is not a permanent storage site of such memory traces [105], but rather interacts with these other brain regions in regulating memory consolidation of different kinds of information [104].

A growing body of evidence indicates that BLA neuronal activity is closely regulated by the medial prefrontal cortex (mPFC) [45,68,138]. The mPFC is primarily involved in higher-order cogni-

tive functions such as thought, decision-making and working memory [61,140] and exerts a strong inhibitory (i.e., fear-reducing) influence on behaviors [7]. Stress exposure is known to impair this mPFC-dependent inhibitory control [94] and to block long-term potentiation in the BLA–mPFC pathway [98]. We found that post-training infusions of a GR agonist administered into the mPFC enhanced memory consolidation of inhibitory avoidance training (Roosendaal et al., unpublished observation). In support of the view that a GR agonist administered into the mPFC enhances memory consolidation via a top-down regulation of BLA activity [100,167], the GR agonist infused into the mPFC after inhibitory avoidance training, but not in non-trained rats, increased BLA levels of phosphorylated extracellular signal-regulated kinase 1/2 (pErk1/2). Blockade of this pErk1/2 activity in the BLA prevented the memory enhancement induced by intra-mPFC administration of the GR agonist. Thus, these findings suggest that GR activation within the mPFC may enhance memory consolidation via a stimulatory influence on BLA activity, i.e., a loss of inhibitory control.

3. Glucocorticoid effects on memory retrieval

Studies showing that stress induces memory impairment when retention is tested shortly after learning (before memory consolidation took place) and glucocorticoid levels are still elevated [54,81], indicated that glucocorticoids possibly interfere with memory retrieval processes. In the first study investigating the specific effects of stress and glucocorticoids on memory retrieval [51], we reported that 30 min after exposure to footshock stress, rats had impaired retrieval of spatial memory of a water-maze task they had acquired 24 h earlier. Interestingly, memory performance was not impaired when rats were tested either 2 min or 4 h after the footshock. These time-dependent effects on retrieval processes corresponded to the circulating corticosterone levels at the time of testing, which suggested that the retrieval impairment was directly related to increased adrenocortical function. In support of this idea, we found that suppression of corticosterone synthesis with metyrapone blocked the stress-induced retention impairment. In addition, systemic corticosterone administered to non-stressed rats 30 min before retention testing induced dose-dependent retention impairment. Because similar footshock stress or corticosterone administration did not affect acquisition or immediate recall on the water-maze task, the corticosterone-induced impairment in retention performance is likely attributable to a selective influence on memory retrieval. In a next step, we have translated these findings to healthy humans and found that a single administration of 25 mg cortisone impaired the recall of words learned 24 h earlier [52]. Several further studies from different laboratories have indicated that glucocorticoids impair memory retrieval of spatial or contextual memory in rats and declarative (mostly episodic) memory in humans [29,39,49,72,83,142,152,153,170,197,198]. Moreover, increased cortisol levels after psychological stress exposure have also been shown to impair declarative memory retrieval [28,58,84].

The memory retrieval impairment induced by a single administration of glucocorticoids appears to be of temporary nature. Corticosterone injection impairs the acute retrieval of contextual fear memory without affecting retrieval performance 48 h later [35]. However, one study reported that a single administration of glucocorticoids not only induces acute, but also prolonged impairment of memory retrieval in humans [190]. In that study, subjects were asked to retrieve previously learned information after ingestion of 35 mg cortisol or placebo. The single administration of cortisol induced an acute impairment in memory recall, which, however, was still observed in an additional recall test after a 1-week washout period. The persistence of this retrieval impairment might be due

to a reduced rehearsal during retrieval under treatment and hence a lower re-encoding of the retrieved material. Alternatively, cortisol might have inhibited memory reconsolidation processes (see discussion under Section 5.4).

An important question is whether glucocorticoid effects on memory retrieval occur only under acutely elevated glucocorticoid levels or also under chronic conditions. There is substantial evidence that sustained endogenous hypercortisolemia, such as found in depression, Cushing's disease or human aging, is often associated with declarative memory impairment [92,168,179,186]. Moreover, prolonged glucocorticoid therapy, which is widely used in clinical practice, is known to induce cognitive deficits [18,24,25,79,199]. Although considerable evidence suggests that long-term glucocorticoid exposure may cause cognitive impairment via cumulative and long-lasting influences on hippocampal function and morphology [24,74,102,172,186], it is possible that also acute hormonal influences on retrieval processes contribute to the memory deficits found with chronic glucocorticoid exposure. To investigate this question, we recently examined memory functions and hippocampal volume in patients with rheumatoid arthritis who were treated either chronically (i.e., for a mean duration of 5 years) with low to moderate doses of prednisone or without glucocorticoids. In both groups, delayed recall of words learned 24 h earlier was assessed under conditions of either elevated or basal glucocorticoid levels in a double-blind, placebo-controlled crossover design. The findings did not indicate harmful effects of a history of chronic prednisone treatment on memory performance or hippocampal volume (i.e., no differences between both groups were found when retention was tested under placebo treatment). However, we found evidence that acute prednisone administration 1 h before retention testing to either the steroid or non-steroid group impaired verbal recall [39]. Thus, these findings indicate that acute and reversible effects of glucocorticoids on retrieval processes contribute to the memory deficits found in conditions of chronically elevated glucocorticoid levels.

3.1. Role of arousal-induced noradrenergic activation in enabling glucocorticoid effects on memory retrieval

Glucocorticoid effects on memory retrieval are highly comparable to those seen in studies investigating memory consolidation in that the effects depend on emotional arousal. Specifically, it has been shown in recent studies in humans that emotionally arousing information is especially sensitive to the retrieval-impairing effects of glucocorticoids (Fig. 3) [48,83,84,182], but see [190], and that emotional arousal during the test situation enables glucocorticoid effects on memory retrieval [85]. Studies of rats investigating the neural mechanisms underlying this selectivity have indicated that glucocorticoid effects on memory retrieval depend critically on noradrenergic activity within the brain. Systemic administration of the β -adrenoceptor antagonist propranolol blocks the memory retrieval impairment of spatial/contextual information induced by a concurrent injection of corticosterone [151]. A β -adrenoceptor antagonist infused into the hippocampus also prevents the retrieval-impairing effect of a GR agonist administered concurrently [153]. As stimulation of β 1-adrenoceptors with systemic injections of the selective agonist xamoterol induces memory retrieval impairment comparable to that seen after corticosterone administration [153], the findings suggest that glucocorticoid effects on memory retrieval involve a facilitation of noradrenergic mechanisms. Together with the finding that norepinephrine is activated by emotional arousal, this could explain why emotionally arousing information or an emotionally arousing test situation is a prerequisite for enabling glucocorticoid effects on memory retrieval. In line with this idea, we recently reported that the β -adrenoceptor antagonist propranolol blocked the impairing effect of




Fig. 3. Glucocorticoid effects on memory retrieval depend on emotional arousal: (A) cortisone administered 1 h before recall testing impaired retrieval of high-arousal words, but not of low-arousal words and (B) concurrent administration of propranolol prevented the impairing effect of cortisone on the retrieval of high-arousal words. Data are presented as mean \pm SEM. *** P = 0.0001 compared with the corresponding cortisone condition. Adapted from de Quervain et al. [48].

cortisone on the retrieval of emotionally arousing verbal material in healthy humans (Fig. 3) [48]. These findings may have important clinical implications as β -adrenoceptor antagonists might prove useful for the prevention of glucocorticoid-induced memory deficits in acute stressful situations [84], as well as for the treatment of memory deficits in conditions associated with chronically elevated glucocorticoid levels, such as depression or medical conditions requiring glucocorticoid treatment [39,150].

3.2. Role of amygdala–hippocampal interaction in enabling glucocorticoid effects on memory retrieval

Extensive evidence from studies in amnesic patients, human imaging studies, and lesion studies in animals indicates that the medial temporal lobe (hippocampus and parahippocampal gyrus) is crucially involved in memory retrieval of spatial and contextual memory in animals and declarative memory in humans [30,119,185]. Systemic administration of glucocorticoids to rats shortly before retention testing induces memory retrieval impairments for contextual and spatial memory [51,151]. Furthermore, we found that local infusions of a GR agonist into the hippocampus of rats induce retrieval impairment on a water-maze spatial task comparable to that seen after systemic administration [152]. In parallel, we performed a PET study in healthy humans and found that acutely administered cortisone reduces blood flow in the medial temporal lobe during memory recall of words, an effect that correlated with the degree of memory retrieval impairment [49]. Moreover, a recent fMRI study found that glucocorticoids decrease hippocampal and prefrontal activation during declarative memory retrieval [121]. Further studies in animals have indicated that the BLA interacts with the hippocampus in mediating glucocorticoid effects on the retrieval of emotionally arousing information. Lesions of the BLA or the infusion of a β -adrenoceptor antagonist into the BLA block the impairing effect of a GR agonist infused into the

hippocampus on memory retrieval of spatial information [152, 153]. In support of the importance of an interaction between the amygdala and the hippocampus during retrieval of emotionally arousing information, human imaging studies have indicated that the degree of interaction between these two brain regions is greater during the retrieval of emotionally arousing declarative information as compared to neutral information [57,183]. Therefore, comparable to the glucocorticoid effects on memory consolidation, the amygdala interacts with the hippocampus in mediating glucocorticoid effects on the retrieval of hippocampus-dependent memory.

4. Glucocorticoid effects on working memory

Working memory is a dynamic process whereby information is updated continuously, providing a temporary storage of information [12,75]. Evidence from lesion, pharmacological, imaging and clinical studies indicates that working memory depends on the integrity of the prefrontal cortex [23,66,126]. Stress exposure is known to impair performance of rats on a delayed alternation task, a task commonly used to assess working memory in rodents [11]. Basal levels of endogenous glucocorticoids are required to maintain prefrontal cortical function [117], but systemic injections of stress doses of corticosterone or administration of the GR agonist RU 28362 into the mPFC impair delayed alternation performance in rats [159]. As similar GR agonist infusions into the mPFC do not impair delayed alternation performance on non-mnemonic control tasks that have similar motivational and motor demands (Barsegyan and Roozendaal, unpublished observation), these findings strongly suggest that glucocorticoids, via GR activation, impair working memory. Additionally, stress or stress-level cortisol treatment is known to impair working memory performance in human subjects during demanding tasks that require a high level of arousal [12,93,178,198,208]. Importantly, like glucocorticoid effects on memory consolidation and retrieval, these hormones interact with noradrenergic mechanisms in inducing working memory impairment. A β -adrenoceptor antagonist administered systemically blocks the impairing effect of corticosterone on working memory in rats [159]. Furthermore, a β -adrenoceptor antagonist or cAMP blocker infused into the mPFC also blocks working memory impairment induced by a GR agonist administered concurrently (Barsegyan and Roozendaal, unpublished observation). Animal studies have shown that glucocorticoid effects on working memory also depend on functional interactions between the BLA and the mPFC. Disruption of BLA activity blocks the effect on working memory of a GR agonist administered into the mPFC [159]. This evidence provides strong support for the hypothesis that BLA activity modulates stress and emotional arousal effects on working memory in other brain regions.

5. Modulatory effects of glucocorticoids on emotional memory: implications for anxiety disorders

From the findings reviewed above we have learned that glucocorticoids enhance memory consolidation but impair memory retrieval and working memory in emotionally arousing situations (Fig. 4). Enhanced memory for emotional events is a well-recognized phenomenon, which helps us to remember important information. Although glucocorticoid-induced temporary impairments of memory retrieval and working memory are certainly unwanted in exam situations, these effects should not *a priori* be regarded as maladaptive. In fact, these effects may actually aid to an accurate storage of emotionally arousing information by blocking, for example, retroactive interference [147]. However, whereas the enhanced memory for emotionally arousing events in most cases




Fig. 4. Effects of stress and glucocorticoids on memory functions. Whereas glucocorticoids enhance memory consolidation, they impair memory retrieval and working memory. All of these hormone effects depend on emotional arousal-induced activation of noradrenergic transmission. NE, norepinephrine.

has a clear adaptive value, in certain circumstances extremely aversive experiences can also lead to highly emotional, traumatic or fearful memories, which contribute to the development and symptoms of anxiety disorders. Therefore, understanding the basic modulatory actions of glucocorticoids on different aspects of cognition may have important implications for understanding and, possibly, treating anxiety disorders. Although we focus in this paper on the implications of memory-modulatory glucocorticoid effects for anxiety disorders, these glucocorticoid effects are likely to have important implications for other psychiatric disorders, such as depression or schizophrenia, as well [17,120,168,187].

5.1. The role of aversive memories in anxiety disorders

Several lines of evidence indicate that after an aversive experience the formation of an aversive memory trace is an important pathogenic mechanism for the development of anxiety disorders, such as PTSD or phobias [89,114,130,131,205]. Neuroimaging studies have shown that while the prefrontal cortex seems to be hypo-responsive, amygdala activity in response to viewing aversive information is exaggerated in patients with PTSD as compared to healthy controls and, importantly, correlates positively with later recall of the aversive information, and with PTSD symptom severity [10,56,65,71,143,180,181]. These findings are in line with the well-known role of the amygdala in the formation of emotional memory, as reviewed above. Furthermore, some evidence indicates that the administration of a β -adrenoceptor blocker, which is known to reduce the consolidation of memory of emotionally arousing experiences, might be preventive with regard to the development of subsequent PTSD [134]. These findings underscore the important pathogenic role of aversive memory formation in the development of PTSD. However, the formation of a strong aversive memory trace is of course not sufficient to develop an anxiety disorder. In fact, building strong memories of an aversive event is a primarily adaptive mechanism and even intrusive thoughts (intrusive memory retrieval) and related symptoms are normal reactions in the first period after an aversive experience. In individuals who do not develop an anxiety disorder, which fortunately is mostly the case, intrusive memory retrieval declines over time, although the aversive memory can still be recalled voluntarily even after a long time. In contrast, in individuals who do develop a chronic anxiety disorder, the aversive memory trace remains easily reactivatable by an aversive cue (e.g. trauma cue or phobic stimulus), or even spontaneously, leading to uncontrollable aversive memory retrieval and related clinical symptoms (re-experiencing in PTSD, fear in phobia).

PTSD is a chronic response to a traumatic event and characterized by the following features: re-experiencing of the traumatic event, avoidance of stimuli associated with the trauma, and hyperarousal. Re-experiencing symptoms include intrusive daytime recollections, traumatic nightmares and flashbacks in which components of the event are relived [8,202]. These re-experiencing symptoms result from excessive retrieval of traumatic memories which often retain their vividness and power to evoke distress for decades or even a lifetime. Importantly, traumatic re-experiencing phenomena are again consolidated (re-consolidated) into memory, which cements the traumatic memory trace (see the discussion of the concept of intrusions in the PTSD research literature, e.g. [22,112,113]). Persistent retrieval, re-experiencing and reconsolidation of traumatic memories is a process that keeps these memories vivid and thereby the disorder alive (see Section 5.4).

Phobic disorders are characterized by marked and persistent fear that is excessive or unreasonable, cued by the presence or anticipation of a specific object or situation [8,13,97]. Exposure to a phobic stimulus almost invariably provokes retrieval of stimulus-associated fear memory which leads to the fear response [42,64,88]. In addition, phobic individuals tend to construct highly negative images of a phobic situation, which substantially contribute to anticipatory anxiety as well as negative postevent processing. Such images are usually associated with explicit fearful memories of past phobic experiences which reinforce negative beliefs that are difficult to suppress and may strengthen the phobic response [60,141]. Thus, retrieval of aversive memories (traumatic memory in PTSD and fear memory in phobias) plays an important role in the symptomatology of these anxiety disorders.

5.2. Glucocorticoids and PTSD

5.2.1. Preventive effects of glucocorticoids with regard to the development of PTSD

From the basic studies discussed above, we have learned that acutely elevated glucocorticoids enhance the consolidation of emotional memories. Based on these findings, it can be assumed that elevated glucocorticoid levels at the time of an aversive experience may contribute to the formation of traumatic and fearful memories. This idea is supported by a recent study on traumatic memories in critically ill patients. These patients often report traumatic memories from intensive care treatment and have a relatively high incidence of chronic stress symptoms and PTSD during follow-up [176]. We found that the number of traumatic memories from the intensive care unit correlated positively with the amount of cortisol acutely administered to patients undergoing cardiac surgery [175]. Theoretically, it might therefore be useful to therapeutically block glucocorticoid signaling immediately after a traumatic incident, as has been proposed for adrenergic signaling [132]. However, to block initial consolidation of aversive memories, the anti-glucocorticoid treatment should be given shortly after the aversive event. This is usually not possible and therefore this approach seems difficult. Moreover, there is evidence suggesting that reduced cortisol excretion in response to a traumatic event is actually associated with an increased risk of developing subsequent PTSD [53,103,206]. These findings suggest that elevated glucocorticoid levels after an aversive event might be preventive with regard to the development of PTSD. This idea is strongly supported by studies showing that prolonged (several days) administration of stress doses of cortisol during intensive care treatment reduces the risk for later PTSD [173,174,177,196]. But how do such findings of preventive effects fit with the idea that glucocorticoids enhance the formation of traumatic memories? After initial consolidation of traumatic experiences, which is likely to be enhanced by glucocorticoids, cortisol levels later on may play a crucial role in controlling the amount of retrieved traumatic memories. Specifically, by the

known reducing effects of glucocorticoids on memory retrieval, high levels of these hormones may partly interrupt the vicious cycle of retrieving, re-experiencing and reconsolidating aversive memories, thereby preventing a further cementation of the aversive memory trace. Studies showing that the preventive effects of glucocorticoid administration are also observed when the treatment started already at the time of the traumatic event [174,196] indicate that such an inhibitory effect of glucocorticoids on memory retrieval prevails the potentially enhancing effect on initial consolidation. Taken together, these findings suggest that elevated glucocorticoid levels (endogenously or pharmacologically) act preventively with regard to the development of PTSD.

5.2.2. Glucocorticoids reduce the retrieval of traumatic memories in chronic PTSD

In addition to individuals at risk for PTSD, also patients with an established PTSD can show low endogenous cortisol levels [19,99,201,203,204,207], but see [133,209]. A recent meta-analysis has shown that low cortisol levels depend on several factors, including gender and trauma type [110]. Low cortisol levels may contribute to a hyper-retrieval of aversive memories. Based on the finding that glucocorticoids impair the retrieval of emotional information, we hypothesized that patients with chronic PTSD might benefit from glucocorticoid treatment. In an initial study, we tested this hypothesis in a small number of patients with chronic PTSD [6]. During a 3-month observation period, low-dose cortisol (10 mg per day) was administered orally for 1 month using a double-blind, placebo-controlled, crossover design. The administration of this low dose of cortisol for 1 month does not cause major side effects and does not suppress endogenous cortisol production [38]. To assess possible treatment effects on the retrieval of traumatic memories, the patients rated daily the intensity and frequency of the feeling of reliving the traumatic event and the physiological distress felt in response to traumatic memories and nightmares (self-administered rating scales from the Clinician Administered PTSD Scale questions). The results of this study indicated that low-dose cortisol treatment had beneficial effects with regard to re-experiencing symptoms and nightmares and we found evidence for cortisol effects that outlasted the treatment period [6].

5.3. Glucocorticoids reduce fear in phobia

In recent clinical studies we found evidence that glucocorticoids may not only have beneficial effects in patients with PTSD but also in patients with phobias. We administered glucocorticoids to 40 subjects with social phobia and 20 subjects with spider phobia in two double-blind, placebo-controlled studies [184]. In the social phobia study, cortisone (25 mg) administered orally 1 h before a socio-evaluative stressor (Trier Social Stress Test [55,80]) significantly reduced self-reported fear during the anticipation-, exposure-, and recovery-phase of the stressor (Fig. 5A). Moreover, the stress-induced release of cortisol in placebo-treated subjects correlated negatively with fear ratings, suggesting that endogenously released cortisol in the context of a phobic situation buffers fear symptoms. This finding also indicates that reduced fear after glucocorticoid administration was not the result of a negative feedback on corticotropin-releasing factor (CRF) release.

In the spider phobia study, repeated oral administration of cortisol (10 mg), but not placebo, 1 h before exposure to a spider photograph induced a progressive reduction of stimulus-induced fear (Fig. 5B). This effect was maintained when subjects were exposed to the stimulus again 2 days after the last cortisol administration, suggesting that cortisol also facilitated the extinction of phobic fear (Fig. 5B). As in phobias retrieval processes cannot be measured directly, it cannot be ruled out that cortisol, perhaps in addition to influencing memory retrieval, may have reduced fear by exerting




Fig. 5. Effects of glucocorticoids on fear in phobia: (A) effects of a single administration of cortisone (25 mg) on fear ratings in social phobia. After substance administration, fear ratings were significantly lower in the cortisone group as compared to the placebo group in the course of the experiment ($P = 0.004$). Asterisks indicate significant differences at a certain time point, $P < 0.05$. T: time (min) in relation to the time point of substance administration at T0. TSST: Trier Social Stress Test and (B) effect of repeated administration of cortisol on stimulus-induced fear in spider phobia. In sessions 2–5, subjects were administered either cortisol (10 mg) or placebo 1 h before exposure to the phobic stimulus (spider photograph), whereas no pharmacological treatment was given in sessions 1 and 6. Fear ratings are depicted as mean \pm SEM. P -values indicate significance of symptom change across sessions for each treatment group. Fear symptoms were assessed using a visual analog scale ranging from 0 (no fear) to 10 (maximal fear). Adapted from Soravia et al. [184].

a direct anxiolytic effect or by modulating other systems involved in the expression of fear. For example, it has been found that acute cortisol administration can influence the startle reflex [26] and reduce preconscious attention to fear in anxious young men [135]. However, in these studies cortisol did not affect subjective fear levels. In addition, in the phobia study, glucocorticoid administration did not affect phobia-unrelated anxiety, mood, wakefulness, or calmness, suggesting that glucocorticoids did not reduce phobic fear by general anxiolytic effects [184]. Moreover, recent findings indicating that acute cortisol elevations cause heightened arousal

ratings of neutral stimuli [1] make a general or direct anxiolytic effect of glucocorticoids unlikely.

5.4. Possible mode of action of glucocorticoids in the reduction of aversive memory

The results of our clinical studies suggest that glucocorticoid administration induces acute effects on clinical symptoms by reducing the retrieval of aversive memories, i.e., traumatic memory in PTSD [6] and fear memory in phobia [184]. Additionally, in both studies we found evidence that symptoms were reduced even after cessation of the treatment period. What might be the underlying mechanism? Let us first review the processes which contribute to the persistence of these disorders: In PTSD, excessive retrieval of traumatic memory, which may be spontaneous or triggered by a trauma cue, leads to re-experiencing of the traumatic event [111]. In phobia, retrieval of fear memory triggered by a fear cue (phobic situation or object) leads to a fear response. (Re)consolidation of such aversive experiences further cements the aversive memory trace and thereby contributes to the persistence of these disorders (Fig. 6A). By inhibiting memory retrieval, cortisol may partly interrupt the vicious cycle of spontaneous retrieving, re-experiencing and reconsolidating traumatic memories in PTSD and, thereby, promote forgetting, a spontaneous process that occurs when memory is not reactivated (Fig. 6B). Furthermore, cortisol may facilitate the extinction of aversive memories, as evidenced by animal studies showing that glucocorticoid signaling promotes memory extinction processes [14,21,200]. Glucocorticoids may facilitate extinction in two ways: (i), because of the cortisol-induced reduction of memory retrieval, an aversive cue is no longer followed by the usual aversive memory retrieval and related clinical symptoms but, instead, becomes associated with a non-aversive experience which is stored as extinction memory; (ii), because elevated glucocorticoid levels are known to enhance the long-term consolidation of memories [27,63,82,86,146], it is possible that glucocorticoids facilitate the storage of corrective experiences. This is supported by recent animal studies showing that postretrieval administration of glucocorticoids is able to enhance the consolidation of extinction memory [4,35]. Theoretically, such postretrieval (or postreactivation) glucocorticoid effects may also be interpreted as an inhibition of reconsolidation [192,195]. However, findings in animals suggest that reconsolidation of aversive memory is disrupted by blocking rather than by activating glucocorticoid signaling [191]. Furthermore, in favor of the memory extinction hypothesis, it has been shown that postretrieval effects of glucocorticoids on memory are of transient nature and are reversed by a reminder, but see [195], which should not occur after inhibited reconsolidation. Although the data currently available rather speak for a facilitating effect of glucocorticoids on memory extinction, it is possible that, perhaps under certain conditions, glucocorticoids may also inhibit memory reconsolidation processes.

6. Conclusion

In the present paper we have reviewed evidence from both animal and human studies, which indicated that glucocorticoids enhance memory consolidation but impair memory retrieval and working memory (Fig. 4). Importantly, these hormone effects depend on emotional arousal-induced activation of noradrenergic transmission within the BLA and on interactions of the BLA with other brain regions, such as the hippocampus and neocortical regions. Therefore, glucocorticoids, via BLA activation, can modulate memory processes of many different kinds of emotionally arousing experiences.




Fig. 6. Model on the role of glucocorticoids in the reduction of aversive memory: (A) excessive retrieval of aversive memories causes re-experiencing symptoms in PTSD and phobic fear in phobia. Reconsolidation of such aversive experiences further cements the aversive memory trace and (B) glucocorticoid-induced reduction of the aversive memory trace. By inhibiting memory retrieval, glucocorticoids partly interrupt this vicious cycle of retrieving (1), re-experiencing (2) and reconsolidating (3) aversive memories, which leads to a weakening of the aversive memory trace (4). Furthermore, because the aversive cue is no longer followed by the usual aversive memory retrieval and related clinical symptoms, the cue becomes associated with a non-aversive experience, which is stored as extinction memory (5). Based on the findings of animal studies, glucocorticoids are likely to enhance long-term consolidation of extinction memory (see text for details).

Enhanced consolidation of emotionally arousing information is an adaptive mechanism, which helps us to retain important information. Reduced memory retrieval and working memory should not *a priori* be regarded as maladaptive as they support this process of retaining important information. In addition, the reduction of memory retrieval may aid to suppressing behaviors that are no more relevant or even maladaptive. This mechanism is especially important in more chronic situations when the organism is forced to adapt to a changed environment (e.g. environmental disaster, war). Under such conditions, also the facilitating effects of glucocorticoids on extinction processes represent an adaptive response, which helps the organism to deal with stressful events [47,101].

Because emotionally aversive memories play an important role in the development and symptomatology of anxiety disorders, we aimed to translate the basic findings on the effects of glucocorticoids on emotional memory in animals and healthy humans to clinical conditions. Specifically, the findings, which indicated that glucocorticoids reduce memory retrieval and enhance extinction of emotional memories led us to hypothesize that these stress hormones might be useful in the treatment of anxiety disorders. Clinical studies and studies of animal models of acquired fear indicate that glucocorticoid treatment indeed reduces the retrieval of traumatic memories and enhances extinction processes. These dual actions of glucocorticoids seem to be especially suited for

the treatment of acquired fear. By inhibiting memory retrieval, glucocorticoids may partly interrupt the vicious cycle of spontaneous retrieving, re-experiencing and (re)consolidating aversive memories. Furthermore, by enhancing extinction processes, glucocorticoids facilitate the storage of corrective experiences. Therefore, the combination of glucocorticoids with exposure techniques in cognitive-behavioral therapy may be a promising approach.

Future research should include large-scale clinical studies to evaluate the therapeutic efficacy of glucocorticoids in the treatment of anxiety disorders and to explore the efficacy of combining glucocorticoid treatment with psychotherapy. More research is also needed to better understand the molecular underpinnings of glucocorticoid actions on different memory processes as well as the role of (epi)genetic differences across individuals [50]. Such research might further promote the understanding of why some individuals become vulnerable to anxiety disorders, whereas others are resilient or even gain strength from stressful experiences.

References

- [1] H.C. Abercrombie, N.H. Kalin, R.J. Davidson, Acute cortisol elevations cause heightened arousal ratings of objectively nonarousing stimuli, *Emotion* 5 (2005) 354–359.
- [2] H.C. Abercrombie, N.H. Kalin, M.E. Thurow, M.A. Rosenkranz, R.J. Davidson, Cortisol variation in humans affects memory for emotionally laden and neutral information, *Behav. Neurosci.* 117 (2003) 505–516.
- [3] H.C. Abercrombie, N.S. Speck, R.M. Monticelli, Endogenous cortisol elevations are related to memory facilitation only in individuals who are emotionally aroused, *Psychoneuroendocrinology* 31 (2006) 187–196.
- [4] K. Abrari, A. Rashidy-Pour, S. Semnani, Y. Fathollahi, Administration of corticosterone after memory reactivation disrupts subsequent retrieval of a contextual conditioned fear memory: dependence upon training intensity, *Neurobiol. Learn. Mem.* 89 (2008) 178–184.
- [5] R. Adolphs, L. Cahill, R. Schul, R. Babinsky, Impaired declarative memory for emotional material following bilateral amygdala damage in humans, *Learn. Mem.* 4 (1997) 291–300.
- [6] A. Aerni, R. Traber, C. Hock, B. Roozendaal, G. Schelling, A. Papassotiropoulos, R.M. Nitsch, U. Schnyder, D.J. De Quervain, Low-dose cortisol for symptoms of posttraumatic stress disorder, *Am. J. Psychiatry* 161 (2004) 1488–1490.
- [7] J. Amat, M.V. Baratta, E. Paul, S.T. Bland, L.R. Watkins, S.F. Maier, Medial prefrontal cortex determines how stressor controllability affects behavior and dorsal raphe nucleus, *Nat. Neurosci.* 8 (2005) 365–371.
- [8] A. American Psychiatric, *Diagnostic and Statistical Manual of Mental Disorders*, fourth ed. (DSM-IV), Washington, DC, 1994.
- [9] I. Arbel, T. Kadar, M. Silbermann, A. Levy, The effects of long-term corticosterone administration on hippocampal morphology and cognitive performance of middle-aged rats, *Brain Res.* 657 (1994) 227–235.
- [10] J.L. Armony, V. Corbo, M.H. Clement, A. Brunet, Amygdala response in patients with acute PTSD to masked and unmasked emotional facial expressions, *Am. J. Psychiatry* 162 (2005) 1961–1963.
- [11] A.F. Arnsten, P.S. Goldman-Rakic, Noise stress impairs prefrontal cortical cognitive function in monkeys: evidence for a hyperdopaminergic mechanism, *Arch. Gen. Psychiatry* 55 (1998) 362–368.
- [12] A. Baddeley, Working memory, *Science* 255 (1992) 556–559.
- [13] D.H. Barlow, M.R. Liebowitz, H.I. Kaplan, B.J. Sadock, Specific and Social Phobias Comprehensive Textbook of Psychiatry, vol. VI, Williams and Wilkins, New York, 1995. pp. 1204–1218.
- [14] D. Barrett, F. Gonzalez-Lima, Behavioral effects of metyrapone on Pavlovian extinction, *Neurosci. Lett.* 371 (2004) 91–96.
- [15] M. Beato, P. Herrlich, G. Schutz, Steroid hormone receptors: many actors in search of a plot, *Cell* 83 (1995) 851–857.
- [16] B.E. Beckwith, T.V. Petros, C. Scaglione, J. Nelson, Dose-dependent effects of hydrocortisone on memory in human males, *Physiol. Behav.* 36 (1986) 283–286.
- [17] J.K. Belanoff, M. Kalezhan, B. Sund, S.K. Fleming Ficek, A.F. Schatzberg, Cortisol activity and cognitive changes in psychotic major depression, *Am. J. Psychiatry* 158 (2001) 1612–1616.
- [18] B. Bermond, S. Surachno, A. Lok, B.I. ten, B. Plasmans, C. Kox, E. Schuller, P.T. Schellekens, R. Hamel, Memory functions in prednisone-treated kidney transplant patients, *Clin. Transplant.* 19 (2005) 512–517.
- [19] L.M. Bierer, L. Tischler, E. Labinsky, S. Cahill, E. Foa, R. Yehuda, Clinical correlates of 24-h cortisol and norepinephrine excretion among subjects seeking treatment following the world trade center attacks on 9/11, *Ann. N.Y. Acad. Sci.* 1071 (2006) 514–520.
- [20] R. Bisaz, L. Conboy, C. Sandi, Learning under stress: a role for the neural cell adhesion molecule NCAM, *Neurobiol. Learn. Mem.* 91 (2009) 333–342.
- [21] B. Bohus, K. Lissak, Adrenocortical hormones and avoidance behaviour of rats, *Int. J. Neuropharmacol.* 7 (1968) 301–306.
- [22] C.R. Brewin, A cognitive neuroscience account of posttraumatic stress disorder and its treatment, *Behav. Res. Ther.* 39 (2001) 373–393.
- [23] G.N. Brito, G.J. Thomas, B.J. Davis, S.I. Gingold, Prelimbic cortex, mediodorsal thalamus, septum, and delayed alternation in rats, *Exp. Brain Res.* 46 (1982) 52–58.
- [24] E.S. Brown, J. Woolston, A. Frol, L. Bobadilla, D.A. Khan, M. Hanczyc, A.J. Rush, J. Fleckenstein, E. Babcock, C.M. Cullum, Hippocampal volume, spectroscopy, cognition, and mood in patients receiving corticosteroid therapy, *Biol. Psychiatry* 55 (2004) 538–545.
- [25] R. Brunner, D. Schaefer, K. Hess, P. Parzer, F. Resch, S. Schwab, Effect of corticosteroids on short-term and long-term memory, *Neurology* 64 (2005) 335–337.
- [26] T.W. Buchanan, A. Brechtel, J.J. Sollers, W.R. Lovallo, Exogenous cortisol exerts effects on the startle reflex independent of emotional modulation, *Pharmacol. Biochem. Behav.* 68 (2001) 203–210.
- [27] T.W. Buchanan, W.R. Lovallo, Enhanced memory for emotional material following stress-level cortisol treatment in humans, *Psychoneuroendocrinology* 26 (2001) 307–317.
- [28] T.W. Buchanan, D. Tranel, R. Adolphs, Impaired memory retrieval correlates with individual differences in cortisol response but not autonomic response, *Learn. Mem.* 13 (2006) 382–387.
- [29] C. Buss, O.T. Wolf, J. Witt, D.H. Hellhammer, Autobiographic memory impairment following acute cortisol administration, *Psychoneuroendocrinology* 29 (2004) 1093–1096.
- [30] R. Cabeza, L. Nyberg, Imaging cognition II: an empirical review of 275 PET and fMRI studies, *J. Cogn. Neurosci.* 12 (2000) 1–47.
- [31] L. Cahill, R. Babinsky, H.J. Markowitsch, J.L. McGaugh, The amygdala and emotional memory, *Nature* 377 (1995) 295–296.
- [32] L. Cahill, L. Gorski, K. Le, Enhanced human memory consolidation with post-learning stress: interaction with the degree of arousal at encoding, *Learn. Mem.* 10 (2003) 270–274.
- [33] L. Cahill, R.J. Haier, J. Fallon, M.T. Alkire, C. Tang, D. Keator, J. Wu, J.L. McGaugh, Amygdala activity at encoding correlated with long-term, free recall of emotional information, *Proc. Natl. Acad. Sci. USA* 93 (1996) 8016–8021.
- [34] L. Cahill, B. Prins, M. Weber, J.L. McGaugh, Beta-adrenergic activation and memory for emotional events, *Nature* 371 (1994) 702–704.
- [35] W.H. Cai, J. Blundell, J. Han, R.W. Greene, C.M. Powell, Postreactivation glucocorticoids impair recall of established fear memory, *J. Neurosci.* 26 (2006) 9560–9566.
- [36] S. Campeau, M.D. Hayward, B.T. Hope, J.B. Rosen, E.J. Nestler, M. Davis, Induction of the c-fos proto-oncogene in rat amygdala during unconditioned and conditioned fear, *Brain Res.* 565 (1991) 349–352.
- [37] T. Canli, Z. Zhao, J. Brewer, J.D. Gabrieli, L. Cahill, Event-related activation in the human amygdala associates with later memory for individual emotional experience, *J. Neurosci.* 20 (2000) RC99.
- [38] A.J. Cleare, E. Heap, G.S. Malhi, S. Wessely, V. O'Keane, J. Miell, Low-dose hydrocortisone in chronic fatigue syndrome: a randomised crossover trial, *Lancet* 353 (1999) 455–458.
- [39] D. Coluccia, O.T. Wolf, S. Kollias, B. Roozendaal, A. Forster, D.J. De Quervain, Glucocorticoid therapy-induced memory deficits: acute versus chronic effects, *J. Neurosci.* 28 (2008) 3474–3478.
- [40] C.D. Conrad, D.D. MacMillan, S. Tsekhanov, R.L. Wright, S.E. Baran, R.A. Fuchs, Influence of chronic corticosterone and glucocorticoid receptor antagonism in the amygdala on fear conditioning, *Neurobiol. Learn. Mem.* 81 (2004) 185–199.
- [41] M.I. Cordero, N.D. Kruyt, J.J. Merino, C. Sandi, Glucocorticoid involvement in memory formation in a rat model for traumatic memory, *Stress* 5 (2002) 73–79.
- [42] B.N. Cuthbert, P.J. Lang, C. Strauss, D. Drobos, C.J. Patrick, M.M. Bradley, The psychophysiology of anxiety disorder: fear memory imagery, *Psychophysiology* 40 (2003) 407–422.
- [43] M.F. Dallman, Fast glucocorticoid actions on brain: back to the future, *Front. Neuroendocrinol.* 26 (2005) 103–108.
- [44] N.A. Datson, J. van der Perk, E.R. De Kloet, E. Vreugdenhil, Identification of corticosteroid-responsive genes in rat hippocampus using serial analysis of gene expression, *Eur. J. Neurosci.* 14 (2001) 675–689.
- [45] R.J. Davidson, Anxiety and affective style: role of prefrontal cortex and amygdala, *Biol. Psychiatry* 51 (2002) 68–80.
- [46] S.F. De Boer, S.J. Koopmans, J.L. Slangen, G.J. Van der, Plasma catecholamine, corticosterone and glucose responses to repeated stress in rats: effect of interstressor interval length, *Physiol. Behav.* 47 (1990) 1117–1124.
- [47] E.R. De Kloet, M.S. Oitzl, M. Joels, Stress and cognition: are corticosteroids good or bad guys?, *Trends Neurosci* 22 (1999) 422–426.
- [48] D.J. De Quervain, A. Aerni, B. Roozendaal, Preventive effect of {beta}-adrenoceptor blockade on glucocorticoid-induced memory retrieval deficits, *Am. J. Psychiatry* 164 (2007) 967–969.
- [49] D.J. De Quervain, K. Henke, A. Aerni, V. Treyer, J.L. McGaugh, T. Berthold, R.M. Nitsch, A. Buck, B. Roozendaal, C. Hock, Glucocorticoid-induced impairment of declarative memory retrieval is associated with reduced blood flow in the medial temporal lobe, *Eur. J. Neurosci.* 17 (2003) 1296–1302.
- [50] D.J. De Quervain, I.T. Kolassa, V. Ertl, P.L. Onyut, F. Neuner, T. Elbert, A. Papassotiropoulos, A deletion variant of the alpha2b-adrenoceptor is related to emotional memory in Europeans and Africans, *Nat. Neurosci.* 10 (2007) 1137–1139.
- [51] D.J. De Quervain, B. Roozendaal, J.L. McGaugh, Stress and glucocorticoids impair retrieval of long-term spatial memory, *Nature* 394 (1998) 787–790.

- [52] D.J. De Quervain, B. Roozendaal, R.M. Nitsch, J.L. McGaugh, C. Hock, Acute cortisone administration impairs retrieval of long-term declarative memory in humans, *Nat. Neurosci.* 3 (2000) 313–314.
- [53] D.L. Delahanty, A.J. Raimonde, E. Spoonster, Initial posttraumatic urinary cortisol levels predict subsequent PTSD symptoms in motor vehicle accident victims, *Biol. Psychiatry* 48 (2000) 940–947.
- [54] D.M. Diamond, M. Fleshner, N. Ingersoll, G.M. Rose, Psychological stress impairs spatial working memory: relevance to electrophysiological studies of hippocampal function, *Behav. Neurosci.* 110 (1996) 661–672.
- [55] S.S. Dickerson, M.E. Kemeny, Acute stressors and cortisol responses: a theoretical integration and synthesis of laboratory research, *Psychol. Bull.* 130 (2004) 355–391.
- [56] E.W. Dickie, A. Brunet, V. Akerib, J.L. Armony, An fMRI investigation of memory encoding in PTSD: influence of symptom severity, *Neuropsychologia* 46 (2008) 1522–1531.
- [57] F. Dolcos, K.S. LaBar, R. Cabeza, Remembering one year later: role of the amygdala and the medial temporal lobe memory system in retrieving emotional memories, *Proc. Natl. Acad. Sci. USA* 102 (2005) 2626–2631.
- [58] G. Domes, M. Heinrichs, U. Rimmel, U. Reichwald, M. Hautzinger, Acute stress impairs recognition for positive words—association with stress-induced cortisol secretion, *Stress* 7 (2004) 173–181.
- [59] M.P. Donley, J. Schulkin, J.B. Rosen, Glucocorticoid receptor antagonism in the basolateral amygdala and ventral hippocampus interferes with long-term memory of contextual fear, *Behav. Brain Res.* 164 (2005) 197–205.
- [60] L. Fehm, J. Margraf, Thought suppression: specificity in agoraphobia versus broad impairment in social phobia?, *Behav. Res. Ther.* 40 (2002) 57–66.
- [61] L.K. Fellows, M.J. Farah, The role of ventromedial prefrontal cortex in decision making: judgment under uncertainty or judgment per se?, *Cereb. Cortex* 17 (2007) 2669–2674.
- [62] B. Ferry, B. Roozendaal, J.L. McGaugh, Basolateral amygdala noradrenergic influences on memory storage are mediated by an interaction between beta- and alpha1-adrenoceptors, *J. Neurosci.* 19 (1999) 5119–5123.
- [63] J.F. Flood, D. Vidal, E.L. Bennett, A.E. Orme, S. Vasquez, M.E. Jarvik, Memory facilitating and anti-amnesic effects of corticosteroids, *Pharmacol. Biochem. Behav.* 8 (1978) 81–87.
- [64] E.B. Foa, M.J. Kozak, Emotional processing of fear: exposure to corrective information, *Psychol. Bull.* 99 (1986) 20–35.
- [65] V. Francati, E. Vermetten, J.D. Bremner, Functional neuroimaging studies in posttraumatic stress disorder: review of current methods and findings, *Depress Anxiety* 24 (2007) 202–218.
- [66] J.M. Fuster, The prefrontal cortex and its relation to behavior, *Prog. Brain Res.* 87 (1991) 201–211.
- [67] R. Galvez, M.H. Mesches, J.L. McGaugh, Norepinephrine release in the amygdala in response to footshock stimulation, *Neurobiol. Learn. Mem.* 66 (1996) 253–257.
- [68] R. Garcia, R.M. Vouimba, M. Baudry, R.F. Thompson, The amygdala modulates prefrontal cortex activity relative to conditioned fear, *Nature* 402 (1999) 294–296.
- [69] P.E. Gold, R. van Buskirk, Facilitation of time-dependent memory processes with posttrial epinephrine injections, *Behav. Biol.* 13 (1975) 145–153.
- [70] S.B. Hamann, T.D. Ely, S.T. Grafton, C.D. Kilts, Amygdala activity related to enhanced memory for pleasant and aversive stimuli, *Nat. Neurosci.* 2 (1999) 289–293.
- [71] C. Heim, C.B. Nemeroff, Neurobiology of posttraumatic stress disorder, *CNS Spectr.* 14 (2009) 13–24.
- [72] S. Het, G. Ramlow, O.T. Wolf, A meta-analytic review of the effects of acute cortisone administration on human memory, *Psychoneuroendocrinology* 30 (2005) 771–784.
- [73] I. Izquierdo, J.A. Quirrellfeldt, M.S. Zanatta, J. Quevedo, E. Schaeffer, P.K. Schmitz, J.H. Medina, Sequential role of hippocampus and amygdala, entorhinal cortex and parietal cortex in formation and retrieval of memory for inhibitory avoidance in rats, *Eur. J. Neurosci.* 9 (1997) 786–793.
- [74] M. Joëls, Corticosteroid actions in the hippocampus, *J. Neuroendocrinol.* 13 (2001) 657–669.
- [75] M.W. Jones, A comparative review of rodent prefrontal cortex and working memory, *Curr. Mol. Med.* 2 (2002) 639–647.
- [76] E.R. Kandel, The molecular biology of memory storage: a dialogue between genes and synapses, *Science* 294 (2001) 1030–1038.
- [77] H. Karst, S. Berger, M. Turiault, F. Tronche, G. Schutz, M. Joëls, Mineralocorticoid receptors are indispensable for nongenomic modulation of hippocampal glutamate transmission by corticosterone, *Proc. Natl. Acad. Sci. USA* 102 (2005) 19204–19207.
- [78] H. Karst, S. Nair, E. Velzing, E.L. Rumpff-van, E. Slagter, P. Shinnick-Gallagher, M. Joëls, Glucocorticoids alter calcium conductances and calcium channel subunit expression in basolateral amygdala neurons, *Eur. J. Neurosci.* 16 (2002) 1083–1089.
- [79] P.A. Keenan, M.W. Jacobson, R.M. Soleymani, M.D. Mayes, M.E. Stress, D.T. Yaloo, The effect on memory of chronic prednisone treatment in patients with systemic disease, *Neurology* 47 (1996) 1396–1402.
- [80] C. Kirschbaum, K.M. Pirke, D.H. Hellhammer, The ‘trier social stress test’—a tool for investigating psychobiological stress responses in a laboratory setting, *Neuropsychobiology* 28 (1993) 76–81.
- [81] C. Kirschbaum, O.T. Wolf, M. May, W. Wippich, D.H. Hellhammer, Stress- and treatment-induced elevations of cortisol levels associated with impaired declarative memory in healthy adults, *Life Sci.* 58 (1996) 1475–1483.
- [82] G.L. Kovacs, G. Telegdy, K. Lissak, Dose-dependent action of corticosteroids on brain serotonin content and passive avoidance behavior, *Horm. Behav.* 8 (1977) 155–165.
- [83] S. Kuhlmann, C. Kirschbaum, O.T. Wolf, Effects of oral cortisol treatment in healthy young women on memory retrieval of negative and neutral words, *Neurobiol. Learn. Mem.* 83 (2005) 158–162.
- [84] S. Kuhlmann, M. Piel, O.T. Wolf, Impaired memory retrieval after psychosocial stress in healthy young men, *J. Neurosci.* 25 (2005) 2977–2982.
- [85] S. Kuhlmann, O.T. Wolf, A non-arousing test situation abolishes the impairing effects of cortisol on delayed memory retrieval in healthy women, *Neurosci. Lett.* 399 (2006) 268–272.
- [86] S. Kuhlmann, O.T. Wolf, Arousal and cortisol interact in modulating memory consolidation in healthy young men, *Behav. Neurosci.* 120 (2006) 217–223.
- [87] J. Kukulja, T.E. Schlapfer, C. Keyser, D. Klingmuller, W. Maier, G.R. Fink, R. Hurlmann, Modeling a negative response bias in the human amygdala by noradrenergic–glucocorticoid interactions, *J. Neurosci.* 28 (2008) 12868–12876.
- [88] P.J. Lang, A.H. Tuma, J.D. Maser, The cognitive psychophysiology of emotion: fear and anxiety, anxiety and the anxiety disorders, Erlbaum, Hillsdale, NJ, 1985, pp. 131–170.
- [89] J. LeDoux, The emotional brain, fear, and the amygdala, *Cell. Mol. Neurobiol.* 23 (2003) 727–738.
- [90] L. Liu, M. Tsuji, H. Takeda, K. Takada, T. Matsumiya, Adrenocortical suppression blocks the enhancement of memory storage produced by exposure to psychological stress in rats, *Brain Res.* 821 (1999) 134–140.
- [91] V.N. Luine, R.L. Spencer, B.S. McEwen, Effects of chronic corticosterone ingestion on spatial memory performance and hippocampal serotonergic function, *Brain Res.* 616 (1993) 65–70.
- [92] S.J. Lupien, M. de Leon, S. De Santi, A. Convit, C. Tarshish, N.P. Nair, M. Thakur, B.S. McEwen, R.L. Hauger, M.J. Meaney, Cortisol levels during human aging predict hippocampal atrophy and memory deficits, *Nat. Neurosci.* 1 (1998) 69–73.
- [93] S.J. Lupien, C.J. Gillin, R.L. Hauger, Working memory is more sensitive than declarative memory to the acute effects of corticosteroids: a dose–response study in humans, *Behav. Neurosci.* 113 (1999) 420–430.
- [94] D.M. Lyons, J.M. Lopez, C. Yang, A.F. Schatzberg, Stress-level cortisol treatment impairs inhibitory control of behavior in monkeys, *J. Neurosci.* 20 (2000) 7816–7821.
- [95] F.S. Maheu, R. Joob, S. Beaulieu, S.J. Lupien, Differential effects of adrenergic and corticosteroid hormonal systems on human short- and long-term declarative memory for emotionally arousing material, *Behav. Neurosci.* 118 (2004) 420–428.
- [96] S. Maren, M.S. Fanselow, Electrolytic lesions of the fimbria/fornix, dorsal hippocampus, or entorhinal cortex produce anterograde deficits in contextual fear conditioning in rats, *Neurobiol. Learn. Mem.* 67 (1997) 142–149.
- [97] I. Marks, Fears, Phobias and Rituals: Panic, Anxiety, and their Disorders, Oxford University Press, New York, 1987.
- [98] M. Maroun, G. Richter-Levin, Exposure to acute stress blocks the induction of long-term potentiation of the amygdala–prefrontal cortex pathway in vivo, *J. Neurosci.* 23 (2003) 4406–4409.
- [99] J.W. Mason, E.L. Giller, T.R. Kosten, R.B. Ostroff, L. Podd, Urinary free-cortisol levels in posttraumatic stress disorder patients, *J. Nerv. Ment. Dis.* 174 (1986) 145–149.
- [100] A.J. McDonald, Organization of amygdaloid projections to the prefrontal cortex and associated striatum in the rat, *Neuroscience* 44 (1991) 1–14.
- [101] B.S. McEwen, Protective and damaging effects of stress mediators, *N. Engl. J. Med.* 338 (1998) 171–179.
- [102] B.S. McEwen, Plasticity of the hippocampus: adaptation to chronic stress and allostatic load, *Ann. N.Y. Acad. Sci.* 933 (2001) 265–277.
- [103] A.C. McFarlane, M. Atchison, R. Yehuda, The acute stress response following motor vehicle accidents and its relation to PTSD, *Ann. N.Y. Acad. Sci.* 821 (1997) 437–441.
- [104] J.L. McGaugh, Memory consolidation and the amygdala: a systems perspective, *Trends Neurosci.* 25 (2002) 456.
- [105] J.L. McGaugh, The amygdala modulates the consolidation of memories of emotionally arousing experiences, *Annu. Rev. Neurosci.* 27 (2004) 1–28.
- [106] J.L. McGaugh, B. Roozendaal, Role of adrenal stress hormones in forming lasting memories in the brain, *Curr. Opin. Neurobiol.* 12 (2002) 205–210.
- [107] J.L. McGaugh, The making of lasting memory, *Mem. Emotion* (2003).
- [108] C.K. McIntyre, T. Hatfield, J.L. McGaugh, Amygdala norepinephrine levels after training predict inhibitory avoidance retention performance in rats, *Eur. J. Neurosci.* 16 (2002) 1223–1226.
- [109] A.C. Medina, J.R. Charles, V. Espinoza-Gonzalez, O. Sanchez-Resendis, R.A. Prado-Alcala, B. Roozendaal, G.L. Quirarte, Glucocorticoid administration into the dorsal striatum facilitates memory consolidation of inhibitory avoidance training but not of the context or footshock components, *Learn. Mem.* 14 (2007) 673–677.
- [110] M.L. Meewisse, J.B. Reitsma, G.J. de Vries, B.P. Gersons, M. Olff, Cortisol and post-traumatic stress disorder in adults: systematic review and meta-analysis, *Br. J. Psychiatry* 191 (2007) 387–392.
- [111] T. Michael, A. Ehlers, Enhanced perceptual priming for neutral stimuli occurring in a traumatic context: two experimental investigations, *Behav. Res. Ther.* 45 (2007) 341–358.
- [112] T. Michael, A. Ehlers, S.L. Halligan, Enhanced priming for trauma-related material in posttraumatic stress disorder, *Emotion* 5 (2005) 103–112.

- [113] T. Michael, A. Ehlers, S.L. Halligan, D.M. Clark, Unwanted memories of assault: what intrusion characteristics are associated with PTSD?, *Behav. Res. Ther.* 43 (2005) 613–628.
- [114] S. Mineka, K. Oehlberg, The relevance of recent developments in classical conditioning to understanding the etiology and maintenance of anxiety disorders, *Acta Psychol. (Amst.)* 127 (2008) 567–580.
- [115] M.I. Miranda, J.L. McGaugh, Enhancement of inhibitory avoidance and conditioned taste aversion memory with insular cortex infusions of 8-Br-cAMP: involvement of the basolateral amygdala, *Learn. Mem.* 11 (2004) 312–317.
- [116] M.I. Miranda, G.L. Quirarte, G. Rodriguez-Garcia, J.L. McGaugh, B. Roozendaal, Glucocorticoids enhance taste aversion memory via actions in the insular cortex and basolateral amygdala, *Learn. Mem.* 15 (2008) 468–476.
- [117] K. Mizoguchi, A. Ishige, S. Takeda, M. Aburada, T. Tabira, Endogenous glucocorticoids are essential for maintaining prefrontal cortical cognitive function, *J. Neurosci.* 24 (2004) 5492–5499.
- [118] R.G. Morris, P. Garrud, J.N. Rawlins, J. O'Keefe, Place navigation impaired in rats with hippocampal lesions, *Nature* 297 (1982) 681–683.
- [119] M.B. Moser, E.I. Moser, Distributed encoding and retrieval of spatial memory in the hippocampus, *J. Neurosci.* 18 (1998) 7535–7542.
- [120] J.W. Newcomer, S. Craft, K. Askins, T. Hershey, M.E. Bardgett, J.G. Csernansky, A.E. Gagliardi, G. Vogler, Glucocorticoid interactions with memory function in schizophrenia, *Psychoneuroendocrinology* 23 (1998) 65–72.
- [121] N.Y.L. Oei, B.M. Elzinga, O.T. Wolf, M.B. Ruiters, J.S. Damoiseaux, J.P.A. Kuijter, D.J. Veltman, P. Scheltens, S.A.R.B. Rombouts, Glucocorticoids decrease hippocampal and prefrontal activation during declarative memory retrieval in young men, *Brain Imag. Behav.* 1 (2007) 31–41.
- [122] M.S. Oitzl, E.R. De Kloet, Selective corticosteroid antagonists modulate specific aspects of spatial orientation learning, *Behav. Neurosci.* 106 (1992) 62–71.
- [123] M.S. Oitzl, H.M. Reichardt, M. Joels, E.R. De Kloet, Point mutation in the mouse glucocorticoid receptor preventing DNA binding impairs spatial memory, *Proc. Natl. Acad. Sci. USA* 98 (2001) 12790–12795.
- [124] S. Okuda, B. Roozendaal, J.L. McGaugh, Glucocorticoid effects on object recognition memory require training-associated emotional arousal, *Proc. Natl. Acad. Sci. USA* 101 (2004) 853–858.
- [125] M. Orchinik, T.F. Murray, F.L. Moore, A corticosteroid receptor in neuronal membranes, *Science* 252 (1991) 1848–1851.
- [126] A.M. Owen, K.M. McMillan, A.R. Laird, E. Bullmore, N-back working memory paradigm: a meta-analysis of normative functional neuroimaging studies, *Hum. Brain Mapp.* 25 (2005).
- [127] M.G. Packard, L. Cahill, J.L. McGaugh, Amygdala modulation of hippocampal-dependent and caudate nucleus-dependent memory processes, *Proc. Natl. Acad. Sci. USA* 91 (1994) 8477–8481.
- [128] M.G. Packard, B.J. Knowlton, Learning and memory functions of the Basal Ganglia, *Annu. Rev. Neurosci.* 25 (2002) 563–593.
- [129] J.G. Pelletier, E. Likhnik, M. Filali, D. Pare, Lasting increases in basolateral amygdala activity after emotional arousal: implications for facilitated consolidation of emotional memories, *Learn. Mem.* 12 (2005) 96–102.
- [130] E.A. Phelps, J.E. LeDoux, Contributions of the amygdala to emotion processing: from animal models to human behavior, *Neuron* 48 (2005) 175–187.
- [131] R.K. Pitman, Post-traumatic stress disorder, hormones, and memory, *Biol. Psychiatry* 26 (1989) 221–223.
- [132] R.K. Pitman, D.L. Delahanty, Conceptually driven pharmacologic approaches to acute trauma, *CNS Spectr.* 10 (2005) 99–106.
- [133] R.K. Pitman, S.P. Orr, Twenty-four hour urinary cortisol and catecholamine excretion in combat-related posttraumatic stress disorder, *Biol. Psychiatry* 27 (1990) 245–247.
- [134] R.K. Pitman, K.M. Sanders, R.M. Zusman, A.R. Healy, F. Cheema, N.B. Lasko, L. Cahill, S.P. Orr, Pilot study of secondary prevention of posttraumatic stress disorder with propranolol, *Biol. Psychiatry* 51 (2002) 189–192.
- [135] P. Putman, E.J. Hermans, H. Koppeschaar, A. van Schijndel, J. van Honk, A single administration of cortisol acutely reduces preconscious attention for fear in anxious young men, *Psychoneuroendocrinology* 32 (2007) 793–802.
- [136] G.L. Quirarte, R. Galvez, B. Roozendaal, J.L. McGaugh, Norepinephrine release in the amygdala in response to footshock and opioid peptidergic drugs, *Brain Res.* 808 (1998) 134–140.
- [137] G.L. Quirarte, B. Roozendaal, J.L. McGaugh, Glucocorticoid enhancement of memory storage involves noradrenergic activation in the basolateral amygdala, *Proc. Natl. Acad. Sci. USA* 94 (1997) 14048–14053.
- [138] G.J. Quirk, D.R. Gehlert, Inhibition of the amygdala: key to pathological states?, *Ann N. Y. Acad. Sci.* 985 (2003) 263–272.
- [139] G.J. Quirk, D. Mueller, Neural mechanisms of extinction learning and retrieval, *Neuropsychopharmacology* 33 (2008) 56–72.
- [140] N. Ramnani, A.M. Owen, Anterior prefrontal cortex: insights into function from anatomy and neuroimaging, *Nat. Rev. Neurosci.* 5 (2004) 184–194.
- [141] R.M. Rapee, R.G. Heimberg, A cognitive-behavioral model of anxiety in social phobia, *Behav. Res. Ther.* 35 (1997) 741–756.
- [142] A. Rashidy-Pour, H. Sadeghi, A.A. Taherain, A.A. Vafaei, Y. Fathollahi, The effects of acute restraint stress and dexamethasone on retrieval of long-term memory in rats: an interaction with opiate system, *Behav. Brain Res.* 154 (2004) 193–198.
- [143] S.L. Rauch, P.J. Whalen, L.M. Shin, S.C. McInerney, M.L. Macklin, N.B. Lasko, S.P. Orr, R.K. Pitman, Exaggerated amygdala response to masked facial stimuli in posttraumatic stress disorder: a functional MRI study, *Biol. Psychiatry* 47 (2000) 769–776.
- [144] J.M. Reul, E.R. De Kloet, Two receptor systems for corticosterone in rat brain: microdistribution and differential occupation, *Endocrinology* 117 (1985) 2505–2511.
- [145] J.M. Revest, F. Di Blasi, P. Kitchener, F. Rouge-Pont, A. Desmedt, M. Turiault, F. Tronche, P.V. Piazza, The MAPK pathway and Egr-1 mediate stress-related behavioral effects of glucocorticoids, *Nat. Neurosci.* 8 (2005) 664–672.
- [146] B. Roozendaal, 1999 Curt P. Richter award. Glucocorticoids and the regulation of memory consolidation, *Psychoneuroendocrinology* 25 (2000) 213–238.
- [147] B. Roozendaal, Stress and memory: opposing effects of glucocorticoids on memory consolidation and memory retrieval, *Neurobiol. Learn. Mem.* 78 (2002) 578–595.
- [148] B. Roozendaal, B. Bohus, J.L. McGaugh, Dose-dependent suppression of adrenocortical activity with metyrapone: effects on emotion and memory, *Psychoneuroendocrinology* 21 (1996) 681–693.
- [149] B. Roozendaal, O. Carmi, J.L. McGaugh, Adrenocortical suppression blocks the memory-enhancing effects of amphetamine and epinephrine, *Proc. Natl. Acad. Sci. USA* 93 (1996) 1429–1433.
- [150] B. Roozendaal, D.J. De Quervain, Glucocorticoid therapy and memory function: lessons learned from basic research, *Neurology* 64 (2005) 184–185.
- [151] B. Roozendaal, D.J. De Quervain, G. Schelling, J.L. McGaugh, A systemically administered beta-adrenoceptor antagonist blocks corticosterone-induced impairment of contextual memory retrieval in rats, *Neurobiol. Learn. Mem.* 81 (2004) 150–154.
- [152] B. Roozendaal, Q.K. Griffith, J. Buranday, D.J. De Quervain, J.L. McGaugh, The hippocampus mediates glucocorticoid-induced impairment of spatial memory retrieval: dependence on the basolateral amygdala, *Proc. Natl. Acad. Sci. USA* 100 (2003) 1328–1333.
- [153] B. Roozendaal, E.L. Hahn, S.V. Nathan, D.J. De Quervain, J.L. McGaugh, Glucocorticoid effects on memory retrieval require concurrent noradrenergic activity in the hippocampus and basolateral amygdala, *J. Neurosci.* 24 (2004) 8161–8169.
- [154] B. Roozendaal, G.K. Hui, I.R. Hui, D.J. Berlau, J.L. McGaugh, N.M. Weinberger, Basolateral amygdala noradrenergic activity mediates corticosterone-induced enhancement of auditory fear conditioning, *Neurobiol. Learn. Mem.* 86 (2006) 249–255.
- [155] B. Roozendaal, J.L. McGaugh, Amygdaloid nuclei lesions differentially affect glucocorticoid-induced memory enhancement in an inhibitory avoidance task, *Neurobiol. Learn. Mem.* 65 (1996) 1–8.
- [156] B. Roozendaal, J.L. McGaugh, The memory-modulatory effects of glucocorticoids depend on an intact stria terminalis, *Brain Res.* 709 (1996) 243–250.
- [157] B. Roozendaal, J.L. McGaugh, Basolateral amygdala lesions block the memory-enhancing effect of glucocorticoid administration in the dorsal hippocampus of rats, *Eur. J. Neurosci.* 9 (1997) 76–83.
- [158] B. Roozendaal, J.L. McGaugh, Glucocorticoid receptor agonist and antagonist administration into the basolateral but not central amygdala modulates memory storage, *Neurobiol. Learn. Mem.* 67 (1997) 176–179.
- [159] B. Roozendaal, J.R. McReynolds, J.L. McGaugh, The basolateral amygdala interacts with the medial prefrontal cortex in regulating glucocorticoid effects on working memory impairment, *J. Neurosci.* 24 (2004) 1385–1392.
- [160] B. Roozendaal, B.T. Nguyen, A.E. Power, J.L. McGaugh, Basolateral amygdala noradrenergic influence enables enhancement of memory consolidation induced by hippocampal glucocorticoid receptor activation, *Proc. Natl. Acad. Sci. USA* 96 (1999) 11642–11647.
- [161] B. Roozendaal, S. Okuda, D.J. De Quervain, J.L. McGaugh, Glucocorticoids interact with emotion-induced noradrenergic activation in influencing different memory functions, *Neuroscience* 138 (2006) 901–910.
- [162] B. Roozendaal, S. Okuda, E.A. van der Zee, J.L. McGaugh, Glucocorticoid enhancement of memory requires arousal-induced noradrenergic activation in the basolateral amygdala, *Proc. Natl. Acad. Sci. USA* 103 (2006) 6741–6746.
- [163] B. Roozendaal, G. Portillo-Marquez, J.L. McGaugh, Basolateral amygdala lesions block glucocorticoid-induced modulation of memory for spatial learning, *Behav. Neurosci.* 110 (1996) 1074–1083.
- [164] B. Roozendaal, G.L. Quirarte, J.L. McGaugh, Glucocorticoids interact with the basolateral amygdala beta-adrenoceptor–cAMP/PKA system in influencing memory consolidation, *Eur. J. Neurosci.* 15 (2002) 553–560.
- [165] B. Roozendaal, G. Schelling, J.L. McGaugh, Corticotropin-releasing factor in the basolateral amygdala enhances memory consolidation via an interaction with the beta-adrenoceptor–cAMP pathway: dependence on glucocorticoid receptor activation, *J. Neurosci.* 28 (2008) 6642–6651.
- [166] B. Roozendaal, C.L. Williams, J.L. McGaugh, Glucocorticoid receptor activation in the rat nucleus of the solitary tract facilitates memory consolidation: involvement of the basolateral amygdala, *Eur. J. Neurosci.* 11 (1999) 1317–1323.
- [167] J.A. Rosenkranz, A.A. Grace, Dopamine-mediated modulation of odour-evoked amygdala potentials during Pavlovian conditioning, *Nature* 417 (2002) 282–287.
- [168] D.R. Rubinow, R.M. Post, R. Savard, P.W. Gold, Cortisol hypersecretion and cognitive impairment in depression, *Arch. Gen. Psychiatry* 41 (1984) 279–283.
- [169] B. Sacchetti, C.A. Lorenzini, E. Baldi, G. Tassoni, C. Bucherelli, Auditory thalamus, dorsal hippocampus, basolateral amygdala, and perirhinal cortex role in the consolidation of conditioned freezing to context and to acoustic conditioned stimulus in the rat, *J. Neurosci.* 19 (1999) 9570–9578.

- [170] A.A. Sajadi, S.A. Samaei, A. Rashidy-Pour, Blocking effects of intra-hippocampal naltrexone microinjections on glucocorticoid-induced impairment of spatial memory retrieval in rats, *Neuropharmacology* 52 (2007) 347–354.
- [171] C. Sandi, S.P. Rose, Corticosterone enhances long-term retention in one-day-old chicks trained in a weak passive avoidance learning paradigm, *Brain Res.* 647 (1994) 106–112.
- [172] R.M. Sapolsky, Glucocorticoids and hippocampal atrophy in neuropsychiatric disorders, *Arch. Gen. Psychiatry* 57 (2000) 925–935.
- [173] G. Schelling, J. Briegel, B. Roozendaal, C. Stoll, H.B. Rothenhauser, H.P. Kapfhammer, The effect of stress doses of hydrocortisone during septic shock on posttraumatic stress disorder in survivors, *Biol. Psychiatry* 50 (2001) 978–985.
- [174] G. Schelling, E. Kilger, B. Roozendaal, D.J. De Quervain, J. Briegel, A. Dagge, H.B. Rothenhauser, T. Krauseneck, G. Nollert, H.P. Kapfhammer, Stress doses of hydrocortisone, traumatic memories, and symptoms of posttraumatic stress disorder in patients after cardiac surgery: a randomized study, *Biol. Psychiatry* 55 (2004) 627–633.
- [175] G. Schelling, M. Richter, B. Roozendaal, H.B. Rothenhauser, T. Krauseneck, C. Stoll, G. Nollert, M. Schmidt, H.P. Kapfhammer, Exposure to high stress in the intensive care unit may have negative effects on health-related quality-of-life outcomes after cardiac surgery, *Crit. Care Med.* 31 (2003) 1971–1980.
- [176] G. Schelling, B. Roozendaal, D.J. De Quervain, Can posttraumatic stress disorder be prevented with glucocorticoids?, *Ann. N. Y. Acad. Sci.* 1032 (2004) 158–166.
- [177] G. Schelling, B. Roozendaal, T. Krauseneck, M. Schmoelz, Q.D. De, J. Briegel, Efficacy of hydrocortisone in preventing posttraumatic stress disorder following critical illness and major surgery, *Ann. N. Y. Acad. Sci.* 1071 (2006) 46–53.
- [178] D. Schoofs, D. Preuss, O.T. Wolf, Psychosocial stress induces working memory impairments in an n-back paradigm, *Psychoneuroendocrinology* 33 (2008) 643–653.
- [179] T.E. Seeman, B.S. McEwen, B.H. Singer, M.S. Albert, J.W. Rowe, Increase in urinary cortisol excretion and memory declines: MacArthur studies of successful aging, *J. Clin. Endocrinol. Metab.* 82 (1997) 2458–2465.
- [180] L.M. Shin, S.L. Rauch, R.K. Pitman, Amygdala, medial prefrontal cortex, and hippocampal function in PTSD, *Ann. N. Y. Acad. Sci.* 1071 (2006) 67–79.
- [181] L.M. Shin, C.I. Wright, P.A. Cannistraro, M.M. Wedig, K. McMullin, B. Martis, M.L. Macklin, N.B. Lasko, S.R. Cavanagh, T.S. Krangel, S.P. Orr, R.K. Pitman, P.J. Whalen, S.L. Rauch, A functional magnetic resonance imaging study of amygdala and medial prefrontal cortex responses to overtly presented fearful faces in posttraumatic stress disorder, *Arch. Gen. Psychiatry* 62 (2005) 273–281.
- [182] T. Smeets, H. Otgaar, I. Candel, O.T. Wolf, True or false? Memory is differentially affected by stress-induced cortisol elevations and sympathetic activity at consolidation and retrieval, *Psychoneuroendocrinology* 33 (2008) 1378–1386.
- [183] A.P. Smith, K.E. Stephan, M.D. Rugg, R.J. Dolan, Task and content modulate amygdala–hippocampal connectivity in emotional retrieval, *Neuron* 49 (2006) 631–638.
- [184] L.M. Soravia, M. Heinrichs, A. Aerni, C. Maroni, G. Schelling, U. Ehlert, B. Roozendaal, D.J. De Quervain, Glucocorticoids reduce phobic fear in humans, *Proc. Natl. Acad. Sci. USA* 103 (2006) 5585–5590.
- [185] L.R. Squire, Memory and the hippocampus: a synthesis from findings with rats, monkeys, and humans, *Psychol. Rev.* 99 (1992) 195–231.
- [186] M.N. Starkman, S.S. Gebarski, S. Berent, D.E. Schteingart, Hippocampal formation volume, memory dysfunction, and cortisol levels in patients with Cushing's syndrome, *Biol. Psychiatry* 32 (1992) 756–765.
- [187] M.N. Starkman, D.E. Schteingart, M.A. Schork, Depressed mood and other psychiatric manifestations of Cushing's syndrome: relationship to hormone levels, *Psychosom. Med.* 43 (1981) 3–18.
- [188] B.A. Strange, R.J. Dolan, Beta-adrenergic modulation of emotional memory-evoked human amygdala and hippocampal responses, *Proc. Natl. Acad. Sci. USA* 101 (2004) 11454–11458.
- [189] J.G. Tasker, S. Di, R. Malcher-Lopes, Minireview: rapid glucocorticoid signaling via membrane-associated receptors, *Endocrinology* 147 (2006) 5549–5556.
- [190] M.S. Tollenaar, B.M. Elzinga, P. Spinhoven, W. Everaerd, Immediate and prolonged effects of cortisol, but not propranolol, on memory retrieval in healthy young men, *Neurobiol. Learn. Mem.* 91 (2009) 23–31.
- [191] S. Tronel, C.M. Alberini, Persistent disruption of a traumatic memory by postretrieval inactivation of glucocorticoid receptors in the amygdala, *Biol. Psychiatry* 62 (2007) 33–39.
- [192] N.C. Tronson, J.R. Taylor, Molecular mechanisms of memory reconsolidation, *Nat. Rev. Neurosci.* 8 (2007) 262–275.
- [193] A.H. van Stegeren, R. Goekoop, W. Everaerd, P. Scheltens, F. Barkhof, J.P. Kuijter, S.A. Rombouts, Noradrenaline mediates amygdala activation in men and women during encoding of emotional material, *Neuroimage* 24 (2005) 898–909.
- [194] A.H. van Stegeren, O.T. Wolf, W. Everaerd, P. Scheltens, F. Barkhof, S.A. Rombouts, Endogenous cortisol level interacts with noradrenergic activation in the human amygdala, *Neurobiol. Learn. Mem.* 87 (2007) 57–66.
- [195] X.Y. Wang, M. Zhao, U.E. Ghitza, Y.Q. Li, L. Lu, Stress impairs reconsolidation of drug memory via glucocorticoid receptors in the basolateral amygdala, *J. Neurosci.* 28 (2008) 5602–5610.
- [196] F. Weis, E. Kilger, B. Roozendaal, D.J. De Quervain, P. Lamm, M. Schmidt, M. Schmolz, J. Briegel, G. Schelling, Stress doses of hydrocortisone reduce chronic stress symptoms and improve health-related quality of life in high-risk patients after cardiac surgery: a randomized study, *J. Thorac. Cardiovasc. Surg.* 131 (2006) 277–282.
- [197] O.T. Wolf, The influence of stress hormones on emotional memory: relevance for psychopathology, *Acta Psychol. (Amst.)* 127 (2008) 513–531.
- [198] O.T. Wolf, A. Convit, P.F. McHugh, E. Kandil, E.L. Thorn, S. De Santi, B.S. McEwen, M.J. de Leon, Cortisol differentially affects memory in young and elderly men, *Behav. Neurosci.* 115 (2001) 1002–1011.
- [199] O.M. Wolkowitz, V.I. Reus, J. Canick, B. Levin, S. Lupien, Glucocorticoid medication, memory and steroid psychosis in medical illness, *Ann. N. Y. Acad. Sci.* 823 (1997) 81–96.
- [200] Y.L. Yang, P.K. Chao, K.T. Lu, Systemic and intra-amygdala administration of glucocorticoid agonist and antagonist modulate extinction of conditioned fear, *Neuropsychopharmacology* 31 (2006) 912–924.
- [201] R. Yehuda, Current status of cortisol findings in post-traumatic stress disorder, *Psychiatr. Clin. North Am.* 25 (2002) 341–368.
- [202] R. Yehuda, Post-traumatic stress disorder, *N. Engl. J. Med.* 346 (2002) 108–114.
- [203] R. Yehuda, L.M. Bierer, Transgenerational transmission of cortisol and PTSD risk, *Prog. Brain Res.* 167 (2008) 121–135.
- [204] R. Yehuda, B. Kahana, K. Binder-Brynes, S.M. Southwick, J.W. Mason, E.L. Giller, Low urinary cortisol excretion in Holocaust survivors with posttraumatic stress disorder, *Am. J. Psychiatry* 152 (1995) 982–986.
- [205] R. Yehuda, J. LeDoux, Response variation following trauma: a translational neuroscience approach to understanding PTSD, *Neuron* 56 (2007) 19–32.
- [206] R. Yehuda, A.C. McFarlane, A.Y. Shalev, Predicting the development of posttraumatic stress disorder from the acute response to a traumatic event, *Biol. Psychiatry* 44 (1998) 1305–1313.
- [207] R. Yehuda, M.H. Teicher, J.R. Seckl, R.A. Grossman, A. Morris, L.M. Bierer, Parental posttraumatic stress disorder as a vulnerability factor for low cortisol trait in offspring of holocaust survivors, *Arch. Gen. Psychiatry* 64 (2007) 1040–1048.
- [208] A.H. Young, B.J. Sahakian, T.W. Robbins, P.J. Cowen, The effects of chronic administration of hydrocortisone on cognitive function in normal male volunteers, *Psychopharmacology (Berl.)* 145 (1999) 260–266.
- [209] E.A. Young, N. Breslau, Cortisol and catecholamines in posttraumatic stress disorder: an epidemiologic community study, *Arch. Gen. Psychiatry* 61 (2004) 394–401.
- [210] M. Zorawski, N.Q. Blanding, C.M. Kuhn, K.S. LaBar, Effects of stress and sex on acquisition and consolidation of human fear conditioning, *Learn. Mem.* 13 (2006) 441–450.