

Figure 15.2
Biochemistry, Seventh Edition
 © 2012 W. H. Freeman and Company

15.23 *Not all alike.* The concentrations of ATP, ADP, and P_i differ with cell type. Consequently, the release of free energy with the hydrolysis of ATP will vary with cell type. Using the following table, calculate the ΔG for the hydrolysis of ATP in liver, muscle, and brain cells. In which cell type is the free energy of ATP hydrolysis most negative?


	ATP (mM)	ADP (mM)	P_i (mM)
Liver	3.5	1.8	5.0
Muscle	8.0	0.9	8.0
Brain	2.6	0.7	2.7

TABLE 15.1 Standard free energies of hydrolysis of some phosphorylated compounds

Compound	kJ mol^{-1}	kcal mol^{-1}
Phosphoenolpyruvate	−61.9	−14.8
1,3-Bisphosphoglycerate	−49.4	−11.8
Creatine phosphate	−43.1	−10.3
ATP (to ADP)	−30.5	− 7.3
Glucose 1-phosphate	−20.9	− 5.0
Pyrophosphate	−19.3	− 4.6
Glucose 6-phosphate	−13.8	− 3.3
Glycerol 3-phosphate	− 9.2	− 2.2

Table 15.1

Biochemistry, Eighth Edition
© 2015 Macmillan Education



Stryer 8th edition: Fig. 15.6

15.14 *A potent donor.* Consider the following reaction:



- (a) Calculate $\Delta G^{\circ'}$ and K'_{eq} at 25°C for this reaction by using the data given in Table 15.1.
- (b) What is the equilibrium ratio of pyruvate to phosphoenolpyruvate if the ratio of ATP to ADP is 10?

Table 15.1 Standard free energies of hydrolysis of some phosphorylated compounds

Compound	kJ mol^{-1}	kcal mol^{-1}
Phosphoenolpyruvate	−61.9	−14.8
1,3-Bisphosphoglycerate	−49.4	−11.8
Creatine phosphate	−43.1	−10.3
ATP (to ADP)	−30.5	− 7.3
Glucose 1-phosphate	−20.9	− 5.0
Pyrophosphate	−19.3	− 4.6
Glucose 6-phosphate	−13.8	− 3.3
Glycerol 3-phosphate	− 9.2	− 2.2




Figure 15.7

Biochemistry, Eighth Edition
© 2015 Macmillan Education

Homework:

solve this yourself

A: Berechnen Sie das ΔG^0 der folgenden Reaktion:



B: Wie muss das Konzentrationsverhältnis $[\text{Kreatin-Phosphat}] / [\text{Kreatin}]$ sein, damit im Gleichgewicht die Konzentration von ATP 12-mal höher als jene von ADP ist? Stellen Sie die entsprechende Gleichung auf und berechnen Sie unter der Annahme von $R = 8.3145 \text{ J K}^{-1} \text{ mol}^{-1}$ und $T = 37^\circ\text{C}$.

Table 15.1 Standard free energies of hydrolysis of some phosphorylated compounds

Compound	kJ mol^{-1}	kcal mol^{-1}
Phosphoenolpyruvate	−61.9	−14.8
1,3-Bisphosphoglycerate	−49.4	−11.8
Creatine phosphate	−43.1	−10.3
ATP (to ADP)	−30.5	− 7.3
Glucose 1-phosphate	−20.9	− 5.0
Pyrophosphate	−19.3	− 4.6
Glucose 6-phosphate	−13.8	− 3.3
Glycerol 3-phosphate	− 9.2	− 2.2

ATP use and regeneration






Figure 15.8

Biochemistry, Eighth Edition
© 2015 Macmillan Education


Most energy \longrightarrow Least energy




Methane



Methanol



Formaldehyde



Formic acid



Carbon dioxide

$\Delta G^{\circ'}$ oxidation
(kJ mol⁻¹)

-820

-703

-523

-285

0

$\Delta G^{\circ'}$ oxidation
(kcal mol⁻¹)

-196

-168

-125


-68

0


Figure 15.9

Biochemistry, Eighth Edition

© 2015 Macmillan Education



Unnumbered 15 p435a
Biochemistry, Eighth Edition
© 2015 Macmillan Education



Unnumbered 15 p435b
 Biochemistry, Eighth Edition
 © 2015 Macmillan Education

Flavine adenine dinucleotide (FAD – FADH₂)




Figure 15.14
Biochemistry, Eighth Edition
© 2015 Macmillan Education

Flavine adenine dinucleotide (FAD – FADH₂)




Figure 15.15

Biochemistry, Eighth Edition

© 2015 Macmillan Education

TABLE 15.2 Some activated carriers in metabolism

Carrier molecule in activated form	Group carried	Vitamin precursor
ATP	Phosphoryl	
NADH and NADPH	Electrons	Nicotinate (niacin) (vitamin B ₃)
FADH ₂	Electrons	Riboflavin (vitamin B ₂)
FMNH ₂	Electrons	Riboflavin (vitamin B ₂)
Coenzyme A	Acyl	Pantothenate (vitamin B ₅)
Lipoamide	Acyl	
Thiamine pyrophosphate	Aldehyde	Thiamine (vitamin B ₁)
Biotin	CO ₂	Biotin (vitamin B ₇)
Tetrahydrofolate	One-carbon units	Folate (vitamin B ₉)
S-Adenosylmethionine	Methyl	
Uridine diphosphate glucose	Glucose	
Cytidine diphosphate diacylglycerol	Phosphatidate	
Nucleoside triphosphates	Nucleotides	

Note: Many of the activated carriers are coenzymes that are derived from water-soluble vitamins.

Table 15.2

Biochemistry, Eighth Edition
© 2015 Macmillan Education




Figure 15.20
Biochemistry, Eighth Edition
© 2015 Macmillan Education

FIGURE 15.20 Energy charge regulates metabolism. When the energy charge is high, ATP inhibits the relative rates of a typical ATP-generating (catabolic) pathway and stimulates the typical ATP-utilizing (anabolic) pathway.