

5-HT and depression: is the glass half-full?

Trevor Sharp¹ and Philip J Cowen²

Mood disorders such as major depression are common illnesses with considerable morbidity and significant mortality. A long-standing theory is that a breakdown in brain serotonin (5-hydroxytryptamine; 5-HT) signalling is critically involved in the symptoms and drug treatment of clinical depression. However, the nature of this 5-HT defect has proved to be frustratingly elusive, and it remains unclear how the 5-HT signalling effects of antidepressant drugs might alter neuropsychological mechanisms to bring about relief of depressed mood. This article highlights recent discoveries that advance our understanding of how 5-HT-evoked changes at molecular, cellular and neuropsychological levels might interact to alleviate the symptoms of clinical depression.

Addresses

¹ University Department of Pharmacology, Mansfield Road, Oxford OX1 3QT, UK

² University Department of Psychiatry, Warneford Hospital, Oxford OX3 7JX, UK

Corresponding author: Sharp, Trevor (trevor.sharp@pharm.ox.ac.uk)

Current Opinion in Pharmacology 2011, **11**:45–51

This review comes from a themed issue on
 Neurosciences
 Edited by Nicholas Barnes

Available online 4th March 2011

1471-4892/\$ – see front matter
 © 2011 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.coph.2011.02.003](https://doi.org/10.1016/j.coph.2011.02.003)

Introduction

Major depression is a common and often severe psychiatric condition characterised by a complex syndrome of affective, vegetative and cognitive symptoms that show a relapse-remitting course over months and often years. All this adds up to a source of considerable morbidity and significant mortality. The World Health Organization forecasts that if current demographic trends continue, unipolar depression will be the greatest source of disability worldwide by the year 2030 [1]. Despite this worrying prediction, modern drug treatments are slow to act, with often poorly tolerated acute adverse effects including anxiety and insomnia, and up to 50% of patients fail to experience an adequate therapeutic response even after a treatment switch [2]. Moreover, only one novel antidepressant drug has been developed in the last 10 years, and major pharmaceutical companies are closing antidepressant drug development programmes [3].

The idea that the symptoms of depression are underpinned by a deficiency in brain 5-HT stretch back over 40 years to the discovery that the first generation of antidepressant drugs blocked 5-HT reuptake or metabolism as part of their pharmacological effect. Since then studies of the link between 5-HT and depression have continued to provide a rich source of discoveries including the development of selective 5-HT reuptake (SSRIs), which are currently the first choice antidepressant drug treatment. However, important questions remain. For instance, even though the interaction of an SSRI with the binding site of the 5-HT transporter (5-HTT; SERT) protein is now known with single amino acid resolution [4], it is unknown how the signalling events triggered by this interaction eventually translate into changes in mood. Equally, whilst the therapeutic effect of SSRIs illustrates how manipulations of 5-HT can have striking effects on mood [5•], it is unclear whether changes in brain 5-HT function are causally linked to the symptoms of depression. This article highlights recent discoveries that impinge on these and related questions.

5-HT abnormalities in depression

In the 1960s and 1970s it was reported that depressed patients had reduced 5-HT binding sites on platelets and low 5-HT metabolite levels in cerebrospinal fluid. These findings were not always reliable [5•], but they provided some of the first evidence for 5-HT changes in depression. This idea was strengthened by later more consistent findings of blunted neuroendocrine responses to 5-HT-selective pharmacological challenges in depressed patients. Interestingly, this effect was observed in patients with a history of depression even when recovered and off drug treatment [6], suggesting a link between reduced 5-HT function and depression vulnerability, and not just the acute depressive state [5•]. The neuroendocrine data accord with more recent findings from positron emission tomography (PET) studies [7], of a widespread decrease of 5-HT_{1A} receptors in the brains of acutely depressed patients as well as patients who have recovered [8,9].

Other PET studies have investigated the 5-HT_{2A} receptor and 5-HTT in depressed patients, but findings to date are inconsistent. If anything, 5-HT_{2A} receptor binding is increased in the cortex of both depressed and recovered subjects, and this neurochemical change correlates with measures of negative beliefs about the self and the world [10,11]. In contrast, some (although not all) imaging studies suggest reduced 5-HTT binding in brain stem regions, which resolves with clinical recovery [12,13]. However, the latter data raise the question of

why depressed patients respond to treatment with drugs like SSRIs that cause a further decrease in 5-HTT availability. It is possible that the drugs are in some way facilitating a prior neuroadaptive response to lowered 5-HT levels; however, this explanation does not seem altogether satisfactory.

The above imaging and neuroendocrine findings combined, provide tantalising evidence that impaired brain 5-HT function is indeed linked to depression and depression vulnerability. These data can be taken as evidence to suggest that the source of the 5-HT deficiency is at the level of postsynaptic 5-HT receptors as well as 5-HT release. In the latter context it should be noted that there are frequent reports of low circulating levels of the 5-HT precursor, L-tryptophan, in severely depressed patients [14]. This deficit has been linked to immune activation that is believed to often accompany the depressive state, and that leads to induction of indoleamine dioxygenase and increased tryptophan degradation [15].

In accord with the above findings, a significant body of data suggests that acute lowering of tryptophan using a tryptophan-free amino acid mixture, elicits a striking return of clinical depression in a high proportion of recovered depressed patients withdrawn from medication [16[•]]. These data do not speak directly to the question of whether depressed patients have low 5-HT release but certainly suggest that maintaining synaptic 5-HT levels is critical for the maintenance of mood in vulnerable individuals, and specifically those who have previously experienced depression and may have an already compromised 5-HT system. This idea accords with recent evidence that deep brain stimulation of the subthalamic nucleus that acutely decreases 5-HT transmission in experimental models [17], can elicit low mood in Parkinson's disease patients who are known to demonstrate preexisting 5-HT deficits at several levels [18].

Interestingly, in people at high familial risk of depression who have not themselves suffered from depression, tryptophan depletion produces a detectable but much milder lowering of mood [16[•]]. This observation suggests that the more profound clinical relapse in mood that follows tryptophan depletion in individuals with a history of depression, may be due to changes that occur as a result of recurrent illness. The nature of these changes, and whether they are best conceptualised at a psychological or neural level, are not known.

The key question of whether 5-HT release is decreased in depression may soon be tested with the development of PET imaging methods to monitor 5-HT release using the radioligand displacement approach, which has been so successful in the dopamine field [19]. There is an ongoing, intensive search for PET radiotracers that are sensitive to

changes in synaptic 5-HT availability, with certain 5-HT_{1A} receptor agonist and 5-HT_{1B} receptor antagonist ligands already showing promise [20[•]], and other putative 5-HT-sensitive PET ligands are on the horizon. For example, studies in depression using the newly available 5-HT₄ receptor PET ligand [¹¹C]SB207145 [21] will be interesting given recent preclinical evidence that 5-HT₄ receptors adapt to enduring (although not acute) changes in synaptic 5-HT availability [22,23].

5-HT genes and depression

Studies of major depression in twins and families have shown moderate to high heritability, and much effort has been made to identify susceptibility genes linked to 5-HT. The 5-HTT gene (slc6a4) is one of the most intensively investigated genetic risk factors for depression. Levels of the 5-HTT vary up to sevenfold between individuals in the human population (e.g. [24]) and this may be due in part to the existence of functional 5-HTT gene variants, many of which continue to be discovered [25^{••},26]. Of particular interest is a 44 bp insertion/deletion polymorphism in the 5-HTT gene-linked polymorphic region (5-HTTLPR) of the 5-HTT gene promoter. This generates a low 5-HTT-expressing allele (s/s) that was found to be associated with factors linked to depression such as neuroticism [7] and increased responsiveness to fearful stimuli [27], as well as an actual increased risk of depression when accompanied by stressful life events in adult as well as early life [28^{••},29[•]].

Interpretation of these findings is complicated by failed replications [30,31[•]], and the view that the effect of individual polymorphisms on the phenotypic expression of complex traits is likely to be small [32]. In addition, PET studies do not indicate that 5-HTT genotype is linked to the density of 5-HTT sites *in vivo* [33]. Intriguingly however, investigations using genetic animal models observe striking phenotypic effects of altered expression of the 5-HTT. Specifically, 5-HTT knockout mice demonstrate anxiety-like behavioural phenotypes whereas 5-HTT overexpressing mice show the opposite [25^{••},34]. Moreover, heterozygote 5-HTT knockout mice lack a noticeable phenotype under baseline conditions but demonstrate a high anxiety phenotype in adulthood following exposure to environmental early life stressors [35,36]. These findings bear a close resemblance to human gene–phenotype association data [28^{••}] and add further support to the idea that an interaction between the 5-HTT gene and early life events are a critical determinant of depression vulnerability [37^{••}].

The neural mechanisms by which environmental adversity in early life might interact with the 5-HTT gene to influence emotionality in adulthood are not known, but likely to include epigenetic programming of gene expression. In this respect it is interesting that the promoter

region of the 5-HTT gene is rich in DNA methylation sites and that changes in the methylation status of these sites have been associated with altered stress sensitivity [38[•],39].

Emerging evidence also associates depression vulnerability as well as antidepressant treatment response with a single nucleotide polymorphism (C(−1019)G) in the promoter region of the 5-HT_{1A} receptor gene, which determines the binding of certain gene repressors and is thereby functional [40[•]]. Interestingly, the putative risk (G/G) allele was recently modelled in mice through a targeted mutation that increased 5-HT_{1A} receptor expression specifically in midbrain raphe neurons [41^{••}]. These mice demonstrated a suppression of 5-HT cell firing due to increased inhibitory 5-HT_{1A} autoreceptor tone. In addition, the mice showed increased sensitivity to stress as well as resistance to behavioural effects of SSRI treatment, thereby recapitulating the phenotype predicted by the human gene-association studies.

Molecular and cellular effects of 5-HT-targeted antidepressants

Although SSRIs rapidly enter the brain and block 5-HT reuptake, as evident in human PET studies of 5-HTT occupancy [42], a course of treatment of several weeks is required for the full antidepressant response. One early hypothesis put forward in the 1980s to explain this paradox was an adaptive desensitization of brain 5-HT₂ receptors and β -adrenoceptors, but this theory was later widely discarded in favour of one that emphasised the importance of 5-HT autoreceptor desensitization [43]. The latter theory posited that **acute inhibition of 5-HT reuptake or metabolism evokes a rise in extracellular 5-HT which is immediately sensed by autoreceptors located on the somatodendrites (5-HT_{1A}) and nerve terminals (5-HT_{1B}) of 5-HT neurons to trigger a fall in 5-HT cell firing and synaptic 5-HT: evidence suggests that with continued antidepressant treatment, 5-HT_{1A} and 5-HT_{1B} autoreceptors desensitize allowing synaptic 5-HT to rise. Recent studies have identified additional mechanisms involved in the feedback control of 5-HT neurons that utilise postsynaptic 5-HT receptors including 5-HT_{2A/2C} receptors [44[•]], which may also contribute to the acute adaptive response of 5-HT neurons to antidepressant treatment.**

The 5-HT autoreceptor desensitization theory instigated drug development programmes that successfully identified molecules with combined 5-HT reuptake/5-HT_{1A/1B} receptor antagonist properties [45]. Although little is currently known about the actions of these dual and triple action drugs in humans, clinical trials have demonstrated an overall beneficial clinical effect of using the 5-HT_{1A}/ β -adrenoceptor ligand pindolol as an adjunct to SSRI treatment in terms of speed of onset of clinical effect [46]. Preclinical and clinical evidence of the SSRI augmenting

properties of 5-HT₂ receptor antagonists (including risperidone) might link to 5-HT₂ receptor feedback mechanisms. For example, in patients who have failed to respond to SSRIs, the augmentation strategy with the best evidence base is the addition of atypical antipsychotic drugs at low doses that would preferentially antagonise 5-HT_{2A/2C} receptors [47].

Recent evidence suggests that the response of 5-HT feedback mechanisms to SSRI administration is just part of a series of therapeutically relevant neuroadaptive events that occur during a course of antidepressant treatment. Current thinking is that increased synaptic 5-HT (presumably following 5-HT autoreceptor desensitization) activates a downstream gene programme that leads to enhanced neuronal plasticity which has failed because of the adverse effects of stress and other environmental and genetic factors [48^{••}]. Much data, including that from genome wide analyses (e.g. see [49]), suggest that the gene programme is complex and likely includes a co-ordinated pattern of expression of many tens if not hundreds of genes involved in multiple processes ranging from synapse formation and strengthening to even neurovascular support. The net effect of these changes is thought to be an activity-dependent structural realignment and functional repair of neural circuitry, which is then better able to adapt to environmental challenges [50^{••},51].

The large diversity of 5-HT-sensitive genes influenced by a course of antidepressant treatment makes it difficult to attribute weight of importance to individual genes and their products, and specific downstream effects. However, there are much data in favour of a key role for brain-derived neurotrophic factor (BDNF) and its receptor TrkB in the actions of antidepressant drug treatment, if not depression pathophysiology [51,52[•]]. Equally, BDNF is a known mediator of neurogenesis, and there is a strong case for the importance of increased neurogenesis in the action of SSRIs although less so for other antidepressant drug classes [53^{••},54].

Identification of the 5-HT receptor subtypes and their signalling pathways that underpin the gene effects of 5-HT-targeted antidepressants is important to guide antidepressant drug development programmes. 5-HT receptor subtypes positively coupled to adenylate cyclase (5-HT₄, 5-HT₆, 5-HT₇) or stimulation of Ca²⁺-dependent kinases (5-HT_{2A}, 5-HT_{2C}) are likely to activate the transcriptional regulator CREB and also MAP kinase, both of which are considered key nodal points in the signalling pathways underlying the gene expression and behavioural effects of SSRI and other antidepressants [55–57]. Interestingly, recent data show that 5-HT₄ and 5-HT₆ receptor agonists demonstrate early onset antidepressant-like molecular and behavioural effects in preclinical models [58,59]. Moreover, there are ongoing clinical trials into the antidepressant potential of psychedelic drug, psilocybin, reinforced by

evidence that a 5-HT_{2A} receptor agonist action may evoke changes in neural plasticity to produce long-lasting improvement in mood [60*].

5-HT and depression: neuropsychological approaches

Recently the idea that 5-HT might mediate adaptive changes in neural plasticity can be considered in the context of an emerging neuropsychological theory of antidepressant drug action (Figure 1). The latter theory emphasises that altered emotional processing is critical to the actions of drug treatments rather than a direct elevation in mood *per se*. In particular, it is proposed that at the neuropsychological level, antidepressants work by inducing a shift away from a cognitive state of negative affective bias, comprising negatively biased attention, memory and appraisal of internal and external emotional stimuli.

That 5-HT is important to this switch in emotional processing is supported by several lines of evidence, including findings that tryptophan depletion caused negative biases in the recognition of facial expressions [61] whereas the SSRI citalopram had the opposite effect [62]. Both effects occur independent of changes in objective or subjective mood. Functional imaging studies suggest that the changes

in biases produced by altered 5-HT neurotransmission are exerted via effects on limbic circuitry involving the amygdala [63**], which plays a key role in the very early non-conscious appraisal of negative emotional stimuli.

Interestingly, the ability of citalopram and other antidepressants to evoke a positive bias in emotional processing is detectable after a single dose [63**]. These observations have led to the idea that at the neuropsychological level antidepressants cause an immediate adjustment in negative affective bias and that this provides a platform for the relearning of new positive emotional associations brought about by the gradual exposure to social cues and reinforcement in this transformed emotional world [63**,64]. Over time this would be expected to lead to conscious improvement in subjective mood as new, positive, emotional contingencies are learned and experienced.

It is noteworthy that acute SSRI administration is also associated with enhanced startle responses and fear recognition [65]. These observations are interpreted as an early increase in threat processing, which is highly relevant to clinical findings of increased anxiety early during a course of SSRI treatment before the onset of anxiolytic effect (for further discussion see [63**]).

Figure 1




Illustration of the proposed sequence of 5-HT-mediated, adaptive neurobiological and neuropsychological processes that occur during a course of several weeks treatment with 5-HT-targeted antidepressant drugs (inhibitors of 5-HT reuptake and metabolism), leading up to the full therapeutic effect. The induction of positive emotional bias occurs after acute treatment and forms the basis for the gradual relearning of positive emotional associations. Similarly, activation and then desensitization of 5-HT feedback occur very early in the course of treatment and are required for the downstream changes in gene expression and increased neural plasticity. As indicated by the shadowed box, it is envisaged that the increase in neural plasticity and formation of neural circuits occurs within emotional processing networks to allow positive emotional stimuli (e.g. social cues) to act and to be relearned and bring about improved mood.

The effects of SSRIs on positive emotional bias and relearning, have an intriguing similarity to cognitive therapy approaches to depression where therapists use reflective discussion with patients to enable them to reframe negative perceptions and appraisals in a more 'realistic' that is, more positive, light. For this reason cognitive therapists are fond of quoting the Greek stoic philosopher, Epictetus (55–135 AD) who observed that, 'Men are disturbed not by things, but by the view which they take of them.' Of course, in contrast to drug treatment, cognitive therapy involves explicit and conscious reframing of experience presumably involving cortical mechanisms, whereas the ability of SSRIs to produce positive emotional bias appears to occur implicitly, probably through actions on non-conscious emotional processing at a limbic level.

The neural plasticity and neuropsychological hypotheses of antidepressant action are not mutually exclusive (Figure 1). Indeed, increased neural plasticity within critical neural circuits might provide the neural substrate through which acute increases in positive emotional processing lead to the relearning of positive emotional associations that ultimately bring about improvements in mood. This idea resonates with recent findings that repeated SSRI treatment brought about the functional (BDNF-dependent) recovery of circuitry of the visual cortex when combined with a visual stimulus operating through this circuitry [66•]. Thus, SSRI treatment may activate plasticity within visual circuits to allow a visual stimulus to act in the same way that SSRIs might activate plasticity within emotional processing circuits to allow a positive emotional stimulus to act and to be relearned. The application of animal models of negative bias [67•] should help provide an experimental means to test this hypothesis.

Conclusion

The evidence that brain 5-HT can influence the expression of depressive symptomatology is very strong, although impaired 5-HT function would appear to be neither necessary nor sufficient to cause clinical depression. In some patients increasing brain 5-HT function through blockade of the 5-HTT (SSRI treatment) seems sufficient to promote clinical recovery, though preclinical studies show that this simple pharmacological action is accompanied by a complex programme of molecular and cellular changes. The ability of SSRIs to both facilitate mechanisms of neural plasticity and remediate negative emotional biases, allows what might be an important convergence of neurobiological and psychological explanations of the role of 5-HT in antidepressant action and depression. Thus, the ability of 5-HT to alter both emotional experience and emotional learning could give this neurotransmitter a pivotal role in the treatment and pathophysiology of emotional disorders.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Organisation WH: *The global burden of disease: 2004 update*. 2008;. Part 4: 51.
2. Rush AJ, Trivedi MH, Wisniewski SR, Nierenberg AA, Stewart JW, Warden D, Niederehe G, Thase ME, Lavori PW, Lebowitz BD *et al.*: **Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report**. *Am J Psychiatry* 2006, **163**:1905–1917.
3. Blier P: **The well of novel antidepressants: running dry**. *J Psychiatry Neurosci* 2010, **35**:219–220.
4. Andersen J, Olsen L, Hansen KB, Taboureau O, Jorgensen FS, Jorgensen AM, Bang-Andersen B, Egebjerg J, Stromgaard K, Kristensen AS: **Mutational mapping and modeling of the binding site for (S)-citalopram in the human serotonin transporter**. *J Biol Chem* 2010, **285**:2051–2063.
5. Cowen PJ: **Serotonin and depression: pathophysiological mechanism or marketing myth?** *Trends Pharmacol Sci* 2008, **29**:433–436.
- A critical account of the role of 5-HT in the symptoms of depression.
6. Bhagwagar Z, Whale R, Cowen PJ: **State and trait abnormalities in serotonin function in major depression**. *Br J Psychiatry* 2002, **180**:24–28.
7. Lesch KP, Bengel D, Heils A, Sabol SZ, Greenberg BD, Petri S, Benjamin J, Muller CR, Hamer DH, Murphy DL: **Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region**. *Science* 1996, **274**:1527–1531.
8. Sargent PA, Kjaer KH, Bench CJ, Rabiner EA, Messa C, Meyer J, Gunn RN, Grasby PM, Cowen PJ: **Brain serotonin1A receptor binding measured by positron emission tomography with [11C]WAY-100635: effects of depression and antidepressant treatment**. *Arch Gen Psychiatry* 2000, **57**:174–180.
9. Drevets WC, Thase ME, Moses-Kolko EL, Price J, Frank E, Kupfer DJ, Mathis C: **Serotonin-1A receptor imaging in recurrent depression: replication and literature review**. *Nucl Med Biol* 2007, **34**:865–877.
10. Meyer JH, Wilson AA, Rusjan P, Clark M, Houle S, Woodside S, Arrowood J, Martin K, Colleton M: **Serotonin2A receptor binding potential in people with aggressive and violent behaviour**. *J Psychiatry Neurosci* 2008, **33**:499–508.
11. Bhagwagar Z, Hinz R, Taylor M, Fancy S, Cowen P, Grasby P: **Increased 5-HT(2A) receptor binding in euthymic, medication-free patients recovered from depression: a positron emission study with [(11C)MDL 100,907]**. *Am J Psychiatry* 2006, **163**:1580–1587.
12. Selvaraj S, Venkatesha Murthy N, Bhagwagar Z, Bose SK, Hinz R, Grasby PM, Cowen PJ: **Diminished brain 5-HT transporter binding in major depression: a positron emission tomography study with [(11C)DASB]**. *Psychopharmacology (Berl)* 2011, **213**:555–562.
13. Bhagwagar Z, Murthy N, Selvaraj S, Hinz R, Taylor M, Fancy S, Grasby P, Cowen P: **5-HTT binding in recovered depressed patients and healthy volunteers: a positron emission tomography study using [11C]DASB**. *Am J Psychiatry* 2007, **164**:1858–1865.
14. Cowen PJ, Parry-Billings M, Newsholme EA: **Decreased plasma tryptophan levels in major depression**. *J Affect Disord* 1989, **16**:27–31.
15. Maes M, Leonard BE, Myint AM, Kubera M, Verkerk R: **The new '5-HT' hypothesis of depression: Cell-mediated immune activation induces indoleamine 2,3-dioxygenase, which leads to lower plasma tryptophan and an increased synthesis of detrimental tryptophan catabolites (TRYCATs), both of which contribute to the onset of depression**. *Prog Neuropsychopharmacol Biol Psychiatry*, in press.

16. Ruhe HG, Mason NS, Schene AH: **Mood is indirectly related to serotonin, norepinephrine and dopamine levels in humans: a meta-analysis of monoamine depletion studies.** *Mol Psychiatry* 2007, **12**:331-359.
A thorough analysis of studies investigating the influence of monoamine depletion in drug-free healthy controls and patients with depression or at risk.
17. Temel Y, Boothman LJ, Blokland A, Magill PJ, Steinbusch HW, Visser-Vandewalle V, Sharp T: **Inhibition of 5-HT neuron activity and induction of depressive-like behavior by high-frequency stimulation of the subthalamic nucleus.** *Proc Natl Acad Sci U S A* 2007, **104**:17087-17092.
18. Tan SKH, Hartung H, Sharp T, Temel Y: **Serotonin-dependent depression in Parkinson's disease: a role for the subthalamic nucleus?** *Neuropharmacology*, in press.
19. Laruelle M: **Imaging synaptic neurotransmission with in vivo binding competition techniques: a critical review.** *J Cereb Blood Flow Metab* 2000, **20**:423-451.
20. Paterson LM, Tyacke RJ, Nutt DJ, Knudsen GM: **Measuring endogenous 5-HT release by emission tomography: promises and pitfalls.** *J Cereb Blood Flow Metab* 2010, **30**:1682-1706.
An up-to-date account of progress in the application of PET radiotracers to detect 5-HT release.
21. Marnier L, Gillings N, Madsen K, Erritzoe D, Baare WF, Svarer C, Hasselbalch SG, Knudsen GM: **Brain imaging of serotonin 4 receptors in humans with [11C]SB207145-PET.** *Neuroimage* 2010, **50**:855-861.
22. Licht CL, Marcussen AB, Wegener G, Overstreet DH, Aznar S, Knudsen GM: **The brain 5-HT₄ receptor binding is down-regulated in the Flinders Sensitive Line depression model and in response to paroxetine administration.** *J Neurochem* 2009, **109**:1363-1374.
23. Vidal R, Valdizan EM, Mostany R, Pazos A, Castro E: **Long-term treatment with fluoxetine induces desensitization of 5-HT₄ receptor-dependent signalling and functionality in rat brain.** *J Neurochem* 2009, **110**:1120-1127.
24. Lundberg J, Halldin C, Farde L: **Measurement of serotonin transporter binding with PET and [11C]MADAM: a test-retest reproducibility study.** *Synapse* 2006, **60**:256-263.
25. Murphy DL, Lesch KP: **Targeting the murine serotonin transporter: insights into human neurobiology.** *Nat Rev Neurosci* 2008, **9**:85-96.
A comprehensive review of the 5-HT transporter gene and the impact of variation of this gene in humans and animal models.
26. Ehli EA, Hu Y, Lengyel-Nelson T, Hudziak JJ, Davies GE: **Identification and functional characterization of three novel alleles for the serotonin transporter-linked polymorphic region.** *Mol Psychiatry* 2011.
27. Pezawas L, Meyer-Lindenberg A, Drabant EM, Verchinski BA, Munoz KE, Kolachana BS, Egan MF, Mattay VS, Hariri AR, Weinberger DR: **5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: a genetic susceptibility mechanism for depression.** *Nat Neurosci* 2005, **8**:828-834.
28. Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, McClay J, Mill J, Martin J, Braithwaite A *et al.*: **Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene.** *Science* 2003, **301**:386-389.
One of the first studies reporting the impact of 5-HT transporter gene x environment interactions on depression risk.
29. Caspi A, Hariri AR, Holmes A, Uher R, Moffitt TE: **Genetic sensitivity to the environment: the case of the serotonin transporter gene and its implications for studying complex diseases and traits.** *Am J Psychiatry* 2010, **167**:509-527.
A critical analysis of human 5-HT transporter gene-phenotype associations.
30. Risch N, Herrell R, Lehner T, Liang KY, Eaves L, Hoh J, Griem A, Kovacs M, Ott J, Merikangas KR: **Interaction between the serotonin transporter gene (5-HTTLPR), stressful life events, and risk of depression: a meta-analysis.** *JAMA* 2009, **301**:2462-2471.
31. Uher R, McGuffin P: **The moderation by the serotonin transporter gene of environmental adversity in the etiology of depression: 2009 update.** *Mol Psychiatry* 2010, **15**:18-22.
A critical analysis of human 5-HT transporter gene-phenotype associations to complement the above review by Caspi *et al.*
32. Flint J, Munafò MR: **The endophenotype concept in psychiatric genetics.** *Psychol Med* 2007, **37**:163-180.
33. Murthy NV, Selvaraj S, Cowen PJ, Bhagwagar Z, Riedel WJ, Peers P, Kennedy JL, Sahakian BJ, Laruelle MA, Rabiner EA *et al.*: **Serotonin transporter polymorphisms (SLC6A4 insertion/deletion and rs25531) do not affect the availability of 5-HTT to [11C] DASB binding in the living human brain.** *Neuroimage* 2010, **52**:50-54.
34. Line SJ, Barkus C, Coyle C, Jennings KA, Deacon RM, Lesch KP, Sharp T, Bannerman DM: **Opposing alterations in anxiety and species-typical behaviours in serotonin transporter overexpressor and knockout mice.** *Eur Neuropsychopharmacol* 2011, **21**:108-116.
35. Bartolomucci A, Carola V, Pascucci T, Puglisi-Allegra S, Cabib S, Lesch KP, Parmigiani S, Palanza P, Gross C: **Increased vulnerability to psychosocial stress in heterozygous serotonin transporter knockout mice.** *Dis Model Mech* 2010, **3**:459-470.
36. Carola V, Frazzetto G, Pascucci T, Audero E, Puglisi-Allegra S, Cabib S, Lesch KP, Gross C: **Identifying molecular substrates in a mouse model of the serotonin transporter x environment risk factor for anxiety and depression.** *Biol Psychiatry* 2008, **63**:840-846.
37. Ansoorge MS, Hen R, Gingrich JA: **Neurodevelopmental origins of depressive disorders.** *Curr Opin Pharmacol* 2007, **7**:8-17.
An interesting account of how deficits in the maturation of brain circuits might impact on emotional function to increase the risk for depression later in life.
38. van IMH, Caspers K, Bakermans-Kranenburg MJ, Beach SR, Philibert R: **Methylation matters: interaction between methylation density and serotonin transporter genotype predicts unresolved loss or trauma.** *Biol Psychiatry* 2010, **68**:405-407.
One of the first studies reporting an association between adverse life events and methylation status of the 5-HT transporter gene.
39. Kinnally EL, Capitanio JP, Leibel R, Deng L, LeDuc C, Haghighi F, Mann JJ: **Epigenetic regulation of serotonin transporter expression and behavior in infant rhesus macaques.** *Genes Brain Behav* 2010, **9**:575-582.
40. Albert PR, Francois BL: **Modifying 5-HT_{1A} receptor gene expression as a new target for antidepressant therapy.** *Front Neurosci* 2010, **4**:1-7.
A review the impact of polymorphic variation and transcriptional regulation of 5-HT_{1A} receptor gene, on antidepressant mechanisms.
41. Richardson-Jones JW, Craig CP, Guiard BP, Stephen A, Metzger KL, Kung HF, Gardier AM, Dranovsky A, David DJ, Beck SG *et al.*: **5-HT_{1A} autoreceptor levels determine vulnerability to stress and response to antidepressants.** *Neuron* 2010, **65**:40-52.
An interesting study demonstrating the power of targeted manipulations of the 5-HT_{1A} receptor gene to investigate the influence of a putative 5-HT_{1A} risk allele on stress vulnerability and antidepressant response.
42. Meyer JH, Wilson AA, Sagrati S, Hussey D, Carella A, Potter WZ, Ginovart N, Spencer EP, Cheek A, Houle S: **Serotonin transporter occupancy of five selective serotonin reuptake inhibitors at different doses: an [11C]DASB positron emission tomography study.** *Am J Psychiatry* 2004, **161**:826-835.
43. Blier P, de Montigny C: **Current advances and trends in the treatment of depression.** *Trends Pharmacol Sci* 1994, **15**:220-226.
44. Sharp T, Boothman L, Raley J, Queree P: **Important messages in the 'post': recent discoveries in 5-HT neurone feedback control.** *Trends Pharmacol Sci* 2007, **28**:629-636.
A review of current knowledge of the pharmacology and physiology of 5-HT feedback mechanisms.
45. Millan MJ: **Dual- and triple-acting agents for treating core and co-morbid symptoms of major depression: novel concepts, new drugs.** *Neurotherapeutics* 2009, **6**:53-77.

46. Whale R, Terao T, Cowen P, Freemantle N, Geddes J: **Pindolol augmentation of serotonin reuptake inhibitors for the treatment of depressive disorder: a systematic review.** *J Psychopharmacol* 2010, **24**:513-520.
47. Nelson JC, Papakostas GI: **Atypical antipsychotic augmentation in major depressive disorder: a meta-analysis of placebo-controlled randomized trials.** *Am J Psychiatry* 2009, **166**:980-991.
48. Krishnan V, Nestler EJ: **The molecular neurobiology of depression.** *Nature* 2008, **455**:894-902.
An interesting review of the neurobiology of depression that combines insights from clinical studies and animal models.
49. Sillaber I, Panhuysen M, Henniger MS, Ohl F, Kuhne C, Putz B, Pohl T, Deussing JM, Paez-Pereda M, Holsboer F: **Profiling of behavioral changes and hippocampal gene expression in mice chronically treated with the SSRI paroxetine.** *Psychopharmacology (Berl)* 2008, **200**:557-572.
50. Castren E: **Is mood chemistry?** *Nat Rev Neurosci* 2005, **6**:241-246.
An lucid account of the neural plasticity hypothesis of depression and antidepressant action.
51. Castren E, Rantamaki T: **The role of BDNF and its receptors in depression and antidepressant drug action: reactivation of developmental plasticity.** *Dev Neurobiol* 2010, **70**:289-297.
52. Groves JO: **Is it time to reassess the BDNF hypothesis of depression?** *Mol Psychiatry* 2007, **12**:1079-1088.
An important critical analysis of the role of BDNF in the cause and treatment of depression.
53. Santarelli L, Saxe M, Gross C, Surget A, Battaglia F, Dulawa S, Weisstaub N, Lee J, Duman R, Arancio O *et al.*: **Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants.** *Science* 2003, **301**:805-809.
This study provided some of the first evidence that an increase in neurogenesis in hippocampus mediates behavioural effects SSRI administration.
54. David DJ, Samuels BA, Rainer Q, Wang JW, Marsteller D, Mendez I, Drew M, Craig DA, Guindard BP, Guilloux JP *et al.*: **Neurogenesis-dependent and -independent effects of fluoxetine in an animal model of anxiety/depression.** *Neuron* 2009, **62**:479-493.
55. Tanis KQ, Duman RS: **Intracellular signaling pathways pave roads to recovery for mood disorders.** *Ann Med* 2007, **39**:531-544.
56. Duric V, Banasr M, Licznarski P, Schmidt HD, Stockmeier CA, Simen AA, Newton SS, Duman RS: **A negative regulator of MAP kinase causes depressive behavior.** *Nat Med* 2010, **16**:1328-1332.
57. Eriksson TM, Delagrangre P, Spedding M, Popoli M, Mathe AA, Ogren SO, Svenningsson P: **Emotional memory impairments in a genetic rat model of depression: involvement of 5-HT/MEK/ Arc signaling in restoration.** *Mol Psychiatry* 2011. In press.
58. Lucas G, Rymar VV, Du J, Mnie-Filali O, Bisgaard C, Manta S, Lambas-Senas L, Wiborg O, Haddjeri N, Pineyro G *et al.*: **Serotonin(4) (5-HT₄) receptor agonists are putative antidepressants with a rapid onset of action.** *Neuron* 2007, **55**:712-725.
59. Svenningsson P, Tzavara ET, Qi H, Carruthers R, Witkin JM, Nomikos GG, Greengard P: **Biochemical and behavioral evidence for antidepressant-like effects of 5-HT₆ receptor stimulation.** *J Neurosci* 2007, **27**:4201-4209.
60. Vollenweider FX, Kometer M: **The neurobiology of psychedelic drugs: implications for the treatment of mood disorders.** *Nat Rev Neurosci* 2010, **11**:642-651.
An interesting account of recent behavioural and neuroimaging studies showing that psychedelic drugs modulate neural circuits implicated in depression and anxiety disorders, and may reduce the symptoms of these disorders.
61. Hayward G, Goodwin GM, Cowen PJ, Harmer CJ: **Low-dose tryptophan depletion in recovered depressed patients induces changes in cognitive processing without depressive symptoms.** *Biol Psychiatry* 2005, **57**:517-524.
62. Harmer CJ, Shelley NC, Cowen PJ, Goodwin GM: **Increased positive versus negative affective perception and memory in healthy volunteers following selective serotonin and norepinephrine reuptake inhibition.** *Am J Psychiatry* 2004, **161**:1256-1263.
63. Harmer CJ, Goodwin GM, Cowen PJ: **Why do antidepressants take so long to work? A cognitive neuropsychological model of antidepressant drug action.** *Br J Psychiatry* 2009, **195**:102-108.
An important account of how antidepressants might elevate mood by acting on negative emotional processing to bring about cognitive and psychological relearning.
64. Pringle A, Browning M, Cowen PJ, Harmer CJ: **A cognitive neuropsychological model of antidepressant drug action.** *Prog Neuropsychopharmacol Biol Psychiatry*, in press.
65. Browning M, Reid C, Cowen PJ, Goodwin GM, Harmer CJ: **A single dose of citalopram increases fear recognition in healthy subjects.** *J Psychopharmacol* 2007, **21**:684-690.
66. Maya Vetencourt JF, Sale A, Viegi A, Baroncelli L, De Pasquale R, O'Leary OF, Castren E, Maffei L: **The antidepressant fluoxetine restores plasticity in the adult visual cortex.** *Science* 2008, **320**:385-388.
An important study reporting that SSRI administration reinstates ocular dominance plasticity in adulthood via a BDNF-dependent mechanism.
67. Enkel T, Gholizadeh D, von Bohlen Und Halbach O, Sanchis-Segura C, Hurlmann R, Spanagel R, Gass P, Vollmayr B: **Ambiguous-cue interpretation is biased under stress- and depression-like states in rats.** *Neuropsychopharmacology* 2009, **35**:1008-1015.
An interesting model of negative cognitive bias in rodents.