

Homework 1 Answers, CS225 | ECE205A, UCSB, 2016 Winter

Copyright:

© Copyright 2016 Wim van Dam (UC Santa Barbara). It is forbidden to copy, distribute, or transmit this work, nor is it allowed to alter, transform, or build upon this work. For example, it is not allowed to upload this document to sites such as coursehero.com.

Question 1

Prove that for each discrete random variable $X \sim p(x)$ it holds that

$$H(X) \leq 2 \log \left(\sum_{x \in \mathcal{X}} \sqrt{p(x)} \right).$$

Answer: We have that

$$\begin{aligned} 2 \log \left(\sum_{x \in \mathcal{X}} \sqrt{p(x)} \right) &= 2 \log \left(\sum_{x \in \mathcal{X}} p(x) \cdot 1/\sqrt{p(x)} \right) \\ &= 2 \log E \left[1/\sqrt{p(X)} \right] \\ &\geq 2 E \left[\log (1/\sqrt{p(X)}) \right] \end{aligned}$$

(by Jensen's Inequality $f(E[Y]) \geq E[f(Y)]$ with the concave $f(Y) = 2 \log(Y)$ and $Y = 1/\sqrt{p(X)}$)

$$\begin{aligned} &= E \left[-\log(p(X)) \right] \\ &= H(X) \end{aligned}$$

Note that Jensen's inequality can be applied (in the direction sought) only when $f(Y)$ is concave, which is indeed the case for $f(Y) = 2 \log(Y)$.

Question 2

Consider two random variables X and Y with alphabets $\mathcal{X} = \{0, 1\}$ and $\mathcal{Y} = \{a, b, c\}$ and the following joint distribution:

$$\begin{array}{lll} \Pr[X = 0, Y = a] = 0.15, & \Pr[X = 0, Y = b] = 0.3, & \Pr[X = 0, Y = c] = 0.05, \\ \Pr[X = 1, Y = a] = 0.25, & \Pr[X = 1, Y = b] = 0.15, & \Pr[X = 1, Y = c] = 0.1. \end{array}$$

Determine the entropic quantities $H(X)$, $I(X; Y)$, et cetera, involved in the Venn diagram of the random variables X and Y .

Answer: The entropy of X is straightforward:

$$H(X) = -\Pr[X = 0] \log \Pr[X = 0] - \Pr[X = 1] \log \Pr[X = 1] = 1.$$

Next, we calculate explicitly the probabilities for $p(Y)$:

$$\begin{aligned} \Pr[Y = a] &= \Pr[Y = a | X = 0] \Pr[X = 0] + \Pr[Y = a | X = 1] \Pr[X = 1] = 0.4 \\ \Pr[Y = b] &= \Pr[Y = b | X = 0] \Pr[X = 0] + \Pr[Y = b | X = 1] \Pr[X = 1] = 0.45 \\ \Pr[Y = c] &= \Pr[Y = c | X = 0] \Pr[X = 0] + \Pr[Y = c | X = 1] \Pr[X = 1] = 0.15. \end{aligned}$$

With these probabilities $p(Y)$ we get the entropy of Y :

$$\begin{aligned} H(Y) &= -\Pr[Y = a] \log \Pr[Y = a] - \Pr[Y = b] \log \Pr[Y = b] - \Pr[Y = c] \log \Pr[Y = c] \\ &= 1.4577 \dots \end{aligned}$$

The conditional entropy $H(Y | X)$ is the last one that we have to calculate explicitly.

$$\begin{aligned} H(Y | X) &= \Pr[X = 0]H(Y | X = 0) + \Pr[X = 1]H(Y | X = 1) \\ &= \frac{1}{2} \sum_{y \in \{a, b, c\}} \Pr[Y = y | X = 0] \log \Pr[Y = y | X = 0] \\ &\quad + \frac{1}{2} \sum_{y \in \{a, b, c\}} \Pr[Y = y | X = 1] \log \Pr[Y = y | X = 1] \\ &= 1.3905 \dots \end{aligned}$$

The remaining entropies then follow in a straightforward manner.

$$\begin{aligned} H(X | Y) &= H(X) + H(Y | X) - H(Y) = 0.9328 \dots \\ H(X, Y) &= H(X) + H(Y | X) = 2.3905 \dots \\ I(X; Y) &= H(X) - H(X | Y) = 0.0672 \dots \end{aligned}$$

In summary:

$$\begin{aligned} H(X) &= 1 \\ H(X | Y) &\approx 0.9328 \\ H(Y) &\approx 1.4577 \\ H(Y | X) &\approx 1.3905 \\ H(X, Y) &\approx 2.3905 \\ I(X; Y) &\approx 0.0672. \end{aligned}$$

Question 3

Let p_0 and p_1 be probability distributions over a finite alphabet $\mathcal{X} = \{1, \dots, D\}$. Consider a 'mixture' of p_0 and p_1 described by $p_\lambda = (1 - \lambda) \cdot p_0 + \lambda \cdot p_1$ with $0 \leq \lambda \leq 1$, such that indeed for $\lambda = 0$ one gets p_0 , for $\lambda = 1$ we get p_1 , for $\lambda = 1/2$ we have the 50/50 mixture of the two probability distributions and so on. What can you prove about the entropy $H(p_\lambda)$ in terms of $H(p_0)$, $H(p_1)$, λ , and D ? Prove your statements.

Answer: Consider the entropy function H on the space of probability distributions $\{(q(1), \dots, q(D)) \in \mathbb{R}_{\geq 0}^D \mid q(1) + \dots + q(D) = 1\}$. The function H is concave and hence we have by Jensen's Inequality:

$$H((1 - \lambda)p_0 + \lambda p_1) \geq (1 - \lambda)H(p_0) + \lambda H(p_1).$$

Note that we used here the concavity of the function $\sum_j -p_j \log p_j$, not the concavity of $\log p$.

An alternative proof of this lower bound goes as follows. The events of the probability distribution p_λ can be thought of as having a hidden bit b that indicates if p_0 was used or p_1 . I.e. with probability $1 - \lambda$ we have $b = 0$ and we sample from p_0 , and with probability λ we have $b = 1$ and we sample from p_1 . If we know this hidden bit, we are dealing with the conditional entropy $H(p_\lambda | b)$. We thus have

$$\begin{aligned} H(p_\lambda) &\geq H(p_\lambda | b) \\ &= \Pr[b = 0]H(p_\lambda | b = 0) + \Pr[b = 1]H(p_\lambda | b = 1) \\ &= (1 - \lambda)H(p_0) + \lambda H(p_1). \end{aligned}$$

In addition this point of view also gives an upper bound through the joint entropy:

$$\begin{aligned} H(p_\lambda) &\leq H(p_\lambda, b) \\ &= H(p_\lambda | b) + H(b) \\ &= (1 - \lambda)H(p_0) + \lambda H(p_1) + H((1 - \lambda, \lambda)). \end{aligned}$$

Of course $H(p_\lambda) \leq \log D$ holds as well, but that is trivial.

Question 4

Prove or disprove the triangle inequality for the relative entropy function D . In other words, does it hold that for all probability distributions p, q, r over the alphabet \mathcal{X} we have $D(p \parallel q) + D(q \parallel r) \geq D(p \parallel r)$?

Answer: We can disprove the Triangle inequality by the following counterexample. Consider $x \in \mathcal{X} = \{0, 1\}$ with

	$x = 0$	$x = 1$
$p(x)$	0.3	0.7
$q(x)$	0.4	0.6
$r(x)$	0.5	0.5

Then, we have

$$D(p \parallel q) = p(0) \log \frac{p(0)}{q(0)} + p(1) \log \frac{p(1)}{q(1)} = 0.0302 \dots$$

$$D(q \parallel r) = q(0) \log \frac{q(0)}{r(0)} + q(1) \log \frac{q(1)}{r(1)} = 0.0290 \dots$$

$$D(p \parallel r) = p(0) \log \frac{p(0)}{r(0)} + p(1) \log \frac{p(1)}{r(1)} = 0.1187 \dots$$

As a result, $D(p \parallel q) + D(q \parallel r) = 0.0592 \dots < 0.1187 \dots = D(p \parallel r)$, hence $D(p \parallel q) + D(q \parallel r) \geq D(p \parallel r)$ does not always hold.