



ACNE MYCOSIS



Our certification



About us.

A Journey of Innovation and Excellence.

Founded in 2013 in Roseto degli Abruzzi, Italy, LCB Srl (Laboratori Chimici Braccili) began its journey as a specialized laboratory for the production of cosmetics and medical devices under private labels. Over the years, LCB has significantly expanded its capabilities, enriching its team with industry experts and investing heavily in research and development. This

strategic focus has enabled LCB to offer clients advanced, market-ready formulations and cutting-edge active ingredients for creating fully customized solutions.



In 2019, a pivotal milestone was achieved with the acquisition of a new facility, which facilitated the expansion of our production machinery and capabilities. This expansion was followed by the launch of a new department in 2021, dedicated to the production of dietary supplements, further diversifying our product range.

Throughout our journey, LCB has proudly collaborated with a multitude of brands, both nationally and internationally. Our commitment to innovation, quality, and customer satisfaction continues to drive us forward, as we strive to be at the forefront of the cosmetic and medical device industry.



BRADERM®






**THE COMPLETE SOLUTION
IN ACNE TREATMENT**

BRADERM[®]

ZOYLAK

30ml

Class IIa medical device for the treatment of mild and moderate facial acne.



BENZOIL PEROXIDE 4%
Antibacterial action

MANDELIC ACID
LACTOBIONIC ACID
RETINOL PALMITATE
Keratolytic action

ZINC SULFATE
Sebum regulating action

GLYCIRRETTIC ACID
PHYTOSPHINGOSINS
Anti-inflammatory action
and moisturizing action

DOSAGE: Apply to the affected area once a day (in the evening), after cleansing with **Zoylak Mousse**.

In case of sensitive skin, apply ZOYLAK every other day. White clothing is recommended as BPO may discolor colored clothing.

BRADERM[®]

AZN CREMA

30ml

Adjuvant cream in the treatment of mild and moderate acne.



SPF 30

AZELOGLYCIN 6%
NICOTINAMIDE 6%

Antibacterial action
and anti-inflammatory

BIOLIN-P[®]
Prebiotic

BURDOCK
Antibacterial action

DOSAGE: Apply AZN CREAM to the affected area 1 or 2 times a day and massage until completely absorbed

TO

In this early stage, the skin phenotype is characterized by numerous **papulopustular and nodular lesions**. Diffuse erythema and marked seborrhea are observed. The lesions show active inflammation and **potential CBA** infection, contributing to the heterogeneity of the skin tissue and the formation of atrophic and hypertrophic scars. Furthermore, **post-inflammatory melanonychia** is present, resulting from the hyperproduction of melanin in response to prolonged inflammatory damage.

T1

During this interval, there is a **quantitative decrease in efflorescence** and a **partial resolution of the erythema**. The density of **nodular lesions** is reduced, and a **decrease in the severity of seborrhea** is observed. These changes suggest therapeutic efficacy in modulating the activity of the sebaceous glands and limiting the local inflammatory response. Although residual skin discoloration persists, the skin begins to reveal areas of normalized texture and a decrease in peri-lesional inflammation.

T2

At the end of the treatment period, there is a **substantial resolution of the acne** with a **marked regression of the papulopustular lesions** and a restructuring of the skin surface. The residual hyperpigmentation shows signs of attenuation, with a normalization of the skin tone and an improvement in the dermal profile. The skin surface appears more homogeneous, with a **significant resolution of the erythema** and a reduction of hypertrophic scars. A normalization of sebaceous secretion and a re-epithelization of the areas previously affected by lesions can be deduced.

THERAPY: AZN CREAM (MORNING) + ZOYLAK (EVENING)
TREATMENT DURATION 3 MONTHS



At the initial stage, a clinical picture of active **inflammatory acne** is visible. The patient presents **multiple erythematopapular lesions** concentrated mainly on the cheek, with some pustules that suggest a bacterial superinfection. **Diffuse erythema** may also be noted, indicating an extensive local inflammatory response. These manifestations are typical of **moderate acne**, which can lead to alterations of the skin barrier and hyperreactivity of the pilosebaceous units.

After 3 months of therapy, an **appreciable reduction in inflammation** and the number of papulopustular lesions was noted. The erythema is reduced, and the skin texture appears more uniform. Although some post-inflammatory macules and superficial scars persist, the number of open and closed comedones seems to have decreased. This evolution demonstrates **a clear response to treatment**, with a normalization of sebum production and an improvement in skin homeostasis.

THERAPY: ZOYLAK APPLIED IN THE EVENING
TREATMENT DURATION 3 MONTHS:

TO

T1



BRADERM[®]

Dorsak

spray

100ml

Class IIa medical device specific for the treatment of back acne.



BENZOIL PEROSSIDO 6%
AZELOGLICINA
NICOTINAMIDE

Antibacterial and
anti-inflammatory action

RETINOL PALMITATE
LACTOBIONIC ACID

Keratolytic action

ZINC SULFATE
Sebum regulating action

DOSAGE: Apply once a day or, in case of sensitive skin, every other day, after washing the back area with a specific product and warm water. We recommend wearing white clothing as benzoyl peroxide may discolor colored clothing. Shake before using.



Inoak

Food supplement

60 capsules of 665mg

Specific supplement for the treatment of
young acne



- **MYO INOSITOL**
■ **D-CHIRO INOSITOL** **40:1**
 - Improved insulin sensitivity
 - Improved ovulation
 - Decreased LH and free testosterone
 - Significant reduction in HOMA index
- **NICOTINAMIDE**
Anti-inflammatory action
- **FOLIC ACID**
■ **BIOTIN**
Improves the appearance of the skin making it elastic and toned

DOSAGE: We recommend taking 2 tablets per day at the same time regardless of meals.

BRADERM[®]

ZOYLAK MOUSSE

100ml

Cleanser mousse for face with acne-prone skin



■ **AZELOGLYCIN
NICOTINAMIDE**
Anti-inflammatory action

■ **BIOLIN-P[®]**
Prebiotic

■ **ZINC SULPHATE**
Sebum-regulating action

■ **BURDOCK**
Antibacterial action

DOSAGE: Massage a small amount of product onto your already damp face, then rinse with warm water.

BRADERM[®]

ROSAC

30ml

Rich cream useful for the treatment of
Rosacea

**GREEN
PIGMENT**



**AZELAIC ACID 10%
NICOTINAMIDE**

Anti-inflammatory action

**HORSE CHESTNUT
RUSCO
TROXERUTIN**

They improve microcirculation

**SODIUM HYALURONATE
PHYTOSPHINGOSINS**

Moisturizing action

ISODECYL SALICYLATE

Barrier against external agents

DOSAGE: Apply the cream to the face 1-2 times a day and massage gently until completely absorbed.



THE COMPLETE SOLUTION IN THE TREATMENT OF MYCOSES





Shampoo DS

200ml

Treatment of **DRY AND OILY DANDRUFF**



- **CICLOPIROXOLAMINE**
- **THIOCONAZOLE**
- **CLIMBAZOLE**

Antifungal and anti-inflammatory action

- **SALICYLIC ACID**
- Keratolytic action

- **UNDECYLENE ACID**
- Fungistatic action

DOSAGE: Apply the product to damp hair, rub in, leave to act for a few minutes and then rinse thoroughly. To be used 2 or 3 times a week.



DS Ker Lozione

50ml

Treatment of **SEBORRHEIC DERMATITIS** of the scalp



- **ITRACONAZOLE**
- **TIOCONAZOLE**
Antifungal action
- **MANDELIC ACID**
Keratolytic action
- **SUPEROXIDE DISMUTASE**
- **EPIGALLOCATECHINS**
Antioxidant and anti-inflammatory action
- **ZINC SULPHATE**
Anti-seborrheic action

DOSAGE: Apply the product to the affected area once or twice a day and rub gently until completely absorbed.

BRADERM[®]

VERSIACTIVE

SPRAY⁺

100ml

Spray emulsion for the treatment of **PITYRIASIS
VERSICOLOR**, immediate results



- **ITRACONAZOLE**
- **TIOCONAZOLE**
- **CYCLOPIROXOLAMINE**
Antifungal action
- **SALICYLIC ACID**
Keratolytic action
- **GLYCYRRHETIC ACID**
Anti-inflammatory action

DOSAGE: Apply the product to the affected area once or twice a day and massage gently until completely absorbed.

BRADERM[®]

VERSIACTIVE

DETERGENTE

200ml

Body and scalp cleanser for the treatment and prevention of **PITIRYASIS VERSICOLOR**



- **CYCLOPIROXOLAMINE**
- **PIROCTONE OLAMINE**

Antifungal action

- **SALICYLIC ACID**

Keratolytic action

- **UNDECYLENE ACID**

Fungistatic action

- **TEA TREE OIL**

Antiseptic action

- **POLIDOCANOL**

Reduces dryness and itching

DOSAGE: Apply the product to damp skin and then rinse thoroughly.

Useful for those who frequent crowded environments such as swimming pools or gyms.



TREATMENT

VERSIACTIVE SPRAY E VERSIACTIVE DETERGENTE

TERAPIA : Treatment with **VERSIACTIVE SPRAY** applied morning and evening for 3 weeks and **VERSIACTIVE DETERGENTE** applied for 2 months

TO: The image shows the back of a patient affected by extensive and chronic Pityriasis Versicolor with evident white spots covering most of the surface.

TO





TREATMENT

VERSIACTIVE SPRAY E VERSIACTIVE DETERGENTE

T1: The patient is treated with Versiactive Detergent, at least once a day in association with Versiactive Spray.

After 20 days of treatments, a clear reduction in Pityriasis Versicolor is noted.

T1



BRADERM[®]

MICOBAT

CREMA

50ml

Emulsion for the treatment of **SEBORROIC DERMATITIS** of the face and trunk



■ **NICOTINAMIDE**
Anti-inflammatory action

■ **ITRACONAZOLE**
■ **CYCLOPIROXOLAMINE**
Antifungal action

■ **BISABOLOL**
Soothing action

■ **SHEA BUTTER**
Emollient action

DOSAGE: Apply small doses to the affected area and massage until completely absorbed, once or twice a day as needed

BRADERM[®]

MICOBAT POLVERE

75gr

Topical powder treatment effective in cases of
INTERTRIGO, TINEA PEDIS and **HYPERHIDROSIS**



- **SILICON DIOXIDE**
Absorbent action
- **TIOCONAZOLE**
■ **CILCOPIROXOLAMINE**
Antifungal action
- **CHLORHEXIDINE**
Antibacterial action
- **ZINC OXIDE**
Anti-inflammatory action
- **SHEA BUTTER**
Emollient and soothing action

DOSAGE: Apply the powder to the affected area, 1 or 2 times a day.

TREATMENT OF **STAPHYLOCOCCUS AUREUS** INFECTION



TO: An intertriginous lesion is shown, with pronounced erythema and possible edema. Dark pigmentation may indicate a subcutaneous hematoma or intense local inflammatory activity, typical of a skin infection with *Staphylococcus Aureus* (a pathogen that can cause severe symptoms due to its toxins and its ability to invade tissues)



T1: The image shows decreased inflammation and erythema, with slight residual discoloration suggesting the hematoma has begun to resolve. The reduction in color intensity and extension of the lesion indicate an effectiveness of the treatment in limiting inflammatory and microbial activity.



TREATMENT OF **STAPHYLOCOCCUS AURES** INFECTION



T2: In the last image, a further improvement of the lesion is noted, with a partial normalization of the skin color. The reduction of the affected area and the appearance of apparently healthy skin suggest that the treatment promoted healing.

THERAPY : topical treatment with **MICOBAT POLVERE** applied locally **2 times a day for 3 days.**

BRADERM[®]

Onycrom

15ml+15ml

Two-phase treatment, effective against
ONYCHOMYCOSIS

PHASE 1

UREA 40%
LACTIC ACID
Keratolytic action

PHASE 2

ITRACONAZOLE
CICLOPIROXOLAMINE
CLIMBAZOLE
Antifungal action

TEA TREE OIL
Anti-inflammatory and
antifungal action



DOSAGE: for the first month use phase 1 in the morning and phase 2 in the evening. In the following months, only use phase 2.

 **BRADERM**[®]

Distrofyd

15ml

Emulsion in tube with applicator brush for the treatment of **NAIL DYSTROPHIES**



■ **ARGAN OIL**
■ **JOJOBA OIL**

They promote nail regrowth

■ **PENTHENOL**
■ **NICOTINAMIDE**

They promote keratinization

■ **RETINOL**

Regenerating and smoothing action

■ **ORGANIC SILICON**
■ **METHIONINE**
■ **ZINC OXIDE**

Strengthening and protective action

DOSAGE: Apply a small amount of product with the appropriate brush on the nail, massaging it with the same, until completely absorbed. Repeat the application twice.

TREATMENT OF ONYCOMYCOSIS



TO: There is a clear state of onychomycosis, characterized by thickening of the nail, dyschromia and nail dystrophy. The surface of the nails appears irregular and brittle, typical of fungal infections that damage the keratin structure.

3 MONTHS



THERAPY: Application of ONYCROM PHASE 1 in the morning for the first month and ONYCROM PHASE 2 in the evening for 3 months.



T1: After the treatment you can notice how the appearance of the nails shows clear improvements. thickening is reduced, and the nail structure appears smoother and tidier. This is indicative of resolution of the onychomycosis and renewed growth of the healthy nail.

3 MONTHS



THERAPY: Application of ONYCROM PHASE 1 in the morning for the first month and ONYCROM PHASE 2 in the evening for 3 months.



TREATMENT OF NAIL DYSTROPHIES



TO: The nails show typical signs of nail dystrophy. Non-uniform thickening is noted, with streaks and areas of opacity, indicating an anomaly in the growth and structure of the nail keratin.

3 MONTHS



T1: After the treatment period, a normalization of the appearance of the nails is observed. Nails are smoother, with reduced thickening and increased transparency, suggesting an improvement in the quality and structure of nail keratin. Streaks are reduced and nails appear more uniform and healthy.

THERAPY: Application of **DISTROFYD**
2 times a day for 3 months.