

CIKAVES

CATEGORY

DERMOCOSMETIC

FUNCTIONAL INGREDIENTS

PANTHENOL 2,5%

It is also said: the skin vitamin or VIT. B5. In nature it is present in brewer's yeast, soy flour, egg yolk, peanuts, etc. In fact, its name derives from: PAN = EVERY and THEMA = PLACE.

It is a water-soluble vitamin so it is easily lost in cooking with water.

Its functions are: to promote epidermal regeneration and facilitate the healing of lesions..

A very important aspect in the scarring process is that panthenol is able to improve skin moisture as it penetrates the deeper layers of the skin and fixes the water there.

GLYCERIN 3%

It has a moisturizing and humectant action that allows the area where it is applied to be more elastic and soft.

TOCOPHERYL ACETATE 0,50%

It helps to increase the synthesis of collagen and glycoproteins, and hyaluronic acid.

Therefore it plays an important moisturizing action contributes to skin elasticity, it also plays an action against free radicals.

SUCRALFATE 0,005%

Cytoprotective agent commonly used as treatment of gastric and duodenal ulcers. Applied to the skin, it helps to improve and accelerate the tissue regeneration process, promoting the recovery of the barrier function.

COLLOIDAL SILVER 0,0005%

Antibacterial and disinfectant properties, it prevents the onset of infections when the skin is irritated, flaky and the barrier function is compromised by the presence of small lesions.

HOW TO USE

It should be used 2/3 times a day by applying a thin layer on the scar. Leave to dry for 4/5 minutes. In case of overdose, remove the excess.

Duration of treatment for at least two months is recommended, longer treatment is required for long-standing scars.

Warnings: do not apply to open or fresh wounds, near eyes and mucous membranes, skin treated with antibiotics or other topical products.

INDICATIONS

It improves the healing phase by promoting re-epithelialization.

Adjuvant in the treatment of Irritative Contact Dermatitis and Allergic Contact Dermatitis of the hands. Bedsores.

Cikaves is useful after minor dermatological surgery, laser, dermabrasion. Reduces the formation of keloids.