

PRODUCT	KERATO 20
CATEGORY	Dermocosmetic
INDICATIONS	The product is indicated for the treatment of skin thickening and for psoriatic forms of the scalp and body.
KERATOLYTIC ACTION	
• <u>UREA 20%</u> Exfoliating and keratolytic action. Present in the epidermis, it essentially comes from the sweat glands and is part of the hygroscopic compounds of the stratum corneum, which constitute the natural hydration factor (Natural Moisturizing Factor). Urea facilitates the elimination of scales, favors the renewal of corneocytes and constitutes a first choice moisturizing and humectant agent for the epidermis since it is able to break the hydrogen bonds of the protein chains and to alternate their configuration and aggregation. Consequently, at high concentrations (above 10%), it has a marked keratolytic effect and increases skin permeability. • <u>MANDELIC ACID 5%</u> Alpha-hydroxy acid extracted from bitter almonds. With a strong exfoliating and moisturizing power, unlike other acids it is not photosensitizing or irritating to the skin. It also has a lightening and depigmenting action.	
ANTI-INFLAMMATORY AND ANTIOXIDANT ACTION	
• <u>NICOTINAMIDE (VITAMIN PP) 2%</u> Water-soluble vitamin well known for its multiple applications, particularly in the dermatological field. It has a marked anti-inflammatory power and it is assumed that the mechanism of action is to be found in its antihistamine activity, scavenger of free radicals and inhibition of the enzyme 3-5 cyclic phosphodiesterase AMP. • <u>ACETYL HEXAPEPTIDE 49 (DELISENS®) 0,5%</u> The innovative peptide DELISENS (Acetyl Hexapeptide-49) improves the skin condition generated following the release from the PAR 2 receptor (Proteinase Activated Receptor 2) of pro-inflammatory mediators, also helping to restore the integrity of the damaged skin barrier. In sensitive skins it increases hydration and improves the smoothness and softness of skin thickening and keloids. The active ingredient has a marked soothing, antipruritic and repairing effect. • <u>FUMARIC ACID 2%</u> Recently also admitted as pharmaceutical active ingredients for oral use, compounds derived from fumaric acid have a certain effectiveness in combating the inflammatory process that develops in psoriatic skin, by blocking the production of proinflammatory cytokines.	
ANTIPRURIGINOUS ACTION	
• <u>ACETYL HEXAPEPTIDE 49 (DELISENS®)</u>	
EMOLLIENTS	
• SWEET ALMOND OIL 1.5% • SHEA BUTTER 1.5% • CERAMIDES and CHOLESTEROL 0.2%	

HOW TO USE	ACUTE FORMS: Apply twice a day on the affected part. CHRONIC FORMS: Apply once a day on the affected part. Warnings: do not apply on open or fresh wounds, close to the eyes and mucous membranes, skin treated with antibiotics or other products for topical use.
-------------------	---