

PRODUCT	BRADERM ONYCROM
PRIMARY PACKAGING	2 TUBES 15ML WITH BRUSH APPLICATOR
CATEGORY	COSMETIC
	This product has been formulated in two separated phases: PHASE 1: UREA and PHASE 2: ANTIMICOTICS as urea 40% together with antifungals, would reduces the effectiveness of the latter. Furthermore, a combined formulation of Urea and Antifungals in a single bottle would lead, given the long period of use of Urea, to an avulsion of the nail. Therefore Onycrom is unique thanks to the separation of the two phases (UREA and ANTIMICOTICI).
ACTIVE INGREDIENTS	PHASE 1: UREA 40% Keratolytic action. In the case of its application on nails affected by mycosis, it is useful to reduce the thickness, induced by the fungus, of the nail plate, facilitating the penetration of the functional active ingredients contained in ONYCROM gel (phase 2). PHASE 2: GEL CICLOPIROXOLAMINE (CPO): 0,3% It belongs to the hydroxypyridone family. The C.P.O. has a strong antifungal and antibacterial action, blocks the production of ATP (adenosine triphosphate), an energy reserve necessary for the vital functions of the fungal cell. Blocks carriers carrying leucine, valine and phenylalanine, with disruption of mRNA and protein synthesis. In addition, CPO binds and subtracts Fe³⁺, inactivating cellular enzymes, causing the death of fungi and bacteria. The CPO is recognized as one important anti-inflammatory action, as it inhibits the inflammatory mediators originating from arachidonic acid by the action of 5α-LIPOXYGENASE. ITRACONAZOLE: 0,10% Its mechanism of action occurs on the cytoplasmic membrane of the fungus, inhibiting the action of cytochrome P450 and consequently blocking the synthesis of ergosterol which represents the main steroid component of the cytoplasmic membrane, with

ACTIVE INGREDIENTS	consequent rupture of the same. It is a latest generation antifungal with broad spectrum activities. CLIMBAZOLE: 0,20% It also acts on the cytoplasmic membrane, damaging it with the inhibition of the enzyme 14-α-DEMETHYLASE responsible for the synthesis of ergosterol. It also carries out an anti-inflammatory activity of lipoxxygenase with consequent reduction of the mediators of inflammation and in synergy with itraconazole. TEA TREE OIL : 0,20% Extracted from an Australian plant, the MELALEUCA ALTERNIFOLIA. It has an antiseptic action and protects against alterations in the skin microbial flora. SALICYLIC ACID 0,5% PIROCTONE OLAMINE 0,5%
HOW TO USE	In the morning: PHASE 1 (UREA 40%) for 1 month In the evening: PHASE 2 (ANTIMICOTIC) for 2-3 month according to doctor suggestion