

Cryptography

Corso di Laurea Magistrale in Informatica

Theoretical Constructions of Pseudorandom Objects and Hash Functions

Ugo Dal Lago



ALMA MATER STUDIORUM
UNIVERSITÀ DI BOLOGNA



Academic Year 2023-2024

Theoretical Constructions of Pseudorandom Objects and Hash Functions

- ▶ As already mentioned, pseudorandom objects and hash functions can be constructed:
 - ▶ **Practically**, as we did in the previous chapter.
 - ▶ **Theoretically**, as we will do in this chapter, but only for pseudorandom objects.


Theoretical Constructions of Pseudorandom Objects and Hash Functions

- ▶ As already mentioned, pseudorandom objects and hash functions can be constructed:
 - ▶ **Practically**, as we did in the previous chapter.
 - ▶ **Theoretically**, as we will do in this chapter, but only for pseudorandom objects.
- ▶ In the theoretical approach, we will show that pseudorandom objects can be constructed **from other objects** whose existence, although not certain, is considered to be highly probable.
 - ▶ Cryptography, pseudorandomness and (partly) computational complexity deal with the **non-existence** of certain polytime algorithms with specific properties.
 - ▶ The work we did in the first part of the course, and which we will continue to do, consists in inferring this non-existence from gradually weaker hypothesis.


Theoretical Constructions of Pseudorandom Objects and Hash Functions

- ▶ As already mentioned, pseudorandom objects and hash functions can be constructed:
 - ▶ **Practically**, as we did in the previous chapter.
 - ▶ **Theoretically**, as we will do in this chapter, but only for pseudorandom objects.
- ▶ In the theoretical approach, we will show that pseudorandom objects can be constructed **from other objects** whose existence, although not certain, is considered to be highly probable.
 - ▶ Cryptography, pseudorandomness and (partly) computational complexity deal with the **non-existence** of certain polytime algorithms with specific properties.
 - ▶ The work we did in the first part of the course, and which we will continue to do, consists in inferring this non-existence from gradually weaker hypothesis.
- ▶ The interest in the constructions we will give is theoretical, but for purely efficiency reasons.

The Situation, in Brief



Where We Want to Go



One-Way Functions

- ▶ *Informally*, a one-way function is a function that is easy to compute, but which is hard to invert.

One-Way Functions

- ▶ *Informally*, a one-way function is a function that is easy to compute, but which is hard to invert.
- ▶ How to *formalise* this idea? We will proceed, in line with this course, by giving an experiment.

$\text{Invert}_{A,f}(n)$:

$x \leftarrow \{0, 1\}^n$;

$y \leftarrow f(x)$;

$z \leftarrow A(1^n, y)$;

Result: $(f(z) = y)$

One-Way Functions

- ▶ *Informally*, a one-way function is a function that is easy to compute, but which is hard to invert.
- ▶ How to *formalise* this idea? We will proceed, in line with this course, by giving an experiment.

Invert_{A,f}(n):
 $x \leftarrow \{0, 1\}^n$;
 $y \leftarrow f(x)$;
 $z \leftarrow A(1^n, y)$;
Result: ($f(z) = y$)

Definition

A function $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ is a **one-way function** iff there exists a polytime and deterministic algorithm which computes f and furthermore for every PPT A there exists a negligible ε such that

$$\Pr(\text{Invert}_{A,f}(n) = 1) \leq \varepsilon(n)$$

One-Way Functions

- ▶ **One-way permutations** are length-preserving one-way functions with the interesting property that $y \in \{0, 1\}^*$ uniquely determines x such that $f(x) = y$.

One-Way Functions

- ▶ **One-way permutations** are length-preserving one-way functions with the interesting property that $y \in \{0, 1\}^*$ uniquely determines x such that $f(x) = y$.
- ▶ *Examples of (Assumed) One-Way Functions:*
 - ▶ **Multiplication Between Natural Numbers**
 - ▶ Consider f_{MULT} defined by $f_{MULT}(x, y) = x \cdot y$: given two strings that we interpret as natural numbers, we return their product.
 - ▶ If we do not put any other constraints on x and y , the function f_{MULT} is easily invertible.
 - ▶ We will study this function in the next chapter.
 - ▶ **Subset-Sum Problem**
 - ▶ Consider instead the function f_{SS} defined by $f_{SS}(x_1, \dots, x_n, J) = (x_1, \dots, x_n, \sum_{j \in J} x_j)$, where $|x_j| = n$ and J is interpreted as a subset of $\{1, \dots, n\}$.
 - ▶ The inverse of f_{SS} corresponds to the so-called subset-sum problem.

Hard-Core Predicates

- ▶ A one-way function f is such that $f(x)$ does not *entirely* reveal x
 - ▶ This does not imply that the same applies to **parts of** x , for example a single bit.
 - ▶ Consider for example a one-way function f , and construct $g(x, y) = (x, f(y))$. g is also a one-way function (if we could invert g , we can also invert f). But $g(x, y)$ reveals an important part of its input, namely x .

Definition

A predicate $hc : \{0, 1\}^* \rightarrow \{0, 1\}$ is called *hard-core predicate* of a function f if and only if hc is polynomial time computable and for every adversary PPT A it holds that

$$\Pr(A(f(x)) = hc(x)) \leq \frac{1}{2} + \varepsilon(n)$$

where ε is negligible.

Hard-Core Predicates

- ▶ It may seem at first sight that hc defined by $hc(x_1 \cdots x_n) = \bigoplus_{i=1}^n x_i$ is a hard-core predicate *for each* function f .
 - ▶ Given g one-way, the function f defined by $f(x) = (g(x), hc(x))$ is also a one-way function, but certainly hc is not hard-core for f , because its value is easily retrievable from the output.
- ▶ For some (non-one-way) functions, it is possible to construct trivial hard-core predicates. For example the function $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ defined by $f(\epsilon) = \epsilon$ and $f(b \cdot s) = s$ for every $b \in \{0, 1\}$ and $s \in \{0, 1\}^*$.
 - ▶ The result $f(s)$ does not depend on the first bit of s , which can then become a hard-core predicate.

The Goldreich-Levin Theorem

Theorem

If there is a one-way function (respectively, a one-way permutation) f , then there exists a one-way function (respectively a one-way permutation) g and a hard-core predicate hc for g .

The Goldreich-Levin Theorem

Theorem

If there is a one-way function (respectively, a one-way permutation) f , then there exists a one-way function (respectively a one-way permutation) g and a hard-core predicate hc for g .

- ▶ This is one of the most important results in the theory of one-way functions, with crucial implications in cryptography.
- ▶ The function g is constructed from f by setting $g(x, r) = (f(x), r)$, while hc is defined by $hc(x, r) = \bigoplus_{i=1}^n x_i \cdot r_i$.

From One-Way Permutations to Pseudorandom Generators

Theorem

Let f be a one-way permutation and let hc be a hard-core predicate for f . Then G defined by $G(s) = (f(s), hc(s))$ is a pseudorandom generator with expansion factor $\ell(n) = n + 1$.


From One-Way Permutations to Pseudorandom Generators

Theorem

Let f be a one-way permutation and let hc be a hard-core predicate for f . Then G defined by $G(s) = (f(s), hc(s))$ is a pseudorandom generator with expansion factor $\ell(n) = n + 1$.

- ▶ This is another crucial result, linking the theory of one-way functions to pseudorandomness.
- ▶ Intuitively, the first $|s|$ bits of G 's output are pseudorandom due to the properties of f , while the last bit is pseudorandom due to the properties of hc .

One-Way Permutations and Pseudorandom Generators



Arbitrary Expansion Factor


Theorem

If there exists a pseudorandom generator G with expansion factor $\ell(n) = n + 1$, then there exists another other pseudorandom generator H , with an arbitrary expansion factor, as long as it is polynomial.

Arbitrary Expansion Factor

Theorem

If there exists a pseudorandom generator G with expansion factor $\ell(n) = n + 1$, then there exists another other pseudorandom generator H , with an arbitrary expansion factor, as long as it is polynomial.



From Generators to Functions


Theorem

If there exists a pseudorandom generator G with expansion factor $\ell(n) = 2n$, then there exists a pseudorandom function.

From Generators to Functions

Theorem

If there exists a pseudorandom generator G with expansion factor $\ell(n) = 2n$, then there exists a pseudorandom function.



Coming Full Circle

Theorem

If a pseudorandom function exists, then there exists a strong pseudorandom permutation.

Coming Full Circle


Theorem

If a pseudorandom function exists, then there exists a strong pseudorandom permutation.

Theorem

If there is a pseudorandom generator, then there is a one-way function.

One-Way Functions and Pseudorandomness



The Overall Picture

