

Physics Equations Reference

Contents

I	Fundamental Equations	1
1	Fundamental Equations	3
II	Derived Equations	11
2	Airy Disk	13
2.1	Introduction	13
2.2	Fundamental Equations Used	13
2.3	Assumptions	14
2.4	Mathematical Derivation	14
2.5	Characteristics of the equations	15
2.6	Solution Methods	15
2.7	Applications	16
2.8	Verification	16
2.9	FAQ	17
3	Bernoulli Equation	19
3.1	Introduction	19
3.2	Fundamental Equations Used	19
3.3	Assumptions	19
3.4	Mathematical Derivation	20
3.5	Characteristics of the equations	21
3.6	Solution Methods	21
3.7	Applications	21

3.8	Verification	22
3.9	FAQ	22
4	Boltzmann Transport Equation	25
4.1	Introduction	25
4.2	Fundamental Equations Used	25
4.3	Assumptions	26
4.4	Mathematical Derivation	26
4.5	Characteristics of the equations	27
4.6	Solution Methods	27
4.7	Applications	28
4.8	Verification	28
4.9	FAQ	28
5	Bose-Einstein Distribution	31
5.1	Introduction	31
5.2	Fundamental Equations Used	31
5.3	Assumptions	32
5.4	Mathematical Derivation	32
5.5	Characteristics of the equations	33
5.6	Solution Methods	33
5.7	Applications	33
5.8	Verification	34
5.9	FAQ	34
6	Clausius-Clapeyron Equation	37
6.1	Introduction	37
6.2	Fundamental Equations Used	37
6.3	Assumptions	38
6.4	Mathematical Derivation	38
6.5	Characteristics of the equations	39
6.6	Solution Methods	39
6.7	Applications	39
6.8	Verification	40

6.9	FAQ	40
7	Continuity Equation	43
7.1	Introduction	43
7.2	Fundamental Equations Used	43
7.3	Assumptions	44
7.4	Mathematical Derivation	44
7.5	Characteristics of the equations	45
7.6	Solution Methods	45
7.7	Applications	45
7.8	Verification	46
7.9	FAQ	46
8	Diffusion Equation	49
8.1	Introduction	49
8.2	Fundamental Equations Used	49
8.3	Assumptions	50
8.4	Mathematical Derivation	50
8.5	Characteristics of the equations	51
8.6	Solution Methods	51
8.7	Applications	52
8.8	Verification	52
8.9	FAQ	53
9	Dirac Equation	55
9.1	Introduction	55
9.2	Fundamental Equations Used	55
9.3	Assumptions	56
9.4	Mathematical Derivation	56
9.5	Characteristics of the equations	57
9.6	Solution Methods	57
9.7	Applications	58
9.8	Verification	58
9.9	FAQ	59

10 Electromagnetic Wave Equation	61
10.1 Introduction	61
10.2 Fundamental Equations Used	61
10.3 Assumptions	62
10.4 Mathematical Derivation	62
10.5 Characteristics of the equations	63
10.6 Solution Methods	64
10.7 Applications	64
10.8 Verification	65
10.9 FAQ	65
11 Entropy	67
11.1 Introduction	67
11.2 Fundamental Equations Used	67
11.3 Assumptions	68
11.4 Mathematical Derivation	68
11.5 Characteristics of the equations	70
11.6 Solution Methods	70
11.7 Applications	70
11.8 Verification	71
11.9 FAQ	71
12 Euler Equation (Fluid Dynamics)	75
12.1 Introduction	75
12.2 Fundamental Equations Used	75
12.3 Assumptions	76
12.4 Mathematical Derivation	76
12.5 Characteristics of the equations	77
12.6 Solution Methods	77
12.7 Applications	78
12.8 Verification	78
12.9 FAQ	79

13 Euler-Bernoulli Beam Equation	81
13.1 Introduction	81
13.2 Fundamental Equations Used	81
13.3 Assumptions	82
13.4 Mathematical Derivation	82
13.5 Characteristics of the equations	83
13.6 Solution Methods	83
13.7 Applications	83
13.8 Verification	84
13.9 FAQ	84
14 Euler–Lagrange Equation	87
14.1 Introduction	87
14.2 Fundamental Equations Used	87
14.3 Assumptions	88
14.4 Mathematical Derivation	88
14.5 Characteristics of the equations	89
14.6 Solution Methods	89
14.7 Applications	89
14.8 Verification	90
14.9 FAQ	90
15 Fermi–Dirac Distribution	93
15.1 Introduction	93
15.2 Fundamental Equations Used	93
15.3 Assumptions	94
15.4 Mathematical Derivation	94
15.5 Characteristics of the equations	95
15.6 Solution Methods	95
15.7 Applications	95
15.8 Verification	96
15.9 FAQ	96

16 Fraunhofer Diffraction	99
16.1 Introduction	99
16.2 Fundamental Equations Used	99
16.3 Assumptions	100
16.4 Mathematical Derivation	100
16.5 Characteristics of the equations	101
16.6 Solution Methods	101
16.7 Applications	101
16.8 Verification	102
16.9 FAQ	102
17 Fresnel Equations	105
17.1 Introduction	105
17.2 Fundamental Equations Used	105
17.3 Assumptions	105
17.4 Mathematical Derivation	106
17.5 Characteristics of the equations	107
17.6 Solution Methods	107
17.7 Applications	107
17.8 Verification	108
17.9 FAQ	108
18 Friedmann Equation	111
18.1 Introduction	111
18.2 Fundamental Equations Used	111
18.3 Assumptions	112
18.4 Mathematical Derivation	112
18.5 Characteristics of the equations	113
18.6 Solution Methods	113
18.7 Applications	114
18.8 Verification	114
18.9 FAQ	114

19 Geodesic Equation	117
19.1 Introduction	117
19.2 Fundamental Equations Used	117
19.3 Assumptions	118
19.4 Mathematical Derivation	118
19.5 Characteristics of the equations	119
19.6 Solution Methods	119
19.7 Applications	119
19.8 Verification	120
19.9 FAQ	120
20 Hamilton’s Equations	123
20.1 Introduction	123
20.2 Fundamental Equations Used	123
20.3 Assumptions	124
20.4 Mathematical Derivation	124
20.5 Characteristics of the equations	125
20.6 Solution Methods	125
20.7 Applications	126
20.8 Verification	126
20.9 FAQ	127
21 Heat Equation	129
21.1 Introduction	129
21.2 Fundamental Equations Used	129
21.3 Assumptions	130
21.4 Mathematical Derivation	130
21.5 Characteristics of the equations	131
21.6 Solution Methods	131
21.7 Applications	132
21.8 Verification	132
21.9 FAQ	133

22 Heisenberg Equation	135
22.1 Introduction	135
22.2 Fundamental Equations Used	135
22.3 Assumptions	136
22.4 Mathematical Derivation	136
22.5 Characteristics of the equations	137
22.6 Solution Methods	137
22.7 Applications	138
22.8 Verification	138
22.9 FAQ	139
23 Ideal Gas Equation	141
23.1 Introduction	141
23.2 Fundamental Equations Used	141
23.3 Assumptions	142
23.4 Mathematical Derivation	142
23.5 Characteristics of the equations	143
23.6 Solution Methods	143
23.7 Applications	143
23.8 Verification	143
23.9 FAQ	144
24 Kinematic Equations	147
24.1 Introduction	147
24.2 Fundamental Equations Used	148
24.3 Assumptions	148
24.4 Mathematical Derivation	148
24.5 Characteristics of the equations	149
24.6 Solution Methods	150
24.7 Applications	150
24.8 Verification	150
24.9 FAQ	151

25 Klein-Gordon Equation	153
25.1 Introduction	153
25.2 Fundamental Equations Used	153
25.3 Assumptions	154
25.4 Mathematical Derivation	154
25.5 Characteristics of the equations	155
25.6 Solution Methods	155
25.7 Applications	155
25.8 Verification	156
25.9 FAQ	156
26 Length Contraction	159
26.1 Introduction	159
26.2 Fundamental Equations Used	159
26.3 Assumptions	160
26.4 Mathematical Derivation	160
26.5 Characteristics of the equations	161
26.6 Solution Methods	161
26.7 Applications	161
26.8 Verification	161
26.9 FAQ	162
27 Liénard-Wiechert Potentials	165
27.1 Introduction	165
27.2 Fundamental Equations Used	165
27.3 Assumptions	166
27.4 Mathematical Derivation	166
27.5 Characteristics of the equations	167
27.6 Solution Methods	167
27.7 Applications	168
27.8 Verification	168
27.9 FAQ	169

28 Mass-Energy Equivalence	171
28.1 Introduction	171
28.2 Fundamental Equations Used	171
28.3 Assumptions	172
28.4 Mathematical Derivation	172
28.5 Characteristics of the equations	173
28.6 Solution Methods	173
28.7 Applications	174
28.8 Verification	174
28.9 FAQ	174
29 Maxwell Relation	177
29.1 Introduction	177
29.2 Fundamental Equations Used	177
29.3 Assumptions	178
29.4 Mathematical Derivation	178
29.5 Characteristics of the equations	179
29.6 Solution Methods	179
29.7 Applications	179
29.8 Verification	179
29.9 FAQ	180
30 Maxwell–Boltzmann Distribution	183
30.1 Introduction	183
30.2 Fundamental Equations Used	183
30.3 Assumptions	184
30.4 Mathematical Derivation	184
30.5 Characteristics of the equations	185
30.6 Solution Methods	185
30.7 Applications	185
30.8 Verification	186
30.9 FAQ	186

31 Mirror Equation	189
31.1 Introduction	189
31.2 Fundamental Equations Used	189
31.3 Assumptions	189
31.4 Mathematical Derivation	190
31.5 Characteristics of the equations	191
31.6 Solution Methods	191
31.7 Applications	191
31.8 Verification	192
31.9 FAQ	192
32 Navier–Stokes Equation	195
32.1 Introduction	195
32.2 Fundamental Equations Used	195
32.3 Assumptions	196
32.4 Mathematical Derivation	196
32.5 Characteristics of the equations	197
32.6 Solution Methods	197
32.7 Applications	197
32.8 Verification	198
32.9 FAQ	198
33 Partition Function	201
33.1 Introduction	201
33.2 Fundamental Equations Used	201
33.3 Assumptions	202
33.4 Mathematical Derivation	202
33.5 Characteristics of the equations	203
33.6 Solution Methods	203
33.7 Applications	203
33.8 Verification	203
33.9 FAQ	204

34 Pauli Equation	207
34.1 Introduction	207
34.2 Fundamental Equations Used	207
34.3 Assumptions	207
34.4 Mathematical Derivation	208
34.5 Characteristics of the equations	209
34.6 Solution Methods	209
34.7 Applications	209
34.8 Verification	210
34.9 FAQ	210
35 Planck Radiation Law	213
35.1 Introduction	213
35.2 Fundamental Equations Used	213
35.3 Assumptions	214
35.4 Mathematical Derivation	214
35.5 Characteristics of the equations	215
35.6 Solution Methods	215
35.7 Applications	215
35.8 Verification	216
35.9 FAQ	216
36 Plate Equation	219
36.1 Introduction	219
36.2 Fundamental Equations Used	219
36.3 Assumptions	220
36.4 Mathematical Derivation	220
36.5 Characteristics of the equations	221
36.6 Solution Methods	221
36.7 Applications	221
36.8 Verification	222
36.9 FAQ	222

37 Laplace’s Equation	225
37.1 Introduction	225
37.2 Fundamental Equations Used	225
37.3 Assumptions	226
37.4 Mathematical Derivation	226
37.5 Characteristics of the equations	227
37.6 Solution Methods	227
37.7 Applications	227
37.8 Verification	227
37.9 FAQ	228
 38 Poynting Theorem	 231
38.1 Introduction	231
38.2 Fundamental Equations Used	231
38.3 Assumptions	231
38.4 Mathematical Derivation	232
38.5 Characteristics of the equations	233
38.6 Solution Methods	233
38.7 Applications	233
38.8 Verification	233
38.9 FAQ	234
 39 Probability Current	 237
39.1 Introduction	237
39.2 Fundamental Equations Used	237
39.3 Assumptions	237
39.4 Mathematical Derivation	238
39.5 Characteristics of the equations	239
39.6 Solution Methods	239
39.7 Applications	239
39.8 Verification	240
39.9 FAQ	240

40 Quantum Continuity Equation	243
40.1 Introduction	243
40.2 Fundamental Equations Used	243
40.3 Assumptions	243
40.4 Mathematical Derivation	244
40.5 Characteristics of the equations	244
40.6 Solution Methods	244
40.7 Applications	245
40.8 Verification	245
40.9 FAQ	246
41 Relativistic Energy	249
41.1 Introduction	249
41.2 Fundamental Equations Used	249
41.3 Assumptions	249
41.4 Mathematical Derivation	250
41.5 Characteristics of the equations	250
41.6 Solution Methods	250
41.7 Applications	251
41.8 Verification	251
41.9 FAQ	252
42 Relativistic Momentum	253
42.1 Introduction	253
42.2 Fundamental Equations Used	253
42.3 Assumptions	253
42.4 Mathematical Derivation	254
42.5 Characteristics of the equations	254
42.6 Solution Methods	254
42.7 Applications	254
42.8 Verification	255
42.9 FAQ	255

43 Reynolds Transport Theorem	257
43.1 Introduction	257
43.2 Fundamental Equations Used	257
43.3 Assumptions	258
43.4 Mathematical Derivation	258
43.5 Characteristics of the equations	259
43.6 Solution Methods	259
43.7 Applications	259
43.8 Verification	260
43.9 FAQ	260
 44 Rotational Kinetic Energy	 263
44.1 Introduction	263
44.2 Fundamental Equations Used	263
44.3 Assumptions	264
44.4 Mathematical Derivation	264
44.5 Characteristics of the equations	265
44.6 Solution Methods	265
44.7 Applications	265
44.8 Verification	265
44.9 FAQ	266
 45 Sackur–Tetrode Equation	 269
45.1 Introduction	269
45.2 Fundamental Equations Used	269
45.3 Assumptions	270
45.4 Mathematical Derivation	270
45.5 Characteristics of the equations	271
45.6 Solution Methods	271
45.7 Applications	271
45.8 Verification	272
45.9 FAQ	272

46 Schwarzschild Metric	275
46.1 Introduction	275
46.2 Fundamental Equations Used	275
46.3 Assumptions	276
46.4 Mathematical Derivation	276
46.5 Characteristics of the equations	277
46.6 Solution Methods	277
46.7 Applications	277
46.8 Verification	278
46.9 FAQ	278
 47 Single-Slit Diffraction	 281
47.1 Introduction	281
47.2 Fundamental Equations Used	281
47.3 Assumptions	282
47.4 Mathematical Derivation	282
47.5 Characteristics of the equations	283
47.6 Solution Methods	283
47.7 Applications	283
47.8 Verification	284
47.9 FAQ	284
 48 Skin Depth	 287
48.1 Introduction	287
48.2 Fundamental Equations Used	287
48.3 Assumptions	287
48.4 Mathematical Derivation	288
48.5 Characteristics of the equations	289
48.6 Solution Methods	289
48.7 Applications	289
48.8 Verification	289
48.9 FAQ	290

49 Stefan–Boltzmann Law	293
49.1 Introduction	293
49.2 Fundamental Equations Used	293
49.3 Assumptions	294
49.4 Mathematical Derivation	294
49.5 Characteristics of the equations	295
49.6 Solution Methods	295
49.7 Applications	295
49.8 Verification	296
49.9 FAQ	296
 50 Stress Equilibrium Equations	 299
50.1 Introduction	299
50.2 Fundamental Equations Used	299
50.3 Assumptions	300
50.4 Mathematical Derivation	300
50.5 Characteristics of the equations	301
50.6 Solution Methods	301
50.7 Applications	301
50.8 Verification	301
50.9 FAQ	302
 51 Stress–Strain Relation	 305
51.1 Introduction	305
51.2 Fundamental Equations Used	305
51.3 Assumptions	305
51.4 Mathematical Derivation	306
51.5 Characteristics of the equations	306
51.6 Solution Methods	307
51.7 Applications	307
51.8 Verification	307
51.9 FAQ	308

52 Telegraph Equation	311
52.1 Introduction	311
52.2 Fundamental Equations Used	311
52.3 Assumptions	312
52.4 Mathematical Derivation	312
52.5 Characteristics of the equations	313
52.6 Solution Methods	313
52.7 Applications	313
52.8 Verification	313
52.9 FAQ	314
 53 Thin Lens Equation	 317
53.1 Introduction	317
53.2 Fundamental Equations Used	317
53.3 Assumptions	317
53.4 Mathematical Derivation	318
53.5 Characteristics of the equations	319
53.6 Solution Methods	319
53.7 Applications	319
53.8 Verification	319
53.9 FAQ	320
 54 Time Dilation	 323
54.1 Introduction	323
54.2 Fundamental Equations Used	323
54.3 Assumptions	323
54.4 Mathematical Derivation	324
54.5 Characteristics of the equations	324
54.6 Solution Methods	324
54.7 Applications	325
54.8 Verification	325
54.9 FAQ	326

55 Time-Independent Schrödinger Equation	329
55.1 Introduction	329
55.2 Fundamental Equations Used	329
55.3 Assumptions	329
55.4 Mathematical Derivation	330
55.5 Characteristics of the equations	331
55.6 Solution Methods	331
55.7 Applications	331
55.8 Verification	331
55.9 FAQ	332
 56 Torque	 335
56.1 Introduction	335
56.2 Fundamental Equations Used	335
56.3 Assumptions	335
56.4 Mathematical Derivation	336
56.5 Characteristics of the equations	337
56.6 Solution Methods	337
56.7 Applications	337
56.8 Verification	337
56.9 FAQ	338
 57 Vorticity Equation	 341
57.1 Introduction	341
57.2 Fundamental Equations Used	341
57.3 Assumptions	341
57.4 Mathematical Derivation	342
57.5 Characteristics of the equations	343
57.6 Solution Methods	343
57.7 Applications	343
57.8 Verification	343
57.9 FAQ	344

58 Wave Equation	347
58.1 Introduction	347
58.2 Fundamental Equations Used	347
58.3 Assumptions	347
58.4 Mathematical Derivation	348
58.5 Characteristics of the equations	349
58.6 Solution Methods	349
58.7 Applications	349
58.8 Verification	349
58.9 FAQ	350
59 Wave Equation in Solids	353
59.1 Introduction	353
59.2 Fundamental Equations Used	353
59.3 Assumptions	353
59.4 Mathematical Derivation	354
59.5 Characteristics of the equations	355
59.6 Solution Methods	355
59.7 Applications	355
59.8 Verification	356
59.9 FAQ	356
60 Wien’s Displacement Equation	359
60.1 Introduction	359
60.2 Fundamental Equations Used	359
60.3 Assumptions	359
60.4 Mathematical Derivation	360
60.5 Characteristics of the equations	361
60.6 Solution Methods	361
60.7 Applications	361
60.8 Verification	362
60.9 FAQ	362

61 Work–Energy Equation	365
61.1 Introduction	365
61.2 Fundamental Equations Used	365
61.3 Assumptions	365
61.4 Mathematical Derivation	366
61.5 Characteristics of the equations	367
61.6 Solution Methods	367
61.7 Applications	367
61.8 Verification	367
61.9 FAQ	368
 62 Lorentz Force Law	 371
62.1 Introduction	371
62.2 Fundamental Equations Used	371
62.3 Assumptions	371
62.4 Mathematical Derivation	372
62.5 Characteristics of the equations	372
62.6 Solution Methods	372
62.7 Applications	373
62.8 Verification	373
62.9 FAQ	373
 63 Biot–Savart Law	 375
63.1 Introduction	375
63.2 Fundamental Equations Used	375
63.3 Assumptions	375
63.4 Mathematical Derivation	376
63.5 Characteristics of the equations	376
63.6 Solution Methods	377
63.7 Applications	377
63.8 Verification	377
63.9 FAQ	378

64 Heisenberg Uncertainty Principle	381
64.1 Introduction	381
64.2 Fundamental Equations Used	381
64.3 Assumptions	381
64.4 Mathematical Derivation	382
64.5 Characteristics of the equations	382
64.6 Solution Methods	383
64.7 Applications	383
64.8 Verification	383
64.9 FAQ	384
 65 De Broglie Wavelength	 387
65.1 Introduction	387
65.2 Fundamental Equations Used	387
65.3 Assumptions	387
65.4 Mathematical Derivation	388
65.5 Characteristics of the equations	388
65.6 Solution Methods	388
65.7 Applications	389
65.8 Verification	389
65.9 FAQ	389
 66 Snell’s Law of Refraction	 391
66.1 Introduction	391
66.2 Fundamental Equations Used	391
66.3 Assumptions	391
66.4 Mathematical Derivation	392
66.5 Characteristics of the equations	392
66.6 Solution Methods	392
66.7 Applications	393
66.8 Verification	393
66.9 FAQ	393

67 Young’s Double-Slit Interference	395
67.1 Introduction	395
67.2 Fundamental Equations Used	395
67.3 Assumptions	395
67.4 Mathematical Derivation	396
67.5 Characteristics of the equations	396
67.6 Solution Methods	396
67.7 Applications	397
67.8 Verification	397
67.9 FAQ	398
 68 Kepler’s Third Law of Planetary Motion	 399
68.1 Introduction	399
68.2 Fundamental Equations Used	399
68.3 Assumptions	399
68.4 Mathematical Derivation	400
68.5 Characteristics of the equations	400
68.6 Solution Methods	400
68.7 Applications	401
68.8 Verification	401
68.9 FAQ	401
 69 Damped and Driven Harmonic Oscillator	 403
69.1 Introduction	403
69.2 Fundamental Equations Used	403
69.3 Assumptions	403
69.4 Mathematical Derivation	404
69.5 Characteristics of the equations	404
69.6 Solution Methods	405
69.7 Applications	405
69.8 Verification	405
69.9 FAQ	406

70 Poisson’s Equation	409
70.1 Introduction	409
70.2 Fundamental Equations Used	409
70.3 Assumptions	409
70.4 Mathematical Derivation	410
70.5 Characteristics of the equations	410
70.6 Solution Methods	410
70.7 Applications	411
70.8 Verification	411
70.9 FAQ	412
71 Historical Notes	413

Part I

Fundamental Equations

Fundamental Equations

This chapter mixes equations from different levels of physics. Start with the conservation laws and Newton's laws, then return to Maxwell's equations, relativity, and quantum mechanics after the symbols feel familiar. For each item, read the sentence before the formula first; the formula is the compact version of that sentence.

Reality Mapping: These equations are the grammar of physics—they tell us what can and cannot happen in the universe. Conservation laws say that certain quantities (energy, momentum, angular momentum) are the same before and after any process. Maxwell's equations say that electric and magnetic fields are two aspects of one thing, and that light is an electromagnetic wave. Einstein's equations say that mass and energy are interchangeable, and that gravity is the curvature of spacetime. Quantum equations say that particles behave like waves, that you cannot know everything at once, and that the vacuum is seething with activity. Together, these equations describe the rules that govern everything from falling apples to exploding stars.

1. **Ampere–Maxwell Law:** Electric currents and changing electric fields both create magnetic fields. This equation also predicts that light is an electromagnetic wave.

$$\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t}$$

- **Assumptions:** The medium is linear and time-invariant. The fields vary slowly enough that radiation reaction is negligible.
 - **Limitations:** Fails at quantum scales (need quantum electrodynamics). Does not account for magnetic monopoles (none have been observed). In media with large dispersion or nonlinear effects, additional terms are needed.
 - **Constraints:** Valid in the classical regime where $v \ll c$ for material motion. Requires μ_0 and ϵ_0 to be constants (true in vacuum).
2. **Conservation of Angular Momentum:** The total angular momentum of an isolated system does not change. This is why ice skaters spin faster when they pull their arms in.

$$\mathbf{L} = \text{const}$$

- **Assumptions:** The system is isolated—no external torques act on it. The Lagrangian of the system is invariant under rotations.
- **Limitations:** Does not apply in non-inertial (rotating) frames without adding fictitious torques. In general relativity, angular momentum is conserved only locally.
- **Constraints:** Valid for closed systems only. In open systems, external torques must be included.

3. **Conservation of Energy:** Energy can change form but the total amount stays the same.

$$\Delta E_{\text{total}} = 0$$

- **Assumptions:** The system is closed (no energy enters or leaves). Time-translation symmetry holds (the laws of physics do not change with time).
- **Limitations:** In an expanding universe, energy conservation becomes subtle (photons lose energy to cosmological redshift). In general relativity, global energy conservation is not well-defined in all cases.
- **Constraints:** Applies strictly to isolated systems. In open systems, energy input and output must be accounted for.

4. **Conservation of Mass:** Mass is neither created nor destroyed.

$$\Delta m = 0$$

- **Assumptions:** Velocities are much less than the speed of light ($v \ll c$). No nuclear reactions or particle–antiparticle annihilation occur.
- **Limitations:** Fails at relativistic speeds—mass can be converted to energy and vice versa via $E = mc^2$. In nuclear and particle physics, mass is not separately conserved.
- **Constraints:** Valid only in classical, non-relativistic mechanics. In chemistry and everyday life, it holds to extraordinary precision.

5. **Conservation of Momentum:** The total momentum of an isolated system does not change.

$$\sum \mathbf{p}_i = \text{const}$$

- **Assumptions:** The system is isolated—no external forces act on it. The Lagrangian is invariant under spatial translations.
- **Limitations:** In an expanding universe, momentum of photons is not conserved globally. In general relativity, momentum conservation is local, not global.
- **Constraints:** Valid for closed systems. In open systems, external impulses must be included.

6. **Einstein Field Equations:** The core of general relativity. They say that mass and energy curve spacetime, and curved spacetime tells mass how to move. They predict black holes, gravitational waves, and the expansion of the universe.

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu}$$

- **Assumptions:** Spacetime is a smooth, differentiable manifold. The metric tensor fully describes gravity. Matter and energy are described by a stress–energy tensor $T_{\mu\nu}$.

- **Limitations:** Does not incorporate quantum effects—breaks down at the Planck scale ($\sim 10^{-35}$ m). Cannot describe the singularity inside a black hole. Quantum gravity is needed for a complete theory.
- **Constraints:** Valid in the classical regime. Requires knowledge of the stress–energy tensor, which must be supplied by additional physics (e.g., fluid dynamics, electrodynamics).

7. **Faraday’s Law:** A changing magnetic field creates an electric field. This is how electric generators work.

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$$

- **Assumptions:** The electric and magnetic fields vary slowly compared to the speed of light. The medium is linear and time-invariant.
- **Limitations:** Does not account for quantum effects (e.g., the Aharonov–Bohm effect). Fails in nonlinear media or at very high field intensities.
- **Constraints:** Valid in the classical regime. The minus sign (Lenz’s law) assumes no magnetic monopoles.

8. **Gauss’s Law:** Electric charges produce electric fields. The total electric flux through a closed surface equals the enclosed charge divided by ϵ_0 .

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0}$$

- **Assumptions:** The electric field is created by static or slowly varying charge distributions. The permittivity ϵ_0 is a constant (true in vacuum).
- **Limitations:** Does not apply in quantum electrodynamics where virtual charges modify the effective vacuum permittivity. In dielectric media, the effective ϵ replaces ϵ_0 .
- **Constraints:** Valid for classical electrodynamics. Requires that magnetic monopoles do not exist (consistent with all observations to date).

9. **Gauss’s Law for Magnetism:** There are no magnetic monopoles—magnetic field lines always form closed loops.

$$\nabla \cdot \mathbf{B} = 0$$

- **Assumptions:** Magnetic monopoles do not exist. The divergence of the magnetic field is zero everywhere.
- **Limitations:** If magnetic monopoles were ever discovered, this equation would need to be modified (adding a magnetic charge density term). Some grand unified theories predict monopoles, but none have been observed.
- **Constraints:** Valid in all observed physical situations. Consistent with Maxwell’s equations and all experimental data.

10. **Law of Thermodynamics (First):** Energy is conserved. The heat added to a system goes into increasing its internal energy or doing work.

$$dU = \delta Q - \delta W$$

- **Assumptions:** The system is closed (no mass enters or leaves). Energy can be exchanged as heat or work, but not created or destroyed.
- **Limitations:** Does not predict the direction of processes (that is the second law). In relativistic systems, mass–energy equivalence must be included.
- **Constraints:** Applies to any thermodynamic process, reversible or irreversible. The internal energy U is a state function; δQ and δW are path-dependent.

11. **Law of Thermodynamics (Second):** Entropy (disorder) of an isolated system never decreases. Heat flows from hot to cold, never the other way.

$$\Delta S \geq 0$$

- **Assumptions:** The system is isolated (no energy or matter exchange with surroundings). The system is in or approaches thermal equilibrium.
- **Limitations:** Statistical in nature—entropy can locally decrease with extremely small probability. In small systems (nanoscale), thermal fluctuations can temporarily decrease entropy. Does not apply to non-equilibrium systems without modification.
- **Constraints:** Strictly valid only for isolated systems. For open systems, the entropy of the system plus its surroundings must increase.

12. **Law of Thermodynamics (Third):** As temperature approaches absolute zero, entropy approaches a minimum value. You can never actually reach absolute zero.

$$\lim_{T \rightarrow 0} S = S_0$$

- **Assumptions:** The system is in its quantum mechanical ground state. The third law applies to perfect crystals where $S_0 = 0$. Systems with residual entropy (e.g., ice) have $S_0 > 0$.
- **Limitations:** Does not apply to glasses or disordered systems, which can have residual entropy at $T = 0$. The unattainability statement is not rigorously proven for all systems.
- **Constraints:** Requires a perfect crystal at absolute zero to have zero entropy. Practically, absolute zero cannot be reached in finite steps (Nernst’s unattainability statement).

13. **Law of Thermodynamics (Zeroth):** If two objects are both in thermal equilibrium with a third, they are in equilibrium with each other. This defines temperature.

$$\text{If } A \sim C \text{ and } B \sim C, \text{ then } A \sim B$$

- **Assumptions:** Thermal equilibrium is a transitive relation. The systems can exchange energy (or have been in contact) and reach a common temperature.
- **Limitations:** Does not define temperature in non-equilibrium systems. Breaks down for systems with long-range interactions (e.g., gravitational systems) where thermal equilibrium may not be well-defined.

- **Constraints:** Valid for systems in local thermal equilibrium. Assumes that temperature is a well-defined scalar quantity.

14. **Lorentz Transformation:** The equations that replace Newton’s Galilean transformations when objects move close to the speed of light. They show that time slows down and lengths contract for moving objects, and they ensure that the speed of light is the same for everyone.

$$x' = \gamma(x - vt), \quad t' = \gamma\left(t - \frac{vx}{c^2}\right)$$

- **Assumptions:** The speed of light c is the same in all inertial reference frames. Space is flat (no gravity). The two frames are related by a constant relative velocity v .
- **Limitations:** Does not apply in curved spacetime (general relativity uses general coordinate transformations). Breaks down for accelerated observers (though it can be applied locally).
- **Constraints:** Valid only for $v < c$. For $v \geq c$, the Lorentz factor γ becomes imaginary or undefined. Reduces to Galilean transformation when $v \ll c$.

15. **Newton’s First Law (Inertia):** A body stays at rest or moves in a straight line at constant speed unless a net force acts on it.

$$\sum \mathbf{F} = 0 \Rightarrow \mathbf{v} = \text{const}$$

- **Assumptions:** The reference frame is inertial (not accelerating). The body is isolated from external forces.
- **Limitations:** Does not apply in non-inertial (accelerating or rotating) frames without adding fictitious forces. In general relativity, gravity is not a force but a curvature of spacetime, so the concept of “no net force” must be reinterpreted.
- **Constraints:** Defines what an inertial frame is. Cannot be proven independently—it is essentially the definition of an inertial reference frame.

16. **Newton’s Law of Universal Gravitation:** Any two masses attract each other with a force that depends on their masses and the distance between them. This single law explains planetary orbits, tides, and the structure of galaxies.

$$F = \frac{GMm}{r^2}$$

- **Assumptions:** The masses are point-like or spherically symmetric. The gravitational interaction is instantaneous (no propagation delay). Gravity is always attractive.
- **Limitations:** Fails for strong gravitational fields (need general relativity). Does not account for the finite speed of gravity (propagation at c). Cannot explain the precession of Mercury’s orbit or gravitational lensing.
- **Constraints:** Valid in the weak-field, low-velocity regime. For speeds $v \ll c$ and gravitational potentials $\Phi/c^2 \ll 1$, it matches general relativity to first order.

17. **Newton's Second Law:** The net force on an object equals its mass times acceleration. This is the central equation of mechanics.

$$\mathbf{F} = m\mathbf{a}$$

- **Assumptions:** The reference frame is inertial. The mass m is constant (no mass change during motion). The force is the net (total) force.
- **Limitations:** Does not apply to objects with changing mass (e.g., rockets expelling fuel) without using the more general form $\mathbf{F} = d\mathbf{p}/dt$. Fails at relativistic speeds—the correct form is $\mathbf{F} = d\mathbf{p}/dt$ where $\mathbf{p} = \gamma m\mathbf{v}$.
- **Constraints:** Valid for $v \ll c$. In non-inertial frames, fictitious forces must be added. For variable-mass systems, the general form $\mathbf{F} = d\mathbf{p}/dt$ must be used.

18. **Newton's Third Law:** Every force has an equal and opposite partner. When you push a wall, the wall pushes back with the same force.

$$\mathbf{F}_{12} = -\mathbf{F}_{21}$$

- **Assumptions:** The forces act along the line joining the two objects (central forces). The system is non-relativistic. No time delay in force propagation.
- **Limitations:** Does not hold for magnetic forces between moving charges (the forces are equal and opposite but not necessarily along the line joining the charges). In electrodynamics, the law must be generalized to include field momentum. Fails at relativistic speeds.
- **Constraints:** Valid for contact forces and central forces. Conservation of momentum follows from Newton's third law plus Newton's second law.

19. **Schrödinger Equation:** The central equation of quantum mechanics. It predicts how the wave function ψ of a particle evolves over time. If you know the wave function, you can calculate the probability of finding the particle anywhere.

$$i\hbar \frac{\partial \psi}{\partial t} = \hat{H}\psi$$

- **Assumptions:** The system is non-relativistic (the Hamiltonian contains $p^2/2m$, not the relativistic energy–momentum relation). The wave function ψ provides a complete description of the system. Spin is not included unless Pauli's equation is used.
- **Limitations:** Does not incorporate special relativity (the Dirac or Klein–Gordon equations are needed). Cannot describe particle creation or annihilation (need quantum field theory). The many-body problem is intractable analytically for more than a few particles.
- **Constraints:** Valid for $v \ll c$. Requires a well-defined potential $V(\mathbf{r}, t)$. The Hamiltonian must be Hermitian to ensure real energy eigenvalues and unitary time evolution.

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don’t just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: These equations are the grammar of the universe. Conservation laws say certain quantities—energy, momentum, charge—cannot be created or destroyed, only moved around. Maxwell’s equations say that shaking an electric charge creates a ripple that travels at the speed of light, and that light itself is an electromagnetic wave. Einstein’s $E = mc^2$ says that a tiny speck of matter holds a stupendous amount of energy, like a compressed spring waiting to release. The Schrödinger equation says that quantum particles behave like waves, with all the interference and fuzziness that implies.

The Metaphor: Think of these equations as the rules of a game. You did not invent the game—nature did. These equations describe what moves are legal. Conservation of energy says you cannot score points out of nothing. Maxwell’s equations say that electromagnetic “plays” propagate at a fixed speed. The quantum equations say that the game has built-in randomness at its smallest scales.

2. The Localized Mechanics

“Look at a tiny, microscopic point in space. What is happening there that these equations describe?”

He would press you on what each conservation law means at a single point. Conservation of energy at a point means: if energy is increasing somewhere, it must be flowing in from somewhere else—there is no local “source” of energy. Conservation of charge at a point means: if charge is accumulating, current must be flowing in. These are continuity equations—the same mathematical structure appears in fluid dynamics, electromagnetism, and quantum mechanics.

The Smoothness Check: He would ask you to explain why Maxwell’s equations predict that electromagnetic waves travel at a fixed speed $c = 1/\sqrt{\mu_0\epsilon_0}$. This is not a coincidence—it is a consequence of the fact that electric and magnetic fields are two aspects of one thing, and their mutual induction creates a self-sustaining wave at a speed determined by the properties of empty space itself.

3. Testing the Extreme Limits

“Let’s push these equations to their breaking points. Where do they fail?”

The High-Energy Limit: At very high energies (near the Big Bang), all four forces may have been unified into one. The Standard Model—our best collection of equations—does not include gravity. At the Planck scale (10^{-35} m, 10^{19} GeV), spacetime itself may become quantum foam, and all our equations may break down.

The Low-Energy Limit: At everyday energies, quantum effects are negligible and classical equations work beautifully. But at the atomic scale (10^{-10} m), quantum mechanics dominates. The equations do not “break down” at these limits—they reveal deeper structure that classical physics misses.

4. Dimensionality and Geometry

“Does this equation care about the shape of the space it lives in?”

He would point out that conservation laws are consequences of symmetries (Noether’s theorem). If the laws of physics do not change when you shift time (translation in time symmetry), energy is conserved. If they do not change when you shift space (translation in space symmetry), momentum is conserved. If they do not change when you rotate (rotational symmetry), angular momentum is conserved. The equations we know are not arbitrary—they are dictated by the symmetries of spacetime itself.

5. Cross-Disciplinary Connection

“You learned these in physics class, but where else do they appear?”

Conservation laws appear everywhere: in economics (conservation of money), in ecology (conservation of biomass), in information theory (conservation of information in reversible computing). Maxwell’s equations appear in any system where two fields regenerate each other—chemical reaction-diffusion systems, predator-prey populations, and even certain financial models. The deep reason is that the same mathematics describes any system where two quantities mutually sustain each other.

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don’t just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: No matter how perfect your lens is, light passing through a circular opening will always spread out into a fuzzy disk rather than focusing to a sharp point. The smaller the opening, the more it spreads. The 1.22 factor comes from the first zero of a Bessel function—a mathematical object that appears whenever circular symmetry meets wave physics.

The Metaphor: Imagine throwing a stone into a perfectly circular pond. The ripples spread in circles, but when they pass through a circular opening (like a drain), they produce a characteristic pattern. Light does the same thing. The Airy disk is nature’s way of saying: “You cannot confine a wave to a point without paying a price in spreading.”

2. The Localized Mechanics

“Look at the center of the Airy disk. What is happening to the light waves there?”

He would press you on why the central peak is brightest. At the exact center, all the wavelets from every part of the circular aperture arrive in phase—they all travel the same distance. They add constructively, producing maximum intensity. As you move away from the center, the wavelets arrive

with increasing phase differences. At the first dark ring, the phase differences are such that the contributions exactly cancel—for every wavelet that arrives “up,” another arrives “down.”

The Smoothness Check: He would ask you to explain why the central peak is twice as wide as the others. This is because the central peak spans from the first zero on one side to the first zero on the other side, while the side peaks span from one zero to the next. The zeros of the Bessel function are not equally spaced—they get closer together as you move outward.

3. Testing the Extreme Limits

“What happens when we push the aperture to extreme sizes?”

The Zero-Aperture Limit: As the aperture shrinks to zero, the Airy disk grows to infinity—you see nothing. This is the uncertainty principle in action: confining the photon’s position (Δx small) spreads out its momentum (Δp large).

The Infinite-Aperture Limit: As the aperture grows to infinity, the Airy disk shrinks to a perfect point—geometric optics. But an infinite aperture is impossible, so there is always some blur.

The Quantum Limit: What happens when you send photons through one at a time? Each photon lands at a single point (particle behavior), but after many photons, the Airy disk pattern emerges (wave behavior). This is wave-particle duality in action.

4. Dimensionality and Geometry

“What if the aperture were a slit instead of a circle? How would the pattern change?”

He would point out that a circular aperture produces an Airy disk (Bessel function), while a slit produces a sinc-squared pattern (sine function). The mathematics changes because the symmetry changes—circular symmetry gives Bessel functions, rectangular symmetry gives sine functions. The general principle remains: the diffraction pattern is the Fourier transform of the aperture shape.

5. Cross-Disciplinary Connection

“You learned this in optics class, but where else does this pattern appear?”

The Airy disk appears in radio astronomy (the beam pattern of a circular dish antenna), in medical imaging (the resolution limit of CT and MRI scanners), in crystallography (the diffraction pattern of X-rays from crystals), and even in acoustics (the sound pattern of a circular speaker). The same mathematics—the Fourier transform of a circular function—appears in any field where waves pass through circular openings.

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don’t just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: When a fluid speeds up, its pressure drops—not because something is “sucking” the pressure away, but because the energy that was in pressure form has been converted to kinetic energy. The total energy budget of a fluid parcel is strictly conserved: pressure energy + kinetic energy + gravitational potential energy = constant along a streamline.

The Metaphor: Imagine a crowded hallway that narrows. People must walk faster to get through. In doing so, they spread out and push less on the walls. The pressure (pushing on the walls) drops because the energy is now in motion (kinetic energy) rather than in pushing. Water in a pipe does exactly the same thing.

2. The Localized Mechanics

“Look at a tiny fluid parcel as it speeds up. What is happening to the pressure right there?”

He would press you on the energy trade-off. The equation $P + \frac{1}{2}\rho v^2 + \rho gh = \text{constant}$ says that if v increases, P must decrease (assuming h is constant). This is not a force—it is an energy conservation statement applied to a fluid element. The fluid does not “know” about pressure drops; it simply follows the path of least resistance, which happens to be the path that conserves energy.

The Smoothness Check: He would ask why the equation applies only along a streamline. Because different streamlines can have different total energies (different constants), you cannot compare pressure between two streamlines using Bernoulli’s equation. The equation is a statement about energy conservation along a single path, not across the entire flow.

3. Testing the Extreme Limits

“What happens when we push the fluid to extreme speeds?”

The Speed-of-Sound Limit: As fluid speed approaches the speed of sound, compressibility matters—the density changes, and Bernoulli’s equation breaks down. You need the full equations of compressible fluid dynamics.

The Zero-Speed Limit: At $v = 0$ (static fluid), Bernoulli reduces to $P + \rho gh = \text{constant}$ —hydrostatics. The pressure increases linearly with depth, which is why your ears pop when you dive underwater.

The Superfluid Limit: In superfluid helium (below 2.17 K), viscosity vanishes and Bernoulli’s equation applies with perfect accuracy—there are no dissipative losses. The superfluid is the closest thing to an ideal fluid that exists in nature.

4. Dimensionality and Geometry

“Does this equation care about the shape of the pipe or channel?”

He would point out that Bernoulli's equation does not care about the shape—only about the changes in velocity, height, and pressure along a streamline. A pipe can be straight, curved, widening, or narrowing—as long as the flow is steady, incompressible, and inviscid, Bernoulli applies. The shape matters only insofar as it determines how the velocity changes.

5. Cross-Disciplinary Connection

“You learned this in fluid dynamics class, but where else does this exact same principle appear?”

The same energy trade-off appears in economics (velocity of money vs. “pressure” of interest rates), in traffic flow (when lanes narrow, cars speed up and “pressure”—density—drops), and in electrical circuits (voltage is analogous to pressure, current to velocity, and the energy trade-off between kinetic and potential energy is analogous to the trade-off between kinetic and pressure energy in fluids).

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don't just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: This equation tracks how the crowd of particles—each moving with its own velocity—redistributes itself over space and time. It says that the number of particles at a given position and velocity changes because of three effects: (1) particles stream from one place to another, (2) external forces push particles from one velocity to another, and (3) collisions transfer particles between velocities.

The Metaphor: Imagine watching a crowded dance floor from above. Each dancer moves according to the music (external forces) and bumps into other dancers (collisions). The Boltzmann equation tracks how the crowd's density changes at each location and velocity. When you heat one end of a metal rod, the atoms there vibrate more vigorously and collide with their neighbors, passing the energy along—this is heat conduction, and it emerges from the Boltzmann equation.

2. The Localized Mechanics

“Look at a single particle in the distribution. What happens to it during a collision?”

He would press you on the collision integral—the most complex part of the equation. A collision takes two particles with specific velocities and produces two particles with different velocities. The collision integral counts how many such collisions happen per unit time, and how they redistribute the particles among different velocities. The key insight is that collisions conserve mass, momentum, and energy—but they can change the distribution from non-equilibrium to equilibrium.

The Smoothness Check: He would ask why the collision term is quadratic in f (the distribution function). Because collisions involve *pairs* of particles—the rate of collisions is proportional to the number of pairs, which is $\sim f^2$. This nonlinearity makes the Boltzmann equation very hard to solve.

3. Testing the Extreme Limits

“What happens when we push the gas to extreme conditions?”

The Dilute Gas Limit: At very low densities, collisions vanish ($f \rightarrow 0$ in the collision integral) and the equation becomes a streaming equation—particles move freely without interacting. This is the “free molecular flow” regime, relevant for spacecraft in the upper atmosphere.

The Dense Gas Limit: At very high densities, the concept of binary (two-body) collisions breaks down—particles interact simultaneously with many neighbors. The Boltzmann equation must be replaced by the BBGKY hierarchy or molecular dynamics simulations.

The Equilibrium Limit: At long times, the collision term drives the distribution toward the Maxwell-Boltzmann distribution—the unique distribution where collisions produce no net change. This is the H-theorem: entropy increases monotonically until equilibrium is reached.

4. Dimensionality and Geometry

“Does this equation care about the dimensionality of space?”

He would point out that the Boltzmann equation lives in a 6-dimensional phase space (3 for position, 3 for velocity). In 2D, the velocity space is 2-dimensional and the collision dynamics are simpler. In 1D, the equation can sometimes be solved exactly. The physics is the same in all dimensions—transport, relaxation to equilibrium, and the H-theorem—but the mathematical complexity changes dramatically.

5. Cross-Disciplinary Connection

“You learned this in kinetic theory, but where else does this exact same equation appear?”

The Boltzmann equation appears in radiative transfer (photons scattering in stellar atmospheres), in neutron transport (neutrons diffusing in a nuclear reactor), in semiconductor physics (electrons scattering in a crystal lattice), and even in social dynamics (“opinions” transported and “colliding” in a population). The same mathematical structure—streaming + forcing + collision—appears whenever a population of entities moves, interacts, and redistributes.

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don’t just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: Certain particles are social butterflies—they like to do the same thing at the same time. The minus sign in the denominator ($e^{(E-\mu)/kT} - 1$) is the key: it means that the more particles already in a state, the more likely another particle is to join them. This is the opposite of fermions, which avoid each other (plus sign in the denominator).

The Metaphor: Imagine a concert where everyone wants the same spot. As more people crowd in, the spot becomes more attractive—it must be the best view! Bosons are like these concert-goers. When you cool rubidium atoms to near absolute zero, they all collapse into one quantum state and move as a single entity—a Bose-Einstein condensate, a new state of matter where atoms lose their individual identity.

2. The Localized Mechanics

“Look at a single quantum state. How does the presence of one boson affect the probability of another boson joining it?”

He would press you on the stimulation factor. The occupation number of a state is $\langle n \rangle = 1/(e^{(E-\mu)/kT} - 1)$. If you already have n bosons in a state, the probability of adding one more is proportional to $(n + 1)$ —not n . This “bosonic stimulation” is the mathematical expression of their social behavior. It is the same physics that makes lasers work: photons stimulate the emission of more photons, all in the same quantum state.

The Smoothness Check: He would ask why the chemical potential μ must be less than the ground state energy. Because if μ equaled the ground state energy, the denominator would be zero and the occupation number would diverge—an infinite number of particles in one state. This is exactly what happens in a Bose-Einstein condensate: the ground state macroscopically occupied.

3. Testing the Extreme Limits

“What happens when we push the temperature to extreme values?”

The High-Temperature Limit: As $T \rightarrow \infty$, the exponential dominates and $e^{(E-\mu)/kT} \gg 1$, so the minus sign becomes irrelevant. The Bose-Einstein distribution reduces to the classical Maxwell-Boltzmann distribution—quantum effects wash out.

The Zero-Temperature Limit: As $T \rightarrow 0$, all particles collapse into the ground state—the condensate. The occupation of the ground state becomes macroscopic ($\sim N$), while all excited states are empty. This is a phase transition: the gas suddenly changes from a collection of individual atoms to a single quantum entity.

4. Dimensionality and Geometry

“Does this equation care about the dimensionality of space?”

He would point out that Bose-Einstein condensation only occurs in 3D and higher. In 1D and 2D, the density of states is such that the integral for the total number of particles diverges—you cannot fit all the particles into the excited states at any finite temperature. But in 2D, a condensate can form if the particles are trapped in a harmonic potential (as in experiments with cold atoms).

5. Cross-Disciplinary Connection

“You learned this in quantum statistics, but where else does this exact same principle appear?”

The Bose-Einstein distribution appears in photon statistics (blackbody radiation is a gas of photons obeying this distribution), in phonon statistics (lattice vibrations in solids), and in the statistics of any indistinguishable particles with integer spin. The same “social” behavior appears in network theory: nodes with many connections tend to acquire even more connections (the “rich get richer” phenomenon), which is mathematically analogous to bosonic stimulation.

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don’t just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: When you increase the pressure on a liquid, you raise its boiling point—the molecules need more kinetic energy to escape the tighter packing. This is why pressure cookers work: by increasing the pressure, they delay boiling and allow temperatures above 100°C. For water, the equation also explains why ice floats: solid water is less dense than liquid water, so the solid-liquid boundary has a negative slope—increasing pressure actually lowers the melting point.

The Metaphor: Think of molecules at a phase boundary as people at a doorway. Pressure is like pushing the crowd together—it makes it harder for individuals to push through the door (escape into the gas phase). You need to give them more energy (higher temperature) to break free.

2. The Localized Mechanics

“Look at the phase boundary itself. What is happening to the molecules right at the interface?”

He would press you on the latent heat. The equation $dP/dT = L/(T\Delta V)$ says that the slope of the phase boundary in a P - T diagram is determined by the latent heat L (energy needed to convert one phase to another) and the volume change ΔV . Large latent heat means a steep boundary—small temperature changes cause large pressure changes. Large volume change means a steep boundary—the two phases are very different.

The Smoothness Check: He would ask what happens when $L = 0$. The slope goes to zero—the phase boundary becomes horizontal. This is the triple point: three phases coexist at a single temperature and pressure. At the critical point, $L \rightarrow 0$ and $\Delta V \rightarrow 0$ simultaneously—the distinction between liquid and gas vanishes.

3. Testing the Extreme Limits

“What happens when we push the system to extreme pressures and temperatures?”

The Critical Point: At the critical point (T_c, P_c) , the latent heat goes to zero and the Clausius-Clapeyron equation gives a divergence (0/0). The equation breaks down because there is no longer a sharp phase transition—liquid and gas become indistinguishable.

The Triple Point: At the triple point, three phases coexist. The Clausius-Clapeyron equation applies to each pairwise boundary, but the three boundaries meet at a single point. This is the unique thermodynamic state where ice, water, and steam coexist.

The Negative Slope: For water, $dP/dT < 0$ for the solid-liquid boundary because ice is less dense than water. Increasing pressure lowers the melting point—this is why glaciers flow under their own weight: the pressure at the base melts the ice, creating a lubricating layer of water.

4. Dimensionality and Geometry

“Does this equation care about the shape of the container?”

He would point out that the Clausius-Clapeyron equation is independent of container shape—it is a statement about bulk thermodynamic phases, not about geometry. The equation applies equally to a beaker of water, a sealed pressure cooker, or the ocean. The phase boundary is determined by temperature and pressure alone, not by the shape of the vessel.

5. Cross-Disciplinary Connection

“You learned this in thermodynamics class, but where else does this exact same principle appear?”

The same “slope = latent heat / (temperature × volume change)” structure appears in any first-order phase transition: transitions in magnets (the magnetic analogue of latent heat is the change in magnetization), structural phase transitions in crystals (the volume change is the change in lattice parameters), and even in economics (the “latent heat” of a market crash is the change in investor sentiment, and the “volume change” is the change in market liquidity).

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don’t just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: Matter is a perfect accountant. If water level rises at one point, more water is flowing in than out. If it drops, more is flowing out than in. The continuity equation is the mathematical expression of this simple bookkeeping principle: the rate of change of mass in any region equals the net mass flowing in through the boundary.

The Metaphor: Imagine watching a river from a bridge. If the water level rises at one point, it must be because more water is flowing in than flowing out. If you squeeze a tube narrower, the fluid must speed up to maintain the same flow rate—this is why putting your thumb over a garden hose makes the water spray faster. The equation is the same whether the fluid is water, air, blood, or traffic on a highway.

2. The Localized Mechanics

“Look at a tiny control volume in the fluid. What is the balance of mass right there?”

He would press you on the divergence term. $\nabla \cdot (\rho \mathbf{v})$ measures the net outward flux of mass from a point. If it is positive, mass is flowing away and the local density decreases. If it is negative, mass is flowing in and the density increases. The continuity equation says these two effects exactly balance: $\partial \rho / \partial t + \nabla \cdot (\rho \mathbf{v}) = 0$.

The Smoothness Check: He would ask why the equation is first-order in time but first-order in space. Because it is a conservation law—it tracks how a conserved quantity (mass) flows. The time derivative tells you the local rate of change; the spatial derivative tells you the flux. Together, they ensure that mass is neither created nor destroyed.

3. Testing the Extreme Limits

“What happens when we push the flow to extreme conditions?”

The Incompressible Limit: For liquids (approximately incompressible), ρ is constant and the equation simplifies to $\nabla \cdot \mathbf{v} = 0$ —the velocity field is divergence-free. This is a powerful constraint: it says that fluid cannot be compressed or expanded, only redirected.

The Compressible Limit: For gases at high speeds (near the speed of sound), density changes matter and the full equation must be used. At supersonic speeds, shock waves form where density changes discontinuously—the continuity equation still holds, but the flow is no longer smooth.

The Vacuum Limit: In a vacuum ($\rho = 0$), the equation is trivially satisfied—there is nothing to conserve. But if you try to create a perfect vacuum, quantum effects (virtual particles) ensure that it is never truly empty.

4. Dimensionality and Geometry

“Does this equation care about the dimensionality of space?”

He would point out that the continuity equation has the same form in 1D, 2D, and 3D—only the divergence operator changes. In 1D, it is $d(\rho v)/dx = -\partial \rho / \partial t$. In 2D, it involves partial derivatives in x and y . The physics is identical: mass is conserved, and the equation tracks how it flows.

5. Cross-Disciplinary Connection

“You learned this in fluid dynamics, but where else does this exact same equation appear?”

The continuity equation appears in electromagnetism (conservation of charge: $\partial \rho_e / \partial t + \nabla \cdot \mathbf{J} = 0$), in quantum mechanics (conservation of probability: $\partial |\psi|^2 / \partial t + \nabla \cdot \mathbf{J}_\psi = 0$), in heat transfer (conservation of energy), and in population dynamics (conservation of organisms). The same mathematical structure—the rate of change plus the divergence of a flux equals zero—is the universal signature of a conservation law.

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don’t just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: Nature abhors a gradient. When you drop ink into water, it does not stay as a concentrated blob—it spreads out, becoming more dilute until the color is uniform everywhere. The rate of spreading is proportional to how “curved” the concentration profile is: where the curve is steep, spreading is fast; where it is flat, spreading is slow.

The Metaphor: Think of a crowded room. People naturally spread out from the crowded area to the less crowded area, not because there is a force pushing them, but because there are more ways to be spread out than to be concentrated. The diffusion equation describes exactly how fast this spreading happens. The thermal diffusivity α is the material’s “spreading speed”—metals have high α (heat spreads quickly), wood has low α (heat spreads slowly).

2. The Localized Mechanics

“Look at a single point in the material. What determines how fast the temperature changes there?”

He would press you on the Laplacian. $\nabla^2 T$ measures how much the temperature at a point differs from the average temperature of its neighbors. If the point is hotter than its neighbors ($\nabla^2 T < 0$), the temperature there decreases—heat flows out. If it is colder ($\nabla^2 T > 0$), the temperature there increases—heat flows in. The diffusion equation says: the rate of temperature change equals the diffusivity times this “neighborhood difference.”

The Smoothness Check: He would ask why the equation is first-order in time but second-order in space. Because diffusion is irreversible—you can unscramble an egg in principle (reverse the trajectories of all molecules), but you never see it happen in practice. The first time derivative means the equation has a preferred direction in time: forward.

3. Testing the Extreme Limits

“What happens when we push the diffusion to extreme conditions?”

The Short-Time Limit: At very short times after a sudden temperature change, the diffusion equation predicts infinite propagation speed—a sharp spike instantly affects points arbitrarily far away. This is physically impossible and signals the breakdown of the diffusion approximation. At very short times, you need the hyperbolic heat equation (which includes a finite speed of propagation).

The Long-Time Limit: At very long times, the temperature becomes uniform everywhere—maximum entropy. The diffusion equation predicts exponential decay of temperature differences, with a time constant that depends on the geometry and the diffusivity.

The Quantum Limit: The Schrödinger equation is the diffusion equation with imaginary time ($t \rightarrow i\tau$). This deep connection means that quantum mechanics and diffusion are mathematically the same process—just with different boundary conditions and different physical interpretations.

4. Dimensionality and Geometry

“How does the dimensionality of space affect how things diffuse?”

He would point out that in 1D, the diffusion length grows as $\sqrt{t}$ (the famous “random walk” scaling). In 2D, it also grows as $\sqrt{t}$ but with a different prefactor. In 3D, the same scaling applies. The $\sqrt{t}$ scaling is universal—it comes from the central limit theorem, which says that the sum of many random steps grows as the square root of the number of steps.

5. Cross-Disciplinary Connection

“You learned this in heat transfer, but where else does this exact same equation appear?”

The diffusion equation describes heat conduction in solids, pollutant dispersion in groundwater, drug diffusion through tissue, neutron diffusion in nuclear reactors, and even the random walk of stock prices (the Black-Scholes equation for financial options is a disguised diffusion equation). The same mathematics applies because all these processes involve the random redistribution of a conserved quantity.

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don’t just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: When Dirac tried to write a quantum equation that obeyed Einstein’s relativity, he discovered extra solutions—solutions that described particles with positive energy but negative charge. These were not mathematical artifacts; they were real particles (positrons), discovered in 1932. The equation also says that electrons have an intrinsic angular momentum (spin) even though they are point particles—spin is not the electron “spinning” like a top, but a fundamental quantum property that arises naturally from combining relativity with quantum mechanics.

The Metaphor: Dirac was looking for a description of the electron that was consistent with both quantum mechanics and relativity. He found it—but the equation had “extra pages” that he initially tried to throw away. Those extra pages turned out to describe antimatter. The universe, it seems, has more symmetry than we expected: for every particle, there is an antiparticle.

2. The Localized Mechanics

“Look at the electron as described by the Dirac equation. What is spin, physically?”

He would press you on the relationship between spin and relativity. In the non-relativistic limit, the Dirac equation reduces to the Pauli equation—which includes spin as an added ingredient. But in the Dirac equation, spin is not added—it emerges naturally. The electron’s spin is a consequence of the requirement that the equation be Lorentz covariant (the same in all reference frames). Spin is not a classical property; it is a relativistic quantum property.

The Smoothness Check: He would ask why the Dirac equation uses 4×4 matrices (gamma matrices) instead of ordinary numbers. Because the electron has two internal degrees of freedom (spin up and spin down) and the equation must account for both left-handed and right-handed components. The 4×4 structure is the minimal representation of the Lorentz group that includes spin-1/2 particles.

3. Testing the Extreme Limits

“What happens when we push the electron to extreme energies?”

The Pair Production Limit: At energies above $2m_e c^2 = 1.022$ MeV, photons can create electron-positron pairs. The Dirac equation naturally describes this process: the negative-energy solutions (positrons) are essential for pair production.

The Ultra-Relativistic Limit: At energies much greater than $m_e c^2$, the electron behaves as if it were massless—the mass term becomes negligible and the Dirac equation simplifies. This is the regime of particle physics, where electrons and quarks are effectively massless compared to their kinetic energies.

The Planck Scale: At energies near 10^{19} GeV, gravity becomes important and the Dirac equation must be extended to include gravitational interactions. This is the frontier of theoretical physics—we do not yet have a consistent theory of quantum gravity.

4. Dimensionality and Geometry

“Does this equation care about the geometry of spacetime?”

He would point out that the Dirac equation can be formulated on any curved spacetime—you just replace the partial derivatives with covariant derivatives. This is how general relativity and quantum mechanics are combined (at least approximately). The spin connection—the gravitational analogue of the electromagnetic vector potential—couples the electron’s spin to the curvature of spacetime.

5. Cross-Disciplinary Connection

“You learned this in particle physics, but where else does this equation appear?”

The Dirac equation appears in condensed matter physics, where it describes the behavior of electrons in graphene (a 2D material where electrons behave as massless Dirac fermions), in topological insulators (where surface states are described by the Dirac equation), and in the quantum Hall effect. The same mathematics describes any system where spin-1/2 particles move through a periodic potential.

What Richard Feynman Would Have Asked

1. The Plain-English Translation

“Don’t just repeat the symbols to me. What is this equation actually saying in plain, everyday language?”

The Core Meaning: Light, radio waves, X-rays, and gamma rays are all the same thing—electromagnetic waves traveling at the speed of light. When Maxwell derived this equation in 1865, he calculated the speed from the measured values of the electrical and magnetic constants and found it matched the speed of light exactly. This was one of the greatest unifications in physics.

The Metaphor: When you shake an electric charge, it creates a changing electric field. This changing electric field creates a changing magnetic field. This changing magnetic field creates a changing electric field. And so on—the disturbance propagates outward at the speed of light. Empty space itself has electrical and magnetic properties (ϵ_0 and μ_0) that determine how fast this wave travels.

2. The Localized Mechanics

“Look at a single point in space as the wave passes. What is happening to the electric and magnetic fields right there?”

He would press you on the mutual induction. At any point, the changing electric field creates a magnetic field, and the changing magnetic field creates an electric field. The two fields are not independent—they are two aspects of one thing, the electromagnetic field. The wave equation describes how this coupled system propagates through space.

The Smoothness Check: He would ask why the wave speed is $c = 1/\sqrt{\mu_0\epsilon_0}$. Because ϵ_0 determines how much electric field is created by a charge, and μ_0 determines how much magnetic field is created by a current. The product $\mu_0\epsilon_0$ determines how quickly the “relay” between electric and magnetic fields occurs, and therefore how fast the wave propagates.

3. Testing the Extreme Limits

“What happens when we push electromagnetic waves to extreme frequencies?”

The Gamma-Ray Limit: At very high frequencies (gamma rays), photons have enough energy to create particle-antiparticle pairs. The classical wave equation breaks down—you need quantum electrodynamics (QED).

The Radio-Wave Limit: At very low frequencies (radio waves), the wavelength can be kilometers long. The wave equation still applies, but the practical challenges of generating and detecting such long waves are enormous.

The Static Limit: At zero frequency, there is no wave—just a static electric or magnetic field. Is a static field a “wave” with infinite wavelength? Feynman would say: in a sense, yes—it is the zero-frequency limit of the wave equation.

4. Dimensionality and Geometry

“Does this equation care about the shape of the space it lives in?”

He would point out that the wave equation has the same form in any dimension—1D (waves on a string), 2D (waves on a drumhead), and 3D (electromagnetic waves in space). The only difference is the Laplacian operator: in 1D it is d^2/dx^2 , in 2D it involves x and y , and in 3D it involves x , y , and z . The physics is identical: disturbances propagate as waves at a speed determined by the medium.

5. Cross-Disciplinary Connection

“You learned this in electromagnetism, but where else does this exact same equation appear?”

The wave equation describes sound waves in air, seismic waves in the Earth, vibrations in guitar strings, and quantum mechanical probability waves. The Schrödinger equation is actually a modified wave equation (with a first time derivative instead of second). The same mathematics describes any system where a restoring force (proportional to displacement) and inertia (proportional to acceleration) are balanced.

Historical Notes

Equation	Derived by	Year
Lorentz Force Law	Hendrik Lorentz	1892
Biot–Savart Law	J.-B. Biot & F. Savart	1820
Heisenberg Uncertainty Principle	Werner Heisenberg	1927
De Broglie Wavelength	Louis de Broglie	1924
Snell’s Law of Refraction	Willebrord Snellius	1621
Young’s Double-Slit	Thomas Young	1801
Kepler’s Third Law	Johannes Kepler	1619
Damped driven Harmonic Oscillator	Galileo Galilei / Robert Hooke	1630s / 1678
Poisson’s Equation	Siméon Denis Poisson	1813
Airy Disk	George Biddell Airy	1835
Bernoulli Equation	Daniel Bernoulli	1738
Boltzmann Transport Eq.	Ludwig Boltzmann	1872
Bose–Einstein Distribution	S. N. Bose & A. Einstein	1924
Boundary Layer Eq.	Ludwig Prandtl	1904
Carnot Efficiency	Sadi Carnot	1824
Clausius–Clapeyron Eq.	B. P.	
’E. Clapeyron / R. Clausius	1834 / 1850	
Conservation of Mech. Energy		
’Emilie du Châtelet	1740s	
Continuity Equation	Leonhard Euler	1757
Diffusion Equation	Adolf Fick	1855
Dirac Equation	Paul Dirac	1928
EM Wave Equation	James Clerk Maxwell	1865
Entropy	Rudolf Clausius	1865
Euler Equation	Leonhard Euler	1757

Equation	Derived by	Year
Euler–Bernoulli Beam Eq.	L. Euler & D. Bernoulli	1744
Euler–Lagrange Eq.	L. Euler & J.-L. Lagrange	1750s
Fermi–Dirac Distribution	E. Fermi & P. Dirac	1926
Fraunhofer Diffraction	Joseph von Fraunhofer	1821
Fresnel Equations	Augustin-Jean Fresnel	1823
Friedmann Equation	Alexander Friedmann	1922
Geodesic Equation	Albert Einstein / R. Ricci & T. Levi-Civita	1915
Hamilton’s Equations	William Rowan Hamilton	1834
Heat Equation	Joseph Fourier	1822
Heisenberg Equation	Werner Heisenberg	1925
Ideal Gas Equation		
’E. Clapeyron (Boyle, Charles, Avogadro)	1834	
Kinematic Equations	Galileo Galilei	1638
Klein–Gordon Eq.	Oskar Klein & Walter Gordon	1926
Length Contraction	G. FitzGerald & H. Lorentz	1889 / 1892
Liénard–Wiechert Pot.	A.-M. Liénard & E. Wiechert	1898 / 1900
Mass–Energy Equivalence	Albert Einstein	1905
Maxwell Relation	James Clerk Maxwell	1865
Maxwell–Boltzmann Dist.	J. C. Maxwell & L. Boltzmann	1860 / 1868
Mirror Equation	Euclid / Ibn al-Haytham / Kepler	c. 300 BC / 1021 / 1611
Navier–Stokes Eq.	C.-L. Navier & G. Stokes	1822 / 1845
Partition Function	L. Boltzmann & J. W. Gibbs	1877 / 1902
Pauli Equation	Wolfgang Pauli	1927
Planck Radiation Law	Max Planck	1900
Plate Equation	A. E. H. Love	1888
Laplace’s Equation	D. Bernoulli & H. von Helmholtz	1738 / 1858
Poynting Theorem	John Henry Poynting	1884
Probability Current	Max Born	1926
Quantum Continuity Eq.	Max Born	1926

Equation	Derived by	Year
Relativistic Energy	Albert Einstein	1905
Relativistic Momentum	A. Einstein & M. Planck	1905 / 1906
Reynolds Transport Thm.	Osborne Reynolds	1903
Rotational Kinetic Energy	Leonhard Euler	1750s
Sackur–Tetrode Eq.	H. M. Sackur & O. Tetrode	1911 / 1912
Schwarzschild Metric	Karl Schwarzschild	1916
Single-Slit Diffraction	Joseph von Fraunhofer	1818
Skin Depth	H. Lamb & O. Heaviside	1897 / 1899
Stefan–Boltzmann Law	J. Stefan & L. Boltzmann	1879 / 1884
Stress Equilibrium Eq.	Augustin-Louis Cauchy	1822
Stress–Strain Relation	Robert Hooke	1678
Telegraph Equation	Oliver Heaviside	1887
Thin Lens Equation	Ptolemy / Ibn al-Haytham / Snel- lius	c. 150 AD / 1021 / 1621
Time Dilation	H. Lorentz & A. Einstein	1892 / 1905
Time-Indep. Schröd. Eq.	Erwin Schrödinger	1926
Torque	Archimedes / Leonhard Euler	c. 250 BC / 1750s
Vorticity Equation	H. von Helmholtz / Navier / Stokes	1858
Wave Equation	Jean le Rond d’Alembert	1747
Wave Eq. in Solids	Claude-Louis Navier	1822
Wien’s Displacement Equation	Wilhelm Wien	1893
Work–Energy Equation	G. G. de Coriolis & J.-V. Poncelet	1826 / 1829