


# MODERN MEDICAL POCKET DICTIONARY

SIMPLE TERMS  
CLEAR DEFINITIONS  
A HEALTHIER TOMORROW

CONCISE  
RELIABLE  
UP-TO-DATE


A COMPREHENSIVE GUIDE TO  
TODAY'S MEDICAL TERMS



ANATOMY



PHYSIOLOGY



DISEASES &  
CONDITIONS



SYMPTOMS  
& SIGNS



DIAGNOSIS  
& PROCEDURES



LABORATORY  
TESTS



IMAGING



BODY SYSTEMS



LIFE STAGES  
& SPECIALTIES



PUBLIC HEALTH  
& MORE

*Knowledge for  
Better Health*



A COMPACT REFERENCE  
FOR STUDENTS, PROFESSIONALS  
AND HEALTH-CONSCIOUS READERS

CHAMAN SINGH VERMA

---

## Preface

This dictionary is a concise reference to medical terms, diseases, investigations, treatments, anatomy, physiology, and related clinical concepts. It is intended to help students, healthcare professionals, educators, and general readers understand unfamiliar medical language.

The entries are arranged alphabetically from A to Z, with each letter presented in its own chapter. This arrangement allows a reader to locate a known term quickly while maintaining a consistent structure throughout the book. Related terms, recognized common names, and alternative names are included when they improve clarity or aid searching.

Definitions are written in simple, formal language and are intended to be clear, accurate, and useful for reference. Medical terminology changes as scientific knowledge and clinical guidance develop; therefore, important information should be checked against current textbooks, primary research, professional guidance, and local clinical protocols.

This book is an educational reference, not a substitute for professional medical advice, diagnosis, or treatment. Readers should seek advice from a qualified healthcare professional when making decisions about health.

*Chaman Singh Verma*

## Medical Disclaimer

**Educational and Reference Purpose Only:** This publication, *The Modern Medical Pocket Dictionary*, is prepared and compiled strictly for educational, academic, historical, and reference purposes. It is intended for medical students, healthcare professionals, educators, and researchers.

**No Medical Advice:** The information contained herein does not constitute professional medical advice, clinical diagnosis, treatment protocols, or formal therapeutic recommendations. It should not be used as a substitute for the professional judgment of qualified physicians, surgeons, or certified healthcare providers.

**Clinical Practice Guidelines:** Healthcare practitioners must always consult primary evidence-based medical literature, institutional protocols, manufacturer prescribing information, and national clinical guidelines (such as WHO, CDC, NICE, or national medical councils) when making diagnostic or therapeutic decisions for individual patients.

**Limitation of Liability:** Neither the author nor the publishers accept any legal responsibility or liability for any errors, omissions, or clinical outcomes resulting from the use or application of any information presented in this work.

# A

## A & E Medicine

Accident and emergency (A&E) medicine provides rapid assessment, stabilization, diagnosis, and initial treatment for sudden illness and injury.

## A Fib

Alternate name; see *Atrial fibrillation*.

## A-T

The adenine–thymine base pair in DNA. The two bases join by hydrogen bonds and help form the structure of DNA.

## A1AT Deficiency

Abbreviation for alpha-1 antitrypsin deficiency, an inherited disorder that can cause early emphysema, liver disease, or both because protective alpha-1 antitrypsin levels are too low or abnormal.

## Aagenaes Syndrome

A rare inherited disorder that causes reduced flow of bile from the liver beginning in infancy and long-lasting swelling from lymph-fluid buildup, usually in the legs. Liver problems may come and go with age, but some people develop vitamin deficiency, poor growth, or liver scarring.

## Aarskog Syndrome

Alternate name; see *Aarskog-Scott Syndrome*.

## Aarskog-Scott Syndrome

A rare inherited developmental disorder, often related to FGD1 variants, characterized by distinctive facial features, short stature, skeletal differences, and genital anomalies.

## Alternate Names:

Aarskog Syndrome

## Aase Syndrome

Alternate name; see *Aase-Smith Syndrome II*.

## Aase-Smith Syndrome I

A rare condition present at birth that may cause excess fluid in the brain, a split in the roof of the mouth, joints that are fixed in bent positions, and unusual hand and facial features. Some people also have heart defects.

## Aase-Smith Syndrome II

A rare inherited disorder that causes reduced production of red blood cells and thumbs with three bones instead of the usual two.

## AAT

Abbreviation for alpha-1 antitrypsin, a protein made mainly in the liver that helps protect lung tissue from enzyme-related damage. Blood testing can measure its amount and type.

## AAT Deficiency

Alternate name for alpha-1 antitrypsin deficiency, an inherited condition associated with emphysema, liver disease, panniculitis, and rarely vasculitis.

## Abadie Sign

Reduced pain sensation when firm pressure is applied to the Achilles tendon, classically described in *tabes dorsalis*.

---

## Abandoned Lead

A wire from a pacemaker, defibrillator, or other implanted electrical device that is no longer connected or used but remains inside the body. A working lead carries electrical signals to the heart and senses its rhythm; its metal core conducts electricity and its silicone or polyurethane covering protects nearby tissue. These materials are chosen to be biocompatible, corrosion-resistant, strong, and flexible, but no implanted material is completely risk-free. A lead may be left in place when removal would be more dangerous because scar tissue can firmly attach it to a vein or the heart. Retained leads can later complicate infection treatment, blood-vessel access, or placement of another device.

### Alternate Names:

Retained Cardiac Device Lead; Abandoned Pacemaker Lead; Abandoned ICD Lead

## Abasia

Inability to walk even though the legs may have enough strength to move in other situations. It may result from a problem with coordination, the nervous system, or the way movement is planned, rather than paralysis.

## Abate

To become less intense or to reduce in severity, frequency, or amount; symptoms, inflammation, pain, or fever may abate.

## Abatement

A reduction in the severity, intensity, frequency, or amount of a symptom, disease process, exposure, or other condition.

## ABCDE Assessment

A systematic emergency assessment of airway, breathing, circulation, disability, and exposure. Life-threatening abnormalities are treated as they are identified, with repeated reassessment.

## ABCDE Rule For Melanoma

A screening mnemonic for suspicious pigmented lesions: asymmetry, border irregularity, color variation, diameter greater than about 6 mm, and evo-

lution. It does not diagnose melanoma; suspicious lesions require clinical assessment.

## Abdomen

The body region between the diaphragm and pelvis. It contains most digestive organs, the kidneys, spleen, pancreas, and major blood vessels and is described clinically by four quadrants or nine regions.

## Abdominal

Relating to the abdomen, the body region between the chest and pelvis that contains organs of the digestive, urinary, and reproductive systems.

## Abdominal Abscess

A localized collection of pus within the abdomen, usually caused by bacterial infection, inflammation, surgery, or organ perforation. It may cause fever and abdominal pain and often requires antibiotics and drainage.

## Abdominal Adhesion

Alternate index label; see *Abdominal Adhesions*.

## Abdominal Adhesions

Bands of scar-like tissue that form inside the abdomen and make two or more organs stick to each other or to the abdominal wall. They most often develop after surgery but may also follow infection, inflammation, endometriosis, or radiation; some are present from birth. They often cause no symptoms, but may cause long-lasting abdominal pain, infertility when they affect pelvic reproductive organs, or a partial or complete intestinal blockage. A blockage may cause pain, bloating, constipation, vomiting, or inability to pass gas.

## Abdominal Angina

Pain in the abdomen after eating, caused by reduced blood flow through the arteries supplying the intestines. It is usually due to hardening and narrowing of these arteries.



---

## Abdominal Aorta

The portion of the aorta below the diaphragm that extends to its bifurcation into the common iliac arteries. It supplies the abdominal organs, pelvic organs, and lower limbs through its branches.

## Abdominal Aortic Aneurysm

A permanent widening of the abdominal aorta, usually defined as a diameter of 3 cm or more. Many cause no symptoms, but pain or rapid enlargement can signal danger. Rupture may cause sudden severe abdominal or back pain, dizziness, fainting, or collapse and requires emergency care.

### Alternate Names:

AAA; Abdominal Aneurysm

## Abdominal Aortic Aneurysm Repair

Surgical or endovascular treatment of an abdominal aortic aneurysm to prevent rupture or treat a rupture.

### Alternate Names:

AAA Repair

## Abdominal Aortic Aneurysm Screening

Ultrasound examination used to detect an abdominal aortic aneurysm before symptoms occur. Screening recommendations depend on age, sex, smoking history, and clinical risk; follow-up intervals depend on aneurysm size.

## Abdominal Aortic Ultrasound

An ultrasound examination of the abdominal aorta used to look for an aneurysm, measure its size, or monitor it over time. It is painless and does not use radiation.

## Abdominal Binder

A wide supportive garment placed around the abdomen to provide gentle compression after surgery, injury, or selected abdominal-wall conditions.

## Abdominal Bloating

A feeling of fullness, pressure, or tightness in the abdomen. It may occur with or without visible enlargement and is commonly caused by intestinal gas,

constipation, food intolerance, or functional gastrointestinal disorders. Persistent or severe bloating may require evaluation.

## Abdominal Bruit

An abnormal blowing or whooshing sound heard over the abdomen with a stethoscope, usually caused by turbulent blood flow through a narrowed or enlarged artery. It may occur with renal-artery stenosis or an abdominal aortic aneurysm. Further assessment depends on symptoms and vascular risk factors.

## Abdominal Cavity

The cavity between the diaphragm and pelvis containing the stomach, liver, gallbladder, spleen, pancreas, kidneys, and intestines.

## Abdominal Circumference

The distance around the abdomen. In pregnancy, fetal abdominal circumference is measured by ultrasound at a standardized level that includes the stomach, portal sinus, and spine; it is used with other measurements to estimate fetal size and assess growth.

## Abdominal Colic

Cramping abdominal pain caused by contractions or obstruction of a hollow organ, such as the intestine, ureter, or bile duct. Pain often comes in waves and may be accompanied by nausea, vomiting, sweating, or altered bowel or urinary function. Severe or persistent pain requires prompt medical assessment.

## Abdominal Compartment Syndrome

A dangerous rise in pressure inside the abdomen, usually above 20 millimeters of mercury, that causes new problems with organ function. Causes include severe fluid buildup, an overly swollen or blocked intestine, internal bleeding, and some conditions after abdominal surgery.

## Abdominal CT

Computed tomography of the abdomen used to evaluate selected causes of abdominal pain, trauma, bowel obstruction, inflammation, bleeding, and

---

masses. The protocol and use of contrast depend on the suspected condition and the patient's clinical status.

### **Abdominal CT Scan**

Alternate name; see *Abdominal CT*.

### **Abdominal Distension**

Visible or measured enlargement of the abdomen, often with a feeling of fullness or tightness. Causes include gas, constipation, fluid, pregnancy, enlarged organs, bowel obstruction, or bleeding; severe pain, vomiting, fever, or inability to pass stool or gas requires urgent assessment.

#### **Alternate Names:**

Abdominal Swelling; Swollen Abdomen; Abdominal Distention

### **Abdominal Distention**

Alternate name; see *Abdominal Distension*.

### **Abdominal Exploration**

Surgical inspection of the abdominal cavity to identify or treat a suspected source of pain, bleeding, infection, obstruction, or other disease.

### **Abdominal Fat**

Adipose tissue stored beneath the skin or around abdominal organs; excess visceral fat is metabolically active and associated with insulin resistance, fatty liver, and cardiovascular risk.

### **Abdominal Fibromatosis**

Alternate name for an abdominal desmoid tumor, a fibrous growth that can invade nearby tissues but does not spread to distant organs; see *Desmoid Tumor*.

### **Abdominal Fluid**

Fluid in the space around the abdominal organs. A small amount may be normal, but excess fluid, called ascites, can cause swelling, discomfort, breathing difficulty, or infection and needs evaluation for its cause.

#### **Alternate Names:**

Peritoneal Fluid

### **Abdominal Girth**

The distance around the abdomen, measured at a specified level. It may be used to monitor abdominal distention, fetal growth, obesity, ascites, or changes in intra-abdominal volume.

### **Abdominal Guarding**

Tensing of the abdominal muscles when the abdomen is touched. It may occur when the lining around the abdominal organs is irritated or inflamed.

### **Abdominal Hernia**

Protrusion of an organ or tissue through a weakness or opening in the abdominal wall. Common types include incisional, umbilical, epigastric, and ventral hernias.

### **Abdominal Hysterectomy**

Surgical removal of the uterus through an incision in the abdominal wall; the cervix, ovaries, and fallopian tubes may or may not also be removed.

### **Abdominal Mass**

A palpable or imaging-detected abnormal collection, enlargement, or lesion within the abdomen. Causes include distended bowel or bladder, organ enlargement, hernia, inflammatory collection, benign tumour, and malignancy; assessment depends on location, mobility, tenderness, systemic features, and appropriate imaging.

### **Abdominal Migraine**

A migraine-related disorder characterized mainly by recurrent episodes of moderate to severe midline abdominal pain, often with nausea or vomiting and little or no headache.

### **Abdominal MRI Scan**

A magnetic resonance imaging study that produces detailed images of the abdomen using magnetic fields and radiofrequency energy.

---

## Abdominal Muscles

The muscles forming the front and sides of the abdominal wall. They support the trunk, help with posture and breathing, and increase intra-abdominal pressure during coughing, lifting, and defecation.

### Alternate Names:

Abdominal Wall Muscles

## Abdominal Neoplasm

An abnormal growth in an organ or tissue of the abdomen. It may be noncancerous or cancerous and can cause pain, swelling, bleeding, changes in bowel function, or no symptoms.

### Alternate Names:

Abdominal Tumor

## Abdominal Neoplasms

Alternate index label; see *Abdominal Neoplasm*.

## Abdominal Pain

Pain or discomfort in the area between the chest and pelvis. It may be felt deep inside, come from the abdominal wall, or be felt in the abdomen even when the cause is elsewhere. Sudden severe pain, rigid muscles, fainting, persistent vomiting, fever, blood in stool, or pregnancy requires prompt medical assessment.

### Alternate Names:

Stomach Pain

## Abdominal Palpation

Part of a physical examination in which a clinician feels the abdomen with the hands to assess tenderness, muscle guarding, masses, or enlarged organs. Findings can guide further evaluation but do not identify a cause on their own.

### Alternate Names:

Palpation of the Abdomen

## Abdominal Point Tenderness

Pain felt in one specific area when the abdomen is gently pressed. It may occur with appendicitis, gallbladder inflammation, or irritation of the abdominal lining; it does not diagnose a condition by itself.

### Alternate Names:

Focal Abdominal Tenderness; Localized Peritoneal Tenderness; Point Pain in Abdomen

## Abdominal Pregnancy

An ectopic pregnancy implanted in the peritoneal cavity rather than the uterus, fallopian tubes, or ovaries. It is rare and can cause severe internal bleeding. Diagnosis requires specialist imaging and management.

## Abdominal Quadrant

One of four topographical anatomical divisions of the abdomen created by perpendicular vertical and horizontal imaginary planes intersecting at the umbilicus (right upper, left upper, right lower, and left lower quadrants), utilized in clinical triage to localize abdominal pain and organ pathology.

## Abdominal Radiography

A regular X-ray of the abdomen. It may be taken while a person is lying down, standing, or lying on their side.

## Abdominal Reflex

Contraction of the abdominal muscles after stimulation of the abdominal skin. An absent or asymmetric reflex may indicate neurologic disease, although it can be absent normally in some people.

## Abdominal Regions

The nine named areas used to describe locations on the abdominal wall: the right and left hypochondriac, epigastric, right and left lumbar, umbilical, right and left iliac, and hypogastric regions.

## Abdominal Retractor

A surgical instrument used to hold back the abdominal wall or organs to improve access and visibility during an operation.

### Alternate Names:

Balfour Retractor; Bookwalter Retractor

## Abdominal Rigidity

Involuntary tensing of the abdominal wall, usually

---

caused by irritation of the peritoneum. It may occur with peritonitis, perforation, severe infection, or intra-abdominal bleeding and requires clinical assessment.

### **Abdominal Sounds**

Sounds produced by movement of gas and fluid through the intestines; changes in frequency or character may support, but do not establish, a diagnosis.

### **Abdominal Swelling**

Alternate name; see *Abdominal Distension*.

### **Abdominal Tap**

Alternate name; see *Abdominocentesis*.

### **Abdominal Thrusts**

A first-aid maneuver for severe choking in a conscious adult or child older than 1 year. Give 5 back blows followed by 5 quick inward and upward pushes above the navel, repeating until the object comes out or the person becomes unconscious. Use chest thrusts for a pregnant or very large person; infants and unconscious people need different first aid.

### **Abdominal Ultrasound**

An ultrasound examination that uses sound waves to produce images of the abdomen and assess its structure and blood flow when applicable.

### **Abdominal Wall**

The layered muscular, fascial, and skin-covered boundary surrounding the abdominal cavity. It supports and protects abdominal organs and helps with posture, breathing, coughing, and straining. Weakness or defects may produce abdominal or incisional hernias; trauma, infection, and tumors may also affect the wall.

### **Abdominal Wall Fat Pad Biopsy**

Removal of a small sample of abdominal fat to look for amyloid, usually with a special Congo-red stain. It is less invasive than biopsy of an affected organ, but a negative result does not rule out amyloidosis.

### **Abdominal Wall Hernia**

Alternate name; see *Abdominal Hernia*.

### **Abdominal Wall Surgery**

Surgery to repair a defect or disease of the abdominal wall, often a hernia. It may be open or laparoscopic, and mesh may or may not be used depending on the defect, location, and patient factors.

### **Abdominal X-Ray**

Alternate name; see *Abdominal Radiography*.

### **Abdominocentesis**

Inserting a needle through the abdominal wall into the space around the abdominal organs to remove or test extra fluid, called ascites. It is also called paracentesis.

### **Abdominopelvic Cavity**

The connected body space below the diaphragm containing abdominal and pelvic organs, including digestive, urinary, and reproductive structures, protected by surrounding muscles, bones, and lining tissues.

### **Abdominoperineal Resection**

An operation that removes the anus, rectum, and usually part of the sigmoid colon. Because the normal anal outlet is removed, the surgeon creates a permanent colostomy, bringing the colon through the abdominal wall to drain stool into a bag. It may be used for selected low rectal or anal cancers.

#### **Alternate Names:**

APR

### **Abdominoplasty**

A surgical procedure that removes excess abdominal skin and may tighten the abdominal wall. It is commonly called a tummy tuck.

#### **Alternate Names:**

Tummy Tuck

### **Abducens Nerve**

The sixth cranial nerve. It controls the muscle that moves the eye outward. Damage to this nerve can

---

prevent outward eye movement and cause double vision.

### **Abducens Nerve Diseases**

Disorders affecting the sixth cranial nerve, which moves the eye outward. Injury or disease can weaken this movement and cause horizontal double vision; causes include diabetes-related nerve injury, raised pressure inside the skull, infection, inflammation, stroke, or trauma.

### **Abducens Nerve Palsy**

Dysfunction of the sixth cranial nerve resulting in paralysis or paresis of the lateral rectus muscle. Clinically characterized by horizontal diplopia that worsens on gaze toward the affected side and convergent strabismus (esotropia) in primary gaze. Common etiologies include microvascular ischemia (diabetes mellitus, hypertension), elevated intracranial pressure (false localizing sign), trauma, and cavernous sinus lesions.

#### **Alternate Names:**

Sixth Nerve Palsy; CN VI Palsy

### **Abducent Nerve**

Alternate name; see *Abducens Nerve*.

### **Abduct**

To move an arm, leg, finger, or other body part away from the center line of the body.

### **Abduction**

Movement of a limb or other body part away from the median plane of the body. In ocular motility, it is movement of an eye outward, away from the nose.

### **Abductor Muscle**

A muscle that moves a body part away from the midline or, for a digit, away from the axis of the hand or foot.

### **Abercrombie Syndrome**

An older name for systemic amyloidosis, in which abnormal amyloid proteins accumulate in organs

and impair their function.

### **Aberrant Conduction**

Transient abnormal intraventricular conduction of a supraventricular electrical impulse through the ventricular conduction system, typically occurring when an impulse reaches bundle branch tissue during its relative refractory period. Most commonly presents as a wide QRS complex with right bundle branch block morphology following a sudden change in cycle length (Ashman phenomenon).

#### **Alternate Names:**

Aberrancy; Ventricular Aberration

### **Aberration**

A deviation from normal structure, function, or appearance; in optics, it is an imperfection that prevents light from forming a sharply focused image.

### **Abetalipoproteinemia**

A rare inherited disorder in which the body cannot make normal particles that carry fat and cholesterol in the blood. It causes poor absorption of dietary fat, deficiencies of vitamins carried with fat, abnormal red blood cells, and possible nerve or eye disease.

### **Abfraction**

Loss of tooth material near the gum line, possibly related to repeated mechanical stress or biting forces. It may cause a notch and tooth sensitivity; a dentist can assess the cause and need for treatment.

### **Abiotrophy**

Progressive loss of function as cells in a tissue or organ gradually degenerate, often because of an inherited or developmental problem. It may cause worsening weakness, loss of sensation, poor coordination, or other problems depending on the affected part of the body.

### **Ablate**

To remove or destroy tissue using surgery, heat, cold, chemicals, or another energy source. Ablation may treat abnormal heart rhythms, tumors, pain pathways, or other selected conditions.



---

## **Ablation**

A procedure that removes or destroys tissue or stops an unwanted body function using surgery, heat, cold, electricity, medicines, or other methods.

## **Ablative Surgery**

Surgery that destroys or removes abnormal tissue, often using excision, heat, cold, electrical energy, or other tissue-destructive methods.

## **Ablepharon**

A rare congenital condition in which one or both eyelids are absent or severely underdeveloped. It can leave the eye exposed and may require protective treatment and reconstructive surgery.

## **Abnormal**

Different from the usual healthy structure, function, or appearance. An abnormal finding may be harmless or may indicate a disease, depending on its type and context.

## **Abnormal (Dysfunctional) Uterine Bleeding**

Bleeding from the uterus that is abnormal in timing, regularity, duration, or amount and is not explained by pregnancy. Causes include structural lesions, ovulatory or hormonal disorders, medicines, bleeding disorders, and endometrial disease; evaluation depends on age and risk factors.

## **Abnormal Color**

A change in the usual color of skin, moist tissues, body fluids, or another structure. Its meaning depends on the color, location, duration, and associated symptoms.

### **Alternate Names:**

Discoloration

## **Abnormal Fissure**

An atypical crack, groove, or cleft in a tissue or organ. It may be congenital or develop from injury, inflammation, scarring, pressure, or disease, and its importance depends on its location and depth.

## **Abnormal Hemoglobins Testing**

Blood testing that looks for unusual forms of hemoglobin, the oxygen-carrying protein in red blood cells. It helps diagnose inherited disorders such as sickle-cell disease and thalassemia; results may require specialized laboratory analysis.

## **Abnormal Labor**

Labor that is unusually slow, prolonged, arrested, or associated with abnormal contractions, fetal position, or fetal intolerance. Evaluation considers cervical change, contraction pattern, pelvic factors, fetal status, infection, and maternal condition before treatment or delivery decisions.

## **Abnormal Pap Test**

A cervical screening result showing abnormal cells or evidence of human papillomavirus. It does not by itself mean cancer; follow-up depends on the cell changes, HPV result, age, and prior screening history.

## **Abnormal Posturing**

An abnormal, sustained position of the body or limbs caused by severe dysfunction of the brain, brainstem, spinal cord, or muscles. Decorticate, decerebrate, and opisthotonic postures can signal a medical emergency.

## **Abnormal Reflex**

A reflex that is absent, reduced, unusually brisk, or different between sides. It can reflect a change in the nerves, spinal cord, brain, or muscle and must be interpreted with the rest of the examination.

## **Abnormal Tooth Shape**

Abnormal tooth shape may result from developmental defects, genetic conditions, trauma, wear, or tumors and can affect function, appearance, and eruption.

## **Abnormal Uterine Bleeding**

Bleeding from the uterus that differs from the usual pattern in amount, duration, frequency, or regularity in a person who is not pregnant. Causes include changes in ovulation or hormones, fibroids, polyps,

---

medicines, infection, and cancer. Bleeding after sex may come from the cervix or vagina rather than the uterus; bleeding during pregnancy is assessed separately.

### **Abnormal Vaginal Bleeding**

Alternate symptom label; see *Abnormal Uterine Bleeding*.

### **Abnormal Virilization**

Development of masculinizing physical features in a person whose expected sex development does not include them, often from excess androgens.

**Alternate Names:**

Virilization

### **Abnormality**

A difference from the usual structure, function, development, or test result. An abnormality may be harmless, temporary, or a sign of disease; its importance depends on the finding, symptoms, and clinical evaluation.

**Alternate Names:**

Anomaly

### **Abnormally Dark Or Light Skin**

Skin that has become darker or lighter than a person's usual skin tone. Darkening is called hyperpigmentation and lightening is called hypopigmentation. Causes include inflammation or injury, sun exposure, medicines, infections, hormonal disorders, and genetic conditions.

### **Abnormally Excessive Sweating**

See *Hyperhidrosis*.

### **ABO Blood Group System**

The major red-cell blood-group system based on whether A and/or B antigens are present on the red-cell surface and the corresponding antibodies are present in plasma. The four principal phenotypes are A, B, AB, and O; correct ABO matching is critical for safe transfusion.

**Alternate Names:**

ABO Blood Group ABO Blood Groups ABO Blood

Types ABO System

### **ABO Blood Typing**

A laboratory test that determines a person's ABO and Rh blood groups. It is performed before transfusion, transplantation, and some obstetric procedures.

### **ABO Incompatibility**

An immune reaction that can occur when blood with an incompatible A, B, or O type is transfused, or when maternal and fetal blood types are incompatible. It can destroy red blood cells, cause a severe transfusion reaction, or affect a newborn.

### **Abortion**

The ending of a pregnancy, either spontaneously or intentionally. An intentional abortion uses medicines or a medical procedure; spontaneous abortion is also called miscarriage.

### **Abortion Complications**

Potential complications after spontaneous or induced abortion, including hemorrhage, retained tissue, infection, uterine injury, cervical trauma, thromboembolism, and emotional or psychological distress.

### **Abortive**

Incomplete, unsuccessful, or stopped before full development; an abortive infection produces limited disease without the usual complete clinical course.

### **Abortive Polio**

A mild, nonparalytic poliovirus infection causing nonspecific fever, sore throat, headache, nausea, or malaise without the neurologic findings of paralytic polio.

### **Above The Knee Amputation**

Amputation of the lower limb through the thigh above the knee joint, usually performed for severe vascular disease, infection, trauma, or cancer.

---

## ABR Test

Auditory brainstem response testing records electrical activity from the auditory nerve and brainstem through scalp electrodes to assess hearing and auditory-pathway function, especially in infants or people unable to provide reliable behavioral responses.

## Abrade

To wear away tissue by friction or erosion; an abrasion is the resulting scrape or superficial loss of skin.

## Abrasion

A superficial injury in which the skin is scraped or rubbed away, usually by friction. In dentistry, abrasion means mechanical wearing away of tooth structure by an external force.

## Abrin

A highly poisonous protein found in rosary pea seeds. It stops cells from making proteins, causing them to die. Swallowing chewed seeds or exposure to abrin powder can cause vomiting, diarrhea, dehydration, breathing problems, organ failure, and death. Suspected exposure requires urgent medical care; there is no routine antidote.

## Abruptio Placenta

Alternate name; see *Abruptio Placentae*.

## Abruptio Placentae

Premature separation of the placenta from the uterine wall before delivery. It can cause vaginal bleeding, abdominal pain, uterine tenderness, fetal distress, and maternal shock and requires urgent obstetric assessment.

## Abs

An informal name for the muscles at the front and sides of the abdomen. They help bend and rotate the trunk, support the spine, protect abdominal organs, and increase pressure for coughing, lifting, breathing, or bowel movements.

## Abscess

A pocket of pus that forms in tissue, an organ, or a body cavity, usually because of an infection. It may cause pain, swelling, warmth, and fever; treatment often requires drainage and sometimes antibiotics.

## Abscessed Tooth

A collection of pus caused by bacterial infection of the tooth or surrounding tissues, usually arising from untreated dental decay, a cracked tooth, or periodontal disease. Pain, facial swelling, fever, and sensitivity may occur. Dental treatment and drainage are usually required; swelling that affects swallowing or breathing is an emergency.

## Abscission

The separation or removal of tissue, either naturally or by cutting. In surgery it may describe removing a growth, damaged tissue, or part of an organ; the exact meaning depends on the medical context.

## Absence Epilepsy

A generalized epilepsy with brief seizures that cause a sudden pause in activity and impaired awareness, often with staring or small repetitive movements. Awareness usually returns quickly without confusion; the seizures may occur many times a day.

### Alternate Names:

Childhood Absence Epilepsy; Pyknolepsy; Petit Mal Epilepsy

## Absence Of Sweating

Anhidrosis is diminished or absent sweating, which may be focal or generalized. It impairs thermoregulation and can result from autonomic dysfunction, peripheral neuropathy, skin disease, medications, or congenital conditions. Severe generalized anhidrosis can cause heat intolerance and exertional hyperthermia.

## Absence Of The Breast

Congenital absence of breast development, called amastia; it may affect one or both sides and can occur with other developmental anomalies.

---

## Absence Of The Nipple

Athelia: a rare congenital absence of one or both nipples, which may occur with hereditary disorders, chromosomal abnormalities, or developmental syndromes.

## Absence Seizure

A brief seizure causing sudden staring and reduced awareness, usually lasting only seconds with rapid recovery. The person may stop speaking or responding and may make small repetitive movements; an EEG helps confirm the diagnosis.

### Alternate Names:

Petit Mal Seizure; Typical Absence Seizure

## Absence Seizure Cross-Reference

Alternate name; see *Absence seizure*.

## Absent Breath Sounds

Lack of audible air movement over part or all of a lung. Causes include pneumothorax, pleural effusion, or airway obstruction; sudden onset requires urgent assessment.

## Absent Eye

Anophthalmia, a congenital absence or severe underdevelopment of the eye globe and ocular tissue; it may affect one or both eyes and may occur with other birth defects.

## Absent Menstrual Periods

Alternate index label; see *Amenorrhea*.

## Absent Pulmonary Valve

A birth defect in which the valve between the heart and the lung arteries is missing or severely underdeveloped. It can enlarge those arteries and cause breathing problems, poor blood flow to the lungs, or other heart defects.

## Absent Reflex

Failure of an expected automatic movement to occur when a tendon or other body area is stimulated. It may suggest a problem in a nerve, nerve root, spinal cord, or muscle, although reflexes can normally differ

between people and sides.

## Absent Thirst

Reduced or absent desire to drink despite a potential need for fluids, which can occur with hypothalamic disease, aging, altered consciousness, medications, or severe illness.

## Absolute Eosinophil Count

A laboratory blood test measuring the exact number of eosinophils per microliter of blood to evaluate allergic disorders, asthma, parasitic infections, drug reactions, or hypereosinophilic syndromes.

### Alternate Names:

AEC; Absolute Eosinophils; Blood Eosinophil Count

## Absolute Neutrophil Count

The number of neutrophils, a type of white blood cell that fights many bacterial and fungal infections, in a volume of blood. A low count increases infection risk, especially when it is severely reduced.

### Alternate Names:

ANC

## Absorb

To take in a substance, such as through the skin or intestine, or to reduce the intensity of radiation or light by taking up its energy.

## Absorption

The process by which a drug enters the bloodstream from its site of administration, such as the digestive tract, skin, or muscle.

## Abstinence

Deliberate avoidance of a substance or behavior, such as alcohol, drugs, tobacco, or sexual activity. Stopping some substances, especially alcohol, sedatives, or opioids, can cause dangerous withdrawal and may require medical supervision.

## Abulia

Marked reduction in motivation, initiative, and spontaneous activity, usually caused by dysfunction of frontal-subcortical brain circuits or related

---

neurologic disease.

### **Abusive Head Trauma**

Injury to an infant or young child caused by inflicted blunt impact, violent shaking, or both. Findings may include subdural hemorrhage, retinal hemorrhage, encephalopathy, fractures, or other injuries; diagnosis requires careful multidisciplinary assessment because no single finding is diagnostic.

#### **Alternate Names:**

Shaken Baby Syndrome; Non-Accidental Head Injury

### **Abutment**

A natural tooth, dental implant, or component that supports a dental prosthesis. Abutments may hold a bridge, denture, or implant crown in position and transfer chewing forces to the supporting tooth, implant, or bone.

### **AC Joint**

The acromioclavicular joint, a gliding joint between the acromion of the scapula and the clavicle at the top of the shoulder.

### **Acalculia**

Loss of the ability to do or understand numbers and calculations that a person could previously manage. It may result from injury or disease affecting areas of the brain involved in calculation.

### **Acalculous Cholecystitis**

Inflammation of the gallbladder without gallstones. It usually occurs suddenly in people who are critically ill, injured, recovering from surgery, unable to eat, or in shock. Reduced bile flow and poor blood supply can contribute, and severe cases may cause tissue death or a hole in the gallbladder. Fever and pain in the upper-right abdomen may occur; ultrasound is commonly used for evaluation.

### **Acanthamoeba**

A free-living amoeba found in soil and water that can cause corneal infection, skin or sinus infection, or rarely granulomatous amoebic encephalitis. Contact-lens exposure to contaminated water in-

creases keratitis risk, while invasive disease is more likely with impaired immunity.

### **Acanthamoeba Keratitis**

A rare but potentially sight-threatening infection of the cornea caused by the free-living amoeba *Acanthamoeba*. It is strongly associated with contaminated contact-lens water or poor lens hygiene and may cause severe eye pain, redness, photophobia, blurred vision, and a ring-shaped corneal infiltrate. Prompt ophthalmic assessment and prolonged anti-amoebic treatment are required.

### **Acanthocyte**

An abnormal red blood cell with irregular, thorny or spike-like projections of varying length and width on its outer membrane. They are seen on a blood smear in advanced liver disease (such as cirrhosis), lipid metabolism disorders, abetalipoproteinemia, or neuroacanthocytosis.

#### **Alternate Names:**

Spur Cell

### **Acanthocytosis**

An abnormal hematologic condition characterized by the presence of circulating red blood cells with irregular, thorny or spiked projections (acanthocytes or spur cells). It is classically observed in abetalipoproteinemia, advanced liver cirrhosis (spur cell anemia), severe malnutrition, and neuroacanthocytosis syndromes.

#### **Alternate Names:**

Spur Cell Formation

### **Acantholysis**

Loss of adhesion between keratinocytes, the main cells of the epidermis (outer skin layer). The cells separate, sometimes forming blisters or surface erosions. Acantholysis is a microscopic finding seen in disorders such as pemphigus and some inherited or infectious skin diseases; it is not a disease by itself.

### **Acanthoma**

A usually benign skin growth made of thickened squamous epidermal cells. Its appearance can resemble other lesions, so a changing or uncertain



---

lesion may need examination or biopsy.

### **Acanthosis Nigricans**

Dark, thickened, velvety skin, usually in folds such as the neck, armpits, or groin. It is often associated with insulin resistance, obesity, type 2 diabetes, hormonal disorders, or medicines; rarely, sudden extensive disease can signal an underlying cancer. Treating the cause may improve the skin changes.

### **Acapnia**

An unusually low carbon-dioxide level in arterial blood, usually caused by breathing too quickly or deeply. It may cause light-headedness, tingling, or a change in the blood's acid level.

### **Acardia**

Absence or severe malformation of the fetal heart, usually occurring in the setting of a rare twin perfusion sequence. The affected fetus cannot sustain independent circulation.

### **Acariasis**

Illness or skin infestation caused by mites or ticks. It may cause itching, a rash, burrows, skin nodules, or deeper tissue inflammation; some mites and ticks can also transmit infection.

### **Acaricide**

A pesticide used to kill mites and related arachnids such as ticks. It may be used on animals, plants, or environments; exposure can irritate skin and may cause poisoning if misused.

### **Acatalasemia**

A rare inherited condition in which catalase, an enzyme that breaks down hydrogen peroxide, is absent or greatly reduced. Many people have no symptoms, but some develop mouth ulcers, tissue damage, or increased sensitivity to oxidative stress.

#### **Alternate Names:**

Acatalasia; Catalase Deficiency

### **Acatalasia**

Alternate name; see *Acatalasemia*.

### **ACC – Adenoid Cystic Carcinoma**

Abbreviation for adenoid cystic carcinoma, a malignant tumor of glandular tissue that may grow slowly but can invade nerves and recur or spread years after treatment.

### **ACC – Agenesis of Corpus Callosum**

Abbreviation for agenesis of the corpus callosum, a congenital absence or incomplete development of the major nerve-fiber bridge connecting the brain's two hemispheres.

### **ACC – Aplasia Cutis Congenita**

Abbreviation for aplasia cutis congenita, a birth defect in which an area of skin, most often on the scalp, is absent or incompletely formed.

### **Accelerated Growth**

An increase in a child's growth velocity above the expected rate for age and developmental stage, determined from serial height or length measurements. It may occur during normal puberty or catch-up growth; abnormal acceleration, especially with early pubertal signs or advanced bone age, may suggest precocious puberty or another endocrine disorder.

### **Accelerated Idioventricular Rhythm**

A rhythm of wide electrical beats arising in the heart's lower chambers, often seen as blood flow returns after a heart attack. It is usually self-limited, but symptoms, the underlying cause, and other heart problems affect its significance.

#### **Alternate Names:**

AIVR

### **Accelerated-Phase Chronic Myeloid Leukemia**

A term used in some classification systems for chronic myeloid leukemia showing features of progression. Criteria vary; the International Consensus Classification includes 10% to 19% blasts, while the 2022 World Health Organization classification no longer recognizes a separate accelerated phase.

---

## Accelerometry

Measurement of movement or acceleration using a sensor. It can assess physical activity, walking, tremor, falls, sleep, or rehabilitation progress, depending on the device and the recording method.

## Acceptable Daily Intake

The estimated amount of a substance that a person can consume every day throughout life without a meaningful health risk, based on available evidence and safety margins. It is usually expressed per kilogram of body weight.

### Alternate Names:

ADI

## Acceptance-Based Therapies

Psychotherapeutic approaches that help a person notice and accept thoughts, feelings, or bodily sensations while reducing unhelpful attempts to control or avoid them.

### Alternate Names:

Acceptance-Based Psychotherapy; Acceptance-Focused Therapy

## Access Needle

A hollow needle used to enter a blood vessel, body cavity, organ, or fluid collection. It may be used to take a sample, drain fluid, give treatment, or help place a catheter.

### Alternate Names:

Introducer Needle; Percutaneous Access Needle; Vascular Access Needle

## Accessory

Additional or supplementary; in anatomy, an accessory structure is an extra part or a structure that assists the usual organ or system.

## Accessory Digestive Organ

An organ that supports digestion but is not part of the alimentary canal, such as the liver, pancreas, gallbladder, or salivary glands.

## Accessory Dwelling Unit

A secondary, independent residential unit on the

same property as a primary home; it is a housing and community-health term rather than a disease or body structure.

## Accessory Muscles

Muscles recruited to assist the primary muscles of respiration or another movement when the normal workload is increased; use of respiratory accessory muscles may indicate respiratory distress.

## Accessory Nerve

The eleventh cranial nerve (CN XI), a primarily motor nerve whose spinal component supplies the sternocleidomastoid and trapezius muscles. It enables head rotation and shoulder elevation; injury may cause weak head rotation, impaired shoulder shrug, and shoulder droop.

### Alternate Names:

Cranial Nerve XI; Spinal Accessory Nerve

## Accessory Pathway

An abnormal electrical connection between the atria and ventricles that can bypass the atrioventricular node and cause pre-excitation or tachyarrhythmias, including atrioventricular reentrant tachycardia.

## Accessory Placenta

An extra placental lobe separate from the main placenta but connected to it by fetal blood vessels; it can increase the risk of retained placental tissue or vessel injury during delivery.

## Accident And Emergency Medicine

Accident and emergency medicine is the specialty concerned with immediate evaluation and treatment of acute illness, injury, poisoning, and life-threatening emergencies.

## Accidental Death

Death caused by an unintentional injury or event rather than disease, suicide, or homicide. The cause is determined through medical, scene, and sometimes legal investigation.

### Alternate Names:

Unintentional Death

---

## Accidental Fall

An unintentional movement to the ground or a lower level. Falls can cause bruises, fractures, head injury, or internal bleeding, especially in older adults; loss of consciousness, severe pain, or confusion requires urgent care.

### Alternate Names:

Fall Injury

## Accommodation

The ability of the eye to change the shape of its lens so that near objects can be focused clearly.

## Accommodation Reflex

The coordinated adjustment of the lens, ciliary muscle, and pupils that shifts the eye from distance to near focus; it includes lens thickening, convergence, and pupillary constriction.

## Accommodative Insufficiency

Inability of the eye to increase lens power adequately for near focusing, causing eyestrain, blurred near vision, headache, or difficulty reading.

## Accouchement

Accouchement is the process of childbirth, including labor, delivery of the infant, and expulsion of the placenta.

## Accretion

Gradual buildup or growth of material that makes an object or body tissue larger. Examples include crystals forming in fluid, extra bone developing after injury, or plaque collecting on teeth.

## ACE Blood Test

A blood test measuring angiotensin-converting enzyme. It may support evaluation of sarcoidosis and some other inflammatory or liver disorders, but an abnormal result is not specific and cannot diagnose a disease by itself.

## ACE Inhibitors

Medicines that reduce formation of angiotensin II, lowering blood pressure and pressure within the kid-

ney's filtering units. They are used for hypertension, heart failure, and selected kidney or cardiovascular conditions; cough, high potassium, kidney-function changes, and rare facial or airway swelling are important risks, and they are generally avoided in pregnancy.

## Acebutolol

Acebutolol is a beta-adrenergic blocker with intrinsic sympathomimetic activity used in some settings for hypertension and cardiac rhythm disorders; it can slow heart rate and conduction.

## Acellular

Without cells. An acellular material may consist of fluid, minerals, or the supporting material around cells and may be natural, diseased, or used in a medical product.

## Acentric Chromosome

A chromosome fragment that lacks a centromere and therefore usually cannot attach normally to the spindle during cell division.

## Acephalostomia

An extremely rare and severe birth defect involving absence or major failure of development of the head and mouth. It is usually incompatible with survival and is identified during prenatal or newborn examination.

## Acetabular Labrum

A fibrocartilaginous rim attached to the margin of the acetabulum that deepens the hip socket, helps maintain a fluid seal, and contributes to hip-joint stability.

## Acetabular Notch

The indentation in the inferior margin of the acetabulum, bridged by the transverse ligament of the acetabulum and forming a passage for structures entering the hip joint.

## Acetabuloplasty

A surgical procedure that reshapes or augments the

---

acetabular rim or socket to improve coverage of the femoral head in selected structural hip disorders.

### Acetabulum

The cup-shaped socket of the hip formed by the fused ilium, ischium, and pubis. It articulates with the femoral head and is deepened by the acetabular labrum.

### Acetaldehyde

A chemical produced when the body metabolizes alcohol. Accumulation may contribute to flushing, nausea, and vomiting.

### Acetaminophen

A widely used medication that relieves mild-to-moderate pain and reduces fever, but lacks significant anti-inflammatory effects. While safe at recommended therapeutic doses, excessive amounts can cause severe, life-threatening liver injury, especially when taken with alcohol or other products containing acetaminophen.

#### Alternate Names:

Paracetamol; APAP

### Acetaminophen And Codeine Overdose

A medical emergency caused by ingesting toxic amounts of a combination pain medicine. It produces dual life-threatening effects: codeine suppresses the central nervous system, causing extreme drowsiness, slowed or stopped breathing, and coma, while acetaminophen depletes protective liver antioxidants, leading to progressive liver failure.

#### Alternate Names:

Co-Codamol Overdose; Tylenol with Codeine Overdose

### Acetaminophen Overdose

Taking more acetaminophen than the body can safely process, causing serious liver damage. A person may have few symptoms at first, followed later by nausea, right-sided abdominal pain, jaundice, or liver failure. Get emergency help or contact a poison center immediately; the antidote N-acetylcysteine works best when started early but may still help when given later.

#### Alternate Names:

Acetaminophen Toxicity; Acetaminophen Poisoning; Paracetamol Toxicity; Paracetamol Overdose; APAP Overdose

### Acetaminophen Poisoning

Alternate name; see *Acetaminophen Overdose*.

### Acetaminophen Toxicity

Alternate name; see *Acetaminophen Overdose*.

### Acetate

A form of acetic acid and a small molecule used by cells in energy and building processes. The body can convert acetate into acetyl-CoA, which helps make or break down fats and other substances.

### Acetazolamide

A medicine that changes salt and fluid handling in the kidneys. It is used for glaucoma, prevention or treatment of altitude illness, and selected fluid, metabolic, or neurologic conditions.

### Acetone

A volatile ketone body produced during increased fat metabolism. It may contribute to the characteristic breath odor of ketosis and ketoacidosis.

### Acetone Body

A ketone produced when the body breaks down fat for energy. Acetone is one of three main ketones and can contribute to the distinctive breath odor of prolonged fasting or diabetic ketoacidosis.

### Acetone Breath

A distinctive sweet, fruity odor on the breath caused by the exhalation of volatile acetone, a ketone body produced during accelerated lipolysis and ketogenesis. It is a characteristic physical examination finding in diabetic ketoacidosis (DKA), as well as in severe starvation, prolonged fasting, and ketogenic metabolic states.

#### Alternate Names:

Fruity Breath; Ketotic Breath; Diabetic Breath Odor

---

## Acetone Poisoning

Harm from swallowing or breathing a large amount of acetone, a solvent found in some products. It may cause irritation, dizziness, vomiting, drowsiness, low blood pressure, or breathing problems; significant exposure is a medical emergency requiring prompt treatment.

## Acetonuria

The presence of acetone or other ketone bodies in the urine, which may occur with fasting, vomiting, uncontrolled diabetes, or other states of increased fat metabolism.

## Acetyl-CoA Carboxylase

An enzyme that begins the process of making fatty acids by changing acetyl-CoA into malonyl-CoA. Its activity responds to the cell's energy supply and to hormones, helping balance fat production with energy needs.

## Acetyl-Coenzyme A

A substance used in many energy and building processes inside cells. It helps release energy from food and provides material for making fatty acids, cholesterol, and ketones.

## Acetylation

A chemical change that adds a small acetyl group to a protein, medicine, or DNA-related molecule. This can alter how the molecule works, how long it lasts, how it is moved, or how strongly a gene is active.

## Acetylcholine

A major neurotransmitter synthesized from choline and acetyl-CoA by choline acetyltransferase. Operating throughout the central and peripheral nervous systems, it mediates neuromuscular transmission at motor endplates, autonomic ganglionic neurotransmission, parasympathetic postganglionic signaling, and central cognitive/memory processing.

### Alternate Names:

ACh

## Acetylcholine Receptor Antibody

Acetylcholine receptor antibody testing detects antibodies associated with autoimmune myasthenia gravis and is interpreted with the neurologic examination and other studies.

## Acetylcholinesterase

A rapid-acting serine hydrolase enzyme localized at synaptic clefts and neuromuscular junctions that hydrolyzes the neurotransmitter acetylcholine into choline and acetate, terminating signal transmission and allowing tissue repolarization.

### Alternate Names:

AChE; Acetylcholine Acetylhydrolase

## Acetylcholinesterase Inhibitors

A group of medicines that slow the breakdown of acetylcholine, a chemical messenger used by nerves. This increases acetylcholine activity and can help manage conditions such as Alzheimer disease and myasthenia gravis or reverse some medicine-related muscle paralysis.

## Acetylcysteine

A medicine that loosens thick mucus so it is easier to clear. It is also the main antidote for acetaminophen overdose because it restores glutathione, a substance the liver uses to neutralize the overdose's toxic breakdown product.

## Acetylsalicylic Acid

Alternate index label; see *Aspirin*.

## ACH

Abbreviation commonly used for achondroplasia, a genetic skeletal disorder causing disproportionate short stature and characteristic changes in the skull, spine, and long bones.

## ACh (Acetylcholine)

Alternate name; see *Acetylcholine*.

## Achalasia

A disorder in which the muscle ring between the esophagus and stomach does not relax properly,



---

and the esophagus cannot move food normally into the stomach. It causes difficulty swallowing and regurgitation.

**Alternate Names:**

Cardiospasm; Esophageal Achalasia

### **Achalasia Of The Cardia**

A disorder in which the valve between the esophagus and stomach does not relax properly and the esophagus loses its normal squeezing movement. Food and liquid can collect in the esophagus, causing progressive difficulty swallowing, regurgitation, chest discomfort, and weight loss.

### **AChE (Acetylcholinesterase)**

Alternate name; see *Acetylcholinesterase*.

### **Achenbach's Syndrome**

A benign condition of recurrent, painful, spontaneous hematoma formation in the pulp of a finger or thumb, typically affecting middle-aged women. It presents with sudden onset of throbbing pain and blue-black discoloration of the fingertip, resolving spontaneously over one to two weeks without treatment.

### **Achilles Tendinitis**

Painful irritation or degeneration of the tendon joining the calf muscles to the heel. It usually follows repeated loading and can cause stiffness, swelling, tenderness, and pain during walking or exercise.

**Alternate Names:**

Achilles Tendinopathy; Calcaneal Tendinitis

### **Achilles Tendinopathy**

Pain and structural disorder of the Achilles tendon, usually caused by repeated loading. It causes posterior heel pain and stiffness and is managed with activity modification, rehabilitation, and treatment of contributing factors.

### **Achilles Tendon**

The thickest and strongest tendon in the human body, connecting the calf muscles (gastrocnemius and soleus) to the heel bone (calcaneus). It trans-

mits muscle force to the foot, enabling the ankle to point downward (plantarflexion) during walking, running, jumping, and standing on tiptoe.

**Alternate Names:**

Calcaneal Tendon; Tendo Calcaneus

### **Achilles Tendon Problems**

Disorders affecting the Achilles tendon, including tendinitis, tendinopathy, bursitis, partial tear, and rupture. Pain, stiffness, swelling, or a sudden loss of push-off strength warrants assessment.

### **Achilles Tendon Repair**

Surgery that reconnects a completely or severely torn Achilles tendon. The ankle is protected in a cast or brace afterward, followed by gradual exercises to restore movement and strength; recovery takes several months.

### **Achilles Tendon Rupture**

A partial or complete tear of the Achilles tendon, usually caused by sudden force. It causes posterior ankle pain, weakness in plantar flexion, and difficulty walking or standing on tiptoe.

### **Achilles Tendonitis**

A commonly used term for pain and inflammation or degeneration of the Achilles tendon. It may affect the tendon at its heel insertion or its mid-portion and is often related to repetitive loading.

### **Achillobursitis**

Inflammation of a bursa near the Achilles tendon, causing pain, tenderness, swelling, or friction-related discomfort at the back of the heel.

### **Achlorhydria**

A condition in which the stomach produces no hydrochloric acid; reduced but not absent acid is called hypochlorhydria. It can occur with autoimmune gastritis or loss of acid-producing stomach cells and may impair absorption of iron or vitamin B12. Acid-suppressing medicines usually reduce acid rather than cause complete achlorhydria.

---

## Achondrogenesis

A group of severe inherited disorders in which cartilage and bone development is profoundly impaired. Affected fetuses have very short limbs and poorly formed bones, and most cases are fatal before or shortly after birth.

## Achondroplasia

An autosomal dominant bone-growth disorder caused by a disease-causing change in the *FGFR3* gene. It causes short arms and legs, short stature, a larger head, and a prominent forehead. One altered gene copy usually causes the condition; two copies cause a severe, often fatal form.

### Alternate Names:

Chondrodystrophy; Achondroplastic Dwarfism

## Achoo Syndrome

The photic sneeze reflex: repeated sneezing triggered by sudden exposure to bright light, especially after entering sunlight from darkness.

## Achromatopsia

An inherited disorder causing absent or greatly reduced color vision, often with light sensitivity and poor visual acuity.

### Alternate Names:

Complete Color Blindness

## Achromia

Achromia is loss or marked reduction of pigment in skin, hair, eye, or another tissue; causes include genetic disorders, autoimmune depigmentation, injury, and chemical exposure.

## Aciclovir

Aciclovir, also called acyclovir, is an antiviral drug that inhibits herpesvirus DNA polymerase and is used for infections such as herpes simplex and varicella-zoster.

## Acid

A substance that increases acidity by releasing hydrogen ions or accepting electrons. Strong acids can damage tissue, while acids also take part in normal

digestion, metabolism, and body-fluid regulation.

## Acid Base Balance

The body's control of acids and bases in its fluids, especially the blood. The lungs remove carbon dioxide and the kidneys adjust acids and bicarbonate to keep blood pH within a narrow safe range.

## Acid Indigestion

Dyspepsia or upper-abdominal discomfort that may include burning, fullness, belching, or nausea; it can accompany reflux, gastritis, ulcer disease, or other conditions.

## Acid Loading Test

A specialized test of how well the kidneys remove acid from the body. After a carefully controlled acid dose, urine is measured for acidity and other substances to investigate suspected kidney-tubule problems.

## Acid Mucopolysaccharides

An older name for long sugar chains found in supporting tissues such as cartilage, skin, and joint fluid. Abnormal breakdown or storage of these substances causes a group of inherited storage disorders.

## Acid Phosphatase

An enzyme that works best in acidic conditions and is found in several tissues, including prostate and blood cells.

### Alternate Names:

ACP

## Acid Reflux

Backward flow of acidic stomach contents into the esophagus, causing heartburn, sour regurgitation, cough, or throat irritation. Frequent reflux may be gastroesophageal reflux disease and can injure the esophagus.

## Acid Soldering Flux Poisoning

Injury from swallowing or inhaling acidic chemicals used in soldering. It may burn the mouth, throat, skin, or eyes and can cause vomiting, breathing

---

difficulty, or dangerous changes in body chemistry; significant exposure requires urgent medical treatment.

### Acid Sphingomyelinase Deficiency

An inherited lysosomal storage disorder caused by deficient acid sphingomyelinase, usually because of disease-causing variants in the *SMPD1* gene. Sphingomyelin accumulates in cells and may cause enlarged liver or spleen, lung disease, bone changes, or neurologic impairment; it is also called Niemann–Pick disease types A and B.

### Acid-Base Analysis

Evaluation of blood pH, bicarbonate, and carbon dioxide to identify metabolic or respiratory disturbances, including those caused by poisoning.

### Acid-Base Disorder

A problem that makes the blood too acidic or too alkaline. It may result from changes in breathing, kidney function, metabolism, or severe illness. Blood tests help identify the cause and whether the body is compensating.

### Acid-Base Imbalance

Alternate name; see *Acid-Base Disorder*.

### Acid-Fast Bacilli

Rod-shaped bacteria that resist decolorization by acid-alcohol solutions during laboratory staining procedures due to the high lipid and mycolic acid content of their cell walls. The group classically includes *Mycobacterium tuberculosis* and *Mycobacterium leprae*.

#### Alternate Names:

AFB; Acid-Fast Bacteria

### Acid-Fast Bacillus Smear

Microscopic examination of a specimen, often sputum, for acid-fast organisms using a special stain. A positive result supports mycobacterial infection but does not by itself identify the species or establish drug susceptibility; culture or molecular testing is also required.

### Acid-Fast Stain

A special dye test used to find mycobacteria and a few related organisms in samples such as sputum. A positive result does not identify the species or show which medicines will work.

### Acidemia

An abnormally low arterial blood pH (below 7.35), indicating an increased hydrogen ion concentration in circulating blood. It is the measurable laboratory state resulting from an underlying respiratory or metabolic acidosis.

### Acidophil

A cell or tissue part that absorbs acidic laboratory dyes strongly, especially the pink dye eosin. The pattern helps pathologists recognize certain cells in tissue samples, including some pituitary cells.

### Acidophilus

*Lactobacillus acidophilus*, a lactic-acid-producing bacterium found in the intestinal and genital microbiota and used in some fermented foods and probiotic products.

### Acidosis

A process that raises acid or removes base in the body, tending to lower blood pH. Acidemia means the measured blood pH is actually low; compensation or mixed disorders can affect the final pH. Causes include breathing problems, kidney disease, severe infection, diabetes, and poisoning.

### Aciduria

Aciduria means an abnormally acidic urine or the presence of excess organic acids in urine; it may reflect diet, dehydration, medication, or an inherited metabolic disorder.

### Acinar Cell

A cell in a small cluster that makes and releases a fluid or substance. In the pancreas, acinar cells release digestive enzymes into small ducts.

---

## Acinar Cell Carcinoma

A rare cancer of cells that resemble enzyme-secreting acinar cells, most often arising in the pancreas. It can form a mass and cause nonspecific abdominal symptoms.

## Acinetobacter Baumannii Resistance

Resistance means that *Acinetobacter baumannii* is not killed by one or more antibiotics. It can cause difficult hospital infections, especially in intensive-care units, so culture and susceptibility testing guide treatment.

## Acinus

A small working unit of a gland made of cells that release their product into a duct. Examples include the secretory units of the pancreas and salivary glands.

## ACL Injury

A sprain or tear of the strong ligament inside the knee that helps prevent the shinbone from sliding too far forward. It commonly follows twisting or sudden stopping and may cause a pop, swelling, pain, or a feeling that the knee gives way.

## ACL Reconstruction

Surgery that replaces a torn anterior cruciate ligament inside the knee with a tendon graft. It may restore stability after injury, especially when the knee gives way or the person plans activities requiring pivoting; rehabilitation is required afterward.

## Acne

A common inflammatory disorder of hair follicles and oil glands. Blocked follicles form blackheads or whiteheads, while inflammation can cause papules, pustules, nodules, scarring, and dark marks, especially on the face, chest, and back.

### Alternate Names:

Acne Vulgaris; Pimples

## Acne Conglobata

A rare, severe nodulocystic form of acne with groups of inflammatory nodules, cysts, interconnected ab-

scesses, and draining sinuses, usually on the trunk, buttocks, or limbs. It can be persistent and leave atrophic or hypertrophic scars, so early dermatologic treatment is important.

## Acne Fulminans

A sudden, severe ulcerating form of acne associated with fever, painful joints, malaise, and sometimes weight loss or abnormal blood counts. It requires urgent specialist care; isotretinoin can precipitate or worsen it if introduced too quickly.

## Acne Inversa

See *Hidradenitis suppurativa*.

## Acne Keloid

A chronic inflammatory condition causing firm scar-like bumps around hair follicles, usually on the back of the neck.

### Alternate Names:

Acne Keloidalis Nuchae

## Acne Keloidalis Nuchae (AKN)

Alternate name; see *Acne Keloid*.

## Acne Rosacea

Acne rosacea is a chronic inflammatory facial skin disorder causing flushing, persistent redness, visible vessels, papules, pustules, or eye symptoms; it is distinct from acne vulgaris.

## Acne Vulgaris

A common disorder of hair follicles involving follicular plugging, increased oil production, and inflammation in which *Cutibacterium acnes* may contribute. It causes blackheads or whiteheads and may also cause inflamed papules, pustules, or deep nodules on the face, chest, back, and upper arms; deeper disease can scar.

## Acneiform Eruptions

Monomorphic follicular papules and pustules that resemble acne but usually lack comedones. They may be caused by medicines, occupational or chemical exposures, infections, or other disorders; timing,

---

distribution, and trigger help distinguish them from acne vulgaris.

### **Acoustic**

Relating to sound, hearing, or how sound waves are produced, transmitted, measured, or received. Acoustic testing can assess hearing, rooms, medical devices, tissues, or signals used during diagnosis.

### **Acoustic Nerve**

The acoustic nerve, properly called the vestibulocochlear nerve, carries signals for hearing and balance from the inner ear to the brain. It is the eighth cranial nerve.

### **Acoustic Neuroma**

A benign, slow-growing neoplasm arising from Schwann cells of the vestibular division of the vestibulocochlear nerve (cranial nerve VIII) within the internal auditory canal or cerebellopontine angle. It classically presents with insidious, progressive unilateral sensorineural hearing loss, asymmetric high-pitched tinnitus, and dysequilibrium.

#### **Alternate Names:**

Vestibular Schwannoma; Acoustic Neurinoma; Acoustic Neurilemmoma

### **Acoustic Reflex**

An automatic tightening of a small middle-ear muscle in response to a loud sound. Testing it helps assess the middle ear, hearing pathways, and facial nerve.

#### **Alternate Names:**

Stapedial Reflex; Stapedius Reflex; Middle-Ear Muscle Reflex

### **Acoustic Shock**

Auditory and other symptoms after exposure to a sudden or intense sound, potentially including ear pain, tinnitus, hyperacusis, dizziness, or temporary hearing change.

### **Acoustic Trauma**

Acute damage to the auditory system from a brief, intense sound such as an explosion or firearm. It

may cause tinnitus, temporary or permanent sensorineural hearing loss, and tympanic or ossicular injury; urgent audiologic assessment is indicated when hearing loss is sudden.

### **Acquired**

Developed after birth rather than being present at birth or inherited; an acquired condition may result from infection, exposure, injury, aging, or another later event.

### **Acquired C1-Inhibitor Deficiency Angioedema**

A bradykinin-mediated disorder caused by acquired deficiency or dysfunction of C1 inhibitor, often associated with lymphoproliferative or autoimmune disease. Recurrent swelling of the skin, bowel wall, or upper airway usually occurs without hives or itching; low C1q and C1-inhibitor testing help distinguish it from hereditary or histamine-mediated angioedema.

### **Acquired Cystic Kidney Disease**

Development of multiple fluid-filled cysts in kidneys with long-standing chronic kidney disease, especially in people receiving dialysis. It can cause blood in the urine, pain, infection, or rarely kidney tumors and is monitored with clinical assessment and imaging.

### **Acquired Feminization**

Development of typically female secondary sexual characteristics after birth in a person not expected to develop them, often from hormonal change.

### **Acquired Immunity**

Immune protection that develops after infection, vaccination, or transfer of antibodies, involving adaptive immune responses directed toward specific antigens.

### **Acquired Immunodeficiency Syndrome**

See *HIV/AIDS*.

### **Acquired Lipodystrophy**

A rare disorder characterized by selective, progres-

---

sive loss of adipose tissue (lipoatrophy) occurring after birth without an inherited genetic defect. Frequently associated with autoimmune diseases, pancreatitis, or antiretroviral therapy (e.g., Lawrence syndrome, Barraquer-Simons syndrome), it leads to ectopic lipid accumulation, severe insulin resistance, hypertriglyceridemia, and hepatic steatosis.

**Alternate Names:**

Acquired Generalized Lipodystrophy; AGL; Lawrence Syndrome; Acquired Partial Lipodystrophy

### Acquired Methemoglobinemia

An acquired hematologic disorder in which hemoglobin iron is oxidized from the ferrous ( $\text{Fe}^{2+}$ ) to the ferric ( $\text{Fe}^{3+}$ ) state by exogenous oxidizing agents (such as dapsone, benzocaine, lidocaine, nitrites, or sulfonamides), producing clinical cyanosis, chocolate-brown arterial blood, and cellular hypoxia refractory to supplemental oxygen.

**Alternate Names:**

Toxic Methemoglobinemia; Drug-Induced Methemoglobinemia; Chemically Induced Methemoglobinemia

### Acquired Nasolacrimal Duct Obstruction

Blockage of the tear-drainage system that develops after birth. It may cause excessive tearing, discharge, or repeated inflammation of the lacrimal sac and can result from aging, infection, injury, inflammation, scarring, or a mass.

### Acquired Platelet Function Defect

Impaired platelet adhesion or aggregation that develops after birth. Causes include antiplatelet medicines, kidney failure, hematologic disease, and other systemic disorders; platelet counts may remain normal, but mucocutaneous bleeding can occur.

### Acquired Tracheomalacia

Acquired tracheomalacia is weakening of the tracheal wall after injury, inflammation, compression, or prolonged ventilation, causing airway collapse and breathing symptoms.

### Acquisition

The process by which new information or a skill is learned and initially encoded, especially in discussions of memory and learning.

**Alternate Names:**

Learning Acquisition; Memory Acquisition

### Acquisition Duration

The time required to collect a measurement, image, or body signal. Longer acquisition may improve detail or accuracy but can increase motion problems, discomfort, radiation exposure, or equipment time.

### Acquisition Time

The time during which a medical imaging system collects data. The required time depends on the imaging method, body region, protocol, and patient movement.

### Acral Lentiginous Melanoma

A subtype of melanoma arising on acral skin (palms, soles, or under nails), appearing as a dark, irregularly pigmented patch or plaque. It is more common in people with darker skin tones and often presents at a later stage due to less frequent surveillance of these sites. Prognosis depends on thickness and stage at diagnosis.

### Acro-Osteolysis

Resorption or destruction of the terminal bones of the fingers or toes. It can occur with inherited disorders, autoimmune disease, severe vascular disease, trauma, or repeated occupational exposure and may cause shortened, painful, or deformed digits.

### Acrocentric Chromosome

A chromosome whose central joining point lies close to one end, giving it a very short section and a longer section. In humans, chromosomes 13, 14, 15, 21, and 22 have this shape.

### Acrochordon

Alternate name; see *Skin Tag*.



---

## Acrocyanosis

Persistent bluish or purplish discoloration and coolness of the hands, feet, nose, or ears, usually without pain, caused by narrowing of small skin blood vessels. It is often worse in cold conditions and may be primary or due to another disorder.

## Acrodermatitis

Inflammation affecting skin on the hands, feet, or other body extremities. Causes vary and include infections, nutritional deficiency, inherited disorders, medicines, and immune reactions; appearance and treatment depend on cause.

## Acrodermatitis Chronica Atrophicans

A late cutaneous manifestation of Lyme borreliosis, usually caused by persistent *Borrelia afzelii* infection in Europe. It begins with unilateral reddish-violet swelling on an extremity and can progress over months or years to thin, atrophic, fibrotic skin; diagnosis uses clinical findings, serology, and sometimes biopsy.

## Acrodermatitis Enteropathica

A zinc-deficiency disorder classically causing periorificial and acral dermatitis, diarrhea, and alopecia. The primary inherited form results from impaired intestinal zinc absorption; acquired deficiency may follow malnutrition, malabsorption, or other illness, and untreated disease can impair growth, healing, and immunity.

## Acrodynea

A painful condition with red, swollen, sweaty hands and feet, classically associated with mercury poisoning and historically called pink disease in children.

## Acrodysostosis

A rare inherited disorder of bone development causing short, broad fingers and toes, a small nose and middle face, and short stature. Some people have learning difficulties or reduced response to parathyroid and other hormones.

## Acrokeratosis Paraneoplastica

A rare skin condition causing thick, wart-like

growths on the hands and feet, often around the nails, as a reaction to an internal cancer. It is most often linked with cancers of the digestive or respiratory tract and needs medical evaluation.

## Acromegaly

A disorder caused by excess growth hormone after height growth has ended, usually from a noncancerous pituitary adenoma. It enlarges the hands, feet, jaw, and soft tissues and may cause headaches, joint problems, diabetes, or vision changes.

## Acromioclavicular Joint

The small joint at the top of the shoulder where the collarbone meets the shoulder blade. It allows limited gliding as the arm moves and is held stable mainly by two groups of strong ligaments.

## Acromioclavicular Joint Injuries

Sprains or dislocations of the joint between the acromion and clavicle, usually caused by a fall or direct shoulder impact. Severity ranges from ligament sprain to complete displacement and is assessed clinically and with imaging.

### Alternate Names:

Acromioclavicular Joint Injury

## Acromion

A bony projection from the shoulder blade that forms the highest point of the shoulder. It joins the collarbone and forms a roof over the shoulder joint, with a tendon and lubricating sac passing beneath it.

## Acroparaesthesia

Acroparaesthesia is tingling, pins-and-needles, burning, or altered sensation in the hands or feet, commonly related to nerve compression, neuropathy, vascular disease, or metabolic causes.

## Acrophobia

An excessive, persistent fear of heights that causes marked anxiety, avoidance, or impairment beyond the ordinary caution associated with high places.

---

## Acropustulosis of Infancy

A recurrent itchy eruption of tiny vesicles and pustules, mainly on the palms and soles of infants and young children. Crops become less frequent and usually resolve by early childhood; scabies may precede or mimic it.

## Acrosin

An enzyme released from the cap-like structure at a sperm's tip. It helps sperm pass through the egg's surrounding layers during fertilization, although successful fertilization requires many additional steps.

## Acrosome

A cap-like structure on the head of a sperm cell containing enzymes that help it attach to and enter the egg during fertilization.

## Acrosome Reaction

The release of enzymes from the cap at the front of a sperm cell after it contacts an egg. These enzymes help the sperm pass through the egg's outer coating; other steps are also needed for fertilization.

## ACTH Blood Test

A blood test that measures the level of ACTH, a hormone made by the pituitary gland. It is usually checked with a cortisol test to help find problems of the adrenal or pituitary glands.

## ACTH Stimulation Test

A test that shows whether the adrenal glands can make cortisol. A synthetic form of ACTH is given, and cortisol levels are measured before and after it is given. A small rise may suggest adrenal insufficiency.

## Actigraphy

Recording movement over days or weeks with a small wearable sensor, usually worn on the wrist. The pattern of movement helps estimate when a person is asleep or awake and how active they are, but it does not measure sleep directly and cannot by itself diagnose every sleep problem.

**Alternate Names:**

Actigraphic Monitoring

## Actin

A protein that helps cells keep their shape and move. In muscle, actin works with myosin to produce contraction; in other cells, it supports movement, internal transport, and changes in cell structure.

**Alternate Names:**

Actin Filament; Microfilament Protein

## Actin Myofilament

A thin muscle fiber made mainly of actin that interacts with myosin, allowing muscle cells to shorten and produce force during contraction and movement.

**Alternate Names:**

Thin Filament

## Actin-Binding Protein

A protein that attaches to actin filaments and helps build, organize, move, or anchor the cell's internal framework. Examples include myosin, gelsolin, and alpha-actinin.

## Acting Out

Expressing emotional conflict or distress through behavior rather than words.

## Actinic

Relating to ultraviolet radiation from sunlight or UV lamps; chronic actinic exposure can cause sunburn, photoaging, and actinic keratoses.

## Actinic Cheilitis

A precancerous sun-damage condition of the lips, usually the lower lip, caused by long-term ultraviolet exposure. It may cause persistent dryness, scaling, whitening, thickening, or blurring of the lip border and can progress to squamous cell carcinoma.

## Actinic Granuloma

A usually harmless inflammatory skin condition that occurs on sun-damaged skin. It causes ring-shaped patches with a thinner center and a raised red border, most often on the neck, arms, or backs

---

of the hands.

### **Actinic Keratosis**

A rough, scaly, sandpaper-like lesion caused by cumulative ultraviolet damage, usually on sun-exposed skin such as the face, scalp, ears, neck, forearms, or hands. It is a keratinocytic precancer and may progress to cutaneous squamous-cell carcinoma, so changing, thick, tender, or bleeding lesions need assessment and treatment.

#### **Alternate Names:**

Solar Keratosis; AK; Senile Keratosis

### **Actinic Keratosis Pathology**

Atypical keratinocytes with disordered maturation in the lower epidermis, often with parakeratosis, alternating orthokeratosis, and solar elastosis. It is a premalignant squamous lesion that can progress to cutaneous squamous cell carcinoma.

### **Actinic Purpura**

Bruising caused by fragile, sun-damaged skin and connective tissue, usually on the backs of the hands and forearms in older adults. The purple patches may appear after minor trauma and are generally harmless, although medicines such as anticoagulants can worsen them. New, widespread, or unexplained bruising warrants assessment.

### **Actinomycetoma**

A slowly worsening bacterial infection of the skin and deeper tissue, usually affecting the foot after soil contamination. It causes swelling, firm lumps, draining channels, and grains in the discharge; untreated disease can damage bone.

### **Actinomycin D**

Actinomycin D, or dactinomycin, is an antineoplastic antibiotic that blocks DNA-dependent RNA synthesis and is used for selected cancers; myelosuppression and tissue injury are important risks.

### **Actinomycosis**

A long-lasting infection caused by a type of bacteria called *Actinomyces* that normally lives in the mouth, intestines, and female genital area. It starts when

these bacteria get past a damaged surface, such as after a dental problem, dental work, abdominal surgery, or use of an intrauterine device. It most often affects the face and neck, chest, or belly and pelvis, where it forms hard, growing lumps and draining tunnels that contain small yellow grains called sulfur granules. It can spread to nearby tissue and look like cancer. Diagnosis uses cultures or tissue samples, and treatment is a long course of antibiotics, usually high-dose penicillin.

### **Action Potential**

A brief electrical signal that travels along a nerve or muscle cell. It begins when the cell's electrical charge changes enough to trigger the signal and allows nerves to communicate and muscles to contract.

### **Action Tremor**

An involuntary rhythmic oscillation of a body part occurring during active voluntary muscle contraction. Subtypes include postural tremors (maintaining a position against gravity), kinetic tremors (moving through an arc), and intention tremors (worsening near the target).

### **Activated Charcoal**

A porous form of carbon that adsorbs some substances in the gastrointestinal tract and can reduce their absorption after selected poisonings. It is not effective for every toxin and can cause aspiration or other complications.

### **Activated Partial Thromboplastin Time**

A blood test that measures clotting through the intrinsic and common coagulation pathways. It may be prolonged by unfractionated heparin, clotting-factor deficiencies, inhibitors, or some liver and systemic disorders.

### **Activated Protein C Resistance**

An inherited thrombophilic disorder characterized by an impaired anticoagulant response of plasma to activated protein C (APC). Approximately 90 to 95 percent of cases are caused by the factor V Leiden mutation (G1691A), which alters the APC cleavage

---

site on factor V, predisposing affected individuals to recurrent deep venous thrombosis and pulmonary embolism.

**Alternate Names:**

APC Resistance; Factor V Leiden Thrombophilia

**Active Caries**

Tooth decay that is continuing to progress. The affected surface is often soft, rough, or chalky and requires dental assessment and management.

**Active Management Of Third Stage**

A set of measures used after birth to reduce postpartum hemorrhage, usually including a uterotonic medicine, controlled cord traction when appropriate, and assessment of uterine tone and bleeding.

**Active Phase Of Labor**

The part of the first stage of labor in which the cervix dilates progressively, generally beginning at about 6 cm and continuing to full dilation. Progress varies between individuals.

**Active Site**

The part of an enzyme where a target molecule attaches and a chemical reaction takes place. Its shape and chemical properties help the enzyme recognize particular molecules and change them into products.

**Active Surveillance**

Close, structured monitoring of a cancer or precancerous condition without immediate treatment. It includes scheduled examinations, laboratory tests, imaging, or biopsies, with treatment started if evidence of progression develops.

**Active Transport**

The movement of a substance across a cell membrane using cellular energy. It often moves the substance against a concentration or electrical gradient and helps cells control salts, nutrients, waste products, and other substances.

**Activities Of Daily Living**

Routine tasks required for personal care and independent living, including basic activities and more complex instrumental activities. Performance is assessed to describe functional status and support care planning.

**Acupressure**

A complementary treatment in which firm pressure is applied to selected points on the body, often with the fingers. It may provide short-term relief of symptoms such as nausea or pain, but evidence of benefit varies by condition.

**Acupuncture**

A treatment originating in traditional Chinese medicine in which thin needles are inserted into the skin and sometimes moved or electrically stimulated. It may help some pain conditions or treatment-related nausea, but benefits vary.

**Acute**

Describing symptoms or a condition that begins or worsens quickly and usually lasts a limited time. It is commonly contrasted with chronic, which develops or continues over a longer period.

**Acute Abdomen**

Sudden or severe abdominal pain with clinical features suggesting a potentially serious intra-abdominal disorder that may require urgent investigation or surgery.

**Acute Acalculous Cholecystitis**

Acute gallbladder inflammation without gallstones, usually occurring in critically ill, injured, postoperative, or fasting patients. Ischaemia, bile stasis, and infection may contribute; it can progress rapidly to necrosis or perforation and requires urgent assessment.

**Acute Adrenal Crisis**

Acute adrenal crisis is life-threatening cortisol deficiency causing hypotension, vomiting, abdominal pain, electrolyte disturbance, hypoglycemia, or shock; urgent glucocorticoid and fluid treatment is

---

required.

### **Acute Agitation**

A state of heightened motor activity, emotional tension, irritability, and impaired behavioral control that may threaten the patient or others. Management begins with de-escalation and assessment for delirium, intoxication, withdrawal, medical illness, and psychosis.

### **Acute Alcohol Intoxication**

Neurobehavioral and physiologic impairment caused by recent ethanol exposure. Severity varies with concentration, tolerance, co-ingestants, and comorbidity; dangerous effects include respiratory depression, aspiration, hypoglycemia, hypothermia, trauma, and coma.

### **Acute Angle-Closure Glaucoma**

An ophthalmic emergency in which sudden closure of the anterior-chamber angle causes a rapid rise in intraocular pressure. Severe eye pain, blurred vision, halos, headache, nausea, vomiting, and a red eye may occur; urgent ophthalmic treatment is required to protect vision.

### **Acute Aortic Syndrome**

A group of sudden, life-threatening problems in the wall of the aorta, the body's largest artery. The wall may tear, bleed within itself, or develop an ulcer, causing severe chest or back pain and requiring emergency assessment.

### **Acute Bacterial Prostatitis**

A sudden bacterial infection of the prostate causing fever, chills, pain between the scrotum and anus or in the lower back, painful urination, and sometimes inability to pass urine. It requires prompt antibiotics and medical care; a forceful prostate examination is avoided because it can spread bacteria into the bloodstream.

### **Acute Bacterial Rhinosinusitis**

Bacterial infection of the paranasal sinuses suspected when symptoms persist without improvement for at least 10 days, become substantially

worse after initial improvement, or are severe with high fever and purulent discharge. Treatment decisions depend on severity, complications, resistance risk, and local guidance.

### **Acute Bilirubin Encephalopathy**

Brain dysfunction in a newborn caused by very high bilirubin levels. Poor feeding, sleepiness, weak muscle tone, irritability, a high-pitched cry, unusual movements, seizures, or coma may develop. It is an emergency requiring rapid treatment to prevent permanent brain injury.

#### **Alternate Names:**

Bilirubin Neurotoxicity; Kernicterus Spectrum Disorder

### **Acute Blood Loss**

Rapid loss of blood that reduces the amount circulating through the body. It can cause weakness, fast heartbeat, low blood pressure, confusion, organ injury, or shock and requires urgent control of bleeding and replacement of lost volume.

### **Acute Bronchitis**

Short-term inflammation of the large airways in the lungs, usually caused by a virus such as a cold or flu. It causes a cough that may bring up clear, yellow, or green mucus and may cause chest soreness, mild fever, tiredness, or wheezing. The cough often lasts 1 to 3 weeks. Because viruses cause more than 90% of cases, antibiotics usually do not help; care focuses on rest, fluids, and symptom relief.

#### **Alternate Names:**

Chest Cold; Tracheobronchitis

### **Acute Bursitis**

Sudden, acute inflammation of a synovial bursa, typically triggered by direct trauma, acute crystal deposition (gout/pseudogout), bacterial infection (most commonly *Staphylococcus aureus*), or unaccustomed repetitive strain. It presents with rapid-onset localized swelling, erythema, warmth, exquisite focal tenderness, and restricted active joint motion.

#### **Alternate Names:**

Acute Bursal Synovitis; Traumatic Bursitis; Septic Bursitis



---

### Acute Caries

Tooth decay that progresses rapidly, often producing a soft lesion or cavity. Risk is increased by factors such as frequent sugar exposure, reduced saliva, or poor oral hygiene.

### Acute Cerebellar Ataxia

Sudden or rapidly developing incoordination caused by cerebellar dysfunction. In children it may follow an infection, but stroke, intoxication, inflammation, tumor, and other urgent causes must be considered.

### Acute Chest Syndrome

A potentially life-threatening pulmonary complication of sickle cell disease characterized by a new lung infiltrate plus fever or respiratory symptoms such as chest pain, cough, hypoxemia, or breathing difficulty.

### Acute Cholangitis

Alternate name; see *Ascending Cholangitis*.

### Acute Cholecystitis

Acute inflammation of the gallbladder, most often caused by persistent cystic-duct obstruction by a gallstone. It commonly causes sustained right-upper-quadrant pain, fever, nausea, vomiting, and a positive Murphy sign; diagnosis combines clinical, laboratory, and imaging findings, usually including ultrasound. Early laparoscopic cholecystectomy is generally preferred when appropriate, with drainage considered when surgery is unsafe.

### Acute Colonic Pseudo-Obstruction

Severe widening of the large intestine without a physical blockage, usually in people who are seriously ill, injured, or recovering from surgery. It can cause pain, swelling, nausea, and vomiting. If it is not treated, the bowel may lose its blood supply or tear.

#### Alternate Names:

Ogilvie Syndrome; Colonic Pseudo-Obstruction

### Acute Compartment Syndrome

A rapid rise in pressure inside a closed muscle com-

partment that reduces blood flow and can permanently damage muscles and nerves. It usually follows a fracture, crush injury, or severe swelling and requires emergency assessment.

#### Alternate Names:

Compartment Syndrome; Intracompartmental Hypertension

### Acute Coronary Syndrome

A clinical spectrum of acute myocardial ischemia caused most often by coronary plaque disruption and thrombosis. It includes unstable angina, non-ST-elevation myocardial infarction (NSTEMI), and ST-elevation myocardial infarction (STEMI); ECG findings and serial cardiac troponin distinguish the major presentations and guide urgent treatment.

### Acute Coronary Syndromes

Another name for *Acute Coronary Syndrome*.

### Acute Cough

A cough lasting less than three weeks. Common causes include viral respiratory infection, acute bronchitis, and pneumonia.

### Acute Cutaneous Lupus Erythematosus (ACLE)

A lupus-specific inflammatory skin eruption that commonly presents as a transient malar or generalized erythematous rash, often triggered by sunlight and associated with systemic lupus activity. The malar rash usually spares the nasolabial folds; evaluation should consider systemic symptoms and organ involvement.

### Acute Cystitis

Acute bacterial infection and inflammation confined to the lower urinary tract and bladder mucosa, typically caused by uropathogenic *Escherichia coli* and presenting with acute dysuria, urinary frequency, urgency, suprapubic pain, and hematuria.

#### Alternate Names:

Acute Bacterial Cystitis; Bladder Infection; Acute UTI



---

### **Acute Dacryocystitis**

Acute infection and inflammation of the lacrimal sac, most commonly caused by obstruction of the nasolacrimal duct leading to stasis and secondary bacterial infection. It presents with painful swelling, redness, and tenderness at the medial canthus with purulent discharge.

### **Acute Decompensated Heart Failure**

Sudden or rapidly worsening heart failure, causing breathlessness, tiredness, leg swelling, or fluid in the lungs. It may be triggered by infection, a heart attack, rhythm problems, uncontrolled blood pressure, or missed treatment and can require urgent hospital care.

### **Acute Diarrhea**

Passage of loose or watery stools, usually three or more times daily, lasting less than two weeks. Infection, food-related illness, medicines, and other intestinal disorders can cause it. Oral rehydration is important; blood in stool, severe dehydration, persistent vomiting, or severe pain requires prompt assessment.

### **Acute Disease**

An illness that begins suddenly or develops over a short period. It often lasts a limited time, although it can be mild, severe, or life-threatening and may sometimes lead to a longer-term condition.

#### **Alternate Names:**

Acute Illness

### **Acute Disseminated Encephalomyelitis**

A rare sudden inflammation of the brain and spinal cord, usually after an infection and less often after vaccination. It can cause confusion, weakness, poor coordination, vision problems, or seizures, especially in children.

### **Acute Ear Infection**

A sudden, rapid-onset bacterial or viral infection of the middle ear space behind the eardrum, characterized by earache, fever, hearing muffled sounds, and bulging of the tympanic membrane.

#### **Alternate Names:**

Acute Otitis Media; AOM; Middle Ear Infection

### **Acute Endometritis**

Acute inflammation of the endometrium, usually caused by ascending infection or retained tissue after pregnancy-related events or uterine instrumentation; fever, pelvic pain, and abnormal bleeding may occur.

### **Acute Epididymitis**

Sudden inflammation of the epididymis, usually caused by infection and sometimes involving the testis. It causes unilateral scrotal pain and swelling and must be distinguished urgently from testicular torsion.

### **Acute Esophageal Necrosis**

Ischemic necrosis of the esophageal mucosa, most commonly caused by low-flow states such as shock, sepsis, or thromboembolism, presenting with hematemesis and chest pain. Endoscopy shows circumferential black necrotic mucosa, typically in the distal esophagus.

### **Acute Exacerbation**

A rapid worsening of a chronic pulmonary condition, with increased breathlessness, cough, oxygen need, or other symptoms; infection, aspiration, embolism, and other triggers may contribute.

### **Acute Exacerbation Of Asthma**

A progressive increase in dyspnea, cough, wheezing, or chest tightness with reduced expiratory airflow compared with the patient's baseline. Severe attacks may cause exhaustion, silent chest, hypercapnia, respiratory failure, and need for urgent treatment.

### **Acute Exacerbation Of COPD**

A sudden worsening of chronic obstructive pulmonary disease, with more breathlessness, cough, or sputum than usual. It is often triggered by a respiratory infection, air pollution, or another stress on the lungs.

### **Acute Fatty Liver Of Pregnancy**

A rare, potentially life-threatening liver disorder

---

that usually develops late in pregnancy. Fat builds up in liver cells and may cause nausea, abdominal pain, jaundice, low blood sugar, abnormal bleeding, kidney failure, or other organ failure.

### **Acute Febrile Illness**

A sudden illness characterized by fever, often with headache, chills, muscle aches, gastrointestinal symptoms, or other signs of infection; the cause may be infectious or noninfectious.

### **Acute Febrile Neutrophilic Dermatositis (Sweet Syndrome)**

An inflammatory neutrophilic skin disease causing abrupt tender red or violaceous papules, plaques, or nodules with fever and neutrophilia. It may be triggered by infection, inflammatory disease, medicines, pregnancy, or an underlying malignancy; biopsy confirms the characteristic neutrophilic infiltrate.

### **Acute Fever in Adults**

A fever in an adult with recent onset, commonly caused by infection, inflammation, medication, heat illness, or other conditions. Severity, vital signs, immune status, source, and associated symptoms guide evaluation; confusion, breathing difficulty, stiff neck, shock, or severe localized pain requires urgent care.

### **Acute Flaccid Myelitis**

A rare serious illness, usually in children, causing sudden weakness and limpness of one or more arms or legs. Facial weakness, swallowing difficulty, or breathing failure may occur; immediate hospital assessment, imaging, and breathing monitoring are needed.

### **Acute Gastritis**

Sudden inflammation or irritation of the stomach lining. It may cause upper-abdominal pain, nausea, vomiting, or bleeding and can result from anti-inflammatory medicines, alcohol, severe illness, infection, or other irritants.

#### **Alternate Names:**

Acute Erosive Gastritis

### **Acute Generalized Exanthematous Pustulosis**

A sudden skin reaction, usually to a medicine, that causes many small pus-filled bumps on widespread red skin. Fever may occur. It often starts within days of exposure and improves after the triggering medicine is stopped.

### **Acute Glomerulonephritis**

Acute immune or infectious inflammation of the glomeruli producing a nephritic syndrome: hematuria with dysmorphic red cells or red-cell casts, variable proteinuria, edema, hypertension, and reduced kidney function. Causes include postinfectious disease, IgA nephropathy, vasculitis, and lupus; complement, serology, and sometimes kidney biopsy help identify the cause.

### **Acute Granulocytic Leukemia**

See *Acute myelogenous leukemia*.

### **Acute Hemorrhagic Edema of Infancy**

A usually self-limited cutaneous small-vessel vasculitis of infants, characterized by fever, edema, and large bruise-like or targetoid purpuric lesions on the face and extremities. Internal-organ involvement is uncommon, but serious causes of purpura must be excluded; most cases resolve within one to three weeks.

### **Acute Hemorrhagic Leukoencephalitis**

A rare, fulminant inflammatory demyelinating disease of the brain, considered a severe form of acute disseminated encephalomyelitis. It can cause rapidly worsening headache, fever, confusion, seizures, weakness, coma, and brain swelling and requires emergency neurologic care.

### **Acute Hepatitis**

Sudden inflammation and injury of the liver, developing over days to months. It may cause tiredness, nausea, pain under the right ribs, dark urine, jaundice, or no symptoms. Causes include viruses, alcohol, medicines, toxins, and autoimmune disease.

---

### Acute HIV Syndrome

A flu-like illness that may occur about 2–6 weeks after HIV infection, when the virus is multiplying rapidly. Fever, sore throat, swollen glands, rash, muscle aches, headache, nausea, diarrhea, or mouth ulcers may occur, but some people have no symptoms.

### Acute Hypercapnic Respiratory Failure

A sudden inability to breathe out enough carbon dioxide, causing it to build up in the blood. Severe lung disease, asthma, weak breathing muscles, sedative medicines, or chest problems can cause confusion, sleepiness, or dangerous breathing failure.

### Acute Hypoxemic Respiratory Failure

A sudden inability of the lungs to put enough oxygen into the blood. Pneumonia, fluid in the lungs, severe inflammatory lung injury, collapsed lung tissue, or a blood clot in the lungs may cause severe breathlessness, bluish skin, confusion, or organ injury.

### Acute Idiopathic Polyneuritis

Guillain-Barré syndrome, an acute immune-mediated disorder of peripheral nerves that usually causes progressive symmetric weakness and reduced reflexes and may impair breathing.

### Acute Illness

A disease or health problem that begins suddenly or over a short period and generally lasts a limited time, although its severity may range from mild to life-threatening.

### Acute Inflammation

A rapid immune response to injury or infection that usually lasts hours to days. It can cause redness, warmth, swelling, pain, and reduced function as extra blood, fluid, and immune cells enter the affected area.

### Acute Inflammatory Demyelinating Polyneuropathy

See *Guillain-Barre syndrome*.

### Acute Inflammatory Demyelinating Polyradiculoneuropathy

The most common form of Guillain-Barré syndrome. The immune system damages the protective covering of nerves, causing progressive weakness, tingling, and reduced reflexes that may spread to the breathing muscles.

### Acute Intermittent Porphyria

An inherited disorder of haem production caused by reduced hydroxymethylbilane synthase activity. Attacks may cause severe abdominal pain, nausea, constipation, dark urine, nerve symptoms, confusion, or seizures; medicines, hormonal changes, fasting, and illness can trigger them.

### Acute Interstitial Nephritis

Sudden inflammation of the tissue between the kidney's filtering structures, most often caused by a reaction to a medicine. Infections and autoimmune disease can also cause it, leading to blood or protein in urine and a sudden decline in kidney function.

### Acute Ischemic Stroke

Sudden loss of brain, spinal-cord, or retinal function because a blood vessel is blocked and the tissue does not receive enough oxygen. Emergency imaging distinguishes it from bleeding and determines urgent treatment.

### Acute Kidney Injury

An abrupt decline in kidney function, defined by a serum-creatinine rise of at least 0.3 mg/dL within 48 hours, a rise to at least 1.5 times baseline within 7 days, or urine output below 0.5 mL/kg/hour for 6 hours. Causes are commonly prerenal hypoperfusion, intrinsic kidney injury, or postrenal obstruction; complications include azotemia, fluid and electrolyte disorders, and acidosis.

#### Alternate Names:

AKI; Acute Kidney Failure; Acute Renal Failure

### Acute Kidney Injury Staging

A standardized classification based on changes in serum creatinine and urine output. KDIGO stages 1 through 3 quantify the severity of abrupt kidney-

---

function decline and support prognosis, monitoring, and treatment decisions.

### **Acute Leukemia**

A fast-growing blood cancer in which immature abnormal blood cells crowd the bone marrow and interfere with normal blood-cell production. It can cause anemia, infections, and bleeding and needs urgent treatment.

### **Acute Limb Ischemia**

Sudden reduction in arterial blood flow to a limb that threatens tissue viability, usually caused by embolism, thrombosis, or graft occlusion. The six classic findings are pain, pallor, pulselessness, paresthesia, paralysis, and poikilothermia; urgent vascular assessment and revascularization are required.

### **Acute Limb Ischemia Six Ps**

Six warning findings of suddenly reduced artery blood flow to a limb: pain, paleness, absent pulse, tingling or numbness, weakness or paralysis, and a cold limb. This is an emergency requiring urgent vascular assessment.

### **Acute Liver Failure**

Rapid deterioration of liver function in a previously healthy individual without prior liver disease. Causes include acetaminophen toxicity, viral hepatitis, drug reactions, autoimmune hepatitis, and Wilson disease. Features include coagulopathy, encephalopathy, and multiorgan failure. Presents with jaundice, confusion, and bleeding.

#### **Alternate Names:**

Fulminant Hepatic Failure; Acute Hepatic Necrosis

### **Acute Low Back Pain**

Lumbosacral pain lasting less than 4 to 6 weeks, most commonly caused by mechanical musculoligamentous strain, facet joint irritation, or acute lumbar disc herniation. Management emphasizes conservative therapy, staying active, and monitoring for neurological red flags.

#### **Alternate Names:**

Acute Lumbago; Mechanical Low Back Strain; Lumbosacral Strain

### **Acute Lymphangitis**

Acute inflammation of lymphatic vessels, typically resulting from streptococcal or staphylococcal infection spreading from a distal wound. It presents with red, tender streaks extending proximally from the site of infection, along with fever, malaise, and regional lymphadenopathy. Treatment requires systemic antibiotics.

### **Acute Lymphoblastic Leukemia**

Acute lymphoblastic leukemia is a rapidly progressive cancer of immature lymphoid precursor cells that crowds the bone marrow and can involve blood, lymph nodes, or the central nervous system.

### **Acute Lymphocytic Leukemia**

A malignant proliferation of immature lymphoid precursors in the bone marrow and blood, commonly presenting with cytopenias, fever, bone pain, or lymphadenopathy.

#### **Alternate Names:**

ALL; Acute Lymphoblastic Leukemia

### **Acute Lymphoid Leukemia**

See *Acute lymphocytic leukemia*.

### **Acute Megakaryoblastic Leukemia**

A rare subtype of acute myeloid leukemia characterized by the proliferation of megakaryoblasts in the bone marrow and peripheral blood.

### **Acute Mesenteric Ischemia**

A vascular surgical emergency caused by an abrupt reduction in intestinal blood flow (most commonly from superior mesenteric artery thromboembolism), classically presenting with severe abdominal pain out of proportion to physical examination findings, rapidly progressing to bowel necrosis and peritonitis.

#### **Alternate Names:**

Acute Mesenteric Infarction; AMI

### **Acute Mitral Regurgitation**

Sudden backward flow of blood from the left ventricle into the left atrium during systole, caused by acute leaflet, chordal, papillary-muscle, or ventricu-

---

lar injury. It can rapidly produce pulmonary edema, low cardiac output, or shock.

### **Acute Mountain Sickness**

A high-altitude illness caused by inadequate acclimatization, usually after ascent above about 2,500 m (8,200 ft). It is characterized by headache plus symptoms such as nausea, loss of appetite, dizziness, fatigue, or disturbed sleep. Worsening symptoms, confusion, inability to walk normally, or breathlessness at rest indicate progression to life-threatening high-altitude cerebral or pulmonary edema.

### **Acute Myeloblastic Leukemia**

See *Acute Myeloid Leukemia*.

### **Acute Myelogenous Leukemia**

Alternate name; see *Acute Myeloid Leukemia*.

### **Acute Myeloid Leukemia**

A clonal hematopoietic neoplasm characterized by the rapid proliferation and accumulation of abnormal, poorly differentiated myeloid progenitor cells (blasts) in the bone marrow and peripheral blood. This suppresses normal hematopoiesis, leading to progressive cytopenias manifested as fatigue from anemia, recurrent or opportunistic infections from neutropenia, and petechiae, ecchymoses, or mucosal bleeding from thrombocytopenia.

#### **Alternate Names:**

AML; Acute Myelogenous Leukemia; Acute Granulocytic Leukemia; Acute Non-Lymphocytic Leukemia

### **Acute Myocardial Infarction**

Sudden death of heart muscle caused by prolonged inadequate coronary blood flow, usually from an acutely obstructed coronary artery; it is a medical emergency.

#### **Alternate Names:**

AMI

### **Acute Myocardial Infarction Pathology**

Histopathologic evolution of myocardial infarction, from early ischemic injury and coagulative necrosis

through inflammation, granulation tissue, and scar formation. The findings depend on the interval between infarction and examination.

### **Acute Necrotizing Ulcerative Gingivitis**

A painful, rapidly developing infection of the gums characterized by punched-out gingival papillae, ulceration, spontaneous bleeding, gray necrotic tissue, and foul breath. It is associated with anaerobic oral bacteria and risk factors such as poor oral hygiene, smoking, stress, malnutrition, or immunosuppression; urgent dental evaluation, cleaning, and treatment are needed.

#### **Alternate Names:**

ANUG; Trench Mouth; Vincent Stomatitis

### **Acute Nephritic Syndrome**

A clinical syndrome of glomerular inflammation marked by hematuria, variable proteinuria, edema, hypertension, and reduced kidney function.

### **Acute Nonlymphocytic Leukemia**

See *Acute myelogenous leukemia*.

### **Acute Obstructive Uropathy**

Sudden impairment of urinary flow caused by obstruction of the urinary tract, potentially producing hydronephrosis, pain, infection, and acute kidney injury.

### **Acute Onset Pain**

Pain that begins suddenly, often over seconds, minutes, or hours.

#### **Alternate Names:**

Sudden-Onset Pain

### **Acute Otitis Media**

Infection of the middle ear space with acute onset of symptoms and signs of middle ear inflammation. Diagnosis requires middle ear effusion plus acute symptoms (ear pain, fever, irritability) or signs (bulging tympanic membrane; erythema alone is not diagnostic). Most common pathogen: *Streptococcus pneumoniae*.



---

### Acute Pain

Pain that begins suddenly and usually lasts for a limited period, often because of injury, illness, or a procedure.

### Acute Pancreatitis

Sudden inflammation of the pancreas, most often caused by gallstones or alcohol. It usually causes severe upper-abdominal pain that may reach the back, with nausea or vomiting; complications can include fluid loss, infection, organ failure, or pancreatic collections.

### Acute Pericarditis

Rapid-onset inflammation of the sac around the heart, often idiopathic or viral and sometimes related to autoimmune disease, heart injury, kidney disease, cancer, or medicines. It typically causes sharp chest pain that worsens with breathing or lying down and improves when sitting up and leaning forward; a friction rub, diffuse ST-segment elevation with PR-segment depression, or a pericardial effusion may occur.

### Acute Poisoning

Harmful effects that develop after a short-term exposure to a toxic substance by swallowing, inhaling, injection, skin contact, or another route. Effects depend on the substance, amount, route, and time of exposure.

### Acute Proctitis

Sudden inflammation of the rectal mucosa causing rectal pain, urgency, tenesmus, bleeding, mucus, or discharge. Infectious, inflammatory, ischemic, traumatic, and radiation-related causes are possible.

### Acute Promyelocytic Leukemia

A subtype of acute myeloid leukemia that can quickly cause severe bleeding or abnormal clotting. It is linked to a specific chromosome change and needs immediate specialist treatment.

### Acute Pulmonary Edema

Acute or rapidly worsening accumulation of fluid in the lungs; see *Pulmonary Edema*.

### Acute Pulmonary Histoplasmosis

An acute lower respiratory tract infection caused by inhaling microconidia of the dimorphic fungus *Histoplasma capsulatum*, commonly found in soil enriched with bird or bat guano. Manifestations range from mild flu-like symptoms to severe pneumonia with mediastinal lymphadenopathy.

#### Alternate Names:

Primary Pulmonary Histoplasmosis; Acute Histoplasmosis Pneumonia; Cave Disease

### Acute Pyelonephritis

A sudden bacterial infection of the kidney, usually spreading upward from the bladder. It can cause fever, chills, pain in the side or back, nausea, and painful or frequent urination. Prompt medical treatment is important because the infection can spread to the bloodstream and damage the kidney.

### Acute Radiation Sickness

Illness caused by receiving a high dose of penetrating radiation over a short time, potentially damaging bone marrow, intestines, skin, and other organs.

### Acute Radiation Syndrome

See *Radiation sickness*.

### Acute Rehabilitation

Multidisciplinary rehabilitation provided during or soon after hospitalization for conditions such as stroke, spinal-cord injury, traumatic brain injury, or major trauma. Treatment is tailored to functional needs.

### Acute Renal Artery Occlusion

Sudden obstruction of blood flow through one or both main renal arteries or their primary branches, most commonly caused by thromboembolism, in situ thrombosis, or trauma. It presents with severe flank or abdominal pain, hematuria, refractory hypertension, and rapid deterioration of kidney function.

#### Alternate Names:

Acute Renal Infarction; Renal Artery Thrombosis; Renal Artery Embolism



---

### Acute Renal Failure

See *Acute kidney injury*.

### Acute Respiratory Disease

A general term for an illness affecting breathing that begins suddenly or over a short time. It may include infections, asthma attacks, or other lung problems and is not one specific diagnosis.

**Alternate Names:**

ARD

### Acute Respiratory Distress Syndrome

A sudden, severe lung injury that develops after a serious illness or injury. Fluid leaks into the lungs' air sacs, making it hard for oxygen to enter the blood. It usually begins within a week of the trigger and causes widespread changes in both lungs on imaging; heart failure or fluid overload alone does not explain it. Treatment addresses the cause and supports breathing; some people need a ventilator.

### Acute Respiratory Failure

A sudden inability of the respiratory system to maintain adequate oxygenation, carbon-dioxide removal, or both. It may result from lung disease, airway obstruction, neuromuscular weakness, or impaired respiratory drive and requires urgent evaluation.

### Acute Retinal Necrosis

A rapidly progressive necrotizing viral retinitis, usually caused by herpes viruses, with vitreous inflammation, peripheral retinal necrosis, occlusive vasculitis, and risk of retinal detachment.

### Acute Retroviral Syndrome

The acute illness that may occur shortly after HIV infection, commonly involving fever, rash, sore throat, lymph-node enlargement, muscle aches, or headache.

### Acute Rhinitis

Sudden inflammation of the nasal lining, commonly caused by a viral infection or an irritant, producing congestion, nasal discharge, and sneezing.

### Acute Rhinosinusitis

See *Acute sinusitis*.

### Acute Sinusitis

Acute inflammation of the mucosa of one or more paranasal sinuses, usually following a viral upper-respiratory infection. It causes nasal obstruction, purulent or discoloured discharge, facial pressure or pain, and reduced smell; persistent, severe, or worsening symptoms may indicate acute bacterial rhinosinusitis or a complication.

**Alternate Names:**

Acute Rhinosinusitis; Paranasal Sinus Infection

### Acute Stress Disorder

A trauma-related mental-health condition beginning after exposure to actual or threatened death, serious injury, or sexual violence and lasting from three days to one month. Symptoms may include intrusive memories, avoidance, negative mood, dissociation, and increased arousal.

### Acute Symptomatic Hyponatremia

A recently developing low blood sodium concentration accompanied by symptoms, particularly neurologic symptoms such as headache, confusion, seizures, or reduced consciousness. Clinical significance depends on the sodium level, rate of change, underlying cause, and associated illness.

### Acute Transverse Myelitis

Inflammation across a segment of the spinal cord that can cause rapidly developing weakness, sensory changes, and bladder or bowel dysfunction.

### Acute Tubular Necrosis

Sudden injury and death of cells in the kidney's tiny tubes, which normally help filter and balance the blood. It can follow prolonged low blood flow, severe illness, or toxic effects from medicines and other substances, causing acute kidney injury.

**Alternate Names:**

ATN

---

### Acute Urinary Retention

Sudden inability to pass urine despite a full bladder, often causing severe lower-abdominal pain and swelling. Causes include blockage, medicines, nerve problems, or impaired bladder contraction.

### Acute Viral Hepatitis

A short-term infection that inflames the liver, caused by a hepatitis virus. Hepatitis A and E usually cause acute infection; hepatitis B, C, and D can also become chronic, and hepatitis D occurs only with hepatitis B. Symptoms may include tiredness, nausea, right-upper-abdominal discomfort, dark urine, or jaundice, but some people have none.

### Acute Viral Illness

A short-duration illness caused by a virus. Symptoms depend on the affected organ system and may include fever, fatigue, cough, vomiting, diarrhea, or rash.

### Acute Vision Loss

Sudden reduction or loss of vision in one or both eyes caused by retinal vascular occlusion, retinal detachment, optic-nerve disease, stroke, acute glaucoma, inflammation, trauma, or other emergency. It requires immediate medical or ophthalmic assessment.

### Acute Weeping Dermatitis

A form of acute dermatitis with inflamed, often blistered skin that releases clear or yellowish fluid and may form crusts. It is a descriptive pattern that can occur with contact dermatitis, eczema, infection, or other inflammatory skin disorders.

### Acute-On-Chronic Liver Failure

Acute deterioration of pre-existing chronic liver disease associated with organ failure and high short-term mortality. It is commonly precipitated by infection, alcohol-related hepatitis, bleeding, or other acute insults and requires urgent multidisciplinary management.

### Acute-Phase Reactants

A group of plasma proteins whose circulating concen-

trations increase or decrease by at least 25 percent during inflammatory states, tissue injury, or infection. Positive reactants include C-reactive protein, ferritin, fibrinogen, and hepcidin; negative reactants include albumin and transferrin.

#### Alternate Names:

Acute-Phase Proteins

### Acyclovir

An antiviral drug that inhibits herpesvirus DNA polymerase and is used for infections caused by herpes simplex virus and varicella-zoster virus.

### Ad Lib

A prescription or clinical instruction meaning as desired or as tolerated, from the Latin *ad libitum*.

### ADA Deficiency

A rare inherited disorder in which cells lack enough adenosine deaminase, an enzyme needed to break down certain cell chemicals. Toxic substances then build up and damage immune cells, often causing severe infections from early infancy.

### Adactyly

Absence of one or more fingers or toes from birth.

### Adam's Apple

The common name for the laryngeal prominence, the visible projection at the front of the neck formed by the angle where the two laminae of the thyroid cartilage meet.

### Adamantine

Very hard or resembling diamond; in dentistry, relating to tooth enamel or enamel-like tissue. The term may describe an unusually hard structure or the appearance of a lesion containing enamel-forming tissue.

### Adamantinoma

A rare, low-grade malignant primary bone neoplasm characterized by a distinctive biphasic histology comprising epithelial cells embedded within an osteofibrous stroma. It shows a dramatic anatomical

---

predilection for the anterior cortex of the mid-shaft (diaphysis) of the tibia (accounting for over 85% to 90% of cases), occasionally involving the adjacent fibula. Most commonly diagnosed in young adults aged 20 to 40 years, it presents as a slow-growing, painful pretibial swelling or pathological fracture. Plain radiographs characteristically show eccentric, multilocular, radiolucent cortical lesions with surrounding sclerosis producing a “soap-bubble” appearance. Because adamantinoma is resistant to chemotherapy and radiotherapy, standard treatment requires wide en bloc surgical resection with negative margins and bone reconstruction.

**Alternate Names:**

Adamantinoma of Long Bones; Tibial Adamantinoma; Malignant Angioblastoma of Bone

**Adams-Stokes Syndrome**

A clinical syndrome characterized by sudden, transient episodes of syncope or loss of consciousness caused by abrupt bradyarrhythmias, high-grade or complete atrioventricular block, or ventricular asystole leading to acute cerebral hypoperfusion.

**Alternate Names:**

Stokes-Adams Syndrome; Adams-Stokes Attack

**Adaptation**

Adaptation is the adjustment of a cell, organism, or person to a changing internal or external condition; in medicine it may involve physiologic compensation, behavior, or rehabilitation.

**Adaptive Design**

A clinical-study design allowing planned changes based on accumulating data while preserving scientific validity. Changes may involve sample size, treatment groups, or enrollment and must follow a prespecified plan.

**Adaptive Equipment**

Tools or devices modified to help a person perform daily activities safely despite disability, weakness, pain, or sensory loss. Examples include bathroom rails, reachers, adapted utensils, wheelchairs, and communication aids.

**Adaptive Immunity**

The part of the immune system that learns to recognize specific germs or abnormal cells. B cells make targeted antibodies, T cells coordinate or destroy affected cells, and memory cells help produce a faster response to a later exposure.

**ADD/ADHD**

Attention-deficit/hyperactivity disorder, a neurodevelopmental condition involving persistent inattention, hyperactivity, impulsivity, or combinations of these symptoms that impair functioning.

**Addiction**

A commonly used, non-diagnostic term for compulsive substance use or recurrent behavior despite harm, with impaired control, craving, or difficulty stopping. In clinical practice, the specific diagnosis is usually named as a substance-use disorder or a recognized disorder due to addictive behavior; physical dependence alone does not establish addiction.

**Addison Anemia**

An older name for pernicious anemia, a vitamin B12-deficiency megaloblastic anemia caused by inadequate intrinsic factor and impaired B12 absorption.

**Alternate Names:**

Addisonian Anemia; Addison’s Anemia

**Addison Disease**

Primary adrenal insufficiency caused by damage to the adrenal glands, so they produce too little cortisol and often too little aldosterone. It may cause fatigue, weight loss, low blood pressure, darkened skin, and adrenal crisis.

**Alternate Names:**

Addison’s Disease Chronic Adrenal Insufficiency Primary Adrenal Insufficiency

**Addison’s Disease**

Destruction or dysfunction of the adrenal cortex causing deficiency of cortisol and aldosterone, presenting with fatigue, hyperpigmentation, weight loss, and salt craving. Lifelong hormone replacement is required.

---

**Alternate Names:**

Hypoadrenocorticism; Chronic Adrenocortical Insufficiency

**Addisonian Anemia**

Alternate name; see *Addison Anemia*.

**Addisonian Crisis**

Life-threatening emergency resulting from acute cortisol deficiency in patients with adrenal insufficiency. Precipitated by infection, trauma, surgery, or abrupt discontinuation of chronic glucocorticoid therapy.

**Additive Genetic Effects**

The contributions of individual alleles that combine approximately additively to influence a measurable trait, without requiring interaction between those alleles.

**Adduct**

To move an arm, leg, finger, toe, or other body part toward the center line of the body.

**Adduction**

Movement of a body part toward the body's midline or, for a finger or toe, toward the hand or foot's central axis. For example, bringing an arm back toward the side is shoulder adduction.

**Adductor Muscle**

A muscle that moves a body part toward the body's midline or, for a digit, toward the axis of the hand or foot.

**Adductor Muscles**

A group of muscles on the inner thigh that pull the leg toward the body's midline and help stabilize the hip. Strain of these muscles is a common cause of groin pain, especially during sports.

**Adenine**

A purine nitrogenous nucleobase that forms complementary base pairs with thymine in DNA (via two hydrogen bonds) and with uracil in RNA. It also forms essential components of adenosine, cellular

energy carriers (ATP, ADP, AMP), second messengers (cAMP), and key metabolic cofactors (NAD<sup>+</sup>, FAD).

**Alternate Names:**

6-Aminopurine; Ade

**Adenitis**

Inflammation of a gland or lymph node, usually caused by infection or an immune disorder. It may cause swelling, tenderness, redness, or fever; treatment depends on the affected site and cause.

**Adenocarcinoma**

A cancer that begins in cells lining or forming glands. It can arise in organs such as the colon, breast, lung, pancreas, prostate, stomach, or uterus and may invade or spread.

**Adenocarcinoma In Situ Of Cervix**

A noninvasive glandular precursor lesion of the cervix, usually associated with high-risk human papillomavirus; abnormal glandular cells remain confined to the epithelium but may progress to invasive adenocarcinoma.

**Adenocarcinoma Of The Colon**

A malignant gland-forming epithelial tumor of the colon, commonly presenting with altered bowel habits, occult bleeding, anemia, or an obstructing mass.

**Adenocarcinoma Of The Lung**

The most common type of non-small-cell lung cancer. It begins in gland-forming cells, often toward the outer lung, and can occur in smokers or nonsmokers. Some tumors have gene changes, such as EGFR, ALK, or KRAS, that guide treatment choices.

**Alternate Names:**

Pulmonary Adenocarcinoma; Bronchial Adenocarcinoma

**Adenohypophysis**

The front part of the pituitary gland, a small hormone-producing organ at the base of the brain. It releases hormones that regulate growth, thyroid

---

activity, adrenal function, reproduction, milk production, and other body processes.

### Adenoid Basal Carcinoma

A rare, usually indolent cervical carcinoma composed of small basaloid nests, often associated with squamous intraepithelial lesions and requiring distinction from more aggressive tumors.

### Adenoid Cystic Carcinoma

A rare malignant cancer that arises in secretory glands, most commonly the major and minor salivary glands of the head and neck, as well as the breast, skin, and airways. It tends to grow slowly but is notable for tracking along nerve pathways (perineural invasion), causing pain or numbness, and recurring or spreading to the lungs years after initial treatment.

#### Alternate Names:

Cylindroma

### Adenoid Cystic Carcinoma Of The Vulva

A rare malignant tumor of vulvar Bartholin-type glands, often causing pain because of perineural invasion. It is usually slow-growing but may recur locally or metastasize after long intervals.

### Adenoid Hypertrophy

Enlargement of the adenoid tissue located in the nasopharynx behind the nasal cavity. Commonly occurring in children due to recurrent respiratory infections or allergies, it can obstruct nasal breathing, leading to persistent mouth breathing, loud snoring, obstructive sleep apnea, a nasal tone of voice, and recurrent middle-ear fluid accumulation (otitis media with effusion).

#### Alternate Names:

Enlarged Adenoids; Adenoidal Hyperplasia

### Adenoid Removal

Alternate name; see *Adenoidectomy*.

### Adenoidectomy

Surgical removal of the adenoids (pharyngeal tonsil) from the back of the nasal passage. It is commonly

performed in children to treat chronic nasal obstruction, pediatric obstructive sleep apnea, recurrent ear infections, or persistent rhinosinusitis that does not improve with medical therapy.

#### Alternate Names:

Adenoid Excision; Adenoid Removal

### Adenoiditis

Acute or chronic inflammation and infection of the adenoid tissue behind the nose. It causes persistent nasal congestion, purulent nasal discharge, mouth breathing, snoring, sore throat, and muffled speech.

### Adenoids

A collection of lymphatic tissue located in the upper airway at the back of the nasopharynx, forming part of Waldeyer's ring. In infants and young children, they help sample and trap inhaled germs to build immunity, but typically shrink in size during late childhood and adolescence.

#### Alternate Names:

Pharyngeal Tonsil; Nasopharyngeal Tonsil

### Adenolipoma

A benign tumor composed of glandular cells and mature fat cells. It may occur in the skin, breast, thyroid, parathyroid, or salivary glands.

### Adenolymphoma

A benign salivary-gland tumor, also called Warthin tumor, composed of oncocytic epithelium and lymphoid stroma and occurring most often in the parotid gland.

### Adenoma

A usually noncancerous gland-forming tumor arising from epithelial tissue. Some adenomas produce hormones or other secretions, and some—especially certain colon and rectal adenomas—contain precancerous changes and may later become cancerous.

### Adenoma Malignum

A rare, slow-looking but invasive cancer of the cervix that produces mucus. Persistent watery vaginal discharge or abnormal bleeding may occur. Diagnosis



---

requires examination of cervical tissue by an experienced pathologist because the cells may look almost normal under the microscope.

### Adenoma-Carcinoma Sequence

The usual step-by-step pattern in which some colon polyps develop genetic changes, become increasingly abnormal, and eventually turn into colon cancer over several years. Removing suitable polyps can interrupt this progression.

### Adenomas

Plural of adenoma; see *Adenoma*.

### Adenomatosis

A condition in which multiple gland-forming tumors called adenomas develop in one organ or tissue. The tumors are often benign, but their number and location may affect cancer risk.

### Adenomatous Polyp

A benign gland-forming polyp, most commonly in the colon, that may contain dysplasia and can progress through the adenoma-carcinoma sequence. Tubular, villous, and tubulovillous types are recognized.

### Adenomatous Polyposis

An autosomal dominant genetic disorder caused by germline mutations in the *APC* tumor suppressor gene, characterized by the progressive development of hundreds to thousands of adenomatous polyps throughout the colon and rectum, carrying a nearly 100 percent lifetime risk of colorectal cancer if left untreated.

#### Alternate Names:

Familial Adenomatous Polyposis; FAP

### Adenomyoepithelioma Of The Breast

A rare biphasic breast tumor composed of proliferating epithelial and myoepithelial cells. Most are benign, but local recurrence and malignant transformation can occur, so complete excision is the usual treatment.

### Adenomyoma

A benign uterine lesion composed of endometrial glands and stroma within a proliferation of smooth muscle; it may cause abnormal uterine bleeding or mimic other uterine tumors.

### Adenomyosis

A condition in which tissue normally lining the uterus grows into its muscular wall. It can enlarge the uterus and cause heavy or painful periods, ongoing pelvic pain, or no symptoms; it is more common during the later reproductive years.

### Adenopathy

Disease or enlargement of a gland, most often used for lymph-node enlargement or abnormal lymph-node involvement.

### Adenosine

A purine nucleoside made of adenine linked to ribose that participates in energy transfer and cell signaling. Given rapidly by injection, it can briefly slow conduction through the atrioventricular node and is used for selected supraventricular tachycardias.

### Adenosine Deaminase Deficiency

An inherited purine-metabolism disorder in which toxic deoxyadenosine metabolites accumulate and damage lymphocytes, commonly causing severe combined immunodeficiency.

### Adenosine Deaminase Severe Combined Immunodeficiency (ADA-SCID)

A rare inherited immune deficiency caused by adenosine deaminase deficiency. Toxic metabolites prevent normal development of lymphocytes, leaving infants highly vulnerable to severe infections.

### Adenosine Diphosphate

A nucleotide formed when ATP loses one phosphate group. ADP can be converted back to ATP to support cellular energy use, and released ADP also activates nearby platelets and promotes platelet aggregation during clot formation.

#### Alternate Names:

---

ADP

### Adenosine Monophosphate

A small molecule made of adenine, ribose, and one phosphate group. It is formed when cells break down or recycle energy-carrying molecules and helps regulate energy use and cell signaling.

#### Alternate Names:

AMP; Adenylic Acid

### Adenosine Triphosphate

The main short-term energy source inside cells. Cells make it from nutrients and use it to power muscle contraction, movement of substances across cell membranes, chemical reactions, and many other essential processes; it is commonly called ATP.

### Adenosis

An increase in the number or size of glands or gland-like tissue. It is often noncancerous, but its meaning depends on the organ involved, such as the breast or lung, and the specific pattern.

### Adenovirus

A group of DNA viruses that commonly cause respiratory illness, including cold-like symptoms, sore throat, fever, bronchitis, croup, or pneumonia. Depending on the type, adenoviruses can also cause conjunctivitis, gastroenteritis, cystitis, or rarely neurologic disease. They spread through respiratory secretions, close contact, contaminated surfaces, and fecal-oral exposure. Most infections are mild, but severe disease is more likely in infants, people with heart or lung disease, and people with weakened immune systems.

### Adenovirus DNA PCR (Feces)

A stool test that uses polymerase chain reaction (PCR) to detect adenovirus genetic material. It can help identify adenovirus in someone with a compatible illness; a positive result is interpreted with symptoms because viral DNA may sometimes be detected without illness.

#### Alternate Names:

Stool Adenovirus PCR; Adenovirus PCR, Stool

### Adenovirus Infection

Illness caused by an adenovirus, commonly with sore throat, cough, fever, pink eye, diarrhea, or vomiting, especially in children. The virus spreads through droplets, close contact, contaminated surfaces, or infected stool. One pattern, pharyngoconjunctival fever, brings sore throat, red eyes, and fever together and spreads easily in daycare or swimming-pool settings. Most infections clear on their own, but severe breathing or widespread disease can occur in infants, older adults, people with heart or lung disease, and people with weak immune systems. Treatment is supportive, with fluids and fever relief; severe illness with breathing difficulty or dehydration may require hospital care.

### Adenoviruses

Plural of adenovirus; see *Adenovirus*.

### Adenylate Cyclase

An enzyme that changes ATP into cyclic AMP, a chemical messenger inside cells. It is often activated by signals received at the cell surface and helps control many cell functions.

#### Alternate Names:

Adenylyl Cyclase

### Adequate Intake

A formal nutritional reference standard within the Dietary Reference Intakes (DRI) system. It represents the recommended average daily nutrient intake level based on observed or experimentally determined approximations in healthy populations, used when scientific evidence is insufficient to calculate an Estimated Average Requirement (EAR) and Recommended Dietary Allowance (RDA) (such as for vitamin K, potassium, or infant nutrition).

#### Alternate Names:

AI; Dietary Reference Intake: Adequate Intake

### ADHD in Children

Attention-deficit/hyperactivity disorder in childhood, characterized by developmentally inappropriate inattention, hyperactivity, or impulsivity across settings. Diagnosis requires persistent impairment and information from caregivers, teachers, and clin-

---

ical assessment.

### Adherence

The extent to which a person's behavior corresponds with agreed healthcare recommendations, including taking medications, following diets, or executing lifestyle changes. Non-adherence may stem from adverse effects, regimen complexity, cognitive impairment, financial constraints, or lack of support.

#### Alternate Names:

Compliance; Treatment Adherence; Medication Adherence

### Adherens Junction

A connection that helps neighboring cells stick together. It links proteins on one cell to the internal support fibers of another, helping tissues keep their shape, strength, and protective barrier.

### Adhesiolysis

A procedure that cuts or removes bands of scar tissue that abnormally join organs or body surfaces. It may relieve pain or release a blocked intestine, but the bands can form again after surgery.

### Adhesion

A band of scar-like tissue that abnormally joins internal surfaces or organs, often after surgery, infection, inflammation, or injury, sometimes causing pain or blockage.

### Adhesion Molecules

Proteins on cells that help cells attach to one another or to the material supporting them. They help organize tissues, guide immune cells to inflamed areas, and can influence how cancer spreads.

### Adhesive Capsulitis

Alternate name; see *Frozen Shoulder*.

### Adhesive Tape

A material used to secure dressings, devices, or tubing to the skin. It is available in paper, cloth, plastic, and other forms.

### Adie Tonic Pupil

A pupil that is unusually large and reacts slowly to light but constricts better when focusing on a near object. It usually affects one eye and results from injury to nerves controlling the pupil; the cause may be unknown or related to another disorder.

### Adipocere

A grayish-white, waxy, soap-like substance formed in decomposing bodies exposed to moisture and lack of oxygen (such as damp soil or submerged water). It is produced by the bacterial hydrolysis and hydrogenation of body fats into fatty acids (saponification). This process retards putrefaction and can preserve body contours, facial features, and traumatic injuries for months or years, assisting forensic identification.

#### Alternate Names:

Grave Wax; Saponification Wax; Mortuary Wax

### Adipocyte

A fat cell that stores energy and releases hormones and other signals affecting appetite, blood sugar, inflammation, and metabolism. White fat mainly stores energy, while brown and beige fat can produce heat; enlarged or excessive fat cells can contribute to insulin resistance.

### Adipocytes

Fat cells that store energy mainly as triglycerides and also release hormones and signaling molecules. White adipocytes primarily store energy, while brown and beige adipocytes can produce heat.

#### Alternate Names:

Fat Cell

### Adipogenesis

The process by which immature cells develop into fat cells. It helps form and maintain body-fat tissue, which stores energy and produces hormones; abnormal increases in the number or size of fat cells can contribute to obesity and metabolic disease.

### Adipokine

A chemical messenger released by body fat. Adipokines help control appetite, blood sugar, fat

---

use, inflammation, blood pressure, and immune activity; abnormal levels may contribute to obesity, diabetes, and heart or blood-vessel disease.

### **Adiponectin**

A hormone produced mainly by fat cells that helps regulate fatty-acid use, glucose metabolism, insulin sensitivity, and inflammation. Levels are often lower with obesity and insulin resistance, but a blood level is not diagnostic by itself.

### **Adipose**

Relating to body fat or composed of adipose tissue, which stores energy and also functions as an endocrine and immune-active organ.

### **Adipose Tissue**

A specialized connective tissue composed mainly of adipocytes that store energy as triglycerides. It provides thermal insulation, cushioning for organs, and endocrine regulation through hormones and signaling molecules such as leptin and adiponectin.

### **Adiposis Dolorosa**

Adiposis dolorosa, or Dercum disease, is a rare disorder characterized by painful fatty deposits, often with obesity, fatigue, and other systemic symptoms.

### **Adiposity**

The amount and location of body fat. Too much body fat, especially around the abdomen, can increase the risk of diabetes, heart disease, high blood pressure, and joint or movement problems.

### **Adipsia**

Little or no feeling of thirst despite the body needing fluid. It can lead to dehydration and dangerously high blood sodium, especially when caused by illness affecting the brain, medicines, aging, or severe illness.

### **Adjacent**

Next to, close to, or sharing a boundary with another structure, tissue, organ, or lesion. The term helps describe location, spread of disease, or rela-

tionships seen on imaging or examination.

### **Adjustment Disorder**

Emotional or behavioral symptoms in response to an identifiable stressor, beginning within three months of the stressor. The symptoms cause significant distress or impairment and do not meet criteria for another mental disorder. They generally do not persist for more than six months after the stressor or its consequences have ended.

### **Adjustment Disorder With Anxiety**

An emotional or behavioral response to an identifiable stressor that causes distress or impairment but does not meet criteria for another mental disorder.

### **Adjustment Disorders**

Alternate index label; see *Adjustment Disorder*.

### **Adjuvant**

An added treatment or substance that enhances the effect of a primary treatment; in vaccines, an adjuvant strengthens the immune response to an antigen.

### **Adjuvant Chemotherapy**

Systemic antineoplastic drug therapy administered following definitive primary treatment, such as complete surgical tumor resection, designed to eradicate occult micrometastatic disease and reduce the risk of cancer recurrence.

### **Adjuvant Therapy**

Additional treatment given after the main treatment to reduce the chance of disease returning. Examples include chemotherapy, radiation, hormone treatment, or immune treatment after surgery for selected cancers.

### **Adlerian Theory**

A psychological theory developed by Alfred Adler, also called individual psychology. It emphasizes feelings of inferiority, goal-directed striving, a person's style of life, and social interest—the capacity to belong to and contribute to the community. Early

---

family experiences and relationships are considered important influences on development.

**Alternate Names:**

Individual Psychology

**ADME**

The four processes that describe what the body does to a medicine: absorption into the bloodstream, distribution through tissues, chemical breakdown, and excretion from the body. These processes influence a medicine's dose, timing, and effects.

**Alternate Names:**

Pharmacokinetics

**Administered Activity**

The measured amount of radioactive material given to a patient for a scan or treatment. The amount is recorded with the radioactive substance, method, date, and time so the exposure and treatment can be checked safely.

**Administration Procedure**

A method used to give a medicine, fluid, vaccine, nutrition, or other treatment, such as by mouth, injection, inhalation, skin application, or infusion. The method affects speed, dose, and risks.

**Adnexa**

Structures next to an organ. In gynecology, the uterine adnexa are the ovaries, fallopian tubes, and supporting ligaments and tissues around the uterus.

**Adnexal Dysplasia**

Abnormal cell development in tissue beside the uterus, such as an ovary or fallopian tube. The change may be related to development or may represent a precancerous process, so its importance depends on the organ and tissue findings.

**Adnexal Mass**

A growth or enlargement involving structures beside the uterus, such as an ovary or fallopian tube.

**Adnexal Torsion**

Twisting of an ovary, fallopian tube, or both around

their supporting tissues. The twist can reduce or stop blood flow. It usually causes sudden one-sided pelvic pain, often with nausea or vomiting, and requires urgent medical assessment and usually surgery.

**Adnexal Tumors**

Tumors arising in structures beside the uterus, especially the ovaries, fallopian tubes, or supporting tissues. They may be benign, borderline, or malignant and are evaluated with examination, imaging, laboratory tests, and sometimes surgery or biopsy.

**Adnexal Tumors And Masses**

Abnormal growths or masses involving the ovary, fallopian tube, or nearby supporting tissues; they may be benign or malignant.

**Adnexitis**

Inflammation of one or more uterine adnexa, usually the fallopian tubes and sometimes the ovaries. It commonly occurs as part of pelvic inflammatory disease and may cause lower abdominal pain, fever, abnormal discharge, or pain during sex; untreated infection can lead to an abscess, infertility, or ectopic pregnancy.

**Alternate Names:**

Salpingo-Oophoritis

**Adolescence**

The developmental period between childhood and adulthood, involving physical maturation, puberty, and major psychological and social changes.

**Adolescent Development**

The physical, cognitive, emotional, and social changes that occur as a person develops from childhood into adulthood, including puberty and the development of greater independence.

**Adolescent Health**

The health and wellbeing of young people (10-19 years), addressing the physical, psychological, and social transitions of puberty and adolescence.



---

## Adolescent Medicine

A medical specialty focused on the physical, mental, reproductive, and social health needs of adolescents and young adults, including prevention, risk behavior, chronic disease, and confidentiality-sensitive care.

## Adolescent Obesity

Alternate index label; see *Pediatric Obesity*.

## Adolescent Schizophrenia

See *Childhood schizophrenia*.

## Adolescent Substance Use

Harmful or risky use of drugs by an adolescent, including nonmedical use of prescription medicines or use of illicit substances. It can impair brain development, increase overdose and injury risk, and signal a substance use disorder requiring confidential, age-appropriate care.

## Adoptive Cell Therapy

A personalized cellular immunotherapy in which a patient's own lymphocytes are expanded or engineered ex vivo and reinfused to attack the tumor; examples include tumor-infiltrating lymphocyte (TIL) therapy and chimeric antigen receptor (CAR) T-cell therapy.

## Adoptive Immunotherapy

Treatment in which immune cells are collected, activated or genetically modified, and returned to the patient to attack disease, especially cancer. Fever, inflammation, and other immune reactions may occur.

## ADP-Ribosylation

A chemical change in which a small molecule is attached to a protein or other cell molecule. This can switch the molecule's activity on or off and is used by normal cell processes and by some bacterial toxins.

## Adrenal Cancer

A rare cancer that begins in one of the adrenal

glands above the kidneys. Some tumors start in the outer cortex and others in the inner medulla; some produce excess hormones, while others cause pain or other symptoms as they grow.

## Adrenal Carcinoma

A rare malignant tumor arising from the adrenal cortex. It may produce excess cortisol, aldosterone, androgens, or estrogens, or present as an incidental mass or pressure symptom; hormonal evaluation and imaging help guide diagnosis and staging.

## Adrenal Cortex

The outer part of each adrenal gland that produces steroid hormones, including cortisol, aldosterone, and adrenal androgens. These hormones help regulate stress responses, blood pressure, salt balance, and metabolism.

## Adrenal Cortex Physiology

The outer layer of each adrenal gland makes hormones that control salt and blood pressure, stress responses, metabolism, and some sexual development. Its activity is regulated by the kidneys, blood potassium, and the brain's hormone-control system.

## Adrenal Cortical Carcinoma

Alternate name; see *Adrenocortical Carcinoma*.

## Adrenal Crisis

A sudden, life-threatening shortage of adrenal hormones, especially cortisol. It may occur with severe adrenal disease or after abruptly stopping long-term steroid treatment, particularly during infection, injury, or surgery. Symptoms can include severe weakness, vomiting, abdominal pain, confusion, low blood pressure, and collapse; emergency treatment is required.

## Adrenal Failure

Inadequate production of adrenal hormones, especially cortisol and sometimes aldosterone; severe acute deficiency can cause adrenal crisis with hypotension and shock.

---

## Adrenal Gland

One of two hormone-producing glands above the kidneys. The outer cortex makes aldosterone, cortisol, and adrenal androgens; the inner medulla makes adrenaline and noradrenaline. Adrenal hormones help regulate blood pressure, metabolism, stress responses, and salt balance.

## Adrenal Gland Cancer

Alternate name for adrenal cancer, a malignant tumor arising in an adrenal gland.

## Adrenal Gland Removal

Surgical excision of an adrenal gland, performed for tumors, hyperplasia, or other adrenal disorders. Approaches include open, laparoscopic, or retroperitoneal; perioperative management requires assessment of hormone production and potential adrenal insufficiency.

## Adrenal Gland Tumor

A growth in one of the adrenal glands above the kidneys. It may be noncancerous or cancerous and may or may not produce hormones; a pheochromocytoma is one possible type.

## Adrenal Glands

Adrenal glands sit above the kidneys and produce cortisol, aldosterone, adrenal androgens, and catecholamines that regulate stress, blood pressure, salt balance, and metabolism.

## Adrenal Hemorrhage

Bleeding into an adrenal gland, often associated with severe infection, serious illness, a bleeding disorder, blood-thinning medicines, or injury. Bleeding into both glands can cause sudden loss of adrenal hormones, very low blood pressure, and shock.

## Adrenal Hyperplasia

Enlargement or overgrowth of adrenal tissue that may increase production of adrenal hormones. Causes include congenital adrenal hyperplasia, chronic stimulation by ACTH, and some adrenal tumors. Symptoms depend on the hormone affected and may include abnormal genital development,

early puberty, excess body hair, high blood pressure, or electrolyte disturbances.

## Adrenal Incidentaloma

An unexpected mass found in an adrenal gland during imaging for another reason. It may be benign or cancerous and may or may not produce hormones, so evaluation usually includes imaging review and selected hormone tests.

## Adrenal Insufficiency

A condition in which the body does not make enough cortisol and sometimes not enough aldosterone. Primary adrenal insufficiency results from damage or dysfunction of the adrenal glands; secondary and tertiary forms result from inadequate pituitary or hypothalamic signals, including suppression after prolonged steroid treatment. Features may include fatigue, low blood pressure, weight loss, salt imbalance, or adrenal crisis.

## Adrenal Mass

An abnormal growth or enlargement in or near an adrenal gland. It may be benign or cancerous and may or may not produce hormones; evaluation usually involves imaging and, when appropriate, hormone tests and specialist review.

## Adrenal Medulla

The inner part of each adrenal gland, which sits above a kidney. It releases adrenaline and noradrenaline during stress, helping increase heart rate, blood pressure, alertness, and blood flow to muscles.

## Adrenal Medulla Physiology

The inner part of each adrenal gland releases adrenaline and noradrenaline when the body's stress-response nerves are activated. These hormones quickly increase alertness, heart activity, blood pressure, and blood flow to muscles.

## Adrenal Myelolipoma

A usually benign adrenal tumor composed of mature fat and blood-forming tissue. It is often found incidentally on imaging and causes no symptoms, but larger lesions may produce mass effect or bleeding.

---

Management depends on size, symptoms, growth, and diagnostic certainty.

### **Adrenal Nodule**

A localized lump in an adrenal gland. It may be a harmless nonfunctioning growth, a hormone-producing tumor, or rarely cancer; assessment depends on its size, imaging features, growth, and hormone production.

### **Adrenal Tumors**

Tumors arising in the adrenal cortex or medulla may be benign or malignant and may or may not produce hormones. Evaluation uses clinical assessment, hormone testing when indicated, and imaging to assess size and appearance; treatment depends on the diagnosis, hormone secretion, symptoms, growth, and risk of malignancy.

### **Adrenal Venous Sampling**

A catheter-based test that measures hormones in blood from each adrenal gland. It helps determine whether one gland is producing too much aldosterone, which can guide treatment for difficult-to-control high blood pressure.

### **Adrenalectomy**

Surgery to remove one or both adrenal glands, usually for a tumor or severe hormone overproduction. Removing one gland may leave enough hormone production; removing both requires lifelong hormone replacement and an emergency steroid plan.

### **Adrenaline**

Also called epinephrine, a hormone and neurotransmitter released during stress. It increases heart rate and contractility, raises blood pressure in many vascular beds, relaxes airway muscle, and is the first-line emergency medicine for anaphylaxis.

### **Adrenarche**

An early stage of sexual development, typically occurring around age six to eight, marked by increased production and secretion of weak adrenal androgens (such as dehydroepiandrosterone) by the zona reticularis of the adrenal cortex. It contributes to early

pubic hair growth, axillary odor, and oily skin.

### **Adrenergic Agent**

A medicine or substance that stimulates or blocks receptors responding to adrenaline (epinephrine) and noradrenaline (norepinephrine). These agents can change heart rate, blood pressure, airway size, and pupil size.

### **Adrenergic Agonist**

A medicine or substance that copies the effects of adrenaline or noradrenaline by activating specific cell receptors. Depending on the receptor, it may raise blood pressure, speed the heart, open the airways, or improve heart pumping.

#### **Alternate Names:**

Sympathomimetic

### **Adrenergic Antagonists**

Medicines or other substances that block adrenaline-like signaling at alpha or beta adrenergic receptors. They can lower blood pressure, slow the heart, reduce cardiac workload, or relax smooth muscle; effects and risks depend on the receptor targeted and the medicine used.

#### **Alternate Names:**

Adrenergic Blockers

### **Adrenergic Bronchodilator Overdose**

Excess exposure to a medicine that opens the airways by stimulating the body's stress-response system. It may cause shaking, agitation, headache, fast heartbeat, low potassium, chest pain, or abnormal rhythms and needs prompt medical assessment.

### **Adrenergic Receptor**

A cell-surface receptor activated by adrenaline (epinephrine), noradrenaline (norepinephrine), or related medicines. Alpha and beta receptors help control heart rate, blood pressure, airway size, pupil size, and other body responses.

### **Adrenergic Receptors**

Cell-surface receptors activated by adrenaline and noradrenaline. Alpha and beta subtypes regulate

---

cardiovascular, respiratory, metabolic, and other physiological responses.

### **Adrenocortical Carcinoma**

A rare cancer that begins in the outer layer, or cortex, of an adrenal gland, one of the glands above the kidneys. Some tumors are functioning and make too much of hormones such as cortisol, aldosterone, or adrenal androgens; others are nonfunctioning and make no excess hormones. Hormone-producing tumors may cause weight gain, high blood pressure, muscle weakness, unusual hair growth, or other changes in sexual development. A nonfunctioning tumor may cause no early symptoms, but a growing or spreading tumor may cause an abdominal lump, abdominal or back pain, or a feeling of fullness.

### **Adrenocorticotrophic Hormone**

A hormone made by the pituitary gland that tells the adrenal glands to make cortisol. Cortisol helps the body respond to stress and helps control blood pressure, blood sugar, and inflammation.

#### **Alternate Names:**

Adrenocorticotrophic Hormone (ACTH); ACTH; Corticotropin

### **Adrenogenital Syndrome**

A group of disorders, usually caused by congenital adrenal hyperplasia, in which excess adrenal androgens cause atypical genital development or early puberty.

### **Adrenoleukodystrophy**

A group of inherited disorders, most often X-linked, in which very-long-chain fatty acids accumulate and damage myelin, the adrenal glands, or both. Childhood cerebral disease can cause behavior and learning changes, seizures, and loss of neurologic function; adrenal insufficiency may also occur.

#### **Alternate Names:**

ALD

### **Adrenomedullin**

A hormone-like substance made by many tissues that widens blood vessels and helps regulate blood pressure, circulation, fluid balance, kidney activity,

and responses to injury.

### **Adrenomyeloneuropathy**

An adult-onset form of X-linked adrenoleukodystrophy that mainly affects the spinal cord and peripheral nerves. It may cause leg stiffness, weakness, walking difficulty, bladder or bowel problems, and sometimes adrenal insufficiency.

### **Adson Test**

A physical examination manoeuvre used to assess for thoracic outlet syndrome and compression of the subclavian artery by a cervical rib or scalene muscles. The examiner palpates the radial pulse while the patient abducts and extends the arm, extends the neck, and rotates the head toward the ipsilateral side with deep inspiration; diminution or obliteration of the radial pulse indicates a positive test.

#### **Alternate Names:**

Adson Manoeuvre; Adson's Test

### **Adsorption**

The sticking of molecules to the outside surface of a solid or liquid rather than entering its substance. Activated charcoal works partly by adsorbing some poisons in the digestive tract.

### **Adult Day Services**

Community-based daytime programs providing supervision, social activities, health-related support, or assistance with daily activities for adults who cannot safely remain alone throughout the day.

#### **Alternate Names:**

Adult Day Care; Adult Day Program

### **Adult Hypertension**

Blood pressure that stays too high over time. The exact level used for diagnosis depends on the guideline and the person's health. Untreated high blood pressure raises the risk of heart disease, stroke, and kidney disease.

#### **Alternate Names:**

Essential Hypertension; Primary Hypertension; Systemic Hypertension

---

## Adult Immunisation

The practice of administering vaccines to adults to protect against preventable infectious diseases, maintain waning immunity from childhood vaccinations, and reduce illness in high-risk groups. Essential adult vaccines include annual influenza, tetanus-diphtheria-pertussis (Td/Tdap) boosters every 10 years and during pregnancy, pneumococcal vaccines for adults aged 65 and older or those with chronic medical conditions, the recombinant herpes zoster vaccine for adults aged 50 and older to prevent shingles, and targeted vaccines for COVID-19, hepatitis B, HPV, and RSV.

### Alternate Names:

Adult Immunization; Adult Vaccination; Preventive Adult Vaccines

## Adult Informed Consent

The ethical and legal clinical communication process whereby a competent adult patient is provided with comprehensive, understandable information regarding the nature, indication, benefits, material risks, and alternatives of a proposed medical intervention or procedure, enabling an autonomous and voluntary decision.

### Alternate Names:

Informed Medical Consent; Surgical Consent; Patient Autonomous Decision-Making

## Adult Nutritional Support in Illness

Dietary strategies providing increased caloric and protein intake during acute or chronic illness, convalescence, or hypermetabolic states to prevent involuntary weight loss and muscle wasting.

### Alternate Names:

High-Calorie Diet in Illness; Nutritional Therapy in Recovery

## Adult Obstructive Sleep Apnea

Recurrent nocturnal collapse or partial obstruction of the pharyngeal airway in adults, producing repetitive episodes of hypopnea or apnea, transient arterial oxyhemoglobin desaturations, sympathetic surge, and fragmented sleep architecture leading to daytime somnolence and increased cardiovascular morbidity.

### Alternate Names:

Adult OSA; Obstructive Sleep Apnea Syndrome; Upper Airway Sleep Apnea

## Adult Primary Brain Tumor

A neoplasm arising de novo within the brain parenchyma, cranial nerves, or meninges in adults, encompassing diffuse gliomas (astrocytoma, oligodendroglioma, glioblastoma), meningiomas, vestibular schwannomas, and primary central nervous system lymphomas.

### Alternate Names:

Adult Primary Brain Tumor; Primary Intracranial Neoplasm; Adult Glioma

## Adult Spinal Deformity

Abnormal curvature or alignment of the spine in an adult, including scoliosis, kyphosis, or loss of normal sagittal balance. It may cause back pain, nerve compression, difficulty walking, or functional disability.

## Adult T-Cell Leukemia

A rare and aggressive peripheral T-cell neoplasm caused by infection with the human T-lymphotropic virus type 1 (HTLV-1). Characterized by generalized lymphadenopathy, hepatosplenomegaly, osteolytic bone lesions, severe hypercalcemia, skin infiltration, and atypical multi-lobed circulating lymphocytes known as flower cells.

### Alternate Names:

Adult T-Cell Leukemia-Lymphoma; ATLL

## Adult-Onset Asthma

Adult-onset asthma is variable airway inflammation and narrowing that begins in adulthood, producing episodic wheezing, cough, chest tightness, or shortness of breath. Diagnosis uses symptoms and lung-function testing, and treatment reduces triggers and airway inflammation.

### Alternate Names:

Late-Onset Asthma; Adult-Onset Reactive Airway Disease

## Adult-Onset Still Disease

A rare inflammatory disease that causes recurring



---

high fevers, joint pain or swelling, a salmon-colored rash, and sore throat. It may also inflame the liver, heart, lungs, or other organs.

### **Advance Directive**

A legal document stating a person's wishes about future medical care and, in some cases, naming a health-care proxy to make decisions if the person cannot decide or communicate. It may address resuscitation, life support, artificial nutrition, and other treatments and is reviewed as circumstances change.

#### **Alternate Names:**

Living Will; Medical Power of Attorney; Healthcare Directive

### **Advance Statements**

Alternate name; see *Living Will*.

### **Advanced Cancer**

Cancer that has grown beyond its original site. It may be locally advanced when it has invaded nearby tissues or metastatic when it has spread to distant organs.

### **Advanced Cardiovascular Life Support**

A standardized set of advanced resuscitation measures for cardiac arrest and other life-threatening cardiovascular emergencies. It includes high-quality CPR, defibrillation when indicated, airway management, medicines, and treatment of reversible causes.

#### **Alternate Names:**

ACLS

### **Advanced Glycation End-Products**

Compounds formed when sugar attaches to proteins or fats without the usual enzyme-controlled process. They increase with aging and persistently high blood sugar and may promote inflammation and damage to blood vessels, nerves, kidneys, and other tissues.

### **Advanced Prostate Cancer**

Prostate cancer that has grown beyond the prostate or spread to distant parts of the body. It may involve nearby tissues, lymph nodes, bones, or other

organs.

### **Advanced Sleep-Wake Phase Disorder**

A body-clock disorder in which a person becomes sleepy much earlier than desired and wakes unusually early. The person may sleep normally on this early schedule but cannot stay asleep until a more convenient time.

### **Adventitia**

The outer layer of connective tissue around a blood vessel or hollow organ. It supports and anchors the structure and contains small nerves and blood vessels that nourish the wall.

#### **Alternate Names:**

Tunica Adventitia

### **Adventitious**

Abnormal, unexpected, or arising from an unusual location or process; adventitious lung sounds include crackles, wheezes, and other added breath sounds.

### **Adverse Effect**

An unintended and harmful or bothersome effect of a medicine or other treatment given at normal doses or for an intended purpose.

### **Adverse Event**

Any new or worsening health problem that happens during or after a medicine, procedure, or other treatment, whether or not the treatment caused it. Clinicians record its timing, severity, and likely cause separately.

### **Adverse Reaction**

An unintended and harmful response to a medicine, treatment, or other exposure. It may be predictable from the medicine's effects, such as bleeding with an anticoagulant, or unpredictable, such as an allergic reaction. Severity ranges from mild nausea or rash to serious organ injury, breathing difficulty, or shock; suspected severe reactions need urgent medical attention.

#### **Alternate Names:**

Adverse Drug Reaction

---

## **Aedes Aegypti**

*Aedes aegypti* is a mosquito that can transmit dengue, yellow fever, chikungunya, Zika, and other viruses.

## **Aegophony**

An altered voice sound heard through a stethoscope when a person says “E” but it sounds like “A.” It can occur over lung tissue made denser by infection or fluid, but it does not identify the cause by itself.

## **Aerobe**

A microorganism that grows or produces energy using oxygen. Some aerobes cannot live without oxygen, while others can grow either with or without it. The distinction helps laboratories understand how an organism may behave in the body.

## **Aerobic**

Requiring or using oxygen. In microbiology, an aerobic organism grows with oxygen; in the body, aerobic energy production uses oxygen to release energy from nutrients for muscles and other cells.

## **Aerobic Bacteria**

Bacteria that need oxygen to grow. They may cause infection in tissues exposed to air, such as the lungs, skin, or surface of a wound; laboratory tests identify the specific species.

## **Aerobic Bacterium**

An aerobic bacterium grows using oxygen, although some species can tolerate or use alternative metabolic pathways; oxygen requirements help guide culture and infection assessment.

## **Aerobic Exercise**

Physical activity using large muscles continuously while the body produces energy with oxygen. Walking, cycling, swimming, and dancing can improve heart, lung, blood-vessel, muscle, mood, and endurance health.

### **Alternate Names:**

Cardio Exercise

## **Aerobiosis**

Life or biological activity that occurs when oxygen is available. Many human cells use oxygen to release energy from food, and some microorganisms need oxygen to grow.

## **Aerophagia**

Excessive swallowing of air, which may cause frequent belching, abdominal bloating, discomfort, or increased intestinal gas. It can occur with rapid eating, chewing gum, anxiety, or swallowing disorders.

## **Aerophagy**

Repeated swallowing of too much air, which can cause frequent belching, abdominal fullness, bloating, or discomfort. It may occur with rapid eating, chewing gum, anxiety, or difficulty swallowing.

### **Alternate Names:**

Aerophagia

## **Aerophobia**

An excessive fear of flying or air travel that causes marked anxiety, avoidance, or interference with daily life. It is a specific phobia and may improve with structured psychological treatment.

## **Aerosol**

Very small solid particles or liquid drops floating in a gas such as air. Aerosols can carry inhaled medicines, pollutants, or infectious particles and may remain in the air or settle on surfaces.

## **Aerospace Medicine**

The field of medicine concerned with maintaining the health, safety, and performance of people who travel or work in aircraft and spacecraft. It addresses effects such as changes in pressure, acceleration, low oxygen, radiation, isolation, and microgravity.

## **Aerotitis**

Ear pain or middle-ear injury caused by failure to equalize pressure during rapid changes in altitude, such as during aircraft ascent or descent.

---

## Aesthetic Dentistry

Dental care intended to improve the appearance of the teeth, gums, or smile. Procedures may include tooth-colored restorations, veneers, whitening, or selected crowns.

## Aesthetic Medicine

Medical care intended to improve appearance or reduce visible signs of aging through procedures, devices, medicines, or skin care; appropriate assessment includes risks, contraindications, and realistic expectations.

### Alternate Names:

Cosmetic Medicine; Aesthetic Dermatology

## Aetiology

The study of causation, or the specific originating cause or combination of pathogenic factors (such as genetic, environmental, infectious, or autoimmune mechanisms) that lead to the development of a disease or clinical disorder.

### Alternate Names:

Etiology; Causation; Pathogenesis

## Affect

The observable expression of a person's emotional state, including facial expression, voice, posture, and emotional responsiveness.

## Affective Disorder

Alternate name; see *Mood Disorders*.

### Alternate Names:

Mood Disorder

## Affective Flattening

Markedly reduced outward expression of emotion, including limited facial expression, voice modulation, and gestures. It may occur in schizophrenia and other psychiatric or neurologic conditions.

## Afferent

Carrying signals toward a central organ or structure; afferent nerves transmit sensory information from the body toward the brain and spinal cord.

## Afferent Arteriole

A small branch of the renal arterial circulation that conveys blood directly into the glomerular capillary tuft of an individual nephron, playing a pivotal role in regulating glomerular filtration pressure and renal blood flow through autoregulatory and tubuloglomerular mechanisms.

## Afferent Loop Syndrome

A blockage or poor emptying in a section of small intestine made during some stomach operations. Digestive juices build up behind the blockage, causing upper-abdominal pain, nausea, vomiting, or poor absorption of nutrients.

## Afferent Lymphatic Vessel

A lymphatic vessel that carries lymph into a lymph node.

## Afferent Neuron

A sensory nerve cell carrying signals from receptors in the body toward the spinal cord and brain, where information is processed and appropriate responses begin.

## Afferent Pathway

A sensory nerve route carrying information from the skin, muscles, organs, or other tissues toward the spinal cord or brain, where the signals are processed and responses can begin.

### Alternate Names:

Sensory Pathway

## Afferent Pupillary Defect

An asymmetry in pupillary light reactivity caused by a unilateral or asymmetric lesion in the anterior visual afferent pathway (optic nerve or retina). Identified during the swinging-flashlight test when swinging the light beam from the unaffected eye to the affected eye results in paradoxical bilateral pupillary dilation rather than constriction.

### Alternate Names:

Marcus Gunn Pupil; Relative Afferent Pupillary Defect; RAPD

---

## Affinity

The strength of attraction or binding between two molecules, such as an antibody and antigen, receptor and ligand, or enzyme and substrate.

## Affinity Chromatography

A laboratory method for separating and purifying a substance by allowing it to stick to a partner designed to recognize it. Other substances are washed away, and the desired substance is then released.

## AFib

Alternate name; see *Atrial Fibrillation*.

## Afibrinogenaemia

Alternate spelling; see *Afibrinogenemia*.

## Afibrinogenemia

A rare congenital or severe acquired coagulation disorder characterized by the complete or near-complete absence of circulating fibrinogen (factor I, <0.1 g/L). Inherited in an autosomal recessive pattern (mutations in *FGA*, *FGB*, or *FGG*), it manifests with umbilical cord bleeding in neonates, spontaneous mucosal and intracranial hemorrhages, hemarthroses, and prolonged bleeding times.

### Alternate Names:

Congenital Afibrinogenemia; Fibrinogen Deficiency; Afibrinogenaemia

## Aflatoxin

A toxin made by certain molds that can contaminate foods such as grains, nuts, and spices. Long-term exposure can damage the liver and increase the risk of liver cancer.

## African Sleeping Sickness

An infection caused by *Trypanosoma brucei* parasites and spread by infected tsetse flies in sub-Saharan Africa. It may cause fever and swollen lymph nodes, followed by sleep changes and nervous-system problems; untreated disease can be fatal.

## African Trypanosomiasis

An infection caused by *Trypanosoma* parasites and

spread by infected tsetse flies in sub-Saharan Africa. It may cause fever and swollen lymph nodes, followed by sleep changes and nervous-system problems; untreated disease can be fatal.

### Alternate Names:

African Sleeping Sickness

## After-Cataract

Clouding of the thin membrane behind the artificial lens after cataract surgery. It can develop months or years later and cause blurred or hazy vision, glare around lights, or reduced sharpness of vision.

### Alternate Names:

Posterior Capsule Opacification; Secondary Cataract

## Afterbirth

The placenta and the membranes that surrounded the baby, which leave the womb after childbirth. If any part remains inside, it can cause heavy bleeding or infection.

## Afterdepolarization

An abnormal electrical oscillation occurring during or immediately following myocardial action potential repolarization, capable of reaching threshold and triggering cardiac arrhythmias. Early afterdepolarizations occur during plateau or repolarization phases, while delayed afterdepolarizations occur after repolarization is complete.

## Afterload

The pressure or resistance the heart must overcome to push blood out of its ventricles. Higher blood pressure or narrowing of blood vessels increases the heart's workload and can make pumping less efficient.

## Afterpains

Cramp-like pains in the lower abdomen during the first few days after childbirth, caused by the womb tightening and shrinking. They may become stronger during breastfeeding and help reduce bleeding.

---

### Aftershave Poisoning

Harm from swallowing aftershave, which may contain alcohol, fragrance, or other chemicals. It can cause vomiting, drowsiness, low blood sugar, breathing problems, or coma; significant ingestion is an emergency requiring prompt poison-center assessment and medical treatment.

### Agammaglobulinaemia

A serious immune deficiency in which the blood contains very few or no antibodies, usually because antibody-producing B cells do not develop normally. It causes repeated bacterial infections, often beginning in infancy or early childhood.

### Agammaglobulinemia

A serious immune deficiency in which the blood contains very few or no antibodies. It often results from abnormal B-cell development and causes repeated bacterial infections of the ears, sinuses, lungs, or bloodstream.

### Aganglionic Megacolon

Congenital absence of nerve cells in part of the large intestine, causing Hirschsprung disease. The affected segment cannot relax normally, leading to intestinal obstruction, abdominal swelling, and delayed passage of stool.

### Agar

A jelly-like substance made from red seaweed. Laboratories add it to nutrient mixtures to grow and identify bacteria and fungi on solid surfaces, such as culture plates. It is also used in some bulk laxatives.

### Agatston Score

A number calculated from a special CT scan that measures calcium in the heart's arteries. A higher score usually means more coronary plaque and higher heart-disease risk, but it does not directly show blockage. Doctors interpret it with age, sex, blood pressure, cholesterol, diabetes, smoking, family history, and other health conditions.

#### Alternate Names:

Coronary Artery Calcium (CAC) Score; Agatston Calcium Score

### Age Regression

A mental state or behavior involving a temporary return to thoughts, feelings, behaviors, or mannerisms associated with an earlier stage of life. It may occur in dissociative disorders, hypnosis, trauma-related states, or some therapeutic contexts and is not itself a diagnosis.

### Age Spots

Flat, tan, brown, or black spots that develop on chronically sun-exposed areas of the skin, most commonly the face, backs of the hands, forearms, and shoulders. Caused by localized proliferation of pigment cells (melanocytes) from years of ultraviolet (UV) radiation exposure and aging, they are benign and harmless. Despite the colloquial name “liver spots,” they have no connection to liver function or liver disease.

#### Alternate Names:

Liver Spots; Solar Lentigines; Senile Lentigines; Sun Spots

### Age-Related Hearing Loss

Gradual hearing loss in both ears caused by aging. It often affects high-pitched sounds first and can make speech harder to understand.

### Age-Standardized Rate

A disease or death rate recalculated as if different populations had the same age distribution. This makes comparisons fairer when one population has more older or younger people than another.

### Ageing

Ageing is the natural, lifelong process of physical and biological change. Over time, body reserves may decrease and the risk of some diseases may increase, but the effects vary between people.

#### Alternate Names:

Aging

### Agenesis

Failure of an organ or body structure to form before birth, so it is absent. Agenesis may affect one organ, such as a kidney, or a specific body structure, such as the corpus callosum.



---

### **Agenesis of Commissura Magna Cerebri**

An older anatomic name for agenesis of the corpus callosum, in which the major commissural fiber tract connecting the cerebral hemispheres is absent or underdeveloped.

### **Agenesis Of The Corpus Callosum**

A birth condition in which the band of nerve fibres connecting the brain's two halves is partly or completely absent. Effects vary widely and may include seizures, developmental delay, learning difficulties, or no noticeable symptoms.

### **Ageusia**

Complete loss of the ability to taste. It may result from infections, mouth or nose problems, medicines, chemotherapy, head injury, nerve damage, nutritional deficiency, or neurological disease. Loss of smell can also make food seem tasteless.

### **Agger Nasi**

The most anterior and superior ethmoidal air cell, located immediately anterior to the attachment of the middle nasal turbinate and serving as an important anatomical landmark in functional endoscopic sinus surgery.

### **Agglutination**

Visible clumping that occurs when antibodies or other binding substances join cells or particles together. Laboratory workers use this reaction to help identify blood groups, infections, and other substances in patient samples.

### **Agglutination Test**

A laboratory test that mixes a patient sample with a known antibody or antigen. Visible clumping suggests that the matching substance is present and may help identify blood groups or infections.

### **Agglutinin**

An antibody or other substance that causes cells or particles to stick together in visible clumps. Agglutinins are used in laboratory tests, including blood-group testing and some infection tests.

### **Agglutinogen**

An antigen on a cell or particle that can react with a matching antibody and cause visible clumping. Agglutinogens are especially important in identifying blood groups.

### **Aggression**

Words or actions intended to threaten, harm, or intimidate another person or damage property. Aggression may result from emotional distress, pain, mental illness, brain disorders, medicines, or substance use.

### **Aggressive**

Describing hostile or harmful behavior, or a disease that grows, damages nearby tissue, or spreads quickly. The meaning depends on whether it describes a person's behavior or a medical condition.

### **Aggressive Angiomyxoma**

A rare, usually slow-growing soft-tissue tumor that most often develops in the vulva, perineum, or pelvis of women of reproductive age. It can grow into nearby tissue and often returns after treatment, but rarely spreads to distant organs.

### **Aggressive Cancer**

A cancer that is likely to grow, invade nearby tissue, or spread quickly. Doctors assess its aggressiveness using the cancer type, cell appearance, genetic features, stage, growth rate, and response to treatment.

### **Agility**

The ability to move, stop, or change direction quickly while staying balanced and controlled. Doctors and therapists may assess agility when evaluating movement, coordination, recovery from injury, or physical function.

### **Agitation**

An uncomfortable state of inner tension, restlessness, or increased activity. It may cause pacing, irritability, or inability to remain calm and can result from medical illness, pain, medicines, substance use, or mental health conditions.

---

## **Agnathia**

A rare birth condition in which the lower jaw is partly or completely absent. It may cause major facial abnormalities and serious problems with breathing, swallowing, and feeding.

## **Agnogenic Myeloid Metaplasia (AMM)**

An older name for primary myelofibrosis, a chronic bone-marrow disorder in which abnormal scar tissue and blood-cell production outside the marrow cause anemia, enlarged spleen, constitutional symptoms, and abnormal blood counts.

## **Agnosia**

An inability to recognize familiar objects, people, sounds, or other sensations even though the related sense works normally. It usually results from damage to areas of the brain that interpret sensory information.

## **Agonist**

A drug or other ligand that binds to a receptor and activates it, producing a biological response. A full agonist can produce the maximum response in a given system, whereas a partial agonist produces a submaximal response even when occupying receptors.

## **Agonist Muscle**

The main muscle responsible for producing a particular movement. It is also called the prime mover.

## **Agoraphobia**

An anxiety disorder involving persistent, intense fear of places or situations where escape may be difficult, help may be unavailable, or panic-like symptoms may feel embarrassing. People may avoid crowds, public transportation, shops, or leaving home alone; the fear and avoidance cause significant distress or interfere with daily functioning.

## **Agostini's Reaction**

A historical chemical test for glucose in urine. It used a gold-chloride reagent and an alkali; when heated with glucose, the mixture turned red. It is not a routine test in modern clinical practice.

## **Alternate Names:**

Agostini's Test

## **Agranulocytes**

A group of white blood cells that do not have prominent granules when viewed under a microscope. The group mainly includes lymphocytes and monocytes, which help defend the body against infection.

## **Agranulocytosis**

A serious condition in which the blood has an extremely low number of granulocytes, especially neutrophils, white blood cells that fight infection. It may be caused by medicines, chemotherapy, infection, autoimmune disease, or bone-marrow problems and can cause fever, sore throat, and severe infections.

## **Agraphesthesia**

An inability to recognize a number or letter traced on the skin with the eyes closed, despite normal basic touch sensation. It may indicate damage to the opposite parietal lobe of the brain.

## **Agraphia**

Loss of a previously learned ability to write because of brain damage or disease. It may follow a stroke, head injury, brain tumour, or neurodegenerative disease and may occur with problems speaking or reading.

## **Aicardi Syndrome**

A rare disorder affecting mainly girls, usually involving abnormal or absent tissue connecting the brain's two halves, distinctive retinal defects, and seizures beginning in infancy. Other brain and eye abnormalities may also occur.

## **Aicardi-Goutieres Syndrome**

A rare inherited disorder in which abnormal immune activity causes inflammation and damage mainly in the brain, skin, and immune system. It may begin in infancy with developmental regression, seizures, brain changes, and painful skin lesions.

---

## AIDS

The most advanced stage of HIV infection, when the immune system is severely weakened. AIDS is diagnosed when the CD4 count is below 200 cells/mm<sup>3</sup> or certain serious infections or cancers develop.

### Alternate Names:

Acquired Immunodeficiency Syndrome; Stage 3 HIV Infection

## AIDS Dementia

A severe form of HIV-associated brain disease occurring mainly in advanced HIV infection. It may cause worsening memory and thinking, slowed movement, difficulty walking, and changes in behaviour.

## AIDS-Associated Nephropathy

Kidney disease caused by HIV, usually in untreated or poorly controlled infection. It often causes large amounts of protein in the urine and may rapidly worsen kidney function, leading to kidney failure.

### Alternate Names:

HIV-Associated Nephropathy

## Ainhum

A rare disorder in which a painful fibrous band slowly tightens around a toe, usually the little toe. The band can damage bone and blood supply and may eventually cause loss of the affected part.

### Alternate Names:

Dactylolysis Spontanea

## Air Abrasion

A dental technique that uses compressed air to spray fine abrasive particles onto a tooth. It can remove limited decay, stains, or old filling material and prepare the tooth for a restoration.

## Air Bag

An inflatable vehicle safety device that rapidly expands during a crash to cushion impact and reduce serious injury. It works with, but does not replace, a properly worn seat belt.

## Air Bronchogram

A branching pattern of air-filled airways seen within

an area of lung that has lost its normal air, usually because the air sacs contain fluid, pus, blood, or cells. It is a sign on a chest X-ray or CT scan.

## Air Contaminants

Harmful substances in indoor or outdoor air, including smoke, dust, gases, chemicals, allergens, and germs. Health effects depend on the substance, amount breathed, duration of exposure, and a person's health.

## Air Embolism

A blockage of blood flow caused by air or another gas bubble entering a blood vessel. It can occur during medical procedures, injury, or rapid pressure changes and may cause sudden breathing, heart, or brain problems.

## Air Passages

The connected tubes that carry air from the nose and mouth to the lungs. They include the throat, voice box, windpipe, and branching airways; swelling, mucus, or an object can narrow or block them.

## Air Polishing

A dental procedure that directs air, water, and a fine powder onto teeth to remove plaque and surface stains. Common powders include glycine, erythritol, and sodium bicarbonate.

## Air Pollution

Air contaminated with harmful gases, particles, smoke, chemicals, or germs. Breathing polluted air can irritate the eyes and lungs and increase the risk of asthma attacks, infections, heart disease, stroke, and other health problems.

## Air Quality

A measure of how clean or polluted the air is, based on the amounts of harmful substances it contains. Poor air quality can affect breathing and heart health, especially in people with existing medical conditions.

---

## Air Sac

A small, thin-walled space that contains air. In the human lung, an air sac is called an alveolus and is where oxygen enters the blood and carbon dioxide leaves it; birds also have larger air sacs that help breathing.

### Alternate Names:

Air Space

## Air Trapping

A condition in which extra air remains in the lungs after breathing out because narrowed or damaged airways prevent complete exhalation. It commonly occurs in asthma, chronic obstructive pulmonary disease, and bronchiolitis.

## Air-Conditioner Lung

Lung inflammation caused by an immune reaction to mold, fungi, bacteria, or other particles released from contaminated air-conditioning or humidifying systems. Symptoms may include cough, shortness of breath, fever, and tiredness.

## Air-Sickness

Motion sickness caused by the movement of an aircraft. Mismatched signals from the eyes and inner ear can cause nausea, dizziness, sweating, headache, and sometimes vomiting.

## Airborne Disease

An infection that can spread when a person breathes in tiny particles containing germs that remain in the air. The risk depends on the germ, ventilation, distance, and length of exposure.

## Airborne Precautions

Measures used to prevent infections that remain suspended in the air from spreading. They include a special single-patient room with filtered exhaust air, limiting movement outside the room, and a properly fitted respirator such as an N95. They are used for tuberculosis, measles, and chickenpox.

## Airplane Ear

Ear pain, fullness, popping, or temporary hearing

loss caused by unequal air pressure on the two sides of the eardrum during takeoff or landing. It is more likely with a blocked nose or ear infection.

### Alternate Names:

Barotitis Media

## Airsickness

Motion sickness caused by travel through the air. Conflicting signals from the eyes and inner ear can produce nausea, dizziness, sweating, headache, and vomiting.

## Airway

Any passage that carries air between the outside of the body and the lungs, including the nose, mouth, throat, voice box, windpipe, and branching tubes. Swelling, mucus, or an object can narrow or block an airway and restrict breathing.

## Airway Clearance Techniques

Methods used to loosen and remove mucus from the breathing tubes. They include controlled coughing, breathing exercises, drainage positions, devices that create resistance during exhalation, and vibrating vests. The choice depends on the illness and the person's strength and ability.

## Airway Management

The methods and equipment used to keep a person's airway open and support breathing when airflow is threatened. They may include positioning, suction, airway devices, or a breathing tube connected to a ventilator.

## Airway Obstruction

Partial or complete blockage of airflow through the breathing passages. Causes include an inhaled object, mucus, swelling, injury, tumour, or relaxed airway muscles. It may cause noisy breathing, wheezing, low oxygen, severe breathing difficulty, or respiratory arrest.

## Airway Resistance

The difficulty air encounters while moving through the breathing tubes. Narrowing from muscle tightening, swelling, mucus, or loss of support increases

---

resistance and makes the breathing muscles work harder.

### **Akathisia**

A movement disorder causing an uncomfortable inner restlessness and a strong urge to move, often with pacing, rocking, or repeated shifting of the legs. It is commonly a side effect of antipsychotic or other medicines.

### **Akinesia**

An inability to initiate or perform voluntary movement, or a marked lack of spontaneous movement. It may occur in Parkinson disease and related disorders, severe depression, or other conditions affecting the brain's movement-control systems.

### **Akinetic**

Describing a person or condition with little or no voluntary movement. It may result from a neurological disease, medicine effect, severe illness, or damage to the brain's movement-control systems.

### **Akinetic Mutism**

A state in which a person is awake but makes little or no voluntary movement or speech. It usually results from severe damage or dysfunction in brain areas that control motivation, attention, and movement.

### **Akinetopsia**

A rare neuropsychological visual deficit characterized by an inability to perceive visual motion, resulting in moving objects appearing as a succession of static freeze-frames. It typically results from bilateral cortical lesions in the middle temporal visual area (area MT/V5).

#### **Alternate Names:**

Motion Blindness; Visual Motion Agnosia

### **Alagille Syndrome**

A genetic disorder in which too few or abnormal bile ducts can cause liver disease and jaundice. It may also affect the heart, kidneys, eyes, blood vessels, bones, and facial features. Most cases involve changes in the JAG1 gene.

### **Alanine**

An amino acid, or protein-building molecule, made by the body and obtained from food. Cells use it to make proteins and to help move nitrogen and produce glucose for energy.

### **Alanine Aminotransferase**

An enzyme found mainly inside liver cells. When liver cells are injured, alanine aminotransferase can leak into the blood, causing a raised test result. The result helps assess liver injury but does not identify its cause by itself.

#### **Alternate Names:**

ALT

### **Alanine Transaminase Blood Test**

A blood test that measures alanine aminotransferase, an enzyme released when liver cells are damaged. A high result may occur with hepatitis, medicine-related injury, fatty liver disease, or other conditions and must be interpreted with other information.

### **ALARA Principle**

A foundational radiation safety standard mandating that occupational and medical ionizing radiation exposures be kept as low as reasonably achievable, utilizing dose optimization, exposure time reduction, increased distance, and protective shielding.

#### **Alternate Names:**

As Low As Reasonably Achievable; ALARA; Radiation Safety Principle

### **Alarmin**

A signal released by injured or severely stressed cells that alerts the immune system. It helps start inflammation and attract protective cells to damaged tissue, even before the body identifies whether infection is present.

### **Albendazole**

An antiparasitic medicine that damages the internal structures parasites need to absorb nutrients and survive. It treats several worm infections, including some intestinal and tissue infections, when prescribed for the appropriate parasite.



---

### Albers-Schönberg Disease

A usually inherited form of osteopetrosis in which bones become unusually dense because old bone is not removed normally. The bones can still break easily and may cause pain or pressure on nerves, especially in the skull.

#### Alternate Names:

Autosomal Dominant Osteopetrosis Type II (ADO II); Marble Bone Disease

### Albinism

A group of inherited genetic conditions in which the body produces little or no melanin, the pigment responsible for skin, hair, and eye coloration. Characterized by pale skin, white or light blond hair, and light-colored irises, it frequently causes visual impairments—including reduced visual sharpness, rapid involuntary eye movements (nystagmus), extreme sensitivity to bright light (photophobia), and abnormal optic nerve development. People with albinism have an elevated risk of severe sunburn and skin cancer, requiring strict sun protection.

#### Alternate Names:

Oculocutaneous Albinism; OCA; Congenital Achromia; Hypopigmentation Disorder

### Albinismus

A genetic condition involving reduced or absent melanin production, causing lighter skin, hair, or eyes and often visual problems such as reduced acuity, nystagmus, and light sensitivity.

### Albino

An older noun for a person or animal with albinism, an inherited condition involving reduced melanin. For people, “person with albinism” is usually preferred because it describes the condition without defining the person by it.

### Albright Hereditary Osteodystrophy

A genetic condition that may cause short stature, a round face, obesity, dental problems, extra bone under the skin, and shortening of some finger bones. It is linked to changes in the *GNAS* gene and may occur with hormone-resistance disorders.

#### Alternate Names:

AHO; Pseudohypoparathyroidism Type 1A Phenotype

### Albright’s Hereditary Osteodystrophy

A group of inherited features that may include short stature, a round face, shortened fingers or toes, extra bone under the skin, and obesity. Some people also have resistance to parathyroid hormone or other hormones.

### Albumen

The white part of an egg, made mostly of water and proteins. Albumen is a food substance and is different from albumin, the main protein in blood plasma.

### Albumin

The main protein in blood plasma, made by the liver. It helps keep fluid inside blood vessels and carries substances such as hormones, fatty acids, and some medicines through the bloodstream.

#### Alternate Names:

Serum Albumin

### Albumin Clearance

A measure of how quickly the blood protein albumin leaves the bloodstream or another body space. It is used mainly in specialized kidney, liver, or research tests to study leakage, transport, or removal of the protein.

### Albumin-To-Creatinine Ratio

A urine test that estimates how much albumin, a blood protein, is leaking through the kidneys by comparing it with creatinine in the same sample. A persistent result above 30 mg/g may indicate kidney damage; results above 300 mg/g indicate severe leakage.

### Albuminocytologic Dissociation

A characteristic cerebrospinal fluid finding marked by an elevated protein concentration in the absence of a commensurate increase in white blood cell count. Classically seen in Guillain-Barré syndrome after the first week of illness, as well as in chronic inflammatory demyelinating polyneuropathy and spinal

---

cord tumors.

**Alternate Names:**

Albumino-Cytological Dissociation; Protein-Cell Dissociation

**Albuminuria**

An abnormally high amount of albumin in the urine, showing that the kidneys are allowing this blood protein to pass through. It may be temporary or persistent; ongoing albuminuria can indicate kidney damage and increased cardiovascular risk.

**ALCAPA Syndrome**

A rare congenital cardiovascular anomaly in which the left coronary artery arises abnormally from the pulmonary artery instead of the aorta. Because the pulmonary artery carries desaturated blood at low pressure, myocardial perfusion is compromised, leading to retrograde coronary flow (coronary steal), infantile heart failure, myocardial infarction, and mitral regurgitation if uncorrected.

**Alternate Names:**

ALCAPA; Bland-White-Garland Syndrome; Anomalous Left Coronary Artery from the Pulmonary Artery

**Alcohol**

Ethanol, a substance in beer, wine, and spirits that affects the central nervous system. Excessive consumption impairs motor coordination and cognitive judgment, produces acute intoxication, and can lead to physical dependence, metabolic toxicity, and chronic organ damage.

**Alcohol Dehydrogenase**

An enzyme found mainly in the liver that begins breaking down alcohol. It changes alcohol into acetaldehyde, a harmful intermediate that is then processed into less toxic substances.

**Alternate Names:**

ADH; Ethanol Dehydrogenase

**Alcohol Intolerance**

An unpleasant reaction after drinking alcohol, often caused by inherited difficulty breaking down alcohol or sensitivity to an ingredient in the drink. Sym-

toms may include flushing, headache, nausea, rapid heartbeat, or worsening asthma.

**Alcohol Intoxication**

Temporary impairment caused by drinking alcohol. It can cause poor judgment, confusion, slurred speech, loss of coordination, vomiting, reduced consciousness, slowed breathing, or coma.

**Alternate Names:**

Alcoholic Intoxication

**Alcohol Poisoning**

A life-threatening overdose of alcohol that severely depresses brain function and can cause vomiting, inability to wake, seizures, slow or irregular breathing, low body temperature, coma, or death.

**Alcohol Swab**

A single-use pad moistened with alcohol, usually used to clean intact skin before an injection or other procedure. The skin is allowed to dry before the procedure.

**Alcohol Use Disorder**

A medical condition in which repeated alcohol use causes difficulty controlling drinking, strong cravings, or problems at home, work, school, or in relationships. Symptoms may be mild, moderate, or severe.

**Alternate Names:**

Alcohol Abuse Alcohol Abuse and Alcoholism Alcohol Addiction Alcohol Dependence Alcohol Dependence (Alcohol Abuse and Alcoholism) Alcohol Dependency Alcoholism

**Alcohol Withdrawal**

Symptoms that can occur after a person who drinks heavily for a long time suddenly stops or greatly reduces alcohol. Symptoms include shaking, sweating, anxiety, nausea, hallucinations, seizures, and delirium; severe withdrawal can be life-threatening.

**Alternate Names:**

Alcohol Withdrawal Syndrome

---

### Alcohol-Associated Liver Disease

Liver injury associated with alcohol use. It ranges from steatosis and inflammation to fibrosis, cirrhosis, liver failure, or liver cancer; severity depends on alcohol exposure, genetics, nutrition, and other health factors.

### Alcohol-Related Psychosis

Hallucinations, delusions, or other psychotic symptoms associated with alcohol use, intoxication, withdrawal, or a coexisting alcohol-related disorder. Other medical and psychiatric causes must be excluded.

### Alcoholic Cardiomyopathy

A disease of the heart muscle caused by long-term heavy alcohol use. The heart becomes enlarged and weak, reducing its ability to pump blood and potentially causing heart failure or abnormal heart rhythms.

### Alcoholic Foetor

A characteristic odor on the breath and body resulting from recent ethanol consumption and the pulmonary excretion of volatile alcohol and its primary metabolite, acetaldehyde. It serves as an immediate clinical sign of acute alcohol exposure or intoxication during physical assessment.

#### Alternate Names:

Alcoholic Fetor; Alcohol Breath; Fetor Alcoholicus

### Alcoholic Hepatitis

An acute inflammation and injury of the liver caused by heavy alcohol use. It commonly causes jaundice, fatigue, fever, nausea, and abdominal pain and can become severe enough to cause liver failure.

#### Alternate Names:

Alcohol-Associated Hepatitis; Alcohol-Induced Hepatitis

### Alcoholic Ketoacidosis

A dangerous buildup of acids in the blood that can occur after prolonged heavy alcohol use, poor food intake, and repeated vomiting. It may cause abdominal pain, nausea, dehydration, rapid breathing, and a normal or low blood-sugar level.

### Alcoholic Liver Disease

Alternate name; see *Alcohol-Associated Liver Disease*.

### Alcoholic Myopathy

Muscle damage caused by heavy alcohol use. It may cause muscle pain, tenderness, weakness, or wasting; severe acute injury can destroy muscle tissue and damage the kidneys.

#### Alternate Names:

Alcohol-Induced Myopathy; Ethanol-Induced Myopathy

### Alcoholic Neuropathy

Nerve damage associated with long-term heavy alcohol use and related nutritional deficiency. It commonly causes burning pain, numbness, tingling, or weakness, especially in the feet and lower legs.

### Alcuronium

A medicine that temporarily relaxes or paralyzes skeletal muscles during anesthesia. Because it can stop the muscles needed for breathing, it is given with controlled airway and breathing support.

### ALD

Abbreviation commonly referring to adrenoleukodystrophy, an inherited disorder of fatty-acid metabolism that can damage the nervous system and adrenal glands.

### Aldolase Blood Test

A blood test that measures aldolase, an enzyme found mainly in muscle and liver. A high result may occur with muscle injury or disease, but it is not specific and is interpreted with symptoms and other tests.

### Aldosterone

A hormone made by the adrenal glands that helps the kidneys retain sodium and water and remove potassium. It helps control blood volume and blood pressure.

---

### **Aldosterone Blood Test**

A blood test that measures the hormone aldosterone. It helps investigate high or low blood pressure and abnormal potassium levels and is usually interpreted together with a renin test.

### **Aldosterone Escape**

A physiological phenomenon in primary hyperaldosteronism where persistent mineralocorticoid excess causes initial sodium and water retention, but subsequent increased atrial natriuretic peptide secretion and pressure natriuresis prevent progressive hypervolemia and severe edema.

### **Aldosterone-to-Renin Ratio**

A clinical screening blood test used in the evaluation of secondary hypertension to detect primary aldosteronism (Conn syndrome). A significantly elevated ratio of plasma aldosterone concentration to plasma renin activity prompts confirmatory testing.

**Alternate Names:**

ARR

### **Aldosteronism With Normal Blood Pressure**

A form of primary aldosteronism in which excess aldosterone is present without obvious hypertension. It may still cause low potassium, metabolic alkalosis, or cardiovascular and kidney risk and requires evaluation of the cause.

### **Aleukemic Leukemia**

A presentation of acute or chronic leukemia in which malignant blasts or abnormal leukocytes are absent or present in extremely low numbers in the peripheral blood smear, despite extensive malignant infiltration of the bone marrow.

### **Alexander Disease**

A rare genetic disorder in which abnormal GFAP protein damages the brain's supporting cells and white matter. It may begin in infancy, childhood, or adulthood and can cause enlarged head size, developmental problems, movement difficulty, seizures, or progressive weakness.

**Alternate Names:**

Fibrinoid Leukodystrophy; GFAP Leukodystrophy

### **Alexia**

Loss of the ability to understand written words despite adequate eyesight, usually because of brain damage from a stroke, injury, tumour, or degenerative disease. It may occur with other language or reading problems.

### **Alexia Without Agraphia**

A classical neurological disconnection syndrome in which a patient can write spontaneously and to dictation but is completely unable to read printed or handwritten text (including their own writing). Caused by an infarction of the left posterior cerebral artery affecting the left visual cortex and the splenium of the corpus callosum, isolating the intact right visual cortex from the language centers of the left hemisphere.

**Alternate Names:**

Pure Alexia; Dejerine Syndrome; Pure Word Blindness

### **Alexithymia**

Difficulty recognizing, understanding, or describing one's own emotions. It is a trait or symptom rather than a diagnosis and may occur with autism, depression, trauma, stress, or other conditions.

### **Alezzandrini Syndrome**

An exceptionally rare acquired disorder of unilateral melanocyte destruction involving ipsilateral facial vitiligo and poliosis with retinal or visual abnormalities and sometimes hearing loss. The cause is uncertain; ophthalmic and audiologic follow-up is important because vision or hearing may progressively worsen.

### **Alfacalcidol**

A medicine related to vitamin D that the liver changes into an active form. It increases absorption of calcium and phosphate and is used for selected disorders of bones, kidneys, or mineral balance.

### **Algia**

A word ending that means pain. It appears in

---

terms such as neuralgia for nerve pain, myalgia for muscle pain, and arthralgia for joint pain; it does not identify the cause.

### **Alglucerase**

An older enzyme-replacement medicine for Gaucher disease, a disorder in which fatty substances build up in cells. It replaces an enzyme the body lacks; newer enzyme preparations are used more often.

### **Algodystrophy**

Older term; see *Complex Regional Pain Syndrome*.

### **Algoneurodystrophy**

Older term; see *Complex Regional Pain Syndrome*.

### **Algor Mortis**

The gradual cooling of the body after death toward the surrounding temperature. The amount of cooling can help estimate the time since death, but it is affected by the environment and other conditions.

### **Aliasing Artifact**

An ultrasound error in which blood flow moving faster than the selected display range appears to change direction or speed. It reflects a limitation of the image display, not necessarily abnormal blood flow.

### **Alien Hand Syndrome**

A neurological condition in which an upper limb performs autonomous, purposeful-appearing actions without conscious voluntary control, often accompanied by the patient's sensation that the limb is foreign or acting independently. It is associated with lesions of the corpus callosum, supplementary motor area, or medial frontal cortex.

#### **Alternate Names:**

Alien Limb Phenomenon; Anarchic Hand

### **Alimentary**

Relating to eating, digestion, or the movement of food through the body. In medical language, it usually refers to the digestive tract and the organs that help break down and absorb food.

### **Alimentary Canal**

The continuous muscular tube through which food passes from the mouth to the anus. It includes the throat, oesophagus, stomach, small intestine, and large intestine.

### **Alkalemia**

An abnormally elevated arterial blood pH (greater than 7.45), indicating a decreased hydrogen ion concentration in circulating blood. It represents the net systemic state caused by underlying metabolic or respiratory alkalosis.

### **Alkali**

A basic substance that can neutralize acids. Strong alkalis, such as some cleaning chemicals, can burn the skin, eyes, and digestive tract, while weaker alkalis have food, laboratory, or medical uses.

### **Alkaline Diet**

A diet based mainly on foods claimed to make the body more alkaline. The lungs and kidneys tightly control blood acidity, so food does not make the blood alkaline, although many such diets include nutritious foods.

### **Alkaline Phosphatase**

An enzyme found mainly in the liver, bile ducts, bones, intestine, and placenta. A high blood level may reflect liver or bile-duct disease, increased bone activity, pregnancy, or another condition and must be interpreted with other tests.

### **Alkaline Phosphatase Test**

A quantitative serum laboratory assay measuring alkaline phosphatase activity to evaluate suspected cholestatic liver disease, biliary obstruction, or increased bone turnover such as Paget disease or bone metastases.

#### **Alternate Names:**

ALP Blood Test; Serum ALP; Alkaline Phosphatase Level

### **Alkaloid**

A naturally occurring chemical containing nitrogen,



---

often made by plants, that can strongly affect the body. Examples include morphine, quinine, nicotine, and atropine.

**Alternate Names:**

Plant Alkaloid; Basic Nitrogenous Organic Compound

**Alkalosis**

A condition in which the blood becomes too alkaline. It may result from repeated vomiting, certain medicines, hormone disorders, or breathing too rapidly, and can cause muscle cramps, tingling, confusion, or abnormal heart rhythms.

**Alkaptonuria**

A rare inherited metabolic disorder caused by reduced activity of homogentisate 1,2-dioxygenase, usually because of disease-causing variants in the *HGD* gene. Homogentisic acid accumulates, darkens urine, and can deposit in cartilage, tendons, skin, and other tissues in a process called ochronosis. Over time it may cause arthritis and stiffness of the spine and large joints, urinary stones, or changes involving heart valves.

**Alkylating Agents**

Drugs or chemicals that damage DNA by attaching chemical groups to it. Some are used to treat cancer by stopping cells from multiplying, but they can also injure healthy cells and cause side effects.

**Allantoin**

A compound used in some skin products to soften, moisturize, and protect irritated or rough skin. Its effect and safety depend on the product and how it is used.

**Allantois**

A temporary structure in the early embryo that helps with blood-vessel development and contributes to the umbilical cord and part of the urinary system.

**Allele**

One of two or more versions of a gene. A person usually inherits one allele from each parent; different alleles can influence traits or the risk of inherited

disease.

**Allelic**

Relating to an allele, one of the different versions of a gene inherited from a parent.

**Allen Test**

A bedside test that checks whether blood can reach the hand through both of its main arteries. It may be done before taking blood from or placing a catheter in the radial artery.

**Allergen**

A usually harmless substance that triggers an allergic immune response in a sensitive person. Common allergens include foods, pollen, medicines, animal proteins, insect venom, and natural rubber.

**Allergen Immunotherapy**

Treatment that gives gradually increasing amounts of a known allergen by injections or under the tongue. Over time, it can reduce allergic symptoms for selected allergies, but it can rarely cause a serious reaction.

**Allergic and Environmental Asthma**

Asthma triggered or worsened by allergens or environmental exposures such as dust mites, molds, pollens, animals, smoke, air pollution, or workplace agents. Management combines exposure reduction, controller therapy, trigger planning, and evaluation for occupational or allergic disease when indicated.

**Allergic Bronchopulmonary Aspergillosis**

An allergic reaction to the fungus *Aspergillus* in the airways, occurring mainly in people with asthma or cystic fibrosis. It can cause worsening breathing symptoms, mucus plugs, airway widening, and high levels of allergy-related cells or antibodies.

**Allergic Cascade**

The chain of immune events that follows exposure to an allergen. Immune cells release substances such as histamine, which can cause itching, swelling, mucus, wheezing, hives, or anaphylaxis.

---

## Allergic Conjunctivitis

Alternate name; see *Eye Allergies*.

## Allergic Contact Dermatitis

A delayed type-IV hypersensitivity reaction after skin contact with an allergen to which the immune system has been sensitized. It causes itching, redness, swelling, blisters, scaling, or chronic fissuring, often extending beyond the contact area; patch testing can help identify the trigger.

## Allergic Dermatitis

Skin inflammation caused by an immune reaction to an allergen. It commonly causes itching, redness, swelling, blisters, or scaling and may occur after contact with a triggering substance.

## Allergic Diseases

Conditions in which the immune system reacts abnormally to a usually harmless substance called an allergen, such as pollen, dust mites, mold, food proteins, medicines, insect venom, or animal proteins. Some reactions involve immunoglobulin E (IgE) antibodies, while others do not. Allergic diseases may affect the nose, eyes, skin, lungs, or digestive system and can cause sneezing, itching, hives, swelling, vomiting, wheezing, or anaphylaxis. Examples include allergic rhinitis, food allergy, drug allergy, allergic conjunctivitis, atopic dermatitis, urticaria, and allergic asthma. Food intolerance is different because it does not involve an immune reaction.

## Allergic Proctocolitis

A benign, non-IgE-mediated food protein-induced inflammatory disorder of the distal colon and rectum, seen predominantly in young infants exposed to cow's milk or soy protein in formula or breast milk, classically causing blood- and mucus-streaked stools in an otherwise thriving baby.

### Alternate Names:

Food Protein-Induced Allergic Proctocolitis; FPIP

## Allergic Reaction

An immune response to a usually harmless substance. It may cause sneezing, itching, hives, swelling, vomiting, or breathing difficulty and can

sometimes become life-threatening anaphylaxis.

### Alternate Names:

Allergy

### Alternate Names:

Allergic Reactions

## Allergic Rhinitis

Inflammation inside the nose caused by an allergic reaction to inhaled substances such as pollen, dust mites, mold, or animal proteins. It causes sneezing, itching, a blocked or runny nose, and sometimes itchy, watery eyes.

### Alternate Names:

Hay Fever; Pollinosis; Allergic Rhinoconjunctivitis

## Allergic Urticaria

An acute, IgE-mediated type I hypersensitivity skin reaction triggered by specific external allergens such as foods, drugs, insect venoms, or latex. Cross-linking of allergen-specific IgE on the surface of dermal mast cells triggers rapid degranulation and release of histamine, prostaglandins, and leukotrienes, causing localized vasodilation and increased vascular permeability that manifests as intensely pruritic, transient, erythematous wheals that resolve within 24 hours.

### Alternate Names:

Acute Allergic Urticaria; IgE-Mediated Urticaria; Allergic Hives

## Allergies to Insect Stings

Allergic reactions to venom from bees, wasps, hornets, or fire ants. Severe reactions can cause hives, throat swelling, wheezing, low blood pressure, or collapse and require immediate epinephrine and emergency care.

## Allergy

An immune response to a usually harmless substance such as pollen, food, medicine, insect venom, or animal proteins. Symptoms range from sneezing, itching, and hives to swelling, breathing difficulty, or life-threatening anaphylaxis.

### Alternate Names:

Allergic Hypersensitivity; Type I Hypersensitivity

---

## Allergy Shots

Injections used for allergen immunotherapy. They contain gradually increasing amounts of a specific allergen to reduce sensitivity over time and are given in a setting where a serious allergic reaction can be recognized and treated.

## Allergy Skin Test

An in vivo diagnostic procedure in which small amounts of suspected allergens are introduced into or onto the skin via prick, puncture, scratch, or intradermal injection to assess immediate IgE-mediated hypersensitivity through localized wheal-and-flare reactions.

### Alternate Names:

Skin Prick Test; Epicutaneous Test; Allergy Skin Testing; Scratch Test

## Allergy Test

A skin or blood test used to look for sensitivity to a suspected allergen. A positive result shows sensitization but does not always prove that exposure causes symptoms, so the result must be considered with the person's history.

### Alternate Names:

Skin Prick Test; Specific IgE Test; Allergen Challenge

## Allergy to Natural Rubber (Latex)

An immune reaction to natural rubber latex that may cause contact dermatitis, hives, wheezing, or anaphylaxis. Avoidance and clearly documented latex precautions are important for people with significant reactions.

## Allis Clamp

A surgical instrument with small teeth used to hold firm tissue during an operation. It provides a strong grip but can injure delicate tissue if applied with too much pressure.

## Allocheiria

A neurological sign in which a person feels or identifies a touch on the opposite side of the body from where it was applied. It reflects impaired localization of sensation and may occur with disorders

affecting the parietal lobe.

### Alternate Names:

Allochiria

## Allodynia

Pain caused by something that normally should not hurt, such as light touch, clothing, or gentle pressure. It can occur with nerve injury, migraine, shingles, fibromyalgia, or other pain conditions.

## Allogeneic

Coming from another person of the same species who is genetically different, as in an allogeneic blood, tissue, stem-cell, or organ transplant.

## Allograft

An organ, tissue, or group of cells transplanted from one genetically different individual to another individual of the same species. The recipient's immune system may reject it, so immunosuppressive medicines may be needed.

## Allosteric Regulation

Control of a protein or enzyme when another molecule attaches away from its main working site. The attachment changes the protein's shape and can increase or decrease how well it works.

## AloBar Holoprosencephaly

A severe form of holoprosencephaly in which the developing forebrain fails to divide and the cerebral hemispheres remain largely fused. It is usually associated with profound developmental impairment and major facial or other congenital abnormalities.

## Aloe

A plant used in some skin products, medicines, and supplements. Aloe gel applied to skin may soothe minor irritation, while products containing aloe latex taken by mouth can cause cramps, diarrhoea, and dangerous fluid or salt loss.

## Aloe Vera

A plant whose clear inner-leaf gel is used in some skin products and supplements. The gel may soothe

---

minor irritation, while oral latex or whole-leaf products can cause diarrhoea, fluid or salt loss, medicine interactions, and toxicity.

### **Alogia**

Markedly reduced speech or speech that conveys little information. It may occur as a negative symptom of schizophrenia and can also be associated with depression, brain disease, or other conditions.

### **Alongshan Virus**

A virus spread by some ticks and reported mainly in parts of Asia and Europe. It may cause fever, headache, fatigue, nausea, or neurological symptoms, but its ability to cause human disease is still being studied.

### **Alopecia**

Hair loss from the scalp or another part of the body. It may be nonscarring, in which the hair follicles remain and hair may regrow, or scarring, in which inflammation or injury destroys the follicles and can cause permanent hair loss. Causes include inherited pattern hair loss, autoimmune disease, hormonal changes, illness, medicines, nutritional problems, infection, repeated pulling, or other injury. The amount, pattern, and speed of hair loss vary with the cause.

### **Alopecia Areata**

An autoimmune, usually nonscarring hair-loss disorder in which immune cells attack anagen hair follicles. It often causes sharply demarcated smooth patches with exclamation-mark hairs and may involve the scalp, beard, eyebrows, or nails; regrowth is possible but relapses and extensive forms can occur.

#### **Alternate Names:**

AA; Patchy Hair Loss; Autoimmune Alopecia

### **Alopecia Celsi**

An older name for alopecia areata, an autoimmune disorder causing sharply defined patches of nonscarring hair loss.

### **Alopecia Cicatrisata**

Older term; see *Scarring Alopecia*.

### **Alopecia Circumscripta**

An older name for alopecia areata, characterized by one or more circumscribed patches of nonscarring hair loss.

### **Alopecia Mucinosa (Follicular Mucinosi)**

A skin disorder in which mucin accumulates within hair follicles and sebaceous glands, causing itchy follicular papules or plaques with hair loss. It may be a primary self-limited condition or a cutaneous sign of lymphoma, especially mycosis fungoides, so persistent or widespread disease needs evaluation.

### **Alopecia Totalis**

Complete loss of hair from the scalp, usually caused by alopecia areata. Hair may regrow, remain absent, or fall out again, and other autoimmune conditions may occur.

### **Alopecia Universalis**

Complete or near-complete loss of hair from the scalp and body, usually caused by extensive alopecia areata. Hair may regrow or recur unpredictably.

### **ALP Blood Test**

Alternate index label; see *Alkaline Phosphatase Test*.

### **ALP Isoenzyme Test**

A blood test that helps identify whether a high alkaline phosphatase level comes mainly from the liver, bones, intestine, or placenta. It can help distinguish liver or bile-duct disease from increased bone activity.

### **Alpha Blockers**

Medicines that relax certain blood vessels and smooth muscles by blocking alpha receptors. They may lower blood pressure or improve urine flow in an enlarged prostate, but can cause dizziness or fainting.

---

## Alpha Cells

Cells in the pancreas that make glucagon. Glucagon raises blood glucose when it falls too low or when the body needs energy between meals.

### Alternate Names:

Pancreatic Alpha Cells; Glucagon-Secreting Cells

## Alpha Hydroxy Acids

A group of acids used in skin products to loosen the outer layer of skin and improve roughness, fine lines, or uneven colour. Common examples are glycolic acid and lactic acid; they can irritate skin and increase sun sensitivity.

### Alternate Names:

AHAs; Fruit Acids

## Alpha Waves

A regular pattern of electrical activity in the brain, usually eight to thirteen cycles per second. It is strongest when a relaxed, awake person has closed eyes and decreases with attention or eye opening.

### Alternate Names:

Alpha Rhythm; Posterior Dominant Rhythm

## Alpha-1

A shortened name commonly used for alpha-1 antitrypsin deficiency or alpha-1 antitrypsin testing; the intended meaning should be confirmed from context.

## Alpha-1 Adrenergic Antagonist

A medicine or other substance that blocks alpha-1 adrenergic receptors. This reduces selected effects of noradrenaline on vascular smooth muscle and other tissues; the precise effects depend on the receptor distribution and the substance used.

### Alternate Names:

Alpha-1 Adrenergic Blocker

## Alpha-1 Antitrypsin Blood Test

A blood test that measures alpha-1 antitrypsin, a protein made by the liver that helps protect lung tissue. A low level may suggest inherited alpha-1 antitrypsin deficiency, which can affect the lungs and liver.

## Alpha-1 Antitrypsin Deficiency

An inherited condition in which low or abnormal alpha-1 antitrypsin increases the risk of emphysema, especially at a young age, and liver disease. Some people also develop inflammation of fatty tissue under the skin.

### Alternate Names:

AATD; AAT Deficiency; Alpha-1 Deficiency

## Alpha-Agonists

Medicines that activate alpha receptors on cells. Depending on the medicine and tissue, they can narrow blood vessels, raise blood pressure, reduce eye fluid, or affect the nervous system.

## Alpha-Fetoprotein

A protein made mainly by a developing fetus. Measuring it in maternal blood can help screen for some fetal conditions, while measuring it in adults can help assess selected liver or germ-cell tumours. An abnormal result does not establish a diagnosis.

## Alpha-Gal Syndrome

An allergy to a sugar found in most mammals but not humans or other primates. It can cause delayed hives, stomach symptoms, or anaphylaxis after eating mammalian meat or using some animal-derived products.

### Alternate Names:

Alpha-Gal Allergy

## Alpha-Globulin

One of several groups of proteins in the liquid part of blood. These proteins carry substances and help with inflammation, clotting, and immune defence; their levels may change with liver disease, inflammation, or other conditions.

## Alpha-Glucosidase Inhibitors

Diabetes medicines that slow the digestion and absorption of carbohydrates in the small intestine. They mainly reduce the rise in blood glucose after meals and may cause gas, bloating, or diarrhoea.



---

### Alpha-Ketoglutarate

A substance made during the breakdown of nutrients for energy. It also helps the body process amino acids and supports several chemical reactions inside cells.

### Alpha-Synuclein

A protein found mainly in nerve cells that helps regulate nerve-cell functions. Abnormal forms can collect into clumps in Parkinson disease, Lewy-body dementia, and related disorders, although the clumps are not the only cause of these diseases.

### Alpha-Thalassemia

An inherited blood disorder, usually autosomal recessive, in which the body makes too little alpha-globin, a part of hemoglobin that carries oxygen. The condition may cause no symptoms, mild anemia, or severe anemia before or shortly after birth, depending on the number of affected alpha-globin genes.

#### Alternate Names:

Alpha-Globin Deficiency; HbH Disease Carrier

### Alpha1-Antitrypsin (AAT) Deficiency

An inherited deficiency or dysfunction of alpha-1 antitrypsin that predisposes to early emphysema, especially with smoking, and sometimes liver disease from abnormal protein accumulation. Testing is considered in early or unexplained COPD, basilar emphysema, or unexplained liver disease.

### Alport Syndrome

An inherited disorder of collagen that can cause progressive kidney disease, hearing loss, and changes in the eyes. It may be caused by changes in the *COL4A3*, *COL4A4*, or *COL4A5* genes and may follow X-linked, autosomal dominant, or autosomal recessive inheritance.

### Alprazolam

A benzodiazepine medicine used for anxiety and panic disorder. It calms brain activity but can cause sleepiness, impaired coordination, dependence, and dangerous slowing of breathing, especially with opioids or alcohol.

### Alstrom Syndrome

A rare inherited disorder caused by changes in the *ALMS1* gene. It may cause progressive vision and hearing loss, obesity, diabetes or insulin resistance, heart disease, and kidney or liver problems.

#### Alternate Names:

Alström Syndrome

### Alternating Hemiplegia Of Childhood

A rare neurological disorder beginning in infancy or early childhood, causing recurrent episodes of temporary paralysis that may affect either side of the body. Episodes may also involve abnormal movements, eye-movement problems, breathing difficulty, seizures, and developmental impairment; symptoms typically disappear during sleep.

### Altitude Headache

Headache related to ascent to high altitude, often as part of acute mountain sickness. It may occur with nausea, dizziness, fatigue, or sleep disturbance and can signal more serious altitude illness when severe or accompanied by neurologic or breathing symptoms.

### Altitude Illness

Illness caused by rapid ascent to high altitude without enough time to adjust. It includes acute mountain sickness, high-altitude brain swelling, and high-altitude lung swelling; severe forms can be fatal.

#### Alternate Names:

Altitude Sickness

### Aluminum Phosphide

A highly poisonous pesticide that releases phosphine gas when it contacts moisture or stomach acid. Poisoning can rapidly damage the heart, lungs, digestive system, and other organs and requires emergency treatment.

### Aluminum Toxicity

Toxicity from excessive aluminum exposure or accumulation, especially in advanced kidney disease or with contaminated dialysis products. Chronic exposure can cause encephalopathy, speech or movement changes, microcytic anemia, and bone pain or

---

osteomalacia; evaluation includes exposure review, kidney function, and specialist-directed aluminum testing.

### **Alvarado Score**

A clinical score used to estimate the likelihood of acute appendicitis from symptoms, examination findings, and blood-test results. It supports assessment but does not by itself confirm or exclude appendicitis.

### **Alveolar**

Relating to an alveolus, especially one of the tiny air sacs in the lungs where oxygen enters the blood and carbon dioxide leaves it.

### **Alveolar Abnormalities**

Changes in the lung air sacs that can interfere with oxygen and carbon-dioxide exchange. They may occur with infection, fluid buildup, emphysema, scarring, or a developmental condition.

### **Alveolar Bone**

The part of the jawbone that surrounds and supports the roots of the teeth. It can shrink after tooth loss and may need rebuilding before some dental implants.

### **Alveolar Bone Loss**

Loss of the bone that supports the teeth, most often caused by periodontitis. It can make teeth loose and may lead to tooth loss; dental treatment can slow further loss.

### **Alveolar Dead Space**

The volume of inhaled air that reaches alveoli that are adequately ventilated but lack sufficient pulmonary capillary blood perfusion, preventing gas exchange from taking place.

### **Alveolar Echinococcosis**

A rare, serious infection caused by the young form of the *Echinococcus multilocularis* tapeworm, usually after swallowing contaminated food or water. It creates slowly growing, tumor-like areas in the liver

that can invade nearby tissue or spread.

### **Alternate Names:**

Alveolar Hydatid Disease

### **Alveolar Gas Equation**

A formula used to estimate the oxygen level in the lung air sacs from the inhaled oxygen level and carbon-dioxide level. Comparing this estimate with arterial blood oxygen helps assess how well the lungs exchange gases.

### **Alveolar Hypoventilation**

Deficient ventilation of the pulmonary alveoli resulting in reduced carbon dioxide elimination, leading to arterial hypercapnia and respiratory acidosis.

### **Alveolar Macrophage**

An immune cell that lives in the lung air sacs and removes inhaled dust, microbes, and damaged cells. It also helps regulate inflammation in the lungs.

### **Alternate Names:**

Dust Cell

### **Alveolar Osteitis**

Painful inflammation of a tooth socket after extraction, usually caused by loss or breakdown of the blood clot that normally protects the socket. It is also called dry socket.

### **Alternate Names:**

Dry Socket; Fibrinolytic Alveolitis

### **Alveolar Proteinosis**

Alternate form; see *Pulmonary Alveolar Proteinosis*.

### **Alveolar Rhabdomyosarcoma**

A cancer of immature muscle-forming cells, usually affecting children or adolescents. It may arise in the limbs, trunk, head, or neck and can spread early to lymph nodes, lungs, bones, or other tissues.

### **Alveolar Ventilation**

The amount of fresh air reaching the lung air sacs each minute and available for oxygen and carbon-dioxide exchange. It depends on breathing rate, breath size, and the amount of air remaining in the

---

breathing passages.

### **Alveolar Ventilation Equation**

A relationship showing how breathing affects blood carbon-dioxide levels. When the amount of fresh air reaching the lung air sacs falls, carbon dioxide rises; when effective breathing increases, carbon dioxide falls, if production stays constant.

### **Alveolar-Arterial Oxygen Gradient**

The difference between oxygen pressure in the lung air sacs and oxygen pressure in arterial blood. It is calculated as  $PAO_2$  minus  $PaO_2$ . A high value may indicate a problem with oxygen transfer, blood flow through the lungs, or lung tissue. A normal value with low blood oxygen may suggest slow breathing or low oxygen in the surrounding air.

#### **Alternate Names:**

A-A Gradient; A-a Gradient; A-a Oxygen Difference

### **Alveolar-Capillary Barrier**

The delicate biological membrane formed by the alveolar type I pneumocyte, capillary endothelial cell, and their intervening fused basement membranes, through which oxygen and carbon dioxide diffuse between alveolar air and capillary blood.

#### **Alternate Names:**

Alveolar-Capillary Membrane; Blood-Air Barrier

### **Alveolitis**

Inflammation of the lung air sacs, usually causing cough and shortness of breath, or inflammation of a tooth socket after extraction, called dry socket. The body site identifies the meaning.

### **Alveolus**

One of the microscopic air sacs at the ends of the lung airways. Oxygen passes from an alveolus into the blood and carbon dioxide passes from the blood into it.

#### **Alternate Names:**

Alveoli

### **Alzheimer Disease**

A progressive disease of the brain and the most com-

mon cause of dementia. It slowly damages nerve cells and the connections between them. Early symptoms often include problems with recent memory, finding words, or thinking through tasks. As the disease advances, it can affect language, judgment, orientation, behavior, and the ability to carry out everyday activities. Abnormal proteins called amyloid plaques and tau tangles build up in the brain and are linked to this damage. Alzheimer disease is not a normal part of aging, although the risk increases with age, and its symptoms and rate of progression vary.

#### **Alternate Names:**

Alzheimer's Disease; Alzheimer's Alzheimer's Alzheimer's Disease

### **Alzheimer's**

A common possessive form of Alzheimer disease, a progressive neurodegenerative disorder causing worsening memory, thinking, behavior, and ability to perform daily activities.

### **Amalgam**

A dental filling material made by combining mercury with metals such as silver, tin, and copper. It is strong and durable but has a metallic appearance and may not be suitable for every person or tooth.

### **Amastia**

A birth condition in which one or both breasts do not develop. The nipple, chest muscle, or other nearby structures may also be absent or underdeveloped.

### **Amaurosis**

Partial or complete loss of vision, which may be temporary or permanent and may result from disease of the eye, retina, optic nerve, or visual pathways in the brain. Sudden vision loss requires emergency assessment.

### **Amaurosis Fugax**

Sudden, temporary loss of vision in one eye, often described as a curtain or shade, caused by briefly reduced blood flow to the retina or optic nerve. It can warn of vascular disease or stroke risk.

---

## Ambidextrous

Able to use the left and right hands with similar skill. It describes a pattern of handedness, not a disease.

## Ambient

Surrounding or present in the immediate environment, such as ambient temperature, light, noise, or air.

## Ambiguous Genitalia

External genital anatomy that does not have typical male or female features. It usually reflects a difference in sex development, which may involve chromosomes, hormones, or the formation of reproductive organs.

## Amblyopia

Reduced vision in one eye (rarely both) resulting from abnormal brain visual processing during early childhood, occurring despite a structurally normal eye. When the brain receives conflicting, blurry, or misaligned visual signals, it suppresses input from the weaker eye and favors the dominant eye. Common causes include crossed or misaligned eyes (strabismic amblyopia), unequal refractive focusing power between the eyes (anisometropic amblyopia), or physical blockage of the visual axis (deprivation amblyopia from congenital cataract or severe ptosis). Because neural visual pathways develop rapidly in early life, treatment is time-sensitive and most effective during the critical period before age 7 to 8. Management involves correcting refractive errors with glasses, clearing anatomical obstructions, and patching the stronger eye (or using blurring atropine drops) to force the brain to strengthen neural pathways in the amblyopic eye.

### Alternate Names:

Lazy Eye; Functional Amblyopia; Visual Suppression Amblyopia

## Ambu Bag

A self-inflating manual resuscitation bag used with a mask or airway device to push air into the lungs when a person is not breathing adequately.

## Ambulatory Blood Pressure Monitoring

Measurement of blood pressure at regular intervals, usually for twenty-four hours, with a portable cuff worn during normal activities and sleep. It can identify blood-pressure patterns that may not appear during a clinic visit.

## Amebiasis

An infection caused by the parasite *Entamoeba histolytica*, usually spread by swallowing contaminated food, water, or material from contaminated hands. It may cause no symptoms, diarrhea, abdominal pain, or bloody diarrhea. In some cases, the infection can spread from the intestine to the liver.

### Alternate Names:

*Entamoeba Histolytica* Infection; Amoebic Dysentery

## Amebic Colitis

Inflammation of the large intestine caused by *Entamoeba histolytica*. It may cause abdominal pain, diarrhoea, bloody stools, or fever.

### Alternate Names:

Amoebic Colitis

## Amebic Dysentery

An acute, invasive intestinal infection caused by the protozoan parasite *Entamoeba histolytica*, characterized by extensive ulceration of the colonic mucosa ("flask-shaped" ulcers). Patients typically present with severe lower abdominal cramping, tenesmus, and frequent passage of bloody, mucoid stools, with potential complications including fulminant necrotizing colitis, toxic megacolon, bowel perforation, and secondary amebic liver abscess.

### Alternate Names:

Amoebic Dysentery; Intestinal Amebiasis; Amebic Colitis

## Amebic Liver Abscess

A collection of infected or damaged tissue in the liver caused by the parasite *Entamoeba histolytica*. It often affects the right side of the liver and may cause fever, pain under the right ribs, and an enlarged or tender liver. Ultrasound or another scan and blood tests can support the diagnosis, but results must be

---

interpreted with the person's symptoms and history. Treatment usually includes a medicine that kills the parasite followed by a medicine that clears it from the intestine; drainage is needed in selected cases.

**Alternate Names:**

Amoebic Liver Abscess; Hepatic Amebiasis; Entamoeba Liver Abscess

**Ameboma**

A chronic, localized inflammatory granulomatous pseudotumor of the bowel wall that develops in a minority of patients with untreated or inadequately treated *Entamoeba histolytica* infection, most commonly affecting the cecum or ascending colon. It presents as a palpable, tender right-lower-quadrant abdominal mass associated with intermittent bloody diarrhea, obstruction, or weight loss, frequently mimicking colonic adenocarcinoma on colonoscopy and cross-sectional imaging.

**Alternate Names:**

Amoeboma; Amebic Granuloma

**Amelanotic Melanoma**

A melanoma that produces little or no dark pigment and may appear pink, red, or skin-coloured instead of black or brown. Its appearance can delay recognition; diagnosis requires examination of a tissue sample.

**Amelia**

A rare birth condition in which one or more arms or legs are completely absent. It may occur alone or with other developmental differences.

**Amelioration**

Improvement or reduction in the severity, frequency, or effect of a symptom, disease, impairment, or other harmful condition.

**Ameloblastoma**

A usually slow-growing tumour that begins in cells involved in tooth development, most often in the lower jaw. It can grow into nearby bone, deform the jaw, and return after treatment.

**Amelogenesis**

The process by which developing teeth form and harden enamel, the tough outer covering of each tooth. Problems with this process can make enamel thin, weak, discoloured, or easily damaged.

**Amelogenesis Imperfecta**

A group of inherited conditions in which tooth enamel does not form normally. The enamel may be thin, soft, rough, discoloured, or easily worn on most or all teeth.

**Amenorrhea**

Absence of menstrual periods. Primary amenorrhea means menstruation has not started by the expected age; secondary amenorrhea means periods stop after they have begun. It may be normal during pregnancy, breastfeeding, or menopause, or may result from hormonal, reproductive, medical, nutritional, or exercise-related conditions.

**American Trypanosomiasis**

Alternate name; see *Chagas disease*.

**Ametropia**

A focusing problem in which the eye does not bring incoming light to a sharp focus on the retina. Near-sightedness, farsightedness, and astigmatism are common forms and can cause blurred vision.

**Amino Acid**

A small molecule used to build proteins. Amino acids also support energy production and other chemical processes in the body.

**Alternate Names:**

Protein Building Block; Amino Monocarboxylic Acid

**Amino Acid Catabolism**

The body's breakdown of amino acids after proteins are digested. Their nitrogen part is removed and safely processed, while the remaining part is used to make energy, glucose, ketones, or other substances.



---

## Amino Acid Symbols

The standard one-letter and three-letter abbreviations used to identify the 20 amino acids most commonly used to build proteins. They make genetic, laboratory, and biochemical information shorter and easier to compare.

## Aminoaciduria

Abnormally large amounts of amino acids in the urine, usually because the kidney tubules cannot reabsorb them properly or because of an inherited metabolic disorder.

## Aminoglycoside Nephrotoxicity

Kidney injury caused by aminoglycoside antibiotics, which can build up in kidney cells. Risk increases with high doses, prolonged treatment, existing kidney disease, or other kidney-toxic medicines.

## Aminoglycoside Ototoxicity

Inner-ear damage caused by aminoglycoside antibiotics, which may lead to hearing loss, ringing in the ears, dizziness, or balance problems. Risk increases with high or prolonged exposure.

## Aminoglycosides

A group of antibiotics that kill certain bacteria by stopping them from making essential proteins. Examples include gentamicin, tobramycin, amikacin, and streptomycin; they can damage the kidneys and inner ear.

## Aminophylline Overdose

Poisoning caused by too much aminophylline, a medicine that opens narrowed airways. It can cause vomiting, tremor, rapid or irregular heartbeat, low blood pressure, seizures, or dangerous changes in blood chemistry.

## Aminotransferase

An enzyme that helps cells process amino acids, the building blocks of proteins. Blood levels of ALT and AST may rise when liver, muscle, or other cells are injured, but the result does not identify the cause alone.

## Amiodarone Pulmonary Toxicity

A serious, potentially life-threatening adverse reaction to chronic amiodarone therapy, manifesting as subacute chronic interstitial pneumonitis, organizing pneumonia, or acute respiratory distress syndrome. Pathologically marked by intra-alveolar foamy lipid-laden macrophages (phospholipidosis) and diffuse alveolar damage.

### Alternate Names:

Amiodarone Lung; Amiodarone-Induced Pneumonitis

## Amiodarone-Induced Skin Discoloration

A blue-grey or slate-grey darkening of sun-exposed skin caused by long-term amiodarone treatment. It is more likely with higher doses and may fade partly after the medicine is stopped.

## Amiodarone-Induced Thyroid Dysfunction

An overactive or underactive thyroid caused by amiodarone. The medicine contains iodine and can either increase thyroid-hormone production or inflame and damage the thyroid.

## Amiodarone-Induced Vortex Keratopathy

Whorl-shaped deposits in the surface layer of the cornea caused by amiodarone. They usually cause no symptoms but may produce halos around lights and generally improve after the medicine is stopped.

## Amitriptyline

A tricyclic antidepressant that changes serotonin and noradrenaline signalling in the brain. It is also used for some nerve pain and migraine prevention; common effects include sleepiness, dry mouth, and constipation.

## Amitriptyline And Perphenazine Overdose

Poisoning caused by taking too much amitriptyline and perphenazine. It can cause severe sleepiness, confusion, seizures, low blood pressure, abnormal heart rhythms, coma, or breathing problems.

---

### **Amitriptyline Hydrochloride Overdose**

Poisoning caused by taking too much amitriptyline. It can cause severe sleepiness, confusion, seizures, low blood pressure, abnormal heart rhythms, coma, or breathing problems.

### **Ammonia**

A nitrogen-containing waste product made when the body processes proteins. The liver normally changes it into urea for removal; high blood levels can affect brain function, especially in severe liver disease.

### **Ammonia Blood Test**

A blood test that measures ammonia, a waste product normally cleared by the liver. A high result may occur with severe liver disease, inherited metabolic conditions, some medicines, or other illnesses and must be interpreted with symptoms.

### **Ammonia Poisoning**

Injury caused by breathing, swallowing, or getting concentrated ammonia on the skin or eyes. It can burn tissues and cause coughing, breathing difficulty, chest pain, or serious lung damage.

### **Ammonia Toxicity**

Irritant or corrosive injury after inhalation, ingestion, or skin or eye contact with ammonia. Concentrated exposure can cause burning, cough, bronchospasm, laryngeal injury, pulmonary edema, or chemical burns; treatment emphasizes immediate decontamination, airway and respiratory support, and specialist toxicology guidance.

### **Ammonium Hydroxide Poisoning**

Injury caused by exposure to ammonium hydroxide, a corrosive chemical found in some cleaners and industrial products. It can burn the skin, eyes, mouth, throat, lungs, and digestive tract.

### **Amnesia**

Loss or impairment of memory beyond ordinary forgetfulness. It may prevent a person from forming new memories, recalling past events, or doing both, and may result from brain injury, illness, medicines,

or emotional trauma.

### **Amnestic Syndrome**

A pattern of memory impairment that significantly affects learning, storing, or recalling information. It may occur with brain injury, stroke, dementia, epilepsy, substance use, or other neurological conditions.

### **Amniocentesis**

A procedure in which a thin needle removes a small sample of fluid from around the developing baby. The sample can be tested for genetic, chromosomal, or infection-related conditions.

### **Amnioinfusion**

A procedure that adds sterile fluid to the sac around the developing baby, usually during labour, to reduce pressure on the umbilical cord or dilute thick meconium.

### **Amnion**

The thin inner membrane forming the fluid-filled sac around the developing baby. It holds the amniotic fluid and helps protect the fetus during pregnancy.

### **Amnion Nodosum**

Small deposits of fetal skin material or other debris attached to the inner membrane of the pregnancy sac. They are usually linked with too little amniotic fluid, often because the developing baby is making less urine.

### **Amnioscope**

An instrument formerly inserted through the cervix to examine the amniotic fluid and membranes. It is rarely used now because ultrasound is safer and more practical.

### **Amniotic Band**

A strand of amniotic membrane that can wrap around a developing body part, restricting growth or blood flow and causing a deformity or loss of tissue.

**Alternate Names:**

---

Amniotic Band Sequence

### Amniotic Band Sequence

A group of birth differences caused when strands of the amniotic membrane wrap around or attach to developing fetal parts. Effects may include constriction rings, missing digits, limb defects, or facial and body-wall abnormalities.

### Amniotic Band Syndrome

Another name for a group of birth differences caused by strands of amniotic membrane constricting or entangling developing fetal parts. The effects vary widely and may involve the limbs, face, or body wall.

#### Alternate Names:

Amniotic Constriction Band Sequence; ADAM Complex; Streeter Dysplasia

### Amniotic Fluid

The liquid surrounding the developing baby inside the amniotic sac. It cushions the baby, allows movement and growth, and helps maintain a stable environment during pregnancy. It is continually produced, swallowed, and replaced.

### Amniotic Fluid Embolism

A rare, life-threatening complication of pregnancy or childbirth in which amniotic material enters the mother's bloodstream and triggers sudden breathing or heart problems, often with severe bleeding.

### Amniotic Fluid Index

An ultrasound estimate of the amount of fluid around the developing baby. It adds the deepest fluid measurements from four areas of the uterus and helps identify unusually low or high fluid levels.

#### Alternate Names:

AFI; Amniotic Fluid Volume Assessment

### Amniotic Sac

The thin, fluid-filled membrane around the developing baby in the uterus. It contains amniotic fluid, cushions the fetus, and usually breaks near or during labour when the waters break.

### Amok

An old, culture-related term for a sudden episode of uncontrolled, often violent behaviour followed by exhaustion or memory loss. Modern assessment considers medical, neurological, psychiatric, substance-related, and cultural factors.

### AMP-Activated Protein Kinase

A master cellular energy-sensing serine/threonine kinase activated during metabolic stress when intracellular AMP-to-ATP or ADP-to-ATP ratios rise (such as during exercise, hypoxia, or fasting). Once activated, it restores energy homeostasis by promoting ATP-generating catabolic processes (such as glucose uptake via GLUT4, fatty acid oxidation, glycolysis, and autophagy) while suppressing ATP-consuming anabolic pathways (such as lipid synthesis, protein synthesis via mTOR inhibition, and gluconeogenesis). It is a primary molecular target of the antidiabetic drug metformin.

#### Alternate Names:

AMPK; 5'-AMP-Activated Protein Kinase

### Amphetamine

A stimulant medicine that increases certain chemical signals in the brain. It is used for selected attention, activity, or sleep disorders; misuse can cause dependence, high blood pressure, abnormal heart rhythms, or psychosis.

### Amphetamine Substance Use

Use of amphetamine-type stimulants, which can be associated with insomnia, anxiety, psychosis, high blood pressure, arrhythmias, overheating, dependence, and overdose.

### Amphetamine Toxicity

A sympathomimetic toxidrome caused by amphetamine or related stimulants, with agitation, mydriasis, sweating, tachycardia, hypertension, hyperthermia, chest pain, seizures, or psychosis. Severe cases may cause stroke, rhabdomyolysis, or cardiac ischemia; benzodiazepines, cooling, and treatment of complications are central.

### Amphetamine-Related Psychiatric

---

## Disorders

Psychiatric syndromes associated with amphetamine exposure, including intoxication, withdrawal, substance-induced psychosis, mood symptoms, anxiety, or a stimulant use disorder.

## Amplified Musculoskeletal Pain Syndrome

A condition, often affecting children or teenagers, in which the nervous system increases normal pain signals. Pain may be severe or widespread and can occur with sensitivity, fatigue, swelling, or difficulty using the affected area.

## Ampulla

A widened part of a tube-shaped body structure. Examples include the ampulla where the bile and pancreatic ducts join the intestine and the ampulla of a fallopian tube.

## Ampulla of Vater

An anatomical dilation located within the medial wall of the descending duodenum, formed by the union of the common bile duct and the main pancreatic duct, draining into the duodenal lumen via the major duodenal papilla.

### Alternate Names:

Hepatopancreatic Ampulla

## Ampullary Carcinoma

A malignant epithelial tumor arising in or around the ampulla of Vater, where the bile duct and pancreatic duct enter the duodenum. It can block bile flow and cause jaundice, abnormal liver tests, pancreatitis, or gastrointestinal bleeding.

### Alternate Names:

Ampullary Cancer

## Amputation

Surgical removal of all or part of a limb, finger, toe, or other body part, often because of severe injury, infection, poor blood flow, or cancer. Traumatic amputation is tissue loss caused by an injury. The level removed depends on healthy tissue, blood supply, healing, and expected function.

### Alternate Names:

Surgical Amputation; Limb Removal

## Amputation Stump

The part of a limb left after amputation. It may develop wound problems, infection, poor circulation, nerve pain, phantom-limb pain, or skin irritation from a prosthesis.

## Amsler Grid

A small square chart used to check central vision. Wavy, missing, or distorted lines may indicate a problem in the macula and are evaluated with a complete eye examination.

## Amusia

Difficulty recognizing, understanding, or producing music despite adequate hearing. It may be present from birth or develop after brain injury, stroke, or another neurological condition.

## Amygdala

A group of nerve cells deep in each side of the brain that helps process emotions, especially fear, and contributes to learning, memory, and responses to important or threatening events.

### Alternate Names:

Amygdaloid Nucleus

## Amylase

An enzyme that begins breaking down starch. It is made mainly by the salivary glands and pancreas and can be measured in blood or urine when pancreatic or salivary-gland disease is suspected.

## Amylase And Lipase

Blood tests for digestive enzymes used when pancreatic disease is suspected. Lipase is generally more specific for pancreatic injury, but neither test alone confirms acute pancreatitis.

## Amylase Test

A blood or urine test that measures the amount of amylase, an enzyme made mainly by the salivary glands and pancreas. It may help assess pancreatitis

---

and other problems affecting the pancreas or salivary glands.

**Alternate Names:**

Serum Amylase; Blood Amylase Test; Amylase Blood Test

**Amylin Analogues**

Medicines that imitate amylin, a hormone released by the pancreas with insulin. They slow stomach emptying, reduce the rise in blood glucose after meals, and reduce glucagon release.

**Amyloid**

An abnormal protein material that forms insoluble deposits in tissues. These deposits can interfere with the function of organs such as the kidneys, heart, nerves, liver, or brain.

**Alternate Names:**

Amyloid Protein

**Amyloid Nephropathy**

Renal involvement occurring in systemic amyloidosis, characterized by extracellular accumulation of insoluble amyloid fibrils in glomerular, vascular, and interstitial compartments, typically leading to heavy proteinuria, nephrotic syndrome, and progressive kidney failure.

**Amyloid Neuropathy**

Nerve damage caused by amyloid deposits in or around peripheral nerves. It may cause numbness, tingling, burning pain, weakness, or problems with automatic body functions such as blood-pressure control.

**Amyloid PET**

A brain scan using a small radioactive tracer that attaches to amyloid deposits. It can help assess selected cognitive disorders, but the result must be interpreted with symptoms and other tests.

**Amyloid Plaque**

A deposit of amyloid-beta protein outside nerve cells in the brain. Plaques are associated with Alzheimer disease, but their presence alone does not explain

every person's memory or thinking problem.

**Amyloid Precursor Protein**

A normal protein found in nerve-cell membranes. Enzymes can cut it into smaller pieces, including amyloid-beta; abnormal accumulation of amyloid-beta in the brain is associated with Alzheimer disease.

**Alternate Names:**

APP; Amyloid-Beta Precursor Protein

**Amyloid Purpura**

Purple or bruise-like spots, often around the eyes, caused by amyloid deposits weakening small blood vessels. It may occur with systemic amyloidosis and can be accompanied by easy bruising elsewhere.

**Amyloidosis**

A group of diseases in which proteins fold abnormally, clump together, and form deposits called amyloid in tissues or organs. Amyloidosis may be localized to one site or systemic, affecting several organs such as the kidneys, heart, liver, nerves, or digestive tract. Major forms include light-chain (AL), serum amyloid A (AA), and transthyretin (ATTR) amyloidosis; effects vary with the amyloid protein and organs involved.

**Alternate Names:**

Amyloid Disease; Amyloid Infiltration

**Amyotrophic Lateral Sclerosis**

A progressive disease in which the nerve cells controlling voluntary muscles gradually die. It causes worsening muscle weakness, stiffness, cramps, twitching, and wasting, often beginning in an arm, leg, or the muscles used for speaking and swallowing. It may lead to difficulty walking, speaking, swallowing, or breathing. Sensation is usually preserved, but some people develop changes in thinking or behavior. Most cases are not inherited and have no known cause. Also called ALS or Lou Gehrig disease.

**Alternate Names:**

Lou Gehrig's Disease ALS

**Anabolic Steroid Induced Hypogonadism**

Reduced testicular hormone production and fertility



---

caused by misuse of anabolic steroids. Extra steroid hormones suppress the body's normal hormone signals and may lead to testicular shrinkage, low sex drive, erectile problems, infertility, or depression.

**Alternate Names:**

ASIH; Anabolic-Androgenic Steroid-Induced Hypogonadism

### Anabolic Steroid Use and Abuse

Use of anabolic-androgenic steroids without a medical indication or at excessive doses. It can suppress natural hormone production and affect fertility, mood, blood pressure, cholesterol, liver function, heart structure, and growth in adolescents.

### Anabolic Steroids

Synthetic medicines related to testosterone that promote muscle and bone growth. They have limited medical uses but are sometimes misused for appearance or performance; misuse can harm the heart, liver, fertility, mood, and hormone system.

### Anabolism

The set of body processes that use energy to build larger molecules from smaller ones, such as making proteins from amino acids or storing glucose as glycogen.

### Anaemia

A condition in which the blood has too few red blood cells or too little haemoglobin. This reduces the blood's ability to carry oxygen and may cause tiredness, weakness, breathlessness, or dizziness.

### Anaerobe

A microorganism that can grow without oxygen. Some anaerobes are harmed or killed by oxygen, while others tolerate oxygen but do not use it to produce energy.

### Anaerobic

Able to live, grow, or function without oxygen. Anaerobic bacteria can cause infections in deep wounds, abscesses, the mouth, abdomen, or pelvis.

### Anaerobic Bacteria

Bacteria that can grow without oxygen. Obligate anaerobes cannot use oxygen and may be harmed by it, while facultative anaerobes can grow with or without oxygen. Anaerobic bacteria normally live in places such as the mouth, intestine, and vagina. They can cause infections in deep wounds, abscesses, the abdomen, pelvis, or other tissues with low oxygen.

### Anaerobic Glycolysis

The metabolic breakdown of glucose into lactate in the absence or insufficiency of cellular oxygen, producing a net yield of two molecules of adenosine triphosphate per glucose molecule and regenerating oxidized nicotinamide adenine dinucleotide (NAD<sup>+</sup>) to maintain energy generation.

### Anagen

The active growth phase of the hair cycle, when cells in the hair root make the hair shaft. Scalp hairs usually remain in this phase for several years before entering a resting or shedding phase.

### Anagen Effluvium

An abrupt, extensive shedding of hair during the active growth (anagen) phase, caused by acute toxic disruption of mitotic activity in the follicular matrix cells. It typically begins within days to weeks of exposure to cytotoxic chemotherapy, ionizing radiation, or acute heavy metal poisoning.

### Anal Abscess

A collection of pus near or within the anus, usually caused by an infected anal gland. It can cause throbbing pain, swelling, fever, and difficulty sitting.

### Anal Canal

The final part of the digestive tract, extending from the rectum to the anus. It contains muscles that control the passage of stool and has different nerve and blood supplies along its length.

### Anal Cancer

Cancer arising in the anal canal or perianal skin, most often squamous-cell carcinoma associated with

---

persistent high-risk HPV infection. Bleeding, pain, a lump, itching, or altered bowel function may occur; examination and biopsy establish the diagnosis.

### **Anal Carcinoma**

Cancer arising in the anal canal or anal margin, most often squamous-cell cancer associated with persistent high-risk human papillomavirus infection. It may cause bleeding, pain, itching, or a lump.

#### **Alternate Names:**

Anal Cancer

### **Anal Condyloma**

A wart-like growth on or around the anus, usually caused by low-risk types of human papillomavirus. It is usually not cancerous, but multiple or recurring warts may occur with other abnormal anal cells.

### **Anal Fissure**

A small tear in the lining of the anus, usually caused by hard stools, constipation, or repeated diarrhea. It can cause sharp pain during bowel movements and bright-red blood on the toilet paper.

### **Anal Fistula**

An abnormal tunnel between the anal canal and nearby skin, often developing after an anal abscess. It may cause repeated discharge, pain, itching, or a small opening near the anus.

### **Anal Intercourse**

Sexual activity involving insertion into the anus. Because the anal lining can tear and does not provide natural lubrication, it can transmit infections and cause injury if precautions are not used.

### **Anal Intraepithelial Neoplasia**

Abnormal cells in the lining of the anus, usually associated with persistent high-risk human papillomavirus infection. Low-grade changes may resolve, while high-grade changes can progress to anal cancer.

### **Anal Itching**

Itching around the anus. Common causes include

moisture, irritation, dermatitis, haemorrhoids, infection, and occasionally pinworms or another disease.

#### **Alternate Names:**

Pruritus Ani; Perianal Itching; Rectal Itching

### **Anal Mucositis**

Inflammation and open sores of the anal lining, often caused by radiation or chemotherapy. It can cause pain, burning, bleeding, discharge, and difficulty passing stool.

### **Anal Necrosis**

Death of tissue around the anus, anal canal, or nearby skin. It may result from severe infection, poor blood flow, injury, or radiation and can cause severe pain, skin discolouration, blisters, fever, or rapid worsening.

### **Anal Pain**

Pain around the anus or nearby skin. Common causes include a fissure, abscess, thrombosed haemorrhoid, muscle spasm, infection, or inflammation.

#### **Alternate Names:**

Proctalgia

### **Anal Papilla**

A small, usually normal projection of the anal lining near the junction between the upper and lower types of lining in the anal canal.

### **Anal Sphincter**

A muscular structure that controls passage of stool and gas from the anal canal. It includes the involuntary internal anal sphincter and the voluntary external anal sphincter.

### **Analgesia**

Relief or absence of pain without necessarily causing unconsciousness. It may result from medicines, a nerve block, or another treatment; touch, pressure, or movement may still be felt.

### **Analgesics**

Medicines that reduce or relieve pain. Non-opioid analgesics reduce pain and fever. Nonsteroidal anti-

---

inflammatory drugs (NSAIDs) reduce pain, fever, and inflammation by lowering the production of substances involved in these responses. Opioid analgesics reduce the brain's and spinal cord's response to pain but may cause sleepiness, constipation, slowed breathing, tolerance, and dependence. Adjuvant analgesics are medicines mainly used for other conditions that can help with certain types of pain, such as nerve pain. The choice depends on the type and severity of pain, the person's health, and possible risks.

**Alternate Names:**

Painkillers; Pain-Relieving Medicines

**Analog**

A substance, structure, or process that resembles another and has a similar function but is not identical in chemical makeup or structure.

**Analogous**

Similar in function or purpose to another structure or process, even though its origin or physical structure is different.

**Analysis**

A careful examination of information, samples, measurements, symptoms, or clinical problems to identify patterns, understand causes, or support a medical decision.

**Analysis Of Urine**

Laboratory examination of urine for blood, cells, protein, glucose, germs, and other substances. Results can help assess urinary infection, kidney disease, diabetes, bleeding, dehydration, and other conditions.

**Analytic Sensitivity**

The smallest amount of a substance that a laboratory method can reliably detect. It describes the test's measurement ability and is different from clinical sensitivity, which describes how well a test detects disease.

**Anaphia**

Complete loss of touch sensation in a body area. It

may result from nerve injury, neurological disease, reduced blood flow, or anaesthetic treatment and can allow burns, wounds, or pressure injuries to go unnoticed.

**Alternate Names:**

Anaptic

**Anaphylactic Shock**

A severe form of anaphylaxis in which a sudden allergic reaction causes dangerously low blood pressure and poor blood flow to organs. It may occur with airway swelling, breathing difficulty, collapse, or cardiac arrest.

**Anaphylactoid Reaction**

An immediate, severe systemic hypersensitivity reaction that closely mimics IgE-mediated anaphylaxis in its clinical manifestations, but occurs via direct non-immune or non-IgE-mediated mast cell and basophil degranulation (such as that triggered by radiocontrast media or certain drugs).

**Alternate Names:**

Non-Allergic Anaphylaxis; Pseudoallergic Reaction

**Anaphylatoxin**

A small peptide fragment (predominantly C3a, C4a, and C5a) generated during activation of the complement cascade. Anaphylatoxins induce mast cell degranulation, smooth muscle contraction, increased vascular permeability, and chemotaxis of phagocytic leukocytes.

**Anaphylaxis**

A sudden, potentially life-threatening systemic hypersensitivity reaction that can involve the skin, airways, lungs, circulation, or gastrointestinal tract. It may cause hives or swelling, breathing difficulty, vomiting, dizziness, dangerously low blood pressure, or collapse; skin findings may be absent. Immediate intramuscular epinephrine and emergency medical care are required.

**Anaphylaxis During Anesthesia**

A severe allergic reaction occurring during or soon after anaesthesia. It may cause sudden low blood pressure, airway narrowing, breathing difficulty, collapse,

---

or skin flushing; triggers can include medicines, antiseptics, or latex.

### **Anaplasia**

Loss of the normal mature features and specialized function of cells. Anaplastic cells may vary greatly in size and shape, have abnormal nuclei, and divide rapidly or unusually. Anaplasia is most often seen in high-grade cancers and usually suggests more aggressive behavior. It is different from metaplasia, in which one mature cell type changes into another cell type.

### **Anaplasmosis**

A tick-borne infection caused by the bacterium *Anaplasma phagocytophilum*. It may cause fever, headache, muscle aches, low platelet and white-cell counts, and raised liver enzymes. PCR or paired blood-antibody tests can help confirm it. Morulae seen in white blood cells can support the diagnosis but are not always present and do not confirm it alone.

#### **Alternate Names:**

Human Granulocytic Anaplasmosis; HGA

### **Anaplastic**

Describing cells or a tumor that has lost many features of its normal mature form. Anaplastic tumors often grow quickly, invade nearby tissue, or spread, although the outlook depends on the specific cancer and its treatment.

### **Anaplastic Astrocytoma**

An older term for a high-grade tumour arising from star-shaped supporting cells in the brain. Modern classification relies on its genetic features as well as its appearance; symptoms depend on its location.

### **Anaplastic Ependymoma**

A high-grade tumour arising from cells lining the brain's fluid-filled spaces or the spinal canal. It can grow quickly and may obstruct the flow of cerebrospinal fluid.

### **Anaplastic Large Cell Lymphoma**

A rare type of non-Hodgkin lymphoma arising from

abnormal T cells. The cells are usually large and carry the CD30 marker; the disease may affect the skin, lymph nodes, or other organs.

#### **Alternate Names:**

ALCL

### **Anaplastic Large-cell Lymphoma**

A usually aggressive T-cell non-Hodgkin lymphoma characterized by large anaplastic cells and CD30 expression. It may involve lymph nodes, skin, bone, or other organs and is classified as systemic or cutaneous.

### **Anaplastic Thyroid Cancer**

A rare, fast-growing thyroid cancer that may appear as a rapidly enlarging neck lump and cause difficulty swallowing, speaking, or breathing.

### **Anaplastic Thyroid Carcinoma**

An extremely aggressive, undifferentiated malignancy of the thyroid follicular epithelium characterized by rapid local invasion into adjacent cervical structures and early distant metastasis. Presenting symptoms include a rapidly enlarging hard neck mass, hoarseness, dysphagia, and dyspnea, carrying a very poor median survival despite multimodality therapy.

#### **Alternate Names:**

Undifferentiated Thyroid Carcinoma; ATC

### **Anarithmetia**

A primary neuropsychological impairment in basic arithmetic reasoning, computational ability, and understanding of numerical concepts, distinguishing it from secondary calculation deficits caused by spatial or linguistic impairments. It is typically associated with lesions of the dominant inferior parietal lobule (angular gyrus).

### **Anasarca**

Severe, widespread swelling caused by excess fluid under the skin and sometimes around organs. It can occur with serious heart, kidney, or liver disease or very low blood protein and may cause rapid weight gain or breathing difficulty.

---

## Anastomosis

A surgical, traumatic, or pathological connection created between two hollow organs, tubular structures, or blood vessels (such as an arterial anastomosis, arterio-venous fistula, or bowel resection re-connection), establishing communication and allowing continuous fluid or luminal flow.

### Alternate Names:

Surgical Anastomosis; Vascular Anastomosis; Enteroanastomosis

## Anastomotic Leak

Leakage from a connection made during surgery, often where two sections of intestine or another hollow organ were joined. An incomplete seal can allow contents into nearby tissue and cause pain, fever, infection, or life-threatening illness.

## Anastrozole

A medicine that lowers the body's production of oestrogen. It is used mainly for hormone-sensitive breast cancer, especially after menopause.

## Anatomic Orientation Terms

Standard words used to describe the position of one body structure relative to another, including above, below, front, back, near, far, inside, and outside.

## Anatomical Axis

An imaginary line around which a body part rotates or moves. Anatomical axes help describe movements such as rotation of the shoulder, hip, spine, or other joints.

## Anatomical Conduit

A natural or surgically created passage that carries air, fluid, tissue, or another structure between body regions, such as an airway, blood vessel, or surgical channel.

## Anatomical Orifice

A natural opening into or out of the body or through a body structure, such as the mouth, nostril, anus, urethra, or an opening of a blood vessel.

### Alternate Names:

## Body Opening

## Anatomical Planes

Imaginary flat surfaces used to divide the body into sections for describing anatomy and interpreting medical scans. Common planes are sagittal, coronal, and transverse.

## Anatomical Position

The standard reference posture for describing the body: standing upright, facing forward, arms at the sides, hands turned forward, and feet directed forward.

## Anatomical Snuffbox

A triangular hollow on the thumb side of the back of the wrist that appears when the thumb is extended. It overlies the scaphoid bone and radial artery; tenderness there after a fall may indicate a scaphoid fracture.

### Alternate Names:

Anatomic Snuffbox

## Anatomist

A scientist or health professional who studies the structure of the human or animal body, including its organs, tissues, blood vessels, nerves, and their relationships.

## Anatomy

The study of body structure and how its parts relate to one another. It includes visible structures, tissues and cells seen with magnification, and how the body develops.

## ANCA-Associated Vasculitis

A group of rare diseases in which the immune system attacks small blood vessels. It can affect the kidneys, lungs, nerves, skin, eyes, and other organs. A blood test for antineutrophil cytoplasmic antibodies (ANCA) can help with diagnosis but cannot confirm the disease on its own.

## Ancillary Pathology Studies

Supplementary diagnostic modalities—including im-



---

munohistochemistry, special histochemical stains, flow cytometry, in situ hybridization, and molecular genetic assays—performed in conjunction with standard light microscopy to achieve precise pathological classification.

**Alternate Names:**

Ancillary Diagnostic Testing; Diagnostic Immunohistochemistry; Special Pathology Stains

### Ancylostomiasis

A parasitic nematode infection caused by species of the genus *Ancylostoma* (such as *Ancylostoma duodenale*), acquired through transcutaneous penetration of infective larvae from contaminated soil, characteristically leading to chronic intestinal blood loss, iron deficiency anemia, and hypoproteinemia.

**Alternate Names:**

Hookworm Infection; Old World Hookworm Disease; Ancylostoma Duodenale Infection

### Andersen-Tawil Syndrome

A rare inherited disorder causing episodes of muscle weakness, abnormal heart rhythms, and distinctive physical features such as a small lower jaw, low-set ears, or widely spaced eyes.

**Alternate Names:**

ATS; Periodic Paralysis Type 3; Andersen Syndrome

### Anderson Disease

A rare inherited disorder in which abnormal glycogen builds up in cells because of changes in the *GBE1* gene. It usually begins in infancy with poor growth and an enlarged liver and may progress to scarring, high blood pressure in the liver, and liver failure.

**Alternate Names:**

Glycogen Storage Disease Type IV (GSD IV); Amylopectinosis

### Androgen

A hormone such as testosterone that helps regulate sexual development, fertility, muscle, bone, and hair growth in people of all sexes.

### Androgen Deficiency

Too little production or effect of androgen hormones, especially testosterone. In males, it may cause reduced sexual desire, erectile difficulty, infertility, loss of muscle or bone, low energy, or reduced facial and body hair.

### Androgen Excess

Abnormally high levels or effects of androgen hormones. It may cause acne, increased facial or body hair, scalp hair loss, irregular periods, infertility, or a deeper voice, especially in females.

**Alternate Names:**

Hyperandrogenism

### Androgen Insensitivity Syndrome

An inherited condition in which cells cannot respond fully to testosterone and related hormones. A person with XY chromosomes may therefore have female-appearing external genitals, incomplete male development, undescended testes, and no uterus.

### Androgen Insufficiency

A level or effect of androgen hormones that is too low for normal body function. Symptoms may include reduced sexual desire, erectile difficulty, infertility, low energy, reduced muscle mass, or weaker bones.

**Alternate Names:**

Androgen Deficiency

### Androgen Receptor

A protein in cells that binds testosterone and related hormones. This allows the hormones to influence sexual development, reproduction, muscle, bones, hair, and some cancers.

### Androgen Suppression

Lowering the production of androgen hormones or blocking their action in the body. It is used mainly for some hormone-sensitive cancers, especially prostate cancer, and can cause hot flashes, sexual problems, bone loss, weight change, and metabolic effects.

**Alternate Names:**

Androgen Deprivation

---

## Androgenetic Alopecia

A common, inherited form of gradual hair loss caused by sensitivity of hair follicles to androgen hormones. It usually produces a receding hairline or crown thinning in males and diffuse thinning over the crown in females.

## Androgenic

Relating to androgen hormones, such as testosterone, or producing their effects. These effects may include facial or body hair, oily skin, a deeper voice, muscle growth, and development of male reproductive features.

## Android Pelvis

A relatively narrow, heart-shaped or funnel-shaped female pelvis with reduced space in the middle. During childbirth, this shape may make passage of the baby more difficult than some other pelvic shapes.

## Andropause

Alternate informal term; see *Male Hypogonadism*.

## Anejaculation

Inability to release semen during orgasm or sexual activity. It may result from nerve or spinal disease, pelvic surgery, medicines, diabetes, hormonal problems, or psychological factors and may cause infertility.

## Anemia

A lower-than-normal amount of haemoglobin in the blood, often with too few red blood cells. Because haemoglobin carries oxygen, anemia can reduce the oxygen delivered to body tissues. It may cause tiredness, weakness, shortness of breath, or dizziness, though mild anemia may cause no symptoms. Causes include blood loss, reduced red-cell production, or increased red-cell breakdown. Evaluation usually starts with a complete blood count, and treatment depends on the cause.

### Alternate Names:

Anaemia

## Anemia and Thrombocytopenia in

## Pregnancy

Low red-cell or platelet counts during pregnancy may reflect iron deficiency, dilution, nutritional disease, immune disorders, hypertensive disease, bleeding, infection, or marrow disease. Severity, cause, symptoms, and trends guide treatment and delivery planning.

## Anemia In Pregnancy

Anemia during pregnancy, commonly caused by iron deficiency, folate deficiency, vitamin B12 deficiency, blood loss, or the normal increase in the liquid part of blood. It can cause tiredness, weakness, breathlessness, or dizziness.

### Alternate Names:

Gestational Anemia; Pregnancy-Associated Anemia

## Anemia Of Chronic Disease

Anemia associated with long-term inflammation, infection, cancer, kidney disease, or another chronic illness. Inflammation limits the use of stored iron and reduces red-cell production; mild cases may cause no symptoms.

## Anemia Of Inflammation

Anemia caused by long-lasting infection, inflammatory disease, cancer, or another condition that activates the immune system. Inflammation traps iron in storage and slows red-cell production.

## Anemia of Malignancy

A multifactorial normocytic or microcytic anemia developing in patients with neoplastic disease, resulting from cytokine-mediated blunting of erythropoiesis, impaired iron utilization, marrow infiltration, hemorrhage, or myelosuppressive antineoplastic therapies.

### Alternate Names:

Cancer-Related Anemia; Tumor-Induced Anemia; Anemia of Cancer

## Anemic

Having anemia, meaning that the blood carries less oxygen than the body needs because it has too few red blood cells or too little haemoglobin.

---

## Anencephaly

A serious neural tube birth defect in which the upper part of the neural tube fails to close during the first weeks of pregnancy. As a result, much of the brain, especially the forebrain, and parts of the skull and scalp do not develop. It is usually detected before birth and is fatal; most affected babies are stillborn or die shortly after birth. There is no treatment that can correct the defect.

## Anergy

A state in which an immune cell does not respond to a substance that would normally activate it. The cell remains present but is functionally quiet; this can help prevent attacks against the body's own tissues.

## Anesthesia

A medically controlled loss of feeling caused by medicines during a procedure. Anesthetic medicines mainly block nerve signals, especially pain signals, from reaching the brain. Local anesthesia numbs a small area while the person remains awake. Regional anesthesia blocks nerves supplying a larger part of the body and may be used while the person remains awake. General anesthesia also changes brain activity to cause controlled unconsciousness. Before modern anesthesia, surgery was extremely painful and had to be very quick; anesthesia made longer and more complex operations possible. Breathing, blood pressure, heart rate, and oxygen levels are monitored.

### Alternate Names:

Anaesthesia

## Anesthesia Gas Analyzers

Devices that measure the amounts of oxygen, carbon dioxide, and inhaled anesthetic gases delivered to and breathed out by a patient during anesthesia. They help detect problems with breathing or gas delivery.

## Anesthesia Informatics

The use of electronic records, monitoring data, and computer systems to support anesthesia care. It can improve documentation, identify medicine interac-

tions or dosing errors, and help evaluate outcomes.

## Anesthesia Machine

Equipment that supplies controlled mixtures of oxygen, air, and anesthetic gases, and supports breathing during anesthesia. It includes systems that remove carbon dioxide and safety devices that warn of delivery problems.

## Anesthesia Monitoring

Continuous assessment of vital functions during anesthesia, including oxygen levels, breathing, blood pressure, heart rate, temperature, and level of unconsciousness as appropriate for the procedure.

## Anesthesia Pharmacogenomics

The study of how inherited genetic differences affect responses to anesthetic and pain medicines. Examples include prolonged muscle paralysis from pseudocholinesterase deficiency and inherited susceptibility to malignant hyperthermia.

## Anesthesiologist

A doctor trained to provide anesthesia and manage pain, breathing, blood pressure, and other vital functions during procedures. Anesthesiologists also assess patients before surgery and may treat critical illness or severe pain.

## Anesthesiology

The medical specialty covering anesthesia, monitoring and support of vital functions during procedures, care before and after surgery, critical care, and treatment of acute or chronic pain.

## Anetoderma

A rare disorder caused by loss or weakening of elastic fibers in the dermis, producing soft, wrinkled, depressed, or pouch-like macules and plaques with a characteristic button-hole feel. It may be primary or follow infection, inflammation, autoimmune disease, or another skin lesion; biopsy and selected systemic evaluation may be needed.

---

## Aneuploidy

An abnormal number of whole chromosomes in a cell, caused by an error during cell division. A cell may have an extra or missing chromosome, as in trisomy 21, and this can affect development or pregnancy.

## Aneurysm

An abnormal bulge or widening in an artery caused by weakness in its wall. Many aneurysms cause no symptoms. Depending on their location and size, they may enlarge, press on nearby structures, or rupture and cause serious bleeding.

### Alternate Names:

Aneurysms

## Aneurysm Clip

A small surgical device placed across the opening of a brain aneurysm to stop blood from entering it. It is usually placed during open brain surgery; rarely, it may move, block a nearby vessel, or allow bleeding to recur.

## Aneurysm Clipping

Brain surgery to place a small clip across the opening of a brain aneurysm, a weak bulge in a brain artery. The clip stops blood from entering the bulge and lowers the risk of bleeding, but the procedure has its own risks and cannot guarantee that a stroke or future bleeding will not occur.

### Alternate Names:

Surgical Clipping; Cerebral Aneurysm Clipping; Microvascular Clipping

## Aneurysm Coils

Soft metal coils placed inside an aneurysm through a catheter. The coils promote clotting within the bulge and reduce blood flow into it, lowering the chance of rupture.

## Aneurysm Repair

A procedure that reinforces, closes, or replaces a weakened section of a blood vessel to prevent rupture or other complications. It may be performed through open surgery or through a catheter inside the vessel.

## Angelman Syndrome

A genetic disorder affecting brain development and function, usually because the maternal copy of the *UBE3A* gene is missing or inactive. It causes severe developmental delay, very limited speech, movement and balance problems, seizures, and a characteristic happy or excitable manner.

## Anger

An emotional response to frustration, unfairness, danger, or a perceived threat. It may range from irritation to intense rage; persistent or uncontrollable anger can affect relationships, safety, and health.

## Angiitis

Pathological inflammation and necrosis of blood-vessel walls, leading to lumen stenosis, ischemia, aneurysm formation, or hemorrhage in affected organ systems.

### Alternate Names:

Vasculitis; Blood Vessel Inflammation; Arteritis

## Angina

Chest discomfort that occurs when the heart muscle does not get enough oxygen-rich blood, usually because of coronary artery disease, but sometimes because of spasm or problems in the heart's small blood vessels. It may feel like pressure, squeezing, tightness, burning, or indigestion and may spread to the arm, jaw, neck, shoulder, or back. Breathlessness, nausea, sweating, or unusual tiredness may occur, sometimes without chest pain. Stable angina usually follows a predictable pattern, often with activity, and improves with rest or prescribed medicine. New, worsening, prolonged, or rest symptoms may be unstable angina or a heart attack and need emergency assessment.

### Alternate Names:

Angina Pectoris

## Angina Bullosa Hemorrhagica

A usually benign, sudden blood-filled blister of the oral mucosa, most often the soft palate. It may follow minor trauma, inhaled or topical steroids, or occur with diabetes; blisters usually rupture and heal without treatment, but recurrent or atypical

---

lesions need evaluation to exclude autoimmune blistering disease.

### Angina Symptoms

Symptoms of reduced blood flow to the heart may include pressure, squeezing, burning, or tightness in the chest, with discomfort spreading to the arm, jaw, back, or shoulder. Breathlessness, sweating, nausea, or unusual tiredness may also occur.

#### Alternate Names:

Ischemic Chest Pain; Anginal Chest Pain; Anginal Discomfort

### Angioblastoma

A usually slow-growing, highly vascular tumour of the central nervous system, most often found in the cerebellum or spinal cord. It may cause headache, balance problems, weakness, or other symptoms from pressure on nearby tissue.

### Angiocardiology

Imaging of the heart and major blood vessels, usually after contrast material is given. X-ray, CT, or MRI can show the heart chambers, valves, narrowed vessels, leaks, abnormal connections, or other structural problems.

### Angiodysplasia

A common acquired vascular malformation characterized by fragile, dilated, tortuous submucosal blood vessels in the gastrointestinal tract, most frequently found in the cecum and right colon of adults over age 60. It is a major cause of recurrent, painless lower gastrointestinal bleeding and unexplained iron-deficiency anemia. It is strongly associated with aortic valve stenosis (Heyde's syndrome, due to acquired von Willebrand factor deficiency) and chronic renal failure. Diagnosis is established on colonoscopy, where lesions appear as small, bright-red, spider-like mucosal patches; active or high-risk bleeding is treated endoscopically using argon plasma coagulation (APC) or electrocautery.

#### Alternate Names:

Colonic Angiodysplasia; Arteriovenous Malformation of the Colon; Vascular Ectasia of the Colon

### Angiodysplasia Of The Colon

Alternate name; see *Angiodysplasia*.

### Angioedema

Sudden, localized swelling of deeper skin or mucosal tissues, often affecting the lips, eyelids, tongue, throat, hands, feet, or bowel wall. Histamine-mediated episodes may occur with hives and itching, whereas bradykinin-mediated forms usually lack hives; tongue or throat swelling, voice change, or breathing difficulty is an emergency.

### Angioedema And Hives

A combination of raised, itchy skin patches called hives and deeper swelling called angioedema. Common triggers include foods, medicines, insect stings, infections, and physical exposures; breathing difficulty is an emergency.

### Angiofibroma

A usually noncancerous growth made of blood vessels and fibrous tissue. It commonly appears as a small facial papule in adolescents or, in a different form, as a vascular tumour in the back of the nose that may cause blockage or nosebleeds.

### Angiofollicular Lymph Node Hyperplasia

An uncommon disorder in which lymph nodes contain excess small blood vessels and abnormal immune-cell growth. It is also called Castleman disease and may affect one lymph-node region or many parts of the body.

### Angiogenesis

The formation of new blood vessels from existing vessels. It is normal during growth, menstruation, and wound healing, but abnormal angiogenesis can help cancers grow and can damage vision in some eye diseases.

### Angiogenesis Inhibition

Preventing or slowing the growth of new blood vessels. This can limit a tumor's blood supply or reduce harmful vessel growth in some eye diseases. Treatment may increase the risk of high blood pressure, bleeding, or delayed wound healing.



---

**Alternate Names:**

Antiangiogenic Therapy

**Angiogenesis Inhibitor**

A therapeutic agent that inhibits the formation of new blood vessels, predominantly by antagonizing vascular endothelial growth factor (VEGF) or its receptors, used in oncology to restrict tumor blood supply and in ophthalmology to treat neovascular macular degeneration.

**Alternate Names:**

Antiangiogenic Agent

**Angiogram**

An image of blood vessels, usually made after contrast material is given. It can show narrowing, blockage, aneurysm, tearing, abnormal connections, or bleeding and may be produced by catheter X-ray, computed tomography, or magnetic resonance imaging.

**Angiography**

Imaging of blood vessels to show their shape and blood flow. It may use X-rays with injected contrast material, computed tomography (CT), magnetic resonance imaging (MRI), or a catheter placed inside a blood vessel. Angiography can show narrowing, blockage, aneurysms, abnormal connections, bleeding, or other vascular abnormalities; catheter angiography may also allow treatment during the same procedure.

**Angioid Streak**

Irregular dark lines caused by breaks in a tough layer beneath the retina. They may bleed or allow abnormal blood vessels to grow, reducing vision, and can be associated with some inherited disorders.

**Angioimmunoblastic T-Cell Lymphoma**

An aggressive subtype of peripheral mature T-cell lymphoma characterized by generalized lymphadenopathy, constitutional symptoms (fever, night sweats, weight loss), hepatosplenomegaly, skin rashes, and polyclonal hypergammaglobulinemia.

**Alternate Names:**

AITL

**Angiokeratoma of the Scrotum**

A benign vascular papule caused by superficial capillary dilation and overlying thickened skin, usually appearing as multiple dark-red, blue, or purple papules on the scrotum. Lesions may bleed after friction and can be mistaken for melanoma or other lesions; examination establishes the diagnosis.

**Angiolymphoid Hyperplasia With Eosinophilia**

An uncommon benign vascular and inflammatory lesion, also called epithelioid hemangioma, that produces solitary or grouped red-brown papules or nodules, usually on the head and neck near the ears or scalp. Eosinophilia may occur; biopsy distinguishes it from Kimura disease and other vascular or neoplastic lesions.

**Angiomyofibroblastoma**

A usually noncancerous, slow-growing tumour of blood-vessel and connective-tissue cells, most often occurring as a small, well-defined, painless mass in the vulva or vagina.

**Angiomyolipoma**

A usually noncancerous kidney tumour made of blood vessels, muscle, and fat. It may occur alone or with tuberous sclerosis and can bleed, especially when large.

**Angiopathy**

Any disease or abnormal change in blood vessels. Causes include diabetes, inflammation, fatty deposits, injury, infection, and inherited disorders; effects depend on the vessels and organs involved.

**Angioplasty**

A catheter procedure used to widen a narrowed or blocked blood vessel, most often an artery. A catheter with a small, deflated balloon is guided to the narrowed area, usually with imaging. The balloon is inflated to widen the vessel, then deflated and removed. A stent may be placed to help keep the vessel open. Angioplasty is used in coronary and other arteries and, in some cases, veins.

**Alternate Names:**

---

Balloon Angioplasty; Percutaneous Transluminal Angioplasty

### **Angioplasty Balloon**

An inflatable cylindrical balloon catheter designed to dilate stenotic vascular segments, fracture atherosclerotic plaques, or deploy and seat endovascular stents.

#### **Alternate Names:**

Balloon Catheter; Dilatation Balloon; Angioplasty Catheter

### **Angiosarcoma**

A rare, fast-growing cancer that begins in cells lining blood or lymph vessels. It may occur in the skin, breast, liver, or bone and can spread to other organs.

### **Angiosarcoma Of The Breast**

A rare cancer arising from blood-vessel cells in the breast. It may cause an enlarging breast mass or bruise-like skin colour change and can develop after breast radiation or long-term lymphoedema.

### **Angiosarcoma Of The Heart**

A rare, aggressive cancer arising from blood-vessel cells in the heart, most often in the right upper chamber. It may obstruct blood flow, cause fluid around the heart or abnormal rhythms, and spread to other organs.

### **Angiotensin**

A group of hormones that help regulate blood pressure and the body's salt and water balance. Angiotensin II narrows blood vessels and signals the kidneys and adrenal glands to retain salt and water.

### **Angiotensin Converting Enzyme**

An enzyme that changes angiotensin I into angiotensin II, which raises blood pressure by narrowing blood vessels. The enzyme also breaks down bradykinin, a substance that can widen vessels.

### **Angiotensin I**

A mostly inactive hormone made from angiotensinogen. Angiotensin-converting enzyme changes it into

angiotensin II, which narrows blood vessels and promotes salt and water retention.

### **Angiotensin II**

A hormone that narrows blood vessels and causes the body to retain salt and water. These effects increase blood pressure and help maintain blood flow when circulating volume is low.

### **Angiotensin Receptor Blockers**

Medicines that prevent angiotensin II from activating its main receptor. They relax blood vessels and reduce salt and water retention, lowering blood pressure and reducing strain on the heart and kidneys.

### **Angiotensin-Converting Enzyme 2**

A cell-surface zinc metalloprotease that cleaves angiotensin II into the vasodilatory and cardioprotective peptide angiotensin-(1-7), acting as an essential counter-regulatory enzyme within the renin-angiotensin-aldosterone system; also serves as the host cell receptor for the SARS-CoV-2 spike protein.

#### **Alternate Names:**

ACE2

### **Angle-Closure Glaucoma**

A type of glaucoma in which the outer edge of the iris blocks the eye's drainage angle. Fluid then builds up, causing eye pressure to rise and potentially damaging the optic nerve and vision. It may develop suddenly with severe eye pain, blurred vision, halos, headache, nausea, or vomiting, or develop slowly without symptoms.

### **Angular Cheilitis**

Inflammation, soreness, and cracking at one or both corners of the mouth. Moisture and saliva, yeast or bacterial infection, dentures, irritation, or iron and vitamin deficiencies can contribute.

### **Anhedonia**

Reduced or absent ability to feel pleasure or interest in activities that were previously enjoyable. It can occur with depression, schizophrenia, substance use, and other conditions.

---

## Anhidrosis

Absent or greatly reduced sweating. It may result from nerve damage, skin disease, dehydration, medicines, or an inherited condition. Without enough sweat, the body may overheat dangerously, especially during exercise or hot weather.

## Animal-Assisted Therapy

A planned healthcare intervention in which a trained animal is used with a qualified professional to support goals such as improving mood, communication, movement, or participation. Safety, hygiene, and allergies must be considered.

## Anion

A particle with a negative electrical charge. Important anions in body fluids include chloride, bicarbonate, and phosphate, which help maintain fluid balance, electrical activity, and the blood's acid level.

## Anion Gap

A value calculated from several blood salts to estimate acids that were not directly measured. A high result can suggest excess acid in the blood, but it must be interpreted with albumin, other tests, symptoms, and the laboratory's method.

## Anion Gap Metabolic Acidosis

A subtype of metabolic acidosis characterized by an elevated serum anion gap (exceeding 12 mEq/L), signifying the accumulation of unmeasured organic or inorganic fixed acids in plasma, such as ketoacids in diabetic ketoacidosis, lactic acid, or exogenous toxic metabolites.

### Alternate Names:

AGMA; High Anion Gap Metabolic Acidosis

## Aniridia

A congenital, bilateral, complete or partial absence of the iris, typically caused by mutations in the PAX6 gene. It is accompanied by photophobia, nystagmus, reduced visual acuity, foveal hypoplasia, and an increased risk of early-onset glaucoma and cataracts, and may occur as part of the WAGR syndrome.

## Anisocoria

An unequal size of the pupils, the black centers of the eyes. In about 20% of healthy people, a slight difference is natural and harmless. When abnormal, it is caused by eye injury, certain eye drops, or nerve disorders that prevent a pupil from widening or shrinking normally. Doctors test how the pupils react in bright light and darkness to identify the underlying cause.

## Anisocytosis

Greater-than-usual variation in the size of red blood cells. It is a laboratory finding rather than a disease and may occur with iron deficiency, vitamin deficiency, or other types of anemia.

## Anisometropia

A difference in the focusing power of the two eyes. It can cause unequal image size, blurred vision, eyestrain, or reduced development of vision in one eye during childhood.

## Ankle

The joint connecting the lower leg and foot. It is formed mainly by the tibia, fibula, and talus and allows the foot to move up and down while ligaments provide stability.

## Ankle Arthroplasty

Surgical replacement of the ankle joint with an artificial joint, usually for severe painful arthritis. It aims to reduce pain while preserving movement; suitability depends on bone, alignment, activity, and surrounding-joint condition.

## Ankle Arthroscopy

A procedure using a small camera and instruments inserted through small cuts around the ankle to diagnose or treat problems such as loose bone or cartilage, inflamed lining, or tissue impingement.

## Ankle Bone

Any of the bones forming the ankle joint, chiefly the lower ends of the tibia and fibula and the upper part of the talus.

---

## Ankle Dislocation

Complete displacement of the ankle bones from their normal joint position, usually after a severe injury. It often occurs with a fracture or ligament injury and can damage blood vessels or nerves, so it requires urgent reduction and imaging.

### Alternate Names:

Dislocated Ankle; Talotibial Dislocation

## Ankle Fracture

A break in one or more bones forming the ankle, usually the tibia, fibula, or talus. It may cause pain, swelling, bruising, and inability to bear weight; displaced or unstable fractures need prompt assessment.

## Ankle Fractures

Breaks in one or more bones forming the ankle joint. They may also injure ligaments, blood vessels, or nerves; the seriousness depends on bone alignment, joint stability, and whether the skin is broken.

## Ankle Joint

The hinge-like joint between the lower ends of the tibia and fibula and the talus. It mainly moves the foot upward and downward and is supported by surrounding ligaments.

## Ankle Pain

Pain in or around the ankle, commonly caused by a sprain, fracture, tendon problem, arthritis, bursitis, or infection. Deformity, marked swelling, fever, numbness, or inability to bear weight requires prompt assessment.

## Ankle Replacement

Surgery that replaces the damaged surfaces of the ankle joint with an artificial joint, usually for severe arthritis causing persistent pain and disability. It aims to reduce pain and preserve ankle movement.

## Ankle Sprain

An injury in which one or more ankle ligaments are stretched or torn, most often on the outer side after the foot twists inward. Pain, swelling, bruising, and

a feeling of instability may occur.

## Ankle Swelling

Enlargement around the ankle caused by injury, inflammation, infection, vein or lymph problems, or general fluid retention. Sudden one-sided swelling with pain, warmth, fever, or breathlessness needs urgent assessment.

## Ankle-Brachial Index

A comparison of blood pressure measured at the ankle with blood pressure measured at the arm. A low result may indicate narrowed leg arteries; a very high result may mean the arteries are too stiff to measure reliably.

## Ankle-Foot Orthosis

A brace that supports the ankle and foot and controls alignment or movement during standing and walking. It may improve walking or prevent foot drop caused by weakness, spasticity, instability, or deformity.

## Ankyloblepharon

Abnormal sticking or joining of the upper and lower eyelid edges. It may be present at birth or follow inflammation, burns, injury, or surgery and can partly block vision when severe.

## Ankyloglossia

Alternate name; see *Tongue-Tie*.

## Ankylose

To become stiff or unable to move at a joint, sometimes because scar tissue or bone joins the joint surfaces.

## Ankylosing

Causing or involving progressive stiffness or loss of movement in a joint, sometimes because scar tissue or bone joins the joint surfaces.

## Ankylosing Spondylitis

A long-term inflammatory arthritis that mainly affects the joints between the spine and pelvis (sacroil-

---

iac joints) and the spine itself. It usually causes lower-back or hip pain and stiffness that is worse after rest or inactivity and may improve with movement. Inflammation can reduce spinal movement and, in severe cases, cause vertebrae to fuse. The hips, ribs, knees, or feet may also be affected. Some people develop eye inflammation (uveitis), psoriasis, or inflammatory bowel disease. HLA-B27 increases risk but does not by itself cause the disease.

**Alternate Names:**

Bekhterev Disease; Radiographic Axial Spondyloarthritis; Marie-Strümpell Disease

### Ankylosis

Abnormal reduction or complete loss of movement in a joint. It may result from fibrous scar tissue, bone fusion, injury, infection, chronic inflammation, or surgery. Fibrous ankylosis involves scar tissue, while bony ankylosis involves fusion of neighboring bones. In dentistry, ankylosis may describe fusion of a tooth root to the surrounding jawbone.

### Ankyrin

A protein that connects selected cell-surface proteins to the cell's internal support framework. It helps red blood cells keep their shape; inherited changes can make the cells fragile and cause anemia.

### Annihilation

The conversion of a particle and its opposite partner into energy when they collide. In PET scanning, a positron meets an electron and produces two gamma rays that travel in opposite directions to create the image.

### Annular Pancreas

A birth defect in which pancreatic tissue forms a ring around part of the duodenum, the first section of the small intestine. The ring may narrow the intestine and cause vomiting, abdominal swelling, or feeding problems.

### Annulus

A ring-shaped structure or band in the body. Examples include the tough outer ring of an intervertebral disc and the fibrous rings surrounding the

heart valves.

### Annulus Fibrosus

The tough, outer fibrocartilaginous ring of an intervertebral disc composed of concentric lamellae of collagen fibers, which encircles and restrains the central gel-like nucleus pulposus and absorbs rotational and compressive mechanical loads.

**Alternate Names:**

Anulus Fibrosus

### ANOCA

Chest pain or other angina symptoms when the major heart arteries do not have a severe blockage. The symptoms may result from poor function of the heart's small vessels or temporary spasm of a coronary artery.

### Anodontia

Complete absence of teeth from birth, caused by failure of tooth development. It is rare and may occur with inherited conditions affecting the skin, hair, nails, and teeth.

### Anogenital Tract

The connected region containing the anus, tissues around the anus, genital organs, and nearby skin and moist linings.

### Anomalous Anatomy

A difference from the usual structure, number, position, origin, or pathway of an organ, blood vessel, duct, or other body part. It may cause no symptoms or may affect health, diagnosis, or surgery.

### Anomalous Coronary Artery

A birth defect in which one of the heart's arteries has an unusual starting point or path. Some arteries pass between the aorta and the pulmonary artery and may be squeezed during hard exercise. This can reduce blood flow to the heart and may cause chest pain, fainting, abnormal heart rhythms, or rarely sudden cardiac death.

**Alternate Names:**

AAOCA; Anomalous Aortic Origin of a Coro-



---

nary Artery; Congenital Coronary Artery Anomaly; ACAOS

### **Anomaly**

A difference from the usual structure, development, function, or measurement of the body. It may be harmless, or it may cause symptoms or affect health.

### **Anomaly Scan**

A detailed ultrasound examination usually performed around 18–22 weeks of pregnancy to assess fetal growth and organs, the placenta, and amniotic fluid, and to screen for major structural birth defects.

#### **Alternate Names:**

Mid-Pregnancy Ultrasound; 20-Week Scan; Fetal Anomaly Scan

### **Anomia**

Difficulty finding or saying a familiar word, especially the name of an object or person, even though its meaning is known. It can occur after a stroke or with other brain or language disorders.

### **Anophthalmia**

A birth defect in which one or both eyes fail to develop. It may occur with other birth defects and can affect growth of the eye socket and face.

### **Anorchia**

Complete absence of testicular tissue, present from birth or caused by loss of testes that had formed. It causes very low or absent testosterone production and infertility unless appropriate specialist care is provided.

### **Anorectal Abscess**

A collection of pus in tissues around the anus or rectum, usually caused by an infected anal gland. It may cause severe pain, swelling, fever, or difficulty sitting and often needs prompt drainage.

### **Anorectal Fistula**

An abnormal tunnel between the anus or rectum and nearby skin or another organ. It often develops after

an abscess or infection and may cause persistent drainage, pain, or repeated abscesses.

#### **Alternate Names:**

Anal Fistula

### **Anorectal Infection**

Infection of the anus, rectum, or nearby tissues. It may cause pain, swelling, discharge, fever, or difficulty passing stool and can lead to an abscess that requires drainage.

### **Anorectal Malformation**

A birth defect in which the anus or rectum does not form normally or does not connect correctly to the skin. The opening may be absent, misplaced, or abnormally connected to another organ.

### **Anorectal Manometry**

A test that measures pressure and sensation in the rectum and anal canal. A thin pressure-sensitive tube assesses how well the muscles relax and tighten and may help investigate constipation or loss of bowel control.

### **Anorexia**

Loss or reduction of appetite. It can result from illness, pain, medicines, emotional distress, or other causes and is different from anorexia nervosa, an eating disorder.

### **Anorexia Nervosa**

A serious eating disorder involving persistent restriction of food, intense fear of gaining weight, and a distorted view of body size or shape. It can cause severe weight loss, malnutrition, slow heart rate, low blood pressure, and organ damage.

#### **Alternate Names:**

Anorexia; Restrictive Eating Disorder

### **Anorgasmia**

Persistent or repeated difficulty reaching orgasm or inability to reach orgasm despite adequate sexual stimulation. It may result from medicines, illness, hormonal changes, pain, stress, trauma, or relationship factors.

---

## Anorgasmy

Another spelling of anorgasmia: persistent or repeated inability to reach orgasm despite adequate sexual stimulation. Possible causes include medicines, illness, hormonal changes, stress, trauma, or relationship factors.

## Anoscopy

Examination of the anal canal using a short, lighted tube called an anoscope. It helps investigate rectal bleeding, haemorrhoids, fissures, fistulas, pain, and other problems near the anus.

## Anoscopy And High-Resolution Anoscopy

Examination of the anal canal and lower rectum with a short viewing tube. High-resolution anoscopy adds magnification and special lighting to identify and assess abnormal tissue, including changes related to HPV.

## Anosmia

Complete loss of the sense of smell. It may follow a viral infection, blocked or inflamed nasal passages, head injury, medicines, toxic exposure, or a neurological or inherited disorder.

## Anosodiaphoria

A neurological state of apparent emotional indifference or unconcern toward severe physical disability, such as hemiplegia, despite intellectual awareness and verbal acknowledgement of the deficit. It is most commonly associated with non-dominant (right) hemisphere parietal lobe lesions.

## Anosognosia

Lack of awareness of, or inability to recognize, one's own illness or neurological impairment. It can occur after certain brain injuries or strokes and is not simply deliberate denial.

## Anotia

A birth defect in which the outer ear is completely absent. The ear canal and middle-ear structures may also be abnormal, causing reduced hearing on the affected side.

## Anovulation

Failure to release an egg from an ovary during a menstrual cycle. It can cause irregular or absent periods, abnormal bleeding, or infertility and may result from hormonal disorders, low body weight, stress, excessive exercise, or polycystic ovary syndrome.

## Anoxia

Complete lack of oxygen reaching body tissues. It can quickly damage or kill cells, especially in the brain, heart, and kidneys, and may result from blocked blood flow, breathing failure, or severe poisoning.

## Anrep Effect

An intrinsic physiological autoregulation mechanism of the myocardium whereby an abrupt increase in ventricular afterload causes an initial stretch followed by a sustained, positive inotropic increase in ventricular contractility over the ensuing several minutes.

## Anserine Bursa

A synovial fluid-filled sac situated on the proximal medial aspect of the tibia, interposed between the tibial collateral ligament and the conjoined tendons of the sartorius, gracilis, and semitendinosus muscles; inflammation causes pes anserine bursitis.

### Alternate Names:

Pes Anserinus Bursa

## Antacids

Medicines that neutralise stomach acid and provide short-term relief from heartburn, indigestion, and acid reflux. Different products contain different minerals and may interact with other medicines or alter bowel movements.

## Antagonist

A drug or other substance that reduces or blocks the effect of another substance. A receptor antagonist binds to a receptor without activating it and reduces the response to an agonist, often by preventing the agonist from binding or signaling.

---

## Antagonist Muscle

A muscle that produces the opposite movement to the main working muscle and helps control or reverse that movement.

## Antalgic Gait

An abnormal, compensatory walking pattern adopted to avoid or minimize pain in a weight-bearing lower limb or spinal structure. It is characterized by a significantly shortened stance phase on the affected, painful limb and a rapid, hurried transition to the contralateral unaffected limb, frequently accompanied by a compensatory trunk lean toward the affected side to reduce joint contact forces.

### Alternate Names:

Pain-Relieving Gait; Limping Gait; Antalgic Walk

## Antebrachium

The anatomical region of the upper limb situated between the elbow and the wrist, consisting of the radius and ulna surrounded by anterior and posterior muscular compartments.

### Alternate Names:

Forearm

## Antecubital Fossa

The shallow hollow at the front of the elbow. It contains the biceps tendon, a major artery, and a major nerve and is commonly used to draw blood or measure blood pressure.

## Antegrade

Moving or occurring in the normal forward direction, such as blood flowing away from the heart or an electrical signal travelling through its usual pathway.

## Antemortem Injury

An injury that occurred before death. It may show bleeding into living tissue, clot formation, inflammation, swelling, or early healing, which can distinguish it from damage occurring after death.

## Antenatal

Relating to the period before birth or to care during pregnancy. Antenatal care monitors the pregnant

person and fetus, screens for complications, and provides health advice and support.

### Alternate Names:

Prenatal; Antepartum

## Antepartum Fetal Surveillance

Tests used before birth to check fetal well-being, especially in higher-risk pregnancies. They may include monitoring fetal movements, the fetal heart rate, ultrasound assessment, and blood-flow measurements.

## Antepartum Hemorrhage

Bleeding from the vagina during the second half of pregnancy and before birth. Causes include placenta previa, placental abruption, and problems with the cervix or vagina; any significant bleeding needs urgent assessment.

## Anterior

Relating to the front of the body or positioned nearer the front than another structure. The term is used to describe body parts, scan findings, injuries, and disease locations.

## Anterior Cerebral Artery Syndrome

A stroke syndrome resulting from occlusion or ischemia of the anterior cerebral artery, classically causing contralateral motor weakness and sensory loss predominantly affecting the lower extremity more than the upper extremity and face. Additional findings may include abulia, urinary incontinence, and frontal release signs due to medial frontal lobe involvement.

### Alternate Names:

ACA Syndrome; ACA Infarction

## Anterior Chamber

The fluid-filled space between the cornea and iris at the front of the eye. It contains aqueous fluid, which nourishes the cornea and lens and helps maintain eye pressure; impaired drainage can contribute to glaucoma.

---

### Anterior Chamber Angle

The area where the cornea and iris meet at the front of the eye. It contains the main drainage tissue for aqueous fluid; blockage here can raise eye pressure and cause glaucoma.

### Anterior Commissure

A transverse bundle of myelinated nerve fibers crossing the midline anterior to the columns of the fornix and above the lamina terminalis, interconnecting the olfactory bulbs, amygdalae, and middle and inferior temporal gyri across the cerebral hemispheres.

### Anterior Cord Syndrome

A spinal-cord injury that weakens movement and reduces pain and temperature sensation below the injury, while vibration and awareness of limb position may be relatively preserved.

### Anterior Cruciate Ligament

A strong band of tissue inside the knee that connects the thigh bone to the shin bone. It helps prevent the shin bone from sliding too far forward and controls knee rotation.

### Anterior Cruciate Ligament Injury

Damage or tearing of the main stabilizing ligament inside the knee, often after sudden turning, stopping, landing, or overextension. It may cause a popping sensation, rapid swelling, pain, and a feeling that the knee gives way.

### Anterior Cruciate Ligament Tear

A partial or complete tear of the ligament inside the knee that limits forward sliding and rotation of the shin bone. It commonly causes a pop, rapid swelling, pain, and knee instability after a sudden twist or landing.

### Anterior Drawer Sign

A physical check for a torn knee ligament (anterior cruciate ligament). With the knee bent, the doctor gently pulls the shin forward; moving too far forward means the ligament is likely torn or stretched.

### Anterior Drawer Test

A clinical orthopedic physical examination maneuver used to assess the integrity of the anterior cruciate ligament (ACL) of the knee or the anterior talofibular ligament of the ankle. Excessive anterior translation of the tibia relative to the femur with a soft or mushy endpoint indicates an ACL tear.

#### Alternate Names:

Anterior Drawer Sign

### Anterior Interosseous Nerve

A pure motor branch of the median nerve arising in the proximal forearm that courses deeply along the anterior surface of the interosseous membrane to innervate the flexor pollicis longus, the lateral half of flexor digitorum profundus, and the pronator quadratus; damage produces the classic inability to make the "OK" sign.

#### Alternate Names:

AIN

### Anterior Interosseous Nerve Syndrome

A pure motor neuropathy of the forearm resulting from compression or neuritis of the anterior interosseous nerve (a motor branch of the median nerve). Causes weakness of the flexor pollicis longus, pronator quadratus, and the radial half of the flexor digitorum profundus, classically producing an inability to form the OK sign (pinching index finger and thumb pulp-to-pulp instead of tip-to-tip).

#### Alternate Names:

Kiloh-Nevin Syndrome; AIN Neuropathy

### Anterior Knee Pain

Pain at the front of or behind the kneecap, often worse with stairs, squatting, running, or prolonged sitting. Common causes include irritation around the kneecap, tendon problems, arthritis, and injury.

### Anterior Mediastinum

The front part of the lower mediastinum, located behind the sternum and in front of the heart sac. It contains fat, connective tissue, lymph nodes, and part of the thymus.

---

### Anterior Midline

The imaginary vertical line running down the center of the front of the body, separating the right and left sides.

### Anterior Myocardial Infarction

A heart attack affecting the front wall of the main pumping chamber, usually because the left anterior descending coronary artery is blocked. It can seriously weaken the heart and requires emergency treatment.

#### Alternate Names:

Anterior Heart Attack; Anterior MI

### Anterior Pituitary

The front part of the pituitary gland. It releases hormones that control the adrenal and thyroid glands, growth, milk production, egg or sperm production, and reproductive function.

### Anterior Talofibular Ligament

A short, flat ligament on the lateral aspect of the ankle that spans from the anterior margin of the lateral malleolus of the fibula to the neck of the talus. It is the primary stabilizer against anterior talar displacement and excessive inversion, and represents the most frequently torn ligament in lateral ankle sprains.

#### Alternate Names:

ATFL

### Anterior Tibial Artery

An artery that carries blood down the front of the leg and continues onto the top of the foot as the dorsalis pedis artery. Its pulse can usually be felt near the front of the ankle.

### Anterior Uveitis

Inflammation in the front part of the eye, usually involving the iris and sometimes nearby tissue. It can cause eye pain, redness, light sensitivity, blurred vision, and floating spots.

#### Alternate Names:

Iritis; Iridocyclitis

### Anterior Vaginal Prolapse

Bulging of the anterior vaginal wall into the vaginal vault, commonly resulting from herniation of the urinary bladder (cystocele) due to attenuation or tearing of the pubocervical fascia and pelvic floor musculature. Symptoms may include a palpable vaginal bulge, pelvic heaviness, urinary frequency, urgency, incomplete emptying, or stress incontinence.

#### Alternate Names:

Cystocele; Bladder Prolapse; Prolapsed Bladder; Anterior Wall Prolapse

### Anterior Vaginal Wall Repair

Surgery to strengthen the front wall of the vagina when the bladder bulges into it. The procedure restores support using the person's own tissues; the approach and risks depend on the prolapse and overall health.

### Anterior/Posterior Drawer Test

A physical examination of knee stability. Pulling the shin bone forward tests the anterior cruciate ligament; pushing it backward tests the posterior cruciate ligament.

### Anterograde Amnesia

A selective impairment of memory in which an individual is unable to form, encode, or consolidate new long-term declarative (episodic and semantic) memories after the onset of a neurological insult, while remote memories formed prior to the event remain largely intact. It typically results from bilateral damage to the medial temporal lobes, hippocampi, mammillary bodies, or diencephalic structures following traumatic brain injury, stroke, herpes simplex encephalitis, or Korsakoff syndrome.

#### Alternate Names:

Post-Traumatic Amnesia; Anterograde Memory Loss; Forward-Acting Amnesia

### Anterolisthesis

Forward slipping of one spinal bone relative to the bone below it. It may result from wear, a stress fracture, injury, or a problem present from birth and may cause back pain or nerve symptoms.



---

## Anteroposterior

Directed from the front of the body toward the back. In an anteroposterior X-ray view, the beam enters through the front and leaves through the back.

## Anthocyanin

A natural plant pigment that gives many fruits, vegetables, and flowers red, purple, or blue colours. It has antioxidant effects in laboratory studies and is found in foods such as berries, grapes, and red cabbage.

## Anthracosis

A benign, non-fibrotic form of pneumoconiosis characterized by the asymptomatic accumulation of inhaled carbon and soot pigment in the lung parenchyma, alveolar macrophages, and draining hilar lymph nodes, commonly seen in urban dwellers and cigarette smokers.

### Alternate Names:

Black Lung Pigmentation; Carbon Pigmentation

## Anthrax

A serious infection caused by the bacterium *Bacillus anthracis*. People can become infected through contact with infected animals or animal products, breathing in spores, or eating contaminated food; skin, lung, or intestinal disease may result.

## Anthrax Blood Test

A laboratory test that looks for the anthrax bacterium or the body's immune response to it in blood or another sample. It helps confirm suspected anthrax, which can affect the skin, lungs, or digestive tract.

## Anthropometry

The measurement of body size and proportions, such as height, weight, waist size, limb length, and skinfold thickness. These measurements help assess growth, nutrition, and body composition.

## Anti-Anxiety Agents

Medicines used to reduce excessive anxiety or panic. Different medicines work in different ways and may

cause sleepiness, dependence, or other side effects; the choice depends on the person and the condition.

### Alternate Names:

Anxiolytics

## Anti-CCP

A blood-test antibody marker associated with rheumatoid arthritis. A positive result supports the diagnosis when symptoms fit, but it does not prove that rheumatoid arthritis is present or predict its course by itself.

## Anti-CCP Antibody

An antibody measured in blood that targets proteins changed by a process called citrullination. It can support a diagnosis of rheumatoid arthritis, but results must be interpreted with symptoms and other tests.

## Anti-Citrullinated Peptide Antibody

Alternate index label; see *Anti-Cyclic Citrullinated Peptide Antibody*.

## Anti-Cyclic Citrullinated Peptide Antibody

An autoantibody directed against citrulline-containing peptide antigens, possessing high diagnostic specificity (greater than 95 percent) for rheumatoid arthritis and correlating with aggressive, erosive joint disease.

### Alternate Names:

Anti-CCP Antibody; ACPA; Anti-Citrullinated Peptide Antibody; Anti-Cyclic Citrullinated Peptide; Cyclic Citrullinated Peptide Antibody

## Anti-D

An injection containing antibodies given to an RhD-negative pregnant person after certain pregnancy events, or around birth, to prevent the person's immune system making antibodies against RhD-positive fetal blood. This helps protect a current or future pregnancy from fetal red-cell destruction.

## Anti-DNase B Blood Test

A blood test that measures the body's antibodies

---

against a protein made by group A streptococcal bacteria. It can support evidence of a recent strep infection when doctors investigate rheumatic fever, kidney inflammation, or another complication.

### **Anti-Double-Stranded DNA Antibody**

An antinuclear autoantibody specifically directed against the native double-helical structure of deoxyribonucleic acid, highly specific for systemic lupus erythematosus and closely correlating with lupus nephritis disease activity and flares.

#### **Alternate Names:**

Anti-dsDNA Antibody; dsDNA Antibody; Anti-Double-Stranded DNA; Double-Stranded DNA Antibody

### **Anti-GBM Disease**

An autoimmune disease in which antibodies attack supporting layers in the kidneys and sometimes the lungs. It can rapidly cause kidney failure, coughing up blood, or bleeding into the lungs and requires urgent specialist care.

### **Anti-Glomerular Basement Membrane Blood Test**

A blood test that detects antibodies attacking the kidney's filtering membranes and sometimes the lungs' supporting membranes. A positive result supports anti-GBM disease when symptoms and other test results are consistent.

### **Anti-Glomerular Basement Membrane Disease**

An autoimmune disease in which antibodies damage the kidney's filtering membranes and sometimes the lungs. It may cause rapidly worsening kidney function, blood in the urine, coughing up blood, or lung bleeding.

### **Anti-HMGCR Myopathy**

A rare autoimmune muscle disease in which antibodies attack a muscle enzyme. It causes ongoing weakness and muscle injury, especially around the hips, thighs, shoulders, and upper arms, and may begin after treatment with a statin medicine.

#### **Alternate Names:**

Statin-Induced Immune-Mediated Necrotizing Myopathy; Anti-HMG-CoA Reductase Myopathy

### **Anti-IgE Therapy**

Treatment using an antibody that binds free immunoglobulin E, an allergy-related immune protein. This reduces activation of allergy cells and may lessen symptoms and attacks in selected allergic asthma and other allergic conditions.

### **Anti-IL4 Receptor Therapy**

Treatment that blocks a cell receptor used by interleukin-4 and interleukin-13, immune signals that promote allergic inflammation. It can reduce airway inflammation in selected asthma and other allergic diseases.

### **Anti-IL5 Therapy**

Treatment that blocks interleukin-5 or its receptor, reducing the production or survival of eosinophils, a type of white blood cell. It is used for selected eosinophilic diseases, including some severe asthma.

### **Anti-Inflammatory**

Pertaining to any drug, substance, or therapeutic intervention that reduces, counteracts, or suppresses inflammation, vascular permeability, tissue edema, and associated pain. Major therapeutic classes include nonsteroidal anti-inflammatory drugs and glucocorticoids.

### **Anti-Insulin Antibody Test**

A blood test that detects antibodies against insulin. These antibodies can bind insulin and release it unpredictably, causing low blood sugar; the test helps investigate unexplained hypoglycaemia and insulin autoimmune syndrome.

### **Anti-MDA5 Dermatomyositis**

A form of dermatomyositis associated with antibodies against a protein called MDA5. It often causes distinctive skin changes with little muscle weakness but carries a high risk of rapidly worsening inflammation and scarring in the lungs.

---

### Anti-Mullerian Hormone

A hormone made by developing ovarian follicles and by cells in the testes before birth. Its blood level can help estimate ovarian follicle number or assess some reproductive and developmental conditions, but it does not measure fertility by itself.

### Anti-NMDA Receptor Encephalitis

A serious immune attack on the brain involving a receptor important for communication between nerve cells. It may cause sudden behavior or psychiatric changes, memory loss, seizures, abnormal movements, reduced consciousness, and unstable heart rate or blood pressure. Urgent hospital treatment is needed.

#### Alternate Names:

Anti-N-Methyl-D-Aspartate Receptor Encephalitis; NMDA Receptor Encephalitis

### Anti-Reflux Surgery

An operation to reduce the backward flow of stomach contents into the oesophagus by strengthening or reshaping the valve between the oesophagus and stomach. It may be considered for selected people with severe reflux or a hiatus hernia.

### Anti-Rust Product Poisoning

Illness caused by swallowing, breathing, or skin exposure to a product used to remove or prevent rust. It may burn the mouth and digestive tract or damage the lungs and other organs; significant exposure requires urgent medical treatment.

### Anti-Smooth Muscle Antibody

An antibody that reacts with proteins in smooth muscle cells. A blood test may support autoimmune hepatitis, but low levels can also occur with other autoimmune diseases or infections and are not diagnostic alone.

### Anti-SRP Myopathy

A rare autoimmune muscle disease caused by antibodies against a protein involved in making new proteins. It causes rapidly progressive weakness, especially around the hips, shoulders, and neck, and may cause difficulty swallowing.

#### Alternate Names:

Anti-Signal Recognition Particle Myopathy; SRP-Associated Necrotizing Myopathy

### Anti-Thyroid Peroxidase Antibody

An autoantibody directed against thyroid peroxidase, the key enzyme required for thyroid hormone synthesis. Elevated titers are the hallmark laboratory finding in Hashimoto thyroiditis and are also present in Graves disease.

#### Alternate Names:

Anti-TPO Antibody; Thyroid Peroxidase Antibody

### Anti-VEGF Injections

Injections, usually given into the eye, that block vascular endothelial growth factor, a signal that promotes abnormal blood-vessel growth and leakage. They can improve or preserve vision in several retinal diseases.

### Antiandrogen

A medicine or substance that lowers the production or blocks the effects of androgen hormones such as testosterone. It is used for selected prostate conditions, androgen-related disorders, and some gender-affirming treatments.

### Antianginal

A medicine used to prevent or relieve angina, the chest discomfort caused by reduced blood flow to heart muscle. Different medicines lower the heart's workload, widen blood vessels, or improve coronary blood flow.

### Antiarrhythmic

A medicine used to prevent or treat certain abnormal heart rhythms. Some medicines aim to restore or maintain a normal rhythm, while others control the heart rate without restoring normal rhythm. The choice depends on the rhythm, heart condition, and risks, because these medicines can also cause abnormal rhythms.

#### Alternate Names:

Antiarrhythmics; Antiarrhythmic Medicines

---

## Antibacterial

Able to kill bacteria or stop them from growing. The term may describe a medicine, disinfectant, surface treatment, or part of the body's immune defence.

## Antibiogram

A report showing which antibiotics are likely to work against bacteria grown from patient samples in a laboratory. It helps clinicians choose treatment and monitor local patterns of antibiotic resistance.

## Antibiotic Misuse in Viral URIs

The inappropriate prescription or self-administration of antibacterial agents for acute viral upper respiratory tract infections (such as the common cold or influenza), which confers no therapeutic benefit and drives antimicrobial resistance and adverse drug events.

### Alternate Names:

Antibiotic Stewardship in URI; Inappropriate Antibiotic Use in Colds; Antibiotic Overuse in Flu

## Antibiotic-Resistant Bacterial Infection

An infection caused by bacteria that are not adequately stopped by one or more antibiotics normally active against them. Resistance may be intrinsic or acquired and can limit effective treatment options.

## Antibiotics

Medicines that kill bacteria or stop them from multiplying. Different antibiotics work in different ways, such as damaging the bacterial cell wall or blocking the production of proteins or DNA. They treat susceptible bacterial infections but do not treat viral infections such as colds or influenza. Incorrect or unnecessary use can cause side effects and antibiotic resistance.

## Antibodies

Proteins made by immune cells that recognize and attach to specific foreign or abnormal substances. They can block germs or toxins, mark them for destruction, and activate other parts of the immune system.

## Antibody

A protein made by the immune system that attaches specifically to a foreign substance, germ, toxin, or abnormal cell and may block it or mark it for removal.

## Antibody Engineering

Designing or changing antibodies so they bind more accurately, last longer, cause fewer unwanted immune reactions, or work better as medicines. Researchers may alter their target, size, immune effects, or attachment to another medicine.

## Antibody Therapy

Treatment using specially prepared antibodies to block, activate, or target a particular molecule or cell. It can be used for cancer, autoimmune disease, infection, inflammation, and other conditions.

### Alternate Names:

Immunotherapy

## Antibody Titer Blood Test

A blood test that estimates the amount of a specific antibody by repeatedly diluting the sample until the antibody can no longer be detected. Results may help show previous exposure, immunity, or a recent infection, depending on the antibody tested.

## Antibody-Dependent Cell-Mediated Cytotox.

An immune process in which antibodies coat an infected, abnormal, or cancer cell and attract killer immune cells. The killer cells then damage and destroy the marked cell.

## Antibody-Drug Conjugate

A treatment in which an antibody carrying a linked medicine binds to a target on a cell and delivers the medicine near or inside that cell. It is used for selected cancers and can still damage healthy tissues or cause medicine-specific toxicities.

## Anticholinergic

Blocking the action of acetylcholine, a chemical messenger used by nerves. These medicines can

---

reduce secretions and muscle spasms but may cause dry mouth, blurred vision, constipation, difficulty urinating, fast heartbeat, or confusion.

### **Anticholinergic Burden**

The combined anticholinergic effect of one or more medicines. A high burden can cause dry mouth, constipation, blurred vision, difficulty urinating, confusion, or delirium, especially in older adults.

### **Anticholinergic Toxicity**

Excess blockade of muscarinic acetylcholine receptors, producing a hot, flushed, dry, tachycardic, mydriatic, delirious patient with urinary retention and reduced bowel sounds. Treatment is supportive with cooling and benzodiazepines; physostigmine is reserved for selected severe cases after toxicology assessment.

### **Anticholinergic Toxidrome**

A pattern of poisoning caused by excessive blockage of acetylcholine signals. It may cause dry skin and mouth, large pupils, fast heartbeat, inability to urinate, reduced bowel activity, overheating, agitation, or delirium.

### **Anticoagulants and Thrombolytics in Pregnancy**

Medicines that prevent clot formation or dissolve selected clots during pregnancy or after delivery. Choice and timing depend on maternal indication, placental transfer, bleeding risk, delivery plans, and whether neuraxial anesthesia may be used; specialist management is required.

### **Anticoagulation**

Treatment that reduces the formation or growth of blood clots by slowing the clotting process. It is used for conditions such as clots in veins, irregular heart rhythm, and selected heart or blood-vessel disorders.

### **Anticodon**

A group of three RNA building blocks on a transfer-RNA molecule. It pairs with a matching sequence on messenger RNA so the cell adds the correct amino

acid while making a protein.

### **Anticonvulsants**

Medicines, also called antiseizure medicines, used to prevent or control seizures. Some are also used for nerve pain, migraine, or mood disorders; their effects, interactions, and side effects differ between medicines.

#### **Alternate Names:**

Antiseizure Medicines; Antiseizure Drugs; Anticonvulsant Drugs

### **Antideformity Position**

A planned position for a limb or joint that helps prevent or limit shortening, stiffness, or deformity after injury, surgery, burns, nerve damage, or prolonged immobility.

### **Antidepressants**

Medicines used mainly to treat depression. Different types change the activity of chemical messengers in the brain and may also treat anxiety, nerve pain, obsessive thoughts, or other conditions.

### **Antidiabetic**

Relating to a medicine or treatment used to control diabetes or lower blood glucose; see *Antidiabetic Drugs*.

### **Antidiabetic Drugs**

Medicines used to manage diabetes by replacing insulin, increasing insulin release or action, reducing glucose production or absorption, or helping the kidneys remove glucose. The appropriate medicine depends on the type of diabetes, blood glucose, other health conditions, and treatment goals.

### **Antidiuretic Hormone**

A hormone made in the brain and released by the pituitary gland that helps the kidneys conserve water. It also narrows blood vessels when blood volume or blood pressure is low. It is also called vasopressin.

### **Antidiuretic Hormone Blood Test**

A blood test that measures vasopressin, the hor-



---

none that helps the kidneys conserve water. It may help investigate excessive thirst, very dilute urine, inappropriate water retention, or other disorders of water balance.

### **Antidiuretic Hormone Physiology**

Vasopressin is made in the hypothalamus and released from the posterior pituitary when the blood becomes too concentrated or blood volume falls. It makes kidney tubules retain water and can narrow blood vessels.

### **Antidiuretic Hormone Resistance**

Failure of the kidneys to respond properly to vasopressin. It causes production of large amounts of dilute urine and intense thirst and may result from medicines, mineral imbalance, kidney disease, or inherited changes.

### **Antidote**

A substance that counteracts or limits the harmful effects of a poison. The correct antidote depends on the substance involved; some antidotes block the poison, remove it, or restore a chemical process disrupted by it.

### **Antiemetics**

Medicines used to prevent or relieve nausea and vomiting. Different types act on chemical signals in the brain or digestive tract, and the choice depends on the cause, age, pregnancy, other medicines, and risk of side effects.

### **Antiepileptic Drugs**

Medicines used to prevent or control seizures by reducing excessive electrical activity in the brain. The best choice depends on the seizure type, age, other conditions, medicines, and possible side effects.

### **Antifreeze Poisoning**

Poisoning after swallowing or otherwise being exposed to antifreeze, especially products containing ethylene glycol. It may first cause intoxication and vomiting, then severe acid buildup, kidney failure, brain injury, or death; severe poisoning is an emergency requiring prompt medical treatment.

### **Antifungal Prophylaxis**

Use of an antifungal medicine to prevent a serious fungal infection in someone at high risk, such as a person with prolonged low white-cell counts, a stem-cell transplant, or severe immune suppression.

### **Antifungal Resistance**

The ability of a fungus to survive a medicine that would normally stop its growth or kill it. Resistance may be present naturally or develop after genetic changes or repeated exposure to antifungal medicines.

### **Antigen**

A substance or part of a substance that the immune system can recognize. It may come from a germ, toxin, transplanted tissue, or abnormal cell and can trigger antibodies or other immune responses.

### **Antigen Detection Test**

A test that looks for a protein or other marker from a specific germ in a patient sample. It is often rapid, but it may miss an infection when the amount of germ material is small.

### **Antigen Presentation**

The process by which an immune or other cell displays small pieces of proteins on its surface for T cells to inspect. This helps the immune system recognize infections, abnormal cells, transplanted tissue, and sometimes the body's own tissues.

### **Antigen Receptor**

A protein on an immune cell that recognizes a particular substance or protein fragment. B-cell receptors can bind whole substances, while T-cell receptors recognize fragments displayed by other cells.

### **Antigen-Presenting Cell**

An immune cell that breaks proteins into small pieces and displays them on its surface so T cells can recognize them. Important antigen-presenting cells include dendritic cells, macrophages, and B cells.

**Alternate Names:**



---

APC

### **Antigenic Drift**

The gradual change in a virus's surface proteins as it multiplies. These small changes can help viruses, especially influenza viruses, partly evade existing immunity and contribute to repeated seasonal outbreaks.

### **Antigenic Shift**

A sudden major change in a virus's surface proteins, usually when two different influenza A viruses exchange genetic material inside the same cell. It can produce a virus to which many people have little immunity.

### **Antigenic Variation**

Changes in a germ's surface markers that help it avoid recognition by the immune system. Variation may happen gradually through mutations or suddenly when genetic material is exchanged between different strains.

### **Antigout**

A medicine used to prevent or treat gout. Some lower uric-acid levels over time, while others reduce the inflammation and pain of an acute gout attack.

### **Antihistamines**

Medicines that block histamine, a chemical involved in allergy symptoms. They can relieve itching, sneezing, runny nose, watery eyes, or hives. Older types may cause drowsiness; medicines that reduce stomach acid by blocking H2 receptors are a separate group.

### **Antihistamines For Allergies**

Medicines that block histamine and relieve allergy symptoms such as itching, sneezing, runny nose, watery eyes, or hives. Some cause drowsiness and interactions or side effects vary between medicines.

### **Antihypertensives In Pregnancy**

Medicines used to lower high blood pressure during pregnancy. The choice depends on blood-pressure

severity and pregnancy stage; some commonly used blood-pressure medicines are avoided because they can harm the fetus.

### **Antileukotriene**

A medicine that blocks leukotrienes, chemical signals that can cause airway narrowing and inflammation. It is used in selected asthma and allergy-related conditions, often as part of a broader treatment plan.

#### **Alternate Names:**

Leukotriene Modifier; Leukotriene Pathway Inhibitor

### **Antimetabolite**

A substance that resembles a chemical needed by cells and interferes with the cell's normal chemical reactions or DNA production. Some are used to treat cancer or suppress an overactive immune system.

### **Antimicrobial**

A substance that kills or limits the growth of microscopic organisms, including bacteria, viruses, fungi, or parasites. Antibiotics are antimicrobials, but they act only against bacteria.

### **Antimicrobial Adverse Effects**

Unwanted effects caused by medicines that treat infections. Depending on the medicine, they may include allergy, diarrhoea, liver or kidney injury, hearing problems, tendon injury, or harmful effects on heart rhythm.

### **Antimicrobial Agent**

A substance that kills microorganisms or stops them from growing. Antimicrobial agents include antibacterial, antiviral, antifungal, and antiparasitic medicines.

### **Antimicrobial Desensitization**

A carefully supervised process of giving gradually increasing doses of an infection-fighting medicine to temporarily reduce an immediate allergy response. It is considered only when the medicine is essential and suitable alternatives are unavailable.

---

## Antimicrobial Dosing Optimization

Choosing the infection medicine, dose, timing, route, and duration so enough medicine reaches the infection without unnecessary harm. Kidney function, liver function, body size, infection severity, and resistance are considered.

## Antimicrobial Hypersensitivity Reactions

Immune reactions to an infection-fighting medicine, ranging from an itchy rash to life-threatening anaphylaxis. The timing and symptoms vary with the medicine and type of immune response.

## Antimicrobial Pharmacodynamics

How an infection-fighting medicine affects a microorganism, including how much medicine and how much exposure are needed to stop or kill it. This information helps select an effective dose and schedule.

## Antimicrobial Pharmacokinetics

How the body absorbs, spreads, changes, and removes an infection-fighting medicine. Kidney or liver disease, age, body size, and other medicines can change how much medicine is needed and how often it is given.

## Antimicrobial Resistance

The ability of a microorganism to survive or grow despite a medicine that would normally kill it or stop its growth. Resistance can affect bacteria, viruses, fungi, or parasites and can spread between organisms.

### Alternate Names:

AMR; Drug Resistance

## Antimicrobial Resistance Mechanisms

Ways microorganisms avoid being killed by infection medicines. They may destroy or change the medicine, alter its target, prevent it entering, or actively pump it out of the cell.

## Antimicrobial Resistance Surveillance

Ongoing collection and analysis of resistance results from patient samples. It shows how resistance

changes over time and helps hospitals and public-health services choose effective treatment and limit spread.

## Antimicrobial Sensitivity Test

A laboratory test that shows which infection-fighting medicines can stop or kill a microorganism grown from a patient sample. It helps guide treatment for a confirmed infection.

## Antimicrobial Stewardship

Coordinated efforts to use infection-fighting medicines only when needed and to choose the right medicine, dose, route, and duration. This improves outcomes and limits resistance and side effects.

## Antimicrobial Susceptibility Testing

Laboratory testing that estimates whether a microorganism is likely to respond to particular infection-fighting medicines. Methods include measuring growth around medicine-containing discs or in liquid; results help select treatment.

## Antimicrobial Therapy

Use of antibacterial, antiviral, antifungal, or antiparasitic medicines to prevent or treat an infection. The choice depends on the microorganism, infection site, severity, resistance, and the person's health.

## Antimitochondrial Antibody

An antibody that reacts with structures inside mitochondria, the parts of cells that produce energy. It is strongly associated with primary biliary cholangitis, an autoimmune bile-duct disease, but the result must be interpreted with symptoms and liver tests.

## Antimony

A metallic chemical element whose compounds can be poisonous. Some antimony compounds have been used to treat leishmaniasis; exposure can affect the heart, liver, kidneys, or nervous system.

## Antineoplastic

See *Antineoplastic Drugs*.

### Alternate Names:

---

Anticancer Agent; Antitumor Agent

### Antineoplastic Drugs

Medicines used to treat cancer. They include chemotherapy, targeted medicines, immunotherapies, and hormone therapies, which work in different ways. Treatment is selected according to the cancer and the person's health; benefits and adverse effects vary by drug.

#### Alternate Names:

Antineoplastic Agents; Anticancer Drugs; Anticancer Medicines; Antitumor Drugs

### Antinuclear Antibody

A diverse family of autoantibodies directed against nucleic acids, histones, ribonucleoproteins, and other nuclear and cytoplasmic components, evaluated via indirect immunofluorescence as a primary serological screen for systemic lupus erythematosus, systemic sclerosis, mixed connective tissue disease, and Sjogren syndrome.

#### Alternate Names:

ANA; Antinuclear Antibodies; Antinuclear Autoantibodies; FANA

### Antinuclear Antibody Panel

A comprehensive laboratory panel of specific autoantibody assays (such as anti-dsDNA, anti-Smith, anti-Ro/SSA, anti-La/SSB, and anti-RNP) utilized to characterize specific autoimmune rheumatic diseases following a positive antinuclear antibody screen.

### Antioxidant

A substance that helps neutralize or limit damage from reactive molecules, including free radicals and some reactive oxygen or nitrogen species. The body makes antioxidant enzymes and obtains other antioxidants from food; high-dose supplements are not automatically beneficial and may be harmful or interact with treatment.

### Antiparietal Cell Antibody Test

A blood test that detects antibodies against cells in the stomach lining that make acid and help absorb vitamin B12. A positive result may support autoimmune gastritis or pernicious anemia but does not

prove either condition alone.

### Antiparkinson

Relating to medicines used to reduce symptoms of Parkinson disease. These medicines may increase or imitate dopamine activity or change other brain signals that influence movement.

### Antiphospholipid Antibodies

Autoantibodies that attach to proteins linked with cell membranes. Persistent antibodies may increase the risk of blood clots and pregnancy complications, but they can also appear temporarily after an infection or with some medicines.

### Antiphospholipid Antibody

An antibody directed against proteins that bind to cell-membrane fats. Persistent antibodies may be associated with blood clots or pregnancy complications, especially when clinical features of antiphospholipid syndrome are present.

#### Alternate Names:

APA; aPL Antibody

### Antiphospholipid Antibody Syndrome

Alternate name; see *Antiphospholipid Syndrome*.

### Antiphospholipid Syndrome

An autoimmune condition in which persistent antiphospholipid antibodies occur together with blood clots or certain pregnancy complications. Assessment uses the person's medical history and blood tests; antibody positivity alone does not establish the syndrome, and positive tests generally need confirmation at least 12 weeks later.

#### Alternate Names:

APS; Antiphospholipid Antibody Syndrome; Hughes Syndrome; aPL Syndrome

### Antiphospholipid Syndrome and Pregnancy

An autoimmune clotting disorder associated with recurrent pregnancy loss, placental insufficiency, fetal growth restriction, preeclampsia, thrombosis, or pregnancy morbidity. Diagnosis requires clinical

---

events plus persistent laboratory antiphospholipid antibodies; pregnancy care is specialist-led.

### **Antiplatelet**

A medicine that reduces platelet activation or aggregation and lowers the formation of arterial blood clots; see *Antiplatelet Agents*.

### **Antiplatelet Agents**

Medicines that reduce the ability of platelets to stick together and form clots. They are used in selected people to lower the risk of clots in arteries, including after some heart or blood-vessel events.

#### **Alternate Names:**

Antiplatelet Drugs; Antiplatelet Medicines

### **Antiplatelet Therapy**

Treatment with medicines that make platelets less likely to stick together. It is used in selected people to reduce clots in arteries, including after some heart attacks, strokes, or stent procedures.

### **Antiprotozoal**

A medicine or substance that kills protozoa or stops them growing. Protozoa are microscopic parasites that can cause infections such as malaria, giardiasis, or amoebiasis.

### **Antipsychotic Medicine**

See *Antipsychotic*.

### **Antipyretic**

A medicine that lowers fever and may ease related discomfort. Acetaminophen and ibuprofen are common examples. Aspirin is generally avoided in children and teenagers with viral illnesses because of the risk of Reye syndrome.

#### **Alternate Names:**

Antipyretics; Fever-Reducing Medicines

### **Antiretroviral**

A medicine that acts against retroviruses, including HIV. Different antiretroviral medicines block different steps by which HIV enters cells, copies its genetic material, or makes new virus particles.

### **Antiretroviral Therapy**

Treatment of HIV infection with a combination of medicines that block several steps in HIV reproduction. It lowers the amount of virus, protects immune function, reduces illness, and greatly lowers the chance of sexual transmission when the virus remains undetectable.

### **Antisepsis**

Use of a chemical product on living skin or tissue to reduce harmful microorganisms and lower infection risk. It does not make tissue completely sterile and must be used in a way that avoids injury.

### **Antiseptic**

A substance used on skin or other living tissue to kill or slow harmful microorganisms. Antiseptics can help reduce infection risk but may damage tissue if used incorrectly.

### **Antiserum**

Blood serum containing antibodies against a specific germ, toxin, or other substance. It can provide temporary immune protection or help neutralize a toxin.

### **Antisocial Personality**

A long-term pattern of disregarding other people's rights, safety, rules, and responsibilities, often including deceit, impulsive behaviour, aggression, or lack of remorse.

#### **Alternate Names:**

Antisocial Personality Disorder

### **Antisocial Personality Disorder**

A mental-health disorder involving a persistent pattern of violating other people's rights and social rules, deceitfulness, impulsivity, aggression, irresponsibility, or lack of remorse. The pattern begins by adolescence and continues into adult life.

#### **Alternate Names:**

ASPD; Dissocial Personality Disorder; Sociopathy

### **Antispasmodic Drugs**

Medicines that relax involuntary muscle and reduce

---

cramping or spasms, most often in the digestive tract. Some are also used for bladder or bile-duct spasms; side effects depend on the medicine.

**Alternate Names:**

Antispasmodics; Spasmolytics

### Antistreptolysin O Titer

A blood test that measures antibodies against a toxin made by group A streptococcal bacteria. A raised or rising result may show a recent strep infection when investigating rheumatic fever or kidney inflammation, but does not prove an active infection.

### Antisynthetase Syndrome

An autoimmune disease in which the immune system attacks proteins needed to make other proteins. It may cause muscle inflammation and weakness, arthritis, fever, colour changes in the fingers, rough cracked skin on the hands, or lung inflammation and scarring.

### Antithrombin

A natural blood protein that slows clotting by blocking thrombin and other clotting proteins, especially factor Xa. Too little antithrombin, whether inherited or acquired, increases the risk of clots in veins.

### Antithrombin Deficiency

A condition in which the blood has too little or too little active antithrombin, a natural anti-clotting protein. It may be inherited or acquired and increases the risk of clots, especially in the veins.

### Antithrombin III Blood Test

A blood test that measures the amount or activity of antithrombin, a protein that limits clotting. A low result may indicate inherited deficiency or may occur with liver disease, kidney protein loss, severe illness, or some treatments.

### Antithrombin III Deficiency

Reduced amount or activity of antithrombin, causing increased risk of blood clots in veins. It may be inherited or result from liver disease, kidney disease, severe illness, or other causes.

### Antithymocyte Globulin

A preparation of antibodies that removes or suppresses T cells, immune cells involved in transplant rejection and some autoimmune reactions. It is used in selected transplants and blood-cell disorders and can cause allergic reactions, low blood counts, and infections.

**Alternate Names:**

Rabbit Antithymocyte Globulin rATG Thymoglobulin

### Antithyroglobulin Antibody Test

A blood test that detects antibodies against thyroglobulin, a protein used to make thyroid hormones. Raised levels may support Hashimoto thyroiditis or other autoimmune thyroid disease but do not determine thyroid function by themselves.

### Antitoxin

An antibody or antibody preparation that binds to and neutralizes a toxin. It can provide temporary protection or treatment for selected poisonings and infections that produce toxins.

### Antitubercular Agents

Medicines used to treat tuberculosis, an infection caused by *Mycobacterium tuberculosis*. Several medicines are used together for months to prevent resistance, with monitoring for liver injury, nerve problems, vision changes, and other side effects.

### Antitussive

A medicine that reduces coughing. Some act on cough-control areas in the brain, while others reduce irritation in the airways. They may not be suitable for every cough, and opioid cough medicines can cause sleepiness and slow breathing.

**Alternate Names:**

Cough Suppressant

### Anton Syndrome

A rare neurological condition characterized by cortical blindness accompanied by visual anosognosia (complete denial of blindness) and visual confabulation. Patients genuinely behave and report as if they can see, despite profound vision loss, classically

---

caused by bilateral occipital lobe infarctions.

**Alternate Names:**

Anton-Babinski Syndrome; Visual Anosognosia

**Antral Follicle Count**

The number of small developing egg-containing sacs seen in both ovaries, usually measured by vaginal ultrasound early in the menstrual cycle. It helps estimate ovarian reserve and response to fertility medicines but does not by itself predict natural fertility.

**Alternate Names:**

AFC; Basal Antral Follicle Count

**Antral Web**

A thin, circumferential mucosal diaphragm in the gastric antrum, usually located one to two centimetres proximal to the pylorus. It can be congenital or acquired and may cause gastric outlet obstruction, nausea, postprandial vomiting, and early satiety.

**Alternate Names:**

Gastric Antral Diaphragm; Antral Septum

**Antrochoanal Polyp**

A usually noncancerous growth that starts in the lining of a cheek sinus and extends into the back of the nose. It is more common in children and young adults and may cause blockage or drainage on one side.

**Antrum**

A cavity or chamber within an organ. The term commonly refers to the lower part of the stomach near its outlet or the air-filled sinus in the cheek.

**Anuria**

Almost complete absence of urine production, often defined in an adult as less than about 100 millilitres in 24 hours. It can signal severe kidney failure, blockage, shock, or another emergency.

**Anus**

The opening at the end of the digestive tract through which stool leaves the body. Two ring-shaped muscles help control when stool and gas are released.

**Anxiety**

A feeling of worry, fear, or unease that can be a normal response to danger or stress. It becomes a disorder when it is excessive, persistent, difficult to control, or interferes with daily life.

**Anxiety & Panic**

Anxiety disorders involve excessive fear, worry, or physiological arousal; panic attacks are sudden episodes of intense fear with symptoms such as palpitations, breathlessness, chest discomfort, trembling, or fear of dying.

**Anxiety Disorder**

Alternate index label; see *Anxiety Disorders*.

**Anxiety Disorders**

A group of mental-health conditions involving excessive fear, worry, or anxiety that persists, is difficult to control, or causes significant distress or impairment. Major types include generalized anxiety disorder, panic disorder, social anxiety disorder, specific phobia, agoraphobia, separation anxiety disorder, and selective mutism. Physical symptoms may include a racing heart, sweating, muscle tension, poor sleep, or stomach upset.

**Aorta**

The body's largest artery, carrying oxygen-rich blood from the left side of the heart to the rest of the body. It passes through the chest and abdomen and divides into the arteries supplying the pelvis and legs.

**Aortal**

Relating to the aorta, the large artery that carries blood from the heart to the body.

**Aortic**

Relating to the aorta, the main artery carrying blood from the heart to the body.

**Aortic Aneurysm**

An abnormal bulge or widening in the aorta, the large artery that carries blood from the heart to



---

the body, caused by weakness in its wall. It may occur in the chest or abdomen and often causes no symptoms. If it enlarges, it may rupture or the aortic wall may tear (dissect); either can be life-threatening.

### **Aortic Aneurysm Rupture**

A break through the wall of an aortic aneurysm that causes severe internal bleeding. It may produce sudden chest, back, or abdominal pain, collapse, and shock and is a life-threatening emergency.

### **Aortic Aneurysm Screening**

Testing people at increased risk for an aortic aneurysm before symptoms appear. Abdominal ultrasound is commonly used; eligibility depends on age, sex, smoking history, and family history.

### **Aortic Angiography**

Imaging of the aorta after contrast material is given, usually with X-ray, computed tomography, or magnetic resonance imaging. It can show narrowing, blockage, aneurysm, tearing, or abnormal blood flow.

### **Aortic Arch**

The curved part of the aorta between its upward and downward sections. Three major arteries branch from it to supply the head, neck, and arms; disease here can affect blood flow or press on nearby structures.

### **Aortic Arch Syndrome**

Reduced blood flow caused by narrowing or blockage of the aortic arch or its major branches. It may cause unequal pulses or blood pressure, arm fatigue, dizziness, or neurological symptoms.

### **Aortic Balloon Pump**

A temporary heart-support device. A catheter with a balloon is placed in the main artery in the chest; the balloon inflates and deflates with each heartbeat to improve blood flow to the heart and reduce the heart's workload.

#### **Alternate Names:**

Intra-Aortic Balloon Pump; IABP

### **Aortic Body**

A small group of sensory cells near the aortic arch that detects changes in blood oxygen, carbon dioxide, and acidity. It sends signals to the brain to help adjust breathing and circulation.

### **Aortic Coarctation**

A narrowing of the aorta present from birth, usually just beyond the branches that supply the head and arms. It can cause higher blood pressure in the arms than in the legs, weak leg pulses, poor growth, or heart strain. Some people have no symptoms until later in life.

### **Aortic Disease**

Any disorder affecting the aorta, the body's main artery, including aneurysm, dissection, narrowing, inflammation, or fatty deposits. Some cases cause no symptoms; sudden severe chest, back, or abdominal pain is an emergency.

### **Aortic Dissection**

A tear in the inner lining of the aorta that allows blood to split the layers of its wall. It can suddenly cause severe chest or back pain, stroke, organ damage, or death and needs emergency assessment.

#### **Alternate Names:**

Dissecting Aortic Aneurysm; Dissecting Aneurysm; Aortic Tear

### **Aortic Embolism**

Blockage of an artery by clot, fatty material, or other debris that breaks away from the aorta and travels downstream. It can suddenly reduce blood flow to a limb or organ.

### **Aortic Graft Infection**

Infection involving an artificial graft used to repair or replace part of the aorta. It can cause fever, bloodstream infection, bleeding, or a new abnormal connection with the intestine and requires urgent specialist care.

---

## Aortic Injury

Damage to the aorta, often caused by severe chest trauma or a medical procedure. A tear can cause internal bleeding, reduced blood flow, shock, and a life-threatening emergency.

## Aortic Insufficiency

Leakage through the aortic valve that allows blood to flow backward into the heart's main pumping chamber. It can make the heart work harder and may eventually cause heart failure.

## Aortic Regurgitation

Leakage through an aortic valve that does not close completely, allowing blood to flow from the aorta back into the left ventricle during diastole. Acute regurgitation can cause sudden pulmonary edema or shock; chronic regurgitation may enlarge and weaken the left ventricle and cause breathlessness, chest discomfort, palpitations, or heart failure. Echocardiography assesses the cause and severity.

## Aortic Regurgitation Murmur

A blowing sound heard during the part of the heart-beat when the heart relaxes, caused by blood leaking backward through the aortic valve. It is often loudest along the left side of the breastbone.

## Aortic Rupture

A life-threatening break through the wall of the aorta, often following an aneurysm or dissection. It can cause sudden severe chest, back, or abdominal pain, internal bleeding, collapse, and shock.

## Aortic Sclerosis

Calcification, fibrosis, and thickening of the aortic valve leaflets without significant hemodynamic obstruction or restriction of valve opening. It is a common echocardiographic finding in older adults and is associated with increased cardiovascular risk.

## Aortic Stenosis

Narrowing of the aortic valve that obstructs blood leaving the left ventricle. Common causes include age-related calcification, a bicuspid valve, and rheumatic disease; severe disease can cause

exertional breathlessness, chest pain, fainting, left-ventricular thickening, or heart failure. Echocardiography assesses severity.

## Aortic Stenosis Grading

Assessment of how severely the aortic valve is narrowed using ultrasound measurements of valve area and blood-flow speed or pressure difference. Results are generally described as mild, moderate, or severe and help guide monitoring and treatment.

## Aortic Valve

A valve between the heart's left pumping chamber and the aorta. It opens to let blood leave the heart and closes to stop blood flowing backward.

## Aortic Valve Disease

Narrowing, leakage, or another abnormality of the valve between the heart's left pumping chamber and the aorta. It may cause breathlessness, chest pain, tiredness, fainting, or no symptoms at first.

## Aortic Valve Insufficiency

Leakage of the aortic valve that allows blood to flow backward into the left pumping chamber between heartbeats. The extra workload can enlarge or weaken the heart over time.

## Aortic Valve Regurgitation

Backward leakage of blood through an aortic valve that does not close completely. The heart must pump harder, and severe or long-standing leakage can cause enlargement or heart failure.

### Alternate Names:

Aortic Regurgitation; Aortic Insufficiency; Aortic Incompetence

## Aortic Valve Replacement

A cardiac surgical or catheter-directed interventional procedure in which a severely diseased or calcified aortic valve is excised or displaced and replaced with a mechanical prosthesis or a bioprosthetic tissue valve.

### Alternate Names:

AVR; TAVR; TAVI

---

## Aortic Valve Stenosis

Alternate index label; see *Aortic Stenosis*.

## Aortitis

Inflammation of the aorta, the body's largest artery. It may result from an autoimmune disease or infection and can weaken the artery or cause narrowing, aneurysm, dissection, or leakage of the aortic valve.

## Aortocaval Compression Syndrome

Hemodynamic compromise occurring in late pregnancy when the gravid uterus compresses the inferior vena cava and abdominal aorta while the mother is in the supine position. Leads to decreased venous return, reduced cardiac output, hypotension, dizziness, and fetal distress, rapidly relieved by placing the patient in the left lateral decubitus position.

### Alternate Names:

Supine Hypotensive Syndrome of Pregnancy

## Aortoenteric Fistula

A life-threatening direct communication between the aorta and the gastrointestinal tract, most frequently involving the third or fourth portion of the duodenum. It may occur primarily from an eroding abdominal aortic aneurysm or secondarily following prosthetic aortic reconstruction, classically presenting with minor herald gastrointestinal bleeding followed by catastrophic exsanguination.

## Aortography

Imaging of the aorta after contrast material is injected, often through a catheter. It can show aneurysms, tears, narrowing, blockages, or abnormal blood-vessel connections; computed tomography or magnetic resonance imaging is often used instead.

## Aortoiliac Occlusive Disease

Severe atherosclerotic stenosis or complete occlusion of the distal abdominal aorta and common iliac arteries. It is clinically characterized by the triad of buttock and thigh claudication, diminished or absent femoral pulses, and erectile dysfunction in men.

### Alternate Names:

Leriche Syndrome

## Aortopulmonary Window

A rare birth defect in which an opening connects the aorta and pulmonary artery. Too much blood may flow to the lungs, causing breathing problems, poor growth, or heart failure in infancy.

## AP X-Ray

An X-ray view in which the beam enters through the front of the body and exits through the back. It is called an anteroposterior view.

## APACHE II Score

A hospital scoring system that combines vital signs, laboratory results, age, and serious long-term illnesses to estimate the severity of critical illness. A higher score generally indicates greater risk, but it does not predict an individual outcome with certainty.

## Apathy

Reduced motivation, interest, or emotional response. It can occur with depression, dementia, Parkinson disease, brain injury, other neurological illness, or medicines and is different from ordinary tiredness or sadness.

## APC Tumor Suppressor

A protein made from the APC gene that helps control cell growth and the removal of damaged cells. Harmful changes in this gene can allow excess intestinal polyps to form and increase colorectal-cancer risk.

## Apert Syndrome

A rare inherited condition in which some skull bones join too early. It causes an unusually shaped head and face and often joined fingers or toes; breathing, vision, hearing, or development may also be affected.

## Apex

The tip or pointed end of a body structure. Examples include the lower pointed end of the heart, formed mainly by the left ventricle, and the upper

---

tip of a lung.

### **Apex Beat**

The outermost and lowest point on the precordium at which the cardiac impulse can be palpated or inspected during ventricular systole, typically located in the left fifth intercostal space along the midclavicular line.

#### **Alternate Names:**

Apical Impulse

### **Apgar Score**

A quick assessment of a newborn's colour, heart rate, reflex response, muscle tone, and breathing, usually recorded one and five minutes after birth. It describes immediate adaptation and helps guide care but does not predict long-term health by itself.

#### **Alternate Names:**

Apgar Assessment; Newborn Apgar Score

### **Aphagia**

Complete inability to swallow. It may result from a blockage, neurological disease, severe pain, or loss of normal swallowing-muscle function and can lead to choking, dehydration, or aspiration.

### **Aphakia**

Absence of the eye's natural lens, usually after cataract surgery, injury, or a birth difference. It causes major focusing problems and is usually corrected with an intraocular lens, contact lens, or strong glasses.

### **Aphantasia**

Markedly reduced or absent ability to voluntarily form mental visual images. It is a cognitive phenomenon rather than, by itself, a disease and may occur with otherwise normal vision, memory, and imagination; distress or functional impairment warrants clinical assessment.

### **Aphasia**

A brain-related disorder that impairs speaking, understanding, reading, or writing language. It most often follows a stroke but can also result from head

injury, brain tumour, infection, or degeneration.

### **Apheresis**

A procedure in which blood passes through a machine that removes a selected component, such as plasma, platelets, red cells, or abnormal cells, and returns the remaining blood to the person.

### **Apheresis Donation**

A blood donation in which a machine collects one component, such as platelets or plasma, and returns the other blood components to the donor. The process may allow more of the needed component to be collected than whole-blood donation.

### **Aphonia**

Complete loss of the ability to produce a normal voice, although whispering may remain possible. Causes include inflammation, vocal-cord paralysis, injury, growths, nerve disease, or psychological factors.

### **Aphrasia**

Loss or impairment of the ability to speak. The term is older and less precise than aphasia, which can affect speaking, understanding, reading, or writing.

### **Aphthous Stomatitis**

Repeated painful ulcers on the moist lining of the mouth, commonly called canker sores. They are usually shallow, heal without scarring, and are not contagious.

### **Aphthous Ulcer**

A small, painful ulcer on the moist lining of the mouth, commonly called a canker sore. It usually heals within one to two weeks and is not contagious.

### **Apical**

Relating to or located at the tip of a body structure, such as the tip of a lung, heart, or tooth.

### **Apical Ballooning Syndrome**

A temporary weakening and ballooning of the main pumping chamber of the heart, often after severe

---

emotional or physical stress. It can resemble a heart attack but usually occurs without a blocked coronary artery.

### **Apical Periodontitis**

Inflammation or infection around the tip of a tooth root, usually caused by infected or dead pulp. It may cause pain when biting, swelling, or no symptoms and is assessed with dental examination and imaging.

### **Apical Pulse**

The heartbeat counted by listening over the tip of the heart, usually on the left side of the chest. It may be used when the pulse at the wrist is difficult to feel or when an accurate heart rate is needed.

### **Apicoectomy**

Dental surgery that removes the tip of a tooth root and nearby infected tissue, usually when infection persists after root-canal treatment. The end of the root is then sealed.

### **Aplasia**

Failure of an organ, tissue, or group of cells to develop, causing it to be absent or severely underdeveloped. It may be present from birth or result from inherited disease, injury, infection, or severe damage to the cells that produce it.

### **Aplasia Cutis Congenita**

A birth defect in which a localized area of skin, most often on the scalp, is absent or incompletely formed. It may heal as a scar or hairless patch, sometimes with an underlying skull or vascular abnormality; large, ulcerated, or atypical lesions need specialist evaluation.

### **Aplastic**

Describing a tissue that has failed to develop or produce its normal cells. The term is often used for bone-marrow failure, in which too few red cells, white cells, and platelets are made.

### **Aplastic Anemia**

A rare, serious bone-marrow failure disorder in which the marrow cannot make enough blood cells, usually because blood-forming stem cells are damaged. It lowers red cells, white cells, and platelets—not just red cells—so it can cause tiredness, infections, and easy bruising or bleeding. Causes include immune attack, some medicines, toxic chemicals, radiation, infections, and inherited conditions; sometimes no cause is found.

### **Aplastic Crisis**

A sudden, temporary stop in red-cell production that causes a rapid worsening of anemia. It most often follows parvovirus B19 infection in people whose red cells already have a shortened lifespan, such as those with sickle-cell disease.

### **Apley Compression Test**

A physical examination manoeuvre used to evaluate for meniscal tears of the knee joint. With the patient prone and the knee flexed to ninety degrees, the examiner applies downward axial force to the heel while rotating the tibia internally and externally; pain or joint line clicking indicates meniscal pathology.

#### **Alternate Names:**

Apley Grinding Test; Apley's Compression Test

### **Apley Distraction Test**

A clinical examination technique used to differentiate collateral ligament injuries from meniscal tears. With the patient prone and the knee flexed to ninety degrees, the examiner stabilizes the posterior thigh and applies upward traction to the lower leg with internal and external rotation; localized pain suggests ligamentous sprain rather than meniscal damage.

#### **Alternate Names:**

Apley's Distraction Test

### **Apley Grind Test**

A physical examination maneuver used in orthopedics to evaluate meniscus tears of the knee. With the patient prone and the knee flexed to 90 degrees, the examiner stabilizes the thigh, applies downward axial compressive force onto the heel while externally

---

and internally rotating the tibia; pain or mechanical clicking indicates a torn meniscus.

**Alternate Names:**

Apley Compression Test; Apley Meniscal Test

**Apley Scratch Test**

A rapid physical examination maneuver used in musculoskeletal medicine to assess the active range of motion of the shoulder joint complex. The patient is asked to reach behind the head and touch the superior medial angle of the opposite scapula (evaluating glenohumeral abduction and external rotation) and reach behind the back to touch the inferior angle of the opposite scapula (evaluating adduction and internal rotation). Asymmetry or restricted reach indicates rotator cuff pathology, adhesive capsulitis, or glenohumeral arthritis.

**Alternate Names:**

Apley Scratch Maneuver; Apley Shoulder Test

**APLS**

Abbreviation for antiphospholipid syndrome, an autoimmune disorder associated with venous or arterial thrombosis and pregnancy complications, with persistent antiphospholipid antibodies.

**Apnea**

A temporary pause in breathing. Prolonged or repeated episodes can lower oxygen, raise carbon dioxide, slow the heartbeat, cause loss of consciousness, or lead to death.

**Apnea of Infancy**

Repeated pauses in breathing in an infant. Causes and significance vary with age and clinical context; apnea of prematurity is a distinct diagnosis in preterm infants.

**Apnea Of Prematurity**

Repeated pauses in breathing in a premature infant because breathing control is not fully developed. Episodes may lower oxygen or slow the heartbeat and require monitoring and, when necessary, breathing support.

**Apnea-Hypopnea Index**

The average number of complete or partial breathing interruptions during each hour of sleep, measured in a sleep study. It helps describe sleep-apnea severity, but symptoms, oxygen levels, and other findings also matter.

**Apneic**

Relating to a period when breathing has stopped temporarily. Apneic episodes may occur during sleep, illness, poisoning, seizures, or severe airway blockage and can quickly lower oxygen.

**Apneustic Breathing**

An abnormal pattern with an unusually long pause after breathing in. It usually indicates serious injury to the brainstem and may occur with other signs of severe neurological damage.

**Apnoea**

The temporary stopping of breathing; the spelling is commonly used in British English. It may result from airway blockage, reduced breathing signals from the brain, illness, medicines, or sleep-related breathing disorders.

**Apocrine Hidrocystoma**

A benign cyst of an apocrine sweat gland, commonly presenting as a translucent or bluish, dome-shaped papule near the eyelid or elsewhere on the skin. It may enlarge with heat or sweating and is usually diagnosed clinically.

**Apoenzyme**

The protein part of an enzyme before its required helper substance has attached. It is usually inactive or less active until the helper, called a cofactor, binds to it.

**Apolipoprotein B100**

A protein on cholesterol-carrying particles such as low-density and very-low-density lipoproteins. Its blood level estimates the number of particles that can contribute to fatty buildup in artery walls.



---

### **Apolipoprotein CII**

A protein on fat-carrying particles in the blood that activates an enzyme needed to break down triglycerides. Too little or faulty apolipoprotein CII can cause triglycerides to rise to very high levels.

### **Apolipoproteins**

Proteins attached to fats in the blood that help transport and process cholesterol and triglycerides. Different apolipoproteins are found on different fat-carrying particles and have different effects.

### **Aponeurosis**

A broad, flat sheet of tough connective tissue that attaches a muscle to bone, another muscle, or other tissues and helps transmit the muscle's force.

### **Apophenia**

Seeing meaningful patterns or connections in unrelated or random events. Mild experiences can occur normally, while marked or distressing experiences may occur with some psychiatric or neurological conditions.

### **Apophyses**

Bony projections that grow from or arise near a bone and usually provide attachment for muscles or ligaments. Unlike the main growth plate of a long bone, an apophysis does not normally increase the bone's length.

### **Apoptosis**

A controlled process of cell death in which a cell dismantles itself and is removed with little inflammation. It helps shape developing organs and remove old, damaged, or abnormal cells; failure of this process can contribute to cancer.

### **Apoptosis Extrinsic Pathway**

A controlled cell-death process that begins when an outside signal attaches to a death receptor on a cell's surface. Internal enzymes then take the cell apart in an orderly way, usually without causing the swelling and inflammation of an injury.

### **Apoptosis Intrinsic Pathway**

A controlled cell-death process that begins with severe stress inside a cell, such as DNA damage, lack of growth signals, or lack of oxygen. The cell's energy-producing structures then release signals that activate enzymes to dismantle it.

### **Apoptotic Body**

A small membrane-bound piece of a cell that has begun controlled cell death. Immune or neighbouring cells can recognize and remove these pieces without causing much inflammation.

### **Appalachian Type Amyloidosis**

A rare hereditary transthyretin amyloidosis variant associated with abnormal amyloid deposition, often causing neuropathy, autonomic symptoms, or heart involvement. Diagnosis requires genetic and tissue or specialized laboratory evaluation.

### **Apparent Mineralocorticoid Excess**

A rare inherited disorder in which cortisol acts too strongly on the kidneys because it is not changed into an inactive form normally. It can cause severe high blood pressure, low potassium, and unusually low levels of renin and aldosterone.

### **Appendageal**

Relating to structures that develop from the skin, including hair follicles, sweat glands, and oil-producing glands.

### **Appendectomy**

Surgery to remove the appendix, usually to treat appendicitis. It may be performed laparoscopically through small incisions using a camera or through a larger open incision; the approach depends on the condition and clinical needs.

#### **Alternate Names:**

Appendicectomy; Laparoscopic Appendectomy; Appendix Removal; Open Appendectomy

### **Appendiceal**

Relating to the appendix, a narrow, finger-shaped pouch attached to the beginning of the large intes-

---

tine.

### **Appendiceal Adenocarcinoma**

A cancer arising from gland-forming cells in the appendix. It may be found after appendicitis or unexpectedly and can spread through the abdominal lining; outlook depends on its type, size, and spread.

### **Appendiceal Cancer**

Cancer arising in the appendix. Types include neuroendocrine tumours, mucin-producing tumours, and adenocarcinoma; some can rupture and spread tumour cells or mucus through the abdomen.

### **Appendiceal Rupture**

A tear or hole in an inflamed appendix that allows infected material into the abdomen. It can cause peritonitis, an abscess, sepsis, or severe abdominal pain and requires urgent care.

### **Appendiceal Tumors**

Tumours arising in the appendix, including neuroendocrine tumours, mucin-producing tumours, and adenocarcinoma. Some cause no symptoms, while others cause appendicitis, pain, blockage, or abdominal swelling.

### **Appendicitis**

Inflammation of the appendix, often after its opening becomes blocked by hardened stool, mucus, or swollen tissue. It commonly causes pain that moves to the lower right abdomen, nausea, loss of appetite, and fever and can lead to rupture.

#### **Alternate Names:**

Epityphlitis; Acute Appendicitis

### **Appendicular Skeleton**

The bones of the arms and legs and the shoulder and pelvic girdles that attach them to the central skeleton.

### **Appendix**

A small, finger-shaped pouch attached to the beginning of the large intestine. It contains immune tissue and may help support healthy gut bacteria,

although its exact function is not fully understood and it is not essential for digestion. Inflammation of the appendix is called appendicitis.

### **Appendix Cancer**

A malignant neoplasm arising within the appendix, including neuroendocrine tumors, goblet cell carcinoids, mucinous cystadenocarcinomas, and colonic-type adenocarcinomas.

### **Appendix Testis**

A small remnant of tissue near the upper part of the testis, left over from early development. It is usually harmless but can sometimes twist and cause pain in boys.

### **Appetite**

The psychological desire to eat food, distinct from physiological hunger, modulated by central nervous system pathways, neuroendocrine signaling, sensory stimuli, emotional states, and metabolic demands.

### **Appetite Regulation**

The control of hunger and fullness by the brain, digestive tract, hormones, emotions, habits, sleep, and surroundings. Signals such as ghrelin promote hunger, while leptin and other signals help promote fullness.

### **Applied Behavior Analysis**

A structured approach that observes behaviour and uses learning principles, practice, and reinforcement to build useful skills or reduce behaviour that interferes with daily life. Goals should be individualized and respectful of the person.

### **Apposition**

Placement of two structures next to or against each other. In wound care, good apposition means that the edges of a cut are accurately aligned so they can heal.

### **Apprehension Test**

A provocative orthopedic examination maneuver designed to detect joint instability, most commonly

---

applied to the shoulder (glenohumeral joint) or knee (patellofemoral joint). In the anterior shoulder apprehension test, the examiner places the arm in 90 degrees of abduction and maximal external rotation; a positive test occurs when the patient exhibits visible fear, alarm, or resistance due to a feeling of impending anterior subluxation or dislocation. In the patellar apprehension test (Fairbank's sign), lateral displacement of the patella elicits anxiety and quadriceps contraction.

**Alternate Names:**

Anterior Apprehension Test; Crank Test; Fairbank's Apprehension Test

**Appropriate For Gestational Age**

Describing a fetus or newborn whose estimated size or weight is within the expected range for the number of weeks of pregnancy, commonly between the 10th and 90th percentiles.

**Apraxia**

Difficulty performing a learned, purposeful movement even though strength, sensation, coordination, and understanding are sufficient. It results from a problem in the brain's planning of movement rather than simple weakness.

**Apraxia of Speech**

A neurogenic motor speech disorder characterized by an impaired ability to plan, sequence, and program the sensorimotor commands necessary for phonetically normal speech production in the absence of significant neuromuscular weakness or paralysis. Marked by inconsistent articulatory errors, effortful groping, and disrupted prosody.

**Alternate Names:**

Verbal Apraxia; Speech Apraxia

**Apraxic Gait**

A higher-level neurological gait disorder characterized by the loss of the ability to coordinate voluntary stepping and leg movements in the absence of primary sensory loss, weakness, or cerebellar ataxia. Patients exhibit hesitation in initiating walking, a wide-based stance, short shuffling steps, and a tendency for the feet to appear glued to the floor

(magnetic gait), commonly seen in normal pressure hydrocephalus, bilateral frontal lobe lesions, and advanced subcortical vascular encephalopathy.

**Alternate Names:**

Gait Apraxia; Frontal Gait Disorder; Magnetic Gait; Bruns Apraxia

**Aprosodia**

A neurological speech and language disorder characterized by an inability to express or comprehend the emotional tone, rhythm, pitch, and melody (prosody) of verbal communication. It is classically associated with lesions of the non-dominant (right) cerebral hemisphere in areas homologous to Broca and Wernicke expressive and receptive language regions.

**Alternate Names:**

Dysprosody; Affective Agnosia of Speech

**APS**

Abbreviation for antiphospholipid syndrome; it is also used for other conditions in different contexts, so the intended meaning should be confirmed.

**Aquagenic Keratoderma**

Temporary painful whitening, thickening, and wrinkling of the palms or soles after contact with water. It may cause prickling or burning, is often associated with cystic fibrosis, and usually settles after the skin dries.

**Aquaporin**

A channel in a cell membrane that lets water pass through. Aquaporin-2 in kidney tubules responds to vasopressin and helps the kidneys return water to the blood instead of losing it in urine.

**Aquatic Therapy**

Exercise or rehabilitation performed in water. Water buoyancy reduces load on joints while water resistance strengthens muscles and supports balance, movement, and conditioning.

**Aqueous Flare**

A hazy beam seen in the clear fluid at the front

---

of the eye during slit-lamp examination. It means inflammation has allowed proteins to leak into the eye and may occur with uveitis, infection, injury, or surgery.

### **Aqueous Humor**

The clear fluid in the front part of the eye. It nourishes the cornea and lens, removes waste, and helps maintain pressure and the shape of the eye.

### **Arachidonic Acid Pathway**

A major cellular lipid metabolic cascade in which arachidonic acid, released from cell membrane phospholipids by phospholipase A2, is converted into bioactive eicosanoids (prostaglandins, prostacyclin, thromboxane A2, and leukotrienes) via cyclooxygenase and lipoxygenase enzymes.

### **Arachnitis**

An older spelling of arachnoiditis, chronic inflammation and scarring of the arachnoid membrane around spinal or cranial nerves that can cause persistent pain, sensory symptoms, weakness, or bladder and bowel dysfunction.

### **Arachnodactyly**

Abnormally long, thin fingers or toes. It may occur alone or as a feature of inherited connective-tissue disorders such as Marfan syndrome.

### **Arachnoid Cyst**

A usually noncancerous, fluid-filled sac between layers of the membrane covering the brain or spinal cord. Many cause no symptoms, but a large cyst may cause headache, seizures, weakness, or pressure on nearby tissue.

### **Arachnoid Cysts**

Fluid-filled sacs between layers of the membrane covering the brain or spinal cord. Many are discovered by chance and cause no symptoms; larger cysts can press on nearby tissue and cause neurological problems.

### **Arachnoid Granulations**

Macroscopic tufted outgrowths of the arachnoid membrane that project through the dura mater into the dural venous sinuses, predominantly the superior sagittal sinus, serving as the primary structural pathway for the passive reabsorption of cerebrospinal fluid into venous blood.

#### **Alternate Names:**

Pacchionian Bodies; Arachnoid Villi

### **Arachnoid Mater**

The middle of three thin membranes covering the brain and spinal cord. The space beneath it contains the fluid that cushions the nervous system and carries important blood vessels.

### **Arachnoidal Fibroblastoma**

A rare benign or low-grade tumor arising from fibroblastic cells of the arachnoid coverings. Symptoms depend on location and may result from compression of the brain, spinal cord, or nerves.

### **Arachnoiditis**

Inflammation of the arachnoid, one of the membranes covering the brain and spinal cord, most often in the spine. Inflammation may cause thickening, scarring, and bands that make nerve roots stick together or interfere with the normal flow of cerebrospinal fluid. Possible causes include infection, bleeding around the spinal cord, spinal injury, chronic nerve compression, surgery or another spinal procedure, and chemical irritation. Symptoms may include severe or burning back and limb pain, numbness, tingling, weakness, abnormal reflexes, or problems with bladder or bowel function. The course may remain stable or worsen over time.

### **Arbovirus**

A virus spread by a biting insect or tick, such as a mosquito, tick, or sandfly. Arboviruses can cause fever, rash, joint pain, bleeding, or inflammation of the brain, depending on the virus.

### **Arbovirus A Chikungunya Type**

An older classification term for chikungunya virus, a mosquito-borne alphavirus that causes fever and

---

often severe inflammatory joint pain, rash, headache, and muscle pain.

### **ARBs**

Medicines that block the main receptor for angiotensin II, relaxing blood vessels and reducing salt and water retention. They lower blood pressure and can protect the heart and kidneys in selected conditions, but are usually not combined with ACE inhibitors.

### ***Arcanobacterium haemolyticum* Infection**

A bacterial infection that commonly causes pharyngitis in adolescents and young adults, with fever, sore throat, tonsillar exudate, cervical lymph-node enlargement, and sometimes a scarlatiniform or morbilliform rash. It can resemble group A streptococcal infection, so throat testing and clinical assessment guide treatment.

### **Arciform**

Having a curved or arch-like shape. The term may describe the appearance of a skin lesion, rash, blood vessel, or other body structure.

### **Arcuate Uterus**

A mild variation in the shape of the inside top of the uterus, present from birth, that creates a shallow curved indentation. It is usually considered a normal variant and generally has little or no effect on fertility or pregnancy.

### **Arcus Senilis**

A greyish-white, circular lipid deposition ring appearing in the peripheral corneal stroma, commonly seen in elderly individuals as a benign finding, or in younger patients (*arcus juvenilis*) associated with familial hyperlipidemia.

#### **Alternate Names:**

Corneal Arcus; Arcus Lipoides; Gerontoxon

### **ARDS**

Abbreviation for acute respiratory distress syndrome, a life-threatening inflammatory lung injury causing fluid leakage into the air sacs, severe oxygenation failure, and often the need for ventilatory

support.

### **Area Under The Concentration-Time Curve**

A measure of the total amount of a medicine in the blood over a period of time. It helps compare how medicines are absorbed and removed and can guide dosing for selected treatments.

### **Aregenerative Anemia**

Anemia with an inappropriately low reticulocyte response, indicating inadequate red-cell production by the bone marrow. Causes include iron, vitamin B12, or folate deficiency, kidney disease, inflammation, endocrine disease, marrow failure, and some medicines.

### **Arenaviruses**

A family of viruses carried mainly by rodents. Some cause serious human diseases, including viral haemorrhagic fevers, after exposure to infected rodent urine, droppings, saliva, or contaminated dust.

### **Areola**

The darker circular area surrounding the nipple. It contains small glands that help lubricate and protect the skin, especially during breastfeeding.

### **ARG Deficiency**

Abbreviation for arginase deficiency, an inherited urea-cycle disorder in which arginine accumulates and can cause progressive spasticity, developmental problems, seizures, and elevated ammonia.

### **Argentine Hemorrhagic Fever**

A serious viral illness caused by Junín virus, an arenavirus carried by rodents in parts of Argentina. It can cause fever, weakness, bleeding, and nervous-system problems and requires urgent medical care.

### **Argininosuccinate Lyase Deficiency**

An inherited urea-cycle disorder caused by deficient argininosuccinate lyase. Argininosuccinic acid and ammonia can accumulate, causing poor feeding, vomiting, liver problems, developmental impair-

---

ment, or life-threatening hyperammonemia.

### **Argininosuccinic Aciduria**

A rare inherited disorder in which the body cannot safely remove waste nitrogen produced when protein is broken down. Ammonia and argininosuccinic acid build up and may cause vomiting, sleepiness, seizures, liver problems, or developmental impairment.

### **Argon Plasma Coagulation**

A procedure that uses electrically charged argon gas to deliver heat to the surface of tissue. It can stop bleeding or destroy selected abnormal tissue, often during an endoscopic examination.

#### **Alternate Names:**

APC; Argon Plasma Hemostasis

### **Argyll Robertson Pupil**

A small, irregular pupil that becomes smaller when focusing on a nearby object but reacts poorly or not at all to light. It is classically associated with late syphilis affecting the nervous system, but other causes are possible.

### **Arias-Stella Reaction**

A benign change in cells lining the uterus or cervix caused by pregnancy hormones or hormone medicines. Under a microscope it can resemble cancer, so the clinical setting and other findings are important.

### **Arm**

The part of the upper limb between the shoulder and elbow. In everyday language, “arm” may also mean the entire upper limb, including the forearm and hand.

### **Arm CT Scan**

A scan that uses X-rays and computer processing to create cross-sectional images of the arm. It can show fractures, bleeding, inflammation, masses, and other structural problems.

### **Arm Fracture**

A break in one or more bones of the upper limb, usually the upper arm bone or one of the forearm bones. It may cause pain, swelling, bruising, deformity, or inability to move the arm normally.

### **Arm MRI Scan**

A scan that uses a magnetic field and radio waves to create detailed images of the arm. It can show muscles, tendons, ligaments, nerves, bone marrow, blood vessels, and masses without X-ray radiation.

### **Arm Pain**

Pain in the shoulder, upper arm, elbow, forearm, or nearby tissues. Causes include muscle or tendon injury, nerve compression, arthritis, and blood-vessel disease; sudden severe pain with chest discomfort or breathlessness is an emergency.

#### **Alternate Names:**

Brachialgia; Upper-Limb Pain

### **Armpit Lump**

A lump or swelling in the armpit, commonly caused by an enlarged lymph node, skin infection, cyst, or noncancerous growth. A hard, fixed, enlarging, persistent, or unexplained lump needs medical assessment.

### **Arnold-Chiari Malformation**

A birth defect in which part of the lower brain extends through the opening at the base of the skull into the spinal canal. It may block the normal flow of brain fluid and cause headache, neck pain, balance problems, swallowing difficulty, or weakness.

### **Aromatase**

An enzyme that changes androgen hormones into oestrogen. It is found in tissues including the ovaries, fat, and breast and helps regulate hormone levels.

### **Aromatase Excess Syndrome**

A rare inherited condition in which too much aromatase causes excess oestrogen production. It may cause breast development in males and early puberty, rapid growth followed by short adult height,



---

or other hormone-related changes.

### **Arousal Index**

The average number of brief awakenings or shifts to lighter sleep during each hour of a sleep study. A high result suggests repeatedly interrupted sleep, but its importance depends on symptoms and other sleep-study findings.

### **Arrector Pili**

A small involuntary muscle attached to a hair follicle. When it contracts, it raises the hair and produces goosebumps.

### **Arrector Pili Muscle**

The small involuntary muscle attached to each hair follicle. Nerve signals can make it contract, raising the hair and producing temporary goosebumps.

#### **Alternate Names:**

Piloerector Muscle; Arrectores Pili

### **Arrested Caries**

Tooth decay that has stopped progressing, often because the affected surface has become hard, smooth, and easier to clean. It may remain stable but still requires dental assessment and prevention of new decay.

### **Arrhythmia**

An abnormal heart rate or rhythm. The heartbeat may be too fast, too slow, irregular, or start from the wrong place; some arrhythmias are harmless, while others reduce blood flow or cause cardiac arrest.

### **Arrhythmias**

Abnormal heart rates or rhythms caused by changes in the heart's electrical signals. They range from harmless extra beats to rhythms that cause fainting, poor circulation, or cardiac arrest.

### **Arrhythmogenic Right Ventricular Cardiomyopathy**

An inherited disease in which heart muscle, especially in the right pumping chamber, is replaced by scar tissue and sometimes fat. This can cause

dangerous irregular rhythms, fainting, or sudden cardiac arrest, particularly during exercise.

### **Arsenic**

A naturally occurring element that is poisonous in many forms. Exposure can cause vomiting, diarrhoea, nerve problems, heart problems, skin changes, and increased cancer risk; some compounds have limited medical uses.

### **Arsenic Poisoning**

Illness caused by sudden or long-term exposure to arsenic. Acute poisoning can cause severe vomiting, abdominal pain, and diarrhea. Long-term exposure, especially to inorganic arsenic in drinking water or food, can cause skin darkening and thickening, nerve and other organ problems, and increased risk of skin, lung, or bladder cancer.

#### **Alternate Names:**

Arsenic Toxicity; Arsenicosis

### **Arsenic Toxicity**

Harm caused by arsenic exposure. It may affect the digestive system, heart, nerves, blood, skin, and nails; severe exposure can be fatal.

### **Artemisinin**

A compound originally obtained from the sweet-wormwood plant and used, with other medicines, to treat malaria. Combination treatment is important because malaria parasites can become resistant when artemisinin is used alone.

### **Arteria Femoralis**

The femoral artery, the main artery supplying the thigh and lower limb. It continues from the external iliac artery below the groin and can be felt or accessed near the groin.

### **Arterial Blood Gas**

A blood test, usually from an artery at the wrist, that measures acidity, oxygen, carbon dioxide, and bicarbonate. It helps assess breathing, oxygen delivery, and acid-base disorders.

---

### Arterial Blood Gas Analyzer

A machine that measures acidity, oxygen, carbon dioxide, and related values in arterial blood. The results help assess how well a person is breathing and whether the blood is too acidic or alkaline.

### Arterial Catheter

A thin tube placed in an artery, often at the wrist, to measure blood pressure continuously and take repeated blood samples. Possible complications include bleeding, infection, clotting, or reduced blood flow.

### Arterial Catheterization

Placement of a thin tube in an artery, usually at the wrist, for continuous blood-pressure monitoring or repeated arterial blood tests. It is used when close monitoring is needed and can cause bleeding, infection, clotting, or reduced circulation.

### Arterial Dissection

A pathological process in which a disruption or tear in the tunica intima allows circulating blood under pressure to enter and cleave the media, creating a false lumen that tracks longitudinally along the arterial wall and may lead to vessel occlusion or rupture.

### Arterial Duplex

An ultrasound test that shows the structure of arteries and measures blood flow through them. It can detect narrowing, blockage, aneurysm, or whether a vascular graft remains open.

### Arterial Embolism

Sudden blockage of an artery by material carried in the bloodstream, usually a blood clot. It can abruptly cut off blood flow to a limb, brain, kidney, intestine, or other organ and is an emergency.

### Arterial Embolization

A procedure that deliberately blocks a selected artery, usually to stop bleeding, reduce blood supply to a tumour, or treat an abnormal blood-vessel connection.

### Arterial Injury

Damage to an artery that may cause severe bleeding, a clot, narrowing, reduced blood flow, tissue loss, or life-threatening complications. It can result from trauma, surgery, or a medical procedure.

### Arterial Insufficiency

Reduced blood flow through an artery to a tissue or limb, usually because the artery is narrowed or blocked. It may cause pain with walking, pain at rest, cool skin, slow-healing wounds, or weak pulses.

### Arterial Line

A thin tube placed in an artery, usually at the wrist, for continuous blood-pressure monitoring and repeated arterial blood sampling. It is used in critical illness or major surgery and can cause bleeding, infection, clotting, or reduced circulation.

### Arterial Resistance

The opposition blood encounters as it flows through arteries, especially the small arteries that can narrow or widen. It helps control blood pressure and how much blood reaches each tissue.

#### Alternate Names:

Vascular Resistance; Systemic Vascular Resistance

### Arterial Spin Labeling

An MRI method that uses magnetically labeled blood as a natural tracer to measure blood flow, especially in the brain, without injecting a tracer.

### Arterial Stenosis

Narrowing of an artery that reduces blood flow to the tissues beyond it. Common causes include fatty plaque, inflammation, injury, or an inherited abnormality of the artery wall.

### Arterial Stick

Puncture of an artery with a needle, usually to obtain an arterial blood sample. It may cause temporary pain or bruising and, rarely, bleeding, arterial injury, or reduced blood flow.

---

## Arterial Stiffness

Reduced ability of arteries to expand and recoil with each heartbeat. It is associated with ageing, high blood pressure, diabetes, kidney disease, and fatty artery disease and is linked with higher cardiovascular risk.

## Arterial Switch Operation

Open-heart surgery usually performed in newborns whose two main arteries leave the wrong pumping chambers. The arteries and coronary arteries are reconnected so blood flows normally; lifelong heart follow-up is needed.

### Alternate Names:

Jatene Procedure; ASO

## Arterial Tension

A measurement taken from arterial blood. Depending on the context, it may mean blood pressure inside an artery or the amount of oxygen or carbon dioxide dissolved in the blood.

## Arterial Thromboembolism

Blockage of an artery by a blood clot that formed elsewhere and travelled through the bloodstream. It can suddenly reduce blood flow to the brain, heart, limb, kidney, intestine, or another organ.

## Arterial Thrombosis

Formation of a blood clot inside an artery, often where a fatty plaque has damaged the vessel lining. The clot can block blood flow to the heart, brain, limb, intestine, or another organ.

## Arterial Ulcer

A slow-healing wound caused by poor blood flow through an artery, usually on a toe, heel, or other pressure point. It may be deep and sharply outlined, with a pale or dead-looking base and cool, shiny, hairless surrounding skin.

## Arteries

Blood vessels that carry blood away from the heart. Most carry oxygen-rich blood, but the pulmonary and umbilical arteries carry oxygen-poor blood;

strong elastic and muscular walls help withstand and regulate pressure.

### Alternate Names:

Arterial Vessels

## Arteriogram

An image of an artery, usually made after contrast material is injected. It can show narrowing, blockage, aneurysm, abnormal connections, or other changes in blood flow.

## Arteriole

A small branch of an artery that delivers blood to a network of tiny vessels called capillaries. Its muscular wall can narrow or widen to control blood flow and blood pressure.

## Arterioles

Small branches of arteries that control blood flow into capillaries by narrowing or widening their muscular walls. They are important in regulating blood pressure and the supply of blood to tissues.

### Alternate Names:

Resistance Vessels; Small Arteries

## Arteriolosclerosis

A form of vascular disease affecting small arteries and arterioles, characterized by mural thickening, luminal narrowing, and loss of elasticity, classically manifesting as hyaline arteriolosclerosis (associated with aging and diabetes) or hyperplastic arteriolosclerosis (associated with severe hypertension).

## Arteriosclerosis

Thickening, hardening, and loss of elasticity in artery walls. It includes several processes, including age-related stiffening and plaque buildup, and can reduce blood flow to the heart, brain, legs, or other organs.

## Arteriosclerosis / Atherosclerosis

Arteriosclerosis means hardening or thickening of artery walls. Atherosclerosis is a type in which fatty, inflammatory plaque builds up inside the walls and may restrict blood flow or rupture.

---

## Arteriosclerotic Retinopathy

Damage or changes in the retina caused by thickened, narrowed, or hardened small blood vessels, most often after long-standing high blood pressure. Severe disease can reduce vision.

## Arteriovenous Fistula

An abnormal connection between an artery and a vein that bypasses the normal capillary network. It may be present from birth, follow injury or a procedure, or be created to provide access for haemodialysis.

## Arteriovenous Fistula Creation

Surgery to connect an artery to a nearby vein to create blood access for hemodialysis. Over time, the vein may become wider and stronger so it can be used for dialysis needles. It often has advantages over a graft or catheter, but its success, risks, and long-term usefulness vary from person to person.

### Alternate Names:

AV Fistula Creation; Arteriovenous Fistula Surgery; Hemodialysis Fistula Placement; AVF Construction

## Arteriovenous Graft

An artificial tube connecting an artery and vein to provide access for haemodialysis, usually when a person's veins cannot form a suitable fistula. It can clot or become infected and needs regular monitoring.

## Arteriovenous Malformation

An abnormal tangle of arteries and veins joined without the usual capillaries between them. It can bleed, cause seizures or other neurological symptoms, or put extra strain on the heart.

## Arteriovenous Malformation Resection

A complex neurosurgical operation performed to remove an arteriovenous malformation (AVM)—an abnormal, high-pressure tangle of blood vessels connecting arteries directly to veins in the brain without intervening capillaries. Because the thin-walled veins cannot tolerate direct arterial pressure, AVMs carry high risks of catastrophic brain bleeding (hemorrhagic stroke), seizures, and chronic headaches.

Under high-magnification operating microscopes and continuous image guidance, a neurosurgeon enters through a craniotomy, painstakingly identifies and seals off every feeding artery supplying the malformation, coagulates and detaches the draining veins, and completely excises the vascular nidus from surrounding healthy brain tissue to eliminate the danger of hemorrhage permanently.

### Alternate Names:

AVM Resection; Surgical Excision of Brain AVM; Cerebral AVM Removal

## Arteriovenous Nicking

A clinical fundoscopic finding in chronic hypertensive retinopathy characterized by the apparent tapering, deflection, or constriction of a retinal venule where it is crossed by a thickened, sclerotic retinal arteriole (Gunn's sign). Because the arteriole and venule share a common adventitial sheath at crossing points, progressive arteriolosclerosis compresses the thinner-walled venule, indicating sustained moderate-to-severe systemic vascular hypertension and microvascular end-organ damage.

### Alternate Names:

AV Nicking; Gunn's Sign; Arteriovenous Crossing Changes

## Arteritis

Inflammation of an artery wall. The inflammation can narrow, weaken, or block the artery, reducing blood flow and causing pain or organ damage in the affected area.

## Artery

A blood vessel that carries blood away from the heart. Most arteries carry oxygen-rich blood, except the pulmonary arteries, which carry oxygen-poor blood to the lungs. Their elastic, muscular walls withstand and regulate pressure.

## Artery Occlusion

A blockage inside an artery, often caused by a clot, fatty plaque, or material that travelled from elsewhere. It can cut off blood flow; sudden pain, paleness, coldness, numbness, weakness, or absent pulse is an emergency.

---

## Arthralgia

Pain in one or more joints without clear signs of inflammation, such as swelling, warmth, or redness. Causes include overuse, injury, infection, autoimmune disease, and early joint wear.

## Arthritis

A group of conditions that affect joints and the tissues around them; it is not one single disease. Depending on the type, arthritis may involve inflammation, gradual breakdown of joint tissues, infection, or crystal deposits. It can cause joint pain, swelling, warmth, stiffness, and reduced movement. Some types can also affect the skin, eyes, heart, lungs, or other organs. Examples include osteoarthritis, rheumatoid arthritis, gout, and ankylosing spondylitis.

### Alternate Names:

Joint Inflammation; Articular Inflammation

## Arthritis (Osteoarthritis)

Osteoarthritis, a degenerative joint disorder involving cartilage loss, bone remodeling, pain, stiffness, and reduced function. It commonly affects the knees, hips, hands, spine, and feet.

## Arthritis of the Wrist

Inflammation or degenerative disease affecting one or more wrist joints. Pain, swelling, stiffness, reduced grip, and limited motion may result from osteoarthritis, rheumatoid arthritis, crystal disease, injury, infection, or other inflammatory disorders.

## Arthritis Urethritica

An older name for reactive arthritis, inflammatory arthritis occurring after certain gastrointestinal or genitourinary infections and sometimes accompanied by urethritis, conjunctivitis, or skin findings.

## Arthrocentesis

A procedure in which a needle removes fluid from a joint. The fluid can be tested for infection, blood, or crystals and may be removed to reduce pain and pressure.

## Arthrodesis

Surgery that permanently joins the bones of a damaged joint so they heal as one solid unit. It can relieve severe pain and improve stability, but the fused joint no longer moves.

## Arthrogram

An image of a joint taken after contrast material is placed inside or around it. It can show tears or problems in cartilage, ligaments, the joint capsule, or other joint structures.

## Arthrography

Imaging of a joint after contrast material is injected into it. It can help examine the shoulder, hip, wrist, knee, or another joint when ordinary imaging does not show enough detail.

## Arthrogryposis

A group of conditions present from birth in which two or more joints are permanently stiff or limited in movement. It usually results from reduced movement before birth and may involve muscles, nerves, or connective tissue.

## Arthrogryposis Multiplex Congenita

A group of birth conditions causing stiffness and limited movement in several joints. The affected joints may be in the arms, legs, jaw, or spine, and muscle weakness or abnormal nerves and connective tissue may also be present.

## Arthrolysis

A procedure that releases scar tissue or other restrictions around a stiff joint to improve movement. It may be performed through small cuts with a camera or through open surgery.

## Arthropathy

Any disease or abnormal condition affecting a joint. It may result from inflammation, wear, injury, infection, bleeding, nerve damage, or crystals such as those that cause gout.

### Alternate Names:

Joint Disease

---

## Arthroplasty

Surgery to repair, reshape, or replace a damaged joint to reduce pain and improve movement or alignment. It may replace part or all of the joint and is used for severe arthritis, fractures, or other joint damage.

## Arthropod Vector

An insect, tick, or related animal that carries and transmits an infectious organism between hosts. Examples include mosquitoes spreading malaria or ticks spreading Lyme disease.

### Alternate Names:

Disease Vector

## Arthroscope

A thin instrument with a light and camera used to look inside a joint through a small cut. It can help diagnose damage and guide keyhole surgery to repair or remove abnormal tissue.

## Arthroscopy

Keyhole surgery in which a camera and small instruments are inserted through small cuts to examine or treat a joint. It may be used to repair cartilage or ligaments, remove loose tissue, or treat an inflamed joint lining.

## Arthrosis

Degenerative damage or wear of a joint, involving loss of cartilage and changes in the underlying bone. Osteoarthritis is the most common example and can cause pain, stiffness, and reduced movement.

## Arthus Reaction

A local immune reaction that causes redness, swelling, pain, and sometimes tissue damage several hours after exposure to a substance. It occurs when antibodies and the substance form deposits that activate inflammation in the affected tissue.

## Articular Cartilage

Smooth, tough tissue covering the ends of bones inside a joint. It allows bones to move with little friction and helps spread forces across the joint.

## Articular Cartilage Injuries

Damage to the smooth cartilage covering the ends of bones in a joint, sometimes extending into the underlying bone. It may cause pain, swelling, catching, or reduced movement and can follow injury, repeated stress, or joint disease.

## Articular Ligament

A strong band of connective tissue joining bones around a joint. It stabilizes the joint and limits excessive bending, sliding, or twisting.

### Alternate Names:

Joint Ligament

## Articulation

A joint or connection between bones. In speech, articulation also means forming clear speech sounds using the tongue, lips, teeth, and other speech organs.

## Artifacts

Distortions or false features in a medical image that do not represent real body structures. They may result from movement, metal, air, the imaging beam, or limits of the scanner and image-processing method.

## Artificial Abortion

Older term; see *Induced Abortion*.

## Artificial Heart

A mechanical device that supports or replaces the heart's pumping action. It may support one or both lower chambers temporarily or long term and requires management of power, infection, bleeding, and blood clots.

## Artificial Insemination

A fertility procedure in which prepared sperm is placed into the cervix or uterus to increase the chance of pregnancy. The sperm may come from a partner or a donor.

## Artificial Insemination By Donor

Fertility treatment in which sperm from a screened



---

donor is placed into the uterus or cervix to attempt pregnancy. It may be used when partner sperm is unavailable or unlikely to result in pregnancy.

### **Artificial Insemination By Husband**

Fertility treatment in which prepared sperm from a woman's husband or male partner is placed into the uterus or cervix to increase the chance of fertilization and pregnancy.

### **Artificial Intelligence**

Computer systems that learn from data to recognize patterns, make predictions, or support healthcare decisions. Medical AI can assist diagnosis, treatment, and research, but it must be tested, monitored for bias, and supervised by qualified professionals.

**Alternate Names:**

AI

### **Artificial Kidney**

A dialysis device that removes metabolic waste and excess fluid from blood and helps correct electrolyte disturbances when renal function is inadequate.

### **Artificial Pacemaker**

An implanted device that sends electrical signals to the heart when its natural rhythm is too slow or irregular. It helps maintain an adequate heart rate and may have one or more leads.

### **Artificial Rupture Of Membranes**

Deliberate breaking of the amniotic sac during labour or an examination. It may be used to start or strengthen labour, assess the amniotic fluid, or attach internal monitors; it carries risks such as infection and umbilical-cord prolapse.

### **Artificial Saliva**

A product that moistens the mouth and temporarily relieves discomfort from dry mouth. It may improve speaking, swallowing, and comfort but does not replace fluids, dental care, or treatment of the cause.

**Alternate Names:**

Saliva Substitute

### **Artificial Skin**

A biological or manufactured covering placed over a wound to protect it and support growth of new skin. It may reduce fluid loss and contamination, but some wounds later need a permanent skin graft.

**Alternate Names:**

Skin Substitute

### **Artificial Sweeteners**

Sweet-tasting substances that provide few or no calories and are used instead of sugar. They can reduce added sugar for some people but do not by themselves make a diet healthy; effects and safety depend on the product and amount.

### **Artificial Urinary Sphincter**

An implanted device that controls urine leakage by gently closing and releasing the urethra. It is used for selected severe stress incontinence and requires surgery and follow-up to operate and maintain the device.

### **Artificial Ventilation**

Support or replacement of breathing with a machine that moves air and oxygen into and out of the lungs when a person cannot breathe well enough alone. It may use a mask or a tube and requires close monitoring.

### **Artificial Ventricle**

A mechanical device that helps or replaces the pumping action of one or both lower heart chambers in severe heart failure. The exact device and treatment goal vary; it is not the same as every ventricular assist device or a total artificial heart.

**Alternate Names:**

Total Artificial Heart; TAH; Ventricular Assist Device

### **Aryepiglottic Fold**

A triangular mucosal fold enclosing ligamentous and muscular fibers, extending bilaterally from the lateral margins of the epiglottis to the arytenoid cartilages, forming the lateral anatomical boundaries of the laryngeal inlet.

---

## Arytenoid Cartilage

One of a pair of small cartilages at the back of the voice box. They anchor the vocal cords and help the laryngeal muscles open, close, and adjust the cords for breathing and speech.

## ASA Physical Status Classification System

A scale used before anesthesia to summarize a patient's overall health and major medical conditions, from healthy to severely ill. It helps the care team communicate risk but does not by itself predict complications or outcome.

## Asbestos-Related Disorders

Diseases that may develop years after breathing asbestos fibres, including pleural plaques, lung scarring, lung cancer, and mesothelioma. Risk increases with greater exposure and smoking; symptoms may include breathlessness, cough, or chest pain.

## Asbestosis

Permanent scarring of the lungs caused by long-term inhalation of asbestos fibres. It can gradually cause breathlessness, dry cough, and reduced lung function and increases the risk of lung cancer and mesothelioma.

## Ascariasis

Infection with a roundworm acquired by swallowing eggs from contaminated soil, food, or water. It may cause no symptoms, abdominal pain, poor nutrition, intestinal blockage, or cough while larvae pass through the lungs.

## Ascending Aorta

The first section of the aorta, rising from the heart's left pumping chamber. It gives rise to the coronary arteries and continues into the aortic arch; an aneurysm or dissection here can be life-threatening.

## Ascending Aortic Aneurysm

An enlarged, weakened area in the part of the aorta that rises from the heart. It may be associated with a bicuspid aortic valve, inherited connective-tissue disease, high blood pressure, or artery-wall disease

and may rupture or dissect.

## Ascending Cholangitis

A life-threatening bacterial infection of the biliary tree caused by a combination of bile duct obstruction (most commonly a gallstone in the common bile duct or a tumor) and bacterial ascent from the duodenum into stagnant bile under pressure. It characteristically presents with Charcot's triad (fever with chills, right-upper-quadrant pain, and jaundice), which can rapidly progress to Reynolds' pentad (adding confusion and septic shock/hypotension in acute suppurative cases). Blood tests show leukocytosis and hyperbilirubinemia, and imaging (ultrasound, CT, or MRCP) confirms biliary duct dilation. Emergency management consists of resuscitation with intravenous fluids, broad-spectrum IV antibiotics, and urgent biliary decompression via endoscopic retrograde cholangiopancreatography (ERCP).

### Alternate Names:

Acute Cholangitis; Acute Suppurative Cholangitis; Biliary Sepsis

## Ascending Colon

The first part of the large intestine, running upward on the right side of the abdomen from the appendix region to the bend beneath the liver. It absorbs water and helps form stool.

## Ascites

Abnormal buildup of fluid inside the abdomen. It commonly results from advanced liver disease but may also be caused by cancer, heart failure, kidney disease, infection, or pancreatitis and can cause swelling, discomfort, or breathlessness.

## Ascitic Fluid

Fluid collected from the abdomen in a person with ascites. Laboratory testing can help identify liver disease, infection, cancer, pancreatitis, or other causes of the fluid buildup.

### Alternate Names:

Peritoneal Fluid; Ascitic Effusion

## Ascorbic Acid

Vitamin C, a water-soluble vitamin needed to make

---

collagen, heal wounds, absorb iron, protect cells, and support normal immune and connective-tissue function.

### **Ascus**

A Pap-test result meaning that some cells from the cervix look slightly unusual, but it is unclear whether the change is important. Follow-up depends on age, HPV testing, previous results, and clinical guidance; it is not a cancer diagnosis.

### **Asepsis**

The condition or practice of preventing contamination by disease-causing microorganisms, especially during medical or surgical procedures. It includes sterile technique, environmental controls, hand hygiene, and appropriate handling of instruments and materials.

### **Aseptic**

Free from disease-causing contamination, or performed using methods that prevent microorganisms from entering a wound, specimen, medicine, or body site. Aseptic does not necessarily mean completely sterile.

### **Aseptic Loosening**

Loss of firm attachment of an artificial joint or other implant without an active infection. Repeated stress or a reaction to tiny wear particles can cause pain, instability, and eventual implant failure.

### **Aseptic Meningitis**

Inflammation of the membranes around the brain and spinal cord without the usual evidence of bacterial meningitis. Viruses are common causes, but medicines, autoimmune disease, and other infections can also be responsible.

### **Aseptic Necrosis**

See *Avascular necrosis (osteonecrosis)*.

#### **Alternate Names:**

Avascular Necrosis; Osteonecrosis; Ischemic Bone Necrosis

### **Aseptic Technique**

Practices used to prevent harmful microorganisms contaminating a wound, body site, medicine, or specimen. They include hand hygiene, clean or sterile equipment, surface cleaning, and limiting contact with exposed sterile items.

### **Asherman Syndrome**

Scar tissue inside the uterus that partly or completely joins the uterine walls. It most often follows a procedure inside the uterus or complications of pregnancy and may cause light or absent periods, infertility, or repeated miscarriage.

### **Ashman Phenomenon**

A temporary wide-looking heartbeat that occurs when a short beat follows a long pause, often during atrial fibrillation. It can resemble a dangerous rhythm on an ECG, so clinicians interpret it with the full tracing and the person's symptoms.

#### **Alternate Names:**

Ashman's Phenomenon; Aberrant Intraventricular Conduction

### **ASL**

American Sign Language, a complete visual language used by many Deaf and hard-of-hearing people. Healthcare communication should respect each person's language preference and provide qualified interpreters when needed.

### **ASOT**

An abbreviation for the antistreptolysin O test, a blood test measuring antibodies against a toxin made by group A streptococcal bacteria. It can help show a recent strep infection when investigating later complications.

### **Asp**

The standard three-letter abbreviation for aspartic acid, an amino acid used to build proteins and involved in energy production and other cell reactions.

### **Aspartame**

A low-calorie sweetener made from two amino-acid

---

components. It is not suitable for people with phenylketonuria because it releases phenylalanine when digested.

### **Aspartate Aminotransferase**

An enzyme found mainly in the liver, heart, muscles, and red blood cells. When these tissues are injured, the enzyme can enter the blood and its level may rise.

### **Aspartate Aminotransferase Blood Test**

A blood test measuring the level of the enzyme aspartate aminotransferase. A high result can indicate injury to the liver, heart, muscles, or red blood cells and must be interpreted with other tests.

### **Aspartic Acid**

An amino acid that the body can make and uses to build proteins and carry out cell reactions. It is one of the components of aspartame but is not the same substance as the sweetener.

### **Asperger Syndrome**

A former diagnostic classification for a neurodevelopmental profile within the autism spectrum characterized by qualitative differences in social communication and social reciprocity, accompanied by restricted, repetitive patterns of behavior and intense focal interests, without significant delays in general language acquisition or cognitive development.

#### **Alternate Names:**

Asperger Disorder; Asperger's Syndrome; Asperger's Disorder; Autistic Psychopathy

### **Aspergillosis**

A group of infections and allergic conditions caused by *Aspergillus* molds. It may affect the airways, sinuses, or lungs and can include allergic disease, a fungal growth in an existing lung cavity, or invasive infection.

#### **Alternate Names:**

Aspergillus Infection; Aspergillus Disease

### **Aspergillosis Precipitin**

A qualitative or semi-quantitative serological test detecting circulating precipitating IgG antibodies against *Aspergillus* antigens, useful in diagnosing chronic pulmonary aspergillosis and allergic bronchopulmonary aspergillosis.

#### **Alternate Names:**

Aspergillus Antibodies; Aspergillus IgG Precipitins

### **Aspergillus**

A genus of saprophytic molds whose airborne conidia are ubiquitous in the environment, capable of causing allergic bronchopulmonary aspergillosis, aspergilloma formation in pre-existing cavitary lung lesions, or severe invasive tissue infections in immunocompromised hosts.

### **Aspergillus Infection**

Alternate index label; see *Aspergillosis*.

### **Asphalt Cement Poisoning**

Illness caused by swallowing, inhaling, or having significant skin contact with hot or solvent-containing asphalt cement. Burns, breathing problems, dizziness, or chemical injury can occur; significant exposure requires urgent medical treatment.

### **Asphyxia**

A life-threatening lack of oxygen caused by inability to breathe or by blocked blood oxygen delivery, such as from choking, drowning, strangulation, or severe airway blockage. It can quickly cause unconsciousness, brain injury, or death.

### **Asphyxia Neonatorum**

Failure of a newborn to start or maintain effective breathing at or soon after birth. It can reduce oxygen to the brain and other organs and requires immediate assessment and breathing support.

### **Asphyxial Death Pathology**

Autopsy findings associated with impaired oxygenation or ventilation, which may include congestion, petechiae, pulmonary edema, and visceral hypoxia. These findings are usually nonspecific and must be

---

interpreted with scene, history, and toxicology.

### **Asphyxiating Thoracic Dysplasia**

A rare inherited disorder in which short ribs create a narrow, stiff chest that restricts breathing, especially in infancy. Kidney, liver, and limb abnormalities may also occur.

#### **Alternate Names:**

Jeune Syndrome

### **Aspirate**

Fluid, cellular material, tissue fragments, or particulate matter evacuated from a cavity, cyst, organ, or lesion via negative-pressure suction; or foreign liquid and solid material accidentally inhaled into the tracheobronchial tree.

#### **Alternate Names:**

Aspirated Material; Biopsy Aspirate

### **Aspiration**

Breathing food, liquid, saliva, vomit, blood, or another substance into the voice box or lungs instead of swallowing it into the oesophagus. It can cause choking, airway blockage, inflammation, or pneumonia.

### **Aspiration Pneumonia**

A lung infection that develops after inhaling saliva, food, liquid, or stomach contents. It is more likely with swallowing difficulty, reduced consciousness, stroke, or tube feeding and may cause cough, fever, breathlessness, or chest pain.

### **Aspiration Pneumonitis**

Chemical inflammation of the lungs after inhaling stomach contents, especially acidic fluid. It can cause coughing, wheezing, low oxygen, breathing difficulty, and fluid or collapse in parts of the lungs.

### **Aspiration Pneumonitis and Pneumonia**

Aspiration pneumonitis is an acute chemical lung injury after inhaling gastric contents; aspiration pneumonia is infection after inhaling contaminated material. Fever, cough, hypoxia, or infiltrates after an aspiration event require assessment, with treat-

ment guided by severity and evidence of infection.

### **Aspiration Risk**

The likelihood that food, liquid, saliva, or stomach contents will enter the airway or lungs. Risk increases with swallowing difficulty, stroke, reduced alertness, nerve or muscle disease, reflux, and some dental or feeding problems.

### **Aspirin**

A medicine that can relieve pain, lower fever, reduce inflammation, and make platelets less likely to form blood clots. Low-dose aspirin is often used after a heart attack, stroke, or some heart procedures, but it is not suitable for everyone because it can cause serious bleeding. It should not be taken daily for prevention without medical advice.

#### **Alternate Names:**

Acetylsalicylic Acid; ASA

### **Aspirin Overdose**

Acute or chronic poisoning resulting from excessive salicylate ingestion. Characteristic features include tinnitus, nausea, vomiting, hyperventilation causing early respiratory alkalosis, followed by severe uncoupled oxidative phosphorylation, high anion gap metabolic acidosis, hyperthermia, confusion, seizures, and noncardiogenic pulmonary edema.

#### **Alternate Names:**

Salicylate Toxicity; Aspirin Poisoning; Salicylism

### **Aspirin Poisoning**

Alternate name; see *Aspirin Overdose*.

### **Aspirin-Exacerbated Respiratory Disease**

A condition involving asthma, chronic nasal or sinus inflammation with polyps, and respiratory reactions to aspirin or some similar pain medicines. Exposure may trigger wheezing, nasal blockage, cough, or shortness of breath.

### **Assay**

A laboratory method used to detect or measure a substance, biological activity, genetic feature, or response in a sample.



---

### **Assist-Control Ventilation**

A ventilator setting that gives a full preset breath whenever the patient starts one and also gives breaths at a minimum rate if the patient does not breathe enough. Each breath is set by volume or pressure.

### **Assisted Breech Delivery**

A vaginal birth in which the baby's buttocks or feet come first and a trained clinician helps deliver the body and head. It is considered only in carefully selected circumstances because breech birth can be dangerous for the baby or pregnant person.

### **Assisted Circulation**

Mechanical or procedural support that maintains blood flow when the heart cannot pump enough on its own. It may use a pump, oxygenator, or other device for a short time or longer-term support.

#### **Alternate Names:**

Mechanical Circulatory Support

### **Assisted Delivery With Forceps**

A vaginal birth in which a trained clinician places specially shaped instruments around the baby's head to help complete delivery. It may be considered when labour is prolonged or the baby or pregnant person needs delivery sooner.

### **Assisted Living**

A residential setting where people receive help with daily activities such as bathing, dressing, meals, medicines, or transport while retaining some independence. It generally does not provide the continuous skilled nursing care of a hospital or nursing home.

### **Assisted Reproductive Technology**

Clinical, surgical, and laboratory procedures that involve the in vitro manipulation of human oocytes, spermatozoa, or embryos to achieve pregnancy, including in vitro fertilization (IVF), intracytoplasmic sperm injection (ICSI), gamete intrafallopian transfer (GIFT), and frozen embryo transfer.

#### **Alternate Names:**

ART; Assisted Reproduction; Assisted Conception

### **Assistive Device**

Equipment that helps a person move, communicate, perform daily activities, or remain independent. Examples include a cane, wheelchair, hearing aid, communication tool, brace, walker, or adapted household equipment.

### **Assistive Device Training**

Instruction in choosing, fitting, using, and maintaining equipment such as walkers, canes, crutches, braces, or wheelchairs, including safe walking, transfers, and stair use.

### **Association**

A relationship in which two findings, events, or conditions occur together more often than expected. An association alone does not prove that one causes the other.

### **Association Cortex**

Areas of the brain's outer layer that combine information from different senses. They support memory, language, planning, judgment, learning, and other complex behavior rather than controlling one simple movement or sensation.

### **Association Study**

A research study examining whether a genetic variant, exposure, or other factor occurs more often with a particular disease or finding. Such a relationship does not prove that the factor causes the disease.

### **Asteatotic Eczema**

A dry-skin eczema pattern, also called eczema craquelé, in which fine fissures create a cracked or porcelain-like appearance, usually on the shins, arms, or trunk of older adults. Cold, dry weather, frequent washing, and diuretics can worsen it; emollients and treatment of inflammation help.

### **Astereognosis**

Inability to recognize a familiar object by touch alone even though basic touch sensation works. It usually results from damage to a brain area that combines sensory information, often after a stroke



---

or other brain injury.

### **Asterixis**

Brief, repeated lapses in holding a posture that cause a flapping movement, often when the arms are stretched out and the wrists are bent back. It can occur with liver or kidney failure, breathing disorders, or some medicines.

### **Asteroid Hyalosis**

A usually harmless eye condition in which tiny calcium-and-fat particles float in the gel inside the eye and appear as bright spots. It often affects one eye and causes few symptoms, but dense particles can obscure the retina.

### **Asthenia**

Abnormal lack of physical strength or energy. It is a symptom rather than a disease and may occur with infection, chronic illness, malnutrition, depression, or medicine effects.

### **Asthenic**

Describing a person or condition marked by weakness, or sometimes a slender body build. In a clinical note, the surrounding context shows which meaning is intended; asthenia specifically means reduced strength or energy.

### **Asthenopia**

Eye strain or visual fatigue that may cause aching, burning, blurred vision, headache, or difficulty focusing. Refractive errors, dry eyes, poor lighting, and prolonged screen use can contribute.

### **Asthenozoospermia**

Abnormally reduced movement of sperm, which can make fertilization more difficult. It is assessed by semen analysis and may result from infection, a varicocele, toxins, illness, or an unknown cause.

#### **Alternate Names:**

Asthenospermia

### **Asthma**

A long-term inflammatory disease of the airways in

the lungs. The airways are overly sensitive and can narrow when triggered because their lining swells, the surrounding muscles tighten, and extra mucus forms. This variable narrowing makes it harder for air to move, especially when breathing out. Symptoms may include wheezing, cough, chest tightness, and shortness of breath; they can come and go and may be triggered by allergens, infections, smoke, air pollution, exercise, certain medicines, or cold air.

### **Asthma Attack**

A sudden worsening of asthma in which inflamed, narrowed airways make breathing difficult. It may cause wheezing, cough, chest tightness, or breathlessness; trouble speaking, blue lips, severe exhaustion, or poor response to a reliever is an emergency.

### **Asthma Controller Medications**

Medicines used regularly to reduce airway inflammation and prevent asthma symptoms and attacks. Inhaled corticosteroids are the main controller medicine. Other medicines may be added when needed; long-acting bronchodilators should not be used alone for asthma.

#### **Alternate Names:**

Asthma Controllers; Long-Term Control Medications; Inhaled Corticosteroids for Asthma

### **Asthma Exacerbation**

A sudden or gradual worsening of asthma symptoms and airflow through the lungs. It may cause increasing wheeze, cough, chest tightness, or breathlessness and can become life-threatening.

### **Asthma in Older Adults**

Asthma in later life may be underdiagnosed because symptoms overlap with COPD, heart disease, deconditioning, and medication effects. Diagnosis uses history and objective airflow testing when possible; inhaler technique, comorbidities, cognition, and treatment adverse effects need special attention.

### **Asthma in Pregnancy**

Asthma may improve, worsen, or remain unchanged during pregnancy. Maintaining control with appropriate inhaled treatment is generally safer than al-

---

lowing maternal hypoxia; worsening wheeze, breathlessness, or poor response to rescue medicine requires prompt care.

### **Asthma Medications**

Medicines for asthma include regular treatments that reduce airway inflammation and reliever treatments that quickly ease narrowed airways. The regimen depends on symptoms, lung function, attack risk, age, pregnancy, inhaler technique, and response.

### **Asthma Quick-Relief Medications**

Medicines used to ease asthma symptoms quickly by opening narrowed airways. They may include a short-acting bronchodilator or, for some people, an inhaled corticosteroid–formoterol inhaler. The correct reliever depends on the person’s treatment plan.

#### **Alternate Names:**

Asthma Relievers; Rescue Inhalers; Short-Acting Beta-Agonists; SABA

### **Asthma-COPD Overlap**

A pattern in which a person has features of both asthma and chronic obstructive pulmonary disease, such as variable symptoms together with persistent airflow limitation. Diagnosis and treatment require individual assessment because the conditions and risks differ.

### **Astigmatism**

A focusing problem caused by an unevenly curved cornea or lens. Light is not focused at one clear point on the retina, causing blurred or distorted vision at near or far distances.

### **Astringent**

A substance that makes surface tissues contract and may reduce minor secretions or bleeding. It is used in some skin, mouth, or wound products but can irritate tissue and does not replace assessment of persistent bleeding or serious injury.

### **Astrocyte**

A star-shaped support cell in the brain and spinal

cord. It helps nourish nerve cells, maintain the blood–brain barrier, regulate the chemical environment, and respond to injury.

### **Astrocytoma**

A tumour arising from astrocytes, the star-shaped support cells of the brain or spinal cord. Astrocytomas vary from slow-growing to aggressive, and symptoms depend on their location and size.

### **Astrovirus Infection**

An acute viral infection of the gastrointestinal tract caused by human astroviruses, a leading cause of non-bacterial gastroenteritis primarily in infants, young children, the elderly, and immunocompromised patients. It presents with watery diarrhea, mild abdominal pain, low-grade fever, and vomiting, typically resolving spontaneously within 2 to 4 days with oral rehydration therapy.

#### **Alternate Names:**

Astrovirus Gastroenteritis; Human Astrovirus Infection

### **Asymmetrical**

Unequal in shape, size, distribution, or appearance between corresponding sides. Asymmetry in a pigmented skin lesion can be a warning sign that needs assessment, especially with other changes.

### **Asymptomatic**

Having a disease, infection, or abnormal finding without noticeable symptoms. A person may still have detectable disease, develop complications, or transmit an infection; this differs from presymptomatic illness, in which symptoms appear later.

### **Asymptomatic Adults**

Adults who have no noticeable symptoms of the condition being assessed. Absence of symptoms does not exclude early or hidden disease, risk factors, or the ability to transmit an infection.

### **Asymptomatic Bacteriuria**

Bacteria detected in a properly collected urine sample when the person has no urinary symptoms. It does not always require treatment, but management

---

is important in selected situations such as pregnancy or before some urinary-tract procedures.

### **Asymptomatic Bacteriuria In Pregnancy**

Bacteria growing in a pregnant person's urine without burning, urgency, or other urinary symptoms. It is identified by urine culture and is important because untreated infection increases the risk of kidney infection and pregnancy complications.

### **Asymptomatic Carrier**

A person who carries an infectious organism without symptoms but may still transmit it to others. Whether transmission occurs depends on the organism, body site, and contact.

### **Asymptomatic HIV Infection**

HIV infection without noticeable symptoms at present. HIV can still multiply and gradually weaken the immune system, and the person may transmit it without effective treatment.

### **Asymptomatic Infections**

Infections in which the person has no noticeable symptoms. The organism may still be detectable, cause later complications, or spread to others; this differs from presymptomatic infection, in which symptoms develop later.

#### **Alternate Names:**

Subclinical Infections

### **Asystole**

Absence of effective electrical activity in the heart's lower chambers and absence of useful pumping. It causes cardiac arrest and requires immediate resuscitation; a flat-looking ECG line must first be checked to ensure it is not an equipment or lead problem.

### **At-Home COVID-19 Testing**

At-home testing for COVID-19 commonly uses a rapid antigen test from a nasal sample. A negative result does not always exclude early infection; repeat testing or a molecular test may be needed when suspicion remains high.

### **Ataxia**

Loss of smooth control and coordination of movement. It is a sign rather than one specific disease and may cause an unsteady walk, poor balance, clumsy movements, slurred speech, shaking, or abnormal eye movements. It can result from a problem in the cerebellum, spinal cord, sensory nerves, or balance organs and is different from muscle weakness alone. Causes include inherited disorders, stroke, multiple sclerosis, infection, vitamin deficiency, medicines, alcohol, or other toxins. The onset may be sudden or gradual, and the course depends on the cause.

### **Ataxia Telangiectasia**

A rare autosomal-recessive DNA-repair disorder causing progressive cerebellar ataxia, telangiectasias of the eyes or skin, immune deficiency, radiosensitivity, and increased risk of leukemia, lymphoma, or other cancers. Respiratory infections and elevated alpha-fetoprotein may occur.

#### **Alternate Names:**

A-T; Ataxia-Telangiectasia

### **Ataxic Breathing**

An irregular, unpredictable pattern in which the rate and depth of breaths vary randomly. It usually indicates severe injury to the lower brainstem and may precede breathing failure.

### **Ataxic Gait**

An abnormal, unsteady, and poorly coordinated walking pattern resulting from dysfunction in the motor coordination or sensory balance pathways. In cerebellar ataxia, the gait is characteristically wide-based, irregular, lurching, and clumsy, with difficulty maintaining a straight line and inability to perform tandem walking. In sensory ataxia (caused by dorsal column or peripheral proprioceptive loss), the gait is wide-based with high-stepping stamping of the feet, markedly worsening with eye closure (positive Romberg sign).

#### **Alternate Names:**

Cerebellar Gait; Sensory Ataxic Gait; Uncoordinated Gait

---

### **Atelectasis**

Partial or complete collapse of a region of lung, reducing the amount of air it contains. It may result from blocked airways, pressure from outside the lung, shallow breathing, or scarring and can reduce oxygen exchange.

#### **Alternate Names:**

Lung Collapse

### **Athelia**

Absence of one or both nipples, present from birth or acquired after surgery, injury, or disease. It may occur alone or with other differences in breast or chest-wall development.

### **Atherectomy**

A minimally invasive endovascular catheter technique utilizing directional, rotational, orbital, or laser devices to physically excise, shave, vaporize, or pulverize calcified or atheromatous plaque from the intraluminal wall of diseased arteries.

#### **Alternate Names:**

Endovascular Atherectomy; Directional Atherectomy; Rotational Atherectomy; Orbital Atherectomy

### **Atheroembolic Renal Disease**

Kidney injury caused when cholesterol crystals break from an artery plaque and block small kidney vessels, often after a vascular procedure. It may cause worsening kidney function, mottled skin, painful toes, or raised eosinophils.

### **Atheroma**

A fatty, inflamed buildup within an artery wall that forms part of an atherosclerotic plaque. It may narrow the artery or rupture and trigger a blood clot.

#### **Alternate Names:**

Atheromatous Plaque; Arterial Plaque

### **Atheromatous Plaque**

An asymmetrical, fibrofatty lesion developing within the tunica intima of medium and large arteries, comprising a necrotic lipid-rich core of oxidized

cholesterol crystals and apoptotic debris covered by a fibrous cap of collagen and vascular smooth muscle cells.

#### **Alternate Names:**

Atheroma; Atherosclerotic Plaque

### **Atherosclerosis**

A disease in which fatty, cholesterol-rich plaques build up inside artery walls. Plaques can narrow arteries or rupture and form clots, reducing blood flow to the heart, brain, legs, or other organs.

### **Atherosclerosis Pathology**

A chronic arterial intimal disease characterized by lipid-rich plaques with a fibrous cap, inflammation, smooth-muscle proliferation, calcification, and possible ulceration or thrombosis.

### **Atherosclerotic**

Relating to fatty plaque buildup and hardening inside artery walls. Atherosclerotic changes can narrow an artery or encourage a clot to form.

### **Atherosclerotic Heart Disease**

A disease in which cholesterol-containing plaque builds up inside the heart's arteries, narrowing them and reducing blood flow. It may cause chest pain, heart attack, or other heart problems.

### **Atherosclerotic Plaque**

A buildup of cholesterol, fat, inflammatory cells, and scar-like tissue inside an artery wall. It can gradually narrow the artery or rupture and cause a blood clot.

### **Atherothrombotic Stroke**

An ischemic stroke caused by a clot forming on fatty plaque in a large artery supplying the brain or travelling from that plaque into a brain artery. It deprives brain tissue of oxygen.

#### **Alternate Names:**

Large-Artery Atherosclerotic Stroke; Atherothrombotic Ischemic Stroke

---

## Athetosis

Slow, involuntary, writhing movements that may affect the hands, fingers, feet, face, neck, or trunk. They result from abnormal function in brain areas that help control movement and may occur with inherited or acquired brain injury.

## Athlete Foot

Alternate name; see *Athlete's Foot*.

## Athlete's Foot

A common superficial fungal infection (dermatophytosis) of the skin of the feet, most commonly caused by *Trichophyton rubrum* or *Trichophyton interdigitale*. Classically presents with pruritus, erythema, scaling, maceration, and fissuring, especially in the interdigital web spaces.

### Alternate Names:

Tinea Pedis; Foot Ringworm; Foot Dermatophytosis

## Athlete's Heart

Normal physiological cardiac remodeling induced by intensive, prolonged physical endurance or strength training, manifested as physiological left ventricular hypertrophy, chamber dilation, and resting sinus bradycardia, without diastolic dysfunction.

### Alternate Names:

Athletic Heart Syndrome; Athlete Heart Remodeling

## Atlanto-Occipital Dislocation

A severe injury in which the first neck bone separates from the base of the skull. It is extremely unstable and can damage the spinal cord or brainstem, so the neck must be immobilized and emergency care provided immediately.

## Atlanto-Occipital Joints

The paired joints between the upper surface of the atlas, the first cervical vertebra, and the occipital condyles at the base of the skull. They mainly allow nodding and small side-to-side movements while helping stabilize the head and upper neck.

## Atlantoaxial Instability

Excessive movement between the first and second cervical vertebrae, sometimes caused by trauma, rheumatoid arthritis, congenital conditions, or ligament weakness. It may cause neck pain, occipital headache, neurologic symptoms, or spinal-cord compression.

## Atlantoaxial Instability in Down Syndrome

Excessive movement between the first and second cervical vertebrae in a person with Down syndrome, often related to ligament laxity or altered bone development. Many people have no symptoms, but neck pain, weakness, gait change, or other neurologic signs require prompt assessment before activities that stress the neck.

## Atlantoaxial Joint

The complex synovial articulation between the first cervical vertebra (atlas, C1) and the second cervical vertebra (axis, C2). Comprising one median pivot joint (dens with anterior arch) and two lateral facet joints, it provides approximately 50% of total cervical rotational range of motion. Instability may arise from trauma (odontoid fractures), rheumatoid arthritis, or Down syndrome.

### Alternate Names:

Atlas-Axis Joint; C1-C2 Articulation

## Atlas

The first bone of the neck, supporting the skull and helping the head nod. It is ring-shaped and has openings through which important blood vessels pass.

## Atom

The smallest unit of a chemical element that retains the element's properties. It contains a central nucleus surrounded by electrons.

## Atonic

Lacking normal muscle tone or ability to contract. For example, an atonic bladder does not squeeze effectively to empty urine.

---

### Atonic Seizure

A seizure causing sudden loss of muscle tone, often for only a few seconds. The person may drop their head, knees, or whole body and fall; injuries can occur during the fall.

#### Alternate Names:

Drop Attack; Akinetic Seizure

### Atopic

Having a tendency to develop allergic conditions such as eczema, allergic rhinitis, or asthma. This tendency may run in families and reflects increased sensitivity of the immune system.

### Atopic Dermatitis

A chronic, relapsing, intensely itchy inflammatory skin disease involving epidermal-barrier dysfunction and immune dysregulation. It causes dry, inflamed, scaly, or lichenified lesions whose distribution varies with age; scratching can produce infection, excoriations, and sleep disturbance.

#### Alternate Names:

Atopic Eczema; Eczema; Infantile Eczema; Allergic Dermatitis

### Atopic Eruption Of Pregnancy

An itchy eczema-like rash that begins or worsens during pregnancy, often on the face, neck, chest, or skin folds. It usually improves after delivery and is not harmful to the fetus.

### Atopic Rhinitis

Allergy-related inflammation inside the nose. Exposure to pollen, dust mites, mould, animals, or other triggers can cause sneezing, itching, a blocked or runny nose, and reduced smell.

#### Alternate Names:

Allergic Rhinitis

### Atopy

A genetic tendency to develop allergic conditions such as eczema, allergic rhinitis, asthma, or food allergy after exposure to common triggers.

### Atorvastatin

A medicine that lowers low-density lipoprotein cholesterol by reducing cholesterol production in the liver. It lowers the risk of heart attack and stroke in appropriate people; muscle symptoms and liver problems are uncommon but important side effects.

### Atovaquone–Proguanil

A combination of medicines used to prevent or treat some forms of malaria. The medicines interfere with energy production and folate use in malaria parasites; suitability depends on the type of malaria and the person's health.

### ATP

Adenosine triphosphate, the main short-term energy-carrying molecule in cells. Cells release usable energy when ATP is broken down and remake it during metabolism.

### ATP Synthase

A protein machine in mitochondria and some other cell membranes that uses the movement of hydrogen ions to make ATP, the main molecule cells use for immediate energy.

### Atransferrinemia

A very rare inherited disorder in which the blood has little or no transferrin, the protein that carries iron. It can cause severe anemia, poor growth, and harmful iron buildup in organs.

### Atresia

A birth defect in which a normal opening or passage is absent, closed, or severely narrowed. Examples include intestinal, bile-duct, heart-valve, and ear-canal atresia; effects depend on the structure involved.

### Atresia Of Bile Ducts

See *Biliary atresia*.

### Atria

The two upper chambers of the heart. The right atrium receives blood from the body and the left



---

atrium receives blood from the lungs; each passes blood to the lower pumping chamber on its side.

### **Atrial**

Relating to either of the heart's two upper chambers, which receive blood and pass it to the lower pumping chambers.

### **Atrial Ablation**

A procedure that uses heat, cold, or another form of energy to create small scars in heart tissue responsible for an abnormal atrial rhythm. The scars block unwanted electrical signals.

#### **Alternate Names:**

Cardiac Ablation

### **Atrial Arrhythmia**

An abnormal heart rhythm that begins in one of the heart's upper chambers. It may cause palpitations, breathlessness, dizziness, chest discomfort, or fainting; examples include atrial fibrillation and atrial flutter.

### **Atrial Fib**

Alternate name; see *Atrial Fibrillation*.

### **Atrial Fibrillation**

An abnormal heart rhythm in which the heart's upper chambers (atria) beat chaotically instead of contracting in an organized way. The heartbeat is often irregular and may be fast, but some people have no symptoms. Blood may pool in the atria and form clots, increasing the risk of stroke.

#### **Alternate Names:**

AF; AFib; Auricular Fibrillation

### **Atrial Fibrillation Ablation**

A catheter procedure that creates small scars, usually around the pulmonary veins, to block electrical signals that trigger atrial fibrillation. It may reduce symptoms, but recurrence and procedure-related risks remain possible.

### **Atrial Fibrillation Medicines**

Medicines used for atrial fibrillation may slow the

heart rate, restore or maintain a regular rhythm, or reduce stroke risk by preventing clots. Choice depends on symptoms, heart structure, kidney function, bleeding risk, and other conditions.

### **Atrial Flutter**

A supraventricular arrhythmia characterized by a rapid, organized, macro-reentrant electrical circuit within the atria, typically circulating around the cavotricuspid isthmus of the right atrium. Manifested on ECG as rapid, regular sawtooth flutter waves (typically 250–350 bpm) with variable AV nodal conduction block (often 2:1), it presents with palpitations, fatigue, and elevated thromboembolic risk.

#### **Alternate Names:**

AFlutter; Cavotricuspid Isthmus-Dependent Flutter

### **Atrial Myxoma**

A usually noncancerous tumour arising inside the heart, most often in the left atrium. It may obstruct blood flow, cause breathlessness or fainting, or release fragments that block blood vessels.

### **Atrial Natriuretic Peptide**

A hormone released by the heart's upper chambers when they stretch from extra blood. It tells the kidneys to remove more salt and water and helps relax blood vessels, lowering blood volume and pressure.

### **Atrial Septal Defect**

A hole present from birth in the wall between the heart's upper chambers. A small defect may cause no problems. A larger one can increase blood flow to the lungs and enlarge or strain the right side of the heart; it may also cause abnormal rhythms or, rarely, allow a clot to pass into an artery and cause a stroke.

### **Atrial Septum**

The wall separating the heart's right and left upper chambers. A hole in this wall is called an atrial septal defect.

### **Atrial Standstill**

Failure of the heart's upper chambers to produce

---

normal electrical signals or contractions. It can cause a very slow heartbeat, dizziness, fainting, poor circulation, or heart failure.

### **Atrial Switch Operation**

An older operation for transposition of the major heart arteries that redirects blood inside the atria using a tunnel-like passage. It restores circulation but leaves the right pumping chamber supporting the body and may cause later rhythm or heart-muscle problems.

#### **Alternate Names:**

Mustard Procedure; Senning Procedure

### **Atrial Tachycardia**

A fast heart rhythm that starts in an area of an upper chamber outside the normal pacemaker. It may cause palpitations, breathlessness, dizziness, chest discomfort, or fainting.

### **Atrial Thrombosis**

Formation of a blood clot inside one of the heart's upper chambers, especially during atrial fibrillation. The clot can break loose and block blood flow to the brain or another organ.

### **Atrioventricular**

Relating to the connection between a heart's upper chamber and the lower chamber on the same side. The term includes the mitral and tricuspid valves and the electrical pathway that coordinates the timing of their contractions.

### **Atrioventricular Block**

Delayed or interrupted electrical signals between the heart's upper and lower chambers. It may cause a slow heartbeat, tiredness, dizziness, fainting, or no symptoms; severe block can require urgent treatment and pacing.

### **Atrioventricular Bundle**

A bundle of specialized electrical fibers that carries signals from the heart's central relay point into the lower chambers. Its branches help both pumping chambers contract in a coordinated way.

#### **Alternate Names:**

Bundle of His

### **Atrioventricular Canal Defect**

A birth defect in which the wall between the heart chambers and the valves between upper and lower chambers do not form normally. It can allow abnormal blood flow and often causes heart failure or poor growth in infancy.

#### **Alternate Names:**

Atrioventricular Septal Defect (AVSD); Endocardial Cushion Defect

### **Atrioventricular Dissociation**

A cardiac rhythm abnormality in which the atria and ventricles depolarize and contract independently under the control of separate pacemakers, occurring when ventricular pacing is faster than atrial rate or during complete atrioventricular block.

#### **Alternate Names:**

AV Dissociation

### **Atrioventricular Nodal Reentrant Tachycardia**

A sudden, regular, fast heartbeat caused by an electrical loop near the heart's central relay point. Episodes may cause pounding heartbeat, chest discomfort, breathlessness, dizziness, or fainting and may stop as suddenly as they begin.

#### **Alternate Names:**

AV Nodal Reentrant Tachycardia; AVNRT

### **Atrioventricular Nodal Reentry Tachycardia**

A rapid, regular heartbeat caused by a looping electrical signal near the atrioventricular node. It commonly begins and ends suddenly and may cause palpitations, dizziness, breathlessness, or fainting.

### **Atrioventricular Node**

A small area of specialized tissue near the junction of the heart's upper and lower chambers that receives electrical signals and briefly delays them before passing them to the lower chambers. This delay allows the upper chambers to empty first.

---

### Atrioventricular Septal Defect

Alternate name; see *Atrioventricular canal defect*.

### Atrioventricular Valves

The two heart valves between the upper and lower chambers: the tricuspid valve on the right and the mitral valve on the left. They open to let blood pass forward and close to prevent backflow.

### Atrium

One of the heart's two upper chambers, which receive returning blood and pass it to the lower pumping chambers. The right atrium receives blood from the body; the left receives oxygen-rich blood from the lungs.

### Atrophic Gastritis

See *Chronic Atrophic Gastritis*.

### Atrophic Glossitis

An inflammatory condition of the tongue characterized by the diffuse atrophy and loss of lingual filiform and fungiform papillae, resulting in a smooth, shiny, erythematous, and beefy red appearance often accompanied by lingual burning, pain, or altered taste. It is a key physical finding in nutritional deficiencies including vitamin B12 (Hunter's glossitis), folate, iron (Plummer-Vinson syndrome), niacin, and riboflavin, as well as systemic conditions such as celiac disease and severe protein-calorie malnutrition.

#### Alternate Names:

Hunter's Glossitis; Bald Tongue; Smooth Tongue; Moeller-Hunter Glossitis

### Atrophic Vaginitis

Alternate name; see *Vaginal Atrophy*.

### Atrophoderma Maculatum

A rare skin condition causing flat, slightly sunken patches in which the skin becomes thinner and changes colour. The patches usually have no open sore or active infection.

### Atrophy

Shrinkage and loss of function of a tissue, organ,

or muscle because its cells become smaller or fewer. Causes include disuse, nerve damage, poor blood supply, malnutrition, hormone deficiency, chronic disease, and ageing; it may be partly reversible.

### Atropine

A medicine that blocks acetylcholine signals at certain nerve endings. It can increase heart rate, reduce secretions, widen the pupils, and counteract selected poisonings or dangerously slow heartbeats.

### Atropine In Organophosphate Poisoning

A medicine used in organophosphate poisoning to block excessive acetylcholine effects, such as wet secretions, narrowed airways, slow heartbeat, vomiting, and diarrhoea. Severe poisoning requires emergency treatment and monitoring.

### Attachment Disorder

See *Reactive Attachment Disorder*.

### Attack Rate

The proportion of people at risk who develop a particular illness during a defined outbreak or exposure period. It is commonly used to describe foodborne outbreaks and other limited events.

### Attention

The ability to select information, focus on it, and maintain that focus while limiting distraction. Attention can be affected by sleep loss, stress, illness, medicines, or brain disorders.

### Attention Deficit Hyperactivity Disorder

Alternate index label; see *Attention-Deficit/Hyperactivity Disorder*.

### Attention-Deficit/Hyperactivity Disorder

A neurodevelopmental disorder involving persistent inattention, hyperactivity, and impulsive behaviour that affects functioning or development. Presentations may be mainly inattentive, mainly hyperactive-impulsive, or combined.

#### Alternate Names:

ADHD; Hyperkinetic Disorder

---

## Attenuated

Weakened or reduced in force, severity, concentration, or biological activity. An attenuated vaccine contains a weakened form of a microorganism that is intended to stimulate immunity without usually causing disease.

## Attenuation Correction

A computer adjustment for the way body tissues weaken signals during medical imaging. It helps make PET, SPECT, and combined scan images more accurate by compensating for signal loss.

## Attributable Risk

The difference in disease risk between people exposed and not exposed to a possible cause. It estimates the extra risk associated with the exposure, but does not by itself prove causation.

## Atypia

An unusual appearance of cells or tissue seen under a microscope. It may result from irritation, inflammation, ageing, a precancerous change, or cancer and must be interpreted with the tissue pattern and clinical context.

## Atypia Morphology

An unusual appearance or arrangement of cells or tissue under a microscope. It may result from irritation or inflammation, or represent a precancerous or cancerous change; its meaning depends on the whole sample and clinical setting.

### Alternate Names:

Cytologic Atypia

## Atypical

Different from the usual appearance, behaviour, structure, or course of a condition. Atypical does not by itself mean that a finding is harmless, precancerous, or cancerous.

## Atypical (Dysplastic) Nevus

A melanocytic mole with unusual clinical or microscopic features such as asymmetry, irregular border, varied color, or larger size. Most are benign,

but multiple atypical nevi increase melanoma risk; a changing, bleeding, or distinctly different lesion should be examined and may require biopsy.

## Atypical Antipsychotics

A group of medicines used mainly for schizophrenia, bipolar disorder, and selected other conditions. They change several brain signals and can cause sleepiness, weight gain, movement problems, high blood sugar, or other metabolic changes.

## Atypical Endometrial Hyperplasia

An overgrowth of abnormal cells in the lining of the uterus. It is not cancer, but it can develop into uterine-lining cancer or occur beside cancer, so medical assessment and follow-up are important.

## Atypical Facial Pain

Persistent or recurring pain in the face that does not fit a clear nerve, dental, sinus, or other recognized cause. Evaluation looks for treatable causes before considering persistent facial-pain disorders.

## Atypical Features

A pattern sometimes used to describe depression in which mood can improve temporarily in response to positive events, with increased sleep, increased appetite or weight, heavy-feeling limbs, and marked sensitivity to rejection.

## Atypical Fibroxanthoma

A low-grade cutaneous sarcoma that usually arises as a rapidly growing red, pink, or flesh-coloured nodule on sun-damaged head or neck skin of an older adult. Diagnosis requires biopsy because it can mimic squamous-cell carcinoma or melanoma; complete excision or Mohs surgery is typical treatment.

## Atypical Genitalia

External genital anatomy that is not typically classified as male or female. It may result from differences in chromosomes, hormones, gonads, or genital development and requires respectful specialist assessment.

---

### Atypical Hemolytic Uremic Syndrome

A rare disease in which inherited or acquired abnormalities of the complement system cause uncontrolled immune activity and damage to small blood vessels, especially in the kidneys. It can destroy red blood cells, lower platelets, and cause sudden kidney injury, high blood pressure, or neurological problems. Unlike typical hemolytic uremic syndrome, it is not primarily caused by a toxin-producing diarrheal infection.

#### Alternate Names:

aHUS; Complement-Mediated HUS

### Atypical Hyperplasia Of The Breast

An overgrowth of unusual cells in a breast duct or lobule that is not cancer. It can increase the future risk of breast cancer and may require closer screening or preventive discussion.

### Atypical Lymphocyte

A lymphocyte, a type of white blood cell, with an unusual size or appearance under a microscope. It often reflects immune activity during a viral infection, but the finding alone does not identify a specific disease.

### Atypical Mole

A mole with unusual clinical or microscopic features, also called a dysplastic nevus. Most are non-cancerous, but multiple atypical moles can increase melanoma risk; a changing, bleeding, or distinctly different mole needs examination.

### Atypical Nevi

Alternate plural name; see *Dysplastic Nevi*.

### Atypical Pneumonia

A lung infection that often develops gradually and may cause more general symptoms than chest symptoms, such as headache, tiredness, sore throat, or muscle aches. Causes include certain bacteria and viruses.

### Atypical Spitz Nevus

An unusual mole made of pigment-producing cells

that resembles a Spitz nevus but has concerning features. It can resemble melanoma under a microscope, so expert skin examination and tissue review may be needed.

### Atypical Teratoid Rhabdoid Tumor

A rare, fast-growing cancer of the brain or spinal cord that occurs mainly in infants and young children. It may cause vomiting, headache, sleepiness, balance problems, seizures, or developmental changes.

#### Alternate Names:

ATRT; Rhabdoid Brain Tumor

### Audiogram

A chart that shows a person's ability to hear at different pitches or frequencies.

### Audiologist

A health professional who assesses hearing and fits hearing aids.

### Audiology

The healthcare field concerned with hearing, balance, and related communication problems. Audiologists test hearing and balance, provide hearing aids and other devices, offer training and support, and identify when medical or surgical care may be needed.

### Audiometry

Measurement of hearing sensitivity at different sound frequencies and intensities, usually using headphones or insert earphones.

### Auditory Acuity

The sensitivity or sharpness of hearing, commonly assessed by determining the softest sounds a person can detect at different frequencies.

### Auditory Area

A region of the cerebral cortex that processes sound information.

#### Alternate Names:

Auditory Cortex

---

## Auditory Fatigue

A temporary decrease in hearing sensitivity or listening ability after prolonged loud sound exposure; rest usually helps, but repeated exposure can cause lasting damage.

## Auditory Hallucinations

False sensory perceptions of sound occurring in the absence of any external acoustic stimulus. They may be non-verbal (such as clicks, ringing, or music) or verbal (such as single words, conversations, running commentary, or command voices). Verbal auditory hallucinations are a hallmark feature of psychotic disorders such as schizophrenia, but can also occur in severe mood disorders with psychotic features, delirium, temporal lobe epilepsy, sensory deprivation, and substance withdrawal.

### Alternate Names:

Paracusia; Phonemes; Auditory Sensory Hallucinations

## Auditory Information Processing Disorder

See *Auditory processing disorder*.

## Auditory Nerve

The portion of the vestibulocochlear nerve that carries hearing signals from the inner ear to the brain.

## Auditory Neuropathy Spectrum Disorder

A form of sensorineural hearing impairment characterized by preserved cochlear outer hair cell function (demonstrated by normal otoacoustic emissions or cochlear microphonics) but abnormal, dyssynchronous neural transmission along the eighth cranial nerve.

### Alternate Names:

ANSD; Auditory Neuropathy

## Auditory Ossicle

One of three small middle-ear bones—the malleus, incus, and stapes—that transmit and amplify vibrations from the eardrum to the inner ear. Disease, injury, or fixation of these bones can cause conductive hearing loss.

### Alternate Names:

Ear Ossicle

## Auditory Pathway

A chain of nerve structures carrying sound signals from the inner ear through the brainstem to the auditory cortex for recognition and interpretation.

## Auditory Perception

The brain's process of detecting, recognizing, and interpreting sounds received through the auditory system.

## Auditory Processing Disorder

Difficulty recognizing or interpreting spoken sounds even though basic hearing sensitivity may be normal. Affected people may struggle to understand speech in noise, follow rapid instructions, or distinguish similar sounds; assessment requires specialized testing and management may include communication strategies and support.

### Alternate Names:

APD

## Auditory Threshold

The quietest sound a person can hear at a particular pitch or frequency. It is measured during a hearing test and helps describe the degree and type of hearing loss in each ear.

### Alternate Names:

Hearing Threshold

## Auer Rods

Needle-shaped structures sometimes seen inside immature blood cells under a microscope. They are made from fused cell granules and strongly suggest a cancer of immature myeloid blood cells, especially acute myeloid leukemia.

## Auerbach Plexus

An extensive intrinsic neural network of unmyelinated nerve fibers and postganglionic parasympathetic ganglia situated in the muscularis externa between the circular and longitudinal smooth muscle layers throughout the gastrointestinal tract, coordi-



---

nating peristalsis and motor activity.

**Alternate Names:**

Myenteric Plexus

**Auger Electrons**

Very low-energy electrons released by some radioactive atoms after a change inside the atom. They travel only a tiny distance and can damage nearby DNA, a property being studied for precisely targeted radiation treatment.

**Augmentation Mammaplasty**

An operation to increase breast size or change breast shape, usually with implants or transferred fat. Possible risks include infection, bleeding, changes in sensation, scarring, and the need for further surgery.

**Augmentation Mammoplasty**

An operation to increase breast size or change breast shape, usually with implants or transferred fat. Possible risks include infection, bleeding, changes in sensation, scarring, and the need for further surgery.

**Alternate Names:**

Breast Augmentation

**Augmentation Of Labor**

Treatment used when labour has started but contractions are not strong or frequent enough for the cervix to continue opening. It may involve oxytocin or deliberately breaking the waters, with monitoring of parent and baby.

**Aura**

A temporary change in vision, sensation, speech, movement, or awareness that may occur before or during a migraine or seizure. Symptoms usually develop gradually and then resolve, but a new or prolonged aura needs urgent assessment.

**Aural Fullness**

A feeling of pressure, blockage, or fullness in the ear. It can result from poor pressure equalization, fluid, earwax, inner-ear disease, or jaw-joint problems and may occur with hearing loss, ringing, or dizziness.

**Alternate Names:**

Ear Pressure; Plugged-Ear Sensation; Blocked-Ear Sensation

**Aural Polyps**

Soft, abnormal tissue growths in the ear canal or middle ear. They often result from long-lasting infection or inflammation, but persistent discharge, pain, or bleeding needs specialist examination to exclude a more serious cause.

**Auricle**

The visible outer part of the ear attached to the side of the head. Its ridges collect and direct sound into the ear canal, helping the brain judge where sounds come from.

**Auricular Hematoma**

A collection of blood between the cartilage and skin covering of the outer ear, usually after a blow or sports injury. Prompt drainage and compression help prevent permanent thickening and deformity of the ear.

**Alternate Names:**

Pinna Hematoma; Subperichondrial Auricular Hematoma

**Auscultation**

Listening to sounds produced inside the body, usually with a stethoscope. Clinicians use it to assess the heart, lungs, abdomen, and blood vessels for findings such as murmurs, wheezing, crackles, or abnormal blood-flow sounds.

**Auscultatory Gap**

A temporary disappearance of the sounds heard while a blood-pressure cuff is slowly deflated, followed by their return. If it is missed, the top blood-pressure number may be recorded falsely low.

**Auspitz Sign**

A skin check for psoriasis where tiny pinpoint bleeding spots appear when the silvery scales of a rash are gently scraped away.

---

### **Austin Flint Murmur**

A low-pitched sound heard during the middle part of the heartbeat over the heart's apex. It is associated with severe leakage of the aortic valve and altered filling of the mitral valve.

### **Autism**

Alternate name; see *Autism Spectrum Disorder*.

### **Autism Spectrum Disorder**

A lifelong neurodevelopmental condition involving persistent differences in social communication and interaction, along with restricted or repetitive behaviors, interests, or sensory patterns. Features begin during the developmental period, vary widely among individuals, and may affect communication, learning, relationships, daily activities, or the type and amount of support needed.

#### **Alternate Names:**

ASD; Autistic Spectrum Disorder; Pervasive Developmental Disorder

### **Autistic Savant**

An autistic person with an unusually strong, specific ability—such as music, art, calculation, or memory—compared with their other abilities. The term describes a pattern of skills and is not itself a medical diagnosis.

### **Autoantibodies**

Immune proteins that mistakenly recognize and attach to the body's own cells, tissues, or proteins. They may cause disease or serve as test clues; their presence alone does not prove that a person has an autoimmune disease.

### **Autoantibody**

An immune protein that mistakenly recognizes and binds a substance from the person's own body. It may damage tissues or interfere with their function, or it may be present only as a laboratory sign without causing disease.

### **Autoantigen**

A substance normally found in the body that is mis-

takenly recognized as foreign by the immune system. This mistaken response can lead to inflammation or damage to the person's own cells or tissues.

### **Autoclave**

A machine that sterilizes equipment by exposing it to pressurized steam at a controlled temperature and time. It destroys bacteria, viruses, fungi, and spores when the correct cycle is used.

### **Autocrine Signaling**

A mode of intercellular and intracellular communication in which a cell synthesizes and secretes a messenger molecule (such as a cytokine, hormone, or growth factor) that binds to surface receptors on the very same cell that produced it, initiating a feedback response.

### **Autoerythrocyte Sensitivity**

A rare condition causing painful bruises or raised purple patches, thought to involve an abnormal reaction to a person's own red-cell components. Other bleeding disorders, medicines, and causes of bruising must be excluded first.

### **Autogenic Drainage**

A controlled-breathing method that moves mucus from the smaller airways toward the throat so it can be coughed out. It uses different breathing depths and requires instruction and practice.

### **Autograft**

Tissue moved from one part of a person's body to another part of the same body. Because the tissue is genetically the person's own, rejection is unlikely, but the original site may have pain, scarring, or other complications.

### **Autoimmune**

Describing an immune response that mistakenly targets the body's own cells, tissues, or proteins. It can cause inflammation, tissue damage, or impaired organ function.

---

## Autoimmune Associated Epilepsy

See *Autoimmune epilepsy*.

## Autoimmune Autonomic Ganglionopathy

A rare, severe autoimmune disorder of the autonomic nervous system characterized by widespread sympathetic and parasympathetic failure. Manifests clinically with severe orthostatic hypotension, anhidrosis, dry mouth and eyes, tonic pupils, gastrointestinal dysmotility (gastroparesis, intestinal pseudo-obstruction), and urinary retention; strongly associated with autoantibodies directed against the ganglionic nicotinic acetylcholine receptor ( $\alpha_3\beta_4$  subtype).

### Alternate Names:

AAG; Subacute Autonomic Neuropathy

## Autoimmune Disease

A disease in which the immune system mistakenly attacks the body's own cells, tissues, or organs. It may cause inflammation, pain, tissue damage, or loss of organ function.

## Autoimmune Diseases

Diseases caused by immune responses against the body's own tissues. They may affect one organ, such as the thyroid, or many parts of the body and can produce inflammation and lasting damage.

## Autoimmune Disorder

A condition in which the immune system mistakenly attacks the body's own cells, tissues, or organs. It may be limited to one organ or affect many organs, causing inflammation and impaired function.

## Autoimmune Encephalitis

Inflammation of the brain caused by an immune attack on nerve cells or their proteins. It may cause seizures, confusion, memory problems, unusual movements, behaviour changes, sleep problems, or reduced consciousness.

## Autoimmune Encephalopathy

See *Autoimmune Encephalitis*.

## Autoimmune Endocrine Disease

An autoimmune disorder that damages a hormone-producing gland or hormone-producing cells. Examples include autoimmune thyroid disease, Addison disease, and type 1 diabetes; it may cause too little or, less often, abnormal hormone production.

## Autoimmune Enteropathy

A rare disease in which the immune system damages the lining of the small intestine, causing ongoing diarrhoea, poor nutrient absorption, weight loss, or growth problems. It may occur with other immune disorders.

## Autoimmune Epilepsy

Recurrent seizures caused by an immune attack on the brain. Seizures may begin suddenly or be difficult to control and can occur with memory problems, confusion, movement changes, or other signs of brain inflammation.

## Autoimmune Gastritis

Long-term inflammation that damages acid- and vitamin-B12-producing cells in the stomach. It can cause low stomach acid, iron deficiency, vitamin B12 deficiency, anaemia, digestive symptoms, and an increased risk of certain stomach growths.

## Autoimmune Hemolytic Anemia

Anaemia caused when immune proteins attach to and destroy red blood cells faster than the body can replace them. It may cause tiredness, pale or yellow skin, dark urine, rapid heartbeat, or shortness of breath.

## Autoimmune Hepatitis

Long-term liver inflammation caused by an immune attack on liver cells. It may cause tiredness, nausea, joint pain, or jaundice, while some people have no symptoms; untreated disease can lead to scarring and liver failure.

## Autoimmune Keratitis

Inflammation of the clear front surface of the eye caused by or associated with immune disease. It may cause eye pain, redness, light sensitivity, blurred

---

vision, or a corneal scar and needs prompt eye assessment.

### **Autoimmune Liver Disease Panel**

A group of blood tests that looks for immune proteins linked with autoimmune liver diseases. Results are interpreted with liver tests, symptoms, examination, and sometimes a liver biopsy; the panel alone cannot establish a diagnosis.

### **Autoimmune Lymphoproliferative Syndrome**

An inherited immune disorder in which lymphocytes do not die off normally after an immune response, causing noncancerous enlargement of lymph nodes and spleen from early childhood. Autoimmune destruction of blood cells may cause anemia, low platelets, or low white-cell counts. It is often due to a defect in the Fas cell-death pathway; diagnosis uses blood counts, specialized immune-cell testing, and gene testing, and there is an increased risk of lymphoma.

### **Autoimmune Pancreatitis**

A rare form of pancreas inflammation caused by an abnormal immune response. It may cause painless jaundice, abdominal pain, weight loss, or a mass that resembles cancer and often improves with immune-suppressing treatment.

### **Autoimmune Polyendocrine Syndromes**

A group of inherited or acquired disorders in which immune attacks affect several hormone-producing glands. Depending on the type, features may include adrenal failure, low parathyroid hormone, thyroid disease, diabetes, chronic fungal infection, or skin changes.

### **Autoimmune Polyendocrinopathy**

A disorder in which immune attacks affect several hormone-producing glands. One inherited form can cause chronic fungal infections of skin or mouth, low parathyroid hormone, adrenal failure, and other autoimmune conditions.

### **Autoimmune Polyglandular Syndrome**

A group of disorders in which the immune system damages two or more hormone-producing glands. Possible effects include adrenal failure, thyroid disease, low parathyroid hormone, diabetes, or problems with reproductive hormones.

### **Autoimmune Thyroid Disease and Pregnancy**

Thyroid autoimmunity can cause hypothyroidism, hyperthyroidism, pregnancy loss, preterm birth, or fetal thyroid effects. Thyroid function and medicine doses are monitored closely before and during pregnancy and after delivery.

### **Autoimmune Thyroiditis**

Immune-mediated inflammation and damage of the thyroid gland. See *Hashimoto Thyroiditis*.

### **Autoimmunity**

An immune response directed against the body's own cells, tissues, or proteins. It may cause no harm, or it may produce inflammation and disease when it damages tissues or interferes with organ function.

### **Autoimmunity: Mechanisms**

Autoimmunity can develop when the immune system fails to remove or control self-reactive cells, or when an infection triggers responses that also recognize body tissues. Genetic factors, immune regulation, and environmental triggers may all contribute.

### **Autoinflammatory Diseases**

Conditions in which the body's early, nonspecific immune response becomes overactive and causes repeated or ongoing inflammation. They may cause fever, rash, joint pain, abdominal pain, or organ damage, often without disease-specific autoantibodies.

### **Autoinoculation**

The spread of an infection from one part of a person's body to another, often by touching or scratching. This can transfer germs to broken skin, the eyes, the mouth, or another vulnerable site.

---

## Autologous

Taken from and returned to the same person. Examples include storing a person's own blood for later transfusion or collecting and returning their own stem cells during treatment.

## Autologous Blood Donation

Collection and storage of a person's own blood before an operation for possible transfusion during or after surgery. It avoids incompatibility with donor blood but may not be suitable or necessary for everyone.

## Autologous Breast Reconstruction

Surgical reconstruction of the breast contour utilizing the patient's own vascularized tissue flaps harvested from donor sites such as the abdomen (e.g., deep inferior epigastric perforator [DIEP] or transverse rectus abdominis myocutaneous [TRAM] flap), back (latissimus dorsi flap), or gluteal/thigh regions.

### Alternate Names:

Autologous Breast Reconstruction; Flap Breast Reconstruction; Autogenous Tissue Reconstruction; DIEP Flap Reconstruction

## Autologous Fat Transplant

A procedure that removes fat from one area of a person's body and injects it into another area to restore volume or change shape. Some of the transferred fat may be reabsorbed, and repeat procedures may be needed.

### Alternate Names:

Autologous Fat Grafting; Fat Transfer; Lipofilling

## Autologous Stem Cell Transplantation

A clinical procedure in which a patient's own hematopoietic stem cells are mobilized, harvested from peripheral blood or bone marrow, cryopreserved, and reinfused following high-dose myeloablative chemotherapy or radiation to rescue bone marrow hematopoiesis.

### Alternate Names:

Autologous Bone Marrow Transplant; Auto-SCT

## Autolysis

The breakdown of cells or tissues by their own enzymes, usually after the cells die. This natural process contributes to tissue decomposition and can affect specimens examined after death.

## Automated External Defibrillator

A portable device that checks the heart rhythm during cardiac arrest and gives an electrical shock when an abnormal rhythm may respond to one. It provides spoken or visual instructions for using it safely.

## Automaticity

The physiological capacity of specialized cardiac pacemaker myocytes (principally in the sinoatrial node, atrioventricular junction, and His-Purkinje system) to undergo spontaneous diastolic phase 4 depolarization and generate action potentials without external neurohumoral stimulation.

## Autonomic Disorder

A problem affecting nerves that control automatic body functions such as blood pressure, heart rate, digestion, sweating, bladder emptying, sexual function, and pupil size. Causes include diabetes, nerve disease, infection, immune disease, and medicines; symptoms depend on the affected functions.

## Autonomic Dysreflexia

A sudden, potentially life-threatening rise in blood pressure in a person with a spinal-cord injury, usually at or above the upper chest. A problem such as a full bladder or bowel below the injury can trigger pounding headache, sweating, flushing, or heart-rate changes.

## Autonomic Ganglion

A small group of nerve-cell bodies outside the brain and spinal cord that passes signals between the autonomic nervous system and organs such as the heart, intestines, glands, and bladder.

## Autonomic Nervous System

The part of the nervous system that controls body functions that usually happen without conscious ef-

---

fort, including heart rate, blood pressure, digestion, sweating, pupil size, bladder activity, and sexual responses.

### Autonomic Neuropathy

Damage to nerves that control automatic functions such as blood pressure, heart rate, digestion, sweating, bladder emptying, and sexual function. Diabetes is common, but other diseases, toxins, and medicines can also cause it.

#### Alternate Names:

Dysautonomia; Autonomic Nerve Dysfunction

### Autophagy

A normal process in which cells break down and recycle worn-out proteins and cell parts. It helps cells survive stress and remove damaged material, but abnormal autophagy may contribute to some cancers and other diseases.

### Autopsy

A medical examination of a body after death. It may include external examination, examination of internal organs, tissue testing, and sometimes toxicology to help determine the cause of death or study disease.

#### Alternate Names:

Post-Mortem Examination; Necropsy; Obduction

### Autoradiography

A laboratory method that uses a radioactive label and a photographic or digital detector to show where a substance is located in a tissue, cell, or biological sample.

### Autosomal Dominant

A pattern of inheritance in which one altered copy of a gene on a non-sex chromosome can cause a condition. A person with one altered copy may pass it to about half of their children.

### Autosomal Dominant Inheritance

Inheritance in which one altered copy of a gene on a non-sex chromosome can cause a condition. Each child of a parent with one altered copy commonly

has a one-in-two chance of inheriting it.

### Autosomal Dominant PKD

An inherited disease in which many fluid-filled cysts gradually develop in the kidneys and may also occur in the liver. Cysts can cause high blood pressure, pain, blood in the urine, infections, and progressive kidney failure.

### Autosomal Dominant Polycystic Kidney Disease

An autosomal dominant inherited disease, usually caused by a change in the *PKD1* or *PKD2* gene, in which many cysts gradually enlarge the kidneys and can reduce kidney function. It may cause high blood pressure, pain, blood in the urine, kidney infections, liver cysts, or rarely brain-artery aneurysms.

#### Alternate Names:

ADPKD

### Autosomal Dominant Tubulointerstitial Kidney Disease

An inherited disease that gradually damages the kidney's small tubules and surrounding tissue. Kidney function declines, often without much protein or blood in the urine; some forms cause early gout, low blood pressure, or anaemia.

### Autosomal Inheritance

The passing of a genetic trait through one of the 22 pairs of chromosomes that are not sex chromosomes. Such traits can affect people of any sex and may follow dominant or recessive patterns.

### Autosomal Recessive

A pattern of inheritance in which a condition usually develops only when both copies of a gene are altered. Parents with one altered copy are often healthy carriers but can pass the condition to a child.

### Autosomal Recessive Disorders

Conditions that usually occur when a person inherits an altered copy of the same gene from each parent. People with one altered copy are often healthy carriers.



---

### Autosomal Recessive Inheritance

Inheritance in which a condition usually requires altered copies of a gene from both parents. When both parents are carriers, each pregnancy commonly has a one-in-four chance of producing an affected child.

### Autosomal Recessive PKD

An inherited disease that usually causes enlarged, poorly functioning kidneys in babies or children and scarring of the liver. Severe disease can interfere with lung development before birth; milder forms may appear later.

### Autosomal Recessive Polycystic Kidney Disease

An inherited disease that usually affects babies or children, causing enlarged kidneys with small cysts and progressive kidney problems. It can also cause liver scarring, high blood pressure, and breathing problems in severe early disease.

#### Alternate Names:

ARPKD

### Autosome

Any chromosome that is not a sex chromosome. Humans usually have 22 pairs of autosomes. Genes on these chromosomes can cause inherited conditions in people of any sex.

#### Alternate Names:

Non-Sex Chromosome

### Avascular Necrosis

See *Osteonecrosis*.

### Avastin

The brand name for bevacizumab, a monoclonal antibody that blocks vascular endothelial growth factor (VEGF). It is used for several cancers and, in ophthalmology, by injection into the eye for selected retinal conditions; risks include high blood pressure, bleeding, clotting, impaired wound healing, and rare gastrointestinal perforation.

### Avian Flu

Alternate name; see *Avian Influenza*.

### Avian Influenza

An infection caused by avian-adapted influenza A virus strains (such as H5N1, H5N6, or H7N9) transmitted to humans primarily via direct contact with infected poultry or contaminated environments. Clinical manifestations range from conjunctivitis and upper respiratory symptoms to fulminant viral pneumonia, acute respiratory distress syndrome (ARDS), multi-organ failure, and death.

#### Alternate Names:

Bird Flu; Avian Flu; Highly Pathogenic Avian Influenza; HPAI

### AVM

Abbreviation for arteriovenous malformation, an abnormal direct connection between arteries and veins that bypasses capillaries. AVMs can bleed, cause seizures or neurologic deficits, or produce symptoms from high-flow effects.

### AVN

Alternate name; see *Avascular Necrosis*.

### Avoidance Behavior

Actions taken to stay away from a feared or distressing situation, thought, place, person, or activity. Avoidance may briefly reduce anxiety but can keep the fear going and interfere with treatment, work, relationships, or daily life.

### Avoidant Personality Disorder

A long-term pattern of avoiding social contact because of strong feelings of inadequacy and fear of criticism, embarrassment, or rejection. It can limit relationships and work; psychological treatment can help.

### Avoidant/Restrictive Food Intake Disorder

An eating disorder characterized by persistent failure to meet appropriate nutritional or energy needs, resulting in significant weight loss, nutritional defi-

---

ciency, dependence on enteral feeding/supplements, or marked interference with psychosocial functioning. Unlike anorexia nervosa, the restriction is driven by sensory sensitivity, lack of interest in eating, or fear of aversive consequences (e.g., choking or vomiting), rather than body-image disturbance.

**Alternate Names:**

ARFID; Selective Eating Disorder

**Avolition**

A marked reduction in motivation or ability to begin and complete purposeful activities. A person may neglect hygiene, household tasks, work, or social contact. It can occur in schizophrenia, severe depression, and other conditions.

**Avulsed Tooth**

A tooth completely knocked out of its socket by an injury. It is a dental emergency; prompt treatment may allow the tooth to be replaced in its socket if it is handled by the crown and kept moist.

**Avulsion**

An injury in which tissue is partly or completely torn away by force. It may involve skin, muscle, a nail, a fingertip, or a tooth and can cause severe bleeding, tissue loss, infection, and loss of function.

**Avulsion Fracture**

A break in which a tendon or ligament pulls off a small piece of attached bone, usually during a sudden movement or injury. Treatment depends on the bone, fragment position, joint involvement, age, and associated damage.

**Avuncular**

Relating to an uncle. In medical family-history or genetic writing, it describes the relationship between a person and their parent's brother or sister's husband.

**Awake Craniotomy**

A specialized neurosurgical procedure in which the patient is deliberately kept awake, alert, and responsive during portions of open brain surgery. Performed primarily to remove tumors (such as gliomas)

or epileptic seizure foci located within or adjacent to critical functional brain areas (the "eloquent cortex," which controls speech, language, motor function, and sensory perception), the operation begins under general anesthesia or conscious sedation while the skull bone flap is temporarily opened. The neurosurgical team then awakens the patient and applies gentle, low-voltage electrical stimulation to the exposed brain surface (cortical mapping) while a speech-language pathologist or neuropsychologist asks the patient to name pictures, read words, or move fingers and toes. If stimulating a particular brain spot temporarily stops speech or movement, surgeons know to spare that exact pathway. This real-time functional mapping maximizes tumor resection while safeguarding essential neurological abilities, reducing post-operative paralysis or loss of speech.

**Alternate Names:**

Awake Brain Surgery; Intraoperative Brain Mapping; Functional Craniotomy

**Awake Fiberoptic Intubation**

A method of placing a breathing tube while the person remains awake and breathes independently. Numbing medicine and a flexible camera are used to guide the tube when a difficult airway is expected.

**Awareness**

Conscious recognition of oneself, surroundings, thoughts, or sensations. Clinicians assess it along with alertness, attention, memory, and orientation when examining brain function.

**Alternate Names:**

Consciousness

**Awareness Under Anesthesia**

Unintended awareness or memory of events during general anaesthesia. A person may hear sounds, feel pressure or pain, or recall part of the operation; although uncommon, it can cause significant distress and needs support.

**Axial Skeleton**

The bones forming the central framework of the body: the skull, spine, ribs, breastbone, and as-

---

sociated small bones of the neck. It supports the body and protects the brain, spinal cord, heart, and lungs.

### **Axial Spondyloarthritis**

Long-term inflammation affecting the joints between the spine and pelvis and other parts of the spine. It commonly causes back pain and stiffness that improve with movement and may eventually limit spinal flexibility.

### **Axilla**

The hollow area beneath the shoulder, commonly called the armpit. It contains major blood vessels, nerves, lymph nodes, and tissues connecting the chest and neck with the arm.

### **Axillary**

Relating to the armpit or to structures within it, including the axillary blood vessels, nerves, and lymph nodes.

#### **Alternate Names:**

Armpit-Related; Pertaining to the Axilla

### **Axillary Artery**

The main artery passing through the armpit and supplying blood to the shoulder, chest wall, and arm. It continues as the brachial artery near the lower border of the armpit.

### **Axillary Dissection**

Surgery that removes selected lymph nodes and nearby tissue from the armpit, often for cancer assessment or treatment. It may cause numbness, shoulder stiffness, fluid collection, nerve injury, or swelling of the arm.

#### **Alternate Names:**

Axillary Node Dissection

### **Axillary Lymph Node**

A lymph node in the armpit that drains lymph from the arm, chest wall, and parts of the breast. Its size or involvement by cancer, infection, or inflammation may be assessed by examination or imaging.

### **Axillary Lymphadenectomy**

An operation to remove lymph nodes from the armpit, usually to assess or treat breast cancer or melanoma. Possible complications include pain, numbness, shoulder stiffness, a fluid collection, nerve injury, and arm swelling.

### **Axillary Lymphadenopathy**

An abnormal enlargement, induration, or matted consistency of the lymph nodes within the axilla. It receives lymphatic drainage from the ipsilateral breast, upper extremity, and chest wall. Major causes include metastatic breast carcinoma, lymphomas, melanoma, acute pyogenic infections of the upper limb, cat-scratch disease (*Bartonella henselae*), tuberculosis, and systemic autoimmune diseases.

#### **Alternate Names:**

Axillary Adenopathy; Enlarged Axillary Nodes

### **Axillary Nerve**

A nerve from the neck that passes around the shoulder and supplies muscles that lift the arm. It also provides feeling over the outer shoulder and may be injured by shoulder dislocation or upper-arm fracture.

### **Axillary Nerve Dysfunction**

Impairment of the axillary nerve, often from shoulder trauma, dislocation, compression, or surgery. It can cause weakness of shoulder abduction, deltoid wasting, and numbness over the lateral shoulder.

### **Axillary Tail Of The Breast**

The extension of breast tissue toward the armpit. It is also called the tail of Spence.

### **Axillary Vein**

A large vein in the armpit that drains blood from the arm and shoulder. It continues beneath the collarbone as the subclavian vein and returns blood toward the heart.

### **Axis**

The second bone in the neck. Its tooth-like projec-

---

tion acts as a pivot for the first neck bone, allowing the head and upper neck to rotate.

### **Axis Vertebra**

The second neck vertebra, whose tooth-like projection allows the first vertebra and head to rotate while protecting the spinal cord.

#### **Alternate Names:**

C2; Axis

### **Axolemma**

The thin outer membrane surrounding a nerve cell's axon. It maintains the electrical difference needed for nerve impulses and contains channels that allow those impulses to travel along the axon.

### **Axon**

The long extension of a nerve cell that carries electrical signals away from the cell body toward another nerve cell, muscle, or gland.

### **Axon Fasciculation**

The process during development in which growing nerve fibres group together to form organized nerve pathways. Chemical guidance signals and cell connections help them reach their correct targets.

### **Axon Guidance**

The process by which growing nerve fibres find and connect with their correct targets during development. Chemical signals and surrounding cells help direct their path.

### **Axon Hillock**

The cone-shaped part where a nerve cell's main body joins its axon, the long fiber that carries signals away. It combines incoming electrical signals and usually starts the nerve impulse when those signals are strong enough.

### **Axon Terminal**

The end of a nerve-cell axon where chemical messengers are released to communicate with another nerve cell, muscle, or gland.

### **Axonal Transport**

The movement of proteins, cell parts, and other materials along a nerve cell's axon between the cell body and the axon terminal. It supports nerve-cell maintenance and communication.

### **Axoneme**

The internal framework of tiny tubes inside a cilium or sperm tail. Together with motor proteins, it produces the bending movements that help move fluid over cells or propel sperm.

### **Axonotmesis**

A moderate degree of peripheral nerve injury in Seddon's classification characterized by disruption of the internal axon and its myelin sheath with subsequent Wallerian degeneration distal to the lesion site, but with anatomical preservation of the surrounding endoneurium, perineurium, and epineurium, allowing spontaneous axonal regeneration.

### **Ayurveda**

A traditional system of health care from India that uses ideas about bodily balance and may include diet, lifestyle practices, massage, and herbal products. Some products can interact with medicines or contain harmful contaminants.

### **Ayurvedic Medicine**

Health care based on traditional Indian theories of bodily balance, using individualized diet, lifestyle practices, physical treatments, and herbal products. Evidence and safety vary, and some products may interact with medicines or contain toxic ingredients.

### **Azole Antifungals**

A group of medicines that stop fungi making ergosterol, a substance needed for strong cell membranes. This weakens or kills the fungus. They can interact with other medicines and may affect the liver or heart rhythm.

### **Azoospermia**

The absence of sperm in semen on laboratory examination. It may result from a blockage that prevents sperm from leaving the reproductive tract or from

---

reduced or absent sperm production in the testes.

### **Azotaemia**

Alternate spelling; see *Azotemia*.

### **Azotemia**

A build-up of nitrogen-containing waste products in the blood, usually because the kidneys are filtering less effectively. It may occur with dehydration, reduced kidney blood flow, or kidney disease.

### **Azygos Vein**

A large vein running along the right side of the spine that drains blood from the chest wall into the superior vena cava. It provides an alternate route for blood when major veins are blocked.

### **Azygous Vein**

Alternate spelling; see *Azygos Vein*.





# B

## B And T Cell Screen

A blood test that measures B cells and T cells, two major groups of immune cells. It may help evaluate suspected immune deficiency, unusual blood-cell counts, or some blood cancers, but results must be interpreted with other tests.

## B And T Cells

B cells make antibodies, while T cells coordinate immune responses or destroy infected and abnormal cells. Both are types of lymphocytes, a group of white blood cells that protect against infection and disease.

## B And T Lymphocytes

Two major types of immune cells. B lymphocytes can make antibodies and immune memory, while T lymphocytes help control immune responses or destroy infected and abnormal cells.

## B Cell

A type of white blood cell that develops in the bone marrow. After activation, it can become an antibody-producing cell or a memory cell that helps the immune system respond faster to a later infection.

### Alternate Names:

B Lymphocyte; Bursa-Equivalent Lymphocyte

## B Hemophilia

Alternate name; see *Hemophilia B*.

## B Lymphocyte

Alternate name; see *B Cell*.

## B Virus

A herpes virus carried mainly by macaques. It can cause a rare but severe infection of the brain and spinal cord in people after a bite, scratch, or contact with infected animal tissue.

## B-Cell Activation

The process in which a B cell receives signals after recognizing a foreign substance, then multiplies and develops into antibody-producing cells or long-lasting memory cells.

## B-Cell Acute Lymphoblastic Leukemia

A rapidly progressive cancer of immature B-lymphocyte precursors in the bone marrow. Abnormal cells crowd out normal blood-forming cells and may enter the blood, lymph nodes, central nervous system, or other organs; diagnosis and treatment require specialist hematology care.

### Alternate Names:

B-Cell ALL; B-Cell Acute Lymphocytic Leukemia; Adult B-Cell Acute Lymphoblastic Leukemia

## B-Cell Acute Lymphoblastic Leukemia for Adults

Acute lymphoblastic leukemia arising from immature B-cell precursors in an adult. It can cause anemia, infection, bleeding, bone pain, and abnormal blood counts and requires urgent hematology care.

## B-Cell Leukemia/Lymphoma Panel

A laboratory test that examines cells from blood, bone marrow, or tissue for abnormal groups of B cells. It uses cell features and marker proteins to

---

help identify or classify some blood cancers.

### **B-Cell Lymphoma**

A group of cancers that begin in B lymphocytes. Different types grow at different speeds and may affect lymph nodes or organs; diagnosis and treatment depend on the exact type, extent, and person's health.

### **B-Cell Receptor**

A protein on the surface of a B cell that binds a particular foreign substance. This binding sends signals that can make the cell multiply, mature, and produce antibodies.

### **B-Lymphocyte**

Alternate spelling; see *B Cell*.

### **B-Lynch Suture**

A surgical stitch placed around the uterus to control severe bleeding after childbirth when the uterus remains relaxed and medicines have not stopped the bleeding. It compresses the uterus while usually preserving the possibility of future pregnancy.

### **B-Mode Ultrasound**

The usual two-dimensional ultrasound image created from echoes of sound waves sent into the body. It shows the size, shape, and structure of organs and soft tissues without using radiation.

### **B-Type Natriuretic Peptide**

A hormone released mainly by the heart when its chambers are stretched or under pressure. A blood test for BNP or the related NT-proBNP can support assessment of heart failure, but age, kidney disease, rhythm, body size, and other illnesses affect the result.

#### **Alternate Names:**

BNP; Brain Natriuretic Peptide; NT-proBNP

### **Ba**

The chemical symbol for barium, a metallic element used in compounds such as barium sulfate. Barium sulfate can be swallowed or given as an enema to

outline parts of the digestive tract on X-rays.

### **Babcock Clamp**

A surgical instrument with smooth, rounded jaws used to hold delicate tubular tissue gently. It may be used on the intestine, fallopian tubes, or other tissues that could be injured by a crushing clamp.

### **Babesia**

A group of parasites transmitted mainly by certain ticks. They infect red blood cells and can cause fever, tiredness, anaemia, low platelets, jaundice, or dark urine; severe illness is more likely without a spleen or with weak immunity.

### **Babesia Divergens**

A species of parasite that infects red blood cells and can cause babesiosis, especially in Europe. It may cause severe destruction of red cells, particularly in people without a spleen or with a weakened immune system.

### **Babesia Microti**

A parasite and the most common cause of babesiosis in North America. It is spread by black-legged ticks, infects red blood cells, and may cause no symptoms or severe anaemia, jaundice, and organ failure.

### **Babesiosis**

An infection caused by Babesia parasites, usually spread by ticks. The parasites enter red blood cells and may cause fever, tiredness, anaemia, low platelets, jaundice, or dark urine; severe disease can affect several organs.

### **Babinski Reflex**

A reflex tested by stroking the sole of the foot. In babies, the big toe normally moves upward and the other toes spread; in older children and adults, the toes usually curl downward. An upward response later in life may indicate a brain or spinal-cord problem.

### **Babinski Response**

Alternate name; see *Babinski Reflex*.

---

## Babinski Sign

Alternate name; see *Babinski Reflex*.

## Baby Acne

Small red or white bumps, usually on an infant's face, caused by temporary effects of hormones. It commonly improves without treatment; gentle washing is sufficient, and adult acne products are not used on infant skin unless a clinician advises.

## Baby Blues

A common short period of tearfulness, irritability, anxiety, or mood changes beginning in the first days after childbirth. It usually improves within two weeks; persistent, severe, or dangerous thoughts may indicate postnatal depression or an emergency.

## Baby Health

An infant's physical growth, development, feeding, sleep, immunizations, screening, and protection from illness or injury. Regular care also monitors hearing, vision, movement, communication, and social development.

## Baby Holding Techniques

Safe ways to support an infant's head, neck, back, and body while carrying, feeding, or moving the baby. Newborns need full head and neck support; never shake an infant or leave one unattended on a raised surface.

## Babyhood

The period of infancy, from birth through the first year of life. During this time, babies undergo rapid growth and development of movement, communication, feeding skills, immunity, and social responses.

## Bacillaceae

A family of rod-shaped bacteria that includes species able to form resistant spores. Some members are harmless or useful, while others can cause infections such as anthrax or food poisoning.

## Bacillary Angiomatosis

A vascular infection caused mainly by *Bartonella*

*henselae* or *Bartonella quintana*, usually in people with weakened immunity. It produces friable red-to-purple papules or nodules and may involve lymph nodes, liver, spleen, bone, or other organs; biopsy and special stains or molecular testing may help distinguish it from Kaposi sarcoma. Antibiotic treatment is usually effective.

## Bacillary Dysentery

An intestinal infection, commonly caused by *Shigella* bacteria, that produces fever, abdominal cramps, and frequent diarrhoea that may contain blood or mucus. Severe dehydration or complications require medical care.

## Bacille Calmette-Guérin

A weakened form of a mycobacterium used as a vaccine against tuberculosis in many countries. It is also placed into the bladder as an immune treatment for some early bladder cancers.

## Bacille Calmette-Guérin Vaccine

A weakened vaccine made from a related type of germ. In many countries it helps protect children from severe tuberculosis. A related treatment is also placed into the bladder for some early bladder cancers.

### Alternate Names:

BCG Vaccine

## Bacilluria

The presence of rod-shaped bacteria in urine, detected by microscopy or culture. It may represent contamination, harmless colonization, or a urinary infection; symptoms, the amount of bacteria, and the organism determine its significance.

## Bacillus

A rod-shaped bacterium. The word can describe bacterial shape or members of the *Bacillus* genus, many of which form resistant spores; some are harmless, while others cause disease.

## Bacillus Anthracis

A spore-forming bacterium that causes anthrax in people and animals. Infection may affect the skin,

---

digestive tract, or lungs and can rapidly become life-threatening, requiring urgent specialist treatment and public-health action.

### **Bacillus Cereus**

A spore-forming bacterium that can cause food poisoning after contaminated food is eaten. It may cause sudden vomiting or watery diarrhoea and, rarely, serious infections in people with weakened immunity.

### **Bacillus Cereus Food Poisoning**

A foodborne illness caused by toxins produced by *Bacillus cereus*, manifesting in two distinct clinical forms: an emetic syndrome characterized by acute nausea and vomiting developing 1 to 5 hours after ingestion (classically linked to fried or reheated rice), and a diarrheal syndrome with abdominal cramps and watery diarrhea developing 8 to 16 hours after ingestion (associated with contaminated meats, vegetables, or sauces).

### **Bacitracin**

An antibiotic applied to the skin that prevents susceptible bacteria from building their protective cell wall. It is used for selected superficial infections, but it can cause skin irritation or allergic reactions.

### **Bacitracin Overdose**

Harm caused by too much exposure to bacitracin. Skin use usually causes local irritation, while swallowing a large amount or receiving excessive injected medicine can cause vomiting, allergic reactions, or kidney injury.

### **Bacitracin Zinc Overdose**

Harm caused by excessive exposure to bacitracin zinc. The effects depend on the amount and route; large exposures may cause vomiting, allergic reactions, or kidney injury and require prompt medical treatment.

### **Back Injury**

Damage to the muscles, ligaments, joints, discs, bones, or nerves of the back after trauma, lifting, or repeated strain. Pain and stiffness are common;

weakness, numbness, loss of bladder or bowel control, or major-trauma pain requires urgent assessment.

### **Back Labor**

Severe back pain during labor, often associated with the baby's position, especially occiput-posterior positioning. It may occur with or without typical abdominal contractions and is managed with position changes, movement, support, analgesia, and standard obstetric assessment.

### **Back Pain**

Pain or discomfort in any part of the back, from the neck to the lower back. It is often caused by muscles, joints, or discs, but may also arise from nerves or internal organs.

#### **Alternate Names:**

Dorsalgia; Backache; Spinal Pain

### **Back Pressure**

Pressure that builds up behind an obstruction or against the normal direction of flow. It may occur in an airway, blood vessel, duct, or hollow organ and can impair movement of air, blood, or fluid.

### **Back Spasms**

Sudden or sustained involuntary tightening of back muscles, causing pain, stiffness, or restricted movement. They may follow strain, awkward movement, spinal or joint problems, or irritation of nearby nerves; weakness, numbness, fever, or loss of bladder or bowel control requires assessment.

### **Back Strains and Sprains**

Injuries involving overstretching or tearing of back muscles, tendons, or ligaments. They commonly cause localized pain, stiffness, and reduced movement after lifting, twisting, overuse, or trauma; neurologic symptoms or major trauma require prompt evaluation.

### **Back-Blows**

Firm blows delivered between the shoulder blades as first aid for severe choking in a responsive person. They are used with age-appropriate choking procedures and do not replace emergency help.

---

## Backache

Pain or discomfort in the back, from the neck to the buttocks. It is a symptom rather than a disease and may result from muscles, joints, discs, nerves, or pain referred from an internal organ.

## Backwash Ileitis

Mild, superficial inflammation of the terminal ileum that occurs in patients with extensive ulcerative colitis (pancolitis). It results from an open, incompetent ileocecal valve that allows inflamed colonic contents and bacteria to reflux backward into the small intestine. Unlike Crohn's disease, the inflammation is continuous with the cecum, is confined to the mucosa without strictures or granulomas, and resolves after surgical removal of the colon. Its presence is associated with severe colitis and an increased long-term risk of colorectal dysplasia.

### Alternate Names:

Reflux Ileitis; Ulcerative Colitis Backwash Ileitis

## Baclofen

A medicine that reduces muscle spasm by lowering nerve signals in the spinal cord. It can cause sleepiness, weakness, dizziness, or confusion; stopping it suddenly after regular use can cause dangerous withdrawal symptoms.

## Bacteraemia

Alternate spelling; see *Bacteremia*.

## Bacteremia

The presence of living bacteria in the bloodstream. It may be temporary, intermittent, or continuous. Some cases cause no illness, while others produce fever and can progress to sepsis or infection of distant organs.

## Bacteria

Microscopic single-celled organisms. Many live harmlessly or help the body, while others cause infections; bacteria can live alone or in communities and may develop resistance to antibiotics.

## Bacterial Adherence

The process by which bacteria attach to body cells or surfaces, often the first step in infection. Surface proteins and tiny hair-like structures help bacteria stay in place and resist removal.

## Bacterial Arthritis

Infection of a joint by bacteria, causing pain, swelling, warmth, restricted movement, and often fever. It is a medical emergency because cartilage can be rapidly damaged and bloodstream infection may develop.

## Bacterial Biofilm

A community of bacteria attached to a surface and enclosed in a protective material they produce. Biofilms can resist immune defenses and antibiotics and cause persistent infections on wounds, catheters, implants, or teeth.

## Bacterial Capsule

A protective outer coating around some bacteria, often made of sugars. It can help bacteria avoid being swallowed by immune cells, survive in the body, and cause disease.

## Bacterial Cell Wall

A strong outer layer that gives bacteria their shape and prevents them from bursting. Many bacterial cell walls contain peptidoglycan, a material made from linked sugars and short protein chains.

## Bacterial Classification

The grouping of bacteria according to their genetic features, appearance, chemistry, and evolutionary relationships. Classification helps identify organisms, understand relatedness, and guide diagnosis, infection control, and treatment.

## Bacterial Conjugation

The transfer of DNA from one bacterium to another through direct contact. The transferred DNA may include genes that help bacteria resist antibiotics or cause disease.

---

## **Bacterial Conjunctivitis**

A contagious bacterial infection of the thin surface covering the eye and inner eyelids. It commonly causes redness, irritation, gritty discomfort, and thick discharge that sticks the eyelids together. Eye pain, light sensitivity, reduced vision, or contact-lens use needs prompt assessment.

## **Bacterial Dermatitis**

Skin inflammation caused by bacterial infection. The skin may become red, warm, swollen, painful, or covered with pus or crusts. It can start in healthy skin or develop when bacteria infect an existing rash or wound.

## **Bacterial Dormancy**

A low-activity state that allows bacteria to survive conditions in which they cannot grow. Some form highly resistant spores; others remain as ordinary cells that temporarily tolerate stress and may become active again when conditions improve.

## **Bacterial Endocarditis**

A serious infection of the heart's inner lining or valves. It may cause fever, chills, tiredness, a new or changing murmur, or weight loss. Bacteria can damage valves or break off and block blood vessels; diagnosis usually requires blood cultures and heart imaging.

## **Bacterial Endocarditis Vegetations**

Clumps of bacteria, clotting material, and platelets that form on a heart valve or the heart's inner lining during bacterial endocarditis. They can develop when bacteria in the blood attach to a damaged or abnormal heart surface.

## **Bacterial Endospores**

Highly resistant resting structures made by some bacteria, including *Bacillus* and *Clostridium*, when nutrients are scarce. They can survive drying, heat, chemicals, and long periods of time, then grow into active bacteria when conditions improve.

## **Bacterial Flagella**

Thread-like structures that help some bacteria move

through liquid or across surfaces. A rotating base works like a motor and turns the attached filament; flagella can also influence how bacteria attach and spread.

## **Bacterial Gastroenteritis**

Acute inflammation of the stomach and intestines caused by pathogenic bacteria such as *Salmonella*, *Campylobacter*, *Shigella*, or pathogenic *Escherichia coli*. It commonly causes diarrhea, abdominal cramps, fever, and sometimes bloody stools or vomiting.

## **Bacterial Genetics**

The study of bacterial genes, DNA, inheritance, and genetic change. Bacteria usually have one main chromosome and may carry extra DNA rings called plasmids that can include antibiotic-resistance genes.

## **Bacterial Growth Phases**

The stages bacteria pass through when growing in a culture. They first adapt, then multiply rapidly, later reach a balance as nutrients fall and waste builds up, and finally decline as conditions become unsuitable.

## **Bacterial Infection**

An illness caused by bacteria entering, multiplying, or spreading in body tissues. Many harmless bacteria normally live on the skin and in the gut, but disease-causing bacteria can invade the body or enter places that are usually free of germs, such as the bloodstream. They can damage tissue, make harmful substances, or trigger inflammation. Examples include strep throat, urinary infections, pneumonia, skin infections, and some food poisoning. Some bacterial infections need antibiotics, but not all do; the correct treatment depends on the infection and the test results.

### **Alternate Names:**

Bacterial Disease; Bacterial Illness; Bacteriosis

## **Bacterial Infections and Pregnancy**

Bacterial infections during pregnancy can affect the urinary tract, vagina, cervix, placenta, membranes, fetus, or newborn. Diagnosis and antibiotic selection



---

account for symptoms, culture results, gestational age, resistance, allergies, and maternal-fetal safety.

### **Bacterial Meningitis**

A serious infection of the membranes around the brain and spinal cord. It can cause fever, severe headache, stiff neck, vomiting, confusion, seizures, or a rash and requires immediate hospital treatment.

### **Bacterial Meningococcal Meningitis**

Meningitis caused by invasive *Neisseria meningitidis*. Fever, severe headache, neck stiffness, confusion, vomiting, or a rapidly spreading purpuric rash requires immediate emergency treatment and public-health contact tracing.

### **Bacterial Metabolism**

The chemical reactions bacteria use to obtain energy, build cell material, grow, and survive. Different bacteria use oxygen, fermentation, light, or other substances, and some of these processes are targets for antibiotics.

### **Bacterial Morphology**

The size, shape, and arrangement of bacteria seen under a microscope. Common forms are spheres, rods, and spirals; bacteria may occur singly, in pairs, chains, clusters, or other patterns.

### **Bacterial Overgrowth**

Small-intestinal bacterial overgrowth is an excessive number of bacteria in the small intestine, where relatively few normally live. It can cause bloating, abdominal pain, diarrhoea, weight loss, or vitamin deficiencies and may result from poor intestinal movement or structural changes.

#### **Alternate Names:**

Small Intestinal Bacterial Overgrowth; SIBO

### **Bacterial Pathogenesis**

The ways bacteria enter the body, attach to cells, invade tissues, avoid immune defenses, and cause illness. Damage may result from direct invasion, toxins, inflammation, or disruption of normal body functions.

### **Bacterial Peritonitis**

A bacterial infection of the lining of the abdominal cavity. It may follow bowel perforation, surgery, another abdominal infection, or infection of abdominal fluid and can cause severe pain, fever, vomiting, and sepsis.

### **Bacterial Pharyngitis**

Bacterial infection and inflammation of the throat, often caused by group A *Streptococcus*. It may cause sudden sore throat, fever, painful swallowing, swollen neck glands, and sometimes white patches on the tonsils.

### **Bacterial Pili**

Tiny hair-like structures on the surface of some bacteria. They help bacteria attach to body cells or surfaces; some also help them move or take up DNA from their surroundings.

### **Bacterial Pneumonia**

A bacterial infection of the lungs that causes inflammation and may fill air spaces with fluid or pus. It can cause fever, cough, phlegm, chest pain, shortness of breath, and low oxygen levels.

### **Bacterial Prostatitis**

A bacterial infection of the prostate gland. It may cause pain in the pelvis or lower back, painful or frequent urination, fever, chills, or difficulty passing urine. Treatment usually requires medical assessment and antibiotics.

#### **Alternate Names:**

Acute Bacterial Prostatitis; Chronic Bacterial Prostatitis; Infectious Prostatitis

### **Bacterial Protein**

A protein produced by a bacterium that may form part of its structure, support its metabolism, help it attach to host cells, or contribute to disease.

### **Bacterial Spore**

A dormant, highly resistant form made by some bacteria when conditions are harsh. Spores can survive drying, heat, and many disinfectants and

---

may later grow into active bacteria when conditions improve.

### **Bacterial Toxemia**

The presence or effects of harmful bacterial substances in the bloodstream. It can occur during severe infection and may cause widespread inflammation, low blood pressure, shock, or organ injury.

### **Bacterial Toxin**

A harmful substance made or released by bacteria. It may kill or injure cells, interfere with body functions, or trigger severe inflammation; toxins may be secreted or released when bacteria break apart.

### **Bacterial Toxins**

Harmful substances produced or released by bacteria. Some are secreted proteins that target particular cells, while others are parts of certain bacterial outer membranes that can trigger widespread inflammation when released.

### **Bacterial Tracheitis**

A bacterial infection and swelling of the windpipe. It can cause fever, a harsh cough, noisy or difficult breathing, and rapid narrowing of the airway, making urgent treatment necessary.

### **Bacterial Transduction**

The transfer of DNA from one bacterium to another by a virus that infects bacteria. The transferred DNA may give the receiving bacterium new traits, including toxin production or antibiotic resistance.

### **Bacterial Transformation**

The uptake of loose DNA from the environment by a bacterium able to absorb it. The new DNA may change the bacterium's features, including how it causes disease or responds to antibiotics.

### **Bacterial Vaginosis**

A common change in the vaginal bacteria in which protective lactobacilli decrease and other bacteria increase. It may cause a thin, grey or white discharge and a fish-like smell, but many people have

no symptoms.

### **Bacterial Virulence Factors**

Features that help bacteria cause disease, such as attaching to cells, entering tissues, avoiding immune defenses, producing toxins, or obtaining nutrients. Examples include surface proteins, protective coatings, enzymes, and toxins.

### **Bactericidal**

Pertaining to an antimicrobial agent or physical process capable of directly killing bacteria, rather than merely inhibiting their cellular growth or replication. Classic bactericidal classes include beta-lactams, aminoglycosides, and fluoroquinolones.

#### **Alternate Names:**

Bactericide

### **Bactericide**

A substance that kills bacteria rather than only stopping them from multiplying. Whether it works depends on the organism, medicine, dose, infection site, and the person's immune defenses.

### **Bacteriocin**

A protein-like substance made by some bacteria that inhibits or kills other susceptible bacteria nearby. It gives the producing bacterium a competitive advantage, but its medical importance depends on the organism and setting.

### **Bacteriology**

The study of bacteria, including their structure, genes, growth, spread, effects on health, and role in disease. It includes laboratory methods used to identify bacteria and test antibiotic sensitivity.

### **Bacteriophage**

A virus that infects bacteria and uses bacterial cells to make more copies of itself. Some phages destroy the bacteria they infect and are being studied for selected antibiotic-resistant infections.

### **Bacteriophage Lambda**

A virus that infects the bacterium *Escherichia coli*.

---

It may multiply and burst the bacterial cell or insert its genetic material into the cell's DNA and remain inactive until later.

### **Bacteriophage T4**

A virus that infects *Escherichia coli*, makes many copies inside the bacterium, and usually breaks the cell open to release them.

### **Bacteriophages**

Viruses that infect bacteria and use them to make more viruses. They may destroy the bacterial cell or insert their genetic material into it and remain inactive for a time.

### **Bacteriorhodopsin**

A light-sensitive protein in some microorganisms. It uses light energy to move charged particles across the cell membrane, helping the organism produce energy.

### **Bacteriostasis**

The stopping or slowing of bacterial growth without necessarily killing the bacteria. The immune system can then remove the remaining organisms; this differs from an agent that directly kills bacteria.

### **Bacteriostatic**

Describes a substance that stops or slows bacteria from multiplying without directly killing them. The body's immune system can then help clear the remaining bacteria.

### **Bacterium**

A single bacterial cell. It usually lacks a nucleus, has genetic material in the cell interior, and may have an outer cell wall; some bacteria are helpful, while others cause infection.

### **Bacteriuria**

The presence of bacteria in urine detected by testing. It may occur with urinary symptoms as part of an infection or without symptoms; the result must be interpreted with the sample quality, amount of bacteria, pregnancy, and other medical factors.

### **Bacteroidaceae**

A family of bacteria that commonly live in the intestine and grow without oxygen. They are usually harmless in the gut but can cause abscesses or other infections if they enter wounds or body areas that should be free of bacteria.

### **Bacteroides**

A group of bacteria that normally live in the intestine and grow without oxygen. They are usually harmless there but can cause abscesses or bloodstream infection if bowel contents enter wounds or other sterile areas.

### **Bacteroides Caccae**

A bacterium that normally lives in the intestine without oxygen. It is usually harmless there but may be found in mixed abdominal or bloodstream infections when intestinal bacteria escape into other tissues.

### **Bacteroides Dorei**

A bacterium that normally lives in the human intestine and grows without oxygen. It is generally considered a harmless member of the gut microbiota.

### **Bacteroides Fragilis**

A bacterium that normally lives in the intestine without oxygen. If bowel contents leak into the abdomen, it is an important cause of abscesses, abdominal infection, bloodstream infection, and sepsis.

### **Bacteroides Gingivalis**

An older name for *Porphyromonas gingivalis*, a bacterium that grows without oxygen in dental plaque. It is associated with inflammation and destruction of the tissues supporting the teeth.

### **Bacteroides Ovatus**

An anaerobic Gram-negative bacterium that is part of the intestinal microbiota and can contribute to opportunistic infection if it enters damaged or normally sterile tissue.

---

### **Bacteroides Thetaiotaomicron**

An anaerobic Gram-negative bacterium that is an important member of the human gut microbiota and helps break down complex dietary carbohydrates; it can cause opportunistic infection outside the intestine.

### **Bacteroides Ureolyticus**

A bacterial name historically applied to an anaerobic, urease-producing organism now generally classified as *Campylobacter ureolyticus*, which can be associated with wound or bloodstream infection.

### **Bacteroides Vulgatus**

An anaerobic Gram-negative bacterium commonly found in the human intestine that can contribute to infection when the intestinal barrier is disrupted.

### **Bad Breath**

An unpleasant odor from the mouth, medically called halitosis. It most often results from bacteria, food debris, gum disease, tooth decay, dry mouth, or tonsil problems; less commonly, it may be related to nose, throat, digestive, or other medical conditions.

#### **Alternate Names:**

Halitosis; Oral Malodor; Fetor Oris

### **BAER**

Alternate name; see *Brainstem Auditory Evoked Response*.

### **Bag-Valve Mask**

A hand-operated resuscitation device, also called a manual resuscitator, consisting of a self-inflating bag, one-way valves, a face mask or airway connector, and often an oxygen reservoir. It delivers positive-pressure ventilation and may be used with a face mask, supraglottic airway, or endotracheal tube.

#### **Alternate Names:**

BVM Manual Resuscitator Bag-Valve-Mask Device

### **Bag-Valve-Mask Ventilation**

A method of giving breathing support with a hand-squeezed bag and face mask. It is used when a

person is not breathing or is not breathing well enough. The rescuer keeps the airway open and gives gentle breaths using room air or added oxygen until normal breathing returns or more advanced airway support is provided.

#### **Alternate Names:**

BVM Ventilation; Bag-Mask Ventilation; Ambu Bag Ventilation; Manual Resuscitator Ventilation

### **Bagassosis**

An allergic lung inflammation caused by breathing dust from mouldy bagasse, the fibrous waste left after sugarcane juice is extracted. It may cause cough, fever, breathlessness, and fatigue after workplace exposure.

### **Bags Under Eyes**

Mild puffiness or loose skin beneath the eyes. It commonly develops with aging, inherited facial features, lack of sleep, allergies, or fluid retention and is usually harmless.

#### **Alternate Names:**

Puffy Eyes

### **Bainbridge Reflex**

An increase in heart rate that can occur when extra blood stretches the upper chambers of the heart. It helps the heart move increased venous blood forward.

### **Bak Protein**

A protein on the outer membrane of mitochondria, the cell's energy-producing structures, that can help start controlled cell death. When activated, it makes openings that release cell-death signals; abnormal control may contribute to cancer.

### **Baker Cyst**

A fluid-filled swelling behind the knee caused by fluid from the knee joint collecting in a small sac. It often occurs with arthritis, a meniscus injury, or another condition causing knee inflammation.

#### **Alternate Names:**

Popliteal Cyst; Baker's Cyst

---

### **Baker's Cyst**

A fluid-filled lump behind the knee, usually caused by excess fluid from an irritated knee joint. It may cause tightness or pain and can occasionally rupture, producing calf swelling that can resemble a blood clot.

### **Baking Powder Overdose**

Illness after swallowing a large amount of baking powder. Depending on its ingredients and amount, it may cause vomiting, abdominal swelling, excess sodium, disturbed body salts, or an abnormal blood-alkali level.

### **Balamuthia**

A genus of free-living amoebas found in soil and water. Some species can rarely infect people, usually causing slowly progressive inflammation of the brain and sometimes skin disease.

### **Balamuthia Mandrillaris**

A free-living amoeba found in soil and water that can rarely infect humans. It may enter through broken skin or the nose and cause severe, gradually worsening infection of the brain or skin.

### **Balamuthia Mandrillaris Infection**

A rare, often severe and sometimes fatal infection caused by the free-living amoeba *Balamuthia mandrillaris*. It may begin as a skin lesion and progress to granulomatous amoebic encephalitis with headache, confusion, seizures, or focal neurologic deficits. Diagnosis is difficult and requires specialized testing; urgent infectious-disease and neurologic evaluation is essential.

### **Balance**

The ability to keep the body steady while sitting, standing, or moving. It depends on information from the inner ear, eyes, muscles, joints, and brain; problems with any of these systems can increase the risk of falls.

### **Balance Assessment**

An evaluation of steadiness during sitting, standing, walking, and changing position. It may include

movement tests and review of vision, medicines, strength, sensation, and dizziness to identify fall risk and guide support.

### **Balance Disorders**

Problems with steadiness or coordinated movement that may cause dizziness, unsteadiness, or falls. Causes include inner-ear disease, vision problems, brain or nerve disorders, muscle weakness, medicines, and blood-pressure changes.

### **Balance Disorders And Vertigo**

Conditions that cause unsteadiness, dizziness, or a false feeling of movement. Causes may involve the inner ear, brain, nerves, vision, muscles, medicines, or blood pressure and require assessment when severe or new.

### **Balance Impairment**

Reduced ability to remain steady while sitting, standing, or walking. It may cause swaying, stumbling, or falls and can result from inner-ear disease, vision problems, weakness, nerve disease, medicines, or brain disorders.

### **Balance Problems**

Difficulty maintaining body stability while standing or moving, often caused by disorders of the inner ear, nervous system, vision, or muscles.

### **Balance Training**

Exercises that improve steadiness and reduce falls by practising safe posture, weight shifting, walking, and responses to movement. A clinician or therapist can adapt exercises to a person's strength, dizziness, and medical conditions.

### **Balanced Study**

A clinical study design in which participant characteristics are evenly distributed between comparison groups at baseline, either by design or by randomization, to reduce confounding and allow fair comparison of interventions.

---

## Balanitis

Inflammation of the glans penis, often involving the prepuce (balanoposthitis). Causes include poor hygiene, irritants, infection, and dermatologic conditions such as lichen planus, psoriasis, and lichen sclerosus. It may cause erythema, swelling, discharge, pruritus, and discomfort; treatment depends on the cause.

## Balanitis Xerotica Obliterans

A long-term inflammatory skin disorder affecting the foreskin and head of the penis. It may cause pale tight skin, itching, pain, painful erections, difficulty passing urine, and narrowing of the opening; it needs medical treatment and follow-up.

### Alternate Names:

BZO; Penile Lichen Sclerosus

## Balanoposthitis

Inflammation of both the glans penis and foreskin. It may cause redness, swelling, itching, pain, discharge, unpleasant odour, or difficulty retracting the foreskin. Causes include irritants, poor moisture control, Candida or bacterial infection, sexually transmitted infections, diabetes, and inflammatory skin disease; recurrent or persistent cases need examination, and inability to return a retracted foreskin over the glans is an emergency.

## Balantidiasis

Balantidiasis is intestinal infection caused by the protozoan *Balantidium coli*, producing diarrhea, abdominal pain, or colitis, especially after fecal exposure.

## Balantidium

A genus of large, single-celled parasites that can infect the large intestine. Infection may cause diarrhoea, abdominal pain, or inflammation of the colon, although many infected people have no symptoms.

## Balantidium Coli

A single-celled parasite that can infect the large intestine, usually after contact with food or water contaminated by animal or human stool. It may cause diarrhoea, abdominal pain, or colitis.

## Balbani Body

A temporary cluster of cell parts and genetic messages near the nucleus of a developing egg cell. It helps organize materials that the early embryo needs after fertilization.

## Baldness

Commonly used term for substantial hair loss, especially androgenetic alopecia, also called pattern hair loss. In this condition, genetically susceptible hair follicles gradually become smaller under the influence of androgens.

### Alternate Names:

Pattern Hair Loss

## Balint's Syndrome

A rare disorder caused by damage to both back portions of the brain. It can cause difficulty seeing more than one object at a time, reaching accurately for objects while looking at them, and directing visual attention.

## Ball-And-Socket Joint

A joint in which the rounded end of one bone fits into a cup-shaped part of another bone. It allows movement in many directions, as in the hip and shoulder.

## Ballance Sign

A physical examination finding characterized by fixed dullness to percussion in the left flank and shifting dullness in the right flank. It indicates a ruptured spleen with a localized left subcapsular or perisplenic hematoma and free blood in the peritoneal cavity.

## Ballard Score

A newborn examination used to estimate how many weeks of pregnancy had passed at birth. It assesses physical features and muscle tone or movement; the estimate is useful when pregnancy dates or early ultrasound information are uncertain.

## Ballism

A movement disorder causing sudden, involuntary,



---

large, flinging movements of an arm or leg. It usually results from damage or dysfunction in deep movement-control areas of the brain.

### **Ballismus**

Alternate name; see *Ballism*.

### **Balloon Aortic Valvuloplasty**

A catheter procedure in which a balloon is inflated across a narrowed aortic valve to improve blood flow from the heart. The benefit may be temporary, and the valve can narrow again; some people later need valve replacement.

### **Balloon Brachytherapy**

A form of internal radiation treatment in which a balloon catheter is placed in a body space, often the cavity left after breast-cancer surgery, and radioactive material is temporarily put inside it to treat nearby tissue.

### **Balloon Catheter**

A thin flexible tube with an inflatable tip. It can widen a narrowed blood vessel or body passage, open a stent, stop bleeding by applying pressure, or help perform another procedure.

### **Balloon Dilation**

A procedure that widens a narrowed blood vessel or body passage by inflating a small balloon at the narrowed area. The result and risks depend on the site and underlying cause.

#### **Alternate Names:**

Balloon Dilatation; Balloon Angioplasty When Performed in a Blood Vessel

### **Balloon Occlusion**

Temporary blockage of a blood vessel or other passage using an inflatable balloon on a catheter. It may control bleeding, test blood flow, or support another procedure; possible complications include reduced blood supply, clotting, and vessel injury.

### **Balloon Tamponade**

Temporary control of bleeding or blockage by inflat-

ing a balloon inside or against a body passage or blood vessel. It applies pressure to the bleeding area but can injure tissue and requires close monitoring.

### **Balneology**

The study of mineral waters, bathing, and spa treatments and their possible health effects. These treatments may ease some symptoms but do not replace proven care, and heat or water exposure can be unsafe for some people.

### **Balneotherapy**

Treatment using mineral or thermal water, baths, muds, or related spa methods. It may relieve symptoms in selected conditions, but evidence varies and heat, infection, falls, or heart strain can make it unsafe for some people.

### **Balo Disease**

A rare inflammatory demyelinating disease of the brain and spinal cord, also called Balo concentric sclerosis. MRI typically shows lesions made of alternating rings of damaged and relatively preserved myelin. It may cause progressive weakness, speech or thinking problems, seizures, or other neurologic deficits and requires specialist assessment.

#### **Alternate Names:**

Balo Concentric Sclerosis

### **Balsam**

A thick, resin-containing substance from certain plants, used in some fragrances, skin products, and traditional remedies. It can irritate the skin or cause allergic contact dermatitis in sensitive people.

### **Bamboo Spine**

An X-ray appearance in advanced ankylosing spondylitis in which new bone bridges the spinal bones, making the spine look like a bamboo stalk. The spine may become stiff and more likely to break after injury.

#### **Alternate Names:**

Poker Spine

---

## Band Cell

An immature neutrophil, a white blood cell that helps fight infection, whose nucleus has a curved band shape. Increased numbers may occur when the bone marrow responds to infection, inflammation, or severe stress.

## Band Heterotopia

A rare birth-related brain-development condition in which a band of extra brain tissue forms below the brain's outer layer. It can cause seizures, learning problems, and developmental delay. Some cases are linked to changes in the *DCX* gene.

### Alternate Names:

Double Cortex Syndrome; Subcortical Band Heterotopia

## Band Keratopathy

A horizontal strip of calcium deposits in the clear front surface of the eye. It may result from long-term inflammation or high calcium levels and can cause irritation, blurred vision, or reduced sight.

## BANF

Alternate name; see *Neurofibromatosis Type 2*.

## Bankart Lesion

A tear in the front-lower rim of the shoulder socket, usually after the upper arm has dislocated. It can make the shoulder unstable and increase the chance of repeated dislocation.

## Bankart Repair

An open or arthroscopic surgical procedure performed to reattach and reconstruct a torn anterior-inferior glenoid labrum and stretched joint capsule (Bankart lesion) to the glenoid rim to restore joint stability following recurrent anterior shoulder dislocations.

## Bannayan-Riley-Ruvalcaba Syndrome

A rare inherited condition that can cause many noncancerous growths, a large head size, intestinal polyps, extra freckles around the penis, and increased risk of certain cancers. It results from

changes in the PTEN gene.

## Banti Syndrome

Older term; specify the cause of portal hypertension and hypersplenism.

### Alternate Names:

Banti Disease; Banti's Syndrome

## Bar Graph

A chart that represents values with rectangular bars whose lengths or heights correspond to the quantities being compared.

## Barakat Syndrome

An inherited disorder that can cause low parathyroid hormone, hearing loss, and kidney abnormalities. These problems may occur together because of genetic changes affecting the development of several organs.

## Barber Itch

An inflammatory infection of hair follicles in the beard area, often caused by bacteria or dermatophyte fungi and associated with itchy bumps or pustules.

### Alternate Names:

Barber's Itch

## Barber Say Syndrome

A rare congenital disorder caused by changes in the TWIST2 gene, characterized by excessive facial hair growth (hypertrichosis), loose, atrophic skin, a wide mouth, distinctive facial features, and eyelid abnormalities such as ectropion.

## Barbiturate

A group of medicines that slow brain activity and can cause relaxation, sleep, anaesthesia, or seizure control. They can suppress breathing, cause dependence, and be fatal in overdose, so use requires careful medical supervision.

## Barbiturate Intoxication And Overdose

Poisoning caused by too much of a barbiturate medicine. It can cause extreme sleepiness, confu-

---

sion, slow or stopped breathing, low blood pressure, coma, or death and requires emergency treatment.

### **Barbiturate Toxicity**

Dose-related central nervous system depression from a barbiturate, ranging from ataxia and slurred speech to coma, respiratory failure, hypotension, hypothermia, and loss of protective reflexes. Management is supportive with airway and cardiovascular support, glucose assessment, and selected decontamination or enhanced elimination for severe long-acting-drug poisoning.

### **Barbiturates**

Medicines that reduce activity in the brain and spinal cord. They may be used for anaesthesia or selected seizure disorders, but their risk of breathing suppression, dependence, and overdose means safer alternatives are often preferred.

### **Bardet-Biedl Syndrome**

A rare inherited disorder that may cause retinal degeneration, obesity, extra fingers or toes, kidney disease, and developmental or reproductive problems.

#### **Alternate Names:**

BBS

### **Bare-Metal Stent**

An expandable metal mesh scaffold without an antiproliferative drug coating, deployed via endovascular catheterization to maintain vascular patency in coronary or peripheral arteries following balloon angioplasty.

#### **Alternate Names:**

BMS; Uncoated Vascular Stent; Bare Stent

### **Bariatric Embolization**

A specialized catheter procedure that reduces blood flow to selected stomach arteries to alter hunger-related signaling and support weight loss. It remains limited or investigational in many settings and may cause bleeding, stomach injury, or other vascular complications.

### **Bariatric Surgery**

Operations that reduce stomach capacity or change digestion to support substantial weight loss and improve obesity-related disease. They require long-term changes in eating, supplements, follow-up, and monitoring for complications.

### **Bariatrics**

The medical field focused on preventing and treating obesity and related conditions. Care may include nutrition, physical activity, psychological support, medicines, and weight-loss surgery.

### **Barium**

A metallic element used medically mainly as barium sulfate, a substance that outlines the digestive tract during certain X-ray examinations. Soluble barium compounds are poisonous and are different from barium sulfate.

### **Barium Enema**

An X-ray examination of the large intestine after barium is put into the rectum. Sometimes air is added to show the bowel lining more clearly. It may be used to examine blockage, diverticula, or changes after surgery; other tests are now preferred for routine colon-cancer screening.

### **Barium Meal**

An X-ray examination in which swallowed barium coats the swallowing tube, stomach, and first part of the small intestine. It can show narrowing, ulcers, abnormal movement, or other structural changes.

### **Barium Study**

An X-ray examination that uses barium to outline parts of the digestive tract. Examples include a barium swallow, barium enema, and upper-GI series. It can show the shape and movement of the digestive tract.

### **Barium Sulfate**

A liquid or paste swallowed or placed in the rectum to outline the digestive tract during X-ray examinations. It is not absorbed normally, but breathing it into the lungs or leakage through a bowel hole can

---

cause serious complications.

### Barium Swallow

An X-ray examination of the swallowing tube after a person drinks barium. It can show narrowing, abnormal shape, or problems with movement. A barium follow-through continues the examination into the small intestine.

### Barium-133

A radioactive form of barium used mainly to check and calibrate radiation detectors and imaging equipment. It is not ordinarily used as a diagnostic scan or treatment for patients.

### Barlow Maneuver

A clinical physical examination test used to detect developmental dysplasia of the hip in infants. The examiner adducts the flexed hip and applies gentle posterior pressure along the shaft of the femur; a palpable clunk or sensation of the femoral head slipping out of the acetabulum indicates hip instability.

#### Alternate Names:

Barlow Test; Barlow Dislocation Test

### Barlow Sign

A check for hip problems in newborns where the doctor gently presses the baby's bent thigh backward; feeling the hip slip out of place shows the joint is unstable or loose.

### Barlow Test

A physical examination maneuver for detecting hip instability in infants. With the hip flexed, gentle pressure is applied to see whether the femoral head can be displaced out of the socket, suggesting developmental dysplasia of the hip.

### Barlow's Disease

Barlow's disease is an older name for severe vitamin-C deficiency, or scurvy, causing bleeding gums, bruising, poor wound healing, and bone or joint symptoms.

### Barognosis

The ability to recognize and compare the weight of objects by holding them. Losing this ability despite normal basic touch sensation may indicate a problem in brain areas that interpret sensory information.

### Baroreceptor

A pressure sensor in the walls of major arteries, especially in the neck and chest. It detects blood-pressure changes and signals the brain to adjust heart rate and blood-vessel narrowing.

### Baroreceptor Reflex

A rapid automatic response that helps keep blood pressure stable. Pressure sensors in major arteries signal the brain to change heart rate and blood-vessel tightness when blood pressure rises or falls.

### Baroreceptors

Stretch-sensitive nerve endings in major arteries that detect changes in blood pressure. They send signals to the brain, which adjusts heart rate and blood-vessel narrowing to help maintain stable pressure.

### Baroreflex

A rapid reflex that helps regulate blood pressure by adjusting heart rate and blood-vessel tone in response to changes in arterial pressure.

### Barotitis Media

Alternate name; see *Airplane Ear*.

### Barotrauma

Injury caused by a sudden or excessive difference in pressure, affecting the lungs, ears, sinuses, or other air-filled spaces. Lung barotrauma can cause a collapsed lung or air escaping into tissues.

### Barr Body

The tightly packed, inactive copy of an X chromosome seen inside the nucleus of many cells that contain more than one X chromosome. It helps balance gene activity between cells with different

---

numbers of X chromosomes.

### **Barraquer-Simons Disease**

A rare acquired disorder in which fat gradually disappears from the face, arms, and upper body while remaining or increasing around the hips and legs. It may be associated with immune disease and kidney complications.

### **Barrel Chest**

A rounded, enlarged appearance of the chest caused by increased lung volume or changes in the chest wall. It may occur with chronic obstructive pulmonary disease, severe asthma, or other long-standing lung conditions.

### **Barrett Esophagus**

A change in the lower lining of the swallowing tube caused by long-term acid reflux. The changed lining can increase the risk of precancerous changes and cancer.

### **Barrett Esophagus Surveillance**

Planned repeat examinations of the lower swallowing tube, usually with an endoscope and tissue samples, to look for precancerous cell changes or early cancer. Timing depends on whether abnormal cell changes are present and their severity.

### **Barrett's Esophagus**

Alternate spelling; see *Barrett Esophagus*.

#### **Alternate Names:**

Columnar Epithelium Lined Lower Esophagus

### **Barrett's Oesophagus**

Alternate spelling; see *Barrett Esophagus*.

### **Barrier Creams**

Skin products that form a protective layer against moisture, friction, irritants, and body fluids. They can help prevent or soothe irritation, but infected, broken, or worsening skin may need medical assessment.

### **Barrier Nursing**

Care practices used to stop infections from spreading between patients, staff, and visitors. They may include hand cleaning, protective clothing, masks, gloves, and keeping a person with a contagious infection separate when needed.

#### **Alternate Names:**

Isolation Nursing; Barrier Precautions; Transmission-Based Precautions

### **Bart's Hemoglobinopathy**

A severe inherited blood disorder caused when all four genes needed to make alpha-globin are missing or inactive. The fetus cannot make enough normal haemoglobin, causing severe anaemia, swelling, and often death before or soon after birth.

### **Barth Syndrome**

A rare inherited disorder of mitochondrial energy production caused by changes in the TAZ gene. It may cause heart-muscle disease, low muscle tone, weakness, poor growth, low neutrophil counts, and recurrent infections.

### **Barthel Index**

A scoring tool that measures how much help a person needs with daily activities such as eating, bathing, dressing, using the toilet, moving between surfaces, walking, and climbing stairs. It is used to track function and rehabilitation progress.

### **Bartholin Abscess**

A painful collection of pus caused by infection of a Bartholin gland near the vaginal opening. It can cause severe swelling and pain when sitting or walking and usually needs drainage; antibiotics may be needed in selected cases.

#### **Alternate Names:**

Infected Bartholin Cyst; Bartholin Gland Abscess

### **Bartholin Cyst Or Abscess**

A blockage of a Bartholin gland duct near the vaginal opening can form a usually painless, fluid-filled cyst; infection can cause a painful abscess. A new or persistent lump, especially in someone aged 40 or older, needs medical assessment because rare gland

---

tumors can look similar.

**Alternate Names:**

Bartholin Cyst; Bartholin Duct Cyst

**Bartholin Gland**

One of two small glands beside the vaginal opening that release lubricating fluid. A blocked duct can cause a cyst, and infection can cause a painful abscess with swelling and difficulty sitting or walking.

**Bartholin Gland Cyst**

See *Bartholin Cyst Or Abscess*.

**Bartholin Gland Hyperplasia**

Enlargement caused by an increase in the cells of a Bartholin gland. It may appear as a lump near the vaginal opening and can resemble a cyst or tumour, so a persistent or unusual mass needs assessment.

**Bartholin Gland Mucinous Cyst**

See *Bartholin Cyst Or Abscess*.

**Bartholin's Cyst**

Alternate spelling; see *Bartholin Cyst Or Abscess*.

**Bartholin's Glands**

Alternate spelling; see *Bartholin Gland*.

**Bartholinitis**

Inflammation or infection of a Bartholin gland near the vaginal opening. It may cause pain, redness, swelling, or an abscess and can make sitting or walking uncomfortable.

**Bartonella**

A group of bacteria usually spread by fleas, lice, ticks, or contact with animals. Different species can cause cat-scratch disease, trench fever, skin lesions, heart-valve infection, or bloodstream infection.

**Bartonella Elizabethae**

A species of Bartonella bacteria that has rarely been linked to bloodstream infection and inflammation of the heart valves, especially in people with underlying health problems.

**Bartonella Henselae**

The bacterium that most commonly causes cat-scratch disease. It is usually spread by a kitten scratch or bite and may cause a small skin bump, swollen nearby lymph nodes, fever, or, rarely, disease of other organs.

**Bartonella Infections**

Infections caused by Bartonella bacteria. They may produce swollen lymph nodes, fever, skin lesions, anaemia, heart-valve infection, or inflammation of organs; symptoms and severity depend on the species and immune status.

**Bartonella Quintana**

A Bartonella species spread mainly by body lice. It causes trench fever, with recurring fever, headache, bone pain, and weakness, and can sometimes cause bloodstream or heart-valve infection.

**Bartonellaceae**

A family of bacteria that includes Bartonella species. Members can cause cat-scratch disease, trench fever, skin and blood-vessel lesions, bloodstream infection, or infection of the heart valves.

**Bartonellosis**

Infection caused by a Bartonella bacterium. Depending on the species, it may cause swollen lymph nodes, recurring fever, skin lesions, anaemia, heart-valve infection, or disease affecting several organs.

**Bartter Disease**

Alternate name; see *Bartter Syndrome*.

**Bartter Syndrome**

An inherited kidney disorder causing excessive loss of salt and potassium in urine. It may lead to dehydration, low blood pressure, muscle cramps or weakness, frequent urination, excessive thirst, and poor growth in children.

**Basal**

Relating to the base of a structure or to a person's resting level of activity, such as resting metabolism



---

or temperature measured after sleep.

### **Basal Body**

A small structure at the base of a cilium or flagellum, such as a cell hair or sperm tail. It anchors the structure to the cell and helps organize the parts that produce movement.

### **Basal Cell**

A cell in the deepest layer of an epithelial tissue, where it can divide and produce new cells. In the epidermis, basal cells help renew the skin and contain keratinocyte stem and progenitor cells.

### **Basal Cell Carcinoma**

The most common skin cancer, arising from basal keratinocytes and associated mainly with cumulative ultraviolet exposure. It may appear as a pearly or translucent papule, a sore that repeatedly bleeds or does not heal, or a scar-like patch; local invasion can destroy nearby tissue, although distant spread is uncommon. Diagnosis requires biopsy, and treatment commonly involves excision, Mohs surgery, or selected nonsurgical or targeted therapies.

### **Basal Cell Skin Cancer**

Alternate name; see *Basal Cell Carcinoma*.

### **Basal Ganglia**

Deep brain structures that help control movement, habits, learning, and emotions. Problems in these structures can cause tremor, stiffness, slowed movement, or unwanted movements.

### **Basal Ganglia Dysfunction**

Abnormal function of deep brain structures that help control movement and learned habits. It may cause tremor, stiffness, slowed movement, twisting movements, or other involuntary movements, as in Parkinson disease or Huntington disease.

### **Basal Joint Arthritis**

Alternate name; see *Thumb Arthritis*.

### **Basal Lamina**

A thin layer beneath many lining cells that anchors them to supporting tissue. It also helps control movement of substances between the lining and the tissue below.

### **Basal Metabolic Rate**

The amount of energy the body uses at complete rest to maintain essential functions such as breathing, circulation, temperature, and cell activity. It varies with body size, age, muscle mass, hormones, illness, and other factors.

### **Basal Nuclei**

Groups of nerve cells deep within the brain that help start, stop, and control movement. They also contribute to learning, habits, motivation, and other thinking processes.

### **Basal Tear**

A tear that is continuously present on the eye's surface. It keeps the cornea moist, smooth, and protected and helps wash away dust and microorganisms.

### **Basal Temperature**

Body temperature measured immediately after waking and before eating, exercise, or other activity. A small sustained rise often occurs after ovulation and can help track menstrual cycles.

### **Base**

A substance that can accept hydrogen ions or reduce acidity. In water, a base usually has a pH above 7; strong bases can burn tissue, while weaker bases are important in body fluids and chemical reactions.

### **Base Excision Repair**

A way cells repair small areas of DNA damage. Enzymes remove the faulty DNA building block, fill the gap with the correct one, and seal the DNA strand.

### **Base Pair**

Two matching building blocks joined inside DNA or

---

RNA. In DNA, adenine pairs with thymine and cytosine pairs with guanine; in RNA, adenine usually pairs with uracil.

### Base Pairing

The matching of complementary DNA or RNA building blocks. This matching helps copy genetic information accurately and allows RNA molecules to fold into useful shapes or attach to other molecules.

### Base Sequence

The exact order of DNA or RNA building blocks in a molecule. It carries genetic information and can be examined to identify inherited changes, mutations, or infectious organisms.

### Basedow's Disease

Older name; see *Graves disease*.

### Baseline

A starting measurement or description of a person's health before treatment, illness, or observation. Later findings are compared with it to identify improvement, worsening, or change.

### Baseline EKG

An electrocardiogram recorded before treatment, a procedure, or a change in health. It provides a starting record for comparison with later heart tracings.

#### Alternate Names:

Baseline ECG; Reference Electrocardiogram

### Basement Membrane

A thin supporting layer beneath many cells that line organs and body surfaces. It anchors the cells, helps control exchange with underlying tissue, and can act as a barrier to tumour spread.

### Basic First Aid

Immediate help given for illness or injury until professional care arrives. It may include making the area safe, calling emergency services, controlling bleeding, helping a choking person, starting resuscitation, and monitoring breathing.

### Basic Life Support

Emergency care for a person who is unresponsive and not breathing normally. It includes calling for help, chest compressions, rescue breaths when appropriate, and use of an automated defibrillator if advised.

### Basic Metabolic Panel

A routine blood test measuring eight vital substances: blood sugar (glucose), calcium, four key electrolytes (sodium, potassium, chloride, and bicarbonate), and two waste products filtered by the kidneys (BUN and creatinine). It provides a rapid overview of fluid and electrolyte balance, kidney function, blood sugar control, and blood acid-base status.

#### Alternate Names:

BMP; Chem-7; Chemistry Screen; Electrolyte and Kidney Panel

### Basic Reproduction Number

The average number of people one infected person would infect in a population where everyone is susceptible, under specified conditions. It helps describe spread but changes with behavior, immunity, treatment, and the environment.

### Basidiomycota

A major fungal group that reproduces by forming spores on club-shaped structures called basidia. It includes mushrooms, rusts, smuts, and some fungi that can cause human disease.

### Basilar

Relating to the base or lowest part of a body structure. For example, a basilar finding may be near the bottom of the skull, lung, or another organ.

### Basilar Artery

The artery formed by the union of the vertebral arteries that supplies the brainstem, cerebellum, and posterior brain.

### Basilar Artery Aneurysm

A weak, widened area in the artery supplying the

---

brainstem and back of the brain. It may press on nearby nerves, cause double vision or weakness, or rupture and produce life-threatening bleeding around the brain.

### **Basilar Artery Migraine**

A variant of migraine with aura originating from brainstem or bilateral occipital cortex dysfunction, presenting with symptoms such as vertigo, bilateral visual disturbances, dysarthria, ataxia, tinnitus, and decreased consciousness followed by headache.

#### **Alternate Names:**

Migraine with Brainstem Aura; Basilar Migraine

### **Basilar Invagination**

A craniocervical junction anomaly in which the margins of the foramen magnum and upper cervical spine (odontoid process) migrate superiorly into the posterior cranial fossa, potentially compressing the brainstem, upper cervical cord, and lower cranial nerves.

#### **Alternate Names:**

Basilar Impression

### **Basilar Membrane**

A flexible membrane in the cochlea that supports the organ of Corti and vibrates in response to sound. Different parts respond best to different sound frequencies.

### **Basilar Skull Fracture**

A traumatic fracture involving the bones forming the base of the skull (ethmoid, sphenoid, temporal, or occipital bones). Clinical hallmarks include cerebrospinal fluid rhinorrhea or otorrhea, hemotympanum, Battle sign, and raccoon eyes.

### **Basilic Vein**

A superficial vein of the upper limb that ascends along the medial side of the arm and joins the brachial veins to form the axillary vein. It is used for venous access in selected settings.

### **Basiliximab**

A monoclonal antibody that blocks the interleukin-

2 receptor alpha chain on activated T cells. It is used as induction immunosuppression, especially in kidney transplantation, and increases susceptibility to infection.

### **Basis Pontis**

The basis pontis is the front portion of the pons containing descending motor fibers and pontine nuclei that relay signals to the cerebellum.

### **Basket Cell**

A basket cell is an inhibitory interneuron that forms synaptic contacts around the cell body or axon initial segment of another neuron.

### **Basophil**

A rare type of white blood cell involved in allergic and inflammatory reactions. It releases substances such as histamine and can increase in number with some allergies, infections, and blood disorders.

### **Basophil Adenoma**

A usually noncancerous tumour of hormone-producing cells in the pituitary gland. It may release too much hormone or press on nearby structures, causing headache, vision problems, or other hormone-related symptoms.

### **Basophilia**

An unusually high number of basophils in the blood. It may occur with allergies, inflammation, infection, an underactive thyroid, or disorders in which the bone marrow makes too many blood cells.

### **Basophilic Focus**

An area in a tissue sample that stains blue or purple with a laboratory dye. The finding describes appearance under the microscope and must be interpreted with the surrounding tissue and other tests.

### **Basophilic Stippling**

The abnormal presence of multiple fine or coarse, dark-blue granules scattered throughout erythrocytes on a Romanowsky-stained peripheral blood smear. The granules represent precipitated riboso-

---

mal ribonucleic acid (RNA) and are characteristically seen in lead poisoning, thalassemias, sideroblastic anemia, and myelodysplastic syndromes.

**Alternate Names:**

Punctate Basophilia

**Basosquamous Carcinoma**

A skin cancer with features of both basal-cell and squamous-cell carcinoma. It usually develops on sun-damaged skin and may behave more aggressively than ordinary basal-cell cancer, so complete treatment and follow-up are important.

**Bassen-Kornzweig Syndrome**

A rare inherited disorder in which the body cannot make certain particles that carry fat in the blood. It causes poor absorption of dietary fat, deficiencies of fat-soluble vitamins, unusual red cells, and progressive nerve or eye problems.

**Bateman Purpura**

Alternate name; see *Actinic Purpura*.

**Bath Salts Abuse And Addiction**

Harmful use of synthetic stimulant drugs sold as “bath salts.” They can cause agitation, paranoia, hallucinations, seizures, dangerous overheating, heart problems, dependence, or death; severe symptoms require emergency care.

**Bather’s Itch**

An itchy rash caused by tiny parasite larvae released by infected snails in fresh or salt water. The larvae cannot mature in people, so the rash usually settles on its own but can be intensely uncomfortable.

**Bathing A Patient In Bed**

Washing a person who cannot safely get out of bed. Care includes keeping water comfortably warm, protecting privacy and skin, preventing falls, checking for pressure injuries, and helping only as much as needed.

**Bathing An Infant**

Washing a baby safely in shallow water while keeping

one hand on the baby at all times. The water is comfortably warm, the room warm, and all supplies are ready before bathing; a baby is never left alone near water.

**Batrachotoxin**

A very powerful poison found in some frogs and birds. It keeps nerve and heart sodium channels open, causing abnormal nerve activity, dangerous heart rhythms, paralysis, and potentially death.

**Batroxobin**

An enzyme derived from snake venom that changes fibrinogen, a blood protein, into fibrin, the material that strengthens a blood clot. It has been studied or used in selected bleeding and clotting settings and can alter bleeding risk.

**Batson Plexus**

A valveless network of epidural and paravertebral veins that connects the deep pelvic and thoracic veins to the internal vertebral venous system. Because of the lack of valves and low intravascular pressures, it provides an alternate route for the retrograde hematogenous spread of pelvic infections and metastases (especially prostate and breast carcinoma) to the spine and central nervous system.

**Alternate Names:**

Batson’s Venous Plexus; Vertebral Venous Plexus

**Batson Venous Plexus**

A valveless network of epidural and vertebral veins connecting the deep pelvic and thoracic veins to the internal vertebral venous plexuses and intracranial venous sinuses. Because it lacks valves, retrograde flow occurs during increases in intra-abdominal or intrathoracic pressure (such as coughing or straining), providing a direct pathway for the hematogenous dissemination of infections and metastases (notably prostate and breast carcinoma) to the vertebral column and brain.

**Alternate Names:**

Batson Plexus; Vertebral Venous Plexus

**Batten Disease**

A group of inherited disorders in which waste mate-

---

rial builds up in nerve cells. They usually begin in childhood and cause worsening seizures, vision loss, movement problems, learning difficulties, and loss of abilities.

### **Battered Child Syndrome**

Older term; see *Child Physical Abuse* or *Abusive Head Trauma*.

### **Batley Bacillus**

An older name for the group of nontuberculous mycobacteria now called *Mycobacterium avium* complex. These organisms can cause lung disease, swollen lymph nodes, or widespread infection, especially when immunity is severely weakened.

### **Battle Sign**

Bruising behind the ear over the bone, which is a warning sign of a break in the base of the skull.

### **Bauhin Valve**

The valve between the last part of the small intestine and the first part of the large intestine. It helps regulate passage of intestinal contents and limits backward movement from the colon.

### **Bauxite-Workers' Lung**

A lung disorder caused by long-term workplace inhalation of bauxite dust. It may produce scarring or inflammation in the lungs and can cause cough, breathlessness, or reduced lung function.

### **Bax Protein**

A protein that helps start programmed cell death when a cell is damaged or no longer needed. Excessive or insufficient Bax activity can affect cell survival and has been studied in cancer and other diseases.

### **Bayes Theorem**

A rule for updating the likelihood of a diagnosis after new evidence. It combines the chance of the disease before the test with how accurately the test result supports or argues against that disease.

### **Bazin Disease**

A chronic, nodular, and often ulcerating form of lobular panniculitis predominantly affecting the posterior lower legs of young or middle-aged women. It represents a delayed-type hypersensitivity reaction associated with *Mycobacterium tuberculosis* or other systemic antigens.

#### **Alternate Names:**

Erythema Induratum of Bazin; Erythema Induratum

### **Bazin's Disease**

An older name for tender, inflamed lumps in the fatty tissue under the skin of the lower legs. It is sometimes linked with tuberculosis but can have other causes and may ulcerate as it heals.

### **Bcgosis**

An infection caused by the weakened mycobacteria used in BCG treatment or vaccination. It may remain local or spread to organs, especially in people with weakened immunity, and can cause fever, cough, or inflammation.

### **Beach Chair Position**

A semi-sitting position used during some operations, especially shoulder surgery. The head, neck, and pressure points must be supported, and blood pressure and brain blood flow require careful monitoring.

### **Beacopp Regimen**

A combination chemotherapy regimen used mainly to treat Hodgkin lymphoma; it includes bleomycin, etoposide, doxorubicin, cyclophosphamide, vincristine, procarbazine, and prednisone, with dosing and supportive care determined by the oncology team.

### **Beals Syndrome**

A rare inherited connective-tissue disorder present from birth. It may cause permanently bent joints, long slender fingers and toes, a curved spine, and ears with a crumpled appearance; severity varies.

---

### Beam Hardening Artifact

A false dark band or streak on a CT image caused when bone or metal removes some parts of the X-ray beam before it reaches the detector. It can obscure nearby body structures.

### Beat Number

The number assigned to a particular heartbeat or rhythmic event when beats are counted or analysed. It can help describe timing, rate, or changes in a recorded heart rhythm.

### Beau Lines

Transverse depressions or grooves across the nail plate caused by temporary interruption of nail-matrix growth. They may follow severe illness, fever, trauma, malnutrition, chemotherapy, or systemic inflammatory disease.

### Beau's Lines

Transverse grooves or depressions across a fingernail or toenail caused by temporary interruption of nail-matrix growth. They may follow severe illness, injury, high fever, nutritional stress, or local trauma; the timing of the groove can help estimate when the event occurred.

### Becaplermin

Becaplermin is a recombinant platelet-derived growth factor gel used in selected diabetic foot ulcers with appropriate wound care.

### Bechterew Disease

An older name for ankylosing spondylitis, a long-term inflammatory disease affecting the joints between the spine and pelvis and often the spine itself. It causes back pain and stiffness that may improve with movement.

#### Alternate Names:

Ankylosing Spondylitis; Marie-Strümpell Disease

### Beck Triad

Three findings that may occur when fluid or blood compresses the heart: low blood pressure, swollen neck veins, and quiet heart sounds. The triad may

be absent, so suspected compression needs urgent examination and imaging.

### Becker Muscular Dystrophy

An X-linked recessive muscle disease caused by reduced or abnormal dystrophin, a protein that supports muscle fibres. It usually causes slowly progressive weakness beginning in childhood or adulthood and may also affect the heart.

### Becker Myotonia

An inherited muscle condition causing stiffness because muscles relax slowly after contraction. It often begins in childhood, may improve after repeated movement, and can cause muscle enlargement, difficulty starting movement, or brief weakness.

#### Alternate Names:

Autosomal Recessive Myotonia Congenita; Recessive Generalized Myotonia

### Becker Nevus

A usually harmless, irregularly shaped patch of darker skin that often becomes noticeable during adolescence and may develop increased hair growth. It commonly affects one shoulder or the upper trunk and is not usually cancerous.

### Becker's Muscular Dystrophy

An inherited muscle disease causing progressive weakness, usually beginning later and advancing more slowly than Duchenne muscular dystrophy. It can also weaken the heart and requires regular heart assessment.

### Becker's Nevus

A usually harmless darkened patch of skin, often with extra hair, that commonly appears on one shoulder or the upper trunk during adolescence. It is not cancer, but changes or uncertainty require assessment.

### Beckwith-Wiedemann Spectrum

An epigenetic overgrowth disorder classically characterized by the clinical triad of macrosomia, macroglossia, and abdominal wall defects (omphalocele or umbilical hernia), accompanied by hemi-



---

hyperplasia, neonatal hypoglycemia secondary to hyperinsulinism, and an elevated predisposition to pediatric embryonal malignancies (notably Wilms tumor and hepatoblastoma).

**Alternate Names:**

BWS; Beckwith-Wiedemann Syndrome

**Beckwith-Wiedemann Syndrome**

An overgrowth condition present from birth that may cause a large body or tongue, abdominal-wall defects, enlarged organs, low blood sugar in newborns, and increased risk of certain childhood cancers. Regular specialist monitoring is needed.

**Beclomethasone**

Beclomethasone is a corticosteroid used by inhalation, nasal spray, or topical routes to reduce inflammation in asthma, rhinitis, or skin disease.

**Becquerel**

The International System (SI) derived unit of radionuclide activity, defined as one nuclear decay or disintegration per second ( $1 \text{ Bq} = 1 \text{ s}^{-1}$ ). Clinical diagnostic and therapeutic radioisotope doses are typically quantified in megabecquerels (MBq) or gigabecquerels (GBq).

**Alternate Names:**

Bq; SI Unit of Radioactivity

**Bed Rest**

Staying in bed or remaining mostly reclined to limit activity. It is used only for selected medical reasons because prolonged inactivity can cause muscle loss, blood clots, constipation, pressure injuries, and reduced independence.

**Bedbug**

A small, flat, blood-feeding insect that hides in bedding, furniture, and cracks. Its bites can cause itchy skin bumps or allergic reactions, but bedbugs are not known to routinely spread human disease.

**Bedbug Bites**

Itchy skin reactions caused by bedbugs feeding at night. Bites may appear as red, swollen, itchy marks

in clusters or lines on exposed skin, sometimes days after exposure; some people have no visible reaction. Bedbugs are not known to spread disease, but scratching can cause secondary infection and rare allergic reactions need medical care.

**Bedbugs**

Small blood-feeding insects that cause itchy bites and skin reactions and can spread through bedding, clothing, and furniture.

**Bednar Aphthae**

Shallow sores on the back of a baby's palate caused by pressure or rubbing, often during feeding. They may make feeding painful and usually improve when the source of irritation is corrected.

**Bedpan**

A container used by a person who cannot safely get out of bed to urinate or pass stool. It is positioned securely, privacy is maintained, and the skin is cleaned and dried afterward.

**Bedsore**

A pressure injury, also called a pressure sore or pressure ulcer, caused by prolonged pressure or pressure combined with shear and friction. It usually develops over a bony area such as the heel, hip, elbow, or tailbone; limited movement, moisture, poor nutrition, and reduced sensation increase risk.

**Bedsore**

Plural of bedsore; see *Bedsore*.

**Alternate Names:**

Pressure Ulcers; Decubitus Ulcers; Pressure Sores; Decubiti

**Bedwetting**

Alternate name; see *Enuresis*.

**Bee And Wasp Sting**

A sting from a bee or wasp that injects venom and usually causes immediate pain, redness, swelling, and itching. Trouble breathing, throat swelling, widespread hives, faintness, or vomiting may indi-

---

cate anaphylaxis and requires emergency treatment.

### **Bee Sting**

A venom-injecting injury from a bee that usually causes local pain, redness, and swelling. A severe allergic reaction can cause widespread hives, breathing difficulty, faintness, or swelling of the mouth and throat.

#### **Alternate Names:**

Apis Envenomation; Honeybee Sting

### **Bee Stings**

Injuries caused by bees injecting venom into the skin. They may cause local pain and swelling, a large reaction extending beyond the sting, or life-threatening anaphylaxis in an allergic person.

### **Beef or Pork Tapeworm Infection**

Intestinal infection with *Taenia saginata* or *Taenia solium* acquired from undercooked beef or pork; *T. solium* eggs can also cause cysticercosis in tissues.

### **Beer-Drinker's Heart**

Historical and informal term; see *Alcoholic Cardiomyopathy*.

### **Beers Criteria**

A regularly updated list of medicines that may cause extra harm in some older adults, such as confusion, falls, or bleeding. It supports medication review but does not replace individualized clinical judgment.

### **Beeswax Poisoning**

Illness after ingesting or inhaling a large amount of beeswax. Beeswax is generally of low toxicity, but large ingestions may cause gastrointestinal blockage or aspiration, especially in children.

### **Behavior**

The observable actions, responses, and activities of an organism, influenced by genetic, neurobiological, psychological, and environmental factors. In medicine, behavioral assessment helps diagnose psychiatric, developmental, and neurological conditions.

### **Behavior Disorder**

A persistent pattern of actions or emotional responses that causes significant distress, interferes with daily life, or harms relationships. Causes may be developmental, psychological, medical, or environmental.

### **Behavior Test**

A structured assessment of behavior, mood, attention, thinking, or social functioning. It may use observation, interviews, questionnaires, or standardized tasks to help identify developmental, psychiatric, or neurologic concerns.

### **Behavior Therapy**

A form of psychotherapy based on learning principles that uses reinforcement, extinction, and other behavioral techniques to modify maladaptive behaviors. It is effective for anxiety disorders, phobias, obsessive-compulsive disorder, and habit disorders.

### **Behavioral Abnormality**

A deviation from typical patterns of behavior that may indicate a psychiatric, neurodevelopmental, or neurologic condition. It is assessed by history, observation, standardized measures, and developmental norms to determine clinical significance.

### **Behavioral Disorder**

A persistent pattern of behavior that causes clinically significant distress or impairment in daily functioning, relationships, learning, or safety. The term is broad; interpretation considers age, development, mental health, medical conditions, environment, trauma, and substance exposure.

### **Behavioral Genetics**

The study of how genes and environmental factors influence behavior and mental traits. Most behaviors reflect many genes interacting with life experiences, not one gene alone.

### **Behavioral Rehabilitation**

Structured therapy that helps a person change behaviors, build coping skills, and regain daily function after illness, injury, addiction, or disability. It

---

may include counseling, goal setting, and practice of useful routines.

### **Behcet Disease**

A long-term inflammatory illness that repeatedly causes painful mouth sores and genital sores. It may also inflame the eyes, skin, joints, blood vessels, brain, or digestive tract and can threaten vision or other organs.

### **Behcet Syndrome**

Another name for Behcet disease, an inflammatory illness that causes repeated mouth and genital sores and may affect the eyes, skin, joints, blood vessels, brain, or digestive tract.

### **Behcet's Syndrome**

Behcet syndrome is a relapsing inflammatory disorder that can cause recurrent mouth and genital ulcers, eye inflammation, skin lesions, and vascular, neurologic, or gastrointestinal disease. Treatment is tailored to the organs involved and may include immunosuppression.

#### **Alternate Names:**

Behçet Disease; Silk Road Disease; Adamantiades-Behçet Syndrome

### **Behr Syndrome**

A rare inherited neurological disorder causing progressive optic-nerve damage, poor vision, muscle stiffness, unsteady movement, and sometimes weakness or bladder problems.

### **Behçet Disease**

Standard spelling of Behcet disease, a relapsing inflammatory disorder that can affect the mouth, genitals, eyes, skin, joints, blood vessels, nervous system, or digestive tract.

### **Behçet's Syndrome**

Behçet's syndrome is a relapsing inflammatory disorder causing recurrent oral and genital ulcers, eye inflammation, skin lesions, and sometimes vascular, neurologic, or gastrointestinal disease.

### **Bejel**

Bejel is endemic treponematoses, a chronic infection caused by *Treponema pallidum pertenue* or related treponemes and spread mainly by close nonsexual contact in endemic areas.

### **Bekam**

A Malay name for cupping, a practice in which cups create suction on the skin; wet cupping may also involve superficial skin incisions.

### **Belching**

The release of swallowed air or gas from the stomach through the mouth. Frequent belching may occur with aerophagia, gastroesophageal reflux, functional dyspepsia, or carbonated-drink consumption.

### **Belinostat**

A cancer medicine that blocks enzymes involved in turning genes on and off. It is used for some peripheral T-cell lymphomas that have returned or resisted earlier treatment.

### **Bell Palsy**

Sudden weakness or paralysis of one side of the face caused by dysfunction of the facial nerve. The eyelid may not close fully, and taste or hearing may change. Most cases improve, but urgent assessment is needed to distinguish it from stroke and to protect the eye.

#### **Alternate Names:**

Bell's Palsy; Idiopathic Facial Palsy; Facial Nerve Palsy

### **Bell's Palsy**

Alternate spelling; see *Bell Palsy*.

### **Bell's Phenomenon**

Upward and outward movement of the eyes when a person closes the eyelids forcefully. This normal protective reflex helps shield the cornea and may become visible when facial-nerve weakness prevents complete eyelid closure.

---

### Bell-Clapper Deformity

A birth-related difference in how a testicle is attached inside the scrotum, allowing it to move and twist more freely than usual. It is often present on both sides and increases the risk of testicular torsion, a sudden emergency that cuts off blood flow.

#### Alternate Names:

Bell-Clapper Anomaly; High Tunica Vaginalis Insertion

### Belladonna

*Atropa belladonna*, the deadly nightshade plant, whose tropane alkaloids such as atropine and scopolamine can cause anticholinergic poisoning.

### Bells Palsy

Alternate spelling without an apostrophe; see *Bell Palsy*.

### Belly

The abdomen, the region of the body between the chest and pelvis containing much of the digestive, urinary, and reproductive organs.

### Below-Knee Amputation

Surgery to remove the lower leg while preserving the knee joint. It may be needed for severe injury, infection, poor blood flow, or cancer; rehabilitation focuses on healing, strength, mobility, and a prosthesis when suitable.

### Bence Jones Protein

An abnormal antibody fragment made by some plasma-cell disorders. It may appear in blood or urine and help identify conditions such as multiple myeloma, but it can also damage the kidneys.

### Bence-Jones Protein

An abnormal antibody fragment made by a single group of plasma cells. Finding it in urine or blood may support evaluation for multiple myeloma or another plasma-cell disorder and possible kidney injury.

### Benchmarking

Comparing healthcare results, safety, costs, or processes with a standard or similar service. The comparison can identify areas for improvement but does not by itself prove that one service causes better outcomes.

### Bendamustine

A cancer medicine that damages tumour-cell DNA and is used for selected blood cancers. It can also damage healthy bone-marrow cells, causing low blood counts, infection, tiredness, nausea, or other side effects.

### Bendamustine Hydrochloride

The hydrochloride form of bendamustine, a chemotherapy medicine used for selected blood cancers. It damages DNA and may cause low blood counts, infection, infusion reactions, nausea, tiredness, or other side effects.

### Bendiocarb

A carbamate insecticide that can overstimulate nerves after significant exposure. Poisoning may cause sweating, salivation, vomiting, diarrhoea, small pupils, muscle twitching, breathing difficulty, or weakness and requires urgent medical treatment.

### Bendopnea

The onset of shortness of breath occurring within thirty seconds of bending forward at the waist (such as when putting on socks or tying shoes). It serves as a specific physical examination sign of heart failure, reflecting an acute increase in ventricular filling pressures and pulmonary capillary wedge pressure superimposed on already elevated baseline hemodynamics.

#### Alternate Names:

Forward-Bending Dyspnea

### Bendroflumethiazide

A diuretic medicine that helps the kidneys remove salt and water. It is used for high blood pressure or fluid retention and may lower potassium or sodium and increase uric acid or blood sugar.

---

## Bends

A form of decompression sickness caused by gas bubbles forming in the body when pressure falls too quickly, usually during or after diving. It commonly causes deep joint or muscle pain but can also injure the brain, spinal cord, lungs, or circulation.

## Benedikt Syndrome

A brainstem vascular stroke syndrome resulting from a paramedian lesion of the midbrain tegmentum involving the oculomotor nerve fascicles, red nucleus, and substantia nigra. It is clinically characterized by ipsilateral third cranial nerve palsy (ptosis, dilated pupil, downward-and-outward eye) and contralateral involuntary choreoathetosis, intention tremor, and ataxia.

### Alternate Names:

Paramedian Midbrain Syndrome

## Benedikt's Syndrome

A rare pattern of neurological problems caused by damage in the midbrain. It may cause a drooping eyelid or double vision on one side and tremor, poor coordination, or involuntary movements on the opposite side.

## Beneficence And Non-Maleficence

Two principles of medical ethics: beneficence means acting to benefit the patient, while non-maleficence means avoiding unnecessary harm. Good decisions balance expected benefits, risks, patient values, and available alternatives.

## Benfotiamine

A fat-soluble form of thiamine, or vitamin B1, sold in some supplements and studied for diabetic nerve damage. Evidence of benefit is limited and it does not replace proven treatment for diabetes or neuropathy.

## Benfuracarb

A carbamate insecticide that can overstimulate nerves after significant exposure. It may cause sweating, salivation, vomiting, diarrhoea, small pupils, muscle twitching, breathing difficulty, or weakness and requires urgent medical treatment.

## Benign

Not cancerous. A benign growth does not invade nearby tissues or spread to other parts of the body, but it can still cause problems because of its size, location, or hormone production.

### Alternate Names:

Nonmalignant; Noncancerous

## Benign Adrenal Tumors

Noncancerous growths in an adrenal gland. Some are found by chance and cause no problems, while others release excess hormones or become large enough to cause pressure symptoms.

## Benign Ameloblastoma

A usually slow-growing tumour arising from cells involved in tooth development, most often in the jaw. It is not cancer but can expand, damage bone, and displace teeth, so treatment usually requires specialist surgery.

## Benign Brain Tumor

A noncancerous growth in or around the brain. Although it does not spread like cancer, it can cause headache, seizures, vision or movement problems, or fluid blockage by pressing on nearby structures.

## Benign Chondroblastoma

A noncancerous cartilage-producing bone tumour that usually develops near the end of a long bone in an adolescent or young adult. It often causes pain, swelling, or reduced movement near a joint.

## Benign CNS Cyst

A noncancerous, fluid-filled sac in or around the brain or spinal cord. Many cause no symptoms, but a large or strategically located cyst can press on nerves or block the normal flow of brain fluid.

## Benign Ear Cyst Or Tumor

A noncancerous lump or fluid-filled sac in the outer, middle, or inner ear. It may cause no symptoms or lead to pain, discharge, hearing loss, dizziness, or pressure depending on its location and size.

---

## Benign Epithelioma

A noncancerous growth arising from cells that line the skin or an organ. It does not invade or spread, but removal or testing may be needed if it bleeds, grows, causes symptoms, or is difficult to distinguish from cancer.

## Benign Esophageal Stricture

A noncancerous narrowing of the esophageal lumen, most commonly resulting from chronic acid reflux injury (peptic stricture), corrosive ingestion, radiation, or prior surgery, causing progressive dysphagia.

### Alternate Names:

Peptic Stricture; Esophageal Stenosis, Benign; Esophageal Stricture

## Benign Fasciculation Syndrome

A neurological condition characterized by persistent, involuntary muscle twitching (fasciculations) in the absence of motor weakness, muscle denervation, sensory neuropathy, or an underlying motor neuron disease such as amyotrophic lateral sclerosis.

## Benign Hemangiopericytoma

Older term; see *Solitary Fibrous Tumor*.

## Benign Lesions of the Ovaries

Noncancerous ovarian growths or cysts, including functional cysts, endometriomas, dermoids, fibromas, and cystadenomas. Evaluation uses symptoms, examination, pregnancy testing, ultrasound, and sometimes tumor markers or surgery to distinguish benign disease from malignancy.

## Benign Liver Tumors

Noncancerous growths in the liver, including hemangiomas, focal nodular hyperplasia, and hepatocellular adenomas. Most are found by chance on ultrasound, CT, or MRI and do not require treatment. Some may cause pain, liver enlargement, bleeding, or rarely change into cancer.

## Benign Melanocytic Lesion

An older or imprecise term for a benign melanocytic growth, such as a melanocytic nevus or mole; unlike

malignant melanoma, it is not cancerous.

## Benign Melanocytic Nevi Pathology

Well-circumscribed proliferations of benign melanocytes arranged in nests or cords at the dermoepidermal junction and/or dermis. Typical lesions show maturation with depth and lack significant cytologic atypia or destructive growth.

## Benign Meningioma

A usually slow-growing, noncancerous tumour arising from the membranes around the brain or spinal cord. It may cause seizures, headache, weakness, or vision problems by pressing on nearby structures.

## Benign Mesenchymoma

A rare noncancerous tumour containing more than one type of tissue such as fat, cartilage, or bone. Its symptoms depend on the site and size, and diagnosis requires examination of tissue.

## Benign Migratory Glossitis

Alternate name; see *Geographic Tongue*.

## Benign Mixed Tumor

A usually noncancerous salivary-gland tumour containing more than one tissue type. It often presents as a slow-growing, painless lump, but long-standing or recurrent growths require assessment because rare malignant change can occur.

## Benign Mullerian Papilloma

A rare noncancerous, wart-like growth in the vulva or vagina formed from Müllerian-type lining cells. It often appears as a small polyp and is diagnosed by examining a tissue sample.

## Benign Myoepithelioma

A noncancerous tumour of myoepithelial cells, often arising in a salivary gland or other glandular tissue. It usually grows slowly and may cause a painless lump; diagnosis depends on tissue examination.

## Benign Neoplasm

A noncancerous growth of cells that remains at its



---

original site and does not spread to distant organs. It may still cause harm by pressing on nearby structures, blocking a passage, bleeding, or producing hormones.

### **Benign Orgasmic Headache**

A sudden severe headache during sexual activity or orgasm after bleeding, stroke, vessel disease, and other dangerous causes have been excluded. A first or unusually severe headache needs urgent medical assessment.

#### **Alternate Names:**

Primary Headache Associated with Sexual Activity; Sexual-Activity Headache

### **Benign Paraganglioma**

A usually slow-growing tumour of specialized nerve-related cells. It may release adrenaline-like hormones or press on nearby nerves and blood vessels; symptoms and treatment depend on its site and activity.

### **Benign Paroxysmal Positional Vertigo**

A common inner-ear disorder causing brief spinning attacks when the head changes position. Tiny calcium particles move into a balance canal and trigger dizziness, often lasting less than a minute.

### **Benign Peripheral Nerve Tumor**

A noncancerous tumour arising from or beside a nerve outside the brain and spinal cord. It may cause a lump, pain, tingling, numbness, weakness, or no symptoms at all.

### **Benign Positional Vertigo**

Alternate name; see *Benign Paroxysmal Positional Vertigo*.

### **Benign Prostatic Hyperplasia**

Noncancerous growth of prostate tissue that can narrow the urine channel. It is not prostate cancer and does not itself turn into cancer. It may cause a weak stream, difficulty starting, urgency, frequent night-time urination, dribbling, or incomplete emptying.

#### **Alternate Names:**

BPH

### **Benign Prostatic Obstruction**

Blockage at the bladder outlet caused by an enlarged prostate or increased muscle tone in the prostate. It may cause a weak or interrupted stream, straining, incomplete emptying, urinary retention, or recurrent infection.

### **Benign Rolandic Epilepsy**

A childhood seizure disorder, now often called self-limited epilepsy with centrotemporal spikes. Brief seizures commonly affect the face or mouth during sleep and may cause twitching, tingling, drooling, or temporary trouble speaking; it usually ends by adolescence.

#### **Alternate Names:**

Self-Limited Epilepsy with Centrotemporal Spikes; SeLECTS; Benign Childhood Epilepsy with Centrotemporal Spikes; Rolandic Epilepsy

### **Benign Skin Lesions**

Noncancerous growths or other localized changes in the skin. Examples include skin tags, epidermoid cysts, dermatofibromas, and seborrheic keratoses. Their appearance and cause vary; some can resemble skin cancer, so a new or changing lesion may need professional assessment.

#### **Alternate Names:**

Benign Cutaneous Lesions; Noncancerous Skin Lesions

### **Benign Tumor**

A noncancerous growth that remains localized and does not invade nearby tissue or spread to distant organs. It may still cause problems by pressing on structures, blocking a passage, bleeding, or making hormones.

### **Benign Urethral Lesions**

Noncancerous growths or structural changes in the urethra, such as polyps, caruncles, papillomas, or inflammatory lesions. They may cause bleeding, discharge, pain, narrowing, or difficulty passing urine, but some cause no symptoms.

---

**Alternate Names:**

Urethral Polyps; Urethral Caruncle; Benign Urethral Tumors

**Benign Vulvar Lesions**

Noncancerous growths or skin disorders of the vulva, including cysts, warts, inflammatory lesions, fibromas, and pigmentary lesions. Persistent, changing, bleeding, painful, or atypical lesions may require biopsy.

**Bennett Fracture**

An intra-articular fracture-subluxation of the base of the first metacarpal bone. The smaller medial articular fragment remains attached to the trapezium by the anterior oblique ligament, while the main metacarpal shaft is displaced proximally, laterally, and dorsally by the pulling force of the abductor pollicis longus tendon.

**Benserazide**

A medicine combined with levodopa in some Parkinson treatments. It blocks levodopa from being changed into dopamine outside the brain, allowing more to reach the brain and reducing some unwanted effects.

**Bent Penis**

Curvature of the penis that may be present from birth or develop later. It can be painless or cause pain, difficulty with erection or intercourse, and sometimes problems passing urine.

**Bent Spine Syndrome**

Severe forward bending of the trunk that improves when lying down. It results from weakness or abnormal contraction of back muscles and can occur with muscle, nerve, or movement disorders.

**Alternate Names:**

Camptocormia

**Bentall Procedure**

Open-heart surgery that replaces a diseased aortic valve, the first part of the aorta, and the aortic root with an artificial valve and tube graft. The coronary arteries are reattached to the graft; lifelong

monitoring and sometimes blood-thinning treatment are needed.

**Alternate Names:**

Bentall Operation; Composite Aortic Root Replacement

**Benzaldehyde**

A chemical with an almond-like smell used in flavourings, fragrances, and manufacturing. Concentrated exposure can irritate the eyes, skin, or lungs; it is not a routine medical treatment.

**Benzazepines**

A family of chemical structures made from joined benzene and seven-membered nitrogen-containing rings. Individual compounds may have different biological or medicinal effects, so the family name alone does not identify a treatment.

**Benzene**

An industrial chemical and solvent found in some fuels and smoke. High short-term exposure can affect the brain and heart; repeated exposure can damage bone marrow and increase the risk of leukaemia.

**Benzene Poisoning**

Illness caused by breathing, swallowing, or absorbing benzene. Short-term exposure may cause dizziness, drowsiness, or irregular heartbeat; repeated exposure can damage bone marrow and increase the risk of leukaemia.

**Benzhexol**

Benzhexol, or trihexyphenidyl, is an anticholinergic drug used mainly to reduce tremor and rigidity in Parkinson disease or drug-induced extrapyramidal symptoms.

**Benzo[a]pyrene**

A polycyclic aromatic hydrocarbon (PAH) formed during the incomplete combustion of organic matter, including tobacco smoke, vehicle exhaust, and grilled meats. Its active metabolite, benzo[a]pyrene-7,8-dihydrodiol-9,10-epoxide (BPDE), forms mutagenic DNA adducts that drive chemical carcinogenesis.

---

**Alternate Names:**

BaP; 3,4-Benzopyrene; Benzo(a)pyrene

**Benzoates**

Salts or chemical forms of benzoic acid used mainly as food preservatives or ingredients in some products. Their effects and safety depend on the specific compound and amount; some people develop irritation or sensitivity.

**Benzocaine**

A local anaesthetic applied to skin or moist body surfaces to reduce sensation. Rarely, especially after excessive use, it can cause methemoglobinaemia, a condition in which blood cannot carry oxygen normally.

**Benzodiazepine**

A medicine that enhances the effect of the inhibitory chemical messenger GABA in the brain. It may reduce anxiety, induce sleep, relax muscles, or control seizures, but can cause sedation, impaired coordination, dependence, and dangerous breathing suppression when combined with opioids or other sedatives.

**Benzodiazepine Toxicity**

Excess benzodiazepine effect causing drowsiness, slurred speech, ataxia, impaired coordination, and sometimes coma or respiratory depression, especially with opioids or alcohol. Treatment is usually airway and supportive care; flumazenil is reserved for carefully selected patients because it can precipitate seizures or withdrawal.

**Benzodiazepine Use Disorder**

A pattern of benzodiazepine use that causes loss of control, harm, or problems at work, home, or in relationships. Regular use can cause dependence, and sudden stopping can produce life-threatening withdrawal.

**Benzodiazepine Withdrawal**

Symptoms after reducing or stopping regular benzodiazepine use. Anxiety, insomnia, shaking, sweating, and sensitivity to light may occur; severe withdrawal can cause seizures, hallucinations, or delirium and

requires supervised tapering.

**Benzopyran**

A chemical ring structure formed by joining benzene and pyran rings. It occurs in some natural substances and medicines; the term describes chemistry rather than one specific disease or treatment.

**Benzopyrene**

A group of chemicals produced by incomplete burning and found in tobacco smoke, vehicle exhaust, and heavily charred food. Some forms can damage DNA and increase cancer risk after the body changes them chemically.

**Benzopyrenes**

Chemicals with a benzopyrene structure that form during incomplete combustion. Several occur in smoke and pollution, can damage DNA, and are linked with increased cancer risk.

**Benzoyl Peroxide**

A topical medicine that kills acne-causing bacteria and helps remove blocked skin cells. It can cause dryness, peeling, irritation, or bleaching of hair and fabrics.

**Benzphetamine**

A sympathomimetic amine and stimulant formerly used in some countries as a short-term adjunct for weight reduction; it can cause dependence, cardiovascular effects, and insomnia and is restricted or unavailable in many jurisdictions.

**Benztropine**

An anticholinergic medicine used mainly for movement problems caused by some medicines and for acute muscle spasms called dystonia. It can cause dry mouth, blurred vision, constipation, urinary retention, confusion, or worsen glaucoma.

**Benzydamine**

A topical anti-inflammatory pain-relieving medicine used for painful inflammation in the mouth, throat, or skin. It may cause temporary numbness, irrita-

---

tion, or an allergic reaction.

### **Benzyl Benzoate**

A topical chemical used to kill mites causing scabies and, in some settings, lice. It can irritate the skin and is applied as directed, with care to avoid the eyes, mouth, and broken skin.

### **Benzylamines**

Chemical compounds containing a benzyl group joined to an amino group. Some are used to make medicines or other chemicals, while their effects depend on the individual compound.

### **Benzylpenicillin**

A penicillin antibiotic used for serious infections caused by susceptible bacteria, including some streptococcal infections, meningitis, and syphilis. Allergic reactions can be severe.

### **BEP Regimen**

A chemotherapy combination of bleomycin, etoposide, and cisplatin, commonly used for some germ-cell cancers. It can cause low blood counts, infection, kidney damage, hearing changes, nerve symptoms, or lung injury and requires monitoring.

### **Berberine**

A plant compound sold in some supplements and studied for blood-sugar, cholesterol, and antimicrobial effects. Evidence varies, and it can cause stomach symptoms, low blood sugar, drug interactions, or harm during pregnancy.

### **Bereavement**

The emotional response to the death of someone important. Grief may include sadness, yearning, anger, numbness, sleep or appetite changes, and waves of distress; it varies widely and can sometimes lead to prolonged grief or depression.

### **Bereavement Reaction**

Emotional and physical responses after someone dies, such as sadness, disbelief, anger, poor sleep, or loss of appetite. These reactions are common and

do not automatically indicate a mental disorder.

### **Berger Disease**

An older name for IgA nephropathy, a kidney disease in which an immune protein builds up in kidney filters. It may cause episodes of blood in the urine after infections, protein in the urine, high blood pressure, and kidney failure.

### **Berger's Disease**

Alternate name; see *IgA Nephropathy*.

### **Beriberi**

Illness caused by severe vitamin B1 deficiency. It may damage nerves and cause weakness, numbness, or difficulty walking, or affect the heart and cause swelling, fast heartbeat, and heart failure; urgent vitamin replacement is needed.

### **Berloque Dermatitis**

A phototoxic skin reaction caused when perfume or cosmetics containing psoralens, classically bergamot oil, are followed by ultraviolet-A exposure. It may produce burning, redness, or blisters and then a pendant- or drip-shaped area of hyperpigmentation on the face or neck.

### **Bernard-Soulier Disease**

A rare inherited bleeding disorder in which platelets are unusually large and do not stick properly to damaged blood vessels. It causes easy bruising, nosebleeds, heavy periods, and prolonged bleeding after injury or procedures.

### **Bernard-Soulier Syndrome**

A rare autosomal recessive bleeding disorder caused by a deficiency or qualitative defect in the platelet glycoprotein Ib-IX-V complex, the primary receptor for von Willebrand factor. Patients present with mucocutaneous hemorrhage, prolonged bleeding time, thrombocytopenia, and abnormally large giant platelets on peripheral blood smear, with impaired platelet aggregation to ristocetin that does not correct with normal plasma.

#### **Alternate Names:**

Giant Platelet Syndrome; Glycoprotein Ib Defi-

---

ciency

### **Bernard–Soulier Syndrome**

A rare inherited bleeding disorder in which platelets are unusually large and cannot attach normally to injured blood vessels. It causes easy bruising, nosebleeds, heavy periods, and prolonged bleeding after injury or procedures.

#### **Alternate Names:**

Giant Platelet Disorder; Platelet-Type Bleeding Disorder 1

### **Berry Aneurysm**

A small, rounded balloon-like bulge at a branching point of an artery at the base of the brain. If it bursts, it can cause life-threatening bleeding around the brain.

### **Berylliosis**

A long-term lung disease caused by sensitivity to inhaled beryllium dust or fumes. It can cause cough, breathlessness, fatigue, and scarring or inflammation that resembles sarcoidosis.

### **Beryllium**

A light metal used in some industrial products. Breathing its dust or fumes can cause allergic lung inflammation, long-term lung disease, and increased cancer risk, especially after repeated workplace exposure.

### **Best Disease**

An inherited macular disorder, usually caused by variants in BEST1, with abnormal accumulation of yellow material at the macula. Vision loss varies and may progress to atrophy, neovascularization, or scarring.

### **Beta 2-Microglobulin**

A small protein released by most cells. Blood or urine levels may rise with inflammation, some blood cancers, or reduced kidney removal; the result is interpreted with symptoms and other tests.

### **Beta Amyloid**

Small protein fragments made from amyloid precursor protein. They can build up into plaques in the brain and are associated with Alzheimer disease, although plaques alone do not explain every symptom.

#### **Alternate Names:**

A $\beta$  Peptide; Amyloid-Beta Protein

### **Beta Carotene**

A yellow-orange plant pigment that the body can convert into vitamin A. It is found in foods such as carrots and leafy vegetables; excessive supplements can turn the skin yellow-orange.

#### **Alternate Names:**

Provitamin A Carotenoid; Carotenoid Provitamin

### **Beta Cell**

A cell in the pancreas that makes and releases insulin when blood glucose rises. Insulin helps body cells use glucose and keeps blood glucose from becoming too high.

#### **Alternate Names:**

Beta Cells; Pancreatic Beta Cell; Pancreatic Beta Cells

### **Beta Globulin**

A group of blood proteins identified together during a protein blood test. They include proteins that carry iron or fats and support immunity; abnormal levels can occur with inflammation, liver disease, or immune disorders.

### **Beta Glucans**

Complex carbohydrates found in oats, barley, yeast, fungi, and some bacteria. Oat and barley forms can modestly lower cholesterol when eaten regularly; effects on immunity depend on the source and preparation.

### **Beta Interferons**

Immune-modifying medicines used to reduce relapses and new brain or spinal-cord lesions in some forms of multiple sclerosis. Common effects include flu-like symptoms and injection-site reactions; mon-

---

itoring may detect liver or blood-count abnormalities.

### **Beta Particle**

A fast-moving electron or positively charged equivalent released by some radioactive atoms. It can damage nearby tissue and is therefore used carefully in selected radiation treatments and laboratory applications.

### **Beta Radiation**

Radiation made of fast electrons or their positively charged counterparts released by unstable atoms. It can injure living tissue but can also be directed at selected tissues for treatment or used in medical testing.

### **Beta Thalassemia**

An inherited blood disorder in which the body makes too little beta-globin, a part of haemoglobin. It can cause mild or severe anaemia, tiredness, jaundice, bone changes, or an enlarged spleen.

### **Beta Tocopherol**

One naturally occurring form of vitamin E. It acts as a fat-soluble antioxidant, but the body generally uses alpha-tocopherol more effectively; its specific clinical importance is still being studied.

### **Beta-Adrenoceptor-Blocking Drugs**

Medicines that block adrenaline-like signals at beta receptors. They can slow the heart, lower blood pressure, reduce tremor, and protect the heart in selected conditions, but may cause fatigue, low pulse, or breathing problems in susceptible people.

### **Beta-Agonist**

A pharmacological agent that stimulates beta-adrenergic receptors. Beta-2 selective agonists (such as albuterol and salmeterol) induce bronchial smooth muscle relaxation and bronchodilation in asthma and chronic obstructive pulmonary disease.

#### **Alternate Names:**

Beta-Adrenergic Agonist

### **Beta-Alanine**

An amino acid that helps muscles make carnosine, which buffers acid during intense activity. Supplements may modestly improve some short, high-intensity exercise but commonly cause temporary tingling of the skin.

### **Beta-Blocker Toxicity**

Overdose causing bradycardia, hypotension, atrioventricular block, cardiogenic shock, hypoglycemia, bronchospasm, seizures, or altered mental status. Propranolol may widen the QRS or cause seizures; severe poisoning may require vasopressors, glucagon, high-dose insulin with glucose, bicarbonate for sodium-channel blockade, and intensive monitoring.

### **Beta-Blockers**

See *Beta Blockers*.

### **Beta-Blockers Overdose**

Poisoning from taking too much of a beta-blocker. It can cause a very slow heartbeat, low blood pressure, wheezing, low blood sugar, confusion, seizures, or shock and requires emergency assessment.

### **Beta-Carotene**

A plant pigment that the body can convert into vitamin A. It occurs in foods such as carrots and leafy greens; high-dose supplements may increase lung-cancer risk in smokers and are not a substitute for a balanced diet.

### **Beta-Carotene Blood Test**

A blood test measuring beta-carotene, a plant pigment that can become vitamin A. A low result may suggest poor nutrition or fat absorption; a high result may cause harmless yellow-orange skin.

### **Beta-Chemokines**

Small immune signals that attract white blood cells to areas of infection or inflammation. They help coordinate defense, but excessive activity can contribute to long-lasting inflammation and tissue damage.



---

### Beta-Endorphin

A natural pain-relieving substance made in the brain and pituitary gland. It acts on opioid receptors and also influences stress responses, mood, and some hormone functions.

### Beta-Glucans

Complex carbohydrates found in foods such as oats, barley, mushrooms, and yeast. Their effects vary by source; some provide fibre and may modestly affect cholesterol, blood glucose, or immune activity.

### Beta-Haemolytic Streptococcus

A streptococcus that completely breaks down red blood cells around its colonies in a laboratory culture. Important group A and group B species can cause throat, skin, bloodstream, newborn, or other serious infections.

### Beta-Human Chorionic Gonadotrophin

The measurable part of the pregnancy hormone hCG. Blood or urine testing helps detect pregnancy and can help assess some pregnancy problems or selected tumours when interpreted with other findings.

### Beta-Lactam Antibiotics

A large group of antibiotics that stop susceptible bacteria building their protective cell wall. They include penicillins, cephalosporins, carbapenems, and related medicines; allergic reactions and bacterial resistance can limit their use.

### Beta-Lactamase

An enzyme made by some bacteria that breaks down certain penicillin-like antibiotics and makes them ineffective. Laboratory testing can identify this resistance, and some medicines block selected beta-lactamases.

### Beta-Lactamase Inhibitor

A compound that binds irreversibly to and inactivates bacterial beta-lactamase enzymes, thereby protecting co-administered beta-lactam antibiotics from enzymatic hydrolysis. Examples include clavulanic acid, sulbactam, tazobactam, and avibactam.

### Beta-Mannosidosis

A rare autosomal recessive disorder in which cells cannot break down certain complex sugars. The stored material can cause developmental delay, hearing loss, repeated infections, seizures, or skeletal and skin problems.

### Beta-Oxidation

Alternate form; see *Fatty Acid Oxidation*.

### Beta-Sheet

A common folded shape in proteins in which extended strands lie beside one another and are held by chemical bonds. Excessive beta-sheet folding can contribute to abnormal protein deposits in some diseases.

### Beta-Thalassemia

An autosomal recessive disorder in which the body makes too little or no beta-globin, a part of haemoglobin. Severity ranges from mild carrier status to severe anaemia requiring regular transfusions and monitoring for iron overload.

### Beta-Thalassemia Intermedia

A moderately severe inherited form of beta-thalassemia causing ongoing anaemia, enlarged spleen, jaundice, or bone changes. People may need occasional transfusions, but usually do not require regular lifelong transfusions.

### Beta-Thalassemia Major

The severe, transfusion-dependent form of beta-thalassemia resulting from homozygous or compound heterozygous mutations that completely abolish beta-globin chain production, leading to profound microcytic anemia, extramedullary hematopoiesis, skeletal deformities, and secondary hemochromatosis.

#### Alternate Names:

Cooley Anemia

### Beta-Thalassemia Minor

The carrier form of beta-thalassemia, caused by one altered beta-globin gene. It usually causes no symp-

---

toms or mild small-cell anaemia and is not treated with iron unless iron deficiency is also confirmed.

**Alternate Names:**

Beta-Thalassemia Trait; Cooley Trait

**Beta-Thalassemia Trait**

The heterozygous carrier state for a beta-globin gene variant. It is usually asymptomatic or causes mild microcytosis with mild or no anemia; hemoglobin electrophoresis often shows increased HbA2. It is also called beta-thalassemia minor.

**Alternate Names:**

Beta-Thalassemia Minor; Thalassemia Minor; Cooley Trait

**Betacoronavirus**

A group of coronaviruses that includes viruses causing respiratory illnesses in people and animals. Individual viruses differ in spread and severity; some cause mild colds, while others can cause severe pneumonia.

**Betaherpesvirinae**

A subgroup of herpes viruses that includes cytomegalovirus and human herpesviruses 6 and 7. They remain in the body after infection and can reactivate, especially when immunity is weakened.

**Betaine**

A natural substance involved in processing homocysteine and helping cells maintain their water balance. It is also sold as a supplement, but high doses can cause side effects or interact with medicines.

**Betamethasone**

A strong corticosteroid medicine that reduces inflammation and immune activity. It may be applied to skin or given in other forms; prolonged or extensive use can thin the skin, worsen infections, or suppress the body's steroid production.

**Betamethasone Valerate**

A corticosteroid applied to the skin to reduce inflammation and itching in conditions such as eczema or psoriasis. Excessive, prolonged, or covered applica-

tion can thin the skin and increase absorption into the body.

**Betaxolol**

A relatively beta-1-selective blocker used orally for some cardiovascular conditions and topically to lower intraocular pressure in glaucoma. It may cause bradycardia, hypotension, or bronchospasm.

**Bethanechol**

Bethanechol is a muscarinic cholinergic agonist that stimulates bladder and gastrointestinal smooth muscle, used selectively for urinary retention or impaired motility.

**Bethesda System For Reporting Thyroid Cytopathology**

A six-tier standardized system for reporting thyroid fine-needle aspiration results, from nondiagnostic and benign categories to follicular-pattern, suspicious, and malignant categories; each tier guides risk assessment and management.

**Bethlem Myopathy**

A usually slowly progressive inherited muscle disease caused by changes in collagen VI. It causes weakness, often beginning in childhood, together with tight joints or shortened muscles, especially in the fingers, elbows, ankles, or spine; many people remain able to walk into adulthood.

**Alternate Names:**

Collagen VI-Related Myopathy (Mild Form); Benign Congenital Myopathy with Contractures

**Bevacizumab**

A monoclonal antibody that blocks vascular endothelial growth factor and inhibits new blood-vessel formation. It is used for selected cancers and some eye diseases; important risks include hypertension, bleeding, thrombosis, proteinuria, impaired wound healing, and gastrointestinal perforation.

**Bexarotene**

An oral retinoid X-receptor agonist used mainly for cutaneous T-cell lymphoma; it commonly causes hypertriglyceridaemia, hypothyroidism, and liver

---

abnormalities and is teratogenic.

### **Bezafibrate**

Bezafibrate is a fibrate lipid-lowering drug that activates PPAR-alpha and lowers triglycerides; muscle and liver effects require monitoring.

### **Bezoar**

A compact mass of material that cannot be digested and becomes trapped in the stomach or intestine. It may contain hair, plant matter, medicines, or other substances and can cause pain, vomiting, blockage, or bleeding.

### **Bezold-Jarisch Reflex**

An inhibitory autonomic cardiovascular reflex characterized by the triad of profound bradycardia, peripheral vasodilation with severe hypotension, and transient apnea. It is triggered by the stimulation of non-myelinated vagal sensory C-fibers in the inferoposterior wall of the left ventricle by mechanical deformation, chemical agents, or acute ischemia during inferior myocardial infarction.

### **BH4 Deficiency**

An inherited deficiency of tetrahydrobiopterin, a cofactor needed to process phenylalanine and make several neurotransmitters. It can cause high phenylalanine, seizures, movement or developmental problems, and neurologic disease; treatment depends on the specific defect.

### **Biacetyl**

Biacetyl, or diacetyl, is a diketone compound with a buttery odor used in flavoring and industry; inhalational exposure can injure the small airways.

### **Bias**

A systematic deviation that distorts a measurement, study result, clinical judgment, or decision away from the truth or from an intended fair comparison.

### **Bibliotherapy**

The use of reading materials as a therapeutic intervention to support mental health, educate patients

about conditions, or promote behavioral change. It may be used alone or alongside other treatments for depression, anxiety, or chronic disease self-management.

### **Bicalutamide**

A non-steroidal androgen-receptor antagonist used in prostate cancer, usually with androgen-deprivation therapy; adverse effects include hot flashes, breast enlargement, sexual dysfunction, and possible liver injury.

### **Bicarbonate**

A substance in the blood that helps keep the body's acid level balanced. The lungs and kidneys regulate it, and a blood bicarbonate result can help identify acid-base or metabolic problems.

### **Bicarbonate Buffer System**

The primary extracellular chemical buffer system in the human body, consisting of carbonic acid and bicarbonate ion in equilibrium with dissolved carbon dioxide, regulated by the lungs (via carbon dioxide excretion) and the kidneys (via bicarbonate reabsorption and excretion).

### **Biceps**

The biceps is a two-headed muscle, especially the biceps brachii of the upper arm, that flexes the elbow and supinates the forearm.

### **Biceps Brachii Muscle**

The two-part muscle on the front of the upper arm. It bends the elbow, turns the palm upward, and helps lift the arm. Injury or tendon rupture can cause weakness, pain, bruising, or a bulging "Pop-eye" appearance.

### **Biceps Reflex**

Bending of the elbow after the biceps tendon is gently tapped. This reflex tests the nerve supplying the biceps and the related spinal-cord pathways, mainly at the fifth and sixth neck levels.

---

## **Bicipital**

Bicipital means relating to a two-headed muscle, especially the biceps tendon, groove, or muscle of the upper arm.

## **Bicipital Groove**

The intertubercular groove of the humerus that transmits the tendon of the long head of the biceps brachii, with the transverse humeral ligament and adjacent rotator-cuff structures contributing to tendon stability.

## **Bicipital Tendinitis**

Irritation and degeneration of the long-head biceps tendon near the shoulder, causing pain in the front of the shoulder that may worsen with lifting, reaching overhead, or resisted elbow flexion.

## **Bicipital Tuberosity**

The radial tuberosity, a roughened prominence on the proximal radius where the biceps brachii tendon inserts and produces flexion and supination of the forearm.

## **Bickerstaff Brainstem Encephalitis**

A rare autoimmune post-infectious central nervous system disorder closely related to the Miller Fisher variant of Guillain-Barré syndrome. It is characterized by the clinical triad of acute progressive ophthalmoplegia, severe ataxia, and central nervous system disturbance (altered consciousness or hyperreflexia), typically accompanied by serum anti-GQ1b IgG autoantibodies.

## **Bicolor**

Having two distinct colours. In medicine, the term may describe a skin lesion, eye, specimen, or imaging finding; its significance depends on the structure and clinical context.

## **Biconcave**

Curved inward on both sides. Normal red blood cells have this disc shape, which makes them flexible and provides a large surface area for carrying oxygen.

## **Biconvex**

Curved outward on both sides. The term may describe the shape of a lens, implant, anatomical structure, or finding on a scan.

## **Bicornuate**

Having two horn-like cavities or projections. A bicornuate uterus is a congenital difference in which the upper uterus forms two partly separated spaces.

## **Bicornuate Uterus**

A uterus with two upper cavities caused by incomplete joining during development before birth. Many people have no symptoms, but it can increase the risk of miscarriage, preterm birth, or abnormal baby position.

## **Bicuspid**

Having two points or flaps. The term describes some teeth, which have two main points, and some heart valves, which have two flaps instead of the usual three.

## **Bicuspid Aortic Valve**

A congenital aortic valve with two leaflets instead of the usual three. It can lead to aortic stenosis, aortic regurgitation, enlargement of the aorta, or other congenital heart abnormalities.

## **Bicuspid Valve**

A heart valve with two leaflets rather than the usual three. The term most often refers to congenital bicuspid aortic valve, which can cause aortic stenosis, regurgitation, infective endocarditis, or ascending-aortic disease.

## **Bid Protein**

A protein that helps connect signals from outside a cell with programmed cell death inside it. When activated, it can help mitochondria release signals that lead to destruction of a damaged or unwanted cell.

## **Bielschowsky Head Tilt Test**

A clinical diagnostic test used to evaluate trochlear

---

nerve (cranial nerve IV) palsy causing superior oblique muscle paresis. Hypertropia and vertical diplopia worsen when the patient tilts the head toward the ipsilateral (paretic) side and improve when tilted toward the contralateral side, as the intact vestibular ocular reflex attempts intorsion.

**Alternate Names:**

Bielschowsky Phenomenon; Head Tilt Test

**Bier Spots**

Small, pale macules surrounded by an erythematous background on the extremities, typically elicited when the limbs are placed in a dependent position. They represent a benign physiological vascular anomaly caused by exaggerated local vasoconstriction in response to increased venous pressure and resolve upon limb elevation.

**Biermer's Anemia**

Another name for pernicious anaemia, in which the body cannot absorb enough vitamin B12 because the stomach does not provide the needed absorption factor. It causes anaemia and can damage nerves if untreated.

**Bietti's Crystalline Dystrophy**

A rare inherited eye disease in which tiny crystals and progressive damage develop in the retina and sometimes the cornea. It may cause difficulty seeing at night, loss of side vision, and gradual loss of central vision.

**Bifascicular Block**

A cardiac intraventricular conduction defect in which two of the three main conduction fascicles below the bundle of His are blocked, characteristically presenting as complete right bundle branch block (RBBB) combined with either left anterior fascicular block (LAFB) or left posterior fascicular block (LPFB).

**Bifocal**

Having two points of focus. Bifocal glasses contain one area for distance vision and another for near vision, allowing both prescriptions in one lens.

**Bifurcation**

The point where one body structure, blood vessel, airway, nerve, or pathway divides into two branches.

**Big Toe Sign**

Upward movement of the big toe, sometimes with spreading of the other toes, when the sole is stroked. In an adult, this may suggest a problem in pathways connecting the brain and spinal cord.

**Bigeminy**

A heart-rhythm pattern in which each normal heart-beat is followed by an early extra beat. The extra beat often begins in the lower heart chambers and may cause palpitations or no symptoms.

**Biglycan**

A protein-and-sugar component of the tissue framework around cells. It helps organize collagen and may influence inflammation, wound healing, and blood-vessel changes.

**Biguanide**

A group of blood-glucose-lowering medicines that reduce glucose production by the liver and improve the body's response to insulin. Kidney function, side effects, and other health conditions guide their use.

**Biguanides**

A group of medicines that lower blood glucose mainly by reducing glucose production in the liver and improving insulin sensitivity. They are used chiefly in type 2 diabetes.

**Alternate Names:**

Biguanide Drugs; Biguanide Antidiabetics

**Bilateral**

Affecting both sides of the body or both members of a paired structure, such as both eyes, ears, knees, lungs, or kidneys.

**Bilateral Acoustic Neurofibromatosis**

Alternate name; see *Neurofibromatosis Type 2*.

---

### **Bilateral Acoustic Neuroma**

Tumours affecting both vestibular nerves, usually representing bilateral vestibular schwannomas associated with NF2-related schwannomatosis. They may cause hearing loss, tinnitus, balance problems, or other cranial-nerve symptoms and require specialist neurologic and genetic assessment.

#### **Alternate Names:**

Bilateral Vestibular Schwannoma; Bilateral Acoustic Neurinoma

### **Bilateral Ankle Swelling**

Alternate form; see *Ankle Swelling*.

### **Bilateral Gynecomastia**

Enlargement of glandular breast tissue on both sides in a male. It may result from normal hormone changes, medicines, obesity, liver or kidney disease, or rarely cancer.

### **Bilateral Hydronephrosis**

Enlargement of the urine-collecting spaces in both kidneys because urine cannot drain normally. Causes include blockage, reflux, stones, or bladder problems; prolonged obstruction can damage kidney function.

### **Bilateral Microphthalmos**

A birth condition in which both eyes are unusually small. Vision may be reduced, and other eye or developmental differences may occur; assessment by an eye specialist is needed.

### **Bilateral Oophorectomy**

Surgery to remove both ovaries, sometimes performed for cancer, severe endometriosis, or high inherited cancer risk. Before natural menopause it causes immediate loss of ovarian hormones and infertility, so long-term effects need discussion.

### **Bilateral Peritonsillar Abscesses**

Collections of pus beside both tonsils, usually after a throat infection. They can cause severe sore throat, difficulty swallowing, trouble opening the mouth, muffled speech, and airway narrowing, requiring

urgent hospital treatment.

### **Bilateral Pneumonia**

Infection or inflammation affecting both lungs. It may cause fever, cough, chest discomfort, breathlessness, or low oxygen; severity depends on the cause, extent, and person's health.

### **Bilateral Polycystic Ovarian Syndrome**

A nonstandard patient-facing name for polycystic ovary syndrome, a hormonal and metabolic disorder characterized by ovulatory dysfunction, androgen excess, and sometimes polycystic ovarian morphology. The syndrome is not defined simply by cysts in both ovaries.

### **Bilateral Tonic-Clonic Seizure**

A seizure causing loss of awareness followed by stiffening and rhythmic jerking of both sides of the body. First aid aims to protect the person from injury without restraining them, and emergency help is needed if the seizure lasts five minutes or more.

### **Bile**

A fluid made by the liver and stored in the gallbladder that helps digest fats. It carries bile salts, cholesterol, pigments, and waste products into the intestine.

### **Bile Acid Sequestrants For Cholesterol**

Medicines that trap bile acids in the intestine so they leave in stool. The liver then uses cholesterol to make more bile acids, lowering LDL cholesterol; constipation and medicine interactions are common concerns.

### **Bile Acids**

Substances made by the liver and released into the intestine to help digest and absorb fats and vitamins A, D, E, and K. Abnormal blood levels can indicate liver or bile-flow problems.

### **Bile Culture**

A laboratory test that grows microorganisms from a bile sample to identify infection in the gallbladder



---

or bile ducts and help select effective treatment.

### **Bile Duct**

A tube that carries bile from the liver and gallbladder into the first part of the small intestine. Blockage or inflammation can prevent bile flow and cause pain, jaundice, infection, or pancreatitis.

### **Bile Duct Cancer**

Cancer that begins in a bile duct. It can block bile flow and cause yellow skin, dark urine, pale stools, itching, abdominal pain, or weight loss; treatment depends on its location and spread.

#### **Alternate Names:**

Cholangiocarcinoma; Biliary Tract Cancer

### **Bile Duct Carcinoma**

Alternate name; see *Bile Duct Cancer*.

### **Bile Duct Obstruction**

A blockage that prevents bile moving from the liver or gallbladder into the intestine. It may cause jaundice, dark urine, pale stools, itching, abdominal pain, infection, or pancreatitis.

### **Bile Duct Stricture**

Narrowing of a bile duct that slows or blocks bile flow. It may result from surgery, inflammation, stones, or cancer and can cause jaundice, pain, infection, or abnormal liver tests.

### **Bile Ducts**

Tubes that carry bile from the liver and gallbladder to the small intestine. A blockage or narrowing can cause jaundice, abdominal pain, infection, or inflammation of the pancreas.

### **Bile Pigment**

A coloured substance, mainly bilirubin, made when haemoglobin from old red blood cells is broken down. The liver processes it and excretes it in bile; build-up causes yellowing of the skin and eyes.

### **Bile Reflux**

Backward flow of bile and other intestinal contents

into the stomach and sometimes the swallowing tube. It may cause upper-abdominal pain, nausea, bitter fluid, or inflammation and can occur after surgery or when normal stomach movement is impaired.

#### **Alternate Names:**

Duodenogastric Reflux; Biliary Gastritis

### **Bile Reflux Gastritis**

Inflammation and mucosal erythema of the stomach caused by the retrograde flow of duodenal contents, including alkaline bile acids and pancreatic enzymes, into the gastric cavity, frequently occurring following gastric resection or cholecystectomy.

#### **Alternate Names:**

Alkaline Reflux Gastritis

### **Bile Salt Emulsification**

The process in which bile salts break dietary fat into tiny droplets. This increases the surface area available to digestive enzymes and helps the intestine absorb fat and fat-soluble vitamins.

### **Bile Salt Malabsorption**

Failure to reabsorb enough bile salts in the lower small intestine. Excess bile salts enter the colon and draw in water, causing chronic watery diarrhoea, urgency, and abdominal discomfort.

### **Bile Salts**

Substances made from cholesterol in the liver and released in bile. They help break fats into small droplets so the intestine can digest and absorb fats and vitamins A, D, E, and K.

### **Bilevel Positive Airway Pressure**

Breathing support delivered through a mask that gives higher pressure when breathing in and lower pressure when breathing out. It can improve oxygen and carbon-dioxide levels in selected illnesses and requires monitoring.

### **Bilharzia**

An infection caused by *Schistosoma* parasites acquired when contaminated freshwater contacts the skin. It may affect the intestine, liver, urinary tract,

---

or other organs and can cause long-term inflammation and scarring.

### **Bili Lights**

Special blue lights used to treat jaundice in newborns. The light changes bilirubin in the skin into forms the body can remove more easily, helping prevent bilirubin from damaging the brain.

### **Biliary**

Relating to bile, the gallbladder, or the tubes that carry bile from the liver to the intestine.

### **Biliary Atresia**

A serious disease in newborns in which the bile ducts outside the liver are absent, scarred, or blocked. Bile cannot drain normally, causing persistent jaundice, pale stools, dark urine, liver injury, and progressive cirrhosis without prompt treatment.

#### **Alternate Names:**

Congenital Biliary Atresia Extrahepatic Biliary Atresia

### **Biliary Calculi**

Stones formed from hardened bile in the gallbladder or bile ducts. They may cause no symptoms or block bile flow, producing pain, jaundice, infection, or inflammation of the gallbladder or pancreas.

### **Biliary Cirrhosis**

Permanent liver scarring caused by long-term injury or blockage of the bile ducts. The term has also been used for primary biliary cholangitis, an immune disease affecting small bile ducts.

### **Biliary Colic**

Steady pain in the upper right abdomen or upper middle abdomen caused by a gallstone temporarily blocking the gallbladder outlet. It often follows a meal and lasts minutes to hours; fever or persistent pain suggests a complication.

### **Biliary Decompression**

A procedure that drains blocked bile ducts to restore bile flow. A doctor may place a stent or drainage

tube through an endoscope or through the skin, depending on the cause and location of the blockage.

### **Biliary Dyskinesia**

Gallbladder pain without gallstones or another obvious structural problem, often associated with abnormal emptying. Diagnosis requires typical symptoms and exclusion of other causes; a scan may help but is not decisive alone.

### **Biliary Fever**

Fever caused by infection or inflammation in the gallbladder or bile ducts. Fever with upper-abdominal pain or jaundice may indicate a rapidly worsening infection and needs urgent medical care.

### **Biliary Fistula**

An abnormal passage allowing bile to leak from the gallbladder or bile ducts into another organ, the abdominal cavity, or the skin. It may cause pain, infection, bile drainage, or skin irritation.

### **Biliary Sand**

Fine crystals or thick sludge suspended in bile, usually containing cholesterol, calcium salts, or pigment. It may disappear, form gallstones, block a duct, or cause gallbladder inflammation.

### **Biliary Scintigraphy**

A specialized nuclear medicine scan that tracks the production and flow of bile from the liver into the gallbladder, bile ducts, and small intestine using an injected radioactive tracer (such as technetium-99m labeled iminodiacetic acid). It is the most accurate imaging test for diagnosing acute cholecystitis (indicated by failure of the gallbladder to fill due to cystic duct blockage). It is also used to detect post-surgical bile leaks, evaluate biliary dyskinesia by measuring gallbladder emptying after cholecystokinin administration, and differentiate biliary atresia from neonatal hepatitis in jaundiced infants.

#### **Alternate Names:**

Cholescintigraphy; HIDA Scan; Hepatobiliary Scintigraphy; Hepatobiliary Iminodiacetic Acid Scan

---

## Biliary Stasis

Impairment, delay, or cessation of normal bile flow through intrahepatic or extrahepatic biliary channels, predisposing to biliary sludge, cholelithiasis, jaundice, and secondary bacterial cholangitis.

### Alternate Names:

Cholestasis

## Biliary Stent

A small tube placed inside a bile duct to keep it open and allow bile to drain when it is narrowed or blocked. Stents can move, clog, or cause infection or pancreatitis and require follow-up.

## Biliary System

The liver, gallbladder, and bile ducts that make, store, and carry bile into the small intestine to help digest fats.

## Biliary Tract

The network of ducts carrying bile from the liver and gallbladder to the small intestine. Blockage or inflammation can cause upper-abdominal pain, jaundice, infection, or pancreatitis.

## Biliary Tract Biopsy

The acquisition of tissue specimens or exfoliated cells from the extrahepatic or intrahepatic bile ducts or gallbladder for histopathologic examination, commonly performed via endoscopic retrograde cholangiopancreatography (ERCP), intraductal brush cytology, or image-guided percutaneous needle biopsy.

### Alternate Names:

Bile Duct Biopsy; Gallbladder Biopsy; Biliary Cytology

## Biliary Tract Injuries

Damage to the gallbladder or bile ducts, often after abdominal surgery or trauma. Bile may leak or drainage may become blocked, causing abdominal pain, jaundice, infection, or later narrowing.

## Biliary Tree

The anatomical conduit system of ducts and vessels that channels bile from intrahepatic bile canali-

culi through segmental ducts, right and left hepatic ducts, common hepatic duct, cystic duct, and common bile duct into the second part of the duodenum.

### Alternate Names:

Biliary Tract

## Bilious

Containing or relating to bile. Bilious vomit is usually yellow or green because bile has entered the stomach contents.

## Bilious Vomiting

Vomiting green or yellow fluid containing bile. In a newborn or infant, clearly green vomit can indicate bowel blockage and requires immediate medical assessment.

## Bilirubin

A yellow substance made mainly when old red blood cells are broken down. The liver changes it and sends it into bile; too much bilirubin in the blood causes yellowing of the skin and eyes.

## Bilirubin Blood Test

A blood test measuring total and direct bilirubin. High results can occur when red blood cells break down too quickly, the liver cannot process bilirubin, or bile flow is blocked; newborn levels help assess risk of brain injury.

## Bilirubin Encephalopathy

Brain injury caused by very high levels of unconjugated bilirubin, especially in newborns. Early signs may include poor feeding, sleepiness, weak or stiff muscles, and seizures; permanent injury is called kernicterus.

## Bilirubin Metabolism

The process by which the body makes bilirubin from old red blood cells, carries it to the liver, changes it into a water-soluble form, and removes it in bile and stool.

## Bilirubinuria

The presence of water-soluble bilirubin in urine. It

---

may occur with liver disease or blocked bile flow and can make urine appear unusually dark.

### **Biliverdin**

A green pigment formed when haemoglobin is broken down. The body normally converts it into bilirubin, which the liver processes and removes in bile.

### **Billroth I**

A surgical procedure for partial gastrectomy in which the distal stomach (antrum and pylorus) is resected and the remaining proximal gastric remnant is anastomosed directly to the duodenum (gastro-duodenostomy).

#### **Alternate Names:**

Billroth I Reconstruction; Gastroduodenostomy

### **Billroth I Operation**

A surgical reconstruction procedure following partial gastrectomy (antrectomy) in which the remaining gastric remnant is anastomosed directly to the duodenal stump (gastroduodenostomy).

### **Billroth II**

A surgical procedure for partial gastrectomy in which the distal stomach is resected, the duodenal stump is closed, and the remaining gastric remnant is anastomosed to a loop of the proximal jejunum (gastrojejunostomy), leaving a blind duodenal limb for biliopancreatic drainage.

#### **Alternate Names:**

Billroth II Reconstruction; Gastrojejunostomy

### **Billroth II Operation**

A surgical reconstruction procedure following partial gastrectomy in which the duodenal stump is oversewn closed and the gastric remnant is anastomosed directly to the side of a loop of proximal jejunum (gastrojejunostomy).

### **Billroth's Operation**

An older type of partial stomach-removal surgery. In Billroth I the remaining stomach is joined to the duodenum; in Billroth II it is joined to the jejunum,

a later part of the small intestine.

### **Bilophila**

A group of bacteria that can live without oxygen in the human intestine. Most are harmless there, but *Bilophila wadsworthia* has been found in some appendicitis, abdominal, wound, and bloodstream infections.

### **Bilophila Wadsworthia**

A bacterium that tolerates bile and normally occurs in small numbers in the colon. If intestinal bacteria enter damaged or normally sterile tissue, it may contribute to appendicitis, abdominal infection, or wound infection.

### **Bimanual**

Performed using both hands. A bimanual pelvic examination uses one hand inside the vagina and the other on the abdomen to assess pelvic organs.

### **Bimatoprost**

Bimatoprost is a prostaglandin analogue used as eye drops to lower intraocular pressure and, in some formulations, promote eyelash growth.

### **Bimedial**

Relating to or located on the inner side of two structures. The exact structures involved must be specified because the term alone does not identify a disease or diagnosis.

### **Bimellar**

Made of two thin layers or plates. The term describes the structure of a membrane, tissue, or other object and has no disease meaning by itself.

### **Binary Fission**

A form of reproduction in which one cell divides into two genetically similar cells. It is common among bacteria and many single-celled organisms.

### **Binder Excipient**

An inactive ingredient added during tablet manufacture to help powder particles stick together. It has

---

no intended treatment effect but affects the tablet's strength and breakdown.

### **Binding Protein**

A protein that attaches specifically to another molecule, such as a hormone, medicine, nutrient, DNA segment, or cell-surface signal. Binding can transport, store, activate, or block the other molecule.

### **Binding Site**

The particular area of a molecule where another molecule attaches. The interaction can change activity, transmit a signal, allow transport, or enable a laboratory test to detect the target.

### **Binet-Simon Scale**

An early intelligence test for children that compared performance with tasks expected at different ages. It influenced modern cognitive testing but is no longer the standard tool for clinical assessment.

### **Binge Drinking**

A pattern of consuming enough alcohol over a short period to produce a high blood alcohol concentration and increased risk of injury or acute toxicity.

#### **Alternate Names:**

Episodic Heavy Drinking; Heavy Episodic Drinking

### **Binge Eating Disorder**

Alternate form; see *Binge-Eating Disorder*.

### **Binge-Eating Disorder**

A mental-health condition involving repeated episodes of eating large amounts of food with loss of control and distress, without regular behaviors intended to compensate. Weight alone does not establish the diagnosis.

#### **Alternate Names:**

BED

### **Binge-Purge Disorder**

An older or patient-facing name for bulimia nervosa, an eating disorder involving recurrent binge eating followed by compensatory behaviors such as self-

induced vomiting, fasting, or excessive exercise.

### **Binocular Vision**

The use of both eyes together to produce one image and judge depth. Misalignment or poor coordination between the eyes can cause double vision, eyestrain, or reduced depth perception.

#### **Alternate Names:**

Two-Eyed Vision; Binocular Visual Function

### **Binocular Visual Loss**

Reduced vision affecting both eyes. Causes may involve the eyes, optic nerves, brain, or visual pathways; sudden loss, weakness, speech trouble, or severe headache requires emergency assessment.

### **Binswanger Disease**

A form of small-vessel disease in the brain that damages deep white matter. It can cause gradually worsening thinking, walking difficulty, mood changes, and urinary problems.

### **Binswanger's Dementia**

A form of vascular cognitive impairment caused by diffuse small-vessel disease affecting the brain's deep white matter.

#### **Alternate Names:**

Subcortical Ischemic Vascular Disease; Subcortical Arteriosclerotic Encephalopathy

### **Bio-Informatics**

The use of computers and statistics to store, compare, and interpret biological information such as DNA sequences, genes, proteins, and medical data.

### **Bioabsorbable Vascular Stent**

A temporary endovascular scaffold constructed from biodegradable polymers (such as poly-L-lactic acid) or bioresorbable metallic alloys (such as magnesium) engineered to provide vessel patency during vascular remodeling before undergoing hydrolytic degradation and systemic resorption.

#### **Alternate Names:**

Bioabsorbable Vascular Scaffold; BVS; Biore-sorbable Stent; BRS

---

## Bioaccumulation

The gradual build-up of a substance in an organism when absorption is faster than removal. Pollutants, medicines, and metals can accumulate in tissues and may cause harm when levels become high.

## Bioactive

Having an effect on living cells, tissues, or organisms, such as a bioactive drug, hormone, nutrient, or signaling molecule.

## Bioassay

A test that measures the effect or activity of a substance by observing its action on living cells, tissues, organisms, or a biological system. It can be used to estimate potency, toxicity, or biologic activity.

## Bioavailability

The amount of a medicine that reaches the bloodstream in an active form and how quickly it gets there. It depends on how the medicine is given, absorbed, and broken down; intravenous medicines have complete availability.

## Bioburden

The number and types of living microorganisms present on a surface, medical device, tissue, or medicine before sterilization. Testing it helps confirm that cleaning and sterilization methods are adequate.

## Biochanin A

A naturally occurring isoflavone found mainly in legumes and other plants. It is studied for biologic effects but is not established as a treatment for disease.

### Alternate Names:

Biochanin a Isoflavone; 5,7-Dihydroxy-4'-Methoxyisoflavone

## Biochemical

Relating to the chemical substances and reactions that occur in living organisms, including metabolism and cellular signaling.

## Biochemical Assessment

Evaluation using laboratory measurements such as blood proteins, salts, glucose, fats, vitamins, and minerals. Results can help assess nutrition, metabolism, organ function, or disease but must be interpreted with the clinical picture.

## Biochemical Reaction

A chemical change occurring in a living cell or body, usually controlled by enzymes. It may build or break down molecules and supports energy production, growth, repair, and normal organ function.

## Biochemical Recurrence

A rise in a laboratory marker after treatment that suggests disease remains or has returned, even though symptoms or scans may not yet show it. The meaning depends on the disease and the test used.

### Alternate Names:

Biochemical Relapse; Biochemical Failure

## Biochemistry

The study of the chemical substances and reactions occurring in living organisms. Clinical biochemistry uses laboratory measurements to assess health and disease.

## Biocompatibility

The ability of a material or device to perform its intended function without causing an unacceptable harmful reaction in body tissues.

## Biodegradable Implant

An implant designed to provide support temporarily and then gradually break down into substances the body can process or remove. Its safety depends on how quickly it breaks down and how tissues respond.

## Biodistribution

The way a medicine, tracer, or other substance spreads through the blood and different organs over time. It affects where treatment acts, how well an image is produced, and radiation or toxicity to healthy tissue.



---

## Bioengineering

The application of engineering principles to biology and medicine, including the design of medical devices, biomaterials, tissue-engineering systems, prostheses, and diagnostic technologies.

## Bioequivalence

A finding that two versions of a medicine enter the bloodstream at a similar rate and amount when given at the same dose. It supports the expectation of comparable effects and safety for approved uses.

## Bioethics

The study of moral questions in medicine, biology, research, and healthcare technology. It considers consent, privacy, fairness, benefit, harm, quality of life, and respect for patients and communities.

## Biofeedback

A technique that shows information about a body function, such as muscle tension, heart rate, or breathing, so a person can learn to control it. It may support treatment of selected pain, stress, or movement problems.

## Biofilm

A community of microorganisms attached to a surface and surrounded by a protective material they produce. Biofilms can form on teeth, wounds, catheters, and implants and may be difficult to remove.

## Biofilm-Associated Infections

Infections caused by microorganisms living in a protective layer on tissue, wounds, teeth, or medical devices. They can persist despite antibiotics and may require removal or replacement of an infected device.

## Biofluid

A fluid from a living body, such as blood, urine, saliva, spinal fluid, or joint fluid. Biofluids can be tested for cells, chemicals, microorganisms, medicines, or signs of disease.

## Biologic Therapy

Treatment using a medicine made from living cells or biological substances to target a specific immune or disease pathway. It is used for selected inflammatory, autoimmune, cancer, and other conditions.

## Biologic Therapy In Dermatology

Treatment with targeted biological agents, usually monoclonal antibodies or receptor proteins, that inhibit selected immune pathways. Biologics are used for diseases such as psoriasis, atopic dermatitis, hidradenitis suppurativa, and urticaria; infection risk, vaccination status, screening, and monitoring must be considered.

## Biologic Variation

Normal change in a body measurement within the same person over time or between different people. It must be considered when deciding whether a laboratory change represents real disease or treatment-related change.

## Biological

Relating to living organisms, their parts, functions, processes, or products. In medicine, a biological product is made from or resembles a substance from a living source.

## Biological Availability

The amount and speed at which a medicine or other substance becomes available to produce an effect in the body. It depends on the route, absorption, breakdown, and removal of the substance.

## Biological Clock

The body's internal timing system that helps coordinate daily sleep, alertness, hormone release, body temperature, and other repeating functions in response to light and other environmental signals.

## Biological Half-Life

The time needed for the body to remove half of a medicine or other substance through metabolism and excretion. It differs from the physical half-life of a radioactive substance and helps guide dosing or safety decisions.

---

## Biological Model

A cell, tissue, organism, computational system, or other representation used to study biological mechanisms, disease, or treatment responses.

## Biological Process

A coordinated series of molecular, cellular, or physiological events in a living organism, such as metabolism, cell division, inflammation, or tissue repair.

## Biological Relative

A person who shares genetic ancestry with someone. Family medical history uses biological relatives to assess inherited risks, while relatives by marriage or legal relationship do not share that genetic information.

## Biological Therapy

Treatment using biological substances, cells, vaccines, antibodies, or similar products to change immune activity or target a specific disease pathway. It is used in selected cancers, immune disorders, and other conditions.

## Biological Transport

The movement of substances such as water, salts, nutrients, medicines, or cells within the body or across cell membranes. It may occur by diffusion, channels, pumps, filtration, or transport vesicles.

## Biological Valve

A prosthetic cardiac valve fabricated from animal or human donor tissue (such as porcine aortic valves or bovine pericardial leaflets), which carries a low thrombogenicity and does not require lifelong anticoagulation, though with limited long-term durability compared to mechanical valves.

### Alternate Names:

Bioprosthetic Valve; Tissue Valve

## Biological Variability

Natural differences in a body measurement between people or changes in the same person over time, separate from laboratory error. It can affect how

test results are interpreted.

### Alternate Names:

Biologic Variation; Physiologic Variability

## Biological Warfare

The deliberate use of microorganisms, toxins, or other biological agents to cause illness, death, or disruption in people, animals, or plants.

## Bioluminescence

Light produced by a living organism through a chemical reaction. It is used by some organisms for communication or defense and is used in laboratories to detect biological activity.

## Biomarker

A measurable feature that gives information about a normal body process, disease, likely outcome, or response to treatment. It may be measured in blood, tissue, genes, images, or other samples.

## Biomarker Testing

Laboratory testing of genes, proteins, metabolites, or other measurable features to help identify disease, estimate likely outcome, choose treatment, or monitor response.

## Biomaterial

A natural or manufactured material designed to contact or interact with the body. Examples include implant surfaces, artificial joints, wound dressings, prostheses, and materials that release medicines.

## Biomechanical Compliance

How easily a body structure changes size or shape when pressure or force is applied. Lung and blood-vessel compliance affect breathing and circulation, while joint compliance affects movement and stability.

## Biomechanics

The study of how forces and movement affect the body. It examines walking, posture, joints, muscles, and loads to understand injury, improve rehabilitation, or design supportive devices.

---

## Biomedical Cement

A medical material used to fill gaps, stabilize broken bone, anchor an implant, or repair tissue. Its use and strength depend on the procedure and the tissue being treated.

## Biomedical Device

An instrument, implant, software system, or other product intended to diagnose, monitor, prevent, or treat disease or injury. Its safety and usefulness depend on proper design, testing, maintenance, and correct use.

## Biomedical Engineering

The use of engineering, materials, computing, and design to create or improve medical devices, diagnostic tests, implants, prostheses, imaging systems, and other healthcare technologies.

## Biomedical Tape

Adhesive tape designed for medical use, such as securing dressings, tubes, sensors, or small devices. Different types vary in strength, flexibility, water resistance, and risk of skin irritation.

## Biomedicine

Medicine grounded in the biological sciences, including the study of disease mechanisms, diagnosis, prevention, and treatment through laboratory and clinical research.

## Biometry

Measurement of body structures or functions. In eye care, optical biometry measures the eye to help calculate the strength of an artificial lens before cataract surgery.

## Biomicroscopy

Examination of living tissue with a microscope. In eye care, a slit lamp provides magnified views of the cornea, front chamber, lens, and other eye structures.

## Biomolecule

A molecule made by or found in living organ-

isms, such as a protein, fat, sugar, DNA, or RNA. Biomolecules provide structure, energy, information, and control of cell functions.

## Biomonitoring

Repeatedly measuring substances or body changes over time using blood, urine, breath, hair, or other samples. It can assess exposure to poisons or medicines, treatment levels, hormones, or disease activity.

## Biomphalaria

A group of freshwater snails that can carry the parasite *Schistosoma mansoni*. The parasite develops in the snails and can later infect people who contact contaminated freshwater.

## Bionics

The use of electronic or mechanical devices to replace or support body functions, such as hearing, movement, or sensation. Examples include cochlear implants, powered prostheses, and other assistive systems.

## Biophotonics

The use of light to study, diagnose, monitor, or treat living tissues and biological processes. Examples include optical imaging, fluorescence testing, laser surgery, and photodynamic therapy.

## Biophysical Profile

A pregnancy assessment combining fetal-heart-rate monitoring with ultrasound. It checks fetal movement, breathing movements, muscle tone, and amniotic-fluid volume to help assess fetal well-being.

## Biopsy

Removal of a small sample of cells or tissue so it can be examined under a microscope to diagnose disease, including cancer. Types include fine-needle aspiration (cells or fluid), core-needle (a tissue cylinder), incisional (part of an area), excisional (the whole area), punch (a circular piece of skin), and endoscopic biopsy (through a viewing tube). Possible complications include pain, bleeding, and infection; risks vary with the body site and procedure.

---

**Alternate Names:**

Biopsies

**Biopsy Lung**

Removal of a small lung-tissue sample using a needle, bronchoscope, or operation. Examination can investigate infection, inflammation, scarring, or cancer; the method depends on the lesion and procedural risks.

**Biopsy Needle**

A needle used to remove cells or tissue for laboratory examination. A fine needle usually collects loose cells, while a core needle removes a small cylinder that preserves the tissue's arrangement.

**Biopsy Site**

The body location from which cells or tissue are removed for testing. It is chosen to sample the abnormal area while limiting injury to nearby organs, nerves, and blood vessels.

**Biopsy Specimen**

Cells or tissue removed from the body for laboratory examination. Microscopy, special stains, immune tests, or genetic tests may be used to identify cancer, infection, inflammation, or another condition.

**Biosafety**

Practices, equipment, and containment methods used to prevent accidental exposure to or release of infectious organisms, biological materials, or toxins.

**Biosafety Level**

A classification describing the laboratory facilities, equipment, work practices, and protective measures needed to handle biological agents safely according to their risk.

**Biosecurity**

Measures that prevent biological agents or toxins from being accidentally or deliberately released, stolen, or misused. It includes secure facilities, controlled access, staff training, risk assessment, and responsible handling of biological materials.

**Biosensor**

A device that uses a biological material to recognize a substance and a detector to measure it. Medical examples include sensors for glucose, hormones, infectious organisms, or heart-related markers.

**Biosimilar**

A biological medicine highly similar to an already approved biological medicine, with no meaningful differences in safety or effectiveness for approved uses. Because these medicines are complex, similarity is shown through comparative testing.

**Biospecimen**

A sample of human tissue, cells, blood, urine, or another body fluid collected for testing or research. Correct collection, labelling, storage, and handling help keep the sample reliable.

**Biostatistics**

The use of statistics to plan medical studies, analyse health data, measure uncertainty, and interpret evidence about disease, diagnosis, prevention, and treatment.

**Biosynthesis**

The process by which living cells use enzymes and energy to build complex molecules from simpler substances, such as making proteins from amino acids.

**Biot Breathing**

An irregular breathing pattern with groups of normal-sized breaths interrupted by unpredictable pauses in breathing. It may occur with serious damage affecting the lower brainstem and requires urgent assessment.

**Biot's Breathing**

An abnormal pattern of irregular breaths interrupted by unpredictable pauses in breathing. It suggests disruption of the brain's breathing control and may occur with severe neurological illness.

---

## **Biotechnology**

The use of living cells, organisms, or biological molecules to develop products and processes such as vaccines, biological medicines, genetic tests, and gene therapies.

## **Biotelemetry**

Measuring and sending body information from a person or animal to a distant receiver. It can be used for heart-rhythm, temperature, breathing, glucose, or intensive-care monitoring.

## **Bioterrorism**

The deliberate release of infectious organisms, toxins, or other biological agents to cause illness, death, fear, or disruption. Response involves rapid detection, public-health coordination, protection, testing, isolation when needed, and treatment or prevention.

## **Bioterrorism Agents**

Infectious organisms or toxins that could be deliberately released to cause widespread illness or panic. Public-health authorities prepare for agents causing anthrax, smallpox, botulism, plague, tularemia, or severe viral haemorrhagic fever.

## **Biotherapy**

Treatment using living organisms, cells, genes, or biological substances to prevent or treat disease. Immunotherapy, some cell therapies, and selected biological medicines are forms of biotherapy.

## **Biotin**

A water-soluble B vitamin needed for enzymes involved in processing fats, sugars, and amino acids. Deficiency is uncommon but can cause a skin rash, hair loss, weakness, or neurological symptoms.

## **Biotin Vitamin B7**

Alternate name; see *Biotin*.

## **Biotinidase Deficiency**

A rare inherited disorder in which the body cannot recycle biotin efficiently. Without treatment it can cause seizures, developmental problems, hearing or

vision loss, skin rash, and weakness; lifelong biotin usually prevents symptoms.

## **Biotinylation**

The attachment of biotin to another molecule, usually a protein, so it can be found or separated in a laboratory test. The biotin acts as a tag that binds to special substances, making the target easier to detect or collect. Biotinylation can occur naturally or be done artificially.

## **Biotransformation**

The natural process by which the body changes drugs and other foreign substances, mainly through enzymes in the liver but also in other tissues. It often makes them less active and easier to remove. Phase I reactions modify the substance; phase II reactions may attach another molecule to help eliminate it. These steps are not always sequential, and the process can produce an active or harmful substance instead.

## **Biotrauma**

Inflammation caused by mechanical injury to the lungs during breathing-machine treatment. Inflammatory signals can spread through the body and contribute to organ injury; careful, lung-protective settings reduce the risk.

## **BiPAP**

Bilevel Positive Airway Pressure (BiPAP), a form of noninvasive breathing support delivered through a mask. It provides higher pressure during inhalation and lower pressure during exhalation. It may be used for conditions such as hypercapnic COPD exacerbation, sleep-related breathing disorders, and cardiogenic pulmonary edema.

### **Alternate Names:**

Bilevel Positive Airway Pressure

## **BiPAP Machine**

A device that delivers two levels of breathing pressure through a mask—higher during inhalation and lower during exhalation. Settings are prescribed and adjusted according to the person's breathing problem and response.

---

## Biparental

Involving both biological parents. In genetics, biparental inheritance means that a child receives relevant genetic material from both the mother and the father.

## Biparietal Diameter

The width of a baby's head measured by ultrasound from one side to the other. It helps estimate pregnancy age and monitor growth, but is interpreted with other measurements because head shape can affect the result.

## Bipartite Ossification

Formation or persistence of two separate ossification centers in a bone. If they do not fuse, a bipartite bone, such as a bipartite patella, may result. This is usually an incidental developmental variant but can become painful after injury or repetitive stress.

## Biphenyl Compounds

Chemicals made of two linked benzene rings. Polychlorinated biphenyls (PCBs) are long-lasting industrial pollutants; exposure can harm the liver, immune system, and hormone system and may increase cancer risk.

## Bipolar

Alternate name; see *Bipolar Disorder*.

## Bipolar Affective Disorder

Alternate name; see *Bipolar Disorder*.

## Bipolar Disorder

A mental health condition involving episodes of mania, hypomania, and/or major depression. Bipolar I is defined by at least one manic episode; depressive episodes are common but not required. Bipolar II includes at least one hypomanic episode and at least one major depressive episode, without a history of mania. Episodes can affect mood, sleep, energy, judgment, and daily functioning.

### Alternate Names:

Manic-Depressive Illness

## Bipolar Disorder in Child and Teens

Bipolar disorder beginning during childhood or adolescence, with episodes of unusually elevated or irritable mood and increased energy alternating with depression. Diagnosis requires careful assessment because symptoms overlap with other developmental and psychiatric conditions.

## Bipolar I Disorder

A mood disorder defined by at least one manic episode lasting at least seven days or requiring hospitalization. A manic episode is a distinct period of abnormally elevated, expansive, or irritable mood and increased goal-directed activity or energy.

## Bipolar II Disorder

A mood disorder characterized by at least one hypomanic episode and at least one major depressive episode, with no history of a full manic episode. The combination of hypomania and depression can cause substantial distress or impairment even though hypomania is less severe than mania.

## Bipolar Neuron

A bipolar neuron has one dendrite and one axon extending from opposite sides of its cell body, as found in the retina and sensory pathways.

## Bipolaris

A genus of fungi found mainly in soil and plants. Some species can rarely cause eye, sinus, skin, or invasive infections, particularly in people with reduced immunity. Identification requires laboratory culture or molecular testing because treatment depends on the species and site of infection.

## Birbeck Granules

Small rod- or tennis-racket-shaped structures found inside Langerhans cells. They are associated with these cells but are not by themselves proof of Langerhans cell histiocytosis.

## Bird Fancier's Lung

A form of hypersensitivity pneumonitis caused by breathing proteins from bird feathers or droppings. It can cause cough and shortness of breath; repeated



---

exposure may lead to permanent lung scarring.

### **Bird Flu**

An infectious zoonotic illness caused by avian influenza type A viruses (notably H5N1, H5N6, H7N9) that naturally circulate among wild waterfowl and domestic poultry. Transmission to humans occurs via direct contact with infected birds or contaminated environments, producing acute respiratory distress syndrome (ARDS), multi-organ dysfunction, and high case-fatality rates.

#### **Alternate Names:**

Avian Influenza; Avian Flu; Bird Flu; H5N1 Infection

### **Birt-Hogg-Dube Syndrome**

An autosomal dominant genodermatosis caused by germline mutations in the *FLCN* gene, characterized by benign cutaneous fibrofolliculomas, pulmonary cysts predisposing to spontaneous pneumothorax, and an elevated risk of renal cell carcinomas.

#### **Alternate Names:**

BHD Syndrome

### **Birt-Hogg-Dube Syndrome (BHDS)**

Alternate spelling and abbreviation; see *Birt-Hogg-Dubé Syndrome*.

### **Birt-Hogg-Dubé Syndrome**

An inherited condition caused by a change in the *FLCN* gene. It can cause small, usually harmless skin growths, cysts in the lungs with repeated collapsed lung, and an increased risk of kidney tumors. It is autosomal dominant, meaning one altered copy of the gene can cause the condition.

### **Birth**

The process by which a baby leaves the uterus, including labour, delivery of the baby, and delivery of the placenta.

### **Birth Asphyxia**

Failure to establish or maintain effective breathing and oxygen supply around birth. It can injure the brain or other organs and needs immediate newborn

assessment and resuscitation; it may cause neonatal encephalopathy but is not the same diagnosis.

### **Birth Control**

Methods used to prevent pregnancy, including condoms, pills, implants, injections, intrauterine devices, fertility-awareness methods, and permanent procedures. Choice depends on effectiveness, health, side effects, future plans, access, and personal preference.

### **Birth Control And Family Planning**

Planning whether and when to have children using contraception, fertility care, pregnancy advice, and reproductive health services. Care respects personal choices and considers health conditions, safety, goals, and informed consent.

### **Birth Control Methods**

Birth-control methods include condoms, pills, implants, injections, intrauterine devices, patches, rings, fertility-awareness approaches, and permanent contraception. Effectiveness, sexually transmitted infection protection, side effects, reversibility, and medical eligibility vary by method.

### **Birth Control Pill Overdose**

Illness after taking too many contraceptive tablets. Most exposures cause little or no serious harm, but nausea, vomiting, drowsiness, or other symptoms may occur, and the product and amount taken help guide management.

### **Birth Control Pills**

Tablets taken on a schedule to prevent pregnancy, usually by stopping ovulation, thickening cervical mucus, or changing the uterine lining. They do not protect against sexually transmitted infections and work best when taken correctly.

### **Birth Defect**

A problem in the body's structure or function that is present at birth. It may result from genetic or chromosome changes, infections, medicines, alcohol, poor nutrition, or other causes; sometimes no cause is found.

---

**Alternate Names:**

Congenital Anomaly; Congenital Malformation; Birth Defects; Congenital Disorder

**Birth Defects**

Structural or functional problems present at birth that may affect development or health. Causes include genetic or chromosomal changes, infections, medicines, nutritional deficiencies, and environmental exposures; sometimes no cause is found.

**Birth Defects Testing**

Prenatal or newborn testing used to screen for or diagnose structural, genetic, chromosomal, metabolic, or functional conditions present at birth. Tests may include ultrasound, serum screening, cell-free DNA, diagnostic procedures, and newborn screening.

**Birth Injury**

Physical harm to a newborn during labour or delivery. It may affect the skin, bones, nerves, muscles, or brain and can result from difficult delivery, the baby's position, or medical complications.

**Birth Order**

The sequence in which siblings are born within a family. While birth order has been studied for psychological effects, its direct medical significance is limited to obstetric considerations such as risks associated with grand multiparity.

**Birth Pains**

Birth pains are painful uterine contractions and related pressure during labor as the cervix dilates and the baby moves through the birth canal.

**Birth Rate**

The number of live births in a population during a specified period, usually expressed per 1,000 people per year.

**Birth Spacing**

The practice of planning the time between giving birth and becoming pregnant again. Recommended spacing varies by person and health guidance, but

waiting about 18 months or longer may reduce some risks for the parent and baby.

**Alternate Names:**

Interpregnancy interval; Child spacing; Pregnancy spacing.

**Birth Weight**

The first weight recorded after birth; for a liveborn infant, it is preferably measured within the first hour, before significant postnatal weight loss. Low birth weight is less than 2,500 g regardless of gestational age; very low birth weight is less than 1,500 g, and extremely low birth weight is less than 1,000 g. These are overlapping categories, and lower birth weight is associated with greater risk of neonatal complications.

**Birthmark**

A localized mark on the skin or in blood vessels that is present at birth or appears soon afterward. Examples include pigmented moles, haemangiomas, and port-wine stains.

**Birthmarks**

Marks on the skin present at birth or appearing early in life. Most are harmless, but rapid growth, colour or shape change, bleeding, ulceration, pain, or a mark affecting vision or breathing needs medical assessment.

**Birthmarks And Other Skin Pigmentation Problems**

Birthmarks or changes in skin colour may result from blood vessels, pigment cells, inflammation, medicines, hormones, or inherited conditions. A rapidly changing, bleeding, ulcerated, painful, or distinctly different mark requires examination.

**Bis In Die**

Latin for "twice daily," traditionally abbreviated b.i.d. Clear prescriptions state when doses are to be taken rather than relying only on this abbreviation.

**Bisacodyl**

A stimulant laxative that increases bowel movement and helps empty the bowel. It is used short term

---

for constipation or before some procedures; cramps, diarrhoea, and dehydration can occur.

### **Bisexual**

Bisexual describes sexual or romantic attraction to more than one gender; it is an identity or orientation, not a disease or disorder.

### **Bisexuality**

Bisexuality is a sexual orientation involving attraction to more than one gender; it is not a disease or disorder.

### **Bishop Score**

A score used before inducing labour to describe how ready the cervix is for birth. It considers opening, thinning, softness, position, and the baby's head position; a higher score generally predicts an easier induction.

### **Bismuth**

A metal used in some medicines and industrial products. Certain bismuth compounds can soothe diarrhoea or stomach symptoms, but excessive or prolonged exposure can cause nervous-system or kidney toxicity.

### **Bisoprolol**

A beta-1-selective blocker used to treat hypertension, angina, and selected patients with heart failure. It lowers heart rate and blood pressure and may cause bradycardia, hypotension, or fatigue.

### **Bispecific Antibody Therapy**

Treatment using an engineered antibody that binds two different targets, such as a cancer-cell marker and an immune-cell marker, to bring cells together or block more than one disease pathway. It is used for selected cancers and can cause immune reactions or other treatment-specific effects.

### **Bisphenol A**

An endocrine-disrupting industrial chemical used predominantly in the synthesis of polycarbonate plastics and epoxy resins lining food and beverage

containers. It exhibits weak estrogenic and antiandrogenic activity and has been implicated in metabolic and reproductive disturbances.

#### **Alternate Names:**

BPA; 4,4'-(Propane-2,2-diyl)diphenol

### **Bisphosphonate**

A medicine that slows the removal of old bone. It is used mainly for osteoporosis, Paget disease, and some cancer-related bone problems. Rare but important risks include jaw-bone damage and unusual thigh-bone fractures.

### **Bisphosphonate-Related Osteonecrosis of the Jaw**

A severe adverse reaction characterized by exposed, necrotic bone in the maxillofacial region persisting for more than eight weeks in patients receiving bisphosphonate therapy without prior craniofacial radiation therapy.

#### **Alternate Names:**

BRONJ; Medication-Related Osteonecrosis of the Jaw

### **Bisphosphonates**

Medicines that slow the breakdown of bone. They are used mainly to prevent fractures in osteoporosis and to treat Paget disease and some bone problems caused by cancer.

#### **Alternate Names:**

Bisphosphonate

### **Bitemporal Hemianopia**

Loss of side vision on the outer side of both eyes. It usually occurs when a growth or other problem presses on the place where the two optic nerves cross, often near the pituitary gland.

### **Bitemporal Hemianopsia**

A visual field deficit in which vision is lost in the outer (temporal) halves of both the right and left visual fields, classically caused by compression of crossing decussating nerve fibers at the optic chiasm by a pituitary adenoma or craniopharyngioma.

---

## Bitewing

A dental X-ray showing the crowns of upper and lower teeth and the bone around them in one image. It is especially useful for finding decay between teeth and early loss of supporting bone.

## Bitewing Radiograph

An X-ray taken inside the mouth that shows the crowns of back teeth and the nearby gum-supporting bone. It helps detect decay between teeth and bone loss from gum disease.

## Bitot's Spots

Foamy, white or grey patches on the white part of the eye, usually near the cornea. They are commonly linked to vitamin A deficiency and may occur with dry eyes or poor night vision. They need medical assessment because untreated vitamin A deficiency can damage the eye; vitamin A treatment should follow clinical advice.

### Alternate Names:

Bitot Spots; Conjunctival Foamy Patches; Keratinized Conjunctival Plaques

## Bituminosis

A work-related breathing disorder caused by repeated exposure to bitumen or asphalt fumes and dust. It may irritate the airways and contribute to chronic cough, breathlessness, or other lung disease.

## Biuret Test

A laboratory test that detects protein by turning violet when protein reacts with a copper-containing solution. It is mainly used in teaching, research, and some chemical analyses rather than routine patient diagnosis.

## Bivalve

Bivalve means having two shells or valves; bivalve mollusks may be relevant to food allergy, poisoning, or parasitic exposure.

## Bizarre Parosteal Osteochondromatous Proliferation

A rare benign but locally recurrent bone-surface le-

sion, also called Nora Lesion, composed of cartilage, bone, and spindle cells; it most often affects the hands or feet.

## BK Virus

A common virus that usually remains inactive after childhood infection. It can reactivate when immunity is suppressed, especially after organ transplantation, and damage the transplanted kidney or urinary tract.

## BK Virus Nephropathy

Kidney inflammation and damage caused by reactivation of BK virus, usually in a kidney-transplant recipient. It can reduce transplant function and is managed mainly by carefully reducing immune-suppressing treatment and monitoring viral levels.

## Black Death

The major plague pandemic of the fourteenth century, caused mainly by the bacterium *Yersinia pestis*. It caused widespread severe illness and death and is distinct from the individual forms of plague seen today.

## Black Eye

Bruising and swelling around the eye, usually caused by blunt trauma. Vision may remain normal, but eye pain, double vision, reduced vision, an irregular pupil, severe headache, vomiting, or loss of consciousness can indicate an eye or orbital injury and requires urgent assessment.

### Alternate Names:

Periorbital Hematoma; Shiner

## Black Fly

A small biting fly that can cause painful or itchy skin reactions. In some tropical regions, black flies spread the parasite that causes river blindness.

## Black Hairy Tongue

A usually harmless dark, furry appearance of the tongue caused by overgrown surface projections trapping debris or pigment. It may improve with tongue cleaning, good oral hygiene, and addressing smoking or dry mouth.

---

## Black Lung

Lung disease caused by long-term inhalation of coal dust, usually in miners. It can cause cough and breathlessness and may progress from small areas of dust-related damage to extensive permanent scarring.

## Black Lung Disease

A dust-related lung disease caused by prolonged coal-dust exposure. It may cause no symptoms at first but can lead to cough, breathlessness, reduced lung function, and severe permanent scarring.

## Black Nightshade Poisoning

Poisoning after eating black nightshade or its berries, which may contain toxic plant chemicals. It can cause nausea, vomiting, abdominal pain, confusion, enlarged pupils, seizures, or abnormal heart rhythms.

## Black Or Tarry Stools

Black or tarry stools may indicate upper gastrointestinal bleeding, but can also result from iron or bismuth. New or persistent findings require medical assessment.

## Black Stain

Dark discoloration on the tooth surface, often near the gumline, caused by external deposits or chromogenic bacteria. Dental assessment determines the cause and appropriate removal.

## Black Triangle

A triangular gap between neighboring teeth near the gumline, usually caused by loss or shrinkage of the gum papilla after gum disease, injury, or dental treatment. It may trap food and can be treated cosmetically or orthodontically.

## Black Widow Spider

The black widow spider is a venomous spider whose bite can cause severe muscle pain, abdominal cramping, sweating, hypertension, and autonomic symptoms.

**Alternate Names:**

Black Widow Spider Bite; Latrodectism

## Black Widow Spider Bite

Envenomation by a *Latrodectus* spider. It may cause local pain followed by severe muscle cramps, abdominal rigidity, sweating, nausea, hypertension, or autonomic symptoms; severe cases need urgent medical care.

## Blackflesh

A nontechnical description of tissue that looks dark or black. Causes include bruising, severe loss of blood supply, dead tissue, infection, or staining. New, painful, cold, or spreading discoloration needs urgent medical assessment.

## Blackhead

An open blocked hair follicle filled with oil and dead skin cells. The material looks black because it reacts with air, not because the skin is dirty.

## Blackheads

Open comedones formed when a hair follicle becomes blocked with sebum and keratin and the surface material oxidizes. They are a common non-inflammatory lesion of acne, especially on the face, chest, and back.

## Blackout

A period for which a person cannot remember events, often after heavy alcohol use. Memory loss can also occur with drugs, seizures, head injury, fainting, or other causes of altered consciousness.

**Alternate Names:**

Amnestic Episode; Alcohol-Related Amnesia When Caused by Alcohol

## Blackwater Fever

A severe complication of malaria in which red blood cells break down inside blood vessels. It can cause dark urine, severe anaemia, jaundice, kidney failure, and shock and requires emergency treatment.

## Bladder

A muscular hollow organ that stores urine from

---

the kidneys and contracts to release it through the urethra during urination.

### **Bladder Augmentation**

Surgery that enlarges the bladder, usually by adding a section of bowel or another tissue. It can increase urine storage when the bladder is too small or has high pressure.

### **Bladder Biopsy**

Removal of a small piece of bladder lining, usually during cystoscopy, for laboratory examination. It can investigate a visible growth, unexplained blood in the urine, inflammation, or suspected bladder cancer.

### **Bladder Calculi**

Alternate name; see *Bladder Stones*.

### **Bladder Cancer**

A malignant tumor arising in the urinary bladder, most often from the urothelial lining. Blood in the urine is a common presenting feature, although symptoms vary.

#### **Alternate Names:**

Urinary Bladder Neoplasm; Urothelial Carcinoma

### **Bladder Carcinoma**

Cancer that begins in the lining of the urinary bladder, most often in cells called urothelial cells. Blood in the urine is common; smoking and some workplace chemical exposures increase risk.

### **Bladder Diary**

A record of drinks, urination times, urine amounts, urgency, leakage, and nighttime voiding over several days. It helps describe bladder habits and patterns of urinary symptoms.

### **Bladder Diseases**

Disorders of the bladder, including infection, stones, tumors, neurogenic dysfunction, inflammation, obstruction, and impaired emptying.

### **Bladder Disorders**

A broad spectrum of pathological conditions affecting the urinary bladder, encompassing infectious cystitis, interstitial cystitis (bladder pain syndrome), urothelial carcinoma, neurogenic bladder dysfunction, calculi, and bladder outlet obstruction.

#### **Alternate Names:**

Bladder Diseases; Urinary Bladder Diseases; Diseases of the Bladder

### **Bladder Diverticula**

Pockets that push outward from the bladder wall. They may be present from birth or develop when urine repeatedly faces resistance leaving the bladder, and can trap urine, leading to infection or stones.

### **Bladder Diverticulum**

A pouch extending from the bladder wall. It may cause no symptoms or trap urine and lead to infection, stones, or incomplete emptying; treatment addresses the cause and symptoms.

### **Bladder Exstrophy**

Bladder Exstrophy is a rare birth defect in which the bladder is open and exposed outside the body through a defect in the lower abdominal wall. It is part of the exstrophy–epispadias complex and may also affect the urethra, pelvic bones, pelvic floor, abdominal muscles, and genitals. It can cause difficulty storing urine, urinary leakage, urine-control problems, and risks to the kidneys.

### **Bladder Exstrophy Repair**

Reconstructive surgery for bladder exstrophy that places the bladder inside the body and repairs the lower abdomen and genitals. Treatment is often staged and aims to protect the kidneys and improve urine control.

### **Bladder Fistula**

An abnormal passage between the bladder and another organ or the skin, often the vagina, bowel, or uterus. It may cause constant urine leakage, repeated infection, gas or stool in urine, or unusual vaginal wetness.



---

## Bladder Health And Disorders

Healthy bladder function stores urine and releases it at an appropriate time. Common problems include infection, urgency, leakage, inability to empty, stones, inflammation, and nerve-related bladder dysfunction.

### Bladder Infection

An infection of the bladder, usually caused by bacteria. It may cause burning urination, frequent or urgent urination, lower-abdominal discomfort, cloudy or bloody urine, and sometimes fever.

### Bladder Infections in Children

Urinary tract infections involving a child's bladder, usually caused by bacteria. Symptoms may include painful urination, frequent urination, new wetting, abdominal or back pain, fever, or vomiting; infants may have only fever or poor feeding.

### Bladder Infections in Teens and Adults

Bacterial infection of the bladder, commonly called cystitis. Burning urination, urgency, frequency, lower-abdominal discomfort, or blood in the urine are typical; fever, flank pain, or vomiting may indicate kidney involvement.

### Bladder Inflammation

Alternate name; see *Interstitial Cystitis*.

### Bladder Neck

The narrow lower part of the bladder where it joins the urethra. Its muscles help control the release of urine and can contribute to blockage if they scar or fail to relax.

### Bladder Neck Contracture

Scar-related narrowing of the bladder outlet, often after prostate surgery or other treatment. It can restrict urine flow and cause a weak stream, difficulty emptying, or urinary retention.

### Bladder Neck Obstruction

Narrowing or blockage where the bladder joins the urethra, making it difficult for urine to leave the

bladder. It may cause a weak stream, straining, incomplete emptying, urinary retention, or recurrent infection.

### Bladder Neoplasm

An abnormal growth arising in the bladder. It may be noncancerous or cancerous; diagnosis requires examination, imaging, cystoscopy, and often a tissue sample to determine its type and extent.

### Bladder Outlet Obstruction

A blockage where urine leaves the bladder, caused by an enlarged prostate, scarred urethra, stone, tumour, or pelvic-organ prolapse. It may cause a weak stream, difficulty starting, retention, infection, or kidney damage.

### Bladder Pain

Pain or pressure felt in the lower abdomen or pelvis that may relate to infection, stones, inflammation, bladder pain syndrome, injury, or cancer. Blood in urine, fever, severe pain, or inability to urinate needs prompt assessment.

### Bladder Pain Syndrome

Long-lasting bladder or pelvic pain, pressure, or discomfort with urinary urgency or frequency when infection and other causes have been excluded. Symptoms often flare and settle and require individualized treatment.

### Bladder Perforation

A tear or hole in the bladder wall caused by trauma, pelvic surgery, instrumentation, severe inflammation, or obstruction. Urine may leak into surrounding tissues, causing abdominal pain, distention, blood in the urine, or infection. It requires urgent medical assessment and may need catheter drainage or surgical repair.

### Bladder Problems

Conditions that interfere with storing or passing urine, such as leakage, urgency, repeated infection, stones, or inability to empty. Causes include infection, weak pelvic muscles, nerve disease, prostate enlargement, or blockage.

---

## Bladder Prolapse

Alternate name; see *Anterior Vaginal Prolapse*.

## Bladder Retraining

A planned urination schedule that gradually increases time between bathroom visits to improve bladder control and reduce urgency.

## Bladder Spasm

An involuntary contraction of the bladder muscle that can cause sudden urgency, cramping, urine leakage, or pain. It may occur with infection, catheter use, bladder irritation, neurological conditions, or after surgery. Treatment addresses the cause and may include bladder training or medicines.

## Bladder Stones

Hard mineral deposits in the bladder that may cause pain, blood in urine, frequent urination, interrupted flow, or difficulty emptying. Some cause no symptoms and are found during tests for another problem.

### Alternate Names:

Bladder Calculi

## Bladder Trauma

Injury to the bladder caused by a blow, pelvic fracture, penetrating wound, surgery, or a medical instrument. It may cause blood in the urine, lower-abdominal pain, difficulty urinating, or leakage of urine into nearby tissues.

## Blalock-Taussig Operation

Surgery that creates a connection between a major artery and a lung artery to increase blood flow to the lungs in selected blue-baby heart defects. Modern versions usually use a small artificial tube.

## Blalock-Taussig Shunt

A palliative surgical procedure designed to augment pulmonary blood flow in infants with cyanotic congenital heart defects characterized by restricted pulmonary flow (such as tetralogy of Fallot or pulmonary atresia). The classic procedure anastomosed the subclavian artery directly to the pulmonary

artery, whereas the modified technique uses a synthetic Gore-Tex graft between the subclavian and pulmonary arteries.

### Alternate Names:

BT Shunt; Blalock-Thomas-Taussig Shunt

## Bland Diet

A temporary eating pattern using soft, mildly seasoned foods and avoiding items that worsen a person's symptoms. It may help some stomach or bowel problems but does not treat their underlying cause.

## Blaschko Lines

Invisible patterns formed by the movement of skin cells before birth. Some birthmarks and skin diseases follow these lines as stripes or swirls; they usually do not follow nerves or blood vessels.

## BLAST

An immature cell that normally develops into a blood cell. A high number of blasts in blood or bone marrow may indicate acute leukaemia or another serious marrow disorder.

## Blast Cell

An immature precursor cell, especially one in the bone marrow that will normally develop into a blood cell. Too many blasts can indicate acute leukaemia or another marrow disorder.

## Blast Crisis

An advanced, rapidly worsening phase of some blood cancers in which immature blast cells fill the bone marrow or blood and behave like acute leukaemia. It requires urgent specialist treatment.

## Blast Injuries

Injuries caused by an explosive pressure wave and its associated fragments, heat, displacement, or toxic exposure. They may affect the lungs, ears, brain, abdomen, eyes, bones, and soft tissues.

## Blast Injury

Harm caused by an explosion. The pressure wave can injure lungs, ears, or organs; flying objects can

---

cause wounds; being thrown can cause fractures; burns, smoke, toxins, and crushing can add further injuries.

### **Blast Phase**

An advanced phase of some blood cancers in which many immature blast cells appear in the blood or marrow. The disease becomes more aggressive and may resemble acute leukaemia.

### **Blastema**

A mass of undifferentiated, highly proliferating embryonic cells capable of growth and cellular differentiation into organs, tissues, or specialized structures; in nephroblastoma (Wilms tumor), blastemal cells constitute a primitive malignant component.

### **Blastic Plasmacytoid Dendritic-Cell Neoplasm**

A rare, fast-growing blood cancer arising from immature immune cells. It often affects the skin, bone marrow, lymph nodes, or nervous system and requires urgent specialist treatment.

### **Blastocyst**

An early embryo formed about five to six days after fertilization. It is a hollow group of cells with an inner group that can become the fetus and an outer layer that helps form the placenta.

### **Blastocyst Transfer**

An assisted-reproduction procedure in which an embryo grown in the laboratory for about five or six days is placed inside the uterus. The decision to transfer one or more embryos considers success rates, patient factors, and the risk of multiple pregnancy.

### **Blastocystis**

A common microscopic organism that may be found in the digestive tract after contaminated food or water is swallowed. Many people have no symptoms, and it is not always clear whether it causes illness. Diarrhoea, stomach cramps, bloating, or nausea may occur. Stool testing can detect it, but a positive result does not prove that it caused the symptoms; treatment is considered only in selected persistent

cases.

#### **Alternate Names:**

Blastocystis Hominis; Blastocystis spp.; Blastocystosis; Intestinal Blastocystis

### **Blastocystis Hominis**

Historical name; see *Blastocystis*.

### **Blastocystosis**

Alternate name; see *Blastocystis*.

### **Blastoma**

A cancer that develops from immature or embryonic cells. The specific name identifies the organ and tumour type, such as a kidney, nerve, liver, or eye blastoma.

### **Blastomyces Dermatitidis**

A fungus found in soil and damp organic material that can cause blastomycosis when its spores are inhaled. Infection usually affects the lungs but can spread to skin, bones, joints, or other organs.

### **Blastomycosis**

A fungal infection acquired by inhaling *Blastomyces* spores, usually from soil. It commonly causes pneumonia and may spread to the skin, bones, joints, or other organs; symptoms can develop gradually.

### **Blasts**

Immature cells in blood-forming tissue that normally develop into mature blood cells. An excess in blood or bone marrow may indicate acute leukaemia or another serious marrow disorder.

### **Blautia Producta**

A bacterium that normally lives in the human intestine and grows without oxygen. It is usually harmless but has occasionally been found in infections, especially in people with serious illness.

### **Bleb**

A small thin-walled pocket containing air or fluid. In the lung, a bleb can rupture and cause a collapsed lung; on the skin, the term may describe a blister-

---

like lesion.

### **Bleeder's Disease**

Alternate name; see *Hemophilia*.

### **Bleeding**

The escape of blood from blood vessels. It may be visible outside the body or occur inside a tissue, organ, or body cavity; severe, unexplained, or persistent bleeding needs medical attention.

### **Bleeding Disorder**

A condition causing unusually heavy, prolonged, or spontaneous bleeding because of problems with clotting proteins, platelets, blood vessels, or medicines. Examples include hemophilia, von Willebrand disease, platelet disorders, and some forms of disseminated intravascular coagulation.

### **Bleeding Disorders**

Alternate name; see *Bleeding Disorder*.

### **Bleeding Esophageal Varices**

Severe bleeding from enlarged, fragile veins in the lower swallowing tube. These veins usually develop when liver disease raises pressure in veins entering the liver. Vomiting blood or passing black stools is an emergency.

### **Bleeding Gums**

Bleeding from the gum tissue, often caused by plaque-related gingivitis, periodontitis, trauma, or vigorous brushing. It can also occur with blood disorders or medicines that affect clotting. Persistent bleeding, swelling, loose teeth, or spontaneous bleeding requires dental or medical evaluation.

### **Bleeding Into A Thyroid Nodule**

Alternate name; see *Hemorrhagic Thyroid Nodule*.

### **Bleeding Tendency**

An increased susceptibility to bleeding, which may result from platelet disorders, coagulation factor deficiencies, vascular abnormalities, or anticoagulant medications. Evaluation includes personal and fam-

ily history, complete blood count, and coagulation studies.

### **Bleeding Time**

An older test measuring how long a small skin puncture takes to stop bleeding. It mainly reflected platelet function and small blood vessels and has largely been replaced by more reliable laboratory tests.

### **Bleeding Varices**

Enlarged fragile veins, usually in the swallowing tube or stomach, that have ruptured and are bleeding. Vomiting blood, black stools, weakness, or faintness indicates a life-threatening emergency.

### **Blennorrhea**

Thick discharge containing mucus and pus from a moist body surface, especially the urethra or eye. Causes include infection and inflammation, so the affected site and other symptoms guide testing and treatment.

### **Bleomycin**

A cancer medicine that damages tumour-cell DNA and is used for selected cancers. Its most important serious risk is lung injury; fever, skin changes, low blood counts, and allergic reactions can also occur.

### **Bleomycin Sulfate**

The sulphate form of bleomycin, a cancer medicine that damages DNA. It can cause potentially permanent lung injury, so breathing symptoms and lung function may need monitoring during treatment.

### **Blepharitis**

A common, usually chronic inflammation of the eyelid margins that can cause itchy or burning eyelids, redness, tearing, dry-eye symptoms, and crusting at the base of the eyelashes. Anterior blepharitis affects the skin, lash follicles, and lash bases; posterior blepharitis affects the meibomian oil glands and often accompanies rosacea or dry eye. Lid hygiene and warm compresses may help, but pain, marked swelling, light sensitivity, or vision change needs prompt eye assessment.

---

**Alternate Names:**

Eyelid Margin Disease; Meibomian Gland Dysfunction

**Blepharoconjunctivitis**

Inflammation of the eyelid edges and the surface covering the eye. It can cause crusting, itching, redness, tearing, and irritation due to infection, allergy, dry eye, or blocked eyelid glands.

**Blepharophimosis**

Abnormally short horizontal openings between the eyelids. It may occur with drooping eyelids and other facial or developmental features and can interfere with vision.

**Blepharoplasty**

Surgery to reshape or repair an eyelid, sometimes removing or repositioning excess skin, muscle, or fat. It may improve vision blocked by heavy lids, restore an injured lid, or be performed for appearance.

**Blepharoptosis**

Drooping of the upper eyelid because of a weak muscle, damaged nerve, stretched tendon, birth difference, ageing, or a heavy eyelid. It can obstruct vision and may signal a neurological emergency if sudden.

**Blepharospasm**

Involuntary tightening of the muscles around the eyes, causing repeated blinking or forceful eyelid closure. It may occur alone or with eye irritation, medicines, or a movement disorder.

**Blind**

Having severe or complete loss of vision that substantially limits the ability to see. The cause and remaining level of vision are described because blindness varies in degree and may affect one or both eyes.

**Blind Loop Syndrome**

Poor absorption caused by excess bacteria growing in a slow-moving or bypassed part of the small

intestine. It may cause bloating, diarrhoea, weight loss, and vitamin deficiencies and can follow surgery or bowel disease.

**Blind Spot**

An area within the field of vision where sight is absent. Everyone has a normal small blind spot where the optic nerve leaves the retina, but a new or enlarged blind spot may indicate eye or nerve disease.

**Blinded**

In a research study, describing a participant, clinician, assessor, or analyst who does not know the assigned treatment or relevant result. This helps reduce expectations from influencing care or measurements.

**Blinding**

Keeping treatment assignments or study information hidden from selected participants, clinicians, assessors, or analysts to reduce expectation and measurement bias. A study states exactly who is unaware and what is concealed.

**Blindness**

Severe or complete loss of vision in one or both eyes caused by disease, injury, or a developmental condition affecting the eye, optic nerve, or brain. Sudden vision loss is an emergency.

**Blindness And Vision Loss**

Reduced or absent ability to see caused by problems in the eye, optic nerve, brain, injury, or development. It may be sudden or gradual, partial or complete, and temporary or permanent.

**Blinking**

Repeated brief closing and reopening of the eyelids that spreads tears, protects the cornea, and clears particles. It may increase with irritation, stress, tics, or vision problems and decrease during prolonged screen use.

---

## Blister

A raised pocket of fluid within or beneath the outer layer of skin. Small blisters are called vesicles; larger ones are called bullae. Blisters can result from rubbing, burns, heat, infection, allergy, or inflammatory skin disease and commonly affect the hands and feet.

## Blisters

Plural of blister; see *Blister*.

### Alternate Names:

Bulla; Vesicle When Small

## Bloating

A feeling of abdominal fullness, tightness, or visible swelling. Gas, constipation, food intolerance, or digestive disorders are common causes; persistent bloating with weight loss, vomiting, pain, or early fullness needs assessment.

## Block

An obstruction or interruption of flow, movement, electrical signals, or communication. The meaning depends on the structure involved, such as an airway, nerve, blood vessel, or heart.

## Blockage Of Upper Airway

Obstruction of airflow through the nose, mouth, throat, or voice box. It can rapidly cause suffocation; noisy breathing, inability to speak, blue skin, or severe distress requires immediate emergency help.

## Blocked Tear Duct

Partial or complete blockage of the passage that drains tears from the eye into the nose. It can cause constant watering, discharge, or repeated infection and is common in babies.

## Blocked Tear Ducts

Obstruction of the nasolacrimal drainage system, causing excessive tearing, discharge, or recurrent infection. In infants it often improves as the duct develops; persistent or painful obstruction may require ophthalmic treatment.

## Blocked Ureter

Alternate name; see *Ureteral Obstruction*.

## Blocking Agent

A substance that prevents or reduces a biological process by blocking a receptor, channel, enzyme, or other target. Its effects and risks depend on the target and the tissue affected.

### Alternate Names:

Blocker; Antagonist When It Acts by Receptor Antagonism

## Blocking Agents

Substances that reduce a body process by blocking a receptor, enzyme, ion channel, nerve signal, or other pathway. Their effects and risks vary according to the target and medicine.

## Blocking Antibody

An immune protein that prevents another molecule from attaching to its target or interferes with a biological interaction. It may be used as treatment, protect against disease, or contribute to an autoimmune disorder.

## Blood

A body fluid made of plasma, red blood cells, white blood cells, and platelets. It carries oxygen, nutrients, hormones, and waste and helps fight infection, stop bleeding, and regulate temperature.

## Blood Alcohol Concentration

The amount of alcohol in a person's blood, usually expressed as a percentage or mass per volume. Effects depend on the level, timing, tolerance, other substances, and health; one result does not fully predict impairment.

## Blood Alcohol Level

A measurement of alcohol concentration in blood. Its effects depend on the amount, timing, tolerance, other substances, and individual health, so the result must be interpreted with the person's condition.



---

## Blood Bank

A specialized medical facility and laboratory service that collects, tests, separates, and safely stores donated blood and its components (such as red blood cells, platelets, and plasma). It performs vital compatibility testing, blood typing, and antibody screening (crossmatching) before transfusions to prevent severe transfusion reactions and ensure patient safety.

### Alternate Names:

Blood Banking; Transfusion Service; Immunohematology Laboratory

## Blood Banking

Alternate name; see *Blood Bank*.

## Blood Biomarker

A measurable sign of a disease or condition that can be isolated or detected in a blood sample, such as a disease-associated antibody, hormone, protein, or other substance.

## Blood Blister

A raised pocket under the skin containing blood, usually caused by rubbing, pinching, pressure, or another injury that breaks small blood vessels.

## Blood Brain Barrier

A protective filter formed by specialized blood vessels and nearby brain cells. It controls what can pass from the bloodstream into the brain, limiting many medicines, chemicals, germs, and immune cells.

## Blood Cancer

A cancer affecting blood-forming tissue, blood cells, or the lymphatic system. Abnormal cells can crowd out normal cells, impair immunity, cause anaemia or bleeding, or spread to organs.

## Blood Cell

A cellular part of blood, including red blood cells that carry oxygen, white blood cells that help fight infection, and platelets that help stop bleeding. Most are made in the bone marrow.

## Blood Cholesterol

Cholesterol carried in the blood. The body needs it to build cell membranes and make hormones and bile acids. Too much LDL cholesterol can form fatty buildup in the arteries and increase the risk of heart disease and stroke. Blood cholesterol is usually checked as part of a lipid profile.

## Blood Cholinesterase Test

A clinical enzymatic assay measuring erythrocyte acetylcholinesterase (true cholinesterase) or plasma butyrylcholinesterase (pseudocholinesterase) activity to diagnose organophosphate or carbamate pesticide toxicity and evaluate inherited pseudocholinesterase deficiency.

### Alternate Names:

Pseudocholinesterase Test; Plasma Cholinesterase; Serum Butyrylcholinesterase

## Blood Chromium Test

A specialized trace element assay quantifying whole blood or serum chromium concentrations to assess occupational exposure, evaluate metal-on-metal orthopedic implant wear and systemic metallosis, or monitor trace element balance in total parenteral nutrition.

### Alternate Names:

Serum Chromium; Whole Blood Chromium; Chromium Exposure Test

## Blood Circulation

The continuous movement of blood from the heart through the lungs and body and back again. It delivers oxygen and nutrients and carries away carbon dioxide and other waste.

## Blood Cleaner

A method used to reduce certain infectious organisms in donated blood or blood components while keeping them suitable for transfusion. It may use chemicals or light, but it does not remove every possible infection risk.

## Blood Clot

A mass of thickened blood formed during clotting. It is protective when it seals an injury, but a clot

---

inside a blood vessel can block flow or travel to the lungs, brain, heart, or another organ.

### **Blood Clot in Lung**

Alternate name; see *Pulmonary Embolism*.

### **Blood Clots**

Clumps of thickened blood. Clots can stop bleeding, but abnormal clots in veins or arteries may block blood flow and cause deep-vein thrombosis, pulmonary embolism, stroke, heart attack, or tissue damage.

#### **Alternate Names:**

Thrombi; Thrombosis; Blood Clot

### **Blood Coagulation**

The linked process that changes blood from a liquid to a firm clot after injury. Too little clotting causes bleeding, while excessive or misplaced clotting can cause dangerous vessel blockage.

### **Blood Coagulation Disorder**

A condition in which blood clots too little, too much, or at the wrong time. Bleeding disorders result from inadequate clot formation, whereas thrombophilic disorders cause excessive or misplaced clotting. Causes include genetic variants, medicines, liver disease, immune conditions, cancer, and other illness. Examples include hemophilia, von Willebrand disease, thrombocytopenia, thrombophilia, and disseminated intravascular coagulation.

### **Blood Count**

A blood test measuring the number and features of red cells, white cells, and platelets. It helps assess anaemia, infection, inflammation, bleeding risk, and disorders of the bone marrow.

### **Blood Culture**

A test in which blood is placed in special bottles to see whether bacteria or fungi grow. It helps investigate bloodstream infection, sepsis, or infection of the heart valves.

### **Blood Culture Contamination**

Accidental entry of skin or environmental organisms into a blood-culture bottle during collection or handling, rather than true infection in the blood. It can lead to unnecessary antibiotics, so results are interpreted with symptoms and other culture sets.

### **Blood Differential Test**

A blood test measuring the numbers and proportions of different white blood cells, including neutrophils, lymphocytes, monocytes, eosinophils, and basophils. It can help investigate infection, inflammation, allergy, leukaemia, or marrow disease.

### **Blood Donation**

Voluntary collection of whole blood or selected blood components from a screened donor for transfusion or manufacture of medical products. Eligibility checks, infection testing, sterile collection, and post-donation recovery protect donor and recipient.

### **Blood Donation Before Surgery**

Giving and storing a person's own blood before an operation for possible later transfusion. It is suitable only in selected situations and can cause anaemia or delays if the operation is postponed.

### **Blood Donor**

A person who gives blood or blood components for transfusion or medical use. Screening protects the donor and helps ensure that donated blood is safe for the recipient.

### **Blood Draw**

Collection of blood, usually with a needle placed in a vein, for testing, donation, or treatment. Temporary pain, bruising, dizziness, or rarely infection may occur.

### **Blood Dyscrasia**

A nonspecific clinical term designating any abnormal condition, imbalance, or disease involving the cellular elements (erythrocytes, leukocytes, platelets) or clotting factors of the blood, such as aplastic anemia, leukopenia, or thrombocytopenia.

---

## Blood Film

A thin layer of peripheral blood spread onto a glass microscope slide, stained with a Romanowsky stain (e.g., Wright-Giemsa), and examined under microscopy for quantitative and qualitative evaluation of erythrocyte morphology, leukocyte differentials, platelet adequacy, and identification of bloodborne parasites.

### Alternate Names:

Peripheral Blood Smear; Blood Smear; Peripheral Smear

## Blood Flow

The movement of blood through vessels and organs. It depends mainly on the pressure difference and resistance in the vessels and delivers oxygen and nutrients while removing waste.

## Blood Flow Velocity

The speed at which blood moves through a vessel. It is usually fastest in large arteries and slowest in capillary networks, allowing time for exchange with tissues.

## Blood Gas Analysis

A test measuring oxygen, carbon dioxide, acidity, and related values in blood. It helps assess breathing, oxygen delivery, and acid-base balance in serious illness.

## Blood Gases

Measurements of oxygen, carbon dioxide, acidity, and related substances in blood. They help determine how well a person is breathing, exchanging gases, and maintaining acid-base balance.

## Blood Glucose

The amount of sugar called glucose in the blood. The body regulates it using food absorption, liver production, insulin, and other hormones; levels that are too low or high can cause illness.

## Blood Glucose Monitoring

The regular measurement of sugar (glucose) levels in the blood, primarily used to manage diabetes

and detect dangerously high (hyperglycemia) or low (hypoglycemia) blood sugar. Testing is commonly performed using a handheld glucometer with a small drop of blood from a fingertip prick, or with a wearable continuous glucose monitor (CGM) that checks glucose in the fluid under the skin around the clock. It tracks how meals, physical activity, illness, and medications (such as insulin) affect glucose levels throughout the day.

### Alternate Names:

Blood Sugar Monitoring; Self-Monitoring of Blood Glucose; SMBG; Continuous Glucose Monitoring; CGM

## Blood Glucose Monitors

Portable devices that measure capillary blood glucose, usually with a lancet, test strip, and meter. Correct sampling, strip storage, calibration, and interpretation are important because readings can be affected by technique, circulation, or device limitations.

## Blood Group

A classification of blood based on specific inherited markers (antigens) on the surface of red blood cells. The two most vital systems are the ABO system (types A, B, AB, and O) and the Rhesus (Rh) system (positive or negative based on the RhD marker). Accurate blood grouping and crossmatching are essential before transfusions to prevent life-threatening immune reactions and during pregnancy to prevent Rh disease (hemolytic disease of the newborn).

### Alternate Names:

Blood Type; Blood Groups; ABO and Rh Grouping; Blood Typing

## Blood Group Incompatibility

An immunological mismatch between the blood types of a donor and recipient, or between mother and fetus (such as ABO or Rh incompatibility), leading to antibody-mediated hemolysis, acute hemolytic transfusion reactions, or hemolytic disease of the newborn.

## Blood Groups

Alternate name; see *Blood Group*.

---

### Blood In Semen

Blood in the fluid released during ejaculation. It is often temporary and may follow infection, inflammation, injury, or a urinary-tract procedure. Persistent or recurrent blood, pain, fever, or urinary symptoms needs medical assessment.

#### Alternate Names:

Hematospermia; Hemospermia

### Blood in Stool

Alternate name; see *Rectal Bleeding*.

### Blood In The Eye

Bleeding on the eye surface or inside the eye. A small surface bleed may be harmless, but bleeding inside the eye, eye pain, injury, or any vision change requires prompt eye assessment.

### Blood in Urine

The abnormal presence of red blood cells in the urine, detected either macroscopically as pink, red, or tea-colored urine (gross hematuria) or microscopically on urinalysis ( $\geq 3$  erythrocytes per high-power field). Common etiologies include urinary tract infections, nephrolithiasis, glomerular nephritis, benign prostatic hyperplasia, and urothelial malignancies.

#### Alternate Names:

Hematuria; Gross Hematuria; Microscopic Hematuria; Blood in the Urine

### Blood Lancet

Alternate name; see *Lancet*.

### Blood Lead Level Test

A quantitative diagnostic laboratory assay measuring venous or capillary whole-blood lead concentration in micrograms per deciliter ( $\mu\text{g}/\text{dL}$ ). Utilized for pediatric screening, occupational exposure monitoring, and evaluating symptomatic plumbism.

#### Alternate Names:

BLL Assay; Venous Blood Lead Screen; Plumbism Blood Test

### Blood Loss

Loss of blood from the circulation because of injury, surgery, childbirth, internal bleeding, or disease. Severe loss can reduce oxygen delivery and cause shock, requiring urgent control of bleeding and replacement of blood or fluids.

### Blood Low Red Cell Count

Alternate name; see *Anemia*.

### Blood PH

A measure of how acidic or alkaline the blood is. The lungs regulate carbon dioxide and the kidneys regulate acids and bicarbonate; normal arterial pH is about 7.35–7.45.

### Blood Plasma

The liquid part of blood, made mostly of water and containing proteins, salts, nutrients, hormones, gases, and waste products. It carries blood cells and dissolved substances throughout the body.

### Blood Preservation

The controlled collection, processing, cooling, freezing, and storage of donated blood or its components so they remain safe and effective until transfusion.

### Blood Pressure

The force of circulating blood against artery walls, recorded as two numbers in millimetres of mercury. The systolic number is the pressure when the heart contracts; the diastolic number is the pressure between beats.

### Blood Pressure Cuff

The inflatable part of a blood-pressure measuring device that wraps around a limb and briefly compresses an artery. Cuffs come in different sizes and may be used with a manual sphygmomanometer or an automated monitor.

### Blood Pressure Measurement

Measurement of arterial pressure with an inflatable cuff or another approved device. The result records systolic pressure when the heart contracts and dias-

---

tolic pressure when it relaxes; correct cuff size and positioning improve accuracy.

### **Blood Pressure Monitors For Home**

Devices that measure blood pressure outside a clinic, usually with an inflatable upper-arm cuff. A validated monitor, correctly sized cuff, proper positioning, and repeated readings give more reliable results.

### **Blood Pressure Regulation**

The body's control of blood pressure through the heart, blood vessels, kidneys, hormones, and nervous system. These systems adjust heart pumping, blood-vessel width, and body-fluid balance.

### **Blood Product**

A component made from donated blood for medical treatment, such as red blood cells, platelets, plasma, or cryoprecipitate. The product is selected according to the person's bleeding, anaemia, or clotting problem.

### **Blood Products**

Components prepared from donated blood for transfusion, including red blood cells, platelets, plasma, and cryoprecipitate. Each replaces a different blood function, and the choice depends on the person's symptoms, bleeding, and laboratory results.

### **Blood Protein**

A protein dissolved in blood plasma. These proteins help transport substances, maintain fluid balance, support immunity, and control clotting; abnormal amounts can occur with inflammation, liver or kidney disease, immune disorders, or some cancers.

### **Blood Results**

Measurements obtained from a blood sample, such as cell counts, glucose, salts, hormones, or organ-function markers. They must be interpreted using the laboratory's reference range, symptoms, medicines, and other findings.

### **Blood Sedimentation**

The settling of red blood cells in a tube of treated blood. The erythrocyte sedimentation rate is a nonspecific sign that may rise with inflammation, infection, anaemia, pregnancy, or other conditions.

### **Blood Smear**

A thin layer of blood spread on a glass slide, stained, and examined under a microscope. It shows the size, shape, and appearance of blood cells and can reveal abnormal cells, parasites, or cell fragments.

### **Blood Spatter Analysis**

The forensic examination of bloodstain size, shape, and distribution to help reconstruct events. It may distinguish drops caused by gravity, contact transfer, impact, or forceful expulsion, but conclusions have limits.

### **Blood Substitutes**

Fluids or artificial products intended to replace some functions of blood. Most restore circulating volume; some are designed to carry oxygen. They do not replace every function of blood and are used only in selected situations.

### **Blood Sugar**

The amount of glucose in the blood. Insulin, glucagon, the liver, food, activity, illness, and other hormones affect it. Persistently high or very low levels can cause symptoms and serious organ injury.

### **Blood Sugar Test**

A test measuring glucose in the blood. It may be done while fasting, at a random time, or after a glucose drink, and helps diagnose or monitor diabetes and other disorders of glucose control.

### **Blood Test**

Analysis of blood cells or substances dissolved in blood, including chemicals, proteins, hormones, antibodies, or genetic material. Results help investigate symptoms, diagnose disease, assess organ function, and monitor treatment.

---

## Blood Thinner

A common name for medicines that lower the chance of harmful blood clots forming or growing. The term includes anticoagulants, which slow the body's clotting process, and antiplatelet medicines, which make platelets less likely to clump together. They do not literally thin the blood or usually dissolve an existing clot, and they can cause serious bleeding.

## Blood Transfusion

The administration of whole blood or a specific blood component, such as red blood cells, platelets, plasma, or cryoprecipitate, to treat blood loss, anaemia, thrombocytopenia, or a clotting-factor deficiency. Red cells improve oxygen-carrying capacity; platelets, plasma, and cryoprecipitate support haemostasis. ABO/Rh typing, antibody screening, and compatibility testing help reduce transfusion reactions. Risks include febrile or allergic reactions, acute haemolysis, circulatory overload, and rare transfusion-transmitted infection; the decision depends on the patient's symptoms, laboratory results, bleeding, and overall clinical context.

## Blood Typing

A laboratory test identifying markers on red blood cells, especially the ABO and Rh systems. It helps select compatible blood for transfusion and identifies some pregnancy-related risks from blood-group incompatibility.

## Blood Urea Nitrogen

The amount of nitrogen from urea in the blood. Urea is made when the body breaks down protein and is removed mainly by the kidneys; the result is interpreted with creatinine, hydration, and the clinical situation.

## Blood Urea Nitrogen Test

A clinical serum chemistry test measuring the amount of nitrogen derived from urea, the principal end product of protein catabolism formed in the liver. Elevated serum concentrations (azotemia) occur in acute or chronic renal impairment, prerenal hypoperfusion, severe dehydration, and upper gastrointestinal hemorrhage.

## Alternate Names:

BUN Test; Serum Blood Urea Nitrogen; Urea Nitrogen Test

## Blood Vessel

A tube that carries blood through the body. Arteries carry blood away from the heart, veins return it, and tiny capillaries allow oxygen, nutrients, and waste products to pass between blood and tissues.

## Blood Vessels

The arteries, arterioles, capillaries, venules, and veins that carry blood through the body. They deliver oxygen and nutrients, remove waste, and help control blood pressure, temperature, and fluid exchange.

## Blood Viscosity

How thick blood is and how much it resists flowing. It is influenced mainly by the number of red blood cells, but also by blood proteins and temperature. Excessively thick blood can slow circulation and increase clot risk.

## Blood Volume

The total amount of blood circulating in the body, including plasma and blood cells. Too little can cause low blood pressure and shock; too much can strain the heart and lungs.

## Blood-Air Barrier

The thin anatomical partition in the lung consisting of alveolar type I epithelium, fused capillary endothelial basement membrane, and capillary endothelium across which alveolar gases diffuse into and out of pulmonary blood flow.

## Alternate Names:

Alveolar-Capillary Membrane

## Blood-Borne Disease

An infection that can spread through infected blood or certain body fluids, for example through shared needles, needlestick injuries, sexual exposure, or contaminated medical equipment. Examples include HIV and hepatitis B and C.



---

## Blood-Borne Pathogens

Germs that can spread through infected blood or certain body fluids. Examples include HIV and hepatitis B and C viruses; prevention includes vaccination where available, gloves, safe sharps handling, and appropriate exposure testing.

## Blood-Brain Barrier

A protective filter formed by specialized cells lining brain blood vessels and their supporting tissues. It controls which medicines, chemicals, germs, and immune cells can move from the blood into brain tissue.

## Blood-Retinal Barrier

A physiological and anatomical barrier consisting of retinal capillary endothelial tight junctions (inner barrier) and retinal pigment epithelial tight junctions (outer barrier), regulating the microenvironment of neural retinal tissue.

### Alternate Names:

BRB

## Blood-Testis Barrier

A specialized physiological barrier formed by tight junctions between adjacent Sertoli cells near the base of seminiferous tubules, partitioning spermatogonia from developing spermatocytes and preventing autoimmune destruction of haploid germ cells.

### Alternate Names:

BTB

## Bloodborne Pathogens

Infectious agents spread through infected blood or certain body fluids. Important examples include HIV and hepatitis B and C viruses; exposure may require urgent medical assessment, testing, and preventive treatment.

## Bloom Syndrome

A rare autosomal-recessive disorder caused by pathogenic changes in both copies of the *BLM* gene, which encodes a DNA helicase. It causes marked short stature, a sun-sensitive telangiectatic facial rash, recurrent infections, and a greatly increased risk of multiple cancers, often at unusually young

ages; specialist surveillance and strong sun protection are important.

## Blount Disease

A growth disorder in which the upper part of the shinbone grows unevenly, causing worsening bowing of one or both legs. It can affect young children or adolescents and may cause an abnormal gait, unequal leg length, or knee damage.

## Blue Baby

An infant whose skin, lips, or nail beds look blue because the blood carries too little oxygen or circulation is inadequate. This can signal heart, lung, airway, blood, or other serious disease and needs urgent assessment.

## Blue Bottle

A common name for a bluebottle jellyfish. Its tentacles can inject venom that causes immediate stinging pain, raised marks, redness, and sometimes muscle cramps, breathing difficulty, or a severe allergic reaction.

## Blue Discoloration Of The Skin

A blue or gray tint of the skin, lips, or nail beds caused by low oxygen, reduced blood flow, or abnormal haemoglobin. Sudden or widespread discoloration, especially with breathing difficulty or confusion, is an emergency.

## Blue Nevus

Usually benign melanocytic lesions formed by pigment-producing cells deep in the skin, giving them a blue, blue-gray, or blue-black colour. Common blue nevi are usually small and stable; cellular blue nevi may be larger or deeper and rarely undergo malignant change. A new, enlarging, bleeding, or changing lesion requires examination and sometimes biopsy.

## Blue Nevus

A usually harmless blue-gray or blue-black mole caused by pigment-producing cells located deeper than those in most moles. A new, changing, bleeding, or painful lesion requires examination.

---

### Blue Nightshade Poisoning

Illness caused by eating blue nightshade or a related poisonous plant containing glycoalkaloids. Symptoms may include vomiting, diarrhoea, abdominal pain, confusion, weakness, seizures, or abnormal heart rhythm; it is an emergency requiring urgent medical treatment.

### Blue NæVus

A usually harmless blue-gray or blue-black mole formed by pigment-producing cells deep in the skin. It requires assessment if it changes in size, shape, colour, or symptoms.

### Blue Rubber Bleb Nevus Syndrome

A rare disorder with multiple slow-flow venous malformations, typically soft blue or purple compressible skin nodules and lesions in the gastrointestinal tract. Gastrointestinal bleeding can cause iron-deficiency anaemia, and other organs may be involved; evaluation may include imaging, endoscopy, and blood counts.

### Blue Sclera

A blue-gray appearance of the normally white outer layer of the eye, usually caused by thinning that allows deeper pigment to show through. It can occur in inherited connective-tissue or bone disorders and needs clinical assessment.

### Blue Sclerae

A distinct bluish or grayish discoloration of the ocular sclera caused by abnormal thinness or translucency of the scleral collagen fibers, which allows the pigment of the underlying uveal tract and choroid to be visible. It is a hallmark physical finding of osteogenesis imperfecta and can also occur in Ehlers-Danlos syndrome, Marfan syndrome, and severe iron deficiency anemia.

### Blue Toe Syndrome

The acute onset of localized cyanosis, purplish discoloration, and painful ischemia of one or more toes in the presence of intact distal pulses, caused by microembolization of atheromatous debris or small cholesterol crystals from proximal arterial sources.

### Alternate Names:

Trash Foot; Digital Atheroembolism

### Blue-Green Algae

Blue-green algae are cyanobacteria that can grow rapidly in water. Some release toxins that cause skin irritation, vomiting, liver injury, breathing problems, or neurologic illness when people or animals are exposed.

### Blumberg Sign

Sudden sharp pain when a doctor presses down gently on the belly and then quickly lets go, showing that the lining of the abdomen is swollen and irritated.

### Blunt Abdominal Trauma

An injury to the abdomen without an object piercing the skin, often caused by a crash, fall, or blow. It can damage the spleen, liver, kidneys, bowel, or pancreas, sometimes without early visible bruising.

### Blunt Force Trauma

Injury caused by a blow from a blunt object or by striking a blunt surface, without penetration by a sharp edge. It can produce bruises, grazes, torn skin, internal bleeding, fractures, or organ damage.

### Blunted Affect

Reduced outward expression of emotion. A person may show less facial movement, voice variation, or body language than expected, although their internal feelings may be stronger. It can occur in mental or neurologic disorders.

### Blurred Vision

Vision that appears out of focus, hazy, or unclear. Causes include an incorrect glasses prescription, dry eye, eye disease, medicines, migraine, or neurologic illness. Sudden blurring, pain, weakness, or vision loss needs urgent assessment.

### Blush

Temporary facial redness or warmth caused by increased blood flow near the skin surface. It

---

may follow emotion, heat, exercise, fever, alcohol, medicines, hormonal changes, or some medical conditions.

### **Blushing**

Temporary redness or warmth of the face caused by increased blood flow in the skin. Common triggers include emotion, heat, exercise, alcohol, hormonal changes, medicines, fever, and some illnesses.

### **Board-Like Rigidity**

Severe involuntary tightening of the abdominal muscles, making the abdomen feel hard. It can indicate irritation of the abdominal lining from serious inflammation, infection, bleeding, or perforation and requires urgent assessment.

### **Boas' Sign**

Pain or unusual skin sensitivity below the right shoulder blade, traditionally associated with gallbladder disease. It is not specific enough to establish a diagnosis and must be interpreted with other findings.

### **Bobble-Head Doll Syndrome**

A rare childhood movement disorder causing repeated forward-and-back head movements. It is often linked to a fluid-filled cyst or other lesion near the third ventricle of the brain and requires neurologic evaluation and imaging.

### **Bocavirus**

A group of small viruses that can infect humans and animals. Human bocavirus most often affects the airways of young children, although infection may cause illness in other tissues or people with weakened immunity.

### **Bocavirus Infection**

A viral infection, usually of the airways in young children. It may cause a runny nose, cough, fever, wheezing, bronchiolitis, or pneumonia, often together with another infection. Treatment is usually supportive; breathing difficulty or dehydration needs prompt care.

### **Bochdalek Hernia**

A birth defect in which abdominal organs move through an opening in the back and side of the diaphragm into the chest. It can prevent normal lung development and cause severe breathing problems in a newborn.

### **Bock's Disease**

An older name for sarcoidosis, an inflammatory disease in which clusters of immune cells form in organs. It most often affects the lungs and lymph nodes but can also affect the skin, eyes, heart, or nervous system.

### **BODE Index**

A scoring tool for chronic obstructive pulmonary disease that combines body weight, breathing-test results, breathlessness, and walking distance. A higher score suggests more severe disease and a greater risk of hospital admission or death; it does not predict an individual outcome with certainty.

#### **Alternate Names:**

BODE Score; COPD Multidimensional Index

### **Bodily Secretion**

A substance made and released by cells or glands, such as sweat, saliva, mucus, tears, digestive juices, or hormones. Secretions protect tissues, aid digestion, regulate body functions, or help remove waste.

### **Body**

The physical organism, including its cells, tissues, organs, and interacting body systems. Health depends on the structure and function of these parts and how they respond to internal and external conditions.

### **Body Aches**

Widespread muscle or body pain that can occur with infection, fever, exercise, inflammation, medicines, or other illness. Severe pain with weakness, dark urine, breathing difficulty, chest pain, or high fever needs prompt assessment.

---

## Body Cavity

An enclosed space inside the body that contains and protects organs. Major cavities include the skull, spinal canal, chest, abdomen, and pelvis; their linings may produce small amounts of lubricating fluid.

## Body Cell Mass

The living, metabolically active cells of the body, including cells in muscles, organs, blood, and other tissues. It excludes stored fat and water outside cells and is sometimes estimated in nutrition assessment.

## Body Composition

The relative amounts of fat, muscle and other lean tissue, bone mineral, and water in the body. It provides more information about health and nutrition than body weight alone, although estimates vary by method.

## Body Dysmorphia

Alternate name; see *Body Dysmorphic Disorder*.

## Body Dysmorphic Disorder

A mental-health disorder involving persistent distress about one or more perceived flaws in appearance that are minor or not noticeable to others. Repetitive checking, grooming, comparison, or avoidance interferes with daily life.

### Alternate Names:

BDD

## Body Fat

Adipose tissue that stores energy, cushions organs, insulates the body, and releases hormones and other signals. Too much or too little body fat can affect metabolism, movement, fertility, immunity, and other aspects of health.

## Body Fluid

The water and dissolved substances (such as electrolytes, nutrients, proteins, and gases) that make up approximately 60% of human body weight. Body fluids are organized into two major compartments: intracellular fluid (fluid inside cells, making up

about two-thirds) and extracellular fluid (fluid outside cells, including interstitial fluid between tissues and blood plasma), along with specialized fluids such as cerebrospinal fluid, synovial joint fluid, lymph, and digestive secretions. They are essential for transporting nutrients and waste, regulating body temperature, and lubricating tissues and joints.

### Alternate Names:

Body Fluids; Bodily Fluids; Total Body Water; Body Fluid Compartments

## Body Fluid Exposures

Occupational or accidental contact with blood or potentially infectious body fluids through the eyes, mouth, broken skin, or a penetrating injury. Risk assessment determines testing, wound care, vaccination, and postexposure prophylaxis.

## Body Fluids

Alternate name; see *Body Fluid*.

## Body Habitus

A person's general build, including body size, shape, muscle, fat distribution, and proportions. Clinicians may record it during an examination, but it does not by itself diagnose disease or predict health.

## Body Height

The vertical distance from the soles of the feet to the top of the head, measured while standing when possible. It helps assess growth and nutrition and is used in medication, kidney-function, and other calculations.

## Body Identification

The process of establishing who a deceased person is. It may use fingerprints, dental records, medical information, DNA, personal effects, or visual recognition, with the method chosen according to the circumstances.

## Body Image

A person's thoughts, feelings, and beliefs about their body and appearance. Distress or inaccurate beliefs about appearance can occur in body dysmor-

---

phic disorder, eating disorders, depression, or other mental-health conditions.

### Body Lice

Infestation with human body lice, which live mainly in clothing and move to the skin to feed. Bites cause intense itching and scratching; treatment includes washing or heat-treating clothing and bedding and prescribed insecticide when needed.

#### Alternate Names:

Pediculosis Corporis; Body Louse Infestation

### Body Mass Index

A screening measure calculated by dividing weight in kilograms by height in meters squared. It helps assess weight in relation to height, but it does not directly measure body fat or provide a complete assessment of health. Results are interpreted with age and other clinical factors; children and teens are assessed using age- and sex-specific percentiles.

#### Alternate Names:

BMI; Quetelet Index; BMI Calculation

### Body Odor

Smell produced when skin secretions, especially sweat, are broken down by skin microorganisms. A sudden or unusual change may be related to diet, medicines, infection, or a metabolic disorder.

### Body Odor and Sweating

Alternate index label; see *Sweating and Body Odor*.

### Body Odour

Smell from the breakdown of skin secretions by microorganisms. The smell can change with sweating, hygiene, diet, medicines, infection, or some metabolic diseases.

### Body Of The Epididymis

The middle part of the epididymis, where sperm continue to mature as they move from the head toward the tail.

### Body Of The Sternum

The broad middle part of the breastbone between

the manubrium and xiphoid process. It joins most of the ribs through their costal cartilages.

### Body Piercing

Creation of an opening through skin or a mucous membrane to insert jewellery. Possible complications include pain, bleeding, infection, scarring, allergic reaction, and damage to teeth, nerves, or cartilage.

### Body Plethysmography

A breathing test performed inside a sealed cabin to measure lung volumes, including air that cannot be exhaled. It helps identify restricted lungs, over-expanded lungs, and some airway problems.

### Body Position

The arrangement of the body or a body part, such as standing, lying on the back, lying face down, or lying on the side. Position can affect breathing, circulation, examination results, and medical procedures.

### Body Region

A named area of the body used to describe the location of an organ, symptom, injury, or examination finding, such as the chest, abdomen, pelvis, or arm.

### Body Ringworm

A fungal infection of the skin on the body. It usually causes an expanding, ring-shaped, scaly rash with a more active edge and clearer centre; it spreads by contact and is treatable with antifungal medicine.

#### Alternate Names:

Tinea Corporis; Dermatophytosis of Glabrous Skin

### Body Space

A normal or potential compartment inside the body, such as the space around the lungs or abdominal organs. Air, blood, fluid, or infection can collect there and interfere with organ function.

### Body Substance

Material produced by or found in the body, including blood, saliva, urine, faeces, mucus, tissue, and

---

other secretions. Some substances can carry infection and require appropriate handling precautions.

### **Body Surface**

The outside of the body, mainly the skin and accessible mucous membranes. Clinicians may estimate its area when assessing burns, calculating some medicine doses, or making physiological measurements.

### **Body Surface Area**

An estimate of the body's external area, usually calculated from height and weight. It is used in burn assessment, some medicine doses, and selected medical calculations; it is an estimate rather than a direct measurement.

### **Body Temperature**

The degree of heat in the body, determined by the balance between heat production and heat loss. It varies with age, activity, time of day, illness, and measurement site; fever and hypothermia require clinical interpretation.

### **Body Temperature Norms**

Expected temperature ranges vary with age, time of day, activity, illness, and whether temperature is measured by mouth, ear, armpit, skin, or rectum. The measurement method and symptoms are considered when interpreting fever or low temperature.

### **Body Wall**

The layers of skin, connective tissue, muscle, and sometimes bone that surround and protect a body compartment, such as the chest or abdomen. They also help movement, breathing, and support.

### **Body Weight**

The measured mass of a person, including fat, muscle, bone, water, and organs. It helps assess growth, nutrition, and fluid changes and is used with other information for some medical calculations.

### **Boeck's Sarcoid**

An older name for sarcoidosis, an inflammatory

disease in which clusters of immune cells form in tissues. It commonly affects the lungs and nearby lymph nodes but may also involve the skin, eyes, heart, or nervous system.

### **Boerhaave Syndrome**

A sudden full-thickness tear of the food pipe, usually after forceful vomiting. Leakage into the chest can cause severe chest or upper-abdominal pain, infection, shock, and breathing problems; it is a surgical emergency.

### **Boerhaave's Syndrome**

A spontaneous full-thickness tear of the food pipe, most often following forceful vomiting. Contents can leak into the chest and cause severe infection, chest pain, breathing difficulty, and shock.

### **Bogros' Space**

A layer of tissue behind the groin region and in front of the abdominal lining. Surgeons use this natural plane when repairing some groin hernias.

### **Bohn Nodules**

Small, harmless cysts containing keratin that appear on a newborn's gums or the roof of the mouth. They usually disappear without treatment and do not interfere with feeding.

### **Bohr Effect**

The tendency of haemoglobin to release more oxygen when tissues become more acidic or contain more carbon dioxide. This helps active tissues receive oxygen where energy use and carbon-dioxide production are greatest.

### **Bohring Syndrome**

A rare genetic disorder associated with severe developmental delay, poor growth before birth, distinctive facial features, reduced movement, and frequent feeding or breathing difficulties.

### **Boil**

A painful, pus-filled skin abscess centered on an infected hair follicle and surrounding tissue, most



---

commonly caused by *Staphylococcus aureus* bacteria. It typically begins as a tender, red nodule that progressively enlarges, develops a central yellowish head of pus, and eventually ruptures to drain. Multiple or recurrent boils are termed furunculosis, whereas a deeper, connected cluster of several boils is classified as a *Carbuncle*.

**Alternate Names:**

Furuncle; Furunculosis; Boils; Skin Abscess

**Boils**

Alternate name; see *Boil*.

**Boils and Carbuncles**

Alternate name; see *Boil* and *Carbuncle*.

**Bolt Device**

A bolt is a threaded fastener used to hold parts together. In medicine, the term may describe a device component or a specific product name, so its meaning depends on the clinical context.

**Bolus**

A single measured amount of medicine or fluid given over a short time, usually by injection or infusion. In digestion, it means food that has been chewed and is ready to swallow.

**Bolus Gastrostomy Feeding**

The administration of prescribed liquid enteral formula or hydration directly into the stomach through a gastrostomy tube over a short period using a syringe or gravity drip. Involves tube flushing before and after each feeding to maintain patency.

**Alternate Names:**

Bolus G-Tube Feeding; Bolus Enteral Nutrition; Syringe Feeding

**Alternate Names:**

G-Tube Bolus Feeding; Enteral Bolus Nutrition

**Bolus Infusion**

Delivery of a measured amount of fluid or medicine into a vein over a short period. The dose and speed depend on the medicine, the person's condition, and the intended effect; errors can cause serious harm.

**Bombay Blood Group**

A rare inherited blood type in which red blood cells lack the H marker needed to form A or B markers. It may look like type O but is usually compatible only with blood from another Bombay-type donor.

**Bombesin Receptor**

A protein on or inside a cell that binds bombesin-like signalling substances and changes cell activity. These receptors help regulate processes such as hormone release, digestion, and cell growth and are studied in some cancers.

**Bone**

A strong living tissue that forms the skeleton, supports the body, protects organs, enables movement with muscles, stores minerals, and contains marrow that makes blood cells. Bone is continually broken down and rebuilt.

**Bone And Mineral Disorders**

Diseases affecting bone strength, bone formation, or the body's balance of calcium and phosphate. Examples include osteoporosis, rickets, osteomalacia, parathyroid disorders, and inherited bone diseases.

**Bone Callus**

New tissue that forms around a broken bone during healing. It first consists of a soft blood clot and connective tissue, then cartilage and immature bone, which are gradually reshaped into stronger bone.

**Bone Cancer**

Cancer that starts in bone, such as osteosarcoma or chondrosarcoma, or cancer that has spread there from another organ. Persistent focal pain, swelling, or a fracture after minor injury may occur; diagnosis uses imaging and tissue testing.

**Alternate Names:**

Bone Sarcoma; Primary Bone Malignancy

**Bone Cement**

A self-curing acrylic biomaterial composed of polymethylmethacrylate (PMMA) utilized in orthopedic and spine surgery to anchor prosthetic joint

---

components, fill osseous voids, and stabilize vertebral compression fractures during vertebroplasty or kyphoplasty.

**Alternate Names:**

Polymethylmethacrylate; PMMA; Acrylic Bone Cement

**Bone Conduction**

Hearing sound through vibrations carried by the skull to the inner ear, bypassing the outer and middle ear. Bone-conduction testing helps identify whether hearing loss comes from the outer or middle ear or from the inner ear and nerve.

**Bone Cyst**

A fluid-filled or blood-filled space within bone. Many bone cysts are benign, but they can weaken the bone and cause pain, swelling, or fracture. Common types include simple bone cysts and aneurysmal bone cysts.

**Alternate Names:**

Bone Cysts

**Bone Densitometry**

See *Bone Density Screening*.

**Bone Density**

The amount of mineral packed into a measured area of bone. It is usually estimated with a DXA scan; lower density is associated with greater fracture risk, but results must be interpreted with age and other risk factors.

**Bone Density Screening**

A DXA scan that measures bone mineral at sites such as the hip and spine. It helps identify osteoporosis and estimate fracture risk, especially in people with age-related or other risk factors.

**Bone Development**

The formation, growth, and shaping of bones from before birth through adulthood. Bones lengthen and widen as new tissue is made and old tissue is removed; hormones, nutrition, activity, and injury influence this process.

**Bone Disease**

Any condition that weakens, deforms, inflames, infects, or otherwise damages bone. Examples include osteoporosis, fractures, infections, tumours, inherited skeletal disorders, and diseases caused by abnormal mineral or hormone levels.

**Bone Fracture**

A partial or complete break in a bone caused by injury, repeated stress, or weakened bone. Pain, swelling, bruising, deformity, and loss of function may occur. An open fracture or loss of sensation or circulation needs emergency care.

**Bone Fractures**

Alternate name; see *Bone Fracture*.

**Bone Graft**

Bone tissue or a substitute placed in a bone defect to support repair, join bones together, rebuild an area, or help a fracture heal. The material may come from the person, a donor, or a synthetic source.

**Bone Health**

The strength, structure, and ability of bones to grow, repair, and resist fractures. It is influenced by age, nutrition, hormones, physical activity, medicines, smoking, alcohol, and medical conditions.

**Bone Infarction**

Death of bone and bone-marrow tissue caused by loss of blood supply. The term often refers to an affected area in the middle or shaft of a long bone; it is related to osteonecrosis, which can occur in different parts of bone. Some bone infarcts cause no symptoms and are found on imaging done for another reason.

**Alternate Names:**

Bone Infarct

**Bone Infection**

Infection of bone, usually caused by bacteria. It may reach bone through the bloodstream, spread from nearby tissue, or enter through surgery or an open injury, causing pain, swelling, fever, or poor

---

healing.

### **Bone Lesion Biopsy**

Removal of a small sample from an abnormal area of bone with a needle or operation for laboratory examination. Testing can distinguish infection, non-cancerous growth, and cancer; the route must be planned with any later surgery in mind.

### **Bone Markers in Osteoporosis**

Blood or urine measurements reflecting bone formation or resorption. They can help assess turnover and response or adherence to therapy in selected patients, but they do not replace fracture-risk assessment or bone-density testing.

### **Bone Marrow**

Soft tissue inside many bones. Red marrow makes red cells, white cells, and platelets; yellow marrow stores fat and can produce more blood cells when the body needs them.

### **Bone Marrow Aspiration**

Removal of a small amount of liquid marrow, usually from the back of the hipbone, with a needle. It is examined to assess blood-cell development, abnormal cells, infection, or genetic changes.

### **Bone Marrow Aspiration And Biopsy**

Two ways of sampling marrow, usually from the back of the hipbone. Aspiration collects liquid marrow for cell and genetic testing; biopsy removes a small solid core to assess marrow structure, cell production, scarring, infection, or cancer.

### **Bone Marrow Biopsy**

A procedure that removes marrow, usually from the back of the hipbone, for examination. It may include liquid aspiration and a solid core to assess blood-cell production, abnormal cells, scarring, infection, or cancer.

### **Bone Marrow Culture**

A laboratory test in which marrow is placed in special conditions to grow and identify bacteria,

fungi, or other organisms when an infection of the marrow or bloodstream is suspected.

### **Bone Marrow Depression**

Reduced production of blood cells by the marrow. It can cause anaemia, increased infection risk from too few white cells, bleeding from too few platelets, or a combination of these problems.

### **Bone Marrow Disorder**

Any disease that affects the marrow's production of red cells, white cells, or platelets. It may cause anaemia, frequent infections, bleeding, abnormal blood counts, or symptoms from abnormal marrow cells.

### **Bone Marrow Donation**

The harvesting of hematopoietic stem and progenitor cells (HSPCs) from an allogeneic or autologous donor via surgical iliac crest bone marrow aspiration or apheresis of granulocyte colony-stimulating factor (G-CSF)-mobilized peripheral blood stem cells for hematopoietic stem cell transplantation.

#### **Alternate Names:**

Hematopoietic Stem Cell Donation; Stem Cell Donation; Bone Marrow Harvest

### **Bone Marrow Failure**

A condition in which the marrow cannot make enough red cells, white cells, or platelets. Causes include inherited disorders, aplastic anaemia, medicines, toxins, infections, marrow cancers, and other diseases.

#### **Alternate Names:**

Marrow Failure Bone Marrow Failure Syndrome

### **Bone Marrow Hyperplasia**

An increase in the number of marrow cells because blood-cell production has accelerated. It may be a response to ongoing red-cell destruction, blood loss, low oxygen, or another increased demand for blood cells.

### **Bone Marrow Necrosis**

Death of the blood-forming tissue inside bones. It

---

can occur with severe sickle-cell disease, blood cancers, or infection and may cause severe bone pain, anaemia, low white-cell or platelet counts, and fever.

### **Bone Marrow Transplantation**

A treatment in which healthy blood-forming stem cells are infused to replace or restore bone-marrow cells damaged by disease or treatment. The cells may come from the patient or a donor. It is used for selected blood cancers, marrow failure, and some inherited disorders; risks include infection, graft failure, organ toxicity, and graft-versus-host disease after a donor transplant.

#### **Alternate Names:**

BMT; Bone Marrow Transplant

### **Bone Mass**

The amount of bone tissue in the skeleton or in a particular area. Greater bone mass generally provides more strength, while low bone mass can increase the risk of osteoporosis and fractures.

#### **Alternate Names:**

Bone Mineral Content; Skeletal Bone Mass

### **Bone Matrix**

The material surrounding bone cells that gives bone its shape and strength. It contains a flexible framework, mainly type I collagen, reinforced with mineral crystals made largely from calcium and phosphate.

### **Bone Metastases**

Areas of cancer that have spread to bone from another part of the body. They may cause bone pain, weaken the bone and lead to fracture, raise blood calcium, or press on the spinal cord or nerves.

#### **Alternate Names:**

Metastatic Bone Disease; Skeletal Metastases

### **Bone Metastasis**

See *Bone Metastases*.

### **Bone Mineral Density**

The amount of mineral, mainly calcium, in a measured area of bone. A DXA scan estimates bone strength, helps diagnose osteoporosis, assesses frac-

ture risk, and monitors treatment together with age and other risk factors.

#### **Alternate Names:**

BMD; Bone Density; Bone Mineral Content

### **Bone Mineral Density Test**

See *Bone Density Screening*.

### **Bone Morphogenetic Protein**

A family of growth factors belonging to the transforming growth factor-beta (TGF-beta) superfamily that orchestrate osteogenesis, chondrogenesis, and bone remodeling, utilized therapeutically in spinal fusion and non-union fracture repair.

#### **Alternate Names:**

BMP

### **Bone Necrosis**

Death of bone tissue, usually because its blood supply is reduced or because of severe injury, infection, radiation, or prolonged corticosteroid use. It can cause pain, bone collapse, joint damage, and fracture.

### **Bone Neoplasm**

An abnormal growth of cells arising in bone. It may be noncancerous or cancerous; imaging and often a biopsy determine its type, behaviour, extent, and appropriate treatment.

### **Bone Pain**

Aching or other discomfort arising from bone. Causes include injury, fracture, infection, reduced blood supply, metabolic disease, overuse, or cancer. Persistent, severe, or unexplained pain requires medical assessment.

### **Bone Pain Or Tenderness**

Pain when a bone is used, pressed, or touched. It may result from injury, fracture, infection, inflammation, metabolic disease, or cancer; persistent or unexplained tenderness needs medical assessment.

### **Bone Plate**

A metal or other body-compatible plate fixed to

---

bone with screws. It holds broken pieces in position, supports healing, or helps correct a deformity during orthopaedic surgery.

### **Bone Regeneration**

Formation of new bone to repair or replace bone that is damaged, missing, or diseased. It occurs naturally during fracture healing and may be supported by grafts, implants, or other treatments.

### **Bone Remodeling**

The lifelong replacement of old bone with new bone. Cells remove worn bone and other cells build replacement bone, helping maintain strength, shape, and the body's calcium balance.

### **Bone Resorption**

The removal and breakdown of bone tissue, which releases minerals into the blood. It is normal during bone renewal, but excessive resorption can weaken bones and contribute to osteoporosis.

### **Bone Sarcoma**

Alternate name; see *Bone cancer*.

### **Bone Saw**

A surgical instrument used to cut bone during procedures such as amputation, joint replacement, fracture repair, or orthopaedic reconstruction.

### **Bone Scan**

A nuclear medicine diagnostic imaging technique in which technetium-99m-labeled bisphosphonates (e.g., <sup>99m</sup>Tc-MDP) are administered intravenously and imaged with a gamma camera to identify areas of osteoblastic activity, osteomyelitis, occult fractures, or skeletal metastases.

#### **Alternate Names:**

Bone Scintigraphy; Skeletal Scintigraphy; Radionuclide Bone Scan

### **Bone Screws**

Screws used during orthopaedic surgery to hold broken bone pieces together or attach plates, joint components, or other implants. Their size, mate-

rial, and placement depend on the bone and the operation.

### **Bone Spurs**

Extra projections of bone that usually form near joints after long-term wear, arthritis, injury, or abnormal stress. They may cause no symptoms or may irritate nearby tissues and restrict movement.

#### **Alternate Names:**

Osteophytes

### **Bone Transplantation**

Transfer of bone tissue or a bone substitute to repair a defect, support fracture healing, or assist reconstruction of the spine, jaw, or a joint. The graft may come from the person, a donor, or a manufactured material.

### **Bone Tumor**

An abnormal mass that starts in bone. It may be noncancerous or cancerous; examination, scans, and often a biopsy determine its type, whether it is growing or spreading, and the best treatment.

### **Bone Tumor-Epidermoid Cyst-Polyposis**

A historical descriptive name for Gardner syndrome, an inherited adenomatous polyposis condition associated with numerous colorectal polyps, osteomas, epidermoid cysts, and a high risk of colorectal cancer.

### **Bone Turnover Markers**

Substances measured in blood or urine that reflect how quickly bone is being built or broken down. They may support assessment of some metabolic bone disorders or treatment response but do not replace bone-density testing.

### **Bone Type**

A way of grouping bones by shape. Long bones support movement, short bones provide stability, flat bones protect organs or provide broad muscle attachment, irregular bones have complex shapes, and sesamoid bones lie within tendons.

---

## **Bone X-Ray**

An X-ray picture of a bone or group of bones used to look for fractures, abnormal shape, arthritis, infection, bone loss, or some tumours.

## **Bone-Anchored Hearing Implant**

A hearing device fixed to the skull bone that transmits sound vibrations to the inner ear, bypassing the outer or middle ear. It may be used for selected conductive, mixed, or single-sided hearing loss.

## **Bone-Marrow Transplantation**

A treatment that replaces damaged or diseased blood-forming marrow with healthy stem cells from the patient or a donor. It is used for selected blood cancers, marrow failure, and some inherited disorders.

## **Bones And Joints**

Bones provide support and protection, while joints connect bones and allow movement. Joints may be fixed, slightly movable, or freely movable and are held together by tissues such as cartilage, ligaments, and joint fluid.

## **Bones of the Skeleton**

The body's framework of bones and cartilage, providing support, protection, leverage for movement, mineral storage, and marrow-based blood-cell production.

## **Bongkreik Poisoning**

Severe poisoning caused by bongkreikic acid in contaminated fermented foods, especially coconut or corn products. It can cause vomiting, abdominal pain, weakness, liver failure, neurologic injury, and death; urgent treatment is required.

## **Bonnevie-Ullrich Syndrome**

An older name for the physical features sometimes associated with Turner syndrome, a condition involving an absent or altered X chromosome. Features may include short stature, a webbed neck, and reduced ovarian function.

## **Bony Exostosis**

A localized extra growth of bone. It may occur on the jaw or palate and is often harmless, but removal may be considered if it repeatedly causes injury, interferes with speech or dentures, or needs diagnosis.

## **Bony Labyrinth**

A system of channels inside the dense part of the skull bone that houses the inner ear. It contains the cochlea for hearing and the vestibule and semi-circular canals for balance.

## **Borborygmi**

Rumbling or gurgling sounds made when gas and liquid move through the intestines. They are often normal, but major changes together with pain, vomiting, swelling, or inability to pass stool may signal bowel disease.

## **Borborygmus**

A single rumbling or gurgling intestinal sound caused by movement of gas and liquid through the bowel. Its significance depends on symptoms and the overall abdominal examination.

## **Border Definition**

The sharpness and shape of the edge of a skin lesion or abnormal growth. A poorly defined or irregular edge can be concerning, but the border alone cannot distinguish harmless from cancerous disease.

## **Borderline**

Describes a result or condition near the boundary between normal and abnormal or between diagnostic categories. It usually requires interpretation with symptoms, repeat testing, or other findings; the meaning depends on the clinical context.

## **Borderline Ovarian Cancer**

An ovarian epithelial tumor with atypical proliferating cells but no destructive stromal invasion on examination. It generally behaves less aggressively than invasive ovarian cancer, but can recur and requires appropriate staging and long-term follow-up.



---

## **Borderline Personality Disorder**

A long-term mental-health condition involving unstable emotions, self-image, relationships, and behavior. It may include intense fear of abandonment, impulsive actions, self-harm, rapidly changing moods, and difficulty managing feelings.

### **Alternate Names:**

BPD; Emotionally Unstable Personality Disorder (EUPD)

## **Bordetella**

A group of small bacteria that can infect the airways of humans or animals. *Bordetella pertussis* causes whooping cough; other species usually affect animals but can rarely cause opportunistic human infection.

### ***Bordetella Avium***

A bacterium that mainly causes respiratory disease in turkeys and other birds. Human infection is very rare and is usually reported in people with weakened immunity or serious underlying illness.

### ***Bordetella Bronchiseptica***

A bacterium that commonly causes respiratory infection in dogs and other animals. It rarely infects humans, but may cause pneumonia or bloodstream infection, especially in people with weakened immunity.

### ***Bordetella Hinzii***

A bacterium found mainly in poultry. It rarely causes respiratory or bloodstream infection in humans, usually in people with weakened immunity or significant underlying disease.

### ***Bordetella Parapertussis***

A bacterium that can cause a whooping-cough-like respiratory infection. Illness is often milder than infection with *Bordetella pertussis*, but prolonged coughing can still occur and the infection can spread to others.

### ***Bordetella Pertussis***

The bacterium that causes whooping cough, a highly contagious respiratory infection. It attaches to air-

way lining, produces toxins that disrupt airway function, and can cause repeated coughing, vomiting, breathing pauses, and serious illness in infants.

## ***Bordetella Petrii***

An environmental bacterium that rarely causes human disease. When it is found in a specimen, clinicians decide whether it represents true infection by considering the body site, symptoms, and other test results.

## **Boric Acid**

A chemical used in some industrial products, insecticides, and limited topical preparations. Swallowing it or absorbing a large amount can poison the body; significant exposure requires prompt medical assessment.

## **Boric Acid Poisoning**

Illness caused by swallowing or absorbing too much boric acid. It may cause vomiting, diarrhoea, abdominal pain, skin changes, kidney injury, seizures, shock, or death and requires urgent poison or emergency care.

## **Borna Disease**

An uncommon viral infection of the nervous system caused by Borna disease virus. It is mainly recognized in animals; human infection is rare and may cause severe inflammation of the brain.

## **Bornholm Disease**

A contagious viral illness causing sudden attacks of severe chest or upper-abdominal muscle pain, often with fever, headache, or vomiting. It is usually caused by an enterovirus and generally resolves with supportive care.

## **Boron**

A chemical element found naturally in food, water, soil, and industrial materials. Small dietary amounts are generally tolerated, but excessive exposure can cause digestive symptoms, skin problems, reproductive effects, or developmental toxicity.

---

## **Borrelia**

A group of spiral-shaped bacteria spread mainly by ticks or lice. Different species cause Lyme disease, relapsing fever, or other infections, with symptoms and treatment depending on the species and exposure.

## **Borrelia Burgdorferi**

A spiral-shaped bacterium spread by infected black-legged ticks and the main cause of Lyme disease in North America and parts of Europe. Infection may affect the skin, joints, nervous system, or heart.

## **Borrelia Infection**

Infection with a *Borrelia* bacterium. Depending on the species, it may cause Lyme disease or relapsing fever, with symptoms such as fever, headache, fatigue, rash, muscle pain, or joint and nerve problems.

## **Borrelia Miyamotoi**

A bacterium spread by some hard-bodied ticks that causes hard-tick relapsing fever. Illness commonly includes repeated episodes of fever, headache, fatigue, chills, and muscle or joint aches.

## **Borreliosis**

Infection caused by a *Borrelia* bacterium. Human forms include Lyme disease and relapsing fever; symptoms differ according to the species, stage of infection, route of spread, and organs involved.

## **Bortezomib**

A cancer medicine that blocks proteasomes, cell structures that remove unwanted proteins. This can kill susceptible cancer cells and is used mainly for multiple myeloma and some lymphomas; nerve damage, low blood counts, digestive symptoms, and infections can occur.

## **Bosentan**

A medicine that blocks endothelin, a substance that narrows blood vessels. It is used for pulmonary arterial hypertension, but can injure the liver and harm a developing fetus, so liver tests and pregnancy prevention are important.

## **Bosniak Classification**

A system for describing kidney cysts on contrast CT or MRI according to their walls, septations, mineral deposits, and solid parts. Categories help estimate cancer risk and guide monitoring, further imaging, or surgery.

## **Boston Criteria**

Clinical and brain-imaging rules used to estimate the likelihood of cerebral amyloid angiopathy in living people. Age, symptoms, and MRI findings such as bleeding in certain parts of the brain are considered. The criteria classify cases as possible or probable; definite diagnosis generally requires examination of brain tissue.

### **Alternate Names:**

Boston Criteria for Cerebral Amyloid Angiopathy

## **Bot Fly**

A fly whose larvae can live in animal or human tissue, causing myiasis. In people, a larva may produce a painful boil-like skin lump with a small breathing opening; removal is performed by a clinician to avoid complications.

## **Botfly**

A fly whose larvae can infest skin or wounds in animals and occasionally people. The infestation is called myiasis and may cause a painful lump, discharge, itching, or tissue irritation.

## **Bothriocephalus**

An older name for a group of fish tapeworms. Related tapeworm infections may cause abdominal symptoms and, when prolonged, can reduce vitamin B12 absorption and lead to anaemia.

## **Botryomycosis**

A chronic bacterial infection that forms firm granules within lesions and can resemble a fungal infection. It most often affects the skin and tissue beneath it but may involve internal organs, especially in people with weakened immunity.

---

## **Bottle Feeding**

Feeding an infant expressed breast milk or infant formula from a bottle. Safe preparation, clean equipment, correct formula mixing, and responsive feeding help prevent infection, choking, overfeeding, and poor nutrition.

## **Botulinum Toxin**

A substance made by *Clostridium botulinum* that temporarily blocks nerve signals to muscles and some glands. Carefully measured injections treat selected muscle spasms, movement disorders, migraine, sweating, and cosmetic concerns; too much can cause weakness or breathing failure.

## **Botulism**

A rare, life-threatening illness caused by botulinum toxin, which blocks nerve signals. It causes blurred or double vision, drooping eyelids, difficulty swallowing, progressive weakness, and sometimes breathing failure; urgent antitoxin and intensive care may be needed.

## **Bouchard Nodes**

Firm enlargements of the middle joints of the fingers, usually caused by osteoarthritis. They may be painless or may cause stiffness, pain, reduced movement, and finger deformity.

## **Bouchard's Node**

A hard enlargement of one middle finger joint, usually caused by osteoarthritis. It results from changes in the joint and surrounding bone and may cause pain, stiffness, or reduced movement.

## **Bouchard's Nodes**

Hard enlargements of the middle finger joints caused most often by osteoarthritis. They can limit movement or cause pain and stiffness, although some people have no symptoms.

### **Alternate Names:**

Proximal Interphalangeal Osteophytes; PIP-Joint Nodes

## **Bougie**

A slender medical instrument used to examine or gradually widen a narrowed passage, such as the food pipe or urethra. It must be used carefully because it can cause pain, bleeding, perforation, or infection.

## **Bougienage**

Gradual widening or examination of a narrowed body passage with a bougie. It may relieve swallowing or urinary obstruction, but the cause of narrowing must be assessed and the procedure can cause bleeding or perforation.

## **Bouin's Solution**

A laboratory liquid containing picric acid, formaldehyde, and acetic acid that preserves tissue for microscopic examination. It is hazardous to handle and can interfere with some later genetic tests.

## **Bounding Pulse**

A hyperdynamic, high-amplitude arterial pulse characterized by a rapid, forceful upstroke and wide pulse pressure (Corrigan or water-hammer pulse), classically associated with aortic regurgitation, thyrotoxicosis, severe anemia, fever, patent ductus arteriosus, or septic shock.

### **Alternate Names:**

Hyperdynamic Pulse; Water-Hammer Pulse; Corrigan Pulse

## **Bourbon Virus**

A rare tick-associated viral infection that can cause fever, tiredness, poor appetite, nausea, and low white-cell or platelet counts. No specific treatment or vaccine is established; care is supportive and other serious infections must be excluded.

## **Boutonneuse Fever**

A tick-borne infection caused usually by *Rickettsia conorii*. It commonly causes fever, headache, rash, and a dark sore at the bite site; prompt antibiotic treatment reduces the risk of serious illness.

---

### **Boutonniere Deformity**

A finger shape problem where the middle joint stays bent inward and the tip bends backward, often caused by an injury or rheumatoid arthritis.

### **Bouveret Syndrome**

A rare blockage near the stomach outlet caused by a gallstone that has entered the intestine through an abnormal connection from the gallbladder. It can cause persistent vomiting, upper-abdominal pain, bloating, and inability to eat.

### **Bovine**

Relating to cattle or to material obtained from cattle. In medicine it may describe an animal disease, a biological product, tissue used in treatment, or an infection that can pass between cattle and people.

### **Bovine Anaplasmosis**

A tick- or blood-spread infection of cattle caused by *Anaplasma marginale*. The bacterium infects red blood cells and destroys them, causing anaemia, weakness, fever, reduced milk production, or death.

### **Bovine Papillomavirus**

A group of papillomaviruses that infect cattle and can cause skin warts or other growths. Different strains vary in their effects; human disease is not the usual concern.

### **Bovine Spongiform Encephalopathy**

A fatal disease of cattle caused by misfolded proteins called prions. It progressively damages the brain and nervous system; people can rarely develop variant Creutzfeldt–Jakob disease after eating contaminated cattle tissue.

### **Bow Legs**

An angular deformational alignment of the lower extremity characterized by outward lateral bowing of the tibia and femur such that the knees remain abnormally separated when the medial malleoli of the ankles are approximated.

#### **Alternate Names:**

Genu Varum; Bowlegs; Bandy Legs

### **Bowditch Effect**

The intrinsic autoregulatory property of cardiac muscle whereby an increase in heart rate leads to a progressive, stepwise increase in the force of myocardial contraction due to intracellular accumulation of calcium ions.

#### **Alternate Names:**

Treppe Phenomenon; Frequency-Dependent Inotropy

### **Bowel**

The small and large intestines. They receive partly digested food, absorb nutrients and water, move contents forward, and form and store stool before it leaves the body.

### **Bowel Burn**

Injury to the bowel lining caused by heat, chemicals, radiation, reduced blood flow, or another damaging exposure. It can cause pain, bleeding, perforation, infection, or narrowing and requires assessment according to the cause.

### **Bowel Cancer**

Cancer arising in the colon or rectum. It may cause a persistent change in bowel habit, blood in stool, anaemia, weight loss, abdominal pain, or no early symptoms; screening and timely assessment improve detection.

### **Bowel Death**

Death of part of the intestine, also called bowel necrosis or intestinal infarction. It usually results from severely reduced blood flow but may also follow severe infection, inflammation, blockage, or twisting of the bowel. It may cause severe abdominal pain, vomiting, bloody stool, fever, or a swollen and tender abdomen. The dead bowel can tear and cause peritonitis, sepsis, or shock; urgent surgery is often needed.

### **Bowel Disorders**

Pathological conditions affecting the small or large intestine, encompassing inflammatory bowel disease (Crohn disease, ulcerative colitis), infectious enterocolitis, ischemic bowel disease, bowel obstruction,

---

and colorectal neoplasms.

**Alternate Names:**

Intestinal Diseases; Bowel Diseases; Intestinal Disorders

**Bowel Habit Changes and Bloating**

A group of digestive symptoms consisting of diarrhea, constipation, gas, and abdominal bloating. They may result from diet, infection, medicines, food intolerance, or disorders such as irritable bowel syndrome; persistent, severe, bloody, or unexplained symptoms require assessment.

**Bowel Habits**

A person's usual pattern of passing stool, including how often it happens and the stool's form and ease of passage. Changes may include new constipation, diarrhea, urgency, or a different pattern.

**Bowel Incontinence**

Inability to control the passage of stool or liquid from the anus. It can result from diarrhoea, constipation, muscle or nerve injury, childbirth, bowel disease, or neurologic illness and may often be improved with treatment.

**Alternate Names:**

Fecal Incontinence

**Bowel Infarction**

Death of part of the intestine because its blood supply is blocked or severely reduced. It can cause severe abdominal pain, bloody stool, vomiting, and abdominal tenderness and is a life-threatening emergency.

**Bowel Ischemia**

Reduced blood flow to the intestine, causing injury from too little oxygen. It may cause sudden or worsening abdominal pain, vomiting, or bloody stool and can progress to bowel death if circulation is not restored.

**Bowel Movement**

The passing of stool from the rectum through the anus. Stool frequency and form vary among healthy

people.

**Bowel Obstruction**

A blockage that prevents food, fluid, and gas from moving through the intestine. It can cause cramping pain, swelling, vomiting, and inability to pass stool or gas and may require urgent hospital treatment.

**Bowel Preparation**

Emptying and cleansing the colon before a colonoscopy or other bowel procedure, usually with a special diet and a prescribed laxative solution. Following the instructions is important because poor preparation can hide lesions.

**Alternate Names:**

Bowel Prep; Colon Cleansing; Colonic Preparation

**Bowel Program**

A planned routine using food, fluids, medicines, movement, and scheduled toilet visits to produce regular bowel movements and manage constipation or loss of bowel control.

**Bowel Retraining**

A structured routine that uses regular toilet times, suitable food and fluids, physical activity, and helpful posture to establish predictable bowel movements and improve constipation or bowel-control problems.

**Bowel Sounds**

Gurgling or clicking sounds made as the intestines move gas and liquid. Their presence and pattern can provide clues during an abdominal examination but cannot diagnose a disorder by themselves.

**Bowel Transit Time**

The time food and other contents take to travel through the digestive tract. Testing it can help investigate unusually slow or fast movement when a bowel-movement or digestive disorder is suspected.

**Bowel Urgency**

A sudden, hard-to-delay need to pass stool. It can occur with diarrhoea, infection, irritable bowel syn-

---

drome, inflammatory bowel disease, or disorders of the rectum and may lead to accidents.

### **Bowen's Disease**

Squamous cell carcinoma in situ, an early skin cancer confined to the epidermis. It usually appears as a slowly enlarging, persistent scaly red or brown patch or plaque, often on sun-exposed skin; biopsy confirms the diagnosis, and untreated disease can progress to invasive squamous cell carcinoma.

### **Bowenoid Papulosis**

An HPV-associated squamous intraepithelial lesion of genital or perianal skin, usually presenting as multiple reddish-brown or violaceous papules. Biopsy shows atypical cells resembling squamous-cell carcinoma in situ, but many lesions behave indolently; diagnosis, treatment, and follow-up are important because progression can occur.

### **Bowen's Disease**

Squamous cell carcinoma in situ, a cancer limited to the epidermis. It usually appears as a persistent scaly or crusted red patch and can progress to invasive squamous cell carcinoma if untreated.

### **Bowlegs**

Alternate name; see *Bow Legs*.

### **Bowler's Thumb**

Irritation or scarring of the digital nerve along the thumb caused by repeated pressure from a bowling ball. It may produce pain, numbness, tingling, or a tender lump and may improve with rest or equipment changes.

### **Bowman Capsule**

A cup-shaped structure at the beginning of each kidney filtering unit. It surrounds a knot of tiny blood vessels and collects the fluid filtered from blood before it passes through the rest of the kidney tubule.

### **Bowman Membrane**

A thin, strong layer near the front of the cornea

that supports and protects the eye's clear surface. Significant injury can scar it because it does not normally grow back.

### **Bowman's Capsule**

The cup-shaped beginning of a kidney filtering unit. It surrounds a cluster of small blood vessels and receives the fluid filtered from blood.

### **Bowman's Gland**

A small gland high inside the nose that releases watery fluid onto the smell-sensing lining. The fluid helps dissolve scent molecules and clear the surface so odours can be detected.

### **Boxed Warning**

The strongest safety warning required in a medicine's prescribing information when there is a serious or life-threatening risk. It explains a major hazard but does not necessarily mean the medicine is unsuitable for every patient.

### **Boxer Fracture**

A transverse or comminuted fracture through the distal metaphysis or neck of the fifth metacarpal bone (and occasionally the fourth metacarpal), characteristically accompanied by volar angulation of the metacarpal head resulting from direct axial loading with a clenched fist.

### **Boxer's Ear**

A deformity of the outer ear caused by repeated injury or untreated blood collection around the ear cartilage. It can produce thickened, scarred tissue with a cauliflower-like appearance.

### **Boxer's Fracture**

A break near the knuckle end of the fifth metacarpal, the hand bone below the little finger. It commonly follows punching and causes pain, swelling, tenderness, or a changed knuckle shape.

### **BPH (Benign Prostatic Hyperplasia)**

Enlargement of the prostate that can obstruct urine flow. Symptoms include weak stream, hesitancy,



---

incomplete emptying, urgency, frequency, and night-time urination; severe retention or kidney effects require medical care.

### **Braces and Retainers**

Alternate name; see *Dental Braces*.

### **Brachial**

Relating to the upper arm, including its bones, muscles, blood vessels, and nerves. The brachial artery supplies the arm, and the brachial plexus carries nerve signals to and from the arm and hand.

### **Brachial Artery**

The main artery of the upper arm. It carries blood from the armpit toward the elbow, where it divides into the radial and ulnar arteries, and it is commonly used to measure a pulse or blood pressure.

### **Brachial Plexopathy**

Damage or dysfunction of the nerve network supplying the shoulder, arm, and hand. It can cause pain, weakness, numbness, or paralysis after injury, pressure, inflammation, a tumour, or radiation.

### **Brachial Plexus**

A network of nerves from the lower neck and upper chest that supplies movement and sensation to the shoulder, arm, forearm, and hand. Its branches include the musculocutaneous, axillary, radial, median, and ulnar nerves.

### **Brachial Plexus Cords**

Three parts of the nerve network supplying the arm, named lateral, posterior, and medial because of their position around an artery in the armpit. They join to form nerves controlling movement and feeling in the upper limb.

### **Brachial Plexus Injuries**

Damage to the nerve network supplying the shoulder and upper limb, caused by stretching, pressure, or a penetrating injury. Symptoms range from pain or numbness to weakness or paralysis, depending on the nerves affected.

### **Brachial Plexus Injury**

Damage to the nerve network of the shoulder and arm, sometimes occurring during a difficult birth or after trauma. It may cause weakness, loss of movement, numbness, or paralysis; recovery depends on the severity and nerves involved.

### **Brachial Plexus Injury In Newborns**

Damage to the nerves supplying a newborn's shoulder, arm, and hand, usually after a difficult delivery. The affected arm may be weak or still, and recovery ranges from spontaneous improvement to requiring specialist treatment.

### **Brachial Plexus Neuropathies**

Disorders affecting the nerve network that supplies the shoulder, arm, and hand. They can cause pain, weakness, numbness, or loss of movement because of injury, pressure, inflammation, tumours, or radiation.

### **Brachial Plexus Roots**

The first portions of the nerve network that supplies the arm, arising from the lower neck and upper chest. Damage to different roots produces characteristic patterns of weakness, numbness, and pain.

### **Brachial Plexus Trunks**

Three sections of the brachial plexus—upper, middle, and lower—that carry nerve fibres toward the arm. Injury to the upper trunk can cause weakness of the shoulder and elbow, while lower-trunk injury can mainly affect the hand.

### **Brachial Vein**

Deep veins accompanying the brachial artery that drain the arm and join the basilic vein to form the axillary vein.

### **Brachialgia**

Pain that travels from the neck or shoulder down the arm. It often results from irritation of a neck nerve or the brachial plexus and may occur with numbness, tingling, or weakness.

---

## Brachialis Muscle

A strong muscle at the front of the upper arm, beneath the biceps. It is the main muscle that bends the elbow and works whether the palm faces up, down, or sideways.

## Brachiocephalic Artery

The first and largest arterial branch arising from the aortic arch, which ascends obliquely behind the manubrium of the sternum before bifurcating into the right common carotid and right subclavian arteries.

### Alternate Names:

Innominate Artery

## Brachiocephalic Trunk

The first large branch of the aorta in the chest. It carries blood toward the right side of the head, neck, and arm and divides into the right common carotid and right subclavian arteries.

## Brachiocephalic Veins

Two large veins formed by the joining of veins from the head, neck, and arm. They drain blood from these areas into the superior vena cava, which returns it to the heart.

## Brachioplasty

An operation to remove or reshape excess skin and tissue of the upper arm, sometimes called an arm lift. Possible complications include scars, altered sensation, bleeding, fluid collection, infection, and wound problems.

## Brachioradial Pruritus

A localized neuropathic itch, burning, or tingling affecting the outer forearms, usually related to irritation or compression of cervical nerve roots, often C4–C6. The skin may look normal or show scratch marks; cooling can temporarily relieve symptoms, but persistent cases need evaluation for cervical spine or other neurologic causes.

## Brachioradialis Reflex

A deep tendon stretch reflex elicited by briskly tap-

ping the tendon of the brachioradialis muscle near the styloid process of the radius with a reflex hammer, producing forearm flexion and slight pronation. It is innervated by the radial nerve and tests the integrity of the C5 and C6 cervical spinal cord segments and nerve roots.

### Alternate Names:

Supinator Reflex; Radial Reflex

## Brachium

The part of the upper limb between the shoulder and elbow. It contains the upper-arm bone, muscles that move the shoulder and elbow, and major nerves and blood vessels.

## Brachycephalic

Describes a skull that is relatively short from front to back and broad from side to side. It may be a normal family characteristic or part of a condition affecting skull growth.

## Brachycephaly

A broad, short skull shape. It may develop from pressure on an infant's head or from early joining of skull bones; assessment determines whether brain growth or treatment is affected.

## Brachydactyly

A birth condition in which one or more fingers or toes are shorter than usual because of altered development of their bones or surrounding tissues. It may occur alone or as part of an inherited syndrome.

## Brachytherapy

Internal radiation treatment in which a radioactive source is placed inside or close to a tumour. It delivers a concentrated dose to a limited area and may use a temporary applicator or a source left in place permanently.

### Alternate Names:

Brachy Therapy Internal Radiation Therapy

## Brachytherapy Seed

A tiny radioactive source placed in or near a tumour to deliver local radiation. Seeds may be temporary

---

or permanent and are used most often for selected cancers, including prostate cancer.

### **Bradyarrhythmia**

Any cardiac rhythm disorder characterized by an abnormally slow heart rate (typically fewer than 60 beats per minute in adults), resulting from sinus node dysfunction, sinus arrest, sinoatrial block, or high-grade atrioventricular block.

#### **Alternate Names:**

Bradycardia Arrhythmia

### **Bradycardia**

A slower-than-usual heart rate, often below 60 beats per minute in adults. It can be normal during sleep or in trained athletes, but may result from heart-conduction disease or medicines and cause dizziness, fainting, tiredness, or breathlessness.

#### **Alternate Names:**

Slow Heartbeat

### **Bradycardia (Slow Heart Rate)**

A slower-than-expected heart rate. It may be normal in some people or result from conduction disease, medicines, metabolic problems, or heart disease; fainting, chest pain, breathlessness, or confusion requires urgent assessment.

### **Bradycardia Of Prematurity**

A slow heart rate in a premature infant, often occurring with pauses in breathing and low oxygen. It commonly reflects immature breathing and heart control and is monitored and treated according to its frequency and severity.

### **Bradycardia-Tachycardia Syndrome**

Alternate name; see *Sick Sinus Syndrome*.

### **Bradykinesia**

Slowness and reduced size of voluntary movements. It can make starting or repeating actions difficult and may occur with stiffness, tremor, reduced arm swing, a less expressive face, or a shuffling walk in Parkinson disease and related disorders.

### **Bradykinesia and Athetosis**

Hypokinetic and dyskinetic movement patterns characterized by generalized slowness of movement initiation (bradykinesia) or continuous slow, writhing, twisting involuntary movements (athetosis), frequently observed in Parkinson disease, secondary parkinsonism, or basal ganglia encephalopathies.

#### **Alternate Names:**

Slowness of Movement; Hypokinesia and Athetosis; Extrapyramidal Slowness

### **Bradykinetic**

Describes abnormally slow movement, especially the reduced speed and size of movement seen in Parkinson disease and related neurologic disorders.

### **Bradykinin**

A natural chemical messenger that widens some blood vessels and allows more fluid to leave them. It contributes to inflammation, swelling, pain, and rare reactions to certain blood-pressure medicines.

### **Bradyphrenia**

Unusually slow thinking or mental processing. It may cause delayed responses and difficulty completing tasks and can occur with neurologic disease, mental illness, medicines, depression, or other conditions.

### **Bradypnea**

Abnormally slow breathing, often fewer than 12 breaths per minute in an adult. It may result from sedative or opioid effects, brain disease, low body temperature, hormone disorders, or severe illness and can lead to respiratory failure.

### **Bradypnoea**

Abnormally slow breathing. Causes include medicines, neurologic disease, metabolic disturbance, low body temperature, or severe illness; breathing that is difficult, shallow, or associated with confusion requires urgent care.

### **BRAF Gene**

A gene that encodes a signaling protein in the

---

RAS-RAF-MEK-ERK pathway, which regulates cell growth and division. Activating BRAF variants can drive cancers such as melanoma, thyroid cancer, and colorectal cancer and may guide targeted treatment.

### Braille System

A raised-dot writing and reading system used by people who are blind or have low vision. Patterns represent letters, numbers, punctuation, and symbols and can be read by touch.

### Brain

The main organ of the nervous system, protected inside the skull. It processes senses, controls movement and body functions, and supports thinking, memory, emotions, language, and consciousness.

### Brain Abscess

A pocket of pus and infection inside the brain. It can develop when infection spreads from the blood, nearby ear or sinus disease, surgery, or an injury and may cause headache, fever, seizures, or weakness.

### Brain Aneurysm

A weakened bulge in an artery supplying the brain. If it bursts, it can cause sudden severe headache, vomiting, neck stiffness, collapse, or other neurological problems.

### Brain Aneurysm Repair

Treatment to prevent a brain aneurysm from bleeding or bleeding again. A surgeon may place a clip or a specialist may seal the aneurysm from inside the artery with coils or another device.

### Brain Arteriovenous Malformation

A congenital vascular dysplasia consisting of a direct high-flow, low-resistance communication between cerebral arteries and veins through an abnormal vascular nidus without an intervening capillary bed, carrying a risk of intracranial hemorrhage, focal seizures, and neurological deficits.

#### Alternate Names:

Cerebral Arteriovenous Malformation; Brain AVM; Intracranial AVM

### Brain AVM

Alternate name; see *Brain Arteriovenous Malformation*.

### Brain Cancer

Cancer affecting brain tissue or its coverings. Symptoms depend on the location and may include seizures, headache, weakness, speech or vision changes, personality change, or pressure-related vomiting; diagnosis usually requires brain imaging and sometimes tissue testing.

#### Alternate Names:

Malignant Brain Tumor; Primary Brain Neoplasm

### Brain Concussion

Alternate name; see *Concussion*.

### Brain Death

Permanent loss of all brain and brainstem function, including consciousness, reflexes, and the ability to breathe independently. When formally confirmed according to law and medical standards, it means the person has died.

### Brain Disorder

Any illness or abnormality affecting the brain's structure or function, including stroke, infection, tumour, epilepsy, injury, degenerative disease, or a condition present from birth.

### Brain Edema

Swelling caused by extra fluid in brain tissue. It can increase pressure inside the skull after injury, stroke, infection, bleeding, tumour, or severe illness and may cause headache, vomiting, seizures, confusion, or reduced consciousness.

### Brain Embolism

Alternate name; see *Embolic Stroke*.

### Brain Fog

An informal description of difficulty concentrating, thinking clearly, remembering, or finding words. It is a symptom rather than a diagnosis and may occur with poor sleep, stress, illness, medicines, depression,

---

or other conditions.

### **Brain Fornix**

A curved bundle of nerve fibres deep in the brain that helps connect the hippocampus with other areas involved in memory. Damage may cause difficulty learning or remembering information.

### **Brain Haemorrhage**

Bleeding within or around the brain. It may result from high blood pressure, injury, an aneurysm or other blood-vessel problem, clotting disorders, or medicines and can cause sudden neurologic symptoms.

### **Brain Health**

The ability of the brain to support thinking, memory, learning, movement, emotions, communication, and consciousness. It is supported by controlling vascular risks, sleeping well, physical activity, balanced nutrition, social connection, and treatment of illness.

### **Brain Hemorrhage**

Bleeding within or around the brain, often caused by high blood pressure, an aneurysm, a blood-vessel malformation, injury, or blood-thinning treatment. Sudden severe headache, weakness, speech trouble, vomiting, seizure, or reduced consciousness requires emergency care.

### **Brain Herniation**

Dangerous displacement of brain tissue caused by severe swelling, bleeding, a tumour, or another increase in pressure inside the skull. It can compress the brainstem, impair breathing and consciousness, and requires emergency treatment.

### **Brain Imaging**

Tests that create pictures of brain structure or activity. CT and MRI show tissue, bleeding, swelling, tumours, and strokes; blood-vessel or nuclear scans answer more specific questions.

#### **Alternate Names:**

Neuroimaging; Cerebral Imaging

### **Brain Injuries**

Damage to the brain ranging from a concussion to severe injury, commonly caused by falls, crashes, sports impacts, or penetrating wounds. Effects depend on the area and severity and may include headache, confusion, memory loss, seizures, or unconsciousness.

### **Brain Injury**

Damage or impaired function of the brain caused by trauma, lack of oxygen, stroke, infection, toxins, or another disease. Effects vary with the location and severity and may affect movement, sensation, thinking, behaviour, or consciousness.

### **Brain Lesions**

Areas of brain tissue that look or function abnormally because of a tumour, stroke, infection, inflammation, injury, loss of nerve insulation, or a blood-vessel disorder. Symptoms depend mainly on the lesion's location and size.

### **Brain Malformations**

Structural differences in the brain that develop before birth. They may result from genetic changes, disrupted fetal development, infection, or other prenatal influences and can affect movement, learning, seizures, or other neurologic functions.

### **Brain Mapping**

Methods used to identify where brain functions and structures are located. Imaging, electrical recordings, or direct stimulation may guide research or help surgeons avoid important areas during selected operations.

### **Brain Metastases**

Cancer deposits that have spread to the brain from another organ. They may cause headache, seizures, weakness, speech or vision changes, confusion, or no symptoms and are assessed with brain imaging.

#### **Alternate Names:**

Secondary Brain Cancer

---

## Brain Metastasis

A cancer deposit that has spread to the brain from another organ. Symptoms may include headache, seizures, weakness, speech or vision changes; treatment depends on the number, location, original cancer, and overall health.

## Brain Monitoring

Recording brain activity or pressure to assess neurologic function, seizures, sleep, or the depth of anaesthesia. The method used depends on the clinical question; a monitor result is interpreted with examination findings and other tests.

## Brain Natriuretic Peptide Test

A blood test measuring BNP or NT-proBNP, substances released when the heart is stretched. A raised result can support a diagnosis of heart failure, but it must be interpreted with symptoms, examination, age, kidney function, and other findings.

## Brain Neoplasm

An abnormal growth of cells in the brain or its coverings. It may be noncancerous or cancerous; imaging and sometimes biopsy determine its type, behaviour, extent, and treatment options.

## Brain Neoplasms

Tumors arising in the brain, its coverings, or nearby central nervous system structures. They may be benign or malignant and can cause headache, seizures, cognitive or behavioral change, weakness, sensory loss, or hormonal or visual problems; MRI and tissue diagnosis guide classification.

## Brain PET Scan

A scan that uses a small amount of radioactive tracer to show chemical activity in the brain. It can help investigate selected tumours, memory disorders, seizures, or brain function when other scans do not answer the question.

## Brain Sand

Brain sand is an older name for corpora arenacea, tiny calcified deposits commonly found in the pineal

gland and sometimes other brain regions.

## Brain SPECT

A nuclear-imaging scan that shows blood flow or selected brain activity using a radioactive tracer. It may help evaluate seizures, stroke-related blood-flow changes, dementia, or other specialist questions when interpreted with other findings.

## Brain Stem

The lower part of the brain connecting the cerebrum with the spinal cord and controlling essential functions such as breathing and consciousness.

## Brain Stem Glioma

A tumor arising from glial cells in the brainstem, the part of the brain controlling important functions such as breathing, swallowing, eye movement, and balance. Symptoms and treatment depend on its exact location and behavior.

## Brain Stimulation

Use of electrical, magnetic, or other energy to change brain activity. Methods include electroconvulsive therapy, transcranial magnetic stimulation, and deep-brain stimulation; their uses, benefits, and risks differ.

## Brain Surgery

An operation on the brain or nearby structures to remove or sample a tumor, relieve pressure, treat bleeding, repair an abnormal blood vessel, or control selected seizures. Risks depend on the area and procedure and can include bleeding, infection, or neurologic loss.

### Alternate Names:

Brain Tumor Surgery

## Brain Tumor

An abnormal growth of cells in the brain or its coverings. It may be benign or malignant, begin in the brain, or spread there from cancer elsewhere. It can cause symptoms by affecting nearby tissue, blocking fluid flow, bleeding, or raising pressure inside the skull; symptoms may include seizures, headache, vomiting, weakness, or changes in thinking.



---

## Brain Tumors in Adults

Tumors arising in or near the adult brain, including primary gliomas, meningiomas, metastases, and other lesions. Symptoms depend on location and may include headache, seizures, weakness, personality change, vision problems, or cognitive decline.

## Brain Tumour

An abnormal growth in or near the brain that may be noncancerous or cancerous. It can cause seizures, headache, vomiting, changes in thinking, weakness, or other problems by affecting nearby brain tissue.

## Brain Ventricles

Four connected spaces inside the brain that contain and circulate cerebrospinal fluid. This fluid cushions the brain and spinal cord, carries substances, and helps remove waste; blockage can cause pressure and hydrocephalus.

## Brain Wave

A pattern of electrical activity made by groups of nerve cells and recorded with an electroencephalogram. Patterns change with sleep, alertness, seizures, and some disorders of brain function.

## Brain Waves

Patterns of electrical activity produced by groups of nerve cells and recorded with an electroencephalogram. Their timing and shape can help assess sleep, seizures, brain function, and some disorders affecting consciousness.

### Alternate Names:

Cerebral Electrical Activity; EEG Rhythms

## Brainstem

The lower part of the brain connecting the cerebrum with the spinal cord. It helps control breathing, heart rate, blood pressure, swallowing, eye movements, alertness, and pathways carrying signals between the brain and body.

## Brainstem Auditory Evoked Potential

An electrophysiological test measuring the neural transmission of acoustic signals from the eighth

cranial nerve through the brainstem pathways to the auditory cortex, used to assess hearing acuity and brainstem integrity.

### Alternate Names:

BAEP; Brainstem Auditory Evoked Response; BAER

## Brainstem Auditory Evoked Response

An electrodiagnostic test that records electrical potentials generated by the auditory nerve and brainstem pathways in response to acoustic click or tone stimuli. It is utilized to assess hearing acuity in newborns and uncooperative patients, and to evaluate auditory pathway lesions such as acoustic neuromas.

### Alternate Names:

Brainstem Auditory Evoked Potential; BAEP; Brainstem Auditory Evoked Response; BAER; Auditory Brainstem Response; ABR

## Brainstem Death

Irreversible cessation of all functions of the brainstem, certified clinically by persistent deep coma, total absence of brainstem reflexes, and documented apnea during formal apnea testing in the presence of a known, irreversible intracranial cause.

## Brainstem Gliomas

Gliomas arising in the midbrain, pons, or medulla. They may affect eye movements, facial function, swallowing, balance, strength, or breathing; imaging and tumor biology guide diagnosis and management, which differs between children and adults.

## Brainstem Tumor

A growth arising in the brainstem, which connects the brain with the spinal cord and controls important functions. Symptoms may include problems with eye movement, balance, swallowing, speech, strength, or breathing.

## Branchial

Relating to structures that develop from the paired tissue arches in an embryo's early head and neck. These arches contribute to parts of the face, throat, voice box, muscles, bones, blood vessels, and nerves.

---

## Branchial Arch

One of several paired structures in an embryo that develop into parts of the face, throat, voice box, muscles, bones, blood vessels, and nerves of the head and neck.

## Branchial Cleft

A groove between developing head-and-neck arches in an embryo. If part of it remains after birth, it may form a cyst, a narrow passage, or an opening, usually on the side of the neck.

## Branchial Cleft Anomalies

Congenital developmental defects resulting from incomplete obliteration of the embryonic branchial apparatus, manifesting as cervical cysts, sinuses, or fistulae, most commonly arising along the anterior border of the sternocleidomastoid muscle from the second branchial cleft.

## Branchial Cleft Anomaly

A birth-related remnant of developing head-and-neck tissue that may form a cyst, narrow passage, or skin opening. Repeated infection, swelling, or drainage from the side of the neck needs medical assessment.

## Branchial Cleft Cyst

A congenital, usually lateral neck cyst caused by persistence of embryonic branchial-arch tissue, most often a second-branchial remnant. It may enlarge or become painful after an upper-respiratory infection and can cause drainage or swallowing difficulty; a new lateral neck mass in an adult requires evaluation to exclude malignancy before it is assumed to be a cyst.

## Branchial Cyst

A usually harmless, birth-related fluid-filled lump on the side of the neck caused by leftover tissue from early development. It may become larger or painful during an infection. A new cyst-like neck lump in an adult needs careful assessment because cancer can look similar. Treatment depends on the cause and may include surgical removal.

**Alternate Names:**

Branchial Cleft Cyst; Second Branchial Cleft Cyst; Lateral Cervical Cyst

## Branchio-Oto-Renal Syndrome

An inherited disorder that can cause neck cysts or openings, hearing loss, and kidney abnormalities. The combination and severity vary, and some affected people have only one or two of these features.

## Branchioma

A rare cyst or growth arising from tissue related to early head-and-neck development. It often presents as a lump in the neck and requires examination and imaging to establish its nature.

## Brass-Founder's Ague

An older name for metal-fume fever, a short-lived flu-like illness after breathing fumes from heated metals such as zinc or brass. Symptoms may include fever, chills, headache, muscle aches, cough, and nausea.

## Brassicasterol

A plant sterol found especially in rapeseed and related oils. It is mainly used to identify the composition of plant oils and has no routine role as a medical treatment.

## Brassicidic Acid

A long-chain fatty acid found in some plant oils. It is mainly a chemistry term and is not an established medicine or routine clinical test.

## Braxton Hicks Contractions

Irregular tightening of the uterus during pregnancy, often becoming more noticeable later in pregnancy. They usually do not steadily strengthen or open the cervix and may ease with rest, fluids, or a change of activity.

**Alternate Names:**

False Labor; Practice Contractions

## Brazilian Pemphigus Foliaceus

An endemic form of superficial pemphigus seen especially in parts of Brazil, also called fogo selvagem. Autoimmune attack on epidermal cell connections

---

causes fragile superficial blisters and erosions without the deep mucosal disease typical of pemphigus vulgaris.

### **Brazilian Trypanosomiasis**

Chagas disease, a parasitic infection caused by *Trypanosoma cruzi*. It is transmitted mainly by triatomine insects or congenitally and may later cause cardiomyopathy, arrhythmias, megaesophagus, or megacolon.

### **BRCA**

BRCA1 and BRCA2 are genes that help repair damaged DNA. An inherited harmful change in either gene increases the risk of breast, ovarian, fallopian-tube, primary peritoneal, prostate, and pancreatic cancers.

### **BRCA1 And BRCA2**

Genes that help repair breaks in DNA and limit abnormal cell growth. An inherited harmful variant raises the risk of several cancers and may affect screening, preventive surgery, and treatment choices for the person and relatives.

### **BRCA1 And BRCA2 Gene Testing**

A blood or saliva test looking for harmful inherited changes in BRCA1 or BRCA2. Results can guide cancer screening, preventive options, and treatment and may provide important information for biological relatives.

### **BRCA1 Gene**

A gene involved in repairing damaged DNA and limiting tumor formation. An inherited harmful variant increases the risk of breast, ovarian, fallopian-tube, peritoneal, prostate, and pancreatic cancers; genetic counseling helps interpret test results.

### **BRCA2 Gene**

A gene that helps repair serious DNA damage. An inherited harmful variant increases the risk of breast, ovarian, prostate, pancreatic, and some other cancers and may affect relatives; genetic counseling helps explain testing and options.

### **Breakbone Fever**

An older name for dengue fever, reflecting the severe muscle and joint pain that may occur with this mosquito-borne viral infection. Fever, headache, rash, nausea, or dangerous bleeding may also develop.

### **Breakthrough Bleeding**

Unscheduled, unexpected uterine bleeding or spotting occurring during the active hormonal phase of combined oral contraceptive use or hormone replacement therapy, frequently seen during early months of initiation.

### **Breakthrough Infection**

An infection that occurs despite vaccination or another preventive measure intended to reduce its risk. Protection and illness severity vary with the germ, intervention, timing, and person's immune system.

### **Breakthrough Pain**

A short flare of pain that occurs despite otherwise controlled persistent pain. It may happen unexpectedly or after activity; treatment depends on the cause, pattern, regular pain medicines, and safety risks.

### **Breast**

One of the two organs on the front of the chest, made of milk-producing tissue, ducts, fat, connective tissue, blood vessels, nerves, and skin. It can be affected by benign conditions and cancer.

### **Breast Abscess**

A pocket of pus in breast tissue, often developing after mastitis during breastfeeding. It may cause a painful lump, redness, warmth, and fever; ultrasound can confirm it, and drainage plus antibiotics may be needed.

### **Breast Anatomy**

The breast contains skin, fat, milk-producing lobules, ducts, connective tissue, blood vessels, nerves, and lymph vessels that extend toward the armpit. Its appearance changes with age, hormones, pregnancy, and breastfeeding.

---

## Breast And Thoracic Cancer Surgery

Operations for cancers of the breast or chest, including selected cancers of the lung, pleura, food pipe, mediastinum, or chest wall. Surgery may remove a tumour and nearby lymph nodes, with further treatment based on the cancer type and stage.

### Alternate Names:

Breast and Thoracic Oncologic Surgery

## Breast Augmentation

A procedure to increase breast size or change its shape, usually with an implant or transferred fat. Risks include infection, bleeding, scarring, altered sensation, implant problems, and the need for further surgery.

### Alternate Names:

Breast Enlargement; Augmentation Mammoplasty

## Breast Augmentation Surgery

An operation that enlarges or reshapes the breasts using implants or transferred tissue. The surgeon discusses expected results, alternatives, risks, recovery, and possible later procedures before treatment.

## Breast Bone

A common name for the breastbone, or sternum, a flat bone in the centre of the chest. It joins the ribs and helps protect the heart and lungs.

## Breast Calcifications

Small calcium deposits seen in breast tissue on a mammogram. Most are harmless, but their size, shape, and arrangement may require additional imaging or biopsy to exclude an early cancer.

## Breast Cancer

Cancer that begins in breast tissue. It may appear as a lump, skin or nipple change, or abnormal imaging, although early cancer may cause no symptoms; treatment depends on its type, stage, and biology.

## Breast Cancer And Lymphedema

Breast-cancer surgery or radiation can damage lymph drainage and cause lasting swelling, heaviness, tightness, or skin changes in the arm, breast,

or chest. Early assessment and specialist treatment can reduce progression and complications.

## Breast Cancer and Pregnancy

Breast cancer diagnosed during pregnancy or shortly after delivery. Evaluation and treatment are individualized to the cancer stage and gestational age; some surgery and chemotherapy can be given during pregnancy, while radiation and certain targeted medicines are generally deferred.

## Breast Cancer and Your Genes

The role of inherited genetic variants in breast-cancer risk, including changes in *BRCA1*, *BRCA2*, *PALB2*, and other genes. Genetic counseling helps determine whether testing and enhanced screening are appropriate.

## Breast Cancer Genetics

The study of inherited gene changes that raise breast-cancer risk. Important genes include *BRCA1*, *BRCA2*, *PALB2*, *TP53*, *PTEN*, and *CHEK2*; genetic counselling helps decide whether testing and extra screening are appropriate.

## Breast Cancer Grade

A description of how abnormal breast-cancer cells look under a microscope and how quickly they are likely to grow. Grade is different from cancer stage, which describes the cancer's size and spread.

## Breast Cancer in Men

Breast cancer arising in male breast tissue. It is uncommon but may present as a painless lump, nipple retraction, discharge, or skin change and requires prompt diagnostic imaging and biopsy.

## Breast Cancer Prevention

Ways to lower, but not eliminate, breast-cancer risk include physical activity, limiting alcohol, avoiding tobacco, appropriate screening, and discussing inherited risk. Preventive medicine or surgery may be considered when risk is substantially high.

---

## Breast Cancer Recurrence

Return of breast cancer after treatment, either near the original site, in nearby lymph nodes, or in distant organs. New persistent symptoms or a lump require assessment, although many such symptoms have noncancerous causes.

## Breast Cancer Screening

Testing to find breast cancer before symptoms appear, usually with mammography. Ultrasound or MRI may be added for selected people according to age, breast density, inherited risk, and other factors.

## Breast Cancer Staging

Assessment of how far breast cancer has grown or spread, including the tumour's features, nearby lymph nodes, and distant organs. The stage helps estimate outlook and choose treatment.

## Breast Carcinoma

A cancer that begins in breast tissue, usually in the ducts or milk-producing lobules. Its treatment depends on the cell type, hormone and other markers, tumour size, lymph nodes, and spread.

## Breast Cyst

A benign, fluid-filled sac within breast tissue, most common in women between the ages of 35 and 50. It typically presents as a smooth, round, movable lump that can fluctuate in size and tenderness with the menstrual cycle. Ultrasound imaging distinguishes fluid-filled cysts from solid masses. Simple cysts are noncancerous and do not increase the risk of breast cancer; they generally require no intervention unless fluid is drained with a needle to relieve discomfort.

## Breast Cysts

Alternate name; see *Breast Cyst*.

## Breast Cytology

Examination of individual cells from a breast lump, nipple discharge, or needle sample. It can suggest infection, a harmless change, or cancer, but a core biopsy is often needed to determine the full tissue pattern.

## Breast Density

The relative amount of milk-gland and connective tissue compared with fat on a mammogram. Dense breast tissue can hide small cancers and is linked with higher breast-cancer risk; screening plans depend on individual risk and local guidance.

## Breast Development

Growth and maturation of breast tissue, beginning mainly at puberty and changing with hormones, pregnancy, breastfeeding, ageing, and some medicines. Uneven development is common, but a new persistent change requires assessment.

## Breast Development in Boys

Enlargement or development of breast tissue in a boy, most often gynecomastia caused by a temporary imbalance between estrogen and androgen effects. Persistent, painful, asymmetric, or rapidly enlarging tissue needs assessment.

## Breast Duct

A channel that carries milk from milk-producing areas toward the nipple. Blockage, inflammation, or abnormal cells in a duct can cause a lump, pain, or nipple discharge.

## Breast Engorgement

Overfilling and swelling of the breasts with milk and extra tissue fluid, usually a few days after birth. Both breasts may feel firm, warm, painful, and make the nipples harder for an infant to grasp.

## Breast Enlargement In Males

Enlargement of glandular breast tissue in a male, called gynecomastia. It may result from normal hormonal changes, medicines, obesity, liver or kidney disease, or other hormone disorders; a hard one-sided lump needs assessment.

## Breast Examination

Inspection and gentle examination of the breasts, nipples, and nearby lymph nodes to assess lumps, pain, skin or nipple changes, and other findings. It may be followed by imaging or biopsy when needed.



---

## Breast Feed

To feed an infant directly from the breast with human milk. Human milk provides nutrients and immune protection, but feeding may need support when the parent or infant has illness, difficulty latching, or medicine-related concerns.

## Breast Finding

Any noticeable breast change, such as a lump, pain, skin change, nipple inversion, or discharge. Its significance depends on the examination and imaging and sometimes requires a biopsy.

## Breast Health

Care and awareness aimed at recognising breast changes and detecting disease early. It includes awareness of what is normal, screening appropriate for age and risk, and prompt reporting of a new lump, discharge, or persistent skin change.

## Breast Imaging-Reporting And Data System

A standard system for describing mammogram, breast ultrasound, and breast MRI findings. Its numbered categories indicate whether results are normal, benign, probably benign, suspicious, or already known to be cancer and help guide follow-up.

## Breast Infection

Infection and inflammation of breast tissue, often occurring during breastfeeding. It can cause local pain, warmth, redness, swelling, and fever; worsening symptoms, a pus-filled lump, or feeling very unwell needs medical care because an abscess may develop.

## Breast Infection (Mastitis)

Inflammation or infection of breast tissue, often during breastfeeding but also possible in other settings. Pain, redness, warmth, swelling, fever, and a lump may occur; an abscess can develop if infection persists.

## Breast Lift

Surgery that raises and reshapes sagging breast tissue by removing extra skin and, when needed,

repositioning the nipple. It changes shape rather than necessarily increasing breast size and can leave scars or alter sensation.

## Breast Lobe

One of the larger sections of a breast's milk-producing gland. It contains smaller lobules and ducts that carry milk toward the nipple.

## Breast Lobule

A small milk-producing unit within the breast, made of tiny milk glands and ducts. Lobules enlarge and become more active during pregnancy and breastfeeding.

## Breast Lump

A swelling, mass, or thickened area in breast tissue. Most breast lumps (around 80% to 90%) are harmless and noncancerous, most often caused by fluid sacs (cysts), noncancerous growths (fibroadenomas), hormone changes, or past injury. Cancerous lumps are less common and usually feel hard, uneven, and fixed in place. Doctors identify the cause through physical examination, imaging such as ultrasound or mammograms, and a needle test (biopsy) if needed.

## Breast Lump Removal

An operation to remove a breast lump, usually so it can be examined or because it causes symptoms or concern. The need for surgery depends on examination, imaging, biopsy results, and the person's preferences.

## Breast Lumps

Alternate name; see *Breast Lump*.

## Breast Lumps (Benign)

Noncancerous breast masses such as cysts, fibroadenomas, fat necrosis, or fibrocystic changes. A new, persistent, hard, fixed, enlarging, or otherwise concerning lump requires clinical evaluation and appropriate imaging.

## Breast Milk

Milk made by the breasts after childbirth. It pro-



---

vides energy, nutrients, fluid, and immune substances that support an infant's growth and help protect against some infections.

### **Breast Milk Expression and Storage**

The manual or mechanical extraction of human breast milk and its subsequent storage following strict hygiene, refrigeration, and freezing protocols to preserve bioactive immunoglobulins, micronutrients, and enzymatic properties while preventing microbial contamination.

#### **Alternate Names:**

Pumping and Storing Breast Milk; Breast Milk Storage; Milk Expression

### **Breast Milk Jaundice**

Yellowing of the skin and eyes that begins after the first week in an otherwise well, feeding, and growing breastfed infant. It is usually temporary, but every newborn with jaundice needs assessment to exclude dangerous causes; feeding is generally continued unless clinicians advise otherwise.

#### **Alternate Names:**

Human Milk Jaundice; Late-Onset Neonatal Jaundice

### **Breast MRI**

An imaging test using a strong magnet and radio waves to create detailed pictures of breast tissue. Contrast is often used; MRI is added for selected screening, implant, or diagnostic questions rather than replacing mammography for everyone.

### **Breast MRI Scan**

A scan that uses magnetic fields and radio waves, sometimes with injected contrast, to show breast tissue in detail. It can help assess high cancer risk, an abnormality, implants, or the extent of known cancer.

### **Breast Neoplasm**

An abnormal growth of cells in breast tissue. It may be noncancerous or cancerous; imaging and often a biopsy determine its type, behaviour, extent, and treatment.

### **Breast Pain**

Pain or tenderness in one or both breasts. Common causes include hormonal changes, cysts, infection, injury, medicines, or chest-wall problems; persistent focal pain or pain with a lump needs assessment.

#### **Alternate Names:**

Mastalgia; Mastodynia

### **Breast Pain (Mastalgia)**

Pain or tenderness in one or both breasts. It may be cyclic, related to hormones, injury, cysts, infection, medicines, or chest-wall structures; a new mass, skin change, discharge, or persistent focal pain warrants assessment.

### **Breast Quadrant**

One of four areas used to describe where a finding is located in the breast: upper or lower, and inner or outer. The location helps describe examination and imaging findings.

### **Breast Rash**

A change such as redness, scaling, itching, or irritation of breast skin. Causes include dermatitis, allergy, infection, or rarely inflammatory breast cancer; a persistent, one-sided, painful, or nipple-associated rash needs assessment.

### **Breast Reconstruction**

Breast Reconstruction is surgery to rebuild the shape and appearance of a breast, most often after mastectomy. It may use a breast implant, tissue transferred from another part of the body, or both. Reconstruction may begin during mastectomy (immediate reconstruction) or be performed later (delayed reconstruction). It may also include reconstruction of the nipple and areola and may require several procedures.

### **Breast Reduction**

Surgery that removes breast tissue, fat, and skin to reduce breast size and relieve symptoms such as back, neck, or shoulder pain. Possible effects include scars, altered sensation, breastfeeding difficulty, and uneven shape.

---

## Breast Screening

Testing people without breast symptoms to find breast cancer early, most commonly with mammography. The recommended starting age and interval depend on local guidance, personal risk, family history, and previous findings.

## Breast Self-Examination

Looking at and feeling one's own breasts to notice a new lump, skin or nipple change, or discharge. It may improve familiarity with normal changes but does not replace recommended screening or clinical assessment.

## Breast Skin And Nipple Changes

Changes such as dimpling, redness, scaling, inversion, crusting, or discharge may result from benign conditions, infection, hormonal changes, or cancer. New, persistent, one-sided, or bloody changes require assessment.

## Breast Surgeon

A surgeon who diagnoses and treats breast conditions, including benign disease, breast cancer, reconstruction, and operations that reduce cancer risk.

## Breast Surgery

An operation on the breast to remove or examine a lump, treat cancer, reduce size, reconstruct shape, or address another condition. The type of surgery depends on the diagnosis, extent of disease, and person's goals.

## Breast Ultrasound

Imaging that uses sound waves to examine breast tissue. It helps assess a lump, distinguish fluid from solid tissue, investigate a mammogram finding, guide a biopsy, and evaluate selected implants or dense breasts.

## Breast-Conserving Treatment

An oncological treatment approach for early-stage breast cancer that aims to treat the cancer effectively while preserving the natural appearance of the breast. It consists of breast-conserving surgery (a lumpectomy, wide local excision, or quadrantec-

tomy to completely remove the tumor with clear microscopic tissue margins and assess sentinel lymph nodes) followed by adjuvant whole-breast radiation therapy to destroy any remaining microscopic cancer cells. Large randomized clinical trials demonstrate that breast-conserving therapy yields long-term survival rates equivalent to complete mastectomy. Contraindications include multicentric disease across multiple quadrants, extensive diffuse microcalcifications, prior chest wall radiation, or persistently positive surgical margins.

### Alternate Names:

Breast-Conserving Therapy; BCT; Breast-Conserving Surgery; BCS; Lumpectomy with Radiotherapy

## Breastfeeding

Feeding an infant human milk from the breast. It provides nutrition and immune protection; many health organisations recommend exclusive breastfeeding for about six months, followed by complementary foods while breastfeeding continues as desired and feasible.

## Breastfeeding With Rheumatoid Arthritis

A person with rheumatoid arthritis may be able to breastfeed. Disease activity, fatigue, hand function, and medicine safety are reviewed because some treatments are compatible and others require avoidance or careful timing.

## Breastfeeding-Associated Nipple and Skin Changes

Dermatologic and mucosal complications of the nipple-areolar complex occurring during lactation, including painful erythema, skin fissures, milk blisters (blebs), contact dermatitis, Raynaud phenomenon of the nipple, and infectious mastitis or candidiasis.

### Alternate Names:

Nipple Trauma in Breastfeeding; Sore Nipples; Nipple Fissures in Lactation

## Breath

Air moved into and out of the lungs. Its smell,

---

depth, rate, pattern, and oxygen or carbon-dioxide content can provide clues about breathing, infection, metabolism, or other health problems.

### **Breath Alcohol Test**

A test measuring alcohol in exhaled air to estimate alcohol concentration in the blood. Results depend on timing, the device, sampling, and health factors and may be used in clinical or legal settings.

### **Breath Analysis**

Measurement of gases or other chemicals in exhaled air. Different tests can estimate alcohol, assess airway inflammation, detect infection, or study metabolism; results depend on the method and usually complement rather than replace standard testing.

### **Breath Focus**

A mindfulness or relaxation exercise that directs attention to the feeling of breathing. It may help some people reduce stress or emotional arousal but does not replace treatment for a medical or mental-health condition.

#### **Alternate Names:**

Breath-Focused Meditation; Breathing-Focused Mindfulness

### **Breath Odor**

An unpleasant smell from the mouth or exhaled air. Common causes include dental or gum disease, dry mouth, tobacco, food, medicines, and infection; persistent odor requires assessment by a dentist or clinician.

### **Breath Sounds**

Sounds made as air moves through the airways and lungs, heard with a stethoscope. Reduced, absent, or unusual sounds can provide clues about mucus, narrowing, fluid, collapse, or other lung problems.

### **Breath Stacking**

A technique in which several small breaths are taken without fully exhaling, increasing the amount of air in the lungs. It may improve cough strength in selected people with weak breathing muscles and is

taught by a professional.

### **Breath Test**

Analysis of exhaled air to detect a substance made by the body or by microorganisms. Examples include tests for *Helicobacter pylori* and tests measuring hydrogen or methane after a sugar drink to assess digestion and absorption.

### **Breath-Holding Spell**

A brief involuntary episode in a young child, usually after crying, pain, fear, or anger, in which breathing pauses and the child may change colour or briefly faint before recovering.

### **Breath-Holding Spells**

Brief involuntary episodes, usually between about six months and six years, triggered by upset, pain, or fear. A child may cry, exhale, change colour, stiffen or faint briefly, then recover; a first or atypical episode needs assessment.

### **Breathalyzer**

A device that estimates alcohol concentration in the blood by measuring alcohol in exhaled air. Results can be affected by timing, device accuracy, recent drinking, mouth alcohol, and sampling conditions.

### **Breathing**

Movement of air into and out of the lungs. It brings oxygen into the body and removes carbon dioxide through coordinated activity of the brain, nerves, muscles, and airways.

### **Breathing Difficulty**

Uncomfortable or difficult breathing, also called breathlessness. Causes include lung or heart disease, airway blockage, anaemia, infection, anxiety, and metabolic illness; sudden, severe, or worsening difficulty needs urgent assessment.

### **Breathing Difficulty First Aid**

Immediate emergency supportive care provided to an individual experiencing acute respiratory distress, including positioning the patient upright in

---

a tripod or sitting posture, loosening restrictive clothing, assisting with prescribed bronchodilator or epinephrine autoinjector administration, and promptly contacting emergency medical services.

**Alternate Names:**

First Aid for Dyspnea; Emergency Respiratory Support; Respiratory Distress Management

**Breathing Exercises**

Deliberate changes in breathing pattern used to support relaxation, lung expansion, airway clearance, breath control, or rehabilitation. The safest technique depends on the person's condition and is taught when needed.

**Breathlessness**

Alternate name; see *Dyspnea*.

**Breech Baby**

A fetus positioned in the womb with the buttocks or feet pointing downward toward the birth canal instead of the head. Most babies turn into a head-down (vertex) position before late pregnancy, but those remaining breech near term may be delivered by cesarean section or planned vaginal breech birth, or turned via external cephalic version.

**Alternate Names:**

Breech fetus; Breech presentation.

**Breech Birth**

Birth in which the baby is positioned bottom or feet first rather than head first. The safest delivery plan depends on the exact position, pregnancy stage, baby and parent factors, and the team's experience.

**Breech Delivery**

Birth when the fetus presents buttocks or feet first rather than head first. The safest plan depends on fetal type and size, gestational age, maternal and pelvic factors, provider skill, and whether vaginal breech birth or cesarean delivery is appropriate.

**Breech Presentation**

A fetal position in which the buttocks or feet point toward the birth canal. Types include frank, com-

plete, and footling breech; risks include cord prolapse, difficulty delivering the head, and birth injury, so delivery needs careful planning.

**Alternate Names:**

Breech position; Malpresentation.

**Bregma**

The point on the skull where the main seam between the frontal and parietal bones meets the seam between the two parietal bones. In infants it lies beneath the soft spot at the front of the skull.

**Bremsstrahlung**

Radiation produced when fast electrons slow down or change direction near matter. It is relevant to some imaging and radiation treatments and requires appropriate shielding and safety controls.

**Brenner Tumor**

A rare ovarian growth made of cells resembling those lining the urinary tract. Most are small, one-sided, noncancerous, and found incidentally, but uncommon borderline or cancerous forms also occur.

**Brenner Tumour**

A usually noncancerous ovarian growth whose cells resemble those lining the urinary tract. Rare borderline and cancerous forms occur and are distinguished by tissue examination.

**Breslau Bacillus**

An older name for *Salmonella Typhimurium*, a bacterium that can cause foodborne gastroenteritis with diarrhoea, abdominal cramps, fever, and vomiting.

**Breslow Thickness**

The depth, measured in millimetres, to which a melanoma has grown into the skin. It is an important part of staging and helps estimate the risk of spread and guide treatment.

**Bretylum Tosylate**

An older medicine that can suppress certain dangerous abnormal heart rhythms, especially ventricular fibrillation. It is now rarely used because of limited

---

availability, side effects, and newer treatments.

### **Brevibacillus**

A group of spore-forming bacteria found in soil, water, and food. Human infection is uncommon but can occur opportunistically, especially in people with weakened immunity or medical devices.

### **Brevibacterium**

A group of bacteria found on skin, in soil, and in food. They commonly contribute to body or cheese odour; rare species can cause opportunistic infection, particularly in people with serious illness or devices.

### **Brevibacterium Casei**

A species of *Brevibacterium* found in the environment and occasionally on people. It rarely causes bloodstream or device-related infection, usually in a person with major underlying illness or an indwelling device.

### **Brevundimonas**

A group of environmental water-associated bacteria that rarely cause human disease. Opportunistic infections, including bloodstream or wound infection, are more likely in people with weakened immunity or medical devices.

### **Brevundimonas Diminuta**

An environmental water-associated bacterium that rarely causes opportunistic human infection. Reported illnesses include bloodstream, wound, or device-related infection, mainly in people with serious underlying disease.

### **Brevundimonas Vesicularis**

An environmental bacterium that rarely causes human infection. When it is isolated from a clinical sample, symptoms, specimen quality, and other results determine whether it represents true disease or contamination.

### **Bricklayer's Disease**

An older occupational term for lung disease caused by breathing dust during masonry work, especially

silica-containing dust. Long-term exposure can cause silicosis, which scars the lungs and impairs breathing.

### **Bridge To Transplant**

Temporary treatment or mechanical support that keeps a person with advanced heart failure alive and stable until a heart transplant can be performed. The device, timing, and risks depend on heart function and overall health.

### **Bridge Work**

A fixed dental replacement for one or more missing teeth. Artificial teeth are attached to prepared neighbouring teeth or implants, restoring some chewing function and appearance.

### **Brief Psychotic Disorder**

A sudden episode of delusions, hallucinations, severely disorganized speech, or markedly disorganized behavior lasting at least one day but less than one month, followed by a return toward usual functioning.

#### **Alternate Names:**

Brief Reactive Psychosis; Acute Transient Psychosis

### **Brief Resolved Unexplained Event**

A sudden, brief (under 1 minute), fully resolved episode in an infant under one year of age characterized by one or more of: cyanosis or pallor, absent, decreased, or irregular breathing, marked change in tone (hypertonia or hypotonia), and altered level of responsiveness, with no explanatory underlying cause identified following a thorough history and physical examination.

#### **Alternate Names:**

BRUE; Apparent Life-Threatening Event; ALTE

### **Brill's Disease**

Recurrence of epidemic typhus years after the first infection because dormant *Rickettsia prowazekii* becomes active again. It is usually milder than the original illness but can still be spread by body lice.



---

### Brill-Zinsser Disease

Recurrent epidemic typhus caused by reactivation of dormant *Rickettsia prowazekii* years after the original infection. The illness is usually milder but can be transmitted by body lice.

### Brimonidine Tartrate

A medicine that reduces certain nerve signals and blood-vessel activity. Eye drops lower pressure inside the eye, and skin preparations can reduce some facial redness; side effects and interactions may require medical attention.

### Brine Bath

Bathing in salty water, sometimes used for comfort or certain skin conditions. It does not treat every skin problem, and hot water or salt can worsen irritation, dry skin, wounds, or some medical conditions.

### Brittle Asthma

An uncommon term for severe asthma with sudden, unpredictable, or difficult-to-control attacks despite appropriate treatment. Specialist review is needed to confirm asthma, identify triggers or other diagnoses, and reduce risk.

### Brittle Bone Disease

An inherited disorder, usually osteogenesis imperfecta, in which bones break easily. It can also cause short stature, loose joints, blue-gray eye whites, hearing loss, dental problems, or other skeletal changes.

### Brittle Nail Syndrome

A pattern of nails that split, peel, crack, or break easily, often because of repeated wetting and drying, detergents, trauma, ageing, or inflammatory nail disease. Iron deficiency, thyroid disease, fungal infection, and inherited disorders can contribute; persistent, painful, or widespread changes need assessment.

### Brivudine

An antiviral medicine used in some countries for shingles. It is not used with fluoropyrimidine cancer

medicines or certain related treatments because the interaction can be fatal.

### Broad Casts

Wide, waxy or granular urinary microscopic casts that form in dilated, atrophic collecting ducts of chronically damaged nephrons with sluggish urinary flow. Their presence in the urine sediment is a hallmark of advanced chronic kidney disease and end-stage renal disease.

#### Alternate Names:

Renal Failure Casts; Wide Casts

### Broad Ligament

A sheet of tissue extending from the sides of the uterus to the pelvic walls. It helps support the uterus, fallopian tubes, and ovaries and contains blood vessels, nerves, and connective tissue.

### Broad Nasal Bridge

An unusually wide upper part of the nose between the eyes. It may be a normal family feature or part of a condition present from birth or an inherited syndrome.

### Broad-Based Gait

An abnormal walking pattern in which the feet are placed abnormally far apart to increase the base of support and maintain balance. It is a cardinal clinical physical sign of midline cerebellar vermis lesions (cerebellar ataxia) or severe loss of lower limb proprioception (sensory ataxia).

### Broad-Complex Tachycardia

A fast heart rhythm with wide QRS beats on an ECG, usually 120 milliseconds or longer. It may come from the lower heart chambers or from an upper-chamber rhythm conducted abnormally. Because it may be dangerous, it is generally treated as ventricular tachycardia until proven otherwise.

#### Alternate Names:

Wide-Complex Tachycardia

### Broca Aphasia

A language disorder in which speech is slow, effortful,



---

and difficult to produce, while understanding may be relatively preserved. It usually follows damage to the dominant frontal language area, often from a stroke.

### **Broca's Aphasia**

A nonfluent language disorder caused by damage to the dominant frontal language region. Speech is effortful and limited, while understanding may be better preserved than speaking or repeating words.

### **Broca's Area**

A region in the dominant frontal lobe that helps plan and produce spoken language. Damage can cause slow, effortful speech and difficulty forming words, while understanding may be relatively preserved.

### **Brodie Abscess**

A chronic, well-circumscribed subacute pyogenic bone abscess that represents a localized form of osteomyelitis. It characteristically develops in the metaphysis of long bones (most frequently the proximal or distal tibia) in children and young adults, presenting as a lytic bone lesion with a surrounding ring of sclerotic bone on radiography.

### **Brodie's Abscess**

A localized, slowly developing pocket of bone infection, often near the end of a long bone. It commonly causes persistent, sometimes night-time, bone pain with little fever and may require prolonged antibiotics or surgery.

### **Brodman's Area**

A region of the cerebral cortex anatomically defined and numbered (1 through 52) based on its distinct cellular architecture (cytoarchitectonics), with specific areas corresponding to distinct functional neurological roles (such as area 4 for primary motor cortex).

### **Brody Disease**

A rare inherited muscle disorder causing painless stiffness and slow muscle relaxation after strenuous or repeated exercise. It often begins in childhood and may affect the face, eyelids, or hands; symptoms

can resemble myotonia but arise from a different muscle problem.

#### **Alternate Names:**

Brody Myopathy; SERCA1-Related Muscle Disease

### **Broken Ankle**

A break in one or more bones forming the ankle joint. It usually causes pain, swelling, bruising, and difficulty bearing weight; deformity, an open wound, numbness, or poor circulation requires emergency care.

### **Broken Arm**

A break in one or more arm bones, usually causing pain, swelling, bruising, deformity, or loss of function. An open injury, severe deformity, numbness, or reduced circulation requires urgent care.

### **Broken Blood Vessel in Eye**

Alternate name; see *Subconjunctival Hemorrhage*.

### **Broken Bone**

A partial or complete break in a bone caused by injury, repeated stress, or weakened bone. Pain, swelling, bruising, deformity, or loss of function may occur. Examination and imaging are often needed, but some fractures are not visible on the first X-ray.

### **Broken Clavicle**

Alternate name; see *Broken Collarbone*.

### **Broken Collarbone**

A break in the clavicle, usually after a fall or direct blow. It causes pain, swelling, tenderness, and reduced shoulder movement; numbness, breathing difficulty, or a skin-threatening deformity requires urgent care.

#### **Alternate Names:**

Broken Clavicle

### **Broken Finger**

A break in a finger bone caused by a blow, crush, or twist. Pain, swelling, bruising, deformity, and reduced movement may occur; assessment checks alignment, tendon function, sensation, and circula-

---

tion and guides splinting or surgery.

**Alternate Names:**

Finger Fracture

**Broken Foot**

A break in one or more foot bones after a fall, blow, twist, or repeated stress. Pain, swelling, bruising, and difficulty walking are common; deformity, an open wound, numbness, or inability to walk needs urgent assessment.

**Broken Hand**

A break in one or more hand bones, causing pain, swelling, bruising, deformity, or reduced movement. Assessment includes alignment, tendon function, sensation, and circulation because some breaks require reduction or surgery.

**Broken Heart Syndrome**

Temporary weakening of the heart muscle, often after severe emotional or physical stress. It can resemble a heart attack with chest pain or breathlessness, but the coronary arteries usually have no blocking clot.

**Broken Hip**

Alternate name; see *Hip Fracture*.

**Broken Leg**

A break in one or more leg bones, usually causing pain, swelling, bruising, deformity, or inability to bear weight. An open injury, numbness, or loss of circulation requires emergency care.

**Alternate Names:**

Leg Fracture

**Broken Little Finger**

Alternate name; see *Broken Finger*.

**Broken Nose**

A break in one or more nasal bones that may cause pain, swelling, bruising, bleeding, deformity, or difficulty breathing through the nose. Persistent bleeding, clear fluid leakage, or severe deformity needs urgent assessment.

**Broken Or Dislocated Jaw**

A break or displacement of the jaw that can cause pain, swelling, an altered bite, limited movement, loose teeth, or difficulty speaking, swallowing, or breathing. It needs urgent medical and dental assessment.

**Broken Or Knocked Out Tooth**

A tooth damaged or completely forced out by injury. A knocked-out permanent tooth is a dental emergency; the tooth is handled by the crown, kept moist, and the root is not scrubbed.

**Broken Ribs**

Breaks in one or more ribs, usually causing local chest pain that worsens with breathing, coughing, or movement. Severe breathing difficulty, increasing pain, coughing blood, or major injury requires urgent assessment.

**Broken Toe**

A break in a toe bone, usually after a direct blow or crush. Pain, swelling, bruising, and tenderness are common; deformity, an open wound, numbness, or poor circulation needs prompt assessment.

**Alternate Names:**

Toe Fracture; Phalanx Fracture

**Broken Wrist**

A break in one or more bones near the wrist, often causing pain, swelling, bruising, deformity, or reduced hand function. Numbness, pale or cold fingers, an open wound, or severe deformity needs urgent care.

**Bromelain**

A mixture of enzymes from pineapple. It is sold as a supplement and studied for pain and inflammation, but evidence and product strength vary; it can cause allergy, stomach upset, or interactions with medicines.

**Bromhexine**

A medicine that thins sticky mucus in the airways so it may be coughed up more easily. It can cause

---

stomach upset or rash and rarely causes a serious skin reaction.

### **Bromhidrosis**

Unusually strong or unpleasant body odour produced when skin microorganisms break down apocrine or eccrine sweat and other secretions. It commonly affects the armpits, feet, or groin and may be worsened by hyperhidrosis, occlusion, infection, or retained skin debris; treatment addresses hygiene, moisture, and any underlying cause.

### **Bromides**

Salts or compounds containing bromide. Excessive or prolonged exposure can cause bromism, with symptoms affecting the nervous system, mood, digestion, or skin.

### **Bromine**

A reactive chemical used in industry and some flame-resistant materials. Bromine and many bromine compounds can irritate or burn tissue, while substantial exposure can cause poisoning affecting several organs.

### **Bromism**

Poisoning caused by prolonged or excessive exposure to bromide compounds. It may cause confusion, drowsiness, poor coordination, mood or behaviour changes, digestive symptoms, acne-like eruptions, or other neurologic problems.

### **Bromobenzenes**

Industrial chemicals made by attaching one or more bromine atoms to benzene. Exposure can irritate the eyes, skin, and airways or cause other toxic effects, depending on the specific compound and amount.

### **Bromocriptine**

A medicine that stimulates dopamine receptors and reduces prolactin release. It is used for high prolactin and selected movement or hormone disorders; nausea, dizziness, low blood pressure, hallucinations, and other effects may occur.

### **Bromoderma**

An inflammatory skin eruption linked to bromine or bromide exposure. It may appear as acne-like, pustular, or other irritated lesions and usually improves when the source is identified and removed.

### **Bromodosis**

Very unpleasant foot odour caused when sweat, moisture, and skin debris are broken down by bacteria or fungi. Washing, drying, breathable footwear, and treatment of any infection may help.

### **Bromophos**

An organophosphate insecticide. Exposure can overstimulate nerves and cause sweating, excess saliva, vomiting, breathing difficulty, weakness, seizures, or collapse and requires urgent poison or emergency care.

### **Brompheniramine Overdose**

Poisoning from taking too much brompheniramine, an antihistamine. It may cause severe sleepiness or agitation, confusion, dry mouth, rapid heartbeat, seizures, or dangerous heart rhythms; urgent help is needed.

### **Bromthymol Blue**

A chemical indicator that changes colour according to acidity and alkalinity. It is used in laboratories and is not a medicine or treatment.

### **Bronchi**

The two main airways carrying air from the windpipe into the lungs. Each divides into smaller branches; the right main bronchus is wider and more vertical, so inhaled objects are more likely to enter it.

### **Bronchial**

Relating to the bronchi, the large airways branching from the windpipe. Bronchial disease may cause cough, wheezing, mucus production, narrowing, or reduced airflow.

---

## Bronchial Artery

An artery from the body's general circulation that supplies the walls of the bronchi and supporting lung tissues. It differs from the pulmonary arteries, which carry blood to the lungs for oxygen exchange.

## Bronchial Artery Embolization

A catheter procedure that blocks bleeding arteries supplying the airways, usually to control severe or repeated coughing of blood. Imaging guides the catheter and the material used; careful technique is needed to avoid blocking arteries supplying the spinal cord.

### Alternate Names:

BAE; Bronchial Embolization; Embolization of Bronchial Arteries

## Bronchial Asthma

See *Asthma*.

## Bronchial Biopsy

Removal of a small tissue sample from an airway during bronchoscopy. Laboratory examination can investigate inflammation, infection, abnormal growths, or cancer; bleeding and breathing complications are possible.

## Bronchial Breath Sounds

Harsh, high-pitched, hollow breath sounds with a prolonged expiratory phase and an audible pause between inspiration and expiration. Normally auscultated over the trachea and main bronchi, their presence over peripheral lung fields indicates abnormal transmission through consolidated lung tissue, as in lobar pneumonia.

## Bronchial Carcinoid

A usually slow-growing neuroendocrine tumour arising in an airway. Typical carcinoids are generally less aggressive than atypical carcinoids, but either can block an airway, cause coughing or bleeding, or spread to other sites.

## Bronchial Carcinoma

See *Lung Cancer*.

## Bronchial Circulation

The systemic arterial blood supply providing oxygenated blood and nutrients to the conducting airways, supporting stroma, and visceral pleura of the lungs, arising from bronchial arteries branching off the thoracic aorta or intercostal arteries.

## Bronchial Fistula

An abnormal passage between an airway and another structure, such as the space around the lung, food pipe, or skin. It can cause persistent air leakage, coughing, infection, or breathing failure.

## Bronchial Hemorrhage

Bleeding from airway blood vessels, usually seen as coughing up blood. Heavy bleeding can block the airway or cause breathing failure and requires emergency treatment.

## Bronchial Hyperresponsiveness

An abnormal state of airway sensitivity in which conducting airways undergo exaggerated bronchoconstriction in response to low concentrations of non-allergic chemical stimuli (such as methacholine or histamine) or physical triggers; a physiological hallmark of asthma.

### Alternate Names:

Airway Hyperreactivity

## Bronchial Obstruction

Partial or complete blockage of an airway by mucus, an inhaled object, a tumour, swelling, or pressure from outside. It can reduce airflow and lead to lung collapse, repeated infection, wheezing, or breathlessness.

## Bronchial Stenosis

Narrowing of an airway caused by scarring, inflammation, infection, injury, a tumour, or a previous procedure. It can cause localised wheezing, breathlessness, repeated infection, or poor mucus clearance and may require widening or surgery.

## Bronchial Stricture

Abnormal narrowing of an airway caused by scar-

---

ring, inflammation, infection, a tumour, or previous instrumentation. It may produce localised wheezing, cough, repeated infection, or collapse of the lung beyond the narrowing.

### **Bronchial Tree**

The branching network of airways from the windpipe through the main, lobar, and smaller bronchi and bronchioles toward the air sacs. It conducts and distributes air throughout the lungs.

### **Bronchial Tubes**

The two main bronchi and their branches, which carry air from the windpipe into the lungs and divide into progressively smaller airways.

#### **Alternate Names:**

Bronchi; Bronchial Tree When Referring to the Branching Airway System

### **Bronchial Vein**

A vein that drains blood from the walls of the larger airways and nearby lung tissue. It empties partly into veins returning blood to the heart and partly into pulmonary veins.

### **Bronchiectasis**

A long-term condition in which parts of the airways become permanently widened and damaged. The airways may not clear mucus well, so mucus builds up and can lead to a persistent cough, repeated chest infections, and breathlessness. Causes include severe or repeated infections, other lung or immune conditions, and airway blockage; sometimes the cause is unknown. Some people also cough up blood.

### **Bronchiole**

A small airway without cartilage that branches from a bronchus and carries air toward the lung's air sacs. Its smooth muscle can narrow or widen the passage and regulate airflow.

### **Bronchiolectasis**

Permanent widening of the small airways, usually caused by chronic inflammation, infection, or blockage. It may occur with bronchiectasis and contribute to poor mucus clearance and repeated infec-

tion.

### **Bronchioles**

Small branching airways that carry air from the bronchi toward the air sacs and regulate airflow with their surrounding muscle.

### **Bronchiolitis**

Inflammation and blockage of the smallest airways, usually from a viral infection in infants and young children. It can cause cough, wheezing, fast breathing, chest retractions, poor feeding, and low oxygen.

### **Bronchiolitis Obliterans**

A rare disease in which inflammation and scarring permanently narrow or close the smallest airways. It can follow severe infection, inhaled injury, autoimmune disease, or transplantation and causes persistent cough and breathlessness.

### **Bronchitis**

Inflammation of the larger airways. Acute bronchitis is usually caused by a virus and causes cough, with or without mucus, while chronic bronchitis means mucus-producing cough for at least three months in two successive years.

### **Bronchoalveolar Lavage**

A bronchoscopy procedure in which sterile fluid is put into a small lung area and then collected. The sample can be tested for infection, inflammation, abnormal cells, or other lung disease.

### **Bronchocele**

Abnormal widening of a bronchus filled with mucus, usually because an airway is blocked or developed abnormally. It may cause cough, repeated infection, or no symptoms and is assessed with imaging.

### **Bronchocentric Granulomatosis**

A rare, non-necrotizing or necrotizing granulomatous inflammatory disorder centered primarily on conducting bronchi and bronchioles, frequently associated with tissue eosinophilia and severe asthma or allergic bronchopulmonary aspergillosis.

---

## Bronchoconstriction

Narrowing of the airways because their muscle tightens, the lining swells, or mucus builds up. It increases resistance to airflow and commonly occurs in asthma, allergic reactions, or irritant exposure.

## Bronchoconstrictor Agents

Substances that narrow the airways by tightening airway muscle or causing inflammation. Examples include histamine, leukotrienes, irritant gases, and some medicines.

## Bronchodilator

A medicine that relaxes muscles around the airways, widening them and making breathing easier. It can relieve or prevent wheezing and breathlessness in asthma, chronic obstructive pulmonary disease, and other conditions with narrowed airways.

### Alternate Names:

Bronchodilators

## Bronchodilator Response

The change in airflow measured by spirometry after a bronchodilator is given. Improvement can support variable airway narrowing, but no response does not exclude asthma and a response does not prove one specific disease.

## Bronchodilator Reversibility

Improvement in measured airflow after an inhaled bronchodilator. It can support variable airway narrowing, but one result alone cannot confirm or exclude asthma or chronic obstructive pulmonary disease.

## Bronchoesophageal Fistula

An abnormal connection between an airway and the food pipe. It may be present from birth or develop from cancer, infection, injury, or treatment and can cause coughing during swallowing, repeated pneumonia, or breathing problems.

## Bronchogenic Carcinoma

See *Lung Cancer*.

## Bronchogenic Cyst

A birth-related fluid-filled cyst that develops from early airway or food-pipe tissue, often near the wind-pipe or central chest. It may cause no symptoms or press on nearby structures and sometimes requires removal.

## Bronchomalacia

Weakness of an airway wall that allows it to collapse too much, especially during breathing out. It may cause noisy breathing, cough, breathlessness, repeated infections, or difficulty clearing mucus.

## Bronchophony

An examination finding in which spoken sounds are heard unusually clearly through a stethoscope over part of the lung. It can occur when lung tissue is solidified by infection or another process but is not diagnostic by itself.

## Bronchopleural Fistula

An abnormal connection between an airway and the space around the lung, often after lung surgery or severe infection. It can cause a persistent air leak, breathlessness, infection, or air under the skin.

## Bronchopneumonia

Infection causing scattered areas of inflammation and consolidation around the small airways in one or more lung lobes. It may cause fever, cough, mucus, chest pain, fast breathing, and low oxygen.

## Bronchoprovocation Test

A breathing test used to look for airways that narrow too easily, often when asthma is suspected but routine spirometry is normal. A substance such as methacholine is inhaled in increasing amounts while breathing is measured. A significant fall in airflow supports airway sensitivity, but does not by itself prove asthma.

### Alternate Names:

Methacholine Challenge Test; Bronchial Provocation Test



---

## Bronchopulmonary Dysplasia

A chronic lung disease in some premature infants whose developing lungs have been affected by immaturity, inflammation, oxygen, or breathing support. It may cause ongoing oxygen need, fast breathing, or wheezing.

## Bronchopulmonary Segments

Natural sections of a lung lobe, each supplied by its own airway branch and pulmonary artery branch. Their separation helps doctors describe disease and allows some operations to remove only an affected segment.

## Bronchoscope

A thin tube with a light and camera used to examine the windpipe and airways. Instruments passed through it can remove an object or collect tissue, fluid, or mucus for testing.

## Bronchoscopic Culture

Laboratory testing of fluid or tissue collected during bronchoscopy to look for bacteria, fungi, or mycobacteria when sputum or other ordinary samples are not sufficient.

## Bronchoscopy

Examination of the windpipe and airways with a flexible or rigid camera tube. It can investigate persistent cough, infection, bleeding, or a lung mass and can remove foreign material or collect samples; sedation or anaesthesia is used.

## Bronchospasm

Sudden tightening of airway muscles that narrows the breathing passages. It can cause wheezing, cough, chest tightness, and difficulty breathing and may be triggered by asthma, allergy, infection, irritants, or airway procedures.

## Bronchostenosis

Narrowing of a bronchus that restricts airflow. Causes include scarring, inflammation, infection, a tumour, an inhaled object, pressure from outside, or a previous procedure and may cause wheezing or repeated infection.

## Bronchus

One of the two main branches of the windpipe that carries air into a lung. Each divides into smaller branches; the right main bronchus is wider and more vertical, making it a more common site for inhaled objects to lodge.

## Bronchus Disease

Any condition affecting one or more bronchi, the air passages leading into the lungs. Infection, asthma, chronic inflammation, blockage, scarring, or a tumour may reduce airflow and cause cough, wheezing, or repeated infection.

## Bronchus Neoplasm

An abnormal growth arising in a bronchus. It may be noncancerous or cancerous; imaging, bronchoscopy, and tissue testing help determine its type, extent, and treatment.

## Brooke Disease

A rare inherited skin disorder involving abnormal thickening of the outer skin or hair follicles. Because the eponym has been used for more than one condition, the original source and clinical features are checked.

## Brooke-Spiegler Syndrome

An inherited disorder causing multiple usually non-cancerous skin tumours, often on the head and neck. It is commonly linked to changes in the CYLD gene and may include cylindromas, spiradenomas, or trichoepitheliomas.

## Broselow Tape

A colour-coded tape that estimates a child's weight from body length during an emergency. It helps select medicine doses, fluid amounts, equipment sizes, and defibrillator settings, but clinical assessment and an actual weight are used when available.

### Alternate Names:

Broselow Pediatric Emergency Tape; Pediatric Length-Based Resuscitation Tape

---

### **Brow Presentation**

A fetal position in which the forehead leads because the head is partly extended. The head presents a wide diameter, so persistent brow presentation at term may prevent vaginal birth and requires specialist delivery planning.

### **Brown Adipose Tissue**

A specialized type of adipose tissue characterized by multilocular lipid droplets, abundant mitochondria, and high levels of uncoupling protein 1 (UCP1, thermogenin), specialized for non-shivering thermogenesis in newborns and hibernating mammals.

**Alternate Names:**

Brown Fat; BAT

### **Brown Fat**

A type of body fat containing many energy-producing structures that generate heat. It is especially important for keeping newborns warm and can also be activated by cold in adults.

### **Brown Lung Disease**

An occupational breathing disorder caused by repeated exposure to cotton, flax, or hemp dust, traditionally called byssinosis. It can cause chest tightness, cough, wheezing, and work-related narrowing of the airways.

### **Brown Recluse Spider**

A venomous spider whose bite may initially be mild but can later cause painful skin damage, blistering, or tissue death. Rarely, its venom causes fever, red-cell destruction, kidney injury, or other systemic illness.

### **Brown Recluse Spider Bite**

Alternate name; see *Spider Bites*.

### **Brown Sequard Syndrome**

A spinal-cord injury affecting mainly one side. It typically causes weakness and loss of vibration or position sense on the injured side and loss of pain and temperature sensation on the opposite side below the injury.

### **Brown Stain**

A brown deposit on teeth caused by pigments from food, drinks, tobacco, or bacteria. Professional cleaning may remove surface stains, but decay or enamel disease needs separate treatment.

### **Brown Syndrome**

Limited ability to lift the eye when it is turned inward, because the superior oblique tendon cannot glide freely through its pulley, the trochlea. The affected eye appears lower than the other in that gaze direction and double vision may occur. It can be present at birth or develop from inflammation, injury, sinus disease or surgery, or rheumatic disease such as rheumatoid arthritis. Some cases resolve on their own; persistent double vision is treated by an eye specialist.

### **Brown Teeth**

Brown discolouration of teeth caused by surface stains, decay, enamel defects, excess fluoride, medicines, injury, or other conditions. A dental examination identifies the cause and appropriate treatment.

### **Brown-Sequard Syndrome**

A neurological syndrome caused by hemisection or unilateral damage to the spinal cord, characterized by ipsilateral loss of motor function (corticospinal tract) and proprioception/vibration (dorsal column), and contralateral loss of pain and temperature sensation (spinothalamic tract) below the lesion level.

### **Brown-Séquard Syndrome**

A spinal-cord injury affecting mainly one side. It causes weakness and loss of position or vibration sense on the injured side and loss of pain and temperature sensation on the opposite side below the injury.

### **Bruce Protocol**

A graded treadmill test in which speed and slope increase at set stages. It assesses exercise capacity, symptoms, heart-rate and blood-pressure responses, and signs of reduced blood flow to the heart.

---

## Brucella

A group of bacteria that usually infect animals but can infect people through unpasteurised dairy products, inhalation, or contact with infected animals or tissues. Human infection causes brucellosis.

## Brucella Abortus

A Brucella species mainly associated with cattle. It can infect people exposed to infected animals, tissues, or unpasteurised dairy products and cause prolonged fever and other brucellosis symptoms.

## Brucella Canis

A Brucella species associated with dogs. It rarely infects people after contact with infected dogs or reproductive material and may cause prolonged fever or other brucellosis symptoms.

## Brucella Melitensis

A Brucella species mainly associated with goats and sheep and a frequent cause of severe human brucellosis, usually acquired through unpasteurised dairy products or infected animals.

## Brucella Suis

A Brucella species associated with pigs and some wildlife. It can infect people through animal contact or contaminated food and may cause prolonged illness or infection in bones and organs.

## Brucellaceae

A family of bacteria that includes Brucella and related animal-associated species. Some members cause brucellosis, an infection that can pass from animals to people.

## Brucellosis

An infection caused by Brucella bacteria, usually acquired from infected animals, animal tissues, or unpasteurised dairy products. It may cause prolonged fever, sweats, tiredness, headache, and muscle or joint pain.

## Bruch's Membrane

A thin supporting layer between the retina and the

blood-rich tissue behind it. It helps nourish and maintain the retina and can become damaged in ageing, diabetes, and some retinal diseases.

## Bruck's Syndrome

A rare inherited disorder combining fragile bones with joint stiffness or fixed contractures present from birth. It results from changes affecting connective tissue and may cause repeated fractures and limited movement.

## Brudzinski Sign

Involuntary bending of the hips and knees when the neck is gently bent forward. It can occur with irritation of the tissues around the brain and spinal cord, including meningitis, but does not confirm the diagnosis.

## Brugada ECG Pattern Type 1

A characteristic ECG pattern with raised ST segments and an inverted T wave in certain right-sided chest leads. It may occur with Brugada syndrome, fever, medicines, or other conditions. The ECG pattern alone does not establish the inherited syndrome or predict the same risk in every person.

### Alternate Names:

Type 1 Brugada Pattern; Coved ST Elevation Pattern

## Brugada Phenocopy

A temporary ECG pattern resembling Brugada syndrome but caused by another problem, such as fever, abnormal blood salts, reduced heart blood flow, medicines, or poisoning. The pattern resolves when the cause is corrected and is distinguished from inherited disease.

### Alternate Names:

Acquired Brugada Pattern; BrP

## Brugada Syndrome

An inherited heart-rhythm disorder that can cause dangerous ventricular arrhythmias and sudden cardiac death, often during rest, sleep, or fever. Diagnosis relies on a characteristic ECG pattern and clinical assessment.

---

## Brugia

A group of parasitic worms spread by mosquitoes that can cause lymphatic filariasis. The infection may damage lymph vessels and lead to recurrent fever, swelling, or long-term enlargement of limbs or genital tissues.

## Bruise

Bleeding under unbroken skin after small blood vessels are damaged, usually by a blow, pressure, or injury. It commonly changes colour as it heals; unexplained or frequent bruising needs medical assessment.

## Bruises

Areas of bleeding beneath the skin after small blood vessels are damaged. Frequent, large, spontaneous, painful, or unusually persistent bruises may result from medicines, platelet or clotting problems, liver disease, nutritional deficiency, or injury.

## Bruising

Purplish, blue, green, or yellow skin discolouration caused by blood leaking into tissue, usually after injury. Colour changes as the blood is broken down; widespread or unexplained bruising needs assessment.

## Bruit

An abnormal whooshing sound heard over a blood vessel, usually with a stethoscope, caused by turbulent blood flow. It may suggest narrowing but does not by itself establish the cause or severity.

## Brunner Gland

A gland in the first part of the small intestine that releases mucus and alkaline fluid. These secretions protect the intestinal lining and help neutralise acid arriving from the stomach.

## Brunner Glands

Compound branched tubuloalveolar submucosal glands located exclusively in the duodenum, concentrated primarily in the bulb and proximal descending portions. They secrete an alkaline, bicarbonate-rich, mucin-containing fluid that protects the duode-

nal mucosa from gastric acid erosion and optimizes luminal pH for pancreatic digestive enzymes.

### Alternate Names:

Duodenal Submucosal Glands

## Brunner's Gland

A mucus- and bicarbonate-producing gland in the duodenum, the first part of the small intestine. Its alkaline secretion protects the lining from stomach acid and supports digestion.

## Brunner's Glands

Glands in the duodenum that release mucus and bicarbonate-rich fluid to protect the intestinal lining from stomach acid and help create a suitable environment for digestive enzymes.

## Bruns Syndrome

Episodic, sudden attacks of severe headache, vomiting, vertigo, and syncope triggered by abrupt neck movements, caused by transient obstruction of cerebrospinal fluid flow in the fourth ventricle by a mobile mass, such as a cysticercal cyst or ependymoma.

## Brush Burns

Superficial friction injuries caused by scraping skin against a rough surface. They cause pain, redness, and grazes; cleaning and protecting the area usually allow healing, but deep, extensive, or infected injuries need medical care.

## Brushfield's Spots

Small gray or white spots arranged around the coloured part of the eye. They are common in people with Down syndrome but can also occur in healthy people and usually do not affect vision.

## Bruton Agammaglobulinemia

An X-linked primary immunodeficiency disorder caused by mutations in the Bruton tyrosine kinase (*BTK*) gene, resulting in failure of pre-B cells to differentiate into mature B lymphocytes, profound pan-hypogammaglobulinemia, and recurrent pyogenic bacterial infections beginning after maternal antibodies wane.

---

**Alternate Names:**

X-Linked Agammaglobulinemia; XLA

**Bruxism**

Repeated clenching or grinding of the teeth, often during sleep. It can wear or fracture teeth and cause jaw, face, ear, or headache pain; dental review can assess damage and possible sleep, stress, or medicine-related causes.

**Alternate Names:**

Teeth Grinding; Tooth Clenching

**Bryant's Traction**

An older treatment in which an infant's or young child's legs are gently elevated with traction to treat selected hip or thigh conditions. It is rarely used and requires careful specialist monitoring.

**Bubo**

A tender, enlarged lymph node, usually in the groin, armpit, or neck, caused by infection or inflammation. Buboes are classically linked with plague but also occur in sexually transmitted and other infections.

**Bubonic Plague**

A serious infection caused by *Yersinia pestis*, usually spread by infected flea bites. It causes fever and painful swollen lymph nodes and can spread to the blood or lungs without prompt antibiotic treatment.

**Buccal**

Relating to the cheek or its inner lining. A buccal medicine is placed between the cheek and gum so it dissolves and is absorbed through the mouth.

**Buccal Gland**

A small salivary gland in the cheek that releases mucus into the mouth. The secretion helps keep the cheek lining moist and supports chewing and swallowing.

**Buccal Mucosa**

The non-keratinized stratified squamous mucous membrane lining the inside of the cheeks, bounded by the labial commissures anteriorly, the retromolar

trigone posteriorly, and the alveolar sulci superiorly and inferiorly. It contains the opening of the parotid duct (Stensen duct) opposite the maxillary second molar and serves as a site for clinical examination of exanthems (such as Koplik spots in measles), oral lichen planus (Wickham striae), aphthous stomatitis, leukoplakia, and drug absorption.

**Buccal Smear**

A sample of cells gently collected from the inside of the cheek for laboratory testing. It may be used for genetic analysis or, less commonly, to investigate infection or abnormal cells.

**Buccalmucosa**

A spelling variant of buccal mucosa, the moist lining on the inside of the cheeks.

**Buccinator**

A muscle in the cheek that presses the cheek against the teeth. It helps keep food between the teeth and assists sucking, blowing, chewing, and speaking.

**Buckle Fracture**

Alternate name; see *Torus Fracture*.

**Budd-Chiari Syndrome**

Blockage of blood leaving the liver through the hepatic veins, sometimes extending to the large vein that carries blood to the heart (inferior vena cava). It is most often caused by blood clots and may cause upper-abdominal pain, an enlarged liver, fluid in the abdomen, jaundice, or liver failure.

**Alternate Names:**

Chiari-Budd Syndrome

**Budesonide**

A corticosteroid that reduces inflammation. It is used in selected inhaled, nasal, digestive, or rectal treatments for conditions such as asthma, chronic obstructive lung disease, allergic symptoms, and inflammatory bowel disease; side effects depend on the route and dose.

---

## **Budvicia**

A group of water-associated bacteria related to *Yersinia*. They rarely cause human disease, but some isolates have been associated with urinary or other opportunistic infections.

## **Budvicia Aquatica**

A water-associated bacterium that has rarely been isolated from human urinary infections, including infections in people with urinary catheters. Its significance depends on symptoms and specimen quality.

## **Buerger Disease**

An inflammatory disease that blocks small and medium blood vessels, especially in the hands and feet. It is strongly linked to tobacco use and can cause pain, ulcers, poor circulation, and loss of tissue.

## **Buerger Test**

A bedside physical examination assessment used to evaluate the severity of peripheral arterial occlusive disease in the lower extremities. The patient's leg is elevated to 45 or 90 degrees until maximum pallor develops (Buerger's angle of circulatory sufficiency), and is then placed in a dependent sitting position; a prolonged delay in the return of pink skin color followed by a deep red or purple reactive erythema (rubor of dependency) indicates advanced arterial insufficiency.

## **Buerger's Disease**

A tobacco-associated inflammatory disease of small and medium arteries and veins, also called thromboangiitis obliterans. Reduced blood flow can cause pain, colour changes, ulcers, gangrene, and loss of fingers or toes.

## **Buerger's Disease**

Thromboangiitis obliterans, an inflammatory clotting disease of small and medium arteries and veins, strongly associated with tobacco or nicotine exposure. It can cause painful fingers or toes, ulcers, and tissue loss; complete nicotine cessation is essential.

## **Buffalo Hump**

A pad of fat at the back of the neck and upper back. It may occur with obesity, long-term corticosteroid excess, some HIV treatments, or Cushing syndrome and is assessed together with other findings.

## **Buffer**

A chemical solution that resists a change in acidity when small amounts of acid or base are added. Buffers help maintain stable conditions in blood, cells, medicines, and laboratory tests.

## **Buffer System**

A chemical solution consisting of a weak acid and its conjugate base (or a weak base and its conjugate acid) that resists changes in hydrogen ion concentration and pH upon the addition of small amounts of acidic or basic substances.

## **Buffy Coat**

The thin pale layer of white blood cells and platelets that forms between plasma and red blood cells when a blood sample is spun. It can be collected for laboratory testing.

## **Bufonidae**

The family of amphibians commonly called true toads. Some species produce bufotoxins in their skin glands; swallowing or absorbing these toxins can affect the heart and nervous system.

## **Buformin**

An older biguanide medicine for diabetes that has been withdrawn or restricted in many countries because it can cause dangerous lactic-acid buildup, especially when kidney function is poor.

## **Bug Spray Poisoning**

Illness after exposure to an insecticide. Effects depend on the chemical and may include skin or eye irritation, vomiting, breathing difficulty, unusual sweating, confusion, seizures, or other nervous-system problems; severe poisoning requires urgent medical treatment.



---

## **Bulbar**

Relating to the lower brainstem or to the nerves controlling swallowing, speaking, chewing, coughing, and movement of the tongue and soft palate. Bulbar weakness can allow food or saliva to enter the airway.

## **Bulbar Conjunctiva**

The part of the eye's clear moist covering that lies over the white of the eye. It joins the inner surface of the eyelids around the front of the eye and can become inflamed or injured.

## **Bulbar Palsy**

Weakness or paralysis of muscles controlled by lower cranial nerves. It can cause slurred speech, difficulty swallowing, weak coughing, poor handling of saliva, and sometimes breathing problems.

## **Bulbar Paralysis**

Weakness or paralysis of muscles used for speaking, swallowing, chewing, coughing, and moving the tongue or palate. It may result from brainstem, motor-neuron, or neuromuscular disease and can threaten breathing.

## **Bulbocavernosus**

A paired muscle in the pelvic floor that helps empty the urethra and contributes to erection and ejaculation. It surrounds structures at the base of the penis or the vaginal opening.

## **Bulbospongiosus**

A paired pelvic-floor muscle that helps empty the urethra and assists erection and the release of genital secretions during sexual activity.

## **Bulbourethral Gland**

One of two small glands below the prostate that release a slippery, alkaline fluid into the urethra during sexual arousal. The fluid helps lubricate the passage and reduce acidity.

## **Bulbourethral Glands**

Two small male reproductive glands, also called Cowper glands, that release mucus-like fluid into the

urethra during sexual arousal. The fluid lubricates the urethra and may reduce acidity.

## **Bulging**

Projecting or swelling outward beyond the usual shape. A bulge may involve a disc, membrane, organ, skin area, or body wall, and its significance depends on the structure and cause.

## **Bulging Eyes**

Alternate name; see *Exophthalmos*.

## **Bulging Fontanelle**

An outward protrusion or tense swelling of an infant's soft spot (fontanelle) when the baby is calm and upright, indicating elevated intracranial pressure from conditions such as meningitis, encephalitis, hydrocephalus, or intracranial hemorrhage.

### **Alternate Names:**

Tense Fontanelle; Fontanelle Protrusion; Raised Intracranial Pressure in Infants

## **Bulimia**

Alternate name; see *Bulimia Nervosa*.

## **Bulimia Nervosa**

An eating disorder involving recurrent binge-eating episodes, a sense of loss of control, and recurrent inappropriate compensatory behaviors such as vomiting, laxative misuse, fasting, or excessive exercise. Self-evaluation is unduly influenced by body shape or weight, and the pattern causes clinically significant distress or impairment.

### **Alternate Names:**

Bulimia; Bulimic Disorder; Binge-Purge Disorder

## **Bulla**

A large fluid-filled blister on the skin or mucous membrane, or a large air-filled space in an organ such as the lung. The cause and treatment depend on its location.

## **Bullae**

Large fluid-filled blisters on or beneath the skin or a mucous membrane. They can result from friction,

---

burns, infection, allergy, medicines, or autoimmune disease and may need care if widespread or infected.

### **Bullectomy**

An operation to remove one or more large air-filled spaces in damaged lung tissue. It may be considered when a giant bulla compresses healthier lung, causes severe breathlessness, or contributes to repeated pneumothorax.

#### **Alternate Names:**

Excision of Pulmonary Bullae; Resection of Giant Bulla

### **Bullet Track**

The path a firearm projectile takes through the body, from the entry wound through injured tissues to an exit wound or a retained bullet. The path may damage skin, organs, blood vessels, nerves, or bone.

### **Bullet Wounds**

Penetrating or perforating injuries caused by firearm projectiles. They may damage skin, organs, blood vessels, nerves, and bones and require urgent assessment even when the external wound looks small.

### **Bullosa Hereditaria**

Historical term; see *Epidermolysis Bullosa*.

### **Bullous**

Describes a condition involving large, raised, fluid-filled blisters. Bullous disease can result from infection, burns, medicines, allergy, or an autoimmune reaction.

### **Bullous Disease of Diabetes (Bullosis Diabeticorum)**

A rare diabetes-related blistering disorder producing spontaneous, usually painless tense blisters on the feet or lower legs. Lesions often heal without scarring, but wound care and evaluation for infection or another blistering disease are needed.

### **Bullous Emphysema**

A severe manifestation of chronic obstructive pul-

monary disease characterized by the formation of large, thin-walled air-filled spaces (bullae) exceeding one centimeter in diameter within the lung parenchyma, predisposing to compression of healthy lung and pneumothorax.

#### **Alternate Names:**

Vanishing Lung Syndrome

### **Bullous Impetigo**

A contagious superficial bacterial skin infection, usually caused by *Staphylococcus*, that produces fragile fluid-filled blisters and yellow or varnish-like crusts. It is especially common in infants and young children.

### **Bullous Pemphigoid**

An autoimmune skin disease, usually in older adults, that often begins with an itchy, hive-like rash followed by firm, fluid-filled blisters. The blisters are less fragile than those of pemphigus, and mouth involvement is less common. Diagnosis may involve a skin biopsy and blood tests for related antibodies.

#### **Alternate Names:**

BP

### **Bullous Systemic Lupus Erythematosus (BSLE)**

A rare autoimmune subepidermal blistering disease occurring in people with systemic lupus erythematosus, often producing tense vesicles or bullae on normal or inflamed, sometimes sun-exposed skin. Diagnosis uses skin biopsy with direct immunofluorescence and serologic testing to distinguish it from bullous pemphigoid and other blistering disorders; dapsone and treatment of SLE are commonly considered by specialists.

### **Bully**

A person who repeatedly harms, threatens, or intimidates someone with less power. Bullying can cause injury, anxiety, depression, sleep problems, school avoidance, and other health effects.

### **Bullying**

Repeated physical, verbal, social, or online aggression involving a power imbalance. It can harm

---

mental and physical health; safety planning, support, and intervention by responsible adults are important.

### **Bumetanide**

A diuretic that makes the kidneys remove extra salt and water in urine. It is used for swelling related to heart, liver, or kidney disease; excessive fluid loss can cause dehydration and abnormal blood salts.

### **Bumps**

Raised or palpable areas on or under the skin. Causes range from harmless cysts, blocked follicles, and insect bites to injury, allergy, infection, or tumours; a growing, painful, hard, or persistent bump requires assessment.

### **BUN**

Alternate name; see *Blood Urea Nitrogen Test*.

### **BUN-to-Creatinine Ratio**

The ratio of blood urea nitrogen to serum creatinine, used clinically to differentiate etiologies of acute kidney injury. A ratio greater than 20:1 suggests prerenal azotemia, whereas a ratio less than 15:1 is characteristic of intrinsic renal damage (such as acute tubular necrosis).

### **Bundle Branch Block**

Delay or interruption of electrical signals through one of the heart's lower-chamber conduction branches. It may cause no symptoms or reflect heart disease and is diagnosed on an ECG.

### **Bundle Branch Reentry Tachycardia**

A form of monomorphic ventricular tachycardia occurring in patients with severe underlying structural heart disease and conduction system disease, in which the reentrant electrical circuit involves the bundle of His and the right and left bundle branches.

### **Bundle Of His**

A short pathway of specialised heart tissue that carries electrical signals from the atria to the ventricles. It divides into right and left branches so the lower

chambers contract in a coordinated way.

### **Bundle of Kent**

An abnormal, congenital accessory muscular conduction pathway directly connecting atrial myocardium to ventricular myocardium, bypassing the physiological delay of the atrioventricular node; the anatomical substrate of Wolff-Parkinson-White syndrome.

#### **Alternate Names:**

Accessory Atrioventricular Pathway

### **Bundle-Branch Block**

Delayed or interrupted electrical signalling through one of the heart's lower-chamber conduction branches. It can change the ECG and heartbeat pattern and may be harmless or related to underlying heart disease.

### **Bungarotoxins**

Poisons in the venom of krait snakes that disrupt nerve signals to muscles. A bite can cause drooping eyelids, weakness, progressive paralysis, and breathing failure and requires emergency treatment.

### **Bunion**

A change in the joint at the base of the big toe that pushes the toe toward the others and creates a bump on the inner foot. It may cause pain, redness, swelling, stiffness, and difficulty wearing shoes.

### **Bunion Removal**

An operation to correct a painful bunion by realigning bone and, when needed, balancing nearby soft tissues. It is considered when symptoms persist despite suitable shoes and other nonsurgical care.

### **Bunionectomy**

Surgery to correct a painful bunion by cutting or shifting bone and adjusting surrounding tissues. The operation is chosen according to the deformity, symptoms, activity, other health conditions, and response to nonsurgical care.

### **Bunionette**

A bony prominence at the outer side of the joint

---

below the little toe. It may become painful, red, or swollen from pressure, tight shoes, or altered foot mechanics.

**Alternate Names:**

Tailor's Bunion; Bunion of the Fifth Toe

**Bunions**

Deformities at the base of the big toe that push the toe toward the others and create a bump on the inner foot. They may cause pain, swelling, stiffness, and difficulty finding comfortable shoes.

**Bunionoplasty**

Surgery to correct a bunion by realigning the toe joint and, when needed, balancing nearby soft tissues. The operation is selected according to the deformity and symptoms.

**Bunyaviridae**

An older virus-family grouping that included several viruses spread by insects, ticks, or rodents. Modern classification has changed; individual viruses differ in the illnesses they cause.

**Bunyavirus Infection**

Infection caused by a bunyavirus, a group that includes viruses responsible for illnesses such as hantavirus disease, Rift Valley fever, and some mosquito- or tick-borne febrile syndromes. Symptoms and treatment depend on the specific virus.

**Bunyaviruses**

Alternate name; see *Viral Hemorrhagic Fevers*.

**Alternate Names:**

Bunyavirus Infection

**Buphthalmos**

Enlargement of an infant's or young child's eye caused by high pressure inside the eye while its outer coat is still flexible. It is usually associated with congenital or infantile glaucoma and needs urgent eye care.

**Bupivacaine**

A long-acting local anaesthetic that temporarily

blocks nerve signals to prevent pain during local, nerve-block, or epidural procedures. Too much in the bloodstream can cause seizures, dangerous heart effects, or cardiac arrest.

**Bupivacaine Hydrochloride**

The hydrochloride form of bupivacaine, a long-acting local anaesthetic used to block pain signals. Accidental injection into a blood vessel can cause severe neurologic or heart toxicity.

**Buprenorphine**

A medicine that partly activates opioid receptors and is used to treat opioid-use disorder and some pain. It can cause dependence and overdose, especially with other sedatives, although respiratory suppression is limited compared with full opioid agonists.

**Buprenorphine/Naloxone Toxicity**

Opioid toxicity from excessive buprenorphine exposure, characterized by somnolence, miosis, hypoventilation, hypoxia, or coma; risk increases with alcohol, benzodiazepines, or other sedatives. Buprenorphine binds tightly to opioid receptors, so airway support and repeated or prolonged naloxone treatment may be required.

**Bupropion**

A medicine used for depression and, in some settings, to help stop smoking. It changes norepinephrine and dopamine signalling and may cause insomnia, dry mouth, or anxiety; it increases seizure risk and is unsuitable for some seizure or eating disorders.

**Buried Penis**

A normally developed penis that is concealed by surrounding skin, scar tissue, or excess suprapubic fat. It may cause hygiene problems, recurrent inflammation, difficulty urinating, or sexual difficulties; causes can be congenital, obesity-related, inflammatory, or associated with circumcision complications.

**Burkholderia**

A group of environmental bacteria found in soil and water. Some species cause melioidosis or glan-

---

ders, while others cause difficult-to-treat lung, bloodstream, or wound infections, especially in people with chronic lung disease or weakened immunity.

### **Burkholderia Cenocepacia**

A bacterium in the *Burkholderia cepacia* complex that can cause severe, persistent lung infection, especially in people with cystic fibrosis or other chronic lung disease.

### **Burkholderia Cepacia**

An environmental bacterium in the *Burkholderia cepacia* complex. It can cause opportunistic lung, bloodstream, wound, or device-related infection, particularly in people with cystic fibrosis or weakened immunity.

### **Burkholderia Gladioli**

An environmental bacterium that can rarely cause lung, bloodstream, or wound infection, especially in people with cystic fibrosis or weakened immunity.

### **Burkholderia Mallei**

The bacterium that causes glanders, an infection mainly affecting horses and related animals. People can become infected through contact with infected animals or tissues.

### **Burkholderia Pseudomallei**

The bacterium that causes melioidosis. Infection follows contact with contaminated soil or water and may cause pneumonia, abscesses, bloodstream infection, or long-term illness.

### **Burkholderiaceae**

A family of mainly environmental bacteria that includes *Burkholderia*. Some members cause serious infections in people or animals, while others are harmless or used in environmental research.

### **Burkitt Lymphoma**

A very fast-growing cancer of B lymphocytes, often involving the abdomen, jaw, bone marrow, or nervous system. It requires prompt intensive treatment; the pattern differs between endemic, sporadic, and

immunodeficiency-associated forms.

#### **Alternate Names:**

Burkitt's Tumor

### **Burkitt's Lymphoma**

A very fast-growing cancer of B lymphocytes. It may affect the abdomen, jaw, bone marrow, or brain and spinal cord and needs prompt intensive treatment with careful supportive care.

### **Burkitt's Tumor**

Alternate name; see *Burkitt Lymphoma*.

### **Burn**

An injury to the skin or deeper tissue caused by heat, chemicals, electricity, radiation, friction, or extreme cold. Severity depends on depth, size, location, and associated injury; extensive or deep burns need urgent medical care.

### **Burn Injury**

Alternate name; see *Burns*.

### **Burn Shock**

A complex form of combined distributive, hypovolemic, and cellular shock developing after extensive cutaneous burn injury (typically greater than 20 percent total body surface area), characterized by massive capillary leak, progressive plasma extravasation into interstitial tissues, and systemic hypoperfusion.

### **Burn Surgery**

Surgical care for serious burns, including assessing depth and area, removing dead tissue, covering wounds, performing skin grafts, and treating scars or movement-limiting contractures.

### **Burn Wound Infections**

Infection of damaged skin or deeper tissue after a burn. Concerning changes include increasing pain, redness or swelling spreading beyond the burn, pus, foul odour, delayed healing or graft loss, fever, or other systemic illness; prompt assessment may require wound sampling, cleaning or debridement, and

---

topical or systemic antimicrobial treatment.

### **Burning Feet**

Burning, tingling, or painful sensations in the feet, often caused by nerve damage from diabetes, nutritional deficiency, alcohol, medicines, or nerve compression. New, progressive, or weakness-associated symptoms need assessment.

### **Burning Mouth Syndrome**

A chronic oral pain disorder causing daily burning, scalding, or tingling discomfort in the mouth, often involving the tongue, lips, or palate, usually with a normal-appearing mucosa. Altered taste or dry mouth may occur. It may be primary, with no identifiable cause, or secondary to conditions or exposures such as nutritional deficiency, oral infection, medication effects, allergy, diabetes, or hormonal change; evaluation excludes treatable causes.

#### **Alternate Names:**

Scalded Mouth Syndrome Stomatodynia

### **Burning Thigh Pain**

Alternate name; see *Meralgia Paresthetica*.

### **Burnout**

A work-related state of emotional exhaustion, mental distance or cynicism, and reduced effectiveness caused by chronic workplace stress. It is not the same as depression, although both may occur together.

### **Burns**

Tissue injuries caused by heat, chemicals, electricity, radiation, friction, or extreme cold. Seriousness depends on depth, area, and location; burns affecting the face, airway, hands, genitals, major joints, or large areas need prompt care.

#### **Alternate Names:**

Thermal Injury; Burn Trauma

### **Burnt-Out Joint**

An older description of a joint in which active inflammation has subsided but permanent damage, deformity, pain, or restricted movement remains.

### **Burping**

Release of swallowed air or gas from the stomach through the mouth. Occasional burping is normal; frequent, painful, or newly worsening burping may result from swallowed air, reflux, food intolerance, or another digestive disorder.

### **Burr Cells**

Erythrocytes that exhibit numerous (usually 10 to 30) short, blunt, evenly spaced, and symmetrical spicules across their entire surface membrane. They are typically observed in end-stage renal disease and uremia, severe hypomagnesemia, and pyruvate kinase deficiency.

#### **Alternate Names:**

Echinocytes; Crenated Red Cells

### **Burr Hole Evacuation**

An emergency or urgent neurosurgical procedure in which one or two small holes (approximately 10 to 14 mm in diameter) are drilled through the skull bone to drain an abnormal fluid or blood collection compressing the brain. Most frequently performed to treat chronic subdural hematomas (which commonly occur in older adults after minor, often forgotten head injuries), a neurosurgeon incises the scalp and dura mater, washes out the liquefied old blood and clots with warm sterile saline, and places a temporary flexible drain to prevent fluid reaccumulation. By rapidly decompressing the brain, burr hole drainage reverses life-threatening brain herniation, confusion, focal weakness, and lethargy with significantly less surgical trauma, blood loss, and recovery time than a formal craniotomy.

#### **Alternate Names:**

Burr Hole Drainage; Trephination; Chronic Subdural Hematoma Drainage; Burr Hole Craniostomy

### **Burroughs**

An inconsistent surname or eponym in medical sources rather than a precise diagnosis; the associated condition or person must be identified from context.

### **Bursa**

A small fluid-filled sac that reduces friction between



---

moving tissues, such as a tendon and bone or skin and bone, especially near a joint. Inflammation of a bursa is called bursitis.

**Alternate Names:**

Bursae Synovial Bursae

**Bursa Inflammation**

Alternate name; see *Bursitis*.

**Bursectomy**

Surgical removal of a bursa, usually considered for persistent painful bursitis, recurrent infection, or a structural problem that has not improved with appropriate nonsurgical treatment.

**Bursitis**

Inflammation of a bursa, the small fluid-filled sac that cushions movement between tendons, muscles, bones, and skin. It causes localized pain, tenderness, swelling, or reduced movement and may result from pressure, injury, infection, or arthritis.

**Bursitis Of The Heel**

Inflammation of a bursa near the heel, causing localized tenderness, swelling, and pain when walking or when pressure is applied. It may follow repeated friction, injury, inflammatory arthritis, or infection.

**Alternate Names:**

Retrocalcaneal Bursitis

**Bursitis of the Knee**

Alternate name; see *Knee Bursitis*.

**Bursopathy**

Any disorder of a bursa, including inflammation, infection, bleeding, thickening, or degeneration. Affected bursae may cause localized pain, swelling, tenderness, and restricted movement around a joint.

**Burton Line**

A thin, purplish-blue or black discoloration line along the gingival margin at the base of the teeth. Caused by the precipitation of lead sulfide following the chemical reaction between circulating lead and hydrogen sulfide produced by oral bacterial decay;

it represents a classical physical sign of chronic lead poisoning (plumbism).

**Alternate Names:**

Burtonian Line; Lead Line

**Buruli Ulcer**

A chronic skin infection caused by *Mycobacterium ulcerans*, usually affecting an arm or leg. Its toxin, mycolactone, causes a painless nodule, plaque, or swelling that can progress to an undermined ulcer with extensive tissue destruction and little early inflammation; antibiotic treatment, wound care, and selected surgery can prevent disability.

**Burundanga**

A colloquial term for scopolamine (hyoscyne hydrobromide), a tropane alkaloid and muscarinic receptor antagonist that induces central anticholinergic effects, anterograde amnesia, sedation, and severe toxicity.

**Alternate Names:**

Scopolamine; Hyoscyne; Hyoscyne Hydrobromide

**Buserelin**

A synthetic gonadotropin-releasing hormone analogue. Continuous treatment first briefly increases and then suppresses pituitary reproductive hormones, lowering ovarian or testicular hormone production in selected fertility and hormone-sensitive conditions.

**Buspirone**

A nonbenzodiazepine medicine used mainly for generalized anxiety disorder. It acts on serotonin signaling and usually takes days to weeks to help; it does not provide immediate relief from an acute panic attack.

**Busulfan**

An alkylating chemotherapy medicine used mainly to prepare the body for blood-forming stem-cell transplantation and sometimes to treat chronic myeloid leukemia. It can suppress bone marrow and may cause seizures, infertility, lung injury, or liver damage.

---

## **Butanol**

A flammable industrial alcohol used as a solvent. Swallowing it or substantial exposure to its vapor can irritate the eyes, skin, and airways and may cause dizziness, vomiting, drowsiness, or other nervous-system effects.

## **Butazolidin Overdose**

Toxic effects from excessive phenylbutazone, an older nonsteroidal anti-inflammatory drug. Severe exposure may cause gastrointestinal bleeding, kidney injury, bone-marrow suppression, metabolic disturbances, or seizures.

## **Butocarboxim**

A carbamate insecticide that inhibits acetylcholinesterase. Significant exposure can cause cholinergic poisoning, with sweating, salivation, vomiting, diarrhea, wheezing, muscle weakness, seizures, or respiratory failure.

## **Buttiauxella**

A genus of environmental gram-negative bacteria in the Enterobacterales; human infection is uncommon and usually opportunistic.

## **Buttiauxella Agrestis**

A species of environmental Gram-negative bacteria found in water, soil, and other natural settings. It is rarely reported from human clinical samples; when detected, the specimen source and the patient's symptoms determine whether it represents infection or harmless contamination.

## **Buttiauxella Ferragutiae**

A species of environmental Gram-negative bacteria. It is uncommon in human disease, and a laboratory finding requires interpretation with the specimen source, repeat cultures, symptoms, and other evidence before deciding whether treatment is needed.

## **Buttiauxella Gaviniae**

A species of *Buttiauxella* bacteria found mainly in environmental samples. Human infection is uncommon; identification from a clinical specimen should be reviewed with the microbiology laboratory and

interpreted according to the patient's condition.

## **Buttiauxella Noackiae**

A species of environmental Gram-negative bacteria that is rarely associated with human infection. Its significance depends on where it was found, whether it was recovered repeatedly, and whether the patient has compatible symptoms or signs of infection.

## **Buttock**

The buttock is the rounded posterior region overlying the gluteal muscles, used for movement, sitting, examination, and selected injections.

## **Buttocks**

The paired fleshy regions overlying the gluteal muscles at the back of the pelvis, formed mainly by muscle and subcutaneous fat.

## **Button Batteries**

Button batteries are small batteries that can cause severe tissue injury if swallowed or inserted into the nose or ear. Suspected ingestion requires emergency evaluation.

## **Button Magnets In The Nasal Cavity**

Small magnets placed in or inhaled into the nose can attract across the nasal partition and cut off its blood supply. They may cause tissue death or a hole and require urgent medical removal.

## **Buttonhole Deformity**

Buttonhole deformity is a finger deformity with flexion of the proximal interphalangeal joint and hyperextension of the distal interphalangeal joint, often due to central slip injury or rheumatoid disease.

## **Buttressing**

Buttressing is reinforcement or thickening that supports a weakened structure; in bone it may describe cortical stress adaptation or surgical structural support.

---

## **Butylamine**

Butylamine is an irritating industrial amine whose vapor or liquid can injure the eyes, skin, and respiratory tract after exposure.

## **Butyric Acid**

Butyric acid is a short-chain fatty acid produced by gut bacteria during fiber fermentation and used by colon cells as an energy source; concentrated exposure irritates tissue.

## **Bx**

A common medical abbreviation for biopsy, the removal of cells or tissue for microscopic or laboratory examination.

## **By-Pass**

A route made around a blocked or narrowed passage. In medicine, a bypass may redirect blood, food, bile, urine, or another fluid through a natural or surgically created pathway.

## **Bypass**

Creation of an alternative route around a blocked or narrowed passage. A bypass can restore blood flow or redirect body contents, using a natural pathway, tube, vein, or artificial graft depending on the organ involved.

## **Bypass Graft**

A vein or artificial tube surgically connected to route blood around a blocked or weakened artery. Grafts are used in coronary, leg, and aortic operations; blood flow, infection, clotting, and long-term narrowing affect their durability.

## **Bypass Surgery**

An operation that creates a new route around a blocked or diseased passage. Examples include coronary and leg-artery bypasses and gastric bypass; possible complications include bleeding, infection, clotting, and narrowing of the connection.

## **Byssinosis**

An occupational lung disease caused by repeated

inhalation of dust from cotton, flax, hemp, or other textile fibers. It can cause chest tightness, cough, and breathing difficulty that worsen during or after work exposure.

---

# C

## C Fiber

A thin nerve fiber that carries slow, aching pain, warmth, and itch from the skin and other tissues to the spinal cord and brain.

## C-11 Hydroxylase Deficiency

A form of congenital adrenal hyperplasia caused by impaired 11-beta-hydroxylase activity. Reduced cortisol synthesis increases adrenal androgens and deoxycorticosterone, which may cause androgen excess, virilization, hypertension, and low potassium.

## C-17 Hydroxylase Deficiency

A form of congenital adrenal hyperplasia caused by impaired 17-alpha-hydroxylase activity. It reduces cortisol and sex-steroid production and may cause hypertension, low potassium, undervirilization, or absent puberty.

## C-Arm Fluoroscopy

A movable X-ray system that gives live images during a medical procedure. It helps show bones, instruments, blood vessels, or other body parts while the procedure is being performed.

## C-Peptide

A small protein fragment released when the pancreas makes insulin. Measuring it helps show how much insulin a person's body is making, rather than how much injected insulin is present. Kidney function and blood-glucose levels can affect the result.

## C-Reactive Protein

A protein made by the liver that rises when there is inflammation or infection. It can help monitor

inflammatory diseases and possible infection, but it does not identify the cause by itself. A high-sensitivity CRP test may be used with other information to assess heart-disease risk.

## C-Reactive Protein Test

A blood test that measures C-reactive protein. The level may rise with inflammation or infection, but the test cannot identify the cause by itself and must be interpreted with symptoms and other findings.

## C-Section

Alternate index label; see *Cesarean Section*.

## C. Diff

Alternate index label; see *C. difficile infection*.

## C. Difficile

*Clostridioides difficile* is a spore-forming bacterium that can multiply in the bowel, especially after antibiotics disrupt normal bacteria. Its toxins cause watery diarrhea and colon inflammation, which can become severe or recur.

## C. Difficile Colitis

Alternate index label; see *Pseudomembranous colitis*.

## C. Difficile Infection

An infection caused by *Clostridioides difficile*, a bacterium that can grow excessively in the bowel after antibiotic use. It commonly causes watery diarrhea and inflammation of the colon and may lead to dehydration, severe colitis, or recurrence.

**Alternate Names:**

---

C. Diff Clostridioides Difficile (C. Diff) Infection

### **C1-INH**

Abbreviation for C1 esterase inhibitor, a complement-regulating protein. Inherited or acquired deficiency or dysfunction can cause recurrent episodes of bradykinin-mediated angioedema affecting the skin, abdomen, or airway.

### **C1Q Nephropathy**

A rare kidney disorder in which an immune protein called C1q collects in the kidney's filtering units. It may cause blood or protein in the urine, swelling, high blood pressure, or reduced kidney function.

### **C282Y Mutation**

A change in the HFE gene that is the most common genetic cause of hereditary hemochromatosis, a condition in which the body stores too much iron. Having two copies of this mutation increases the risk, although not everyone with two copies develops the disease.

### **CA 19-9**

A substance measured in blood that may be increased in pancreatic and other digestive-system cancers. It can also rise with blocked bile ducts or inflammation, so it cannot screen for or diagnose cancer by itself.

### **CA 27-29**

A blood test that measures a protein sometimes released by breast-cancer cells. It may help monitor selected people with known breast cancer but cannot by itself screen for or diagnose cancer.

### **CA-125**

A blood protein that may rise in ovarian and some other cancers. It can also increase with menstruation, pregnancy, endometriosis, infection, liver disease, or other noncancerous conditions, so it cannot diagnose ovarian cancer alone.

#### **Alternate Names:**

Cancer Antigen 125; Carbohydrate Antigen 125; MUC16

### **CA-125 Blood Test**

See *CA-125*.

### **Cabazitaxel**

A taxane chemotherapy medicine used with prednisone for metastatic castration-resistant prostate cancer after docetaxel. It can cause low white-cell counts, infection, diarrhea, and serious allergic reactions.

### **Cabergoline**

A dopamine D2-receptor agonist used to suppress prolactin secretion and treat hyperprolactinemia; it is also used in selected Parkinson disease settings and requires monitoring for adverse effects.

### **CABG**

Coronary artery bypass grafting, surgery that creates a new path around a narrowed or blocked heart artery to improve blood flow to the heart muscle.

#### **Alternate Names:**

Coronary Artery Bypass Grafting; Coronary Bypass Surgery

### **Cabot Ring**

A thread-like or ring-shaped remnant sometimes seen inside a red blood cell on a blood smear. It may occur when red blood cells are being made abnormally, including in severe anemia, but it is not a diagnosis by itself.

### **Cachectic**

Affected by cachexia, a severe wasting syndrome involving involuntary weight loss, muscle loss, weakness, and often reduced appetite in chronic disease.

### **Cachetic**

An alternate spelling of cachectic, describing severe unintentional loss of body weight and muscle caused by a serious long-term illness.

### **Cachexia**

A wasting syndrome caused by serious long-term illness, marked by ongoing unintentional loss of weight, muscle, and body fat. It may cause poor



---

appetite, weakness, and inflammation and cannot be corrected by nutrition alone.

### **Cacosmia**

A distorted sense of smell in which ordinary odors seem unusually foul or unpleasant. It may occur with nose or sinus disease, infection, seizures, head injury, or another nervous-system disorder.

### **CAD**

Coronary artery disease, narrowing or blockage of the arteries supplying the heart, usually caused by fatty plaque. It can reduce oxygen to heart muscle and cause chest pain, heart attack, heart failure, or dangerous rhythms.

### **CAD (Coronary Artery Disease)**

Abbreviation for coronary artery disease, narrowing or blockage of the heart's arteries by atherosclerotic plaque. It can cause angina, heart attack, heart failure, arrhythmia, or sudden death.

### **CADASIL Syndrome**

An inherited disorder of small brain blood vessels caused by changes in the NOTCH3 gene. It may cause migraine with aura, repeated strokes, mood changes, and gradually worsening memory or thinking problems.

### **Cadherin**

A cell-surface protein that helps neighboring cells attach to one another and keeps tissues organized. Loss or alteration of cadherins can allow some cancer cells to invade nearby tissue and spread.

### **Cadmium**

A toxic metal found in tobacco smoke, some batteries and pigments, workplace dust, and contaminated food or soil. Long-term exposure can damage the kidneys, lungs, and bones and increases the risk of some cancers.

### **Caduceus**

A winged staff with two entwined snakes, traditionally associated with Hermes. It is often used

incorrectly as a medical symbol and differs from the single-snake Rod of Asclepius.

### **Cadusafos**

An organophosphate pesticide that blocks acetylcholinesterase, causing excess nerve signaling after exposure. Poisoning may produce sweating, salivation, vomiting, diarrhea, small pupils, muscle weakness, breathing difficulty, or seizures.

### **Caecal**

Relating to the caecum, the pouch at the beginning of the large intestine where the small intestine joins the colon and the appendix begins.

### **Caecitis**

Inflammation of the caecum, the first part of the large intestine. It may result from infection, inflammatory bowel disease, reduced blood flow, neutropenia, or spread of nearby inflammation and can cause right-sided abdominal pain and fever.

### **Caecopexy**

An operation that attaches the caecum, the first pouch of the large intestine, to nearby tissue to prevent or treat abnormal movement or twisting.

### **Caecostomy**

An operation that creates an opening from the caecum to the abdominal wall. It may temporarily release pressure, drain the bowel, or divert stool when the normal route is blocked or unsafe.

### **Caecotomy**

An operation that opens the caecum, usually to remove an object, access the bowel, or assist another abdominal procedure. The opening may be closed during the same operation.

### **Caecum**

The first pouch-like part of the large intestine, located where the ileum joins the colon and attached to the appendix.

---

## Caesarean Birth

Delivery of a baby through surgical openings in the abdomen and uterus. It may be planned or urgent when problems such as fetal distress, labor complications, abnormal presentation, placenta previa, or multiple pregnancy make vaginal birth unsafe.

## Caesarean Section

A caesarean section is surgical delivery of a baby through incisions in the abdomen and uterus when vaginal birth is unsafe or not feasible.

## Cafe-Au-Lait Macules

Alternate spelling; see *Café-Au-Lait Macules*.

## Cafe-Au-Lait Spot

Alternate name; see *Café-Au-Lait Macules*.

## Caffeine

A stimulant found in coffee, tea, chocolate, soft drinks, and energy drinks. It increases alertness and can raise heart rate, cause shakiness, or disturb sleep.

## Caffeine Intoxication

A toxic reaction to excessive caffeine, causing restlessness, anxiety, tremor, insomnia, nausea, vomiting, rapid heartbeat, or agitation. Severe poisoning can cause abnormal heart rhythms, seizures, or dangerous metabolic changes.

## Caffeine Overdose

A dangerous amount of caffeine can cause severe agitation, tremor, vomiting, rapid heartbeat, abnormal heart rhythms, low blood pressure, seizures, or disturbed body chemistry. Severe symptoms require urgent medical care.

## Caffeine Toxicity

Stimulant poisoning causing nausea, vomiting, agitation, tremor, tachycardia, hypertension, hypokalemia, hyperglycemia, and sometimes seizures or ventricular dysrhythmias. Severe cases require ECG and electrolyte monitoring, benzodiazepines for agitation or seizures, supportive care, and oc-

asionally activated charcoal or extracorporeal removal.

## Caffeine Withdrawal

A group of symptoms after suddenly reducing or stopping regular caffeine use. Headache, tiredness, low mood, poor concentration, irritability, nausea, or flu-like feelings usually begin within a day and improve within several days.

## Café Au Lait Spot

A flat, clearly outlined light-brown patch of skin caused by extra pigment. One or two are often harmless; many patches, especially with other findings, may indicate an inherited disorder.

## Café Coronary

Sudden blockage of the airway by food, usually during a meal. Complete blockage prevents breathing and can rapidly cause collapse or death unless emergency choking first aid and medical treatment restore the airway.

## Café-Au-Lait Macules

Flat, light-brown skin patches with relatively uniform pigmentation. Multiple lesions may be associated with genetic disorders such as neurofibromatosis type 1.

## Caisson Disease

Decompression sickness caused by gas bubbles forming in tissues or blood after a rapid fall in surrounding pressure, such as after diving. It can cause joint pain, skin changes, breathing problems, brain or spinal-cord injury, or collapse.

## Caladium Plant Poisoning

Poisoning from chewing or swallowing caladium, whose needle-like calcium oxalate crystals irritate the mouth and throat. Burning, pain, drooling, and swelling may occur; trouble breathing or swallowing is an emergency.

## Calamine

A skin preparation containing zinc compounds used

---

to soothe itching and irritation from minor rashes, insect bites, and sunburn. It should not be applied to deep wounds, infected skin, or the eyes.

### **Calbindin**

A protein that binds calcium inside cells and helps control calcium levels and cell signaling, especially in nerve and intestinal cells. Laboratories may also use it as a marker to identify certain tissues.

### **Calcaneal Spur**

A bony growth from the heel bone, usually where the plantar fascia or Achilles tendon attaches. It may cause no symptoms; when heel pain occurs, the associated soft-tissue inflammation is often more important than the spur itself.

### **Calcaneal Tendon**

Alternate name; see *Achilles Tendon*.

### **Calcaneal Valgus**

An outward tilting of the heel so that the foot rolls toward the body's midline. It may occur with flat feet, altered lower-limb alignment, or developmental and neuromuscular conditions and can affect walking or cause pain.

### **Calcaneum**

The calcaneum, or calcaneus, is the large bone forming the heel. It supports body weight, connects the ankle to the foot, and transmits force from the calf muscles through the Achilles tendon.

### **Calcaneus**

The largest bone in the ankle and foot, forming the heel. It supports body weight and connects with the ankle bone and cuboid; fractures commonly follow falls from height and may require imaging and specialist care.

### **Calcarine Sulcus**

An anatomical fissure located on the medial surface of the occipital lobe of the brain, running horizontally from the parieto-occipital sulcus to the occipital pole; its upper and lower banks constitute the

primary visual cortex (Brodmann area 17).

#### **Alternate Names:**

Calcarine Fissure

### **Calcifediol**

The main circulating storage form of vitamin D, also called 25-hydroxyvitamin D. A blood test for calcifediol is used to assess vitamin-D status before the body converts it to the active hormone calcitriol.

### **Calciferol**

A fat-soluble form of vitamin D that helps the body absorb calcium and phosphate and maintain healthy bones, muscles, and immune function. The body changes it into active vitamin-D hormones.

### **Calcific Bursitis**

A condition in which calcium deposits form in or around a bursa, often near the shoulder, causing pain, tenderness, swelling, and restricted movement. Treatment depends on severity and may include rest, physical therapy, medicines, drainage, or deposit removal.

#### **Alternate Names:**

Calcifying Bursitis; Hydroxyapatite Bursitis

### **Calcific Tendinitis**

A condition in which calcium crystals build up inside a tendon, most often a rotator-cuff tendon in the shoulder. It can cause sudden severe pain, stiffness, and restricted movement, sometimes followed by gradual improvement as deposits break down.

### **Calcific Uremic Arteriolopathy**

A rare, life-threatening complication, usually of advanced kidney disease, in which calcium and clots block tiny blood vessels in the skin and fatty tissue. It causes severe pain, purple or black skin, wounds, tissue death, and a high risk of infection.

### **Calcification**

Deposition of calcium salts in body tissue. It may be a normal part of bone formation, a response to injury or aging, or a sign of abnormal calcium or phosphate balance; its importance depends on the

---

site and pattern.

### **Calcified Granuloma**

A small area of past inflammation that has healed and become hardened by calcium deposits. It is often found in the lung after an old infection and usually does not represent active disease.

### **Calcineurin**

A protein that helps activate T cells, which are immune cells, after they receive signals. Medicines that block it reduce immune activity and are used to prevent transplant rejection and treat selected immune diseases.

### **Calcinosis**

An older or shortened term for calcinosis, in which calcium salts build up in the skin or tissues beneath it. Firm white or yellow lumps may form, sometimes ulcerate, and release chalky material.

### **Calcinosis**

Abnormal deposition of insoluble calcium salts in the skin or deeper soft tissues, often associated with autoimmune connective-tissue disease, kidney disease, trauma, or an abnormal calcium-phosphate balance. Firm white or yellow nodules or plaques may be painful, restrict movement, ulcerate, or release chalky material.

### **Calcinosis Cutis**

Deposition of calcium salts in the skin or subcutaneous tissue, producing firm white or yellow papules, nodules, or plaques. Lesions may be painless or tender, restrict movement, ulcerate, or discharge chalky material; the cause and blood calcium or phosphate levels guide evaluation.

### **Calciphylaxis**

A life-threatening vascular calcification syndrome predominantly affecting patients with end-stage renal disease on dialysis or secondary hyperparathyroidism. Progressive medial calcification and intimal fibroplasia of small dermal and subcutaneous arterioles lead to luminal thrombosis, severe tissue ischemia, intensely painful violaceous plaques

with livedoid patterns, and rapidly progressive non-healing ischemic skin necrosis with black eschar formation.

#### **Alternate Names:**

Calciphylaxis Cutis; Calcific Uremic Arteriopathy; CUA; Vascular Calcification-Uremic Syndrome

### **Calcitonin**

A hormone made by specialized cells in the thyroid gland. It helps lower blood calcium by reducing calcium release from bone and increasing calcium loss through the kidneys, although it has a smaller role than parathyroid hormone in adults.

### **Calcitonin Blood Test**

A blood test that measures calcitonin, a hormone produced by thyroid C cells. It is used mainly to evaluate or monitor medullary thyroid carcinoma and may be affected by other thyroid, kidney, gastrointestinal, or medicine-related factors.

### **Calcitriol**

The active hormone made from vitamin D, mainly by the kidneys. It increases intestinal absorption of calcium and phosphate and helps regulate parathyroid hormone, bone formation, and blood-mineral levels.

### **Calcitriol And Analogues**

Medicines related to the active form of vitamin D that reduce excessive parathyroid-hormone release and help correct mineral imbalance in some kidney disorders. They can raise calcium or phosphate, so blood levels require monitoring.

### **Calcium**

A mineral needed for strong bones and teeth, muscle movement, nerve signals, blood clotting, and normal cell function. Blood calcium exists in an active free form and forms attached to proteins or other substances.

### **Calcium Blood Test**

A blood test that measures calcium, usually as total calcium. The result may be interpreted with albumin, a blood protein, or measured as ionized

---

calcium when the active form is needed. It helps assess parathyroid, bone, and kidney disorders.

### **Calcium Carbonate**

A calcium supplement and antacid that neutralizes stomach acid. Excessive use can cause constipation, high blood calcium, or kidney problems and may interfere with medicine absorption.

### **Calcium Carbonate Overdose**

Overdose of calcium carbonate that may cause gastrointestinal symptoms, hypercalcemia, alkalosis, kidney injury, or kidney stones.

### **Calcium Carbonate With Magnesium Overdose**

Taking too much of these antacid ingredients can cause nausea, vomiting, constipation or diarrhea, weakness, confusion, abnormal calcium or magnesium levels, and kidney injury. Severe symptoms or reduced kidney function require urgent medical advice.

### **Calcium Channel**

A protein in the cell membrane that controls calcium entry into cells. Calcium channels help regulate muscle contraction, the release of substances, nerve signals, and heart rhythm.

### **Calcium Channel Blocker Toxicity**

Overdose causing vasodilatory or cardiogenic shock with hypotension, bradycardia, atrioventricular block, hyperglycemia, and sometimes pulmonary edema or cardiac arrest. Treatment may include intravenous calcium, vasopressors, high-dose insulin with glucose, and selected lipid emulsion or mechanical support under toxicology guidance.

### **Calcium Chloride**

A calcium salt given by injection for severe low blood calcium, dangerously high potassium, or certain toxic exposures. It can cause serious tissue injury if it leaks outside the vein and therefore requires careful administration.

### **Calcium Citrate**

A calcium supplement that is absorbed even when stomach acid is low. It may prevent or treat calcium deficiency but can cause constipation and, in some people, contribute to kidney stones.

### **Calcium Deficiency**

Too little calcium intake or availability for the body's needs. Long-term deficiency can weaken bones; severe deficiency can lower blood calcium and cause tingling, muscle cramps, spasms, or abnormal heart rhythms.

### **Calcium for Osteoporosis**

Calcium intake is part of osteoporosis prevention and treatment, together with vitamin D, resistance exercise, fall prevention, and medicines when indicated. Excess supplementation can cause kidney stones, constipation, or other harm.

### **Calcium Gluconate**

A calcium salt taken by mouth or given by injection to treat calcium deficiency and some emergencies, including dangerously high potassium or magnesium. Intravenous treatment requires medical monitoring because it can affect the heart and veins.

### **Calcium Hydroxide Poisoning**

Exposure to a strongly alkaline chemical that can burn the mouth, throat, esophagus, skin, or eyes. Do not induce vomiting; eye, breathing, or swallowing problems require immediate emergency assessment.

### **Calcium Phosphate**

A mineral that forms much of the hard structure of bones and teeth. Excess calcium phosphate can also deposit in soft tissues, especially after tissue damage or when blood calcium or phosphate levels are abnormal.

### **Calcium Pyrophosphate Arthritis**

Inflammation caused by calcium pyrophosphate crystals in cartilage or a joint. Sudden attacks, often called pseudogout, commonly affect the knee or wrist; chronic deposits can cause ongoing arthritis

---

and joint damage.

### Calcium Pyrophosphate Deposition

Alternate index label; see *Calcium Pyrophosphate Deposition Disease*.

### Calcium Pyrophosphate Deposition Disease

A disorder in which calcium pyrophosphate crystals collect in cartilage and joints. It may cause sudden painful swelling of one joint, usually the knee or wrist, or a chronic arthritis that resembles osteoarthritis or rheumatoid arthritis.

#### Alternate Names:

Calcium Pyrophosphate Deposition

### Calcium Sensing Receptor

A protein in the parathyroid glands and kidneys that detects the blood calcium level. It helps control parathyroid hormone and influences how much calcium and phosphate the kidneys keep or remove.

### Calcium Signaling

A way cells communicate using changes in calcium concentration. These signals help control muscle contraction, secretion of substances, energy use, movement, and activity of genes.

### Calcium Stones

The most common type of kidney stone, made mainly of calcium oxalate or calcium phosphate. Low urine volume and certain mineral imbalances increase risk; stones may cause severe side pain, blood in the urine, blockage, or infection.

### Calculi

Hardened mineral concretions, such as gallstones, kidney stones, or salivary stones, formed when dissolved substances precipitate and aggregate within a body organ or duct.

### Calculus

A hard deposit formed when minerals collect and solidify in the body. Examples include a kidney stone, gallstone, or hardened dental plaque.

#### Alternate Names:

Tartar

### Calculus Removal

Professional dental treatment that removes hardened plaque from teeth and below the gumline using hand or ultrasonic instruments. It helps control gum inflammation and may include deep cleaning of affected tooth roots.

### Caldwell-Luc Operation

An operation that reaches the maxillary sinus through the upper gum and jawbone. It is used selectively to treat or examine certain sinus lesions, chronic disease, or foreign bodies when less invasive routes are unsuitable.

### Calf

The back part of the lower leg between the knee and ankle. It contains the gastrocnemius, soleus, blood vessels, nerves, and other tissues and helps point the foot downward and propel walking.

### Calf Pseudohypertrophy

An enlarged, firm appearance of the calf muscles caused mainly by replacement of damaged muscle with fat and fibrous tissue rather than increased muscle strength. It is seen especially in Duchenne and Becker muscular dystrophies and some related disorders.

#### Alternate Names:

Pseudohypertrophy of the Calves; Muscular Pseudohypertrophy

### Calibration

Checking and, when necessary, adjusting a medical or laboratory instrument against a known standard so its measurements are accurate and reliable. Regular calibration supports correct diagnosis, treatment, and patient safety.

### Caliciviridae

A family of viruses that includes norovirus and sapovirus, which commonly cause vomiting and diarrhea. Other members infect animals; illness depends on the particular virus, the person or ani-



---

mal infected, and the route of exposure.

### **Calipers**

An instrument with two arms used to measure distance, thickness, or body dimensions. In health care, calipers may measure skinfold thickness, joint movement, wounds, or the size of body structures.

### **Calla Lily Poisoning**

Poisoning from chewing or swallowing calla lily, whose needle-like calcium oxalate crystals irritate the mouth and throat. Burning, pain, drooling, and swelling may occur; breathing or swallowing difficulty requires emergency care.

### **Callosal**

Relating to the corpus callosum, the broad bundle of nerve fibers connecting the left and right halves of the brain. Callosal abnormalities can affect communication between the two sides.

### **Callosity**

A small area of thick, hardened skin caused by repeated pressure or rubbing. It commonly develops on the hands or feet and may become painful if the pressure continues.

### **Callosotomy**

Surgical division of the corpus callosum to reduce the spread of certain severe, treatment-resistant seizures between the cerebral hemispheres.

### **Callus**

A thick, hardened area of skin formed by repeated pressure or rubbing. Calluses commonly occur on hands or feet, usually cause no harm, and may become painful or crack when extensive.

### **Calluses and Corns**

Localized areas of thickened skin caused by repeated pressure or friction. Calluses are broader; corns are more focal and often painful. Diabetes, poor circulation, or infection requires professional foot care rather than self-cutting.

### **Calmodulin**

A protein inside cells that senses calcium and passes its signals to other proteins. It helps control muscle contraction, movement of substances through cell membranes, chemical reactions, and activity of genes.

### **Calnexin**

A protein inside cells that helps newly made proteins fold into the correct shape. It checks proteins before they are sent to other parts of the cell or broken down.

### **Calor**

Abnormal warmth of a body part, usually caused by increased blood flow during inflammation. It may occur with redness, swelling, pain, or reduced function.

### **Caloric Restriction**

Deliberately eating fewer calories without causing malnutrition. It may help with weight loss and some metabolic risks, but it should provide enough protein, vitamins, minerals, and other nutrients.

### **Caloric Stimulation**

A balance test in which warm or cool air or water is placed in the ear canal. The temperature briefly stimulates the inner-ear balance organs and helps compare the balance nerve on each side.

### **Calorie**

A unit of energy. In nutrition, one Calorie with a capital C equals one kilocalorie (kcal), the energy needed to raise the temperature of one kilogram of water by one degree Celsius.

#### **Alternate Names:**

Kilocalorie; Kcal; Calories

### **Calorimetry**

Measurement of heat production or gas exchange to estimate how much energy the body uses. Indirect calorimetry uses oxygen consumption and carbon-dioxide production to help assess metabolism and nutritional needs.

---

## Calot Triangle

An anatomical space in the porta hepatis bounded inferiorly by the cystic duct, medially by the common hepatic duct, and superiorly by the inferior surface of the liver (or cystic artery in modern descriptions). It serves as an essential surgical landmark during cholecystectomy for identifying the cystic artery and cystic duct.

### Alternate Names:

Triangle of Calot; Cystohepatic Triangle

## Calpain

A group of enzymes activated by calcium that cut selected proteins inside cells. They help regulate cell movement, signaling, programmed cell death, and platelet activity; abnormal calpain activity contributes to some muscle and nervous-system diseases.

## Calpainopathy

An inherited muscle disease caused by harmful changes in both copies of the CAPN3 gene. It slowly weakens hip, pelvic, shoulder, and trunk muscles, often causing shoulder-blade winging, back curvature, difficulty climbing, and walking problems while usually sparing facial and heart muscles.

### Alternate Names:

Limb-Girdle Muscular Dystrophy R1 (LGMD R1); LGMD 2A; Calpain-3 Deficiency

## Calponin

A protein that binds to actin, a component of the cell's internal framework, and helps regulate contraction in smooth muscle. Laboratory staining for calponin can help identify muscle-related cells in tissue samples.

## Calprotectin

A protein released mainly by activated white blood cells. Measuring it in stool helps detect inflammation in the intestines, particularly in inflammatory bowel disease, and can help distinguish it from some noninflammatory conditions.

## Calreticulin

A protein inside cells that helps newly made proteins

fold correctly and helps regulate calcium storage and immune signaling. Harmful changes in its gene are found in some blood cancers involving excessive production of blood cells.

## Calretinin

A calcium-binding protein found in certain nerve cells and cells lining body cavities. Pathologists may use calretinin staining to help distinguish mesothelioma from some other cancers in a tissue sample.

## Calvaria

The dome-shaped upper part of the skull that covers and protects the brain. It is formed mainly by the flat bones of the forehead, top and sides of the head, and back of the skull.

## Calvarium

The dome-like portion of the skull formed mainly by the frontal, parietal, and occipital bones. It encloses and protects the upper parts of the brain.

## Calyx

A cup-shaped body structure that supports tissue or collects fluid. In the kidney, renal calyces collect urine from the pyramids and carry it toward the renal pelvis.

## Cambridge Classification

A grading system used with endoscopic retrograde cholangiopancreatography to describe chronic pancreatitis. It rates abnormalities of the main pancreatic duct and its branches, from normal or uncertain findings to mild, moderate, or severe disease.

## Cameron Lesions

Long, shallow erosions or ulcers in the stomach where it is squeezed by the diaphragm, usually in people with a large hiatal hernia. They may cause chronic blood loss, iron-deficiency anemia, or visible gastrointestinal bleeding.

## Campho-Phenique Overdose

Poisoning after swallowing or excessive exposure to a topical product containing camphor and other ingre-

---

dients. It may cause nausea, vomiting, drowsiness, agitation, or seizures and requires poison-control or emergency advice.

### **Camphor**

A strong-smelling substance used in some topical products and traditional remedies. Swallowing even a relatively small amount can cause nausea, agitation, drowsiness, or rapidly developing seizures and may be life-threatening.

### **Camphor Overdose**

Toxic exposure to too much camphor, causing nausea, vomiting, agitation, drowsiness, seizures, coma, or slowed breathing. Seizures may begin quickly; suspected ingestion requires immediate poison-control or emergency assessment.

### **Campimetry**

Measurement and mapping of the area a person can see while looking straight ahead. It helps detect and describe visual-field loss caused by disease of the retina, optic nerve, or brain pathways.

### **Campomelic Dysplasia**

A rare inherited disorder caused by changes affecting the SOX9 gene. It causes bowed long bones, poorly developed shoulder blades, characteristic facial features, and often serious problems with breathing or other organs.

### **Camptocormia**

Severe forward bending of the trunk that worsens while standing or walking and improves when lying down. It may result from muscle disease, Parkinson disease, dystonia, or another nerve or muscle disorder.

### **Camptodactyly**

A fixed bending deformity of one or more fingers, usually at the middle finger joint. It may be present from birth or develop later because of shortened tendons, muscles, connective tissue, or joint structures.

### **Campylobacter**

A group of bacteria that commonly cause foodborne intestinal infection. Illness may include fever, abdominal cramps, diarrhea, or bloody stools; some species can invade the bloodstream or cause later nerve or joint problems.

### **Campylobacter Coli**

A *Campylobacter* species that can cause foodborne intestinal infection, usually after contaminated meat, water, or other food. Symptoms include fever, abdominal cramps, diarrhea, and sometimes bloody stools.

### **Campylobacter Enteritis**

An acute bacterial foodborne or waterborne diarrheal illness caused predominantly by *Campylobacter jejuni*, presenting with fever, severe cramping abdominal pain mimicking appendicitis, and inflammatory bloody diarrhea, with potential post-infectious Guillain-Barre syndrome.

#### **Alternate Names:**

Campylobacteriosis

### **Campylobacter Fetus**

A *Campylobacter* species that can cause bloodstream infection and infections of blood vessels, joints, the brain, or other tissues. Serious disease is more likely in older adults and people with weakened immunity.

### **Campylobacter Food Poisoning**

A foodborne infection caused by *Campylobacter* bacteria, often from undercooked poultry, unpasteurized milk, or contaminated water. It causes diarrhea, cramps, and fever and may rarely trigger bloodstream infection or nerve inflammation.

### **Campylobacter Infection**

A bacterial infection, usually caused by *C. jejuni* or *C. coli*, acquired from contaminated poultry, food, water, or unpasteurized milk. It causes fever, cramps, and diarrhea that may be bloody; rare complications include bloodstream infection, arthritis, or Guillain-Barré syndrome.

---

## **Campylobacter Jejuni**

A major cause of foodborne intestinal infection, usually acquired from undercooked poultry, contaminated water, or unpasteurized milk. It commonly causes fever, abdominal cramps, and diarrhea that may contain blood.

## **Campylobacter Serology Test**

A blood test that detects antibodies produced after Campylobacter infection. It may support evaluation of selected complications after infection, but stool molecular testing or culture is generally used to diagnose active intestinal disease.

## **Campylobacteriosis**

Infection caused by Campylobacter bacteria, usually producing fever, abdominal pain, and diarrhea that may be bloody. Dehydration is common; rare complications include bloodstream infection, reactive arthritis, and Guillain-Barré syndrome.

## **Canagliflozin**

A medicine that lowers blood glucose by causing the kidneys to remove more glucose in urine. It may also reduce some heart-failure and kidney risks, but can cause genital infections, dehydration, low blood pressure, or ketoacidosis.

## **Canal Of Schlemm**

A circular drainage channel at the edge of the cornea that carries aqueous fluid from the front of the eye into the bloodstream. Poor drainage can raise eye pressure and contribute to glaucoma.

## **Canaliculus**

A small anatomical channel. The tear canaliculi are tiny passages in the eyelids that carry tears from the eye surface into the lacrimal sac and then the nose.

## **Canalith Repositioning Maneuver**

A series of guided head and body movements that moves displaced inner-ear crystals out of a semicircular canal. It is commonly used to treat posterior-canal benign paroxysmal positional vertigo.

## **Canalith Repositioning Maneuvers**

Head and body movements that move loose crystals from an inner-ear semicircular canal back to the utricle, where they no longer trigger false movement signals. The maneuver is selected according to the affected canal and side.

## **Canalization**

Formation or restoration of a channel through tissue, a blood vessel, or an embryonic structure. In a blood clot, canalization means that small channels form through the clot and partly restore blood flow.

## **Canavan Disease**

A rare inherited disorder in which deficiency of the ASPA enzyme damages the brain's white matter. It usually begins in infancy with poor muscle tone, delayed development, increasing head size, feeding problems, and later severe movement and communication impairment.

## **Cancellous Bone**

Porous bone made of thin supporting plates or struts. It is found inside many bones, especially near joints and in the vertebrae, and contains marrow while helping distribute forces.

## **Cancer**

A group of diseases in which abnormal cells grow and divide without normal control and may invade nearby tissues or spread to other parts of the body. Spread through the blood or lymph system to form a new tumor is called metastasis. Cancer usually begins after changes in genes that control cell growth, division, DNA repair, or cell death. It can start in almost any tissue. Many cancers form solid tumors, but cancers such as leukemia usually affect blood-forming tissues without forming a solid tumor. A benign tumor is not cancer and does not invade or spread in the way a malignant tumor does.

## **Cancer And Lymph Nodes**

Cancer cells may travel from a primary tumor to nearby lymph nodes through lymphatic vessels. Examination, imaging, or biopsy of these nodes helps determine the cancer's extent and guide treatment.

---

## Cancer Care Team

The health professionals who work together to diagnose, treat, and support a person with cancer. Depending on the illness, the team may include medical and radiation oncologists, surgeons, pathologists, nurses, pharmacists, therapists, and palliative-care specialists.

## Cancer Cell

A cell with changes that allow it to grow and survive without normal control. Cancer cells may lose their specialized function, resist cell death, invade nearby tissues, enter blood or lymph, and form tumors elsewhere.

## Cancer Cluster

A group or area with more cancer cases than expected during a specified time. Investigation compares the number and types of cancers with reliable background rates and examines possible shared exposures while considering chance variation.

## Cancer Genetics

The study of inherited and acquired genetic changes that affect cancer risk, behavior, or treatment. Testing may identify inherited risk syndromes or changes in a tumor that help guide treatment and family counseling.

## Cancer Grading

Assessment of how abnormal cancer cells and their tissue pattern look under a microscope. The grade gives information about likely growth behavior and is different from staging, which describes how far the cancer has spread.

## Cancer Immunotherapy

Cancer treatment that helps the immune system recognize or attack cancer cells. Methods include checkpoint inhibitors, engineered immune cells, immune-stimulating proteins, and some vaccines; use and side effects depend on the cancer and treatment.

## Cancer in Lung

Alternate name; see *Lung Cancer*.

## Cancer Mutations

Genetic changes in cancer cells that can alter growth, survival, or spread. Most develop during life and are not inherited. Some drive cancer development, while others have little effect; testing selected changes can help guide prognosis or targeted treatment.

## Cancer Nutrition

Nutrition care for people with cancer, who may lose appetite or weight because of the disease or treatment. Individual plans may address calories, protein, fluids, swallowing, nausea, diarrhea, and cachexia while respecting the person's goals and treatment.

## Cancer of the Ear

Cancer arising in the outer ear, ear canal, middle ear, or nearby tissues. Persistent sores, bleeding, pain, discharge, hearing loss, or a growing lump require examination; treatment depends on the site and tumor type.

## Cancer of the Tongue

Malignant tumor arising in the tongue, most often squamous cell carcinoma. A persistent sore, lump, bleeding area, numbness, difficulty swallowing, or unexplained tongue pain needs prompt examination.

## Cancer Of Unknown Origin

Alternate index label; see *Carcinoma of unknown primary*.

## Cancer Of Unknown Primary

Cancer that has spread to another site but whose original starting point cannot be identified after appropriate examination and testing. Tissue studies, scans, and selected molecular tests may help guide treatment.

## Cancer Pain

Pain caused by a tumor, its spread, or its treatment. It may be constant or intermittent and can arise from bone, nerves, organs, surgery, radiation, or medicines; treatment combines control of the cause with suitable pain relief and supportive care.

---

**Alternate Names:**

Malignancy-Related Pain; Oncologic Pain

**Cancer Prevention**

Actions that lower the chance of developing cancer or find precancerous changes early. They include avoiding tobacco, limiting alcohol, vaccination against cancer-causing infections, protecting skin from ultraviolet light, reducing harmful exposures, and recommended screening.

**Cancer Progression**

Growth or spread of cancer or worsening of its effects over time. Progression may occur at the original site, in nearby lymph nodes, or in distant organs and is assessed with symptoms, examination, imaging, laboratory tests, or biopsy.

**Cancer Recurrence**

Cancer that returns after a period when it could not be detected. It may recur near the original site or in another part of the body and may be called local, regional, or distant recurrence.

**Cancer Risk**

The likelihood that a person will develop cancer. It is influenced by age, inherited variants, family history, tobacco, alcohol, certain infections, radiation, environmental exposures, body weight, and chance; having a risk factor does not guarantee cancer.

**Cancer Screening**

Testing people who have no symptoms to find selected cancers or precancerous changes early, when treatment may work better. Each test has specific age, risk, interval, benefits, and possible harms.

**Cancer Staging**

Describing how far a cancer has grown or spread. Assessment may include the primary tumor, nearby lymph nodes, distant metastases, scans, and tissue examination; stage helps guide treatment and estimate prognosis.

**Cancer Surveillance**

Ongoing monitoring for cancer recurrence, progression, treatment effects, or new cancers after diagnosis or treatment. It may include examinations, imaging, laboratory tests, and symptom review according to the cancer and treatment.

**Cancer Survivor**

A person living with cancer from diagnosis through treatment and afterward. The term includes people whose cancer is controlled or cured and those with ongoing disease; follow-up may address recurrence, late effects, emotional health, and quality of life.

**Cancer Susceptibility Genes**

Genes in which inherited harmful variants increase the chance of developing one or more cancers. Examples include BRCA1, BRCA2, APC, and mismatch-repair genes; carrying a risk variant does not guarantee that cancer will develop.

**Cancer Treatment**

Medical, surgical, radiation, systemic, or supportive care used to control cancer, remove it, slow its growth, prevent recurrence, or relieve symptoms. Treatment is selected according to cancer type, stage, biomarkers, overall health, and patient goals.

**Cancer Treatment–Related Bleeding**

Bleeding or easy bruising that develops during cancer treatment, often because chemotherapy, radiation, or stem-cell transplantation lowers the platelet count. Signs may include nosebleeds, bleeding gums, unusual bruises, tiny red or purple skin spots, blood in urine or stool, or unusually heavy menstrual bleeding. Persistent or heavy bleeding needs prompt medical attention.

**Cancer Treatment-Induced Premature Menopause**

The premature and often irreversible cessation of ovarian endocrine and reproductive function (iatrogenic premature ovarian insufficiency) resulting from gonadotoxic chemotherapy, pelvic radiotherapy, or surgical bilateral oophorectomy.



---

**Alternate Names:**

Treatment-Induced Early Menopause; Iatrogenic Premature Ovarian Insufficiency; Chemotherapy-Induced Menopause

**Cancer Treatments**

Cancer treatment may include surgery, radiation, chemotherapy, immunotherapy, targeted medicines, hormone therapy, stem-cell transplantation, or supportive and palliative care. The choice depends on cancer type, location, stage, molecular features, overall health, and patient goals.

**Cancer-Associated Thrombosis**

A blood clot in a vein or artery occurring with active cancer or its treatment. Cancer, surgery, immobility, central lines, and some treatments increase risk; anticoagulant treatment is chosen according to clot location, bleeding risk, and cancer status.

**Cancer-Related Fatigue**

Persistent physical, emotional, or mental exhaustion caused by cancer, its treatment, or related problems. It may occur even after little activity and may not improve fully with sleep or rest.

**Cancers of the Oral Mucosa**

Malignant tumors arising in the moist lining of the mouth, most commonly squamous cell carcinoma. Persistent ulcers, red or white patches, lumps, bleeding, pain, numbness, or swallowing difficulty require prompt examination and biopsy; tobacco, alcohol, HPV, and other factors can contribute.

**Cancrum Oris**

A rapidly destructive infection of the mouth and face, seen mainly in severely malnourished or medically vulnerable children. It begins with painful tissue death around the gums and can spread through the cheek and other facial tissues.

**Candela**

The international unit used to measure the brightness of a light source in a specified direction. It is a physical unit, not a measure of a medicine's light dose or the amount of light reaching tissue.

**Candida**

A group of yeasts that may normally live on the skin, in the mouth, or in the digestive and genital tracts. Overgrowth can cause mouth, genital, skin, or, in vulnerable people, bloodstream and organ infections.

**Candida Albicans**

The *Candida* species most often causing human infection. It can cause oral thrush, vaginal yeast infection, moist skin-fold rash, or serious bloodstream and organ infection in people with weakened defenses.

**Candida Albicans Infection**

Infection caused by *Candida albicans*. It may be limited to the mouth, vagina, or skin, or may invade the bloodstream and organs, especially in people who are hospitalized or have weakened immunity.

**Candida Auris**

A yeast that can live on skin without symptoms or cause bloodstream, wound, or ear infection. It often resists several antifungal medicines, spreads readily in health-care settings, and requires specialized testing and strict infection-control measures.

**Candida Auris Infection**

Infection or symptom-free colonization caused by *Candida auris*. Invasive infection may affect the blood or organs, especially in hospitalized or immunocompromised people; treatment is guided by susceptibility testing because resistance is common.

**Candida Dubliniensis**

A yeast related to *Candida albicans* that can live in the mouth and cause oral thrush or, rarely, invasive infection. It is more likely to cause disease when immune defenses are impaired.

**Candida Esophagitis**

Yeast infection of the esophagus, usually caused by *Candida*. It commonly causes painful or difficult swallowing and chest discomfort, especially in people with weakened immunity or after certain medicines.

---

### **Candida Glabrata**

A *Candida* species that can cause genital, urinary, bloodstream, or other invasive infection. It may resist some commonly used antifungal medicines, so laboratory susceptibility testing can help guide treatment.

### **Candida Infection Of The Skin**

A superficial yeast infection, often in warm, moist skin folds. It causes a red, itchy, sore rash that may have small separate spots around its edge and is more common with moisture, diabetes, obesity, or immune suppression.

### **Candida Parapsilosis**

A *Candida* species that can cause bloodstream and catheter-related infection, especially in newborns, hospitalized people, and those receiving intravenous treatment. It may spread between patients through hands or contaminated equipment.

### **Candida Tropicalis**

A *Candida* species that can cause mouth or genital infection and may invade the bloodstream or organs. Serious infection is more likely in people with neutropenia, cancer, or other immune impairment.

### **Candidal Intertrigo**

A *Candida* yeast infection in warm, moist skin folds such as the armpits, groin, under the breasts, or beneath an abdominal fold. It causes a red, moist, itchy or burning rash, sometimes with small spots at the edge.

### **Candidal Vaginitis**

Inflammation of the vagina and vulva caused by overgrowth of *Candida* yeast. It usually causes intense itching, soreness, redness, pain with sex or urination, and a thick white discharge without a strong odor.

### **Candidal Vulvovaginitis**

Alternate index label; see *Vulvovaginal Candidiasis*.

### **Candidemia**

*Candida* yeast infection of the bloodstream. It can spread to organs and is more common with central lines, surgery, intensive care, recent antibiotics, or weakened immunity; it requires prompt medical treatment and evaluation for spread.

### **Candidiasis**

Alternate name; see *Yeast Infection*.

### **Canefield Fever**

An older name for leptospirosis, a bacterial infection acquired through water or soil contaminated with urine from infected animals. It can cause fever, headache, muscle pain, jaundice, kidney injury, meningitis, or bleeding.

### **Canicola Fever**

A form of leptospirosis caused by a strain of *Leptospira* bacteria commonly carried by dogs. It may cause fever, headache, muscle aches, red eyes, kidney injury, or liver involvement after contact with infected urine or contaminated water.

### **Canine Teeth**

The pointed teeth between the incisors and premolars. They help tear food and guide the jaws during biting and chewing.

### **Canker**

A nontechnical word often used for a painful, shallow ulcer on the moist lining of the mouth. Because the word can have other meanings, the sore's exact site and appearance should be identified.

### **Canker Sore**

A painful, shallow ulcer on the moist lining of the mouth, usually round or oval with a red border. It is not contagious and commonly heals within one to two weeks; persistent, severe, or unusual sores need assessment.

#### **Alternate Names:**

Aphthous Ulcer; Recurrent Aphthous Stomatitis (RAS)

---

## Canker Sores

Alternate name; see *Canker Sore*.

## Cannabidiol

A cannabinoid from the cannabis plant that usually does not produce the “high” caused by THC. A purified form treats selected severe seizure disorders; other products vary in strength, purity, side effects, and medicine interactions.

## Cannabinoid

A substance that acts on cannabinoid receptors or related signaling systems. Cannabinoids may come from plants, the body, or laboratories; effects can include pain relief, appetite changes, sedation, impaired memory, or intoxication.

## Cannabinoid Hyperemesis Syndrome

Repeated episodes of severe nausea, vomiting, and abdominal pain associated with long-term frequent cannabis use. Hot showers may temporarily ease symptoms, but lasting improvement usually requires stopping cannabis.

## Cannabinoid Receptor

A cell-surface protein, mainly CB1 or CB2, that responds to cannabinoids made by the body or found in cannabis and medicines. It helps regulate pain, appetite, mood, memory, immune activity, and inflammation.

## Cannabinol

A cannabinoid formed as THC breaks down with age or exposure to air. It may have mild psychoactive effects, but its medical benefits and safety are less well established than those of some other cannabinoids.

## Cannabis

A plant containing cannabinoids such as THC and CBD. Use can alter perception, memory, coordination, mood, and heart rate; it may have selected medical uses but can also cause dependence, anxiety, psychosis, or other harms.

## Cannabis Use Disorder

A pattern of cannabis use that causes significant difficulty controlling use or problems at home, work, school, relationships, or health. Tolerance and withdrawal may occur, with irritability, anxiety, sleep problems, or reduced appetite after stopping.

## Cannabis-Related Disorders

Clinically significant problems related to cannabis use, including intoxication, withdrawal, cannabis use disorder, and substance-induced psychiatric or medical effects.

## Cannula

A thin hollow tube placed in a blood vessel, body cavity, or tissue to deliver medicines or fluids, remove material, measure pressure, provide oxygen, or allow access for another device.

## Cannula Hub

The connector at the outer end of a cannula that attaches it to tubing, a syringe, or another device. It may include a port or valve for giving fluids, taking samples, or flushing the cannula.

## Cannulation

Placement of a cannula into a blood vessel, body cavity, duct, or other passage. It provides access for giving fluids or medicines, removing material, monitoring, collecting samples, or performing a procedure.

## Canola Oil

A vegetable oil relatively low in saturated fat and containing mostly monounsaturated and polyunsaturated fats. Replacing solid fats with canola oil can support a heart-healthy eating pattern when total calories remain appropriate.

## Canrenone

A metabolite related to spironolactone that blocks aldosterone’s action. It promotes loss of sodium and water while reducing potassium loss and has been used in some countries for conditions involving fluid retention or excess aldosterone.

---

## Cantharides

Preparations made from blister beetles that contain cantharidin, a powerful irritant. Swallowing or applying it can cause severe mouth, stomach, kidney, and urinary-tract injury and may be life-threatening.

## Canthotomy

A surgical cut through the outer corner of the eyelids, usually performed urgently to relieve dangerous pressure around the eye and protect vision after severe orbital swelling or injury.

## Canthus

The corner where the upper and lower eyelids meet. The inner canthus is beside the nose, and the outer canthus is toward the temple.

## Cantlie Line

An imaginary line across the liver from the gallbladder to the large vein behind the liver. It follows the middle hepatic vein and marks the functional division between the right and left liver, including their blood supply and bile drainage.

## Cantu Syndrome

A rare genetic condition, usually caused by an ABCC9 or KCNJ8 gene change. It may cause excessive hair growth, distinctive facial features, loose skin, enlarged bones, skeletal differences, and an enlarged or abnormal heart.

## Capacitation

The final physiological maturation process that mammalian spermatozoa undergo within the female reproductive tract, involving the removal of glycoprotein coats and seminal plasma proteins from the acrosome, enabling the sperm to undergo the acrosome reaction and fertilize an oocyte.

## Capacity

A person's ability, at a particular time, to understand information about a medical decision, appreciate how it applies to them, weigh the options, and communicate a choice.

## Capacity And Competence

Capacity is the clinical ability to make a specific decision by understanding, using, and communicating relevant information. Competence is a legal determination; the two terms are related but not interchangeable.

## Capecitabine

An oral chemotherapy medicine that the body converts to fluorouracil, which interferes with cancer-cell DNA production. It is used for selected cancers and can cause diarrhea, mouth sores, hand-foot syndrome, low blood counts, or heart problems.

## Capgras Delusion

A psychiatric misidentification syndrome in which an individual maintains a fixed, unshakeable delusion that a close relative, spouse, or companion has been replaced by an identical-looking imposter, double, or clone. Hypothesized to stem from a neuroanatomical disconnection between visual facial recognition pathways in the fusiform gyrus and emotional familiarity processing in the amygdala.

### Alternate Names:

Capgras Syndrome; Illusion of Doubles

## Capgras Syndrome

A fixed false belief that someone familiar has been replaced by an identical impostor. It may occur with psychotic illness, dementia, brain injury, seizures, or other neurologic disorders and needs clinical assessment.

## Capillaria

A group of parasitic roundworms. Different species can cause intestinal infection, liver disease, or infection of other tissues; symptoms and treatment depend on the species and affected organ.

## Capillaria Philippinensis

A parasitic roundworm acquired by eating raw or undercooked freshwater fish. It infects the intestine and can cause persistent watery diarrhea, abdominal pain, weight loss, poor absorption of nutrients, and loss of blood proteins.

---

## Capillaroscopy

Examination of tiny blood vessels, usually in the skin beside a fingernail, with magnification. It can show widened or missing vessels, bleeding, and abnormal patterns that support evaluation of connective-tissue and small-vessel disease.

## Capillary

The smallest type of blood vessel, forming networks that connect arterioles with venules. Its thin wall allows exchange of oxygen, nutrients, fluid, and waste between blood and tissues.

### Alternate Names:

Capillaries

## Capillary Bed

A network of capillaries within a tissue that connects small arteries to small veins. It provides a large surface for exchange of oxygen, carbon dioxide, nutrients, hormones, fluid, and waste.

## Capillary Blood Gas

A test of blood acidity and carbon dioxide using a small sample from a heel or finger. It is often used in babies and children when an arterial sample is difficult; oxygen results may not match an arterial blood gas.

### Alternate Names:

CBG

## Capillary Electrophoresis

A laboratory method that separates charged molecules in a very thin tube using an electric field. It can analyze proteins, medicines, DNA, or other substances in a sample, depending on the test.

## Capillary Exchange

The movement of oxygen, carbon dioxide, nutrients, fluid, hormones, and waste between blood in capillaries and nearby tissues.

## Capillary Hemangioma

A usually harmless growth made of small blood vessels, common in infancy. It often appears during the first weeks of life, grows for a time, and then

slowly becomes smaller. Some lesions near the eye, airway, or other important areas need treatment. Medicines such as propranolol may be used for high-risk lesions, while small superficial lesions may need only observation or local treatment.

### Alternate Names:

Infantile Hemangioma; Strawberry Hemangioma; Strawberry Nevus; Benign Capillary Hemangioma

## Capillary Leak Syndrome

A condition in which fluid and proteins escape from small blood vessels into surrounding tissues. It can cause sudden swelling, low blood pressure, thickened blood, shock, and organ failure; severe infection, medicines, immune disease, or rare unexplained episodes may trigger it.

## Capillary Malformation

A congenital difference in small skin blood vessels that produces a flat pink, red, or violaceous patch, such as a port-wine stain. It usually persists and may darken or thicken over time; location and associated findings determine whether eye, brain, or other specialist evaluation is needed.

## Capillary Microscope

A microscope used to magnify tiny blood vessels, especially those beside the fingernails. It helps assess vessel shape, size, bleeding, and areas without vessels in some circulation and connective-tissue disorders.

## Capillary Nail Refill Test

A bedside check of circulation in which a nail bed is pressed until it turns pale and the time for color to return is observed. Temperature, lighting, age, skin color, shock, and vascular disease affect the result, so it is not used alone.

## Capillary Permeability

How easily fluid and dissolved substances pass through the walls of small blood vessels. Increased permeability can allow extra fluid and proteins to enter nearby tissues and cause swelling.

---

### Capillary Plasma

The liquid component of blood obtained from small vessels in skin or tissue, containing water, electrolytes, proteins, hormones, nutrients, and waste products.

### Capillary Refill

A bedside assessment of peripheral perfusion performed by briefly pressing a nail bed or skin and measuring the time for normal color to return. Delayed refill may indicate poor circulation, shock, hypothermia, or vascular disease.

### Capillary Refill Time

The time required for color to return after pressure briefly blanches a nail bed or area of skin. A delayed result may suggest impaired peripheral perfusion, but interpretation depends on age, temperature, technique, and the clinical setting.

### Capillary Sample

A small blood specimen obtained by puncturing the skin, usually a finger or heel, for bedside glucose, hemoglobin, newborn screening, or other point-of-care tests.

### Capillary Vessel

A microscopic vessel with a thin endothelial wall where blood exchanges oxygen, carbon dioxide, nutrients, fluid, and waste with surrounding tissue.

### Capitate

The largest wrist bone, located centrally in the distal row of carpal bones. It connects with several other wrist bones and the third metacarpal, helps transmit force through the wrist, and can rarely develop blood-supply loss after fracture.

### Capitation

A health-care payment arrangement in which a clinician or organization receives a fixed amount per enrolled person for specified services over a set period, whether or not that person uses the services.

### Caplan Syndrome

The coexistence of rheumatoid arthritis with occupational pneumoconiosis (most commonly coal workers' pneumoconiosis or silicosis), characterized by the development of multiple discrete, round, peripheral pulmonary nodules measuring 0.5 to 5 centimeters on chest radiography. Histologically, the nodules resemble enlarged rheumatoid nodules with central necrotic debris surrounded by chronic inflammatory cells and dust-laden macrophages.

#### Alternate Names:

Rheumatoid Pneumoconiosis; Silicoarthritis

### Caplan's Syndrome

A lung disorder combining rheumatoid arthritis with pneumoconiosis caused by long-term inhalation of coal or mineral dust. It produces multiple lung nodules and may cause cough or breathing difficulty.

### Capnocytophaga

A group of bacteria found mainly in the mouths of dogs, cats, and people. Some species can cause serious human infection after animal bites or scratches, especially in people without a functioning spleen or with weakened immunity.

### Capnocytophaga Canimorsus

A bacterium commonly found in the mouths of dogs and cats. It can cause rapidly severe bloodstream infection, meningitis, or skin infection after a bite or scratch, especially in older adults or people without a functioning spleen or with weakened immunity.

### Capnocytophaga Infection

Infection caused by *Capnocytophaga* bacteria, often after a dog or cat bite, scratch, or contact with animal saliva. It may cause a wound infection, fever, bloodstream infection, or meningitis, with greatest risk in people with impaired immunity or no spleen.

### Capnocytophaga Ochracea

A bacterium found mainly in the human mouth and associated with dental plaque and gum disease. Infection outside the mouth is uncommon but can occur when immune defenses are severely weakened.



---

## Capnograph

A monitor that measures and displays carbon dioxide in exhaled air, often showing both the end-tidal value and a waveform. It is used to assess ventilation and to monitor selected patients during anesthesia, sedation, transport, or resuscitation.

## Capnography

Continuous measurement and display of carbon dioxide in exhaled air. The waveform and end-tidal value help assess breathing, airway blockage, circulation, and correct placement of a breathing tube during anesthesia, sedation, or emergency care.

## Caps

A prescription abbreviation for capsules, oral medicines enclosed in a shell that dissolves in the digestive tract. Abbreviations can be misread, so complete dosage-form wording is safer on prescriptions.

## Capsaicin

The substance that makes chili peppers feel hot. Applied in controlled amounts to the skin, it can reduce some nerve and joint pains by temporarily lowering pain signaling; it commonly causes initial burning or redness.

## Capsid

The protein shell surrounding a virus's genetic material. It protects the genome and helps the virus attach to and enter host cells; its structure influences which cells can be infected and how the immune system responds.

## Capsid Proteins

Proteins that assemble around a virus's genetic material to form its protective shell. Some also help the virus attach to and enter cells and are recognized by the immune system.

## Capsular Contracture

Abnormal tightening and hardening of scar tissue around an implanted device, especially a breast implant. It may make the area firm, painful, distorted, or displaced and sometimes requires correc-

tive surgery.

## Capsular Warning Syndrome

A clinical syndrome characterized by recurrent, stereotyped episodes of transient focal motor or sensory deficits affecting the face, arm, and leg without cortical signs (such as aphasia or visual neglect). Occurring repeatedly within a 24- to 72-hour window, it indicates impending lacunar infarction of the internal capsule due to branch atheromatous disease of a lenticulostriate artery.

### Alternate Names:

Internal Capsule Warning Syndrome; Fluctuating Lacunar TIA

## Capsule

A protective covering around an organ, joint, or other structure, or an oral medicine enclosed in a dissolvable shell. The exact meaning depends on whether the term is used in anatomy or pharmacology.

## Capsule Endoscope

A swallowable capsule containing a tiny camera and transmitter. As it travels through the digestive tract, it sends images, especially of the small intestine, where conventional endoscopes may not reach easily.

## Capsule Endoscopy

A test in which a person swallows a small wireless camera that takes pictures of the digestive tract, especially the small intestine. The capsule usually passes naturally but can become stuck where the bowel is narrowed.

## Capsule Of Glisson

The layer of connective tissue surrounding the liver. It extends inward around the liver's blood vessels, bile ducts, and nerves and helps support and protect these structures.

## Capsule Of Tenon

A thin connective-tissue sheath surrounding the eyeball and separating it from the fat of the eye socket. It allows smooth eye movement and creates a space used for some eye injections and operations.

---

## Capsulectomy

Surgical removal of some or all of the scar-tissue capsule around an organ, implant, or other structure. It may be performed for painful tightening, infection, abnormal tissue, or revision of a reconstruction.

## Capsulotomy

A procedure that makes an opening in a capsule. In the eye, a laser may open a cloudy capsule after cataract surgery; in other surgery, a capsule around an implant or joint may be opened to reduce tightness. The method and risks depend on the body part.

### Alternate Names:

Laser Capsulotomy; YAG Capsulotomy; Posterior Capsulotomy; Lens Capsulotomy

## Captopril

An angiotensin-converting enzyme inhibitor used for high blood pressure, heart failure, and some kidney problems. It can cause cough, high potassium, low blood pressure, or swelling.

## Captopril Renal Scan

A kidney imaging test using a radioactive tracer before and after captopril. It compares kidney blood flow and function and may help evaluate high blood pressure caused by narrowing of a kidney artery.

## Caput Medusae

Enlarged veins spreading out from the navel, caused by increased pressure in the portal vein system, usually from liver disease. They are a physical sign of abdominal collateral circulation rather than a disease by themselves.

## Caput Succedaneum

Soft swelling of a newborn's scalp caused by pressure during labor. It can cross the skull's suture lines and usually disappears within hours to a few days; rapidly enlarging swelling or pallor requires urgent assessment.

## CAR T-Cell

A T lymphocyte genetically changed to carry a

receptor that recognizes a specific marker on cancer cells. These cells can be used as treatment for selected blood cancers but may cause severe immune reactions or nervous-system effects.

## CAR T-Cell Therapy

A cancer treatment in which a person's T cells are collected, genetically changed to recognize cancer cells, and returned to the body. It is used for selected cancers and can cause serious immune or nervous-system reactions.

## Carbachol

A medicine that mimics acetylcholine and stimulates muscarinic receptors. In selected eye procedures, it constricts the pupil and can lower pressure inside the eye; effects may include sweating, slow heartbeat, or breathing problems.

## Carbamate

A chemical group found in some medicines and pesticides. Carbamate insecticides can temporarily block acetylcholinesterase, causing excess nerve activity with sweating, salivation, vomiting, weakness, wheezing, or seizures.

## Carbamazepine

A medicine used for focal seizures, trigeminal neuralgia, and selected bipolar disorder. Important risks include severe skin reactions, low blood counts, liver injury, low sodium, dizziness, and interactions with many other medicines.

## Carbamazepine Toxicity

Overdose causing drowsiness, ataxia, nystagmus, anticholinergic effects, hypotension, seizures, coma, or sodium-channel-mediated QRS widening and dysrhythmias. Serial drug levels and ECGs help guide care; activated charcoal, sodium bicarbonate for cardiotoxicity, and hemodialysis may be used in severe cases.

## Carbamide

Another name for urea, a waste product made when the body breaks down protein and removed mainly by the kidneys. Urea is also used in some skin

---

creams to soften and hydrate thick or dry skin.

### **Carbamoyl Phosphate Synthetase Deficiency**

An inherited urea-cycle disorder caused by deficient carbamoyl phosphate synthetase I. The liver cannot efficiently convert ammonia into urea, so hyperammonemia may cause poor feeding, vomiting, sleepiness, seizures, coma, or developmental injury.

### **Carbapenem**

A group of powerful beta-lactam antibiotics that kill bacteria by blocking construction of their cell wall. They are reserved mainly for serious infections because resistance and important side effects can limit treatment choices.

### **Carbapenem-Resistant Enterobacteriaceae**

Alternate name; see *CRE Infection*.

### **Carbapenem-Resistant Enterobacteriaceae Infection**

Alternate name; see *CRE Infection*.

### **Carbapenem-Resistant Organisms**

Bacteria that cannot be reliably treated with carbapenem antibiotics, often because they produce enzymes that destroy these medicines. They can cause difficult infections and spread in health-care settings, where strict infection control is important.

### **Carbaryl**

A carbamate insecticide that temporarily blocks acetylcholinesterase. Substantial exposure can cause sweating, excess saliva, vomiting, diarrhea, small pupils, muscle weakness, wheezing, seizures, or breathing failure.

### **Carbidopa**

A medicine given with levodopa for Parkinson disease. It blocks conversion of levodopa to dopamine outside the brain, allowing more levodopa to reach the brain and reducing nausea, vomiting, and low blood pressure.

### **Carbimazole**

An antithyroid medicine that the body converts to methimazole. It reduces production of thyroid hormones and can rarely cause dangerously low white-cell counts or liver injury.

### **Carbofuran**

A highly toxic carbamate pesticide that blocks acetylcholinesterase. Ingestion, inhalation, or skin exposure can rapidly cause sweating, vomiting, muscle weakness, seizures, breathing failure, and death.

### **Carbohydrate**

One of the three main macronutrients, found in sugars, starches, fruits, grains, vegetables, legumes, and milk. Sugars and most starches are digested into simple sugars, mainly glucose, for energy; dietary fiber is a carbohydrate that is not fully digested.

### **Carbohydrate Counting**

A way to plan meals by estimating the grams of carbohydrate in food. People using insulin may match their insulin dose to the amount eaten; accurate reading of food labels and advice from a diabetes team are important.

### **Carbohydrate Metabolism**

The processes cells use to break down, store, and make sugars. These processes provide energy and help keep blood glucose stable; problems can occur in diabetes and inherited disorders of sugar storage.

### **Carbohydrates**

Sugars, starches, and fiber that form one of the three main macronutrients in food. Sugars and most starches are broken down into simple sugars, mainly glucose, for energy, while fiber is not fully digested and can support bowel function, fullness, cholesterol levels, and blood-glucose control.

### **Carbolic Acid**

Phenol, a chemical once used as a surgical antiseptic and still found in some industrial products. It kills microorganisms by damaging proteins and cell membranes but can severely burn skin, eyes, and internal organs.

---

**Alternate Names:**

Phenol

**Carbolic Acid Poisoning**

Exposure to phenol, a corrosive and systemically toxic chemical that can cause burns, altered consciousness, metabolic acidosis, and organ injury.

**Carbon**

A chemical element forming the backbone of organic molecules such as carbohydrates, fats, proteins, and nucleic acids.

**Carbon Dioxide**

A gas produced when cells use energy and removed mainly by the lungs during exhalation. The body controls its blood level through breathing; excess can cause headache, confusion, drowsiness, or respiratory failure.

**Carbon Dioxide Absorber**

A device in some anaesthesia breathing systems that removes carbon dioxide from exhaled air. It allows selected gases to be reused while helping prevent carbon-dioxide build-up during controlled breathing.

**Carbon Dioxide Transport**

The movement of carbon dioxide from body tissues to the lungs. Blood carries it dissolved in plasma, attached to haemoglobin, and mainly as bicarbonate; the lungs remove it during exhalation.

**Carbon Ion Therapy**

A form of particle radiation treatment that uses carbon ions to damage tumor DNA. It may deliver concentrated energy and has higher biological effectiveness than conventional X-rays for some tumors, but availability is limited.

**Carbon Monoxide**

A colorless, odorless gas that binds hemoglobin and prevents blood from delivering enough oxygen. Poisoning may cause headache, dizziness, nausea, confusion, collapse, or death and requires immediate

fresh air and emergency care.

**Carbon Monoxide Poisoning**

Poisoning from breathing carbon monoxide, which prevents blood from carrying and delivering oxygen. It may cause headache, dizziness, nausea, confusion, fainting, seizures, or death; leave the area and obtain emergency care immediately.

**Carbon Monoxide Toxicity**

Tissue hypoxia caused by inhaled carbon monoxide binding hemoglobin and impairing cellular oxygen use. Headache, dizziness, nausea, confusion, syncope, myocardial ischemia, seizures, or coma may occur; pulse oximetry can be misleading, so co-oximetry and exposure history are important. Give 100% oxygen and consider hyperbaric therapy for selected severe cases.

**Carbon Tetrachloride**

A toxic industrial solvent that can damage the liver and kidneys when swallowed or inhaled. Suspected exposure requires urgent contact with poison-control or emergency services.

**Carbonate**

A chemical form of carbon dioxide found in minerals and body fluids. Carbonate salts are used in some antacids and supplements; their effects, absorption, and risks depend on the particular salt.

**Carbonic Anhydrase Inhibitors**

Medicines that block carbonic anhydrase, an enzyme involved in fluid and acid balance. They reduce fluid production in the eye to lower eye pressure and have selected uses for swelling, seizures, or metabolic disorders.

**Carbophenothion**

An organophosphate insecticide that blocks acetylcholinesterase. Significant exposure can cause sweating, excess saliva, vomiting, muscle weakness, confusion, seizures, breathing failure, and death.

---

## Carboplatin

A platinum chemotherapy medicine that damages cancer-cell DNA and prevents it from being copied. It can lower blood counts, especially platelets, and may cause nausea, anemia, kidney effects, nerve symptoms, or allergic reactions.

## Carboprost

A prostaglandin medicine that strongly contracts the uterus. It is used in selected cases of severe bleeding after childbirth and for some pregnancy-related procedures; it can cause diarrhea, vomiting, fever, and dangerous airway tightening in people with asthma.

## Carbosulfan

A carbamate insecticide that blocks acetylcholinesterase and overstimulates nerves. Exposure can cause sweating, salivation, vomiting, diarrhea, small pupils, muscle weakness, wheezing, seizures, or breathing failure.

## Carboxy-Lyases

Enzymes that break a carbon-carbon bond and release carbon dioxide. They participate in many biochemical reactions, including processes that make or break down amino acids and other molecules.

## Carboxyhaemoglobin

Hemoglobin bound to carbon monoxide. This reduces the blood's ability to carry and release oxygen; an increased level supports recent carbon-monoxide exposure but must be interpreted with symptoms and timing.

## Carboxyhemoglobin

A compound formed when carbon monoxide binds to hemoglobin. It prevents hemoglobin from carrying and releasing oxygen normally, causing headache, dizziness, confusion, collapse, or death in severe poisoning.

## Carboxyhemoglobinemia

An abnormally high level of carboxyhemoglobin in the blood after carbon-monoxide exposure. It reduces oxygen delivery and may cause headache, nau-

sea, confusion, fainting, heart injury, brain injury, or death.

## Carboxyl Group

A chemical group containing carbon, oxygen, and hydrogen that gives many molecules acidic properties. It is found in amino acids, fatty acids, and numerous medicines and metabolic compounds.

## Carboxylesterase

An enzyme that breaks ester bonds in medicines, toxins, and natural fats. Differences in its activity can change how quickly some substances are broken down and can affect their effects, interactions, or toxicity.

## Carboxylic Acid

An organic compound containing a carboxyl group, which gives it acidic properties. Many important fatty acids, amino-acid products, and substances involved in metabolism are carboxylic acids.

## Carboxypeptidase H

An enzyme inside cells that helps turn newly made precursor proteins into active hormones and nerve-signalling substances by trimming amino acids from one end.

## Carbuncle

A group of connected boils, usually caused by *Staphylococcus aureus*, forming a painful swollen area with several pus-draining openings. Fever may occur, and risk is increased by diabetes or a weakened immune system.

## Carbuncles and Boils

Alternate index label; see *Boils and Carbuncles*.

## Carcinoembryonal Antigen

Another name for carcinoembryonic antigen, or CEA, a blood marker that may rise in colorectal and other cancers. It can also increase for noncancerous reasons and is not suitable for general cancer screening.

---

## Carcinoembryonic Antigen

A protein measured in blood that may be increased in colorectal and some other cancers. It can also rise with smoking, liver disease, bowel inflammation, or other noncancerous conditions, so it is mainly used to monitor selected known cancers rather than to screen for cancer.

## Carcinoembryonic Antigen Tumor Marker

Carcinoembryonic antigen, a substance measured mainly in blood to help monitor some colorectal and other cancers. Levels can also rise from smoking or noncancerous conditions, so CEA cannot diagnose cancer by itself.

## Carcinogen

A substance, form of radiation, or infection that can cause cancer by damaging cells or changing how they grow. Risk depends on the type and amount of exposure.

## Carcinogenesis

The multistep process by which normal cells acquire changes that allow uncontrolled growth and become cancer. Changes may arise from inherited traits, random errors, infections, tobacco, radiation, chemicals, or other exposures and often accumulate over time.

## Carcinogenic

Capable of causing or promoting cancer by damaging genetic material or altering how cells grow, divide, repair themselves, or avoid cell death.

## Carcinoid

A usually slow-growing neuroendocrine tumor that can arise in the digestive tract, appendix, or lungs. Some release hormones or other substances that cause flushing, diarrhea, wheezing, or heart-valve disease.

## Carcinoid Flushing

Sudden, transient cutaneous erythema and warmth, predominantly affecting the face, neck, and upper chest, provoked by systemic release of serotonin,

bradykinin, and histamine from a metastatic neuroendocrine (carcinoid) tumor with hepatic metastases.

### Alternate Names:

Carcinoid Flush

## Carcinoid Heart Disease

Damage to the inner lining and valves of the heart caused by hormone-like substances released by a neuroendocrine tumor. It mainly affects the tricuspid and pulmonary valves and can lead to right-sided heart failure.

## Carcinoid Syndrome

A group of symptoms caused by hormone-like substances released by some neuroendocrine tumors, especially after spread to the liver. It may cause facial flushing, diarrhea, wheezing, rapid heartbeat, and eventually heart-valve disease.

## Carcinoid Tumor

A usually well-differentiated neuroendocrine tumor arising in hormone-producing cells, commonly in the appendix, small intestine, rectum, or lungs. Its behavior ranges from slow-growing to capable of spreading and causing hormone-related symptoms.

## Carcinoid Tumors

Neuroendocrine tumors that often grow slowly and may release hormones or other active substances. They commonly arise in the digestive tract or lungs and can sometimes invade or spread to distant organs.

### Alternate Names:

Neuroendocrine Tumors (NETs); Carcinoid Neoplasms

## Carcinoma

A cancer that begins in epithelial cells, which cover body surfaces and line organs and glands. Common types include adenocarcinoma, squamous-cell carcinoma, basal-cell carcinoma, and urothelial carcinoma; behavior depends on site and cell features.



---

### **Carcinoma En Cuirasse**

An advanced form of breast cancer in which cancer cells spread through lymphatic vessels in the skin, causing widespread hardening and thickening that can make the chest wall resemble armor.

### **Carcinoma Erysipeloides**

Spread of an internal cancer through lymph vessels in the skin, producing a red, firm, sharply outlined area that can resemble the infection erysipelas. It most often occurs with cancers of the breast or digestive tract.

### **Carcinoma In Situ**

Abnormal cancer-like cells occupying the full thickness of an epithelial lining but not yet invading through its supporting boundary. It is an early, noninvasive stage that may progress if untreated.

### **Carcinoma Of Unknown Primary**

Cancer found outside its original site when the starting organ cannot be identified despite evaluation. Treatment uses the tumor's appearance, genetic features, spread pattern, symptoms, and likely tissue of origin.

#### **Alternate Names:**

Cancer of Unknown Origin Occult Primary Cancer

### **Carcinomatosis**

Widespread spread of carcinoma to multiple surfaces or organs, often along the lining of the abdomen, chest, or other body cavities. It usually indicates advanced cancer and may cause fluid buildup, pain, bowel problems, or breathing difficulty.

### **Carcinosarcoma**

An uncommon cancer containing both gland- or surface-lining cancer cells and connective-tissue cancer cells. It most often begins in the uterus, tends to grow aggressively, and is diagnosed by examining a tissue sample.

### **Cardia**

The upper part of the stomach surrounding the opening where the esophagus enters. It helps re-

ceive swallowed food and can be affected by reflux, inflammation, hiatal hernia, or stomach cancer.

### **Cardiac**

Relating to the heart. The word may describe the heart muscle, chambers, valves, electrical system, blood supply, rhythm, or diseases such as coronary disease, cardiomyopathy, and heart failure; in “cardia,” it instead refers to the upper stomach.

#### **Alternate Names:**

Heart-Related; Cardiac-Related

### **Cardiac Ablation**

A procedure that uses heat, cold, or another form of energy to create a small scar in heart tissue causing an abnormal rhythm. The scar blocks faulty electrical signals and may restore a regular rhythm.

### **Cardiac Ablation Procedures**

Procedures that use heat, cold, or another energy source to treat heart-rhythm disorders by creating controlled scars that block abnormal electrical signals. Risks include bleeding, blood-vessel injury, recurrence, and damage to nearby heart structures.

### **Cardiac Allograft**

A donor heart transplanted into another person, usually for end-stage heart failure. Lifelong immune-suppressing treatment is needed to reduce rejection; complications include infection, rejection, coronary-vessel disease, cancer, and graft failure.

### **Cardiac Allograft Vasculopathy**

Progressive narrowing of the heart's blood vessels after transplantation caused by abnormal thickening of their inner walls. It may silently reduce blood flow and cause graft failure, rhythm problems, heart attack, or sudden death.

### **Cardiac Amyloidosis**

Buildup of abnormal amyloid protein in the heart, making the heart muscle stiff and impairing filling. It can cause breathlessness, swelling, rhythm problems, or heart failure; treatment depends on whether the protein is light-chain or transthyretin.

---

## Cardiac Amyloidosis Diagnosis

Evaluation combines symptoms, examination, blood and urine tests, heart imaging, and sometimes tissue biopsy. Specialized nuclear imaging may support transthyretin amyloidosis only after testing has excluded light-chain amyloidosis.

## Cardiac Amyloidosis Staging

A system for estimating the severity and outlook of transthyretin cardiac amyloidosis using markers of heart strain and injury, sometimes with other findings. Higher stages indicate more advanced disease; treatment and prognosis must be individualized.

## Cardiac Arrest

Sudden stopping of effective heart pumping, causing collapse, unresponsiveness, and absent or abnormal breathing. Immediate emergency action includes calling for help, starting chest compressions, and using an automated defibrillator if available.

### Alternate Names:

Cardiopulmonary Arrest; Circulatory Arrest

## Cardiac Arrhythmia

Alternate name; see *Arrhythmia*.

## Cardiac Arrhythmias

Abnormal heart rates or rhythms caused by problems forming or conducting electrical impulses. They include rhythms that are too fast, too slow, or irregular and may cause palpitations, dizziness, fainting, or no symptoms.

## Cardiac Axis

The average direction in which the heart's lower chambers become electrically activated, estimated from an electrocardiogram. A leftward or rightward shift may reflect normal variation or heart, conduction, or lung disease.

## Cardiac Biomarkers

Substances measured in blood to detect or assess heart-muscle injury. Troponin is the preferred marker; results must be interpreted with symptoms, electrocardiography, imaging, and changes

in repeated tests because many conditions can raise levels.

## Cardiac Cachexia

Severe unintentional loss of weight and muscle associated with advanced heart failure or other serious heart disease. It can cause weakness and reduced function and reflects inflammation, altered metabolism, poor appetite, and increased nutritional needs.

## Cardiac Catheter

A thin flexible tube inserted through a blood vessel and guided to the heart. It can measure pressures, take blood samples, inject contrast for imaging, or deliver treatment.

## Cardiac Catheterization

A procedure in which thin tubes are guided through a blood vessel to the heart to measure pressures, assess oxygen levels, inject contrast, image coronary arteries, or perform treatment. Possible risks include bleeding, rhythm problems, vessel injury, and kidney effects.

## Cardiac Conduction System

Specialized heart tissue that creates and carries electrical signals so the chambers contract in the correct order. It includes the sinoatrial node, atrioventricular node, bundle of His, bundle branches, and Purkinje fibers.

## Cardiac Contractility

The strength of heart-muscle contraction at a given level of filling and resistance. It affects how much blood the heart can pump with each beat.

## Cardiac CT

Computed-tomography imaging of the heart and nearby blood vessels. Depending on the protocol, it can assess coronary calcium, coronary narrowing, heart structure, valves, and selected aspects of heart function.

---

## Cardiac CT Angiography

A noninvasive scan using contrast and computed tomography to show the coronary arteries and heart structures. It is useful for assessing selected people with chest pain and can show narrowing, plaque, and coronary calcium.

## Cardiac Cycle

The events in one complete heartbeat. During diastole the heart chambers relax and fill with blood; during systole the ventricles contract and pump blood into the lungs and the rest of the body.

## Cardiac Defibrillation

Delivery of an electrical shock to stop ventricular fibrillation or pulseless ventricular tachycardia during cardiac arrest. It must be combined with high-quality CPR and advanced resuscitation according to the rhythm and situation.

## Cardiac Device Infections

Infection involving a pacemaker, implantable defibrillator, its pocket, or a lead. It may cause redness, pain, drainage, fever, bloodstream infection, or infection of the heart lining and often requires antibiotics and removal of infected hardware.

## Cardiac Diet

An eating pattern emphasizing vegetables, fruits, whole grains, beans, fish, nuts, and unsaturated fats while limiting excess sodium, added sugar, saturated fat, and trans fat. It should be adapted for heart failure, kidney disease, diabetes, and nutrition needs.

## Cardiac Electrophysiology Study

An electrophysiology study in which thin tubes are placed through a blood vessel into the heart to record and stimulate its electrical activity. It helps find the cause of some abnormal rhythms or unexplained fainting.

## Cardiac Enzymes

Older term; see *Cardiac Biomarkers*.

## Cardiac Event Monitors

Portable devices that record or send an electrocardiogram when symptoms occur or at planned times. They help detect intermittent abnormal rhythms that may be missed during a short clinic ECG.

### Alternate Names:

Cardiac Event Monitor; Ambulatory ECG

## Cardiac Failure

Alternate name; see *Heart Failure*.

## Cardiac Fibroma

A rare noncancerous tumor made of fibrous tissue in the heart, usually in a ventricle or septum. It may obstruct blood flow, interfere with electrical signals, or cause rhythm problems and heart failure.

## Cardiac Gated SPECT

A heart scan using a small radioactive tracer while images are synchronized with the heartbeat. It can assess blood flow to heart muscle, pumping strength, wall movement, and areas that are damaged or still viable.

## Cardiac Glycoside

A medicine or plant substance that can strengthen heart contraction and slow electrical conduction through the heart. Digoxin is the main clinical example; the safe dose is narrow and toxicity can cause dangerous rhythms.

## Cardiac Glycoside Overdose

Toxic exposure to a cardiac glycoside such as digoxin. It may cause nausea, vomiting, confusion, visual changes, slow or fast abnormal rhythms, high potassium, seizures, or cardiac arrest and requires urgent assessment.

## Cardiac Glycoside Plant Poisoning

Poisoning by oleander, foxglove, or similar plants containing digoxin-like cardiac glycosides. Nausea, vomiting, visual disturbances, bradycardia, atrioventricular block, ventricular dysrhythmias, and hyperkalemia may occur; ECG, potassium, and renal function should be monitored, and digoxin-specific

---

antibody fragments are used for life-threatening toxicity.

### Cardiac Hemangioma

A rare noncancerous tumor made of blood-vessel tissue in or around the heart. Many cause no symptoms, but a large or strategically located tumor may obstruct flow, disturb rhythm, or cause fluid around the heart.

### Cardiac Imaging Tests

Tests that show the heart's structure, movement, blood flow, or tissue. They include echocardiography, CT, MRI, and nuclear scans; the appropriate test depends on symptoms, the suspected condition, kidney function, and the clinical question.

### Cardiac Index

The amount of blood the heart pumps each minute adjusted for body size. It helps assess pumping function, especially in seriously ill people, but a single value must be interpreted with blood pressure, symptoms, organ function, and the overall clinical situation.

### Cardiac Intravascular Ultrasound

An ultrasound probe on a catheter that creates cross-sectional images from inside a coronary artery. It helps measure plaque, vessel size, narrowing, tears, calcium, and stent expansion during selected procedures.

### Cardiac Ischemia

Alternate name; see *Myocardial Ischemia*.

### Cardiac Markers

Blood tests used to detect myocardial injury or cardiac stress, including troponin, creatine kinase-MB, and natriuretic peptides. Results are interpreted with symptoms, electrocardiography, imaging, timing, and other causes of abnormal values.

### Cardiac Metastases

Cancer that has spread to the heart or its surrounding tissues from another organ. Lung and breast

cancers, melanoma, lymphoma, and kidney cancer are common sources; spread may cause fluid around the heart, obstruction, rhythm problems, or heart failure.

### Cardiac MRI

Magnetic-resonance imaging that shows heart structure, pumping, blood flow, scar, swelling, infiltrative disease, and congenital abnormalities without x-rays. Contrast may be avoided or used cautiously in selected people with severe kidney disease.

### Cardiac Muscle

Involuntary muscle that forms the myocardium, the main muscular layer of the heart. Its cells are electrically connected so the chambers contract in a coordinated way to pump blood.

### Cardiac Myocyte

A specialized heart-muscle cell that contracts to pump blood. Electrical signals and movement of calcium inside the cell coordinate its contraction and relaxation.

### Cardiac Myocytes

The muscle cells that make up most of the heart and contract with each heartbeat. Damage to them can reduce pumping function and release proteins such as troponin into the blood.

### Cardiac Myxoma

The most common primary noncancerous tumor of the heart, usually arising in the left atrium. It can obstruct blood flow, cause breathlessness or constitutional symptoms, or release clots; surgical removal is usually recommended.

### Cardiac Output

The volume of blood pumped by the heart each minute. It depends mainly on heart rate and stroke volume, the amount ejected with each beat, and helps indicate whether circulation is adequate.

### Cardiac Output Measurement

Measurement of blood pumped by the heart each

---

minute using methods such as echocardiography, a pulmonary-artery catheter, oxygen-consumption calculations, or noninvasive monitors. The method's accuracy depends on the patient and testing conditions.

### **Cardiac Pacemaker**

An implanted or temporary device that sends electrical impulses to maintain an adequate heartbeat when the heart's natural rhythm is too slow or unreliable. It may be placed inside the body or used externally during emergencies.

### **Cardiac Pacemaker Types**

Single-chamber pacemakers stimulate one heart chamber; dual-chamber devices coordinate the atrium and ventricle; biventricular or resynchronization devices stimulate both ventricles to improve timing in selected heart-failure patients.

### **Cardiac Perforation**

A hole or tear through the heart wall, sometimes caused by injury or a medical procedure. It can cause bleeding around the heart, cardiac tamponade, shock, and death and requires immediate treatment.

### **Cardiac PET**

A nuclear-imaging test using a positron-emitting tracer to assess heart-muscle blood flow, metabolism, and living or scarred tissue. Selected tracers can also help evaluate inflammation or infiltrative heart disease.

### **Cardiac PET Imaging**

Positron-emission tomography using a radioactive tracer to measure blood flow and metabolism in heart muscle. It can quantify blood flow and help distinguish living from scarred tissue, but availability, radiation exposure, and tracer choice affect its use.

### **Cardiac Power Output**

A measure combining blood pressure and cardiac output to estimate how effectively the heart is pumping. It is useful in severe heart failure or cardiogenic shock, but clinicians interpret it with examination

findings, organ function, and other circulation measurements.

### **Alternate Names:**

CPO

### **Cardiac Radiation Injury**

Damage to the heart or its blood vessels that may appear months or years after radiation to the chest. It can cause inflammation or scarring around the heart, coronary disease, valve disease, heart-muscle weakness, or rhythm and conduction problems.

### **Cardiac Radionuclide Imaging**

Heart imaging performed after a small radioactive tracer is given, usually with SPECT or PET. It can assess blood flow, metabolism, living or scarred heart muscle, inflammation, and, when synchronized with the heartbeat, pumping function.

### **Cardiac Rehab Outcomes**

Cardiac rehabilitation can improve exercise ability, blood-pressure and cholesterol control, confidence, quality of life, and psychological well-being after selected heart conditions. Benefits vary with the program, health condition, and participation.

### **Cardiac Rehabilitation**

A medically supervised program after conditions such as heart attack, coronary procedures, bypass surgery, or heart failure. It combines gradual exercise, education, risk-factor reduction, medicine support, nutrition, and emotional care to improve recovery and reduce future risk.

### **Cardiac Rehabilitation Phases**

Cardiac rehabilitation usually has three phases: hospital-based care soon after a cardiac event, supervised outpatient exercise and education, and long-term activity and risk-factor management. The program is adjusted to the person's condition and recovery.

### **Cardiac Resynchronization Therapy**

A pacing treatment that coordinates contraction of the heart's lower chambers. It is used in selected people with symptomatic heart failure and delayed



---

electrical activation and can improve symptoms, pumping, and survival when appropriate.

### **Cardiac Rhabdomyoma**

A usually noncancerous tumor of heart muscle, most often found in children and associated with tuberous sclerosis. It may shrink on its own but can obstruct blood flow or disturb rhythm and pumping if large.

### **Cardiac Sarcoidosis**

Sarcoidosis affecting the heart, where clusters of inflammatory cells can form in heart muscle and the electrical system. It may cause heart block, dangerous ventricular rhythms, heart failure, or sudden cardiac arrest.

### **Cardiac Sarcoma**

A rare cancer arising from supporting or connective tissue of the heart. It may obstruct blood flow, disturb heart rhythm, impair pumping, or release clots that travel to other parts of the body.

### **Cardiac Septum**

The wall separating the right and left chambers of the heart, consisting of atrial and ventricular portions and preventing normal mixing of oxygenated and deoxygenated blood.

### **Cardiac Skeleton**

A strong connective-tissue framework around the heart valves and parts of the wall between the chambers. It supports the valves, provides attachment for heart muscle, and helps electrically separate the upper and lower chambers.

### **Cardiac SPECT**

A nuclear scan that shows blood flow to the heart muscle and can help identify reduced flow, prior damage, or living muscle. When synchronized with the heartbeat, it can also assess chamber movement and pumping function.

### **Cardiac Stress Test**

A test that observes the heart while it works harder, usually during exercise or after a medicine increases

its workload. It may include an ECG or imaging and can help detect reduced blood flow or abnormal rhythms.

### **Cardiac Stress Testing**

Assessment of the heart during exercise or medicine-induced stress to look for reduced blood flow, abnormal rhythms, or limited pumping reserve. The method is chosen according to the person's ability to exercise and the clinical question.

### **Cardiac Syncope**

Fainting caused by a heart problem that briefly reduces blood flow to the brain. It may result from a very slow or fast rhythm, blocked blood flow, or severe heart weakness. Fainting during exercise or with heart disease needs urgent assessment.

### **Cardiac Syndrome X**

Angina-like chest pain with signs of reduced blood flow to the heart on stress testing but no significant blockage in the main coronary arteries on angiography. It is attributed to dysfunction of the tiny vessels that supply the heart muscle, called coronary microvascular dysfunction, and is more common in women. Symptoms may include chest pressure, burning, or discomfort with activity or at rest; diagnosis also requires excluding other causes of chest pain, including noncardiac ones.

#### **Alternate Names:**

Microvascular Angina

### **Cardiac Tamponade**

A life-threatening state in which pressure from fluid, blood, or pus in the sac around the heart impairs cardiac filling and reduces cardiac output. It can cause low blood pressure, breathlessness, fainting, raised neck veins, shock, or cardiac arrest. Echocardiography supports the diagnosis, but clinical assessment and the cause are also important; urgent drainage with a needle or surgery is often required.

#### **Alternate Names:**

Pericardial Tamponade; Hemopericardium with Tamponade; Cardiac Compression



---

### Cardiac Tamponade Physiology

Pressure from blood or fluid around the heart prevents the chambers from relaxing and filling. This lowers cardiac output and may cause low blood pressure, a marked breathing-related change in pulse pressure, and shock.

### Cardiac Transplantation

Replacement of a failing heart with a donor heart in carefully selected people with end-stage heart failure. Lifelong immune-suppressing treatment and regular monitoring are needed for rejection, infection, coronary-vessel disease, and other complications.

### Cardiac Tumors

Tumors involving the heart may begin there or spread from another organ. They are rare and may obstruct blood flow, disturb rhythm, cause fluid around the heart, or release clots; treatment depends on type, location, size, and spread.

### Cardiac Ventriculography

Imaging of the heart's lower chambers to assess their size, wall movement, and ability to pump blood. It may use x-rays with contrast or another imaging method and is now often replaced by echocardiography or MRI.

### Cardiac Volume

The amount of blood in a heart chamber at a particular time. The amount before contraction and after contraction helps assess chamber size, filling, and pumping function.

### Cardiac Wall

The three-layered wall of the heart: the inner endocardium, muscular myocardium, and outer epicardium. It supports the heart's structure, electrical activity, contraction, and interaction with surrounding tissues.

### Cardinal Body Part

A major body region or structure used in anatomical description or clinical examination, such as the head, neck, chest, abdomen, pelvis, limbs, or an important internal organ.

### Cardinal Cell Part

A main component of a cell, such as its membrane, cytoplasm, nucleus, or an organelle. Each part has a role in maintaining the cell, processing information, or performing specialized functions.

### Cardinal Ligament

A band of strong connective tissue extending from the cervix and upper vagina to the sidewall of the pelvis. It helps support the uterus and vagina and contains the uterine blood vessels, which cross over the ureter.

#### Alternate Names:

Mackenrodt's Ligament; Lateral Cervical Ligament; Transverse Cervical Ligament

### Cardinal Signs of Inflammation

The four classic signs of inflammation are heat (calor), pain (dolor), redness (rubor), and swelling (tumor). Inflamed tissue may also lose some normal function.

### Cardio-Oncology

The medical field focused on preventing and treating heart and blood-vessel problems in people with cancer. It monitors effects of chemotherapy, targeted or immune treatment, and chest radiation while helping cancer therapy continue safely when possible.

### Cardiobacteriaceae

A family of bacteria that includes *Cardiobacterium*. Some members can enter the bloodstream and infect heart valves, causing infective endocarditis.

### Cardiobacterium

A slow-growing bacterium in the HACEK group that can enter the bloodstream and infect heart valves. It may cause prolonged fever, fatigue, weight loss, a new murmur, or embolic complications.

### Cardiobacterium Hominis

The *Cardiobacterium* species most often associated with human infection. It can cause slowly developing infective endocarditis with fever, fatigue, weight

---

loss, a new heart murmur, valve damage, or clots traveling to other organs.

### **Cardiofaciocutaneous Syndrome**

A rare genetic disorder affecting the heart, facial appearance, skin, hair, growth, and development. It is usually caused by a change in a gene involved in cell-growth signals.

#### **Alternate Names:**

CFC Syndrome

### **Cardiogenic Shock**

A life-threatening state in which a severely weakened heart cannot supply enough blood to the body. It may cause low blood pressure, confusion, cold skin, little urine, breathing difficulty, and failure of important organs.

### **Cardiolipin**

A fat-like molecule found mainly in the inner membrane of mitochondria, the cell's energy-producing structures. Antibodies against cardiolipin can be associated with antiphospholipid syndrome and abnormal blood clotting.

### **Cardiologist**

A doctor trained to diagnose and treat diseases of the heart and blood vessels, including coronary disease, rhythm disorders, valve disease, heart failure, and high blood pressure.

### **Cardiology**

The branch of medicine concerned with the heart and blood vessels. It covers prevention, diagnosis, and treatment of coronary disease, heart failure, rhythm and valve disorders, cardiomyopathy, congenital heart disease, and high blood pressure.

### **Cardiomegaly**

Enlargement of the heart seen on examination or imaging. It may result from enlarged chambers, thickened heart muscle, fluid around the heart, or a combination; causes include high blood pressure, valve disease, cardiomyopathy, and congenital disease.

### **Cardiometabolic**

Relating to the interaction between heart and blood-vessel disease and metabolic conditions such as obesity, insulin resistance, diabetes, and abnormal blood lipids.

### **Cardiomyopathy**

A disease of the heart muscle that can make it enlarged, thickened, stiff, or weak. These changes can make it harder for the heart to pump or fill with blood and may cause heart failure or abnormal heart rhythms. Major types include dilated, hypertrophic, and restrictive cardiomyopathy.

### **Cardiomyopathy Classification**

A system for describing cardiomyopathy by its heart structure and function, other organs involved, genetic pattern, cause, and functional severity. This information helps communicate diagnosis, guide testing, and estimate prognosis.

### **Cardiopathy**

Any disease or disorder of the heart, including problems of the heart muscle, valves, coronary arteries, electrical rhythm, or structure present from birth.

### **Cardioplegia**

Temporary stopping and protection of the heart during open-heart surgery, usually by delivering a cold solution that changes electrical and metabolic activity. It allows the surgeon to operate while limiting heart-muscle injury.

### **Cardiopulmonary**

Relating to the heart and lungs or to their combined function. For example, cardiopulmonary exercise testing assesses how the heart, lungs, blood, and muscles respond during physical activity.

#### **Alternate Names:**

Cardiorespiratory

### **Cardiopulmonary Bypass**

A heart-lung machine temporarily takes over circulation and oxygen exchange during some operations. It pumps and oxygenates blood while the heart is

---

stopped or opened; possible complications include bleeding, inflammation, clots, stroke, and organ injury.

### **Cardiopulmonary Bypass Machine**

A machine that temporarily pumps and oxygenates blood during selected heart or major-vessel operations. It allows surgeons to work while the heart is stopped or the lungs are not carrying out their usual gas exchange.

#### **Alternate Names:**

Heart-Lung Machine; Cardiopulmonary Bypass Pump

### **Cardiopulmonary Exercise Testing**

A monitored treadmill or bicycle test that measures how the heart, lungs, blood, and muscles respond as exercise becomes harder. It helps assess unexplained breathlessness, exercise limitation, and severity of heart or lung disease.

### **Cardiopulmonary Resuscitation**

An emergency life-saving procedure performed when a person has suffered cardiac arrest—meaning their heart has suddenly stopped beating effectively, causing them to collapse, lose consciousness, and stop breathing normally. CPR combines rapid, rhythmic chest compressions (pressing hard and fast in the center of the chest at a rate of 100–120 compressions per minute) to manually pump oxygen-rich blood to the brain and vital organs, with artificial ventilation (rescue breaths) to deliver oxygen into the lungs. In out-of-hospital emergencies, untrained bystanders can perform “hands-only” CPR by giving continuous high-quality chest compressions without rescue breaths until professional help arrives. Immediate CPR, combined with the prompt use of an automated external defibrillator (AED) to shock lethal irregular rhythms, can double or triple a person’s chance of surviving sudden cardiac arrest.

#### **Alternate Names:**

CPR; Basic Life Support; BLS; Closed-Chest Cardiac Massage

### **Cardiorenal Syndrome**

A condition in which heart and kidney dysfunction

worsen one another. Sudden or long-term heart disease can injure the kidneys, and sudden or long-term kidney disease can strain the heart; treatment must address both organs.

### **Cardiorespiratory**

Relating to the heart, blood vessels, and respiratory system considered together, as in cardiorespiratory fitness, monitoring, or failure.

### **Cardiorespiratory Endurance**

The ability of the heart, lungs, blood, and muscles to deliver and use oxygen during sustained physical activity. It is influenced by fitness, heart and lung disease, anemia, muscle condition, and other factors.

#### **Alternate Names:**

Cardiorespiratory Fitness; Aerobic Endurance

### **Cardioselective Beta-Blockers**

Beta-blocker medicines that mainly block beta-1 receptors in the heart, slowing the heart rate and reducing its workload. They may still narrow airways at higher doses and are used for selected heart, blood-pressure, and rhythm conditions.

### **Cardiospasm**

Older term; see *Achalasia*.

### **Cardiotocography**

Continuous recording of a fetus’s heart rate and the mother’s uterine contractions during pregnancy or labor. The pattern helps assess fetal well-being, but results must be interpreted with the stage of labor and the overall clinical situation.

### **Cardiotoxicity**

Injury or impaired function of the heart caused by a medicine, toxin, radiation, or disease. It may produce rhythm problems, reduced heart-muscle strength, chest pain from poor blood flow, or heart failure.

### **Cardiotoxicity From Chemotherapy**

Heart injury caused by some chemotherapy medicines. It may weaken heart muscle, reduce

---

pumping, or cause rhythm problems during or after treatment; risk depends on the medicine, dose, other treatments, and existing heart disease.

### **Cardiotoxin**

A substance that directly injures heart muscle or disrupts the heart's electrical activity or blood supply. It may cause abnormal rhythms, reduced pumping, heart failure, or heart-muscle injury.

### **Cardiovascular**

Relating to the heart and blood vessels, including circulation, blood pressure, and diseases affecting the heart or vascular system.

### **Cardiovascular Agents**

Medicines that act on the heart or blood vessels. They include treatments for high blood pressure, abnormal rhythms, angina, heart failure, blood clots, and abnormal cholesterol levels.

### **Cardiovascular Changes In Pregnancy**

During pregnancy, blood volume and cardiac output increase to support the uterus and developing baby. The rise in plasma is greater than the rise in red cells, causing normal dilutional anemia; heart rate and blood-vessel resistance also change.

### **Cardiovascular Disease**

Disease of the heart or blood vessels, including coronary artery disease, heart failure, abnormal rhythms, valve disease, stroke-related vascular disease, and disorders of arteries or veins.

### **Cardiovascular Disease and Pregnancy**

Heart disease may worsen during pregnancy because blood volume and cardiac output increase, while pregnancy can reveal previously unrecognized disease. Risk depends on the specific lesion and ventricular function; coordinated cardiology and maternal-fetal care guides monitoring, medicines, and delivery.

### **Cardiovascular Health**

The condition of the heart and blood vessels and

the habits that protect them. Regular activity, not using tobacco, healthy eating, adequate sleep, and control of blood pressure, cholesterol, diabetes, and weight can lower cardiovascular risk.

### **Cardiovascular Infection**

An infection involving the bloodstream, heart, heart valves, or blood vessels. Examples include infective endocarditis, myocarditis, infected vascular grafts, and infected pacemakers or other devices.

### **Cardiovascular Injury**

Damage to the heart or blood vessels caused by trauma, reduced blood flow, inflammation, toxins, surgery, or another disease. Effects may include bleeding, impaired circulation, heart-muscle injury, or abnormal rhythms.

### **Cardiovascular Neoplasm**

A noncancerous or cancerous tumor arising in the heart, blood vessels, or related tissues. Its symptoms and treatment depend on the tissue type, location, size, effect on circulation, and whether it has spread.

### **Cardiovascular Pharmacology**

The study of how medicines affect the heart, blood vessels, blood pressure, rhythm, blood clotting, and cholesterol or lipid metabolism, including their uses, interactions, benefits, and risks.

### **Cardiovascular Prevention in Diabetes**

Medical and lifestyle strategies to reduce the high risk of heart attacks and strokes in diabetic patients, including blood pressure control, statin therapy, glucose management, smoking cessation, and healthy diet.

#### **Alternate Names:**

Diabetic Cardiovascular Risk Reduction; Heart Disease Prevention in Diabetes

### **Cardiovascular Risk Reduction**

Actions that lower the chance of heart attack, stroke, heart failure, and vascular death. They include avoiding tobacco, controlling blood pressure, cholesterol, and diabetes, staying active, eating well, maintaining a healthy weight, and taking indicated

---

medicines.

### **Cardiovascular Screening**

Assessment for cardiovascular risk or disease, often before symptoms develop. It may include blood-pressure measurement, cholesterol and diabetes testing, review of family history and tobacco use, and targeted tests based on age and risk.

### **Cardiovascular Surgery**

Operations on the heart or major blood vessels, including bypass surgery, valve repair or replacement, aortic surgery, and heart transplantation. Many open procedures require a heart-lung machine and specialized care before and after surgery.

### **Cardiovascular System**

The heart, blood, and network of arteries, arterioles, capillaries, venules, and veins that circulate blood through the pulmonary and systemic circuits. It delivers oxygen, nutrients, hormones, and immune cells, carries carbon dioxide and other waste for removal, and helps regulate temperature and blood pressure.

### **Cardiovascular-Kidney-Metabolic (CKM) Syndrome**

A framework describing interconnected cardiovascular, kidney, and metabolic dysfunction, often involving obesity, diabetes, hypertension, dyslipidemia, or chronic kidney disease. Risk is assessed across stages to guide prevention and coordinated care.

### **Cardioversion**

Restoration of a more regular heart rhythm using a synchronized electrical shock or medicine. Before cardioversion for some atrial rhythms, clinicians assess clot risk and may give anticoagulant treatment to reduce the risk of stroke.

### **Cardiovirus**

A group of viruses in the picornavirus family. Some infect people or animals and may cause fever, muscle inflammation, heart inflammation, or other illness; effects depend on the species and the person's defenses.

### **Carditis**

Inflammation of the heart muscle, inner lining, valves, or surrounding sac. It may result from infection, an immune reaction, medicines, or another inflammatory disease and can cause chest pain, breathlessness, rhythm problems, or heart failure.

### **Care Coordination**

Organizing communication and health-care activities among clinicians, services, patients, and caregivers so care remains safe, continuous, and appropriate when people move between settings or receive several types of treatment.

### **Caregiver**

A family member, friend, or paid worker who helps another person with health care, personal activities, daily tasks, transportation, medicines, or emotional support.

### **Caregiver Burden**

The emotional, physical, social, or financial strain of providing ongoing care for another person. It may lead to exhaustion, low mood, sleep problems, isolation, or difficulty meeting one's own health needs.

### **Caregiver Stress**

Strain caused by caring for another person, often leading to tiredness, worry, irritability, sleep problems, or reduced well-being. Support, respite, counseling, and practical help can reduce its effects.

### **Caries**

A biofilm-mediated, multifactorial disease in which acids from bacterial metabolism of fermentable carbohydrates cause net demineralization of enamel, dentin, or cementum. Early lesions may be arrested or remineralized, but progressive disease forms cavities and can reach the pulp, causing pain or infection.

### **Caries Risk Assessment**

Evaluation of a person's chance of developing tooth decay by reviewing past and current cavities, plaque, diet, fluoride exposure, saliva, oral hygiene, tooth



---

structure, medicines, and medical conditions. Results guide prevention and follow-up.

### **Carina**

The ridge at the bottom of the windpipe where it divides into the right and left main bronchi. It is a landmark during bronchoscopy and breathing-tube placement; inhaled objects more often enter the wider, more vertical right bronchus.

### **Cariogenic**

Likely to promote tooth decay, especially by supplying sugars that plaque bacteria convert into acids or by supporting conditions that remove minerals from teeth.

### **Cariogram**

A computer-based tool that combines factors such as diet, plaque bacteria, fluoride exposure, saliva, previous cavities, and social or clinical history to estimate tooth-decay risk and show where prevention may help.

### **Carisoprodol**

A prescription muscle relaxant used short term for acute painful muscle conditions. It can cause drowsiness, dizziness, impaired coordination, misuse, dependence, and dangerous breathing suppression, especially with alcohol or sedatives.

### **Carminative**

A substance traditionally used to ease intestinal gas or bloating. It may relax intestinal muscle or help gas pass, but persistent pain, vomiting, swelling, or altered bowel habits need medical assessment.

#### **Alternate Names:**

Antiflatulent; Gas-Relieving Agent

### **Carmine**

A red coloring made from cochineal insects and used in some foods, cosmetics, and medicines. Rarely, it causes allergy, wheezing, hives, or anaphylaxis in sensitive people.

### **Carmustine**

A chemotherapy medicine that damages DNA and is used for selected brain tumors, blood cancers, and other cancers. It can lower blood counts and cause nausea, infertility, liver problems, or delayed lung injury.

### **Carnett Sign**

A clinical diagnostic test used to distinguish somatic abdominal wall pain from visceral intra-abdominal pain. The examiner palpates the point of maximal tenderness while the patient tenses the abdominal wall musculature by raising their head and shoulders or lifting both straight legs. Increased or unchanged pain indicates abdominal wall origin (positive Carnett sign), whereas diminished pain suggests an intra-abdominal visceral source.

#### **Alternate Names:**

Carnett's Sign; Abdominal Wall Tenderness Test

### **Carney Complex**

A rare inherited disorder, usually caused by a *PRKAR1A* gene change, that can produce multiple noncancerous skin, heart, and endocrine tumors. It may also cause excess hormones and requires lifelong, organ-specific monitoring.

### **Carney Triad**

A rare condition involving combinations of a stomach gastrointestinal stromal tumor, a benign cartilage tumor in the lung, and a paraganglioma outside the adrenal gland. It usually occurs sporadically and more often affects young women.

### **Carnitine**

A substance that transports long-chain fats into mitochondria, the cell structures that make energy, so the fats can be used as fuel. The body usually makes enough; severe deficiency can cause weakness, low blood sugar, or heart and muscle disease.

### **Carnitine Deficiency**

An abnormally low amount of carnitine, which transports long-chain fatty acids into mitochondria for energy production. Primary inherited or secondary deficiency can cause muscle weakness, low blood



---

sugar, cardiomyopathy, or episodes of metabolic decompensation.

### **Carnitine Palmitoyltransferase II Deficiency**

A rare inherited disorder in which muscle cells cannot use long-chain fats normally for energy. Exercise, fasting, cold, or infection can trigger muscle pain, stiffness, muscle breakdown, dark urine, and kidney injury; symptoms may be absent between attacks.

#### **Alternate Names:**

CPT II Deficiency; CPT2 Deficiency

### **Carnosine**

A small molecule made from beta-alanine and histidine, found mainly in muscle and nervous tissue. It helps buffer acidity; claims that supplements prevent disease or improve performance remain uncertain.

### **Carnosinemia**

A rare inherited disorder in which deficient carnosinase causes excess carnosine in blood and urine. Children may have developmental delay, low muscle tone, seizures, or other neurologic problems of varying severity.

### **Caroli Disease**

A rare inherited disorder in which the larger bile ducts inside the liver are widened in segments. It can cause repeated bile-duct infections, stones, abdominal pain, jaundice, and, in some people, liver scarring or kidney disease.

### **Carotene**

An orange-yellow plant pigment found in foods such as carrots and leafy vegetables. Some forms, especially beta-carotene, can be converted to vitamin A; excess intake may harmlessly turn the skin yellow-orange.

### **Carotenemia**

Yellow-orange discoloration of the skin caused by increased carotene in the blood, usually from eating large amounts of carotene-rich foods or supplements. The whites of the eyes remain white, which helps distinguish it from jaundice; persistent or unexplained

discoloration needs assessment.

#### **Alternate Names:**

Carotenoid Pigments; Provitamin-a Pigments

### **Carotenoid**

A family of fat-soluble yellow, orange, or red pigments found mainly in plants and some microorganisms. Some, such as beta-carotene, can form vitamin A; others, such as lutein, support eye tissues.

### **Carotenoids**

Plant pigments that give many fruits and vegetables their yellow, orange, or red color. Some can be converted to vitamin A, while others have roles in eye and cellular function.

### **Carotid**

Relating to either of the two major arteries in the neck that carry blood from the heart toward the brain, face, and other head and neck structures.

### **Carotid Artery**

One of the major arteries supplying the head and neck. Each common carotid divides into an internal branch that supplies the brain and an external branch that supplies much of the face, scalp, and neck.

### **Carotid Artery Disease**

Narrowing or blockage of a carotid artery, usually from fatty plaque. It can reduce blood flow to the brain or release a clot and cause a transient ischemic attack or ischemic stroke.

### **Carotid Artery Dissection**

A tear in the inner lining of a carotid artery that lets blood split the arterial wall and form an intramural hematoma. It may cause sudden neck or head pain, partial Horner syndrome, transient ischemic attack, or ischemic stroke; urgent vascular imaging is needed.

### **Carotid Artery Stenosis**

Narrowing of an artery in the neck that supplies the brain, usually from fatty plaque. It may reduce

---

blood flow or release a clot and increase the risk of a transient ischemic attack or stroke.

### **Carotid Artery Stenting**

A procedure that places a small mesh tube inside a narrowed carotid artery to keep it open. It is used selectively to reduce stroke risk; clot release, stroke, bleeding, and re-narrowing are possible complications.

### **Carotid Body**

A small group of sensing cells near the division of the common carotid artery. It detects changes in blood oxygen, carbon dioxide, and acidity and sends signals that help adjust breathing.

### **Carotid Body Tumor**

A rare, highly vascular, usually benign growth arising from the oxygen-sensing cells (carotid body) at the fork of the main neck artery. It typically presents as a painless, pulsating lump below the jaw that moves side-to-side but not up-and-down; as it grows, it can press on nearby nerves and cause hoarseness or swallowing difficulties, and surgical removal is the standard treatment.

#### **Alternate Names:**

Carotid body tumour; Chemodectoma; Carotid paraganglioma; Glomus caroticum tumor.

### **Carotid Bruit**

A blowing sound heard over a carotid artery when blood flow is turbulent. It can occur with plaque but does not reliably show how narrow the artery is; ultrasound or other vascular imaging is needed when disease is suspected.

### **Carotid Cavernous Fistula**

An abnormal connection between blood vessels near the carotid artery and a large vein behind the eye. It may cause a bulging or red eye, a pulsing noise, double vision, high eye pressure, or reduced vision. Brain and blood-vessel imaging is used for diagnosis. Treatment depends on the type and may include endovascular closure.

#### **Alternate Names:**

Caroticocavernous Sinus Fistula; CCF; Carotid-

Cavernous Sinus Fistula; Cavernous Sinus Fistula

### **Carotid Duplex Ultrasound**

An ultrasound test that shows plaque and measures blood flow through the carotid arteries. It helps estimate narrowing without radiation and may guide decisions about medicines, surgery, or stenting.

### **Carotid Endarterectomy**

Surgery to open a carotid artery and remove fatty plaque from its inner wall. It can reduce stroke risk in selected people with significant narrowing, especially after symptoms, but carries risks such as stroke, heart attack, and nerve injury.

### **Carotid Glomus**

A specialized peripheral chemoreceptor sensory organ located in the adventitia of the carotid bifurcation, containing glomus type I cells that detect arterial hypoxemia, hypercapnia, and acidosis to reflexively stimulate respiration via the glossopharyngeal nerve.

#### **Alternate Names:**

Carotid Body

### **Carotid Occlusion**

Complete blockage of an artery supplying the brain, usually from plaque or a blood clot. Some people have no symptoms because other vessels compensate, but sudden weakness, speech trouble, or vision loss is an emergency.

### **Carotid Sinus Hypersensitivity**

An exaggerated reflex response to tactile mechanical stimulation of the carotid sinus baroreceptors, producing marked bradycardia (cardioinhibitory type), profound hypotension (vasodepressor type), or both, predisposing to presyncope or syncope.

#### **Alternate Names:**

Carotid Sinus Syndrome

### **Carotid Sinus Massage**

A medical test in which a clinician gently presses over a neck artery to see how the heart rhythm responds. It may help assess or stop some fast heart

---

rhythms, but it can be dangerous and must only be done by a trained clinician after checking for risks.

**Alternate Names:**

Carotid Sinus Stimulation; Carotid Baroreceptor Massage

**Carotid Stenosis**

Narrowing of a carotid artery, usually from fatty plaque. It may reduce blood flow to the brain or release a clot, causing a transient ischemic attack or stroke; treatment depends on symptoms, severity, and overall risk.

**Carpal Bones**

The eight small bones forming the wrist: scaphoid, lunate, triquetrum, pisiform, trapezium, trapezoid, capitate, and hamate. They connect the forearm to the hand and work together to allow wrist movement and transmit force.

**Alternate Names:**

Carpals

**Carpal Joints**

The joints between the wrist bones and between the wrist, forearm, and hand bones. Together they allow the wrist to bend, straighten, move side to side, and rotate with the forearm.

**Carpal Tunnel**

A narrow passage on the palm side of the wrist formed by the wrist bones and a strong ligament. It contains the median nerve and finger-flexing tendons; pressure on the nerve causes carpal tunnel syndrome.

**Carpal Tunnel Biopsy**

Removal of a small tissue sample from the carpal tunnel or its contents when an unusual infiltrative, inflammatory, infectious, or tumor-related process is suspected. It is not a routine test for ordinary carpal tunnel syndrome.

**Carpal Tunnel Release**

Surgery that cuts the ligament forming the roof of the wrist tunnel to relieve pressure on the median

nerve. It may improve numbness, tingling, pain, and weakness when symptoms do not respond to suitable nonsurgical care.

**Carpal Tunnel Syndrome**

Pressure on the median nerve as it passes through the narrow wrist tunnel. It commonly causes numbness, tingling, burning, or pain in the thumb, index, middle, and part of the ring finger, sometimes with thumb weakness.

**Alternate Names:**

CTS; Median Nerve Entrapment

**Carpal Tunnel Syndrome Tests**

Examination and nerve tests used to assess suspected median-nerve compression. Wrist-position or tapping tests may reproduce symptoms, while nerve-conduction studies and ultrasound can support diagnosis and assess severity when needed.

**Carpometacarpal Joint**

A joint between a wrist bone and the base of a hand bone. The thumb joint is a mobile saddle-shaped joint that allows opposition and is commonly affected by osteoarthritis.

**Carpus**

The group of eight small bones forming the wrist between the forearm and the metacarpals of the hand. The carpus provides stability while allowing controlled wrist movement.

**Carrageenan**

A substance extracted from red seaweed and used to thicken or stabilize some foods and medicines. Different forms have different properties; evidence about effects on human inflammation depends on the type and amount consumed.

**Carrier**

A person who has one altered copy of a gene for a recessive condition and usually has no symptoms. A carrier can pass the altered gene to a child; risk depends on the other parent's genes and the inheritance pattern.

---

### Carrier Excipient

An inactive ingredient that supports an active medicine in a formulation. It may help control the medicine's dose, stability, absorption, appearance, or delivery and can occasionally cause allergy or intolerance.

### Carrier Screening

Genetic testing for people without symptoms to find whether they carry a harmful gene variant that could be passed to a child. Results can guide reproductive choices and may identify the need to test a partner or relatives.

### Carrier State

A condition in which a person carries and may spread a microorganism without symptoms. Some carriers need testing, treatment, vaccination of contacts, or other measures to prevent transmission, depending on the infection.

### Carrier-Mediated Transport

Movement of a substance across a cell membrane with help from a specific carrier protein. It may use no direct energy when moving down a concentration gradient or use energy to move against that gradient.

### Carry-Over Effect

Persistence of an earlier treatment's effect into a later measurement or treatment period. It can distort results in crossover studies or make a person's current clinical state difficult to interpret.

### Carteolol

A nonselective beta-blocker used mainly as eye drops to lower pressure inside the eye in open-angle glaucoma or ocular hypertension. It can sometimes slow the heartbeat, lower blood pressure, or worsen breathing problems.

### Cartilage

A tough, flexible tissue found in joints, the ear, nose, rib cage, and spinal discs. It cushions joints, supports body structures, and resists pressure, but it has limited blood supply and heals slowly after injury.

### Cartilage Disease

A disorder that damages cartilage in joints or supporting structures. It can cause pain, stiffness, swelling, deformity, impaired growth, or loss of smooth movement and may result from injury, aging, inflammation, or inherited disease.

### Cartilaginous Joint

A joint in which bones are connected by cartilage and allow little or limited movement. Examples include joints between spinal bones and the joint joining the two front pelvic bones.

### Caruncles

Small fleshy anatomical projections. Examples include the lacrimal caruncle at the inner corner of the eye and the urethral caruncle near the opening of the urethra.

### Carvallo Sign

A physical examination finding in which the pansystolic murmur of tricuspid regurgitation increases in intensity during inspiration. The accentuation is caused by increased venous return to the right heart during the inspiratory phase of respiration.

#### Alternate Names:

Carvallo's Sign; Rivero-Carvallo Sign

### Carvedilol

A beta-blocker that also relaxes blood vessels. It slows the heart and reduces its workload and is used for selected cases of high blood pressure, heart failure, and after a heart attack; it can lower blood pressure and heart rate.

### CAS

Alternate name; see *Childhood Apraxia of Speech*.

### Casanthranol

A stimulant laxative that increases bowel movement and secretion of water into the intestine. Prolonged or excessive use can cause diarrhea, dehydration, electrolyte abnormalities, and reliance on laxatives.

---

## Cascade

A series of linked reactions in which one activated step triggers the next. Examples include the blood-clotting, complement, and inflammatory cascades.

## Cascara

A plant-derived stimulant laxative that increases bowel movement. Prolonged or excessive use can cause diarrhea, dehydration, electrolyte imbalance, and reliance on laxatives; its use is restricted in some places.

## Case Definition

Standardized clinical, laboratory, and exposure criteria used to classify a person as a suspected, probable, or confirmed case during surveillance or an outbreak investigation. It may differ from the criteria used to diagnose one patient.

## Case Fatality Rate

The proportion of people diagnosed with a disease who die from it during a stated period. It describes severity in a particular group, but depends on accurate diagnosis, follow-up, treatment, and the population studied.

## Case Management

Coordinated planning, referral, and follow-up of health and social services to help a person receive appropriate care, avoid gaps or duplication, and meet clinical, functional, and personal goals.

## Case-Fatality Rate

The proportion of diagnosed people who die from a disease during a stated period, calculated as deaths from that disease divided by diagnosed cases. It estimates severity in the group studied and is not the same as population mortality.

## Casein

The main family of proteins in cow's milk and other mammalian milk. It supplies amino acids and minerals and forms curds during digestion and cheese-making; it can trigger reactions in people with cow's-milk protein allergy.

## Casein Kinase

A family of enzymes that attach phosphate groups to proteins and thereby change their activity. Casein kinases help regulate cell signaling, growth, DNA repair, development, and the body's internal clock.

## Caseous

Having a soft, crumbly, cheese-like appearance. The term often describes dead tissue in granulomas caused by tuberculosis or certain other long-lasting infections.

## Caseous Necrosis

A form of tissue death producing soft, crumbly, white or yellow material that resembles cheese. It is classically seen in granulomas caused by tuberculosis, though similar changes can occur in some other chronic infections.

## Cashew Allergy

An immune reaction to cashew proteins that may cause itching, hives, swelling, vomiting, wheezing, or life-threatening anaphylaxis. People with this allergy should avoid cashews and follow an emergency action plan.

## Caspase

A group of enzymes that cut specific proteins inside cells. Some initiate orderly programmed cell death, while others activate inflammation and immune defense against infection.

## Caspase Activation

Conversion of inactive caspase enzymes into their active form inside a cell. Depending on the caspase and triggering signal, this can start programmed cell death or activate inflammation and immune defense.

## Caspofungin

An intravenous antifungal medicine that blocks construction of an important fungal cell-wall component. It is used for serious *Candida* infections and selected *Aspergillus* infections and may cause liver-test abnormalities, infusion reactions, or low potassium.

---

## Caspofungin Acetate

The acetate salt form of caspofungin, an intravenous antifungal medicine that blocks fungal cell-wall construction. It is used for selected serious *Candida* and *Aspergillus* infections under medical supervision.

## Cassette

A holder that contains recording or imaging material. In traditional radiography, it held the film or image receptor during exposure; the word may also describe holders used in laboratory or monitoring equipment.

## Cast

A rigid or semi-rigid support placed around an injured limb or joint to limit movement while a bone, ligament, tendon, or other tissue heals. It must be monitored for swelling, pressure injury, numbness, or poor circulation.

## Cast Nephropathy

A classic renal manifestation of multiple myeloma in which excessive circulating monoclonal immunoglobulin free light chains (Bence-Jones proteins) precipitate with Tamm-Horsfall mucoprotein in distal tubules, forming dense casts that obstruct tubules and induce giant cell inflammation.

### Alternate Names:

Myeloma Kidney

## Castleman Disease

A group of uncommon disorders causing abnormal enlargement and activity of lymph nodes. Disease may affect one lymph-node area or the whole body and can cause fever, fatigue, anemia, weight loss, and organ problems.

## Castor Oil

A vegetable oil from the seeds of *Ricinus communis*. Taken by mouth, it acts as a stimulant laxative; excessive use can cause cramping, diarrhea, dehydration, and electrolyte disturbances.

## Castor Oil Overdose

Excessive ingestion of castor oil, causing intestinal

irritation, cramping, nausea, vomiting, and diarrhea. Severe exposure can cause dehydration, low blood pressure, or fluid and electrolyte imbalance.

## Castration

Removal or functional suppression of the testes or ovaries, reducing production of sex hormones. It may be surgical or achieved with medicines and can affect fertility, sexual function, mood, and bone health.

## Cat Allergy

An immune reaction to proteins in cat saliva, skin flakes, or other secretions. It can cause sneezing, itchy eyes, wheezing, hives, or asthma symptoms; reducing exposure and using appropriate allergy treatment may help.

## CAT Scan

Older name; see *Computed Tomography*.

## Cat Scratch Disease

An infection caused by *Bartonella henselae*, usually after a scratch or bite from a kitten or cat. It commonly causes a small skin lesion and nearby swollen lymph nodes and is usually self-limited, though serious complications can occur.

## Cat's Eye Reflex

A bright white reflection from the pupil, called leukocoria, rather than the usual red reflection in photographs. It can signal cataract, retinal disease, or retinoblastoma and requires urgent eye assessment, especially in a child.

## Cat-Nap

A short period of sleep, usually during the day. Frequent unintended sleep may indicate inadequate sleep, a medicine effect, or a sleep disorder and should be assessed if it affects safety or daily life.

## Cat-Scratch Disease

An infection caused by *Bartonella henselae*, usually after a scratch or bite from an infected cat. It commonly causes a small skin bump, nearby swollen



---

lymph nodes, fever, and fatigue; rarely it affects the eye, liver, spleen, brain, or heart.

**Alternate Names:**

Cat-Scratch Fever

**Catabolism**

The set of chemical processes that break down complex molecules such as carbohydrates, fats, and proteins into simpler substances, releasing energy that cells can use. Examples include digestion-related breakdown, glycolysis, and fat oxidation.

**Catagen**

The short transition phase of the hair cycle between active growth and rest. The hair follicle shrinks, hair growth stops, and the hair remains attached until the later shedding phase.

**Alternate Names:**

Catagen Phase; Regression Phase of Hair Growth

**Catalase**

An enzyme that breaks down hydrogen peroxide into water and oxygen. It protects cells from oxidative damage and helps some bacteria survive inside immune cells.

**Catalepsy**

A state of greatly reduced movement and responsiveness in which a person may remain rigid or hold an imposed posture. It can occur with catatonia, narcolepsy, neurologic disease, or some psychiatric conditions.

**Catalyst**

A substance that speeds a chemical reaction without being used up. It works by providing an easier reaction pathway; enzymes are biological catalysts, and catalysts change reaction speed without changing the final balance of products and reactants.

**Catamenial Epilepsy**

A pattern in which seizures become more frequent or severe at certain times in the menstrual cycle. Hormone changes may contribute, and a seizure diary can help guide neurologic treatment.

**Catamenial Pneumothorax**

Repeated collapse of a lung occurring around the start of menstruation, usually because of thoracic endometriosis. It often affects the right side and causes sudden chest pain or breathlessness; recurrent cases need specialist evaluation and may require surgery or hormone treatment.

**Alternate Names:**

Endometriosis-Related Pneumothorax; Catamenial Thorax

**Cataplexy**

A sudden, brief loss of muscle strength triggered by strong emotion such as laughter, surprise, or anger. Awareness usually remains clear; it is strongly associated with narcolepsy and may cause the head, jaw, knees, or whole body to buckle.

**Cataract**

Clouding of the eye's natural lens, which reduces the passage of light to the retina. It can cause blurred or faded vision, glare, poor contrast, and difficulty seeing at night.

**Cataract Removal**

Surgery that removes a cloudy natural lens from the eye and usually replaces it with an artificial intraocular lens. It is performed when the cataract interferes with vision, work, driving, reading, or other daily activities.

**Cataract Surgery**

Removal of a clouded eye lens and replacement with an artificial lens to improve vision. Possible complications include infection, bleeding, retinal problems, increased eye pressure, or a need for later treatment, though serious complications are uncommon.

**Cataracts**

Alternate index label; see *Cataract*.

**Alternate Names:**

Adult Cataracts; Cataracts in Adults

**Catastrophic Aging**

A rapid loss of physical or mental function in an

---

older person after illness, an operation, a fall, or another stress. Limited ability to recover can make decline harder to reverse; early treatment and rehabilitation may help.

### **Catastrophic Reaction**

An unusually intense emotional or behavioral response to a situation that may seem minor to others. It can occur with dementia, brain injury, or other neurologic disease and may involve shouting, fear, crying, or agitation.

#### **Alternate Names:**

Catastrophic Response; Catastrophic Emotional Reaction

### **Catatonia**

A serious syndrome involving marked changes in movement, speech, and response to the environment. Features may include immobility, mutism, staring, unusual postures, rigidity, repetitive movements, or copying speech or actions; causes include psychiatric and medical disorders.

### **Catbite**

A puncture wound caused by a cat's teeth. Cat bites can push bacteria deep into tissue and may infect tendons, joints, or bone, so prompt cleaning and medical assessment are often needed.

### **Catch-Up Immunization**

Vaccination of a person who is behind the recommended schedule using age- and risk-appropriate intervals. A delayed series usually does not need to be restarted, but minimum ages, intervals, and medical precautions must be followed.

### **Catechin**

A group of plant compounds found in tea, cocoa, some fruits, and other foods. They show antioxidant effects in laboratory studies, but catechin supplements are not proven treatments and may cause side effects or interact with medicines.

### **Catechol**

A chemical compound with two neighboring hydroxyl groups, used in industry and involved in

making some biologically active substances. Contact or inhalation can irritate tissues and substantial exposure may be toxic.

### **Catecholamine**

A group of hormones and nerve-signaling chemicals that includes epinephrine, norepinephrine, and dopamine. They help control the body's stress response, heart rate, blood pressure, alertness, and metabolism.

### **Catecholamine Blood Test**

A blood test measuring epinephrine, norepinephrine, and dopamine. It may help evaluate unusual hormone release from a pheochromocytoma or related tumor, although metanephrine testing is often preferred for initial screening.

### **Catecholamine Crisis**

A sudden, life-threatening surge of stress hormones causing very high blood pressure, rapid or irregular heartbeat, sweating, headache, and possible injury to the heart, brain, kidneys, or other organs. It can occur with certain adrenal or nerve tumors.

### **Catecholaminergic Polymorphic Ventricular Tachycardia**

A rare inherited rhythm disorder in which exercise or strong emotion triggers dangerous fast rhythms from the heart's lower chambers. The heart may look normal and the resting ECG may be normal; fainting or cardiac arrest can occur, especially in children and young adults.

#### **Alternate Names:**

CPVT; Bidirectional Ventricular Tachycardia; Stress-Induced Polymorphic VT

### **Catecholamines**

Adrenaline, noradrenaline, and dopamine are catecholamines. These hormones and nerve-signaling chemicals help regulate stress responses, blood pressure, heart function, alertness, and metabolism.

### **Category Five Hurricane**

A Saffir–Simpson hurricane with sustained winds of at least 157 miles per hour (252 km/h), capa-

---

ble of catastrophic damage and prolonged loss of habitability.

### **Category I Fetal Heart Tracing**

A reassuring fetal heart-rate pattern with a baseline of 110–160 beats per minute, moderate variability, and no late or variable decelerations. It is usually consistent with adequate fetal oxygenation at that time.

### **Category II Fetal Heart Tracing**

A fetal heart-rate pattern that is not clearly normal or abnormal. It requires continued monitoring and assessment of the mother, fetus, labor, and changes over time because it may become reassuring or concerning.

### **Category III Fetal Heart Tracing**

An abnormal fetal heart-rate pattern that may indicate inadequate fetal oxygenation or acid buildup. It includes absent variability with recurrent late or variable decelerations or a true sinusoidal pattern and requires prompt clinical evaluation and action.

### **Category Three Hurricane**

A Saffir–Simpson hurricane with sustained winds of 111–129 miles per hour (178–208 km/h), capable of devastating damage.

### **Category Two Hurricane**

A Saffir–Simpson hurricane with sustained winds of 96–110 miles per hour (154–177 km/h), capable of extensive damage to structures and vegetation.

### **Catenin**

A family of proteins that helps neighboring cells attach and communicate. Beta-catenin also carries growth signals inside cells; abnormal activity can allow some cancers to grow or spread.

### **Caterpillar Dermatitis**

An itchy or painful skin reaction after contact with irritating or venomous caterpillar hairs or spines. Washing the area and removing visible hairs may help; eye exposure, breathing difficulty,

or widespread symptoms need medical care.

### **Caterpillars**

The larval stage of butterflies and moths. Hairs or spines on some species can cause painful skin inflammation, eye injury, or allergic reactions after contact or inhalation.

### **Cathartic**

A substance that promotes bowel emptying, usually by increasing stool water, intestinal movement, or both. Excessive or prolonged use can cause diarrhea, dehydration, electrolyte imbalance, or dependence on laxatives.

#### **Alternate Names:**

Purgative; Laxative

### **Cathepsin**

A family of enzymes that break down proteins inside cells, especially in lysosomes, the cell's recycling compartments. They help with immune activity, tissue remodeling, and bone turnover; abnormal activity may contribute to inflammation, bone loss, or cancer.

### **Cathepsin D**

An enzyme in lysosomes that breaks down proteins in an acidic environment. It also participates in cell death and immune processes; abnormal activity has been studied in cancer and nervous-system disease.

### **Cathepsin H**

An enzyme in lysosomes that breaks down proteins and helps activate some biologically active peptides. It contributes to normal cell maintenance and may be altered in certain diseases.

### **Cathepsin K**

An enzyme made by osteoclasts, the cells that remove old bone. It breaks down the protein framework of bone; excessive activity can contribute to bone loss and osteoporosis.

### **Cathepsin L**

An enzyme in lysosomes that breaks down proteins

---

and helps process antigens for immune recognition. It also participates in tissue remodeling and can assist entry or spread of some infectious agents.

### **Cathepsin Z**

An enzyme found mainly in lysosomes that cuts proteins into smaller pieces. It helps cells process materials and may contribute to immune responses and inflammation.

### **Cathepsins B**

A group of protein-cutting enzymes, including cathepsin B, that help recycle proteins inside cells. Abnormal activity can contribute to inflammation, tissue remodeling, invasion of nearby tissue by tumors, and spread of cancer.

### **Catheter**

A thin flexible tube placed in a blood vessel, body cavity, or duct to deliver medicines or fluids, drain urine or other material, measure pressure, collect samples, or guide a procedure.

### **Catheter-Associated Urinary Tract Infection**

A hospital-acquired infection of the urinary tract occurring in a patient with an indwelling urinary catheter within the prior 48 hours, characterized by bacteriuria, pyuria, fever, or suprapubic tenderness.

#### **Alternate Names:**

CAUTI

### **Catheter-Associated UTI**

A urinary infection associated with an indwelling catheter. Risk rises with prolonged use, and symptoms may include fever, lower abdominal or flank pain, or confusion; avoiding unnecessary catheters and removing them early lowers risk.

### **Catheter-Directed Thrombolysis**

A minimally invasive image-guided procedure used to rapidly dissolve dangerous blood clots in major arteries or veins. Under real-time X-ray (fluoroscopy), an interventional radiologist or vascular surgeon threads a specialized catheter through a small skin puncture in the groin or arm directly into

the blood clot (thrombus). Clot-dissolving medication (such as tissue plasminogen activator [tPA]) is infused directly into the clot over several hours, often augmented by ultrasonic vibrations or mechanical aspiration to break up the blockage. Because the medicine is concentrated directly inside the clot, much lower doses are required compared to standard intravenous therapy, significantly reducing the risk of bleeding in other organs while effectively treating severe deep vein thrombosis (DVT), massive pulmonary embolism, and acute limb ischemia.

#### **Alternate Names:**

CDT; Endovascular Thrombolysis; Catheter-Directed Clot Lysis

### **Catheter-Related Infections**

Infections caused by microorganisms entering along or contaminating an indwelling catheter. They may cause local redness, pain, drainage, fever, or bloodstream infection; treatment may require antibiotics and catheter removal or replacement.

### **Catheter-Related UTI**

A urinary tract infection associated with an indwelling urinary catheter, usually caused by bacteria entering or traveling along it. Diagnosis requires compatible symptoms and appropriately collected urine testing because bacteria may be present without infection.

### **Catheterization**

Placement of a catheter into a blood vessel, body cavity, duct, or other passage to drain, infuse, measure, collect samples, or perform a procedure. Risks include pain, bleeding, infection, blockage, and injury to nearby tissue.

### **Catheterized Urine Culture**

A urine culture performed on a specimen collected through urinary catheterization, usually when a clean-catch sample cannot be obtained or when catheter-associated infection is being assessed. Collection technique and timing affect interpretation because colonization and contamination may cause bacterial growth without symptomatic infection.

---

## Cathexis

The investment of emotional or psychological energy in a person, idea, object, or activity, a term used mainly in psychology and psychoanalytic theory.

## Cation

An atom or molecule with a positive electrical charge, such as sodium, potassium, calcium, or hydrogen. These charged particles help control fluid balance, nerve signals, muscle contraction, and blood acidity.

### Alternate Names:

Positively Charged Ion

## Cauda Equina

The bundle of lumbar, sacral, and coccygeal nerve roots that continues below the end of the spinal cord. These nerves control sensation and movement in the legs and help control the bladder, bowel, and sexual organs.

### Alternate Names:

Cauda Equina Nerve Roots; Horse-Tail Nerve Bundle

## Cauda Equina Syndrome

An emergency caused by severe pressure on the bundle of nerves at the bottom of the spine, often from a large slipped disc. Warning signs include numbness around the groin or buttocks, difficulty urinating, loss of bladder or bowel control, severe leg pain, or leg weakness. Urgent assessment and MRI are needed; some people need emergency surgery to reduce the risk of lasting nerve damage.

### Alternate Names:

CES; Cauda Equina Compression; Spinal Nerve Root Compression

## Cauda Equina Syndrome Cross-Reference

Alternate name; see *Cauda equina syndrome*.

## Caudal

Toward the tail or lower end of the body; in human anatomy, generally toward the feet or away from the head.

## Caudal

Positioned toward the tail or lower part of the body; in human anatomy, usually toward the feet or away from the head.

## Caudal Dysplasia

A group of birth defects in which the lower spine and nearby structures do not develop fully. It may cause abnormalities of the legs, pelvis, bowel, bladder, or genital organs and is more common with maternal diabetes.

## Caudate Nucleus

A curved group of nerve cells deep in the brain that helps control movement, learning, memory, motivation, and behavior. Disease affecting it may contribute to movement or cognitive problems.

## Cauliflower Ear

A thick, irregular deformity of the outer ear caused when blood or fluid collects between its cartilage and covering, usually after blunt injury or repeated friction. Prompt drainage and compression can protect the cartilage and prevent permanent change.

### Alternate Names:

Auricular Hematoma Deformity; Boxer's Ear; Wrestler's Ear

## Caulking Compound Poisoning

Poisoning or injury after contact with or swallowing a caulking product. Symptoms depend on its ingredients and may include eye, skin, or airway irritation, vomiting, drowsiness, or aspiration into the lungs.

## Causalgia

Severe persistent burning pain, sensitivity, and other sensory changes after injury to a peripheral nerve. It is now usually considered complex regional pain syndrome type II and may include swelling, color or temperature changes, and movement difficulty.

### Alternate Names:

Complex Regional Pain Syndrome Type II; Causalgic Pain

---

## Cause Of Death

The disease, injury, or toxic exposure that starts the chain of events leading directly to death. It differs from the immediate mechanism, such as respiratory failure, and from contributing conditions.

## Causes of Cancer

Cancer develops after changes in cells allow abnormal growth and survival. Risk can be increased by inherited variants, tobacco, alcohol, certain infections, ultraviolet or other radiation, harmful chemicals, chronic inflammation, and random changes during cell division.

### Alternate Names:

Carcinogenesis; Cancer Etiology; Oncogenic Factors

## Caustic

Able to chemically burn or destroy living tissue, as strong acids and alkalis can. Swallowing or contact with a caustic substance can injure the mouth, esophagus, stomach, skin, eyes, or airway and may be an emergency.

## Cauterization

Intentional destruction or sealing of tissue with heat, electricity, chemicals, or another energy source. It may control bleeding, remove abnormal tissue, or close small vessels and wounds.

## Cautery

Use of heat, electricity, chemicals, or another energy source to destroy or seal tissue, commonly to control bleeding or remove a small abnormal area.

## Cautery Device

A medical instrument that uses heat, electricity, or another energy source to seal bleeding vessels or destroy unwanted tissue. Use can cause burns or damage surrounding structures if poorly controlled.

## Caveola

A small pocket-like fold in a cell membrane that helps organize signals, move materials into or out of the cell, and respond to stretching or mechanical stress.

## Caveolins

Proteins that form or regulate caveolae, small pockets in cell membranes. They help organize cell signals, handle fats, move materials across membranes, and respond to mechanical stress.

## Cavernous Hemangioma

A cluster of enlarged, thin-walled blood vessels that may occur in the skin, brain, spinal cord, or other tissues. Symptoms depend on location and may result from bleeding, pressure, pain, or altered function.

## Cavernous Malformations

Clusters of unusually wide, fragile blood vessels, usually in the brain or spinal cord. They may cause no symptoms, but bleeding or pressure on nerve tissue can produce headaches, seizures, weakness, numbness, vision problems, or other neurologic changes.

### Alternate Names:

Cavernoma; Cerebral Cavernoma

## Cavernous Sinus

A large venous channel beside the pituitary gland that drains blood from the eye and face. It contains important eye-movement nerves and the internal carotid artery; infection or clotting there can be life-threatening.

## Cavernous Sinus Thrombosis

A dangerous blood clot, often infected, in the venous channel behind the eyes. It usually follows infection of the face, nose, sinuses, teeth, or orbit and may cause severe headache, fever, bulging eyes, double vision, or vision loss.

## Cavities

Holes or damaged areas in teeth caused by acids made by plaque bacteria from sugars and other carbohydrates. Decay can reach the inner pulp and cause pain or infection; prevention and treatment may include fluoride, cleaning, fillings, or root-canal care.

### Alternate Names:

Tooth Decay; Dental Caries; Dental Cavitation



---

## Cavities And Tooth Decay

Progressive loss of tooth structure caused by acids produced when plaque bacteria use sugars. Decay can move from enamel into dentin and the pulp, causing sensitivity, pain, infection, abscess, or tooth loss.

## Cavity

Alternate name; see *Cavities*.

## Cavum

An anatomical cavity or hollow space in the body. The word is usually followed by a location or structure name; for example, the cavum septi pellucidi is a normal fluid-filled space in the brain.

## CBC

See *Complete Blood Count*.

### Alternate Names:

Complete Blood Count; Full Blood Count

## CBC Blood Test

See *Complete Blood Count*.

## CCP

Cyclic citrullinated peptide, a target used in a blood test for antibodies associated with rheumatoid arthritis. A positive anti-CCP result supports the diagnosis and may indicate greater risk of joint damage, but it is not diagnostic alone.

## CCTA

Alternate name; see *Coronary Computed Tomography Angiography*.

## CCU

A hospital unit providing continuous heart monitoring and specialized treatment for people with serious acute cardiac problems, such as heart attack, dangerous rhythm disturbance, or severe heart failure.

### Alternate Names:

Coronary Care Unit; Cardiac Care Unit

## Cd Antigens

Proteins or other markers on the surface of cells that help identify cell types, especially immune cells. Laboratory antibodies against these markers can classify cells, diagnose blood disorders, and guide selected treatments.

## CD4 Count

The number of CD4 T lymphocytes in a blood sample. It helps assess immune function and the risk of opportunistic infection in people with HIV; results are interpreted with viral load, symptoms, treatment, and trends over time.

## CD4-Positive T Cell

A type of white blood cell that coordinates immune responses. Different CD4 cells help activate other immune cells, support antibody production, defend mucosal surfaces, or limit excessive inflammation.

## CD8-Positive T Cell

A type of white blood cell that recognizes abnormal signals on infected or altered cells. Cytotoxic CD8 cells can destroy virus-infected, cancerous, or otherwise abnormal cells.

## Cea

An abbreviation for carcinoembryonic antigen, a blood marker used mainly to monitor selected cancers rather than to screen people without symptoms.

## CEA Assay

A laboratory test measuring carcinoembryonic antigen in a specimen, usually blood. It is mainly used to monitor selected known cancers and cannot diagnose or exclude cancer by itself.

## CEA Blood Test

A blood test measuring carcinoembryonic antigen, used mainly to monitor selected cancers such as colorectal cancer and assess treatment response or recurrence. Smoking, inflammation, and other non-cancerous conditions can also raise it.

---

### Cecal Hemorrhage

Bleeding from the caecum, the first part of the large intestine. Causes include inflammation, infection, reduced blood flow, polyps, diverticular disease, or cancer and may produce blood in the stool or anemia.

### Cecal Infection

Infection or inflammation of the caecum caused by microorganisms, reduced blood flow, inflammatory bowel disease, or nearby disease. It may cause right-sided abdominal pain, fever, diarrhea, or tenderness.

### Cecal Neoplasm

A noncancerous or cancerous growth arising in the caecum. It may cause bleeding, anemia, abdominal pain, altered bowel habits, or blockage and is usually evaluated with colonoscopy and tissue sampling.

### Cecal Volvulus

Twisting of the caecum and nearby colon around the tissue that supports them, causing large-bowel blockage. It can produce sudden abdominal pain, swelling, vomiting, and inability to pass stool or gas and usually requires urgent treatment.

### Cecostomy

A surgically created opening from the caecum to the abdominal wall to release pressure, drain bowel contents, or divert stool. It may be temporary or permanent depending on the underlying problem.

### Cecum

The first pouch-like part of the large intestine, in the lower right abdomen where the small intestine joins the colon. The appendix arises from it; disease can include inflammation, tumors, bleeding, or twisting that blocks the bowel.

### Cedar Leaf Oil Poisoning

Toxic exposure to cedar-leaf oil, usually after swallowing or inhaling it. It may irritate the mouth, stomach, lungs, or nervous system and can cause vomiting, dizziness, seizures, or breathing problems; urgent poison-control advice is needed.

### Cefaclor

A second-generation cephalosporin antibiotic used for selected susceptible bacterial infections. It can cause allergic reactions, nausea, diarrhea, or antibiotic-associated colitis and should be used only when appropriate.

### Cefadroxil

An oral first-generation cephalosporin antibiotic used for selected susceptible skin, throat, urinary, and other infections. Possible effects include allergy, nausea, diarrhea, and antibiotic-associated colitis.

### Cefamandole

A second-generation cephalosporin antibiotic used in some settings for susceptible bacterial infections. It can cause allergy or diarrhea and may increase bleeding risk, especially in people with poor nutrition or liver disease.

### Cefatrizine

An older oral cephalosporin antibiotic used in some countries for susceptible bacterial infections. Its usefulness depends on local availability, bacterial resistance, allergy risk, and current treatment guidance.

### Cefazolin

An injectable first-generation cephalosporin antibiotic used for selected susceptible infections and commonly given before surgery to prevent infection. Allergy, diarrhea, and dose adjustment in kidney disease may be important.

### Cefdinir

An oral third-generation cephalosporin antibiotic used for selected susceptible respiratory, skin, and other infections. It may cause diarrhea, nausea, allergic reactions, or antibiotic-associated colitis and can interact with iron-containing products.

### Cefepime

An injectable fourth-generation cephalosporin used for serious infections caused by susceptible bacteria, including some hospital-acquired infections. Kidney-dose adjustment is important because accumulation

---

can cause confusion, seizures, or other neurologic toxicity.

### **Cefixime**

An oral third-generation cephalosporin used for selected susceptible urinary, respiratory, ear, and other bacterial infections. It can cause diarrhea or allergy, and unnecessary use promotes antibiotic resistance.

### **Cefmenoxime**

An injectable third-generation cephalosporin used in some countries for susceptible bacterial infections. Possible effects include allergic reactions, diarrhea, and antibiotic-associated colitis.

### **Cefmetazole**

An injectable cephamycin antibiotic used in some countries for selected abdominal, pelvic, and other infections caused by susceptible bacteria. Allergy, diarrhea, and antibiotic-associated colitis are possible.

### **Cefonicid**

A long-acting second-generation cephalosporin antibiotic used for selected susceptible infections. Its use is limited in many places; dose selection depends on kidney function, local resistance, and the infection site.

### **Cefoperazone**

A third-generation cephalosporin active against many gram-negative bacteria and sometimes combined with a beta-lactamase inhibitor. It can cause allergy, diarrhea, bleeding, or impaired bile flow and requires appropriate monitoring.

### **Cefotaxime**

An injectable third-generation cephalosporin used for serious susceptible infections, including meningitis, pneumonia, bloodstream infection, and sepsis. Allergy, diarrhea, and antibiotic-associated colitis are possible.

### **Cefotetan**

An injectable cephamycin antibiotic used for se-

lected serious infections and surgical prevention. It covers some anaerobic bacteria but can cause allergy, diarrhea, bleeding, or a reaction if alcohol is consumed around treatment.

### **Cefotiam**

A second-generation cephalosporin used in some countries for susceptible respiratory, urinary, skin, and other bacterial infections. Allergy, gastrointestinal effects, and antibiotic-associated colitis are possible.

### **Cefoxitin**

An injectable cephamycin antibiotic used for selected abdominal, pelvic, skin, and other infections, including some involving anaerobic bacteria. Allergy, diarrhea, and antibiotic-associated colitis can occur.

### **Ceftazidime**

An injectable third-generation cephalosporin with activity against *Pseudomonas aeruginosa* and other susceptible gram-negative bacteria. It is used for selected serious infections and can cause allergy, diarrhea, or neurologic effects when it accumulates in kidney disease.

### **Ceftazidime-Avibactam**

A combination antibiotic in which avibactam blocks some bacterial enzymes that destroy ceftazidime. It is used for selected difficult gram-negative infections, including some resistant organisms; kidney-dose adjustment and susceptibility testing are important.

### **Ceftibuten**

An oral cephalosporin antibiotic used for selected susceptible respiratory, urinary, and other bacterial infections. It may cause diarrhea or allergy, and inappropriate use promotes resistance and antibiotic-associated colitis.

### **Ceftizoxime**

An injectable third-generation cephalosporin used for selected serious infections caused by susceptible bacteria, including some abdominal, pelvic, urinary, and anaerobic infections. Allergy, diarrhea, and

---

antibiotic-associated colitis are possible.

### **Ceftriaxone**

An injectable third-generation cephalosporin used for many serious susceptible bacterial infections, including meningitis, pneumonia, bloodstream infection, and gonorrhea. Allergy, diarrhea, biliary problems, and antibiotic-associated colitis are possible.

### **Cefuroxime**

A second-generation cephalosporin available by mouth or injection for selected respiratory, ear, urinary, skin, and other bacterial infections. It can cause allergy, nausea, diarrhea, or antibiotic-associated colitis.

### **Celebrex**

A brand name for celecoxib, a selective cyclooxygenase-2 nonsteroidal anti-inflammatory drug used for pain and inflammation. It can cause stomach, kidney, cardiovascular, allergic, or bleeding complications and should be used as directed.

### **Celiac Artery**

A major branch of the abdominal aorta that supplies the upper digestive organs, including the stomach, liver, gallbladder, spleen, pancreas, and first part of the small intestine. It divides into the left gastric, splenic, and common hepatic arteries.

### **Celiac Crisis**

A rare, life-threatening severe flare of celiac disease causing profuse diarrhea or vomiting, dehydration, major electrolyte disturbances, and sometimes shock. It requires urgent hospital treatment and strict control of gluten exposure.

### **Celiac Disease**

A chronic autoimmune disease in which eating gluten, a protein found in wheat, barley, and rye, causes the immune system to damage the lining of the small intestine. The damage can flatten the intestine's tiny absorbing structures, called villi, and reduce the absorption of nutrients. Symptoms may

include diarrhea, constipation, bloating, abdominal pain, nausea, weight loss, tiredness, anemia, or poor growth in children. Some people have no digestive symptoms but may develop weak bones, nerve symptoms, or an itchy blistering rash called dermatitis herpetiformis. Celiac disease is different from wheat allergy and non-celiac gluten sensitivity, which does not damage the small intestine.

#### **Alternate Names:**

Nontropical Sprue

### **Celiac Plexus**

A network of nerves in the upper abdomen that carries signals to and from the stomach, liver, pancreas, intestines, and other organs. It can be targeted with an injection for selected severe abdominal pain.

### **Celiac Sprue**

An autoimmune enteropathy triggered by the ingestion of dietary gluten in genetically predisposed individuals expressing HLA-DQ2 or HLA-DQ8, leading to chronic villous atrophy, crypt hyperplasia, and malabsorption.

#### **Alternate Names:**

Celiac Disease; Gluten-Sensitive Enteropathy; Nontropical Sprue

### **Celiac Trunk**

The first major branch of the abdominal aorta, supplying the upper digestive organs. It divides into the left gastric, splenic, and common hepatic arteries and is surrounded by nerves of the celiac plexus.

### **Celiac Trunk Stenosis**

Narrowing of the celiac trunk, caused by plaque or, in some people, compression by the diaphragm's median arcuate ligament. It may cause upper abdominal pain after meals and weight loss, although imaging findings do not always explain symptoms.

### **Cell**

The smallest living unit that can perform life processes. Human cells usually contain DNA in a nucleus and specialized internal structures that produce energy, make proteins, communicate, and maintain the body's tissues.

---

## Cell Adhesion

The attachment of cells to one another or to the supporting material around them. It helps maintain tissue structure, transmit signals, guide movement, and organize healing and development.

## Cell Aggregation

Clumping of cells through interactions between their surfaces or surrounding proteins. It is important in development, immune responses, wound healing, and formation of blood clots.

## Cell Body

The main part of a cell containing the nucleus and most internal structures. In a nerve cell, it excludes long extensions such as the axon and dendrites.

## Cell Count

The number of cells in a sample of blood, body fluid, tissue, or laboratory culture. It helps assess blood-cell levels, inflammation, infection, immune function, or abnormal cell growth.

## Cell Culture

Growth of cells outside the body in a controlled nutrient environment. It is used for research, medicine testing, diagnosis, production of biological products, and study of how cells respond to disease or treatment.

## Cell Cycle

The series of stages through which a cell grows, copies its DNA, checks for damage, and divides into two cells. Disruption of cycle control can contribute to developmental disorders and cancer.

## Cell Death

The loss of a cell's ability to remain alive. It may be a controlled process that removes old or damaged cells, or an uncontrolled result of injury, infection, toxins, lack of oxygen, or severe disease.

## Cell Dedifferentiation

Loss of some specialized features by a mature cell, making it more like an immature cell. It can occur

during repair or development and may contribute to abnormal growth in some cancers.

## Cell Density

The number of cells in a given area or volume of tissue or fluid. It can change with growth, inflammation, healing, infection, or cancer and is measured in laboratory samples and tissue studies.

## Cell Differentiation

The process by which an unspecialized cell develops the structure and function of a specialized cell type, such as a nerve, muscle, blood, or skin cell.

## Cell Division

The process by which one cell produces new cells. Mitosis makes most body cells for growth and repair, while meiosis makes reproductive cells with half the usual number of chromosomes.

## Cell Fusion

Joining of two cells or their outer membranes to form one larger cell, sometimes with several nuclei. It occurs in normal development and repair and can also occur during infection or disease.

## Cell Growth

Increase in a cell's size and contents through production of proteins, membranes, energy stores, and other components. Abnormal growth signals can contribute to tissue overgrowth or cancer.

## Cell Maturation

Development of an immature cell into a mature cell with a specialized structure and function. It is essential for producing healthy blood, nerve, muscle, immune, and other body cells.

## Cell Migration

Movement of cells through tissue or across a surface in response to chemical, mechanical, or environmental signals. It is important in development, immune defense, wound healing, and cancer spread.

---

## Cell Motility

The ability of cells to move through or across a surface using their internal framework, attachment structures, or specialized projections such as cilia and flagella. It supports development, immunity, wound healing, and cancer spread.

### Alternate Names:

Cell Locomotion

## Cell Movement

Directed or random movement of a cell through tissue or across a surface. It is important in development, immune responses, wound repair, inflammation, and the spread of cancer.

## Cell Nucleus

The membrane-covered compartment containing most of a cell's DNA. It helps control gene activity, growth, division, repair, and production of proteins needed for normal cell function.

## Cell Pellet

The concentrated layer of cells collected at the bottom of a tube after spinning a sample in a centrifuge. It may be removed for microscopic, genetic, chemical, or other laboratory testing.

## Cell Physiology

The study of how cells obtain energy, transport substances, communicate, respond to signals, repair themselves, and perform specialized functions.

## Cell Polarity

The unequal arrangement of structures and activities within a cell, giving it distinct sides or directions. It enables organized transport, signaling, movement, development, and tissue structure.

## Cell Proliferation

Increase in cell number through repeated division. Normal proliferation supports growth and repair, while excessive or uncontrolled proliferation can cause tissue overgrowth, benign tumors, or cancer.

## Cell Secretion

Release of substances made by cells, such as hormones, enzymes, nerve chemicals, mucus, or immune signals. Secretion supports communication, digestion, protection, movement, and regulation of body processes.

## Cell Senescence

A state in which a cell remains active but permanently stops dividing, often because of aging, stress, or DNA damage. Senescent cells can help prevent cancer but may contribute to inflammation when they accumulate.

### Alternate Names:

Cellular Senescence; Replicative Senescence

## Cell Signaling

The process by which cells detect information through receptors and transmit signals inside the cell to change functions such as growth, movement, metabolism, or immune activity.

## Cell Sorting

Separation of cells into groups according to features such as size, density, surface markers, or fluorescence. It is used in research, diagnosis, blood-cell analysis, and preparation of selected cells for treatment.

## Cell Structure

The arrangement of a cell's membrane, cytoplasm, nucleus, and internal structures. These parts work together to produce energy, make proteins, communicate, divide, and carry out specialized functions.

## Cell Surface

The outer membrane of a cell and the molecules attached to it. It controls movement of substances, attachment, recognition, communication, and interactions with immune cells and surrounding tissue.

## Cell Survival

The ability of a cell to remain alive and functional despite injury, lack of nutrients, toxins, infection, or other stress. It depends on the cell type, the severity and duration of the stress, and the cell's



---

repair systems.

### Cell Therapy

Treatment that uses living cells to replace damaged cells, restore a function, or change an immune response. Uses range from established blood-forming stem-cell transplantation to experimental treatments; benefits and risks depend on the cells and disease.

### Cell Transplantation

Transfer of living cells into a person to replace missing cells, restore tissue function, or modify immunity. Blood-forming stem-cell transplantation is established for selected diseases; rejection, infection, graft-versus-host disease, and other risks may occur.

### Cell Wall

A rigid outer layer surrounding plant, fungal, and bacterial cells. It provides shape and protection; human and animal cells have a flexible membrane but no cell wall.

### Cell-Cell Junction

A specialized connection between neighboring cells that provides attachment, communication, or controlled passage of substances. Examples include tight junctions, desmosomes, and gap junctions.

### Cell-Cell Signaling

The ways cells send and receive messages through direct contact, electrical activity, or chemicals such as hormones and neurotransmitters. Signaling coordinates growth, movement, immune responses, repair, and other body functions.

### Cell-Free DNA

Non-encapsulated extracellular DNA fragments circulating freely in blood plasma or other biofluids, released from apoptotic or necrotic cells; analyzed in non-invasive prenatal screening (NIPT) and oncology liquid biopsies.

**Alternate Names:**

cfDNA; Circulating Cell-Free DNA

### Cell-Free Fetal DNA Screening

A prenatal blood test that analyzes small DNA fragments from the placenta to estimate the chance of selected chromosome conditions. It is a screening, not a diagnostic, test; a high-risk or unclear result needs counseling and confirmatory testing.

### Cell-Matrix Junction

A specialized connection anchoring a cell to the supporting material around it. It also transmits signals that influence cell shape, movement, survival, and tissue organization.

### Cell-Mediated Immune Response

Alternate name; see *Cell-Mediated Immunity*.

### Cell-Mediated Immunity

Immune defense carried out mainly by T lymphocytes and other cells rather than by antibodies. It is important for controlling infected cells, cancer cells, and organisms that live inside cells.

**Alternate Names:**

Cell-Mediated Immune Response

### Cellophane

A thin transparent film made from cellulose. In medicine it may be used in some dressings, packaging, or laboratory techniques; its clinical meaning depends on the context.

### Cellular

Relating to cells or to processes occurring within or between cells.

### Cellular Adaptation

A reversible change in a cell's size, number, structure, or function in response to altered demands or mild injury. Adaptation may help a cell survive but can become harmful if the stress continues.

### Cellular Immunity

Immune defense carried out mainly by T cells, natural killer cells, macrophages, and other cells rather than antibodies alone. It helps destroy infected or abnormal cells and control inflammation.

---

### Cellular Metabolic Process

A chemical reaction or linked pathway in a cell that produces energy, builds cell components, or breaks down nutrients and waste.

### Cellular Quiescence

A reversible, non-dividing physiological state in which a cell exits the active cell division cycle and enters the G0 phase in response to anti-mitogenic signals or nutrient deprivation, while retaining the capacity to re-enter the proliferative cycle upon appropriate growth factor stimulation.

### Cellular Respiration

The process cells use to release energy from food. Most cells use oxygen to convert nutrients into ATP, the cell's usable energy, while limited energy can be made without oxygen.

### Cellular Swelling

Early reversible cell injury in which failure of membrane pumps causes sodium and water to collect inside the cell. It commonly follows reduced blood flow, low oxygen, toxins, or other stresses.

#### Alternate Names:

Hydropic Change; Hydropic Degeneration; Cloudy Swelling

### Cellular Vesicle

A small membrane-bound compartment that transports, stores, or releases substances inside or outside a cell. Vesicles help move proteins, fats, hormones, and waste.

### Cellulite

A common harmless dimpled or uneven appearance of skin, usually on the thighs, hips, or buttocks, caused by the arrangement of fat, connective tissue, and skin. It is different from cellulitis, a bacterial infection.

### Cellulitis

A bacterial infection of the deeper skin and tissue beneath it, causing spreading redness, warmth, swelling, tenderness, and sometimes fever or chills.

It commonly follows a break in the skin and can spread rapidly or enter the bloodstream.

### Cellulomonas

A group of bacteria found mainly in soil, plants, and other environmental materials. They break down cellulose; human infection is rare and usually occurs opportunistically in people with severe illness or weakened immunity.

### Cellulomonas Hominis

A rare environmental bacterium that has occasionally caused human infection, usually in people with weakened immunity or serious underlying illness. Its significance depends on symptoms, the body site, and laboratory identification.

### Cellulose

A complex carbohydrate forming the structure of plant cell walls. Humans cannot digest it, but it is dietary fiber that adds bulk to stool and can support regular bowel movements when enough fluid is consumed.

### Cellulosimicrobium

A group of environmental bacteria found mainly in soil. Human infection is rare but may affect people with serious illness, implanted devices, or weakened immune defenses and can involve the blood, heart, or other tissues.

### Cellulosome

A group of enzymes joined together by some microorganisms to break down cellulose in plant cell walls. It helps the organism digest plant material and is studied for biotechnology and biofuel production.

### Celosomy

A rare birth defect involving incomplete closure of the body wall, allowing abdominal or chest organs to protrude outside the body. The term is used for severe omphalocele-like malformations and requires specialist newborn care.

---

## **Cement**

A material used to fill spaces or attach an artificial joint or other prosthesis to bone. Dental cement bonds or protects dental work; the appropriate type depends on the procedure.

## **Cemento-Enamel Junction**

The boundary where tooth enamel covering the crown meets cementum covering the root. Its shape varies, and an exposed junction may be sensitive or more vulnerable to decay.

## **Cementoma**

A usually noncancerous jaw lesion containing cementum-like mineralized tissue. Its behavior and treatment depend on the specific subtype, location, size, symptoms, and relationship to nearby teeth.

## **Cementum**

A thin mineralized layer covering a tooth root. Fibers of the periodontal ligament attach to it and help anchor the tooth in its socket; it can be lost with gum disease.

## **Cenesthesia**

The general awareness of one's body, including its condition, position, and internal sensations. Disturbance may occur in some neurologic or psychiatric illnesses and can alter how bodily sensations are experienced.

## **Centenarian**

A person aged 100 years or older. Centenarians are a diverse group; some remain independent, while others need substantial support, and age alone does not determine health or cognitive ability.

## **Center**

A central point, region, or facility organized around a function. In medicine, the term may describe an anatomical location, a treatment facility, or the middle of a measured structure.

## **Centigram**

A metric unit of mass equal to one hundredth of a

gram, or 10 milligrams, used for small quantities of medicines or specimens.

## **Centimeter**

A metric unit of length equal to one hundredth of a meter, commonly used in clinical measurements such as wound size, tumor dimensions, and anatomical distances.

## **Centipede**

A many-legged arthropod whose bite can inject venom and cause local pain, swelling, and redness; severe allergic reactions are uncommon but possible.

## **Centipoise**

A unit of dynamic viscosity equal to one millipascal-second, used to describe the flow resistance of blood, fluids, medicines, and other materials.

## **Central**

Located near the middle of the body or an organ, rather than near its edge. The word may describe a body structure, nervous-system area, blood line, symptom, or pattern of disease.

## **Central Apnea**

A pause in breathing caused by absent or reduced respiratory effort from impaired brainstem signaling, rather than upper-airway obstruction.

## **Central Auditory Processing Disorder**

Difficulty interpreting sounds in the brain despite adequate hearing. People may struggle to understand speech in noise, follow rapid instructions, identify sounds, or distinguish similar words.

## **Central Cord Syndrome**

The most common form of incomplete traumatic spinal cord injury, typically resulting from hyperextension trauma in patients with preexisting cervical spondylosis. It is characterized by disproportionately greater motor weakness and impairment in the upper extremities compared to the lower extremities, accompanied by variable sensory loss and bladder dysfunction.

---

**Alternate Names:**

Cervical Central Cord Syndrome; Schneider Syndrome

**Central Core Disease**

An inherited muscle disorder, often caused by an RYR1 gene change, that may cause low muscle tone, delayed motor development, and weakness. Some affected people can develop life-threatening overheating with certain anesthetics.

**Alternate Names:**

CCD; Core Myopathy; RYR1-Related Central Core Disease

**Central Cyanosis**

A bluish or purplish discoloration of the skin, mucous membranes, tongue, and lips resulting from an abnormally elevated concentration of deoxygenated hemoglobin (greater than 5 g/dL) in systemic arterial blood, typically reflecting severe respiratory compromise or right-to-left cardiac shunting.

**Central Deafness**

Severe hearing loss caused by damage to sound-processing pathways or areas of the brain, rather than the outer, middle, or inner ear. Hearing tests may show poor understanding despite detected sounds.

**Central Diabetes Insipidus**

Excessive urination and intense thirst caused by too little antidiuretic hormone from the brain. The kidneys cannot conserve enough water, so dehydration and high blood sodium may develop.

**Central Fatigue**

A sense of reduced mental energy, concentration, alertness, or motivation arising primarily from central nervous-system processes.

**Alternate Names:**

Central Fatigue; Mental Fatigue

**Central Hypothyroidism**

A condition with too little thyroid hormone because the brain's pituitary gland or hypothalamus does

not provide enough stimulation to the thyroid.

**Central Line**

A catheter inserted into a large vein with its tip in or near the heart, used to give medications or nutrition, draw blood, or monitor central venous pressures.

**Central Line Placement**

Insertion of a catheter into a large vein, commonly in the neck, chest, or groin, with its tip near the heart. It allows long-term medicines, nutrition, fluids, blood sampling, or monitoring.

**Central Line-Associated Bloodstream Infection**

A primary nosocomial bloodstream infection occurring in a patient with a central vascular catheter in place for > 48 hours that is not related to an infection at another anatomic site.

**Alternate Names:**

CLABSI; Central Venous Catheter Infection; Catheter-Related Bloodstream Infection; CRBSI

**Central Nervous System**

The brain and spinal cord. It receives and interprets information, controls movement and many automatic body functions, and supports thought, memory, emotion, and regulation of hormones.

**Central Nervous System Germinoma**

A malignant germ-cell tumor usually arising in the brain, often near the pineal or suprasellar region. It may cause headache, visual or endocrine abnormalities, hydrocephalus, or impaired puberty; MRI, tumor markers, and specialist evaluation are important.

**Central Nervous System Lymphoma**

A lymphoma involving the brain, spinal cord, cerebrospinal fluid, or eye that can cause neurologic or visual symptoms.

**Central Nervous System Vascular**

---

## Malformations

Abnormal groups or connections of blood vessels in or around the brain or spinal cord. Some cause no symptoms, while others can cause headaches, seizures, weakness, or bleeding; treatment depends on the type, size, location, and risk.

## Central Neurofibromatosis

Alternate name; see *Neurofibromatosis Type 2*.

## Central Pain Syndrome

Chronic pain caused by damage or dysfunction in the brain or spinal cord. Pain may be burning, aching, electric, or unusually sensitive to touch and may occur after stroke, spinal-cord injury, multiple sclerosis, or other lesions.

## Central Pontine Myelinolysis

Brain injury caused most often by correcting very low blood sodium too quickly. Damage in the pons may cause weakness, trouble speaking or swallowing, confusion, reduced consciousness, or locked-in syndrome.

## Central Precocious Puberty

Central precocious puberty begins when the hypothalamus and pituitary activate the normal puberty pathway unusually early. It causes early secondary sexual characteristics and accelerated bone maturation; evaluation identifies causes and treatment may delay progression when appropriate.

## Central Retinal Artery Occlusion

A sudden blockage of the main artery supplying the retina, causing severe, usually painless loss of vision in one eye. It is an eye stroke requiring emergency assessment for related vascular disease.

## Central Retinal Vein Occlusion

Blockage of the vein that drains blood from the retina, causing sudden blurring or loss of vision, usually in one eye and typically less severe than a central arterial occlusion. It most often results from thrombosis linked to aging, hypertension, diabetes, open-angle glaucoma, or other vascular disease and may cause retinal hemorrhages, swelling, and macu-

lar edema. Vision may recover substantially or be permanently impaired, especially in ischemic forms; urgent eye assessment is recommended, and treatment targets underlying risk factors plus, in selected cases, intravitreal anti-VEGF therapy or laser to control macular edema.

## Central Serous Chorioretinopathy

A condition in which fluid collects beneath the retina, usually near the macula, causing blurred or distorted central vision. It may be linked to corticosteroids or stress and often needs eye-specialist follow-up.

## Central Serous Choroidopathy

A disorder in which fluid accumulates beneath the neurosensory retina, usually near the macula, causing blurred or distorted central vision. It is associated with corticosteroid exposure and stress-related factors; persistent or recurrent disease requires ophthalmic evaluation.

## Central Sleep Apnea

Repeated pauses in breathing during sleep because the brain temporarily stops sending signals to breathe, without physical airway blockage. It may occur with heart failure, high altitude, or opioid use.

### Alternate Names:

CSA; Cheyne-Stokes Respiration

## Central Sleep Apnea Syndromes

Repeated pauses or reductions in breathing during sleep caused by diminished respiratory drive rather than upper-airway blockage. Causes include heart failure, opioids, neurologic disease, high altitude, and treatment-emergent patterns; evaluation and therapy depend on the underlying mechanism.

## Central Sulcus

A prominent anatomical landmark on the lateral cerebral cortex separating the precentral gyrus of the frontal lobe (primary motor cortex) from the postcentral gyrus of the parietal lobe (primary somatosensory cortex).

### Alternate Names:

---

Rolandic Fissure; Sulcus of Rolando

### Central Venous Catheter

A thin tube ending in a large vein near the heart. It provides reliable access for prolonged or irritating treatments and monitoring but can cause infection, blood clots, bleeding, or injury.

### Central Venous Catheter Dressing Change

A standardized sterile nursing procedure involving the aseptic removal of existing catheter dressings, chlorhexidine-based antiseptics of the percutaneous exit site, and application of a sterile transparent semipermeable occlusive dressing.

#### Alternate Names:

CVC Dressing Change; Central Line Dressing Care; Catheter Site Dressing

### Central Venous Catheter Flushing

The pulsatile instillation of sterile preservative-free 0.9% sodium chloride or heparinized lock solution through a central venous catheter lumen using positive pressure to prevent intraluminal thrombus formation, fibrin buildup, and catheter occlusion.

#### Alternate Names:

CVC Flushing; Central Line Heparin Lock; Catheter Patency Flushing

### Central Venous Line

A vascular catheter whose tip lies in a major central vein, commonly the superior vena cava, for reliable long-term or high-concentration intravenous access.

### Central Venous Pressure

Pressure measured in or near the right atrium, usually through a central venous catheter. It reflects right-sided filling conditions but is affected by ventilation, heart function, vascular tone, and measurement technique; it should not be used alone to guide fluid treatment.

### Central Vertigo

Dizziness or an illusion of motion caused by dysfunction in the brainstem, cerebellum, or central

vestibular pathways. It may occur with neurologic signs, severe imbalance, or less position-dependent symptoms.

### Central Vision

Detailed vision directed at the point of gaze, provided primarily by the macula and fovea; it is needed for reading and recognizing faces.

### Centriacinar Emphysema

A form of emphysema in which damage begins in the small airways and nearby air sacs, usually in the upper lungs. It is strongly linked to smoking and can cause cough, wheezing, breathlessness, and reduced exercise ability.

### Centric-Fusion Translocation

A Robertsonian translocation in which the long arms of two acrocentric chromosomes fuse near their centromeres, leaving a combined chromosome and usually losing the short arms.

### Centrifugation

Spinning a specimen at high speed to separate components by density, such as cells from plasma or precipitated material from a liquid.

### Centrifuge

A machine that spins samples rapidly to separate their parts by density. Laboratories use it to separate blood cells from plasma, concentrate urine material, or prepare tissue samples for testing; speed is recorded in revolutions or relative force.

### Centrilobular Emphysema

A type of permanent lung damage in which the small airways and nearby air sacs are destroyed, usually from long-term smoking or inhaled irritants. It often affects the upper lungs and can cause breathlessness, cough, wheezing, and reduced ability to exercise.

#### Alternate Names:

Centriacinar Emphysema; Proximal Acinar Emphysema



---

### Centriole

A cylindrical microtubule structure in the centrosome that helps organize the mitotic spindle and, in some cells, forms the basal body of a cilium or flagellum.

### Centromere

The constricted chromosome region where sister chromatids remain attached and the kinetochore forms for accurate chromosome segregation during cell division.

### Centronuclear Myopathy

A group of inherited muscle diseases in which muscle fibers develop abnormally. Symptoms range from severe weakness and breathing problems at birth to milder adult-onset weakness, often affecting the shoulders, hips, eyelids, eye movements, face, or feet.

#### Alternate Names:

CNM; Myotubular Myopathy (General)

### Centrosome

The main microtubule-organizing center of an animal cell, containing a pair of centrioles and coordinating the spindle during mitosis.

### Centrum

The main body of a vertebra, located in front of the spinal canal. It bears much of the body's weight and joins neighboring vertebrae through shock-absorbing discs.

### Cephacetrile

An older first-generation cephalosporin antibiotic that inhibits bacterial cell-wall synthesis; it is not commonly used in current practice.

### Cephalalgia

Headache or pain in the head, including pain arising from cranial or related structures. It is a symptom with many possible primary and secondary causes.

### Cephalexin

An oral first-generation cephalosporin that inhibits

bacterial cell-wall synthesis and treats selected skin, respiratory, urinary, and other susceptible infections.

### Cephalgia

Alternate spelling; see *Headache*.

### Cephalhaematoma

A collection of blood beneath the membrane covering a newborn's skull bone, usually caused by birth-related pressure or injury. It is limited by skull sutures and usually resolves without treatment.

### Cephalhematoma

A traumatic subperiosteal collection of blood over a neonatal cranial bone, commonly the parietal bone, resulting from rupture of periosteal blood vessels during delivery; clinically distinguished from caput succedaneum by remaining strictly bounded by cranial suture lines.

### Cephalic

Relating to the head or toward the head; in obstetrics, a cephalic presentation means the fetus presents head first.

### Cephalic Vein

A superficial vein on the lateral side of the upper limb that drains the hand and forearm and joins the axillary vein near the shoulder.

### Cephalohematoma

A collection of blood trapped under the outer layer of a newborn's skull bone, usually caused by pressure during a long labor or delivery with vacuum suction or forceps. Because the blood is held tightly to a single bone, the swelling stays on one side of the head and does not cross the skull's bone seams (sutures). It is usually harmless and shrinks away on its own within a few weeks, though the breaking down of trapped blood can sometimes cause mild newborn jaundice.

#### Alternate Names:

Cephalhematoma; Cephalohaematoma.

---

## Cephalometry

Measurement of the skull and facial structures, usually on standardized radiographs, used in orthodontic planning and assessment of craniofacial growth.

## Cephalopelvic Disproportion

A situation in which the fetal head cannot pass easily through the mother's pelvis because of their relative size or shape. It may prevent vaginal delivery, but cannot always be confirmed before labor.

## Cephaloridine

An older injectable first-generation cephalosporin antibiotic whose use declined because of nephrotoxicity and the availability of safer alternatives.

## Cephalosporin

A beta-lactam antibiotic that kills susceptible bacteria by blocking construction of their protective cell wall. Different generations act against different bacteria; allergy, diarrhea, and resistance are possible.

## Cephalosporins

Cephalosporins are beta-lactam antibiotics that inhibit bacterial cell-wall synthesis; generations differ in spectrum, and allergy, diarrhea, and resistance are important considerations.

## Cephalothin

An older injectable first-generation cephalosporin that inhibits bacterial cell-wall synthesis and was used for susceptible serious infections.

## Cephapirin

A first-generation cephalosporin antibiotic formerly used for susceptible bacterial infections; current availability and indications vary by country.

## Cephradine

A first-generation cephalosporin antibiotic related to cephalexin that inhibits bacterial cell-wall synthesis and treats selected susceptible infections.

## Ceramide

A type of fat that forms part of cell membranes and helps the outer skin retain water and block irritants. Ceramides also participate in signals controlling cell growth and survival.

## Cercaria

A cercaria is a free-swimming larval stage of a trematode parasite; contact with some species can cause swimmer's itch or other infection.

## Cercarial Dermatitis

Alternate name; see *Swimmer's Itch*.

## Cerclage

A procedure in which a strong stitch is placed around the cervix to help prevent it opening too early during pregnancy. It may be placed through the vagina or abdomen and requires obstetric follow-up.

## Cerebellar

Relating to the cerebellum, the part of the brain that helps coordinate balance, posture, eye movements, and accurate movement. Cerebellar disease may cause clumsiness, tremor, or unsteady walking.

## Cerebellar Ataxia

Poor coordination caused by disease or injury affecting the cerebellum or its connections. It may cause an unsteady walk, clumsy movements, slurred speech, tremor, or abnormal eye movements.

## Cerebellar Cortex

The folded outer layer of the cerebellum containing Purkinje and granular neurons that coordinates balance, posture, timing, and precision of movement.

## Cerebellar Disease

A disorder of the cerebellum causing impaired coordination, gait ataxia, dysmetria, tremor, abnormal eye movements, or speech changes.

## Cerebellar Dysmetria

Inaccurate control of the distance or force of a move-

---

ment because of cerebellar dysfunction. A person may reach past a target, stop short, or have difficulty judging movement size.

### **Cerebellar Hemangioblastoma**

A usually benign, highly vascular tumor of the cerebellum that may cause headache, imbalance, vomiting, or hydrocephalus and can occur with von Hippel-Lindau disease.

### **Cerebellar Hypoplasia**

Incomplete development of the cerebellum, usually beginning before birth. It may cause poor coordination, abnormal eye movements, seizures, delayed development, or problems with balance and movement.

### **Cerebellar Neoplasm**

A benign or malignant tumor involving the cerebellum; symptoms may include poor coordination, imbalance, abnormal eye movements, headache, vomiting, or hydrocephalus.

### **Cerebellar Nuclei**

Deep clusters of cerebellar neurons that receive processed signals from the cerebellar cortex and send coordinated motor output to the brainstem and thalamus.

### **Cerebellar Peduncle**

One of three pairs of large, compact white matter fiber tracts (superior, middle, and inferior cerebellar peduncles) that physically and functionally connect the cerebellum to the midbrain, pons, and medulla oblongata, respectively.

### **Cerebellar Tonsillar Herniation**

The life-threatening downward displacement of the cerebellar tonsils through the foramen magnum into the upper cervical spinal canal, causing direct compression of the cardiorespiratory and vasomotor centers of the medulla oblongata.

#### **Alternate Names:**

Coning; Tonsillar Herniation

### **Cerebellar Tumor**

A growth in or near the cerebellum that may cause poor coordination, unsteady walking, abnormal eye movements, vomiting, or pressure buildup.

### **Cerebellar Vermis**

The midline portion of the cerebellum that helps control trunk posture, gait, balance, and coordination of eye and head movements.

### **Cerebellitis**

Inflammation of the cerebellum, often after an infection and sometimes from an immune reaction. It may cause sudden poor coordination, unsteady walking, slurred speech, abnormal eye movements, headache, or reduced alertness.

### **Cerebellopontine Angle**

The angle-shaped region at the junction of the cerebellum, pons, and petrous temporal bone. Lesions in this space may affect cranial nerves and commonly include vestibular schwannoma.

### **Cerebellopontine Angle Tumor**

A growth in the space between the cerebellum and brainstem. It may press on nerves that control hearing, balance, facial movement, or sensation and cause hearing loss, dizziness, facial weakness, headache, or poor coordination.

### **Cerebellum**

A region at the back of the brain, behind the brainstem, that coordinates balance, posture, eye movements, and accurate voluntary movement. Injury or disease can cause unsteady walking, clumsy movements, tremor, or slurred speech.

### **Cerebral**

Relating to the cerebrum or, more broadly, to the brain.

### **Cerebral Amyloid Angiopathy**

A disorder, more common in older adults, in which amyloid, usually amyloid-beta, builds up in the walls of small and medium-sized blood vessels in

---

the brain, especially near the brain's surface. The deposits weaken the vessels and can cause tiny areas of bleeding, small leaks, or a larger bleed in the outer parts of the brain. It may also cause brief neurological episodes, seizures, problems with thinking, or dementia. It can occur with Alzheimer disease, but the two conditions are not the same.

### **Cerebral Aneurysm**

An abnormal focal enlargement of an artery in or around the brain. It may remain asymptomatic or rupture and cause subarachnoid hemorrhage. Risk factors include hypertension, smoking, certain inherited disorders, and family history.

### **Cerebral Angiography**

An X-ray test in which contrast dye is put into an artery to make the blood vessels of the brain visible. It can show aneurysms, abnormal artery-to-vein connections, narrowing, or blockage and is usually done with a thin tube called a catheter.

### **Cerebral Aqueduct**

A narrow channel through the midbrain connecting the third and fourth brain ventricles. It allows cerebrospinal fluid to flow, and blockage can enlarge the ventricles and increase pressure.

### **Cerebral Arteriovenous Malformation**

An abnormal direct connection between cerebral arteries and veins without a normal capillary bed. It may cause intracranial hemorrhage, seizures, headache, or focal neurological deficits and is evaluated with brain imaging and angiography.

### **Cerebral Artery**

An artery that carries blood to part of the brain. A blockage can cause an ischemic stroke, while a break in the vessel can cause bleeding in or around the brain; the effects depend on which artery is involved.

### **Cerebral Atrophy**

Loss or shrinkage of brain tissue caused by aging, neurodegenerative disease, injury, infection, stroke, or another condition. Effects vary from none to

memory loss, weakness, seizures, or impaired thinking.

### **Cerebral Autoregulation**

The brain's ability to keep its blood flow relatively steady when blood pressure changes. This control can fail after severe brain injury, stroke, low oxygen, high carbon dioxide, or effects from some medicines.

### **Cerebral Calcification**

Abnormal calcium deposition in brain tissue, with causes including inherited, metabolic, infectious, vascular, and age-related disorders.

### **Cerebral Cavernous Malformation**

A low-flow vascular malformation composed of dilated thin-walled vascular spaces in the brain or spinal cord. It may cause seizures, headaches, or hemorrhage, although some lesions remain asymptomatic.

### **Cerebral Cavernous Malformations**

Groups of unusually widened, thin-walled blood vessels in the brain or spinal cord. They may cause seizures, headaches, weakness, or bleeding; some cause no symptoms. Multiple lesions can run in families because of inherited gene changes.

### **Cerebral Cortex**

The folded outer layer of the cerebrum containing nerve cells that support thinking, sensation, movement, language, memory, and awareness. Its folds increase the amount of working brain tissue within the skull.

### **Cerebral Crus**

A bundle of nerve fibers in the midbrain that carries movement signals from the brain's outer layer toward the brainstem and spinal cord. Damage can cause weakness or poor control on the opposite side.

### **Cerebral Diplegia**

A form of cerebral palsy in which stiffness and weakness affect both legs more than the arms. It can make standing, walking, balance, and other move-

---

ments difficult.

### **Cerebral Edema**

Swelling caused by excess fluid in or around brain tissue. It can raise pressure inside the skull, reduce blood flow, and cause headache, vomiting, confusion, seizures, coma, or life-threatening brain displacement.

### **Cerebral Embolism**

Alternate name; see *Embolic Stroke*.

### **Cerebral Haemorrhage**

Bleeding in or around the brain caused by a damaged blood vessel. It can injure brain tissue, increase pressure inside the skull, and cause sudden headache, weakness, confusion, seizures, or loss of consciousness.

### **Cerebral Hemisphere**

One of the two large halves of the cerebrum, each containing specialized but interconnected cortical regions.

### **Cerebral Hemorrhage**

Bleeding into brain tissue, usually from a damaged blood vessel. It can injure nearby brain cells, cause swelling and pressure, and produce sudden weakness, speech trouble, headache, reduced alertness, or death.

### **Cerebral Hypoxia**

Too little oxygen reaching the brain, often because of cardiac arrest, severe breathing failure, low blood pressure, or choking. Prolonged oxygen shortage can permanently injure brain cells.

### **Cerebral Infarction**

Permanent brain injury caused by blocked blood flow in a brain artery. The blockage may result from a clot formed in a narrowed artery or a clot traveling from the heart or another vessel.

### **Cerebral Ischemia**

Reduced blood flow and oxygen reaching brain tis-

sue. It may cause temporary symptoms, such as a transient ischemic attack, or permanent injury if the interruption lasts long enough.

### **Cerebral Lobes**

Major regions of the cerebrum with different but overlapping functions: the frontal lobe supports planning and movement, the parietal sensation and spatial awareness, the temporal hearing and memory, and the occipital vision.

### **Cerebral Malaria**

A life-threatening form of malaria in which parasites cause brain dysfunction, often with confusion, seizures, abnormal behavior, or coma. It is usually caused by *Plasmodium falciparum* and requires urgent treatment.

### **Cerebral Palsy**

A group of lifelong disorders, abbreviated CP, caused by abnormal development or damage to the developing brain. It affects movement, muscle control, balance, coordination, and posture. The brain problem usually does not worsen, but a person's symptoms and abilities may change with age. Some people also have seizures or problems with learning, speech, swallowing, vision, hearing, or joint movement. Common types include spastic, dyskinetic, ataxic, and mixed cerebral palsy.

#### **Alternate Names:**

CP; Spastic Diplegia; Infantile Cerebral Paralysis

### **Cerebral Palsy Entry**

Alternate name; see *Cerebral palsy*.

### **Cerebral Peduncle**

One of two bundles of nerve fibers in the midbrain that carry movement signals from the cerebrum to the brainstem and spinal cord. Damage can cause weakness or loss of coordinated movement on the opposite side.

### **Cerebral Perfusion Pressure**

The pressure pushing blood through the brain, estimated from average arterial pressure minus pressure inside the skull. It falls when blood pressure drops

---

or brain swelling raises skull pressure.

### **Cerebral Salt Wasting**

Excessive loss of sodium and water in urine associated with brain injury or disease. It can cause low blood sodium and dehydration and requires careful replacement guided by clinical and laboratory monitoring.

### **Cerebral Salt-Wasting Syndrome**

A clinical disorder of hyponatremia and extracellular fluid volume depletion occurring in acute central nervous system disease (such as subarachnoid hemorrhage or head trauma), characterized by renal natriuresis, polyuria, and true intravascular hypovolemia.

**Alternate Names:**

CSWS

### **Cerebral Thrombosis**

Formation of a blood clot within a cerebral artery, reducing blood flow and potentially causing an ischemic stroke.

### **Cerebral Toxoplasmosis**

A severe, life-threatening opportunistic central nervous system infection caused by the reactivation of latent *Toxoplasma gondii* tissue cysts, occurring almost exclusively in patients with advanced cellular immunodeficiency (such as individuals with AIDS with CD4 counts below 100 cells/ $\mu$ L or recipients of organ transplants). It presents with progressive headache, fever, altered mental status, seizures, and focal neurological deficits, with contrast neuroimaging characteristically revealing multiple ring-enhancing brain lesions with surrounding vasogenic edema predominantly in the basal ganglia and corticomedullary junction.

**Alternate Names:**

Toxoplasma Encephalitis; CNS Toxoplasmosis; Toxoplasmic Encephalitis

### **Cerebral Tumor**

An abnormal growth in or involving the brain. Symptoms depend on its location, size, growth, and effect on pressure inside the skull and may include

headache, seizures, weakness, or changes in thinking.

### **Cerebral Tumours**

Abnormal growths in the cerebrum, the largest part of the brain. They may be noncancerous or cancerous and can cause seizures, headache, changes in thinking, weakness, speech problems, or vision changes.

### **Cerebral Vasospasm**

Abnormal narrowing of cerebral arteries that can reduce brain perfusion and cause delayed cerebral ischemia, especially after aneurysmal subarachnoid hemorrhage.

### **Cerebral Venous Sinus Thrombosis**

A blood clot in a large vein or drainage channel of the brain. It can cause headache, seizures, visual problems, weakness, brain bleeding, or increased pressure inside the skull.

### **Cerebral Ventricle**

One of four connected spaces inside the brain that contain cerebrospinal fluid. The fluid cushions the brain and spinal cord, carries nutrients and waste, and flows through the ventricles before being absorbed around the brain.

### **Cerebral Ventricles**

One of four connected spaces inside the brain that contain and help circulate cerebrospinal fluid. This fluid cushions the brain and spinal cord and helps remove waste.

**Alternate Names:**

Cerebral Ventricle; Ventricular System

### **Cerebral Ventriculography**

Imaging of the brain's fluid-filled ventricles to assess their size, shape, connections, or blockage. Modern scans usually use computed tomography or magnetic resonance imaging rather than older invasive methods.



---

## Cerebral Ventriculomegaly

Enlargement of one or more fluid-filled spaces inside the brain. It may result from blocked fluid drainage, loss of brain tissue, or developmental variation; its importance depends on the cause and severity.

## Cerebral Visual Impairment

Reduced vision caused by damage to the brain's visual pathways or visual-processing areas rather than the eyes. It may cause poor visual attention, difficulty recognizing objects, or inconsistent use of vision.

## Cerebriform Nevus Sebaceous

A usually harmless birthmark made of overgrown oil glands, often on the scalp or face, with a soft, folded surface resembling the brain. It can rarely develop another tumor, so persistent or changing lesions need review.

## Cerebritis

Inflammation and infection of brain tissue before a definite abscess forms, usually caused by bacteria, fungi, or parasites. It can cause headache, fever, seizures, or neurologic deficits and requires urgent imaging and antimicrobial treatment.

## Cerebroplacental Ratio

A number calculated from fetal Doppler ultrasound by comparing blood flow in a brain artery with blood flow in the umbilical artery. A low result may suggest that the fetus is adapting to reduced placental supply and must be interpreted with other findings.

## Cerebrosides

Fat-containing substances found especially in the membranes and protective coverings of nerve cells. They help maintain nerve structure and are broken down abnormally in some inherited storage diseases.

## Cerebrospinal Fluid

A clear fluid surrounding the brain and spinal cord and filling spaces inside the brain. It cushions the nervous system, helps carry nutrients and waste, and is continuously produced and reabsorbed by

the body.

### Alternate Names:

Spinal Fluid

## Cerebrospinal Fluid Analysis

Laboratory examination of cerebrospinal fluid, usually obtained by lumbar puncture. It may assess infection, bleeding, inflammation, demyelinating disease, or malignant cells. Analysis can include opening pressure, appearance, cell counts, protein, glucose, microbiology, and other tests selected for the clinical question.

## Cerebrospinal Fluid Collection

A procedure, usually lumbar puncture, that collects cerebrospinal fluid for laboratory analysis; it can help evaluate infection, bleeding, inflammation, malignancy, and other disorders of the central nervous system.

## Cerebrospinal Fluid Culture

A laboratory test that attempts to grow bacteria, fungi, or other microorganisms from cerebrospinal fluid to investigate meningitis or another central nervous system infection. Results depend on prior antimicrobial treatment, specimen volume, transport, and the clinical context.

## Cerebrospinal Fluid Cytology

Microscopic examination of cells in cerebrospinal fluid to evaluate suspected malignancy, infection, inflammation, or hemorrhage. It is especially useful for detecting leptomeningeal tumor involvement when interpreted with clinical and imaging findings.

## Cerebrospinal Fluid Leak

Escape of cerebrospinal fluid through a tear in the membranes surrounding the brain or spinal cord, often causing positional headache.

### Alternate Names:

Spinal Fluid Leak

## Cerebrotendinous Xanthomatosis (CTX)

A rare inherited disorder of bile-acid synthesis, usually caused by *CYP27A1* variants. Cholesterol-

---

derived substances accumulate in the brain, tendons, and other tissues and may cause cataracts, chronic diarrhea, tendon xanthomas, seizures, ataxia, or progressive neurologic disease.

### **Cerebrovascular**

Relating to the brain's blood vessels and blood flow. Cerebrovascular problems can reduce oxygen delivery and cause strokes, brief warning episodes, bleeding, or lasting problems with movement, speech, vision, or thinking.

#### **Alternate Names:**

Brain-Vascular; Cerebral Vascular

### **Cerebrovascular Accident**

A stroke caused by a blocked brain artery or bleeding in the brain. It can suddenly cause weakness, numbness, trouble speaking, vision loss, confusion, severe headache, or loss of balance and requires emergency care.

### **Cerebrovascular Accident Prevention**

Measures to reduce the risk of stroke, including controlling blood pressure, diabetes, and lipids, avoiding tobacco, treating atrial fibrillation when indicated, and using appropriate antithrombotic therapy.

### **Cerebrovascular Disease**

A group of disorders affecting blood vessels and blood flow in the brain, including stroke, transient ischemic attack, aneurysm, and vascular malformations.

### **Cerebrovascular Disorder**

A disorder of brain blood vessels or cerebral blood flow, including ischemia, hemorrhage, aneurysm, malformation, and venous thrombosis.

### **Cerebrovascular Occlusion**

Blockage or closure of a blood vessel supplying the brain, reducing blood flow and potentially causing brain injury or an ischemic stroke.

### **Cerebrum**

The largest part of the brain, made of two connected halves. It supports conscious movement, sensation, language, memory, thinking, emotions, and vision; different areas specialize in different functions.

### **Certification**

A formal process confirming that a person, service, laboratory, or facility meets specified training, quality, safety, or professional standards.

### **Certified Nurse-Midwife**

An advanced-practice nurse trained and credentialed to provide reproductive, pregnancy, birth, postpartum, and newborn care, including management of normal pregnancy and referral of complications.

### **Certolizumab Pegol**

A biologic medicine that blocks tumor necrosis factor and treats selected inflammatory diseases, including rheumatoid arthritis and Crohn disease. It can increase the risk of serious infections.

### **Ceruloplasmin**

The main protein that carries copper in the blood. It is made mostly by the liver; unusually low or high levels can occur in Wilson disease, inflammation, liver disease, or other conditions.

#### **Alternate Names:**

Caeruloplasmin

### **Ceruloplasmin Blood Test**

A blood test measuring ceruloplasmin, the main copper-carrying protein in plasma; it is used with other copper studies to evaluate Wilson disease and disorders of copper metabolism.

### **Cerumen**

Earwax, a waxy substance made in the ear canal. It traps dust and helps protect and moisturize the canal; too much can block hearing or cause discomfort.

### **Cerumen Impaction**

Earwax packed tightly enough to block the ear canal

---

or prevent examination of the eardrum. It may cause blocked hearing, pressure, ringing, itching, pain, or dizziness.

### **Ceruminous Adenocarcinoma**

A malignant tumor of the ceruminous glands in the external ear canal that may cause pain, discharge, hearing loss, or a persistent ear-canal mass.

### **Cervical**

Relating to the neck or to the cervix, the lower opening of the uterus. The meaning depends on context, such as cervical spine, cervical pain, or cervical cancer.

### **Cervical Adenocarcinoma**

A malignant gland-forming tumor arising in the uterine cervix; it may be HPV-associated or HPV-independent and commonly presents with abnormal bleeding or discharge.

### **Cervical Arthritis**

Degenerative or inflammatory disease of the neck joints that can cause neck pain, stiffness, reduced motion, and sometimes nerve or spinal-cord compression.

### **Cervical Atlas**

The first bone of the neck, called C1. It is a ring-shaped vertebra that supports the skull and moves around the second neck vertebra when the head turns.

### **Cervical Biopsy**

Removal of a small piece of tissue from the cervix for laboratory examination. It is commonly performed after an abnormal screening result to look for precancerous or cancerous changes.

### **Cervical Cancer**

Cancer arising in the cervix, most often after persistent infection with a high-risk human papillomavirus type. Early disease may cause no symptoms; later signs include unusual bleeding, discharge, or pelvic pain.

### **Cervical Cancer Screening and Prevention**

Secondary and primary preventive interventions aimed at eliminating invasive cervical carcinoma, encompassing prophylactic human papillomavirus (HPV) vaccination, routine cervical cytology (Pap smear), and high-risk HPV nucleic acid co-testing.

#### **Alternate Names:**

Cervical Cancer Screening; Pap Smear Screening; Pap Smear; HPV Co-Testing; Cervical Cytology Screening

### **Cervical Cap**

A flexible contraceptive barrier placed over the cervix to keep sperm from entering the uterus. It is used with spermicide and must remain in place for the recommended time after sex.

### **Cervical Carcinoma**

A malignant tumor of the uterine cervix, most commonly associated with persistent high-risk human papillomavirus infection.

### **Cervical Cerclage**

A procedure that places a strong stitch around the cervix during pregnancy to help keep it closed when early opening raises the risk of pregnancy loss or premature birth. It may be placed through the vagina or abdomen.

### **Cervical Change**

A change in the uterine cervix or cervical spine, depending on context; clinical significance is determined by the specific structural, inflammatory, or pregnancy-related finding.

### **Cervical Dilatation**

Opening of the cervix during labor, measured from closed to about 10 centimeters. The cervix also becomes thinner and shorter, allowing the baby's head to pass into the birth canal.

### **Cervical Dilation**

The opening of the cervical canal during labor, progressing from closed to about 10 centimeters. It

---

occurs with thinning and shortening of the cervix and is needed for vaginal birth.

### **Cervical Disc Herniation**

A neck disc bulges or tears and presses on a nearby nerve root or the spinal cord. It may cause neck pain, arm pain, tingling, numbness, or weakness.

### **Cervical Dislocation**

Abnormal displacement of one cervical vertebra relative to another, usually from trauma, that can destabilize the neck and injure the spinal cord.

### **Cervical Dysplasia**

See *Cervical Intraepithelial Neoplasia*.

### **Cervical Dystocia**

Failure of the uterine cervix to dilate normally during labor, which can delay or prevent vaginal delivery.

### **Cervical Dystonia**

Involuntary tightening of neck muscles that pulls the head into an abnormal position. It may cause neck pain, twisting, jerky movements, or tremor.

#### **Alternate Names:**

Spasmodic Torticollis; Wry Neck

### **Cervical Ectropion**

A common condition in which cells normally lining the cervical canal extend onto the outer cervix. It may cause increased discharge or bleeding after sex, but it is usually benign and is not the same as cervical cancer.

### **Cervical Effacement**

Thinning and shortening of the cervix during labor, measured from 0% to 100%. Complete effacement leaves the cervix very thin and usually occurs with opening of the cervix.

### **Cervical Endometriosis**

Endometrial-like tissue in or on the cervix that may cause bleeding after sex, pelvic pain, or an abnormal cervical lesion.

### **Cervical Exam In Labor**

A vaginal examination during labor that checks how open and thin the cervix is and how low and positioned the baby's head is. Repeated examinations help assess progress, but are limited when infection or bleeding is a concern.

### **Cervical Ganglion**

A cluster of nerve-cell bodies in the sympathetic chain of the neck that relays autonomic signals to the head, neck, and upper chest.

### **Cervical Incompetence**

Older term; see *Cervical Insufficiency*.

### **Cervical Insufficiency**

Painless shortening or opening of the cervix during pregnancy before term, increasing risk of second-trimester loss or spontaneous preterm birth.

### **Cervical Intraepithelial Neoplasia**

Biopsy-confirmed abnormal changes in the surface squamous cells of the cervix, usually associated with persistent high-risk human papillomavirus infection. CIN is graded from 1 to 3 by how abnormal the cells look and how much tissue is affected. CIN is not cancer; many low-grade changes clear, while high-grade changes have a greater chance of becoming cancer if not monitored or treated.

#### **Alternate Names:**

CIN

### **Cervical Length Screening**

Ultrasound measurement of the cervix, usually through the vagina, to estimate the risk of premature birth. A shorter cervix can increase risk, but interpretation depends on gestational age, pregnancy history, symptoms, and the measurement method.

### **Cervical Lymphadenopathy**

Enlarged lymph nodes in the neck, commonly caused by infection but sometimes reflecting malignancy, autoimmune disease, or other systemic illness.

---

### Cervical MRI Scan

Magnetic resonance imaging of the cervical spine that uses a magnetic field and radio waves to produce detailed images of the neck vertebrae, discs, spinal cord, nerves, and surrounding soft tissues.

### Cervical Mucus

Fluid made by glands in the cervix. It becomes thinner and more slippery around ovulation, which can help sperm move, and thicker at other times. During pregnancy, it helps form a protective plug in the cervical canal.

### Cervical Myelopathy

Dysfunction of the spinal cord caused by compression in the cervical spine, often producing hand clumsiness, gait disturbance, weakness, sensory changes, or abnormal reflexes.

### Cervical Osteoarthritis

Alternate name; see *Cervical Spondylosis*.

### Cervical Pain

Alternate name; see *Neck Pain*.

### Cervical Plexus

A network of nerves in the neck that carries sensation from the neck and part of the head and helps control some neck muscles and the diaphragm.

### Cervical Polyp

A usually benign polypoid overgrowth of endocervical or ectocervical tissue that may cause intermenstrual or postcoital bleeding or may be found incidentally during examination.

### Cervical Polyps

Usually noncancerous growths arising from the cervix or its canal. They may cause bleeding between periods, bleeding after sex, or discharge and are often removed for laboratory examination.

### Cervical Pregnancy

A rare ectopic pregnancy implanted in the cervical canal rather than the main uterine cavity. It may

cause painless early-pregnancy bleeding and can lead to severe hemorrhage, requiring urgent specialist care.

### Cervical Radiculopathy

Irritation or compression of a nerve leaving the neck, often from a disc problem or arthritis. It can cause neck pain spreading into the arm, with tingling, numbness, or weakness.

### Cervical Rib

An extra rib growing from a neck vertebra, usually the lowest one. It often causes no symptoms but can narrow the space for nerves or blood vessels and produce arm pain, numbness, or swelling.

### Cervical Ripening

Softening, thinning, and early opening of the cervix before or during labor. It may occur naturally or be induced with medicines or mechanical methods when labor needs to be started; the method depends on cervical findings, pregnancy factors, and uterine-scar history.

### Cervical Ripening Agents

Medicines or mechanical devices used to soften and prepare the cervix before labor induction. Options include prostaglandins and cervical balloons; selection and dosing depend on the cervix, pregnancy, contraindications, and local obstetric protocol.

### Cervical Screening

Testing used to detect high-risk human papillomavirus infection and precancerous or cancerous changes of the cervix. The method and interval depend on age, prior results, vaccination, immune status, and local guidelines.

### Cervical Spine

The seven bones in the neck, numbered C1 through C7. They support the head, protect the upper spinal cord, allow neck movement, and contain openings for important blood vessels.

---

### Cervical Spine CT Scan

Computed tomography of the cervical spine that uses x-rays and computer reconstruction to show the bones and alignment of the neck; it is particularly useful for detecting fractures and other bony abnormalities.

### Cervical Spine Involvement in Rheumatoid Arthritis

Rheumatoid inflammation affecting the upper cervical joints, especially the atlantoaxial joint. Instability or bone erosion can cause neck pain, occipital headache, nerve symptoms, or spinal-cord compression and may require imaging before procedures or surgery.

### Cervical Spine Pain

Alternate name; see *Neck Pain*.

### Cervical Spine Trauma

An injury to the neck bones, discs, ligaments, spinal cord, or exiting nerves. It may cause pain, weakness, numbness, or paralysis; suspected serious injury requires immobilization and urgent assessment.

### Cervical Spondyloarthropathy

Wear or inflammation affecting the joints and discs of the neck. It may cause neck stiffness and pain, or press on nerves or the spinal cord, producing arm symptoms, weakness, or walking problems.

### Cervical Spondylosis

Age-related wear of the neck, including changes in discs, bones, joints, and ligaments. It may cause neck pain and stiffness and, if nerves or the spinal cord are compressed, limb symptoms.

#### Alternate Names:

Cervical Osteoarthritis; Degenerative Cervical Spine Disease

### Cervical Spondylotic Myelopathy

Progressive spinal cord compression resulting from age-related degenerative osteoarthritic changes in the cervical spine (intervertebral disc herniation, osteophytic spurring, ligamentum flavum hypertro-

phy), presenting with clumsy hands, spastic gait ataxia, hyperreflexia, and Hoffman sign.

#### Alternate Names:

CSM; Cervical Spondylotic Myelopathy

### Cervical Squamous Cell Carcinoma

A cancer that begins in the flat surface cells of the cervix, most often after persistent infection with a high-risk human papillomavirus type. It may cause unusual bleeding or discharge, but early disease may have no symptoms.

### Cervical Squamous Intraepithelial Lesion

Alternate name; see *Cervical dysplasia*.

### Cervical Stenosis

Narrowing or complete closure of the passage through the cervix, the lower part of the uterus. It may be present from birth or develop after menopause, cervical surgery, radiation, endometrial procedures, or cervical or uterine cancer. It often causes no symptoms, but may cause painful or absent periods, abnormal bleeding, trapped blood or pus in the uterus, or rarely infertility. This term refers to the cervix; narrowing of the spinal canal is called cervical spinal stenosis.

### Cervical Strain (Whiplash)

Injury to the neck muscles, ligaments, and other soft tissues caused by rapid flexion and extension, commonly during a motor-vehicle collision. Neck pain, stiffness, headache, and shoulder or arm symptoms may occur.

### Cervical Vertebrae

The seven bones of the neck, numbered C1 through C7. They support the head, protect the upper spinal cord, and allow neck movement; openings in several of them carry the vertebral arteries.

### Cervicitis

Inflammation of the cervix, commonly caused by a sexually transmitted infection, changes in normal vaginal bacteria, chemical irritation, or allergy. It may cause discharge, bleeding after sex, or pelvic discomfort, but can be symptomless.



---

## Cervicogenic Headache

Head pain caused by a problem in the neck, such as a joint, disc, muscle, or nerve disorder. Pain often starts in the neck and spreads to one side of the head and may worsen with neck movement.

### Alternate Names:

Cervical Headache; Neck-Related Headache

## Cervix

The lower part of the uterus that opens into the vagina. It lets menstrual blood leave the uterus, allows sperm to enter, produces cervical mucus, and widens during childbirth.

## Cervix Carcinoma

Cancer arising from the cervix, most often related to persistent high-risk human papillomavirus infection; screening and HPV vaccination reduce risk, and treatment depends on stage.

## Cervix Cryosurgery

Freezing abnormal surface cells of the cervix with a very cold probe. It may treat selected precancerous changes after appropriate testing; watery discharge and cramping are common temporarily.

## Cervix Disease

Any disorder affecting the lower part of the uterus, including infection, polyps, precancer, cancer, scarring, or pregnancy-related changes. Symptoms and treatment depend on the specific condition.

## Cervix Neoplasm

A benign, precancerous, or malignant growth of the cervix; evaluation may include pelvic examination, HPV testing, cytology, colposcopy, and biopsy.

## Cervix Uteri

The cervix uteri is the lower portion of the uterus that projects into the vagina and forms the passage between the uterus and birth canal.

## Cesarean Delivery

Alternate name; see *Cesarean section*.

## Cesarean Section

A surgical procedure used to deliver a baby through incisions made in the mother's abdomen and uterus. It may be scheduled in advance (planned or elective) or performed urgently when vaginal delivery becomes unsafe for the mother or baby. Common reasons include fetal distress, abnormal fetal position (such as breech), placenta covering the cervix (placenta previa), prolonged or stalled labor, and prior uterine surgeries. Most modern procedures utilize a low transverse incision in the lower segment of the uterus under regional anesthesia (spinal or epidural), allowing the mother to remain awake during delivery.

### Alternate Names:

C-Section; Caesarean Section; Cesarean Delivery; Caesarean Delivery; Lower Segment Cesarean Section; LSCS

## Cesium

A chemical element whose radioactive isotope cesium-137 can cause serious external or internal radiation injury after contamination or exposure.

## Cessation of Menstruation

Cessation of menstrual periods, called amenorrhea; it is a normal part of menopause but may also result from pregnancy, illness, endocrine disorders, medications, stress, excessive exercise, or inadequate nutrition.

## Cestoda

The tapeworm class of flatworms; human infection follows ingestion of eggs or larval tissue and can cause intestinal disease or tissue cysts.

## Cestode

A cestode is a tapeworm, a segmented parasitic flatworm that can infect humans or animals through contaminated food, water, or intermediate hosts.

## Cestodiasis

Infection or infestation caused by cestodes, commonly called tapeworms. Disease may involve the intestine or tissues, depending on the species and life-cycle stage.

---

**Alternate Names:**

Tapeworm Infection; Cestode Infection

**Cetirizine**

A second-generation H1 antihistamine used for allergic rhinitis and urticaria; it is less sedating than older antihistamines but can still cause drowsiness.

**Cetrimide**

An antiseptic and detergent used in some skin preparations and laboratories. It reduces microorganisms by disrupting cell membranes, but concentrated or swallowed amounts can irritate or burn tissues.

**Cetrorelix**

A hormone-blocking medicine used during assisted reproduction to prevent premature ovulation. It temporarily blocks gonadotropin-releasing hormone receptors; injection-site reactions, headache, nausea, or allergy may occur.

**Cetuximab**

A monoclonal antibody that blocks the epidermal growth factor receptor. It is used for selected colorectal, head-and-neck, and other cancers and commonly causes acne-like rash.

**Cevimeline**

A muscarinic receptor agonist used to stimulate salivary secretion and relieve dry mouth, especially in selected patients with Sjögren syndrome.

**Cevimeline Hydrochloride**

A medicine that stimulates muscarinic receptors to increase saliva. It is used mainly for dry mouth in Sjögren syndrome and can cause sweating, nausea, or blurred vision.

**CF**

Alternate name; see *Cystic Fibrosis*.

**CF (Cystic Fibrosis)**

Abbreviation for cystic fibrosis, an inherited disorder caused by CFTR variants that produce thick secretions, chronic lung infection, pancreatic insuffi-

ciency, infertility in many males, and other organ complications.

**CFS**

Chronic fatigue syndrome, a condition involving substantial persistent fatigue with post-exertional worsening and associated symptoms that are not explained by another disorder.

**Alternate Names:**

Chronic Fatigue Syndrome; Myalgic Encephalomyelitis/Chronic Fatigue Syndrome

**CFS (Chronic Fatigue Syndrome)**

Abbreviation for myalgic encephalomyelitis/chronic fatigue syndrome, a complex illness involving substantial reduction in activity, post-exertional malaise, unrefreshing sleep, and often cognitive impairment or orthostatic intolerance.

**CFTR Modulator Therapy**

Medicines that improve the amount or function of defective CFTR protein in people with cystic fibrosis. The appropriate combination depends on the person's CFTR variants; treatment can improve breathing, nutrition, and salt-water balance.

**CHA2DS2-VASc Score**

A score estimating stroke risk in a person with atrial fibrillation. It considers heart failure, high blood pressure, age, diabetes, previous stroke, blood-vessel disease, and sex; the result helps clinicians decide whether a blood thinner may offer more benefit than harm.

**Alternate Names:**

CHA2DS2-VASc; Atrial Fibrillation Stroke Risk Score

**Chaddock Reflex**

An alternative neurological test for corticospinal tract upper motor neuron dysfunction, elicited by stroking the skin of the lateral foot around the lateral malleolus from posterior to anterior, producing an extensor plantar response (great toe dorsiflexion).

**Alternate Names:**

Chaddock Sign

---

### Chaddock Sign

A pathological neurological reflex elicited by firmly stroking the skin beneath and around the lateral malleolus from posterior to anterior, which provokes extension (dorsiflexion) of the great toe and fanning of the other toes, signaling corticospinal tract (upper motor neuron) dysfunction.

#### Alternate Names:

Chaddock Reflex; Chaddock's Sign

### Chadwick Sign

A bluish-purple discoloration of the cervix, vagina, and vulval mucosa caused by local venous engorgement and heightened vascularity in early pregnancy, typically observable by speculum examination around 6 to 8 weeks of gestation.

#### Alternate Names:

Chadwick's Sign; Jacquemier Sign

### Chafing

Skin irritation caused by repeated friction, moisture, heat, or tight clothing, producing redness, burning, tenderness, and sometimes erosions in opposing or rubbing areas.

### Chagas Cardiomyopathy

The chronic, progressive inflammatory and fibrosing cardiac disease resulting from long-standing infection with the protozoan parasite *Trypanosoma cruzi* (Chagas disease), emerging 10 to 30 years after primary infection in up to 30% of chronically infected individuals. It is characterized by biventricular enlargement, apical left ventricular aneurysms, conduction system disease (notably right bundle branch block and left anterior fascicular block), malignant ventricular tachyarrhythmias, heart failure, and systemic thromboembolism.

#### Alternate Names:

Chagasic Cardiomyopathy; Chronic Chagasic Cardiomyopathy; Chagas Heart Disease; *Trypanosoma Cruzi* Cardiomyopathy

### Chagas Disease

An infection caused by the parasite *Trypanosoma cruzi*, usually spread by infected kissing bugs. It can also pass from mother to baby or through con-

taminated blood, organs, or food. Early illness may be mild or absent; years later, some people develop heart disease or enlargement of the oesophagus or colon.

### Chagasic Megacolon

A severe gastrointestinal complication of chronic Chagas disease (*Trypanosoma cruzi* infection) caused by parasite-induced destruction of intramural autonomic ganglia in the myenteric (Auerbach) plexus of the large intestine. The resulting denervation produces loss of coordinated peristalsis, progressive organ dilatation, and muscular hypertrophy predominantly affecting the sigmoid colon and rectum, clinically manifesting with severe intractable constipation, abdominal distention, fecaloma impaction, and risk of sigmoid volvulus.

#### Alternate Names:

Mega-Megacolon in Chagas Disease; Chagas Megacolon

### Chagoma

A red, swollen lump where the parasite causing Chagas disease enters the body. It may be painful and accompanied by swollen nearby lymph nodes during the early stage of infection.

### Chain Of Custody

A dated record showing who collected, stored, transferred, tested, and finally disposed of an item or specimen. It helps prove that medical, forensic, or laboratory evidence was identified correctly and was not altered or mixed up.

### Chain-Termination Codon

A codon that signals the end of protein translation; the three standard stop codons are UAA, UAG, and UGA in messenger RNA.

### Chalazion

A usually painless, firm eyelid lump caused by blockage and inflammation of a meibomian oil gland; persistent or atypical lesions require ophthalmic assessment.

---

### Challenge Agent

A substance, organism, allergen, or physiologic stimulus deliberately administered in a controlled test to provoke and measure a response.

### Challenge Testing

A controlled exposure to a suspected trigger, such as a food or medication, to assess whether it produces a reproducible reaction.

#### Alternate Names:

Provocation Testing; Challenge Test

### Chamberlain Procedure

An anterior mediastinotomy surgical technique involving an incision through the second or third left anterior intercostal space with resection of cartilage, designed to biopsy anterior mediastinal masses or aortopulmonary window lymph nodes that are inaccessible via cervical mediastinoscopy.

#### Alternate Names:

Anterior Mediastinotomy

### Chamomile Extract

An extract of chamomile flowers used in traditional and complementary products; allergic reactions and medicine interactions are possible, while evidence varies by indication.

### Chanarin-Dorfman Disease

A rare inherited lipid-storage disorder, usually caused by changes in the *ABHD5* gene, in which triglycerides accumulate in skin, liver, muscle, blood cells, and other tissues. It commonly causes congenital ichthyosis and may also cause liver disease, cataracts, hearing loss, muscle weakness, or developmental problems.

#### Alternate Names:

Chanarin-Dorfman Syndrome; Dorfman-Chanarin Disease; Neutral Lipid Storage Disease With Ichthyosis

#### Alternate Names:

American Trypanosomiasis; Trypanosoma Cruzi Infection

### Chance Fracture

A horizontal flexion-distraction injury of the thoracolumbar spine involving a transverse fracture line through the vertebral body, pedicles, laminae, and spinous process. Classically associated with lap-type seatbelt restraint deceleration injuries in motor vehicle collisions, it is frequently accompanied by blunt intra-abdominal organ trauma.

#### Alternate Names:

Seatbelt Fracture; Flexion-Distraction Spine Fracture

### Chancre

A usually painless ulcer at the site of inoculation of an infection, especially the primary lesion of syphilis. It may be accompanied by regional lymph-node enlargement and heals without treatment, although infection persists.

### Chancroid

A sexually transmitted infection caused by *Haemophilus ducreyi*, producing one or more painful, ragged genital ulcers with a purulent base and tender, sometimes suppurative groin lymph nodes. Clinical diagnosis requires excluding syphilis and genital herpes; antibiotic treatment and partner evaluation are important.

### Chandelier Sign

Severe lower belly pain felt when a doctor moves the neck of the womb during an examination, usually pointing to an infection in the female reproductive organs.

### Chandipura Virus Infection

An acute, fulminant pediatric encephalitis caused by Chandipura virus, a vesiculovirus belonging to the family Rhabdoviridae and transmitted by phlebotomine sandflies (*Phlebotomus papatasi*). Predominantly affecting children under 15 years of age in central and western India, it is characterized by the rapid onset of high fever, vomiting, and altered sensorium, progressing within 24 to 48 hours to generalized convulsions, coma, and respiratory failure, carrying an exceptionally high case-fatality rate.

#### Alternate Names:

---

Chandipura Encephalitis; Chandipura Viral Encephalitis

### **Change in Bowel Habits**

A new change in stool frequency, consistency, caliber, or pattern, including persistent diarrhea, constipation, or alternating bowel habits. A persistent or unexplained change, especially with bleeding, weight loss, anemia, or severe pain, requires medical evaluation.

### **Channelopathy**

A disorder caused by abnormal proteins that control movement of charged particles through cell membranes. It can affect nerves, muscles, or the heart and may cause seizures, weakness, or abnormal rhythms.

### **Chaplain**

A trained spiritual-care professional who supports patients and families with religious, existential, emotional, or ethical concerns in healthcare settings.

### **Chapped Hands**

Dry, cracked, inflamed skin of the hands caused by irritants, cold, low humidity, frequent washing, eczema, or allergy; fissures can become painful or infected.

### **Chapped Lips**

Dry, scaling, fissured lips caused by weather, licking, dehydration, irritants, medicines, or cheilitis; persistent lesions need assessment for infection or another disorder.

### **Char Syndrome**

A rare genetic disorder, usually caused by variants in *TFAP2B*, characterized by distinctive facial features, a heart defect such as patent ductus arteriosus, and abnormalities of the fifth fingers or toes.

### **Charcoal**

A black carbon-rich material made by heating organic matter with little oxygen. In medicine, specially prepared charcoal may be given in selected

poisonings to reduce absorption of some substances, but it is not useful for every poisoning.

### **Charcot Arthropathy**

Progressive joint damage caused by loss of protective sensation, so repeated injuries go unnoticed. The joint may become warm, swollen, unstable, and deformed, especially in people with diabetic nerve damage.

### **Charcot Foot**

Progressive weakening and deformity of the foot or ankle caused by repeated unnoticed injury in a person with reduced sensation, commonly from diabetes. Early warmth and swelling require urgent off-loading to prevent collapse and ulcers.

### **Charcot Joint**

Severe destruction of a joint caused by loss of pain and position sensation. Repeated unnoticed injury can fragment bone, loosen or dislocate the joint, and produce major deformity.

### **Charcot Triad**

The combination of fever, pain in the upper-right abdomen, and jaundice, traditionally associated with acute infection of blocked bile ducts. This pattern is important but may be absent even in serious disease.

### **Charcot Triad of Cholangitis**

The combination of fever, pain in the upper-right abdomen, and jaundice. It is a classic pattern of infection in blocked bile ducts, but many people with serious infection do not have all three signs.

### **Charcot's Joints**

Charcot's joints are severely damaged, unstable joints caused by loss of protective sensation, classically in diabetic neuropathy or other neurologic disease.

### **Charcot's Triad**

Charcot's triad is fever, jaundice, and right-upper-quadrant pain, a classic but incomplete clinical pat-



---

tern of acute ascending cholangitis.

### **Charcot-Bouchard Aneurysm**

A minute microaneurysm arising on small (100 to 300 micrometer) lenticulostriate penetrating arteries of the basal ganglia and thalamus in patients with chronic hypertension; its rupture is the classic cause of primary intracerebral hemorrhage.

### **Charcot-Leyden Crystals**

Slender, microscopic hexagonal bipyramidal crystals composed of eosinophil lysophospholipase (galectin-10) protein. They are found in sputum, stool, tissues, and bodily fluids in conditions characterized by eosinophilic inflammation, notably bronchial asthma, allergic rhinitis, and invasive parasitic infestations.

#### **Alternate Names:**

Charcot-Leyden Crystal; Galectin-10 Crystals

### **Charcot-Marie-Tooth Disease**

A group of inherited peripheral-nerve disorders that damage nerves supplying the feet, legs, hands, and arms. Symptoms usually include gradually worsening weakness, muscle wasting, numbness, foot deformity, and difficulty walking. It is genetically heterogeneous and may follow autosomal dominant, autosomal recessive, or X-linked inheritance.

#### **Alternate Names:**

CMT; Hereditary Motor and Sensory Neuropathy (HMSN); Peroneal Muscular Atrophy

### **Charcot-Wilbrand Syndrome**

A rare neuropsychiatric syndrome after brain injury characterized by loss of visual imagery and visual dreaming, sometimes with difficulty recognizing or interpreting visual information.

### **Chargaff Rule**

The observation that double-stranded DNA contains approximately equal amounts of adenine and thymine and of guanine and cytosine, reflecting complementary base pairing.

### **CHARGE Syndrome**

A genetic condition, usually involving a new change in the CHD7 gene, that can affect the eyes, ears, heart, breathing passages, growth, development, and reproductive system. Features vary widely.

### **Charles Bonnet Syndrome**

Visual hallucinations occurring in a person with significant vision loss who otherwise understands that the images are not real. The images may be simple patterns or detailed scenes; new hallucinations still require medical assessment.

### **Charley Horse**

A sudden painful muscle cramp, commonly in the calf or thigh. It may follow exercise, dehydration, prolonged positioning, nerve problems, or certain medicines and usually lasts from seconds to minutes.

### **Chasing The Dragon**

Inhalation of vapor produced by heating heroin, a method of drug use that can cause dependence, overdose, toxic exposures, and neurological injury.

### **Chaulmoogra Oil**

Chaulmoogra oil is a historical plant oil once used to treat leprosy and some skin diseases; it has largely been replaced by effective antimicrobial therapy.

### **CHD**

Alternate index label; see *Congenital Heart Disease*.

### **Checkpoint Inhibitor**

A class of monoclonal antibody immunotherapies that block negative inhibitory regulators of T-cell activation (such as PD-1, PD-L1, or CTLA-4), thereby relieving immune suppression and unleashing anti-tumor cytotoxic T-lymphocyte responses.

#### **Alternate Names:**

Immune Checkpoint Inhibitor; ICI

### **Checkpoints**

Regulatory signals that help prevent immune cells from attacking healthy tissues. Some cancers use checkpoint pathways to evade immunity; checkpoint-



---

inhibitor medicines block selected signals to restore antitumor immune activity.

### **Chediak-Higashi Syndrome**

A rare inherited immune disorder caused by changes in both copies of the *LYST* gene. It can cause unusually light or silvery hair, repeated infections, easy bruising, nerve problems, and a severe inflammatory crisis.

### **Cheek**

The facial region forming the side wall of the mouth, containing skin, muscle, fat, salivary structures, nerves, and blood vessels.

### **Cheilectropion**

An eversion deformity of the lip, usually involving the lower lip. It may result from scarring, congenital malformation, or loss of tissue support.

### **Cheilitis**

Inflammation of one or both lips causing dryness, scaling, cracks, swelling, pain, or sores. Causes include irritation, allergy, infection, nutritional deficiency, sun exposure, and skin disease; persistent or unusual changes need assessment.

### **Cheilitis Glandularis**

A rare inflammatory disorder of minor salivary glands in the lips, usually the lower lip, causing enlarged gland openings, swelling, mucus or saliva discharge, and sometimes eversion of the lip. Persistent lesions need biopsy or specialist assessment because chronic inflammation can mimic or coexist with neoplasia.

### **Cheilitis Granulomatosa**

Persistent swelling of one or both lips caused by granulomatous inflammation. It may occur alone or as part of Melkersson–Rosenthal syndrome or Crohn disease; recurrent swelling, facial palsy, or a fissured tongue requires evaluation.

### **Cheilodynia**

Pain of the lips, which may result from inflammation,

infection, trauma, allergy, dryness, nerve pain, or another oral or dermatologic disorder.

### **Cheilognathopalatoschisis**

A birth defect in which the upper lip, the gum-bearing part of the jaw, and the roof of the mouth do not join completely before birth. It can interfere with feeding, speech, hearing, and tooth development.

### **Cheilognathoschisis**

A congenital cleft of the lip and alveolar part of the upper jaw, with or without associated palatal involvement.

### **Cheiloplasty**

Plastic or reconstructive surgery of the lip, performed to repair congenital clefting, trauma, tumors, scars, or other deformity.

### **Cheiloschisis**

Cleft lip, a congenital fissure of the upper lip caused by incomplete fusion of facial processes during embryonic development. It may occur alone or with cleft palate.

### **Cheilosis**

A disorder of the lips characterized by dryness, scaling, fissuring, or inflammation. Causes include irritation, nutritional deficiency, infection, and systemic disease.

### **Cheiromegaly**

Abnormal enlargement of one or both hands, classically associated with excess growth hormone but also possible with local overgrowth or other disorders.

### **Cheiopompholyx**

A type of hand eczema that causes intensely itchy, small fluid-filled blisters on the palms or sides of the fingers. The skin may later peel, crack, or become painful.

---

## Chelating Agent

A compound that binds a metal ion through multiple chemical sites, allowing the metal to be measured, mobilized, or removed in selected toxic exposures.

## Chelating Agents

Chelating agents bind specific metal ions to form complexes that can facilitate their removal or reduce toxicity; examples include agents for lead, iron, or copper poisoning.

## Chelation

The formation of a stable complex between a metal ion and a molecule with multiple binding sites. In medicine, chelation can be used under supervision to bind and promote elimination of selected toxic metals.

## Chelation Study

A test that uses a metal-binding substance and imaging or blood measurements to assess how the body handles a mineral or to locate abnormal deposits. The exact method depends on the clinical question.

## Chelation Therapy

Treatment using a medicine that binds a specific harmful or excess metal so the body can remove it in urine or stool. It is used only for selected poisonings or disorders because it can harm the kidneys or alter essential minerals.

## Chemexfoliation

Application of a chemical solution to remove controlled layers of skin, improving selected scars, pigmentation, wrinkles, or superficial lesions.

## Chemical

A substance defined by its molecular composition and properties; medical effects depend on dose, route, exposure duration, metabolism, and biological target.

## Chemical Ablation

Destruction of unwanted tissue by injecting or applying a chemical irritant or sclerosant, used in selected

lesions, vessels, cysts, or other conditions.

## Chemical Burn

Damage to skin, eyes, or internal tissues caused by a corrosive chemical. Injury may worsen while the chemical remains present; contaminated clothing should be removed and the area flushed promptly with plenty of water.

## Chemical Burn Or Reaction

Tissue injury caused by contact with a corrosive or irritating chemical, commonly affecting the skin or eyes; severity depends on the substance, concentration, contact time, and tissue involved.

## Chemical Burns

Tissue injury caused by contact with corrosive acids, alkalis, oxidizers, or other chemicals. Damage may continue until the substance is removed and can involve the skin, eyes, airway, or gastrointestinal tract.

## Chemical Carcinogenesis

Development of cancer after exposure to chemicals that cause DNA damage, abnormal cell signaling, chronic tissue injury, or altered gene regulation.

## Chemical Cleavage

Breaking a chemical bond with a reagent, used in laboratory analysis, molecular biology, synthesis, or controlled degradation of a larger molecule.

## Chemical Cofactor

A nonprotein chemical helper required for an enzyme or biologic reaction, such as a metal ion, vitamin-derived coenzyme, or other small molecule.

## Chemical Eye Injury

Damage to the eye from an acidic or alkaline substance. Rinse the eye immediately with plenty of clean water or saline, remove contact lenses if easy, and obtain urgent eye care because especially alkaline chemicals can keep penetrating tissue.

---

## Chemical Fractionation

Separation of a mixture into fractions according to chemical or physical properties such as solubility, charge, size, or binding affinity.

## Chemical Impurity

An unintended chemical present in a medicine, specimen, or material, potentially affecting safety, stability, measurement accuracy, or biologic activity.

## Chemical Injury

Tissue damage caused by contact with a corrosive or irritating chemical. Acids and alkalis can destroy tissue differently, and immediate thorough rinsing, removal of contaminated clothing, and medical assessment may be needed.

## Chemical Peel

Application of a chemical solution to the skin to produce controlled exfoliation and regeneration of superficial or deeper skin layers.

### Alternate Names:

Chemexfoliation; Chemical Resurfacing

## Chemical Peels

Treatments that apply a chemical solution to remove a controlled amount of skin. Superficial peels affect the outer layer; deeper peels reach the underlying dermis and have greater risks of scarring, pigment changes, infection, and prolonged healing.

## Chemical Pleurodesis

A therapeutic procedure in which a sclerosing chemical agent (such as sterile talc slurry, doxycycline, or bleomycin) is instilled into the pleural space via a chest tube to induce intense parietal and visceral pleural inflammation, obliterating the pleural cavity to prevent recurrent pneumothorax or malignant pleural effusion.

## Chemical Pneumonitis

Lung inflammation and injury after inhaling or breathing in irritating chemicals, stomach contents, fuels, or other toxic substances. It may cause cough, shortness of breath, wheezing, low oxygen, and chest

imaging changes.

## Chemistry

The study of substances, their composition, properties, reactions, and transformations, forming the basis of biochemistry, pharmacology, toxicology, and laboratory medicine.

## Chemo

Informal shorthand for chemotherapy, treatment using drugs that destroy cancer cells, inhibit their growth, or suppress abnormal immune cells.

## Chemo Brain

Problems with memory, concentration, word-finding, or mental processing that may occur during or after cancer treatment. Symptoms can affect work and daily activities and may have several contributing causes.

### Alternate Names:

Post-Chemotherapy Cognitive Impairment

## Chemoattractant

A chemical signal that guides cells, especially immune cells, toward its source. Such signals help direct cells to areas of infection, inflammation, tissue repair, or normal development.

## Chemodectoma

A usually slow-growing tumour arising from specialized hormone-sensing nerve tissue, most often near the carotid artery in the neck. It may cause a neck lump or press on nearby nerves and blood vessels.

## Chemoembolization

A treatment that delivers cancer medicine through an artery feeding a tumour and then blocks that artery. It is used mainly for selected liver cancers to concentrate treatment and reduce the tumour's blood supply.

## Chemokine

A small signaling protein that directs immune cells to sites of infection, inflammation, or tissue repair. Different chemokines attract different cell types by

---

binding to matching receptors.

### **Chemokine Receptor**

A cell-surface receptor that binds chemokines and directs immune-cell movement; some also serve as entry cofactors for viruses such as HIV.

### **Chemokinesis**

Increased or altered movement of a cell in response to a chemical, without movement specifically toward or away from the chemical source. It differs from chemotaxis, which is directional.

### **Chemoprevention**

The use of a medicine or other chemical substance to lower the chance of cancer developing or returning. It may be considered for selected high-risk people, but benefits, side effects, and evidence differ by cancer and medicine.

### **Chemoradiation**

Treatment that gives chemotherapy and radiation during the same treatment period. The medicine may make cancer cells more sensitive to radiation and may treat cancer elsewhere, but combined side effects can be greater.

### **Chemoradiotherapy**

Cancer treatment that combines chemotherapy with radiation treatment. The medicines may make cancer cells more sensitive to radiation or treat cells elsewhere, but combined treatment can increase tiredness, nausea, skin changes, and other side effects.

### **Chemoreceptor**

A sensor that detects chemical changes such as oxygen, carbon dioxide, or acidity in the blood or surrounding fluid. These sensors help regulate breathing, heart activity, and other automatic body responses.

### **Chemoreceptor Cell**

A sensory cell that detects chemical changes such as oxygen, carbon dioxide, pH, taste molecules, or

odorants and converts them into neural or physiologic signals.

### **Chemoreceptor Reflex**

An autonomic homeostatic reflex initiated by peripheral (carotid and aortic bodies) or central (medullary) chemoreceptors in response to hypoxia, hypercapnia, or acidosis, driving increases in minute ventilation, sympathetic vasomotor tone, and blood pressure.

### **Chemoreceptor Trigger Zone**

A brain area that detects vomiting-provoking chemicals in the blood or cerebrospinal fluid and activates brain circuits that produce nausea and vomiting.

### **Chemoreceptors**

Chemoreceptors detect chemical changes such as oxygen, carbon dioxide, pH, or odor molecules and trigger neural or physiologic responses.

### **Chemoresistance**

Reduced response of cancer cells or other target cells to a medicine that previously inhibited or destroyed them. It may be present from the start or develop during treatment.

### **Chemosenitivity**

How strongly cells or a tumor respond to a chemical treatment. A sensitive tumor may shrink or stop growing, but laboratory sensitivity tests do not always predict a person's treatment response.

### **Chemosenitization**

Making cancer cells or microorganisms more vulnerable to treatment by adding an agent that improves delivery, blocks resistance, or increases treatment-related damage.

### **Chemosis**

Swelling of the clear membrane covering the white part of the eye. It may result from allergy, infection, irritation, injury, surgery, or fluid retention and can make the eye look blistered.

---

## Chemosurgery

Chemosurgery uses a chemical agent to destroy selected abnormal tissue, historically including chemical treatment of skin lesions; modern use is specialized.

## Chemotactic Factors

Chemical signals that attract or repel cells by forming a concentration gradient. During infection or tissue injury, they help direct immune cells to the affected area.

## Chemotaxis

Directed movement of a cell or organism toward or away from a chemical stimulus. It is important in immune-cell recruitment and inflammatory responses.

## Chemotherapy

Treatment using medicines to kill cancer cells or slow their growth, often by disrupting cell division or damaging DNA. It may be used alone or with surgery or radiation. Healthy fast-growing cells can also be affected, causing nausea, hair loss, anemia, or reduced immunity.

## Chemotherapy Extravasation

Leakage of an intravenous cancer medicine into tissue around the vein. Some medicines can cause severe pain, blistering, or tissue death; the infusion must stop immediately and the affected area needs urgent protocol-based care.

## Chemotherapy Regimen

A specific combination of anticancer drugs, dosing schedule, and treatment cycles designed for a particular cancer type and stage. Regimens are standardized by evidence-based protocols and may be modified based on response, toxicity, and patient factors.

## Chemotherapy Treatment

Chemotherapy uses medicines that damage or inhibit rapidly dividing cancer cells. It may be given before or after local treatment, for advanced disease, or for symptom control; effects and monitoring

depend on the drugs, cancer, treatment goal, and organ function.

## Chemotherapy-Induced Nausea and Vomiting

A common and debilitating adverse effect of antineoplastic therapy mediated by central neurokinin-1 (NK1) and peripheral/central serotonin (5-HT<sub>3</sub>) receptor pathways, categorized as acute, delayed, anticipatory, or breakthrough emesis.

**Alternate Names:**

CINV

## Chemotherapy-Induced Oral Mucositis

Painful inflammation and ulceration of the mouth caused by injury to rapidly dividing oral lining cells during chemotherapy or some cancer treatments. It can impair eating, drinking, and speech and increase infection risk; oral care, pain control, nutrition, and treatment of infection are important.

## Chemotherapy-Induced Peripheral Neuropathy

Nerve damage caused by some cancer medicines. It commonly produces tingling, numbness, burning pain, or unusual sensitivity in the hands and feet and may affect balance, movement, or daily activities.

## Cherry Angioma

A common harmless growth made of small blood vessels, usually a bright-red or purple spot or bump on the trunk or limbs. It may bleed after injury but usually needs no treatment unless the diagnosis is uncertain or removal is wanted.

## Cherry Eye

Prolapse of the gland of the third eyelid, producing a rounded red mass at the inner corner of the eye. The term is used especially in veterinary ophthalmology.

## Cherry-Red Spot

A bright-red area at the center of the retina surrounded by pale retina. It occurs when the surrounding retina becomes swollen or loses blood flow and may indicate a serious metabolic, vascular, or storage disorder.

---

## **Cherubism**

A rare inherited condition in which both sides of the jaw enlarge because of overgrowth of bone-removing tissue. It usually begins in childhood and often improves after puberty, but it can affect teeth, chewing, speech, or appearance.

## **Chest**

The thorax, the region between the neck and abdomen containing the heart, lungs, major vessels, and other structures protected by the ribs and sternum.

## **Chest Cold**

Alternate name; see *Acute Bronchitis*.

## **Chest CT**

Computed tomography of the chest using x-rays and computer reconstruction to produce cross-sectional images of the lungs, heart, blood vessels, mediastinum, pleura, and chest wall.

## **Chest Infection**

An infection of the airways or lung, commonly caused by a virus or bacterium. It may cause cough, mucus, fever, wheezing, chest discomfort, or shortness of breath; severe breathing difficulty, confusion, blue lips, or persistent high fever requires urgent care.

## **Chest MRI**

Magnetic resonance imaging of the chest using a magnetic field and radio waves to evaluate soft tissues, blood vessels, heart structures, mediastinum, chest wall, and selected lung or pleural abnormalities.

## **Chest Pain**

Pain, pressure, burning, or discomfort in the chest. Causes include heart, lung, digestive, muscle, bone, or anxiety-related problems. New severe pain, breathlessness, sweating, faintness, or spreading pain requires emergency care.

### **Alternate Names:**

Angina Pectoris; Thoracalgia; Chest Discomfort

## **Chest Physiotherapy**

Techniques that help move mucus out of the lungs, including positioning, gentle percussion, breathing exercises, coughing techniques, and selected breathing devices. It may help some people with bronchiectasis, cystic fibrosis, or recovery after surgery.

## **Chest Tightness**

A sensation of pressure, constriction, or heaviness in the chest that may arise from heart, lung, gastrointestinal, musculoskeletal, or anxiety-related causes.

## **Chest Tube**

A flexible tube inserted into the pleural space to remove air, blood, or fluid and help the lung re-expand.

## **Chest Tube Insertion**

Placement of a flexible tube through the chest wall into the space around a lung to remove air, blood, pus, or fluid and help the lung expand. It is connected to a drainage system and requires monitoring for bleeding, infection, or persistent air leak.

## **Chest Tube Thoracostomy**

Placement of a tube through the chest wall into the space around a lung to drain air, blood, pus, or fluid. It may be needed for a collapsed lung, bleeding, infected fluid, or certain problems after chest or heart surgery.

## **Chest Wall**

The layers of skin, muscle, ribs, breastbone, spine, and supporting tissues that surround the chest. They protect the heart and lungs and move with the breathing muscles to help expand and contract the chest.

## **Chest Wall Deformity**

An abnormal shape of the ribs, breastbone, spine, or surrounding tissues. A sunken or protruding breastbone is often mainly cosmetic, but severe deformity or scoliosis can restrict breathing or affect the heart.



---

## Chest Wall Pain

Alternate index label; see *Costochondritis*.

## Chest X-Ray

An imaging test using a small amount of radiation to show the lungs, heart, ribs, and chest structures. It can detect pneumonia, collapsed lung, fluid, some fractures, and changes in heart size.

## Chewing

The coordinated use of teeth, jaw muscles, and joints to break food into smaller pieces for swallowing and digestion; pain or missing teeth can interfere.

## Chewing Tobacco

Alternate name; see *Smokeless tobacco*.

## Cheyne-Stokes Respiration

A repeating pattern in which breathing gradually becomes deeper and faster, then shallower, followed by a pause. It may occur during sleep with heart failure, stroke, brain disease, or at high altitude.

## CHF

Heart failure in which the heart cannot pump or fill well enough for the body's needs. Reduced circulation can cause breathlessness, tiredness, leg swelling, and fluid buildup in the lungs or elsewhere.

## CHH

Abbreviation for congenital or central hypogonadotropic hypogonadism, in which deficient hypothalamic or pituitary signaling leads to low sex-hormone production and absent or incomplete puberty. Some forms also cause reduced or absent sense of smell.

## Chiari I Malformation

A condition in which the lower part of the cerebellum extends through the opening at the skull base. It may cause headaches worsened by straining, neck pain, balance problems, or no symptoms.

## Chiari Malformation

A structural abnormality in which part of the cere-

bellum extends downward through the opening at the base of the skull. Different types vary in severity and may cause headache, balance problems, swallowing difficulty, or no symptoms.

## Chiari-Frommel Syndrome

Continued milk production and absent menstrual periods after pregnancy, usually caused by persistently high prolactin levels. Evaluation looks for pituitary disorders, medicines, thyroid disease, and other causes.

## Chicken Soup And Sickness

A supportive-care topic concerning the use of warm fluids and nutrition during illness; chicken soup may ease hydration or congestion but does not replace diagnosis or treatment.

## Chickenpox

An acute, highly contagious primary infection with varicella-zoster virus. An itchy generalized rash appears in successive crops as spots, blisters, and crusts, often with fever and malaise; adults, pregnant people, newborns, and immunocompromised people have higher risk of pneumonia or other severe complications.

## Chickenpox (Varicella)

A contagious infection caused by varicella-zoster virus, producing an itchy blistering rash, fever, and malaise. Complications include bacterial skin infection, pneumonia, cerebellitis, and severe disease in pregnancy or immunocompromise.

## Chief Cell

A cell in the stomach lining that makes pepsinogen, an inactive substance that becomes the protein-digesting enzyme pepsin in stomach acid.

## Chigger Bites

Itchy skin reactions caused by larval mites. Small red bumps or blisters often develop where clothing is tight and may itch for days. Washing the skin, avoiding scratching, and itch treatment usually provide relief.

---

## Chiggers

Larval harvest mites whose bite causes intensely itchy grouped red papules or welts; symptoms result from skin irritation, not from the mite burrowing into human tissue.

## Chignon

Temporary swelling of a baby's scalp where a vacuum device was applied during assisted delivery. It results from the suction and usually disappears within a day or two without treatment.

## CHIK

Alternate name; see *Chikungunya*.

## Chikungunya

A mosquito-borne viral disease caused by chikungunya virus, usually producing abrupt fever and severe, often symmetric joint pain. Headache, muscle pain, joint swelling, nausea, fatigue, and rash may occur; joint symptoms can persist for months or years.

### Alternate Names:

Chikungunya Fever; Chikungunya Virus Disease; CHIKV Infection

## Chikungunya Fever

An acute arboviral febrile illness caused by the Chikungunya virus (genus *Alphavirus*) transmitted by *Aedes* mosquitoes, characterized by high fever, maculopapular rash, and severe, often incapacitating bilateral polyarthralgia and polyarthritis that may persist chronically.

## Chikungunya Virus

An alphavirus transmitted primarily by infected female *Aedes aegypti* and *Aedes albopictus* mosquitoes. Infection can cause abrupt fever, severe polyarthralgia, joint swelling, rash, and prolonged inflammatory joint symptoms.

## Chilaiditi Syndrome

The symptomatic clinical presentation of Chilaiditi sign, defined as the anatomical interposition of a loop of colon or small intestine between the superior

surface of the liver and the right hemidiaphragm. Manifests with intermittent abdominal pain, constipation, nausea, vomiting, or abdominal distention, occasionally mimicking a perforated hollow viscus with pseudo-pneumoperitoneum on abdominal radiography.

### Alternate Names:

Hepatodiaphragmatic Interposition Syndrome

## Chilblain

An inflammatory skin lesion caused by an abnormal response to cold and damp conditions. It produces itchy, red or violaceous swollen areas, usually on fingers, toes, ears, or nose.

## Chilblains

Itchy, painful, red or purple swellings caused by an abnormal skin response after exposure to cold and damp conditions. They usually improve with warmth but can blister, ulcerate, or become infected.

### Alternate Names:

Pernio

## Child Abuse

Physical, sexual, or emotional harm, or serious neglect, inflicted on a child by a caregiver or another person. Warning signs may include unexplained injuries, changing accounts, fear, developmental changes, or unmet basic needs; suspected abuse requires safeguarding action under local law.

## Child Adoption

Child adoption is the legal process by which parental rights and caregiving responsibility are transferred to adoptive parents; medical care includes records, consent, and family history.

## Child Behavior

The observable actions, responses, and conduct of children, influenced by developmental stage, temperament, environment, and neurobiological factors. Behavioral assessment helps identify developmental disorders, emotional difficulties, and parenting needs.

---

## Child Development

Progressive physical, cognitive, language, emotional, and social changes from infancy through adolescence, assessed against developmental milestones and individual variation.

## Child Health

The physical, developmental, mental, and social well-being of children, including prevention, nutrition, immunization, safety, and treatment of illness.

## Child Health Services

Child health services provide preventive care, immunization, growth and developmental monitoring, screening, treatment, and support for children and families.

## Child Maltreatment

Abuse or neglect of a child, including physical, sexual, or emotional abuse and failure to provide needed care or supervision. It can harm a child's safety, health, growth, and development.

### Alternate Names:

Child Abuse

## CHILD Naevus

A congenital skin and limb disorder named for congenital hemidysplasia with ichthyosiform erythroderma and limb defects. It causes one-sided scaly skin changes with asymmetrical limb or skeletal abnormalities and may involve other organs.

## Child Neglect

Failure to provide a child with adequate food, shelter, clothing, supervision, education, emotional care, or medical care. It can impair growth, development, safety, and health and requires safeguarding assessment when suspected.

## Child Neglect And Emotional Abuse

Failure to provide a child's basic needs or repeated behavior that harms emotional development, safety, or wellbeing; suspected abuse requires safeguarding assessment and appropriate reporting under local law.

## Child Physical Abuse

Intentional physical injury or harm inflicted on a child by a caregiver or another person, including hitting, burning, shaking, or fractures; suspected abuse requires immediate safeguarding assessment.

## Child Poisoning

Exposure of a child to a medicine, chemical, plant, or other substance that may cause harm. Symptoms can be delayed, so suspected poisoning requires immediate advice from a poison service or emergency clinician rather than waiting for symptoms.

## Child Psychiatry

The medical specialty concerned with assessment and treatment of emotional, behavioral, developmental, and psychiatric disorders in children and adolescents.

## Child Psychology

The study of children's cognitive, emotional, social, and behavioral development and the effects of relationships, learning, biology, and environment.

## CHILD Syndrome

Congenital hemidysplasia with ichthyosiform nevus and limb defects, a rare disorder usually caused by an *NSDHL* variant. It causes one-sided skin abnormalities with limb, skeletal, and sometimes neurologic or internal-organ differences.

## Child-Pugh Classification

A scoring system used to estimate the severity of chronic liver disease. It combines blood bilirubin, albumin, and clotting results with fluid in the abdomen and liver-related confusion; the score supports, but does not replace, individual assessment.

## Child-Pugh Score

A widely used clinical staging system assessing the prognosis, surgical mortality, and necessity of liver transplantation in patients with cirrhosis, incorporating total bilirubin, serum albumin, prothrombin time/INR, ascites severity, and hepatic encephalopathy grade.

---

**Alternate Names:**

Child-Turcotte-Pugh Score; CTP Score

**Childbed Fever**

An older name for fever caused by infection after childbirth, miscarriage, or abortion. The infection often affects the uterus and may cause lower abdominal pain, foul-smelling discharge, or worsening illness requiring prompt treatment.

**Childbirth**

The process of giving birth. Labour usually includes cervical opening, birth of the baby, and delivery of the placenta; timing and the need for medical assistance vary between pregnancies.

**Childhood**

The developmental period between infancy and adolescence, during which physical growth, brain development, learning, and social skills progress.

**Childhood Absence Epilepsy**

An epilepsy syndrome that usually begins in school-aged children with frequent brief pauses in awareness. During a seizure, the child may stare and stop responding for seconds, then quickly return to normal without confusion.

**Childhood ADHD**

Attention-deficit/hyperactivity disorder beginning in childhood, with persistent developmentally inappropriate inattention, hyperactivity, or impulsivity that causes impairment across one or more settings.

**Childhood Angiosarcoma**

A rare malignant tumor arising from blood-vessel or lymphatic-endothelial cells in a child; diagnosis and treatment require specialist pediatric oncology care.

**Childhood Apraxia Of Speech**

A speech-sound disorder in which a child has difficulty planning and coordinating the movements needed for clear speech.

**Alternate Names:**

CAS

**Childhood Asthma**

Asthma beginning during childhood, involving variable airway narrowing and inflammation that cause wheezing, cough, chest tightness, or shortness of breath.

**Childhood Cataracts**

Cataracts occurring during infancy, childhood, or adolescence. They may be inherited, present at birth, caused by injury or inflammation, or associated with another condition, and can interfere with visual development.

**Childhood Cyclic Vomiting**

Recurrent stereotyped episodes of intense nausea and vomiting separated by periods of normal health. It is a functional gut-brain disorder diagnosed after other causes are considered and may be associated with migraine.

**Childhood Depression**

Depressive illness in a child, which may present as persistent sadness, irritability, loss of interest, withdrawal, sleep or appetite change, low energy, poor concentration, or physical complaints. Suicidal thoughts require immediate emergency help.

**Childhood Disintegrative Disorder**

A rare neurodevelopmental condition characterized by a marked regression and profound loss of previously acquired motor, language, social, cognitive, and adaptive skills after at least two years of apparently normal development.

**Alternate Names:**

CDD; Heller's Syndrome; Heller's Dementia; Disintegrative Psychosis

**Childhood Ependymoma**

A tumor arising from ependymal cells lining the brain ventricles or spinal canal in children; symptoms depend on location and obstruction of cerebrospinal-fluid flow.

---

### Childhood Fibrosarcoma

A malignant tumor of fibroblastic connective tissue in children, usually presenting as a growing mass; treatment depends on location, extent, and tumor biology.

### Childhood Ganglioglioma

A usually slow-growing mixed neuronal and glial brain tumor in children or young adults, often presenting with long-standing focal seizures.

### Childhood Giant Axonal Neuropathy

The childhood-onset form of giant axonal neuropathy, a rare inherited neurodegenerative disorder causing progressive peripheral neuropathy, ataxia, weakness, developmental or cognitive changes, and characteristic tightly curled hair in many affected people.

### Childhood Interstitial Lung Disease

A group of rare lung disorders in infants and children that affect the tissue around the air sacs and may interfere with oxygen exchange. They can cause rapid breathing, cough, low oxygen, poor growth, or limited exercise ability.

### Childhood Leiomyosarcoma

A rare malignant tumor of smooth muscle in a child; it may arise in soft tissue or an organ and requires biopsy, staging, and specialist treatment.

### Childhood Leukemia

Cancer of blood-forming cells in a child, most often acute lymphoblastic or acute myeloid leukemia; symptoms may include pallor, bruising, fever, bone pain, or recurrent infection.

### Childhood Liposarcoma

A rare malignant tumor derived from fat-forming tissue in a child; diagnosis requires imaging and biopsy, and treatment is individualized by a sarcoma team.

### Childhood Muscular Dystrophy

A group of inherited disorders causing progressive

muscle weakness and wasting beginning in childhood. Different types vary in inheritance, muscles affected, heart and breathing involvement, and rate of progression.

### Childhood Obesity

Alternate index label; see *Pediatric Obesity*.

### Childhood Osteosarcoma

A malignant bone tumor that commonly develops near the ends of long bones during adolescence; pain, swelling, or a fracture may occur and treatment combines chemotherapy and surgery.

### Childhood Rhabdomyosarcoma

A malignant tumor showing skeletal-muscle differentiation, occurring mainly in children and adolescents; it may arise in the head and neck, urinary or reproductive tract, or limbs.

### Childhood Schizophrenia

A rare psychotic disorder beginning in childhood, involving disturbances in thinking, perception, behavior, and emotional functioning.

#### Alternate Names:

Very Early-Onset Schizophrenia (VEOS); Early-Onset Schizophrenia

### Children

Humans in the developmental period between infancy and adolescence; medical assessment must account for age-specific anatomy, physiology, growth, communication, and consent.

### Children And Grief

The emotional and behavioral response of children to death or major loss, which varies by developmental stage and may include sadness, regression, anger, worry, or physical symptoms.

### Children's Cancer Centers

Specialized services that coordinate diagnosis, treatment, rehabilitation, psychosocial support, survivorship care, and research for children and adolescents with cancer.

---

**Alternate Names:**

Pediatric Oncology Centers; Childhood Cancer Clinics

**Chill**

A chill is a sensation of cold, often with shivering, that can accompany fever, infection, exposure, anxiety, or other illness.

**Chills**

A sensation of cold with involuntary shivering, commonly accompanying fever or infection; chills without measured fever may also occur with anxiety, hypoglycaemia, or environmental cold.

**Chilomastix Mesnili**

A usually harmless single-celled organism that may be found in stool. Its presence generally does not cause illness or require treatment, but it can indicate exposure to contaminated food or water.

**Chimera**

An organism or tissue containing genetically distinct cell populations derived from different zygotes or embryos, or a laboratory construct combining genetic material from different sources.

**Chimeric Antigen Receptor**

A synthetic receptor engineered to recognize a selected target on a cell surface. It is placed into immune cells, especially T cells, to direct them against cancer cells in some cellular immunotherapies.

**Chimeric Antigen Receptor T-Cell Therapy**

A cancer treatment in which a person's T cells are collected, genetically changed in a laboratory to recognize a cancer marker, multiplied, and returned to the body. Treatment can cause serious inflammation, infection risk, or neurologic effects.

**Chimerism**

The presence in one person of groups of cells with different genetic origins. It may occur naturally, after an organ or stem-cell transplant, or when cells

pass between a pregnant person and fetus.

**Chin**

The lower facial prominence formed mainly by the mandible and overlying soft tissue, contributing to facial contour, speech, chewing, and airway anatomy.

**Chin Augmentation**

A cosmetic procedure that increases chin projection with an implant, injectable filler, or bone surgery to change facial contour; risks include infection, asymmetry, nerve injury, and implant problems.

**Chinese Herbal Medicine**

The use of plant, mineral, or animal-derived preparations within traditional Chinese medicine. Products vary in composition and may interact with medicines or contain contaminants, so reliable sourcing and medical guidance are important.

**Chinese Restaurant Syndrome**

An older, controversial term for short-lived symptoms such as headache, flushing, or palpitations attributed to eating restaurant food; evidence does not show that monosodium glutamate causes these symptoms in most people at usual dietary exposures.

**Chionablepsia**

Temporary eye pain and blurred vision caused by ultraviolet light reflected from snow or ice, commonly called snow blindness. Symptoms may appear hours after exposure and usually improve as the cornea heals.

**Alternate Names:**

Snow Blindness; Photokeratitis

**CHIP**

Clonal hematopoiesis of indeterminate potential: an age-related finding in which blood-forming cells acquire a gene change and expand without meeting criteria for a blood cancer. It can increase the future risk of myeloid cancer and cardiovascular disease and may require monitoring.



---

## Chiropody

Chiropody is care of the feet, including assessment and treatment of skin, nail, pressure, and biomechanical problems; it is also called podiatry.

## Chiropractic

A healthcare practice that mainly uses hands-on treatment for muscle, joint, and spine problems, especially spinal manipulation. Benefits and risks vary by the condition, technique, and practitioner's training.

## Chiropractic Care For Back Pain

Manual therapy, primarily spinal manipulation and mobilization, provided for acute and chronic low back pain when indicated. Evidence supports its use for acute low back pain in particular, though it is not appropriate for all causes of back pain.

## Chiropractor

A healthcare practitioner who provides manual or manipulative treatment, especially for musculoskeletal conditions, according to the practitioner's training and local regulation.

### Alternate Names:

Chiropractic Practitioner; Chiropractic Clinician

## Chitin

A strong natural substance forming the cell walls of fungi and the outer coverings of insects, spiders, and shellfish. Humans do not digest it, but exposure to chitin-containing material can contribute to allergic or airway irritation in some people.

## Chlamydia

In common clinical use, an infection with *Chlamydia trachomatis*, a sexually transmitted bacterium that often causes no symptoms but may cause discharge, painful urination, pelvic inflammatory disease, infertility, eye infection, or newborn infection. The word can also refer to the bacterial genus, which includes species causing respiratory or bird-associated infections.

## Chlamydia (Chlamydial Genitourinary

## Infections)

Genitourinary infection caused by *Chlamydia trachomatis*, often without symptoms. Untreated infection can cause cervicitis, pelvic inflammatory disease, ectopic pregnancy, infertility, or neonatal infection; testing, treatment, and partner management are important.

## Chlamydia Pneumoniae Infection

A respiratory infection caused by *Chlamydia pneumoniae*, formerly called *Chlamydophila pneumoniae*. It commonly causes gradual-onset atypical pneumonia, bronchitis, or upper-respiratory symptoms.

## Chlamydia Psittaci

A bacterium that causes psittacosis, a zoonotic infection usually acquired by inhaling organisms from infected birds.

## Chlamydia Trachomatis

A bacterium that commonly causes sexually transmitted infection of the urogenital tract and can also cause conjunctivitis or pneumonia.

## Chlamydia Trachomatis Infection

A bacterial infection spread mainly through sexual contact. It often causes no symptoms but may cause discharge or painful urination and can lead to pelvic inflammatory disease, infertility, eye infection, or newborn infection.

## Chlamydiaceae

A family of bacteria that includes *Chlamydia* species. Some cause sexually transmitted infections, eye disease, or respiratory infection; diagnosis depends on the species, specimen, symptoms, and laboratory testing.

## Chlamydial Pneumonia

A lung infection caused by *Chlamydia* bacteria, usually with gradually worsening cough, fever, tiredness, headache, or sore throat. Diagnosis and treatment depend on the species, severity, and other medical conditions.

---

### Chlamydial Pneumonias

Pneumonia caused by *Chlamydia* species, including *C. pneumoniae* and *C. psittaci*. Illness is often gradual with cough, fever, headache, and atypical chest findings; exposure history and targeted testing help distinguish it from other community-acquired pneumonias.

### Chlamydiosis

Infection caused by bacteria of the genus *Chlamydia*. Depending on the species and site, it may cause sexually transmitted infection, conjunctivitis, pneumonia, or other disease.

### Chloasma

Patchy brown or gray-brown darkening of the skin, usually on the face. It is often related to pregnancy, hormones, and sunlight and is commonly called melasma.

#### Alternate Names:

Melasma; Mask of Pregnancy

### Chloracne

An acne-like eruption caused by exposure to certain halogenated aromatic compounds, such as dioxins and related industrial chemicals.

### Chloral Hydrate

A sedative-hypnotic compound with limited modern use because of toxicity, dependence, respiratory depression, and cardiac or gastrointestinal adverse effects.

### Chlorambucil

An oral chemotherapy medicine that damages DNA and is used mainly for some chronic leukemias and lymphomas. It can lower blood counts and increase infection risk.

### Chloramphenicol

An antibiotic that stops bacteria from making essential proteins. It is reserved for selected serious infections when safer or more suitable medicines are unavailable because it can rarely cause dangerous bone-marrow suppression and other severe effects.

### Chloranemia

An older term for anemia in which red blood cells contain too little hemoglobin and appear unusually pale. Iron deficiency is a common cause; modern evaluation uses blood counts and iron studies.

### Chlordane

A persistent organochlorine insecticide formerly used for termite control; exposure can affect the nervous system and liver and has environmental persistence.

### Chlordecone

A persistent organochlorine pesticide that can cause neurologic, reproductive, and liver effects after significant exposure and remains in the environment for long periods.

### Chlordiazepoxide

A benzodiazepine medicine that slows brain activity and may be used short term for alcohol-withdrawal symptoms or severe anxiety. It can cause drowsiness, dependence, impaired coordination, and dangerous breathing suppression with alcohol or opioids.

### Chlordiazepoxide Overdose

Overdose of chlordiazepoxide, a benzodiazepine, that can cause drowsiness, impaired coordination, respiratory depression, and coma, especially with other sedatives.

### Chlorethoxyfos

An organophosphate insecticide that can block a nerve enzyme, causing sweating, vomiting, weakness, breathing problems, seizures, or other poisoning symptoms after exposure.

### Chlorfenvinphos

An organophosphate insecticide that can cause cholinergic toxicity by inhibiting acetylcholinesterase; use is restricted or prohibited in many places.

### Chlorhexidine

A broad-spectrum antiseptic used for skin, mucosal,

---

and oral preparation; it can cause irritation, allergy, or rare severe anaphylaxis.

### Chloride

A negatively charged mineral in blood and other body fluids. It works with sodium and other electrolytes to control fluid balance, nerve and muscle function, and the body's acid level.

### Chloride Channel

A protein forming a passage for chloride ions across a cell membrane. Chloride channels help control electrical signals, fluid movement, and muscle or nerve function.

### Chloride In Diet

Chloride consumed in food, usually as part of salt. It helps maintain fluid balance, nerve signals, and stomach acid; excessive intake often accompanies excess sodium.

### Chloride Ion

The negatively charged form of chlorine and a major extracellular electrolyte that helps maintain fluid balance, osmotic pressure, acid-base status, and electrical neutrality.

### Chloride Shift

An electrochemical exchange process in red blood cells mediated by the anion exchanger 1 (band 3 protein), wherein bicarbonate generated by carbonic anhydrase exits the erythrocyte in exchange for an incoming plasma chloride ion in systemic capillaries, reversed in pulmonary capillaries.

#### Alternate Names:

Hamburger Phenomenon

### Chlorinated Lime

Chlorinated lime, or bleaching powder, releases chlorine and is used as a disinfectant; concentrated exposure can burn skin, eyes, and airways.

### Chlorinated Lime Poisoning

Exposure to chlorinated lime, a hypochlorite-containing corrosive that can injure the skin, eyes,

airways, or gastrointestinal tract.

### Chlorine

A reactive halogen element and disinfectant; inhalation of chlorine gas can injure the eyes and airways and cause chemical pneumonitis or pulmonary edema.

### Chlorine Poisoning

Inhalation or contact exposure to chlorine gas, which irritates the eyes and airways and can cause cough, bronchospasm, chemical pneumonitis, or pulmonary edema.

### Chlormadinone Acetate

A synthetic progestogen used in some hormonal medicines; effects and risks include changes in bleeding, mood, metabolism, and thrombotic risk depending on the formulation.

### Chlormethine

Chlormethine, or mechlorethamine, is an alkylating anticancer drug used systemically or topically for selected cancers and cutaneous T-cell lymphoma.

### Chlorodeoxyadenosine

Another name for cladribine, a purine-analog medicine used for selected blood cancers and multiple sclerosis; prolonged lymphocyte suppression and infection risk are important considerations.

### Chloroform

A volatile solvent and environmental contaminant that can irritate the lungs and nervous system and damage the liver or kidneys after significant exposure.

### Chlorogenic Acid

A natural plant compound found in coffee, fruits, and vegetables. It has antioxidant effects in laboratory studies, but evidence is insufficient to use supplements to prevent or treat disease.

### Chloroleukemia

A rare form of myeloid leukaemia associated with

---

greenish masses of abnormal immature blood-forming cells outside the bone marrow. These masses are also called myeloid sarcoma or chloroma.

**Alternate Names:**

Granulocytic Sarcoma; Chloroma; Myeloid Sarcoma

**Chloroma**

A mass of immature myeloid blood-forming cells growing outside the bone marrow, usually associated with acute myeloid leukaemia. It may affect skin, bone, lymph nodes, or internal organs and is also called myeloid sarcoma.

**Alternate Names:**

Myeloid Sarcoma; Granulocytic Sarcoma; Extramedullary Myeloid Tumor

**Chlorophyll Poisoning**

Ingestion of chlorophyll-containing products, which usually causes mild gastrointestinal effects; product ingredients and dose determine clinical risk.

**Chlorophyllin**

A water-soluble, copper-containing derivative of chlorophyll used in some products and studied for effects such as odor control; clinical uses depend on evidence and context.

**Alternate Names:**

Chlorophyllin Copper Complex; Sodium Copper Chlorophyllin

**Chloropicrin**

A pungent soil fumigant and irritant that can injure the eyes, skin, and respiratory tract after exposure.

**Chloroprene**

An industrial chemical used to make chloroprene rubber; repeated or high exposure can irritate tissues and has carcinogenic concern.

**Chloropsia**

A visual disturbance in which objects appear green or greenish. It may result from retinal or visual-pathway disease, medicines, toxins, migraine, or other conditions and needs assessment if persistent or sudden.

**Chloroquine**

An antimalarial medicine used for susceptible Plasmodium species and selected autoimmune conditions; retinal toxicity, cardiotoxicity, and drug interactions require attention.

**Chlorothiazide**

A thiazide diuretic used to treat high blood pressure and fluid retention. It increases urine production and may cause low sodium, low potassium, dehydration, or high blood sugar.

**Chlorpheniramine**

A first-generation antihistamine used for allergy symptoms; drowsiness, dry mouth, blurred vision, urinary retention, and other anticholinergic effects may occur.

**Chlorphentermine**

An older stimulant appetite suppressant related to amphetamines; it is not routinely used because of cardiovascular, psychiatric, and dependence risks.

**Chlorpromazine**

A first-generation antipsychotic and phenothiazine used for schizophrenia, severe nausea, and selected other conditions; adverse effects include sedation, hypotension, movement disorders, and QT prolongation.

**Chlorpromazine Overdose**

Overdose of chlorpromazine that may cause sedation, hypotension, anticholinergic effects, seizures, cardiac conduction abnormalities, or neuroleptic malignant syndrome.

**Chlorpropamide**

A first-generation sulfonylurea that lowers blood glucose by stimulating pancreatic insulin release; prolonged hypoglycemia and hyponatremia are important risks.

**Chlorpyrifos**

An organophosphate insecticide that inhibits acetylcholinesterase; significant exposure can cause cholin-

---

ergic toxicity with secretions, bronchospasm, bradycardia, weakness, or seizures.

### **Chlortalidone**

Chlortalidone, also called chlorthalidone, is a long-acting thiazide-like diuretic used mainly for hypertension and fluid retention.

### **Chlorthalidone**

A long-acting thiazide-like diuretic used for hypertension and edema; it can cause hyponatremia, hypokalemia, hyperuricemia, and kidney-related electrolyte changes.

### **Chlorzoxazone**

A centrally acting muscle relaxant used for painful muscle spasm in some countries; drowsiness and rare serious liver injury are important risks.

### **CHO Cell**

A Chinese hamster ovary cell line widely used to produce recombinant proteins, antibodies, vaccines, and other biologic medicines.

### **Choanal Atresia**

A birth defect in which tissue blocks one or both passages from the nose to the throat. Both sides can cause severe breathing difficulty in a newborn, especially during feeding, and require urgent treatment.

### **Chocolate Cyst**

A chocolate cyst is an ovarian endometrioma filled with old blood from endometriosis, which can cause pelvic pain, infertility, or an adnexal mass.

### **Choice Behavior**

The process of selecting among alternatives, studied in psychology and neuroscience to understand decision-making, impulsivity, and reward processing. Abnormal choice behavior may be relevant to addiction, eating disorders, and psychiatric conditions.

### **Choke**

Airway blockage by food or another object that pre-

vents normal airflow and can rapidly cause inability to speak, cyanosis, loss of consciousness, or death.

### **Choking**

Blockage of the airway by food or another object, making breathing difficult or impossible. A person who cannot speak, cough effectively, or breathe needs immediate first aid and emergency medical help.

### **Choking First Aid for Adults and Children**

Immediate emergency management for complete foreign-body airway obstruction (FBAO) in conscious adults and children over 1 year of age, utilizing subdiaphragmatic abdominal thrusts (Heimlich maneuver) or alternating back blows and chest thrusts.

#### **Alternate Names:**

Heimlich Maneuver; Foreign Body Airway Obstruction Management; FBAO First Aid

### **Choking First Aid for an Unresponsive Person**

Advanced resuscitation algorithm for an unresponsive patient with foreign body airway obstruction, involving immediate initiation of cardiopulmonary resuscitation (CPR), visualization of the oral cavity prior to rescue breaths, and avoidance of blind finger sweeps.

#### **Alternate Names:**

Unconscious Choking Management; Obstructed Airway CPR; Foreign Body CPR

### **Choking First Aid for Infants**

Emergency rescue intervention for complete foreign-body airway obstruction in conscious infants under 1 year of age, consisting of 5 prone back blows delivered between the scapulae followed by 5 supine chest thrusts while firmly supporting the head.

#### **Alternate Names:**

Infant Choking Management; Infant Foreign Body Airway Relief; Infant FBAO

### **Cholagogue**

A substance that encourages the gallbladder to re-

---

lease bile into the small intestine. Bile helps the body digest fats; medicines or foods described as cholagogues should be used carefully when gallstones or blocked bile ducts are possible.

**Alternate Names:**

Gallbladder Stimulant; Biliary Evacuant

**Cholangiitis**

Inflammation of the tubes that carry bile from the liver and gallbladder to the intestine. Infection, a blockage, autoimmune disease, or another bile-duct disorder may cause it.

**Cholangiocarcinoma**

A malignant neoplasm originating from the epithelial cells lining the intrahepatic or extrahepatic biliary tract. It typically presents with painless obstructive jaundice, pruritus, pale acholic stools, dark tea-colored urine, right upper quadrant abdominal pain, and constitutional weight loss.

**Alternate Names:**

Bile Duct Cancer; Bile Duct Carcinoma; Cholangiocellular Carcinoma

**Cholangiography**

Imaging of the bile ducts after contrast material makes them visible. It can show gallstones, narrowing, leaks, or blockage and may be performed during surgery, through the skin, or during endoscopy.

**Cholangiohepatitis**

Inflammation of the bile ducts and liver, often caused by infection, blockage, parasites, or repeated bile-duct inflammation. It may cause fever, pain under the right ribs, jaundice, abnormal liver tests, and sometimes serious infection.

**Cholangitis**

Inflammation or infection of the tubes that carry bile from the liver and gallbladder to the intestine. A blockage from a gallstone or narrowing can cause fever, pain, jaundice, and rapidly worsening bloodstream infection.

**Cholate**

The ionized form of cholic acid, a primary bile acid that helps emulsify dietary fat and is conjugated and secreted by the liver.

**Cholecalciferol**

Vitamin D3, produced in skin by ultraviolet light and obtained from food or supplements, then converted to active metabolites that regulate calcium and phosphate.

**Cholecyst**

The gallbladder, a small sac beneath the liver that stores and concentrates bile. When food enters the intestine, it contracts and releases bile to help digest dietary fats.

**Cholecystectomy**

Surgery to remove the gallbladder, a small organ under the liver that stores bile. It may be needed for painful gallstones, gallbladder inflammation, gallstone-related pancreatitis, or selected polyps. It is often done through small cuts using a camera, but open surgery may be needed when this is not safe. Bile still flows from the liver into the intestine after surgery.

**Alternate Names:**

Gallbladder Removal; Laparoscopic Cholecystectomy; Lap Chole; Open Cholecystectomy

**Cholecystitis**

Inflammation of the gallbladder, usually caused by a gallstone persistently blocking the cystic duct. It may also occur without gallstones, especially in critically ill, injured, postoperative, fasting, or shocked patients. Acute cholecystitis commonly causes persistent right-upper-quadrant pain, tenderness or a positive Murphy sign, fever, nausea, and vomiting. Diagnosis typically combines clinical findings, blood tests, and imaging, usually ultrasound. Treatment may include supportive care, antibiotics when indicated, early laparoscopic cholecystectomy, or gallbladder drainage when surgery is unsafe.

**Alternate Names:**

Inflamed Gallbladder



---

## Cholecystography

An imaging test of the gallbladder after contrast material is given to show how it fills, concentrates bile, and empties. Ultrasound, CT, or other modern tests are usually preferred for current gallbladder evaluation.

## Cholecystokinin

A hormone released by the small intestine after food, especially fat and protein, enters it. It makes the gallbladder release bile, stimulates digestive enzymes, and helps create a feeling of fullness.

## Cholecystoses

Non-inflammatory disorders causing deposits, wall thickening, or other structural changes in the gallbladder, including cholesterosis and adenomyomatosis. Many are incidental, but symptoms and imaging findings determine whether further evaluation is needed.

## Cholecystostomy

The creation of an opening into the gallbladder lumen for tube decompression and drainage of infected bile, frequently performed percutaneously under image guidance in critically ill patients with acute cholecystitis who are unfit for urgent cholecystectomy.

### Alternate Names:

Percutaneous Cholecystostomy

## Choledochal Cyst

An abnormal widening of one or more bile ducts, usually present from birth. It may cause abdominal pain, jaundice, pancreatitis, or infection and can increase later cancer risk, so specialist evaluation is needed.

## Choledochal Cysts

Choledochal cysts are congenital or acquired dilations of the bile ducts that can cause abdominal pain, jaundice, pancreatitis, cholangitis, or stones. Imaging establishes the anatomy, and surgical removal is often recommended because of recurrent complications and cancer risk.

### Alternate Names:

Choledochal Cyst; Congenital Bile Duct Dilatation

## Choledocholithiasis

A gallstone within the common bile duct, which may cause biliary pain, jaundice, cholangitis, or pancreatitis and is usually evaluated with liver tests and imaging.

## Cholelithiasis

The presence of stones in the gallbladder or biliary tract. Stones may contain cholesterol, pigment, or mixed material and may cause no symptoms or lead to biliary pain and complications.

## Cholera

An acute intestinal infection caused by toxigenic *Vibrio cholerae*, usually acquired through contaminated water or food. It can produce sudden profuse watery diarrhoea, vomiting, severe dehydration, shock, and death within hours if untreated; prompt oral or intravenous rehydration is central treatment. Safe water, sanitation, food hygiene, and vaccination in selected settings help prevent spread.

## Choleresis

The production and release of bile by liver cells. Bile helps digest fats and carries some waste products from the liver into the intestine.

## Choleretic

A substance that encourages the liver to make or release bile. Bile helps digest fats and remove some waste products; a choleretic should be used carefully when gallstones or blocked bile ducts may be present.

### Alternate Names:

Bile-Secretory Agent; Bile-Flow Stimulant

## Cholestasis

Reduced production or flow of bile. It may cause itching, yellow skin or eyes, dark urine, pale stools, and abnormal liver tests; a blockage and liver-cell causes must be distinguished.

---

## Cholestasis Of Pregnancy

A pregnancy-related liver disorder in which bile acids build up in the blood and often cause intense itching without a rash, especially on the palms and soles. It can increase risks to the fetus, so testing and obstetric monitoring are important.

### Alternate Names:

Obstetric Cholestasis

## Cholesteatoma

A keratin-filled sac of abnormal squamous epithelium in the middle ear or mastoid that can erode bone and cause chronic discharge, hearing loss, or intracranial complications.

## Cholesterol

A lipid required for cell membranes and synthesis of steroid hormones and bile acids. Excess atherogenic cholesterol can contribute to arterial plaque.

### Alternate Names:

Serum Cholesterol; Blood Cholesterol

## Cholesterol And Lifestyle

Diet, physical activity, weight, smoking, and alcohol use can affect blood cholesterol and cardiovascular risk. Lifestyle changes complement medicines when cholesterol remains high.

## Cholesterol And Lipids

Fatty substances in the blood, including LDL cholesterol, HDL cholesterol, and triglycerides, that play critical roles in cellular function and energy metabolism. Abnormal lipid levels, particularly high LDL and triglycerides with low HDL, promote atherosclerosis and increase the risk of heart attack and stroke.

## Cholesterol Crystal Embolization

Small pieces of cholesterol plaque breaking loose from a large artery and blocking smaller arteries. It may damage the kidneys, skin, eyes, brain, or intestines, often after a vascular procedure.

## Cholesterol Crystals

Needle-shaped cholesterol deposits that can form

inside fatty artery plaques. If pieces break loose, they may block smaller vessels and reduce blood flow to the kidneys, skin, eyes, brain, or intestines.

## Cholesterol Emboli

Cholesterol crystals released from an atherosclerotic plaque that lodge in small arteries and cause ischemia or inflammation in downstream tissues.

## Cholesterol Embolism

Blockage of small arteries by cholesterol crystals released from an artery wall, often after vascular procedures or disruption of atherosclerotic plaque.

## Cholesterol Embolization

A condition in which tiny pieces of fatty material from an artery wall break loose and travel into smaller blood vessels. It can damage the kidneys, skin, eyes, brain, or intestines, sometimes after a vascular procedure or blood-thinning treatment.

## Cholesterol Lowering

Alternate name; see *Hyperlipidemia*.

## Cholesterol Synthesis

The process by which cells, especially liver and intestinal cells, make cholesterol from smaller molecules. Cholesterol is needed for cell membranes, hormones, and bile acids, but excess production contributes to high blood cholesterol.

## Cholesterol Testing And Results

Laboratory assessment of blood lipids, usually including total cholesterol, LDL cholesterol, HDL cholesterol, and triglycerides, used to estimate cardiovascular risk and guide prevention or treatment.

## Cholesterolemia

The presence and concentration of cholesterol in the blood. Abnormal elevation is a risk factor for atherosclerotic cardiovascular disease.

## Cholesterosis

Deposition of cholesterol esters in the gallbladder mucosa, producing yellow speckling or a "strawberry

---

gallbladder." It is often incidental but may coexist with gallstones or chronic gallbladder inflammation.

### **Cholestyramine**

Cholestyramine is a bile-acid sequestrant used to lower LDL cholesterol and sometimes to treat bile-acid diarrhea or itching from cholestasis.

### **Choline**

Choline is an essential nutrient used to make cell membranes and acetylcholine and to support liver and nervous-system function.

### **Cholinergic**

Relating to acetylcholine, a chemical messenger used by nerves. Cholinergic signals affect muscles, glands, the heart, digestion, breathing, and parts of the brain.

### **Cholinergic Crisis**

A life-threatening excess of acetylcholine, often after organophosphate or carbamate poisoning. It can cause sweating, salivation, vomiting, diarrhea, pinpoint pupils, wheezing, muscle weakness, seizures, and respiratory failure.

### **Cholinergic Neuron**

A nerve cell that releases acetylcholine as its principal neurotransmitter.

#### **Alternate Names:**

Acetylcholine-Releasing Neuron; Cholinergic Nerve Cell

### **Cholinergic Toxidrome**

A poisoning pattern caused by excessive acetylcholine activity, often producing salivation, tearing, urination, diarrhea, vomiting, bronchial secretions, slow heart rate, muscle fasciculations, weakness, or respiratory failure.

### **Cholinergic Urticaria**

A chronic inducible urticaria triggered by increased body temperature or sweating, such as during exercise, hot bathing, spicy food, or emotional stress. It causes numerous small, short-lived itchy wheals,

sometimes with flushing or angioedema; rare breathing or circulatory symptoms require emergency care.

### **Cholinesterase**

Cholinesterase enzymes break down acetylcholine; inhibition increases cholinergic activity and is used therapeutically but can also cause poisoning.

### **Chondritis**

Inflammation of cartilage, such as the cartilage of the ear, nose, ribs, or larynx. It may result from infection, trauma, or immune-mediated disease.

### **Chondroblastic Osteosarcoma**

A type of bone cancer in which the cancer cells make cartilage-like material. It often affects the widened ends of long bones in adolescents and young adults and is diagnosed from imaging and a tissue sample.

### **Chondroblastoma**

A chondroblastoma is an uncommon usually benign cartilage-forming bone tumor, often arising near the end of a long bone in adolescents or young adults.

### **Chondrocalcinosis**

Calcium-crystal deposits in cartilage or other joint tissues that appear on imaging. They are often linked to calcium pyrophosphate deposition disease and may cause joint pain and swelling or be found without symptoms.

### **Chondrocyte**

A mature cell that makes and maintains cartilage, the firm flexible tissue found in joints, airways, and other places. Chondrocytes help keep cartilage smooth and strong, but cartilage usually heals slowly because it has little direct blood supply.

#### **Alternate Names:**

Cartilage Cell

### **Chondrodermatitis Nodularis Helicis**

A painful inflammatory nodule affecting the skin and cartilage of the ear's helix or antihelix, often related to repeated pressure, sleeping on one side,

---

cold, sun, or local trauma. A firm tender crusted lesion can resemble skin cancer, so persistent or atypical lesions may need biopsy; pressure relief is an important treatment.

### **Chondrodysplasia**

A disorder of cartilage and bone development that causes abnormal skeletal growth. Depending on the type, it may produce short limbs, short stature, enlarged joints, or other bone abnormalities.

### **Chondroitin Sulfate**

A sulfated glycosaminoglycan found in cartilage and other connective tissues; it is also marketed in supplements for osteoarthritis, with variable evidence of benefit.

### **Chondroma**

A usually harmless tumour made of mature cartilage. It most often forms inside a bone, especially in the hands and feet, or on the bone surface. Many cause no symptoms, but pain, growth, or fracture needs assessment.

### **Chondromalacia**

Softening or breakdown of cartilage, especially the cartilage under the kneecap. It may cause pain at the front of the knee, grinding, or discomfort with stairs, squatting, or prolonged sitting.

### **Chondromalacia Patellae**

A musculoskeletal disorder characterized by softening, fibrillation, and degeneration of the hyaline articular cartilage on the posterior articulating surface of the patella. It presents with anterior retro-patellar knee pain aggravated by climbing stairs, kneeling, squatting, or prolonged sitting.

#### **Alternate Names:**

Patellofemoral Syndrome; Runner's Knee

### **Chondromyxoid Fibroma**

A rare benign cartilage-forming bone tumor, usually arising in the metaphysis of a long bone and presenting with localized pain or an incidental radiographic lesion.

### **Chondrosarcoma**

A cancer that forms cartilage, usually within a bone. It is more common in adults and often grows slowly, though some types can grow faster or spread. Rarely, it begins in soft tissue rather than bone.

### **Chondrosarcoma Of The Breast**

An extremely rare breast cancer whose cells produce cartilage-like tissue. It usually appears as a growing breast mass and is diagnosed by examining tissue; treatment generally requires complete surgical removal by a specialist team.

### **Chopart's Joint**

The combined joint across the middle of the foot, between the hindfoot and midfoot. It includes the talonavicular and calcaneocuboid joints and helps the foot adapt to surfaces and transfer force during walking.

### **Chorangioma**

A benign vascular tumor of the placenta, usually small and incidental but occasionally associated with fetal anemia, hydrops, growth restriction, or other complications when large.

### **Chordae Tendineae**

Strong, thin cords inside the heart that connect the mitral and tricuspid valves to small muscles. They help keep the valves from turning inside out when the heart contracts.

### **Chordee**

Chordee is abnormal downward or other curvature of the penis, commonly associated with hypospadias or scar tissue and sometimes requiring surgical correction.

### **Chordoma**

A rare cancer arising from leftover cells of early spinal development. It usually begins near the tailbone, base of the skull, or spine and can cause pain, weakness, or nerve problems as it grows.

---

## Chorea

Uncontrolled, brief, irregular movements that flow unpredictably from one body part to another. It may result from inherited disease, autoimmune illness, medicines, infection, or damage to movement-control areas of the brain.

## Chorea and Myoclonus

Sudden, rapid, irregular, jerky, and non-rhythmic involuntary muscle contractions or shock-like jerks. Chorea flows unpredictably from one muscle group to another, whereas myoclonus represents brief shock-like twitches, observed in metabolic encephalopathies, post-hypoxic syndromes, and neurodegenerative disorders.

### Alternate Names:

Jerky Involuntary Movements; Choreiform Activity; Myoclonic Jerks

## Choreoathetosis

A movement disorder combining brief, unpredictable jerks with slower twisting movements. It can result from injury or disease affecting movement-control areas deep in the brain, inherited conditions, or certain medicines.

## Chorioamnionitis

Infection or inflammation of the amniotic fluid, membranes, placenta, or decidua, usually due to ascending polymicrobial infection. Maternal fever, uterine tenderness, fetal tachycardia, and maternal leukocytosis may occur; treatment requires broad-spectrum antibiotics and delivery planning.

### Alternate Names:

Intraamniotic Infection

## Choriocarcinoma

A fast-growing cancer arising from cells that normally help form the placenta. It usually develops after a pregnancy or molar pregnancy and produces high levels of the pregnancy hormone hCG; it can spread but often responds well to chemotherapy.

## Chorion

The outer membrane surrounding an early embryo. Its finger-like projections help form the fetal part

of the placenta, where oxygen, nutrients, and waste are exchanged between fetal and maternal blood supplies.

## Chorion Villus Sampling

Chorion villus sampling removes placental tissue for genetic or chromosomal testing, usually early in pregnancy; it carries a small procedure-related miscarriage risk.

## Chorionic Gonadotrophin

Chorionic gonadotrophin, especially hCG, is a placental hormone that maintains early pregnancy and is measured in pregnancy tests and selected tumor evaluations.

## Chorionic Villus Sampling

A prenatal test usually performed around 10 to 13 weeks by removing a small placental sample through the abdomen or cervix. It can identify many chromosome and inherited disorders but carries a small miscarriage risk.

## Chorioretinitis

Inflammation involving the choroid and retina, often caused by infection, autoimmune disease, or systemic inflammatory illness. It can produce floaters and impaired vision.

## Choroid

The blood-vessel-rich layer between the retina and the white outer coat of the eye. It supplies oxygen and nutrients to the outer retina and helps absorb stray light inside the eye.

## Choroid Plexus

A network of tissue inside the brain's ventricles that produces most of the cerebrospinal fluid.

## Choroid Plexus Carcinoma

A cancer arising from the choroid plexus, tissue that produces cerebrospinal fluid, potentially causing pressure buildup, headaches, vomiting, or neurologic problems.

---

### **Choroid Plexus Cyst**

A fluid-filled cyst within the choroid plexus of a fetal cerebral ventricle. An isolated cyst with otherwise reassuring prenatal assessment is usually benign, but its significance depends on other ultrasound findings and screening results.

### **Choroid Plexus Tumor**

A growth arising from the tissue that makes cerebrospinal fluid inside the brain. It occurs most often in children and may block fluid flow or produce too much fluid, causing hydrocephalus, headache, vomiting, or increasing head size.

### **Choroidal Dystrophies**

Inherited disorders causing progressive degeneration of the choroid, retinal pigment epithelium, and retina, leading to night blindness, visual-field loss, and eventual vision impairment.

### **Choroidal Melanoma**

A cancer arising from pigment-producing melanocytes in the choroid, the vascular layer beneath the retina. It may cause visual changes or be found incidentally and can spread, especially to the liver; treatment depends on size and location.

### **Choroidal Neovascularization**

Growth of fragile abnormal blood vessels beneath or through the retina. Leakage or bleeding can damage central vision, especially in wet age-related macular degeneration and some inflammatory or inherited eye disorders.

### **Choroideremia**

Choroideremia is an X-linked inherited retinal degeneration causing progressive night blindness and loss of peripheral then central vision, mainly in males.

### **Choroiditis**

Choroiditis is inflammation of the choroid, the vascular layer behind the retina, often causing blurred vision, floaters, or visual-field disturbance.

### **Christmas Disease**

An inherited bleeding disorder, also called hemophilia B, caused by low or abnormal clotting factor IX. It can cause prolonged bleeding or bleeding into joints and muscles and is treated with replacement or other specialist therapy.

### **Chromaffin**

Describes cells that contain stored adrenaline-like hormones. They are found mainly in the inner part of the adrenal glands and in related nerve tissue and release these hormones during stress or in response to nerve signals.

### **Chromaffinoma**

A tumor of cells that can produce adrenaline-like hormones, usually in the adrenal gland or nearby nerve tissue. Hormone release may cause episodes of high blood pressure, headache, sweating, and palpitations.

### **Chromate**

Chromate is a hexavalent chromium compound that can cause irritant or allergic dermatitis, nasal injury, respiratory disease, and cancer after occupational exposure.

### **Chromatic**

Chromatic means relating to color or wavelengths of light; chromatic vision depends on cone photoreceptors and neural processing.

### **Chromatids**

Chromatids are the two copies of a replicated chromosome joined at the centromere before separating during cell division.

### **Chromatin**

The combination of DNA and proteins that packages genetic material inside the cell nucleus. Its arrangement controls which genes are available for use and helps regulate DNA copying and repair.

### **Chromatography**

A laboratory separation method in which substances



---

partition between a stationary phase and a moving phase, allowing identification or measurement of components in a mixture.

### **Chromatopsia**

A visual disturbance in which objects appear abnormally colored. It may result from retinal disease, toxic exposure, medication effects, or other disorders of the visual system.

### **Chromhidrosis**

A rare disorder in which coloured sweat, often yellow, green, blue, or black, appears because pigment is produced in apocrine glands. Similar discoloration can result when ordinary sweat becomes coloured by surface bacteria or chemicals, a process called pseudochromhidrosis; evaluation distinguishes these causes and looks for an underlying medicine or skin disorder.

### **Chromium**

Chromium is a metallic element; trivalent chromium is a trace nutrient, while hexavalent compounds can be toxic, corrosive, mutagenic, and carcinogenic.

### **Chromium In Diet**

Reference information on chromium as a trace element involved in carbohydrate and fat metabolism. Chromium deficiency is rare; most people obtain adequate chromium from normal dietary intake, and routine chromium supplementation is generally not recommended.

### **Chromoblastomycosis**

A chronic implantation infection caused by pigmented fungi, usually after minor trauma introduces the organism into skin. It produces slowly enlarging warty, scaly, or tumorous plaques, most often on the legs; biopsy may show muriform bodies, and prolonged antifungal treatment is often required.

### **Chromopertubation**

A fertility test performed during laparoscopy in which colored fluid is passed through the uterus while the fallopian tubes are viewed. Fluid spilling from the tube ends suggests that they are open;

no spill may indicate blockage or surrounding scar tissue.

### **Chromosomal Abnormality**

A change in the number or structure of chromosomes. It may involve an extra or missing chromosome, or a missing, repeated, rearranged, or exchanged chromosome segment, and can affect development, health, or fertility.

### **Chromosomal Breakage Syndromes**

A group of inherited DNA-repair disorders in which chromosomes are unusually fragile or break easily. They may cause congenital anomalies, growth or developmental problems, bone-marrow failure, immune abnormalities, and increased cancer risk.

### **Chromosomal Microarray**

A genetic test that detects gains and losses of chromosome material, including copy-number variants and some regions of homozygosity; it is commonly used for developmental delay, intellectual disability, congenital anomalies, or autism-spectrum presentations.

### **Chromosome**

A thread-like structure made of DNA and proteins that carries genetic information. Most human body cells normally contain 46 chromosomes arranged in 23 pairs.

### **Chromosome 1**

The largest human chromosome, containing many genes involved in development, cell function, and disease. Extra, missing, or rearranged material can cause genetic disorders.

### **Chromosome 13**

A human chromosome containing genes important for development and cell function. Abnormal chromosome 13 number or structure can cause disorders such as trisomy 13.

### **Chromosome 2**

A large human chromosome containing many genes

---

involved in growth, development, and normal body function. Changes in its number or structure can cause genetic disease.

### **Chromosome 3**

A human chromosome containing genes involved in development, immune function, and cell regulation. Deletions, duplications, or other changes may cause genetic disorders or cancer.

### **Chromosome 4**

A human chromosome containing many genes involved in development and cellular function. Structural or numerical abnormalities can cause congenital or inherited conditions.

### **Chromosome 5**

A human chromosome containing genes involved in development, immunity, and cell regulation. Loss of part of its short arm causes the genetic condition cri-du-chat syndrome.

### **Chromosome Complement**

The complete set of chromosomes in a cell, including their number and sex-chromosome composition, usually described by a karyotype.

### **Chromosome Duplication**

A structural chromosome abnormality in which a segment of genetic material is copied one or more additional times, potentially altering gene dosage and development.

### **Chromosome X**

One of the human sex chromosomes. Most females have two X chromosomes and most males have one X and one Y; genes on the X chromosome can cause X-linked disorders when altered.

### **Chromosome Y**

The sex chromosome typically present in most males and absent in most females. It contains the sex-determining *SRY* gene and other genes important for testis development and sperm production.

### **Chronic**

Persisting or recurring over a long period, in contrast to acute; the appropriate duration depends on the disease or clinical context.

### **Chronic Adrenal Insufficiency**

Alternate name; see *Addison's Disease*.

### **Chronic Allograft Nephropathy**

Gradual loss of function in a transplanted kidney over months or years, caused by immune injury, high blood pressure, recurrent disease, medicines, or other damage. Testing and biopsy help identify the cause.

### **Chronic Atrophic Gastritis**

Long-term inflammation that destroys glands in the stomach lining. It may reduce stomach acid and intrinsic factor, impair iron or vitamin B12 absorption, and increase the risk of anemia or stomach tumors.

### **Chronic Autoimmune Hashimoto Thyroiditis**

A long-term immune-system attack on the thyroid gland that gradually damages thyroid tissue and often causes too little thyroid hormone. It may produce tiredness, cold intolerance, weight change, or an enlarged thyroid.

### **Chronic Autoimmune Thyroiditis**

Long-term inflammation caused by the immune system attacking the thyroid, usually called Hashimoto thyroiditis. It commonly leads to low thyroid hormone and may be associated with thyroid antibodies.

### **Chronic Bacterial Prostatitis**

A recurring bacterial infection of the prostate that may cause repeated urinary infections, burning or urgent urination, pelvic discomfort, or painful ejaculation. Diagnosis uses appropriate urine tests, and treatment often lasts several weeks.

#### **Alternate Names:**

NIH Category II Prostatitis; Recurrent Bacterial

---

Prostatitis

### Chronic Bronchitis

A cough producing mucus on most days for at least three months in each of two consecutive years, after other causes are excluded. It reflects long-term airway irritation and commonly occurs with chronic obstructive pulmonary disease.

#### Alternate Names:

Airway Hypersecretion; Chronic Mucous Bronchitis

### Chronic Bronchitis With Cor Pulmonale

Chronic mucus-producing airway disease complicated by high blood pressure in the lung arteries and strain or failure of the right side of the heart (cor pulmonale). Chronic obstructive pulmonary disease is a common underlying cause.

### Chronic Bronchitis With Emphysema

Coexisting chronic mucus-producing airway disease and emphysema, in which air sacs in the lungs are damaged. They commonly occur together in chronic obstructive pulmonary disease, but spirometry is needed to confirm persistent airflow obstruction.

### Chronic Caries

Tooth decay that develops slowly over time. It may cause a firm dark area, a cavity, sensitivity, or no symptoms; without dental treatment, decay can reach the inner tooth, causing pain, infection, or tooth loss.

### Chronic Cholecystitis

Repeated or persistent inflammation of the gallbladder, usually related to gallstones and impaired emptying. It may cause episodic right-upper-abdominal pain, nausea, or food intolerance and can lead to scarring or a contracted gallbladder.

### Chronic Compartment Syndrome

Alternate name; see *Chronic Exertional Compartment Syndrome*.

### Chronic Coronary Syndrome

Long-term coronary artery disease with stable or

recurring symptoms caused by reduced blood flow to the heart muscle. It may occur with exertion or emotional stress and can also be present without symptoms.

### Chronic Cough

A cough lasting longer than eight weeks. Common causes include asthma, nasal drainage, acid reflux, smoking, and some blood-pressure medicines; persistent cough requires assessment for its cause.

#### Alternate Names:

Persistent Cough; Intractable Cough

### Chronic Daily Headaches

Headaches occurring on at least 15 days each month for more than three months. They may reflect chronic migraine, tension-type headache, frequent pain-medicine use, or another disorder and require assessment for the cause.

### Chronic Disease

A disease or condition that persists for a prolonged period, often months or years, and may require continuing monitoring or management.

### Chronic Ear Infection

A persistent, recurrent, or long-standing infection of the middle ear, often associated with a persistent eardrum perforation and painless or foul-smelling ear drainage.

#### Alternate Names:

Chronic Otitis Media; Chronic Suppurative Otitis Media; CSOM

### Chronic Endometritis

Long-lasting inflammation of the inner lining of the uterus. It may be linked to infection or tissue left after pregnancy and can cause irregular bleeding, pelvic discomfort, or difficulty becoming pregnant; some people have no symptoms.

### Chronic Eosinophilic Leukaemia

A rare blood cancer in which the bone marrow makes too many eosinophils, a type of white blood cell, because of abnormal cell growth. It may cause

---

tiredness, fever, weight loss, an enlarged spleen, or damage to the heart, lungs, nerves, or skin.

### **Chronic Eosinophilic Pneumonia**

A usually noninfectious lung inflammation in which eosinophils, a type of white blood cell, collect in lung tissue. It often causes weeks of cough, breathlessness, fever, night sweats, or weight loss and may recur after treatment.

### **Chronic Exertional Compartment Syndrome**

Exercise-related increased pressure within a muscle compartment that causes predictable pain, tightness, or neurologic symptoms during activity.

#### **Alternate Names:**

Chronic Compartment Syndrome

### **Chronic Fatigue Syndrome**

A long-term illness involving substantial reduced ability to function, worsening after physical or mental activity, unrefreshing sleep, and often difficulty thinking or standing. Symptoms are not explained by another condition and need careful evaluation.

#### **Alternate Names:**

CFS; Myalgic Encephalomyelitis (ME); Systemic Exertion Intolerance Disease (SEID)

### **Chronic Fever**

A fever that continues or repeatedly returns over a long period. Causes include infection, inflammatory disease, cancer, and medicines; persistent unexplained fever needs medical evaluation based on its pattern and accompanying symptoms.

### **Chronic Gastritis**

Chronic, progressive inflammation of the gastric mucosa leading over time to mucosal atrophy, loss of specialized parietal and chief cells, and intestinal metaplasia. Major causes include persistent *Helicobacter pylori* infection and autoimmune gastritis targeting parietal cells and intrinsic factor.

### **Chronic Glomerulonephritis**

Long-standing inflammation and scarring of the

kidney glomeruli that can progressively reduce filtration and lead to chronic kidney disease or kidney failure.

### **Chronic Granulomatous Disease**

An inherited immune disorder in which certain white blood cells cannot effectively kill some bacteria and fungi. It causes repeated serious infections, swollen lymph nodes, pneumonia, skin infections, and inflammation called granulomas.

### **Chronic Hypertension**

Persistently elevated systemic arterial blood pressure present over time, whether due to an identified underlying cause or without an identifiable secondary cause. It is distinct from hypertension arising only during pregnancy.

### **Chronic Hypertension In Pregnancy**

Hypertension that predates pregnancy or is diagnosed early in pregnancy. It increases the risk of preeclampsia, fetal growth problems, preterm birth, and placental complications and requires ongoing maternal and fetal monitoring.

### **Chronic Idiopathic Constipation**

Long-lasting difficult or infrequent bowel movements without an identified structural or metabolic cause. Symptoms may include hard stools, straining, incomplete emptying, or abdominal discomfort; treatment includes diet, fluids, activity, and selected medicines.

### **Chronic Inflammation**

Inflammation that continues for weeks, months, or years. It can occur with persistent infection, autoimmune disease, long-term irritation, or obesity and may cause ongoing tissue damage while the body attempts repair.

### **Chronic Inflammatory Demyelinating Polyneuropathy**

An immune disorder that damages the insulating covering of nerves outside the brain and spinal cord. It causes gradually worsening or recurring weakness, numbness, poor balance, and reduced reflexes in the

---

limbs.

### **Chronic Inflammatory Response Syndrome**

A proposed multisystem illness involving persistent inflammatory symptoms after an exposure such as certain water-damaged building environments. Reported symptoms are nonspecific and may include fatigue, headache, cognitive difficulty, and respiratory complaints. Symptoms require a careful medical evaluation for established infectious, allergic, toxic, autoimmune, and other causes.

### **Chronic Interstitial Nephritis**

Long-term inflammation and scarring of the kidney tissue between the tubules. It may result from medicines, repeated infection, urinary obstruction, toxins, autoimmune disease, or inherited disorders and can gradually reduce kidney function.

### **Chronic Intestinal Pseudo-Obstruction**

Repeated symptoms of bowel blockage without a physical obstruction, caused by poor nerve or muscle function in the intestine. It may cause abdominal swelling, pain, vomiting, constipation, and inability to absorb nutrition.

### **Chronic Kidney Disease**

Kidney damage, a kidney-structure problem, or reduced kidney function that lasts at least three months. The kidneys may then have difficulty filtering waste and extra fluid from the blood and balancing salts, acids, and other body functions. Diabetes and high blood pressure are common causes. Early disease often causes no symptoms; advanced disease may cause swelling, tiredness, poor appetite, nausea, itching, anemia, mineral and bone problems, or kidney failure. Chronic kidney disease is different from a sudden loss of kidney function, called acute kidney injury.

#### **Alternate Names:**

CKD; Chronic Renal Disease; Chronic Renal Failure

### **Chronic Kidney Disease Progression**

Worsening loss of kidney function or increasing kidney damage over time. Progression is assessed with

trends in estimated glomerular filtration rate, albumin in urine, blood pressure, symptoms, and other findings.

### **Chronic Kidney Disease Staging**

Classification of kidney disease by cause, eGFR category, and albuminuria category. G1 through G5 represent eGFR from normal or high to kidney failure; staging is applied when abnormalities persist for at least 3 months and helps estimate progression and complications.

### **Chronic Kidney Disease-Mineral and Bone Disorder**

A systemic disorder of mineral and bone metabolism resulting from chronic kidney disease, characterized by abnormalities of calcium, phosphorus, parathyroid hormone, or vitamin D metabolism; abnormalities in bone turnover, mineralization, linear growth, or strength (renal osteodystrophy); and extra-skeletal vascular or soft-tissue calcification.

#### **Alternate Names:**

CKD-MBD; Renal Mineral and Bone Disorder

### **Chronic Kidney Failure**

Alternate name; see *Chronic Kidney Disease*.

### **Chronic Lead Poisoning**

Illness caused by long-term or repeated exposure to lead. It may cause abdominal pain, constipation, tiredness, anemia, nerve damage, high blood pressure, or kidney injury. In children, it can impair learning, attention, growth, or development. Treatment begins with stopping the exposure; some people need specialist-directed chelation.

### **Chronic Leukemia**

A blood cancer that develops gradually as abnormal white blood cells multiply and remain in the blood or bone marrow. Some types progress slowly but still require regular medical monitoring.

### **Chronic Limb-Threatening Ischemia**

Severe, long-lasting reduction of blood flow to a limb, usually from narrowed or blocked arteries. It causes pain at rest, ulcers that do not heal, or



---

gangrene and requires urgent vascular assessment to protect the limb.

### **Chronic Liver Disease**

A broad term for long-term liver damage or disease. Causes include viral infection, alcohol, fatty-liver disease, autoimmune disease, blocked bile flow, medicines, and inherited disorders. It may occur with or without cirrhosis; progressive scarring can lead to cirrhosis or liver failure.

### **Chronic Low Back Pain**

Persistent or recurring pain in the lumbar or lumbosacral region lasting longer than 12 weeks. Multifactorial etiologies include degenerative disc disease, lumbar spinal stenosis, spondylosis, facet arthropathy, and central pain sensitization.

#### **Alternate Names:**

Chronic Lumbago; Degenerative Lumbar Spine Disease; Long-Term Back Pain

### **Chronic Lymphocytic Leukemia**

A usually slow-growing blood cancer in which abnormal B lymphocytes build up in the blood, bone marrow, and lymph nodes. It may cause enlarged lymph nodes, tiredness, repeated infections, anaemia, or no symptoms at first.

#### **Alternate Names:**

CLL; Chronic Lymphoid Leukemia

### **Chronic Lymphocytic Thyroiditis**

An autoimmune disease in which immune attack gradually damages the thyroid, often causing hypothyroidism, fatigue, cold intolerance, constipation, dry skin, or goiter.

### **Chronic Megacolon**

Persistent abnormal enlargement and impaired propulsion of the colon. Causes include severe constipation, aganglionosis, neurologic disease, metabolic disorders, medications, or chronic intestinal obstruction.

### **Chronic Meningitis**

Persistent inflammation of the membranes around

the brain and spinal cord, usually lasting four weeks or longer. Causes include tuberculosis, fungal or other infections, cancer, autoimmune disease, and medication reactions; symptoms may develop gradually or fluctuate.

### **Chronic Mesenteric Ischemia**

Reduced blood flow to the intestines, usually from hardening and narrowing of abdominal arteries. It often causes cramping after meals, food avoidance, and weight loss and needs vascular evaluation.

### **Chronic Migraine**

Headache on at least 15 days each month for more than three months, with migraine features on at least eight days. Pain may be one-sided or throbbing and may include nausea or sensitivity to light or sound.

### **Chronic Mitral Regurgitation**

Long-standing leakage through the mitral valve during left-ventricular contraction. It may be primary, such as from prolapse, rheumatic disease, or endocarditis, or secondary to left-ventricular enlargement or ischemia; chronic volume overload can enlarge the left atrium and ventricle and lead to atrial fibrillation, pulmonary hypertension, or heart failure.

### **Chronic Motor Or Vocal Tic Disorder**

A neurodevelopmental disorder with persistent motor or vocal tics lasting more than one year; symptoms often fluctuate and may be treated with education, behavioral therapy, or medicines when impairment is significant.

### **Chronic Mucocutaneous Candidiasis**

Persistent or recurrent *Candida* infections of the skin, nails, and mucous membranes, often beginning in childhood. It may reflect a specific inherited T-cell or IL-17 pathway defect, or occur with autoimmune or endocrine disease; persistent disease requires evaluation for an underlying immune problem.

### **Chronic Myelogenous Leukemia**

A blood cancer in which the bone marrow makes too



---

many abnormal white blood cells, usually because of the BCR::ABL1 gene change. It may progress over time but is often controlled with targeted medicines.

### **Chronic Myeloid Leukemia**

A blood cancer usually caused by the Philadelphia chromosome, which creates the BCR::ABL1 gene. The bone marrow produces too many abnormal white blood cells, and targeted treatment can often control the disease.

### **Chronic Myelomonocytic Leukemia**

A rare blood cancer in which the bone marrow makes too many monocytes, a type of white blood cell, along with abnormal blood-forming cells. It can cause anemia, infections, bleeding, fatigue, or an enlarged spleen.

### **Chronic Neutrophilic Leukaemia**

A rare blood cancer in which the bone marrow continually makes too many mature neutrophils, a type of white blood cell, without infection being the main cause. It may cause tiredness, fever, weight loss, or an enlarged spleen.

### **Chronic Obstructive Pulmonary Disease**

A long-term lung disease that causes persistent air-flow blockage and breathing difficulty. Many people have emphysema, chronic bronchitis, or both, but COPD patterns vary and can involve other airway and lung-tissue changes. Symptoms may include breathlessness, cough, mucus, and wheezing. Smoking is a common cause, but other harmful inhaled exposures can also contribute; spirometry is used to confirm persistent airflow obstruction.

#### **Alternate Names:**

COPD

### **Chronic Otitis Media**

Long-standing or recurrent inflammation of the middle ear, most often from persistent fluid and negative pressure after repeated infection or eustachian-tube dysfunction. It may be largely silent or cause pain, discharge, hearing loss, and eardrum changes, and it can damage the eardrum or middle-ear bones; forms with continuous discharge carry higher risk of

cholesteatoma and intracranial complications. Management controls infection and ventilation problems, and referral is warranted when discharge or hearing loss persist.

### **Chronic Pain**

Pain that persists or recurs for longer than three months. It may have an identifiable ongoing cause or persist after healing, and is influenced by complex interactions among biological, psychological, and social factors.

### **Chronic Pain Syndrome**

Long-lasting pain accompanied by substantial effects on movement, sleep, mood, relationships, or daily activities. Treatment addresses pain and these related effects rather than relying only on pain medicines.

### **Chronic Pancreatitis**

Repeated or long-term inflammation that permanently scars the pancreas. It may cause upper-abdominal pain, poor digestion, weight loss, diabetes, or oily stools; alcohol, smoking, inherited disorders, and blocked ducts are possible causes.

### **Chronic Paroxysmal Hemicrania**

A primary headache disorder causing frequent, short-lasting, severe one-sided headaches with autonomic symptoms and a characteristic response to indomethacin.

#### **Alternate Names:**

CPH; Chronic Paroxysmal Hemicrania Syndrome

### **Chronic Pelvic Pain**

Pain in the pelvic area that lasts for six months or longer. It may be constant or come and go, and may be related to reproductive, urinary, digestive, muscle, nerve, or pain-processing conditions. Sometimes more than one cause is present, or no single cause can be found.

#### **Alternate Names:**

CPPS; Persistent Pelvic Pain Syndrome

### **Chronic Pelvic Pain in Women**

Pain in the pelvis lasting or recurring for months

---

and affecting daily life, sexual activity, urination, bowel function, or mood. Causes may include endometriosis, pelvic floor disorders, bladder or bowel disease, adhesions, neuropathic pain, or no single identifiable cause.

### **Chronic Pelvic Pain Syndrome**

Persistent or recurring pain in the pelvis, area between the genitals and anus, lower abdomen, back, or scrotum without a proven ongoing bacterial infection. Urinary symptoms, painful ejaculation, muscle tension, and sexual problems may also occur.

#### **Alternate Names:**

CPPS; NIH Category III Prostatitis; Chronic Non-bacterial Prostatitis

### **Chronic Progressive External Ophthalmoplegia**

A slowly worsening muscle disorder causing drooping eyelids and limited movement of both eyes. It often begins in adulthood and may occur alone or with weakness of other muscles; inherited mitochondrial disease is a common cause.

#### **Alternate Names:**

CPEO; Progressive External Ophthalmoplegia (PEO)

### **Chronic Pyelonephritis**

Long-term inflammation and scarring of the kidneys, often after repeated infections or backward flow of urine. It can gradually reduce kidney function and may cause high blood pressure, poor growth, or kidney failure.

### **Chronic Recurrent Multifocal Osteomyelitis**

A rare autoinflammatory bone disorder, often beginning in children or adolescents, causing recurrent sterile inflammation and pain at multiple bone sites; it is also called CRMO.

### **Chronic Renal Failure**

Alternate name; see *Chronic Kidney Disease*.

### **Chronic Respiratory Failure**

Long-term inability of the lungs to maintain enough oxygen in the blood, remove enough carbon dioxide, or both. It may result from COPD, interstitial lung disease, neuromuscular weakness, chest-wall disorders, or other chronic conditions.

#### **Alternate Names:**

Chronic Ventilatory Failure; Long-Term Respiratory Insufficiency

### **Chronic Rhinitis**

Long-lasting inflammation or irritation of the nose causing congestion, runny nose, sneezing, itching, or mucus draining into the throat. Causes include allergy, irritants, medicines, and structural problems; treatment is chosen after identifying the cause.

#### **Alternate Names:**

Chronic Non-Allergic Rhinitis; Vasomotor Rhinitis; Persistent Rhinitis

### **Chronic Rhinosinusitis**

Persistent inflammation of the nose and paranasal sinuses lasting at least 12 weeks, usually with nasal obstruction or discharge and sometimes facial pressure or reduced smell. It may occur with or without nasal polyps.

### **Chronic Rhinosinusitis With Nasal Polyps**

Long-lasting inflammation of the nose and sinuses with soft, swollen growths called nasal polyps. It can cause blocked breathing through the nose, drainage, reduced smell, facial pressure, or recurrent sinus symptoms.

### **Chronic Sinusitis**

Persistent inflammation of the paranasal sinuses for at least 12 weeks, causing nasal obstruction, discharge, facial pressure, and reduced smell.

#### **Alternate Names:**

Chronic Rhinosinusitis; CRS

### **Chronic Spontaneous Urticaria**

The persistent occurrence of itchy, erythematous wheals (hives), angioedema, or both for six weeks

---

or longer without an identifiable external physical or allergic trigger, frequently driven by autoimmune autoantibodies against IgE or its high-affinity receptor (Fc-epsilon-RI).

**Alternate Names:**

CSU; Chronic Idiopathic Urticaria

### **Chronic Stiff Neck**

Persistent reduced movement or tightness in the neck. Muscle strain, arthritis, disc disease, inflammation, or nerve problems may be responsible; fever, injury, weakness, numbness, or a severe headache requires prompt medical assessment.

### **Chronic Subdural Hematoma**

A collection of blood and breakdown products between the dura and arachnoid that develops over weeks, often after minor trauma in older adults, causing headache, confusion, weakness, or gait change.

### **Chronic Suppurative Otitis Media**

Persistent inflammation and infection of the middle ear with recurrent or continuous ear discharge, usually through a perforated tympanic membrane. It can cause conductive hearing loss and may lead to cholesteatoma or other complications.

**Alternate Names:**

CSOM; Chronic Otitis Media with Discharge; Chronic Suppurative Middle-Ear Infection

### **Chronic Tension-Type Headache**

A recurring headache occurring on at least 15 days per month for more than three months, usually causing bilateral pressing or tightening pain rather than throbbing. It may be associated with scalp or neck tenderness but typically lacks prominent nausea; a new or changing pattern requires clinical assessment.

### **Chronic Thromboembolic Pulmonary Hypertension**

High blood pressure in the lung arteries caused by old blood clots that remain attached and block blood flow after a pulmonary embolism. It can strain the right side of the heart and may be treatable with

surgery, catheter procedures, or medicines.

### **Chronic Toxicity**

Harm caused by repeated or long-term exposure to a substance, even when each individual dose is small. Damage may build up over time and affect organs such as the brain, liver, kidneys, lungs, or nerves.

### **Chronic Traumatic Encephalopathy**

A brain disorder associated with repeated blows or jolts to the head over time. It may be linked with later changes in memory, thinking, mood, behavior, or movement; diagnosis cannot be confirmed during life by a single test.

### **Chronic Urticaria**

The occurrence of recurrent, transient, pruritic wheals, angioedema, or both for six weeks or longer. It encompasses chronic spontaneous urticaria (which occurs without an identifiable physical trigger, frequently driven by IgG autoantibodies against IgE or the high-affinity IgE receptor FcεRI on dermal mast cells) and chronic inducible urticaria (triggered by physical stimuli such as cold, heat, pressure, vibration, or sunlight). Individual wheals characteristically resolve within 24 hours without leaving permanent residual pigmentation or scarring.

**Alternate Names:**

Chronic Spontaneous Urticaria; Chronic Hives; Chronic Idiopathic Urticaria; CSU

### **Chronic Venous Disease**

Long-term difficulty returning blood from the legs to the heart because veins leak, are blocked, or do not receive enough help from leg muscles. It can cause varicose veins, swelling, skin changes, and ulcers.

### **Chronic Venous Insufficiency**

Poor return of blood from the legs because the vein valves do not close properly or the veins are blocked. It can cause swelling, aching, varicose veins, dark or hardened skin, and ulcers near the ankle.

---

### Chronic Venous Stasis

Persistent pooling of blood in the lower limbs caused by impaired venous return. It can produce edema, skin discoloration, dermatitis, and venous ulcers.

### Chronic Vulvar Pain

Alternate index label; see *Vulvodynia*.

### Chronicity

How long a disease, symptom, or process has lasted or how long it tends to continue. A condition may be described as acute when it is recent and short-lasting, or chronic when it persists over a long period.

### Chronobiology

Chronobiology studies biological timing and rhythms, including circadian sleep-wake cycles, seasonal rhythms, and their effects on health.

### Chronotropic

Chronotropic describes an influence on heart rate; a positive chronotropic effect increases rate, while a negative effect decreases it.

### Churg-Strauss Disease

Former name; see *Eosinophilic Granulomatosis With Polyangiitis*.

### Churg-Strauss Syndrome

A rare disease in which inflamed small and medium blood vessels are associated with asthma and too many eosinophils, a type of white blood cell. It can damage the lungs, nerves, skin, heart, or kidneys.

#### Alternate Names:

EGPA; Eosinophilic Granulomatosis with Polyangiitis

### Chvostek Sign

Twitching of facial muscles after tapping over the facial nerve near the ear. It can occur with low blood calcium but is not reliable enough by itself to diagnose that condition.

### Chvostek's Sign

Chvostek's sign is twitching of facial muscles after tapping near the facial nerve, sometimes associated with hypocalcemia but neither sensitive nor specific.

### Chylangioma

A usually noncancerous growth made of widened lymph vessels, often in the neck or chest. It may appear as a soft swelling and can press on the airway, food pipe, nerves, or other nearby structures; it is also called a lymphangioma.

### Chyle

A milky fluid formed in intestinal lymph vessels after a meal. It contains lymph and absorbed dietary fat and normally travels through the thoracic duct into the bloodstream.

### Chylomicron

A particle made by the intestine after eating fat. It carries dietary fat and cholesterol through the lymph and blood, then leaves remnants for the liver to remove.

### Chylomicronemia Syndrome

Severe elevation of triglyceride-rich chylomicrons in blood, usually from genetic or acquired impairment of triglyceride clearance; it can cause abdominal pain, eruptive xanthomas, and recurrent acute pancreatitis.

### Chylothorax

A buildup of chyle in the space around a lung, usually after injury or blockage of the thoracic duct. It can cause breathlessness and requires treatment of the leak, often with drainage and dietary or surgical measures.

### Chylous Ascites

Accumulation of lymphatic fluid rich in triglycerides within the abdominal cavity. It can result from lymphatic obstruction, malignancy, trauma, surgery, infection, or congenital disease and may cause abdominal distention, malnutrition, or immune-cell loss.

---

## Chyluria

Passage of lymphatic fluid into the urine, giving it a milky appearance. It usually results from an abnormal communication between lymphatic vessels and the urinary tract.

## Chyme

A semifluid mixture of partially digested food and gastric or intestinal secretions that passes from the stomach into the small intestine.

### Alternate Names:

Digestive Chyme; Gastric Chyme

## Chymotrypsin

Chymotrypsin is a pancreatic digestive enzyme that cleaves proteins in the small intestine; abnormal pancreatic release can contribute to tissue injury.

## CIC

Commonly, clean intermittent catheterization: periodic insertion of a catheter to empty the bladder, followed by removal; other expansions are possible by context.

## Cicatricial Alopecia

Hair loss caused by destruction and scarring of hair follicles, usually leading to permanent loss in affected areas.

### Alternate Names:

Scarring Alopecia; Cicatricial Hair Loss

## Cicatrization

Cicatrization is healing by formation of a scar after tissue injury, with replacement of damaged tissue by fibrous connective tissue.

## Ciclopia

A rare severe congenital malformation in which the eyes are fused or a single median eye develops, usually as part of holoprosencephaly.

## CICO Emergency

A life-threatening airway emergency in which a breathing tube cannot be placed and enough oxygen cannot be delivered by mask or another airway de-

vice. It requires immediate emergency action, often including an emergency surgical airway.

### Alternate Names:

Cannot Intubate, Cannot Oxygenate; CICO; Failed Airway Crisis; Emergency Front-of-Neck Airway Scenario

## Cidofovir

Cidofovir is an antiviral nucleotide analogue used for selected serious DNA-virus infections; kidney toxicity requires careful dosing and monitoring.

## CIDP

Chronic inflammatory demyelinating polyneuropathy is a chronic, immune-mediated, peripheral neuropathy characterized by progressive or relapsing symmetrical weakness and sensory loss in all four limbs, with areflexia or hyporeflexia, lasting for more than eight weeks.

## Cigarette

A small cylinder of tobacco or other material wrapped in paper for smoking; smoke exposure delivers nicotine and many toxic and carcinogenic substances.

## Cigarette Smoking

Alternate form; see *Smoking*.

## Ciguatera

Food poisoning caused by eating reef fish containing ciguatoxins. It may cause gastrointestinal illness followed by neurologic symptoms such as paresthesias and temperature-sensation disturbance.

## Ciguatera Poisoning

Ciguatera poisoning follows eating reef fish containing ciguatoxins. Gastrointestinal symptoms may be followed by neurologic sensations such as temperature reversal, tingling, itching, weakness, or abnormal heart rate; treatment is supportive and prevention depends on avoiding high-risk fish.

### Alternate Names:

Ciguatera Fish Poisoning (CFP); Ciguatoxicity

---

## Ciguatera Toxicity

Food poisoning from ciguatoxins in contaminated reef fish. It typically causes early nausea, vomiting, diarrhea, and abdominal pain followed by perioral or distal paresthesias, itching, weakness, and temperature reversal; bradycardia or hypotension can occur. Treatment is supportive because no reliably proven antidote exists.

## Ciguatera Toxin

Alternate name; see *Ciguatera poisoning*.

## Ciguatoxin

A heat-stable toxin produced by marine microorganisms and accumulated in some reef fish, causing ciguatera poisoning with gastrointestinal, neurologic, and temperature-sensation symptoms.

## Cilia

Tiny hair-like structures on cells that move fluid, mucus, or particles across a tissue surface. In the airways, they help clear inhaled material; some cilia also sense signals and help move an egg cell.

## Ciliary Body

A ring-shaped part of the eye between the iris and choroid. Its muscle changes the lens shape for focusing, and its processes make the fluid that fills the front of the eye.

## Ciliary Body Melanoma

A melanoma arising in the uveal tract behind the iris. It may be asymptomatic or cause blurred vision, floaters, pain, or a visible change in the pupil or iris; ophthalmic examination and imaging guide diagnosis and eye-preserving or surgical treatment.

## Ciliary Dyskinesia

Abnormal movement of cilia that impairs clearance of mucus from the airways and can cause recurrent respiratory infections.

## Ciliary Flush

A violaceous, ring-like injection of deep episcleral vessels encircling the corneal limbus, serving as a

cardinal physical sign of active intraocular inflammation such as iritis, keratitis, or acute angle-closure glaucoma.

### Alternate Names:

Perilimbal Injection; Ciliary Injection

## Ciliary Muscle

A ring of smooth muscle in the eye that changes the lens shape for focusing at different distances and helps regulate aqueous humor outflow through ciliary-body structures.

## Ciliary Neuralgia

Pain arising from or felt along the ciliary nerves around the eye, often described as deep orbital or eye pain; the cause requires clinical evaluation.

## Ciliopathy

A disorder caused by abnormal cell cilia, the tiny structures that move material or sense signals. It may affect development, vision, kidney function, breathing, or several organs.

## Cimetidine

Cimetidine is an H<sub>2</sub>-receptor antagonist that reduces gastric acid secretion; it has numerous drug interactions and is used less often than newer agents.

## Cinacalcet

A calcimimetic drug that activates calcium-sensing receptors and lowers parathyroid hormone, used for selected forms of hyperparathyroidism and hypercalcemia.

## Cineangiocardiology

Radiographic recording of the heart and blood vessels during passage of contrast material, used to visualize cardiac chambers, valves, and coronary or great vessels.

## Cineangiography

Motion-picture X-ray imaging of the heart or blood vessels while contrast material passes through them, used to show blood flow and moving cardiac structures.



---

## Cinefluorography

Motion-picture recording of fluoroscopic images, used to document movement of organs or contrast material during a diagnostic examination.

## Cineplasty

Surgical creation of a muscle-controlled stump or related reconstruction, historically used to improve control of a prosthetic limb.

## Cingulotomy

A brain operation that interrupts selected nerve fibers involved in emotion and pain processing. It is reserved for rare, severe conditions that have not improved with other treatments and can cause bleeding, seizures, personality changes, or other brain injury.

## Cinnarizine

Cinnarizine is an antihistamine with calcium-channel-blocking properties used in some countries for motion sickness and vestibular symptoms.

## Ciprofloxacin

Ciprofloxacin is a fluoroquinolone antibiotic used for selected bacterial infections; tendon injury, nerve effects, QT effects, and resistance limit indiscriminate use.

## Circadian Rhythm

An internal cycle of about 24 hours that helps regulate sleep, alertness, hormones, body temperature, and metabolism. Light and darkness reset the cycle each day through signals from the eyes to the brain.

## Circadian Rhythm Sleep Disorder

A family of sleep disturbances arising from a persistent misalignment between an individual's endogenous biological circadian rhythm and external societal light-dark cycles, including delayed sleep phase disorder, advanced sleep phase disorder, shift-work disorder, and jet lag.

## Circadian Rhythms

Approximately 24-hour biological cycles that regu-

late sleep and wakefulness, hormone release, body temperature, metabolism, and other physiological functions in response chiefly to light and darkness.

## Circle of Willis

A ring-shaped network of arteries at the base of the brain that connects the front and back blood supplies. It may provide an alternate route for blood if an artery narrows or becomes blocked, but its structure varies and it may not provide enough backup flow.

## Circulating Tumor Cell

A cancer cell that has detached from a tumor and entered the bloodstream. Testing may help study cancer spread or treatment response, but clinical use depends on the specific validated test and cancer.

## Circulating Tumor DNA

Small fragments of DNA released by cancer cells into the blood. Testing them may help identify tumour changes, monitor treatment, or detect remaining disease, but it does not replace tissue diagnosis in every situation.

## Circulation

Continuous movement of blood through the heart and blood vessels, or of lymph through lymphatic vessels, to transport oxygen, nutrients, hormones, and waste.

## Circulatory

Relating to the movement of blood or lymph and to the organs and vessels that carry them.

## Circulatory System

The heart and blood-vessel network that pumps and distributes blood through pulmonary and systemic circuits; broadly, the term may include the lymphatic system.

## Circumcision

Surgical removal of the foreskin covering the head of the penis. It may be performed for cultural, religious, preventive, or medical reasons; possible

---

complications include bleeding, infection, pain, and injury, although serious problems are uncommon with proper care.

### **Circumduction**

A compound, sequential circular joint movement combining flexion, abduction, extension, and adduction such that the distal end of an extremity moves in a circle while the proximal shaft delineates a cone, seen at ball-and-socket joints such as the shoulder and hip.

### **Circumflex**

Curving around; the circumflex coronary artery is a branch of the left coronary artery that travels around the heart's lateral and posterior surface.

### **Circumflex Artery**

A branch of the left coronary artery that runs around the left side and back of the heart. It supplies blood to the left atrium and much of the outer and back wall of the left ventricle.

### **Circumvallate Papillae**

Circumvallate papillae are large taste papillae arranged near the back of the tongue and surrounded by a trench containing taste buds and serous glands.

### **Circumvallate Placenta**

An extrachorial placental configuration in which the chorionic plate is smaller than the basal plate, producing a raised circumferential edge; it may be associated with bleeding, abruption, or growth restriction.

### **Cirrhosis**

Severe scarring of the liver caused by long-term liver damage. Scar tissue replaces healthy liver tissue and can reduce liver function and blood flow. It may cause no symptoms at first, but later may cause tiredness, poor appetite, weight loss, itching, jaundice, swelling of the abdomen or legs, easy bruising or bleeding, or confusion. Severe cirrhosis may cause internal bleeding, liver failure, or liver cancer.

### **Cirrhosis Of The Liver**

Alternate name; see *Cirrhosis*.

### **Cisplatin**

A platinum-based chemotherapy drug that forms DNA cross-links and is used to treat several cancers; important toxicities include kidney injury, hearing loss, neuropathy, and nausea.

### **Cisterna Chyli**

A widened, sac-like part of the lymphatic system in the upper abdomen. It receives lymph from the lower body and intestines and continues upward as the thoracic duct.

### **Cisterna Magna**

The largest subarachnoid cistern, located in the posterior cranial fossa between the inferior surface of the cerebellum and the dorsal aspect of the medulla oblongata, which receives cerebrospinal fluid directly from the fourth ventricle via the median aperture (foramen of Magendie).

#### **Alternate Names:**

Cerebellomedullary Cistern

### **Citalopram**

Citalopram is a selective serotonin-reuptake inhibitor used to treat depression and some anxiety disorders; nausea, sexual effects, and QT prolongation are important considerations.

### **Citrate**

Citrate is a salt or ion of citric acid that participates in metabolism and binds calcium; it is used in anticoagulant blood collection and some stone-prevention treatments.

### **Citrate Synthase**

An enzyme that begins an important energy-producing process inside cells. It joins acetyl-CoA with oxaloacetate to make citrate, the first step of the citric acid cycle, which helps release energy from food.

---

### Citric Acid Cycle

A series of chemical reactions inside mitochondria, the cell's energy-making structures, that breaks down food-derived molecules and produces energy. It also makes substances needed to build proteins, fats, and other cell components.

### Citric Acid Urine Test

A urine test measuring citrate excretion, used mainly in the evaluation of recurrent kidney stones because low urinary citrate increases the risk of calcium stone formation.

### Citrullinaemia

An inherited disorder of the urea cycle causing citrulline and ammonia buildup, which can lead to vomiting, confusion, seizures, coma, or death.

### Citrullinemia

An inherited disorder in which the body cannot properly remove nitrogen through the urea cycle. Citrulline and ammonia build up, causing vomiting, poor feeding, sleepiness, confusion, seizures, coma, or death without urgent treatment.

### CJD

Alternate name; see *Creutzfeldt-Jakob Disease*.

### CK

Creatine kinase, an enzyme found in skeletal muscle, heart muscle, and brain whose blood level may rise after tissue injury.

#### Alternate Names:

Creatine Phosphokinase; Creatine Kinase Enzyme

### CKD-Mineral Bone Disorder

A complication of chronic kidney disease in which abnormal calcium, phosphate, vitamin D, and parathyroid hormone levels weaken bones and may cause calcium deposits in blood vessels and other tissues.

### Cladribine

Cladribine is a purine analogue used for selected blood cancers and multiple sclerosis; lymphocyte depletion and infection risk require monitoring.

### Clagett Procedure

A staged surgical procedure for chronic pleural empyema with bronchopleural fistula, consisting of open thoracic fenestration (open window thoracostomy) to drain the pleural space, followed by sequential cavity debridement, antibiotic irrigation, and eventual secondary chest wall closure.

#### Alternate Names:

Open Window Thoracostomy Closure; Clagett Window

### Clammy Skin

Skin that feels cool, damp, or unusually sweaty. It may occur with heat, anxiety, pain, low blood sugar, shock, or other illness.

### Clamp

A surgical instrument used to grasp, compress, occlude, or hold tissue, blood vessels, tubing, or other structures.

### Clang Association

Clang association is speech in which word choice is driven by sound, rhyme, or pun rather than meaning, seen particularly in mania or psychosis.

### Clarithromycin

Clarithromycin is a macrolide antibiotic used for selected bacterial and mycobacterial infections; interactions and QT prolongation are important risks.

### Clark Level Of Invasion

An older system describing how deeply a melanoma has grown through the skin layers. Modern melanoma assessment relies more on tumor thickness, ulceration, spread, and other features.

### Class IA Antiarrhythmics

Heart-rhythm medicines that slow electrical conduction and lengthen the time heart cells need before they can fire again. They treat selected abnormal rhythms but can themselves cause dangerous rhythm disturbances.

---

### **Class IB Antiarrhythmics**

Heart-rhythm medicines that reduce abnormal electrical activity, especially in injured heart muscle. They are used mainly for selected fast rhythms arising from the lower chambers and require monitoring.

### **Class IC Antiarrhythmics**

Strong heart-rhythm medicines that slow electrical conduction without greatly changing recovery time. They can treat selected upper-chamber rhythms but may be dangerous in people with structural heart disease.

### **Class III Antiarrhythmics**

Heart-rhythm medicines that lengthen the recovery period between electrical beats. They treat selected abnormal rhythms but may slow the heart, affect organs, or prolong the electrical interval and cause dangerous rhythms.

### **Classic Galactosemia**

A serious inherited disorder caused by severe deficiency of galactose-1-phosphate uridylyltransferase. Milk feeding can rapidly cause jaundice, liver dysfunction, hypoglycemia, infection, and neurologic injury; lifelong galactose restriction is required.

### **Classic Migraine**

A migraine attack preceded or accompanied by an aura, such as temporary visual, sensory, or speech changes. Headache may follow, with nausea and light or sound sensitivity, although patterns vary.

### **Classic Type Pfeiffer Syndrome**

A genetic craniosynostosis syndrome, usually related to an *FGFR2* variant, causing early fusion of skull sutures, broad or deviated thumbs and great toes, and variable facial, airway, neurologic, or developmental problems.

### **Clathrin**

A major polymeric coat protein composed of three heavy and three light chains forming a three-legged triskelion, which self-assembles into a polyhedral basket-like lattice that coats and internalizes endocytic vesicles during receptor-mediated endocytosis.

### **Claude Syndrome**

A brainstem stroke syndrome caused by an ischemic or hemorrhagic infarction of the dorsal midbrain tegmentum affecting the oculomotor nerve fascicles, red nucleus, and cerebellothalamic tract. Clinical features include ipsilateral third cranial nerve palsy (ptosis, dilated pupil, and ophthalmoplegia) and contralateral cerebellar ataxia, tremor, and dysmetria.

#### **Alternate Names:**

Claude's Syndrome; Inferior Rubro-Spinal Syndrome

### **Claudication**

Pain, aching, cramping, or weakness in a limb brought on by walking or other exertion. Vascular claudication, usually caused by reduced arterial blood flow, improves with rest; neurogenic claudication, often caused by lumbar spinal stenosis, typically improves with sitting or leaning forward.

#### **Alternate Names:**

Vascular Claudication

### **Claudication Distance**

The distance a person can walk before exertional limb pain begins. It is used to describe symptoms and monitor change.

### **Claustrophobia**

An intense fear of enclosed or confined spaces that may cause anxiety, panic, or avoidance. Symptoms can occur in places such as elevators, tunnels, or MRI scanners.

### **Clavicle**

The collarbone, a curved bone running from the breastbone to the shoulder blade. It supports the shoulder and commonly breaks after a fall or direct blow, causing pain, swelling, or visible deformity.

### **Clavicle Fracture**

A break in the collarbone that may occur during birth or after trauma. A newborn may show asymmetric arm movement or tenderness; older children and adults may have pain, swelling, or deformity. Diagnosis is usually made by examination and imag-

---

ing.

### **Clavus**

A clavus is a localized painful thickening of skin, or corn, caused by repeated pressure or friction.

### **Claw Foot**

A toe deformity in which the toes bend upward at their bases and downward at their middle and end joints. It may result from muscle imbalance, nerve damage, arthritis, or unsuitable footwear.

### **Claw Hand**

A deformity in which the large finger joints bend backward while the smaller joints bend inward. It is commonly caused by ulnar-nerve damage and may impair grip and hand function.

### **Claw Toe**

A toe deformity characterized by extension at the metatarsophalangeal joint and flexion at the proximal and distal interphalangeal joints.

### **Clay-Colored Stool**

Pale, gray, or white stool caused by reduced bile reaching the intestine. It can occur with bile-duct obstruction, liver disease, or pancreatic disease and requires evaluation, especially when accompanied by jaundice or dark urine.

#### **Alternate Names:**

Acholic Stool; Pale Stool

### **Clean Catch Urine Sample**

A midstream urine specimen collected after cleaning the genital area to reduce contamination and improve interpretation of urinalysis or culture.

### **Clean-Catch Specimen**

A urine sample collected after cleansing the surrounding area and beginning urination, with the middle portion collected in a sterile container to reduce contamination.

### **Cleaning Supplies And Equipment**

Materials and devices used to remove soil and mi-

croorganisms from surfaces or equipment; selection and disinfection level depend on the item and infection risk.

### **Clear Cell Carcinoma**

Clear cell carcinoma is a cancer whose cells appear clear because of glycogen or lipid; it can arise in organs such as kidney, ovary, or endometrium.

### **Clear Cell Carcinoma Of The Ovary**

A type of ovarian cancer whose cells look clear under a microscope. It is often associated with endometriosis and may be found at an earlier stage, but can resist some standard treatments and increase the risk of blood clots; treatment depends on stage and overall health.

### **Clear Cell Renal Cell Carcinoma**

The most common type of kidney cancer in adults. The cancer cells look clear under a microscope because they contain stored fat and sugar; it may cause blood in the urine, pain, or a mass, but early disease may cause no symptoms.

#### **Alternate Names:**

CcRCC; Conventional Renal Cell Carcinoma

### **Clear Cell Renal Cell Carcinoma Pathology**

A renal carcinoma composed of clear cells containing lipid and glycogen, commonly arranged in nests or sheets separated by a delicate branching vasculature. Diagnosis and grading depend on architecture, cytology, necrosis, and invasion.

### **Clear Liquid Diet**

A limited diet consisting of transparent liquids such as water, clear broth, apple juice, and black coffee that leave minimal residue. It is used for short periods before procedures, during acute illness, or after surgery to maintain hydration while resting the gastrointestinal tract.

### **Clear Margins**

A laboratory finding that no abnormal or cancer cells are seen at the edge of tissue removed during a procedure. It suggests the abnormal area may have

---

been completely removed, but it does not by itself rule out disease elsewhere.

### **Clearance**

The rate at which the body removes a medicine or other substance from the blood. The liver, kidneys, and sometimes other organs contribute; clearance helps determine how much medicine is needed and how often it should be given.

### **Cleft Lip**

A birth defect in which the upper lip does not join completely before birth. The opening may be small or extend toward the nose and may occur alone or with a cleft palate, affecting feeding, teeth, or speech.

### **Cleft Lip And Cleft Palate**

Birth defects in which the upper lip, the roof of the mouth, or both do not join together fully before birth. They can cause problems with feeding, speech, hearing, or teeth and may need treatment by several specialists.

### **Cleft Lip And Palate Repair**

Surgery to close a cleft in the lip, palate, or both and improve feeding, speech, hearing, breathing, or appearance. Repair is usually staged during childhood and requires long-term dental and specialist follow-up.

### **Cleft Palate**

A birth defect in which there is an opening in the roof of the mouth. It may occur alone or with a cleft lip and can cause problems with feeding, speech, hearing, or teeth.

### **Cleidocranial Dysostosis**

An inherited bone disorder, usually caused by a RUNX2 gene change, in which the collarbones may be small or absent and skull bones close late. Extra teeth, delayed tooth eruption, short stature, and joint problems are common.

### **Cleidocranial Dysplasia**

An inherited disorder of bone development affecting the collarbones and skull. It can cause delayed closure of skull joints, unusual teeth, short stature, and other skeletal differences.

### **Clemastine**

Clemastine is a first-generation antihistamine used for allergic symptoms; sedation, anticholinergic effects, and impaired driving are concerns.

### **Click-Murmur Syndrome**

Alternate name; see *Mitral Valve Prolapse*.

### **Climacteric**

The period of physiologic change around the end of reproductive capacity, including the menopausal transition. It may involve changes in menstruation, vasomotor symptoms, and other endocrine effects.

### **Climate Change**

Long-term changes in temperature, precipitation, and weather patterns, largely driven by human-produced greenhouse gases. Health effects include heat illness, worsened air quality, infectious-disease changes, injuries, and effects on food, water, and mental health.

### **Clindamycin**

Clindamycin is a lincosamide antibiotic used for selected infections; diarrhea and *Clostridioides difficile* colitis are important adverse effects.

### **Clinical**

Relating to direct observation, examination, diagnosis, or treatment of a patient, rather than solely to laboratory or basic research findings.

### **Clinical Communication**

Exchange of information between healthcare professionals, patients, and caregivers to support diagnosis, treatment, safety, and informed decisions.

### **Clinical Counseling**

Guidance provided in healthcare to help patients



---

make informed decisions about health, prevention, treatment, and self-management.

### **Clinical Depression**

A depressive illness causing persistent sadness or loss of interest, together with changes in sleep, appetite, energy, thinking, or self-worth that significantly interfere with daily life.

### **Clinical Deterioration**

Worsening health that increases the risk of organ failure, disability, or death. Warning signs may include new confusion, breathing difficulty, abnormal vital signs, reduced urine, worsening pain, or poor circulation.

### **Clinical Diagnostics**

The use of history, examination, laboratory tests, imaging, pathology, and other methods to identify or evaluate a health condition and guide care.

### **Clinical Disease**

Disease that produces recognizable signs or symptoms or other findings detectable in a patient, as opposed to a silent or subclinical state.

### **Clinical Frailty Scale**

A scale used by a healthcare professional to describe how fit, independent, or frail an older person is. It considers activity, daily function, and illness burden and supports care planning.

### **Clinical Handover**

The transfer of responsibility for a patient's care from one healthcare professional or team to another. A safe handover states the diagnosis, current condition, risks, pending tasks, and what to do if the patient worsens.

### **Clinical Imaging**

Medical imaging used to visualize anatomy, physiology, or disease, including radiography, ultrasound, computed tomography, magnetic resonance imaging, and nuclear medicine.

### **Clinical Nutrition**

The use of nutrition science to prevent, identify, and manage health problems. It includes assessment of food intake and nutritional status, dietary treatment, and nutrition support when ordinary eating is not enough or is unsafe.

### **Clinical Pharmacology**

The medical science of medicines and their effects in people. It includes how medicines work, how the body absorbs and removes them, how they are used, and how their benefits, interactions, and risks are assessed.

### **Clinical Presentation**

The symptoms, signs, examination findings, and test results seen in a person with a health problem. The presentation can differ with the cause, severity, age, other illnesses, and the time since the problem began.

### **Clinical Psychologist**

A licensed mental-health professional who assesses and treats psychological, emotional, behavioral, and cognitive problems using psychotherapy and other evidence-based interventions within local scope of practice.

### **Clinical Reasoning**

The process healthcare professionals use to combine a person's symptoms, examination, test results, history, and risks when deciding what diagnosis is most likely and what care is appropriate.

### **Clinical Trial**

A planned study in which people receive or are observed for an intervention to evaluate its safety, effects, or outcomes. Trials may be conducted in phases and for many types of health interventions.

### **Clinical Trial Phases**

Stages for testing a medical treatment: Phase I mainly examines safety and dose, Phase II studies early effectiveness and safety, Phase III compares it with usual care in larger groups, and Phase IV monitors longer-term use.

---

## Clinicopathologic Correlation

Combining a person's symptoms, examination, scans, laboratory tests, and tissue findings to reach the most accurate diagnosis, especially when a small sample may not show the whole disease.

## Clinistix

Clinistix is a reagent-strip method historically used to detect glucose in urine through an enzymatic color reaction.

## Clinitest

Clinitest is a tablet-based copper-reduction test historically used to detect reducing substances in urine, including glucose; it is less specific than modern methods.

## Clinodactyly

A finger or toe that curves sideways from birth or during development, most often the little finger. It is usually mild but may interfere with movement or occur with a genetic condition.

## Clinoid Process

A small bony projection at the base of the skull near the eye socket and major nerves and blood vessels. It helps form the support around the pituitary region and optic canal.

## CLIP

A device used to hold tissue or structures together; surgical clips can close blood vessels, mark tissue, or occlude structures such as the vas deferens.

## Clitoral

Relating to the clitoris, an erectile organ with a dense concentration of sensory nerves at the front of the vulva.

## Clitorectomy

Surgical removal of all or part of the clitoris, sometimes with removal of adjacent vulvar tissue; nonconsensual forms constitute female genital mutilation and can cause serious harm.

## Clitoridectomy

Surgical removal of part or all of the clitoris. In medicine it may be performed for selected tumors or severe disease; nonmedical removal is a harmful practice with no health benefit.

## Clitoris

A small, sensitive erectile organ located at the anterior junction of the labia minora, beneath the clitoral hood (prepuce). The clitoris is the primary source of female sexual pleasure and is homologous to the male penis.

## Clitorodynia

Persistent or recurrent pain, burning, or marked sensitivity of the clitoris. Causes may include infection, skin disease, nerve pain, trauma, pelvic-floor dysfunction, or another vulvar pain disorder.

## Clitoromegaly

Clitoromegaly is enlargement of the clitoris, which may be congenital or acquired from androgen excess, endocrine disease, medication, or other causes.

## Clivus

A central, sloped bony surface on the floor of the posterior cranial fossa formed by the fusion of the basilar part of the sphenoid bone and the basiocciput, extending anterior to the foramen magnum and providing a bed for the pons and basilar artery.

## CLL

Alternate name; see *Chronic Lymphocytic Leukemia*.

## Cloacal Abnormalities

A group of rare birth-related conditions in which the urinary, genital, and intestinal structures do not separate or develop normally. They may affect the bladder, bowel, spine, external genitalia, and reproductive organs.

## Cloacal Exstrophy

A very rare birth defect in which the lower abdominal wall, bladder, bowel, spine, and reproductive or urinary organs do not form normally and may be

---

exposed. Newborns need urgent specialist surgery and long-term coordinated care.

### **Cloacitis**

Inflammation of a cloaca or cloacal region, particularly in disorders affecting the common terminal outlet of the urinary, genital, and intestinal tracts.

### **Clodronate**

Clodronate is a bisphosphonate that reduces bone resorption and is used in selected bone and cancer-related conditions.

### **Clofazimine**

Clofazimine is a riminophenazine antimicrobial used mainly in multidrug treatment of leprosy and selected mycobacterial infections; skin discoloration is common.

### **Clomifene**

Clomifene, or clomiphene, is a selective estrogen-receptor modulator used to induce ovulation in selected infertility settings.

### **Clomipramine**

Clomipramine is a tricyclic antidepressant particularly effective for obsessive-compulsive disorder; anticholinergic, cardiac, and overdose toxicity are important.

### **Clonal Evolution**

The gradual change of a cancer as groups of tumor cells acquire different genetic changes. Groups that grow faster, spread more easily, or resist treatment can become dominant and make the cancer progress or recur.

### **Clonal Expansion**

Multiplication of cells descended from one original cell. It may occur when immune cells respond to a trigger or when an abnormal cell population grows in cancer or another disorder.

#### **Alternate Names:**

Clonal Proliferation; Clone Expansion

### **Clonazepam**

Clonazepam is a benzodiazepine anticonvulsant and anxiolytic that enhances GABA activity; sedation, dependence, and respiratory depression can occur.

### **Clonic Seizure**

A seizure causing repeated, rhythmic jerking of muscles. It may affect one part of the body or both sides, with or without loss of awareness; the cause and seizure type are determined from the episode and medical assessment.

### **Clonidine**

Clonidine stimulates central alpha-2 receptors to reduce sympathetic outflow and is used for hypertension and selected other conditions; abrupt withdrawal can cause rebound hypertension.

### **Clonidine Toxicity**

Alpha-2-adrenergic agonist poisoning causing somnolence, miosis, bradycardia, hypotension, respiratory depression, and sometimes transient hypertension early after ingestion. Management is supportive with airway and cardiovascular monitoring; naloxone may improve opioid-like central depression in selected cases.

### **Clonorchiasis**

Infection with the Chinese liver fluke, *Clonorchis sinensis*. It affects the biliary tract and may cause inflammation, obstruction, and long-term hepatobiliary complications.

### **Clonus**

Repeated, rhythmic muscle jerks triggered by suddenly stretching a muscle, often at the ankle. It commonly indicates overactive reflexes from a problem affecting the brain or spinal cord.

### **Clopidogrel**

Clopidogrel is an antiplatelet drug that irreversibly blocks the platelet P2Y<sub>12</sub> receptor, reducing arterial thrombosis risk after selected cardiovascular events or procedures.

---

### **Closed Comedo**

A small skin lesion formed when a follicular opening is blocked by sebum and keratin but remains covered by the epidermis. It is commonly called a whitehead.

#### **Alternate Names:**

Whitehead

### **Closed Fracture**

A fracture in which the broken bone does not communicate through a wound with the external environment.

### **Closed Reduction**

Nonoperative realignment of a fractured or dislocated bone or joint by manipulation without surgically opening the site.

### **Closed Reduction Of A Fractured Bone**

A nonsurgical procedure that realigns a broken bone by manipulating it through the skin without opening the fracture site, usually followed by immobilization and imaging to confirm alignment.

### **Closed Suction Drain With Bulb**

A flexible drain connected to a compressible bulb that creates gentle negative pressure to remove blood or fluid from a surgical or injured area; the bulb is emptied and output recorded as directed.

### **Closed Surgical Wound Care**

Care of an incision whose edges are closed with sutures, staples, adhesive, or strips, including keeping it clean and dry as instructed and watching for infection or separation.

### **Closed-Angle Glaucoma**

Glaucoma caused by blockage of the eye's drainage angle, so fluid cannot leave and pressure rises. Sudden attacks can cause severe eye pain, blurred vision, halos, headache, nausea, and permanent sight loss.

### **Closed-Loop Insulin Delivery**

Automated insulin delivery systems that combine continuous glucose monitoring with an insulin pump and control algorithm. The algorithm adjusts basal

insulin delivery based on sensor glucose values in real time. Hybrid closed-loop systems require user-initiated meal boluses but automatically manage basal rates.

### **Clostridial Infection**

Infection caused by bacteria in the Clostridiaceae group, including *Clostridium* and *Clostridioides* species. These infections may cause antibiotic-associated colitis, food poisoning, botulism, tetanus, or rapidly destructive wound and muscle infections; symptoms, transmission, and treatment depend on the species and toxins involved.

#### **Alternate Names:**

Clostridial Infections; *Clostridium* Infection

### **Clostridial Myonecrosis**

A rapidly progressive, life-threatening necrotizing infection of skeletal muscle caused primarily by *Clostridium perfringens* or other histotoxic clostridia. Typically arising after deep traumatic injuries, compound fractures, or surgery, it is characterized by severe pain out of proportion to exam findings, rapid tissue edema, bronze skin discoloration, bullae, palpable gas crepitus, and systemic toxemia.

#### **Alternate Names:**

Gas Gangrene; *Clostridial* Gangrene

### ***Clostridioides Difficile***

A spore-forming bacterium that can overgrow after antibiotics disturb normal gut bacteria. Its toxins inflame the colon and cause watery diarrhea, abdominal pain, fever, and sometimes life-threatening colitis.

### ***Clostridioides Difficile* Infection**

Diarrhea and colon inflammation caused by overgrowth of *Clostridioides difficile*, often after antibiotic use. It may be mild or severe and can recur; diagnosis requires appropriate stool testing in a person with symptoms.

### ***Clostridioides Difficile* Pathogenesis**

Antibiotics can remove protective bowel bacteria, allowing *C. difficile* to multiply. Its toxins injure

---

the colon and cause inflammation and diarrhoea, while hardy spores can survive in the environment and spread infection.

### **Clostridium**

A genus of anaerobic, usually spore-forming bacteria that includes species capable of causing diseases such as tetanus, botulism, gas gangrene, and antibiotic-associated colitis.

### **Clostridium Botulinum**

A bacterium that produces botulinum toxin, which blocks nerve signals to muscles and causes weakness or paralysis. Food, wounds, or swallowed spores can cause botulism, a medical emergency affecting breathing and swallowing.

### **Clostridium Difficile**

Former taxonomic name; see *Clostridioides Difficile*.

### **Clostridium Difficile Colitis**

*Clostridioides difficile* colitis is antibiotic-associated inflammation of the colon caused by toxin-producing bacteria. It causes watery diarrhea, abdominal pain, fever, and sometimes severe colitis; diagnosis and treatment depend on symptoms, testing, recurrence, and severity.

#### **Alternate Names:**

C. Diff Colitis; *Clostridioides Difficile*-Associated Diarrhea (CDAD); Pseudomembranous Colitis

### **Clostridium Difficile Infection**

Alternate name; see *C. Difficile Infection*.

### **Clostridium Perfringens**

A bacterium that can cause rapidly destructive muscle infection with gas in tissues or food poisoning with sudden diarrhea and abdominal cramps. Severe wound infection requires emergency surgery and antibiotics.

### **Clostridium Perfringens Food Poisoning**

An acute foodborne toxico-infection caused by ingestion of large numbers of vegetative cells of *Clostridium perfringens* type A that sporulate in the intes-

tine and produce enterotoxin. It typically develops 6 to 24 hours after eating contaminated cooked meats, poultry, or gravies held at warm temperatures, causing intense abdominal cramps and watery diarrhea that usually subside within 24 hours.

### **Clostridium Tetani**

A soil bacterium whose toxin causes tetanus after entering a wound. It produces painful muscle stiffness and spasms, often beginning in the jaw, and can interfere with swallowing and breathing; vaccination prevents it.

### **Clot**

A semisolid mass of blood formed during clotting. A clot can stop bleeding, but one that blocks a blood vessel can reduce blood flow or travel elsewhere and cause serious injury.

### **Clot Buster**

A medicine that breaks down a blood clot, also called a thrombolytic medicine. It may be used urgently for selected strokes, heart attacks, or lung clots, but it can cause dangerous bleeding and is not safe for everyone.

### **Cloth Dye Poisoning**

Exposure to cloth dyes that may cause irritation, gastrointestinal symptoms, allergic reactions, or systemic toxicity depending on the chemical composition.

### **Clotrimazole**

Clotrimazole is an azole antifungal used topically or orally in selected formulations for fungal infections of skin, mouth, or vagina.

### **Clotting**

The body's process of stopping bleeding by narrowing a damaged vessel, forming a platelet plug, and producing a protein mesh that strengthens the plug. Excessive or inappropriate clotting can block blood vessels.

---

### Cloudy Cornea

Loss of corneal transparency from edema, scarring, infection, dystrophy, trauma, or deposits, potentially causing blurred vision, pain, photophobia, or glare.

### Clove Oil

Clove oil contains eugenol and is used in some dental and topical products; concentrated ingestion or exposure can cause irritation, liver injury, or seizures.

### Clozapine

Clozapine is an atypical antipsychotic for treatment-resistant schizophrenia and selected suicidality risk; neutropenia, myocarditis, seizures, constipation, and metabolic effects require monitoring.

### Club Drugs List And Side Effects

Club drugs include substances such as MDMA, ketamine, GHB, and others used recreationally in social settings. Effects may include agitation, hallucinations, seizures, dangerous temperature elevation, respiratory depression, cardiovascular toxicity, impaired consent, and overdose.

### Club Foot

A congenital foot deformity in which the foot is turned inward and downward and the calf may be smaller. It is usually treated early with serial casting, bracing, and sometimes surgery.

### Clubbing

Enlargement and rounding of the fingertips or toes with more curved nails and loss of the normal nail angle. It may accompany long-term lung, heart, bowel, liver, or other disease and should be medically assessed.

### Clubbing Of The Fingers Or Toes

Bulbous enlargement of fingertip or toe soft tissue with increased nail curvature caused by chronic changes in the nail bed; it can accompany lung, heart, gastrointestinal, or other disease.

### Clubbing of the Nails

Rounding and widening of the fingertips or toes with increased nail curvature and loss of the normal nail-fold angle. It may be inherited or associated with lung, heart, gastrointestinal, liver, or other disease and warrants assessment when new or progressive.

### Clubfoot

A birth defect in which a baby's foot is turned inward and downward. The foot may be stiff and difficult to move into a normal position.

### Clubfoot Repair

Treatment to gradually straighten a baby's turned-in foot, usually with gentle stretching and casts, followed by a brace. Some children need a small tendon procedure or later surgery if the deformity returns or is severe.

### Clubhand

A birth defect in which the hand and wrist curve toward the thumb or little-finger side because forearm bones or other tissues developed abnormally. Treatment depends on function and may include therapy or surgery.

### Clumping

The gathering of cells, particles, or proteins into groups. In a laboratory sample, clumped red blood cells or platelets can affect test results; in the body, clumping can sometimes block vessels.

### Cluster

A group of similar events, findings, cells, or cases occurring together in space, time, or clinical pattern.

### Cluster Headache

Repeated attacks of extremely severe pain around or behind one eye, lasting about 15 minutes to three hours. The same side may tear, redden, swell, or feel congested, and the person often becomes restless.

### Cluster Headaches

Alternate name; see *Cluster Headache*.



---

### Clutton's Joints

Painless swelling of both knees or other large joints caused by long-term inflammation in children with congenital syphilis. Joint movement is usually preserved, but the underlying infection requires treatment.

### CMTC

Abbreviation for cutis marmorata telangiectatica congenita, a congenital vascular skin anomaly causing persistent reticulated red or purple mottling. It may be isolated or associated with limb asymmetry, ulcers, or other vascular abnormalities.

### CMV

Cytomegalovirus, a common herpesvirus that may cause mild or asymptomatic infection but can cause serious disease in fetuses and people with weakened immunity.

#### Alternate Names:

Cytomegalovirus (CMV)

### CMV Blood Test

A blood test that looks for cytomegalovirus antibodies, viral proteins, or viral genetic material. Antibodies usually show past or recent exposure, while viral material can help detect active infection, especially in people with weakened immunity.

### CMV Colitis

Inflammation of the colon caused by cytomegalovirus, usually in someone with severely weakened immunity. It may cause bloody diarrhea, abdominal pain, fever, or weight loss and is confirmed with appropriate testing, often including biopsy.

### CMV Pneumonia

Pneumonia caused by cytomegalovirus, usually in people with substantial immune suppression; fever, cough, shortness of breath, and low oxygen may occur and require targeted evaluation and antiviral treatment.

### CMV Retinitis

Infection and inflammation of the retina caused by cytomegalovirus, usually in people with severely weakened immunity. It may cause floaters, blurred or missing areas of vision, and blindness unless treated urgently.

### CNS Infection

Infection of the brain, spinal cord, meninges, or cerebrospinal-fluid spaces caused by bacteria, viruses, fungi, parasites, or other organisms; presentation and treatment depend on the pathogen and anatomic site.

### CNS Lymphoma

A lymphoma involving the central nervous system, most often diffuse large B-cell lymphoma. It may involve brain parenchyma, meninges, eyes, or cerebrospinal fluid and requires urgent hematologic and neurologic evaluation.

### CO Poisoning

Illness caused by breathing carbon monoxide, an odorless gas that prevents blood from carrying oxygen normally. It may cause headache, dizziness, nausea, confusion, fainting, seizures, or death and requires immediate emergency care.

### CO<sub>2</sub> Blood Test

A blood chemistry measurement that usually estimates serum total carbon dioxide, which largely reflects bicarbonate. It helps assess acid-base and electrolyte disorders but must be interpreted with sodium, chloride, kidney function, and, when needed, blood-gas results.

### Coagulate

To change from a liquid to a semisolid or solid state, especially the formation of a blood clot through coagulation.

#### Alternate Names:

Clot; Clot Formation

### Coagulation

The body's process for stopping bleeding by forming

---

a stable blood clot. Platelets first gather at an injured vessel, then blood proteins create a mesh of fibrin that strengthens the plug; too little clotting causes bleeding, while too much can cause harmful clots.

### Coagulation Cascade

A linked series of reactions involving blood-clotting proteins that ends by forming fibrin, the mesh that strengthens a platelet plug. Problems at different steps can cause excessive bleeding or harmful clotting.

### Coagulation Defects

Problems in blood vessels, platelets, or clotting proteins that make bleeding last too long or cause harmful clots to form. Causes may be inherited, acquired, medicine-related, or due to liver or other disease.

### Coagulation Factor

A blood protein required for the coagulation cascade and formation of a stable fibrin clot; deficiencies or inhibitors can cause bleeding, while excessive activation can contribute to thrombosis.

### Coagulation Factor Deficiencies

Inherited or acquired disorders in which one or more clotting factors are reduced or function abnormally, causing excessive bleeding. The clinical pattern and laboratory findings depend on the affected factor.

### Coagulation Factor V Leiden

An autosomal dominant genetic variant (G1691A point mutation in the factor V gene) that confers resistance to proteolytic cleavage and inactivation by activated protein C (APC resistance), representing the most common inherited thrombophilia in Caucasian populations.

### Coagulation Profile

A group of blood tests that assess how quickly blood clots, commonly including prothrombin time/INR and activated partial thromboplastin time. Results help investigate bleeding, clotting, liver disease, vitamin K deficiency, and monitor selected medicines.

### Coagulation Time

The time a blood sample takes to form a clot under specified laboratory conditions. Modern tests usually measure particular clotting pathways more precisely to investigate bleeding or abnormal clotting.

### Coagulative Necrosis

A type of tissue death, usually caused by loss of blood flow, in which the tissue's basic shape remains for a while. It can occur in most solid organs but usually not in the brain.

### Coagulopathy

A problem with the body's ability to form or control blood clots. It may cause excessive bleeding or harmful clots and can result from inherited conditions, medicines, liver disease, low platelets, or severe illness.

### Coal Worker's Pneumoconiosis

Lung disease caused by breathing coal dust over many years. It may begin with small areas of dust damage and progress to extensive scarring, causing cough, breathlessness, and reduced lung function.

### Coal Workers' Pneumoconiosis (Black Lung Disease)

An occupational lung disease caused by long-term inhalation of coal mine dust. It can produce small nodules and progressive massive fibrosis, leading to cough, breathlessness, impaired lung function, pulmonary hypertension, or respiratory failure; exposure prevention is central.

### Coarctation

A localized narrowing of a tubular anatomical structure, especially coarctation of the aorta, which obstructs blood flow and produces characteristic pressure differences.

### Coarctation Of The Aorta

Alternate name; see *Aortic Coarctation*.

---

## Coarse Crackles

Wet, bubbling sounds heard through a stethoscope as air moves through fluid or secretions in larger airways. They may occur with pneumonia, bronchiectasis, chronic lung disease, or fluid in the lungs and are not specific to one condition.

## Coats Disease

An uncommon retinal disorder characterized by abnormal retinal blood vessels, leakage, exudation, and sometimes retinal detachment, usually affecting one eye in childhood.

## Cobalamin

Vitamin B12, a nutrient needed for DNA production, healthy red blood cells, nerve function, and normal processing of certain fats and proteins.

## Cobalamin Deficiency

Alternate index label; see *Vitamin B12 Deficiency*.

## Cobalt

Cobalt is a metal present in vitamin B12 and some industrial materials; excess exposure can injure the heart, thyroid, lungs, or skin.

## Cobalt Poisoning

Toxic exposure to cobalt that may cause gastrointestinal, neurologic, thyroid, cardiac, or lung effects after significant or prolonged exposure.

## Cobalt-57

A radioactive form of cobalt used as a test source for checking and calibrating gamma cameras. It is mainly used for equipment quality control and is not normally given to patients as a diagnostic or treatment medicine.

## Cobalt-Chromium

A strong, corrosion-resistant metal mixture used for cast dental restorations, partial dentures, crowns, and selected medical devices. Some people may develop sensitivity, and prolonged exposure to cobalt can be harmful.

## Cocaine

A stimulant drug that blocks monoamine reuptake and can also act as a local anesthetic; use can cause dependence, cardiovascular complications, seizures, stroke, and overdose.

## Cocaine Cardiotoxicity

Cocaine overstimulates the heart and blood vessels, raising heart rate and blood pressure and sometimes tightening the heart's arteries. It can cause chest pain, heart attack, dangerous rhythms, heart failure, or sudden death, even without longstanding artery disease.

## Cocaine Intoxication

A toxic reaction to cocaine that may cause agitation, overheating, high blood pressure, seizures, abnormal rhythms, heart attack, stroke, or sudden death.

## Cocaine Substance Use

Use of cocaine, a stimulant that can cause dependence, agitation, seizures, heart attack, stroke, dangerous overheating, and severe psychiatric or cardiovascular complications.

## Cocaine Toxicity

A toxic state caused by cocaine, often producing agitation, high blood pressure, rapid heart rate, hyperthermia, seizures, coronary vasospasm, myocardial infarction, or stroke. Treatment is supportive and may include sedation, temperature control, and management of cardiovascular or neurologic complications.

## Cocaine Use Disorder

A pattern of cocaine use that becomes difficult to control or causes problems with health, work, relationships, or safety. Craving, repeated use despite harm, tolerance, and withdrawal may occur.

### Alternate Names:

Cocaine Dependence; Cocaine Abuse; Cocaine Addiction

## Cocaine Withdrawal

Physical and emotional symptoms after stopping

---

or reducing cocaine, including fatigue, depressed mood, increased appetite, sleep changes, irritability, and cravings. Severe depression or suicidal thoughts require urgent help.

### **Cocaine-Associated Cardiomyopathy**

Cardiac muscle dysfunction associated with cocaine exposure, which may involve ischemia, myocarditis, arrhythmia, hypertension, or direct myocardial toxicity. Presentation ranges from chest pain and reduced ejection fraction to heart failure or sudden death.

### **Coccarboxylase**

Coccarboxylase is thiamine pyrophosphate, the active coenzyme form of vitamin B1 required for several oxidative carbohydrate and amino-acid reactions.

### **Coccidioides Complement Fixation**

A blood test that detects complement-fixing antibodies to *Coccidioides*; antibody levels can support diagnosis and help assess the severity or progression of coccidioidomycosis.

### **Coccidioides Immitis**

A dimorphic soil fungus and a principal cause of coccidioidomycosis, or Valley fever, acquired by inhaling airborne arthroconidia. Infection is usually pulmonary but can disseminate, especially in people with impaired cellular immunity.

### **Coccidioides Precipitin Test**

A serologic test that detects antibodies to *Coccidioides*, typically early immunoglobulin M antibodies, and supports evaluation of suspected coccidioidomycosis when interpreted with symptoms and other testing.

### **Coccidioidomycosis**

A fungal infection caused by *Coccidioides* fungi, usually acquired by breathing spores from disturbed soil. It most often affects the lungs and causes cough, fever, tiredness, or chest pain; rarely, it spreads to the skin, bones, joints, or brain.

**Common name:** Valley fever

### **Coccidioidomycosis and Valley Fever**

Infection caused by *Coccidioides* fungi, acquired by inhaling spores from disturbed soil in endemic regions. It may cause a self-limited respiratory illness, but dissemination to skin, bone, joints, or meninges can occur, especially in pregnancy or immunosuppression.

### **Coccidiosis**

Infection caused by microscopic parasites called coccidia. It usually affects the intestine and may cause watery diarrhea, stomach cramps, vomiting, dehydration, or weight loss; severe or widespread disease is more likely when immunity is weakened.

### **Coccus**

A round or nearly round bacterium. These bacteria may occur alone, in pairs, chains, or clusters; their shape helps describe them but does not identify the species or illness by itself.

### **Coccydynia**

Pain in the region of the coccyx, often worsened by sitting. Causes include trauma, childbirth, degenerative change, and less commonly infection or tumor.

#### **Alternate Names:**

Coccygodynia; Tailbone Pain

### **Coccygectomy**

Surgical removal of the coccyx, reserved for selected cases of severe persistent coccygeal pain or a lesion that has not responded to conservative treatment.

### **Coccyx**

The small pointed bone at the bottom of the spine, commonly called the tailbone. It connects to the sacrum, supports pelvic-floor tissues, and may become painful after a fall, childbirth, or prolonged sitting.

### **Cochlea**

The spiral-shaped part of the inner ear that changes sound vibrations into nerve signals. Fluid movement inside it stimulates sensory hair cells, allowing the

---

brain to recognize pitch and loudness.

### Cochlear Duct

The coiled, endolymph-filled membranous tube of the inner ear, situated between the scala vestibuli and scala tympani, housing the organ of Corti along its basilar membrane for auditory sensory transduction.

#### Alternate Names:

Scala Media

### Cochlear Implant

An electronic device for severe hearing loss that bypasses damaged inner-ear hair cells and stimulates the hearing nerve. It can provide useful sound perception but does not restore normal hearing and requires training.

### Cochlear Implants

Surgically implanted devices that convert sound into electrical signals delivered to the auditory nerve. They can provide hearing access for selected people with severe or profound sensorineural hearing loss when hearing aids are insufficient.

### Cockayne Syndrome

A rare inherited disorder in which cells cannot repair some types of DNA damage normally. It causes poor growth, developmental delay, worsening nerve problems, sensitivity to sunlight, hearing or vision loss, and an appearance of premature aging.

### Cockcroft-Gault Formula

A standardized mathematical equation used to estimate creatinine clearance (CrCl) and assess glomerular filtration rate from serum creatinine concentration, age, total body weight, and biological sex:

$$\text{CrCl} = \frac{(140 - \text{Age}) \times \text{Weight (kg)}}{72 \times \text{Serum Creatinine (mg/dL)}} \times (0.85 \text{ if female})$$

#### Alternate Names:

Cockcroft-Gault Equation; Creatinine Clearance Estimation Formula

### Code Black

A hospital emergency alert for a serious security or safety threat. The exact meaning differs between hospitals, so staff follow the facility's emergency policy.

### Code Blue

A hospital emergency signal indicating that a person may be in cardiopulmonary arrest and requires immediate resuscitation response.

### Codeine Overdose

Overdose of codeine, an opioid, that can cause respiratory depression, impaired consciousness, pinpoint pupils, and coma. Naloxone may be required in emergency care.

### Codman Triangle

A triangular area of new bone seen on an X-ray when a rapidly growing bone lesion lifts the outer covering of bone. It suggests aggressive bone disease but does not identify a specific diagnosis.

### Codon

A group of three genetic building blocks in messenger RNA that specifies one amino acid or signals the end of protein production. Several different codons can specify the same amino acid.

### Coefficient of Inbreeding

The probability that the two copies of a gene at a locus in an individual are identical by descent from a common ancestor; it is used in population and medical genetics.

### Coenzyme

A small organic molecule that helps an enzyme carry out a chemical reaction. Many coenzymes come from vitamins and temporarily carry chemical groups or electrons.

### Coenzyme Q

A fat-soluble molecule in mitochondrial membranes that transfers electrons during cellular energy production. It also acts as an antioxidant, and the body

---

makes it naturally, although levels may change with age or illness.

### **Coenzyme Q10**

CoQ10, a lipid-soluble compound in mitochondrial electron transport and an antioxidant; it is also sold as a dietary supplement, although benefits vary by condition.

### **Cofactor**

A nonprotein helper, such as a mineral ion or vitamin-derived molecule, that an enzyme needs to perform a chemical reaction. Without its cofactor, the enzyme may not work normally.

### **Coffee-Ground Emesis**

Vomit containing dark brown or black granular material from blood altered by stomach acid. It may indicate upper gastrointestinal bleeding and requires urgent medical assessment.

### **Coffin-Lowry Syndrome**

A rare inherited developmental disorder, usually linked to an RPS6KA3 gene change. It can cause intellectual disability, distinctive facial features, short stature, spinal curvature, tapered fingers, and sudden falls triggered by surprise.

**Alternate Names:**

CLS

### **Cogan'S Syndrome**

A rare immune disorder causing inflammation of the cornea and inner ear. It may produce eye pain, reduced vision, dizziness, ringing, or hearing loss and can sometimes inflame large blood vessels.

### **Cognition**

The mental processes involved in acquiring, storing, retrieving, and using information, including attention, memory, language, reasoning, and judgment.

### **Cognitive**

Relating to mental processes such as attention, memory, learning, language, reasoning, perception, judgment, problem solving, and decision-making.

### **Cognitive Behavior Therapy**

A talking treatment that helps a person notice unhelpful thoughts and behaviors and develop more useful ways to respond. It is used for problems such as anxiety, depression, pain, insomnia, and some eating disorders.

### **Cognitive Behavioral Therapy**

A structured talking therapy that helps a person identify unhelpful thoughts and behaviors and practice more useful responses. It is used for depression, anxiety, insomnia, pain, and other conditions.

### **Cognitive Behavioral Therapy For Back Pain**

A structured psychological treatment that addresses pain-related thoughts, behaviors, sleep, activity, and coping to reduce disability and improve function in chronic back pain.

### **Cognitive Behavioral Therapy For Insomnia**

A structured talking treatment for persistent insomnia. It changes unhelpful beliefs and habits about sleep and uses methods such as a regular sleep schedule, limiting time awake in bed, relaxation, and stimulus control.

### **Cognitive Behavioural Therapy**

A structured psychological treatment that helps a person identify and change unhelpful thoughts and behaviors. It is used for conditions such as anxiety, depression, insomnia, pain, and some eating disorders.

### **Cognitive Decline**

A reduction over time in one or more cognitive abilities, such as memory, attention, language, reasoning, or visuospatial function.

### **Cognitive Distortions**

Repeated patterns of inaccurate or unhelpful thinking, such as assuming the worst, seeing situations as all good or all bad, or believing one knows what others think. They can worsen anxiety or depression and may be addressed in therapy.



---

## Cognitive Fitness

The active maintenance of mental sharpness and brain health through stimulating activities, physical exercise, social engagement, and proper nutrition. Regular challenges such as learning new skills, puzzles, reading, and social interaction support neuroplasticity and cognitive reserve.

## Cognitive Function

The mental abilities used for attention, memory, language, learning, reasoning, judgment, planning, problem solving, and understanding information.

### Alternate Names:

Cognition; Cognitive Ability

## Cognitive Health And Memory

Cognitive health encompasses the ability to think clearly, learn, remember, and make decisions, with memory being a core component that involves encoding, storing, and retrieving information. Age-related changes in processing speed and recall are normal, but persistent decline may signal mild cognitive impairment or dementia.

## Cognitive Impairment

A decline in thinking abilities such as memory, attention, language, judgment, or problem solving. It may be mild with independence preserved or severe enough to interfere with everyday activities.

## Cognitive Rehabilitation

Structured therapy to improve cognitive function or help a person compensate for difficulties with memory, attention, language, or executive skills after illness or injury.

## Cognitive Reserve

The brain's ability to use alternative strategies or networks to maintain function despite injury or disease.

## Cognitive Screening

A brief assessment used to identify possible problems with memory, attention, language, or other cognitive functions. It does not by itself diagnose dementia

or another disorder.

## Cognitive Symptoms

Changes in thinking abilities such as attention, memory, processing speed, language, judgment, planning, or problem solving. They may occur with neurologic disease, mental-health conditions, medicines, sleep problems, substance use, or other medical conditions.

## Cognitive Therapy

A psychological treatment that helps a person identify and change unhelpful thoughts and beliefs and develop more adaptive behaviors; cognitive behavioral therapy combines it with behavioral methods.

## Cogwheel Rigidity

A type of muscle stiffness where moving an arm or leg feels jerky and catches like moving gears, commonly seen in Parkinson's disease.

## Cohort

A group of people who share a defined characteristic or experience during a specified period and are studied over time or compared with another group.

## Cohort Study

An observational study that identifies groups based on an exposure or other characteristic and compares the incidence of disease or other outcomes between them over time. It may be prospective or retrospective and commonly estimates relative risk.

## Coincidence Detection

The PET-scanning method of recording two gamma rays that reach opposite detectors at nearly the same time after a radioactive tracer changes inside the body. Their path helps the scanner locate where the tracer is concentrated.

## Coitus

Sexual intercourse, traditionally meaning penile-vaginal penetration; the term does not include every form of sexual activity or intimate contact.

---

### **Coitus Interruptus**

The withdrawal method of contraception, in which the penis is removed from the vagina before ejaculation; it is less reliable than many other contraceptive methods.

### **Coke**

Slang for cocaine, a stimulant drug; the word can also mean a carbon fuel product, so context is essential.

### **Cold**

Alternate name; see *Common Cold*.

### **Cold Abscess**

A pocket of pus that develops slowly with little redness, warmth, or pain. It may occur with tuberculosis or other long-term infections and still requires medical evaluation and treatment.

### **Cold Agglutinin**

An antibody that makes red blood cells stick together in cool conditions. When present in excess or unusually active, it can cause red-cell destruction and anemia.

### **Cold Agglutinin Disease**

An autoimmune blood disorder in which antibodies become active in cooler parts of the body and cause red blood cells to stick together and break down. It may cause anemia, fatigue, jaundice, or cold-related color changes.

### **Cold Agglutinin Syndrome**

A condition in which antibodies become active in cool parts of the body, making red blood cells clump and break down. It may cause anemia or blue, painful fingers and toes.

### **Cold Agglutinins**

Antibodies that attach to red blood cells in cooler conditions and may make them clump or break down. Small amounts can be harmless; high activity may cause anemia and cold-related circulation symptoms.

### **Cold Allergy**

Alternate name; see *Cold Urticaria*.

### **Cold Exposure**

Alternate index label; see *Hypothermia*.

### **Cold Hands**

Hands that feel unusually cold, which may result from environmental exposure, reduced blood flow, Raynaud phenomenon, anemia, thyroid disease, or other medical conditions.

### **Cold Injury**

Harm caused by low temperature, including hypothermia, frostbite, chilblains, and nonfreezing tissue injury. Risk increases with prolonged exposure, wetness, wind, or poor circulation and may affect the whole body or a local area.

### **Cold Intolerance**

Unusual sensitivity or inability to tolerate cold temperatures, which may occur with hypothyroidism, anemia, low body weight, vascular disease, or other conditions.

### **Cold Ischemia Time**

The duration during organ transplantation between the initiation of hypothermic organ preservation perfusion in the donor and the re-establishment of warm arterial blood reperfusion in the recipient, an important determinant of primary graft function.

### **Cold Medicines And Children**

Safety guidance for giving medicines to children with cold symptoms; many combination products provide little benefit in young children and can cause dosing errors or serious adverse effects.

### **Cold Panniculitis**

Inflammation of subcutaneous fat after exposure to cold, producing firm, red or violaceous, sometimes tender plaques or nodules. It is most often seen on the cheeks of infants or on exposed areas and usually improves with warming and avoidance of further cold.

---

## Cold Sore

A contagious infection of the lips or skin around the mouth, usually caused by herpes simplex virus type 1 and occasionally by HSV-2. It often begins with tingling or burning, followed by grouped fluid-filled blisters that crust and heal in about 1–2 weeks. The virus remains in the body and can reactivate, sometimes after illness, stress, fever, or sunlight exposure.

### Alternate Names:

Fever Blister; Herpes Labialis

## Cold Sores

Recurrent fluid-filled sores, usually on or around the lips, caused by herpes simplex virus, most often HSV-1. They are contagious from tingling or blister formation until healing and can be severe in immunocompromised people or newborns.

## Cold Urticaria

Raised, itchy patches or swelling that appear after the skin is exposed to cold. A large area of cold exposure, such as swimming in cold water, can rarely cause fainting or a life-threatening allergic reaction.

### Alternate Names:

Cold Allergy

## Colectomy

Surgery to remove part or all of the colon. It may treat cancer, severe inflammatory bowel disease, diverticular complications, blocked bowel, or inherited high cancer risk; the amount removed depends on the disease.

## Colibacillosis

Disease caused by pathogenic strains of *Escherichia coli*, which may produce intestinal infection, urinary-tract infection, sepsis, or other illness.

## Colic

Severe wave-like pain caused by contraction or blockage of a hollow organ, such as the intestine, gallbladder, or urinary tract. The term also describes repeated crying in an otherwise healthy young infant.

## Colicky Pain

Intermittent, wave-like cramping pain caused by contraction, distension, or obstruction of a hollow organ; common examples include biliary, renal, and intestinal colic.

## Coliform

A group of bacteria used as an indicator of possible contamination in water or food. Finding them does not prove that disease-causing bacteria are present, because many coliforms are harmless environmental organisms.

## Colitis

Inflammation of the colon, which may cause diarrhea, abdominal pain, blood or mucus in stool, fever, or urgency. Causes include infection, inflammatory bowel disease, reduced blood flow, medicines, and radiation.

## Collagen

A strong protein that forms much of the body's connective tissue. Different types support and strengthen skin, bone, cartilage, tendons, ligaments, blood vessels, and many internal structures.

## Collagen Disease

An older, broad term for diseases that affect connective tissues and often involve abnormal immune activity. Modern medical writing usually names the specific disease instead.

## Collagen Implant

A medical material made from or based on collagen, a structural protein, used to support tissue repair or add soft-tissue volume. Sources may be animal-derived or manufactured. Its uses, durability, and risks depend on the product and the treated body area.

## Collagenase

An enzyme that breaks down collagen, a major structural protein in skin, bone, cartilage, and connective tissue. Natural collagenases help remodel tissue, while medicines or bacterial enzymes can be used in selected treatments or cause tissue damage.

---

**Alternate Names:**

Collagen-Digesting Enzyme; Collagen-Degrading Enzyme

**Collagenous Colitis**

A type of microscopic colitis in which a thick layer of collagen develops beneath the colon lining. It usually causes persistent, watery, nonbloody diarrhea and may require colon biopsy for diagnosis.

**Collapse Of The Lung**

See *Lung Collapse*.

**Collapsed Arch**

Alternate index label; see *Flatfeet*.

**Collapsed Lung**

A common description of pneumothorax, in which air in the pleural space compresses the lung. It may cause sudden chest pain and shortness of breath and can be life-threatening when tension develops. Atelectasis is a different cause of loss of lung expansion, usually involving blockage or compression of lung tissue.

**Collapsing Glomerulopathy**

An aggressive and histologically distinctive variant of focal segmental glomerulosclerosis (FSGS) characterized by global or segmental collapse of the glomerular capillary tuft accompanied by marked hypertrophy and hyperplasia of overlying visceral epithelial cells (podocytes). Associated with human immunodeficiency virus infection (HIV-associated nephropathy), high-risk *APOL1* genotypes, interferon therapy, and severe parvovirus B19 infection.

**Alternate Names:**

Collapsing FSGS; Collapsing Focal Segmental Glomerulosclerosis

**Collapsing Pulse**

A bounding arterial pulse characterized by a rapid, forceful upstroke followed by an abrupt, sudden collapse, best appreciated by elevating the patient's arm while grasping the radial artery with the palmar aspect of the fingers. It is characteristic of aortic regurgitation and high-output states such as severe

anaemia, thyrotoxicosis, or arteriovenous fistulas.

**Alternate Names:**

Corrigan Pulse; Water-Hammer Pulse

**Collateral**

Providing an alternative or parallel route, such as collateral blood vessels that supply tissue around an obstruction; in other contexts it means accompanying or secondary.

**Collateral Circulation**

Alternate blood-vessel routes that develop or enlarge when a main artery becomes narrowed or blocked. They may partly maintain blood flow to tissue, but they may not prevent symptoms or injury if the blockage is severe.

**Collateral Ligament Injury**

A sprain or tear of a strong ligament on the inner or outer side of the knee. It may cause pain, swelling, tenderness, or looseness after a twisting or sideways force; treatment depends on severity and may include a brace and rehabilitation.

**Collecting Duct**

Small kidney tubes that receive fluid from several filtering units and adjust its water and salt content before it becomes urine. Antidiuretic hormone helps them return water to the blood.

**Colles Fracture**

A break near the wrist end of the larger forearm bone, with the broken hand-side fragment displaced backward. It usually follows a fall onto an outstretched hand and causes wrist pain, swelling, and deformity.

**Collimator**

A device in an imaging camera that allows radiation traveling in selected directions to reach the detector. It helps form a clearer image, but its design affects the balance between detail and the ability to detect weak signals.

---

## Colloids

Fluids containing large particles that tend to stay in blood vessels longer than simple salt solutions. Some are used to support circulation, but they can cause fluid overload, allergic reactions, or kidney problems and are chosen according to the clinical situation.

## Coloboma

A birth defect caused by incomplete formation of part of the eye, leaving a notch or gap in the iris, retina, choroid, or optic nerve. Vision problems vary with the structure and amount affected.

## Coloboma Of The Iris

A congenital keyhole- or notch-shaped defect in the iris caused by incomplete embryonic closure, sometimes associated with colobomas in the retina, optic nerve, or other organs.

## Colocolonic Intussusception

Telescoping of one part of the colon into the next section. It is uncommon in adults and may be caused by a polyp, tumor, or other lesion, producing intermittent pain, bloating, vomiting, or altered bowel movements.

## Cologne Poisoning

Ingestion of cologne containing alcohol and fragrances that may cause intoxication, hypoglycemia, respiratory depression, or gastrointestinal irritation.

## Cologuard

A stool-based colorectal-cancer screening test combining DNA-marker analysis with a fecal blood test; a positive result requires diagnostic colonoscopy.

## Colon

The main part of the large intestine between the small intestine and rectum. It absorbs water and salts and stores and moves stool through its ascending, transverse, descending, and sigmoid sections.

## Colon Cancer

A malignant tumor arising in the colon, usually

from gland-forming cells in the bowel lining. Many cases develop from precancerous polyps, but some arise through other pathways. Symptoms may include blood in the stool, a persistent change in bowel habits, abdominal discomfort, unexplained weight loss, fatigue, or iron-deficiency anemia; early cancer may cause no symptoms. Diagnosis usually involves colonoscopy and biopsy, followed by staging to guide treatment. Management may include surgery, chemotherapy, targeted therapy, or immunotherapy depending on the stage and tumor features.

### Alternate Names:

Colon Carcinoma; Cancer of the Colon; Bowel Cancer (informal or regional term)

## Colon Cancer Prevention

Colorectal-cancer risk can be reduced through recommended screening, physical activity, healthy weight, limiting alcohol, avoiding tobacco, and a diet rich in fiber-containing plant foods. Screening can detect precancerous polyps and early cancer before symptoms develop.

## Colon Cancer Screening

Testing intended to detect colorectal cancer or precancerous polyps before symptoms develop in people at average risk. Options include stool-based tests and direct visualization such as colonoscopy; the appropriate method and interval depend on age, risk, prior findings, and local guidelines. People with symptoms or increased inherited or personal risk may need diagnostic or earlier specialist evaluation rather than routine screening.

## Colon Polyps

Growths from the inner lining of the colon. Many are harmless, but some can slowly become colorectal cancer; removing polyps during colonoscopy can prevent some cancers, and follow-up depends on their number, size, and type.

### Alternate Names:

Colonic Polyps; Colorectal Adenomas

## Colonic Diverticulitis

Alternate name; see *Diverticulitis*.

---

### Colonic Inertia

Abnormally slow movement of stool through the colon, causing persistent constipation, bloating, abdominal discomfort, and infrequent bowel movements. It is diagnosed after other causes of constipation have been considered.

### Colonic Ischemia

Reduced blood flow to the colon causing tissue injury. It may produce sudden abdominal pain, bloody diarrhea, and tenderness and can result from low blood pressure, narrowed arteries, clots, medicines, or inflammation of blood vessels.

### Colonic Mucosa

The inner lining of the colon. It absorbs water and salts, makes mucus, and contains cells that help protect the colon.

### Colonic Polyps

Alternate name; see *Colon polyps*.

### Colonic Pseudo-Obstruction

Massive clinical and radiological dilation of the colon, predominantly involving the cecum and right colon, in the complete absence of a mechanical or physical obstructing lesion; associated with severe non-operative medical illnesses or recent surgery.

#### Alternate Names:

Ogilvie Syndrome; Acute Colonic Pseudo-Obstruction

### Colonic Stenosis

Narrowing of part of the large intestine that makes it harder for stool and gas to pass. Causes include scar tissue, inflammation, cancer, or reduced blood supply; symptoms may include pain, bloating, constipation, vomiting, or bowel blockage.

### Colonic Tissue Culture

Microbiologic cultivation and identification of biopsy tissue harvested from the colonic mucosa during colonoscopy to detect fastidious bacterial pathogens, mycobacteria (e.g., *Mycobacterium tuberculosis*), or fungal organisms.

#### Alternate Names:

Colon Biopsy Culture; Colonic Mucosal Culture

### Colonization

The presence and growth of microorganisms on or in the body without tissue damage or illness. Colonization can occur before infection, but it does not by itself explain symptoms or require antibiotics.

### Colonoscope

A flexible, lighted instrument with a camera that is inserted through the anus to examine the rectum and colon. It can contain channels for air, irrigation, biopsy tools, or devices used to remove polyps.

### Colonoscopy

An examination of the rectum and colon using a flexible camera. It can find inflammation, bleeding, tumors, and polyps and allows tissue sampling or removal of some polyps during the same procedure.

### Colony-Forming Unit

A laboratory estimate of the number of living microorganisms in a sample, based on the colonies that grow in culture. One colony may come from one organism or from a group of organisms.

### Colony-Stimulating Factors

Natural or manufactured substances that signal the bone marrow to make more blood cells. Some are used to prevent or treat low white-cell counts caused by cancer treatment or other illness.

### Color Blindness

Reduced or absent ability to distinguish certain colors under normal lighting. It is often inherited, especially through X-linked variants, but may also result from eye disease, neurologic disease, aging, or medicines.

#### Alternate Names:

Deficient Color Vision Poor Color Vision

### Color Doppler

An ultrasound technique that uses colors to show the direction and relative speed of blood flow over



---

a regular gray-scale image. It helps assess vessels, heart valves, and circulation.

### **Color Vision Deficiency**

Reduced ability to distinguish some colors, commonly red and green. It is often inherited because of differences in the eye's color-sensing cells, but it can also develop from eye disease, optic-nerve or brain disorders, medicines, or toxins. The type and severity vary.

### **Color Vision Test**

An eye test that assesses the ability to distinguish colors or identify color patterns. It helps detect inherited color-vision deficiency and acquired abnormalities caused by eye, optic-nerve, brain, medicine, or toxic disorders.

### **Colorado Tick Fever**

A viral infection spread by certain ticks, usually causing sudden fever, headache, tiredness, and muscle aches. Some people develop a rash, nausea, or low blood counts; most recover with supportive care, but severe illness can occur.

### **Colorectal Adenocarcinoma**

The most common cancer of the colon and rectum, arising from gland-forming cells in the bowel lining. It may cause bleeding, anemia, altered bowel habits, pain, or no early symptoms and can spread to other organs.

### **Colorectal Adenoma**

A usually noncancerous gland-forming growth in the colon or rectum that may gradually develop into colorectal cancer.

### **Colorectal Cancer**

Cancer arising in the colon or rectum, including colon cancer and rectal cancer. It may cause bleeding, altered bowel habits, iron-deficiency anemia, abdominal pain, weight loss, or no early symptoms. Diagnosis usually involves colonoscopy and biopsy; evaluation and treatment depend on the tumor's location, stage, and molecular features.

### **Colorectal Cancer Genetics**

The study of inherited variants that increase the risk of colorectal cancer. Important inherited syndromes include Lynch syndrome, involving DNA mismatch-repair genes, and familial adenomatous polyposis, involving the APC gene.

### **Colorectal Cancer Screening**

Public health strategy to detect colorectal cancer and precancerous polyps in asymptomatic individuals. Current guidelines recommend screening beginning at age forty-five.

### **Colorectal Cancer Screening Tests**

Tests used to detect colorectal cancer or precancerous polyps in people without symptoms. They include stool-based tests and examinations that visualize the colon, such as colonoscopy, CT colonography, and flexible sigmoidoscopy. The appropriate test and interval depend on age, risk, and local guidance.

### **Colorectal Polyp**

A growth arising from the lining of the colon or rectum. Many are benign, but some adenomatous or serrated polyps can become cancerous over time and are removed during colonoscopy.

### **Colorectal Polyps**

Growths arising from the lining of the colon or rectum. They may be non-neoplastic, adenomatous, or serrated; some can progress to colorectal cancer, so removal and microscopic examination guide surveillance.

#### **Alternate Names:**

Bowel Polyp; Bowel Polyps

### **Colorectal Surgery**

Surgical treatment of diseases of the colon, rectum, and anus. Procedures may include segmental bowel resection, restorative reconstruction, stoma formation, or local procedures, depending on the disease and its extent.

---

## Colostomy

Surgery that brings part of the large intestine through an opening in the abdomen, called a stoma. Stool and gas leave through the stoma into a pouch. A colostomy may be temporary or permanent and may be needed for cancer, bowel blockage, severe infection, injury, or other bowel disease. Stoma nurses can teach people how to care for it and return to usual activities.

### Alternate Names:

Bowel Stoma; Large Intestinal Stoma; Colic Stoma; Ostomy

## Colostrum

The thick yellowish breast fluid made during late pregnancy and the first days after birth. It provides concentrated nutrition and immune protection, especially through antibodies, to the newborn.

## Colpitis

Inflammation of the vagina caused by infection, altered vaginal bacteria or yeast, irritation, allergy, or hormonal changes. Symptoms may include discharge, itching, burning, pain, odor, or painful intercourse.

## Colporrhaphy

Surgery that repairs a weakened front or back wall of the vagina. It may support the bladder or bowel when pelvic tissues bulge into the vagina and cause pressure, discomfort, or difficulty emptying.

## Colposcope

A lighted magnifying instrument used to examine the cervix, vagina, or vulva closely and guide biopsy of suspicious areas.

## Colposcopy

An examination using a magnifying instrument to inspect the cervix, vagina, and vulva after a screening test shows an abnormality. The clinician may apply solutions and take tissue samples from suspicious areas.

## Colposcopy-Directed Biopsy

A specialized gynecologic diagnostic procedure in which acetic acid- or Lugol iodine-enhanced cervical, vaginal, or vulvar epithelial abnormalities visualized under colposcopic magnification are biopsied for histopathologic evaluation of dysplasia or malignancy.

### Alternate Names:

Colposcopic Biopsy; Cervical Punch Biopsy; Directed Cervical Biopsy

## Columnar Epithelium Lined Lower Esophagus

Alternate index label; see *Barrett's Esophagus*.

## Coma

A state of profound unresponsiveness in which a person cannot be awakened and does not show purposeful responses. It is a medical emergency with many possible causes.

### Alternate Names:

Comatose State; Unconsciousness

## Coma Scale

A standardized tool for describing level of consciousness using eye opening, verbal response, and motor response.

## Combination Inhalers

Inhalers containing two or more medicines, such as an inhaled corticosteroid with a long-acting bronchodilator. They may simplify regular treatment for asthma or chronic obstructive pulmonary disease; the prescribed device and schedule must be followed.

## Combination Therapy

Use of two or more treatments together to address a disease or mechanism. The benefits and risks depend on the condition and the specific combination.

## Combined Electrolyte Imbalances

Abnormal blood levels of calcium, phosphate, and potassium. This combination does not have one single cause; each result must be confirmed and

---

interpreted with kidney function, medicines, symptoms, and the rest of the electrolyte panel.

### **Combined Hormone Therapy**

Hormone therapy containing both estrogen and a progestogen, generally used when endometrial protection is needed in a person with a uterus.

#### **Alternate Names:**

Estrogen-Progestogen Therapy; Combined Menopausal Hormone Therapy

### **Combined Hyperlipidemia**

A blood-fat disorder with high cholesterol and triglycerides, often increasing harmful lipoproteins and raising the long-term risk of artery disease.

#### **Alternate Names:**

Mixed Hyperlipidemia; Combined Hyperlipoproteinemia

### **Combined Oral Contraceptive Pill**

Oral contraceptive formulations containing both an estrogen (typically ethinylestradiol) and a progestin that prevent pregnancy primarily through feedback inhibition of gonadotropins to suppress ovulation, accompanied by thickening of cervical mucus and endometrial thinning.

#### **Alternate Names:**

Combined Oral Contraceptives; COCP; Combination Birth Control Pills; The Pill

### **Combined Spinal-Epidural Analgesia**

A form of regional pain relief that gives medicine into the spinal fluid for rapid relief and places a catheter outside it for continued dosing. It is often used during labor or surgery and requires monitoring.

### **Comedocarcinoma**

A form of ductal carcinoma in situ of the breast characterized by central necrosis within malignant ducts, which may produce a comedo-like appearance.

### **Comedonal Acne**

A form of acne in which open comedones (blackheads) and closed comedones (whiteheads) predominate, with little inflammatory redness or swelling.

It results from blockage of hair follicles by sebum and keratin.

### **Comedones**

Plugs of sebum and keratin within hair follicles, appearing as open blackheads or closed whiteheads and representing a primary lesion of acne.

### **Comfort Measures Only**

A care plan focused on relief of pain, breathlessness, distress, and other symptoms rather than treatments intended to prolong life.

### **Commando Procedure**

Extensive heart surgery for severe infection destroying both the aortic and mitral valve areas and the tissue between them. The surgeon removes infected tissue, rebuilds the damaged area, and replaces both valves when necessary.

#### **Alternate Names:**

Aortomitral Fibrous Body Reconstruction; Interventricular Fibrosa Reconstruction

### **Commensal**

Describing a microorganism that lives on or in the body and benefits from the relationship without normally harming the person. Many harmless skin and bowel bacteria are commensal.

### **Comminuted**

Describing a fracture in which one bone breaks into several pieces. These fractures may be caused by high-energy injury and can be unstable or involve nearby tissues, so treatment depends on the bone, alignment, skin, nerves, and blood vessels.

### **Comminuted Fracture**

A broken bone divided into several pieces, usually by a high-energy injury. It may be unstable and damage nearby skin, muscles, nerves, or blood vessels, so treatment depends on the site and severity.

### **Commissurotomy**

Surgical incision or separation of a commissure, especially a fused cardiac valve commissure, to relieve

---

obstruction.

### **Commode**

A portable toilet or chair placed near a bed for a person who cannot easily reach a bathroom.

### **Common Bile Duct**

The tube formed when ducts from the liver and gallbladder join. It carries bile into the first part of the small intestine, where bile helps digest dietary fat.

### **Common Carotid Artery**

One of two major arteries carrying blood from the heart toward the head and neck. Each divides into an internal branch supplying the brain and an external branch supplying the face and scalp.

### **Common Cold**

A short-term viral infection of the upper respiratory tract, especially the nose and throat. More than 200 viruses can cause it; rhinoviruses are the most common, and seasonal human coronaviruses and several other respiratory viruses can also cause colds. Symptoms may include a runny or blocked nose, sneezing, sore throat, cough, headache, mild body aches, and sometimes a low-grade fever. The viruses spread mainly through respiratory droplets and close contact and can also spread from contaminated hands or surfaces. Symptoms usually peak within two or three days and improve in less than a week, although a cough or runny nose may last 10 to 14 days.

#### **Alternate Names:**

Acute Viral Rhinopharyngitis; Coryza; Head Cold

### **Common Cold In Babies**

A viral infection of a baby's nose and throat causing a runny or blocked nose, cough, sneezing, and feeding difficulty. Babies may become dehydrated or have breathing trouble, so poor feeding, fewer wet diapers, or fast breathing needs prompt care.

### **Common Fibular Nerve**

A major nerve that passes around the outer side of the knee into the lower leg and foot. It helps lift

and turn the foot and carries feeling from much of the lower leg and top of the foot.

#### **Alternate Names:**

Common Peroneal Nerve

### **Common Iliac Arteries**

Two large arteries formed where the abdominal aorta divides. Each artery supplies blood to the pelvis and then divides into arteries serving the pelvic organs and lower limb.

### **Common Iliac Vein**

A vein formed by the union of the internal and external iliac veins. The right and left common iliac veins unite to form the inferior vena cava.

### **Common Medical Abbreviations And Terms**

Medical abbreviations and terms summarize diagnoses, tests, medicines, anatomy, and treatment instructions. Because abbreviations can be ambiguous or unsafe, patients should ask clinicians or pharmacists to explain unfamiliar wording rather than infer its meaning.

#### **Alternate Names:**

Medical Acronyms; Clinical Terminology

### **Common Migraine**

A migraine attack without a preceding aura. It usually causes moderate or severe throbbing pain lasting four to 72 hours, often with nausea or sensitivity to light or sound and worsening with routine activity.

#### **Alternate Names:**

Migraine Without Aura; Migraine Without Neurologic Aura

### **Common Peroneal Nerve Dysfunction**

Impairment of the common peroneal nerve, often from compression near the fibular head, trauma, surgery, or systemic nerve disease. It may cause foot drop, weakness of ankle eversion or dorsiflexion, and sensory loss over the lateral leg or dorsum of the foot.

---

### **Common Peroneal Neuropathy**

A peripheral nerve disorder causing weakness of ankle dorsiflexion and eversion, sensory loss over the anterolateral leg and dorsum of the foot, and possible foot drop.

### **Common Uterine and Ovarian Growths**

Three usually noncancerous types of growth: fibroids arise from uterine muscle, cysts are fluid-filled sacs often found in the ovaries, and polyps are tissue projections from the uterine lining or cervix.

### **Common Variable Immunodeficiency**

A primary antibody-deficiency disorder characterized by low immunoglobulin G with low immunoglobulin A and/or M and impaired responses to vaccines. It causes recurrent sinus, ear, and lung infections and may also cause chronic gastrointestinal disease, autoimmunity, enlarged lymph nodes, granulomas, or increased lymphoma risk.

### **Commotio Cordis**

Sudden cardiac arrest after a blunt blow to the chest at a vulnerable moment in the heartbeat, usually without visible damage to the heart. It is an emergency requiring immediate CPR and defibrillation when available.

### **Communicable Disease**

An illness caused by a specific infectious agent (such as a bacterium, virus, parasite, or fungus) or its toxic products that is transmissible directly or indirectly from an infected individual, animal, or environmental reservoir to a susceptible host. Transmission occurs via respiratory droplets, direct contact, bodily fluids, vectors (such as mosquitoes or ticks), contaminated water, food, or fomites; prevention and control rely on immunization, sanitation, antimicrobial therapy, and public-health surveillance with mandatory disease notification.

#### **Alternate Names:**

Communicable Diseases; Transmissible Disease; Infectious and Communicable Disease

### **Communicating Hydrocephalus**

A buildup of cerebrospinal fluid that enlarges the

brain's fluid-filled spaces because the fluid is not absorbed normally after it circulates. It can cause headache, vomiting, walking difficulty, sleepiness, or thinking problems.

### **Communicating With Patients**

The clinical exchange of information, concerns, preferences, and decisions between healthcare professionals and patients, using clear language, listening, empathy, and confirmation of understanding.

### **Communicating With Someone With Aphasia**

Communication strategies for a person with impaired language from brain injury or disease, including short sentences, extra time, visual cues, and supported conversation rather than assuming impaired intelligence.

### **Communicating With Someone With Dysarthria**

Communication strategies for a person whose speech muscles are weak or poorly coordinated, including reducing background noise, confirming messages, and using writing or augmentative aids when needed.

### **Community Acquired Infection**

An infection caught outside a healthcare facility or present when a person is admitted, rather than acquired during recent healthcare treatment. Examples include some lung, urinary-tract, and skin infections.

### **Community Health Worker**

A trained member of a community who helps connect people with health services, education, prevention, and social support.

### **Community Mental-Health Treatment**

Mental-health assessment, treatment, rehabilitation, and support delivered outside a hospital, such as in clinics, homes, schools, or community programs. It may include medication management, psychotherapy, crisis support, peer support, and help with housing, work, or daily functioning.

---

## Community Rehabilitation

Therapy and support delivered at home, in clinics, or in community settings to improve movement, communication, independence, participation, and daily functioning after illness or injury.

## Community Reintegration

The process of returning to everyday life and taking part in work, education, family, social, and leisure activities after illness or injury. Support may include rehabilitation, changes to the home or workplace, transport help, and practical or emotional support.

## Community-Acquired

Acquired outside a healthcare facility or before healthcare exposure. For example, community-acquired pneumonia begins outside a hospital and may involve different organisms and resistance patterns from hospital-acquired disease.

## Community-Acquired Pneumonia

Pneumonia acquired outside a hospital or long-term-care facility. It may cause cough, fever, chest pain, breathlessness, and low oxygen; diagnosis and treatment depend on examination, imaging, severity, likely organisms, comorbidities, and local antibiotic guidance. CURB-65 or PSI can help assess the need for hospital care but does not replace clinical judgment.

### Alternate Names:

CAP

## Community-Acquired Pneumonia In Adults

A lung infection that begins outside a hospital or long-term-care facility. It may cause cough, fever, chest pain, breathlessness, or low oxygen; treatment and the need for hospital care depend on severity and other health conditions.

## Comorbidity

The presence of two or more health conditions in the same person. The conditions may affect one another and can make symptoms, medicine choices, and treatment planning more complicated.

## Compact Bone

Dense outer bone tissue arranged in strong cylindrical units called osteons. It provides most of the skeleton's strength and protects inner marrow while supporting movement and weight-bearing.

### Alternate Names:

Cortical Bone; Dense Bone

## Comparison

A reference in an imaging report to earlier studies used to assess whether a finding has changed.

## Compartment Pressure

The pressure inside a closed group of muscles, nerves, and blood vessels. If it rises too much, blood flow can fall and cause permanent tissue damage; measurements support but do not replace examination.

## Compartment Syndrome

Dangerous pressure buildup inside a confined muscle group that reduces blood flow and can permanently damage muscle and nerves. Severe pain, swelling, numbness, or weakness after injury requires emergency assessment and often surgery.

## Compartment Syndrome Cross-Reference

Alternate name; see *Compartment syndrome*.

## Compensated Heart Failure

Heart failure in which symptoms and fluid buildup are stable or controlled with treatment. The heart failure is still present and can worsen, especially if a new illness or other problem develops.

## Competence

A legal determination that a person can make a particular decision or manage a matter. In clinical care, decision-making capacity is assessed for each specific choice.

## Competitive Antagonist

A medicine or substance that occupies the same receptor as another substance but does not activate it. By competing for that site, it reduces the other substance's effect; increasing the other substance



---

may overcome the competition.

### Competitive Inhibition

Reduced activity of an enzyme when another substance competes with its usual target for the same binding site. The effect is often reversible, and more of the usual target may reduce the inhibitor's effect.

### Complement

A group of blood proteins that strengthens immune defense. When activated, complement can mark germs for removal, attract immune cells, increase inflammation, or directly damage some microbes.

### Complement C1Q Deficiency

A rare inherited lack of a complement protein needed for immune defense and removal of antibody-bound material. It can cause repeated infections and autoimmune illness resembling lupus, including rash, joint, kidney, or blood problems.

### Complement C2/C4 Deficiency

Inherited or acquired lack of complement C2 or C4, proteins needed especially for the classical complement pathway. It can increase susceptibility to infections with encapsulated bacteria and autoimmune disease resembling lupus. Testing may show low CH50 with low C2 or C4; care includes vaccination, prompt infection treatment, and specialist management of autoimmune disease or recurrent infections.

### Complement C3

A major blood protein in the complement immune system. When activated, it helps mark microbes for removal and promotes inflammation; severe deficiency can cause repeated serious infections.

### Complement Cascade

A chain reaction involving many immune proteins in blood and tissues. It marks microbes for destruction, attracts immune cells, increases inflammation, and can directly break holes in some microbes.

### Complement Component 4

A blood test measuring complement component C4, part of the classical complement pathway; low levels may occur with immune-complex disease, hereditary angioedema, or complement consumption.

### Complement Component C3

A blood test measuring complement component C3, a major protein of the complement immune system; abnormal levels can support evaluation of immune-complex disease, infection, or complement deficiency.

### Complement Component Deficiency

A genetically determined deficiency in one of the proteins of the complement cascade, predisposing to recurrent pyogenic infections (early classical components), severe autoimmune immune-complex disorders (C1q, C2, C4), or disseminated *Neisseria* infections (terminal membrane attack complex C5-C9).

### Complement Deficiencies

Conditions caused by too little or absent activity of one or more complement proteins. They can increase the risk of particular infections and autoimmune diseases; the effects depend on the missing protein.

### Complement Deficiency

Reduced or absent activity of one or more complement proteins, increasing susceptibility to some infections or autoimmune disease.

### Complement Fixation Test For C Burnetii

A serologic test detecting complement-fixing antibodies to *Coxiella burnetii*, the bacterium that causes Q fever; antibody phase patterns and titers help distinguish recent from persistent infection.

### Complement System

A group of blood proteins that supports immune defense by destroying microbes and promoting inflammation and phagocytosis.

---

### **Complementary Sequence**

A DNA or RNA sequence whose building blocks pair with those of another sequence. This matching allows genetic strands to bind and is used in copying, testing, and regulating genetic information.

### **Complete Blood Count**

A blood test that measures red blood cells, white blood cells, platelets, hemoglobin, and hematocrit. It may also measure red-cell size and hemoglobin content and list the different types of white blood cells. It helps assess anemia, infection, inflammation, bleeding, and some bone-marrow disorders.

### **Complete Breech**

A type of breech presentation in which the fetus has both hips and knees flexed (in a cross-legged sitting posture) so that the buttocks and feet present together at the maternal pelvic inlet. It accounts for approximately 5–10% of breech presentations.

#### **Alternate Names:**

Flexed breech; Complete breech presentation.

### **Complete Cystectomy**

Surgery to remove the entire urinary bladder, usually for invasive bladder cancer. Because urine can no longer leave through the bladder, the surgeon creates a urinary diversion such as an ileal conduit, a continent reservoir, or a new bladder from bowel.

### **Complete Denture**

A removable artificial set of teeth replacing all teeth in the upper jaw, lower jaw, or both. It rests on the gums and may improve chewing, speech, and appearance but can require adjustment for comfort and stability.

### **Complete Miscarriage**

Pregnancy loss in which all pregnancy tissue has passed from the uterus, usually followed by easing bleeding and cramps. A clinician may use examination, hormone testing, or ultrasound to confirm that no tissue remains and to exclude an ectopic pregnancy; heavy bleeding, severe pain, dizziness, or fever needs urgent care.

#### **Alternate Names:**

### **Complete Spontaneous Abortion**

### **Complete Response**

No detectable evidence of cancer after treatment on the tests used. It does not prove that every cancer cell has been eliminated, so continued follow-up is needed.

### **Complex Carbohydrates**

Carbohydrates made of many linked sugar units, including starch and dietary fiber. They occur in foods such as grains, beans, vegetables, and potatoes; fiber is not fully digested and supports bowel function.

### **Complex Febrile Seizure**

A seizure triggered by fever in a young child that lasts more than 15 minutes, affects only part of the body, or happens more than once during the same illness. It needs medical assessment to exclude brain infection or another cause. Most children recover fully, but the risk of another seizure is higher than with a simple febrile seizure.

#### **Alternate Names:**

Atypical Febrile Seizure; Complicated Febrile Seizure

### **Complex Regional Pain Syndrome**

A long-lasting pain condition, usually affecting one arm or leg after an injury, surgery, or another event. It is thought to involve abnormal signaling or damage in the peripheral and central nervous systems. Pain may be severe and continue beyond the usual healing period, with increased sensitivity to touch or movement. The affected area may also develop swelling, changes in skin color or temperature, unusual sweating, stiffness, weakness, or difficulty moving. CRPS type I occurs without a confirmed major nerve injury and was formerly called reflex sympathetic dystrophy; CRPS type II follows an identified peripheral nerve injury and was formerly called causalgia. Symptoms and recovery vary.

#### **Alternate Names:**

CRPS; Reflex Sympathetic Dystrophy; Complex Regional Dystrophy Syndrome

---

## Compliance

The extent to which a person follows a treatment plan. The term is often replaced by adherence, which recognizes that treatment decisions should be discussed and agreed with the patient.

## Complicated Bereavement

Alternate name; see *Complicated Grief*.

## Complicated Grief

Persistent, intense grief that substantially interferes with daily functioning and lasts longer than expected for the person's cultural and social context.

### Alternate Names:

Complicated Bereavement Traumatic Grief

## Complicated Migraine

An older, nonspecific term for migraine accompanied by prolonged or prominent neurologic symptoms; the specific syndrome should be identified when possible.

### Alternate Names:

Migraine with Prolonged Aura; Complex Migraine

## Complicated Urinary Tract Infection

A urinary infection associated with functional or structural abnormality, obstruction, catheterization, renal impairment, male anatomy, pregnancy, or another factor increasing treatment failure or complication risk. Management requires culture-directed therapy and correction of the underlying problem when possible.

## Complication

A new problem or harmful effect that develops during or after an illness, treatment, or procedure. Examples include infection, bleeding, organ damage, medicine reactions, and blood clots; seriousness depends on the problem and its effect.

## Composite Resin

A tooth-colored material used to repair cavities, chipped teeth, or other defects. It is bonded to the tooth and shaped to match it, but may wear or stain sooner than some harder restorations.

## Compound Heterozygote

A person who has two different disease-causing changes in the same gene, one inherited in each copy of the gene. Together, the changes can cause an inherited disorder.

## Compound Muscle Action Potential

The summated electrical voltage recorded over a muscle belly during a motor nerve conduction study, reflecting the near-simultaneous depolarization of all innervated muscle fibers following supramaximal electrical stimulation of its motor nerve.

### Alternate Names:

CMAP; Motor Response

## Compound Presentation

A birth position in which a hand, arm, foot, or another limb lies beside the baby's head or buttocks. The limb may move back during labor, but the position can obstruct delivery or increase injury risk.

## Compound Presentations

A fetal presentation in which an extremity, usually a hand or arm, lies beside the presenting head or breech. It may resolve spontaneously, but persistent obstruction, cord prolapse, or fetal distress can require urgent obstetric intervention.

## Compounding

Preparation, mixing, or altering of a medicine to meet an individual patient's needs when an appropriate approved product is unavailable.

## Compounding Pharmacy

A pharmacy that prepares customized formulations for an individual patient when a suitable commercially manufactured product is unavailable or unsuitable.

### Alternate Names:

Compounding Pharmacy Practice; Extemporaneous Pharmacy

## Comprehensive Geriatric Assessment

A detailed review of an older person's illnesses,

---

medicines, movement, daily activities, memory, mood, nutrition, and support at home. The findings are used to create a coordinated care plan based on the person's goals.

### **Comprehensive Metabolic Panel**

A routine 14-substance blood test combining the eight basic metabolic tests (glucose, calcium, electrolytes, BUN, and creatinine) with six liver and protein tests (ALT, AST, ALP, bilirubin, albumin, and total protein). Commonly ordered before surgery and during acute medical care, it evaluates whether the liver and kidneys can safely clear anesthetics and medications, verifies electrolyte stability to prevent irregular heart rhythms during surgery, and checks protein stores necessary for post-surgical wound healing.

#### **Alternate Names:**

CMP; Chem-14; Comprehensive Chemistry Screen

### **Compress**

To press together or apply pressure, as with a bandage or device, or to reduce the volume of something.

### **Compressed Nerve**

Alternate name; see *Pinched Nerve*.

### **Compression**

Pressure applied to tissue or a structure, which may reduce swelling or bleeding but can also impair nerves, blood vessels, or organs when excessive.

### **Compression Bandage**

A bandage that applies controlled pressure to a limb or body part. It may reduce swelling, support injured tissue, or help manage venous disease, but improper use can impair circulation or damage skin.

### **Compression Fracture**

Collapse of a vertebra or other bone because of compression, often related to osteoporosis or trauma.

### **Compression Fractures Of The Back**

Collapse of one or more vertebral bodies caused

by weakened bone, often from osteoporosis, or by substantial trauma. It can cause sudden back pain, height loss, kyphosis, and rarely nerve or spinal-cord compression.

### **Compression Stockings**

Elastic garments that apply graduated pressure to the legs, supporting venous return and reducing swelling or selected risks of venous pooling; correct fit and arterial circulation matter.

### **Compression Therapy**

Use of external pressure, usually from garments or bandages, to support venous or lymphatic flow and reduce swelling in selected conditions.

### **Compression Ultrasound**

An ultrasound test in which gentle pressure is applied to a vein. A healthy vein usually collapses; failure to collapse may indicate a blood clot, especially in the deep veins of a limb.

### **Compressive Atelectasis**

Collapse of part of a lung because fluid, air, a tumor, or another process presses on it from outside. The lung may re-expand when the pressure is relieved.

### **Compton Scattering**

A change in the direction and energy of an X-ray or other high-energy photon after it strikes an electron. It creates scattered radiation that can reduce image quality and add unnecessary exposure during imaging.

### **Compulsions**

Repeated actions or mental rituals that a person feels unable to resist, often to reduce distress caused by unwanted thoughts. They take up time or interfere with daily life and may not realistically prevent the feared event.

### **Compulsive Gambling**

Alternate, nonpreferred label; see *Gambling Disorder*.

---

### Compulsive Obsessive Disorder

Alternate name; see *Obsessive-Compulsive Disorder*.

### Compulsive Overeating

Alternate name; see *Binge-Eating Disorder*.

### Compulsive Sexual Behavior

Recurrent sexual thoughts or behaviors that are difficult to control and cause significant distress or impairment.

#### Alternate Names:

Compulsive Sexual Behavior Disorder

### Compulsive Sexual Behavior Disorder

A persistent pattern of failure to control intense, repetitive sexual impulses or behaviors, resulting in marked distress or significant impairment and continuing despite adverse consequences. It is not defined simply by a high sex drive or moral disapproval; assessment considers control, distress, consent, and coexisting mental-health or substance-use conditions.

#### Alternate Names:

Compulsive Sexual Behavior; Problematic Sexual Behavior

### Compulsive Stealing

Alternate index label; see *Kleptomania*.

### Computed Tomography

An imaging test that uses rotating X-rays and computer processing to create detailed cross-sectional pictures. It is fast and can show bones, bleeding, organs, blood vessels, tumors, infection, and internal injury.

### Computed Tomography Pulmonary Angiography

A CT scan performed after injected contrast highlights the lung arteries. It is commonly used to look for a pulmonary embolism and may also show strain on the heart or other chest abnormalities.

#### Alternate Names:

CTPA; CT Pulmonary Angiogram; Pulmonary CT Angiography

### Computed Tomography Scan

An X-ray scan processed by a computer to produce cross-sectional images of the body. It can rapidly detect bleeding, fractures, infection, blocked bowel, blood clots, tumors, and many other urgent conditions.

### Computed Tomography Urography

A CT scan of the kidneys, ureters, and bladder, usually using contrast material to show the urinary tract. It can show stones, blockage, bleeding, tumors, or structural changes.

### Computerized Axial Tomography

Computed tomography, an imaging method that uses X-rays and computer processing to produce cross-sectional images of the body; it is also called a CT or CAT scan.

### Concato's Disease

An old name for progressive polyserositis, inflammation of several serous membranes, such as the pleura, pericardium, and peritoneum, often with fluid accumulating in the chest or abdomen. It may complicate tuberculosis, cancer, or connective-tissue disease and can cause shortness of breath, fatigue, and abdominal swelling.

### Concealed Conduction

A heart rhythm event in which an electrical impulse enters part of the heart's conduction system but stops before producing a visible heartbeat. The affected tissue may then delay or block the next impulse and alter the heartbeat pattern.

### Concealed Pregnancy

A pregnancy that is deliberately hidden or not recognized until late in pregnancy or childbirth. It may occur because symptoms are mild, pregnancy tests are misleading, or the person is unable or unwilling to recognize the pregnancy; medical assessment is still important.

### Concentration

The amount of a substance in a given volume, such as the level of glucose or medicine in blood. It is

---

reported in units such as milligrams per deciliter or millimoles per liter.

### **Concentric Muscle Action**

Muscle action in which a muscle produces force while becoming shorter, such as when lifting an object.

### **Conception**

The joining of a sperm and egg, usually in a fallopian tube, to form the first cell of an embryo. The resulting cells divide as they travel toward the uterus, where implantation may occur several days later.

#### **Alternate Names:**

Fertilization

### **Conchae**

Curved bony structures covered by mucosa inside the nasal cavity that warm, humidify, and filter inhaled air.

### **Conclusion**

The part of a report that summarizes the main findings and their clinical significance or recommended next steps.

### **Concomitant**

Existing or occurring at the same time as another condition, treatment, or event. Concomitant medicines or illnesses may change diagnosis, treatment choices, expected effects, side effects, or monitoring needs.

### **Concordance**

The frequency with which two people, especially twins, show the same trait or diagnosis. Higher agreement in identical than fraternal twins may suggest a genetic contribution, but shared environment also matters.

### **Concrescence**

A dental abnormality in which two neighboring teeth become joined by the hard tissue covering their roots, while their crowns remain separate. It

most often affects back teeth and may complicate extraction.

### **Concretio Cordis**

An abnormal joining or fusion of tissues involving the heart or its surrounding sac. It may result from inflammation, scarring, or a birth defect and can restrict heart movement or interfere with normal function.

### **Concussion**

A mild traumatic brain injury caused by a blow, jolt, or force that briefly changes brain function. Symptoms may include headache, dizziness, confusion, memory trouble, nausea, or sensitivity to light; routine scans may be normal.

### **Conditioning**

Learning in which repeated association or reinforcement changes a behavior or response; in medicine, conditioning can also mean preparing a patient or tissue for a treatment.

### **Conditions Caused By Deletion Mutations**

Deletion mutations remove DNA segments and can disrupt one gene or several neighboring genes. The clinical effect depends on size, location, inheritance, and whether the remaining copy compensates; genetic counseling and testing may clarify diagnosis and recurrence risk.

### **Condom**

A thin barrier sheath worn on the penis or inserted into the vagina that helps prevent pregnancy and reduces transmission of sexually transmitted infections when used correctly.

#### **Alternate Names:**

Male Condom; Barrier Contraception; External Condom

### **Conduct Disorder**

A childhood or adolescent disorder involving a persistent pattern of behavior that violates the rights of others or major age-appropriate rules. Examples include aggression, property destruction, deceitful-



---

ness, theft, and serious rule violations.

### **Conduction Aphasia**

A fluent aphasic syndrome resulting from a lesion disrupting the arcuate fasciculus connecting Broca area and Wernicke area, characterized by intact auditory comprehension and fluent spontaneous speech, but profound impairment in verbal repetition and prominent phonemic paraphasias.

### **Conduction Deafness**

Hearing loss caused by a problem in the outer or middle ear that prevents sound from reaching the inner ear normally. Earwax, fluid, a damaged eardrum, or changes in the small ear bones may be responsible.

### **Conductive Deafness**

Hearing loss resulting from a problem in the external auditory canal, tympanic membrane, or middle ear that reduces conduction of sound to the inner ear.

### **Conductive Hearing Loss**

Hearing loss caused by a problem in the outer or middle ear that prevents sound from reaching the inner ear normally. Earwax, fluid, a damaged eardrum, bone changes, or ear-bone disease may be responsible.

### **Condylar Process**

The rounded articular projection of a bone, especially the mandibular process that articulates with the temporal bone to form the temporomandibular joint.

### **Condyle**

A rounded projection at the end of a bone that forms part of a joint and helps guide movement against another bone.

### **Condyloid Joint**

A joint in which an oval surface of one bone fits into an oval depression in another bone. It allows bending, straightening, and side-to-side movement, but little rotation.

### **Condyloma**

A wart-like growth, often in the genital or anal area and commonly caused by human papillomavirus; condyloma lata are lesions of secondary syphilis.

### **Condyloma Acuminata**

Soft, wart-like growths around the genitals or anus caused by certain types of human papillomavirus. They spread through close skin-to-skin sexual contact; treatment removes visible warts but may not remove the virus, and vaccination helps prevent many infections.

### **Condyloma Acuminatum**

A soft, wart-like growth around the genitals or anus caused by certain human papillomavirus types. It spreads through intimate skin contact; treatment removes visible warts, but the virus may remain and recur.

### **Condyloma Lata**

Broad, flat, moist, whitish or grayish-pink plaques and papules occurring in warm intertriginous skin and mucosal areas (particularly the perianal, vulvar, and scrotum regions) during secondary syphilis, containing abundant highly infectious *Treponema pallidum* spirochetes.

#### **Alternate Names:**

Condylomata Lata; Syphilitic Condyloma

### **Condyloma Latum**

A broad, flat, moist, highly infectious lesion of secondary syphilis, usually around the genitals or anus. It differs from a genital wart and indicates syphilis requiring testing and antibiotic treatment.

### **Condylomata Acuminata**

Alternate name; see *Genital Warts*.

### **Condylomata Lata**

Broad, flat, moist, grayish-white papules occurring in secondary syphilis, typically found in the anogenital region and other intertriginous areas. They are highly infectious and contain abundant *Treponema pallidum* organisms. Diagnosis is by darkfield mi-

---

croscopy or serological testing for syphilis.

### **Cone Biopsy**

A surgical procedure that removes a cone-shaped portion of the cervix for diagnosis or treatment of high-grade cervical precancer and to determine whether invasive cancer is present.

### **Cone-Beam Computed Tomography**

A dental or facial scan that uses a cone-shaped X-ray beam to create three-dimensional images of teeth, jawbone, and nearby structures. It should be used when the extra detail will affect care.

### **Cones**

Light-sensitive retinal cells responsible for color vision and sharp detail, working best in daylight or other relatively bright conditions.

#### **Alternate Names:**

Cone Photoreceptors; Cone Cells

### **Confabulation**

Unintentional production of inaccurate memories or stories without an attempt to deceive. It may occur with memory disorders or injury affecting the front part of the brain.

### **Confidentiality**

The duty of healthcare professionals and organizations to protect a person's health information from unauthorized access or disclosure. Information may be shared when the patient agrees or when law permits or requires it.

### **Confirmatory Test**

A test used to verify a preliminary or screening result, often with a different method or greater specificity. Confirmation should be selected according to the condition, specimen, timing, and accepted clinical guidelines.

### **Confluent and Reticulated Papillomatosis**

An uncommon skin disorder causing asymptomatic or mildly itchy brown, papillomatous papules and plaques that merge centrally and form a net-like

pattern, usually on the upper trunk, neck, or axillae. Diagnosis is clinical after excluding fungal infection and other causes of hyperpigmentation; treatment may include specialist-prescribed antibiotics.

### **Conformal Radiation Therapy**

Radiation treatment planned so that the high-dose region conforms closely to the tumor's three-dimensional shape while limiting exposure of nearby normal tissue.

### **Confounding**

A distortion that occurs when another factor is related to both an exposure and an outcome, making their relationship appear stronger, weaker, or different from reality. For example, age may affect both an exposure and disease risk. Study design and statistical adjustment can reduce confounding.

### **Confusion**

Impaired ability to think clearly, orient, attend, remember, or respond appropriately; sudden confusion is delirium and may indicate infection, medication effect, metabolic illness, stroke, or another emergency.

### **Confusion Assessment Method**

A bedside assessment for delirium. It checks for a sudden or fluctuating change, poor attention, disorganized thinking, and an altered level of alertness; the pattern helps identify delirium but does not replace finding its cause.

### **Congenital**

Present at birth, whether inherited, caused by an early developmental difference, or acquired before birth through infection, exposure, or another condition.

### **Congenital Adrenal Hyperplasia**

A group of usually autosomal recessive disorders in which the adrenal glands cannot make hormones normally. The most common form can cause salt loss, excess androgen effects, early puberty, or genital differences and may be life-threatening in newborns.

---

### **Congenital Antithrombin III Deficiency**

An inherited condition in which the blood lacks enough working antithrombin, a natural substance that limits clotting. It increases the risk of clots in deep veins or the lungs, sometimes from a young age.

### **Congenital Biliary Atresia**

Alternate name; see *Biliary Atresia*.

### **Congenital Birth Defect**

A structural or functional abnormality present before or at birth. Causes may include genetic changes, infections, nutritional deficiencies, medicines, environmental exposures, or unknown factors.

### **Congenital Cataract**

Clouding of the eye lens present at birth or developing soon afterward. It can interfere with visual development and may require early eye assessment and treatment to prevent lasting reduced vision.

### **Congenital CMV**

Cytomegalovirus infection acquired before birth. Some babies have small head size, hearing or vision loss, seizures, jaundice, an enlarged liver or spleen, or developmental problems; others appear well initially but develop later hearing loss.

### **Congenital CNS Malformation**

A difference in the structure of the brain, spinal cord, or related tissues that develops before birth. Effects range from none to problems with movement, learning, seizures, breathing, or other functions, depending on the area involved.

### **Congenital Contractural Arachnodactyly**

An inherited condition causing unusually long, slender fingers and toes, stiff or bent joints from birth, thin muscles, and sometimes a curved spine. Joint stiffness often improves with age, but movement and heart or aortic findings may need monitoring.

### **Congenital Cytomegalovirus**

Cytomegalovirus infection present at birth after

transmission during pregnancy. Some infants have hearing loss, vision problems, small head size, seizures, or developmental disability; others have no symptoms initially.

### **Congenital Cytomegalovirus Infection**

Cytomegalovirus infection acquired before birth. It may cause hearing loss, developmental problems, microcephaly, or no symptoms.

### **Congenital Diaphragmatic Hernia**

A hole in the muscle separating the chest and abdomen that allows abdominal organs to move into the chest before birth. The crowded lungs may be underdeveloped, causing severe breathing problems after delivery.

### **Congenital Diaphragmatic Hernia Repair**

Surgery to return abdominal organs to the abdomen and close the opening in the diaphragm. It is performed after the newborn is stable enough for surgery; breathing support may be needed before and afterward.

### **Congenital Disorder**

A medical condition or structural abnormality present in an individual from birth, which may be caused by genetic factors or environmental influences in the womb.

### **Congenital Fibrinogen Deficiency**

An inherited absence or abnormality of fibrinogen that can cause bleeding, thrombosis, or both depending on the specific molecular defect.

### **Congenital Glaucoma**

High pressure in a baby's eye caused by abnormal development of the eye's drainage system. It can enlarge the eye and cause watering, light sensitivity, or cloudy vision; early treatment helps protect sight.

### **Congenital Heart Defect**

A structural abnormality of the heart or major vessels present at birth. It may alter blood flow and range from a small defect needing monitoring to

---

a serious condition requiring urgent treatment or surgery.

### **Congenital Heart Defect Corrective Surgery**

Surgical or transcatheter interventions designed to anatomically or physiologically correct congenital cardiovascular anomalies (such as atrial/ventricular septal defects, tetralogy of Fallot, or transposition of the great arteries).

#### **Alternate Names:**

Congenital Heart Surgery; Pediatric Cardiac Surgery; Corrective Cardiac Repair

### **Congenital Heart Defects**

Alternate form; see *Congenital Heart Defect*.

### **Congenital Heart Defects In Children**

Alternate index label; see *Congenital Heart Defects*.

### **Congenital Heart Disease**

A heart or major blood-vessel problem present from birth. It may change blood flow, cause no symptoms, or lead to breathing difficulty, blue color, poor growth, heart failure, or abnormal rhythms. Treatment ranges from monitoring and medicines to catheter procedures or surgery.

### **Congenital Heart Disease In Adults**

A heart defect present from birth that continues into adulthood; see *Congenital Heart Defects*.

### **Congenital Heart Murmur**

A heart murmur heard in a newborn or infant. It may be innocent or may indicate a structural heart defect; assessment considers the murmur's characteristics and associated symptoms or examination findings.

### **Congenital Hemolytic Jaundice**

Jaundice present from or caused by an inherited condition that increases red-blood-cell destruction, producing excess unconjugated bilirubin; examples include hereditary spherocytosis and some enzymatic defects.

### **Congenital Hip Dislocation**

Abnormal development or displacement of the hip joint present at birth or developing in infancy, now generally called developmental dysplasia of the hip.

### **Congenital Hydrocephalus**

Hydrocephalus present at birth or caused by a developmental abnormality, genetic condition, or congenital infection; see *Hydrocephalus*.

### **Congenital Hyperinsulinism**

An inherited or developmental disorder in which a baby's pancreas releases too much insulin. This can cause dangerously low blood sugar, poor feeding, seizures, brain injury, or death unless recognized and treated promptly.

### **Congenital Hyperthyroidism**

Excess thyroid hormone in a newborn, most often caused by thyroid-stimulating antibodies passing from a mother with Graves disease. It may cause a fast heartbeat, irritability, poor feeding, poor weight gain, or an enlarged thyroid.

### **Congenital Hypertrichosis Lanuginosa**

A very rare form of congenital hypertrichosis in which fine, soft lanugo-like hair covers much of the body from birth or early infancy. It may occur alone or as part of a genetic syndrome; examination looks for associated dental, skeletal, neurologic, or developmental findings.

### **Congenital Hypothyroidism**

Too little thyroid hormone from birth, most often because the thyroid is missing, misplaced, or underdeveloped, or cannot make hormone normally. Newborn screening and early treatment are essential because untreated deficiency can impair growth and brain development.

### **Congenital Malaria**

The transmission of Plasmodium parasites (predominantly Plasmodium falciparum or Plasmodium vivax) from an infected mother to the fetus across the placenta or during delivery. Although relatively uncommon due to the protective barrier of mater-

---

nal antibodies and placental syncytiotrophoblast, affected newborns can present in the neonatal period with fever, irritability, severe hemolytic anemia, jaundice, hepatosplenomegaly, thrombocytopenia, and poor feeding.

**Alternate Names:**

Congenital Plasmodium Infection; Neonatal Malaria

### **Congenital Malformation**

A structural abnormality that develops before birth and is present at birth, caused by genetic, environmental, developmental, or unknown factors.

### **Congenital Megacolon**

A birth condition in which part of the intestine lacks the nerve cells needed to move stool. It causes blockage and enlargement of the bowel above it, with delayed first stool, swelling, vomiting, or severe constipation.

### **Congenital Melanocytic Nevus**

A dark mole present at birth or appearing soon afterward, caused by a collection of pigment-producing cells. Most remain harmless, but larger lesions have a higher melanoma risk and need specialist monitoring.

### **Congenital Mitral Valve Anomalies**

Birth differences in the mitral valve, which controls blood flow from the upper to the lower left heart chamber. The valve may leak or be narrowed, causing no symptoms or leading to breathlessness, poor growth, or heart failure.

### **Congenital Muscular Dystrophy**

A group of inherited muscle disorders causing weakness or low muscle tone at birth or during early infancy. Some types also affect the brain, eyes, breathing, heart, or joints; severity varies widely.

**Alternate Names:**

CMD; Congenital Dystrophy

### **Congenital Myasthenic Syndromes**

A group of inherited disorders that disrupt communication between nerves and muscles. They cause

weakness that worsens with activity and may affect the eyes, swallowing, breathing, or limb movement.

### **Congenital Myopathies**

A group of inherited muscle disorders whose weakness or low muscle tone is present at birth or begins in infancy or childhood. They differ in their genetic cause and biopsy features and may affect movement, breathing, swallowing, the spine, or the heart.

### **Congenital Nasolacrimal Duct Obstruction**

Blockage of the tear-drainage channel present at birth, commonly causing persistent watering or discharge from one or both eyes. Most cases occur because a membrane at the lower end of the duct has not opened normally.

### **Congenital Nephrotic Syndrome**

Severe kidney disease beginning before birth or during the first three months of life, causing large amounts of protein loss in urine, swelling, low blood protein, poor growth, infections, and increased clotting risk.

### **Congenital Nevi**

Multiple melanocytic nevi present at birth or appearing soon afterward. Most are benign, but melanoma risk rises with larger or numerous lesions, so changes in colour, size, surface, or symptoms should prompt specialist review.

### **Congenital Nevus**

A dark or raised mole present at birth or soon after. Most are harmless, but larger or unusual moles have a higher risk of melanoma and should be checked from time to time. Imaging or removal is considered only for selected high-risk moles.

**Alternate Names:**

Congenital Melanocytic Nevus; Congenital Melanocytic Naevus; CMN; Congenital Pigmented Nevus; Bathing Trunk Nevus

### **Congenital Penile Curvature**

Curvature of the penis present from birth, usually caused by unequal development of the erectile tis-



---

sues. Mild curvature may need no treatment; significant curvature causing functional difficulty may require specialist assessment.

### **Congenital Plagiocephaly**

A difference in the shape of a baby's head that is present at birth or develops soon afterward. It may result from position before or after birth and differs from craniosynostosis, in which skull bones join too early.

### **Congenital Platelet Function Defects**

Inherited disorders in which platelets do not work normally even though their number may be normal. They can cause easy bruising, nosebleeds, heavy menstrual bleeding, or excessive bleeding after injury or surgery.

### **Congenital Pneumonia**

A lung infection present at birth, acquired before or during delivery. A newborn may have rapid or difficult breathing, temperature instability, poor feeding, or sepsis and needs urgent evaluation and treatment.

### **Congenital Protein C Or S Deficiency**

An inherited lack or abnormality of protein C or protein S, natural substances that limit clotting. It increases the risk of blood clots in veins, sometimes beginning in infancy or childhood.

### **Congenital Rubella**

Fetal infection with rubella virus acquired during pregnancy, potentially causing hearing loss, cataracts, congenital heart disease, developmental impairment, and other findings.

### **Congenital Rubella Syndrome**

Birth defects caused by rubella infection during pregnancy, especially early pregnancy. Affected babies may have hearing loss, cataracts, heart defects, small head size, developmental problems, enlarged organs, or a low platelet count.

### **Congenital Scoliosis**

A sideways curve of the spine caused by vertebrae that formed abnormally before birth. The curve may worsen as the child grows, so its pattern and progression require regular assessment.

### **Congenital Syphilis**

Syphilis infection passed from a pregnant person to the fetus. It can cause miscarriage, stillbirth, premature birth, rash, liver or bone disease, hearing loss, or problems that appear later; newborns need prompt testing and treatment.

### **Congenital Toxoplasmosis**

Infection with the parasite *Toxoplasma gondii* passed to a fetus during pregnancy. It may cause eye inflammation, brain changes, enlarged fluid spaces, seizures, hearing or vision problems, or developmental difficulties; some babies have no signs at birth.

### **Congenital Tracheomalacia**

Congenital tracheomalacia is weakness of the infant tracheal cartilage causing dynamic airway collapse, noisy breathing, cough, or recurrent respiratory symptoms.

### **Congenital Vaginal Obstruction**

A blockage of the vaginal canal present at birth, caused by missing or abnormal vaginal tissue, an obstructing membrane, or another developmental difference. It may prevent menstrual blood from leaving the body after puberty and cause pelvic pain.

#### **Alternate Names:**

OEIS Complex

### **Congenital Varicella Syndrome**

A rare congenital disorder occurring in infants born to mothers who contracted primary varicella zoster virus infection (chickenpox) during the first or early second trimester of pregnancy (typically 8 to 20 weeks gestation). Clinical hallmarks include zig-zag cicatricial skin scarring in a dermatomal distribution, unilateral limb hypoplasia, chorioretinitis, optic atrophy, microcephaly, and autonomic or central nervous system dysfunction.



---

**Alternate Names:**

Fetal Varicella Syndrome; Congenital Varicella Embryopathy

**Congenital Zika Syndrome**

A group of birth problems caused by Zika virus infection during pregnancy. Babies may have a small head, abnormal brain development, eye or hearing problems, stiff joints, or developmental difficulties; care focuses on the child's specific needs because there is no single cure.

**Congestion**

Excess blood or fluid in a tissue because drainage is reduced. It may cause swelling, heaviness, breathlessness, or impaired organ function, depending on the affected area and the underlying cause.

**Congestive Heart Failure**

Older term; see *Heart Failure*.

**Congestive Hepatopathy**

Liver injury caused by chronically elevated venous pressure, usually from right-sided heart failure or constrictive pericardial disease. It may cause liver enlargement, abnormal liver tests, ascites, and eventually cardiac cirrhosis.

**Conization**

Surgical removal of a cone-shaped portion of the cervix for diagnosis or treatment of cervical dysplasia or selected early cervical cancers.

**Conjoined Twins**

Identical twins who are physically joined because the early embryo did not completely separate. They may share organs or body structures, so detailed imaging and care by specialists are needed; the outlook depends on what is shared.

**Conjugated Equine Estrogens**

A mixture of estrogenic compounds historically derived from the urine of pregnant mares and used in some hormone-therapy formulations.

**Alternate Names:**

CEE; Conjugated Estrogens

**Conjugated Hyperbilirubinemia**

An elevation of direct, water-soluble bilirubin in the blood, usually caused by impaired bile formation or excretion, bile-duct obstruction, or hepatocellular injury. Jaundice, dark urine, pale stools, or itching may occur.

**Conjugated Linoleic Acid**

A group of related fats found mainly in meat and dairy products and sold as supplements. Claims about weight loss or muscle improvement are inconsistent, and supplements may adversely affect blood sugar or liver function.

**Conjunctiva**

The thin, clear tissue lining the inside of the eyelids and covering the white part of the eye. It protects and lubricates the eye and helps the eyelids move smoothly across its surface.

**Conjunctival**

Relating to the conjunctiva, the thin mucous membrane covering the front of the eye and lining the inner eyelids.

**Conjunctival Melanoma**

A rare malignant melanoma of the conjunctiva, often appearing as a pigmented or occasionally amelanotic lesion on the eye surface. It can recur locally or spread to lymph nodes and distant organs, so ophthalmic examination, biopsy, and surveillance are important.

**Conjunctival Squamous-Cell Carcinoma**

A cancer arising from the surface cells of the conjunctiva, often near the border of the cornea. It may appear as a persistent growth, redness, or irritation and can invade nearby eye tissue if not treated.

**Conjunctivitis**

Inflammation of the clear membrane covering the front of the eye and lining the eyelids. It may cause redness, itching, watering, irritation, or discharge

---

and can result from infection, allergy, or chemical exposure.

### Conjunctivoplasty

Plastic or reconstructive surgery of the conjunctiva, performed to repair defects, scarring, adhesions, or other abnormalities of the ocular surface.

### Conn Syndrome

Primary aldosteronism caused by autonomous overproduction of aldosterone by one or both adrenal glands. It usually causes high blood pressure and may cause low blood potassium, although potassium can be normal.

### Conn's Syndrome

Alternate name for primary aldosteronism, in which one or both adrenal glands make too much aldosterone.

### Connective Tissue

A diverse type of tissue that supports, binds, protects, and insulates organs and structures throughout the body.

### Connective Tissue Disease

A disorder affecting tissues that support and connect organs, often because the immune system attacks the body's own tissues. It may involve skin, joints, muscles, blood vessels, lungs, kidneys, or other organs. Examples include lupus, systemic sclerosis, rheumatoid arthritis, and Sjögren syndrome.

### Connexin 26 Deficiency

An inherited change in the *GJB2* gene that can cause hearing loss present from birth. The severity and whether one or both ears are affected can vary.

#### Alternate Names:

GJB2-Related Deafness; DFNB1 Hearing Loss

### Consanguinity

A biological relationship between people who share a recent common ancestor; it can increase the chance that both parents carry the same recessive disease variant.

### Conscious Sedation For Surgical Procedures

Medicines given during a procedure to make a person relaxed and sleepy while allowing them to respond to voice or touch and usually breathe independently. Continuous monitoring is needed because breathing, blood pressure, and alertness can become dangerously reduced.

### Consolidation

An imaging finding in which air in part of the lung's air spaces is replaced by fluid, pus, blood, or cells, making that area appear denser. It may occur with pneumonia, fluid buildup, bleeding, or a tumor; consolidation describes an appearance, not a specific disease.

### Consolidation Therapy

Additional treatment given after an initial response or remission to destroy remaining disease and lower the chance of relapse. It may include medicines, radiation, surgery, or a transplant depending on the illness.

### Constipation

A condition involving fewer than three bowel movements per week, hard or lumpy stools, difficult or painful passage, straining, or a feeling that the bowel has not emptied completely. Causes include diet, medicines, dehydration, functional bowel disorders, and structural or systemic disease.

### Constitutional Delay In Puberty

A common pattern in which puberty starts later than average but eventually progresses normally. Growth and bone development may also be delayed, often with a family history of late puberty.

### Constitutional Growth Delay

A normal variant in which height growth and bone maturation are temporarily behind average, often with a family history of late puberty. Most affected children continue growing and enter puberty later than peers but reach a typical adult height.

#### Alternate Names:

Constitutional Delay of Growth and Puberty; CGD

---

## **Constraint-Induced Movement Therapy**

Rehabilitation that limits use of a less-affected limb while providing structured practice with an affected limb to improve functional use in selected people.

## **Constrictive Pericarditis**

A chronic condition in which a thickened or scarred, noncompliant sac around the heart restricts diastolic filling. It can cause breathlessness, fatigue, leg or abdominal swelling, enlarged neck veins, and liver congestion; causes include infection, surgery, radiation, or unknown disease.

## **Consultation-Liaison Psychiatry**

Psychiatric care for people who are being treated for a physical illness. The psychiatrist helps assess and treat problems such as confusion, depression, anxiety, substance withdrawal, or difficulty coping during medical care.

## **Contact Overdose**

Overdose of a combination cold medicine that may cause acetaminophen toxicity, antihistamine or decongestant effects, sedation, seizures, or cardiac complications depending on ingredients.

## **Contact Allergen Testing**

Patch testing checks whether a substance causes allergic skin inflammation. Small amounts of suspected allergens are placed on the skin, usually the back, and the area is examined after one or more days for a delayed reaction.

## **Contact Dermatitis**

Red, itchy, burning, or blistered skin caused by contact with an irritating or allergy-triggering substance. Common triggers include soaps, chemicals, metals, fragrances, plants, and medicines.

### **Alternate Names:**

Allergic Contact Dermatitis; Irritant Contact Dermatitis

## **Contact Lenses**

Thin lenses placed directly on the eye to correct vision or deliver selected treatment. Proper fit-

ting, handwashing, cleaning, and replacement are essential because misuse can cause corneal infection, inflammation, or vision-threatening injury.

## **Contact Precautions**

Steps used to prevent spread of germs by touch. They may include gloves and a gown, hand hygiene, a private room, and equipment kept for one patient, depending on the infection and local policy.

## **Contact Tracing**

Identifying and monitoring persons exposed to infectious disease. Critical for outbreak control. Examples: COVID-19, Ebola, TB.

## **Contact Urticaria**

Hives, itching, or swelling that develop within minutes to about an hour after a substance touches the skin or mucosa. It may be immunoglobulin-E-mediated, such as with latex or foods, or caused by direct irritation; widespread symptoms, wheezing, throat swelling, or faintness suggest anaphylaxis.

## **Contagious Disease**

An infectious disease that can be transmitted directly or indirectly from one person, animal, or source to another.

## **Contamination**

Unwanted microorganisms or other material entering a specimen, surface, medicine, or normally sterile area without causing infection in the person. It can produce misleading test results and must be distinguished from true disease.

## **Continuing Medical Education**

Education after professional training that updates a healthcare worker's knowledge, skills, judgment, and ability to provide safe, effective patient care.

## **Continuity Of Care**

Consistent and coordinated care over time and across clinicians or settings, integrating the patient's history, current needs, treatment plan, and preferences. It supports safety, adherence, prevention,

---

and chronic-disease outcomes.

### **Continuous Bladder Irrigation**

Continuous flow of sterile fluid through a urinary catheter into and out of the bladder. It helps prevent blockage and wash out blood clots or debris, often after urinary-tract surgery.

### **Continuous Combined Hormone Therapy**

A hormone treatment regimen taking estrogen and progestogen continuously, often to manage menopausal symptoms while reducing regular withdrawal bleeding.

#### **Alternate Names:**

Continuous Combined HRT; Continuous Estrogen-Progestogen Therapy

### **Continuous Electronic Fetal Monitoring**

Continuous recording of the baby's heart rate and the pregnant person's contractions during labor. It may identify patterns suggesting that the baby needs assessment or changes in care.

### **Continuous Glucose Monitoring**

A wearable sensor that measures glucose in fluid beneath the skin at regular intervals and shows current levels, trends, and alerts. It helps manage diabetes but may need confirmation with a blood-glucose test when readings do not fit symptoms.

### **Continuous Kidney Replacement Therapy**

A slow, continuous form of blood purification running 24 hours a day in intensive care units for severe acute kidney injury. It gradually removes excess fluid and metabolic toxins, making it suitable for critically ill patients who are too unstable to tolerate standard hemodialysis.

#### **Alternate Names:**

CKRT; Continuous Renal Replacement Therapy; CRRT

### **Continuous Murmurs**

Heart sounds heard during both the heart's contraction and relaxation because blood flows continuously through an abnormal opening or connection.

A persistent murmur needs assessment to identify its cause.

### **Continuous Passive Motion Machine**

A device that gently moves a limb through a set range of motion while the person relaxes. It may be used after selected operations, especially knee surgery, but does not replace active rehabilitation.

### **Continuous Positive Airway Pressure**

Noninvasive respiratory support that delivers a constant positive pressure throughout the breathing cycle to keep the upper airway or alveoli open.

### **Continuous Renal Replacement Therapy**

A continuous, 24-hour-a-day blood filtration and dialysis therapy performed in intensive care units for critically ill patients with acute kidney failure who are hemodynamically unstable (such as those with severe septic shock, multiple organ failure, or very low blood pressure). Unlike standard intermittent hemodialysis, which cleanses blood rapidly over three to four hours and can trigger dangerous drops in blood pressure, CRRT cleanses blood and removes excess fluid very slowly and gently through an extracorporeal circuit. Blood is pumped through a synthetic filter (hemofilter) where metabolic waste products, toxins, and fluid are removed by diffusion and convection before the blood is returned to the body. Modalities include continuous venovenous hemofiltration (CVVH), hemodialysis (CVVHD), and hemodiafiltration (CVVHDF).

#### **Alternate Names:**

CRRT; Continuous Kidney Replacement Therapy; CKRT; Continuous Hemodiafiltration

### **Continuous Subcutaneous Insulin Infusion**

Delivery of rapid-acting insulin through a pump and a subcutaneous infusion set to provide basal insulin and mealtime doses.

### **Contraception**

Any method used to prevent pregnancy, including intrauterine devices, implants, pills, injections, condoms, fertility-awareness methods, and sterilization.

---

Methods differ in effectiveness, duration, side effects, cost, and protection against sexually transmitted infections. The best choice depends on health, preferences, reproductive plans, and access to care.

### **Contraception And Birth Control**

Contraception encompasses methods used to prevent pregnancy, including hormonal options like oral contraceptives, patches, rings, injections, and implants, as well as non-hormonal methods like copper IUDs, barrier methods, diaphragms, and permanent sterilization.

### **Contraception For Adolescents**

Methods to prevent unintended pregnancy. Options: combined oral contraceptives, progestin-only pills, depot medroxyprogesterone acetate, contraceptive implant (Nexplanon), intrauterine devices (IUDs), condoms, and abstinence.

### **Contraceptive Implant**

A small medicine-releasing device placed under the skin, usually in the upper arm, to prevent pregnancy for an extended period. It releases a progestogen and can cause irregular bleeding or other hormone-related effects.

### **Contraceptives**

Methods used to prevent pregnancy, including hormonal medicines, implants, intrauterine devices, condoms, and permanent procedures. Methods differ in effectiveness, duration, and risks; most do not protect against sexually transmitted infections, while condoms reduce but do not eliminate that risk.

### **Contracted Pelvis**

A pelvis whose size or shape is reduced or abnormal enough to interfere with passage of the fetus during labor.

### **Contractile Proteins**

Proteins, especially actin and myosin, that interact inside muscle cells to generate force, shorten fibers, and produce body movement.

**Alternate Names:**

Muscle Contractile Proteins; Actomyosin Proteins

### **Contraction**

Shortening or tightening of a muscle; uterine contractions during labor thin and dilate the cervix and help deliver the fetus.

### **Contraction Stress Test**

A pregnancy test that records the baby's heart rate while contractions occur, sometimes after gentle stimulation or medicine. Repeated slowing of the heart rate may suggest that the placenta cannot supply enough oxygen during labor.

### **Contractions**

Rhythmic tightening of the uterus during labor that thins and opens the cervix and helps move the baby through the birth canal. Before labor, Braxton Hicks contractions are often irregular and do not progressively open the cervix.

### **Contracture**

A lasting inability to fully move a joint because nearby muscles, tendons, ligaments, skin, or other tissues have shortened or stiffened. It may follow injury, burns, paralysis, inflammation, or prolonged immobility.

### **Contracture Deformity**

A fixed limitation of movement caused by shortening or tightening of muscle, tendon, ligament, joint capsule, skin, or other soft tissue. It may follow neurologic disease, prolonged immobility, injury, burns, or inflammation.

### **Contracture Prevention**

Measures such as positioning, stretching, exercise, splinting, orthoses, and serial casting that help preserve joint movement in people at risk of fixed shortening.

### **Contraindication**

A condition or circumstance in which a treatment, procedure, or medication should not be used because the potential harm outweighs the expected benefit.



---

**Alternate Names:**

Contraindications

**Contralateral**

On or affecting the side opposite a stated body part or injury. For example, a stroke on the left side of the brain may cause weakness on the right side of the body.

**Contrast**

A difference that makes structures distinguishable; in imaging, contrast material is administered to improve visualization of organs, vessels, or tissues.

**Contrast Agent**

A substance given or placed in the body to make organs, tissues, blood vessels, or body spaces easier to see during imaging. Different agents are used for X-ray, CT, MRI, ultrasound, or digestive-tract studies.

**Contrast Echocardiography**

An ultrasound examination of the heart using tiny gas-filled bubbles in the bloodstream to make heart chambers, blood flow, or selected areas of heart muscle easier to see.

**Contrast Medium**

A substance administered or placed in the body to make tissues, organs, or body spaces more visible during imaging.

**Alternate Names:**

Contrast Agent; Radiographic Contrast Material

**Contrast Radiology**

Imaging of the gastrointestinal tract using contrast material, such as barium, to outline the esophagus, stomach, intestine, or colon on radiographs or fluoroscopy.

**Contrast Reaction**

An unwanted response after an imaging contrast agent, ranging from nausea, itching, or a rash to breathing difficulty, low blood pressure, or a severe allergic-like reaction. Imaging staff should be told

about previous reactions.

**Contrast-Associated Acute Kidney Injury**

Acute worsening of kidney function occurring after administration of contrast, whether or not contrast is causal. Risk assessment includes baseline kidney function, hemodynamic status, diabetes, nephrotoxins, urgency, and the need to avoid unnecessary contrast exposure.

**Contrast-Enhanced Computed Tomography**

A CT scan performed after contrast material makes selected organs, blood vessels, or abnormal areas easier to see.

**Contrast-Enhanced Magnetic Resonance Imaging**

An MRI scan performed after contrast material makes selected tissues, blood vessels, or abnormal areas easier to see.

**Contrast-Enhanced Ultrasound**

An ultrasound examination using tiny gas-filled bubbles in the bloodstream to improve pictures of blood flow and help characterize selected liver, heart, blood-vessel, or other lesions.

**Contrast-Induced Nephropathy**

A sudden decline in kidney function after an imaging test using iodinated contrast, although the contrast is not always the cause. Risk is higher with severe kidney disease, dehydration, low blood pressure, or other kidney stresses.

**Contrecoup**

An injury occurring on the side of an organ opposite the site of impact, especially a brain injury produced when the brain moves within the skull after trauma.

**Control Group**

The comparison group in a study, which may receive usual care, a placebo, or no intervention according to the study design.



---

### **Controlled Cord Traction**

Gentle pulling on the umbilical cord after the placenta has separated, while supporting the uterus. A trained birth attendant may use it to help deliver the placenta and reduce blood loss.

### **Controlled Ovarian Hyperstimulation**

Use of hormone medicines to make several ovarian follicles mature at the same time for fertility treatment. Ultrasound and blood tests guide dosing and help reduce the risk of ovarian hyperstimulation, a potentially serious complication.

#### **Alternate Names:**

COH; Ovarian Stimulation

### **Controlled-Substance Schedule I**

A United States controlled-substance category for drugs with high misuse potential and no accepted medical use under federal law.

### **Controlled-Substance Schedule II**

A United States legal category for medicines with accepted medical uses but high misuse, dependence, or overdose potential. Prescribing, dispensing, refills, and storage are subject to strict rules.

### **Controlled-Substance Schedule III**

A United States legal category for medicines with accepted medical uses and less misuse potential than Schedule II substances, although dependence and misuse remain possible. Rules vary by medicine and jurisdiction.

### **Controlled-Substance Schedule IV**

A United States controlled-substance category for drugs with accepted medical uses and lower misuse potential than Schedule III drugs.

### **Controlled-Substance Schedule V**

A United States controlled-substance category for drugs with the lowest level of misuse potential among the federal schedules.

### **Controllers**

Long-term asthma medicines taken regularly to re-

duce airway inflammation, prevent symptoms and attacks, and lower the need for quick-relief inhalers.

#### **Alternate Names:**

Controller Medicines; Maintenance Asthma Therapy

### **Contusion**

A bruise caused by a blunt blow that damages small blood vessels without breaking the skin. It may cause pain, swelling, and discoloration; severe injuries can damage muscle, organs, or deeper tissues.

#### **Alternate Names:**

Contusions

### **Conus Medullaris Syndrome**

A group of symptoms caused by damage to the lower end of the spinal cord. It may cause back pain, weakness or numbness in the legs and pelvic area, and loss of bladder, bowel, or sexual function.

### **Convergence**

The inward movement of both eyes so that they are directed toward the same nearby object.

### **Convergence Insufficiency**

Difficulty turning both eyes inward together to focus on something nearby. It can cause eyestrain, headaches, blurred or double vision, and reading difficulty, especially in children and young adults.

### **Conversion Disorder**

A condition causing genuine problems such as weakness, abnormal movements, loss of sensation, or seizure-like episodes because of altered nervous-system functioning. The symptoms are not intentionally produced and can interfere with daily life.

#### **Alternate Names:**

Functional Neurological Symptom Disorder

### **Convulsion**

A sudden episode of involuntary, often rhythmic muscle contractions that may occur with a seizure or other disturbance of brain function. It describes the motor activity, and not all seizures cause convulsions; some produce staring, unusual sensations,

---

altered awareness, or other non-motor symptoms.

**Alternate Names:**

Convulsive Episode; Seizure-Related Motor Activity

**Cooking Without Salt**

Preparing food with little or no added sodium to reduce dietary sodium intake, often useful in blood-pressure or fluid-retention management when combined with an individualized eating plan.

**Cooley's Anemia**

Alternate index label; see *Thalassemia*.

**Coombs Test**

A blood test that detects antibodies affecting red blood cells. It helps investigate immune-related destruction of red cells and checks compatibility before some blood transfusions.

**Cooper's Ligament**

Fibrous connective-tissue septa extending from the deep surface of the breast toward the skin and pectoral fascia. Tumor invasion or fibrosis may cause skin retraction.

**Coordination**

The ability to control movements smoothly, accurately, and in the correct order. It depends on the brain, nerves, muscles, vision, and balance and may be tested with simple hand or leg movements.

**COPD**

See *Chronic Obstructive Pulmonary Disease*.

**COPD And Other Health Problems**

The interaction between chronic obstructive pulmonary disease and conditions such as cardiovascular disease, osteoporosis, depression, diabetes, and lung cancer, which can complicate symptoms and care.

**COPD Exacerbation**

Acute worsening of respiratory symptoms beyond normal day-to-day variation in a patient with COPD that leads to additional therapy. Common triggers

include viral or bacterial infection, air pollution, and cardiopulmonary comorbidity.

**COPD Flare-Ups**

Acute worsening of breathlessness, cough, or sputum beyond usual daily variation in chronic obstructive pulmonary disease, often triggered by infection, pollution, or other irritants.

**COPD Maintenance Medications**

Pharmacologic agents administered on a daily, scheduled regimen to prevent exacerbations, reduce symptoms, and slow airflow limitation in chronic obstructive pulmonary disease, including long-acting muscarinic antagonists (LAMAs), long-acting beta-2 agonists (LABAs), and inhaled corticosteroids (ICS).

**Alternate Names:**

COPD Controller Medications; Long-Acting COPD Bronchodilators; COPD Maintenance Therapy

**COPD Rehabilitation**

A supervised program combining exercise, breathing strategies, education, nutrition, and support to help people with COPD breathe and function better, remain active, and reduce the effects of flare-ups.

**COPD Rescue Medications**

Short-acting inhaled bronchodilators administered on an as-needed basis to rapidly relieve acute dyspnea and bronchoconstriction during COPD symptom flares, encompassing short-acting beta-2 agonists (SABAs) and short-acting muscarinic antagonists (SAMAs).

**Alternate Names:**

COPD Quick-Relief Drugs; COPD Reliever Inhalers; Fast-Acting COPD Bronchodilators

**Copper**

An essential trace element required for iron transport, connective-tissue cross-linking, energy metabolism, neurotransmitter synthesis, and antioxidant enzymes. Deficiency can cause anemia, neutropenia, impaired neurologic function, and bone abnormalities; excess is toxic and accumulates in Wilson disease.

---

## Copper Deficiency

A lack of copper, a trace element needed for iron transport, connective tissue, nerve function, and pigment production. It may follow malnutrition, malabsorption, gastrointestinal surgery, excess zinc, or an inherited disorder such as Menkes disease, and can cause anemia, low white-cell count, bone changes, and nerve or spinal-cord damage. Copper replacement treats deficiency when intake or absorption is inadequate.

## Copper In Diet

Copper obtained from foods such as shellfish, nuts, seeds, beans, whole grains, and organ meats. The body needs small amounts, while excessive supplements or exposure can be harmful; intake should be tailored for deficiency or disease.

## Copper Poisoning

Toxic exposure to copper or copper salts that may cause vomiting, abdominal pain, hemolysis, liver injury, kidney injury, or shock.

## Coprolalia

Involuntary utterance of obscene, taboo, or socially inappropriate words, most often described as a complex vocal tic in Tourette syndrome.

## Copy Number Variant

A genetic change in which a section of DNA is missing or present in extra copies. Whether it affects health depends on its size, location, genes, inheritance, and available evidence.

## Copy Number Variation

A deletion or duplication of a section of DNA that changes how many copies are present. Some variations are harmless, while others alter gene activity and cause developmental, neurologic, or other disorders.

## CoQ10

A fat-soluble substance made by the body that helps mitochondria produce energy and also acts as an antioxidant. It is sold as a supplement, but benefit varies by condition and it can interact with

medicines.

## Cor

A Latin anatomical word meaning heart, appearing in medical terms that describe the heart, heart-related structures, or diseases affecting circulation.

## Cor Pulmonale

Enlargement and weakening of the right side of the heart caused by high pressure in the lung arteries, usually from long-term lung disease. It may cause breathlessness, leg swelling, enlarged neck veins, or liver congestion.

## Cor Triatriatum

A rare birth defect in which a membrane divides the upper left heart chamber into two spaces. If the membrane blocks blood returning from the lungs, it can cause breathlessness, poor growth, or heart failure; some people have no symptoms.

## Coracoid Process

A hook-shaped bony projection from the shoulder blade that points forward. Several muscles and ligaments attach to it, and injury may occur with a direct blow or shoulder dislocation.

## Cord

A long, narrow structure such as the spinal cord, which contains the central nervous tissue, or the umbilical cord, which connects a fetus to the placenta.

## Cord Blood Stem Cells

Blood-forming stem cells collected from the umbilical cord and placenta after birth. They can make different types of blood and immune cells and may be used for stem-cell transplantation.

## Cord Blood Testing

Laboratory testing of blood collected from the umbilical cord at birth, which may include blood type, complete blood count, blood gases, glucose, or tests for infection or inherited disease when clinically indicated.

---

## Cord Presentation

A position in which the umbilical cord lies between the baby's presenting part and the cervix while the membranes are intact. If the membranes break, the cord may slip down and become compressed, threatening the baby's oxygen supply.

## Cordocentesis

A prenatal procedure using ultrasound guidance to take a small blood sample from the umbilical cord. It can assess fetal anemia, infection, or selected genetic conditions but carries risks including bleeding and pregnancy loss.

## Cordotomy

A procedure that cuts selected pain-sensing pathways in the spinal cord to relieve severe pain when other treatments have failed. It may cause numbness, weakness, breathing problems, or other lasting nerve complications.

## Core Biopsy

Removal of one or more cylindrical tissue samples with a hollow cutting needle for microscopic examination. It preserves more tissue architecture than fine-needle aspiration and is used for breast, liver, prostate, and other lesions.

## Core Needle Biopsy

Removal of small cylinders of tissue with a hollow cutting needle for laboratory examination. Imaging may guide the needle, and the samples preserve tissue structure better than fluid or cell aspiration alone.

## Cori Cycle

A process in which muscles send lactate to the liver, where it is changed into glucose. The glucose returns to the blood and can be used again by muscles for energy.

## Cori Disease

A rare inherited disorder in which the body cannot break down stored glycogen normally. It can cause low blood sugar, an enlarged liver, muscle weakness, and sometimes heart disease.

## Alternate Names:

Glycogen Storage Disease Type III (GSD III); Forbes Disease; Debrancher Deficiency

## Corn

A localized thickening of skin caused by repeated pressure or friction, often occurring on the toes or feet.

## Alternate Names:

Clavus; Heloma

## Corn-Cockle Poisoning

Poisoning caused by ingestion of corn cockle (*Agrostemma githago*) seeds, which may contaminate grain and cause stomach upset, weakness, or other toxic effects.

## Alternate Names:

Githism (obsolete historical term); Agrostemma Githago Intoxication

## Cornea

The clear, curved front surface of the eye. It protects the eye and bends light so images focus on the retina; scratches, infection, swelling, or scarring can cause pain and blurred vision.

## Corneal Abrasion

A scratch or loss of the outer layer of the cornea, often from a fingernail, foreign body, contact lens, makeup tool, or other injury. It can cause pain, tearing, redness, light sensitivity, and blurred vision.

## Corneal Cross-Linking

A treatment that uses riboflavin and ultraviolet light to strengthen the cornea. It can slow or stop worsening thinning and bulging, especially in keratoconus, but may cause pain during healing.

## Corneal Disease

Corneal disease affects the transparent front surface of the eye through infection, inflammation, trauma, dystrophy, degeneration, dryness, or abnormal shape. Pain, redness, light sensitivity, discharge, or reduced vision can signal urgent disease because scarring may impair sight.

---

## Corneal Dystrophy

A group of usually inherited disorders in which abnormal material builds up in one or more layers of the cornea. They may cause blurred vision, glare, pain, or recurrent corneal erosions and often affect both eyes.

## Corneal Edema

Swelling of the cornea from excess fluid, making it cloudy and causing blurred vision, glare, halos, or pain. Causes include damage to its inner cells, glaucoma, inflammation, injury, or eye surgery.

## Corneal Endothelium

The innermost layer of cells in the cornea. It moves extra fluid out of the cornea and helps keep it clear.

## Corneal Epithelium

The thin outer layer of the cornea that protects the eye and helps keep its surface smooth.

## Corneal Implant

A device or tissue substitute placed in or on the cornea, the clear front surface of the eye, to change its shape, replace damaged tissue, or improve vision. Benefits and risks depend on the implant and eye condition.

## Corneal Inflammation

Alternate name; see *Keratitis*.

## Corneal Injury

Damage to the clear front surface of the eye, including scratches, foreign bodies, chemical burns, infections, or cuts that may threaten vision.

## Corneal Laceration

A cut through part or all of the cornea, usually caused by sharp trauma or a foreign object. It can cause pain, tearing, light sensitivity, reduced vision, and leakage of aqueous humor.

## Corneal Reflex

An involuntary protective reflex resulting in bilateral blinking upon mechanical or tactile stimulation

of either cornea. The sensory afferent limb is mediated by the ophthalmic division of the trigeminal nerve (cranial nerve V1) and the motor efferent limb is carried by the facial nerve (cranial nerve VII) supplying the orbicularis oculi muscles.

### Alternate Names:

Blink Reflex; Corneal Blink Reflex

## Corneal Stroma

The thick middle layer of the cornea. It gives the cornea much of its strength and helps keep it clear.

## Corneal Transplant

Surgery that replaces part or all of a damaged or diseased cornea with healthy donor tissue. It may restore vision or relieve pain; risks include rejection, infection, increased eye pressure, and need for further treatment.

## Corneal Transplantation

Surgical replacement of part or all of a diseased or damaged cornea with donor corneal tissue to restore vision or relieve pain.

### Alternate Names:

Corneal Transplant

## Corneal Ulcer

A loss of tissue forming an open sore on the cornea, the clear front surface of the eye. It is most often caused by infection with bacteria, viruses, fungi, or parasites, but may also follow an injury, severe dryness, poor eyelid closure, or contact-lens problems. It may cause eye pain, redness, tearing, discharge, sensitivity to light, or blurred vision. Healing can leave a corneal scar, and a deep or central ulcer can seriously damage vision or perforate the eye.

## Corneal Ulcers And Infections

Infection or inflammation of the cornea that may produce an epithelial defect or ulcer, pain, redness, tearing, discharge, and light sensitivity. Bacterial, fungal, protozoal, and viral causes can threaten vision and require urgent ophthalmic assessment.

---

## Corneitis

Inflammation of the cornea, the clear front surface of the eye. It may cause pain, redness, watering, light sensitivity, or blurred vision and can result from infection, injury, immune disease, or chemical exposure.

## Cornelia De Lange Syndrome

A genetic developmental disorder that can cause distinctive facial features, slow growth, intellectual disability, limb differences, feeding problems, hearing loss, and heart or other organ abnormalities. Severity varies widely.

## Corns

Small, hard areas of thickened skin caused by repeated pressure or rubbing, usually on the toes or other pressure points. They may hurt when pressed; people with diabetes or poor circulation should avoid cutting them themselves.

## Corns And Calluses

Corns are small, localized areas of thickened skin caused by repeated pressure or friction, usually on the toes. Calluses are broader areas of thickened, hardened skin, usually on the soles of the feet or palms of the hands. Both are usually harmless but may become painful.

## Cornual Pregnancy

An ectopic pregnancy that implants in the part of the fallopian tube passing through the muscular wall of the uterus. It may grow silently before rupturing and causing life-threatening internal bleeding.

### Alternate Names:

Cornual Pregnancy; Interstitial Ectopic Pregnancy

## Coronal

Relating to a plane that divides the body into front and back portions, or to the crown or top of a structure.

## Coronal Caries

Tooth decay affecting the crown, the visible part of a tooth above the gum. It can begin in pits,

grooves, or smooth surfaces when bacterial acids dissolve enamel and dentin.

## Coronal Plane

A vertical plane that divides the body or an organ into front (anterior) and back (posterior) portions.

## Coronary

Relating to the arteries and blood flow that supply the heart muscle. Narrowing or blockage of these arteries can reduce oxygen to the heart and cause chest pain or a heart attack.

### Alternate Names:

Cardiac Blood-Vessel-Related; Heart-Coronary

## Coronary Aneurysm

An abnormal widening or ballooning of a coronary artery. It may result from atherosclerosis, inflammation, infection, injury, or a congenital condition and can cause clotting, blockage, or rarely rupture.

## Coronary Angiography

An X-ray examination of the heart arteries after contrast is injected through a thin catheter. It shows narrowed or blocked areas and can guide treatment, but the procedure can cause bleeding, stroke, kidney injury, or other complications.

## Coronary Angioplasty

A procedure that opens a narrowed or blocked artery supplying the heart. A thin tube with a small balloon is guided to the blockage. The balloon is inflated to widen the artery, and a small mesh tube called a stent is often left in place to help keep it open.

### Alternate Names:

Percutaneous Coronary Intervention; PCI; Coronary Balloon Angioplasty; PTCA

## Coronary Arteries

The blood vessels that arise from the aorta and supply oxygen to the heart muscle. The right and left arteries divide into branches, and blockage in one can cause angina or a heart attack.



---

## Coronary Artery

One of the blood vessels branching from the aorta that supplies oxygen-rich blood directly to the heart muscle.

## Coronary Artery Aneurysm

A weakened section of an artery supplying the heart that has become abnormally widened. It may form a blood clot, reduce blood flow, or rarely rupture; causes include artery disease, inflammation, infection, or injury.

## Coronary Artery Anomaly

A birth difference in where a heart artery begins, how it travels, or how it ends. Some cause no problems, while others reduce blood flow or increase the risk of chest pain, dangerous rhythms, or sudden cardiac arrest.

## Coronary Artery Bypass

A surgical procedure that creates a new route for blood flow around a narrowed or blocked coronary artery using a vascular graft.

## Coronary Artery Bypass Grafting

Surgery that uses a blood-vessel graft to create a route around severely narrowed or blocked heart arteries. It can improve blood flow and symptoms and may improve survival in selected people with extensive disease.

## Coronary Artery Bypass Surgery

An operation that uses a blood vessel from another part of the body to create a new route around narrowed or blocked heart arteries and improve blood flow to heart muscle.

### Alternate Names:

CABG; Coronary Bypass Surgery

## Coronary Artery Calcium Score

A non-invasive computed tomography measurement of calcified atheromatous plaque within the epicardial coronary arteries, quantified using the Agatston scoring method. It serves as an independent predictor of future major adverse cardiovascular

events and aids in cardiovascular risk stratification for borderline- or intermediate-risk asymptomatic adults.

### Alternate Names:

Agatston Score; CAC Score

## Coronary Artery Disease

A condition in which fatty plaque builds up in the arteries that supply blood to the heart muscle, narrowing them and reducing blood flow. It may cause no symptoms, chest discomfort (angina), or a heart attack, and can lead to heart failure or abnormal heart rhythms.

### Alternate Names:

Coronary Heart Disease

## Coronary Artery Fistula

An unusual connection between an artery supplying the heart and a heart chamber or another blood vessel. It may cause no symptoms or lead to breathlessness, chest pain, abnormal rhythms, heart failure, or infection of the heart lining.

## Coronary Artery Spasm

Temporary tightening of a heart artery that reduces blood flow and may cause chest pain, often at rest. Smoking, stimulant drugs, stress, or other vessel problems can trigger it.

## Coronary Calcium Score

A CT measurement of calcium in the arteries supplying the heart. More calcium usually means more plaque and higher future cardiovascular risk, but a zero score does not exclude soft plaque or eliminate all risk.

## Coronary Calcium Scoring

A computed tomography scan measuring calcium-containing plaque in heart arteries. The score helps estimate future cardiovascular risk in selected people but does not show all plaque or replace clinical assessment.

## Coronary Care Unit

A hospital unit for people with serious or unstable

---

heart conditions. Staff continuously monitor heart rhythm, blood pressure, oxygen, and other vital functions and provide urgent treatment for problems such as heart attack, dangerous rhythms, or heart failure.

**Alternate Names:**

CCU

### Coronary Circulation

The network of arteries and veins that supplies the heart muscle with oxygen and removes waste. Blood flow increases when the heart works harder and falls when an artery is narrowed or blocked.

### Coronary Computed Tomography Angiography

A noninvasive CT scan performed after injected contrast highlights the heart arteries. It can show plaque, narrowing, or blockage and is useful for selected people with suspected coronary artery disease.

**Alternate Names:**

CCTA; Coronary CT Angiography; Cardiac CTA

### Coronary Dominance

The pattern describing which heart artery gives rise to the vessel supplying the back part of the heart. Most people have right dominance; left or shared patterns are normal variations.

### Coronary Endarterectomy

Surgery removing hardened plaque from the inner lining of a heart artery. It may improve blood flow in selected complex blockages but is uncommon and carries risks of heart attack, bleeding, or vessel injury.

**Alternate Names:**

Coronary Plaque Endarterectomy; Coronary Endarterectomy Procedure

### Coronary Insufficiency

Blood flow through the heart arteries is not enough to meet the heart muscle's need for oxygen, usually because the arteries are narrowed. It may cause chest pain, especially during exertion, or a heart attack.

### Coronary Microvascular Angina

Chest pain caused by poor blood-flow regulation in the heart's smallest blood vessels. The main heart arteries may look open on testing, but the small vessels do not deliver enough oxygen during stress or sometimes at rest.

### Coronary Microvascular Disease

Alternate name; see *Small Vessel Disease*.

### Coronary Microvascular Dysfunction

Abnormal function of the small heart vessels that limits blood flow even when the major arteries are not severely narrowed. It may cause chest pain, especially in women and people with diabetes or high blood pressure.

### Coronary Occlusion

Obstruction of a coronary artery, commonly by atherosclerotic plaque and thrombus, that can cause myocardial ischemia or infarction.

### Coronary Perfusion Pressure

The physiological net pressure gradient driving myocardial blood flow through coronary capillary beds, calculated as the difference between aortic diastolic pressure and left ventricular end-diastolic pressure.

**Alternate Names:**

CPP

### Coronary Revascularization

Restoration of blood flow through the coronary arteries, usually by percutaneous coronary intervention (angioplasty with or without stenting) or coronary artery bypass grafting. It is used to treat selected obstructive coronary artery disease and acute coronary syndromes.

### Coronary Sinus

A large vein on the back of the heart that collects de-oxygenated blood from the heart muscle and drains it into the right atrium.

### Coronary Slow Flow Phenomenon

Delayed movement of blood through the small heart

---

vessels despite no major blockage on angiography. It may cause repeated chest pain at rest or with activity and can occasionally lead to abnormal rhythms or heart injury.

**Alternate Names:**

CSFP; Slow Coronary Flow; Primary Coronary Slow Flow

### Coronary Spasm

Alternate name; see *Coronary Artery Spasm*.

### Coronary Steal Phenomenon

A situation in which blood is diverted toward healthier heart muscle and away from an area already receiving too little blood. It can worsen chest pain during certain medicines or stress tests.

**Alternate Names:**

Coronary Steal Syndrome; Myocardial Steal

### Coronary Steal Syndrome

A hemodynamic phenomenon in which pharmacologically induced vasodilation (such as with dipyridamole or adenosine) diverts blood flow away from an already maximally dilated, stenotic coronary vascular bed toward well-perfused non-stenotic vessels, worsening regional myocardial ischemia.

### Coronary Stent

A small mesh tube placed in a narrowed heart artery to help keep it open after angioplasty. Medicines are usually needed afterward to reduce the risk of clotting.

### Coronary Thrombosis

Formation of a blood clot in a coronary artery, potentially obstructing blood flow and causing myocardial infarction.

### Coronavirus

A family of RNA viruses that infect people and many animals. Their surface has crown-like spike proteins, which give the family its name. Four common human coronaviruses usually cause mild or moderate upper-respiratory illnesses such as colds, while other human coronaviruses can cause severe diseases, in-

cluding severe acute respiratory syndrome (SARS), Middle East respiratory syndrome (MERS), and COVID-19. The virus that causes COVID-19 is called SARS-CoV-2; it is different from the viruses that cause SARS and MERS.

### Coronavirus (COVID-19) and Asthma

COVID-19 in a person with asthma may worsen breathing symptoms or trigger an exacerbation, although risk varies with asthma control and severity. Usual prescribed controller treatment should not be stopped without medical advice.

### Coronavirus (COVID-19) and High Blood Pressure

COVID-19 in a person with hypertension may carry higher risk when other cardiovascular or age-related risk factors are present. Prescribed blood-pressure medicines should generally be continued unless the treating clinician advises otherwise.

### Coronavirus (COVID-19) and Pregnancy

SARS-CoV-2 infection during pregnancy can increase the risk of severe maternal illness, hospitalization, preterm birth, and other complications. Vaccination, prompt assessment, and individualized treatment are important.

### Coronavirus Disease 2019

Alternate index label; see *COVID-19*.

### Coronavirus Disease COVID-19

Alternate index label; see *COVID-19*.

### Coronaviruses

Coronaviruses are a family of viruses that infect humans and animals. Human disease ranges from common-cold syndromes to severe illnesses caused by SARS-CoV, MERS-CoV, and SARS-CoV-2.

### Corporeal

Relating to the physical body rather than thoughts, feelings, or the spirit. In medical writing, the word may describe a bodily symptom, physical examination, or treatment directed at the body.

---

## Corpus

Latin for body; in anatomy it refers to the main body of an organ or structure, such as the corpus callosum or corpus uteri.

## Corpus Albicans

A dense, white fibrous scar tissue mass within the ovary formed by the regression and involution of an unfertilized corpus luteum, consisting of hyalinized collagen and degenerated luteal cells.

### Alternate Names:

White Body

## Corpus Callosum

A broad band of nerve fibers connecting the brain's left and right halves. It allows the two sides to share information needed for movement, sensation, language, vision, and thinking.

## Corpus Cavernosum

One of two columns of erectile tissue in the penis that fill with blood during erection.

### Alternate Names:

Cavernous Body; Penile Corpus Cavernosum

## Corpus Luteum

The temporary endocrine structure formed from an ovarian follicle after ovulation; it produces progesterone during the luteal phase and early pregnancy.

### Alternate Names:

Yellow Body; Luteal Body

## Corpus Luteum Cyst

A usually harmless ovarian cyst that forms after ovulation when the normal hormone-producing structure fills with fluid or blood. It often disappears within a few menstrual cycles but may cause pelvic pain, delayed menstruation, or bleeding if it ruptures.

## Corpus Spongiosum

A column of erectile tissue surrounding the penile urethra and expanding distally to form the glans.

### Alternate Names:

Spongy Body; Corpus Spongiosum Penis

## Corrected Age

The developmental age of a premature baby after allowing for the weeks born before 40 weeks of pregnancy. It helps interpret growth and milestones, usually during the first two years.

## Corrigan Pulse

A strong pulse that rises quickly and then collapses rapidly. It is classically associated with aortic-valve leakage but can occur with other high-flow states and requires clinical evaluation.

## Corrosive

A chemical that destroys tissue on contact, such as a strong acid or alkali. Exposure can burn skin, eyes, airways, or the digestive tract; prompt prolonged rinsing and urgent medical advice may be needed.

## Corrosive Ingestion

Swallowing an acid, alkali, or other substance that can burn or destroy the mouth, throat, esophagus, or stomach. It may cause pain, drooling, vomiting, difficulty swallowing, or breathing problems and can lead to perforation.

## Corrugator Muscle

A small facial muscle that pulls the eyebrows downward and inward, helping create frowning and other concentrated facial expressions.

### Alternate Names:

Corrugator Supercilii; Glabellar Muscle

## Cortex

The outer layer of an organ or structure. Examples include the brain's cerebral cortex, the outer part of the adrenal gland, and the outer layer of a hair.

### Alternate Names:

Cortical Layer; Outer Cortex

## Corti's Ganglion

A group of nerve-cell bodies inside the cochlea that carries hearing signals from the inner ear toward the brain. Damage to this pathway can contribute to sensorineural hearing loss.

---

## Cortical

Relating to the cortex, the outer layer of an organ such as the brain, kidney, or adrenal gland.

## Cortical Bone

Dense bone forming the outer shell of most bones and providing much of the skeleton's mechanical strength.

### Alternate Names:

Compact Bone; Cortical Tissue

## Cortical Cataract

A cataract that begins in the outer part of the eye's lens and forms spoke-like cloudy areas extending inward. It can cause glare, halos, and blurred vision, especially in bright light.

## Corticobasal Degeneration

A rare, progressive brain disease that usually causes worsening movement and thinking problems, often more on one side. Features may include stiffness, slow movement, poor coordination, muscle spasms, speech trouble, or difficulty using a limb.

## Corticospinal Tract

The primary descending somatic motor pathway originating from pyramidal neurons in layer V of the cerebral motor cortex, descending through the internal capsule, brainstem, and medullary pyramids (where 85 to 90 percent decussate) to synapse on spinal lower motor neurons for voluntary fine motor control.

### Alternate Names:

Pyramidal Tract

## Corticosteroid

A steroid hormone from the adrenal cortex or a synthetic analogue; glucocorticoids reduce inflammation and immune activity, while mineralocorticoids regulate salt and water balance.

## Corticosteroids

Medicines similar to hormones made by the adrenal glands that reduce inflammation and suppress immune activity. They can help many conditions

but may cause infection, high blood sugar, mood changes, bone loss, or adrenal suppression.

## Corticosteroids Mechanism

Corticosteroids enter cells and switch genes that control inflammation and immune activity. They reduce the release of inflammatory chemicals and limit the movement and activation of several immune cells, easing swelling, pain, and tissue damage.

## Corticosteroids Overdose

Overdose of corticosteroids that may cause gastrointestinal symptoms, mood changes, hyperglycemia, fluid retention, infection risk, or adrenal suppression.

## Corticotropin-Releasing Factor

A hypothalamic hormone that stimulates the pituitary gland to release adrenocorticotrophic hormone and helps coordinate the stress response.

### Alternate Names:

Corticotropin-Releasing Hormone; CRH

## Cortisol

A hormone made by the adrenal glands that helps the body respond to stress, maintain blood pressure and blood sugar, regulate metabolism, and control inflammation. Too much or too little can cause serious illness.

## Cortisol Blood Test

A blood test measuring cortisol, a hormone produced by the adrenal cortex. Results help evaluate adrenal insufficiency or cortisol excess, but vary substantially with time of day, stress, illness, medicines, and the assay used.

## Cortisol Urine Test

A test measuring cortisol excreted in a timed urine collection, commonly 24-hour urine free cortisol, used to evaluate suspected excess cortisol production such as Cushing syndrome.

## Cortisone

A corticosteroid that is converted in the body to

---

cortisol and can be used to reduce inflammation or replace glucocorticoid activity.

### **Coryza**

Acute inflammation of the nasal mucous membranes with nasal discharge and obstruction, commonly occurring in the common cold or other upper-respiratory infection.

### **Cosmeceutical**

A cosmetic product marketed as having biologically active, drug-like benefits; the term has no uniform formal regulatory category in many jurisdictions.

### **Cosmetic Allergies**

Cosmetic products can cause irritant dermatitis or allergic contact dermatitis, producing itching, redness, swelling, scaling, or blistering where applied. Stopping the suspected product and medical assessment, sometimes with patch testing, help identify the responsible ingredient.

### **Cosmetic Dermatology**

Dermatologic care intended to change appearance rather than treat disease, including procedures for wrinkles, scars, pigmentation, hair, or facial volume. Risks include infection, scarring, pigment changes, allergic reactions, and injury from improperly performed treatment.

### **Cosmetic Ear Surgery**

Surgery to change the size, shape, or position of the outer ear, often to correct prominent ears or a structural difference. Risks include bleeding, infection, scarring, altered sensation, and an unsatisfactory result.

### **Cosmetic Surgery**

Surgery intended primarily to change or improve appearance rather than treat disease or restore function. It still carries risks such as infection, bleeding, anesthesia complications, scarring, and unsatisfactory results.

### **Costal Cartilage**

Flexible cartilage that connects the front ends of the ribs to the sternum or to the cartilage of the rib above. It gives the chest wall flexibility during breathing.

### **Costal Groove**

A shallow groove along the lower inner border of a rib that contains an intercostal nerve and blood vessels.

### **Costal Margin**

The lower curved edge of the rib cage, formed mainly by the cartilage of the lower ribs.

### **Costello Syndrome**

A rare genetic condition, usually caused by an HRAS gene change, affecting growth, development, skin, heart, and muscles. It may cause feeding difficulty, short stature, distinctive facial features, heart disease, and increased risk of certain tumors.

### **Costochondritis**

Inflammation of the cartilage where one or more ribs join the breastbone (sternum), causing localized chest-wall pain and tenderness that may worsen with movement, deep breathing, coughing, or pressing the area. Because similar pain can result from heart or lung emergencies, new or severe chest pain needs prompt assessment.

#### **Alternate Names:**

Costosternal Inflammation; Chest-Wall Pain Costosternal Chondrodynia Costosternal Syndrome

### **Costochondritis And Tietze Syndrome**

Costochondritis is inflammation or irritation at the junction of ribs and the breastbone, causing reproducible chest-wall pain. Tietze syndrome is a less common related condition with visible swelling at one or a few joints; heart and lung emergencies must still be excluded.

#### **Alternate Names:**

Costosternal Syndrome; Anterior Chest Wall Syndrome



---

### **Costosternal Chondrodynia**

Alternate index label; see *Costochondritis*.

### **Costosternal Syndrome**

Alternate index label; see *Costochondritis*.

### **Costotransverse Joint**

A joint where the tubercle of a rib meets the transverse process of a thoracic vertebra. It helps guide rib movement during breathing.

### **Costovertebral Joint**

A joint where a rib meets a vertebra in the spine. It helps the ribs move during breathing.

### **Costovertebral-Angle Tenderness**

Pain felt when a doctor gently thumps the lower back just below the ribs, which points to a kidney infection.

### **Cot Death**

Sudden infant death syndrome, the sudden unexplained death of an apparently healthy infant, usually during sleep, that remains unexplained after investigation.

### **Cotard Syndrome**

A rare syndrome involving a fixed belief that one is dead, missing body parts, or no longer exists, sometimes accompanied by severe depression, psychosis, or other psychiatric or neurologic illness. It is a descriptive syndrome rather than a standalone diagnosis.

### **Cotton-Wool Spots**

Small, fluffy, whitish-yellow retinal lesions representing localized micro-infarcts of the retinal nerve fiber layer caused by occlusion of precapillary arterioles and subsequent axoplasmic stasis. They are commonly observed on fundoscopic examination in diabetic retinopathy, severe hypertension, and retinal vein occlusion.

#### **Alternate Names:**

Cytoid Bodies; Retinal Nerve Fiber Layer Infarcts

### **Cough**

A protective reflex that clears mucus, irritants, or foreign material from the airways. Coughs may be short-lived or persistent and can result from infection, allergy, asthma, reflux, medicines, heart disease, or other conditions.

### **Cough Augmentation**

Techniques that help a person with weak breathing muscles clear mucus. They may include assisted coughing, taking several stacked breaths, or using a device that pushes air in and then rapidly pulls it out.

### **Cough Headaches**

Head pain triggered by coughing, sneezing, straining, laughing, or another sudden pressure increase. New, severe, or persistent attacks need assessment because some are caused by a structural brain problem.

### **Cough In Adults**

A cough in an adult may be acute, subacute, or chronic and may result from infection, asthma, chronic obstructive pulmonary disease, reflux, upper-airway cough syndrome, medication, or other disease. Coughing blood, breathing difficulty, chest pain, low oxygen, or persistent unexplained cough requires medical assessment.

### **Cough Suppressant**

An antitussive medicine or measure that reduces the urge or reflex to cough; treatment should address the underlying cause and avoid suppressing a productive cough when clearance is needed.

### **Cough Syncope**

A brief faint caused by coughing, which raises chest pressure, reduces venous return and cardiac output, and transiently lowers cerebral blood flow.

### **Coughing Up Blood**

Expectoration of blood or blood-streaked sputum from the lower respiratory tract, called hemoptysis; heavy bleeding or breathing difficulty is an emergency.

---

## Councilman Body

A rounded, intensely eosinophilic apoptotic hepatocyte globule with condensed pyknotic or fragmented nuclear material seen on microscopic examination of liver biopsies, characteristic of acute viral hepatitis, yellow fever, and severe toxic hepatitis.

### Alternate Names:

Councilman-Rocha Lima Body; Apoptotic Hepatocyte

## Count Rate

The number of detected events each second during measurement. In medical imaging, very high rates can cause missed events because equipment needs recovery time, reducing accuracy unless corrected.

## Countercurrent Multiplication

A kidney process that creates a concentrated salt gradient deep in the kidney. This gradient allows the collecting ducts to reclaim more water when antidiuretic hormone is present.

## Counterfeit Drug

A product falsely labeled as an approved medicine or made to imitate one. It may contain the wrong ingredient, too little or too much active drug, contaminants, or no active drug and can cause treatment failure or poisoning.

## Counterpulsation

Timed inflation and deflation of a balloon or pressure cuffs with each heartbeat to improve blood flow to the heart or reduce the heart's workload. It is used only in selected serious cardiac conditions.

## Countertransference

A healthcare professional's emotional responses to a patient, influenced by the professional's experiences or by the interaction. Recognizing these responses helps maintain safe, objective, and effective care.

## Counting Carbohydrates

Estimating the grams of carbohydrate in foods and drinks to help plan meals, match insulin, and manage blood glucose, especially in diabetes.

## Coup-Contrecoup Injury

Brain injury in which the brain is damaged both beneath the point of impact and on the opposite side as it moves inside the skull. Falls and vehicle crashes are common causes.

## Courvoisier Sign

A painless, enlarged gallbladder in a person with jaundice, traditionally suggesting blockage of the bile duct by a tumor rather than a gallstone. It is a clue, not a diagnosis.

## Cover Test

An objective ocular alignment test used to identify and measure manifest strabismus (tropia). While the patient fixes their gaze on a target, one eye is covered and the uncovered eye is observed for any compensatory refixation movement.

## Covered Vascular Stent

A vascular stent lined with a graft material that supports the vessel and excludes a leak or aneurysm from the bloodstream.

### Alternate Names:

Stent Graft; Endovascular Graft; Endoprosthesis

## COVID Toe

A chilblain-like skin lesion reported in association with COVID-19, usually affecting the toes or fingers and appearing red, purple, or swollen.

## COVID-19

Coronavirus disease caused by infection with SARS-CoV-2. Illness ranges from asymptomatic or mild disease to pneumonia, respiratory failure, thrombosis, or death; common symptoms include fever, cough, shortness of breath, fatigue, muscle aches, and altered taste or smell. Some people develop persistent symptoms after the acute infection.

## COVID-19 Antibody Test

A blood test that detects antibodies produced in response to SARS-CoV-2 infection or vaccination; it does not reliably diagnose current infection or establish complete immunity.

---

## COVID-19 Complications

Complications of SARS-CoV-2 infection can include pneumonia, hypoxemic respiratory failure, blood clots, myocarditis, acute kidney injury, neurologic disease, multisystem inflammatory syndrome, and persistent post-COVID symptoms.

## COVID-19 in Infants and Children

COVID-19 in children and infants is often mild but can cause pneumonia, dehydration, croup-like illness, or multisystem inflammatory syndrome in children. Breathing difficulty, poor feeding, unusual sleepiness, or dehydration requires urgent assessment.

## COVID-19 In Pregnancy

COVID-19 during pregnancy can be more severe than in nonpregnant adults and increases risks such as hospitalization, preterm birth, and stillbirth. Fetal infection before birth is uncommon but possible; vaccination and prompt care reduce risk.

## COVID-19 Recovery

Recovery after COVID-19 varies from days to months. Persistent fatigue, breathlessness, cognitive symptoms, loss of smell or taste, and post-exertional worsening may represent post-COVID condition and warrant appropriate evaluation.

## COVID-19 Symptoms

Symptoms of SARS-CoV-2 infection, which may include fever, cough, sore throat, congestion, fatigue, muscle aches, headache, loss of smell or taste, vomiting, or diarrhea.

## COVID-19 Test

A test for current SARS-CoV-2 infection using a respiratory sample. Molecular tests detect viral genetic material and antigen tests detect viral proteins; results depend on timing, sample quality, and the amount of virus present.

## COVID-19 Testing

Testing for SARS-CoV-2 uses antigen or nucleic-acid methods on respiratory samples. Test timing, symptoms, exposure, and the purpose of testing

affect interpretation and the need for repeat testing.

## COVID-19 Vaccines

Vaccines that prime immune responses against SARS-CoV-2 and reduce severe disease, hospitalization, and death; products, schedules, effectiveness, and recommendations vary by age and risk.

## COVID-19 Virus Test

A test for current SARS-CoV-2 infection, usually using a nucleic acid amplification test or antigen test on a respiratory specimen; results must be interpreted with symptoms, timing, and test performance.

## COVID-19-Associated Anosmia and Ageusia

COVID-19 can temporarily impair smell and taste through infection or inflammation of nasal and olfactory pathways. Most people improve, but persistent loss, distortion, poor nutrition, or neurologic symptoms warrants follow-up and smell-retraining advice.

## Cow's Milk Allergy

Alternate name for cow's milk protein allergy; see *Milk Allergy*.

## Cow's Milk And Children

Nutritional guidance on cow's milk for children, balancing protein, calcium, vitamin D, fat, iron, allergy, and total intake so milk does not displace a varied diet.

## Cow's Milk Introduction in Infancy

Pediatric nutritional guidance establishing that unmodified whole cow's milk should be withheld as a primary beverage until 12 months of age due to its high renal solute load, low iron bioavailability, risk of occult gastrointestinal blood loss, and inadequate micronutrient profile.

## Alternate Names:

Cow's Milk in Infancy; Infant Milk Guidelines; Whole Milk Introduction

---

## Cowden Disease (Multiple Hamartoma Syndrome)

Alternate names; see *Cowden Syndrome*.

## Cowden Syndrome

An inherited PTEN-hamartoma tumour syndrome, usually caused by a pathogenic *PTEN* variant. It may cause multiple hamartomas, characteristic mucocutaneous lesions, and increased risks of breast, thyroid, endometrial, kidney, and colorectal cancers; lifelong organ-specific surveillance is important.

## Cowper's Gland

One of a pair of small male accessory sex glands below the prostate. Its secretion contributes mucus and fluid to the urethra before ejaculation.

## Cowpox

A zoonotic viral infection caused by cowpox virus or related orthopoxviruses, producing localized pustular or ulcerative skin lesions and sometimes systemic illness.

## Cox-2

Cyclooxygenase-2, an inducible enzyme that produces inflammatory prostaglandins; selective COX-2 inhibitors reduce pain and inflammation but may have cardiovascular, renal, and gastrointestinal risks.

## COX-2 Inhibitors

Medicines that reduce the action of an enzyme involved in pain and inflammation. They may cause less stomach irritation than some older anti-inflammatory medicines, but can still cause bleeding, kidney problems, high blood pressure, or heart risks.

### Alternate Names:

Cyclooxygenase-2 Inhibitors; Selective COX-2 Inhibitors

## Coxa Vara

A hip deformity in which the upper thigh bone has an unusually narrow angle, making the leg shorter or changing how the hip bears weight. It may cause a limp, pain, or limited movement.

## Coxalgia

Pain arising from the hip or nearby tissues, often felt in the groin and sometimes spreading to the thigh or knee. Causes vary with age and include infection, injury, growth problems, arthritis, cartilage damage, stress fractures, and reduced blood supply.

## Coxiella Burnetii

A bacterium that causes Q fever, usually spread by breathing dust contaminated by infected farm animals. It can cause fever, headache, hepatitis, or pneumonia and may persist as a serious infection of heart valves.

## Coxsackie

Alternate name; see *Coxsackievirus*.

## Coxsackie Virus Type A16

A virus that commonly causes hand, foot, and mouth disease, especially in children. Fever, painful mouth sores, and a rash or blisters on the hands and feet usually resolve without serious complications.

## Coxsackievirus

A group of viruses that can cause fever, sore throat, mouth sores, rashes, or hand, foot, and mouth disease. Some types can also inflame the heart, its surrounding sac, the muscles, or the coverings of the brain.

## Cozen Test

A physical examination manoeuvre used to assess for lateral epicondylitis (tennis elbow). The patient extends and radially deviates the wrist against examiner resistance while the elbow is stabilized in flexion; sharp pain at the lateral epicondyle constitutes a positive finding.

### Alternate Names:

Cozen's Test; Tennis Elbow Test

## CP

A commonly used abbreviation for cerebral palsy, a group of permanent disorders of movement and posture caused by injury or abnormal development of the developing brain. Associated problems may in-

---

volve sensation, communication, cognition, seizures, or feeding.

## **CPAP**

Continuous positive airway pressure delivered through a mask to splint open the upper airway during sleep. It is commonly used for obstructive sleep apnea and can reduce apneas and related symptoms when used consistently.

## **CPAP Machine**

A device that sends a steady stream of air through a mask to keep the upper airway open during sleep. It is commonly used for obstructive sleep apnea; benefit depends on correct mask fit and regular use.

## **CpG**

A DNA sequence in which cytosine is followed by guanine on the same strand, often occurring in regions whose methylation helps regulate gene expression.

## **CPK Isoenzymes Test**

A blood test that separates forms of creatine kinase to help identify whether an increased level comes mainly from skeletal muscle, heart muscle, or brain. Troponin testing is usually preferred when a heart attack is suspected.

## **CPR**

Alternate index label; see *Cardiopulmonary Resuscitation*.

## **CPVT**

Alternate name; see *Catecholaminergic Polymorphic Ventricular Tachycardia*.

## **Crabs**

Alternate colloquial name; see *Pubic Lice*.

## **Crabs (Pubic Lice)**

Infestation of coarse body hair by pubic lice, causing itching and visible nits or lice. It is usually sexually transmitted but can spread through close contact; sexual partners and possible coexisting infections

should be assessed.

## **Cracked Heels**

Dry, thickened skin at the heel that develops fissures, sometimes deep enough to bleed or become infected. Moisturizers, gentle filing, and treatment of contributing skin disease can help; painful or infected cracks need assessment.

## **Cracked Tooth Syndrome**

A partial crack in a tooth that may cause sharp pain when biting or when pressure is released, and sensitivity to hot or cold. The crack can be difficult to locate and may worsen over time. Dental treatment depends on its depth and whether the inner pulp is involved.

## **Crackles**

Brief popping or bubbling sounds heard in the lungs, usually while breathing in. They may occur when small airways reopen or when air moves through fluid, as in pneumonia, heart failure, or scarring.

## **Alternate Names:**

Rales

## **Cradle Cap**

A common form of seborrheic dermatitis in infants, causing greasy or flaky scales on the scalp with mild redness. It is usually harmless and improves with gentle scalp care; persistent inflammation or signs of infection require assessment.

## **Cramp**

A sudden, involuntary, often painful muscle contraction. Cramps may result from exertion, dehydration, electrolyte changes, pregnancy, medicines, nerve disease, or reduced blood flow.

## **Cramping Leg Pain**

Pain caused by involuntary, often painful muscle contraction in the leg; contributors include exertion, dehydration, electrolyte abnormalities, medications, nerve disease, and impaired circulation.

---

### **Cramps But No Period**

Pelvic cramps without menstruation can result from ovulation, pregnancy, implantation, constipation, urinary disease, infection, endometriosis, cysts, or other pelvic conditions. Severe one-sided pain, faintness, fever, vomiting, or possible pregnancy requires urgent assessment.

### **Cranberries**

Red fruits containing fiber and plant compounds. Cranberry products may modestly reduce recurrent urinary infections in some people but do not treat an established infection; they can interact with warfarin and other medicines.

### **Cranial**

Relating to the skull or toward the head, as opposed to caudal or toward the lower end of the body.

### **Cranial Arteritis**

Inflammation of arteries, usually in the scalp and head, that can cause a new headache, scalp tenderness, jaw pain while chewing, fever, or vision loss. Sudden visual symptoms require emergency treatment.

### **Cranial Cavity**

The hollow space inside the skull that contains and protects the brain, its coverings, blood vessels, and cerebrospinal fluid. A rise in pressure inside this space can compress the brain and become life-threatening.

### **Cranial Fasciitis**

A usually harmless but rapidly growing lump in the scalp or nearby soft tissue, most often in infants and young children. It can look like cancer on examination or scans, so a tissue sample may be needed to confirm the diagnosis.

### **Cranial Implant**

An artificial material or device placed in or on the skull to repair a defect, protect the brain, drain fluid, stimulate nerves, or support treatment. Risks include infection, bleeding, and implant failure.

### **Cranial Mononeuropathy III**

Damage to the third cranial nerve, which controls several eye muscles, the eyelid, and the pupil. It can cause double vision, a drooping eyelid, abnormal eye position, or an enlarged pupil and needs prompt evaluation.

### **Cranial Nerve**

One of 12 pairs of nerves arising from the brain or brainstem. They control smell, vision, eye movement, facial sensation and movement, hearing, balance, swallowing, voice, and other head and neck functions.

### **Cranial Nerve Palsy**

Weakness or loss of function in one or more nerves arising from the brain. It may cause double vision, facial weakness, hearing loss, trouble swallowing, or tongue weakness, depending on the nerve involved.

### **Cranial Nerve X**

The vagus nerve, the tenth cranial nerve, which carries parasympathetic and sensory fibers and contributes to swallowing, voice, and regulation of thoracic and abdominal organs.

### **Cranial Nerves**

The 12 pairs of nerves arising from the brain or brainstem that provide sensory, motor, and autonomic functions to the head, neck, and parts of the trunk.

### **Cranial Sutures**

Fibrous joints between skull bones that permit molding in infancy and accommodate brain growth before progressively fusing; premature fusion causes craniosynostosis.

### **Craniectomy**

A life-saving neurosurgical procedure in which a large section of the skull bone (bone flap) is surgically removed and deliberately left off to create space for a severely swollen brain to expand. Frequently performed as an emergency decompressive craniectomy following massive traumatic brain injury, large hemispheric ischemic stroke (malignant



---

middle cerebral artery infarction), or severe intracerebral hemorrhage, the procedure prevents dangerous surges in intracranial pressure that would otherwise compress brainstem centers and cause fatal brain herniation. The removed bone is either frozen in a tissue bank or stored subcutaneously in the patient's abdominal wall until brain swelling resolves, after which the skull is reconstructed in a subsequent cranioplasty.

**Alternate Names:**

Decompressive Craniectomy; Hemicraniectomy; Suboccipital Craniectomy

### **Craniocerebral Injury**

An injury involving the scalp, skull, brain, or several of these structures, ranging from minor wounds to life-threatening traumatic brain injury.

### **Craniofacial**

Relating to the skull and face. Craniofacial conditions may involve bones, soft tissues, teeth, eyes, ears, or airway and may be congenital, traumatic, developmental, or acquired.

### **Craniofacial Abnormalities**

Abnormalities of the skull, face, or related tissues that may be present at birth or develop later. Causes include genetic conditions, disrupted development, trauma, infection, and tumors.

### **Craniofacial Disorder**

A broad category of congenital anomalies, genetic syndromes, or acquired conditions that affect the structural development, growth, and morphology of the skull, facial bones, jaws, and associated soft tissues. Major congenital forms arise from abnormal cranial neural crest cell migration or pharyngeal arch morphogenesis and include craniosynostosis (premature cranial suture fusion, as in Crouzon, Apert, and Pfeiffer syndromes), orofacial clefts (cleft lip and palate), branchial arch malformations (Treacher Collins syndrome, craniofacial microsomia, Goldenhar syndrome), and sequence disorders (Pierre Robin sequence). Clinical consequences frequently involve upper airway obstruction, feeding difficulty, elevated intracranial pressure, vision and hearing

impairment, dental malocclusion, and speech articulation defects. Optimal management requires coordinated multidisciplinary care combining specialized reconstructive surgery, neurosurgery, orthodontics, audiology, and speech-language therapy.

**Alternate Names:**

Craniofacial Anomaly; Craniofacial Malformation; Craniofacial Condition

### **Craniofacial Dysostosis**

A birth disorder in which some skull bones join too early and the face develops unusually. It can change head shape, affect the eyes or airway, and sometimes increase pressure around the brain; severity varies.

### **Craniofacial Microsomia**

A birth condition in which one side of the face develops less fully than the other. It may affect the jaw, ear, eye, facial muscles, and nearby tissues, with severity ranging from mild to complex.

### **Craniopharyngioma**

A usually slow-growing tumor near the pituitary gland and hypothalamus. It may cause headache, vision loss, excessive thirst or urination, hormone problems, growth changes, or increased pressure in the skull.

### **Cranioplasty**

Surgery to repair a missing or damaged part of the skull, often after a head injury or earlier brain surgery. A stored piece of bone or an artificial material may be used. It protects the brain and may improve skull shape or symptoms, but results and risks vary.

**Alternate Names:**

Skull Reconstruction; Cranial Reconstruction; Skull Defect Repair

### **Craniosynostosis**

Craniosynostosis is premature fusion of one or more skull sutures, altering head shape and sometimes increasing pressure on the developing brain.

**Alternate Names:**

Craniostenosis

---

## Craniosynostosis Repair

Surgery to separate prematurely fused skull bones and reshape the skull when needed. It aims to improve head shape and make room for brain growth; timing and technique depend on the child's age and anatomy.

## Craniotabes

Soft areas of an infant's skull that bend or indent more easily than expected. They can be part of normal newborn molding but may also result from rickets or another bone disorder and should be interpreted with the baby's examination and growth.

## Craniotomy

An operation in which a surgeon temporarily removes a piece of the skull to reach the brain or its coverings. It may be used to remove a tumor, stop bleeding, repair an aneurysm, or treat another brain problem. The bone is usually put back and secured at the end of the operation.

### Alternate Names:

Open Brain Surgery; Bone Flap Surgery; Craniotomy Bone Flap

## Cranium

The part of the skull that encloses and protects the brain, consisting of the cranial bones and their joints.

## Crapulence

An unpleasant state of sickness or discomfort caused by excessive eating or drinking, especially excessive alcohol use.

## Craving

A strong, often difficult-to-control desire for a substance, behavior, or experience, commonly seen in substance-use disorders.

### Alternate Names:

Strong Urge; Compulsive Desire

## Crazy Paving Pattern

A lung-imaging pattern of hazy areas crossed by thick lines, resembling paving stones. It can occur

with several conditions, including infection, bleeding, inflammation, or protein buildup, so it is not a diagnosis by itself.

## CRC

Colorectal cancer, malignant disease arising in the colon or rectum; screening and evaluation depend on age, risk, symptoms, and test results.

## CRE

Enterobacterales bacteria resistant to carbapenem antibiotics, important medicines for serious infections. They may colonize without illness or cause difficult-to-treat urinary, wound, abdominal, lung, or bloodstream infections.

## CRE Infection

Carbapenem-resistant Enterobacterales are bacteria resistant to important antibiotics and can cause urinary, bloodstream, abdominal, wound, or lung infections. Infection-control precautions, culture-based treatment, and specialist advice are important because therapeutic options may be limited.

### Alternate Names:

Carbapenem-Resistant Enterobacteriaceae Infection; CRE Colonization

## Cream

A spreadable semisolid preparation applied to the skin, often containing a medicine. Creams are usually less greasy than ointments and may be useful on moist or broad areas, but the correct product depends on the skin condition.

## Creatine

A nitrogen-containing compound stored mainly in skeletal muscle that helps supply energy for short, high-intensity activity.

## Creatine Kinase

An enzyme found mainly in muscle and released into blood when muscle cells are damaged. Levels may rise after strenuous exercise, injury, seizures, muscle disease, or some medicines; other tests help identify the source.

---

### **Creatine Phosphokinase Test**

A blood test measuring creatine kinase, an enzyme released with muscle injury; elevated levels may occur in skeletal-muscle disease, strenuous exercise, rhabdomyolysis, or myocardial injury.

### **Creating A Family Health History**

Collecting a record of relatives' diseases, ages at diagnosis, ancestry, and causes of death to identify inherited risk and guide screening, prevention, and genetic counseling.

### **Creatinine**

A breakdown product of muscle creatine phosphate produced at a constant rate, filtered by the glomerulus and minimally reabsorbed; serum creatinine is the most used marker of kidney function.

### **Creatinine Blood Test**

A blood test measuring serum creatinine, a waste product generated largely by skeletal muscle. It is used with age, body size, and other factors to estimate kidney filtration, but interpretation is affected by muscle mass, diet, medicines, and acute illness.

### **Creatinine Clearance Test**

A test estimating kidney filtration from creatinine in a timed urine collection and blood sample. It can be affected by incomplete urine collection and creatinine secretion, so estimated glomerular filtration rate is often preferred for routine assessment.

### **Creatinine Test**

A blood or urine test measuring creatinine to help assess kidney filtration and muscle-related creatinine production.

#### **Alternate Names:**

Creatinine Measurement; Renal Function Creatinine Test

### **Creatinine Urine Test**

A urine test measuring creatinine concentration or excretion, used to assess kidney handling, interpret other urine measurements, or judge the completeness of a timed collection. Results depend on muscle

mass, diet, collection quality, and kidney function.

### **Creeping Eruption**

An intensely itchy winding skin track caused by larvae, usually hookworms from contaminated soil or sand, migrating within the superficial skin; it is also called cutaneous larva migrans.

### **Cremasteric Reflex**

A superficial neurological reflex elicited by lightly stroking or scratching the superior and medial aspect of the inner thigh, causing prompt contraction of the cremaster muscle and upward retraction of the ipsilateral testis, mediated through spinal cord segments L1 and L2 via the genitofemoral and ilioinguinal nerves.

#### **Alternate Names:**

Cremaster Reflex; Genitofemoral Reflex

### **Crepitation**

Alternate name; see *Crepitus*.

### **Crepitus**

A crackling, grating, or popping sensation or sound caused by movement of tissue, bone, joint surfaces, or air in soft tissue.

#### **Alternate Names:**

Crepitation; Crackling Sensation

### **Crescentic Glomerulonephritis**

A rapidly worsening inflammation of the kidney's filtering units. It can cause blood or protein in urine, swelling, high blood pressure, and kidney failure and usually requires urgent specialist treatment.

### **CREST Syndrome**

A limited cutaneous form of systemic sclerosis named for Calcinosis, Raynaud phenomenon, Esophageal dysfunction, Sclerodactyly, and Telangiectasia. It may also affect the lungs, heart, kidneys, or digestive tract; monitoring for pulmonary hypertension and other organ involvement is important.

### **Creutzfeld-Jakob Disease**

Alternate spelling; see *Creutzfeldt-Jakob Disease*.

---

## Creutzfeldt Jakob Disease

Alternate spelling; see *Creutzfeldt-Jakob Disease*.

## Creutzfeldt-Jakob Disease

A rare, rapidly progressive brain disease caused by misfolded prion proteins that damage and destroy nerve cells. It usually causes worsening dementia, often with poor coordination, involuntary jerking movements, vision problems, stiffness, or loss of movement. About 85% of cases are sporadic, meaning no cause is found; about 5–15% are inherited, and a small number are acquired through specific medical exposures. Variant Creutzfeldt-Jakob disease is a separate form linked to exposure to the prion that causes bovine spongiform encephalopathy. CJD is fatal.

### Alternate Names:

CJD Variant CJD VCJD

## Cri Du Chat Syndrome

A genetic condition caused by loss of part of chromosome 5. Babies may have a high-pitched cry, poor growth, low muscle tone, distinctive facial features, and developmental or intellectual difficulties.

## Cri-Du-Chat Syndrome

A genetic disorder caused by a deletion of the short arm of chromosome 5 (5p-), with the size of the deletion correlating with severity. Incidence 1 in 15,000-50,000.

## CRIA Syndrome

A rare inherited inflammatory disorder caused by a change in the RIPK1 gene. Repeated attacks of fever, swollen lymph nodes, muscle or joint pain, and inflammation often begin during childhood.

## Crib Death

Alternate name; see *Sudden Infant Death Syndrome*.

## Cribriform Carcinoma

A carcinoma with a sieve-like microscopic growth pattern. The term may describe several tumors, including adenoid cystic carcinoma or cribriform breast carcinoma, so the organ of origin and pathol-

ogy report determine its meaning.

## Cribriform Plate

The horizontal, sieve-like bony lamina of the ethmoid bone situated in the floor of the anterior cranial fossa, perforated by numerous small foramina transmitting the olfactory nerve (CN I) fascicles from the nasal olfactory mucosa into the anterior cranial cavity.

## Cricoid Cartilage

A complete ring of firm tissue at the lower part of the voice box, just above the windpipe. It supports the airway and connects with other structures involved in breathing and voice.

## Cricoid Pressure

Pressure applied to the cricoid cartilage during some emergency airway procedures. Its use is intended to reduce stomach contents entering the airway, but it can make ventilation or intubation more difficult and is clinician-dependent.

## Cricothyroid Membrane

A tough fibroelastic ligament located in the anterior neck between the inferior border of the thyroid cartilage and the superior border of the cricoid cartilage; the target anatomical landmark for emergency airway cricothyroidotomy.

### Alternate Names:

Cricothyroid Ligament; Conus Elasticus

## Cricothyroidotomy

An emergency surgical airway procedure in which an incisional opening is made through the skin and cricothyroid membrane to introduce an endotracheal or tracheostomy tube, indicated in catastrophic "cannot intubate, cannot ventilate" scenarios.

### Alternate Names:

Cricothyroidectomy; Emergency Surgical Airway

## Cricothyrotomy

An emergency operation creating an opening through the front of the neck into the airway when a person cannot be intubated or oxygenated by other

---

means. It provides temporary access for breathing and has serious risks.

### **Crigler-Najjar Syndrome**

A rare inherited disorder in which the liver cannot properly process bilirubin. It causes persistent yellowing of the skin and eyes; severe forms can damage the brain unless treated urgently.

### **Crime Scene Investigation**

The systematic documentation, collection, preservation, and analysis of physical evidence from a suspected crime scene, often supporting forensic medical and legal conclusions.

### **Crimean-Congo Hemorrhagic Fever**

A serious viral disease caused by Crimean-Congo hemorrhagic fever virus. It is usually spread by the bite of an infected tick or by contact with infected animal blood or tissues, especially during or soon after slaughter. Infected human blood or body fluids can also spread it, particularly in health-care settings. Symptoms usually begin suddenly and may include fever, headache, muscle aches, dizziness, neck or back pain, sore eyes, vomiting, diarrhea, or abdominal pain. Severe illness can cause a rash from bleeding, bleeding from body openings, liver or kidney failure, shock, and death.

### **Criminal Abortion**

Abortion performed illegally, outside permitted legal indications and procedures (e.g., unsafe methods, unqualified practitioners, or non-medical settings), carrying high maternal risk.

### **CRISPR**

A gene-editing method that uses a guide molecule and an enzyme to find and change a chosen DNA sequence. It is used in research and some treatments but can make unintended changes and requires careful monitoring.

### **CRISPR Therapy**

A treatment that uses CRISPR-based genome-editing tools to make a targeted change to DNA in a patient's cells. Cells may be edited outside the

body and returned to the patient, or edited directly inside the body. It is used or studied for selected genetic and other diseases; possible concerns include unintended genetic changes, immune reactions, and effects that require long-term monitoring.

### **CRISPR-Cas9**

A laboratory gene-editing tool that uses a guide molecule to direct an enzyme to a chosen DNA sequence. The enzyme cuts the DNA so the cell can remove, disable, or replace genetic material.

### **Crista Galli**

A vertical, crest-like bony projection of the ethmoid bone rising upward from the cribriform plate into the anterior cranial fossa, serving as the anterior intracranial attachment site for the falx cerebri.

### **Critical Care**

Specialized care for people with life-threatening illness or injury who need close monitoring and support for failing organs.

### **Critical Care Nursing**

Specialized nursing care for critically ill patients requiring close monitoring, organ support, rapid assessment, complex therapies, and coordination with multidisciplinary intensive-care teams.

### **Critical Findings**

Test or examination findings that may indicate a life-threatening problem and require immediate communication and action. Examples include a collapsed lung, brain pressure, or a tear in the aorta.

### **Critical Illness Myopathy**

A muscle weakness disorder that develops during severe illness, especially sepsis, organ failure, prolonged intensive-care treatment, or use of certain medicines. It can affect arms, legs, and breathing muscles, making movement and removal from a ventilator difficult. Rehabilitation supports recovery.

#### **Alternate Names:**

CIM; Acute Quadriplegic Myopathy; Critical Care Myopathy



---

### Critical Illness Polyneuropathy

Damage to many peripheral nerves that can develop during severe critical illness, especially sepsis or multiorgan failure. It causes weakness, reduced reflexes, and difficulty coming off a ventilator and may occur together with critical illness myopathy.

### Critical Illness-Related Corticosteroid Insufficiency

Inadequate corticosteroid activity during critical illness, caused by impaired adrenal production, altered hormone metabolism, or reduced tissue response. It may contribute to low blood pressure, shock, inflammation, or poor response to usual supportive care.

### Critical Limb Ischemia

Severely reduced blood flow to an arm or leg causing pain even at rest, a wound that will not heal, or gangrene. It threatens the limb and requires urgent vascular assessment.

### Critical Results

Test findings suggesting a potentially life-threatening problem that require prompt communication, assessment, treatment, and documentation by the responsible healthcare team.

### Critical Value

A laboratory result indicating a potentially life-threatening condition that requires prompt communication to a responsible clinician or care team. The threshold is test- and laboratory-specific and should be defined by institutional policy.

### Crohn Disease

Alternate spelling; see *Crohn's Disease*.

### Crohn Disease Cobblestoning

A bumpy appearance of the bowel lining seen during an internal examination in some people with Crohn disease. Deep ulcers surround areas of less-inflamed tissue, creating a pattern resembling cobblestones.

### Crohn Disease Pathology

Patchy, chronic, often transmural intestinal inflammation with architectural distortion, fissures, creeping fat, and possible noncaseating granulomas. The disease may involve any part of the gastrointestinal tract and commonly produces skip lesions.

### Crohn Ileitis

Crohn disease involving the ileum, the final part of the small intestine, causing chronic or relapsing inflammation that may produce abdominal pain, diarrhea, weight loss, strictures, or fistulas.

### Crohn Ileocolitis

Crohn disease affecting both the ileum and colon, causing patchy chronic inflammation that may lead to abdominal pain, diarrhea, weight loss, strictures, fistulas, or perianal disease.

### Crohn's Disease

A chronic inflammatory bowel disease caused by abnormal immune reactions in the digestive tract. It can affect any part of the tract, from the mouth to the anus, but most often affects the end of the small intestine or the beginning of the colon. Inflammation is often patchy and can extend through the full thickness of the bowel wall. Symptoms may include diarrhea, cramping or abdominal pain, weight loss, tiredness, fever, or anemia. Symptoms may flare and then improve during remission. Complications can include narrowed bowel segments, fistulas, abscesses, and inflammation of the joints, eyes, or skin.

#### Alternate Names:

Regional Enteritis; Granulomatous Enteritis; Crohn's

### Cronkhite-Canada Syndrome

A rare, non-familial gastrointestinal polyposis syndrome characterized by multiple juvenile-like hamartomatous polyps throughout the stomach, small bowel, and colon, accompanied by distinctive ectodermal changes including generalized alopecia, cutaneous hyperpigmentation, and dystrophic onycholysis.

#### Alternate Names:



---

CCS; Gastrointestinal Polyposis with Ectodermal Changes

### **Cross-Eyes (Strabismus)**

Misalignment of the eyes so they do not point in the same direction. Persistent strabismus in a child can impair visual development and cause amblyopia; sudden adult onset may indicate neurologic or ocular disease.

### **Cross-Matching**

Laboratory testing that checks whether donor blood is compatible with a recipient before transfusion. It helps prevent dangerous immune reactions by detecting antibodies that could attack transfused red cells.

### **Cross-Section**

A slice or plane made through a structure, or an image showing that slice, such as a cross-sectional CT image.

### **Cross-Sectional Study**

A study that measures possible causes and health outcomes at one point in time. It can show how common a condition is and whether factors are linked, but it usually cannot show which came first or prove cause and effect.

### **Cross-Tolerance**

Reduced response to one substance because the body has adapted to another substance with a similar effect. It can increase overdose risk when a person changes substances or combines them.

### **Crossed Extensor Reflex**

A contralateral polysynaptic spinal reflex in which noxious or painful stimulation causing flexion and withdrawal of an injured limb simultaneously triggers contraction of extensor muscles and inhibition of flexor muscles in the opposite limb to maintain upright balance and posture.

#### **Alternate Names:**

Crossed Extension Reflex; Contralateral Extensor Reflex

### **Crossmatch**

Laboratory test mixing donor and recipient blood to detect preformed antibodies that could cause transfusion or transplant reactions.

### **Crosti-Gianotti Syndrome**

A self-limited papular rash of childhood, also called papular acrodermatitis of childhood. It may follow viral infection or vaccination and causes symmetric papules on the face, buttocks, and limbs, usually resolving over weeks.

### **Crotch**

The region where the inner thighs meet the trunk, including the groin and nearby perineal structures.

### **Croup**

An acute viral infection causing inflammation and narrowing of the upper airway, typically producing a barking cough, hoarseness, and inspiratory stridor in young children.

#### **Alternate Names:**

Laryngotracheobronchitis; Acute Laryngotracheitis

### **Crouzon Syndrome**

A craniosynostosis syndrome, usually caused by a pathogenic *FGFR2* variant, in which skull sutures fuse too early. It changes the shape of the skull and face and may cause shallow eye sockets, vision problems, airway obstruction, dental abnormalities, hearing loss, or raised pressure around the brain.

### **Crow-Fukase Syndrome**

Another name for POEMS syndrome, a rare plasma-cell disorder associated with polyneuropathy, organ enlargement, endocrinopathy, monoclonal protein, and characteristic skin changes.

### **Crowe Sign**

Small freckles in the armpits or groin. They are one finding that can support neurofibromatosis type 1, but they are not enough to make the diagnosis alone.

#### **Alternate Names:**

Crowe's Sign; Axillary Freckling Sign

---

## Crown

The visible part of a tooth above the gum line, or a dental restoration that covers and protects a damaged tooth.

### Alternate Names:

Dental Crown; Anatomic Tooth Crown

## Crown Lengthening

A dental procedure that removes or reshapes gum tissue, and sometimes a small amount of bone, to expose more tooth. It may allow a damaged tooth to be restored or change a gummy smile.

## Crown-Rump Length

The ultrasound measurement from the top of an embryo's head to its bottom. It is the most accurate way to estimate pregnancy age during early pregnancy and helps monitor early growth.

## Crowning

The stage of childbirth when the baby's head remains visible at the vaginal opening between contractions because it has stretched the surrounding tissues. It occurs shortly before the head is born.

## CRP

C-reactive protein, a substance made by the liver that rises during inflammation or infection. A CRP test can help assess inflammation and response to treatment, but it does not identify the cause by itself.

## Cruciate Ligament

One of two strong bands crossing inside the knee that stabilize forward, backward, and twisting movement. The anterior cruciate ligament is commonly torn during sports or a sudden change in direction.

## Crush Injury

Damage caused when a body part is squeezed or trapped under pressure. It can cause broken bones, muscle death, nerve and vessel injury, dangerous chemical changes in blood, shock, or kidney failure.

## Crush Syndrome

A life-threatening illness after prolonged muscle compression, often when someone is trapped. Damaged muscle releases substances that can cause shock, dangerous heart rhythms, and sudden kidney failure, especially after pressure is removed.

## Crushing Injury

Damage caused when a body part is compressed between heavy objects. It can produce fractures, nerve and blood-vessel injury, tissue death, and rhabdomyolysis. Severe limb compression or sudden release after prolonged entrapment requires emergency treatment.

## Crust

Dried serum, blood, pus, or other exudate on the skin surface, often covering an erosion, ulcer, vesicle, pustule, or inflamed lesion.

## Crusted Scabies

A severe form of scabies causing thick crusts full of very large numbers of mites. It is extremely contagious and occurs mainly in people with weakened immunity, older adults, or people with reduced sensation.

## Cruveilhier-Baumgarten Sign

A continuous epigastric or periumbilical vascular venous hum and palpable thrill resulting from turbulent collaterals in recanalized umbilical or paraumbilical veins in patients with severe portal hypertension and cirrhosis.

### Alternate Names:

Cruveilhier-Baumgarten Murmur; Cruveilhier-Baumgarten Syndrome

## Cry For Help

A verbal or behavioral expression of distress or need for assistance; in a mental-health setting it should be taken seriously and assessed for immediate safety concerns.

## Crying In Childhood

Vocal expression of distress or emotion in a child;

---

persistent, inconsolable, unusual, or pain-associated crying may indicate illness, injury, abuse, or unmet developmental needs.

### **Crying In Infancy**

Crying is an infant's normal communication of hunger, discomfort, fatigue, or need for contact, but sudden inconsolable crying with lethargy, fever, vomiting, or injury needs urgent assessment.

### **Cryoablation**

A treatment that uses controlled freezing to destroy selected abnormal tissue, such as a tumor or abnormal heart pathway. Imaging or specialized equipment guides the treatment, which may cause bleeding, injury, or recurrence.

### **Cryobiopsy**

A biopsy method that uses a freezing instrument to remove a larger tissue sample, often from the lung. It can improve diagnosis in selected cases but carries risks such as bleeding and a collapsed lung.

### **Cryoglobulinemia**

A condition in which abnormal blood proteins thicken or clump when cold. They can inflame small blood vessels and cause purplish skin spots, joint pain, nerve problems, kidney injury, or poor circulation.

### **Cryoglobulinemic Vasculitis**

Inflammation and injury of small blood vessels caused by abnormal blood proteins that become active in the cold. It may cause purple spots, joint pain, nerve problems, kidney disease, or poor circulation.

### **Cryoglobulins**

Immunoglobulins that precipitate at low temperatures and dissolve on warming; circulating cryoglobulins can cause vasculitis, purpura, neuropathy, kidney disease, or hyperviscosity.

### **Cryopexy**

A procedure that uses controlled freezing to destroy

or seal abnormal tissue, especially retinal breaks or small retinal detachments.

### **Cryoprecipitate**

A blood-product concentrate containing mainly fibrinogen and some clotting proteins. It is given in selected serious bleeding disorders when a person's fibrinogen level is too low.

### **Cryopreservation**

Preserving cells, tissues, sperm, eggs, or embryos by cooling them to very low temperatures with protective chemicals. The material can later be thawed for treatment, research, or reproduction.

### **Cryopyrin-Associated Autoinflammatory Syndromes**

Alternate index label; see *Cryopyrin-Associated Periodic Syndromes*; the group is also called CAPS.

### **Cryopyrin-Associated Periodic Syndromes**

A group of inherited inflammatory disorders causing repeated fever, rash, joint symptoms, eye inflammation, or hearing loss because immune activity is overactive.

#### **Alternate Names:**

Cryopyrin-Associated    Autoinflammatory    Syndromes

### **Cryostat**

An instrument that maintains very low temperatures and cuts thin frozen sections of tissue for rapid microscopic examination.

### **Cryosurgery**

Surgery that destroys abnormal tissue by controlled freezing, used in selected skin lesions, tumors, and other conditions.

### **Cryotherapy**

Treatment using controlled cold to destroy or reduce abnormal tissue. It may treat selected skin lesions, tumors, eye problems, or prostate disease; pain, blistering, scarring, and damage to nearby tissue

---

are possible.

### **Cryotherapy For Prostate Cancer**

A treatment that destroys prostate tissue by repeated freezing and thawing, used selectively for localized or recurrent prostate cancer.

### **Cryotherapy For The Skin**

Destruction of selected skin lesions by controlled freezing, usually with liquid nitrogen, for conditions such as warts, actinic keratoses, or some superficial tumors.

### **Crypt**

A small cavity, gland, or recessed space in body tissue, such as an intestinal crypt or tonsillar crypt.

### **Cryptitis**

Inflammation of a small gland or recessed space in body tissue, most often the tiny glands and spaces lining the anal canal. It may cause pain, bleeding, or discharge.

### **Crypto**

Clinical shorthand that may refer to cryptococcosis or Cryptosporidium infection; the intended meaning depends on the clinical context.

### **Cryptococcal Infection**

Alternate name; see *Cryptococcosis*.

### **Cryptococcal Meningitis**

A fungal infection of the coverings and fluid around the brain, usually in people with weakened immunity. It often develops gradually with headache, fever, nausea, confusion, vision changes, or unusual sleepiness and requires urgent treatment.

#### **Alternate Names:**

Cryptococcic Meningitis

### **Cryptococcic Meningitis**

Meningitis caused by *Cryptococcus* fungi, especially in people with advanced HIV or other immune suppression. Headache, fever, confusion, nausea, and visual symptoms may develop gradually and require

urgent antifungal treatment.

### **Cryptococcosis**

Cryptococcosis is fungal infection caused by *Cryptococcus*, often affecting the lungs or brain in people with immune suppression.

#### **Alternate Names:**

Cryptococcal Infection; Torulosis

### **Cryptococcus Neoformans**

A yeast found in soil and bird droppings that can cause lung or brain infection after it is inhaled. Serious disease is most likely in people with weakened immunity and may require prolonged antifungal treatment.

### **Cryptogenic Organizing Pneumonia**

An inflammatory lung illness in which healing tissue blocks small airways and air sacs, without a known cause. It may cause cough, fever, and breathlessness and often improves with specialist-directed treatment.

### **Cryptorchidism**

A condition in which one or both testicles have not moved into the scrotum by birth. Some descend during the first months of life; a testicle that remains undescended or later moves back up needs medical assessment. Persistent undescended testicles increase the risk of reduced fertility, twisting, injury, and testicular cancer.

#### **Alternate Names:**

Undescended Testicle; Undescended Testis; Cryptorchism; Maldescended Testis

### **Cryptosporidiosis**

An intestinal infection caused by microscopic *Cryptosporidium* parasites. It spreads when contaminated water, food, surfaces, or material from infected people or animals is swallowed. It may cause watery diarrhea, stomach cramps, nausea, and loss of fluids. Some people have no symptoms, while illness may be prolonged or severe in people with weakened immune systems.

---

### **Cryptosporidium Enteritis**

Diarrheal intestinal infection caused by *Cryptosporidium* parasites, transmitted through contaminated water, food, or contact; illness can be prolonged or severe in immunocompromised people.

### **Crystal Arthropathy**

Joint inflammation caused by crystals collecting in or around a joint. Uric-acid crystals cause gout, while calcium-pyrophosphate crystals cause pseudogout; sudden pain, swelling, warmth, and redness are common, and joint-fluid testing can confirm the cause.

### **Crystal-Induced Synovitis**

Acute or chronic joint inflammation caused by deposition of crystals such as monosodium urate, calcium pyrophosphate, or basic calcium phosphate in or around the joint.

### **Crystalline Lens**

The clear, flexible structure inside the eye that focuses light onto the retina. It changes shape for near vision and becomes cloudy in cataracts, causing blurred or reduced vision.

### **Crystalloids**

Sterile fluids containing water and small dissolved substances such as salts or sugar. They can be given through a vein to replace lost fluid, but the choice and amount depend on the person's condition.

### **Crystalluria**

The presence of crystals in urine, which may reflect precipitation of salts, medication effects, metabolic disease, or a tendency to form urinary stones.

### **CSF Analysis**

Alternate name; see *Cerebrospinal Fluid Analysis*.

### **CSF Cell Count**

Laboratory measurement of red and white blood cells in cerebrospinal fluid, used to evaluate meningitis, encephalitis, subarachnoid hemorrhage, inflammation, malignancy, and other central nervous

system disorders.

### **CSF Coccidioides Complement Fixation Test**

A cerebrospinal-fluid test for complement-fixing antibodies to *Coccidioides*, used with other findings to evaluate suspected coccidioidal meningitis.

### **CSF Glucose Test**

A laboratory measurement of glucose in cerebrospinal fluid, usually interpreted with a simultaneous blood-glucose value. Low cerebrospinal-fluid glucose may occur in bacterial, fungal, or tuberculous meningitis and other disorders, but results require the complete clinical and cerebrospinal-fluid profile.

### **CSF Leak**

Escape of cerebrospinal fluid through a defect in the skull base or spinal dura, causing positional headache, clear nasal or ear drainage, wound leakage, meningitis risk, or intracranial hypotension.

### **CSF Myelin Basic Protein**

A test measuring a protein released when the insulating covering of nerve fibers is damaged. An increased level may support evaluation of multiple sclerosis or other nervous-system injury but is not specific by itself.

### **CSF Oligoclonal Banding**

A laboratory test looking for immune-protein bands in the fluid around the brain and spinal cord and comparing them with blood. The pattern may support multiple sclerosis or another inflammatory nervous-system disorder but is not diagnostic by itself.

### **CSF Rhinorrhea**

Leakage of the fluid surrounding the brain and spinal cord through a skull-base defect into the nose. It may cause a persistent clear, watery drip, often worse when bending forward, and can increase the risk of meningitis.

---

### CSF Smear

Microscopic examination of cerebrospinal fluid for cells, organisms, or abnormal material, supporting evaluation of meningitis, hemorrhage, malignancy, or inflammation.

### CSF Total Protein

A measurement of protein in cerebrospinal fluid, usually obtained by lumbar puncture. High or low levels may support evaluation of infection, bleeding, inflammation, nerve disorders, or other brain and spinal-cord disease.

### CSF-VDRL Test

A test of the fluid around the brain and spinal cord for antibodies associated with syphilis. A positive result strongly supports neurosyphilis, but a negative result does not completely exclude it and must be interpreted with symptoms and other tests.

### CT

Computed tomography, an imaging method that uses multiple X-ray measurements and computer reconstruction to produce cross-sectional images.

#### Alternate Names:

Computed Tomography; CAT Scan

### CT Angiography

A CT scan performed after injected contrast highlights blood vessels. It can show narrowing, blockage, aneurysm, dissection, bleeding, or abnormal vessel connections in areas such as the brain, heart, chest, abdomen, or limbs.

### CT Angiography of Abdomen and Pelvis

High-resolution contrast-enhanced computed tomography optimized for vascular visualization of the abdominal aorta, visceral branches, renal arteries, and iliac vessels to diagnose aneurysms, dissections, mesenteric ischemia, and hemorrhage.

#### Alternate Names:

CTA Abdomen and Pelvis; Abdominal CT Angiography; Mesenteric CTA

### CT Angiography of Head and Neck

Rapid contrast-enhanced computed tomographic angiography evaluating the extracranial carotid/vertebral arteries and intracranial cerebral arterial circulation in acute ischemic stroke, intracranial aneurysms, and vascular dissections.

#### Alternate Names:

CTA Head and Neck; Carotid CTA; Cerebral CT Angiography; Intracranial CTA

### CT Angiography of the Extremities

Non-invasive contrast-enhanced computed tomographic imaging of upper or lower extremity arterial vasculature to assess peripheral arterial disease, traumatic vascular disruption, arterial emboli, or bypass graft patency.

#### Alternate Names:

CTA Extremities; Peripheral CTA; Runoff CTA; Lower Extremity CTA

### CT Angiography – Chest

Computed tomography angiography of the chest uses intravenous contrast and computed tomography to visualize the thoracic aorta, pulmonary arteries, or other chest vessels. It can evaluate embolism, aneurysm, dissection, bleeding, and vascular malformation.

### CT Chest

A CT scan of the chest that shows the lungs, heart, blood vessels, mediastinum, pleura, ribs, and chest wall. It can evaluate infection, tumors, injury, fluid, blood clots, and other abnormalities.

### CT Colonography

A CT scan of the colon after bowel cleansing and gentle inflation with air or carbon dioxide. It can look for polyps or cancer when colonoscopy is incomplete or unsuitable, but an abnormal result usually requires colonoscopy for tissue removal or biopsy.

#### Alternate Names:

Virtual Colonoscopy

### CT Contrast

A substance administered during computed tomog-



---

raphy to make selected tissues, vessels, or body spaces more visible.

### **CT Coronary Angiography**

A CT scan taken after injected contrast outlines the arteries supplying the heart. It can show plaque or narrowing and is a non-invasive option for selected people with possible coronary artery disease.

### **CT Dose-Length Product**

A measure of the radiation output from a CT scan that combines the scan settings with the length of the body examined. It helps imaging teams compare and improve protocols but does not directly predict one person's cancer risk.

### **CT Enterography**

A CT scan of the small intestine after drinking contrast to expand the bowel and receiving injected contrast. It helps evaluate Crohn disease, inflammation, tumors, obstruction, and unexplained intestinal bleeding.

### **CT Perfusion**

A CT technique that estimates blood flow and related perfusion parameters in tissue, commonly to assess selected cases of acute stroke or tumors.

### **CT Pulmonary Angiography**

CT imaging performed after intravenous contrast to show the pulmonary arteries and evaluate suspected pulmonary embolism or other vascular abnormalities.

### **CT Scan**

An imaging test that uses rotating X-rays and a computer to create detailed cross-sectional pictures (slices) of the body. By taking pictures from multiple angles, it provides far more detail than a standard flat X-ray, quickly showing bones, internal bleeding, blood clots, organ injuries, and tumors. Scans can be done plain or with a contrast dye swallowed or injected into a vein to highlight blood vessels and internal organs.

#### **Alternate Names:**

Computed Tomography; CAT Scan

### **CTD-Associated Interstitial Lung Disease**

Interstitial lung disease occurring with autoimmune connective-tissue disease such as systemic sclerosis, rheumatoid arthritis, myositis, or Sjögren disease. Symptoms, lung function, high-resolution CT, antibodies, and disease activity guide monitoring and immunomodulatory or antifibrotic treatment.

### **Ctenocephalides Flea Bite**

Alternate name; see *Flea Bites in Humans*.

### **CTL**

Cytotoxic T lymphocyte, an immune cell that recognizes and kills infected, abnormal, or target cells through antigen-specific mechanisms.

### **CTLA-4**

A protein on some immune cells that acts as a brake on T-cell activity. Blocking this brake can strengthen an immune attack against cancer but may also cause inflammation in healthy organs.

### **CTPA**

Alternate name; see *Computed Tomography Pulmonary Angiography*.

### **CTS**

Commonly, carpal tunnel syndrome: compression of the median nerve at the wrist causing numbness, tingling, pain, or weakness in the hand.

### **Cubital**

Relating to the elbow or nearby forearm. Cubital structures include the joint, bones, muscles, nerves, and blood vessels, and problems there may cause pain, restricted movement, numbness, or weakness.

### **Cubital Tunnel Syndrome**

Pressure or stretching of the ulnar nerve at the elbow, causing tingling or numbness in the little and ring fingers, hand weakness, or pain. Symptoms often worsen when the elbow stays bent.

### **Cubitus**

The elbow or the forearm; the term is also used

---

in names of structures and conditions around the elbow.

### **Cubitus Valgus**

An outward angle of the forearm at the elbow that is larger than usual when the arm is straight. It may be present from birth or develop after an elbow injury. In some people, it can put pressure on the ulnar nerve and cause tingling, numbness, or weakness in the ring and little fingers.

### **Cubitus Varus**

An abnormal inward angulation of the forearm relative to the upper arm when the elbow is extended, often developing after a childhood elbow fracture.

### **Cuboid**

A cube-shaped bone on the outer side of the midfoot. It helps form the foot arch and connects several foot bones; injury or displacement can cause outer-foot pain, especially during walking.

### **Cul-De-Sac**

A space or pouch that ends without an opening on the other side. In the pelvis, it may refer to the space behind the uterus or between the bladder and rectum.

### **Culdocentesis**

A procedure that places a needle through the back of the vagina to sample or drain fluid from the lowest part of the pelvis. Ultrasound and less-invasive tests have replaced it in many situations.

### **Cullen Sign**

Bluish bruising in the skin around the belly button, which suggests bleeding inside the abdomen or severe inflammation of the pancreas.

### **Cullen's Sign**

Blue or purple discoloration around the navel caused by blood leaking into tissues inside the abdomen and tracking to the skin. It can signal serious internal bleeding, including bleeding from severe pancreatitis or pregnancy-related emergencies.

## **Cultural Psychiatry**

The study and clinical practice of how cultural factors influence mental health, illness expression, diagnosis, and treatment. Cultural concepts of distress, idioms of distress, and cultural syndromes must be considered. The DSM-5 includes a Cultural Formulation Interview.

## **Culture**

A laboratory test in which a clinical specimen is placed in conditions that allow microorganisms to grow and be identified; susceptibility testing may help guide treatment.

## **Culture And Sensitivity**

A laboratory test that grows microorganisms from a sample and checks which medicines can stop them. It helps identify the cause of an infection and select effective treatment.

## **Culture Media**

Nutrient materials prepared to grow microorganisms in a laboratory, with different formulas chosen for specific organisms, tests, or identification purposes.

## **Culture Swab**

A sterile swab used to collect material from skin, wounds, the throat, genital tract, or another site for laboratory culture. The test may identify bacteria or fungi and guide antibiotic choice; viral testing usually uses separate molecular or antigen tests.

## **Culture-Negative Endocarditis**

Infective endocarditis in which routine blood cultures do not identify an organism, often because of prior antibiotics, fastidious bacteria, fungi, or noninfectious mimics; diagnosis requires integrated clinical, imaging, and laboratory assessment.

## **Cumulative Dose**

The total amount of a substance or radiation received over time. Cumulative exposure matters for medicines, toxins, radiation, carcinogens, and pregnancy-related risks because effects may build gradually.

---

## Cumulative Trauma Disorder

Pain or injury that develops after repeated forceful, awkward, or prolonged movements. It may affect muscles, tendons, nerves, or other soft tissues in the hands, arms, neck, shoulders, or back.

## Cup-To-Disc Ratio

A comparison between the central hollow of the optic nerve and the whole visible optic nerve head. An enlarged or changing ratio may suggest glaucoma but must be interpreted with eye pressure and other findings.

## Cupping

Enlargement or deepening of the central depression in the optic disc, assessed during glaucoma evaluation and interpreted with eye pressure and other findings.

### Alternate Names:

Optic-Disc Cupping; Glaucomatous Cupping

## Cupping Therapy

A complementary practice that places cups on the skin to create suction, sometimes after small skin cuts. It may leave bruises or marks and can cause burns, infection, or blood loss; evidence of benefit varies by condition.

## CURB-65 Score

A five-point tool estimating the severity of community-acquired pneumonia using confusion, blood urea, breathing rate, blood pressure, and age 65 or older. It helps guide hospital decisions but does not replace clinical judgment.

## Curcumin

A yellow compound in turmeric that has anti-inflammatory effects in laboratory studies. Evidence that supplements treat disease is limited or variable, and concentrated products can interact with medicines or cause stomach or liver problems.

## Cure

Complete elimination or permanent resolution of a disease. The meaning of cure varies by condition;

remission or no detectable disease does not always mean that a disease has been cured.

## Curettage

Removal or scraping of tissue with a curette, a spoon-shaped or ring-shaped instrument. It may be used to treat or diagnose problems in the uterus, skin, bone, teeth, or another body area.

## Curettage Specimen

Tissue scraped or scooped from a body surface, cavity, or bone lesion for laboratory examination. Because the pieces are often small and fragmented, they may not show the full structure or depth of disease.

## Curette

A spoon-shaped or ring-shaped surgical instrument used to scrape, remove, or clean tissue from a cavity, bone, skin lesion, or tooth-supporting area. Its use and risks depend on the procedure and body site.

## Curie

A unit of radioactivity equal to 3.7 billion nuclear disintegrations per second; the SI unit is the becquerel.

## Curling Ulcer

An acute ulcer in the stomach or duodenum that can develop after severe burns. Reduced blood flow and major physical stress weaken the digestive lining and may cause dangerous bleeding.

## Curling's Ulcer

An acute peptic ulcer associated with severe burns, often occurring in the duodenum or stomach during physiologic stress.

## Currant Jelly Sputum

A distinctive thick, viscous, dark-red, blood-tinged mucoid sputum characteristic of acute destructive lobar pneumonia caused by *Klebsiella pneumoniae*, produced by extensive alveolar necrosis, hemorrhage, and abundant capsular polysaccharide secretion.

### Alternate Names:

---

Klebsiella Sputum; Red Currant Jelly Sputum

### **Curschmann Spirals**

Microscopic corkscrew-shaped casts of condensed mucus and desquamated bronchial epithelial cells found in sputum or bronchoalveolar lavage fluid, representing plugs formed in small distal bronchioles in patients with bronchial asthma and chronic bronchitis.

#### **Alternate Names:**

Curschmann's Spirals; Bronchial Mucus Casts

### **Curvature Of The Penis**

A bend of the erect or flaccid penis caused by congenital tissue asymmetry, scar tissue such as Peyronie disease, trauma, or other conditions; it may impair intercourse or cause pain.

#### **Alternate Names:**

Penile Curvature

### **Curvature of the Spine (Scoliosis)**

Abnormal sideways curvature and rotation of the spine. It may be idiopathic, congenital, neuromuscular, or degenerative and can cause asymmetry, pain, reduced function, or impaired breathing when severe.

### **Curvularia Alcornii**

A dark-pigmented environmental fungus that rarely causes infection. It may affect the skin, sinuses, or brain, mainly in people with weakened immunity, and requires specialist laboratory identification and treatment.

### **Cushing Disease**

Cushing syndrome caused by an adrenocorticotrophic hormone-secreting pituitary tumor. Excess cortisol can cause central weight gain, rounded facial features, skin changes, muscle weakness, high blood pressure, high blood glucose, and menstrual or mood changes.

### **Cushing Syndrome**

Illness caused by prolonged exposure to too much cortisol or steroid medicine. It can cause central

weight gain, easy bruising, muscle weakness, high blood pressure, diabetes, bone loss, infections, and mood or menstrual changes.

#### **Alternate Names:**

Hypercortisolism; Cortisol Excess Syndrome

### **Cushing Syndrome Complications**

Long-term excess cortisol can cause high blood pressure, diabetes, infections, blood clots, weak bones, muscle weakness, and heart or blood-vessel disease. Treating the cause and monitoring these problems can reduce complications.

### **Cushing Syndrome Due To Adrenal Tumor**

Excess cortisol caused by an adrenal tumor, producing Cushing syndrome with features such as central weight gain, hypertension, diabetes, muscle weakness, and skin changes.

### **Cushing Triad**

A classic constellation of physiological responses indicating life-threateningly elevated intracranial pressure and impending transtentorial or tonsillar brainstem herniation, consisting of systolic hypertension with widened pulse pressure, bradycardia, and irregular respirations.

#### **Alternate Names:**

Cushing's Triad; Cushing Reflex

### **Cushing Ulcer**

An acute stomach or duodenal ulcer associated with severe brain disease, head injury, or neurosurgery. Excess nerve stimulation can increase acid production and cause abdominal pain or gastrointestinal bleeding.

### **Cushing's Disease**

A pituitary tumor that makes too much adrenocorticotrophic hormone (ACTH), causing the adrenal glands to produce too much cortisol. Features may include weight gain around the trunk, a round face, easy bruising, purple stretch marks, muscle weakness, high blood pressure, high blood glucose, and menstrual or mood changes.

---

## Cushingoid

Resembling the physical features of excess glucocorticoid exposure, such as a round face, central weight gain, thin skin, easy bruising, and muscle weakness.

## Cusp

A pointed projection, such as a projection of a tooth or one of the leaflets of a heart valve.

## Cusp Fracture

A break in one of the raised points of a tooth, often around a large filling or weakened tooth. It may cause pain when biting or sensitivity and may require a bonded repair, crown, or other dental treatment.

## Cut

A break or incision made in tissue, usually by a sharp object; the depth, contamination, and location determine the required care.

## Cutaneous

Relating to the skin. The skin forms a protective barrier, helps control temperature, senses touch and pain, and supports vitamin D production. Skin findings can provide clues to infections, immune diseases, hormone disorders, and skin cancers.

## Cutaneous Abscess

A collection of pus in the skin or tissues beneath it, usually caused by bacterial infection. It may be painful, red, warm, and swollen; treatment may require drainage and sometimes antibiotics.

## Cutaneous and Oral Melanocytic Nevus

Usually benign moles composed of melanocytes in the skin or oral mucosa. Most remain stable, but a lesion that changes in size, colour, border, surface, or symptoms should be examined for atypia or melanoma.

## Cutaneous Angiosarcoma

A fast-growing cancer of blood-vessel cells in the skin, usually on the scalp or face of older adults. It may look like a bruise, ulcerate, and spread.

## Cutaneous Anthrax

A skin infection caused by *Bacillus anthracis*, usually producing a painless black sore surrounded by swelling.

## Cutaneous B-Cell Lymphoma

A group of non-Hodgkin lymphomas in which clonal B lymphocytes primarily involve the skin. Primary cutaneous follicle-centre and marginal-zone lymphomas often behave indolently, whereas diffuse large B-cell lymphoma, leg type, is more aggressive; biopsy, staging, and immunophenotyping guide treatment.

## Cutaneous Bleeding

Leakage of blood into the skin or the tissue just beneath it. It may appear as tiny red or purple spots called petechiae, larger areas called purpura, or bruises called ecchymoses. Causes include injury, low platelet levels, clotting disorders, fragile blood vessels, and medicines such as anticoagulants.

## Cutaneous Candidiasis

A *Candida* yeast infection of the skin, often affecting warm, moist folds such as the groin, armpits, or under the breasts. It causes red, itchy, moist plaques with small satellite spots; diabetes, antibiotics, occlusion, and immune suppression increase risk.

## Cutaneous Diphtheria

A skin infection caused by *Corynebacterium diphtheriae* bacteria. It usually produces a long-lasting ulcer or sore that may have a gray membrane and can spread toxin or infection to other people.

## Cutaneous Drug Reaction

A skin reaction caused by a medicine taken by mouth, given by injection, or applied to the skin. It may be allergic or nonallergic and can cause a rash, itching, hives, blisters, peeling, or pain; widespread blistering, mouth sores, fever, facial swelling, or breathing difficulty requires urgent care.

## Cutaneous Horn

A cone-shaped, hard projection of compacted ker-

---

atin arising from the skin. The underlying lesion may be benign, precancerous, or cancerous, especially when the base is broad, indurated, painful, or inflamed, so removal or biopsy is usually needed.

### **Cutaneous Langerhans Cell Histiocytosis**

An uncommon disorder in which abnormal immune cells build up in the skin. It may cause scaly or crusted spots, rash, or purple marks; testing checks whether other organs are involved.

### **Cutaneous Larva Migrans**

An intensely itchy, winding skin rash caused by hookworm larvae from contaminated soil or sand entering bare skin. The larvae move beneath the skin and usually die without reaching internal organs.

### **Cutaneous Leiomyoma**

A usually benign smooth-muscle tumour of the skin, often presenting as firm, painful papules or nodules that may be triggered by cold or pressure. Multiple lesions can be associated with hereditary leiomyomatosis and renal-cell cancer, so extensive or familial disease needs genetic and renal-risk assessment.

### **Cutaneous Leishmaniasis**

A skin infection caused by *Leishmania* parasites transmitted by sandflies, usually producing one or more chronic ulcers.

### **Cutaneous Lipomas**

Multiple benign tumors of mature fat cells arising in or beneath the skin, usually soft, mobile, and painless. A rapidly growing, painful, fixed, or deep mass needs imaging or biopsy because not every fatty-appearing lump is a simple lipoma.

### **Cutaneous Lupus**

A form of lupus erythematosus limited to the skin, manifesting as discoid lesions, subacute cutaneous plaques, or acute malar erythema. It may occur in isolation or as a feature of systemic lupus erythematosus.

### **Cutaneous Lupus Erythematosus**

An immune disorder causing inflamed skin patches or sores that may worsen with sunlight, scar, or change skin color. It may remain limited to skin or occur with systemic lupus and requires appropriate evaluation.

### **Cutaneous Lymphoma**

A cancer of immune cells that mainly affects the skin. It may cause persistent patches, plaques, lumps, or widespread redness. The exact type determines whether it behaves slowly or aggressively and how it is treated.

### **Cutaneous Manifestations of HIV**

Skin findings associated with HIV infection may reflect acute HIV, opportunistic infections, inflammatory disease, medicine reactions, or cancers such as Kaposi sarcoma. Conditions including herpes zoster, molluscum contagiosum, candidiasis, and extensive or recurrent eruptions may become more severe as immune function declines; effective antiretroviral therapy often improves HIV-related skin disease.

### **Cutaneous Melanoma**

A malignant tumor of melanocytes, the pigment-producing cells of skin. It may arise in a new lesion or a changing mole and can spread if not treated; the ABCDE changes, an evolving lesion, bleeding, or a nonhealing pigmented spot requires prompt dermatologic assessment and biopsy.

### **Cutaneous Metastases**

Secondary deposits of cancer in the skin, originating from a primary tumor elsewhere in the body. They may present as nodules, plaques, or ulcers, and often indicate advanced or recurrent disease.

### **Cutaneous Metastasis**

A deposit of malignant cells in the skin from a primary tumor elsewhere or, less often, another cutaneous site; it may present as a firm nodule, plaque, or inflammatory-appearing lesion.



---

### Cutaneous Polyarteritis Nodosa

Inflammation of small and medium blood vessels limited mainly to the skin. It may cause painful lumps, a net-like purple pattern, or ulcers, usually on the legs, without the internal-organ disease seen in systemic forms.

### Cutaneous Pseudolymphoma

A benign reactive lymphoid proliferation in the skin that can resemble cutaneous lymphoma clinically or under the microscope. It may follow an insect bite, tattoo, medicine, infection, or other stimulus; biopsy and sometimes additional studies are needed to distinguish it from true lymphoma.

### Cutaneous Skin Tag

A small, soft, usually harmless growth of extra skin, often on a narrow stalk in the neck, armpit, groin, or another skin fold. It may catch on clothing or become irritated.

### Cutaneous Small-Vessel Vasculitis

Inflammation of tiny blood vessels in the skin, often causing groups of raised purple spots, especially on the lower legs. Medicines, infections, and immune diseases can trigger it; tests may check whether other organs are involved.

### Cutaneous Squamous Cell Carcinoma

A malignant tumor of keratinocytes, often arising on chronically sun-damaged skin. It may appear as a rough, scaly, crusted, tender, or ulcerated growth and can invade locally or metastasize, particularly when high-risk features are present.

### Cutaneous T-Cell Lymphoma

A cancer of immune cells that mainly affects the skin. It may begin as itchy, scaly patches or thicker plaques and later form lumps or widespread redness; some types also involve the blood, lymph nodes, or other organs.

### Cutaneous Vasculitis

Inflammation of small blood vessels in the skin, causing red or purple spots, raised purpura, blisters, sores, or pain. Medicines, infections, autoimmune

disease, or inflammation elsewhere in the body may be responsible.

### Cuticle

The thin outermost layer of a hair shaft formed by overlapping cells that protect the inner cortex.

#### Alternate Names:

Hair Cuticle; Cuticular Layer

### Cuticle Remover Poisoning

Exposure to cuticle-remover products, which may contain alkaline or acidic chemicals that can injure the skin, eyes, mouth, or gastrointestinal tract.

### Cutis Anserina

Gooseflesh or piloerection, in which contraction of tiny muscles around hair follicles makes the skin appear rough or dimpled.

### Cutis Laxa

A group of inherited or acquired connective-tissue disorders characterized by loose, inelastic skin caused by abnormal elastic fibers. Some forms also affect the lungs, blood vessels, heart, gastrointestinal tract, or skeleton.

### Cutis Marmorata

A transient or persistent physiological mottling of the skin presenting as a reticular, bluish-purple, or erythematous marble-like vascular network, triggered by cold exposure due to vasomotor constriction of deep cutaneous arterioles and dilatation of superficial capillary venules.

#### Alternate Names:

Cutis Marmorata Telangiectatica Congenita; CMTC

#### Alternate Names:

Mottled Skin; Livedo Reticularis Physiologica

### Cutis Verticis Gyrata

A rare condition in which thickened scalp skin forms deep folds and furrows resembling the surface of the brain. It may be primary or associated with neurologic, endocrine, inflammatory, or genetic conditions; persistent scalp changes warrant assessment.

---

## Cuts And Puncture Wounds

Breaks in the skin caused by sharp, blunt, or penetrating trauma; evaluation considers depth, contamination, retained foreign bodies, tendon or nerve injury, vascular damage, and tetanus protection.

### Alternate Names:

Lacerations; Minor Open Traumatic Wounds

## Cuts and Wounds

Alternate name; see *Wound Care*.

## Cutting Balloon Catheter

A balloon catheter with tiny blades or scoring surfaces that make controlled cuts in hardened plaque while the balloon widens a narrowed blood vessel. It is used in selected vascular procedures.

### Alternate Names:

Cutting Balloon Angioplasty

## CVC

Central venous catheter, a catheter whose tip lies in a large central vein and provides intravenous access for medications, fluids, nutrition, or monitoring.

## Cyanide

A highly poisonous chemical that prevents cells from using oxygen. Exposure can rapidly cause headache, confusion, seizures, breathing failure, cardiovascular collapse, and death.

## Cyanide Poisoning

A medical emergency in which cyanide stops cells from using oxygen. It may follow smoke inhalation or industrial, accidental, or intentional exposure and can cause headache, confusion, seizures, collapse, and death within minutes.

## Cyanide Toxicity

Cellular asphyxiation caused by inhibition of mitochondrial oxidative phosphorylation. Rapid agitation or confusion, seizures, cardiovascular collapse, severe lactic acidosis, and coma may develop after smoke inhalation or ingestion; treatment must not await confirmatory testing when suspicion is high, and hydroxocobalamin is the preferred antidote in

many settings.

## Cyanoacrylates Poisoning

Exposure to cyanoacrylate adhesives, commonly known as instant glue. Contact may bond tissues or irritate the skin, eyes, or airways; ingestion usually causes local adhesion or irritation.

## Cyanosis

A blue or gray-blue color of the skin, lips, or moist linings caused by too little oxygen in blood or abnormal blood flow. Sudden cyanosis, especially with breathing difficulty, requires emergency care.

## Cyanotic

Having a bluish or slate-gray discoloration of skin or mucous membranes, usually reflecting increased deoxygenated hemoglobin or abnormal blood flow.

## Cyanotic Heart Disease

A congenital or acquired heart disorder that reduces systemic oxygen saturation and causes cyanosis, often through right-to-left shunting or inadequate pulmonary blood flow.

## CyberKnife Radiosurgery

CyberKnife is a robotic system that delivers many precisely aimed radiation beams to a tumor or other target without an open operation.

## Cycle Time

The time needed to complete one cycle of a medical process, such as collecting an image or recording a repeating body signal. The length depends on the test, equipment, and purpose.

## Cyclic Citrullinated Peptide

A substance used in a blood test for antibodies associated with rheumatoid arthritis. A positive result supports the diagnosis but must be interpreted with symptoms, examination, and other tests.

## Cyclic Edema

Recurrent swelling, often of the legs or hands, that changes with the menstrual cycle or other regu-

---

lar hormonal pattern. The diagnosis requires exclusion of heart, kidney, liver, venous, lymphatic, medication-related, and other causes of edema.

### **Cyclic Guanosine Monophosphate**

A chemical messenger inside cells that helps relax smooth muscle, widen blood vessels, process visual signals, and control other body functions. Some medicines work partly by increasing or preserving its effects.

#### **Alternate Names:**

CGMP; Cyclic GMP

### **Cyclic Hormone Therapy**

Hormone therapy in which estrogen is given continuously or for part of a cycle and progesterone is added for a defined number of days.

#### **Alternate Names:**

Sequential Hormone Therapy; Cyclic Estrogen-Progesterone Therapy

### **Cyclic Migraines**

Migraine attacks that recur in a predictable pattern, sometimes around menstruation or at regular intervals. They may include headache, nausea, light or sound sensitivity, and aura; preventive treatment may be considered when attacks are frequent or disabling.

### **Cyclic Neutropenia**

Repeated episodes of low neutrophils, the white blood cells that fight many infections. Episodes often recur at regular intervals and may cause fever, mouth ulcers, sore throat, or infections.

### **Cyclic Vomiting Syndrome**

Repeated, similar attacks of severe nausea and vomiting separated by periods of normal health. It is often related to migraine biology; dehydration and other causes must be considered during evaluation.

#### **Alternate Names:**

Cyclic Vomiting

### **Cyclic Vomiting Syndrome**

### **Cross-Reference**

Alternate name; see *Cyclic vomiting syndrome*.

### **Cyclin-Dependent Kinase**

An enzyme that helps control cell growth and division by working with a cyclin protein. Abnormal activity can contribute to cancer, and some targeted medicines block these enzymes.

### **Cyclist's Syndrome**

Pain, numbness, tingling, or altered sensation in the perineum caused by prolonged pressure on the pudendal nerve during cycling. It is a form of pudendal nerve irritation or entrapment and may improve with changes in saddle fit, riding position, activity, and specialist treatment.

#### **Alternate Names:**

Cyclist Syndrome; Pudendal Neuralgia Associated With Cycling

### **Cyclooxygenase**

An enzyme family that helps convert arachidonic acid into prostaglandins and related mediators involved in pain, inflammation, and platelet function.

#### **Alternate Names:**

COX Enzyme; Prostaglandin-Endoperoxide Synthase

### **Cyclooxygenase-2**

An inducible cyclooxygenase enzyme that generates prostaglandins involved in inflammation, pain, and fever and is inhibited by COX-2-selective medicines.

### **Cyclops**

A term that may refer to cyclopia, a rare severe congenital malformation with a single central eye, or to a fibrous nodule causing loss of extension after knee surgery; context determines the meaning.

### **Cyclospora Infection**

An intestinal infection caused by *Cyclospora* parasites, usually acquired from contaminated food or water. It can cause prolonged or recurring watery diarrhea, cramps, nausea, tiredness, and weight loss.

---

## Cyclosporiasis

An intestinal infection caused by the parasite *Cyclospora cayetanensis*. It is usually spread through contaminated food or water and may cause prolonged or repeated watery diarrhea, abdominal cramps, nausea, tiredness, loss of appetite, and weight loss.

### Alternate Names:

*Cyclospora Cayetanensis* Enteritis; *Cyclospora* Infection

## Cyclothymia

Historical or alternate term; see *Cyclothymic Disorder*.

## Cyclothymic Disorder

A chronic mood disorder involving repeated periods of hypomanic symptoms and depressive symptoms that do not meet full criteria for mania, hypomania, or a major depressive episode. Symptoms persist for at least two years in adults or one year in children and adolescents and cause clinically significant distress or impairment.

## Cylindroma

A usually benign adnexal tumor composed of basoid cells, often presenting as a firm pink or skin-coloured nodule on the scalp or face; multiple lesions may occur in Brooke-Spiegler syndrome.

## Cylindroma and Spiradenoma

Usually benign skin appendage tumors arising from sweat-gland-related structures. Cylindromas are often firm scalp or facial nodules, while spiradenomas are characteristically painful; multiple lesions may occur in Brooke-Spiegler syndrome and a rapidly enlarging or ulcerated lesion needs review.

## CYP2C19 Polymorphism

An inherited difference in a liver enzyme that breaks down some medicines. It can make a medicine act too weakly or too strongly, so genetic or medicine review may guide treatment for selected drugs.

## CYP2C9 Polymorphism

An inherited variation in the CYP2C9 drug-metabolizing enzyme that can make some medicines, especially warfarin, break down more slowly or quickly. Genotype may help inform dosing in selected settings, alongside clinical factors and monitoring.

## CYP2D6 Polymorphism

An inherited difference in a liver enzyme that breaks down many medicines. Some people process certain drugs slowly or quickly, which can change their benefits and side effects.

## CYP450 System

A family of enzymes, found mainly in the liver, that breaks down many medicines and foreign chemicals. Genetic differences and other medicines can speed up or slow down this process and alter drug effects.

## Cyproheptadine Overdose

Overdose of cyproheptadine, an antihistamine, that may cause sedation, agitation, anticholinergic effects, seizures, respiratory depression, or cardiac effects.

## Cyprus Fever

An older regional name for brucellosis, a bacterial infection acquired from infected animals or unpasteurised animal products. It may cause prolonged or recurring fever, sweats, fatigue, joint pain, and involvement of other organs.

### Alternate Names:

Brucellosis; Malta Fever; Mediterranean Fever

## Cyst

A closed sac containing fluid, air, cells, or semisolid material. Cysts can occur in the skin, organs, glands, or other tissues and may be harmless, infected, or related to another disease.

## Cystadenoma

A usually noncancerous gland-forming tumor containing fluid-filled spaces or cyst-like structures. It can occur in several organs, may enlarge or cause

---

symptoms, and sometimes requires removal or monitoring.

### **Cystectomy**

Surgery to remove part or all of the urinary bladder. Partial cystectomy is used for selected localized disease; complete or radical cystectomy removes the bladder and requires another route for urine to leave the body.

### **Cystic Acne**

Severe acne with deep, painful lumps that can leave permanent scars. It often needs clinician-directed treatment; squeezing or picking the lesions increases scarring and infection risk.

### **Cystic Diseases of the Kidney**

Disorders in which fluid-filled cysts develop in one or both kidneys. They may be inherited or acquired and can cause enlargement, pain, infection, bleeding, stones, high blood pressure, or progressive loss of kidney function.

### **Cystic Echinococcosis**

A parasitic infection in which slowly growing cysts form, most often in the liver or lungs, after swallowing *Echinococcus* eggs from infected dogs or contaminated food, water, or soil. Cysts may cause pain, cough, blockage, or serious allergic reactions if they rupture.

### **Cystic Fibrosis**

An autosomal recessive disorder caused by disease-causing changes in both copies of the *CFTR* gene. The abnormal CFTR protein disrupts the movement of salt and water across cell surfaces, making mucus thick and sticky. This mucus can block the airways and cause repeated lung infections, inflammation, and permanent lung damage. It can also block pancreatic ducts, preventing digestive enzymes from reaching the intestine and causing poor digestion, poor absorption of nutrients, or poor growth. Cystic fibrosis may also affect the liver, intestines, sweat glands, and reproductive system. People with cystic fibrosis usually have unusually salty sweat.

**Alternate Names:**

CF

### **Cystic Fibrosis Carrier Screening**

Genetic testing to identify a person who carries one disease-causing CFTR variant and usually has no cystic-fibrosis symptoms. Results help estimate reproductive risk and guide partner testing or genetic counseling.

### **Cystic Hygroma**

A congenital or early-childhood lymphatic malformation consisting of thin-walled, fluid-filled spaces, usually in the neck or upper chest. It may be detected before birth and can cause swelling, difficulty breathing, or feeding problems when large.

**Alternate Names:**

Cystic Lymphangioma; Lymphatic Malformation

### **Cystic Kidney Disease**

A group of disorders in which fluid-filled cysts develop in one or both kidneys. Some forms are inherited, while others develop later; cysts may enlarge the kidneys or reduce their function.

### **Cystic Lung Disease**

A group of disorders in which multiple air- or fluid-filled cysts develop in the lungs. Causes include lymphangioleiomyomatosis, pulmonary Langerhans-cell histiocytosis, Birt-Hogg-Dubé syndrome, lymphoid interstitial pneumonia, congenital abnormalities, infection, and smoking-related disease; high-resolution CT helps classify the pattern.

**Alternate Names:**

Multiple Cystic Lung Disease; Pulmonary Cystic Disease

### **Cystic Medial Necrosis**

An older term for weakening of the middle layer of the aortic wall. The weakened aorta may stretch into an aneurysm or tear, especially in people with inherited connective-tissue disorders, high blood pressure, or aging-related disease.

### **Cystic Teratoma**

A germ-cell tumor containing tissues derived from



---

more than one embryonic layer, commonly occurring in the ovary as a mature cystic teratoma or dermoid cyst. Most are benign, but torsion, rupture, infection, or rare malignant transformation can occur.

### Cysticercosis

Infection caused by the larval form of the pork tapeworm, *Taenia solium*, after a person swallows its eggs from contaminated food, water, or hands. Cysts may form in muscle, skin, the eyes, or the brain. Brain infection, called neurocysticercosis, may cause seizures, headaches, or other nervous-system problems. Cysticercosis is different from intestinal taeniasis and is not caused by eating undercooked pork.

### Cystine and Struvite Urinary Crystals

Two types of crystals that may appear in urine. Cystine crystals can indicate an inherited tendency to form stones, while struvite crystals may occur with some urinary infections; results need clinical interpretation.

#### Alternate Names:

Hexagonal Cystine and Magnesium Ammonium Phosphate Crystals; Urinary Calculi Crystals

### Cystine Stones

Kidney stones formed when inherited cystinuria causes excess cystine in urine. They often recur and may be difficult to dissolve; abundant fluids, medical treatment, and sometimes procedures help prevent or remove them.

### Cystinosis

A rare inherited disorder in which cystine, a building block of proteins, builds up inside cells and damages organs. It often affects the kidneys and eyes and may also impair growth, the thyroid, muscles, or other tissues.

### Cystinuria

An inherited disorder in which the kidneys release too much cystine into urine, allowing recurrent cystine kidney stones to form. It requires long-term fluid intake, monitoring, and sometimes medicines

to prevent stones.

### Cystitis

Inflammation of the bladder that may be caused by infection, irritation, medicines, radiation, or other conditions. It can cause burning or discomfort with urination, frequent or urgent urination, lower-abdominal pressure, or blood in the urine; some forms cause bladder or pelvic pain without infection.

### Cystitis Pathology

Inflammatory injury of the urinary bladder mucosa and wall. Acute cases may show edema, congestion, and neutrophils, while chronic or specialized forms show lymphocytes, fibrosis, ulceration, or characteristic epithelial changes.

### Cystocele

Bulging of the bladder into the front wall of the vagina because the supporting pelvic tissues have weakened. It may cause pelvic pressure, a vaginal bulge, urinary leakage, difficulty emptying the bladder, or no symptoms.

### Cystography

X-ray or other imaging of the urinary bladder after contrast is placed in it. It can show bladder injury, leaks, fistulas, structural abnormalities, or urine flowing backward toward the kidneys.

### Cystoid Macular Edema

Swelling at the center of the retina caused by fluid collecting in small cyst-like spaces. It can blur or distort central vision and may follow eye surgery, inflammation, diabetes, or retinal blood-vessel disease.

### Cystoisosporiasis

An intestinal infection caused by the parasite *Cystoisospora belli*. It is usually spread by swallowing contaminated food or water and may cause watery diarrhea, abdominal cramps, nausea, vomiting, loss of appetite, and fever. The illness may be prolonged or severe in people with weakened immune systems.



---

## Cystometric Study

A bladder test that measures pressure, sensation, and urine storage while the bladder fills and empties. It helps investigate urgency, leakage, difficulty emptying, weak bladder muscles, or nerve-related urinary problems.

## Cystometrogram

A recording of bladder pressure and sensation during filling and emptying. It is part of urodynamic testing used to investigate leakage, urgency, blockage, weak contractions, or nerve-related bladder problems.

## Cystometry

A bladder-function test measuring pressure and sensation while the bladder fills and empties. It helps assess urgency, leakage, obstruction, weak contractions, nerve problems, or other urinary difficulties.

## Cystoscope

A thin, lighted instrument used to examine the urethra and bladder and, when needed, perform procedures.

## Cystoscopy

Cystoscopy uses a camera passed through the urethra to inspect the bladder and urethra and may allow biopsy or treatment.

## Cystostomy

A procedure that creates an opening from the bladder to the skin or another passage so urine can drain when it cannot leave normally through the urethra. It may be temporary or long term.

## Cystotome

A small surgical instrument used to cut the thin covering of the eye's lens during selected cataract procedures. Modern instruments and techniques vary according to the operation.

## Cystourethrography

An X-ray examination of the bladder and urethra after contrast material is placed in the bladder. Im-

ages during filling and urination can show leaks, narrowing, structural problems, or urine flowing backward toward the kidneys.

## Cysts

Closed sacs or tissue cavities containing fluid, air, pus, blood, cells, or semisolid material. They can occur in the skin, organs, glands, or other tissues; many are harmless, but some result from infection, developmental or inherited disease, or a tumor. A new, growing, painful, changing, or infected cyst-like lump should be medically assessed.

## Cytochrome B5 Reductase

An enzyme that keeps hemoglobin in the chemical form needed to carry oxygen. An inherited deficiency can cause methemoglobinemia, in which the skin looks blue and blood cannot deliver oxygen normally.

## Cytochrome C

A protein inside mitochondria that carries electrons during energy production. When released from damaged mitochondria, it can also help start the controlled death of a cell.

## Cytochrome P450

A family of enzymes, found mainly in liver cells, that chemically changes many medicines, toxins, and hormones so the body can remove them. Genetic differences and drug interactions can alter their activity.

## Cytogenetic Karyotyping

A laboratory examination that arranges a person's chromosomes to check their number and large structural changes. It can help investigate inherited conditions, developmental differences, infertility, pregnancy concerns, and some cancers.

### Alternate Names:

Karyotype

## Cytogenetics

The study of chromosomes, including their number, structure, and arrangement. Laboratory tests can help identify inherited conditions, developmental

---

differences, some pregnancy problems, leukemia, and other cancers.

### **Cytokine**

A small protein messenger that allows cells to communicate during immunity, inflammation, infection, healing, and blood-cell production. Cytokines can protect tissues, but excessive or poorly controlled activity can cause disease.

### **Cytokine Release Syndrome**

A potentially serious whole-body inflammatory reaction caused by a sudden release of immune signaling proteins, sometimes after immune-based cancer treatment or infection. Fever, low blood pressure, breathing difficulty, and organ problems may occur.

### **Cytokine Storm**

A dangerous overreaction of the immune system in which large amounts of inflammatory messenger proteins are released. It can cause widespread tissue injury, low blood pressure, breathing failure, organ damage, and shock.

### **Cytokines**

Small signaling proteins released by immune and other cells that regulate inflammation, immunity, blood-cell production, and tissue responses.

#### **Alternate Names:**

Immune Signaling Proteins; Cytokine Mediators

### **Cytologic Evaluation**

Microscopic examination of individual cells or small cell groups from a fluid, brushing, scraping, or fine-needle sample to detect infection, inflammation, or malignancy.

### **Cytology**

The study of individual cells under a microscope. Cells may come from a body fluid, scraping, brushing, or needle sample and can be examined for cancer, infection, inflammation, or other disease.

### **Cytology Exam Of Pleural Fluid**

Microscopic examination of fluid around the lung

for malignant cells and other cellular changes, performed with chemical and microbiologic testing when evaluating an effusion.

### **Cytology Exam Of Urine**

Microscopic examination of shed urinary-tract cells, used mainly to detect high-grade urothelial malignancy and selected abnormal cells; a negative result does not exclude cancer.

### **Cytomegalovirus**

A common herpesvirus that often causes no symptoms or a mild illness in healthy people. It can cause serious disease in unborn babies and people with weakened immunity, affecting organs such as the lungs, eyes, brain, or digestive tract.

#### **Alternate Names:**

CMV; Human Betaherpesvirus 5; Human Cytomegalovirus (HCMV)

### **Cytomegalovirus Gastroenteritis and Colitis**

An opportunistic viral gastrointestinal infection caused by cytomegalovirus (CMV) reactivation in severely immunocompromised individuals (e.g., advanced HIV/AIDS, solid-organ transplantation), characterized by mucosal ulceration, hematochezia, abdominal pain, and potential bowel perforation.

#### **Alternate Names:**

CMV Colitis; Cytomegalovirus Colitis; CMV Enteritis; Gastrointestinal Cytomegalovirus

### **Cytomegalovirus Infection**

Infection with cytomegalovirus, which may cause no symptoms or a fever-and-fatigue illness in healthy people. Infection during pregnancy or in someone with weakened immunity can cause hearing loss, organ disease, or other serious complications.

### **Cytomegalovirus Retinitis**

A severe, sight-threatening opportunistic infection of the retina caused by cytomegalovirus, occurring almost exclusively in patients with advanced cellular immunodeficiency (typically individuals with AIDS and CD4 counts below 50 cells/ $\mu$ L or recipients of organ transplants). It manifests as full-thickness

---

necrotizing retinitis with perivascular retinal hemorrhages and fluffy white infiltrates ("pizza-pie" appearance), causing floaters, scotomas, decreased visual acuity, and a high risk of rhegmatogenous retinal detachment.

**Alternate Names:**

CMV Retinitis; Cytomegaloviral Chorioretinitis

### Cytopathology

The examination of individual cells under a microscope to help diagnose cancer, infection, inflammation, or other disease. Cells may come from fluid, a brushing, a smear, or a needle sample.

### Cytopenia

A lower-than-normal number of one or more blood-cell types: red blood cells, white blood cells, or platelets. The effects depend on which cells are low and how severe the reduction is.

### Cytoplasm

The jelly-like material inside a cell outside its nucleus. It contains the cell's working structures and is where many chemical reactions, transport processes, and energy-producing activities occur.

### Cytoreductive Surgery

Surgery that removes as much visible cancer as can be taken safely. It may not remove every cancer cell and is often combined with chemotherapy, radiation, or other treatment; its value depends on the cancer's type, site, and spread.

### Cytoskeleton

A network of protein fibers inside a cell that maintains its shape, moves materials, enables movement, and helps the cell divide. It constantly changes in response to the cell's needs.

### Cytotoxic

Toxic to cells or capable of killing cells, as with some chemotherapy drugs, toxins, or cytotoxic immune cells.

### Cytotoxic Alopecia

Hair loss caused by exposure to a cytotoxic treatment or other toxic injury to actively growing hair follicles.

**Alternate Names:**

Drug-Induced Alopecia; Chemotherapy-Related Hair Loss

### Cytotoxic T Cell

An immune cell that recognizes and kills infected, abnormal, or cancerous cells. It identifies small protein signals displayed on the target cell and can trigger that cell to die.

### Cytotoxicity

The ability of a substance or cell to damage or kill another cell. This may be part of normal immune defense or may result from medicines, toxins, radiation, or other harmful exposures.

### Differential Diagnosis of Viral Respiratory Infection and Allergic Rhinitis

Colds, influenza, and allergies can all cause nasal symptoms and cough but differ in cause, fever pattern, duration, and contagiousness. Viral illness usually improves with supportive care, while allergies respond to trigger reduction and appropriate antihistamine or nasal treatment.

**Alternate Names:**

Upper Respiratory Conditions; Allergic and Viral Rhinopathy

---

# D

## 1p36 Deletion Syndrome

A genetic disorder caused by loss of a piece of chromosome 1. It may cause developmental and intellectual disability, small head size, low muscle tone, seizures, hearing loss, heart problems, and distinctive facial features; severity varies.

### Alternate Names:

Monosomy 1p36; Chromosome 1p36 Deletion Syndrome

## 2,3-Bisphosphoglycerate

A substance made in red blood cells that helps hemoglobin release oxygen to body tissues.

### Alternate Names:

2,3-DPG; 2,3-Bisphosphoglycerate

## 22q11.2 Deletion Syndrome

A genetic condition caused by a missing piece of chromosome 22. It may cause heart defects, palate problems, reduced immune defenses, low calcium, distinctive facial features, learning or developmental difficulties, and increased risk of mental illness. The effects differ widely between people.

### Alternate Names:

DiGeorge Syndrome Spectrum; Velocardiofacial Syndrome (VCFS); 22q11.2DS

## D & C

Dilation and curettage, a procedure in which the cervix is widened and tissue is removed from the uterine cavity for treatment or diagnosis.

## D And C

Dilation and curettage, a procedure that opens the cervix and removes tissue from the uterine lining

for treatment or diagnosis.

## D And E

Dilation and evacuation, a procedure that gently opens the cervix and removes pregnancy tissue from the uterus using suction and instruments. It is performed by trained clinicians for selected pregnancy-related indications.

## D Trisomy Syndrome

An older name for trisomy 13 (Patau syndrome), a chromosome disorder caused by an extra chromosome 13. It can cause severe developmental impairment, brain and facial differences, heart defects, and abnormalities of other organs.

## D-Dimer

A substance released into the blood when a blood clot breaks down. A high level can occur with a blood clot, but also with infection, pregnancy, recent surgery, cancer, or other conditions. A normal result can help rule out a clot in carefully selected patients whose initial risk is low or moderate.

## D-Dimer Test

A blood test measuring a substance released when blood clots are broken down. A normal result can help rule out a clot in carefully selected low-risk people; a high result has many possible causes and does not prove a clot.

### Alternate Names:

D-Dimer Assay; Fibrin Degradation Fragment Test

## D-Lactic Acidosis

A rare metabolic encephalopathy and high anion gap

---

metabolic acidosis occurring in patients with short bowel syndrome, resulting from colonic bacterial fermentation of unabsorbed carbohydrates into the D-isomer of lactic acid, which humans metabolize slowly.

**Alternate Names:**

D-Lactate Encephalopathy

**D-Xylose Absorption**

An older test of how well the small intestine absorbs a simple sugar after it is swallowed. Low amounts in blood or urine may suggest intestinal lining disease, but kidney function and collection quality affect results.

**D.O.**

Doctor of Osteopathic Medicine, a physician degree whose graduates are licensed to practice medicine and surgery like MD graduates in the United States.

**D2 Receptor**

A protein on cells that responds to dopamine, a brain chemical involved in movement, motivation, and reward. Some medicines work partly by blocking or stimulating this receptor.

**D&C Instruments**

Instruments used to dilate the cervix and remove tissue from the uterus during dilation and curettage.

**Da Costa's Syndrome**

Older term; specify the underlying cause of chest pain, palpitations, fatigue, or breathlessness.

**Dacarbazine**

A chemotherapy medicine used for Hodgkin lymphoma, melanoma, and some other cancers. It may cause severe nausea, low blood counts, or liver injury.

**Daclizumab**

A monoclonal antibody that blocks the interleukin-2 receptor on activated immune cells. It was used for selected immune conditions but was withdrawn or restricted in many settings because of serious liver

and other risks.

**Dacomitinib**

A targeted medicine that blocks several members of the epidermal-growth-factor receptor family and is used for selected EGFR-mutated lung cancers. Diarrhea, skin reactions, mouth sores, and lung or liver problems may occur.

**Dacrocytes**

Teardrop-shaped red blood cells seen on a blood smear. They may occur when red blood cells pass through scarred or crowded bone marrow, and can be seen in myelofibrosis, bone-marrow cancers, and severe thalassemia.

**Alternate Names:**

Teardrop Cells

**Dacryoadenitis**

Inflammation of a tear-producing gland, causing painful swelling near the outer upper eyelid. It may be caused by infection, inflammation, or a condition affecting the whole body.

**Dacryocystitis**

Infection or inflammation of the tear sac near the inner corner of the eye, usually because the tear-drainage passage is blocked. It can cause pain, redness, swelling, and pus and may need prompt treatment.

**Dacryocystocele**

A fluid-filled swelling near the inner corner of the eye caused by blockage and widening of the tear-drainage system. In a newborn it may interfere with breathing if it extends into the nose and needs prompt assessment.

**Dacryocystography**

An imaging test in which contrast material is put into the tear-drainage passages to show narrowing, blockage, abnormal connections, or leakage. It may help plan treatment for persistent tearing or repeated tear-sac infection.



---

### **Dacryocystorhinostomy**

Surgery creating a new passage from the tear sac into the nose to bypass a blocked tear duct. It is used for persistent tearing or repeated tear-sac infections.

### **Dacryocyte Analysis**

Examination or counting of teardrop-shaped red blood cells on a blood smear; numerous dacryocytes may occur with marrow fibrosis, infiltration, severe anemia, or splenic disease.

### **Dacryolithiasis**

Formation of a stone or concretion within the lacrimal drainage system, which may cause excessive tearing, discharge, recurrent dacryocystitis, or obstruction.

### **Dacryostenosis**

Narrowing or blockage of the passages that drain tears from the eye into the nose. It can cause constant tearing, sticky discharge, or repeated infection of the tear sac.

### **Dactinomycin**

An anticancer medicine that binds DNA and interferes with gene activity and cell growth. It is used for selected cancers and can cause bone-marrow suppression, mouth sores, nausea, and tissue injury if it leaks outside a vein.

### **Dactylitis**

Swelling and inflammation of an entire finger or toe, producing a sausage-like appearance. It can occur with sickle-cell disease, psoriatic arthritis, infection, or other inflammatory conditions.

### **Dactylomegaly**

Abnormal enlargement of a finger or toe, which may occur as an isolated finding or as part of an endocrine, skeletal, vascular, or inflammatory disorder.

### **Dactylosymphysis**

A birth difference in which two neighboring fingers

or toes are joined together by skin, bone, or both. The connection may limit movement or affect growth and hand or foot function.

### **Daidzein**

A naturally occurring isoflavone found in soy and other legumes that can be converted by intestinal bacteria into related compounds.

### **Daily Bowel Care Program**

A planned routine for preventing and treating constipation or fecal incontinence, often including timing, fluids, diet, activity, medication, and rectal measures when prescribed.

### **Dakin Solution**

A dilute sodium hypochlorite solution used as a topical antiseptic for selected contaminated or infected wounds. Concentration and exposure time must be controlled because stronger solutions can injure viable tissue.

### **Dalrymple Sign**

A clinical ophthalmologic sign characterized by abnormal widening of the palpebral fissure and retraction of the upper eyelid, revealing a strip of white sclera above the superior corneal limbus in primary gaze, classically seen in Graves ophthalmopathy and thyrotoxicosis.

#### **Alternate Names:**

Dalrymple's Sign; Thyrotoxic Eyelid Retraction

### **Dalteparin**

An injectable blood thinner used to prevent or treat clots in veins and the lungs. It reduces clotting through the body's natural anticoagulant system; bleeding is the main risk, and kidney disease may require dose adjustment.

### **Dalteparin Sodium**

The sodium salt of dalteparin, a low-molecular-weight heparin used to prevent or treat venous thromboembolism; bleeding is the principal serious adverse effect.

---

## Daltonism

Color-vision deficiency, especially difficulty distinguishing certain colors because of abnormal or absent retinal cone function.

## DALY

Disability-adjusted life year, a measure combining years of healthy life lost because of premature death and disability.

## Damage To The Amygdala

Damage to the amygdala can affect fear processing, emotional learning, memory, threat recognition, and social behavior. Causes include trauma, stroke, infection, tumor, epilepsy, or degenerative disease, and effects depend on the side and extent of injury.

## Damage-Control Resuscitation

Emergency treatment for severe injury that quickly controls bleeding, gives appropriate blood products, prevents dangerous cooling, and corrects clotting problems while avoiding harmful excess fluid.

## Damage-Control Surgery

A planned series of operations for a critically ill or severely injured person who cannot safely tolerate a long repair. The first operation controls bleeding or contamination; after stabilization, a later operation completes the repair.

## Damus-Kaye-Stansel Procedure

A heart operation for selected children with a single working ventricle and a blocked or narrowed route from the ventricle to the aorta. It connects the pulmonary artery to the aorta to create a wider route for blood leaving the heart.

### Alternate Names:

DKS Procedure; Damus-Stansel-Kaye Operation

## Danazol

A synthetic androgen with antigonadotropic effects, used selectively for endometriosis, hereditary angioedema, or other conditions; androgenic, lipid, liver, and thrombotic adverse effects limit use.

## Dance Sign

A clinical physical examination finding in pediatric intussusception characterized by a palpable emptiness or flattening in the right lower quadrant (ileocecal fossa), resulting from the invagination and distal displacement of the cecum and terminal ileum into the ascending colon.

### Alternate Names:

Dance's Sign; Signe de Dance

## Dandruff

A common scalp condition causing small white or yellow flakes and sometimes itching. It is often related to oily skin, seborrheic dermatitis, or the scalp's reaction to the normal yeast *Malassezia*; it is not usually contagious. Persistent severe flaking, redness, sores, or hair loss may indicate another scalp disorder.

## Dandruff And Scalp Conditions

Scalp flaking and itching may result from dandruff, seborrheic dermatitis, psoriasis, contact allergy, fungal infection, or head lice. Persistent redness, sores, pain, or hair loss needs medical assessment.

## Dandy Fever

An older name for dengue fever, a mosquito-borne viral infection causing sudden fever, headache, severe muscle or joint pain, nausea, or rash. Severe dengue can cause bleeding, shock, and organ failure.

## Dandy-Walker Malformation

A condition present at birth in which part of the cerebellum does not develop normally and one of the brain's fluid spaces becomes enlarged. It may cause fluid buildup in the brain, delayed development, balance problems, or seizures.

## Dandy-Walker Syndrome

A birth difference in the cerebellum and nearby fluid spaces of the brain. It may cause enlarged head size from fluid buildup, delayed development, balance problems, seizures, or no symptoms.

---

## Dane Particle

The complete infectious form of hepatitis B virus. It contains the virus's genetic material inside a protein shell and outer envelope, allowing it to enter liver cells and reproduce.

## Dangerous Abbreviations

Abbreviations that can be misread and cause medication or clinical errors; safer practice is to write the complete term when an abbreviation is ambiguous.

## Danon Disease

A rare inherited disorder, usually caused by an LAMP2 gene change, in which cells cannot properly recycle materials. It may cause heart-muscle disease, muscle weakness, intellectual disability, or eye problems.

## Dantrolene

A muscle relaxant that reduces the release of calcium inside muscle cells. It is the main treatment for malignant hyperthermia and may cause weakness or liver injury.

## Dapsone

A sulfone antimicrobial used for leprosy, selected dermatitis, and other infections; haemolysis, especially with G6PD deficiency, methaemoglobinaemia, agranulocytosis, and severe skin reactions are important risks.

## Daptomycin

An antibiotic used for selected serious infections caused by gram-positive bacteria, including resistant strains. It can damage muscles and kidneys, so treatment requires appropriate testing and monitoring.

## Daptomycin Resistance

Reduced or absent response of bacteria to daptomycin, often because changes in the bacterial membrane prevent the antibiotic from working. It can develop during treatment and limits options for serious infections.

## Darbepoetin Alfa

A long-acting medicine that stimulates the bone marrow to make more red blood cells. It is used for selected anemia related to kidney disease or cancer treatment, but can raise blood pressure and the risk of blood clots.

## Darier Sign

A skin reaction in which rubbing a lesion causes it to become red, swollen, and itchy within minutes. It strongly suggests cutaneous mastocytosis, but the finding must be considered with the rest of the examination.

### Alternate Names:

Darier's Sign; Mastocytoma Stroking Sign

## Darier's Disease

An inherited skin disorder that causes persistent, firm, often greasy or crusted bumps, usually on the chest, back, scalp, or skin folds. It may also change the nails or affect the mouth. Heat, sweating, friction, and sunlight can trigger flares.

### Alternate Names:

Darier Disease; Darier-White Disease; Keratosis Follicularis

## Dark Adaptation

The gradual adjustment that allows the eyes to see better after moving from bright light into darkness. Poor dark adaptation may result from retinal disease, vitamin A deficiency, aging, or some medicines.

## Dartos

A thin muscle layer beneath the skin of the scrotum. It tightens or relaxes the scrotal skin in response to temperature and other signals, helping regulate the position of the testicles.

## Darunavir

An antiretroviral medicine that blocks HIV protease, an enzyme needed to make new viruses. It is used with other HIV medicines and has important interactions and possible liver or skin reactions.

---

### **Darwin's Tubercle**

A harmless small bump on the upper rim of the outer ear. It is a common normal variation in ear shape and usually needs no treatment.

### **Dasatinib**

A targeted cancer medicine that blocks abnormal tyrosine-kinase signals, especially those caused by BCR::ABL1. It treats selected blood cancers and may cause low blood counts, fluid around the lungs, bleeding, or abnormal heart rhythms.

### **DASH Diet**

An eating pattern designed to lower blood pressure by emphasizing vegetables, fruits, whole grains, beans, nuts, and low-fat dairy while limiting salt, saturated fat, and sugary foods. Individual needs may require adjustment.

### **Dating Ultrasound**

An ultrasound examination, usually performed in early pregnancy, used to estimate gestational age and calculate the expected date of delivery. First-trimester crown-rump length is generally the most accurate ultrasound measurement for dating.

#### **Alternate Names:**

Pregnancy Dating Scan; Gestational-Age Ultrasound

### **Daubenton's Plane**

A craniometric reference plane passing through the opisthion, the midpoint of the posterior margin of the foramen magnum, and the lower margins of the orbits. It is used in describing skull orientation.

### **Daunorubicin**

A chemotherapy medicine that damages cancer-cell DNA and is used mainly for selected blood cancers. It can lower blood counts and, especially after repeated doses, weaken or damage the heart.

### **Daunorubicin Hydrochloride**

The hydrochloride salt of daunorubicin, an anthracycline chemotherapy drug that interferes with DNA and is used mainly for acute leukemias. Important

risks include bone-marrow suppression and heart damage.

### **Dav Regimen**

An abbreviation used for more than one cancer-treatment combination. In older leukemia reports, DAV meant daunorubicin, cytarabine, and etoposide; in newer reports, it may mean daunorubicin, cytarabine, and venetoclax. The source must identify the drugs and cancer before the regimen is used.

### **David Procedure**

An operation that replaces an enlarged or weakened aortic root while keeping the person's own aortic valve. The valve is supported inside a graft and the coronary arteries are reattached; bleeding, stroke, valve leakage, or recurrent aortic disease are possible.

#### **Alternate Names:**

David Operation; Valve-Sparing Root Replacement; Reimplantation of Aortic Valve

### **Daw**

An abbreviation meaning dispense as written, indicating that the pharmacist should dispense the prescribed product rather than automatically substitute a generic equivalent when permitted by law.

### **Dawn Phenomenon**

A rise in blood glucose during the early morning caused by normal hormone changes that help the body wake up. It may be more noticeable in people with diabetes whose insulin supply cannot match the rise.

### **Dawson Fingers**

Long, finger-like areas seen on some brain MRI scans in multiple sclerosis. They form around small veins near the brain's fluid spaces and support, but do not by themselves prove, the diagnosis.

#### **Alternate Names:**

Dawson's Fingers; Perivenular MS Plaques

### **Day Hospital**

A facility that provides multidisciplinary medical,

---

psychiatric, rehabilitative, or diagnostic care during the day without overnight admission. It can support intensive treatment while allowing the patient to return home.

### Day-Blindness

Poor vision in bright light, also called hemeralopia. It can result from retinal disease, cone dysfunction, or other eye disorders.

### Daytime Sleepiness

Alternate name; see *Excessive daytime sleepiness*.

### DBA

Abbreviation for Diamond-Blackfan anemia, an inherited bone-marrow failure syndrome that usually causes anemia in infancy or early childhood and may be associated with characteristic physical differences or other congenital abnormalities.

### Dcc Gene

A gene that helps guide developing nerve fibers and may also help control cell growth. Harmful changes in this gene can cause some developmental nerve disorders and may contribute to certain cancers.

### DCIS

Alternate index label; see *Ductal carcinoma in situ*.

### DDH

Alternate index label; see *Hip dysplasia*.

### De Lange Syndrome

A congenital developmental disorder, also called Cornelia de Lange syndrome, involving characteristic facial features, growth and developmental delay, limb abnormalities of variable severity, and intellectual disability.

#### Alternate Names:

Cornelia De Lange Syndrome

### De Musset Sign

Repeated head nodding that matches the heartbeat. It can occur when the aortic valve leaks severely, but it is uncommon and cannot confirm the diagnosis

without examination and heart tests.

### De Novo

Newly arising in an individual rather than inherited from either parent; the term is commonly used for genetic variants or disease presentations.

### De Novo Variant

A genetic change found in a person that was not found in either biological parent. It usually begins in an egg, sperm, or very early embryo and may cause disease; testing parents can help estimate the chance in future pregnancies.

### De Quervain Tendinitis

Alternate name; see *De Quervain Tenosynovitis*.

### De Quervain Tenosynovitis

Painful swelling and narrowing around the tendons on the thumb side of the wrist. It causes pain with thumb movement, gripping, or lifting and may cause tenderness or swelling near the base of the thumb.

#### Alternate Names:

Radial Styloid Tenosynovitis; De Quervain Tendinopathy

### De Quervain Thyroiditis

A self-limiting subacute granulomatous inflammatory disorder of the thyroid gland, typically occurring following a viral upper respiratory infection, presenting with exquisite anterior neck tenderness, thyroid enlargement, fever, elevated erythrocyte sedimentation rate, and transient thyrotoxicosis followed by recovery.

#### Alternate Names:

Subacute Granulomatous Thyroiditis; Giant Cell Thyroiditis

### De Quervain's Tendonitis

Alternate spelling; see *De Quervain Tenosynovitis*.

### De Quervain's Tenosynovitis

Alternate spelling; see *De Quervain Tenosynovitis*.

---

### De Winter ECG Pattern

An ECG pattern that can signal a sudden severe blockage in a major artery supplying the front of the heart, even without the usual ST elevation. It is a heart-attack emergency requiring immediate specialist assessment and treatment; exercise testing must not be used.

#### Alternate Names:

De Winter Pattern; De Winter T Waves; De Winter Sign

### De Winter T-Wave Pattern

An electrocardiographic presentation considered a high-risk ST-segment elevation myocardial infarction (STEMI) equivalent, indicative of acute proximal left anterior descending (LAD) coronary artery occlusion. Characterized by 1- to 3-millimeter upsloping ST-segment depression at the J-point in precordial leads  $V_1$  through  $V_6$  that transitions into tall, prominent, symmetric, hyperacute T waves, often accompanied by 0.5- to 1-millimeter ST-segment elevation in lead aVR.

#### Alternate Names:

De Winter ECG Sign; De Winter Syndrome

### De-Escalation

A set of verbal and nonverbal techniques used to reduce agitation and prevent violence while preserving dignity and safety. It includes calm communication, appropriate personal space, clear limits, environmental modification, and offering choices.

### Deacetylase Inhibitor

A medicine or research compound that blocks enzymes removing chemical groups called acetyl groups from proteins. Some alter gene activity and are used for selected cancers; effects and risks depend on the specific medicine.

### Dead Space

The part of a breath that does not exchange oxygen and carbon dioxide with the blood, including the nose, throat, and larger airways. It increases when air reaches lung areas with too little blood flow.

### Dead Space Ventilation

Breathing that moves air through areas of the lungs where little or no blood is available for gas exchange. It may increase with a pulmonary embolism, low blood flow, or emphysema and can make breathing less effective.

### Dead Time

The brief period after an event during which a detector or recording system cannot register another event, limiting the measured count rate.

### Deafblindness

Combined hearing and vision loss that significantly affects communication, access to information, or moving around independently. The degree of hearing and sight loss varies; support is tailored to each person's remaining senses and communication needs.

#### Alternate Names:

Deaf-Blindness; Dual Sensory Loss

### Deafferentation

Loss or interruption of nerve signals carrying sensation from a body area to the brain or spinal cord. It may cause numbness, unusual sensations, poor awareness of position, or persistent nerve pain.

### Deafness

Complete or severe loss of hearing in one or both ears. The problem may involve sound passing through the outer or middle ear, the inner ear or hearing nerve, or more than one of these areas.

### Deaner

A former brand name for deanol, also called dimethylaminoethanol, an older product once marketed for behavioral or learning problems. It is not a standard modern treatment, and evidence for medical benefit is limited.

### Death

The permanent end of life. It may be confirmed when the heart and breathing stop permanently or when all brain function has permanently stopped,



---

according to medical and legal standards.

### **Death Rate**

The number of deaths in a defined population during a specified period, usually expressed per population size and used in public-health comparisons.

### **Death Rattle**

Noisy, rattling breathing caused by secretions accumulating in the throat or upper airways, often occurring near the end of life when swallowing and coughing weaken.

### **Deauville Criteria**

A five-point scale used with a PET scan to compare lymphoma activity with normal organs. The score helps clinicians judge whether treatment is working, but it must be interpreted with the type of lymphoma and other findings.

### **Deaver Retractor**

A hand-held surgical instrument with a broad, curved blade used to gently move organs or tissues aside. It is often used during chest or abdominal surgery to improve the surgeon's view.

### **Debility**

General physical weakness or reduced ability to function, often caused by illness, aging, poor nutrition, prolonged inactivity, or recovery from major stress. The underlying cause should be assessed when weakness persists.

#### **Alternate Names:**

Weakness

### **Debride**

To remove dead, damaged, infected, or contaminated tissue from a wound or body site to support healing and control infection.

### **Debridement**

Removal of dead, damaged, infected, or contaminated tissue from a wound. It may be done with instruments, irrigation, or other methods to reduce infection, reveal healthy tissue, and support healing.

### **Debulk**

To reduce the size or amount of a tumor or other abnormal tissue, usually by surgery or another treatment, without necessarily removing every abnormal cell.

### **Debulking**

Removal of as much of a tumor as possible when removing all of it is unsafe or impossible. The aim is to reduce symptoms or improve the effect of later treatment; benefit depends on the cancer and situation.

### **Decalcification**

Loss of calcium salts from bone, teeth, or other mineralized tissue. In teeth it may weaken enamel; in bone it can contribute to reduced strength.

### **Decannulation**

Removal of a tube, especially a tracheostomy tube, after a person can breathe safely, protect the airway, and clear secretions without it. The site is monitored for bleeding, narrowing, or breathing difficulty.

### **Decanoic Acid**

A ten-carbon saturated fatty acid, also called capric acid, found in some fats and studied for antimicrobial, metabolic, and antiseizure effects.

### **Decapitation**

Separation of the head from the body, seen in severe trauma (vehicle accidents, train injuries, explosions), industrial accidents, hanging with judicial execution, or homicide. Complete decapitation is almost always rapidly fatal due to exsanguination and brain ischemia.

### **Decay Correction**

A calculation that adjusts a radioactive measurement for the gradual loss of activity over time. It allows imaging or laboratory results obtained at different times to be compared accurately.

---

### Deceased Donor

Individual who has died (usually from brain death or cardiac death) and donated organs, tissues, or both for transplantation.

### Decenoic Acid

An unsaturated ten-carbon fatty acid; its biologic effects depend on the position and geometry of its double bond and on tissue and dietary context.

### Decerebrate Posture

Abnormal straightening of the arms and legs caused by severe injury to the brainstem or nearby brain structures. It is a sign of serious neurologic injury and may occur after head injury, bleeding, or increased pressure in the skull.

### Decerebrate Posturing

Abnormal straightening of the arms and legs, sometimes with inward turning of the arms, caused by severe brain or brainstem injury. It indicates a medical emergency, but the posture alone cannot predict recovery.

### Decerebrate Rigidity

Severe, abnormal straightening of the arms and legs, sometimes with the arms turned inward. It usually indicates serious brain or brainstem injury or dangerously high pressure inside the skull and requires emergency assessment.

### Decerebrate State

A severe neurologic posture marked by extension and inward rotation of the arms and extension of the legs, usually indicating serious injury to the brainstem or structures above it.

### Decerebration

Severe brain dysfunction that causes abnormal straightening of the arms and legs. It usually results from major brain or brainstem injury or increased pressure inside the skull and is a medical emergency.

### Decibel

A unit used to describe sound intensity. Higher

values mean louder sound; repeated or very loud exposure can damage hearing, depending on the level and duration.

### Decidua

The specialized lining of the uterus during pregnancy. It is shed after delivery and helps support the pregnancy and placenta.

### Decidua Basalis

The specialized maternal endometrium directly underlying the implanted blastocyst and interacting with chorionic villi, forming the maternal contribution to the mature placenta.

#### Alternate Names:

Basal Decidua

### Deciduitis

Inflammation of the specialized uterine lining that forms during pregnancy. It may be associated with infection or other pregnancy complications and can require assessment when pain, fever, or bleeding occurs.

### Deciduosis

Pregnancy-related tissue from the uterine lining growing in an unusually large area or outside the uterus, such as on the cervix or pelvic surfaces. It can look like a tumor during examination or surgery but is usually benign.

### Deciduous Teeth

The 20 primary teeth that erupt during childhood and are gradually replaced by permanent teeth. Caries, infection, trauma, or premature loss can affect nutrition, speech, space for permanent teeth, and oral development.

### Deciduous Tooth

A primary or baby tooth that is eventually shed and replaced by a permanent tooth. Early decay or loss can affect chewing, speech, and the space needed for permanent teeth.

---

## Decision Making

The process of comparing information, options, benefits, risks, and preferences to choose an action or plan, including shared decision-making in health-care.

## Decision-Making Capacity

The ability to understand information about a medical decision, consider the possible results, compare choices, and communicate a choice. It applies to a specific decision, may change over time, and is different from legal competence.

## Decitabine

A cancer medicine that changes abnormal gene activity and interferes with growth of blood-forming cancer cells. It is used for selected blood cancers and may cause low blood counts, infection, bleeding, or fatigue.

## Declarative Memory

Memory for facts, information, and personal events that a person can consciously remember and describe.

## Decoction

A preparation made by simmering plant material in water to extract soluble substances. The strength and safety depend on the plant, preparation, dose, contamination, and interactions with medicines.

## Decompensated Cirrhosis

Advanced liver scarring that has begun to cause problems such as abdominal fluid buildup, internal bleeding, jaundice, or confusion from waste products. It requires specialist care and may require assessment for liver transplantation.

## Decomposition

The natural breakdown of a dead body after death. Cells and tissues first break down themselves, followed by the action of microorganisms; the speed and appearance depend on temperature, moisture, and other conditions.

## Decompression

Reduction of pressure on a tissue or body compartment, such as removing pressure from a spinal nerve, draining an obstructed organ, or treating pressure changes after diving.

## Decompression Sickness

Illness caused by nitrogen bubbles forming in the body when pressure falls too quickly, usually after diving. It may cause joint pain, skin changes, breathing problems, weakness, confusion, or paralysis and needs emergency care.

## Deconditioning

Loss of strength, fitness, balance, and function after inactivity, illness, or prolonged bed rest. It increases fall risk but can often improve gradually with safe movement, exercise, nutrition, and rehabilitation.

## Decongestant

A medicine that reduces swelling inside the nose and relieves nasal congestion. Some oral decongestants can raise blood pressure or speed the heartbeat; prolonged use of some nasal sprays can cause rebound congestion.

### Alternate Names:

Decongestants; Nasal Decongestants

## Decontamination

Steps that remove or reduce a harmful substance from the skin, eyes, clothing, or digestive tract and prevent further absorption. The correct method depends on the substance; some exposures require washing, irrigation, or specialist treatment.

## Decorin

A small protein-sugar molecule found around cells. It helps organize collagen, influences tissue growth and repair, and may affect scar formation.

## Decorticate Posture

Abnormal bending of the arms with straightening of the legs, caused by severe injury to the brain. It is a sign of serious brain damage and reduced consciousness.

---

### **Decorticate Posturing**

Abnormal bending of the arms toward the chest with straightening of the legs, caused by severe brain injury. It signals impaired consciousness and requires emergency assessment; posture alone cannot predict outcome.

### **Decorticate Rigidity**

An abnormal stereotyped motor posturing in comatose patients characterized by flexion and adduction of the upper extremities with extension and internal rotation of the lower extremities, reflecting damage to the cerebral hemispheres or internal capsule above the level of the red nucleus (midbrain).

#### **Alternate Names:**

Decorticate Posturing; Flexor Posturing

### **Decreased Alertness**

Decreased alertness is reduced awareness or responsiveness that may result from metabolic illness, intoxication, infection, seizure, stroke, trauma, or sleep-related causes.

### **Decreased Appetite**

A pathological or temporary reduction in the desire to ingest food, commonly caused by systemic illness, chronic infection, malignancy, psychological stress, depression, or medications.

#### **Alternate Names:**

Anorexia; Hyporexia; Loss of Appetite; Poor Appetite

### **Decreased Breath Sounds**

Softer-than-normal sounds heard when listening to the lungs, suggesting reduced air movement. Causes include fluid or air around the lung, partial lung collapse, blockage, or shallow breathing; sudden one-sided loss can be an emergency.

### **Decreased Concentration**

Reduced ability to sustain attention or mental focus; causes include sleep loss, stress, medications, mood disorders, substance use, and neurologic or medical illness.

### **Decreased Dentin**

Reduced thickness or formation of dentin, the mineralized tissue beneath tooth enamel; causes include developmental defects, wear, erosion, caries, or dental treatment.

### **Decreased Libido**

Reduced sexual desire that may be related to relationships, mood, stress, medications, hormones, chronic illness, pain, or other health factors.

### **Decreased Tear Production**

Alternate index label; see *Dry eyes*.

### **Decubitus**

A lying or reclining position; the term also appears in decubitus ulcer, a pressure injury caused by prolonged pressure and shear.

### **Decubitus Ulcer**

Damage to skin and deeper tissue caused by prolonged pressure, sliding, or rubbing, usually over a bony area. It may begin with persistent redness and progress to an open wound, especially in people with limited movement.

### **Deep**

Located farther inside the body or a structure, away from its surface. For example, deep muscles lie beneath muscles closer to the skin.

### **Deep Brain Stimulation**

A treatment in which implanted electrodes deliver controlled electrical impulses to selected brain regions. It is used for conditions such as Parkinson disease, essential tremor, and some forms of dystonia.

### **Deep Femoral Artery**

The profunda femoris artery, a major branch of the femoral artery that supplies the deep muscles of the thigh and gives rise to perforating branches.

### **Deep Infiltrating Endometriosis**

Endometriosis that grows deeply into tissues around

---

the uterus, bowel, bladder, or other pelvic structures. It may cause severe period pain, pain with sex or bowel movements, urinary symptoms, infertility, and sometimes requires complex specialist treatment.

**Alternate Names:**

DIE; Deep Endometriosis

### Deep Neck Space Infection

A bacterial infection in one or more of the tissue spaces deep in the neck. It may cause neck swelling, severe throat pain, fever, difficulty swallowing, or breathing problems and can spread to the chest or threaten the airway.

### Deep Sleep

The deepest stage of normal non-REM sleep, when a person is difficult to awaken and the body carries out important physical restoration. Too little may contribute to daytime sleepiness and poor function.

### Deep Tendon Reflex

An automatic muscle response caused by tapping a tendon with a reflex hammer. It helps assess nerves and the spinal cord. Common examples are the knee, ankle, biceps, and triceps reflexes.

### Deep Tendon Reflexes

Involuntary monosynaptic stretch reflexes elicited by tapping a muscle tendon briskly with a reflex hammer (including the biceps, triceps, brachioradialis, patellar, and Achilles reflexes), used clinically to evaluate the structural and functional integrity of peripheral sensory and motor nerve roots and central upper motor neuron pathways.

**Alternate Names:**

DTRs; Muscle Stretch Reflexes

### Deep Vein Thrombosis

A blood clot in a deep vein, usually in the leg, caused by reduced blood flow, increased clotting tendency, or injury to the vein wall. It may cause one-sided leg swelling, pain, warmth, or discoloration, although some clots cause no symptoms. Part of the clot can break off and travel to the lungs, causing a potentially life-threatening pulmonary embolism.

**Alternate Names:**

DVT; Deep Venous Thrombosis; Deep Venous Thrombosis (DVT); Phlebothrombosis

### Deep Vein Thrombosis Ultrasound

An ultrasound examination that checks whether deep veins compress normally and whether blood flows through them. A vein that does not compress may contain a clot; results are interpreted with symptoms and other tests.

### Deep Venous Reflux

Backward flow of blood through weakened valves in deep veins. It can cause long-term leg swelling, heaviness, skin changes, pain, and ulcers from increased pressure in the veins.

### Deerfly Fever

An older name for tularemia, an infection caused by *Francisella tularensis*. It can spread through tick or deerfly bites, infected animals, contaminated water, or inhalation and may cause fever, ulcers, swollen lymph nodes, pneumonia, or severe systemic illness.

### Defaecation

The expulsion of stool from the rectum through the anus. Difficulty, pain, urgency, incontinence, bleeding, or a sustained change in bowel habit may indicate constipation, anorectal disease, neurologic dysfunction, or another condition.

### Defecation

The process of passing stool from the rectum through the anus. It involves signals from a full rectum, contraction of the bowel, and relaxation of the anal muscles, followed by voluntary control of timing.

### Defecation Syncope

Fainting during or immediately after defecation, usually a situational reflex involving straining, reduced venous return, and a temporary fall in blood pressure.

---

## Defecography

An imaging test that records the rectum and pelvic floor while a person passes a soft contrast material. It can show prolapse, a bulge, inward telescoping, or poor coordination causing difficult stool passage.

## Defective Virus

A virus missing genetic information needed to reproduce independently. It can multiply only with help from another virus or host system and may influence infection without always causing disease on its own.

## Defence Mechanisms

Mental processes that help a person manage conflict, anxiety, or distress, often without conscious awareness. They may be helpful or harmful depending on how rigidly they are used and how they affect daily life.

## Defense Mechanism

A psychological process, often unconscious, that helps a person manage conflict, anxiety, or distress; its usefulness depends on context and flexibility.

## Defense Mechanisms

Psychological processes, often outside conscious awareness, that help a person manage distress or conflict. Examples include denial, projection, repression, humor, and sublimation.

## Defense Wounds

Injuries to the hands, wrists, arms, or other areas that may occur when a person tries to protect against an attack. Their appearance and location can support a forensic interpretation but cannot prove how an injury happened by themselves.

## Defenses

The body's protective mechanisms against infection, injury, and other threats, including physical barriers and immune responses.

### Alternate Names:

Host Defenses; Immune Defenses

## Defensin

A small protective protein made by skin, lining tissues, and immune cells. It can damage microbes and is part of the body's early defense against infection.

## Deferasirox

An oral iron-chelating medicine used to reduce chronic iron overload when clinically indicated; kidney, liver, hearing, and gastrointestinal toxicity require monitoring.

## Deferiprone

An oral iron chelator used for transfusional iron overload; neutropenia or agranulocytosis is an important adverse effect requiring blood-count monitoring.

## Deferoxamine

A parenteral iron chelator used for severe or chronic iron overload and sometimes aluminum toxicity; treatment requires monitoring for ocular, auditory, renal, and infectious complications.

## Defibrillation

Delivery of an electrical shock to stop ventricular fibrillation or pulseless ventricular tachycardia during cardiac arrest. The shock can restore an organized rhythm and must be combined with high-quality CPR and emergency care.

## Defibrillator

A device that delivers an electrical shock to treat certain life-threatening abnormal heart rhythms. External defibrillators are used during cardiac arrest; implanted devices can detect and treat dangerous rhythms automatically.

## Defibrillator Implant

An implanted device that continuously monitors heart rhythm and can use pacing or an electrical shock to stop dangerous fast rhythms. It may be recommended for selected people at high risk of sudden cardiac arrest.

## Deficiency Diseases

Illnesses caused by too little of a needed nutrient



---

or other substance, such as iron, iodine, vitamin D, or vitamin B12. Symptoms depend on the missing substance.

### **Deficient Color Vision**

Alternate index label; see *Color blindness*.

### **Deformation**

An alteration in the shape or structure of a body part, tissue, or material caused by developmental forces, injury, disease, pressure, or mechanical stress.

### **Deformity**

An abnormal shape, alignment, or structure of a body part caused by development, injury, disease, pressure, or unusual growth. It may affect appearance, movement, comfort, or organ function.

### **Degarelix**

An injected hormone-blocking medicine that rapidly lowers testosterone. It is used for advanced prostate cancer; injection-site reactions, hot flushes, bone loss, and other hormone-related effects may occur.

### **Degeneration**

Progressive structural or functional deterioration of cells, tissues, or organs caused by aging, disease, injury, inherited factors, or other processes.

### **Degenerative Arthritis**

Alternate index label; see *Osteoarthritis*.

### **Degenerative Cervical Myelopathy**

Spinal-cord dysfunction caused by age-related changes in the neck that narrow the spinal canal and compress the cord. It may cause hand numbness or clumsiness, limb weakness, balance or walking problems, and sometimes bladder or bowel changes; worsening symptoms need prompt medical assessment.

#### **Alternate Names:**

Cervical Spondylotic Myelopathy; DCM

### **Degenerative Chorea**

Progressive involuntary, dance-like movements

caused by degeneration of brain circuits that control movement. The term is most closely associated with Huntington disease, which may also cause changes in thinking, mood, behavior, speech, or swallowing.

#### **Alternate Names:**

Huntington's Chorea; Huntington Disease

#### **Alternate Names:**

Degenerative Disorder; Neurodegenerative Disease

### **Degenerative Disc**

Degenerative disc disease describes age- or wear-related changes in spinal discs that may cause neck or back pain, stiffness, or nerve compression. Imaging changes can occur without symptoms; treatment usually begins with activity, exercise-based rehabilitation, and pain management.

#### **Alternate Names:**

DDD; Degenerative Disc Disease; Spondylosis

### **Degenerative Disc Disease**

Wear, drying, or cracking of the cushioning discs between spinal bones, often related to aging or injury. It may cause back or neck pain, stiffness, or nerve pressure; imaging changes can occur without symptoms.

### **Degenerative Disease**

A condition in which a body tissue or organ gradually becomes damaged or loses function. Causes include ageing, inherited problems, injury, and other disease; examples include osteoarthritis and some disorders of the brain or nerves.

### **Degenerative Disk Disease**

Alternate spelling; see *Degenerative Disc*.

### **Degenerative Disorder**

A disease in which progressive cellular or tissue deterioration causes loss of structure or function, as in neurodegenerative, joint, retinal, or muscle disorders.

### **Degenerative Joint Disease**

Gradual loss of joint cartilage with changes in nearby bone and tissues, commonly meaning osteoarthritis.

---

tis. It can cause pain, stiffness, swelling, reduced movement, and difficulty with daily activities.

**Alternate Names:**

Osteoarthritis

**Degenerative Spine Disease**

Wear-related changes in the spinal discs, joints, or bones that can cause back or neck pain, stiffness, reduced movement, or pressure on nerves. Symptoms may include leg or arm pain, numbness, weakness, or walking difficulty.

**Degenerative Spondylolisthesis**

Forward slipping of one spinal bone over another because joints and disks have worn down, sometimes causing back pain, leg pain, or nerve compression.

**Deglutition**

The coordinated process of moving food, liquid, or saliva from the mouth through the throat and esophagus into the stomach. Problems can cause choking, coughing, or food entering the airway.

**Alternate Names:**

Swallowing

**Degos Disease (Malignant Atrophic Papulosis)**

A rare occlusive small-vessel disease causing multiple porcelain-white, atrophic skin papules with a surrounding pink or violaceous rim. It may remain limited to the skin or involve the intestine, brain, or other organs; new lesions with abdominal pain, headache, or neurologic symptoms require urgent specialist assessment.

**Degranulation**

Release of stored chemicals from cells such as mast cells, white blood cells, and platelets. The chemicals help respond to injury or infection but can also cause allergy, inflammation, blood-vessel changes, or clotting.

**Degrees Fahrenheit**

A unit on the Fahrenheit temperature scale. In clinical care, body temperature may be reported

in Fahrenheit; 98.6 degrees Fahrenheit is an approximate average normal oral temperature, not a universal threshold.

**Dehiscence**

Partial or complete separation of a healing wound or surgical closure. It may involve only the skin or deeper tissues; increasing pain, drainage, or exposed tissue requires prompt medical assessment.

**DEHP**

Di(2-ethylhexyl) phthalate, a chemical added to some plastics to make them flexible. Repeated or high exposure may affect reproduction, development, or hormone function, so medical exposure is limited when possible.

**Dehydration**

Loss of more water and electrolytes than are replaced, often from vomiting, diarrhea, fever, sweating, or poor fluid intake. It can cause thirst, dry mouth, reduced urination, dizziness, fast heartbeat, abnormal electrolyte levels, or shock.

**Dehydroascorbic Acid**

The oxidized form of vitamin C. Cells can take it up and convert it back to active vitamin C, allowing it to participate in antioxidant and other chemical reactions.

**Dehydroepiandrosterone**

A hormone made mainly by the adrenal glands that the body can convert into other sex hormones. Levels change with age and illness; supplements may cause acne, unwanted hair, mood changes, or medicine interactions.

**Deinstitutionalize**

To help a person leave long-term institutional care and live in the community with suitable housing, healthcare, and support. The plan should protect safety, treatment needs, and independence.

**Deiodinase**

An enzyme that removes iodine from thyroid hor-

---

mones and changes how active they are. Different tissues use these enzymes to help control metabolism, growth, brain development, and responses to thyroid hormone.

### **Deja Vu**

A brief feeling that a new situation has happened before. Occasional episodes are common, but frequent episodes with loss of awareness, unusual movements, or confusion may occur with seizures and need assessment.

### **Dejerine-Roussy Syndrome**

Persistent burning, aching, or painful abnormal sensation on one side of the body after a stroke damages a deep brain sensory area. Normally harmless touch or temperature may become painful.

### **Delavirdine**

An older HIV medicine that blocks reverse transcriptase, an enzyme HIV needs to copy its genetic material. It was used with other HIV medicines, but resistance and drug interactions limit its use.

### **Delayed Cord Clamping**

Waiting, commonly 30–60 seconds or longer, before clamping the umbilical cord after birth. This allows extra blood to reach the newborn and may improve early circulation and iron stores; timing depends on both patients' condition.

### **Delayed Ejaculation**

Persistent difficulty or inability to ejaculate despite adequate sexual stimulation, causing distress. Medicines, nerve or hormone problems, chronic illness, psychological factors, and relationship circumstances may contribute.

### **Delayed Emergence**

Failure to wake and respond as expected after anesthesia. Causes include lingering anesthetic medicines, low body temperature, abnormal blood sugar or salts, breathing problems, stroke, or other complications and require prompt assessment.

### **Delayed Gastric Emptying**

Alternate index label; see *Gastroparesis*.

### **Delayed Graft Function**

A manifestation of acute kidney injury following renal allograft transplantation, classically defined as the requirement for dialysis within the first seven days post-transplantation, commonly attributable to prolonged cold ischemia and acute tubular necrosis.

#### **Alternate Names:**

DGF

### **Delayed Growth**

Slower-than-expected increase in a child's height, weight, or head size. Nutrition, chronic disease, hormone problems, genetic conditions, and family growth pattern can contribute; growth should be assessed over time.

### **Delayed Hemolytic Transfusion Reaction**

An extravascular hemolytic immune response occurring typically 3 to 14 days following blood transfusion, mediated by anamnestic IgG alloantibodies (frequently against Kidd or Rh antigens) targeting transfused donor erythrocytes.

#### **Alternate Names:**

DHTR

### **Delayed Hypersensitivity**

An immune reaction that develops hours or days after exposure rather than immediately. It can cause contact dermatitis, some medicine rashes, or a reaction to a tuberculosis skin test.

### **Delayed or Absent Tooth Formation**

Delayed or absent tooth formation means teeth develop or erupt later than expected or fail to form, due to local, genetic, endocrine, nutritional, or systemic causes.

### **Delayed Orgasm**

Persistent difficulty or inability to reach orgasm despite adequate stimulation, causing distress. Medicines, illness, hormonal or nerve problems,

---

anxiety, trauma, and relationship factors may contribute.

### **Delayed Puberty**

Puberty that has not begun or is not progressing at the expected age. Family pattern, poor nutrition, chronic illness, hormone disorders, or problems involving the ovaries, testes, brain, or pituitary may contribute.

### **Delayed Puberty In Boys**

Lack of testicular enlargement or other expected pubertal changes by the usual age. Causes include normal late family development, poor nutrition, chronic illness, low testosterone, or hypothalamus or pituitary disorders.

### **Delayed Puberty In Girls**

Lack of breast development or menstruation by the usual age, or puberty that stops progressing. Causes include normal late family development, low energy availability, chronic illness, ovarian failure, or hypothalamus or pituitary disorders.

### **Delayed Sleep Phase**

A body-clock pattern in which a person naturally falls asleep and wakes much later than desired. Sleep may be normal at the preferred times but interfere with school, work, or social activities.

**Alternate Names:**

DSWPD; Delayed Sleep Phase Syndrome (DSPS)

### **Delayed Sleep Phase Syndrome**

A persistent body-clock disorder in which sleep begins and ends much later than desired, despite the ability to sleep normally on that schedule. It causes difficulty meeting work, school, or social obligations.

### **Delayed Sleep-Wake Phase Sleep Disorder**

Alternate name; see *Delayed Sleep Phase Syndrome*.

### **Deletion**

Loss of a section of DNA or part of a chromosome. The effects depend on the size, location, and genes

involved; deletions may cause no problems or lead to inherited, developmental, or medical disorders.

### **Deletion Mutation**

A genetic change in which one or more DNA building blocks are missing. Its effect depends on the location and size of the deletion and whether it changes a gene or its control region.

### **Delftia**

A group of oxygen-using bacteria found mainly in soil and water. Human infection is uncommon but can occur as an opportunistic infection, especially in people with weakened immunity or medical devices.

### **Delirium**

A sudden change in attention, awareness, and thinking that fluctuates over hours or days. It usually results from illness, infection, medicines, substance use or withdrawal, dehydration, or another medical problem.

### **Delirium In Anesthesia**

Sudden confusion, poor attention, and changing awareness during recovery from anesthesia or surgery. Possible causes include medicines, infection, pain, low oxygen, dehydration, or abnormal body chemistry and require prompt assessment.

### **Delirium Superimposed On Dementia**

Delirium occurring in a person who already has dementia. The sudden, fluctuating worsening in attention and awareness differs from the person's usual cognitive impairment and may be triggered by illness, medicines, dehydration, or hospitalization.

### **Delirium Tremens**

A life-threatening form of alcohol withdrawal causing severe confusion, agitation, hallucinations, shaking, sweating, fever, fast heartbeat, and high blood pressure. It commonly begins two to four days after heavy drinking stops and requires emergency treatment.

---

## **Delirium Tremens (DTs)**

A severe, potentially fatal form of alcohol withdrawal characterized by delirium, agitation, tremor, autonomic overactivity, hallucinations, and sometimes seizures, usually beginning several days after reduction or cessation of heavy drinking.

## **Delivery**

Childbirth, ending with birth of the baby and delivery of the placenta. It may occur through the vagina, sometimes with forceps or vacuum assistance, or through cesarean surgery.

## **Delivery Of Posterior Arm**

A maneuver for shoulder dystocia in which the clinician brings the baby's back arm through the birth canal. This reduces the width of the shoulders and may help complete delivery.

## **Delivery Presentations**

The fetal body part positioned to enter the birth canal first. Common presentations are head-first, breech, and shoulder; the presentation affects delivery planning and risks.

## **Delta Agent**

An older name for hepatitis D virus, which can reproduce only when hepatitis B virus is present. Infection may cause severe acute hepatitis or worsen long-term liver disease.

## **Delta Waves**

Very slow electrical brain waves that are normal during deep non-REM sleep. Excessive delta activity while awake may indicate brain dysfunction and must be interpreted with symptoms and other tests.

### **Alternate Names:**

Delta Rhythm

## **Delta-ALA Urine Test**

A urine test measuring a substance used to make hemoglobin. Increased levels may occur with acute porphyria or lead poisoning and are interpreted with symptoms and other tests.

## **Delta-Tocopherol**

One form of vitamin E found in foods and supplements. It has antioxidant activity, but alpha-tocopherol is the main form maintained in human blood and used to assess vitamin E status.

## **Deltacoronavirus**

A group of coronaviruses identified by their genetic features. Some infect birds or mammals, and their ability to infect people or cause disease depends on the specific virus and host.

## **Deltaretrovirus**

A group of viruses that includes human T-cell leukemia viruses. They copy their RNA into DNA inside infected cells; some can persist for years and cause blood-cell disorders or cancer.

## **Deltoid**

The large, triangular muscle covering the shoulder. It helps lift the arm and stabilize the shoulder joint; injury or nerve damage can cause pain and weakness.

## **Deltoid Ligament**

A strong, fan-shaped triangular ligament complex on the medial aspect of the ankle joint, composed of superficial and deep components originating from the medial malleolus and attaching to the talus, calcaneus, and navicular bones to resist excessive eversion.

### **Alternate Names:**

Medial Collateral Ligament of the Ankle

## **Deltoid Muscle**

A triangular shoulder muscle attached to the collarbone, shoulder blade, and upper arm. Its front, middle, and back fibers move the arm forward, outward, and backward; the axillary nerve supplies it.

## **Delusion**

A firmly held false belief that remains despite clear evidence to the contrary and is not explained by the person's cultural background. It may involve

---

persecution, grandiosity, illness, or other themes.

### **Delusional Disorder**

A mental disorder with one or more persistent delusions lasting at least one month, while overall functioning is relatively preserved. Hallucinations, if present, relate closely to the delusional belief.

### **Delusional Infestation**

A fixed belief that living or inanimate pathogens are infesting the body despite no convincing medical or microbiological evidence. It may be primary or secondary to a psychiatric, neurologic, medical, medication-related, or substance-related condition; evaluation should exclude real infestation and other causes while avoiding dismissive care.

### **Demarcation**

A clear boundary between two areas or conditions, such as the line separating healthy tissue from tissue that has died. It may guide diagnosis or treatment.

### **Demeclocycline**

A tetracycline antibiotic that blocks bacterial protein production. It was also used to reduce excess water retention in some patients with inappropriate antidiuretic hormone secretion; adverse effects include sun sensitivity and kidney or liver problems.

### **Demecolcine**

A colchicine-related substance that disrupts microtubules, structures needed for cell division. It has limited medical use and can be highly toxic, causing gastrointestinal, blood-cell, nerve, or organ injury.

### **Dementia**

A lasting decline in memory, thinking, language, judgment, or other mental abilities that interferes with independent daily life. Causes include brain diseases, blood-vessel disease, infections, medicines, and some treatable medical problems.

#### **Alternate Names:**

Major Neurocognitive Disorder; Cognitive Decline

### **Dementia And Driving**

The effect of cognitive impairment on driving safety. Assessment considers memory, judgment, visuospatial skills, reaction time, medications, and local licensing requirements.

### **Dementia Behavioral and Sleep Changes**

Common behavioral disturbances and sleep-wake cycle disruptions occurring in individuals with dementia, including wandering, sundowning, aggression, agitation, and nighttime restlessness.

#### **Alternate Names:**

Behavioral and Psychological Symptoms of Dementia; BPSD; Sundowning in Dementia

### **Dementia Due To Metabolic Causes**

Cognitive impairment caused or worsened by metabolic problems such as thyroid disease, vitamin deficiency, organ failure, electrolyte disturbance, or medication toxicity; some causes are treatable.

### **Dementia with Lewy Bodies**

A progressive neurodegenerative dementia characterized pathologically by extensive intracellular alpha-synuclein inclusions (Lewy bodies) in cortical and subcortical neurons, presenting with fluctuating cognitive performance, recurrent well-formed visual hallucinations, REM sleep behavior disorder, and spontaneous parkinsonism.

#### **Alternate Names:**

DLB; Lewy Body Dementia

### **Demethylation**

Removal of a small chemical group called a methyl group from a molecule such as DNA or a protein. In DNA, this change can alter which genes are active and affect cell development or disease.

### **Demineralization**

Loss of minerals from a tissue such as tooth enamel or bone. It can weaken the tissue, causing early tooth decay or contributing to fragile bones, depending on the cause.



---

## Demulcent

A substance that coats and soothes irritated moist body linings, temporarily reducing discomfort from dryness, inflammation, or minor injury.

## Demyelinating CNS Disease

A disorder in which the protective covering of nerves in the brain, spinal cord, or vision pathways is damaged. Causes include immune disease, infection, inherited conditions, toxins, and metabolic problems.

## Demyelinating Disease

A disease in which the protective covering around nerve fibers is damaged. Nerve signals then slow or fail, causing problems such as weakness, numbness, vision changes, or poor coordination.

## Demyelination

Loss or damage of myelin, the protective coating that helps nerve fibers send signals quickly. It can disrupt movement, sensation, vision, balance, or thinking and may result from immune, infectious, inherited, or toxic disorders.

## Denaturant

A substance added to another product to change its taste, smell, or properties, often to make it unsuitable for drinking. Some denaturants are poisonous or irritating, so health effects depend on the product and amount swallowed or contacted.

## Dendrite

A branching extension of a nerve cell that receives signals from other nerve cells or sensory receptors and carries those signals toward the cell body.

## Dendrites

Branching extensions of nerve cells that receive signals from other neurons or sensory receptors and carry those messages toward the cell body.

## Dendritic Cell

An immune cell that captures foreign material and shows pieces of it to other immune cells. This helps

start a targeted immune response against infections and abnormal cells.

## Dendritic Cells

Immune cells that collect material from germs or damaged cells and show identifying pieces to T cells. This helps the immune system start a targeted response and develop longer-lasting protection.

## Dendritic Spine

A tiny branch on a nerve cell that receives signals from other nerve cells. Its shape and number can change with learning, development, and some brain diseases.

## Dendritic Ulcer

A branching epithelial ulcer of the cornea, classically caused by herpes simplex virus. It may cause a red eye, pain, light sensitivity, tearing, or reduced vision and requires ophthalmic assessment.

## Denervate

To remove or interrupt the nerve supply to a tissue or organ, either deliberately during treatment or because of injury or disease. The change may reduce movement, sensation, pain, or automatic organ function.

## Denervation

Denervation is loss of nerve supply to a muscle or tissue, causing weakness, atrophy, altered sensation, or autonomic dysfunction.

## Dengue

A mosquito-borne viral infection spread mainly by daytime-biting *Aedes* mosquitoes. It commonly causes sudden fever, severe headache, pain behind the eyes, muscle and joint aches, nausea, and sometimes a rash. Some people develop severe dengue with bleeding, fluid leakage, shock, or organ injury. NS1, PCR, or antibody tests can support the diagnosis depending on the timing, but results must be interpreted with symptoms and exposure history. Care focuses on careful fluids and avoiding aspirin or ibuprofen unless a clinician advises otherwise.

**Alternate Names:**

---

Breakbone Fever; Dengue Fever; DENV Infection

## Dengue Fever

Alternate name; see *Dengue*.

## Dengue Hemorrhagic Fever

A severe form of dengue infection in which blood vessels leak and bleeding or shock may develop. Warning signs include severe abdominal pain, repeated vomiting, bleeding, extreme tiredness, cold skin, or reduced urine; urgent hospital care and careful fluids are needed.

## Dengue Shock Syndrome

Life-threatening circulatory shock caused by severe dengue with plasma leakage and reduced effective blood volume, often accompanied by bleeding or organ dysfunction.

## Dengue Virus

The virus that causes dengue fever and is spread mainly by infected *Aedes* mosquitoes. Four major types exist; protection after infection is strongest against the same type and does not reliably prevent later infection by other types.

## Denial

A psychological response in which a person cannot or will not accept a distressing fact, diagnosis, or situation. It may briefly reduce distress but can delay needed decisions or treatment when persistent.

## Denileukin Difitox

A laboratory-made protein combining part of interleukin-2 with diphtheria toxin. It was used for selected cutaneous T-cell lymphomas by delivering the toxin to cells bearing the interleukin-2 receptor; toxicity limits use.

## Dennie-Morgan Fold

A prominent, symmetrical single or double skin crease or infraorbital fold immediately below the margin of the lower eyelid, recognized as a characteristic cutaneous hallmark of atopic dermatitis and allergic diathesis.

## Alternate Names:

Dennie-Morgan Line; Dennie Fold

## Denominator

The number below the line in a fraction. In a medical rate or proportion, it represents the total population or group being compared with the number above it.

## Denosumab

A medicine that blocks a protein needed to activate bone-destroying cells. It is used for osteoporosis and some cancer-related bone problems; stopping it without follow-up can cause rapid bone loss and spine fractures.

## Denosumab For Osteoporosis

An injection given every six months that blocks a signal activating bone-destroying cells. It lowers fracture risk, including at the spine and hip, but calcium must be monitored and another treatment is needed when it is stopped.

## Dens

The tooth-shaped projection of the second neck vertebra. It fits into the first neck vertebra and acts as a pivot that allows the head to turn.

## Dens Evaginatus

An extra bump or cusp on a tooth, most often a premolar. It may contain an extension of the tooth's nerve and can wear, break, or develop decay and pain.

## Dens Invaginatus

An inward folding of a tooth's enamel and dentin that may reach the nerve. The pocket can trap bacteria and food, increasing the risk of early decay, nerve infection, and tooth loss.

## Densitometry

Measurement of how dense a tissue is, most often the mineral content of bone. A bone-density scan helps diagnose osteoporosis, estimate fracture risk, and monitor treatment.

---

## Density

The amount of mass in a given volume. In medicine, density measurements help describe bone mineral content, tissue composition, fluid collections, or findings on imaging.

## Dent Disease

An inherited disorder of the kidney's filtering tubes, usually affecting boys. It causes loss of small proteins in urine, excess urinary calcium, kidney stones or calcium deposits, and sometimes progressive kidney failure.

## Dental Abscess

A pocket of pus caused by infection in a tooth or surrounding gum. It may cause severe toothache, swelling, fever, or difficulty opening the mouth; spreading facial or neck swelling needs urgent care.

## Dental Age

An estimate of a child's developmental age based on tooth formation, hardening, and eruption. It can help assess growth but does not exactly determine the child's actual age.

## Dental Alloy

A mixture of metals used to make dental fillings, crowns, bridges, dentures, orthodontic appliances, or other restorations. Its strength, appearance, corrosion resistance, and possible allergies depend on its composition.

## Dental Alveolus

The part of the upper or lower jawbone that forms a tooth socket and holds the tooth root. It can shrink after tooth loss or severe gum disease.

## Dental Amalgam

A durable filling material made by mixing liquid mercury with powdered metals, mainly silver, tin, and copper. It is strong but has a silver color and may be unsuitable in some situations.

## Dental Anesthesia

Use of local anesthetic, sedation, or occasionally

general anesthesia to prevent pain or reduce anxiety during dental treatment. The method depends on the procedure, health, age, and patient preference.

## Dental Anxiety

Fear or distress about dental treatment that may cause avoidance, delayed care, or difficulty during appointments. Reassurance, communication, pain control, and behavioral or medicine-based support may help.

## Dental Arch

The curved arrangement of teeth and the bone supporting them in the upper or lower jaw. Its shape affects tooth alignment, bite, speech, chewing, and space for dental restorations.

## Dental Articulator

A device that holds models of the upper and lower teeth and imitates jaw movements. It helps dental professionals design and adjust crowns, bridges, dentures, and other restorations so they fit the bite.

## Dental Avulsion

Complete displacement of a tooth from its socket, usually caused by an injury. A permanent tooth that is knocked out needs immediate dental care; hold it by the crown, keep it moist, and do not scrub the root.

## Dental Bone Graft

An oral and maxillofacial surgical procedure involving the placement of autologous bone, allograft, xenograft, or synthetic osteoconductive matrix into the maxilla or mandible to reconstruct alveolar ridge deficiencies or augment the maxillary sinus prior to dental implant placement.

### Alternate Names:

Dental Bone Graft; Alveolar Ridge Augmentation; Maxillofacial Bone Grafting

## Dental Braces

Fixed or removable appliances that apply gentle, controlled force to move teeth and improve their alignment or bite. Treatment requires cleaning and follow-up; possible problems include soreness, decay,

---

gum disease, and root shortening.

**Alternate Names:**

Orthodontic Braces; Orthodontic Appliances

**Dental Bridge**

A fixed dental appliance that replaces one or more missing teeth by attaching artificial teeth to nearby teeth or implants. It can improve chewing and appearance but requires healthy supports and careful cleaning.

**Dental Calculus**

Hardened plaque formed when minerals in saliva or gum fluid collect on teeth. It cannot be removed reliably by brushing alone and can contribute to gum inflammation and disease.

**Dental Caries**

Tooth decay caused when bacteria use sugars and produce acids that remove minerals from enamel and dentin. Untreated decay can cause sensitivity, pain, infection, and tooth loss.

**Dental Cavities**

Permanent holes or weakened areas caused when plaque bacteria use sugars and make acids that dissolve a tooth. Decay can spread from enamel into the deeper tooth, causing sensitivity, pain, infection, or tooth loss.

**Dental Cement**

A dental material used to attach crowns, bridges, or orthodontic bands, fill small spaces, or protect exposed tooth tissue. Its strength and use depend on the material and treatment.

**Dental Ceramic**

A tooth-colored material used for crowns, veneers, bridges, fillings, and implant restorations. It provides good appearance and resistance to wear, but its strength and suitability vary by type and location.

**Dental Clasp**

A part of a removable partial denture that rests

against or around supporting teeth to help keep the denture in place. Poor fit or excessive pressure can irritate teeth or gums.

**Dental Crown**

A custom-made covering placed over a damaged or weakened tooth to restore its shape, strength, chewing function, or appearance. It may be made from ceramic, metal, resin, or a combination of materials.

**Dental Crowns**

Custom-made coverings placed over most or all of a damaged or weakened tooth. A crown restores shape, strength, chewing function, and appearance after decay, fracture, or root-canal treatment.

**Dental Deposit**

Material that collects on a tooth, such as plaque, hardened tartar, food debris, or stain. Some deposits irritate the gums or support decay and may require professional cleaning.

**Dental Disease**

A disorder of the teeth, gums, jaw, or supporting tissues, including decay, gum disease, infection, and tooth loss. Fluoride, daily cleaning, limited frequent sugar exposure, and dental visits reduce risk.

**Dental Drill**

A powered dental handpiece with a rotating bur, driven by air or an electric motor, used to remove decayed tooth structure, shape teeth, prepare teeth for fillings or crowns, and perform other dental procedures. Different burs and speeds are selected for the material and procedure.

**Alternate Names:**

Dental Handpiece

**Dental Dysplasia**

Abnormal development of a tooth or its supporting tissue. It may change the tooth's shape, color, strength, position, or eruption and can increase the risk of wear, decay, or infection.

---

## Dental Erosion

Gradual loss of tooth enamel caused by acid rather than bacteria. Acidic foods or drinks, reflux, vomiting, and some medicines may contribute, causing sensitivity or worn teeth; reducing acid exposure helps prevent progression.

## Dental Extraction

Removal of a tooth from its jaw socket because of severe decay, infection, gum disease, impaction, crowding, trauma, or treatment planning. Risks include bleeding, infection, nerve injury, fracture, and dry socket.

## Dental Fillings

Materials placed in a tooth after decay or damage is removed to restore its shape and function. Resin, glass ionomer, metal amalgam, or other materials may be used depending on the tooth and clinical situation.

## Dental Fissure

A narrow natural groove or developmental defect in a tooth, especially on a chewing surface. Food and bacteria can collect there, increasing the risk of decay.

## Dental Floss

A thin thread used to remove plaque and food between teeth and just below the gumline. Daily cleaning between teeth helps prevent gum inflammation but does not replace brushing or dental care.

## Dental Fluorosis

Changes in tooth enamel caused by too much fluoride while the teeth are developing. It may appear as white lines or patches and, in more severe cases, brown discoloration or pits.

## Dental Hygiene

Daily care of the teeth, gums, tongue, and mouth, including brushing with fluoride toothpaste and cleaning between teeth. It removes plaque and helps prevent decay, gum disease, and bad breath.

## Dental Implant

A small biocompatible fixture placed in the jawbone to support a replacement tooth, bridge, or denture. Bone may grow around it for stability; risks include infection, poor healing, nerve injury, and implant failure.

## Dental Impression Material

A material placed in a tray to make an accurate mold of the teeth and mouth tissues. The mold helps make crowns, dentures, orthodontic appliances, restorations, or study models.

## Dental Injuries

Damage to teeth, gums, jawbone, or mouth tissues caused by a fall, blow, sports injury, accident, or biting force. Prompt assessment may save displaced or knocked-out teeth; severe bleeding or head injury is an emergency.

### Alternate Names:

Dental Trauma; Traumatic Dental Injury

## Dental Inlay

A custom-made filling fitted inside the grooves of a damaged tooth and bonded in place. It restores areas too large for a simple filling but does not cover the tooth's cusps like an onlay or crown.

## Dental Material

A substance used in dental care, such as composite, amalgam, ceramic, metal, cement, bonding material, impression material, or implant material. Choice depends on the treatment, location, strength, appearance, and patient factors.

## Dental Occlusion

The way the upper and lower teeth meet when the mouth closes or the jaw moves. An abnormal bite may contribute to difficulty chewing, tooth wear, jaw pain, or muscle strain.

## Dental Overjet

The horizontal distance between the upper and lower front teeth when the teeth are closed. A large overjet may increase the risk of trauma to the upper front

---

teeth and affect biting.

### **Dental Papilla**

Soft embryonic tissue beneath a developing tooth. Its cells form the dentin and living pulp, which contains the tooth's nerves and blood vessels.

### **Dental Pellicle**

A thin protective film of salivary proteins that forms on tooth enamel soon after cleaning. Bacteria can attach to it and build dental plaque.

### **Dental Plaque**

A sticky film of bacteria, saliva, and food material on teeth and along the gums. If not removed by brushing and cleaning between teeth, it can cause decay, gum inflammation, and gum disease.

### **Dental Polishing**

Smoothing restorative surfaces to reduce plaque accumulation. Essential for longevity.

#### **Alternate Names:**

Dental Polishing; Coronal Polish

### **Dental Porcelain**

A tooth-colored ceramic used for crowns, veneers, bridges, and other restorations. It provides a natural appearance but may chip or fracture under heavy force.

### **Dental Precision Attachment**

A precision mechanical or magnetic interlocking device used to secure, stabilize, and retain a removable partial denture or overdenture to retained tooth roots, natural abutments, or dental implants.

#### **Alternate Names:**

Denture Attachment; Precision Attachment; Overdenture Attachment; Dental Attachment

### **Dental Procedure**

An examination, preventive service, or treatment involving the teeth, gums, jaw, or mouth, such as cleaning, filling, extraction, root-canal treatment, or restoration. Benefits and risks depend on the

procedure.

### **Dental Prosthesis**

An artificial replacement for one or more missing teeth or nearby mouth tissues, such as a denture, bridge, crown, or implant-supported restoration. It restores appearance, speech, and chewing.

### **Dental Public Health**

Efforts to prevent and control oral disease in communities. They include fluoride programs, oral-health education, screening, disease monitoring, access to dental care, and policies that reduce differences in oral health.

### **Dental Pulp**

The living soft tissue at the center of a tooth. It contains nerves, blood vessels, and connective tissue and can become inflamed or infected when decay or injury reaches it.

### **Dental Pulp Necrosis**

Death of the soft tissue inside a tooth, usually after untreated inflammation, deep decay, trauma, or loss of blood supply. The tooth may become discolored and develop infection around its root.

### **Dental Pulpitis**

Inflammation of the soft tissue inside a tooth, commonly caused by decay, trauma, a crack, or dental treatment. Brief pain may be reversible; spontaneous or lingering pain may require root-canal treatment or extraction.

### **Dental Radiographs**

X-ray images of the teeth and jaws used to find decay, bone loss, infection, impacted teeth, fractures, and developmental problems. Types include bitewing, periapical, occlusal, and panoramic views.

### **Dental Radiography**

Use of X-rays to examine teeth and jaws. Bitewing, periapical, panoramic, and three-dimensional scans answer different questions; exposure is kept as low as reasonably achievable while obtaining useful images.



---

## Dental Restorative Materials

Materials used to repair teeth, including composite resin, glass ionomer, amalgam, ceramic, and metal alloys. Selection depends on tooth location, size of damage, strength, appearance, moisture control, cost, and patient needs.

## Dental Root

The part of a tooth below the gum that anchors it in the jawbone. It contains a canal with nerves and blood vessels and is covered by cementum and supported by a ligament. Infection, decay, cracks, or loss of bone around the root can cause pain and tooth loosening.

## Dental Sac

A layer of tissue surrounding a developing tooth. It forms the tissues that support the tooth, including the periodontal ligament, cementum, and part of the surrounding jawbone.

## Dental Screening

A check of the teeth, gums, and mouth to look for decay, gum disease, oral lesions, or risk factors before serious symptoms develop. Screening does not replace a complete dental examination.

## Dental Sealant

A thin protective coating placed in the grooves of teeth, especially molars. It blocks food and bacteria from collecting in deep pits and lowers the risk of decay.

## Dental Sealants

Thin resin or glass-ionomer coatings placed in the pits and grooves of teeth, especially molars. They reduce plaque and food retention and lower the risk of decay but do not replace brushing or fluoride.

## Dental Sleep Apnea Therapy

Alternate name; see *Oral appliance therapy*.

## Dental Splint

Fixation of mobile teeth or fractured tooth. Wire and composite, orthodontic bracket splint. A dental

device or wire used to stabilize loose teeth or a fractured tooth while healing occurs.

### Alternate Names:

Occlusal Guard; Dental Nightguard; Bite Splint

## Dental Trauma

Alternate name; see *Dental Injuries*.

## Dental Traumatology

The prevention, assessment, and treatment of injuries to teeth and surrounding mouth tissues. Injuries may involve cracks, fractures, displacement, root damage, or complete tooth loss and often need prompt care.

## Dental X-Rays

X-ray images of teeth and jaws used to detect decay, bone loss, infection, impacted teeth, fractures, developmental problems, and other disease not visible during an examination.

## Dentate Gyrus

A curved part of the hippocampus that helps process incoming information and form memories. It also produces some new nerve cells during adulthood.

## Dentate Line

A zigzag transverse anatomical boundary in the anal canal separating the upper two-thirds (endodermal origin, simple columnar epithelium, autonomic visceral innervation) from the lower one-third (ectodermal origin, stratified squamous epithelium, somatic pudendal innervation).

### Alternate Names:

Pectinate Line

## Dentatorubral-Pallidoluysian Atrophy

A rare inherited brain disorder caused by an extra-long repeated section of the ATN1 gene. Symptoms vary with age and may include seizures, poor coordination, involuntary movements, behavior changes, and worsening thinking ability.

### Alternate Names:

DRPLA; Haw River Syndrome; Naito-Oyanagi Disease

---

## Dentifrice

A preparation, such as toothpaste or tooth powder, used to clean teeth. Fluoride dentifrice strengthens enamel and helps prevent tooth decay when used regularly.

## Dentigerous Cyst

A fluid-filled sac that forms around the crown of an unerupted tooth, usually in the jaw. It may cause swelling, move nearby teeth, or damage bone if it becomes large or infected.

## Dentin

The hard tissue beneath tooth enamel that forms most of the tooth. Tiny channels within it can transmit sensitivity, especially when dentin is exposed by decay, wear, gum recession, or injury.

## Dentin Hypersensitivity

Alternate term; see *Dentin Sensitivity*.

## Dentin Sensitivity

Brief, sharp tooth pain caused by cold, heat, sweet foods, touch, or air reaching exposed dentin. Decay, worn enamel, gum recession, cracks, or aggressive brushing can expose it.

## Dentinogenesis

Formation of dentin by specialized cells during tooth development and repair. Abnormal dentin formation may produce weak, discolored, sensitive teeth that wear or fracture easily.

## Dentinogenesis Imperfecta

An inherited disorder in which dentin forms abnormally, causing teeth that may look blue-gray or amber and break, wear, or discolor easily. Some forms occur with fragile bones.

## Dentist

A licensed healthcare professional who examines, prevents, diagnoses, and treats diseases and injuries of the teeth, gums, jaw, and mouth.

## Dentistry

The healthcare field concerned with preventing, diagnosing, and treating disorders and injuries of the teeth, gums, jaw, and mouth, including restoration, replacement, and oral health education.

## Dentition

The teeth present in the mouth, including their number, arrangement, development, and eruption. Children usually have primary teeth that are replaced by permanent teeth; disease, injury, or development can alter either set.

## Denture

A removable replacement for some or all missing teeth and nearby mouth tissues. Partial dentures replace selected teeth; complete dentures replace all teeth in one jaw. Correct fit, cleaning, and review help prevent sores and infection.

## Denture Base

The part of a denture that rests on the gums and mouth tissues and supports the artificial teeth. Its shape and fit distribute chewing pressure and help prevent soreness.

## Denture Problems

Difficulties caused by a removable denture, including poor fit, looseness, pain, sores, breakage, difficulty chewing or speaking, or infection. A dental review may correct the cause or determine whether replacement is needed.

## Dentures

Removable appliances that replace some or all missing teeth and nearby gum tissue. They can improve chewing, speech, and appearance but need proper fitting, daily cleaning, gradual adjustment, and regular dental review.

## Denturist

A dental professional authorized in some regions to make, fit, adjust, and repair removable dentures within the local scope of practice. Dentists may also assess oral disease and provide related treatment.

---

## Denver Developmental Screening Test

A screening questionnaire and task-based assessment for young children that checks personal-social, fine-motor, language, and gross-motor skills. It identifies possible developmental delay but does not diagnose its cause.

## Denys-Drash Syndrome

A rare genetic disorder caused by changes in the *WT1* gene. It can cause severe kidney disease, differences in sex development, and a high risk of Wilms kidney tumor in childhood.

## Deoxycoformycin

An anticancer medicine that blocks an enzyme needed by some cells to make DNA. It was used for selected blood cancers and can suppress bone marrow, weaken immunity, and increase the risk of serious infections.

## Deoxycytidine

A DNA building block made of the base cytosine joined to the sugar deoxyribose. Cells use it to make and repair DNA.

## Deoxycytidine Kinase

An enzyme that chemically activates deoxycytidine and some similar medicines. The activated medicines can interfere with the DNA of cancer cells or viruses and stop them from multiplying.

## Deoxyglucose

A molecule resembling glucose that is used mainly in research and PET imaging. When labeled with a radioactive form of fluorine, it can show tissues using unusually large amounts of glucose, including many tumors and areas of inflammation.

## Deoxyguanosine

A DNA building block made of the base guanine joined to the sugar deoxyribose. Cells use it to make and repair DNA.

## Deoxyhemoglobin

Hemoglobin that is not carrying oxygen. A high

amount in blood near the skin can make the lips, skin, or nail beds appear blue, a finding called cyanosis.

## Deoxyribonuclease

An enzyme that breaks DNA into smaller pieces. An inhaled laboratory-made form can thin thick mucus in some people with cystic fibrosis by breaking down DNA released from damaged cells.

## Deoxyribonucleic Acid

DNA, the molecule that stores hereditary information in cells. Its sequence helps instruct protein production and cell control; changes may be inherited or acquired during life and can affect health.

### Alternate Names:

DNA

## Deoxyribonucleosides

Molecules made of one of the four DNA bases joined to the sugar deoxyribose. Cells use them as starting materials when making or repairing DNA.

## Deoxyribonucleotides

DNA building blocks made of a base, deoxyribose sugar, and phosphate groups. The four forms—dATP, dGTP, dCTP, and dTTP—are assembled into DNA strands.

## Dependence

Physical adaptation to repeated exposure to a medicine or substance, so that reducing or stopping it can cause withdrawal symptoms. Dependence can occur with or without addiction.

## Dependent Personality Disorder

A long-term pattern of excessive need to be cared for, difficulty making decisions independently, fear of separation, and submissive behavior that causes distress or impairs relationships and daily life.

## Dependovirus

A group of small viruses that usually need another virus, such as an adenovirus or herpesvirus, to reproduce. They are generally not known to cause

---

serious disease in healthy people.

### **Depersonalization**

A feeling of being detached from or observing one's own body, thoughts, or emotions as if from outside. It can occur with anxiety, trauma, panic, substance use, or other mental health conditions.

### **Depersonalization-Derealization Disorder**

A disorder causing repeated or persistent feelings that one's body, thoughts, emotions, or surroundings are unreal or distant, while the person knows these feelings are not literally true. Symptoms cause significant distress or impairment.

### **Dephosphorylation**

Removal of a phosphate group from a molecule. This can switch proteins and other cell components on or off and change how cells communicate, use energy, grow, or respond to signals.

### **Depigmentation**

Loss or reduction of normal pigment, especially melanin in the skin, hair, or mucous membranes. It may result from vitiligo, inflammation, injury, infection, chemical exposure, or treatment.

### **Depilation**

Removal of hair at or near the skin surface by shaving, cutting, waxing, or a chemical product. It differs from epilation, which removes hair together with its root.

### **Depilatory Poisoning**

Harmful exposure to a hair-removal product, usually by swallowing or contact with the eyes, skin, or airways. Ingredients may cause irritation, chemical burns, vomiting, or breathing problems.

### **Depolarization**

A change that makes the inside of a cell less electrically negative. In nerve and muscle cells, it can start or transmit an electrical signal that produces movement or sensation.

### **Depolarizing Neuromuscular Blocker**

A paralytic agent, exemplified exclusively by succinylcholine, that acts as an acetylcholine receptor agonist at the neuromuscular junction, inducing persistent membrane depolarization, transient muscle fasciculations, and subsequent flaccid motor paralysis.

#### **Alternate Names:**

Depolarizing Muscle Relaxant

### **Depot Implant**

A small device placed in the body that releases medicine slowly over an extended period. It can provide steady treatment but may require a procedure for insertion and removal.

### **Deprescribing**

A planned, supervised reduction or stopping of a medicine when its harms, burden, or limited benefit outweigh expected benefit. The plan considers withdrawal, return of illness, safer alternatives, and patient goals.

### **Depressed Fracture**

A break in which bone is pushed inward, most often in the skull. It can injure the brain or other underlying tissue and requires urgent assessment, especially after head injury or with neurologic symptoms.

### **Depression**

A group of mental health disorders involving a persistent sad, empty, or irritable mood, or a loss of interest or pleasure in activities. It is more than the short-lived sadness or low mood that most people experience at times. Other symptoms may include changes in sleep, appetite, energy, concentration, movement, self-worth, or hope, as well as physical aches or digestive problems. Symptoms cause significant distress or interfere with daily life and may include thoughts of death or suicide. Major depression involves symptoms most of the day, nearly every day, for at least two weeks; persistent depressive disorder involves less severe symptoms lasting at least two years.

#### **Alternate Names:**

Major Depressive Disorder (MDD); Unipolar De-

---

pression

### Depression Screening

Use of questions or a validated questionnaire to identify possible depression. A positive screen is not a diagnosis; a clinician must assess symptoms, duration, impairment, medical causes, treatment needs, and safety.

### Depressive Disorder

A mood disorder involving persistent low mood, loss of interest, or both, with possible changes in sleep, appetite, energy, concentration, movement, or self-worth. Diagnosis depends on the pattern, duration, severity, and functional impact of symptoms.

### Depressive Illness

Broad, non-diagnostic term; see *Depressive Disorder*.

### Depth Perception

The ability to judge how far away objects are and where they are in three-dimensional space. It depends mainly on using both eyes together and also on movement, size, shading, and perspective cues.

### Depurination

Loss of the DNA bases adenine or guanine, leaving a gap in the DNA strand. If the gap is not repaired correctly, it can cause a genetic mutation.

### Derbyshire Neck

Older term; see *Goiter*.

### Dercum Disease (Adiposis Dolorosa)

Alternate names; see *Dercum's Disease*.

### Dercum's Disease

A rare disorder characterized by multiple painful fatty growths, often with obesity, fatigue, and widespread tenderness. The cause is uncertain, and diagnosis is based on symptoms and examination after other causes are excluded.

### Derealization

A feeling that the outside world is unreal, distant, dreamlike, or visually changed, while the person knows it is not literally unreal. It may occur with anxiety, panic, trauma, dissociation, or substance use.

### Dermabrasion

A procedure that uses a rapidly rotating instrument to remove outer skin layers. It may improve selected scars, wrinkles, sun damage, or lesions but can cause pain, pigment changes, infection, or scarring.

### Dermal

Relating to the dermis, the skin layer beneath the epidermis. The dermis contains connective tissue, blood vessels, nerves, hair follicles, and sweat glands that support and nourish the skin.

### Dermal Fillers

Injectable materials placed under or within the skin to restore volume or reduce wrinkles or scars. Common effects include swelling and bruising; infection, lumps, skin damage, or rare blockage of a blood vessel can be serious.

### Dermal Melanocytosis

A usually harmless blue-gray or slate-colored skin patch caused by pigment-producing cells located deep in the skin. It is often present at birth and commonly affects the lower back or buttocks.

### Dermal Papilla

A small structure at the base of a hair follicle that supplies signals and nutrients to developing hair cells. Its activity helps control hair growth and the hair-growth cycle.

### Dermal Structure

The dermis has a thin superficial layer and a deeper supportive layer. It contains collagen, elastic fibers, blood and lymph vessels, nerves, hair follicles, sweat glands, and oil glands.

---

## **Dermatan Sulfate**

A long chain of sugar molecules found in skin, connective tissue, blood vessels, and heart valves. It helps organize tissues and can influence wound healing and blood clotting.

## **Dermatitis**

Inflammation of the skin causing some combination of itching, redness, swelling, scaling, dryness, or blisters. It may result from irritation, allergy, infection, medicines, genetics, or another disease.

## **Dermatitis Herpetiformis**

A long-lasting, intensely itchy rash with groups of small bumps or blisters, commonly on the elbows, knees, buttocks, or scalp. It is strongly linked to celiac disease and usually improves with a strict gluten-free diet.

### **Alternate Names:**

DH; Duhring-Brocq Disease

## **Dermatofibroma**

A common harmless, firm skin nodule, often on the legs. It may be brown or pink and often dimples when pinched. It usually needs no treatment, but a changing or bleeding lesion should be examined.

## **Dermatofibrosarcoma**

A rare, slow-growing cancer of connective tissue in the skin. It usually begins as a firm plaque or nodule and can grow into nearby tissue, but it rarely spreads to distant organs.

## **Dermatofibrosarcoma Protuberans**

A rare cancer arising from connective tissue in the skin. It often appears as a slowly enlarging firm plaque or nodule, commonly on the trunk; it can recur locally but rarely spreads. Complete surgical removal is the usual treatment.

### **Alternate Names:**

DFSP

## **Dermatofibrosarcoma Protuberans Of The Breast**

A rare, usually slow-growing skin cancer involv-

ing the breast skin or nearby superficial tissue. It can extend into surrounding tissue and recur if incompletely removed; treatment aims for complete surgical removal.

## **Dermatoglyphics**

The study of ridge patterns on the fingers, palms, toes, and soles. Patterns help with identification and may differ in some genetic or developmental conditions, but they cannot diagnose a condition by themselves.

## **Dermatographia**

A form of hives in which light scratching, rubbing, or pressure causes raised, red, itchy lines within minutes. It is usually harmless, though symptoms may be troublesome.

## **Dermatographic Urticaria**

A form of hives in which scratching, rubbing, or pressure causes raised, itchy lines on the skin within minutes. The reaction results from release of histamine by skin immune cells.

## **Dermatographism**

Hives that appear as raised linear welts within minutes after the skin is stroked or scratched. They result from an exaggerated release of histamine and may occur with other allergic or hive disorders.

## **Dermatologic**

Relating to the skin, hair, nails, or medical care for these structures. Dermatologic problems include rashes, infections, wounds, hair loss, nail changes, acne, and skin cancers.

## **Dermatologic Disorder**

A disease affecting the skin, hair, nails, or skin glands. It may cause rash, scaling, pigment change, lumps, ulcers, pain, itching, hair loss, or abnormal nail growth.

## **Dermatologic Manifestations of Hematologic Disease**

Skin, hair, nail, or mucosal findings caused by a



---

blood or bone-marrow disorder, its treatment, or an associated infection. Examples include pallor, petechiae, purpura, bruising, pruritus, ulcers, nodules, and infiltrative lesions; new or unexplained findings may require blood tests and biopsy.

### **Dermatologist**

A physician who specializes in diagnosing and treating disorders of the skin, hair, nails, and related mucous membranes, including infections, inflammatory diseases, tumors, and cosmetic concerns.

### **Dermatology**

The medical specialty concerned with preventing, diagnosing, and treating diseases and injuries of the skin, hair, nails, and related mucous membranes.

### **Dermatomal**

Following or relating to a dermatome—a strip of skin supplied by sensory fibers from a single spinal nerve root. Dermatomal patterns can help localize nerve-root disease and are characteristic of some conditions, such as herpes zoster.

### **Dermatome**

Either a surgical tool that removes a thin layer of skin for a graft or a strip of skin supplied by one spinal nerve. The meaning depends on whether the term is being used in surgery or nerve anatomy.

### **Dermatomycosis**

A fungal infection of the skin, hair, or nails. It may cause scaling, itching, discoloration, cracking, or hair loss; treatment depends on the site and may require a topical or oral antifungal medicine.

### **Dermatomyositis**

An autoimmune disease that causes muscle inflammation and often weakness, especially in muscles near the hips and shoulders, along with skin changes such as a purple rash around the eyes or raised patches over the knuckles. It may affect swallowing or breathing muscles. In adults, it is associated with an increased risk of cancer, particularly around the time the disease begins.

### **Dermatopathology**

The study of skin disease by examining tissue samples with a microscope and related laboratory methods. It helps distinguish inflammatory, infectious, harmless, and cancerous skin disorders.

### **Dermatophyte**

A fungus that grows in keratin, the tough protein in skin, hair, and nails. It causes tinea infections such as athlete's foot, ringworm, and fungal nail disease.

### **Dermatophytosis**

A superficial fungal infection of the skin, hair, or nails. It can cause ring-shaped rashes, scaling, itching, cracked skin, or thickened discolored nails.

### **Dermatoplasty**

Surgical repair or reconstruction of damaged skin, often using tissue rearrangement or a skin graft. It may restore coverage after injury, burns, removal of a lesion, or another operation.

### **Dermatoscope**

A handheld optical instrument that magnifies and illuminates skin lesions, often with polarized or nonpolarized light. It reveals structures below the skin surface and supports clinical assessment of pigmented and other lesions.

### **Dermatosis**

Any disease or disorder of the skin, including inflammatory, infectious, allergic, inherited, or tumor-related conditions. It may cause rash, itching, scaling, pigment changes, lumps, ulcers, or pain.

### **Dermis**

The strong, flexible skin layer beneath the epidermis. It contains collagen and elastic fibers, blood vessels, nerves, hair follicles, and sweat and oil glands that support and nourish the skin.

### **Dermoid**

A usually benign cyst or tumor containing tissues such as skin, hair, oil glands, fat, or teeth. It may occur in the ovary or other sites and can cause

---

pressure, rupture, infection, or twisting of an organ.

### **Dermoid Cyst**

A benign congenital choristoma resulting from the entrapment of ectodermal and mesodermal elements along embryonic suture lines during fetal development. The cyst wall is lined by keratinized stratified squamous epithelium containing dermal appendages such as hair follicles, sebaceous glands, and sweat glands, filled with oily keratin debris. In ophthalmology and pediatric surgery, it is the most common congenital orbital mass in children, typically presenting as a painless, firm, smooth, subcutaneous nodule fixed to the underlying bone, most frequently at the superolateral orbital rim along the frontozygomatic suture. A key surgical principle is complete intact en bloc excision; rupture of the cyst wall releases irritant lipid contents that trigger intense, painful granulomatous foreign-body inflammation. The term also encompasses mature cystic teratomas of the ovary containing hair, teeth, and sebaceous material.

#### **Alternate Names:**

Orbital Dermoid Cyst; Periorbital Dermoid; Benign Cystic Teratoma; Epidermoid and Dermoid Choristoma

### **Dermoid Cyst Of The Ovary**

A usually benign ovarian cyst that can contain skin, hair, fat, glands, or teeth. It may cause pain, pressure, rupture, or twisting of the ovary and is assessed with examination and imaging.

### **Dermoscopy**

Examination of a skin lesion with magnification and special lighting that reveals structures not visible to the unaided eye. It helps assess pigmented and other lesions but does not replace biopsy when cancer is suspected.

### **Descemet Membrane Endothelial Keratoplasty**

An advanced partial-thickness corneal transplantation procedure in which only the diseased recipient corneal endothelium and Descemet membrane are stripped and replaced by an isolated donor Descemet

membrane-endothelium sheet. It offers rapid visual recovery and very low allograft rejection rates in corneal endothelial disorders such as Fuchs endothelial dystrophy.

#### **Alternate Names:**

DMEK; Endothelial Keratoplasty

### **Descending Colon**

The left-sided part of the large intestine, extending downward from the transverse colon toward the sigmoid colon. It absorbs water and salts and helps compact stool before it moves toward the rectum.

### **Descending Thoracic Aorta**

The part of the body's main artery that runs from the aortic arch through the chest to the diaphragm. It gives branches to the chest wall and organs and continues as the abdominal aorta.

### **Desensitization**

A treatment that gradually exposes a person to increasing or controlled amounts of a specific allergen or medicine. It can reduce immune reactions but must be supervised because serious reactions may occur.

### **Desferrioxamine**

A medicine that binds excess iron and helps remove it from the body. It is used for serious iron overload and selected aluminum toxicity; prolonged treatment can affect hearing, vision, kidneys, or growth.

### **Desflurane**

An inhaled medicine used to maintain general anesthesia. It acts quickly but may irritate the airway, lower blood pressure, cause nausea, or rarely trigger malignant hyperthermia in susceptible people.

### **Desiccate**

To dry out or remove moisture from a tissue, substance, surface, or specimen. Excessive drying can damage skin or wounds and can alter medicines or laboratory samples.

---

## Desiccated Thyroid

A thyroid-hormone medicine made from dried animal thyroid tissue. It contains mainly thyroxine and triiodothyronine; excess treatment can cause palpitations, bone loss, or other effects, so blood tests and monitoring are needed.

### Alternate Names:

Desiccated Thyroid Extract; Natural Desiccated Thyroid

## Desipramine

A tricyclic antidepressant that increases norepinephrine activity. It is used for selected mental health or pain conditions but may cause dry mouth, constipation, drowsiness, abnormal heart rhythms, or dangerous overdose.

## Desipramine Hydrochloride Overdose

Taking too much desipramine hydrochloride can cause severe drowsiness, confusion, seizures, low blood pressure, coma, and life-threatening heart-rhythm or conduction problems. It requires emergency assessment, even if symptoms are initially mild.

## Desmin

A structural protein that links parts of muscle cells and helps maintain their alignment. Changes in the desmin gene or abnormal desmin buildup can cause muscle weakness and heart-muscle disease.

## Desmoid Tumor

A locally aggressive growth of fibrous tissue that usually does not spread to distant organs but can invade nearby structures and recur. It may be associated with familial adenomatous polyposis; treatment depends on location and symptoms.

## Desmoid Tumors

Fibrous growths that usually do not spread to distant organs but can invade nearby tissues and recur. Some remain stable; observation, medicine, or surgery may be considered according to symptoms and location.

## Desmoid Tumour

A growth of fibrous connective tissue that usually does not spread elsewhere but can grow into nearby structures and return after treatment. Its management depends on size, site, growth, and symptoms.

## Desmoplasia

Formation of dense, scar-like tissue around a tumor, long-lasting inflammation, or an injury. Around some cancers it can make a lump feel firm or fixed, but its significance depends on the cells, location, and other findings.

## Desmoplastic

Describing the formation of dense, scar-like fibrous tissue, often around a tumor or in response to chronic inflammation or injury.

## Desmoplastic Fibroblastoma

A rare, usually harmless tumor made of collagen-producing cells. It is typically a slow-growing, painless lump in superficial or deep soft tissue and is treated according to its size, site, and symptoms.

## Desmoplastic Fibroma

A rare noncancerous but locally aggressive fibrous growth in bone. It can cause pain, swelling, or fracture and may return after treatment.

## Desmoplastic Infantile Ganglioglioma

A rare, usually slow-growing brain tumor of infancy containing both nerve-cell and support-cell components. It may cause seizures, increasing head size, vomiting, or other signs of pressure in the brain.

## Desmoplastic Melanoma

A rare form of skin melanoma with abundant scar-like tissue. It often develops on sun-damaged skin, may have little or no pigment, and can grow deeply or recur locally.

## Desmoplastic Reaction

Formation of dense, scar-like fibrous tissue in response to injury, chronic inflammation, or invasion by a cancer. It may make affected tissue firm and

---

can complicate interpretation of imaging or biopsy.

### **Desmoplastic Small Round Cell Tumor**

A rare and aggressive soft-tissue sarcoma that usually develops in the abdomen or pelvis of children, adolescents, or young adults, especially males. It often forms several masses on the lining of the abdomen and can spread within the abdomen or to other sites. Symptoms may include abdominal swelling or pain, nausea, vomiting, diarrhea, or constipation. Tumor cells typically have an *EWSR1-WT1* gene fusion, which helps confirm the diagnosis.

### **Desmoplastic Small Round Cell Tumors**

Plural form; see *Desmoplastic small round cell tumor*.

#### **Alternate Names:**

DSRCT

### **Desmoplastic Trichoepithelioma**

A rare harmless tumor arising from a hair follicle, usually appearing as a small firm facial bump or ring-shaped lesion. It can resemble a skin cancer and may require biopsy for confirmation.

### **Desmopressin**

A medicine that mimics vasopressin, a hormone that reduces urine production and helps control bleeding. It is used for central diabetes insipidus, some bleeding disorders, and selected nighttime urination problems; excess fluid can dangerously lower blood sodium.

### **Desmosome**

A strong junction that fastens neighboring cells together, especially in the skin and heart muscle. It links internal support fibers and helps tissues resist stretching and friction.

### **Desogestrel**

A progestin used in some hormonal contraceptives. It prevents pregnancy mainly by suppressing ovulation and thickening cervical mucus; irregular bleeding, mood changes, breast tenderness, or other hormonal effects may occur.

### **Desonide**

A low-strength corticosteroid applied to the skin to reduce redness, itching, and inflammation from conditions such as eczema or dermatitis. Excessive or prolonged use can thin the skin or cause other local effects.

### **Desoximetasone**

A corticosteroid applied to the skin to reduce inflammation and itching in responsive rashes. Strength, body site, amount, and duration affect the risk of skin thinning and absorption into the body.

### **Desoxycorticosterone**

A steroid hormone with mineralocorticoid effects. It makes the kidneys retain sodium and water and lose potassium, thereby influencing blood volume and blood pressure.

### **Desquamation**

Shedding or peeling of the outer layer of skin or another body lining. It may be normal or result from dryness, irritation, infection, inflammation, burns, medicines, or another skin disorder.

### **Desquamative Interstitial Pneumonia**

A rare inflammatory lung disease strongly associated with smoking. Immune cells collect in the air sacs, causing cough and breathlessness; stopping smoking is essential, and additional treatment may be needed.

### **Desulfovibrio**

A group of bacteria that can live without oxygen and use sulfate during metabolism. They occur in the environment and intestine; invasive human infection is uncommon and must fit the patient's symptoms and specimen findings.

### **Desvenlafaxine Succinate**

A form of desvenlafaxine, an antidepressant that increases serotonin and norepinephrine signaling. It is used for major depressive disorder and may cause nausea, insomnia, increased blood pressure, or withdrawal symptoms if stopped suddenly.

---

## Desynchronosis

Alternate name; see *Jet Lag*.

## Detached

Separated or no longer connected. In medicine it may describe a retina, tissue, device, or feeling of emotional distance; the clinical meaning depends on the surrounding term and findings.

## Detached Retina

Separation of the light-sensitive retina from the tissue beneath it. Flashes, new floaters, or a curtain-like shadow may occur; it is an emergency because delayed treatment can cause permanent vision loss.

## Detergent

A cleaning substance that helps remove oils and dirt from surfaces. Some detergents can irritate or burn the skin, eyes, airways, or digestive tract, especially when concentrated or swallowed.

## Deterministic Effects

Tissue injuries from radiation that usually occur only after exposure exceeds a threshold. Higher exposure generally causes more severe injury, such as skin damage, cataracts, or tissue death.

## Detoxification

Medical treatment that reduces the effects of a poison or helps the body remove it. Care may include supportive treatment, an antidote, activated charcoal, or dialysis; stomach washing is rarely appropriate.

## Detritic Synovitis

Inflammation of the lining of a joint caused by loose particles from damaged cartilage, bone, blood, crystals, or a worn implant. It can cause swelling, pain, stiffness, and reduced movement.

## Detrusor Instability

Involuntary contractions of the bladder muscle while the bladder is filling. They can cause sudden urinary urgency, frequent urination, nighttime urination, or urge leakage.

## Detrusor Muscle

The muscle forming the bladder wall. It relaxes while urine collects and contracts to empty the bladder; abnormal activity can cause urgency or leakage, while weak activity can make emptying difficult.

## Detrusor Sphincter Dyssynergia

A neurological voiding disorder occurring in patients with suprasacral spinal cord lesions, characterized by involuntary, simultaneous contraction of the urinary bladder detrusor muscle and the external urethral sphincter during micturition, producing functional bladder outlet obstruction.

### Alternate Names:

DSD

## Detumescence

Return of erectile tissue to its non-erect state after sexual stimulation or an erection ends. It occurs when blood leaves the erectile tissue and pressure falls.

## Deubiquitinating Enzymes

Enzymes that remove a small regulatory tag from other proteins. This can change whether a protein is broken down, where it works, and how cells control signals, repair DNA, and respond to infection.

## Deubiquitination

Removal of the ubiquitin tag from a protein. This process can prevent the protein from being broken down or can change its activity, location, or role in cell signaling.

## Deuteranopia

An inherited red-green color-vision deficiency caused by missing green-sensitive pigment in the retina. People may have difficulty distinguishing some shades of green, red, brown, and orange.

## Deuterium

A naturally occurring form of hydrogen whose nucleus contains one proton and one neutron. Small amounts occur in water; very high exposure to heavy water can disrupt cell function.



---

## Development

The progressive growth and change by which a person, organ, or ability acquires its structure and function. It includes physical, brain, language, emotional, social, and adaptive changes.

### Developmental Coordination Disorder

A childhood condition in which movement skills develop more slowly or less accurately than expected and interfere with dressing, writing, sports, school, or daily activities. It is not explained only by another medical or neurologic condition.

### Developmental Delay

A child's acquisition of movement, language, thinking, social, or self-care skills is slower than expected for age. Delay may affect one or several areas and can result from genetic, neurologic, sensory, medical, or environmental causes.

### Developmental Disabilities

Lifelong conditions that begin during childhood and affect movement, learning, thinking, language, behavior, or ability to manage daily activities. Support needs vary and may include medical, educational, communication, and social services.

### Developmental Disorder

A condition that begins during childhood and affects physical, intellectual, language, learning, behavioral, or social development. It may interfere with school, relationships, communication, or daily independence.

### Developmental Disorders

Conditions beginning during childhood that affect thinking, language, movement, learning, behavior, or social functioning. Assessment considers developmental history, daily impact, coexisting conditions, strengths, and needed educational or clinical support.

### Developmental Dysplasia Of The Hip

Abnormal development of an infant's or child's hip, ranging from a shallow socket to partial or complete dislocation. Early detection and treatment help

the hip develop normally and reduce later pain or arthritis.

### Developmental Expressive Language Disorder

A childhood disorder causing persistent difficulty expressing ideas with spoken words, sentences, or grammar. Hearing, development, autism, and other possible causes should be assessed before diagnosis.

### Developmental Genital Tract Anomalies

Differences in the formation or structure of the uterus, cervix, vagina, fallopian tubes, or external genitalia that arise before birth. Effects range from none to infertility, pain, menstrual problems, or pregnancy complications.

### Developmental History

A record of a child's developmental milestones and abilities, including motor, language, cognitive, social, and adaptive skills, used to identify possible delays or regression.

### Developmental Milestone

Alternate name; see *Developmental Milestones*.

### Developmental Milestones

A set of functional skills or age-specific behaviors that most children achieve within a predictable developmental window as they grow. Milestones are clinically categorized across four primary domains: gross motor (such as head control, rolling, sitting, and walking), fine motor and adaptive skills (such as pincer grasp, stacking blocks, and drawing), language and communication (such as cooing, single words, phrases, and storytelling), and cognitive/social-emotional development (such as social smiling, pretend play, and interactive sharing). Routine pediatric monitoring tracks milestones from infancy (head steady at 2–4 months, unsupported sitting at 6 months, first steps and single words at 12 months) through toddlerhood and early childhood (running and 2-word phrases at 2 years, conversational sentences at 3 years, and independent dressing and hopping at 4–5 years) to identify developmental delays or regression.



---

**Alternate Names:**

Developmental Milestone; Child Development Milestones; Pediatric Milestones; Developmental Milestones Record

**Developmental Milestones Record**

Alternate name; see *Developmental Milestones*.

**Developmental Process**

The physical, brain, emotional, social, and functional changes through which a person or organ grows and acquires abilities. Development is influenced by genes, health, nutrition, relationships, learning, and environment.

**Developmental Psychology**

The study of how thinking, emotions, relationships, behavior, and abilities change from infancy through old age. It examines the effects of genes, health, learning, experience, relationships, and environment.

**Developmental Reading Disorder**

A learning disorder causing persistent difficulty learning accurate, fluent reading. Problems may include recognizing words, linking letters to sounds, spelling, or understanding written language despite appropriate teaching.

**Developmental Regression**

Loss of skills a child previously acquired, such as speech, movement, social interaction, or self-care. Regression is a warning sign requiring prompt evaluation for neurologic, genetic, metabolic, infectious, or other causes.

**Developmental Stage**

A period of growth characterized by typical physical, thinking, emotional, social, language, or self-care changes. Children vary, so stages describe broad patterns rather than exact deadlines.

**Developmental Surveillance**

Ongoing monitoring of a child's development through caregiver concerns, observation, history,

and milestone review during routine care. It can identify concerns early but differs from formal screening with a standardized tool.

**Deviated Nasal Septum**

Alternate name for deviated septum, in which the wall dividing the nasal passages is displaced from the midline.

**Alternate Names:**

Deviated Septum; Nasal Septal Deviation

**Deviated Septum**

A nasal partition displaced away from the center, often from normal development or injury. A marked deviation can block one side, causing difficulty breathing through the nose, congestion, nosebleeds, sinus problems, or sleep symptoms.

**Deviation**

A difference from an expected position, direction, value, or pattern. In medicine, it may describe anatomy, movement, a laboratory result, or a measurement and must be interpreted in context.

**Devic Disease**

An autoimmune demyelinating central nervous system disorder characterized by recurrent, severe bilateral optic neuritis and longitudinally extensive transverse myelitis spanning three or more vertebral segments, typically mediated by pathogenic anti-aquaporin-4 IgG autoantibodies (AQP4-IgG).

**Alternate Names:**

Neuromyelitis Optica; NMO

**Device Embolisation**

Movement of a medical device or broken piece of one through the bloodstream until it lodges in and blocks a blood vessel. It can reduce blood flow and may cause pain, organ injury, or an emergency requiring removal.

**Devices For Hearing Loss**

Equipment that improves access to sound, including hearing aids, cochlear implants, bone-conduction devices, alerting systems, and communication acces-

---

sories. Choice depends on the type and severity of hearing loss and personal needs.

### **Dexamethasone**

A strong, long-acting corticosteroid that reduces inflammation and immune activity. It is used for selected allergic, brain-swelling, cancer-related, endocrine, and other conditions; risks include infection, high blood sugar, mood changes, and adrenal suppression.

### **Dexamethasone Suppression Test**

A test of cortisol control. After taking a small dose of dexamethasone, blood or urine cortisol is measured; failure to suppress may suggest excess cortisol but can have other causes and needs clinical interpretation.

### **Dexamphetamine**

A stimulant that increases dopamine and norepinephrine activity in the brain. It is used for selected attention or sleep disorders and may cause reduced appetite, insomnia, fast heartbeat, high blood pressure, or dependence.

### **Dexibuprofen**

A nonsteroidal anti-inflammatory medicine used for short-term pain and inflammation. It can irritate the stomach and may cause ulcers, bleeding, kidney injury, fluid retention, or cardiovascular problems.

### **Dexketoprofen**

A nonsteroidal anti-inflammatory medicine used for short-term pain. It reduces substances that cause pain and inflammation but can cause stomach bleeding, kidney injury, fluid retention, heart problems, or allergic reactions.

### **Dexlansoprazole**

A proton-pump inhibitor that reduces stomach-acid production. It is used for acid reflux and related esophageal injury; prolonged use may contribute to infections, low magnesium or vitamin B12, fractures, or kidney inflammation.

### **Dexmedetomidine**

A sedative that activates alpha-2 receptors and is given by injection during monitored care or intensive care. It usually depresses breathing less than many sedatives but can slow the heart or lower blood pressure.

### **Dexmethylphenidate**

A stimulant used to treat attention-deficit/hyperactivity disorder. It increases dopamine and norepinephrine activity and may cause reduced appetite, insomnia, headache, increased heart rate or blood pressure, or misuse.

### **Dexrazoxane**

A medicine used in selected cancer treatments to reduce heart damage from anthracyclines or tissue injury after leakage of an anthracycline from a vein. It can suppress bone marrow and increase infection risk.

### **Dexrazoxane Hydrochloride**

A salt form of dexrazoxane used in selected settings to reduce anthracycline-related heart damage or tissue injury after medicine leakage from a vein. Its use requires specialist dosing and monitoring.

### **Dextrin**

A group of shorter carbohydrate chains formed when starch breaks down. Dextrins are used in foods, laboratory products, and medicines and are generally digested as carbohydrates.

### **Dextroamphetamine**

A stimulant that increases dopamine and norepinephrine activity in the brain. It is used for selected attention or sleep disorders and may cause insomnia, reduced appetite, fast heartbeat, high blood pressure, or dependence.

### **Dextrocardia**

A congenital condition in which the heart points or lies mainly on the right side of the chest. It may occur alone or with reversed organs and structural heart defects.

---

## Dextromethorphan

A cough suppressant that reduces activity in brain pathways controlling the cough reflex. Excessive amounts can cause drowsiness, agitation, hallucinations, seizures, breathing depression, or serotonin toxicity.

## Dextromethorphan Overdose

Taking too much dextromethorphan can cause drowsiness, confusion, agitation, hallucinations, fast heartbeat, seizures, breathing depression, or serotonin toxicity. It requires urgent poison or emergency assessment.

## Dextroposition Of The Heart

The heart is pushed toward the right side of the chest by nearby organs, the diaphragm, lung disease, or another space-occupying problem. The heart itself may be normally formed and is not necessarily reversed as in dextrocardia.

## Dextrose

D-glucose, a simple sugar used for energy. Oral or intravenous dextrose can treat or prevent low blood glucose, but excessive amounts can raise blood sugar and disturb body fluids.

## DFSP

Alternate name; see *Dermatofibrosarcoma Protuberans*.

## DH

Abbreviation commonly used for dermatitis herpetiformis, an intensely itchy blistering skin disorder associated with gluten-sensitive enteropathy and celiac disease. The abbreviation can have other meanings and should be interpreted in context.

## DHA

Docosahexaenoic acid, an omega-3 fatty acid that is an important part of brain and retinal cell membranes. It is obtained mainly from oily fish, some algae, and supplements.

## DHEA

Dehydroepiandrosterone, a hormone precursor made mainly by the adrenal glands. The body can convert it into male and female sex hormones; levels vary with age, illness, and some hormone disorders.

## DHEA-Sulfate Test

A blood test measuring a hormone made mainly by the adrenal glands. It can help investigate early puberty, excess body hair, some hormone disorders, congenital adrenal hyperplasia, or an adrenal tumor; results vary with age and sex.

## Diabetes And Alcohol

Alcohol can cause delayed low blood glucose, interact with medicines, and make glucose control harder. Risk depends on amount, timing, food intake, diabetes type, liver health, and treatment.

## Diabetes And Exercise

Physical activity can improve glucose control, fitness, and heart health. People using insulin or medicines that lower glucose may need to check levels and adjust food or treatment with professional guidance to prevent low glucose.

## Diabetes And Eye Disease

Diabetes can damage retinal blood vessels, causing retinopathy and macular swelling that may reduce vision. Good glucose and blood-pressure control and regular dilated eye examinations help prevent or limit sight loss.

## Diabetes And Eye Problems

Diabetes may cause retinopathy, macular swelling, cataracts, glaucoma, and vision loss. Glucose and blood-pressure control, regular dilated eye examinations, and timely laser, injection, or surgical treatment reduce the risk of severe loss.

## Diabetes And Kidney Disease

Kidney damage caused by diabetes. It may begin with albumin leaking into urine and progress to reduced kidney function or kidney failure; glucose, blood-pressure, and kidney monitoring can slow progression.

---

## Diabetes And Metabolic Health

Diabetes is linked with insulin resistance, excess body fat, abnormal cholesterol, and high blood pressure. Together these problems increase the risk of heart disease and can worsen blood-glucose control.

## Diabetes And Nerve Damage

Long-term diabetes can injure nerves, causing burning pain, tingling, numbness, weakness, digestive or bladder problems, sexual difficulties, or loss of protective feeling in the feet.

## Diabetes Complications

Problems caused by diabetes, including low or high glucose emergencies and long-term damage to the eyes, kidneys, nerves, heart, blood vessels, teeth, and feet. Regular monitoring and risk-factor control reduce risk.

## Diabetes Eye Exams

Eye examinations that look for diabetic retinopathy, macular swelling, cataracts, glaucoma, and other sight-threatening problems. Dilated examination or retinal photography may detect changes before vision is affected.

## Diabetes in Children: Dealing with Low Blood Sugar

Recognition and treatment of hypoglycemia in a child with diabetes. Symptoms may include sweating, trembling, hunger, irritability, confusion, seizure, or loss of consciousness; fast-acting glucose is used when the child is awake and able to swallow, with emergency glucagon for severe episodes.

## Diabetes in Children: Preventing Low Blood Sugar

Measures to reduce hypoglycemia in children with diabetes, including individualized insulin dosing, regular meals and snacks, glucose monitoring, planning around activity, and education about symptoms and emergency glucagon.

## Diabetes in Pregnancy

Alternate name; see *Gestational Diabetes*.

## Diabetes in Pregnancy (Gestational Diabetes)

Diabetes first recognized during pregnancy, caused by inadequate insulin response to pregnancy-related metabolic changes. It increases risks for excessive fetal growth, birth complications, preeclampsia, and later type 2 diabetes and requires monitoring and treatment.

## Diabetes Insipidus

A disorder causing large amounts of very dilute urine and intense thirst because vasopressin is lacking or the kidneys do not respond to it. It is unrelated to diabetes mellitus and can cause dehydration.

## Diabetes Medications

Medicines that lower blood glucose or replace insulin. They work in different ways, and the choice depends on the type of diabetes, kidney and heart health, other medicines, cost, and treatment goals.

## Diabetes Mellitus

A group of diseases in which blood glucose, also called blood sugar, stays too high. Insulin is a hormone made by the pancreas that helps glucose move from the blood into cells for energy. Diabetes develops when the body makes too little or no insulin, cannot use insulin properly, or both. The main types are type 1 diabetes, type 2 diabetes, and gestational diabetes, which develops during pregnancy; other forms can result from genetic conditions, pancreatic disease, or medicines. Persistently high blood glucose can damage the eyes, kidneys, nerves, heart, and blood vessels. Diabetes mellitus is different from diabetes insipidus, a disorder of water balance that is not caused by high blood glucose.

### Alternate Names:

Diabetes

## Diabetes Mellitus and Pregnancy

Diabetes before or first recognized during pregnancy can increase risks of miscarriage, fetal overgrowth or growth restriction, congenital anomalies, hypertensive disease, stillbirth, and neonatal metabolic problems. Glucose control, retinal and kidney assessment, medication review, and fetal surveillance

---

are important.

### **Diabetes Mellitus Type 1**

An autoimmune disorder in which the immune system destroys insulin-producing pancreatic cells, causing little or no insulin. It can begin at any age and requires insulin; untreated deficiency can cause ketoacidosis.

### **Diabetes Mellitus Type 2**

A disorder in which the body resists insulin and, over time, may not make enough of it. Risk is increased by family history, excess body weight, and inactivity; it can occur at any age and may have no early symptoms.

### **Diabetes Prevention**

The risk of type 2 diabetes can often be reduced through regular physical activity, nutritious meals, avoiding tobacco, adequate sleep, and weight loss when appropriate. People at increased risk may benefit from structured prevention programs and regular testing.

### **Diabetes Treatment**

Diabetes treatment combines nutrition, physical activity, glucose monitoring, education, and medicines chosen for the type of diabetes and individual risks. Care also addresses blood pressure, lipids, kidneys, eyes, nerves, feet, vaccines, and cardiovascular health.

### **Diabetic Amyotrophy**

Alternate name; see *Diabetic Neuropathy*.

### **Diabetic Angiopathy**

Damage to small or large blood vessels caused by diabetes. It can reduce circulation, injure the eyes, kidneys, heart, brain, or feet, and increase the risk of wounds, heart attack, or stroke.

### **Diabetic Autonomic Neuropathy**

Diabetes-related damage to nerves that control automatic body functions. It may cause dizziness on standing, fast heartbeat, sweating changes, de-

layed stomach emptying, bladder problems, sexual dysfunction, or reduced awareness of low glucose.

### **Diabetic Cardiomyopathies**

Heart-muscle disease associated with diabetes that is not fully explained by blocked heart arteries or high blood pressure. It can make the heart relax or pump poorly and may lead to heart failure.

### **Diabetic Cardiomyopathy**

Heart-muscle damage associated with diabetes, after other causes such as blocked arteries and high blood pressure are considered. It may cause poor relaxation, weak pumping, heart failure, or rhythm problems.

### **Diabetic Coma**

Profound unconsciousness caused by a severe diabetes-related problem, such as extremely low or high blood glucose, ketoacidosis, severe dehydration, or abnormal body chemistry. It is a medical emergency.

### **Diabetic Complications**

Acute or long-term problems caused by diabetes, including low or high glucose emergencies and damage to the eyes, kidneys, nerves, heart, blood vessels, teeth, and feet.

### **Diabetic Diet**

An individualized eating pattern for diabetes that balances carbohydrate portions, fiber, protein, and healthy fats with medicines, activity, culture, and health goals. It supports glucose control without requiring the same foods or schedule for everyone.

#### **Alternate Names:**

Diabetic Meal Plan; Medical Nutrition Therapy for Diabetes

### **Diabetic Eye Disease**

Eye damage caused by diabetes, including diabetic retinopathy, diabetic macular edema, cataract, glaucoma, and cranial-nerve or optic-nerve complications. Regular dilated eye examinations can detect treatable disease before vision is lost.



---

## Diabetic Focal Neuropathy

Alternate name; see *Diabetic Neuropathy*.

## Diabetic Foot

Foot problems associated with diabetes, including loss of sensation, deformity, ulcers, infection, poor circulation, and slow healing. Daily inspection, proper footwear, glucose control, and early care help prevent serious injury.

## Diabetic Foot Disease

Foot damage caused by a combination of diabetes-related nerve loss, poor circulation, deformity, and impaired healing or immunity. A small injury can progress to an ulcer, deep infection, or amputation without early care.

## Diabetic Foot Infections

Infection involving the skin, soft tissue, or bone of a diabetic foot. Neuropathy, ischemia, ulceration, and impaired immunity can conceal severity or delay healing, so assessment includes depth, perfusion, systemic illness, and need for antimicrobial therapy or source control.

## Diabetic Foot Ulcer

A chronic non-healing sore on the foot in a person with diabetes, usually caused by a combination of peripheral neuropathy, poor blood circulation, and localized pressure or minor trauma.

### Alternate Names:

Diabetic Ulcer; Neuropathic Foot Ulcer; Diabetic Foot Sore

## Diabetic Gastroparesis

A complication of diabetes in which damage to the stomach's nerves slows the movement of food from the stomach into the small intestine. It may cause early fullness, nausea, vomiting, bloating, abdominal pain, and difficult-to-control blood sugar levels.

## Diabetic Hyperglycemic Hyperosmolar Syndrome

A life-threatening diabetes emergency with extremely high blood glucose, severe dehydration, and

concentrated blood, usually without major ketone buildup. It can cause confusion, seizures, coma, and shock and requires hospital treatment.

## Diabetic Hyperosmolar Hyperglycemic State

A severe diabetes emergency marked by extremely high glucose, severe dehydration, and increased blood concentration with little or no ketoacidosis. Infection, illness, or inadequate treatment may trigger it; urgent hospital care is required.

## Diabetic Hypoglycemia

Blood glucose that is too low in a person with diabetes, often because of insulin, certain medicines, missed food, alcohol, or unusual exercise. Shaking, sweating, confusion, seizures, or unconsciousness may occur.

## Diabetic Ketoacidosis

A life-threatening hyperglycemic emergency caused by insufficient effective insulin, with diabetes or hyperglycemia, elevated ketones, and metabolic acidosis. Diagnostic criteria generally include glucose at least 200 mg/dL or known diabetes, beta-hydroxybutyrate at least 3 mmol/L or significant ketonuria, and pH below 7.30 or bicarbonate below 18 mmol/L. Polyuria, dehydration, nausea, vomiting, abdominal pain, Kussmaul breathing, and altered mental status may occur; euglycemic DKA is possible, especially with SGLT2 inhibitors.

### Alternate Names:

DKA

## Diabetic Ketoacidosis Monitoring

Monitoring during treatment includes frequent glucose checks, blood salts, kidney function, ketones, acidity, fluid balance, urine output, and mental status. Close monitoring helps guide fluids, insulin, and electrolyte replacement safely.

## Diabetic Kidney Disease

Kidney disease occurring in a person with diabetes, often attributed to diabetes when no other cause is evident. It may involve albumin in the urine, reduced kidney filtering, or both, and can progress



---

to kidney failure. Other causes of chronic kidney disease should also be considered.

**Alternate Names:**

DKD; Diabetic Nephropathy

**Diabetic Macular Edema**

Swelling of the macula, the central part of the retina, caused by leaking blood vessels in diabetes. It can blur or distort central vision and may occur at any stage of diabetic retinopathy.

**Diabetic Mastopathy**

A rare, noncancerous, firm breast thickening associated mainly with long-standing diabetes. It can feel or look like breast cancer, so examination and imaging—and sometimes a biopsy—may be needed to establish the diagnosis.

**Diabetic Microvascular Complications**

Diabetes-related damage to small blood vessels, mainly affecting the eyes, kidneys, and nerves. Good glucose, blood-pressure, and cholesterol control, along with regular screening, lowers the risk of progression.

**Diabetic Neuropathy**

Nerve damage caused by diabetes. It may cause burning pain, tingling, numbness, weakness, loss of protective feeling in the feet, or problems with digestion, blood pressure, sweating, bladder control, or sexual function.

**Alternate Names:**

Diabetic Nerve Damage; Peripheral Diabetic Neuropathy

**Diabetic Neuropathy Entry**

Alternate name; see *Diabetic Neuropathy*.

**Diabetic Peripheral Neuropathy**

Diabetes-related damage to nerves, usually beginning in the toes and feet and later affecting the hands. It can cause numbness, burning pain, weakness, and loss of protective sensation that increases injury risk.

**Diabetic Proximal Neuropathy**

Alternate name; see *Diabetic Neuropathy*.

**Diabetic Retinopathy**

Damage to retinal blood vessels caused by diabetes. Early disease may have no symptoms; advanced disease can cause macular swelling, bleeding, retinal detachment, and vision loss. Regular dilated eye examinations are important.

**Alternate Names:**

Diabetic Eye Disease; Diabetic Microangiopathy of Retina

**Diabetic Third Nerve Palsy**

An ischemic microvascular cranial mononeuropathy affecting the third cranial (oculomotor) nerve secondary to diabetes mellitus, classically presenting with acute-onset diplopia, ptosis, and extraocular muscle weakness with characteristic pupil sparing.

**Alternate Names:**

Diabetic Oculomotor Neuropathy; Pupil-Sparing Third Nerve Palsy; Diabetic Cranial Neuropathy III

**Diabetogenic**

Able to cause or promote diabetes or high blood glucose. The term may describe a disease, hormone, medicine, genetic change, or other factor that reduces insulin action or increases glucose production.

**Diabetologist**

A physician specializing in diabetes prevention, diagnosis, and treatment, including glucose management and complications involving the heart, kidneys, nerves, eyes, and feet.

**Diabrosis**

An old medical term for erosion or corrosion that eats through tissue and may cause an ulcer or perforation, especially in a blood-vessel or organ wall. Modern reports usually name the specific cause and injury.

**Diacetoxyscirpenol**

A toxin produced by some *Fusarium* molds that can contaminate food or animal feed. Exposure

---

may irritate the digestive tract, suppress blood-cell production, and impair immune function.

### **Diacylglycerol**

A fat-like molecule made from cell membranes that acts as an internal chemical signal. It helps control cell growth, movement, secretion, metabolism, and responses to hormones or other signals.

### **Diagnosis**

Identification of a disease or health condition using symptoms, medical history, examination, and appropriate tests. A clinician may compare several possible causes before reaching the most likely diagnosis.

### **Diagnostic**

Relating to identifying a disease, condition, or cause of symptoms. A diagnostic test provides information that must be interpreted with the person's history, examination, and other findings rather than alone.

### **Diagnostic Imaging**

Use of images to detect, locate, or monitor disease or injury. Methods include X-ray, ultrasound, computed tomography, magnetic resonance imaging, and nuclear imaging; findings must be interpreted with the clinical situation.

### **Diagnostic Laparoscopy**

A minimally invasive procedure in which a camera is inserted through small abdominal incisions to inspect pelvic or abdominal organs. The clinician may take tissue samples or perform treatment during the same procedure.

### **Diagnostic Microbiology**

Laboratory testing used to identify microorganisms that may cause disease. Methods include microscopy, culture, antigen or antibody tests, and genetic tests; results must be interpreted with symptoms and the specimen site.

### **Diagnostic Peritoneal Lavage**

An older emergency test for serious internal bleeding

or injury after major abdominal trauma. A small tube is placed into the abdomen and fluid is put in and then removed for testing. Ultrasound and CT scans are now used more often, but this test may still be used when those scans are not available.

#### **Alternate Names:**

DPL; Peritoneal Lavage; Abdominal Lavage

### **Diagnostic Procedure**

An examination or procedure used to identify, exclude, locate, or monitor a disease or injury. The purpose, preparation, risks, accuracy, and follow-up depend on the specific procedure and patient.

### **Diagnostic Radiology**

The medical specialty that uses imaging to diagnose and characterize disease or injury. It includes X-ray, ultrasound, computed tomography, magnetic resonance imaging, and selected nuclear-medicine studies.

### **Diagnostic Service**

A healthcare service that collects or interprets information to detect, identify, characterize, or monitor a disease or health condition, such as laboratory testing, imaging, or specialist assessment.

### **Diagnostic Test**

A test used to evaluate symptoms, signs, or an abnormal screening result and help confirm, exclude, or characterize a disease. Its value depends on accuracy, timing, the patient's condition, and how the result will guide care.

### **Diagnostic Trial**

A clinical study that evaluates how accurately a test detects, rules out, or predicts a disease or important health outcome. Researchers may compare it with an established test or another accepted way of determining the diagnosis.

### **Diagnostic Uncertainty**

A situation in which available information does not establish one diagnosis with enough confidence. It is managed by explaining uncertainty, arranging follow-up, watching for warning signs, and ordering

---

targeted further tests when appropriate.

### **Diagnostics**

The tests, examinations, and clinical reasoning used to detect, identify, classify, or monitor a health condition. Results must be interpreted in context because no test is perfect.

### **Diakinesis**

The final stage before the first division that produces eggs or sperm. Paired chromosomes become tightly packed and remain connected at points where they exchanged genetic material.

### **Dialectical Behavior Therapy**

A structured talking treatment that combines behavior-change skills with mindfulness and acceptance. It teaches ways to manage intense emotions, tolerate distress, improve relationships, and reduce harmful behaviors.

### **Dialister Pneumosintes**

A bacterium found in the mouth and respiratory tract that grows without oxygen. It may contribute to mixed infections, but a positive sample must be interpreted with symptoms and collection site.

### **Dialysis**

Treatment that replaces part of the kidneys' work by removing waste, excess water, and some chemicals from the blood. The main types are hemodialysis and peritoneal dialysis.

### **Dialysis Access**

A route used to connect a person's blood circulation to a hemodialysis machine. Options include an arteriovenous fistula, graft, or central venous catheter; each has different risks of infection, clotting, and inadequate flow.

### **Dialysis Adequacy**

How well dialysis removes waste and extra fluid and keeps body salts and acid levels within a safe range. Blood tests, treatment records, symptoms, nutrition, and weight help determine whether dialysis is

sufficient.

### **Dialysis Complications of Chronic Renal Failure**

Acute and chronic problems related to hemodialysis or peritoneal dialysis, including hypotension, disequilibrium, access infection or thrombosis, bleeding, electrolyte shifts, peritonitis, and amyloidosis.

### **Dialysis Disequilibrium Syndrome**

A rare brain reaction during or soon after starting or intensifying hemodialysis, caused by rapid changes in dissolved substances. Headache, nausea, confusion, seizures, or coma may occur; gradual treatment reduces risk.

### **Dialysis Machine**

Equipment that moves blood or dialysis fluid across a filter to remove waste, excess water, and selected salts when the kidneys cannot do so adequately.

### **Dialysis Solutions**

Specially prepared fluids used during hemodialysis or peritoneal dialysis. They allow waste and excess water to move out of the blood; their salt, glucose, and other contents are adjusted to the patient's needs.

### **Dialysis Therapy**

Use of hemodialysis or peritoneal dialysis to replace part of kidney function by removing waste and excess fluid and correcting some salt and acid imbalances. Treatment is individualized according to kidney function, symptoms, health, and goals.

### **Dialysis-Related Beta-2m Amyloidosis**

Amyloid deposition caused by accumulation of beta-2 microglobulin in people receiving long-term dialysis, especially with inadequate clearance. It can cause shoulder or joint pain, carpal tunnel syndrome, bone cysts, and destructive arthropathy.

### **Diamine**

A molecule containing two amino groups. Diamines occur naturally in metabolism and are used in lab-

---

oratory, pharmaceutical, and industrial chemistry; health effects depend on the specific compound and exposure.

### **Diamond-Blackfan Anemia**

A rare inherited disorder in which the bone marrow makes too few red blood cells, usually beginning in infancy. It may also cause short stature, thumb or facial differences, or heart defects.

#### **Alternate Names:**

DBA; Diamond-Blackfan Syndrome; Congenital Hypoplastic Anemia

### **Diamond-Gardner Syndrome**

A rare condition in which painful bruises appear repeatedly without a clear injury, most often in women. It is a diagnosis made after other bleeding causes are excluded; old skin tests are not reliable confirmation.

#### **Alternate Names:**

Autoerythrocyte Sensitization Syndrome; Painful Bruising Syndrome

### **Diamorphine**

A strong opioid medicine used in selected settings for severe pain or during medical procedures. It can cause sleepiness, nausea, constipation, dependence, and dangerous breathing depression, especially in overdose or with other sedatives.

### **Dianisidine**

An industrial chemical used in some dyes and laboratory reagents. Significant or repeated exposure can irritate tissues and may damage blood or increase cancer risk; workplace protection is important.

### **Diapedesis**

Movement of blood cells, especially infection-fighting white cells, through the walls of tiny blood vessels into nearby tissue. This allows them to reach and respond to infection, injury, or inflammation.

### **Diaper Rash**

Inflammation of skin covered by a diaper, commonly caused by moisture, urine, stool, friction, or yeast

infection. It causes redness and soreness; a persistent, severe, blistering, or spreading rash needs medical assessment.

#### **Alternate Names:**

Diaper Dermatitis; Napkin Dermatitis

### **Diaphoresis**

Excessive sweating. It may be a normal response to heat or exercise, or a sign of fever, pain, low blood glucose, anxiety, infection, heart problems, or another condition.

### **Diaphoretic**

Describing heavy sweating or something that causes sweating. Sudden cold sweating with chest pain, faintness, breathlessness, confusion, or weakness may signal a serious medical problem and needs urgent assessment.

### **Diaphragm**

The dome-shaped muscle separating the chest from the abdomen and the main muscle used for breathing. When it contracts, it flattens and draws air into the lungs; when it relaxes, air leaves.

### **Diaphragmatic Crus**

One of two musculotendinous bundles (right and left crura) originating from the anterior aspects of the upper lumbar vertebral bodies and fibrocartilaginous discs, ascending to insert into the central tendon of the diaphragm and encircling the esophageal hiatus.

### **Diaphragmatic Eventration**

Abnormal thinning or weakness of part or all of the diaphragm, causing it to sit higher than usual. It may cause no symptoms or lead to breathlessness, especially if severe or present from birth.

### **Diaphragmatic Hernia**

Movement of abdominal organs through a hole or weak area in the diaphragm into the chest. It may be present from birth or follow injury and can impair breathing or cause bowel obstruction.

---

## Diaphragmatic Paralysis

Loss of normal diaphragmatic movement from phrenic-nerve injury, neuromuscular disease, surgery, trauma, or idiopathic causes. Symptoms range from none to breathlessness when lying flat and sleep-related hypoventilation; bilateral disease is more likely to cause respiratory failure.

## Diaphragmatic Rupture

A tear in the diaphragm, usually caused by a severe blunt or penetrating injury to the chest or abdomen. Abdominal organs may move into the chest, causing breathing difficulty, obstruction, or strangulation.

## Diaphysis

The long central shaft of a bone such as the femur, humerus, or tibia. It contains a hollow marrow cavity and is surrounded by strong compact bone; fractures there may require prolonged healing.

## Diarrhea

Passing loose or watery stools more often than usual. Causes include infections, medicines, food intolerance, inflammation, malabsorption, or other illnesses. Diarrhea can lead to dehydration and, when prolonged, malabsorption.

## Diarrhea-Predominant IBS

Alternate name; see *Irritable Bowel Syndrome*.

## Diarrhoea

Alternate spelling; see *Diarrhea*.

## Diarthrosis

A freely movable joint in which the bones are separated by a space containing lubricating fluid. The knee, hip, shoulder, and elbow are examples.

## Diascopy

A clinical examination technique in which a transparent glass slide or plastic lens is pressed firmly against a skin lesion to assess whether it blanches, allowing differentiation between intravascular blood (erythema, telangiectasias, spider naevi) and extravasated red blood cells (purpura, petechiae, ec-

chymoses).

## Diastase

An older name for enzymes that break starch into smaller sugars. Modern medical writing usually uses the term amylase for the related digestive enzyme activity.

## Diastasis

Separation or widening between structures that normally lie together. It commonly describes separation of the abdominal muscles after pregnancy and is not itself a true hernia, although both can occur together.

## Diastasis Recti

Widening and thinning of the tissue between the two front abdominal muscles, commonly during or after pregnancy or major weight change. It can cause a midline bulge when the abdomen is strained and is not a true hernia.

## Diastematomyelia

A birth defect in which the spinal cord is divided into two parts, often by bone or fibrous tissue. It may occur with abnormal vertebrae, a skin mark, weakness, altered sensation, or bladder problems.

## Diastole

The part of each heartbeat when the heart muscle relaxes and its chambers fill with blood. The lower blood-pressure number is the pressure in the arteries during this phase.

## Diastolic

Relating to diastole, when the heart relaxes and fills. Diastolic blood pressure is the lower number in a blood-pressure reading.

## Diastolic Blood Pressure

The pressure in the arteries while the heart relaxes between beats. It is the lower number in a blood-pressure reading and is interpreted with the upper number, symptoms, age, and other health conditions.

---

## Diastolic Dysfunction

A condition where the heart muscle becomes stiff and cannot relax properly between beats. Although the heart still squeezes with normal strength, the stiffness makes it difficult for the heart chambers to fill with blood. This causes pressure inside the heart to rise, which can make fluid back up into the lungs and legs. Commonly caused by aging, long-term high blood pressure, and diabetes, it typically produces shortness of breath during activity, tiredness, and swollen ankles. It is usually detected using an ultrasound scan of the heart (echocardiogram).

### Alternate Names:

Stiff Heart; Left Ventricular Diastolic Dysfunction; LVDD; Impaired Heart Relaxation

## Diastolic Heart Failure

An older term usually referring to heart failure caused by a stiff heart that does not fill easily, often with preserved pumping percentage. Diagnosis also requires symptoms, signs, and evidence of raised filling pressure.

## Diastolic Murmur

An extra or unusual heart sound heard while the heart relaxes between beats. It often reflects abnormal flow through a heart valve or nearby blood vessel and needs medical evaluation.

## Diastolic Murmurs

Extra heart sounds heard during the relaxation phase between beats. They commonly result from abnormal flow through a heart valve or major blood vessel and require clinical assessment.

## Diathermy

Use of high-frequency electrical or electromagnetic energy to produce heat in tissue. It may be used during surgery to cut tissue or stop bleeding, or in selected treatments to warm deeper tissues.

## Diathesis

A tendency to develop a particular disease, symptom, or body response. For example, a bleeding diathesis means a person is more likely to bleed than usual.

## Diatom Test

A forensic test that looks for microscopic algae in tissues after suspected drowning. It may support an investigation but cannot prove drowning by itself because algae can enter the body from other sources.

## Diatomaceous Earth

A powder made from the fossilized remains of microscopic algae. Inhalation of fine particles can irritate the lungs, and some forms containing crystalline silica can cause serious lung disease with repeated exposure.

## Diatrizoate Meglumine

An iodinated contrast substance used for selected X-ray or imaging studies. It improves visibility of body structures but may cause allergic-like reactions, kidney injury, aspiration problems, or thyroid effects.

## Diazepam

A benzodiazepine that calms brain activity and is used for selected anxiety, muscle-spasm, seizure, and alcohol-withdrawal conditions. It can cause drowsiness, falls, dependence, withdrawal, and dangerous breathing depression with opioids or alcohol.

## Diazepam Overdose

Taking too much diazepam can cause extreme sleepiness, confusion, poor coordination, low blood pressure, slowed breathing, coma, or death, especially when combined with alcohol, opioids, or other sedatives.

## Diazinon

An organophosphate insecticide that blocks acetylcholinesterase, causing excess nerve signals. Exposure can lead to sweating, salivation, vomiting, diarrhea, breathing difficulty, muscle weakness, seizures, or respiratory failure.

## Diazinon Poisoning

Poisoning by diazinon, causing excess cholinergic activity with sweating, salivation, vomiting, diarrhea, wheezing, slow heartbeat, muscle weakness, seizures, or breathing failure. It requires emergency



---

treatment and decontamination.

### **Diazonium Compounds**

Chemicals containing a diazonium group, used in dyes, synthesis, and laboratory reactions. Some are unstable or toxic and may irritate tissues or release hazardous products.

### **Diazoxide**

A medicine that opens potassium channels and can reduce insulin release or relax blood vessels. It is used in selected cases of low blood sugar from excess insulin and severe high blood pressure; fluid retention and high glucose may occur.

### **Dibekacin**

An aminoglycoside antibiotic used in some regions for serious bacterial infections. It blocks bacterial protein production but can damage the kidneys, hearing, or balance, especially with prolonged or high-dose treatment.

### **Dibenzepin**

A tricyclic antidepressant used in some countries for depressive disorders. It can cause drowsiness, dry mouth, constipation, blurred vision, abnormal heart rhythms, and dangerous toxicity in overdose.

### **Dibucaine**

A long-acting local anesthetic applied to skin or mucous membranes to reduce pain or itching. Excessive use or absorption can cause numbness beyond the treatment area, seizures, heart problems, or other toxicity.

### **Dicarboxylic Acid**

An organic molecule containing two acid groups. These molecules take part in metabolism and may accumulate in some inherited or acquired metabolic disorders.

### **Dichlorophen**

A chemical formerly used in some disinfectants and medicines. Exposure can irritate the skin, eyes, or digestive tract; serious effects depend on the amount,

route, and duration of exposure.

### **Dichlorvos**

An organophosphate insecticide that can block acetylcholinesterase. Poisoning may cause sweating, salivation, vomiting, diarrhea, muscle weakness, breathing difficulty, seizures, or respiratory failure.

### **Dichorionic Diamniotic Twin Placenta**

A twin pregnancy in which each fetus has its own outer membrane and amniotic sac, usually with a separate placenta. It generally has fewer shared-placenta complications, but routine maternal and fetal monitoring remains important.

### **Dichorionic Twins**

Twins that each have a separate outer membrane and amniotic sac. Most fraternal twins and some identical twins are dichorionic; risks still include premature birth, growth problems, and pregnancy complications.

### **Diclofenac**

A nonsteroidal anti-inflammatory medicine used for pain and inflammation. It can cause stomach irritation or bleeding, kidney or liver injury, fluid retention, and increased cardiovascular risk.

### **Diclofenac Sodium**

A salt form of diclofenac used for pain and inflammation. It can irritate or bleed the stomach and may affect the kidneys, blood pressure, liver, or heart.

### **Diclofenac Sodium Overdose**

Taking too much diclofenac sodium can cause nausea, vomiting, stomach bleeding, kidney injury, drowsiness, seizures, breathing problems, or abnormal body chemistry and requires urgent medical assessment.

### **Dicloxacillin**

A penicillin-type antibiotic that blocks bacterial cell-wall production. It is used for susceptible infections, especially some staphylococcal infections; allergy,

---

diarrhea, and liver injury can occur.

### **DICOM**

An international format and communication standard for storing, sending, and viewing medical images and related information. It allows images from X-ray, CT, MRI, ultrasound, and other systems to be shared.

### **Dicoumarol**

An older oral anticoagulant that blocks vitamin K recycling and reduces production of several clotting factors. It has largely been replaced by medicines with more predictable effects and remains important mainly historically.

### **Dicrotic**

Describing a second small wave or notch in an artery's pressure pulse, usually produced when the aortic valve closes. It may be visible on a monitor or arterial pressure tracing.

### **Dicrotophos**

An organophosphate insecticide that can inhibit acetylcholinesterase. Poisoning may cause sweating, salivation, vomiting, diarrhea, muscle weakness, breathing difficulty, seizures, or respiratory failure.

### **Dicyclomine**

An anticholinergic medicine that relaxes intestinal muscle and may reduce cramping in selected bowel disorders. It can cause dry mouth, blurred vision, constipation, fast heartbeat, confusion, or difficulty urinating.

### **Dieffenbachia Poisoning**

Poisoning after chewing or swallowing parts of the Dieffenbachia plant. Needle-like calcium oxalate crystals can cause immediate burning, pain, drooling, and swelling of the mouth and throat; airway swelling requires emergency care.

### **Diencephalon**

A central part of the brain containing the thalamus and hypothalamus. It helps process sensory informa-

tion and regulate hormones, temperature, appetite, sleep, movement, and other automatic functions.

### **Dientamoebiasis**

An infection of the human large intestine caused by *Dientamoeba fragilis*, a non-invasive, flagellated trichomonad protozoan that exists solely in a trophozoite stage. It frequently presents as chronic or intermittent gastrointestinal distress characterized by watery or loose diarrhea, abdominal pain, flatulence, anorexia, nausea, and fatigue, particularly in children, travelers, and institutionalized populations.

#### **Alternate Names:**

Dientamoeba Fragilis Infection; Dientamoeba Colitis

### **DIEP Flap**

A reconstructive surgery using skin and fat from the lower abdomen, supplied by small blood vessels, to rebuild a breast or another body area while preserving most abdominal muscle.

### **Diesel Oil**

A petroleum fuel whose fumes or liquid can irritate the skin, eyes, and lungs. Swallowing or aspiration into the lungs can cause severe chemical pneumonia; suspected exposure requires prompt poison or medical advice.

### **Diestrus**

A phase of the reproductive cycle in many mammals between periods of sexual receptivity. It is characterized by activity and later regression of the corpus luteum.

### **Diet Therapy**

Use of an individualized eating plan to prevent, manage, or treat a health problem. It may address diabetes, kidney disease, food allergy, malnutrition, digestive disorders, or other needs and is tailored to the person's health and goals.

### **Dietary Assessment**

Systematic review of a person's food, drink, nutrient, and supplement intake using methods such as a

---

food diary, recall, or questionnaire. It helps identify deficiencies, excesses, patterns, and changes needed for health.

### **Dietary Calcium**

Calcium obtained from food or drinks, needed for bones, teeth, muscles, nerves, and blood clotting. Requirements vary with age and health; excess supplements may cause kidney stones or other problems.

### **Dietary Carbohydrate**

Carbohydrate in foods and drinks, including sugars, starches, and fiber. Digestible carbohydrate provides energy and affects blood glucose; fiber is only partly or not digested and supports bowel function.

### **Dietary Fat**

A nutrient made mainly of triglycerides and fatty acids that provides energy, supports cell membranes, and helps absorb vitamins A, D, E, and K. The type and amount of fat affect cardiovascular health.

### **Dietary Fats**

Fats obtained from food, including saturated, monounsaturated, polyunsaturated, and trans fats. They provide energy and support cell membranes and vitamin absorption; replacing saturated or trans fats with unsaturated fats is generally healthier.

### **Dietary Fats and Oils**

Edible fats and oils, including butter, margarine, and cooking oils, provide energy and essential fatty acids and help the body absorb vitamins A, D, E, and K. Their health effects depend on the type and amount consumed and the overall eating pattern.

### **Dietary Fiber**

Plant material that is not fully digested. It helps form stool, supports regular bowel movements, may improve cholesterol and glucose control, and can increase fullness.

### **Dietary Fibre**

Plant-derived carbohydrate that is not fully digested

in the small intestine. Soluble and insoluble fiber support bowel function and may improve cholesterol, glucose control, and fullness; increase intake gradually with adequate fluids unless medically restricted.

### **Dietary Inflammation**

Inflammatory activity that may be influenced by long-term eating patterns, excess body fat, and overall health. Diets rich in vegetables, legumes, whole grains, and unsaturated fats generally support lower inflammation than diets high in heavily processed foods.

#### **Alternate Names:**

Diet-Induced Inflammation; Diet-Associated Systemic Inflammation

### **Dietary Iron**

Iron from food or supplements, needed to make hemoglobin, muscle proteins, and enzymes. Too little can cause iron-deficiency anemia; too much can be toxic, especially in children or people with iron-storage disorders.

### **Dietary Phospholipid**

A food-derived fat containing phosphate that helps form cell membranes and transports fats in the blood. Its digestion and handling depend on food intake, absorption, and liver function.

### **Dietary Phosphorus**

Phosphorus from food, needed for bones, teeth, cell membranes, and energy use. People with reduced kidney function may need to limit highly absorbable sources to prevent high blood phosphorus.

### **Dietary Protein**

Protein from food that is digested into amino acids and small peptides. These support muscle and tissue repair, enzymes, hormones, immunity, and energy when needed.

### **Dietary Reference Intakes**

A set of nutrient-intake reference values used to plan and assess diets. They include the recommended allowance, adequate intake, estimated average re-

---

quirement, and tolerable upper intake level, which vary with age, sex, and life stage.

### **Dietary Sterol**

A cholesterol-like substance found in foods, including cholesterol and plant sterols. Absorption and metabolism affect blood cholesterol; plant sterols can reduce intestinal cholesterol absorption in some people.

### **Dietary Sugars**

Sugars are simple carbohydrates that provide energy but can raise blood glucose and contribute to dental decay or excess calories when consumed in excess.

### **Dietary Supplement**

A product taken by mouth to provide vitamins, minerals, herbs, amino acids, or other dietary ingredients. Supplements may be useful for a documented need but can interact with medicines or cause toxicity.

### **Dietary Supplementation**

Deliberate addition of nutrients or other substances to the diet to correct a deficiency, meet increased needs, or support a health goal. The dose and need should be assessed because excess can be harmful.

### **Dietary Supplements**

Products containing vitamins, minerals, herbs, amino acids, or other substances intended to add to the diet. They do not replace balanced food and may interact with medicines or cause adverse effects.

#### **Alternate Names:**

Nutritional Supplements

### **Dietetics**

The science and professional practice of using food and nutrition to promote health, prevent disease, and manage medical conditions through individualized dietary advice.

### **Diethylcarbamazine**

An antiparasitic medicine used mainly for lymphatic

filariasis and some other filarial infections. Fever, rash, swelling, or inflammation may occur as the immune system reacts to dying parasites.

### **Diethylhexylphthalate**

Another name for DEHP, a chemical added to some plastics to make them flexible. Repeated or high exposure may affect reproduction, development, or hormone function.

### **Diethylpropion**

A stimulant appetite suppressant used short term in selected weight-management plans. It can cause insomnia, anxiety, fast heartbeat, high blood pressure, misuse, and dependence; medical supervision is required.

### **Diethylstilbestrol**

A synthetic estrogen formerly given to some pregnant people. Exposure before birth can increase the risk of reproductive-tract abnormalities, fertility problems, and certain rare cancers later in life.

### **Dietitian**

A nutrition professional who assesses food and nutrient needs and provides individualized advice for health, disease prevention, and medical conditions. Professional titles and licensing rules vary by region.

### **Dietl Crisis**

Severe, episodic, colicky flank or abdominal pain accompanied by nausea, vomiting, and oliguria, caused by transient kinking or obstruction of the ureter when a mobile or ptotic kidney shifts position into the pelvis upon standing (nephroptosis), relieved upon lying supine.

#### **Alternate Names:**

Renal Ptosis Crisis; Dietl Syndrome

### **Dieulafoy Lesion**

An unusually large artery close to the surface of the stomach or another digestive organ that can suddenly cause severe bleeding, sometimes without a visible ulcer. It may produce vomiting of blood or black stools.

---

## Diff Dx

An abbreviation for differential diagnosis, the structured comparison of possible causes of a person's symptoms, signs, or test results.

## Differences Of Sex Development

Variations present from birth in chromosomes, gonads, hormones, or internal or external genital anatomy. Medical evaluation explains the variation, addresses health needs, and supports informed decisions and well-being.

## Differential Blood Count

A blood test measuring the numbers or percentages of different white blood cells, including neutrophils, lymphocytes, monocytes, eosinophils, and basophils. Results help assess infection, inflammation, allergy, and blood disorders.

## Differential Diagnosis

The process of comparing possible diseases or conditions that could explain the same symptoms or findings. Clinicians consider serious causes first, use examination and targeted tests, and revise the possibilities as new information appears.

## Differentiated Cancer

A cancer whose cells still resemble the normal tissue from which they arose. The degree of resemblance, called differentiation, helps pathologists describe tumor grade and may provide information about behavior.

## Differentiation

The process by which cells develop specialized structures and functions. In cancer pathology, it describes how closely abnormal cells resemble the normal tissue from which they originated.

## Differentiation Antigens

Molecules on a cell's surface that indicate its tissue type or stage of development. Laboratory tests for them help identify blood-cell disorders, classify cancers, and guide selected treatments.

## Difficult Airway

A situation in which a clinician may have trouble keeping a person breathing or placing a device into the airway. Causes include unusual anatomy, swelling, injury, illness, or limited neck movement and require careful planning.

## Difficult Intubation

Difficulty placing a breathing tube in the windpipe despite appropriate technique and equipment. Limited mouth or neck movement, unusual anatomy, swelling, injury, obesity, or a previous difficult intubation may increase risk.

## Difficult Urination

Trouble starting or maintaining urine flow, a weak stream, pain, straining, or a feeling of incomplete emptying. Causes include infection, blockage, prostate or pelvic disorders, medicines, and nerve or bladder problems.

## Difficult-to-Control Asthma

Asthma that remains poorly controlled despite appropriate treatment. Possible reasons include ongoing triggers, incorrect inhaler use, another illness, an incorrect diagnosis, severe airway inflammation, or a different response to medicines. Persistent symptoms need clinical review.

## Difficulty Speaking

Alternate name; see *Voice Disorders*.

## Difficulty Swallowing

Alternate name; see *Dysphagia*.

## Difficulty Swallowing (Dysphagia)

Trouble moving food, liquid, or saliva safely from the mouth to the stomach. Causes may involve the throat, esophagus, nerves, muscles, or obstruction; choking, aspiration, weight loss, or progressive symptoms require prompt evaluation.

## Difficulty Urinating

Trouble starting, maintaining, or emptying urine, often described as hesitancy, weak stream, straining,

---

or incomplete emptying; causes include obstruction, infection, medicines, and nerve or muscle disorders.

### **Difficulty Walking**

Trouble walking safely or normally because of pain, weakness, poor balance, numbness, joint disease, neurologic illness, or heart or lung limitations. Sudden onset requires urgent assessment.

### **Diffuse**

Spread broadly or throughout an area rather than limited to one small, clearly defined focus. The term may describe a disease, pain, swelling, imaging finding, or other process.

### **Diffuse Alopecia**

Alternate name; see *Diffuse Hair Loss*.

### **Diffuse Alveolar Hemorrhage**

A serious condition in which blood leaks into many of the lung's air sacs. It may cause coughing up blood, breathlessness, low oxygen, falling blood counts, and widespread shadows on a chest scan. Bronchoscopy with increasingly bloody samples can support the diagnosis, but the full clinical picture and cause must be assessed.

#### **Alternate Names:**

DAH; Pulmonary Alveolar Hemorrhage

### **Diffuse Astrocytoma**

An older name for an infiltrating brain tumor made of astrocyte-like cells. Modern classification depends on molecular findings, including IDH status, which help determine diagnosis, outlook, and treatment.

### **Diffuse Axonal Injury**

Widespread damage to brain nerve fibers caused by rapid acceleration, deceleration, or rotation of the head. It can cause prolonged unconsciousness or severe brain dysfunction, and MRI may show the injury.

### **Diffuse Esophageal Spasm**

A disorder in which the esophagus contracts too early or unevenly, causing intermittent chest pain

and difficulty swallowing liquids or solids. Testing helps distinguish it from blockage, reflux, and heart disease.

### **Diffuse Hair Loss**

A generalized, non-scarring reduction in hair density or widespread shedding affecting the entire scalp, distinguishing it from focal or patchy alopecia. Major causes include telogen effluvium following physiologic stress or childbirth, anagen effluvium induced by cytotoxic chemotherapy, diffuse androgenetic alopecia, systemic endocrinopathies such as hypothyroidism, and nutritional deficiencies in iron or zinc.

#### **Alternate Names:**

Diffuse Alopecia; Generalized Alopecia; Diffuse Effluvium

### **Diffuse Idiopathic Skeletal Hyperostosis**

A condition in which extra bone forms along spinal ligaments and where tendons attach to bone. It can cause stiffness, pain, swallowing difficulty, or fractures after minor injury.

#### **Alternate Names:**

DISH; Forestier Disease

### **Diffuse Large B-Cell Lymphoma**

A fast-growing cancer of B lymphocytes, a type of white blood cell. It may cause painless swollen lymph nodes, fever, night sweats, or weight loss and usually requires prompt specialist treatment.

### **Diffuse Midline Glioma**

An aggressive brain or spinal-cord cancer arising in a midline structure, often in children. Symptoms depend on location and may include weakness, movement problems, vision changes, vomiting, or seizures.

#### **Alternate Names:**

DMG

### **Diffuse Parenchymal Lung Disease**

Alternate name; see *Interstitial Lung Disease*.



---

## Diffuse Parenchymal Lung Diseases

Alternate index label; see *Interstitial Lung Disease*.

## Diffuse Proliferative Glomerulonephritis

A pattern of glomerular inflammation in which many glomeruli show increased cellularity and tissue injury. It often produces hematuria, proteinuria, hypertension, and reduced kidney function and may result from immune-complex disease such as lupus or infection.

## Diffuse Scleroderma

A form of systemic sclerosis in which skin thickening extends beyond the hands and face onto the trunk or limbs. It carries a higher risk of lung, kidney, heart, blood-vessel, and digestive complications.

## Diffusing Capacity

The ability of the lungs to move oxygen and other gases from the air sacs into the blood. It may be reduced by emphysema, interstitial lung disease, pulmonary vascular disease, or anemia.

## Diffusing Capacity Of The Lung

A measure of how efficiently gas crosses from the lung air sacs into nearby blood vessels. It is commonly measured as DLCO and helps evaluate lung tissue, blood vessels, and hemoglobin-related problems.

## Diffusion

Movement of molecules from an area where they are more concentrated to an area where they are less concentrated, caused by their random motion. It helps exchange gases and substances in the body.

## Diffusion Anisotropy

A difference in the direction water moves through tissue, especially along nerve fibers in the brain. MRI can use this pattern to study nerve-fiber structure and injury.

## Diffusion Capacity

A measure of how efficiently gases pass from lung air sacs into blood. It is commonly reported as DLCO

and may be reduced in interstitial lung disease, emphysema, pulmonary vascular disease, or anemia.

## Diffusion Restriction

An area where water movement appears reduced on a special MRI sequence. It can occur with an acute stroke, abscess, densely packed tumor, seizure, or other problem and must be interpreted with the full examination.

## Diffusion Tensor Imaging

An MRI technique that maps the direction of water movement along nerve fibers. It helps show the paths and condition of important nerve connections in the brain.

## Diffusion-Weighted Imaging

An MRI technique that shows how water molecules move within tissue. Areas with restricted movement may occur in acute stroke, abscess, densely packed tumors, seizures, or other conditions.

## Diffunisal

A salicylate nonsteroidal anti-inflammatory medicine used for pain and inflammation. It can cause stomach irritation or bleeding, kidney injury, fluid retention, allergic reactions, or cardiovascular problems.

## Digastric Muscle

A two-part muscle beneath the jaw that helps open the mouth, lower the jaw, and raise the hyoid bone during swallowing and speech.

## DiGeorge Syndrome

A genetic disorder usually caused by loss of a small part of chromosome 22. It can affect the heart, immune system, calcium regulation, palate, hearing, learning, and development.

### Alternate Names:

22q11.2 Deletion Syndrome

## Digestion

The process of breaking food into smaller substances that the body can absorb and use. It begins in

---

the mouth and continues through the stomach and intestines with movement, acid, bile, and digestive enzymes.

### **Digestive Diseases**

Disorders affecting the digestive tract, liver, gallbladder, or pancreas. Examples include reflux, ulcers, infections, inflammatory bowel disease, blockage, malabsorption, and cancer.

### **Digestive Disorder**

A disease affecting the digestive tract or the process of digestion. It may cause abdominal pain, nausea, reflux, diarrhea, constipation, bleeding, vomiting, or poor nutrient absorption.

### **Digestive Health**

Healthy function of the digestive tract in breaking down food, absorbing nutrients and water, and removing waste. It is influenced by diet, fluids, medicines, movement, illness, and the gut lining and microbiome.

#### **Alternate Names:**

Gut Health

### **Digestive System**

The gastrointestinal tract and accessory organs that mechanically and chemically process food, absorb nutrients and water, and eliminate solid waste. It includes the mouth, pharynx, esophagus, stomach, small and large intestines, rectum, anus, liver, gallbladder, and pancreas.

### **Digestive Tract**

The continuous hollow passage through which food travels from the mouth to the anus. It includes the throat, esophagus, stomach, small intestine, large intestine, rectum, and anus.

#### **Alternate Names:**

Gastrointestinal Tract; Alimentary Canal

### **Digit**

A finger or toe. Most fingers and toes contain three small bones, while the thumb and great toe usually contain two; muscles, tendons, nerves, and blood

vessels allow movement and sensation.

### **Digital Anorectal Examination**

A physical examination in which a clinician uses a lubricated, gloved finger to evaluate the anal canal, distal rectum, sphincter muscle tone, and adjacent structures such as the prostate.

#### **Alternate Names:**

DARE; Digital Rectal Examination; DRE; Rectal Exam

### **Digital Breast Tomosynthesis**

A breast X-ray test that takes pictures from several angles and combines them into thin, layered images. It can show some breast findings more clearly than a single flat image.

### **Digital Clubbing**

Enlargement and rounding of the fingertips with curved nails and loss of the normal angle at the nail base. It may be associated with lung, heart, digestive, liver, or other diseases and needs evaluation.

### **Digital Fibrokeratoma**

A rare, harmless, firm skin growth usually found on a finger or toe. It may resemble a wart; removal or microscopic examination may be needed when its appearance is unusual or it causes pain or interference.

### **Digital Ischemia**

Too little blood flow to a finger or toe, causing pain, pallor, coolness, numbness, or tissue death. It may result from arterial blockage, severe Raynaud phenomenon, inflammation, or an embolus and can be an emergency.

### **Digital Mammography**

Breast imaging that records X-ray pictures electronically. It helps detect masses, distortions, and calcium deposits and allows images to be enlarged, adjusted, stored, and compared.

### **Digital Mucous Cyst**

A usually harmless fluid-filled lump near the last

---

joint of a finger, often linked to osteoarthritis. It may press on the nail, cause pain, or occasionally become infected.

### **Digital Papillary Adenocarcinoma**

A rare cancer of a sweat gland, usually on a finger, hand, or foot. It often appears as a slow-growing lump but can return locally or spread to other organs.

### **Digital Radiography**

X-ray imaging in which the picture is captured electronically instead of on film. The image can be viewed, stored, adjusted, and shared on a computer.

### **Digital Rectal Exam**

A physical examination using a lubricated, gloved finger to feel the anal canal and rectum and, when relevant, the prostate. It may detect tenderness, bleeding, lumps, hemorrhoids, or muscle problems.

### **Digital Rectal Examination**

An examination in which a clinician gently feels inside the rectum with a lubricated, gloved finger. It may check for pain, bleeding, lumps, hemorrhoids, prostate changes, or muscle problems after explanation and consent.

### **Digital Subtraction Angiography**

An X-ray study in which computer processing removes background structures from contrast images so blood vessels are clearer. It can diagnose narrowing or blockage and guide selected vascular procedures.

### **Digitalis Preparation**

A preparation containing a cardiac glycoside such as digoxin or digitoxin. It can strengthen heart contraction and slow electrical conduction, but the dose range is narrow and toxicity can cause digestive, visual, or rhythm problems.

### **Digitalis Toxicity**

Harmful effects from digitalis or related cardiac glycosides, including nausea, vomiting, visual changes,

confusion, slow or fast abnormal heart rhythms, and high potassium. Kidney disease and drug interactions increase risk.

### **Digitate Warts**

Finger-like or thread-like warts, usually on the face or neck, caused by human papillomavirus infection. They are generally harmless but can spread or recur.

#### **Alternate Names:**

Verruca Digitata

### **Digitoxin**

A cardiac glycoside that can strengthen heart contraction and slow electrical conduction. Its safe and toxic doses are close together, so abnormal rhythms, digestive symptoms, visual changes, and drug interactions are important concerns.

### **Digoxin**

A cardiac medicine used in selected people with heart failure or atrial fibrillation. It can strengthen contraction and slow the heart rate, but kidney function, electrolytes, other medicines, and blood levels affect toxicity risk.

### **Digoxin Immune Fab**

Antibody fragments given by injection that bind digoxin and related cardiac glycosides, reducing their effects. They are used for life-threatening abnormal rhythms, severe toxicity, or a potentially very large ingestion.

### **Digoxin Test**

A blood test measuring digoxin concentration to help guide dosing or assess possible toxicity. The result must be interpreted with timing, symptoms, kidney function, electrolytes, and interacting medicines; the number alone does not prove toxicity.

### **Digoxin Toxicity**

Poisoning from digoxin or a related cardiac glycoside. Nausea, vomiting, confusion, visual changes, slow or fast abnormal rhythms, and dangerous potassium changes may occur; severe cases require emergency treatment.

---

**Alternate Names:**

Digoxin Toxicity

**Dihydrocodeine**

An opioid medicine used for selected pain or cough conditions. It can cause constipation, nausea, drowsiness, dependence, dangerous breathing depression, and overdose, especially with alcohol or other sedatives.

**Dihydroergotamine**

An ergot medicine used for selected acute migraine attacks. It narrows certain blood vessels and must be avoided in several vascular, heart, liver, kidney, and pregnancy-related conditions.

**Dihydropyrimidine Dehydrogenase Deficiency (DPYD-Related)**

Reduced activity of dihydropyrimidine dehydrogenase, usually caused by harmful variants in the DPYD gene. It can impair breakdown of fluorouracil and related medicines and greatly increase the risk of severe or life-threatening toxicity.

**Dihydrotestosterone**

A potent androgen made from testosterone. It contributes to development of male external genitalia, facial and body hair, prostate growth, and other androgen effects.

**Dihydroxyacetone**

A three-carbon sugar involved in metabolism. Its phosphate form is an intermediate used when cells break down glucose or make glucose.

**Dilantin Overdose**

Taking too much phenytoin can cause involuntary eye movements, poor coordination, slurred speech, drowsiness, confusion, low blood pressure, abnormal heart rhythms, seizures, or coma.

**Dilatation**

Widening or expansion of a hollow organ, blood vessel, duct, opening, or other body structure. It may be normal, caused by disease, or deliberately

produced during a procedure.

**Dilate**

To become wider or make a passage, blood vessel, opening, or body structure wider. Dilation may occur naturally, result from medicines or disease, or be created during a procedure.

**Alternate Names:**

Expand; Enlarge

**Dilated Cardiomyopathy**

A heart-muscle disease in which one or more chambers enlarge and pump weakly. Causes include inherited disease, infection, toxins, medicines, pregnancy, blocked arteries, and inflammation; it can lead to heart failure or abnormal rhythms.

**Alternate Names:**

DCM; Congestive Cardiomyopathy

**Dilated Cardiomyopathy Prognosis**

Outlook varies with heart-pumping strength, symptoms, cause, abnormal rhythms, other illnesses, and response to treatment. Medicines and devices can improve function for some people; severe disease may require advanced heart-failure care.

**Dilated Pore of Winer**

A benign, enlarged opening of a hair follicle that appears as a solitary black or dark keratin-filled pore, usually on the face, neck, or trunk of an adult. It may resemble a skin tumour; complete removal is diagnostic and usually curative.

**Dilation**

Widening or expansion of a hollow organ, blood vessel, duct, pupil, opening, or other body structure. It may be normal, caused by disease, or deliberately produced during a procedure.

**Dilation and Curettage**

A procedure that opens the cervix and removes tissue from inside the uterus using suction or an instrument. It may treat miscarriage or retained pregnancy tissue, investigate bleeding, or remove abnormal tissue; risks include bleeding, infection,

---

and perforation.

### **Dilation And Evacuation**

A procedure that opens the cervix and removes pregnancy tissue from the uterus using suction and instruments. It is performed by trained clinicians for selected pregnancy-related indications, with risks including bleeding, infection, and uterine injury.

### **Dilator**

An instrument or device used to widen a body opening or passage gradually or temporarily. It may be used during examinations, surgery, childbirth care, dental treatment, or treatment of a narrowing.

### **Diltiazem**

A calcium-channel blocker used for high blood pressure, angina, and heart-rate control in some abnormal rhythms. It can slow the heart, lower blood pressure, cause constipation, or worsen heart failure in susceptible people.

### **Dilute Russell Viper Venom Time**

A blood-clotting test used mainly to detect lupus anticoagulant, an antibody associated with antiphospholipid syndrome. Anticoagulant medicines and other clotting problems can affect the result, so confirmatory testing is needed.

### **Dilution**

Lowering the concentration of a substance by adding solvent or another fluid. Accurate dilution is important when preparing medicines, laboratory reagents, standards, or specimens.

### **Dilution Anemia**

A pseudoanemia caused by an expansion of plasma volume out of proportion to red blood cell mass, resulting in reduced hemoglobin concentration and hematocrit despite normal total circulating erythrocyte volume, typically seen during pregnancy and aggressive intravenous fluid administration.

#### **Alternate Names:**

Hypervolemic Anemia; Hemodilution

### **Dimenhydrinate**

An antihistamine used to prevent or treat motion sickness and nausea. It can cause drowsiness, dry mouth, blurred vision, constipation, confusion, or difficulty urinating.

### **Dimenhydrinate Overdose**

Taking too much dimenhydrinate can cause severe drowsiness or agitation, confusion, hallucinations, seizures, abnormal heart rhythms, breathing depression, or coma and requires urgent medical assessment.

### **Dimercaprol**

An injected medicine that binds certain toxic metals, including arsenic, mercury, and gold, so the body can remove them. It can cause high blood pressure, fast heartbeat, nausea, pain, and other adverse effects.

### **Dimesna**

A sulfur-containing compound studied as a protective or detoxifying medicine, including in some chemotherapy-related settings. Its clinical role depends on the specific treatment and available evidence.

### **Dimethoate**

An organophosphate insecticide that blocks acetylcholinesterase. Poisoning can cause sweating, excess saliva, vomiting, diarrhea, small pupils, muscle weakness, breathing problems, seizures, or coma.

### **Dimethoxon**

A poisonous breakdown product of some organophosphate pesticides. It can cause sweating, excess saliva, vomiting, diarrhea, small pupils, muscle twitching, weakness, breathing difficulty, or seizures and requires urgent poisoning advice.

### **Dimethyl Fumarate**

An oral medicine used for multiple sclerosis and selected skin conditions. It can cause flushing, digestive symptoms, low white-cell counts, liver injury, allergic reactions, or rare serious infections.

---

### Dimethyl Fumarate Allergy

An allergic or hypersensitivity reaction to dimethyl fumarate, a medicine used for some forms of multiple sclerosis. It may cause rash, itching, swelling, breathing difficulty, or other immune symptoms and requires medical assessment.

### Dimethyl Sulfoxide

An organic solvent used in laboratories and some specialized medical or investigational applications. Contact or ingestion can irritate tissues, and it may carry other substances through the skin.

### Dimethyl Sulphoxide

Another spelling of dimethyl sulfoxide, an organic solvent used in laboratories and selected medical or investigational applications. Exposure may cause irritation, a characteristic odor, or other adverse effects.

### Dimethylhydrazines

A group of highly toxic chemicals formerly used mainly in rocket fuel. Exposure can irritate or burn tissues and damage the nervous system, liver, blood, or other organs.

### Dimetilan

A poisonous carbamate insecticide that disrupts nerve signals by blocking acetylcholinesterase. Exposure can cause sweating, vomiting, diarrhea, small pupils, weakness, breathing difficulty, seizures, or coma and needs urgent help.

### Diminished Ovarian Reserve

Fewer remaining eggs than expected for a person's age, which may reduce response to fertility treatment and shorten the time available for conception. Tests estimate reserve but cannot precisely predict natural fertility.

#### Alternate Names:

Decreased Ovarian Reserve; DOR

### Dimocillin

A penicillin-type antibiotic used historically or in selected settings for susceptible bacterial infections.

Choice depends on the organism, resistance testing, infection site, allergies, and local treatment guidance.

### Dimorphic

Existing in two distinct forms. Some fungi are dimorphic, growing as a mold in the environment and as a yeast in body tissues; this ability can contribute to infection.

### Dinophyceae

A group of aquatic microorganisms, including dinoflagellates. Some produce toxins during harmful algal blooms that can contaminate seafood or water and cause neurologic, digestive, or liver illness.

### Dinoprost

A prostaglandin medicine that stimulates uterine contraction and other smooth-muscle effects. It is used mainly in veterinary medicine and selected reproductive-health settings; pain, diarrhea, breathing difficulty, and blood-pressure changes may occur.

### Dinoprostone

A prostaglandin medicine used to soften the cervix and help start labor or manage selected pregnancy-related conditions. It requires obstetric monitoring because it can cause contractions that are too frequent, pain, fever, nausea, or fetal heart-rate changes.

### Dinucleoside Phosphates

Molecules containing two nucleosides joined by phosphate groups. They occur in nucleotide metabolism and may take part in cellular signaling, energy transfer, or DNA and RNA-related reactions.

### Dinutuximab

An antibody-based cancer medicine that attaches to a marker found on many neuroblastoma cells and helps the immune system destroy them. It can cause severe nerve pain, allergic reactions, fever, and fluid leakage from blood vessels.



---

### Diocetyl Sodium Sulphosuccinate

A surfactant and stool-softening medicine also called docusate sodium. It helps water mix with stool and is used in selected constipation settings, although evidence of benefit is limited.

### Diopter

A unit of optical power equal to the reciprocal of focal length in meters. Eyeglass and contact-lens prescriptions use diopters to state the strength needed to focus light clearly.

### Dioxin

A group of long-lasting environmental chemicals produced by some industrial processes and burning. They can build up in body fat; long-term exposure may affect development, reproduction, immunity, hormones, and cancer risk.

### Dipeptide

A molecule made of two amino acids joined together. Dipeptides are produced during protein digestion and metabolism and can be absorbed or used in further chemical reactions.

### Dipeptidyl Peptidase-4 Inhibitors

Oral diabetes medicines that prolong the action of natural gut hormones. They help the body release insulin after meals and reduce glucagon, usually without causing weight gain or low blood glucose when used alone.

### Dipetalonema Infection

Infection with a *Dipetalonema* parasitic worm, usually acquired through an arthropod bite. It may affect skin or tissue beneath the skin; diagnosis and treatment depend on the species and clinical findings.

### Diphallia

A rare birth condition in which the penis is partly or completely duplicated. Abnormalities of the urinary tract, anus, spine, or other organs may also occur and require specialist assessment.

### Diphenhydramine

An antihistamine with sedating and anticholinergic effects used in some allergy or sleep products. It can impair alertness and cause dry mouth, confusion, urinary retention, abnormal rhythms, or toxicity in overdose.

### Diphenhydramine Overdose

Taking too much diphenhydramine can cause severe drowsiness or agitation, confusion, hallucinations, seizures, abnormal heart rhythms, breathing depression, coma, or death and requires urgent medical assessment.

### Diphenoxylate

An opioid-related medicine that slows intestinal movement to reduce diarrhea, usually combined with atropine. Excessive use can cause drowsiness, constipation, dependence, breathing depression, or poisoning.

### Diphosphate

A chemical group containing two linked phosphate units. Diphosphate is involved in energy use, nucleotide formation, and other chemical reactions in cells.

### Diphosphonate

A chemical group found in compounds that can bind strongly to bone mineral. Some related medicines are used or studied for disorders of bone breakdown and calcium balance.

### Diphtheria

A serious infection caused by toxin-producing *Corynebacterium* bacteria. It can form a thick coating in the throat and cause airway blockage, heart inflammation, nerve damage, or death; vaccination prevents most cases.

### Diphtheria Toxin

A poisonous protein made by some *Corynebacterium* bacteria. It blocks protein production inside cells, causing local tissue injury and potentially severe heart, nerve, and airway complications.

---

### **Diphtheroid Bacillus**

A rod-shaped bacterium resembling *Corynebacterium* under the microscope. Many are normal skin bacteria or contaminants, but some can cause opportunistic infection in people with serious illness or medical devices.

### **Diphylobothriasis**

An intestinal infection caused by a fish tapeworm acquired from raw or undercooked freshwater fish. It may cause abdominal symptoms and, rarely, vitamin B12 deficiency and anemia.

### **Diphylobothrium Latum**

A fish tapeworm acquired by eating raw or undercooked freshwater fish. It lives in the intestine and may cause abdominal symptoms or compete for vitamin B12, occasionally leading to anemia.

### **Diplacusis**

Hearing one sound as two different pitches or qualities. It often reflects unequal hearing between the ears or inner-ear dysfunction and may follow infection, noise injury, or other ear disease.

### **Diplegia**

Severe weakness or paralysis affecting matching parts on both sides of the body. In spastic diplegia, the legs are affected more than the arms, often because of early brain injury.

### **Diplococcus**

A bacterium that appears as pairs of round cells under a microscope. The term describes shape, not one species; some diplococci cause infections in people.

### **Diploid Cell**

A cell containing two complete sets of chromosomes, one inherited from each parent. Most human body cells are diploid; eggs and sperm normally contain only one set.

### **Diploidy**

Having two complete sets of chromosomes, usually

one from each parent. This is the normal chromosome state of most human body cells.

### **Diplopia**

Seeing two images of one object. It may occur with one eye or only when both eyes are open; causes include dry eye, cataract, eye misalignment, nerve disease, muscle disease, or brain disease.

#### **Alternate Names:**

Double Vision

### **Dipole**

A pair of equal and opposite electrical or magnetic charges separated by a distance. The concept is used in physiology, medical imaging, and physics to describe electrical or magnetic fields.

### **Dipper**

A person whose blood pressure falls during sleep, usually by about 10 percent or more compared with daytime readings. The pattern is measured with a portable 24-hour monitor and may help assess cardiovascular risk.

### **Diprosopus**

A rare birth defect in which part or all of the face is duplicated. It usually occurs with severe abnormalities of the brain, heart, or other organs and often causes major developmental problems.

### **Dipsosis**

Excessive thirst or a persistent urge to drink. Causes include dehydration, high blood glucose, high blood concentration, medicines, dry mouth, or disorders affecting thirst regulation.

### **Dipstick Device**

A strip or small device containing chemical pads that change color when placed in a specimen, commonly urine. It provides a rapid screen for substances such as blood, protein, glucose, or infection markers.

### **Diptera**

The insect order containing flies and mosquitoes. Some species spread infections through bites or con-

---

tamination, while others can cause infestations such as myiasis.

### **Dipylidium Caninum**

A tapeworm of dogs and cats that can occasionally infect people, especially children, after an infected flea is swallowed. Infection may cause mild digestive symptoms or visible worm segments in stool.

### **Dipyridamole**

A medicine that reduces platelet clumping and widens blood vessels. It is used in selected stroke-prevention plans and heart stress testing; headache, dizziness, low blood pressure, and bleeding may occur.

### **Direct Acting Antivirals**

A modern class of oral antiviral pharmacological agents that target specific non-structural viral proteins (such as NS3/4A protease, NS5A replication complex, and NS5B polymerase inhibitors in hepatitis C virus), achieving sustained virologic cure rates exceeding 95 percent.

#### **Alternate Names:**

DAAs

### **Direct Action**

An effect produced through an intervention's immediate target, without requiring an intermediate pathway or mediator. The term may describe a medicine, hormone, nerve signal, or other biological process.

### **Direct Antiglobulin Test**

A blood test that detects antibodies or complement attached to a person's red blood cells. A positive result may occur with autoimmune hemolysis, a transfusion reaction, or hemolysis affecting a newborn.

### **Direct Coombs Test**

Another name for the direct antiglobulin test, which detects antibodies or complement attached to red blood cells. It can help investigate autoimmune hemolysis, transfusion reactions, medicine-related hemolysis, or newborn hemolysis.

### **Direct Hernia**

An inguinal hernia in which abdominal tissue pushes through a weak area in the back wall of the groin canal. It may cause a groin lump or discomfort and can occasionally become trapped.

### **Direct Inguinal Hernia**

An abdominal wall hernia in which peritoneal sac contents protrude directly forward through a weakened transversalis fascia within Hesselbach triangle, situated medial to the inferior epigastric vessels.

### **Direct Oral Anticoagulants**

Oral blood thinners that directly block a clotting protein, such as thrombin or factor Xa. They prevent and treat clots but can cause bleeding and require attention to kidney function, interactions, and procedures.

### **Direct Transmission**

Spread of an infectious organism through immediate contact or short-range respiratory droplets from an infected person or animal, without an intermediate object, food, water source, or vector.

### **Direct Veneer**

A tooth-colored composite layer shaped and hardened directly on a tooth to improve appearance or repair minor damage. It usually preserves more tooth than some alternatives but may stain, chip, or need replacement.

#### **Alternate Names:**

Direct Composite Veneer; Composite Dental Veneer

### **Directly Observed Therapy**

A healthcare worker or trained supporter watches a person take each dose of medicine. It can support completion of complex treatment, especially for tuberculosis, and helps detect adverse effects or missed doses.

### **Dirofilaria Repens**

A mosquito-transmitted parasitic worm that rarely infects people. It usually causes a single lump beneath the skin or near the eye because humans are

---

an accidental host; it does not usually mature or spread in the body.

### **Dirt-Eating**

Repeatedly eating soil or other nonfood substances, a form of pica. It may be associated with iron deficiency, pregnancy, nutritional problems, or mental health conditions and can expose a person to infection or toxins.

### **Dirty Necrosis**

Dead tumor tissue mixed with broken cell material and many white blood cells. It is often seen in rapidly growing cancers and is interpreted by a pathologist with the rest of the tissue sample.

### **Disability**

A difficulty with body function, activity, or participation caused by a health condition and influenced by the person's environment. Disability may be temporary or lifelong and varies in severity and support needs.

### **Disability Evaluation**

An assessment of how a health condition affects physical, mental, or daily functioning. History, examination, records, and appropriate tests may inform decisions about work, benefits, insurance, rehabilitation, or support.

### **Disability-Adjusted Life Year**

A population-health measure combining years lost from early death with years lived in less than full health. One DALY represents one lost year of healthy life under the method used.

### **Disaccharidase**

An intestinal enzyme that breaks a double sugar into simple sugars that can be absorbed. Lactase, for example, breaks lactose in milk into glucose and galactose.

### **Disaccharide**

A carbohydrate made of two linked simple sugars, such as lactose, sucrose, or maltose. It must be split

by digestive enzymes before the intestine can absorb its component sugars.

### **Disarticulation**

Amputation performed through a joint rather than by cutting through the shaft of a bone. The level is chosen according to the injury, disease, circulation, function, and possibility of healing.

### **Disc**

A thin, round structure. In the spine, an intervertebral disc is a tough, flexible cushion between neighboring vertebrae that absorbs load and allows movement.

### **Discectomy**

Surgery to remove part of a damaged spinal disc that is pressing on a nerve. It may relieve leg or arm pain and weakness, but risks include infection, bleeding, nerve injury, and recurrent disc problems.

### **Discitis**

Inflammation or infection of a spinal disc, often involving the nearby vertebral bones. It may cause severe localized back or neck pain, stiffness, fever, or reduced movement and usually requires imaging and medical evaluation.

### **Discography**

An imaging procedure in which contrast is injected into a spinal disc to examine it and see whether the injection reproduces pain. Because it has limitations and risks, it is used selectively.

### **Discoid**

Coin-shaped or round. In medicine, it often describes a skin lesion or rash, such as the round plaques of discoid lupus; the associated condition must be identified separately.

### **Discoid Eczema**

A long-lasting type of eczema that causes itchy, round or oval patches of inflamed skin, which may blister, ooze, crust, or become dry and cracked. It can affect different parts of the body and sometimes

---

becomes infected; it is also called nummular eczema.

**Alternate Names:**

Nummular Eczema; Discoid Dermatitis; Nummular Dermatitis

**Discoid Lupus**

A long-lasting form of skin lupus causing round, inflamed, scaly plaques. Lesions may scar, change skin color, or cause permanent hair loss on the scalp; some people later develop systemic lupus.

**Discoid Lupus Erythematosus**

A chronic autoimmune skin disease causing persistent round, inflamed plaques that may scale, scar, or change pigment. It commonly affects the face, ears, and scalp and may cause permanent hair loss.

**Discoloration**

An unusual change in the color of skin, mucous membranes, nails, or another tissue. Causes include bruising, low oxygen, jaundice, inflammation, infection, or altered pigment; sudden, painful, widespread, or unexplained change needs assessment.

**Discomfort**

An unpleasant physical or emotional sensation that may be milder or less clearly located than pain. Its importance depends on the site, duration, associated symptoms, and underlying cause.

**Discordance**

Lack of agreement between findings, measurements, genetic results, or the medical conditions of two related people. The significance depends on what is being compared and the clinical setting.

**Discordant**

Not agreeing, matching, or corresponding. In medicine it may describe conflicting test results or different medical findings in related people, twins, or paired organs.

**Discordant Couple**

A couple whose partners have different results or

medical statuses relevant to their care, such as one partner having HIV and the other not. Counseling can reduce transmission risk and support reproductive decisions.

**Disease**

An abnormal condition that affects the body or mind and changes normal structure or function. Diseases may be infectious, inflammatory, inherited, degenerative, metabolic, immune-related, or cancerous.

**Disease Progression**

Change in a disease over time, often meaning worsening. It is assessed through symptoms, examination, laboratory results, imaging, function, complications, and response to treatment.

**Disease Reporting**

Notification of specified diseases, infections, or health events to public-health authorities. Reports help detect outbreaks, protect contacts, arrange follow-up, and monitor health patterns; requirements vary by location.

**Disease Reservoir**

A person, animal, environment, or material in which an infectious organism normally lives and from which it can spread to a susceptible host.

**Disease Response**

The body's biological or clinical reaction to a disease or its treatment. It is assessed using symptoms, signs, laboratory results, imaging, physical function, or survival.

**Disease Surveillance**

Ongoing collection, analysis, interpretation, and reporting of health information to detect, monitor, and respond to disease patterns, outbreaks, and changes in population health.

**Disease Susceptibility**

Increased likelihood of developing a disease because of genes, age, immune status, exposures, behavior, other illnesses, or environmental factors. Suscep-

---

tibility raises risk but does not mean disease is certain.

### **Disease Vector**

A living carrier, commonly a mosquito, tick, flea, or other arthropod, that transfers an infectious organism between hosts. The vector may transmit infection without becoming ill itself.

### **Disease-Free Survival**

The length of time after treatment during which a person remains alive without detectable evidence of the disease. The exact definition and assessment method depend on the disease and study.

### **Disfigurement**

A visible change that substantially alters appearance, caused by injury, disease, surgery, a birth difference, or scarring. Its physical, emotional, and social effects vary between people.

### **Disinfectant**

A chemical or physical agent used on nonliving surfaces to destroy or inactivate many microorganisms. Disinfectants are not necessarily safe for skin, wounds, food, or internal use.

### **Disinfection**

Cleaning that destroys or inactivates most disease-causing microorganisms on nonliving surfaces. It may not destroy bacterial spores; the required level depends on the object, organism, and intended use.

### **Disk**

A flat, round structure. In the spine, an intervertebral disk lies between neighboring vertebrae, absorbs shock, and helps the back move.

#### **Alternate Names:**

Disc

### **Disk Battery Ingestion**

Ingestion of a button or coin battery, which can cause rapid electrical and chemical tissue injury, especially when lodged in the esophagus. It is a medical emergency requiring urgent localization and

removal.

### **Disk Diffusion Susceptibility Test**

A laboratory test of bacterial antibiotic sensitivity. Drug-containing disks are placed on growing bacteria, and the clear areas around them are measured and compared with standards to guide treatment.

### **Discectomy**

Surgery to remove part or all of a damaged spinal disk pressing on a nerve or the spinal cord. It may relieve pain or weakness caused by disk herniation; risks include infection, bleeding, nerve injury, and recurrence.

### **Diskitis**

Inflammation or infection of a spinal disk and nearby vertebral bone. It commonly causes severe localized back pain, stiffness, fever, or reluctance to move and requires medical assessment and imaging.

### **Dislocated Ankle**

Alternate name; see *Ankle Dislocation*.

### **Dislocated Elbow**

Loss of normal alignment of the elbow joint, usually after a fall or collision. It causes severe pain, deformity, and limited movement and needs urgent reduction with checks for fractures, nerve injury, and impaired circulation.

### **Dislocated Hip**

Displacement of the upper thighbone from its hip socket, usually after major trauma or around an artificial joint. It is an emergency because nearby nerves and the blood supply to the bone may be damaged.

### **Dislocated Knee**

Loss of alignment between the thighbone and shinbone, usually after severe injury. Major artery or nerve damage may be hidden, so urgent reduction and circulation checks are needed even if the knee returns to position.



---

## Dislocated Shoulder

Displacement of the upper arm bone from the shoulder socket. It causes pain, deformity, and inability to move normally and requires reduction with assessment of nerves, blood vessels, and fractures.

## Dislocation

Complete loss of normal contact between the surfaces of a joint, usually from injury or disease. It can cause severe pain and deformity and may occur with fractures or nerve and blood-vessel injury.

## Dislocation Of The Lens

Partial or complete movement of the eye's natural lens away from its normal position because its supporting fibers are weak or torn. It may cause blurred or double vision, eye pain, or glaucoma.

## Disofenin Scan

A nuclear-imaging test that follows bile production and flow through the liver, gallbladder, and bile ducts. It can help detect blockage, leakage, inflammation, or selected causes of abdominal pain.

## Disopyramide

An antiarrhythmic medicine used selectively for certain abnormal heart rhythms. It can slow electrical conduction and weaken heart contraction and may cause dry mouth, urinary retention, low blood pressure, or new arrhythmias.

## Disorder

A disturbance of normal body or mind structure or function that causes symptoms, impairment, or important health risk. The term does not by itself indicate how severe the problem is.

## Disordered Thinking

A disturbance in the organization, clarity, logic, or flow of thoughts. It may occur with mental illness, brain disease, delirium, substance use, medicines, or other medical conditions.

## Disorders of Fluid, Electrolyte, and

## Acid-Base Balance

Disorders that disturb body-fluid balance, electrolyte concentrations, or blood acidity. They may arise from kidney, heart, endocrine, gastrointestinal, medication, or critical-illness problems and can affect the brain, muscles, and heart.

## Disorders of Oral Pigmentation

Conditions causing focal or diffuse dark, brown, blue, or black discoloration of the mouth lining. Causes include normal variation, trauma, inflammation, medicines, smoking, endocrine or systemic disease, and melanocytic or other tumors; new, changing, solitary, or symptomatic lesions need oral examination.

## Disorganized Symptoms in Schizophrenia

Markedly disorganized speech, behavior, or emotional expression occurring in schizophrenia. Modern classifications describe these features within schizophrenia-spectrum disorders rather than as a separate subtype.

### Alternate Names:

Disorganized Schizophrenia (deprecated historical subtype label)

## Disorientation

Difficulty knowing the time, place, situation, or one's identity. It may occur with delirium, infection, abnormal body chemistry, intoxication or withdrawal, brain injury, or dementia; sudden onset needs urgent assessment.

## Dispensary

A place or service that stores, prepares, and supplies medicines and may provide basic medication advice. Safe dispensing includes checking the prescription, labeling the medicine, and explaining how to use it.

## Dispensed Amount

The quantity of medicine supplied to a patient. It is determined by the prescribed dose, treatment duration, formulation, refill rules, and applicable safety or legal requirements.

---

## Dispensing

The professional process of checking a prescription, selecting and preparing the medicine, labeling it, and supplying it with instructions for safe use.

## Displacement

Movement of an anatomical structure away from its normal position, as in a displaced fracture, joint injury, organ shift, or device migration. Effects depend on the structure involved and the amount of movement.

## Disruption

An interruption or breakdown of normal structure, function, continuity, or process. In medicine, it may describe damaged tissue, a broken barrier, disturbed development, or an interruption in the usual work of an organ or system.

## Disruptive, Impulse-Control, and Conduct Disorders

A group of disorders involving persistent problems with emotional or behavioral self-control that violate the rights of others, major age-appropriate rules, or social norms. Examples include oppositional defiant disorder, conduct disorder, intermittent explosive disorder, kleptomania, and pyromania.

## Dissect

To separate tissue layers or structures by cutting or careful blunt separation for surgery, examination, or anatomical study.

## Dissecting Aneurysm

Alternate name; see *Aortic Dissection*.

## Dissection

Separation of tissue layers, either during surgery or because of disease. In an artery, a tear in the inner lining lets blood split the wall layers and can block blood flow or cause the artery to rupture.

## Disseminate

To spread widely through the body, a tissue, or a population. Infection, cancer, inflammation, or

another process may disseminate rather than remain in one limited area.

## Disseminated Coccidioidomycosis

A fungal infection in which *Coccidioides* spreads beyond the lungs to skin, bones, joints, brain coverings, or other organs. It can be severe and is more likely in people with impaired immunity.

## Disseminated Cutaneous Blastomycosis

A *Blastomyces* infection that spreads from the lungs or another site to multiple areas of skin. It may also involve bones or other organs and requires antifungal treatment.

## Disseminated Gonococcal Infection

A systemic complication of *Neisseria gonorrhoeae* infection resulting from bacteremic dissemination from mucosal surfaces. It classically manifests as an arthritis-dermatitis syndrome (migratory polyarthralgias, tenosynovitis, and painless pustular or hemorrhagic skin lesions) or as acute purulent septic monoarthritis.

### Alternate Names:

DGI; Gonococcal Bacteremia; Gonococcal Arthritis-Dermatitis Syndrome

## Disseminated Intravascular Coagulation

A life-threatening disorder in which clotting is activated throughout the bloodstream. Small clots can injure organs while using up platelets and clotting factors, causing serious bleeding at the same time.

### Alternate Names:

DIC

## Disseminated Mycobacterium Avium–Intracellulare Infection

A widespread infection with *Mycobacterium avium* complex, usually in people with severe immune suppression. It may involve blood, bone marrow, lymph nodes, liver, spleen, or other organs.

## Disseminated Neuroblastoma

Neuroblastoma that has spread from its original site to distant organs, bone marrow, bones, liver, or

---

lymph nodes. Treatment and outlook depend on age, spread, tumor biology, and response to specialist care.

### **Disseminated Sclerosis**

An older name for multiple sclerosis, an immune-mediated disease in which areas of damage develop in the brain, spinal cord, or optic nerves and cause symptoms that vary over time.

### **Disseminated Tuberculosis**

Tuberculosis that has spread through the bloodstream or lymphatic system to multiple organs. It may cause fever, weight loss, fatigue, organ damage, meningitis, or widespread small lesions and needs urgent treatment.

### **Disseminated Varicella Infection**

Chickenpox infection that spreads beyond the skin and may involve the lungs, brain, liver, or other organs. It is more likely and more severe in adults, pregnant people, and those with weakened immunity.

### **Disseminated Zoster**

A severe, widespread reactivation of varicella-zoster virus characterized by greater than 20 cutaneous vesicles outside the primary and adjacent dermatomes, occurring predominantly in immunocompromised individuals and carrying a high risk of visceral organ dissemination (pneumonitis, encephalitis, hepatitis).

### **Dissociation**

A disruption in the usual connection between consciousness, memory, identity, emotions, perceptions, behavior, or sense of self. Mild forms may occur during daydreaming; severe or persistent forms can impair daily life.

### **Dissociative Amnesia**

Inability to recall important autobiographical information, often related to trauma or severe stress, beyond ordinary forgetfulness. Memory loss may involve a specific event, period, or broader part of a person's life and is not better explained by ordi-

nary forgetting, substances, a neurologic disorder, or another medical condition.

### **Dissociative Disorder**

Alternate name; see *Dissociative Disorders*.

### **Dissociative Disorders**

A group of mental disorders involving disruption in the usual integration of identity, memory, consciousness, perception, emotion, or behavior. Examples include dissociative amnesia, dissociative identity disorder, and depersonalization-derealization disorder. Temporary dissociative experiences can occur without constituting a disorder; a dissociative disorder causes significant distress or impairment.

### **Dissociative Identity Disorder**

A mental health disorder involving two or more distinct identity states and repeated gaps in memory for everyday events or personal information. It can cause changes in behavior, emotions, perception, and daily functioning.

### **Distal**

Farther from the point where a body part attaches or begins, or farther from the center of the body. For example, the hand is distal to the elbow.

### **Distal Convoluted Tubule**

The segment of the renal nephron located between the macula densa and the cortical collecting duct, responsible for calcium reabsorption under parathyroid hormone regulation and thiazide-sensitive sodium-chloride cotransport.

#### **Alternate Names:**

DCT

### **Distal Median Nerve Dysfunction**

Abnormal function of the median nerve near the forearm, wrist, or hand, causing numbness, tingling, pain, or weakness in its area of supply. Compression at the wrist is a common cause.

### **Distal Pancreatectomy**

Surgery removing the body or tail of the pancreas,

---

usually while leaving the head. It may be used for selected tumors, cysts, injury, or inflammation; diabetes and leakage from the pancreas are possible risks.

### **Distal Radius Fractures**

Breaks near the wrist end of the forearm bone, often after a fall. They may cause pain, swelling, deformity, or numbness; treatment depends on alignment, joint involvement, bone quality, and nerve or blood-vessel injury.

### **Distal Renal Tubular Acidosis**

A kidney disorder in which the kidneys cannot remove enough acid into the urine. Acid builds up in the blood and may cause low potassium, kidney stones, weak bones, muscle weakness, or poor growth in children.

### **Distal Splenorenal Shunt**

Surgery connecting the splenic vein to the left kidney vein to lower pressure in veins around the esophagus. It may reduce bleeding from varices while preserving some blood flow through the liver.

### **Distal Tubule**

A small kidney tube that adjusts the amount of salt, calcium, and water left in forming urine. Its activity helps regulate blood pressure and body chemistry and can be affected by some diuretic medicines.

### **Distamycins**

A group of antibiotic-like compounds that bind to DNA and have been studied for antimicrobial or anticancer effects. Their clinical use is limited and depends on the specific compound and evidence.

### **Distant**

Far from a reference point or original site. In cancer, distant disease means that cancer has spread to remote organs or lymph nodes rather than remaining near the primary tumor.

### **Distended**

Abnormally stretched, swollen, or enlarged by gas,

fluid, blood, tissue, or pressure. The term may describe the abdomen, bladder, stomach, bowel, veins, or another body structure.

### **Distension**

Enlargement or stretching of a body part or hollow organ, often from gas, fluid, urine, stool, blood, pressure, or blockage. Abdominal distension with severe pain, vomiting, fever, or inability to pass stool or gas needs prompt assessment.

### **Distention**

Enlargement or stretching of an organ, cavity, or body part, such as abdominal distention from gas, fluid, swelling, or blockage.

### **Distichia**

Growth of an extra row of eyelashes, usually from openings of glands along the eyelid. Misguided lashes may rub the eye and cause irritation, tearing, corneal scratches, or ulcers.

### **Distortion**

An abnormal change in the shape, structure, appearance, or representation of tissue or an image. New breast, facial, or visual distortion may indicate disease and should be assessed with appropriate examination or testing.

### **Distraction Osteogenesis**

A surgical method for growing new bone by slowly separating two bone sections after a controlled cut. A device gradually widens the gap to lengthen or reshape the jaw, limbs, or other bones.

### **Distress**

Significant physical or emotional suffering, discomfort, or strain related to illness, injury, treatment, or difficult circumstances. Severe distress may affect sleep, functioning, decisions, or safety and may need support.

### **Distributive Shock**

Shock caused by marked vasodilation and abnormal distribution of blood flow, producing low systemic

---

vascular resistance and inadequate effective tissue perfusion despite normal or high cardiac output early in the course. Sepsis, anaphylaxis, neurogenic shock, and some toxins are causes; skin may be warm initially, with hypotension, tachycardia or bradycardia, altered mentation, oliguria, and rising lactate.

### District Nurse

A community nurse who provides care in homes, clinics, or other local settings. Duties may include wound care, chronic-disease support, health education, medicine monitoring, prevention, and coordination with other services.

### Disulfiram Toxicity

Poisoning from excessive disulfiram, which may cause nausea, vomiting, abdominal pain, drowsiness, confusion, hypotension, seizures, coma, peripheral neuropathy, or liver injury. A separate disulfiram-alcohol reaction causes flushing, throbbing headache, vomiting, tachycardia, and hypotension; treatment is withdrawal of exposure and supportive care.

### Disulfiram-Like Reaction

An unpleasant reaction after alcohol is taken with disulfiram or another interacting substance. Flushing, headache, nausea, vomiting, palpitations, chest discomfort, or low blood pressure may occur.

### Disulfoton

A highly toxic organophosphate insecticide that inhibits acetylcholinesterase. Exposure can cause sweating, excess saliva, vomiting, diarrhea, muscle weakness, breathing failure, seizures, or coma.

### Dithiolethione

A sulfur-containing compound studied for activating cellular defense and detoxification pathways. It may influence inflammation and protection from chemical injury, but clinical uses depend on evidence for the specific compound.

### Ditiocarb

A term for dithiocarbamate compounds, some of which are used as pesticides or metal-binding chem-

icals. Toxic effects vary by compound and may involve the nervous system, liver, kidneys, or skin.

### Diuresis

Increased production of urine. It may result from excess fluid, high blood glucose, diabetes insipidus, medicines called diuretics, or other causes; excessive loss can lead to dehydration and electrolyte problems.

### Diuretic Drug

A medicine that makes the kidneys remove more sodium and water in urine. It is used for high blood pressure, heart failure, and swelling but can cause dehydration, low blood pressure, electrolyte changes, or kidney problems.

### Diuretic Resistance

A clinical failure to achieve expected therapeutic natriuresis, fluid excretion, and decongestion despite escalating doses of loop diuretics, commonly encountered in decompensated heart failure, nephrotic syndrome, and cirrhosis with refractory ascites.

### Diurnal

Relating to daytime or occurring during a daily 24-hour cycle. The term may describe symptoms, hormone levels, body functions, or patterns of activity and sleep.

### Diverticula

Alternate name; see *Diverticulosis*.

### Diverticular Bleeding

The painless, abrupt passage of large volumes of maroon or bright red blood per rectum resulting from transmural erosion or rupture of a weakened vasa recta mucosal arteriole at the neck of a colonic diverticulum; the most frequent cause of massive lower gastrointestinal hemorrhage in adults.

#### Alternate Names:

Diverticular Hemorrhage

### Diverticular Disease

A condition involving small pouches in the wall of

---

the colon. Diverticulosis means the pouches are present, while diverticular disease generally refers to symptoms or complications such as abdominal pain, altered bowel habits, bleeding, inflammation, infection, blockage, or perforation. Diverticulitis is inflammation or infection of one or more pouches and commonly causes lower-abdominal pain and fever.

### **Diverticulitis**

Inflammation of one or more small pouches in the wall of the colon. It usually causes steady pain in the lower-left abdomen, tenderness, fever, and changes in bowel movements. Severe cases may cause an abscess, a tear in the bowel, blockage, or a serious infection.

### **Diverticulosis**

The presence of multiple small, outward-protruding pouches (pseudodiverticula) formed by mucosal herniation through the muscular layer of the colon wall, most commonly occurring in the sigmoid and descending colon. Associated with aging and low-fiber diets, it is typically asymptomatic but represents the leading cause of sudden, painless, massive lower gastrointestinal bleeding (diverticular hemorrhage) when adjacent nutrient blood vessels rupture. It can also progress to diverticulitis if a pouch becomes obstructed, inflamed, or infected. Prevention and management focus on a high-fiber diet, hydration, and maintaining regular bowel habits.

#### **Alternate Names:**

Colonic Diverticulosis; Diverticular Pouching; Diverticular Disease of the Colon

### **Diverticulosis And Diverticulitis**

Diverticulosis means small pouches have formed in the colon wall. Diverticulitis occurs when one or more pouches become inflamed or infected, causing abdominal pain, fever, and bowel changes.

### **Diverticulum**

A pouch-like projection from the wall of a hollow organ, especially the intestine. It may cause no symptoms or become inflamed, infected, blocked, bleed, or rarely perforate.

### **Dix-Hallpike Maneuver**

A bedside test for a common inner-ear cause of brief positional vertigo. The clinician moves the head and body into a specific position and watches for dizziness and characteristic eye movements.

### **Dizygotic Twin**

One of a pair of fraternal twins formed when two separate eggs are fertilized by two separate sperm. The twins share about half their genetic variation on average, like ordinary siblings.

### **Dizygotic Twins**

Twins formed when two separate eggs are fertilized by two separate sperm. They are genetically like ordinary siblings and may have separate or shared-appearing placentas; the pregnancy still requires appropriate monitoring.

### **Dizziness**

A sensation of spinning, lightheadedness, imbalance, faintness, or being unsteady. Causes include inner-ear, heart, blood-pressure, vision, medicine, blood, metabolic, or neurologic problems.

### **Dizzy**

Alternate name; see *Dizziness*.

### **DKD**

Alternate index label; see *Diabetic Nephropathy*.

### **DLCO**

A pulmonary function test parameter measuring the ease with which inhaled carbon monoxide transfers across the alveolar-capillary barrier into pulmonary erythrocyte hemoglobin, reduced in emphysema, interstitial lung disease, and pulmonary vascular disorders.

#### **Alternate Names:**

DLCO; Transfer Factor

### **DLI**

Donor lymphocyte infusion, in which immune cells from a stem-cell donor are given after transplantation to strengthen an immune attack against return-



---

ing cancer. It can cause graft-versus-host disease and other complications.

## **DM**

An abbreviation most commonly meaning diabetes mellitus, a group of disorders with persistently high blood glucose. Because DM can have other meanings, the surrounding clinical context is important.

## **DM2**

An abbreviation for type 2 diabetes mellitus, in which the body resists insulin and may gradually produce too little of it. It can cause long-term damage to the eyes, kidneys, nerves, heart, and blood vessels.

## **DMD**

Abbreviation for Duchenne muscular dystrophy, an X-linked muscle disease caused by deficient dystrophin. It usually begins in childhood with progressive proximal weakness, difficulty running or climbing, calf enlargement, and later heart and respiratory complications.

## **DMFT Index**

A dental measure counting permanent teeth that are decayed, missing because of decay, or filled. It estimates a person's lifetime experience of tooth decay but does not show whether a current problem is active.

## **DMG**

Alternate name; see *Diffuse Midline Glioma*.

## **DMSA Scan**

A nuclear-imaging test using a small amount of labeled dimercaptosuccinic acid to show kidney tissue. It can identify differences in kidney function, scars, or injury, especially in children with repeated urinary infections.

## **DNA**

Deoxyribonucleic acid, the molecule that stores hereditary information in humans and most organisms. Its sequence helps control cell function and

protein production; changes may be inherited or acquired during life.

## **DNA Adduct**

A chemical group attached to DNA after exposure to a reactive substance. If not repaired, it can alter DNA copying, cause mutations, and potentially contribute to cancer.

## **DNA Alkylation**

Attachment of an alkyl group to DNA. This can change base pairing, block copying or repair, cause mutations, or kill a cell; some anticancer medicines work partly this way.

## **DNA Amplification**

Production of many copies of a selected DNA sequence, naturally in cells or in the laboratory by methods such as polymerase chain reaction. It helps detect, identify, or study genes and microorganisms.

## **DNA Binding**

Attachment of a protein or other molecule to a DNA sequence. This can control gene activity, DNA copying, repair, chromosome organization, or the action of some medicines.

## **DNA Damage**

A chemical or structural change in DNA that can interfere with gene function or copying. Cells usually repair such damage, but persistent or incorrectly repaired damage can cause mutations, cell death, or cancer.

## **DNA Demethylation**

Removal or loss of small chemical groups called methyl groups from DNA. It can change how accessible a gene is and alter gene activity during development, cell reprogramming, or disease.

## **DNA Double Helix**

The twisted ladder-like structure of DNA, formed by two strands running in opposite directions. Sugar and phosphate form the outside rails, while paired bases join the strands inside.

---

## DNA Fragmentation

Breaking DNA strands into smaller pieces. It occurs normally during programmed cell death and can also result from injury, radiation, chemicals, disease, or laboratory processing.

## DNA Gyrase

A bacterial enzyme that twists and untwists DNA so it can be copied and separated. Fluoroquinolone antibiotics block this enzyme, stopping bacterial growth and survival.

## DNA Helicase

An essential adenosine triphosphate-dependent motor enzyme that unwinds and separates double-stranded DNA into single strands ahead of the replication fork, enabling DNA polymerases access for replication, transcription, and recombination.

## DNA Integration

Insertion of foreign or copied DNA into a cell's genome. It occurs in some viral infections, evolution, and gene therapy and can alter nearby gene activity if the insertion affects a control region.

## DNA Ligases

Enzymes that join breaks in DNA strands during copying, repair, and genetic recombination. They help preserve the continuous structure of chromosomes.

## DNA Ligation

Joining DNA fragments by forming a chemical bond between them. Cells use it during DNA copying and repair, and laboratories use it in genetic testing and cloning.

## DNA Methylation

Addition of small chemical groups called methyl groups to DNA, usually at cytosine bases. It can change how strongly a gene is expressed without changing the DNA sequence.

## DNA Packaging

Arrangement of long DNA molecules around his-

tone proteins and into compact structures called chromosomes. Packaging allows DNA to fit inside the nucleus and helps control access to genes.

## DNA Polymerase

A family of cellular enzymes that synthesize new DNA strands by polymerizing deoxynucleotides complementary to a single-stranded template in a 5-prime to 3-prime direction, with many isoforms possessing 3-prime to 5-prime exonucleolytic proof-reading activity.

## DNA Primase

An enzyme that makes short RNA starting pieces so other enzymes can begin copying DNA. It is required during cell division and DNA repair.

## DNA Primers

Short nucleic-acid sequences that provide a starting point for copying DNA. Cells make them during replication, and laboratories use designed primers to amplify selected genetic sequences.

## DNA Probe

A labeled single-stranded DNA sequence designed to bind a matching target. It can help detect a gene, mutation, chromosome region, or microorganism in a laboratory sample.

## DNA Profiling

A laboratory test that compares selected parts of a person's DNA. It can help identify a person, assess biological relationships, identify remains, or check whether cells came from a donor or another person. Results are very informative but must be interpreted with the sample quality, testing method, and legal or medical setting.

### Alternate Names:

DNA Fingerprinting; Genetic Fingerprinting; DNA Typing; STR Analysis

## DNA Repair

Cellular processes that detect and correct damage or copying errors in DNA. Repair pathways address small base changes, bulky chemical damage, mismatched bases, and breaks in one or both DNA

---

strands.

### **DNA Repair Mechanisms**

Cellular systems that correct DNA damage caused by copying errors, chemicals, radiation, or reactive oxygen. Failure of some pathways increases mutation risk and can contribute to inherited cancer syndromes.

### **DNA Replication**

The process by which DNA copies itself before cell division. Each new DNA molecule contains one original strand and one newly made strand, allowing genetic information to pass to daughter cells.

### **DNA Sequencing**

Determination of the order of DNA building blocks in a sample. It can examine a single gene, selected genes, all coding regions, or the whole genome to identify genetic changes.

### **DNA Virus**

A virus whose genetic material is DNA. DNA viruses vary widely in transmission, tissues affected, and disease; some cause persistent infection, while others cause short-term illness.

### **Do Not Resuscitate**

A medical order stating that cardiopulmonary resuscitation should not be attempted if a person's breathing or heartbeat stops, according to the person's informed wishes or legally authorized decision-maker.

### **Do-Not-Resuscitate Order**

A medical order instructing clinicians not to perform cardiopulmonary resuscitation if a person's breathing or heartbeat stops. It does not automatically refuse other treatments or comfort care.

#### **Alternate Names:**

DNR Order; DNAR Order; Allow Natural Death (AND)

### **DOACs**

Direct oral anticoagulants are oral blood thinners

that directly block thrombin or factor Xa. They prevent and treat blood clots but can cause bleeding and require attention to kidney function, interactions, and planned procedures.

### **DOB**

An abbreviation for date of birth, the day, month, and year a person was born. It helps confirm identity and calculate age for records, screening, medicine doses, and treatment decisions.

### **Dobutamine**

An intravenous medicine that increases heart contraction and, sometimes, heart rate. It is used short term for selected cases of severe low-output heart failure or shock and requires close monitoring.

### **Dobutamine Stress Echocardiography**

An ultrasound heart test using dobutamine to increase heart workload when a person cannot exercise adequately. New areas of poor wall movement may suggest reduced blood flow, but results require clinical interpretation.

### **Doc**

Informal shorthand for doctor, a person qualified to practice medicine or another health profession. The exact role depends on the person's training and professional license.

### **Docetaxel**

An anticancer medicine that interferes with microtubules, structures needed for cell division. It is used for several cancers and can cause low blood counts, infection, hair loss, nerve problems, or allergic reactions.

### **Doconexent**

A pharmaceutical name for docosahexaenoic acid, an omega-3 fatty acid found in oily fish, algae, and cell membranes. Its effects depend on the product, dose, and reason for use.

### **Docosadienoic Acid**

A 22-carbon fatty acid with two double bonds,

---

present in small amounts in body and dietary lipids. It is studied mainly for possible effects on cell membranes and metabolism.

### **Docosahexaenoic Acid**

An omega-3 fatty acid that is an important structural part of brain and retinal cell membranes. It comes mainly from oily fish, algae, or supplements and is especially important during brain development.

### **Docosapentaenoic Acid**

A 22-carbon polyunsaturated fatty acid with five double bonds. It occurs as omega-3 and omega-6 forms in foods and tissues; its health effects depend on the form and amount.

### **Docosatrienoic Acid**

A 22-carbon fatty acid with three double bonds, found in small amounts in body and dietary lipids. It is studied mainly for its possible roles in cell membranes and metabolism.

### **Docusate**

A stool-softening medicine that helps water mix with stool. It may be used when avoiding straining is important, but evidence for treating ordinary constipation is limited; diarrhea and abdominal cramps can occur.

### **Dog Allergy**

An immune reaction to proteins from a dog's skin flakes, saliva, urine, or other secretions. It can cause sneezing, blocked or itchy nose, watery eyes, hives, cough, or asthma symptoms.

### **Dog Bite**

An injury caused by a dog's teeth. Wash the wound promptly and seek assessment for tissue damage, infection, tetanus protection, and rabies risk, especially for deep bites or bites to the face or hand.

#### **Alternate Names:**

Canine Bite Trauma; Dog Bite Wound

### **Dolasetron**

A medicine that blocks serotonin 5-HT<sub>3</sub> receptors and helps prevent nausea and vomiting in selected settings. It can cause headache, constipation, or dangerous heart-rhythm changes.

### **Dolasetron Mesylate**

A salt form of dolasetron, a medicine that blocks serotonin 5-HT<sub>3</sub> receptors to prevent nausea and vomiting in selected settings. It can cause constipation, headache, or dangerous heart-rhythm changes.

### **Dolichocephalic**

Having a relatively long, narrow skull. It may be a normal family or positional variation, but marked or worsening skull narrowing in an infant may indicate early fusion of a skull suture and needs assessment.

### **Dolichol**

A long-chain fatty alcohol used inside cells to assemble sugar chains attached to proteins. This process is important for protein folding, transport, and function.

### **Dolichostenomelia**

Abnormally long, slender limbs, often used to describe a marfanoid body habitus. It may occur in connective-tissue disorders such as Marfan syndrome and should be interpreted with the person's other skeletal and cardiovascular findings.

### **Doll Eye Reflex**

The oculocephalic reflex in which rapid passive rotation or flexion and extension of the head in an unconscious or comatose patient produces conjugately opposed compensatory eye movements when the brainstem pathways linking cranial nerves III, VI, and VIII through the medial longitudinal fasciculus are functionally intact.

#### **Alternate Names:**

Oculocephalic Reflex; Doll's Eyes Phenomenon

### **Doll's Eye Sign**

A physical examination finding of the oculocephalic reflex in comatose patients, in which rapid hori-

---

zontal or vertical rotation of the head produces conjugate eye deviation in the direction opposite to head movement, demonstrating intact midbrain and pontine brainstem pathways.

**Alternate Names:**

Oculocephalic Reflex

**Dolor**

The Latin word for pain, one of the classic signs of inflammation. Pain may result from irritated nerves, swelling, injury, infection, or another cause and is interpreted with other findings.

**Domain**

A defined region or area. In biology, a protein domain is a part with a particular structure or function; in genetics, a domain may be a genomic or regulatory region.

**Domestic Violence**

A pattern of physical, sexual, emotional, psychological, or financial abuse or coercive control by an intimate partner or household member. It can cause injury, fear, isolation, and serious health effects; confidential support is available.

**Dominant**

Describing a genetic variant whose effect can appear when only one copy is present. An affected parent with an autosomal dominant condition may pass the variant to each child with a one-in-two chance.

**Domperidone**

A medicine that blocks dopamine receptors outside the brain and is used in some countries for nausea or slow stomach movement. It can prolong the heart rhythm and cause dangerous arrhythmias.

**Donath-Landsteiner Antibody**

An antibody that can attach to red blood cells in cold conditions and destroy them when the blood warms. It is associated with paroxysmal cold hemoglobinuria, a rare cause of red-cell breakdown.

**Alternate Names:**

Donath-Landsteiner      Hemolysin;      Biphasic

Hemolysin

**Donath-Landsteiner Test**

A blood test that detects an antibody associated with paroxysmal cold hemoglobinuria. The antibody binds red cells in the cold and destroys them when the blood warms.

**Donepezil**

A medicine that increases acetylcholine activity in the brain and may temporarily improve symptoms of Alzheimer disease. It does not cure the disease and can cause nausea, diarrhea, slow heartbeat, fainting, or sleep problems.

**Donnan Equilibrium**

The distribution of charged particles across a partly permeable membrane when large charged particles cannot cross. It helps explain cell volume, membrane electrical charge, and movement of salts and water.

**Donohue Syndrome**

A rare inherited disorder in which the body cannot respond normally to insulin. It causes severe growth and feeding problems, very little body fat, unusual facial features, and serious metabolic complications beginning in infancy.

**Donor**

A person who provides blood, cells, tissue, an organ, sperm, eggs, or another biological material for another person or for research. Screening, informed consent, compatibility, and traceability support safety.

**Donor Insemination**

A fertility treatment in which sperm from a screened donor is placed in the reproductive tract to attempt pregnancy. The donor may be known or anonymous, and counseling and legal requirements vary by location.

**Donor Suitability Assessment**

Evaluation of a potential donor's medical history,

---

examination, laboratory tests, infection risks, organ function, and compatibility to determine whether donation is safe and suitable for transplantation.

### **Donor Testing**

Laboratory and clinical evaluation of a potential donor for infections, cancer, organ function, blood type, and other factors that affect donation safety and recipient compatibility.

### **Donovan Bodies**

Intracellular clusters of deeply staining, encapsulated, pleomorphic gram-negative rod bacteria (*Klebsiella granulomatis*) located within the cytoplasm of large mononuclear histiocytes, identified on Giemsa or Wright-stained tissue smears from lesions of granuloma inguinale (donovanosis).

#### **Alternate Names:**

Donovan Body; Leishmanoid Granules

### **Donovanosis**

A sexually transmitted infection caused by *Klebsiella granulomatis*. It produces slowly enlarging, usually painless genital ulcers that may bleed easily and requires medical diagnosis and antibiotic treatment.

#### **Alternate Names:**

Granuloma Inguinale; Calymmatobacterium Infection

### **Dopa**

A chemical precursor that the body can convert into dopamine. Levodopa, a form that reaches the brain, is used with another medicine to improve movement problems in Parkinson disease.

### **Dopa-Responsive Dystonia**

An inherited movement disorder causing sustained muscle contractions and abnormal postures, sometimes with Parkinson-like symptoms. It often improves markedly with a small dose of levodopa.

### **Dopamine**

A chemical messenger used by nerve cells and some hormone-producing cells. It helps control movement,

motivation, reward, attention, learning, blood-vessel tone, and release of certain hormones.

### **Dopamine Agonist**

A medicine that stimulates dopamine receptors and mimics some effects of dopamine. It is used for selected movement, hormone, or other disorders and may cause nausea, sleepiness, low blood pressure, hallucinations, or impulse-control problems.

### **Dopamine Agonists**

Medicines that stimulate dopamine receptors and are used for selected Parkinson, restless-legs, or hormone disorders. Possible effects include nausea, sleepiness, low blood pressure, hallucinations, and impulse-control problems.

### **Dopamine Antagonist**

A medicine or compound that blocks dopamine receptors. It is used in selected psychiatric, nausea, or digestive treatments but can cause movement problems, drowsiness, hormone changes, or other adverse effects.

### **Dopamine Receptor**

A protein on or inside a cell that responds to dopamine. Different receptor types affect movement, mood, hormone release, blood vessels, and other functions and are targets of some medicines.

### **Dopamine Transporter**

A protein that clears dopamine from the space between nerve cells by moving it back into the releasing nerve cell. Stimulants and some imaging tracers affect or measure its activity.

### **Dopaminergic Neuron**

A nerve cell that makes and releases dopamine. These cells help control movement, motivation, reward, attention, and hormone regulation; loss of some contributes to Parkinson disease.

### **Dopaminergic Pathways**

Four main nerve pathways that use dopamine to send signals. They help control movement, reward,



---

thinking, emotions, and the release of some hormones.

### **Doppler Echocardiography**

A heart ultrasound technique that measures the direction and speed of blood flow through the heart and major vessels. It helps assess valves, chambers, pressure, and abnormal flow.

### **Doppler Ultrasound**

An ultrasound technique that measures moving blood to assess its direction and speed. It can identify narrowed or blocked vessels, abnormal flow, blood clots, and circulation problems without ionizing radiation.

### **Doppler Ultrasound of a Limb**

An ultrasound examination measuring blood flow through the arteries and veins of an arm or leg. It can help identify narrowing, blockage, a clot, aneurysm, or other circulation problem without ionizing radiation.

### **Doppler Velocimetry**

Ultrasound measurement of blood-flow speed and resistance in fetal or placental vessels. It helps assess circulation and fetal well-being when poor growth, placental problems, or other pregnancy risks are suspected.

### **Dor Procedure**

A heart operation that removes or excludes a scarred, bulging part of the left ventricle after a large heart attack. It reshapes the chamber in selected people with severe ischemic heart-muscle disease.

#### **Alternate Names:**

Endoventricular Circular Patch Plasty (EVCPP);  
Surgical Ventricular Restoration

### **Doravirine**

An HIV medicine that blocks reverse transcriptase, an enzyme HIV needs to copy its genetic material. It is used with other medicines for HIV-1 infection; interactions and resistance affect its choice.

### **Dorea**

A group of bacteria commonly found in the human intestine and able to grow without oxygen. Finding it usually reflects normal gut organisms unless it comes from a normally sterile site with compatible illness.

### **Doripenem**

An intravenous antibiotic that damages bacterial cell walls and is used for selected serious infections. It can cause allergy, diarrhea, seizures, or resistance and should be chosen according to the organism and patient.

### **Dormant**

Inactive or not multiplying. The term may describe a latent infection, quiet tumor cell, inactive disease process, or other biological state that can become active under certain conditions.

### **Dorsal**

Relating to the back or upper surface of a body part. In human anatomy it usually means toward the back, although in the hand or foot it refers to the opposite surface from the palm or sole.

### **Dorsal Blocking Splint**

A hand splint that limits wrist and finger extension to protect healing tendons, commonly after flexor-tendon repair. Position and duration are set by the treating surgical and therapy team.

### **Dorsal Body Cavity**

The body cavity toward the back. It includes the cranial cavity, which contains the brain, and the spinal cavity, which contains the spinal cord.

### **Dorsal Column-Medial Lemniscal Pathway**

The major ascending sensory pathway conveying conscious proprioception, discriminative fine touch, and vibratory sensation from peripheral mechanoreceptors through the spinal cord fasciculus gracilis and cuneatus to the medulla, where fibers decussate in the medial lemniscus to ascend to the thalamus.

---

**Alternate Names:**

DCML Pathway; Dorsal Column System

**Dorsal Funiculus**

A back column of the spinal cord containing nerve fibers that carry vibration, fine touch, and awareness of body position to the brain.

**Dorsal Kyphosis**

Excessive forward curvature of the upper spine, causing a rounded upper back. It may result from posture, spinal wear, osteoporosis, or vertebral fractures and can cause pain, height loss, or breathing difficulty.

**Dorsal Root**

The back root of a spinal nerve. It carries sensory signals from the body into the spinal cord.

**Dorsal Root Ganglion**

A swelling on the back root of a spinal nerve containing the cell bodies of sensory nerve cells. It relays information about touch, pain, temperature, and body position to the spinal cord.

**Dorsalis Pedis Artery**

An artery running across the top of the foot as a continuation of the anterior tibial artery. Its pulse helps assess blood flow to the foot.

**Dorsiflexion**

Upward movement of the foot at the ankle, bringing the toes toward the shin. It helps clear the foot during walking; weakness can cause foot drop and tripping.

**Dorsocervical Fat Pad**

A visible accumulation of fat at the back of the neck and upper shoulders, associated with obesity, corticosteroid excess, some medicines, or lipodystrophy; it differs from spinal kyphosis.

**Dorsolateral**

Describing a position toward the back and side of a body structure. For example, a dorsolateral area

is behind and to one side rather than toward the front or middle.

**Dorsum**

The back or upper surface of a body part, such as the back of the hand, top of the foot, or upper surface of the tongue.

**Dorzolamide**

An eye medicine that lowers pressure inside the eye by reducing fluid production. It is used for glaucoma or raised eye pressure and may cause stinging, blurred vision, or a bitter taste.

**Dosage**

The amount, frequency, route, and duration of a medicine given to achieve benefit while limiting harm. It may need adjustment for age, weight, kidney or liver function, interactions, and the condition treated.

**Dosage Forms**

The physical form in which a medicine is supplied, such as a tablet, capsule, liquid, cream, inhaler, injection, patch, suppository, or implant. The form affects how and how quickly it is given.

**Dose**

The amount of a medicine, radiation, or other substance given or received at one time. The appropriate dose depends on the substance, purpose, age, weight, organ function, interactions, and individual risk.

**Dose Rate**

The amount of a medicine, radiation, or other exposure delivered during a specified time, such as milligrams per hour or grays per minute. It affects both benefit and toxicity.

**Dose Titration**

Gradual adjustment of a medicine dose according to its effects, treatment goals, blood tests, or side effects. The dose may be increased, decreased, or kept the same as the person's response is assessed.

---

### **Dose-Limiting Toxicity**

A treatment-related adverse effect severe enough to prevent increasing the dose or require reducing it. The term is used especially in early cancer trials to estimate a tolerable dose.

### **Dose-Response Curve**

A graph showing how the size of a biological effect changes as the dose of a medicine or substance increases. It helps describe potency, maximum effect, and the dose range that produces benefit or harm.

### **Dose-Response Relationship**

The relationship between the amount of a medicine or substance and the size of its effect. Increasing the dose may increase benefit, toxicity, or both, but the pattern depends on the substance and person.

### **Dosimeter**

An instrument worn or placed near a person or object to measure or record exposure to ionizing radiation. It helps monitor occupational or treatment-related radiation.

### **Dosimetry**

Measurement and calculation of radiation dose received by a person, tissue, or object. In treatment, it helps plan and verify that the intended dose reaches the target while limiting exposure elsewhere.

### **DOTATATE**

A substance that binds somatostatin receptors found on many neuroendocrine tumors. When linked to a radioactive tracer, it is used in PET imaging to locate and stage tumors and assess possible targeted treatment.

### **DOTS**

A tuberculosis-control strategy involving reliable diagnosis, standardized combination treatment, medicine-supply support, monitoring, and sometimes directly observed doses. It aims to cure infection and limit drug resistance and transmission.

### **Double Aortic Arch**

A birth defect in which two aortic arches form a ring around the windpipe and esophagus. It may cause noisy breathing, cough, feeding difficulty, vomiting, or trouble swallowing.

### **Double Blind**

A study design in which participants and the researchers assessing outcomes do not know which treatment each participant receives during the blinded phase. It reduces expectation and assessment bias.

#### **Alternate Names:**

Double-Masked Study; Double-Blind Protocol

### **Double Blind Trial**

A clinical trial in which participants and the investigators measuring outcomes do not know the assigned treatment during the blinded phase. This helps reduce expectations from influencing results.

### **Double Bubble Sign**

An imaging finding in a newborn showing two enlarged, air- or fluid-filled areas representing the stomach and first part of the small intestine. It strongly suggests a blockage near the start of the small intestine, such as duodenal atresia, but other causes are possible.

#### **Alternate Names:**

Duodenal Atresia Sign; Double Bubble Radiographic Appearance

### **Double Inlet Left Ventricle**

A rare birth defect in which both upper heart chambers connect mainly to one lower chamber. It produces single-ventricle circulation and mixing of oxygen-rich and oxygen-poor blood and usually needs staged specialist treatment.

### **Double Outlet Right Ventricle**

A birth defect in which both major arteries arise mainly from the right lower heart chamber. An opening between the lower chambers allows blood from the left chamber to reach the arteries; anatomy and symptoms vary.

---

## Double Uterus

A birth difference in which the uterus develops as two separate cavities or two uteruses. Some people have no symptoms; others may have menstrual problems, infertility, miscarriage, or pregnancy complications.

## Double Vision

Seeing two images of one object. Causes include eye-surface or lens disease, eye misalignment, muscle or nerve problems, medicines, migraine, or brain disease; sudden onset with neurologic symptoms or eye pain is an emergency.

## Double-Balloon Enteroscopy

An endoscopic procedure using two balloons to move a flexible camera through much of the small intestine. It can locate bleeding or lesions, take biopsies, and perform selected treatments.

## Double-Blind Method

A research method in which treatment assignments are hidden from participants and relevant study personnel during the blinded phase to reduce expectation and assessment bias.

## Double-Blind Study

A clinical study in which participants and the investigators assessing outcomes do not know the assigned treatment group during the blinded phase.

## Double-Contrast Barium Enema

An X-ray examination in which barium coats the inside of the colon and air expands it. It can show polyps, masses, narrowing, or changes in the bowel lining.

### Alternate Names:

Air-Contrast Barium Enema

## Double-Outlet Right Ventricle

A birth defect in which both major arteries arise mainly from the right lower heart chamber. An opening between the lower chambers allows blood to reach the arteries; symptoms and treatment depend on the exact anatomy.

## Douche

A stream of water or solution directed into a body cavity, most often the vagina. Vaginal douching is not needed for hygiene and can irritate tissue or disturb normal bacteria and increase infection risk.

## Dowager Hump

An exaggerated posterior thoracic hyperkyphosis causing an abnormal dorsal hump at the base of the neck, caused by multiple anterior wedge compression fractures of osteoporotic thoracic vertebral bodies in elderly patients.

### Alternate Names:

Senile Kyphosis; Postmenopausal Thoracic Hyperkyphosis

## Dowager's Hump

An exaggerated forward curve of the upper spine that creates a rounded back. It may result from osteoporosis and collapsed vertebrae; new pain, height loss, breathing difficulty, or worsening curvature needs assessment.

### Alternate Names:

Kyphosis

## Dowling-Meara Syndrome

A severe inherited form of epidermolysis bullosa simplex, usually caused by variants in the *KRT5* or *KRT14* gene. Widespread grouped blisters may begin at birth or in infancy and can affect the skin, mouth, nails, and airway.

### Alternate Names:

Epidermolysis Bullosa Simplex, Dowling-Meara Type; EBS-DM

## Down Syndrome

A genetic condition caused by an extra copy of chromosome 21. It may affect learning, muscle tone, growth, and appearance and is associated with heart, hearing, vision, thyroid, sleep, and digestive problems.

### Alternate Names:

Trisomy 21; Down's Syndrome

---

## Downregulation

A decrease in the number, sensitivity, or activity of receptors, genes, or signaling pathways. It may occur after prolonged stimulation, medicine exposure, inflammation, or another regulatory change.

## Doxapram

A medicine that stimulates breathing and is used rarely in selected situations. It can cause agitation, high blood pressure, fast or abnormal heart rhythms, seizures, or other serious effects.

## Doxazosin

An alpha-1 blocker that relaxes blood vessels and prostate or bladder-neck muscle. It is used for high blood pressure or urinary symptoms from an enlarged prostate and can cause dizziness or fainting.

## Doxazosin Mesylate

A salt form of doxazosin, an alpha-1 blocker used for high blood pressure and urinary symptoms from an enlarged prostate. It can cause dizziness, low blood pressure, fainting, or falls, especially after the first dose.

## Doxepin

A tricyclic antidepressant also used in low doses for insomnia or skin itching. It can cause drowsiness, dry mouth, constipation, blurred vision, abnormal heart rhythms, and dangerous toxicity in overdose.

## Doxepin Overdose

Taking too much doxepin can cause severe drowsiness, confusion, seizures, low blood pressure, coma, and life-threatening heart-rhythm or conduction problems, especially with alcohol or other medicines.

## Doxercalciferol

A vitamin D-like medicine used in selected people with chronic kidney disease to reduce excess parathyroid hormone. Calcium and phosphate must be monitored because levels can become too high.

## Doxorubicin

An anthracycline cancer medicine that damages

DNA and blocks cell division. It treats many cancers but can cause low blood counts, infection, nausea, hair loss, tissue injury after leakage, and dose-related heart damage.

## Doxorubicin Hydrochloride

A salt form of doxorubicin, an anthracycline cancer medicine used for many cancers. Important risks include low blood counts, infection, tissue injury after leakage, and dose-related heart damage.

## Doxycycline

A tetracycline antibiotic that blocks bacterial protein production. It treats many bacterial and tick-borne infections and can cause nausea, sun sensitivity, esophageal irritation, diarrhea, or tooth and bone effects in developing children.

## Doxycycline Hyclate

A salt form of doxycycline, a tetracycline antibiotic that blocks bacterial protein production. It is used for susceptible infections; nausea, sun sensitivity, esophageal irritation, diarrhea, and allergic reactions may occur.

## Doxylamine

A sedating antihistamine used in some sleep aids and pregnancy-related nausea treatments. It can cause drowsiness, dry mouth, blurred vision, constipation, urinary retention, confusion, or impaired driving.

## DPP-4 Inhibitor

An oral diabetes medicine that prolongs the action of hormones released after meals, increasing glucose-dependent insulin release and reducing glucagon. It is used mainly in type 2 diabetes and usually has a low risk of low glucose.

## DPP-4 Inhibitors

Oral medicines for type 2 diabetes that prolong hormones released after meals, helping the pancreas release insulin and reducing glucagon. They usually do not cause weight gain or low glucose when used alone but have modest glucose-lowering effects.

---

## DPT Immunization

Vaccination that protects against diphtheria, pertussis (whooping cough), and tetanus. Current childhood and booster vaccines use age-appropriate formulations and are given as a series according to the recommended schedule.

## DR

Alternate name; see *Diabetic Retinopathy*.

## Dracunculiasis

Guinea-worm disease caused by a parasite acquired from unsafe drinking water containing infected tiny water fleas. About a year later, a painful blister usually forms on a leg as the adult worm emerges through the skin.

## Dracunculosis

Infection with the Guinea-worm parasite after drinking unsafe water containing infected water fleas. It causes a painful blister, usually on a leg, followed by emergence of a long adult worm through the skin.

## Dracunculus Medinensis

The parasitic worm that causes Guinea-worm disease. People become infected by drinking water containing tiny infected water fleas; roughly a year later, an adult female worm emerges through the skin.

## Draf Procedure

A standardized series of endoscopic endonasal frontal sinus drainage techniques (Draf type I, IIa, IIb, and III) utilized in rhinology. Draf type III (modified endoscopic Lothrop procedure) involves complete resection of the frontal sinus floor, superior nasal septum, and interfrontal sinus septum to create a single large drainage neo-ostium into both frontal sinuses.

### Alternate Names:

Endoscopic Frontal Sinusotomy; Modified Lothrop Procedure

## Drain

A tube or other device used to remove blood, pus,

fluid, or air from a wound, body cavity, or organ. Drainage may prevent pressure, infection, or impaired healing after injury or surgery.

## Drain Opener Poisoning

Exposure to a caustic drain-cleaning product that can cause severe burns of the skin, eyes, airway, or gastrointestinal tract.

## Drainage

Removal of blood, pus, fluid, or air from a body site, wound, or abnormal collection. It may occur naturally or use a needle, catheter, tube, or surgical opening to relieve pressure and aid healing.

## Drainage Procedures

Treatments that remove pus, blood, or other collected fluid, sometimes with ultrasound or CT guidance. They may use a needle, catheter, tube, or surgery to treat abscesses and other fluid collections.

## Drainpipe Cleaners

Strong chemical products used to clear blocked pipes, commonly containing caustic alkalis or acids. Swallowing, inhaling, or splashing these products can cause severe burns to the mouth, throat, eyes, skin, or lungs and requires urgent poison-control or medical advice.

## Drawer Test

An orthopedic manual examination of the flexed knee joint assessing the integrity of the cruciate ligaments; forward tibial translation tests the anterior cruciate ligament (anterior drawer test) and backward tibial translation tests the posterior cruciate ligament (posterior drawer test).

### Alternate Names:

Knee Drawer Sign; Anterior and Posterior Drawer Tests

## Dream

A sequence of images, thoughts, sensations, or emotions experienced during sleep. Dreams are often most vivid during rapid-eye-movement sleep and may be influenced by stress, medicines, illness, or sleep disorders.



---

## Dreams

Experiences involving images, thoughts, sensations, or emotions during sleep, often vivid during rapid-eye-movement sleep. Dreams are normal, but distressing, violent, or disruptive dreams may occur with stress, medicines, or sleep disorders.

## Drepanocytosis

Sickle-cell disease or a related condition in which an abnormal form of hemoglobin makes red blood cells become rigid and sickle-shaped. This can cause anemia, painful blockages of blood flow, and damage to organs.

## DRESS Syndrome

A serious delayed allergic reaction to a medicine, usually beginning two to eight weeks after exposure. It may cause fever, widespread rash, facial swelling, abnormal blood counts, hepatitis, kidney injury, or other organ inflammation.

## Dressing

Material placed over a wound to protect it from contamination, absorb drainage, control minor bleeding, maintain moisture, or deliver local treatment. It should be changed as directed and monitored for infection.

## Dressler Syndrome

An immune-related inflammation of the sac around the heart that occurs weeks after a heart attack, heart surgery, or other heart injury. It can cause fever, sharp chest pain, and fluid around the heart or lungs.

### Alternate Names:

Post-Cardiac Injury Syndrome Postmyocardial Infarction Syndrome

## Dressler's Syndrome

Alternate spelling; see *Dressler Syndrome*.

## DRI

Dietary reference values used to estimate the amounts of vitamins, minerals, protein, and other nutrients needed by healthy people. They help

assess diets but are not individualized treatment targets for every illness.

### Alternate Names:

Dietary Reference Intakes

## Dried Specimen

A biological sample collected and dried on a prepared card or other surface for transport and laboratory testing. Dried blood spots are commonly used for screening, but correct collection and handling are essential.

## Drinking Water

Water safe for human consumption. It should be free of disease-causing organisms and harmful chemicals; boiling, filtration, chlorination, or bottled water may be needed when the supply is contaminated or uncertain.

## Drip

A drop-by-drop flow of liquid, or fluid given slowly into a vein through tubing and a catheter. The infusion rate is controlled to deliver medicines, fluids, blood products, or nutrients safely.

## Driving And Older Adults

Evaluation of whether an older person can drive safely, considering vision, attention, memory, hearing, movement, health conditions, and medicines. Age alone does not determine driving ability.

## Driving Evaluation

An assessment of driving ability after illness, injury, or a health change. It may examine vision, attention, memory, reaction time, strength, and vehicle control in a clinic and on the road; adaptive equipment may help.

## Driving Rehabilitation

Assessment and training to help a person resume driving safely or use adaptive vehicle controls after illness or injury. It may address vision, attention, reaction time, strength, vehicle operation, and road safety.

---

## Dronabinol

A medicine related to the active substance in cannabis, used for selected cases of chemotherapy-related nausea or severe appetite loss. It may cause dizziness, sleepiness, confusion, mood changes, or impaired coordination.

## Drool

Saliva that unintentionally escapes from the mouth. It may result from excess saliva, difficulty swallowing, poor lip closure, dental problems, or a neurologic condition affecting mouth control.

## Drooling

Unintentional loss of saliva from the mouth caused by excess saliva, difficulty swallowing, poor lip closure, or reduced control of the mouth. Persistent drooling may need assessment for dental, medication, or neurologic causes.

## Drooping Eyelids

Ptosis, or abnormal lowering of one or both upper eyelids. Causes include age-related tendon stretching, nerve or muscle disease, congenital weakness, trauma, or a mass; sudden ptosis with double vision, headache, or unequal pupils is urgent.

## Drop Attack

A sudden, unprovoked collapse to the ground resulting from abrupt loss of postural tone in the lower extremities without loss of consciousness, occurring in vertebrobasilar insufficiency, cryptogenic falls of older adults, or inner ear otolithic hydrops (Tumarkin otolithic crisis).

## Drop Foot

Alternate name; see *Foot Drop*.

## Drop Hand Syndrome

Weakness or paralysis of the muscles that straighten the wrist and fingers, usually from injury or pressure on the radial nerve. The wrist hangs downward and gripping or releasing objects becomes difficult.

## Droperidol

A medicine used in selected clinical settings to treat severe nausea or agitation. It can prolong the heart's electrical recovery time and rarely cause dangerous arrhythmias, so heart-risk factors and monitoring may be important.

## Droplet Precautions

Infection-control measures for infections spread by respiratory droplets released during coughing, sneezing, or talking. They may include a medical mask, hand hygiene, patient separation, and limiting transport according to the infection.

## Dropper Bottle

A bottle with a device that releases liquid one drop at a time. The medicine's concentration, prescribed number of drops, route, and measuring device must be followed because drop size can vary.

## Drowning

Breathing impairment caused by submersion or immersion in liquid. It may be fatal or nonfatal. Lack of oxygen can injure the brain and other organs; complications include aspiration, lung injury, and respiratory failure.

## Drowning Signs

Alternate name; see *Drowning*. (The terms 'dry drowning' and 'wet drowning' are non-standard clinical descriptions).

## Drowsiness

An increased tendency to fall asleep or reduced alertness. It may result from too little sleep, medicines, alcohol or other substances, metabolic illness, sleep disorders, infection, or neurologic disease; sudden severe drowsiness needs urgent assessment.

## Drug

A substance that changes body function or is used to prevent, diagnose, or treat disease. Drugs include prescription medicines, nonprescription medicines, biological products, and substances used recreationally or illegally.

---

## Drug Caution Code

A label or code that warns about an important medicine risk, such as allergy, bleeding, interaction, overdose, or a need for special monitoring. Its meaning depends on the coding system and product information.

## Drug Delivery

The way a medicine is prepared and given so its active ingredient reaches the intended body site at an effective dose. Routes include swallowing, inhalation, skin application, injection, implantation, and targeted systems.

## Drug Desensitization

A supervised process that gives gradually increasing doses of a medicine to temporarily reduce an allergic reaction when that medicine is essential and no suitable alternative exists. The tolerance usually ends when doses are interrupted.

## Drug Discovery

The process of finding and developing substances that may prevent, diagnose, or treat disease. It includes laboratory research, safety testing, dose studies, clinical trials, manufacturing, and review by a regulatory authority.

## Drug Eruptions

Adverse skin or mucosal reactions caused by medicines, ranging from mild morbilliform rashes or hives to fixed drug eruptions and severe cutaneous adverse reactions. Blistering, skin pain or peeling, mouth or eye involvement, fever, facial swelling, or organ symptoms require urgent assessment.

### Alternate Names:

Fixed Drug Eruption; FDE

## Drug Hypersensitivity

An abnormal reaction in which the body's immune system mistakenly treats a medicine as a harmful invader. It is different from common side effects, such as mild stomach upset or drowsiness. Reactions can happen quickly (within minutes to an hour of taking the medicine), causing hives, itchy skin, facial or throat swelling, trouble breathing, or a se-

vere whole-body allergic reaction called anaphylaxis. Other reactions develop slowly over several days or weeks, causing widespread red skin rashes, fever, or painful blistering. Common medicines that can trigger these reactions include antibiotics (such as penicillin and sulfa drugs), pain relievers (such as aspirin and ibuprofen), and seizure medications.

### Alternate Names:

Drug Allergy; Medication Allergy; Drug Hypersensitivity Reaction; Allergic Drug Reaction

## Drug Implant

A solid drug-containing pellet, container, or delivery device placed under the skin or in another tissue to release medicine continuously or slowly over an extended period. It may provide local or whole-body treatment; the product determines whether it biodegrades or requires removal and how it is monitored.

## Drug Implants

Plural of drug implant; see *Drug Implant*.

## Drug Repurposing

Studying or using a medicine developed for one condition to treat a different condition. Earlier safety information may help, but the new use still needs evidence about effectiveness, dose, interactions, and risks.

## Drug Safety

The prevention, detection, and management of harmful effects, interactions, errors, and misuse involving medicines. It includes choosing the right medicine and dose, checking allergies and interactions, following instructions, monitoring effects, and reporting suspected problems.

## Drug Screening

Testing a sample such as urine, blood, saliva, or hair for selected drugs or their breakdown products. Initial results can be falsely positive or negative and may require confirmation with a more specific test.

## Drug Testing

Laboratory analysis of blood, urine, saliva, hair,

---

or another sample to detect medicines, recreational drugs, their breakdown products, or poisons. The test's purpose, detection limits, timing, and medicines taken affect interpretation.

### **Drug- and Nephrotoxin-Associated Kidney Disorders**

Kidney injury caused by medicines, contrast agents, industrial chemicals, supplements, or other toxins. Patterns include acute tubular injury, interstitial nephritis, glomerular disease, crystals, obstruction, and chronic fibrosis; medication review and prompt withdrawal of a culprit may limit damage.

### **Drug-Eluting Stent**

A vascular stent coated with a medicine that reduces tissue growth and helps keep the treated vessel open. It is used in selected narrowed or blocked arteries and requires appropriate antiplatelet management.

### **Drug-Food Interaction**

A change in a medicine's effect, absorption, or safety caused by food or drink. Food may increase or decrease absorption, while some foods can alter medicine metabolism or raise the risk of adverse effects.

### **Drug-Induced Bullous Disorders**

Medication-triggered blistering diseases, including bullous pemphigoid, pemphigus, linear IgA disease, bullous fixed drug eruption, and severe reactions such as Stevens–Johnson syndrome or toxic epidermal necrolysis. New widespread blisters, skin tenderness, peeling, fever, or mouth, eye, or genital involvement is an emergency.

### **Drug-Induced Gingival Hyperplasia**

Enlargement or overgrowth of gum tissue associated especially with phenytoin, cyclosporine or tacrolimus, and calcium-channel blockers such as nifedipine or amlodipine. It can impair brushing, trap plaque, and promote bleeding or periodontal disease; dental care and review of the medicine with the prescriber are needed, and treatment should not be stopped abruptly.

### **Drug-Induced Glaucoma**

Glaucoma or raised intraocular pressure caused or worsened by a medicine, commonly corticosteroids and less often agents that produce pupillary block or secondary inflammation. Treatment involves removing the trigger when possible and controlling pressure.

### **Drug-Induced Hepatotoxicity**

Liver injury caused by a medicine, supplement, or other therapeutic exposure. Patterns include hepatocellular injury, cholestasis, or a mixed picture; diagnosis requires exposure review and exclusion of competing causes.

### **Drug-Induced Liver Injury**

Acute or chronic hepatic damage provoked by exposure to prescribed medications, herbal products, or dietary supplements, manifesting as hepatocellular, cholestatic, or mixed injury patterns via intrinsic direct hepatotoxicity or idiosyncratic immune mechanisms.

#### **Alternate Names:**

DILI; Toxic Hepatitis

### **Drug-Induced Lupus**

A lupus-like autoimmune reaction caused by certain medicines, often producing joint pain, muscle pain, fever, and inflammation around the lungs or heart. Symptoms and abnormal antibodies usually improve after the causative medicine is stopped under medical supervision.

#### **Alternate Names:**

Drug-Induced Lupus Erythematosus; DILE

### **Drug-Induced Lupus Erythematosus**

An autoimmune lupus-like syndrome provoked by prolonged exposure to certain pharmacological agents (most classically procainamide, hydralazine, and isoniazid), characterized by arthralgias, serositis, anti-histone antibodies, and resolution following drug discontinuation.

#### **Alternate Names:**

DILE; Drug-Induced Lupus

---

## Drug-Induced Pemphigus

An autoimmune blistering disorder triggered by a medicine, with antibodies disrupting adhesion between epidermal cells and causing fragile blisters or erosions on skin or mucous membranes. Diagnosis uses biopsy and immunofluorescence; the suspected medicine is stopped only with medical guidance, and the eruption may improve after withdrawal.

## Drug-Induced Photosensitivity

An exaggerated skin reaction after exposure to sunlight or other ultraviolet light while taking or applying a photosensitizing medicine. Phototoxic reactions resemble an intense sunburn and may occur quickly, whereas photoallergic reactions are delayed and eczematous; sun protection and clinician-guided review of the medicine are important.

## Drug-Induced Pigmentation

Darkening or altered colour of the skin, nails, teeth, or mucous membranes caused by a medicine or its metabolites. Causes include increased melanin, drug or metabolite deposition, and iron deposition; examples include minocycline, amiodarone, clofazimine, antimalarials, and some chemotherapy medicines.

## Drug-Induced Pulmonary Toxicity

Respiratory injury caused by a medicine, radiation, or treatment-related immune reaction. It may appear as pneumonitis, organizing pneumonia, fibrosis, edema, hemorrhage, bronchospasm, or pulmonary vascular disease; diagnosis requires temporal review and exclusion of infection and other causes.

## Drugs That May Cause Erection Problems

Some medicines can contribute to difficulty getting or keeping an erection by affecting blood flow, hormones, nerves, or mood. A clinician should review the timing and all medicines before changing treatment.

## Drusen

Yellow-white deposits that form beneath the retina, often with aging. Small deposits may cause no symptoms; many or larger deposits can be asso-

ciated with age-related macular degeneration and need eye examination.

## Dry Age-Related Macular Degeneration

Alternate name; see *Dry Macular Degeneration*.

## Dry AMD

Alternate name; see *Dry Macular Degeneration*.

## Dry Beriberi

The predominantly neurologic form of severe thiamine deficiency. It can cause symmetric peripheral neuropathy, burning or numb feet, muscle weakness, wasting, impaired reflexes, and difficulty walking.

### Alternate Names:

Tumarkin Drop Attack; Cryptogenic Drop Fall

## Dry Cell Battery Poisoning

Injury or poisoning caused by swallowing, leaking, or otherwise contacting a dry-cell battery. A swallowed button battery can rapidly burn the esophagus and is an emergency; do not induce vomiting and seek urgent medical help.

## Dry Cough

A cough that produces little or no mucus. Common causes include viral infection, asthma, acid reflux, allergies, smoking, certain medicines, and other airway or lung conditions; persistent cough needs assessment.

## Dry Cupping

A complementary treatment in which cups create suction on intact skin without making cuts. It may temporarily relieve some muscle pain, but evidence is limited; bruising, burns, skin injury, infection, and worsening of skin disease can occur.

## Dry Drowning

Alternate name; see *Drowning*.

## Dry Eye

A disorder in which tears are insufficient or evaporate too quickly, causing burning, grittiness, redness, watering, light sensitivity, or blurred vision. Long-

---

lasting disease can damage the eye surface.

**Alternate Names:**

Dry-Eye Disease; Keratoconjunctivitis Sicca; Dry Eyes

**Dry Eye Disease**

Alternate name; see *Dry Eye*.

**Dry Eye Syndrome**

Alternate name; see *Dry Eye*.

**Dry Gangrene**

A form of tissue death (coagulative necrosis) resulting from severe, prolonged insufficiency of arterial blood supply without superimposed bacterial infection. Most commonly affecting the toes, feet, or fingers in individuals with advanced peripheral artery disease, diabetes, or severe frostbite, the tissue becomes cold, dry, shrunken, and black (mummified). A sharp, distinct line of demarcation typically separates the dead tissue from adjacent healthy skin. Unlike wet gangrene, there is minimal swelling or foul odor unless a secondary bacterial infection develops. Management focuses on keeping the area dry and protected, performing urgent vascular evaluation and revascularization to restore blood flow, and elective surgical amputation when necessary.

**Alternate Names:**

Ischemic Gangrene; Mummification; Senile Gangrene; Arterial Gangrene

**Dry Hair**

Hair that feels rough, brittle, dull, or frizzy because its outer layer is damaged or lacks moisture. Heat, chemical treatments, weather, frequent washing, nutritional problems, and thyroid disease may contribute.

**Dry Ice**

Solid carbon dioxide that changes directly into a cold gas. Contact can cause severe cold burns, and gas in an enclosed space can displace oxygen and cause unconsciousness or suffocation; never seal it in a container.

**Dry Macular Degeneration**

The more common form of age-related macular degeneration, in which light-sensing cells in the macula gradually deteriorate without bleeding from new abnormal vessels. It can cause slow loss of detailed central vision.

**Alternate Names:**

Atrophic AMD; Dry AMD; Geographic Atrophy

**Dry Mouth**

A sensation of insufficient saliva or oral dryness, medically called xerostomia. It may result from medicines, dehydration, mouth breathing, anxiety, salivary-gland disease, radiation, or systemic illness and can impair speaking, swallowing, taste, oral comfort, and dental health.

**Alternate Names:**

Cottonmouth

**Dry Mouth Xerostomia**

Alternate name; see *Xerostomia*.

**Alternate Names:**

Asialia; Hyposalivation; Oral Dryness

**Dry Powder Inhaler**

A breath-powered device that delivers powdered medicine into the lungs. It does not require pressing and breathing at the same time, but the user must inhale strongly and correctly for the dose to reach the airways.

**Dry Skin**

Skin that is rough, scaly, tight, or itchy because it has lost moisture or its protective barrier is damaged. Aging, cold or dry air, frequent washing, eczema, medicines, and some illnesses can contribute.

**Dry Skin and Itching**

Xerosis with pruritus, often caused by low humidity, irritating products, eczema, medicines, or systemic disease. Persistent, widespread, severe, or sleep-disrupting itch warrants evaluation for dermatologic, liver, kidney, thyroid, blood, or neurologic causes.



---

## Dry Socket

Painful inflammation after tooth extraction when the protective blood clot is lost or breaks down too early. Severe pain may begin a few days later and spread to the ear or jaw; dental treatment is needed.

## Dry-Powder Inhaler

A device that delivers powdered medicine to the lungs when a person inhales through it. Correct preparation, a strong breath, and proper technique are needed; the device should not usually be used with a spacer.

## DSM

The Diagnostic and Statistical Manual of Mental Disorders, a reference that describes recognized mental-health diagnoses and their criteria. Clinicians use it with history and examination; it does not replace individual clinical judgment.

## DSPS

Alternate name; see *Delayed Sleep Phase Syndrome*.

## DSRCT

Alternate name; see *Desmoplastic Small Round Cell Tumor*.

## DSWPD

Alternate name; see *Delayed Sleep Phase Syndrome*.

## DTaP

A vaccine for infants and young children that protects against diphtheria, tetanus, and pertussis (whooping cough). It is given as a series, with later booster vaccines used as children grow older.

## Dual Antiplatelet Therapy

The simultaneous clinical administration of two distinct antiplatelet agents with complementary mechanisms of action (typically aspirin paired with a P2Y<sub>12</sub> receptor antagonist such as clopidogrel, ticagrelor, or prasugrel), indicated following acute coronary syndrome or percutaneous coronary intervention with stent placement.

**Alternate Names:**

DAPT

## Dual Energy X-Ray Absorptiometry

See *Bone Mineral Density Test*.

**Alternate Names:**

DXA; DEXA

## Dual-Energy CT

A CT scan that uses two X-ray energy levels to help distinguish materials in the body. It can improve evaluation of kidney stones, gout, blood vessels, bleeding, and some tissues while using the usual CT principles and risks.

## Dual-Energy X-Ray Absorptiometry

Alternate spelling; see *Dual Energy X-Ray Absorptiometry*.

## Dual-Photon Absorptiometry

An older test that used two photon energy levels to estimate bone mineral content. It has largely been replaced by dual-energy X-ray absorptiometry for measuring bone density and fracture risk.

## Duane Retraction Syndrome

A birth-related eye-movement disorder caused by abnormal nerve control of an eye muscle. Turning the eye inward or outward may be limited, and the eyeball may pull backward and the eyelids narrow.

**Alternate Names:**

DR Syndrome; Duane Syndrome; Stilling-Turk-Duane Syndrome

## Dubin-Johnson Syndrome

An inherited liver-transport disorder in which conjugated bilirubin is not moved normally into bile. It causes intermittent mild jaundice, often triggered by illness or stress, but usually does not cause serious liver damage.

## Dubowitz Syndrome

A rare developmental condition associated with poor growth, a small head, characteristic facial features, developmental or learning difficulties, and variable immune, skin, or behavioral problems. Features

---

differ among affected people.

### **Duchenne Muscular Dystrophy**

An X-linked recessive muscle disease, usually affecting boys, caused by disease-causing changes in the dystrophin gene. Progressive muscle weakness begins in childhood and may affect walking, breathing, the heart, and learning or behavior.

#### **Alternate Names:**

DMD; Duchenne Dystrophy

### **Duck Embryo Vaccine**

An older rabies vaccine made by growing inactivated rabies virus in duck embryos. It has largely been replaced by safer, more consistent cell-culture vaccines and is not commonly used today.

### **Duct**

A tube or channel that carries fluid within the body, such as bile, pancreatic juice, saliva, tears, or secretions from a gland. Blockage or injury can cause pain, swelling, infection, or impaired organ function.

### **Duct Ectasia**

Dilation and chronic periductal inflammation of a glandular duct system, most commonly referring to mammary duct ectasia of the subareolar breast ducts. See *Mammary Duct Ectasia*.

### **Ductal Carcinoma**

A cancer that begins in cells lining a duct. The term most often describes breast cancer or pancreatic cancer; its behavior and treatment depend on the organ, spread, and tumor features.

### **Ductal Carcinoma In Situ**

Abnormal cancer-like cells confined to the milk ducts of the breast without invasion into nearby breast tissue. It has not spread elsewhere, but some cases can become invasive if untreated.

#### **Alternate Names:**

DCIS Intraductal Carcinoma

### **Ductal-Dependent Lesions**

Serious congenital heart defects in which blood flow

to the lungs or body depends on the fetal ductus arteriosus remaining open after birth. Affected newborns may rapidly become blue, breathless, pale, or shocked and need urgent specialist care.

### **Ductless Glands**

Glands that release hormones directly into the bloodstream rather than through a tube. Examples include the pituitary, thyroid, adrenal glands, and pancreatic islets; their hormones regulate growth, metabolism, stress, and reproduction.

### **Ductus Arteriosus**

A blood vessel in the fetus connecting the pulmonary artery to the aorta, allowing blood to bypass the lungs before birth. It normally closes soon after birth; failure to close can strain the heart and lungs.

### **Ductus Venosus**

A fetal blood vessel that directs oxygen-rich blood from the umbilical vein toward the heart while partly bypassing the liver. It normally closes after birth; abnormal fetal flow can indicate circulatory or placental problems.

### **Duffy-Null Associated Neutrophil Count**

A naturally lower measured neutrophil count associated with the Duffy-null blood-group pattern, common in some populations. It usually does not increase infection risk and should not automatically be treated as disease.

### **Dugas Sign**

A clinical orthopedic test for anterior glenohumeral dislocation; the sign is positive if the patient is unable to touch the opposite shoulder with the hand of the injured extremity while the elbow remains resting in contact with the chest wall.

#### **Alternate Names:**

Dugas' Sign; Anterior Shoulder Dislocation Test

### **Duke Criteria**

A set of clinical, blood-culture, and heart-imaging findings used to classify suspected infective endocarditis. The criteria combine major and minor findings; the result supports diagnosis but must be

---

interpreted with the patient's history and examination.

**Alternate Names:**

Modified Duke Criteria; Duke Endocarditis Criteria

**Dukes Classification**

An older system for describing how far colorectal cancer has spread through the bowel wall, into nearby lymph nodes, or to distant organs. Modern staging systems are more detailed and are generally preferred.

**Dull Pain**

Aching, heavy, or pressure-like pain that is not sharp or stabbing. It can arise from muscles, organs, inflammation, or other tissues, and its significance depends on location, duration, associated symptoms, and examination findings.

**Duloxetine**

A medicine that increases serotonin and norepinephrine signaling in the brain. It treats depression, anxiety, some nerve pain, fibromyalgia, and chronic muscle or bone pain; nausea, sleep changes, sweating, and withdrawal symptoms may occur.

**Duloxetine Hydrochloride**

The salt form of duloxetine, a medicine used for depression, anxiety, some nerve pain, fibromyalgia, and chronic muscle or bone pain. It may cause nausea, sleep changes, sweating, blood-pressure changes, or withdrawal symptoms.

**Dumbbell Ossification**

Bone formation or a bone lesion shaped like a dumbbell, often extending through an opening between the spinal canal and surrounding tissues. Imaging is needed to identify its cause and relationship to nerves or other structures.

**Dumdum Fever**

An older name for visceral leishmaniasis, a parasitic infection transmitted by sandflies. It can cause prolonged fever, weight loss, enlarged spleen or liver, anemia, low blood counts, and life-threatening illness without treatment.

**Dumping Syndrome**

Symptoms caused by food moving too quickly from the stomach into the small intestine, often after stomach surgery. Early symptoms include cramps, diarrhea, flushing, sweating, dizziness, and a fast heartbeat; later low blood sugar may occur.

**Dunbar Syndrome**

Alternate name; see *Median Arcuate Ligament Syndrome*.

**Dunphy Sign**

A clinical sign of acute appendicitis and localized peritoneal inflammation characterized by a sharp exacerbation of right lower quadrant abdominal pain triggered by coughing.

**Alternate Names:**

Dunphy's Sign; Cough Sign of Appendicitis

**Duodenal**

Relating to the duodenum, the first part of the small intestine immediately beyond the stomach. This segment receives bile and pancreatic juice and begins digestion and absorption of nutrients.

**Duodenal Adenocarcinoma**

A cancer arising from gland-forming cells in the duodenum. It may cause abdominal pain, vomiting, bleeding, anemia, jaundice, or blockage; diagnosis usually requires imaging, endoscopy, and tissue sampling.

**Duodenal Atresia**

A birth defect in which the duodenum is completely blocked, preventing food and digestive fluid from passing. Newborns usually develop green vomiting and feeding difficulty; it is often associated with other birth defects and requires surgery.

**Duodenal Bulb**

The widened first part of the duodenum just beyond the stomach outlet. It receives partly digested food and is a common site for peptic ulcers; inflammation or scarring can narrow it and delay passage.

---

## Duodenal Disease

Any disorder affecting the duodenum, such as inflammation, ulcer, infection, bleeding, blockage, poor nutrient absorption, or a growth. Symptoms may include upper-abdominal pain, vomiting, anemia, or weight loss.

## Duodenal Diverticulum

A pouch that bulges outward from the duodenal wall, often near the openings of the bile and pancreatic ducts. It commonly causes no symptoms but can rarely lead to bleeding, inflammation, blockage, pancreatitis, or bile-duct infection.

## Duodenal Fistula

An abnormal connection between the duodenum and another organ, bowel segment, or skin. Digestive fluid may leak through it, causing infection, skin damage, dehydration, electrolyte loss, or poor nutrition.

## Duodenal Hemorrhage

Bleeding from the duodenum, often caused by a peptic ulcer, inflammation, abnormal blood vessel, injury, or cancer. It may cause black stools, vomiting blood, weakness, dizziness, anemia, or shock.

## Duodenal Infection

Infection affecting the duodenum, caused by bacteria, parasites, viruses, or spread from nearby digestive disease. Symptoms can include upper-abdominal pain, nausea, vomiting, diarrhea, fever, or bleeding.

## Duodenal Neoplasm

An abnormal growth in the duodenum that may be benign or cancerous. It can cause pain, bleeding, vomiting, blockage, anemia, or jaundice; evaluation depends on its size, location, appearance, and tissue type.

## Duodenal Obstruction

A partial or complete blockage that prevents food and digestive fluid from passing through the duodenum. It can cause repeated vomiting, abdominal swelling, pain, and dehydration and may need ur-

gent imaging and treatment.

## Duodenal Papilla

A small raised opening in the duodenal wall where bile and pancreatic juice enter the intestine. Narrowing, blockage, or inflammation at this site can cause jaundice, pancreatitis, or digestive symptoms.

## Duodenal Perforation

A hole through the full thickness of the duodenal wall, usually caused by a severe ulcer, injury, or a medical procedure. Sudden severe abdominal pain, rigid muscles, and infection of the abdominal lining require emergency treatment.

## Duodenal Stenosis

Narrowing of the duodenum that partly blocks passage of food and digestive fluid. It may cause vomiting, abdominal swelling, pain, early fullness, poor feeding, or dehydration; treatment depends on the cause and severity.

## Duodenal Tissue Culture

Microbiologic culture of endoscopic mucosal biopsy specimens obtained from the duodenum to diagnose atypical bacterial, mycobacterial, or fungal infections of the proximal small intestine.

### Alternate Names:

Duodenal Biopsy Culture; Small Intestine Mucosal Culture

## Duodenal Ulcer

An open sore in the duodenal lining, commonly related to *Helicobacter pylori* infection or anti-inflammatory medicines. It may cause upper-abdominal pain and can bleed, perforate the bowel, or obstruct food passage.

## Duodenitis

Inflammation of the duodenal lining, which may result from *Helicobacter pylori*, excess acid, anti-inflammatory medicines, infection, celiac disease, or inflammatory bowel disease. It can cause upper-abdominal pain, nausea, vomiting, or bleeding.

---

## Duodenoscope

A flexible viewing tube passed through the mouth and stomach to examine the duodenum and reach the openings of the bile and pancreatic ducts. It may be used during procedures to remove stones or place stents.

## Duodenostomy

A surgically created opening from the duodenum to the skin or another part of the intestine. It may be used in selected cases for feeding, drainage, or bypassing a blockage.

## Duodenum

The first part of the small intestine, between the stomach and jejunum. It receives bile and pancreatic juice, mixes them with stomach contents, and begins much of the digestion and absorption of nutrients.

## Duplex Doppler Ultrasound Scanning

An ultrasound test that shows the structure of blood vessels and measures the direction and speed of blood flow. It can detect narrowing, blockage, clots, aneurysms, or abnormal connections without radiation.

### Alternate Names:

Duplex Ultrasound

## Duplex Surveillance

Planned examinations and ultrasound checks of a blood vessel, graft, stent, aneurysm, or dialysis access. Monitoring can detect recurrent narrowing, clotting, leakage, or failure before serious symptoms develop.

## Duplex Ultrasound

An ultrasound examination that combines pictures of body structures with Doppler measurement of blood flow. It is commonly used to assess blood vessels and can identify narrowing, blockage, clots, or abnormal flow.

## Duplication

A genetic change in which a section of DNA or a chromosome is copied one or more extra times. The

additional genetic material can alter how much of a gene is active and may cause a health condition, although some duplications have little or no effect.

## Dupuytren Contracture

Progressive thickening and shortening of tissue in the palm that pulls one or more fingers toward the palm and can limit hand function.

### Alternate Names:

Dupuytren Disease; Palmar Fibromatosis

## Dupuytren's Disease

A usually progressive disorder in which the palmar fascia, a layer of fibrous tissue beneath the skin of the palm and fingers, becomes thick and forms nodules and cords. The cords can gradually prevent one or more fingers, especially the ring or little finger, from straightening and may limit hand function. It usually begins in middle age or later and is more common in men, although anyone can develop it. The condition begins in the palm and is not primarily a disease of the finger joints; the later bending deformity is called Dupuytren contracture.

## Dupuytren's Disease

A progressive thickening and shortening of tissue in the palm that pulls one or more fingers into flexion. It may limit hand function; treatment ranges from observation and therapy to needle procedures or surgery.

## Dura

The tough outermost layer of protective tissue surrounding the brain and spinal cord. It helps contain cerebrospinal fluid, supports blood vessels, and can become inflamed, torn, thickened, or infected.

## Dura Mater

The tough outer layer of tissue surrounding and protecting the brain and spinal cord. It helps support blood vessels and contains the fluid-filled coverings around the central nervous system.

## Durable Power Of Attorney

A legal authorization allowing a designated person to act or make decisions for another person when

---

specified conditions are met. In healthcare, it may appoint a healthcare proxy.

### **Dural Arteriovenous Fistula**

An abnormal direct connection between an artery and a vein in the tough covering around the brain or spinal cord. It may cause a pulsing sound in the ear, headache, seizures, vision or other nerve problems, or bleeding.

#### **Alternate Names:**

DAVF

### **Dural Arteriovenous Fistulas**

Abnormal links between arteries and veins in the protective covering around the brain or spinal cord. They can cause headaches, pulsating sounds, seizures, bleeding, or neurologic problems and may need vascular treatment.

### **Dural Sac**

The tough, tubular covering around the spinal cord, its surrounding fluid, and the spinal nerve roots inside the spinal canal. A tear can allow spinal-fluid leakage and cause positional headaches.

### **Dural Venous Sinuses**

Channels between layers of the brain's tough covering that collect blood and cerebrospinal fluid and drain them toward the neck veins. A clot in these channels can cause headache, seizures, vision problems, or stroke-like symptoms.

### **Duramorph**

A brand name for preservative-free morphine used by injection, including in epidural or intrathecal analgesia; respiratory depression is a serious risk.

### **Durapatite**

Another name for hydroxyapatite, the calcium-phosphate mineral that gives bones and teeth much of their hardness. Its amount and arrangement contribute to bone strength and can change in metabolic bone disease.

### **Duroziez Sign**

An auscultatory vascular sign of severe aortic regurgitation consisting of a double murmur heard over the femoral artery—a systolic murmur generated by proximal compression and a diastolic murmur generated by distal compression with the stethoscope diaphragm.

#### **Alternate Names:**

Duroziez's Sign; Duroziez Double Murmur

### **Durules**

A formulation designed to release a medicine slowly over an extended period. Such tablets or pellets must be used exactly as directed because crushing or chewing may release too much medicine at once.

### **Dust**

Small solid particles suspended in air or settled on surfaces. Inhaled dust can irritate the nose and lungs; some types can cause allergy, asthma, infection, poisoning, or long-term lung disease.

### **Dust Disease**

Lung or other illness caused by repeated exposure to harmful dust, often at work. Effects depend on the dust and exposure and may include asthma, scarring of lung tissue, chronic cough, or cancer.

### **Dust Mite Allergy**

An immune reaction to proteins from tiny mites living in household dust. It can cause sneezing, blocked or itchy nose, watery eyes, cough, wheezing, eczema flares, or asthma symptoms.

#### **Alternate Names:**

House Dust Mite Hypersensitivity; Dust Mite Sensitization

### **Dutasteride**

A medicine that lowers the hormone dihydrotestosterone and gradually shrinks an enlarged prostate. It improves urinary symptoms and lowers retention risk but may reduce sexual function and lowers the blood-test level used to screen for prostate cancer.



---

## DVT

A blood clot in a deep vein, usually in the leg. It can cause swelling, pain, warmth, or redness and may travel to the lungs, causing a life-threatening pulmonary embolism.

## DVT Prevention

Steps to reduce the risk of a deep-vein blood clot, including early movement, leg exercises, compression devices, and preventive anticoagulant medicine when appropriate. The choice depends on clot risk and bleeding risk.

## Dwarfism

A medical condition causing unusually short adult height, often defined as about 4 feet 10 inches (147 cm) or less. Causes include skeletal disorders, growth-hormone problems, genetic conditions, and chronic disease; features vary by cause.

## DWI FLAIR Mismatch

A pattern on brain MRI in which an area is abnormal on diffusion images but not yet clearly abnormal on FLAIR images. In selected people with an unknown stroke onset time, it may help identify potentially treatable recent stroke.

## Dx

A medical abbreviation for diagnosis or diagnostic assessment. Because abbreviations can be misunderstood, the full word should be used in patient instructions and other documents where clarity matters.

## DXA Scan

Alternate name; see *Dual Energy X-Ray Absorptiometry*.

## Dyad

A pair of related people, structures, cells, or measurements considered together. Medical examples include two linked findings or two parts of a cellular or body structure that function together.

## Dye

A colored substance used to stain tissue, identify cells, make structures visible, or provide contrast during a medical test. Some dyes can cause allergic reactions or affect the kidneys, especially when injected.

## Dye Lasers

Lasers that use a liquid organic dye to produce adjustable wavelengths of light. In medicine, selected wavelengths can treat blood-vessel or pigmented skin lesions; eye and skin protection is essential.

## Dye Remover Poisoning

Harm from swallowing, inhaling, or getting dye-removing chemicals on the skin or eyes. Depending on the ingredients, exposure can cause burns, breathing problems, vomiting, poisoning, or organ injury and needs poison-control advice.

## Dying

The physical process of approaching death, often involving increasing weakness, sleepiness, reduced appetite and drinking, changes in breathing, and less responsiveness. Emotional, social, cultural, and spiritual support may help the person and family.

## Dynamin Gtpase

A family of GTP-binding proteins that constrict membranes during vesicle budding, endocytosis, and other cell-membrane processes.

## Dynamometer

An instrument that measures force, commonly hand-grip or muscle strength. Clinicians use the result to compare sides, identify weakness, monitor recovery, and assess physical function during rehabilitation or neurologic examination.

## Dynein Atpase

The energy-using activity of dynein, a motor protein that moves materials inside cells along microtubules. Dynein also powers the movement of cilia and flagella, which help move fluid or cells.

---

## **Dynorphin**

A naturally occurring opioid peptide that mainly activates kappa-opioid receptors in the nervous system. It influences pain, stress, mood, movement, and hormone regulation.

## **Dyphylline**

A bronchodilator used in some countries to relax airway muscles in asthma or chronic obstructive lung disease. It may cause nausea, tremor, headache, nervousness, or a fast heartbeat.

## **Dysaesthesia**

An unpleasant abnormal sensation, such as burning, electric-shock pain, pins and needles, or excessive sensitivity to light touch. It commonly results from nerve injury, diabetes, medicines, or other nerve disease.

## **Dysania**

Persistent or excessive difficulty getting out of bed or waking up, often associated with sleep disorders, depression, medication effects, or other medical causes. It is a descriptive symptom rather than a formal diagnosis by itself.

## **Dysarthria**

Slurred, weak, slow, or poorly coordinated speech caused by a problem in the brain, nerves, or muscles used for speaking. Sudden dysarthria can be a sign of stroke.

## **Dysautonomia**

A disorder of the automatic nervous system that controls blood pressure, heart rate, sweating, temperature, digestion, bladder function, and sexual function. Causes include nerve disease, diabetes, medicines, and inherited conditions.

## **Dysbarism**

Illness caused by a change in surrounding pressure, usually during diving, flying, or hyperbaric treatment. It includes decompression sickness and pressure injury to the ears, sinuses, lungs, or other tissues.

## **Dysbiosis**

An imbalance or change in the community of microorganisms living in the body, especially the intestine. Antibiotics, infection, diet, and illness may contribute; dysbiosis is associated with some digestive and inflammatory disorders, but it is not always a proven cause.

### **Alternate Names:**

Gut Microbiome Imbalance

## **Dyscalculia**

A persistent difficulty understanding numbers, quantities, or arithmetic that is not explained by inadequate teaching or general intellectual disability. It may begin during development or follow brain injury and can affect school or daily tasks.

## **Dyschezia**

Difficult or painful passage of stool, often with excessive straining or a feeling of incomplete emptying. Causes include constipation, pelvic-floor coordination problems, anal disorders, and other bowel disease.

## **Dyscrasia**

An older broad term for an abnormal blood or bone-marrow condition. It may describe abnormal blood cells, clotting, or proteins in the blood, so the specific disorder should be named when possible.

### **Alternate Names:**

Blood Dyscrasia

## **Dyscrasias**

An older plural term for disorders affecting blood, bone marrow, or body-fluid composition. It may refer to abnormal blood cells, clotting, or abnormal proteins, so the specific disorder must be identified.

## **Dysdiadochokinesia**

Difficulty making quick, repeated movements in opposite directions, such as rapidly turning the hands over and back. It can occur with problems in the cerebellum or its nerve pathways.

### **Alternate Names:**

Dysdiadochokinesis; Adiadochokinesia

---

### **Dysdiadochokinesis**

Difficulty making quick, repeated movements in opposite directions, such as rapidly turning the hands over and back. It can occur with problems in the cerebellum or its nerve pathways.

### **Dysembryoplastic Neuroepithelial Tumor**

A usually slow-growing, low-grade brain tumor that develops most often in the outer part of the brain in children or young adults. It commonly causes long-standing focal seizures and is often treatable with surgery.

### **Dysentery**

Diarrhea with visible blood, often with cramps, fever, mucus, and an urgent need to pass stool. Several bacteria and parasites can cause it, so the cause needs medical assessment.

### **Dysesthesia**

An unpleasant abnormal sensation, such as burning, electric-shock pain, or painful sensitivity to light touch. It may result from injury or disease of nerves in the brain, spinal cord, or limbs.

### **Dysferlinopathy**

A group of inherited muscle diseases caused by changes in both copies of the DYSF gene, which makes a muscle-membrane repair protein. Weakness often begins in adolescence or early adulthood, affecting the hips, shoulders, or calves.

#### **Alternate Names:**

DYSF-Related Muscular Dystrophy; Dysferlin Deficiency

### **Dysfibrinogenemia**

An inherited or acquired condition in which fibrinogen, a blood-clotting protein, has abnormal structure or function. It may cause unusual bleeding, excessive clotting, or no symptoms and is diagnosed with clotting and specialized blood tests.

### **Dysfunction**

Abnormal, reduced, or poorly controlled function of an organ, tissue, body system, or process. It

may be temporary or permanent and can result from disease, injury, medicines, inherited conditions, stress, or aging.

### **Dysfunctional Uterine Bleeding**

Older term; see *Abnormal Uterine Bleeding*.

### **Dysfunctional Voiding**

Abnormal coordination between the bladder muscle and the muscles around the urethra during urination. It may cause a weak or interrupted stream, straining, incomplete emptying, urinary infections, wetting, or constipation, especially in children.

### **Dysgammaglobulinemia**

An abnormal level or pattern of antibodies in the blood, with some antibody types low or absent and others normal or increased. It may cause repeated infections, autoimmune problems, or no symptoms and is identified by blood tests.

### **Dysgerminoma**

A cancer of the ovary arising from germ cells, the cells that can develop into eggs. It occurs most often in adolescents and young adults and may cause pelvic pain, swelling, or an abdominal mass.

### **Dysgeusia**

An altered sense of taste, including reduced, distorted, or unpleasant taste. Causes include viral infection, mouth or nose disease, nerve injury, nutritional deficiency, dry mouth, medicines, and other illnesses.

### **Dysgraphia**

A persistent difficulty with handwriting, spelling, or expressing ideas in writing despite appropriate learning opportunities. It may reflect problems with motor skills, language, attention, or brain function and can occur with other learning disorders.

### **Dyshidrosis**

A skin condition causing small, intensely itchy blisters on the palms, sides of the fingers, or soles. The skin may peel, crack, or become sore; triggers in-

---

clude sweating, irritation, allergy, and stress.

**Alternate Names:**

Dyshidrotic Eczema; Pompholyx; Vesicular Hand Dermatitis

**Dyshidrotic Eczema**

A recurrent, intensely pruritic dermatosis affecting the lateral margins of the fingers, palms, and soles, characterized by deep-seated, "tapioca-like" intraepidermal vesicles that subsequently resolve with desquamation and fissuring.

**Alternate Names:**

Pompholyx; Dyshidrosis

**Dyskeratosis Congenita**

A rare hereditary telomere biology disorder caused by mutations in telomere maintenance genes, characterized by the classic triad of reticulated reticular skin hyperpigmentation, dystrophic nails, and oral mucosal leukoplakia, associated with a high risk of progressive bone marrow failure and pulmonary fibrosis.

**Alternate Names:**

Zinsser-Cole-Engman Syndrome; Telomere Biology Disorder

**Dyskinesia**

Abnormal movement that is often involuntary, excessive, repetitive, twisting, or irregular. It may result from neurologic disease or medicines, including long-term treatment for movement disorders, and can interfere with walking, speaking, or daily activities.

**Dyslexia**

A learning disorder involving persistent difficulty recognizing words accurately or fluently, decoding them, spelling, or linking letters with speech sounds. It is not caused by low intelligence or poor motivation and may continue into adulthood.

**Alternate Names:**

Specific Reading Disorder; Developmental Dyslexia

**Dyslipidemia**

An unhealthy level or pattern of fats in the blood,

such as high LDL cholesterol, high triglycerides, low HDL cholesterol, or a combination. It can promote artery disease and, when triglycerides are very high, pancreatitis.

**Dyslipidemias**

Conditions involving abnormal blood fats, including high LDL cholesterol, high triglycerides, low HDL cholesterol, or mixed changes. They can increase the risk of artery disease and pancreatitis when triglycerides are very high.

**Dysmenorrhea**

Painful menstrual periods, usually causing cramping in the lower abdomen and sometimes back pain, nausea, diarrhea, headache, or dizziness. Severe, new, or worsening pain may indicate an underlying pelvic disorder.

**Dysmenorrhoea**

Painful menstrual periods. Primary pain occurs without another pelvic disease, while secondary pain results from a condition such as endometriosis, fibroids, adenomyosis, pelvic infection, or an intrauterine device.

**Dysmetria**

Inability to judge or control the distance, force, or speed of a voluntary movement. A person may overshoot or undershoot a target; it commonly reflects dysfunction of the cerebellum or its connecting pathways.

**Dysmyelogenic Leukodystrophy**

An older or nonspecific term for a leukodystrophy in which myelin develops abnormally. Modern diagnosis generally identifies the specific genetic or acquired disorder and its pattern of hypomyelination, dysmyelination, or demyelination.

**Alternate Names:**

Dysmyelinating Leukodystrophy

**Dysorexia**

An abnormal appetite, including little desire to eat, excessive hunger, or disordered eating. It may result from illness, medicines, emotional distress,

---

metabolic disease, or an eating disorder and can affect nutrition and weight.

### **Dysostoses**

Inherited or developmental conditions in which one or more bones form or grow abnormally. They may affect the skull, face, spine, ribs, arms, or legs and can alter body shape, movement, growth, hearing, breathing, or other functions.

### **Dyspareunia**

Recurrent or persistent genital or pelvic pain before, during, or after sexual activity, especially attempted vaginal penetration. Pain may be at the vaginal opening or deeper in the pelvis and can result from dryness, infection, skin disease, pelvic-floor muscle spasm, endometriosis, inflammation, injury, or emotional factors.

### **Dyspepsia**

The medical term for indigestion, describing persistent or recurring discomfort centered in the upper stomach area. Common symptoms include upper abdominal pain or burning, feeling uncomfortably full after normal meals, and feeling full too soon while eating. In the majority of people (around 70% to 80%), tests reveal no ulcers or visible physical damage, a condition termed functional dyspepsia. Identifiable causes include peptic ulcers, acid reflux, *Helicobacter pylori* bacterial infection, and medications such as nonsteroidal anti-inflammatory drugs (NSAIDs). Doctors evaluate persistent symptoms through bacterial breath or stool tests, medication trials, or an upper endoscopy camera examination.

#### **Alternate Names:**

Indigestion; Functional Dyspepsia; Upset Stomach

### **Dyspeptic**

Describing indigestion symptoms such as upper-abdominal discomfort or burning, early fullness, bloating, belching, or nausea, especially after eating.

### **Dysphagia**

The medical term for difficulty swallowing, describing trouble moving food, liquids, or saliva from the mouth to the stomach. It is divided into two main

types: oropharyngeal dysphagia (a mouth or throat problem, often caused by nerve or muscle disorders like stroke or Parkinson's disease, leading to coughing or choking right as swallowing begins) and esophageal dysphagia (a food-pipe problem, caused by narrowings, growths, or muscle spasms, where food feels stuck behind the breastbone). Potential complications include food entering the airways (aspiration) and weight loss. Doctors identify the cause using specialized X-ray video swallow studies (barium swallow) or an upper endoscopy.

#### **Alternate Names:**

Difficulty Swallowing; Deglutition Disorder

### **Dysphagia Diet**

Food and drinks altered in texture or thickness to make swallowing safer after a swallowing assessment. The plan should reduce aspiration risk while providing enough calories, protein, fluids, and enjoyable foods.

### **Dysphagia Lusoria**

Difficulty swallowing caused by extrinsic esophageal compression from an aberrant right subclavian artery (arteria lusoria) that arises abnormally as the last branch of the aortic arch and crosses retroesophageally to reach the right arm.

#### **Alternate Names:**

Transfer Dysphagia; High Dysphagia

### **Dysphagia Rehabilitation**

Therapy to improve safe swallowing, nutrition, and hydration. It may include swallowing exercises, posture or pacing strategies, food-texture changes, and caregiver education based on the cause and assessment.

### **Dysphasia**

Difficulty using or understanding language because of a problem in the brain's language areas. It may affect speaking, understanding, reading, or writing; many clinicians use the clearer term aphasia.

### **Dysphonia**

An abnormal voice, such as hoarseness, breathiness, roughness, strain, weakness, or loss of voice. Causes

---

include infection, vocal-cord swelling or growths, nerve problems, voice overuse, reflux, and cancer; persistent symptoms need evaluation.

### **Dysphonia Spastica**

Alternate name; see *Spasmodic Dysphonia*.

### **Dysphoria**

A state of marked emotional distress, unease, or dissatisfaction. Gender dysphoria specifically means distress related to a difference between experienced gender and sex assigned at birth; it is not itself a mental illness.

### **Dysplasia**

Abnormal size, shape, or arrangement of cells in a tissue. It is not cancer, but some forms—especially when severe or persistent—can increase the chance of cancer and may need monitoring or treatment.

### **Dysplasia Epiphysealis Hemimelica (Trevor Disease)**

A rare developmental overgrowth of cartilage in one side of an epiphysis, usually affecting a lower-limb joint in childhood. It can cause asymmetric bony masses, pain, stiffness, deformity, or restricted movement and is evaluated with radiographs or MRI.

### **Dysplastic Melanocytic Nevi Pathology**

Melanocytic nevi with architectural disorder and variable cytologic atypia, often showing bridging of rete ridges, lamellar fibroplasia, and atypical melanocytes. The degree of atypia is graded and must be distinguished from melanoma.

### **Dysplastic Nevi**

Atypical moles with unusual size, shape, color, border, or microscopic features. Most do not become melanoma, but having many can increase melanoma risk; a changing, bleeding, or suspicious mole should be examined.

### **Dysplastic Nevus**

An atypical mole that may be larger, unevenly col-

ored, or have an irregular border. Most do not become cancer, but having many atypical moles can increase melanoma risk; changing lesions should be examined.

### **Dyspnea**

The feeling of difficult, uncomfortable, or insufficient breathing. Causes include asthma, lung or heart disease, anemia, infection, muscle weakness, anxiety, and other conditions.

### **Dyspnea On Exertion**

Shortness of breath that occurs or worsens with physical activity. It may result from heart or lung disease, anemia, poor fitness, or other conditions; new, severe, or rapidly worsening symptoms need prompt assessment.

### **Dyspnoea**

Alternate name; see *Dyspnea*.

### **Dyspraxia**

A developmental difficulty planning and coordinating movements that interferes with daily activities, learning, or motor skills. It is not explained by muscle weakness, loss of sensation, or inadequate opportunity to learn.

**Alternate Names:**

Developmental Coordination Disorder

### **Dysraphism**

A birth defect caused by incomplete joining of developing body structures, especially the spinal cord or spinal bones. Effects range from a small skin mark to weakness, loss of sensation, or bladder and bowel problems.

### **Dysrhythmia**

An abnormal heart rhythm that may be too fast, too slow, or irregular. Some cause no symptoms; others cause palpitations, dizziness, fainting, chest pain, stroke, or inadequate blood flow.

### **Dyssomnia**

Broad older term; see the specific diagnosis, such



---

as *Insomnia*, *Hypersomnia*, or *Circadian Rhythm Sleep-Wake Disorder*.

### **Dyssomnias**

Sleep disorders that mainly cause difficulty falling asleep, staying asleep, waking at the desired time, or staying awake. Causes include sleep schedules, medical conditions, medicines, mood disorders, and abnormal sleep regulation.

### **Dysthyroid Eye Disease**

An autoimmune inflammatory disorder of the orbit most commonly associated with Graves disease (and occasionally Hashimoto thyroiditis or euthyroid states). Autoantibodies targeting TSH receptors on orbital fibroblasts stimulate excessive production of hyaluronic acid and adipogenesis, causing marked enlargement and edema of extraocular muscle bellies (characteristically sparing their tendons) and orbital fat expansion. Symptoms include forward protrusion of the eyes (proptosis), upper eyelid retraction, lid lag, grittiness, double vision (diplopia), and, in severe cases, vision loss from corneal ulceration or compressive optic neuropathy at the orbital apex. It is strongly aggravated by smoking. Management includes smoking cessation, artificial tears, intravenous corticosteroids, targeted biologic therapy (teprotumumab), orbital radiation, and surgical orbital decompression.

#### **Alternate Names:**

Thyroid Eye Disease; Graves' Ophthalmopathy; Graves' Orbitopathy; Dysthyroid Ophthalmopathy; TED

### **Dystocia**

Difficult or unusually slow labor caused by ineffective contractions, the baby's size or position, or the mother's pelvis or soft tissues. It can increase risks for both parent and baby and may require assisted delivery or cesarean birth.

### **Dystonia**

A neurological movement disorder in which changes in the brain's movement signals make muscles tighten or contract without the person intending them to. This can cause slow, repeated or twisting

movements, tremor, or abnormal and sometimes painful postures. It may affect one body part, several connected areas, or most of the body, and symptoms may be triggered by movement, stress, or tiredness. Dystonia may be inherited, have no known cause, or result from brain disease or injury, infection, poisoning, or certain medicines. It can begin at any age and may remain stable or worsen over time.

### **Dystonia Disorder**

A disorder causing repeated or sustained involuntary muscle contractions, which produce twisting movements, tremor, or abnormal postures. Symptoms may be limited to one area or involve the whole body.

### **Dystonin**

A protein made from the DST gene that helps attach the internal framework of cells to other cell structures. Harmful gene changes can cause inherited nerve disease, muscle weakness, and fragile or blistering skin.

### **Dystrophia Unguium**

Abnormal nail growth, shape, color, or texture. Causes include injury, fungal infection, inflammation, poor nutrition, medicines, reduced blood flow, and inherited disorders; treatment depends on the cause.

### **Dystrophic Calcification**

Buildup of calcium salts in damaged, dead, or chronically inflamed tissue even though the blood calcium level is normal. It can occur in scars, diseased valves, tumors, or injured muscles and may impair function.

### **Dystrophin**

A large sub-sarcolemmal cytoskeletal structural protein that anchors the internal actin cytoskeleton of skeletal and cardiac muscle fibers to the extracellular matrix through the dystrophin-associated glycoprotein complex; mutations in the *DMD* gene lead to Duchenne and Becker muscular dystrophies.

---

### **Dystrophin-Glycoprotein Complex**

A group of proteins that connects the internal framework of muscle cells to their surrounding support tissue and protects the cell membrane during contraction. Changes in its components can cause several inherited muscular dystrophies.

#### **Alternate Names:**

DGC; Dystrophin-Associated Glycoprotein Complex (DAGC)

### **Dystrophy**

Abnormal development, maintenance, or degeneration of a tissue or organ, often caused by an inherited or metabolic condition. Effects depend on the tissue and may include muscle weakness, poor vision, fragile structures, or loss of function.

### **Dysuria**

Pain, burning, or discomfort during urination. Common causes include bladder or urethral infection, vaginal or penile inflammation, stones, irritation, and sexually transmitted infections; fever, flank pain, pregnancy, or blood in urine needs medical assessment.

#### **Alternate Names:**

Painful Urination

### **Déjà Vu**

A feeling that a current situation has happened before, even though it is probably new. Occasional episodes are common; frequent episodes with impaired awareness or unusual movements may occur with seizures.

# E

## **E Coli Enteritis**

Infection and inflammation of the intestine caused by disease-producing strains of *Escherichia coli*. It may cause watery or bloody diarrhea, cramps, vomiting, and dehydration; some strains can injure the kidneys.

## **E Coli Infection**

Infection caused by *Escherichia coli* bacteria. Depending on the strain and site, it may cause diarrhea, urinary infection, wound or abdominal infection, newborn meningitis, or bloodstream infection; some strains cause kidney injury.

## **E-Cigarettes And E-Hookahs**

Electronic devices that heat a liquid to produce an inhaled aerosol. Many contain nicotine, which can cause dependence; the aerosol may contain irritants and toxic substances and is not harmless water vapor.

## **E-D Syndrome**

Alternate name; see *Ehlers-Danlos Syndrome*.

## **E. Coli**

A group of bacteria commonly found in the intestines of people and animals. Most are harmless, but some strains cause diarrhea, urinary infection, or serious illness such as bloodstream infection or kidney injury.

## **E. coli Infection**

Alternate spelling; see *E Coli Infection*.

## **E. Coli O157:H7**

A strain of *Escherichia coli* that produces Shiga toxin and can cause severe cramps and bloody diarrhea. It may injure the kidneys, especially in children; urgent medical advice is needed for bloody diarrhea or reduced urination.

## **Eagle Syndrome**

A rare cause of throat, face, jaw, or neck pain related to an unusually long bony projection near the skull base or a hardened nearby ligament. Pain may worsen with swallowing or turning the head.

## **Eales Disease**

An uncommon inflammatory disorder of small retinal veins, usually affecting otherwise healthy young adults. Blocked vessels can cause abnormal new vessels and bleeding inside the eye, leading to floaters or reduced vision.

## **Ear**

The organ for hearing and balance. The outer ear collects sound, the middle ear transmits vibrations through the eardrum and tiny bones, and the inner ear converts sound and movement into signals sent to the brain.

## **Ear Barotrauma**

Ear injury caused by unequal pressure between the middle ear and surrounding air or water, often during flying or diving. It may cause pain, fullness, blocked hearing, ringing, dizziness, bleeding, or a torn eardrum.

---

## Ear Canal

The passage between the outer ear and eardrum. Its skin and earwax protect the middle ear, but infection, injury, foreign objects, or wax buildup can cause pain, discharge, or reduced hearing.

## Ear Cartilage

Flexible connective tissue that forms much of the outer ear and gives it shape. Injury or infection can cause pain, swelling, deformity, or damage that permanently changes the ear.

## Ear Disease

Any disorder of the outer, middle, or inner ear. It may cause pain, discharge, hearing loss, ringing, dizziness, infection, or balance problems; diagnosis requires examination of the ear and sometimes hearing or imaging tests.

## Ear Disorders

Disorders affecting the outer ear, ear canal, eardrum, middle-ear bones, pressure-equalizing tube, cochlea, or balance organs. They include infection, injury, blockage, inflammation, hearing disorders, and growths. Disorders of the outer, middle, or inner ear. Symptoms, examination findings, causes, and treatment depend on the affected structure and may include pain, discharge, hearing loss, ringing, dizziness, or imbalance.

## Ear Drainage Culture

A laboratory test of fluid draining from the ear to identify bacteria or fungi. Results help guide treatment but must be interpreted with the examination because normal organisms, contamination, or colonization can affect the result.

## Ear Examination

Inspection of the outer ear and ear canal and viewing of the eardrum, usually with an otoscope. Hearing, balance, or additional tests may be added when symptoms suggest middle- or inner-ear disease.

## Ear Foreign Body

An object lodged in the ear canal, most often in a child. It may cause pain, reduced hearing, bleeding,

discharge, or injury; removal should be done with direct visualization and suitable instruments.

### Alternate Names:

Auditory Canal Foreign Body; Aural Foreign Body

## Ear Infection

Infection or inflammation affecting part of the ear, including the outer ear, ear canal, middle ear, or inner ear. It may cause ear pain, fever, discharge, reduced hearing, dizziness, or irritability in children.

## Ear Microbiome

The community of microorganisms living on or in the ear, especially the ear canal. They interact with skin and earwax and may influence infection risk; moisture, cleaning, hearing devices, inflammation, and illness can change the community.

## Ear Neoplasm

An abnormal growth in the outer, middle, or inner ear that may be benign or cancerous. It can cause a persistent lump, pain, discharge, hearing loss, dizziness, or facial weakness and may require imaging or biopsy.

## Ear Speculum

A small cone-shaped tip fitted to an otoscope to direct light into the external ear canal and permit examination of the canal and tympanic membrane. Disposable and reusable types are available in different sizes.

## Ear Tag

A small, skin-covered projection on or near the outer ear, usually present from birth. It is generally harmless, but examination may be appropriate if it causes problems or occurs with other birth findings.

## Ear Thermometer

A device that estimates body temperature by measuring heat in the ear canal near the eardrum. Correct positioning is important; earwax, a small ear canal, or an ear infection can affect the reading.

---

## Ear Tube Insertion

A procedure placing a small ventilation tube through the eardrum to allow air into the middle ear and fluid to drain. It may help repeated infections or persistent fluid but can cause drainage, blockage, scarring, or a persistent opening.

## Ear Wax

A mixture of gland secretions and shed skin that protects and cleans the ear canal. A buildup can cause reduced hearing, fullness, ringing, itching, or discomfort; cotton swabs and inserted objects can worsen blockage or injure the ear.

### Alternate Names:

Cerumen; Earwax

## Earache

Pain felt in the ear or nearby area. It may arise inside the ear or be referred from the teeth, jaw, throat, or neck; identifying the source requires an examination when pain persists or is severe.

### Alternate Names:

Otalgia

## Eardrum

A thin membrane separating the ear canal from the middle ear. Sound makes it vibrate, and the vibration moves the tiny middle-ear bones; infection, pressure, or injury can tear it or reduce hearing.

## Eardrum Repair

Treatment to close a hole or repair damage in the eardrum, often using a tissue graft or other surgical technique. It may improve hearing and prevent infection, but healing is not always complete.

## Earliest Signs Of Increased Intracranial Pressure

Early warning signs may include worsening headache, vomiting, blurred or double vision, drowsiness, slowed thinking, or behavior change. Causes include bleeding, swelling, infection, a growth, or excess brain fluid; progressive symptoms need urgent assessment.

## Earlobes

The soft lower parts of the outer ears, made mainly of skin and fatty or connective tissue without cartilage. They may be affected by piercing injury, infection, cysts, allergic reactions, or tears.

## Early Afterdepolarization

An extra abnormal electrical signal during the heart's recovery phase after each beat. It is more likely when the QT interval is prolonged and can trigger dangerous abnormal rhythms.

### Alternate Names:

EAD; Early After-Depolarization

## Early Childhood Tooth Decay

Early-childhood tooth decay is bacterial acid destruction of primary teeth, promoted by frequent sugars, poor brushing, enamel defects, and prolonged bottle or nursing exposure.

## Early Decelerations

A gradual temporary slowing of the fetal heart rate that begins and ends with a uterine contraction, usually from pressure on the baby's head. When isolated, it is generally reassuring, but the complete tracing and clinical situation matter.

## Early Detection

Finding a disease at an early stage, before it causes major complications or spreads. It may result from screening or prompt evaluation of symptoms and can improve treatment options for some conditions.

## Early Diagnosis

Identifying a disease before it has progressed or caused major complications. Earlier diagnosis may provide more treatment choices, but its benefit depends on the disease and the accuracy and usefulness of available tests.

## Early Endosome

A small membrane-bound compartment inside a cell that receives material brought in from the cell surface. It sorts that material for return to the surface, delivery elsewhere in the cell, or breakdown.

---

### Early Gastric Cancer

Stomach cancer limited to the inner lining or the layer just beneath it, whether or not nearby lymph nodes are involved. It may be found early because of screening or testing for digestive symptoms.

#### Alternate Names:

EGC

### Early Infantile Epileptic Encephalopathy

A group of severe seizure disorders beginning in infancy that can disrupt brain development. Causes may be genetic, structural, metabolic, or unknown; frequent seizures and developmental concerns require specialist evaluation.

### Early Intervention

Support provided to infants and young children with developmental delays or disabilities. It may include physical, occupational, speech, behavioral, educational, and family services to improve development, function, and participation.

### Early Pregnancy Loss

Loss of a pregnancy in the first trimester, most often because of chromosomal abnormalities in the embryo. Evaluation and treatment depend on symptoms, ultrasound, bleeding, infection risk, pregnancy location, and the person's preferences.

### Early Puberty

Alternate name; see *Precocious Puberty*.

### Early Repolarization Pattern

An electrocardiogram pattern with a small rise or notch near the end of the QRS wave, often in the lower or side leads. It is often harmless, but its meaning depends on symptoms, family history, and the rest of the tracing.

#### Alternate Names:

Early Repolarization ECG Pattern; BER

### Early Repolarization Syndrome

A rare heart-rhythm condition associated with a characteristic ECG pattern and, in some people, an increased risk of dangerous ventricular arrhythmias.

Risk depends on the pattern, symptoms, and personal or family history.

#### Alternate Names:

ERS; Malignant Early Repolarization

### Early Satiety

Feeling full sooner than expected while eating, sometimes after only a small amount of food. It may result from stomach or digestive disease and should be assessed if persistent or associated with weight loss.

### Early-Stage Prostate Cancer

Prostate cancer that remains confined to the prostate or nearby tissues and has not spread to distant organs. The term includes cancers with different grades and risks of progression.

#### Alternate Names:

Early Prostatic Carcinoma; Localized Prostate Cancer

### Earwax

A waxy mixture made in the outer ear canal that lubricates and protects the skin and traps dust and microorganisms. It usually clears naturally, but buildup can block hearing or cause discomfort.

### Earwax Blockage

A buildup of wax that partly or completely blocks the ear canal. It may cause reduced hearing, fullness, ringing, itching, pain, or cough; safe removal may require a clinician.

### Earwax Removal

Removal of excess or impacted earwax by a clinician after examining the ear. Instruments, gentle suction, or irrigation may be used when safe; inserting objects at home can cause injury or push wax deeper.

### Eating Disorder

A group of mental-health conditions involving harmful eating patterns that cause distress, medical risk, or impaired functioning. Some involve intense concern about food, weight, or body shape, but



---

avoidant/restrictive food intake disorder (ARFID) does not require body-image concerns. Examples include anorexia nervosa, bulimia nervosa, binge-eating disorder, and ARFID.

## Eating Disorders

Alternate name; see *Eating Disorder*.

## Eaton-Lambert Syndrome

An uncommon condition in which the immune system interferes with nerve signals to muscles. It causes weakness, often in the hips and shoulders, reduced reflexes, dry mouth, and other automatic-body problems. It can be linked to small-cell lung cancer.

### Alternate Names:

Lambert-Eaton Myasthenic Syndrome; LEMS

## EB

Alternate name; see *Epidermolysis Bullosa*.

## EBCT

Electron-beam computed tomography, an older rapid CT method that was used especially to measure calcium in the coronary arteries. Newer CT technology is now more commonly used for heart and vessel imaging.

### Alternate Names:

Electron-Beam Computed Tomography; Ultrafast CT

## Ebola

Alternate name; see *Ebola virus disease*.

## Ebola Hemorrhagic Fever

Alternate name; see *Ebola Virus Disease*.

## Ebola Virus

A virus that causes Ebola disease. It spreads mainly through direct contact with infected blood or body fluids, usually after symptoms begin; illness may include fever, vomiting, diarrhea, bleeding, shock, and organ failure.

## Ebola Virus Disease

A severe, often fatal viral disease caused by Ebola viruses and affecting humans and other primates. After an incubation period of about 2–21 days, it may begin with fever, tiredness, muscle pain, headache, or sore throat and progress to vomiting, diarrhea, rash, bleeding, shock, or organ failure. It spreads mainly when blood or other body fluids from an infected person or animal, or contaminated objects such as clothing or bedding, enter another person through broken skin or the eyes, nose, or mouth. People generally cannot spread the disease before symptoms begin, but can spread it after becoming ill.

## Ebola Virus Infection

A severe viral infection caused by Ebola virus, characterized by fever and systemic illness that may progress to gastrointestinal symptoms, vascular leak, shock, bleeding, and multiorgan failure. Diagnosis and infection control require specialized care.

## Ebolavirus

A group of viruses that includes those causing Ebola disease. Infection can produce severe fever, vomiting, diarrhea, bleeding, shock, and organ damage and spreads through contact with infected body fluids.

## Ebstein Anomaly

A birth defect in which the tricuspid valve is positioned too low in the right ventricle and does not close normally. It can cause leakage of blood, enlargement of the right heart, blue coloring, heart failure, or abnormal rhythms.

## Ebstein's Anomaly

A birth defect in which the tricuspid valve is displaced downward and leaks. Effects range from no symptoms to blue coloring, heart failure, enlargement of the right heart, or abnormal heart rhythms.

## EBUS-TBNA

A procedure that uses a flexible breathing-tube camera and ultrasound to locate lymph nodes or masses beside the airways, then obtains tissue with a needle.

---

It helps diagnose and stage lung cancer, sarcoidosis, lymphoma, and tuberculosis.

**Alternate Names:**

Endobronchial Ultrasound Transbronchial Needle Aspiration; Convex Probe EBUS

**Ecbolics**

Medicines that stimulate contractions of the uterus. They may be used to start or strengthen labor or control heavy bleeding after birth; dosing requires medical supervision because excessive contractions can harm the parent or baby.

**Eccentric**

Positioned away from the center. In exercise, an eccentric muscle action occurs when a muscle produces force while lengthening, such as lowering a weight; it can improve strength but may cause soreness or injury.

**Eccentric Action**

Muscle contraction that produces force while the muscle lengthens, such as the controlled lowering of an object. It helps control movement and build strength but may cause muscle soreness when new or intense.

**Alternate Names:**

Eccentric Contraction

**Eccentrocyte Count**

The number or proportion of red blood cells whose hemoglobin is pushed toward one side of the cell. This smear finding may indicate oxidative injury or a hemolytic disorder but must be interpreted with other blood results.

**Ecchymosis**

A flat purple, blue, or brown skin mark caused by blood leaking into tissues, commonly after an injury. Easy or unexplained bruising can also result from medicines, low platelets, clotting disorders, fragile vessels, or illness.

**Eccrine Carcinoma**

A rare malignant tumor arising from eccrine sweat-

gland structures. It often presents as a slowly growing plaque or nodule, but may invade nearby tissue or recur; biopsy is required and treatment usually involves complete surgical removal.

**Eccrine Gland**

A small sweat gland found throughout the skin that releases watery sweat onto the surface. It helps cool the body and is especially numerous on the palms, soles, and forehead.

**Eccrine Glands**

Sweat glands distributed through the skin that release watery sweat directly onto the surface. They help control body temperature; disorders can cause excessive sweating, reduced sweating, blockage, or tumors.

**Eccrine Porocarcinoma**

A rare skin cancer arising from cells of sweat-gland ducts. It may appear as a slowly enlarging lump or patch and can recur or spread; biopsy and complete removal are often needed.

**Eccrine Poroma**

A usually harmless growth arising from a sweat-gland duct, often appearing as a flesh-colored, pink, or reddish lump on a hand or foot. It may bleed or be tender and can be removed if needed.

**ECG**

A recording of the heart's electrical activity made with skin electrodes. It can show heart rate, rhythm, conduction, reduced blood flow, prior heart damage, or other cardiac problems.

**Alternate Names:**

Electrocardiogram; EKG

**ECG/EKG Machine**

A device that records the heart's electrical activity through electrodes placed on the skin. A standard 12-lead recording helps assess rhythm, conduction, heart strain, reduced blood flow, and prior heart injury.

---

## Echinacea

A group of plants used in some herbal products for colds or claimed immune support. Evidence of benefit is inconsistent; allergic reactions and interactions with medicines can occur, especially in people with plant allergies.

## Echinocandins

A group of antifungal medicines that stop fungi building their protective cell wall. They are given mainly by injection for serious *Candida* infections and selected *Aspergillus* infections; liver reactions and infusion effects can occur.

## Echinococcal Cysts

Fluid-filled cysts caused by the larval stage of *Echinococcus* tapeworms, most often in the liver or lungs. They may remain silent or cause pressure, pain, breathing symptoms, or a severe allergic reaction if they rupture.

## Echinococcosis

A parasitic infection caused by larval *Echinococcus* tapeworms. Cystic disease usually forms cysts in the liver or lungs; alveolar disease grows invasively in the liver and may spread, requiring specialist treatment.

## Echinococcus

A group of tapeworms that can cause disease when people swallow eggs from contaminated soil, food, water, or contact with infected animals. Larvae can form cysts or invasive lesions, especially in the liver or lungs.

## Echinococcus Granulosus

A tapeworm whose larvae cause cystic echinococcosis. Dogs commonly carry the adult worm, and people become infected by swallowing eggs from contaminated material; cysts most often develop in the liver or lungs.

## Echinococcus Multilocularis

A tapeworm species whose larvae cause alveolar echinococcosis. The infection mainly affects the liver, where it grows slowly and invasively and may

spread to other organs.

## Echinococcus Vogeli

A tapeworm species that causes polycystic echinococcosis, a disease producing multiple parasitic cysts mainly in the liver and sometimes other abdominal organs.

## Echinocytes

Red blood cells with many small, evenly spaced projections around their edges. They may reflect kidney disease or an inherited blood disorder, but can also be created by improper storage or preparation of the blood sample.

### Alternate Names:

Burr Cells

## Echo

A common abbreviation for echocardiography, an ultrasound test that examines the heart's chambers, valves, pumping action, and blood flow. The full term is preferable in patient instructions.

## Echocardiogram

An ultrasound picture of the heart that shows its chambers, valves, pumping movement, and blood flow without ionizing radiation. It helps evaluate heart failure, valve disease, congenital defects, clots, and other heart problems.

## Echocardiography

Ultrasound imaging of the heart. It shows the chambers, valves, pumping action, and blood flow; the probe usually scans through the chest and may be passed into the esophagus for closer views.

## Echoencephalography

Ultrasound imaging of the brain, historically used mainly in infants to look for enlarged fluid spaces or displacement of brain structures. CT or MRI is now usually preferred for detailed brain imaging.

## Echogenic Bowel

An ultrasound finding in which a fetus's bowel appears unusually bright, sometimes as bright as bone.

---

It may be a normal variation or may be associated with infection, chromosome conditions, cystic fibrosis, bleeding, or other fetal problems.

### **Echogenic Intracardiac Focus**

A small bright spot, usually in a muscle inside the fetal heart, seen on ultrasound. When isolated and all other findings and screening are reassuring, it is commonly a harmless variation; its significance depends on the full evaluation.

### **Echography**

Medical ultrasonography, which uses high-frequency sound waves to create images of tissues, organs, blood flow, or a developing fetus.

### **Echolalia**

Repeating words or phrases spoken by someone else, either immediately or after a delay. It can occur in autism, developmental conditions, brain disorders, or some psychiatric illnesses and must be interpreted with other symptoms.

### **Echophenomena**

Automatic, non-purposeful imitation or repetition of actions, words, or sounds observed in others without explicit awareness or intent. Subtypes include echolalia (vocal repetition) and echopraxia (imitation of movements), commonly observed in Tourette syndrome, catatonia, and certain neurodevelopmental or neurodegenerative disorders.

### **Echopraxia**

Involuntary imitation or repetition of another person's movements. It may occur in some neurologic, developmental, or psychiatric conditions and is interpreted with other symptoms.

### **Echovirus**

A group of enteroviruses spread mainly by the fecal-oral route and sometimes by respiratory secretions. Infection is often mild or silent but can cause rash, respiratory illness, meningitis, encephalitis, heart inflammation, or severe newborn infection.

### **Eclampsia**

A new seizure during pregnancy or after delivery related to preeclampsia or pregnancy-related high blood pressure. It is an emergency requiring protection of breathing, seizure treatment, blood-pressure control, and urgent obstetric care.

### **Eclampsia in Pregnancy**

Alternate name; see *Eclampsia*.

### **ECMO**

Alternate name; see *Extracorporeal Membrane Oxygenation*.

### **ECOG Performance Status**

A scale from 0 to 5 describing how much a person's illness limits daily activity. It is used to estimate prognosis and decide whether someone may tolerate certain cancer treatments; 0 means fully active and 5 means death.

### **Econazole**

A topical antifungal that damages the fungal cell membrane and treats ringworm, athlete's foot, jock itch, and some yeast skin infections. It may cause local irritation, burning, itching, or allergic contact dermatitis.

### **Economy Class Syndrome**

Alternate informal name; see *Deep Vein Thrombosis*.

### **Ecstasy**

A street name for MDMA, an illegal stimulant and hallucinogenic drug. It can cause overheating, dehydration or dangerously low sodium, fast heartbeat, seizures, confusion, liver injury, and death.

### **Ectasia**

Abnormal widening of a blood vessel, duct, tube, or organ. Its importance depends on the structure involved; it may cause pressure, poor flow, bleeding, blockage, or rupture.

### **Ecthyma**

A bacterial skin infection that extends deeper than

---

a superficial sore and causes painful, punched-out ulcers covered by thick crusts. It commonly affects the legs and may need wound care and antibiotic treatment.

### **Ecthyma Gangrenosum**

A serious skin infection causing rapidly developing dark or black ulcers from tissue death, often due to *Pseudomonas* bloodstream infection. It occurs mainly in people with severely weakened immunity or very low white-cell counts and requires emergency treatment.

### **Ectoderm**

The outer layer of cells in an early embryo. It develops into the skin, hair, nails, nervous system, eyes, ears, and other sense-related structures.

### **Ectodermal Dysplasia**

A group of inherited disorders affecting tissues such as hair, teeth, nails, skin, and sweat glands. Features may include sparse hair, abnormal teeth, reduced sweating, overheating, dry skin, and repeated infections.

### **Ectodermal Dysplasias**

Inherited disorders affecting structures such as hair, teeth, nails, skin, and sweat glands. Depending on the type, they may cause abnormal teeth or hair, reduced sweating, overheating, dry skin, and repeated infections.

### **Ectodermosis Erosiva Pluriorificialis**

Historical name; see *Stevens-Johnson Syndrome*.

### **Ectomesenchymoma**

A rare cancer containing both nerve-related and connective-tissue parts, usually occurring in children. It may appear as a growing mass or cause pain; diagnosis requires tissue examination and specialist cancer treatment.

### **Ectoparasite**

A parasite that lives on the outer surface of a host, such as a louse, flea, tick, or mite. It may cause

itching, bites, skin disease, anemia, or transmission of infections.

### **Ectopia Cordis**

A rare birth defect in which the heart lies partly or completely outside the chest because the chest wall did not form normally. It is usually life-threatening and requires immediate specialist care.

### **Ectopia Lentis**

Displacement of the eye's natural lens from its normal position. Injury or inherited connective-tissue disorders may cause it, leading to blurred or double vision, unequal focusing, or increased eye pressure.

### **Ectopic**

Located or occurring in an abnormal position. For example, an ectopic pregnancy implants outside the main cavity of the uterus, most often in a fallopian tube.

### **Ectopic ACTH Syndrome**

Excess cortisol production caused by a tumor outside the pituitary making adrenocorticotrophic hormone (ACTH). It can cause weight gain, muscle weakness, high blood pressure, diabetes, low potassium, infections, and other features of cortisol excess.

### **Ectopic Beat**

An extra or early heartbeat that starts outside the heart's usual pacemaker. It may feel like a skipped beat, flutter, or thump and can occur in healthy people or with heart disease.

### **Ectopic Cushing Syndrome**

Excess cortisol caused by a tumor outside the pituitary producing ACTH or, rarely, a related hormone. It may cause high blood pressure, diabetes, low potassium, muscle weakness, infections, and rapid physical changes.

### **Ectopic Heartbeat**

A heartbeat that starts outside the heart's usual pacemaker. Early beats may feel like a skipped beat,

---

flutter, or sudden thump and can occur in healthy people or with structural heart disease.

### **Ectopic Heartbeats**

Extra or early beats arising outside the heart's usual pacemaker. They may feel like a skipped beat, thump, or flutter and are often harmless, but frequent episodes, fainting, chest pain, or breathlessness need assessment.

### **Ectopic Kidney**

A kidney located somewhere other than its usual position in the upper back of the abdomen. It may work normally or be associated with abnormal drainage, urine reflux, stones, or infection.

### **Ectopic Pacemaker**

An excitable cardiac focus or group of pacemaker cells located outside the sinoatrial node (in the atria, atrioventricular junction, or ventricles) that prematurely initiates cardiac depolarizations and rhythm, frequently producing ectopic beats or tachycardia.

#### **Alternate Names:**

Ectopic Focus; Latent Pacemaker

### **Ectopic Pregnancy**

A pregnancy implanted outside the main cavity of the uterus, most often in a fallopian tube. It may cause missed periods, abdominal pain, or vaginal bleeding; rupture can cause life-threatening internal bleeding and requires emergency care.

#### **Alternate Names:**

Tubal Pregnancy; Extrauterine Gestation

### **Ectopic Thyroid**

Thyroid tissue located outside its usual position in the lower front of the neck, most often at the base of the tongue. It may be the only functioning thyroid tissue and can cause swallowing, voice, or breathing problems.

### **Ectopic Thyroid Tissue**

Thyroid tissue that developed outside the usual thyroid location because it did not move normally during development. It is often at the tongue base

and may be the only working thyroid tissue, sometimes causing swallowing or airway symptoms.

### **Ectopic Tissue**

Tissue located in an unusual position, usually because it developed there, failed to move normally during growth, or became implanted elsewhere. Effects depend on the tissue and whether it causes pressure, bleeding, blockage, or abnormal function.

### **Ectopic Ureter**

A ureter that does not enter the bladder in its normal position. It may drain into the urethra, vagina, or another abnormal site and can cause urine leakage, infection, blockage, or kidney damage.

### **Ectopion**

Outward turning or eversion of a body structure, most often the lower eyelid or the cervix. The effect depends on the site and may include irritation, exposure, bleeding, or abnormal discharge.

### **Ectrodactyly**

A birth difference in which one or more central fingers or toes are missing, leaving a deep cleft in the hand or foot. It may occur alone or with other limb or genetic differences and can affect function.

### **Ectromelia**

A birth defect in which one or more limbs are absent or severely underdeveloped. In laboratory medicine, the same word names a mouse poxvirus infection; these are separate uses and should not be confused.

### **Ectropion**

Outward turning of an eyelid, usually the lower lid, so the eye may not close or drain tears normally. It can cause tearing, dryness, irritation, and corneal injury; aging, weakness, scarring, or paralysis may contribute.

### **Ectropion (Drooping Eyelids)**

Alternate name; see *Ectropion*.



---

## Eculizumab

An antibody medicine that blocks part of the immune system called complement. It treats selected blood, kidney, and nerve disorders but greatly increases the risk of meningococcal and other serious infections, so vaccination and monitoring are essential.

## Eczema

A group of skin conditions that cause itchy, dry, inflamed, red, scaly, cracked, or sometimes oozing skin. The skin barrier is weakened, so the skin loses moisture and becomes easier to irritate. Atopic dermatitis is the most common type. Symptoms may worsen because of irritants, dry weather, heat, sweating, stress, or skin infection. Eczema is not contagious.

## Eczema Herpeticum

A rapidly spreading herpes simplex infection in skin affected by eczema or another barrier disorder. It causes many similar painful blisters or punched-out sores, often with fever, and needs urgent antiviral treatment because the eyes and other organs may be affected.

## Eczematous Dermatitis

A pattern of inflammatory skin disease characterized by itching, redness, scaling, dryness, oozing, or vesicles. Causes include atopic, allergic contact, irritant contact, and other forms of dermatitis.

## ED

Erectile dysfunction, difficulty getting or keeping an erection sufficient for sexual activity. It may reflect blood-vessel, nerve, hormone, medicine, psychological, or relationship factors and can signal broader health problems.

### Alternate Names:

Erectile Dysfunction; Impotence

## Edema

Swelling caused by extra fluid in body tissues. It may leave a dent when pressed or feel firm. Common causes include heart, kidney, or liver disease, poor vein drainage, blocked lymph flow, and some

medicines.

## Edema Trunk

Abnormal accumulation of fluid in the soft tissues of the chest or abdomen, producing swelling from heart, kidney, liver, lymphatic, venous, inflammatory, or medication-related causes.

## Edentulism

Complete or partial absence of natural teeth. It can impair mastication, speech, nutrition, facial support, and quality of life and may be managed with removable, fixed, or implant-supported prostheses.

## Edentulous

Having some or all natural teeth missing. Tooth loss may result from decay, gum disease, injury, extraction, or a developmental condition and can affect chewing, speech, nutrition, jawbone support, and appearance; dentures or implants may restore function.

## Edentulousness

Complete or partial absence of natural teeth. It may result from dental disease, trauma, developmental abnormalities, or tooth extraction and can impair chewing, speech, and oral function.

## Edetic Acid

Another name for EDTA, a chemical that binds certain metals. It is used in blood-collection tubes, laboratories, and selected treatments for heavy-metal poisoning.

## EDMD

Abbreviation; see *Emery-Dreifuss Muscular Dystrophy*.

## EDTA

A chemical that binds metals. It is used in some blood-collection tubes and, in selected cases, as a supervised treatment for heavy-metal poisoning; unnecessary use can harm the kidneys and lower essential minerals.

---

## Edward's Syndrome

Alternate spelling; see *Edwards Syndrome*.

### Alternate Names:

Trisomy 18; Trisomy E

## Edwards Syndrome

A chromosome condition caused by an extra chromosome 18 in some or all cells. It commonly causes poor growth, developmental impairment, feeding difficulty, heart defects, and other birth abnormalities; medical needs and survival vary widely.

## Edwardsiella

A group of bacteria associated with water and animals. Some species, especially *E. tarda*, can rarely cause diarrhea, wound infection, bloodstream infection, or other illness in people.

## Edwardsiella Tarda

A bacterium found in fish, reptiles, and water that can rarely infect people. It may cause diarrhea, wound infection, gallbladder or abdominal infection, or bloodstream infection.

## EEG

A test that records the brain's electrical activity through sensors placed on the scalp. It helps evaluate seizures, unexplained episodes, altered awareness, certain sleep disorders, and some brain illnesses.

### Alternate Names:

Electroencephalogram

## EFAST Examination

A rapid bedside ultrasound used after serious injury to look for internal bleeding, fluid around the heart, and air or blood around the lungs. A normal scan does not exclude every internal injury, so repeat assessment or CT may be needed.

## Efavirenz

An antiretroviral medicine that blocks HIV reverse transcriptase and is used in combination treatment for HIV infection. It can cause vivid dreams, dizziness, mood or nervous-system effects, rash, and

important medicine interactions.

## Effacement

Thinning and shortening of the cervix during labor, measured as a percentage from 0 to 100. Effacement helps the cervix open and usually occurs along with dilation as contractions progress.

## Effective Dose

An estimate that combines radiation exposure to different organs into one number. It helps compare the approximate risk of imaging procedures but cannot predict an individual person's exact risk.

## Effective Half-Life

The time needed for the amount of a radioactive substance in the body or a tissue to fall by half through both natural decay and removal by the body. It helps estimate how long radiation exposure lasts.

## Effective Reproduction Number

The average number of people one infected person infects at a particular time, taking account of immunity, behavior, and control measures. Continued spread generally occurs when the number is above 1.

### Alternate Names:

R<sub>t</sub>

## Effectiveness

How well a treatment, test, or public-health intervention achieves its intended result in usual real-world practice. It differs from efficacy, which describes performance under ideal, controlled study conditions.

## Effector

A cell, molecule, organ, or mechanism that carries out a response after receiving a signal. For example, an immune cell may destroy an infected cell after activation.

## Efferent

Carrying signals away from a central structure. In

---

the nervous system, efferent pathways carry movement or automatic-control signals from the brain or spinal cord to muscles, glands, or organs.

### **Efferent Arteriole**

The small blood vessel that carries blood away from the kidney's filtering unit. It helps control pressure in the filter and leads to the small vessels around the kidney tubules.

### **Efferent Ductules**

A series of 10 to 20 delicate, ciliated convoluted tubules that penetrate the tunica albuginea of the testis to transport spermatozoa from the rete testis into the head (caput) of the epididymis, reabsorbing the majority of testicular luminal fluid.

#### **Alternate Names:**

Ductuli Efferentes

### **Efferent Lymphatic Vessel**

A lymphatic vessel that carries lymph out of a lymph node.

### **Efferent Nerve**

A nerve pathway carrying signals away from the brain or spinal cord to muscles, glands, or organs. Damage can cause weakness, impaired automatic function, or loss of control in the affected area.

### **Efferent Neuron**

A nerve cell that carries movement or automatic-control signals from the brain or spinal cord to muscles, glands, or organs. Damage can cause weakness or impaired control of body functions.

### **Efferent Pathway**

A nerve route carrying signals away from the brain or spinal cord to muscles, glands, or organs. It controls movement and automatic functions such as heart rate, digestion, sweating, and bladder activity.

### **Effervescent Tablets**

Tablets dissolved in water before use, releasing bubbles as they form a drinkable solution. Some contain substantial sodium or other ingredients, so peo-

ple with heart, kidney, or blood-pressure problems should check the label.

### **Efficacy**

How well a treatment produces its intended result under ideal, carefully controlled study conditions. It differs from effectiveness, which describes how well the treatment works in ordinary clinical practice.

### **Effort Angina**

A classic presentation of stable ischemic heart disease characterized by substernal chest discomfort, tightness, or pressure predictably provoked by physical exertion, emotional stress, or cold exposure, and consistently relieved within minutes by rest or sublingual nitroglycerin.

#### **Alternate Names:**

Classic Angina; Exertional Angina

### **Effort Syncope**

Transient loss of consciousness occurring abruptly during or immediately following physical exertion, serving as a cardinal red-flag symptom of severe cardiac outflow obstruction (such as critical aortic stenosis or hypertrophic obstructive cardiomyopathy) or exertion-induced ventricular arrhythmia.

### **Effusion**

Abnormal buildup of fluid in a body space, such as around a joint, lung, or heart. It may cause swelling, pain, stiffness, or breathing difficulty; the cause may be injury, infection, inflammation, heart failure, or cancer.

### **Effusive-Constrictive Pericarditis**

A form of pericardial disease in which fluid around the heart coexists with constrictive physiology. Signs that the heart remains restricted after the fluid is drained support the diagnosis.

### **EGD**

An examination of the esophagus, stomach, and duodenum using a flexible camera passed through the mouth. It can investigate pain, swallowing problems, bleeding, or anemia and can obtain samples or treat selected lesions.

---

## **EGFR**

A protein on the cell surface that receives growth signals. Changes that overactivate EGFR can help some cancers grow; testing the tumor may guide use of treatments that block this pathway.

## **EGFR Inhibition**

Cancer treatment that blocks growth signals sent through the epidermal growth factor receptor. It may slow selected colorectal, lung, or head-and-neck cancers when the tumor has suitable features; rash, diarrhea, and other effects may occur.

## **Egg**

A nutrient-rich food containing protein, fat, vitamins, and minerals, but also a common allergen. Raw or undercooked eggs can transmit Salmonella; thorough cooking and individual allergy or dietary needs guide safe use.

## **Egg Allergy**

An immune reaction to egg proteins that may cause hives, swelling, vomiting, stomach pain, wheezing, or anaphylaxis. It is more common in children; reactions vary in severity and require an individualized avoidance and emergency plan.

### **Alternate Names:**

Egg Hypersensitivity; Hen's Egg Allergy

## **Egg Shell**

The hard calcium-containing outer covering of an egg. In medical imaging, "eggshell calcification" describes a thin rim of calcium around a lesion or lymph node.

## **Egg-Cell**

A mature female reproductive cell, also called an oocyte, that can unite with sperm during fertilization. It contains half the usual genetic material and can develop into an embryo after fertilization.

## **Eggerthella**

A group of bacteria that can live in the human intestine without causing disease. Rarely, especially in people with serious illness, they can cause blood-

stream infection, abscesses, or other opportunistic infections.

## **Eggshell Nail**

A nail that becomes unusually thin, soft, flexible, and sometimes pale or concave. Possible causes include nutritional deficiency, chronic illness, aging, or repeated trauma; persistent changes should be medically assessed.

## **Egophony**

A change in voice sounds heard through a stethoscope in which a spoken "E" sounds like an "A." It may occur when lung tissue becomes dense or filled with fluid, such as in pneumonia.

## **EHBA**

Abbreviation for extrahepatic biliary atresia, a serious infant disorder in which the bile ducts outside the liver are absent, scarred, or blocked, causing persistent jaundice and progressive liver injury.

## **Ehlers-Danlos Syndrome**

A group of inherited connective-tissue disorders that can cause unusually flexible joints, stretchy or fragile skin, easy bruising, and poor wound healing. Some types also weaken blood vessels or internal organs and require specialist care.

### **Alternate Names:**

EDS; Hypermobility Syndrome

## **EHR**

Electronic health record: a digital collection of a person's health information, such as diagnoses, medicines, allergies, test results, and clinical notes, that can be shared among authorized healthcare professionals.

## **Ehrlichia**

A group of bacteria spread mainly by ticks that infect certain white blood cells and cause ehrlichiosis. Infection may produce fever, headache, muscle aches, low blood counts, and serious illness if untreated.

---

## Ehrlichia And Anaplasma

Tick-borne bacteria that infect white blood cells. They can cause fever, headache, muscle aches, weakness, low blood counts, and abnormal liver tests; severe disease is more likely in older or immunocompromised people.

## Ehrlichiosis

A tick-borne infection caused by Ehrlichia bacteria. It commonly causes fever, headache, muscle aches, weakness, low white-cell or platelet counts, and raised liver enzymes; untreated infection can damage the brain, lungs, or other organs.

## Ehrlichiosis And Anaplasmosis

Tick-borne bacterial infections that may cause fever, headache, muscle aches, weakness, low blood counts, and abnormal liver tests. Prompt medical assessment and appropriate antibiotics reduce the risk of severe complications.

## Eicosanoid

A signaling substance made from certain fats that helps regulate inflammation, blood-vessel width, platelet activity, pain, and muscle contraction. Prostaglandins, leukotrienes, and thromboxanes are examples.

## Eicosanoid Degradation

Breakdown and inactivation of eicosanoids, the body's short-lived fatty-acid signals. This limits how long and how strongly they affect inflammation, blood vessels, platelets, airways, and other muscles.

## Eicosanoid Metabolism

The body's production, modification, and breakdown of prostaglandins, leukotrienes, thromboxanes, and related fatty-acid signals that influence inflammation, blood vessels, platelets, pain, and airway function.

## Eicosanoid Receptor

A protein on or inside a cell that responds to prostaglandins, leukotrienes, thromboxanes, or related lipid signals. Activation can change inflam-

mation, blood-vessel tone, platelet activity, pain, or smooth-muscle contraction.

## Eicosapentaenoic Acid

The omega-3 fat EPA, found in oily fish and some supplements. It forms part of cell membranes and can be changed into signaling substances involved in inflammation and blood clotting.

## Eicosatetraenoic Acid

A 20-carbon fatty acid with four double bonds. Arachidonic acid is the best-known form and is used by cells to make prostaglandins, leukotrienes, thromboxanes, and other signaling substances.

## Eighth Cranial Nerve

The vestibulocochlear nerve, which carries hearing and balance signals from the inner ear to the brain. Damage can cause hearing loss, ringing, dizziness, imbalance, or abnormal eye movements.

### Alternate Names:

Vestibulocochlear Nerve; Auditory Nerve

## Eikenella

A group of bacteria found in the mouth. Some species can cause deep wound, bone, joint, or bloodstream infection, especially after human bites or injuries involving a person's teeth.

## Eikenella Corrodens

A slow-growing bacterium normally found in the mouth that can cause infection after human bites or clenched-fist injuries. It may produce abscesses, bone or joint infection, or rarely bloodstream infection.

## Eisenmenger Complex

Alternate form; see *Eisenmenger syndrome*.

## Eisenmenger syndrome

A form of advanced pulmonary vascular disease caused by a longstanding congenital heart defect that initially produces excessive blood flow to the lungs. Progressive elevation of pulmonary vascular resistance leads to right-to-left or bidirectional

---

shunting, reduced blood oxygen, and cyanosis.

### Eisenmenger's Syndrome

Alternate form; see *Eisenmenger syndrome*.

### Ejaculation

The release of semen through the penis, usually during sexual climax. It involves coordinated contractions of reproductive ducts, glands, pelvic muscles, and nerves; pain, blood, or major change in ejaculation needs assessment.

### Ejaculatory Duct

One of two small tubes formed where the vas deferens joins a seminal-vesicle duct. Each passes through the prostate and opens into the urethra, carrying semen during ejaculation.

### Ejaculatory Ducts

Two small tubes within the prostate that carry semen from the vas deferens and seminal vesicles into the urethra. Blockage or inflammation can contribute to painful ejaculation, infertility, or reduced semen volume.

### Ejaculatory Dysfunction

Difficulty ejaculating, such as delayed ejaculation, absent ejaculation, painful ejaculation, or ejaculation into the bladder instead of out through the penis. Causes include medicines, nerve or pelvic disorders, diabetes, prostate procedures, and psychological factors.

### Ejection Click

A sharp heart sound heard just after the heart begins to contract, caused by opening of an abnormal aortic or pulmonary valve. It may occur with a bicuspid valve or valve narrowing and is assessed with other findings.

### Ejection Fraction

The percentage of blood pumped out of a heart chamber with each beat. It is used to describe the heart's pumping strength, but a normal value does not rule out heart failure.

### Ejection Murmur

A midsystolic crescendo-decrescendo heart murmur that begins shortly after the first heart sound (S1), peaks in midsystole, and ends before the second heart sound (S2), produced by turbulent blood flow across an obstructed or narrowed ventricular outflow tract or semilunar valve, as seen in aortic stenosis and pulmonic stenosis.

#### Alternate Names:

Systolic Ejection Murmur; Midsystolic Murmur

### EKG

A recording of the heart's electrical activity made through electrodes on the skin. It can show heart rate, rhythm, conduction problems, reduced blood flow, prior heart injury, or signs of heart enlargement.

#### Alternate Names:

Electrocardiogram; ECG

### Ekman-Lobstein Disease

Alternate name; see *Osteogenesis Imperfecta*.

### Elastance

A physical and physiological measure of the stiffness and tendency of a hollow organ or vascular structure to recoil toward its original resting dimensions after being distended by pressure, defined mathematically as the change in pressure divided by the change in volume (the inverse of compliance).

### Elastic Cartilage

Flexible supporting tissue containing many elastic fibers. It helps structures such as the outer ear, epiglottis, and pressure-equalizing ear tube keep their shape while bending.

### Elastic Fiber

A thin connective-tissue fiber that allows skin, lungs, blood vessels, and other structures to stretch and recoil. Damage or loss of elastic fibers can reduce flexibility or weaken tissues.

### Elastic Skin

Alternate name; see *Ehlers-Danlos Syndrome*.



---

## Elastic Tissue

Connective tissue containing fibers that let structures stretch and return toward their original shape. It helps arteries pulse, lungs expand, skin recoil, and some ligaments support movement.

## Elasticity

The ability of a tissue or material to return toward its original shape after stretching or compression. It contributes to the function of skin, lungs, blood vessels, muscles, and ligaments.

## Elastics

Elastic bands used in medical or dental care, such as orthodontic bands, compression supports, wound devices, or surgical ties. Their purpose and safe use depend on the specific treatment.

## Elastin

A flexible protein in connective tissue that lets skin, lungs, blood vessels, and other structures stretch and recoil. Aging, inflammation, and chronic sun exposure can damage elastin.

## Elastofibroma

A harmless, slowly growing mass made of fatty and elastic tissue, most often beneath the lower shoulder blade in older adults. It may cause swelling or movement-related discomfort but often needs no treatment.

## Elastography

An imaging method that estimates how stiff tissue is. It is commonly used to assess liver scarring and may help evaluate breast, thyroid, prostate, or other abnormal areas without tissue removal.

## Elastomer

A flexible rubber-like material that can stretch and return toward its original shape. It is used in some medical tubing, seals, gloves, devices, and implants; biocompatibility depends on the material.

## Elastosis

Abnormal buildup, breakdown, or alteration of elas-

tic fibers in tissue. It is commonly associated with long-term sun exposure and can occur in some degenerative, inherited, or lung disorders.

## Elastosis Perforans Serpiginosa

A rare perforating dermatosis characterized by transepidermal elimination of abnormal elastic fibers, presenting clinically as small, keratotic, umbilicated papules arranged in a serpiginous or annular configuration on the neck, face, or upper extremities, often associated with systemic disorders such as Down syndrome, Ehlers-Danlos syndrome, and pseudoxanthoma elasticum.

### Alternate Names:

EPS; Lutz Disease

## Elbow

The joint connecting the upper arm and forearm, formed by the humerus, radius, and ulna. It allows bending and straightening and works with nearby joints to rotate the forearm.

## Elbow (Radial Head or Neck) Fracture

A break in the radial head or neck, the upper end of the radius at the elbow, usually after a fall onto the hand or a direct blow. It can cause pain, swelling, tenderness, and difficulty bending, straightening, or rotating the forearm. Assessment may include X-rays; treatment depends on the fracture and any associated injury, and stable fractures often benefit from early, gentle movement to reduce stiffness.

### Alternate Names:

Radial Head or Neck Fracture; Radial Head/Neck Fracture; Radial Neck Fracture

## Elbow Dislocation

Displacement of the radius and ulna from their normal alignment with the humerus, usually after a fall or forceful injury. It causes severe pain, deformity, and loss of motion and requires prompt neurovascular assessment.

## Elbow Extensor

A muscle that straightens the elbow, mainly the triceps and anconeus. These muscles help push, reach, lift, throw, and control lowering of the arm.

---

## Elbow Flexors

Muscles that bend the elbow, mainly the biceps, brachialis, and brachioradialis. They help lift the forearm and turn the palm upward; injury, pain, nerve problems, or tendon damage can weaken movement.

## Elbow Joint

A joint complex between the humerus, radius, and ulna that allows the arm to bend and straighten and the forearm to rotate. Cartilage, ligaments, muscles, and fluid-filled coverings provide stability and movement.

## Elbow Pain

Pain in or around the elbow caused by tendon injury, arthritis, bursitis, nerve pressure, fracture, instability, infection, or referred pain from the neck or shoulder. Swelling, deformity, weakness, or injury needs assessment.

### Alternate Names:

Elbow Arthralgia; Cubital Pain

## Elbow Replacement

Surgery that removes damaged parts of the elbow and replaces them with artificial components to relieve pain and improve movement. It may be used for severe arthritis or injury; infection, loosening, and nerve or bone problems can occur.

## Elder Abuse

An intentional act or failure to provide care that harms an older person. It includes physical, emotional, sexual, or financial abuse and neglect; warning signs require sensitive assessment and safeguarding according to local law.

## Elective

Planned in advance rather than performed immediately for an emergency. An elective procedure may still be medically necessary; its timing follows assessment of benefits, risks, alternatives, and the person's preferences.

## Elective Abortion

A planned ending of a pregnancy using medicines or a procedure, usually before an immediate medical emergency develops. The method depends on pregnancy duration, health, personal choice, and local clinical and legal requirements.

## Elective Surgery

Surgery scheduled in advance because it is not needed immediately to prevent serious harm. It may still be important or medically necessary, and timing follows discussion of benefits, risks, alternatives, and preferences.

## Electric Axis of the Heart

The net average direction and magnitude of the electrical electromotive forces traveling through the ventricular myocardium during ventricular depolarization, represented on the frontal plane electrocardiogram and normally oriented between -30 degrees and +90 degrees.

### Alternate Names:

Mean QRS Axis

## Electric Impedance

The opposition a material or tissue offers to alternating electrical current, measured in ohms. Medical devices use impedance changes to assess body composition, monitor signals, or detect contact and function.

## Electric Shock

Injury caused when electrical current passes through the body. It can cause burns, muscle damage, falls, abnormal heart rhythms, breathing failure, nerve injury, or internal damage even when skin burns are small.

## Electrical Alternans

A beat-to-beat change in the size or direction of ECG complexes. It can occur when the heart moves within a large fluid collection around it and may indicate cardiac tamponade, an emergency.

---

## Electrical Cardioversion

A controlled electrical shock delivered at a specific point in the heartbeat to restore a normal rhythm, often for atrial fibrillation or flutter. Sedation, clot risk, heart stability, and follow-up treatment must be considered.

## Electrical Conductance

The ability of a material or tissue to carry electrical current. It depends on how easily charged particles move and is important in nerve, muscle, heart, and medical-device measurements.

## Electrical Injury

Damage caused by electrical current passing through the body. Severity depends on voltage, current, duration, pathway, and skin resistance; effects may include burns, heart-rhythm problems, muscle breakdown, nerve injury, falls, or breathing failure.

## Electrical Stimulation

Use of controlled electrical currents to activate nerves or muscles. It may help selected pain, weakness, movement, or rehabilitation problems; benefits and risks depend on the device, body site, and underlying condition.

## Electrical Synapse

A direct connection between cells through tiny channels that let charged particles and electrical signals pass rapidly from one cell to another. It helps coordinate some brain, heart, and other tissue activity.

## Electrocardiogram

A recording of the heart's electrical signals from skin electrodes. It can help identify abnormal rhythms, conduction problems, reduced blood flow, heart attack, chamber strain, and some effects of medicines or electrolyte changes.

**Alternate Names:**

ECG; EKG

## Electrocardiographic Gating

An imaging synchronization technique used in cardiac computed tomography, magnetic resonance

imaging, and nuclear cardiology that synchronizes image acquisition with specific phases of the patient's cardiac cycle (prospective or retrospective gating) to eliminate cardiac motion artifacts.

**Alternate Names:**

ECG Gating

## Electrocardiography

The process of recording the heart's electrical activity through skin electrodes, usually as a 12-lead ECG. It assesses rate, rhythm, conduction, chamber strain, reduced blood flow, prior injury, and some medicine or electrolyte effects.

## Electrocauterization

A procedure using controlled electrical energy to heat tissue, destroy selected tissue, or seal bleeding vessels. It can cause burns, scarring, smoke exposure, or injury to nearby tissue and requires appropriate precautions.

## Electrocautery

Use of heat produced by electrical current to cut tissue, destroy unwanted tissue, or stop bleeding during a procedure. Possible complications include burns, scarring, smoke inhalation, and damage to nearby structures.

## Electrocoagulation

A surgical technique using electrical energy to heat tissue, seal blood vessels, or destroy selected tissue. It is used to control bleeding and may cause burns, scarring, or unintended injury.

**Alternate Names:**

Surgical Diathermy; Electrocautery

## Electroconvulsive Therapy

A treatment that causes a brief controlled seizure while the person is under general anesthesia. It can rapidly treat severe depression, catatonia, or mania; temporary confusion and memory problems are common risks.

## Electrocution

Serious injury or death caused by electrical current

---

passing through the body. It may produce burns, abnormal heart rhythms, muscle or nerve damage, breathing failure, falls, or hidden internal injury and requires emergency evaluation.

### **Electrode**

A conductive contact that carries electrical signals or current to or from the body, a device, or a laboratory sample. Electrodes are used in tests such as ECG and EEG and in selected treatments.

### **Electrodesiccation**

A procedure using electrical energy to dry and destroy selected superficial tissue or control bleeding. It may treat some skin lesions, often with scraping, and can cause pain, scarring, infection, or pigment change.

### **Electroencephalogram**

A recording of brain electrical activity through scalp electrodes. It helps evaluate seizures, prolonged altered awareness, encephalopathy, selected sleep disorders, and some intensive-care conditions; a normal result does not exclude epilepsy.

### **Electroencephalography**

The process of recording brain electrical activity with electrodes on the scalp. It can help identify abnormal patterns associated with seizures, sleep disorders, or diffuse brain dysfunction.

### **Electrogastrogram**

A recording of the stomach's electrical activity, usually through electrodes on the abdominal skin. It measures stomach rhythm and is used mainly in specialist assessment of suspected motility disorders.

### **Electrolysis**

Hair removal using electrical energy delivered through a fine probe into individual hair follicles. Repeated treatment may permanently reduce hair growth, but pain, irritation, pigment changes, scarring, or infection can occur.

### **Electrolyte**

A mineral that carries an electrical charge when dissolved in body fluids, such as sodium, potassium, calcium, magnesium, chloride, or phosphate. Electrolytes help control nerves, muscles, heart rhythm, fluid balance, and acid-base balance.

### **Electrolyte Balance**

Maintenance of appropriate levels of minerals such as sodium, potassium, calcium, magnesium, chloride, and phosphate in body fluids. The kidneys, hormones, intestines, and lungs regulate them; imbalance can affect nerves, muscles, heart rhythm, and hydration.

### **Electrolyte Imbalance**

An abnormal level of minerals such as sodium, potassium, calcium, magnesium, chloride, or phosphate in the blood. It may cause weakness, cramps, confusion, seizures, or abnormal heart rhythms; causes include fluid loss, kidney disease, medicines, and hormone disorders.

### **Electrolyte Panel**

A blood test measuring selected electrolytes, commonly sodium, potassium, chloride, and bicarbonate, and sometimes related minerals. It helps assess fluid balance, kidney function, acid-base disorders, and causes of weakness or confusion.

### **Electrolytes**

Electrically charged minerals in body fluids, including sodium, potassium, chloride, bicarbonate, calcium, magnesium, and phosphate. They support nerve conduction, muscle contraction, fluid balance, and acid-base regulation.

### **Electromagnetic Navigation Bronchoscopy**

A computer-guided form of bronchoscopy that helps a flexible camera reach small areas near the outer lung. It can guide a biopsy of a lesion not visible with ordinary bronchoscopy, but bleeding, infection, or a collapsed lung can occur.

---

### **Electromechanical Dissociation**

A cardiac arrest in which the monitor shows electrical activity but the heart does not pump enough blood to produce a pulse. It requires immediate CPR and treatment of the underlying cause.

#### **Alternate Names:**

EMD; Pulseless Electrical Activity; PEA

### **Electromyography**

A test that records the electrical activity of skeletal muscles, often together with nerve-conduction studies, to evaluate peripheral nerve, nerve-root, neuromuscular-junction, and muscle disorders.

### **Electron Microscope**

A microscope that uses electrons rather than visible light to produce highly magnified images of cells, tissues, microorganisms, or materials.

### **Electron Transport Chain**

A series of protein systems inside mitochondria that transfer energy from nutrients to oxygen and use it to make ATP, the cell's main energy source. Problems in this process can impair organs that need substantial energy, such as the brain and muscles.

### **Electron-Beam Computed Tomography**

A rapid CT technique that uses an electronically steered X-ray beam to image the heart and measure calcium deposits in coronary arteries. It has largely been replaced by newer CT methods.

#### **Alternate Names:**

EBCT

### **Electronic Cigarette**

A battery-powered device that heats liquid into an inhaled aerosol, often containing nicotine and other chemicals. It can cause dependence, airway irritation, lung injury, or poisoning from liquid nicotine and is not harmless.

### **Electronic Fetal Monitoring**

Continuous or intermittent recording of the fetal heart rate and uterine contractions during pregnancy or labor. It helps identify patterns suggesting

that the baby may not be receiving enough oxygen, but results require clinical interpretation.

### **Electronystagmography**

A test that records eye movements with sensors around the eyes, often while the balance system is stimulated. It helps evaluate involuntary eye movements and inner-ear or balance-nerve disorders; video testing is now often preferred.

### **Electrophoresis**

A laboratory method that separates charged molecules by how they move through a gel or liquid in an electric field. It can analyze proteins, hemoglobin, DNA, and other substances.

### **Electrophysiologic Testing**

Testing that records or deliberately stimulates the heart's electrical system, usually with catheters, to investigate abnormal rhythms and guide treatment. It is invasive and carries risks such as bleeding, infection, or rhythm disturbance.

### **Electrophysiology**

The study of electrical activity in living cells and tissues. Clinical electrophysiology includes tests and treatments involving heart rhythm, nerves, muscles, and the connections between them.

### **Electrophysiology Study**

An invasive test in which thin electrode catheters are placed in the heart to map electrical pathways and trigger or measure abnormal rhythms. It can guide ablation or other treatment; bleeding, infection, and rhythm complications are possible.

### **Electroretinography**

A test that records the retina's electrical response to flashes or patterned light. It helps evaluate inherited or acquired retinal disorders when examination and imaging do not fully explain visual loss.

### **Electrosurgery**

Use of electrical energy during a procedure to cut tissue, destroy tissue, or control bleeding. Risks in-

---

clude burns, scarring, smoke exposure, unintended tissue injury, and interference with implanted electronic devices.

### **Electrosurgical Unit**

A generator that delivers high-frequency electrical energy through an electrode to cut tissue or stop bleeding. Safe use requires attention to current pathways, burns, nearby structures, and possible interference with implanted electronic devices.

### **Elejalde Syndrome**

A very rare autosomal-recessive neurocutaneous disorder characterized by silvery or gray hair, abnormal melanin distribution, and severe neurologic disease such as seizures, hypotonia, and developmental impairment. It requires specialist genetic and neurologic evaluation.

### **Element**

A pure substance made of atoms with the same number of protons. Some elements, such as iron and iodine, are essential nutrients; others can be toxic, and health effects depend on dose and exposure.

### **Elephant Ear Poisoning**

Irritation or poisoning after eating or handling elephant-ear or taro plants, which contain sharp calcium-oxalate crystals. It can cause intense burning and swelling of the mouth and throat and may obstruct breathing.

### **Elephantiasis**

Severe, long-lasting swelling and thickening of skin, usually in a limb or the scrotum, caused by blocked lymph drainage. Lymphatic filariasis is one cause; other causes include repeated infection, surgery, cancer, or injury.

### **Elevated Blood Pressure**

Blood pressure readings that are repeatedly above the healthy range. Persistent elevation increases the risk of heart disease, stroke, kidney disease, and blood-vessel problems, although a single high reading does not establish a diagnosis.

### **Elevated Hemidiaphragm**

An imaging finding in which one side of the diaphragm sits higher than usual. Causes include reduced lung volume, nerve or muscle weakness, abdominal disease, or a normal variation; significance depends on symptoms and other findings.

### **Elevated Liver Enzymes**

Higher-than-expected blood levels of enzymes released by stressed or injured liver cells. Causes include fatty liver, medicines, alcohol, viral hepatitis, blocked bile flow, and muscle injury; the pattern and degree guide further evaluation.

#### **Alternate Names:**

Transaminitis; Elevated Aminotransferases

### **Elevation**

Upward movement or position of a body part, such as raising the arm, shoulder, jaw, or foot. The term can also describe raising an injured limb to reduce swelling.

### **Elevator**

A surgical or dental instrument used to lift, separate, or loosen tissue, bone, or a tooth from nearby structures. The exact design and use depend on the procedure.

### **Eligible**

Meeting the requirements for a treatment, clinical trial, screening program, benefit, or healthcare service. Eligibility does not guarantee that an intervention is appropriate or that it will be available.

### **Elimination**

Removal of waste products or medicines from the body, mainly through the kidneys, liver, lungs, skin, or intestines. Reduced elimination can make a medicine remain longer in the body and increase its effects or toxicity.

### **Elimination Diets**

Short-term eating plans that remove suspected foods and later reintroduce them in a planned way to identify food-related symptoms. They should be su-



---

pervised when many foods are excluded or nutrition could be compromised.

**Alternate Names:**

Exclusion Diet; Diagnostic Elimination Diet

**Elimination Disorders**

A group of childhood-onset disorders involving inappropriate urination or defecation, including enuresis and encopresis. These disorders are diagnosed in a developmentally inappropriate context and are distinguished from incontinence caused by a known medical, neurologic, structural, infectious, or substance-related condition.

**Elimination Half-Life**

The time required for the amount of a medicine in the blood to fall by half. It helps guide dosing intervals and predicts accumulation and clearance; kidney or liver disease, age, and interactions can change it.

**Elimination Patterns**

The timing, frequency, amount, and appearance of urination and bowel movements. Changes can provide clues about hydration, kidney or bowel function, continence, infection, bleeding, or other illness.

**ELISA Blood Test**

A blood test that uses an enzyme-linked reaction to detect or measure an antibody, antigen, hormone, protein, or other substance. Results depend on the test design, timing, sample, and likelihood of disease before testing.

**Elixir**

A sweetened liquid preparation, historically sometimes containing alcohol, used to carry a medicine. The exact ingredients and strength determine its medical effects, interactions, and poisoning risk.

**Elizabethkingia**

A group of environmental bacteria that can rarely cause severe infection, especially in newborns, older adults, or people with weakened immunity. Illness may involve the blood, lungs, brain, or other organs.

**Elliptocyte Count**

The number or proportion of oval or elongated red blood cells seen on a blood smear. Increased elliptocytes may occur in an inherited red-cell disorder or some anemias and must be interpreted with other results.

**Elliptocytosis**

A condition in which many red blood cells are oval or elongated rather than round. Inherited forms may cause red cells to break down early and produce anemia, while some people have no symptoms.

**Alternate Names:**

Ovalocytosis

**Ellis-Van Creveld Syndrome**

An inherited skeletal condition causing short limbs, extra fingers or toes, a narrow chest, and abnormal teeth or nails. Congenital heart disease is common and may affect breathing, growth, or physical function.

**Ellsworth-Howard Test**

A specialized test of how the kidneys respond to parathyroid hormone. It measures changes in urine after the hormone is given and may help distinguish pseudohypoparathyroidism from other causes of low parathyroid hormone effects.

**Alternate Names:**

PTH Responsiveness Test; Parathyroid Hormone Infusion Test

**Elongin**

A protein complex that helps RNA polymerase II continue copying genetic information into RNA. It contributes to gene regulation and cell function and is mainly a molecular-biology term.

**Elotuzumab**

An antibody medicine that helps the immune system recognize and attack myeloma cells. It is used with other medicines for selected multiple-myeloma patients and can cause infusion reactions, infection, fatigue, or low blood counts.

---

## Eltrombopag

An oral medicine that stimulates the thrombopoietin receptor in bone marrow and increases platelet production. It is used for selected low-platelet conditions and can affect the liver, clotting risk, or blood counts.

## Eltrombopag Olamine

The salt form of eltrombopag, an oral medicine that increases platelet production in selected blood disorders. Liver tests, blood counts, medicine interactions, and clotting risk may require monitoring.

## Emaciation

Extreme loss of body weight, muscle, and fat caused by severe malnutrition, chronic disease, cancer, infection, or other illness. It can weaken immunity and function and requires assessment of causes and nutritional needs.

## Embolectomy

Removal of a traveling blood clot or other blockage from a blood vessel with surgery or a catheter procedure to restore blood flow. It may be urgent when a limb or organ is threatened.

### Alternate Names:

Arterial Embolectomy; Surgical Embolectomy

## Emboli

Particles or material carried through the bloodstream that can lodge in a smaller vessel and block blood flow. Emboli may be clots, fat, air, infected material, or rarely tumor fragments.

## Embolic Stroke

A stroke caused by a clot or other material traveling to and blocking an artery that supplies the brain. Emergency treatment may restore blood flow and limit permanent brain injury.

### Alternate Names:

Brain Embolism Cerebral Embolism

## Embolic Stroke of Undetermined Source

A non-lacunar ischemic stroke detected on neuroimaging without proximal arterial stenosis exceed-

ing 50 percent, a major cardioembolic source (such as atrial fibrillation), or other established specific cause despite comprehensive diagnostic evaluation.

### Alternate Names:

ESUS

## Embolism

Blockage of a blood vessel by material that traveled through the bloodstream from another site. The material is most often a clot but may be fat, air, infected material, or rarely tumor tissue.

## Embolization

Deliberate blockage of a blood vessel or abnormal vessel connection using a catheter and materials such as coils, particles, or plugs. It can control bleeding, reduce tumor blood flow, or treat selected aneurysms and malformations.

## Embolization Therapy

An image-guided procedure that blocks selected blood vessels with coils, particles, plugs, or other materials. It can control bleeding, reduce blood supply to a tumor, treat aneurysms, or close abnormal vessel connections.

## Embolus

Material traveling through the bloodstream that may lodge in a smaller vessel and block it. It may be a blood clot, fat, air bubble, infected material, or rarely tumor tissue.

## Embryo

A developing human from fertilization through the end of the eighth week after fertilization. From the ninth week until birth, the developing human is called a fetus.

## Embryo Neoplasm

A tumor arising from primitive cells resembling developing embryonic tissue, usually in a child. Diagnosis and treatment depend on the tissue of origin, microscopic and molecular features, location, and spread.

---

## Embryo Transfer

A fertility procedure in which one or more laboratory-grown embryos are placed into the uterus through a thin catheter. It may be done soon after fertilization or after freezing and thawing; implantation and pregnancy are not guaranteed.

### Alternate Names:

ET; Intrauterine Embryo Transfer

## Embryogenesis

The process by which an embryo develops after fertilization through cell division, movement, specialization, and formation of organs and body structures.

## Embryology

The study of development from fertilization through the embryonic and fetal stages, including how organs and body structures form, grow, and become specialized.

## Embryoma

A rare tumor made of primitive cells resembling early embryonic tissue. The term can refer to different tumor types, so diagnosis depends on the organ involved and microscopic examination.

## Embryonal Carcinoma

An aggressive germ-cell tumor composed of primitive malignant cells, occurring in the testis, ovary, or other sites and often treated with chemotherapy and surgery.

## Embryonal CNS Tumor

A highly cellular malignant pediatric CNS tumor with primitive or embryonal morphology. Diagnosis depends on integrated histologic and molecular classification, and treatment usually combines surgery, radiation when appropriate, and chemotherapy.

## Embryonal Rhabdomyosarcoma

A malignant skeletal-muscle-lineage tumor that commonly affects young children and may arise in the head and neck, urinary tract, or reproductive tract.

## Embryonal Tumors

Cancers that arise from cells resembling developing embryonic tissue, often occurring in children and commonly affecting the nervous system.

## Embryonic

Relating to an embryo, the developing human or animal during the early period after fertilization. In humans, the embryonic period ends at about eight weeks after fertilization, followed by the fetal period.

## Embryonic Cell

A cell of an early embryo capable of division and differentiation into specialized tissues; embryonic stem cells have broad developmental potential.

## Embryonic Induction

A developmental process in which one group of embryonic cells signals nearby cells to develop into a particular tissue or structure.

## Embryonic Mosaic

A developing embryo containing two or more groups of cells with different genetic makeups, produced by a change after fertilization. Effects depend on the cells involved and the proportion of each cell group.

## Embryonic Stage

The early developmental period after fertilization during which the basic body plan and major organs begin to form. In human pregnancy it lasts through about eight weeks after fertilization.

## Emergence Agitation

Temporary confusion, restlessness, fear, or inconsolable crying while waking from anesthesia. It is more common in children and usually resolves with reassurance and monitoring, but pain, low oxygen, or another complication must be excluded.

## Emergency Airway Puncture

An emergency procedure that creates an airway through the neck when severe upper-airway blockage prevents breathing and a breathing tube cannot

---

be placed. It is a temporary, high-risk lifesaving measure performed only by trained professionals.

### **Emergency Care**

Immediate assessment and treatment for a serious or time-sensitive illness or injury. Clinicians first protect breathing, circulation, brain function, and control of bleeding, pain, or infection, then arrange continued treatment or transfer.

### **Emergency Code**

A hospital alert that summons a designated response to an urgent event, such as cardiac arrest, fire, or security threat. The meaning of each code varies by institution and should be confirmed locally.

### **Emergency Contraception**

Birth control used after unprotected sex or contraceptive failure to reduce the chance of pregnancy. Options include an oral medicine or copper intrauterine device; effectiveness depends mainly on the method and how soon it is used.

### **Emergency Department**

The hospital service that evaluates and treats sudden illness and injury. Staff prioritize people by seriousness, stabilize life-threatening problems, diagnose causes, provide initial treatment, and arrange discharge, admission, or transfer.

### **Emergency Medicine**

The medical specialty focused on rapid assessment, stabilization, diagnosis, and initial treatment of sudden illness and injury at any age. It includes resuscitation, urgent procedures, poisoning care, and decisions about admission or transfer.

### **Emergency Procedures**

Planned actions for responding safely to urgent events such as cardiac arrest, airway blockage, contamination, equipment failure, or radiation exposure while protecting patients, staff, and the public.

### **Emerging Infectious Disease**

An infection that is newly recognized, has recently

increased, or is spreading into a new population or region. Causes may include animal-to-human transmission, genetic change, travel, climate, or loss of effective control.

### **Emery-Dreifuss Muscular Dystrophy**

A rare inherited muscle disease causing early tightness around the elbows, ankles, and neck followed by slowly progressive weakness. Heart conduction problems, abnormal rhythms, and cardiomyopathy may occur, so regular heart monitoring is essential.

### **Emesis**

Forceful emptying of stomach contents through the mouth, commonly called vomiting. It may result from infection, medicines, poisoning, pregnancy, blockage, brain disease, or other illness and can cause dehydration.

### **Emetic**

A substance or medicine that causes vomiting. Emetics are generally not used for poisoning because vomiting can cause aspiration, worsen burns or injury, and delay safer treatment.

### **Emetics**

Substances that induce vomiting. They are generally avoided for poisoning because vomit can enter the lungs or cause additional injury, and vomiting does not reliably remove a dangerous substance.

### **EMG**

A test that records electrical activity in muscles, usually with nerve-conduction testing. It helps investigate weakness, numbness, nerve injury, trapped nerves, muscle disease, and disorders of the nerve-muscle connection.

### **Alternate Names:**

Electromyography

### **Eminentia**

An anatomical term for a raised or prominent part of a body structure. It may provide attachment for a muscle or ligament, form part of a joint, or serve as a landmark during examination or surgery.

---

## Emission

The discharge or release of a substance, usually a fluid. In male reproduction, emission is the movement of semen into the urethra before ejaculation.

## Emmenagogue

A substance traditionally claimed to start or increase menstrual bleeding. Evidence and safety vary, and some products can cause poisoning, miscarriage, or interactions; missed or abnormal periods need medical evaluation rather than self-treatment.

## Emmetropia

The usual focusing state of the eye in which incoming parallel light focuses on the retina without extra effort. Nearsightedness, farsightedness, or astigmatism causes light to focus away from the retina.

## Emodin

A plant chemical studied for laxative, anti-inflammatory, and anticancer effects. Evidence and safety are not established for routine treatment, and concentrated products may cause diarrhea, dehydration, or medicine interactions.

## Emollient

A substance applied to skin to soften and moisturize it, reduce water loss, and strengthen the protective barrier. Emollients are commonly used for dry skin and eczema; greasy products can make floors or clothing slippery.

## Emotion

A feeling state involving subjective experience, brain and body responses, behavior, and interpretation of events. Emotions are normal, but persistent or extreme changes may occur with mental or physical illness.

## Emotional Adjustment

The process of adapting thoughts, feelings, and behavior to illness, loss, stress, or a major life change. Difficulty adjusting may cause persistent distress or impaired functioning and may benefit from support or treatment.

## Emotional Dependency

Reliance on another person for reassurance, self-worth, or emotional stability to a degree that limits independence or causes distress. It is not a diagnosis by itself and may reflect anxiety, low self-esteem, trauma, or relationship difficulties.

## Emotional Disorders

Mental-health conditions involving persistent or severe changes in mood or emotional responses that cause distress or interfere with daily life, relationships, school, or work. Examples include depressive, anxiety, and trauma-related disorders.

## Emotional Dysregulation Disorder

Alternate name; see *Borderline Personality Disorder*.

## Emotional Eating

Eating in response to stress, boredom, sadness, anger, or other feelings rather than mainly physical hunger. Occasional emotional eating is common, but frequent episodes can affect health and may benefit from behavioral or psychological support.

## Empathy

The ability to understand another person's feelings and point of view. It supports respectful communication and compassionate care but does not require agreeing with the person or experiencing the same feelings.

## Emphysema

A type of chronic obstructive pulmonary disease in which air sacs are permanently damaged and enlarged, reducing air exchange. Smoking is a common cause; breathlessness, cough, and reduced exercise capacity may develop gradually.

## Emphysematous Aortitis

A rare, life-threatening infection of the aortic wall that produces gas within the artery, usually caused by organisms such as *Clostridium septicum*. It may cause severe back or abdominal pain, sepsis, bleeding, or an abnormal connection with the intestine.

---

### **Emphysematous Cholecystitis**

A severe gallbladder infection in which gas-forming bacteria produce gas in the gallbladder wall or cavity. It is more common in people with diabetes and can rapidly cause tissue death, perforation, or sepsis.

### **Emphysematous Hepatitis**

A rare, life-threatening gas-forming infection of liver tissue, usually caused by bacteria in people with diabetes or weakened immunity. It can cause fever, abdominal pain, sepsis, and liver failure.

### **Emphysematous Prostatitis**

A rare, severe prostate infection in which gas-forming bacteria produce gas within prostate tissue. It may cause fever, pelvic or perineal pain, difficulty urinating, urinary retention, and sepsis.

### **Emphysematous Pyelonephritis**

A severe kidney infection in which gas-forming bacteria destroy kidney tissue. It occurs mainly in people with diabetes and can cause fever, flank pain, shock, and sepsis; urgent drainage, antibiotics, and sometimes surgery are needed.

### **Empiric Antibiotic Therapy**

Starting an antibiotic before the exact germ and its medicine sensitivity are known. The first choice is based on the symptoms, likely source of infection, local resistance patterns, and the person's health; it may be changed when test results return.

### **Empiric Risk**

An estimate of risk based on observed data from a population or clinical setting. Its usefulness depends on how similar the studied population, exposure, and outcome are to the person or situation being assessed.

### **Empirical**

Based on observation, measurement, or experience rather than theory alone. Empirical treatment begins before the exact cause is confirmed and is revised when test results or the clinical course provide more information.

### **Empirical Treatment**

Treatment started on the basis of symptoms and likely causes before a diagnosis is confirmed. It should be reassessed when test results or the response to treatment becomes available.

### **Employee**

A person who works for an employer under an agreement. In occupational health, assessment may consider workplace exposures, safety, physical or mental function, and ability to perform duties safely.

### **Empty Calories**

A nontechnical term for foods or drinks that provide energy but little protein, fiber, vitamins, or minerals, such as many sugary drinks and highly processed snacks. Frequent intake can displace more nutritious foods.

### **Empty Can Test**

A clinical examination manoeuvre used to evaluate the integrity of the supraspinatus muscle and tendon. The patient's arms are abducted ninety degrees in the scapular plane with full internal rotation (thumbs pointing downward), and the examiner applies downward pressure against active resistance; weakness or pain indicates supraspinatus tendinopathy or tear.

#### **Alternate Names:**

Jobe Test; Supraspinatus Test

### **Empty Nose Syndrome**

A poorly understood, controversial condition that can follow extensive surgery on the nasal turbinates, in which the nose feels blocked even though the airway is open. People may describe a dry, suffocating or breathless feeling, inability to sense airflow, crusting, headache, fatigue, and poor concentration, with little relief from decongestants. Findings of obstruction may be absent, so diagnosis is clinical; management is supportive and includes moisturizing, saline care, and consultation with clinicians experienced with the condition.

### **Empty Sella Syndrome**

A condition in which fluid presses the pituitary



---

gland flat within its bony space. It is often found incidentally and causes no symptoms, but some people have headaches, vision problems, or reduced pituitary-hormone production.

### **Empyema**

A collection of pus in a normally sterile body space, most often infected fluid around a lung. It can cause fever, chest pain, cough, breathlessness, and sepsis and usually requires antibiotics plus drainage.

### **Empyema Necessitans**

Extension of infected pus from the space around a lung through the chest wall, forming a painful swelling. It is a serious complication requiring imaging, antibiotics, and drainage.

### **Empyema Necessitatis**

The direct extension and burrowing of an intrapleural empyema through the parietal pleura and chest wall tissues into the subcutaneous tissue and skin, creating an extrapleural abscess, classically caused by *Mycobacterium tuberculosis*, *Actinomyces israelii*, or neglected bacterial empyema.

#### **Alternate Names:**

Empyema Necessitatis; Pleurocutaneous Fistula

### **EMR**

Electronic medical record: a digital record of health information maintained within a particular health-care organization or practice. It may include diagnoses, notes, medicines, allergies, test results, and treatment plans.

### **Emtricitabine**

An antiretroviral medicine that blocks HIV reverse transcriptase and is used with other medicines to treat or prevent HIV infection. Kidney function and hepatitis B status are important because stopping it can worsen hepatitis B.

### **Emulsion**

A mixture of two liquids that normally do not blend, such as oil and water, with one dispersed in the other. Emulsions are used in some medicines, nutrition products, cosmetics, and laboratory preparations.

### **Enalapril**

A medicine that blocks formation of angiotensin II, relaxing blood vessels and reducing strain on the heart. It treats high blood pressure and heart failure but can cause cough, high potassium, kidney problems, or rarely facial swelling.

### **Enalaprilat**

The active injectable form related to enalapril that blocks angiotensin II formation and lowers blood pressure. It is used in selected settings and can cause low blood pressure, high potassium, kidney problems, or facial swelling.

### **Enamel**

The hard, mineral-rich outer covering of a tooth crown. It protects the sensitive inner tooth but cannot repair itself after major damage; acids and bacteria can cause erosion and decay.

### **Enamel Hypoplasia**

Incomplete formation of tooth enamel while a tooth is developing, leaving it thin, grooved, pitted, or discolored. Affected teeth are more vulnerable to sensitivity, wear, and decay.

### **Enamel Organ**

The developing tissue of a tooth that produces enamel and helps shape the crown. It is present during tooth formation and later contributes to the protective covering around the developing tooth.

### **Enamel Pearl**

A small lump of enamel on a tooth root, often near where the roots divide. It can make cleaning difficult and interfere with gum attachment, increasing the risk of localized gum disease.

### **Enanthem**

A rash, spot, ulcer, or other lesion on a mucous membrane, especially inside the mouth. It may occur with infection, inflammation, an allergic or medicine reaction, or a systemic illness.

---

## **Enanthema**

A rash or lesion on a moist membrane lining the mouth, throat, genitals, or another body surface. Infection, inflammation, medicines, or systemic disease may cause it.

## **Enarthrosis**

A ball-and-socket joint that allows movement in many directions, including bending, straightening, rotation, and movement away from or toward the body. The hip and shoulder are examples.

## **Encapsulated**

Enclosed by a capsule or clearly defined layer of tissue. This may describe a cyst, growth, infection, or organ; being encapsulated does not by itself prove that a lesion is harmless or cancerous.

## **Encapsulated Peritoneal Sclerosis**

A rare serious complication of long-term peritoneal dialysis in which scar tissue forms around the intestines. It can block the bowel and cause repeated pain, vomiting, and poor nutrition.

### **Alternate Names:**

Sclerosing Encapsulating Peritonitis; EPS

## **Encapsulating Peritoneal Sclerosis**

A rare condition in which thick scar tissue surrounds the intestines and can cause repeated blockage, pain, vomiting, and poor nutrition. It is most associated with long-term peritoneal dialysis but can have other causes.

## **Encephalitis**

Inflammation of brain tissue, often caused by an infection or an abnormal immune reaction. It may cause fever, headache, confusion, seizures, behavior change, weakness, or movement problems and requires urgent medical care.

## **Encephalitis Lethargica**

A rare inflammatory brain disorder described mainly during the early twentieth century, causing sleepiness, abnormal movements, eye-movement problems, and other neurologic symptoms. Its cause remains

uncertain.

## **Encephalitis Virus**

A virus capable of infecting the brain and causing encephalitis. It may spread through mosquitoes, ticks, animals, respiratory secretions, food, or other routes depending on the virus.

## **Encephalitozoon**

A group of tiny spore-forming parasites that can infect the intestine, eyes, kidneys, or several organs. Serious disease occurs mainly in people with weakened immunity.

## **Encephalocele**

A birth defect in which the coverings of the brain and sometimes brain tissue protrude through an opening in the skull. It may cause a visible head swelling and can be associated with seizures, fluid buildup, developmental problems, or other brain abnormalities.

## **Encephalofacial Angiomatosis**

Sturge-Weber syndrome, a congenital neurocutaneous disorder involving capillary malformations of the face and leptomeninges. It may cause seizures, glaucoma, stroke-like episodes, headaches, and developmental or visual problems.

## **Encephalomalacia**

Softening or loss of brain tissue after injury, stroke, bleeding, infection, or reduced blood flow. The resulting permanent damage may cause seizures, weakness, movement problems, memory changes, or other neurologic effects.

## **Encephalomyelitis**

Inflammation affecting both the brain and spinal cord, caused by infection or an abnormal immune response. It may cause confusion, seizures, weakness, numbness, balance problems, or bladder changes and requires urgent neurologic assessment.

## **Encephalopathy**

A disorder that broadly affects brain function, caus-

---

ing confusion, drowsiness, personality change, difficulty thinking, or coma. Causes include infection, toxins, organ failure, medicines, metabolic problems, seizures, and reduced oxygen.

### **Encephalotrigeminal Angiomatosis**

A birth condition involving abnormal blood vessels in the skin of the face and the coverings of the brain. It may cause a port-wine stain, seizures, weakness, learning problems, or glaucoma, but symptoms vary.

#### **Alternate Names:**

Sturge-Weber Syndrome; SWS

### **Enchondroma**

A usually harmless cartilage growth inside a bone, often in the hands or long bones and found by chance. Pain, enlargement, fracture, or unusual imaging features may require evaluation for a more serious tumor.

### **Enchondromatosis**

A disorder in which multiple cartilage growths develop inside bones. It can cause uneven growth, limb deformity, pain, or fractures; a small number of lesions may become cancerous and require monitoring.

### **Encoding**

The process by which sensory or other information is changed into a form the brain can store in memory. Attention, sleep, stress, illness, and brain injury can affect encoding.

### **Encopresis**

Repeated passage of stool in inappropriate places by a child at an age or developmental level when bowel control is expected. It is often related to long-term constipation and retained stool but may have other medical, developmental, or emotional contributors.

### **End Point**

A predefined outcome or measurement used to judge a clinical study, treatment, diagnostic test, or disease process. It may be a symptom, laboratory result, event, survival measure, or patient-reported outcome.

### **End Stage**

The advanced phase of a disease in which normal function is severely impaired. Meaning, criteria, expected course, and treatment options depend on the disease, such as advanced kidney, liver, lung, or heart failure.

### **End-Of-Life Care**

Care for a person nearing death that focuses on comfort, symptom relief, dignity, communication, and personal goals. It also supports family and caregivers and may include hospice or palliative services.

### **End-Stage Kidney Disease**

Advanced chronic kidney disease in which the kidneys can no longer adequately remove waste or control fluid, electrolytes, and acid levels. Treatment options include dialysis, kidney transplantation, or supportive care based on health and preferences.

### **End-Stage Renal Disease**

The most advanced stage of chronic kidney disease, usually when kidney filtration is below 15 or dialysis is needed. Waste and fluid can accumulate, causing nausea, itching, confusion, swelling, anemia, or heart complications.

#### **Alternate Names:**

ESRD

### **Endangiitis**

Inflammation of the inner lining of a blood vessel. Swelling or scarring may narrow the vessel and reduce blood flow; the effects depend on the cause and the vessels involved.

### **Endarterectomy**

Surgery that removes fatty plaque and diseased inner lining from an artery to improve blood flow, most often a carotid artery in the neck. Possible complications include bleeding, stroke, heart attack, or nerve injury.

### **Endarteritis**

Inflammation of the inner lining of an artery that

---

can narrow the vessel and reduce blood flow. Causes include infection, autoimmune disease, injury, and other vascular disorders.

## **Endemic**

Regular presence of a disease or infectious agent in a particular population or region at an expected baseline level. Endemic does not mean that every person is infected or that cases cannot increase.

## **Endemic Diseases**

Diseases that remain present at a predictable level in a particular population or region. Their frequency is shaped by local environment, immunity, behavior, exposures, healthcare access, and prevention measures.

## **Endemic Fungi**

Fungi naturally found in particular geographic regions whose spores can cause infection when inhaled. Disease may affect the lungs and sometimes spread to other organs, especially in people with weakened immunity.

## **Endemic Goiter**

Thyroid enlargement occurring commonly in a population, traditionally because of insufficient iodine intake. It may occur with normal, low, or high thyroid function depending on the cause.

## **Endergonic**

Describing a chemical reaction that requires an input of energy. In cells, energy from food or another reaction drives processes such as building proteins, DNA, or other complex molecules.

## **Ending Pregnancy With Medicines**

Ending a pregnancy with prescribed medicines that cause the uterus to empty. The appropriate method depends on pregnancy duration, health, symptoms, and local clinical requirements; follow-up confirms completion and checks for complications.

## **Endobronchial Foreign Body**

An object lodged in a bronchus, partly or completely

blocking an airway. It may cause sudden coughing, one-sided wheezing, collapse of part of the lung, or pneumonia and often requires urgent bronchoscopy for removal.

## **Endobronchial Tumor**

A growth inside a bronchus that can narrow the airway and cause cough, wheezing, coughing blood, lung collapse, or repeated pneumonia. It may be benign or cancerous; diagnosis usually requires imaging and tissue sampling.

## **Endobronchial Ultrasound**

A procedure combining bronchoscopy with ultrasound to examine structures beside the airways and guide needle sampling of lymph nodes or masses. It is commonly used to diagnose and stage lung cancer and other chest diseases.

## **Endobronchial Valve**

A one-way valve placed through a bronchoscope into a selected airway to reduce air trapping in severe emphysema. It lets air and mucus leave but limits new air entering; pneumothorax and infection are possible complications.

## **Endocannabinoid**

A signaling molecule made naturally by the body that activates cannabinoid receptors. Endocannabinoids help regulate pain, appetite, mood, memory, movement, and immune activity.

## **Endocardial Cushion Defect**

A birth defect in which the central wall of the heart and the valves between its chambers do not form normally. It can allow abnormal blood flow and cause heart failure, high pressure in lung vessels, poor growth, or breathing difficulty.

## **Endocardial Fibroelastosis**

Thickening of the inner lining of the heart from excess fibrous and elastic tissue, usually in infants or children. It can stiffen the heart or weaken its pumping and may lead to heart failure.

---

## Endocarditis

Infection or inflammation of the heart's inner lining, usually involving a valve. It can cause fever, fatigue, a new or changed murmur, stroke or other emboli, valve destruction, or heart failure.

### Alternate Names:

Infective Endocarditis; Valvular Endocarditis

## Endocarditis Prophylaxis

Preventive antibiotics given before selected dental procedures to people at highest risk of serious infective endocarditis complications. They are not needed for everyone; eligibility depends on the heart condition and procedure.

## Endocarditis Surgical Indications

Reasons surgery may be needed for endocarditis include heart failure from valve damage, uncontrolled infection or abscess, infection of a prosthetic valve, fungal infection, or a high risk of emboli. Decisions require a specialist heart team.

## Endocardium

The thin inner lining of the heart chambers and valves. It provides a smooth surface for blood flow and continues into the inner lining of blood vessels; inflammation or infection can damage it.

## Endocervical

Relating to the inner canal or lining of the cervix, the lower part of the uterus that connects the uterine cavity with the vagina.

## Endocervical Culture

A laboratory test that attempts to grow bacteria or fungi from a sample collected from the inner cervix. It may help investigate cervicitis or selected infections, although molecular tests are preferred for many sexually transmitted infections.

## Endocervical Gram Stain

A microscope test that stains cells and organisms from the inner cervix. It may show inflammation or some bacteria but cannot by itself reliably diagnose every sexually transmitted infection.

## Endocrine

Relating to hormones released into the bloodstream by glands or specialized cells. Endocrine organs include the thyroid, pituitary, adrenal glands, pancreas, and reproductive glands.

## Endocrine Disease

A disorder of a hormone-producing gland or hormone action. It may affect growth, metabolism, reproduction, stress responses, blood pressure, or fluid and electrolyte balance.

## Endocrine Disorders

Diseases in which hormone-producing glands make too much or too little hormone, or the body does not respond to hormones properly. They may affect growth, metabolism, reproduction, mood, blood pressure, and fluid balance. Examples include thyrotoxicosis, hypothyroidism, hypogonadism, Cushing syndrome, and Addison's disease.

## Endocrine Disrupting Chemicals

Chemicals that can interfere with hormone production, transport, breakdown, or action. Some exposures are associated with reproductive, developmental, thyroid, metabolic, or cancer outcomes, but risk depends on the substance, dose, and timing.

## Endocrine Disruptor

A chemical that interferes with hormone systems. Possible effects include altered development, reproduction, thyroid function, or metabolism; health risk depends on the chemical, exposure level, duration, and timing.

## Endocrine Finding

A symptom, examination result, imaging result, or laboratory result that may suggest abnormal hormone production, hormone action, or an endocrine-gland disorder.

## Endocrine Function

The production, release, transport, and action of hormones that regulate metabolism, growth, reproduction, stress responses, blood pressure, and fluid or electrolyte balance.

---

## Endocrine Gland

An organ that releases hormones directly into the bloodstream. Its chemical messages regulate growth, metabolism, reproduction, stress responses, blood pressure, and other body functions.

## Endocrine Glands

Ductless organs that release hormones into the bloodstream to regulate distant tissues. They include the pituitary, thyroid, adrenal glands, pancreas, and reproductive glands.

## Endocrine Hypertension

High blood pressure caused by excess or abnormal hormone activity. Causes include primary aldosteronism, excess cortisol, or excess adrenaline-like hormones from an adrenal or related tumor.

## Endocrine Myopathies

Muscle disorders caused by hormonal disease, such as thyroid, adrenal, parathyroid, or pituitary dysfunction. Weakness, exercise intolerance, cramps, or muscle pain may improve when the underlying endocrine disorder is treated.

## Endocrine Neoplasm

A benign or cancerous growth arising in a hormone-producing gland or cell. It may release too much hormone or cause symptoms by pressing on nearby structures; evaluation depends on its location and behavior.

## Endocrine Pancreas

The hormone-producing part of the pancreas, made mainly of pancreatic islets. Its cells release insulin, glucagon, somatostatin, and other hormones that help regulate blood glucose and digestion.

## Endocrine System

The network of glands and hormone-producing cells and tissues that release chemical messengers into the blood. It includes structures such as the hypothalamus, pituitary, thyroid, parathyroids, adrenal glands, pancreatic islets, and gonads; its hormones regulate growth, metabolism, reproduction, stress responses, sleep, blood pressure, fluid balance, and

other aspects of homeostasis.

## Endocrine Testing

Blood, urine, saliva, stimulation, suppression, and sometimes imaging tests used to assess hormone production and action. Results depend on timing, medicines, age, sex, illness, and the test method.

## Endocrine Therapy

Treatment that lowers, blocks, or changes hormone signals to control hormone-sensitive cancer. It may be used for breast, prostate, or other cancers; effects depend on the medicine and can include hot flashes, sexual changes, or bone loss.

## Endocrine Tissues

Cells or groups of cells that produce and release hormones into the bloodstream or nearby tissues. They are found in glands and in organs such as the pancreas, ovaries, testes, kidneys, and intestine.

## Endocrinologist

A physician who diagnoses and treats disorders of hormones, endocrine glands, and metabolism, including diabetes, thyroid disease, adrenal disease, osteoporosis, and reproductive hormone problems.

## Endocrinologists

Physicians specializing in hormone, endocrine-gland, and metabolic disorders, including diabetes, thyroid disease, adrenal disease, osteoporosis, and reproductive hormone problems.

## Endocrinology

The medical specialty concerned with hormones, endocrine glands, and metabolism. It diagnoses and treats conditions affecting diabetes, thyroid function, growth, reproduction, bones, adrenal glands, and fluid or electrolyte balance.

## Endocrinopathy

A disorder of an endocrine gland or hormone system that causes too much, too little, or abnormal hormone action. Effects depend on the hormone and may involve growth, metabolism, reproduction,



---

mood, or fluid balance.

### **Endocytosis**

The process by which a cell folds its outer membrane around material and brings it inside in a small sac. Cells use it to absorb nutrients, remove particles, and control surface signals.

### **Endoderm**

The innermost of the three early embryonic cell layers. It develops into the linings of the digestive and respiratory systems and contributes to organs such as the liver, pancreas, thyroid, and bladder.

### **Endodontics**

The dental specialty concerned with the soft tissue inside teeth, root canals, and tissues around tooth roots. Endodontists diagnose and treat tooth infection, inflammation, injury, and related pain.

### **Endodontist**

A dentist with advanced training in diagnosing and treating disease or injury of the tooth's inner tissue, root canals, and tissues around tooth roots. They commonly perform complex root-canal treatment.

### **Endogenous**

Originating within the body or produced by its own cells or tissues, rather than introduced from outside. Endogenous hormones, chemicals, infections, or substances may be contrasted with external sources.

### **Endogenous Cushing Syndrome**

Hypercortisolism caused by internal overproduction of cortisol or its stimulatory hormones, including pituitary, adrenal, or ectopic sources. Features may include central weight gain, hypertension, diabetes, proximal weakness, bruising, and bone loss.

### **Endogenous Opiates**

Opioid-like substances naturally made by the body, including endorphins, enkephalins, and dynorphins. They act on opioid receptors and influence pain, stress, mood, reward, movement, and other nervous-system functions.

### **Endogenous Opioids**

Opioid-like peptides made naturally by the body, including endorphins, enkephalins, and dynorphins. They activate opioid receptors to influence pain, stress, mood, reward, movement, and hormone regulation.

#### **Alternate Names:**

Endorphins and Enkephalins; Endogenous Opioid Peptides

### **Endogenous Retrovirus**

A piece of viral DNA inherited in the human genome from an ancient virus that infected a reproductive cell. Most such sequences are inactive, although some can influence gene control or normal development.

### **Endoleak**

Continued blood flow into the weakened part of an artery after endovascular aneurysm repair. Some types indicate an inadequate seal or device problem and may need urgent repair; others are monitored according to sac growth and risk.

### **Endolimax**

A group of amoebas that can live in the intestine. They are usually harmless colonizers rather than causes of disease, and a stool finding should be interpreted with symptoms and other test results.

### **Endolimax Nana**

An amoeba that can live harmlessly in the intestine and appear in stool tests. It usually does not cause illness; its presence may indicate exposure to contaminated food or water and should not alone explain digestive symptoms.

### **Endolymph**

Fluid inside the membranous part of the inner ear that helps sensory organs detect sound and head movement. Abnormal pressure or composition can contribute to dizziness, hearing changes, or ringing.

### **Endolymphatic Hydrops**

Pathological distension and increased volume of the

---

endolymphatic fluid compartment of the membranous inner ear (scala media and vestibular apparatus) resulting from impaired fluid resorption in the endolymphatic sac, constituting the underlying pathophysiologic hallmark of Ménière disease.

**Alternate Names:**

Meniere Hydrops; Inner Ear Hydrops

## Endometrial

Relating to the endometrium, the inner lining of the uterus. This lining changes during the menstrual cycle and can support implantation of a pregnancy; abnormal growth may cause bleeding or affect fertility.

## Endometrial Ablation

A procedure that destroys much of the uterine lining to reduce heavy menstrual bleeding. It is not contraception, pregnancy afterward can be dangerous, and it is generally intended for people who do not plan future pregnancy.

## Endometrial Adenocarcinoma

A gland-forming cancer of the uterine lining. It often causes abnormal bleeding, especially after menopause; treatment depends on the cancer's type, grade, stage, molecular features, and overall health.

## Endometrial Biopsy

Removal of a small sample of the uterine lining for laboratory examination, often to investigate abnormal bleeding or suspected overgrowth or cancer. Cramping and light bleeding are common; occasionally the sample is insufficient for diagnosis.

## Endometrial Cancer

Cancer arising from the inner lining of the uterus. Abnormal bleeding—especially after menopause—is a common warning sign; evaluation usually includes examination, imaging, and a sample of the uterine lining.

**Alternate Names:**

Uterine Endometrial Neoplasm; Endometrial Carcinoma

## Endometrial Carcinoma

Cancer arising from the uterine lining, often causing abnormal bleeding, particularly after menopause. Diagnosis requires examination of uterine-lining tissue, and treatment depends on type and spread.

## Endometrial Disorder

Any condition affecting the uterine lining, including inflammation, polyps, abnormal overgrowth, endometriosis-related changes, or cancer. It may cause heavy or irregular bleeding, pelvic pain, infertility, or no symptoms.

## Endometrial Hyperplasia

Overgrowth of glands in the uterine lining, often after prolonged estrogen exposure without enough progesterone. It may cause heavy, irregular, or postmenopausal bleeding. It is classified as hyperplasia without atypia or atypical hyperplasia, also called endometrial intraepithelial neoplasia (EIN). Hyperplasia without atypia is not precancer; atypical hyperplasia/EIN is precancerous and can sometimes be found alongside endometrial cancer, so it needs prompt evaluation.

## Endometrial Implants

Alternate name; see *Endometriosis*.

## Endometrial Intraepithelial Neoplasia

A precancerous overgrowth of abnormal glands in the uterine lining. It can exist with or progress to endometrial cancer, so treatment may involve removal of the uterus or hormone treatment when preserving fertility is important.

**Alternate Names:**

EIN; Atypical Endometrial Hyperplasia

## Endometrial Neoplasm

An abnormal growth of the uterine lining that may be benign, precancerous, or cancerous. Symptoms, tissue examination, imaging, and—if cancer is present—stage determine treatment.

## Endometrial Polyp

A localized growth of the uterine lining that projects

---

into the uterine cavity. It may cause irregular or heavy bleeding, infertility, or no symptoms; some polyps contain precancerous or cancerous cells.

### **Endometrial Polyps**

Localized growths of the uterine lining projecting into the uterine cavity. They may cause irregular or heavy bleeding, infertility, or no symptoms, and some require removal for tissue examination.

### **Endometrial Receptivity Analysis**

A test that examines activity of genes in a sample of the uterine lining to estimate when it may best support embryo implantation. Its ability to improve pregnancy rates is uncertain, so results need specialist interpretation.

#### **Alternate Names:**

ERA; Endometrial Receptivity Assay

### **Endometrial Stromal Nodule**

A rare, usually harmless, well-defined growth of connective-tissue cells in the uterus. It differs from cancer by not invading nearby tissue or blood vessels, but microscopic examination is needed for reliable classification.

### **Endometrial Stromal Sarcoma**

A rare cancer arising from connective-tissue cells of the uterine lining. Some forms grow slowly and others aggressively; abnormal bleeding or pelvic symptoms may occur, and treatment depends on grade, spread, hormone receptors, and molecular findings.

#### **Alternate Names:**

ESS

### **Endometrioid Carcinoma Of The Vulva**

A rare gland-forming cancer of the vulvar skin or nearby tissue that resembles cancer of the uterine lining. Diagnosis requires tissue examination and confirmation that it did not spread from the uterus or ovary.

### **Endometrioid Ovarian Carcinoma**

A type of ovarian cancer whose cells resemble glands

of the uterine lining. It may be associated with endometriosis; symptoms, treatment, and outlook depend on stage, grade, and tumor features.

### **Endometrioma**

An ovarian cyst caused by endometriosis in which old blood collects. It may cause pelvic pain, painful periods, pain during sex, or infertility, although some people have no symptoms.

### **Endometriosis**

A chronic inflammatory condition in which tissue resembling the lining of the uterus grows outside the uterus, most often in the pelvis, ovaries, fallopian tubes, or tissue around the pelvic organs. It can also affect the bowel or bladder. This tissue responds to hormones and can cause repeated inflammation, bleeding, scar tissue, adhesions, or ovarian cysts. Symptoms may include severe period pain, ongoing pelvic pain, heavy or irregular bleeding, pain during sex or bowel movements, painful urination, bloating, nausea, tiredness, or infertility. Some people have no symptoms.

#### **Alternate Names:**

Endometrioma; Pelvic Endometriosis

### **Endometritis**

Inflammation or infection of the uterine lining, often after childbirth, miscarriage, abortion, or a procedure entering the uterus. It may cause fever, pelvic pain, tenderness, abnormal discharge, or bleeding and needs treatment.

### **Endometrium**

The inner lining of the uterus. It thickens each month under hormone control to prepare for pregnancy and sheds during menstruation; abnormal thickening, inflammation, polyps, or cancer can cause bleeding or fertility problems.

### **Endomyocardial Biopsy**

A procedure that removes tiny samples from the heart muscle through a catheter in a vein. The samples can help check for rejection after a heart transplant or investigate some heart-muscle diseases.

#### **Alternate Names:**

### **Endomyocardial Fibrosis**

Scarring and thickening of the inner heart lining and nearby muscle that makes the heart stiff and reduces filling. It can cause breathlessness, swelling, abnormal rhythms, or heart failure.

### **Endomysium**

A delicate layer of reticular connective tissue that individually invests each single skeletal muscle fiber, containing an intricate network of capillaries and nerve terminals essential for metabolic exchange and muscle contraction.

### **Endoneurium**

A delicate layer of loose connective tissue and collagen fibrils that directly surrounds each individual myelinated or unmyelinated axon and its accompanying Schwann cell sheath within a peripheral nerve fascicle.

### **Endoparasite**

A parasite that lives inside its host, such as a worm in the intestine or a microscopic parasite in blood or tissue. It may cause no symptoms or lead to inflammation, anemia, organ damage, or infection.

### **Endopeptidase**

An enzyme that cuts proteins or peptides at internal points rather than removing amino acids only from an end. Digestive enzymes such as pepsin and trypsin are examples.

### **Endophthalmitis**

A severe infection or inflammation inside the eye, usually after eye surgery, an injury, or spread through the bloodstream. It can cause severe pain, redness, swelling, and rapid vision loss and requires emergency eye treatment.

### **Endoplasmic Reticulum**

A network of tiny membrane-lined spaces inside cells. It helps make and transport proteins and fats, stores calcium, and processes substances needed for

cell structure and function.

### **Endoprosthesis**

An artificial device placed inside the body to replace, support, or restore a body structure. Examples include joint replacements, vascular grafts or stents, and some devices used to keep an airway open.

### **Endoribonuclease**

An enzyme that cuts RNA at internal chemical bonds. It helps process RNA into usable forms or break down RNA that is damaged, unnecessary, or produced by viruses.

### **Endorphin**

A natural chemical made by the body that acts on opioid receptors in the brain and other tissues. It can reduce the feeling of pain and influence stress, mood, and other body responses.

### **Endorphins**

Natural opioid-like chemicals made by the body that reduce pain and influence mood. They are released during stress, exercise, injury, and other situations and act through opioid receptors in the nervous system.

#### **Alternate Names:**

Endogenous Opioid Peptides; Beta-Endorphin

### **Endoscope**

A flexible or rigid instrument with a light and camera used to view an internal body passage or cavity. It can sometimes collect tissue, stop bleeding, remove growths, or perform other treatments.

### **Endoscopic Endonasal Surgery**

A minimally invasive operation performed through the nostrils and nasal passages using an endoscope, commonly to reach selected skull-base, pituitary, sinus, or nearby lesions. Risks include bleeding, infection, cerebrospinal-fluid leak, and injury to nearby structures.

### **Endoscopic Mucosal Resection**

An endoscopic procedure that removes a superficial

---

growth from the digestive tract after lifting it away from deeper tissue. The removed tissue is examined to determine whether it was completely removed and whether cancer is present.

### **Endoscopic Retrograde Cholangiopancreatography**

A procedure combining an endoscope and X-ray imaging to examine and treat the bile and pancreatic ducts. It can remove stones, open narrowed ducts, place stents, or obtain samples; pancreatitis, bleeding, and infection are possible risks.

### **Endoscopic Submucosal Dissection**

An endoscopic procedure that removes a larger or flatter superficial growth from the digestive tract in one piece by dissecting beneath it. It allows detailed tissue examination but carries risks of bleeding and perforation.

### **Endoscopic Third Ventriculostomy**

A brain procedure used to treat some types of hydrocephalus, in which fluid builds up because its normal path is blocked. A small camera is used to make an opening in one of the brain's fluid spaces so the fluid can drain. It may reduce the need for a shunt, but not everyone is a suitable candidate and the opening can close later.

#### **Alternate Names:**

ETV; Third Ventriculostomy; Endoscopic Ventriculostomy

### **Endoscopic Thoracic Sympathectomy**

A minimally invasive operation that interrupts selected nerves in the chest, usually for severe sweating of the hands. Possible complications include increased sweating elsewhere, drooping eyelid, collapsed lung, nerve injury, and heart or blood-pressure changes.

### **Endoscopic Ultrasound**

A procedure combining an endoscope with ultrasound to view the digestive-tract wall and nearby organs. It can guide needle sampling of lymph nodes, pancreas, or other masses and help assess the depth or spread of disease.

### **Endoscopy**

Examination of an internal body passage or cavity with a camera-equipped instrument. Depending on the site, it can directly inspect tissue, collect samples, remove growths, widen narrow areas, or stop bleeding.

### **Endosome**

A small membrane-bound compartment inside a cell that receives material taken in from the cell surface and sorts it for recycling, transport, or breakdown.

### **Endosteum**

A thin membrane lining the inner surfaces of bones, including the marrow cavity and small internal channels. It contains cells involved in bone formation, remodeling, and repair.

### **Endostosis**

Abnormal formation of bone on or within an existing bone. It may be a localized developmental finding or part of a broader bone disorder; the significance depends on its location, extent, and symptoms.

### **Endothelial**

Relating to the thin layer of cells lining blood and lymphatic vessels and some internal surfaces. These cells help control blood flow, vessel leakiness, inflammation, clotting, and new-vessel growth.

### **Endothelial Cell**

A cell lining a blood or lymphatic vessel. It regulates movement between blood and tissues and helps control vessel widening, inflammation, clotting, and formation of new blood vessels.

### **Endothelial Nitric Oxide Synthase**

A calcium- and calmodulin-dependent enzyme constitutively expressed in vascular endothelial cells that synthesizes nitric oxide from L-arginine, mediating basal vasodilation, inhibition of leukocyte adhesion, and suppression of platelet aggregation.

#### **Alternate Names:**

eNOS; NOS3

---

## Endothelin

A small protein made mainly by cells lining blood vessels that strongly narrows vessels and affects blood pressure, circulation, inflammation, and tissue growth. Excess signaling contributes to some cases of pulmonary arterial hypertension.

## Endothelin Receptor Antagonists

Medicines that block signals that narrow blood vessels, especially in the lungs. They are used for selected pulmonary arterial hypertension; liver injury, anemia, swelling, and serious fetal harm are important risks.

## Endothelium

The thin layer of specialized cells lining blood and lymphatic vessels. It controls exchange between blood and tissues and helps regulate vessel diameter, inflammation, clotting, and new-vessel growth.

## Endothelium-Derived Relaxing Factor

A substance released by vessel-lining cells that relaxes blood vessels. It is largely identified with nitric oxide, which also helps reduce platelet sticking and vessel-wall inflammation.

### Alternate Names:

EDRF; Nitric Oxide

## Endotoxin

A component of the outer membrane of certain bacteria, especially Gram-negative bacteria. When released into the body, it can trigger fever, inflammation, low blood pressure, organ injury, and life-threatening sepsis.

## Endotracheal Intubation

Placement of a breathing tube through the mouth or nose into the windpipe to keep the airway open, deliver ventilator support, give anesthetic gases, or protect the lungs. Incorrect placement can cause severe harm and must be confirmed.

## Endotracheal Tube

A breathing tube passed through the mouth or nose into the windpipe to maintain an airway and support

ventilation. An inflatable cuff may seal the airway; irritation, injury, infection, or incorrect placement can occur.

## Endourologist

A urologist specializing in minimally invasive procedures performed through the urinary tract or small access sites. Common work includes treating urinary stones, narrowing, blockage, and selected tumors.

## Endovaginal Ultrasound

Ultrasound imaging performed with a covered probe placed gently in the vagina to examine the uterus, uterine lining, ovaries, cervix, and early pregnancy. It uses sound waves rather than radiation and may cause temporary discomfort.

## Endovascular

Located inside a blood vessel or performed from within it, usually with a catheter passed through a small skin puncture. Endovascular procedures can treat aneurysms, clots, narrowing, bleeding, or abnormal vessels.

## Endovascular Aneurysm Repair

A catheter procedure that places a fabric-covered graft inside an aortic aneurysm to reinforce the artery and reduce pressure on its weakened wall. It usually recovers faster than open surgery but requires lifelong imaging for leakage or device problems.

### Alternate Names:

EVAR; Endovascular Stent-Graft Repair; Endovascular Aortic Repair

## Endovascular Aortic Aneurysm Repair

A minimally invasive procedure in which an expandable fabric-covered stent graft is deployed within the diseased aortic segment via femoral or iliac arterial access to exclude the aneurysm sac from systemic arterial pressure and prevent rupture.

### Alternate Names:

Endovascular Aneurysm Repair; EVAR; TEVAR; Endovascular Aortic Repair



---

## Endovascular Coiling

A minimally invasive, catheter-based procedure performed to treat brain aneurysms from within the blood vessels without opening the skull. Guided by real-time X-ray fluoroscopy, an interventional neurosurgeon or neuroradiologist threads a long, micro-thin catheter through an artery in the groin (femoral artery) or wrist (radial artery) all the way up into the blood vessels of the brain and directly inside the bulging aneurysm sac. Soft, ultra-fine platinum coils (each as thin as a human hair) are deployed through the catheter into the aneurysm, where they pack together and induce natural blood clot formation (thrombosis). The resulting clot seals the aneurysm from the inside, preventing blood from entering and drastically reducing the risk of aneurysm rupture and fatal intracranial bleeding.

### Alternate Names:

Endovascular Aneurysm Coiling; Guglielmi Detachable Coiling; GDC; Aneurysm Embolization Coils

## Endovascular Embolization

A catheter procedure that deliberately blocks a selected blood vessel using coils, particles, plugs, or another material. It may control bleeding or treat an aneurysm, malformation, or tumor blood supply; non-target blockage is a possible risk.

## Endovascular Graft

A fabric-covered artificial tube placed inside a diseased blood vessel through a catheter. It can reinforce a weak artery or exclude an aneurysm; suitability and follow-up depend on anatomy, symptoms, and imaging.

## Endovascular Repair

A minimally invasive procedure that uses a catheter-delivered graft, balloon, stent, or other device to repair a blood vessel from inside. Bleeding, clotting, infection, leakage, or need for further treatment can occur.

## Endovenous Laser Ablation

A treatment for some varicose veins in which a thin laser fiber is placed inside the vein through a small needle opening. Heat closes the treated vein, and

blood flows through healthier veins. It is usually done with local numbing medicine; bruising, pain, clots, nerve irritation, or return of the vein can occur.

### Alternate Names:

EVLA; Endovenous Laser Therapy; EVLT; Endovenous Thermal Ablation

## Endurance

The ability to continue physical or mental activity without becoming tired quickly. It depends on heart and lung function, muscle conditioning, energy, sleep, and overall health and can be assessed with activity or exercise tests.

## Enema

Introduction of liquid into the rectum through the anus to soften stool, stimulate bowel emptying, deliver selected treatment, or assist a test. Overuse can cause irritation, dehydration, or electrolyte problems.

## Energy

The capacity needed for body functions, movement, growth, and temperature control, obtained mainly from food. Persistent low energy may reflect poor sleep, inadequate nutrition, anemia, hormone disease, mood disorder, infection, or another illness.

## Energy Conservation

Strategies that reduce unnecessary effort and fatigue during daily activities, such as planning, pacing, prioritizing, resting, sitting for tasks, changing position, and using helpful equipment.

## Energy Expenditure

The energy the body uses for basic functions, movement, digestion, growth, pregnancy, and temperature control. It varies with body size, muscle mass, age, activity, illness, and other conditions.

## Energy Increased

A noticeable rise in energy or activity. It may be normal, but unusually high energy with little need for sleep, racing thoughts, impulsive behavior, or irritability can occur with stimulant use, thyroid

---

disease, mania, or hypomania.

### **Energy Intake**

The amount of energy obtained from food and drink, usually measured in calories. Appropriate intake depends on age, body size, activity, illness, pregnancy, and nutritional goals.

### **Energy Metabolism**

The chemical processes that convert nutrients into usable energy for cells. They include breaking down carbohydrates, fats, and proteins and storing or using the resulting energy for body functions.

### **Energy Resolution**

The ability of a radiation detector to distinguish radiation particles with different energies. Better resolution allows more accurate identification and separation of radioactive substances during imaging or laboratory measurement.

### **Energy Window**

The range of radiation energies accepted by a detector or gamma camera to create an image. Choosing it appropriately improves image quality by accepting desired signals and reducing scattered radiation.

### **Enfuvirtide**

An injectable HIV medicine that blocks fusion of HIV with human immune cells. It is used with other antiretroviral medicines for resistant infection; injection-site reactions and allergic reactions can occur.

### **Engagement**

Descent of the widest part of the baby's presenting head or body into the parent's pelvis. It may occur before labor or during labor and indicates that the baby has entered the birth passage.

### **Enhanced Elimination**

Medical methods used to remove an absorbed poison more quickly, such as dialysis or changing urine chemistry. They are used only for selected toxins because benefits, timing, and risks depend on the

substance and the person's condition.

### **Enhanced Pause**

An unusually long pause in breathing or another measured body signal. Its significance depends on the test and setting and may require interpretation with symptoms and the complete recording.

### **Enhanced Recovery After Surgery**

A coordinated care plan before, during, and after surgery that supports recovery through education, appropriate nutrition and fasting, pain control, early movement, early eating, and prevention of complications.

### **Enhancement Artifact**

An imaging effect in which tissue behind a fluid-filled or low-density structure appears brighter than it really is, especially on ultrasound. It is a technical artifact, not necessarily a sign of increased blood flow or tissue density.

### **Enhancer**

A stretch of DNA that increases activity of a gene when regulatory proteins bind to it. It may be far from the gene and can work in either direction, helping control when and where the gene is active.

### **Enkephalin**

A natural opioid-like peptide made by the body that activates mainly delta and mu opioid receptors. It helps regulate pain, stress, mood, movement, and other nervous-system functions.

#### **Alternate Names:**

Leu-Enkephalin; Met-Enkephalin

### **Enlarged Adenoids**

Overgrowth of immune tissue behind the nose that can block nasal breathing and cause mouth breathing, snoring, repeated ear-fluid problems, or sleep-disordered breathing. Symptoms and age determine whether treatment is needed.

### **Enlarged Fontanelle**

A soft spot on an infant's skull that is significantly

---

larger than normal for age, which may be associated with conditions such as congenital hypothyroidism, rickets, skeletal dysplasias, or increased intracranial pressure.

**Alternate Names:**

Large Fontanelle; Delayed Fontanelle Closure

**Enlarged Heart**

An increase in heart size caused by conditions such as high blood pressure, valve disease, heart-muscle disease, anemia, or athletic adaptation. It may cause no symptoms or lead to breathlessness, swelling, rhythm problems, or heart failure.

**Alternate Names:**

Cardiomegaly

**Enlarged Liver**

Alternate name; see *Hepatomegaly*.

**Alternate Names:**

Hepatomegaly; Hepatic Enlargement

**Enlarged Prostate**

Alternate name; see *Benign Prostatic Hyperplasia*.

**Enlarged Spleen**

Increase in spleen size caused by infection, liver or blood disease, inflammation, cancer, or congestion. It may cause fullness or pain and can increase rupture risk; evaluation considers symptoms, blood counts, and the underlying cause.

**Alternate Names:**

Splenomegaly; Splenic Enlargement

**Enophthalmos**

Abnormal backward sinking of an eyeball within the eye socket. It may follow injury, loss of tissue around the eye, dehydration, aging, or a nerve disorder and can cause unequal eye appearance or double vision.

**Enoxaparin**

A low-molecular-weight heparin that reduces clotting by enhancing antithrombin activity, especially against factor Xa. It prevents or treats blood clots but can cause bleeding, low platelets, or injection-site bruising.

**Enrichment**

Adding nutrients to food after processing has removed some of them. Enriched flour, for example, commonly has certain B vitamins and iron added back to help prevent nutritional deficiencies.

**ENT Physician**

An otolaryngologist, a physician who diagnoses and treats disorders of the ear, nose, throat, and related head and neck structures.

**Entamoeba**

A group of single-celled amoebas that live in the intestine or other tissues. Some species are harmless, while *Entamoeba histolytica* can cause diarrhea, intestinal ulcers, or liver abscesses.

**Entamoeba Coli**

A usually harmless amoeba found in the human intestine and spread by fecal contamination. Its presence in stool indicates exposure to contaminated food or water but usually does not cause illness or require treatment.

**Entamoeba Hartmanni**

A small, usually harmless intestinal amoeba spread through fecal contamination. It generally causes no disease; identification in stool mainly indicates possible exposure to contaminated food, water, or hands.

**Entamoeba Histolytica**

A parasite spread when contaminated food, water, or hands are swallowed. It can cause diarrhea, abdominal pain, and bloody stools and may invade the intestine or spread to the liver, causing an abscess.

**Entamoeba Polecki**

An intestinal amoeba that can infect people after fecal contamination, often through contact with animals or contaminated food and water. It is usually harmless but may occasionally be associated with diarrhea.

---

## Enteral

Relating to or delivered through the digestive tract. Enteral nutrition provides food and fluids by mouth or through a feeding tube into the stomach or intestine.

## Enteral Formula

A liquid nutrition preparation given by mouth or feeding tube. It may provide complete nutrition or selected nutrients and is chosen according to digestion, absorption, fluid needs, kidney or liver disease, and metabolic requirements.

## Enteral Nutrition

Delivery of food, fluids, and nutrients through the digestive tract, by mouth or feeding tube. It may be used when swallowing or eating is unsafe or inadequate; aspiration, diarrhea, blockage, and tube displacement are possible complications.

## Enteric

Relating to the intestine or digestive tract. The term can also describe organisms, infections, medicines, or coatings associated with the intestines.

## Enteric Bacteria

Bacteria that live in or affect the intestinal tract. Some are harmless normal residents, while others cause diarrhea or infections outside the intestine, such as urinary, wound, or bloodstream infection.

## Enteric Duplication Cyst

A birth-related cyst attached to or within the digestive tract, most often the small intestine. It may cause pain, blockage, bleeding, infection, or no symptoms and can require surgery.

## Enteric Fever

A serious infection caused by *Salmonella* Typhi or *Salmonella* Paratyphi, usually spread through food or water contaminated with human waste. It can cause gradually rising fever, headache, stomach pain, weakness, diarrhea or constipation, and sometimes a rash. Untreated illness can cause bleeding or a hole in the intestine; testing and antibiotic treatment are needed.

## Alternate Names:

Typhoid Fever; Paratyphoid Fever; Salmonella Enteric Infection

## Enteric Nervous System

The network of nerve cells within the digestive tract that helps control movement, secretion, blood flow, and communication with the brain and spinal cord.

## Enteric-Coated

Covered with a protective layer that resists stomach acid and dissolves in the small intestine. This can protect the stomach or protect a medicine from stomach acid; such tablets should not usually be crushed.

## Enteritis

Inflammation of the small intestine caused by infection, inflammatory disease, radiation, reduced blood flow, medicines, or other injury. It may cause abdominal pain, diarrhea, vomiting, fever, bleeding, or dehydration.

## Enterobacter

A group of Gram-negative bacteria that may live in the intestine but can cause healthcare-associated urinary, bloodstream, wound, respiratory, or abdominal infection, especially in vulnerable people.

## Enterobacter Aerogenes

A former name for *Klebsiella aerogenes*, a Gram-negative bacterium that can cause urinary, lung, wound, abdominal, or bloodstream infection, especially in healthcare settings. It may resist several antibiotics.

## Enterobacter Cloacae

A Gram-negative bacterium that can cause healthcare-associated urinary, bloodstream, lung, wound, or abdominal infection. It may resist several antibiotics, so laboratory susceptibility testing guides treatment.

## Enterobacter Hormaechei

A bacterium in the *Enterobacter cloacae* group that

---

can cause opportunistic healthcare-associated infection of blood, urine, lungs, wounds, or devices. Testing for antibiotic resistance is often important.

### **Enterobacteriaceae**

A family of Gram-negative bacteria that includes *Escherichia*, *Klebsiella*, *Enterobacter*, *Salmonella*, and *Shigella*. Some normally live in the intestine, while others cause intestinal, urinary, wound, or bloodstream infection.

### **Enterobacteriaceae Carbapenem Resistance**

Resistance in certain Gram-negative bacteria to carbapenem antibiotics, often caused by enzymes that destroy these medicines. Such infections can be difficult to treat and require laboratory testing, infection control, and specialist guidance.

### **Enterobiasis**

Infection with pinworms, small intestinal worms whose eggs spread easily between people. It commonly causes intense itching around the anus at night, disturbed sleep, and sometimes irritation in the genital area.

### **Enterobius**

A group of pinworms that live in the intestine. *Enterobius vermicularis* causes the common pinworm infection, which spreads when eggs are swallowed and often causes nighttime anal itching.

### **Enterobius Vermicularis**

A small intestinal worm commonly called the pinworm or threadworm. Adult females lay eggs around the anus, usually at night, causing itching and allowing infection to spread through contaminated hands or surfaces.

#### **Alternate Names:**

Pinworm; Threadworm; *Enterobius*

### **Enterocoele**

A pelvic organ prolapse in which a pouch of abdominal lining and sometimes small intestine bulges into the space between the vagina and rectum. It may cause pelvic pressure, a vaginal bulge, back or rectal

discomfort, or no symptoms.

#### **Alternate Names:**

Small Bowel Prolapse; Douglas Pouch Hernia

### **Enterochromaffin Cells**

Specialized enteroendocrine cells located throughout the mucosal epithelium of the gastrointestinal tract that synthesize, store, and release serotonin (5-hydroxytryptamine) and substance P to modulate gastrointestinal motility, secretion, and visceral pain perception.

#### **Alternate Names:**

EC Cells; Kulschitzky Cells

### **Enterochromaffin-Like Cells**

Specialized neuroendocrine cells situated within the oxyntic mucosa of the gastric fundus and body that synthesize and secrete histamine in response to gastrin and acetylcholine stimulation, directly stimulating parietal cell H<sup>+</sup>/K<sup>+</sup>-ATPase proton pumps.

#### **Alternate Names:**

ECL Cells

### **Enteroclysis**

An imaging test in which contrast material is delivered through a tube into the small intestine to look for inflammation, narrowing, tumors, or blockage. CT or MRI enterography is now more commonly used in many settings.

### **Enterococcaceae**

A family of bacteria that includes *Enterococcus* species. These bacteria may normally live in the intestine but can cause urinary, wound, bloodstream, or heart-valve infections when they enter protected body sites.

### **Enterococcal Infection**

An infection caused by *Enterococcus* bacteria, which normally live in the intestine but can cause urinary-tract, wound, abdominal, bloodstream, or heart-valve infections. Enterococcal infections are more common after hospitalization, surgery, antibiotic exposure, or use of medical devices. Some strains resist multiple antibiotics, so culture and susceptibility testing guide treatment.

---

**Alternate Names:**

Enterococcosis; Enterococcal Sepsis

**Enterococcus**

A group of bacteria normally found in the intestine that can cause urinary, wound, abdominal, bloodstream, or heart-valve infection. Some strains resist several antibiotics and require laboratory susceptibility testing.

**Enterococcus Durans**

A bacterium that may live in the bowel but can rarely cause infection in seriously ill or immunocompromised people. It may affect blood, urine, wounds, or other sites, and testing guides antibiotic treatment.

**Enterococcus Faecalis**

A common intestinal bacterium that can cause urinary, wound, abdominal, bloodstream, or heart-valve infection. Disease is more likely after surgery, hospitalization, medical-device use, or when immunity is weakened.

**Enterococcus Faecium**

An *Enterococcus* species that can cause healthcare-associated urinary, wound, abdominal, bloodstream, or heart-valve infection. Some strains resist vancomycin and other antibiotics, making laboratory testing essential.

**Enterocolitis**

Inflammation of the small intestine and colon caused by infection, immune disease, reduced blood flow, toxins, or other injury. It may cause abdominal pain, diarrhea, fever, vomiting, bleeding, dehydration, or severe illness.

**Enterocyte**

A cell lining the small intestine that digests and absorbs nutrients, water, and electrolytes. It also helps form the intestinal barrier and participates in immune defense.

**Enterocytozoon**

A group of tiny spore-forming parasites that can cause persistent diarrhea and intestinal disease, especially in people with weakened immunity. Some species can also affect other organs.

**Enteroendocrine Cell**

A specialized cell in the intestinal lining that senses food and releases hormones regulating digestion, appetite, gallbladder and pancreatic activity, and blood-glucose control.

**Enterohepatic Circulation**

Recycling of a substance from the liver into bile and the intestine, then back to the liver through the bloodstream. This can prolong the effects of some medicines and natural body chemicals.

**Enteromonas Hominis**

A usually harmless microorganism that may live in the intestine and appear in stool samples. It generally does not cause disease, so its presence should be interpreted with symptoms and other test results.

**Enteropathic**

Relating to disease of the intestine or caused by intestinal disease. The term may describe arthritis, skin changes, protein loss, or other problems occurring with inflammatory bowel disease or impaired intestinal absorption.

**Enteropathic Arthritis**

Inflammatory joint disease associated with inflammatory bowel disease such as Crohn disease or ulcerative colitis. It may affect the spine, joints between the spine and pelvis, or joints of the arms and legs.

**Enteropathy**

Any disease or disorder affecting the intestine. Causes include infection, inflammation, gluten sensitivity, poor blood flow, toxins, medicines, inherited conditions, or impaired absorption.

**Enteropathy-Associated T-Cell**



---

## Lymphoma

A rare, fast-growing cancer of immune cells in the intestine. It can be linked to long-standing celiac disease and may cause severe abdominal pain, weight loss, poor nutrient absorption, blockage, or a hole in the bowel.

### Alternate Names:

EATL; Enteropathy-Type Intestinal T-Cell Lymphoma

## Enteroscopy

Examination of the small intestine with a camera-equipped instrument. It can identify disease, obtain tissue samples, treat bleeding, or remove selected growths beyond the reach of routine upper endoscopy.

## Enterospasm

An involuntary tightening or contraction of the intestine that may cause cramping, abdominal pain, bloating, or altered movement of intestinal contents.

## Enterostomy

A surgically created opening that brings part of the intestine to the abdominal surface so stool or intestinal contents can leave the body or a feeding solution can enter.

## Enterostomy Procedure

Surgery that creates an opening from the intestine to the abdominal wall, such as an ileostomy or colostomy. It may divert stool or support feeding; bleeding, infection, blockage, and skin problems can occur.

## Enterotoxin

A toxin made by some microorganisms that acts on the intestinal lining and can cause vomiting, cramps, watery diarrhea, dehydration, or other intestinal illness.

## Enterovesical Fistula

An abnormal connection between the intestine and urinary bladder. It can cause repeated urinary infections, air or stool in the urine, abdominal pain,

or urine leaking into the intestine and may result from inflammation, cancer, or surgery.

## Enterovirulent E. Coli

A group of *Escherichia coli* strains that cause intestinal disease by making toxins, attaching to the bowel, or invading it. Illness may include watery or bloody diarrhea, cramps, vomiting, or dehydration.

## Enterovirus

A group of viruses spread mainly by respiratory secretions or contact with infected stool. Many infections are mild or cause no symptoms, but some cause hand-foot-and-mouth disease, rash, meningitis, heart inflammation, or severe illness in newborns.

## Enterovirus D68

A virus that usually causes a respiratory illness but can rarely cause severe pneumonia or sudden weakness of the limbs, especially in children. It spreads mainly through respiratory secretions and contaminated surfaces.

## Enterovirus Infection

Alternate name; see *Enterovirus*.

## Enthesis

The place where a tendon, ligament, or joint capsule attaches to bone. Repeated stress or inflammation there can cause localized pain and stiffness and occurs in several inflammatory joint diseases.

## Enthesitis

Inflammation where a tendon, ligament, or joint capsule attaches to bone. It can cause localized pain, tenderness, and stiffness and is common in spondyloarthritis, including psoriatic and inflammatory-bowel-disease-related arthritis.

## Enthesitis And Enthesopathy

Enthesitis is inflammation where a tendon, ligament, or fascia attaches to bone. Enthesopathy includes inflammation, wear, injury, or other disease at that site and may cause localized pain and stiffness.

---

**Alternate Names:**

Enthesopathy; Insertion Tendinopathy

**Enthesitis-Related Arthritis**

A type of arthritis that begins in childhood and causes inflammation where tendons or ligaments attach to bone. It often affects the legs, feet, or lower back and may also cause swelling in joints or pain where the heel tendon attaches.

**Alternate Names:**

ERA; Juvenile Spondyloarthritis

**Entorhinal Cortex**

A region of the brain that links the hippocampus with other memory and navigation areas. It helps process spatial information and new memories and is affected early in many cases of Alzheimer disease.

**Entrance Wound**

The wound where a projectile enters the body. Its appearance may help estimate the direction and distance of the shot, but findings vary with the weapon, clothing, body site, and circumstances and require forensic examination.

**Entropion**

Inward turning of an eyelid, usually the lower lid, so the eyelashes rub against the eye. It can cause irritation, tearing, pain, corneal scratches, infection, and reduced vision.

**Enucleation**

The surgical removal of an entire organ or well-defined encapsulated tumor in one piece without cutting into surrounding healthy tissue. In ophthalmology, it specifically refers to the complete surgical removal of the eyeball (globe) and optic nerve stump while preserving the eyelids, orbital fat, and extraocular muscles. It is performed for intraocular malignancies (e.g., retinoblastoma, uveal melanoma), severe irreparable ocular trauma, endophthalmitis, or a blind, painfully disfigured eye. An orbital implant is placed and attached to the eye muscles to support a prosthetic eye (artificial eye). In general surgery, it describes the clean “shelling out” of an intact benign tumor (such as a uterine

fibroid, insulinoma, or cyst).

**Alternate Names:**

Ocular Enucleation; Eye Enucleation; Surgical Enucleation

**Enucleation Procedure**

Alternate name; see *Enucleation*.

**Enuresis**

Repeated involuntary urination, usually during sleep in a child expected to remain dry. It may be primary or begin again after dryness; constipation, infection, diabetes, sleep problems, stress, or urinary disorders may contribute.

**Environment**

The physical, chemical, biological, and social surroundings in which a person lives or works. Environmental factors such as air, water, housing, noise, climate, and pollutants can influence health.

**Environmental Exposure**

Contact with a chemical, pollutant, organism, physical hazard, or other factor in the surroundings. Health effects depend on the substance, dose, route, duration, timing, and individual susceptibility.

**Environmental Health**

The field concerned with how surroundings affect human health. It includes air and water quality, food safety, housing, workplace hazards, climate, radiation, pollution, and prevention of harmful exposures.

**Environmental Illness**

Illness caused or worsened by an environmental exposure. Evaluation considers the timing, dose, setting, symptoms, other possible causes, and whether symptoms improve when exposure stops.

**Enzyme**

A protein, or occasionally an RNA molecule, that speeds a specific chemical reaction without being used up. Enzymes are essential for digestion, energy production, DNA processes, and many other cell

---

functions.

### **Enzyme Activity**

The rate at which an enzyme carries out a chemical reaction under specified conditions. It is affected by temperature, acidity, inhibitors, helper molecules, substrate amount, and enzyme concentration.

### **Enzyme Deficiency**

Reduced or absent activity of an enzyme, usually inherited or sometimes acquired. It can cause buildup of substances, lack of needed products, and characteristic metabolic or cellular disease.

### **Enzyme Induction**

Increased production or activity of an enzyme, often after exposure to a medicine or chemical. If liver drug-metabolizing enzymes are induced, some medicines may be broken down faster and become less effective.

### **Enzyme Inhibition**

Reduced activity of an enzyme, sometimes caused by a medicine or chemical. If a drug-metabolizing enzyme is inhibited, another medicine may accumulate and cause stronger effects or toxicity.

### **Enzyme Kinetics**

The study of how quickly enzyme reactions occur and how their rate changes with substrate amount, temperature, acidity, inhibitors, helper molecules, and enzyme concentration.

### **Enzyme Markers**

Enzymes measured in blood or another body fluid to help assess tissue injury or organ function. Abnormal levels can suggest liver, bile-duct, muscle, heart, bone, or pancreatic disease but are not a diagnosis by themselves.

### **Enzyme Precursors**

Inactive or weakly active proteins that become working enzymes after being cut or chemically changed. This allows the body to activate digestive and other enzymes at the correct place and time.

### **Enzyme Therapy**

Treatment that replaces, supplements, blocks, or uses an enzyme to correct a deficiency or alter a disease process. The purpose, dose, benefits, and risks depend on the specific enzyme and condition.

### **Enzyme-Linked Immunosorbent Assay**

A laboratory test that uses antibodies and an enzyme-generated color signal to detect or measure a target such as an antibody, antigen, hormone, or infection marker.

### **Eosinophil**

A type of white blood cell involved in allergic inflammation and defense against some parasites. Eosinophils normally make up a small fraction of blood cells; high levels can occur with allergy, infection, medicines, or other disease.

### **Eosinophilia**

An abnormally high number of eosinophils in the blood, commonly caused by allergy, parasites, medicines, autoimmune disease, or some blood disorders. Severe or persistent elevation can injure the heart, lungs, skin, nerves, or other organs.

### **Eosinophilia-Myalgia Syndrome**

A potentially severe illness characterized by intense muscle pain and increased eosinophils, historically associated with contaminated L-tryptophan supplements. It may also cause rash, swelling, nerve symptoms, breathing problems, or internal-organ injury.

### **Eosinophilic Asthma**

Asthma in which eosinophils, a type of white blood cell, contribute to airway inflammation. It may cause recurring wheezing, cough, chest tightness, or breathlessness and can be difficult to control in some people.

### **Eosinophilic Bronchitis**

A cause of long-lasting cough in which eosinophils inflame the airways without the variable narrowing typical of asthma. The cough may resemble asthma or reflux and needs assessment to identify the cause.

---

## Eosinophilic Esophagitis

A chronic immune-related disease in which eosinophils inflame the esophagus. It may cause difficulty swallowing, food becoming stuck, chest or upper-abdominal pain, reflux-like symptoms, and narrowing of the esophagus.

## Eosinophilic Esophagitis Pathology

Esophageal mucosal disease with increased intraepithelial eosinophils, basal-zone hyperplasia, spongiosis, and sometimes lamina-propria fibrosis. Histologic findings are interpreted with symptoms and endoscopic features.

## Eosinophilic Fasciitis

A rare inflammatory disease causing painful swelling and hardening of the deeper skin and tissue around muscles, usually in the arms or legs. Blood eosinophils may be high, and joint stiffness or limited movement can develop.

### Alternate Names:

Shulman Syndrome; Diffuse Fasciitis with Eosinophilia

## Eosinophilic Gastroenteritis

A rare immune-related disease in which eosinophils inflame the stomach or intestines. Symptoms may include abdominal pain, nausea, vomiting, diarrhea, bleeding, poor absorption, or weight loss.

## Eosinophilic Gastrointestinal Disorders

Conditions in which eosinophils, a type of white blood cell, build up in one or more parts of the digestive tract and cause inflammation. Symptoms may include pain, vomiting, swallowing difficulty, diarrhea, or poor growth; diagnosis usually requires endoscopy and biopsy.

## Eosinophilic Granuloma

A localized form of Langerhans-cell disease in which abnormal immune cells collect, often in bone or skin. Bone lesions can cause pain, swelling, or fractures and may require imaging and tissue examination.

## Eosinophilic Granuloma (Pulmonary

## Histiocytosis X)

Historical or imprecise label; see *Pulmonary Langerhans Cell Histiocytosis*.

## Eosinophilic Granulomatosis With Polyangiitis

A rare disease in which inflamed blood vessels occur with asthma and high levels of eosinophils, a type of white blood cell. It can damage the lungs, skin, nerves, heart, kidneys, or digestive organs.

### Alternate Names:

Allergic Granulomatosis Allergic Granulomatosis and Angiitis

## Eosinophilic Pneumonia Pathology

Lung inflammation rich in eosinophils involving alveolar spaces, interstitium, or both. Additional findings may include organizing pneumonia, edema, and eosinophilic infiltrates around vessels or airways.

## Eosinophils

White blood cells that help control allergic inflammation and defend against some parasites. High levels may occur with allergies, asthma, infection, medicines, autoimmune disease, or blood disorders; low levels usually have little significance.

### Alternate Names:

Eosinophil Granulocytes

## Ependyma

A specialized layer of cells lining the brain's fluid-filled ventricles and the spinal cord's central canal. It helps separate nervous tissue from cerebrospinal fluid and supports movement of that fluid.

## Ependymal Cell

A supporting cell lining the brain ventricles and spinal cord central canal. It helps maintain the cerebrospinal-fluid boundary and, in some areas, helps move or produce the fluid.

## Ependymal Tumor

A growth arising from cells lining the brain ventricles or spinal canal. It may block cerebrospinal-fluid flow and cause headache, vomiting, balance problems,

---

weakness, or hydrocephalus; treatment depends on its type and extent.

### **Ependymoblastoma**

A rare, aggressive brain tumor of childhood made from very immature cells. It may cause headache, vomiting, seizures, or changes in movement or development; tissue and genetic testing guide treatment and outlook.

### **Ependymoma**

A tumor arising from cells lining the brain ventricles or spinal canal. It may block fluid flow and cause headache, vomiting, seizures, balance problems, weakness, or hydrocephalus, especially in children.

### **Ephedrine**

A stimulant that activates the sympathetic nervous system and can widen airways and raise blood pressure and heart rate. It has limited medical uses and may cause palpitations, insomnia, anxiety, high blood pressure, or misuse.

### **Ephedrinum**

Another name for ephedrine, a stimulant that can increase heart rate, blood pressure, alertness, and airway diameter. Excessive use may cause dangerous heart-rhythm, blood-pressure, or nervous-system effects.

### **Ephelides**

Small, flat freckles caused by increased pigment production, often inherited. They become darker with sunlight and usually fade when exposure decreases; sun protection helps prevent further pigment change and skin damage.

### **Ephrins**

Proteins on the surface of cells that help guide cell position, nerve growth, blood-vessel development, and tissue organization. Abnormal ephrin signaling can contribute to some diseases and cancers.

### **Epiblast**

A layer of cells in the early embryo that develops into the three main cell layers of the body and ultimately forms most tissues and organs.

### **Epicanthal Folds**

Skin folds of the upper eyelid that partly cover the inner corner of the eye. They are normal in many people but can also occur with some chromosome or developmental conditions.

### **Epicardium**

The thin outer layer of the heart wall and the inner layer of its surrounding protective sac. It contains connective tissue, fat, blood vessels, and nerves and helps reduce friction as the heart beats.

### **Epichlorohydrin**

An industrial chemical used to make resins and other materials. Exposure can irritate or burn the skin, eyes, and airways; swallowing or substantial exposure can cause serious poisoning and requires urgent medical advice.

### **Epicondyle**

A bony projection beside a rounded joint surface where muscles and ligaments attach. Repeated strain or injury around an elbow epicondyle can cause localized pain and weakness.

### **Epicondylitis**

Pain and tenderness where a tendon attaches near an epicondyle, usually from repeated strain rather than infection. At the elbow it can weaken gripping and wrist movement and is commonly called tennis or golfer's elbow.

#### **Alternate Names:**

Tennis Elbow; Golfer's Elbow

### **Epicondylitis (Tennis Elbow)**

Overuse-related pain at the outer elbow, usually involving tendons that attach forearm muscles to the lateral epicondyle. Despite the name, it is generally tendinopathy rather than acute inflammation and may cause pain with gripping or wrist extension.

---

## Epidemic

An increase in disease cases above the number normally expected in a population, place, or period. An epidemic may result from infection, environmental exposure, behavior, or another shared cause.

## Epidemic Cerebrospinal Meningitis

Historical name; see *Meningococcal Meningitis*.

## Epidemic Cholera

Alternate name; see *Cholera*.

### Alternate Names:

EKC; Adenoviral Keratoconjunctivitis; Shipyard Eye

## Epidemic Keratoconjunctivitis

A very contagious viral infection of the eye. It can cause red eyes, pain, watering, swollen eyelids, light sensitivity, blurred vision, and swollen glands near the ear; symptoms may last for weeks.

## Epidemiologic Method

A structured way to measure how often a health problem occurs and to study whether exposures, behaviors, or other factors are associated with it in a population. Results guide prevention and policy.

## Epidemiologic Studies

Research that examines the distribution, causes, risk factors, prevention, or outcomes of health events in populations.

## Epidemiological Study

A study of health and disease in groups of people. It may measure how often a condition occurs, compare possible causes or exposures, or examine outcomes over time to help guide prevention and healthcare planning.

### Alternate Names:

Epidemiologic Study

## Epidemiologist

A scientist or public-health professional who studies the distribution, causes, risk factors, and prevention of health conditions in populations.

## Epidemiology

The study of how often diseases and other health events occur, who is affected, and which factors influence them. It uses population data to identify causes, guide prevention, investigate outbreaks, and evaluate healthcare.

## Epidermal

Relating to the epidermis, the outermost layer of skin. An epidermal problem affects this protective surface layer rather than the deeper skin tissues.

## Epidermal Acanthosis

Thickening of the epidermis caused by increased growth of its cells, seen in conditions such as psoriasis, chronic rubbing, and some inflammatory or genetic skin disorders.

## Epidermal Cells

Cells forming the outer layer of skin, chiefly keratinocytes, which create a protective barrier; melanocytes, immune cells, and touch-associated cells are also present.

## Epidermal Growth Factor

A signaling protein that binds the epidermal-growth-factor receptor and promotes cell growth, survival, and tissue repair; abnormal signaling can contribute to cancer.

## Epidermal Inclusion Cyst

A common, usually harmless lump beneath the skin filled with keratin, a protein found in skin. It often has a small central opening and may become swollen, painful, infected, or recur if its wall remains.

## Epidermal Layers

The epidermis has several layers of cells that make and protect the skin. From deepest to most superficial they are basal, spinous, granular, clear, and horny layers; the clear layer is mainly in the palms and soles.

## Epidermal Nevus

A usually benign, often present-from-birth or child-



---

hood skin lesion caused by an overgrowth of epidermal cells. It commonly appears as a localized, linear, or warty thickened patch and may occur with other developmental abnormalities.

### Epidermal Nevus Syndrome

A group of mosaic developmental disorders in which extensive or systematized epidermal nevi occur with abnormalities of the brain, eyes, skeleton, limbs, or other organs. Evaluation is guided by the distribution of the nevi and associated neurologic, ocular, or musculoskeletal findings.

### Epidermis

The outer layer of skin that forms a protective barrier against injury, germs, and water loss. It contains cells that make keratin and pigment, as well as immune and touch-sensing cells, and it is replaced continuously.

### Epidermodysplasia Verruciformis

A rare inherited or acquired disorder of cutaneous immunity in which susceptibility to specific beta-human papillomaviruses causes persistent flat wart-like or pityriasis-versicolor-like lesions. Sun-exposed lesions can develop actinic keratosis and cutaneous squamous-cell carcinoma, so lifelong skin surveillance and sun protection are important.

### Epidermoid Cyst

Alternate index label; see *Epidermoid Cysts*.

### Epidermoid Cysts

Slow-growing, usually harmless sacs beneath the skin filled with keratin. They often appear as movable lumps and may become painful, red, swollen, or infected if they rupture.

#### Alternate Names:

Epidermal Inclusion Cysts; Infundibular Cysts

### Epidermolysis Bullosa

A group of inherited disorders in which fragile skin blisters or tears after minor rubbing or injury because of defects in proteins that join skin layers. Severity varies; some forms heal without scars, while others cause scarring, tight joints, poor nutrition,

infection, or increased skin-cancer risk.

#### Alternate Names:

Bullosa Hereditaria; Hereditary Epidermolysis Bullosa

### Epidermolysis Bullosa Acquisita

An uncommon autoimmune blistering disease in which antibodies target type VII collagen, a protein anchoring the epidermis to the dermis. Tense blisters and erosions develop after minor trauma, often heal with scarring or milia, and may affect mucous membranes.

### Epidermolytic Ichthyosis

An inherited keratinization disorder, usually caused by variants in KRT1 or KRT10, in which newborns may have widespread redness and superficial blistering that later evolves into thick, scaling skin and sometimes palmoplantar keratoderma.

#### Alternate Names:

Epidermolytic Hyperkeratosis; Bullous Congenital Ichthyosiform Erythroderma

### Epidermophyton

A group of fungi that grow in keratin, a tough protein in skin, hair, and nails. They can cause ringworm infections, including itchy, scaly skin disease and fungal infection of the nails, but do not usually infect living hair shafts.

### Epidermophyton Floccosum

A dermatophyte fungus that commonly causes tinea of the body, groin, and feet and can infect nails.

### Epididymal Cyst

A usually benign, fluid-filled cyst arising in the epididymis. It may be painless and may require treatment only if it causes significant symptoms or complications.

### Epididymal Neoplasm

A benign or malignant growth arising in the epididymis; most palpable epididymal masses are benign, but persistent or enlarging lesions require evaluation.

---

## Epididymides

The paired coiled ducts on the posterior surface of the testes where sperm mature and are stored before entering the vasa deferentia.

### Alternate Names:

Epididymal Ducts; Convolutd Seminiferous Ducts

## Epididymis

A tightly coiled tube along the back of each testicle where sperm mature, gain the ability to move, and are stored before passing into the vas deferens. It has a head, body, and tail.

## Epididymitis

Inflammation of the epididymis, usually caused by infection. It commonly causes gradually developing pain and swelling on one side of the scrotum. Causes include sexually transmitted infections, urinary bacteria, urine-flow problems, and less commonly injury or medicines.

## Epididymitis And Orchitis

Alternate name for epididymo-orchitis, inflammation of the epididymis and testicle.

## Epididymo-Orchitis

Inflammation of the epididymis and testicle, usually caused by an infection. It can cause one-sided scrotal pain, swelling, redness, painful urination, and fever. The cause may be a sexually transmitted infection or bacteria from the urinary tract, depending on the person and situation. Sudden severe pain needs emergency assessment to rule out testicular torsion; treatment depends on the cause.

### Alternate Names:

Epididymo-orchitis; Testicular and Epididymal Inflammation; Orchiepididymitis

## Epidural

Pain relief or regional anesthesia given through a small catheter placed in the lower back, near the spinal nerves. It is commonly used during labor and some operations; possible effects include temporary low blood pressure, itching, headache, or incomplete relief.

## Epidural Abscess

A pocket of pus between the tough outer covering of the brain or spinal cord and nearby bone. It can compress the brain, spinal cord, or nerves, causing fever, severe pain, weakness, numbness, or loss of bladder control and requires emergency treatment.

## Epidural Analgesia

Pain relief produced by delivering numbing medicine, sometimes with an opioid, through a small catheter into the space around the spinal nerves. It may be used during labor or surgery while the person remains awake. Low blood pressure, itching, headache, infection, or incomplete relief can occur.

## Epidural Analgesia For Labor

Pain relief during labor provided through a small catheter placed in the lower back. Numbing medicine, sometimes with an opioid, is given gradually to reduce pain while allowing the laboring person to remain awake and participate in birth. Temporary low blood pressure, itching, headache, or incomplete relief can occur.

## Epidural Anesthesia

Regional anesthesia produced by injecting numbing medicine, sometimes with an opioid, into the space around the spinal nerves. It blocks pain from part of the body while the person stays awake; low blood pressure, headache, infection, or nerve injury can rarely occur.

### Alternate Names:

Epidural Anaesthesia

## Epidural Blood Patch

A procedure that injects a small amount of the person's own blood into the space around the spinal nerves to seal a leak after spinal anesthesia or an accidental puncture. It usually treats a severe positional headache; back pain, infection, or another puncture can occur.

## Epidural Hematoma

A collection of blood between the skull and the brain's tough outer covering, usually after a head injury. It can rapidly compress the brain, caus-

---

ing headache, vomiting, confusion, weakness, loss of consciousness, or death and requires emergency treatment.

### **Epidural Neoplasm**

A tumor in the space outside the dura surrounding the spinal cord or brain; spinal epidural tumors can compress nerves or the spinal cord and cause pain, weakness, or sensory loss.

### **Epidural Space**

The space between the tough outer covering of the spinal canal and the canal wall. Medicines injected there can reduce pain during childbirth, surgery, or some pain treatments.

### **Epidural Steroid Injection**

A minimally invasive outpatient spine procedure used to relieve severe radiating back, neck, arm, or leg pain (such as sciatica or lumbar radiculopathy) caused by an inflamed or compressed spinal nerve. Under live X-ray guidance (fluoroscopy) and local anesthesia, a pain specialist or spine clinician carefully directs a thin needle into the epidural space surrounding the spinal nerves and dural sac, injects a tiny amount of contrast dye to verify accurate placement, and administers a powerful long-acting corticosteroid combined with a local anesthetic. The medication coats the irritated nerve roots, flushing away inflammatory chemicals and dramatically reducing swelling caused by herniated discs, spinal stenosis, degenerative disc disease, or bone spurs. Epidural steroid injections can be performed through translaminar, transforaminal, or caudal routes, offering substantial pain relief that allows patients to engage in active physical therapy and frequently avoid spinal surgery.

#### **Alternate Names:**

ESI; Epidural Injections for Back Pain; Translaminar Epidural Injection; Transforaminal Epidural Injection; Caudal Epidural Steroid Injection

### **Epigastric**

Relating to the upper middle area of the abdomen between the ribs. Pain there may arise from the stomach, esophagus, pancreas, gallbladder, intestine,

or other organs.

### **Epigastric Artery**

An artery supplying the front abdominal wall. The superior and inferior epigastric arteries provide blood to abdominal muscles and skin and are important landmarks during some procedures.

### **Epigastric Pain**

Pain or discomfort in the upper middle abdomen, below the ribs and above the navel. Causes include indigestion, ulcer, reflux, gallbladder disease, pancreatitis, and other conditions; severe or persistent pain needs assessment.

### **Epigastric Region**

The upper middle section of the abdomen between the rib margins and above the navel. It overlies parts of the stomach, duodenum, pancreas, and liver; pain there can have several digestive or vascular causes.

### **Epigastrium**

The upper middle part of the abdomen, below the breastbone and between the ribs. It overlies parts of the stomach, liver, pancreas, and small intestine; pain there has many possible causes.

### **Epigenetics**

Changes in how genes are switched on or off without changing the DNA sequence itself. Chemical tags and DNA-packaging changes can be influenced by development, environment, and illness and may sometimes be reversible.

### **Epigenome**

The collection of chemical tags and DNA-packaging features that help control which genes are active in a cell without changing the DNA sequence.

### **Epigenomics**

The study of chemical and structural changes that control which genes are active across the genome without changing the DNA sequence. These changes can influence development, disease, and treatment response.

---

## Epiglottic Vallecula

A shallow mucosal depression situated in the glossoepiglottic space between the base of the tongue and the anterior surface of the epiglottis, bounded by median and lateral glossoepiglottic folds, serving as the anatomical seat for the tip of a curved Macintosh laryngoscope blade during endotracheal intubation.

### Alternate Names:

Vallecula Epiglottica; Vallecular Fossa

## Epiglottis

A leaf-shaped flap of flexible tissue at the base of the tongue that helps direct food and liquid into the esophagus during swallowing and away from the airway. Swelling can rapidly obstruct breathing.

## Epiglottitis

A potentially life-threatening infection or inflammation causing swelling around the epiglottis and upper airway. Fever, severe sore throat, drooling, trouble swallowing, noisy breathing, or sitting forward requires emergency care; the throat should not be examined forcefully.

## Epilepsia Partialis Continua

Repeated or nearly continuous twitching in one part of the body, often an arm, leg, or side of the face. Awareness may remain normal, but the condition can last for hours or longer and needs urgent medical assessment.

### Alternate Names:

EPC; Kojewnikoff Syndrome

## Epilepsy

A chronic brain disorder that causes a lasting tendency to have recurrent unprovoked seizures. A seizure is a brief episode caused by excessive, abnormal electrical activity in the brain and may change movement, sensation, awareness, behavior, or consciousness. Seizures may begin in one area of the brain (focal seizures) or involve both sides from the start (generalized seizures). Causes include genetic conditions, brain injury or malformation, stroke, infection, metabolic or immune disease, or no known cause. A single seizure, especially one triggered by

a temporary illness or abnormal body chemistry, does not always mean epilepsy.

## Epilepsy and Seizures

Alternate name; see *Epilepsy*.

### Alternate Names:

Seizure Disorder

## Epilepsy Surgery

Surgery or a procedure used for selected people whose seizures continue despite appropriate medicines and arise from an area that can be safely treated. Detailed testing identifies the seizure source and estimates benefits and risks.

## Epileptic Spasms

Brief seizures, most common in infants, in which the head, body, arms, or legs suddenly bend or stiffen. They often occur in clusters and need prompt assessment because they can affect development.

### Alternate Names:

Infantile Spasms

## Epileptiform

Resembling a seizure or the electrical pattern associated with seizure activity. It may describe movements or an EEG finding but does not by itself prove that a person has epilepsy.

## Epimysium

A substantial sheath of dense irregular collagenous connective tissue that completely envelops the entire exterior surface of a skeletal muscle, continuous with tendons, muscular fascial planes, and the periosteum of bone.

## Epinephrin

Alternate spelling; see *Epinephrine*.

## Epinephrine

A hormone and neurotransmitter released during stress that increases heart activity, opens the airways, and narrows many blood vessels. As a medicine, it treats anaphylaxis and selected cardiac or breathing emergencies.

---

**Alternate Names:**

Adrenaline

**Epineurium**

The thick, outermost fibrous connective tissue sheath that encloses an entire peripheral nerve, surrounding multiple nerve fascicles, longitudinal blood vessels (vasa nervorum), and adipose tissue to provide tensile mechanical protection.

**Epiphora**

Excessive tearing caused by blocked tear drainage or irritation that triggers extra tears. Causes include dry eye, allergy, infection, eyelid position problems, and blockage of the tear duct.

**Epiphyseal Plate**

The growth plate, a layer of cartilage near the end of a growing bone that allows the bone to lengthen. It gradually closes after puberty; injury can cause pain, deformity, or unequal growth.

**Alternate Names:**

Growth Plate; Physis

**Epiphyseal Plate Disorders**

Conditions affecting a growing bone's growth plate, including fractures, inflammation, infection, abnormal blood supply, premature closure, and disorders in which an epiphysis shifts or separates. They may cause pain, swelling, limping, deformity, or unequal limb growth.

**Epiphysis**

The expanded end of a long bone, usually covered by smooth cartilage where it forms a joint. It contains mostly spongy bone and helps determine the shape and movement of the joint.

**Epiphysitis**

Inflammation or injury involving a growth plate or the area near the end of a growing bone. It may cause localized pain, swelling, or altered growth, and the cause depends on age and the affected bone.

**Epiploic Foramen**

The narrow anatomical communication aperture connecting the greater peritoneal sac with the lesser sac (omental bursa), bounded anteriorly by the hepatoduodenal ligament (containing the portal triad), posteriorly by the inferior vena cava, superiorly by the caudate lobe of the liver, and inferiorly by the first part of the duodenum.

**Alternate Names:**

Foramen of Winslow

**Epipodophyllotoxin**

A group of cancer medicines derived from a plant compound that damage DNA by blocking an enzyme needed for cell division. They treat selected cancers but may cause low blood counts, nausea, hair loss, infection, or rarely a later blood cancer.

**Epi-retinal Membrane**

A thin layer of scar-like tissue growing on the retina, usually over the macula. It may cause blurred or distorted central vision and sometimes needs surgery if visual impairment is significant.

**Epirubicin**

An anthracycline cancer medicine that damages DNA and blocks cell division. Important risks include low blood counts, infection, nausea, hair loss, tissue injury after leakage, and dose-related heart damage.

**Epirubicin Hydrochloride**

The salt form of epirubicin, an anthracycline cancer medicine that damages tumor-cell DNA. It is used for selected cancers but can cause low blood counts, tissue injury after leakage, and dose-related heart damage.

**Episcleritis**

Usually short-lived inflammation of the thin tissue over the white of the eye, causing a localized red patch and mild tenderness or discomfort. Severe pain, light sensitivity, or vision loss suggests deeper eye disease.



---

## Episiotomy

A cut made in the tissue between the vagina and anus during vaginal birth to enlarge the opening. Routine use is not recommended; selected use may help urgent or assisted delivery but can cause pain, bleeding, infection, or deeper tearing.

## Episiotomy Repair

Closing an episiotomy or related tear with stitches after birth. The area is examined to identify deeper injury, and care includes pain relief, hygiene, monitoring for infection, and follow-up of healing and pelvic-floor function.

## Episodic Ataxia

An inherited condition causing repeated attacks of poor coordination and balance. Episodes may last seconds to hours and can include dizziness, slurred speech, muscle tightening, or abnormal eye movements, with improvement between attacks.

## Epispadias

A birth defect in which the urethral opening is on the upper surface of the penis or in an unusual position in a female. It may affect urine control and is sometimes associated with bladder-exstrophy conditions.

## Epistasis

A genetic interaction in which one gene changes or masks the effect of another gene. This can alter inherited traits and complicate how a disease risk or laboratory finding is predicted from genetic testing.

## Epistaxis

Medical term; see *Nosebleeds*.

## Epithalamic

Relating to the epithalamus, a small region near the center of the brain that includes the pineal gland and structures involved in sleep timing, hormonal rhythms, and responses to reward or stress.

## Epithalamus

A dorsal posterior division of the diencephalon com-

prising the pineal gland (epiphysis cerebri), the habenular nuclei, the stria medullaris thalami, and the posterior commissure, involved in circadian melatonin rhythmicity and limbic-midbrain connections.

## Epithelial

Relating to cells that cover the skin, line body passages and organs, or form glands. These cells protect surfaces and may absorb substances, release fluids, or help detect changes.

## Epithelial Cells

Cells that cover the skin, line body passages and organs, and form many glands. They create a protective barrier and may absorb substances, release fluids, move material, or sense changes in their surroundings.

### Alternate Names:

Epithelial Tissue Cells

## Epithelial Cyst

A usually benign, enclosed sac lined by surface-type cells and filled with fluid, mucus, or a skin protein called keratin. It may enlarge, become painful, or become infected and sometimes requires removal.

## Epithelial Tissue

Tissue made of closely packed cells that covers the skin, lines internal spaces and tubes, and forms glands. It provides protection and may absorb, secrete, transport, or exchange substances.

## Epithelioid Cell

A rounded cell that resembles a surface-lining cell. The term often describes an activated immune cell in a granuloma, but it can also describe the appearance of some tumors.

## Epithelioid Histiocyte

An activated macrophage that has undergone structural transformation under the influence of interferon-gamma in chronic granulomatous inflammation, acquiring abundant pale eosinophilic cytoplasm, elongated slipper-shaped nuclei, and indistinct cell borders resembling epithelial cells.



---

### **Epithelioid Sarcoma**

A rare cancer of connective or soft tissue, often beginning in the skin or deeper tissues of the hands, forearms, feet, or lower limbs. It may recur or spread and requires specialist treatment.

### **Epithelioid Trophoblastic Tumor**

A rare pregnancy-related cancer arising from cells that normally help form the placenta. It may appear months or years after pregnancy, cause abnormal bleeding, and requires specialist cancer treatment.

### **Epithelioma**

An older or broad term for a tumor arising from epithelial cells; the specific diagnosis should identify whether it is benign, precancerous, or malignant.

### **Epitheliopathy**

Disease or injury affecting epithelial cells, which cover body surfaces and line organs. The term is nonspecific and may describe conditions of the cornea, skin, gastrointestinal tract, or other epithelial tissues.

### **Epithelium**

A layer or sheet of closely packed cells covering the skin, lining body spaces and tubes, or forming glands. It protects surfaces and may absorb substances, release fluids, transport material, or allow exchange.

### **Epitope**

The small, specific part of an antigen that is recognized by an antibody or immune-cell receptor. Recognition can trigger an immune response; epitopes are important in infections, vaccines, allergies, and antibody-based treatments.

### **Epitope Mapping**

The process of identifying an epitope, the small part of an antigen recognized by an antibody or specific immune-cell receptor. It helps researchers understand immune responses and develop vaccines, diagnostic tests, and antibody-based treatments.

### **Eplerenone**

A medicine that blocks aldosterone, a hormone that promotes salt retention and heart remodeling. It treats high blood pressure and selected heart-failure patients; kidney function and blood potassium require monitoring.

### **Epley Maneuver**

A sequence of head and body movements that returns displaced inner-ear particles to their usual position. It can relieve brief spinning triggered by head movement in posterior-canal positional vertigo.

### **EPO Test**

A blood test measuring erythropoietin, a kidney-made hormone that stimulates red-blood-cell production. It may help investigate anemia or an abnormally high red-cell count alongside blood counts and other tests.

### **Epoetin Alfa**

A laboratory-made form of erythropoietin that stimulates the bone marrow to produce red blood cells. It treats selected anemias, but can raise blood pressure and the risk of blood clots if hemoglobin rises too high.

### **Epoetin Beta**

A laboratory-made form of erythropoietin used for selected anemias, including some caused by chronic kidney disease. Treatment is individualized and requires monitoring because it may cause high blood pressure or blood clots.

### **Eponym**

A medical name derived from a person's name or another proper name, such as the name of a disease, sign, procedure, or body structure. Eponyms may be replaced by descriptive terms to improve clarity.

### **Epoprostenol**

A short-acting medicine that widens blood vessels in the lungs and reduces platelet clumping. Continuous intravenous infusion treats severe pulmonary arterial hypertension; interruption can cause rapid worsening.

---

## Epoxide Hydrolase

An enzyme that changes reactive chemicals called epoxides into less reactive substances. It helps process some medicines, toxins, and natural fats and can influence their effects and the risk of tissue injury.

## Epoxy Compounds

Chemicals containing a reactive group used in adhesives, coatings, plastics, and other materials. Uncured components can irritate the skin and eyes, cause allergic skin inflammation, or irritate the airways; workplace exposure may cause asthma.

## Epsom Salts

Magnesium sulfate crystals used in some baths and, occasionally, as a laxative. Drinking them can cause diarrhea or magnesium poisoning, especially in people with kidney disease, so medical advice is important.

## Epstein Pearls

Small, white, harmless cysts containing keratin that appear on a newborn's gums or roof of the mouth. They usually disappear within weeks without treatment and should not be squeezed.

## Epstein-Barr Virus

A common virus spread mainly through saliva. It can cause infectious mononucleosis and remains dormant in the body; it is also linked to some lymphomas, other cancers, and immune disorders.

## Epstein-Barr Virus Antibody Test

A blood test that measures antibodies produced against Epstein-Barr virus. The pattern of antibodies, symptoms, and timing can help distinguish recent infection from past infection, but results need clinical interpretation.

## Eptifibatide

An injectable antiplatelet medicine used during selected urgent coronary treatments. It prevents platelets from clumping by blocking a platelet-binding receptor, but can cause serious bleeding and requires blood-count and kidney monitoring.

## Epulis

A localized growth arising from the gum or tissues around a tooth. Most are reactive and benign, but persistent, painful, bleeding, or enlarging growths need dental examination and sometimes biopsy.

## Equilibrium Receptors

Sensory cells in the inner ear that detect head movement and position. Signals from them, the eyes, and muscles help the brain maintain balance and coordinate posture and eye movements.

## Equilin

An estrogen hormone found mainly in pregnant mares and included in some mixtures of conjugated estrogens. Estrogen treatment can increase the risk of blood clots and other adverse effects and requires individualized medical advice.

## Equine Morbillivirus

A virus related to measles that has caused severe respiratory or neurologic disease in horses and can infect other mammals. Suspected cases require veterinary investigation and infection-control measures.

## Equinovarus Deformity

A deformity in which the foot points downward and turns inward. It may be present at birth or develop from muscle, nerve, joint, or bone problems, limiting walking and requiring cause-specific treatment.

## Equipment Safety

Measures that reduce harm from medical devices by using the correct equipment, checking it before use, cleaning it properly, maintaining it, and training users. Faulty equipment should be removed from service.

## Equivalence Trial

A clinical study designed to determine whether a new treatment and an established treatment have effects within a predefined clinically acceptable range of one another. The range must be set before results are analyzed.

---

## Eradication

Complete removal of an infection or disease from a person or population. For an individual infection, eradication may require a full treatment course and a follow-up test to confirm that the organism is gone.

## Erb Palsy

A peripheral obstetric or traumatic traction injury to the upper trunk of the brachial plexus involving spinal nerve roots C5 and C6 (and sometimes C7), causing paralysis of the deltoid, supraspinatus, infraspinatus, and biceps brachii muscles, resulting in an adducted, internally rotated shoulder and extended, pronated forearm ('waiter's tip' posture).

### Alternate Names:

Erb-Duchenne Palsy; Upper Trunk Brachial Plexus Palsy

## Erb's Palsy

Weakness or paralysis of the shoulder and arm caused by injury to the upper nerves of the brachial plexus, often during difficult birth. The arm may hang inward with the elbow straight and palm facing backward.

## ErbB Receptors

A family of cell-surface proteins that receive growth signals and regulate cell division, survival, and specialization. Abnormal activity in some members can promote cancers and may guide targeted treatment.

## Erbium

A chemical element used in specialized lasers and materials. Erbium lasers can precisely remove or remodel selected skin and dental tissue; treatment may cause pain, burns, pigment changes, or scarring.

## ERCP

A procedure using a flexible camera and X-ray guidance to diagnose or treat blocked, narrowed, or leaking bile or pancreatic ducts. It can remove stones or place a stent, but may cause pancreatitis, bleeding, infection, or perforation.

## Erdheim-Chester Disease

A rare non-Langerhans cell histiocytic neoplasm driven by activating kinase mutations in the MAPK pathway (predominantly BRAF V600E), characterized pathologically by lipid-laden foamy histiocyte infiltration causing symmetric diaphyseal osteosclerosis of long bones, periaortic fibrosis ('coated aorta'), and perinephric infiltration ('hairy kidney').

### Alternate Names:

ECD; Non-Langerhans Histiocytosis

## Erectile Disorder

Alternate name; see *Erectile Dysfunction*.

## Erectile Dysfunction

Repeated difficulty getting or keeping an erection firm enough for satisfactory sexual activity. It may result from blood-vessel disease, diabetes, nerve or hormonal disorders, medicines, penile conditions, anxiety, depression, or mixed causes and can signal cardiovascular risk.

## Erection

Temporary enlargement and stiffening of the penis when blood fills erectile tissue and drainage through veins decreases. It can occur with sexual stimulation, during sleep, or spontaneously and normally subsides afterward.

## Erector Spinae

A group of muscles running along both sides of the spine. They straighten and stabilize the back and help bend the spine sideways during standing, walking, and lifting.

### Alternate Names:

Sacrospinalis Muscle; Spinalis and Longissimus Complex

## Ergocalciferol

Vitamin D2, a form of vitamin D used to prevent or treat deficiency. It helps the body absorb calcium and maintain bones, but excessive doses can cause high blood calcium, nausea, confusion, or kidney injury.

---

## Ergolines

A family of compounds related to chemicals made by ergot fungi. Some affect serotonin, dopamine, or adrenaline receptors and are used or studied for migraine, hormone disorders, or neurologic conditions.

## Ergometrine

A medicine that contracts the uterus and narrows some blood vessels. It may help control severe bleeding after childbirth, but can dangerously raise blood pressure and is unsuitable for some people with hypertension or heart disease.

## Ergonomics

Designing work, tools, furniture, and environments to match the body's abilities and limits. Good ergonomics can reduce strain, awkward posture, repetitive-injury risk, fatigue, and errors.

## Ergosterol

A fat-like substance that helps maintain fungal cell membranes. Many antifungal medicines damage fungi by binding ergosterol or blocking its production; human cells use cholesterol instead.

## Ergot

A fungus that can infect grains such as rye and produce toxic chemicals. Eating contaminated grain or taking excessive ergot-derived medicines can cause severe blood-vessel narrowing, neurologic symptoms, or uterine contractions.

## Ergotamine

A medicine that narrows certain blood vessels and can stop some migraine attacks. It has many interactions and should not be overused because it can cause dangerous loss of blood flow to the limbs or organs.

## Ergotism

Poisoning caused by ergot toxins or excessive ergot-derived medicines. Severe blood-vessel narrowing can cause burning pain, numbness, cold or discolored limbs, confusion, seizures, tissue death, or dangerous uterine contractions.

## Ergots

Chemicals produced by ergot fungi, some of which are used in medicines for migraine or bleeding. They can strongly narrow blood vessels and interact with other medicines, causing serious circulation problems or poisoning.

## Eribulin Mesylate

A cancer medicine that interferes with microtubules, structures needed for cell division. It is used for selected advanced breast cancers and liposarcomas; common risks include low blood counts, nerve damage, fatigue, and nausea.

## Erlotinib Hydrochloride

A cancer medicine that blocks growth signals from the epidermal growth-factor receptor. It treats selected cancers with suitable molecular changes and may cause rash, diarrhea, liver injury, or lung inflammation.

## Erosion

Loss of only the surface layer of skin or a moist lining, without the deeper tissue loss of an ulcer. Friction, inflammation, infection, injury, or chemicals may cause it and may produce pain or bleeding.

## Erosion Lesion

A shallow area where the surface layer of skin or a moist lining has been lost. It may result from rubbing, inflammation, infection, injury, or chemicals and usually heals when the cause is removed.

## Erosive Esophagitis

Inflammation of the swallowing tube with visible breaks in its lining, usually from repeated stomach-acid reflux. It can cause painful swallowing, heartburn, or bleeding and may lead to narrowing if untreated.

## Erosive Gastritis

Irritation and inflammation of the stomach lining with shallow areas of damage. Common causes include anti-inflammatory pain medicines, alcohol, severe illness, and stress on the body; it may cause pain, vomiting, or gastrointestinal bleeding.

---

### Erosive Lichen Planus

A long-lasting immune-mediated inflammation that causes painful raw areas, burning, bleeding, or soreness in the mouth or genital tissues. Scarring and narrowing may develop, so examination and treatment can help protect function.

#### Alternate Names:

Vulvovaginal Gingival Syndrome; Erosive Vulvovaginal Lichen Planus

### Erratic Results

Test results that change unexpectedly or do not agree with repeated measurements. Possible causes include sample problems, device error, normal biological variation, medicines, or a test that does not fit the clinical question.

### Ertapenem

A once-daily antibiotic for selected serious infections caused by susceptible bacteria. It does not reliably treat *Pseudomonas* or enterococcal infections and can cause diarrhea, allergic reactions, seizures, or antibiotic-resistant organisms.

### Erucic Acid

A fatty acid found in some plant oils, including mustard and older varieties of rapeseed oil. Modern edible oils are usually bred to contain little of it; unusually high exposure may harm the heart.

### Erucic Acids

Long-chain fatty acids found mainly in certain plant oils. The amount consumed depends on the oil, and unusually high exposure may be harmful; ordinary low-erucic edible oils provide much less.

### Eructation

Release of swallowed air or gas from the stomach through the mouth, commonly called belching. Frequent troublesome belching may be related to swallowed air, reflux, food intolerance, or other digestive problems.

#### Alternate Names:

Belching; Burping

### Eruption Cyst

A soft, fluid-filled swelling over a tooth that is about to come through the gum, often blue or purple. It usually disappears when the tooth erupts, but painful or infected swelling needs dental review.

### Eruptive Fever

Fever occurring with a widespread rash. Viral infections are common causes, but bacteria, medicines, allergic reactions, and immune disorders can also be responsible; urgent assessment is needed for severe illness or breathing difficulty.

### Eruptive Vellus Hair Cysts

Benign follicular cysts containing fine vellus hairs that appear as multiple small, smooth, red-brown or skin-colored papules, often on the chest, abdomen, or limbs. They may be inherited or sporadic and usually need treatment only for symptoms or cosmetic reasons.

### Eruptive Xanthomas

Sudden crops of small yellow-red bumps, often on the buttocks, elbows, knees, or trunk, caused by very high blood triglycerides. They signal a serious lipid disorder requiring prompt evaluation and treatment.

### Eruptive Xanthomatosis

A sudden outbreak of many small yellow-red skin bumps caused by fat collecting in skin immune cells. It is usually linked to very high triglycerides, sometimes from uncontrolled diabetes or an inherited lipid disorder.

### Eryptosis

A controlled process by which aging or damaged red blood cells shrink and are marked for removal from the bloodstream. It helps maintain blood-cell balance but may increase in some diseases or during oxidative stress.

### Erysipelas

A sudden bacterial infection of the upper skin and lymph vessels, usually causing a painful, raised, sharply bordered red patch with fever. Prompt antibiotic treatment is important to prevent spread

---

or complications.

### **Erysipeloid**

A skin infection caused by bacteria acquired through cuts while handling fish, meat, poultry, or animals. It usually causes a slowly expanding painful purple-red patch on a finger or hand and needs antibiotic treatment.

### **Erysipelothrix**

A group of bacteria found in animals and animal products that can enter broken skin and cause a painful localized infection. Rarely, infection spreads to the bloodstream or heart valves.

### **Erythema**

Redness of the skin or a moist body lining caused by increased blood flow near the surface. It may result from inflammation, infection, heat, allergy, or another illness and often fades briefly when pressed.

### **Erythema Ab Igne**

A net-like red-brown skin discoloration caused by repeated exposure to moderate heat, such as heating pads, hot-water bottles, or heaters. Stopping the exposure may allow improvement; persistent lesions need examination.

### **Erythema Annulare Centrifugum**

A reactive gyrate erythema presenting with slowly expanding, polycyclic, annular erythematous plaques characterized by central clearing and a distinctive trailing rim of fine scale along the inner border of the advancing erythematous margin.

#### **Alternate Names:**

EAC; Centrifugal Annular Erythema

### **Erythema Chronicum Migrans**

An expanding skin rash that can appear days to weeks after a tick transmits Lyme disease bacteria. It may be evenly red or ring-shaped and does not always look like a classic bull's-eye.

### **Erythema Dose**

The amount of ultraviolet or other radiation needed

to produce visible skin redness. The minimal erythema dose is the smallest amount that causes a defined response and varies with skin type, medicines, and exposure conditions.

### **Erythema Dyschromicum Perstans**

An acquired pigmentary disorder causing slowly spreading, sharply defined gray or blue-gray macules and patches, often on the face, neck, and trunk. Early lesions may have a reddish raised border; biopsy may help distinguish it from other causes of dermal hyperpigmentation, and treatment is often difficult.

### **Erythema Elevatum Diutinum**

A rare chronic cutaneous leukocytoclastic vasculitis presenting as persistent, symmetrical, red-violaceous to yellowish-brown papules, nodules, and plaques over the extensor surfaces of joints, associated with IgA paraproteinemia, hematologic disorders, and chronic infections.

#### **Alternate Names:**

EED; Chronic Extensor Leukocytoclastic Vasculitis

### **Erythema Gyrratum Repens**

A rapidly spreading, ringed skin rash with a wood-grain pattern. It is often associated with an internal cancer, especially of the lung, so new or persistent cases require prompt medical evaluation.

### **Erythema Induratum**

A form of nodular vasculitis causing recurrent tender, sometimes ulcerating nodules on the backs of the lower legs. Biopsy shows lobular panniculitis with vasculitis; tuberculosis and other infections, inflammatory disease, venous problems, and medicines may be associated, so the cause should be investigated.

### **Erythema Infectiosum**

A usually mild infection caused by parvovirus B19, also called fifth disease. A short flu-like prodrome may be followed by bright-red cheeks and a lacy rash on the trunk and limbs; adults may have joint pain, while infection can cause severe anemia in people with certain blood disorders or immunosuppression



---

and complications in pregnancy.

### **Erythema Infectiosum (Fifth Disease)**

Alternate name; see *Erythema Infectiosum*.

#### **Alternate Names:**

Fifth Disease; Slapped Cheek Disease

### **Erythema Marginatum**

A temporary, non-itchy, ring-shaped or wavy red rash that usually appears on the trunk and upper limbs. It is a recognized sign of acute rheumatic fever and may come and go over days or weeks.

### **Erythema Migrans**

An expanding rash that develops days to weeks after a tick transmits Lyme disease bacteria. It may be uniformly red or ring-shaped and is usually diagnosed from its appearance and exposure history.

### **Erythema Multiforme**

A short-lived immune reaction, often triggered by herpes simplex infection or, less commonly, another infection or medicine. It causes symmetrical target-shaped spots, usually on the hands, feet, arms, or face; mouth involvement may occur.

### **Erythema Nodosum**

Inflammation of the fat beneath the skin (panniculitis) that causes tender red lumps, usually on both shins. Infections, medicines, inflammatory bowel disease, sarcoidosis, and other conditions may trigger it. The lumps usually fade like bruises without ulcerating or scarring.

### **Erythema Toxicum**

A common harmless newborn rash with red spots, small bumps, or tiny pus-filled areas. It often appears during the first days of life and disappears on its own without treatment.

### **Erythema Toxicum Neonatorum**

A harmless newborn rash that usually appears during the first few days as red blotches, bumps, or tiny pustules on the face, trunk, or limbs. It usually spares the palms and soles and clears within one to

two weeks.

#### **Alternate Names:**

Erythema Neonatorum; Toxic Erythema of the Newborn

### **Erythrasma**

A superficial bacterial infection in warm, moist skin folds that causes well-defined reddish-brown patches, often in the groin, armpits, or between toes. It usually responds to prescribed antibacterial treatment and keeping folds dry.

### **Erythroblast**

An immature red-blood-cell precursor made in the bone marrow. It develops through several stages before becoming a mature red blood cell; abnormal numbers may indicate altered blood-cell production.

### **Erythroblastosis Fetalis**

A condition in which antibodies from the pregnant person destroy a fetus's or newborn's red blood cells. It can cause anemia, jaundice, widespread swelling, heart failure, or death and is often prevented by blood-group screening and preventive treatment.

### **Erythroblastosis Foetalis**

Destruction of fetal or newborn red blood cells by antibodies from the mother. It can cause anemia, jaundice, widespread fluid buildup, heart failure, or death. Pregnancy blood-group testing and preventive treatment can reduce the risk.

### **Erythrocyte**

A mature red blood cell that carries oxygen from the lungs to tissues and helps return carbon dioxide to the lungs. Its flexible shape allows it to pass through very small blood vessels.

### **Erythrocyte Alloimmunization and Pregnancy**

Maternal antibodies directed against fetal red-cell antigens, commonly RhD, can cross the placenta and cause fetal or newborn hemolytic anemia. Blood-group screening, antibody testing, fetal surveillance, and preventive anti-D immunoglobulin when indicated reduce risk.

---

## Erythrocyte Casts

Tube-shaped groups of red blood cells seen in urine. They form inside the kidney's small tubes and strongly suggest bleeding or inflammation in the kidney's filtering units, such as glomerulonephritis or vasculitis.

### Alternate Names:

Red Blood Cell Casts; RBC Casts

## Erythrocyte Enzyme Deficiency

An inherited or acquired lack of a protein that red blood cells need for energy or protection. The cells may break down too early and cause anemia; examples include G6PD and pyruvate-kinase deficiency.

## Erythrocyte Membrane

The flexible outer layer of a red blood cell. It maintains the cell's shape, allows it to bend through small vessels, and controls movement of water and substances into and out of the cell.

## Erythrocyte Sedimentation Rate

A blood test measuring how quickly red blood cells settle in a tube over one hour. A faster rate can occur with inflammation, but it does not identify the cause and is affected by anemia, age, pregnancy, and other conditions.

## Erythrocytosis

An abnormally high number or volume of red blood cells, reflected by high hemoglobin or hematocrit. It may result from dehydration, low oxygen, excess erythropoietin, or a bone-marrow disorder and can increase clot risk.

## Erythroderma

Intense redness and usually scaling affecting about 90

### Alternate Names:

Exfoliative Dermatitis; Exfoliative Erythroderma; Generalized Exfoliative Dermatitis

## Erythrodermic Psoriasis

A rare, potentially life-threatening form of psoriasis causing nearly all the skin to become intensely red,

painful, and scaly. Fever, chills, dehydration, infection, and unstable blood pressure may occur and require urgent care.

## Erythrodysplasia

Abnormal development or maturation of red-blood-cell precursors in the bone marrow. It can produce ineffective blood formation and anemia and may occur with a marrow disorder or other illness.

## Erythroid Hyperplasia

An increase in immature red-blood-cell precursors in the bone marrow, usually because the body is responding to anemia, blood loss, red-cell destruction, or low oxygen. The underlying cause needs evaluation.

## Erythrokeratoderma Variabilis et Progressiva

A rare inherited keratinization disorder characterized by persistent, often symmetrical red and thickened scaly plaques together with more variable migrating erythematous patches. Onset is usually in infancy or childhood, and treatment aims to reduce scaling and inflammation with emollients, keratolytics, or specialist-directed retinoids.

## Erythroleukemia

A rare acute myeloid leukemia characterized by prominent abnormal erythroid precursors in the bone marrow, often with dysplastic or immature myeloid cells. Diagnosis requires blood and marrow evaluation with immunophenotypic and genetic classification.

## Erythroleukoplakia

A mouth lesion containing both red and white areas. It has a substantial risk of advanced precancerous change or cancer, so a dentist or specialist should examine it and usually perform a biopsy.

## Erythromelalgia

Repeated attacks of burning pain, warmth, and redness in the feet, hands, or sometimes face, often triggered by heat or exercise and eased by cooling. It may be inherited or linked to blood or nerve

---

disorders.

### **Erythromycin**

An antibiotic that blocks bacterial protein production. It treats selected infections and can be used when some other antibiotics are unsuitable; nausea, diarrhea, liver injury, and abnormal heart rhythm are possible.

### **Erythromycin Estolate**

An oral form of the antibiotic erythromycin. It blocks bacterial protein production and treats selected infections; stomach upset, liver inflammation, allergic reactions, and medicine interactions can occur.

### **Erythromycin Ophthalmic Prophylaxis**

Antibiotic ointment placed in a newborn's eyes soon after birth to reduce the risk of severe eye infection from gonorrhea acquired during delivery. It does not replace testing and treatment of infections during pregnancy.

### **Erythroplakia**

A persistent, sharply defined red patch on a moist lining, especially inside the mouth. It may contain advanced precancerous change or cancer and requires prompt examination, usually with biopsy.

### **Erythroplasia**

A sharply bordered red patch on skin or a moist lining. Persistent lesions, especially in the mouth or genital area, may represent a serious precancerous change or cancer and need medical assessment.

### **Erythroplasia Of Queyrat**

Squamous-cell carcinoma in situ affecting the head of the penis or foreskin. It usually appears as a persistent, well-defined red or velvety patch and can progress to invasive cancer without treatment.

### **Erythropoiesis**

The process of making mature red blood cells, mainly in the bone marrow. It is stimulated by erythropoietin when oxygen is low and requires healthy

marrow plus adequate iron, vitamin B12, and folate.

### **Erythropoiesis-Stimulating Agents**

Medicines that signal the bone marrow to make more red blood cells. They are used for selected anemias, including some caused by kidney disease or cancer treatment, but may increase blood pressure and clot risk.

### **Erythropoietin**

A hormone made mainly by the kidneys when body tissues receive too little oxygen. It signals the bone marrow to produce red blood cells; kidney disease can reduce this signal and contribute to anemia.

### **Erythropoietin Deficiency**

An insufficient amount of erythropoietin for the body's need, resulting in reduced stimulation of red-blood-cell production in the bone marrow. It commonly contributes to anemia in chronic kidney disease; reduced response to available erythropoietin is a separate phenomenon.

### **Erythropoietin Receptor**

A protein on immature red-blood-cell precursors that receives the erythropoietin signal. Activation helps these cells survive, multiply, and mature into red blood cells.

### **Erythropoietin Test**

A blood test measuring the kidney-made hormone that stimulates red-blood-cell production. It can help investigate selected anemias or high red-cell counts and is interpreted with the blood count, oxygen status, and kidney function.

### **Erythrosine**

A red dye used in some foods, medicines, and laboratory procedures. Exposure is usually small, but regulations vary and people with a known sensitivity should follow product and medical advice.

### **ESBL**

An enzyme made by some bacteria that destroys many penicillin and cephalosporin antibiotics. In-

---

fections caused by these bacteria can be harder to treat and require laboratory testing to select an effective medicine.

### **ESBL-Producing Organisms**

Bacteria that make enzymes capable of breaking down many penicillin and cephalosporin antibiotics. They commonly include certain intestinal bacteria such as *E. coli* and *Klebsiella*; treatment depends on susceptibility testing and infection site.

### **Escape Reaction**

A body response that restores an organ's activity after it has been reduced by a medicine, hormone, nerve signal, or feedback system. The response may lessen the effect of continued treatment or stimulation.

### **Eschar**

A firm layer of dead tissue, often black or brown, over a burn, pressure injury, wound, infection, or area with poor blood supply. It can hide deeper damage and may need specialist assessment or removal.

### **Escharotomy**

A surgical cut through stiff dead tissue around a deep burn to release pressure. It may restore circulation to a limb or allow the chest to expand, and is performed urgently when swelling threatens function.

### **Escherichia**

A group of bacteria that includes *E. coli* and related species found in intestines or the environment. Some strains are harmless, while others cause diarrhea, urinary infection, wound infection, or bloodstream infection.

### **Escherichia Albertii**

An uncommon intestinal bacterium that can cause diarrheal illness. It may be mistaken for certain *E. coli* strains, so specialized laboratory identification may be needed when infection is suspected.

### **Escherichia Blattae**

A rarely reported *Escherichia* species associated mainly with insects and the environment. Its significance in a human sample is uncertain and must be judged from symptoms, the specimen source, and other test results.

### **Escherichia Coli**

A bacterium that normally lives in the intestine. Some strains are harmless, while others cause urinary, intestinal, wound, or bloodstream infections, including severe bloody diarrhea and hemolytic uremic syndrome.

### **Escherichia Coli Infection**

Infection caused by a harmful strain of *E. coli*. It may affect the intestine, urinary tract, wounds, or bloodstream; symptoms, contagiousness, and treatment depend on the strain and site.

### **Escherichia Coli O157:H7 Infection**

Infection with a toxin-producing *E. coli* strain, usually acquired from contaminated food, water, animals, or people. It commonly causes severe cramps and bloody diarrhea and can lead to kidney failure; urgent medical advice is important.

### **Escherichia Fergusonii**

An uncommon bacterium that can occasionally cause urinary, wound, intestinal, or bloodstream infection, especially in people with serious illness or weakened immunity. Laboratory identification and susceptibility testing guide treatment.

### **Escherichia Hermannii**

An uncommon *Escherichia* species that rarely causes opportunistic infection, including urinary or bloodstream infection. Its importance depends on the patient's symptoms, immune status, and the specimen in which it is found.

### **Escherichia Vulneris**

An uncommon bacterium rarely linked to wound, urinary, or bloodstream infection, usually in people with other serious illnesses. A laboratory result requires clinical correlation to distinguish infection

---

from contamination.

### **Escitalopram**

An antidepressant that increases available serotonin in the brain. It treats depression and generalized anxiety disorder; nausea, sleep changes, sexual problems, withdrawal symptoms, bleeding risk, and rarely serotonin toxicity may occur.

### **ESKD**

Severe permanent loss of kidney function, usually meaning kidney filtration below about 15 milliliters per minute or a need for dialysis or transplantation. It can cause fluid, electrolyte, anemia, and toxin problems and needs kidney-specialist care.

### **Esomeprazole**

A medicine that reduces stomach-acid production by blocking the acid pump in stomach cells. It treats reflux, ulcers, and acid-related disorders; prolonged or inappropriate use may cause infections, low magnesium, or vitamin deficiencies.

### **Esomeprazole Magnesium**

A magnesium salt of esomeprazole that reduces stomach-acid production by blocking the stomach's acid pump. It treats reflux and ulcers; long-term use may cause diarrhea, infections, low magnesium, or nutrient deficiencies.

### **Esophageal**

Relating to the esophagus, the muscular tube that carries swallowed food and liquid from the throat to the stomach.

### **Esophageal Achalasia**

A disorder in which the lower esophageal valve does not relax properly and the esophagus cannot push food normally into the stomach. It causes difficulty swallowing, regurgitation, chest discomfort, and weight loss.

### **Esophageal Adenocarcinoma**

A cancer arising from gland-forming cells, usually in the lower esophagus and often linked to Barrett

esophagus. Reflux, obesity, and smoking increase risk; progressive swallowing difficulty or weight loss requires evaluation and biopsy.

### **Esophageal Atresia**

A birth defect in which the esophagus ends in a blind pouch instead of connecting normally to the stomach. A connection to the windpipe may also occur, causing drooling, choking, coughing, and breathing problems during feeds.

### **Esophageal Bleeding**

Bleeding from the esophagus, which may result from enlarged veins, inflammation, ulcers, tears, injury, or cancer. Vomiting blood, passing black stools, dizziness, or fainting requires emergency medical care.

### **Esophageal Cancer**

A malignant tumor of the swallowing tube. The main types are squamous-cell cancer and adenocarcinoma; progressive difficulty swallowing, weight loss, pain, or bleeding may occur, and treatment depends on the stage.

#### **Alternate Names:**

Esophageal Carcinoma; Esophagus Malignancy

### **Esophageal Candidiasis**

A yeast infection of the esophagus, more common with weakened immunity, diabetes, or certain medicines. It often causes painful or difficult swallowing and is treated with antifungal medicine, sometimes after endoscopy confirms it.

### **Esophageal Carcinoma**

Cancer arising in the esophagus, most often squamous-cell carcinoma or adenocarcinoma. Progressive difficulty swallowing, weight loss, chest pain, or bleeding may occur; endoscopic biopsy and staging guide treatment.

### **Esophageal Culture**

Laboratory growth testing on a sample from the esophagus to identify bacteria or fungi when infection is suspected. Results depend on how the sample was collected and must be interpreted with



---

symptoms and endoscopic findings.

### **Esophageal Dilation**

A procedure that gently stretches a narrowed area of the esophagus with a tube or balloon passed through an endoscope. It can improve swallowing caused by scarring, inflammation, injury, or other narrowing; tearing and bleeding are uncommon but serious risks.

### **Esophageal Diseases**

Disorders affecting the swallowing tube between the throat and stomach. They include reflux, inflammation, narrowing, movement problems, cancer, infection, and birth defects, often causing swallowing difficulty, pain, heartburn, or regurgitation.

### **Esophageal Diverticulum**

A pouch that bulges outward from the wall of the esophagus. It may cause difficulty swallowing, food coming back up, coughing, bad breath, or no symptoms; treatment depends on its size, location, and effects.

### **Esophageal Duplication**

A rare birth defect consisting of a cyst or extra segment beside the esophagus. It may press on the esophagus or airway, causing swallowing difficulty, breathing problems, infection, or no symptoms.

### **Esophageal Fistula**

An abnormal connection between the esophagus and another organ, often the airway. It can allow food or liquid into the lungs, causing coughing during swallowing, repeated pneumonia, poor nutrition, or severe illness.

### **Esophageal Foreign Body**

Food or another object stuck in the esophagus, causing pain, difficulty swallowing, drooling, or breathing problems. A battery, sharp object, complete blockage, or suspected tear requires emergency removal.

### **Esophageal Infection**

Infection of the esophagus caused by a yeast, virus, or bacterium, usually in people with weakened immunity or damaged lining. It may cause painful swallowing, difficulty swallowing, chest pain, or ulcers.

### **Esophageal Intramural Pseudodiverticulosis**

A rare condition in which small drainage ducts of glands within the esophageal wall become widened. It may cause repeated swallowing difficulty, food impaction, or narrowing and is diagnosed with imaging or endoscopy.

### **Esophageal Leiomyoma**

A usually benign smooth-muscle tumor arising in the esophageal wall. It may cause difficulty swallowing, chest discomfort, or no symptoms and is distinguished from other submucosal tumors with imaging, endoscopy, and tissue assessment when needed.

### **Esophageal Leukoplakia**

A persistent white patch on the esophageal lining caused by thickened surface cells. It is uncommon but may contain precancerous change, so evaluation and biopsy are often needed to determine its significance.

### **Esophageal Manometry**

A test that measures pressure and muscle coordination in the esophagus during swallowing. A thin pressure-sensitive tube passes through the nose into the stomach and helps diagnose achalasia, spasm, and other movement disorders.

### **Esophageal Motility Disorder**

A disorder of the muscle contractions or nerve control that moves food through the esophagus. It may cause difficulty swallowing, chest pain, regurgitation, or food sticking; testing may include esophageal manometry.

### **Esophageal Necrosis**

Death of esophageal lining, sometimes producing



---

a black appearance. It can result from severe low blood flow, shock, acid injury, or caustic substances and may lead to bleeding or perforation.

### **Esophageal Perforation**

A hole through the full thickness of the esophageal wall, caused by forceful vomiting, an instrument, injury, or disease. It can release contents into the chest, causing severe infection and requiring emergency treatment.

### **Esophageal pH Monitoring**

A test that records how much acid enters the esophagus, usually for 24 hours, using a thin tube or a small wireless sensor. It helps determine whether acid reflux explains persistent symptoms when the diagnosis is uncertain.

### **Esophageal Spasm**

A movement disorder in which esophageal muscles contract too strongly or without normal coordination. It can cause intermittent chest pain and difficulty swallowing; pressure testing helps distinguish it from other disorders.

### **Esophageal Spasms**

Uncoordinated or forceful contractions of the esophagus that may cause sudden chest pain, food sticking, or difficulty swallowing. Symptoms can resemble heart pain and should be assessed when severe or new.

### **Esophageal Squamous Cell Carcinoma**

A cancer arising from the flat cells lining the esophagus, often in its upper or middle part. Smoking, heavy alcohol use, certain infections, and nutritional factors increase risk; diagnosis requires endoscopic biopsy.

### **Esophageal Stenosis**

Narrowing of the passage through the esophagus caused by scarring, inflammation, a tumor, radiation, or a birth defect. It commonly causes progressively worsening difficulty swallowing, especially solid foods.

### **Esophageal Stent**

A tube placed inside a narrowed or leaking esophagus to keep it open and allow swallowing. It is often used for cancer or injury; movement, blockage, bleeding, pain, or perforation can occur.

### **Esophageal Stricture**

A narrowed area of the esophagus, commonly caused by acid reflux, caustic injury, inflammation, radiation, or procedures. It usually causes gradually worsening difficulty swallowing solids and may require stretching during endoscopy.

### **Esophageal Surgery**

An operation on the esophagus or the area where it meets the stomach. It may treat cancer, severe narrowing, perforation, reflux, or movement disorders; the procedure and risks depend on the disease and overall health.

### **Esophageal Ulceration**

An open sore in the esophageal lining caused by reflux, infection, medicines, caustic injury, inflammation, or cancer. It may cause painful swallowing, chest pain, or bleeding and often requires endoscopic evaluation.

### **Esophageal Varices**

Enlarged veins in the esophageal wall, usually caused by high pressure in the portal vein from advanced liver disease. They may rupture without warning and cause life-threatening vomiting of blood.

#### **Alternate Names:**

Esophageal Dilated Veins; Portosystemic Varices

### **Esophageal Varices Cross-Reference**

Alternate name; see *Esophageal varices*.

### **Esophageal Web**

A thin shelf of tissue projecting into the upper esophagus. It may cause difficulty swallowing solid food and can be associated with iron-deficiency anemia; persistent symptoms may require endoscopy.

---

## Esophagectomy

Surgery to remove part or all of the esophagus, usually for cancer or severe irreversible damage. The remaining digestive tract is reconstructed, commonly using the stomach or a segment of intestine.

## Esophagitis

Inflammation or injury of the esophageal lining. Common causes include acid reflux, infection, medicines that irritate the esophagus, eosinophilic or other immune-mediated disease, swallowed irritants, and radiation. Symptoms may include heartburn, painful or difficult swallowing, chest pain, food sticking, or bleeding; persistent inflammation can cause ulcers or narrowing.

## Esophagogastric Bypass

Surgery that creates a new route between the esophagus and stomach to restore passage of food, usually after esophageal removal or severe blockage. The method depends on anatomy, disease, and surgical goals.

## Esophagogastric Junction

The area where the esophagus joins the stomach. Its muscular valve and surrounding structures help food enter the stomach while limiting backward flow of stomach contents.

## Esophagogastric Tamponade

Temporary control of severe bleeding from esophageal or stomach veins by inflating a balloon device to apply pressure. It is a rescue measure used while definitive treatment is arranged and can obstruct breathing or cause aspiration.

## Esophagogastroduodenoscopy

An upper endoscopic diagnostic and therapeutic procedure in which a thin, flexible video endoscope is passed through the mouth to examine the lining of the esophagus, stomach, and duodenum.

### Alternate Names:

EGD; Upper Endoscopy; Upper Gastrointestinal Endoscopy

## Esophagoscope

A flexible or rigid instrument used to look inside the esophagus. It may allow a clinician to take tissue samples, remove a foreign object, treat bleeding, or perform another procedure.

## Esophagoscopy

Examination of the esophagus with a camera-equipped instrument. It may investigate swallowing difficulty, bleeding, reflux damage, narrowing, foreign objects, infection, or suspected cancer and may allow treatment or biopsy.

## Esophagram

An X-ray examination performed while a person swallows contrast liquid. It shows the shape and movement of the esophagus and can reveal narrowing, leaks, pouches, blockage, or abnormal swallowing.

## Esophagus

A muscular tube that carries food and liquid from the throat to the stomach. Coordinated muscle contractions move swallowed material downward, while valves at each end help limit reflux and air entry.

## Esophagus Disease

A disorder of the esophagus causing swallowing difficulty, pain, reflux, regurgitation, bleeding, obstruction, inflammation, motility disturbance, or cancer.

## Esophagus Neoplasm

A benign, precancerous, or malignant growth in the esophagus; evaluation may require endoscopy, biopsy, imaging, and staging.

## Esotropia

A condition in which one eye turns inward toward the nose, causing the eyes to point in different directions. It may begin in infancy or result from focusing problems, illness, or injury; glasses, patching, or surgery may help depending on the cause.

---

## ESR

The erythrocyte sedimentation rate, a nonspecific blood test measuring how quickly red cells settle in a tube; it can reflect inflammation but cannot identify its cause.

### Alternate Names:

SED Rate

## ESRD

Alternate index label; see *End-Stage Renal Disease*.

## Essential

Necessary or fundamental; in a diagnosis such as essential hypertension, it indicates that no single secondary cause has been identified.

## Essential Fats

Fatty acids the body cannot make in sufficient amounts and must obtain from food. They support cell membranes, skin, brain function, and production of signaling molecules involved in inflammation.

### Alternate Names:

Essential Fatty Acids

## Essential Fatty Acid

A fatty acid the body cannot make in sufficient amounts and therefore must obtain from food; the main human examples are linoleic acid and alpha-linolenic acid.

## Essential Hypertension

Long-term high blood pressure with no single identifiable underlying cause. Many genetic, kidney, blood-vessel, metabolic, and lifestyle factors contribute, and untreated disease raises the risk of stroke, heart disease, kidney damage, and vision problems.

## Essential Medicines

Medicines considered necessary to meet the most important health needs of a population. They are selected for proven benefit, safety, affordability, and practical availability and should be reliably accessible through healthcare services.

## Essential Mixed Cryoglobulinemia

An immune disorder in which abnormal proteins in the blood thicken or clump in the cold and inflame small blood vessels. It may cause purple spots, joint pain, weakness, nerve problems, or kidney disease.

## Essential Nutrient

A nutrient the body cannot make in sufficient amounts and must obtain from food or supplements. Essential nutrients include certain amino acids, fatty acids, vitamins, minerals, and water.

## Essential Thrombocythemia

A long-term bone-marrow disorder that produces too many platelets. It can cause abnormal clotting or bleeding and may cause headaches, vision changes, burning pain in fingers or toes, or no symptoms.

## Essential Thrombocytosis

A chronic myeloproliferative neoplasm with persistent overproduction of platelets, often driven by JAK2, CALR, or MPL mutations. It can cause thrombosis, bleeding, microvascular symptoms, or splenomegaly.

## Essential Tremor

A nervous-system disorder causing rhythmic shaking, usually of both hands during movement or when holding a position. It may also affect the head or voice, gradually worsen, and interfere with writing, eating, or other tasks. It can run in families but also occur without a family history.

### Alternate Names:

Benign Essential Tremor; Familial Essential Tremor; Hereditary Essential Tremor

## Essiac

A herbal mixture promoted as a treatment for cancer and other illnesses. Reliable evidence has not shown that it cures cancer, and its ingredients may interact with medicines or cause liver, kidney, or other harm.

## Estazolam

A sedative medicine sometimes used short-term for

---

insomnia. It slows brain activity but can cause daytime drowsiness, falls, memory problems, slowed breathing, dependence, and withdrawal; alcohol and opioids increase its dangers.

### **Ester**

A chemical formed when an acid combines with an alcohol. Esters occur naturally in fats, medicines, and flavorings; their health effects depend on the specific compound, amount, route of exposure, and duration.

### **Esterase**

An enzyme that breaks ester chemical bonds. Esterases help process fats, medicines, and other substances; abnormal activity or inhibition can alter drug effects and may contribute to poisoning.

### **Esterified Estrogens**

A mixture of estrogen hormones used for selected menopausal symptoms or estrogen deficiency. Treatment can cause bleeding and increase risks of blood clots, stroke, and breast or uterine cancer, so benefits and risks must be individualized.

### **Esthesia**

The ability to sense touch, pain, temperature, movement, or body position. Reduced, increased, or altered sensation may result from nerve injury, brain disease, medicines, or changes in consciousness.

### **Esthesiometer**

An instrument used to measure sensitivity to touch, pressure, or other sensations. Eye clinicians may use one to assess corneal sensation when nerve injury or an eye-surface problem is suspected.

### **Esthesioneuroblastoma**

A rare cancer arising from nerve-related cells high in the nasal cavity. It may cause blocked or bleeding nose, reduced smell, facial pain, or vision symptoms and is diagnosed with imaging and biopsy.

### **Estimated Average Glucose**

An estimate of a person's average blood-glucose

level calculated from the hemoglobin A1c result. It is reported in familiar glucose units but does not show daily highs, lows, or sudden changes.

### **Estimated Date Of Confinement**

The approximate date when a pregnancy is expected to reach delivery, calculated from the last menstrual period or an early ultrasound. Only a small proportion of births occur on the estimated date.

### **Estimated Due Date**

The approximate date of delivery, usually calculated as 40 weeks from the first day of the last menstrual period and adjusted when needed using an early ultrasound. Most births occur before or after this date.

### **Estimated Fetal Weight**

An ultrasound-based estimate of a fetus's weight using measurements of the head, abdomen, and thigh bone. It helps assess growth and plan care, but commonly differs from actual birth weight by about 10–15 percent.

### **Estimated Glomerular Filtration Rate**

An estimate of how well the kidneys filter blood, calculated mainly from blood creatinine and other factors. It helps assess kidney disease and adjust medicines but is less reliable during rapidly changing illness or unusual body size.

### **Estimation of Fetal Weight**

An ultrasound-based estimate of fetal size using measurements such as head, abdominal, and femur dimensions. It helps assess growth and delivery planning but has an important margin of error and is interpreted with gestational age and clinical findings.

### **Estradiol**

The main active estrogen hormone during most reproductive years. It helps regulate ovulation and menstruation and supports the vagina, uterus, bones, brain, and other tissues; levels fall after menopause.

**Alternate Names:**

---

17 $\beta$ -Estradiol

### **Estradiol Blood Test**

A blood test measuring estradiol. It may help evaluate puberty, menstrual or fertility problems, ovarian or testicular function, menopause, or selected hormone disorders; results vary with age, sex, cycle timing, pregnancy, and medicines.

### **Estramustine**

A cancer medicine combining an estrogen-like component with a compound that interferes with cell division. It has been used for selected advanced prostate cancers; blood clots, fluid retention, nausea, and heart effects may occur.

### **Estriol**

An estrogen produced mainly by the placenta during pregnancy from substances made by the fetus and mother. Maternal blood estriol may be included in prenatal screening, but an abnormal result does not by itself diagnose a problem.

### **Estrogen**

A group of hormones involved in reproductive development, menstruation, fertility, bone strength, and many other body functions. The main forms and amounts vary with age, sex, pregnancy, and ovarian function.

### **Estrogen Overdose**

Taking or receiving more estrogen than prescribed. It may cause nausea, vomiting, breast tenderness, headache, or abnormal bleeding and can increase the risk of blood clots; urgent advice is needed after a large exposure.

### **Estrogen Receptor**

A cell protein that binds estrogen and changes cell activity. Testing for estrogen receptors helps classify some cancers, especially breast cancer, and can guide hormone-based treatment.

### **Estrogen Replacement Therapy**

Treatment supplying estrogen to relieve selected

menopausal symptoms or replace deficient hormone. If the uterus is present, a progestogen is usually added to protect its lining; risks and benefits must be reviewed individually.

### **Estrogen Test**

A blood, urine, or saliva test measuring one or more estrogen hormones. It may help evaluate puberty, menstrual problems, fertility, menopause, pregnancy, or hormone disorders; interpretation depends on the test type and timing.

#### **Alternate Names:**

Estrogen Levels Test Estrogen Blood Test

### **Estrogen-Replacement Therapy**

Treatment that supplies estrogen when natural levels are low, such as after menopause or ovarian failure. It may relieve hot flashes and protect bone, but can cause bleeding and increase some health risks.

#### **Alternate Names:**

Estrogen Therapy

### **Estrogens**

Hormones, including estradiol, that regulate reproductive development, menstruation, fertility, bones, and other tissues. Before menopause they are produced mainly by the ovaries; after menopause, levels are lower and come from several tissues.

### **Estrone**

A naturally occurring estrogen made partly by converting other hormones in body tissues. It is the main estrogen present after menopause, although estradiol is usually more active before menopause.

### **Estrous Cycle**

The recurring reproductive cycle of many female mammals other than humans. It includes hormonal and physical changes that prepare for ovulation, mating, and possible pregnancy.

### **Eszopiclone**

A sleep medicine that enhances inhibitory signaling in the brain. It can help insomnia but may

---

cause drowsiness, impaired coordination, memory problems, dependence, unusual sleep behaviors, or a bitter taste.

### **Etacrynic Acid**

A diuretic that makes the kidneys remove more salt and water, reducing swelling and sometimes blood pressure. It can cause dehydration, abnormal electrolytes, kidney problems, or hearing damage, especially at high doses.

### **Etanercept**

A biologic medicine that blocks tumor-necrosis factor, an inflammation signal. It treats selected arthritis, psoriasis, and other inflammatory diseases but can increase serious infection risk, so tuberculosis and hepatitis screening may be needed.

### **Ethacrynic Acid**

A loop diuretic that increases kidney excretion of salt and water. It can reduce fluid overload but may cause dehydration, low electrolytes, kidney injury, or dose-related hearing damage.

### **Ethambutol**

An antibiotic used with other medicines to treat tuberculosis and selected infections caused by related bacteria. It can damage the optic nerve, causing blurred vision or trouble distinguishing colors, so vision testing is important.

### **Ethambutol Toxicity**

A dose- and duration-dependent adverse pharmacological effect of the first-line antimycobacterial agent ethambutol, characterized by retrobulbar or axial optic neuritis, presenting with bilateral painless reduction in visual acuity, central scotomas, and loss of red-green color discrimination.

### **Ethane**

A colorless, highly flammable gas used mainly to make industrial chemicals. It is not usually poisonous through direct action, but high concentrations can displace oxygen and cause suffocation or fire hazards.

### **Ethanol**

The alcohol in alcoholic drinks and some disinfectants, medicines, and fuels. Drinking too much can impair judgment, suppress breathing, lower blood sugar, cause dependence, and damage the liver and other organs.

### **Ethanol Metabolism**

The body's breakdown of alcohol, mainly in the liver, into acetaldehyde and then acetate. The process affects blood-alcohol levels, drug interactions, intoxication, hangovers, and long-term alcohol-related organ damage.

### **Ethanol Poisoning**

Harm from drinking a dangerous amount of alcohol, causing confusion, poor coordination, vomiting, slow or irregular breathing, low blood sugar, seizures, coma, or death. An unconscious or poorly breathing person needs emergency care.

### **Ethanolamine**

A chemical used in some industrial products, medicines, and cosmetic ingredients. Concentrated exposure can irritate or burn the skin, eyes, and airways; health effects depend on the product and amount of exposure.

### **Ether**

A family of chemicals containing an oxygen atom between carbon groups. Some are volatile solvents; diethyl ether was once an anesthetic but is highly flammable, and inhalation can cause irritation, dizziness, or unconsciousness.

### **Ethers**

A group of chemicals containing an oxygen atom between two carbon groups. Some are used as solvents or fuels; exposure can cause irritation, dizziness, unconsciousness, or fire and explosion hazards.

### **Ethics**

Principles used to decide what is right, fair, respectful, and responsible in healthcare and research. Medical ethics includes informed choice, privacy, avoiding harm, helping patients, and treating peo-



---

ple fairly.

### **Ethinyl Estradiol**

A synthetic estrogen used in many hormonal contraceptives and some other treatments. It can cause nausea or irregular bleeding and increases blood-clot risk, especially with smoking, migraine with aura, or certain medical conditions.

### **Ethiodized Oil**

An iodine-containing oil used as a contrast material in selected imaging or treatment procedures, including some lymphatic and liver procedures. It can affect thyroid function or rarely enter blood vessels and cause an embolism.

### **Ethion**

An organophosphate pesticide that can poison the nervous system by blocking acetylcholinesterase. Exposure may cause sweating, vomiting, pinpoint pupils, muscle twitching, weakness, breathing failure, or death and requires urgent treatment.

### **Ethionamide**

An antibiotic used with other medicines for drug-resistant or difficult-to-treat tuberculosis. It interferes with mycobacterial cell-wall production and may cause nausea, liver injury, nerve symptoms, or thyroid problems.

### **Ethmocephaly**

A severe birth defect in which the eyes are very close together and the nose is replaced or accompanied by a tube-like structure. It usually occurs with major brain-development abnormalities and is often life-limiting.

### **Ethmoid Bone**

A light, complex bone between the eye sockets that forms part of the nasal septum, nasal roof, and inner eye socket. It contains air cells and openings for smell nerves, so injury can affect smell or the eye.

### **Ethmoid Sinus**

One of several small air spaces in the ethmoid bone

between the nose and eye sockets. Inflammation or infection can cause nasal blockage, headache, or eye-area pain and may spread to nearby tissues.

### **Ethmoid Sinusitis**

Inflammation or infection of the air spaces between the nose and eyes. It may cause congestion, discharge, fever, headache, or pain around the eyes; eye swelling, vision changes, or severe illness requires urgent care.

### **Ethmoidal Air Cells**

Multiple small air-filled spaces within the ethmoid bone between the nose and eye sockets. They are part of the sinus system and can become blocked or infected, causing sinus pressure and discharge.

### **Ethmoidal Sinus**

A group of small, thin-walled air spaces in the ethmoid bone between the eye sockets and above the nasal cavity. They are divided into front and back groups according to where they drain.

#### **Alternate Names:**

Ethmoid Air Cells; Paranasal Ethmoid Sinus

### **Ethmoiditis**

Inflammation or infection of the ethmoid air spaces between the nose and eyes. It may cause fever, nasal blockage, discharge, facial pain, or swelling and can rarely spread to the eye socket or brain.

### **Ethnicity**

A social and cultural identity shaped by shared ancestry, history, language, traditions, or community; it is not a biological diagnosis and should not be used as a substitute for individual risk assessment.

### **Ethoglucid**

An older sclerosing substance used historically to treat selected vascular lesions or hemorrhoids; modern use is limited by tissue-injury risks and alternatives.

### **Ethosuximide**

An anticonvulsant used mainly for absence seizures;

---

common adverse effects include gastrointestinal symptoms, fatigue, and headache, with rare blood or skin reactions.

### **Ethoxzolamide**

A carbonic-anhydrase-inhibiting medicine that reduces formation of certain body fluids. It has been used for selected eye and metabolic conditions; possible effects include tingling, nausea, kidney stones, and electrolyte changes.

### **Ethyl Chloride**

A volatile liquid sprayed on skin to produce brief cooling and numbness before minor procedures, but misuse can cause cold injury or toxicity.

### **Ethyl Ether**

A highly flammable liquid formerly used as an anesthetic and still used as a laboratory solvent. Breathing its vapor can cause irritation, dizziness, drowsiness, or loss of consciousness; swallowing it can injure the lungs if it is aspirated. Exposure requires fresh air, fire precautions, and medical assessment when symptoms are severe.

### **Ethylene**

A colorless, flammable gas used in chemical manufacturing and to ripen fruit. High concentrations can displace oxygen and cause headache, dizziness, drowsiness, unconsciousness, or suffocation. Remove a person from exposure only when it is safe and seek emergency help for breathing or consciousness problems.

### **Ethylene Chlorohydrin**

A toxic industrial chemical used in some manufacturing processes. Exposure can irritate the eyes, skin, and airways and may affect the nervous system, liver, or kidneys. It can be absorbed through skin, so contaminated clothing should be removed safely and urgent poison-control advice obtained.

### **Ethylene Dibromide**

A toxic chemical formerly used as a soil fumigant and fuel additive. It can irritate the eyes and lungs and may damage the liver, kidneys, or nervous sys-

tem; long-term exposure is associated with cancer risk. Suspected exposure needs urgent poison-control or occupational-health assessment.

### **Ethylene Glycol**

A toxic alcohol found in some antifreeze and industrial products. Ingestion can cause severe metabolic acidosis, neurologic impairment, kidney injury, and death.

### **Ethylene Glycol Blood Test**

A blood test measuring ethylene glycol, a toxic alcohol found in some antifreeze products. Detection supports diagnosis of poisoning, but treatment must not wait for the result when the exposure and metabolic findings are strongly suggestive.

### **Ethylene Glycol Poisoning**

Poisoning after swallowing ethylene glycol, found in some antifreeze products. Early effects may resemble alcohol intoxication, followed by severe acid buildup, seizures, coma, and kidney failure; emergency antidote and sometimes dialysis are needed.

### **Ethylene Glycol Toxicity**

Toxic alcohol poisoning that progresses from inebriation to high-anion-gap metabolic acidosis and then acute kidney injury from calcium oxalate deposition. An osmolar gap, hypocalcemia, and urinary oxalate crystals may support the diagnosis; give fomepizole promptly and use hemodialysis for severe acidosis, renal failure, or significant toxic burden.

### **Ethylene Oxide**

A flammable gas used to sterilize heat-sensitive medical equipment and make other chemicals. It can irritate the skin and airways, cause nerve injury, and with repeated exposure increase cancer risk. Sterilized equipment must be adequately aerated before use, and workplace exposure requires strict controls.

### **Ethinodiol Diacetate**

A synthetic progestogen formerly used in some oral contraceptives. It helps prevent pregnancy mainly by suppressing ovulation and changing cervical mu-

---

cus and the uterine lining.

### **Etidronic Acid**

A medicine related to bisphosphonates that can slow the breakdown of bone by attaching to bone mineral. It has also been used to bind minerals in some nonmedical products; effects and safety depend on the use and dose.

### **Etilefrine**

A medicine that stimulates some of the body's adrenaline-like receptors and can raise blood pressure. It increases blood-vessel tightening and heart activity, so it may cause fast heartbeat, headache, or other effects and requires medical supervision.

### **Etiology**

The cause or set of causes of a disease or health condition, including genetic, infectious, environmental, immune, traumatic, and behavioural factors.

### **Etodolac**

A nonsteroidal anti-inflammatory drug used for pain and inflammation; gastrointestinal bleeding, kidney injury, cardiovascular events, and hypersensitivity are important risks.

### **Etomidate**

A short-acting intravenous anesthetic used for induction, particularly when cardiovascular stability is desired; it can cause myoclonus, nausea, and transient adrenal suppression.

### **Etonogestrel**

A man-made form of the hormone progesterone used in a small contraceptive implant placed under the skin of the upper arm. It mainly prevents ovulation and also thickens cervical mucus; irregular bleeding and other hormone-related effects can occur.

### **Etoposide**

A chemotherapy medicine that interferes with DNA repair and cell division. It treats several cancers but can cause low blood counts, infection, hair loss, nausea, mouth sores, or a later risk of leukemia.

### **Etoposide Phosphate**

A water-soluble form of etoposide that changes into etoposide inside the body. It provides another method of giving the chemotherapy medicine, with similar cancer treatment effects and potential toxicities.

### **Etoricoxib**

An anti-inflammatory pain medicine that selectively blocks an enzyme involved in pain and swelling. It may help arthritis or acute pain but can raise blood pressure and increase heart, kidney, fluid-retention, and stomach risks.

### **Eubacterium**

A group of bacteria that usually live without oxygen in the mouth and intestine. Most are part of normal microbial communities, but some can rarely cause opportunistic infection in seriously ill people.

### **Eubacterium Limosum**

A bacterium that lives without oxygen in the intestine and helps break down nutrients. It is rarely associated with human infection, and its importance in a sample depends on symptoms and the collection site.

### **Eubacterium Rectale**

A common intestinal bacterium that helps ferment dietary fiber and produce short-chain fatty acids. These products support the colon and contribute to normal gut microbial activity.

### **Eucalyptus Oil**

A concentrated plant oil containing eucalyptol and other chemicals. Swallowing it can poison the nervous system and cause vomiting, drowsiness, seizures, or breathing problems; it should not be taken internally without medical advice.

### **Eucalyptus Oil Overdose**

Poisoning after swallowing too much eucalyptus oil. Burning, vomiting, drowsiness, confusion, seizures, and breathing problems may occur; do not induce vomiting and seek emergency poison-control advice.

---

## Euchromatin

A relatively open form of DNA packaging inside the cell nucleus. Its structure makes nearby genes more accessible for reading and activity than DNA in tightly packed regions.

## Eugenicist

A person who supports eugenics, a discredited belief that society should control reproduction to alter inherited traits in a population. Eugenic policies caused coercion, discrimination, and forced sterilization and have no place in ethical healthcare.

## Eugenics

A discredited ideology that seeks to change a population by encouraging some people to reproduce and preventing others from doing so. Its history includes racism, forced sterilization, institutional abuse, and genocide.

## Eugenol Oil Overdose

Poisoning after excessive exposure to clove-derived eugenol oil. It may irritate the mouth and stomach and cause liver injury, bleeding, seizures, low blood pressure, slowed breathing, or coma.

## Euglycemic Diabetic Ketoacidosis

A severe metabolic complication of diabetes mellitus characterized by profound anion gap metabolic acidosis and elevated serum ketones occurring with normal or only mildly elevated blood glucose levels (less than 250 mg/dL), frequently triggered by SGLT2 inhibitor therapy, restricted carbohydrate intake, or acute physiological stress.

### Alternate Names:

EuDKA; Normal Glycemic Ketoacidosis

## Euphoria

An unusually intense feeling of happiness or well-being. It may occur normally, with mania, brain disease, or some medicines and drugs; sudden extreme euphoria with unsafe behavior needs clinical assessment.

## Eustachian Tube

A passage connecting the middle ear to the back of the nose. It opens during swallowing or yawning to equalize pressure, drain fluid, and help protect the middle ear from throat secretions.

## Eustachian Tube Dysfunction

Poor opening or closing of the passage between the middle ear and back of the nose. It can cause ear pressure, popping, muffled hearing, pain, or fluid after colds, allergies, or pressure changes.

## Eustachian Tube Patency

The ability of the passage between the middle ear and back of the nose to remain open enough to equalize pressure and drain fluid. Poor patency can cause fullness, muffled hearing, or middle-ear fluid.

## Eustachian Tube Problems

Problems with the ear-to-nose passage can cause fullness, popping, pain, muffled hearing, or repeated middle-ear fluid or infection. Colds, allergies, inflammation, and anatomy may contribute; persistent hearing loss needs examination.

### Alternate Names:

Eustachian Tube Dysfunction (ETD); Auditory Tube Obstruction

## Eustachian Valve

A crescentic embryological flap of tissue at the junction of the inferior vena cava and the inferior right atrium that in fetal life directs oxygenated umbilical venous blood across the foramen ovale into the left atrium; usually persists as a benign vestigial remnant in adults.

### Alternate Names:

Valve of the Inferior Vena Cava

## Eustachian-Tube Dysfunction

Failure of the passage between the middle ear and back of the nose to equalize pressure or clear fluid. It may cause fullness, popping, reduced hearing, or ear pain after infection, allergies, or pressure changes.

---

## Euthanasia

The deliberate ending of a person's life by another person to relieve suffering. It is ethically and legally distinct from allowing a natural death, stopping unwanted treatment, or providing comfort-focused palliative care.

## Euthanasia And Assisted Dying

Terms describing legally regulated practices intended to end life or help a person end life to relieve suffering. Laws and safeguards differ widely; they are distinct from palliative care and withdrawal of unwanted treatment.

## Euthyroid

Having normal thyroid function, usually shown by thyroid-stimulating hormone and thyroid-hormone levels within the laboratory's reference ranges. A person can be euthyroid even when another illness temporarily changes some thyroid test results.

## Euthyroid Goiter

An enlarged thyroid gland with normal thyroid-hormone production. It may result from iodine imbalance, autoimmune disease, nodules, medicines, or inherited factors and should be assessed if persistent, enlarging, or causing swallowing or breathing problems.

## Euthyroid Hyperthyroxinemia

Elevated circulating thyroxine with an otherwise euthyroid clinical state. Binding-protein changes, inherited transport abnormalities, assay interference, or medication effects can produce the laboratory pattern without true thyroid hormone excess.

## Euthyroid Sick Syndrome

Temporary changes in thyroid blood tests during severe illness, starvation, or major surgery without a primary thyroid disorder. Treatment usually focuses on the underlying illness rather than thyroid hormone.

## Eutocia

Normal labor and birth progressing without major obstruction or complication. The term describes

the course of delivery, not a separate disease or treatment.

## Euvolemic Hyponatremia

Low blood sodium with no obvious clinical signs of excess or deficient body fluid volume. Common mechanisms include inappropriate antidiuretic hormone activity, adrenal insufficiency, hypothyroidism, and excess water intake.

## Evacuation

Removal of people from danger or removal of material from a body cavity, organ, or wound. In clinical care it may refer to clearing a hematoma, bowel, uterus, or other collection; the context determines the meaning.

## Evacuation Of Uterus

Removal of pregnancy tissue or other retained material from the uterus using medicines, suction, or a surgical procedure. It may treat miscarriage, abortion, retained placenta, or bleeding and carries risks of infection, bleeding, or uterine injury.

## Evaluation of Fetal Death

Assessment after suspected or confirmed fetal death includes ultrasound confirmation, maternal examination, laboratory testing, review of medical and pregnancy history, and discussion of placental or fetal examination when appropriate. Care also addresses delivery, bleeding or infection risk, grief, and future-pregnancy planning.

## Evans Syndrome

An autoimmune disorder characterized by the simultaneous or sequential development of autoimmune hemolytic anemia (positive direct antiglobulin test) and immune thrombocytopenic purpura, occasionally accompanied by autoimmune neutropenia.

### Alternate Names:

Evans's Syndrome; AIHA and ITP Syndrome

## Event Monitor

A portable device that records the heart's electrical rhythm when symptoms occur or at scheduled intervals over days or weeks. It can help detect

---

intermittent abnormal rhythms that a brief ECG may miss.

### Event-Free Survival

The length of time after treatment begins or a study starts before a defined event occurs, such as disease progression, relapse, a complication, or death. The exact event must be stated for comparisons to be meaningful.

### Everolimus

A medicine that suppresses a cell-growth pathway and immune activity. It is used for selected cancers, transplant care, and some genetic disorders; mouth sores, infection, high blood sugar, lung inflammation, and low blood counts may occur.

### Eversion

Turning the sole of the foot outward, away from the body's midline. The movement uses several ankle and foot joints and can be limited by injury, arthritis, or nerve and muscle disorders.

### Evidence-Based Practice

Care that combines the best available research, professional clinical judgment, and the patient's goals, values, and circumstances. It supports informed decisions while recognizing that evidence and individual needs may differ.

### Evisceration

Protrusion of internal organs through a surgical abdominal wound that has opened. This is a surgical emergency; the wound should be covered with sterile moist material, and urgent assessment is needed for repair and infection prevention.

### Evoked Potential

An electrical response recorded from the nervous system after a visual, sound, or touch stimulus. The test assesses whether signals travel normally along sensory pathways and may reveal nerve damage or slowed transmission.

### Evoked Potentials

Tests that record nervous-system responses to visual, sound, or touch stimuli. They can assess pathways for sight, hearing, or sensation and may help identify slowed signaling from disorders affecting nerves, the brain, or spinal cord.

#### Alternate Names:

VEP; BAER; SSEP; Evoked Responses

### Evolution

Gradual change in inherited traits of a population across generations. In clinical writing, the term can also describe how a disease, symptom, or treatment response changes over time.

### Evolutionarily Conserved Gene

A gene whose sequence or function remains similar across different species. This similarity suggests that the gene performs an important biological role, although it does not by itself prove a disease association.

### Ewart Sign

A physical-examination finding of a large fluid collection around the heart: dullness, reduced breath sounds, and bronchial breathing over the lower left back. Modern imaging is more reliable for confirming the cause.

### Ewing Sarcoma

A rare, aggressive cancer of bone or soft tissue, most often affecting children, teenagers, and young adults. It may cause persistent pain, swelling, fever, or a fracture and requires biopsy and specialist treatment.

### Ewing's Family of Tumors

Alternate name; see *Ewing Sarcoma*.

### Ewing's Sarcoma

A malignant tumor of bone or soft tissue, most common in children and young adults. It may cause localized pain, swelling, or fever and is diagnosed with imaging, biopsy, and molecular testing; treatment usually combines chemotherapy with surgery



---

and/or radiation.

### **Ewing's Tumour**

A rare aggressive cancer that usually begins in bone or nearby soft tissue in children or young adults. Persistent pain, swelling, fever, or an unexplained fracture may occur; diagnosis requires imaging and biopsy.

### **Ewing's Family of Tumors**

A group of aggressive small round-cell cancers, including Ewing sarcoma of bone and soft tissue, most often affecting children, adolescents, and young adults. Treatment commonly combines chemotherapy with surgery and/or radiation.

### **Exacerbate**

To make a disease, symptom, or medical problem worse. Infection, stress, exertion, allergens, medicines, or delayed treatment may exacerbate symptoms, depending on the condition.

### **Exacerbation**

A worsening or flare of a disease or its symptoms after a period of relative stability. Triggers may include infection, allergens, treatment interruption, exertion, or another physical or emotional stress.

### **Exacerbation Of COPD**

A sudden worsening of breathlessness, cough, or sputum beyond usual daily variation in chronic obstructive pulmonary disease. Infections and air pollution are common triggers; severe breathlessness, confusion, or blue lips requires urgent care.

### **Exaggerated Placental Site**

A noncancerous increase in placental cells at the place where the placenta attaches to the uterus. It may resemble a pregnancy-related cancer in a tissue sample, so an experienced pathologist must distinguish it from cancer.

### **Examination**

A systematic assessment of health using observation, touch, listening, measurements, and selected tests.

It may be general or focused on a complaint and helps identify normal findings, illness, injury, and urgency.

### **Examination Table**

A padded, adjustable surface that supports a person during examination, testing, minor procedures, or treatment. It should be stable, cleanable, accessible, and used with appropriate safety measures.

### **Exanthem**

A widespread skin rash, often caused by an infection or medicine. Its appearance, timing, fever, mucosal involvement, and other symptoms help clinicians determine whether it may be contagious or serious.

### **Exanthem Subitum**

A common viral illness in infants and toddlers, usually caused by human herpesvirus 6 or 7. It often causes three to five days of high fever followed by a pink rash as the fever settles.

#### **Alternate Names:**

Roseola Infantum; Sixth Disease

### **Exanthema**

A widespread skin eruption associated with an illness affecting the body. Viral infections are common causes, but medicines, bacteria, and immune disorders may also cause it; blisters, skin pain, mouth sores, or severe illness need prompt assessment.

### **Exanthema Subitum**

A common viral illness in infants and toddlers, usually caused by human herpesvirus 6 or 7. It often causes three to five days of high fever followed by a pink rash as the fever settles.

#### **Alternate Names:**

Roseola Infantum; Sixth Disease

### **Exanthematous Pustulosis**

A sudden medicine-related eruption of many small, noninfectious pustules on red skin, often with fever. It usually improves after the responsible medicine is stopped under medical guidance, but severe reactions require urgent assessment.

---

### Excavation

A hollow, depression, or removal of tissue or material. In medicine it may describe a normal body feature, a disease-related defect, or the surgical creation or clearing of a space.

### Exceptional Release

Permission to use a donated organ that does not meet usual acceptance criteria when clinicians judge it suitable for a particular recipient. The decision requires careful risk assessment, documentation, and informed consent.

### Excess Post-Exercise Oxygen Consumption

Increased oxygen use after exercise as the body returns to its usual metabolic and physiologic state.

### Excessive Bruising

Bruises that occur often, are unusually large, appear without clear injury, or last unusually long. Causes include medicines, fragile skin, low platelets, clotting disorders, liver disease, or other illness.

### Excessive Daytime Sleepiness

An excessive tendency to feel drowsy or fall asleep during waking hours, despite an opportunity for adequate nighttime sleep. It may occur with insufficient sleep, sleep disorders, medicines, medical or neurologic conditions, or mood disorders.

### Excessive Menstrual Bleeding

Alternate name; see *Heavy Menstrual Bleeding*.

### Excessive Scarring

Alternate name; see *Keloid*.

### Excessive Sweating

Sweating more than needed to cool the body, either all over or in specific areas such as the hands, feet, or armpits. It may result from hyperhidrosis, medicines, infection, hormones, or another illness.

### Excessive Tearing

Tears overflowing onto the face because the eyes

make too many tears or drainage is blocked. Irritation, allergy, infection, dry eye, eyelid problems, or a blocked tear duct may be responsible.

### Excessive Thirst

Polydipsia, an abnormally strong or persistent desire to drink, commonly associated with dehydration, diabetes mellitus, diabetes insipidus, medicines, or psychiatric conditions.

### Excessive Yawning

Excessive yawning is unusually frequent or difficult-to-control yawning. It may accompany sleep deprivation, excessive daytime sleepiness, medication effects, vasovagal reactions, or less commonly neurologic or cardiopulmonary disease; persistent unexplained episodes merit clinical evaluation.

### Exchange Transfusion

A procedure that repeatedly removes small amounts of a baby's blood and replaces them with compatible donor blood. It can rapidly reduce dangerous bilirubin or antibody-related anemia when other treatment is insufficient.

### Excise

To remove tissue, a growth, or a lesion by cutting it out. The removed material may be examined in a laboratory to identify the cause and determine whether abnormal cells reach the edges.

### Excision

Surgical removal of part or all of a lesion, tissue, or organ. The specimen may be examined to establish a diagnosis, and removal may also treat disease when the affected area can be safely removed.

### Excision Biopsy

Alternate name; see *Excisional Biopsy*.

### Excisional Biopsy

Removal of an entire lump or abnormal area, sometimes with a small margin of nearby tissue, so a laboratory can examine it. It may provide a diagnosis and, for selected small lesions, complete

---

treatment.

### **Excitation**

Activation or increased responsiveness of a cell, nerve, muscle, molecule, or body system after a stimulus. In nerves and muscles, it can start electrical signals that produce movement or other functions.

### **Excitation-Contraction Coupling**

The chain of events that links an electrical signal in a muscle cell to contraction. The signal causes calcium release inside the cell, allowing muscle proteins to interact and generate force.

### **Excitatory Neurotransmitter**

A chemical messenger that makes a receiving nerve cell more likely to generate an electrical signal. Examples include glutamate and, in some settings, acetylcholine; excessive activity can injure nerve cells.

### **Excitatory Synapse**

A junction between nerve cells where chemical signaling makes the receiving cell more likely to produce an electrical impulse. Balanced excitatory and inhibitory signals are needed for normal brain and nerve function.

### **Excitotoxicity**

The pathological process of neuronal injury and cell death resulting from excessive or prolonged synaptic stimulation by excitatory neurotransmitters, primarily glutamate, triggering massive intracellular calcium influx, mitochondrial dysfunction, reactive oxygen species generation, and apoptotic protease activation.

### **Excitotoxin**

A substance that overstimulates nerve cells and can damage or kill them. Excessive signaling may allow too much calcium into cells, contributing to injury in some brain diseases or poisoning states.

### **Exclamation Point Hair**

A short broken hair that is thinner at its outer end

and wider near the scalp. It is a characteristic finding of alopecia areata but can occasionally appear in other hair disorders.

### **Exclusion Diet**

A planned diet that removes a suspected food and later reintroduces it to see whether symptoms recur. It should be supervised when possible and must not replace emergency allergy evaluation when severe reactions or anaphylaxis are possible.

### **Excoriate**

To remove or injure the skin surface by scratching, rubbing, or picking. Repeated excoriation can cause open sores, infection, scarring, and difficulty healing.

### **Excoriation**

A superficial skin injury caused by scratching, rubbing, or picking. It may cause pain, bleeding, crusting, infection, or scarring and usually heals when the irritation or behavior stops.

### **Excoriation Disorder**

An obsessive-compulsive and related disorder involving recurrent skin picking that causes lesions, scarring, infection, or other tissue damage. Repeated attempts to reduce or stop the picking are unsuccessful, and the behavior causes significant distress or impairment; it is not better explained by a skin disease, substance, or another mental disorder.

#### **Alternate Names:**

Skin-Picking Disorder

### **Excreta**

Material expelled from the body, including stool, urine, sweat, vomit, and respiratory secretions. Some excreta carry infectious organisms, so hand-washing, protective equipment, cleaning, and safe disposal help prevent spread.

### **Excretion**

Removal of waste, medicines, or other substances from the body, mainly through urine, bile, breath, stool, sweat, or breast milk. Kidney and liver function strongly affect how long many substances remain in the body.

---

## Executive Function

Mental skills used to plan, focus, remember instructions, control impulses, solve problems, make decisions, and switch between tasks. Problems may follow brain injury, dementia, developmental conditions, or mental illness.

## Executive Functions

Mental abilities that support planning, organization, attention, memory, decision-making, self-control, and flexible behavior. Difficulties can affect work, school, relationships, and self-care and may have neurologic, developmental, or psychiatric causes.

### Alternate Names:

Executive Control

## Exemestane

A medicine that lowers estrogen production by blocking an enzyme used after menopause. It treats selected hormone-sensitive breast cancers and may cause hot flashes, joint pain, bone loss, fatigue, or abnormal cholesterol.

## Exenatide

An injectable medicine that imitates a gut hormone, increasing insulin release when glucose is high, reducing glucagon, and slowing stomach emptying. It treats type 2 diabetes but may cause nausea, dehydration, or pancreatitis.

## Exencephaly

A severe birth defect in which much of the developing brain lies outside the skull because the skull and protective coverings did not form normally. It usually progresses to anencephaly and is not compatible with long-term survival.

## Exenteration

Extensive surgery removing organs and tissues from a body compartment, such as the pelvis or eye socket. It is reserved for selected advanced cancers or destructive disease and can cause major functional and emotional effects.

## Exercise

Planned physical activity intended to improve or maintain strength, endurance, flexibility, balance, function, or health. Aerobic, resistance, and balance activities should be chosen and increased according to a person's abilities and medical risks.

### Alternate Names:

Physical Exercise

## Exercise And Age

The way physical activity needs and responses may change with age; safe exercise should match fitness, health, and ability.

## Exercise And Immunity

The relationship between physical activity and immune function; regular moderate activity generally supports health, while excessive exertion may increase short-term illness risk.

## Exercise Headaches

Headaches occurring during or after strenuous physical activity. Primary exercise headaches are recurrent and usually not caused by an underlying disorder, but a first, abrupt, severe, or atypical exertional headache requires prompt medical assessment to exclude secondary causes such as subarachnoid hemorrhage, arterial dissection, or reversible cerebral vasoconstriction syndrome.

## Exercise Intolerance

An inability to do as much physical activity as expected because of unusual tiredness, breathlessness, chest discomfort, leg pain, weakness, or dizziness. Heart, lung, blood-vessel, muscle, nerve, blood, metabolic, and conditioning problems can cause it.

## Exercise Prescription

An individualized physical-activity plan specifying the type, frequency, intensity, duration, and progression of exercise. It is adapted to a person's goals, fitness, medical conditions, symptoms, and safety needs.

---

### Exercise Stress Test

A test that records heart rate, rhythm, blood pressure, and symptoms while a person walks on a treadmill or pedals a stationary bicycle. It assesses exercise capacity and may suggest reduced heart blood flow or an abnormal rhythm.

### Exercise Tolerance

The ability to perform physical activity before symptoms such as breathlessness, fatigue, chest discomfort, or pain limit activity.

### Exercise-Induced Anaphylaxis

A rare form of anaphylaxis triggered by physical exertion, sometimes only when exercise follows ingestion of a specific food or medicine. Symptoms may include flushing, hives, swelling, wheezing, gastrointestinal symptoms, low blood pressure, or collapse.

### Exercise-Induced Arrhythmias

Abnormal heart rhythms that begin or worsen during physical exertion. They may result from inherited electrical disorders, heart muscle disease, or reduced blood flow; fainting, chest pain, or palpitations during exercise needs urgent evaluation.

### Exercise-Induced Asthma

Airway narrowing during or soon after exercise, causing cough, wheezing, chest tightness, or breathlessness. It can occur with or without persistent asthma and is managed with an individualized prevention and treatment plan.

### Exercise-Induced Bronchoconstriction

Temporary narrowing of the airways during or after exercise, causing cough, wheezing, chest tightness, or shortness of breath. Cold, dry air can worsen it; diagnosis and preventive inhaler use should be guided clinically.

#### Alternate Names:

Exercise-Induced Asthma; Exercise-Induced Bronchoconstriction (Exercise-Induced Asthma)

### Exercises To Help Prevent Falls

Activities that improve leg strength, balance, flexi-

bility, walking, and confidence. A program should match ability and medical conditions; supervised practice may be safest for people with previous falls or severe weakness.

### Exfoliation Syndrome

An age-related eye disorder in which abnormal material collects on the lens and other front-eye structures. It increases the risk of glaucoma and can make cataract surgery more complicated, so regular eye checks are important.

### Exfoliative Cytology

Examination of cells that naturally shed from a body surface or are gently collected. It may help investigate infection, inflammation, or cancer, but abnormal or uncertain results may require tissue biopsy.

### Exfoliative Dermatitis

Alternate name; see *Erythroderma*.

### Exhalation

Breathing air out of the lungs. It is usually passive at rest as the lungs recoil, but muscles actively assist exhalation during exercise, coughing, or breathing difficulty.

### Exhaled Nitric Oxide

A breath test that estimates one type of airway inflammation. A raised result can support asthma assessment and suggest likely benefit from inhaled steroids, but smoking, infection, allergies, and current treatment can affect the result.

### Exhaustion

Extreme physical or mental tiredness that reduces the ability to function. It may follow exertion, poor sleep, illness, dehydration, anemia, hormone disorders, medicines, or stress; severe or persistent symptoms need evaluation.

### Exhibitionism

Sexual arousal from exposing one's genitals to an unsuspecting person. Repeated nonconsensual ex-

---

posure can harm others and becomes a diagnosable disorder when urges or actions cause significant distress, impairment, or risk.

### **Exhibitionistic Disorder**

A mental-health disorder involving repeated sexual arousal from exposing the genitals to an unsuspecting person, with action on the urges or clinically significant distress or impairment.

### **Exhumation**

Lawful removal of a buried body or remains for identification, further testing, investigation, or repatriation. It requires legal authorization and respectful handling; testing may include toxicology, DNA analysis, or examination of injuries.

### **Existential Distress**

Emotional suffering related to meaning, identity, purpose, freedom, mortality, or loss of control. It can occur during serious illness and may improve with empathetic communication, counseling, spiritual care, and symptom relief.

### **Exit Wound**

The wound where a projectile leaves the body. Its size and shape vary with the projectile, tissue, clothing, and bone, so appearance alone cannot prove the direction or identify every injury; complete examination is required.

### **Exocervix**

The part of the cervix that projects into the vagina and is covered mainly by flat squamous cells. It is examined during pelvic examination and cervical screening.

### **Exocrine**

Relating to glands that release substances through ducts onto a body surface or into a cavity. Examples include sweat, salivary, tear, and digestive glands.

### **Exocrine Gland**

A gland that sends its secretion through a duct to a body surface or internal cavity. Sweat, salivary, tear,

and digestive glands are examples; their secretions help protect, lubricate, or digest.

### **Exocrine Glands**

Glands that release substances through ducts onto a surface or into a body cavity. They include sweat, salivary, tear, and digestive glands and support protection, lubrication, digestion, and temperature control.

### **Exocrine Pancreas**

The part of the pancreas that releases digestive enzymes and bicarbonate into the small intestine. These secretions help break down fats, proteins, and carbohydrates and neutralize stomach acid.

### **Exocrine Pancreatic Insufficiency**

Failure of the pancreas to release enough digestive enzymes, causing poor digestion and absorption. It may lead to greasy stools, bloating, diarrhea, weight loss, and vitamin deficiency.

### **Exocrine Pancreatic Insufficiency EPI**

Alternate index label; see *Exocrine Pancreatic Insufficiency*.

#### **Alternate Names:**

EPI; Pancreatic Exocrine Insufficiency

### **Exocytosis**

The process by which a cell releases substances outside itself. Small sacs inside the cell join its outer membrane and release hormones, nerve signals, digestive enzymes, or other proteins.

### **Exogenous**

Coming from outside the body rather than being produced within it. An exogenous substance may be a medicine, hormone, nutrient, toxin, infection, or other externally introduced material.

### **Exogenous Cushing Syndrome**

A condition caused by prolonged or excessive corticosteroid treatment, producing effects similar to too much natural cortisol. Features may include weight gain, easy bruising, high blood pressure, diabetes,



---

weakness, and bone loss.

### **Exomphalos**

A birth defect in which abdominal organs protrude through the belly button into the base of the umbilical cord and are covered by a membrane. Newborns need protection of the sac and specialist surgical care.

### **Exon**

A segment of a gene that remains in the mature messenger RNA after unused sections are removed. It may contain instructions for making part of a protein or, in some genes, help regulate gene activity.

### **Exopeptidase**

An enzyme that removes amino acids from the end of a protein. Digestive exopeptidases help break dietary proteins into smaller units that the intestine can absorb.

### **Exophiala**

A group of darkly pigmented environmental fungi that can rarely cause skin, lung, eye, or widespread infection. Disease is more likely in people with weakened immunity or implanted medical devices.

### **Exophthalmic Goitre**

An enlarged thyroid associated with Graves disease, in which immune stimulation causes excess thyroid hormone. Graves disease may also cause eye dryness, pain, swelling, or bulging.

### **Exophthalmos**

The abnormal forward protrusion or displacement of one or both eyeballs from the orbit, most frequently caused by Graves disease (thyroid eye disease), orbital tumors, or inflammation.

#### **Alternate Names:**

Bulging Eyes; Proptosis; Ocular Proptosis

### **Exosomes**

Very small membrane-covered particles released by cells that carry proteins, fats, and genetic material to other cells. They help cells communicate and are

being studied for diagnosis and drug delivery, but many medical uses remain experimental.

### **Exostosis**

A benign growth of bone. It may be harmless, but location can cause pain, pressure, restricted movement, or blockage; repeated cold-water exposure can produce bony growths in the ear canal.

#### **Alternate Names:**

Surfer's Ear; Osteoma of External Auditory Canal

### **Exotoxins**

Poisonous proteins released by some bacteria that damage particular cells or disrupt body functions. They cause diseases such as botulism, tetanus, diphtheria, or cholera and may require urgent treatment.

### **Exotropia**

A form of misaligned eyes in which one eye turns outward. It may be occasional or constant and can cause double vision, reduced depth perception, or loss of vision development in children; eye assessment guides treatment.

### **Expanded Rubella Syndrome**

Alternate name; see *Congenital Rubella Syndrome*.

### **Expectant Treatment**

Careful observation with regular review instead of immediate medicine or surgery when immediate intervention is not necessary. It includes clear follow-up and warning signs because treatment may be needed if symptoms worsen or tests change.

### **Expectorants**

Medicines intended to loosen or increase airway mucus so it can be coughed up more easily. Benefits vary, and fluids, humidification, and treatment of the underlying illness may be more important than an expectorant.

### **Expectoration**

Coughing up and bringing mucus or phlegm out of the respiratory tract. Changes in amount, color, smell, blood, fever, or breathlessness may indicate

---

infection or another lung problem.

### Experience

A person's direct knowledge, perception, or emotional response to an event. In healthcare, patient experience includes how care is accessed, communicated, coordinated, and received.

#### Alternate Names:

Subjective Experience; Patient-Reported Experience

### Expiration

Movement of air out of the lungs, usually caused by relaxation of the breathing muscles and natural recoil of the lungs. During forceful breathing, abdominal and chest muscles actively push air out.

### Expiratory Grunting

A grunting sound made during breathing out, caused by partially closing the voice box. It can briefly help keep small air spaces open but, especially in a baby or child, often signals respiratory distress and needs assessment.

### Expiratory Reserve Volume

The maximum additional volume of air that can be actively and forcefully exhaled from the lungs at the end of a normal, quiet tidal expiration, normally measuring approximately 1.0 to 1.2 liters in healthy adults.

#### Alternate Names:

ERV

### Explant

To remove an implanted device, prosthesis, or tissue. The term can also mean tissue removed from an organism and maintained in laboratory culture for study.

### Explantation

Surgical removal of an implanted device or prosthesis. It may be needed because of infection, malfunction, migration, breakage, tissue injury, intolerance, or completion of temporary treatment.

### Exploratory Laparotomy

Open surgery to enter the abdomen and directly find or treat a serious problem such as bleeding, perforation, blocked bowel, or loss of blood flow when less-invasive tests or treatment are insufficient.

### Exploratory Operation

Surgery performed to inspect an area directly when scans, tests, or less-invasive procedures cannot establish the cause or extent of disease. Tissue may be sampled and treatment may be performed during the operation.

### Exposure

Contact with a physical, chemical, biological, or psychological agent. Health risk depends on what the agent is, how much contact occurred, how it entered the body, and how long exposure lasted.

### Exposure Route

The way a substance enters the body, such as by breathing, swallowing, skin contact, injection, or the eyes. The route affects how quickly and how much of the substance is absorbed and how toxic it may be.

### Expression

In genetics, the process by which a gene's information is used to make RNA or protein. In clinical examination, expression can also mean the outward display of emotion or symptoms; altered gene activity can contribute to disease.

### Expression Hopping

A shift from one form of addictive behavior to another, such as replacing substance use with compulsive gambling. The underlying loss of control and harm may persist even when the specific behavior changes.

### Expressivity

The degree to which a genetic variant's effects appear in a person. Variable expressivity means that people with the same variant may have different features or different severity of the associated condition.

---

## Exsanguination

Life-threatening loss of blood from severe internal or external bleeding. It can rapidly cause shock, organ failure, and death; immediate pressure or bleeding control, emergency care, and blood replacement may be needed.

## Exserohilum

A group of darkly pigmented molds that can rarely cause sinus, eye, skin, or invasive infection. Serious disease is more likely in people with weakened immunity and requires specialist testing and treatment.

## Extend

To straighten a joint or increase the angle between two bones. For example, extending the elbow moves the forearm away from the upper arm toward a straight position.

### Alternate Names:

Extension

## Extension

Movement that straightens a joint or increases the angle between bones, such as straightening the knee or elbow. Excessive or forced extension can strain or tear muscles, tendons, ligaments, cartilage, or the joint.

## Extensively Drug-Resistant Tuberculosis

Tuberculosis caused by bacteria resistant to several key medicines, including isoniazid, rifampin, a fluoroquinolone, and at least one other important medicine. It is difficult to treat and requires specialist, susceptibility-guided therapy.

### Alternate Names:

XDR-TB; XDR Tuberculosis

## Extensor

A muscle or other structure that straightens a joint or increases the angle between bones. Extensor muscles commonly act opposite flexor muscles and help position limbs and fingers.

## Extensor Digitorum

A forearm muscle whose tendons straighten the

fingers and help extend the wrist. Overuse near its origin can contribute to outer-elbow pain commonly called tennis elbow.

## Extensor Plantar Response

An abnormal neurological reflex in which firm tactile stroking of the lateral plantar border of the foot produces slow extension (dorsiflexion) of the great toe accompanied by fanning of the other toes, indicating corticospinal (pyramidal) tract impairment.

### Alternate Names:

Positive Babinski Sign; Babinski Reflex

## Extent

The size, severity, range, or spread of a disease, injury, symptom, or treatment effect. Describing extent helps determine prognosis, further testing, and treatment.

## External Auditory Canal

The passage from the outer ear to the eardrum. It is lined with skin, hair, and wax-producing glands that help protect the ear, but it can become blocked, irritated, or infected.

## External Beam Partial Breast Irradiation

An accelerated conformal radiation therapy technique delivered via external beam linac that targets exclusively the lumpectomy cavity margin following breast-conserving surgery, sparing the remainder of the ipsilateral breast tissue in selected early-stage breast cancer patients.

### Alternate Names:

EB-APBI; Accelerated Partial Breast Irradiation; Targeted Breast Radiotherapy

## External Beam Radiation Therapy

Cancer treatment delivered from a machine outside the body to a carefully planned area. It can damage cancer cells while limiting nearby tissue exposure; fatigue, skin changes, and organ-specific effects may occur.

## External Capsule

A thin bundle of nerve fibers in the brain near

---

the basal ganglia. It carries signals between the cerebral cortex and deeper brain regions involved in movement, attention, and other functions.

### **External Carotid Artery**

A major branch of the neck artery that supplies blood to much of the face, scalp, neck, and parts of the brain coverings through several branches.

### **External Cephalic Version**

A procedure in which a clinician applies gentle pressure through the maternal abdomen to turn a fetus from a breech or transverse presentation into a head-down (cephalic) position for vaginal delivery. It is typically performed near term with continuous ultrasound and fetal heart rate monitoring.

#### **Alternate Names:**

ECV; Cephalic version.

### **External Ear**

The visible part of the ear and the ear canal leading to the eardrum. It collects sound, directs it inward, and helps protect the middle and inner ear.

### **External Fixation**

Stabilization of a broken bone or injured limb with pins or wires connected to a frame outside the body. It may be temporary or definitive, especially when swelling, wounds, or contamination prevent internal fixation.

### **External Iliac Artery**

A major artery carrying blood from the pelvis toward the leg. It becomes the femoral artery below the groin and gives branches to the lower abdominal wall and pelvic region.

### **External Iliac Vein**

A continuation of the femoral vein above the inguinal ligament that joins the internal iliac vein to form the common iliac vein.

### **External Incontinence Devices**

Products worn or placed outside the body to collect urine or stool or protect clothing from leakage. Ex-

amples include urinary sheaths, absorbent products, and fecal collection systems; skin care and correct fitting help prevent injury.

### **External Jugular Vein**

A visible vein running along the side of the neck that drains blood from the scalp and face toward the chest. Its fullness can provide clues about pressure in the right side of the heart.

### **External Nose**

The visible nose formed by bone, cartilage, skin, and soft tissue. It shapes the nasal passages, helps warm and filter inhaled air, supports smell, and contributes to facial structure.

### **External Oblique Muscle**

The broad, superficial muscle on each side of the abdomen. It compresses abdominal contents and helps bend and rotate the trunk; its lower border contributes to the groin ligament and canal.

### **External Otitis**

Inflammation or infection of the ear canal outside the eardrum, often called swimmer's ear. Moisture, scratching, or skin disease may contribute, causing pain, itching, discharge, swelling, or reduced hearing.

#### **Alternate Names:**

Otitis Externa; Swimmer's Ear

### **External Resorption**

Loss of tooth root or other tooth structure caused by a process starting outside the tooth. Injury, pressure, inflammation, or replacement by bone may contribute; treatment depends on location and severity.

### **External Ventricular Drain**

A critical neurosurgical bedside procedure in which a flexible plastic catheter is surgically placed through a small burr hole in the skull directly into one of the fluid-filled cavities (lateral ventricles) of the brain. The catheter connects to an external closed drainage and pressure-transducer system. An external ventricular drain serves two vital functions:

---

continuously monitoring intracranial pressure (the pressure inside the skull) and therapeutically draining excess cerebrospinal fluid or blood. It is essential in managing acute hydrocephalus, severe traumatic brain injury, subarachnoid hemorrhage, intraventricular hemorrhage, or intracranial hypertension, preventing brain herniation and death.

**Alternate Names:**

EVD; Ventriculostomy; External Ventricular Drainage; Ventricular Catheter

**Extinction**

In learning, the fading of a learned response when reinforcement stops. In neurology, it can mean failing to notice one side of the body or space when both sides are stimulated, suggesting an attention disorder.

**Extirpation**

Complete surgical removal of a tissue, organ, growth, or foreign body. The term describes the extent of removal rather than a particular disease or surgical technique.

**Extra-Dural**

Located outside the dura mater, the tough protective membrane around the brain and spinal cord. The term may describe a space, collection of blood, injection, or other structure outside this membrane.

**Extra-Gastrointestinal Stromal Tumor**

A tumor with features of a gastrointestinal stromal tumor that begins outside the digestive tract, often in abdominal supporting tissues. Diagnosis uses imaging, tissue examination, and molecular tests to guide cancer treatment.

**Extraamniotic**

Located outside the amniotic sac surrounding a fetus. The term may describe a procedure, device, or fluid placed between the amnion and nearby tissues during pregnancy care.

**Extracapsular Cataract Surgery**

Cataract surgery that removes the cloudy lens while leaving most of its surrounding capsule in place. An

artificial lens is usually inserted into the remaining capsule to restore focusing ability.

**Extracellular**

Located outside cells. Extracellular fluid, proteins, ions, and signals surround cells and help provide nutrients, remove waste, support tissues, and coordinate cell activity.

**Extracellular Exosome**

A tiny membrane-covered vesicle released by a cell that can carry proteins, fats, and genetic material to other cells. Exosomes help cells communicate and are being studied as disease markers and treatment vehicles.

**Extracellular Fluid**

Fluid outside cells, including blood plasma and the fluid between cells. It carries nutrients, hormones, and waste and is regulated mainly by the kidneys, circulation, and hormones.

**Extracellular Matrix**

A network of proteins and other substances outside cells that supports tissues and helps cells attach, communicate, grow, and repair injury. Its composition differs among bone, skin, muscle, and other organs.

**Extracellular Pathogens**

Microorganisms that live and multiply outside human cells. Examples include Streptococcus, Staphylococcus, and some Escherichia coli. They can damage tissues directly or through toxins, and the immune system usually fights them with antibodies and other defenses.

**Extracellular Region**

The area outside cells. It contains fluid, minerals, proteins, and signals that support tissues and influence how nearby cells grow, move, and function.

**Extracellular Space**

The fluid- and matrix-filled area outside cells. Nutrients, waste products, hormones, and signaling

---

molecules move through it, and its composition helps control cell survival, movement, and function.

### **Extracellular Vesicle**

A membrane-bound particle released by a cell that transfers molecular cargo between cells and is studied in physiology, disease, biomarkers, and drug delivery.

### **Extracorporeal**

Occurring or performed outside the body; extracorporeal circulation and hemodialysis route blood through an external machine.

### **Extracorporeal Circulation**

Movement of a person's blood through equipment outside the body and back again. Heart-lung machines, dialysis, and some life-support systems use it to add oxygen, remove waste, or support circulation; bleeding, clotting, infection, and organ injury are possible.

### **Extracorporeal Membrane Oxygenation**

An advanced, temporary mechanical life-support technique that drains deoxygenated blood from the body, circulates it through an external artificial membrane lung to add oxygen and remove carbon dioxide, and pumps it back into the bloodstream. It is indicated for severe, life-threatening respiratory failure or cardiogenic shock refractory to standard mechanical ventilation and intensive care. It is utilized in two main configurations: veno-venous (VV-ECMO) for isolated respiratory support, and veno-arterial (VA-ECMO) for combined hemodynamic and respiratory support. Major risks include hemorrhage from systemic anticoagulation, thromboembolism, hemolysis, cannula-site infection, and limb ischemia.

#### **Alternate Names:**

ECMO; Extracorporeal Life Support; ECLS; VV-ECMO; VA-ECMO

### **Extracorporeal Shock Wave Lithotripsy**

A treatment that uses focused shock waves from outside the body to break kidney or ureter stones into smaller pieces that can pass in urine. Bruis-

ing, bleeding, pain, blockage, or incomplete stone removal can occur.

### **Extracranial**

Located or occurring outside the skull or cranial cavity. For example, extracranial arteries supply the scalp and face, while extracranial injury affects tissues outside the skull and brain.

### **Extracranial Hematoma**

A collection of blood outside the skull or cranial cavity, usually caused by bleeding into soft tissues after injury.

### **Extract**

A concentrated preparation made by separating active substances from a plant, herb, food, or other source. Strength, purity, interactions, and safety vary widely, especially in unregulated supplements.

#### **Alternate Names:**

Botanical Extract; Pharmacological Extract

### **Extraction**

Removal of a tooth, tissue, foreign object, or other material from the body using a procedure. The method and risks depend on what is removed, where it is located, and the person's health.

### **Extragenital Germ Cell Tumors**

Germ-cell tumors arising outside the ovaries or testes, commonly in the mediastinum, retroperitoneum, or central nervous system. They may be seminomatous or nonseminomatous and are evaluated with imaging, tumor markers, pathology, and assessment for a gonadal primary.

### **Extrahepatic Biliary Atresia**

Alternate name; see *Biliary Atresia*.

### **Extrahepatic Cholestasis**

Reduced or blocked bile flow in the ducts outside the liver. Gallstones, scarring, inflammation, or a tumor may cause yellow skin, dark urine, pale stools, itching, pain, or infection and may need urgent treatment.



---

### **Extramammary Paget Disease**

A rare skin cancer usually affecting the vulva, anus, scrotum, or nearby skin. It often appears as a persistent itchy, sore, red, or scaly patch and requires biopsy to confirm diagnosis and assess for related cancer.

### **Extramammary Paget's Disease**

A rare skin cancer involving gland-bearing skin outside the breast, often around the vulva, anus, or penis. It may appear as a persistent itchy, red, scaly, or oozing patch and requires biopsy.

### **Extramedullary Hematopoiesis**

The making of blood cells outside the bone marrow, usually in the liver or spleen. It can happen when the bone marrow is damaged, crowded, or unable to make enough blood cells.

#### **Alternate Names:**

EMH; Extramedullary Blood Formation

### **Extranodal**

Located outside the lymph nodes. An extranodal disease or lymphoma affects an organ or other tissue, such as the stomach, skin, lung, bone, or brain.

### **Extranodal Marginal Zone Lymphoma**

A usually slow-growing cancer of B lymphocytes arising in tissues such as the stomach, salivary glands, thyroid, lungs, or eye. Symptoms depend on the organ involved and treatment may include observation, infection treatment, or cancer therapy.

#### **Alternate Names:**

Extranodal Marginal Zone Lymphoma MALT Lymphoma Mucosa-Associated Lymphoid Tissue Lymphoma

### **Extranodal Natural Killer T-Cell Lymphoma**

An aggressive lymphoma arising from immune cells, often affecting the nose and upper throat. It is strongly associated with Epstein-Barr virus and may cause blockage, bleeding, swelling, fever, or tissue destruction.

### **Extraocular**

Outside the eyeball or involving the muscles, nerves, and tissues around it. Extraocular structures move the eye, protect it, support vision, and connect it with the brain and face.

### **Extraocular Muscle**

One of the muscles outside the eyeball that moves the eye in different directions. Weakness or nerve injury can cause double vision, eye misalignment, or difficulty looking in a particular direction.

### **Extraocular Muscle Function Testing**

Examination of eye movements to assess the muscles and nerves that move the eyes. The person follows targets in different directions; abnormal movement can indicate muscle, nerve, brain, or eye-socket disease.

### **Extraocular Muscles**

Six muscles outside each eyeball that move the eyes up, down, inward, outward, and in rotation. Weakness or nerve injury can cause double vision, eye misalignment, or difficulty looking in a particular direction.

### **Extrapulmonary Manifestation Of Tuberculosis**

Tuberculosis affecting tissues outside the lungs, such as lymph nodes, bones, kidneys, the brain coverings, abdomen, or reproductive organs. Symptoms vary by site and treatment requires multiple antibiotics.

### **Extrapyramidal Disorder**

A disorder of brain circuits that regulate movement, posture, and muscle tone. It may cause stiffness, slowness, tremor, abnormal movements, or restlessness and can result from disease or medicines.

### **Extrapyramidal Symptoms**

Movement problems caused by medicines or diseases affecting movement circuits. They include stiffness, tremor, slowed movement, restlessness, sudden spasms, or involuntary movements, commonly after dopamine-blocking medicines.

---

## Extrapyramidal System

Interconnected brain structures and pathways, including the basal ganglia, that help control automatic movement, posture, muscle tone, and coordination. Damage or medicine effects can produce abnormal movements.

## Extrapyramidal Tracts

Nerve pathways outside the main voluntary movement tract that influence posture, muscle tone, automatic movements, and coordination. Disturbance can cause stiffness, tremor, slowness, or involuntary movements.

## Extraskkeletal Chondroma

A benign cartilage-forming lump arising in soft tissue rather than from bone, often in the hands or feet. It may calcify, cause pain or pressure, and sometimes needs removal to confirm the diagnosis.

## Extraskkeletal Mesenchymal Chondrosarcoma

A rare aggressive cancer of soft tissue that contains malignant cartilage. It may arise outside bone and requires specialist biopsy, staging, and combined cancer treatment.

## Extraskkeletal Osteosarcoma

A rare aggressive cancer that produces bone-like material in soft tissue rather than beginning inside bone. It usually affects adults and requires imaging, biopsy, staging, and specialist treatment.

## Extrasystole

Alternate name; see *Premature Ventricular Contractions*.

## Extrauterine Pregnancy

A pregnancy implanted outside the normal uterine cavity, most often in a fallopian tube. It can rupture and cause life-threatening internal bleeding; one-sided pain, fainting, or bleeding during pregnancy requires urgent care.

## Extravasation

Leakage of blood, medicine, urine, or another body fluid from a vessel or duct into nearby tissues. It can cause swelling, pain, inflammation, infection, or tissue death; treatment depends on the substance, location, and severity.

## Extravascular Hemolysis

The early removal and breakdown of red blood cells mainly in the spleen and liver. It can cause anemia, yellowing of the skin or eyes, and an enlarged spleen.

### Alternate Names:

Reticuloendothelial Hemolysis; Splenic Erythrocyte Clearance

## Extraventricular Drain

A temporary tube placed through the skull into a fluid-filled space in the brain. It can measure pressure and drain extra brain fluid or blood, especially when fluid buildup is dangerous.

### Alternate Names:

EVD; Ventriculostomy Catheter

## Extremities

The arms and legs, including their bones, joints, muscles, nerves, blood vessels, and skin. Problems in an extremity may affect movement, sensation, circulation, or function.

## Extremity

A limb of the body. The upper extremity includes the shoulder, arm, forearm, wrist, and hand; the lower extremity includes the hip, thigh, leg, ankle, and foot.

## Extremity Angiography

Imaging of blood vessels in an arm or leg, usually after contrast is introduced. It can show narrowing, blockage, aneurysm, abnormal connections, or injury and may sometimes be combined with treatment.

## Extremity Hardware Removal

Surgical extraction of orthopedic implants such as plates, screws, intramedullary rods, or wires from

---

an arm or leg. Typically indicated for persistent localized pain, implant infection, mechanical irritation, soft-tissue impingement, nonunion, or planned subsequent procedures.

**Alternate Names:**

Orthopedic Hardware Removal; Implant Removal; Plate and Screw Removal

**Extremity X-Ray**

An X-ray image of an arm or leg used to look for fractures, joint alignment problems, arthritis, infection, tumors, foreign objects, or other bone abnormalities.

**Extrinsic**

Coming from outside a structure, organ, or person. An extrinsic cause may be an external injury, medicine, infection, toxin, environmental exposure, or other outside influence.

**Extrinsic Allergic Alveolitis**

An immune-mediated lung disease caused by repeated inhalation of organic dusts, molds, bird proteins, or other environmental substances. It can cause cough, breathlessness, fever, and, with ongoing exposure, permanent lung scarring; it is also called hypersensitivity pneumonitis.

**Extrinsic Factor**

An influence from outside a person, tissue, or body process. Examples include medicines, infections, toxins, temperature, nutrition, stress, and environmental exposure; effects depend on the type and amount.

**Extrinsic Ureteral Obstruction**

Blockage of a ureter caused by pressure from outside the ureter, such as a tumor, scar tissue, enlarged lymph nodes, blood-vessel abnormality, or nearby inflammation. It can reduce urine drainage and cause kidney swelling or loss of kidney function.

**Alternate Names:**

Extrinsic Ureteral Compression; Secondary Ureteral Obstruction

**Extubate**

To remove a breathing tube after a person can breathe adequately, protect the airway, clear secretions, and maintain safe oxygen levels without the ventilator.

**Extubation**

Removal of a breathing tube from the windpipe after a person can breathe independently, keep the airway open, clear secretions, and maintain adequate oxygen and carbon-dioxide levels.

**Extubation Criteria**

Findings used to judge whether a person can safely have a breathing tube removed. They include adequate breathing and oxygenation, stable circulation, alertness, airway protection, effective cough, and recovery from muscle-paralyzing medicines.

**Exudate**

Protein-rich fluid that leaks from inflamed or injured blood vessels into tissues or a body cavity. Its presence usually indicates inflammation, infection, or tissue injury rather than pressure alone.

**Exudative Pleural Effusion**

A protein-rich collection of fluid between the lung and chest wall caused by local inflammation or injury. Pneumonia, cancer, tuberculosis, and blood clots are possible causes; testing the fluid helps guide treatment.

**Eye**

The organ that detects light and forms vision. It includes the cornea, lens, iris, retina, optic nerve, and supporting tissues within the eye socket; disease can affect sight, movement, comfort, or appearance.

**Eye Allergies**

An allergic reaction affecting the eyes and often the inner eyelids. It usually causes itching, redness, watering, burning, or swollen eyelids and is commonly called allergic conjunctivitis.

**Alternate Names:**

Eye Allergy Ocular Allergy

---

## Eye And Orbit Ultrasound

An ultrasound scan of the eye and surrounding socket. It can assess the retina, optic nerve, eye muscles, masses, bleeding, or other structures when the view through the front of the eye is obscured.

## Eye Bank

An organization that recovers, evaluates, stores, and distributes donated corneas or other eye tissue for transplantation, research, or education. Donor screening and tissue testing help protect recipients.

## Eye Bleed

Bleeding on, inside, or around the eye. A bright-red painless patch on the white of the eye is often superficial, but bleeding inside the eye, pain, injury, or vision change requires prompt assessment.

## Eye Cancer

A cancer that begins in the eyelid, surface of the eye, uvea, retina, or tissues around the eye. Warning signs can include a growing or changing spot, persistent redness, vision changes, eye pain, bulging, or a new pupil abnormality; diagnosis usually requires specialist examination and imaging or biopsy.

## Eye Color

The visible color of the iris, determined mainly by pigment and light scattering. A new, one-sided, uneven, or painful color change may indicate injury, inflammation, medicine effects, or another eye disorder.

## Eye Development

Formation and maturation of the eyes, optic nerves, and visual pathways before and after birth. Disruption can cause structural abnormalities, reduced vision, misalignment, or delayed development of visual skills.

## Eye Disease

Any disorder affecting the eye or visual pathways, including refractive errors, cataract, glaucoma, retinal disease, infection, inflammation, and tumors. Symptoms such as sudden vision loss, severe pain, flashes, or a curtain-like shadow require urgent assessment.

## Eye Disorder

A disease or structural problem affecting the eye or visual pathways. It may cause changed vision, pain, redness, discharge, double vision, field loss, or other findings and may require routine or urgent assessment.

## Eye Disorders

Diseases or structural problems affecting the eye, optic nerve, or visual pathways. They may cause blurred or lost vision, pain, redness, discharge, double vision, or field loss and can require routine or emergency assessment. Diseases affecting the eye or visual pathways, including refractive problems, cataracts, glaucoma, retinal disease, infection, inflammation, injury, and tumors. Symptoms may involve vision, pain, redness, discharge, or eye movement.

## Eye Drops

Liquid medicines, lubricants, or diagnostic preparations placed on the eye surface. They may relieve dryness or treat disease; contaminated bottles, shared drops, or incorrect use can cause injury or infection.

## Eye Emergencies

Conditions needing immediate eye assessment include sudden vision loss, chemical exposure, penetrating injury, severe pain, new neurologic symptoms, or rapidly spreading infection. Do not rub an injured eye or press on it.

## Eye Floaters

Alternate name; see *Floaters*.

## Eye Health

Maintaining vision and preventing avoidable eye disease through appropriate examinations, injury and ultraviolet protection, control of diabetes and blood pressure, safe contact-lens care, and prompt attention to new symptoms.

## Eye Hemorrhage

Bleeding on, inside, or around the eye. A small surface bleed may be painless, while bleeding in the

---

retina, vitreous, or eye socket can impair vision or cause pressure and needs medical assessment.

### **Eye Infection**

Infection of the eyelid, conjunctiva, cornea, internal eye, or surrounding tissues caused by bacteria, viruses, fungi, or parasites. Pain, light sensitivity, reduced vision, or swelling around the eye needs prompt care.

### **Eye Injury**

Damage to the eye or nearby tissues from blunt or penetrating trauma, chemicals, foreign objects, radiation, or heat. Vision change, severe pain, bleeding, or an embedded object requires urgent assessment.

### **Eye Irritation**

Burning, itching, redness, or watering caused by dryness, allergy, smoke, chemicals, infection, or a foreign particle. Chemical exposure requires immediate prolonged rinsing with clean water and urgent medical advice.

### **Eye Melanoma**

A cancer arising from pigment-producing cells in the eye, most often in the uvea. It may cause blurred vision, flashes, a dark spot, or no symptoms; treatment depends on size, location, spread, and vision potential.

#### **Alternate Names:**

Uveal Melanoma; Ocular Melanoma; Choroidal Melanoma

### **Eye Movement**

Coordinated movement of the eyes that directs gaze and keeps images aligned. It depends on the eye muscles, their nerves, and brain control; abnormal movement may cause double vision or imbalance.

### **Eye Muscle Repair**

Treatment to correct a damaged or misaligned eye muscle or its attachment, usually by eye-muscle surgery. It may improve alignment or double vision but does not always restore normal movement and carries surgical risks.

### **Eye Neoplasm**

A benign or malignant growth arising in the eye or surrounding tissues. Symptoms and treatment depend on the tissue involved, size, location, effect on vision, and whether the growth has spread.

### **Eye Pain**

Pain in or around the eye caused by surface irritation, infection, injury, inflammation, glaucoma, or disease behind the eye. Severe pain with vision loss, light sensitivity, nausea, or trauma requires urgent assessment.

### **Eye Prosthesis**

An artificial eye or implant used to restore appearance after an eye is lost or severely damaged. It generally does not restore sight; fitting, cleaning, and periodic specialist care are needed.

### **Eye Redness**

Redness caused by enlarged or inflamed blood vessels on or inside the eye. Irritation and conjunctivitis are common causes, but pain, light sensitivity, discharge, injury, or vision change may indicate serious disease.

### **Eye Retraction Syndrome**

Alternate name; see *Duane Retraction Syndrome*.

### **Eye Trauma**

Injury ranging from a surface scratch to a penetrating wound or ruptured eyeball. Rinse chemical exposure immediately, avoid pressure or rubbing, shield penetrating injuries, and obtain urgent care for vision change or severe pain.

### **Eye Twitching**

Repeated involuntary eyelid or eye-area spasms, commonly linked to fatigue, stress, caffeine, irritation, or eye strain. Persistent spasms, facial weakness, eye closure, or other neurologic symptoms need assessment.

### **Eyeball**

The roughly spherical organ of vision containing the

---

cornea, sclera, iris, lens, retina, internal fluids, and other tissues. It focuses light onto the retina, which sends visual signals through the optic nerve.

**Alternate Names:**

Globe

**Eyebrow**

A strip of hair above the eye socket that helps divert sweat, water, and particles away from the eye and contributes to facial expression.

**Eyeglass**

A lens mounted in a frame to correct focusing problems such as nearsightedness, farsightedness, astigmatism, or presbyopia. The prescription is selected after measuring vision and the eyes.

**Eyelash**

A hair growing from the eyelid edge that helps shield the eye from particles and triggers blinking when touched. Misrouted or infected lashes can irritate the eye surface.

**Eyelashes**

Short curved hairs along the eyelid edges that help protect the eye from dust, moisture, and bright light. Inward-growing lashes can rub the cornea and require treatment.

**Eyelid**

A movable fold of skin, muscle, and glands that protects the eye, spreads tears across its surface, and helps regulate light. Swelling, malposition, or weakness can impair protection and vision.

**Eyelid Bump**

A localized eyelid swelling, often a painful infected gland called a stye or a firmer blocked oil gland called a chalazion. Warm compresses may help; persistent, recurrent, or vision-affecting bumps need examination.

**Eyelid Cancer**

Cancer arising in the skin or other tissues of the eyelid, most commonly basal cell carcinoma. A

persistent or changing lump, sore, bleeding area, loss of eyelashes, or eyelid distortion needs ophthalmic or dermatologic evaluation.

**Eyelid Coloboma**

A birth defect causing a notch or missing section of the eyelid. It may leave the eye exposed, impair closure, and occur with other eye or facial abnormalities requiring specialist assessment.

**Eyelid Disorder**

A disease or structural problem of the eyelid, such as infection, inflammation, a blocked gland, malposition, weakness, or tumor. It may cause pain, swelling, poor eye protection, irritation, or visual problems.

**Eyelid Drooping**

Alternate name; see *Ptosis*.

**Eyelid Inflammation**

Alternate name; see *Blepharitis*.

**Eyelid Lift**

Alternate name; see *Blepharoplasty*.

**Eyelid Neoplasm**

A benign, precancerous, or cancerous growth of the eyelid. A persistent, enlarging, changing, bleeding, crusted, or ulcerated lesion needs examination by an eye or skin specialist.

**Eyelid Problems**

Disorders affecting eyelid skin, glands, muscles, nerves, position, or movement, including blepharitis, styes, chalazia, ptosis, entropion, ectropion, and tumors. Pain, vision change, severe swelling, or sudden onset requires prompt assessment.

**Eyelid Swelling**

Puffiness of one or both eyelids caused by allergy, infection, inflammation, injury, a blocked gland, or fluid accumulation. Fever, severe pain, impaired eye movement, bulging eye, or vision change may indicate an emergency.



---

## **Eyelid Twitch**

Alternate name; see *Eye Twitching*.

## **Eyelids**

Movable folds of skin, muscle, and glands that protect the eyes, spread tears, and help control light. Their position and movement are important for keeping the eye surface moist and safe.

## **Eyesight**

Alternate name; see *Vision*.

## **Eyestrain**

Tired, dry, sore, or uncomfortable eyes, sometimes with blurred vision or headache, after prolonged visual effort. Screen use, glare, poor lighting, dry eye, or an uncorrected focusing problem may contribute; rest often helps.

## **Ezetimibe**

An oral medicine that reduces absorption of cholesterol from the intestine. It lowers blood cholesterol alone or with another medicine and may cause diarrhea, abdominal discomfort, or rarely liver or muscle problems.

## **Idiopathic Edema**

Persistent swelling, usually of the legs, occurring without an identified heart, kidney, liver, venous, lymphatic, or medication-related cause. The diagnosis is one of exclusion and requires evaluation for more common causes before treatment.

## **Optical Aids**

Eyeglasses correct focusing errors, sunglasses reduce glare and ultraviolet exposure, and magnifiers enlarge nearby targets. Correct prescription, fit, lens protection, and the underlying eye condition determine the appropriate aid.

### **Alternate Names:**

Corrective Eyewear; Eyeglasses; Eyeglasses and Sunglasses; Eyeglasses, Sunglasses, and Magnifiers; Magnifiers; Sunglasses; Visual Aids



# F

## Fab Immunoglobulins

The target-binding part of an antibody. A laboratory-made Fab fragment can recognize and block a harmful substance or help remove it from the bloodstream in selected treatments.

## Fabry Disease

An inherited lysosomal storage disorder caused by reduced activity of alpha-galactosidase A because of a disease-causing variant in the *GLA* gene. Globotriaosylceramide and related substances accumulate in cells, especially in blood vessels, the kidneys, heart, nerves, and skin. Features may include burning pain in the hands and feet, reduced sweating, small dark-red skin spots, kidney disease, heart disease, or stroke. The condition is usually X-linked and can affect people of any sex.

## Fabry's Disease

Alternate spelling; see *Fabry Disease*.

## Face

The front part of the head, including the eyes, nose, mouth, cheeks, and jaw. Its muscles produce facial expression, its nerves provide movement and sensation, and its structures support breathing, eating, speech, and vision.

## Face and Brow Presentation

Abnormal fetal head presentations in which the face or brow leads instead of the vertex. Labor progress and delivery safety depend on fetal position, flexion, pelvis, gestational age, and whether the presentation converts; persistent brow or mentum-posterior face presentation may obstruct vaginal birth.

## Face Pain

Pain in the face that may arise from teeth, sinuses, nerves, eyes, jaw joints, skin, blood vessels, or other tissues. Location, duration, triggers, swelling, fever, and neurologic signs help identify the cause.

## Face Powder Poisoning

Illness after swallowing or inhaling face powder or its ingredients. Coughing, choking, vomiting, breathing difficulty, drowsiness, or persistent symptoms require poison-control or emergency advice; do not induce vomiting.

## Face Presentation

A birth position in which the baby's neck is extended so the face, rather than the top of the head, leads through the birth canal. The baby's chin direction helps determine whether vaginal birth is safe.

## Facelift

Cosmetic surgery that removes or repositions selected facial and neck skin and deeper tissues to reduce sagging. It does not stop aging and may cause bleeding, infection, nerve injury, scarring, or temporary swelling and numbness.

## Alternate Names:

Rhytidectomy

## Facet Joint

A small joint at the back of the spine connecting neighboring vertebrae and guiding movement. Arthritis, injury, or inflammation can cause localized neck or back pain and stiffness.

---

## Facet Joints

Small paired joints at the back of the spine that guide bending and rotation and help stabilize vertebrae. Wear, inflammation, or injury may cause localized neck or back pain.

## Facet Rhizotomy

A procedure that uses heat or another method to interrupt small nerves carrying pain signals from joints at the back of the spine. It may reduce selected chronic neck or back pain, but relief may be temporary.

## Facial Anatomy in Cutaneous Surgery

The layered arrangement of facial skin, fat, muscles, nerves, vessels, glands, and connective tissue that guides safe dermatologic and reconstructive procedures. Knowledge of relaxed skin tension lines and danger zones helps preserve function and appearance.

## Facial Asymmetry

A difference in the size, position, movement, or shape of the two sides of the face. It may be present from birth or result from injury, nerve weakness, muscle disease, swelling, or a growth.

## Facial Cleft

A birth defect caused by incomplete joining of facial tissues during development. It may affect the lip, nose, mouth, or other facial structures and can interfere with feeding, speech, breathing, hearing, or appearance.

## Facial Colliculus

An anatomical elevation on the floor of the fourth ventricle in the dorsal pons, formed by motor fibers of the facial nerve (cranial nerve VII) looping over the underlying abducens nucleus (cranial nerve VI); a lesion produces ipsilateral facial paralysis and lateral gaze palsy.

## Facial Edema

Swelling of the face caused by excess fluid in tissues. Allergy, infection, injury, kidney or heart disease, and blocked veins may cause it; sudden lip, tongue,

or throat swelling with breathing difficulty is an emergency.

## Facial Expression

The visible movement or position of facial muscles that communicates emotion, pain, intention, or other information. Reduced or unequal expression can indicate facial-nerve or muscle dysfunction.

## Facial Fractures

Breaks in one or more facial bones, such as the nasal, orbital, cheek, maxillary, or mandibular bones, usually caused by trauma. Pain, swelling, deformity, bleeding, double vision, malocclusion, or airway problems may occur.

## Facial Muscle

A muscle that moves the skin of the face for expression and helps with speaking, eating, breathing through the mouth, or closing the eyes. Weakness may result from nerve, muscle, or brain disease.

## Facial Neoplasm

A benign or malignant growth arising in facial skin, soft tissue, bone, salivary glands, nerves, or other structures. Symptoms and treatment depend on its tissue of origin, size, location, and spread.

## Facial Nerve

The seventh cranial nerve, which controls facial expression and carries taste from part of the tongue. It also contributes to tear and saliva production and helps protect the ear from loud sounds.

## Facial Nerve Palsy

Weakness or paralysis of facial muscles caused by dysfunction of the seventh cranial nerve. It may affect eye closure, smiling, taste, tears, or saliva; sudden weakness with arm or speech symptoms may indicate a stroke.

## Facial Nerve Palsy From Birth Trauma

One-sided facial weakness in a newborn caused by pressure on the facial nerve during birth. It often improves as the nerve recovers, but eye protection,

---

feeding, and other causes of weakness must be assessed.

### **Facial Nerve Problems**

Disorders affecting the nerve that controls facial movement and some taste, tear, saliva, and sound-protection functions. Causes include Bell palsy, stroke, infection, injury, or a tumor; sudden weakness needs emergency assessment.

#### **Alternate Names:**

Facial Nerve Disorder; Facial Nerve Dysfunction

### **Facial Nucleus**

A group of nerve cells in the pons, part of the brainstem, whose fibers form the facial nerve and control facial-expression muscles. Injury can cause weakness of the face and related functions.

### **Facial Pain**

Pain in the face that may arise from dental or sinus disease, jaw-joint problems, nerve disorders, infection, injury, or other causes. The pattern and associated symptoms guide evaluation.

### **Facial Palsy**

Weakness or paralysis of facial muscles, usually on one side, caused by facial-nerve dysfunction. Bell palsy is one cause, but stroke, infection, injury, Lyme disease, and tumors are also possible; sudden onset with other neurologic signs is urgent.

#### **Alternate Names:**

Bell's Palsy; Facial Paralysis

### **Facial Paralysis**

Partial or complete loss of facial-muscle movement caused by disease or injury of the facial nerve or its brain pathways. Sudden onset, especially with limb weakness or speech trouble, requires urgent stroke assessment.

### **Facial Paresis**

Weakness of muscles supplied by the facial nerve, causing reduced facial movement and sometimes poor eye closure, altered taste, or abnormal tearing. The cause may be nerve inflammation, injury,

infection, stroke, or a tumor.

### **Facial Soft Tissue Injuries**

Lacerations, abrasions, contusions, punctures, or avulsions involving the skin and underlying soft tissues of the face. Evaluation should consider injury to facial nerves, ducts, vessels, the eye, and deeper facial structures.

### **Facial Spasm**

Involuntary repeated tightening of facial muscles. It may affect one side around the eye or mouth and can result from nerve irritation, medicines, stress, or another neurologic disorder; persistent spasms need assessment.

### **Facial Swelling**

Enlargement of part or all of the face caused by infection, allergy, injury, dental or salivary disease, sinus disease, fluid retention, or a growth. Rapid swelling of the mouth or throat with breathing difficulty is an emergency.

### **Facial Tics**

Sudden, repeated, involuntary facial movements or sounds, such as blinking, grimacing, or nose wrinkling. They may increase with stress and occur in tic disorders; persistent or disruptive tics can be treated.

### **Facial Trauma**

Injury affecting facial skin, soft tissue, teeth, nerves, blood vessels, eyes, nose, or bones. Assessment must consider airway, bleeding, vision, brain injury, and structural damage.

### **Facial Vein**

A vein that drains blood from the superficial face toward the internal jugular vein. Connections with deeper veins mean facial infections can rarely spread toward the eye socket or brain.

### **Facies**

A characteristic facial appearance associated with a particular disease, inherited condition, or physio-

---

logic state. It can provide clues but is not diagnostic by itself and must be interpreted with other findings.

### **Facies Hippocratica**

The classic physical appearance of the face seen in patients near death, after severe prolonged exhaustion, or during terminal peritonitis, characterized by sunken hollow eyes, pinched nose, collapsed temples, cold clammy sweat, and a leaden, dusky or pallid complexion.

#### **Alternate Names:**

Hippocratic Facies; Hippocratic Face

### **Facilitated Diffusion**

Movement of a substance across a cell membrane through a channel or carrier protein, from higher to lower concentration. The process does not directly use cellular energy and helps move substances that cannot cross the membrane unaided.

### **Facio-Scapulo-Humeral Dystrophy**

Alternate name; see *Facioscapulohumeral Muscular Dystrophy*.

### **Facioscapulohumeral Dystrophy**

An inherited muscle disease causing gradually progressive weakness of the face, shoulder blades, upper arms, and sometimes legs or trunk. Onset and severity vary, and breathing, hearing, and eye complications may occur.

### **Facioscapulohumeral Muscular Dystrophy**

An inherited disorder causing progressive, often uneven weakness of facial, shoulder-blade, and upper-arm muscles. Symptoms may begin in adolescence or adulthood and can later involve the trunk, hips, or legs.

### **Factitious Disorder**

A mental-health disorder in which a person deliberately creates, exaggerates, or falsely reports illness to assume the sick role, without a clear external reward.

#### **Alternate Names:**

Munchausen Syndrome; Pathomimicry

### **Factitious Disorder by Proxy**

Alternate name; see *Factitious Disorder Imposed on Another*.

### **Factitious Disorder Imposed on Another**

A form of medical child abuse or caregiver falsification in which a parent or guardian deliberately exaggerates, fabricates, or directly induces physical or psychological symptoms in a dependent child, presenting the victim for repeated medical evaluations, invasive testing, or surgeries without apparent external gain other than assuming the role of a devoted caregiver.

#### **Alternate Names:**

Munchausen Syndrome by Proxy; MSBP; Pediatric Symptom Falsification

### **Factitious Disorders**

Alternate name; see *Factitious Disorder*.

### **Factitious Hyperthyroidism**

A state of excessive thyroid hormone in the body caused by ingestion of thyroid hormone or another external source rather than increased thyroid-gland production.

### **Factor**

A substance that contributes to a biological process or clinical result. In hematology, the term usually refers to blood-clotting proteins that work together to stop bleeding.

### **Factor H Deficiency**

A rare inherited or acquired problem in which a protein that keeps part of the immune system under control is missing or faulty. It can cause repeated infections or inflammation that damages small blood vessels, especially in the kidneys.

### **Factor I Deficiency**

An ambiguous term. In immunology, complement factor I deficiency causes uncontrolled alternative-pathway activation, low C3, recurrent pyogenic infections, and sometimes glomerulonephritis or atypical hemolytic uremic syndrome. In coagulation termi-



---

nology, factor I is fibrinogen; congenital fibrinogen deficiency causes bleeding. The intended meaning must be specified.

### **Factor II Assay**

A blood test measuring factor II activity, which helps assess the clotting pathway and investigate abnormal bleeding, prolonged prothrombin time, or suspected prothrombin deficiency or inhibition.

### **Factor IX**

A vitamin-K-dependent blood-clotting protein that works with factor VIII to form clots. Inherited deficiency causes hemophilia B, leading to prolonged or spontaneous bleeding, especially into joints and muscles.

### **Factor IX Assay**

A blood test measuring factor IX activity, used to diagnose or monitor factor IX deficiency, the cause of hemophilia B, and to guide replacement treatment.

### **Factor V**

A blood-clotting protein that helps generate thrombin and fibrin, the mesh that strengthens a clot. Too little can cause bleeding; an inherited change affecting its regulation can increase venous-clot risk.

### **Factor V Assay**

A blood test measuring how well factor V supports clotting. It helps investigate abnormal bleeding or clotting-test results and diagnose inherited deficiency or an acquired reduction from liver disease or other illness.

### **Factor V Deficiency**

An inherited or acquired bleeding disorder caused by too little active factor V. It may cause easy bruising, nosebleeds, heavy menstrual bleeding, or excessive bleeding after injury, dental work, surgery, or childbirth.

### **Factor V Leiden**

An inherited variant in the *F5* gene that makes factor V less responsive to activated protein C, a

natural protein that normally limits clotting. This leaves clotting active for longer and increases the tendency to form clots in veins, especially deep-vein thrombosis and pulmonary embolism. Many people who carry one copy never develop a clot; having two copies generally increases the risk further. Other factors, such as family history, pregnancy, estrogen exposure, surgery, injury, smoking, obesity, and immobility, can also affect risk.

### **Factor V Leiden Mutation**

Alternate name; see *Factor V Leiden*.

### **Factor VII**

A liver-made blood-clotting protein that starts clot formation when tissue is injured. Inherited or acquired deficiency can cause easy bruising, nosebleeds, or serious bleeding after injury or procedures.

### **Factor VII Assay**

A blood test measuring factor VII activity. It helps investigate a prolonged prothrombin time and identify inherited deficiency, liver disease, vitamin-K deficiency, or medicine-related reduction.

### **Factor VII Deficiency**

A bleeding disorder caused by too little or poorly functioning factor VII. Bleeding severity varies and may include bruising, nosebleeds, heavy periods, bleeding after surgery, or severe hemorrhage; prothrombin time is usually prolonged.

### **Factor VIII**

A blood-clotting protein that works with factor IX to form a stable clot. Inherited deficiency causes hemophilia A, with prolonged or spontaneous bleeding, often into joints, muscles, or other tissues.

### **Factor VIII Assay**

A blood test measuring factor VIII activity. It helps diagnose or monitor hemophilia A and investigate acquired reductions caused by inhibitors, liver disease, or other conditions.

---

## Factor X

A blood-clotting protein that helps start the final steps of clot formation. Too little factor X can cause prolonged or unexpected bleeding and is assessed with clotting tests and other blood studies.

## Factor X Assay

A blood test measuring factor X activity, a key clotting step shared by two clotting pathways. It helps investigate abnormal bleeding and diagnose or monitor inherited or acquired factor X deficiency.

## Factor X Deficiency

A rare inherited or acquired bleeding disorder caused by too little active factor X. Bleeding severity varies from bruising to serious hemorrhage, and both major clotting-time tests may be prolonged.

## Factor XI Deficiency

An inherited or acquired bleeding disorder caused by reduced factor XI activity. Bleeding is variable and is often noticed after surgery, dental procedures, childbirth, or trauma rather than occurring spontaneously.

## Factor XII

A blood-clotting protein that starts one pathway measured by laboratory clotting tests. Factor XII deficiency can greatly prolong the activated partial-thromboplastin time but usually does not cause abnormal bleeding.

## Factor XII Assay

A blood test measuring factor XII activity. It may help explain an isolated prolonged activated partial-thromboplastin time, especially when the person has no unusual bleeding.

## Factor XII Deficiency

A deficiency of factor XII that can greatly prolong one laboratory clotting test but usually does not cause bleeding. It is often discovered unexpectedly during testing for another reason.

## Factor XIII

The clot-stabilizing protein that strengthens the fibrin mesh after a clot forms. Factor XIII deficiency can cause delayed bleeding, poor wound healing, or repeated pregnancy loss despite initially normal routine clotting tests.

## Factor XIII Deficiency

A rare autosomal recessive bleeding disorder caused by a deficiency of plasma factor XIII (fibrin-stabilizing factor). Because standard coagulation tests (PT, PTT, platelet count) are completely normal, the disorder is diagnosed via abnormal clot solubility assays in 5M urea or quantitative factor XIII antigen assays; characteristically manifests with delayed umbilical stump hemorrhage, recurrent soft-tissue hematomas, and recurrent spontaneous miscarriages.

### Alternate Names:

Fibrin-Stabilizing Factor Deficiency; FXIII Deficiency

## Facultative Anaerobe

A microorganism that can grow with or without oxygen. It uses oxygen when available but can switch to other energy-producing processes when oxygen is absent; many intestinal bacteria have this ability.

## Faecal Fistula

An abnormal connection between the intestine and another organ, the skin, or the body surface that allows stool or intestinal fluid to leak. It can cause infection, dehydration, skin injury, and poor nutrition.

## Faecalibacterium

A group of oxygen-sensitive bacteria commonly found in the intestine. They can produce butyrate, a substance that supports the colon lining; their presence is studied in relation to gut health and inflammation.

## Faeces

The solid waste passed from the intestine through the anus, also called stool. Its color, consistency,

---

frequency, and contents can provide clues about digestion, bleeding, infection, inflammation, or other illness.

### **Faget Sign**

The unusual clinical physical finding of fever accompanied by a paradoxically normal or slow pulse rate (sphygmothermic dissociation / relative bradycardia), classically seen in yellow fever, typhoid fever, Legionnaires' disease, and tularemia.

#### **Alternate Names:**

Faget's Sign; Sphygmothermic Dissociation

### **Fahr Disease**

A rare disorder involving abnormal calcium deposits in parts of the brain. It may cause movement problems, seizures, cognitive changes, or psychiatric symptoms; similar deposits can also result from other metabolic or genetic conditions.

### **Fahr Syndrome**

A rare neurological disorder characterized by abnormal, symmetrical, non-atherosclerotic calcification of the basal ganglia, thalamus, and cerebral cortex, leading to progressive parkinsonism, chorea, ataxia, and neuropsychiatric disturbances.

#### **Alternate Names:**

Idiopathic Basal Ganglia Calcification; IBGC

### **Fahr's Disease**

Alternate spelling; see *Fahr Disease*.

### **Fahr's Syndrome**

A rare disorder involving calcium deposits in deep brain regions, sometimes causing movement problems, seizures, thinking changes, or psychiatric symptoms. Doctors look for inherited causes and treatable problems with calcium or parathyroid hormones.

### **Failed Induction Of Labor**

Failure of labor to progress after medicines or procedures intended to start it. The decision to continue induction or perform cesarean birth depends on cervical change, contractions, time, the mother's

condition, and fetal well-being.

### **Failed Intubation**

Inability to place a breathing tube safely in the windpipe. It is an airway emergency requiring prompt help, continued oxygen delivery, an alternative airway method, and a plan to avoid repeated traumatic attempts.

### **Failure To Thrive**

Older term for faltering weight, a pattern of slower-than-expected weight gain in a child. See *Faltering Weight*.

### **Fainting**

A brief loss of consciousness and muscle tone caused by temporarily reduced blood flow to the brain, followed by spontaneous recovery. Reflex episodes are common, but fainting during exercise, with chest pain, or without warning needs urgent assessment.

#### **Alternate Names:**

Syncope; Vasovagal Episode; Passing Out

### **Falciform Ligament**

A fold of tissue attaching the liver to the front abdominal wall and diaphragm. It marks the division between the liver's right and left lobes and contains the remnant of a fetal umbilical vein.

### **Falciparum Malaria**

Malaria caused by *Plasmodium falciparum* parasites transmitted by infected mosquitoes. It can worsen rapidly and cause brain involvement, severe anemia, kidney injury, breathing failure, shock, or death, so urgent treatment is required.

### **Fall**

An unplanned movement to the ground or another lower level. Falls may result from poor balance, weakness, vision problems, medicines, dizziness, low blood pressure, or environmental hazards and can cause serious injury.

### **Fall Prevention**

Steps to reduce falling, including strength and bal-

---

ance training, medication review, vision and hearing checks, safe footwear, adequate lighting, removal of hazards, and support with walking when needed.

### **Fall Risk**

The likelihood of falling based on factors such as previous falls, balance, walking ability, strength, vision, medicines, dizziness, blood-pressure changes, cognition, and environmental hazards.

### **Fall Screening**

A brief assessment of previous falls, balance, walking, strength, medicines, vision, dizziness, cognition, and surroundings to identify fall risk and guide prevention or further evaluation.

### **Fallen Arches**

Flattening of the foot's inner arch, also called flat feet. It may be flexible or rigid and may cause no symptoms or lead to foot, ankle, knee, or back discomfort; treatment depends on symptoms and cause.

#### **Alternate Names:**

Pes Planus; Flat Feet; Fallen Foot Arches

### **Fallen Bladder**

A common term for bladder prolapse, in which weakened pelvic-support tissues allow the bladder to bulge into the front wall of the vagina. It may cause pressure, leakage, incomplete emptying, or no symptoms.

### **Fallopian Tube**

One of two tubes between the ovaries and uterus. An egg usually travels through a tube, where fertilization may occur, before reaching the uterus; blockage or scarring can affect fertility.

### **Fallopian Tube Cancer**

A rare cancer beginning in a fallopian tube. It may cause pelvic pain, a mass, abnormal or watery discharge, or no early symptoms; diagnosis, staging, and treatment resemble those for ovarian cancer.

### **Fallopian Tube Disorders**

Disorders affecting the tubes that transport the ovum, including infection, inflammation, blockage, ectopic pregnancy, hydrosalpinx, torsion, endometriosis, and cancer. They may cause pelvic pain, infertility, abnormal bleeding, or acute abdominal symptoms.

### **Fallopian Tube Torsion**

Twisting of a fallopian tube, sometimes around a cyst or fluid-filled tube, that can cut off blood flow. Sudden severe pelvic pain, nausea, or vomiting requires urgent surgical assessment.

### **Fallopian Tubes**

The two tubes connecting the ovaries and uterus. They transport eggs and are common sites of fertilization; blockage, inflammation, scarring, pregnancy, or tumors can affect them.

### **Fallopian-Tube Endometriosis**

Endometriosis involving or surrounding a fallopian tube. It may cause pelvic pain, adhesions, or infertility, or be found unexpectedly; treatment depends on symptoms, disease extent, and fertility goals.

### **Fallot's Tetralogy**

A birth defect combining four heart abnormalities that reduce blood flow to the lungs and may cause bluish skin, breathing difficulty, poor growth, or fainting. Specialist care and usually surgery are required.

### **Falls**

Unplanned events in which a person comes to rest on the ground or another lower surface. Falls may cause fractures or head injury and often involve balance, weakness, vision, medicines, dizziness, or environmental hazards.

### **False Labor**

Contractions that may be uncomfortable but do not progressively open and thin the cervix. They are often irregular and may lessen with rest, hydration, or a change in activity, unlike established labor.

---

### False Negative

A test result that says a condition is absent when it is actually present. It can occur because testing was too early, the sample was inadequate, the condition level was low, or the test was not sensitive enough.

### False Passages

Abnormal channels made through tissue during instrumentation, surgery, or injury instead of following the normal body pathway. They can cause bleeding, perforation, infection, or failure to place a tube correctly.

### False Positive

A test result that indicates a condition is present when it is not. It may result from cross-reaction, contamination, technical error, or testing in a group where the condition is uncommon.

### False Ribs

The lower ribs that do not attach directly to the breastbone. Their cartilage joins the cartilage above or ends freely, allowing flexibility but leaving them vulnerable to injury in some areas.

### False-Negative Result

A negative result when the tested condition is actually present. Early testing, an inadequate sample, prior treatment, low levels of the target, technical error, or limited test sensitivity can cause it.

### False-Positive Result

A positive result when the tested condition is actually absent. Cross-reaction, contamination, technical error, limited specificity, or testing in a low-risk population can produce it.

### Faltering Weight

Slower-than-expected weight gain or a concerning downward change in a child's growth pattern. The 2026 AAP and NASPGHAN guideline uses this term in place of failure to thrive; feeding difficulty, inadequate intake, poor absorption, chronic disease, or other factors may contribute and require assessment.

### Alternate Names:

Failure To Thrive; Growth Faltering

### Falx Cerebelli

A small, triangular crescent-shaped fold of dura mater situated in the posterior cranial fossa inferior to the tentorium cerebelli, projecting vertically forward between the right and left cerebellar hemispheres.

### Falx Cerebri

A large, sickle-shaped vertical fold of dura mater that dips vertically into the longitudinal cerebral fissure between the right and left cerebral hemispheres, attaching anteriorly to the crista galli and frontal crest and posteriorly to the tentorium cerebelli and internal occipital protuberance.

### Famciclovir

An antiviral medicine that changes into penciclovir in the body and limits multiplication of herpes viruses. It can treat selected shingles, cold-sore, or genital-herpes episodes but does not remove the virus permanently.

### Familial

Occurring more often within a family because of shared genes, environment, behaviors, or a combination. Familial does not necessarily mean that one specific gene causes the condition.

### Familial Adenomatous Polyposis

An autosomal dominant inherited condition, usually caused by a disease-causing change in the *APC* gene, causing numerous precancerous polyps in the colon and rectum. Without preventive treatment, colorectal cancer is very likely, so lifelong specialist screening and risk-reducing surgery are often needed.

### Familial Alobar Holoprosencephaly

An inherited or familial form of alobar holoprosencephaly, a severe developmental defect in which the embryonic forebrain fails to divide normally. It may be associated with major midline facial abnormalities and requires genetic and specialist evaluation.

---

### **Familial Benign Pemphigus (Hailey-Hailey Disease)**

An inherited disorder caused by impaired calcium handling between epidermal cells, producing recurrent fragile blisters, erosions, and painful fissures in skin folds. Heat, sweating, friction, and infection can trigger flares; it is not contagious.

### **Familial Chordoma**

A familial predisposition to chordoma, a rare malignant tumor arising from remnants of the embryonic notochord. A family history may warrant genetic counseling and evaluation for tumors involving the skull base or spine.

### **Familial Combined Hyperlipidemia**

An inherited tendency to have high LDL cholesterol, high triglycerides, or both, often affecting several relatives. It increases early cardiovascular risk and is managed with lifestyle measures and lipid-lowering treatment when indicated.

### **Familial Congestive Cardiomyopathy**

Alternate name; see *Dilated Cardiomyopathy*.

### **Familial Cutaneous Amyloidosis**

An inherited tendency to develop amyloid deposits limited mainly to the skin, often producing itchy lichen or rippled macular lesions. The specific pattern and associated systemic risk require dermatologic assessment.

### **Familial Dilated Cardiomyopathy (FDC)**

Alternate name; see *Dilated Cardiomyopathy*.

### **Familial Dysautonomia**

An inherited nerve disorder affecting sensation and automatic functions such as blood pressure, heart rate, sweating, tears, digestion, and swallowing. Symptoms may include dizziness, temperature problems, repeated lung infections, and poor growth.

### **Familial Dysbetalipoproteinemia**

An inherited disorder in which the body clears certain cholesterol- and triglyceride-rich particles

poorly. It can cause high blood fats, yellowish skin deposits, and premature artery disease, especially with other metabolic risk factors.

### **Familial Emphysema**

Emphysema caused or strongly predisposed by an inherited disorder, especially alpha-1 antitrypsin deficiency. It can occur at a younger age and may be worsened by smoking or occupational exposures.

### **Familial Hypercholesterolaemia**

An inherited disorder causing very high LDL cholesterol from childhood. Without effective treatment, cholesterol can build up in artery walls and cause early heart attack, stroke, or disease of other arteries.

### **Familial Hypercholesterolemia**

Usually an autosomal dominant inherited disorder causing persistently high LDL cholesterol because the body removes it from blood poorly. Family history, cholesterol levels, physical findings, and sometimes genetic testing support the diagnosis; rare severe forms may involve changes in both gene copies.

### **Familial Hypertriglyceridemia**

An inherited tendency toward high blood triglycerides, often worsened by diabetes, obesity, alcohol, or certain medicines. Very high levels can cause abdominal pain, eruptive skin bumps, and acute pancreatitis.

### **Familial Hypocalciuric Hypercalcemia**

An inherited condition causing lifelong, usually mild high blood calcium with low urinary calcium. People often have no symptoms, and recognizing it prevents unnecessary parathyroid surgery; specialist testing may confirm the diagnosis.

### **Familial Jaundice**

Jaundice caused by an inherited disorder of bilirubin production, conjugation, transport, or excretion. Examples include Gilbert syndrome, Crigler-Najjar syndrome, Dubin-Johnson syndrome, and Rotor syndrome.



---

### **Familial Lichen Amyloidosis (Type IX)**

A hereditary form of primary localized cutaneous amyloidosis in which amyloid deposits in the skin cause intensely itchy, thickened, rippled, or hyperpigmented lesions, usually on the shins or upper back.

### **Familial Lipoprotein Lipase Deficiency**

A rare inherited disorder in which triglycerides cannot be broken down normally. It causes extremely high triglycerides, recurrent pancreatitis, abdominal pain, and yellow-red skin bumps and requires specialist dietary and medical care.

### **Familial Mediterranean Fever**

Usually an autosomal recessive inherited inflammatory disorder, often associated with changes in the *MEFV* gene, causing repeated short attacks of fever with abdominal, chest, or joint pain. Without treatment, ongoing inflammation can damage the kidneys; preventive medicine can reduce attacks and complications.

### **Familial Multiple Lipomatosis**

An inherited tendency to develop many soft, slow-growing, usually painless fatty lumps under the skin, often on the trunk and limbs. They are generally benign, but painful, rapidly growing, or unusual lumps need assessment.

### **Familial Multiple Polyposis**

An older name for familial adenomatous polyposis, an inherited disorder causing numerous adenomatous colorectal polyps and a very high risk of colorectal cancer without appropriate surveillance and preventive treatment.

### **Familial Multiple Trichoepithelioma**

An inherited condition causing many benign tumors from hair follicles, usually small skin-colored facial bumps. They may increase over time and can affect appearance; treatment is chosen according to number, size, and location.

### **Familial Paroxysmal Peritonitis**

An inherited inflammatory disorder causing re-

peated short attacks of fever with severe abdominal, chest, or joint pain. Without treatment, ongoing inflammation can damage the kidneys; preventive treatment can reduce attacks and complications.

#### **Alternate Names:**

Familial Mediterranean Fever

### **Familial Primary Pulmonary Hypertension**

Heritable pulmonary arterial hypertension caused by pathogenic variants, most often involving the *BMPR2* pathway. Progressive narrowing of pulmonary arteries increases right-heart strain and can cause breathlessness, fainting, or right-heart failure.

### **Familial Renal Amyloidosis**

An inherited amyloidosis in which a variant protein, often transthyretin or another hereditary amyloid precursor, deposits in the kidneys and sometimes the nerves, heart, or other organs. Proteinuria and progressive kidney dysfunction may occur; genetic and specialist evaluation guides care.

### **Familial Testotoxicosis**

An inherited condition in which boys produce excess testosterone without normal brain stimulation. It causes early sexual development, rapid growth, and advanced bone age but may reduce adult height without treatment.

### **Familial Tremor**

A tremor affecting more than one family member, often reflecting inherited essential tremor or another genetic movement disorder. The pattern, age of onset, medicines, and associated neurologic findings help identify the cause.

### **Family Involvement**

Participation of family members or chosen support people in healthcare planning, treatment, rehabilitation, or daily care, with the patient's permission and attention to privacy, safety, preferences, and caregiver burden.

### **Family Planning**

Personal and healthcare decisions about whether

---

and when to have children. It includes contraception, fertility care, preconception health, pregnancy planning, and support for informed reproductive choices.

### **Family Study**

A structured review of health information in relatives to understand how a disease or trait occurs in a family. It may include family history, medical records, examination, genetic counseling, or testing.

### **Family Therapy**

Counseling involving family members or other close relationships to improve communication, manage conflict, and support recovery. It may help with mental illness, addiction, chronic disease, caregiving stress, or family adjustment.

### **Famophos**

An organophosphate insecticide that can poison the nervous system by blocking acetylcholinesterase. Exposure may cause sweating, salivation, vomiting, pinpoint pupils, muscle weakness, breathing failure, or death and requires emergency care.

### **Famotidine**

A medicine that blocks histamine signals to reduce stomach-acid production. It treats reflux, heartburn, and ulcers; dose reduction may be needed in kidney disease, and headache, diarrhea, or confusion can occur.

### **Fanconi Anemia**

An inherited disorder affecting DNA repair and bone-marrow blood-cell production. It may cause low blood counts, infections, bleeding, short stature, birth differences, organ problems, and a high risk of leukemia or other cancers.

### **Fanconi Syndrome**

A kidney-tubule disorder causing excess loss of glucose, phosphate, bicarbonate, amino acids, and other substances in urine. It can lead to dehydration, abnormal blood acidity, weak bones, poor growth, or kidney disease.

### **Fanconi-Bickel Syndrome**

A rare autosomal recessive glycogen storage disease caused by mutations in the *SLC2A2* gene encoding the GLUT2 glucose/galactose transporter, characterized by hepatomegaly, glycogen accumulation, proximal renal tubular dysfunction (renal Fanconi syndrome), and severe growth failure.

### **FAP**

An inherited condition causing many precancerous polyps in the colon and rectum. Colorectal cancer is very likely without preventive treatment, so regular specialist surveillance and risk-reducing surgery are often needed.

#### **Alternate Names:**

Familial Adenomatous Polyposis

### **Farad**

The standard unit of electrical capacitance, equal to one coulomb per volt. It is used in medical equipment and electrophysiology to describe how much electrical charge a component can store.

### **Farber Lipogranulomatosis**

A rare inherited storage disorder caused by deficient acid ceramidase. Fatty material accumulates and may cause painful skin lumps, joint stiffness, hoarseness, bone changes, breathing problems, and neurologic disease.

### **Farmer Lung**

A form of hypersensitivity pneumonitis (extrinsic allergic alveolitis) caused by repeated inhalation of thermophilic actinomycetes (such as *Saccharopolyspora rectivirgula*) spores from moldy hay or compost, presenting with acute or chronic dyspnea, dry cough, fever, and bilateral pulmonary infiltrates.

#### **Alternate Names:**

Farmer's Lung; Moldy Hay Hypersensitivity Pneumonitis

### **Farmer's Lung**

Lung inflammation caused by repeatedly inhaling moldy hay, grain, or other farm dust. It can cause fever, cough, chills, and breathlessness hours after exposure and may lead to permanent scarring if

---

exposure continues.

### **Farsightedness**

A focusing problem in which close objects look blurred because light would focus behind the retina. It may cause eyestrain or headaches and is usually corrected with glasses or contact lenses.

#### **Alternate Names:**

Hyperopia; Hypermetropia

### **Fascia**

Sheets or bands of connective tissue that surround, separate, support, and connect muscles, organs, nerves, and blood vessels. Injury or inflammation of fascia can cause pain, stiffness, or restricted movement.

### **Fascia Adherens**

A strong connection joining neighboring cells, especially heart-muscle cells. It links the internal contractile framework of one cell to the membrane of another so force can pass through the tissue.

### **Fascia Lata**

The deep, dense fibrous connective tissue investing sheath of the thigh, which encloses thigh musculature and thickens laterally to form the iliotibial tract, continuous superiorly with the inguinal ligament and fascia of the gluteus maximus.

### **Fascial Loop**

A loop or sling made from a person's tissue or synthetic material to support a body structure. For example, a sling may support the urethra in selected treatment of stress urinary incontinence.

### **Fasciculation**

A brief, involuntary twitch visible under the skin from contraction of a small group of muscle fibers. It may be harmless, but widespread or persistent twitching with weakness needs neurologic evaluation.

### **Fasciculus Cuneatus**

The lateral ascending white matter sensory tract

of the posterior (dorsal) column of the spinal cord, present above thoracic level T6, carrying conscious proprioception, vibratory sense, and fine discriminative touch from the upper trunk, neck, and upper extremities to the nucleus cuneatus of the medulla.

### **Fasciculus Gracilis**

The medial ascending white matter sensory tract of the posterior (dorsal) column of the spinal cord, present throughout the entire length of the spinal cord, conveying conscious proprioception, vibration, and fine discriminative touch from the lower extremities and lower trunk to the nucleus gracilis of the medulla.

### **Fasciitis**

Inflammation of the connective tissue around muscles and other structures. It may be localized, as in plantar fasciitis, or rapidly destructive, as in necrotizing infection requiring emergency surgery and antibiotics.

### **Fasciola**

A genus of parasitic flatworms, or liver flukes, that infect some animals and people. Infection is acquired by swallowing immature parasites on contaminated water plants or water and may affect the liver and bile ducts.

### **Fasciola Hepatica**

A liver fluke that infects people after contaminated aquatic plants or water are swallowed. Immature worms migrate through the liver, while adult worms live in bile ducts and can cause pain, fever, or obstruction.

### **Fascioliasis**

Infection with liver flukes after eating contaminated aquatic plants or drinking contaminated water. It may cause fever, right-upper-abdominal pain, enlarged liver, or high eosinophils and later cause bile-duct pain, infection, or blockage.

### **Fasciolopsis**

A genus of large intestinal flukes that can infect people who eat contaminated aquatic plants.

---

Heavy infection may cause abdominal pain, diarrhea, swelling, poor nutrition, or intestinal obstruction.

### **Fasciolopsis Buski**

A large intestinal fluke acquired by eating aquatic plants carrying immature parasites. Heavy infection may cause abdominal pain, diarrhea, intestinal inflammation, malnutrition, or swelling.

### **Fasciotomy**

Emergency surgery that cuts tight fascia to release dangerous pressure in a muscle compartment. It may prevent permanent nerve and muscle damage after crush injury, fracture, burns, or loss and return of blood flow.

### **Fast Heart Rate**

A heart rate faster than expected for a person's age and activity. Exercise, fever, pain, stress, dehydration, anemia, medicines, or an abnormal rhythm may cause it; chest pain or fainting is urgent.

### **Fast-Twitch Fiber**

A muscle fiber capable of quick, powerful contraction but more rapid fatigue than a slow-twitch fiber. It supports sprinting, jumping, lifting, and other brief high-intensity activity.

#### **Alternate Names:**

Type II Muscle Fiber; Fast-Glycolytic Fiber

### **Fasting**

Going without food, and sometimes without drinks, for a defined period. It may be required before blood tests, anesthesia, surgery, or certain procedures, or undertaken for personal or religious reasons.

### **Fasting Blood Glucose**

Blood sugar measured after no caloric intake, usually for at least eight hours. It helps screen for and diagnose diabetes or impaired glucose regulation, but results must be interpreted with the clinical context.

### **Fasting Lipid Profile**

A blood test measuring cholesterol and triglycerides

after a period without food. It helps assess cardiovascular risk and guide treatment; fasting is particularly useful when triglycerides need accurate measurement.

#### **Alternate Names:**

Lipid Panel; Blood Lipid Screen

### **Fasting Plasma Glucose Test**

A blood test measuring blood glucose after fasting, usually at least eight hours. It helps identify diabetes or prediabetes and is interpreted with repeat testing or other results when needed.

### **Fat**

Energy-rich substances that provide fuel, insulation, and components for cell membranes and hormones. Dietary fats also help absorb vitamins A, D, E, and K; types include saturated, unsaturated, and trans fats.

#### **Alternate Names:**

Lipid; Adipose Substance

### **Fat Embolism**

Entry of fat droplets into the bloodstream, usually after major trauma, a thigh or pelvic fracture, or orthopedic surgery. It may cause breathing difficulty, confusion, rash, or low oxygen and can become life-threatening.

### **Fat Embolism Syndrome**

A severe, life-threatening systemic complication occurring 24 to 72 hours following long bone fractures (such as femur or tibia) or pelvic trauma, characterized by the clinical triad of acute respiratory distress with hypoxemia, neurological dysfunction (confusion, encephalopathy), and a petechial skin rash distributed across the anterior chest, neck, and axillae.

#### **Alternate Names:**

FES; Traumatic Fat Embolism

### **Fat Measurement**

Estimation of body-fat amount or distribution using methods such as skinfolds, bioimpedance, DEXA, imaging, or laboratory analysis. Results vary by method and should be interpreted with overall

---

health, strength, and metabolic findings.

### **Fat Necrosis**

Death of fat cells, often after pancreatitis, injury, or breast surgery. It can cause pain, inflammation, or a firm lump that resembles cancer on examination or imaging and may require testing to distinguish the cause.

### **Fat Pad**

A localized cushion of fatty tissue that protects, supports, insulates, or permits movement around an organ or joint. Inflammation or injury of a fat pad can cause localized pain and swelling.

### **Fat-Soluble Toxins**

Poisonous chemicals that dissolve in body fat and may remain for a long time. Some pesticides, industrial chemicals, polychlorinated biphenyls, and dioxins can accumulate after repeated exposure and harm several organs.

### **Fat-Soluble Vitamin**

A vitamin that dissolves in fat and can be stored in body tissues. Vitamins A, D, E, and K are fat-soluble; deficiency or excessive supplementation can cause illness.

### **Fat-Soluble Vitamins**

Vitamins A, D, E, and K, which dissolve in fat and can accumulate in the body. They support vision, bones, immunity, cell protection, and blood clotting; excessive supplements can be toxic.

### **Fatal Catatonia**

A severe, life-threatening form of catatonia with extreme motor disturbance, fever, unstable blood pressure or heart rate, dehydration, and risk of organ failure. It requires emergency treatment.

### **Fatal Familial Insomnia**

A very rare inherited prion disease, usually caused by a disease-causing variant in one copy of the *PRNP* gene. Misfolded prion protein damages the thalamus and other brain areas involved in sleep and

automatic body functions. It causes progressively severe loss of normal sleep, along with problems such as sweating, abnormal heart rate or blood pressure, temperature changes, weight loss, impaired movement, and changes in thinking. The disease usually begins in adulthood, progresses over time, and is fatal.

### **Fatigue**

Persistent tiredness or reduced energy that limits physical or mental activity and may not improve fully with rest. Sleep problems, anemia, infection, hormone or organ disease, medicines, mood disorders, and neurologic conditions can cause it.

### **Fatty Acid**

An organic molecule with a hydrocarbon chain and a terminal carboxyl group. Fatty acids may circulate freely or be joined to glycerol and other molecules in fats and cell membranes; they can provide energy and serve as starting materials for signaling chemicals. They may be saturated, monounsaturated, or polyunsaturated.

### **Fatty Acid Oxidation**

The metabolic breakdown of fatty acids to acetyl-CoA and other products that can be used for energy production. Much of this process occurs through mitochondrial beta-oxidation, although other cellular pathways also contribute.

#### **Alternate Names:**

Beta-Oxidation

### **Fatty Acid Oxidation Disorder**

An inherited disorder in which cells cannot use certain fats normally for energy. Fasting or illness may cause low blood sugar, vomiting, muscle breakdown, weakness, heart problems, or sudden metabolic illness.

### **Fatty Acid Synthase**

A group of linked enzymes that helps cells make fatty acids, especially palmitate. These fatty acids are used for energy storage, cell membranes, and other cellular functions.

---

## Fatty Acid Synthesis

The metabolic production of fatty acids from smaller carbon-containing molecules, mainly through acetyl-CoA and malonyl-CoA. It supports the formation of energy stores, cell membranes, and signaling molecules.

## Fatty Acids

A family of organic molecules that are components of fats and cell membranes and can also circulate freely in the body. They provide or store energy and serve as precursors for signaling molecules; their biological and dietary effects depend on their structure, type, and amount.

## Fatty Degeneration

Abnormal buildup of fat inside cells, especially liver cells, caused by metabolic problems, alcohol, toxins, poor nutrition, obesity, or other injury. The change may reverse or progress to inflammation and scarring.

## Fatty Liver

Alternate name; see *Fatty Liver Disease*.

## Fatty Liver Disease

Excess fat stored in liver cells. It may be related to metabolic dysfunction, obesity, diabetes, abnormal cholesterol, alcohol use, medicines, or other causes. Metabolic dysfunction-associated steatotic liver disease (MASLD) is the current term for fatty liver associated with cardiometabolic risk factors; inflammation and cell injury are called metabolic dysfunction-associated steatohepatitis (MASH). Disease may remain mild or progress to fibrosis, cirrhosis, liver failure, or liver cancer.

### Alternate Names:

Hepatic Steatosis; Steatotic Liver Disease; MASLD

## Fatty Streak

An early artery-wall change in which cholesterol-rich immune cells collect beneath the lining. It can be an early stage of atherosclerosis, although some fatty streaks never progress to a harmful plaque.

## Favism

Rapid destruction of red blood cells after eating fava beans, usually in a person with glucose-6-phosphate dehydrogenase deficiency. It may cause jaundice, dark urine, weakness, breathlessness, or severe anemia and needs medical care.

## Favre-Racouchot Syndrome

A benign chronic-sun-damage disorder in which solar elastosis is accompanied by open or closed comedones and small epidermal cysts, usually around the temples, outer eyes, cheeks, or lateral neck. It is associated with photoaging and sometimes smoking.

## Favus

A long-lasting fungal infection of the scalp, usually causing yellow cup-shaped crusts around hairs. Without treatment it can destroy follicles and cause permanent scarring hair loss.

## Fc Receptor

A protein on an immune cell that binds the tail end of an antibody. This helps the cell recognize, ingest, or destroy the antibody's target and can also influence inflammation.

## Fear

An emotional and body response to a perceived threat that prepares a person to protect themselves. Persistent or excessive fear can cause avoidance, panic, sleep problems, and impaired daily functioning.

## Fears

Feelings of threat or apprehension. Fears are normal when proportionate to danger, but persistent, excessive, or disabling fears may indicate an anxiety disorder and can often be treated.

## Febrile

Relating to fever, a temporary rise in body temperature usually caused by infection, inflammation, or another illness. Fever differs from heat-related hyperthermia; severe illness, dehydration, confusion, or fever with very low white cells is urgent.



---

## Febrile Convulsion

A seizure during fever in a young child without evidence of brain infection or another immediate cause. Most are brief and generalized, but a prolonged, repeated, or one-sided seizure requires emergency assessment.

## Febrile Convulsions

Seizures triggered by fever in young children, usually between six months and five years, without evidence of infection or another direct brain problem. Most resolve quickly, but first or prolonged seizures require medical assessment.

## Febrile Neutropenia

Fever in a person with a dangerously low neutrophil count, the white cells that fight infection. Infection may be present without obvious symptoms, so urgent examination, cultures, and prompt antibiotics are often needed.

## Febrile Seizure

A seizure during fever in a child, usually between six months and five years, without evidence of brain infection or a previous seizure without fever. Most are brief, but prolonged, repeated, or one-sided events require urgent assessment.

### Alternate Names:

Fever-Induced Seizure; Febrile Convulsion

## Febrile Seizure Cross-Reference

Alternate name; see *Febrile seizure*.

## Febrile Seizures

Seizures associated with fever in young children without evidence of brain infection or a previous unprovoked seizure. Most do not cause lasting harm, but the first seizure or any prolonged event needs medical evaluation.

### Alternate Names:

Febrile Convulsions; Pyrexial Seizures

## Febuxostat

A medicine that lowers uric-acid production for long-term prevention of gout attacks. It does not relieve

sudden gout pain immediately and may require monitoring for liver problems and cardiovascular risk.

## FEC

Abbreviation; see *FECD*.

## Fec Protocol

A chemotherapy regimen combining fluorouracil, epirubicin, and cyclophosphamide for selected cancers. It can cause low blood counts, infection, nausea, hair loss, mouth sores, infertility, and heart or bladder-related toxicity.

## Fecal Calprotectin

A stool test measuring a protein released during intestinal inflammation. A raised result can support evaluation for inflammatory bowel disease, but infection, medicines, age, and other conditions can also raise it.

## Fecal Culture

Laboratory testing that grows bacteria from a stool sample to investigate infectious diarrhea. It may identify a treatable cause and guide antibiotics, but some infections require toxin, molecular, or parasite testing instead.

## Fecal Elastase

A stool test estimating whether the pancreas releases enough digestive enzymes. A low result may suggest pancreatic insufficiency, causing poor digestion, greasy stools, weight loss, or vitamin deficiency.

## Fecal Fat

Measurement of excess fat in stool. High fecal fat suggests poor digestion or absorption from pancreatic disease, reduced bile, intestinal disease, or another condition and may explain greasy, difficult-to-flush stools.

## Fecal Immunochemical Test

A stool screening test that detects human blood, mainly from the lower intestine, using antibodies. A positive result does not diagnose cancer but usually

---

requires colon evaluation, often colonoscopy.

### **Fecal Impaction**

A large, hard mass of stool trapped in the rectum or colon. It may cause pain, abdominal swelling, nausea, urinary symptoms, bowel blockage, or watery leakage around the mass and may require professional removal.

### **Fecal Incontinence**

Inability to control passage of stool or gas, ranging from soiling to complete loss of bowel control. Causes include diarrhea, constipation, sphincter injury, nerve disease, rectal prolapse, and cognitive impairment.

#### **Alternate Names:**

Accidental Bowel Leakage; Encopresis (Pediatric)

### **Fecal Lactoferrin**

An iron-binding glycoprotein released by degranulating neutrophils in the intestinal mucosa; its elevated concentration in stool serves as a sensitive, non-invasive biomarker to detect active mucosal inflammation in inflammatory bowel disease (Crohn disease, ulcerative colitis) and differentiate it from irritable bowel syndrome.

### **Fecal Microbiota Transplant**

Transfer of carefully screened and prepared donor stool or its microbial contents into the intestine to restore helpful microbes, mainly for selected recurrent *C. difficile* infection. Infection transmission remains a possible risk.

### **Fecal Microbiota Transplantation**

Transfer of screened donor stool or microbial material into a patient's intestine to restore microbial balance. It is used mainly for selected recurrent *C. difficile* infections after standard therapy and requires regulated donor screening.

### **Fecal Occult Blood Test**

A stool test that detects blood too small to see. It may screen for colorectal cancer or investigate digestive-tract bleeding, but it does not identify the cause; a positive result usually requires further

colon evaluation.

### **Fecal Smear**

Examination of a thin stool sample with a microscope or stain. Depending on the test, it may help identify parasites, inflammatory cells, blood, fat, or other abnormalities.

### **Fecal Transplant**

A procedure placing carefully screened donor stool into the intestine to restore helpful gut microbes. It is mainly used for selected recurrent *C. difficile* infection and has small but important infection and procedure-related risks.

### **Fecalith**

A hard mass of dried stool, often in the appendix or colon. It may contribute to appendicitis, blockage, inflammation, or perforation.

### **Fecaluria**

Stool, gas, or a fecal smell in urine, usually indicating an abnormal connection between the intestine and urinary tract. It often accompanies recurrent urinary infection and requires medical evaluation.

### **FECD**

An abbreviation for Fuchs endothelial corneal dystrophy, a progressive loss of corneal cells that normally remove fluid. It can cause morning blur, glare, painful swelling, and reduced vision; advanced disease may need a corneal transplant.

#### **Alternate Names:**

Fuchs Dystrophy; Fuchs Endothelial Corneal Dystrophy

### **Feces**

Solid or semisolid waste passed from the bowel, containing water, undigested material, bacteria, shed cells, and metabolic waste. Changes in stool can provide clues about digestion, bleeding, infection, or inflammation.

### **Fecolith**

A hard mass of dried stool that can block the ap-

---

pendix or bowel and contribute to inflammation, appendicitis, or perforation.

### **Fecundity**

The biological ability to conceive and produce a live birth. It depends on age, ovulation, ovarian reserve, fallopian tubes, uterus, sperm health, and other factors; it differs from the chance of pregnancy in one cycle.

### **Feedback**

Information about a system's output that changes later activity. Negative feedback stabilizes functions such as temperature, glucose, and hormones, while positive feedback strengthens processes such as labor contractions.

### **Feedback Inhibition**

A control process in which the end product of a chemical pathway slows an earlier step. This prevents unnecessary buildup of the product and helps cells use nutrients and energy efficiently.

### **Feedback Mechanism**

A regulatory process in which a system's result influences what happens next. Negative feedback maintains stability, while positive feedback amplifies a response until a defined endpoint is reached.

### **Feeding Jejunostomy**

Placement of a feeding tube into the jejunum, part of the small intestine, when eating or stomach feeding is unsafe or impossible. It provides nutrition but can cause infection, blockage, leakage, or bowel injury.

### **Feeding Methods**

Ways of providing food or liquid, including breast or bottle feeding, oral feeding, tube feeding, and intravenous nutrition. The safest method depends on swallowing ability, age, illness, and nutritional needs.

### **Feeding Poor**

Taking too little food or fluid for age or health needs. In an infant it may signal illness, swallow-

ing difficulty, breathing problems, pain, reflux, or developmental difficulty and requires assessment if persistent.

### **Feeding Tube**

A flexible tube used to deliver nutrition, fluids, or medicines to the stomach or intestine when eating is inadequate or unsafe. Risks include blockage, leakage, infection, aspiration, and injury.

### **Feingold Diet**

A restrictive diet that avoids certain food additives and some naturally occurring chemicals, promoted for behavior or attention problems. Evidence of benefit is limited, and restriction can cause nutritional problems without professional guidance.

### **Felbamate**

An antiseizure medicine reserved for selected severe or drug-resistant epilepsy. It can rarely cause life-threatening bone-marrow failure or liver failure, so specialist prescribing and regular blood and liver tests are essential.

### **Felodipine**

A calcium-channel blocker that relaxes artery walls and lowers blood pressure. It may cause ankle swelling, flushing, headache, fast heartbeat, or gum enlargement and should be taken as prescribed.

### **Felon**

A painful bacterial infection in the fleshy fingertip, often after a small puncture. Pus and pressure can damage tissue, bone, or tendons, so prompt assessment and sometimes drainage and antibiotics are needed.

### **Felty Syndrome**

A rare complication of long-standing rheumatoid arthritis involving a very low neutrophil count and often an enlarged spleen. It increases infection risk and requires specialist treatment of arthritis and neutropenia.

---

## Felty's Syndrome

A rare complication of long-standing rheumatoid arthritis causing neutropenia and often an enlarged spleen. It raises the risk of serious infection and may require disease-modifying treatment, growth factors, or other specialist care.

## Female

Describing the sex category generally associated with ovaries, eggs, and female reproductive anatomy. Biological traits vary among individuals, and sex, gender identity, and gender expression are distinct concepts.

## Female Athlete Triad

A pattern of low energy availability, menstrual disturbance, and reduced bone strength in physically active females. Inadequate nutrition or excessive exercise can increase the risk of stress fractures and other health problems.

## Female Breast

A mammary gland and its skin, fat, connective tissue, ducts, and lobules on the front of the chest. Its structure changes with age, hormones, pregnancy, breastfeeding, and body composition.

## Female Circumcision

Removal or injury of part of the external female genitalia for nonmedical reasons. It has no health benefit and can cause severe pain, bleeding, infection, urinary and sexual problems, childbirth complications, and lasting psychological harm.

## Female Condom

A soft internal sheath placed in the vagina before sex that forms a barrier against semen and can reduce pregnancy and transmission of some sexually transmitted infections when used correctly.

## Female Condoms

Flexible internal barriers placed in the vagina before sex to help prevent pregnancy and reduce exposure to semen and some sexually transmitted infections. They should not be used at the same time as an external condom.

## Female Genital Mutilation

Cutting or removing external female genital tissue for nonmedical reasons. It has no health benefit and can cause pain, bleeding, infection, urinary and sexual problems, childbirth complications, and long-term psychological harm.

## Female Genital Sores

Painful or painless ulcers, blisters, or eroded lesions located on the vulva, vagina, cervix, or perineum. Common etiologies include sexually transmitted infections (herpes simplex virus, syphilis, chancroid, lymphogranuloma venereum), non-infectious inflammatory disorders (Behçet disease, lichen planus), contact dermatitis, trauma, or malignancy.

### Alternate Names:

Female Genital Ulcers; Vulvar Lesions; Vulvar Ulcers

### Alternate Names:

Female Genital Ulcers; Vulvar Ulcerations

## Female Genitalia

The external and internal reproductive organs, including the vulva, vagina, cervix, uterus, fallopian tubes, and ovaries. These structures support urination, sexual function, menstruation, fertility, pregnancy, and birth.

## Female Gonad

An ovary, the reproductive gland that stores and releases eggs and produces hormones including estrogen and progesterone.

## Female Infertility

Difficulty achieving pregnancy because of problems with ovulation, ovarian reserve, fallopian tubes, uterus, hormones, or other factors. Evaluation considers both partners and may be recommended after a defined period of trying or sooner with risk factors.

### Alternate Names:

Female Sterility; Subfertility in Women

## Female Orgasmic Disorder

Persistent or recurrent difficulty achieving orgasm that causes personal distress. It may be lifelong

---

or acquired and can relate to health conditions, medicines, mood, trauma, sexual pain, stimulation, or relationship factors.

**Alternate Names:**

Anorgasmia; Inhibited Female Orgasm

**Female Pattern Baldness**

Gradual scalp hair thinning, usually widest near the part or crown, influenced by genes and hormones. Iron deficiency, thyroid disease, medicines, and other causes should be considered when loss is new or rapid.

**Female Pelvis**

The pelvic bones and soft tissues in a person with typical female anatomy. It is often broader with a wider outlet than typical male anatomy, but substantial individual variation exists.

**Female Phenotype**

The observable anatomical, physiologic, and developmental traits associated with female sex. These traits vary with genes, hormones, development, age, and individual biology.

**Female Reproductive Organs**

Female reproductive organs include the ovaries, fallopian tubes, uterus, cervix, vagina, and external genitalia, which support hormones, ovulation, menstruation, pregnancy, and birth.

**Female Reproductive System**

The ovaries, fallopian tubes, uterus, cervix, vagina, vulva, and related hormones. It supports egg release, menstruation, pregnancy, birth, and sexual function and can be affected by developmental, hormonal, infectious, structural, or cancerous disorders.

**Female Self-Catheterization**

Intermittent insertion and removal of a catheter through the female urethra to empty the bladder when normal urination is inadequate, reducing retention and protecting kidney function.

**Female Sexual Dysfunction**

A broad term for persistent or recurrent difficulty with sexual desire, arousal, lubrication, orgasm, or pain that causes clinically significant distress. Medical conditions, medicines, mood, trauma, relationships, and life changes may contribute.

**Alternate Names:**

FSD; Hypoactive Sexual Desire Disorder

**Female Sexual Interest/Arousal Disorder**

Persistent or recurrent low sexual interest or arousal causing personal distress. Medical conditions, medicines, mood, trauma, pain, relationship factors, hormonal changes, and other social or psychological factors may contribute.

**Female Urethral Meatus**

The external opening of the female urethra, through which urine leaves the body; it is located anterior and superior to the vaginal opening.

**Female Urethral Opening**

The opening through which urine leaves the female urethra, located in the vulvar vestibule in front of the vaginal opening. Its position and appearance may vary normally.

**Feminization**

Development of physical or hormonal traits commonly associated with female puberty, such as breast growth or reduced body hair. It may be normal, part of gender-affirming treatment, or caused by hormones, medicines, or disease.

**Femoral**

Relating to the femur or thigh, including the thigh bone, artery, vein, nerve, or neck of the femur.

**Femoral Anteversion**

An inward twist of the thigh bone that turns the knees or feet inward, often noticed in childhood. It commonly improves with growth; persistent pain, severe asymmetry, or walking problems need orthopedic assessment.

---

## **Femoral Artery**

A major artery carrying blood through the groin and thigh to the leg. Its pulse can be felt in the groin, and it is used for some catheter procedures; injury or blockage can threaten the limb.

## **Femoral Canal**

The medial, smallest compartment of the femoral sheath, measuring approximately 1.3 cm in length, containing loose connective tissue, lymphatic vessels, and the deep inguinal lymph node of Cloquet; site of femoral herniation below the inguinal ligament.

## **Femoral Hernia**

A bulge of abdominal tissue through an opening below the groin ligament and beside the femoral vessels. It has a relatively high risk of trapping or losing blood supply to the contents and often needs surgical assessment.

## **Femoral Hernia Repair**

Surgery that returns trapped tissue to the abdomen and closes or reinforces the opening in the groin. Repair is often recommended because femoral hernias can become trapped or strangulated.

## **Femoral Nerve**

A large nerve from the lower back that supplies muscles that flex the hip and straighten the knee and provides sensation to the front thigh and inner leg. Injury can cause weakness, numbness, or falls.

## **Femoral Nerve Dysfunction**

Reduced function of the nerve supplying the front thigh and knee-straightening muscles. It can cause hip or knee weakness and altered sensation and may result from diabetes, compression, injury, surgery, or pelvic disease.

## **Femoral Nerve Stretch Test**

A neurological examination manoeuvre used to assess for upper lumbar nerve root irritation (L2, L3, or L4) or femoral neuropathy. With the patient prone, the examiner passively flexes the knee and extends the hip; reproduction of pain radiating down the anterior thigh indicates a positive test.

## **Alternate Names:**

Prone Knee Bending Test; Reverse Straight Leg Raise

## **Femoral Neuropathy**

Damage to the femoral nerve causing weakness in hip flexion or knee extension and numbness or pain in the front thigh or inner leg. Causes include diabetes, injury, bleeding, compression, and pelvic disease.

## **Femoral Pulse**

The pulse felt over the femoral artery in the groin. Comparing it with pulses elsewhere helps assess blood flow to the leg and can provide clues about arterial blockage or shock.

## **Femoral Shaft Fractures**

Breaks in the middle portion of the thigh bone, often from high-energy trauma but sometimes from weakened bone. They can cause major blood loss, deformity, and inability to walk and usually require urgent stabilization and specialist treatment.

## **Femoral Sheath**

A funnel-shaped fascial sleeve formed by downward prolongation of the transversalis fascia (anteriorly) and iliac fascia (posteriorly) beneath the inguinal ligament into the femoral triangle, compartmentalized into lateral (femoral artery), intermediate (femoral vein), and medial (femoral canal) spaces.

## **Femoral Triangle**

A triangular area at the upper front of the thigh bordered by the groin ligament and nearby muscles. It contains the femoral nerve, artery, vein, and lymph nodes and is an important examination and access site.

## **Femoral Vein**

The major deep vein of the thigh, carrying blood from the leg toward the pelvis and heart. A clot in it can cause deep-vein thrombosis and pulmonary embolism.



---

### **Femtogram**

A unit of mass equal to one quadrillionth of a gram, or  $10^{-15}$  gram. It is used in laboratory measurements of extremely small quantities.

### **Femtoliter**

A unit of volume equal to one quadrillionth of a liter, or  $10^{-15}$  liter. It is used to describe very small laboratory or cellular volumes.

### **Femur**

The thigh bone and the longest, strongest bone in the body. It joins the hip to the knee and includes a head, neck, shaft, and lower end; fractures can threaten mobility and blood supply.

### **Femur Fracture**

A break in the thigh bone caused by trauma or weakened bone. It may cause severe pain, deformity, blood loss, and inability to bear weight; treatment depends on location, alignment, and overall health.

### **Femur Length**

The ultrasound-measured length of a fetus's thigh bone. It helps estimate pregnancy age and assess growth, but interpretation depends on other measurements, family size, pregnancy dating, and possible skeletal or chromosomal conditions.

### **Fencer's Pose**

A posture in which turning the head makes the arm and leg on that side straighten while the opposite arm and leg bend. It is a normal reflex in young infants but may indicate a brain or nerve problem if it persists or appears unexpectedly.

### **Fenfluramine**

Fenfluramine is a serotonergic medicine used in some jurisdictions for selected severe epileptic encephalopathies; historical appetite-suppressant use was limited by valvular heart disease and pulmonary hypertension, so cardiac monitoring is required.

### **Fenitrothion**

A poisonous organophosphate insecticide. Exposure

can cause sweating, excess saliva, vomiting, diarrhea, small pupils, muscle twitching, breathing difficulty, seizures, or weakness because it disrupts nerve signals; suspected poisoning needs urgent medical help.

### **Fenobucarb**

A carbamate insecticide that can cause nerve poisoning after significant absorption, producing sweating, vomiting, small pupils, muscle weakness, breathing difficulty, or seizures.

### **Fenofibrate**

A medicine that lowers blood triglycerides and can modestly change cholesterol levels by increasing the liver's processing of fats. It is used for selected lipid disorders; liver, kidney, muscle, and gallbladder risks require monitoring.

### **Fenoldopam**

A short-acting dopamine D1-receptor agonist used intravenously for severe hypertension in selected settings; it can cause hypotension, flushing, headache, and increased intraocular pressure.

### **Fenoprofen**

Fenoprofen is a nonsteroidal anti-inflammatory drug used for pain and inflammatory arthritis; gastrointestinal bleeding, renal injury, fluid retention, and cardiovascular events are important risks.

### **Fenoprofen Calcium Overdose**

Taking too much fenoprofen, an anti-inflammatory pain medicine, can cause nausea, vomiting, stomach bleeding, drowsiness, kidney injury, breathing or acid-base problems, seizures, or coma. Suspected overdose requires immediate poison-control or emergency advice; do not induce vomiting.

### **Fenoterol**

A short-acting beta-2 bronchodilator that relaxes airway muscle and relieves bronchospasm. It may cause tremor, nervousness, low potassium, or a fast or irregular heartbeat and is not approved everywhere.

---

## **Fenoxycarb**

An insect-growth regulator that imitates juvenile hormone and disrupts development of immature insects. It is used for pest control, and exposure should be minimized because products can irritate skin, eyes, or airways.

## **Fentanyl Citrate**

The citrate salt of fentanyl, a potent opioid analgesic used for severe pain and anesthesia; respiratory depression, sedation, dependence, and overdose are major risks.

## **Fentanyl Derivatives**

Strong synthetic opioid medicines related to fentanyl, including sufentanil, remifentanil, and alfentanil. They may relieve severe pain or support anesthesia but can rapidly slow breathing, cause overdose, and produce dependence, so close monitoring is essential.

## **Fenthion**

A poisonous organophosphate insecticide. Exposure can cause sweating, excess saliva, vomiting, diarrhea, small pupils, muscle twitching, breathing difficulty, seizures, or weakness because it disrupts nerve signals; suspected poisoning needs urgent medical help.

## **Fermentation**

A metabolic process in which microorganisms or cells convert organic substrates without using oxidative respiration, producing products such as acids, alcohols, or gases.

## **Ferriman-Gallwey Score**

A standardized clinical scoring system used to quantify the severity of hirsutism in women by visually grading terminal hair density from zero (absence) to four (extensive) across nine androgen-sensitive body areas, with a total score of eight or higher generally indicating clinical hirsutism.

### **Alternate Names:**

Modified Ferriman-Gallwey Scale; mFG Score

## **Ferritin**

A protein that stores iron inside cells and releases it when needed. Blood ferritin often reflects body iron stores, but inflammation, infection, liver disease, or cancer can raise it even when iron is low.

## **Ferritin Blood Test**

A blood test measuring stored iron indirectly. Low ferritin strongly supports iron deficiency; high ferritin may reflect iron excess, inflammation, infection, liver disease, or cancer and must be interpreted with other tests.

## **Ferroptosis**

A controlled form of cell death caused by iron-dependent damage to cell fats and failure of antioxidant protection. It differs from apoptosis and ordinary tissue necrosis and is being studied in cancer and other diseases.

## **Ferrous Sulfate**

An oral iron supplement used to prevent or treat iron-deficiency anemia. It may cause nausea, constipation, abdominal discomfort, or dark stools; accidental overdose can be life-threatening, especially in children.

## **Fertile Period**

The several days in a menstrual cycle when sex can result in pregnancy, including the days before and shortly after ovulation. Timing varies, and calendar estimates are not reliable contraception for everyone.

## **Fertilisation**

Union of a sperm and an egg to form a zygote, the first cell of an embryo. It usually occurs in a fallopian tube before the developing embryo moves toward the uterus.

## **Fertility**

The biological ability to conceive and produce offspring. It depends on ovulation, sperm production, reproductive anatomy, hormones, age, general health, and other factors affecting either partner.

---

## Fertility Domain

The part of health concerned with reproductive function, ability to conceive, fertility goals, infertility, preservation of fertility, and related medical, emotional, and social care.

## Fertility Drugs

Medicines used to stimulate ovulation, support reproductive treatment, or improve the chance of pregnancy. They can cause side effects and multiple pregnancy, so selection and monitoring require fertility-specialist care.

## Fertility Preservation

Steps taken to protect the possibility of future pregnancy before treatment or disease may damage reproductive function. Options include freezing eggs, embryos, sperm, or selected reproductive tissue, depending on age, timing, and goals.

## Fertility Treatment

Medical or procedural care intended to help achieve pregnancy. Options include treating underlying disease, ovulation medicines, surgery, insemination, or in-vitro fertilization, chosen according to cause, age, health, and reproductive goals.

## Fertilization

Fusion of sperm and egg genetic material to form a zygote, usually in a fallopian tube. This begins embryonic development, which may continue after the zygote reaches the uterus and implants.

## Ferumoxylol

An intravenous iron preparation used for selected adults with iron-deficiency anemia. It can cause allergic reactions, low blood pressure, or iron overload, so monitoring is required during and after infusion.

## FESS

Functional endoscopic sinus surgery uses instruments passed through the nostrils to open blocked sinus drainage pathways and remove selected diseased tissue. It may improve chronic sinus disease but can cause bleeding, infection, eye injury, or fluid leakage.

## Festinating Gait

Walking with increasingly short, quick steps and a forward-leaning posture, commonly associated with Parkinson disease or other movement disorders. It increases fall risk and may improve with specialist treatment and physical therapy.

## Fetal Acid-Base Status

An assessment of a fetus's oxygen supply and blood acidity, often using umbilical-cord blood gases at birth. Results must be interpreted with the sampling site, timing, fetal condition, and complete clinical record.

## Fetal Alcohol Spectrum Disorder

A group of lifelong conditions that can occur after alcohol exposure before birth. Effects vary widely and may involve growth, physical development, brain development, learning, memory, attention, behavior, speech, coordination, or everyday functioning. Some people have characteristic facial features, birth defects, or problems with organs such as the heart or kidneys, but these are not present in everyone. The effects can occur even when the person does not have the facial features or growth problems associated with fetal alcohol syndrome.

### Alternate Names:

FASD; Prenatal Alcohol Exposure Syndrome

## Fetal Alcohol Spectrum Disorders

Alternate plural form; see *Fetal Alcohol Spectrum Disorder*.

## Fetal Alcohol Syndrome

The most severe recognized form of fetal alcohol spectrum disorder, involving characteristic facial features, poor growth, and brain-development problems after prenatal alcohol exposure. No known safe amount of alcohol exists during pregnancy.

### Alternate Names:

FAS; Fetal Alcohol Spectrum Condition

## Fetal Anatomy Survey

A comprehensive second-trimester obstetric ultrasound examination typically performed between 18 and 22 weeks of gestation to systematically evaluate

---

fetal organ anatomy, biometry, placental position, amniotic fluid volume, and detect congenital structural anomalies.

**Alternate Names:**

Level II Ultrasound; Fetal Anomaly Scan; Second-Trimester Ultrasound; Fetal Anatomy Survey

**Fetal Attitude**

The position of a fetus's head, arms, legs, and trunk in relation to one another. Normal flexion helps birth; unusual extension or flexion can change the presenting part and affect labor.

**Fetal Bradyarrhythmia**

An unusually slow or irregular fetal heart rhythm. Causes include conduction block, extra beats, maternal or fetal disease, or medicines; persistent abnormalities require prompt specialist assessment.

**Fetal Bradycardia**

A fetal heart-rate baseline below 110 beats per minute for at least ten minutes. Medicines, maternal illness, fetal heart disease, low oxygen, cord problems, or placental separation may cause it and persistent cases require urgent obstetric assessment.

**Fetal Circulation**

Blood flow before birth in which the placenta supplies oxygen and nutrients and special connections direct much blood around the lungs, which are not yet used for breathing. These connections normally close after birth.

**Fetal Death**

Death of a fetus before delivery. Medical and legal definitions vary by gestational age and location; care includes confirming the diagnosis, supporting the family, and discussing delivery and follow-up.

**Fetal Development**

Growth and maturation from the end of the embryonic period until birth. Organs enlarge and mature, movements develop, and body systems gradually acquire the functions needed after delivery.

**Fetal Disease**

A disease, structural abnormality, or developmental problem affecting a fetus before birth. Causes include genetic changes, infection, medicines, toxins, and poor placental function; ultrasound and prenatal tests may identify it.

**Fetal Distress**

A nonspecific term for concerning fetal findings during labor, especially abnormal heart-rate patterns that may indicate low oxygen, cord compression, infection, or placental problems. Clinicians assess the whole pattern and may act urgently.

**Fetal Echocardiography**

A specialized ultrasound examination of the fetal heart used to assess cardiac structure, rhythm, and function, including possible congenital heart disease.

**Fetal Fibronectin**

An extracellular matrix glycoprotein present at the decidual-chorionic interface; detection in cervicovaginal secretions between 22 and 34 weeks of gestation serves as a valuable negative predictive biomarker for imminent preterm labor and delivery within 7 to 14 days.

**Alternate Names:**

fFN

**Fetal Growth Restriction**

A fetus that is smaller than expected because it may not be reaching its growth potential, often because the placenta is not supplying enough oxygen or nutrients. Repeat ultrasound, fluid, blood-flow, and well-being checks guide care.

**Fetal Heart**

The developing heart of a fetus, whose rate and rhythm can be assessed by ultrasound or electronic monitoring to evaluate fetal well-being.

**Fetal Heart Monitoring**

Continuous electronic or intermittent auscultatory surveillance of the fetal heart rate and its dynamic response to uterine contractions during labor and

---

delivery (cardiotocography), evaluated for baseline rate, variability, accelerations, and decelerations (early, late, variable).

**Alternate Names:**

Cardiotocography; CTG; Electronic Fetal Monitoring

**Fetal Heart Rate**

The number of times a fetus's heart beats each minute, usually about 110–160 during pregnancy and labor. It can be checked with a Doppler or electronic monitor; changes may reflect normal activity, medicines, or reduced oxygen.

**Fetal Hemoglobin**

The main oxygen-carrying protein in a fetus's red blood cells. It binds oxygen more strongly than adult hemoglobin, helping the fetus obtain oxygen from the placenta; its amount normally falls after birth.

**Fetal Hydantoin Syndrome**

A distinct pattern of congenital malformations resulting from maternal gestational exposure to hydantoin anticonvulsants (predominantly phenytoin). Hallmarks include craniofacial dysmorphism (cleft lip/palate, broad depressed nasal bridge, hypertelorism), hypoplasia of distal phalanges and nails, intrauterine growth restriction, microcephaly, and neurodevelopmental delay.

**Alternate Names:**

Dilantin Embryopathy; Phenytoin Syndrome

**Fetal Hydrops**

Severe fluid buildup in at least two parts of a fetus, such as under the skin, around the lungs or heart, or in the abdomen. Causes include anemia, heart disease, infection, chromosome conditions, and blood-group incompatibility and require specialist care.

**Fetal Hypoxia**

Too little oxygen reaching a fetus before or during birth. Placental disease, maternal breathing or circulation problems, cord compression, or prolonged labor can contribute; abnormal heart-rate patterns

require prompt obstetric assessment.

**Fetal Kick Counts**

Observation and recording of fetal movements by a pregnant person. A noticeable reduction from the usual pattern should prompt timely obstetric advice.

**Alternate Names:**

Kick Counting; Fetal Activity Assessment

**Fetal Lie**

The direction of a fetus's long body axis compared with the pregnant person's spine. The fetus may lie lengthwise, sideways, or diagonally; a sideways or diagonal position near birth can affect delivery planning.

**Fetal Lung Maturity**

The developmental state of the fetal pulmonary system characterized by sufficient production of pulmonary surfactant by type II pneumocytes to prevent neonatal respiratory distress syndrome; assessed in amniotic fluid via the lecithin-to-sphingomyelin (L/S) ratio or lamellar body count.

**Alternate Names:**

FLM

**Fetal Macrosomia**

A fetus that is larger than expected, often estimated at 4,000–4,500 grams or more at birth depending on the clinical definition. Diabetes and previous large babies increase risk; complications include difficult birth, injury, and cesarean delivery.

**Fetal Malpresentation**

A fetal presentation other than the usual vertex presentation, including breech, transverse, or oblique lie, which may affect labor and delivery planning.

**Fetal Monitor**

A device that records fetal heart rate and, when configured for electronic fetal monitoring, uterine contractions. An external Doppler ultrasound sensor detects the fetal heartbeat, while a tocodynamometer detects uterine contractions. In selected labor

---

settings, a fetal scalp electrode can provide an internal fetal heart-rate signal.

### **Fetal Monitoring**

Assessment of fetal heart rate and, sometimes, uterine contractions to evaluate fetal wellbeing during pregnancy or labor.

### **Fetal Presentation**

The part of the baby positioned to enter the pregnant person's pelvis first. The usual presentation is head first; breech, shoulder, face, brow, and mixed presentations can affect labor and delivery planning.

### **Fetal Pyelectasis**

Mild widening of the urine-collecting area of a fetus's kidney seen on an ultrasound scan. It may go away before birth or be linked to a urinary blockage or another condition, so follow-up depends on its size and other findings.

### **Fetal Scalp Electrode**

A small electrode placed on the fetus's scalp through the cervix to record the heart rate directly during labor. It may give a clearer signal when external monitoring is inadequate but can cause infection, bleeding, or scalp injury.

### **Fetal Scalp PH Testing**

A test that analyzes a small blood sample from the baby's scalp during labor to assess whether reduced oxygen may be causing acid buildup. It may clarify an abnormal fetal heart-rate pattern, but use is limited and depends on local expertise and clinical circumstances.

### **Fetal Scalp Stimulation**

A test during labor in which the baby's scalp is gently touched during a vaginal examination. A brief rise in the baby's heart rate may be reassuring, but the result must be interpreted with the full heart-rate recording and clinical situation.

### **Fetal Stage**

The fetal stage begins after the embryonic period

and continues until birth, characterized mainly by growth and maturation of established organs.

### **Fetal Tachycardia**

A sustained fetal heart-rate baseline above 160 beats per minute for at least 10 minutes. Causes may include maternal fever, infection, medicines, fetal hypoxia, or fetal arrhythmia.

### **Fetal Tissue**

Fetal tissue is tissue from a developing fetus and may be relevant to pregnancy care, pathology, research, transplantation, or genetic testing under applicable ethical and legal rules.

### **Fetal Ultrasound**

Imaging during pregnancy using sound waves to assess the fetus, placenta, fluid, growth, movement, and anatomy. It can estimate gestational age, identify some structural problems, assess growth or blood flow, and guide selected procedures.

### **Fetal Viability**

The possibility that a fetus could survive outside the uterus with intensive medical care. It depends on pregnancy age, fetal condition, treatment available, and other factors and is not defined by one fixed week for every pregnancy.

### **Fetal Weight**

An estimate of a fetus's mass, usually calculated from ultrasound measurements. It helps assess growth and identify possible growth restriction or unusually large size, but it is not exact.

### **Fetal-Maternal Erythrocyte Distribution Blood Test**

A test that estimates the amount of fetal red blood cells in maternal circulation, helping guide Rh immune globulin dosing after fetomaternal hemorrhage.

### **Fetishism**

A recurrent sexual interest in a nonliving object or a specific non-genital body part. The interest alone



---

is not a mental disorder.

### **Fetishistic Disorder**

A mental-health disorder involving repeated, intense sexual arousal from a nonliving object or a specific non-genital body part, with clinically significant distress or impairment or involvement of harm or lack of consent.

### **Fetomaternal Hemorrhage**

Passage of fetal blood into the pregnant person's bloodstream through the placenta. Small amounts can occur normally, but heavy bleeding across the placenta after trauma, placental separation, or a procedure can cause fetal anemia and affect preventive treatment.

### **Fetomaternal Transfusion**

Movement of fetal blood cells across the placenta into the pregnant person's bloodstream. Small amounts are common, but heavy transfer after trauma, placental problems, or procedures can cause fetal anemia and may require urgent testing and treatment.

### **Fetor Hepaticus**

A musty or sweet breath odor that can occur with severe liver dysfunction when certain substances are not adequately cleared. It is a warning sign requiring evaluation for advanced liver disease and complications.

### **Fetoscope**

An instrument used to hear or assess the fetal heart-beat through the pregnant person's abdominal wall, especially later in pregnancy.

### **Fetus**

A developing human from the start of the ninth week after fertilization until birth. During this period, established organs grow and mature; medical descriptions may instead use pregnancy weeks counted from the last menstrual period.

### **FEV1/FVC Ratio**

The amount of air forcefully exhaled in the first second divided by the total amount exhaled. A lower-than-expected ratio suggests narrowed airways, but interpretation uses age, sex, height, reference values, and bronchodilator response.

### **Fever**

A regulated rise in body temperature caused by changes in the body's temperature-control system, commonly in response to infection or inflammation. It may also occur with certain medicines, cancers, autoimmune disorders, or other conditions. Fever differs from hyperthermia, in which body temperature rises without a normal change in the temperature set point.

### **Fever And Headache**

Fever with headache may occur with viral illness, influenza, sinus disease, dehydration, meningitis, encephalitis, or other infection. Severe sudden headache, neck stiffness, confusion, rash, seizure, weakness, or reduced consciousness requires emergency assessment.

### **Fever Blister**

Alternate name; see *Cold Sore*.

### **Fever Of Unknown Origin**

Persistent or recurrent fever whose cause remains unexplained after an appropriate initial evaluation. Causes may include infection, inflammation, cancer, or medicines and require systematic reassessment.

### **Fever-Induced Seizure**

Alternate name; see *Febrile seizures*.

### **Fexofenadine**

An antihistamine used to relieve sneezing, itching, runny nose, and watery eyes from allergies and itching from hives. It usually causes less drowsiness than older antihistamines, but headache or nausea can occur.

---

## Fiber

Dietary fiber is plant material that the small intestine cannot digest. It adds bulk to stool, supports regular bowel movements, and some types help lower blood cholesterol and slow rises in blood glucose.

## Fiber Optic

A technology that sends light through thin glass or plastic fibers. Flexible fiber-optic instruments are used in endoscopy to view body passages and guide some minimally invasive procedures.

## Fibrates

A group of medicines that mainly lower high triglyceride levels by changing how the liver and tissues handle fats. They can cause indigestion, gallstones, liver problems, or muscle injury, especially with some other lipid medicines.

## Fibre

A threadlike structure in the body or another material. In nutrition, fibre is plant material that is not digested in the small intestine and helps regulate bowel movements.

## Fibrillary Glomerulonephritis

A rare non-amyloid glomerular disease characterized by glomerular deposition of randomly arranged, non-branching fibrils (16 to 24 nm in diameter) that do not stain with Congo red, strongly positive for DNAJB9 by immunohistochemistry, presenting clinically with proteinuria, hematuria, and renal insufficiency.

### Alternate Names:

FGN; DNAJB9 Glomerulopathy

## Fibrillation

Rapid, disorganized activity of muscle fibers. In the heart, atrial fibrillation causes an irregular rhythm and can lead to blood clots; ventricular fibrillation stops effective pumping.

## Fibrin

An insoluble, threadlike protein that forms a mesh at an injured blood vessel. The mesh traps blood cells

and strengthens the clot that helps stop bleeding.

## Fibrin Degradation Products Blood Test

A blood test that measures fragments released when blood clots are broken down. High levels may occur with widespread clotting, severe infection, liver disease, recent surgery, or pregnancy, so results require clinical interpretation.

## Fibrin Sealant

A surgical adhesive containing clotting proteins that form a fibrin clot when applied to tissue. It can help control bleeding or seal selected tissues, but it does not replace repair of major blood-vessel injuries.

## Fibrin Split Products

Soluble protein fragments released into the bloodstream during plasmin-mediated enzymatic cleavage and degradation of cross-linked fibrin polymers and fibrinogen, elevated in disseminated intravascular coagulation, deep vein thrombosis, and acute pulmonary embolism.

### Alternate Names:

FDPs; Fibrin Degradation Products

## Fibrinogen

A blood protein made mainly by the liver that is changed into fibrin during clot formation. Low levels can cause bleeding; levels may rise with inflammation and fall in severe liver disease or widespread clotting.

## Fibrinogen Assay

A laboratory test that measures the amount or clotting activity of fibrinogen in blood. It helps investigate unexplained bleeding, abnormal clotting, liver disease, or suspected widespread clotting disorders.

## Fibrinogen Blood Test

A blood test measuring fibrinogen, a liver-made protein needed to form blood clots. Low levels may contribute to bleeding, while high levels commonly accompany inflammation, infection, pregnancy, or cardiovascular risk.

---

### **Fibrinogen Decreased**

A lower-than-normal blood fibrinogen level, which can weaken clot formation and cause bleeding. Causes include inherited deficiency, severe liver disease, major blood loss or transfusion, and widespread activation of clotting.

### **Fibrinoid Leukodystrophy**

An older name associated with Alexander disease, a rare astrocyte disorder in which Rosenthal fibers accumulate and white matter is damaged. Clinical features vary by age and may include developmental regression, seizures, movement problems, or bulbar dysfunction.

### **Fibrinoid Necrosis**

Severe damage to a blood-vessel wall in which blood proteins and immune material collect within the wall. It can occur in some inflammatory vessel diseases and is identified by examining affected tissue.

### **Fibrinolysis**

The natural process that dissolves a blood clot after healing begins. An enzyme called plasmin breaks down fibrin; excessive breakdown can cause bleeding, while insufficient breakdown can allow harmful clots to persist.

### **Fibrinolytic Agent**

A medicine that dissolves a blood clot by helping the body convert plasminogen into plasmin, which breaks down fibrin. It may treat selected emergencies but can cause serious bleeding, including bleeding in the brain.

### **Fibrinolytic Therapy**

Treatment that uses clot-dissolving medicine to restore blood flow in selected emergencies, such as certain strokes, heart attacks, or pulmonary embolisms. The main risk is serious bleeding, so eligibility is assessed urgently.

#### **Alternate Names:**

Thrombolytic Therapy; Clot-Dissolving Therapy

### **Fibrinopeptide A Blood Test**

A blood test that measures a protein fragment released when the body begins converting fibrinogen into fibrin. It can indicate clotting activity but is not specific enough to diagnose a condition by itself.

### **Fibrinous Pericarditis**

Inflammation of the sac surrounding the heart, with a sticky protein called fibrin coating its surfaces. It may cause sharp chest pain, a scratchy heart sound, and fluid around the heart.

### **Fibroadenoma**

A common noncancerous breast lump made of glandular and connective tissue. It is usually firm, smooth, movable, and painless, especially in adolescents and young adults; imaging helps distinguish it from other breast masses.

### **Fibroadenoma Of The Breast**

A noncancerous breast lump containing glandular and connective tissue. It is commonly smooth, firm, movable, and painless, but any new or changing breast lump should be examined and appropriately imaged.

### **Fibroblast**

A cell that makes collagen and other supporting materials in connective tissue. Fibroblasts help maintain skin, tendons, and organs and become active during wound healing and scar formation.

### **Fibrocartilage**

A tough type of cartilage that combines strength with limited flexibility. It helps absorb pressure and resist pulling in the discs between spinal bones, the pubic joint, and the knee menisci.

### **Fibrocystic Breast Changes**

Alternate name; see *Fibrocystic Breast Changes*.

### **Fibrocystic Breast Condition**

Alternate name; see *Fibrocystic Breast Changes*.

---

## Fibrocystic Breasts

Common noncancerous changes in breast tissue that may cause lumpiness, thickening, cysts, swelling, tenderness, or discomfort. Symptoms may change during the menstrual cycle and usually improve after menopause. Fibrocystic breast changes do not increase the risk of breast cancer.

### Alternate Names:

Fibrocystic Breast Disease (deprecated historical term)

## Fibrodysplasia Ossificans Progressiva

A rare inherited disorder in which muscles, tendons, ligaments, and other connective tissues gradually turn into bone. This extra bone forms outside the normal skeleton and can make joints stiff or fixed, restricting movement. Episodes of soft-tissue swelling and inflammation may be followed by permanent bone formation, and injury or invasive procedures can trigger these episodes. The disorder usually becomes noticeable in early childhood and may later limit opening of the mouth or expansion of the rib cage. Most cases are caused by a change in the *ACVR1* gene. A characteristic feature present from birth is a malformed, shortened, or inward-turned big toe.

### Alternate Names:

FOP; Stone Man Syndrome

## Fibroepithelial Polyp Of The Vulva

A usually noncancerous, soft tissue growth on the vulva or nearby vaginal tissue. It may enlarge during pregnancy or the reproductive years and can cause irritation, bleeding, or discomfort if large.

## Fibrofolliculoma

A small, usually noncancerous skin growth arising around a hair follicle, often on the face, neck, or upper trunk. Multiple lesions may occur with an inherited disorder that also raises kidney-tumor risk.

## Fibroid

A noncancerous growth of smooth muscle, most often in the uterus. It may cause heavy menstrual bleeding, pelvic pressure, pain, frequent urination, fertility problems, or no symptoms.

### Alternate Names:

Uterine Leiomyoma; Myoma

## Fibroid Tumor

A noncancerous growth of the uterus made from smooth muscle and connective tissue. It may cause heavy or painful periods, pelvic pressure, frequent urination, infertility, or no symptoms.

## Fibroids

Noncancerous growths in the muscle of the uterus. They may cause heavy or painful menstrual bleeding, pelvic pressure, frequent urination, constipation, fertility problems, or no symptoms.

## Fibrolamellar Carcinoma

A rare primary liver cancer that usually affects adolescents and young adults without cirrhosis. It may cause abdominal pain, a mass, weight loss, or abnormal liver tests; imaging and biopsy establish diagnosis and resection is considered when feasible.

## Fibrolamellar Hepatocellular Carcinoma

A distinct primary malignant hepatic neoplasm occurring predominantly in adolescents and young adults without pre-existing cirrhosis or viral hepatitis, histologically characterized by large polygonal eosinophilic hepatocytes embedded within dense, lamellar collagenous stroma; associated with a *DNAJB1-PRKACA* fusion gene.

## Fibrolipoma

A usually noncancerous, slow-growing lump made of mature fat and fibrous tissue. It is often painless and lies beneath the skin, but a new or enlarging lump should be assessed.

## Fibroma

A usually benign tumor made mainly of fibrous connective tissue. Fibromas can develop in skin, soft tissue, bone, or organs; their effects depend on their location and size, and some site-specific forms may recur or interfere with organ function.

---

### **Fibroma Of Tendon Sheath**

A usually noncancerous, slow-growing lump attached to the covering of a tendon, most often in a finger, hand, or wrist. It is firm and may recur after removal.

### **Fibromas**

Plural of fibroma; see *Fibroma*.

### **Fibromatoses**

A group of connective-tissue growth disorders that do not spread to distant organs but may grow into nearby structures and return after treatment. Effects depend on the affected body site.

### **Fibromatosis**

An overgrowth of connective-tissue cells that does not spread to distant organs but can invade nearby tissue and recur. Desmoid-type disease may cause pain, a firm mass, or impaired organ function.

### **Fibromuscular Dysplasia**

An abnormal structure of medium-sized artery walls, most often in the kidney or neck arteries. It can cause high blood pressure, headache, or stroke, or produce an artery bulge or tear.

### **Fibromyalgia**

A long-term condition that causes widespread body pain and increased sensitivity to pain. It commonly causes fatigue, unrefreshing sleep, stiffness, and difficulty concentrating or remembering. The exact cause is not fully understood, but altered pain processing in the nervous system may contribute. Diagnosis is based mainly on the pattern of symptoms and examination; no single laboratory or imaging test confirms it.

### **Fibromyalgia Syndrome**

A long-term pain condition causing widespread body pain, fatigue, poor-quality sleep, and difficulty thinking or concentrating. Doctors diagnose it from the symptom pattern and examination, while treatment combines activity, sleep support, and selected medicines.

### **Alternate Names:**

FMS

### **Fibromyxosarcoma**

A rare cancer of soft tissue containing fibrous and jelly-like material. It may present as a growing lump or pain; diagnosis requires tissue examination, and treatment depends on its site and spread.

### **Fibronectin**

A protein found in blood and connective tissue that helps cells attach, move, and organize during tissue repair. It supports wound healing and the structure of tissues around cells.

### **Fibroproliferation**

Excess growth of connective-tissue cells with increased production of scar tissue. It can thicken or stiffen organs, including the lungs, and may develop after injury, inflammation, or surgery.

### **Fibrosarcoma**

A rare cancer arising from cells that make fibrous connective tissue. It usually develops in deep soft tissue or bone and may cause a growing mass, pain, or reduced movement.

### **FibroScan**

A painless ultrasound-based test that estimates liver stiffness. Greater stiffness can suggest scarring, but inflammation, congestion, obesity, or fluid in the abdomen can affect accuracy and results may need other tests.

### **Fibrosing Alopecia**

Hair loss caused by inflammation and permanent scarring of hair follicles. The affected skin may look smooth or shiny, and regrowth is unlikely once follicles are destroyed; early treatment may limit progression.

### **Fibrosing Cholangitis**

Chronic inflammation and scarring of the bile ducts, most commonly referring to primary sclerosing cholangitis. It can cause cholestasis, itching, jaun-

---

dice, recurrent bacterial cholangitis, cirrhosis, and increased risk of bile-duct cancer.

### **Fibrosis**

Excess formation of scar tissue after ongoing injury or inflammation. Fibrosis can make an organ stiff and reduce its function, such as causing breathlessness in the lungs or impaired filtration in the kidneys.

### **Fibrous Cavernositis**

Inflammation and fibrosis of the erectile tissue of the penis, which may cause pain, swelling, erectile dysfunction, or curvature. Infection, trauma, injections, and other inflammatory conditions can contribute.

### **Fibrous Dysplasia**

A noncancerous bone disorder in which normal bone is replaced by weaker abnormal tissue. It may affect one or several bones and cause pain, deformity, fractures, or pressure on nearby nerves.

### **Fibrous Dysplasia Pathology**

A fibro-osseous lesion containing irregular strands of woven bone in a cellular fibrous stroma, classically without prominent osteoblastic rimming. The lesion may be monostotic or polyostotic.

### **Fibrous Interstitial Pneumonia**

A pattern of lung disease in which inflammation and scar tissue thicken the interstitium and impair oxygen transfer. Causes include idiopathic pulmonary fibrosis, autoimmune disease, medication toxicity, and environmental exposure.

### **Fibrous Joint**

A joint in which bones are held together by strong connective tissue and have little or no movement. Skull sutures and syndesmoses are examples.

### **Fibrous Plaque**

A thickened area in an artery wall containing scar-like tissue, cholesterol, and other material. It can narrow blood flow or rupture, allowing a blood clot to form and block the artery.

### **Alternate Names:**

Atherosclerotic Plaque

### **Fibrous Plaques of the Penis**

Fibrous scar-like plaques within the penile tunica albuginea, characteristic of Peyronie disease. They may cause penile curvature, pain during erection, shortening, or difficulty with sexual activity.

### **Fibrous Sclerosis of the Penis**

Another descriptive name for Peyronie disease, a disorder in which fibrous plaques form in the erectile covering and may bend or shorten the penis.

### **Fibula**

The slender bone on the outer side of the lower leg. It provides muscle attachment and forms the outer ankle. The nerve near its upper end can be injured by pressure or fracture.

### **Fibular Artery**

Also called the peroneal artery, it is a branch of the leg's main artery that supplies the outer and back lower leg and contributes blood flow around the ankle.

### **Fick Principle**

A method for estimating blood flow from how much oxygen the body uses and the difference in oxygen content between blood entering and leaving the lungs. It can estimate cardiac output during specialized heart testing.

### **Alternate Names:**

Fick Method; Fick Cardiac Output Determination

### **Fiducial Markers**

Small visible markers placed on or inside the body to identify a precise location during imaging, radiation treatment, or surgery. They help clinicians align equipment and target the intended tissue accurately.

### **Field Fever**

An older name for leptospirosis, a bacterial infection acquired through water or soil contaminated with urine from infected animals. It may cause fever,



---

headache, muscle pain, jaundice, kidney injury, or meningitis.

### Field Of View

The area included in a medical image or examination. A wider field shows more of the body, while a narrower field provides greater detail about a smaller area.

### Fifth Disease

A usually mild infection caused by parvovirus B19. Children may develop fever, cold-like symptoms, bright-red cheeks, and a lacy rash. It can cause serious fetal anemia during pregnancy and prolonged anemia in some vulnerable people.

#### Alternate Names:

Erythema Infectiosum; Slapped Cheek Syndrome; Parvovirus B19 Infection

### FIGO Leiomyoma Classification

A system that describes where a uterine fibroid lies in relation to the uterine cavity and outer surface. The location category helps clinicians predict symptoms and choose treatment.

### Filamentous Fungus

A fungus that grows as branching threads called hyphae, which form a network called mycelium. Some filamentous fungi cause skin, nail, lung, or invasive infections, especially in people with weakened immunity.

### Filarial Elephantiasis

Long-lasting swelling and thickening of skin and underlying tissues caused by lymphatic blockage from filarial worms. It commonly affects the legs or genitals and may require antiparasitic treatment, skin care, and swelling management.

### Filarial Lymphedema

Chronic progressive swelling, dermal fibrosis, and non-pitting lymphedema of the limbs, genitalia, or breasts resulting from adult filarial worm infection (*Wuchereria bancrofti*, *Brugia malayi*) causing obstructive damage to the lymphatic drainage pathways, progressing to irreversible elephantiasis.

#### Alternate Names:

Lymphatic Filariasis; Elephantiasis Tropica

### Filariasis

Infection with threadlike parasitic worms spread by mosquitoes, flies, or other insects. Depending on the species, worms may affect lymph vessels, skin, or eyes, causing swelling, inflammation, itching, or vision problems.

### Filler

A material injected or placed in tissue to restore volume, smooth wrinkles, change contours, or replace lost structure. Risks include pain, infection, lumps, and accidental blockage of a blood vessel.

### Filler Excipient

An inactive ingredient added to a medicine to provide bulk, improve stability, or help form a tablet, capsule, or liquid. It may rarely cause allergy or intolerance.

### Filopodium

A thin, finger-like projection from a cell surface. It senses nearby surroundings and helps cells attach, move, communicate, or form connections, including connections made by developing nerve cells.

### Filoviridae

A family of viruses that includes Ebola and Marburg viruses. They can cause severe fever, bleeding, organ failure, and death, and are mainly spread through direct contact with blood or other infected body fluids.

### Filoviruses

Viruses such as Ebola and Marburg that can cause severe hemorrhagic fever. Symptoms may include fever, headache, muscle pain, vomiting, diarrhea, bleeding, and organ failure; spread occurs through infected body fluids.

### Filtered Back Projection

A computer method for creating an image from scan measurements, especially in computed tomography.

---

It is fast, but images may contain more noise or artifacts than those made with newer reconstruction methods.

### **Filtering Procedure**

Eye surgery that creates a new drainage route for fluid to lower pressure inside the eye. It may treat glaucoma when medicines or other procedures do not control pressure adequately.

### **Filtration**

Separation of particles or dissolved substances from a fluid by passing it through a barrier. In the kidneys, tiny filters remove water and small waste products from blood to begin urine formation.

### **Filtration Fraction**

The percentage of plasma flowing through the kidneys that is filtered to begin urine formation. It helps describe kidney blood flow and filtering function and is calculated from kidney filtration compared with plasma flow.

### **Filum Terminale**

A thin strand of connective tissue that anchors the lower end of the spinal cord to the tailbone. Abnormal attachment can restrict cord movement and cause symptoms of tethered spinal cord.

### **Fimbria**

A fringe-like structure. In the fallopian tube, the fimbria are finger-like projections near the ovary that help guide a released egg into the tube; the term is also used for bacterial attachment structures.

### **Fimbriae**

Finger-like projections at the end of each fallopian tube that help collect an egg released from the ovary and guide it into the tube. Their movement and tiny surface hairs help direct the egg toward the uterus.

### **Fimbriae Proteins**

Bacterial surface proteins that form hair-like projections used to attach bacteria to human cells, medical

devices, or other surfaces. Attachment can help bacteria form communities and begin infection.

### **Fine Crackles**

Brief, high-pitched, popping sounds heard mainly during breathing in. They may occur when small airways or air sacs open in conditions such as lung scarring, fluid congestion, or other lung disease.

### **Fine Motor Control**

The coordinated use of small muscles for accurate movements, especially of the hands and fingers. It supports tasks such as writing, buttoning clothes, using utensils, and handling small objects.

### **Fine Motor Skills**

Abilities that use small muscles for controlled movements, especially in the hands and fingers. Children develop skills such as grasping, feeding themselves, drawing, dressing, and writing at different individual rates.

### **Fine Needle Aspiration**

A procedure using a thin needle to remove cells or fluid from a lump or organ for laboratory examination. It usually causes little tissue injury and may be guided by touch or ultrasound.

### **Fine Needle Aspiration Of The Thyroid**

A procedure using a thin needle, often guided by ultrasound, to collect thyroid cells from a nodule. Laboratory examination helps determine whether the nodule is benign, malignant, or needs further testing.

### **Fine-Needle Aspiration**

A biopsy in which a thin needle removes cells or fluid from a lump or cyst for laboratory examination. It may be guided by touch or imaging and is usually less invasive than a surgical biopsy.

#### **Alternate Names:**

FNA; Fine-Needle Aspiration Cytology; FNAC

### **Finegoldia**

A group of bacteria that grow without oxygen and

---

normally live on skin or mucous membranes. They can occasionally cause wound, soft-tissue, bone, joint, or bloodstream infections.

### **Finegoldia Magna**

A bacterium normally found on skin and mucous membranes that grows without oxygen. It can cause infections after wounds or surgery, including abscesses and deeper soft-tissue infections.

### **Finger**

One of the hand's digits, made of bones, joints, muscles, tendons, nerves, blood vessels, and skin. Fingers support grasping, touch, balance, and precise movements.

### **Finger Dislocation**

An injury in which the bones of a finger joint are forced out of their normal alignment. Pain, swelling, deformity, and limited movement are common; fractures, tendon damage, and impaired circulation must also be checked.

### **Finger Flexors**

Muscles and tendons that bend the fingers at their joints. Injury can cause pain or inability to bend a finger and may require prompt assessment, especially after a cut or forceful injury.

### **Finger Injury**

Damage to a finger's bone, joint, tendon, ligament, nerve, blood vessel, nail, or skin, usually from trauma. Assessment checks alignment, movement, sensation, blood flow, and possible hidden fractures.

### **Finger Joint**

A connection between the bones of a finger that allows bending, straightening, and limited side-to-side movement. Finger joints can be affected by injury, arthritis, inflammation, infection, or dislocation.

### **Finger Pain**

Discomfort in one or more fingers caused by injury, arthritis, tendon problems, nerve compression, infection, inflammation, or reduced blood flow. Sudden

severe pain, deformity, numbness, or discoloration needs prompt care.

### **Finger Stick**

A small fingertip puncture used to collect a drop of capillary blood. The sample can be tested immediately, commonly for blood glucose or other point-of-care measurements.

### **Finger Sucking**

Persistent sucking of a finger or thumb beyond early childhood. Prolonged habit can alter tooth position, the palate, bite, and facial growth; dental assessment may be useful when it persists or causes injury.

### **Finger-Nose Test**

A clinical coordination test in which the patient alternately touches their own nose and the examiner's outstretched finger using the index finger. Dysmetria (overshooting or undershooting) or intention tremor during the movement indicates ipsilateral cerebellar hemisphere dysfunction.

### **Fingernail**

The hard protective plate covering the end of a finger, made mainly of keratin and growing from the nail root. Changes in color, shape, or growth can sometimes signal injury or disease.

### **Fingers That Change Color**

Fingers that become white, blue, purple, or red because of changes in blood flow, temperature, or oxygen. Recurrent attacks may be Raynaud phenomenon; persistent changes, pain, numbness, or sores need medical assessment.

### **Finkelstein Test**

A wrist examination that gently stresses the tendons on the thumb side. Pain there may support de Quervain tenosynovitis, but the result is not specific and must be interpreted with symptoms and examination.

### **Finnish Type Amyloidosis (Type V)**

A hereditary gelsolin amyloidosis caused by vari-

---

ants in the *GSN* gene. It commonly causes slowly progressive corneal lattice dystrophy, cranial-nerve weakness, and lax or aging-looking skin, with variable systemic involvement.

### **Fire Ants**

Ants whose stings inject venom and cause immediate burning pain, redness, and itchy pus-filled bumps. Multiple stings can cause severe swelling, vomiting, breathing difficulty, or a life-threatening allergic reaction.

### **Firesetting Behavior**

Deliberately starting fires. Repeated fire-setting may be linked to an impulse-control disorder, mental illness, substance use, developmental problems, or social stress and requires assessment of safety and risk to others.

### **First Adult Seizure**

The initial suspected unprovoked or provoked seizure in an adult, requiring assessment for immediate causes, recurrence risk, injury, medication or substance effects, and mimics such as syncope.

### **First Aid**

Immediate basic care for injury or sudden illness before professional treatment arrives. It includes ensuring safety, calling emergency services, checking breathing, starting resuscitation when needed, and controlling severe bleeding.

### **First Aid for Burns**

Cool a recent thermal burn under cool running water, remove nearby jewelry or clothing that is not stuck to the skin, and cover it loosely with a clean dressing. Do not apply ice or break blisters. Large, deep, chemical, electrical, facial, or airway burns need urgent medical care.

### **First Aid Kit**

A group of supplies for treating minor injuries and giving temporary care during emergencies. Useful items include gloves, dressings, bandages, antiseptic, scissors, and emergency contact information.

### **First Menstrual Period**

Menarche, the first menstrual bleeding, usually occurring during puberty. Timing varies, but absence by the expected developmental stage or bleeding with severe pain or other concerning symptoms warrants medical assessment.

### **First Stage Of Labor**

The time from regular labor contractions that cause cervical change until the cervix is fully open. It includes early and active phases, with progress varying according to the person, fetus, and clinical circumstances.

### **First Trimester**

The first twelve weeks of pregnancy, when the embryo's organs begin forming and the placenta develops. Nausea, fatigue, and breast tenderness are common; prenatal care includes dating and early health assessment.

### **First-Degree Atrioventricular Block**

A cardiac conduction delay in which every atrial impulse is conducted to the ventricles but with a uniformly prolonged PR interval exceeding 200 milliseconds (0.20 seconds) on electrocardiogram, typically benign and asymptomatic.

#### **Alternate Names:**

First-Degree AV Block

### **First-Degree AV Block**

A mild delay in the electrical signal traveling from the heart's upper chambers to its lower chambers. Every signal still reaches the lower chambers; it is often symptomless but may reflect medicines or heart disease.

### **First-Degree Burn**

A superficial burn affecting only the outer skin layer. It causes redness, pain, and dryness without blisters, as with mild sunburn, and usually heals within several days without scarring.

### **First-Degree Heart Block**

A delay in the electrical signal between the heart's

---

upper and lower chambers, shown on an electrocardiogram by a prolonged but constant interval. It is often harmless, though medicines or heart disease may contribute.

### **First-Episode Psychosis**

The first recognized period of losing contact with reality, such as hearing or seeing things that are not present or holding fixed false beliefs. Urgent assessment looks for medical, medication, substance, and psychiatric causes.

### **First-Pass Effect**

The reduction of an orally swallowed medicine by the intestine and liver before it reaches the bloodstream. This can reduce the amount available to act and may make another route necessary.

### **First-Pass Metabolism**

Breakdown of a swallowed medicine by the intestinal wall and liver before it enters general circulation. Extensive breakdown lowers the active amount reaching the body and can influence dose or route of administration.

### **Fish Odour Syndrome**

A rare metabolic disorder in which trimethylamine builds up and causes a fish-like smell in sweat, breath, or urine. Symptoms may vary with diet; testing, dietary guidance, and emotional support can help.

### **Fish Oil**

Oil containing omega-3 fatty acids, usually obtained from oily fish. Supplements may lower triglycerides at prescribed doses, but benefits vary and side effects or medicine interactions should be reviewed with a clinician.

### **Fish Poisoning**

Illness caused by toxins or germs in contaminated or improperly stored fish. Symptoms may include vomiting, diarrhea, flushing, tingling, weakness, or breathing difficulty; severe symptoms require urgent medical care.

### **Fish Tank Granuloma**

A chronic localized cutaneous mycobacterial infection caused by *Mycobacterium marinum*, contracted through contact of broken skin with non-chlorinated aquarium water or infected fish. It presents as persistent red-to-violaceous papules, nodules, or ulcers typically on the hands or fingers, occasionally ascending along lymphatic channels in a sporotrichoid pattern.

#### **Alternate Names:**

Aquarium Granuloma; Swimming Pool Granuloma

### **Fish Tapeworm Infection**

An intestinal infection acquired by eating raw or undercooked fish containing tapeworm larvae. It may cause abdominal discomfort, diarrhea, weight loss, or vitamin B12 deficiency and is usually treatable with antiparasitic medicine.

### **Fish Venom**

Poison produced by certain fish, usually delivered through spines or sharp body parts. It can cause immediate severe pain, swelling, numbness, tissue injury, or rarely whole-body poisoning.

### **Fish-Handler's Disease**

A skin infection caused by *Erysipelothrix* bacteria after handling infected fish, shellfish, meat, or animals. It usually causes a slowly expanding, painful purple-red lesion on the hand or fingers.

### **Fisher Syndrome**

An acute autoimmune variant of Guillain-Barré syndrome characterized by the classic clinical triad of acute ophthalmoplegia, ataxia, and areflexia, strongly associated with anti-GQ1b IgG ganglioside antibodies following an upper respiratory or gastrointestinal infection.

#### **Alternate Names:**

Miller Fisher Syndrome; MFS

### **Fisher's Syndrome**

A rare nerve disorder, usually after an infection, causing eye-muscle weakness, poor coordination, and reduced or absent reflexes. It is a variant of Guillain-Barré syndrome and needs prompt medical

---

assessment.

### **Fishhook Removal**

Removal of a fishhook from skin or deeper tissue. The method depends on the hook's location and depth; clinicians assess nerves, blood vessels, tendons, joints, tetanus protection, and infection risk.

### **Fissure**

A narrow crack, groove, or split in a body structure. An anal fissure is a painful tear in the anal lining, often causing sharp pain and bright-red bleeding during bowel movements.

### **Fissured Tongue**

A usually harmless condition in which grooves develop on the tongue's surface. Food may collect in the grooves and cause irritation or odor; gentle tongue cleaning and routine oral care can help.

### **Fistula**

An abnormal passage between two body parts, such as two organs, a blood vessel and another structure, or an organ and the skin. It may result from infection, inflammation, injury, surgery, or cancer. Symptoms and treatment depend on its location; some close with medical treatment, while others need a procedure or surgery.

#### **Alternate Names:**

Pathologic Fistula; Fistulous Tract; Abnormal Organ Tunnel

### **Fistula-In-Ano**

An abnormal tunnel between the anal canal and skin near the anus. It commonly follows an anal abscess and may occur with Crohn disease, causing ongoing drainage, discomfort, or recurrent abscesses.

### **Fistulotomy**

Surgery that opens an anal fistula, an abnormal tunnel between the anal canal and nearby skin, so it can heal from the inside out. It is mainly used for simple fistulas. More complex fistulas may need another treatment to protect bowel control.

#### **Alternate Names:**

Anal Fistulotomy; Fistula-in-Ano Surgery; Laying Open of Fistula

### **Fit**

A seizure or sudden episode of altered movement or awareness; this everyday term is also used to mean physically healthy. In medical writing, the specific symptom or diagnosis should be stated instead.

### **Fitz Hugh Curtis Syndrome**

Alternate spelling; see *Fitz-Hugh-Curtis Syndrome*.

### **Fitz-Hugh-Curtis Syndrome**

Inflammation and scarring around the liver caused by pelvic inflammatory disease, usually from gonorrhea or chlamydia. It can cause sharp right-upper-abdominal pain, fever, and pain that worsens with movement or breathing.

### **Five-Alpha Reductase**

An enzyme that changes testosterone into dihydrotestosterone (DHT). DHT supports male sexual development and contributes to prostate growth and male-pattern hair loss. Medicines that block this enzyme are used for an enlarged prostate and some types of hair loss.

#### **Alternate Names:**

5-Alpha Reductase; 5- $\alpha$ -Reductase; Steroid 5-Alpha Reductase

### **Five-Year Survival Rate**

The percentage of people alive five years after a diagnosis or treatment. It summarizes outcomes for groups, not an individual's exact prognosis, which also depends on cancer type, stage, health, and treatment.

### **Fixation**

Holding a body part, tissue, or broken bone in the correct position while it heals. Methods include stitches, wires, screws, plates, rods, casts, splints, or external frames.

### **Fixative**

A chemical used to preserve cells or tissue for labo-



---

ratory examination by preventing decay and maintaining structure. Common fixatives can irritate skin, eyes, and airways and must be handled safely.

### **Fixator Muscle**

A muscle that holds one part of the body steady while another part moves.

### **Fixed Drug Eruptions**

Recurrent sharply demarcated red or dusky patches caused by a medicine, often at the same sites each time exposure occurs. Lesions may blister or ulcerate and leave persistent dark pigmentation; stopping the culprit drug prevents recurrence.

### **Fixed Joint**

A joint that permits little or no movement because the bones are joined by bone or dense connective tissue. Some fixed joints are normal; others result from injury, disease, or surgical fusion.

#### **Alternate Names:**

Synarthrosis; Immovable Joint

### **Fixed Macrophage**

An immune cell that stays within a particular tissue instead of circulating in blood. It removes dead cells and microbes and helps coordinate inflammation and repair.

### **Flaccid**

Soft and lacking normal firmness or muscle tone. It may describe a limb, muscle, bladder, or penis; sudden flaccid weakness, especially on one side, can signal a neurologic emergency.

### **Flagellin**

The main protein in a bacterial flagellum, the whip-like structure that helps some bacteria move. Human immune cells can recognize flagellin and respond with inflammation.

### **Flagellum**

A long, whip-like structure that helps a cell or microorganism move. Human sperm and some bacteria have flagella, which differ in structure and function.

### **Flail Chest**

A serious chest injury in which several ribs break in more than one place, leaving part of the chest wall unstable. Severe pain and difficult breathing can result, often with lung bruising and need for urgent care.

### **Flakka**

A street name for illicit synthetic stimulant drugs, especially alpha-PVP. Overdose can cause extreme agitation, hallucinations, seizures, dangerously high temperature, muscle breakdown, irregular heart-beat, or death and requires emergency treatment.

### **Flanged Needle**

A needle with a widened rim or attached flange that helps stabilize it, limit how far it enters, or connect it to another device. The flange may reduce movement and tissue injury during sampling, injection, drainage, or infusion.

### **Flank**

The side of the body between the lower ribs and the hip, including part of the abdomen and back. Pain in this area may arise from the kidney, muscles, spine, bowel, or nearby organs.

### **Flank Pain**

Pain between the lower ribs and hip, often caused by kidney stones, kidney infection, muscle strain, spine problems, or nearby abdominal disease. Fever, vomiting, blood in urine, injury, or severe pain needs prompt evaluation.

### **Flap Operation**

Surgery that moves a piece of skin, tissue, or muscle, while preserving its blood supply, to cover a wound or reconstruct a damaged body part. Risks include infection, bleeding, and loss of the flap.

### **Flapping Tremor**

An involuntary, irregular jerky movement (asterixis) characterized by brief, sudden lapses of posture in the dorsiflexed hands when the arms are held extended with the wrists hyperextended, characteristic of hepatic encephalopathy, uremic encephalopathy,

---

and severe hypercapnia.

**Alternate Names:**

Asterixis; Liver Flap

**Flare**

A temporary increase in symptoms or inflammation after a disease has been stable or less active. Flares may be triggered by infection, stress, injury, or treatment changes; severe or unusual symptoms need assessment.

**Alternate Names:**

Exacerbation; Disease Flare-Up

**Flare-Up**

A sudden or gradual worsening of a long-term disease after a period of relative stability. The cause may be infection, stress, missed treatment, or disease progression and should be assessed if severe or persistent.

**Flashes**

Brief flickers or streaks of light seen without an outside light source. They may result from changes in the eye's gel, migraine, or inflammation; sudden flashes with floaters or missing vision require urgent eye assessment.

**Flashing**

Seeing brief flashes or streaks of light without an external source. New flashing, especially with floaters, a curtain-like shadow, eye pain, or reduced vision, can indicate a retinal emergency.

**Flat Affect**

Very little outward expression of emotion, such as reduced facial movement, gestures, or voice variation. It is a clinical sign, not a diagnosis, and may occur with psychiatric, neurologic, or medication-related conditions.

**Flat Bones**

Broad, relatively thin bones such as the skull, ribs, breastbone, and shoulder blades. They protect organs, provide surfaces for muscle attachment, and contain marrow that produces blood cells.

**Flat Feet**

A lowered or absent arch along the inner side of the foot, causing more of the sole to touch the ground. It is often painless, but rigid or painful flat feet may reflect tendon, joint, or bone problems.

**Flatfoot**

A foot with a reduced or collapsed inner arch. It may be flexible and painless or rigid and painful; causes include normal development, loose ligaments, tendon problems, injury, arthritis, or bone shape.

**Flatfoot (Pes Planus)**

Alternate name; see *Fallen Arches*.

**Flatulence**

Passage of gas through the anus, produced by swallowed air and bacterial digestion of food in the intestine. It may increase with diet, constipation, swallowing air, or digestive disorders; persistent troublesome symptoms need assessment.

**Flatus**

Gas released through the anus after air is swallowed or intestinal bacteria break down food. Passing gas is normal; inability to pass gas with increasing abdominal swelling, vomiting, or pain may indicate bowel obstruction.

**Alternate Names:**

Intestinal Gas; Expelled Gas

**Flatus and Bloating**

Alternate name; see *Intestinal Gas*.

**Flavin Adenine Dinucleotide**

A helper molecule made from vitamin B2 that allows enzymes to transfer electrons during energy production and other chemical reactions. It is commonly abbreviated FAD.

**Alternate Names:**

FAD; Riboflavin Dinucleotide

**Flaviviridae**

A family of single-stranded RNA viruses that includes dengue, yellow fever, Zika, hepatitis C,

---

and West Nile viruses. Different members spread through insects, blood, or other routes and cause different illnesses.

### **Flavivirus**

A group of RNA viruses that includes dengue, yellow fever, West Nile, Japanese encephalitis, and Zika viruses. Many spread through mosquitoes or ticks and can cause fever, inflammation, or nervous-system disease.

### **Flaviviruses**

Viruses in the flavivirus group, including dengue, Zika, yellow fever, and West Nile viruses. They are often spread by mosquitoes or ticks and may cause fever, bleeding, liver disease, birth defects, or brain inflammation.

### **Flavobacteriaceae**

A family of mainly gram-negative bacteria found in water, soil, and other environments. Most do not infect people, but some can cause opportunistic infections, especially in hospitalized or immunocompromised patients.

### **Flavobacterium**

A group of gram-negative bacteria commonly found in water and soil. A few species can cause opportunistic infections in people or animals, particularly when immunity is weakened.

### **Flavoxate**

A medicine used to relieve bladder spasms, urgency, frequency, or painful urination. It treats symptoms rather than infection or blockage and can cause dry mouth, blurred vision, dizziness, or constipation.

### **Flaxseed Extract**

A preparation made from flaxseed that may contain omega-3 fat, lignans, and soluble fiber. Effects vary by product; it may interact with medicines affecting clotting, blood sugar, or blood pressure.

### **Flea**

A small wingless insect that feeds on blood. Flea

bites can cause itchy skin reactions, and fleas can transmit infections between animals or, less commonly, to people.

### **Flea Bites**

Itchy red bumps caused by fleas feeding on the skin, often clustered around the ankles or lower legs. Scratching can cause infection; treating pets and the environment helps prevent new bites.

### **Flea Bites In Humans**

Small itchy bumps, often around the ankles or lower legs, caused by flea feeding. Allergic reactions may produce larger areas of swelling; control requires treating infested animals, bedding, rooms, and symptoms.

#### **Alternate Names:**

Pulicosis; Flea-Borne Dermatitis

### **Fleas**

Small wingless insects that feed on blood and infest animals, homes, or clothing. They can cause itchy bites, allergic skin disease, anemia in heavily infested animals, and transmission of some infections.

### **Flecainide**

A prescription antiarrhythmic medicine that slows electrical signals in the heart to prevent selected abnormal rhythms. It is not suitable for everyone and can worsen dangerous rhythms, especially with structural heart disease.

### **Fleischner Sign**

A chest radiographic sign of acute pulmonary embolism consisting of a prominent, enlarged central pulmonary artery terminating abruptly, caused by distension of the vessel by a large impacted thromboembolus and downstream vasoconstriction.

#### **Alternate Names:**

Fleischner's Sign; Dilated Central Pulmonary Artery Sign

### **Flesh-Eating Bacteria**

Informal name for bacteria that can cause necrotizing fasciitis, a rapidly progressive deep soft-tissue

---

infection. Severe pain, swelling, fever, skin discoloration, blisters, or serious illness requires emergency treatment.

### **Flesh-Eating Bug**

An informal name for necrotizing fasciitis, a rapidly spreading infection of deep skin and soft tissue. Severe pain out of proportion to visible changes, swelling, fever, blisters, or skin discoloration requires emergency care.

### **Flex**

To bend a joint so the angle between connected bones becomes smaller. For example, bending the elbow brings the forearm toward the upper arm; pain, swelling, stiffness, or weakness can limit this movement.

#### **Alternate Names:**

Flexion; Joint Bending

### **Flexed Breech**

Alternate name; see *Complete breech*.

### **Flexibility**

The range of movement available at a joint or group of joints. It depends on bones, muscles, tendons, ligaments, joint health, nervous-system control, age, training, and pain.

### **Flexible Nasolaryngoscopy**

An examination using a thin flexible camera passed through the nose to view the nasal passages, throat, and voice box. It helps investigate hoarseness, swallowing difficulty, blockage, lesions, and vocal-cord movement.

### **Flexible Sigmoidoscopy**

An examination using a flexible camera to view the rectum and lower colon. It can investigate bleeding, diarrhea, or bowel symptoms and monitor known disease; preparation and biopsy may be needed.

### **Flexion**

Movement that bends a joint and decreases the angle between connected bones, such as bending

the elbow, knee, hip, or fingers.

### **Flexor**

A muscle or tendon that bends a joint or body part. Flexors in the arm and hand bend the fingers and wrist; injury can cause pain, weakness, or loss of movement.

### **Flexor Digitorum Profundus**

A deep forearm muscle whose tendons bend the end joints of the fingers and also assist wrist movement. Its inner part is supplied by the ulnar nerve and its outer part by the median nerve.

### **Flexor Digitorum Superficialis**

A forearm muscle whose tendons bend the middle joints of the fingers and assist wrist movement. Damage to the muscle, tendon, or median nerve can cause pain and difficulty bending the fingers.

### **Flexor Tendon Anatomy**

The arrangement and function of the tendons that bend the fingers, thumb, and wrist. Flexor tendons pass from forearm muscles through the wrist and hand in defined zones and sheaths; injury can impair finger flexion and requires assessment of tendon, nerve, and blood-vessel function.

### **Flexor Tenosynovitis**

A serious infection around a tendon that bends a finger, often after a cut or puncture. The finger may become swollen, stay partly bent, and hurt when straightened; urgent hand assessment and treatment are needed.

#### **Alternate Names:**

Suppurative Tenosynovitis; Kanavel Tenosynovitis

### **Flexural**

Relating to a skin fold or the inner surface of a bent joint, such as the elbow crease, back of the knee, armpit, groin, or fold beneath the breast.

### **Flibanserin**

A prescription medicine for persistent, distressing low sexual desire in some premenopausal women. It

---

is taken regularly rather than as needed and can cause dizziness, sleepiness, low blood pressure, or fainting, especially with alcohol.

### **Flight-Or-Fight Response**

The body's rapid response to perceived danger, releasing stress hormones that increase heart rate, breathing, alertness, and muscle readiness so a person can escape or confront a threat.

#### **Alternate Names:**

Sympathoadrenal Response; Acute Stress Response

### **Floater**

Alternate name; see *Floaters*.

### **Floaters**

Moving spots, threads, cobwebs, or rings in the field of vision caused by shadows from material in the eye's vitreous gel. They are often age-related, but sudden onset can indicate bleeding or a retinal tear.

### **Floating Kidney**

Unusual downward movement of a kidney when a person stands, sometimes called nephroptosis. It is often symptomless but may cause position-related flank pain, urinary symptoms, or rarely impaired urine drainage.

### **Floating Rib**

One of the eleventh or twelfth ribs, attached to the spine but not connected to the breastbone. These ribs are less protected and can be injured by direct trauma.

### **Floating Ribs**

The eleventh and twelfth pairs of ribs, which attach to the spine but not the breastbone. They are relatively mobile and less protected; fractures can cause pain and may injure nearby organs.

### **Flocculation**

The joining of tiny particles suspended in a liquid into larger clumps. Laboratory tests may use this visible reaction to detect antibodies or other substances; it is also used in water treatment and

pharmaceutical preparation.

### **Floor Of The Mouth Cancer**

Cancer arising in the tissues beneath the tongue, most often from surface cells. A persistent ulcer, lump, pain, bleeding, numbness, or difficulty swallowing or speaking needs prompt examination.

#### **Alternate Names:**

Oral Floor Squamous Cell Carcinoma; Sublingual Malignancy

### **Floppy Baby Syndrome**

An informal description of an infant with unusually low muscle tone and a limp posture. Causes include brain, spinal cord, nerve, muscle, genetic, or metabolic disorders and require medical evaluation.

### **Floppy Iris Syndrome**

A cataract-surgery complication in which the iris becomes unusually loose, moves with fluid currents, and narrows the working space. It is associated with some prostate medicines and should be anticipated by the surgeon.

### **Floppy Valve Syndrome**

A condition in which the mitral valve's leaflets bulge backward into the upper heart chamber when the heart contracts. It is often harmless but may cause palpitations, chest discomfort, or leakage through the valve.

#### **Alternate Names:**

Mitral Valve Prolapse; Barlow Syndrome

### **Flora**

The microorganisms living normally or temporarily in a body site or environment. Some protect against harmful organisms, but they can cause infection if they enter another body site or defenses are weakened.

### **Flossing**

Cleaning between teeth with dental floss or another interdental device to remove plaque and food that a toothbrush cannot reach. Gentle daily cleaning can help prevent gum disease and cavities.

---

## Flotation

A laboratory method that separates material by density. For example, a flotation test can concentrate parasite eggs from stool so they are easier to identify under a microscope.

## Flow Cytometry

A laboratory test that examines cells or particles one at a time as they pass through a laser beam. It can count cells and identify their features, helping diagnose blood cancers and immune disorders.

## Flow-Volume Loop

A graph showing how much air moves in and out of the lungs at different speeds during forceful breathing. Its shape can help identify airway narrowing, reduced lung size, or an upper-airway blockage.

## Flowmeter

A device that measures the rate at which a gas or liquid moves. In medicine, it may measure oxygen delivery, intravenous fluid flow, or breathing flow such as peak expiratory flow.

## Floxacin

An antibacterial medicine used in some countries to treat infections caused by susceptible bacteria, especially staphylococci. It can cause allergic reactions, diarrhea, liver problems, or other antibiotic-related effects.

## Floxuridine

An anticancer medicine that interferes with DNA production, slowing or killing rapidly dividing cells. It is used only in selected treatment plans and may cause low blood counts, mouth sores, diarrhea, or other toxicity.

## Flu

An acute respiratory infection caused by influenza viruses. It commonly causes sudden fever, cough, sore throat, headache, muscle aches, and fatigue and can lead to pneumonia or worsen chronic diseases.

## Flu (Influenza)

Alternate name; see *Flu*.

### Alternate Names:

Influenza; Grippe; Viral Flu

## Flucloxacillin

An antibiotic used for selected bacterial infections, especially those caused by susceptible staphylococci. It can cause nausea, diarrhea, allergy, or liver injury and should be taken only as prescribed.

## Fluconazole

An antifungal medicine used for selected yeast and other fungal infections. It can interact with many medicines and may cause nausea, rash, or liver injury; treatment depends on the infection and patient.

## Flucytosine

An antifungal medicine converted inside susceptible fungi into substances that interfere with genetic-material production. It is used for selected serious infections, usually with another antifungal, and requires blood-count and liver monitoring.

## Fludarabine

A chemotherapy medicine that interferes with DNA production in abnormal blood cells. It treats selected blood cancers but can severely suppress immunity and blood counts, increasing the risk of serious infections.

## Fludarabine Phosphate

A form of fludarabine used in selected blood-cancer treatment plans. It interferes with DNA production and can cause prolonged low blood counts, immune suppression, infections, and neurologic or other serious effects.

## Fludeoxyglucose F-18

A radioactive form of glucose used in positron-emission tomography scans. Active tissues absorb more of it, helping doctors assess some cancers, brain disorders, and heart muscle; uptake is not specific for cancer.



---

## Fludrocortisone

A medicine that helps the body retain salt and water and raise blood pressure. It treats some adrenal-hormone deficiencies and selected orthostatic disorders; monitoring is needed for swelling, high blood pressure, and low potassium.

## Fluid Around The Heart

Extra fluid in the sac surrounding the heart. Small amounts may cause no symptoms, while rapid or large collections can restrict heart filling and cause breathlessness, low blood pressure, or life-threatening tamponade.

### Alternate Names:

Pericardial Effusion; Hydropericardium

## Fluid Balance

The relationship between water entering and leaving the body. The kidneys, hormones, thirst, breathing, sweat, and stool help regulate it; disruption can cause dehydration, swelling, or abnormal blood salts.

## Fluid Imbalance

Too much or too little body water or an abnormal level of salts such as sodium or potassium. Causes include vomiting, diarrhea, kidney or heart disease, hormone disorders, dehydration, or excess fluid treatment.

## Fluid in Lungs

Alternate name; see *Pleural Effusion*.

## Fluid Management

Planning and monitoring fluids, salts, and urine output to maintain circulation and organ function. Treatment may involve drinking, intravenous fluids, diuretics, or fluid restriction according to the underlying problem.

## Fluid Overload

Excess water and salt in the body, causing swelling, rapid weight gain, high blood pressure, or fluid in the lungs. It may result from heart, kidney, or liver disease or too much fluid treatment.

## Fluid Therapy

Treatment using fluids, often through a vein, to restore hydration, blood volume, or salt balance. The type and amount depend on the cause; excess fluid can worsen swelling, heart failure, or lung congestion.

### Alternate Names:

Intravenous Fluid Therapy; Volume Resuscitation

## Fluid Wave

A physical test for fluid buildup in the belly (ascites); tapping one side of the stomach causes a ripple or wave that can be felt with the other hand on the opposite side.

## Fluke

A flat, leaf-shaped parasitic worm. Different species infect the liver, lungs, intestines, or blood vessels and are acquired through contaminated food, water, or contact with freshwater.

## Flumazenil

An antidote that can reverse sedation and slowed breathing caused by some benzodiazepine medicines. It can trigger seizures or sudden withdrawal and is not suitable for every overdose, especially mixed overdoses.

## Flunarizine

A medicine used in some countries to prevent migraine and treat selected balance disorders. It can cause sleepiness, weight gain, depression, or movement problems and should be prescribed with appropriate monitoring.

## Flunisolide

An inhaled or nasal corticosteroid used to reduce inflammation in asthma or allergic rhinitis. It may cause throat irritation, nosebleeds, or oral thrush; regular use does not relieve a sudden severe asthma attack.

## Fluocinolone

A corticosteroid applied to skin, scalp, or other affected areas to reduce inflammation and itching.

---

Prolonged or extensive use can thin the skin or rarely cause effects throughout the body.

### **Fluocinolone Acetonide**

A corticosteroid medicine applied to the skin or scalp to reduce redness, swelling, and itching. Overuse can thin the skin, worsen infections, or rarely affect the body's hormone system.

### **Fluocinonide**

A strong corticosteroid applied to selected inflammatory skin conditions to reduce itching, redness, and swelling. It should be used only as directed because prolonged or extensive use can damage skin or cause systemic effects.

### **Fluocortolone**

A corticosteroid used in selected skin or other inflammatory conditions. It reduces inflammation but can cause skin thinning, infection, or other steroid effects, especially when used for a long time or over large areas.

### **Fluorescein**

A bright fluorescent dye used in eye examinations and some imaging tests. In the eye, it highlights scratches, ulcers, and abnormal blood-vessel leakage; rare allergic reactions can occur after injection.

### **Fluorescein Angiography**

An eye test in which fluorescent dye is injected into a vein and photographs track its movement through retinal blood vessels. It can show leakage, blockage, or abnormal vessels in retinal disease.

### **Fluorescein Eye Stain**

A test in which dye is placed on the eye and viewed under blue light. It highlights scratches, ulcers, foreign-body injuries, and some dry-eye surface damage.

### **Fluorescence**

The emission of visible light by a substance after it absorbs light or other energy. Clinicians and researchers use it to identify cells, chemicals, microbes,

or structures in laboratory and imaging tests.

### **Fluorescence In Situ Hybridization**

A laboratory test that uses glowing DNA probes to find specific chromosome regions or gene changes in cells. It can detect selected missing, extra, rearranged, or amplified genetic material.

#### **Alternate Names:**

FISH; Molecular Cytogenetic Assay

### **Fluorescence Microscopy**

A microscope method that uses fluorescent labels to make selected cells, molecules, or microbes visible. It helps examine tissue structure, cell proteins, genetic material, and infectious organisms.

### **Fluorescent Dye**

A substance that absorbs light of one color and emits light of another. It can label cells, tissues, molecules, or microbes so they can be detected during laboratory or imaging examinations.

### **Fluoridation**

Adding a controlled amount of fluoride to water or dental products to strengthen tooth enamel and reduce cavities. Excessive fluoride during tooth development can cause dental fluorosis, which appears as enamel discoloration or pits.

### **Fluoride**

A mineral that strengthens tooth enamel, helps repair early damage, and reduces acid production by cavity-causing bacteria. It is found in some water supplies, toothpaste, mouth rinses, and professional dental treatments.

### **Fluoride In Diet**

Fluoride obtained from drinking water and foods or drinks made with fluoridated water. The right amount helps protect teeth, while excessive intake, especially in children, can damage developing enamel.

### **Fluoride Overdose**

Excessive fluoride swallowed over a short time. It

---

can cause nausea, vomiting, abdominal pain, low blood calcium, muscle spasms, abnormal heart rhythms, or seizures and requires urgent poison-control or medical advice.

### **Fluoride Poisoning**

Illness caused by swallowing too much fluoride. Early symptoms include nausea, vomiting, and abdominal pain; severe poisoning can lower blood calcium and cause muscle spasms, seizures, or dangerous heart rhythms.

### **Fluoride Toxicity**

Poisoning that can cause corrosive gastrointestinal pain, vomiting, hypocalcemia, hyperkalemia, tetany, seizures, hypotension, and ventricular dysrhythmias. Serum calcium, potassium, magnesium, and ECG should be monitored; urgent treatment may include calcium replacement, airway support, and toxicology consultation.

### **Fluoride Varnish**

A concentrated fluoride coating painted onto teeth by a dental professional to prevent or slow cavities. It hardens quickly and is used at intervals based on age, tooth condition, and cavity risk.

### **Fluorimetry**

A laboratory method that measures light emitted by a fluorescent substance. It can detect or estimate very small amounts of chemicals, medicines, proteins, or biological material.

### **Fluorine**

A highly reactive chemical element found in compounds such as fluoride. Small controlled amounts of fluoride help protect teeth, while excessive exposure can poison the body or damage developing bones and teeth.

### **Fluorine-18**

A radioactive form of fluorine used to label tracers for positron-emission tomography scans. Its short half-life allows useful imaging while limiting how long radioactivity remains in the body.

### **Fluorometholone**

A corticosteroid eye medicine used for selected inflammatory conditions. Prolonged use can raise eye pressure, delay healing, worsen infection, or increase cataract risk, so it requires medical supervision.

### **Fluorophotometry**

A method that measures fluorescent light from the eye or another sample. It can help estimate substances or tissue properties, but results must be interpreted with the test method and clinical findings.

### **Fluoropyrimidine**

A class of anticancer medicines that resembles substances used to build DNA and RNA. They disrupt genetic-material production in rapidly dividing cells and can cause mouth sores, diarrhea, low blood counts, and hand-foot symptoms.

### **Fluoroquinolone**

An antibiotic that kills bacteria by blocking enzymes needed for bacterial DNA copying. It is reserved for selected infections because it can cause tendon injury, nerve symptoms, heart-rhythm changes, blood-sugar changes, or severe diarrhea.

### **Fluoroquinolone-Associated Disability**

A term used for disabling, sometimes long-lasting or potentially permanent adverse effects linked to fluoroquinolone antibiotics. Reported effects may involve tendons, muscles, joints, peripheral nerves, the central nervous system, or other organs and may occur together. It is not a formally established single diagnosis, so the individual adverse effects should be identified and recorded separately.

#### **Alternate Names:**

FQAD; Fluoroquinolone Toxicity Syndrome

### **Fluoroquinolones**

A group of antibiotics that kill bacteria by blocking enzymes required for bacterial DNA replication. They treat selected infections but are avoided when safer options work because of potentially serious tendon, nerve, heart, and blood-sugar effects.

---

## Fluoroscopy

An X-ray method that produces moving images inside the body. It helps guide procedures and examine swallowing, blood vessels, joints, and digestive or urinary passages; contrast dye may be used and radiation exposure occurs.

## Fluorosis

A change in developing tooth enamel caused by too much fluoride during childhood. It usually appears as white streaks or spots; more severe cases cause brown discoloration or pits.

## Fluorouracil

An anticancer medicine that interferes with DNA production. It is applied to selected precancerous skin lesions or given for some cancers; possible effects include mouth sores, diarrhea, low blood counts, and serious toxicity in susceptible people.

## Fluorouracil Side Effect

An unwanted effect of fluorouracil, such as mouth sores, diarrhea, nausea, low blood counts, hand-foot skin reactions, or heart problems. Severe symptoms require prompt medical review and treatment adjustment.

## Fluoxetine

An antidepressant that increases available serotonin in the brain. It treats depression and several other conditions, but may cause nausea, sleep or sexual problems, agitation, or rarely dangerous serotonin excess or suicidal thoughts.

## Fluoxymesterone

A synthetic androgen medicine used in selected hormone-related conditions. It can affect the liver, cholesterol, mood, fertility, prostate, and blood count and should be used only with medical supervision.

## Flupentixol

An antipsychotic medicine used in some countries for schizophrenia and other psychiatric conditions. It can reduce hallucinations and delusions but may cause movement problems, sleepiness, hormonal

changes, or rare serious reactions.

## Fluphenazine

An antipsychotic medicine used to treat schizophrenia and related psychotic symptoms. It can cause stiffness, tremor, restlessness, sleepiness, hormonal changes, and rare but serious involuntary movements or fever.

## Fluphenazine Hydrochloride

The hydrochloride salt form of fluphenazine, an antipsychotic medicine for selected psychotic disorders. Monitoring is needed because it can cause movement disorders, sedation, hormonal effects, and other serious adverse reactions.

## Flurandrenolide

A corticosteroid applied to the skin to reduce inflammation, redness, and itching. Prolonged, extensive, or covered use can thin the skin, worsen infection, or allow steroid effects throughout the body.

## Flurazepam

A long-acting sedative medicine used for short-term treatment of insomnia in some settings. It can cause next-day drowsiness, falls, memory problems, dependence, and dangerous breathing suppression with alcohol or opioids.

## Flurbiprofen

A nonsteroidal anti-inflammatory medicine used for pain and swelling and, in some forms, eye inflammation. It can cause stomach bleeding, kidney problems, heart risks, or allergic reactions and should be used as directed.

## Flush

A sudden feeling of warmth or visible reddening caused by increased skin blood flow. In healthcare, flush can also mean washing a tube, body passage, or intravenous line with fluid.

## Flushable Reagent Stool Blood Test

Alternate name; see *Fecal Occult Blood Test*.

---

## Flushing

Temporary redness and warmth of the skin caused by increased blood flow. Triggers include heat, exercise, emotion, alcohol, medicines, fever, or hormone disorders; flushing with wheezing, diarrhea, faintness, or palpitations needs evaluation.

## Flutamide

A medicine that blocks male-hormone effects and is used with other hormone treatment for prostate cancer. It can seriously injure the liver and may cause nausea, breast tenderness, or reduced blood counts.

## Flutter

A rapid, regular, repetitive movement or contraction. Atrial flutter is an abnormal heart rhythm caused by a fast electrical circuit in the upper chambers and can increase the risk of stroke.

## Flutter Valve

A handheld breathing device that creates gentle pressure and vibrations during exhalation. It can help loosen and move mucus in some people with bronchiectasis, cystic fibrosis, or chronic lung disease.

## Fluvastatin

A statin medicine that lowers low-density lipoprotein cholesterol by reducing cholesterol production in the liver. It lowers cardiovascular risk but can cause muscle symptoms, liver abnormalities, or rare muscle breakdown.

## Fluvoxamine

An antidepressant that increases serotonin signaling and is used mainly for obsessive-compulsive disorder and selected other conditions. It can cause nausea, sleep changes, sexual problems, medicine interactions, or rarely serotonin toxicity.

## Flux

An old, nonspecific term for diarrhea or an excessive discharge, especially from the bowel. Modern medical descriptions should identify the actual symptom, duration, severity, and suspected cause.

## FM

An abbreviation sometimes used for fibromyalgia, a long-term condition causing widespread pain, fatigue, poor sleep, and concentration problems. Abbreviations can be unclear, so the full term is preferred in patient records.

### Alternate Names:

Fibromyalgia; Fibromyalgia Syndrome

## FMRI

Functional magnetic resonance imaging maps changes in brain blood flow and oxygen use while a person performs tasks or rests. It can help identify language or movement areas and does not use X-ray radiation.

## FMS

An abbreviation sometimes used for fibromyalgia syndrome, a long-term condition with widespread pain, fatigue, poor sleep, and cognitive symptoms. The full term is preferred because FMS can have other meanings.

### Alternate Names:

Fibromyalgia; Chronic Widespread Pain

## FNA

An abbreviation for fine-needle aspiration, a procedure using a thin needle to collect cells or fluid from a lump or organ for laboratory examination. The full term is clearer in patient information.

## FND

Functional neurological disorder causes real weakness, abnormal movements, seizure-like episodes, sensory changes, or other nervous-system symptoms because nerve signaling is disrupted. Symptoms are not imagined; diagnosis is based on characteristic examination findings and may require tests to exclude other causes.

### Alternate Names:

Functional Neurological Disorder; Conversion Disorder

## Foam Cell

A cell containing many fat droplets, giving it a

---

foamy appearance under a microscope. Foam cells, often immune cells that have taken up cholesterol, are common in early artery-wall plaque.

### **Foam Cells**

Cells filled with fat droplets, often immune cells that have taken up altered cholesterol. They commonly form early in artery-wall plaque and contribute to inflammation and atherosclerosis.

### **Focal**

Limited to one specific area rather than spread widely. A focal finding may involve an image, seizure, infection, injury, or neurologic problem and must be interpreted according to its location.

### **Focal Adhesion**

A specialized attachment site where a cell connects to its surrounding support material. It helps cells stick, sense physical forces, move, and send signals during growth and tissue repair.

### **Focal Aware Seizures**

Seizures arising from a localized brain network in which awareness is preserved during the event. Symptoms may be motor, sensory, autonomic, cognitive, emotional, or experiential.

### **Focal Cortical Dysplasia**

A localized congenital malformation of neocortical architecture and development characterized by disrupted cortical lamination, bizarre dysmorphic neurons, and abnormal balloon cells (in type IIb). It represents the most frequent histopathological etiology of medically refractory focal epilepsy in children and young adults requiring surgical resective therapy.

#### **Alternate Names:**

FCD; Taylor Dysplasia

### **Focal Dermal Hypoplasia Syndrome**

A rare X-linked disorder, usually in females, with thin or absent areas of skin following Blaschko lines and variable abnormalities of limbs, teeth, eyes, hair, and other organs. Management is multidisciplinary and tailored to the findings.

### **Focal Glomerulosclerosis**

Scarring affecting some kidney filters or only parts of affected filters. It can cause protein to leak into urine, swelling, high blood pressure, and gradual loss of kidney function.

### **Focal Impaired Awareness Seizures**

Focal-onset seizures in which awareness is impaired during part or all of the seizure. Automatism, behavioral arrest, confusion, and postictal amnesia may occur.

### **Focal Neurologic Deficits**

Loss of a particular function, such as strength, sensation, speech, vision, or coordination, because of a localized problem in the brain, spinal cord, or a nerve. Sudden onset may indicate stroke.

### **Focal Neurological Deficit**

An isolated, discrete impairment of central or peripheral nervous system function attributable to damage within a specific anatomical neuroaxis location, such as unilateral hemiparesis, localized sensory loss, cranial nerve palsy, or aphasia.

### **Focal Neuropathy**

Damage or irritation affecting one peripheral nerve or a small group of nerves. It may cause pain, tingling, numbness, weakness, or altered movement in the area supplied by those nerves.

#### **Alternate Names:**

Mononeuropathy; Isolated Nerve Lesion

### **Focal Nodular Hyperplasia**

A usually harmless overgrowth of liver tissue related to an unusual local blood supply. It often causes no symptoms, is found on imaging, and rarely needs treatment unless its diagnosis is uncertain.

### **Focal Sclerosis**

Hardening or scarring limited to a particular area of tissue. In the kidney, it may describe scarring of some filters or parts of filters, which can impair filtration and cause protein in urine.



---

### **Focal Segmental Glomerular Sclerosis**

A kidney disorder in which scars affect some filters or parts of them. It can cause protein in urine, swelling, high blood pressure, and declining kidney function and may result from genetic, immune, or other conditions.

### **Focal Segmental Glomerulosclerosis**

A kidney disorder in which scars affect some filters or parts of them. It can cause heavy urine protein, swelling, high blood pressure, and kidney failure; causes include genetic changes, infections, medicines, and excess filtration.

### **Focal Seizure**

A seizure that begins in one area of the brain. It may cause unusual movement, sensation, behavior, emotion, or awareness and can spread to involve both sides of the brain.

### **Focal-Onset Seizure**

A seizure that begins in one area of the brain. Awareness may remain normal or become impaired, and symptoms depend on the affected area; the seizure may spread and cause whole-body stiffening and jerking.

### **Focus Words**

Words or short phrases used during relaxation or stress-management exercises to direct attention and support calm breathing. They may help coping but are not a treatment for an underlying medical disorder.

### **Focused Assessment with Sonography for Trauma**

A rapid bedside point-of-care ultrasonographic examination performed in acute trauma resuscitation to screen for free intraperitoneal blood (Morison pouch, splenorenal recess, pelvis) and hemopericardium (subxiphoid view).

#### **Alternate Names:**

FAST Exam; Trauma Ultrasound

### **FODMAP**

An abbreviation for fermentable oligosaccharides, disaccharides, monosaccharides, and polyols—short-chain carbohydrates that may be poorly absorbed and cause gas, bloating, pain, or diarrhea. A supervised reduction followed by gradual reintroduction may help some people with irritable bowel syndrome.

### **FODMAP Diet**

A short-term eating plan that reduces certain poorly absorbed carbohydrates that can cause gas, bloating, pain, or diarrhea. Foods are gradually reintroduced with professional guidance to identify personal triggers.

### **FODMAPs**

A group of short-chain carbohydrates that may be poorly absorbed in the small intestine and fermented by gut bacteria. In susceptible people they can cause gas, bloating, abdominal pain, or diarrhea.

### **Foetal Acidosis**

Excess acidity in fetal blood, usually caused by inadequate oxygen delivery or impaired gas exchange. Fetal heart-rate monitoring and, in selected situations, blood testing help assess its likelihood and severity.

### **Foetal Alcohol Syndrome**

A lifelong condition caused by alcohol exposure before birth, with characteristic facial features, poor growth, and brain-development problems affecting learning, behavior, or coordination. No amount of alcohol is known to be safe during pregnancy.

### **Foetor Hepaticus**

A distinctive, sweet, musty, or breath odor resembling rotten eggs and garlic, caused by the exhalation of volatile dimethyl sulfide and mercaptans that bypass hepatic metabolism due to portosystemic shunting and severe liver failure.

#### **Alternate Names:**

Fetor Hepaticus; Breath of the Dead

---

## Foetus

The developing human offspring from the end of the embryonic period, about nine weeks after fertilization, until birth. During this period, organs grow and mature and body functions become more coordinated.

## Foix-Alajouanine Syndrome

A subacute necrotizing myelopathy caused by a slow-flow spinal dural arteriovenous fistula (dAVF), resulting in chronic spinal venous hypertension, progressive venous congestion, ischemic myelopathy, and lower-extremity paraparesis with sphincter impairment.

### Alternate Names:

Subacute Necrotic Myelopathy; Spinal Dural AV Fistula Myelopathy

## Folate

Vitamin B9, needed for DNA production, cell division, and healthy red blood cells. Adequate folate before and during early pregnancy lowers neural-tube-defect risk; high-dose supplements can hide vitamin B12 deficiency.

## Folate Antagonist

A medicine that blocks folate-related processes needed to make DNA. Depending on the drug, it may treat cancer, inflammatory disease, or bacterial infection, but can cause low blood counts, mouth sores, or liver problems.

## Folate Deficiency

Too little vitamin B9, caused by poor intake, increased need, malabsorption, alcohol use, or some medicines. It can cause large abnormal red blood cells, anemia, fatigue, sore tongue, and pregnancy complications.

## Folate Deficiency Anemia

Anemia caused by too little folate, or vitamin B9. Red blood cells become unusually large and immature, causing fatigue, weakness, pale skin, shortness of breath, or a sore tongue. Blood tests help confirm the cause, and folate replacement is given after vitamin B12 deficiency is considered.

### Alternate Names:

Vitamin B9 Deficiency Anemia; Megaloblastic Folate Anemia

## Folate Vitamin B9

Alternate name; see *Folate*.

## Folate-Deficiency Anemia

Alternate spelling; see *Folate Deficiency Anemia*.

## Foley Catheter

A flexible tube placed through the urethra into the bladder to drain urine. A small balloon holds it in place; risks include discomfort, blockage, urinary infection, and urethral injury.

## Folic Acid

A manufactured form of vitamin B9 needed for DNA production and healthy blood cells. Supplements prevent many fetal neural-tube defects when taken before conception and treat folate deficiency.

## Folic Acid And Birth Defect Prevention

Taking folic acid before conception and during early pregnancy reduces the risk of neural-tube defects, which affect development of the fetal brain and spinal cord. Higher-risk people may need a prescribed dose.

## Folic Acid Deficiency Anemia

Alternate name; see *Folate Deficiency Anemia*.

## Folic Acid In Diet

Folate occurs naturally in leafy vegetables, beans, citrus fruits, and liver, while folic acid is added to some fortified foods. Intake supports blood formation and fetal development; needs increase during pregnancy.

## Folic Acid Supplementation

Taking vitamin B9 before conception and during early pregnancy to reduce neural-tube-defect risk. Dose depends on individual risk; supplementation also treats deficiency and supports DNA production and red-cell formation.

---

### **Folic Acid Test**

A clinical laboratory blood test measuring serum or red blood cell folate concentrations to evaluate megaloblastic anemia, malnutrition, malabsorption syndromes, or dietary deficiency.

#### **Alternate Names:**

Serum Folate Test; RBC Folate Assay; Folate Blood Test

### **Folie a Deux**

A psychiatric syndrome in which a delusional belief or paranoid ideation is transmitted from an individual with an established primary psychotic disorder to one or more emotionally close, dependent individuals who share the delusion.

#### **Alternate Names:**

Shared Psychotic Disorder; Shared Delusional Disorder

### **Folinic Acid**

An active form of folate used to treat or prevent folate-related toxicity and to support selected chemotherapy treatments. It can also treat certain folate deficiencies; its use depends on the clinical situation.

### **Follicle**

A small sac or organized tissue structure. A hair follicle produces a hair, while an ovarian follicle contains a developing egg; other follicles can occur in glands or lymphatic tissue.

### **Follicle Stimulating Hormone**

A hormone released by the pituitary gland. In females it helps ovarian follicles grow and produce estrogen; in males it supports sperm production. Levels vary with age, sex, and reproductive stage.

### **Follicle-Stimulating Hormone**

A pituitary hormone that supports egg-containing follicle development and estrogen production in females and sperm production in males.

### **Follicle-Stimulating Hormone**

### **Abnormalities**

Abnormally high or low FSH levels may reflect altered ovarian or testicular function, pituitary or hypothalamic disease, menopause, pregnancy, medicines, or a changing menstrual cycle. Interpretation requires age, sex, reproductive stage, and related hormone tests.

### **Follicle-Stimulating Hormone Blood Test**

A blood test measuring a hormone involved in ovarian function and sperm production. It helps evaluate infertility, menstrual changes, puberty problems, menopause, testicular or ovarian failure, and pituitary disorders.

### **Follicular**

Relating to a follicle, especially a hair follicle or an ovarian follicle. A follicular skin problem is centered around a hair and may appear as a bump, pustule, plug, or inflamed area.

### **Follicular Cyst**

A usually harmless ovarian cyst that forms when a follicle does not release an egg or does not shrink normally. It often causes no symptoms and disappears within a few menstrual cycles without treatment.

### **Follicular Fluid**

The liquid inside an ovarian follicle surrounding a developing egg. It contains hormones and other substances that support follicle growth and help coordinate ovulation.

### **Follicular Hyperkeratosis**

Excess keratin blocking hair follicles and causing small rough bumps. It commonly occurs with keratosis pilaris and, less often, with nutritional deficiency or other skin conditions.

### **Follicular Lymphoma**

A usually slow-growing cancer of immune cells that often causes painless swelling of lymph nodes. It may involve bone marrow or other organs and can sometimes change into a faster-growing lymphoma.

---

### **Follicular Thyroid Carcinoma**

A thyroid cancer that begins in follicle-forming cells and can spread through the bloodstream, especially to bone or lungs. Tissue examination is needed to show invasion and distinguish it from a benign growth.

### **Follicular Unit**

A natural group of one or more hair follicles along with nearby skin glands, nerves, and tiny muscles. Hair-transplant procedures may move these units from a fuller area to a thinning area.

### **Follicular Unit Transplantation**

Hair-restoration surgery that moves natural groups of hair follicles from a donor area to thinning or bald areas. Results take months, and possible problems include scarring, infection, and continued hair loss.

#### **Alternate Names:**

FUT; Strip Hair Restoration

### **Folliculitis**

Inflammation or infection around a hair follicle, causing small itchy, red, or tender bumps and sometimes pus. Causes include bacteria, yeast, friction, blocked skin, shaving, and ingrown hairs.

### **Folliculitis Decalvans**

A rare inflammatory scalp disorder causing repeated pustules, crusting, hair loss, and permanent scarring. Early dermatologic treatment aims to control inflammation and infection and limit further follicle destruction.

### **Fomentation**

Applying a warm, moist cloth or material to part of the body to soothe pain, relax muscles, or improve comfort. Excessive heat can burn skin and should be avoided.

### **Fomepizole**

An antidote for poisoning with methanol or ethylene glycol. It blocks formation of their toxic breakdown products and may be combined with intensive supportive care and dialysis.

### **Fomite**

An object or surface contaminated with an infectious organism that can transfer it to another person. Examples include towels, equipment, toys, or needles, although the importance of surface spread varies by organism.

### **Fomites**

Objects or surfaces that may carry infectious organisms and transfer them through contact. Examples include shared towels, medical equipment, toys, and other contaminated items; cleaning and hand hygiene reduce risk.

### **Fondaparinux**

An injectable anticoagulant that prevents formation or extension of blood clots by blocking activated factor X. It is used for selected clotting disorders and prevention after surgery; bleeding is its main risk.

### **Fondaparinux Sodium**

The sodium salt form of fondaparinux, an injectable medicine that reduces clotting by blocking activated factor X. It treats or prevents selected blood clots but can cause serious bleeding and requires dose adjustment in kidney disease.

### **Fong Disease**

Alternate name; see *Nail-Patella Syndrome*.

### **Fonofos**

A highly poisonous organophosphate insecticide. Exposure can cause sweating, excess saliva, vomiting, small pupils, muscle twitching, breathing difficulty, seizures, or weakness and requires urgent poison-control or emergency care.

### **Fonsecaea**

A group of darkly pigmented fungi that can cause slow-growing skin and soft-tissue infections called chromoblastomycosis. Infection usually follows injury and may produce warty plaques or nodules, especially in tropical regions.

---

## Fontan Procedure

An operation for some children born with only one effective pumping chamber. It directs blood returning from the body to the lungs without a separate pumping chamber, improving circulation but requiring lifelong follow-up for rhythm, heart, liver, clotting, and protein-loss problems.

### Alternate Names:

Fontan Operation; Fontan Circulation; Total Cavopulmonary Connection (TCPC)

## Fontanel

A soft gap between an infant's skull bones that allows the brain and skull to grow and helps the skull mold during birth. Its size and firmness can provide clues about hydration or pressure inside the skull.

## Fontanelle

A soft membranous space between an infant's skull bones. Fontanelles allow growth and molding during birth; the front one usually closes during the second year, while the back one closes earlier.

## Food Addiction

A nonformal term for intense cravings or loss of control around eating. It is not a universally accepted diagnosis; assessment should consider binge-eating disorder, mood, stress, medicines, sleep, and access to food.

## Food Additives

Substances added to food to preserve it or change its flavor, color, texture, or stability. Most are safe in approved amounts, but some people may have intolerance or allergy to particular additives.

## Food Allergies

Immune reactions to particular food proteins. They may cause hives, swelling, vomiting, wheezing, or life-threatening anaphylaxis; strict avoidance and an emergency plan are important for people with severe reactions.

### Alternate Names:

Food Allergy; Food Hypersensitivity

## Food Allergy

An immune reaction to a food protein. It may cause hives, swelling, vomiting, abdominal symptoms, wheezing, or anaphylaxis. Common triggers include milk, egg, peanut, tree nuts, fish, shellfish, soy, and wheat. Severe reactions require an emergency plan.

### Alternate Names:

Egg Hypersensitivity; Ovomucoid Allergy

### Alternate Names:

Cow's Milk Protein Allergy (CMPA); Dairy Hypersensitivity

## Food Contamination

The presence of harmful germs, toxins, chemicals, allergens, or foreign materials in food. Contamination can occur during production, storage, preparation, or serving and may cause infection, poisoning, allergic reactions, or injury.

## Food Hygiene

Practices that prevent food contamination and food-borne illness, including washing hands, separating raw and cooked foods, cooking thoroughly, chilling promptly, and using safe water.

## Food Intolerance

A troublesome reaction to a food that does not involve the immune system. Lactose intolerance is common; symptoms may include gas, bloating, cramps, or diarrhea and depend on the food and amount.

## Food Jags

A child's repeated preference for only a few foods or refusal of foods previously accepted. It is common during development but needs assessment if it limits nutrition, growth, social eating, or causes distress.

## Food Poisoning

Illness from eating food contaminated with germs, toxins, or harmful chemicals. Nausea, vomiting, diarrhea, abdominal cramps, and fever are common.

## Food Protein-Induced Allergic

---

## Proctocolitis

A non-IgE-mediated food reaction, usually in an infant, causing inflammation of the lower bowel and visible blood or mucus in otherwise well-appearing stools. Cow's-milk or soy protein is often involved.

## Food Protein-Induced Enterocolitis Syndrome

A delayed, non-IgE-mediated food allergy that usually causes repeated vomiting, diarrhea, pallor, lethargy, or dehydration several hours after eating a trigger food. It is most common in infants and young children.

## Food Safety

Handling, storing, preparing, and labeling food in ways that prevent germs, toxins, harmful chemicals, and allergens from causing illness.

## Food Supplement

A product taken to add nutrients or other substances to the diet, such as vitamins, minerals, herbs, or protein. Supplements can interact with medicines and are not automatically safe or effective.

## Food-Borne Illness

Illness caused by eating food contaminated with bacteria, viruses, parasites, toxins, or chemicals. Symptoms commonly include nausea, vomiting, diarrhea, cramps, and fever; safe handling and cooking help prevent it.

### Alternate Names:

Food Poisoning; Acute Alimentary Toxicosis

## Foodborne Disease

Illness acquired by eating or drinking material contaminated with germs, toxins, or harmful chemicals. Severity ranges from brief diarrhea to dehydration, organ damage, or life-threatening infection.

## Foot

The part of the lower limb below the ankle, made of bones, joints, muscles, tendons, ligaments, nerves, and blood vessels. Its shape and arches support standing, balance, shock absorption, and walking.

## Foot Drop

Difficulty lifting the front of the foot because of weakness in the ankle and foot muscles. It may cause tripping or a high-stepping walk and can result from nerve injury, spinal disease, muscle disease, or stroke.

### Alternate Names:

Peroneal Nerve Palsy; Steppage Gait

## Foot Fracture

A break in one or more foot bones caused by sudden injury, repeated stress, or weakened bone. Pain, swelling, bruising, and difficulty bearing weight are common; treatment ranges from protection and immobilization to surgery.

### Alternate Names:

Metatarsal Fracture; Tarsal Fracture

## Foot Pain

Pain in the foot caused by injury, overuse, arthritis, nerve problems, infection, or poor circulation. Persistent pain, inability to bear weight, numbness, discoloration, fever, or an open wound requires assessment.

## Foot Pain (Plantar Fasciitis)

Alternate name; see *Plantar Fasciitis*.

## Footling Breech

A variation of incomplete breech presentation in which one or both feet (single or double footling) lie below the buttocks and present first at the maternal pelvic inlet. It carries the highest risk of umbilical cord prolapse and fetal head entrapment among all breech presentations, usually requiring delivery by cesarean section.

### Alternate Names:

Incomplete breech; Footling breech presentation.

## FOP

Abbreviation; see *Fibrodysplasia Ossificans Progressiva*.

## Foramen

A natural opening or passage through a bone or



---

other body structure, usually allowing a nerve, blood vessel, or other tissue to pass.

### **Foramen Lacerum**

A triangular, irregular aperture in the middle cranial fossa base bounded by the sphenoid, apex of the petrous temporal, and basilar occipital bones, filled with fibrocartilage in life and crossed superiorly by the internal carotid artery as it enters the cavernous sinus.

### **Foramen of Monro**

A paired, crescent-shaped aperture situated between the column of the fornix and the anterior pole of the thalamus connecting each lateral cerebral ventricle with the midline third ventricle, permitting the physiological circulation of cerebrospinal fluid.

#### **Alternate Names:**

Interventricular Foramen

### **Foramen Ovale**

An opening between the upper chambers of the fetal heart that directs blood away from the unused lungs. It usually closes after birth; persistence is called a patent foramen ovale.

#### **Alternate Names:**

PFO

### **Foramen Rotundum**

A circular bony canal in the greater wing of the sphenoid bone that connects the middle cranial fossa with the pterygopalatine fossa, transmitting the maxillary division of the trigeminal nerve (cranial nerve V2).

### **Foramen Spinosum**

A small, round aperture located posterolateral to the foramen ovale in the greater wing of the sphenoid bone, transmitting the middle meningeal artery, middle meningeal vein, and the nervous spinosus (meningeal branch of CN V3) into the cranial cavity.

### **Foraminotomy**

Surgery that enlarges an opening where a spinal nerve leaves the backbone, relieving pressure caused

by arthritis, a disc problem, or other narrowing. Risks include infection, bleeding, nerve injury, and persistent symptoms.

### **Forbes-Albright Syndrome**

A clinical syndrome characterized by hyperprolactinemia, galactorrhea, and amenorrhea in women in the absence of recent pregnancy, classically caused by a prolactin-secreting pituitary adenoma (prolactinoma).

### **Force Plates**

Platforms that measure forces beneath the feet during standing, walking, or other movement. Clinicians and researchers use them to assess balance, gait, symmetry, and how weight is distributed.

### **Forced Expiratory Volume**

The amount of air forcefully exhaled during a measured time after taking a full breath. The first-second value helps assess narrowed or restricted airways and is interpreted with other lung-test results.

### **Forced Oscillation Technique**

A breathing test in which a device sends gentle pressure waves through the airways while a person breathes normally. It measures how easily air moves and can help assess lung disease when forceful breathing tests are difficult.

### **Forced Vital Capacity**

The total amount of air forcefully exhaled after a person breathes in fully. A low value may occur when the lungs are small or stiff or when airways prevent complete emptying.

### **Forceps**

A handheld instrument used to grasp, hold, or move tissue or medical materials. Different designs have smooth or toothed tips and are chosen according to the procedure.

### **Forceps Delivery**

An assisted vaginal birth in which forceps guide the

---

baby's head out of the birth canal. It may be used for prolonged labor, concerning fetal findings, or inability to push safely; tears and newborn injury are possible.

### **Forceps-Assisted Delivery**

A vaginal delivery assisted by placing forceps around the fetal head. It is considered only when the cervix is fully open, the head's position is known, and vaginal birth is judged safe; maternal and newborn injuries can occur.

### **Forchheimer Spots**

Small, pinhead-sized, reddish macules or petechiae appearing on the soft palate and uvula during the prodromal phase of rubella (German measles).

#### **Alternate Names:**

Forchheimer's Spots; Rubella Enanthem

### **Forcible Intercourse**

Sexual intercourse without freely given consent, involving force, threats, coercion, or inability to consent. It is sexual violence; survivors may need immediate safety support, medical care, emergency contraception, infection prevention, and counseling.

### **Fordyce Granules**

Ectopic, asymptomatic sebaceous glands that appear as clusters of tiny, painless yellowish-white papules on the buccal mucosa and the vermilion border of the lips, representing a common normal anatomical variant.

#### **Alternate Names:**

Fordyce Spots; Fordyce's Granules

### **Fordyce's Angiokeratomas**

Small, usually harmless dark-red or purple spots caused by widened tiny blood vessels, commonly on the scrotum or vulva. They may bleed after injury; unusual or changing lesions should be examined.

### **Forearm**

The part of the arm between the elbow and wrist, containing the radius and ulna bones, muscles, tendons, nerves, and blood vessels.

### **Forearm Circumference**

The distance around the forearm at a specified point. Measurement can help monitor growth, muscle loss, swelling, nutrition, or changes during treatment and rehabilitation.

### **Forehead**

The part of the face above the eyebrows and in front of the frontal skull bone. It contains skin, muscles, nerves, and blood vessels and helps with facial expression and protection.

### **Forehead Lift**

Cosmetic surgery that raises drooping eyebrows or forehead skin and may reduce wrinkles. Risks include scarring, numbness, asymmetry, hairline changes, bleeding, infection, and nerve injury.

#### **Alternate Names:**

Brow Lift; Corrugator Resection

### **Foreign Body**

An object that is not normally part of the body and becomes lodged in tissue or a body passage, such as glass, metal, food, or a splinter. It may cause injury, blockage, infection, or inflammation.

### **Foreign Body Airway Obstruction**

Partial or complete blockage of the breathing passages by an inhaled object or material. Severe choking with inability to speak, cough, or breathe is life-threatening and requires immediate first aid and emergency help.

### **Foreign Body Aspiration**

Inhalation of an object or food into the airway, especially in young children. Choking, persistent cough, wheezing, noisy breathing, or reduced breath sounds may occur; complete blockage is an emergency.

### **Foreign Body in Ear**

Alternate name; see *Ear Foreign Body*.

### **Foreign Body In The Nose**

An object lodged in a nasal passage, often in a child. One-sided blockage, foul discharge, bleeding,

---

pain, or difficulty breathing may occur; safe removal prevents tissue damage and infection.

### **Foreign Body Reaction**

Inflammation around material that remains in tissue, such as a splinter, stitch, implant, or device. It may cause a lump, redness, pain, drainage, or scar tissue and can resemble infection.

### **Foreign-Body Aspiration**

Entry of an object or food into the voice box, windpipe, or bronchial tubes, especially in young children. Sudden choking, cough, wheezing, or noisy breathing may occur; complete blockage is an immediate emergency.

### **Forensic**

Relating to the use of scientific or medical knowledge in legal investigations, court proceedings, identification, injury assessment, or determining the cause and manner of death.

### **Forensic Anthropology**

The examination of human remains, especially bones, for legal investigations. It can help determine whether remains are human, estimate biological characteristics, assess injury, and support identification.

### **Forensic Autopsy**

A specialized post-mortem examination performed under statutory or judicial authority by a forensic pathologist or medical examiner to determine the cause and manner of violent, sudden, suspicious, unnatural, or unattended deaths and preserve legal evidence.

#### **Alternate Names:**

Medico-Legal Autopsy; Coroner Post-Mortem

### **Forensic Autopsy of Blunt Force Trauma**

Postmortem examination of injuries caused by blunt impact, including abrasions, contusions, lacerations, fractures, and internal organ damage. Wound patterns, distribution, vitality, and associated evidence help reconstruct the event.

### **Forensic Autopsy of Sharp Force Injuries**

Postmortem examination of incised, stab, chop, or other sharp-force wounds. The examiner documents wound morphology, depth, direction, vital reaction, clothing damage, and associated injuries to help determine mechanism and manner of death.

### **Forensic Defense Wounds**

Injuries to the hands, arms, or other body areas that may occur when a person tries to protect against an assault. Their meaning must be assessed with the history and all other evidence.

#### **Alternate Names:**

Forensic Defense Injuries

### **Forensic Evidence**

Physical, biological, digital, or documentary material collected and analyzed during a legal investigation. Examples include blood, DNA, fibers, fingerprints, toxicology samples, photographs, and medical records.

### **Forensic Examination**

A structured medical or scientific assessment performed for legal purposes. It may document injuries, collect evidence after assault, assess impairment, examine a death, or analyze a scene or specimen.

### **Forensic Genetics**

The use of DNA testing to identify people, establish biological relationships, or compare biological evidence in legal investigations. Results require proper collection, chain of custody, and expert interpretation.

### **Forensic Laceration**

A torn wound caused by blunt force, usually with irregular bruised edges and tissue strands crossing the wound. These features help distinguish it from a clean cut caused by a sharp object.

#### **Alternate Names:**

Blunt Force Tear; Tears and Ruptures

### **Forensic Odontology**

The use of dental knowledge in legal investigations.

---

Comparing teeth, dental records, and radiographs can help identify remains and document dental injuries or other evidence.

### **Forensic Pathology**

The medical study of deaths and injuries for legal purposes. Forensic pathologists investigate sudden, unexpected, violent, or suspicious deaths and help determine the cause and manner of death.

### **Forensic Pathology of Firearm Wounds**

Examination of firearm-related injuries, including entrance and exit wounds, soot, stippling, bullet paths, projectile effects, clothing damage, and range-of-fire findings, integrated with imaging, scene evidence, and recovered bullets.

### **Forensic Pathology of Thermal Injuries**

Examination of burns and other heat-related injuries, including burn depth and distribution, soot or airway injury, heat artifacts, and evidence of vital reaction. Findings are interpreted with scene information and toxicology.

### **Forensic Psychiatry**

The application of psychiatry to legal questions, such as ability to stand trial, mental state during an alleged offense, risk assessment, involuntary treatment, and decision-making capacity.

### **Forensic Radiography**

The use of X-rays, computed tomography, magnetic resonance imaging, or other imaging in legal investigations. It can identify people, document injuries, locate objects, assess trauma, and support death investigations.

### **Forensic Toxicology**

The testing of blood, urine, hair, tissue, or other specimens for medicines, alcohol, drugs, poisons, and their breakdown products to assess possible roles in illness, impairment, poisoning, or death.

### **Forensic Toxicology Screen**

An initial laboratory test of biological samples for

medicines, alcohol, drugs, poisons, or their breakdown products. A positive screen usually needs confirmation and measurement before legal or medical conclusions are made.

### **Forensics**

The use of scientific methods to answer legal questions, including identification, toxicology, injury analysis, and interpretation of biological or physical evidence. Reliable findings require validated methods, documentation, and expert interpretation.

### **Foreskin**

The fold of skin covering and protecting the head of the penis. It normally becomes retractable over time; forceful retraction can cause pain, tearing, or scarring.

#### **Alternate Names:**

Prepuce; Penile Prepuce

### **Formaldehyde**

A strong-smelling chemical used in manufacturing and tissue preservation. It can irritate the eyes, skin, and airways, cause allergic reactions, and increase cancer risk with substantial or long-term exposure.

### **Formalin**

A water-based solution of formaldehyde used to preserve tissue for laboratory examination. It can irritate or burn the skin, eyes, and airways and must be handled with appropriate ventilation and protection.

### **Formalin-Fixed Paraffin-Embedded Tissue**

Tissue preserved in formalin, processed, and embedded in wax so thin sections can be examined. These blocks support microscopic, protein, and many genetic tests, although preservation can alter some results.

### **Formate**

A salt or ester of formic acid. In medicine, formate is important because it is a toxic breakdown product of methanol and can contribute to severe acidosis and vision loss after poisoning.

---

## Formed Elements Of Blood

The cells and cell fragments suspended in blood: red blood cells carry oxygen, white blood cells help defend against infection, and platelets help form clots and stop bleeding.

## Formetanate

A carbamate insecticide that can poison the nervous system by disrupting acetylcholine breakdown. Exposure may cause sweating, vomiting, small pupils, muscle weakness, breathing difficulty, or seizures and requires urgent medical advice.

## Formication

The feeling that insects are crawling on or under the skin when none are present. It may occur with nerve disease, menopause, substance withdrawal, medicines, or psychiatric illness and should be medically evaluated if persistent.

## Formoterol Fumarate

A long-acting inhaled medicine that relaxes airway muscles and helps prevent symptoms of asthma or chronic obstructive lung disease. It is not a rescue medicine for sudden severe breathing difficulty. When used for asthma, it should be combined with an inhaled corticosteroid rather than used alone.

## Formothion

An organophosphate insecticide that can poison the nervous system. Exposure may cause sweating, excess saliva, vomiting, muscle twitching, breathing difficulty, seizures, or weakness and requires urgent emergency or poison-control advice.

## Formparanate

A pesticide-related chemical whose health effects depend on the formulation, dose, and route of exposure. Suspected poisoning should be treated as urgent and assessed through poison-control or emergency services.

## Formula Feed

Prepared infant nutrition given by bottle, cup, or feeding tube when breast milk is unavailable, insufficient, or not chosen. The formula and preparation

method should match the infant's age and medical needs.

## Formula Feeding

Feeding an infant with commercially prepared formula instead of or in addition to breast milk. Safe preparation, clean equipment, correct dilution, and age-appropriate amounts help prevent infection, dehydration, and poor nutrition.

## Formulae

The plural of formula. In healthcare, the word may mean mathematical equations, medicine formulations, or prepared nutritional products such as infant formula.

## Formulary

A list of medicines approved or preferred by a health system, insurer, or institution. It may show coverage, restrictions, alternatives, and requirements for approval, but does not replace a clinician's prescribing judgment.

## Fornix

A recess or arch-shaped structure. In the eye, the conjunctival fornix is the fold where the eyelid lining joins the surface covering the eyeball; the term also describes parts of the brain and vagina.

## Forrest Classification

A standardized, internationally recognized endoscopic classification system used to stratify peptic ulcer hemorrhage into risk categories for rebleeding and determine the necessity of endoscopic hemostasis. Subdivided into active spurting hemorrhage (Ia), active oozing (Ib), visible non-bleeding vessel (IIa), adherent clot (IIb), pigmented flat spot (IIc), and clean-based ulcer (III).

### Alternate Names:

Forrest Endoscopic Bleeding Criteria; Forrest Staging

## Forrester Classification

A system that groups people with severe heart failure or shock according to blood flow and congestion. "Warm" means adequate flow and "cold" reduced

---

flow; “dry” means little congestion and “wet” significant congestion, helping guide urgent treatment.

**Alternate Names:**

Forrester Hemodynamic Subsets; Forrester Score

**Fortification**

Adding vitamins or minerals to foods to improve nutritional intake and prevent deficiency. Examples include iodine in salt, vitamin D in some dairy products, and folic acid in many grain products.

**Fortified Wine**

Wine with added distilled alcohol, giving it a higher alcohol content than ordinary wine. Excess use can cause intoxication, dependence, liver disease, injuries, and interactions with medicines.

**Fosaprepitant Dimeglumine**

An intravenous medicine converted to aprepitant that blocks a brain signaling pathway involved in nausea and vomiting. It helps prevent chemotherapy-related sickness and can interact with other medicines.

**Foscarnet Sodium**

An antiviral medicine used for selected serious or resistant cytomegalovirus and herpesvirus infections. It can damage the kidneys and disturb blood salts, so hydration and laboratory monitoring are important.

**Fosfomycin**

An antibiotic that blocks an early step in bacterial cell-wall production. It is used for selected urinary or difficult-to-treat infections; diarrhea, allergic reactions, and resistance can occur.

**Fosinopril**

An angiotensin-converting enzyme inhibitor used for high blood pressure and some forms of heart failure. It relaxes blood vessels and reduces fluid strain; cough, high potassium, kidney problems, or swelling can occur.

**Fossa**

A shallow hollow or depression in a bone or other body structure. It may provide space for a joint, muscle, blood vessel, nerve, gland, or part of an organ.

**Fossa Navicularis**

An expanded, boat-shaped anatomical dilation of the penile urethra situated within the glans penis immediately proximal to the external urethral meatus, lined by non-keratinized stratified squamous epithelium.

**Alternate Names:**

Navicular Fossa of the Urethra

**Fossula**

A small pit or shallow depression in an anatomical structure. Examples include small depressions in a bone or on a tooth; the term describes shape and location rather than a disease.

**Foster Kennedy Syndrome**

A neuro-ophthalmic syndrome characterized by ipsilateral optic atrophy and anosmia (due to direct compression of the optic nerve and olfactory tract by an anterior cranial fossa tumor, such as an olfactory groove meningioma) and contralateral papilledema (due to raised intracranial pressure).

**Alternate Names:**

Kennedy Syndrome; Gowers-Paton-Kennedy Syndrome

**Fotemustine**

A chemotherapy medicine that can cross into the brain and is used in some countries for selected advanced cancers. It can suppress bone marrow, increasing the risk of infection and bleeding.

**Fothergill Sign**

A physical examination test used to differentiate an abdominal wall mass (such as a rectus sheath hematoma) from an intra-abdominal mass; a hematoma or mass within the rectus sheath does not cross the midline and remains fixed and palpable during contraction of the rectus abdominis muscles when the patient raises their head.



---

**Alternate Names:**

Fothergill's Sign; Rectus Sheath Hematoma Sign

**Foul-Smelling Stools**

Strongly odorous feces that may result from diet, infection, bleeding, malabsorption, altered gut bacteria, or rapid transit; persistent change with weight loss or diarrhea warrants review.

**Found Dead**

A circumstance in which a person is discovered deceased without the death having been observed. Medical or forensic examination may determine identity, timing, cause, and manner of death.

**Four-Dimensional Ultrasound**

Three-dimensional ultrasound shown as movement over time, creating a live view. It may help assess fetal anatomy or other structures, but image quality and diagnostic value depend on the examination and equipment.

**Fourchette**

The small fold of tissue at the back of the opening of the vulva, where the inner labial folds meet. It may stretch or tear during childbirth or sexual injury.

**Fourier Analysis**

A mathematical method that separates a complex signal into simpler repeating frequencies. It is used in medical imaging, heart and brain signal analysis, and other measurements to identify patterns.

**Fournier Disease**

Alternate name; see *Fournier Gangrene*.

**Fournier Gangrene**

A rapidly spreading, life-threatening infection that destroys tissue around the genitals, perineum, or anus. Severe pain, swelling, fever, skin discoloration, or feeling very ill requires immediate surgery and antibiotics.

**Fourth Finger**

The ring finger, located between the middle finger

and little finger. It contains bones, joints, tendons, nerves, blood vessels, and skin that support movement and grasping.

**Fourth Heart Sound**

An extra, low heart sound heard right before the main heartbeat, caused by the heart working harder to push blood into a stiff heart muscle.

**Fourth Stage Of Labor**

The first hours after the placenta is delivered, when the birthing person is closely observed for heavy bleeding, abnormal blood pressure, poor uterine contraction, pain, and other early complications.

**Fourth Venereal Disease**

An old name for lymphogranuloma venereum, a sexually transmitted infection caused by particular Chlamydia strains. It may cause a small genital sore followed by painful groin swelling and, untreated, rectal complications.

**Fourth-Degree Tear**

A severe childbirth tear extending from the vaginal area through the anal sphincter and into the rectal lining. It needs careful repair, antibiotics when appropriate, follow-up, and monitoring for bowel-control problems.

**Fovea**

A tiny central depression in the retina responsible for the sharpest detailed and color vision. Damage to it can impair reading, face recognition, and central vision.

**Fovea Centralis**

The central pit of the retina with the greatest concentration of cone cells, providing the sharpest central and color vision. It has no major blood vessels crossing directly over its center.

**Foville Syndrome**

An inferior medial pontine stroke syndrome typically resulting from occlusion of paramedian perforating branches of the basilar artery. Characterized by

---

ipsilateral facial paralysis (cranial nerve VII), ipsilateral horizontal conjugate gaze palsy (abducens nucleus or PPRF), and contralateral hemiplegia of the upper and lower extremities (corticospinal tract).

**Alternate Names:**

Foville Inferior Pontine Syndrome

**Fowler Position**

A semi-upright position with the head and upper body raised, commonly about 45 to 60 degrees. It may improve comfort and breathing or be used during examinations and procedures.

**Fowler's Syndrome**

A rare cause of persistent difficulty emptying the bladder, usually in young women, related to abnormal relaxation of the urethral sphincter. Evaluation excludes blockage and nerve disease; treatment may require drainage or neuromodulation.

**Fox-Fordyce Disease**

A rare chronic skin disorder caused by blocked and inflamed sweat glands. It produces intensely itchy small bumps, commonly in the armpits, pubic region, or around the nipples.

**Foxglove Poisoning**

Poisoning from foxglove plants or medicines containing cardiac glycosides. It can cause nausea, vomiting, confusion, visual changes, dangerously slow or irregular heart rhythms, or cardiac arrest and requires emergency care.

**Fractional Excretion Of Sodium**

The percentage of filtered sodium leaving the body in urine. It can help assess kidney function and distinguish reduced kidney blood flow from some tubular injuries, but diuretics and other illnesses can make it misleading.

**Fractional Excretion of Urea**

The percentage of filtered urea excreted in urine. It can support assessment of kidney blood flow when diuretics affect sodium testing, but infection, bleeding, nutrition, and chronic kidney disease also influ-

ence the result.

**Alternate Names:**

FEUrea; Urinary Urea Excretion Index

**Fractional Exhaled Nitric Oxide**

A breath test that estimates type 2, often eosinophilic, airway inflammation. The result is interpreted with symptoms, age, treatment, and other clinical findings.

**Alternate Names:**

Fractional Exhaled Nitric Oxide; Exhaled NO Marker

**Fractional Flow Reserve**

A pressure measurement made during coronary angiography to determine whether a narrowed heart artery significantly limits blood flow. The result helps guide decisions about stents or medical treatment.

**Fractional IPV**

A smaller-than-usual dose of inactivated polio vaccine given into the skin rather than the muscle. When used in an approved schedule, it helps protect against polio; the number of doses and timing follow local immunization guidance.

**Fractional Shortening**

An echocardiographic measurement of left ventricular systolic contractile function representing the percentage change in the left ventricular internal diameter between end-diastole (LVIDd) and end-systole (LVIDs). Calculated as the difference divided by end-diastolic diameter multiplied by 100 percent, with normal values typically ranging from 25 to 45 percent.

**Alternate Names:**

LV Fractional Shortening; FS

**Fracture**

A break or crack in a bone caused by injury, repeated stress, or weakened bone. Pain, swelling, bruising, deformity, or inability to use the part may occur; treatment restores alignment and supports healing.

---

## Fracture Displacement

Movement of broken bone pieces out of their normal alignment, including sideways shift, angling, shortening, or rotation. The need for reduction or surgery depends on the bone, joint involvement, age, and function.

## Fracture Fixation

Stabilization of a broken bone with a cast, splint, plate, screw, nail, wire, or external frame. The goal is to maintain alignment while the bone heals and, when possible, preserve movement.

## Fracture Liaison Service

A coordinated healthcare program for people with fragility fractures. It evaluates bone strength and underlying causes, arranges treatment and fall prevention, and follows patients to reduce future fractures.

## Fracture Malunion

Healing of a broken bone in an incorrect position, such as with angling, shortening, or rotation. It may cause deformity, pain, weakness, altered movement, or abnormal stress on nearby joints.

## Fracture Nonunion

Failure of a broken bone to heal within the expected time, with no continuing progress toward healing. Pain, movement at the fracture site, or persistent weakness may occur, and causes include poor blood supply, infection, instability, and smoking.

## Fracture Of The Temporal Bone

A break in a skull bone containing parts of the ear and nearby nerves and blood vessels. It may cause hearing loss, dizziness, facial weakness, bleeding, or leakage of brain fluid and needs urgent assessment.

## Fracture Prevention

Measures that lower the chance of broken bones, including treating osteoporosis, improving strength and balance, preventing falls, correcting vitamin or mineral deficiency, and reviewing medicines that increase risk.

## Fracture-Fixation Device

A plate, screw, rod, pin, wire, or external frame used to hold broken bone pieces in alignment during healing. Choice depends on the fracture, bone quality, soft-tissue injury, and expected movement.

## Fractured Clavicle In The Newborn

A broken collarbone in a newborn, usually related to birth stress. The baby may move one arm less or cry when handled; most heal well with gentle support and follow-up.

## Fractured Elbow

A break involving one of the bones forming the elbow. It can cause pain, swelling, deformity, and limited movement and may injure nearby nerves or blood vessels.

## Fractured Ilium

A break in the broad upper part of the pelvic bone, usually from major trauma or weakened bone. It may cause severe pain and can accompany injury to pelvic organs or blood vessels.

## Fractured Nose

A break in the nasal bones or supporting cartilage, usually after a blow. Pain, swelling, bleeding, and deformity may occur; a collection of blood inside the septum requires urgent treatment.

### Alternate Names:

Nasal Bone Fracture; Nasal Trauma

## Fractured Rib

A break in one or more ribs, usually from injury or forceful coughing. Sharp pain worsens with breathing or movement; lung collapse, bleeding, or injury to nearby organs must be considered after significant trauma.

### Alternate Names:

Costal Fracture; Broken Ribs

## Fractures

Broken bones caused by injury, repeated stress, weakened bone, or disease. Assessment considers location, alignment, skin injury, blood supply, nerve

---

function, and associated damage; treatment may include immobilization, reduction, or surgery.

### **Fragile X Syndrome**

An X-linked inherited condition caused by a CGG-repeat expansion in the *FMR1* gene. It can affect learning, speech, behavior, social development, and physical features and is a common inherited cause of intellectual disability; expression varies because of repeat size, methylation, and sex.

### **Fragile X-Associated Tremor/Ataxia Syndrome**

A late-onset progressive neurodegenerative disorder affecting premutation carriers (55 to 200 CGG repeats) of the *FMR1* gene, typically manifesting in men older than fifty years. Characterized by progressive intention tremor, cerebellar gait ataxia, parkinsonism, cognitive executive decline, and symmetric hyperintensities of the middle cerebellar peduncles ("MCP sign") on T2-weighted MRI.

**Alternate Names:**  
FXTAS

### **FRAIL Scale**

A brief questionnaire assessing fatigue, ability to climb stairs, walking ability, illness burden, and unintentional weight loss. It screens for frailty risk but does not replace a complete medical and functional assessment.

### **Frailty**

A state of reduced physical reserve that makes an older or seriously ill person more vulnerable to falls, infection, hospitalization, disability, and other stress. Signs include weakness, slow walking, exhaustion, inactivity, and unintentional weight loss.

### **Frambesia**

Alternate name; see *Yaws*.

**Alternate Names:**  
Treponematoses; Pian; Bouba

### **Framboesia**

Another name for yaws, a chronic infection caused

by *Treponema pallidum pertenue*. It spreads mainly through skin contact in tropical areas and can cause skin lesions, bone pain, and deformity if untreated.

### **Frameshift Mutation**

A genetic change caused by inserting or deleting DNA bases in numbers that are not multiples of three. This shifts how the genetic code is read and often produces a severely altered or nonworking protein.

### **Framycetin**

An aminoglycoside antibiotic applied to selected bacterial skin, ear, or eye infections. It can cause local allergy or irritation; prolonged or extensive use may allow absorption and antibiotic-related toxicity.

### **Francisella**

A group of small bacteria that includes *Francisella tularensis*, which causes tularemia. People can acquire infection through ticks, animals, contaminated water, food, or inhaled particles.

### **Francisella Tularensis**

A bacterium that causes tularemia, usually acquired through ticks, deer flies, infected animals, contaminated water or food, or inhalation. It can cause fever, an ulcer with swollen lymph nodes, pneumonia, or severe systemic illness.

### **Francisellaceae**

A family of small bacteria that includes *Francisella* species. Some cause tularemia or related infections in animals and people; disease type depends on the species and route of exposure.

### **Francois Dyscephaly Syndrome**

Alternate name; see *Hallermann's Syndrome*.

### **Frank Breech**

The most common breech presentation (accounting for 50–70% of all breech births), in which the fetus is positioned with hips flexed and knees extended so that the feet lie near the head and the buttocks present first to the birth canal. Compared to com-

---

plete or footling breech, it has the lowest risk of umbilical cord prolapse.

**Alternate Names:**

Extended breech; Frank breech presentation.

**Frank Haemoptysis**

Alternate form; see *Haemoptysis*.

**Frank Sign**

A diagonal skin crease running across the lobule of the ear from the tragus to the posterior margin of the auricle, recognized as an independent clinical physical sign associated with coronary artery disease and peripheral vascular atherosclerosis.

**Alternate Names:**

Frank's Sign; Diagonal Earlobe Crease

**Frank-Starling Law**

The fundamental cardiac physiological principle stating that, within physiological limits, the stroke volume and contractile force of the ventricles increase in direct proportion to the volume of blood filling the heart at end-diastole (preload), resulting from optimized myocardial actin-myosin overlap.

**Alternate Names:**

Starling Law of the Heart; Heterometric Autoregulation

**Frank-Starling Mechanism**

The heart normally contracts more strongly when its chambers fill with more blood, up to a limit. This helps match the amount pumped with the amount returning to the heart.

**Frasier Syndrome**

A rare inherited disorder caused by changes in the WT1 gene. It can cause abnormal sexual development, progressive kidney disease, and increased risk of tumors in the gonads.

**Fraternal Twins**

Twins formed when two separate eggs are fertilized by two separate sperm. They are genetically like ordinary siblings and may be the same or different sex.

**FRAX Tool**

A calculator estimating a person's ten-year probability of a major osteoporosis-related fracture or hip fracture. It uses factors such as age, sex, bone density, previous fracture, and other risks and supports treatment decisions.

**Alternate Names:**

FRAX Fracture-Risk Calculator

**FRAXA**

Alternate genetic name; see *Fragile X Syndrome*.

**Freckle**

A small, flat, tan or light-brown area of increased skin pigment that often darkens with sunlight. Freckles are usually harmless, but a changing, bleeding, irregular, or unusual spot needs examination.

**Freckles**

Small, flat pigmented spots that often become darker after sun exposure and reflect inherited skin characteristics. They are usually harmless; changes in size, shape, color, symptoms, or bleeding require assessment.

**Free Fragment**

A piece of spinal-disc material that has broken away from the main disc and may press on a nearby nerve. It can cause back or leg pain, numbness, weakness, or bladder and bowel problems.

**Alternate Names:**

Sequestered Disc Fragment; Extruded Herniated Fragment

**Free Radical**

A highly reactive molecule with an unpaired electron. The body makes some during normal metabolism, while excessive amounts can damage cell membranes, proteins, and DNA as part of oxidative stress.

**Free Radical Injury**

Cell damage caused by excessive reactive molecules that alter fats, proteins, or DNA. It can contribute to inflammation and tissue injury, although the importance varies with the disease and exposure.

---

### Free T4 Test

A blood test measuring unbound thyroxine, the thyroid hormone available to tissues. It is usually interpreted with thyroid-stimulating hormone to assess overactive or underactive thyroid function and treatment response.

### Free Thyroxine Index

A calculated estimate of the thyroid hormone available to tissues, using total thyroxine and a measure of hormone binding. It can help assess thyroid function when binding-protein levels affect total hormone results.

### Free Water Clearance

A kidney measurement showing whether urine contains more or less water than blood plasma. Positive values indicate excretion of dilute urine; negative values indicate water retention and concentrated urine.

### Freeze Drying

A preservation process that freezes material and removes ice under low pressure. It helps preserve medicines, vaccines, biological samples, and foods while limiting damage from heat and moisture.

### Freezing

Exposure to very low temperature that can freeze tissue, cause frostbite, or dangerously lower body temperature. The term also describes preserving food, medicines, or samples by storing them at low temperature.

### Freiberg Disease

An idiopathic avascular necrosis (osteochondrosis) affecting the head of a metatarsal bone (most commonly the second metatarsal head), typically occurring in adolescent and young adult females, presenting with forefoot pain and stiffness exacerbated by weight-bearing.

#### Alternate Names:

Freiberg Infraction; Metatarsal Head Osteochondrosis

### Fremitus

A vibration felt or heard through body tissues. During a chest examination, transmitted voice vibrations can change with fluid, air, or solid tissue in or around the lungs.

### French Disease

An old name for syphilis, a sexually transmitted infection caused by *Treponema pallidum*. Untreated infection can progress from sores to widespread disease and later damage the brain, nerves, heart, or other organs.

### Frenectomy

A procedure that removes or releases a frenulum, a fold of tissue that may restrict tongue movement, pull on the gums, separate teeth, or interfere with dentures. Bleeding, infection, and scarring are possible.

### Frenulum

A fold of skin or mucous membrane that anchors a body part and limits its movement. Examples include the tongue fold, lip folds, and penile frenulum.

#### Alternate Names:

Frenulum; Lingual or Labial Fold

### Frenulum Breve

An abnormally short penile frenulum that may cause pain, tearing, or bleeding during erection or sexual activity. Treatment depends on symptoms and may include observation or a minor surgical procedure.

### Frenum

A narrow fold of tissue that attaches one body surface to another and limits movement. Examples include the folds beneath the tongue, inside the lips, and beneath the foreskin.

### Frequency

Urinating more often than usual, sometimes with only small amounts passed. It differs from producing a large total urine volume and may result from infection, pregnancy, diabetes, medicines, or bladder



---

disorders.

### **Frequency Of Urination**

Needing to urinate more often than usual. Causes include increased fluid intake, urinary infection, pregnancy, diabetes, diuretic medicines, bladder irritation, prostate enlargement, or other urinary disorders.

### **Frequency Range**

The lowest to highest sound or signal frequencies that a device can detect, produce, or amplify effectively. In hearing care, it helps describe which pitches a hearing aid can process.

### **Frequency Response**

How strongly a device or body system responds to different sound or signal frequencies. In a hearing aid, it describes the amount of amplification provided for low, middle, and high pitches.

### **Frequent Bowel Movements**

Passing stool more often than is usual for a person. Causes include diet, infection, medicines, inflammation, poor absorption, or functional bowel disorders; blood, fever, dehydration, or weight loss needs assessment.

### **Frequent Or Urgent Urination**

Urinating often or having sudden urges that are difficult to postpone. Infection, bladder irritation, diabetes, pregnancy, medicines, obstruction, or overactive bladder can cause these symptoms.

### **Frequent Urination**

Needing to urinate more often than usual. It may result from drinking more, infection, diabetes, pregnancy, medicines, prostate enlargement, overactive bladder, or another urinary condition.

### **Fresh Frozen Plasma**

A blood transfusion component prepared from whole blood or apheresis and frozen within eight hours of collection, containing all physiological coagulation factors and plasma proteins in normal physiolog-

ical concentrations, used to treat complex coagulopathies and massive transfusion.

### **Alternate Names:**

FFP

### **Fresh Specimen**

A recently collected biological sample tested promptly or preserved correctly before changes affect its cells, organisms, chemistry, or diagnostic accuracy.

### **Frey Syndrome**

An abnormal gustatory sweating and facial flushing over the parotid and preauricular region occurring in response to mastication or tasting food, caused by aberrant misdirected regeneration of severed parasympathetic secretomotor fibers of the auriculotemporal nerve into sympathetic sweat glands following parotid gland surgery or trauma.

### **Alternate Names:**

Auriculotemporal Syndrome; Gustatory Sweating

### **Frey's Syndrome**

Sweating, warmth, or flushing over the cheek or temple while eating, caused by nerve regrowth after parotid-gland surgery or injury. Symptoms may be treated with topical medicines, injections, or other specialist care.

### **Friable**

Describes tissue that is unusually soft, fragile, and easily torn. Friable tissue may bleed readily when touched and can occur with inflammation, infection, cancer, or poor healing.

### **Friction**

Rubbing between surfaces. Skin friction can cause irritation, blisters, or sores; reducing moisture, pressure, and rubbing helps prevent injury.

### **Friction Blisters**

Fluid-filled blisters caused by repeated rubbing or pressure, commonly on the feet or hands. Protecting the area, reducing friction, and keeping a ruptured blister clean helps healing; extensive, infected, or

---

recurrent blisters need assessment.

### **Friedlander Pneumonia**

A severe, acute lobar pneumonia caused by *Klebsiella pneumoniae*, classically presenting in alcohol-dependent or immunocompromised patients with high fever, dense upper lobe cavitory consolidation, and thick, gelatinous "current-jelly" bloody sputum.

### **Friedman Curve**

An older graph describing cervical opening over time during labor. Modern labor care recognizes wider normal variation and does not use this historical curve alone to diagnose slow labor or decide on intervention.

### **Friedreich Ataxia**

An inherited disorder that progressively damages nerves and coordination. It can cause unsteady walking, weakness, reduced sensation, spinal curvature, heart muscle disease, and sometimes diabetes.

### **Friedreich's Ataxia**

An inherited condition that gradually affects balance, coordination, sensation, and strength. Some people also develop heart muscle disease, spinal curvature, or diabetes and need regular neurologic and cardiac care.

### **Friedreichs Ataxia Syndrome**

An inherited disorder that gradually damages nerves and coordination. It can cause unsteady walking, weakness, reduced sensation, spinal curvature, heart muscle disease, and diabetes and usually requires long-term specialist care.

### **Froin Syndrome**

The triad of clear xanthochromia, elevated protein content, and rapid spontaneous clotting of cerebrospinal fluid (CSF) obtained below a complete anatomical obstruction of the spinal subarachnoid space (e.g., from a compressive spinal cord tumor or severe spinal stenosis).

#### **Alternate Names:**

Froin's Syndrome; Spinal Subarachnoid Blockade Syndrome

### **Froment Sign**

A clinical sign of ulnar nerve palsy; when attempting to firmly pinch a piece of paper between the thumb and index finger, paralysis of the adductor pollicis forces the patient to compensate by flexing the interphalangeal joint of the thumb using the flexor pollicis longus (innervated by the anterior interosseous nerve).

#### **Alternate Names:**

Froment's Paper Sign; Ulnar Pinch Sign

### **Frontal**

Relating to the forehead, frontal skull bone, or frontal lobe of the brain. The frontal lobe helps control voluntary movement, planning, attention, speech, behavior, and decision-making.

### **Frontal Bone**

The skull bone forming the forehead, roof of the eye sockets, and part of the nasal region. It contains the frontal sinuses and protects the front of the brain.

### **Frontal Bossing**

An unusually prominent forehead caused by forward bulging of the frontal skull bones. It may be familial or associated with bone disease, excess pressure in the skull, or skeletal disorders.

### **Frontal Lobe**

The front portion of the brain's outer layer. It helps control movement, planning, attention, judgment, behavior, personality, and speech production in the language-dominant hemisphere.

### **Frontal Lobe Disorder**

A condition affecting brain areas involved in planning, judgment, movement, attention, speech, and behavior. Causes include stroke, injury, tumor, dementia, infection, or other neurologic disease.

### **Frontal Lobe Epilepsy**

Epilepsy caused by seizures beginning in the frontal lobe. Episodes may involve sudden movements, unusual behavior, impaired awareness, or events during sleep and can be difficult to distinguish from

---

sleep disorders.

**Alternate Names:**

FLE; Frontal Lobe Epilepsy Syndrome

**Frontal Lobe Seizures**

Seizures beginning in the front part of the brain, often causing brief unusual movements, behaviors, sensations, speech changes, or impaired awareness. Symptoms depend on the exact brain area involved.

**Alternate Names:**

Frontal Lobe Ictal Episodes; Hypermotor Seizures

**Frontal Lobe Syndrome**

A clinical neuropsychiatric disorder resulting from traumatic, ischemic, or neurodegenerative injury to the frontal lobes, manifesting as executive dysfunction, behavioral disinhibition, apathy, abulia, impaired social judgment, utilization behavior, and loss of emotional control.

**Alternate Names:**

Frontal Dysexecutive Syndrome

**Frontal Sinus**

An air-filled cavity within the frontal skull bone above the eye. It connects with the nasal passages and can become inflamed or blocked during sinusitis.

**Frontal Sinusitis**

Inflammation of the air-filled cavities above the eyes, usually from a viral infection or blocked drainage. It can cause forehead pain, congestion, and discharge; severe headache, eye swelling, fever, confusion, or neurologic symptoms require urgent care.

**Frontalis Suspension**

An operation connecting the upper eyelid to the forehead muscle with a sling to lift a drooping eyelid. It is used when the eyelid-lifting muscle is weak, especially in congenital or childhood ptosis.

**Frontotemporal Dementia**

A group of progressive brain disorders affecting the frontal and temporal lobes. Early changes often involve personality, behavior, judgment, language, or movement rather than memory alone.

**Alternate Names:**

FTD; Pick Complex; Behavioral Variant FTD

**Frontotemporal Lobar Degeneration**

Progressive loss of nerve cells in the frontal and temporal brain regions. It can cause early changes in behavior, personality, language, judgment, or movement and may occur at a relatively young age.

**Frostbite**

A freezing injury to the skin and deeper tissues that can cause numbness, pale or waxy skin, swelling, blisters, and tissue death.

**Frotteuristic Disorder**

A mental disorder involving persistent sexual arousal from touching or rubbing against a person without consent, with clinically significant distress or impairment or acting on the urges. Nonconsensual touching is sexual assault.

**Frozen Section**

A rapid tissue examination performed during surgery. A specimen is frozen, cut into thin slices, and examined to provide immediate information about a lesion or surgical margin, although permanent sections are often more accurate.

**Frozen Shoulder**

A painful condition in which the shoulder capsule tightens, progressively limiting movement. It often improves slowly over months or years; diabetes, thyroid disease, and prolonged immobility increase risk.

**Alternate Names:**

Adhesive Capsulitis; Periarthritis of Shoulder

**Fructosamine**

A blood measurement of sugar attached to blood proteins, reflecting average blood glucose over roughly the previous two to three weeks. It can help when the longer-term hemoglobin A1c test is unreliable.

---

## Fructose

A natural sugar found in fruit, honey, some vegetables, and many sweetened foods. It is processed mainly by the liver; poor absorption can cause gas, bloating, cramps, or diarrhea after eating it.

### Alternate Names:

Fruit Sugar; Levulose

## Fructose Intolerance

Difficulty digesting or processing fructose, depending on the condition. Poor intestinal absorption causes gas, bloating, and diarrhea, while inherited inability to process fructose can cause low blood sugar and liver injury.

## Fructose Malabsorption

Incomplete absorption of fructose in the small intestine. Unabsorbed sugar is fermented by bacteria and can cause gas, bloating, cramps, or diarrhea; reducing personal dietary triggers may help.

## Fructose Metabolism

The body's processing of fructose, mainly in the liver, into substances used for energy or stored. Rare enzyme defects can cause harmless fructose in urine or serious low blood sugar and liver injury after intake.

## Fructose-1,6-Bisphosphatase Deficiency

An inherited disorder of gluconeogenesis caused by deficient fructose-1,6-bisphosphatase. Fasting or illness can lead to low blood glucose, lactic acidosis, ketosis, vomiting, and marked weakness.

## Fruity Breath

A sweet, acetone-like breath odor caused by ketones. In someone with diabetes, especially with high glucose, vomiting, abdominal pain, deep breathing, or confusion, it may signal life-threatening ketoacidosis.

## Frustration

An emotional response to blocked goals, unmet needs, or difficult circumstances. It may cause irritability, anger, distress, withdrawal, or behavior

changes and becomes clinically important when persistent or impairing.

## Fryns Syndrome

A rare inherited condition present from birth, often involving a hole in the diaphragm, characteristic facial features, and abnormal fingers or toes. Brain, heart, kidney, or genital abnormalities may also occur.

## FSH

An abbreviation for follicle-stimulating hormone, which supports ovarian follicle development and sperm production. Blood levels help assess infertility, puberty, menopause, and disorders of the pituitary gland, ovaries, or testes.

### Alternate Names:

Follicle-Stimulating Hormone; Follitropin

## Ft.

An abbreviation for foot or feet, a length of 12 inches or about 30.48 centimeters. In medical records, the surrounding context should distinguish it from other abbreviations.

## FTA-ABS Blood Test

A blood test detecting antibodies against *Treponema pallidum*, the bacterium that causes syphilis. It supports diagnosis but often stays positive after treatment and cannot by itself show whether infection is active.

## FTA-ABS Test

A syphilis blood test that detects antibodies against *Treponema pallidum*. Because antibodies may remain for life after infection, clinicians interpret the result with other tests, symptoms, treatment history, and exposure risk.

## Fuchs Endothelial Corneal Dystrophy

A usually bilateral disorder in which the cornea's inner pumping cells are lost and tiny deposits develop. Fluid then accumulates, causing morning blur, glare, swelling, pain, and reduced vision.

### Alternate Names:

---

Fuchs Dystrophy

### **Fuchs Endothelial Dystrophy**

A progressive, bilateral, hereditary dystrophic disorder of the corneal endothelium characterized by accelerated endothelial cell loss and the formation of extracellular collagenous excrescences (guttae) on Descemet membrane. As endothelial pump function fails, the cornea develops progressive stromal edema, epithelial microcysts, painful bullae (bullous keratopathy), and painless loss of visual acuity, classically worse upon awakening.

#### **Alternate Names:**

FECD; Fuchs Corneal Dystrophy

### **Fuchs' Dystrophy**

A usually inherited, slowly progressive disease of the inner layer of the cornea. The damaged endothelial cells cannot remove enough fluid, so the cornea becomes swollen and cloudy. It may cause blurred or hazy vision, glare, halos, light sensitivity, and vision that is worse in the morning. In advanced disease, painful blisters and corneal scarring may develop. It usually affects both eyes, although one eye may be worse than the other.

### **Fucose**

A sugar found in some cell-surface proteins and fats. It helps cells recognize and attach to one another and participates in immune signaling and other biological processes.

### **Fucosidosis**

A rare inherited disorder in which cells cannot break down certain sugar-containing substances. Their buildup progressively affects development, movement, bones, growth, sweating, and organ function.

### **Fucosyltransferase**

An enzyme that attaches the sugar fucose to proteins or fats. This changes how cells recognize one another, develop, interact with microbes, and display some blood-group markers.

### **Fuel Oil Poisoning**

Illness after swallowing, inhaling, or having fuel

oil on the skin. Inhalation can damage the lungs and cause coughing or breathing difficulty; vomiting should not be induced, and urgent poison-control advice is needed.

### **Fugue**

A rare dissociative state involving unexpected travel or wandering with loss of memory about personal identity or the past. It needs assessment for trauma, mental illness, substances, neurologic disease, and safety risks.

### **Fukuda Stepping Test**

A bedside vestibular examination test in which the patient marches in place for 50 to 100 steps with eyes closed and arms extended; a rotation of greater than 30 to 45 degrees towards one side indicates an ipsilateral unilateral peripheral vestibular hypofunction.

#### **Alternate Names:**

Unterberger-Fukuda Test; Vestibular Stepping Test

### **Fukuyama Congenital Muscular Dystrophy**

A rare inherited muscle disorder beginning in infancy, causing low muscle tone, weakness, contractures, and delayed development. Abnormalities of the brain and eyes, seizures, and heart disease may also occur.

#### **Alternate Names:**

FCMD; Fukuyama Type Congenital Muscular Dystrophy

### **Fulguration**

Destruction or sealing of tissue using electrical energy during a procedure. It may control bleeding or treat selected abnormal tissue, but nearby tissue can be burned or injured.

### **Full Agonist**

A medicine or chemical that binds to a cell receptor and produces the strongest response that receptor can generate. The actual effect still depends on dose, receptor number, tissue, and individual response.

---

## Full Blood Count

See *Complete Blood Count*.

### Alternate Names:

FBC; Complete Blood Count (CBC)

## Full Liquid Diet

A diet containing liquids and foods that become liquid at room temperature, such as milk, strained soups, yogurt, and custard. It may be used temporarily, but prolonged use can lack fiber and some nutrients.

## Full Sibling

A person who shares both biological parents with another person. This relationship is relevant to family medical history, genetic-risk assessment, and some tissue or organ-donor matching.

## Full-Thickness Skin Graft

A graft containing the outer skin layer and the entire dermis, moved to cover a wound or reconstruct a defect. It usually gives durable coverage but requires a suitable donor site and blood supply.

## Fulminant

Beginning suddenly and becoming severe very quickly. The term describes a dangerous pattern of illness, such as rapidly progressive liver failure, infection, or inflammation requiring urgent treatment.

## Fulminant Hepatitis

Rapid, severe liver injury causing loss of liver function, abnormal bleeding, and sometimes confusion or coma from toxin buildup. It is a medical emergency that may require intensive care and liver transplantation.

## Fulminant Myocarditis

Sudden, severe inflammation of the heart muscle causing poor pumping, dangerous rhythms, or shock. It requires intensive care and treatment of the cause and complications; temporary mechanical heart support may be needed.

## Fulminating Hyperpyrexia

Alternate name; see *Malignant Hyperthermia*.

## Fulvestrant

An injectable hormone treatment that blocks and helps destroy estrogen receptors in selected hormone-sensitive breast cancers. It can cause injection-site pain, hot flushes, joint symptoms, fatigue, and liver-test changes.

## Fumigation

Treating an enclosed area with gas, vapor, or smoke to kill pests or microorganisms. People and animals must leave beforehand and return only after the area is declared safe and adequately ventilated.

## Functional Asplenia

Poor or absent spleen function even though the spleen is present. It increases the risk of severe infections, especially from certain bacteria, so vaccination, prompt fever assessment, and sometimes preventive antibiotics are important.

## Functional Assessment

Evaluation of how well a person moves, communicates, thinks, and performs daily activities. It identifies support needs, tracks change, guides rehabilitation, and helps plan safe care.

## Functional Capacity

The level of physical, mental, communication, and self-care activity a person can safely perform under particular conditions. It depends on health, environment, assistance, and personal goals.

## Functional Constipation

Long-lasting constipation without an identified blockage or other clear disease causing it. Stools may be hard or difficult to pass, or a person may feel that the bowel has not emptied fully. It may be related to diet, low fluid intake, inactivity, or regularly ignoring the urge to pass stool.

## Functional Decline

A reduction in a person's ability to move, think,



---

communicate, or perform daily activities compared with their previous level. In older adults it may result from illness, medicines, hospitalization, pain, poor nutrition, or lack of activity.

### **Functional Diarrhea**

Long-lasting loose or watery stools without an identified infection or other clear disease causing them. Unlike irritable bowel syndrome, belly pain is not the main symptom. Diet, medicines, stress, and changes in bowel movement can contribute.

### **Functional Disorder**

A condition in which symptoms or impaired body function arise from altered processing or regulation without a structural abnormality sufficient to explain them. The symptoms are real and can cause substantial disability.

### **Functional Dyspepsia**

Ongoing upper-abdominal discomfort, burning, early fullness, or unpleasant fullness after meals without an ulcer or other structural cause explaining symptoms. It may involve altered stomach movement or gut sensitivity.

#### **Alternate Names:**

Nonulcer Dyspepsia; Nonulcer Stomach Pain

### **Functional Electrical Stimulation**

Electrical impulses applied to nerves or muscles to produce a useful movement, such as lifting the foot during walking. It may assist rehabilitation after stroke, spinal injury, or other nerve damage.

### **Functional Endoscopic Sinus Surgery**

A minimally invasive surgical procedure performed through the nostrils to treat chronic rhinosinusitis, recurrent sinus infections, or nasal polyposis that fail to respond to medical treatment. Using specialized high-definition fiberoptic endoscopes and micro-debriding instruments, an ear, nose, and throat (ENT) surgeon visualizes the narrow paranasal sinus drainage pathways without making facial or external skin incisions. The surgeon gently widens the natural sinus openings (ostia), clears diseased mucosal tissue or obstructing polyps, and straight-

ens anatomical obstructions. By restoring natural mucus clearance and airflow ventilation while preserving healthy normal mucous membranes, FESS relieves chronic facial pressure, nasal obstruction, and headache, allowing topical steroid rinses to reach the sinus cavities effectively.

#### **Alternate Names:**

FESS; Endoscopic Sinus Surgery; ESS; Functional Sinus Surgery

### **Functional Gastrointestinal Disorders**

Digestive symptoms caused by altered gut movement or sensitivity without an obvious structural disease. Symptoms may include pain, bloating, constipation, diarrhea, nausea, or early fullness.

#### **Alternate Names:**

Disorders of Gut-Brain Interaction; FGIDs

### **Functional Hypothalamic Amenorrhea**

Missing menstrual periods because stress, inadequate energy intake, excessive exercise, or disordered eating suppresses reproductive hormones. It can lower estrogen, impair fertility, and weaken bones, but often improves when the underlying imbalance is corrected.

#### **Alternate Names:**

Hypothalamic Amenorrhea; FHA

### **Functional Impairment**

Difficulty performing usual work, study, self-care, relationships, or other daily activities because of symptoms, illness, injury, or treatment. The degree of impairment is considered separately from the diagnosis and may change over time.

### **Functional Incontinence**

Urine leakage because a person cannot reach, recognize, or use the toilet in time, even when bladder storage may be adequate. Mobility, memory, vision, clothing, communication, or environmental barriers may contribute.

### **Functional Independence Measure**

A structured tool that rates how much help a person needs with self-care, toileting, transfers, walking, communication, and thinking. It is used mainly in

---

rehabilitation to track function and progress.

### **Functional MRI**

A magnetic-resonance scan that estimates brain activity from changes in blood flow and oxygen use. It can map language or movement areas before selected brain procedures and does not use X-ray radiation.

### **Functional Murmur**

An extra heart sound caused by increased blood flow through a structurally normal heart. It may occur with fever, anemia, pregnancy, or exercise and is assessed to exclude valve or heart disease.

### **Functional Neurologic Disorder**

A disorder in which altered nervous-system function causes genuine symptoms such as weakness, abnormal movements, functional seizures, or sensory changes. Diagnosis is based on characteristic positive clinical features showing incompatibility with recognized neurologic disease, rather than solely on the absence of structural abnormalities.

#### **Alternate Names:**

FND; Conversion Disorder; Functional Neurological Symptom Disorder

### **Functional Neurologic Disorder/Conversion Disorder**

Alternate name; see *Functional Neurologic Disorder*.

### **Functional Neurological Disorder**

Alternate name; see *Functional Neurologic Disorder*.

### **Functional Neurological Symptom Disorder**

Alternate name; see *Functional Neurologic Disorder*.

### **Functional Ovarian Cysts**

Ovarian cysts related to the normal menstrual cycle, usually a follicular cyst or corpus-luteum cyst. They commonly resolve without treatment but may cause pain, bleeding, rupture, or ovarian torsion.

### **Functional Residual Capacity**

The amount of air left in the lungs after a normal relaxed breath out. It helps keep air sacs open and may be reduced by obesity or restrictive lung disease or increased by air trapping.

### **Functional Screening**

A brief check of a person's ability to move safely, communicate, perform daily activities, and manage basic self-care. Abnormal results may lead to a fuller medical, occupational, or rehabilitation assessment.

### **Functional Status**

A person's ability to manage daily activities such as bathing, dressing, eating, toileting, walking, shopping, cooking, medicines, finances, and transportation. Decline may signal illness or a need for support.

### **Functional Urinary Incontinence**

Urine leakage that happens because a person cannot reach, recognize, or use the toilet in time, even when the bladder may work normally. Problems with walking, memory, vision, clothing, communication, or the environment can contribute; treatment addresses these barriers and other causes of leakage.

### **Functioning Carcinoid**

Alternate name; see *Carcinoid Tumor*.

### **Fundal Height**

The distance from the pubic bone to the top of the uterus, measured during pregnancy. It roughly follows weeks of pregnancy after about 20 weeks; differences may reflect dating error, growth problems, twins, or fluid changes.

### **Fundoplication**

Surgery that wraps the upper stomach around the lower esophagus to strengthen the valve and reduce acid reflux. It may also repair a hiatal hernia; swallowing difficulty, gas, or recurrent reflux can occur afterward.

---

## Fundoscopy

Examination of the back of the eye using an ophthalmoscope or camera. It views the retina, optic nerve, macula, and blood vessels to assess conditions such as diabetes, high blood pressure, or retinal disease.

## Fundus

The rounded or farthest part of a hollow organ. Examples include the upper part of the stomach, the top of the uterus, and the inside back surface of the eye.

## Fundus Oculi

The inside back surface of the eye visible through the pupil. It includes the retina, optic nerve, macula, and retinal blood vessels.

## Fundus Photography

Photographing the back of the eye to record the retina, optic nerve, macula, and blood vessels. Images help detect and monitor retinal disease and allow comparison over time.

## Fundus Uteri

The rounded upper part of the uterus above the openings of the fallopian tubes. Its position and size can be assessed during pelvic examination and pregnancy.

## Fungal

Relating to fungi or caused by a fungus. Fungal infections may affect the skin, nails, mouth, lungs, bloodstream, or other organs.

## Fungal Arthritis

A fungal infection inside a joint, usually after surgery, injury, injection, bloodstream spread, or with weakened immunity. It often develops slowly with persistent pain, swelling, and reduced movement and may need prolonged treatment and surgery.

## Fungal Corneal Ulcer

A serious fungal infection and open sore on the cornea, the clear front surface of the eye. It may

follow an injury involving plant material, contaminated contact lenses, or prolonged steroid eye-drop use. Severe pain, redness, light sensitivity, blurred vision, or a cloudy spot may occur. Eye examination and testing of a corneal sample can support the diagnosis; urgent specialist treatment is needed to protect vision.

### Alternate Names:

Fungal Keratitis; Keratomycosis; Mycotic Corneal Ulcer

## Fungal Endophthalmitis

A fungal infection inside the eye, often after bloodstream infection, surgery, injury, or weakened immunity. It can rapidly damage vision and requires urgent specialist treatment with antifungal medicine and sometimes surgery.

## Fungal Infection

An illness caused by fungi—microscopic organisms such as yeasts and molds—entering, growing, or multiplying in or on the body. Fungi are distinct from both bacteria and viruses. Many harmless fungi live naturally in the environment (such as soil and plants) or quietly on human skin and mucous membranes. An infection happens when fungi overgrow or break through the body's defenses. Fungal infections generally fall into two main groups: superficial or mild infections (such as athlete's foot, ringworm, jock itch, fungal nail infections, and oral or vaginal thrush, which affect the outer skin, hair, nails, or mucous membranes and cause itching, redness, flaking, or white patches) and deep or invasive infections (such as aspergillosis, cryptococcal meningitis, or mucormycosis, where fungal spores enter the lungs or bloodstream and spread to vital organs like the brain, lungs, heart, or kidneys). Deep infections are life-threatening and primarily strike individuals with weakened immune systems, such as people undergoing cancer treatment, taking immunosuppressive drugs, or living with uncontrolled diabetes or HIV. Fungal infections do not respond to antibiotics or antivirals; they are treated with antifungal medicines applied topically or taken orally or intravenously. Keeping the skin clean, dry, and practicing good hygiene helps prevent many common fungal infections.

---

**Alternate Names:**

Mycosis; Mycotic Infection; Fungal Disease

**Fungal Nail Infection**

A fungal infection of a fingernail or toenail causing thickening, discoloration, brittleness, or separation from the nail bed. Diagnosis may require testing because injury and skin disease can look similar.

**Alternate Names:**

Onychomycosis; Tinea Unguim; Nail Mycosis

**Fungal Nail Infections**

Alternate name; see *Fungal Nail Infection*.

**Fungal Nails**

Nails infected by fungi may become thick, yellow or brown, brittle, crumbly, or separated from the nail bed. Testing may be needed before treatment, which can be topical or oral.

**Alternate Names:**

Onychomycosis; Tinea Unguim

**Fungal Pathogenesis**

The ways fungi cause disease, including attaching to cells, growing into tissue, avoiding immune defenses, and releasing damaging substances. The process varies with the fungus, body site, and person's immunity.

**Fungal Pneumonia**

Pneumonia caused by fungi, including endemic fungi, *Aspergillus*, *Pneumocystis*, and other opportunistic organisms. Risk depends on geography, immune status, neutropenia, transplantation, and inhaled exposures; diagnosis may require cultures, antigen or molecular tests, imaging, and bronchoscopy.

**Fungal Resistance Mechanisms**

Changes that allow fungi to survive antifungal medicines, such as altering a drug target, pumping medicine out of the cell, or changing cell-wall production. Resistance can make infections harder to treat.

**Fungal Skin Infections**

Infections of the skin caused by dermatophytes or yeasts, including ringworm, athlete's foot, and candidiasis. They often cause red, itchy, scaly, or ring-shaped areas and thrive in warm, moist conditions.

**Fungating**

Describing a mass or wound that grows outward with an irregular, ulcerated, dead, or foul-smelling surface. It may occur with advanced cancer or severe chronic wounds and can bleed, leak, and become infected.

**Fungemia**

The presence of living fungi in the bloodstream, most often *Candida* in hospitalized patients. It can spread to organs and requires urgent cultures, source evaluation, and appropriate antifungal treatment.

**Fungi**

Organisms including yeasts, molds, and mushrooms. Some live harmlessly in the environment or body, while others cause infections of the skin, mouth, lungs, bloodstream, or other organs.

**Fungicide**

A chemical used to kill or inhibit fungi. Some fungicides are used on crops or surfaces, while selected medicines act against fungi in people; toxicity depends on the substance and exposure.

**Fungoid**

Resembling a fungus or mushroom in appearance. In medicine it may describe an outward-growing, irregular lesion or mass, but appearance alone does not establish its cause.

**Fungus**

A type of organism including yeasts and molds, with distinctive cell walls and membranes. Fungi reproduce in various ways and can cause superficial or invasive infection, especially when immunity is weakened.

---

## **Funiculus**

A cord-like bundle of nerve fibers or other structures. In the spinal cord, a funiculus contains nerve pathways; in the spermatic cord, it is the bundle carrying the vas deferens, blood vessels, nerves, and lymph vessels.

## **Funisitis**

Inflammation of the umbilical cord, usually associated with infection or inflammation of the membranes around the fetus. It may increase newborn infection risk and requires assessment of the placenta, mother, and baby.

## **Funnel Chest**

A chest-wall shape present from birth in which the breastbone and nearby ribs curve inward. It is often mild, but severe deformity may affect exercise, breathing, body image, or heart function.

### **Alternate Names:**

Pectus Excavatum; Sunken Chest

## **Funnel-Web Spider Bite**

A potentially life-threatening bite from a funnel-web spider. Venom may cause pain, sweating, tingling, muscle spasms, breathing difficulty, or collapse; keep the person still, apply pressure immobilization, and seek emergency care.

## **Furan**

A chemical compound used in manufacturing and found in small amounts in some heated foods. High or repeated exposure may harm the liver and other organs, so occupational exposure should be controlled.

## **Furocoumarin**

A plant chemical found in some citrus and other plants that can make skin unusually sensitive to ultraviolet light. Contact followed by sunlight may cause redness, blistering, or dark pigmentation.

## **Furrowed Tongue**

Alternate name; see *Fissured Tongue*.

## **Fursultiamin**

A fat-soluble form of thiamine, or vitamin B1, used in some countries as a vitamin preparation. Its clinical value and safety depend on the indication, dose, and other medicines.

## **Furuncle**

A painful, pus-filled infection of a hair follicle, usually caused by *Staphylococcus aureus*. It forms a red lump and may drain; larger, recurrent, facial, or fever-associated lesions need medical assessment.

## **Furuncle And Carbuncle**

A furuncle is a deep infected hair follicle, or boil. A carbuncle is a connected group of boils forming a larger, deeper infection; both may require drainage and sometimes antibiotics.

## **Furuncular Myiasis**

A fly-larva infestation forming a boil-like skin lump with a central opening through which the larva breathes. It may cause pain, movement sensations, or drainage and requires safe removal.

## **Furunculosis**

Repeated or multiple boils caused by infection of hair follicles, often with *Staphylococcus aureus*. Risk factors include close contact, skin disease, diabetes, and weakened immunity; treatment may require drainage, antibiotics, and prevention measures.

## **Fusariosis**

An opportunistic infection caused by *Fusarium* fungi, mainly affecting people with severely weakened immunity or prolonged low neutrophil counts. It can cause skin lesions, bloodstream infection, pneumonia, and disease in multiple organs.

## **Fusarium**

A group of environmental molds that can cause nail, eye, skin, sinus, lung, or widespread infections. Severe disease occurs mainly in people with weakened immunity, especially prolonged neutropenia.

---

### **Fusarium Infection**

An infection caused by *Fusarium* molds, which may affect the nails, eyes, skin, sinuses, lungs, or bloodstream. Invasive disease mainly affects people with severe immune suppression and requires urgent specialist antifungal care.

### **Fusarium Solani**

An environmental mold that can cause eye, nail, skin, or invasive infections, especially in people with weakened immunity. Its clinical importance depends on the specimen, body site, symptoms, and laboratory confirmation.

### **Fused Placenta**

A placenta in which areas that normally form separate lobes are joined. It is often an anatomical variation, but careful examination after delivery helps identify any retained placental tissue or abnormal blood vessels.

### **Fused Teeth**

Alternate name; see *Dental Fusion*.

### **Fusidic Acid**

An antibiotic that blocks bacterial protein production and is used for selected staphylococcal infections. Resistance can develop, so it is often combined with another antibiotic and used according to local guidance.

### **Fusiform**

Shaped like a spindle, wider in the middle and narrowing toward both ends. The term may describe a muscle, cell, swelling, or blood-vessel enlargement such as an aneurysm.

### **Fusion**

Joining of two structures that are normally separate. It may describe bones, joints, teeth, organs, genes, or tissues and can occur naturally, during development, after injury, or through surgery.

#### **Alternate Names:**

Dental Fusion; Synodontia

### **Fusion Gene**

A gene formed when parts of two separate genes become joined. It may produce an abnormal protein that drives some cancers and can sometimes be identified by genetic testing.

### **Fusion Of The Ear Bones**

Abnormal joining or stiffening of the tiny middle-ear bones, which can reduce sound transmission and cause conductive hearing loss. Evaluation identifies the cause and whether surgery or hearing support may help.

### **Fusion Protein**

A protein made when parts of two different proteins become joined, usually because their genes have fused. Some fusion proteins drive cancer; others are deliberately engineered for research or treatment.

### **Fusobacterium**

A group of bacteria that grow without oxygen and commonly live in the mouth and intestine. Some species can cause dental, throat, abdominal, wound, bloodstream, or other invasive infections.

### **Fusobacterium Infection**

Infection caused by *Fusobacterium* bacteria, which may spread from the mouth or intestine to the throat, lungs, abdomen, skin, joints, or bloodstream. Symptoms and treatment depend on the site and severity.

### **Fusobacterium Necrophorum**

A bacterium that can cause severe throat infection and Lemierre syndrome, in which infection spreads to neck veins and may produce infected clots, lung complications, and bloodstream infection.

### **Fusobacterium Nucleatum**

A common mouth bacterium associated with gum disease. It can occasionally cause bloodstream or other invasive infection and has also been studied for possible links with intestinal and pregnancy-related disease.



---

### **Fusospirochetal Gingivitis**

A painful gum infection involving fusobacteria and spirochetes, causing bleeding, ulcers, swelling, and foul breath. It is associated with poor oral hygiene, stress, smoking, or illness and needs dental treatment.



## G

### **G-Protein Coupled Receptors**

Proteins on the cell surface that receive signals from hormones, medicines, and other substances and pass those signals into the cell. They help control heart rate, vision, digestion, mood, and inflammation.

### **G6PD Deficiency**

An inherited condition in which red blood cells can break down after infections, fava beans, or certain medicines. Episodes may cause anemia, jaundice, fatigue, shortness of breath, or dark urine.

### **G6PD Deficiency in the Newborn**

Glucose-6-phosphate dehydrogenase deficiency presenting during the neonatal period, most often with jaundice from hemolysis. Early recognition helps avoid oxidant triggers and supports monitoring or treatment of significant hyperbilirubinemia.

### **Gaba Receptor**

A brain-cell protein that responds to GABA, a chemical messenger that usually reduces nerve activity. GABA receptors help regulate sleep, anxiety, muscle control, and seizures and are targets of several medicines.

### **Gabapentin**

A medicine used for certain seizures and nerve-related pain. It can cause sleepiness, dizziness, swelling, or impaired coordination; the dose often needs adjustment in kidney disease and stopping suddenly can cause problems.

### **GAD**

An abbreviation for generalized anxiety disorder,

involving excessive, difficult-to-control worry about several areas of life for at least six months, with symptoms such as restlessness, tension, fatigue, poor concentration, or sleep disturbance.

#### **Alternate Names:**

Generalized Anxiety Disorder

### **Gadolinium**

A metal used in a chemically bound form to improve some MRI images. Serious reactions are uncommon, but kidney function, pregnancy, prior reactions, and possible tissue retention are considered before use.

### **Gadolinium-Based Contrast Agent**

A substance injected during some MRI scans to make organs, blood vessels, or abnormal tissue easier to see. The agent and dose are selected according to kidney function, pregnancy, allergies, and the scan's purpose.

### **Gadolinium-Induced Nephrogenic Systemic Fibrosis**

A rare illness causing thick, tight, hardened skin and sometimes damage to internal organs after certain MRI contrast exposure, mainly in severe kidney disease. Safer contrast agents have made it very uncommon.

### **Gadopentetate Dimeglumine**

An injectable gadolinium contrast agent used to improve selected MRI images. It can cause allergic reactions and is used cautiously or avoided in severe kidney disease because of rare serious complications.

---

## **Gag Gene**

A retroviral gene that encodes structural proteins forming the virus's inner support, shell, and core. These proteins help assemble new virus particles inside infected cells.

## **Gag Protein**

A large retroviral protein that is cut into components forming the virus's inner support, protective shell, and genetic-material core. These parts are required to assemble new virus particles.

## **Gag Reflex**

A protective response in which touching the back of the throat causes throat-muscle contraction and sometimes retching. It helps prevent food or objects from entering the airway.

### **Alternate Names:**

Pharyngeal Reflex

## **Gagging**

Retching caused by stimulation of the throat or back of the tongue. It may occur with swallowing difficulty, reflux, infection, examination, aspiration, or distress.

## **Gain**

The amount by which a hearing aid or amplifier increases sound intensity, measured in decibels. Correct gain improves hearing while limiting excessive loudness, feedback, discomfort, and further hearing damage.

## **Gain-Of-Function Mutation**

A genetic change that makes a gene or its protein more active, gives it a new activity, or causes activity at the wrong time or place. Its effects range from harmless variation to inherited disease or cancer.

## **Gaisbock Syndrome**

A form of relative or stress polycythemia characterized by an elevated hematocrit resulting from decreased plasma volume rather than increased red cell mass, typically occurring in middle-aged, hypertensive, anxious, and overweight individuals.

### **Alternate Names:**

Stress Polycythemia; Polycythemia Hypertonica

## **Gait**

A person's pattern of walking, including speed, balance, step length, posture, arm movement, and turning. Gait reflects the combined function of the brain, nerves, muscles, joints, vision, and balance system.

## **Gait Abnormality**

A change from a person's usual walking pattern caused by problems with muscles, joints, nerves, balance, vision, heart function, or general health. Assessment examines speed, stability, symmetry, strength, sensation, and pain.

## **Gait Apraxia**

Difficulty starting or carrying out walking even though basic muscle strength is adequate. It may occur with disease or injury affecting the front of the brain and can cause short steps, freezing, or instability.

## **Gait Assessment**

An examination of how a person walks, including speed, steps, balance, posture, turning, and use of aids. It can reveal problems with strength, sensation, coordination, joints, vision, or thinking.

## **Gait Ataxia**

Unsteady, poorly coordinated walking caused by problems in the cerebellum, sensation pathways, balance organs, or other parts of the nervous system.

## **Gait Cycle**

The complete sequence of walking movements from one heel strike to the next heel strike of the same foot. It includes the stance phase when the foot bears weight and the swing phase when it moves forward.

## **Gait Disturbance**

An abnormal walking pattern involving speed, balance, coordination, strength, or step movement.

---

Causes include neurologic disease, joint or muscle problems, inner-ear disorders, pain, medicines, and systemic illness.

### **Gait Speed**

Walking speed measured over a set distance. It is a simple indicator of mobility and physical reserve, helps estimate fall or disability risk, and can track changes during illness or rehabilitation.

### **Gait Training**

Therapy that teaches or retrains safer walking after injury, illness, weakness, or balance problems. It may use exercises, parallel bars, walkers, canes, stairs, or other aids to improve independence.

### **Gaits**

Patterns of walking. Different gaits may reflect pain, weakness, joint disease, balance problems, or neurologic conditions; assessment considers the person's usual walking pattern and associated findings.

### **Galactocele**

A milk-filled breast cyst caused by blockage of a milk duct, usually during breastfeeding or after weaning. It commonly feels like a painless lump; examination and ultrasound distinguish it from other breast masses.

### **Galactocerebrosidase (GALC) Deficiency**

Alternate name; see *Galactocerebrosidase Deficiency*.

### **Galactocerebrosidase Deficiency**

An inherited lack of an enzyme needed to break down certain fats in nerve tissue. Severe deficiency causes Krabbe disease, which progressively damages the brain and nerves.

#### **Alternate Names:**

Krabbe Disease; Galactosylceramidase Deficiency

### **Galactocerebroside Beta-Galactosidase Deficiency**

Alternate name; see *Galactocerebrosidase Deficiency*.

### **Galactagogue**

A medicine, herb, food, or other substance claimed to increase breast-milk production. Evidence and safety vary, and effects may depend on feeding technique, milk removal, medical causes, and interactions.

### **Galactokinase Deficiency**

An autosomal recessive inborn error of galactose metabolism caused by mutations in the *GALK1* gene, which encodes galactokinase. The resulting accumulation of galactose in tissues leads to reduction by aldose reductase into galactitol in the crystalline lens, causing the progressive development of bilateral infantile cataracts without the hepatic, renal, or cognitive damage seen in classic galactosemia.

#### **Alternate Names:**

GALK Deficiency; Type II Galactosemia

### **Galactorrhea**

The inappropriate or spontaneous secretion of milk or milk-like fluid from one or both breasts unrelated to normal pregnancy or breastfeeding. It is most commonly caused by hyperprolactinaemia secondary to pituitary prolactinomas, dopamine-antagonist medications (such as antipsychotics and antiemetics), primary hypothyroidism (via elevated thyrotropin-releasing hormone), chronic kidney disease, or local chest wall stimulation.

#### **Alternate Names:**

Galactorrhoea; Non-Puerperal Lactation; Hyperprolactinemic Discharge

### **Galactorrhea-Amenorrhea Syndrome**

Alternate name; see *Forbes-Albright Syndrome*.

### **Galactorrhoea**

Alternate name; see *Galactorrhea*.

### **Galactosaemia**

A rare inherited disorder in which the body cannot properly process galactose, a sugar in milk. Untreated infants may develop feeding problems, jaundice, liver injury, cataracts, serious infection, or developmental problems.

---

## Galactose

A simple sugar that forms part of lactose in milk. The body normally processes it for energy, but inherited galactosemia can make milk and other galactose-containing foods dangerous for infants.

## Galactose Metabolism

The body's process for changing galactose, part of milk sugar, into energy or other useful substances. Inherited enzyme problems can block this process and cause galactosemia.

## Galactose-1-Phosphate Uridyl Transferase Deficiency

A deficiency of the GALT enzyme that causes classic galactosemia, impairing galactose processing and potentially causing liver injury, jaundice, cataracts, poor feeding, infection, and developmental problems in untreated infants.

## Galactose-1-Phosphate Uridyltransferase Blood Test

A blood test measuring an enzyme needed to process galactose. It helps screen for or diagnose classic galactosemia, especially in newborns or infants with symptoms after milk feeding.

## Galactosemia

An inherited disorder in which the body cannot properly process galactose. Classic disease can cause vomiting, jaundice, liver injury, cataracts, serious infection, and developmental problems after milk feeding.

## Galactosialidosis

A rare inherited storage disorder in which cells accumulate certain sugar-containing substances. It can cause developmental delay, muscle weakness, bone changes, kidney disease, and eye problems, with severity varying widely.

## Galactoside

A compound in which the sugar galactose is joined to another molecule, such as another sugar, a fat, or a protein. Galactosides occur in cells and in some foods.

## Galactosyltransferase

An enzyme that attaches galactose to proteins, fats, or other sugars while cells build complex molecules. These molecules support cell structure, recognition, signaling, and nerve insulation.

## Galant Reflex

A newborn reflex in which stroking one side of the back causes the infant's trunk to curve toward that side. It normally fades during early infancy; marked asymmetry or persistence can occur with nervous-system problems.

## Galeazzi Fracture

A fracture of the middle to distal third of the radial shaft associated with dislocation or subluxation of the distal radioulnar joint (DRUJ), typically requiring anatomic open reduction and internal fixation to restore forearm pronation and supination.

### Alternate Names:

Galeazzi Fracture-Dislocation; Piedmont Fracture

## Galeazzi Fracture-Dislocation

A forearm injury involving a break in the lower part of the radius and dislocation of the nearby joint between the radius and ulna. It can cause wrist pain, instability, and difficulty turning the forearm.

### Alternate Names:

Galeazzi Fracture; Reverse Monteggia Fracture

## Galeazzi Sign

A finding in an infant lying with hips and knees bent, in which one knee is lower than the other. It may suggest hip dislocation, developmental hip dysplasia, or unequal thigh-bone length.

### Alternate Names:

Allis Sign; Galeazzi Test

## Gall Bladder Disease

Any disorder of the gallbladder, including stones, inflammation, infection, polyps, abnormal movement, or cancer. Symptoms may include right-upper-abdominal pain, nausea, vomiting, fever, or jaundice.



---

## Gall-Bladder

A small pear-shaped sac beneath the liver that stores and concentrates bile. Bile is released into the small intestine to help digest fats.

## Gallavardin Phenomenon

A finding in severe aortic-valve narrowing where the heart murmur sounds harsh near the upper chest but musical near the apex. It reflects different sound frequencies transmitted through the chest.

## Gallbladder Absence

Having no gallbladder because it did not develop or was surgically removed. Bile then flows directly from the liver into the intestine; most people can digest normally, though some develop digestive symptoms.

## Gallbladder Adenomyomatosis

A benign, non-inflammatory hyperplastic alteration of the gallbladder wall characterized by excessive epithelial proliferation with smooth muscle hypertrophy and deep, pouch-like invaginations of the luminal mucosa extending into the muscularis propria (termed Rokitansky-Aschoff sinuses). Often identified on ultrasound by intramural diverticula containing cholesterol crystals that produce characteristic V-shaped comet-tail reverberation artifacts.

### Alternate Names:

Adenomyoma of Gallbladder; Diverticular Gallbladder Disease

## Gallbladder Cancer

A cancer arising in the gallbladder, sometimes associated with long-term inflammation or stones. It may cause abdominal pain, jaundice, weight loss, or blocked bile flow, but can also be found unexpectedly after surgery.

### Alternate Names:

Gallbladder Carcinoma; Biliary Malignancy

## Gallbladder Carcinoma

A cancer of the gallbladder, most often arising from its lining cells. It may be found incidentally or cause right-upper-abdominal pain, jaundice, weight loss, or bile-duct blockage; treatment depends on its stage and spread.

## Gallbladder Diseases

Conditions affecting the gallbladder, including stones, inflammation, infection, polyps, abnormal emptying, and cancer. Symptoms and treatment depend on the exact disorder and whether bile flow is blocked.

## Gallbladder Duplication

A rare birth variation in which two gallbladders develop, with separate or shared drainage ducts. It may cause no symptoms but can be associated with stones or inflammation and is important to identify before surgery.

## Gallbladder Hydrops

Severe, non-inflammatory distension and dilation of the gallbladder filled with clear, sterile, watery or mucoid transudate resulting from chronic, complete obstruction of the cystic duct (most often by an impacted gallstone in Hartmann's pouch) in the absence of acute infection.

### Alternate Names:

Mucocele of Gallbladder; Hydrops Vesicae Felleae

## Gallbladder Infection

Informal term for inflammation or infection of the gallbladder, often after a stone blocks the cystic duct. Right-upper-quadrant pain, fever, nausea, vomiting, and tenderness may occur; treatment depends on severity and may include antibiotics, drainage, or surgery.

## Gallbladder Inflammation

Inflammation of the gallbladder, most often caused by a gallstone blocking its drainage. It can cause persistent right-upper-abdominal pain, fever, nausea, and vomiting.

### Alternate Names:

Cholecystitis

## Gallbladder Mucocele

Enlargement of the gallbladder with mucus or watery fluid because its outlet is blocked, often by an impacted stone. It may cause pain or infection; drainage or removal may be used.

---

## **Gallbladder Neck**

The narrow, tapered S-shaped constriction of the gallbladder that connects the gallbladder body to the cystic duct, characterized internally by spiral mucosal folds (valves of Heister); an impacted gallstone here leads to acute cholecystitis or Mirizzi syndrome.

### **Alternate Names:**

Collum Vesicae Biliaris

## **Gallbladder Necrosis**

Death of gallbladder tissue caused by severe inflammation or inadequate blood supply. It can lead to perforation, abdominal infection, and sepsis.

## **Gallbladder Pain**

Pain from temporary blockage of the gallbladder outlet, usually by a stone. It is often steady, severe, and felt under the right ribs after meals and may include nausea or vomiting.

### **Alternate Names:**

Biliary Colic; Gallstone Pain

## **Gallbladder Perforation**

A hole through the gallbladder wall that allows bile or infected contents into the abdomen. It can cause severe inflammation, abscess, peritonitis, or sepsis.

## **Gallbladder Polyp**

A mucosal projection extending into the lumen of the gallbladder, categorized as pseudopolyps (cholesterol polyps, inflammatory polyps, adenomyomatosis) or true epithelial neoplasms (adenomas); polyps larger than 10 mm carry an increased malignancy risk warranting cholecystectomy.

## **Gallbladder Radionuclide Scan**

A nuclear medicine test that follows a tracer through the liver, gallbladder, and bile ducts. It helps assess suspected gallbladder inflammation, blockage, or abnormal bile flow.

## **Gallbladder Volvulus**

Twisting of the gallbladder around its supporting tissues and duct, cutting off blood flow. It causes

sudden severe right-upper-abdominal pain, nausea, vomiting, and abdominal irritation.

## **Gallium**

A metallic element used in some radioactive tracers, medical compounds, and technology. Its medical effects depend on the isotope or compound and how it is administered and distributed in the body.

## **Gallium Nitrate**

A gallium-containing compound formerly used in selected cancer settings to reduce high calcium levels. Its use is limited because it can damage the kidneys and has largely been replaced by other treatments.

## **Gallium Scan**

A nuclear medicine scan using radioactive gallium to locate some infections, inflammation, or tumors. It is used less often now because newer imaging tests may provide more specific or faster information.

## **Gallium-68**

A radioactive isotope used to label tracers for PET scans. Linked to a suitable targeting molecule, it can help locate some neuroendocrine tumors or prostate-cancer lesions; it is used for diagnosis, not direct treatment.

## **Gallop Rhythm**

An extra heart sound creating a three- or four-beat rhythm. It may reflect abnormal heart filling, stiff heart muscle, or heart failure and is interpreted with symptoms and other examination findings.

## **Gallstone Ileus**

A blockage of the intestine caused by a large gallstone that has entered through an abnormal connection from the gallbladder. It can cause pain, vomiting, swelling, and inability to pass stool or gas.

## **Gallstone Pancreatitis**

Sudden inflammation of the pancreas caused by a gallstone blocking the shared opening of the bile and pancreatic ducts. It usually causes severe persistent

---

upper-abdominal pain and may be associated with complications or remaining duct stones.

### **Gallstones**

Solid deposits that form in the gallbladder when bile contains too much cholesterol or bilirubin, too little bile salt, or remains in the gallbladder without emptying fully. They may cause no symptoms or block bile flow, producing severe upper-abdominal pain, inflammation, jaundice, infection, or pancreatitis.

#### **Alternate Names:**

Cholelithiasis; Biliary Calculi

### **GALT Deficiency**

Abbreviation; see *Galactose-1-Phosphate Uridyl Transferase Deficiency*.

### **Galvanocautery**

Use of heat generated by electric current to destroy selected tissue or control bleeding during a procedure. It can cause burns or damage nearby structures if not carefully controlled.

### **Gambling Disorder**

A persistent inability to control gambling despite financial, relationship, work, health, or emotional harm. The disorder can cause debt, secrecy, distress, and suicide risk and benefits from professional behavioral and mental-health support.

#### **Alternate Names:**

Pathological Gambling; Compulsive Gambling; Ludomania

### **Gamete**

A mature reproductive cell, such as a sperm or egg, containing one set of chromosomes. Two gametes combine during fertilization to form a cell with genetic material from both biological parents.

### **Gamete Intrafallopian Transfer**

An assisted-reproduction procedure in which eggs and sperm are placed into a fallopian tube so fertilization may occur inside the body. It is now used less often than in-vitro fertilization.

### **Gametogenesis**

The formation and maturation of sperm or eggs in the testes or ovaries. It includes cell division that reduces chromosome number so gametes can combine during fertilization.

### **Gamma Camera**

A device used in some nuclear-medicine scans to detect radiation from a small tracer inside the body and create pictures showing an organ's structure or activity.

### **Gamma Globulin**

The part of blood plasma containing most immunoglobulins, or antibodies, which help defend against infections and can be measured or given as immune treatment.

### **Gamma Hydroxybutyrate**

A central-nervous-system depressant that can cause drowsiness, confusion, slowed breathing, coma, or death in overdose. It is also misused recreationally and can cause dependence and withdrawal.

### **Gamma Knife**

A noninvasive radiation treatment that aims many focused beams at selected brain tumors or other lesions. It avoids an open operation but can cause swelling, bleeding, or injury to nearby brain tissue.

### **Gamma Knife Radiosurgery**

Gamma Knife is a focused radiation system that delivers multiple precisely aimed beams, mainly to selected brain tumors and vascular or other brain lesions.

### **Gamma Rays**

High-energy electromagnetic radiation produced by radioactive decay or nuclear reactions. It can damage cells and is used in controlled amounts for some cancer treatments and medical sterilization.

### **Gamma-Aminobutyric Acid**

A chemical messenger that usually reduces nerve activity in the brain and spinal cord. It helps regulate

---

sleep, anxiety, muscle activity, and seizures through two main receptor types.

**Alternate Names:**

GABA

**Gamma-Benzene Hexachloride**

An insecticide, also called lindane, that can cause nervous-system toxicity and seizures after excessive exposure. Medical use is restricted in many places because safer treatments are available.

**Gamma-Chemokines**

Signaling proteins that attract or activate immune cells through specific receptors. They help coordinate inflammation, immune surveillance, and movement of defensive cells into affected tissue.

**Gamma-Globulin**

A group of blood proteins that includes antibodies. The term may refer to antibody-containing plasma fractions or immunoglobulin preparations used to provide temporary immune protection or treatment.

**Gamma-Glutamyl Transferase Blood Test**

A blood test measuring an enzyme found mainly in the liver and bile ducts. It can support evaluation of liver or bile-duct disease and help explain a high alkaline-phosphatase result.

**Gamma-Hydroxybutyrate Toxicity**

Rapid-onset central nervous system depression from GHB or related compounds, with drowsiness, ataxia, vomiting, bradycardia, respiratory depression, coma, and sometimes agitation or delirium during recovery. There is no specific antidote; protect the airway, support ventilation, and monitor until mental status and breathing recover.

**Gamma-Interferon**

A cytokine made by activated immune cells that strengthens macrophage activity and cell-mediated defense against certain infections. It also contributes to inflammation and immune regulation.

**Gamma-Ray Therapy**

Treatment that uses controlled gamma radiation to damage the DNA of selected abnormal cells. It can treat some cancers, but nearby healthy tissue may also be affected and side effects depend on the treated area.

**Gammaherpesvirinae**

A subgroup of herpesviruses that establish long-lasting infection in lymphoid or related cells. Some members, including Epstein-Barr virus, can cause human disease and contribute to certain cancers.

**Gamna-Gandy Bodies**

Small, firm, yellow-brown or reddish-brown siderofibrotic nodules in the splenic red pulp containing deposits of hemosiderin, iron, and calcium encrusting connective tissue and elastic fibers, resulting from organized focal hemorrhages in chronic congestive splenomegaly and sickle cell disease.

**Alternate Names:**

Gamna-Gandy Nodules; Siderofibrotic Bodies

**GAN**

Abbreviation; see *Giant Axonal Neuropathy*.

**Ganciclovir**

An antiviral medicine that interferes with viral DNA production and is used mainly for serious cytomegalovirus infection. It can suppress bone marrow, affect the kidneys, and cause other serious toxicity.

**Ganglia**

Groups of nerve-cell bodies outside the brain and spinal cord. They relay and organize signals between the central nervous system and the rest of the body.

**Ganglioglioma**

A usually slow-growing tumor containing nerve cells and supporting brain cells. It often causes seizures and may occur in the brain or spinal cord; treatment commonly involves surgical removal.

---

## Ganglion

A cluster of nerve-cell bodies outside the brain and spinal cord. It acts as a relay point for signals between the central nervous system and body tissues.

## Ganglion Cell

A nerve cell located in a ganglion. Retinal ganglion cells collect visual signals and send them through the optic nerve to the brain.

## Ganglion Cyst

A noncancerous, fluid-filled lump near a joint or tendon, most often at the wrist or hand. It may be painless or cause pressure, pain, numbness, or limited movement.

### Alternate Names:

Synovial Cyst

## Ganglionectomy

Surgical removal of a ganglion, a cluster of nerve-cell bodies, or occasionally a ganglion cyst. The purpose, benefits, and risks depend on the exact structure and body site being treated.

## Ganglioneuroblastoma

A tumor made of both mature and immature nerve-related cells, usually in the adrenal or tissue beside the spine. It mainly affects children and may cause a mass, pain, high blood pressure, or other symptoms.

## Ganglioneuroma

A usually noncancerous tumor arising from nerve tissue of the autonomic nervous system. It often causes no symptoms but may produce a mass, pain, or hormone-related effects depending on its location.

## Ganglioneuromatosis

A condition in which nerve-related ganglion tissue grows diffusely or in multiple areas. Effects depend on the organs involved and may include masses, pain, bowel symptoms, or hormone-related changes.

## Ganglionic Blockers

Medicines that interrupt nerve signals at autonomic ganglia, affecting blood pressure, heart rate, diges-

tion, and other automatic functions. They are rarely used because of widespread side effects.

## Ganglions

Alternate plural form; see *Ganglion*.

## Ganglioside

A fat-containing molecule in cell membranes, especially nerve cells. Gangliosides help cells communicate and recognize one another; abnormal accumulation occurs in several inherited storage disorders.

## Gangliosidosis

A group of inherited disorders in which gangliosides accumulate inside cells because a breakdown enzyme is missing. The buildup often damages the nervous system and may also affect the eyes, bones, organs, and development.

## Gangliosidosis GM2 Type 2

Alternate name; see *Sandhoff Disease*.

## Gangrene

Death of body tissue caused by severely reduced blood flow, infection, injury, or a combination of these factors. It may affect skin, muscle, or an internal organ. The area may become discolored, cold, swollen, painful, numb, or foul-smelling; fever, confusion, or feeling very ill may occur with spreading infection.

## Ganser Syndrome

A rare dissociative or psychiatric syndrome characterized by approximate answers, confusion, altered awareness, and sometimes hallucinations or neurologic-like symptoms. It may occur with severe stress, trauma, other mental disorders, or medical and substance-related conditions.

## Gap Junction

A tiny connection between neighboring cells that allows ions and small molecules to pass directly between them. Gap junctions help coordinate heartbeats, nerve activity, and tissue growth.

---

## Gardasil

A brand of vaccine that protects against infection with selected human papillomavirus (HPV) types. It helps prevent HPV-related cancers and genital warts and does not treat an existing HPV infection.

## Gardner Syndrome

An inherited condition caused by an APC gene change, producing many colon polyps and increasing colorectal-cancer risk. It may also cause bone growths, dental abnormalities, skin cysts, and desmoid tumors.

### Alternate Names:

Bone Tumor-Epidermoid Cyst-Polyposis; Familial Adenomatous Polyposis With Osteomas

## Gardnerella

A group of bacteria that may live in the genital tract. Overgrowth of Gardnerella, especially *G. vaginalis*, can contribute to bacterial vaginosis with discharge, odor, and altered vaginal bacteria.

## Gardnerella Vaginalis

A bacterium associated with bacterial vaginosis, in which normal vaginal bacteria become imbalanced. It may be present without illness; when symptomatic, discharge, odor, irritation, or burning can occur.

## Gargle

To rinse the mouth and throat with liquid while making a bubbling sound. Gargling may soothe irritation or help deliver a prescribed treatment.

## Garré's Osteomyelitis

A chronic inflammatory reaction in the jawbone, often near an infected tooth, causing firm swelling. Dental infection, persistent swelling, or pain may be present.

## Gartner Duct

A remnant of an embryonic duct in the female reproductive tract. It can persist harmlessly or form a cyst along the side wall of the vagina or broad ligament.

## Gartner Duct Cyst

A usually harmless fluid-filled cyst arising from an embryonic duct remnant along the side wall of the vagina. It often causes no symptoms, but a large cyst may cause pressure, pain, or difficulty with intercourse.

## Gas

A state of matter without a fixed shape or volume. In medicine, gas may mean air or another gas in the digestive tract, blood, tissues, or a body cavity.

## Gas (Flatus)

Alternate name; see *Gas* – *Flatulence*.

### Alternate Names:

Flatulence; Flatus Expulsion

## Gas And Gas Pains

Fullness, bloating, cramps, or discomfort from swallowed air or gas made by intestinal bacteria. Foods, constipation, food intolerance, and digestive disorders can contribute.

## Gas Chromatography

A laboratory method that separates vaporized chemicals so they can be identified or measured. It is often combined with mass spectrometry to test medicines, poisons, pollutants, or biological samples.

## Gas Exchange

Movement of oxygen from inhaled air into blood and carbon dioxide from blood into exhaled air. Oxygen then moves to tissues while carbon dioxide returns to the lungs for removal.

## Gas Gangrene

A life-threatening infection in which bacteria produce toxins and gas inside damaged tissue, rapidly destroying muscle. Severe pain, swelling, blisters, discoloration, or a crackling feeling may occur.

## Gas Gangrene (Clostridial Myonecrosis)

A rapidly progressive, often fatal infection in which *Clostridium* species destroy muscle and produce



---

toxins and gas in damaged tissue. Severe pain, swelling, skin discoloration, systemic toxicity, and shock require emergency surgery, antibiotics, and intensive care.

### **Gas Poisoning**

Illness caused by inhaling a toxic gas, with effects determined by the agent: carbon monoxide and cyanide cause cellular or functional hypoxia, while chlorine, ammonia, and similar irritants injure the airways and lungs. Headache, confusion, cough, bronchospasm, pulmonary edema, seizures, or collapse may occur; identify the gas and provide immediate fresh air, oxygen, and respiratory support.

### **Gas Scavenger**

A substance or device that removes unwanted gases from a fluid, container, or breathing system. In healthcare it may help prevent toxic exposure or maintain safe gas delivery.

### **Gas Transport**

Movement of oxygen from the lungs to body tissues and carbon dioxide back to the lungs through the blood. Hemoglobin carries most oxygen and part of the carbon dioxide.

### **Gas – Flatulence**

Passage of gas from the digestive tract through the anus, caused by swallowed air and bacterial digestion of food. It is normal, but excess gas with bloating or pain may reflect diet, constipation, or digestive disease.

### **Gaskin Maneuver**

A maneuver for shoulder dystocia in which the birthing person moves onto hands and knees. The position can change pelvic dimensions and help release the baby's shoulder.

#### **Alternate Names:**

All-Fours Position; Hands-and-Knees Posture

### **Gasoline Poisoning**

Harm from swallowing, inhaling, or getting gasoline on the skin. It can cause nausea, dizziness, drowsiness, seizures, abnormal heart rhythms, or chemical

lung injury, especially if vomited and inhaled.

#### **Alternate Names:**

Petrol Poisoning; Hydrocarbon Toxicity

### **Gasserian Ganglion**

A cluster of nerve cells near the base of the skull that carries sensation from the face and controls some chewing muscles. It is involved in trigeminal neuralgia and other facial-nerve disorders.

### **Gastralgia**

Pain in the upper abdomen or stomach area. Causes include indigestion, reflux, ulcers, gallbladder disease, pancreatic inflammation, infection, or other conditions; severe pain may occur with bleeding, vomiting, fever, or weight loss.

### **Gastrectomy**

Surgery removing part or all of the stomach, usually for cancer or severe disease. Food passage is reconstructed, and possible effects include early fullness, weight loss, nutritional deficiencies, reflux, or dumping symptoms.

### **Gastric Acid**

Strong acid made by the stomach lining that helps break down food, activate digestive enzymes, and kill many swallowed microbes. Excess can contribute to reflux or ulcers; too little can impair digestion and nutrient absorption.

### **Gastric Adenocarcinoma**

The most common stomach cancer, arising from cells lining the stomach. It may cause indigestion, pain, early fullness, vomiting, bleeding, or weight loss; tissue examination and staging identify its type and extent.

### **Gastric Analysis**

Examination of stomach contents for volume, acidity, or selected substances. It is used only in specific situations because endoscopy, imaging, and blood tests are more useful for many stomach disorders.

---

### Gastric Antral Vascular Ectasia

Enlarged fragile blood vessels in the lower stomach that can cause slow internal bleeding and iron-deficiency anemia. It may appear as red stripes or spots during endoscopy and is associated with some liver and autoimmune diseases.

#### Alternate Names:

GAVE; Watermelon Stomach

### Gastric Antrum

The distal, non-acid-secreting muscular portion of the stomach located between the incisura angularis and the pylorus, containing G cells that secrete gastrin and D cells that secrete somatostatin, serving as the site of mechanical food grinding and trituration.

### Gastric Balloon

A temporary balloon placed inside the stomach to create fullness and support weight loss. Nausea, vomiting, reflux, deflation, blockage, bleeding, or perforation can occur.

### Gastric Bezoar

A mass of material that cannot be digested and collects in the stomach. It may contain plant fiber, hair, or medicines and can cause fullness, pain, vomiting, blockage, or poor nutrition.

### Gastric Biopsy

Removal of a small sample of stomach lining, usually during endoscopy, for laboratory examination. It can help assess inflammation, infection, precancerous changes, or cancer.

### Gastric Body

The main middle part of the stomach between the fundus and the antrum. It stores and mixes food and contains glands that release acid and digestive substances.

### Gastric Bypass

An operation that makes the stomach smaller and connects it to a lower part of the small intestine. It limits food intake and nutrient absorption and can improve severe obesity and related diseases.

### Gastric Bypass Surgery

Surgery that creates a small stomach pouch and reroutes food to the small intestine. It produces substantial weight loss and may improve diabetes, high blood pressure, sleep apnea, and other obesity-related conditions.

#### Alternate Names:

Roux-en-Y Gastric Bypass; Bariatric Bypass

### Gastric Cancer

See *Stomach Cancer*.

### Gastric Carcinoid Tumor

A neuroendocrine tumor arising in the stomach, often from cells that make digestive hormones. Some are linked to chronic gastritis and high gastrin levels; behavior ranges from slow-growing to aggressive.

### Gastric Cardia

The anatomical transition zone of the stomach immediately surrounding the gastroesophageal junction where the tubular esophagus empties into the stomach, lined by simple columnar mucous-secreting glands without parietal or chief cells.

### Gastric Culture

A laboratory test that grows bacteria from a stomach sample, usually collected during endoscopy. It can identify *Helicobacter pylori* and help select antibiotics, although other tests are more commonly used.

#### Alternate Names:

*Helicobacter Pylori* Culture; Gastric Biopsy Culture

### Gastric Dysplasia

Abnormal cells in the stomach lining that have some features of cancer but have not invaded deeper tissue. Low- or high-grade changes may increase the risk of stomach cancer.

### Gastric Emptying

The passage of food and liquid from the stomach into the small intestine through coordinated stomach contractions and opening of the pylorus. Abnormally slow or rapid emptying can cause digestive

---

symptoms.

### **Gastric Emptying Study**

A test that measures how quickly food leaves the stomach. The patient eats a meal containing a small tracer, and a scanner records the remaining food over several hours to detect unusually slow or rapid emptying.

### **Gastric Fistula**

An abnormal passage between the stomach and another organ or the skin. Stomach contents may leak through it, causing drainage, infection, skin irritation, dehydration, or poor nutrition.

### **Gastric Fundus**

The upper, rounded part of the stomach, located above the main stomach body and near the diaphragm. It can expand to store food and contains glands that produce digestive secretions.

### **Gastric Glands**

Glands in the stomach lining that make acid, digestive enzymes, mucus, intrinsic factor, and hormones.

### **Gastric Hemorrhage**

Bleeding from the stomach, commonly caused by an ulcer, a tumor, or blood-vessel damage. It may cause vomiting blood, black stools, dizziness, anemia, or shock. It is one possible source of upper gastrointestinal bleeding, which can also come from the esophagus or duodenum.

#### **Alternate Names:**

Gastric Bleeding

### **Gastric Intestinal Metaplasia**

See *Intestinal Metaplasia*.

### **Gastric Juice**

Fluid made by the stomach containing acid, digestive enzymes, mucus, water, salts, and intrinsic factor. It helps break down food, protect the stomach lining, and absorb vitamin B12 later in digestion.

### **Gastric Lavage**

A procedure that flushes the stomach through a tube passed through the mouth or nose. It is rarely used because it can cause aspiration or injury and is reserved for selected emergencies.

### **Gastric Leiomyosarcoma**

A rare cancer of smooth muscle in the stomach wall. It may cause pain, bleeding, vomiting, or a mass and is evaluated with imaging and biopsy; treatment usually involves surgery when possible.

### **Gastric Lymphoma**

A lymphoma that starts in the stomach. Some types, especially early-stage mucosa-associated lymphoid tissue (MALT) lymphoma, are linked to *Helicobacter pylori* and may improve after the infection is treated; other types may need cancer treatment.

### **Gastric Malignancy**

See *Stomach Cancer*.

### **Gastric Motility**

The coordinated muscle activity that mixes stomach contents and moves them toward the small intestine.

### **Gastric Mucosa**

The moist inner lining of the stomach. It contains glands that produce acid, digestive enzymes, mucus, and intrinsic factor and is protected by a mucus barrier; inflammation or injury can cause pain or bleeding.

### **Gastric Muscularis**

The muscle layers in the stomach wall that mix food and move it toward the pylorus. The stomach has an extra inner muscle layer that helps with grinding.

### **Gastric Necrosis**

Death of stomach tissue caused by severely reduced blood flow, corrosive injury, extreme stretching, infection, or another serious insult. It can lead to perforation, severe infection, and shock.

---

## Gastric Obstruction

Partial or complete blockage that prevents food and liquid from leaving the stomach normally. It may cause repeated vomiting, upper-abdominal pain, fullness, swelling, dehydration, and weight loss.

## Gastric Outlet Obstruction

Blockage where the stomach empties into the small intestine, usually at the pylorus or duodenum. Causes include ulcer scarring, tumors, inflammation, and infantile pyloric stenosis; vomiting can cause dehydration.

## Gastric Perforation

A hole through the stomach wall that allows stomach contents to leak into the abdomen. It can cause severe inflammation, infection, and shock.

## Gastric Pneumatosis

Gas within the stomach wall. It may result from pressure-related injury and be relatively mild, or from severe infection, which can destroy tissue and cause sepsis.

## Gastric Polyps

Growths projecting from the stomach lining. Most are harmless, but some types are linked to chronic inflammation, inherited conditions, or long-term acid-suppressing treatment and may contain precancerous changes.

## Gastric Rugae

Folds in the inner lining of the empty stomach that flatten as the stomach fills. They allow the stomach to expand and increase the lining's surface area.

## Gastric Secretion

The release of acid, enzymes, mucus, intrinsic factor, and hormones by the stomach lining.

## Gastric Serosa

The thin outer covering of the stomach, formed by the visceral peritoneum. It reduces friction as the stomach moves inside the abdomen.

## Gastric Stenosis

Abnormal narrowing of the stomach or its outlet that slows or blocks emptying. It may result from birth defects, ulcer scarring, inflammation, tumors, or thickening of the outlet muscle.

## Gastric Submucosa

The connective-tissue layer beneath the stomach lining. It contains blood vessels, lymph vessels, nerves, and supporting tissue.

## Gastric Suction

Removal of stomach contents through a tube passed through the nose or mouth. It may relieve severe distension or vomiting from obstruction and is used only when clinically indicated because complications can occur.

### Alternate Names:

Nasogastric Aspiration; Stomach Pumping

## Gastric Surgery

An operation on the stomach to remove disease, repair damage, reduce its size, or bypass a blockage. The exact procedure and risks depend on the condition and may include bleeding, infection, leakage, or altered digestion.

## Gastric Tissue Biopsy And Culture

Removal of small stomach-lining samples during endoscopy for microscopic examination and, when needed, laboratory testing for infection. It can investigate inflammation, *Helicobacter pylori*, precancerous changes, or tumors.

### Alternate Names:

Endoscopic Gastric Biopsy; Mucosal Culture

## Gastric Ulcer

An open sore in the stomach lining, commonly caused by *Helicobacter pylori* infection or anti-inflammatory medicines. It may cause burning upper-abdominal pain, bleeding, anemia, or perforation.

## Gastric Varices

Enlarged, fragile veins in the stomach caused by

---

increased pressure in the portal circulation, often from advanced liver disease. Rupture can cause sudden, life-threatening vomiting of blood.

### **Gastric Volvulus**

Twisting of the stomach that can block food passage and cut off blood supply. It may cause sudden severe upper-abdominal pain, retching without vomiting, bloating, and inability to pass a tube.

### **Gastric Wall**

The layered wall of the stomach, made up of the mucosa, submucosa, muscle layers, and outer covering. It allows the stomach to store, mix, and move food.

### **Gastrin**

A hormone made mainly in the lower stomach and upper small intestine. It stimulates stomach acid production and helps maintain the stomach lining; levels rise after eating and may become excessive in certain diseases.

### **Gastrin Blood Test**

A blood test measuring the hormone gastrin. It helps investigate excessive stomach acid, recurrent ulcers, or loss of acid-producing cells. Acid-suppressing medicines and fasting can significantly affect the result.

#### **Alternate Names:**

Serum Gastrin Level; Fasting Gastrin Assay

### **Gastrin Secretion**

Release of the hormone gastrin, mainly from cells in the lower stomach after food enters. Gastrin increases stomach acid and digestive-enzyme release and supports growth and function of the stomach lining.

### **Gastrin-Releasing Peptide**

A regulatory neuropeptide released by postganglionic parasympathetic vagal nerve endings in the gastric antrum that binds to receptors on G cells, stimulating gastrin release and subsequent gastric acid secretion.

#### **Alternate Names:**

GRP; Mammalian Bombesin

### **Gastrinoma**

A tumor that makes too much gastrin, a hormone that causes the stomach to produce acid. It usually starts in the pancreas or duodenum and may cause Zollinger–Ellison syndrome, repeated peptic ulcers, diarrhea, and acid reflux. Some cases are linked to MEN1 and may spread to the liver or nearby lymph nodes.

### **Gastritis**

Inflammation of the stomach lining. It may begin suddenly (acute) or last a long time (chronic), and may cause no symptoms or upper-abdominal pain, nausea, vomiting, or bleeding. *H. pylori* infection and autoimmune disease are important causes. Some medicines, alcohol, and severe illness can also injure the lining; when injury occurs without prominent inflammation, it is called gastropathy.

#### **Alternate Names:**

Type A Gastritis; Autoimmune Metaplastic Atrophic Gastritis

### **Gastrocnemius**

The large muscle at the back of the calf. It helps point the foot downward and bend the knee during standing, walking, running, and jumping; injury can cause calf pain, swelling, or weakness.

### **Gastrocnemius Muscle**

The large superficial calf muscle that joins the Achilles tendon. It points the foot downward and helps bend the knee and push the body forward during walking, running, and jumping.

### **Gastroduodenal Artery**

A major arterial branch arising from the common hepatic artery, descending posterior to the first part of the duodenum before dividing into the right gastro-omental and superior pancreaticoduodenal arteries; erosion by a posterior duodenal ulcer produces massive upper gastrointestinal hemorrhage.

#### **Alternate Names:**

GDA

---

### Gastroduodenal Crohn Disease

Crohn disease affecting the stomach or first part of the small intestine. It may cause upper-abdominal pain, nausea, vomiting, weight loss, ulcers, narrowing, or blockage; treatment varies with symptoms and disease severity.

### Gastroduodenostomy

An operation creating a new opening between the stomach and duodenum so food can pass directly into the small intestine. It may bypass a blockage or restore passage after part of the stomach is removed.

### Gastroenteritis

Infection or inflammation of the stomach and intestines, commonly caused by viruses and sometimes by bacteria, parasites, or toxins. It usually causes diarrhea, vomiting, cramps, and fever and can lead to dehydration.

#### Alternate Names:

Acute Gastroenteritis

### Gastroenteritis Bacterial

Inflammation of the stomach or intestines caused by bacteria, usually after contaminated food or water. It can cause diarrhea, cramps, fever, vomiting, and dehydration; blood in stool or severe illness may occur.

### Gastroenteritis Viral

Alternate name; see *Viral gastroenteritis*.

### Gastroenterologist

A physician specializing in disorders of the digestive tract, liver, gallbladder, bile ducts, and pancreas. They diagnose and treat digestive problems using medicines, endoscopy, procedures, lifestyle care, and surgery referrals.

### Gastroenterology

The medical specialty concerned with the digestive tract, liver, gallbladder, bile ducts, and pancreas. Gastroenterologists diagnose and treat these conditions with medicines, endoscopy, procedures, and referrals for surgery.

### Gastroenterostomy

Surgical creation of an opening between the stomach and a segment of the small intestine, usually the jejunum, to bypass or replace a blocked or diseased gastric outlet.

### Gastroepiploic Artery

The gastroepiploic arteries run along the greater curve of the stomach and supply blood to the stomach and nearby omentum.

### Gastroesophageal

Relating to both the stomach and esophagus, especially their joining point or the backward movement of stomach contents into the esophagus.

### Gastroesophageal Junction Adenocarcinoma

A gland-forming cancer arising where the esophagus joins the stomach. It may cause difficulty swallowing, reflux, weight loss, anemia, or bleeding; endoscopy, biopsy, and scans determine treatment and extent.

### Gastroesophageal Reflux

Backward flow of stomach contents into the esophagus. It may cause heartburn, sour fluid in the throat, cough, chest discomfort, or no symptoms. Spitting up is common in healthy infants and usually resolves with age; reflux is called disease when it causes troublesome symptoms or complications.

### Gastroesophageal Reflux Disease

Reflux that causes troublesome symptoms or complications, such as persistent heartburn, pain, feeding difficulty, or injury to the esophagus. In infants, ordinary spit-up without distress or poor growth is common and does not by itself mean reflux disease.

### Gastroesophageal Reflux Disease (GERD)

Alternate name; see *Gastroesophageal Reflux Disease*.

#### Alternate Names:

GERD; Acid Reflux Disease



---

### **Gastroesophageal Z-Line**

The boundary where the lining of the lower food pipe changes to the lining of the stomach. Its position and appearance can help identify conditions such as Barrett esophagus.

#### **Alternate Names:**

Squamocolumnar Junction; SCJ; Ora Serrata of the Esophagus

### **Gastrogastic Intussusception**

A rare condition in which one part of the stomach folds into an adjoining part, sometimes because of a polyp or tumor. It may cause pain, vomiting, blockage, or bleeding.

### **Gastrografen Study**

An X-ray study using a water-soluble contrast liquid to show the digestive tract. It may be chosen when a leak or perforation is possible, because the contrast is safer outside the tract than barium.

### **Gastrointestinal**

Relating to the stomach and intestines, or to the complete digestive tract from the mouth to the anus. These organs digest food, absorb nutrients and water, move contents forward, and remove waste.

### **Gastrointestinal Agents**

Medicines or other substances that act on the digestive tract to relieve or treat problems such as nausea, vomiting, diarrhea, constipation, ulcers, reflux, or intestinal spasms. Their effects and risks depend on the specific agent.

### **Gastrointestinal Amyloidosis**

Buildup of abnormal protein deposits in the digestive tract. It can impair movement, absorption, or bleeding control and may occur with disease affecting other organs such as the heart, kidneys, or nerves.

### **Gastrointestinal Bleeding**

Bleeding anywhere in the digestive tract. It may be acute or chronic and may come from the upper digestive tract, small intestine, or lower digestive

tract. It can cause vomiting blood, black stools, or red blood in the stool.

### **Gastrointestinal Bleeding Scan**

A nuclear-medicine scan that uses a small radioactive tracer in the blood to find active bleeding in the digestive tract. It can help localize bleeding when endoscopy does not identify the source.

### **Gastrointestinal Cancer**

Cancer arising in the digestive tract or related organs, including the esophagus, stomach, intestine, rectum, liver, pancreas, and biliary tract. Symptoms and treatment depend on the organ, tumor type, and stage.

### **Gastrointestinal Decontamination**

Methods used after some poisonings to reduce absorption, such as activated charcoal or whole-bowel irrigation. The choice depends on the substance, timing, symptoms, and risk of aspiration; stomach washing is rarely used.

### **Gastrointestinal Disease**

Any disorder affecting the digestive tract or related organs. Examples include reflux, ulcers, inflammatory bowel disease, infection, malabsorption, liver disease, and cancer.

### **Gastrointestinal Disease and Pregnancy**

Pregnancy can alter gastrointestinal motility and symptoms, while disorders such as inflammatory bowel disease, reflux, gallbladder disease, liver disease, or bowel obstruction may affect nutrition and pregnancy care. Severe pain, persistent vomiting, bleeding, fever, or dehydration requires assessment.

### **Gastrointestinal Disorder**

A disease affecting any part of the digestive tract, from the esophagus to the anus. It may cause pain, nausea, vomiting, diarrhea, constipation, bleeding, blockage, or poor nutrient absorption, depending on the affected structure.

---

### **Gastrointestinal Fistula**

An abnormal connection between the digestive tract and another organ, the skin, or another section of bowel. Digestive fluid may leak through it, causing infection, dehydration, skin damage, or poor nutrition.

### **Gastrointestinal Foreign Bodies**

Objects swallowed or inserted into the gastrointestinal tract that may pass spontaneously or become lodged, cause obstruction, perforation, bleeding, or poisoning. Evaluation and removal depend on the object, location, symptoms, and risk of injury.

### **Gastrointestinal Gas**

Air or other gas within the stomach or intestines, usually from swallowed air or bacterial digestion of food. Excess gas may cause belching, bloating, cramps, or passing gas and can increase with constipation or food intolerance.

### **Gastrointestinal Health**

The normal structure and function of the digestive system, including digestion, absorption, bowel movement, and a balanced relationship with gut microorganisms. Problems may include bleeding, persistent pain, vomiting, or unexplained weight loss.

### **Gastrointestinal Hemorrhage**

Bleeding from any part of the digestive tract, from the esophagus to the anus. It may cause vomiting blood, black stools, red blood in stool, anemia, dizziness, or shock.

### **Gastrointestinal Hormone**

A chemical messenger made by cells in the digestive tract that helps regulate stomach acid, intestinal movement, digestion, appetite, gallbladder activity, or blood-glucose control.

### **Gastrointestinal Motility Disorder**

Abnormal movement of food, liquid, or stool through the digestive tract. It can cause difficulty swallowing, nausea, vomiting, bloating, abdominal pain, constipation, or diarrhea; causes may involve

digestive muscles, nerves, hormones, or a blockage.

### **Gastrointestinal Neoplasm**

An abnormal growth arising in the digestive tract. It may be a harmless growth, a precancerous change, or cancer; symptoms and treatment depend on its tissue, location, and behavior.

### **Gastrointestinal Obstruction**

Partial or complete blockage of food, fluid, or stool through the digestive tract. It can cause cramping pain, vomiting, swelling, constipation, dehydration, loss of blood supply, perforation, or sepsis.

### **Gastrointestinal Pain**

Pain arising from the digestive tract or related organs. Causes include infection, ulcers, inflammation, blockage, gallstones, and diseases of the liver or pancreas. It may be severe, persistent, or worsening.

### **Gastrointestinal Perforation**

A hole through the wall of the digestive tract that allows its contents to leak into surrounding tissue. It can cause severe abdominal pain, peritonitis, sepsis, and shock.

### **Gastrointestinal Stromal Tumor**

A tumor that begins in supporting cells in the wall of the digestive tract, most often in the stomach or small intestine. Its behavior varies; treatment may include surgery or medicines designed to block specific tumor signals.

### **Gastrointestinal Stromal Tumors**

Alternate name; see *Gastrointestinal Stromal Tumors (GISTs)*.

### **Gastrointestinal Stromal Tumors (GISTs)**

Mesenchymal tumors arising most often in the wall of the stomach or small intestine and commonly driven by KIT or PDGFRA pathway abnormalities. They may cause bleeding, pain, obstruction, or an incidental mass; risk assessment uses size, site, mitotic activity, and rupture.

---

## Gastrointestinal System

The connected organs that move and digest food, absorb nutrients and water, and remove solid waste. It includes the mouth, esophagus, stomach, intestines, rectum, and anus, with help from the liver, gallbladder, and pancreas.

## Gastrointestinal Tract

The continuous digestive passage from the mouth through the esophagus, stomach, small and large intestines, rectum, and anus. It moves food, digests it, absorbs nutrients and water, and removes waste.

## Gastrointestinal Villi

Tiny finger-like projections lining the small intestine that greatly increase the area available to absorb nutrients and water. Each contains small blood vessels and a lymph vessel that carry absorbed substances away.

## Gastrojejunostomy

An operation creating a new connection between the stomach and jejunum, the middle part of the small intestine. It may bypass a blocked stomach outlet or replace a damaged section of digestive tract.

## Gastromalacia

Abnormal softening of the stomach wall. It is a rare descriptive finding that may result from severe inflammation, poor blood supply, or tissue injury and can increase the risk of weakness or perforation.

## Gastroparesis

A disorder in which the stomach empties too slowly, or sometimes stops emptying, even though there is no physical blockage. The stomach muscles or the nerves that control them do not move food normally. It may cause early fullness, nausea, vomiting, bloating, belching, upper-abdominal pain, heartburn, and poor appetite. Diabetes is the most common known cause; other causes include stomach surgery, nerve or muscle disorders, some autoimmune or viral illnesses, although the cause is often unknown. Some medicines can slow stomach emptying or cause similar symptoms without causing gastroparesis itself. Severe disease can lead to dehydration, poor

nutrition, and difficult-to-control blood glucose.

### Alternate Names:

Delayed Gastric Emptying; Gastric Hypomotility

## Gastropathy

Damage or disease of the stomach lining without the prominent inflammation seen in gastritis. Causes include medicines, alcohol, poor blood flow, bile reflux, and severe illness; symptoms may include pain, nausea, or bleeding.

## Gastroplasty

Surgery that changes the size or shape of the stomach, usually to support weight loss or repair a defect. Possible complications include bleeding, infection, leakage, reflux, and nutritional deficiencies.

## Gastroptosis

Downward displacement of the stomach from its usual position. It may be seen on imaging and may or may not cause symptoms; the finding alone does not establish the cause of digestive complaints.

### Alternate Names:

Gastric Prolapse

## Gastroschisis

A birth defect in which the intestine, and sometimes other organs, protrudes through an opening beside the umbilicus without a protective covering. The exposed bowel may be swollen or damaged.

## Gastroschisis Repair

Surgical repair of gastroschisis, a congenital abdominal wall defect where intestines protrude through an opening near the umbilicus. The bowel is returned to the abdomen and the defect is closed, sometimes in stages using a temporary covering.

### Alternate Names:

Abdominal Wall Defect Repair; Congenital Evisceration Closure

## Gastroscope

A thin, flexible instrument with a light and camera used to view the esophagus, stomach, and first part of the small intestine. It can also pass tools for

---

biopsy or treatment.

### Gastroscopy

Examination of the esophagus, stomach, and duodenum using a flexible camera tube passed through the mouth. It can identify bleeding, ulcers, inflammation, narrowing, or tumors and allows biopsies or selected treatments.

### Gastrospenic Fistulization

An abnormal connection between the stomach and spleen, usually caused by a severe stomach ulcer or a splenic abscess. It can cause upper-abdominal pain, digestive-tract bleeding, severe infection, and sepsis.

### Gastrostomy

A surgically created opening through the abdominal wall into the stomach. A tube placed through it can provide food, fluids, or medicines or remove stomach contents when normal feeding is unsafe or impossible.

### Gastrostomy Tube

A tube passed through the abdominal wall into the stomach to provide food, fluids, or medicines when swallowing is unsafe or inadequate. Problems can include skin infection, leakage, blockage, bleeding, and movement of the tube.

### Gastrostomy Tube Insertion

A surgical, laparoscopic, or endoscopic procedure in which a feeding tube is placed directly through the abdominal wall into the stomach for long-term enteral nutrition.

#### Alternate Names:

G-Tube Placement; Percutaneous Endoscopic Gastrostomy; PEG Tube Insertion

### Gastrula

An early embryo stage formed after gastrulation, when cell layers reorganize into ectoderm, mesoderm, and endoderm that give rise to body tissues.

### Gastrulation

An early stage of embryo development in which cells move and reorganize into three layers that later form all major tissues and organs.

### Gated Blood Pool Scan

A radionuclide cardiac study, also called radionuclide ventriculography or MUGA, that uses ECG-gated images of labeled blood to measure ventricular ejection fraction and assess ventricular wall motion. It is used selectively when quantitative ventricular function is needed.

### Gatifloxacin

Gatifloxacin is a fluoroquinolone antibiotic that inhibits bacterial DNA gyrase and topoisomerase IV; systemic use is restricted in many settings because of dysglycaemia and other class toxicities, while ophthalmic use remains available in some regions.

### Gaucher Disease

An autosomal recessive lysosomal storage disorder caused by reduced activity of the enzyme glucocerebrosidase, usually because of disease-causing variants in the *GBA1* gene. Glucosylceramide accumulates in macrophages and other cells, affecting the spleen, liver, bone marrow, bones, lungs, and sometimes the nervous system. It can cause enlargement of the spleen or liver, anemia, low platelets, easy bruising, bone pain or fractures, and neurologic problems. Type 1 is usually non-neuronopathic, while types 2 and 3 involve the nervous system.

### Gaucher Splenomegaly

Alternate name; see *Gaucher disease*.

### Gaucher's Disease

Alternate spelling; see *Gaucher Disease*.

### Gaucher-Schlagenhauser

Alternate name; see *Gaucher Disease*.

### Gaultheria Oil

Oil of wintergreen, an essential oil containing concentrated methyl salicylate. It may be used in small

---

amounts in topical products, but swallowing even a small amount can cause serious salicylate poisoning, especially in children.

### **Gauss**

A unit of magnetic flux density equal to  $10^{-4}$  tesla, used in physics, imaging, and descriptions of magnetic fields.

### **Gauze**

A woven or nonwoven material used to clean, dress, pack, or absorb fluid from a wound. It may be sterile or nonsterile and comes in several forms.

### **Gavage**

Giving liquid food or medicine through a tube into the stomach. It is used when swallowing is unsafe or oral intake is inadequate, including for premature infants who cannot coordinate sucking, swallowing, and breathing.

### **Gayet-Wernicke Syndrome**

Alternate name; see *Wernicke Encephalopathy*.

### **Gaze Palsy**

An inability to move both eyes conjugately in a particular direction (horizontal, vertical, or upgaze), caused by lesions involving the cortical frontal eye fields, the paramedian pontine reticular formation (PPRF), or the midbrain rostral interstitial nucleus of the medial longitudinal fasciculus (riMLF).

#### **Alternate Names:**

Conjugate Gaze Palsy; Ocular Conjugate Palsy

### **GBM**

An abbreviation for glioblastoma, an aggressive cancer that starts in supporting cells of the brain. It can grow quickly and cause headache, seizures, weakness, speech or memory problems, or personality changes. Treatment may include surgery, radiation, and medicines.

#### **Alternate Names:**

Glioblastoma Multiforme; Grade IV Astrocytoma

### **Gefitinib**

An oral epidermal growth factor receptor tyrosine-kinase inhibitor used for selected advanced non-small-cell lung cancers with activating EGFR mutations.

### **Gegenhalten**

A velocity-independent, variable increase in resistance to passive limb movement in which the patient involuntarily opposes the examiner's manipulation with a force proportional to that applied by the examiner, commonly observed in diffuse frontal lobe disease, dementia, and metabolic encephalopathy.

#### **Alternate Names:**

Paratonia; Oppositional Paratonia

### **Geiger Counter**

An instrument that detects and measures ionizing radiation. It can identify radioactive contamination or exposure, but its readings do not by themselves show how much biological harm the radiation caused.

### **Gel**

A semisolid preparation in which liquid is held in a soft network. Gels may be applied to the skin, eyes, mouth, or other areas to soothe tissue, deliver medicine, or form a protective layer.

### **Gel Electrophoresis**

Gel electrophoresis separates DNA, RNA, or proteins by moving them through a gel in an electric field, usually according to size and charge.

### **Gelastic Seizure**

A rare seizure that causes sudden, repeated bursts of laughter without feeling amused. It is often linked to a small growth called a hypothalamic hamartoma, but other brain conditions can also cause it.

### **Gelatin**

Gelatin is a protein made from collagen and used in foods, medicines, capsules, and some medical materials.

---

## Gelfoam

An absorbable sponge made from purified animal gelatin that helps control bleeding during surgery. It provides a surface for clot formation and is gradually absorbed; swelling or infection can rarely occur.

## Gelineau Syndrome

A sleep disorder causing repeated, uncontrollable daytime sleep episodes. Some people also have sudden muscle weakness triggered by emotion, sleep paralysis, or vivid dream-like experiences while falling asleep or waking.

## Gemcitabine

A chemotherapy medicine that becomes active inside cells and interferes with DNA production, slowing or killing rapidly dividing cancer cells. It is used for several cancers and may cause low blood counts, nausea, rash, or organ toxicity.

## Gemcitabine Hydrochloride

The hydrochloride form of gemcitabine, a chemotherapy medicine that interferes with DNA production in dividing cells. It is used for several cancers; important risks include infection from low blood counts and treatment-related organ injury.

## Gemella

Gemella is a group of bacteria that normally may inhabit the mouth and upper airway but can rarely cause serious infections, including bloodstream infection.

## Gemfibrozil

A medicine that lowers very high blood triglycerides and may reduce the risk of pancreatitis. It changes how the liver processes fats but can interact with statins and increase the risk of serious muscle injury.

## Gemifloxacin

An antibiotic in the fluoroquinolone group used for selected bacterial respiratory infections. It can cause nausea and, rarely, tendon injury, nerve symptoms, abnormal heart rhythm, severe diarrhea, or dangerous allergic reactions.

## Gemination

An attempt by one developing tooth bud to divide, producing a large tooth with a split or partly divided crown. The total number of teeth is usually normal, and dental assessment guides treatment.

## Gemistocytic Astrocytoma

A former histologic variant of diffuse astrocytoma composed partly of gemistocytic astrocytes. The term is now used mainly descriptively because integrated molecular classification determines the modern tumor diagnosis and grade.

## Gemtuzumab Ozogamicin

An antibody-linked chemotherapy medicine used for selected acute myeloid leukemia. The antibody attaches to CD33-positive leukemia cells and delivers a toxic drug; liver injury, infusion reactions, and low blood counts are important risks.

## Gender

A social, cultural, and personal concept involving identity, roles, expression, and expectations. Gender is related to, but distinct from, sex characteristics and sexual orientation.

## Gender Dysphoria

Clinically significant distress or impairment associated with incongruence between a person's experienced or expressed gender and sex assigned at birth. Having a gender identity different from sex assigned at birth does not by itself constitute gender dysphoria.

## Gender Identity

A person's internal sense of being a woman, man, both, neither, or another gender, which may or may not correspond to sex assigned at birth.

## Gender Incongruence

A difference between a person's experienced or expressed gender and sex assigned at birth. Gender incongruence is a descriptive term and does not by itself imply distress, impairment, or a mental disorder.



---

## Gene

A section of DNA that contains instructions for making a functional product, usually a protein or RNA. Genes are arranged at specific locations on chromosomes and influence inherited traits and cell function.

## Gene Amplification

An increase in the number of copies of a gene or genomic region, which can raise gene expression and drive some inherited conditions or cancers.

## Gene Conversion

A nonreciprocal transfer of sequence information from one DNA region to a homologous region, potentially changing an allele during recombination or repair.

## Gene Deletion

Loss of part or all of a gene from DNA, potentially abolishing its product or altering dosage and causing disease when clinically significant.

## Gene Dosage

The number of functional copies of a gene and the amount of product they produce, influenced by deletions, duplications, chromosome number, and regulation.

## Gene Duplication

An extra copy of a gene created by replication or rearrangement, which may have no effect, alter dosage, or provide material for new gene function.

## Gene Expression

The process by which a cell uses information in a gene to make RNA or protein. Cells control when and how strongly genes are expressed, allowing different tissues to develop and perform specialized functions.

## Gene Frequency

The proportion of all gene copies in a population that carry a particular version, or allele, of that gene. It describes population genetics and does not

by itself predict an individual's disease risk.

## Gene Fusion

A hybrid gene formed when segments of two separate genes join, sometimes producing an abnormal protein that drives cancer or other disease.

## Gene Modification

An intentional or naturally occurring alteration of DNA sequence or gene regulation that changes cellular function or inherited traits.

## Gene Mutation

A permanent change in the DNA sequence of a gene, which may be harmless, beneficial, or pathogenic depending on its effect and inheritance.

## Gene Panel

A genetic test that examines a selected group of genes linked to a person's symptoms or suspected condition. It can provide focused results but may miss changes in genes or DNA regions not included in the panel.

## Gene Rearrangement

Reorganization of DNA segments within or between chromosomes that can alter gene structure, expression, immune-receptor diversity, or cancer biology.

## Gene Silencing

Reduction or stopping of a gene's activity through DNA changes, chromatin control, RNA interference, or other processes that limit protein production.

## Gene Targeting

A molecular method that changes a chosen genomic locus through homologous recombination or engineered editing to study or alter gene function.

## Gene Therapy

Treatment that adds, removes, replaces, or edits genetic material in a patient's cells to treat disease. It may correct a missing gene function or alter harmful activity; risks include immune reactions and unintended genetic effects.

---

## Gene Transfer

Movement of genetic material into a cell or organism, naturally or through laboratory vectors, potentially changing gene expression or correcting a defect.

## Gene-Environment Interaction

A gene-environment interaction occurs when an environmental exposure changes disease risk differently according to a person's genotype, so inherited susceptibility and exposure jointly influence phenotype.

## Gene-Modified

Describing cells or organisms whose genetic material has been deliberately altered using recombinant DNA or genome-editing methods.

## General Anesthesia

A controlled, temporary state of unconsciousness produced by medicines so a person does not feel or remember a procedure. Breathing, blood pressure, heart rate, and other body functions are monitored throughout.

## General Anesthetic

A medicine used to produce a controlled, reversible state of unconsciousness and lack of awareness during surgery or another procedure. It may be combined with separate medicines for pain relief or muscle relaxation; breathing and other vital functions require monitoring.

### Alternate Names:

General Anesthetics

## General Paresis

A late form of syphilis affecting the brain and spinal cord. It can gradually cause memory loss, personality or mood changes, poor judgment, weakness, tremor, or difficulty walking. Blood and spinal-fluid tests help identify the infection.

### Alternate Names:

General Paresis of the Insane; Paralytic Dementia; Neurosyphilis

## Generalised

Widespread or involving multiple regions of the body

rather than being confined to a single localized site.

## Generalized

Distributed widely or involving most or all of a body region, system, or population rather than being confined to one focus.

## Generalized Anxiety Disorder

A mental-health disorder involving excessive, difficult-to-control worry about several areas of life on most days for at least six months. Restlessness, tiredness, poor concentration, irritability, muscle tension, and sleep problems may occur.

## Generalized Anxiety Disorder In Children

Excessive, difficult-to-control worry in children about multiple topics (school performance, health, family safety) occurring more days than not for at least 6 months. Associated with restlessness, fatigue, difficulty concentrating, irritability, muscle tension, and sleep disturbance. Treatment includes cognitive behavioral therapy and possibly medication.

### Alternate Names:

Pediatric GAD; Childhood Generalized Anxiety

## Generalized Essential Telangiectasia

Widespread fine superficial dilated vessels without an identifiable underlying systemic disease or localized cause. They are usually asymptomatic, but a new or rapidly progressive pattern should be assessed for medicines, connective-tissue disease, liver disease, or other secondary causes.

## Generalized Granuloma Annulare

A widespread skin condition causing many small, skin-colored, red, or purple bumps over the trunk and limbs. It may be associated with diabetes and can last longer than localized granuloma annulare.

## Generalized Lipodystrophy

A disorder with near-total loss of adipose tissue, severe insulin resistance, diabetes, high triglycerides, fatty liver, and abnormal fat deposition in other tissues.

---

## Generalized Lymphadenopathy

Pathological enlargement of lymph nodes involving two or more non-contiguous anatomical lymph node chains. Unlike localized adenopathy, it reflects systemic, hematologic, infectious, or autoimmune disease. Major etiologies include lymphomas, leukemias (such as chronic lymphocytic leukemia), systemic viral infections (HIV, Epstein-Barr virus, cytomegalovirus), secondary syphilis, disseminated tuberculosis, systemic lupus erythematosus, sarcoidosis, and drug-induced hypersensitivity reactions (such as DRESS syndrome).

### Alternate Names:

Diffuse Lymphadenopathy; Systemic Lymphadenopathy

## Generalized Seizure

A seizure arising from or rapidly involving both cerebral hemispheres, commonly producing impaired awareness, bilateral motor activity, or brief absence episodes.

## Generalized Tonic-Clonic Seizure

A seizure causing loss of consciousness, stiffening of the whole body, and then repeated jerking of the limbs. Confusion and sleepiness often follow; a seizure lasting five minutes or longer is prolonged.

## Generalized Weakness

Reduced strength affecting multiple body regions, caused by systemic illness, metabolic or electrolyte disturbance, medication, infection, muscle disease, nerve disease, neuromuscular-junction disease, or brain or spinal-cord disease. It may occur with speech, facial, breathing, or walking difficulty.

## Generalized-Onset Seizure

A seizure that begins in networks on both sides of the brain. It may cause staring, brief muscle jerks, stiffening, loss of muscle tone, or whole-body stiffening and jerking, depending on the type.

## Generator

A device that produces a short-lived radioactive substance from a longer-lived one for nuclear-medicine tests. The product is separated, checked for purity

and sterility, and then used to make a tracer.

## Generic

Describing a medicine by its active ingredient rather than a brand name; an approved generic drug must meet applicable standards for quality and therapeutic equivalence to its reference product.

## Genes

Sections of DNA that contain instructions for making proteins or functional RNA. They are arranged on chromosomes and influence inherited traits, cell activity, disease risk, and responses to some treatments.

## Genesis

The origin, formation, or development of something, such as a disease, organ, symptom, or process. Understanding its genesis can help explain causes, risk factors, progression, prevention, and treatment.

## Genetic Anticipation

Earlier onset or increasing severity of a hereditary disorder in successive generations, often caused by expansion of unstable repeated DNA sequences.

## Genetic Carrier

A person with one pathogenic variant for a recessive disorder who is usually unaffected but can transmit the variant to children.

## Genetic Code

The rules cells use to read genetic instructions in groups of three DNA or RNA building blocks. Each group directs the cell to add a particular amino acid to a protein or to stop making the protein.

## Genetic Counseling

A professional discussion that explains how inherited factors may affect a person or family. It may cover risk, testing choices, results, reproductive options, emotional support, and implications for relatives.

## Genetic Counselling

A process in which a trained professional explains in-

---

herited conditions, test results, health implications, and reproductive choices to a person or family.

### Genetic Counselor

A trained health professional who explains inherited conditions, estimates genetic risk, discusses testing, and helps patients and families understand results and choices. They may coordinate testing of relatives when appropriate.

### Genetic Disease

Alternate name; see *Genetic Disorder*.

### Genetic Disorder

A disease or disorder caused wholly or partly by a pathogenic change in genetic material. The change may involve one or more genes, chromosome number or structure, mitochondrial DNA, or gene regulation. It may be inherited through germline DNA, arise de novo, or occur only in certain tissues as a somatic change. Some conditions also result from interactions among multiple genetic variants and environmental factors; not every genetic change causes disease.

### Genetic Drift

Random changes in allele frequencies from one generation to the next, especially in small populations, independent of natural selection.

### Genetic Engineering

Deliberate changing, adding, removing, or transferring genetic material using laboratory methods. It supports research, diagnosis, treatment development, agriculture, and biotechnology but requires careful assessment of safety, ethics, and unintended effects.

### Genetic Heterogeneity

The situation in which similar features or the same disease can result from different genetic causes. Different genes or different changes within one gene may produce overlapping effects.

### Genetic Linkage

The tendency of nearby genes or DNA markers on the same chromosome to be inherited together because crossing over is less likely between them.

### Genetic Marker

A detectable DNA variation or feature used to track inheritance, identify a chromosome region, assess disease risk, or distinguish samples.

### Genetic Polymorphism

A common DNA variation with at least two alleles in a population, which may influence traits, drug response, or disease susceptibility.

### Genetic Profile

A set of genetic variants or DNA markers describing an individual, tumor, organism, or sample for diagnosis, risk assessment, identification, or research.

### Genetic Recombination

Exchange or rearrangement of DNA segments that creates new allele combinations during meiosis, repair, immune-cell development, or laboratory manipulation.

### Genetic Screening

Testing people without symptoms to estimate the chance of a genetic condition. Examples include newborn screening, carrier screening before or during pregnancy, and prenatal screening; screening results are not always diagnostic.

### Genetic Sequencing

Analysis of the order of DNA bases to identify variants associated with genetic disorders. The method is selected according to the gene, variant, and clinical question.

#### Alternate Names:

DNA Sequencing; Next-Generation Sequencing (NGS)

### Genetic Sonogram

A detailed pregnancy ultrasound, usually performed around 18–22 weeks, that examines fetal anatomy

---

and may look for findings associated with chromosome conditions. It cannot detect every genetic disorder.

### **Genetic Susceptibility**

Inherited tendency to develop a disease or respond to an exposure, which usually modifies risk rather than determining outcome alone.

### **Genetic Technique**

A laboratory method used to detect, amplify, sequence, edit, compare, or otherwise analyze DNA, RNA, chromosomes, or gene expression.

### **Genetic Testing**

Analysis of DNA or chromosomes to assess disease risk, explain symptoms, identify carriers, or guide treatment. Results may also provide information about inherited traits and conditions affecting relatives.

### **Genetic Transcription**

The first step in using a gene: an enzyme copies information from DNA into RNA. The RNA may then guide protein production or perform another function inside the cell.

### **Genetic Variant Of Uncertain Significance**

A DNA change whose effect on health is not yet known. It does not by itself show that a person has a disease or predict future disease and may later be classified as harmless or disease-causing as evidence improves.

### **Genetics**

Genetics is the study of genes, heredity, genetic variation, and how DNA and gene regulation influence traits, disease risk, diagnosis, and treatment response.

### **Geniculate Body**

One of two paired relay structures in the brain. The lateral geniculate body processes visual signals, while the medial geniculate body processes hearing

signals before sending them to the cerebral cortex.

### **Geniculate Ganglion**

An L-shaped sensory ganglion of the facial nerve (cranial nerve VII) located within the facial canal of the temporal bone, housing the primary pseudo-unipolar sensory neuronal cell bodies conveying special taste sensation from the anterior two-thirds of the tongue (via chorda tympani).

### **Genital**

Relating to the reproductive organs or external sexual structures. Genital symptoms may involve skin, mucosa, nerves, glands, or the urinary and reproductive systems.

### **Genital Edema**

Swelling of genital tissues from fluid accumulation caused by infection, inflammation, allergy, trauma, venous or lymphatic obstruction, or systemic disease.

### **Genital Herpes**

An infection of the genital or anal area caused by herpes simplex virus type 1 or type 2 (HSV-1 or HSV-2). It spreads mainly through sexual skin-to-skin contact and can be transmitted even when no sores are present. It may cause painful blisters or ulcers, burning urination, swollen glands, or no symptoms; outbreaks can recur. Infection during pregnancy can threaten the newborn.

### **Genital Herpes in Pregnancy**

Genital herpes caused by HSV-1 or HSV-2 can be acquired, recur, or be transmitted to the newborn during pregnancy or delivery. Management considers timing of infection, lesions or prodrome at labor, antiviral therapy, and delivery planning to reduce neonatal transmission.

### **Genital Herpes in Women**

Herpes simplex virus infection of the female genitalia, causing painful blisters or ulcers. It may have no symptoms. Antiviral treatment can reduce symptoms and transmission; infection during pregnancy can affect the newborn.



---

**Alternate Names:**

Female Genital Herpes; HSV-2 Infection in Women

**Genital Injury**

Damage to the external or internal genital tissues from childbirth, accidents, sexual contact, assault, medical procedures, or disease. It may cause pain, swelling, bleeding, or difficulty urinating.

**Genital Mycoplasmas**

Bacteria that may live in or infect the genital or urinary tract. Some cause inflammation of the urethra or cervix, pelvic inflammatory disease, or few symptoms; testing helps distinguish infection from harmless colonization.

**Genital Prolapse**

The dropping or bulging of one or more pelvic organs (such as the bladder, uterus, or rectum) into or through the vagina due to weakening of the pelvic floor muscles and supporting tissues. It commonly causes a feeling of a vaginal bulge or heaviness, lower back discomfort, and urinary or bowel difficulties; risks include vaginal childbirth, aging, menopause, and chronic straining.

**Alternate Names:**

Pelvic organ prolapse; Uterovaginal prolapse; Urogenital prolapse.

**Genital System**

The organs involved in reproduction and sexual function, including the gonads, reproductive ducts, accessory glands, and external genital structures. Anatomy differs between individuals and changes with development and age.

**Genital Ulcer Disease**

A clinical syndrome characterized by single or multiple ulcerative breaches of the genital or perianal skin and mucosa, predominantly caused by sexually transmitted pathogens including herpes simplex virus (HSV-2), *Treponema pallidum* (syphilis chancre), and *Haemophilus ducreyi* (chancroid).

**Alternate Names:**

GUD

**Genital Wart**

A wart in the genital or anal area caused by certain types of human papillomavirus. Warts may be flat or raised and can recur; treatment removes visible lesions but may not eliminate the virus immediately.

**Genital Warts**

Benign anogenital growths caused most often by low-risk HPV types 6 and 11. They may be flat, raised, or cauliflower-like and may recur after treatment. They spread through intimate skin-to-skin contact; evaluation may include screening for other sexually transmitted infections, and HPV vaccination prevents many causative types.

**Alternate Names:**

Condylomata Acuminata; Anogenital Warts

**Genitalia**

The external and internal reproductive organs. The term includes the vulva and vagina, or the penis, scrotum, and associated structures, together with the gonads and reproductive ducts when used broadly; anatomy varies with sex, age, and developmental stage.

**Genitals**

The external and internal reproductive organs. Female genitals: external (vulva — mons pubis, labia majora, labia minora, clitoris, vaginal opening, Bartholin glands) and internal (vagina, cervix, uterus, fallopian tubes, ovaries).

**Genito-Pelvic Pain/Penetration Disorder**

Persistent or recurrent difficulty with vaginal penetration, including pain, fear or anxiety about pain, involuntary pelvic-floor tightening, or pain during attempted penetration, causing clinically significant distress. Causes may include pelvic-floor problems, infection, hormonal changes, injury, or emotional factors.

**Genito-Urinary Medicine**

The medical specialty concerned with urinary and reproductive health. It includes evaluation and treatment of urinary infections, sexual health problems, sexually transmitted infections, and some re-



---

productive conditions.

### **Genitofemoral Nerve**

A branch of the lumbar plexus (arising from L1 and L2 nerve roots) that pierces the anterior surface of the psoas major muscle and divides into a genital branch (innervating the cremaster muscle and scrotal/labial skin; mediating the cremasteric reflex) and a femoral branch (cutaneous sensation to the anterior upper thigh).

### **Genitourinary**

Relating to the urinary system and the reproductive organs, which share developmental and anatomical relationships.

### **Genitourinary Fistula**

Alternate name for urogenital fistula, an abnormal connection between the urinary and genital tracts.

### **Genitourinary Neoplasm**

An abnormal growth in the urinary or reproductive organs, such as the kidney, bladder, prostate, testis, ovary, or uterus. It may be benign or cancerous; examination, imaging, and sometimes biopsy identify its nature.

### **Genitourinary Syndrome Of Menopause**

A group of changes in the genital and urinary tissues after menopause, caused mainly by lower estrogen levels. It may include vaginal dryness, burning, pain during sex, urinary urgency, painful urination, or repeated urinary infections.

### **Genitourinary Tract**

The combined urinary and reproductive organs and their associated ducts.

### **Genodermatoses**

Inherited disorders that mainly affect the skin, hair, nails, sweat glands, or related organs. Examples include inherited scaling disorders, blistering diseases, and some neurocutaneous syndromes; severity varies widely.

### **Genome**

The complete genetic material of an organism, including genes and DNA that does not directly code proteins. It contains instructions influencing development, body function, inherited traits, disease risk, and treatment response.

### **Genomic Imprinting**

A pattern in which a gene's activity depends on whether its copy came from the mother or father. Abnormal imprinting can cause conditions such as Prader–Willi or Angelman syndrome.

### **Genomic Instability**

An increased tendency for DNA changes, chromosome damage, and mutations to accumulate, contributing to cancer development, treatment resistance, and disease progression.

### **Genomics**

The study of an organism's complete genetic material, including its DNA sequence, variation, activity, and interaction with disease and treatment. It supports research, diagnosis, risk assessment, and personalized care.

### **Genotoxicity**

The ability of a substance or physical agent to damage genetic material. Genotoxicity may cause mutations or chromosome damage, but a positive test does not by itself establish that an agent causes cancer in humans.

### **Genotype**

A person's genetic makeup, meaning the versions of one or more genes they carry. The genotype helps influence observable traits and disease risk together with other genes and environmental factors.

### **Genotype Determination**

Genotype determination is the use of laboratory testing to identify an individual's genetic variants or alleles at one or more locations.

---

## Genotype-Phenotype Correlation

The relationship between a person's genetic makeup and the observable features, severity, age of onset, or clinical course of a condition; correlations may be incomplete because other genetic and environmental factors contribute.

## Gentamicin

An aminoglycoside antibiotic that interferes with bacterial protein production. It can injure the kidneys or hearing and is used with dose and blood-level monitoring.

## Gentamicin Sulfate

The sulfate salt of gentamicin, an aminoglycoside antibiotic that interferes with bacterial protein production. It can injure the kidneys or hearing and is used with careful dosing and monitoring.

## Gentian

A plant of the *Gentiana* genus whose roots or other parts have been used in herbal preparations, traditionally as a bitter digestive; clinical evidence and product quality vary.

## Gentian Violet

A purple antiseptic and dye with antifungal and antibacterial activity, used in selected topical applications but capable of irritating tissue and staining skin.

## Genu

The knee or a knee-shaped part of an anatomical structure; the term also describes a bend, such as the genu of the corpus callosum.

## Genu Recurvatum

Hyperextension of the knee beyond the usual straight position, caused by laxity, deformity, muscle imbalance, neurologic disease, or injury.

## Genu Valgum

An alignment abnormality in which the knees angle inward and may touch while the ankles remain apart. It may be developmental or caused by bone, joint,

or metabolic disease.

### Alternate Names:

Knock-Knees; Tibial Valgus

## Genu Varum

Outward bowing of the legs, commonly called bowlegs. It is often normal in young children and improves with growth, but persistent, worsening, unequal, painful, or severe bowing may result from rickets, growth-plate disease, or another condition.

### Alternate Names:

Bowlegs; Bandy Legs

## Geographic Atrophy

An advanced form of dry age-related macular degeneration in which areas of light-sensing and supporting cells in the center of the retina are permanently lost. It causes slowly worsening blind spots and loss of detailed central vision.

## Geographic Tongue

A harmless condition in which smooth red patches with pale borders appear and change location on the tongue. It usually causes no symptoms but may make spicy or acidic foods sting.

### Alternate Names:

Benign Migratory Glossitis

## Geophagia

Persistent eating of soil or other nonfood substances; it may cause poisoning, infection, intestinal obstruction, or iron-deficiency anemia.

## Geophagy

The deliberate ingestion of soil or clay; it may cause lead or parasite exposure, gastrointestinal obstruction, or iron-deficiency anemia and can occur with pica.

## Geotrichum

*Geotrichum* is a genus of fungi that may colonize humans and can rarely cause superficial or invasive infection, especially in people with weakened immunity.

---

## Geriatric Assessment

A broad review of an older person's medical conditions, medicines, mobility, memory, mood, daily activities, nutrition, safety, home situation, and goals. It identifies treatable problems and guides an individualized care plan.

## Geriatric Bioethics

The systematic exploration of moral and ethical considerations in the medical care of elderly patients, encompassing decisional capacity, informed consent, advanced healthcare directives, surrogate decision-making, palliative care transitions, and protection of vulnerable older adults.

### Alternate Names:

Geriatric Bioethics; Ethics in Geriatric Medicine; Ethics of Aging

## Geriatric Care Manager

A professional who helps coordinate medical, social, housing, and support services for an older adult. They may assess needs, organize care, communicate with families, and help plan for changing abilities.

### Alternate Names:

Aging-Life Care Manager; Eldercare Coordinator

## Geriatric Dentistry

Dental care adapted to the needs of older adults, including tooth decay, gum disease, dry mouth, dentures, swallowing concerns, medicine effects, and examination for oral cancer.

## Geriatric Depression

Depression in an older adult may cause low mood, loss of interest, fatigue, sleep or appetite changes, slowed thinking, physical complaints, or memory problems. Medical illness, medicines, isolation, and bereavement may contribute.

### Alternate Names:

Late-Life Depression; Depression in the Elderly; Late-Onset Depressive Disorder

## Geriatric Depression Scale

A questionnaire used to screen older adults for symptoms of depression. It can identify people with pos-

sible depression, but a positive result does not by itself establish depression or explain its cause.

## Geriatric Dream Changes

Dreams in older adults may become more vivid or be affected by medicines, sleep disorders, depression, or changes in the sleep cycle. Repeated frightening dreams, acting them out, confusion, or sudden change may occur.

### Alternate Names:

Geriatric Sleep Patterns; Dreaming in Older Adults

## Geriatric Dysphagia

Difficulty swallowing in an older adult. It may cause coughing or choking during meals, food remaining in the mouth, weight loss, dehydration, or aspiration and can result from stroke, dementia, muscle disease, blockage, or medicines.

### Alternate Names:

Presbyphagia; Age-Related Swallowing Dysfunction

## Geriatric Hospitalization

Hospital care for an older adult, with attention to medical illness as well as mobility, medicines, nutrition, hearing, vision, and cognition. Preventing falls, delirium, pressure injuries, and loss of independence supports safer recovery.

### Alternate Names:

Inpatient Geriatric Care; Acute Care for Elders (ACE)

## Geriatric Infection

Infection in an older adult may cause confusion, weakness, reduced appetite, or a fall without causing fever. Common sites include the urine, lungs, skin, and bloodstream; illness can worsen quickly.

### Alternate Names:

Infection in the Elderly; Late-Life Infection

## Geriatric Informed Consent

The process of ensuring that an older patient understands a proposed intervention, its benefits, risks, and alternatives before choosing freely. If decision-making ability is impaired, an authorized decision-maker may help according to local law.

---

**Alternate Names:**

Geriatric Decision-Making Capacity; Surrogate Consent in Older Adults

**Geriatric Medical Ethics**

Ethical decision-making in the care of older adults, including respect for the person's wishes, decision-making capacity, likely benefits and harms, fairness, privacy, and appropriate involvement of family or caregivers.

**Geriatric Nursing**

Nursing care focused on the health, function, medication safety, cognition, mobility, and goals of older adults.

**Geriatric Nutrition**

Nutrition care for older adults that considers aging, illness, medicines, frailty, dental or swallowing problems, appetite, and access to food. Goals include maintaining strength, muscle, hydration, function, and quality of life.

**Geriatric Osteoarthritis**

Osteoarthritis occurring in later life, commonly affecting the knees, hips, hands, or spine and causing joint pain, stiffness, and reduced movement. Management may include exercise, weight management, medicines, assistive measures, or joint replacement when appropriate.

**Alternate Names:**

Age-Related Osteoarthritis; Senile Osteoarthritis

**Geriatric Rehabilitation**

Therapy that helps an older adult regain or maintain movement, strength, independence, and safety after illness, injury, surgery, or functional decline. The plan is adapted to health, abilities, goals, and support needs.

**Geriatric Screening**

Checking older adults who have no symptoms for diseases or risks that may benefit from early detection. Examples include falls, depression, osteoporosis, and selected cancers.

**Alternate Names:**

Geriatric Screening; Senior Health Assessment

**Geriatric Sleep Changes**

Sleep patterns and sleep problems in older adults, including insomnia, sleep apnea, circadian changes, and medication-related disturbances.

**Alternate Names:**

Geriatric Sleep Disorders; Elder Sleep Management

**Geriatric Syndrome**

A common problem in older adults caused by several interacting factors rather than one disease. Examples include falls, frailty, delirium, incontinence, immobility, and pressure injuries; causes may be medical, social, or both.

**Geriatric Trauma**

Injury in an older adult, often involving falls, fractures, head injury, or bleeding. Age-related frailty, osteoporosis, balance problems, and medicines that increase bleeding can make injuries more serious and recovery slower.

**Geriatrician**

A physician specializing in the health care of older adults, including complex illness, multiple conditions, medications, function, and goals of care.

**Geriatrics**

The medical specialty focused on health, function, independence, and quality of life in older adults. Care considers multiple illnesses, medicines, memory, mobility, nutrition, mood, safety, goals, and support.

**Germ**

Alternate informal term; see *Microorganism*.

**Germ Cell**

A reproductive cell or precursor that gives rise to sperm or eggs and carries genetic material to the next generation.

---

## **Germ Cell Tumor**

A tumor arising from cells that normally develop into sperm or eggs. It may occur in the testicles, ovaries, or other sites such as the chest or abdomen; some types are highly treatable with surgery, chemotherapy, or both.

## **Germ Cell Tumor Staging**

Determination of the extent and risk of a germ-cell tumor using examination, imaging, tumor markers, surgical findings, and pathology. Stage and histologic type guide surgery, chemotherapy, surveillance, and prognosis.

## **Germ Cell Tumors**

Tumors arising from cells that normally develop into sperm or eggs. They can occur in the testicles, ovaries, or other body sites and may be benign or cancerous, depending on the type.

## **Germ Layer**

One of the three early layers of an embryo—ectoderm, mesoderm, or endoderm—from which all major tissues and organs develop. Each layer gives rise to characteristic body structures during development.

## **Germ Theory Of Disease**

The evidence-based principle that particular microorganisms can cause particular infectious diseases. It underlies infection control, vaccination, antimicrobial treatment, sterilization, and methods used to identify disease-causing organisms.

## **German Measles**

Rubella: a contagious viral infection caused by rubella virus, characterized by fever, lymph-node enlargement, and a fine rash. Infection during pregnancy can harm the fetus, causing congenital rubella syndrome with cardiac, auditory, and developmental defects.

### **Alternate Names:**

Rubella; Three-Day Measles

## **Germanium**

A chemical element sometimes included in unproven health supplements. Germanium compounds have no established medical benefit and can cause kidney damage, nerve damage, or other serious toxicity when taken internally.

## **Germinoma**

A usually slow-growing germ-cell cancer that most often develops in the pineal or suprasellar area of the brain. Related tumors occur in the testicles or ovaries; treatment commonly uses radiation, chemotherapy, surgery, or combinations.

## **Germline Mutation**

A genetic change present in an egg or sperm and therefore usually present in nearly every cell of the person. Some changes can be passed to children and may increase the risk of inherited disorders or cancer.

## **Germline Testing**

Genetic testing of blood, saliva, or another sample to look for changes that may be inherited or passed to children. Results can affect the person's care, relatives, cancer screening, and reproductive decisions.

## **Germline Variant**

A DNA change present in cells that can give rise to eggs or sperm and therefore may be passed to children. It is usually present in most cells of the person, unlike a change limited to a tumor or other tissue.

## **Gerontologist**

A specialist studying aging and its biological, psychological, social, or economic effects. Gerontologists may work in research, healthcare, policy, counseling, education, housing, or community services for older people.

## **Gerstmann Syndrome**

A neurological cognitive syndrome caused by a lesion of the dominant angular gyrus in the inferior parietal lobule (typically left hemisphere), character-

---

ized by the classic tetrad of agraphia (dysgraphia), acalculia (dyscalculia), finger agnosia, and right-left disorientation.

**Alternate Names:**

Angular Gyrus Syndrome; Dominant Parietal Lobe Syndrome

**Gerstmann-Straussler-Scheinker Syndrome**

A rare, autosomal dominant inherited prion disease caused by mutations in the PRNP gene (most commonly P102L), characterized pathologically by multicentric PrP-amyloid plaques throughout the cerebellum and cerebral cortex, presenting with slowly progressive cerebellar ataxia, spasticity, and late dementia.

**Alternate Names:**

GSS Syndrome; GSS Prion Disease

**Gesell Developmental Schedules**

Standardized developmental assessment schedules that describe expected motor, language, adaptive, and social milestones in children; they support screening but do not by themselves diagnose a disorder.

**Gestation**

The period of pregnancy during which an embryo and fetus develop in the uterus. Pregnancy is usually dated as about 40 weeks from the first day of the last menstrual period, or about 38 weeks from fertilization.

**Gestation Period**

The time from conception to birth or, clinically, from the last menstrual period to delivery; human pregnancy is usually about 40 weeks from the last menstrual period.

**Gestational Age**

The length of a pregnancy measured in weeks from the first day of the last menstrual period. It is usually about two weeks longer than the time since fertilization and helps estimate development, due date, and appropriate care.

**Alternate Names:**

Menstrual Age; Fetal Postmenstrual Age

**Gestational Diabetes**

Carbohydrate intolerance of variable severity with onset or first recognition during pregnancy. It occurs when pancreatic insulin secretion cannot overcome pregnancy-induced insulin resistance driven by placental hormones, including human placental lactogen, progesterone, and cortisol. Maternal risks include preeclampsia and operative delivery; fetal and neonatal risks include macrosomia, shoulder dystocia, birth injury, neonatal hypoglycemia, hyperbilirubinemia, and respiratory distress. Screening is typically performed between 24 and 28 weeks of gestation using an oral glucose challenge or tolerance test. Management centers on medical nutrition therapy, physical activity, and blood glucose monitoring, with insulin or oral agents added when targets are not met. Glucose levels typically normalize after delivery, but affected individuals carry a significantly increased lifetime risk of developing type 2 diabetes mellitus.

**Alternate Names:**

GDM; Gestational Diabetes Mellitus

**Gestational Diabetes Mellitus**

Alternate medical term; see *Gestational Diabetes*.

**Gestational Hypertension**

New-onset high blood pressure after 20 weeks of pregnancy without protein in the urine or other features of preeclampsia. It may later develop into preeclampsia.

**Gestational Pyelonephritis**

A kidney infection occurring during pregnancy, usually causing fever, flank pain, nausea, or urinary symptoms. It can lead to sepsis, preterm labor, or breathing complications.

**Alternate Names:**

Pyelonephritis in Pregnancy; Gravid Kidney Infection

**Gestational Sac**

The first fluid-filled structure usually seen inside the uterus on an early pregnancy ultrasound. Its size



---

and appearance help estimate pregnancy age and assess early development, but an early scan may not yet show an embryo.

### **Gestational Trophoblastic Disease**

A group of uncommon conditions caused by abnormal growth of cells that normally help form the placenta. It includes molar pregnancy and some pregnancy-related cancers and may cause unusually high pregnancy hormone levels, bleeding, or a uterus larger than expected.

### **Gestational Trophoblastic Neoplasia**

Persistent or cancerous growth of placenta-related cells after a pregnancy, miscarriage, abortion, or molar pregnancy. It is often detected by pregnancy-hormone levels that remain high or rise and is usually very responsive to specialist cancer treatment.

### **GFR**

Glomerular filtration rate, an estimate of how well the kidneys filter blood. It is usually calculated from blood creatinine and other information, and a persistently low value may indicate chronic kidney disease.

#### **Alternate Names:**

Glomerular Filtration Rate; Estimated GFR (eGFR)

### **GHD**

Abbreviation; see *Growth Hormone Deficiency*.

### **Ghon Complex**

The combination of a primary tuberculous lung focus (Ghon focus) and caseous granulomatous lymphadenitis in the draining regional hilar or mediastinal lymph nodes, representing the hallmark of primary pulmonary tuberculosis that often undergoes fibrocalcification.

#### **Alternate Names:**

Primary Tuberculous Complex; Ranke Precursor Complex

### **Ghon Focus**

A primary granulomatous tuberculous pulmonary

lesion consisting of a circumscribed area of caseous necrosis typically located subpleurally in the mid-to-lower lung zone, representing the initial site of *Mycobacterium tuberculosis* infection.

#### **Alternate Names:**

Primary Focus of Ghon; Subpleural Tuberculous Focus

### **Ghrelin**

A hormone produced mainly in the stomach that increases hunger and helps regulate energy balance. It also stimulates growth-hormone release and changes with eating, fasting, sleep, and body weight.

### **Gianotti Crosti Syndrome**

Alternate spelling; see *Gianotti-Crosti Syndrome*.

#### **Alternate Names:**

Papular Acrodermatitis of Childhood; Infantile Lichenoid Acrodermatitis

### **Gianotti-Crosti Syndrome**

A childhood skin condition characterized by a papular or papulovesicular rash on the face, buttocks, and extremities, often following viral infections (EBV, hepatitis B, CMV). Self-limited, lasting 2-8 weeks. Also called papular acrodermatitis of childhood.

### **Giant Axonal Disease**

Alternate name; see *Giant Axonal Neuropathy*.

### **Giant Axonal Neuropathy**

A rare inherited disorder in which abnormal giant nerve fibers develop, causing progressive peripheral neuropathy, weakness, loss of sensation, poor coordination, and sometimes changes in hair or skin. Onset and severity vary.

### **Giant Cell Arteritis**

Inflammation of arteries, usually in people over 50, often affecting vessels near the temples and eyes. It can cause new headache, scalp tenderness, jaw pain while chewing, or vision changes and may permanently damage sight.

---

### **Giant Cell Arteritis Emergency**

A severe presentation of suspected giant cell arteritis with visual symptoms, jaw or tongue pain during chewing, or reduced blood flow to the nervous system. It carries a risk of permanent visual loss.

### **Giant Cell Glioblastoma**

A rare histologic variant of glioblastoma characterized by numerous large, often multinucleated tumor cells. It may cause headache, seizures, weakness, or changes in speech, vision, memory, or behavior, depending on location. Diagnosis uses brain imaging and tumor tissue analysis; treatment is coordinated by a neuro-oncology team.

#### **Alternate Names:**

Glioblastoma, Giant Cell; Monstrocellular Sarcoma

### **Giant Cell Interstitial Pneumonia**

A rare severe form of interstitial lung inflammation with multinucleated giant cells, causing cough, breathlessness, and low oxygen. It requires specialist evaluation to identify the cause and guide treatment.

### **Giant Cell Myocarditis**

A rare, severe inflammation of the heart muscle caused by an abnormal immune reaction. It can weaken the heart and cause dangerous heart rhythms or heart failure.

### **Giant Cell Tumor**

See *Giant Cell Tumor Of Bone*.

### **Giant Cell Tumor Of Bone**

A usually noncancerous but locally aggressive bone tumor made up of abnormal cells and many large bone-resorbing cells. It usually occurs near the ends of long bones in adults after skeletal growth is complete. It can damage nearby bone, recur, and rarely spread to the lungs.

#### **Alternate Names:**

GCTB; Osteoclastoma

### **Giant Cell Tumor Of Soft Tissue**

A usually low-grade soft-tissue neoplasm containing numerous osteoclast-like giant cells, most often arising in the extremities of adults and generally having a favorable outcome after excision.

### **Giant Cell Tumor Of Tendon Sheath**

A common benign localized tenosynovial tumor, usually presenting as a slow-growing firm nodule on a finger or hand; recurrence can occur after excision.

### **Giant Cells**

Large cells formed by fusion of macrophages or abnormal cellular growth, seen in granulomatous inflammation, foreign-body reactions, and some tumors.

### **Giant Congenital Nevus**

A very large dark or brown skin mark present at birth. It may grow with the child and has a higher risk of melanoma than smaller birthmarks.

#### **Alternate Names:**

Giant Pigmented Nevus; Garment Nevus; Bathing Trunk Nevus

### **Giant Coronary Aneurysm**

An unusually large bulge in an artery supplying the heart. It can form a clot, reduce blood flow, or rarely rupture.

### **Giant Cyst of the Retina**

A large fluid-filled retinal cystic lesion that can distort or damage the retina and impair vision. Evaluation typically requires a detailed eye examination and retinal imaging.

### **Giant Fibroadenoma**

An uncommon clinical variant of benign breast fibroadenoma exceeding 5 cm in diameter or 500 grams in weight, characterized by rapid encapsulated proliferation of epithelial and stromal elements, frequently occurring during adolescence or pregnancy.

#### **Alternate Names:**

Juvenile Giant Fibroadenoma

---

### Giant Hypertrophic Gastritis

Severe thickening of the stomach lining and its folds, which can cause stomach pain, nausea, swelling, and loss of blood proteins. It may increase stomach-cancer risk.

### Giant Lymph Node Hyperplasia

A rare condition in which lymph-node tissue grows excessively. It may affect one area or many parts of the body and can cause enlarged nodes, fever, fatigue, anemia, weight loss, or organ problems.

#### Alternate Names:

Castleman Disease; Angiofollicular Lymph Node Hyperplasia

### Giant Platelet Syndrome

An inherited platelet disorder caused by deficiency of glycoprotein Ib/IX/V complex, resulting in large platelets and impaired platelet adhesion. Presents with mucocutaneous bleeding, thrombocytopenia, and giant platelets on blood smear.

#### Alternate Names:

Bernard-Soulier Syndrome; Platelet Glycoprotein Ib Deficiency

### Giant Right Atrium

Marked enlargement of the heart's right upper chamber, usually from long-standing valve disease, a congenital heart defect, or abnormal rhythm. It can disturb the heartbeat, compress nearby structures, or allow clots to form and is assessed with heart imaging.

### Giant Urticaria

Alternate name; see *Urticaria*.

### Giardia

A microscopic parasite that infects the small intestine and spreads through contaminated water, food, or hands. It can cause foul-smelling diarrhea, cramps, bloating, nausea, weight loss, or no symptoms.

### Giardia Infection

An intestinal infection caused by *Giardia* parasites,

usually spread through contaminated water, food, or contact with an infected person. It can cause prolonged foul-smelling diarrhea, cramps, bloating, weight loss, and poor nutrient absorption.

### Giardia Lamblia

The parasite that causes giardiasis. It is swallowed in contaminated water, food, or material from an infected person and can cause foul-smelling diarrhea, abdominal cramps, gas, bloating, and weight loss.

### Giardiasis

An intestinal infection caused by the parasite *Giardia duodenalis*. It spreads when contaminated water, food, surfaces, or material from an infected person is swallowed. It may cause diarrhea, foul-smelling or greasy stools, gas, abdominal cramps, nausea, bloating, weight loss, and poor nutrient absorption.

### Gibbus Deformity

A sharp, angular, structural kyphosis of the spine produced by the anterior collapse or destruction of one or more adjacent vertebral bodies, classically seen in tuberculosis of the spine (Pott disease) or severe osteoporotic wedge fractures.

#### Alternate Names:

Pott Gibbus; Angular Kyphosis

### Gibraltar Fever

An old name for brucellosis, an infection caused by *Brucella* bacteria, usually acquired through unpasteurized dairy products or contact with infected animals.

### Giemsa Stain

A laboratory dye used to show blood cells and some bacteria, parasites, and other organisms under a microscope. It can help identify malaria parasites and other infections in blood or tissue samples.

### Gigabecquerel

A unit of radioactivity equal to one billion nuclear disintegrations per second. It measures the activity of a radioactive source, not directly the radiation dose absorbed by a person.

---

## Gigantism

Excessive growth caused by too much growth hormone during childhood, before the bones finish lengthening. It usually results from a pituitary growth and causes unusual height and enlargement of body tissues.

## Gigli Saw

A flexible wire saw used during some orthopedic, neurosurgical, or amputation procedures to cut bone. It is handled carefully to limit injury to nearby soft tissues.

## Gilbert Syndrome

A common inherited condition in which the liver processes bilirubin more slowly than usual. It can cause occasional mild yellowing of the eyes or skin, especially during fasting, illness, dehydration, or stress, and usually does not harm the liver.

## Gilbert's Disease

Alternate spelling; see *Gilbert Syndrome*.

## Gilbert's Syndrome

A common inherited condition in which the liver processes bilirubin slowly, causing occasional mild yellowing of the eyes, especially during illness, fasting, or stress.

## Gilbert-Lereboullet Syndrome

Alternate name; see *Gilbert Syndrome*.

## Gilles De La Tourette Syndrome

A neurological disorder causing repeated involuntary movements and sounds called motor and vocal tics. Symptoms begin in childhood and often change over time.

## Gingiva

The firm pink tissue, commonly called the gums, that surrounds and protects the teeth and the bone around their roots. Healthy gingiva usually does not bleed with gentle brushing; redness, swelling, or bleeding may indicate gum disease.

**Alternate Names:**

Gums; Gingival Tissue

## Gingiva Disease

A disorder of the gums, including plaque-induced gingivitis, periodontitis, infection, immune disease, recession, or tumor.

## Gingival Disease

Inflammation of the gums, usually caused by plaque buildup. The gums may become red, swollen, tender, or bleed during brushing. It usually does not damage the supporting bone and can often improve with daily cleaning and professional dental care.

**Alternate Names:**

Marginal Gingivitis; Periodontal Gum Inflammation

## Gingival Hemorrhage

Bleeding from gum tissue, commonly due to inflammation and plaque but also caused by trauma, periodontal disease, medicines, or bleeding disorders.

## Gingival Hyperplasia

An abnormal enlargement or overgrowth of gingival tissue around the teeth that can impair cleaning, cause bleeding, and increase periodontal risk. Causes include plaque-related inflammation and medicines such as phenytoin, cyclosporine, and some calcium-channel blockers; hereditary and systemic conditions can also contribute.

## Gingival Infections

Infections of the gums, usually involving bacteria in dental plaque. They can cause red, swollen, tender, or bleeding gums, bad breath, pus, and sometimes spreading facial swelling; dental assessment helps prevent tissue and bone loss.

## Gingival Margin

The edge of the gum tissue that surrounds and touches a tooth. It normally forms a shallow seal; inflammation or recession can make it bleed or expose the tooth root.

---

## Gingival Melanoma

A rare, aggressive cancer of pigment-producing cells in the gums. It may appear as a dark or non-pigmented lump, patch, or ulcer; biopsy can identify it.

## Gingival Neoplasm

An abnormal growth in the gums that may be reactive, benign, precancerous, or cancerous. It may cause persistent swelling, ulceration, bleeding, pain, or a lump; examination and sometimes biopsy identify its nature.

## Gingival Overgrowth

Gingival overgrowth is an abnormal increase in gum tissue, which may be caused by medicines, inflammation, or systemic disease.

## Gingival Pocket

The small space between the gum and tooth surface, measured during a dental examination. A persistently deep pocket usually indicates loss of attachment from periodontal disease and can collect plaque and bacteria.

## Gingival Recession

Pulling back of the gums so that part of a tooth root becomes exposed. Gum disease, forceful brushing, tooth position, thin gum tissue, or treatment can cause it and may lead to sensitivity or root decay.

## Gingivectomy

Surgical removal of gum tissue to treat overgrowth, reshape the gums, expose decay, or improve access. Possible complications include bleeding, sensitivity, infection, or gum recession.

## Gingivitis

Inflammation of the gingiva characterized by redness, swelling, and bleeding with brushing or probing, usually due to dental plaque. It is generally reversible with effective oral hygiene and professional cleaning; it can precede periodontitis, but does not inevitably progress to it.

## Gingivitis Transmission

Gingivitis itself is not directly contagious, but the bacteria that cause it (particularly *Porphyromonas gingivalis*) can be transmitted through saliva. Poor oral hygiene is the primary risk factor. Prevention focuses on personal oral care.

## Gingivoplasty

Surgical reshaping or contouring of the gums to improve their form, remove diseased tissue, or make cleaning easier. It may be combined with gingivectomy or other periodontal treatment.

## Gingivostomatitis

Common viral infection of the mouth and gums, most often caused by primary herpes simplex virus infection but also associated with other infections, trauma, or systemic disease. It may cause painful ulcers, swollen bleeding gums, fever, and difficulty eating or drinking.

## Ginkgo Biloba

An herbal product derived from *Ginkgo biloba* leaves and marketed for cognitive or circulatory benefits; evidence is variable and it may increase bleeding risk or interact with medicines.

## Ginseng

A group of plants whose roots are used in traditional medicine and dietary supplements; products vary in composition and may interact with medicines or affect blood glucose and clotting.

## Ginseng Preparation

A product made from ginseng root and sold as a supplement. Products vary in strength and purity; interactions and side effects can occur.

## Girdle Pain

Pain that wraps around the chest or abdomen in a band-like pattern. It may result from nerve irritation, spinal disease, shingles, muscle injury, or internal-organ disease.

---

### Gitelman Syndrome

A rare inherited kidney disorder that causes the body to lose too much salt and magnesium in urine. It may lead to muscle cramps, weakness, tiredness, thirst, or abnormal heart rhythms.

### Glabella

The smooth area of the forehead between the eyebrows and above the bridge of the nose.

### Gland

An organ or group of cells that produces and releases substances. Endocrine glands release hormones into blood, while exocrine glands release substances through ducts or onto a surface.

### Glanders

A serious infection caused by the bacterium *Burkholderia mallei*, which mainly affects horses and can rarely infect people exposed to infected animals or tissues. It may cause skin sores, fever, pneumonia, or bloodstream infection.

### Glandular Fever

Alternate name; see *Infectious Mononucleosis*.

### Glandular Plague

Alternate name; see *Bubonic Plague*.

### Glans Penis

The expanded, sensitive distal end of the penis containing the external urethral opening and covered by the foreskin when present.

### Glanzmann Thrombasthenia

An inherited platelet function disorder caused by deficiency or dysfunction of platelet glycoprotein IIb/IIIa, producing mucocutaneous bleeding despite a normal platelet count.

### Glasgow Coma Scale

A bedside score describing a person's level of consciousness by assessing eye opening, speaking, and movement. Scores range from 3 to 15; lower scores indicate less response, but medicines and injuries

can affect the result.

### Glass Eye

A glass eye is an ocular prosthesis made to replace the appearance of an eye that is missing or severely damaged.

### Glass Ionomer

A tooth-colored dental material that bonds to tooth structure and slowly releases fluoride. It is used for selected fillings, repairs, protective bases, and children's teeth; durability depends on the location and amount of stress.

### Glassy Cell Carcinoma

A rare poorly differentiated adenosquamous carcinoma of the cervix with distinctive glassy cytoplasm, prominent nucleoli, and aggressive clinical behavior.

### Glaucoma

A group of eye diseases in which the optic nerve is damaged, often in association with high pressure inside the eye. Types include open-angle, angle-closure, normal-tension, secondary, and congenital glaucoma. It can cause permanent loss of side or central vision, while early disease may have no symptoms.

### Glaucoma Drainage Implant

A surgically implanted tube-and-plate device that diverts aqueous humor from the anterior chamber to a subconjunctival reservoir, lowering intraocular pressure in selected glaucomas.

### Glaucoma Surgery

Laser or surgical procedures used to lower pressure inside the eye by improving fluid drainage or, less often, reducing fluid production. They may be considered when medicines do not adequately control pressure or are unsuitable; they can slow further optic-nerve damage but cannot restore vision already lost.

### Glaze Poisoning

Toxic exposure to ceramic or pottery glaze, which



---

may contain lead, cadmium, or other heavy metals. Symptoms depend on the metal involved and may include abdominal pain, neurological symptoms, and organ damage. Severe cases may be treated with chelation therapy.

**Alternate Names:**

Ceramic Glaze Poisoning; Lead or Cadmium Glaze Toxicity

**Gleason Grading System**

A histological grading system for prostate cancer based on glandular pattern differentiation. Scores historically ranged from 2 to 10, but current reporting summarizes them as Grade Groups 1 to 5 (scores 6 to 10), with higher groups indicating more aggressive cancer. Used to guide treatment decisions and predict prognosis.

**Alternate Names:**

Gleason Score; Histological Prostate Grade

**Gleason Score**

A grading system estimating prostate cancer aggressiveness by examining tissue patterns under a microscope and combining the two most common patterns.

**Alternate Names:**

Gleason Grade Group; Prostate Adenocarcinoma Score

**Gleason Tumor Grade**

A histopathologic grade of prostate adenocarcinoma based on the architectural pattern of the tumor glands; it contributes to the Gleason score and Grade Group.

**Glenn Procedure**

A staged heart operation for some children born with only one effective pumping chamber. It connects the large vein returning blood from the upper body directly to the lung arteries, allowing blood to pick up oxygen while reducing the workload on the heart.

**Alternate Names:**

Bidirectional Glenn (BDG) Procedure; Cavopulmonary Anastomosis

**Glenohumeral Joint**

The ball-and-socket shoulder joint formed by the upper-arm bone and shoulder blade. It allows the arm to move in many directions but is relatively easy to dislocate because it sacrifices some stability for movement.

**Glenoid**

The shallow socket on the shoulder blade that joins the head of the upper-arm bone to form the shoulder joint. Its shape and surrounding labrum help balance mobility with stability.

**Alternate Names:**

Glenoid Cavity; Scapular Glenoid Fossa

**Glenoid Cavity**

The shallow socket on the shoulder blade that receives the rounded head of the upper-arm bone. A rim of tough tissue deepens the socket, but the joint's wide range of movement also makes it prone to dislocation.

**Glenoid Labrum**

A fibrocartilaginous rim attached to the peripheral margin of the glenoid fossa of the scapula, which deepens the shallow glenoid cavity to enhance glenohumeral joint stability and serves as the superior anchor for the long head of the biceps brachii tendon.

**Alternate Names:**

Glenoid Ligament

**Gliadin**

A group of wheat storage proteins that, together with glutenin, form gluten; gliadin peptides trigger the immune response in celiac disease.

**Glial Cell**

A non-neuronal cell that supports, nourishes, insulates, and regulates neurons; examples include astrocytes, oligodendrocytes, microglia, and Schwann cells.

**Glial Cells**

Support cells of the brain and spinal cord that nourish neurons.

---

ish and protect nerve cells, form insulation, maintain the chemical environment, and help repair tissue. Some brain tumors arise from glial cells.

### **Glial Fibrillary Acidic Protein**

A protein found mainly in support cells of the brain and nervous system. Laboratory tests can use it to help identify some brain tumors or detect injury to support cells after brain trauma.

#### **Alternate Names:**

GFAP

### **Glibenclamide**

An oral sulfonylurea medicine, also called glyburide, used for type 2 diabetes. It stimulates the pancreas to release insulin and can cause prolonged, sometimes severe low blood sugar and weight gain.

### **Gliclazide**

Gliclazide is a sulfonylurea for type 2 diabetes that stimulates pancreatic beta-cell insulin release; hypoglycaemia, weight gain, and dose accumulation in organ impairment are important considerations.

### **GLIM Criteria**

An international framework for diagnosing malnutrition. It considers body changes such as weight or muscle loss together with causes such as poor intake, poor absorption, or illness-related inflammation and grades severity according to the findings.

### **Glimepiride**

An oral sulfonylurea medicine for type 2 diabetes that increases insulin release from functioning pancreatic beta cells. Low blood sugar is the main serious risk, particularly with missed meals, kidney disease, or other glucose-lowering medicines.

### **Glioblastoma**

A fast-growing cancerous tumor of the central nervous system. It forms from glial tissue, which supports nerve cells, and usually develops in the brain of an adult; it can also occur in the spinal cord. The tumor can grow into nearby healthy tissue. Symptoms depend on its location and may include headache, seizures, weakness, changes in speech,

memory, vision, balance, or behaviour. Also called glioblastoma multiforme (GBM).

### **Glioblastoma Multiforme**

An aggressive cancer arising from supporting cells of the brain or spinal cord. It may cause headache, seizures, personality or memory changes, weakness, or speech problems. Treatment often combines surgery, radiation, and medicines.

### **Glioma**

A tumor arising from supporting cells of the brain or spinal cord. Some gliomas grow slowly, while others are aggressive; symptoms and treatment depend on the tumor's type, location, molecular features, and spread.

### **Glioma-Polyposis Syndrome**

Alternate name; see *Turcot Syndrome*.

### **Gliomatosis Cerebri**

A diffuse infiltrative growth pattern of glioma involving multiple brain regions, rather than a separate tumor type in current classifications; it may cause progressive cognitive, motor, or seizure-related symptoms.

### **Gliosarcoma**

A rare aggressive brain cancer containing both glial and connective-tissue-like tumor features. Symptoms depend on location and may include headache, seizures, weakness, or cognitive changes; tissue examination identifies the tumor type.

### **Gliososis**

A reactive increase in glial cells after injury, infection, ischemia, or other damage to the central nervous system.

### **Glipizide**

Glipizide is a short-acting sulfonylurea that lowers blood glucose by stimulating insulin secretion from functioning pancreatic beta cells; hypoglycaemia is the principal adverse effect.

---

### Glisson's Capsule

A thin layer of connective tissue surrounding the liver and extending between its sections. It contains nerves and blood vessels and can cause pain when the liver becomes swollen or inflamed.

### Global Aphasia

A severe language disorder caused by extensive damage to the dominant cerebral hemisphere, impairing speech production, comprehension, reading, and writing.

### Global Developmental Delay

Noticeably delayed development in at least two areas, such as movement, speech, learning, social skills, or daily activities, in a child younger than five years. Causes vary and may include genetic, neurologic, hearing, vision, or environmental factors; assessment guides support and further testing.

#### Alternate Names:

GDD; Neurodevelopmental Delay

### Global Health

The study and practice of improving health across countries and populations. It includes problems that cross borders, such as outbreaks, unequal access to care, pollution, climate change, malnutrition, and differences in life expectancy.

### Global Longitudinal Strain

An echocardiogram measurement showing how much the heart muscle shortens lengthwise as it contracts. It can detect early weakening of the left ventricle before the usual pumping measurement changes and is interpreted with the complete heart scan and clinical findings.

#### Alternate Names:

GLS; 2D Speckle-Tracking GLS; Left Ventricular Longitudinal Strain

### Globe Rupture

A full-thickness disruption of the outer wall of the eye caused by blunt or penetrating trauma. It may result in deformity, pain, hemorrhage, loss of vision, and extrusion or prolapse of intraocular contents.

### Globin

A protein that joins with heme to form hemoglobin and related oxygen-carrying molecules. Different globin chains help determine the type of hemoglobin made at different stages of life.

### Globoid Cell Leukoencephalopathy

Alternate name; see *Krabbe Disease*.

### Globozoospermia

A rare sperm disorder in which sperm heads are round and lack a normal acrosome, causing impaired fertilization and male infertility.

### Globulin

A group of blood proteins that includes antibodies and proteins involved in transport, inflammation, and clotting. Blood tests may divide globulins into several groups; an abnormal level can occur with infection, immune disease, liver disease, kidney disease, or some cancers.

### Globus

A persistent or intermittent sensation of a lump or tightness in the throat without a true obstructing mass, often called globus pharyngeus.

### Globus Hystericus

A sensation of a lump or tightness in the throat without a structural obstruction, now called globus sensation; reflux, muscle tension, and anxiety may contribute.

### Globus Pallidus

The globus pallidus is a group of deep brain structures involved in regulating movement as part of the basal ganglia.

### Globus Sensation

The persistent or intermittent sensation of a lump or foreign body in the throat without an anatomic lesion or dysphagia, also termed globus pharyngeus, typically noticed between meals. There is no impedance to swallowing and no weight loss; the feeling worsens with stress, anxiety, or laryngophary-

---

ryngeal reflux.

### **Glomangiomyoma**

A benign glomus tumor showing both vascular and smooth-muscle differentiation, usually presenting as a small painful subcutaneous nodule.

### **Glomangiosarcoma**

A rare cancerous glomus tumor with abnormal cells and invasive growth. It may occur in the skin or deeper soft tissue.

### **Glomerular Basement Membrane**

The thick extracellular matrix layer of the renal filtration barrier interposed between glomerular endothelial cells and podocyte foot processes, composed of type IV collagen, laminin, and heparan sulfate proteoglycans that impart mechanical structural support and negative charge selectivity.

#### **Alternate Names:**

GBM

### **Glomerular Capsule**

A cup-shaped structure around a tiny group of blood vessels in the kidney. It collects the fluid filtered from blood and passes it into a kidney tubule, where useful water and substances are reclaimed and waste is removed.

### **Glomerular Filtrate**

The watery fluid filtered from blood into the kidney tubules by a glomerulus, a small group of kidney blood vessels. It normally contains water and small dissolved substances but very little protein or no blood cells.

### **Glomerular Filtration**

The first step in urine formation, in which pressure moves water and small dissolved substances from blood through kidney filters into the tubules. Blood cells and most proteins normally remain in the bloodstream.

### **Glomerular Filtration Rate**

An estimate of how much blood fluid the kidneys

filter each minute. It is calculated mainly from blood creatinine and other factors; a persistently low result can indicate chronic kidney disease, but the result is interpreted with other findings.

### **Glomerular Hyperfiltration**

An abnormally elevated single-nephron or whole-kidney glomerular filtration rate exceeding physiological upper limits, representing an early hemodynamic adaptive response in early diabetic kidney disease, obesity, and after unilateral nephrectomy that eventually accelerates glomerulosclerosis.

### **Glomeruli**

Microscopic clusters of capillaries in the kidneys where blood filtration begins. Each glomerulus is surrounded by a capsule that collects filtered fluid and passes it into a kidney tubule.

#### **Alternate Names:**

Renal Glomeruli; Malpighian Corpuscles

### **Glomerulomegaly**

Abnormal enlargement of kidney glomeruli, seen with diabetes, obesity-related hyperfiltration, congenital syndromes, or other causes of glomerular stress.

### **Glomerulonephritis**

Inflammation or other immune-related injury of the kidney's filtering units. It can cause blood or protein in urine, swelling, high blood pressure, and reduced kidney function and may follow an infection or occur with an autoimmune disease.

#### **Alternate Names:**

GN; Glomerular Nephritis

### **Glomerulonephritis Acute**

Sudden inflammation of the kidney glomeruli that can cause blood or protein in urine, edema, hypertension, and reduced kidney function.

### **Glomerulopathy**

Any disease affecting the kidney's filtering units. It may cause blood or protein in urine, swelling, high blood pressure, reduced filtration, or kidney failure

---

and may result from diabetes, infection, immune disease, inherited disorders, or other injury.

### Glomerulosclerosis

Scarring and hardening of the kidney's filtering units. It can reduce filtration and cause protein in urine, swelling, high blood pressure, or kidney failure; causes include diabetes, inflammation, genetic disease, and other injury.

### Glomerulus

A tuft of tiny blood vessels inside a kidney capsule where water and small dissolved substances are filtered from blood to begin urine formation. Damage can allow blood or excess protein into urine and reduce kidney function.

### Glomus Jugulare Tumor

A usually noncancerous but locally invasive growth arising near the jugular vein at the skull base. It may cause pulsating noise in the ear, hearing loss, dizziness, or weakness of nearby nerves.

#### Alternate Names:

Jugular Paraganglioma; Tympanojugular Glomus

### Glomus Tumor

A usually harmless growth arising from cells that help control blood flow and temperature, most often beneath a fingernail or in a finger. It typically causes severe pinpoint pain and sensitivity to cold or pressure.

### Glomus Tumors

Usually benign tumors of glomus cells involved in temperature regulation and blood flow, commonly beneath a fingernail or in a fingertip. They characteristically cause severe localized pain and cold sensitivity; excision is often curative.

### Glomus Tympanicum

A benign vascular tumor arising from the glomus bodies on the promontory of the middle ear, presenting with pulsatile tinnitus, hearing loss, and a reddish mass behind the tympanic membrane.

### Glomus Tympanicum Tumor

A paraganglioma arising from the glomus bodies on the promontory of the middle ear. Benign tumor causing pulsatile tinnitus and conductive hearing loss. Diagnosis by otoscopy and imaging. Treatment includes surgery or observation.

#### Alternate Names:

Middle Ear Paraganglioma; Tympanic Glomus

### Glossal

Relating to the tongue, the muscular organ used for chewing, swallowing, tasting, and speaking.

### Glossalgia

Pain or burning of the tongue, with causes including trauma, infection, nutritional deficiency, neuropathy, medication, or burning-mouth syndrome.

### Glossectomy

Surgical removal of all or part of the tongue, performed for malignancies, severe trauma, or an unusually large tongue. Procedures range from partial to total removal. Speech and swallowing rehabilitation may follow.

### Glossinidae

The family of flies belonging to the genus *Glossina*, commonly known as tsetse flies. These insects are vectors of trypanosomes that cause African trypanosomiasis (sleeping sickness) in humans and nagana in animals.

#### Alternate Names:

Tsetse Flies; *Glossina* Vectors

### Glossitis

Inflammation of the tongue, which can cause pain, swelling, and changes in tongue appearance. Causes include nutritional deficiencies (iron, B12, folate), infections (bacterial, viral, fungal), allergic reactions, and autoimmune conditions. Treatment depends on the underlying cause.

### Glossitis Areata Migrans

Alternate name; see *Geographic Tongue*.

---

## Glossodynia

Persistent burning, soreness, or tingling in the tongue, lips, palate, or other mouth tissues, often without visible changes. It may occur with dry mouth, altered taste, nutritional deficiency, medicines, nerve problems, or other illness.

### Alternate Names:

Burning Mouth Syndrome; Orofacial Dysesthesia

## Glossopharyngeal Nerve

The ninth cranial nerve, carrying sensation and taste from the back of the tongue and throat, signals from blood-pressure and blood-chemistry sensors, and motor control for one throat muscle. It also stimulates saliva production.

## Glossopharyngeal Neuralgia

Recurrent, brief attacks of severe stabbing or electric-shock-like pain in the throat, tonsillar region, base of the tongue, or ear, caused by irritation of the glossopharyngeal nerve. Swallowing, coughing, or talking may trigger attacks.

## Glossoptosis

Backward displacement of the tongue that can narrow the upper airway, particularly in infants with craniofacial disorders.

## Glossopyrosis

Alternate name; see *Burning Mouth Syndrome*.

## Glottis

The part of the voice box containing the vocal folds and the opening between them. It opens for breathing, closes to protect the airway during swallowing, and vibrates to produce the voice.

## GLP-1 Agonists

Medicines that mimic a natural gut hormone. They help lower blood glucose after meals, slow stomach emptying, and reduce appetite. They are used for selected people with diabetes or obesity and commonly cause nausea, vomiting, diarrhea, or constipation.

## GLP-1 Receptor Agonists

A class of medicines that activate the glucagon-like peptide-1 receptor. They increase glucose-dependent insulin release, reduce inappropriate glucagon release, slow gastric emptying, and increase fullness. They are used for selected people with type 2 diabetes or chronic weight management. Examples include semaglutide, liraglutide, and dulaglutide.

### Alternate Names:

GLP-1 Agonists; Incretin Mimetics

## GLS

Alternate name; see *Global Longitudinal Strain*.

## Glu

Commonly an abbreviation for glucose in laboratory or clinical notes; in biochemical notation, Glu can also denote the amino acid glutamic acid, so context matters.

## Glucagon

A hormone made by pancreatic alpha cells that raises blood glucose when it falls too low. It prompts the liver to release stored glucose and can be given for severe hypoglycemia.

## Glucagon Blood Test

Laboratory measurement of glucagon levels in the blood, used to evaluate suspected glucagonoma, hypoglycemia, or other pancreatic endocrine disorders. Levels are affected by fasting, meals, and medications.

### Alternate Names:

Serum Glucagon Level; Plasma Glucagon Assay

## Glucagon Signaling

The chain of cell signals started when glucagon binds its receptor. These signals increase cyclic AMP and activate proteins that help the liver release glucose and the body maintain energy balance.

## Glucagon-like Peptide-1

A hormone released mainly by the intestine after eating. It helps regulate blood glucose by enhanc-



---

ing glucose-dependent insulin secretion, reducing glucagon release, slowing stomach emptying, and influencing appetite.

### Glucagonoma

A rare pancreatic neuroendocrine tumor that secretes excess glucagon, causing diabetes mellitus (mild, new onset), dermatitis (necrolytic migratory erythema, characteristic painful, erythematous, blistering rash on groin, perineum, extremities), diarrhea, and deep vein thrombosis.

### Glucan

A complex carbohydrate made of many linked glucose molecules. Different glucans occur in plants, fungi, bacteria, and foods and may act as dietary fiber or interact with the immune system.

### Glucarpidase

An enzyme that rapidly breaks down methotrexate in the bloodstream and is used for toxic methotrexate concentrations when renal clearance is inadequate.

### Glucocerebrosidase Deficiency

Gaucher Disease: an inherited lysosomal storage disorder caused by deficient glucocerebrosidase, leading to lipid accumulation in cells and possible enlargement of the spleen and liver, anemia, bone disease, or neurologic disease.

#### Alternate Names:

Gaucher Disease; Acid Beta-Glucosidase Deficiency

### Glucocorticoid

A corticosteroid hormone or medicine that regulates inflammation, immunity, metabolism, and stress responses; prolonged treatment has systemic adverse effects.

### Glucocorticoid Remediable Aldosteronism

An autosomal dominant cause of secondary endocrine hypertension caused by an unequal crossover mutation that fuses the ACTH-responsive promoter of the 11-beta-hydroxylase gene (*CYP11B1*) to the aldosterone synthase coding sequence

(*CYP11B2*), rendering aldosterone secretion suppressible by dexamethasone.

#### Alternate Names:

GRA; Familial Hyperaldosteronism Type I

### Glucocorticoid Replacement Therapy

Treatment of adrenal insufficiency with a glucocorticoid medicine to replace the cortisol the body lacks. Doses may vary with symptoms, illness, and physical stress.

### Glucocorticoid-Induced Osteoporosis

Bone loss and increased fracture risk caused or accelerated by prolonged or high-dose glucocorticoid treatment; risk depends on dose, duration, baseline bone health, and other factors.

### Glucocorticoid-Remediable Aldosteronism

An autosomal dominant form of secondary hypertension caused by an unequal crossover genetic recombination fusing the promoter of the 11 $\beta$ -hydroxylase gene (*CYP11B1*) to the coding region of the aldosterone synthase gene (*CYP11B2*). This places aldosterone production under the ectopic regulatory control of adrenocorticotrophic hormone (ACTH), resulting in hyperaldosteronism, hypokalemia, and early-onset hypertension that responds promptly to exogenous glucocorticoid suppression.

#### Alternate Names:

GRA; Familial Hyperaldosteronism Type I

### Glucocorticoid-Sparing Therapy

Use of a non-glucocorticoid treatment strategy to control inflammation while reducing cumulative corticosteroid exposure. It aims to limit osteoporosis, infection, diabetes, hypertension, adrenal suppression, and other glucocorticoid complications.

### Glucocorticoids

Steroid hormones made by the adrenal glands, along with medicines that act in the same way. They reduce inflammation and immune activity but can cause infection, high blood glucose, bone loss, or adrenal suppression with prolonged use.

---

## **Glucometer**

A handheld device that measures blood glucose, usually from a small capillary-blood sample obtained by finger prick.

## **Gluconates**

Salts or esters of gluconic acid used in some medicines and supplements. Examples include calcium gluconate and ferrous gluconate; their effects, dosing, and risks depend on the specific compound.

## **Gluconeogenesis**

A metabolic pathway that generates glucose from non-carbohydrate precursors such as lactate, glycerol, and glucogenic amino acids. It occurs mainly in the liver and, to a variable extent, in the kidneys and other tissues.

## **Glucosamine**

A naturally occurring compound used in cartilage formation and sometimes taken as a supplement for joint symptoms.

## **Glucosamine and Chondroitin**

Dietary supplements marketed for joint symptoms and cartilage support. Evidence of benefit varies, and products may interact with medicines or affect blood sugar or bleeding risk.

## **Glucose**

A simple sugar that is the main energy source for the body's cells. It comes from digesting carbohydrates and can also be released from stored energy or made by the liver. Hormones, especially insulin, help keep blood glucose within a healthy range.

## **Glucose Challenge Test**

A screening test for gestational diabetes mellitus, commonly using a nonfasting 50-g glucose drink followed by a one-hour plasma-glucose measurement. An abnormal screening result is followed by a diagnostic oral glucose tolerance test; the threshold depends on the protocol used.

### **Alternate Names:**

GCT; One-Hour Glucose Challenge; O'Sullivan

### Screening Test

## **Glucose Intolerance**

Impaired handling of glucose resulting in blood-glucose concentrations above normal but below the diagnostic threshold for diabetes in a defined testing context. It often reflects insulin resistance or inadequate insulin secretion and is associated with increased risk of type 2 diabetes and cardiovascular disease.

## **Glucose Management Indicator**

An estimate of the hemoglobin A1c value calculated from average continuous-glucose-monitoring readings. It helps summarize recent glucose exposure but is not the same as a laboratory A1c and may differ from it.

## **Glucose Screening Tests During Pregnancy**

Laboratory tests to detect gestational diabetes mellitus, including the oral glucose tolerance test and glucose challenge test. They are usually performed between 24 and 28 weeks of gestation; abnormal results are followed by additional testing or management.

### **Alternate Names:**

Gestational Diabetes Screening; Maternal Glucose Screen

## **Glucose Sensor**

A small sensor placed under the skin that measures glucose in the fluid between cells. Used in continuous glucose monitoring, it sends repeated readings to a receiver or compatible device and can show trends and alerts.

## **Glucose Test**

A laboratory or point-of-care test that measures glucose in blood or another sample. It is used to screen for, diagnose, or monitor diabetes and low or high blood sugar; interpretation depends on timing, fasting status, and clinical context.

## **Glucose Tolerance Test**

A test of how the body handles a measured glu-

---

cose drink. Blood samples are taken before and at set times afterward to help diagnose diabetes or gestational diabetes; thresholds depend on the protocol.

**Alternate Names:**

Oral Glucose Tolerance Test (OGTT); 3-Hour Diagnostic OGTT

### Glucose Transport

Movement of glucose across cell membranes through specialized transport proteins. Insulin increases glucose entry into muscle and fat cells, while other transporters allow glucose to enter the brain, liver, kidneys, and intestines.

### Glucose Transporter

A membrane protein that moves glucose across a cell membrane; transporter defects or altered regulation can affect glucose metabolism.

### Glucose Urine Test

Laboratory test to detect glucose in the urine (glycosuria). Normally absent; presence indicates blood glucose exceeding renal threshold, suggesting diabetes mellitus or renal tubular dysfunction. May be detected by dipstick or laboratory analysis.

**Alternate Names:**

Urinary Glucose; Glycosuria Screen

### Glucose-6-Phosphatase

An enzyme in the liver and kidneys that releases glucose from glucose-6-phosphate. Inherited deficiency causes glycogen-storage disease type I, which can produce low blood glucose, enlarged liver, and metabolic problems.

### Glucose-6-Phosphate Dehydrogenase Deficiency

An X-linked red-blood-cell enzyme disorder. Infections, fava beans, or certain medicines can cause oxidative damage and sudden destruction of red blood cells, leading to hemolytic anemia, jaundice, dark urine, fatigue, or shortness of breath.

### Glucose-6-Phosphate Dehydrogenase Test

Laboratory test to measure G6PD enzyme activity in red blood cells, used to diagnose G6PD deficiency. Important before prescribing certain medications (primaquine, dapsone) that can cause hemolytic anemia in deficient individuals.

**Alternate Names:**

G6PD Assay; Erythrocyte Enzyme Screen

### Glucosuria

The presence of glucose in urine, most commonly from blood glucose exceeding the renal reabsorption threshold, diabetes, pregnancy, or a proximal tubular reabsorption defect.

### Glucosyl Ceramide Lipidosis

Alternate name; see *Gaucher Disease*.

### Glucosylceramidase

A lysosomal enzyme that breaks down glucosylceramide, a fat-containing cell substance. Inherited deficiency causes Gaucher disease, allowing fatty material to accumulate in the spleen, liver, bone marrow, bones, or nervous system.

### Glucuronate

The anion or salt of glucuronic acid, a sugar-acid component of connective-tissue substances and a molecule used in hepatic conjugation reactions.

### Glucuronic Acid

A sugar acid used by the liver to conjugate bilirubin, drugs, hormones, and toxins, increasing their water solubility for excretion.

### Glucuronide

A compound formed when glucuronic acid is conjugated to a drug, hormone, bilirubin, or other molecule, usually increasing water solubility and facilitating excretion.

### Glucuronyl Transferase

UDP-glucuronosyltransferase: a liver enzyme involved in phase II drug metabolism, catalyzing glucuronidation of bilirubin, drugs, and toxins to make

---

them water-soluble for excretion. Deficiency causes Gilbert syndrome or Crigler-Najjar syndrome.

**Alternate Names:**

UDP-Glucuronosyltransferase; UGT Enzyme

### Glue Ear

Persistent fluid in the middle ear without acute infection, often causing conductive hearing loss and a feeling of blockage. It is common in children.

**Alternate Names:**

Secretory Otitis Media; Otitis Media with Effusion (OME)

### GLUT4 Transporter

The major insulin- and exercise-regulated glucose transporter isoform expressed primarily in striated muscle (skeletal and cardiac) and adipose tissue, which translocates from intracellular vesicles to the plasma membrane upon insulin stimulation to facilitate cellular glucose uptake.

### Glutamate

The main excitatory chemical messenger in the brain and spinal cord. It helps nerve cells communicate, learn, and form memories; excessive glutamate activity can contribute to nerve-cell injury.

### Glutamate Decarboxylase

Glutamate decarboxylase is an enzyme that converts glutamate into the neurotransmitter GABA; antibodies against it are associated with some autoimmune disorders.

### Glutamate Receptor

A neuronal receptor activated by glutamate, the major excitatory neurotransmitter, including NMDA, AMPA, kainate, and metabotropic receptor classes.

### Glutamic Acid

An amino acid that acts as a metabolic intermediate and neurotransmitter precursor; its charged form, glutamate, mediates most fast excitatory signaling in the brain.

### Glutamic Acid Decarboxylase

An enzyme that converts glutamate to GABA; antibodies to it can occur in type 1 diabetes.

**Alternate Names:**

GAD; Glutamate Decarboxylase

### Glutamine

An amino acid used to build proteins, transport nitrogen, and provide energy for some cells, especially cells in the intestine and immune system. The body usually makes enough, but needs may change during severe illness.

### Glutaral

Glutaral is a chemical disinfectant and tissue fixative that can irritate the skin, eyes, and airways.

### Glutaric Acidemia Type 1

A rare inherited disorder in which the body cannot break down certain amino acids, allowing harmful substances to build up. Babies may have a large head and can develop sudden brain injury, abnormal movements, or stiffness during illness; newborn screening and dietary treatment can reduce harm.

**Alternate Names:**

Glutaric Aciduria Type 1 (GA-1); Glutaryl-CoA Dehydrogenase Deficiency

### Glutaric Aciduria

An inherited metabolic disorder in which the body cannot properly break down certain amino acids. Type I can cause brain injury, movement problems, and enlarged head size.

### Glutathione

A small antioxidant molecule made from glutamate, cysteine, and glycine. Reduced glutathione helps control oxidative stress and helps the liver neutralize reactive chemicals, including NAPQI produced from acetaminophen. N-acetylcysteine can restore glutathione after acetaminophen overdose.

### Glutathione Disulfide

The oxidized form of glutathione, produced after glutathione neutralizes reactive oxygen compounds.

---

Cells normally convert it back to active glutathione to maintain antioxidant protection.

### Glutathione Peroxidase

A selenium-dependent antioxidant enzyme family that catalyzes the reduction of toxic hydrogen peroxide and organic lipid hydroperoxides to water and corresponding alcohols, utilizing reduced glutathione (GSH) as an essential electron donor to protect cell membranes from oxidative injury.

**Alternate Names:**

GPx

### Glutathione Synthetase Deficiency

An inherited disorder caused by deficient glutathione synthetase, the enzyme needed to make glutathione. Severity ranges from hemolytic anemia and excess 5-oxoproline to metabolic acidosis, neurologic impairment, recurrent infection, or seizures.

### Glutathione Transferase

A family of enzymes that attaches glutathione to selected medicines, toxins, and products of cell metabolism. This usually makes them less reactive and easier for the body to process or remove.

### Gluteal

Relating to the buttock or the gluteal muscles, which extend and rotate the hip and help stabilize the pelvis.

### Gluteal Gait

An abnormal walking gait caused by weakness or paralysis of the hip abductor muscles (gluteus medius and minimus) innervated by the superior gluteal nerve, manifested by a downward pelvic tilt toward the unsupported swing leg during single-leg stance, compensated by leaning the trunk toward the affected stance side.

**Alternate Names:**

Trendelenburg Gait; Gluteus Medius Gait

### Gluteal Injury

Damage to the muscles, tendons, nerves, or soft tissues of the buttock caused by trauma, overuse,

injection, or falls. Pain, bruising, weakness, difficulty walking, severe swelling, numbness, inability to bear weight, or bowel or bladder changes may occur.

### Gluteal Muscle

One of the large muscles of the buttock, especially the gluteus maximus, medius, or minimus. These muscles extend, rotate, and move the hip and help stabilize the pelvis during standing and walking.

### Gluteal Tendinopathy

Tendon degeneration, microtrauma, and tearing affecting the insertion of the gluteus medius or gluteus minimus tendons onto the greater trochanter of the femur, representing the predominant cause of greater trochanteric pain syndrome in middle-aged and older adults.

### Gluten

A complex of storage proteins (prolamins and glutelins) found in wheat (gliadin), barley (hordein), and rye (secalin). Gluten provides elasticity to dough. In susceptible individuals, gluten triggers an autoimmune response (celiac disease) or non-autoimmune sensitivity (non-celiac gluten sensitivity, NCGS).

### Gluten Enteropathy

Celiac disease, an immune reaction to gluten that damages the small intestine and can cause diarrhea, weight loss, anemia, fatigue, or poor nutrient absorption.

**Alternate Names:**

Celiac Disease; Celiac Sprue

### Gluten Intolerance

Alternate name; see *Non-Celiac Gluten Sensitivity*.

### Gluten Sensitivity

Alternate name; see *Nonceliac gluten sensitivity*.

### Gluten-Free Diet

A diet that avoids wheat, barley, rye, and foods contaminated with them. It is essential for celiac

---

disease and may help selected people with non-celiac gluten sensitivity; poorly planned diets can lack fiber or nutrients.

### **Gluten-Sensitive Enteropathy**

Alternate name; see *Celiac Disease*.

### **Gluteus**

Relating to the buttock muscles. The gluteus maximus, medius, and minimus move and stabilize the hip and pelvis during standing, walking, climbing, and other movements.

### **Gluteus Maximus Muscle**

Largest, most superficial gluteal muscle forming the buttock bulk, from posterior iliac gluteal line, sacrum, coccyx, and sacrotuberous ligament to the iliotibial tract and gluteal tuberosity. Innervated by inferior gluteal nerve (L5-S2). Primary hip extensor and lateral rotator.

### **Gluteus Medius Muscle**

Thick fan-shaped muscle between maximus and minimus, from external ilium to the greater trochanter. Innervated by superior gluteal nerve (L4-S1). Primary hip abductor essential for pelvic stability during single-leg stance. Superior gluteal nerve injury causes Trendelenburg sign with contralateral pelvic drop.

### **Gluteus Minimus Muscle**

The smallest and deepest gluteal muscle, extending from the outer ilium to the greater trochanter. It abducts and medially rotates the hip and helps stabilize the pelvis during walking.

### **Glyburide**

An oral sulfonylurea that stimulates pancreatic insulin release for type 2 diabetes; prolonged or severe hypoglycemia is an important risk.

### **Glycaemia**

The amount of glucose in the blood. Insulin lowers blood glucose, while glucagon and other hormones raise it when needed. Abnormally high or low levels

can cause symptoms and, if persistent, harm organs.

### **Glycated Hemoglobin**

A blood test showing the average blood glucose level over roughly the previous two to three months. It is used to diagnose diabetes and monitor treatment, but results may be misleading when red blood cell survival is abnormal.

#### **Alternate Names:**

Hemoglobin A1c; HbA1c; Glycohemoglobin

### **Glycation**

A chemical process in which sugar attaches to proteins, fats, or genetic material without the usual enzyme control. Excess glycation during prolonged high blood glucose can damage blood vessels, nerves, kidneys, eyes, and other tissues.

### **Glycemia**

The amount of glucose in the blood. It is affected by food, activity, illness, the liver, and hormones such as insulin and glucagon. Too little is hypoglycemia; too much is hyperglycemia.

### **Glycemic Control**

The level and pattern of blood-glucose management over time. It includes glucose values, symptoms, and the effects of food, activity, illness, medicines, pregnancy, and other circumstances.

### **Glycemic Index**

A ranking of carbohydrate-containing foods according to how quickly they raise blood glucose compared with a reference food. It does not account for the amount of carbohydrate eaten in a serving.

### **Glycemic Index And Diabetes**

The glycemic index ranks carbohydrate-containing foods by their effect on blood glucose levels. Low GI foods ( $\leq 55$ ) cause gradual glucose rise; high GI foods ( $\geq 70$ ) cause rapid spikes. Important for diabetes management and dietary planning.

#### **Alternate Names:**

GI; Glycemic Load Assessment



---

## Glycemic Load

A measure that combines a food's glycemic index with the amount of available carbohydrate in a serving to estimate its effect on blood glucose.

## Glyceraldehyde

A small sugar with three carbon atoms that appears as the body breaks down glucose and other carbohydrates. It is quickly changed into other substances during the steps that release energy from food.

## Glyceride

A glyceride is a compound formed when glycerol is linked to one or more fatty acids, including many dietary fats.

## Glycerin

A colorless, odorless, sweet-tasting liquid also known as glycerol. It attracts water, dissolves some substances, and is used in medicines, skin products, foods, and selected treatments.

## Glycerol

A colorless, sweet-tasting alcohol that forms part of fats. It is used in medicines, foods, and skin products and can draw water into the bowel when used in some laxatives.

### Alternate Names:

Glycerine; Trihydroxypropane

## Glycerol Kinase

Glycerol kinase is an enzyme that adds phosphate to glycerol, allowing it to enter pathways for energy production and fat metabolism.

## Glycerophosphates

Compounds containing glycerol and phosphate that help build cell membranes and participate in energy metabolism, chemical signaling, and other cellular processes.

## Glycerophospholipids

The major membrane phospholipids built from glycerol, fatty acids, phosphate, and a polar head group, including phosphatidylcholine and phos-

phatidylethanolamine.

## Glyceryl Trinitrate

A nitrate medicine, also called nitroglycerin, that releases nitric oxide and relaxes blood vessels. It can relieve or prevent angina, but may cause headache, dizziness, or low blood pressure and can interact dangerously with phosphodiesterase-5 inhibitors.

## Glycerylphosphorylcholine

A choline-containing substance formed when certain membrane fats are broken down. It occurs in body fluids and cells and participates in membrane metabolism and the supply of choline.

## Glycine

A small amino acid used to make proteins and heme. It also acts as an inhibitory chemical messenger, especially in the spinal cord and brainstem, helping regulate movement and reflexes.

## Glycine Receptor

A ligand-gated chloride channel activated by glycine that mediates inhibitory neurotransmission, especially in the spinal cord and brainstem.

## Glycocalyx

A sugar-rich coating on the outside of many cells, including cells lining blood vessels. It helps protect surfaces, control passage of substances, support cell attachment, and influence inflammation and clotting.

## Glycogen

The body's stored form of glucose. It is kept mainly in the liver, which releases it between meals, and in muscles, which use it during activity.

## Glycogen Storage Disease

Any of several inherited disorders in which the body cannot make or break down glycogen normally. Depending on the type, these disorders may cause low blood glucose, an enlarged liver, muscle weakness, exercise intolerance, or heart problems.

---

### **Glycogen Storage Disease Type 0 (GSD-0)**

An inherited glycogen-synthase deficiency in which the liver or muscles cannot store normal amounts of glycogen. Liver disease may cause fasting ketotic hypoglycemia and post-meal hyperglycemia, while muscle disease may cause exercise intolerance, weakness, or cardiomyopathy.

### **Glycogen Storage Diseases**

Inherited disorders that prevent cells from making or breaking down glycogen normally. They can cause low blood sugar, enlarged liver, muscle weakness, or heart disease.

### **Glycogenesis**

The process of converting glucose into glycogen for storage, mainly in the liver and muscles. It occurs after eating, when blood glucose and insulin levels rise.

### **Glycogenolysis**

The breakdown of stored glycogen. The liver uses it to release glucose into the blood between meals, while muscles use their glycogen locally for energy during exercise.

### **Glycohemoglobin**

Hemoglobin with glucose chemically attached to it; glycated hemoglobin A1c reflects average blood glucose over roughly the preceding two to three months.

### **Glycolipid**

A lipid containing a carbohydrate group that contributes to cell membranes, recognition, signaling, and myelin structure.

### **Glycolysis**

A series of reactions in the cytosol that breaks one glucose molecule into two pyruvate molecules. It produces ATP and other molecules used in further energy production and can proceed when oxygen is limited.

### **Glycopeptide**

A molecule made of a short protein chain with one or more attached sugar groups. Glycopeptides occur in cell-surface proteins, antibodies, and some medicines, and the attached sugars can affect structure and function.

### **Glycophorin**

A sugar-coated protein in the red-blood-cell membrane that helps maintain the cell surface and carries markers used to distinguish blood-cell types.

### **Glycoprotein**

A protein with covalently attached carbohydrate chains, important in receptors, antibodies, enzymes, mucus, cell adhesion, and extracellular matrix.

### **Glycopyrrolate**

Glycopyrrolate is an anticholinergic medicine used to reduce secretions and, in some inhaled forms, to help control chronic obstructive lung disease; it may cause dry mouth, constipation, or urinary retention.

### **Glycosaminoglycan**

A long, unbranched carbohydrate chain found in the extracellular matrix and connective tissues. Abnormal breakdown or accumulation of GAGs occurs in mucopolysaccharidoses.

### **Glycosaminoglycans**

Glycosaminoglycans are long chains of sugars found in connective tissue and involved in its structure, hydration, and signaling.

### **Glycoside**

A compound in which a sugar is chemically linked to another molecule. Glycosides occur in plants, foods, and medicines; their effects depend on the attached molecule and how the body breaks the bond.

### **Glycosphingolipid**

A membrane lipid containing ceramide and one or more sugars, involved in cell recognition, signaling, and nervous-system structure.

---

## Glycosuria

The presence of glucose in the urine. It most often occurs when blood glucose is high, but it can also result from pregnancy, certain medicines, or a kidney-tubule problem that prevents normal glucose reabsorption.

## Glycosylated Hemoglobin

Hemoglobin with glucose attached; the A1C measurement reflects average blood sugar over roughly two to three months and helps diagnose or monitor diabetes.

### Alternate Names:

Hemoglobin A1c; HbA1c; Glycohemoglobin

## Glycosylation

Attachment of sugar chains to proteins or fats inside cells. This process helps determine their shape, stability, location, and activity; abnormal glycosylation can contribute to inherited disease, cancer, and immune disorders.

## Glycosylphosphatidylinositol

A lipid-and-sugar anchor that attaches certain proteins to the outer surface of a cell membrane. Defects in making or attaching this anchor can cause rare blood, nerve, or developmental disorders.

## Glycosyltransferase

Glycosyltransferase is an enzyme that transfers a sugar to a protein, lipid, or other molecule during glycan synthesis.

## Glycyrrhetic Acid

An active metabolite of licorice glycosides that inhibits 11-beta-hydroxysteroid dehydrogenase; excess exposure can cause hypertension, low potassium, and fluid retention.

## Glycyrrhizic Acid

The major saponin in licorice root that is converted to glycyrrhetic acid and can produce mineralocorticoid-like toxicity when consumed excessively.

## Glyphosate

A broad-spectrum herbicide that inhibits plant EPSP synthase; concentrated exposure can irritate tissues and cause systemic poisoning, especially after ingestion.

## Gm

Abbreviation; see *Gram*.

## GN

Abbreviation; see *Glomerulonephritis*.

## Gnathic

Relating to the jaw, especially the upper jaw or lower jaw and their development, structure, or position.

## Gnathion

The lowest midline point on the outer surface of the lower jaw. It is used as a landmark in dental, orthodontic, facial, and skull measurements.

## Gnathitis

Inflammation of the jaw or jaw tissues. It may cause pain, swelling, tenderness, difficulty opening the mouth, or chewing problems and can result from infection, injury, or inflammatory disease.

## Gnathoplasty

Plastic or reconstructive surgery to reshape or repair the jaw. It may improve function, facial balance, or a structural defect; risks include infection, nerve injury, altered bite, and recurrence.

## Gnathostoma

A group of roundworms that can infect people who eat raw or undercooked freshwater fish, frogs, or other carriers. Larvae may migrate through skin, organs, or the nervous system and cause swelling, pain, or serious neurologic disease.

## Gnathostoma Spinigerum

A roundworm that can cause gnathostomiasis after people eat raw or undercooked freshwater animals. Migrating larvae may produce recurring painful skin swellings and, rarely, eye or brain disease.

---

## Gnathostomiasis

A parasitic infection caused by *Gnathostoma* larvae acquired from undercooked freshwater fish or other hosts, producing migratory skin swellings or deeper neurologic or ocular disease.

## Gnawing Pain

A dull, persistent pain often described as an ache or a feeling that something is eating away at the body. It can arise from muscles, nerves, internal organs, or inflammation. Its location, duration, triggers, and accompanying symptoms help identify the cause.

## GNE Myopathy

A rare inherited muscle disease that usually begins in early adulthood and slowly weakens the muscles of the lower legs, causing foot drop, then the hands and arms. The thigh muscles are often spared until late disease. There is no cure, but rehabilitation helps maintain function.

### Alternate Names:

Nonaka Myopathy; Distal Myopathy with Rimmed Vacuoles (DMRV); Hereditary Inclusion Body Myopathy (HIBM)

## GnRH

Gonadotropin-releasing hormone, a hormone from the brain that signals the pituitary gland to release hormones controlling ovulation, sperm production, and sex-hormone production.

### Alternate Names:

Gonadotropin-Releasing Hormone

## GnRH Agonist

A medicine that first increases and then suppresses signals from the pituitary gland to the reproductive organs. It lowers sex-hormone production and is used in selected fertility treatments, endometriosis, fibroids, and hormone-sensitive cancers.

## GnRH Antagonist

A medicine that directly blocks the brain-to-pituitary signal controlling reproductive hormones. It lowers these hormones quickly without the temporary hormone rise that can occur when a GnRH agonist is started.

## GnRH Antagonists

Medicines that block a brain hormone signal and quickly reduce pituitary release of reproductive hormones. They are used in some fertility treatments and hormone-sensitive cancers, with effects depending on the specific medicine.

## Goal Attainment Scaling

A way to measure progress toward personal health-care goals. The goal is defined in advance with levels ranging from much less than expected to much better than expected, then scored after treatment.

## Goal Setting

Choosing clear, realistic outcomes and deciding how and when to work toward them. In healthcare, goals can reflect a person's priorities, include a way to measure progress, and change as needs or circumstances change.

## Goals-Of-Care Discussion

A clinical conversation that aligns treatment recommendations with a patient's values, priorities, prognosis, acceptable burdens, and preferred outcomes. It may include decisions about hospitalization, resuscitation, artificial nutrition, and comfort-focused care.

## Goblet Cell

A mucus-secreting epithelial cell, abundant in the respiratory and intestinal tracts, that protects surfaces with a mucin-rich layer.

## Goblet Cell Adenocarcinoma

A distinctive appendiceal adenocarcinoma composed of goblet-like mucinous cells, formerly called goblet cell carcinoid. It can spread within the peritoneum or to the ovaries and is managed according to grade, stage, and dissemination.

## Goblet Cells

Cells lining the intestines and airways that make and release mucus. Mucus moistens and protects these surfaces and helps trap particles, germs, and other substances for removal.

---

## Goeckerman Regimen

A dermatological therapy used for extensive or recalcitrant plaque psoriasis combining the daily topical application of crude coal tar with exposure to ultraviolet B (UVB) phototherapy.

### Alternate Names:

Goeckerman Therapy; Coal Tar and UVB Regimen

## Goiter

Enlargement of the thyroid gland, either throughout the gland or as one or more lumps. It may occur with normal, overactive, or underactive thyroid function and can result from iodine deficiency, immune disease, or a tumor.

## Goitre

Enlargement of the thyroid gland. It may occur with normal, low, or high thyroid hormone levels and can result from iodine deficiency, autoimmune disease, nodules, or inflammation.

## Goitrogenic Foods

Foods or food components that can interfere with thyroid hormone production or iodine use under some conditions; risk depends on intake, preparation, iodine status, and thyroid health.

## Goitrous Autoimmune Thyroiditis

Alternate name; see *Hashimoto Thyroiditis*.

### Alternate Names:

Thyromegaly; Struma; Thyroid Enlargement

## Gold

A metal used in selected dental restorations, implants, laboratory applications, and historical medicinal preparations; exposure can cause contact allergy or toxicity depending on the compound and dose.

## Gold Alloy

A mixture of gold and other metals used for some dental restorations. It is durable, fits closely, and is usually well tolerated by body tissues, but it costs more and has a distinctive appearance.

## Gold Alloys

Metal mixtures containing gold and other elements, used in selected dental restorations or medical devices; properties and biocompatibility depend on composition.

## Gold Therapy

Treatment with gold-containing medicines, formerly used mainly for rheumatoid arthritis. Its use is now limited because benefit is slow and serious effects can include skin or mouth inflammation, kidney injury, and reduced blood-cell counts.

## Golden Hour

The golden hour is the early period after a serious injury or medical emergency when rapid assessment and treatment may improve the chance of survival and recovery.

## Goldenhar Syndrome

A birth condition, also called oculo-auriculo-vertebral spectrum, in which one side of the face, ear, eye, or spine develops differently. Features may include facial asymmetry, a small or misshapen ear, hearing loss, eyelid or eye-surface growths, and spinal differences.

## Goldmann Tonometry

A slit-lamp examination that estimates intraocular pressure by measuring the force needed to flatten a defined area of the cornea.

## Golfer Elbow

An overuse injury of the tendons on the inner side of the elbow. Repeated gripping or wrist movement can cause pain and tenderness, usually worsened by lifting, bending the wrist, or making a fist.

## Golfer's Elbow

Painful irritation or degeneration of tendons attached to the inner side of the elbow. Repeated gripping or wrist movement can cause tenderness, reduced strength, and pain during lifting or bending.

---

## **Golgi Apparatus**

A structure inside cells that modifies, sorts, and packages proteins and fats made by the cell, sending them to other parts of the cell or releasing them outside it.

## **Golgi Cisternae**

Flattened membrane-bound sacs that form the Golgi apparatus and modify, sort, and package proteins and lipids for transport within or outside the cell.

## **Golgi Tendon Organ**

A sensory structure where a muscle joins its tendon. It detects tension in the muscle and tendon and helps the nervous system control muscle force and protect against excessive strain.

## **Gomori's Stain**

Gomori's stain is a group of laboratory stains used to highlight structures such as connective tissue, carbohydrates, fungi, or muscle components in tissue samples.

## **Gomphosis**

A fixed joint in which a tooth is held in its socket by a strong ligament.

## **Gonad**

A testis or ovary, an organ producing reproductive cells and sex hormones. Gonadal function influences puberty, fertility, menstrual cycles, sexual development, bone health, and other body processes.

## **Gonad Disorder**

A disease affecting testes or ovaries and causing abnormal reproductive-cell production, hormone secretion, development, fertility, or sexual function.

## **Gonadal Agenesis**

Congenital absence or severe underdevelopment of a gonad, resulting in impaired sex-hormone production and infertility or differences in sexual development.

## **Gonadal Dysgenesis**

Incomplete or abnormal development of the ovaries or testes, often resulting from chromosomal abnormalities, gene variants, or disordered sex development. It may cause atypical genital development, delayed or absent puberty, infertility, and abnormal gonadal hormone production.

## **Gonadal Vein**

A vein draining a testis or ovary. The right gonadal vein usually drains directly into the inferior vena cava, while the left usually drains into the left renal vein.

## **Gonadoblastoma**

A rare tumor that develops in an abnormal or incompletely developed ovary or testicle. It may produce sex hormones and has a risk of becoming a malignant germ-cell tumor, especially when a Y-chromosome change is present.

## **Gonadorelin**

Synthetic gonadotropin-releasing hormone that stimulates pituitary release of luteinizing hormone and follicle-stimulating hormone when given intermittently.

## **Gonadotrophin**

A hormone that acts on the ovaries or testes. The pituitary hormones follicle-stimulating hormone and luteinizing hormone regulate egg development, ovulation, sperm production, and sex-hormone release.

## **Gonadotrophin-Releasing Factors**

Signals from the brain that control release of the pituitary hormones acting on the ovaries or testes. The main regulator is gonadotropin-releasing hormone, released in pulses from the hypothalamus.

## **Gonadotrophins**

Hormones that act on the ovaries or testes. Follicle-stimulating hormone helps develop eggs and sperm, while luteinizing hormone triggers ovulation and supports sex-hormone production.



---

## Gonadotrophs

Pituitary cells that release luteinizing hormone and follicle-stimulating hormone after stimulation by GnRH, regulating the ovaries, testes, fertility, and sex-hormone production.

## Gonadotropin

A hormone, chiefly luteinizing hormone or follicle-stimulating hormone, that acts on the ovaries or testes to regulate gamete production and sex-hormone secretion.

## Gonadotropin Hormone

A hormone that acts on the ovaries or testes. The pituitary gonadotropins follicle-stimulating hormone and luteinizing hormone regulate reproductive development, ovulation, sperm production, and sex-hormone production.

## Gonadotropin-Releasing Hormone

A hormone made in the brain that signals the pituitary gland to release two hormones, FSH and LH. These hormones control the ovaries and testes, including egg development, ovulation, sperm production, and sex-hormone production.

### Alternate Names:

GnRH

## Gonadotropins

Hormones that act on the gonads. In humans, the pituitary gonadotropins are follicle-stimulating hormone and luteinizing hormone, which regulate gamete production and sex-hormone secretion.

## Gonads

The reproductive organs: testes in males and ovaries in females. They produce sperm or eggs and make sex hormones such as testosterone, estrogen, and progesterone. These hormones influence puberty, fertility, menstrual cycles, pregnancy, and bone health.

## Gondoic Acid

A 20-carbon monounsaturated fatty acid found in some seed oils and incorporated into dietary and

tissue lipids.

## Goniometer

An instrument measuring joint angles and range of motion, used to assess stiffness, document impairment, and track progress during rehabilitation.

## Gonioscopy

An eye examination using a special contact lens to view the angle where the iris meets the cornea. It helps determine whether glaucoma is open-angle or angle-closure and guides treatment.

## Gonococcal

Relating to *Neisseria gonorrhoeae*, the bacterium that causes gonorrhea. Gonococcal infection commonly affects the genitals, rectum, or throat and may spread to joints or blood.

## Gonococcal Arthritis

A disseminated infection of joints caused by *Neisseria gonorrhoeae*, often with fever, migrating joint or tendon pain, skin pustules, and inflammation around tendons.

## Gonococcal Pharyngitis

Infection of the throat by *Neisseria gonorrhoeae*, usually acquired through sexual contact. It may cause sore throat or no symptoms.

## Gonococcal Urethritis

Infection and inflammation of the urethra caused by *Neisseria gonorrhoeae*. It may cause burning urination or discharge, but some people have no symptoms; untreated infection can spread.

## Gonococcus

A spherical bacterium of the genus *Neisseria*. *Neisseria gonorrhoeae*, the gonococcus, causes the sexually transmitted infection gonorrhea.

## Gonorrhea

A sexually transmitted infection caused by the bacterium *Neisseria gonorrhoeae*. It may cause discharge, painful urination, pelvic or rectal symptoms,

---

or no symptoms. Untreated infection can lead to infertility, joint infection, or infection in a newborn.

### **Gonorrhoea**

Alternate spelling; see *Gonorrhea*.

### **Good Samaritan**

A person who voluntarily helps someone in an emergency, especially by providing immediate aid before professional responders arrive.

### **Good Samaritan Law**

A law that may protect a person who provides reasonable emergency assistance from certain civil claims, subject to jurisdiction-specific conditions and exceptions such as gross negligence.

### **Goodell Sign**

A clinical sign of early pregnancy characterized by marked softening of the uterine cervix (produced by increased pelvic vascularity and edema), identifiable upon digital pelvic bimanual examination around the sixth to eighth week of gestation.

### **Goodpasture Syndrome**

An autoimmune disease in which antibodies attack the lung air sacs and kidney filters. It can cause coughing blood, breathlessness, blood in the urine, and rapidly worsening kidney failure.

### **Goodpasture's Syndrome**

An autoimmune disease in which antibodies attack the lungs and kidney filters, causing coughing blood, kidney inflammation, blood in urine, and sometimes rapid kidney failure.

### **Goose Bump**

A small raised area around a hair follicle caused by contraction of tiny arrector pili muscles, commonly in response to cold, fear, or strong emotion.

### **Goose-Skin**

Goose skin is the temporary raising of hair follicles, usually caused by cold, fear, or strong emotion.

### **Gooseflesh**

Temporary raised bumps around hair follicles caused by contraction of tiny skin muscles, often in response to cold, fear, or strong emotion. It is usually harmless.

### **Gorlin Syndrome**

Alternate name; see *Gorlin's Syndrome*.

### **Gorlin's Syndrome**

An inherited disorder that increases the risk of multiple basal-cell skin cancers, jaw cysts, and abnormalities of the bones, eyes, and nervous system.

### **Goserelin**

A gonadotropin-releasing hormone agonist that initially stimulates and then suppresses pituitary gonadotropins; it is used in hormone-sensitive prostate or breast cancer and selected gynecologic conditions.

### **Gottron Papules**

Raised red, purple, or scaly patches over the knuckles and other finger joints. They are a characteristic skin finding of dermatomyositis, an inflammatory disease that can also cause muscle weakness.

### **Gottron Sign**

Symmetrical, violaceous, macular or papular erythematous patches and scaling located over the extensor aspects of the joints (particularly the elbows, patellae, and medial malleoli), serving as a characteristic cutaneous hallmark of dermatomyositis.

#### **Alternate Names:**

Gottron's Sign; Dermatomyositis Extensor Erythema

### **Gougerot-Carteaud Syndrome**

A skin disorder causing brown, slightly raised spots that join into net-like patches, usually on the upper chest, back, or neck of young people.

### **Gout**

A type of inflammatory arthritis caused by a buildup of urate, a substance made when the body breaks down purines. When urate levels stay high, needle-

---

shaped crystals can form in and around joints. These crystals cause sudden attacks, or flares, of severe pain, swelling, warmth, redness, and tenderness, often in the big toe but also in other joints. Repeated flares can damage joints. Long-standing gout may cause firm crystal deposits called tophi and uric acid kidney stones.

**Alternate Names:**

Gouty Arthritis; Monosodium Urate Arthropathy

**Gout Flare**

A sudden attack of gout caused by inflammation around urate crystals in a joint. The joint becomes very painful, swollen, warm, and red, often beginning in the big toe. Infection can cause similar symptoms.

**Gout Nodulosis**

A condition in which many firm urate deposits, called tophi, form under the skin or around joints because of long-standing gout. They may occur at the ears, fingers, toes, elbows, or Achilles tendons and can restrict movement or break through the skin.

**Gout Suppressants**

Medicines that lower uric-acid production or increase its excretion to prevent recurrent gout attacks and urate crystal deposition.

**Gouty Tophus**

A nodular, chalky subcutaneous or periarticular deposit of crystallized monosodium urate monohydrate crystals surrounded by chronic foreign-body granulomatous inflammation, occurring in chronic uncontrolled hyperuricemia on the helix of the ear, olecranon bursa, Achilles tendon, and distal joints.

**Alternate Names:**

Tophus; Tophi

**Gower Syndrome**

A form of reflex syncope, also called vasovagal or vasodepressor syncope, in which a trigger causes a fall in blood pressure and sometimes heart rate, briefly reducing brain perfusion.

**Gowers Sign**

A maneuver in which a person uses the hands to push up from the floor or thighs because of proximal muscle weakness, classically seen in muscular dystrophy.

**Graafian Follicle**

A mature, fluid-filled sac in an ovary that contains an egg ready for release. It enlarges during the menstrual cycle and normally opens at ovulation; after release, the remaining tissue forms the corpus luteum.

**GRACE Score**

A risk score used in people with an acute coronary syndrome to estimate the chance of death or another heart attack during the hospital stay and over the next six months. It uses age, vital signs, kidney function, cardiac arrest, ECG changes, heart-failure signs, and heart-injury blood tests to guide treatment intensity.

**Alternate Names:**

Global Registry of Acute Coronary Events Score; GRACE Risk Model

**Gracilis Muscle**

A long, slender strap-like muscle located on the medial aspect of the thigh originating from the inferior pubic ramus and body of the pubis, descending to insert into the pes anserinus on the proximal medial tibia; acts to adduct the thigh and flex and internally rotate the leg.

**Grade**

A measure of how abnormal cells look under a microscope and, in many cancers, how quickly the growth is likely to behave. Lower grades usually look more like normal cells; higher grades usually look more abnormal. The grading system differs by disease.

**Grade IV Astrocytoma**

Older term; see *Glioblastoma*.

**Alternate Names:**

Glioblastoma; GBM

---

## Graded Potentials

Small changes in an electrically active cell caused by a stimulus. Their size depends on the stimulus strength, and they can either increase or decrease the chance that the cell will send a nerve signal.

## Gradenigo Syndrome

A classic clinical triad consisting of suppurative otitis media (otorrhea), retro-orbital or temporal pain in the sensory distribution of the trigeminal nerve (cranial nerve V), and ipsilateral abducens nerve palsy (cranial nerve VI) caused by petrous apicitis.

### Alternate Names:

Gradenigo's Syndrome; Petrous Apicitis Triad

## Graefe's Sign

A delayed downward movement of the upper eyelid during downward gaze, classically associated with thyroid eye disease but not specific to it.

## Graft

Tissue or a synthetic material transferred to another site to replace, repair, reinforce, or bypass a damaged body structure.

## Graft Versus Host

Alternate name; see *Graft Versus Host Disease*.

## Graft Versus Host Disease

A complication of a stem-cell or bone-marrow transplant in which immune cells from the donor attack the recipient's tissues. It commonly affects the skin, liver, and digestive tract and may be short-term or long-lasting; prevention and treatment use immune-suppressing medicines.

### Alternate Names:

GVHD; Transfusion-Associated or Allogeneic GVHD

## Graft-Versus-Host Disease

Alternate spelling; see *Graft Versus Host Disease*.

## Graft-Versus-Host Reaction

An immune reaction in which donor immune cells

attack the recipient's tissues after transplantation, potentially affecting skin, intestines, liver, lungs, or other organs.

## Grafting

The surgical transfer of tissue or a synthetic substitute to repair, replace, or support a body structure. An autograft comes from the same person, whereas an allograft comes from another person of the same species.

## Graham-Little-Piccardi-Lasseur Syndrome

A rare variant of lichen planopilaris characterized by scarring hair loss on the scalp, nonscarring loss of axillary or pubic hair, and follicular papules on the trunk. It is an inflammatory disorder requiring dermatologic assessment and treatment to limit permanent loss.

## Gram

A metric unit used to measure mass, equal to one-thousandth of a kilogram. Grams are commonly used for medicines, laboratory samples, food, and body substances; one thousand milligrams make one gram.

### Alternate Names:

Metric Gram; Unit G

## Gram Mass Unit

A metric unit of mass equal to one-thousandth of a kilogram; medication doses and laboratory measurements are commonly expressed in grams or smaller metric units.

## Gram Metric Unit

A metric unit of mass equal to one-thousandth of a kilogram or approximately 15.432 grains.

## Gram Stain

A laboratory test that colors bacteria to group them as Gram-positive or Gram-negative. It helps identify possible causes of infection quickly and can guide initial treatment while more specific tests are performed.

---

## Gram Stain Of Skin Lesion

Microscopic examination of a skin lesion specimen stained with Gram stain to identify bacteria. Useful for diagnosing bacterial skin infections and guiding antibiotic therapy. Results available within hours.

## Gram Stain Of Tissue Biopsy

Microscopic examination of tissue biopsy specimen stained with Gram stain to identify bacteria. Used to diagnose tissue infections and guide antimicrobial therapy. May be performed alongside culture and histopathology.

### Alternate Names:

Tissue Gram Stain; Histochemical Bacterial Stain

## Gram's Stain

Gram's stain is a laboratory method that classifies bacteria by how their cell walls retain the stain, helping guide identification and initial antibiotic choices.

## Gram-Negative

Describing bacteria with a thin peptidoglycan cell wall and an outer membrane containing lipopolysaccharide; they stain pink or red with the Gram stain.

## Gram-Negative Bacilli

Rod-shaped bacteria that appear pink or red with a Gram stain because they have a thin peptidoglycan layer and an outer membrane. They include harmless body organisms and bacteria that can cause urinary, respiratory, intestinal, bloodstream, wound, or other infections.

## Gram-Negative Bacteria

Bacteria that do not retain crystal violet during Gram staining and therefore appear pink or red. They have a thin peptidoglycan layer and an outer membrane that may contain lipopolysaccharide. Examples include *Escherichia coli*, *Klebsiella*, *Pseudomonas*, and *Neisseria*.

## Gram-Negative Cocci And Coccobacilli

Gram-negative bacteria that are either round or short rod-shaped. Cocci are round, while coccobacilli are short rods that may look almost round.

They include bacteria that can cause respiratory, genital, bloodstream, intestinal, and other infections.

## Gram-Negative Folliculitis

Folliculitis caused by gram-negative bacteria, often after prolonged antibiotic treatment for acne or in people with impaired immunity. It produces papules and pustules, frequently on the face, and requires culture-directed treatment.

## Gram-Negative Meningitis

Meningitis caused by gram-negative bacteria, including *Escherichia coli*, *Klebsiella*, and *Neisseria* species. It can progress rapidly to sepsis, neurologic injury, and death.

## Gram-Negative Toe Web Infection

Infection of the moist spaces between the toes by gram-negative bacteria, causing maceration, erosions, malodor, pain, or discharge. Moisture, occlusive footwear, and existing tinea pedis predispose; drying and appropriate antimicrobial treatment are important.

## Gram-Positive

Describing bacteria with a thick peptidoglycan cell wall that retains crystal violet during Gram staining and therefore appears purple.

## Gram-Positive Bacilli

Rod-shaped bacteria that retain the purple part of the Gram stain because of their thick peptidoglycan cell wall. Some are harmless colonizers, while others can cause wound, intestinal, respiratory, bloodstream, or nervous-system infections.

## Gram-Positive Bacteria

Bacteria that retain crystal violet in the Gram stain (appear purple/blue), characterized by a thick peptidoglycan cell wall and no outer membrane.

## Gram-Positive Cocci

Round bacteria that appear purple with a Gram

---

stain because they have a thick peptidoglycan cell wall. Important groups include staphylococci, streptococci, and enterococci, which may normally live on the body or cause skin, throat, lung, urinary, heart-valve, or bloodstream infections.

### **Gramicidin**

Gramicidin is a topical antibiotic that forms channels in bacterial membranes; it is not generally used systemically because it can be toxic.

### **Grand Multipara**

A person who has given birth five or more times beyond the gestational threshold used to count parity; definitions can vary by obstetric system.

### **Grand Rounds**

An educational medical conference in which clinicians present a case, diagnosis, management, or clinical topic to a professional audience.

### **Granisetron**

Granisetron is a selective 5-HT<sub>3</sub>-receptor antagonist used to prevent chemotherapy-, radiotherapy-, or surgery-related nausea and vomiting; headache, constipation, and QT prolongation can occur.

### **Granisetron Hydrochloride**

The hydrochloride form of granisetron, a medicine that blocks serotonin 5-HT<sub>3</sub> receptors to prevent nausea and vomiting from chemotherapy, radiation, or surgery. Headache, constipation, and heart-rhythm changes may occur.

### **Granular Casts**

Small grainy particles formed inside kidney tubules and seen in urine. They may occur when kidney cells are injured or break down and can be found in acute kidney injury or some chronic kidney diseases.

### **Granular Cell Tumor Of The Breast**

An uncommon tumor showing Schwann-cell differentiation that can occur in the breast and mimic carcinoma clinically or on imaging. Histologic examination distinguishes it from malignancy; most

are benign.

### **Granular Cell Tumors**

Usually slow-growing tumors derived from Schwann-cell-lineage tissue, most often affecting the skin, tongue, or subcutaneous tissue. Most are benign, but some are malignant; persistent or changing nodules require examination and histologic diagnosis.

### **Granular Leukocyte**

A granulocyte, a white blood cell containing cytoplasmic granules; the main types are neutrophils, eosinophils, and basophils.

### **Granular Parakeratosis**

A rare skin condition causing brown or reddish, sometimes itchy patches or papules, often in skin folds such as the armpits. It reflects abnormal keratinization and may be difficult to distinguish from other rashes.

### **Granulation**

The formation of new vascular connective tissue during wound healing, producing moist red or pink granulation tissue made of capillaries, fibroblasts, and extracellular matrix.

### **Granulation Tissue**

New tissue that grows in a healing wound and contains small blood vessels, connective tissue, and immune cells. Healthy granulation tissue is usually pink or red and moist.

### **Granulicatella**

Granulicatella is a group of bacteria that may live in the mouth and can rarely cause bloodstream infection or endocarditis.

### **Granulicatella Adiacens**

A mouth bacterium that can rarely cause serious infection, especially infection of heart valves or the bloodstream. It may be difficult to grow in culture, and treatment depends on susceptibility testing.



---

## Granulicatella Elegans

A bacterium normally found in the mouth that can rarely cause bloodstream infection or infection of a heart valve. It may grow poorly in routine culture, so repeated positive cultures and compatible symptoms are important when judging its significance.

## Granulocyte

A type of white blood cell containing visible granules. Neutrophils, eosinophils, and basophils are granulocytes that help fight infection, trigger inflammation, and respond to allergic or parasitic disease.

## Granulocyte Colony-Stimulating Factor

A hematopoietic glycoprotein growth factor that stimulates the bone marrow to produce and differentiate granulocyte precursors into mature neutrophils and mobilize them into circulating blood; recombinant forms (filgrastim, pegfilgrastim) are used to treat and prevent chemotherapy-induced neutropenia.

**Alternate Names:**

G-CSF

## Granulocyte Count

A granulocyte count measures the number of neutrophils, eosinophils, and basophils in the blood and helps assess infection, inflammation, and bone-marrow function.

## Granulocytic Sarcoma

A mass of immature myeloid blood-forming cells outside the bone marrow, also called myeloid sarcoma. It may occur with acute myeloid leukemia or precede it.

## Granulocytopenia

An abnormally low number of granulocytes, especially neutrophils, which can increase susceptibility to bacterial and fungal infection.

## Granuloma

A small area of organized inflammation formed when immune cells surround material they cannot easily remove, such as certain germs, foreign material, or

substances released by the body.

## Granuloma Annulare

A benign, noninfectious inflammatory skin condition that causes raised red, pink, or skin-colored bumps arranged in rings, often with clearer skin in the center. It commonly affects the hands, feet, or elbows and may cause little or no discomfort. Widespread disease may be associated with diabetes or thyroid disease, but the cause is often unknown.

## Granuloma Fungoides

Alternate spelling; see *Mycosis Fungoides*.

## Granuloma Gluteale Infantum

A sharply defined reddish-purple inflammatory plaque in the diaper area of an infant, often related to prolonged irritant dermatitis, occlusion, or topical products. Removing irritants and treating the underlying diaper dermatitis usually leads to improvement.

## Granuloma Inguinale

A chronic sexually transmitted infection causing painless, slowly enlarging ulcers in the genital or groin region. It is caused by bacteria and spreads through sexual contact; antibiotics are used in treatment.

## Granulomatosis

A disease in which long-lasting inflammation forms granulomas, small organized collections of immune cells. Depending on the cause, granulomas may affect the lungs, skin, lymph nodes, intestines, or other organs.

## Granulomatosis With Polyangiitis

An autoimmune disease that inflames and damages small and medium-sized blood vessels, often affecting the nose, sinuses, lungs, and kidneys. It may cause sinus problems, cough, coughing blood, or kidney injury.

**Alternate Names:**

GPA

---

### Granulomatous Amebic Encephalitis

A rare, severe brain infection caused by free-living amoebae, usually in people with weakened immunity. It develops over days or weeks and may cause headache, confusion, seizures, weakness, or coma.

### Granulomatous Colitis

Inflammation of the colon in which granulomas are found on tissue examination, often associated with Crohn disease but also possible with infections or other inflammatory conditions.

### Granulomatous Inflammation

A pattern of long-lasting inflammation in which immune cells form small clusters around material they cannot easily remove. It can occur with infections, foreign materials, autoimmune disease, or other inflammatory conditions.

### Granulomatous Mastitis

A chronic inflammatory breast disease characterized by noncaseating granulomas after other causes, especially infection and sarcoidosis, have been excluded. It may present as a painful mass, abscesses, or draining sinuses.

### Granulosa Cell

A granulosa cell is a cell surrounding an ovarian egg that supports its development and produces hormones such as estrogen and inhibin.

### Granulosa Cell Tumor

A tumor arising from hormone-producing cells in the ovary. It may release estrogen, causing irregular bleeding or early puberty, and can recur years after treatment even when it initially appears slow-growing.

### Granulosa Cell Tumors

Ovarian sex-cord stromal tumors that may produce estrogen or inhibin. They can cause abnormal uterine bleeding, endometrial thickening, or an abdominal mass; treatment usually involves surgical staging and long-term follow-up because late recurrence can occur.

### Granulosa Tumor

Alternate form; see *Granulosa Cell Tumor*.

### Granzyme

A protease released by cytotoxic T lymphocytes and natural killer cells that enters target cells through perforin-associated pores and promotes programmed cell death.

### Grasp Reflex

An automatic closing of a baby's fingers when the palm is touched. It is normal in early infancy and usually fades as voluntary hand control develops; persistence or asymmetry can occur with nervous-system problems.

### Grass Allergy

Hypersensitivity reaction to grass pollen, causing allergic rhinitis (hay fever) with symptoms including sneezing, nasal congestion, itchy eyes, and throat irritation. Seasonal, typically spring through fall. Treatment includes antihistamines, nasal corticosteroids, and allergen avoidance.

#### Alternate Names:

Grass Pollen Allergy; Gramineae Pollinosis

### Grass And Weed Killer Poisoning

Illness caused by swallowing, inhaling, or getting herbicide on the skin or eyes. Effects range from irritation, vomiting, and breathing problems to severe poisoning, depending on the product and amount.

#### Alternate Names:

Herbicide Intoxication; Weedicide Poisoning

### Grave Wax

Alternate name; see *Adipocere*.

### Graves Disease

An autoimmune disease that causes an overactive thyroid (hyperthyroidism). The immune system makes antibodies that act like thyroid-stimulating hormone and make the thyroid produce too much thyroid hormone. Symptoms may include weight loss, a fast or irregular heartbeat, sweating, heat intolerance, tremor, nervousness, trouble sleeping,

---

frequent bowel movements, muscle weakness, or an enlarged thyroid (goiter). Some people develop Graves eye disease, which can cause dry or gritty eyes, puffiness, bulging eyes, light sensitivity, or double vision. Rarely, the skin over the shins becomes thickened and red.

### Graves Ophthalmopathy

An autoimmune inflammatory disorder of the retro-orbital tissues and extraocular muscles associated with Graves disease, driven by autoantibodies against the TSH receptor leading to fibroblastic proliferation, glycosaminoglycan deposition, proptosis, upper lid retraction, and diplopia.

#### Alternate Names:

Graves Orbitopathy; Thyroid Eye Disease

### Gravida

The number of times a person has been pregnant, regardless of pregnancy outcome; the current pregnancy is included in the count.

### Gravidity

Gravidity is the number of times a person has been pregnant, including the current pregnancy and pregnancies that did not result in a live birth.

### Gray

A unit used to measure the amount of ionizing radiation absorbed by a material or body tissue. One gray equals one joule of radiation energy absorbed per kilogram. It is used in radiation treatment and safety assessment.

#### Alternate Names:

Absorbed Dose Unit Gy; Radiation Unit Gray

### Gray Baby Syndrome

A serious form of chloramphenicol toxicity in newborns, caused by immature drug metabolism and characterized by abdominal distension, vomiting, cyanosis or gray skin color, cardiovascular collapse, and potentially death.

### Gray Matter

Areas of the brain and spinal cord containing many

nerve-cell bodies and connections. Gray matter processes information and helps control movement, sensation, memory, speech, and other functions.

### Great Saphenous Vein

The longest superficial vein of the lower limb, ascending from the medial foot to drain into the femoral vein at the saphenofemoral junction.

### Greater Curvature

The long, rounded outer border of the stomach, extending from its upper opening to its outlet. It provides attachment for the greater omentum and receives blood from several stomach arteries.

### Greater Omentum

A fold of fatty peritoneal tissue that hangs from the stomach and partly covers the abdominal organs.

### Greater Trochanter

The large bony prominence on the outer upper thigh, where several hip muscles attach. It can be felt through the skin and is a common site of pain from tendon problems, bursitis, or injury.

### Greater Trochanteric Pain Syndrome

Pain over the lateral hip centered on the greater trochanter, commonly caused by tendinopathy or bursitis of the gluteal tendons. Presents with lateral hip pain worse with walking, climbing stairs, or lying on the affected side. Treatment includes physical therapy and activity modification.

#### Alternate Names:

GTPS; Trochanteric Bursitis; Gluteal Tendinopathy

### Greater Vestibular Glands

Paired glands near the vaginal opening that secrete mucus into the vestibule, contributing to lubrication. They are also called Bartholin glands.

#### Alternate Names:

Bartholin Glands; Vulvovaginal Glands

### Green Blindness

A form of red-green color-vision deficiency in which distinguishing green from certain other colors is

---

difficult, usually because of an inherited cone-photopigment difference.

### **Green Nail Syndrome**

Green discoloration of one or more nails, usually caused by colonization or infection with *Pseudomonas aeruginosa*, often in an onycholytic or chronically moist nail.

### **Green Stool**

Stool with a green color that may result from diet, food coloring, rapid intestinal transit, medicines, or infection.

### **Greenstick Fracture**

An incomplete fracture in which a young bone bends and cracks rather than breaking all the way through.

### **Grenz Zone**

A narrow, uninvolved zone of normal collagenous dermis located immediately beneath the epidermis, separating the epidermal basement membrane from a dense underlying dermal cellular infiltrate, classically seen in lepromatous leprosy, granuloma faciale, and cutaneous lymphomas.

#### **Alternate Names:**

Subepidermal Grenz Zone; Clear Dermal Zone

### **Grey Ankle Phenomenon**

A gray or dusky discoloration around the ankle associated with chronic venous insufficiency and hemosiderin deposition. It occurs with other signs of venous disease.

### **Grey Matter**

Brain and spinal-cord tissue containing many nerve-cell bodies, connections, and supporting cells. It processes information and helps control movement, sensation, thinking, memory, and other functions.

### **Grey Turner Sign**

Bruising on the sides of the belly between the ribs and the hips, which is a sign of internal bleeding from severe pancreas inflammation or an injured blood vessel.

### **Grey Turner's Sign**

Bruising or blue-purple discoloration of the skin at the sides of the lower back. It can occur when bleeding deep inside the abdomen spreads through the tissues, including in severe pancreatitis.

### **Grid Implant**

A flexible electrode array placed on or near the brain or another tissue surface to record electrical activity or deliver stimulation.

### **Grief**

The natural emotional response to loss, particularly the death of a loved one, but also following other significant losses such as divorce, job loss, or declining health. It encompasses a range of feelings including sadness, anger, guilt, and numbness, often occurring in waves rather than linear stages.

### **Grief and Bereavement**

The natural emotional response to loss, encompassing the process of coping with the death of a loved one or other significant losses. Includes grief (emotional response), loss (the event), and bereavement (the period of mourning). May involve physical, emotional, social, and spiritual dimensions.

#### **Alternate Names:**

Bereavement; Mourning; Complicated Grief

### **Grimontia Hollisae**

A salt-loving bacterium from seawater and seafood, formerly called *Vibrio hollisae*. Eating contaminated raw or undercooked seafood can cause watery diarrhea, abdominal cramps, nausea, and fever; severe infection is uncommon but more likely in people with serious underlying illness.

### **Grip Strength**

The force generated when the hand closes around an object, commonly measured with a dynamometer to assess hand function, muscle strength, nutrition, or rehabilitation progress.

### **Gripping**

Cramping or spasmodic abdominal pain, often

---

caused by contractions, gas, inflammation, or obstruction of a hollow digestive organ.

### Griscelli Syndrome

A rare inherited disorder causing unusually light or silvery hair and skin pigmentation. Some forms also cause neurologic problems or severe immune over-activation, which can lead to fever, enlarged organs, low blood counts, and life-threatening illness.

### Griseofulvin

Griseofulvin is an oral antifungal that deposits in newly formed keratin and disrupts dermatophyte mitosis, treating infections of skin, hair, and nails; treatment is prolonged and hepatic or drug-interaction risks apply.

### Groin

The region where the lower abdomen meets the upper thigh, containing the inguinal canal and related muscles, vessels, nerves, and lymph nodes.

### Groin Lump

A palpable mass in the inguinal region, which may represent enlarged lymph nodes, a hernia, cyst, tumor, or another condition.

#### Alternate Names:

Inguinal Mass; Inguinal Lymphadenopathy

### Groin Pain

Pain in the inguinal region, which may originate from musculoskeletal, genitourinary, gastrointestinal, or vascular causes. Common causes include muscle strain, hernia, enlarged lymph nodes, and hip disease.

#### Alternate Names:

Inguinal Pain; Pubalgia

### Grommet Tube

A miniature, spool-shaped synthetic ventilation tube surgically inserted through an incision in the tympanic membrane (myringotomy) to equalize middle ear pressure and facilitate fluid drainage in chronic otitis media with effusion.

#### Alternate Names:

Tympanostomy Tube; Ventilation Tube

### Groove

A long, narrow depression or channel on a body structure, such as a groove on a bone for a tendon or a groove between folds of the brain. Its exact function depends on its location.

### Grooved Tongue

Alternate name; see *Fissured Tongue*.

### Gross Anatomy

The study of body structures visible without a microscope, including organs, body regions, bones, muscles, blood vessels, and their relationships.

### Gross Examination

Inspection of a tissue or other specimen with the unaided eye before it is examined under a microscope. The pathologist records its size, shape, color, and visible abnormalities to help guide diagnosis.

### Gross Hematuria

Visible blood in the urine, appearing pink, red, or brown. Causes include urinary tract infections, kidney stones, bladder or kidney cancer, trauma, and disease of the kidney filters.

#### Alternate Names:

Macroscopic Hematuria; Visible Hematuria

### Gross Motor Control

The ability to use large muscle groups for coordinated movements such as walking, running, jumping, and maintaining posture. Develops progressively from infancy through childhood and is assessed as part of developmental milestones.

#### Alternate Names:

Large Motor Coordination; Fundamental Motor Skills

### Gross Motor Skills

Movements that use the large muscles for posture, balance, walking, running, jumping, and climbing. Children develop these skills at different rates.

---

## Ground Itch

An itchy skin rash caused by hookworm larvae entering the skin, usually after bare-foot contact with contaminated soil. It produces small, red, intensely itchy bumps.

## Ground Substance

The gel-like, noncellular part of connective tissue surrounding cells and fibers. It contains water, proteoglycans, and other molecules that support tissues, allow nutrient movement, and influence inflammation and repair.

## Ground-Glass Opacity

A hazy area on a chest CT scan in which underlying lung vessels remain partly visible. It is a radiologic finding, not a diagnosis, and may result from infection, inflammation, fluid, bleeding, scarring, or sometimes a tumor; its meaning depends on the pattern and clinical context.

## Group 1 Pulmonary Hypertension (PAH)

Pulmonary arterial hypertension (PAH), the Group 1 form of pulmonary hypertension, is a pre-capillary pulmonary vascular disease in which narrowed or remodeled pulmonary arteries raise vascular resistance and pressure. Causes include idiopathic or heritable disease, drugs or toxins, connective-tissue disease, HIV, portal hypertension, congenital heart disease, and schistosomiasis; diagnosis requires right-heart catheterization and evaluation for other groups.

## Group 2 Pulmonary Hypertension

Pulmonary hypertension caused by left-heart disease, including left-ventricular dysfunction, valvular disease, or congenital left-heart disorders. Treatment focuses on the underlying cardiac condition and volume status; pulmonary arterial hypertension-specific drugs are not routinely interchangeable with this group.

## Group 3 Pulmonary Hypertension

Pulmonary hypertension associated with lung disease or hypoxia, such as COPD, interstitial lung disease, sleep-disordered breathing, or chronic hypoventilation. Management emphasizes treating the

lung disorder, correcting hypoxemia, and evaluating for severe or disproportionate vascular disease.

## Group 4 Pulmonary Hypertension

Pulmonary hypertension caused by pulmonary artery obstruction, most importantly chronic thromboembolic pulmonary hypertension. Patients require evaluation for persistent or recurrent thromboembolism and may benefit from anticoagulation, pulmonary endarterectomy, balloon angioplasty, or specialist medical therapy.

## Group A Strep

A bacterium that commonly causes sore throat and skin infections. It can occasionally cause scarlet fever, pneumonia, severe soft-tissue infection, bloodstream infection, or immune complications affecting the heart or kidneys.

## Group A Streptococcal Infections

Infections caused by *Streptococcus pyogenes*, including pharyngitis, scarlet fever, impetigo, cellulitis, pneumonia, necrotizing soft-tissue infection, and bloodstream infection. Immune-mediated complications can include rheumatic fever and poststreptococcal glomerulonephritis.

## Group A Streptococcus

*Streptococcus pyogenes*, a bacterium that can cause pharyngitis, skin infections, invasive disease, and immune-mediated complications.

## Group B Strep

A bacterium that can live harmlessly in the bowel or genital tract but can cause serious infection in newborns during or soon after birth. It can also cause infection in pregnant people and vulnerable adults. Pregnancy screening and care during labor can reduce newborn risk.

### Alternate Names:

Group B Streptococcus; *Streptococcus Agalactiae*

## Group B Strep Disease

Illness caused by group B *Streptococcus* bacteria, which can cause serious infections in newborns and infections in pregnant people or adults with medical



---

risks.

### Group B Streptococcal Disease

Infection caused by group B *Streptococcus* bacteria, which can be serious in newborns and vulnerable adults.

### Group B Streptococcal Infection In Pregnancy

Maternal colonization or infection with *Streptococcus agalactiae*, which can be transmitted to the newborn during labor and cause serious early-onset infection. Screening and intrapartum management reduce neonatal risk.

### Group B Streptococcal Infections in Newborns

Alternate name; see *Group B Streptococcal Neonatal Sepsis*.

### Group B Streptococcal Neonatal Sepsis

Serious bloodstream infection caused by Group B *Streptococcus* in neonates, typically acquired during birth. It can cause fever, lethargy, poor feeding, and breathing difficulty; treatment includes antibiotics and supportive care.

#### Alternate Names:

Early-Onset GBS Sepsis; Neonatal *Streptococcus Agalactiae* Sepsis

### Group B Streptococcal Septicemia of the Newborn

Alternate name; see *Group B Streptococcal Neonatal Sepsis*.

### Group B Streptococcus

A bacterium that may live harmlessly in the bowel or genital tract but can cause serious infection in newborns during or soon after birth. It can also cause infection in pregnant people and vulnerable adults; pregnancy screening helps reduce newborn risk.

### Group B Streptococcus in Pregnancy

Asymptomatic colonization of the maternal recto-

vaginal tract with *Streptococcus agalactiae*. Routine culture screening is performed between 36 and 37 weeks of gestation so that colonized individuals can receive intrapartum intravenous antibiotic prophylaxis to prevent neonatal early-onset sepsis, pneumonia, and meningitis.

#### Alternate Names:

Maternal GBS Colonization; Intrapartum GBS Prophylaxis; Maternal Group B Strep

#### Alternate Names:

Maternal GBS Colonization; Intrapartum GBS Prophylaxis

### Group Practice

A healthcare organization in which several clinicians share facilities, staff, records, or administrative services. Patients may receive coordinated care from different professionals within the same practice.

### Group Therapy

A form of psychotherapy where a small group of individuals with similar issues meet to discuss, interact, and develop coping strategies under the guidance of a trained therapist. Used for depression, anxiety, substance use disorders, grief, and chronic illness. Benefits include peer support and shared experiences.

### Grover Disease

A transient or recurrent acantholytic dermatosis causing itchy red papules or papulovesicles, usually on the trunk of middle-aged or older adults; heat, sweating, and occlusion may aggravate it.

### Grover's Disease

Alternate spelling; see *Grover Disease*.

### Growing Pains

Recurrent, bilateral leg pain in children ages 3-12, typically occurring in the late afternoon or evening and resolving by morning. Pain is deep, aching in the calves, shins, and thighs (not joints). No limping, fever, swelling, or functional impairment.

---

## Growth

The process of increasing in physical size and development, including physical, cognitive, and emotional maturation. In medicine, growth is monitored using growth charts and anthropometric measurements to assess health and development in children.

### Growth Chart

A graphical representation of growth (weight, height/length, head circumference, BMI) plotted against age, standardized by population (WHO growth standards 0-5 years; CDC growth charts 2-20 years; percentiles/z-scores).

### Growth Cone

A specialized, moving tip at the end of a growing nerve fiber. It senses chemical and physical signals and guides the axon toward its target during nervous-system development or repair.

### Growth Disorder

A condition in which a child or adult grows unusually slowly, quickly, or unevenly because of genetic, hormonal, nutritional, chronic, or other medical causes.

### Growth Factor

A protein that sends signals controlling cell growth, division, specialization, survival, or repair. Different growth factors act on different tissues and are involved in wound healing, blood-vessel formation, immunity, and cancer.

### Growth Failure

Growth that is slower than expected or weight gain that is inadequate for age. It can result from poor nutrition, chronic illness, hormone problems, genetic conditions, or difficulty absorbing nutrients.

### Growth Faltering

Alternate name; see *Faltering Weight*.

### Growth Hormone

A hormone released by the anterior pituitary that promotes growth partly through insulin-like growth

factor 1 and also affects protein, fat, and carbohydrate metabolism.

### Growth Hormone Deficiency

Insufficient secretion of growth hormone from the anterior pituitary, causing short stature in children and metabolic effects in adults. In children: delayed growth, short stature, delayed bone age, and increased fat mass. In adults: decreased muscle mass, increased fat mass, reduced bone density, fatigue, and dyslipidemia.

### Growth Hormone Deficiency in Adults

Inadequate growth-hormone secretion in adulthood, usually due to hypothalamic or pituitary disease or prior treatment. It may affect body composition, bone density, exercise capacity, lipids, and quality of life and requires confirmatory endocrine testing.

### Growth Hormone Physiology

Growth hormone is released in pulses by the pituitary gland, especially during sleep. It acts directly on tissues and also stimulates the liver to produce insulin-like growth factor 1, which supports bone and tissue growth.

### Growth Hormone Stimulation Test

A test for suspected growth-hormone deficiency. A medicine or other controlled stimulus is given, and blood samples are taken over time to see whether the pituitary gland releases enough growth hormone. Results depend on the protocol and laboratory.

#### Alternate Names:

GH Provocation Test; Insulin Tolerance Test for GH

### Growth Hormone Suppression Test

A test for excess growth hormone, usually performed after drinking a glucose solution. In healthy people, glucose lowers growth-hormone levels; failure to suppress them supports a diagnosis such as acromegaly.

#### Alternate Names:

GH Suppression Test; Glucose Loading GH Assay

---

### Growth Hormone Test

Laboratory measurement of growth hormone levels in the blood, used to evaluate GH deficiency or excess. Random GH levels have limited utility; stimulation or suppression tests are more informative.

#### Alternate Names:

Serum Growth Hormone; Somatotropin Assay

### Growth Hormone-Releasing Hormone

A 44-amino-acid hypothalamic peptide hormone synthesized by neurons of the arcuate nucleus that travels via the hypophyseal portal system to bind G-protein coupled receptors on anterior pituitary somatotropes, stimulating growth hormone synthesis and pulsatile secretion.

#### Alternate Names:

GHRH; Somatoliberin

### Growth Medium

A nutrient preparation supporting microbial multiplication in vitro, supplying carbon, nitrogen, minerals, and growth factors in liquid or solid form.

### Growth Plate

A layer of cartilage near the end of a growing bone where the bone lengthens. It gradually closes after puberty.

### Growth Plate Fracture

A break involving the cartilage growth area of a child's or adolescent's bone. It can disturb future growth if displaced or untreated; imaging and specialist management may be used.

#### Alternate Names:

Physeal Fractures; Salter-Harris Fractures

### Growth Plate Fractures

Injuries involving the cartilage growth plates at the ends of growing children's bones, with possible effects on future bone growth.

### Growth Retardation

Older wording for growth that is slower than expected for age or developmental stage. Possible

causes include inadequate nutrition, chronic illness, hormone or genetic disorders, poor absorption, and normal family variation; growth is assessed over time rather than by one measurement.

### Growth Scan

An ultrasound examination used to assess fetal size, growth pattern, amniotic fluid, and sometimes blood flow. Measurements are compared with gestational-age standards to identify possible growth restriction or excessive growth.

### Gtp Binding

GTP binding is the attachment of guanosine triphosphate to a protein, often switching that protein into an active state for cell signaling.

### Gtt.

An abbreviation from the Latin *guttae*, meaning drops. It refers to a liquid dose or the number of drops.

### Guaifenesin

An expectorant used to loosen respiratory mucus and make coughing more productive. It provides symptomatic relief in some respiratory illnesses; dosing depends on the formulation and patient.

### Guanfacine

Guanfacine is a centrally acting alpha-2A adrenergic agonist used for hypertension and, in extended-release form, ADHD; sedation, hypotension, bradycardia, and rebound hypertension after abrupt withdrawal are important risks.

### Guanine

Guanine is a purine nitrogenous base in DNA and RNA; it pairs with cytosine and is present in nucleotides such as GMP and GTP that support energy transfer and signaling.

### Guanosine Triphosphate

A guanine nucleotide that provides energy and acts as a molecular switch in protein synthesis, cytoskeletal regulation, and G-protein signaling.

---

### **Guanosinetriphosphatase**

An enzyme that hydrolyzes guanosine triphosphate to guanosine diphosphate and phosphate, regulating signaling and other cellular processes.

### **Guanylate Cyclase**

Guanylate cyclase is an enzyme that converts GTP to cyclic GMP, a second messenger involved in nitric-oxide signaling, smooth-muscle relaxation, photo-transduction, and intestinal fluid secretion.

### **Guarana**

A plant whose seeds contain caffeine and other stimulants. It is used in energy products and supplements; excessive intake may cause anxiety, insomnia, rapid heartbeat, high blood pressure, or interactions with medicines.

### **Guardianship**

A legal arrangement in which an appointed person makes decisions for someone unable to make them independently.

#### **Alternate Names:**

Legal Guardianship; Conservatorship

### **Guarding**

Involuntary tightening of the abdominal muscles when the abdomen is examined, often reflecting irritation or inflammation of the abdominal lining.

### **Guenther Porphria**

Alternate spelling; see *Gunther Disease*.

### **Guevedoces**

A term used in the Dominican Republic for individuals with 46,XY differences in sex development caused by 5-alpha-reductase type 2 deficiency; they may have female-appearing or ambiguous external genitalia at birth and increased virilization at puberty.

### **Guide Device**

A tool used to direct, position, or stabilize another instrument during a procedure, such as a guidewire, drill guide, or needle guide. It helps ensure accurate

placement and reduces injury to surrounding tissue.

### **Guided Imagery**

Guided imagery is a relaxation technique that uses imagined sights, sounds, or other sensations to promote calm and sometimes help manage pain or stress.

### **Guided Tissue Regeneration**

A dental procedure placing a barrier membrane to encourage regrowth of bone, root covering, and supporting ligament around a tooth. It may repair selected periodontal defects after inflammation is controlled.

### **Guillain-Barre Syndrome**

A rare disorder in which the immune system attacks peripheral nerves, often after an infection. Weakness and tingling usually begin in the legs and may spread upward; severe cases affect breathing or swallowing.

#### **Alternate Names:**

GBS; Acute Inflammatory Demyelinating Polyneuropathy (AIDP)

### **Guillain-Barré Syndrome**

An uncommon immune-mediated disorder in which peripheral nerves are attacked, usually causing rapidly progressive weakness and reduced reflexes. It can impair breathing or swallowing and is monitored closely in severe cases.

### **Guillaume Dupuytren**

A French surgeon whose name is associated with Dupuytren contracture, progressive thickening and shortening of palmar fascia that bends the fingers toward the palm.

### **Guinea Worm**

A parasitic infection caused by *Dracunculus medienensis*, acquired by drinking water containing infected copepods and characterized by emergence of a long worm through the skin.

### **Guinea Worm Infection**

Alternate name; see *Guinea Worm*.

---

## Gulf War Syndrome

A term used for a group of persistent symptoms reported by some Gulf War veterans, such as fatigue, pain, cognitive difficulties, gastrointestinal symptoms, and respiratory problems. It is also called Gulf War illness or chronic multisymptom illness; causes are complex and treatment focuses on individual symptoms and function.

## Gullet

The esophagus, a muscular tube carrying swallowed food and liquid from the throat to the stomach. It moves contents by coordinated squeezing and is protected from injury by its moist inner lining.

### Alternate Names:

Esophagus; Food Pipe

## Gum

The soft tissue, also called gingiva, that surrounds and protects the teeth and covers the jawbone. Healthy gums are usually firm and do not bleed with gentle brushing.

## Gum Biopsy

Histological examination of gingival tissue obtained by biopsy, used to diagnose gingival diseases, infections, or malignancies. May involve excisional or incisional biopsy techniques.

### Alternate Names:

Gingival Biopsy; Periodontal Tissue Sampling

## Gum Disease

Inflammation or infection affecting the gums and tissues supporting the teeth. Early disease may cause bleeding and swelling, while advanced disease can destroy bone, loosen teeth, and cause tooth loss.

### Alternate Names:

Periodontal Disease; Periodontitis

## Gum Problems

Gum problems include inflammation, bleeding, recession, ulcers, infection, trauma, and growths. Causes include plaque, tobacco, dry mouth, medicines, nutritional or immune disease, and poorly fitting appliances.

## Gumma

A soft area of tissue damage caused by late untreated syphilis. It can develop in the skin, bones, liver, or other organs and may cause a lump, pain, or organ damage.

### Alternate Names:

Syphilitic Gumma; Tertiary Syphilis Granuloma

## Gummata

More than one gumma: soft inflammatory masses caused by late untreated syphilis. They can affect skin, bone, liver, or other organs and may damage tissue.

## Gummy Smile

A smile showing more gum than usual. It may result from lip movement, tooth eruption, jaw shape, or gum position; treatment is optional and depends on the cause and the person's goals.

## Gums

The firm pink tissue, also called gingiva, surrounding and supporting the teeth. Healthy gums usually do not bleed with gentle brushing; swelling or bleeding can indicate gum disease.

### Alternate Names:

Gingiva; Gingival Mucosa

## Gunn Sign

An ophthalmologic sign of hypertensive retinopathy in which an arteriosclerotic, rigid retinal arteriole crosses and compresses an underlying retinal venule, producing visible tapering and apparent deflection of the venule on both sides of the crossing.

### Alternate Names:

Gunn's Sign; Arteriovenous Nicking

## Gunshot Wound

An injury caused by a firearm projectile. Damage may include bleeding, tissue destruction, fractures, nerve injury, organ damage, and infection.

### Alternate Names:

Firearm Injury; Ballistic Trauma; Bullet Wound

---

## Gunther Disease

A severe autosomal recessive disorder of heme biosynthesis caused by a deficiency of uroporphyrinogen III synthase, leading to massive accumulation of isomer I porphyrins in tissues, manifesting with extreme cutaneous photosensitivity, blistering, mutilating scar formation, erythrodontia, and hemolytic anemia.

### Alternate Names:

Congenital Erythropoietic Porphyria; CEP

## Gustation

The sense of taste. Special cells in taste buds detect sweet, salty, sour, bitter, and savory flavors. Most taste buds are on the tongue, with some in the soft palate and throat. Signals travel through several head nerves to the brain. Taste can change with infections, medicines, smoking, aging, or damage to the mouth or nerves.

## Gustatory

Relating to taste or the sense of taste, including taste buds, taste nerves, and the brain pathways that interpret flavors.

## Gustatory Lacrimation

Excessive tearing triggered by eating or smelling food, usually after an injury to the facial nerve or recovery from facial paralysis. The nerve signals for saliva and tears become incorrectly linked.

## Gustatory Sweating

Sweating triggered by eating or thinking about food, usually over the cheek or temple after injury or surgery involving the parotid region; it is also called Frey syndrome.

## Gut

Alternate informal term; see *Digestive Tract*.

## Gut Epithelium

The layer of cells lining the intestines. It absorbs nutrients and water, forms a barrier against harmful substances, and interacts with immune cells and gut microorganisms.

## Gut Microbiome

The complex community of trillions of microorganisms, including bacteria, viruses, and fungi, residing in the gastrointestinal tract. The gut microbiome plays a crucial role in digestion, immune function, vitamin synthesis, and protection against pathogens.

## Gut-Brain Axis

The two-way communication between the digestive tract and brain through nerves, hormones, immune signals, and substances made by gut microorganisms. It influences digestion, appetite, stress responses, and mood, but its clinical importance is still being studied.

## Guthrie Test

An older name for a newborn blood-spot screening test, originally used to detect phenylketonuria. Modern screening covers more conditions; an abnormal result is followed by confirmatory testing.

## Guttae

Small wart-like bumps on the inner surface of the cornea, the clear front window of the eye. Numerous or severe guttae can impair fluid removal, causing corneal swelling, blurred vision, glare, or eye pain.

## Guttate Psoriasis

A form of psoriasis causing many small, scaly, drop-shaped spots, often after a streptococcal throat infection. It commonly affects children or young adults and may clear, recur, or later become plaque psoriasis.

## Guttural

Relating to the throat. A guttural sound is produced mainly in the throat or back of the mouth rather than at the front of the tongue and lips.

## Guyon Canal Syndrome

An entrapment neuropathy of the ulnar nerve as it passes through Guyon canal (the fibro-osseous ulnar tunnel at the wrist between the pisiform and hook of the hamate), causing sensory loss over the little finger and medial half of the ring finger and motor weakness of the hypothenar and interosseous



---

muscles.

**Alternate Names:**

Ulnar Tunnel Syndrome; Handlebar Palsy

**GVHD**

Abbreviation; see *Graft Versus Host Disease*.

**Gymnast Wrist**

A chronic overuse repetitive stress injury of the distal radial growth plate (physis) occurring in skeletally immature gymnasts and athletes who subject the upper extremities to intense axial loading and hyperextension. Radiographs typically demonstrate distal radial physeal widening, irregular metaphyseal margins, and cystic changes.

**Alternate Names:**

Distal Radial Physeal Stress Syndrome; Gymnast's Epiphysitis

**Gynaecologist**

A medical doctor who specializes in the health of the female reproductive system (vagina, cervix, uterus, fallopian tubes, ovaries) and breasts.

**Gynaecology**

The medical specialty concerned with the female reproductive system, including the ovaries, fallopian tubes, uterus, cervix, vagina, and vulva, as well as related hormonal health.

**Gynaecomastia**

Alternate name; see *Gynecomastia*.

**Gynatresia**

Absence of or blockage in an opening of the female reproductive tract, present from birth or acquired later. If menstrual blood cannot leave the body, it may cause absent periods, pelvic pain, or enlargement of the uterus or vagina.

**Gynecic**

Relating to the female reproductive system or to gynecology.

**Gynecoid Pelvis**

A commonly described pelvic shape with a rounded inlet, relatively broad dimensions, and a spacious outlet; it is associated with typical female pelvic anatomy but individual variation is substantial.

**Gynecologic Pain**

Pain arising from the vulva, vagina, cervix, uterus, tubes, ovaries, pelvic floor, or related nerves and organs. Causes include infection, endometriosis, cysts, torsion, pregnancy complications, malignancy, musculoskeletal disease, and neuropathic or chronic pain.

**Gynecologic Tumor Markers**

Blood or tissue measurements that may support diagnosis, staging, treatment selection, or monitoring of gynecologic tumors. They are not usually diagnostic alone because benign conditions and other cancers can also alter marker levels.

**Gynecological Examination**

A gynecological examination assesses the female reproductive organs and may include inspection, a pelvic examination, and screening tests when appropriate.

**Gynecologist**

A gynecologist is a physician who provides care for the female reproductive system, including preventive care, diagnosis, and treatment.

**Gynecology**

The medical specialty concerned with the health of the female reproductive system (uterus, ovaries, fallopian tubes, cervix, vagina, and vulva).

**Gynecomastia**

A benign enlargement of male breast tissue caused by proliferation of glandular ductal components and stroma secondary to an increased ratio of estrogen to androgen activity. It commonly occurs as a physiological finding in neonates, pubertal adolescents, and older men. Pathological causes include primary or secondary hypogonadism (such as Klinefelter syndrome), liver cirrhosis, chronic kidney dis-

---

ease, estrogen-producing tumors, and medications such as spironolactone, antiandrogens, and anabolic steroids.

**Alternate Names:**

Gynaecomastia; Male Breast Enlargement

**Gyrate Atrophy**

An inherited retinal disorder causing gradually worsening peripheral and night vision loss. It results from abnormal ornithine metabolism and may be managed with dietary treatment, supplements, and regular eye monitoring.

**Gyrus**

A raised fold on the surface of the brain, separated from neighboring folds by grooves called sulci. These folds increase the cerebral cortex's surface area within the skull and contain areas responsible for different functions.

# H

## H And E Staining

A standard laboratory stain using hematoxylin and eosin to show cell nuclei, cytoplasm, and tissue structure under a microscope. It is widely used to examine biopsies and identify inflammation, infection, or tumors.

## H And H

Abbreviation for hemoglobin and hematocrit, two blood measurements commonly used to assess red-blood-cell levels and anemia.

## H And P

Abbreviation for a patient's history and physical examination. The history covers symptoms, past health, medicines, and relevant background; the examination checks the body for signs of illness.

## H Influenzae Meningitis

A serious infection of the coverings of the brain and spinal cord caused by *Haemophilus influenzae* bacteria. It may cause fever, headache, stiff neck, confusion, seizures, or brain injury, especially in unvaccinated children.

## H-Reflex

An electrophysiological monosynaptic reflex recorded during electromyography and nerve conduction studies, elicited by low-intensity submaximal electrical stimulation of sensory afferent fibers (group Ia) of a mixed peripheral nerve (such as the tibial nerve in the popliteal fossa). It bypasses the muscle spindle to evoke a delayed motor response in the corresponding muscle (soleus), providing a physiological measurement of

S1 radicular and spinal reflex arc integrity.

### Alternate Names:

Hoffmann Reflex; Electrophysiological Monosynaptic Reflex

## H. Pylori

Alternate spelling; see *Helicobacter Pylori*.

## H. Pylori Stool Antigen

A non-invasive enzymatic immunoassay that detects active gastrointestinal colonization by *Helicobacter pylori* by identifying bacterial cell-wall protein antigens in stool, recommended for primary diagnosis and post-treatment eradication confirmation.

### Alternate Names:

HpSA; Stool Antigen Test for H. Pylori

## H1N1 Flu

Alternate name; see *H1N1 Influenza*.

## H1N1 Influenza

Influenza A infection caused by an H1N1 virus. It spreads through respiratory droplets and commonly causes sudden fever, cough, sore throat, muscle aches, headache, and tiredness; some people develop pneumonia or severe breathing problems.

## H1N2

A subtype of influenza A virus that can infect humans and pigs. Human infections are uncommon and usually cause an influenza-like respiratory illness; surveillance and testing help distinguish it from other influenza viruses.

---

## H2 Blockers

Medicines that reduce stomach-acid production by blocking histamine H2 receptors in the stomach. They can relieve heartburn and treat some ulcers; possible adverse effects include headache, diarrhea, constipation, and interactions with other medicines.

### Alternate Names:

H2 Receptor Antagonists; Histamine-2 Receptor Blockers

## H2 Receptor Antagonists Overdose

Taking too much of an H2 blocker medicine. It may cause drowsiness, confusion, agitation, abnormal heartbeat, or rarely seizures.

## H5N1

A subtype of avian influenza A virus that mainly infects birds but can occasionally infect mammals and humans after close contact with infected animals or contaminated environments. Human illness can be severe.

## Habenula

A small brain structure helping process reward, unpleasant experiences, stress, motivation, and learned behavior. It connects several brain regions and may influence mood, sleep, pain, and responses to negative outcomes.

## Haberland Syndrome (Encephalocraniocutaneous Lipomatosis)

A rare congenital mosaic disorder characterized by unilateral scalp and facial lipomas or hairless skin lesions, eye abnormalities, and sometimes seizures, developmental differences, or brain malformations.

## Habit

A repeated behavior or pattern of action that may be ordinary, helpful, or harmful; in clinical assessment, its frequency, control, and effect on functioning determine whether it is a concern.

## Habituation

A gradual reduction in response to a repeated, harmless stimulus. For example, a person may stop

noticing a constant sound; habituation differs from addiction or drug tolerance.

## Habitus

A person's general body build and appearance as observed during a medical examination. Clinicians may describe body habitus when noting nutritional state, muscle mass, or features relevant to a condition.

## HAE

Abbreviation; see *Hereditary Angioedema*.

### Alternate Names:

C1-INH; C1 Inhibitor Deficiency

### Alternate Names:

HAE; Hereditary Angioneurotic Edema; HANE; C1-INH Deficiency Angioedema

## Haem

The iron-containing part of hemoglobin that binds oxygen inside red blood cells. Its structure allows blood to carry oxygen from the lungs to tissues and return carbon dioxide.

## Haemangioma

A usually harmless growth made of extra blood vessels. It is common in babies and may appear as a red or purple mark or lump.

## Haemarthrosis

Bleeding into a joint, causing swelling, pain, and reduced movement. It may follow an injury or occur with hemophilia, blood-thinning medicines, surgery, or certain joint disorders.

## Haematemesis

Vomiting blood or material that looks like coffee grounds, usually from bleeding in the esophagus, stomach, or upper small intestine. It can be life-threatening.

## Haematinics

Nutrients or medicines needed to make healthy red blood cells, especially iron, vitamin B12, and folate. Deficiency can cause anemia, although treatment

---

depends on the specific cause.

### **Haematocele**

A collection of blood in a body cavity or tissue, often in the scrotum after injury, surgery, or bleeding. It may cause swelling, pain, or pressure and sometimes needs imaging or drainage.

### **Haematocolpos**

Accumulation of menstrual blood in the vagina, usually because the vaginal opening is blocked or too narrow. It can cause absent periods, monthly pelvic pain, and a swollen vagina or uterus.

### **Haematocrit**

The proportion of blood volume occupied by red blood cells, expressed as a percentage or decimal fraction. The reference range varies with age, sex, altitude, pregnancy, and laboratory method.

### **Haematogenous**

Spread through the bloodstream. An infection, cancer, or other material that travels this way can reach organs or tissues far from its original site.

### **Haematologist**

A medical doctor who specializes in the diagnosis, treatment, and prevention of diseases of the blood and blood-forming organs.

### **Haematoma**

A collection of blood outside a blood vessel, usually caused by injury, surgery, bleeding disorders, or blood-thinning medicines. A hematoma may cause swelling and pain and can compress nearby organs or nerves.

### **Haematometra**

A collection of menstrual blood in the uterus, usually because the cervix or vagina is blocked or absent. It may cause missed periods, recurring pelvic pain, and uterine enlargement.

### **Haematomyelia**

Bleeding into the spinal cord, usually after severe

injury or because of an abnormal blood vessel or bleeding disorder. It can cause sudden back pain, weakness, numbness, paralysis, or loss of bladder control.

### **Haematopericardium**

Blood within the sac around the heart, usually after chest injury, heart surgery, or a tear in the heart or a blood vessel. It can compress the heart and cause life-threatening tamponade.

### **Haematopoiesis**

The process of making blood cells, mainly inside the bone marrow. It produces red cells that carry oxygen, white cells that fight infection, and platelets that help stop bleeding.

### **Haematoporphyrin**

An older term for a disorder involving porphyrins, substances used to make hemoglobin. Porphyrin disorders may cause sensitivity to sunlight, abdominal or nerve symptoms, or red-brown urine, depending on the type.

### **Haematosalpinx**

Blood collected inside a fallopian tube, often from ectopic pregnancy, endometriosis, pelvic injury, or blockage. It may cause pelvic pain and can affect fertility.

### **Haematuria**

Blood in the urine. It may be visible or found only with a urine test and can have many causes, including infection, stones, medicines, or kidney disease.

### **Haemochromatosis**

Alternate spelling; see *Hemochromatosis*.

### **Haemodialysis**

A treatment for severe kidney failure in which a machine filters waste products and extra fluid from the blood. It may be needed temporarily or regularly when the kidneys cannot do enough of this work.

---

## Haemoflagellate

A parasite with a whip-like structure that lives in blood or tissues. Examples include the parasites that cause sleeping sickness and Chagas disease.

## Haemoglobin

The iron-containing protein inside red blood cells that carries oxygen from the lungs to tissues and helps return carbon dioxide to the lungs. A low level is anemia; an abnormal form can cause disease.

## Haemoglobinopathy

An inherited disorder affecting the amount or structure of hemoglobin, the oxygen-carrying protein in red blood cells. Examples include sickle-cell disease and thalassemia, which can cause anemia or other complications.

## Haemoglobinuria

Free hemoglobin in the urine, often released when red blood cells break apart in the bloodstream. It can make urine red or brown and may occur with infections, medicines, burns, or blood disorders.

## Haemolysis

The breakdown of red blood cells, which releases hemoglobin into the blood. It can happen normally as old cells are removed or too quickly because of disease.

## Haemolytic Anaemia

Anemia caused by red blood cells being destroyed faster than the bone marrow can replace them. It may cause tiredness, jaundice, dark urine, rapid heartbeat, breathlessness, or an enlarged spleen.

## Haemolytic Disease Of The Newborn

Anemia and jaundice in a newborn caused by antibodies from the pregnant person crossing the placenta and destroying the baby's red blood cells. Severe cases can cause brain injury without prompt treatment.

## Haemoperitoneum

Bleeding into the abdominal cavity, commonly after

injury, a ruptured ectopic pregnancy, or a bleeding liver, spleen, artery, or other organ. It can cause shock.

## Haemophilia

Alternate spelling; see *Hemophilia*.

## Haemophilia A

Hemophilia A, an inherited bleeding disorder caused by deficiency or dysfunction of clotting factor VIII. It can cause prolonged bleeding, easy bruising, and bleeding into joints or muscles; treatment replaces or stimulates factor VIII activity.

## Haemophilus

A group of bacteria that may live harmlessly in the nose and throat or cause infection. Depending on the species, illness can affect the ears, sinuses, lungs, blood, joints, or brain coverings.

## Haemophilus Aegyptius

A bacterium associated with eye infections and, rarely, severe bloodstream disease. It can spread through respiratory secretions or direct contact, and laboratory identification helps guide treatment and public-health precautions.

## Haemophilus Ducreyi

A fastidious, facultatively anaerobic Gram-negative coccobacillus that is the causative etiological agent of chancroid, characterized by painful, non-indurated genital ulcers with purulent bases and prominent suppurative inguinal lymphadenopathy (buboes).

## Haemophilus Haemolyticus

A bacterium that commonly lives harmlessly in the upper airway but can occasionally cause respiratory or invasive infection, especially in people with weakened immunity or serious underlying illness.

## Haemophilus Infections

Infections caused by Haemophilus bacteria, which may affect the ears, sinuses, lungs, bloodstream, joints, or brain coverings. Severity depends on the



---

species, infection site, age, vaccination, and immune health.

### **Haemophilus Influenzae**

A type of bacteria that can cause ear infections, pneumonia, meningitis, and other infections. Type b, or Hib, can cause severe disease in children, but vaccination greatly reduces this risk.

### **Haemophilus Parainfluenzae**

A small gram-negative bacterium that normally colonizes the upper respiratory tract but can rarely cause respiratory, bloodstream, or other opportunistic infections.

### **Haemoptysis**

Coughing up blood or blood-stained mucus from the airways or lungs. Causes include infection, bronchiectasis, a blood clot, cancer, or irritation; larger amounts may occur with breathlessness or chest pain.

### **Haemorrhage**

British spelling of Hemorrhage; see Hemorrhage.

### **Haemorrhagic Fever**

A group of severe infections caused by certain viruses that can damage blood vessels and disrupt clotting. Fever, weakness, bleeding, shock, and organ failure may occur.

### **Haemorrhagic Fever With Renal Syndrome**

A hantavirus infection usually acquired by inhaling dust contaminated with infected rodent urine or droppings. It causes fever, headache, bleeding, low blood pressure, and temporary kidney failure and may be severe.

### **Haemorrhagic Stroke**

A stroke caused by a burst blood vessel in the brain or in the space around it. The bleeding injures brain tissue and may cause sudden weakness, speech trouble, severe headache, vomiting, or loss of consciousness.

### **Haemorrhoids**

Alternate spelling; see *Hemorrhoids*.

### **Haemosiderin**

A brownish iron-storage material that forms inside cells when iron accumulates, often after repeated bleeding, transfusions, or excess iron absorption. Large deposits can contribute to tissue damage.

### **Haemosiderosis**

Excess storage of iron-containing haemosiderin in tissues, often after repeated transfusions, bleeding into the lungs, or increased iron absorption. Effects depend on the organs involved and may include organ damage.

### **Haemostasis**

The body's normal process for stopping bleeding. Blood vessels narrow, platelets form a temporary plug, and clotting proteins strengthen it with a stable clot.

### **Haemostat**

A substance, device, or instrument used to control bleeding. It may work by pressing on a vessel, sealing tissue, promoting clotting, or temporarily blocking blood flow during an operation.

### **Haemothorax**

Blood collecting between the lung and chest wall, usually after chest injury, surgery, or a medical procedure. It can compress the lung, cause breathing difficulty or shock, and may require drainage.

### **Haff Disease**

Sudden muscle breakdown and severe muscle pain after eating certain contaminated fish or seafood. It can release muscle proteins that damage the kidneys; urgent treatment focuses on fluids and monitoring for complications.

### **Hafnia**

A genus of gram-negative intestinal bacteria; *Hafnia alvei* is the species most often identified in human specimens.

---

## Hafnia Alvei

A Gram-negative bacterium found in the intestine and environment that rarely causes opportunistic infection, usually in people with serious illness or weakened immunity.

## Hageman Factor

An older name for factor XII, a blood protein that helps start one pathway of clotting. Factor XII deficiency can prolong a laboratory clotting test but usually does not cause abnormal bleeding.

## Haglund Deformity

An anatomical prominence (posterosuperior calcaneal exostosis) of the heel bone causing mechanical friction and chronic inflammation of the retrocalcaneal bursa and Achilles tendon, exacerbated by rigid shoes ('pump bump').

### Alternate Names:

Retrocalcaneal Exostosis; Pump Bump

## Hailey-Hailey Disease

An inherited skin disorder that causes repeated blisters, raw areas, and painful cracks, especially in skin folds such as the neck, armpits, and groin. Heat, sweating, friction, and infection can trigger flares.

## Hair

Keratinized filaments produced by follicles in the skin. Hair protects the scalp, contributes to temperature regulation and sensation, and grows through anagen, catagen, and telogen phases.

## Hair Bleach Poisoning

Illness after swallowing or inhaling hair-bleaching chemicals, which may contain peroxide or persulfates. They can irritate or burn the mouth, throat, skin, eyes, or lungs.

## Hair Bulb

The rounded base of a hair follicle where new hair is produced. It contains growing cells and a small blood-supplied area that provides nutrients and helps determine hair growth and color.

## Hair Cells

Special cells in the inner ear that change sound vibrations and head movement into nerve signals for hearing and balance.

## Hair Disease

A disorder affecting hair growth, the hair shaft, or the scalp; causes include genetics, inflammation, infection, medicines, and nutritional problems.

## Hair Dye Poisoning

Harm caused by swallowing or inhaling hair-dye chemicals. Depending on the ingredients and amount, symptoms may include vomiting, swelling, breathing difficulty, allergic reaction, kidney injury, or muscle damage.

## Hair Follicle

A small tube in the skin that produces a hair. Each follicle has a growth phase, a brief transition phase, and a resting phase.

## Hair Loss

The partial or complete loss of hair from areas where it normally grows, most commonly the scalp. Causes include genetic predisposition (androgenetic alopecia), autoimmune conditions (alopecia areata), hormonal changes, medications, stress, and nutritional deficiencies.

### Alternate Names:

Alopecia; Baldness; Effluvium

## Hair Loss Classification

Hair loss is classified as scarring or nonscarring. Scarring hair loss destroys follicles and is usually permanent; nonscarring loss leaves follicles intact and may improve when its cause, such as illness, medicines, stress, or autoimmune disease, is treated.

## Hair Removal

Removal of unwanted hair by shaving, trimming, waxing, depilatory products, electrolysis, or laser treatment. Possible problems include irritation, burns, ingrown hairs, pigment changes, and infection.

---

## Hair Shaft

The part of a hair that extends above the skin. It is made of dead, hardened cells produced by the follicle and can be affected by cutting, heat, chemicals, friction, or damage to the hair root.

## Hair Spray Poisoning

Illness after swallowing or inhaling hair spray. Small accidental exposures may irritate the mouth or lungs, while larger exposures can cause vomiting, drowsiness, breathing problems, or chemical injury.

## Hair Straightener Poisoning

Harm after swallowing a hair-straightening product, which may contain corrosive chemicals. It can burn the mouth, throat, esophagus, or stomach and cause breathing or swallowing problems.

## Hair Tonic Poisoning

Illness after swallowing or inhaling a hair-tonic product. Ingredients such as alcohols or solvents may cause vomiting, drowsiness, low blood sugar, or breathing problems; the product and amount determine severity.

## Hair Transplant

A procedure that moves hair follicles, usually from a donor area of the scalp, to areas affected by permanent hair loss; results depend on diagnosis, donor supply, technique, and postoperative care.

## Hair-Ball

A hair-ball, or trichobezoar, is a mass of swallowed hair in the stomach or intestine, usually associated with repeated hair eating.

## Hair-Pulling Syndrome

Alternate name; see *Trichotillomania*.

## Hairy Cell Leukaemia

Hairy cell leukaemia is a rare slow-growing blood cancer in which abnormal B lymphocytes accumulate in the bone marrow, spleen, and blood.

## Hairy Cell Leukemia

A rare, usually slow-growing cancer of a type of white blood cell. Abnormal cells build up in the bone marrow, blood, and spleen, causing low blood counts, infections, tiredness, or an enlarged spleen.

### Alternate Names:

HCL; Leukemic Reticuloendotheliosis

## Hairy Leukoplakia

A benign, painless, white, corrugated mucosal lesion occurring predominantly along the lateral margins of the tongue that cannot be scraped off, caused by opportunistic infection of epithelial cells by Epstein-Barr virus (EBV) in immunocompromised patients (especially HIV infection).

### Alternate Names:

Oral Hairy Leukoplakia; OHL

## Hairy Tongue

A usually harmless overgrowth and discoloration of the tongue's papillae, often appearing black, brown, or white. It may be associated with smoking, poor oral hygiene, dry mouth, or antibiotics; gentle tongue cleaning and removal of contributing factors usually help.

## Hajdu-Cheney Syndrome

A rare inherited bone disorder causing fragile bones, short stature, abnormal skull and facial development, and sometimes loose joints or dental problems. Severity varies, and specialist monitoring helps prevent complications.

## Halcinonide

A potent topical corticosteroid used to reduce inflammation and itching in steroid-responsive dermatitis and other inflammatory skin disorders; prolonged or extensive use can cause skin atrophy and systemic effects.

## Haldane Effect

The effect of oxygen binding to hemoglobin in the lungs, which helps release carbon dioxide for exhalation. In body tissues, hemoglobin without oxygen carries more carbon dioxide back toward the lungs.

---

## Half-Life

The time it takes for the amount of a medicine in the body, or its blood level, to decrease by half. It helps determine how often a medicine should be taken and how long it remains in the body.

### Alternate Names:

Terminal Half-Life;  $t_{1/2}$

## Halitosis

Persistent unpleasant breath odor, most often from bacteria on the tongue, teeth, or gums. Dry mouth, smoking, foods, sinus or throat disease, reflux, and some illnesses or medicines can also contribute.

## Hallermann's Syndrome

A rare congenital disorder characterized by distinctive craniofacial development, sparse hair, dental abnormalities, and sometimes eye or skeletal findings; it is also called Hallermann–Streiff syndrome.

## Hallpike Test

A clinical vestibular provocative maneuver used to diagnose benign paroxysmal positional vertigo (BPPV) of the posterior semicircular canal; the patient is rapidly lowered from a sitting position to supine with the head extended and turned 45 degrees, provoking rotational upbeat nystagmus and vertigo.

### Alternate Names:

Dix-Hallpike Test; Nylan-Barany Test

## Hallucination

A sensory experience that feels real but occurs without a matching external stimulus. It may involve seeing, hearing, feeling, smelling, or tasting something that is not present.

## Hallucinations

Hallucinations are perceptions without an external sensory stimulus and can arise from psychiatric illness, neurologic disease, delirium, substances, medications, sensory loss, or sleep transitions.

## Hallucinogen

A psychoactive substance that can alter perception,

thought, and mood and may produce hallucinations; effects and toxicity vary by substance, dose, co-ingestants, and mental state.

## Hallux

The great toe, or big toe. It has two bones and helps the foot stay stable and push off during walking. Common problems include hallux valgus (a bunion), hallux rigidus (a stiff, arthritic joint), turf toe (a sprain), and gout causing sudden pain and swelling at the joint where the toe meets the foot.

## Hallux Rigidus

Arthritis of the joint where the big toe meets the foot, causing pain, stiffness, swelling, and reduced upward movement. A bony bump may develop and make walking or wearing shoes difficult.

## Hallux Valgus

A bunion: a change in the alignment of the joint at the base of the big toe, with the toe pointing inward and a prominent area on the inner forefoot. It may become painful from pressure or inflammation.

### Alternate Names:

Bunion; Hallux Abductovalgus

## Hallux Varus

Abnormal medial deviation of the great toe, often following surgery, trauma, neuromuscular imbalance, or congenital deformity; it can cause shoe difficulty and pressure-related pain.

## Halo Nevus

A usually harmless mole surrounded by a ring of lighter skin. The immune system causes the mole and nearby pigment cells to lose color; it often occurs in children or young adults and may slowly fade.

## Halo Sign

A chest computed tomography finding characterized by a ground-glass attenuation perimeter surrounding a dense pulmonary nodule or consolidation, representing alveolar hemorrhage surrounding a fungal focus, highly suggestive of early invasive aspergillosis in neutropenic patients.

---

## Halogen

An element in group 17 of the periodic table, including fluorine, chlorine, bromine, iodine, and astatine; halogens and their compounds have medical, biologic, and toxicologic importance.

## Haloperidol

Haloperidol is a first-generation antipsychotic that blocks dopamine D2 receptors and is used for psychosis, agitation, delirium, and tics; extrapyramidal reactions, tardive dyskinesia, hyperprolactinaemia, and QT prolongation are important risks.

## Halothane

A volatile inhaled anesthetic formerly used to induce or maintain general anesthesia; it can cause hypotension, arrhythmias, and rare severe immune-mediated liver injury.

## Halushi-Behcet's Syndrome

Alternate name; see *Behcet Disease*.

## Hamamelis

A botanical preparation from witch hazel used topically as an astringent for minor skin or mucosal irritation; evidence and product composition vary.

## Hamartoma

A usually harmless growth made of an abnormal arrangement of tissue types normally found in that part of the body. It may occur in organs such as the lung, skin, brain, or intestine and may cause problems if it grows or presses on nearby structures.

## Hamartomatous Polyposis

A condition in which multiple hamartomatous gastrointestinal polyps develop from disorganized but benign tissue overgrowth, sometimes as part of an inherited cancer-predisposition syndrome.

## Hamate

A small, wedge-shaped bone on the little-finger side of the wrist. Its hook provides attachment for ligaments and tendons; fractures there may cause wrist pain or pressure on a nearby nerve.

## Hamate Bone

The hamate is a wedge-shaped wrist bone on the little-finger side that has a hook serving as an attachment and passage point for tendons and nerves.

## Hamman Sign

A crunching or clicking noise heard with a stethoscope over the chest in time with the heartbeat, caused by air trapped in the center of the chest.

## Hamman's Sign

A crackling or crunching sound heard over the chest in time with the heartbeat, usually caused by air in the central chest around the heart and major vessels. It can occur with pneumomediastinum and needs clinical assessment.

## Hamman-Rich Syndrome

An acute, fulminant idiopathic interstitial pneumonia characterized by sudden onset of cough, rapidly progressive dyspnea, hypoxemic respiratory failure, and diffuse alveolar damage on lung biopsy in the absence of a known cause or preexisting lung disease.

### Alternate Names:

Acute Interstitial Pneumonia; AIP

## Hammer

The common name for the malleus, one of the three small bones of the middle ear.

## Hammer Toe

A deformity in which a lesser toe bends downward at its middle joint and may press against the shoe. It can cause pain, stiffness, corns, or calluses and may be flexible at first or become fixed.

## Hammer Toe Repair

Surgery to correct a fixed or painful flexion deformity of a lesser toe, using soft-tissue release, tendon balancing, bone correction, or temporary fixation as indicated.

## Hammertoe

A deformity in which a toe bends downward at its middle joint, giving it a claw-like appearance. It may

---

cause pain, stiffness, corns, or calluses, especially when wearing shoes.

### **Hammertoes**

Alternate plural form; see *Hammertoe*.

### **Hampton Hump**

Wedge-shaped opacification with its base abutting the pleura on chest radiography, representing pulmonary infarction secondary to Pulmonary Embolism. See Pulmonary Embolism (PE).

### **Hampton Line**

A distinct radiological finding identified on a barium meal examination of the stomach, consisting of a thin, sharply demarcated radiolucent line measuring approximately one millimeter in width that traverses the neck of a benign gastric ulcer crater. It represents the undermined mucosal edge of a benign peptic ulcer, distinguishing it from an ulcerated gastric malignancy.

#### **Alternate Names:**

Hampton's Radiolucent Line; Peptic Hampton Line

### **Hampton's Hump**

Alternate spelling; see *Hampton Hump*.

### **Hamstring**

One of three muscles forming the posterior compartment of the thigh: semitendinosus, semimembranosus, and biceps femoris (long and short heads). Common origin: ischial tuberosity (except short head of biceps femoris which originates from the linea aspera of the femur).

### **Hamstring Injury**

A strain or tear of one or more muscles or tendons at the back of the thigh.

### **Hamstring Muscles**

The muscles at the back of the thigh. They bend the knee and help move the hip, especially during walking, running, and jumping. A sudden stretch or forceful contraction can strain or tear them.

### **Hamstring Strain**

Alternate name; see *Hamstring Injury*.

### **Hand**

The part of the upper limb beyond the wrist, containing the palm, fingers, bones, joints, muscles, nerves, and blood vessels used for grasping and sensation.

### **Hand Deformity**

An abnormal shape, alignment, or loss of motion of the hand or fingers caused by congenital difference, trauma, arthritis, nerve dysfunction, tendon disease, or other conditions.

### **Hand Foot And Mouth Disease**

A contagious viral illness, most common in young children, that causes fever, painful mouth sores, and a rash or small blisters on the hands, feet, and sometimes buttocks. It spreads through close contact and contaminated secretions or stool.

### **Hand Hygiene**

Cleaning the hands with soap and water or an alcohol-based hand rub to remove or kill germs. It is especially important before and after patient contact, after touching body fluids, and when hands are visibly dirty.

### **Hand Infections**

Infections involving the skin, tendon sheaths, joints, bones, or deep spaces of the hand. Pain, swelling, redness, impaired movement, fever, or a flexed finger may indicate a condition requiring urgent assessment.

### **Hand Microsurgery**

Specialized hand surgery performed with magnification and fine instruments to repair or reconstruct small blood vessels, nerves, and other delicate tissues. It may be used for complex injuries, replantation, and restoration of blood flow to a threatened digit or hand.



---

## Hand of Benediction

An abnormal hand posture observed when a patient attempts to make a fist, resulting in extension of the index and middle fingers with flexion of the fourth and fifth fingers. It occurs due to proximal median nerve injury, which paralyses the flexor digitorum superficialis and the lateral half of the flexor digitorum profundus.

### Alternate Names:

Benediction Sign; Preacher's Hand

## Hand Or Foot Spasms

Sudden, involuntary tightening of muscles in a hand or foot. Causes include overuse, dehydration, abnormal calcium or other salts, nerve disease, medicines, and reduced blood flow; repeated or painful spasms need assessment.

## Hand Pain

Pain or discomfort in the hand, fingers, or wrist caused by injury, arthritis, nerve compression, tendon problems, infection, or other conditions. Sudden severe pain, deformity, numbness, or loss of circulation needs prompt assessment.

## Hand Prolapse

A fetal hand or arm lying beside or below the part leading into the birth canal. It can interfere with labor or reduce blood flow to the limb; the position requires prompt obstetric assessment and should not be pushed back at home.

## Hand X-Ray

Radiographic imaging of the hand used to evaluate fractures, dislocations, arthritis, infection, tumors, bone age, and other abnormalities of the bones and joints.

## Hand-Arm Vibration Syndrome

An occupational disorder from prolonged exposure to vibrating tools, causing numbness, tingling, reduced grip, and episodic whitening of the fingers from vascular spasm.

## Hand-Foot Syndrome

A dose-limiting toxic cutaneous reaction triggered by certain systemic antineoplastic and targeted chemotherapy agents (notably capecitabine, 5-fluorouracil, pegylated liposomal doxorubicin, cytarabine, and multi-kinase inhibitors like sorafenib and sunitinib). It is characterized by progressive bilateral burning dysesthesia, tingling, painful symmetric erythema, swelling, and subsequent blistering or desquamation primarily localized to the friction-bearing pressure areas of the palms and soles.

### Alternate Names:

Palmar-Plantar Erythrodysesthesia; PPE; Chemotherapy-Induced Acral Erythema; Burgdorf Reaction

## Hand-Foot-And-Mouth Disease

A contagious enteroviral illness, most common in young children, causing fever, painful mouth sores, and a rash or small blisters on the hands and feet, sometimes the buttocks. It spreads through respiratory secretions, blister fluid, contaminated surfaces, and stool and is usually self-limited.

## Hand-Foot-And-Mouth Syndrome

A usually mild, contagious viral illness, commonly caused by enteroviruses, producing fever, mouth sores, and small spots or blisters on the hands and feet. It mainly affects young children but can occur in adults. Hydration, pain relief, and hygiene are important; severe dehydration, neurologic symptoms, or persistent fever requires care.

## Hand-Foot-Mouth Disease

A common enteroviral infection, usually in young children, causing fever, painful oral ulcers, and a vesicular rash on the hands, feet, and sometimes buttocks. It is highly contagious and usually self-limited.

## Hand-Held Retractor

A handheld surgical instrument used to hold tissue or an organ away from the operative field.

### Alternate Names:

Manual Retractor; Handheld Retractor; Senn Retractor

---

### Hand-Schuller-Christian Disease

A multifocal, chronic form of Langerhans cell histiocytosis occurring predominantly in young children, classically defined by the clinical triad of multifocal lytic bone lesions ('punched-out' skull defects), diabetes insipidus, and exophthalmos.

#### Alternate Names:

Multifocal Unisystem LCH; Schuller-Christian Triad

### Hand-Schüller-Christian Disease

Hand-Schüller-Christian disease is an older name for a multisystem form of Langerhans cell histiocytosis, often involving bone, skin, and the pituitary gland.

### Handling Sharps And Needles

Handling sharps and needles means using, passing, disposing of, and storing sharp medical instruments safely to prevent needlestick injuries and infection transmission.

### Handwashing

Handwashing is cleaning the hands with soap and water or an alcohol-based product to remove germs and reduce transmission of infection.

### HANE

Historical abbreviation for hereditary angioneurotic edema, an older name for hereditary angioedema; see *Hereditary Angioedema*.

### Hang-Nail

A small torn flap of skin beside a fingernail or toenail, often caused by dryness or trauma and vulnerable to local infection if torn or contaminated.

### Hanging And Strangulation Signs

Injuries or other findings caused by pressure around the neck that restricts breathing or blood flow. Findings may include neck marks, swelling, voice changes, trouble swallowing, breathing difficulty, confusion, or loss of consciousness; urgent medical assessment is required.

### Hanging Injuries and Strangulation

Injuries caused by constriction of the neck or sus-

pension of the body, with potential airway, vascular, neurologic, and cervical-spine damage. Serious internal injury may occur even when external findings are limited.

### Hangman Fracture

A traumatic bilateral fracture passing through the neural arch, pedicles, or pars interarticularis of the second cervical vertebra (axis, C2), classically caused by sudden violent hyperextension and axial loading or distraction of the cervical spine.

#### Alternate Names:

Traumatic Spondylolisthesis of the Axis; C2 Pedicle Fracture

### Hangman's Fracture

A break through the second neck bone caused by forceful bending or straightening of the neck, often in a crash or fall. It can make the spine unstable and may injure the spinal cord, so urgent immobilization and imaging are needed.

#### Alternate Names:

Axis Pedicle Fracture; C2 Bilateral Pars Fracture

### Hangover

A group of symptoms after alcohol use, such as headache, nausea, thirst, fatigue, dizziness, and sensitivity to light; symptoms reflect effects of alcohol and its metabolites, dehydration, sleep disruption, and inflammation.

### Hansen Disease

A long-lasting infection caused by *Mycobacterium leprae* that mainly affects the skin and peripheral nerves. Pale or reddish numb patches, weakness, and eye problems may occur; multidrug antibiotics cure infection and prevent disability.

#### Alternate Names:

Leprosy; Hanseniasis; *Mycobacterium Leprae* Infection

### Hantavirus

A group of viruses spread mainly by breathing dust contaminated with infected rodent urine, droppings, or nesting material. They can cause severe lung disease or a fever that damages the kidneys.

---

## Hantavirus Cardiopulmonary Syndrome

A severe zoonotic infection that can progress from fever and muscle aches to capillary leak, pulmonary edema, hypoxemic respiratory failure, shock, and cardiac dysfunction.

## Hantavirus Infection

A zoonotic viral infection acquired mainly through inhalation of contaminated rodent excreta. It may cause hemorrhagic fever with renal syndrome or hantavirus pulmonary syndrome, which can progress rapidly to respiratory failure.

## Hantavirus Pulmonary Syndrome

A severe lung infection acquired mainly by inhaling dust contaminated with infected rodent urine, droppings, or nesting material. Fever, muscle aches, headache, and stomach symptoms can rapidly progress to fluid-filled lungs and respiratory failure.

### Alternate Names:

HPS; Sin Nombre Virus Infection

## Haploid

Having one complete set of chromosomes, as in a human sperm or egg. Haploid cells contain half the chromosome number of most body cells and restore the full number when they join during fertilization.

## Haploidentical Donor

A donor who shares one human leukocyte antigen haplotype with the recipient, commonly used for selected hematopoietic stem-cell or organ transplantation when a fully matched donor is unavailable.

## Haploidy

The state of having one complete set of chromosomes, as in human gametes; haploid cells contain 23 chromosomes.

## Haploinsufficiency

A genetic state in which one functional copy of a gene does not produce enough gene product for normal function, so loss or inactivation of the other copy can cause disease.

## Haplotype

A group of nearby gene versions inherited together on one chromosome. Haplotype information can help study inherited disease risk, identify tissue donors, and trace how genetic traits are passed through families.

## Hapten

A hapten is a small molecule that is not immunogenic by itself but can trigger an immune response after binding to a larger carrier protein, as in some drug allergies.

## Haptoglobin

A blood protein made mainly by the liver that binds free hemoglobin released when red blood cells break down. Low levels can support evaluation of hemolysis, but liver disease and inflammation also affect the result.

## Haptoglobin Analyses

Laboratory tests measuring haptoglobin, a plasma protein that binds free hemoglobin; low values can support evaluation of intravascular hemolysis but are also affected by liver disease and inflammation.

## Haptoglobin Blood Test

A blood test measuring haptoglobin, a protein that binds free hemoglobin; low levels may support intravascular hemolysis but can also reflect liver disease or inflammation.

## Haptoglobin Decreased

A low blood haptoglobin result, commonly caused by intravascular red-cell destruction, reduced hepatic production, or severe liver disease; it is interpreted with hemoglobin, bilirubin, reticulocytes, and a blood smear.

## Hard Palate

The bony, mucosa-covered front portion of the roof of the mouth that separates the oral and nasal cavities and provides a surface for chewing and speech.

---

## Harlequin Color Change

A temporary, harmless change in a newborn's skin color in which one side becomes red while the other remains pale. It often appears when the baby lies on one side and usually fades within minutes.

## Harlequin Foetus

Older term; see *Harlequin Ichthyosis*.

## Harlequin Ichthyosis

A rare severe inherited skin disorder in which newborns have thick plates of skin separated by deep cracks. It can impair temperature control, movement, feeding, breathing, and protection from infection and requires intensive neonatal care.

## Harm Reduction

Evidence-based strategies that decrease health, social, and legal harms associated with substance use without requiring immediate abstinence. Examples include naloxone access, sterile injecting equipment, drug-checking where available, safer-use education, and linkage to treatment.

## Harm Reduction Therapy

Care that reduces illness, injury, infection, overdose, and other harms related to substance use or another risky behavior, even when stopping immediately is not possible. It can include safer-use support and treatment for dependence.

### Alternate Names:

Harm Reduction; Risk Mitigation Strategy

## Harmless Murmur

An innocent or physiologic heart murmur produced by normal blood flow without structural heart disease; it is diagnosed only after appropriate clinical assessment.

## Harrison's Sulcus

Harrison's sulcus is a horizontal groove along the lower chest wall caused by inward pulling of the ribs, classically in long-standing childhood breathing difficulty.

## Hartmann Pouch

An anatomical outpouching or dilatation of the inferior wall of the gallbladder neck adjacent to the junction with the cystic duct; gallstone impaction here may cause acute calculous cholecystitis or compress the adjacent common hepatic duct (Mirizzi syndrome).

### Alternate Names:

Infundibulum of the Gallbladder

## Hartmann Procedure

An operation that removes the diseased lower colon, brings the remaining colon through the abdominal wall as a colostomy, and closes the downstream bowel. It is often used when infection, blockage, perforation, or severe illness makes immediate reconnection unsafe.

## Hartnup Disease

A rare inherited problem absorbing and reusing certain amino acids. It may cause a sun-sensitive rash, poor coordination, or mood and thinking changes, although some people have no symptoms.

## Hartnup Disorder

An inherited disorder of neutral amino-acid transport that can cause pellagra-like skin changes, neurologic symptoms, or no symptoms.

## HAS-BLED Score

A score estimating bleeding risk in a person with atrial fibrillation who may need a blood thinner. It considers high blood pressure, kidney or liver disease, previous stroke or bleeding, unstable clotting results, age, medicines, and alcohol; a high score prompts closer monitoring and correction of changeable risks.

### Alternate Names:

HAS-BLED; Anticoagulant Bleeding Risk Index

## Hashimoto Disease

An autoimmune disorder in which antibodies and lymphocytes progressively damage the thyroid, commonly causing hypothyroidism, goiter, fatigue, cold intolerance, constipation, and weight change.

---

## Hashimoto Encephalopathy

A rare, steroid-responsive neurocognitive encephalopathy associated with autoimmune thyroiditis, characterized by fluctuating cognitive impairment, stroke-like episodes, myoclonus, seizures, and high titers of antithyroid antibodies (anti-TPO) independent of thyroid functional status.

### Alternate Names:

Steroid-Responsive Encephalopathy Associated with Autoimmune Thyroiditis; SREAT

## Hashimoto Thyroiditis

An autoimmune disorder in which immune-mediated inflammation gradually damages the thyroid gland and may cause hypothyroidism. The gland may be enlarged or atrophic, and thyroid autoantibodies are often present.

## Hashimoto's Disease

An autoimmune disorder in which the immune system gradually damages the thyroid, often causing low thyroid-hormone levels. It may produce tiredness, cold intolerance, weight change, constipation, or an enlarged thyroid.

### Alternate Names:

Hashimoto Thyroiditis; Chronic Autoimmune Thyroiditis

## Hashimoto's Thyroiditis

A chronic autoimmune inflammation of the thyroid that commonly causes progressive hypothyroidism and may produce a firm, enlarged thyroid gland.

## Hashish

A concentrated form of cannabis made from resin of the cannabis plant. It contains cannabinoids that can impair memory, coordination, and judgment and may cause anxiety, dependence, or psychosis in susceptible people.

## Hashitoxicosis

A temporary period of excess thyroid hormone caused by release of stored hormone during Hashimoto thyroiditis.

## Hassall Corpuscles

Concentrically laminated, eosinophilic epithelial structures located within the medulla of the human thymus gland, composed of degenerate thymic epithelial cells that secrete thymic stromal lymphopoietin (TSLP) and assist in regulatory T cell maturation.

### Alternate Names:

Thymic Corpuscles; Hassall's Bodies

## Hassall's Corpuscles

Hassall's corpuscles are rounded structures in the thymus made of specialized epithelial cells and involved in the development of immune tolerance.

## Haustra

The series of small pouches along the outside of the large intestine. They form as the bowel muscles contract and help the colon mix and move its contents.

## Haverhill Fever

An illness caused by *Streptobacillus* bacteria, usually after drinking contaminated milk or water. It may cause fever, vomiting, headache, muscle or joint pain, and a rash and requires antibiotic treatment.

## Haversian Canal

A central longitudinal microscopic channel running through the core of each osteon (Haversian system) in compact cortical bone, carrying neurovascular structures (capillaries, postcapillary venules, and unmyelinated nerve fibers) to supply osteocytes embedded within surrounding concentric lamellae.

### Alternate Names:

Central Canal of Bone

## Hawkins Test

A physical examination maneuver used to identify subacromial shoulder impingement; the examiner flexes the patient's arm and elbow to 90 degrees and forcefully performs internal rotation of the humerus, compressing the supraspinatus tendon against the coracoacromial ligament.

### Alternate Names:

---

Hawkins-Kennedy Impingement Test; Hawkins Sign

### Hay Fever

A common name for seasonal allergic rhinitis, an allergic response to airborne outdoor plant pollens and mold spores that causes sneezing, a blocked or runny nose, an itchy palate, and watery, irritated eyes. See also *Seasonal Allergies*.

#### Alternate Names:

Seasonal Allergies; Seasonal Allergic Rhinitis; Pollinosis

### Hazardous Materials

Substances that can cause injury, poisoning, fire, radiation exposure, or environmental harm; safe handling requires labeling, protective equipment, exposure control, and emergency procedures.

### Hazardous Substance

A chemical, biological agent, or other material that can harm health through swallowing, inhalation, skin contact, injection, radiation, or environmental exposure. Risk depends on its properties, amount, route, and duration of exposure.

### Hb

Hemoglobin, the iron-containing protein in red blood cells that carries oxygen and contributes to carbon-dioxide transport and blood's acid-base buffering.

### HbA1c

Abbreviation; see *Glycated Hemoglobin*.

#### Alternate Names:

Glycated Hemoglobin; Hemoglobin A1c; A1c Test

### HCG

Human chorionic gonadotropin is a hormone produced mainly during pregnancy; its measurement is used in pregnancy testing and evaluation of selected tumors.

### HCG In Urine

A urine test detecting human chorionic gonadotropin, commonly used as a pregnancy test;

accuracy depends on timing, urine concentration, and test performance.

### Hct

The percentage of blood volume occupied by red blood cells, measured as part of a complete blood count.

### HCV PCR

A polymerase chain reaction test that detects or measures hepatitis C virus RNA in blood, helping identify active infection and monitor treatment response.

### HCV-Associated Rheumatic Disease

Joint, muscle, blood-vessel, or autoimmune problems associated with hepatitis C virus infection. Possible manifestations include joint pain or inflammation, mixed cryoglobulinemia, vasculitis, fatigue, or muscle symptoms.

### HD

Abbreviation; see *Huntington Disease*.

#### Alternate Names:

Hearing Impairment; Hypoacusis

### HDL

High-density lipoprotein, a class of particles that transports cholesterol and other lipids in the blood. HDL concentration is one component of cardiovascular risk assessment; it should not be interpreted as a protective or harmful measure by itself.

### HDL Cholesterol

Cholesterol carried in high-density lipoprotein particles; HDL participates in transporting cholesterol from tissues toward the liver, but its concentration alone does not determine cardiovascular risk.

### HDL Test

A blood test measuring high-density lipoprotein cholesterol, which is used with the rest of the lipid profile to assess cardiovascular risk.



---

## HDR Syndrome

A genetic disorder characterized by hypoparathyroidism, sensorineural deafness, and renal dysplasia or other kidney abnormalities, usually caused by a pathogenic variant in *GATA3*.

## Head

The body region containing the brain, eyes, ears, nose, mouth, skull, and face, connected to the trunk by the neck and supplied by major cranial nerves and vessels.

## Head And Face Reconstruction

Reconstructive surgery to restore the form or function of the skull, face, or associated soft tissues after trauma, cancer treatment, congenital difference, or other disease.

## Head And Neck Cancer

Cancers arising in the oral cavity, pharynx, larynx, sinonasal tract, and salivary glands, most commonly squamous cell carcinoma. Risk factors include tobacco, alcohol, and HPV (oropharyngeal). Treatment involves surgery, radiation, and chemotherapy, often combined to preserve function.

## Head And Neck Oncologic Surgery

Surgical treatment of cancers of the oral cavity, throat, larynx, nose, sinuses, salivary glands, or neck lymph nodes. Procedures may remove the tumor, treat involved lymph nodes, and reconstruct affected tissues while preserving breathing, swallowing, speech, and appearance when possible.

### Alternate Names:

Head and Neck Surgical Oncology; Maxillofacial Oncologic Surgery

## Head Circumference

The measurement around the largest part of a baby's or child's head, taken above the eyebrows and ears. It is plotted over time to assess brain growth; unusually small, large, or rapidly changing measurements may need evaluation.

## Head CT Scan

Computed tomography of the head using x-rays and computer reconstruction to evaluate the brain, skull, bleeding, stroke, tumors, trauma, and other abnormalities.

## Head Impulse Test

A bedside examination of the vestibulo-ocular reflex (VOR); the clinician delivers rapid, unpredictable, small-amplitude passive rotations of the patient's head toward one side while the patient fixates on a central target; a corrective saccade indicates ipsilateral peripheral vestibular hypofunction.

### Alternate Names:

Halmagyi-Curthoys Test; Video Head Impulse Test (vHIT)

## Head Injury

Damage to the scalp, skull, or brain caused by a blow, fall, collision, or penetrating object. It may cause a wound, skull fracture, concussion, bleeding, or brain damage. Worsening headache, repeated vomiting, seizure, confusion, weakness, or loss of consciousness may occur.

### Alternate Names:

Traumatic Brain Injury (TBI); Craniocerebral Trauma

## Head Injury First Aid

Immediate emergency first-aid protocols following head trauma, including stabilizing the cervical spine, maintaining an open airway, controlling external cranial bleeding with gentle pressure, and continuously monitoring level of consciousness and vital signs while urgently seeking medical evaluation for warning signs such as loss of consciousness, repeated vomiting, or focal neurologic deficits.

### Alternate Names:

Cranial Trauma First Aid; Head Trauma Immediate Care; Concussion First Aid

## Head Lice

Infestation of the scalp by the parasitic insect *Pediculus humanus capitis*. It causes itching and visible nits or lice and spreads mainly through close head-to-head contact; treatment requires an effective

---

tive product and careful follow-up.

### Head Movement

Motion of the head produced mainly by the cervical spine and neck muscles, including flexion, extension, rotation, and side bending; pain or restricted movement may indicate injury or disease.

### Head MRI

Magnetic resonance imaging of the head using a magnetic field and radio waves to produce detailed images of the brain, cranial nerves, blood vessels, and surrounding structures.

### Head Of The Epididymis

The upper part of the epididymis, connected to the testis by small ducts and receiving newly made sperm.

### Head Thrust Test

A bedside neuro-otological examination of the vestibulo-ocular reflex (VOR) performed by rapidly rotating the patient's head horizontally while the patient fixates on the examiner's nose; a corrective catch-up saccade indicates peripheral vestibular hypofunction (such as vestibular neuritis).

#### Alternate Names:

Head Impulse Test; Halmagyi-Curthoys Test

### Head Trauma

An injury to the scalp, skull, brain, or other tissues of the head. Concussion, bleeding, skull fracture, and brain injury can result; worsening headache, repeated vomiting, seizure, confusion, weakness, or loss of consciousness may occur.

### Head-Lice

Head lice are small insects that live on the scalp and cause itching; they spread mainly through close head-to-head contact.

### Headache

Pain or discomfort in the head, scalp, or upper neck. Primary headaches include migraine and tension headache; secondary headaches result from another

problem such as infection, injury, bleeding, very high blood pressure, or a brain disorder.

### Headache and Migraine

Alternate name; see *Headache*.

### Headache Red Flags

Critical clinical warning signs in a patient presenting with headache that indicate potentially life-threatening intracranial pathology (such as subarachnoid hemorrhage, meningitis, intracranial tumor, or temporal arteritis). Warning signs include sudden thunderclap onset, fever with meningismus, new focal neurological deficits, onset after age 50, papilledema, or postural worsening.

#### Alternate Names:

Headache Warning Signs; Red Flag Headaches; Danger Signs in Headache

### Headaches/Migraines

Alternate consumer label for headache disorders, especially *Migraine*; recurrent or new severe headaches require appropriate clinical evaluation.

### Healing

The body's process of repairing damaged tissue. It usually involves stopping bleeding, reducing inflammation, forming new tissue, and remodeling a scar; healing may be delayed by infection, poor blood flow, diabetes, or inadequate nutrition.

### Health Anxiety Disorder

Excessive worry about having or developing a serious illness, often despite reassurance or limited medical findings. It can cause repeated checking or health-care visits and is treated with supportive care and evidence-based psychological therapy.

### Health Behavior

An action or pattern of action that affects health, such as eating, physical activity, tobacco use, alcohol use, sexual practices, or taking medicines.

### Health Behavior Change

Modification of behaviors that affect health, such as

---

tobacco use, diet, physical activity, and adherence to treatment.

### **Health Behavior Motivation**

Factors that influence initiation and maintenance of health-related behaviors, including adherence, physical activity, and smoking cessation.

### **Health Care**

The prevention, diagnosis, treatment, and ongoing support provided to maintain or improve health, delivered by professionals, institutions, public-health systems, and informal caregivers.

### **Health Care Agents**

People legally chosen to make healthcare decisions for someone who cannot make or communicate them. Their authority and responsibilities depend on local law and the document that appoints them.

### **Health Care Facility**

A place where health services are delivered, such as a clinic, hospital, laboratory, nursing facility, or rehabilitation center; its services and level of care determine appropriate referrals.

### **Health Care Personnel**

Professionals and support workers involved in patient care, including clinicians, nurses, therapists, technicians, aides, and public-health staff.

### **Health Care Proxy**

A person designated to make healthcare decisions for someone who cannot make or communicate those decisions. The appointment may be recorded in an advance directive or other legal document.

### **Health Checkup**

A preventive visit that reviews symptoms, medical history, medicines, lifestyle, vaccinations, examination findings, and age- or risk-appropriate screening. The content should be individualized rather than based on unnecessary tests.

### **Health Disparities**

Differences in health outcomes or access to care among groups, often associated with social, economic, environmental, geographic, or structural factors.

### **Health Disparity**

A difference in health outcomes or access to health-care between population groups. It may be linked to income, housing, education, discrimination, environment, language, disability, geography, or unequal treatment.

### **Health Education**

The process of giving people understandable information and skills to prevent illness, recognize symptoms, use treatment safely, and make informed health decisions.

### **Health Inequity**

A systematic, avoidable, and unjust difference in health or access to health resources between social groups. It differs from health variation that is not preventable or not linked to unfair social conditions.

### **Health Information Technology**

The use of computers and digital systems to collect, store, exchange, protect, and use health information. Examples include electronic records, telemedicine, patient portals, decision-support tools, and remote monitoring; safe systems protect privacy and support accurate care.

### **Health Literacy**

The ability to find, understand, judge, and use health information and services to make appropriate decisions. It includes communicating with clinicians, following instructions, reading medicine labels, and recognizing when to seek help; clear communication by healthcare staff also improves health literacy.

### **Health Maintenance Organization**

A health-insurance arrangement that usually uses a network of clinicians and requires a primary-care clinician to coordinate specialist care. Coverage rules, referrals, costs, and services vary by plan.

---

## Health Promotion

Actions that help people improve or protect their health and prevent illness. They may include vaccination, healthy eating, physical activity, sleep, screening, safer environments, social connection, and avoiding tobacco. Advice should reflect a person's age, abilities, culture, risks, and preferences.

## Health Risk

The likelihood that a behavior, exposure, condition, or other factor will cause harm to health, considered together with the severity and timing of possible harm.

## Health Risks Of Alcohol Use

Alcohol use can increase the risk of liver disease, several cancers, high blood pressure, injuries, dependence, and problems affecting mental health and relationships.

## Health Surveillance

The ongoing collection and analysis of health data to identify patterns, monitor disease, and guide public-health action.

## Health Survey

A structured set of questions used to collect information about health, symptoms, behaviors, healthcare use, or population needs.

## Health-Related QoL

A person's physical, emotional, social, and daily functioning as affected by health and illness. It reflects the patient's experience and is assessed separately from laboratory results or survival.

## Healthcare Associated Infection

An infection acquired while receiving healthcare or as a result of a medical procedure. It may occur in hospitals, clinics, nursing facilities, or at home and can involve a wound, urinary tract, lungs, bloodstream, or other site.

## Healthcare Provider

A healthcare professional or organization that pro-

vides medical care, such as a physician, nurse, therapist, or clinic.

## Healthcare Proxy

A person designated to make healthcare decisions for someone who cannot make or communicate those decisions. The appointment may be recorded in an advance directive or other legal document.

## Healthy Aging

Developing and maintaining the physical and mental abilities that support well-being and independence in later life. It is influenced by health conditions, activity, nutrition, relationships, environment, and access to care.

## Hearing

The ability to detect and understand sound. Sound vibrations are converted into nerve signals in the inner ear and interpreted by the brain. Hearing loss may result from the outer or middle ear, inner ear, hearing nerve, or brain pathways.

## Hearing Aid

A small electronic device worn in or behind the ear that makes sounds louder or clearer for someone with hearing loss. It contains a microphone, processor, and speaker and is fitted and adjusted according to hearing-test results.

## Hearing Aids And Devices

Electronic devices worn in or behind the ear that amplify sound for individuals with hearing loss. Modern hearing aids include digital signal processing, directional microphones, and connectivity to smartphones and other audio sources.

## Hearing Disorder

A condition that reduces or distorts hearing through disease of the outer or middle ear, cochlea, auditory nerve, or central pathways; audiometry and examination help distinguish conductive from sensorineural loss.

---

## Hearing Impairment

Partial or complete reduction in the ability to detect or understand sound. It may be conductive, sensorineural, or mixed and can result from congenital disorders, infection, noise, ototoxic injury, trauma, ageing, or disease of the ear or auditory pathways.

## Hearing Loss

Reduced ability to hear. Conductive loss results from a problem in the outer or middle ear; sensorineural loss results from damage to the inner ear or hearing nerve; mixed loss has both types. Causes include aging, noise, infection, injury, and some medicines.

## Hearing Loss And Music

Difficulty hearing or distinguishing musical sounds because of reduced hearing, damage to inner-ear cells, or problems processing sound. Music may seem quieter, distorted, or lacking in certain pitches; hearing protection and individualized treatment may help.

## Hearing Loss: Sudden Deafness

A rapid sensorineural hearing loss, usually developing over hours to a few days, that may affect one or both ears. It is an otologic emergency; urgent hearing testing and evaluation are needed because early treatment may improve recovery.

## Hearing Screening

A quick test used to identify possible hearing loss before symptoms are obvious. Newborns are commonly screened before leaving hospital; an abnormal result requires repeat testing and a full hearing assessment rather than confirming permanent loss by itself.

## Hearing Test

An audiologic assessment that measures hearing thresholds and speech perception across frequencies to identify conductive, sensorineural, or mixed hearing loss.

## Heart

A four-chambered muscular pump (two atria, two

ventricles) located in the mediastinum, responsible for circulating blood throughout the body.

## Heart And Vascular Services

Heart and vascular services include prevention, diagnosis, and treatment of diseases affecting the heart and blood vessels.

## Heart Aneurysm

A localized bulging or outpouching of a weakened heart wall, often after myocardial infarction or from cardiomyopathy, that can impair contraction or form clots and arrhythmias.

## Heart Arrhythmia

An abnormal heart rate or rhythm caused by altered electrical activity in the heart. At rest, the adult heart usually beats regularly at about 60–100 times per minute; an arrhythmia may make it beat too fast, too slowly, or irregularly.

### Alternate Names:

Cardiac Dysrhythmias; Irregular Heart Rhythm

## Heart Attack

A heart attack (myocardial infarction) occurs when blood flow to part of the heart muscle is suddenly reduced or blocked, usually because a blood clot forms in a heart artery. The reduced oxygen supply can injure or kill heart muscle. Symptoms may include chest discomfort, shortness of breath, sweating, or nausea, but they vary and may be mild or absent. A suspected heart attack is a medical emergency.

## Heart Attack First Aid

Heart attack first aid means calling emergency services immediately, helping the person rest, and following dispatcher instructions; aspirin may be appropriate for some adults who can safely take it.

## Heart Attack Prevention

Heart-attack prevention targets blood pressure, atherogenic cholesterol, diabetes, tobacco exposure, diet, physical activity, weight, sleep, and adherence to prescribed medicines. Risk-based preventive treatment is more effective than relying on one supplement or symptom.

---

## Heart Block

A delay or interruption in the electrical signal traveling from the heart's upper chambers to its lower chambers. It may cause a slow or irregular heart-beat, dizziness, fainting, or no symptoms.

## Heart Bypass Surgery

Coronary artery bypass grafting creates new routes around blocked coronary arteries using graft vessels to improve myocardial blood flow and relieve ischemic symptoms or reduce selected risks.

## Heart Catheterization

A procedure in which a catheter is passed through a blood vessel into the heart to measure pressures, assess coronary arteries, inject contrast, obtain samples, or perform treatment.

## Heart Chambers

The four spaces inside the heart: the right and left atria, which receive blood, and the right and left ventricles, which pump blood out.

## Heart Contraction

The squeezing of heart muscle that pushes blood from the lower chambers to the lungs and the rest of the body. Electrical signals coordinate the heart-beat, and abnormal contraction can reduce blood flow or cause rhythm problems.

## Heart CT Scan

Computed tomography of the heart and nearby structures, sometimes with ECG synchronization or contrast, used to evaluate coronary arteries, cardiac anatomy, calcium, or selected masses.

## Heart Defect

An abnormality of the heart or major blood vessels. It may be present from birth or develop later and can affect blood flow, heart rhythm, valve function, or pumping ability.

## Heart Disease

A broad term for conditions affecting the heart or blood vessels, including coronary artery disease,

heart failure, abnormal rhythms, valve disease, and congenital defects. Symptoms and treatment depend on the part affected and may include chest pain, breathlessness, palpitations, or swelling.

## Heart Disease And Depression

Depression can occur with heart disease and may worsen quality of life, treatment adherence, and recovery; both conditions should be assessed and treated.

## Heart Disease And Intimacy

Heart disease or its treatment may affect sexual activity, desire, and confidence; individualized medical advice can help address symptoms and safety.

## Heart Disease And Women

Heart disease in women may involve different risk factors, symptoms, and timing than in men; chest pressure, shortness of breath, nausea, fatigue, or jaw or back pain can be important symptoms.

## Heart Failure

Heart failure is a condition in which the heart cannot pump enough blood to meet the body's needs or cannot fill with blood properly. It does not mean that the heart has stopped. The heart may be too weak to pump effectively or too stiff to fill normally. Symptoms may include shortness of breath, tiredness, reduced ability to exercise, and swelling from fluid buildup.

## Heart Failure Due To Valvular Heart Disease

Heart failure caused by narrowing or leakage of one or more heart valves. The valve problem can strain the heart or limit blood flow, causing breathlessness, tiredness, reduced ability to exercise, or swelling. Assessment identifies the affected valve and how it affects heart function.

## Heart Failure In Children

A condition in which a child's heart cannot pump or fill well enough to meet the body's needs. It may cause fast breathing, sweating or tiring during feeds, poor growth, swelling, or reduced activity.



---

## Heart Failure Stages

A framework describing progression from risk without heart disease, to heart changes without symptoms, to heart failure with symptoms, and finally advanced disease with persistent symptoms. Stages help guide prevention and treatment and are different from scales that describe how limited a person feels during activity.

## Heart Failure With Mildly Reduced Ejection Fraction

Heart failure in which the left ventricle pumps out a somewhat smaller-than-usual proportion of its blood, commonly about 41–49 percent. Symptoms may include breathlessness, tiredness, and fluid-related swelling.

## Heart Failure With Preserved Ejection Fraction

Heart failure in which the left ventricle still squeezes out a usual proportion of its blood but is too stiff to fill normally. Fluid then backs up, causing breathlessness, tiredness, and swelling.

## Heart Failure With Reduced Ejection Fraction

Heart failure in which the left ventricle cannot squeeze strongly enough to supply the body. It may cause breathlessness, tiredness, and swelling and can result from blocked arteries, heart-muscle disease, high blood pressure, or valve disease.

## Heart Injury

Damage to heart muscle or surrounding cardiac structures caused by trauma, ischemia, inflammation, toxins, or procedures; symptoms and tests depend on the injured tissue and severity.

## Heart MRI

Magnetic resonance imaging of the heart that evaluates cardiac structure, function, tissue characteristics, blood flow, and selected congenital, ischemic, inflammatory, or infiltrative disorders.

## Heart Murmur

An extra or unusual sound heard when blood flows

through the heart or nearby blood vessels. Some murmurs are harmless, while others result from a valve problem, heart defect, or altered blood flow.

## Heart Murmur Entry

Alternate name; see *Heart Murmur*.

## Heart Murmurs

Heart murmurs are extra or unusual sounds caused by turbulent blood flow through the heart or nearby vessels; some are harmless and others reflect heart disease.

## Heart Muscle

The myocardium, the specialized cardiac muscle that contracts rhythmically to pump blood through the heart and circulation.

## Heart Neoplasm

An abnormal growth in the heart or its surrounding tissues. It may be benign or cancerous and can interfere with blood flow, heart rhythm, or pumping; evaluation uses imaging and sometimes biopsy.

## Heart Pacemaker

A heart pacemaker is a device that sends electrical impulses to maintain an appropriate heartbeat when the heart's natural rhythm is too slow or unreliable.

## Heart Palpitations

Alternate name; see *Palpitation*.

## Heart PET Scan

Positron-emission tomography of the heart using a radiotracer to assess myocardial perfusion, metabolism, viability, inflammation, or selected cardiac masses.

## Heart Rate

The number of heartbeats per minute. Resting rates vary with age, fitness, activity, illness, medicines, and emotion; a persistently very fast, slow, or irregular rate may need assessment.

---

## Heart Rate Variability

The physiological variation and fluctuation in consecutive RR intervals between heartbeats over time, reflecting the autonomic balance and dynamic regulatory modulation between sympathetic and parasympathetic (vagal) nervous system inputs on the sinoatrial node.

**Alternate Names:**

HRV

## Heart Rhythm

The rate and pattern of cardiac electrical activation and contraction, normally initiated by the sinoatrial node; abnormal rhythms include bradycardia, tachycardia, atrial fibrillation, and ventricular arrhythmias.

## Heart Rhythm Disorders

Arrhythmias, in which the heart beats too fast, too slowly, or irregularly because of abnormal electrical impulse formation or conduction.

**Alternate Names:**

Cardiac Arrhythmias; Dysrhythmias

## Heart Septum

The muscular or membranous wall separating the right and left sides of the heart, including the atrial and ventricular septa; defects can permit abnormal blood flow between chambers.

## Heart Sounds

The sounds made as the heart valves open and close during each heartbeat. The usual two sounds are called “lub-dub”; extra or unusual sounds may suggest a valve problem, heart failure, or another condition.

## Heart Surgery

Surgical procedures performed on the heart or great vessels to treat structural or functional cardiac conditions. Common operations include coronary artery bypass grafting (CABG), valve repair or replacement, and repair of congenital heart defects.

## Heart Tamponade

A life-threatening condition in which fluid or blood builds up around the heart and prevents it from filling normally. It can cause breathlessness, faintness, low blood pressure, and shock and requires emergency treatment.

## Heart Transplant

Replacement of a failing heart with a donor heart, followed by lifelong immunosuppression and surveillance for rejection, infection, coronary allograft disease, kidney injury, and medication complications.

## Heart Transplantation

Surgical replacement of a failing heart with a donor heart, usually for selected patients with advanced heart failure who meet specialist criteria.

## Heart Valve

A flap-like structure that directs one-way blood flow through the heart. The four valves are the tricuspid, pulmonary, mitral, and aortic valves.

## Heart Valve Disease

Damage or abnormality of one or more of the heart’s valves. A valve may become narrowed and restrict blood flow (stenosis), or may not close fully and allow blood to leak backward (regurgitation). Causes include age-related changes, a defect present at birth, rheumatic disease, and infection.

## Heart Valve Disorder

Alternate term; see *Heart valve disease*.

## Heart Valve Prosthesis

An artificial valve used to replace or repair a diseased heart valve. Mechanical valves are durable but usually require long-term blood-thinning medicine. Tissue valves may avoid lifelong blood thinners but can wear out and sometimes need another procedure.

## Heart Valve Surgery

Repair or replacement of a diseased cardiac valve to correct severe narrowing or leakage and improve

---

blood flow; choice of repair, mechanical valve, or tissue valve depends on anatomy and patient factors.

**Alternate Names:**

Heart Valve Repair Or Replacement

**Heart Ventricle**

One of the two lower heart chambers that pumps blood to the lungs or systemic circulation; the right ventricle pumps to the pulmonary artery and the left to the aorta.

**Heart Ventricles**

The two lower chambers of the heart. The right ventricle pumps blood to the lungs, and the left ventricle pumps blood to the body.

**Heart-Lung Machine**

A device that temporarily takes over the pumping and gas-exchange functions of the heart and lungs during selected cardiac operations.

**Alternate Names:**

Cardiopulmonary Bypass Pump; Extracorporeal Circuit

**Heart-Smart Substitutions**

Easy food changes that may support heart health, such as choosing unsaturated oils instead of butter, whole grains instead of refined grains, and less processed or salty food.

**Heartbeat**

One complete contraction and relaxation of the heart, which moves blood through the lungs and the rest of the body. The rate and rhythm can be assessed by feeling a pulse or recording the heart's electrical activity.

**Heartburn**

A burning feeling behind the breastbone caused by stomach contents flowing back into the esophagus. It may be accompanied by a sour taste or regurgitation and is not usually caused by the heart. Frequent symptoms may indicate reflux disease.

**Heartburn/GERD**

Alternate name; see *Gastroesophageal Reflux Disease*.

**Heartland Virus**

A tick-borne virus associated with fever, fatigue, headache, muscle aches, diarrhea, and low white-cell or platelet counts. It can resemble ehrlichiosis and may be severe, especially in older or medically vulnerable people.

**Heat**

Thermal energy that raises tissue or body temperature; excessive heat exposure can cause heat cramps, heat exhaustion, heatstroke, or thermal injury.

**Heat Cramps**

Painful, involuntary muscle spasms occurring during or after strenuous activity in heat, often related to fluid and electrolyte loss. Resting in a cool place and replacing fluids and electrolytes may help. Confusion, fainting, or ongoing symptoms may signal heat illness requiring urgent medical attention.

**Heat Emergencies**

Heat emergencies range from heat cramps and exhaustion to heat stroke, in which dangerous core-temperature elevation causes central nervous system dysfunction and organ injury.

**Heat Exhaustion**

A heat-related illness caused by losing too much water and salt through sweating. It may cause heavy sweating, weakness, dizziness, headache, nausea, or fainting. Confusion, seizures, or very hot skin suggest heat stroke and require emergency care.

**Heat Intolerance**

An abnormal inability to tolerate warm environments or increased body heat, often causing excessive sweating, flushing, fatigue, or dizziness. Common causes include hyperthyroidism, menopause, fever, medications, autonomic disorders, and dehydration.

---

## Heat Pack

A device that applies warmth to the body surface to relieve some muscle pain or stiffness. Excessive heat or prolonged contact can cause burns.

## Heat Prostration

Older term; see *Heat Exhaustion*.

## Heat Rash

An itchy or prickly rash caused by blocked sweat ducts during hot, humid conditions. Small red bumps or clear vesicles often appear in skin folds or areas covered by clothing. Cooling, keeping the skin dry, and avoiding occlusive products usually help; infection or persistent symptoms warrants assessment.

## Heat Shock Protein

A stress-response protein that helps other proteins fold, avoid aggregation, and recover after heat, oxidative, infectious, or other cellular stress; some also influence immunity and cancer biology.

## Heat Stress

The strain placed on the body when it cannot adequately lose heat, often during hot or humid conditions, exertion, or inadequate hydration. It can progress from heat cramps or exhaustion to life-threatening heat stroke.

## Heat Stroke

A life-threatening heat-related illness with markedly elevated body temperature and altered mental status. It may cause confusion, seizures, loss of consciousness, and damage to multiple organs.

## Heavy Menstrual Bleeding

Menstrual bleeding that is unusually heavy, prolonged, or disruptive to daily life. Causes include fibroids, adenomyosis, polyps, changes in ovulation, bleeding disorders, medicines, or other medical conditions. Ongoing blood loss may cause iron-deficiency anemia; bleeding during pregnancy is not menstrual bleeding and needs separate assessment.

**Alternate Names:**

Menorrhagia; Hypermenorrhea

## Heavy Metal Poisoning

Harm from excessive exposure to toxic metals or metalloids such as lead, mercury, arsenic, cadmium, or thallium. Effects vary by substance and dose and may include abdominal pain, anemia, nerve injury, kidney damage, or confusion. Management starts with removing the exposure and may include supportive care or specialist-directed chelation therapy.

## Heberden Node

A firm bony bump at the finger joint nearest the fingertip, usually caused by osteoarthritis. It may be painful or tender while developing and can cause stiffness or reduced finger movement.

## Heberden Nodes

Firm bony enlargements at the finger joints nearest the fingertips, commonly caused by osteoarthritis. They may be painful or tender while forming, then remain as permanent bumps that limit movement in some people.

**Alternate Names:**

Heberden's Arthritic Nodes; DIP Joint Osteophytes

## Heberden's Node

A hard, bony enlargement of the distal interphalangeal joint caused by osteoarthritis, usually affecting the fingers and sometimes producing pain, stiffness, or reduced dexterity.

## Heberden's Nodes

Firm bony enlargements of the distal interphalangeal finger joints, usually caused by osteoarthritis.

## Hecht's Pneumonia

A rare, severe form of measles-related pneumonia, usually affecting people with weakened immunity. It can cause rapidly worsening breathing problems and requires urgent hospital treatment.

## Hedgehog Proteins

Signaling proteins that help guide the growth and or-

---

ganization of tissues before birth and during repair. Abnormal activity in this pathway can contribute to some birth differences and cancers.

## Heel

The heel is the back and underside of the foot, formed mainly by the calcaneus and a cushioning fat pad.

## Heel Pain

Pain beneath or around the heel, commonly caused by plantar fasciitis, Achilles tendinopathy, bursitis, stress fracture, nerve compression, or inflammatory arthritis.

### Alternate Names:

Plantar Fasciitis Discomfort; Calcaneal Spur Pain

## Heel Spur

A bony growth arising from the heel bone, often associated with plantar-fascia strain. The spur itself may cause no symptoms; heel pain is more often related to surrounding soft-tissue inflammation.

## Heel-to-Shin Test

A neurological test of lower limb coordination in which the patient places the heel of one foot on the opposite knee and slides it smoothly down the crest of the tibia to the ankle. Inability to maintain contact or an irregular, jerky trajectory indicates cerebellar ataxia or sensory ataxia.

## Heerfordt Syndrome

A specific clinical presentation of acute sarcoidosis characterized by the classic tetrad of anterior uveitis, bilateral parotitis (parotid gland enlargement), low-grade pyrexia, and unilateral or bilateral lower motor neuron facial nerve (CN VII) palsy.

### Alternate Names:

Uveoparotid Fever; Heerfordt-Waldenström Syndrome

## Hegar's Dilators

Hegar's dilators are graduated metal instruments used to gently widen the cervix or other body openings during selected procedures.

## Hegar's Sign

Softening of the lower part of the uterus, once used as an early sign of pregnancy. It is not specific and is not sufficient to confirm pregnancy without testing.

## Height

The vertical distance from the base of a person to the top of the head, measured for growth assessment, nutritional evaluation, drug dosing, and calculation of body-mass index.

## Heimlich Maneuver

A common name for abdominal thrusts used to help a responsive person with severe choking. First-aid steps differ for adults, children, infants, and pregnant people; emergency services should be called for serious airway obstruction.

## Heimlich Maneuver On Self

A self-administered abdominal-thrust technique sometimes used for severe choking when a person cannot breathe, speak, or cough effectively; emergency services should be activated immediately.

## Heimlich Manoeuvre

An emergency procedure for relieving upper airway obstruction caused by a foreign body (choking). The maneuver is performed on a conscious, choking adult who cannot speak, cough effectively, or breathe.

## Heine-Medin Disease

An older name for polio, a viral infection that can inflame the spinal cord and brain. It may cause fever and weakness and sometimes permanent paralysis; vaccination prevents most cases.

## Heinz Bodies

Clumps of damaged hemoglobin inside red blood cells, usually formed after oxidative injury. They may occur in glucose-6-phosphate dehydrogenase deficiency or after exposure to certain medicines, chemicals, or infections.

---

## Hejama

An Arabic term for cupping, a traditional practice in which suction is applied to the skin; wet cupping also involves superficial skin incisions and carries bleeding and infection risks.

## Helcococcus

A genus of gram-positive cocci that can rarely cause opportunistic skin, wound, bone, or bloodstream infections.

## Helcococcus Kunzii

A Gram-positive round bacterium usually found on skin that rarely causes opportunistic infection. Reported infections may involve wounds, blood, or implanted devices in people with serious illness or weakened defenses.

## Helcococcus Pyogenes

A rarely encountered Gram-positive round bacterium that may cause opportunistic human infection. It has been found in wounds or blood, particularly in people with serious illness, chronic wounds, or weakened immunity.

## Helicase

An enzyme that uses cellular energy to separate the paired strands of DNA or RNA. This allows genetic material to be copied, repaired, or read.

## Helicobacter

A genus of spiral-shaped bacteria that colonize the stomach or intestine; some species cause ulcers, gastritis, or other disease.

## Helicobacter Cinaedi

A *Helicobacter* species that can cause bloodstream, skin, bone, joint, or other invasive infections, especially in people with weakened immunity.

## Helicobacter Fennelliae

An intestinal *Helicobacter* species that rarely causes bloodstream or other opportunistic infection. Disease is more likely in people with weakened defenses, and diagnosis requires specialized laboratory identi-

fication.

## Helicobacter Pylori

A gram-negative, spiral-shaped bacterium adapted to colonize the stomach. It can cause chronic gastritis, peptic ulcer disease, and increase the risk of gastric cancer and gastric MALT lymphoma.

## Helicobacter Pylori Eradication

Pharmacological combination therapy designed to eliminate gastric *H. pylori* infection, traditionally utilizing triple therapy (proton pump inhibitor, clarithromycin, amoxicillin) or bismuth quadruple therapy (PPI, bismuth subsalicylate, metronidazole, tetracycline) to heal peptic ulcers and reduce gastric cancer risk.

## Helicobacter Pylori Infection

Infection of the stomach by *H. pylori*, a spiral-shaped bacterium associated with chronic gastritis, peptic ulcer disease, gastric adenocarcinoma, and gastric MALT lymphoma. It is diagnosed with noninvasive or endoscopic testing and treated with combination eradication therapy.

## Helicobacter Pylori-Associated Active Gastritis Pathology

Chronic active gastritis with lymphoplasmacytic inflammation, neutrophils in pits or surface epithelium, lymphoid aggregates, and curved organisms along the surface mucus or foveolar epithelium.

## Helicobacteraceae

A family of spiral-shaped bacteria that includes *Helicobacter* and related genera, many of which inhabit the digestive tract.

## Helicotrema

The narrow anatomical communication aperture located at the apex of the cochlea where the scala vestibuli meets and continues into the scala tympani, allowing pressure equilibration of perilymph between the two compartments.



---

## Heliotherapy

Treatment using controlled exposure to sunlight or artificial ultraviolet light. It has been used for selected skin disorders, but excessive exposure can cause burns, premature aging, eye injury, and skin cancer.

## Heliotrope Rash

A purple or lilac rash around the eyelids, often with swelling. It is a characteristic sign of dermatomyositis, but other findings and tests are needed for diagnosis.

### Alternate Names:

Heliotrope Erythema; Dermatomyositis Eyelid Sign

## Helium

A chemically inert gas used in respiratory mixtures, cryogenics, imaging, and laboratory applications; inhalation can displace oxygen and cause hypoxia despite its lack of direct toxicity.

## Helix

The curved outer rim of the external ear. Its shape is formed by cartilage covered with skin; swelling, tenderness, or a skin change on the helix may follow injury, infection, sun damage, or other skin disease.

## Heller Myotomy

A surgical procedure that cuts the muscles of the lower esophageal sphincter to relieve obstruction in achalasia, usually performed laparoscopically and often combined with an antireflux procedure.

## Heller's Operation

A surgical procedure that cuts the lower esophageal muscle to relieve achalasia, usually with an antireflux procedure.

## HELLP syndrome

A pregnancy-associated syndrome named for hemolysis, elevated liver enzymes, and low platelet count. It is a form of microangiopathic disease that may occur with preeclampsia and can involve upper-abdominal pain, nausea, headache, or high blood pressure.

## Helminth

A parasitic worm, including roundworms, tapeworms, and flukes, that may live in or on a host.

## Helminthiasis

Infection or infestation caused by parasitic worms, including roundworms, tapeworms, and flukes. Symptoms vary by species and may include abdominal problems, anemia, cough, rash, malnutrition, or organ damage.

### Alternate Names:

Worm Infestation; Helminthic Parasitosis

## Helminths

Parasitic worms that cause disease in humans. The major groups are nematodes (roundworms), cestodes (tapeworms), and trematodes (flukes). Helminths have complex life cycles often involving intermediate hosts. Infection occurs through ingestion of eggs or larvae, penetration of skin, or insect vectors.

## Help Prevent Hospital Errors

Actions that reduce hospital errors include confirming identity, checking medicines, communicating clearly, asking questions, and reporting changes or concerns promptly.

## Helper T Cell

A type of white blood cell that coordinates immune responses by sending chemical signals that activate and guide other immune cells, including antibody-producing cells and cells that destroy infected cells.

## Helper T Cells

A type of white blood cell that helps organize the immune response. These cells release chemical messages that activate and guide other immune cells, including cells that make antibodies or destroy infected cells.

### Alternate Names:

CD4+ T Lymphocytes; T Helper Cells

## Helper Virus

A virus that supplies functions needed for another

---

virus or defective viral genome to replicate.

### **HELPERR Mnemonic**

A memory aid for managing shoulder dystocia during birth: call for help, consider an episiotomy, flex the birthing person's legs, apply suprapubic pressure, perform internal maneuvers, remove the baby's posterior arm, and roll to an all-fours position. It requires trained obstetric staff.

### **Hemagglutination Test**

A test in which antibodies or other agents cause red blood cells to clump; the reaction can help detect antibodies, antigens, or selected infectious agents.

### **Hemagglutinin**

A surface glycoprotein, especially on influenza viruses, that binds sialic-acid receptors and helps the virus attach to and enter host cells; it is a major target of protective antibodies.

### **Hemangioblastoma**

A usually slow-growing tumor made of blood-vessel-forming cells, most often found in the cerebellum or spinal cord. It may cause headache, balance problems, or weakness and can be associated with von Hippel–Lindau disease.

### **Hemangioendothelioma**

A rare tumor arising from cells that line blood vessels. Its behavior ranges from slow-growing to cancerous, depending on the subtype and location, so diagnosis requires tissue examination and specialist assessment.

### **Hemangioma**

A usually noncancerous growth made of blood vessels. It may be present at birth or appear during infancy, and can be a small skin mark or a deeper mass. Some become smaller over time; others can cause bleeding, ulceration, or pressure on nearby structures.

#### **Alternate Names:**

Vascular Hemangioma; Benign Vascular Tumor

### **Hemangiopericytoma**

A rare tumor formerly defined by branching “staghorn” vessels and perivascular cells; many lesions classified this way are now recognized as solitary fibrous tumors.

### **Hemangiosarcoma**

A malignant tumor of vascular endothelial cells, an aggressive sarcoma that may arise in the skin, liver, or bone.

### **Hemarthrosis**

Bleeding into a joint, most often the knee, causing rapid swelling, warmth, pain, and limited movement. Injury and bleeding disorders such as hemophilia are common causes; repeated episodes can damage cartilage and lead to long-term joint disease.

### **Hematemesis**

Vomiting blood or material that looks like coffee grounds, usually from bleeding in the esophagus, stomach, or first part of the small intestine. It can be life-threatening.

### **Hematidrosis**

A very rare condition in which blood-tinged fluid appears through apparently intact skin or other surfaces, sometimes during extreme stress. The diagnosis requires careful examination to exclude injury, skin disease, bleeding disorders, or contamination.

### **Hematocele**

A collection of blood around a testicle, usually after scrotal injury or surgery and sometimes from a tumor. It can cause a heavy, firm, painful swelling; ultrasound is needed to check the testicle and blood flow, and large collections may need surgery.

#### **Alternate Names:**

Scrotal Hematocele; Tunica Vaginalis Blood Collection

### **Hematochezia**

Passing bright-red or maroon blood from the rectum. It often comes from the large intestine, rectum, or anus, but heavy bleeding higher in the digestive

---

tract can look the same.

### **Hematocolpos**

Accumulation of menstrual blood in the vagina, usually because an imperforate hymen or another obstructing anomaly prevents outflow; it may cause cyclic pelvic pain and a bulging vaginal membrane.

### **Hematocrit**

The proportion of blood volume occupied by packed red blood cells, expressed as a percentage. One of the most commonly ordered laboratory tests in clinical medicine.

### **Hematogenous**

Spread through the bloodstream, as when infection, tumor cells, or embolic material travels from one body site to another.

### **Hematogenous Osteomyelitis**

An invasive bone infection arising from bacteremic seeding of the vascular marrow spaces, classically affecting the metaphyses of long bones in growing children (due to slow sinusoidal capillary blood flow) and the vertebral bodies (spondylodiscitis) in older adults, most frequently caused by *Staphylococcus aureus*.

### **Hematologic Agents**

Medicines or biologic therapies that act on blood cells, coagulation, marrow, or immune pathways, including agents used for anemia, thrombosis, leukemia, and lymphoma.

### **Hematologic Changes In Pregnancy**

Pregnancy normally increases the fluid part of blood more than the red-cell mass, so blood counts may show mild dilutional anemia. White-cell counts also commonly rise, mainly because of more neutrophils. Results must be interpreted using pregnancy-specific ranges.

### **Hematologic Malignancy**

Alternate name; see *Blood Cancer*.

### **Hematologic Response**

A measurable improvement or change in blood counts or marrow findings after treatment, such as recovery of neutrophils, platelets, or hemoglobin or reduction of malignant cells.

### **Hematological Disease**

A disorder of blood cells, bone marrow, lymphatic tissue, or coagulation, including anemia, leukemia, lymphoma, bleeding disorders, and clotting disorders.

### **Hematologist**

A physician who diagnoses and treats disorders of blood, bone marrow, lymph nodes, clotting, and hematologic cancers.

### **Hematology**

The branch of medicine concerned with the study, diagnosis, treatment, and prevention of disorders of the blood and blood-forming organs (bone marrow, lymph nodes, spleen).

### **Hematology-Oncology**

The medical specialty concerned with blood disorders and cancers, including anemia, leukemia, lymphoma, myeloma, and clotting disorders.

### **Hematoma**

A collection of blood outside a blood vessel, usually after injury, surgery, or bleeding caused by illness or medicines. It may cause swelling and pain and can press on nearby nerves or organs.

### **Hematometra**

Blood trapped inside the uterus because the normal outlet is blocked or too narrow. It can cause absent periods, monthly pelvic pain, and an enlarged tender uterus; the blockage may be present from birth or develop after scarring, surgery, infection, or radiation.

### **Hematometrocolpos**

A buildup of menstrual blood in the uterus and vagina because the normal outlet is blocked, often

---

by tissue present from birth. It may cause absent periods, monthly pelvic pain, and a lower-abdominal mass.

### **Hematomyelia**

Bleeding inside the spinal cord, usually after a serious injury or from an abnormal blood vessel or bleeding disorder. It may cause sudden neck or back pain, weakness, numbness, or loss of bladder or bowel control.

### **Hematophobia**

An excessive or persistent fear of blood, blood-related sights, or medical procedures involving blood that causes distress or avoidance; some people have a vasovagal fainting response.

### **Hematopoiesis**

The process of making and maturing blood cells, mainly in the bone marrow. It produces red blood cells for oxygen transport, white blood cells for immune defense, and platelets for stopping bleeding.

### **Hematopoietic Growth Factor**

A cytokine or medicine that stimulates production, maturation, or survival of blood-forming cells in bone marrow, such as erythropoietin or granulocyte colony-stimulating factor.

### **Hematopoietic Neoplasm**

A cancer arising from blood-forming cells in the bone marrow or immune tissues. Leukemias, lymphomas, and plasma-cell cancers are examples; symptoms and treatment depend on the exact cell type and extent.

### **Hematopoietic Stem Cell Transplantation**

Infusion of blood-forming stem cells to replace damaged or diseased bone marrow. The cells may come from the patient or a donor. Treatment can cause serious infection, bleeding, infertility, or an immune reaction against donor and recipient tissues.

### **Hematopoietic System**

The bone marrow and related organs and cells that

produce, mature, circulate, and remove blood cells, including red cells, white cells, and platelets.

### **Hematoporphyrin Derivative**

A porphyrin-based photosensitizer that accumulates in some tumors and can be activated by light in photodynamic treatment; it may also cause prolonged photosensitivity.

### **Hematosalpinx**

Accumulation of blood within a fallopian tube, often caused by ectopic pregnancy, endometriosis, pelvic surgery, or obstruction, and potentially producing pelvic pain or infertility.

### **Hematosi**

The exchange of gases in the lungs: oxygen moves from inhaled air into the blood, while carbon dioxide moves from the blood into the air to be breathed out.

### **Hematospermia**

Blood in the semen, often temporary and associated with infection, inflammation, trauma, or a recent urologic procedure. Persistent or recurrent bleeding, pain, fever, urinary symptoms, or bleeding in an older adult requires medical evaluation.

### **Hematoxylin**

A dye used in histology, usually with eosin, to stain cell nuclei blue-purple and help show tissue structure.

### **Hematuria**

Blood in the urine. It may be visible as pink, red, or brown urine or found only under a microscope. Causes include infection, stones, medicines, injury, prostate disease, and kidney or urinary-tract disorders.

### **Heme**

Heme is an iron-containing porphyrin group that binds oxygen in haemoglobin and myoglobin and forms the functional centre of several enzymes; abnormal heme metabolism contributes to porphyrias

---

and jaundice.

### **Hemeralopia**

Hemeralopia is reduced vision in bright light, sometimes associated with retinal disorders or abnormal cone function.

### **Hemianesthesia**

The complete loss of tactile, thermal, or nociceptive somatic sensation affecting one vertical half of the body, classically caused by an ischemic or hemorrhagic lesion disrupting the contralateral primary somatosensory cortex or the ventral posterolateral (VPL) nucleus of the thalamus.

### **Hemianopia**

Loss of vision in half of the area a person can see, affecting one or both eyes. It often results from a problem in the brain's visual pathways, such as a stroke or tumor; sudden onset can occur with a serious brain disorder.

### **Hemianopsia**

Loss of vision in half of the visual field of one or both eyes, usually from a lesion of the optic tract, radiations, or occipital cortex; sudden onset may indicate stroke.

### **Hemiballism**

A movement disorder causing sudden, large-amplitude, flinging movements on one side of the body, usually from a lesion involving the subthalamic region.

### **Hemiballismus**

Hemiballismus is involuntary, rapid, flinging movement of one side of the body, usually caused by a lesion of the opposite subthalamic region.

### **Hemicolectomy**

Surgical removal of the right or left portion of the colon, sometimes with nearby lymph nodes and reconnection of the bowel.

### **Hemidesmosome**

A specialized junction that anchors basal epithelial cells to the underlying basement membrane by linking intracellular keratin filaments with extracellular matrix proteins.

### **Hemidiaphragm**

One of the two muscular halves of the diaphragm separating the chest from the abdomen and assisting breathing.

### **Hemifacial Atrophy**

Progressive wasting of skin, subcutaneous tissue, muscle, and sometimes bone on one side of the face, also called Parry–Romberg syndrome.

### **Hemifacial Microsomia**

A birth difference in which one side of the face is smaller or less developed than the other, often affecting the jaw, ear, cheek, and nearby soft tissues. Hearing, vision, and spinal differences may also occur.

### **Hemifacial Spasm**

Repeated involuntary twitching on one side of the face, usually beginning around the eye and sometimes spreading to the cheek or mouth. Irritation of the facial nerve is a common cause; persistent spasms need evaluation.

### **Hemihypertrophy**

Unequal overgrowth of one side or region of the body, producing asymmetry of limbs, trunk, or face; some childhood cases carry increased risk of embryonal tumors.

### **Hemilaryngectomy**

Hemilaryngectomy is surgery to remove part of the larynx, usually to treat a selected laryngeal cancer while preserving as much speech and swallowing function as possible.

### **Hemimandibulectomy**

Hemimandibulectomy is surgery to remove one side of the lower jaw, usually for a tumor or severe bone

---

disease, sometimes followed by reconstruction.

### **Hemimegalencephaly**

A rare developmental brain malformation in which one cerebral hemisphere is abnormally enlarged and dysplastic, often causing infantile seizures, developmental impairment, and contralateral neurologic signs.

### **Hemimelia**

A congenital limb-reduction defect in which part or all of a limb segment is absent, such as fibular or radial hemimelia.

### **Hemineglect**

A neurological attentional disorder classically resulting from an acute lesion of the non-dominant (usually right) posterior parietal cortex, in which the patient fails to orient toward, respond to, or perceive sensory stimuli presented in the hemispace contralateral to the brain lesion.

#### **Alternate Names:**

Spatial Neglect; Unilateral Neglect

### **Hemiparesis**

Weakness on one side of the body, affecting the face, arm, leg, or a combination. It is less severe than paralysis and may result from stroke, brain injury, a tumor, multiple sclerosis, or another nervous-system disorder.

### **Hemiparesis with Conjugate Gaze Deviation**

Weakness on the left side with both eyes looking to the right. This pattern can occur with a serious injury to the right side of the brain, including stroke, but examination and brain imaging are needed to identify the cause.

### **Hemipelvectomy**

Hemipelvectomy is major surgery that removes part or all of one side of the pelvis, with or without removal of the leg, usually for extensive cancer or severe trauma.

### **Hemiplegia**

Complete or near-complete paralysis of one side of the body, including the face, arm, or leg. It usually results from injury to movement pathways in the brain, commonly a stroke; sudden onset can occur with a serious brain disorder.

### **Hemispatial Neglect**

Hemispatial neglect is reduced awareness of one side of the body or space, usually after injury to the opposite cerebral hemisphere.

### **Hemispherectomy**

Hemispherectomy is surgery that removes or disconnects much of one cerebral hemisphere to treat severe, otherwise uncontrolled epilepsy in selected patients.

### **Hemisternebra**

A congenital or developmental partial segment of the sternum, caused by incomplete formation or fusion of sternal elements and sometimes associated with chest-wall anomalies.

### **Hemithorax**

One side of the thorax, including the lung, pleural space, ribs, and other structures on that side of the chest.

### **Hemivertebrae**

Congenital vertebrae formed from only part of the normal vertebral body, producing wedge-shaped bone and potentially causing congenital scoliosis or spinal-cord abnormalities.

### **Hemizygous**

Having only one copy of a gene or chromosome region rather than the usual pair, as occurs for many X-linked genes in people with one X chromosome.

### **Hemobilia**

Bleeding originating within the hepatobiliary tree that enters the gastrointestinal tract via the ampulla of Vater, classically presenting with the triad of biliary colic, upper gastrointestinal bleeding, and



---

obstructive jaundice (Sandblom triad), frequently following liver trauma, biopsy, or instrumentation.

### **Hemochromatosis**

A disorder in which too much iron builds up in the body and can damage the liver, pancreas, heart, joints, skin, and hormone-producing organs. The common HFE-related form is usually autosomal recessive; other inherited forms can have different patterns, and iron overload can also develop from another illness or repeated transfusions.

#### **Alternate Names:**

Hereditary Hemochromatosis

### **Hemochromatosis Cardiomyopathy**

Heart-muscle damage caused by excess iron buildup. The heart may become stiff or weak and cause breathlessness, swelling, abnormal rhythms, or heart failure; treatment aims to remove excess iron and support heart function.

### **Hemocyanin**

A copper-containing oxygen-transport protein in many mollusks and arthropods; unlike human hemoglobin, it circulates dissolved in their hemolymph.

### **Hemocytometer**

A ruled counting chamber used with a microscope to determine the concentration of cells or other particles in a measured volume of fluid.

### **Hemodialysis**

A treatment for kidney failure in which blood passes through a filter outside the body to remove waste and extra fluid before returning to the person. It usually requires repeated sessions and a blood-vessel access; low blood pressure, infection, clotting, and access problems can occur.

### **Hemodialysis Access Care**

Daily maintenance protocols and hygiene measures for an arteriovenous fistula, arteriovenous graft, or central venous catheter used for hemodialysis. Includes daily palpation for a vascular thrill, avoiding blood pressure cuffs or constrictive clothing on the

access arm, and promptly reporting signs of infection or thrombosis.

#### **Alternate Names:**

Vascular Access Care; Fistula and Graft Maintenance; Dialysis Access Management

### **Hemodialysis Access Procedures**

Procedures that create, maintain, repair, or revise vascular access for hemodialysis, including fistula or graft creation, catheter placement, angioplasty, thrombectomy, and treatment of stenosis or infection.

### **Hemodialysis Clearance**

The volume of plasma from which hemodialysis removes a solute per unit time; it depends on blood and dialysate flow, membrane properties, treatment time, and the substance's distribution.

### **Hemodialysis Solution**

The dialysate fluid circulated across a semipermeable membrane during hemodialysis to remove waste and help correct electrolyte, acid-base, and fluid abnormalities.

### **Hemodynamics**

The movement of blood through the heart and blood vessels, including pressure, flow, heart output, and resistance. Hemodynamic measurements help assess circulation and guide treatment in conditions such as shock, heart failure, and critical illness.

### **Hemofiltration**

A blood-cleaning treatment that removes extra water and dissolved waste through a filter. Replacement fluid is added to maintain the right blood volume; it is used in some forms of temporary or long-term kidney support.

### **Hemoglobin**

The iron-containing protein in red blood cells that carries oxygen from the lungs to tissues and helps carry carbon dioxide back to the lungs. Low or abnormal hemoglobin can cause anemia or other blood disorders.

---

## Hemoglobin A

The main type of hemoglobin in most adults. It is the oxygen-carrying protein in red blood cells and is made from two alpha and two beta protein parts.

## Hemoglobin A1c

A blood test that shows the average blood sugar level over the past 2 to 3 months. It is used to help diagnose and monitor diabetes and prediabetes, but it may be less reliable in some blood disorders, pregnancy, or conditions that change red blood cell life span.

### Alternate Names:

HbA1c; Glycated Hemoglobin; Glycosylated Hemoglobin

## Hemoglobin A1C Variability

Factors that can make hemoglobin A1c less reliable include altered red-cell survival, hemoglobin variants, anemia, pregnancy, kidney disease, transfusion, and some treatments. An alternative measure of glucose may be needed when A1c does not reflect recent glycemia.

## Hemoglobin C

A beta-globin variant caused by a hemoglobin gene mutation; homozygosity can cause mild hemolytic anemia and splenomegaly, while inheritance with hemoglobin S causes a clinically significant sickling disorder.

## Hemoglobin C Disease

An inherited hemoglobinopathy caused by a beta-globin variant that produces hemoglobin C. It may cause mild hemolytic anemia, jaundice, splenomegaly, or, when combined with hemoglobin S, clinically significant sickling disease.

## Hemoglobin Electrophoresis

A blood test that separates the different types of hemoglobin. It helps identify inherited conditions such as sickle-cell disease, sickle-cell trait, and thalassemia. Results are interpreted with age, pregnancy, blood counts, recent transfusion, and other tests.

## Hemoglobin H

An unusual form of hemoglobin made from four beta protein parts when the body makes too little alpha hemoglobin. It is unstable and can cause ongoing breakdown of red blood cells in alpha-thalassemia.

## Hemoglobin Increased

An increased hemoglobin concentration, which may reflect dehydration, chronic hypoxemia, erythrocytosis, or a myeloproliferative neoplasm and must be interpreted with hematocrit and clinical context.

## Hemoglobin M

A group of abnormal hemoglobins that stabilize iron in the ferric state and can cause congenital methemoglobinemia and cyanosis.

## Hemoglobin Normal Values

Reference ranges for blood hemoglobin concentration, which vary with age, sex, pregnancy, altitude, laboratory method, and other factors; results should be interpreted using the reporting laboratory's range.

## Hemoglobin S

An inherited form of hemoglobin that can make red blood cells become stiff and sickle-shaped when oxygen is low. It can cause anemia, painful blockages, and organ damage in sickle-cell disease.

## Hemoglobin SC Disease

An inherited form of sickle cell disease caused by having one hemoglobin S gene and one hemoglobin C gene. It can cause anemia, pain episodes, and organ complications.

### Alternate Names:

HbSC; Sickle-Hemoglobin C Disease

## Hemoglobin Trait

The inherited presence of a hemoglobin variant in a heterozygous state, usually causing no symptoms but potentially affecting screening results or reproductive risk.

---

## Hemoglobinopathies

A group of inherited disorders affecting the structure or production of hemoglobin. They include sickle-cell disorders and thalassemias and may cause anemia, hemolysis, abnormal oxygen transport, or blood-vessel complications.

### Hemoglobinopathies: Thalassemias

Thalassemias are inherited disorders of globin chain synthesis imbalance: alpha-thalassemia (HBA1/HBA2 deletions) and beta-thalassemia (HBB mutations).

### Hemoglobinopathy

An inherited disorder affecting the structure or production of hemoglobin. Examples include sickle-cell disease and thalassemia; effects range from no symptoms to anemia, painful blood-vessel blockages, and organ damage.

### Hemoglobinuria

Haemoglobinuria is free haemoglobin in urine, usually from intravascular haemolysis; urine may be dipstick-positive for blood with few or no red cells, and causes include transfusion reactions, malaria, and paroxysmal nocturnal haemoglobinuria.

### Hemoglobinuria Test

A laboratory test evaluating urine for hemoglobin or related findings, used to investigate intravascular hemolysis, hemoglobinuria, transfusion reactions, or other causes of dark urine.

### Hemolysin

A toxin or immune substance that damages the outer covering of red blood cells and makes them break open. Some bacteria produce hemolysins, and the resulting destruction of red cells is called hemolysis.

### Hemolysis

The early destruction of red blood cells, releasing hemoglobin into the blood. It can occur in the bloodstream or in the spleen and liver and may cause anemia, jaundice, dark urine, tiredness, or shortness of breath.

## Hemolytic

Relating to hemolysis, the premature destruction of red blood cells, which can release hemoglobin and produce anemia or jaundice.

### Hemolytic Anemia

Anemia caused by red blood cells being destroyed faster than the body can replace them. It may result from an inherited cell problem, an immune reaction, infection, medicine, toxin, or another illness and can cause jaundice and dark urine.

### Hemolytic Anemia From Toxins

Anemia caused by accelerated destruction of red blood cells after exposure to a chemical or toxin; evaluation identifies hemolysis and removes or treats the responsible exposure.

### Hemolytic Anemias: Autoimmune

Anemias in which the immune system mistakenly destroys red blood cells. Warm-antibody and cold-antibody forms differ in triggers and temperature sensitivity; symptoms may include tiredness, jaundice, dark urine, and breathlessness.

### Hemolytic Anemias: Hereditary

Inherited conditions in which red blood cells are destroyed too quickly. Hereditary spherocytosis is one example; it can cause anemia, jaundice, an enlarged spleen, and gallstones. Other inherited red-cell or enzyme problems can cause similar anemia.

### Hemolytic Crisis

A sudden increase in red-cell destruction causing worsening anemia, jaundice, dark urine, pain, or other manifestations of hemolysis.

### Hemolytic Disease Of The Newborn

Anemia and jaundice caused when antibodies from the pregnant person cross the placenta and destroy the fetus's or newborn's red blood cells. Severe disease can cause widespread fluid buildup or brain injury without prompt treatment.

---

## Hemolytic Index

A laboratory-quality indicator estimating hemolysis in a blood specimen; substantial in-vitro hemolysis can interfere with measurement of several analytes.

## Hemolytic Jaundice

Jaundice resulting from excessive hemolysis (breakdown of red blood cells) causing overproduction of unconjugated bilirubin that exceeds the liver's capacity for conjugation and excretion. See Jaundice for complete overview.

## Hemolytic Transfusion Reaction

An immune or nonimmune complication in which transfused red blood cells are destroyed, potentially causing fever, pain, hemoglobinuria, kidney injury, shock, or disseminated intravascular coagulation.

## Hemolytic Uremic Syndrome

A serious illness in which tiny blood clots damage small blood vessels, causing red blood cells to break apart, platelets to fall, and the kidneys to become injured. It often follows severe diarrheal infection but can have other causes.

## Hemolytic-Uremic Syndrome

A serious illness in which small blood-vessel clots destroy red blood cells, lower the platelet count, and injure the kidneys. It often follows severe diarrhea caused by certain bacteria, but inherited immune problems and other triggers can also cause it.

## Hemoperfusion

A treatment in which blood passes through a material such as charcoal or resin that traps selected poisons or medicines. It is used only for certain severe overdoses because effectiveness depends on the substance and timing.

## Hemopericardium

Accumulation of blood in the pericardial sac around the heart. If pressure rises and prevents the heart from filling, cardiac tamponade can occur.

## Hemoperitoneum

Blood collecting inside the abdominal cavity, usually from trauma, a ruptured blood vessel or organ, surgery, or a ruptured ectopic pregnancy. It can cause abdominal pain, distention, dizziness, or shock and may require emergency imaging and surgery or another procedure.

## Hemopexin

A plasma glycoprotein that binds free heme and transports it to the liver, helping limit oxidative injury during hemolysis.

## Hemophagocytic Lymphohistiocytosis

A rare, life-threatening illness in which the immune system remains excessively active and injures the body's organs. It may be inherited or triggered by infection, cancer, or immune disease. Persistent fever, enlarged spleen, low blood counts, liver problems, bleeding, or confusion may occur.

## Hemophagocytosis Of Sick Cells

A finding in which immune cells engulf blood cells in the bone marrow or other tissues during severe inflammation, infection, or sickle-cell complications. It can worsen anemia and low platelet counts and requires specialist evaluation.

## Hemophilia

A bleeding disorder, usually inherited, caused by too little or poorly working clotting factor VIII (hemophilia A) or factor IX (hemophilia B). It can cause prolonged bleeding and bleeding into joints or muscles. It affects people of all sexes; some women and girls who carry a hemophilia-related gene have low clotting-factor levels and bleeding symptoms. Rarely, hemophilia develops later in life when the immune system blocks a clotting factor. Treatment depends on the type and severity and may include factor replacement or other medicines that prevent or control bleeding.

### Alternate Names:

Haemophilia; Bleeder's Disease; The Royal Disease; Factor Deficiency Disorder; Classic Hemophilia

---

## Hemophilia A

An X-linked recessive deficiency of coagulation factor VIII, causing prolonged bleeding into joints, muscles, and soft tissues. Severity correlates with factor VIII activity levels.

## Hemophilia B

An X-linked recessive deficiency of coagulation factor IX (Christmas disease), causing prolonged bleeding similar to hemophilia A. Severity correlates with factor IX activity levels.

## Hemophilic Arthritis

Chronic joint damage caused by repeated bleeding into a joint, usually in people with hemophilia. Recurrent bleeds inflame and thicken the joint lining and can progressively damage cartilage and bone, causing pain, stiffness, and reduced movement.

### Alternate Names:

Hemophilic Arthropathy; Haemophilic Arthropathy; Haemophilic Arthritis

## Hemophilus

An older spelling or variant of *Haemophilus*, a group of small gram-negative bacteria that includes respiratory and genital pathogens.

## Hemophilus Ducreyi

A gram-negative bacterium that causes chancroid, a sexually transmitted infection with painful genital ulcers and tender lymph nodes.

## Hemophobia

An excessive or persistent fear of blood, also called hematophobia, that causes significant distress, avoidance, or sometimes a vasovagal faint.

## Hemopoietic Stem Cell

A blood-forming stem cell found mainly in the bone marrow. It can make copies of itself and develop into red blood cells, white blood cells, or platelets.

### Alternate Names:

Hematopoietic Stem Cell (HSC); Blood-Forming Cell

## Hemopoietic Tissue

Tissue that produces blood cells, chiefly bone marrow, containing stem cells and supporting cells that generate erythrocytes, leukocytes, and platelets.

## Hemoprotein

A protein containing a heme group with an iron atom, such as hemoglobin, myoglobin, cytochromes, and many oxidoreductive enzymes.

## Hemoptysis

Coughing up blood or blood-stained mucus from the airways or lungs. Causes include infection, irritated or widened airways, a blood clot, or cancer. Larger amounts may occur with breathlessness or chest pain.

## Hemorrhage

Abnormal escape of blood from a damaged blood vessel or, less commonly, from the heart. It may be external or internal, acute or chronic, and range from minor bleeding to life-threatening blood loss causing anemia, shock, or organ compression; the cause and site determine management.

## Hemorrhagic

Pertaining to or characterized by hemorrhage (extravasation of blood from vessels). Used in contexts such as hemorrhagic stroke, hemorrhagic shock, and hemorrhagic fever.

## Hemorrhagic Anemia

Anemia caused by losing blood faster than the body can replace it. Blood loss may be sudden, such as after an injury, or slow, such as from heavy menstrual or intestinal bleeding.

## Hemorrhagic Disorder

A condition that causes abnormal bleeding because of platelet number or function, clotting-factor deficiency, excessive fibrinolysis, or blood-vessel fragility.

## Hemorrhagic Shock

Circulatory failure caused by severe blood loss.

---

It may produce rapid pulse, low blood pressure, cool skin, reduced urine output, confusion, or collapse. Treatment requires urgent control of bleeding, restoration of circulating volume, blood products when indicated, and correction of associated injuries or clotting abnormalities.

### **Hemorrhagic Stroke**

A stroke caused by a blood vessel breaking and bleeding in or around the brain. It can cause sudden weakness, speech or vision problems, severe headache, vomiting, confusion, or loss of consciousness. The effects depend on the location and amount of bleeding.

### **Hemorrhagic Thyroid Nodule**

A sudden bleed inside a thyroid nodule (a lump in the thyroid gland), usually caused by the rupture of fragile blood vessels within an existing fluid-filled cyst or noncancerous growth. It produces an abrupt, tender swelling in the front of the neck that may radiate pain toward the ear or jaw, sometimes accompanied by throat tightness, hoarseness, or mild difficulty swallowing. Although the sudden enlargement can appear alarming, most hemorrhagic nodules are benign and gradually shrink as the trapped blood is reabsorbed over several weeks. It is diagnosed with a thyroid ultrasound, and needle aspiration can drain the fluid to relieve local pressure and confirm the diagnosis.

#### **Alternate Names:**

Intrathyroidal Hemorrhage; Hemorrhagic Thyroid Cyst; Bleeding Into A Thyroid Nodule; Thyroid Nodule Bleed

### **Hemorrhoid**

Singular form; see *Hemorrhoids*.

### **Hemorrhoid Surgery**

An operation to remove or reduce symptomatic internal or external hemorrhoids when bleeding, prolapse, thrombosis, or pain persists despite appropriate conservative treatment.

### **Hemorrhoidal Hemorrhage**

Rectal bleeding from enlarged or injured hemor-

rhoidal vessels, usually bright red and associated with defecation; heavy, persistent, painful, or unexplained bleeding requires evaluation for other causes.

### **Hemorrhoidectomy**

An operation to remove enlarged hemorrhoids that continue to bleed, prolapse, clot, or cause severe symptoms despite other treatment. Recovery may include significant pain, bleeding, difficulty passing urine, or temporary bowel discomfort.

### **Hemorrhoids**

Swollen, engorged vascular cushions and surrounding connective tissue in the lower rectum and anal canal, commonly triggered by chronic straining, constipation, pregnancy, or heavy lifting. They are classified by their position relative to the pain-sensitive dentate line: internal hemorrhoids (above the dentate line) are typically painless and manifest with bright red rectal bleeding or mucosal prolapse during defecation; external hemorrhoids (below the dentate line) are covered by sensitive skin and can develop acute, exquisitely painful blood clots (thrombosed external hemorrhoids). Management ranges from dietary fiber, stool softeners, and sitz baths to office procedures (rubber band ligation) and surgical hemorrhoidectomy for advanced disease.

#### **Alternate Names:**

Piles; Hemorrhoidal Disease; Haemorrhoids

### **Hemosiderin**

An iron-containing pigment left in cells after hemoglobin from red blood cells is broken down. It can build up after repeated bleeding, excessive red-cell destruction, or too much iron in the body.

### **Hemosiderosis**

Excess deposition of hemosiderin, an iron-storage pigment, in tissues. It may follow repeated bleeding or transfusions and can affect organs such as the lungs or liver.

### **Hemosiderotic Synovitis**

Synovial inflammation with hemosiderin deposition caused by recurrent bleeding into a joint, commonly associated with hemophilia, anticoagulation,



---

trauma, or a vascular synovial lesion.

### **Hemospermia**

The presence of blood in semen, usually caused by inflammation or infection of the genitourinary tract and often self-limited, but persistent or recurrent cases require evaluation.

### **Hemostasis**

The body's process for stopping bleeding after a blood vessel is injured. Blood vessels narrow, platelets form a plug, and clotting proteins strengthen it; later, the body removes the clot as healing occurs.

### **Hemostasis And Coagulation**

The linked processes that stop bleeding. Blood vessels narrow, platelets make a plug, and clotting proteins form a stronger clot. Problems with these processes can cause excessive bleeding or unwanted clots.

### **Hemostasis Valve**

A valve or sealing device used with an intravascular catheter or sheath to limit blood loss while allowing passage of wires or instruments.

### **Hemostat**

A surgical clamping instrument with jaws that compress a blood vessel or tissue to control bleeding during a procedure. Many hemostats have a ratcheted locking mechanism that maintains pressure without continuous hand force.

#### **Alternate Names:**

Hemostatic Forceps; Artery Clamp

### **Hemostatic Agent**

A medicine or material used to stop or reduce bleeding. It may form a physical seal, help the blood clot, slow clot breakdown, or replace a missing clotting protein.

### **Hemothorax**

Blood collecting in the space between the lung and chest wall, usually after injury or surgery. It can

cause chest pain, breathlessness, low blood pressure, and reduced breath sounds and often requires urgent drainage and treatment of the bleeding source.

### **Hemovac Drain**

A closed-suction surgical drain with a compressible reservoir used to remove blood or other fluid from a wound or operative site.

### **Henderson-Hasselbalch Equation**

A mathematical derivation expressing the relationship between blood pH, the dissociation constant (pKa) of carbonic acid (6.1), and the ratio of serum bicarbonate concentration to dissolved arterial carbon dioxide tension (0.03 times PaCO<sub>2</sub>), forming the quantitative framework for clinical acid-base interpretation.

### **Hendra Virus**

A rare zoonotic virus carried by fruit bats that can infect horses and occasionally humans through close contact with infected horses. Human disease may cause severe respiratory or neurologic illness and requires public-health management.

### **Henipavirus**

A genus of paramyxoviruses that includes Nipah and Hendra viruses, which can cause severe encephalitis and respiratory disease.

### **Henle Loop**

The U-shaped hairpin segment of a renal nephron connecting the proximal convoluted tubule to the distal convoluted tubule, consisting of a water-permeable descending limb and an active sodium-potassium-chloride-transporting ascending limb that generates a hypertonic medullary interstitial gradient for urine concentration.

#### **Alternate Names:**

Loop of Henle; Ansa Nephroni

### **Henna**

A plant dye used on skin, hair, and textiles. Natural henna is usually orange-brown; so-called black henna may contain para-phenylenediamine, which can cause severe allergic skin reactions and perma-

---

nent scarring.

### Henoch-Schonlein Purpura

An inflammation of small blood vessels, now called IgA vasculitis, that mainly affects children. It can cause raised purple spots on the legs or buttocks, joint pain, abdominal pain, and kidney inflammation. Most cases improve, but kidney monitoring is important.

### Henoch-Schonlein Purpura (HSP)

Alternate name; see *Henoch-Schonlein Purpura*.

#### Alternate Names:

IgA Vasculitis; Anaphylactoid Purpura; Schönlein-Henoch Syndrome

### Henoch-SchöNlein Purpura

An older name for IgA vasculitis, a disease in which small blood vessels become inflamed. It can cause purple spots on the legs or buttocks, joint pain, abdominal pain, and kidney problems.

### HEPA Filter

A high-efficiency air filter that removes most very small airborne particles. It can reduce dust, allergens, and some infectious particles when properly fitted and maintained, but does not remove every gas or chemical.

#### Alternate Names:

High-Efficiency Particulate Air Filter

### Hepar

Latin for liver; it is used in medical terminology and in names of substances or preparations relating to the liver.

### Heparan Sulfate

A sulfated glycosaminoglycan in cell surfaces and extracellular matrix that binds growth factors, regulates signaling, and contributes to filtration-barrier and connective-tissue structure.

### Heparin

An anticoagulant medicine that reduces blood clotting by enhancing antithrombin activity. It may be

given by injection to prevent or treat blood clots.

### Heparin Binding

The attachment of a protein or other molecule to heparin, a highly charged substance related to blood clotting. Binding can alter the molecule's activity, movement, or laboratory measurement.

### Heparin Resistance

The clinical situation in which unusually high doses of unfractionated heparin (exceeding 35,000 units per 24 hours) are required to achieve a therapeutic activated partial thromboplastin time (aPTT) or anti-factor Xa level, most commonly caused by antithrombin deficiency, acute-phase reactant elevations, or high factor VIII levels.

### Heparin-Induced Thrombocytopenia

An immune reaction to heparin that lowers the platelet count while making dangerous blood clots more likely. It usually begins several days after heparin exposure. Suspected cases require stopping heparin and urgent medical treatment with a different anticoagulant.

### Hepatectomy

Surgery to remove part or, rarely, nearly all of the liver, usually to treat a tumor or severe localized liver disease. The remaining liver must be large and healthy enough to work. Bleeding, bile leakage, infection, and liver failure are important risks.

#### Alternate Names:

Liver Resection

### Hepatic

Relating to the liver. The liver processes nutrients and medicines, makes bile and important blood proteins, stores energy, and helps remove harmful substances; hepatic disease can affect one or several of these functions.

### Hepatic Adenoma

A benign, well-demarcated primary epithelial liver neoplasm occurring predominantly in young women with prolonged exposure to oral contraceptives or anabolic steroids, carrying risks of life-threatening

---

spontaneous rupture with intraperitoneal hemorrhage and malignant transformation into hepatocellular carcinoma.

**Alternate Names:**

Hepatocellular Adenoma; HCA

**Hepatic Artery**

The main artery supplying oxygen-rich blood to the liver. It divides into branches that deliver blood to the right and left parts of the organ.

**Hepatic Artery Embolization**

A minimally invasive transcatheter interventional radiological procedure in which particulate emboli or chemoembolic agents are delivered directly into hepatic arterial branches feeding hypervascular liver tumors (such as hepatocellular carcinoma) to induce selective ischemic tumor necrosis.

**Alternate Names:**

Transarterial Chemoembolization; TACE

**Hepatic Artery Occlusion**

Partial or complete blockage of the hepatic artery, usually caused by a blood clot, embolus, inflammation, injury, or a procedure. It may reduce blood flow to the liver and cause liver infarction or ischemic hepatitis, especially when portal-vein blood flow is also reduced.

**Hepatic Biopsy**

Alternate name; see *Liver Biopsy*.

**Hepatic Coma**

Severe loss of consciousness caused by advanced liver failure and buildup of toxic substances in the blood. It is a medical emergency, often preceded by confusion, sleepiness, personality change, or hand-flapping movements.

**Hepatic Cyst**

A fluid-filled sac in the liver. Most simple liver cysts are harmless, cause no symptoms, and are found by chance on imaging. Large, infected, bleeding, parasitic, or complex cysts may cause pain, pressure, fever, or other complications.

**Alternate Names:**

Simple Liver Cyst; Biliary Cyst

**Hepatic Dose Adjustment**

Changing a medicine or its dose because liver disease may slow its breakdown or removal. The correct adjustment depends on the medicine, liver function, other illnesses, and monitoring of benefit and toxicity.

**Hepatic Duct**

The duct formed by the union of the right and left liver ducts, carrying bile from the liver toward the common bile duct and intestine.

**Hepatic Encephalopathy**

Confusion or changes in alertness, behavior, movement, or consciousness caused by severe liver disease and buildup of substances that affect the brain. Symptoms can progress to coma and may be precipitated by factors such as infection, bleeding, constipation, or medicines.

**Hepatic Fissure**

A natural groove or cleft on the liver's surface. It separates parts of the liver and may contain or provide a route for blood vessels, bile ducts, or supporting ligaments.

**Hepatic Flexure**

The bend where the right-sided ascending colon turns into the transverse colon near the liver. It is a normal anatomical landmark that may be seen during imaging, surgery, or colonoscopy.

**Hepatic Granulomas**

Small collections of immune cells that form in the liver in response to infections, medicines, sarcoidosis, or other inflammatory conditions. They often cause no symptoms, but the underlying condition may cause abnormal liver tests, bile-duct blockage, fibrosis, or portal hypertension.

**Hepatic Hemangioma**

A common, usually harmless growth made from

---

small blood vessels in the liver. Most cause no symptoms and are found by chance on imaging; larger ones may cause discomfort or rarely bleeding.

### **Hepatic Hydrothorax**

A buildup of fluid around a lung caused by severe liver disease and high pressure in the abdomen. It is usually on the right side and can cause breathlessness; treatment focuses on the liver disease and fluid control.

### **Hepatic Ischemia**

Liver injury caused by inadequate hepatic blood flow or oxygen delivery, often during shock, severe hypoxemia, cardiac failure, or hepatic vascular occlusion. It can produce a rapid, marked rise in aminotransferases and may progress to acute liver failure.

### **Hepatic Lamina**

A sheet of cells in a developing embryo that contributes to the liver and its bile ducts.

### **Hepatic Necrosis**

Death of liver cells caused by severe ischemia, toxins, viral infection, autoimmune injury, or other insults; extensive necrosis can produce acute liver failure.

### **Hepatic Portal System**

A network of veins that carries blood from the digestive organs, spleen, and pancreas to the liver before the blood returns to the heart. The liver processes nutrients, medicines, and harmful substances in this blood.

### **Hepatic Steatosis**

Excess accumulation of fat in liver cells, commonly called fatty liver; causes include metabolic disease, alcohol, medicines, and other conditions, and inflammation or fibrosis may develop in some people.

### **Hepatic Surgery**

Surgical treatment of liver disease, including partial liver resection, transplantation, and procedures for tumors or biliary disorders. The operation depends

on the diagnosis, tumor extent, liver function, and overall health.

### **Hepatic Synthetic Dysfunction**

Reduced ability of the liver to make important proteins, including albumin and blood-clotting factors. It can cause swelling, easy bleeding, or abnormal clotting tests, although other problems such as vitamin K deficiency or blood-thinning medicines can also alter these tests.

### **Hepatic Trauma**

Liver injury caused by blunt or penetrating trauma, potentially producing internal bleeding and requiring urgent imaging, monitoring, or surgery based on severity.

### **Hepatic Vein**

One of the veins that carries blood away from the liver and into the large vein leading to the heart. The main hepatic veins drain different regions of the liver.

### **Hepatic Vein Obstruction**

Obstruction of hepatic venous outflow, usually from thrombosis or compression, causing abdominal pain, hepatomegaly, ascites, and potentially liver failure.

### **Hepatitis**

Hepatitis is inflammation of the liver caused by viruses, alcohol, medications, toxins, metabolic disease, or autoimmunity and may be acute or chronic.

### **Hepatitis A**

A short-term liver infection caused by hepatitis A virus. It spreads when virus from infected stool is swallowed, often through close contact or contaminated food or water. It may cause fever, tiredness, nausea, abdominal discomfort, or jaundice. Most people recover completely; chronic infection is not expected. Vaccination helps prevent it.

### **Hepatitis A Virus**

The virus that causes hepatitis A. It spreads when virus from infected stool is swallowed, commonly

---

through contaminated food, water, or close contact. It causes a short-term liver infection that may produce fever, nausea, tiredness, dark urine, pale stools, or jaundice; vaccination helps prevent it.

### **Hepatitis B**

A liver infection caused by the hepatitis B virus (HBV). It spreads when infected blood, semen, or certain other body fluids enter another person's body, including through sex, shared needles or equipment, or from an infected parent to a baby around the time of birth. Infection may be acute (short-term) or chronic (long-term). Many people have no symptoms. When symptoms occur, they may include tiredness, fever, loss of appetite, nausea, abdominal pain, joint pain, dark urine, pale stools, or yellow skin and eyes (jaundice). Chronic infection, especially when acquired in infancy or childhood, can cause liver damage, cirrhosis, liver failure, or liver cancer. Vaccination prevents hepatitis B.

### **Hepatitis B Virus**

The virus that causes hepatitis B. HBV spreads when infected blood, semen, or certain other body fluids enter another person's body, including through sex, shared needles or equipment, or around childbirth. It can cause acute (short-term) or chronic (long-term) liver infection. Chronic infection can cause liver damage, cirrhosis, liver failure, or liver cancer. Vaccination prevents hepatitis B.

### **Hepatitis B Window Period**

The interval during acute resolving hepatitis B virus infection between the disappearance of detectable hepatitis B surface antigen (HBsAg) and the appearance of protective anti-HBs antibodies, during which IgM antibody to hepatitis B core antigen (anti-HBc IgM) serves as the sole serological marker of acute infection.

### **Hepatitis C**

A liver infection caused by the hepatitis C virus (HCV). It spreads mainly when infected blood enters another person's body, especially through shared injection equipment; it can also spread through some other blood exposures and, less often, through sex

or from an infected parent to a baby. Acute infection often causes no symptoms, and some people clear the virus without treatment. In many others it becomes chronic (long-term) and can cause liver damage, cirrhosis, liver failure, or liver cancer. Direct-acting antiviral medicines cure more than 95% of infections in most people; no vaccine is available.

### **Hepatitis C Virus**

The virus that causes hepatitis C. HCV spreads mainly when infected blood enters another person's body, especially through shared needles or other injection equipment. It can also spread through some medical exposures, from an infected parent to a baby, or less often through sex. Many people have no symptoms. The infection may be acute (short-term) or become chronic (long-term), which can cause liver damage, cirrhosis, liver failure, or liver cancer. Modern antiviral medicines cure more than 95% of infections in most people, but no vaccine is available.

### **Hepatitis D**

A liver infection caused by the hepatitis D virus (HDV). HDV can infect only people who also have hepatitis B because it needs hepatitis B virus (HBV) to reproduce. Infection with both viruses at the same time is called coinfection; getting HDV after already having HBV is called superinfection. Superinfection is more likely to cause long-term liver disease, cirrhosis, liver failure, or death.

### **Hepatitis D Virus**

A virus that can multiply only when hepatitis B virus is also present. It spreads through infected blood or body fluids and can make hepatitis B more severe, increasing the risk of cirrhosis and liver cancer. Hepatitis B vaccination also prevents hepatitis D.

### **Hepatitis E**

A liver infection caused by the hepatitis E virus (HEV). It usually spreads when people swallow water or food contaminated with stool, and it can also spread through undercooked meat from infected an-



---

imals. It usually causes a short-term illness, often with tiredness, nausea, vomiting, abdominal pain, dark urine, pale stools, or jaundice. Severe illness is more likely during pregnancy and in some people with weakened immune systems. Rarely, infection becomes chronic in people with weakened immunity. No vaccine is available in the United States.

### **Hepatitis E Virus**

The virus that causes hepatitis E. HEV usually spreads through water or food contaminated with stool and can also spread through undercooked meat from infected animals. It commonly causes short-term liver infection, but it may cause severe illness during pregnancy and can occasionally become chronic in people with weakened immune systems. No vaccine is available in the United States.

### **Hepatitis Serology**

Blood testing for antibodies, antigens, and other markers of hepatitis viruses. The results may show a current or past infection, immunity from vaccination, or whether infection has become chronic. Different viruses require different tests. For example, hepatitis B testing commonly includes hepatitis B surface antigen and antibodies to hepatitis B surface and core antigens; hepatitis C testing usually starts with an antibody test and uses an HCV RNA test to show current infection. Results depend on the virus, timing, vaccination history, and other laboratory findings, so a single result may not give the full answer.

### **Hepatitis Virus**

Any of the viruses that can infect and inflame the liver. The five main hepatitis viruses are hepatitis A, B, C, D, and E. They differ in how they spread and in whether they cause short-term or long-term infection. Hepatitis A usually causes a short-term illness; hepatitis B and C can become chronic and damage the liver; hepatitis D occurs only with hepatitis B; and hepatitis E usually causes short-term illness but can be severe during pregnancy or in people with weakened immunity. Vaccines prevent hepatitis A and B, and hepatitis B vaccination also prevents hepatitis D.

### **Hepatitis Virus Panel**

A group of blood tests, usually done on one blood sample, that looks for signs of hepatitis A, B, and C infection. The tests may detect viral antigens or antibodies and may show a current infection, a past infection, or immunity after vaccination. The usual panel does not automatically test for hepatitis D or E, which require separate tests when appropriate. A positive antibody result does not always mean that the virus is still present; additional tests may be needed to confirm current infection.

### **Hepatitis Viruses**

The five main viruses that infect and inflame the liver: hepatitis A, B, C, D, and E. Hepatitis A and E usually cause short-term infection, although hepatitis E can be severe during pregnancy and may become chronic in some people with weakened immunity. Hepatitis B and C can become chronic and cause cirrhosis or liver cancer. Hepatitis D occurs only in people who also have hepatitis B. Vaccines prevent hepatitis A and B, hepatitis B vaccination also prevents hepatitis D, and effective medicines can cure hepatitis C; no vaccine is available for hepatitis C or E in the United States.

### **Hepatobiliary**

Relating to the liver, gallbladder, and bile ducts. The hepatobiliary system makes bile in the liver, stores and concentrates it in the gallbladder, and carries it through the bile ducts to the small intestine, where it helps digest fats. It also includes the liver's roles in processing nutrients, medicines, and waste products.

### **Hepatobiliary Scintigraphy**

A nuclear-medicine scan, often called a HIDA scan, that uses a small injected radioactive tracer to follow bile movement through the liver, bile ducts, gallbladder, and small intestine. A special camera records the tracer over time. The scan can show blocked bile flow, whether the gallbladder fills and empties normally, bile leaks, and findings that support acute gallbladder inflammation or other biliary disorders.

**Alternate Names:**



---

Biliary Scintigraphy; Cholescintigraphy; HIDA Scan; Hepatobiliary Iminodiacetic Acid Scan

## Hepatobiliary System

The liver, gallbladder, and bile ducts. The liver makes and secretes bile, the gallbladder stores and concentrates it, and the bile ducts carry it to the small intestine, where it helps digest fats and carry some waste products out of the liver. The liver also processes nutrients and many medicines and toxins.

## Hepatoblastoma

A rare cancer that begins in the liver and is the most common type of liver cancer in children. It usually affects children younger than 3 years. A child may have no symptoms until the tumor grows larger; possible findings include a lump or swelling in the abdomen, abdominal pain, poor appetite, nausea, vomiting, or unexplained weight loss. Evaluation may include imaging and blood tests, especially the tumor marker alpha-fetoprotein (AFP); a tissue sample may be used when needed to confirm the diagnosis.

## Hepatocellular Adenoma

A rare, usually noncancerous tumor made of mature liver cells. It occurs most often in women of reproductive age and is associated with estrogen exposure, although it can also occur with anabolic steroid use, certain inherited metabolic diseases, or without a known cause. It is often found by chance on imaging and may cause no symptoms. Larger tumors can bleed or rupture, and a small number can change into liver cancer, especially certain subtypes.

## Hepatocellular Carcinoma

A cancer that begins in liver cells called hepatocytes. It is the most common primary liver cancer in adults. It often develops in a liver affected by long-term disease, especially cirrhosis, chronic hepatitis B or C, fatty liver inflammation, or long-term heavy alcohol use, but it can occur without a known risk factor. Early disease may cause no symptoms. Later findings may include a lump or pain in the upper-right abdomen, abdominal swelling, jaundice, tiredness, poor appetite, or unexplained weight loss.

Diagnosis may use imaging, liver blood tests, and the tumor marker alpha-fetoprotein (AFP); a tissue sample is sometimes needed.

## Hepatocellular Jaundice

Jaundice caused by impaired liver cells. Injured hepatocytes cannot take up, change into a water-soluble form, or remove bilirubin normally, so bilirubin builds up in the blood and makes the skin and whites of the eyes yellow. Urine may become dark and stools may become pale; itching, tiredness, nausea, or abdominal discomfort may also occur. Causes include viral or autoimmune hepatitis, alcohol-related liver disease, medicines or toxins, cirrhosis, hemochromatosis, and Wilson disease.

## Hepatocerebral Degeneration

Usually referring to acquired hepatocerebral degeneration, a rare long-term brain disorder linked to advanced liver disease or a portosystemic shunt, an abnormal connection that lets blood bypass the liver. Harmful substances that are normally cleared by the liver, including excess manganese and other neurotoxins, can affect brain areas involved in movement and thinking. Symptoms may include tremor, slowed movement or parkinsonism, poor coordination, unsteady walking, speech changes, abnormal movements, mood or personality changes, and problems with thinking or memory. It is different from the short-lived episodes of hepatic encephalopathy, although the conditions can overlap.

## Hepatocyte

The main working cell of the liver. Hepatocytes process carbohydrates, fats, proteins, medicines, and other substances from the blood. They make bile, albumin, clotting factors, and other blood proteins; store energy as glycogen; change ammonia into urea; and help remove or break down harmful substances. Injury to hepatocytes can reduce these functions and raise liver enzymes in the blood.

## Hepatoduodenal Ligament

The thickened free right margin of the lesser omentum extending between the liver and the first part of the duodenum, transmitting the portal triad (com-

---

mon bile duct anterolaterally, proper hepatic artery anteromedially, and portal vein posteriorly) and forming the anterior boundary of the epiploic foramen.

### Hepatojugular Reflux

A bedside physical sign, also called abdominojugular reflux. Firm, steady pressure on the upper abdomen briefly increases the amount of blood returning to the heart. In a normal response, the jugular veins in the neck rise briefly and then fall while the pressure continues. A sustained rise in jugular venous pressure suggests elevated right-sided heart filling pressure or limited right-ventricle function, often in heart failure. It is a clinical finding, not a diagnosis by itself.

### Hepatolenticular Degeneration

Another name for Wilson disease, a rare inherited disorder caused by changes in the *ATP7B* gene. The altered gene prevents the liver from removing excess copper into bile, so copper builds up to harmful levels in the liver, brain, eyes, kidneys, and other tissues. It may cause hepatitis, cirrhosis, or liver failure; tremor, stiffness, poor coordination, speech or mood changes; and green, gold, or brown Kayser–Fleischer rings around the cornea. The disorder is inherited in an autosomal recessive pattern.

### Hepatologist

A physician who specializes in diseases of the liver and its connected bile ducts and gallbladder. Hepatologists commonly diagnose and manage hepatitis, fatty liver disease, cirrhosis, jaundice, liver failure, liver tumors, and complications of chronic liver disease. Some also care for related diseases of the pancreas and people being assessed for liver transplantation. Hepatology is usually a subspecialty of gastroenterology.

### Hepatology

The branch of medicine concerned mainly with the liver and also with the gallbladder, bile ducts, and related pancreas disorders. It covers how these organs work, how their diseases are diagnosed and managed, and complications such as hepatitis, fatty

liver disease, cirrhosis, jaundice, bile-flow disorders, liver cancer, and liver failure. Hepatology is usually a subspecialty of gastroenterology.

### Hepatoma

An older, nonspecific term for a liver tumor. In some contexts it is used to mean hepatocellular carcinoma, a cancer arising from liver cells, but the word alone does not show whether a tumor is benign or malignant. *Hepatocellular carcinoma* is the preferred term when a malignant liver-cell tumor is meant.

### Hepatomegaly

An enlarged liver. It may be noticed when the liver is felt below the right ribs during an examination or seen on ultrasound, CT, or MRI. Hepatomegaly is a physical finding, not a disease itself. Causes include acute hepatitis, metabolic dysfunction-associated steatotic liver disease, alcohol-related liver disease, congestion from heart or blood-flow problems, bleeding into the liver, bile-duct blockage, storage or infiltrative diseases, and primary or metastatic cancer. It may cause no symptoms or may be associated with right-upper-abdominal discomfort.

### Hepatopulmonary Syndrome

A complication of liver disease or portal hypertension in which blood vessels inside the lungs become abnormally widened. Blood then passes through the lungs without receiving enough oxygen, causing low oxygen levels in the blood. Symptoms may include shortness of breath, bluish skin, or clubbing of the fingers. Breathing difficulty and low oxygen may become worse when sitting or standing and improve when lying down; this pattern is called platypnea and orthodeoxia. The condition is identified by liver disease or portal hypertension, impaired oxygenation, and evidence of widened vessels inside the lungs.

### Hepatorenal Recess

The deep peritoneal space located posteroinferiorly between the visceral surface of the right hepatic lobe and the right kidney, representing the most gravity-dependent recess of the upper abdomen in

---

the supine position where free intraperitoneal fluid or blood preferentially collects.

**Alternate Names:**

Morison Pouch

### Hepatorenal Syndrome

Serious kidney dysfunction that occurs mainly in advanced cirrhosis, usually with portal hypertension and ascites. Widening of blood vessels in the abdominal circulation lowers the amount of blood effectively reaching the kidneys. The kidneys respond by narrowing their blood vessels and retaining salt and water, so filtration falls even though the kidneys may not initially have a primary structural disease. Hepatorenal syndrome may cause acute kidney injury (HRS-AKI), which develops rapidly, or more persistent kidney dysfunction. Diagnosis requires excluding other important causes of kidney injury, such as shock, dehydration, medicines, infection, or intrinsic kidney disease.

### Hepatosplenomegaly

Enlargement of both the liver and spleen at the same time. It is a physical finding rather than a disease itself and may be found during an examination or on imaging. Causes include infections, chronic liver disease with portal hypertension, blood-cell destruction, blood cancers, inherited storage or metabolic disorders, and other infiltrative or inflammatory diseases. It may cause abdominal fullness or discomfort, or no symptoms; other findings depend on the underlying cause. Blood tests and imaging may help identify the cause.

### Hepatotoxicity

Toxic injury to the liver caused by a medicine, herbal product, dietary supplement, poison, chemical, or other exposure. The injury may mainly damage liver cells (hepatocellular injury), slow or block bile flow (cholestatic injury), or show both patterns. It may cause no symptoms but can lead to tiredness, nausea, poor appetite, abdominal pain, itching, dark urine, or jaundice, with abnormal liver blood tests. Most cases improve after the cause is removed, but severe injury can cause acute liver failure.

### Hepatovirus

A genus in the Picornaviridae family of small RNA viruses. Its best-known human member is hepatitis A virus (HAV), which causes hepatitis A and usually spreads by the fecal–oral route, when virus from stool reaches the mouth through contaminated food, water, hands, or close contact. Other hepatoviruses infect different animal hosts and are not established causes of human disease.

### Hepcidin

A peptide hormone made mainly by liver cells and the main regulator of iron balance. When iron stores are high or inflammation is present, hepcidin binds to ferroportin, the protein that exports iron from intestinal cells, macrophages, and liver cells. Ferroportin is then removed, reducing iron absorption from food and the release of recycled or stored iron into the blood. When iron is low or the bone marrow needs more iron to make red blood cells, hepcidin levels usually fall.

### Heptachlor

A persistent organochlorine insecticide formerly used mainly on crops and for termite control. It breaks down slowly into heptachlor epoxide, which can remain in soil and the food chain for a long time. Exposure can occur through contaminated soil, food, water, air, or contact with old treated materials. Studies, especially in animals, associate exposure with effects on the liver, nervous system, reproductive system, and development; evidence of health effects in humans is limited. Most uses have been canceled or severely restricted in the United States.

### HER2

Short for human epidermal growth factor receptor 2, a protein on the surface of cells that helps control normal cell growth. It is made from the *ERBB2* gene. Some cancer cells have too much HER2 protein or extra copies of the *ERBB2* gene; these cancers may grow more quickly and spread more easily. Testing HER2 protein levels or gene copies can help classify a cancer and identify whether HER2-targeted treatment may be useful.

---

## HER2-Positive

A cancer whose cells have increased expression or amplification of human epidermal growth factor receptor 2 (HER2). Some HER2-targeted medicines may be used according to the cancer type and clinical context.

## HER2-Positive Breast Cancer

Breast cancer cells with unusually high levels of the HER2 growth receptor or extra copies of its gene. This can make the cancer grow faster, but it also allows treatment with medicines that specifically block HER2.

## Herbal Supplement

A concentrated plant-derived product marketed for health or symptom relief; composition and evidence vary, and contamination, drug interactions, liver injury, and pregnancy risks must be considered.

## Herbaspirillum

A genus of gram-negative bacteria found in soil and plants; human infection is rare but can occur opportunistically.

## Herbicide

A chemical used to kill or control unwanted plants; exposure can irritate skin, eyes, lungs, or the gastrointestinal tract, with toxicity determined by the specific compound and dose.

## Herd Immunity

Protection that occurs when enough people in a community are immune to an infection that it spreads less easily. The proportion needed varies widely by infection, and herd immunity does not protect everyone equally.

## Hereditary

Transmitted through genetic material from a parent or arising from an inherited variant; a hereditary trait may be dominant, recessive, X-linked, mitochondrial, or multifactorial.

## Hereditary Amyloidosis

A group of inherited disorders in which mutations cause misfolded proteins to form amyloid deposits in organs such as the nerves, heart, kidneys, or gastrointestinal tract.

## Hereditary and Acquired Ichthyosis Vulgaris

Ichthyosis vulgaris is a scaling disorder that may be inherited, commonly related to filaggrin deficiency, or acquired with systemic disease, medicines, or nutritional problems. It produces dry fish-like scale that is often worse on the limbs and improves with moisturizers and keratolytics.

## Hereditary Angioedema

An inherited, usually autosomal dominant and bradykinin-mediated disorder causing repeated episodes of swelling in the skin, hands, feet, stomach or bowel wall, or throat. Types 1 and 2 commonly result from deficiency or dysfunction of C1 inhibitor, often due to variants in the *SERPINC1* gene; other forms may have normal C1-inhibitor levels. Attacks may cause severe abdominal pain or swelling that narrows the upper airway and often occur without itchy hives.

## Hereditary Cancer

Cancer occurring because a person inherited a disease-causing genetic variant that raises cancer risk. Genetic counseling and appropriate risk-based screening may be recommended.

## Hereditary Coproporphria

An inherited disorder of a chemical pathway used to make heme, the oxygen-carrying part of hemoglobin. Attacks may cause severe abdominal pain, vomiting, nerve or mental symptoms, and sometimes blistering skin after sunlight exposure.

## Hereditary Diffuse Gastric Cancer

An inherited condition, often caused by a harmful CDH1 gene change, that greatly increases the risk of a hard-to-detect stomach cancer and also raises the risk of lobular breast cancer. Genetic counseling, specialist screening, and discussion of risk-reducing

---

stomach surgery may be recommended.

**Alternate Names:**

HDGC; Familial Gastric Cancer Syndrome

### Hereditary Diseases

Alternate name; see *Inherited Disease*.

### Hereditary Elliptocytosis

An inherited red-cell membrane disorder in which erythrocytes are elliptical on the blood smear; most people are asymptomatic, but hemolytic anemia can occur in severe forms.

### Hereditary Fructose Intolerance

An inherited disorder in which the body cannot properly break down fructose, a sugar in fruit, honey, and table sugar. Eating it can cause vomiting, sweating, low blood glucose, abdominal pain, or liver injury.

### Hereditary Hemorrhagic Telangiectasia

An inherited disorder in which fragile, abnormal blood vessels form in the nose, skin, lungs, brain, or liver. Recurrent nosebleeds and anemia are common; bleeding or abnormal vessels in internal organs may be serious.

**Alternate Names:**

HHT; Osler-Weber-Rendu Syndrome

### Hereditary Hyperbilirubinemia

An inherited disorder of bilirubin production, conjugation, or excretion that causes persistent or intermittent jaundice, such as Gilbert, Crigler–Najjar, Dubin–Johnson, or Rotor syndrome.

### Hereditary Leiomyomatosis And Renal Cell Cancer

An inherited condition that can cause painful skin muscle tumors, early or multiple uterine fibroids, and an aggressive type of kidney cancer. It is usually related to an FH gene change; genetic counseling and regular specialist skin, uterine, and kidney assessment are important.

**Alternate Names:**

HLRCC; Reed Syndrome

### Hereditary Motor and Sensory Neuropathy

Alternate name; see *Charcot-Marie-Tooth Disease*.

### Hereditary Multiple Exostoses

An inherited disorder in which many benign, cartilage-capped bone growths called osteochondromas form near the ends of growing bones. They usually develop during childhood or adolescence and stop forming when skeletal growth ends. They may cause pain, limited joint movement, unequal limb lengths, short stature, bone deformity, or pressure on nearby nerves and blood vessels. The condition is usually caused by a change in the *EXT1* or *EXT2* gene and follows an autosomal dominant inheritance pattern. Most growths remain benign, but a small number can change into a cancerous cartilage tumor. Also called hereditary multiple osteochondromas.

**Alternate Names:**

Hereditary Multiple Osteochondromas; Diaphyseal Aclasis

### Hereditary Neuropathy with Liability to Pressure Palsies

An autosomal dominant peripheral neuropathy caused by a 1.5-megabase microdeletion of chromosome 17p11.2 encompassing the *PMP22* gene. Characterized by recurrent, painless, transient mononeuropathies (such as peroneal, ulnar, or radial nerve palsies) triggered by minor compressive trauma or prolonged mechanical pressure, with nerve biopsy showing focal myelin sheath thickening termed tomacula.

**Alternate Names:**

HNPP; Tomaculous Neuropathy; Pressure Palsy Neuropathy

### Hereditary Nonpolyposis Colorectal Cancer (Lynch Syndrome)

An inherited cancer-predisposition syndrome usually caused by impaired DNA mismatch repair. It increases the risk of colorectal, endometrial, ovarian, gastric, urinary-tract, and other cancers; genetic counseling and risk-based surveillance are important.



---

### **Hereditary Nonspherocytic Hemolytic Anemia (HNHA)**

An inherited hemolytic anemia in which red blood cells are destroyed without the spherical shape typical of hereditary spherocytosis. Causes include red-cell enzyme defects, membrane disorders, and hemoglobin abnormalities; severity varies by the underlying genetic defect.

### **Hereditary Ovalocytosis**

An inherited red-cell membrane disorder in which many erythrocytes are oval or elliptical; it is often mild but can cause hemolytic anemia in some individuals.

### **Hereditary Paraganglioma-Pheochromocytoma Syndrome**

An inherited condition that increases the risk of tumors arising from nerve-related cells near blood vessels or in the adrenal glands. Some tumors release hormones that cause high blood pressure, headache, sweating, or palpitations; genetic counseling and lifelong imaging and hormone monitoring may be needed.

#### **Alternate Names:**

Hereditary Paraganglioma; SDH-Associated Pheochromocytoma

### **Hereditary Periodic Fever Syndromes**

A group of inherited autoinflammatory disorders causing recurrent episodes of fever and inflammation without a usual infection. Examples include familial Mediterranean fever, tumor-necrosis-factor-receptor-associated periodic syndrome, mevalonate kinase deficiency, and cryopyrin-associated periodic syndromes.

### **Hereditary Persistence of Fetal Hemoglobin**

An inherited condition in which production of fetal hemoglobin continues into adulthood. It is usually harmless and may lessen the severity of some beta-globin disorders such as sickle-cell disease.

### **Hereditary Spastic Paraplegia**

A group of inherited neurologic disorders causing progressive stiffness and weakness of the legs because of corticospinal-tract degeneration. Some forms also affect speech, vision, sensation, coordination, or other organs.

### **Hereditary Spherocytic Anemia**

An inherited red-cell membrane disorder, usually hereditary spherocytosis, in which fragile spherical erythrocytes are destroyed prematurely, causing hemolytic anemia, jaundice, and splenomegaly.

### **Hereditary Spherocytosis**

Usually an autosomal dominant inherited blood disorder in which red blood cells are round rather than disc-shaped and are broken down too quickly, mainly in the spleen; less often it is autosomal recessive. It may cause anemia, jaundice, gallstones, and an enlarged spleen.

### **Hereditary Urea Cycle Abnormality**

An inherited problem removing ammonia, a waste product, from the blood. Ammonia can build up and cause vomiting, sleepiness, confusion, seizures, coma, and brain injury, especially during illness or fasting.

### **Heredity**

The passing of genetic characteristics from parents to children through genes and chromosomes. Heredity influences appearance, body function, disease risk, medicine response, and susceptibility to some inherited conditions.

### **Heredo-Ataxia**

An inherited disorder causing gradually worsening problems with balance, walking, coordination, or speech. Different gene changes cause different forms, so diagnosis usually includes neurologic assessment and genetic testing.

### **Heredopolyneuritis**

An inherited disorder affecting many peripheral nerves. It may cause gradually worsening weakness, muscle wasting, numbness, reduced reflexes,



---

or difficulty walking, often beginning in the feet and lower legs.

### **Hering-Breuer Reflex**

An automatic lung response that slows or stops breathing in when the lungs are stretched too much. It helps protect the lungs from overinflation, especially during large breaths.

### **Hermanski-Pudlak Syndrome**

Alternate spelling; see *Hermansky-Pudlak Syndrome*.

### **Hermansky-Pudlak Syndrome**

A group of inherited disorders that can cause reduced skin, hair, and eye pigment; easy bleeding because platelets do not work normally; and, in some types, lung scarring, bowel inflammation, or low white-cell counts. Genetic and specialized blood testing help confirm the diagnosis.

### **Hernia**

A bulge formed when an organ or tissue pushes through a weak area in the muscle or connective tissue that normally holds it in place. Hernias commonly occur in the abdominal wall and may become trapped or lose their blood supply, causing severe pain.

### **Hernia Mesh**

A sheet of synthetic or biologic material used to strengthen the body wall during some hernia repairs. It may reduce recurrence, but complications can include infection, persistent pain, movement of the mesh, or another hernia.

### **Hernia Repair**

Surgery that returns the displaced tissue to its proper place and closes or strengthens the weak area in the body wall. Repair may use stitches, mesh, or a minimally invasive approach; the choice depends on the hernia and the person's health.

#### **Alternate Names:**

Herniorrhaphy; Hernioplasty

### **Herniated Disc**

A spinal disk that bulges or tears so that its inner material irritates or presses on a nearby nerve. It can cause back or neck pain, pain spreading into an arm or leg, numbness, tingling, or weakness.

### **Herniated Disk**

Displacement of disk material through a tear in the outer ring of an intervertebral disc, most commonly in the lumbar spine (often at L4-L5 and L5-S1) or the cervical spine, which may press on a nearby nerve and cause pain, numbness, tingling, or weakness.

### **Herniation**

Protrusion of an organ, tissue, or structure through an abnormal opening or weak area, as in a herniated disk or abdominal-wall hernia.

### **Herniorrhaphy**

An operation to repair a hernia, in which tissue or an organ has pushed through a weak area in the body wall. The surgeon returns the tissue and closes or reinforces the weakness, sometimes with mesh; recurrence, pain, infection, or injury to nearby structures are possible.

### **Heroin**

A highly addictive opioid that is converted to morphine in the body. Repeated use can cause opioid dependence, tolerance, and withdrawal; excessive exposure can cause opioid overdose with respiratory depression, coma, and death.

### **Heroin Toxicity**

Opioid poisoning characterized by marked drowsiness or coma, pinpoint pupils, slow or absent breathing, hypoxia, and sometimes noncardiogenic pulmonary edema. Immediate airway support and naloxone are central; recurrent respiratory depression can occur because the opioid effect may outlast a naloxone dose.

### **Herpangina**

A common childhood viral illness causing fever, sore throat, and small blisters or ulcers at the back of

---

the mouth. It spreads through respiratory or stool-contaminated secretions and usually improves on its own.

## Herpes

Infection caused by a herpesvirus. The term commonly refers to herpes simplex infection, which causes oral or genital lesions, but other herpesviruses cause distinct diseases.

## Herpes Encephalitis

Inflammation of the brain caused by herpes simplex virus infection.

## Herpes Gestationis

An older name for pemphigoid gestationis, an uncommon autoimmune blistering rash of pregnancy. It usually causes intensely itchy hives followed by blisters around the abdomen and may recur in later pregnancies.

## Herpes Labialis

Alternate name; see *Cold Sore*.

## Herpes Simplex

Infection with herpes simplex virus type 1 (HSV-1) or type 2 (HSV-2). HSV-1 most often causes oral herpes or cold sores, while HSV-2 most often causes genital herpes, although either type can infect either site. Many infections cause no symptoms; when symptoms occur, painful blisters or ulcers may appear and recur. The virus remains inactive in nerve cells and can reactivate later. It spreads mainly through skin-to-skin or oral contact, including when no sore is visible, and rarely passes from a parent to a baby during delivery. Rare complications include eye infection, brain infection, or severe infection in a newborn or a person with weakened immunity.

## Herpes Simplex Encephalitis

A life-threatening brain infection caused most often by herpes simplex virus type 1. It can cause fever, headache, confusion, seizures, behavior changes, or loss of consciousness and needs urgent antiviral treatment.

## Herpes Simplex Infection

A viral infection caused by herpes simplex virus type 1 (HSV-1) or type 2 (HSV-2). HSV-1 usually causes oral herpes, while HSV-2 usually causes genital herpes, but either type can affect the mouth or genitals. Many infections are symptom-free; painful blisters or ulcers may develop and return because the virus remains inactive in nerve cells and can reactivate. HSV spreads mainly through oral or skin-to-skin contact, including contact with skin that looks normal. Rare complications include infection of the eye or brain and severe infection in newborns or people with weakened immunity.

### Alternate Names:

HSV Infection; Cold Sores and Genital Herpes

## Herpes Simplex Keratitis

Infection or inflammation of the cornea caused by herpes simplex virus, usually producing eye pain, redness, tearing, light sensitivity, or blurred vision. Recurrent disease can scar the cornea and impair sight.

## Herpes Simplex Skin Infection

A herpes simplex infection of the skin or moist body linings that causes grouped blisters or shallow sores, often around the mouth or genitals. Widespread infection in someone with eczema can be severe.

## Herpes Simplex Type 1

HSV-1, a DNA virus commonly causing oral or perioral blisters; it can also cause genital infection and remains latent with possible recurrence.

## Herpes Simplex Virus

A double-stranded DNA virus belonging to the Herpesviridae family that causes oral and genital herpes. HSV has two serotypes: HSV-1, which typically causes oral herpes, and HSV-2, which typically causes genital herpes. The virus establishes latent infections in sensory ganglia and can reactivate periodically.

## Herpes Simplex Virus Infection

Infection with herpes simplex virus type 1 or type 2, commonly causing recurrent oral, genital, skin,

---

eye, or central nervous system disease.

### **Herpes Vegetans**

A rare long-lasting herpes simplex infection that causes thick, wart-like or ulcerated growths, usually on genital or anal skin. It occurs mainly in people with weakened immunity and may be difficult to treat.

### **Herpes Viral Culture Of Lesion**

A laboratory culture of fluid or tissue from a lesion to detect herpes simplex virus or varicella-zoster virus; molecular testing is generally more sensitive and is often preferred.

### **Herpes Zoster**

A reactivation of varicella-zoster virus in a sensory nerve ganglion causing pain, tingling, and a blistering rash in one or two adjacent dermatomes, usually on one side and not crossing the midline. Complications include postherpetic neuralgia, eye or ear disease, encephalitis, and pneumonia; lesions can transmit VZV to susceptible people until crusted.

#### **Alternate Names:**

Shingles; Zoster; Acute Posterior Ganglionitis

### **Herpes Zoster Ophthalmicus**

Shingles affecting the forehead, eyelid, or nose near the eye. It can inflame the eye and threaten vision, so a painful blistering rash in this area needs same-day medical assessment and treatment.

### **Herpesviridae**

A family of DNA viruses that can remain inactive in the body after infection and reactivate later. Human members include herpes simplex viruses, varicella-zoster virus, cytomegalovirus, Epstein–Barr virus, and human herpesviruses 6, 7, and 8. They can cause skin or mouth lesions, chickenpox, shingles, glandular fever, congenital infection, or disease in people with weakened immunity.

### **Herpesviridae Infection**

An infection caused by a herpesvirus; symptoms range from localized lesions to serious disease in newborns or people with weakened immunity.

### **Herpetic Gingivostomatitis**

The classic and most common manifestation of primary herpes simplex virus type 1 (HSV-1) infection in infants and young children aged 6 months to 5 years. It is characterized by high fever, irritability, painful submandibular and cervical lymphadenopathy, and widespread friable, inflamed gums with clusters of painful vesicles that rupture into shallow, coalescing ulcers across the buccal mucosa, tongue, and lips, often severely impairing oral intake.

#### **Alternate Names:**

Primary Herpetic Gingivostomatitis; Oral HSV-1 Stomatitis

### **Herpetic Glossitis**

Painful inflammation and sores of the tongue caused by herpes simplex virus. It may occur with blisters or ulcers elsewhere in the mouth and can be more severe or persistent in people with weakened immunity.

### **Herpetic Keratitis**

A herpes simplex infection of the clear front surface of the eye. It may cause one-sided eye pain, redness, light sensitivity, tearing, or blurred vision. A branching ulcer seen with a special dye supports the diagnosis; repeated or deep infection can scar the cornea and damage vision.

#### **Alternate Names:**

Dendritic Keratitis; Herpes Simplex Corneal Infection

### **Herpetic Stomatitis**

Inflammation and ulceration of the mouth caused by herpes simplex virus, most commonly primary HSV-1 infection, with painful vesicles or ulcers and sometimes fever or gingivitis.

### **Herpetic Whitlow**

A painful herpes simplex infection of a finger, causing redness, swelling, and grouped blisters. It can spread by contact and should not be cut open because this may spread infection.

### **Herpetiform**

Resembling herpes lesions, typically describing

---

grouped vesicles or an arrangement of lesions; the term does not by itself establish herpesvirus infection.

### **Herring Bodies**

Microscopic dilations and neurosecretory terminal swellings of unmyelinated axons within the posterior lobe of the pituitary gland (neurohypophysis) that store oxytocin and antidiuretic hormone (vasopressin) bound to neurophysin carrier proteins.

#### **Alternate Names:**

Neurosecretory Bodies; Neurohypophyseal Axonal Swellings

### **Hertel Exophthalmometer**

An optical instrument used to measure the anterior-posterior distance between the lateral orbital rim and the corneal apex to quantify proptosis or exophthalmos.

### **Hertz**

A unit used to measure frequency, equal to one complete cycle or repetition per second. It is used for sound, electrical signals, brain waves, heart signals, and other repeating events.

### **Herxheimer Reaction**

A temporary worsening of fever, chills, headache, muscle aches, or other symptoms soon after antibiotics begin for some infections, especially syphilis. It usually settles within a day and is caused by the body's response to dying bacteria.

### **Hesitation Wounds**

Multiple superficial, tentative parallel incised cuts surrounding a deeper primary fatal cut wound, classically observed in suicides involving sharp instruments (wrists, neck).

### **Hesselbach Triangle**

A naturally weaker area of the lower front abdominal wall. Direct groin hernias can push through this area, which lies between a central abdominal muscle, blood vessels, and the groin ligament.

### **Heterochromatin**

DNA that is tightly packed inside a cell's nucleus and is usually less active in making proteins. It helps organize chromosomes and control which genes are available for use.

### **Heterochromia**

A difference in iris color between the eyes or within one iris, which may be congenital or acquired from disease, trauma, medication, or inflammation.

### **Heterocyclic Amine**

A nitrogen-containing chemical compound formed mainly when meat or fish is cooked at high temperatures; several heterocyclic amines are mutagens and potential dietary carcinogens.

### **Heterogeneity**

Variation in composition, structure, or behavior within a tissue, sample, lesion, or population; in tumors, heterogeneity can affect biopsy interpretation and treatment response.

### **Heterogeneous Echotexture**

An ultrasound appearance in which different parts of a tissue look unlike one another. It is a finding, not a diagnosis, and may result from inflammation, scarring, fat, bleeding, a tumor, or normal variation.

### **Heterograft**

Alternate name; see *Xenograft*.

#### **Alternate Names:**

Heterologous Graft; Xenotransplant; Heteroplastic Graft

### **Heterologous**

Coming from a different species, source, or type. In medicine, the term may describe a graft, tissue, antibody, hormone, or genetic material obtained from a different organism or cell type.

### **Heterophil Antibody**

A heterophil antibody reacts with antigens from unrelated animal species; testing for these antibodies can help diagnose some infections, including infec-

---

tious mononucleosis.

### **Heterophile Antibody**

A heterophile antibody is an antibody that reacts with antigens from unrelated species and can interfere with some laboratory immunoassays.

### **Heterophyes Heterophyes**

A small intestinal fluke acquired by eating raw or undercooked fish; it can cause abdominal symptoms and occasionally tissue complications.

### **Heteroplasia**

Development of tissue in an abnormal location or of a type not normally found there. The effects depend on the tissue involved and may range from no symptoms to obstruction, pain, or impaired function.

### **Heteroplasmy**

The presence of more than one type of mitochondrial DNA in a cell or person, usually a mixture of normal and changed copies. The proportion of changed copies can influence whether disease develops and how severe it is.

### **Heterosexual**

Describing a person who experiences sexual or romantic attraction mainly toward people of a different sex or gender. Sexual orientation is distinct from gender identity and is not a disease or disorder.

### **Heterotaxy Syndrome**

A birth condition in which organs in the chest and abdomen are arranged differently from usual. Heart defects, abnormal spleen function, and problems with the intestines or other organs may occur.

### **Heterotopic**

Located in an abnormal or unusual place; a heterotopic pregnancy is simultaneous implantation outside the uterus and, usually, inside the uterus.

### **Heterotopic Breast Tissue Of The Vulva**

Breast-like glandular tissue located in the vulva outside the normal breast. It may respond to hormonal

changes and develop benign lesions, inflammation, or rarely malignancy; histopathologic examination may be needed for diagnosis.

### **Heterotopic Pregnancy**

A pregnancy in which one embryo implants inside the uterus while another implants outside it, usually in a fallopian tube. It is uncommon but more likely after fertility treatment; an intrauterine pregnancy does not rule out an ectopic pregnancy.

### **Heterotransplant**

A transplant of cells, tissue, or an organ between members of different species. Because the immune difference is substantial, rejection is a major concern and long-term use requires specialized treatment.

### **Heterozygote**

A heterozygote has two different alleles at a genetic locus, one inherited from each parent; phenotype depends on dominance, the variant's effect, and the surrounding genetic and environmental context.

### **Heterozygous**

Having two different alleles of a particular gene at a genetic locus. The observable effect depends on the gene and the relationship between the alleles.

### **Hexachlorobenzene**

A persistent organochlorine compound that can cause porphyria, liver injury, neurologic effects, and other toxicity after significant exposure.

### **Hexachlorophane**

An antibacterial chemical formerly used in some skin cleansers. Repeated or extensive exposure can damage the nervous system, especially in infants, so its medical use is highly restricted.

### **Hexachlorophene**

A topical antiseptic phenolic compound whose excessive or prolonged use can cause neurotoxicity and skin injury; many uses are restricted.



---

## Hexadactyly

The presence of six fingers or six toes on a hand or foot, also called polydactyly.

## Hexadecadienoic Acid

A long-chain fatty acid containing 16 carbon atoms and two double bonds, occurring in lipid metabolism and studied as a biochemical or signaling molecule.

## Hexadecatetraenoic Acid

A 16-carbon polyunsaturated fatty acid containing four double bonds, present in lipid pathways and investigated for effects on membranes and signaling.

## Hexamethonium

An older medicine that blocks nerve signals at autonomic ganglia and can lower blood pressure. It is rarely used because it also causes blurred vision, dry mouth, constipation, urinary retention, and other widespread effects.

## Hexamine

A chemical used in industry and, in some settings, as methenamine, a urinary antiseptic. It releases formaldehyde in acidic urine and is unsuitable for some kidney or liver conditions.

## Hexane

A solvent found in some industrial products and glue. Long-term inhalation can damage peripheral nerves, causing numbness, weakness, and difficulty walking; swallowing or heavy exposure can also injure the lungs and brain.

## Hexanoic Acid

A six-carbon saturated fatty acid, also called caproic acid, found in some fats and odors and used in chemical synthesis; concentrated exposure irritates skin and mucosa.

## Hexokinase

An enzyme that begins the use of glucose by adding a phosphate group to it inside cells. This traps glucose in the cell and allows it to enter energy-producing or storage pathways. Most tissues use

hexokinase; the liver also uses a related enzyme called glucokinase.

## Hexose

A six-carbon monosaccharide, such as glucose, fructose, or galactose, that serves as an energy substrate or metabolic intermediate.

## Hexose Monophosphate Shunt

A cytosolic metabolic alternative pathway to glycolysis that oxidizes glucose-6-phosphate to generate NADPH (essential for reductive biosynthesis and maintaining reduced glutathione in erythrocytes) and ribose-5-phosphate for nucleotide synthesis.

### Alternate Names:

Pentose Phosphate Pathway; PPP

## Hexylresorcinol

A phenolic antiseptic and local anesthetic used in some throat lozenges and topical products; excessive use or allergy can cause adverse effects.

## Heyde Syndrome

A clinical association and triad comprising calcific aortic valve stenosis, acquired type 2A von Willebrand syndrome (caused by shear stress-induced degradation of high-molecular-weight vWF multimers across the stenotic valve), and gastrointestinal hemorrhage from intestinal angiodysplasias.

### Alternate Names:

Heyde's Syndrome; Aortic Stenosis and Angiodysplasia Triad

## Heyde Syndrome Triad

The clinical association between calcific aortic valve stenosis, acquired type 2A von Willebrand disease, and recurrent gastrointestinal bleeding from intestinal angiodysplasia. High shear stress across the stenotic aortic valve uncoils and cleaves high-molecular-weight von Willebrand factor multimers via the plasma protease ADAMTS13, predisposing mucosal angiodysplasias to recurrent hemorrhage; typically resolves completely following aortic valve replacement.

### Alternate Names:

Heyde Syndrome; Aortic Stenosis-Angiodysplasia



---

Syndrome

## HHT

Abbreviation; see *Hereditary Hemorrhagic Telangiectasia*.

## Hiatal

Relating to a hiatus, an opening or passage in an anatomical structure; a hiatal hernia occurs when abdominal tissue passes through the diaphragm's esophageal hiatus.

## Hiatal Hernia

Protrusion of part of the stomach through the esophageal opening in the diaphragm into the chest. It may be sliding or paraesophageal and can be associated with reflux or other symptoms.

## Hiatus

A natural opening or passage in a body structure. For example, the diaphragm has an opening through which the esophagus passes; tissue can sometimes bulge through such an opening.

## Hiatus Hernia

Another term for hiatal hernia: protrusion of part of the stomach through the esophageal hiatus of the diaphragm into the chest. It may be sliding or paraesophageal.

## Hib

*Haemophilus influenzae* type b, a bacterium that can cause serious infections such as meningitis, pneumonia, bloodstream infection, and swelling of the throat. Routine vaccination greatly reduces these infections in children.

## Hibernoma

A rare benign tumor of brown fat, usually a slow-growing painless soft-tissue mass that may feel warm because of its high metabolic and vascular activity.

## Hiccough

Alternate spelling; see *Hiccups*.

## Hiccup

Singular form; see *Hiccups*.

## Hiccups

Involuntary, spasmodic contractions of the diaphragm followed by sudden closure of the glottis, producing a characteristic sound. Persistent or intractable hiccups may reflect irritation of the diaphragm or phrenic and vagal pathways, metabolic disease, medication exposure, or central nervous system disease.

## Hidradenitis

A chronic inflammatory skin disorder causing repeated painful lumps, abscesses, draining tunnels, and scars, usually in the armpits, groin, buttocks, or under the breasts.

## Hidradenitis Suppurativa

A long-lasting skin disease causing repeated painful lumps, pockets of pus, draining tunnels, and scars, usually in the armpits, groin, buttocks, or under the breasts. It is not caused by poor hygiene and is not contagious.

## Hidradenocarcinoma

A rare malignant sweat-gland tumor that may arise de novo or from a hidradenoma, presenting as a firm nodule and capable of local recurrence and metastasis.

## Hidradenoma

A usually harmless tumor arising from sweat-gland cells, typically appearing as a single slow-growing skin lump. It may be flesh-colored, red, cyst-like, or ulcerated, so examination or removal may be needed to distinguish it from other tumors.

## Hidradenoma Papilliferum

A benign papillary tumor of apocrine or mammary-like glands, usually presenting as a solitary flesh-colored or red nodule in the vulvar or anogenital region.

---

## **Hidrocystoma**

A benign cyst of a sweat gland near the eyelid or skin, appearing as a small translucent or bluish papule that may enlarge with heat or sweating.

## **Hidrosis**

Sweating or perspiration; abnormal hidrosis may mean excessive sweating, reduced sweating, or sweating in an unusual pattern.

## **High Altitude**

Elevation high enough that reduced atmospheric pressure lowers available oxygen, commonly beginning around 2,500 meters and increasing the risk of altitude illness.

## **High Altitude Cerebral Edema**

A severe, life-threatening manifestation of acute altitude illness typically occurring above 2,500 to 3,000 meters, caused by hypoxemia-induced cerebral vasodilation, blood-brain barrier disruption, and vasogenic edema, presenting with ataxia, confusion, altered mental status, and coma.

### **Alternate Names:**

HACE

## **High Altitude Illness**

Health problems caused by traveling to high elevation before the body adapts to lower oxygen levels. Symptoms range from headache and nausea to dangerous brain or lung swelling requiring immediate descent.

## **High Arch**

An unusually high curve along the inside of the foot. It may run in families or result from a nerve or muscle disorder and can cause pain, calluses, ankle instability, or difficulty finding comfortable shoes.

### **Alternate Names:**

Pes Cavus; High-Arched Foot; Cavus Foot

## **High Blood Cholesterol Levels**

Too much cholesterol in the blood, especially LDL cholesterol, which can build up in artery walls and increase the risk of heart attack and stroke. It

usually causes no symptoms and is found with a blood test.

## **High Blood Potassium**

Elevated potassium concentration in the bloodstream; see *Hyperkalemia*.

## **High Blood Pressure**

Persistently elevated arterial blood pressure. Diagnosis generally requires properly obtained readings on more than one occasion or other appropriate confirmation. It often causes no symptoms but increases the risk of cardiovascular, cerebrovascular, kidney, and eye disease. Medicines used to lower it include diuretics, ACE inhibitors, angiotensin-receptor blockers, calcium-channel blockers, beta blockers, and other agents selected according to comorbidities and risk.

## **High Blood Pressure (Hypertension)**

Alternate name; see *High Blood Pressure*.

## **High Blood Pressure And Eye Disease**

Long-standing high blood pressure can damage retinal blood vessels and the optic nerve, causing hypertensive retinopathy or vision loss.

## **High Blood Protein**

An abnormally increased concentration of proteins in the blood, which may result from dehydration, inflammation, infection, or a plasma-cell disorder.

## **High Blood Sugar**

Hyperglycemia, an abnormally elevated concentration of glucose in the blood; persistent hyperglycemia is commonly associated with diabetes but can have other causes.

## **High Cholesterol**

An abnormally high level of cholesterol in the blood, especially low-density lipoprotein cholesterol. It often causes no symptoms but can contribute to plaque buildup in the arteries and cardiovascular disease.

---

### **High HDL Cholesterol (Hyperalphalipoproteinemia)**

An unusually high concentration of high-density lipoprotein cholesterol, caused by genetic traits, medicines, or other conditions. Its effect on cardiovascular risk depends on the underlying cause and the person's complete lipid profile.

### **High Hemoglobin Count**

A hemoglobin level above the expected range, which may reflect dehydration, chronic low oxygen, smoking, medication effects, or a blood disorder.

### **High Output Failure**

High-output heart failure, in which cardiac output is elevated or normal but inadequate for the body's unusually high metabolic or circulatory demand. Causes include severe anemia, thyrotoxicosis, beriberi, and large arteriovenous fistulas; treatment targets the cause and may include heart-failure therapy.

### **High Potassium**

Alternate name; see *Hyperkalemia*.

### **High Potassium Level**

Hyperkalemia is an elevated blood potassium concentration that can cause dangerous cardiac conduction abnormalities; confirmation, medication review, kidney assessment, ECG, and urgent treatment depend on level and symptoms.

### **High Protein Diet**

An eating pattern providing more protein than a person's usual or estimated requirement; suitability depends on nutritional needs, kidney and liver function, and the overall diet.

### **High Red Blood Cell Count**

An increased number or concentration of red blood cells in the blood, which may thicken the blood and result from dehydration, low oxygen, smoking, medications, or a bone marrow disorder.

#### **Alternate Names:**

Erythrocytosis; Polycythemia

### **High Triglycerides**

Alternate name; see *Hypertriglyceridemia*.

### **High Uric Acid Level**

An increased concentration of uric acid in the blood, which may contribute to gout or kidney stones and can result from reduced kidney clearance, diet, medicines, or metabolic disease.

### **High White Blood Cell Count**

An increased number of white blood cells, often associated with infection, inflammation, stress, medicines, or a blood or bone marrow disorder.

### **High-Altitude Cerebral Edema**

A severe, potentially fatal form of acute altitude sickness characterized by vasogenic brain edema, elevated intracranial pressure, and neuronal dysfunction occurring in unacclimatized individuals ascending rapidly to altitudes exceeding 2,500 to 3,000 meters. Cardinal signs include severe progressive headache, truncal ataxia, altered mental status, confusion, and eventually stupor or coma.

#### **Alternate Names:**

HACE; Mal de Montagne Cerebral

### **High-Altitude Pulmonary Edema**

Life-threatening fluid buildup in the lungs after rapid travel to high altitude. It causes breathlessness, cough, weakness, and sometimes pink or frothy sputum; immediate descent and emergency treatment are needed.

#### **Alternate Names:**

HAPE; High Altitude Pulmonary Edema

### **High-Density Lipoprotein**

A particle that carries cholesterol through the blood. Often called "HDL cholesterol," its level is one part of heart-risk assessment and does not by itself determine whether someone will have heart disease.

### **High-Density Lipoprotein Cholesterol**

A lipoprotein class synthesized by the liver and intestine containing apolipoproteins A-I and A-II that mediates reverse cholesterol transport by extracting

---

excess cholesterol from peripheral tissues and vascular macrophages and transporting it to the liver for excretion; inversely correlated with atherosclerotic cardiovascular risk.

**Alternate Names:**

HDL-C; Good Cholesterol

**High-Dependency Unit**

A hospital area for people who need closer observation and more treatment than a general ward provides but do not need the full support of an intensive-care unit. Staff monitor vital signs frequently and provide specialized care.

**High-Fiber Foods**

Foods rich in soluble or insoluble dietary fiber, including legumes, whole grains, vegetables, fruit, nuts, and seeds, support bowel regularity and may improve cholesterol and glucose control when increased with adequate fluid.

**High-Flow Nasal Cannula**

A device that delivers warmed, moist oxygen through small tubes in the nose at a high flow rate. It can improve breathing and oxygen levels in people with serious lung illness and is also used after removal of a breathing tube.

**High-Fructose Corn Syrup**

A liquid sweetener containing varying proportions of glucose and fructose, used in many processed foods and drinks. Like other added sugars, high intake increases total energy exposure and is associated with dental caries and cardiometabolic risk; it is not inherently unique in effect compared with equivalent added sugars.

**High-Grade Endometrial Stromal Sarcoma**

An aggressive cancer arising from supporting tissue in the uterus. It can grow into the uterine wall, blood vessels, or distant organs and requires specialist treatment based on its extent and cell features.

**High-Grade Serous Ovarian Carcinoma**

An aggressive cancer involving the ovary, fallopian

tube, or nearby lining of the abdomen. It often spreads before causing clear symptoms and may cause bloating, abdominal pain, or changes in bowel or bladder habits.

**High-Grade Surface Osteosarcoma**

An aggressive bone cancer that starts on the outer surface of a bone. It may cause pain or swelling and can spread to the lungs or other sites, so treatment usually requires specialist cancer care.

**High-Intensity Focused Ultrasound**

A treatment that concentrates sound-wave energy to heat and destroy a selected area of tissue while limiting damage nearby. Uses include some tumors and other conditions, depending on the device and treatment site.

**High-Level Disinfection**

A cleaning process that kills nearly all microorganisms but may not destroy large numbers of bacterial spores. It is used for some reusable medical equipment that touches moist body surfaces and cannot be heat-sterilized.

**High-Output Heart Failure**

A clinical heart failure syndrome occurring in the setting of an elevated resting cardiac output exceeding 8 liters per minute, driven by severely reduced systemic vascular resistance or excessive metabolic demands (such as thyrotoxicosis, severe chronic anemia, systemic arteriovenous fistulas, beriberi, or Paget disease).

**High-Pitched Cry**

An unusually shrill infant cry that can occur with pain, neurologic dysfunction, drug withdrawal, or serious illness and warrants assessment when persistent or accompanied by abnormal behavior or feeding.

**High-Resolution Computed Tomography**

A CT scan that uses very thin images and special settings to show fine detail, especially in the lungs. It can help detect scarring, inflammation, widened airways, and small-airway disease, but it uses X-

---

rays.

**Alternate Names:**

HRCT; High-Resolution Chest CT

**High-Risk Pregnancy Factors**

Preexisting maternal medical conditions, obstetric complications, or fetal conditions (such as advanced maternal age, chronic hypertension, gestational diabetes, preeclampsia, or fetal growth restriction) that increase morbidity or mortality for mother or child.

**Alternate Names:**

Maternal-Fetal Health Risks; High-Risk Gestation; Pregnancy Complication Risk Factors

**High-Sensitivity Cardiac Troponin**

A blood test that detects very small amounts of proteins released when heart muscle is injured. A raised result can occur with a heart attack or many other illnesses, so clinicians interpret repeated results with symptoms, ECG findings, and other tests.

**Alternate Names:**

hs-cTn; High-Sensitivity Troponin Assay

**Highly Pathogenic Avian Influenza**

Highly pathogenic influenza A infection in birds that can cause severe disease and mortality in poultry; selected strains can rarely infect humans after animal exposure.

**Highly Selective Vagotomy**

A type of vagotomy that cuts the vagus-nerve branches that stimulate acid production in the upper stomach while preserving the branches that help the lower stomach empty.

**Alternate Names:**

Parietal Cell Vagotomy; Proximal Gastric Vagotomy

**Hilar**

Relating to a hilum, the gateway where vessels, nerves, ducts, or bronchi enter or leave an organ, especially the lung or kidney.

**Hilar Cholangiocarcinoma**

A cancer at the place where the liver's main bile

ducts join. It can block bile flow and cause yellow skin or eyes, itching, dark urine, pale stools, weight loss, or repeated bile-duct infections.

**Hilar Lymphadenopathy**

Pathological enlargement of the bronchopulmonary (hilar) lymph nodes located at the hilum of the lungs, identifiable on chest radiography and CT, classically presenting as bilateral symmetrical hilar adenopathy in sarcoidosis or unilateral enlargement in tuberculosis and lung malignancies.

**Alternate Names:**

Hilar Adenopathy; Bronchopulmonary Lymphadenopathy

**Hill's Sign**

Hill's sign is an older physical finding described in some cases of aortic regurgitation; it is not sufficient by itself to diagnose the condition.

**Hill-Sachs Lesion**

A dent or compression fracture in the back of the upper-arm bone caused when the shoulder dislocates forward, sometimes contributing to repeated dislocations.

**Hilum**

A depression or opening where blood vessels, nerves, ducts, or other structures enter or leave an organ.

**Hilus**

The hilus is an indentation or entry point on an organ where vessels, nerves, ducts, or airways enter or leave.

**Hindbrain**

The hindbrain is the lower part of the developing brain, giving rise to the medulla, pons, and cerebellum and helping control vital functions and coordination.

**Hindgut**

The hindgut is the embryonic part of the digestive tract that forms much of the distal colon, rectum, and upper anal canal.

---

## Hinge Joint

A joint that mainly allows bending and straightening in one direction, like a door hinge. The elbow and knee are examples.

## Hip

The joint and surrounding region where the thigh bone meets the pelvis, allowing movement and bearing body weight.

## Hip Arthroscopy

A minimally invasive procedure that uses a camera and instruments inserted through small incisions to diagnose and treat selected disorders inside or around the hip joint.

## Hip Bursitis

Hip bursitis is inflammation of a bursa near the hip, commonly causing pain over the outer hip that worsens with lying on that side, stairs, or prolonged walking. Activity changes, physical therapy, medicines, or injection may help after other causes are considered.

### Alternate Names:

Trochanteric Bursitis; Greater Trochanteric Pain Syndrome

## Hip Circumference

The measurement around the widest part of the hips or buttocks, used in anthropometry and sometimes combined with waist circumference to assess body-fat distribution.

## Hip Dislocation

The ball at the top of the thighbone comes out of the hip socket. It is usually caused by major injury and causes severe pain and inability to move the leg; nearby nerves and the blood supply to the bone may also be injured.

## Hip Disorders

The hip joints are ball-and-socket joints connecting the femur to the pelvis, providing stability and a wide range of motion for walking, running, and weight-bearing activities. Common hip conditions

include osteoarthritis, fractures, bursitis, and labral tears.

## Hip Dysplasia

A problem present from birth or developing early in life in which the hip socket and upper thighbone do not fit together securely. The hip may be unstable or dislocate. Early examination and ultrasound or X-ray help guide treatment and protect hip development.

### Alternate Names:

Developmental Dysplasia of the Hip (DDH); Congenital Hip Dislocation

## Hip Fracture

A break in the upper thighbone, commonly near the femoral neck or just below it. It often follows a fall in someone with fragile bones and causes hip or groin pain, inability to bear weight, and sometimes a shortened or outward-turned leg.

## Hip Fracture In Older Adults

A fracture of the upper thighbone in an older person, commonly after a fall or minor force when bone strength is reduced. It can cause pain, loss of mobility, delirium, blood clots, and loss of independence.

### Alternate Names:

Femoral Neck Fracture; Geriatric Hip Fracture

## Hip Fracture Surgery

An operation to stabilize or replace a fractured proximal femur, using internal fixation, hemiarthroplasty, or total hip arthroplasty according to fracture pattern and patient factors.

## Hip Impingement

Abnormal contact between parts of the hip joint during movement, often due to cam or pincer morphology. It may cause groin pain, stiffness, and reduced motion and can contribute to labral or cartilage damage.

## Hip Injury

Damage involving the hip joint, surrounding muscles, tendons, ligaments, or bones; causes range



---

from strain and bursitis to fracture and dislocation, with inability to bear weight requiring prompt assessment.

### **Hip Joint**

The ball-and-socket joint where the top of the thigh-bone fits into the pelvis. It supports body weight and allows the leg to move in many directions; strong ligaments and a cartilage rim help keep it stable.

### **Hip Joint Injection**

Injection of local anesthetic, corticosteroid, or another prescribed agent into or around the hip joint for diagnosis or temporary relief of pain and inflammation.

### **Hip Joint Replacement**

Arthroplasty replaces damaged parts of the hip with prosthetic components to relieve pain and improve mobility, commonly for advanced osteoarthritis, fracture, or other destructive joint disease.

### **Hip Labral Tear**

A tear in the ring of cartilage surrounding the hip socket. It may cause groin pain, clicking, catching, stiffness, or reduced movement and can result from injury, repeated stress, or abnormal hip shape.

### **Hip Pain**

Pain in or around the hip caused by joint disease, tendon or muscle injury, bursitis, fracture, nerve problems, or pain referred from the back.

### **Hip Prosthesis**

An artificial hip joint used to replace a damaged hip. It usually has a stem placed in the thighbone, a ball, and a socket; parts may be metal, ceramic, or durable plastic and may be fixed with or without bone cement.

### **Hip Replacement**

A surgical procedure in which the diseased hip joint is replaced with a prosthetic implant (arthroplasty). Total hip replacement (THR) replaces both the femoral head and the acetabulum.

### **Alternate Names:**

Total Hip Replacement; Total Hip Arthroplasty; THR

### **Hip-Fracture Rehabilitation**

Recovery care after a broken hip that helps restore walking, strength, balance, and independence. It may include early movement, pain control, exercises, nutrition, prevention of blood clots, delirium prevention, and treatment to reduce future fracture risk.

### **Hippocampal Sclerosis**

Severe selective neuronal cell loss and astrocytic gliosis involving the pyramidal cell layer of the CA1 and CA4 subfields and the dentate gyrus of the hippocampus, representing the most common histopathological finding in medically refractory temporal lobe epilepsy.

### **Alternate Names:**

Mesial Temporal Sclerosis; MTS

### **Hippocampus**

A pair of structures deep in the brain that help form and retrieve new memories and support learning and navigation. Damage to them can cause difficulty forming new memories.

### **Hippocrates**

An ancient Greek physician traditionally regarded as a major founder of Western clinical medicine; the Hippocratic corpus emphasizes observation, prognosis, and professional conduct.

### **Hippocratic Oath**

A traditional promise that doctors make to practice medicine ethically, help patients, protect their privacy, and avoid harm. Modern versions differ by country and institution.

### **Hippotherapy**

A treatment in which a trained therapist uses the movement of a horse to support physical, occupational, or speech therapy. It may help some people improve balance, posture, muscle control, coordination, or communication, but it is tailored to individ-

---

ual goals and safety needs.

## Hippus

Hippus is rhythmic, involuntary widening and narrowing of the pupil, which may occur normally or with neurologic or eye disease.

## Hirata Disease

An uncommon autoimmune disorder characterized by recurrent episodes of severe hyperinsulinemic hypoglycemia triggered by high circulating titers of endogenous polyclonal autoantibodies directed against native human insulin. Occurs without prior exogenous insulin therapy, strongly associated with HLA-DR4 alleles and exposure to sulfhydryl-containing medications.

### Alternate Names:

Insulin Autoimmune Syndrome; IAS

## Hirayama Disease

A condition affecting the lower neck spinal cord, usually in teenage boys and young men. It causes slowly worsening, often one-sided weakness and wasting of the hand and forearm muscles. Neck bending can press the spinal cord and reduce its blood supply. Symptoms commonly stop progressing after several years. A neck MRI taken during bending helps confirm the diagnosis; early neck protection may limit damage.

### Alternate Names:

Monomelic Amyotrophy; Juvenile Cervical Flexion Myelopathy

## Hirschsprung Disease

A condition present from birth in which a section of the large intestine lacks the nerve cells needed to relax and move stool. Newborns may fail to pass stool, develop a swollen abdomen, or have severe constipation.

### Alternate Names:

Congenital Aganglionic Megacolon; Colonic Aganglionosis

## Hirschsprung's Disease

A birth condition in which part of the large intestine lacks the nerve cells needed to relax and move stool.

Newborns may fail to pass stool, develop a swollen abdomen, vomiting, or severe constipation.

## Hirsutism

Excessive growth of coarse, pigmented terminal hair in women or other people with a typically female hair pattern, in androgen-dependent areas such as the face, chest, abdomen, lower back, or inner thighs. Common causes include polycystic ovary syndrome, familial or idiopathic hirsutism, adrenal or ovarian hormone disorders, and androgenic medicines. Sudden onset, rapid progression, severe acne, deepening of the voice, increased muscle mass, or other signs of virilization require medical assessment.

### Alternate Names:

Male-Pattern Hair Growth in Women; Androgenic Hair Excess

## Hirudin

A substance first found in medicinal leech saliva that prevents blood from clotting by blocking an important clotting protein. Related medicines have been used or studied for selected clotting conditions.

## Hirudins

A family of thrombin-inhibiting anticoagulant peptides, including natural and recombinant agents, used or studied to prevent or treat clotting in selected clinical situations.

## Hirudo Medicinalis

The medicinal leech, used in selected medical procedures to relieve venous congestion after reconstructive surgery.

## His

Usually referring to the bundle of His, the atrioventricular conduction bundle that carries electrical impulses from the AV node into the right and left bundle branches.

## His Bundle Electrophysiology

An invasive electrophysiologic recording of electrical activity in the atrioventricular His bundle, used to assess conduction-system disease and localize atrioventricular block.

---

## Histamine

A chemical messenger released during allergic and inflammatory reactions. It can cause itching, redness, swelling, mucus production, narrowing of the airways, and increased stomach acid.

## Histamine Degradation

The body's breakdown and removal of histamine after it has acted. This limits effects such as itching, swelling, airway narrowing, and increased stomach acid.

## Histamine Receptor

A protein on or in a cell that responds to histamine. Different receptor types help control allergic symptoms, stomach-acid production, nerve signaling, and immune responses.

## Histamine Release

Discharge of histamine from mast cells or basophils during allergic, immune, or nonimmune activation, causing effects such as itching, vasodilation, swelling, mucus production, and bronchoconstriction.

## Histidine

Histidine is an amino acid used in protein synthesis and a precursor of histamine; it is essential during growth and can be metabolized to urocanic acid and other products.

## Histidine Decarboxylase

An enzyme that converts the amino acid histidine into histamine. It is found in human cells and some microorganisms; histamine influences allergy, stomach acid, blood-vessel responses, and immunity.

## Histidinemia

An inherited metabolic disorder caused by reduced histidase activity, leading to elevated histidine in blood and urine; many affected people have few or no clinically important symptoms.

## Histiocyte

An immune cell found in tissues that can engulf germs, damaged cells, or foreign material and help

alert other immune cells.

## Histiocytic Sarcoma

A rare aggressive cancer of cells that normally help process immune signals and foreign material. It may arise in lymph nodes, the intestine, skin, or other organs and requires specialist pathology and cancer care.

## Histiocytoma

A tumor or tumor-like proliferation of histiocytes; the term may refer to a benign fibrous skin lesion or, less commonly, a malignant histiocytic neoplasm depending on classification.

## Histiocytosis

A group of uncommon disorders in which certain immune cells build up abnormally in the skin, bones, organs, or blood. Effects range from isolated skin or bone problems to serious disease affecting several organs.

## Histiocytosis X

Historical name; see *Langerhans Cell Histiocytosis*.

## Histochemical

Histochemical means using chemical stains or reactions in tissue sections to identify particular substances or cell components.

## Histocompatibility

The degree to which donor and recipient tissues are immunologically alike. Matching tissue markers, especially HLA markers, can lower rejection risk, but antibody testing, organ type, and the recipient's health also influence transplant success.

## Histocompatibility Antigen Test

A test identifying human leukocyte antigen types, which influence tissue compatibility for transplantation and are relevant to some immune-mediated and genetic associations.

## Histocompatibility Testing

Laboratory typing of HLA and other immune mark-

---

ers to assess donor–recipient compatibility before transplantation and to investigate selected immune or transfusion-related questions.

### **Histogenesis**

The developmental process by which unspecialized cells differentiate and organize into mature tissues with characteristic structures and functions.

### **Histological Techniques**

Laboratory methods for preparing, staining, sectioning, and examining tissue under a microscope to reveal cellular architecture, deposits, organisms, or disease-related changes.

### **Histology**

The study of cells and tissues under a microscope. A laboratory prepares a thin tissue section, applies stains that show its structures, and examines it for normal features or changes caused by disease.

### **Histology Cassette**

A labeled perforated container that holds a tissue specimen during processing, allowing fixation, dehydration, paraffin infiltration, and later embedding for microscopic sectioning.

### **Histone**

A protein around which DNA winds so it can fit inside the cell nucleus. Chemical changes to histones help control which genes are turned on or off.

### **Histone Acetylation**

Addition of acetyl groups to histone proteins that usually loosens DNA packaging and can increase access of transcriptional machinery to genes.

### **Histone Acetyltransferase**

An enzyme that transfers acetyl groups to histone proteins, generally promoting a more open chromatin state and regulating gene transcription.

### **Histone Deacetylases**

Enzymes that remove acetyl groups from histones and other proteins, influencing chromatin structure,

gene expression, cell differentiation, and disease biology.

### **Histone Deacetylation**

Removal of acetyl groups from histones, often tightening chromatin and reducing transcription; histone deacetylases are targets of some anticancer drugs.

### **Histone Modification**

A chemical change to a histone protein that alters how tightly DNA is packed and can change which genes a cell uses. These changes help regulate cell growth, development, and response to disease.

### **Histopathologic Grade**

A microscopic estimate of how abnormal and aggressive tumor cells appear compared with their tissue of origin; grade contributes to prognosis and treatment planning alongside stage.

### **Histopathology**

The diagnosis of disease by examining tissue under a microscope. A pathologist studies a biopsy or surgical specimen for abnormal cells, inflammation, infection, scarring, or cancer.

### **Histoplasma**

A genus of fungi that includes *Histoplasma capsulatum*, which can cause histoplasmosis after inhalation of environmental spores.

### **Histoplasma Capsulatum**

A fungus found in soil contaminated with bird or bat droppings that can cause histoplasmosis when its spores are inhaled.

### **Histoplasma Complement Fixation**

A blood test that detects complement-fixing antibodies to *Histoplasma* and can support evaluation of histoplasmosis, although sensitivity, timing, cross-reactions, and immune status affect interpretation.

### **Histoplasmosis**

A fungal infection acquired by breathing in spores from soil contaminated with bird or bat droppings.

---

It often affects the lungs and may be mild, but can spread to other organs, especially in people with weakened immunity.

### **Histrionic Personality Disorder**

A long-term pattern of intense emotional expression, strong need for attention, and behavior that may seem dramatic, suggestible, or uncomfortable when not the focus. Diagnosis requires a persistent pattern that causes significant problems and is made by a qualified mental-health professional.

### **Hitchhiker's Thumb**

A descriptive term for an unusually flexible thumb that can bend backward at the distal joint. It is usually a harmless inherited variation unless associated with pain, instability, or a broader connective-tissue disorder.

### **HIV**

Abbreviation for human immunodeficiency virus; see *Human immunodeficiency virus*.

### **Hiv Antibodies**

Proteins made by the immune system after HIV infection and detected by blood or oral-fluid tests. A positive screening result needs confirmatory testing, and a negative result may be unreliable soon after exposure.

### **HIV in Pregnancy**

HIV infection during pregnancy can affect maternal health and can be transmitted to the fetus or newborn without effective care. Antiretroviral treatment, viral-load monitoring, obstetric planning, and newborn prophylaxis greatly reduce perinatal transmission; treatment decisions require specialist guidance.

### **HIV Infection**

Infection with human immunodeficiency virus, which progressively damages immune function if untreated.

### **Hiv Seropositivity**

The presence of detectable HIV antibodies or other blood-test evidence of infection. A reactive screening test is not final until confirmatory testing is completed.

### **HIV Test**

A blood or oral-fluid test that looks for HIV infection. Laboratory tests may detect viral material or antibodies; a reactive screening result needs confirmatory testing. Tests can be negative early in infection, so repeat testing may be needed after a recent exposure.

### **HIV Transmission And Prevention**

The spread of HIV through infected blood, semen, vaginal fluids, rectal fluids, or breast milk, prevented by testing, treatment that suppresses viral load, condoms, sterile equipment, and pre- or post-exposure prophylaxis.

### **HIV Viral Load**

The amount of HIV genetic material in the blood. It shows how actively HIV is multiplying and whether treatment is working; an “undetectable” result means the amount is below the test’s detection limit, not that HIV has been cured.

#### **Alternate Names:**

HIV Viral Burden; Quantitative HIV RNA

### **HIV-Associated Rheumatic Disease**

Joint, muscle, bone, or connective-tissue problems associated with HIV infection or its immune effects. They may include inflammatory arthritis, reactive arthritis, muscle disease, osteoporosis, or symptoms related to infection or treatment. Antiretroviral therapy suppresses viral replication and reduces transmission and complications.

### **HIV-Related Kidney Disease**

Kidney disease associated with HIV infection, its immune effects, or treatment. It may include collapsing focal segmental glomerulosclerosis, immune-complex disease, thrombotic microangiopathy, or drug toxicity; antiretroviral therapy and kidney-protective care are important.

---

## HIV/AIDS

Common combined term referring to HIV infection and its most advanced stage, AIDS; see *HIV infection* and *AIDS*.

## HIV/AIDS Immunopathogenesis

HIV enters and gradually destroys or disables important immune cells. Without treatment, immune defenses become weaker and serious infections or cancers may develop. Medicines stop most viral multiplication and allow immune recovery.

## HIV/AIDS In Pregnant Women And Infants

HIV care in pregnancy and infancy uses maternal antiretroviral therapy, viral-load monitoring, delivery and feeding planning, infant prophylaxis, and early infant diagnostic testing to reduce vertical transmission.

## HIV/AIDS Nutrition

Nutritional care for people living with HIV or AIDS. Infection, poor appetite, diarrhea, medicine side effects, or other illnesses may cause weight loss or nutrient shortages; care may include food support, supplements, treatment of infections, and review of medicines.

## Hives

The common term for urticaria, describing raised, intensely itchy welts (wheals) on the skin. They happen when immune cells in the skin release histamine, causing fluid to leak from tiny blood vessels. Individual welts typically appear quickly, change shape, and fade away within 24 hours without leaving a scar. Hives lasting under six weeks are acute, often triggered by viral infections, foods, insect stings, or medications. Hives lasting longer than six weeks are chronic and often occur without a known external trigger.

### Alternate Names:

Urticaria; Urticarial Weals

## Hives And Angioedema

Hives are raised itchy welts; angioedema is deeper swelling of the skin or mucous membranes, often

caused by allergy, medicines, or other immune reactions.

## HLA

A group of cell-surface proteins helping the immune system recognize the body's own cells, reject foreign tissue, and influence transplant compatibility.

### Alternate Names:

Human Leukocyte Antigen System; Major Histocompatibility Complex (MHC)

## Hla Antigen

A marker on the surface of cells that helps immune cells recognize what belongs to the body. HLA markers are important for matching donors and recipients in transplantation and are linked with risk of some immune diseases.

## HLA System

The HLA system is a group of cell-surface proteins that present antigens to immune cells and influence transplant matching and autoimmune risk.

## HLA-B27

An inherited immune-system marker associated with increased risk of some inflammatory conditions affecting the spine, joints, eyes, or bowel. A positive test does not prove that a person has one of these diseases.

## HLA-B27 Antigen

A cell-surface immune marker associated with increased risk of ankylosing spondylitis and related spondyloarthritis, reactive arthritis, and uveitis; a positive result is not diagnostic by itself.

## HLHS

Abbreviation; see *Hypoplastic Left Heart Syndrome*.

## HMG-CoA Reductase

An enzyme the liver uses to make cholesterol. Statin medicines partly lower blood cholesterol by blocking this enzyme.



---

## Hmgb Proteins

Proteins that help organize genetic material inside cells. When released from damaged cells, some can also signal tissue injury and promote inflammation.

## Hmgn Proteins

Proteins that help control how genetic material is arranged and used inside cells. They can influence cell growth, specialization, and repair of genetic material.

## Hoarding

Alternate name; see *Hoarding Disorder*.

## Hoarding Disorder

A mental-health condition involving persistent difficulty discarding possessions, even when they have little value. Saving items causes distress and can fill living areas so that normal activities or safety are impaired.

## Hoarseness

A change in the voice that makes it rough, weak, breathy, or strained. Common causes include a viral infection, voice overuse, smoking, reflux, or a growth on the vocal folds. Hoarseness lasting more than a few weeks needs assessment.

## Hodgkin Lymphoma

A cancer of a type of immune cell, usually beginning in lymph nodes. It often causes painless swelling of the nodes, fever, night sweats, itching, or weight loss and is diagnosed by examining a tissue sample.

## Hodgkin Lymphoma In Children

A malignant lymphoma arising from lymphoid tissue in children, characterized by Hodgkin and Reed-Sternberg cells and treated according to subtype, stage, and risk.

## Hodgkin's And Non-Hodgkin's Lymphoma Comparison

Hodgkin lymphoma is defined by characteristic Reed-Sternberg cells and commonly spreads in an or-

derly pattern, while non-Hodgkin lymphomas comprise many B-cell, T-cell, and other lymphoid cancers. Biopsy, immunophenotyping, and staging determine treatment and prognosis.

## Hodgkin's Disease

Alternate name; see *Hodgkin Lymphoma*.

## Hodgkin's Lymphoma

A B-cell lymphoma characterized by the presence of Reed-Sternberg cells (large, binucleate or multinucleate cells with owl-eye nucleoli) in a background of reactive inflammatory cells. See Hodgkin Lymphoma.

## Hoffmann Sign

A reflex test where flicking the middle fingernail makes the thumb and index finger quickly twitch inward, pointing to irritation or pressure on the spinal cord in the neck.

## Hoffmann Syndrome

An uncommon form of adult muscle weakness caused by severe, untreated underactive thyroid disease. It may cause stiff or painful muscles, cramps, enlarged-looking muscles, slow reflex relaxation, and high muscle-enzyme levels; treating the thyroid problem usually improves it.

### Alternate Names:

Hypothyroid Myopathy with Pseudohypertrophy

## Hofmann's Syndrome

A rare form of adult hypothyroid muscle disease causing muscle enlargement, stiffness, cramps, weakness, and slowed reflexes. Treating the low thyroid hormone level usually improves the muscle problems.

## Holiday Heart Syndrome

An episode of atrial or other cardiac arrhythmia occurring after heavy or binge alcohol intake, often in someone without known structural heart disease. Atrial fibrillation is the most common presentation.

---

## Holistic Medicine

An approach that considers physical, psychological, social, behavioral, and sometimes spiritual factors in health and illness. It may support patient-centered care, but individual therapies should be assessed for evidence, safety, and compatibility with effective conventional treatment.

## Hollenhorst Plaque

A bright, glistening, refractile, golden-orange cholesterol crystal embolus observed on ophthalmoscopic examination lodged at a retinal arterial bifurcation, originating from an ulcerated atheromatous plaque in the ipsilateral carotid artery or aortic arch.

### Alternate Names:

Hollenhorst's Plaque; Retinal Cholesterol Embolus

## Holmium

A rare-earth metal used in some medical lasers and radioactive tracers. Holmium laser energy can cut or vaporize tissue, especially in urologic procedures; radioactive holmium requires strict handling and safety controls.

## Holmium Laser Enucleation Of The Prostate

A procedure performed through the urine passage that uses a laser to separate and remove excess prostate tissue blocking urine flow. The tissue is removed from the bladder and examined; temporary bleeding, burning, leakage, infection, or sexual and urinary changes can occur.

### Alternate Names:

HoLEP; Holmium Prostate Enucleation

## Holoprosencephaly

A birth difference in which the developing brain does not divide fully into two cerebral hemispheres. Severity varies; it may affect facial development, movement, feeding, seizures, and intellectual development.

## Holorachischisis

A severe birth defect in which a long section of the spine and spinal cord fails to close during development. It usually causes major weakness, loss of

sensation, and other serious nervous-system problems.

## Holosystolic Murmur Differential

A heart murmur heard throughout the period when the heart's lower chambers contract. Common causes include leakage of the mitral or tricuspid valve and a hole between the lower chambers; its sound, location, and other findings help identify the cause.

## Holt-Oram Syndrome

An inherited condition causing abnormalities of the bones in one or both arms or hands and heart defects. The heart problem may be a hole between chambers or an abnormal rhythm; the severity varies widely.

## Holter Electrocardiography

Continuous ambulatory ECG recording, usually for 24 hours or longer, used to detect intermittent arrhythmias, correlate symptoms with rhythm, and assess heart-rate patterns.

## Holter Monitor

A small portable recorder that continuously tracks the heart's electrical rhythm, usually for a day or longer. It can detect intermittent abnormal rhythms and compare them with symptoms such as palpitations, dizziness, or fainting.

## Holter Monitoring

Portable, continuous electrocardiographic recording over an extended period, commonly 24 hours or longer, to detect intermittent rhythm abnormalities and correlate them with symptoms.

## Homeobox Gene

A gene that helps control how an embryo's body is laid out and how organs and tissues develop. Changes in some homeobox genes can cause birth differences or disease.

## Homeodomain Proteins

Transcription factors containing a homeodomain that bind regulatory DNA sequences and direct

---

developmental patterning and cell differentiation.

### **Homeopathy**

An alternative-health system that uses highly diluted preparations selected according to “like cures like”; reliable evidence has not shown homeopathic products to treat serious disease beyond placebo effects.

### **Homeostasis**

The body’s ability to keep internal conditions within a safe range despite changes outside it. It regulates temperature, blood sugar, blood pressure, fluid, salts, and acidity through coordinated feedback systems.

### **Homeotherm**

An organism that maintains a relatively stable internal body temperature despite changes in environmental temperature, largely through physiological regulation.

### **Homicidal Ideation**

Thoughts about killing or seriously harming another person. Any associated intent, plan, access to means, or inability to maintain safety requires immediate mental-health or emergency assessment.

### **Homicide**

The killing of one person by another, classified legally according to intent and circumstances; medical assessment may address injury patterns, cause of death, toxicology, and safeguarding.

### **Homo Sapiens**

The scientific name for modern humans, the living human species with shared biological features, genetic variation, and worldwide populations.

### **Homocysteine**

An amino acid formed during methionine metabolism. Persistently elevated blood levels may occur with genetic, nutritional, renal, or other disorders.

### **Homocystinuria**

A rare inherited disorder in which the body cannot properly process homocysteine, an amino acid. It may cause eye problems, abnormal bones, learning or developmental problems, and an increased risk of blood clots.

### **Homograft**

A transplant of tissue or an organ between two people of the same species but with different genetic makeup. The recipient’s immune system may reject it, so matching and immune-suppressing treatment are important.

### **Homologous**

Similar because of a shared origin, structure, or genetic relationship. In medicine, homologous tissue or cells come from the same species, while homologous chromosomes carry corresponding genes.

### **Homologous Recombination**

A DNA-repair and genetic-exchange process that uses a matching or nearly matching sequence as a template to repair breaks or reshuffle genetic material.

### **Homologous Transplantation**

Transplantation of cells, tissue, or an organ between genetically different people of the same species, requiring attention to compatibility, rejection, infection, and immunosuppression.

### **Homonymous Hemianopia**

A visual field deficit characterized by the loss of the same half of the visual field (either both right halves or both left halves) in both eyes. It results from a retrochiasmal lesion affecting the optic tract, lateral geniculate nucleus, optic radiation, or primary visual cortex on the side opposite the visual defect.

### **Homosexual**

Describing a person who experiences enduring romantic or sexual attraction to people of the same sex; sexual orientation is not a mental disorder.

---

## Homosexuality

Sexual or romantic attraction to people of the same sex or gender; it is not a mental disorder or pathology.

## Homovanillic Acid

A breakdown product of dopamine found in cerebrospinal fluid, blood, and urine; levels may reflect dopamine metabolism but are interpreted cautiously because many factors affect them.

## Homozygote

An individual or cell carrying two identical alleles at a particular gene locus, whether the alleles are normal, pathogenic, or another variant.

## Homozygous

Having two copies of the same version of a gene, one inherited from each parent. The effect depends on the gene and whether that version is harmless, protective, or disease-causing.

## Homozygous Alpha-1-Antitrypsin Deficiency

Severe inherited alpha-1 antitrypsin deficiency caused by disease-associated variants in both gene copies, increasing risk of early emphysema and liver disease; see *Alpha-1 Antitrypsin Deficiency*.

## Homozygous Familial Hypercholesterolemia

Severe familial hypercholesterolemia caused by pathogenic variants affecting both copies of a relevant lipid-clearance gene; LDL cholesterol can be extremely high and atherosclerotic disease may occur early.

## Honey-Bee Stings

Injuries caused by a honey-bee injecting venom into the skin. A sting usually causes local pain, redness, and swelling; a severe allergic reaction can cause breathing difficulty, fainting, or shock and requires emergency treatment.

## Honeycomb Lung

The characteristic advanced radiographic and histopathological pattern of end-stage diffuse pulmonary fibrosis, consisting of cystic air spaces (3 to 10 mm in diameter) bounded by thickened fibrotic walls, classic for usual interstitial pneumonia (UIP) and idiopathic pulmonary fibrosis.

### Alternate Names:

Honeycombing; Pulmonary Honeycombing

## Honeycomb Lung Pathology

End-stage fibrotic remodeling with multiple irregular cystic airspaces, often subpleural and lined by bronchiolar-type epithelium, accompanied by architectural distortion and interstitial fibrosis.

## Honolulu Fever

An infection caused by *Streptobacillus* bacteria, usually after contact with rats or contaminated food. It may cause fever, vomiting, headache, muscle or joint pain, and a rash and requires antibiotics.

## Hookworm

An intestinal infection caused by parasitic worms whose larvae live in contaminated soil. The larvae can enter through bare skin, and the adult worms may cause abdominal symptoms, blood loss, anemia, or poor growth.

## Hookworm Infection

Intestinal infection caused by hookworms acquired when larvae in contaminated soil penetrate the skin or are ingested. Chronic infection can cause abdominal symptoms, iron-deficiency anemia, protein loss, and impaired growth in children.

## Hookworm Infestation

Intestinal infection by hookworm nematodes, which attach to the intestinal wall and may cause iron-deficiency anemia, abdominal symptoms, or poor growth.

## Hordeolum

Alternate medical name; see *Stye*.

---

## Horizontal Gene Transfer

The movement of genetic material between microorganisms rather than from parent to offspring. It can spread traits such as antibiotic resistance through direct contact, viruses, or uptake of DNA from the surroundings.

## Horizontal Pursuit

Smooth, coordinated movement of both eyes horizontally to follow a slowly moving visual target while maintaining fixation.

## Hormonal Drugs

Medicines that contain, replace, imitate, block, or change the effects of hormones. Examples include insulin, thyroid hormones, corticosteroids, sex hormones, contraceptives, and medicines that block hormones in some cancers. Their uses and risks vary widely by medicine and condition.

## Hormonal Effects In Newborns

Transient newborn findings caused by maternal hormones or normal neonatal endocrine transitions, such as breast enlargement or vaginal bleeding; persistent or severe findings require evaluation.

## Hormone

A chemical messenger made by glands or other tissues and carried in the blood to target cells. It binds specific receptors to regulate functions such as growth, metabolism, reproduction, stress responses, and fluid balance.

## Hormone Levels

Measurements of hormones in blood, urine, or other specimens used to evaluate endocrine function; interpretation depends on the specific hormone, timing, age, sex, medications, and clinical context.

## Hormone Receptor

A protein that binds a specific hormone and converts the signal into cellular responses, either at the cell surface or inside the cell nucleus.

## Hormone Replacement Therapy

Treatment that supplies a hormone the body does not make enough of, such as thyroid, adrenal, sex, or pituitary hormones. Menopausal hormone therapy can relieve hot flashes and vaginal symptoms for selected people, but benefits and risks depend on the hormone, dose, duration, and health history.

## Hormone Therapy

Treatment that adds, lowers, blocks, or changes the action of hormones. It may be used for hormone disorders, contraception, or hormone-sensitive cancers; the type and effects depend on the condition and medicine.

## Hormone Therapy For Breast Cancer

Treatment that blocks or lowers estrogen or progesterone effects in hormone-sensitive breast cancer. It reduces stimulation of cancer cells and may be used after surgery or for advanced disease.

## Hormone Therapy for Endometriosis

Hormonal treatment used to suppress ovarian cycling and reduce endometriosis-related pain or lesion activity. Options include combined hormonal contraceptives, progestins, or other ovarian-suppressing medicines selected according to symptoms, pregnancy plans, and risks.

## Hormone Therapy For Prostate Cancer

Treatment that lowers androgen production or blocks androgen action to slow growth of hormone-sensitive prostate cancer.

## Hormone-Receptor Positive

A cancer whose cells express receptors for hormones such as estrogen or progesterone. Hormone-blocking treatment may be useful for selected cancers with these receptors.

## Hormone-Sensitive Lipase

An enzyme in fat cells that breaks stored triglycerides into fatty acids and glycerol. Its activity rises when the body needs fuel, including during fasting or exercise, and falls after insulin increases.

---

## Hormones

Chemical messengers released by glands or tissues that travel through blood or nearby spaces. They regulate growth, metabolism, reproduction, stress, fluid balance, mood, sleep, and many other body processes.

## Horn

A horn is a pointed, hard growth of keratinized tissue, usually on the skin or as a normal anatomical projection such as a uterine or spinal horn.

## Horner Syndrome

A group of eye and face changes caused by damage to certain nerves. One side may have a smaller pupil, a drooping eyelid, and less sweating; the cause can range from migraine to a serious neck, chest, or brain problem.

### Alternate Names:

Horner's Oculosympathetic Palsy; Bernard-Horner Syndrome

## Horner's Syndrome

A group of findings caused by disruption of sympathetic nerves, including a drooping eyelid, a small pupil, and reduced sweating on one side of the face.

## Hornet Stings

Hornet stings inject venom and usually cause local pain and swelling; multiple stings or allergy can cause a serious systemic reaction.

## Horripilation

Piloerection, or the temporary raising of hair and development of gooseflesh due to cold, fear, emotion, or autonomic stimulation.

## Horseshoe Kidney

A birth difference in which the lower parts of the two kidneys join across the middle of the body. Many people have no symptoms, but kidney stones, urine-flow problems, infections, or high blood pressure can occur.

## Hospice

Care for a person nearing the end of life when the focus is comfort rather than cure. Hospice supports symptoms, emotional and practical needs, and family caregivers at home, in a hospice facility, or in a hospital; eligibility rules vary by location.

## Hospice Care

Team-based care for a person nearing the end of life that prioritizes comfort, symptom control, communication, and family support rather than disease-directed cure.

## Hospice Rehabilitation

Rehabilitation provided during hospice care to maximize comfort, dignity, and participation rather than to restore function at all costs. It may include positioning, pain relief, safe transfers, equipment, and caregiver education.

## Hospital Code

An announcement used in a hospital to summon staff for an emergency. The number or color and the required response depend on that hospital's policy.

## Hospital Latex Allergy Precautions

Clinical protocols and institutional safety measures implemented to protect sensitized hospital patients and healthcare workers from natural rubber latex exposure. Includes using non-latex gloves, latex-free medical devices, and clear allergy identification band protocols.

### Alternate Names:

Latex Allergy Protocol; Latex-Safe Environment; Natural Rubber Latex Precautions

## Hospital- and Ventilator-Associated Pneumonia

Pneumonia developing during hospitalization, including ventilator-associated pneumonia after mechanical ventilation. Diagnosis considers new infiltrates or respiratory decline plus clinical evidence of infection; antimicrobial selection depends on local resistance, prior antibiotics, and patient risk factors.



---

### **Hospital-Acquired Infection**

An infection developing during or soon after hospital care that was not present on admission. It may involve wounds, urine, lungs, blood, or devices and can be caused by resistant organisms. Examples: CLABSI, CAUTI, surgical site infection.

### **Hospital-Acquired Infections**

Infections that develop during or after hospitalisation. Major types include central line-associated bloodstream infections, catheter-associated urinary tract infections, ventilator-associated pneumonia, and surgical site infections. Risk factors include invasive devices, prolonged hospitalisation, and antimicrobial use.

### **Hospital-Acquired Pneumonia**

Lung infection that begins at least 48 hours after hospital admission and was not present when the person arrived. It may cause fever, cough, breathing difficulty, or low oxygen; treatment depends on severity, cultures, and resistance risk.

### **Hospital-Associated Disability**

New or worsened difficulty with walking, self-care, or other daily activities that develops during or after hospitalization. It may result from illness, bed rest, weakness, delirium, poor nutrition, or treatments and can persist after discharge.

### **Host**

A person or other living organism that can be infected by a virus or other pathogen under natural conditions. The host may be a human, animal, or plant. In medical contexts, the host is the patient in whom an infection or disease develops.

### **Host Cell**

A cell in which an infectious agent, parasite, or symbiont lives, replicates, or uses cellular machinery; the host-cell response influences disease and clearance.

### **Host Response**

The body's reaction to an infection, foreign material, transplanted tissue, or other stimulus. It may

include inflammation, immune defense, healing, scar formation, rejection, or tolerance.

### **Hostility**

An attitude or behavior marked by antagonism, anger, distrust, or intent to oppose another person; persistent severe hostility may occur with psychiatric, neurologic, substance-related, or situational problems.

### **Hot Flash**

Alternate singular form; see *Hot Flashes*.

### **Hot Flashes**

Sudden episodes of intense heat, sweating, flushing, and sometimes a racing heart or chills. They are common during menopause but can also result from hormone changes, medicines, cancer treatment, or other illness.

### **Hot Flushes**

Alternate name; see *Hot Flashes*.

### **Hot Tub Folliculitis**

A temporary bacterial infection of hair follicles, commonly caused by *Pseudomonas aeruginosa* after exposure to contaminated hot tubs or pools, producing itchy follicular papules or pustules.

### **Hounsfield Unit**

A number on a CT scan that describes how strongly a tissue absorbs X-rays. Air has a very low value, water is near zero, and bone has a high value; the numbers help distinguish tissues and lesions.

### **Hounsfield Units**

Numbers used on CT scans to describe how strongly tissues absorb X-rays. Air has a very low value, water is near zero, and bone has a high value; the numbers help distinguish tissues and abnormal areas.

### **Hour**

A unit of time equal to 60 minutes, commonly used in medicine to describe symptom duration, dosing

---

intervals, monitoring periods, and treatment timing.

### Hour-Glass Contraction

An abnormal tight band that divides the uterus into upper and lower sections during difficult labor. It can prevent the baby from being delivered normally and requires urgent assessment by an obstetric team.

### Housemaid's Knee

A common name for prepatellar bursitis, inflammation and fluid accumulation in the bursa over the kneecap, often caused by prolonged kneeling, trauma, infection, or inflammatory disease. Painful swelling, redness, warmth, or fever requires assessment for septic bursitis.

### Howell-Jolly Bodies

Small leftover pieces of red-cell DNA seen on a blood smear. They are usually removed by a working spleen, so their presence may indicate absent or reduced spleen function; they can also occur with severe vitamin-B12 or folate deficiency.

### Howship-Romberg Sign

Pain and paresthesias radiating along the medial and inner aspect of the thigh to the knee, caused by compression of the obturator nerve within the obturator canal by an obturator hernia, exacerbated by extension, abduction, and internal rotation of the hip.

#### Alternate Names:

Howship-Romberg Symptom; Obturator Nerve Compression Sign

### HPA (Hypothalamic-Pituitary-Adrenal Axis)

A hormone system linking parts of the brain with the adrenal glands. It helps the body respond to stress by controlling the release of cortisol, which affects blood pressure, blood sugar, immune activity, sleep, and energy.

#### Alternate Names:

Hypothalamic-Pituitary-Adrenal Axis

### HPLC

A laboratory method that separates substances in a liquid sample so they can be identified and measured. It is used to test medicines, body chemicals, poisons, and other samples.

### HPV (Human Papillomavirus)

Alternate name; see *Human Papillomavirus*.

### HPV Infection

Alternate name; see *Human Papillomavirus Infection*.

### HPV Test

A test that detects high-risk human papillomavirus genetic material or related markers in cervical samples to screen for cervical cancer risk or evaluate abnormal cervical cytology.

### HPV/Genital Warts

Alternate name; see *Human Papillomavirus* and *Genital Warts*.

### HPylori

A shorthand name for *Helicobacter pylori*, a bacterium that can live in the stomach and cause inflammation, ulcers, and increased stomach-cancer risk. Confirmed infection is treated with a combination of medicines and follow-up testing.

### HRCT (High-Resolution Computed Tomography)

Alternate name; see *High-Resolution Computed Tomography*.

### HRT (Hormone Replacement Therapy)

Hormone replacement therapy uses estrogen, with a progestogen when needed, to treat selected menopausal symptoms caused by declining ovarian hormone production. It can relieve vasomotor and genitourinary symptoms and help prevent bone loss, but benefits and risks depend on the person's age, medical history, regimen, and duration of use.

---

### **Hs-cTn**

Alternate name; see *High-Sensitivity Cardiac Troponin*.

### **Ht (Hematocrit)**

Common abbreviation for hematocrit, the percentage of blood volume occupied by red blood cells; in other contexts it may mean height.

### **Huber Needle**

A special non-coring needle used to access an implanted port for giving medicines, fluids, blood products, or taking blood samples. Correct placement and sterile technique reduce infection, bleeding, and port damage.

### **Hughes-Stovin Syndrome**

An extremely rare condition involving blood clots in deep veins and several aneurysms in the lung arteries. It may be related to Behçet disease and can cause life-threatening bleeding in the lungs.

#### **Alternate Names:**

HSS; Pulmonary Artery Aneurysm-Thrombosis Syndrome

### **Human Bites**

Injuries caused by human teeth that can inoculate mixed oral bacteria into skin, joints, tendons, or bone. Bites to the hand, deep punctures, delayed presentation, or signs of infection require urgent medical assessment, wound care, and consideration of antibiotics and tetanus prevention.

### **Human Calcitonin**

A peptide hormone produced by thyroid parafollicular cells that can lower blood calcium by inhibiting osteoclast activity; synthetic calcitonin is used selectively for bone and calcium disorders.

### **Human Chorionic Gonadotropin**

A hormone produced mainly by the placenta and measured in blood or urine to detect and monitor pregnancy. Levels normally rise early in pregnancy, but the result must be interpreted with timing, symptoms, and other findings.

#### **Alternate Names:**

HCG; Beta-HCG

### **Human Chromosome Count**

Most human body cells contain 46 chromosomes arranged in 23 pairs, including one pair of sex chromosomes; mature egg and sperm cells normally contain 23.

### **Human Cyclic Neutropenia**

Alternate name; see *Cyclic Neutropenia*.

### **Human Cytomegalovirus Infection**

Alternate name; see *Cytomegalovirus Infection*.

### **Human Embryo**

The developing human organism from fertilization through approximately the eighth week, during which the major body plan and organ systems are established.

### **Human Gastrin**

A peptide hormone produced mainly by gastric G cells that stimulates gastric acid secretion and supports growth of the gastric mucosa after food enters the stomach.

### **Human Genome**

The complete DNA sequence and genetic content of a human, organized into nuclear chromosomes and mitochondrial DNA; variants influence traits, disease risk, and treatment response.

### **Human Herpesvirus 6**

A herpesvirus that commonly infects children and can cause roseola; it may reactivate and cause serious disease in people with weakened immunity.

### **Human Herpesvirus 8**

A herpesvirus associated with Kaposi sarcoma, primary effusion lymphoma, and multicentric Castleman disease, especially in people with weakened immunity.

---

## Human Immunodeficiency Virus

A virus that attacks immune cells, especially CD4 cells, and can gradually weaken the body's defenses. Without effective treatment it may lead to AIDS and serious infections or cancers. HIV-1 is widespread globally and HIV-2 is less common; daily treatment can suppress the virus and protect health.

## Human Immunodeficiency Virus (HIV)

Alternate name; see *Human Immunodeficiency Virus*.

## Human Leukocyte Antigen

A group of cell-surface proteins encoded by genes in the major histocompatibility complex. HLA molecules present antigens to T cells and are important in immune responses and tissue matching for transplantation.

### Alternate Names:

HLA Complex; Human MHC

## Human Metapneumovirus

A respiratory virus that commonly causes cough, fever, runny nose, and wheezing. Infants, older adults, and people with weakened immunity may develop bronchiolitis, pneumonia, or severe breathing difficulty.

## Human Microbiome

The community of bacteria, viruses, fungi, archaea, and other microorganisms living on and inside the human body. These organisms and their genes can influence digestion, immune activity, metabolism, and disease, although finding a microorganism does not by itself prove that it causes illness.

## Human Nociceptin

A neuropeptide that binds the nociceptin/orphanin FQ receptor and modulates pain, stress, reward, autonomic function, and other brain and spinal-cord processes.

## Human Papillomavirus

A large group of viruses that infect the skin and moist body linings. Some types cause warts; per-

sistent infection with high-risk types can lead to precancerous changes and cancers. See also *Human Papillomavirus Infection*.

## Human Papillomavirus (HPV/Genital Warts)

Alternate name; see *Human Papillomavirus*.

## Human Papillomavirus Infection

Infection with one or more human papillomavirus (HPV) types, spread mainly through intimate skin-to-skin or sexual contact. Most infections cause no symptoms and clear without causing health problems. Some types cause warts; persistent infection with high-risk types can cause precancerous changes and cancers of the cervix, anus, genitals, or throat. Vaccination and recommended cervical screening reduce risk.

## Human Placental Lactogen

A hormone made by the placenta during pregnancy. It helps prepare the breasts for feeding and changes how the pregnant person uses fat and sugar, helping provide nutrients to the developing fetus.

## Humectant

A substance that attracts and holds water, often used in skin creams, lotions, foods, and medicines. In skincare, it draws water into the outer skin layer and can reduce dryness when used properly.

## Humeral Epicondyle

A bony prominence at the distal humerus that provides attachment for forearm muscles and ligaments; the medial and lateral epicondyles are located on opposite sides of the elbow.

## Humeral Fracture

A break in the upper-arm bone, involving the proximal, shaft, or distal humerus; pain, swelling, deformity, and possible nerve or blood-vessel injury guide urgent assessment and treatment.

## Humeroradial Joint

The part of the elbow where the upper-arm bone

---

meets the radius, one of the forearm bones. It helps the elbow bend and straighten and supports rotation of the forearm.

### **Humeroulnar Joint**

The main hinge of the elbow, where the upper-arm bone meets the ulna. It allows the arm to bend and straighten and provides much of the elbow's stability.

### **Humerus**

The long bone of the upper arm, extending from the shoulder to the elbow. It joins the shoulder blade above and the radius and ulna below; fractures can injure nearby nerves or blood vessels.

### **Humerus Fracture**

A break in the upper-arm bone caused by trauma, falls, overuse, or weakened bone. Pain, swelling, deformity, and impaired movement may occur, with assessment for nerve and vascular injury.

### **Humidifier**

A device that adds moisture to air; medical humidifiers require appropriate cleaning and water handling because contaminated equipment can disperse infectious or inflammatory material.

### **Humidifier Fever**

A flu-like illness caused by inhalation of microorganisms or biologic contaminants from a contaminated humidifier or ventilation system, often involving fever, chills, cough, and malaise.

### **Humidity**

The amount of water vapor in air, influencing heat loss, respiratory comfort, skin hydration, and survival or transmission of some microorganisms.

### **Humor**

In anatomy, a fluid of the body, especially the aqueous or vitreous humor of the eye; in ordinary language, it means a sense of amusement or comedy.

### **Humoral Immunity**

The part of the immune system that protects mainly through antibodies made by B cells. Antibodies can block germs and toxins and help other immune cells remove them.

### **Hunchback**

A nontechnical term usually referring to pronounced kyphosis, an excessive forward curvature of the upper spine. Causes include osteoporosis, vertebral compression fractures, Scheuermann disease, degenerative change, and congenital disorders; evaluation depends on age, progression, pain, and neurologic findings.

### **Hunger**

The physical drive to eat, influenced by stomach fullness, blood glucose, hormones such as ghrelin, emotions, habits, illness, and the availability or appeal of food.

### **Hungry Bone Syndrome**

A severe drop in blood calcium after treatment of overactive parathyroid glands, often after surgery. Healing bones rapidly take up calcium and other minerals, causing tingling, cramps, spasms, seizures, or abnormal heart rhythms.

### **Hunter Syndrome**

An inherited condition, usually affecting boys, in which iduronate-2-sulfatase deficiency causes complex sugars to accumulate in tissues. It can cause coarse facial features, enlarged organs, stiff joints, hearing loss, breathing problems, and developmental difficulties.

### **Hunter's Syndrome**

Alternate spelling; see *Hunter Syndrome*.

### **Huntington Disease**

A progressive inherited brain disorder caused by an abnormally long repeated section of DNA in the *HTT* gene. The altered gene produces an abnormal huntingtin protein that damages and eventually kills nerve cells involved in movement, thinking, mood, and behavior. Symptoms may include in-

---

voluntary, dance-like movements (chorea), muscle stiffness, poor coordination, changes in speech or swallowing, depression, irritability, and worsening memory or judgment. It usually begins in adulthood, but a less common juvenile form can begin during childhood or adolescence. The condition is inherited in an autosomal dominant pattern, so each child of a parent with one altered copy has a 50% chance of inheriting it.

**Alternate Names:**

Huntington's Chorea; HD

**Huntington's Chorea (HD)**

Alternate name; see *Huntington Disease*.

**Huntington's Disease (HD)**

Alternate name; see *Huntington Disease*.

**Hurler Disease**

Alternate name for severe mucopolysaccharidosis type I; see *Mucopolysaccharidosis Type I*.

**Hurler Syndrome**

Mucopolysaccharidosis type I (MPS I), the severe form, an autosomal recessive lysosomal storage disorder caused by alpha-L-iduronidase deficiency (IDUA gene mutations). Accumulation of dermatan sulfate and heparan sulfate leads to progressive multisystem involvement.

**Hurthle Cell**

A thyroid follicular cell with abundant granular mitochondria-rich cytoplasm, seen in Hashimoto thyroiditis and Hürthle-cell neoplasms.

**Hurthle Cell Cancer**

A rare thyroid cancer arising from Hurthle cells, with behavior ranging from indolent to aggressive.

**Alternate Names:**

Oxyphilic Thyroid Carcinoma; Oncocytoid Carcinoma

**Hurthle Cell Carcinoma (Oncocytic Carcinoma)**

A differentiated thyroid cancer composed predomi-

nantly of oncocytic, or Hurthle, cells. It may behave more aggressively than some other differentiated thyroid cancers and is evaluated with thyroid imaging, cytology or histology, staging, and molecular findings.

**Hutchinson Sign**

The presence of herpetic cutaneous vesicles on the tip, side, or root of the nose in herpes zoster ophthalmicus, indicating viral involvement of the nasociliary branch of the ophthalmic nerve (cranial nerve V1) and carrying a high risk of ocular complications.

**Alternate Names:**

Hutchinson's Sign; Nasociliary Zoster Sign

**Hutchinson Teeth**

A classic dental developmental abnormality occurring in congenital syphilis, characterized by permanent upper central incisors that are small, peg-shaped, or screwdriver-shaped with a distinctive central semilunar notch on their biting edge.

**Alternate Names:**

Hutchinson's Teeth; Notched Syphilitic Incisors

**Hutchinson Triad**

The classic diagnostic clinical triad of late congenital syphilis, consisting of chronic interstitial keratitis, eighth cranial nerve sensorineural deafness, and Hutchinson notched central incisors.

**Alternate Names:**

Hutchinson's Triad; Congenital Syphilis Triad

**Hutchinson's Teeth**

Hutchinson's teeth are permanently notched, widely spaced upper front teeth classically associated with congenital syphilis.

**Hutchinson-Gilford Progeria Syndrome**

An extremely rare, fatal autosomal dominant progeroid disorder caused by a de novo point mutation in the *LMNA* gene that produces an aberrant, farnesylated lamin A protein called progerin, resulting in accelerated nuclear instability, premature aging, alopecia, atherosclerosis, and death from stroke or myocardial infarction in adolescence.



---

**Alternate Names:**

Progeria; HGPS

**Hutchinson-Gilford Progeria Syndrome (HGPS)**

Alternate name; see *Hutchinson-Gilford Progeria Syndrome*.

**Hutchinson-Gilford Syndrome (HGPS)**

Alternate name; see *Hutchinson-Gilford Progeria Syndrome*.

**Hyaline**

A glassy, translucent material or appearance, commonly describing proteinaceous deposits or homogeneous extracellular material seen microscopically in tissues.

**Hyaline Cartilage**

The most common type of cartilage, containing fine type II collagen fibers in a firm, glassy matrix. It supports airways, covers articular surfaces, and forms much of the fetal skeleton and growth plates.

**Hyaline Casts**

Clear, tube-shaped particles sometimes seen in urine. A small number may be normal, especially after exercise or dehydration; larger numbers can occur when the kidneys are under stress and must be interpreted with other urine and blood tests.

**Hyaline Membrane**

A protein-rich layer that forms along the walls of damaged air sacs in the lungs. It blocks oxygen transfer and is classically seen in newborn respiratory distress caused by inadequate surfactant.

**Hyalitis**

Inflammation of the vitreous body of the eye, usually associated with intraocular infection, autoimmune inflammation, trauma, or another ocular disorder. It may cause floaters, blurred vision, or visual loss and requires ophthalmic assessment.

**Hyaloid Canal**

A narrow channel through the vitreous body of the eye that marks the former path of the hyaloid artery from the optic disc toward the lens during fetal development.

**Hyaluronic Acid**

A naturally occurring carbohydrate polymer that binds water in connective tissue, skin, joints, and the eye.

**Alternate Names:**

Hyaluronan; Sodium Hyaluronate

**Hyaluronidase**

An enzyme that breaks down hyaluronic acid in connective tissue. Medicines containing it can help injected fluids or medicines spread through tissue and may also dissolve certain hyaluronic-acid cosmetic fillers.

**Hyaluronoglucosaminidase**

An enzyme that degrades hyaluronan and related glycosaminoglycans; hyaluronidase preparations can increase tissue permeability and assist dispersion of injected fluids or medicines.

**Hybridoma**

A cell line formed by fusing an antibody-producing B lymphocyte with an immortal myeloma cell, allowing continuous production of a monoclonal antibody.

**Hycanthone**

An older antiparasitic drug formerly used against schistosomiasis; its use was abandoned or restricted because of serious toxicity, including liver injury and carcinogenic concern.

**Hydatid Cyst**

A cyst caused by the larval stage of *Echinococcus granulosus*, a tapeworm transmitted from dogs to humans through ingestion of contaminated food. The liver is the most commonly affected organ, followed by the lungs. Cysts grow slowly and may become symptomatic from mass effect or rupture, causing anaphylaxis.

---

## Hydatid Disease

A parasitic infection caused by *Echinococcus* tapeworms. People usually become infected by swallowing eggs from infected dogs or contaminated food, water, or soil. Slow-growing fluid-filled cysts can form, most often in the liver or lungs. A leaking or burst cyst can cause a severe allergic reaction and spread the infection; treatment depends on the cyst and may include medicine, a procedure, or surgery.

### Alternate Names:

Echinococcosis; Cystic echinococcosis; Hydatidosis

## Hydatidiform Mole

A form of gestational trophoblastic disease in which abnormal trophoblastic tissue grows in the uterus. A complete mole has no viable fetus; a partial mole may contain abnormal fetal tissue. Both require follow-up because persistent disease can occur.

## Hydralazine

Hydralazine is a direct arteriolar vasodilator used for selected hypertension, including severe hypertension in pregnancy; reflex tachycardia, fluid retention, headache, and drug-induced lupus are recognized adverse effects.

## Hydramnios

Excess amniotic fluid during pregnancy, also called polyhydramnios, caused by fetal swallowing impairment, maternal diabetes, multiple gestation, fetal anomalies, infection, or sometimes no identifiable cause.

## Hydranencephaly

A rare brain development disorder in which most of the upper brain is replaced by fluid-filled spaces. It can cause severe developmental and movement problems, seizures, feeding difficulty, and a shortened life expectancy.

## Hydrarthrosis

Abnormal accumulation of clear synovial fluid within a joint, producing swelling and restricted movement; causes include osteoarthritis, inflammatory arthritis, trauma, and infection.

## Hydration

The body's water balance and the process of replacing water and electrolytes lost through urine, stool, sweat, breathing, vomiting, or bleeding.

## Hydrazine

A highly reactive chemical used in fuels and industry that can cause irritant, neurologic, hepatic, and systemic toxicity after significant exposure.

## Hydrazine Sulfate

A hydrazine-containing chemical once promoted experimentally for cancer-related symptoms; evidence of benefit is inadequate and toxicity can include liver injury, neuropathy, and seizures.

## Hydrazines

A family of nitrogen-containing chemical compounds used in industry, agriculture, and pharmaceuticals; some are irritants, toxic, mutagenic, or hepatotoxic.

## Hydrazones

Chemical compounds formed by reaction of hydrazine derivatives with aldehydes or ketones, important in medicinal chemistry, analytical chemistry, and some drug structures.

## Hydroa Vacciniforme

A rare photodermatosis causing recurrent vesicles or papules on sun-exposed skin that heal with variciform scars; some cases are associated with Epstein-Barr virus.

## Hydrocarbon

An organic compound composed only of carbon and hydrogen; ingestion or inhalation of petroleum hydrocarbons can cause chemical pneumonitis and systemic toxicity.

## Hydrocarbon Pneumonia

Lung inflammation caused by inhaling or aspirating petroleum products or other hydrocarbons, which can produce cough, breathing difficulty, and chemical pneumonitis.

---

## Hydrocarbon Toxicity

Poisoning from petroleum distillates or other hydrocarbons, with the greatest danger often being aspiration and chemical pneumonitis after ingestion. Cough, choking, hypoxia, respiratory failure, central nervous system depression, or dysrhythmias may occur; do not induce vomiting or perform routine gastric lavage, and provide respiratory and supportive care.

## Hydrocele

A collection of fluid around a testicle. It may be present from birth or develop after infection, injury, or another problem. It often causes painless swelling; examination and, when needed, ultrasound help confirm the cause.

## Hydrocele Repair

Surgery to close or remove the fluid-containing sac around a testicle when a hydrocele is persistent, large, painful, associated with a hernia, or obscures examination.

## Hydrocephalus

Hydrocephalus is an abnormal buildup of cerebrospinal fluid (CSF) in the brain's fluid-filled spaces, called ventricles. It can enlarge the ventricles and sometimes increase pressure on brain tissue. It may result from blocked CSF flow, reduced absorption, bleeding, infection, a tumor, injury, or a developmental condition. Symptoms vary with age and may include headache, vomiting, sleepiness, vision problems, poor balance, walking difficulty, memory changes, or an increasing head size in infants.

## Hydrocephalus Acquired

Acquired hydrocephalus is abnormal accumulation of cerebrospinal fluid after birth due to obstruction, impaired absorption, or rarely excess production; raised intracranial pressure may cause headache, vomiting, gait or cognitive change, and visual symptoms.

## Hydrocephalus Ex-Vacuo

Enlargement of the brain's ventricles secondary to

loss or shrinkage of surrounding brain tissue, rather than primarily from obstructed cerebrospinal-fluid flow.

## Hydrocephaly

Hydrocephalus: abnormal accumulation or circulation of cerebrospinal fluid that enlarges the ventricles and may increase pressure on the brain.

## Hydrochloric Acid

The strong acid in stomach juice that helps digest proteins and kill many swallowed microorganisms. The stomach's mucus lining and other protective mechanisms normally prevent this acid from damaging its tissues.

## Hydrochloric Acid Poisoning

A corrosive injury after swallowing, inhaling, or contacting hydrochloric acid. It can burn the mouth, throat, esophagus, stomach, skin, or lungs; do not induce vomiting and seek emergency care immediately.

## Hydrochlorothiazide

Hydrochlorothiazide is a thiazide diuretic that inhibits sodium-chloride reabsorption in the distal convoluted tubule and treats hypertension and mild oedema; hypokalaemia, hyponatraemia, hyperuricaemia, and photosensitivity may occur.

## Hydrocodone

An opioid analgesic used for moderate to severe pain and, in some formulations, cough; it can cause respiratory depression, sedation, constipation, tolerance, dependence, and overdose.

## Hydrocodone And Acetaminophen Overdose

Taking too much of this opioid-and-pain-reliever combination can slow breathing and cause coma, while acetaminophen can severely damage the liver. Emergency treatment is needed even before symptoms appear.

---

### Hydrocodone Bitartrate

The bitartrate salt of hydrocodone, an opioid analgesic that can cause sedation, constipation, respiratory depression, tolerance, dependence, and overdose.

### Hydrocodone/Oxycodone Overdose

Taking too much hydrocodone or oxycodone can cause extreme sleepiness, pinpoint pupils, slow or stopped breathing, coma, and death. Call emergency services, give naloxone if available, and provide rescue breathing if trained.

### Hydrocolpos

Distension of the vagina with fluid due to obstruction of the vaginal outlet, often from an imperforate hymen or transverse septum, presenting as a pelvic mass in neonates or adolescents.

### Hydrocortisone

A corticosteroid identical or similar to the body's cortisol. It reduces inflammation and immune activity and treats adrenal hormone deficiency; prolonged or high-dose use can cause infection, high blood sugar, skin thinning, or adrenal suppression.

### Hydrocyanic Acid Poisoning

A life-threatening poisoning that prevents cells from using oxygen. It can cause sudden headache, confusion, seizures, breathing failure, cardiovascular collapse, and death; emergency antidote treatment is required immediately.

### Hydroflumethiazide

An older thiazide diuretic that increases urinary salt and water loss to lower blood pressure or reduce swelling. It can disturb sodium, potassium, and glucose levels and may cause dehydration or increased uric acid.

### Hydrofluoric Acid Exposure

Contact with hydrofluoric acid, which can cause deep tissue injury and systemic fluoride toxicity, including dangerous low calcium and cardiac arrhythmias.

### Hydrofluoric Acid Poisoning

A medical emergency after contact with or inhalation of hydrofluoric acid. It can cause deep tissue burns and dangerous low calcium, heart rhythm changes, or death, even when the skin injury looks minor.

### Hydrogel

A water-rich three-dimensional material that can hold and release substances. Hydrogels are used in wound dressings, contact lenses, drug delivery, tissue engineering, and research, with safety depending on composition and use.

### Hydrogen Bond

A weak attraction between a hydrogen atom attached to an electronegative atom and another electronegative atom. These bonds help maintain the shapes of proteins and DNA and influence water's properties.

### Hydrogen Breath Test

A test for some causes of bloating, diarrhea, or abdominal pain. After drinking a measured sugar solution, the person breathes into collection tubes at intervals. The gases can suggest poor sugar absorption or excess bacteria in the small intestine.

### Hydrogen Cyanide

A rapidly acting cellular toxin that inhibits cytochrome oxidase and prevents oxidative phosphorylation; significant exposure can cause severe lactic acidosis, seizures, cardiovascular collapse, and death.

### Hydrogen Peroxide

An oxidizing chemical used in some antiseptic and industrial products; ingestion or concentrated exposure can injure tissue and release gas, causing embolic or respiratory complications.

### Hydrogen Peroxide Poisoning

Harm after swallowing or inhaling hydrogen peroxide. It may irritate or burn tissues and release gas that blocks blood vessels; concentrated products can cause severe injury, so urgent poison-control or

---

emergency advice is required.

### Hydrogen Sulfide

A toxic, flammable gas with a rotten-egg odor at low concentrations that inhibits cellular respiration and can rapidly cause respiratory failure, neurologic effects, or death.

### Hydrogen-Ion Concentration

The concentration of hydrogen ions in a solution, expressed through pH and central to acid–base physiology, enzyme activity, and interpretation of blood-gas results.

### Hydrogenation

A chemical process that adds hydrogen to a compound; in foods it can convert liquid oils into more solid fats.

### Hydromorphone

Hydromorphone is a potent opioid analgesic for severe pain; sedation, constipation, respiratory depression, tolerance, and dependence are important risks, especially with alcohol or other sedatives.

### Hydromorphone Overdose

Taking too much hydromorphone, an opioid, can cause extreme sleepiness, pinpoint pupils, slow or stopped breathing, coma, and death. Call emergency services and give naloxone if available; urgent monitoring is essential.

### Hydromyelia

Abnormal widening of the central canal of the spinal cord, which may be congenital or associated with obstruction of cerebrospinal-fluid flow and neurologic symptoms.

### Hydronephrosis

Swelling of a kidney because urine cannot drain normally. Causes include a stone, narrowing, reflux, a tumor, or prostate enlargement. Prolonged or severe blockage can damage the kidney and may cause pain, infection, or reduced kidney function.

### Hydronephrosis In Newborns

Swelling of a newborn's kidney caused by widened urine-collecting spaces. It may result from temporary changes, blockage, or urine flowing backward and can affect one or both kidneys.

### Hydronephrosis Of One Kidney

Dilation of one kidney's renal pelvis and calyces caused by impaired urine drainage, often from a stone, stricture, reflux, or obstruction of the ureter or ureteropelvic junction.

### Hydrophobia

An involuntary, agonizing spasm of the throat and vocal cord muscles triggered by attempting to drink or by the sight, sound, or mention of water. It is the hallmark neurological sign of furious (encephalitic) rabies, caused by viral damage to brainstem swallowing centers. The severe choking and pain make swallowing impossible, leading to a terrified avoidance of liquids, excessive drooling (hypersalivation), and often aerophobia (spasms triggered by a draft of air). Once hydrophobia appears, rabies is nearly 100% fatal, highlighting the critical need for immediate post-exposure rabies vaccination following animal bites.

#### Alternate Names:

Rabic Hydrophobia; Rabid Pharyngeal Spasms; Fear of Water in Rabies

### Hydropneumothorax

Air and fluid collected together in the space around a lung. It can compress the lung and cause chest pain or severe breathing difficulty, often requiring urgent hospital treatment and drainage.

### Hydrops

Abnormal accumulation of fluid in tissues or body cavities. The term may refer to generalized fetal edema, a localized collection, or organ-specific fluid distension; the cause depends on the clinical setting.

### Hydrops Fetalis

Severe fluid buildup in an unborn baby, affecting at least two areas such as under the skin, around the lungs, around the heart, or in the abdomen.

---

Causes include blood-group incompatibility, anemia, infection, genetic conditions, heart problems, and complications of twin pregnancy.

### **Hydrops Of The Gallbladder**

Distension of the gallbladder by clear mucus or watery fluid, usually caused by prolonged obstruction of the cystic duct by a gallstone. It may cause right-upper-quadrant pain and can progress to inflammation or infection.

### **Hydrosalpinx**

Dilatation of a fallopian tube caused by accumulation of clear fluid, usually after inflammation, infection, endometriosis, or previous pelvic surgery. It may affect fertility.

### **Hydrotherapy**

Exercise or treatment performed in water. Water supports body weight and provides gentle resistance, which may reduce pain and help movement, strength, and balance during rehabilitation. The program should match the person's condition and safety needs.

### **Hydrothorax**

An abnormal collection of clear fluid in the pleural space around a lung. It may result from heart, liver, kidney, or other disease and can cause breathlessness, cough, or chest pressure.

### **Hydroureter**

Abnormal enlargement of a ureter caused by impaired urine drainage.

### **Hydroxamic Acid**

An organic functional group containing hydroxylamine attached to a carbonyl, found in chelators and some medicines, including histone-deacetylase inhibitors.

### **Hydroxocobalamin**

A cobalt-containing form of vitamin B12 used as an antidote for cyanide poisoning. It binds cyanide to form cyanocobalamin, which is excreted in urine; it

can cause red discoloration of skin and body fluids and interfere with some laboratory tests.

### **Hydroxyapatite**

A calcium-and-phosphate mineral that gives much of their hardness to bones and teeth. Bone cells continually remove and replace it as bone grows, repairs itself, and responds to changing physical forces.

### **Hydroxychloroquine**

Hydroxychloroquine is an antimalarial and immunomodulatory drug used for malaria prevention or treatment and selected autoimmune diseases; cumulative retinal toxicity requires ophthalmic monitoring.

### **Hydroxycholesterols**

Oxidized forms of cholesterol produced in the body or during food processing. They participate in cell signaling and cholesterol regulation and may contribute to inflammation or blood-vessel disease when present in excess.

### **Hydroxylase Deficiency (21-Hydroxylase)**

An inherited condition in which the adrenal glands cannot make enough of certain hormones. It can cause salt loss, low blood pressure, and excess male-type hormones, and is the most common cause of congenital adrenal hyperplasia.

### **Hydroxylation**

An enzymatic or chemical reaction that adds a hydroxyl group to a molecule, often altering the activity, solubility, metabolism, or excretion of a drug or biomolecule.

### **Hydroxyproline**

An amino acid formed mainly by post-translational modification of collagen; its measurement has been used as an indicator of collagen turnover but is not a routine diagnostic test.

### **Hydroxyprolinemia**

A rare inherited metabolic finding caused by im-



---

paired hydroxyproline breakdown, leading to elevated hydroxyproline in blood or urine and usually little clinical illness.

### Hydroxysteroid

A steroid hormone or steroid-related compound containing a hydroxyl group; hydroxysteroids include intermediates and products of adrenal and gonadal steroid metabolism.

### Hydroxyurea

A medicine used for some blood cancers and sickle-cell disease. In sickle-cell disease it can reduce painful episodes and other complications by helping red blood cells remain less likely to sickle; blood counts and other safety tests are required.

### Hydroxyzine

Hydroxyzine is a sedating H1-antihistamine used for itching, anxiety, and preoperative sedation; drowsiness, anticholinergic effects, and QT prolongation can occur.

### Hydroxyzine Overdose

Taking too much hydroxyzine can cause severe drowsiness, confusion, agitation, seizures, low blood pressure, or dangerous heart-rhythm changes. Seek urgent poison-control or emergency advice, especially after a large or intentional dose.

### Hydrozoa

A group of mostly marine animals that includes hydroids and some jellyfish. Contact with certain species can cause painful stings or skin irritation; the medical risk depends on the species and exposure.

### Hygiene

Practices that reduce exposure to pathogens and support health, including hand hygiene, oral care, skin care, food safety, and appropriate environmental cleanliness.

### Hygiene Hypothesis

The theory that limited early-life microbial exposure may increase susceptibility to allergies and immune

disorders.

### Hygroma

A fluid-filled swelling, often caused by leakage or blockage of lymphatic vessels. A cystic hygroma usually appears in the neck of a fetus or child, while other hygromas may occur near joints or after injury.

### Hygroma Colli

Alternate name for a cystic hygroma, a congenital lymphatic malformation usually involving the neck; see *Cystic Hygroma*.

### Hymen

A thin fold of mucous membrane that partially covers the external opening of the vagina. The hymen is a remnant of fetal development, with no known physiological function in postnatal life.

### Hymenectomy

Hymenectomy is a procedure to remove or open part of the hymen, usually to treat an obstructed hymenal opening or another symptomatic abnormality.

### Hymenolepiasis

Intestinal infection with dwarf tapeworms, especially *Hymenolepis nana*, acquired by ingesting eggs; it may cause abdominal symptoms, diarrhea, or be asymptomatic.

### Hymenolepis Diminuta

A tapeworm that can infect people who accidentally swallow infected insects, usually in contaminated food. Infection may cause abdominal discomfort, diarrhea, or no symptoms and is treated with an antiparasitic medicine.

### Hymenolepis Nana

A small tapeworm that spreads when microscopic eggs are swallowed, often through contaminated hands, food, or water. It can cause abdominal pain, diarrhea, poor appetite, or no symptoms and is treatable with antiparasitic medicine.

---

### Hymenolepis Tapeworm Infection

Intestinal infection with *Hymenolepis nana* or *H. diminuta*, acquired by ingesting eggs or infected arthropods and causing variable gastrointestinal symptoms.

### Hymenoptera

An insect order that includes bees, wasps, hornets, and ants; stings or bites can cause pain, allergy, or life-threatening anaphylaxis.

### Hyoid Bone

U-shaped bone in the anterior neck at C3, unique for not directly articulating with any other bone. Suspended by muscles and the stylohyoid ligament. Provides attachment for suprahyoid (digastric, stylohyoid, mylohyoid, geniohyoid) and infrahyoid (sternohyoid, sternothyroid, thyrohyoid, omohyoid) muscles.

### Hyoscine

A medicine that blocks muscarinic acetylcholine receptors. Depending on the preparation, it can prevent motion sickness, reduce spasms, or lessen secretions; dry mouth, blurred vision, constipation, confusion, and urinary retention may occur.

### Hyoscyamine

An antimuscarinic medicine that reduces smooth-muscle spasm and secretions; it may be used for selected gastrointestinal or urinary symptoms and can cause dry mouth, blurred vision, urinary retention, or confusion.

### Hyper IgM Syndrome

A primary immunodeficiency disorder characterized by normal or elevated serum IgM concentrations accompanied by severe deficiencies in IgG, IgA, and IgE, most commonly inherited as an X-linked defect in the *CD40LG* gene encoding CD40 ligand on activated T-cells, preventing B-cell immunoglobulin isotype class switching.

**Alternate Names:**

HIGM

### Hyper-IgE Syndrome

A rare inherited immune disorder marked by very high levels of immunoglobulin E, eczema, and repeated skin or lung infections. Some forms also cause unusual bones, teeth, or blood vessels.

### Hyper-IgM Syndrome

See *Hyper-IgM Syndromes*.

### Hyper-IgM Syndromes

Inherited immune disorders in which the body makes normal or high amounts of IgM antibodies but too little IgG, IgA, or IgE. People may develop repeated bacterial, viral, or fungal infections because antibody defenses do not mature normally.

### Hyperactivity

Excessive or developmentally inappropriate motor activity, restlessness, or difficulty remaining still; persistent impairment may occur in ADHD, mania, substance effects, or other conditions.

### Hyperacusis

Reduced tolerance to ordinary environmental sounds, which may be perceived as excessively loud, painful, or distressing. It can occur with hearing disorders, migraine, head injury, neurologic conditions, or without an identifiable cause.

### Hyperadrenocorticism

Excess production or exposure to adrenal glucocorticoids, especially cortisol; in humans it causes Cushing syndrome, which may result from adrenal, pituitary, ectopic, or medication-related causes.

### Hyperaemia

Increased blood flow to a tissue, making it look red or feel warm. It may be a normal response to activity or a response to inflammation or renewed blood flow after temporary blockage.

### Hyperaesthesia

Unusually strong sensitivity to touch, pain, temperature, or other sensations. It may occur with nerve injury, inflammation, migraine, medicines, or

---

disorders affecting the brain or spinal cord.

### **Hyperaldosteronism**

Excess production of the adrenal hormone aldosterone. It can cause high blood pressure, low potassium, muscle weakness, cramps, thirst, and frequent urination; an adrenal growth or overactivity of both glands may be responsible.

### **Hyperaldosteronism Complications**

Beyond hypertension and hypokalemia, primary hyperaldosteronism causes target organ damage including left ventricular hypertrophy, fibrosis, stroke, coronary artery disease, atrial fibrillation, and chronic kidney disease. These complications occur at higher rates than in essential hypertension of similar severity.

### **Hyperalgesia**

Feeling more pain than expected from something that normally hurts. It can happen with inflammation, nerve injury, changes in nervous-system pain processing, or opioid use; it differs from allodynia, which is pain from a normally painless touch.

### **Hyperammonemia**

An abnormally high level of ammonia in the blood. It may result from severe liver disease, an inherited metabolic disorder, medicines, or other illness and can cause confusion, sleepiness, vomiting, seizures, coma, or brain swelling.

### **Hyperammonemia due to Ornithine Transcarbamylase Deficiency**

An inherited urea-cycle disorder in which ornithine transcarbamylase deficiency causes ammonia to accumulate, potentially producing vomiting, confusion, seizures, brain swelling, or coma.

### **Hyperamylasemia**

An elevated blood amylase concentration caused by pancreatic or salivary-gland disease, reduced clearance, macroamylasemia, or other conditions; it is not diagnostic of pancreatitis by itself.

### **Hyperargininemia**

An inherited urea-cycle disorder caused by arginase deficiency, leading to high arginine and progressive spasticity, developmental impairment, seizures, or liver-related findings.

#### **Alternate Names:**

Arginase Deficiency

### **Hyperbaric**

Having a pressure greater than the surrounding atmospheric pressure; hyperbaric oxygen therapy delivers oxygen at increased pressure for selected medical indications.

### **Hyperbaric Chamber**

A pressurized chamber in which a person breathes oxygen or another gas at pressure above sea level, used for conditions such as decompression sickness, air embolism, and selected wounds.

### **Hyperbaric Oxygen Therapy**

Breathing oxygen inside a chamber pressurized above normal air pressure. The treatment increases oxygen dissolved in the blood and is used for selected problems such as carbon-monoxide poisoning, decompression sickness, and certain severe wounds or infections.

### **Hyperbilirubinaemia**

Hyperbilirubinaemia is an abnormally high level of bilirubin in the blood, which can cause jaundice and may result from increased breakdown of red cells, liver disease, or blocked bile flow.

### **Hyperbilirubinemia**

An abnormally high concentration of bilirubin in blood that may cause jaundice; causes include hemolysis, impaired liver processing, hepatocellular injury, and bile-duct obstruction.

### **Hypercalcaemia**

Alternate British spelling; see *Hypercalcemia*.

### **Hypercalcemia**

An abnormally high level of calcium in the blood.

---

It may be caused by overactive parathyroid glands, cancer, medicines, or other conditions. Symptoms can include thirst, frequent urination, constipation, nausea, weakness, confusion, kidney stones, or abnormal heart rhythms.

**Alternate Names:**

High Blood Calcium; Hypercalcaemia; Elevated Serum Calcium

### **Hypercalcemia Of Malignancy**

A high blood-calcium level caused by cancer. It may occur because the cancer releases substances that raise calcium or damages bone. Severe cases can cause thirst, vomiting, weakness, confusion, kidney injury, or abnormal heart rhythms.

### **Hypercalcemia-Supravalvar Aortic Stenosis**

A combination of high blood calcium and narrowing above the aortic valve, classically associated with Williams syndrome; see *Williams Syndrome*.

### **Hypercalcemic Small Cell Ovarian Carcinoma**

A rare, highly aggressive ovarian tumor in young patients, often associated with alterations in SMARCA4 and sometimes presenting with hypercalcemia.

### **Hypercalciuria**

Excessive urinary calcium excretion, which can promote calcium kidney stones and, in some settings, bone mineral loss; evaluation considers diet, medications, kidney function, and endocrine causes.

### **Hypercapnia**

An abnormally high level of carbon dioxide in the blood, usually because breathing is not removing enough of it. It may cause headache, sleepiness, confusion, flushed skin, tremor, or coma and can occur with severe lung or muscle disease.

### **Hypercementosis**

Excessive deposition of cementum around a tooth root, producing a bulbous root contour that can complicate extraction but is often asymptomatic.

### **Hyperchloremic Acidosis**

A normal-anion-gap metabolic acidosis in which bicarbonate loss is accompanied by a relative rise in chloride. Causes include gastrointestinal bicarbonate loss, renal tubular acidosis, certain medicines, and excessive chloride administration.

### **Hypercholesterolaemia**

An abnormally high level of cholesterol in the blood, especially low-density lipoprotein cholesterol. Over time, it can contribute to fatty buildup in arteries and increase the risk of heart attack or stroke.

### **Hypercholesterolemia**

A high level of cholesterol, especially LDL cholesterol, in the blood. It usually causes no symptoms but can gradually narrow arteries and raise the risk of heart attack and stroke; causes may be inherited or related to other diseases, medicines, or diet.

### **Hypercoagulable State**

A condition in which blood clots more easily than usual, increasing the risk of deep-vein thrombosis, pulmonary embolism, stroke, or other clots. Causes may be inherited or acquired, including cancer, pregnancy, some medicines, inflammation, or prolonged immobility.

### **Hypercyanotic Tetralogy**

A sudden episode of worsening blue or gray color, rapid breathing, distress, or faintness in a child with Tetralogy of Fallot. It results from reduced blood flow to the lungs and is an emergency requiring immediate medical treatment.

### **Hyperelastic Skin**

Skin that stretches farther than usual and recoils abnormally; generalized hyperextensibility may occur in connective-tissue disorders such as Ehlers-Danlos syndromes.

### **Hyperemesis**

Excessive or persistent vomiting that can cause dehydration, abnormal body salts, weight loss, and poor nutrition. It may result from pregnancy, infection, medicines, digestive disease, brain illness, or

---

other causes.

### **Hyperemesis Gravidarum**

Severe nausea and vomiting during pregnancy that causes dehydration, weight loss, or abnormal body salts and prevents adequate food or fluid intake. It often begins early in pregnancy and may require fluids, nutrition, medicines, and hospital care.

### **Hyperemia**

Increased blood flow to a tissue, causing redness and warmth; it may be an active physiologic response to exercise or inflammation or a reactive response after temporary ischemia.

### **Hyperemia Grade**

A graded clinical or endoscopic estimate of tissue redness and vascular congestion, used to describe the severity of inflammation or mucosal irritation.

### **Hyperesthesia**

Increased sensitivity to sensory stimulation—touch (tactile), pain (hyperalgesia), or other stimuli—so that normally non-painful or mildly painful stimuli produce exaggerated discomfort.

### **Hyperextension**

Movement that extends a joint beyond its usual neutral or anatomical range, potentially stretching or tearing ligaments, capsules, muscles, or nerves.

### **Hyperfibrinogenemia**

An abnormally high level of fibrinogen, a blood protein needed for clotting. It often rises with inflammation, infection, pregnancy, obesity, or estrogen exposure and may be associated with increased clotting risk.

### **Hyperflexion**

Excessive bending of a joint beyond its normal range, which can injure ligaments, discs, muscles, or other supporting structures.

### **Hypergammaglobulinemia**

An excess of antibody proteins in the blood. It

may result from long-term infection, inflammation, liver disease, or an abnormal immune-cell growth; testing determines whether many antibody types or one type is increased.

### **Hyperglycaemia**

An abnormally high level of glucose in the blood. It may cause thirst, frequent urination, tiredness, or blurred vision and, if severe or prolonged, can damage organs or cause an emergency.

### **Hyperglycemia**

A higher-than-normal level of glucose in the blood. Severe short-term elevation can cause excessive urination, dehydration, and dangerous changes in body chemistry. Long-term elevation can damage the eyes, kidneys, nerves, heart, and blood vessels.

### **Hyperglycinemia**

An abnormally high glycine concentration in blood, most notably in nonketotic hyperglycinemia, a disorder of glycine cleavage that can cause neonatal hypotonia, seizures, and developmental impairment.

### **Hypergonadotropic Hypogonadism**

Reduced testicular or ovarian hormone production caused by gonad failure, with increased pituitary signals and possible infertility, absent periods, or delayed sexual development.

### **Hypergraphia**

Excessive, compulsive, or unusually prolonged writing or drawing, sometimes associated with temporal-lobe epilepsy, mania, or other neurologic or psychiatric conditions.

### **Hyperhidrosis**

Sweating that is excessive for the temperature or activity. It may mainly affect the palms, soles, underarms, or face without another illness, or may be caused by medicines, hormone disorders, infection, or cancer.

### **Hyperhomocysteinemia**

An elevated blood homocysteine concentration as-

---

sociated with genetic enzyme defects, folate or vitamin B12 deficiency, kidney disease, medicines, and increased vascular or pregnancy-related risk in selected settings.

### **Hypericum Perforatum**

St. John's wort, a plant used in some complementary products for depressive symptoms; it can cause drug interactions by inducing hepatic enzymes and transporters and should not be combined casually with medicines.

### **Hyperimmunization**

Repeated or intensive exposure to an antigen, such as through vaccination or laboratory immunization, intended to produce a high antibody level. The term does not by itself describe an illness or harmful "over-vaccinated" state.

### **Hyperimmunoglobulin D Syndrome**

A hereditary autoinflammatory disorder, usually caused by mevalonate kinase deficiency, characterized by recurrent fever, enlarged lymph nodes, abdominal symptoms, rash, and joint or muscle pain; it is also called HIDS or MKD.

### **Hyperimmunoglobulin E Syndrome**

A rare primary immunodeficiency characterized by very high IgE, eczema, recurrent skin and lung infections, and sometimes skeletal or dental abnormalities.

### **Hyperinsulinemia**

An abnormally high insulin level in the blood, often caused by insulin resistance, excess insulin medicine, or rarely an insulin-producing tumor. It may occur with normal or low glucose.

### **Hyperinsulinism**

Excessive insulin secretion or action causing recurrent or persistent hypoglycemia; causes include congenital disorders, tumors, medications, and critical illness.

### **Hyperkalaemia**

Alternate British spelling; see *Hyperkalemia*.

### **Hyperkalemia**

An abnormally high serum potassium level, commonly from impaired kidney excretion, medicines, tissue breakdown, adrenal insufficiency, or acidosis; pseudohyperkalemia from hemolysis must be excluded. ECG changes may progress from peaked T waves and PR prolongation to QRS widening, sine-wave morphology, ventricular fibrillation, or asystole, making severe or rapidly rising potassium an emergency requiring myocardial stabilization and rapid potassium shifting or removal.

#### **Alternate Names:**

High Blood Potassium; Hyperkalaemia; Elevated Serum Potassium

### **Hyperkalemic Periodic Paralysis**

An inherited muscle disorder causing repeated attacks of weakness or temporary paralysis. Attacks may follow rest after exercise, stress, or potassium-rich food and usually improve between episodes.

### **Hyperkeratosis**

Thickening of the outer keratin layer of skin or mucosa, producing callus, scale, or rough plaques in response to friction, inflammation, infection, inherited disorders, or other causes.

### **Hyperkinesia**

Excessive or involuntary movement, such as chorea, dystonia, tics, or tremor, resulting from abnormal activity in motor-control circuits or from drugs and other diseases.

### **Hyperlipasaemia**

An abnormally high blood lipase concentration, often associated with pancreatic inflammation but also seen with kidney failure, gastrointestinal disease, medications, and other conditions.

### **Hyperlipidaemia**

An abnormally high level of fats in the blood, including cholesterol or triglycerides. It may have no



---

symptoms but can increase the risk of artery disease, heart attack, stroke, or pancreatitis.

### **Hyperlipidemia**

Too much cholesterol or triglyceride fat in the blood. It usually causes no symptoms but can build up in artery walls, increasing the risk of heart attack and stroke; inherited disorders and other illnesses, medicines, diet, or alcohol can contribute.

### **Hyperlipoproteinemia**

An excess of one or more lipoproteins in blood, producing high cholesterol or triglycerides and increasing risk of atherosclerosis, pancreatitis, or inherited lipid disease.

### **Hyperlordosis**

Exaggerated inward curvature of the lumbar spine, often producing a swayback appearance and sometimes associated with pain, muscular imbalance, obesity, pregnancy, or structural spinal disease.

### **Hyperlordotic Posture**

An exaggerated inward curvature of the lumbar spine, often producing increased anterior pelvic tilt and a prominent lower-back arch.

### **Hyperlysinemia**

An inherited metabolic abnormality with elevated lysine in blood caused by altered lysine degradation; many people have few symptoms, but some have developmental or neurologic findings.

### **Hypermagnesemia**

An abnormally high concentration of magnesium in the blood, often related to kidney failure or excessive magnesium intake; severe cases can cause weakness, low blood pressure, depressed reflexes, or arrhythmias.

### **Hypermetropia**

Long-sightedness, in which the eye focuses light behind the retina instead of on it. Close objects may look blurred, and children may develop eye strain or an inward-turning eye.

### **Alternate Names:**

Hyperopia; Farsightedness; Long-sightedness

### **Hypermobile Ehlers-Danlos Syndrome**

An inherited connective-tissue disorder marked by unusually flexible or unstable joints, long-term muscle or joint pain, and other variable features involving the skin, automatic body functions, or digestion.

### **Hypermobile Joints**

Joints that move farther than usual. This may cause no problems, or may lead to pain, repeated dislocations, sprains, fatigue, or joint instability.

### **Hypermobility**

Greater-than-usual movement of one or more joints. It may be a harmless trait, but when associated with pain, instability, dislocations, fatigue, or other features it may be part of a hypermobility spectrum disorder.

### **Hypermobility Spectrum Disorder**

A condition involving joint hypermobility together with symptoms such as pain, joint instability, soft-tissue injury, or fatigue, without meeting the criteria for another defined connective-tissue disorder.

### **Hypermobility Syndrome**

Excessive joint movement associated with pain, instability, fatigue, or other symptoms; evaluation distinguishes generalized joint hypermobility from a defined hypermobility spectrum disorder or hypermobile Ehlers-Danlos syndrome.

### **Hypermotility**

Abnormally rapid or excessive movement of an organ, especially the digestive tract. It may cause cramping, urgency, diarrhea, or poor absorption and can result from infection, medicines, inflammation, or other disorders.

### **Hypernatraemia**

An abnormally high level of sodium in the blood, usually because the body has too little water. It may cause thirst, weakness, confusion, seizures, or

---

coma.

### **Hypernatremia**

A high blood-sodium level, usually because the body has lost too much water or cannot obtain enough. It can cause intense thirst, weakness, confusion, seizures, or coma.

### **Hypernatremia Result**

A high blood sodium result, usually reflecting too little free water relative to body sodium; severity and rate of onset determine neurologic risk and the urgency and speed of correction.

### **Hyperopia**

Alternate name; see *Hypermetropia*.

### **Hyperornithinemia**

An elevated blood ornithine concentration caused by defects in ornithine metabolism or transport, including disorders that can produce retinal degeneration or neurologic disease.

### **Hyperosmolar**

Having more dissolved particles than another fluid. When blood becomes too concentrated, water moves out of cells and can cause dehydration, confusion, or other nervous-system problems.

### **Hyperosmolar Hyperglycemic State**

A life-threatening condition in which very high blood glucose causes severe dehydration and concentrated body fluids, usually with little or no ketone buildup. It may cause extreme thirst, weakness, confusion, seizures, or coma.

### **Hyperosmolar Non-Ketotic Coma**

A life-threatening complication of very high blood glucose, usually in type 2 diabetes, causing severe dehydration and sometimes confusion, seizures, or coma.

### **Hyperostosis**

Abnormal excessive formation or thickening of bone, which may be localized or widespread and may

occur in skeletal dysplasias, inflammation, endocrine disease, or medication effects.

### **Hyperoxaluria**

An excess of oxalate in the urine, which can combine with calcium to form kidney stones or calcium deposits. Severe inherited forms can progressively damage the kidneys.

### **Hyperoxia**

Exposure to an abnormally high oxygen concentration or tissue oxygen level, which can cause oxygen toxicity, absorption atelectasis, retinal injury in premature infants, or lung injury depending on dose and duration.

### **Hyperparakeratosis**

Thickening of the outer skin layer in which cells retain their nuclei instead of fully maturing. It may produce rough scale and can result from irritation, inflammation, infection, or an inherited skin disorder.

### **Hyperparathyroid Bone Disease**

Bone changes caused by prolonged high levels of parathyroid hormone. The hormone can increase bone breakdown, lowering bone strength and raising fracture risk; severe long-standing disease can cause osteitis fibrosa cystica.

### **Hyperparathyroidism**

A disorder in which the parathyroid glands release too much parathyroid hormone. Primary hyperparathyroidism usually begins in a gland and often raises blood calcium; secondary hyperparathyroidism is a response to problems such as chronic kidney disease or vitamin D deficiency and may have normal or low calcium. Tertiary hyperparathyroidism can develop after long-standing secondary disease.

### **Hyperpathia**

An unusually strong and sometimes delayed pain response to a painful stimulus. It may occur after nerve injury or with disorders that change how the nervous system processes pain.

---

### **Hyperphagia**

An unusually strong or excessive appetite. It can occur with diabetes, some genetic or brain disorders, certain medicines, or mental health conditions.

### **Hyperphosphatemia**

A high phosphate level in the blood, most often caused by kidney disease. It can disturb calcium balance, weaken bones, and contribute to calcium deposits in blood vessels or soft tissues.

### **Hyperpigmentation**

Darkening of an area of skin because the body makes or stores more pigment there. Causes include sun exposure, inflammation, injury, medicines, hormone disorders, and some skin or internal diseases.

### **Hyperpituitarism**

Excess production of one or more hormones from the pituitary gland, usually because of a benign pituitary tumor. The effects depend on the hormone involved and may affect growth, menstrual cycles, fertility, metabolism, or stress responses.

### **Hyperplasia**

An increase in the number of cells in a tissue or organ, which may make it larger. It can be a normal response or result from disease, hormones, or other stimuli.

### **Hyperplastic Polyp**

A usually noncancerous overgrowth of lining cells, often in the colon or rectum, whose cancer risk depends on size, number, and location.

### **Hyperpnea**

Breathing that is deeper, and sometimes faster, than normal, increasing air movement and occurring with exercise, illness, metabolic changes, or anxiety.

### **Hyperpnoea**

Breathing that is deeper or faster than usual to meet increased demand for oxygen or remove extra carbon dioxide. It may occur during exercise, fever, anxiety, or lung, heart, or metabolic illness.

### **Hyperpolarization**

An electrical change that makes a nerve or muscle cell less likely to send a signal. It occurs when the inside of the cell becomes more electrically negative after certain ion channels open.

### **Hyperprolactinaemia**

An abnormally elevated concentration of prolactin in the blood. Causes include medicines, pregnancy, hypothyroidism, pituitary tumors, and other disorders.

### **Hyperprolactinemia**

An abnormally high level of the hormone prolactin in the blood. Causes include pregnancy, breastfeeding, medicines, an underactive thyroid, kidney disease, or a pituitary tumor. It may disrupt periods, fertility, sexual function, or milk production.

### **Hyperproteinemia**

An abnormally high total blood-protein concentration, usually from dehydration, chronic inflammation, or a monoclonal immunoglobulin disorder.

### **Hyperpyrexia**

An extreme elevation of body temperature, generally above 41.5 °C, that can result from severe infection, heatstroke, malignant hyperthermia, drug toxicity, or other life-threatening conditions.

### **Hyperreactio Luteinalis**

Bilateral ovarian enlargement with multiple theca-lutein cysts caused by an exaggerated response to human chorionic gonadotropin, usually during pregnancy or with trophoblastic disease.

### **Hyperreflexia**

Exaggerated or overactive deep-tendon (stretch) reflexes. It is a neurologic sign, not a diagnosis, and often reflects impaired descending upper-motor-neuron control; it may occur with spasticity, clonus, or an extensor plantar response. Distribution and associated findings help distinguish central nervous-system disease from systemic or toxic causes.

---

## Hypersensitivity

An exaggerated or inappropriate immune response to an antigen, ranging from a mild reaction to anaphylaxis.

### Hypersensitivity Pneumonitis

Lung inflammation caused by repeated breathing of substances such as mold, bird proteins, or other workplace dusts. It may cause cough, breathlessness, fever, or fatigue and can lead to permanent lung scarring if exposure continues.

### Hypersensitivity Reactions

Inappropriate or excessive immune responses causing tissue damage or disease.

### Hypersensitivity Type I

An immediate allergic reaction in which immunoglobulin E antibodies activate mast cells after exposure to an allergen. Symptoms can include itching, hives, wheezing, vomiting, swelling, or life-threatening anaphylaxis.

### Hypersensitivity Type II

An immune reaction in which IgG or IgM antibodies attach to targets on cells or tissues and recruit other immune mechanisms that injure them. Examples include some drug reactions and autoimmune blood-cell destruction.

### Hypersensitivity Type III

An immune reaction in which clusters of antibodies and foreign or self proteins settle in tissues or blood vessels, causing inflammation and injury. It can affect the skin, joints, kidneys, or blood vessels.

### Hypersensitivity Type IV

A delayed immune reaction driven by T cells rather than antibodies. It usually develops 48 to 72 hours after exposure and can cause contact dermatitis, some drug rashes, or inflammation around an infection.

### Hypersensitivity Vasculitis

Inflammation of small blood vessels, usually in the

skin, often triggered by an infection, medicine, or immune reaction. It commonly causes raised purple spots or bruising on the legs.

### Hypersomnia

Being very sleepy during the day or sleeping much longer than usual even after enough sleep. It can be caused by sleep disorders, medicines, depression, substance use, or other illness; it is different from feeling tired without wanting to sleep.

### Hypersplenism

A condition in which an enlarged or overactive spleen removes blood cells too quickly. It may lower red cells, white cells, platelets, or all three and can cause anemia, infections, or easy bleeding.

### Hypertelorism

An unusually wide distance between the eyes because the bony eye sockets are farther apart than usual. It may be present from birth or occur as part of a genetic or developmental condition.

### Hypertension

Medical term; see *High blood pressure*.

### Hypertension and Pregnancy

High blood pressure before or during pregnancy ranges from chronic or gestational hypertension to preeclampsia and eclampsia. It can affect the brain, kidneys, liver, placenta, and fetal growth; blood-pressure monitoring, appropriate medicines, laboratory tests, and fetal surveillance are essential.

### Hypertension and the Kidney

High blood pressure can damage renal arteries and glomeruli, while kidney disease can activate salt-retaining and hormonal pathways that raise blood pressure. Their interaction can accelerate chronic kidney disease, so both blood pressure and kidney function require monitoring.

### Hypertension In Anesthesia

Abnormally high blood pressure during or around an operation. Pain, inadequate anesthesia, low oxy-

---

gen, high carbon dioxide, shivering, bladder fullness, or stopping certain medicines may contribute and increase the risk of bleeding, heart injury, or stroke.

### **Hypertension With Chronic Kidney Disease**

Alternate wording; see *Hypertension and the Kidney*.

### **Hypertensive**

Having blood pressure that remains above the accepted range for the person's age, health, and measurement setting, increasing risk of heart, brain, kidney, or vessel disease.

### **Hypertensive Cerebellar Hemorrhage**

Bleeding into the cerebellum caused by rupture of a vessel damaged by high blood pressure.

### **Hypertensive Crisis**

A dangerously high blood pressure that needs prompt assessment. If it is causing new organ injury, such as stroke, heart failure, kidney injury, or vision loss, it is an emergency requiring immediate hospital treatment.

### **Hypertensive Emergency**

A severe rise in blood pressure that is causing new injury to the brain, heart, kidneys, eyes, or major blood vessels. Symptoms may include chest pain, breathlessness, weakness, confusion, seizures, or vision loss and require immediate hospital care.

### **Hypertensive Encephalopathy**

Brain dysfunction caused by a sudden, severe rise in blood pressure. Headache, confusion, seizures, vision changes, or reduced consciousness may occur and require immediate blood-pressure treatment.

### **Hypertensive Heart Disease**

Changes or damage to the heart caused by long-term high blood pressure. The heart has to work harder to pump against the increased pressure, which can make its muscle thicken or stiffen and may eventually lead to heart failure or abnormal heart rhythms.

### **Hypertensive Intracerebral Hemorrhage**

Bleeding into brain tissue after a blood vessel weakened by high blood pressure breaks, potentially causing sudden headache, weakness, confusion, unconsciousness, or death.

### **Hypertensive Kidney Disease**

Kidney damage associated with long-term high blood pressure. It may lead to protein in the urine and gradually reduced kidney function; diabetes, other kidney diseases, and inherited risk can cause similar findings and must be considered.

#### **Alternate Names:**

Hypertensive Nephropathy

### **Hypertensive Nephrosclerosis**

Scarring and narrowing of small kidney blood vessels caused by long-term high blood pressure. It can gradually reduce kidney function and may cause protein in the urine; controlling blood pressure helps slow progression.

### **Hypertensive Retinopathy**

Damage to the blood vessels and light-sensitive tissue at the back of the eye caused by high blood pressure. It may include narrowed vessels, bleeding, swelling, or fluid leakage. Severe changes can threaten vision and may indicate dangerously high blood pressure.

### **Hypertensive Urgency**

A markedly high blood pressure without clear evidence of new organ injury. It still requires prompt medical review, especially when symptoms are present; treatment is adjusted carefully rather than lowered suddenly without supervision.

### **Hyperthermia**

Hyperthermia is an abnormally high body temperature caused by excessive heat production or impaired heat loss, without a change in the hypothalamic temperature set point. Causes include heat exposure, exertion, medicines, endocrine disorders, infection-related illness, and impaired sweating. Severe hyperthermia can cause confusion, seizures, organ injury, or death and requires urgent treatment.

---

## Hyperthermia For Treating Cancer

A cancer treatment that deliberately heats a tumor or part of the body, sometimes with radiation or medicines, to damage cancer cells or make other treatment work better.

## Hyperthermia In Anesthesia

An abnormally high temperature during or after anesthesia. Causes include infection, drug reactions, thyroid storm, serotonin syndrome, and malignant hyperthermia, a rare emergency with muscle rigidity, rising carbon dioxide, and dangerous temperature elevation.

## Hyperthermic Intraperitoneal Chemotherapy

Heated cancer medicine circulated through the abdominal cavity during surgery after visible tumor is removed. It is used for selected cancers that have spread across the abdominal lining and can cause infection, bleeding, organ injury, or other serious complications.

## Hyperthyroidism

A condition in which the thyroid makes too much thyroid hormone. It may cause weight loss, sweating, heat intolerance, shaking, anxiety, diarrhea, or a fast or irregular heartbeat. Common causes include Graves disease and overactive thyroid nodules.

### Alternate Names:

Thyrotoxicosis; Overactive Thyroid

## Hyperthyroxinemia

An abnormally high concentration of circulating thyroxine, caused by hyperthyroidism, thyroiditis, medication, binding-protein changes, or assay interference; symptoms and TSH clarify the cause.

## Hypertonia

Abnormally increased muscle tightness or resistance when a limb is moved. It may cause stiffness, abnormal posture, spasms, or limited movement and can result from brain, spinal-cord, or other nervous-system disorders.

## Hypertonic Saline

Saline containing more sodium than normal blood, commonly given through a vein for severe symptoms from dangerously low blood sodium or selected urgent conditions. Close monitoring is required because correcting sodium too quickly can cause serious brain injury.

## Hypertonic Saline Inhalation

Inhaling a salty mist through a nebulizer to draw water into airway mucus and make it easier to clear. It may help some people with cystic fibrosis or bronchiectasis, but can briefly narrow the airways and should be used as instructed.

## Hypertonic Solution

A solution with greater effective osmotic concentration than the fluid with which it is compared, causing water to move out of cells across a semipermeable membrane.

## Hypertrichosis

Excessive hair growth in any body area, in a localized or generalized pattern, that is not limited to the androgen-dependent male pattern seen in hirsutism. It may be present from birth or develop later because of inherited conditions, medicines, nutritional or metabolic disease, skin disorders, or rarely cancer. New, widespread, or rapidly progressive growth should be assessed.

## Hypertriglyceridemia

An abnormally high level of triglycerides, a type of fat, in the blood. Causes include diabetes, obesity, alcohol, some medicines, and inherited disorders. Very high levels increase the risk of acute pancreatitis and require medical management.

## Hypertrophic

Relating to hypertrophy, an enlargement of cells or tissue that increases the size of an organ or structure.

## Hypertrophic Cardiomyopathy

A disease in which the heart muscle, especially the left ventricle, becomes abnormally thick and stiff. It may obstruct blood flow, impair filling, cause



---

chest pain or breathlessness, and increase the risk of arrhythmia and sudden death.

**Alternate Names:**

HCM

**Hypertrophic Cardiomyopathy Genetics**

Many cases result from an inherited change in a gene for a heart-muscle protein, commonly MYH7 or MYBPC3. The condition can run in families, so genetic counseling and screening of close relatives may be recommended.

**Hypertrophic Cardiomyopathy Treatment**

Management may include medicines to relieve symptoms, treatment of arrhythmias or obstruction, an implantable defibrillator for selected patients, and surgery or ablation when indicated. Treatment is guided by symptoms, anatomy, risk, and specialist assessment.

**Hypertrophic Gastritis**

Thickening of the gastric mucosal folds caused by glandular overgrowth or inflammation, as in Ménétrier disease; protein loss, pain, nausea, and increased cancer risk may occur.

**Hypertrophic Myocardopathy**

Abnormal thickening of the heart muscle, often inherited. The thickened muscle may become stiff, obstruct blood leaving the heart, cause chest pain or breathlessness, or trigger dangerous abnormal rhythms.

**Hypertrophic Obstructive Cardiomyopathy**

A form of hypertrophic cardiomyopathy in which thickened heart muscle obstructs blood leaving the left ventricle. The obstruction may be present at rest or become more pronounced during exertion; see *Hypertrophic Cardiomyopathy*.

**Hypertrophic Osteoarthropathy**

A syndrome of digital clubbing, periosteal new-bone formation, and joint pain or swelling. It may be primary and inherited or secondary to conditions such as lung cancer, chronic lung infection, cyanotic

heart disease, or inflammatory disorders.

**Hypertrophic Pyloric Stenosis**

Thickening of the muscle at the outlet of the stomach, usually in young infants. It causes increasingly forceful vomiting after feeds, continued hunger, dehydration, and poor weight gain and is usually corrected with surgery.

**Hypertrophic Scar**

A raised, thick scar that stays within the boundaries of the original wound. It forms when the body makes too much scar tissue during healing, often after burns, surgery, or injury. Unlike a keloid, it does not grow beyond the wound edges and may become flatter over time.

**Hypertrophy**

Enlargement of a tissue or organ because its individual cells become larger. It may develop in response to increased workload, hormones, disease, or other stimulation.

**Hypertropia**

Vertical misalignment of the eyes in which one eye deviates upward relative to the other, causing double vision, eye strain, or abnormal head posture.

**Hyperuricemia**

An abnormally high concentration of uric acid in the blood, which may be asymptomatic or contribute to gout and uric-acid kidney stones.

**Hyperventilating**

Breathing faster or deeper than the body needs, lowering carbon dioxide in the blood. It may cause light-headedness, tingling, chest tightness, or fainting and can result from anxiety or serious illness.

**Hyperventilation**

Abnormally rapid, deep, or otherwise excessive breathing that removes more carbon dioxide than the body produces, often causing respiratory alkalosis. It may cause breathlessness, dizziness, tingling, chest tightness, muscle spasms, or fainting. Anxiety

---

or panic can trigger it, but new or severe hyperventilation may reflect heart or lung disease, infection, bleeding, metabolic illness, or another emergency.

### **Hyperventilation Syndrome**

Repeated overbreathing, often during stress or panic, that lowers carbon dioxide in the blood and causes breathlessness, chest tightness, dizziness, tingling, palpitations, or hand and foot spasms. Other heart and lung causes must be considered.

### **Hypervigilance**

A state of excessive alertness to possible threat, often accompanied by scanning, startle, irritability, and sleep difficulty in anxiety, trauma-related disorders, stimulant use, or medical illness.

### **Hyperviscosity Syndrome**

A set of symptoms and signs caused by blood that is abnormally thick, usually from very high levels of an immunoglobulin paraprotein or of red or white blood cells. It can complicate Waldenström macroglobulinemia, multiple myeloma, polycythemia, leukemia, or severe infection. Features include bleeding from the nose or gums, blurred or lost vision from retinal changes, headache, dizziness, confusion, stroke-like symptoms, or heart failure; severe cases are an emergency managed with plasma exchange while the underlying disease is treated.

### **Hypervitaminosis A**

Illness caused by too much vitamin A, usually from supplements or medicines. It may cause headache, nausea, dry skin, hair loss, liver injury, bone pain, or birth defects during pregnancy.

### **Hypervitaminosis D**

Illness caused by too much vitamin D, leading to high blood calcium. It may cause nausea, vomiting, thirst, frequent urination, weakness, kidney stones, confusion, or kidney damage.

### **Hypervolaemia**

An excess of blood or body fluid. It may cause rapid weight gain, swelling, breathlessness, raised neck veins, or high blood pressure, especially with

heart, kidney, or liver disease.

### **Hypervolemia**

Too much fluid in the bloodstream or body tissues. It may cause rapid weight gain, swelling, breathlessness, raised neck veins, or high blood pressure and can occur with heart, kidney, or liver disease.

### **Hypha**

A branching tubular filament that forms the basic structural unit of most molds and many fungi; networks of hyphae form a mycelium and may invade tissue in fungal disease.

### **Hyphema**

Blood collecting in the front part of the eye, usually after an injury. It may cause pain, blurred vision, light sensitivity, or increased eye pressure and requires prompt eye assessment to protect vision.

### **Hypnagogic**

Occurring during the transition from wakefulness to sleep. Hypnagogic sensations or hallucinations can be benign, but frequent or distressing episodes may occur with narcolepsy, sleep deprivation, medicines, or other sleep and neurologic disorders.

### **Hypnagogic Hallucinations**

Vivid sights, sounds, sensations, or feelings that occur while a person is falling asleep. They are often harmless but may be linked to sleep deprivation, narcolepsy, medicines, or other sleep disorders.

### **Hypnic Headache**

A rare primary headache that repeatedly awakens a person during sleep, usually in later life and often at a similar time each night.

### **Hypnogram**

A chart showing when a person is awake and in each sleep stage during a recording. It is commonly made from a sleep study and helps show sleep timing, awakenings, and stage patterns.

#### **Alternate Names:**

Sleep Architecture Diagram; Polysomnographic

---

Hypnogram

### Hypnosis

A state of focused attention and deep relaxation in which a person may respond more readily to guided suggestions. It is not sleep or loss of control and should be used by a trained professional when offered as part of care.

### Hypnotherapy

Hypnotherapy uses focused attention and guided suggestion as an adjunct to care for selected problems such as pain, anxiety, or unwanted habits.

### Hypnotic

A medicine used mainly to help a person fall asleep or stay asleep. Some hypnotics can cause next-day drowsiness, falls, dependence, or slowed breathing, especially when combined with alcohol or other sedatives. Not all sedatives are used as sleep medicines.

### Hypoactive Sexual Desire Disorder

Persistently low or absent sexual interest that causes personal distress or relationship difficulty. Desire can be affected by medicines, hormones, illness, pain, mood, stress, trauma, or relationship factors, so assessment looks for the underlying cause.

### Hypoadrenalism

Insufficient adrenal-cortex hormone production, especially cortisol and sometimes aldosterone; primary disease can cause fatigue, weight loss, low blood pressure, hyperpigmentation, and adrenal crisis.

### Hypoalbuminemia

An abnormally low blood albumin concentration caused by reduced production, inflammation, kidney or gastrointestinal loss, or severe malnutrition; it can contribute to edema and alter drug binding.

### Hypoalphalipoproteinemias

Disorders with low HDL-related apolipoprotein levels, which may impair reverse cholesterol transport and increase cardiovascular risk depending on the cause and overall lipid profile.

### Hypoblast

The inner cell layer of the early embryonic disc that contributes to the yolk-sac lining and helps establish the embryonic body axes during gastrulation.

### Hypocalcaemia

Alternate British spelling; see *Hypocalcemia*.

### Hypocalcemia

An abnormally low level of calcium in the blood (total calcium below 8.5 mg/dL or 2.1 mmol/L). Calcium is essential for nerve transmission, muscle contraction, and cardiac electrical stability. Low levels cause neuromuscular irritability, presenting with perioral numbness, fingertip tingling, painful muscle cramps, tetany, and hallmark physical signs: Chvostek's sign (facial twitching on tapping the facial nerve) and Trousseau's sign (carpal spasm during blood pressure cuff inflation). Severe cases cause seizures, airway spasm (laryngospasm), and a prolonged QT interval on ECG. Common causes include hypoparathyroidism (especially after thyroid surgery), severe vitamin D deficiency, chronic kidney disease, low magnesium, and pancreatitis. Severe cases require urgent intravenous calcium gluconate.

#### Alternate Names:

Low Blood Calcium; Calcium Deficiency; Hypocalcaemia

### Hypocapnia

An abnormally low level of carbon dioxide in the blood, usually because a person is breathing too fast or deeply. Causes include anxiety, pain, low oxygen, lung disease, pregnancy, or excessive ventilator support.

### Hypochloremic Alkalosis

A chloride-depletion form of metabolic alkalosis, commonly associated with vomiting, gastric suction, or diuretic use. Low chloride reflects loss of hydrogen and chloride ions and helps guide assessment of the cause and response to treatment.

### Hypochlorite

A chemical form of chlorine used in disinfectants and bleach. It can kill many germs, but swallowing it

---

or getting concentrated solutions in the eyes, lungs, or skin can cause serious injury.

### **Hypocholesterolemia**

An unusually low blood cholesterol concentration caused by malnutrition, malabsorption, hyperthyroidism, chronic illness, liver disease, or inherited lipid disorders.

### **Hypochondria**

Older term; see *Illness Anxiety Disorder* or *Health Anxiety*.

### **Hypochondroplasia**

An inherited disorder of bone growth that causes short stature and shortened arms and legs, usually less severely than achondroplasia. People may have broad hands, increased inward curvature of the lower back, or limited elbow movement, while facial appearance and intelligence are often typical. It is usually caused by a change in the *FGFR3* gene.

#### **Alternate Names:**

HCH; Mild Achondroplasia Variant

### **Hypochromia**

Reduced hemoglobin color in red blood cells, seen as increased central pallor on a blood smear and commonly associated with iron deficiency or thalassemia.

### **Hypochromic Anaemia**

Hypochromic anaemia is anemia in which red blood cells contain too little hemoglobin and appear unusually pale, commonly because of iron deficiency or thalassemia.

### **Hypochromic Anemia**

Anemia in which red cells contain too little hemoglobin and appear pale, most often because of iron deficiency, thalassemia, chronic disease, or sideroblastic disorders.

### **Hypocitraturia**

Low urinary citrate excretion, which reduces inhibition of calcium-crystal formation and increases risk

of calcium kidney stones; potassium citrate may be used when clinically appropriate.

### **Hypocupremia**

An abnormally low copper concentration in blood or tissues, caused by inadequate intake, malabsorption, excess zinc, or inherited disorders, and potentially producing anemia, neutropenia, or neurologic disease.

### **Hypodermic Needle**

A hollow needle designed to inject medicines or withdraw fluid through the skin, with gauge and length selected according to the substance, route, body site, and patient.

### **Hypodermis**

The layer of fat and loose tissue beneath the skin. It cushions the body, stores energy, helps conserve heat, and contains blood vessels, nerves, and lymph channels.

### **Hypodontia**

A condition present from birth in which one or more permanent teeth fail to develop. It most often affects wisdom teeth, upper side incisors, or second premolars and may affect chewing, appearance, or tooth alignment.

### **Hypoesthesia**

Reduced sensitivity to touch, temperature, pain, or other sensations. It may result from nerve compression, neuropathy, spinal disease, stroke, injury, or other neurologic conditions; sudden or spreading numbness requires prompt assessment.

### **Hypogalactia**

Insufficient breast-milk production for an infant's needs, caused by ineffective milk removal, hormonal or breast disorders, medications, illness, or other feeding difficulties.

### **Hypogammaglobulinemia**

Abnormally low levels of antibodies in the blood, which can lead to repeated infections, especially of

---

the ears, sinuses, lungs, or bloodstream. It may be inherited or develop because of another disease or treatment.

### **Hypogastric Region**

The lower central area of the abdomen, just above the pubic bone. It overlies the bladder and parts of the bowel and reproductive organs; pain or a lump there may arise from urinary, intestinal, or pelvic disease.

### **Hypoglossal Canal**

A paired bony opening situated in the occipital bone above the anterolateral margin of the foramen magnum, directed laterally and forward to transmit the hypoglossal nerve (cranial nerve XII) and a meningeal branch of the ascending pharyngeal artery out of the posterior cranial fossa.

#### **Alternate Names:**

Anterior Condylar Foramen

### **Hypoglossal Nerve**

Cranial nerve XII, which supplies the muscles that move the tongue; a lesion causes tongue weakness, wasting, and deviation toward the affected side on protrusion.

### **Hypoglossal Nerve Injury**

Damage to the nerve that controls tongue movement. It may follow skull-base injury, surgery, a tumor, or infection and can cause tongue weakness, wasting, or deviation when the tongue is extended.

### **Hypoglossal Nerve Palsy**

Weakness or paralysis of the hypoglossal nerve (CN XII), leading to ipsilateral tongue weakness, atrophy, and fasciculations. Causes include tumors at the skull base or in the hypoglossal canal, surgical injury, or central lesions. Unilateral palsy causes the tongue to deviate toward the affected side on protrusion.

### **Hypoglycaemia**

An abnormally low blood-glucose level. It may cause sweating, shaking, hunger, palpitations, confusion, seizures, or loss of consciousness and is especially associated with diabetes medicines, missed meals,

or heavy alcohol use.

### **Hypoglycemia**

An abnormally low blood-glucose concentration that may cause sweating, tremor, palpitations, hunger, confusion, seizures, or loss of consciousness. Causes include glucose-lowering medicines, missed meals, alcohol, and critical illness. Treatment depends on the person's ability to swallow and the cause; severe or recurrent episodes require urgent assessment.

### **Hypoglycemia In Diabetes**

Blood glucose that falls too low in a person with diabetes, usually below 70 mg/dL. It may cause sweating, shaking, hunger, confusion, or fainting; severe episodes prevent self-treatment and require another person's help.

### **Hypoglycemia Unawareness**

Failure to feel the usual warning signs of low blood glucose, such as shaking, sweating, or a racing heart-beat. It can occur after repeated low readings or long-standing diabetes and increases the risk of severe hypoglycemia.

### **Hypoglycemic**

Relating to an abnormally low blood glucose level, or describing a medicine that lowers blood glucose.

### **Hypogonadism**

Reduced function of the testes or ovaries, causing low sex-hormone levels and impaired production of sperm or eggs. It may lead to delayed puberty, absent periods, infertility, low sexual desire, hot flashes, reduced muscle or bone strength, or other symptoms.

### **Hypogonadism In Men**

Low testosterone caused by reduced testicular function or by a problem in the brain signals that control the testes. It may cause low sexual desire, erectile problems, infertility, low energy, reduced muscle, or weak bones.



---

### Hypogonadism In Women

Reduced ovarian hormone production, which may cause absent or irregular periods, hot flashes, vaginal dryness, infertility, low sexual desire, or loss of bone strength. Causes include ovarian failure, pituitary disease, some medicines, severe illness, or surgery.

### Hypogonadotropic Hypogonadism

Low production of sex hormones because the brain or pituitary gland does not send enough stimulating signals to the testes or ovaries. It may cause delayed puberty, absent periods, infertility, low sexual desire, or reduced testosterone. Causes include genetic conditions, pituitary disease, obesity, medicines, or severe illness.

### Hypohidrosis

Reduced or absent sweating. It can make it difficult for the body to cool itself and increase the risk of overheating. Causes include nerve disorders, skin damage, medicines, dehydration, and inherited conditions.

### Hypokalaemia

British spelling of Hypokalemia; see Hypokalemia.

### Hypokalemia

An abnormally low level of potassium in the blood, often caused by vomiting, diarrhea, diuretics, hormone disorders, or movement of potassium into cells. It may cause weakness, cramps, constipation, or dangerous heart-rhythm changes.

### Hypokalemic Periodic Paralysis

An inherited disorder causing repeated attacks of muscle weakness or paralysis when blood potassium falls. Attacks often begin after rest following exercise or after a high-carbohydrate meal, while strength may be normal between attacks.

### Hypokinesia

Reduced or slowed movement. It may cause small steps, less facial expression, reduced arm swing, or small handwriting and commonly occurs in Parkinson disease or other disorders affecting movement control.

### Hypolipidemic Agents

Medicines that lower cholesterol or triglycerides, including statins, ezetimibe, fibrates, bile-acid sequestrants, niacin, and selected injectable therapies.

### Hypolipoproteinemia

An abnormally low concentration of one or more blood lipoproteins, caused by inherited disorders, malabsorption, malnutrition, liver disease, or other systemic illness.

### Hypomagnesaemia

An abnormally low level of magnesium in the blood. It may cause tremor, muscle cramps, weakness, seizures, or abnormal heart rhythms and can result from diarrhea, poor intake, alcohol use, or certain medicines.

### Hypomagnesemia

A low level of magnesium in the blood, often caused by diarrhea, poor nutrition, alcohol use, kidney loss, or some medicines. It may cause tremor, cramps, weakness, seizures, or abnormal heart rhythms.

### Hypomania

A distinct period of unusually elevated, expansive, or irritable mood with increased energy or activity lasting at least four consecutive days. It may involve reduced need for sleep, rapid speech, racing thoughts, inflated self-esteem, or impulsive behavior; the change is observable by others but is not severe enough to cause marked impairment or require hospitalization and has no psychotic features.

### Hypomanic Episode

Alternate name; see *Hypomania*.

### Hypomelanosis Of Ito

A rare condition causing swirled or streaky areas of lighter skin, often from birth. Some affected children also have seizures, developmental differences, eye problems, or abnormalities of the bones or nervous system.



---

## Hypomethylation

An abnormally low amount of methyl marking on DNA or other molecules. It can change which genes are active and may affect chromosome stability, development, inherited disorders, or cancer cells.

## Hypomnesia

Reduced ability to remember or recall information, which may occur with aging, neurologic disease, psychiatric conditions, medications, or other causes.

## Hyponatraemia

Alternate British spelling; see *Hyponatremia*.

## Hyponatremia

An abnormally low level of sodium in the blood, often caused by excess water, fluid loss, medicines, heart or liver disease, kidney problems, or hormone disorders. It may cause nausea, headache, confusion, seizures, or coma; severe or rapidly developing cases are emergencies.

## Hypoparathyroidism

A disorder in which too little parathyroid hormone causes low blood calcium and high phosphate. It most often follows neck surgery but can be inherited or autoimmune. Tingling, muscle cramps, spasms, seizures, or abnormal heart rhythms may occur and need treatment.

## Hypopharyngeal Neoplasm

An abnormal growth in the lower part of the throat beside the voice box and esophagus. It may cause persistent sore throat, painful swallowing, difficulty swallowing, ear pain, or a neck lump and may be cancerous.

## Hypopharynx

The lower part of the pharynx behind and beside the larynx, extending from the oropharynx to the esophagus. It directs food toward the esophagus during swallowing.

## Hypophosphataemia

Alternate British spelling; see *Hypophosphatemia*.

## Hypophosphatasia (HPP)

A rare inherited disorder caused by deficient tissue-nonspecific alkaline phosphatase, usually because of *ALPL* variants. Poor bone and tooth mineralization can cause rickets or osteomalacia, fractures, bone pain, premature loss of teeth, or respiratory problems in severe infantile disease.

## Hypophosphatemia

An abnormally low level of phosphate in the blood (below 2.5 mg/dL or 0.8 mmol/L). Because phosphate is essential for generating cellular energy (ATP) and oxygen delivery (2,3-DPG), its deficiency starves tissues of energy. It is classically triggered by refeeding syndrome (rapid carbohydrate refeeding in malnourished or alcoholic patients driving phosphate into cells), diabetic ketoacidosis recovery, severe malnutrition, or renal phosphate wasting. Symptoms include profound muscle weakness, respiratory failure from diaphragmatic weakness, acute muscle breakdown (rhabdomyolysis), hemolytic anemia, confusion, and seizures. Treatment consists of oral or intravenous phosphate replacement.

### Alternate Names:

Low Blood Phosphate; Hypophosphataemia; Phosphate Deficiency

## Hypophosphatemic Rickets

Rickets caused by low phosphate availability, often from renal phosphate wasting, leading to defective mineralization, bowed legs, bone pain, growth impairment, and dental abnormalities.

## Hypophysectomy

Surgical removal of all or part of the pituitary gland, usually performed through the nose and sphenoid sinus (transsphenoidal hypophysectomy) or rarely through an open craniotomy to treat pituitary tumors, relieve optic nerve compression, or control hormone hypersecretion.

### Alternate Names:

Pituitary Resection; Transsphenoidal Adenomectomy

## Hypophysis

An endocrine gland at the base of the brain that

---

regulates growth, thyroid and adrenal function, reproduction, lactation, and water balance through its hormones.

**Alternate Names:**

Pituitary Gland; Master Endocrine Gland

**Hypophysitis**

Inflammation of the pituitary gland that can cause headache, visual symptoms, and deficiency or excess of pituitary hormones; it may be autoimmune, related to pregnancy, or triggered by immune-checkpoint therapy.

**Hypopigmentation**

Reduced skin pigmentation producing paler areas than the surrounding skin, caused by decreased melanin, reduced melanocyte function or number, inflammation, injury, infection, or inherited disorders.

**Hypopituitarism**

Reduced production of one or more hormones from the pituitary gland because of disease, injury, surgery, radiation, or a nearby brain disorder. It may affect growth, thyroid or adrenal function, periods, fertility, milk production, or water balance.

**Alternate Names:**

Pituitary Insufficiency; Panhypopituitarism

**Hypoplasia**

Incomplete or underdevelopment of a tissue, organ, or body structure, resulting in reduced size or functional capacity. It may be congenital, developmental, genetic, or acquired and ranges from a mild anatomical variant to severe organ dysfunction. Assessment considers the timing of onset, associated malformations, tissue function, and whether the process is isolated or syndromic.

**Hypoplastic Left Heart Syndrome**

A serious birth defect in which the left side of the heart and the aorta are too small to pump enough blood to the body. Newborns need urgent specialist treatment, often involving staged surgery or transplantation.

**Hypoplastic Left Heart Syndrome (HLHS)**

Alternate name; see *Hypoplastic Left Heart Syndrome*.

**Hypoploidy**

A chromosome number below the normal diploid complement, caused by loss of one or more chromosomes; in tumor cytogenetics it may have prognostic significance.

**Hypopnea**

A period of unusually shallow or reduced breathing, often during sleep. Repeated episodes can lower blood oxygen, disturb sleep, and contribute to obstructive sleep apnea or other breathing disorders.

**Hypoproliferative Anemia**

Anemia caused by inadequate red-cell production, usually with an inappropriately low reticulocyte response. Causes include iron deficiency, inflammation, kidney disease, endocrine or nutritional disorders, and bone-marrow failure or infiltration.

**Hypoproteinaemia**

Hypoproteinaemia is an abnormally low level of proteins in the blood, which may result from malnutrition, liver disease, kidney protein loss, or intestinal loss.

**Hypoproteinemia**

An abnormally low amount of protein in the blood, often involving albumin, which can cause swelling and reflect poor nutrition, liver disease, kidney loss, or illness.

**Hypopyon**

A visible white or yellow level of inflammatory material in the front chamber of the eye. It usually indicates severe eye inflammation or infection and may occur with pain, redness, light sensitivity, and reduced vision.

**Hyporeninemic Hypoaldosteronism**

Reduced aldosterone production associated with

---

inappropriately low renin activity, often occurring with diabetic kidney disease or other chronic kidney disorders. Hyperkalemia and a mild normal-anion-gap metabolic acidosis are common findings.

### **Hyposmia**

A reduced ability to smell, without complete loss of smell. It can follow a cold or COVID-19, nasal blockage, head injury, medicines, aging, or a neurologic disorder.

### **Hypospadias**

A birth condition in which the opening of the urethra is on the underside of the penis rather than at its tip. The penis may curve, and the foreskin may be incomplete; severity depends on the opening's location and other abnormalities.

### **Hypospadias Repair**

Surgery reconstructs the urethral opening and, when needed, straightens the penis in a child born with hypospadias; timing and technique depend on anatomy and surgeon assessment.

### **Hypotension**

Abnormally low blood pressure that may reduce blood flow to the brain and other organs. It can result from bleeding, dehydration, heart problems, severe infection, allergy, or medicines and may cause dizziness, fainting, confusion, or shock.

### **Hypotension In Anesthesia**

Low blood pressure during or soon after anesthesia. It may result from anesthetic medicines, blood loss, dehydration, heart problems, or an allergic reaction; the anesthesia team monitors and treats it according to the cause and severity.

### **Hypotensive**

Having or causing abnormally low blood pressure; symptoms may include dizziness, fainting, weakness, confusion, or shock depending on the cause and severity.

### **Hypothalamic Amenorrhea**

Absence of menstrual periods because the hypothalamus reduces its normal release of reproductive signals. It is often linked to low energy availability, excessive exercise, stress, illness, or weight loss and may cause infertility and bone loss.

### **Hypothalamic Dysfunction**

A problem affecting the brain region that helps control hormones, temperature, appetite, thirst, sleep, stress, and reproduction. Causes include injury, tumors, inflammation, infection, surgery, radiation, or inherited disease.

### **Hypothalamic Hamartoma**

A congenital, noncancerous mass of disorganized hypothalamic tissue that can cause gelastic seizures, precocious puberty, behavioral symptoms, or cognitive decline.

### **Hypothalamic Hormone**

A releasing or inhibiting hormone produced by the hypothalamus that regulates pituitary secretion, including signals controlling thyroid, adrenal, reproductive, growth, and lactation axes.

### **Hypothalamic Neoplasm**

An abnormal growth in or near the hypothalamus, a brain region controlling hormones, temperature, appetite, sleep, and water balance. Effects may include vision changes, hormone problems, headaches, seizures, or altered growth.

### **Hypothalamic Tumor**

Alternate name; see *Hypothalamic Neoplasm*.

### **Hypothalamic-Pituitary-Adrenal Axis**

The hormone system linking the brain's hypothalamus and pituitary gland with the adrenal glands. It controls cortisol release, helping regulate the body's response to stress, illness, blood pressure, metabolism, and immune activity.

#### **Alternate Names:**

HPA Axis; Neuroendocrine Stress Axis

---

### **Hypothalamic-Pituitary-Adrenal Axis Physiology**

The brain signals the pituitary gland, which signals the adrenal glands to release cortisol. Cortisol normally follows a daily pattern and feeds back to reduce further signals when enough is present.

### **Hypothalamic-Pituitary-Gonadal Axis Physiology**

The hormone system linking the hypothalamus, pituitary gland, and ovaries or testes. Signals from the hypothalamus prompt the pituitary to release hormones that control ovulation and menstrual cycles, or testosterone production and sperm development.

### **Hypothalamic-Pituitary-Thyroid Axis Physiology**

The hormone system linking the hypothalamus, pituitary gland, and thyroid. The hypothalamus signals the pituitary to release thyroid-stimulating hormone, which prompts the thyroid to make hormones that regulate metabolism, temperature, and energy use.

### **Hypothalamus**

A small region at the base of the brain that helps control automatic body functions and hormone release through the pituitary gland. It regulates body temperature, hunger, thirst, sleep, stress, and parts of emotional behavior.

### **Hypothermia**

A dangerously low internal body temperature, usually below 35 degrees Celsius, caused by cold exposure or impaired heat production. It may cause shivering, confusion, drowsiness, slow breathing, or an abnormal heart rhythm. Severe cases require urgent medical rewarming.

#### **Alternate Names:**

Accidental Hypothermia; Low Core Body Temperature

### **Hypothermia In Anesthesia**

An abnormally low body temperature during or after anesthesia. Anesthetic medicines reduce the body's ability to control heat and may allow heat loss during surgery; warming devices and tempera-

ture monitoring help prevent complications.

### **Hypothyroidism**

A clinical endocrine syndrome resulting from deficient production or action of thyroid hormones (thyroxine [ $T_4$ ] and triiodothyronine [ $T_3$ ]), causing a generalized slowing of basal metabolism. Primary hypothyroidism (over 95% of cases, characterized by elevated serum TSH and low free  $T_4$ ) most commonly results from autoimmune destruction (Hashimoto thyroiditis), surgical thyroidectomy, radioactive iodine ablation, or severe iodine deficiency; central hypothyroidism stems from pituitary or hypothalamic failure. Clinical hallmarks include profound fatigue, cold intolerance, weight gain, constipation, sinus bradycardia, dry skin, non-pitting dermal edema (myxedema), delayed relaxation of deep tendon reflexes, and cognitive slowing.

#### **Alternate Names:**

Underactive Thyroid; Hypothyroid State; Thyroid Deficiency

### **Hypotonia**

Decreased muscle tone presenting as a floppy or poorly supported posture. It may result from muscle, peripheral nerve, neuromuscular-junction, spinal-cord, brain, genetic, or metabolic disorders and requires evaluation when persistent or associated with weakness or developmental delay.

### **Hypotonia-Hypomentia-Hypogonadism-Obesity Syndrome (HHHO)**

Historical descriptive name for Prader-Willi syndrome, a genetic disorder involving infantile hypotonia, developmental or intellectual disability, hypogonadism, and later hyperphagia with obesity; see *Prader-Willi Syndrome*.

### **Hypotonic Solution**

A liquid containing fewer dissolved particles than the fluid inside cells. Water tends to move into cells when they are exposed to it, so medical use requires careful selection and monitoring.

### **Hypotony Maculopathy**

A vision-threatening complication of persistent low

---

intraocular pressure characterized by chorioretinal wrinkling and macular folds, optic disc edema, and decreased visual acuity.

### **Hypotrichosis**

Abnormally sparse or absent hair caused by inherited follicle disorders, autoimmune disease, endocrine or nutritional problems, medications, or scarring alopecia.

### **Hypotrophy**

Underdevelopment or reduced size of a tissue or organ that is present but smaller or less developed than usual. It may begin before birth or result from poor nutrition, reduced use, impaired blood flow, or disease and may affect function.

### **Hypotropia**

Vertical eye misalignment in which one eye deviates downward relative to the other, potentially causing diplopia, abnormal head posture, or suppression of vision.

### **Hypoventilation**

Breathing too slowly or shallowly to remove enough carbon dioxide. It can lower oxygen levels and cause sleepiness, headache, confusion, or respiratory failure; causes include medicines, lung disease, obesity, and muscle or nerve disorders.

### **Hypoventilation Syndromes**

Disorders in which alveolar ventilation is inadequate, causing elevated carbon dioxide, often with low oxygen. Causes include obesity hypoventilation, neuromuscular weakness, chest-wall disease, medicines, and impaired respiratory control; evaluation includes sleep and blood-gas assessment.

### **Hypovolaemia**

Hypovolaemia is an abnormally low volume of circulating blood or body fluid, commonly caused by bleeding, dehydration, or major fluid loss.

### **Hypovolemia**

Hypovolemia is reduced effective circulating blood

volume, usually from bleeding, fluid loss, or inadequate intake; it can cause tachycardia, hypotension, poor perfusion, kidney injury, and hypovolaemic shock.

### **Hypovolemic Shock**

Circulatory failure from loss of intravascular volume, reducing venous return, cardiac output, and tissue perfusion. Hemorrhage is the most dangerous cause; other causes include vomiting, diarrhea, burns, third-spacing, and severe dehydration. Tachycardia, narrow pulse pressure, cool clammy skin, delayed capillary refill, oliguria, confusion, and elevated lactate may occur.

### **Hypoxaemia**

An abnormally low oxygen level in arterial blood. It can result from lung disease, breathing problems, poor blood flow, or other conditions and may cause shortness of breath, confusion, bluish skin, or organ injury.

### **Hypoxanthine**

A naturally occurring substance made when the body breaks down certain genetic-material building blocks. It is normally reused or changed into other compounds during purine metabolism.

### **Hypoxemia**

An abnormally low oxygen level in the blood. It may result from lung or heart disease, blocked airways, shallow breathing, anemia, poor circulation, or high altitude and can cause breathlessness, confusion, or bluish skin.

### **Hypoxia**

Inadequate oxygen availability at the tissue level. It may result from low arterial oxygen, anemia, reduced blood flow, or impaired cellular oxygen use.

### **Hypoxia And Hypoxemia**

Hypoxemia is low oxygen in arterial blood, while hypoxia is inadequate oxygen at the tissue level. Causes include lung disease, airway obstruction, heart disease, anemia, shock, or high altitude; severe breathlessness, cyanosis, confusion, or collapse is



---

urgent.

### **Hypoxia-Inducible Factor**

A heterodimeric master transcription factor complex (composed of an oxygen-regulated alpha subunit and a constitutive beta subunit) that under cellular hypoxic conditions escapes von Hippel-Lindau-mediated proteasomal degradation to upregulate target genes encoding erythropoietin, VEGF, and glycolytic enzymes.

**Alternate Names:**

HIF; HIF-1

### **Hypoxic Cell Injury**

Damage to cells caused by too little oxygen. It may result from poor blood flow, low blood oxygen, anemia, or poisoning that prevents oxygen use; the outcome depends on how severe and prolonged the oxygen shortage is.

### **Hypoxic Ischemic Encephalopathy**

Brain dysfunction caused by insufficient oxygen and blood flow, usually around birth in newborns but also after cardiac arrest or severe respiratory failure. It can cause seizures, altered consciousness, and long-term neurologic disability.

### **Hypoxic Pulmonary Vasoconstriction**

Narrowing of small blood vessels in an area of the lung that receives too little oxygen. This normally redirects blood toward better-ventilated lung areas, but widespread or prolonged narrowing can strain the heart.

### **Hypoxic-Ischemic Brain Injury**

Brain damage caused by too little oxygen or blood flow. It may follow cardiac arrest, suffocation, shock, severe breathing failure, or very low blood glucose and can cause seizures, coma, memory problems, movement problems, or disability.

### **Hypsarrhythmia**

A very abnormal, disorganized pattern on an electroencephalogram, with large slow waves and spikes arising from many areas of the brain. It is strongly associated with infantile spasms, a serious seizure

disorder in babies, and requires prompt specialist evaluation and treatment.

**Alternate Names:**

Hypsarrhythmic EEG Pattern; Chaotic Dysrhythmic EEG

### **Hysterectomy**

Surgical removal of the uterus. The cervix, fallopian tubes, and ovaries may or may not also be removed, depending on the procedure and clinical indication. Hysterectomy permanently ends menstruation and the ability to carry a pregnancy, but it does not cause immediate menopause when functioning ovaries remain. If the cervix remains, cervical cancer screening may still be needed. Indications include cancer, fibroids, severe bleeding, adenomyosis, endometriosis, chronic pelvic pain, and uterine prolapse.

### **Hysterectomy And Surgical Menopause**

Hysterectomy removes the uterus and ends menstruation and the ability to carry a pregnancy. Surgical menopause occurs when both ovaries are removed or ovarian function otherwise stops abruptly; hysterectomy alone does not necessarily cause menopause if the ovaries remain.

### **Hysterosalpingo-Contrast Sonography**

An ultrasound test using fluid or foam contrast to assess the uterine cavity and whether the fallopian tubes are open. It is often used during fertility evaluation without X-ray radiation.

**Alternate Names:**

HyCoSy; Fallopian Tubal Contrast Ultrasound

### **Hysterosalpingo-Contrast-Sonography**

An ultrasound-based technique for evaluating the uterine cavity and fallopian tube patency. Sterile saline or contrast medium is injected through the cervix while transvaginal ultrasound is performed.

### **Hysterosalpingogram**

An X-ray procedure used to evaluate the uterine cavity and fallopian tube patency. A radiopaque contrast medium is injected through the cervix, and serial X-ray images are taken as the contrast fills the



---

uterine cavity and spills from the fallopian tubes.

and examination of thyroid tissue help determine its nature and treatment.

### **Hysterosalpingography**

An X-ray test in which contrast liquid is placed through the cervix to outline the inside of the uterus and fallopian tubes. It can show blockage, an unusual uterine shape, fibroids, polyps, or scar tissue during infertility evaluation. Cramping, bleeding, infection, or contrast reaction can occur.

### **Hysteroscope**

A thin camera-equipped instrument passed through the cervix to inspect the uterine cavity and, when designed for operative use, remove polyps, treat bleeding, or perform other procedures.

### **Hysteroscopic Metroplasty**

A procedure that uses a small camera and instruments passed through the cervix to divide a wall of tissue separating part of the uterus. It may improve the uterine cavity in selected people with fertility problems or repeated pregnancy loss.

### **Hysteroscopic Myomectomy**

A procedure that removes selected fibroids bulging into the inside of the uterus. A camera and instruments pass through the cervix, avoiding an abdominal incision; the procedure may reduce heavy bleeding and improve fertility.

### **Hysteroscopy**

A procedure that passes a small camera through the cervix to inspect the inside of the uterus and, when needed, obtain tissue or treat conditions such as polyps, fibroids, or intrauterine adhesions.

### **Hysterotomy**

A surgical incision into the uterus performed during selected obstetric or gynecologic procedures. It may be used to deliver a baby or reach the uterus, with risks including bleeding and infection.

### **Hürthle Cell Tumour**

A growth arising from hormone-making cells in the thyroid. It may be benign or cancerous; ultrasound

---

# I

## Iatrogenic

Describing an illness, injury, or other harm caused unintentionally by medical examination, treatment, a medicine, a procedure, or a healthcare device. Examples include side effects, procedure-related bleeding, infection, nerve damage, or radiation injury. The term does not imply negligence.

## Iatrogenic Cushing Syndrome

Cortisol excess caused by prescribed or nonprescribed glucocorticoid exposure, including systemic, injected, inhaled, topical, or potent local preparations. Features may include weight gain, hypertension, diabetes, bruising, muscle weakness, infection risk, and adrenal suppression.

## Iatrogenic Disease

Illness or injury caused unintentionally by a medical examination, treatment, medicine, procedure, or healthcare device. It may be temporary or permanent and does not by itself imply negligence.

## Iatrogenic Injury

Harm caused unintentionally by a medical examination, treatment, procedure, medicine, or healthcare device. It may be temporary or permanent and can include infection, bleeding, medication effects, nerve damage, or injury during surgery.

## Iatrogenic Pneumothorax

A pneumothorax inadvertently provoked by an invasive medical or surgical diagnostic procedure (such as central venous catheterization, transthoracic needle lung biopsy, thoracentesis, transbronchial biopsy, or mechanical positive-pressure ventilation).

## Ibandronic Acid

A bisphosphonate medicine used in some settings to reduce bone loss and complications from cancer affecting bone. It can rarely damage the jaw or cause unusual thigh fractures.

## IBD

Inflammatory bowel disease, including Crohn's disease and ulcerative colitis, causing long-term digestive-tract inflammation, diarrhea, abdominal pain, bleeding, weight loss, or fatigue.

### Alternate Names:

Inflammatory bowel disease

## Ibritumomab Tiuxetan

An antibody linked to a radioactive substance and used in selected B-cell lymphomas. The antibody attaches to CD20-positive cells and delivers radiation; low blood counts, infection, and delayed marrow problems are important risks.

## IBS

Abbreviation; see *Irritable Bowel Syndrome*.

## IBS-D

IBS with diarrhea causes recurrent abdominal pain associated with loose stools without structural disease explaining the symptoms. Management may include diet changes, soluble fiber, gut-directed medicines, behavioral therapy, and evaluation for alarm features or alternative diagnoses.

### Alternate Names:

IBS-D; Diarrhea-Predominant Irritable Bowel Syndrome

---

## Ibuprofen

A medicine that reduces pain, fever, and inflammation. It can irritate or bleed the stomach, harm the kidneys, cause fluid retention, or increase heart risks in some people.

## Ibuprofen Overdose

Toxic exposure to more ibuprofen than recommended, which can cause nausea, vomiting, abdominal pain, gastrointestinal bleeding, kidney injury, metabolic acidosis, drowsiness, seizures, or coma.

## Ibuprofen Toxicity

Acute overdose of the nonsteroidal anti-inflammatory drug ibuprofen, manifesting clinically in mild cases with epigastric pain, nausea, and vomiting, and in severe ingestions with high anion gap metabolic acidosis, acute kidney injury, hypotension, hypocalcemia, and coma.

## ICD

In this dictionary, an implantable cardioverter-defibrillator: a device placed under the skin that monitors heart rhythm and uses pacing or an electrical shock to stop dangerous rhythms. In other contexts, ICD may mean International Classification of Diseases.

### Alternate Names:

Implantable cardioverter defibrillator

## ICD-10

Alternate form; see *International Classification of Diseases*.

## ICD-11

The eleventh revision of the World Health Organization's International Classification of Diseases, in effect internationally from 2022 and designed for digital recording, reporting, and analysis of diseases, injuries, and causes of death.

## ICD-O Neoplasm Morphology Classification

A coding system used by cancer registries and pathology services to record what type of tumor

is present and whether it is benign or cancerous. It does not replace the diagnosis, location, or stage of the cancer.

## ICD Shock Therapy

Treatment delivered by an implanted defibrillator to stop dangerous heart rhythms. The device may use painless pacing or an electrical shock, depending on the rhythm.

## I-Cell Disease

A rare inherited disorder in which cells cannot deliver certain digestive enzymes to their waste-processing compartments. Waste then accumulates in cells, causing severe developmental delay, bone abnormalities, unusual facial features, and progressive disease in several organs.

### Alternate Names:

I-Cell

## Ice Pack

A device that applies cold to a body area to help relieve pain or swelling. Direct or prolonged contact can injure the skin.

## Ice Pick Headache

A sudden, severe, stabbing head pain that usually lasts seconds. Attacks may recur in different head areas without warning; new, persistent, or unusually severe pain can have other causes.

### Alternate Names:

Primary stabbing headache

## ICF Framework

A World Health Organization framework for describing how a health condition affects body functions, daily activities, participation in life, and the person's environment. It supports a broader view of disability than diagnosis alone.

## Ichthyosis

A group of conditions that cause very dry, rough, scaly skin. Some forms are inherited and may be present from birth; others are linked to allergies or another illness.

---

## Ichthyosis Vulgaris

A common inherited skin condition causing dry, fine scaling, especially on the arms and legs. It often improves with age and can be managed with regular moisturizers and prescribed scale-reducing treatments.

## Ictal State

The physiological and behavioral state during which an active epileptic seizure event is occurring in the brain, demarcated from the preceding pre-ictal (aura) and succeeding post-ictal recovery phases.

### Alternate Names:

Ictus

## Icteric

Describing yellowing of the skin, whites of the eyes, or other tissues caused by too much bilirubin in the blood.

## Icteric Index

The icteric index is a laboratory estimate of serum yellow discoloration related mainly to bilirubin and can signal jaundice or interfere with testing.

## Icterus

Yellowing of the skin or whites of the eyes caused by excess bilirubin. It may result from liver disease, blocked bile flow, or rapid breakdown of red blood cells.

## Ictus

A sudden neurologic event, usually a stroke or seizure. The word does not identify the exact illness, so the symptoms, examination, tests, and cause must be determined.

## ICU-Acquired Weakness

Generalized muscle weakness that develops during critical illness or an intensive-care stay. It may result from critical illness myopathy, critical illness polyneuropathy, or both and can delay movement and removal from a ventilator.

## Idarubicin

Idarubicin is an anthracycline chemotherapy drug that intercalates DNA and inhibits topoisomerase II, used mainly in acute myeloid leukaemia; myelosuppression and cumulative cardiotoxicity are important risks.

## IDC

Abbreviation; see *Invasive Ductal Carcinoma*.

## Idea Of Reference

A belief that ordinary events, comments, or media messages have a special personal meaning, often seen in psychotic or mood disorders when held with strong conviction.

## Ideas Of Reference

The belief that ordinary events, comments, media messages, or other people's actions have a special personal meaning. If the belief is held with strong conviction despite evidence, it may occur in a psychotic or mood disorder.

## Ideation

The formation or content of thoughts, especially thoughts related to a particular behavior or symptom. Clinical examples include suicidal ideation and homicidal ideation, which require assessment of intent, plan, means, and immediate safety.

## Identical Twins

Two individuals formed when one fertilized egg separates into two embryos. They usually have nearly identical DNA, but differences can develop through mutations, epigenetic changes, environment, and chance.

## Idiopathic

Describing a condition for which no specific cause has been identified after appropriate evaluation. It does not mean that the condition is imaginary, unimportant, or untreatable; it means that its cause is currently unknown.

---

### Idiopathic Anaphylaxis

Repeated anaphylaxis, a severe allergic reaction, for which no specific trigger is found after evaluation. Possible hidden triggers and mast-cell disorders may also be considered.

### Idiopathic Facial Paralysis

Alternate name; see *Bell's Palsy*.

### Idiopathic Guttate Hypomelanosis

Harmless, small, flat white spots caused by reduced skin pigment, usually on the shins or forearms of older adults. They are not contagious and usually do not need treatment.

### Idiopathic Hypercalciuria

Excess calcium in the urine without an identified cause. It can increase the risk of kidney stones and, less often, bone loss.

### Idiopathic Hypersomnia

A sleep disorder that causes extreme daytime sleepiness, long or unrefreshing sleep, and difficulty waking up, even when a person has enough time to sleep. Other causes must be ruled out.

### Idiopathic Interstitial Pneumonias

A group of interstitial lung diseases with no identified cause, characterized by varying degrees of lung inflammation and fibrosis; diagnosis requires clinical, radiologic, and sometimes pathologic assessment.

### Idiopathic Intracranial Hypertension

Increased pressure around the brain without a tumor or other clear cause. It can cause headaches, a pulsing sound in the ears, and vision problems and may damage sight if untreated.

### Idiopathic Myelofibrosis

Alternate name; see *Primary Myelofibrosis*.

### Idiopathic Pulmonary Arterial Hypertension

Group 1 pulmonary arterial hypertension in which no associated disease, drug or toxin, heritable cause,

or other explanation is found after appropriate evaluation. It increases resistance in the pulmonary arteries and strains the right ventricle, causing exertional breathlessness, fatigue, chest discomfort, dizziness, fainting, or swelling. Diagnosis is confirmed with right-heart catheterization and specialist assessment.

### Idiopathic Pulmonary Fibrosis

A long-term lung disease of unknown cause in which scar tissue gradually builds up in the lungs. It causes increasing shortness of breath and a dry cough.

#### Alternate Names:

IPF

### Idiopathic Thrombocytopenic Purpura

An older name for immune thrombocytopenia, a disorder in which the immune system destroys platelets and may cause easy bruising, tiny skin spots, nosebleeds, heavy menstrual bleeding, or other bleeding.

### Idiopathic Toe Walking

Alternate name; see *Toe Walking*.

### Idiosyncrasy

An unusual individual reaction to a medicine or exposure that is not explained by the usual pharmacologic effect.

### Idiosyncratic Reaction

An uncommon, unexpected, and often genetically determined adverse drug reaction that is unrelated to the drug's known pharmacology and usually dose-independent (a Type B reaction).

### Idioventricular Rhythm

A slow heart rhythm that starts in the lower chambers of the heart instead of the normal pacemaker. It may occur when the usual electrical signal is blocked or when the heart is severely ill.

### Id Reaction (Autoeczematization)

A secondary, usually itchy eczematous eruption that appears at sites distant from an infection or inflammatory skin disorder, often on the hands, feet, or



---

trunk. Treating the primary problem and controlling inflammation usually improves it.

### Ifosfamide

Ifosfamide is an alkylating chemotherapy drug that forms DNA cross-links and is used for selected sarcomas and other cancers; haemorrhagic cystitis, encephalopathy, infertility, and myelosuppression require monitoring and mesna protection.

### iFR

Alternate name; see *Instantaneous Wave-Free Ratio*.

### Ig

Abbreviation for immunoglobulin, an antibody protein produced by B cells and plasma cells; major classes include IgG, IgA, IgM, IgE, and IgD.

### IgA

A type of antibody found mainly in body fluids and linings, including saliva, tears, breast milk, and the digestive and breathing passages. It helps protect these surfaces from infection.

### IgA Nephropathy

A kidney disease caused by deposits of immunoglobulin A in the kidney filters, potentially leading to blood in urine, protein loss, high blood pressure, or kidney failure.

#### Alternate Names:

Berger's Disease

### IgA Pemphigus

A rare autoimmune blistering disorder in which IgA antibodies target epidermal cell-adhesion proteins. It causes superficial pustules or blisters, often in annular or circinate patterns, and diagnosis relies on biopsy and direct immunofluorescence.

### IgA Vasculitis

A disease in which the immune system inflames small blood vessels, usually after an infection. It can cause purple spots on the legs, joint pain, abdominal pain, and kidney problems, especially in children.

#### Alternate Names:

Henoch-Schonlein Purpura

### IgD

A less common type of antibody found mainly on the surface of certain immune cells. It helps those cells recognize signals, but its full role in human disease is not yet clear.

### IgE

A type of antibody involved in allergies and defense against some parasites. When it reacts to an allergen, it can trigger symptoms such as sneezing, hives, asthma, or anaphylaxis.

### IgG

The most common type of antibody in the blood. It helps neutralize germs and toxins, supports immune protection after infection or vaccination, and can cross the placenta to protect a developing baby.

### IgG4-Related Disease

An uncommon immune disorder that causes inflammation and scar-like growths in one or more organs. It may affect the pancreas, salivary glands, eyes, kidneys, lungs, or blood vessels; diagnosis usually requires imaging, blood tests, and sometimes a biopsy.

### IgG4-Related Disease Arteritis

Inflammation of a large or medium-sized artery as part of IgG4-related disease. It may thicken or weaken the artery and can cause narrowing, an aneurysm, pain, or reduced blood flow.

### IgG4-Related Ophthalmic Disease

An immune disorder that causes inflammation and firm swelling around the eye, including the tear glands, eyelids, or eye muscles. It may cause bulging, eyelid swelling, double vision, or reduced vision and may affect other organs.

### IgG and Complement Deposition

A laboratory finding in which immune proteins collect where the outer and deeper layers of skin meet. It can occur in lupus and other conditions, but the

---

pattern and symptoms are needed to interpret its meaning.

### **IgM**

A large type of antibody that is often made early during an infection. It helps clump germs and activate other parts of the immune system; a blood test may suggest recent exposure but must be interpreted with other results.

### **IGRT**

Image-guided radiation therapy uses medical imaging, such as CT, MRI, ultrasound, or X-ray, to position the patient and target radiation accurately before or during treatment. It helps deliver radiation to a tumor while reducing exposure to nearby healthy tissue.

### **IIH**

Abbreviation; see *Idiopathic Intracranial Hypertension*.

### **Ileal Atresia**

A condition present at birth in which part of the ileum, the last section of the small intestine, is closed or missing. It causes bowel blockage in a newborn, with a swollen abdomen, green vomiting, and failure to pass the first stool.

### **Ileal Conduit**

A surgically created urine passage made from a short piece of the ileum, the last part of the small intestine. It carries urine from the ureters to an opening in the abdominal wall and an external bag.

### **Ileal Fistula**

An abnormal connection involving the ileum and another organ, bowel segment, or skin surface, which may cause intestinal contents to leak and lead to infection, malnutrition, or fluid loss.

### **Ileal Hemorrhage**

Bleeding from the ileum, the distal portion of the small intestine, caused by ulceration, inflammation, vascular lesions, tumors, trauma, or other disease.

### **Ileal Neoplasm**

An abnormal growth in the ileum, the last part of the small intestine. It may be noncancerous or cancerous and can cause pain, bleeding, blockage, anemia, or no symptoms.

### **Ileal Obstruction**

Partial or complete blockage of the ileum, the final part of the small intestine. Adhesions, hernias, inflammatory bowel disease, tumors, or twisting may cause cramping pain, vomiting, swelling, and inability to pass stool or gas.

### **Ileal Pouch-Anal Anastomosis**

An operation that removes the colon and rectum and creates a pouch from the end of the small intestine. The pouch is connected to the anus so bowel movements can continue through the usual route.

### **Ileal Stenosis**

Narrowing of the ileum that may impair passage of intestinal contents and result from Crohn disease, inflammation, scarring, ischemia, tumor, or surgery.

### **Ileal Ulcer**

An ileal ulcer is a break in the lining of the ileum caused by inflammation, infection, medicines, ischemia, trauma, or another intestinal disorder.

### **Ileitis**

Inflammation of the ileum, the last part of the small intestine. Causes include Crohn disease, infection, reduced blood flow, medicines, and other disorders; symptoms may include lower-right abdominal pain, diarrhea, fever, or bleeding.

### **Ileocaecal**

Relating to the junction between the ileum, the last part of the small intestine, and the cecum, the first part of the large intestine.

### **Ileocecal Junction**

The point where the ileum, the last part of the small intestine, joins the cecum, the first part of the large intestine.

---

## Ileocecal Valve

A flap-like structure where the last part of the small intestine joins the first part of the large intestine. It helps control passage of intestinal contents and limits backflow from the colon.

## Ileostomy

An operation that brings the end of the small intestine through an opening in the abdomen so stool drains into an external pouch. Output is often liquid, so adequate fluids, salt replacement, and skin care are important.

## Ileum

The final and longest part of the small intestine. It absorbs vitamin B12, bile salts, and remaining nutrients before food passes into the large intestine.

## Ileus

Failure of coordinated intestinal propulsion without a mechanical blockage, usually from postoperative bowel inhibition, inflammation, metabolic disturbance, or medication effect. It causes distension, nausea, vomiting, and reduced passage of flatus or stool.

## Iliac

Relating to the ilium or the iliac region, including the iliac bones and iliac blood vessels.

## Iliac Artery

One of the large arteries that carries blood to the pelvis and lower limb. The common iliac arteries divide into internal branches for the pelvic organs and external branches that continue into the legs.

## Iliac Crest

The iliac crest is the curved upper border of the ilium and a landmark and attachment site for abdominal, back, and pelvic muscles.

## Iliac Crest Bone Graft

The surgical harvesting of cancellous and cortical bone from the anterior or posterior iliac crest of the pelvis to provide an osteoconductive, osteoinductive,

and osteogenic autologous graft for orthopaedic non-union fracture repair or spinal fusion.

### Alternate Names:

ICBG

## Iliac Vein

An iliac vein drains blood from the pelvis and lower limb toward the inferior vena cava; thrombosis can cause swelling and pulmonary embolism.

## Iliopsoas Muscles

The psoas major and iliacus muscles working together to bend the hip. They help lift the thigh, stabilize the spine, and maintain posture, and may cause groin or back pain when strained.

### Alternate Names:

Iliopsoas

## Iliopsoas Tendinitis

Overuse-related irritation of the iliopsoas tendon, causing pain in the groin or front of the hip that may worsen with hip flexion, running, or rising from a chair.

## Iliotibial Band

A strong band of tissue running along the outer side of the thigh from the hip to the knee. It helps stabilize the hip and knee during movement.

## Iliotibial Band Friction Syndrome

An overuse injury causing pain on the outside of the knee, and sometimes the hip, when the iliotibial band becomes irritated. It is common in runners and cyclists.

## Iliotibial Band Syndrome

Pain on the outer side of the knee caused by irritation of the iliotibial band, especially during repeated activities such as running or cycling.

### Alternate Names:

ITBS; IT Band Friction Syndrome; Runner's Knee

## Ilium

The broad, upper part of each hip bone. It joins the sacrum at the back, helps form the hip socket, and

---

provides attachment for muscles of the abdomen, back, pelvis, and thigh. Its upper edge is the iliac crest.

### **Illness Anxiety Disorder**

Persistent fear of having or developing a serious illness despite few or no physical symptoms. The person may repeatedly check the body, seek reassurance, search for health information, or avoid medical care; symptoms last at least six months.

### **Illusion**

A misperception or misinterpretation of a real external stimulus, such as mistaking a shadow for a person. Illusions may occur with visual impairment, delirium, seizures, intoxication, or severe psychiatric illness and differ from hallucinations, which occur without an external stimulus.

### **Iloprost**

A prostacyclin analogue that causes vasodilation and inhibits platelet aggregation; inhaled or infused formulations are used for selected forms of pulmonary hypertension and severe peripheral ischemia.

### **Image-Guided Ablation**

Destruction of selected abnormal tissue with heat, cold, electricity, or chemicals while ultrasound, CT, MRI, or another imaging method guides and monitors the treatment. It may treat tumors, abnormal heart rhythms, or other disease.

### **Image-Guided Biopsy**

Removal of a tissue sample while ultrasound, CT, MRI, mammography, or fluoroscopy guides the needle or instrument to the target and helps avoid critical structures.

### **Image-Guided Drainage**

Placement of a needle or catheter while imaging guides its path to remove pus, blood, bile, urine, or other trapped fluid. It may treat an abscess, infected collection, or symptomatic cyst and can cause bleeding or infection.

### **Image-Guided Therapy**

A procedure in which ultrasound, CT, X-ray, MRI, or another imaging method guides a needle, catheter, or treatment device to a specific area. This can improve accuracy while limiting injury to nearby tissue.

### **Imaging**

The production of pictures of internal body structures using methods such as radiography, ultrasound, computed tomography, magnetic resonance imaging, or nuclear medicine.

### **Imaging And Radiology**

Imaging and radiology use methods such as X-ray, ultrasound, CT, and MRI to create pictures that help diagnose or guide treatment of disease.

### **Imaging-Guided Biopsy**

Alternate name; see *Image-Guided Biopsy*.

### **Imaging Technique**

An imaging technique is a method for producing pictures of body structures or processes, such as ultrasound, CT, MRI, or nuclear imaging.

### **Imatinib Mesylate**

The mesylate salt of imatinib, a tyrosine-kinase inhibitor used to treat BCR-ABL-positive chronic myeloid leukemia, gastrointestinal stromal tumors with KIT alterations, and other selected cancers.

### **Imerslund-Gräsbeck Syndrome**

A rare inherited disorder in which the small intestine cannot absorb vitamin B12 properly. It can cause low B12 levels, anemia, tiredness, numbness, balance problems, or other nerve damage despite adequate dietary intake.

### **Imidazoles**

A group of antifungal medicines, including clotrimazole, miconazole, and ketoconazole, used to treat fungal infections of the skin, mouth, or other body areas.

---

### **Iminoacetic Acid Scan**

A nuclear medicine scan that uses a small amount of radioactive medicine to show how bile moves through the liver, gallbladder, and bile ducts. It is also called a HIDA scan.

### **Imipenem**

A strong antibiotic, usually given with cilastatin, used for serious bacterial infections. It can cause allergic reactions, stomach problems, seizures, or antibiotic resistance.

### **Imipramine**

A medicine used for depression and some bladder problems. It can cause dry mouth, constipation, dizziness, abnormal heart rhythms, and dangerous effects in overdose.

### **Imipramine Overdose**

Toxic ingestion of imipramine, a tricyclic antidepressant, which can cause anticholinergic effects, seizures, hypotension, ventricular arrhythmias, coma, and life-threatening cardiac toxicity.

### **Imiquimod**

Imiquimod is a topical immune-response modifier that activates Toll-like receptor 7 and induces cytokines; it treats selected actinic keratoses, superficial basal-cell carcinomas, and external genital warts, often causing local inflammation.

### **Immature Ovarian Teratoma**

A rare cancer of the ovary made from immature cells that can develop into different body tissues. It occurs most often in teenagers or young adults, and its grade and spread guide treatment.

### **Immediate Hypersensitivity**

A rapid allergic reaction, usually involving IgE antibodies and mast cells. It can cause hives, itching, swelling, wheezing, vomiting, or anaphylaxis within minutes of exposure.

### **Immersion Foot**

A painful cold injury caused by prolonged exposure

of the feet to cold, wet conditions without freezing, leading to numbness, swelling, and skin changes.

### **Alternate Names:**

Trench Foot

### **Immobilization**

Restriction of movement of a body part or the whole patient, using rest, splints, casts, braces, traction, or other methods. It may protect an injury but can cause stiffness, pressure injury, muscle loss, venous thromboembolism, and other complications if prolonged.

### **Immobilize**

To restrict movement of a body part, often with a splint or cast, to support healing.

### **Alternate Names:**

Immobilization

### **Immotile Cilia Syndrome**

A genetically heterogeneous autosomal recessive ciliopathy caused by structural defects in ciliary axonemal dynein arms, leading to impaired mucociliary clearance, recurrent sinopulmonary infections, bronchiectasis, male infertility, and situs inversus in the subgroup known as Kartagener syndrome.

### **Alternate Names:**

Primary Ciliary Dyskinesia; PCD

### **Immune**

Having an immune response that recognizes and helps protect against a particular antigen, infection, or toxin; immunity may be innate or acquired.

### **Immune And Infectious Diseases**

Conditions involving abnormal immune defenses or infections caused by bacteria, viruses, fungi, or parasites. Examples include autoimmune disease, allergy, weakened immunity, and infections that are unusually severe or recurrent.

### **Immune Checkpoint**

A control signal that helps prevent immune cells from becoming too active or damaging healthy tissue. Some cancer medicines block these signals to

---

help immune cells attack tumors, but this can cause inflammation in normal organs.

### **Immune Checkpoint Inhibitor**

A cancer medicine that blocks an immune-checkpoint signal, helping immune cells recognize and attack cancer cells. It can also cause inflammation in normal organs, called an immune-related adverse event.

### **Immune Complex**

A cluster formed when an antibody attaches to a foreign or abnormal substance. If these clusters are not cleared, they can collect in tissues and cause inflammation, blood-vessel damage, or kidney disease.

### **Immune Disorder**

A condition caused by an immune system that is overactive, underactive, misdirected, or otherwise abnormal. Major groups include autoimmune disease, immunodeficiency, allergy and hypersensitivity, immune dysregulation, and some inflammatory disorders.

### **Immune Evasion**

The ability of an infection, tumor, or abnormal cell to avoid being recognized or destroyed by the immune system. It may hide identifying markers, block immune signals, or resist immune-cell attack.

### **Immune Hemolytic Anemia**

Anemia caused when antibodies or other immune proteins destroy red blood cells too quickly. It may cause tiredness, jaundice, dark urine, rapid heart-beat, breathlessness, or an enlarged spleen and may follow illness, medicines, or transfusion.

### **Immune-Mediated Necrotizing Myopathy**

An uncommon autoimmune muscle disease that causes worsening weakness, especially in the hips and shoulders, and often markedly raises muscle enzymes. Some cases are linked to statins or specific antibodies; diagnosis may require blood tests, imaging, and muscle testing.

### **Immune Privileged Sites**

Specialized anatomical tissues and compartments (including the brain, anterior chamber of the eye, testes, and pregnant uterus) that tolerate the introduction of foreign antigens without eliciting an inflammatory immune response, maintained by physical blood-tissue barriers, immunosuppressive cytokines, and Fas ligand expression.

### **Immune Reconstitution Inflammatory Syndrome**

A paradoxical inflammatory reaction and clinical deterioration occurring shortly after the initiation of combination antiretroviral therapy (ART) in HIV patients, caused by rapid recovery of pathogen-specific cellular immunity triggering an exaggerated immune response against occult or partially treated opportunistic pathogens (e.g., *Mycobacterium tuberculosis*, *Cryptococcus neoformans*).

#### **Alternate Names:**

IRIS; Immune Restoration Disease

### **Immune-Related Adverse Event**

An inflammatory or autoimmune toxicity caused by immune checkpoint blockade or another immune-activating cancer treatment. It may affect skin, bowel, liver, endocrine organs, lungs, heart, kidneys, or nervous system and often requires treatment-specific immunosuppression.

### **Immune Response**

The body's reaction to an infection, foreign substance, or abnormal cell. It includes rapid defenses such as inflammation and slower, targeted responses involving antibodies and immune cells that remember previous exposures.

### **Immune Status**

The current strength and activity of a person's immune defenses, including protection from previous infection or vaccination and effects of disease, medicines, nutrition, or cancer treatment. It helps estimate infection risk and vaccine response.

### **Immune System**

The body's network of cells, tissues, organs, and



---

proteins that recognizes and responds to infections, abnormal cells, and harmful substances. It includes rapid innate defenses and learned adaptive defenses; overactivity can cause allergy or autoimmunity, while weakness increases infection risk.

### **Immune Thrombocytopenia**

An autoimmune disorder in which the immune system destroys platelets or reduces their production, resulting in an unusually low platelet count. Platelets help stop bleeding, so immune thrombocytopenia may cause easy bruising, tiny red or purple skin spots (petechiae), nosebleeds, bleeding from the gums, heavy menstrual bleeding, or other bleeding; some people have no symptoms. It may be short-term or long-term and may occur on its own or in association with another illness or medicine.

### **Immune Thrombocytopenia in Pregnancy**

An autoimmune disorder in which antibodies destroy platelets, causing a low platelet count and sometimes bruising or bleeding. Pregnancy may complicate diagnosis and delivery planning; treatment and platelet targets depend on bleeding, counts, timing, and planned anesthesia.

### **Immune Thrombocytopenic Purpura**

An older name for immune thrombocytopenia, in which the immune system destroys platelets. It may cause bruising, pinpoint skin spots, nosebleeds, heavy menstrual bleeding, or no symptoms.

### **Immune Tolerance**

The ability of the immune system to avoid attacking the body's own tissues while still responding to infections. Failure of tolerance can contribute to autoimmune disease; excessive tolerance can allow some cancers or infections to persist.

### **Immunity**

The body's ability to recognize and defend against infections or other foreign substances. Protection may be present from birth, develop after infection or vaccination, or be provided temporarily by transferred antibodies.

### **Immunization**

The process of inducing protection against an infection, usually through vaccination. Recommended vaccines and schedules depend on age, previous doses, pregnancy, immune status, medical conditions, travel, exposure risk, and current public-health guidance.

### **Immunizations For People With Diabetes**

Vaccination is especially important in diabetes because infections may be more severe; recommendations depend on age, health status, vaccines received, and local guidance.

### **Immunoassay**

A laboratory test that uses the specific binding between an antibody and its target to detect or measure substances such as hormones, medicines, proteins, antigens, or other antibodies in a sample.

### **Immunoassays**

Laboratory tests that use specific binding between antibodies and target substances to detect or measure them. Examples include ELISA, chemiluminescent tests, and some rapid tests; results must be interpreted with the clinical situation.

### **Immunoblast Count**

An immunoblast count measures activated lymphocytes, usually in a tissue or blood sample, and may help evaluate immune responses or lymphoid disease.

### **Immunoblotting**

Immunoblotting, or Western blotting, separates proteins by electrophoresis and detects a target with specific antibodies; band pattern and intensity are interpreted with controls and clinical context.

### **Immunochemistry**

The study or laboratory measurement of biological molecules using specific antigen-antibody reactions, including immunoassays and immunostaining.

---

## Immunocompetence

The ability of the immune system to recognize harmful or foreign substances and produce an effective, appropriately controlled response. Reduced immunocompetence can increase infection risk and weaken responses to vaccines.

## Immunocompetent

Having an immune system capable of mounting an appropriate response to common infectious or foreign antigens.

## Immunocompromised

Having weakened or impaired immune defenses because of disease, medication, cancer treatment, transplantation, or another cause, increasing susceptibility to some infections and complications.

## Immunocompromised Host

A person whose immune defenses are weakened by an inherited disorder, infection, cancer, organ failure, malnutrition, or immune-suppressing treatment. Such a person may develop unusual, severe, or repeated infections and may have fewer typical symptoms.

## Immunocompromised Pneumonia

Pulmonary parenchymal infection occurring in patients with congenital or acquired immunodeficiencies (such as HIV, chemotherapy-induced neutropenia, or organ transplantation). Frequently caused by opportunistic pathogens including *Pneumocystis jirovecii*, fungi, and cytomegalovirus.

### Alternate Names:

Opportunistic Pneumonia; Immunodeficient Pulmonary Infection; Immunocompromised Host Pneumonia

## Immunoconjugate

A therapeutic or diagnostic molecule in which an antibody or other targeting protein is chemically linked to a drug, toxin, enzyme, or imaging agent.

## Immunocytochemistry

A laboratory method using labeled antibodies to

identify specific proteins or antigens within cells or tissue sections.

## Immunodeficiency

A condition in which part of the immune system is missing, weakened, or does not work properly. It may be inherited or caused by illness, medicines, cancer treatment, poor nutrition, or infection and can lead to repeated or severe infections.

## Immunodeficiency And Infections

A weakened immune system increases the risk of infections that are unusually severe, repeated, persistent, or caused by organisms that rarely affect healthy people. Causes include inherited disorders, HIV, cancer treatment, transplantation, and immune-suppressing medicines.

## Immunodeficiency Diseases

Alternate name; see *Immunodeficiency*.

## Immunodeficiency Disorders

Alternate name; see *Immunodeficiency*.

## Immuno electrophoresis

A laboratory method that separates proteins by electrophoresis and identifies them with specific antibodies, often to study immunoglobulins.

## Immuno electrophoresis – Blood

A laboratory technique that separates blood proteins by electrophoresis and identifies them with specific antibodies, used historically to evaluate abnormal immunoglobulins and monoclonal proteins.

## Immunofixation Blood Test

A blood test that identifies and types monoclonal immunoglobulins, helping evaluate monoclonal gammopathy, multiple myeloma, and related plasma-cell disorders.

## Immunofluorescence

A laboratory method that uses fluorescently labeled antibodies to show specific proteins, germs, or immune deposits in cells, tissues, or body fluids when

---

viewed with a special microscope.

### **Immunofluorescence Technique**

An immunofluorescence technique uses fluorescently labelled antibodies to locate specific proteins, organisms, or immune deposits in a sample.

### **Immunogenetics**

The study of how genes influence immune-system development, function, and disease. It includes inherited susceptibility to autoimmune disease, differences in immune responses, and genetic matching for transplantation.

### **Immunogenicity**

The ability of an antigen, vaccine, or other substance to provoke an immune response, including antibody or cellular responses.

### **Immunoglobulin**

A protein made by immune cells that recognizes a particular foreign substance, germ, or abnormal cell. Antibodies can block harmful targets, mark them for removal, or activate other immune defenses.

#### **Alternate Names:**

Antibody

### **Immunoglobulin A**

An antibody found mainly in mucus, saliva, tears, breast milk, and intestinal fluid. It protects surfaces exposed to the outside world and helps prevent germs from attaching and entering tissues.

### **Immunoglobulin A Deficiency**

An immune deficiency in which IgA levels are markedly reduced or absent, while other immunoglobulin classes may be normal. Some people have recurrent respiratory or gastrointestinal infections, allergies, or autoimmune disease, while others have few symptoms.

### **Immunoglobulin D**

A type of antibody found in very small amounts in the blood and on the surface of some B cells. Its exact role is not fully understood, but it may help

regulate immune responses in the airways and other tissues.

### **Immunoglobulin E**

An antibody involved in many immediate allergic reactions and defense against some parasites. It activates mast cells, which release chemicals causing itching, swelling, mucus, wheezing, or other allergy symptoms.

#### **Alternate Names:**

IgE

### **Immunoglobulin G**

The most common antibody in blood and tissue fluid, helping protect against many infections. It can remain after infection or vaccination, cross the placenta during pregnancy, and sometimes cause harmful immune reactions.

### **Immunoglobulin G4-Related Disease**

A systemic fibroinflammatory disorder characterized by tumefactive lesions, dense lymphoplasmacytic infiltrates enriched with IgG4-positive plasma cells, storiform fibrosis, obliterative phlebitis, and elevated serum IgG4 concentrations, affecting organs such as the pancreas, salivary and lacrimal glands, biliary tree, and retroperitoneum.

#### **Alternate Names:**

IgG4-RD; IgG4-Related Systemic Disease

### **Immunoglobulin G Deficiency**

Reduced IgG production or function that can impair antibody-mediated defense and lead to recurrent sinopulmonary or other infections. Evaluation usually considers IgG subclasses, vaccine responses, and associated immune abnormalities.

### **Immunoglobulin Isotype**

The class of an antibody, determined by its constant region. The main isotypes are IgG, IgA, IgM, IgD, and IgE; each has different roles and locations in immune defense.

### **Immunoglobulin M**

A large antibody usually produced early after a

---

new infection or other immune challenge. It helps activate immune defenses and can be measured in blood, but a high result does not by itself prove a recent infection.

### **Immunoglobulin M Deficiency**

An uncommon antibody deficiency characterized by persistently low IgM with otherwise variable immunoglobulin levels. It may be associated with recurrent infections, allergy, or autoimmune disease.

### **Immunoglobulin Replacement Therapy**

Treatment that supplies pooled antibodies through a vein or under the skin when a person's immune system makes too few effective antibodies. It can reduce recurrent infections in selected inherited or acquired antibody deficiencies; headache, fever, and infusion reactions may occur.

### **Immunochemolysis**

Destruction of red blood cells by an immune reaction involving antibodies, complement, or both. It may occur in autoimmune hemolytic anemia, incompatible transfusion reactions, or other antibody-mediated disorders.

### **Immunohistochemistry**

A laboratory method that uses specially marked antibodies to find particular proteins in a tissue sample. It helps identify tumor type, infection, or immune disease and can guide treatment choices.

### **Immunologic Adjuvant**

A substance added to some vaccines to strengthen or guide the immune response. It may improve protection but can also increase temporary pain, swelling, fever, or other vaccine reactions.

### **Immunologically Privileged Site**

A body area where immune responses are partly limited, helping protect delicate tissues from inflammation while still allowing important defense against infection.

### **Immunological Memory**

The ability of the immune system to respond faster and more strongly when it meets a substance again. Long-lived memory B cells and T cells formed after infection or vaccination provide this protection.

### **Immunological Synapse**

The organized contact point where an immune cell meets a target or another immune cell. Signals gathered at this contact help decide whether the immune cell should become active.

### **Immunological Tolerance**

The reduced or absent immune response to a specific antigen, including the normal lack of attack on the body's own tissues.

### **Immunologic Factor**

A substance, cell, gene, or process that affects immune-system activity or control. Examples include antibodies, immune cells, signaling proteins, inherited traits, medicines, and conditions that alter defense against disease.

### **Immunologic Memory**

The capacity of the adaptive immune system to respond more rapidly and strongly after re-exposure to an antigen because of long-lived memory lymphocytes.

### **Immunologic Skin Test**

An immunologic skin test places a small amount of an antigen or allergen in or on the skin to assess immune reactivity, such as allergy or tuberculosis exposure.

### **Immunologic Surveillance**

The immune system's ongoing detection and removal of abnormal, infected, or damaged cells. This protection helps control infections and may recognize early cancerous changes, although it is not always successful.

### **Immunologic Technique**

A laboratory or clinical method that uses antibodies,

---

antigens, or immune reactions to detect or measure a substance.

### **Immunologic Test**

An immunologic test examines immune cells, antibodies, antigens, or immune-related proteins to help evaluate infection, allergy, autoimmune disease, or immune deficiency.

### **Immunology**

The medical and biological study of the immune system, including its normal defenses and disorders such as allergy, autoimmunity, immunodeficiency, infection, and transplant rejection.

### **Immunomodulation**

Modification of immune activity to enhance, suppress, or redirect an immune response. Immunomodulatory medicines are used in cancer, autoimmune disease, transplantation, and infection but may increase susceptibility to infection or other adverse effects.

### **Immunomodulators**

Medicines or other treatments that change immune-system activity. Depending on the treatment, they may increase, reduce, or redirect immune responses and are used for conditions such as cancer, autoimmune disease, and some infections. Effects and risks vary by treatment.

#### **Alternate Names:**

Immunomodulator; Immune-Modifying Medicines

### **Immunomodulators In Dermatology**

Medicines that adjust local immune activity to treat conditions such as eczema and other inflammatory skin diseases. Topical calcineurin inhibitors are steroid-sparing options and may cause burning or irritation.

### **Immunomodulatory**

Describing a treatment or substance that changes immune activity. It may stimulate, suppress, or otherwise regulate the immune response, depending on the medicine and condition.

### **Immunophenotyping**

A laboratory method that identifies cells by the proteins on or inside them, usually using flow cytometry or stained tissue. It helps classify leukemias and lymphomas and evaluate some immune disorders.

### **Immunoprotein**

A protein involved in immune defense or immune regulation, such as an antibody, complement protein, cytokine, or antigen-presenting molecule. Immunoproteins identify threats, coordinate immune cells, or help remove damaged or infected material.

### **Immunoradiometric Assay**

A laboratory test using radioactive labels on antibodies to measure a specific substance, such as a hormone, protein, or other marker.

### **Immunoregulation**

Control of the strength, duration, and target of immune responses, including activation, suppression, and tolerance. Abnormal immunoregulation contributes to autoimmunity, allergy, immunodeficiency, and chronic inflammation.

### **Immunosenescence**

Age-related changes in immune defenses that can increase susceptibility to some infections, reduce vaccine responses, and alter inflammation. The effects vary with chronic disease, nutrition, medicines, and overall function.

### **Immunosuppression**

Reduced immune-system activity caused by disease, medicines, cancer treatment, or another condition. It can prevent rejection of a transplant or control autoimmune disease but increases susceptibility to infections and some cancers.

### **Immunosuppression In Transplant**

Treatment that reduces immune activity to prevent rejection of a transplanted organ. It can increase infection and cancer risk, affect kidney or liver function, and cause other medicine-related complications.

---

## Immunosuppression Screening

Checks before immune-suppressing treatment to identify infections, cancer risks, organ problems, pregnancy when relevant, and missing vaccinations. Tests may include tuberculosis, hepatitis B or C, HIV, blood counts, and liver and kidney tests.

## Immunosuppressive

Describing a medicine, treatment, or condition that reduces immune-system activity. Immunosuppression can prevent transplant rejection or control autoimmune disease, but it may increase the risk of infection, some cancers, and reduced vaccine responses.

## Immunotherapy

Cancer treatment that helps the immune system recognize or attack cancer cells, or supplies immune cells or antibodies that do so. Main approaches include checkpoint medicines, engineered immune cells, antibodies, and immune-stimulating treatments. Benefits and side effects depend on the cancer and treatment.

## Immunotherapy And Biologic Treatments

Treatments made from or designed to affect living biological systems, including antibodies and other medicines that adjust immune activity. They are used for selected cancers, autoimmune diseases, and inflammatory disorders and may increase infection or organ-inflammation risk.

## Immunotherapy For Cancer

Cancer treatment that stimulates, restores, redirects, or supplies immune activity to recognize and attack malignant cells, including immune-checkpoint inhibitors, cellular therapies, and antibodies.

## Immunotherapy In Dermatology

Treatment that changes immune activity to manage selected skin cancers or inflammatory skin diseases. The treatment and its risks depend on the diagnosis; immune-based cancer medicines can also inflame normal organs.

## Immunotoxicity

Harm to immune-system function caused by chemicals, medicines, infections, or other exposures. It may weaken defenses, trigger excessive sensitivity, cause autoimmunity, or change responses to vaccines and infections.

## Immunotoxin

A targeted toxin made by linking a toxic protein or drug to an antibody or binding molecule that directs it toward selected cells.

## Immunoturbidimetry

A laboratory method that measures cloudiness produced when antibodies react with a target antigen, allowing estimation of the target's concentration.

## Impacted

Firmly wedged or unable to move from its normal position, as with an impacted tooth or stool. The result may cause pain, blockage, inflammation, infection, or damage to nearby tissues.

## Impacted Fracture

A fracture in which one broken bone fragment is driven firmly into another, often producing relative stability but sometimes causing shortening or deformity.

## Impacted Tooth

A tooth that cannot emerge normally because it is blocked by bone, gum, or another tooth. Wisdom teeth and upper canine teeth are commonly affected and may cause pain, infection, decay, cysts, or damage to neighboring teeth.

## Impacted Wisdom Teeth

Wisdom teeth that cannot emerge normally because they are blocked by bone, gum, or another tooth.

### Alternate Names:

Impacted Third Molars; Third Molar Impaction

## Impacted Wisdom Teeth Cross-Reference

Alternate name; see *Impacted wisdom teeth*.



---

## Impaction

Blockage caused by material that is firmly lodged, such as hardened stool, a foreign object, or an impacted tooth. Symptoms depend on the site and may include pain, swelling, vomiting, or inability to pass contents.

## Impaired Fasting Glucose

A fasting blood-glucose level above normal but below the range used to diagnose diabetes. It indicates increased future diabetes risk and may improve with weight management, activity, and treatment of risk factors.

## Impaired Glucose Tolerance

A blood-glucose level that is higher than normal two hours after drinking a measured glucose solution but below the range used to diagnose diabetes. It indicates increased future risk of type 2 diabetes and cardiovascular disease.

## Impaired Smell

Reduced or absent ability to detect odors, called hyposmia or anosmia, caused by nasal obstruction, viral injury, head trauma, neurodegeneration, medications, or other disease.

## Impaired Taste

Impaired taste is reduced, distorted, or absent ability to taste and may result from infection, medicines, oral disease, nerve injury, or changes in smell.

## IMP Dehydrogenase

An enzyme that helps cells make guanosine nucleotides, building blocks needed for DNA and RNA. Blocking it can slow the growth of immune cells, which is why some immunosuppressive medicines target this pathway.

## Impedance Hearing Testing

A group of tests, including tympanometry, that measure the movement of the eardrum and middle-ear function.

## Impedance Plethysmography

A test that measures changes in a limb's electrical resistance as its blood volume changes. It has been used to assess venous blood flow.

## Impella Device

A small temporary heart pump inserted through a blood vessel or during surgery. It moves blood from the left lower heart chamber into the aorta, supporting circulation and reducing the heart's workload during severe heart failure or some heart procedures.

## Imperforate Anus

A birth difference in which the anus is absent, closed, or not connected normally to the rectum. A newborn may not pass stool and may develop abdominal swelling; other spine, heart, kidney, limb, or digestive differences can occur.

## Imperforate Anus Repair

Surgery to create or reposition an anal opening and connect the rectum so stool can pass normally in a baby born with an absent or blocked anus. Some babies need temporary bowel diversion; follow-up assesses healing, constipation, and bowel control.

## Imperforate Hymen

A birth difference in which the hymen completely covers the vaginal opening. After puberty, menstrual blood may collect behind it, causing absent periods, monthly pelvic pain, difficulty urinating, or a pelvic swelling.

## Imperforate Vagina

A rare birth condition in which the vagina is absent, closed, or blocked. It can prevent menstrual blood from leaving the body and cause repeated monthly pain, a pelvic swelling, difficulty passing urine, or fertility problems.

## Impetigo

A contagious bacterial skin infection causing small blisters, sores, or honey-colored crusts. It is common in children but can affect anyone and is usually treated with hygiene measures and, when needed, prescribed antibiotics.

---

**Alternate Names:**

School Sores; Impetigo Contagiosa

**Impingement Syndrome**

Pain and restricted movement caused when a tendon or other soft tissue is compressed between bones, commonly in the shoulder during elevation of the arm.

**Impingement Syndrome of the Shoulder**

Shoulder pain caused when tendons or a fluid-filled cushion beneath the shoulder blade become crowded during arm movement. Pain and limited movement are common.

**Alternate Names:**

Subacromial Impingement; Rotator Cuff Impingement

**Implant**

A medical device, tissue, or material placed inside the body to replace, support, monitor, or deliver treatment.

**Implantable Cardioverter Defibrillator**

An implanted device that detects dangerous ventricular rhythms and treats them with antitachycardia pacing or an electrical shock.

**Implantable Cardioverter-Defibrillator**

An implanted device that monitors heart rhythm and delivers electrical therapy (antitachycardia pacing or defibrillation) for life-threatening ventricular arrhythmias.

**Implantable Defibrillator**

A surgically implanted device that detects dangerous ventricular arrhythmias and delivers pacing or an electrical shock to restore an effective rhythm.

**Implantable Glucose Sensor**

A sensor placed under the skin that measures glucose in interstitial fluid and sends readings to a receiver, reader, or insulin-delivery system. It supports continuous or intermittently scanned glucose monitoring.

**Implantable Loop Recorder**

A small device placed under the skin of the chest to record heart rhythm for months or years. It helps detect intermittent abnormal rhythms when fainting, palpitations, or a stroke remains unexplained by shorter monitoring.

**Implantable Port**

A small reservoir placed under the skin and connected to a tube in a large vein. It provides repeated access for medicines, fluids, blood products, or nutrition without a new vein puncture each time; infection, clotting, blockage, or damage can occur.

**Implantable Venous Access Port**

A totally subcutaneous long-term central venous access device consisting of a silicone or titanium injection reservoir septum connected to a central catheter, accessed percutaneously using a non-coring Huber needle.

**Alternate Names:**

Port-a-Cath; Subcutaneous Infusion Port; Mediport; Venous Access Port

**Implant-Associated Infection**

Infection involving an implanted device or the tissue around it. Germs may stick to the device and form a protective layer, so treatment may require antibiotics, drainage, or removal or replacement of the device.

**Implantation**

The stage of early pregnancy when a fertilized egg attaches to and begins growing into the lining of the uterus, usually about a week after fertilization. It allows the developing pregnancy to receive support from the placenta.

**Implantation Procedure**

Placement of a device, tissue, medicine-delivery system, or other material inside the body for treatment or support. Examples include joint, dental, heart, and breast implants; risks include bleeding, infection, movement, failure, or rejection of donor tissue.

---

## Implant-Based Breast Reconstruction

Surgical reconstruction of the breast mound following mastectomy or trauma utilizing silicone gel-filled or sterile saline-filled prosthetic implants, often preceded by submuscular or prepectoral placement of a temporary tissue expander.

### Alternate Names:

Prosthetic Breast Reconstruction; Implant Breast Reconstruction; Tissue Expander Breast Reconstruction

## Implant Failure

Loss of an implant's ability to work properly or remain safely in place. Causes include infection, breakage, loosening, wear, movement, blockage, material damage, or problems with the surrounding tissue.

## Implants

Medical devices, materials, or tissues placed in the body to replace, support, monitor, or deliver treatment. Possible problems include infection, rejection, movement, breakage, or failure.

## Implant Sterilization

A validated process that removes or kills all microorganisms, including spores, from an implant or implantable device before use. The method is chosen according to the device's material and design.

## Implant-Supported Prosthesis

A fixed or removable dental prosthesis that is supported or retained by one or more dental implants.

## Importance Weight

An importance weight is a statistical weight used to adjust observations so a sample better represents a target population or distribution.

## Impression

The concluding section of a medical or imaging report that summarizes the most important findings, their likely meaning, and recommended next steps.

## Imprinting

A pattern of gene expression determined by whether a gene copy was inherited from the mother or father.

## Impulse

An impulse is a brief signal or force; in physiology it commonly refers to an electrical signal travelling along a nerve or through the heart.

## Impulse Control Disorder

A class of psychiatric conditions characterized by a persistent failure to resist an impulse, drive, or temptation to perform an act harmful to oneself or others, including kleptomania, pyromania, and intermittent explosive disorder.

## Impulse Oscillometry

A breathing test that applies tiny pressure waves while a person breathes normally. It estimates how easily air moves through the lungs and can help assess asthma, chronic lung disease, or small-airway problems, especially when forceful breathing is difficult.

## Impulsive Behavior

Impulsive behavior is action with limited forethought or inhibition despite possible harm and may occur with psychiatric, neurologic, substance-related, or developmental conditions.

## Impulsivity

A tendency to act quickly without adequate consideration of consequences or difficulty delaying a response; it can occur in ADHD, mania, substance use, and other conditions.

## In

In clinical measurements, an abbreviation for inch or inches; the same letters may have other meanings in different contexts.

## Inactivated Vaccines

Vaccines made from microorganisms that have been killed or from parts of them. They cannot reproduce or cause the infection they are designed to prevent,

---

but they may need several doses or booster doses to maintain protection.

### **Inactivator**

A substance or process that makes a biological agent inactive. Inactivators may disable viruses, toxins, enzymes, or medicines and are used in immunity, laboratory testing, sterilization, and treatment.

### **Inamrinone**

Inamrinone is an intravenous medicine that increases heart contraction and widens blood vessels; it was used short term in severe heart failure but can cause low blood pressure and abnormal rhythms.

### **Inanition**

Severe physical depletion caused by inadequate food or nutrient intake, often accompanied by weight loss, weakness, dehydration, and metabolic disturbance. Causes may include starvation, malabsorption, chronic disease, or inability to eat.

### **Inappropriate Antidiuretic Hormone Secretion**

A metabolic disorder of impaired free-water excretion characterized by non-physiologic, continuous secretion of antidiuretic hormone (ADH) despite low plasma osmolality, resulting in euvolemic hypotonic hyponatremia, high urine osmolality, and elevated urinary sodium excretion.

#### **Alternate Names:**

SIADH; Schwartz-Bartter Syndrome

### **Inappropriate Sinus Tachycardia**

A non-paroxysmal cardiac arrhythmia characterized by an abnormally elevated resting sinus heart rate exceeding 100 beats per minute, or an exaggerated, disproportionate heart rate response (mean 24-hour heart rate exceeding 90 beats per minute) to minimal physiological stress or posture changes in the absence of secondary driving factors (such as fever, anemia, or thyrotoxicosis).

#### **Alternate Names:**

IST; Chronic Sinus Tachycardia

### **Inborn Error Of Metabolism**

An inherited disorder in which an enzyme, transporter, or other protein needed for body chemistry does not work correctly. Harmful substances may accumulate or essential substances may be lacking, causing illness from infancy or later in life.

### **Inborn Errors Of Immunity**

Inherited disorders that impair one or more parts of the immune system. They can cause repeated, unusually severe, or unusual infections, as well as autoimmunity, inflammation, allergy, or increased cancer risk.

### **Inborn Errors of Metabolism**

Alternate name; see *Inborn Errors of Metabolism*.

### **Inbreed**

To reproduce between genetically related individuals; inbreeding can increase the chance that recessive inherited disorders will occur.

### **Inbreeding**

Reproduction between people who share a recent ancestor, increasing the chance that a child inherits two copies of the same recessive disease-causing variant. Genetic counseling can help assess family risk.

### **Incarcerated Hernia**

A hernia whose protruded tissue cannot be returned to the abdominal cavity. It may progress to strangulation and loss of blood supply.

### **Incentive Spirometry**

A breathing device that provides visual feedback while a person inhales slowly and deeply. It may be used with coughing, mobilization, and other measures to encourage lung expansion after illness or surgery; it is a lung-expansion exercise and is not the same as diagnostic spirometry.

### **Incest**

Sexual activity or reproduction between close biological relatives. It can involve abuse or coercion and

---

increases the chance of inherited disorders in children; affected people need safe, confidential medical and safeguarding support.

### **Incidence**

The number of new cases of a disease or event in a defined population during a specified period. It describes the chance of developing the condition during that period and differs from prevalence.

### **Incidental Finding**

An unexpected result discovered during an examination or test performed for another reason. It may be harmless or important; follow-up depends on what was found, its appearance, symptoms, and personal risk factors.

### **Incidentaloma**

An unexpected mass or abnormality found during imaging or another test performed for a different reason. It may be harmless or important, so follow-up depends on its size, appearance, location, growth, and the person's risks.

### **Incised Wound**

A cut made by a sharp object such as a knife, piece of glass, or surgical instrument. It usually has straight, clean edges and may damage skin, blood vessels, nerves, tendons, or deeper organs.

### **Incision**

A deliberate cut made by a healthcare professional through the skin or another tissue, usually during surgery or to allow drainage. The wound is closed or left to heal according to its size, location, contamination, and purpose.

### **Incisional Biopsy**

Surgical removal of only part of a lesion for diagnosis, generally used when the lesion is too large or located where complete removal would be inappropriate as an initial procedure.

### **Incisional Hernia**

A hernia through a weakness in a previous surgical

incision. It may appear as a bulge that becomes more prominent with standing or straining. Risk factors include wound infection, obesity, smoking, poor nutrition, chronic cough, and repeated surgery.

### **Incision and Drainage**

A very common minor surgical procedure performed to treat a localized collection of pus beneath the skin or in soft tissues (an abscess). After the overlying skin is cleansed and numbed with local anesthesia, a clinician makes a deliberate scalpel incision through the skin over the most fluctuant (softest) area of the swelling. The accumulated pus, dead tissue, and bacterial debris are evacuated, internal fibrous compartments (loculations) are gently broken up, and the cavity is flushed with sterile saline. For larger or deep abscesses, a sterile gauze ribbon or rubber drain (such as a Penrose drain) may be placed inside to keep the wound edges open so residual fluid can continue draining freely. Incision and drainage provides rapid relief from throbbing pain and pressure, stops infection from spreading deeper into tissue or the bloodstream, and allows the wound cavity to heal cleanly from the inside out.

#### **Alternate Names:**

I&D; Abscess Drainage; Incision and Drainage of Abscess

### **Incisor**

A front tooth shaped mainly for cutting and biting food. Adults usually have eight incisors, and each has a crown above the gum and a root anchored in the jawbone.

### **Incisors**

The flat front teeth used mainly for biting and cutting food. Adults usually have eight incisors, and their health can be affected by decay, injury, gum disease, wear, or poor alignment.

### **Inclusion Body**

A visible collection of material inside a cell. It may consist of viral parts, stored fats or sugars, or abnormal proteins; its appearance can help laboratory specialists identify infection, cell injury, or another disease.

---

### **Inclusion Body Myositis**

A slowly progressive muscle disease, usually beginning after age 50, that causes uneven weakness of the thigh muscles and finger-bending muscles. It can lead to falls and difficulty gripping or swallowing; treatment focuses on safety, exercise, and support.

### **Inclusion Conjunctivitis**

Conjunctivitis caused by *Chlamydia trachomatis*, usually presenting with persistent discharge, follicular inflammation, and sometimes genital infection. In newborns it can be associated with infection elsewhere and complications.

### **Incompetence**

Failure of a structure to close or function adequately; examples include valvular incompetence of the heart and cervical incompetence during pregnancy.

### **Incompetent Cervix**

Painless shortening or opening of the cervix during the middle or late part of pregnancy, without normal labor. It can cause pregnancy loss or premature birth and is assessed with history and ultrasound.

### **Incomplete Abortion**

A pregnancy loss in which some pregnancy tissue remains in the uterus. It may cause vaginal bleeding, cramping, and an open cervix.

### **Incomplete Heart Block**

A cardiac conduction disturbance in which supraventricular impulses are conducted to the ventricles with delay (first-degree AV block) or intermittent failure (second-degree AV block Mobitz I or II), distinguishing it from complete (third-degree) AV block where no impulses conduct.

#### **Alternate Names:**

Partial Heart Block

### **Incomplete Miscarriage**

Partial loss of a pregnancy in which some pregnancy tissue remains inside the uterus, potentially causing bleeding, pain, infection, or continued pregnancy symptoms.

### **Incomplete Ossification**

Failure of bone or cartilage to mineralize or develop fully. It may be part of normal development or result from genetic, metabolic, nutritional, or skeletal disorders and is interpreted with age, imaging, and laboratory findings.

### **Incontinence**

Involuntary loss of control over urination, bowel movements, or both. It may result from weakened muscles, nerve disease, infection, constipation, childbirth, prostate problems, medicines, mobility limits, or other conditions.

### **Incontinentia Pigmenti**

An X-linked dominant disorder affecting skin, teeth, eyes, and the nervous system, caused by pathogenic variants in *IKBKG*. Skin lesions evolve through vesicular, verrucous, hyperpigmented, and atrophic stages and may be associated with retinal disease or seizures.

### **Incoordination**

Lack of coordination between muscles or body movements; the term is often used in older medical language.

### **Incoordination**

Impaired ability to perform smooth, accurate, coordinated movements, often called ataxia. Causes include cerebellar disease, sensory loss, vestibular disorders, intoxication, medicines, and other neurologic conditions.

### **Increased Appetite**

An abnormally excessive desire for food or marked increase in caloric intake, seen in endocrine disorders (such as diabetes mellitus and hyperthyroidism), hypothalamic injury, psychiatric illness, or medication side effects.

#### **Alternate Names:**

Hyperphagia; Polyphagia; Increased Appetite

### **Increased Head Circumference**

Increased head circumference means head size above



---

the expected range for age and sex and may reflect familial size, hydrocephalus, swelling, or a genetic condition.

### **Increased Intracranial Pressure**

Abnormally elevated pressure within the skull caused by brain swelling, hemorrhage, mass lesion, impaired cerebrospinal-fluid circulation, or venous obstruction. It can cause headache, vomiting, visual disturbance, reduced consciousness, and brain herniation.

### **Increased Libido**

Increased libido is heightened sexual desire that may be normal or associated with hormones, medicines, mood elevation, neurologic disease, or substance use.

### **Increased Osteoid**

Increased osteoid is excess unmineralized bone matrix, seen when bone formation outpaces mineralization as in osteomalacia or some metabolic disorders.

### **Increased Vascular Permeability**

A change during inflammation that makes small blood vessels leaky. Fluid, proteins, and immune cells then move into nearby tissue, causing swelling, redness, warmth, and sometimes pain.

### **Increment**

An increase or stepwise change in an amount, measurement, dose, or treatment intensity. The size and timing of an increment should be chosen carefully to achieve benefit while limiting harm.

### **Incretin Effect**

The physiological augmentation of insulin secretion observed after oral glucose administration compared to an identical intravenous glucose load, mediated by gastrointestinal incretin peptides such as glucagon-like peptide-1 (GLP-1) and glucose-dependent insulinotropic polypeptide (GIP).

#### **Alternate Names:**

Gut Incretin Response; Oral Glucose Insulinotropic Effect

### **Incretins**

Incretins are gut hormones, mainly GLP-1 and GIP, released after meals that enhance glucose-dependent insulin secretion; GLP-1 also suppresses glucagon, slows gastric emptying, and reduces appetite.

### **Incubation**

Incubation is the time between exposure to an infectious agent and the appearance of symptoms.

### **Incubation Period**

The time between exposure to an infectious agent and the beginning of symptoms. It varies with the organism, dose, route of exposure, and person's immune response.

### **Incubator**

A controlled-environment chamber culturing microorganisms, cells, or tissues at defined temperature, humidity, and atmospheric composition.

### **Incus**

The incus, commonly known as the anvil, is the middle ossicle of the three bones in the middle ear, articulating with the malleus at the incudomalleolar joint and with the stapes at the incudostapedial joint.

### **Indapamide**

Indapamide is a thiazide-like diuretic used mainly for hypertension; it promotes renal sodium excretion and may cause hyponatraemia, hypokalaemia, dehydration, hyperuricaemia, or photosensitivity.

### **Indel Mutation**

A genetic change in which one or more DNA building blocks are inserted or deleted. If the number is not a multiple of three, it can disrupt the protein reading frame.

### **Independence**

The ability to perform activities and make decisions without assistance, although a person may still use devices, adaptations, or occasional support.

---

## Indeterminate

Indeterminate means not definitively classified or interpreted because available findings are insufficient, conflicting, or nonspecific.

## India Immunization Schedule

The Universal Immunization Schedule (UIP) of the Government of India, providing free vaccines to all children, adolescents, and pregnant women through the public health system.

## Indian Tick Typhus

A tick-borne rickettsial infection endemic to India and South Asia, caused by *Rickettsia conorii* subsp. *indica* and transmitted by brown dog ticks (*Rhipicephalus sanguineus*). It typically manifests with acute high fever, severe headache, generalized maculopapular rash involving palms and soles, and occasionally a black necrotic inoculation eschar (tache noire) at the bite site.

### Alternate Names:

Indian Spotted Fever; *Rickettsia Conorii Indica* Infection

## Indican

Indican is a compound formed from tryptophan metabolism and excreted in urine; abnormal levels may occur with intestinal bacterial overgrowth or metabolic disease.

## Indicate

To show, suggest, or point to a finding or conclusion; an indication is a clinical reason for using a test, treatment, or procedure.

## Indication

A clinical condition, symptom, or question that provides the reason for performing a test, procedure, or treatment.

## Indigestion

Common term for dyspepsia, discomfort or burning in the upper abdomen that may include early fullness, bloating, belching, or nausea. Persistent symptoms, bleeding, weight loss, vomiting, anemia,

or difficulty swallowing require medical evaluation.

## Indigo Carmine

Indigo carmine is a blue dye used in selected medical procedures to visualize urinary or other fluid flow and in laboratory staining.

## Indirect Antiglobulin Test

A blood test that detects antibodies able to react with red blood cells. It is used before transfusion and during pregnancy to identify antibodies that could harm donated blood or a fetus's red cells.

## Indirect Calorimetry

A test that measures oxygen use and carbon-dioxide production to estimate how much energy the body uses at rest. It can help determine nutrition needs when usual calculations are unreliable.

## Indirect Coombs Test

An in vitro laboratory diagnostic test used to detect circulating, unattached IgG antibodies against red blood cell surface antigens in the patient's serum, commonly utilized in prenatal maternal Rh-D alloimmunization screening and pre-transfusion compatibility crossmatching.

### Alternate Names:

Indirect Antiglobulin Test; IAT

## Indirect Inguinal Hernia

An abdominal wall hernia in which abdominal viscera herniate through the deep inguinal ring lateral to the inferior epigastric vessels into the inguinal canal, resulting from congenital persistence and patent failure of closure of the embryonic processus vaginalis.

### Alternate Names:

Congenital Inguinal Hernia; Lateral Inguinal Hernia

## Indirect Veneer

A thin covering made outside the mouth and bonded to the front of a tooth to change its shape, color, or appearance. It may be made from porcelain or composite material. Tooth preparation, strength, hygiene, and long-term maintenance affect its suit-

---

ability.

## **Indium**

Indium is a metal used in industry and medical radioisotopes; exposure effects depend on the compound, dose, and route.

## **Indium-111**

A radioactive form of indium used as a tracer in selected nuclear-medicine tests. It may label white blood cells to help locate inflammation or infection, or be attached to another substance to help show certain tumors. It emits detectable radiation that is recorded by a special camera.

## **Indium-111 Scan**

A nuclear medicine scan that uses the radioactive tracer indium-111 to make selected tissues visible. It may be attached to white blood cells to look for inflammation or infection, or to a substance that helps find some tumors.

## **Indium-Labelled WBC Scan**

A nuclear medicine study in which a patient's white blood cells are labeled with indium-111 and reinjected to localize selected sites of inflammation or infection.

## **Indocyanine Green**

Indocyanine green is a fluorescent dye injected intravenously to assess liver function, blood flow, and retinal or choroidal circulation.

## **Indocyanine Green Clearance**

A dynamic liver function test measuring the elimination rate of intravenously administered indocyanine green dye from circulating plasma by hepatocytes (without enterohepatic recirculation), widely used preoperatively to calculate remnant liver functional reserve prior to major hepatic resection.

### **Alternate Names:**

ICG Clearance

## **Indole**

A chemical structure containing a nitrogen atom,

found in tryptophan and many natural substances and medicines. Indole-containing compounds can affect the nervous system, blood vessels, digestion, inflammation, or other body processes.

## **Indole-3-Carbinol**

A compound formed from glucosinolates in cruciferous vegetables and studied for effects on estrogen metabolism and cancer biology.

## **Indolent**

Slow to develop, progress, or cause symptoms; it often describes a relatively slow-growing disease or tumor.

## **Indoles**

A group of compounds containing an indole ring, including substances involved in biological signaling, metabolism, and medicines.

## **Indometacin**

Indometacin is a nonsteroidal anti-inflammatory medicine used for pain and inflammation; it can cause stomach bleeding, kidney problems, and cardiovascular complications.

## **Indomethacin**

Indomethacin is a potent NSAID used for inflammatory pain and selected acute conditions such as gout; gastrointestinal ulceration or bleeding, renal injury, cardiovascular events, and headache are important adverse effects.

## **Indomethacin Overdose**

Toxic exposure to excessive indomethacin, which can cause nausea, vomiting, stomach bleeding, kidney injury, seizures, or other serious effects.

## **Indoor Allergens**

Substances inside buildings that can trigger allergic or asthma symptoms, including dust mites, mold, cockroach particles, and animal dander. Exposure may cause rhinitis, conjunctivitis, wheezing, or eczema. Reduction of exposure and appropriate allergy or asthma treatment depend on the identified

---

trigger.

### **Induced Abortion**

Deliberate termination of a pregnancy using medicines or a procedure. Safe care depends on gestational age, patient preference, accurate information, legal context, trained providers, and assessment for complications such as heavy bleeding or infection.

### **Induced Hyperthermia**

Induced hyperthermia is deliberate elevation of body or tissue temperature for a therapeutic purpose, with careful monitoring to prevent heat injury.

### **Induced Labor**

Medical initiation of uterine contractions before labor begins naturally to achieve delivery when continuing pregnancy may pose greater risk.

### **Induced Sputum Analysis**

A test in which inhaled salty mist helps a person produce mucus from the airways. The mucus is examined for infection or inflammatory cells to help assess asthma, chronic lung disease, or bronchiectasis.

### **Inducible Laryngeal Obstruction**

Temporary narrowing of the larynx during breathing, often during exercise or exposure to an irritant. It can cause noisy breathing, throat tightness, cough, or shortness of breath and was formerly often called vocal-cord dysfunction.

### **Inducing Labor**

Starting childbirth artificially with medicines, breaking the amniotic sac, or other methods when continuing pregnancy may pose greater risks than delivery.

### **Induction**

Initiation of a physiologic or therapeutic process, such as induction of labor, anesthesia, remission, or immune tolerance. The method, indication, benefits, and risks depend on the clinical context.

### **Induction Of Labor**

Starting uterine contractions medically before labor begins on its own. It may be recommended when continuing pregnancy is riskier, using medicines or procedures that soften the cervix and encourage contractions.

### **Induction Therapy**

The first, often intensive, part of treatment given to reduce or remove a disease. In acute leukemia, its main aim is remission, meaning that tests find no clear evidence of leukemia; further treatment is usually needed to reduce the chance of return.

### **Indurated**

Abnormally firm or hardened when palpated, usually because of inflammation, edema, fibrosis, infiltration, or a tumor within the skin or underlying tissue.

### **Induration**

Abnormal hardening of tissue, often detected by palpation and caused by inflammation, infection, edema, fibrosis, or a mass.

### **Industrial Bronchitis**

Industrial bronchitis is chronic airway irritation and inflammation related to workplace dusts, fumes, or chemicals, causing cough, phlegm, and breathing limitation.

### **Industrial Disease**

A disease caused or worsened by workplace exposures or conditions, such as chemicals, dust, noise, radiation, infection, or repetitive strain.

### **Industrial Health**

The branch of occupational health concerned with preventing and managing illness and injury related to industrial work, workplace exposures, and working conditions.

### **Indwelling Catheter**

A catheter left in place for ongoing drainage, infusion, monitoring, or access; urinary indwelling

---

catheters require appropriate indications and care because they increase infection and trauma risk.

### **Indwelling Catheter Care**

Care of a tube that remains in the bladder or another body space to drain fluid. It includes hand hygiene, keeping the tube unobstructed, preventing infection, emptying the collection bag safely, and seeking help for fever, pain, blockage, or leakage.

### **Indwelling Pleural Catheter**

A flexible tube placed through the chest wall into the space around the lung to drain fluid that keeps returning. It can allow drainage at home and relieve breathlessness, especially when cancer or a lung that cannot fully expand causes the fluid. Possible problems include infection, blockage, pain, or air leakage.

#### **Alternate Names:**

IPC; Tunneled Pleural Catheter; PleurX Catheter

### **Inert Gas Narcosis**

A reversible state of central nervous system depression, euphoria, cognitive impairment, and perceptual distortion experienced by scuba divers breathing compressed air at ambient depths exceeding 30 meters (approximately 4 atmospheres absolute), produced by the increased partial pressure of dissolved nitrogen in neuronal lipid membranes.

#### **Alternate Names:**

Nitrogen Narcosis; Rapture of the Deep

### **Inertia**

Resistance to a change in motion; in obstetrics, uterine inertia means weak or ineffective contractions that slow labor.

### **Infant**

An infant is a child in the first year of life, a period of rapid growth and changing feeding, immune, neurologic, and developmental needs.

### **Infant Botulism**

A serious illness in which *Clostridium botulinum* bacteria grow in an infant's intestine and make a

toxin. Constipation, poor feeding, weak crying, drooping eyelids, limpness, and breathing weakness can develop.

### **Infant Cardiopulmonary Resuscitation**

Basic life support resuscitation protocol tailored for infants under 1 year of age, utilizing chest compressions at a rate of 100–120/min at a depth of 1.5 inches (4 cm) using the two-finger or two-thumb encircling technique combined with rescue ventilations.

#### **Alternate Names:**

Infant CPR; Pediatric Basic Life Support, Infant; BLS Infant

### **Infant Feeding**

Provision of breast milk, appropriately prepared formula, or both to meet an infant's energy, protein, fluid, and micronutrient requirements. Feeding recommendations depend on age, growth, medical conditions, developmental readiness, and safe preparation practices.

### **Infant Feeding Patterns and Diet**

Age-appropriate nutritional guidance for infants during the first year of life, emphasizing exclusive breastfeeding or iron-fortified infant formula for the first six months, followed by the gradual introduction of complementary solid foods.

#### **Alternate Names:**

Infant Nutrition; Infant Feeding Guidelines; Baby Feeding

### **Infant Formula**

Infant formula is manufactured nutrition designed to provide adequate nutrients when breast milk is unavailable, insufficient, or not chosen; preparation and storage must be safe.

### **Infant Formulas**

Commercial preparations designed to provide nutrition for infants when breast milk is unavailable, insufficient, or not chosen.

---

## Infant Gynecomastia

Infant gynecomastia is temporary breast enlargement in a newborn or young infant caused by maternal or neonatal hormones and usually resolves spontaneously.

## Infant Harlequin Syndrome

A transient, benign cutaneous vascular phenomenon in neonates characterized by asymmetric flushing of one half of the body with sharp midline demarcation, caused by transient asymmetric sympathetic tone. The affected side is red and warm while the contralateral side is pale.

## Infant Hearing Loss

Congenital or early-acquired auditory impairment in infants, resulting from genetic mutations (such as connexin 26 mutations), congenital infections (cytomegalovirus, rubella), prematurity, hyperbilirubinemia, or ototoxic drug exposure. Universal newborn hearing screening enables early amplification and speech-language intervention.

### Alternate Names:

Congenital Hearing Loss; Pediatric Auditory Impairment; Neonatal Deafness

## Infanticide

The killing of a child, in law typically defined as the killing of a newborn or infant (usually within the first year, or specifically within 24 hours of birth in some jurisdictions) by its mother while mentally disturbed by the effects of childbirth or lactation.

## Infantile Bradycardia

An unusually slow heart rate in an infant. It can occur normally during sleep, but may also signal low oxygen, infection, medicines, or a heart or conduction problem; symptoms and the infant's age determine the evaluation.

## Infantile Colic

Repeated periods of intense crying in an otherwise healthy young infant, often beginning in the first weeks and improving by three to four months. The baby may draw up the legs or seem difficult to soothe; persistent vomiting, fever, poor feeding, or

poor growth needs assessment.

## Infantile Cortical Hyperostosis (Caffey Disease)

A self-limited disorder of infancy characterized by irritability, soft-tissue swelling, and cortical thickening of affected bones. It is usually associated with COL1A1-related disease and resolves over time.

## Infantile Fibrosarcoma

A rare cancer of connective or soft tissue that usually appears in a baby or young child as a growing lump. It may involve the limbs or trunk; imaging and biopsy guide treatment, which requires specialist care.

## Infantile Hemangioma

Alternate name; see *Hemangioma*.

## Infantile Hepatic Hemangiomas

Benign growths made of extra blood vessels in an infant's liver. Small lesions may cause no symptoms, while numerous or large lesions can enlarge the liver, strain the heart, lower thyroid hormone, or cause bleeding problems. Ultrasound and specialist follow-up guide monitoring and treatment.

## Infantile Paralysis

Paralytic poliomyelitis caused by poliovirus infection involving motor neurons, which can produce acute flaccid paralysis and respiratory failure; vaccination prevents most cases.

## Infantile Spasm

A brief seizure in an infant, often involving sudden bending or stiffening of the body and occurring in clusters. It may affect development.

## Infantile Spasms

A severe seizure disorder of infancy characterized by brief, sudden flexion or extension spasms, often occurring in clusters and associated with developmental impairment.



---

### **Infant Incubators**

Enclosed devices providing controlled warmth, humidity, oxygen, and monitoring for premature, small, or ill newborns who need extra support after birth.

### **Infant Jaundice**

Yellowing of an infant's skin or eyes caused by excess bilirubin. It is often temporary, but jaundice in the first day, rapid worsening, poor feeding, unusual sleepiness, or pale stools can occur with more serious disease.

### **Infant Leukemia**

Infant leukemia is leukemia diagnosed during infancy, in which abnormal blood-forming cells multiply and interfere with normal blood-cell production.

### **Infant Mortality**

The death of an infant before the first birthday; the infant mortality rate is usually expressed per 1,000 live births.

### **Infant Nutrition**

Nutritional support for infants through breast milk, appropriately prepared formula, and developmentally suitable complementary foods. Requirements change with age, growth, illness, prematurity, and feeding ability; unsafe preparation can cause infection or malnutrition.

### **Infant Of A Substance-Using Mother**

A newborn exposed to alcohol or other substances during pregnancy who may need assessment for withdrawal, toxicity, feeding problems, or developmental effects.

### **Infant Of Diabetic Mother**

A newborn born to a mother with diabetes who may be at increased risk of large size, low blood sugar, breathing problems, jaundice, or congenital abnormalities.

### **Infant Reflex**

An infant reflex is an automatic response present during early development, such as rooting, sucking,

grasping, or startle, whose timing and symmetry help assess neurologic function.

### **Infant Reflexes**

Automatic newborn responses such as rooting, sucking, Moro, grasp, and stepping that help assess neurologic development.

### **Infant Reflux**

Backflow of stomach contents into an infant's esophagus, often causing uncomplicated spitting up. It usually improves with age, but poor growth, breathing problems, blood, forceful vomiting, or distress can occur with more serious disease.

### **Infant Respiratory Distress Syndrome**

A serious breathing disorder in a newborn or young infant, usually caused by insufficient lung surfactant and immature lungs. It is also called neonatal respiratory distress syndrome.

### **Infant Sleep Position**

Placing an infant on the back for every sleep is recommended to reduce the risk of sleep-related death. Infants should sleep on a firm, separate surface without pillows, loose bedding, bumpers, or stuffed toys.

### **Infant Test Procedure Preparation**

Preparing a baby for a test may involve checking feeding or fasting instructions, bringing medicines and records, explaining the procedure to caregivers, and planning comfort, warmth, safe positioning, and monitoring afterward.

### **Infarct**

An area of tissue that has died because it did not receive enough blood and oxygen. An infarct can occur in the heart, brain, intestine, kidney, spleen, or other organs.

### **Infarction**

Tissue death caused by inadequate blood flow and oxygen, usually because a vessel is blocked. Examples include heart attack, brain infarction, and

---

intestinal infarction.

### **Infection**

Entry and multiplication of bacteria, viruses, fungi, parasites, or other microorganisms in the body. Infection may cause tissue damage and illness, although some infections produce no symptoms; the cause and site determine treatment.

### **Infection Control**

Practices that prevent microorganisms from spreading between people, objects, and healthcare settings. They include hand cleaning, appropriate protective equipment, safe injection and device care, cleaning, sterilization, vaccination, and isolation when needed.

### **Infection Imaging**

Imaging used to locate an infection when examination and routine tests do not show its source. Depending on the suspected problem, it may use ultrasound, CT, MRI, or a nuclear-medicine tracer; scan findings must be interpreted with symptoms, cultures, and other tests.

### **Infections in Diabetes**

Infection in a person with diabetes, in whom hyperglycemia, vascular disease, neuropathy, or immune dysfunction may increase susceptibility or complicate healing. Assessment includes the source, severity, glycemic status, organ involvement, and need for source control.

### **Infection Prevention**

Measures that reduce the risk of infection, including hand hygiene, personal protective equipment, environmental cleaning, sterilization, disinfection, vaccination, and appropriate isolation precautions.

### **Infections In Pregnancy**

Infections during pregnancy can affect the pregnant person, placenta, fetus, or newborn. Some, including listeriosis, group B streptococcal infection, urinary infection, and sexually transmitted infections, can cause miscarriage, premature birth, fetal illness, or newborn sepsis. Testing and treatment

depend on the organism and stage of pregnancy.

### **Infectious Arteritis**

Inflammation and damage of an artery caused by infection, which can weaken the vessel, reduce blood flow, or produce aneurysm.

### **Infectious Arthritis**

Inflammation of a joint caused by infection, usually bacterial and sometimes viral or fungal. Acute pain, swelling, warmth, fever, and restricted movement may occur, and joint damage can develop quickly.

### **Infectious Conjunctivitis**

Inflammation of the conjunctiva caused by a virus, bacterium, or less commonly another infectious organism, often causing redness and discharge.

### **Infectious Disease**

Illness caused by a pathogen such as a bacterium, virus, fungus, or parasite. Infection may spread between people, from animals or the environment, or from organisms normally living in the body.

### **Infectious Diseases**

Illnesses caused by bacteria, viruses, fungi, parasites, or prions. They may spread through contact, droplets or air, food or water, insects, blood, or sexual contact; prevention and treatment depend on the organism.

### **Infectious Dose**

The amount of a microorganism that must enter the body to cause infection under particular conditions. It varies with the organism, route of exposure, dose strength, and the person's immune defenses.

### **Infectious Encephalitis**

Brain inflammation caused by an infectious organism, most often a virus. It may cause fever, headache, confusion, seizures, behavior changes, weakness, or reduced consciousness.

### **Infectious Encephalomyelitis**

Infection-related inflammation of the brain and

---

spinal cord that may cause fever, headache, confusion, weakness, seizures, or other neurologic signs.

### **Infectious Enterocolitis**

Infection-related inflammation of the small intestine and colon, commonly causing diarrhea, abdominal pain, fever, or vomiting.

### **Infectious Esophagitis**

Infection and inflammation of the esophagus, most often caused by *Candida* yeast or herpes viruses in people with weakened immunity. It commonly causes painful or difficult swallowing and pain behind the breastbone.

### **Infectious Hepatitis**

Alternate name; see *Viral Hepatitis*.

### **Infectious Mediastinitis**

Infection and inflammation of the tissues in the middle of the chest, often after esophageal injury, surgery, or spread from a nearby infection.

### **Infectious Meningitis**

Infection and inflammation of the membranes surrounding the brain and spinal cord. Bacterial meningitis may cause headache, fever, neck stiffness, vomiting, confusion, or a new rash.

### **Infectious Mononucleosis**

A viral illness, usually caused by Epstein–Barr virus, that causes marked fatigue, sore throat, fever, and swollen lymph nodes. The spleen may enlarge during the illness.

#### **Alternate Names:**

Mono (Infectious Mononucleosis)

### **Infectious Myringitis**

Inflammation and infection of the tympanic membrane, often associated with viral or bacterial respiratory infection. It can cause severe ear pain, fever, hearing reduction, and small fluid-filled blisters on the eardrum.

### **Infective Cervicitis**

Infective cervicitis is inflammation of the cervix caused by organisms such as chlamydia, gonorrhea, trichomonas, or herpes, producing discharge, bleeding, or pelvic discomfort.

### **Infective Cystitis**

Infective cystitis is bacterial or other microbial inflammation of the bladder, causing urinary frequency, urgency, burning, and sometimes blood or fever.

### **Infective Endocarditis**

An infection of the heart's inner lining, most often involving a native or prosthetic heart valve and usually caused by bacteria; fungi and other organisms can also cause it. It may form infected vegetations, destroy valves, cause fever or a new murmur, and send emboli to other organs. Diagnosis integrates blood cultures, echocardiography or other imaging, and clinical findings.

### **Infective Endocarditis Duke Criteria**

A validated clinical, microbiological, and echocardiographic diagnostic scoring system used to establish the diagnosis of infective endocarditis. Major criteria include persistent positive blood cultures for typical organisms and echocardiographic evidence of endocardial involvement (vegetation, abscess, or new dehiscence); minor criteria include predisposing heart condition, fever, vascular phenomena, and immunologic phenomena.

#### **Alternate Names:**

Duke Criteria for Infective Endocarditis; Modified Duke Criteria

### **Infective Endocarditis Pathology**

Infection-associated vegetations composed of fibrin, platelets, inflammatory cells, necrotic debris, and microorganisms on a valve or endocardial surface. Destruction, abscesses, emboli, and healing fibrosis may occur.

### **Infective Phlebitis**

Infective phlebitis is infection and inflammation of a vein, often around an intravenous catheter, causing

---

pain, redness, swelling, and possible bloodstream infection.

### **Infective Vaginitis**

Infective vaginitis is infection or inflammation of the vagina caused by yeast, bacteria, parasites, or viruses and may cause discharge, itching, odor, or pain.

### **Inferior**

Below or nearer the feet compared with another body part. For example, the diaphragm is inferior to the heart, and the inferior vena cava carries blood from the lower part of the body to the heart.

### **Inferior Alveolar Nerve Block**

A local-anesthetic injection near a nerve supplying the lower jaw. It temporarily numbs the lower teeth, jawbone, and lower lip on that side and is commonly used for dental procedures.

### **Inferior Colliculus**

The inferior colliculus is a midbrain auditory relay that integrates sound information and helps orient reflexively to auditory stimuli.

### **Inferior Hemifield Loss**

Loss of vision in the lower half of the area a person can see with one or both eyes. Causes include retinal disease, a stroke, or another problem along the brain's visual pathways; sudden loss can occur with serious disease.

### **Inferior Mediastinum**

The part of the mediastinum below the level of the sternal angle. It is divided into the anterior, middle, and posterior mediastinum.

### **Inferior Mesenteric Artery**

A major artery supplying the lower part of the large intestine, including the descending and sigmoid colon and upper rectum. Blockage or reduced flow can cause abdominal pain and ischemic colitis.

### **Inferior Myocardial Infarction**

A heart attack affecting the lower surface of the heart muscle. It usually results from blocked blood flow in a coronary artery and can cause permanent damage.

### **Inferior Petrosal Sinus Sampling**

A special blood test done through small catheters placed in veins near the pituitary gland. It compares hormone levels from the pituitary area with levels in a regular vein to help find the source of excess ACTH in suspected Cushing disease.

#### **Alternate Names:**

IPSS

### **Inferior Vena Cava**

The body's largest vein, carrying blood from the legs, pelvis, abdomen, and liver to the right side of the heart. A clot or pressure on it can cause swelling of both legs and increase the risk of serious complications.

### **Inferior Vena Cava Filter**

A small device placed in the inferior vena cava to catch clots traveling from the legs or pelvis to the lungs. It is used selectively, usually for acute venous thromboembolism when anticoagulation is contraindicated or has failed; complications include migration, vessel perforation, thrombosis, and difficulty removing retrievable filters.

#### **Alternate Names:**

IVC Filter; Inferior Vena Cava Filters; Greenfield Filter

### **Inferior Vena Cava Syndrome**

A constellation of clinical symptoms caused by partial or complete occlusion, thrombosis, or external tumor compression of the inferior vena cava, characterized by bilateral lower-extremity pitting edema, extensive superficial abdominal wall collateral vein dilation with cephalad blood flow, and dependent ascites.

#### **Alternate Names:**

IVC Obstruction; Caval Syndrome

---

### Inferior Wall Myocardial Infarction

Ischemic myocardial necrosis localized to the inferior (diaphragmatic) aspect of the left ventricle, usually caused by acute thrombotic occlusion of the right coronary artery (RCA) or left circumflex artery, presenting with ST-segment elevation in leads II, III, and aVF, frequently associated with sinus bradycardia or AV block.

#### Alternate Names:

IWMI; Inferior STEMI

### Inferolateral

Located below and toward the side of a reference structure.

### Infertility

Failure to achieve pregnancy after 12 months of regular unprotected intercourse, or sooner when age or medical history warrants evaluation. Causes may involve ovulation, sperm, reproductive organs, age, or unexplained factors.

#### Alternate Names:

Subfertility

### Infestation

Presence of parasites such as lice, mites, or worms on or inside the body. The parasite may live on the skin or in an organ and can cause itching, inflammation, tissue injury, or other symptoms.

### Infiltrate

An area of increased tissue density seen on an image, often in the lung. It is nonspecific and may represent infection, fluid, bleeding, inflammation, collapse, tumor, or another process.

### Infiltrating Cancer

Cancer that has grown beyond its original location into nearby tissue. Such growth may damage normal structures and can allow cancer cells to enter lymph vessels or blood vessels and spread elsewhere.

#### Alternate Names:

Invasive cancer

### Infiltrating Ductal Carcinoma

Alternate name; see *Invasive Ductal Carcinoma*.

### Infiltration

Entry or accumulation of cells, fluid, or another substance within tissue. In clinical care, intravenous infiltration means leakage of nonvesicant fluid into surrounding tissue; extravasation of an irritant or vesicant can cause serious injury.

### Inflamed Gallbladder

Alternate name; see *Cholecystitis*.

### Inflamed Pancreas

Alternate name; see *Pancreatitis*.

### Inflamed Pericardium

Alternate name; see *Pericarditis*.

### Inflammaging

The gradual, low-level increase in inflammation that can occur with aging. It is influenced by chronic illness, immune-system changes, infections, and tissue damage and may contribute to frailty, artery disease, muscle loss, or brain disorders.

### Inflammasomes

Groups of proteins inside immune cells that detect danger signals and activate inflammation. They help release inflammatory messenger proteins and can cause an infected or damaged cell to die. Excessive activation may contribute to autoimmune or inflammatory disease.

### Inflammation

The body's protective response to injury, infection, or irritation. It may cause redness, warmth, swelling, pain, and reduced function. Short-term inflammation helps healing, while persistent inflammation can damage tissues and contribute to chronic disease.

### Inflammation Imaging

Imaging used to find areas of active inflammation in the body. A scan using a sugar-like tracer may show infection or inflammation, but cancer can look

---

similar, so scan findings must be interpreted with symptoms, examination, and other tests.

### **Inflammatory Back Pain**

A pattern of chronic back pain characterized by insidious onset, improvement with exercise, lack of improvement with rest, nocturnal pain, and prolonged morning stiffness. It suggests axial spondyloarthritis but is not diagnostic without supportive clinical, laboratory, and imaging findings.

### **Inflammatory Bowel Disease**

A group of chronic conditions that cause inflammation in the digestive tract, mainly Crohn disease and ulcerative colitis. Symptoms may come and go and can include abdominal pain, diarrhea, rectal bleeding, and weight loss. The exact cause is unknown, but genes, the immune system, the environment, and gut bacteria may all play a role.

### **Inflammatory Bowel Disease Diet**

An eating plan adapted for Crohn disease or ulcerative colitis to maintain nutrition and reduce symptoms. During flares, temporary changes or liquid nutrition may help; food triggers vary, and restrictive diets should be supervised.

### **Inflammatory Bowel Disease: Intestinal Problems**

Inflammatory bowel disease, including Crohn disease and ulcerative colitis, causes chronic immune-mediated intestinal inflammation with diarrhea, pain, bleeding, weight loss, and extraintestinal effects. Treatment aims for sustained remission and prevention of structural and nutritional complications.

#### **Alternate Names:**

IBD Enteropathy; Inflammatory Bowel Disease

### **Inflammatory Breast Cancer**

A rare, aggressive breast cancer causing redness, swelling, warmth, and thickened skin, sometimes without a distinct lump. It requires prompt imaging, biopsy, staging, and coordinated treatment.

#### **Alternate Names:**

IBC; Inflammatory Carcinoma of the Breast

### **Inflammatory Myopathies**

A group of disorders in which inflammation damages muscle, usually causing gradually worsening weakness around the hips, thighs, shoulders, or upper arms. Some forms also affect skin, lungs, swallowing, or other organs. Diagnosis and treatment require specialist assessment.

### **Inflammatory Myopathy**

A disorder in which immune or other inflammatory processes damage muscle, causing weakness and sometimes pain or involvement of the skin, lungs, swallowing muscles, or other organs.

### **Inflammatory Response**

The body's organized reaction to infection, injury, or irritation. Blood flow changes and immune cells move into the affected area, causing signs such as redness, warmth, swelling, pain, or reduced function.

### **Infliximab**

A biologic medicine that blocks tumor necrosis factor, an inflammatory signal. It treats several inflammatory diseases but can increase the risk of serious infections, tuberculosis reactivation, infusion reactions, and heart-failure worsening.

### **Influenza**

A contagious respiratory illness caused by influenza A or B viruses, characterized by sudden onset of fever, myalgia, headache, malaise, nonproductive cough, sore throat, and nasal congestion. Seasonal epidemics occur annually. Complications: pneumonia (primary viral or secondary bacterial), exacerbation of underlying conditions.

### **Influenza A Virus**

An influenza virus with segmented RNA that infects people and animals and can cause seasonal epidemics or, after major genetic changes, pandemics.

### **Influenza Test**

A test for influenza A or B using a respiratory specimen. Rapid antigen tests provide quick results but may miss infection. Molecular tests are generally more sensitive and may identify or subtype influenza



---

viruses; results must be interpreted with symptoms, timing, and local laboratory methods.

### **Influenza Vaccination In Pregnancy**

Influenza vaccination during pregnancy helps protect the pregnant person from severe flu and passes some protection to the newborn. The recommended vaccine is inactivated; timing and product choice should follow current local guidance.

### **Influenza Virus**

A segmented, negative-sense RNA virus belonging to the Orthomyxoviridae family. Influenza causes seasonal epidemics and occasional pandemics of respiratory illness. The virus has two surface glycoproteins: hemagglutinin (HA), which mediates attachment, and neuraminidase (NA), which facilitates release.

### **Influenzavirus A**

A group of influenza A viruses that infect people and animals. They are classified into subtypes according to two surface proteins and can cause seasonal outbreaks, severe illness, or pandemics.

### **Influenzavirus B**

The virus genus containing influenza B viruses, which mainly infect humans and cause seasonal respiratory illness.

### **Informatics**

Informatics applies information science, data, computing, and human factors to organize and improve healthcare, biology, research, or another field.

### **Informed Consent**

A patient's voluntary agreement to a proposed intervention after receiving understandable information about its purpose, benefits, risks, alternatives, and the option of declining, with decision-making capacity or appropriate authorization.

### **Infrared Rays**

Infrared rays are electromagnetic radiation with wavelengths longer than visible red light; excessive

exposure can heat tissue and cause burns.

### **Infrared Spectrometry**

A method for identifying or measuring substances by examining how they absorb infrared light. Different chemical bonds absorb different wavelengths, creating a pattern that can help analyze medicines, tissues, samples, or pollutants.

### **Infrared Therapy**

Treatment using infrared radiation, usually to provide heat or light to tissues. It is used in some physiotherapy and skin treatments, but benefits vary and excessive exposure can cause burns or eye injury.

### **Infrarenal Aorta**

The infrarenal aorta is the portion of the abdominal aorta below the renal arteries and above its division into the iliac arteries.

### **Infraspinatus Muscle**

A thick, triangular rotator cuff muscle originating from the infraspinous fossa of the scapula and inserting onto the middle facet of the greater tubercle of the humerus, innervated by the suprascapular nerve; acts primarily as the chief external (lateral) rotator of the glenohumeral joint.

### **Infratentorial Neoplasm**

An abnormal growth in the lower part of the skull, near the cerebellum and brainstem. Its effects depend on size and location and may include headache, vomiting, balance problems, weakness, or changes in coordination.

### **Infrequent**

Infrequent means occurring less often than usual or expected, such as infrequent bowel movements, symptoms, or events.

### **Infundibular Stenosis**

Subvalvular muscular narrowing or hypertrophy of the right ventricular outflow tract below the pulmonary valve, classically forming the dynamic sub-

---

pulmonary obstruction component in tetralogy of Fallot or isolated subpulmonic stenosis.

### **Infundibulopelvic Ligament**

A peritoneal fold extending from the lateral pelvic wall to the superior pole of the ovary that conveys the ovarian artery, ovarian vein, lymphatic channels, and ovarian nerve plexus, serving as a critical surgical landmark during salpingo-oophorectomy.

#### **Alternate Names:**

Suspensory Ligament of the Ovary; IP Ligament

### **Infundibulum**

A funnel-shaped anatomical structure or passage, specifically referring to the hollow neural stalk connecting the hypothalamic median eminence to the posterior pituitary gland (hypophysis), or the funnel-shaped distal expansion of the fallopian tube.

#### **Alternate Names:**

Pituitary Stalk; Funnel-Shaped Anatomical Passage

### **Infuse**

To introduce a fluid, medicine, or other substance gradually into the body, commonly through an intravenous line; it can also mean to soak a substance in a liquid.

### **Infusion**

Slow delivery of a fluid or medicine into a vein, under the skin, or another body site. It allows controlled dosing over time and may require monitoring for allergic reactions, infection, or fluid overload.

#### **Alternate Names:**

Intravenous infusion

### **Infusion Pump**

A medical device that delivers a measured amount of fluid or medicine at a controlled rate. Depending on its design, it may deliver material intravenously, subcutaneously, or intrathecally. Common types include volumetric infusion pumps and syringe pumps; alarms may indicate occlusion, air, low battery, or other device conditions.

#### **Alternate Names:**

IV Infusion Pump

### **Ingested Foreign Body**

An object accidentally or intentionally swallowed that lodges in the esophagus, stomach, or intestines, such as coins, small toys, button batteries, or sharp objects requiring clinical assessment and possible endoscopic retrieval.

#### **Alternate Names:**

Swallowed Foreign Object; Foreign Body Ingestion; Gastrointestinal Foreign Body

### **Ingestion**

The act of taking a substance into the digestive tract by swallowing. Ingestion of a toxic substance can cause poisoning.

### **Ingredient**

An ingredient is a substance included in a medicine, food, cosmetic, or other preparation, whether active or inactive.

### **Ingrown Hair**

A hair that curls back into or grows sideways beneath the skin, causing an itchy or painful bump. It may become inflamed or infected, especially after shaving, waxing, or close trimming.

### **Ingrown Nail**

An ingrown nail occurs when the nail edge grows into surrounding skin, causing pain, redness, swelling, and sometimes infection or granulation tissue.

#### **Alternate Names:**

Onychocryptosis; Ingrown Toenail

### **Ingrown Toenail (Onychocryptosis)**

An ingrown nail affecting a toe, usually the great toe, in which the nail edge penetrates the lateral nail fold. Persistent pain, pus, spreading redness, or recurrent disease may require clinician treatment or partial nail removal.

### **Ingrown Toenails**

A condition in which the edge of a toenail grows into nearby skin, most often on the big toe. It causes pain, redness, and swelling; infection, diabetes, or severe symptoms may complicate it.

---

## Inguinal

Relating to the groin, the region where the lower abdomen meets the upper thigh and where certain hernias, lymph nodes, and injuries occur.

## Inguinal Canal

A short passage through the lower abdominal wall near the groin. It carries the spermatic cord in males and the uterine round ligament in females; weakness here can allow an inguinal hernia.

## Inguinal Hernia

A bulge formed when fat or intestine pushes through a weak area in the groin. It may cause discomfort or a lump that appears with standing or straining; severe pain, vomiting, or a trapped lump may indicate reduced blood flow or bowel blockage.

## Inguinal Hernia Repair

Surgery that returns tissue to the abdomen and closes or strengthens a weak area in the groin. It may be performed through a larger incision or several small ones, with or without mesh; recurrence, infection, and persistent pain are possible.

## Inguinal Orchiectomy

Inguinal orchiectomy is removal of a testicle through an incision in the groin, commonly used when treating suspected testicular cancer.

## Inguinal Region

The groin area where the lower abdomen meets the upper thigh. It contains important muscles, blood vessels, nerves, and lymph nodes and is a common site for hernias and some injuries.

## Inhalant

A substance breathed into the lungs as a gas, vapor, aerosol, or smoke; inhalants may be medicines, workplace substances, or abused chemicals.

## Inhalant Substance Use

Inhalant misuse involves breathing fumes or gases from household or industrial products and can cause sudden cardiac arrest, brain injury, organ damage,

dependence, or death.

## Inhalation

Drawing air, medicine, particles, or vapors into the lungs. Inhalation is a route for respiratory medicines and anesthetics, while accidental inhalation of food, vomit, smoke, or toxic substances can cause airway obstruction or lung injury.

## Inhalational Anesthesia

General anesthesia given by breathing an anesthetic gas or vapor through a mask or breathing tube. The amount reaching the bloodstream controls unconsciousness and is adjusted while breathing, circulation, and other vital signs are monitored.

## Inhalation Injury

Damage to the airway or lungs from smoke, hot gases, or toxic chemicals. Swelling, cough, wheezing, low oxygen, and poisoning by carbon monoxide or cyanide may develop or worsen over several hours after a fire.

## Inhaled Corticosteroids

Anti-inflammatory medicines breathed into the airways to reduce swelling and mucus and prevent asthma symptoms. They do not provide instant relief; regular use can cause hoarseness or oral thrush, which good inhaler technique and mouth rinsing help reduce.

## Inhaled Foreign Body

An object or food particle aspirated into the larynx, trachea, or bronchial tree, causing sudden choking, coughing, wheezing, stridor, or complete airway obstruction requiring immediate first aid or bronchoscopic removal.

### Alternate Names:

Airway Foreign Body; Foreign Body Aspiration; Tracheobronchial Foreign Body

## Inhaled Nitric Oxide

A gas given through a breathing system to widen blood vessels in the lungs and sometimes improve oxygen levels. It is used short term in selected critical illnesses and requires continuous delivery and

---

monitoring for blood-salt or oxygen-related effects.

### **Inhaler**

A device that delivers medicine into the airways through the mouth. Pressurised inhalers release a measured spray, while dry-powder inhalers release powder during a strong breath in; correct technique is important for the medicine to work.

### **Inherited Disease**

A disease caused wholly or partly by a genetic change inherited from a parent or present in reproductive cells and therefore capable of being passed to offspring. Some disease-causing changes arise de novo in an egg, sperm, or early embryo rather than being inherited from a parent.

### **Inhibin**

A glycoprotein hormone produced mainly by the gonads that suppresses pituitary follicle-stimulating hormone secretion and contributes to regulation of reproductive function.

### **Inhibition**

Inhibition is a reduction or suppression of an activity, such as nerve signalling, enzyme action, muscle contraction, or cell growth.

### **Inhibitory Neurochemical**

A chemical messenger that decreases the activity of nerve cells or muscles. Gamma-aminobutyric acid and glycine are important inhibitory neurotransmitters in the nervous system.

#### **Alternate Names:**

Inhibitory neurotransmitter

### **Inhibitory Synapse**

A connection between nerve cells that reduces the chance that the receiving cell will generate an electrical signal.

### **Iniiencephaly**

A rare, usually lethal neural-tube defect characterized by occipital bone defect, severe retroflexion of the head, and marked shortening or distortion of

the cervical and thoracic spine. It is often identified prenatally and may occur with other congenital anomalies.

### **Initial Insomnia**

Initial insomnia is difficulty falling asleep at the beginning of the sleep period, often related to stress, habits, medicines, mood, or another sleep disorder.

### **Injectable Fillers**

Materials placed under or within the skin to restore volume, improve contours, or soften wrinkles. Effects vary by product and may include swelling, bruising, infection, lumps, or rare blood-vessel blockage.

#### **Alternate Names:**

Dermal fillers

### **Injection**

Introduction of a medication, fluid, or other substance into the body with a needle or other injection device.

### **Injection Drug Use**

Use of medicines or illicit substances by injection. Risks include overdose, blood-borne infection, skin and vascular injury, endocarditis, abscesses, and transmission through shared equipment.

### **Injury**

Damage to the body's tissues or function caused by trauma, force, heat, chemicals, radiation, or another harmful exposure.

### **Injury Dating**

Estimation of the age of an injury using clinical appearance, histopathology, imaging, and the history of symptoms. Exact dating is often uncertain, particularly for bruises and soft-tissue injuries, and findings should be reported with appropriate limitations.

### **Injury Site**

The body location where physical harm occurred or where its effects are being examined, documented,

---

treated, and monitored during recovery.

### **Ink Poisoning**

Illness after swallowing or significant exposure to ink; most small pen-ink exposures cause little toxicity, but symptoms depend on the product and amount.

### **Ink Remover Poisoning**

Poisoning caused by chemicals used to remove ink, which may irritate the skin, eyes, lungs, or digestive tract and may cause systemic toxicity.

### **Innate Immune Evasion**

Mechanisms by which pathogens evade the innate immune system. Strategies include capsules that prevent phagocytosis, biofilms that protect against immune cells, and intracellular survival within macrophages. Some bacteria produce enzymes that degrade complement components. Others modify their surface molecules to avoid recognition.

### **Innate Immunity**

The body's rapid defense against germs and tissue damage, present from birth. It includes skin, mucus, inflammation, cells that engulf germs, and protective proteins and usually responds within minutes or hours.

### **Innate Lymphoid Cells**

Immune cells that respond quickly to tissue damage or infection without recognizing one specific antigen. Different groups help control viruses, parasites, bacteria, fungi, allergic inflammation, and tissue repair.

### **Inner Ear**

The deepest part of the ear, containing the cochlea for hearing and balance organs for movement sensing. Disease there can cause hearing loss, ringing, dizziness, or balance problems.

### **Inner Ear Infection**

Inner-ear inflammation can affect hearing and balance, producing vertigo, nausea, tinnitus, or hearing loss. Causes include viral infection, bacterial dis-

ease, autoimmune inflammation, or other disorders; sudden hearing loss, severe imbalance, or neurologic signs may occur.

### **Innervation**

The nerve supply to an organ, muscle, or area of the body, including the functional control provided by those nerves.

### **Innominate**

Innominate means unnamed; in anatomy it may refer to structures such as the innominate artery or innominate bone, depending on context.

### **INOCA**

Ischemia with non-obstructive coronary arteries: objective myocardial ischemia or ischemic symptoms in a patient whose coronary arteries do not have an obstructive epicardial stenosis. Mechanisms include coronary microvascular dysfunction and vasospastic angina; evaluation should identify the mechanism so treatment can be tailored.

### **Inoculation**

Introduction of microorganisms, cells, or other material into a culture, living body, or normally sterile site. It may be deliberate for laboratory growth, vaccination, or treatment, or accidental and capable of causing infection.

### **Inoperable**

Not safely treatable by surgery, often because a disease is too extensive, located near vital structures, or the person is too medically fragile. Other treatments may still be possible.

### **Inorganic**

Not based mainly on carbon-containing molecular structures. Inorganic substances include water, salts, minerals, and metals, many of which are essential to body function while others can be toxic.

### **Inorganic Chemical**

A chemical compound that is generally not based on carbon chains, such as water, salts, minerals, or

---

metals. In the body, inorganic substances support fluid balance, nerve signals, bone structure, and many chemical reactions.

### Inosculation

The joining and establishment of a connection between blood vessels or vascularized tissues, important in wound healing, grafts, flaps, and transplantation.

### Inosine

A nucleoside made of hypoxanthine linked to ribose that participates in purine metabolism and cellular energy pathways.

### Inosine Monophosphate

A nucleotide containing inosine and one phosphate group; it is an intermediate in purine metabolism and a precursor of other nucleotides.

### Inosine Pranobex

A medicine combination marketed in some countries as an immunomodulator or antiviral; evidence and approved uses vary by jurisdiction.

### Inositol

A sugar-like compound involved in cell membranes and signaling; some forms are studied or used for selected metabolic, reproductive, or mental-health conditions.

### Inositol 1,4,5-Trisphosphate

A second messenger molecule generated when phospholipase C cleaves membrane phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>). It binds calcium-release channels on the endoplasmic reticulum and mobilizes intracellular calcium stores.

#### Alternate Names:

IP<sub>3</sub>; Ins(1,4,5)P<sub>3</sub>; Inositol Trisphosphate

### Inositol Phosphates

Phosphorylated forms of inositol that act in cell signaling, calcium regulation, and other intracellular processes.

### Inotropes In Anesthesia

Medicines used during anesthesia or critical illness to increase the force of heart contraction and support blood flow when the heart is failing. They are given by controlled infusion and require monitoring because they can cause abnormal heart rhythms, low or high blood pressure, and increased oxygen demand.

### Inotropic

Affecting the force of cardiac muscle contraction; a positive inotropic drug increases contractility, while a negative inotropic effect decreases it.

### Inotropic Agent

A pharmacological drug or physiological substance that influences the force or energy of myocardial muscular contractions. Positive inotropes (such as dobutamine, milrinone, and digoxin) enhance contractility to support cardiac output in cardiogenic shock or heart failure.

### Inpatient

An inpatient is a person admitted to a healthcare facility for monitoring, diagnosis, treatment, surgery, or care requiring an overnight or formal stay.

### Inpatient Rehabilitation

Rehabilitation provided while a person stays in a hospital or specialized facility. It combines physical, occupational, speech, nursing, and medical care to restore function after problems such as stroke, brain injury, spinal injury, or major illness.

### INR

A standardized blood-test result showing how quickly blood clots. It is used mainly to adjust warfarin and similar medicines; the desired range depends on why the medicine is prescribed, and too high or too low a result can be dangerous.

### Insecta

The biological class containing insects. Some insects bite or sting, cause allergic reactions, contaminate food, or spread infections and parasites; medical importance depends on the species and exposure.



---

### **Insect Bite**

A skin reaction caused when an insect feeds and leaves saliva or other material in the skin. It usually causes an itchy, red, raised spot, but swelling, infection, or a serious allergic reaction can occur.

### **Insect Bites**

Skin injuries caused by an insect feeding or injecting saliva or venom, producing local inflammation and sometimes allergic, infectious, or systemic reactions.

### **Insect Bites And Stings**

Bites and stings from insects such as mosquitoes, fleas, bedbugs, bees, wasps, hornets, yellow jackets, and ants can cause local pain, itching, swelling, allergic reactions, infection, or toxin effects; breathing difficulty or widespread symptoms may occur with a severe reaction.

### **Insecticide**

A chemical or biological agent used to kill or control insects. Exposure can cause skin, eye, respiratory, neurologic, or cholinergic toxicity depending on the product; prevention and treatment require identifying the active ingredient.

### **Insecticide Poisoning**

Poisoning whose presentation depends on the active ingredient. Organophosphates and carbamates cause cholinergic excess with secretions, miosis, bronchospasm, fasciculations, and weakness; pyrethroids can cause paresthesias, tremor, or seizures; other products have different toxicities. Identify the product and treat the agent-specific syndrome.

### **Insect Sting Allergies**

Immune reactions to venom from insects such as bees, wasps, hornets, and fire ants. Reactions may be local or systemic; generalized hives, breathing difficulty, dizziness, vomiting, or throat swelling may indicate anaphylaxis.

#### **Alternate Names:**

Hymenoptera Venom Allergy; Insect Venom Allergy

### **Insect Virus**

A virus that infects insects or other arthropods; some are studied for ecology or biological pest control rather than human disease.

### **Insemination**

Introduction of sperm into the female reproductive tract, either through intercourse or an assisted-reproduction procedure such as intrauterine insemination. Indications, success rates, infection screening, and risks depend on the method and patient factors.

### **Insensible Fluid Loss**

Obligatory water loss from the body that occurs continuously and imperceptibly through evaporation from the respiratory tract (exhaled vapor) and diffusion through the skin (transdermal evaporation without sweating), normally accounting for approximately 600 to 800 mL of fluid loss per day in adults.

### **Insertion**

Movement of a muscle attachment toward its fixed attachment, or placement of one structure into another.

### **Insidious**

Developing slowly and quietly, often without obvious early symptoms. An insidious illness may be present for a long time before it is recognized, so the term describes its gradual pattern rather than its cause or severity.

### **Insinuate**

To introduce or insert gradually into a space, or to suggest something indirectly; in anatomy, a structure may insinuate between or around other structures.

### **In Situ**

In its original place. In pathology, carcinoma in situ consists of abnormal or cancer-like cells that remain within the surface layer and have not invaded deeper tissue. It has not spread through blood or lymph vessels, but some lesions can become invasive if untreated.

---

**Alternate Names:**

In place

**In Situ Carcinoma**

Abnormal cells that remain within the tissue layer where they began and have not grown into deeper tissue or nearby blood or lymph vessels. Some can become invasive if untreated.

**In Situ Hybridization**

A laboratory test that uses marked probes to find a specific DNA or RNA sequence inside cells or tissue. It can detect some gene changes, viruses, or where a gene is active.

**Insomnia**

Alternate name; see *Insomnia Disorder*.

**Insomnia Disorder**

Difficulty initiating or maintaining sleep, or early-morning awakening with inability to return to sleep, at least three nights per week for at least three months. The individual experiences clinically significant distress or impairment in functioning.

**Alternate Names:**

Insomnia; Sleeplessness

**Inspection**

Systematic visual examination of a person or specimen for abnormalities such as changes in color, swelling, deformity, skin lesions, unusual movement, discharge, breathing effort, or overall appearance.

**Inspiratory Capacity**

Inspiratory capacity is the maximum volume inhaled after a normal exhalation and equals tidal volume plus inspiratory reserve volume.

**Inspiratory Crackles**

Brief clicking or popping sounds heard while breathing in. They may occur when small airways or air sacs open and can be caused by fluid, infection, scarring, lung collapse, or heart failure; the sound alone does not identify the cause.

**Inspiratory Reserve Volume**

The maximum additional volume of gas that can be voluntarily and forcefully inhaled into the lungs at the end of a normal, quiet tidal inspiration, typically measuring approximately 2.5 to 3.0 liters in healthy adults.

**Alternate Names:**

IRV

**Inspiratory Time**

The length of time spent breathing in during a natural breath or mechanical-ventilation cycle. Adjusting it can affect airflow, oxygen delivery, comfort, pressure, and the balance between inhaling and exhaling.

**Inspiratory-to-Expiratory Ratio**

The proportion of time dedicated to inspiratory phase versus expiratory phase during mechanical ventilation or spontaneous breathing, conventionally set at 1:2 in normal physiological states, and prolonged to 1:3 or 1:4 in obstructive lung diseases (asthma, COPD) to prevent dynamic hyperinflation (auto-PEEP).

**Alternate Names:**

I:E Ratio

**Instantaneous Wave-Free Ratio**

A measurement made during heart catheterization to judge whether a narrowed heart artery limits blood flow. A pressure wire compares pressure beyond and before the narrowing while the heart is relaxed. A result of 0.89 or less may support treatment when it fits the symptoms and other findings.

**Alternate Names:**

iFR; Wave-Free Ratio; Resting Coronary Physiology

**Instillation**

Slow placement of a liquid medicine or solution into a body cavity, organ, wound, or tissue, such as eye drops, bladder treatment, or fluid placed into a surgical space.

**Instrument**

A tool or device used by healthcare workers for exam-

---

ination, measurement, diagnosis, surgery, treatment, patient care, or laboratory work.

### **Instrumental Activities Of Daily Living**

Everyday tasks needed for independent living outside basic self-care, including preparing food, shopping, managing medicines and money, using transportation, communicating, and maintaining a home. Difficulty with these tasks may signal functional decline.

### **Instrumental ADL**

A complex activity needed for independent community living, such as shopping, cooking, managing medicines or finances, and using transportation.

### **Instrument Sterilization**

A validated process that destroys all forms of microbial life, including bacterial spores, on or within a medical instrument. Common methods include steam under pressure, ethylene oxide, hydrogen-peroxide plasma, and dry heat, selected according to device material and design.

### **Insufficient Cervix**

Painless shortening and opening of the cervix during pregnancy, usually in the second trimester, without labor contractions. It can cause pregnancy loss or early birth and may require close monitoring or treatment.

#### **Alternate Names:**

Cervical Insufficiency; Cervical Incompetence

### **Insula**

The insula is a cerebral cortical region involved in sensation, taste, emotion, interoception, pain, and integration of internal and external information.

### **Insular Cortex**

A pyramid-shaped cortical region situated deep within the lateral sulcus (Sylvian fissure), covered by the frontal, parietal, and temporal opercula, which coordinates interoceptive sensory awareness, visceral autonomic functions, pain perception, gustation, and emotional empathy.

#### **Alternate Names:**

Insula; Island of Reil

### **Insulin**

A hormone made by the pancreas that helps cells use glucose and lowers blood sugar. It is used to treat diabetes, but too much can cause dangerously low blood sugar with sweating, confusion, seizures, or unconsciousness.

### **Insulin Antagonist**

A hormone, medicine, or other factor that opposes insulin's effects and can raise blood glucose.

### **Insulin Aspart**

A rapid-acting insulin analog (modified amino acid sequence) with onset 10-20 minutes, peak 1-3 hours, duration 3-5 hours. Used to cover prandial glucose (injected at or just after meals) in type 1 and type 2 diabetes; also used in continuous subcutaneous insulin infusion (pump) and IV in acute settings.

### **Insulin Autoantibodies**

Autoantibodies directed against insulin or other pancreatic beta-cell antigens, including GAD, IA-2, and ZnT8 antibodies. Their presence supports autoimmune type 1 diabetes but a negative panel does not exclude it.

### **Insulin Coma**

A state of severe hypoglycemia and unconsciousness caused by excessive insulin; it is a medical emergency and is not a current routine treatment.

### **Insulin C-Peptide Test**

A blood or urine test measuring C-peptide released in equal amounts with endogenous insulin; it helps distinguish native insulin production from injected insulin and evaluate hypoglycemia or diabetes classification.

### **Insulin-Dependent Diabetes**

Type 1 diabetes, in which the pancreas produces little or no insulin and replacement insulin is required.

#### **Alternate Names:**

---

Type 1 diabetes

### **Insulin Detemir**

A long-acting insulin analogue used to provide basal insulin replacement in diabetes mellitus; dosing is individualized and hypoglycemia is the principal serious adverse effect.

### **Insulin Glargine**

A long-acting basal insulin analog with a flat, peakless profile providing near-24-hour coverage, administered once daily (or twice in some patients). Formed by modification of human insulin producing slow, steady absorption from the injection depot. Used for type 1 and type 2 diabetes (basal component).

### **Insulin-Induced Lipohypertrophy**

A common complication of subcutaneous insulin therapy characterized by localized accumulation of adipose tissue at injection sites, caused by the lipogenic effect of insulin on subcutaneous fat. It presents as firm, rubbery, non-tender subcutaneous nodules that can cause unpredictable insulin absorption and glycemic variability.

### **Insulin Injection Technique**

The method of injecting insulin into the fatty tissue under the skin. Common sites include the abdomen, thighs, buttocks, and upper arms; repeated injections in one place can cause fatty thickening and variable absorption.

### **Insulin Isophane**

Intermediate-acting human insulin, also called neutral protamine Hagedorn insulin, used for basal glucose control in diabetes mellitus.

### **Insulin-Like Growth Factor**

A hormone, mainly IGF-1, made largely by the liver in response to growth hormone. It supports growth of bone and other tissues and helps regulate metabolism. Abnormally high or low levels can occur with pituitary, liver, nutritional, or chronic illnesses.

### **Insulin-Like Growth Factor 1**

A 70-amino-acid peptide hormone synthesized primarily by hepatocytes under the transcriptional stimulation of growth hormone (GH), which mediates the systemic anabolic, mitogenic, and linear skeletal growth effects of GH; serves as the primary screening biomarker for acromegaly and gigantism.

#### **Alternate Names:**

IGF-1; Somatomedin C

### **Insulin Lipohypertrophy**

Thickened, rubbery fatty tissue that develops where insulin is injected repeatedly. Insulin absorbed from these areas may act unpredictably, so rotating injection sites helps prevent the problem and improves glucose control.

### **Insulin Lispro**

A rapid-acting insulin analogue used around meals or for correction of high blood glucose in diabetes mellitus.

### **Insulinoma**

An insulinoma is a usually benign pancreatic beta-cell neuroendocrine tumour that secretes excessive insulin, causing recurrent fasting or exertional hypoglycaemia relieved by glucose.

### **Insulin Pen Devices**

Pre-filled or refillable devices that deliver measured insulin through a small needle under the skin. They are portable, but the dose, injection technique, storage, and needle disposal must follow medical instructions.

### **Insulin Preparations**

Forms of insulin used to lower blood glucose by helping body cells take up glucose and reducing glucose release by the liver. They differ in how quickly they start and how long they work, so the type and dose must be individualized.

#### **Alternate Names:**

Insulins; Insulin Medicines

---

## Insulin Pumps

Insulin pumps deliver rapid-acting insulin continuously and through programmed meal or correction doses, allowing flexible management of diabetes while requiring monitoring and device safety.

## Insulin Pump Therapy

Delivery of rapid-acting insulin through a small tube under the skin. The pump gives a continuous background dose and extra doses around meals or to correct high blood glucose. Users need training, monitoring, and a backup plan if the device fails.

## Insulin Receptor

A protein on the surface of many cells that binds insulin and sends signals allowing cells to take up and use glucose. It also influences fat, protein, and cell-growth processes.

## Insulin Resistance

A state in which muscle, fat, and liver cells respond poorly to insulin. The pancreas may produce extra insulin at first, but blood glucose can eventually rise, increasing the risk of type 2 diabetes and metabolic syndrome.

### Alternate Names:

IR (Insulin Resistance)

## Insulin Resistance Syndrome

Alternate name; see *Metabolic Syndrome*.

## Insulin Resistance Syndromes

Rare inherited disorders in which the body responds poorly to insulin, sometimes because of a change in the insulin receptor. Severe forms can cause very high insulin levels, abnormal growth, diabetes, unusual body features, and serious illness early in life.

## Insulin Secretion

Release of insulin from pancreatic beta cells in response primarily to increased plasma glucose, with amplification by incretin hormones and nutrients. Insulin promotes glucose uptake, glycogen synthesis, lipogenesis, and protein synthesis while suppressing

hepatic glucose production.

## Insulin Signaling

The cellular process initiated when insulin binds its receptor and activates pathways that regulate glucose uptake, glycogen formation, protein synthesis, and fat metabolism.

## Insulin Storage and Syringe Safety

Clinical safety guidelines for the handling, temperature preservation, and administration of insulin. Includes refrigeration (2°C to 8°C) of unopened vials, room-temperature storage of in-use pens or vials for up to 28–42 days, single-use needle and syringe protocols, and prompt disposal in puncture-resistant biohazard sharps containers.

### Alternate Names:

Insulin Administration Safety; Diabetes Sharps Disposal; Enteral Insulin Handling

## Insulin Therapy in Diabetes

Treatment with subcutaneous injections or continuous pump infusions of insulin to control blood glucose levels in type 1 diabetes, advanced type 2 diabetes, or gestational diabetes.

### Alternate Names:

Insulin Administration; Exogenous Insulin Therapy; Diabetes Insulin Management

## Insulin Tolerance Test

A specialist test in which insulin is given through a vein to lower blood sugar under close supervision. It measures how the body releases stress hormones such as cortisol and growth hormone. Because very low blood sugar can be dangerous, the test requires careful monitoring.

### Alternate Names:

ITT; Insulin-Induced Hypoglycemia Test

## Integrase

Integrase is an enzyme that inserts viral DNA into a host genome, a process targeted by integrase inhibitors used in HIV treatment.

---

## **Integrative Medicine**

Patient-centered care that combines evidence-based conventional treatment with appropriate complementary approaches, emphasizing whole-person health, prevention, lifestyle, and shared decision-making.

## **Integrative Rehabilitation**

A rehabilitation plan that combines standard physical, occupational, or speech therapy with selected complementary approaches, such as relaxation or mindfulness, when they are safe and supported by evidence.

## **Integrin**

A cell-surface receptor that links cells to the extracellular matrix and helps control adhesion, movement, signaling, and immune-cell activity.

## **Integrin Binding**

The attachment of a molecule or cell to an integrin receptor, which can influence adhesion, migration, signaling, or clot formation.

## **Integron**

A genetic element containing an integrase and an attachment site that can capture and express gene cassettes; integrons commonly contribute to bacterial antimicrobial resistance.

## **Integrons**

Genetic elements that capture and express gene cassettes through site-specific recombination. Integrons contain an integrase gene and a recombination site for gene cassette insertion. They are important vehicles for the spread of antibiotic resistance genes, particularly in gram-negative bacteria.

## **Integumentary System**

The body system comprising the skin and its appendages: hair, nails, sweat glands, and sebaceous glands. It forms a protective barrier, limits fluid loss, contributes to thermoregulation and sensation, supports immune defence, and participates in vitamin D synthesis.

## **Intein**

An intein is a protein segment that excises itself from a precursor protein while joining the surrounding extein sequences.

## **Intellectual Disability**

A developmental condition involving significant difficulty with learning, reasoning, communication, and everyday independent skills. It begins during childhood and varies widely in severity; support needs depend on the person's abilities and environment.

## **Intelligence**

Intelligence is the capacity to learn, reason, solve problems, understand information, and adapt; tests measure selected abilities rather than a person's entire worth or potential.

## **Intelligence Test**

An intelligence test uses standardized tasks to estimate cognitive abilities such as reasoning, memory, language, and problem-solving.

## **Intense Pulsed Light**

A treatment using brief pulses of broad-spectrum light for selected skin conditions and hair reduction. Light is absorbed by pigment or blood vessels, and treatment can cause temporary redness, burns, pigment changes, or eye injury.

### **Alternate Names:**

IPL

## **Intensity**

The degree of effort or physiological demand during physical activity.

## **Intensity-Modulated Radiation Therapy**

A type of radiation treatment that varies the strength of many small radiation beams to match the shape of a tumor. It aims to treat the tumor while reducing radiation to nearby healthy tissue.

## **Intensive Care**

Specialized treatment for people who are seriously ill or at risk of organ failure. It provides continuous



---

monitoring, medicines, procedures, and life support such as a ventilator when needed.

### **Intensive Care Rehabilitation**

Rehabilitation begun during or soon after intensive care to reduce weakness and loss of function. It may include safe positioning, movement, breathing exercises, and help with communication or daily activities.

### **Intensive Care Unit**

A hospital unit for people who are critically ill or at high risk of organ failure. Staff provide continuous monitoring, urgent treatments, and support such as a ventilator, dialysis, or medicines to maintain blood pressure.

### **Intentional Obstetrical Clavicle Fracture**

A deliberate fracture of a baby's collarbone during an emergency delivery when the shoulders are stuck and other measures have failed. It creates more room for birth and usually heals, but is used only as a last-resort obstetric procedure.

#### **Alternate Names:**

Cleidotomy; Obstetric Clavicular Fracture

### **Intention Tremor**

A tremor that becomes more pronounced during a purposeful movement toward a target. It commonly suggests cerebellar or cerebellar-pathway dysfunction, although medicines, toxins, and other neurologic disorders can contribute.

### **Interatrial Septum**

The wall between the heart's right and left upper chambers. It includes the fossa ovalis, a normal remnant of fetal circulation, and contains a thinner area that may be crossed during selected catheter procedures.

### **Intercalated Disc**

An intercalated disc is a specialized junction connecting cardiac muscle cells and transmitting electrical and mechanical force through the heart.

### **Intercalating Agent**

An intercalating agent inserts between DNA base pairs and can disrupt replication or transcription; some are laboratory reagents or anticancer drugs.

### **Intercellular**

Located between cells or occurring in the space between cells. Intercellular material and signals help cells attach, communicate, exchange substances, and organize tissues throughout the body.

### **Intercostal**

Intercostal means located between the ribs, especially the muscles, nerves, and blood vessels in those spaces.

### **Intercostal Muscles**

Muscles between the ribs that help expand and compress the chest during breathing and also assist movement of the trunk.

### **Intercostal Retraction**

Inward movement of the skin and soft tissues between the ribs during inspiration, indicating increased work of breathing. It is common in infants and children with respiratory distress and may accompany airway obstruction, bronchiolitis, pneumonia, asthma, or other pulmonary disease.

### **Intercostal Retractions**

Inward pulling of the skin and soft tissues between the ribs during inspiration, indicating increased work of breathing and potentially significant respiratory distress.

### **Intercostal Space**

The space between two neighboring ribs. It contains intercostal muscles, nerves, and blood vessels.

### **Intercurrent Disease**

An illness or complication that develops during the course of another disease or treatment and may affect its management or outcome.

---

### Interdental Brush

A small brush designed to clean between teeth where a regular toothbrush cannot reach. It helps remove plaque and food and may reduce gum inflammation; the correct size should fit without force.

### Interdigital Neuroma

Alternate name; see *Morton's Neuroma*.

### Interference

A factor that alters the accuracy or interpretation of a laboratory test, measurement, or drug effect.

### Interferometry

A measurement method that uses interference between waves to determine distance, motion, thickness, optical properties, or other physical quantities.

### Interferon

Interferons are cytokines that induce antiviral defenses, alter immune-cell activity, and regulate cell growth; type I, II, and III interferons have different receptors and clinical uses.

### Interferon-Alpha

A type I interferon cytokine with antiviral, antiproliferative, and immune-modulating effects; recombinant forms have been used for selected viral infections and cancers.

### Interferon-Beta

A type I interferon cytokine used mainly as an immunomodulatory treatment for relapsing forms of multiple sclerosis.

### Interferon Gamma Release Assay

An in vitro diagnostic blood test measuring the release of interferon-gamma by memory T-lymphocytes upon exposure to specific *Mycobacterium tuberculosis* antigens (ESAT-6 and CFP-10), unaffected by prior BCG vaccination, utilized for latent tuberculosis infection screening.

#### Alternate Names:

IGRA; QuantiFERON-TB

### Interferon-Gamma Release Assay

A blood test that looks for immune evidence of infection with the bacterium that causes tuberculosis. It can support diagnosis of latent infection but cannot alone distinguish latent infection from active disease.

#### Alternate Names:

IGRA; QuantiFERON-TB; T-SPOT.TB

### Interferon Inducer

A substance that stimulates cells to produce interferons, proteins involved in antiviral and immune responses.

### Interferonopathies

Rare inherited or acquired disorders in which interferon-driven immune activity remains abnormally high. They can cause ongoing inflammation, skin or lung disease, blood-vessel injury, or neurologic problems, often beginning in childhood.

### Interferons

Natural proteins that cells release to warn nearby cells and help control the immune response. They are especially important in the body's defense against viruses and are also used as medicines for some infections, cancers, and immune conditions.

### Interferon Signaling

The process by which interferons bind to cell receptors and activate genes that help defend against viruses, regulate immune cells, and sometimes slow cell growth. Different interferons act mainly on different tissues.

### Interleukin

A family of signaling proteins made mainly by immune cells. Interleukins help immune cells grow, communicate, and control inflammation, fever, allergy, infection defense, and tissue repair; different interleukins have different effects.

### Interleukin 1

A group of proinflammatory cytokines, especially IL-1 alpha and IL-1 beta, that promote fever, leukocyte

---

activation, and inflammatory responses.

### **Interleukin 13**

A cytokine produced by type 2 helper T cells and other immune cells that contributes to allergic inflammation, mucus production, and tissue fibrosis.

### **Interleukin 2**

Interleukin-2 is a T-cell growth factor that promotes proliferation and survival of activated T cells and natural-killer cells; high-dose therapy can cause capillary-leak syndrome and systemic toxicity.

### **Interleukin-6**

A cytokine involved in immune regulation and inflammation that stimulates production of acute-phase proteins by the liver.

### **Interleukin 7**

A cytokine important for development, survival, and homeostasis of T lymphocytes and some B-cell precursors.

### **Interleukins**

Cytokines that carry signals between immune cells and regulate inflammation.

**Alternate Names:**

ILs

### **INTERMACS Profile**

A seven-level system for describing how sick a person is with advanced heart failure. Profile 1 means critical shock with very poor blood flow; profiles 2 and 3 describe worsening or dependence on medicines that strengthen the heart. Profiles 4 through 7 describe repeated hospital stays or increasing limits on activity. The profile helps doctors judge urgency and timing for a heart pump or transplant.

**Alternate Names:**

INTERMACS Staging; INTERMACS Scale

### **Intermediate**

Intermediate means between two stages, positions, or levels, such as an intermediate metabolite, cell, or developmental phase.

### **Intermediate Cuneiform**

The smallest of three cuneiform bones in the middle of the foot. It lies between the other cuneiform bones and joins the navicular bone and second metatarsal, helping support the foot's arch.

### **Intermediate-Density Lipoprotein**

A temporary particle formed when the body removes some fat from very-low-density lipoprotein. It can be changed into LDL, the type of cholesterol particle that can contribute to fatty buildup in artery walls.

**Alternate Names:**

IDL

### **Intermediate Filament**

An intermediate filament is a strong cytoskeletal fiber that supports cell shape, mechanical stability, and tissue integrity.

### **Intermediate Uveitis**

Inflammation mainly affecting the gel inside the eye and nearby tissues at the edge of the retina. It may cause blurred vision, floaters, or little pain and can be linked to autoimmune disease, infection, or no identified cause. Eye examination and imaging guide treatment and monitoring for complications.

### **Intermittent Auscultation**

Checking a baby's heart rate at regular intervals during labor with a handheld Doppler or fetoscope rather than recording it continuously. The timing depends on the stage of labor, pregnancy risks, symptoms, and local clinical practice.

### **Intermittent Claudication**

Cramping, aching, or weakness in a leg muscle that reliably starts with walking or exercise and improves with rest. It usually results from narrowed leg arteries and may indicate peripheral artery disease.

### **Intermittent Explosive Disorder**

A mental-health disorder causing repeated, sudden outbursts of verbal or physical aggression that are far greater than the situation warrants. The outbursts are impulsive, not planned, and cause distress

---

or problems in daily life; diagnosis is made from the pattern and history.

### **Intermittent Fasting**

An eating pattern that alternates planned periods of eating with periods of little or no food. Common schedules limit daily eating hours or use selected fasting days. It may help some adults lose weight, but it is not suitable for everyone, especially during pregnancy or with some medical conditions or eating disorders.

### **Intermittent Fever**

A fever that rises and falls with periods when temperature returns to normal or near normal.

### **Intermittent Mandatory Ventilation**

A mechanical ventilation mode in which preset mechanical volume- or pressure-cycled mandatory breaths are delivered at fixed periodic intervals, while permitting spontaneous unassisted or pressure-supported breaths between mandatory cycles.

#### **Alternate Names:**

IMV; Synchronized Intermittent Mandatory Ventilation; SIMV

### **Intern**

A physician in the early stage of postgraduate medical training, traditionally the first year after medical school; the exact training structure varies by country.

### **Internal**

Internal means located within the body, an organ, a structure, or a system rather than on its outer surface.

### **Internal Bleeding**

Internal bleeding is blood loss into a body cavity, tissue, or organ rather than through an obvious external wound. Trauma, ulcers, aneurysm, pregnancy complications, medicines, and tumors can cause it; faintness, shock, abdominal swelling, black stool, or vomiting blood may occur.

### **Internal Capsule**

A compact bundle of nerve fibers deep inside the brain that carries movement and sensation signals between the brain's outer layer and lower brain and spinal centers. Damage can cause weakness, numbness, or other neurologic deficits.

### **Internal Capsule Stroke**

A classic lacunar infarction or intracerebral hemorrhage involving the posterior limb of the internal capsule, causing pure motor hemiparesis characterized by equal motor weakness of the contralateral face, arm, and leg without sensory deficits, visual field loss, or higher cortical signs.

#### **Alternate Names:**

Posterior Limb Lacunar Infarct; Pure Motor Stroke

### **Internal Carotid Artery**

A branch of the common carotid artery that enters the skull and supplies the brain, eyes, and related structures. It contributes to the cerebral arterial circulation.

### **Internal Carotid Artery Dissection**

A tear in the tunica intima of the extracranial internal carotid artery leading to intramural hematoma formation within the arterial wall, manifesting classically with ipsilateral frontotemporal or retro-orbital headache, neck pain, partial Horner syndrome (ptosis and miosis without anhidrosis), and ischemic stroke in young adults.

#### **Alternate Names:**

ICAD

### **Internal Environment**

The fluid and physiologic conditions inside the body that cells require, including temperature, pH, oxygen, electrolytes, and nutrients.

### **Internal Fixation**

Surgical stabilization of a fracture or unstable bone using implanted hardware such as plates, screws, rods, wires, or pins.

---

### Internal Fixator

An internal fixator is a surgically implanted plate, screw, rod, nail, or frame used to stabilize a fracture or correct skeletal alignment.

### Internal Hernia

A loop of intestine slips through an opening inside the abdominal cavity, often after abdominal surgery. It can cause intermittent or sudden bowel obstruction with abdominal pain, vomiting, and bloating.

### Internal Iliac Artery

A major artery branching from the pelvic arteries. It supplies blood to the pelvic organs, buttocks, pelvic muscles, reproductive organs, bladder, and parts of the rectum.

### Internal Iliac Vein

A vein draining the pelvic organs, pelvic wall, and gluteal region that joins the external iliac vein to form the common iliac vein.

### Internal Jugular Vein

A large vein in the neck that drains blood from the brain, face, and neck back to the heart. It joins another major vein behind the collarbone and is sometimes used for central intravenous access.

### Internal Jugular Vein Cannulation

The percutaneous placement of an indwelling central venous catheter into the internal jugular vein, conventionally performed under real-time ultrasound guidance between the sternal and clavicular heads of the sternocleidomastoid muscle for central venous pressure monitoring, inotropic drug administration, or hemodialysis access.

#### Alternate Names:

IJ Line Placement

### Internal Jugular Vein Thrombosis

Thrombotic occlusion of the internal jugular vein, which may arise secondary to prolonged central venous catheterization, hypercoagulable states, deep neck space infection (as in Lemierre syndrome following oropharyngeal sepsis), local head and neck

malignancy, or intravenous drug use. Clinically manifests with anterior cervical pain, swelling, palpable cord sign, and risk of septic pulmonary embolism.

#### Alternate Names:

IJV Thrombosis; Jugular Vein Phlebothrombosis

### Internal Oblique Muscle

A broad muscle layer in the side of the abdominal wall. It helps compress the abdomen, bend and rotate the trunk, support the spine, and strengthen the groin and abdominal wall.

### Internal Podalic Version

A rare obstetric procedure in which a clinician turns a baby inside the uterus by bringing the feet downward, usually to deliver a second twin in an unusual position. It carries serious risks and is performed only in selected emergencies by an experienced specialist.

### Internal Resorption

Destruction of tooth tissue from the inside of the tooth outward, caused by activity of specialized cells. It may produce a pink area or no symptoms and can weaken the tooth; dental imaging determines whether root-canal treatment can save it.

### International Classification of Diseases

A World Health Organization system for coding diseases, injuries, and other health conditions. Versions such as ICD-10 and ICD-11 support clinical records, statistics, and billing; a code does not replace a clinical diagnosis.

### International Normalized Ratio

The INR, a standardized calculation from the prothrombin time used to assess blood clotting and guide treatment with vitamin K antagonists such as warfarin.

### International Unit

A unit that measures the biological effect of a substance, such as insulin, a vitamin, or a vaccine. The amount represented by one unit differs between substances, so international units cannot be directly converted to milligrams without knowing the sub-

---

stance.

### **Interneuron**

An interneuron connects neurons within the central nervous system and helps integrate, inhibit, or coordinate neural signals.

### **Interneurons**

Neurons within the central nervous system that connect other neurons and help process or coordinate neural signals.

### **Internist**

An internist is a physician specializing in adult internal medicine, including diagnosis and management of complex medical conditions.

### **Internuclear Ophthalmoplegia**

A disorder of coordinated side-to-side eye movement caused by damage to a nerve pathway in the brainstem. One eye may not move inward properly, causing double vision or jerky eye movements; stroke and multiple sclerosis are common causes.

### **Interosseous**

Interosseous means located between bones, such as the membrane or muscles between the forearm or lower-leg bones.

### **Interosseous Nerve Syndrome**

A compressive peripheral neuropathy of either the anterior interosseous nerve (causing weakness of the flexor pollicis longus and flexor digitorum profundus, preventing the 'OK' sign) or the posterior interosseous nerve (causing finger and thumb extension weakness with intact wrist extension and preserved cutaneous sensation).

#### **Alternate Names:**

Kiloh-Nevin Syndrome; AIN / PIN Palsy

### **Interpersonal Psychotherapy**

A time-limited talking therapy, used especially for depression, that links mood symptoms with relationships and life changes. It helps address grief, conflict, role changes, loneliness, communication,

and social support.

### **Interphalangeal Joint**

A joint between the bones of a finger or toe. These joints mainly allow the digit to bend and straighten.

### **Interphalangeal Dislocation**

Displacement of a finger or toe joint between phalanges, usually after hyperextension, twisting, or direct impact. Pain, deformity, swelling, and restricted motion occur; fractures, tendon injury, and neurovascular compromise may accompany it.

### **Interphase**

Interphase is the portion of the cell cycle between divisions, including growth, DNA replication, and preparation for mitosis or meiosis.

### **Interphase Cell**

A cell between divisions that grows, performs normal functions, and copies its DNA before possible division.

### **Interproximal Caries**

Dental caries occurring on the contact surfaces between adjacent teeth. It may be difficult to detect clinically and is commonly assessed with bitewing radiographs when imaging is indicated.

### **Interrupted Aortic Arch**

A rare congenital heart defect in which the aorta has a complete gap between its ascending and descending parts. Newborns can rapidly develop poor circulation and heart failure.

### **Intersexual**

Older term; see *Difference of Sex Development*.

### **Interstitial**

Relating to the spaces between cells or tissues, such as interstitial fluid or interstitial lung disease.

### **Interstitial Cystitis**

A long-lasting condition, also called bladder pain syndrome, that causes pain, pressure, or tenderness



---

felt in the bladder or pelvis together with urinary symptoms. Pain often worsens as the bladder fills and improves after urination; frequent urination and a sudden urge to urinate are common, and sex may be painful. Symptoms may flare and then improve. There is no single known cause. The diagnosis is based on the symptoms and examination while other conditions with similar symptoms, such as a urinary infection, are considered or ruled out.

**Alternate Names:**

IC; Bladder Pain Syndrome; Painful Bladder Syndrome

**Interstitial Fluid**

Interstitial fluid is the fluid surrounding tissue cells, formed from blood plasma and returned to circulation through lymphatic vessels.

**Interstitial Granulomatous Dermatitis**

A rare inflammatory skin disorder that causes red, purple, or brown patches or raised areas, often on both sides of the body. It may be linked to autoimmune disease, medicines, or another illness.

**Interstitial Keratitis**

Inflammation of the corneal stroma without primary involvement of the corneal surface. It may follow infection, autoimmune disease, or trauma and can cause pain, photophobia, redness, and reduced vision.

**Interstitial Lung Disease**

A group of lung conditions that affect the tissue around the air sacs. They can cause inflammation, scarring, or both, making it harder for the lungs to expand and oxygen to reach the blood. Symptoms and how the condition changes over time vary by type, but may include shortness of breath, a dry cough, and low oxygen.

**Alternate Names:**

ILD; Diffuse Parenchymal Lung Disease (DPLD)

**Interstitial Nephritis**

Inflammation of the renal interstitium and tubules, which can be acute or chronic. Acute interstitial nephritis is most commonly caused by medi-

cations, particularly NSAIDs, antibiotics including penicillins and cephalosporins, and proton pump inhibitors.

**Interstitial (Nonidiopathic) Pulmonary Fibrosis**

Fibrotic interstitial lung disease with an identifiable or suspected cause, such as connective-tissue disease, occupational exposure, medicines, radiation, or chronic hypersensitivity pneumonitis. Management combines exposure control, treatment of the cause, monitoring, and selected anti-inflammatory or antifibrotic therapy.

**Interstitial Pneumonia**

Inflammation affecting the supporting tissue around the lung air sacs. It may be caused by infection, autoimmune disease, medicines, or other lung disorders and can cause dry cough, fever, breathlessness, and low oxygen. Examination and imaging help determine the cause and treatment.

**Alternate Names:**

Idiopathic Interstitial Pneumonia; Interstitial Pneumonitis

**Interstitial Pulmonary Fibrosis**

Scarring in the supporting tissue between the lung air sacs, which can reduce oxygen transfer. It may cause progressive breathlessness and dry cough; causes include autoimmune disease, medicines, radiation, or no identified cause.

**Alternate Names:**

Idiopathic Pulmonary Fibrosis (IPF); Interstitial Pneumonitis

**Alternate Names:**

Diffuse Parenchymal Lung Disease; Lung Fibrosis

**Interstitium**

The supporting tissue between and around the lung air sacs, capillaries, and small airways; inflammation or fibrosis of this tissue can impair lung expansion and gas exchange.

**Intertrigo**

Intertrigo is inflammation in opposing skin folds caused by moisture, heat, friction, and maceration;

---

Candida or bacterial infection may complicate the tender erythematous rash.

**Alternate Names:**

Intertriginous Dermatitis

**Intertrochanteric Fracture**

An extracapsular fracture of the proximal femur occurring along the dense trabecular line extending between the greater and lesser trochanters, carrying a low risk of avascular femoral head necrosis compared to femoral neck fractures, commonly treated with a cephalomedullary nail or sliding hip screw.

**Interval**

An interval is the time, distance, or range between two points or events, such as a dosing or follow-up interval.

**Intervening Sequence**

A nucleotide sequence within a gene transcript that lies between coding sequences; in many genes it is an intron removed during RNA processing.

**Intervention**

An action intended to prevent, diagnose, treat, or change a health condition or outcome, such as a medicine, procedure, counseling, or public-health measure.

**Alternate Names:**

Medical Intervention; Clinical Intervention

**Interventional**

Relating to a procedure or active treatment used to diagnose or manage disease, often through minimally invasive access rather than open surgery.

**Interventional Cardiology**

A subspecialty using catheter-based procedures to diagnose and treat coronary, structural, and selected vascular heart conditions, including angioplasty, stenting, and transcatheter valve procedures.

**Interventional Procedure**

A targeted procedure used to diagnose or treat disease, often through a needle, tube, or small inci-

sion and guided by imaging. Examples include biopsy, drainage, vessel treatment, and tumor ablation; risks depend on the body area and technique.

**Interventional Procedures**

Procedures that diagnose or treat disease through a needle, catheter, or small incision, often using imaging for guidance. Examples include biopsy, drainage, vessel opening, embolization, and ablation; less invasive does not mean risk-free.

**Interventional Radiologist**

A physician trained to perform minimally invasive diagnostic and therapeutic procedures using medical imaging for guidance.

**Interventional Radiology**

A specialty using imaging guidance to perform minimally invasive diagnostic and therapeutic procedures, such as biopsies, drainages, embolization, and vascular treatments.

**Interventricular Septum**

The wall between the heart's two lower chambers. It is mostly muscle and helps the heart pump effectively; it also contains part of the electrical system that coordinates the heartbeat.

**Intervertebral Disc**

A strong, flexible cushion between neighboring spinal bones. It absorbs shock and allows limited movement; a bulging or damaged disc can irritate nearby nerves and cause back or neck pain, numbness, or weakness.

**Intervertebral Disk**

A strong, flexible cushion between neighboring spinal bones that absorbs shock and allows movement. A damaged or herniated disk can press on nearby nerves and cause back pain, numbness, tingling, or weakness.

**Alternate Names:**

Intervertebral disc

---

## Intervertebral Foramen

An opening between two neighboring bones of the spine through which a spinal nerve leaves the spinal canal. Narrowing of this opening can press on the nerve and cause pain, numbness, tingling, or weakness in its area of the body.

### Alternate Names:

Neural foramen

## Intestinal Absorption

Movement of digested nutrients, water, vitamins, and salts through the intestinal lining into the blood or lymph. Damage to the intestine, pancreas, liver, or bile flow can reduce absorption and cause deficiency.

## Intestinal Angina

A clinical manifestation of chronic mesenteric ischemia characterized by severe, cramping, periumbilical or epigastric abdominal pain developing 30 to 60 minutes after eating ('food fear'), leading to progressive weight loss, caused by severe atherosclerotic stenosis of two or more mesenteric arteries.

### Alternate Names:

Chronic Mesenteric Ischemia; Postprandial Abdominal Angina

## Intestinal Atresia

A birth defect in which part of the intestine is closed or missing, blocking passage of milk and digestive contents. Newborns may develop swollen abdomen, vomiting, and failure to feed and usually need surgery.

## Intestinal Cancer

A cancer that begins in the small or large intestine. It may cause pain, bleeding, anemia, altered bowel habits, blockage, or weight loss, but early disease may cause no symptoms; diagnosis and treatment depend on its site and stage.

## Intestinal Carcinoid Tumor

A neuroendocrine tumor arising in the intestine, especially the ileum or jejunum. It may be incidental or cause pain, obstruction, bleeding, or hormone-related carcinoid syndrome after liver or

other metastases develop.

## Intestinal Crypts

Small glands in the intestinal lining that make mucus and help replace worn-out lining cells.

## Intestinal Diseases

Diseases of the small or large intestine include infection, inflammation, poor absorption, blockage, polyps, and cancer. Symptoms may include pain, diarrhea, constipation, bleeding, vomiting, weight loss, or nutritional deficiency.

## Intestinal Duplication

A birth difference in which an extra segment or cyst develops beside or within the intestine. It may cause no symptoms or lead to pain, bleeding, infection, blockage, or twisting of the bowel.

## Intestinal Failure

Severe loss of intestinal absorptive function, so the body cannot obtain enough nutrients, water, or electrolytes from food and usually needs intravenous nutrition or fluids to maintain health.

## Intestinal Fistula

An abnormal passage connecting the intestine with another organ, another bowel segment, or the skin. It may develop after inflammation, infection, surgery, or injury and can cause leakage, infection, dehydration, or poor nutrition.

## Intestinal Flukes

Parasitic flatworms that infect the intestine after contaminated food or water is consumed. Infection may cause abdominal pain, diarrhea, malabsorption, or nutritional problems, and diagnosis depends on exposure history and stool or other testing.

## Intestinal Ganglioneuromatosis

A rare, usually noncancerous overgrowth of nerve cells and supporting fibers in the intestinal wall. It may cause pain, diarrhea, constipation, or blockage and can occur with certain inherited tumor syndromes.

---

## Intestinal Gas

Gas within the digestive tract that may cause belching, bloating, abdominal discomfort, passing wind, or symptoms from an underlying digestive disorder.

## Intestinal Ischemia

Too little blood flow reaching the intestines. It may develop suddenly from a clot or blockage, or gradually from narrowed arteries, and can cause severe pain, bloody stools, or bowel damage.

### Alternate Names:

Mesenteric Ischemia; Bowel Ischemia

## Intestinal Leiomyoma

A usually benign smooth-muscle tumor arising in the wall of the intestine; symptoms depend on size and location and may include bleeding, obstruction, or abdominal pain.

## Intestinal Lipodystrophy

Alternate name; see *Whipple's Disease*.

## Intestinal Lipomatosis

A usually noncancerous overgrowth of fatty tissue in the intestinal wall. It often causes no symptoms, but larger areas can lead to bleeding, bowel blockage, or telescoping of one bowel segment into another.

## Intestinal Lymphangiectasia

Dilation or obstruction of intestinal lymphatic vessels, causing leakage of lymph into the bowel and loss of protein, lymphocytes, and fat-soluble vitamins. It may produce edema, diarrhea, abdominal symptoms, or poor growth.

## Intestinal Lymphoma

A cancer of immune-system cells that begins in the small or large intestine. It may cause abdominal pain, weight loss, diarrhea, bleeding, or blockage; diagnosis requires tissue sampling and specialist evaluation.

## Intestinal Malrotation

A birth defect in which the intestines do not move into their usual position before birth. It can allow

the bowel to twist and lose its blood supply, causing green vomiting, pain, swelling, or sudden illness in an infant.

## Intestinal Metaplasia

A change in which the normal lining of the stomach or another area becomes more like the lining of the intestine, usually after long-term irritation or inflammation. Some forms increase the future risk of stomach cancer.

## Intestinal Microbiome

The community of microorganisms living in the intestines. These organisms can affect digestion, vitamin production, immune activity, and protection from harmful germs.

## Intestinal Motility

The muscle movements that mix and push food, fluid, and stool through the intestines.

## Intestinal Mucosa

The inner lining of the intestine, where digestion and nutrient absorption occur. It produces mucus and contains immune defenses that help protect the body from germs and harmful substances in food.

## Intestinal Obstruction

A partial or complete blockage that prevents food, fluid, and gas from moving normally through the intestine. Adhesions, hernias, tumors, inflammation, twisting, or poor bowel movement can cause pain, swelling, vomiting, and inability to pass stool or gas.

## Intestinal Obstruction And Ileus

Impaired passage of intestinal contents caused by a mechanical blockage or by reduced bowel motility without a physical blockage; symptoms commonly include abdominal pain, distension, vomiting, and inability to pass stool or gas.

## Intestinal Obstruction Repair

Surgery or another intervention to relieve an intestinal blockage, remove a diseased segment when

---

necessary, and restore bowel continuity or create a stoma when required.

### Intestinal Parasite

A protozoan or worm that lives in the gastrointestinal tract and may cause diarrhea, abdominal pain, anemia, malnutrition, or no symptoms.

### Intestinal Perforation

A hole through the bowel wall that allows digestive contents and bacteria into the abdomen. It may result from inflammation, infection, blockage, injury, or surgery and can cause severe pain, abdominal rigidity, infection, and shock.

### Intestinal Polyp

A growth that projects from the inner lining of the small or large intestine. Many are harmless, but some can slowly become cancer, so their size, number, type, and location guide follow-up or removal.

### Intestinal Polyposis

The presence of many polyps in the intestine. It may be inherited or acquired and can cause bleeding, anemia, blockage, or increased cancer risk; evaluation often includes colonoscopy and family-history assessment.

### Intestinal Pseudo-Obstruction

Bowel swelling and symptoms of blockage caused by poor intestinal movement despite no physical obstruction. It may be temporary after illness or surgery or result from a long-term nerve or muscle disorder.

#### Alternate Names:

Ogilvie Syndrome; Acute Colonic Pseudo-Obstruction

### Intestinal Resection

Alternate name; see *Bowel Resection*.

### Intestinal Villi

Tiny finger-like projections lining the small intestine that greatly increase its surface area. Their blood and lymph vessels absorb digested nutrients; dam-

age to them can cause diarrhea, weight loss, and nutritional deficiencies.

### Intestinal Wall

The layers of tissue that surround the intestinal passage. They move intestinal contents, absorb nutrients and water, and help protect against harmful substances.

### Intestine

The part of the digestive tract extending from the stomach to the anus, consisting of the small intestine and large intestine.

### Intestine Disorder

Any disease affecting the small or large intestine, including infection, inflammation, poor absorption, blockage, polyps, or cancer. Symptoms may include pain, diarrhea, constipation, bleeding, vomiting, weight loss, or nutritional deficiency.

### Intestine Neoplasm

An abnormal growth in the small or large intestine. It may be noncancerous or cancerous and can cause bleeding, anemia, pain, altered bowel movements, blockage, or no symptoms.

### Intestine Obstruction

Intestinal obstruction is partial or complete blockage of the passage of intestinal contents, causing pain, swelling, vomiting, and failure to pass stool or gas.

### Intimal Dissection

A tear in the inner lining of an artery that allows blood to enter and split the vessel wall, creating a false lumen. Aortic or other arterial dissection can compromise blood flow and is a medical emergency.

### Intimal Hyperplasia

Thickening of the inner lining of an artery after injury, surgery, a stent, or a vascular graft. The narrowing can reduce blood flow and cause a treated artery or dialysis access to become blocked again.

---

### **Intimate Partner Violence In Pregnancy**

Physical, sexual, or psychological abuse by a current or former partner during pregnancy. It can harm the pregnant person and fetus and is associated with complications such as injury, preterm birth, and placental abruption.

### **Intolerance**

An adverse response to a substance that does not involve the immune mechanism of an allergy.

#### **Alternate Names:**

Nonallergic Sensitivity; Food Intolerance

### **Intoxication**

A state of physical or mental impairment caused by recent exposure to a substance, such as alcohol, a medicine, or another drug.

### **Intra-Abdominal Abscess**

A pocket of pus inside the abdomen, often near the bowel, pelvis, or diaphragm. It can cause fever, abdominal pain, vomiting, or feeling unwell and may need antibiotics plus drainage or surgery.

### **Intraabdominal Cancer**

Cancer arising in an organ or tissue within the abdomen, such as the bowel, stomach, liver, pancreas, ovaries, or abdominal lining. Symptoms and treatment depend on the organ, cancer type, extent, and overall health.

### **Intraabdominal Hemorrhage**

Bleeding inside the abdominal cavity, often after trauma, surgery, rupture of an organ or blood vessel, pregnancy complications, or internal disease. It may cause pain, swelling, dizziness, or shock.

### **Intraabdominal Hernia**

A loop of intestine or other abdominal tissue slips through an abnormal opening inside the abdominal cavity rather than through the abdominal wall. It can cause pain, vomiting, and bowel obstruction.

### **Intra-Abdominal Hypertension**

A sustained or repeated pathological elevation of

intra-abdominal pressure equal to or exceeding 12 mmHg (measured via transvesical bladder pressure catheterization), which when exceeding 20 mmHg with new organ failure defines abdominal compartment syndrome.

#### **Alternate Names:**

IAH

### **Intraabdominal Infection**

An infection within the abdominal cavity, such as peritonitis or an abscess, often requiring antibiotics and sometimes drainage or surgery.

### **Intra-Aortic Balloon Pump**

A temporary heart-support device with a balloon inside the body's main artery. It inflates and deflates with each heartbeat to improve blood flow and reduce the heart's workload; bleeding, infection, and reduced limb blood flow can occur.

### **Intra-Arterial**

Intra-arterial means within an artery, including a catheter, injection, infusion, or measurement performed directly in an arterial vessel.

### **Intraarticular**

Intraarticular means within a joint, such as an injection, fracture, loose body, or inflammatory process inside the joint space.

### **Intra-Articular Injection**

An injection into a joint space, such as the knee, shoulder, hip, or ankle. It may deliver medicine or remove joint fluid; corticosteroids, local anesthetics, and other substances may be used for selected joint problems.

#### **Alternate Names:**

Joint Injection; Intraarticular Injection; Intra-Articular Corticosteroid Injection; Joint Infiltration

### **Intrabronchial**

Intrabronchial means within a bronchus, one of the air passages branching from the trachea into the lungs.



---

### **Intracardiac Echocardiography**

An invasive imaging modality in which a steerable catheter-tipped ultrasound probe is introduced via the femoral vein into the cardiac chambers (usually the right atrium) to provide real-time, high-resolution visualization of endocardial structures and guide structural heart and electrophysiology interventions.

**Alternate Names:**

ICE

### **Intracardiac Electrophysiology Study**

An invasive test using electrode catheters inside the heart to record electrical activity and provoke or map arrhythmias, helping guide diagnosis and treatment.

### **Intracartilaginous**

Located within cartilage, such as a lesion, injection, or developing structure. The term identifies position and does not by itself indicate whether the finding is normal, injured, inflamed, or cancerous.

### **Intracavernosal Injection**

An injection of a prescription medicine into the erectile tissue of the penis to treat erectile dysfunction. The medicine relaxes muscle and widens blood vessels, producing an erection that does not require sexual stimulation. Pain, bruising, scar tissue, or a prolonged painful erection can occur.

**Alternate Names:**

ICI; Penile Injection Therapy; Intracavernous Injection

### **Intracavitary**

Intracavitary means within a body cavity, such as a treatment, device, fluid, or lesion located in the thorax, abdomen, uterus, or another cavity.

### **Intracavity Radiotherapy**

Intracavity radiotherapy places a radiation source inside a body cavity near a tumor, commonly to treat selected cancers of the cervix, uterus, or vagina.

### **Intracellular**

Inside a cell or occurring within a cell. Intracellular fluid contains most of the body's potassium and magnesium, whereas sodium is found mainly outside cells. The cell membrane controls movement of water and dissolved substances between the cell and its surroundings.

### **Intracellular Fluid**

The fluid contained within cells, comprising approximately two thirds of total body water in a typical adult. Its major solutes include potassium, magnesium, phosphate, and intracellular proteins, and its composition is maintained by selective membrane transport.

### **Intracellular Membrane**

A membrane inside a cell that separates compartments and controls movement of substances between them.

### **Intracellular Pathogens**

Germs that live and multiply inside the body's cells. Viruses always do this, and some bacteria do as well; living inside cells can make them harder for immune defenses and some medicines to eliminate.

### **Intracellular Transport**

Movement of proteins and other materials within a cell, often in membrane-bound sacs along the cell's internal framework.

### **Intracerebral**

Intracerebral means within the brain tissue, as in an intracerebral hemorrhage, injection, electrode, or lesion.

### **Intracerebral Hemorrhage**

Bleeding directly into brain tissue, usually from a small ruptured artery. It can cause sudden headache, weakness, vomiting, confusion, seizures, or loss of consciousness.

### **Intracisternal**

Intracisternal means within a cistern, especially a

---

cerebrospinal-fluid space at the base of the brain.

### **Intracoronal**

Intracoronal means within the crown of a tooth or within a coronary structure, with the exact meaning determined by context.

### **Intracoronary**

Intracoronary means within a coronary artery or the coronary circulation, such as a catheter, stent, imaging study, or medicine delivery.

### **Intracranial**

Inside the skull. The term may describe the brain, blood vessels, fluid, bleeding, pressure, infection, or a growth within the skull. Extra fluid, blood, or tissue can raise pressure and damage the brain, causing headache, vomiting, confusion, weakness, seizures, or loss of consciousness.

### **Intracranial Aneurysm**

A weakened, bulging area in a blood vessel inside the skull, usually in an artery supplying the brain. Many cause no symptoms, but a leak or rupture can cause bleeding around the brain and a sudden, severe headache.

### **Intracranial Hemorrhage**

Bleeding inside the skull, either within the brain or around it. It may follow injury, high blood pressure, a ruptured aneurysm, or medicines that prevent clotting and can rapidly cause brain pressure, disability, or death.

### **Intracranial Pressure**

The pressure inside the skull produced by the brain, blood, and surrounding fluid. Too much pressure can reduce blood flow to the brain and cause headache, vomiting, confusion, unequal pupils, seizures, unconsciousness, or life-threatening brain injury.

### **Intracranial Pressure Monitoring**

Continuous or intermittent measurement of pressure inside the skull, usually in critically ill patients with

severe brain injury, hemorrhage, hydrocephalus, or other risk of intracranial hypertension.

### **Intracranial Venous Malformations**

Abnormal collections or connections of veins within the skull that may bleed or affect nearby brain tissue.

### **Intractable**

Difficult to control, relieve, or cure despite appropriate treatment, as in intractable pain or seizures.

### **Intracytoplasmic Sperm Injection**

A fertility treatment in which a single sperm is injected directly into an egg in a laboratory to help achieve fertilisation. It is also called ICSI.

### **Intradermal**

Located within the dermis, the layer of skin beneath the epidermis; an intradermal injection places medication or test material into that layer.

### **Intradermal Nevus**

A common, benign melanocytic skin nevus characterized by nests of mature melanocytes located entirely within the dermis without junctional melanocytic proliferation, appearing as a soft, dome-shaped, skin-colored, pedunculated or light brown papule.

#### **Alternate Names:**

Intradermal Melanocytic Nevus; Dermal Mole

### **Intradermal Skin Test**

A small amount of a test substance is injected into the top layer of the skin. The area is then checked for a reaction. This method is used for tests such as tuberculin testing and some allergy tests.

### **Intraductal Carcinoma**

Alternate name; see *Ductal Carcinoma In Situ*.

### **Intraductal Papillary Mucinous Neoplasm**

Alternate name; see *Pancreatic Cysts*.

---

### **Intraductal Papilloma**

A benign, wart-like tumor that grows within the lining of a breast milk duct, composed of delicate fibrovascular finger-like projections covered by epithelial and myoepithelial cells. It is the most common cause of unilateral, spontaneous, clear or bloody (serosanguinous) nipple discharge in women aged 30 to 50 years. Solitary central papillomas arise in large subareolar ducts and have minimal malignant risk, whereas multiple peripheral papillomas arise in smaller peripheral ducts and carry an increased risk of breast cancer. Diagnosis involves ultrasound, mammography, or ductography, and complete surgical excision (microdochectomy) is typically curative.

#### **Alternate Names:**

Ductal Papilloma; Solitary Intraductal Papilloma; Breast Papilloma

### **Intraduodenal**

Intraduodenal means within the duodenum, the first part of the small intestine beyond the stomach.

### **Intraepidermal**

Intraepidermal means within the epidermis, the outermost layer of skin.

### **Intraepithelial**

Located within the layer of cells that covers a body surface or lines an organ. The word may describe infection, inflammation, or abnormal cell changes that have not yet grown through the supporting layer beneath them.

### **Intraesophageal**

Intraesophageal means within the esophagus, such as a probe, pressure sensor, lesion, or medication delivery.

### **Intrahepatic**

Within the liver. For example, intrahepatic bile-duct blockage or inflammation can cause jaundice and abnormal liver tests, while a tumor may begin in liver cells or in bile ducts inside the liver. The exact meaning depends on the condition being described.

### **Intrahepatic Cholestasis**

Reduced bile flow within the liver causing bile substances to build up in blood, often producing itching, jaundice, dark urine, or pale stools.

### **Intrahepatic Cholestasis Of Pregnancy**

A pregnancy-related liver disorder causing intense itching, often on the palms and soles, without a primary rash. Bile acids rise in the blood and can increase fetal risk, so testing and obstetric monitoring are needed.

### **Intralesional**

Delivered directly into a lesion, such as an injection of corticosteroid into a keloid or a medicine into a skin, joint, or tumor lesion. The drug, dose, sterility, and lesion type determine safety.

### **Intralymphatic**

Intralymphatic means within a lymphatic vessel or channel, such as a medicine, tumor spread, infection, or imaging contrast.

### **Intramedullary**

Intramedullary means within the medulla of an organ or the central canal of a bone, especially an implant placed inside a long bone.

### **Intramedullary Nail**

A metal implant inserted into the medullary canal of a long bone to align and stabilize a fracture. It shares load along the bone axis and is commonly used for femoral and tibial shaft fractures.

### **Intramedullary Spinal Cord Abscesses**

Localized collections of pus within the spinal cord, usually caused by infection and associated with severe neurologic deficits, fever, or spinal pain. MRI and urgent antimicrobial and surgical evaluation are typically required.

### **Intramural Hematoma**

Bleeding into the wall of the digestive tract, often in the small intestine. It can follow injury or blood-thinning treatment and may cause abdominal pain,

---

vomiting, or blockage.

### **Intramural Plaque**

An intramural plaque is an atherosclerotic deposit within an artery wall that can narrow the lumen or destabilize and cause thrombosis.

### **Intramuscular**

Within a muscle. An intramuscular injection delivers a medicine into muscle tissue, commonly in the upper arm, thigh, or hip, where it is absorbed into the bloodstream.

### **Intramuscular Injection**

Delivery of a medicine into a muscle with a needle. The injection site and needle size depend on the medicine and the person's age and body size; pain, bleeding, infection, or nerve injury are possible.

### **Intramuscular Injection Technique**

The method used to place medicine into muscle tissue with a needle and syringe. The site, needle, and injection method vary with the medicine and the person.

#### **Alternate Names:**

Intramuscular Injection Technique; Parenteral Administration

### **Intramuscular Myxoma**

A rare, usually noncancerous tumor made of jelly-like tissue within a skeletal muscle. It often appears as a slow-growing painless lump, but imaging or biopsy may be needed to distinguish it from other masses.

### **Intraocular**

Within the eyeball. The term may describe a medicine injection, lens, infection, bleeding, foreign body, pressure, or other finding inside the eye. Increased pressure inside the eye can damage the optic nerve and cause permanent loss of vision.

### **Intraocular Infection**

An infection inside the eye, such as endophthalmitis, that can rapidly threaten vision.

### **Intraocular Lens**

An artificial lens placed inside the eye, usually during cataract surgery, to focus light after the cloudy natural lens is removed. Lens choices can emphasize distance, near vision, or astigmatism correction. Possible complications include glare, clouding behind the lens, infection, or retinal problems.

### **Intraocular Lens Dislocation**

Movement of an artificial lens away from its intended position after eye surgery. It may cause blurred or double vision, glare, or pain and can require repositioning or replacement by an eye specialist.

### **Intraocular Lens Implant**

An artificial lens implanted inside the eye, usually during cataract surgery; see *Intraocular Lens*.

### **Intraocular Lymphoma**

A lymphoma involving the eye, often the jelly-like vitreous or the retina. It may cause blurred vision, floaters, or inflammation and can occur with lymphoma in the brain or other parts of the body.

### **Intraocular Pressure**

The pressure of fluid inside the eye. If it stays too high, it can damage the optic nerve and cause glaucoma, although glaucoma can sometimes occur even when the pressure is within the usual range.

### **Intraocular Tumor**

A growth arising within the eye. It may be noncancerous or cancerous; examples include uveal melanoma in adults and retinoblastoma in children. It can affect vision or eye pressure.

### **Intraoperative**

Occurring during a surgical operation, including monitoring, findings, complications, or treatment decisions made during the procedure.

### **Intraoperative Blood Transfusion**

Blood or blood components given during surgery to replace major blood loss and maintain oxygen delivery. The decision depends on bleeding, symptoms,

---

circulation, hemoglobin, and the person's medical condition.

### **Intraoperative Neuromonitoring**

Electrical tests used during selected operations to watch the function of the brain, spinal cord, nerves, or muscles. Signals can warn the surgical team about a possible change in nerve function, allowing them to reassess the procedure and protect nearby structures.

### **Intraoral**

Located inside the mouth or performed through the mouth, such as an examination, scan, injection, or surgical procedure.

### **Intraosseous Access**

Placement of a special needle into the inner part of a bone to give fluids, blood, or medicines when a vein cannot be reached quickly. It is temporary and can cause pain, leakage, infection, or bone injury.

### **Intraosseous Infusion**

An emergency vascular access technique in which a specialized rigid hollow-bore trocar needle is drilled through the cortical bone into the rich medullary vascular marrow space (commonly the proximal tibia, proximal humerus, or distal femur) to deliver fluids, blood products, and resuscitation medications when peripheral intravenous access fails.

#### **Alternate Names:**

IO Access; Intraosseous Vascular Access

### **Intrapartum**

The period during labor and childbirth, from the onset of regular uterine contractions leading to cervical dilation until delivery of the baby and placenta.

### **Intrapartum Fever**

Fever during labor, which may result from infection, epidural analgesia, dehydration, or other causes and can affect the mother or fetus.

### **Intrapartum Stillbirth**

Intrapartum stillbirth is fetal death occurring during

labor after a live fetus was documented before labor, requiring obstetric investigation and support.

### **Intrapericardial**

Located inside the pericardial sac, the thin membrane surrounding the heart. An intrapericardial mass or fluid collection may affect how the heart fills or beats.

### **Intraperitoneal**

Intraperitoneal means within the peritoneal cavity or administered into it, including selected medicines, dialysis fluid, or procedures.

### **Intraperitonealroute**

A route of treatment in which fluid or medicine is placed into the abdominal cavity. It is used for peritoneal dialysis and selected treatments and requires sterile technique to prevent infection.

### **Intraprostatic**

Located within the prostate gland, describing a lesion, injection, infection, biopsy, procedure, or other finding centered inside prostate tissue.

### **Intrapulmonary**

Located within lung tissue or occurring inside the lung's air spaces, blood vessels, or other structures. Intrapulmonary findings may represent normal anatomy, infection, fluid, bleeding, inflammation, or tumors.

### **Intrapulmonary Shunt**

Blood flow through parts of the lungs that are not receiving enough air, so the blood cannot take up normal oxygen. It can occur with pneumonia, lung collapse, or severe lung injury and may cause low oxygen despite supplemental oxygen.

### **Intrasynovial**

Intrasynovial means within the synovial lining or fluid-filled space of a joint or tendon sheath.

### **Intratendinous**

Located within a tendon, describing a tear, calcium

---

deposit, injection, degenerative change, or other abnormality affecting the tendon's substance.

### **Intratesticular**

Located within testicular tissue. An intratesticular mass may be benign or malignant and is usually evaluated with physical examination and scrotal ultrasound, interpreted in the clinical context.

### **Intrathecal**

Intrathecal means within the fluid-filled space surrounding the brain and spinal cord; medicines may be given there by lumbar puncture.

### **Intrathecal Chemotherapy**

Cancer medicine placed directly into the fluid surrounding the brain and spinal cord, usually through a lumbar puncture or implanted reservoir. It is used for selected leukemias, lymphomas, or cancer spread to this fluid; infection, bleeding, headache, and nervous-system toxicity are possible.

### **Intrathoracic**

Intrathoracic means within the chest cavity, including the lungs, heart, mediastinum, vessels, and thoracic lymph nodes.

### **Intrathoracic Localized Obstruction**

A blockage within the chest affecting an airway, esophagus, or major blood vessel. Tumors, inhaled objects, scarring, or abnormal vessels may cause breathing difficulty, swallowing problems, chest pain, or reduced blood flow and require targeted testing.

### **Intratracheal**

Intratracheal means within the trachea, such as a tube, medicine, foreign body, lesion, or route of administration.

### **Intratympanic**

Intratympanic means delivered through the eardrum into the middle ear, often for selected inner-ear disorders or local treatment.

### **Intrauterine**

Located or occurring within the uterus, including a pregnancy, device, medicine, lesion, or procedure. The term describes location and requires additional information to explain the finding's purpose, cause, or risk.

### **Intrauterine Contraceptive Device**

A small device placed in the uterus to prevent pregnancy; copper IUDs release copper, while hormonal IUDs release a progestin, and both are highly effective reversible contraception.

### **Intrauterine Device**

A small device placed inside the uterus to prevent pregnancy for several years while remaining reversible. Copper devices work without hormones; hormonal devices release a progestogen and often reduce menstrual bleeding. Rare risks include expulsion, perforation during insertion, and infection soon after placement.

### **Intrauterine Devices**

Small devices placed in the uterus to prevent pregnancy; copper and hormonal IUDs are highly effective long-acting reversible contraceptives.

### **Intrauterine Fetal Demise / Stillbirth**

Death of a fetus before complete delivery. The gestational-age definition varies by jurisdiction. Causes may include placental disease, umbilical-cord complications, infection, genetic abnormalities, maternal disease, or unexplained factors.

### **Intrauterine Growth Restriction**

A fetus that is smaller than expected for its stage of pregnancy or is not growing normally. Causes include reduced placental function, maternal illness, fetal genetic conditions, infection, or inaccurate dating; monitoring guides care.

### **Intrauterine Insemination**

A fertility procedure in which prepared sperm is placed directly into the uterus around ovulation. It is used for selected infertility causes and usually depends on at least one open fallopian tube; risks



---

include cramps, infection, and multiple pregnancy.

### **Intrauterine Pressure Catheter**

A thin tube placed inside the uterus during labor to measure contraction strength and timing more directly. It is used in selected situations when external monitoring does not provide enough information and can cause infection or injury.

### **Intrauterine System**

A small T-shaped contraceptive device placed inside the uterus that slowly releases a progestogen hormone to prevent pregnancy. It is also called IUS.

### **Intrauterine Transfusion**

A specialist procedure that gives compatible red blood cells to a fetus through the umbilical cord or, less often, the fetal abdomen. It treats severe fetal anemia and carries risks such as bleeding, infection, early labor, or fetal loss.

### **Intravascular**

Intravascular means within a blood vessel, including a catheter, clot, injection, or process occurring in the circulation.

### **Intravascular Catheter Infections**

Infections caused by microorganisms entering or growing on a catheter placed inside a blood vessel. They may produce fever, chills, low blood pressure, or bloodstream infection. Diagnosis uses examination and blood cultures, and treatment may require antibiotics and catheter removal.

### **Intravascular Filter**

A small device placed in a large vein to catch clots traveling from the legs toward the lungs. It is used selectively when blood-thinning treatment cannot be given or has not worked adequately.

### **Intravascular Hemolysis**

Destruction of red blood cells within the bloodstream, releasing hemoglobin into plasma; causes include immune hemolysis, mechanical trauma, infections, toxins, and transfusion reactions.

### **Intravascular Lymphoma**

A rare, fast-growing blood cancer in which abnormal white blood cells collect inside small blood vessels. It may affect the skin, brain, or other organs, causing fever, skin changes, or neurologic problems.

### **Intravascular Ultrasound**

An ultrasound test done from inside a blood vessel. A tiny ultrasound device on a catheter creates pictures of the vessel wall and any narrowing or fatty plaque, helping guide some treatments.

**Alternate Names:**

IVUS

### **Intravenous**

Into or within a vein; intravenous medicines, fluids, or blood products are delivered directly into the bloodstream.

### **Intravenous Anesthesia**

General anesthesia started or maintained with medicines given through a vein. It acts quickly, but the anesthesia team must adjust the dose and continuously monitor breathing, blood pressure, heart rhythm, and recovery.

### **Intravenous Catheters**

Flexible devices inserted into a vein to deliver fluids, medicines, blood products, nutrition, or contrast, or to obtain blood samples; complications include infection, thrombosis, infiltration, and extravasation.

### **Intravenous Drip**

Delivery of fluid, blood products, nutrients, or medicine directly into a vein through a catheter. The amount and speed are prescribed and monitored for reactions, fluid overload, infection, or vein irritation.

**Alternate Names:**

Intravenous Infusion

### **Intravenous Immunoglobulin**

A purified preparation of pooled antibodies given through a vein (IVIg) for selected immune deficiencies, autoimmune diseases, inflammatory disorders,

---

and some neurologic conditions.

### **Intravenous Leiomyomatosis**

Extension of benign uterine smooth-muscle tumor into venous channels, sometimes reaching the inferior vena cava or right heart. Imaging is important because extensive disease can cause serious cardiovascular complications.

### **Intravenous Lipid Emulsion Therapy**

Emergency treatment using a fat-containing fluid through a vein for selected severe poisonings, especially toxicity from local anesthetics. It can help remove some medicines from vital tissues and requires intensive monitoring.

### **Intravenous Pyelogram**

An X-ray examination of the kidneys, ureters, and bladder after iodine-containing contrast is injected into a vein and passed into the urine. CT urography or ultrasound often provides this information today.

### **Intravenous Pyelography**

An X-ray examination of the urinary tract after intravenous contrast administration; it is now often replaced by CT urography or ultrasound.

### **Intravenous Sedation**

Administration of sedative medicines through a vein to reduce anxiety, discomfort, or awareness during a procedure while maintaining appropriate monitoring and readiness to support breathing.

### **Intravenous Thrombolysis**

Administration of a clot-dissolving medicine through a vein to restore blood flow in selected acute vascular conditions; see *Thrombolysis*.

### **Intraventricular Hemorrhage**

Bleeding into the fluid-filled spaces inside the brain, especially in very premature newborns. The blood can block fluid flow and cause brain injury; ultrasound is used to detect and monitor it.

### **Intraventricular Hemorrhage Of The**

### **Newborn**

Intraventricular hemorrhage in a newborn is bleeding into the brain's ventricular system, most often in premature infants, and can cause hydrocephalus or neurologic injury.

### **Intravesical Therapy**

Treatment delivered directly into the bladder through a catheter. It is used for conditions such as some bladder cancers or interstitial cystitis and may cause bladder irritation or, less commonly, effects elsewhere in the body.

### **Intravitreal Injection**

An eye procedure in which medicine is injected into the jelly-like center of the eye to treat retinal or macular disease. Temporary discomfort or floaters may occur; infection, bleeding, retinal detachment, and raised eye pressure are uncommon but serious risks.

### **Intrinsic**

Belonging naturally to a structure or process and arising from within it, rather than being externally imposed.

### **Intrinsic Factor**

A protein secreted by stomach parietal cells that is required for vitamin B12 absorption in the terminal ileum.

### **Intrinsic Pathway of Coagulation**

The contact activation cascade of blood coagulation initiated when factor XII (Hageman factor), prekallikrein, and high-molecular-weight kininogen encounter exposed subendothelial collagen or negatively charged surfaces, sequentially activating factors XI and IX, forming the tenase complex (IXa, VIIIa, Ca<sup>2+</sup>, phospholipid) to activate factor X; monitored clinically by the activated partial thromboplastin time (aPTT).

### **Intrinsic Renal AKI**

Sudden loss of kidney function caused by damage within the kidney itself. Causes include medicine or toxin injury, inflammation, infection, poor blood

---

supply, and immune disease; treatment depends on the cause and may require hospital care.

### **Intrinsic Sphincter Deficiency**

Weakness of the urinary sphincter that causes stress urinary incontinence.

**Alternate Names:**

ISD

### **Intrinsic Sympathomimetic Activity**

The pharmacological property of certain beta-adrenergic receptor antagonist drugs (such as acebutolol and pindolol) that act as partial agonists, stimulating beta-receptors to a mild degree while simultaneously blocking endogenous catecholamines, resulting in less resting bradycardia.

**Alternate Names:**

ISA

### **Introitus**

An entrance or opening, especially the entrance to the vagina.

### **Intron**

A non-coding segment of a gene that is transcribed into RNA but spliced out before the messenger RNA is translated into a protein.

### **Intrusive Thought**

An unwanted, distressing thought, image, or impulse that enters awareness repeatedly; it does not necessarily reflect intent or desire.

### **Intubate**

To place a tube into a body passage, especially an endotracheal tube through the mouth or nose into the trachea to maintain an airway or support ventilation.

### **Intubation**

Placement of a tube into the airway or another body passage to maintain it or deliver treatment.

### **Intussusception**

A condition in which one segment of the intestine

slides inside an adjacent segment like a folding telescope. It can block the bowel and reduce blood flow, especially in infants and young children, causing cramping pain, vomiting, swelling, or bloody stool. Ultrasound may show a target-like ring, and an air or contrast enema may be used in children.

**Alternate Names:**

Intestinal telescoping; Invagination of the bowel; Ileocolic intussusception.

### **Inulin**

Inulin is a plant fructan used as a dietary fiber and in tests of glomerular filtration because it is freely filtered and minimally handled by the kidney.

### **In Utero**

Within the uterus before birth. The term describes a developing fetus or an event that occurs during pregnancy.

### **Invasive**

Able to enter and spread through tissues; the term may describe an infection, tumor, procedure, or disease that penetrates nearby structures.

### **Invasive Candidiasis**

A serious *Candida* infection that enters the bloodstream or deep organs, most often in people who are critically ill or immunocompromised. It may cause persistent fever or organ dysfunction and requires prompt medical diagnosis and treatment.

### **Invasive Ductal Carcinoma**

The most common type of breast cancer, characterized by malignant cells that have broken through the duct wall and invaded surrounding breast tissue. It can metastasize via lymphatic or hematogenous spread.

### **Invasive Lobular Carcinoma**

A breast cancer whose cells spread through breast tissue in single-file patterns rather than forming a distinct lump. It can be difficult to detect on examination or imaging and may affect more than one area or breast.

---

## **Invasive Mechanical Ventilation**

Breathing support delivered through a tube placed in the trachea. A ventilator moves air and oxygen into the lungs and removes carbon dioxide when a person cannot breathe adequately without support.

## **Invasive Mole**

A hydatidiform mole that invades the uterine myometrium or vascular spaces; it is a form of gestational trophoblastic neoplasia and may persist or metastasize.

## **Invasive Procedure**

A medical examination, test, or treatment that enters the body by piercing the skin or inserting an instrument, tube, needle, or camera into internal tissues or organs. Examples range from needle biopsies and endoscopy to open surgery. Possible risks include bleeding, infection, pain, tissue damage, and anesthesia complications.

### **Alternate Names:**

Invasive Medical Procedure; Invasive Intervention; Invasive Surgery

## **Inverse Agonist**

A medicine that binds to a receptor and reduces its natural baseline activity, producing an effect opposite to that of an agonist. Its clinical effect depends on the receptor and the tissues where it is present.

## **Inverse Psoriasis**

A morphological variant of plaque psoriasis that preferentially involves intertriginous and flexural anatomical regions (axillae, inframammary folds, inguinal creases, and intergluteal cleft), appearing as well-demarcated, smooth, glistening erythematous plaques without typical silvery scales due to maceration.

### **Alternate Names:**

Flexural Psoriasis; Intertriginous Psoriasis

## **Inversion**

A chromosome segment that has broken off and rejoined in the reverse direction. It often causes no health problem, but can affect health if the break

disrupts a gene or interferes with reproduction.

## **Invert**

To turn upside down, reverse, or turn inward; in anatomy, inversion can describe movement of the sole of the foot toward the midline.

## **Inverted Nipple**

Alternate name; see *Nipple Inversion Or Retraction*.

## **Inverted Papilloma**

A usually benign but locally destructive tumor arising most often in the nasal cavity or sinuses; it can recur and rarely become cancerous.

## **Invert Sugar**

A mixture of glucose and fructose produced by splitting sucrose, used as a sweetener in foods and medicines. It has nutritional effects similar to other added sugars and contributes to total carbohydrate and energy intake.

## **Investigational**

Being studied in clinical research and not yet established as safe and effective for a particular use. Investigational medicines and devices require appropriate oversight, informed consent, and clear communication of uncertainty.

## **Inveterate**

Long-standing, deeply established, or difficult to change, often describing a habit, condition, or behavior. The word indicates persistence but does not identify the cause, severity, treatment, or expected outcome.

## **In Vitro**

In vitro means performed outside a living organism, such as in a test tube, culture dish, or laboratory system.

## **In Vitro Fertilisation**

A fertility treatment in which eggs are collected from the ovaries, fertilized with sperm in a laboratory, and one or more resulting embryos are placed in the

---

uterus.

### **In Vitro Fertilization**

An assisted-reproduction treatment in which eggs are collected, fertilized with sperm in a laboratory, and an embryo is transferred to the uterus. Risks include ovarian hyperstimulation, multiple pregnancy, ectopic pregnancy, and procedure-related complications.

### **In Vivo**

Taking place or being studied inside a living person, animal, or other organism. For example, an in vivo test examines how a medicine works in the body rather than only in a laboratory dish.

### **In-Vivo**

Occurring or being studied inside a living person, animal, or other organism, rather than only in a laboratory sample or cell culture.

### **Involucrum**

An involucrum is a sheath of new bone that forms around dead bone, classically in chronic osteomyelitis.

### **Involuntary**

Occurring without conscious control or deliberate choice, such as a reflex, autonomic function, or involuntary muscle movement.

### **Involuntary Movement**

A movement that occurs without conscious control, such as a tremor, tic, chorea, dystonia, or myoclonus.

### **Involuntary Movements**

A diverse group of neurologic motor disorders characterized by uncontrollable, purposeless, or abnormal spontaneous body movements, including tremors, chorea, athetosis, hemiballismus, dystonia, myoclonus, and tics, stemming from basal ganglia, cerebellar, or motor circuit pathology.

#### **Alternate Names:**

Hyperkinetic Movement Disorders; Involuntary Mo-

tor Activity; Dyskinetic Movements

### **Involuntary Muscle**

Muscle that contracts without conscious control, including smooth muscle in internal organs and cardiac muscle in the heart.

### **Involuntary Psychiatric Treatment**

Assessment, hospitalization, or treatment without the patient's consent under applicable law when serious mental disorder creates a substantial risk of harm or severe inability to meet basic needs. The least restrictive alternative, due process, and ongoing reassessment are required.

### **Involute**

To shrink, return toward a former size, or undergo involution; in anatomy, an involute structure may be tightly coiled or rolled inward.

### **Involution**

The normal return of the uterus to its smaller, prepregnancy size after childbirth. This occurs gradually over several weeks as the uterine muscle contracts and the placental attachment heals.

### **Iodamoeba Butschlii**

A usually non-disease-causing intestinal amoeba spread by swallowing cysts in contaminated food or water. Finding it in stool suggests fecal exposure but usually does not require treatment unless another infection is present.

### **Iodide**

The negatively charged form of iodine, used by the thyroid to make hormones and present in medicines, supplements, and contrast compounds.

### **Iodide-Associated Sialadenitis**

Swelling and inflammation of a salivary gland after exposure to a large amount of iodide, sometimes from a medicine or imaging contrast. It can cause painful swelling and dry mouth and usually improves when the cause is addressed.

---

## Iodides

Salts containing iodine, sometimes used in carefully selected thyroid disorders.

## Iodinated Contrast Media

Iodine-containing substances injected for CT scans, angiography, X-rays, or fluoroscopy to make vessels and organs easier to see. They can cause allergic-like reactions, leakage into tissue, thyroid changes, or kidney injury in susceptible people.

## Iodine

An essential mineral used to make thyroid hormones, which regulate growth, brain development, and metabolism. Too little iodine can cause thyroid enlargement and hormone deficiency, especially harmful during pregnancy and childhood; too much can also disturb thyroid function. Main sources include iodized salt, seafood, dairy, and some supplements.

## Iodine-123

A radioactive form of iodine used mainly to create images of the thyroid. It gives off a type of radiation detected by a camera and generally exposes the body to less radiation than iodine-131.

## Iodine-131

A radioactive form of iodine used mainly to diagnose or treat selected thyroid disorders. Thyroid cells absorb it, allowing targeted radiation, but exposure can affect other tissues and requires safety precautions.

## Iodine Deficiency

Too little iodine in the diet. Because iodine is needed to make thyroid hormones, deficiency can cause thyroid enlargement and low thyroid function. Severe deficiency during pregnancy or childhood can impair growth and brain development.

### Alternate Names:

Iodine Deficiency Disorders; IDD; Endemic Iodine Deficiency

## Iodine Perfusion

The distribution of iodine-containing contrast through tissues or blood vessels, measured during imaging such as computed tomography or angiography. Contrast can rarely cause allergic reactions and may affect kidney or thyroid function.

## Iodine Poisoning

Iodine poisoning follows excessive iodine exposure and can irritate the gastrointestinal tract, disrupt thyroid function, and cause systemic toxicity depending on dose and formulation.

## Iododerma

A skin eruption caused by excess iodine, often appearing as acne-like bumps or pus-filled spots on the face, arms, or legs. It usually improves after the iodine source is stopped under medical guidance.

## Iodoform

A yellow iodine-containing compound formerly used as an antiseptic and in some wound or dental preparations. It can irritate tissue and cause toxicity if absorbed or swallowed and is now used only in selected contexts.

## Iohexol

A water-soluble iodine-containing contrast medicine used during CT scans, angiography, and other imaging studies to make vessels and organs easier to see.

## Ion

An atom or molecule carrying a positive or negative electric charge. Physiologically important ions include sodium, potassium, calcium, chloride, and bicarbonate, whose concentrations influence nerves, muscles, fluid balance, and acid-base status.

## Ion Channel

A protein pore in a cell membrane that allows selected electrically charged particles, such as sodium, potassium, calcium, or chloride, to cross. Ion channels are important for nerve signals, muscle contraction, and secretion.



---

## Ionization

The formation of charged particles by gaining or losing electrons or protons. A medicine's degree of ionization affects its solubility, membrane passage, absorption, distribution, and urinary excretion.

## Ionized Calcium Test

A clinical laboratory assay measuring free (unbound), physiologically active calcium ions in serum or whole blood. It provides an accurate assessment of calcium homeostasis independent of serum albumin concentrations, particularly in critical illness, acid-base disorders, and parathyroid pathology.

### Alternate Names:

Ionized Calcium; Free Calcium Test; Serum Free Calcium

## Ionizing Radiation

Radiation with enough energy to remove electrons from atoms and create ions, including X-rays, gamma rays, and certain particulate emissions. It can cause deterministic tissue effects at sufficient doses and stochastic cancer risk at lower doses.

## Ionophore

A molecule that transports ions across a lipid membrane. Ionophores have research, antimicrobial, and veterinary uses; some can disrupt human cellular ion balance and are toxic.

## Ions

Electrically charged atoms or molecules, including the electrolytes that regulate nerve conduction, muscle function, hydration, and acid-base balance. Abnormal ion concentrations can cause weakness, seizures, arrhythmias, or altered consciousness.

## Iontophoresis

Use of a mild electrical current to drive an ionized medicine through the skin or alter sweat-gland activity. It is used in selected dermatologic and rehabilitation settings; skin irritation and burns can occur.

## Ion Transport

Movement of charged particles across a cell membrane through channels, carriers, or pumps. It maintains membrane potentials, nerve signaling, muscle contraction, fluid balance, and epithelial transport.

## Iopamidol

A nonionic iodine-containing contrast medicine used to improve CT, angiography, and other medical images. It can cause warmth, nausea, allergy, kidney problems, or thyroid effects, so risks require assessment beforehand.

## Iothalamate Clearance

A specialized test of kidney filtration that measures how quickly injected iothalamate is removed from the blood, sometimes using timed urine samples. It is used when a more direct estimate of kidney function is needed.

## Ioxaglic Acid

An older iodinated contrast medium used for radiographic imaging, including selected angiographic and urinary-tract studies. It has largely been superseded by newer agents, but its history is relevant when assessing prior contrast reactions.

## IPC

Alternate name; see *Indwelling Pleural Catheter*.

## Ipecac

A vomiting medicine formerly used after poisoning, now not routinely recommended because vomiting can cause aspiration and delay more appropriate emergency treatment.

## Ipilimumab

An immunotherapy antibody that blocks CTLA-4, removing a brake on T cells so they can attack cancer. It can cause immune inflammation in the bowel, skin, liver, lungs, hormone glands, or other organs.

## IPMN

Abbreviation; see *Pancreatic Cysts*.

---

## **Ipratropium**

Ipratropium is an inhaled short-acting muscarinic antagonist that reduces vagally mediated bronchoconstriction and nasal secretion; it is used in COPD and acute asthma with bronchodilators.

## **Ipratropium Bromide**

An inhaled medicine that relaxes narrowed airways and reduces mucus production. It is used mainly for chronic obstructive pulmonary disease and, with other medicines, for severe asthma attacks. Dry mouth and blurred vision can occur.

## **Iproniazid**

Iproniazid is an older monoamine oxidase inhibitor antidepressant that is no longer commonly used because of serious liver toxicity and drug interactions.

## **Ipsilateral**

On or affecting the same side of the body as a stated structure or injury. For example, weakness in the right arm and right leg is ipsilateral to a finding on the right side of the body; the opposite term is contralateral.

## **IQ Testing**

IQ testing uses standardized tasks to estimate intellectual functioning; results should be interpreted with education, language, development, and clinical context.

## **Irbesartan**

Irbesartan is an angiotensin-II receptor blocker used for hypertension and diabetic kidney disease; it can cause hyperkalaemia, hypotension, and renal-function changes and is contraindicated in pregnancy.

## **Iridium**

Iridium is a dense metal whose radioactive isotopes, especially iridium-192, are used in industrial testing and some medical brachytherapy.

## **Iridocyclitis**

Inflammation of the iris and nearby ciliary body

inside the eye. It can cause pain, redness, sensitivity to light, blurred vision, and inflammatory cells in the front of the eye; untreated disease can threaten vision.

## **Iridodialysis**

A traumatic or surgical separation or avulsion of the iris root from its anatomical attachment to the ciliary body, producing a characteristic D-shaped eccentric pupil, monocular diplopia, photophobia, and an anterior chamber hyphema.

## **Iridodonesis**

Shaking or trembling of the iris when the eye moves, often because the lens is absent or its supporting fibers are weak. An eye examination can identify the cause and related lens or eye problems.

## **Iridoschisis**

A rare splitting of the iris tissue into layers, usually in older adults. It may be associated with glaucoma or corneal problems and should be assessed by an eye specialist.

## **Iridotomy**

A laser or surgical procedure that makes a tiny opening in the colored part of the eye to improve fluid drainage. It is used to treat or prevent selected types of angle-closure glaucoma.

## **Irinotecan**

A chemotherapy medicine that inhibits topoisomerase I and is used for colorectal and several other cancers.

## **Irinotecan Hydrochloride**

A chemotherapy medicine that blocks topoisomerase I and damages cancer-cell DNA, used for selected cancers with diarrhea and low blood counts possible.

## **Iris**

The colored, muscular ring between the cornea and lens that controls the size of the pupil and therefore the amount of light entering the eye. Its muscles constrict the pupil in bright light and enlarge it in

---

dim light.

### **Iris Atrophy**

Thinning or loss of iris tissue resulting from trauma, chronic inflammation, ischaemia, or neurodegenerative disease, leading to a translucent or patched iris appearance. It may be sectoral or diffuse and can be associated with decreased pupillary response, photophobia, or secondary glaucoma.

### **Iris Cyst**

An iris cyst is a fluid-filled lesion on or within the iris, usually benign but sometimes affecting vision, pressure, or the appearance of the eye.

### **Iris Disease**

A disorder of the colored part of the eye, including inflammation, injury, infection, developmental abnormalities, or tumors.

### **Iris Leiomyoma**

A rare benign smooth-muscle tumor of the iris that may appear as a pigmented or nonpigmented mass. It can distort the pupil, impair vision, or resemble melanoma, so ophthalmic examination and imaging are needed.

### **Iris Melanoma**

Iris melanoma is a malignant tumor of pigment cells in the iris and may appear as a growing pigmented lesion, pupil distortion, or glaucoma.

### **Iris Neoplasm**

An iris neoplasm is an abnormal growth arising in the colored part of the eye; it may be benign or malignant and can affect vision or eye pressure.

### **Iris Prolapse**

Protrusion of iris tissue through a corneal or scleral wound, most often after penetrating eye trauma or surgery. It may compromise the globe and requires urgent ophthalmic management.

### **Iritis**

Inflammation localized to the iris, the colored ante-

rior portion of the uveal tract. It characteristically presents with deep, aching ocular pain, marked photophobia, blurred vision, ciliary flush (perilimbal injection), a constricted pupil (miosis), and inflammatory cells and flare in the anterior chamber. It is frequently associated with systemic autoimmune conditions, particularly HLA-B27-positive spondyloarthropathies, sarcoidosis, and viral infections.

#### **Alternate Names:**

Anterior Uveitis; Iridocyclitis

### **Iron**

A mineral needed to make hemoglobin, the protein that carries oxygen in red blood cells, and to support muscles and cellular energy. Low iron may result from blood loss, poor intake, increased needs, or poor absorption.

### **Iron Deficiency**

A shortage of iron needed to make hemoglobin and support normal cell function. It may result from blood loss, low intake, increased needs, or poor absorption and can cause tiredness, pale skin, shortness of breath, brittle nails, or unusual cravings.

### **Iron-Deficiency Anaemia**

Alternate spelling; see *Iron Deficiency Anemia*.

### **Iron-Deficiency Anaemia In Adolescents**

Iron-deficiency anaemia during adolescence, commonly related to rapid growth, inadequate dietary iron, or menstrual blood loss. Assessment should identify the cause and guide dietary advice and clinical treatment.

### **Iron Deficiency Anemia**

Anemia caused by too little iron to make normal hemoglobin. It may cause tiredness, weakness, pale skin, breathlessness, headache, or unusual cravings; treatment replaces iron and investigates blood loss, poor intake, or poor absorption.

#### **Alternate Names:**

IDA; Microcytic Hypochromic Anemia

---

## Iron-Deficiency Anemia

Alternate spelling; see *Iron Deficiency Anemia*.

## Iron-Deficiency Anemia in Children

Iron-deficiency anemia in a child caused by too little iron, increased needs during growth, blood loss, or poor absorption. It reduces hemoglobin production and may cause pale skin, tiredness, irritability, poor concentration, or delayed development when severe or long-lasting.

## Iron-Deficiency Anemia in Young Children

Iron-deficiency anemia in an infant or toddler caused by too little iron, rapid growth, excessive cow's milk intake, blood loss, or poor absorption. It may cause pale skin, tiredness, irritability, poor appetite, or developmental problems when severe or long-lasting.

## Iron Overdose

Poisoning after taking too much iron, especially from supplements. It can cause vomiting, abdominal pain, internal bleeding, severe acid imbalance, shock, liver failure, and death.

## Iron Overload

Excess iron stored in the body, which can damage the liver, heart, pancreas, joints, or endocrine organs.

## Iron Toxicity

Dose-related poisoning, usually after ingestion of iron tablets, causing vomiting, abdominal pain, gastrointestinal bleeding, lethargy, shock, high-anion-gap metabolic acidosis, liver failure, or coma. Serial serum iron levels, acid-base status, and organ function guide care; severe systemic poisoning may require deferoxamine and intensive support.

## Irradiated Foods

Irradiated foods are exposed to controlled ionizing radiation to reduce pathogens, insects, or spoilage; the process does not make food radioactive.

## Irradiation

Exposure to or treatment with radiation; irradiation may be used diagnostically or therapeutically, or may cause tissue injury depending on dose and type.

## Irradiation Fibrosis

Late-stage irreversible soft-tissue, pulmonary, or visceral scarring, collagen deposition, and microvascular obliterative endarteritis resulting from ionizing therapeutic radiation injury, mediated by sustained transforming growth factor-beta (TGF-beta) activation.

## Irregular

Not following the usual pattern, rhythm, shape, timing, or order. In medicine, it may describe an uneven heartbeat, an unpredictable menstrual cycle, an unusual skin edge, or a result that does not match the expected pattern.

## Irregular Menstruation

Menstrual bleeding that is unusually early, late, frequent, infrequent, prolonged, heavy, or unpredictable for the person's age and reproductive context. Causes include pregnancy, ovulatory disorders, endocrine disease, medicines, structural lesions, and perimenopause.

## Irregular Respiration

An abnormal breathing pattern in rate, depth, rhythm, or pauses. It may occur with sleep disorders, metabolic disturbance, lung disease, drug effects, or neurologic injury and may be accompanied by cyanosis or reduced consciousness.

## Irregular Sleep-Wake Syndrome

A sleep disorder in which sleep and wakefulness occur in several poorly timed periods throughout the day and night instead of one regular nighttime sleep. It is more common with brain disease, dementia, aging, or limited exposure to daylight and daily routines.

## Irreversible Cell Injury

Damage so severe that a cell cannot recover and eventually dies. It may follow prolonged lack of

---

oxygen, poisoning, infection, heat, or other injury and can lead to loss of function in the affected tissue.

### **Irrigate**

To wash or flush a body part, wound, cavity, or medical device with a fluid to remove debris, contaminants, or secretions.

### **Irrigation**

Flushing a wound, body cavity, or organ with fluid to remove blood, debris, microorganisms, or unwanted material.

### **Irrigation Solution**

A sterile or appropriately prepared liquid used to wash a wound, body cavity, organ, or surgical field. The solution must be suitable for the tissue because contamination, pressure injury, electrolyte shifts, or chemical irritation can cause harm.

### **Irritable Bowel Syndrome**

A disorder of gut–brain interaction that causes repeated abdominal pain and changes in bowel movements, such as diarrhea, constipation, or both. Pain is often related to passing stool and may occur with bloating or a feeling that the bowel has not emptied completely. IBS may be mainly constipation (IBS-C), mainly diarrhea (IBS-D), or a mixture of both (IBS-M). The gut may be unusually sensitive or its muscles may move food too quickly or too slowly. Unlike inflammatory bowel disease, IBS does not cause visible damage or disease in the digestive tract.

#### **Alternate Names:**

IBS

### **Irritable Bowel Syndrome Diet**

An eating plan adapted to reduce symptoms of irritable bowel syndrome while maintaining adequate nutrition. It may include a short trial of reducing fermentable carbohydrates followed by gradual food reintroduction.

### **Irritable Mood**

A mood marked by increased sensitivity, impatience, anger, or a tendency to become upset easily.

### **Irritability**

An increased tendency to become annoyed, impatient, angry, or easily upset. It may occur with stress, sleep loss, pain, mood disorders, substance use, medicines, infection, or other medical conditions.

### **Irritant Contact Dermatitis**

Inflammation of the skin caused by direct chemical or physical injury rather than an immune-mediated allergy. It commonly produces redness, burning, scaling, fissures, or pain at the area of exposure.

### **Irritants**

Substances that inflame or damage tissues and cause symptoms without necessarily triggering a true immune allergy. Examples include smoke, chemicals, soaps, fumes, and pollutants affecting skin, eyes, airways, or digestion.

### **Irritation**

A local unpleasant or inflammatory response to mechanical, chemical, thermal, infectious, or allergic stimuli. It may cause redness, itching, burning, pain, cough, or tearing depending on the tissue involved.

### **Isaacs Syndrome**

A rare peripheral nerve hyperexcitability disorder causing continuous muscle-fibre activity, stiffness, cramps, fasciculations, myokymia, and delayed muscle relaxation. It may be autoimmune, paraneoplastic, or associated with inherited or acquired peripheral-nerve disorders.

### **Isavuconazole**

An antifungal medicine used for selected serious infections, including invasive aspergillosis and mucormycosis. It blocks production of a substance fungi need for their cell membranes and can cause liver problems or medicine interactions.

### **Isazophos**

Isazophos is an organophosphate pesticide that can inhibit acetylcholinesterase and cause cholinergic toxicity after exposure.

---

## Ischaemia

Inadequate blood flow and oxygen delivery to a tissue, usually because of narrowed or blocked blood vessels, causing dysfunction or injury.

## Ischaemic Heart Disease

Heart disease caused by reduced blood supply to the heart muscle, usually as a result of coronary artery disease; it may cause angina or myocardial infarction.

## Ischemia

Reduced blood flow causing too little oxygen for a tissue's needs. It can cause pain, impaired function, permanent injury, or tissue death and may affect the heart, brain, limbs, or bowel.

## Ischemia-Reperfusion Injury

Damage that can occur when blood flow returns to tissue that has been deprived of oxygen. The returning blood can trigger inflammation and cell injury, affecting the heart, brain, limbs, or transplanted organs.

## Ischemic Bone Necrosis

Alternate name; see *Avascular Necrosis*.

## Ischemic Cardiomyopathy

Weakening of the heart muscle caused by long-term reduced blood flow or previous heart attacks from coronary artery disease. Scarring can reduce pumping ability and increase the risk of heart failure and dangerous rhythms.

## Ischemic Cascade

The sequential biochemical and cellular pathophysiological events triggered by acute arterial occlusion and tissue ischemia, beginning with ATP depletion, failure of Na<sup>+</sup>/K<sup>+</sup>-ATPase pumps, membrane depolarization, excessive intracellular calcium influx, phospholipase and protease activation, free radical generation, and apoptotic or necrotic cell death.

## Ischemic Chest Pain

Alternate name; see *Angina Pectoris*.

## Ischemic Cholangiopathy

Damage to the bile ducts caused by inadequate blood flow, usually through the hepatic artery. It can cause bile-duct injury, bile blockage, repeated cholangitis, or narrowing of several bile ducts. It is especially important after liver transplantation or injury to the hepatic artery.

## Ischemic Colitis

Inflammation and injury of the colon caused by reduced blood flow. It often causes sudden abdominal pain, bloody diarrhea, or rectal bleeding and commonly affects areas such as the splenic flexure.

### Alternate Names:

Colon Ischemia; Ischemic Bowel Disease of the Colon

## Ischemic Fasciitis

A noncancerous mass-like reaction in deep soft tissue caused by pressure and reduced blood flow, usually over a bony area in a person who is frail or has limited mobility. It can resemble a sarcoma on examination or imaging.

## Ischemic Gangrene

Death of tissue after a prolonged severe loss of blood supply, most often in a toe, foot, finger, or leg. The tissue may become cold, numb, and black, with risk of infection.

## Ischemic Heart Disease

Heart disease caused by inadequate blood flow through narrowed or blocked coronary arteries. It may cause chest pressure, breathlessness, heart attack, rhythm problems, heart failure, or sudden death.

### Alternate Names:

Coronary artery disease; coronary heart disease

## Ischemic Hepatitis

Widespread liver-cell injury caused by too little blood flow or oxygen. It commonly occurs with severe heart failure, very low blood pressure, respiratory failure, or severe infection. Liver enzymes usually rise suddenly and greatly. Despite its name, it is mainly liver-cell injury rather than inflammation.



---

### Ischemic Optic Neuropathy

Damage to the optic nerve caused by reduced blood flow, often producing sudden, usually painless vision loss. It may be arteritic or non-arteritic; older adults may also have headache, jaw pain, or scalp tenderness.

### Ischemic Preconditioning

An intrinsic cardioprotective phenomenon in which brief, sub-lethal episodes of myocardial ischemia and reperfusion render the myocardium resistant to subsequent prolonged, potentially lethal ischemic insult, significantly reducing ultimate infarct size.

### Ischemic Priapism

Alternate name; see *Priapism*.

### Ischemic Stroke

Sudden brain injury caused by a blocked blood vessel that reduces oxygen to part of the brain. It may cause one-sided weakness, facial drooping, or speech or vision problems.

### Ischiorectal Abscess

A deep perianal suppurative infection occupying the ischioanal (ischiorectal) fossa between the anal sphincter complex and the obturator internus muscle, typically arising from an infected cryptoglandular anal gland and presenting with severe throbbing perianal pain, induration, and systemic fever requiring surgical incision and drainage.

#### Alternate Names:

Ischioanal Abscess; Deep Perianal Abscess

### Ischium

The lower, back part of each hip bone. Its rounded lower area bears weight when sitting, and nearby parts provide attachment for the hamstring muscles and other pelvic structures.

### Ishak Score

A scoring system used on a liver biopsy to describe inflammation and scarring, especially in chronic hepatitis. Higher scores generally indicate more extensive damage, but the result must be interpreted

with other tests and the person's history.

### Islet Cell

A cell in a pancreatic islet that helps control blood glucose; beta cells make insulin and alpha cells make glucagon.

### Islet Cell Cancer

Alternate name; see *Pancreatic Neuroendocrine Tumors*.

### Islets Of Langerhans

Small clusters of hormone-producing cells in the pancreas that release insulin, glucagon, and other hormones regulating blood sugar and metabolism.

### Isoaminile

Isoaminile is an older sympathomimetic appetite suppressant that is no longer widely used because of cardiovascular and other adverse effects.

### Isoantibodies

Antibodies made against antigens from another individual of the same species, such as antibodies involved in blood-group incompatibility.

### Isoantigen

An antigen that differs between individuals of the same species and can trigger an immune response after exposure.

### Isocarboxazid

Isocarboxazid is a monoamine oxidase inhibitor used for depression; it has important interactions with other medicines and tyramine-rich foods.

### Isochromosomes

Abnormal chromosomes with two identical arms caused by division across the wrong axis of the centromere.

### Isocitrate Dehydrogenase

An enzyme involved in how cells produce energy. Changes in the IDH1 or IDH2 genes can alter this enzyme and are found in some brain tumors, blood

---

cancers, and other tumors, helping guide diagnosis and treatment.

### **Isocyanate**

A reactive chemical group found in some industrial products; exposure can irritate the airways and cause occupational asthma or skin reactions.

### **Isoelectric Point**

The acidity level at which a molecule, especially a protein, has no overall electrical charge. This property affects how the molecule dissolves and moves in laboratory tests.

### **Isoenzyme**

One of several forms of an enzyme that catalyze the same chemical reaction but differ in structure, location, or regulation. Measuring selected isoenzymes can help identify the tissue affected by injury.

### **Isofenphos**

A poisonous organophosphate insecticide. Exposure can cause sweating, excess saliva, vomiting, diarrhea, small pupils, muscle twitching, breathing difficulty, seizures, or weakness because it disrupts nerve signals.

### **Isoflurane**

Isoflurane is an inhaled anesthetic used to maintain general anesthesia; it can lower blood pressure and depress breathing.

### **Isogeneic Graft**

A transplant between genetically identical individuals, such as identical twins, so immune rejection risk is relatively low.

### **Isohemagglutinins**

Naturally occurring IgM antibodies present in human serum directed against ABO blood group carbohydrate antigens that are absent on the individual's own red blood cells (anti-A in group B, anti-B in group A, and both in group O), responsible for acute ABO-incompatible hemolytic transfusion reactions.

### **Isokinetic Exercise**

Exercise in which a machine keeps a joint moving at a set speed while the muscles provide resistance. It is used mainly to measure or build strength during rehabilitation.

### **Isolate**

To separate a person, organism, tissue, or substance from others; infection-control isolation separates people who may transmit a pathogen.

### **Isolated Systolic Hypertension**

Elevated systolic blood pressure with normal diastolic pressure, commonly related to arterial stiffness in older adults and associated with increased cardiovascular risk.

### **Isolation**

Separating a person with a contagious infection from others to limit spread, using appropriate rooms, protective equipment, and infection-control procedures.

### **Isolation Precautions**

Measures used with standard precautions to reduce transmission of infection. Additional precautions may be contact, droplet, or airborne, depending on the mode of spread.

### **Isoleucine**

An essential amino acid obtained from food that helps build proteins, support energy production, maintain muscle, and repair body tissues.

### **Isomerase**

An enzyme that rearranges atoms within a molecule to form an isomer. Isomerases participate in metabolism, and inherited defects in related pathways can cause metabolic disease.

### **Isometric**

Describing muscle contraction that produces force without a meaningful change in muscle length or joint movement.

---

## **Isometropia**

The condition in which both eyes have approximately equal refractive power, so neither eye has substantially more near-sightedness, far-sightedness, or astigmatism than the other.

## **Isoniazid**

Isoniazid is a first-line antituberculous drug that inhibits mycolic-acid synthesis in *Mycobacterium tuberculosis*; hepatotoxicity and peripheral neuropathy are important risks, and pyridoxine is often given.

## **Isoniazid Toxicity**

Harm from excessive or unusual sensitivity to isoniazid, especially liver injury, nerve damage, or seizures. Treatment may include vitamin B6, supportive care, activated charcoal, and seizure control.

## **Isoprenaline**

Isoprenaline, also called isoproterenol, is a beta-adrenergic medicine that can increase heart rate and is used in selected emergency cardiac situations.

## **Isoprocab**

Isoprocab is a carbamate insecticide that can cause cholinergic poisoning if swallowed or absorbed in a significant amount.

## **Isopropanol Alcohol Poisoning**

Poisoning from isopropanol, often found in rubbing alcohol, causing intoxication, vomiting, abdominal pain, low blood pressure, and sometimes coma.

## **Isopropyl Alcohol**

A volatile alcohol used as a disinfectant and solvent. Ingestion can cause rapid intoxication, vomiting, abdominal pain, hypotension, and coma; management is supportive and differs from treatment of methanol or ethylene-glycol poisoning.

## **Isoproterenol**

Isoproterenol, also called isoprenaline, is a beta-adrenergic medicine that increases heart rate and contractility and is used in selected emergency set-

tings.

## **Isosorbide**

Isosorbide is a nitrate medicine that relaxes blood vessels and is used mainly to prevent or relieve angina; headache and low blood pressure are common adverse effects.

## **Isosorbide Dinitrate**

A nitrate medicine that widens blood vessels and reduces the heart's workload. It is used mainly to prevent angina attacks; headache, dizziness, and low blood pressure may occur, and it can interact dangerously with erectile-dysfunction medicines.

## **Isospora**

An older name for a group of intestinal coccidian parasites, now generally classified as *Cystoisospora*. Infection can cause watery diarrhea, especially in people with weakened immunity.

## **Isospora Belli**

The former name for *Cystoisospora belli*, a parasite spread by swallowing oocysts in contaminated food or water. It causes watery diarrhea, cramps, nausea, and weight loss and can be severe or prolonged in immunocompromised people.

## **Isosporiasis**

Older name for *Cystoisosporiasis*, an intestinal infection caused by the parasite *Cystoisospora belli*.

## **Iso-Sulfan Blue**

A blue dye used to map lymphatic drainage or identify sentinel lymph nodes during selected surgical procedures.

## **Isotonic**

Describing muscle contraction in which muscle length changes to move a joint under a relatively constant external load.

## **Isotonic Exercise**

Exercise in which muscles change length while producing force against a relatively constant load, as

---

in lifting or lowering a weight.

### Isotonic Solution

A solution with an effective osmotic concentration close to that of plasma, such as normal saline or balanced crystalloid. Isotonic fluids may expand extracellular volume, but choice and amount must be tailored because excess can cause fluid overload or electrolyte disturbance.

### Isotope

One of two or more atoms of the same element with the same number of protons but different numbers of neutrons.

### Isotretinoin

A vitamin-A-related medicine used for severe acne that has not improved with other treatment. It reduces skin-oil production and clogged pores but can cause birth defects, so pregnancy prevention and regular medical monitoring are essential.

### Isovaleric Acidemia

A rare inherited disorder in which the body cannot properly break down the amino acid leucine. Harmful substances build up and may cause poor feeding, vomiting, unusual sleepiness, low muscle tone, or a sweaty-feet odor, especially during illness or fasting. Newborn screening can detect it; treatment uses a controlled diet and prevents prolonged fasting.

#### Alternate Names:

IVA; Isovaleryl-CoA Dehydrogenase Deficiency

### Isovolumetric Contraction

The early phase of cardiac ventricular systole lasting from the closure of the atrioventricular (mitral and tricuspid) valves to the opening of the semilunar (aortic and pulmonic) valves, during which ventricular pressure rises steeply at a constant ventricular volume without ejection of blood.

#### Alternate Names:

Isovolumic Contraction Phase

### Isozyme

One of several molecular forms of an enzyme that

catalyze the same reaction but differ in structure, tissue distribution, or regulation.

### Ispaghula Husk

Ispaghula husk, or psyllium, is a soluble fiber that absorbs water and helps regulate stool consistency; adequate fluid is important when using it.

### Isradipine

Isradipine is a calcium-channel blocker used to lower blood pressure; it may cause flushing, headache, ankle swelling, or dizziness.

### ISUP Grade Group

A five-level system that grades prostate cancer by how abnormal the cancer looks under a microscope. Group 1 is the lowest risk, usually with a Gleason score of 6 or less. Groups 2 and 3 represent intermediate risk, while groups 4 and 5 represent high or highest risk. The result helps guide monitoring, surgery, radiation, or other treatment.

#### Alternate Names:

Grade Group; ISUP Prostate Grading System; WHO/ISUP Grade Group

### Itch

Itch, or pruritus, is an unpleasant skin sensation that provokes scratching and may arise from dermatitis, allergy, infection, medicines, kidney or liver disease, blood disorders, nerve disease, or systemic illness. Persistent generalized itch requires evaluation.

#### Alternate Names:

Pruritus

### Itching

An unpleasant skin sensation that makes a person want to scratch. It may be short-term or persistent and can result from skin disease, allergy, infection, medicines, nerve problems, or illness affecting organs such as the liver or kidneys.

### Itchy Ear

An itching sensation of the external ear or ear canal, which may result from dermatitis, allergy, infection, dryness, or irritation from objects placed in the ear.

---

## Itchy Skin

Alternate name; see *Pruritus*.

## Iterative Reconstruction

A computer method that repeatedly improves medical images by comparing calculated results with measured scan data. It can reduce image noise and sometimes allow lower radiation exposure, but usually requires more processing time.

## ITP

Abbreviation; see *Immune Thrombocytopenia*.

## Itraconazole

Itraconazole is a triazole antifungal that inhibits fungal ergosterol synthesis and treats several superficial and systemic mycoses; absorption, extensive drug interactions, liver injury, and negative inotropic effects require attention.

## IUCD

An intrauterine contraceptive device, usually referring to an intrauterine device placed in the uterus to prevent pregnancy.

## Ivabradine

Ivabradine slows the heart rate by acting on the sinus node and is used for selected patients with chronic heart failure or stable angina.

## IVC Filter

A small device placed in the body's large lower vein to catch clots traveling from the legs to the lungs. It is used selectively when blood thinners cannot be used or have not prevented clots; blockage, movement, or vessel injury can occur.

## Ivermectin

An antiparasitic medicine used for selected worm and mite infections, including strongyloidiasis, onchocerciasis, and scabies, when prescribed appropriately.

## IVF

In vitro fertilization: the assisted reproductive tech-

nology in which oocytes are retrieved, fertilized by sperm outside the body (in vitro), and the resulting embryos transferred to the uterus. Indications: tubal factor, male factor (with ICSI), endometriosis, unexplained, ovulation disorders, genetic testing, and same-sex couples or individuals using donor sperm or eggs.

## IV Fluids

Fluids given through a vein to replace lost water and salts, support circulation, or provide maintenance during illness. The type and amount depend on the person's blood pressure, kidney and heart function, electrolytes, and cause of fluid loss.

## IV Treatment At Home

Intravenous treatment given outside a hospital, usually through a catheter with instructions for safe infusion, line care, monitoring, and follow-up.

## Ixabepilone

Ixabepilone is a chemotherapy medicine used for some advanced breast cancers; important risks include nerve damage, low blood counts, and allergic reactions.

## Ixodes

A genus of hard ticks, including the ticks that transmit Lyme disease and several other infections. A bite may be painless; removing the tick promptly reduces, but does not eliminate, infection risk.

## IBS (Irritable Bowel Syndrome)

Alternate name; see *Irritable Bowel Syndrome*.

## IC

Abbreviation for interstitial cystitis, also called bladder pain syndrome; see *Interstitial Cystitis*.

## Iceland Disease

Historical name for epidemic neuromyasthenia, now generally classified among outbreaks of myalgic encephalomyelitis/chronic fatigue syndrome.

---

### **Icelandic Type Amyloidosis (Type VI)**

Hereditary cerebral amyloid angiopathy caused by cystatin C amyloid deposition in cerebral blood vessels; it can cause recurrent brain hemorrhage and neurologic disability.

### **Icterus (Chronic Familial)**

Alternate name for Gilbert syndrome, an inherited disorder causing intermittent mild unconjugated hyperbilirubinemia; see *Gilbert Syndrome*.

### **Idiopathic Amyloidosis**

Amyloidosis without an identified underlying cause after appropriate evaluation; the clinical manifestations depend on the organs involved and the amyloid protein.

### **Idiopathic Aplastic Anemia**

Aplastic anemia in which no specific cause is identified after evaluation; see *Aplastic Anemia*.

### **Idiopathic Autoimmune Hemolytic Anemia**

Autoimmune hemolytic anemia without an identified associated disease, medicine, or other trigger; see *Autoimmune Hemolytic Anemia*.

### **Idiopathic Basal Ganglia Calcification**

Calcification of the basal ganglia without an identified secondary cause, historically called Fahr disease; symptoms may include movement, psychiatric, cognitive, or seizure disorders.

### **Idiopathic Chronic Erosive Gastritis**

Long-standing erosive inflammation of the stomach lining without an identified cause; see *Erosive Gastritis*.

### **Idiopathic Dilated Cardiomyopathy**

Dilated cardiomyopathy without an identified cause after evaluation; see *Dilated Cardiomyopathy*.

### **Illinois Type Amyloidosis**

Historical or imprecise label for primary, usually immunoglobulin light-chain amyloidosis; the type of

deposited protein should be confirmed rather than inferred from the geographic name.

### **Immunizations**

Alternate plural form; see *Immunization*.

### **Immunoblastic Lymphadenopathy**

Historical term for a systemic immune-reactive lymph-node disorder, now generally classified within other lymphoproliferative or immune disorders according to biopsy findings.

### **Immunodeficiency with Ataxia Telangiectasia**

Alternate name; see *Ataxia Telangiectasia*.

### **Impotence (Erection Problems)**

Alternate name for erectile dysfunction, the persistent inability to obtain or maintain an erection adequate for sexual activity; see *Erectile Dysfunction*.

### **Impotence**

Older term; see *Erectile Dysfunction*.

### **Incision Care After Surgery**

Care of a surgical wound to keep it clean and protected, monitor for infection or separation, and follow instructions about dressing changes, bathing, activity, and when to seek medical advice.

### **Incontinence in Men**

Urinary incontinence in men, which may result from prostate disease or treatment, bladder dysfunction, neurologic disease, infection, or other causes; evaluation depends on symptoms and associated findings.

### **Incontinence/OAB**

Urinary incontinence associated with overactive bladder symptoms such as urgency, frequency, or nocturia; see *Overactive Bladder* and *Urinary Incontinence*.

### **Indiana Type Amyloidosis (Type II)**

Hereditary transthyretin amyloid polyneuropathy



---

type II, also called Indiana/Swiss type, caused by a pathogenic transthyretin variant and often involving peripheral nerves.

### **Indolent Systemic Mastocytosis**

A usually slow-progressing form of systemic mastocytosis in which abnormal mast cells accumulate in organs such as bone marrow, skin, or digestive tract; see *Systemic Mastocytosis*.

### **Infantile Gaucher Disease**

Gaucher disease presenting in infancy, often with severe neurologic and systemic involvement; see *Gaucher Disease*.

### **Infantile Myoclonic Encephalopathy**

A severe early-life epileptic encephalopathy with recurrent myoclonic seizures and developmental impairment; diagnosis and classification require specialist neurologic evaluation.

### **Infantile Sleep Apnea**

Repeated pauses or reductions in an infant's breathing during sleep; causes include immature breathing control, airway obstruction, lung disease, neurologic disease, or other illness and require age-appropriate assessment.

### **Infantile Spinal Muscular Atrophy**

Spinal muscular atrophy beginning in infancy, causing progressive weakness from loss of motor neurons; see *Spinal Muscular Atrophy*.

### **Infantile Spinal Muscular Atrophy Type I**

The severe infantile form of spinal muscular atrophy, usually beginning in the first months of life with marked weakness and low muscle tone; early disease-modifying treatment and respiratory and nutritional support are important.

### **Infectious Chorea**

Chorea caused by an infection or post-infectious immune response, such as Sydenham chorea after group A streptococcal infection; see *Chorea*.

### **Inflammatory Bowel Disease/Crohn's Disease**

Alternate consumer label for inflammatory bowel disease, especially Crohn disease; see *Inflammatory Bowel Disease* and *Crohn Disease*.

### **Inherited Lipodystrophy**

A genetic disorder causing partial or generalized loss or abnormal distribution of body fat, often with severe insulin resistance, diabetes, high triglycerides, and fatty liver; see *Lipodystrophy*.

### **Iron Overload Anemia**

Anemia associated with excess iron accumulation, often in disorders such as sideroblastic anemia or repeated transfusion; see *Iron Overload* and *Sideroblastic Anemia*.

### **Irregular Menstrual Periods**

Menstrual bleeding that is unusually unpredictable, infrequent, frequent, prolonged, or variable in amount; causes include pregnancy, ovulatory disorders, endocrine disease, medications, structural disease, and perimenopause.

### **Irritable Bowel Syndrome (IBS)**

Alternate name; see *Irritable Bowel Syndrome*.

### **Ischemic Necrosis of Bone**

Bone death caused by inadequate blood supply, also called osteonecrosis or avascular necrosis; see *Avascular Necrosis*.

### **Isoleucine 33 Amyloidosis**

A rare hereditary transthyretin amyloidosis variant involving an isoleucine substitution at position 33; clinical features depend on whether nerve, heart, or other organs are affected.



# J

## Jaccoud Arthropathy

A usually non-damaging joint disorder that can pull the fingers toward the little-finger side. The deformity may improve when the hand is moved and is often linked with lupus or past rheumatic fever.

## Jaccoud's Arthropathy

A deforming but usually non-damaging form of arthritis that can cause the fingers to drift toward the little-finger side. The deformity may improve when the hand is moved and is often linked to lupus or past rheumatic fever.

## Jack-In-The-Pulpit Poisoning

Jack-in-the-pulpit poisoning follows ingestion of calcium oxalate crystals from the plant, causing intense burning and swelling of the mouth, throat, and gastrointestinal tract.

## Jackknife Position

A face-down position used during some operations in which the table is bent at the hips. The buttocks are raised while the head and legs are lower, improving access to the rectal, anal, or posterior pelvic regions.

### Alternate Names:

Kraske Position

## Jacksonian Epilepsy

A focal seizure that begins with twitching or stiffening in one body part and spreads in an orderly way to nearby body parts. Awareness may remain normal at first.

### Alternate Names:

Jacksonian March; Focal Motor Seizure

## Jacksonian March

A clinical progression of focal motor seizure activity in which epileptic discharges propagate along contiguous areas of the primary motor cortex (precentral gyrus homunculus), resulting in clonic contractions that begin focally in one anatomical region (such as the thumb or angle of the mouth) and march sequentially to involve adjacent muscle groups of the limb or face.

### Alternate Names:

Jacksonian Seizure; Focal Motor March

## Jacksonian Seizure

A focal motor seizure that begins in one body part and may spread in an orderly pattern to adjacent body regions; awareness may remain intact at first.

## Jacobsen Syndrome

A rare chromosome condition caused by loss of part of chromosome 11. It may cause low platelets, unusual facial features, heart defects, bleeding, and developmental delay.

### Alternate Names:

11q Terminal Deletion Syndrome; Jacobsen Deletion Disorder

## Jadassohn-Lewandowsky Syndrome

A form of pachyonychia congenita, an inherited condition causing very thick, abnormal nails, painful thick skin on the palms and soles, and sometimes white patches in the mouth or rough bumps around hair follicles.

### Alternate Names:

Pachyonychia Congenita Type 1; PC-1

---

### Jadassohn's Naevus

A hairless skin patch or raised lesion present from birth, usually caused by an overgrowth of oil glands. It may change during life and rarely develops another tumor.

### Jaffe-Lichtenstein Syndrome

A form of fibrous dysplasia affecting several bones without the hormone problems or café-au-lait skin marks seen in McCune-Albright syndrome. Affected bones may be weak, painful, or misshapen.

#### Alternate Names:

Polyostotic Fibrous Dysplasia; Jaffe-Lichtenstein Disease

### JAK2

A protein inside cells that passes signals from certain hormones and immune chemicals. An activating change in the JAK2 gene can make the bone marrow produce too many blood cells, as in several myeloproliferative disorders.

### JAK-STAT Pathway

A signaling system cells use to respond to hormones and immune chemicals. It helps control growth, blood-cell production, inflammation, and immune activity; abnormal signaling can contribute to inflammatory disease or cancer.

### Jammed Finger

A finger injury caused by sudden force, often involving a sprain, joint dislocation, fracture, or tendon injury. It may cause pain, swelling, deformity, or difficulty moving the finger.

### Janeway Lesions

Small, usually painless red or purple spots on the palms or soles that can occur with infective endocarditis, an infection of the heart lining or valves. They result from infected particles blocking tiny blood vessels.

### Janus Kinase

One of a family of proteins inside cells that pass signals from hormones and immune chemicals. Janus

kinases help control inflammation, blood-cell production, and immune responses.

### Janus Kinase Inhibitors

Medicines that block Janus kinase proteins and reduce certain immune signals. They are used for selected inflammatory and blood disorders, but can increase the risk of infections, blood clots, and other serious effects.

### Japanese Encephalitis

A viral infection spread by mosquitoes that can inflame the brain. Most infections cause no symptoms, but severe illness can cause fever, confusion, seizures, movement problems, or lasting disability.

### Jarcho-Levin Syndrome

A rare birth condition in which the bones of the spine and ribs do not form normally. It can cause a short, stiff chest, short stature, breathing problems, and spinal curvature.

#### Alternate Names:

Spondylocostal Dysostosis; Spondylothoracic Dysplasia

### Jargon Aphasia

A type of language disorder in which speech is fluent but filled with made-up words, incorrect sounds, or sentences that are hard to understand. Understanding spoken or written language is also impaired.

#### Alternate Names:

Neologistic Aphasia; Word Salad Speech

### Jarisch-Herxheimer Reaction

A short-lived reaction that may occur soon after antibiotics are started for certain infections. It can cause fever, chills, headache, muscle aches, and temporarily worse symptoms.

### Jatene Procedure

An operation for babies born with the major arteries leaving the wrong heart chambers. The aorta and pulmonary artery are switched, and the coronary arteries are reattached.

#### Alternate Names:

---

Arterial Switch Operation; ASO

### **Jaundice**

Yellowish pigmentation of the skin, sclerae, and mucous membranes caused by elevated levels of circulating bilirubin in the blood. Etiologies include pre-hepatic hemolytic processes, intra-hepatic hepatocellular injury (hepatitis, cirrhosis), or post-hepatic biliary obstruction (gallstones, strictures, neoplasms).

#### **Alternate Names:**

Icterus; Hyperbilirubinemic Scleral Icterus; Biliary Pigmentation

### **Jaundice And Breastfeeding**

Jaundice associated with breastfeeding may reflect inadequate milk intake in the early days or persistent breast-milk jaundice after other serious causes have been excluded.

### **Jaundice Causes**

Jaundice can result from rapid breakdown of red blood cells, inherited bilirubin problems, liver disease, medicines or toxins, or blockage of bile flow. Blood tests and imaging help identify the cause and guide treatment.

### **Jaundice In Adults**

Yellowing of the skin or eyes in an adult from increased bilirubin. Causes include red-cell breakdown, liver injury, impaired bile flow, or inherited conditions. Dark urine, pale stool, abdominal pain, fever, confusion, or rapid onset may occur.

### **Jaundice In Neonates**

Yellowing of a newborn's skin or eyes from increased bilirubin. It is often temporary, but jaundice in the first 24 hours, rapid worsening, poor feeding, unusual sleepiness, fever, pale stools, or dark urine may occur with more serious disease.

### **Jaundice In Newborns**

Yellowing of a newborn's skin or eyes caused by increased bilirubin. It is often temporary, but very high or rapidly rising levels can injure the brain.

### **Jaw**

The upper and lower bony structures of the mouth—the maxilla and mandible—which support the teeth and contribute to chewing, speech, facial shape, and the oral airway.

### **Jaw Fracture**

A break in the lower jawbone or upper jawbone, usually after a blow to the face. Signs may include pain, swelling, loose teeth, numbness, a changed bite, or difficulty opening the mouth. Severe breaks can block the airway, especially when the lower jaw is broken in the front on both sides. Treatment may require realignment, wiring, plates, or surgery.

### **Jaw Jerk Reflex**

A brief closing movement of the jaw when the chin is tapped with the mouth slightly open. An unusually strong response may suggest a problem in the brain's motor pathways.

### **Jaw Neoplasm**

An abnormal growth in the upper or lower jawbone or nearby tissues. It may be a cyst, a benign tumor, or cancer and can cause swelling, pain, loose teeth, numbness, or bone destruction.

### **Jaw Pain**

Pain in the jaw, jaw joint, or chewing muscles caused by temporomandibular disorders, dental disease, infection, injury, or less commonly heart or nerve disease. It may occur with chest discomfort, sweating, breathlessness, exertion, restricted movement, or persistent pain.

### **Jaw Thrust Maneuver**

An airway-opening technique that moves the lower jaw forward without bending the neck. It is used when a neck injury is possible.

### **Jaw Tumors And Cysts**

Abnormal growths or fluid-filled sacs developing in the jawbone or nearby tissues. They may be harmless or cancerous and can cause swelling, pain, tooth movement, numbness, or bone destruction.

---

**Alternate Names:**

Odontogenic Tumors And Cysts

**Jaw-Winking Syndrome**

A congenital synkinetic neuro-ophthalmic condition (Marcus Gunn phenomenon) characterized by unilateral congenital ptosis that transiently resolves or retracts upward with movement of the lower jaw during mastication, sucking, or lateral pterygoid contraction, caused by anomalous congenital innervation of the levator palpebrae superioris by the motor branch of the trigeminal nerve (CN V3).

**Alternate Names:**

Marcus Gunn Jaw-Winking Phenomenon; Trigemino-Oculomotor Synkinesis

**Jehovah's Witness**

A member of a Christian religious group whose beliefs may lead them to decline transfusions of whole blood and certain blood components. Individual choices about blood products and medical procedures vary.

**Jejunal Atresia**

A condition present at birth in which part of the jejunum, a section of the small intestine, is closed or missing. It causes bowel blockage in a newborn, with green vomiting, a swollen abdomen, and failure to pass the first stool.

**Jejunal Diverticulum**

A pouch that forms in the wall of the jejunum, the middle part of the small intestine. It is often symptom-free but can be linked with bacterial overgrowth, poor absorption, inflammation, or bleeding.

**Alternate Names:**

Jejunal Diverticulosis

**Jejunal Fistula**

An abnormal connection involving the jejunum and another organ, the skin, or a body surface. It may cause intestinal contents to leak, infection, malnutrition, dehydration, or sepsis.

**Jejunal Hemorrhage**

Bleeding from the jejunum, the middle segment of the small intestine, which may cause maroon or black stool, anemia, abdominal pain, or shock depending on its rate and cause.

**Jejunal Neoplasm**

An abnormal growth in the jejunum, the middle part of the small intestine. It may be benign or cancerous and can cause pain, bleeding, anemia, vomiting, blockage, or weight loss.

**Jejunal Stenosis**

Narrowing of the jejunal lumen that can impede intestinal contents and cause cramping, vomiting, distension, malabsorption, or obstruction; causes include Crohn disease, adhesions, tumors, and ischemic injury.

**Jejunal Ulcer**

An ulcerated break in the lining of the jejunum, caused by inflammation, infection, ischemia, medicines, tumor, or anastomotic disease and potentially producing pain, bleeding, or perforation.

**Jejunitis**

Jejunitis is inflammation of the jejunum, which may result from infection, inflammatory bowel disease, ischemia, medication, or other intestinal disorders.

**Jejunioileal Atresia**

A birth defect in which part of the small intestine is completely closed or missing, so milk and digestive fluid cannot pass. Newborns may develop green vomiting, a swollen abdomen, and failure to pass stool. Imaging shows the blockage, and surgery restores intestinal passage.

**Alternate Names:**

Jejunal Atresia; Ileal Atresia; Small Bowel Atresia

**Jejunojejunostomy**

A surgical anastomosis created between two segments of the jejunum, routinely performed in Roux-en-Y gastric bypass and Roux-en-Y gastrojejunostomy reconstruction to restore continuity between



---

the biliopancreatic limb and the alimentary limb.

### Jejunostomy

A surgically created opening into the jejunum, often used for placement of a feeding tube when nutrition cannot safely be given through the stomach.

### Jejunostomy Feeding Tube

A jejunostomy feeding tube delivers nutrition directly into the jejunum when oral or gastric feeding is unsafe or inadequate, with attention to formula, rate, position, and complications.

### Jejunum

The middle part of the small intestine. It absorbs much of the nutrients from food before the remaining contents pass into the ileum.

### Jellyfish Sting

A painful skin reaction caused when a jellyfish tentacle releases venom into the skin. It may cause burning, redness, and raised lines; severe exposure can cause vomiting, breathing trouble, fainting, or widespread symptoms.

#### Alternate Names:

Jellyfish Envenomation; Sea Sting

### Jellyfish Stings

Alternate name; see *Jellyfish stings*.

### Jendrassik Maneuver

A clinical examination technique used to reinforce and elicit sluggish deep tendon reflexes (such as the patellar tendon reflex) by having the patient hook their flexed fingers together and pull forcefully apart, which increases spinal motoneuron pool excitability by reducing descending central inhibition.

#### Alternate Names:

Jendrassik Reinforcement; Reflex Reinforcement Maneuver

### Jendrassik's Maneuver

A technique used during a reflex examination in which a person pulls the hands apart or tenses another muscle group. The distraction can make a

knee or other tendon reflex easier to observe.

### Jerk Nystagmus

Repeated involuntary eye movements with a slow movement in one direction followed by a quick movement back. It may result from problems in the balance organs, cerebellum, or brainstem.

### Jerusalem Cherry Poisoning

Jerusalem cherry poisoning follows ingestion of *Solanum pseudocapsicum* berries, which contain toxic compounds that may cause gastrointestinal symptoms, confusion, weakness, or cardiac effects.

### Jervell and Lange-Nielsen Syndrome

A rare inherited condition causing severe hearing loss from birth and an abnormal heart rhythm. The rhythm problem can cause fainting, seizures, or sudden death.

### Jet Lag

Temporary circadian rhythm disruption after rapid travel across time zones, causing sleep disturbance, fatigue, impaired concentration, and sometimes gastrointestinal symptoms.

#### Alternate Names:

Desynchronosis; Time Zone Change Syndrome

### Jet Lag Prevention

Jet lag prevention uses timed light exposure, sleep adjustment, hydration, and carefully planned activity or melatonin strategies to shift the circadian rhythm across time zones.

### Jeune Syndrome

A rare inherited bone disorder in which the chest is unusually narrow and the limbs are short. It may cause breathing problems, extra fingers or toes, and kidney or liver disease.

#### Alternate Names:

Asphyxiating Thoracic Dystrophy; ATD

### Jimsonweed Poisoning

Jimsonweed poisoning causes anticholinergic toxicity, including dilated pupils, dry mouth, rapid heart

---

rate, urinary retention, agitation, delirium, seizures, or coma.

### **Job Syndrome**

A rare inherited immune disorder causing eczema, repeated skin and lung infections, and very high levels of an allergy-related antibody. Dental, bone, and facial differences may also occur.

### **Jock Itch**

A fungal infection of the groin and inner thighs, also called tinea cruris. It causes an itchy, scaly, ring-shaped or spreading rash, often worsened by warmth and moisture. Pain, fever, extensive involvement, or persistent rash may occur with another condition.

#### **Alternate Names:**

Tinea Cruris; Ringworm of the Groin

### **Jod-Basedow Phenomenon**

Overactive thyroid function triggered by an excess of iodine, usually in a person with thyroid nodules or other thyroid tissue that works independently of normal control. It may cause weight loss, tremor, sweating, or a fast heartbeat.

### **Joffroy Sign**

A physical examination sign of Graves disease and hyperthyroidism characterized by the absence of normal forehead wrinkling and frowning when the patient looks upward while keeping the head in a fixed forward position.

#### **Alternate Names:**

Joffroy's Sign; Thyrotoxic Forehead Sign

### **Johanson-Blizzard Syndrome**

A rare autosomal recessive congenital malformation syndrome caused by mutations in the UBR1 gene, characterized by exocrine pancreatic insufficiency, hypoplasia or aplasia of the nasal alae ('beaked nose'), scalp vertex defects, oligodontia, sensorineural hearing loss, and growth retardation.

#### **Alternate Names:**

JBS; Pancreatic Insufficiency-Alar Hypoplasia Syndrome

### **Joint**

The place where two or more bones meet. Joints may be fixed, slightly movable, or freely movable, such as the knee, hip, or shoulder.

### **Joint Aspiration**

A needle procedure that removes fluid from a joint for testing or symptom relief, helping assess infection, bleeding, crystals, or inflammation.

### **Joint Capsule**

A tough envelope surrounding a freely movable joint. Its inner lining makes slippery fluid that helps the joint move, while the capsule helps hold the bones together and provides stability.

### **Joint Contracture**

Permanent or long-lasting limitation of joint movement caused by tightening or scarring of muscles, tendons, ligaments, the joint capsule, skin, or nearby tissues. It may follow immobility, injury, burns, inflammation, or nerve disease.

### **Joint Crepitus**

An audible or palpable grating, cracking, or crunching sensation produced by the movement of a joint against resistance, resulting from the friction of roughened articular cartilage surfaces, osteophyte contact, or fractured bone fragments (as in osteoarthritis).

### **Joint Diseases**

Conditions that affect joints, including osteoarthritis, inflammatory arthritis, gout, infection, injury, and tumors. They may cause pain, swelling, stiffness, deformity, or reduced movement; treatment depends on the cause.

### **Joint Effusion**

Extra fluid inside a joint, often causing swelling, stiffness, or restricted movement. Causes include injury, infection, inflammatory arthritis, bleeding, and crystal disease; a clinician may remove fluid for testing.

---

## Joint Fluid Culture

Joint fluid culture tests synovial fluid for bacterial, fungal, or other organisms when septic arthritis is suspected; results are interpreted with cell count, crystals, and clinical findings.

## Joint Fluid Gram Stain

A quick laboratory test that stains fluid removed from a swollen joint to look for bacteria. It can support the evaluation of a serious joint infection but a negative result does not rule infection out.

## Joint Inflammation

Alternate name; see *Arthritis*.

## Joint Injection

A therapeutic or diagnostic procedure in which a medication (such as a long-acting corticosteroid, local anesthetic, or hyaluronic acid viscosupplement) is introduced directly into a synovial joint space under sterile conditions to relieve pain and reduce intra-articular inflammation.

### Alternate Names:

Intra-Articular Injection

## Joint Mouse

A loose piece of cartilage, bone, or other tissue inside a joint that can cause catching, locking, pain, or swelling.

## Joint Pain

Pain or aching in one or more joints or in the tissues around them; the medical term is arthralgia. It is a symptom rather than a single disease and may result from arthritis, inflammation, injury, overuse, infection, crystals such as those in gout, or problems in nearby muscles, tendons, or bursae. It may occur with stiffness, swelling, warmth, redness, tenderness, or reduced movement. The number of joints affected, the timing, and associated symptoms help identify the cause.

## Joint Prosthesis

An artificial joint or implant designed to replace a damaged or diseased biological joint, restoring

mobility and relieving chronic pain.

## Joint Replacement

Surgery that removes damaged joint surfaces and replaces them with artificial parts to reduce pain and improve movement. It is commonly performed for severe arthritis or joint destruction; rehabilitation often follows.

## Joint Replacement Rehabilitation

Exercise, education, and daily-activity training after joint replacement, helping restore safe movement, strength, balance, confidence, and independence.

## Joint Stiffness

Reduced ease or range of joint movement, often worse after rest. Causes include arthritis, injury, inflammation, prolonged immobility, or muscle and tendon disorders. It may occur with swelling, redness, fever, severe pain, or inability to use the joint.

## Joint Swelling

Enlargement of a joint caused by excess synovial fluid, blood, inflammation, infection, trauma, or structural disease. It may be accompanied by pain, warmth, stiffness, or restricted movement and can indicate inflammatory or septic arthritis.

## Joint X-Ray

An X-ray picture of a joint used to look for broken bones, abnormal alignment, arthritis-related wear, bone loss, deposits, or other changes. It uses a small amount of radiation and does not show all soft-tissue injuries.

## Jones Criteria

A set of findings used to classify suspected acute rheumatic fever after group A streptococcal infection. Major findings include heart inflammation, migrating joint inflammation, abnormal movements, a ring-shaped rash, and small lumps under the skin; minor findings provide additional support.

### Alternate Names:

Revised Jones Criteria; AHA Rheumatic Fever Criteria

---

## Jones Fracture

A break across the fifth metatarsal, the long bone leading to the little toe, near its base. This area has a limited blood supply, so healing may be slow.

### Alternate Names:

Fifth Metatarsal Watershed Fracture; Metaphyseal-Diaphyseal Fifth Metatarsal Fracture

## Jones Metatarsal Fracture

A break across the fifth metatarsal near its base, close to the little toe. This area has a limited blood supply, so healing may be slow or incomplete.

### Alternate Names:

Fifth Metatarsal Jones Fracture; Metaphyseal-Diaphyseal Fifth Metatarsal Fracture

## Jones Tube

A specialized Pyrex glass bypass tube surgically implanted from the medial canthus (caruncle) directly into the nasal cavity (conjunctivodacryocystorhinostomy) to establish tear drainage in patients with severe, irreversible obstruction of the lacrimal canaliculi.

## Joubert Syndrome

A rare inherited disorder of brain development that affects breathing control, balance, eye movements, and muscle tone. Babies may have unusual breathing, poor muscle tone, delayed development, and abnormal eye movements. Some also have kidney, liver, or retinal disease, or extra fingers or toes.

### Alternate Names:

JSRD; Joubert-Related Disorder

## Joule-Thomson Effect

The thermodynamic temperature change that occurs when a high-pressure real gas expands adiabatically through a throttling valve or porous plug into a lower-pressure environment without doing external work. In medical cryosurgery, rapid expansion of pressurized argon gas causes rapid cooling down to minus 140 degrees Celsius to induce lethal cellular cryoablation of tumors.

### Alternate Names:

Joule-Kelvin Effect; Joule-Thomson Cryoexpansion

## J Receptor

A small sensory nerve ending in the lung near the tiny blood vessels. It responds to fluid buildup or pressure in the lung and can contribute to rapid, shallow breathing and breathlessness.

### Alternate Names:

Juxtacapillary Receptor

## Jugular

Relating to the neck, especially the jugular veins that carry blood from the head and neck back to the heart.

## Jugular Bulb

A dilated anatomical expansion of the internal jugular vein: the superior bulb occupies the jugular fossa of the temporal bone immediately below the middle ear floor, while the inferior bulb lies near the termination of the internal jugular vein behind the sternal end of the clavicle.

## Jugular Foramen

An opening at the base of the skull through which a major neck vein and several nerves pass. Disease in this area can cause difficulty swallowing, hoarseness, shoulder weakness, hearing symptoms, or other neurologic problems.

## Jugular Foramen Syndrome

A clinical neurological syndrome resulting from compressive or infiltrative lesions (such as glomus jugulare tumors, schwannomas, or skull base metastases) affecting the structures traversing the jugular foramen, characterized by ipsilateral paralysis of the glossopharyngeal (CN IX), vagus (CN X), and accessory (CN XI) cranial nerves, leading to dysphagia, vocal cord paralysis, and trapezius weakness.

### Alternate Names:

Vernet Syndrome; Posterior Lacerated Foramen Syndrome

## Jugular Vein

A major vein that carries blood from the head, face, and neck back to the heart. Each side has an internal jugular vein and an external jugular vein.

---

### Jugular Venous Distension

Pathological engorgement and visible protrusion of the internal or external jugular veins in the neck above the level of the clavicle with the patient positioned at 45 degrees, providing a direct physical reflection of elevated right atrial and central venous pressure, classic for right heart failure, cardiac tamponade, and constrictive pericarditis.

#### Alternate Names:

JVD; Raised Jugular Venous Pressure

### Jugular Venous Distention

Visible elevation and engorgement of the internal jugular venous pulsation above the sternal angle of Louis (normally less than 3 to 4 centimeters vertical height at 45 degrees head elevation). It serves as a cardinal physical sign of elevated central venous pressure and right ventricular volume or pressure overload (as in right heart failure, constrictive pericarditis, or cardiac tamponade).

#### Alternate Names:

JVD; Elevated Jugular Venous Pressure; JVP Elevation

### Jugular Venous Pressure

The visible height of blood in the neck veins. It helps clinicians estimate pressure in the right side of the heart and assess fluid buildup.

#### Alternate Names:

JVP

### Jugular Venous Waveform

The pattern of rises and falls seen in the neck vein during a heartbeat. It reflects contraction and relaxation of the right side of the heart and filling of the vein.

#### Alternate Names:

JVP Waveform

### Jugulodigastric Node

A prominent deep cervical lymph node situated at the junction where the posterior belly of the digastric muscle crosses the internal jugular vein, receiving primary lymphatic drainage from the palatine tonsils, base of tongue, and posterior pharynx, frequently enlarged in acute pharyngotonsillitis and

head and neck neoplasms.

#### Alternate Names:

Tonsillar Lymph Node; Subdigastric Node

### Jumper's Knee

Alternate name; see *Patellar Tendinitis*.

### Junctional Bradycardia

A cardiac arrhythmia originating from the atrioventricular (AV) junction or bundle of His with an inherent firing rate slower than the normal junctional escape rate (fewer than 40 beats per minute), characterized by narrow QRS complexes and absent, inverted, or retrograde P waves.

### Junctional Ectopic Tachycardia

A form of supraventricular tachycardia originating from an abnormally automatic focus within the atrioventricular junction or His bundle, occurring most frequently in pediatric patients following corrective surgery for congenital heart defects (such as tetralogy of Fallot or ventricular septal defect). Characterized by atrioventricular dissociation and a rapid narrow QRS complex rate (170 to 260 beats per minute) that responds poorly to electrical cardioversion.

#### Alternate Names:

JET; Congenital Junctional Tachycardia

### Junctional Epidermolysis Bullosa

A severe inherited blistering disorder in which the layers of skin separate easily. It can cause widespread blisters, skin wounds, poor feeding, and other complications from birth.

#### Alternate Names:

JEB

### Junctional Escape Rhythm

A physiological backup cardiac rhythm originating from pacemaking cells within the atrioventricular (AV) junction and bundle of His at an intrinsic rate of 40 to 60 beats per minute, characterized by narrow QRS complexes with absent, inverted, or retrograde P waves, arising when sinoatrial node pacemaking fails or during complete AV nodal block.

---

**Alternate Names:**

AV Junctional Rhythm; Nodal Escape Rhythm

**Junctional Nevus**

A usually benign, often flat mole formed by pigment cells at the junction between the outer and deeper layers of skin. It may change in size, shape, color, or symptoms.

**Junctional Premature Complex**

An early ectopic cardiac depolarization arising prematurely within the atrioventricular junction, manifesting on electrocardiogram as a normal or narrow QRS complex occurring before the next expected sinus beat, often accompanied by inverted retrograde P waves before, during, or after the QRS.

**Alternate Names:**

PJC; Premature Junctional Contraction

**Junctional Rhythm**

A heart rhythm that starts near the connection between the upper and lower chambers instead of the heart's usual pacemaker. It may be slow or regular and can occur when the usual pacemaker is suppressed or blocked.

**Junctional Tachycardia**

A supraventricular arrhythmia arising from enhanced automaticity or triggered activity within the AV node or bundle of His, characterized by a rapid heart rate (greater than 100 beats per minute), regular narrow QRS complexes, and retrograde or dissociated atrial activation, frequently associated with digitalis toxicity, myocardial ischemia, or pediatric cardiac surgery.

**Alternate Names:**

Nonparoxysmal Junctional Tachycardia; Accelerated Junctional Rhythm

**Jungian Analysis**

Jungian analysis is a psychotherapy approach based on Carl Jung's theories, using exploration of unconscious processes, symbols, dreams, and personality development.

**Juvenile Absence Epilepsy**

An idiopathic generalized epilepsy syndrome with onset in adolescence (typically between 9 and 13 years of age), characterized by absence seizures occurring less frequently than in childhood absence epilepsy, accompanied on EEG by generalized 3 to 4 Hz spike-and-wave discharges, frequently accompanied by generalized tonic-clonic seizures.

**Alternate Names:**

JAE; Adolescent Absence Epilepsy

**Juvenile Angiofibroma**

A noncancerous but highly blood-vessel-rich tumor that usually develops near the back of the nasal cavity in adolescent boys. It may cause worsening nasal blockage, repeated nosebleeds, or facial swelling.

**Juvenile Arthritis**

A general term for inflammatory joint disease beginning in childhood. It may cause persistent swelling, pain, stiffness, reduced movement, fever, rash, or eye inflammation; the subtype determines monitoring and treatment.

**Alternate Names:**

Juvenile Idiopathic Arthritis; JIA

**Juvenile Astrocytoma**

A tumor arising from support cells in a child's or adolescent's brain or spinal cord. Many grow slowly, but symptoms depend on location and may include headache, seizures, vision changes, weakness, balance problems, or hormone disturbances.

**Juvenile Dermatomyositis**

An autoimmune illness in children that weakens muscles, especially around the hips and shoulders, and causes a purple rash around the eyes or raised patches over the knuckles. It may also affect swallowing, breathing, or other organs.

**Juvenile Granulosa Cell Tumor**

A rare, malignant or low-grade sex cord-stromal ovarian neoplasm occurring predominantly in children and young women under age 30, characterized by solid and cystic morphology, luteinized cells, and hyperestrogenism leading to isosexual precocious



---

puberty in young girls.

### Juvenile Hypermobile Spectrum Disorder

A hypermobility spectrum disorder beginning in childhood or adolescence, with excessive joint movement plus pain, instability, soft-tissue injury, fatigue, or related symptoms.

### Juvenile Hypermobility Syndrome

A pattern of unusually flexible joints in a child or adolescent that causes pain, joint instability, tiredness, or repeated sprains. Assessment checks for other connective-tissue disorders; treatment may include strengthening, activity guidance, and joint protection.

### Juvenile Idiopathic Arthritis

A group of inflammatory diseases in which joint inflammation begins before age 16 and persists for at least six weeks after other causes are excluded. Depending on the subtype, children may have joint swelling, stiffness, or pain, as well as fever, rash, or eye inflammation.

### Juvenile Laryngeal Papillomatosis

A benign but aggressive, recurrent viral neoplastic condition of the respiratory tract caused by perinatally acquired human papillomavirus types 6 or 11, characterized by multiple exophytic squamous papillomas on the vocal cords and larynx causing progressive hoarseness, stridor, and airway compromise.

#### Alternate Names:

Recurrent Respiratory Papillomatosis; RRP

### Juvenile Myoclonic Epilepsy

A generalized epilepsy syndrome that typically begins in adolescence and causes myoclonic jerks, often after awakening, with possible generalized tonic-clonic or absence seizures.

### Juvenile Nasopharyngeal Angiofibroma

A benign, highly vascular, unencapsulated mesenchymal tumor occurring almost exclusively in adolescent males, originating in the posterolateral wall of the nasal cavity near the sphenopalatine foramen,

presenting clinically with recurrent painless epistaxis, nasal obstruction, and progressive facial swelling.

#### Alternate Names:

JNA; Nasopharyngeal Fibroangioma

### Juvenile Osteoporosis

Low bone strength during childhood or adolescence that increases the risk of fractures. It may occur on its own or result from chronic illness, poor nutrition, hormone disorders, limited mobility, or medicines.

### Juvenile Polyposis Syndrome

An inherited condition causing many noncancerous growths in the stomach or intestines. The growths can bleed, cause anemia, and increase the risk of bowel and stomach cancer.

### Juvenile Schizophrenia

Alternate name; see *Childhood Schizophrenia*.

### Juvenile Xanthogranuloma

A rare, usually harmless growth of immune cells in the skin of an infant or young child, appearing as a firm red, yellow, or brown bump. It often fades without treatment, but rarely affects the eye or internal organs.

### Jakob-Creutzfeldt Disease

Alternate name; see *Creutzfeldt-Jakob Disease*.

### Jakob's Disease

Alternate name; see *Creutzfeldt-Jakob Disease*.

### Jansen Disease

Alternate name for PTH1R-related Jansen metaphyseal chondrodysplasia, a rare inherited skeletal disorder with short-limb stature, abnormal metaphyses, and parathyroid-hormone-independent hypercalcemia; see *Jansen Metaphyseal Chondrodysplasia*.

### Jansen Metaphyseal Chondrodysplasia

A rare autosomal dominant skeletal disorder caused by activating variants in *PTH1R*. It causes short-limb short stature, metaphyseal bone abnormalities,

---

joint swelling, and often hypercalcemia, hypercalciuria, or nephrocalcinosis.

### Japanese B Encephalitis

Alternate name; see *Japanese Encephalitis*.

### Jaundice in Newborns (Hyperbilirubinemia)

Alternate name for neonatal jaundice caused by increased bilirubin; very early, rapidly worsening, or severe jaundice requires prompt assessment to prevent bilirubin-related brain injury.

### Jonston's Alopecia

Historical or misspelled index form for alopecia, meaning hair loss; see *Alopecia*.

### JRA Juvenile Rheumatoid Arthritis

Older name for juvenile idiopathic arthritis, an inflammatory arthritis beginning in childhood; see *Juvenile Idiopathic Arthritis*.

### Jumping Frenchmen of Maine

A rare historical startle disorder characterized by an exaggerated response to sudden stimuli, sometimes with involuntary jumping or other abrupt movements.

### Juvenile Diabetes

Older name for type 1 diabetes, which usually requires lifelong insulin replacement; see *Type 1 Diabetes*.

### Juvenile Retinoschisis

An inherited retinal disorder, usually X-linked, in which layers of the retina split and impair central or peripheral vision; see *Retinoschisis*.

### Juvenile Rheumatoid Arthritis

Older name for juvenile idiopathic arthritis, a group of inflammatory joint diseases beginning before age 16; see *Juvenile Idiopathic Arthritis*.

### Juxta-Articular

Located next to a joint. The term may describe

bone, soft tissue, swelling, a cyst, or a deposit beside a joint rather than within the joint itself.

### Juxtaglomerular Apparatus

A group of specialized kidney cells that senses blood flow and salt levels and releases renin. It helps regulate blood pressure, blood volume, and the body's salt balance.

### Juxtaglomerular Cells

Specialized, modified vascular smooth muscle cells situated predominantly in the media of the afferent arteriole as it enters the renal glomerulus, which synthesize, store in granules, and secrete the enzyme renin in response to decreased renal perfusion pressure, sympathetic stimulation, or reduced sodium delivery to the macula densa.

#### Alternate Names:

JG Cells; Granular Cells

### Juxtaglomerular Cell Tumor

A rare, benign renal neoplasm arising from the specialized renin-secreting myoepithelial juxtaglomerular cells of the afferent arteriole, characterized by marked hyperreninemia, secondary hyperaldosteronism, severe hypertension, and hypokalemia.

#### Alternate Names:

Reninoma; Juxtaglomerular Renin-Secreting Tumor

### Juxtavesical

Next to the urinary bladder. The term often describes the lower part of a ureter as it approaches the bladder, where a passing stone may become lodged and cause side pain, urinary frequency, or painful urination.

### J Wave

An abnormal, positive deflection or dome-shaped hump occurring at the junction between the QRS complex and the ST segment on an electrocardiogram, classically seen in systemic hypothermia (typically when core body temperature drops below 32 degrees Celsius) or severe hypercalcemia.

#### Alternate Names:

Osborn Wave; Camel-Hump Sign

# K

## Kabuki Syndrome

A rare genetic disorder that may cause distinctive facial features, short stature, developmental or learning difficulties, weak muscle tone, skeletal changes, and heart defects. Hearing loss, seizures, immune problems, and kidney abnormalities may also occur. Most cases result from a change in the *KMT2D* or *KDM6A* gene.

### Alternate Names:

Niikawa-Kuroki Syndrome

## Kahler Disease

Another name for multiple myeloma, a cancer of plasma cells in the bone marrow. It can produce an abnormal protein and cause weak or damaged bones, high calcium, kidney problems, and anemia.

### Alternate Names:

Multiple Myeloma; Plasma Cell Myeloma

## Kahler's Disease

Alternate name; see *Multiple Myeloma*.

## Kala-Azar

A serious parasitic infection caused by *Leishmania* species and transmitted by sandflies. It may cause prolonged fever, weight loss, enlargement of the spleen and liver, and suppression of blood-cell production.

### Alternate Names:

Black Sickness; Dumdum Fever

## Kallikrein-Kinin System

A group of blood and tissue proteins that makes chemical messengers such as bradykinin. These messengers widen small blood vessels, make them more

leaky, and contribute to pain and inflammation.

### Alternate Names:

Kininergetic System; Bradykinin Cascade

## Kallmann Syndrome

An inherited condition that causes delayed or absent puberty and a reduced or absent sense of smell. It occurs when the brain does not send the hormone signals needed to start puberty.

## Kallmann Syndrome and IHH

Conditions in which deficient hypothalamic or pituitary gonadotropin signaling causes low sex hormones and absent or delayed puberty. Kallmann syndrome also includes impaired smell and may be genetic; fertility and sex-hormone replacement require endocrine care.

## Kanamycin

An antibiotic used for some serious bacterial infections. It can permanently damage hearing and harm the kidneys, especially with prolonged or high-dose use.

## Kangaroo Care

Holding a newborn directly against a caregiver's bare chest while monitoring the infant's condition. It supports thermal regulation, physiologic stability, bonding, and feeding, and is particularly useful for many preterm or low-birth-weight infants.

### Alternate Names:

Kangaroo Mother Care; KMC

## Kaolin

A soft clay mineral used in some medicines, labora-

---

tory products, and industrial materials. Breathing large amounts of kaolin dust over time can damage the lungs.

### **Kaposiform Hemangioendothelioma**

A rare blood-vessel tumor that usually affects infants and young children. It can grow into nearby tissue and may cause a serious loss of platelets and problems with blood clotting.

#### **Alternate Names:**

KHE

### **Kaposi Sarcoma**

A vascular neoplasm associated with human herpesvirus 8 (HHV-8). It may cause nonblanching red, purple, brown, or black macules, plaques, or nodules on the skin or oral mucosa and can involve lymph nodes or internal organs; disease is more likely or aggressive with immunosuppression, including HIV or transplantation.

### **Kaposi's Sarcoma**

A vascular tumor associated with human herpesvirus 8, also called Kaposi sarcoma-associated herpesvirus. It may cause purple, red, or brown lesions of the skin or oral mucosa and can involve lymph nodes or internal organs; it is more likely or aggressive when immunity is weakened, such as with HIV infection or after organ transplantation.

### **Karnofsky Performance Status**

A scale used to describe how well a person can perform daily activities and care for themselves. It helps clinicians assess overall function and ability to tolerate treatment, especially in cancer care.

### **Kartagener Syndrome**

An inherited condition in which tiny hair-like structures that clear mucus do not work properly. It causes repeated sinus and lung infections and may place the internal organs in a reversed position.

### **Kartagener Triad**

The combination of reversed internal-organ positions, long-term sinus inflammation, and widened damaged airways. It results from an inherited prob-

lem with microscopic moving hairs that clear mucus.

### **Karyomegaly**

Abnormal enlargement of a cell's nucleus, the part that contains most of its genetic material. It may occur with injury, infection, toxic exposure, tissue repair, or cancer and is interpreted with other findings.

### **Karyotype**

An organized picture of a person's chromosomes, arranged in pairs to show their number and large-scale structure. It can detect some missing, extra, or rearranged chromosomes but not every genetic condition.

### **Karyotype Result**

A laboratory report describing the number and visible structure of a person's chromosomes. It may identify some chromosome differences, such as an extra chromosome, but a normal result does not exclude all genetic disorders.

### **Karyotyping**

A laboratory test that examines chromosomes in dividing cells to detect some missing, extra, or rearranged chromosomes. It may be used in pregnancy, infertility, repeated miscarriage, developmental conditions, or blood cancers.

### **Kasabach-Merritt Phenomenon**

A serious blood-clotting problem linked to certain fast-growing blood-vessel tumors in babies. Platelets become trapped in the tumor, causing a very low platelet count, anemia from broken red blood cells, and low levels of clotting proteins.

#### **Alternate Names:**

KMP; Kasabach-Merritt Syndrome

### **Kasabach-Merritt Syndrome**

Another name for Kasabach-Merritt phenomenon: a serious clotting problem in babies with certain fast-growing blood-vessel tumors. It causes very low platelets, anemia from broken red blood cells, and low levels of clotting proteins.

---

**Alternate Names:**

Kasabach-Merritt Phenomenon; Vascular Tumor Consumptive Coagulopathy

**Kasai Procedure**

An operation for babies with biliary atresia, in which the blocked bile ducts outside the liver are removed and a loop of intestine is connected to the liver so bile can drain. Earlier surgery gives the best chance of restoring bile flow, but many children eventually need a liver transplant.

**Alternate Names:**

Hepatopertoenterostomy; Kasai Operation; Por-toenterostomy

**Kashin-Beck Disease**

A long-term disease of the bones and joints that can cause pain, stiffness, deformity, and poor growth. It is linked to low selenium and environmental factors in some regions.

**Katayama Fever**

An acute systemic reaction to schistosome infection, usually occurring weeks after exposure and causing fever, rash, cough, abdominal symptoms, and eosinophilia.

**Alternate Names:**

Katayama Syndrome; Acute Bilharziasis

**Katz Index**

A scale that assesses independence in six basic activities of daily living: bathing, dressing, toileting, transferring, continence, and feeding.

**Kawasaki Disease**

An acute inflammatory disease of childhood that can affect the coronary arteries. It commonly causes persistent fever, rash, conjunctival redness, mouth or extremity changes, and swollen lymph nodes; diagnosis is clinical and incomplete forms occur.

**Alternate Names:**

Mucocutaneous Lymph Node Syndrome; KD

**Kawasaki Disease Diagnostic Criteria**

The usual clinical pattern used to identify Kawasaki

disease: fever for at least five days plus at least four of five findings—red eyes without pus, a varied skin rash, swollen neck lymph nodes, changes in the lips or mouth, and redness, swelling, or peeling of the hands or feet. Some children have an incomplete form.

**Kayser-Fleischer Rings**

Golden-brown or greenish rings around the colored part of the eye caused by copper buildup. They strongly suggest Wilson disease when other findings fit and are seen with a special eye examination.

**K-Complex**

A large brain-wave pattern seen during stage 2 sleep. It may occur on its own or after a sound or other sensory stimulus and is recorded during an electroencephalogram (EEG).

**Kearns-Sayre Syndrome**

A rare inherited disorder affecting the mitochondria, the parts of cells that make energy. It usually begins in childhood or young adulthood and can cause drooping or limited eye movement, vision problems, heart-rhythm problems, and muscle weakness.

**Keel Breast**

A chest-wall shape in which the breastbone pushes outward, giving the chest a keel-like appearance. It usually reflects pectus carinatum, which develops during growth and may affect appearance, breathing, or exercise tolerance.

**Kegel**

A pelvic-floor muscle contraction exercise used to improve pelvic-floor strength and control of selected types of urinary or fecal incontinence.

**Kegel Exercise**

Repeated tightening and relaxing of pelvic-floor muscles to improve strength and control, helping reduce urine leakage and support pelvic organs.

**Kehr Sign**

Pain felt at the tip of the left shoulder when blood or

---

irritation affects the underside of the diaphragm, the main breathing muscle. It can occur with irritation from a spleen injury but is not specific to it.

### **Keloid**

A raised, firm scar that grows beyond the edges of the original wound because the body makes too much scar tissue. It may itch or hurt and can develop after surgery, injury, burns, or piercings.

#### **Alternate Names:**

Keloid Scar; Cheloid

### **Keloid and Hypertrophic Scar**

Raised scars caused by excessive collagen deposition during wound healing. Hypertrophic scars remain within the original wound, whereas keloids extend beyond it; symptoms may include itch, pain, tightness, and cosmetic concern.

### **Keloid Scar**

A raised, firm scar that grows beyond the original skin injury because of excessive scar formation. It may itch, hurt, or recur after removal and is more common in some families and skin types.

### **Keratan Sulfate**

A complex sugar found mainly in the clear front of the eye, cartilage, and brain. In some inherited disorders it cannot be broken down properly and builds up, damaging the eyes, bones, or other tissues.

### **Keratan Sulphate**

A sulfated glycosaminoglycan found especially in the cornea and cartilage; abnormal accumulation occurs in some mucopolysaccharidoses.

### **Keratotomy**

Removal or reshaping of part of the cornea, the clear front surface of the eye. It may be done to correct vision or remove abnormal tissue, and it can affect eyesight.

### **Keratic Precipitates**

Small groups of inflammatory cells that collect on the inner surface of the cornea, usually during in-

flammation inside the front of the eye. Their size and appearance can help describe the type of inflammation.

### **Keratin**

A strong structural protein forming much of the skin's outer layer, hair, nails, and other protective tissues. It helps resist water loss, friction, chemicals, and physical injury.

### **Keratinocyte**

The main type of cell in the outer layer of skin. It produces keratin and helps form the skin barrier that limits water loss and protects against injury and infection.

### **Keratinocytes**

The predominant cells of the epidermis that produce keratin and form the skin's protective barrier.

### **Keratin Pearl**

A round, layered collection of keratin seen under a microscope in some well-differentiated squamous cell skin cancers.

#### **Alternate Names:**

Squamous Pearl; Horn Pearl

### **Keratitis**

Inflammation or infection of the clear front surface of the eye. It can cause eye pain, redness, tearing, sensitivity to light, blurred vision, or a cloudy spot.

#### **Alternate Names:**

Corneal Ulceration; Corneal Inflammation

### **Keratoacanthoma**

A fast-growing, dome-shaped skin growth with a central plug, usually on sun-exposed skin. It may look like a type of skin cancer, so a clinician often removes or samples it. Treatment depends on its appearance, size, site, and the test results.

#### **Alternate Names:**

Well-Differentiated Squamous Cell Carcinoma, Keratoacanthoma Type; KA; Crateriform Epithelioma



---

## **Keratocele**

Severe thinning or ulceration of the cornea that allows its inner layers to bulge outward. It can lead to a hole in the cornea and permanent loss of vision.

## **Keratoconjunctivitis**

Inflammation of the cornea and the conjunctiva, the clear surface covering the white of the eye. Infection, allergy, dryness, immune disease, or chemical injury may cause redness, pain, discharge, light sensitivity, and blurred vision.

## **Keratoconjunctivitis Sicca**

Long-lasting dryness and inflammation of the eye's front surface, commonly called dry-eye disease. It can cause burning, grittiness, redness, fluctuating vision, or light sensitivity because too few tears are made or tears evaporate too quickly.

### **Alternate Names:**

Dry Eye Syndrome; Ocular Surface Disease

## **Keratoconus**

A condition in which the clear front surface of the eye becomes thin and bulges into a cone shape. It can cause blurred or distorted vision, glare, and sensitivity to light. Glasses, special contact lenses, or treatment to strengthen the cornea may help.

## **Keratocyte Count**

An estimate of the number of living support cells in the middle layer of the cornea. It can help assess corneal health, injury, disease, or changes after eye surgery, although results are interpreted with the complete eye examination.

## **Keratoderma**

Excessive thickening of the skin on the palms or soles. It may be inherited or develop because of skin disease, infection, medicines, or another illness and can cause painful cracks or difficulty walking.

## **Keratoderma Blennorrhagica**

Thickened, scaly, sometimes pustular skin lesions, usually on the soles or palms, associated with reactive arthritis. Treatment focuses on the underlying

infection or inflammation and the skin symptoms.

## **Keratoderma Blennorrhagicum**

Thickened, scaly, or pus-filled skin lesions on the palms and soles associated with reactive arthritis. The lesions may spread to other areas and often appear after an infection affecting the urinary or digestive tract.

## **Keratoderma Blenorrhagicum**

Alternate spelling of keratoderma blennorrhagicum, a cutaneous manifestation of reactive arthritis with waxy, crusty, vesiculopustular lesions on the palms and soles. It is a reactive phenomenon typically occurring in the context of genitourinary or gastrointestinal infection.

## **Keratoderma Blennorrhagicum**

Thickened, scaly, or wart-like skin lesions, often on the palms and soles, associated with reactive arthritis and sometimes infection-triggered inflammation.

## **Keratolysis**

Breakdown or loss of keratinized tissue, especially the superficial layer of the skin; it may occur in some infections and inflammatory skin disorders.

## **Keratolytics**

Substances, such as salicylic acid or urea, that soften and loosen excess hard skin. They may be used for thickened skin, corns, calluses, or some warts.

## **Keratoma**

A localized thickening or tumor-like growth of keratinized skin, such as a callus or corn; the exact type and need for evaluation depend on its appearance and behavior.

## **Keratomalacia**

Softening and ulceration of the cornea caused most often by severe vitamin A deficiency, potentially leading to blindness.

## **Keratometer**

An eye instrument that measures the curvature of

---

the cornea, the clear front surface of the eye. The measurement helps assess astigmatism and fit rigid contact lenses.

### **Keratometry**

Measurement of the curve of the cornea, the clear front surface of the eye. It helps assess focusing problems and choose contact lenses or an artificial lens for cataract surgery.

### **Keratoplasty**

Surgery to replace or reshape part or all of the cornea, the clear front surface of the eye. It can restore vision after corneal damage, but healing problems, infection, and rejection of donor tissue are possible.

### **Keratoprosthesis**

An artificial cornea implanted to restore sight when a donor-corneal transplant is unlikely to work, such as after severe chemical injury or extensive surface damage. It carries risks of infection, glaucoma, and device problems.

### **Keratosis**

A thickened, rough, or otherwise abnormal area of skin caused by excess keratin. Some forms are harmless, while others result from infection, inflammation, sun damage, or a change that can precede skin cancer.

### **Keratosis Follicularis**

Alternate name for Darier disease; see *Darier's Disease*.

### **Keratosis Follicularis (Darier Disease)**

Alternate name; see *Darier's Disease*.

### **Keratosis Obturans**

A buildup of shed skin cells that blocks the ear canal. It can cause ear pain, hearing loss, discharge, and widening or damage of the canal.

### **Keratosis Pilaris**

A common, harmless skin condition that causes

small, rough bumps around hair follicles, usually on the upper arms, thighs, buttocks, or face.

#### **Alternate Names:**

KP

### **Kerion**

An inflamed, boggy, pus-forming mass caused by a fungal infection of hair-bearing skin, usually a severe form of tinea capitis.

### **Kerley A Lines**

Long, fine lines seen on a chest X-ray, usually spreading outward from the central lung areas. They can reflect fluid buildup in the lung tissue.

#### **Alternate Names:**

Kerley's A Lines; Upper Zone Interstitial Edema Lines

### **Kerley B Lines**

Short, thin lines seen near the outer edges and bases of the lungs on a chest X-ray. They commonly reflect fluid buildup in the lung tissue, often related to heart failure.

#### **Alternate Names:**

Kerley's B Lines; Septal Lines

### **Kernicterus**

Permanent brain damage caused by very high levels of bilirubin, a yellow substance that builds up during severe newborn jaundice. It can cause movement problems, hearing loss, problems looking upward, and damage to tooth enamel.

#### **Alternate Names:**

Bilirubin Encephalopathy; Nuclear Jaundice; Chronic Bilirubin Toxicity

### **Kernig Sign**

Pain or resistance when the knee is straightened while the hip is bent. It may suggest irritation of the coverings of the brain and spinal cord, such as with meningitis, but it is not diagnostic by itself.

### **Kernig's Sign**

Alternate spelling; see *Kernig Sign*.

---

## Kernohan Notch Phenomenon

A paradoxical false-localizing neurological sign in which acute uncal transtentorial herniation displaces the midbrain against the contralateral rigid tentorial notch (Kernohan notch), compressing the contralateral cerebral peduncle (corticospinal tract) and producing an unexpected ipsilateral hemiparesis.

### Alternate Names:

Kernohan-Woltman Notch Phenomenon; Kernohan Sign

## Kerosene Poisoning

Harm after swallowing or inhaling kerosene. The greatest danger is entry into the lungs, which can cause chemical pneumonia; coughing, choking, breathing difficulty, drowsiness, or vomiting may occur.

## Keshan Disease

A heart-muscle disease linked to severe selenium deficiency, occurring mainly in regions where soil and food contain little selenium. It can weaken the heart and cause heart failure.

## Ketamine

Ketamine is a dissociative anaesthetic and NMDA-receptor antagonist used for anaesthesia, procedural sedation, and selected treatment-resistant depression; emergence reactions, increased blood pressure, nausea, misuse, and respiratory complications are possible.

## Ketamine Hydrochloride

Ketamine hydrochloride is a dissociative anesthetic used for anesthesia, procedural sedation, and selected pain or depression treatments; it can cause increased blood pressure, hallucinations, and impaired coordination.

## Ketamine Toxicity

Acute adverse effects or overdose of the NMDA receptor antagonist dissociative anesthetic ketamine, characterized by perceptual distortions, hallucinations, severe emergence delirium, profound nystagmus, hypertension, tachycardia, and in chronic abusers, severe ulcerative cystitis ("ketamine bladder").

## Keto Acid

A chemical made when the body breaks down certain amino acids or other nutrients. Keto acids can be used to make energy or new substances, but abnormal amounts may occur in metabolic disease.

## Ketoacidosis

A dangerous buildup of acidic ketones in the blood, usually because there is too little insulin or prolonged starvation. It can cause vomiting, abdominal pain, deep breathing, confusion, coma, and death.

## Ketoconazole

Ketoconazole is an imidazole antifungal that inhibits fungal ergosterol synthesis; topical forms treat superficial mycoses, while oral use is restricted because of potentially severe liver injury, adrenal suppression, and drug interactions.

## Ketogenesis

The production of ketone bodies from fatty acids, mainly in the liver during fasting, prolonged exercise, or inadequate insulin activity.

## Ketogenic Diet

An eating plan very low in carbohydrates and higher in fat that makes the body use ketones for energy. It can help some children with difficult-to-control epilepsy, but may cause nutrient deficiencies, constipation, kidney stones, or abnormal cholesterol without supervision.

## Ketone

A type of chemical compound. In medicine, the term often refers to ketone bodies made when the body uses fat for energy; small amounts can be normal, but high levels may cause ketoacidosis.

## Ketone Bodies

Water-soluble molecules produced mainly by the liver from fatty-acid-derived acetyl-CoA during fasting, starvation, prolonged exercise, or insulin deficiency. The principal ketone bodies are acetoacetate,

---

beta-hydroxybutyrate, and acetone.

## Ketones

Water-soluble substances produced mainly by the liver when the body breaks down fat for energy, such as during fasting, a low-carbohydrate state, or insufficient insulin. The main ketone bodies are beta-hydroxybutyrate, acetoacetate, and acetone. Small amounts can be normal, but high levels may cause ketoacidosis, especially in diabetes; ketones can be measured in blood or urine.

## Ketones Blood Test

A blood test measuring ketone bodies, such as beta-hydroxybutyrate, used to assess ketosis and help diagnose or monitor diabetic ketoacidosis and other causes of ketone production.

## Ketones Urine Test

A urine test detecting ketone bodies produced when the body metabolizes fat; it can support evaluation of diabetic ketoacidosis, starvation, vomiting, or other metabolic states but may lag behind blood ketone levels.

## Ketonuria

Ketonuria is the presence of ketone bodies in urine, occurring with fasting, vomiting, low-carbohydrate intake, pregnancy, or uncontrolled diabetes; marked ketonuria with hyperglycaemia may indicate diabetic ketoacidosis.

## Ketoprofen

A non-steroidal anti-inflammatory drug that inhibits cyclo-oxygenase and reduces pain, inflammation, and fever; gastrointestinal bleeding, kidney injury, cardiovascular events, and hypersensitivity are important risks.

## Ketoprofen Overdose

Harm caused by taking too much ketoprofen, a nonsteroidal anti-inflammatory drug. It may cause nausea, vomiting, stomach bleeding, drowsiness, kidney injury, seizures, or a dangerous buildup of acid in the blood.

## Ketorolac

A potent non-steroidal anti-inflammatory analgesic used for short-term management of moderate-to-severe acute pain; treatment duration is restricted because gastrointestinal bleeding, kidney injury, and impaired healing can occur.

## Ketorolac Tromethamine

The tromethamine salt of ketorolac, a potent non-steroidal anti-inflammatory analgesic used short term for moderate-to-severe acute pain; it is contraindicated in several bleeding, renal, pregnancy, and perioperative situations.

## Ketosis

A state in which the body makes and uses increased amounts of ketones after breaking down fat for energy. It can occur during fasting or a low-carbohydrate diet, but high ketone levels with illness or high blood sugar may signal diabetic ketoacidosis.

## Ketosis-Prone Type 2 Diabetes

A form of diabetes that may present with diabetic ketoacidosis despite features of type 2 diabetes, such as obesity or preserved insulin production. Insulin dependence can change over time according to beta-cell reserve, autoimmunity, and recovery from the initial metabolic stress.

## Ketotic Hypoglycemia

Low blood sugar accompanied by ketones, usually after a young child has gone without food or has been ill and unable to eat. Drowsiness, weakness, vomiting, or seizures may occur. Repeated episodes can have other causes.

## Ketotifen

An antihistamine with mast-cell-stabilizing activity used in some countries for allergic disorders and, in ophthalmic form, allergic conjunctivitis; drowsiness, increased appetite, and irritation may occur.

## Kidney

One of two bean-shaped organs that filter waste from the blood into urine. The kidneys also help

---

control body water, salts, blood pressure, and the balance of acids in the blood.

### **Kidney Abscess**

Alternate name for renal abscess, a localized collection of pus in or around the kidney.

#### **Alternate Names:**

Renal Abscess; Perinephric Abscess

### **Kidney Biopsy**

A procedure in which a small sample of kidney tissue is removed, usually with a needle guided by ultrasound, to identify kidney disease. Bleeding is the main risk.

### **Kidney Calculi**

Alternate term; see *Kidney stones*.

### **Kidney Cancer**

A malignant tumor arising in the kidney. Renal cell carcinoma is the most common type in adults, but other types occur, especially in children.

### **Kidney Cancer In Children**

Cancer arising in a child's kidney. Wilms tumor is the most common type, but other kidney tumors can occur; symptoms may include an abdominal mass, pain, blood in the urine, fever, or high blood pressure.

### **Kidney Cysts**

Fluid-filled sacs in or on the kidneys. Simple cysts are common and usually harmless, while complex or multiple cysts may be associated with infection, bleeding, inherited disease, or cancer risk and may impair kidney function.

#### **Alternate Names:**

Renal Cysts; Renal Cystic Disease

### **Kidney Disease**

Any condition that damages the kidneys or reduces their ability to filter blood, balance fluids and minerals, control blood pressure, or make hormones. Causes include diabetes, high blood pressure, infection, inflammation, inherited disorders,

and medicines.

### **Kidney Disease and Pregnancy**

Kidney disease can affect fertility, blood pressure, fluid balance, medication choices, and pregnancy outcomes, while pregnancy can worsen some kidney disorders. Preconception counseling and coordinated obstetric-nephrology care help monitor kidney function, proteinuria, blood pressure, fetal growth, and medicine safety.

### **Kidney Disease-Related Anemia**

Anemia caused or worsened by kidney disease, mainly because damaged kidneys produce less erythropoietin, the hormone that stimulates red-cell production. Iron deficiency, inflammation, and blood loss may also contribute.

### **Kidney Diseases**

Disorders affecting the kidneys or urinary collecting system that can impair filtration, urine formation, fluid balance, blood pressure, or electrolyte regulation.

### **Kidney Failure**

A condition in which the kidneys can no longer remove enough waste or control fluid and body salts. It may develop suddenly or gradually and can require medicines, dialysis, or a transplant.

#### **Alternate Names:**

End-Stage Renal Disease; ESRD; Renal Failure

### **Kidney Function Tests**

Blood and urine tests used to assess how well the kidneys remove waste and control fluid and salts. Common measures include creatinine, estimated filtration rate, urine protein, and urinalysis.

### **Kidney Infection**

An infection of one or both kidneys, usually caused by bacteria that travel upward from the bladder. It can cause fever, chills, side or back pain, nausea, vomiting, and painful urination and may lead to sepsis.

---

## Kidney Neoplasm

An abnormal growth in the kidney that may be noncancerous or cancerous. It may cause blood in the urine, pain, a lump, weight loss, or no symptoms; imaging and sometimes biopsy identify its type and extent.

## Kidney Pain

Pain felt in the side or back below the ribs that may be caused by kidney stones, infection, obstruction, inflammation, or referred pain.

### Alternate Names:

Flank Pain; Renal Colic; Nephralgia

## Kidney Position

A side-lying surgical position in which the waist is gently flexed and the side of the body is raised. It gives surgeons better access to the kidney and nearby tissues while staff protect pressure points and breathing.

## Kidney Removal

Surgery to remove part or all of a kidney, usually for cancer, severe damage, infection, or a nonfunctioning kidney. The remaining kidney may provide adequate function, but bleeding, infection, injury to nearby organs, or reduced kidney function can occur.

## Kidney Replacement Therapy

Treatment that replaces some functions of severely damaged kidneys. It includes hemodialysis, peritoneal dialysis, continuous kidney replacement therapy, and kidney transplantation.

## Kidney Stone

Singular form; see *Kidney stones*.

## Kidney Stones

Hard mineral deposits formed from substances in urine. They may obstruct the urinary tract and cause colicky flank or abdominal pain, blood in the urine, vomiting, or infection.

## Kidney Stones In Children

Mineral deposits that form in a child's kidney or urinary tract. They may cause side or abdominal pain, blood in the urine, nausea, vomiting, urinary symptoms, or infection, but some cause no symptoms.

### Alternate Names:

Pediatric Nephrolithiasis; Pediatric Urolithiasis; Pediatric Renal Calculi

## Kidney Transplant

Surgical placement of a healthy donor kidney into a person with kidney failure to restore renal function. Lifelong immunosuppressive treatment is generally required to reduce rejection risk.

## Kidney Transplantation

Placement of a donor kidney to replace the filtering work of kidneys that have permanently failed. A living-donor kidney may work sooner, but all recipients need lifelong medicines and monitoring for rejection, infection, blood-vessel problems, and return of kidney disease.

## Kidney Transplant Rejection

An immune attack that damages a transplanted kidney. It may occur suddenly or gradually and can cause reduced urine, swelling, high blood pressure, or worsening blood tests; treatment depends on the type and severity.

## Kidney Trauma

Alternate name for renal trauma, an injury to the kidney caused by blunt or penetrating force.

## Kienbock Disease

Damage and gradual collapse of the lunate, a small bone in the wrist, because its blood supply is reduced. It may cause wrist pain, swelling, stiffness, and weaker grip. Treatment depends on bone damage, wrist alignment, symptoms, and whether arthritis has developed.

## Kieselguhr

Diatomaceous earth containing substantial silica; in-



---

halation of its dust can produce occupational pneumoconiosis and other lung injury.

### **Kiesselbach Plexus**

A network of small arteries on the front part of the nasal septum and a common source of nosebleeds.

### **Kiesselbach Triangle**

An anterior vascular network on the anteroinferior nasal septum (Little's area) formed by anastomoses of the anterior ethmoidal, sphenopalatine, greater palatine, and superior labial arteries, serving as the anatomical site for over 90

#### **Alternate Names:**

Little Area; Kiesselbach's Plexus

### **Kikuchi Disease**

A usually short-term inflammation of the lymph nodes, most often in the neck, with no known cause. It mainly affects young adults and may cause tender swollen nodes, fever, night sweats, and a low white-blood-cell count.

#### **Alternate Names:**

Kikuchi-Fujimoto Disease; Histiocytic Necrotizing Lymphadenitis

### **Kikuchi-Fujimoto Disease**

Another name for Kikuchi disease, a usually short-term inflammation of the lymph nodes, most often in the neck. It may cause painful swollen nodes, fever, night sweats, and a low white-blood-cell count.

#### **Alternate Names:**

Kikuchi Disease; Histiocytic Necrotizing Lymphadenitis

### **Killer Cold Virus**

Alternate name; see *Adenovirus Infection*.

### **Killian's Triangle**

An area of potential weakness in the pharyngeal wall between the thyropharyngeus and cricopharyngeus muscles, predisposing to Zenker diverticulum formation.

### **Killian Triangle**

A triangular anatomical area of muscular weakness in the posterior hypopharyngeal wall situated between the oblique fibers of the thyropharyngeus muscle and the transverse fibers of the cricopharyngeus muscle; the site of mucosal herniation forming a Zenker diverticulum.

#### **Alternate Names:**

Killian Dehiscence

### **Killip Classification**

A four-level system for describing heart failure during a heart attack. The levels range from no signs of heart failure to lung fluid buildup and, at the most severe level, shock caused by the heart's inability to pump enough blood.

### **Kilocalorie**

A unit of energy equal to 1,000 small calories; the dietary Calorie, written with a capital C, is one kilocalorie.

### **Kilojoule**

The SI unit of energy, equal to 1,000 joules; approximately 4.184 kilojoules equal one kilocalorie.

### **Kimmelstiel-Wilson Lesion**

A round area of thickened tissue in a kidney filter that can occur with long-standing diabetes. It is a sign of diabetic kidney damage and may occur with protein in the urine or reduced kidney function.

#### **Alternate Names:**

Nodular Diabetic Glomerulosclerosis; Kimmelstiel-Wilson Nodules

### **Kimmelstiel-Wilson Nodules**

Round areas of thickened tissue seen in the kidney filters of some people with long-standing diabetes. They reflect diabetic kidney damage and may occur with protein in the urine and declining kidney function.

### **Kimmelstiel-Wilson Syndrome**

A pattern of kidney scarring caused by long-standing diabetes. It can make protein leak into the urine

---

and gradually reduce kidney function, sometimes leading to kidney failure.

### **Kimura Disease**

A rare long-term immune disorder causing painless lumps under the skin, usually in the head or neck, enlarged nearby lymph nodes, and high levels of eosinophils or IgE. It can sometimes affect the kidneys.

### **Kinase**

An enzyme that attaches a small chemical group called a phosphate to another molecule. This acts like a switch controlling cell growth, energy use, movement, communication, and many other processes.

### **Kinase Inhibitor**

A targeted therapeutic agent (typically a small molecule or monoclonal antibody) that selectively inhibits the enzymatic phosphorylation activity of specific protein kinases (such as tyrosine kinases, serine/threonine kinases, or cyclin-dependent kinases) involved in oncogenic cellular signaling cascades.

### **Kinesia Paradoxa**

A brief, unexpected ability to move normally or quickly in a person who usually has severe slowness or difficulty moving, as can occur in advanced Parkinson disease. It may happen during a strong emotional experience or in response to an urgent situation.

#### **Alternate Names:**

Paradoxical Kinesia; Paradoxical Movement

### **Kinesiology**

The study of human movement and the forces produced by muscles, joints, and other body structures. It is used in biomechanics, rehabilitation, exercise science, and the assessment of movement and physical function.

### **Kinesthesia**

The sense of how the body's parts are moving and where they are positioned, even when the eyes are closed. Information from muscles, joints, and skin

helps the brain control posture, balance, and coordinated movement.

### **Kinetics**

The study of how a process changes over time. In medicine, pharmacokinetics describes how the body absorbs, distributes, changes, and removes a medicine, helping determine its dose and timing.

### **Kinetochores**

A specialized, disc-shaped multiprotein complex assembled onto the centromeric DNA region of a eukaryotic chromosome that captures spindle microtubules and coordinates chromosome segregation during mitosis and meiosis.

### **Kingella**

Kingella is a genus of bacteria that can live in the mouth and can occasionally cause bloodstream, bone, joint, or heart-valve infections.

#### **Kingella Denitrificans**

Kingella denitrificans is a rare bacterium found mainly in the upper airway; it can occasionally cause invasive infection.

#### **Kingella Kingae**

Kingella kingae is a bacterium that can cause bone or joint infection, especially in young children, and can rarely cause endocarditis.

#### **Kingella Oralis**

Kingella oralis is a mouth-associated bacterium that rarely causes invasive infection, including bloodstream infection or endocarditis.

#### **Kingella Potus**

Kingella potus is a rare bacterium that has been associated with occasional human infections; clinical importance depends on the specimen and evidence of true infection.

### **King's College Criteria**

A set of findings used to estimate the risk of death in acute liver failure and the possible need for a liver

---

transplant. The criteria differ for acetaminophen poisoning and other causes.

### **Kinin**

A short peptide, such as bradykinin, that widens blood vessels, increases vascular permeability, and contributes to pain and inflammation.

### **Kinin-Kallikrein System**

An endogenous enzymatic cascade of blood proteins that produces bioactive peptide kinins (primarily bradykinin and kallidin) from high- and low-molecular-weight kininogens, mediating vasodilation, increased microvascular permeability, tissue inflammation, and pain sensation.

#### **Alternate Names:**

Kallikrein-Kinin Cascade

### **Kinins**

A group of vasoactive peptides, chiefly bradykinin and kallidin, generated from kininogens during inflammation; they promote vasodilation, increased vascular permeability, smooth-muscle effects, and pain.

### **Kinocilium**

A specialized hair-like structure on certain sensory cells, especially in the inner ear. It helps the cell detect the direction of movement and contributes to balance and hearing signals.

### **Kissing Disease**

Informal term; see *Infectious mononucleosis*.

### **Kite Angle**

An anatomical radiographic angle measured on an anteroposterior radiograph of the foot between the longitudinal axis of the talus and the longitudinal axis of the calcaneus. A normal angle ranges from 20 to 40 degrees; convergence or reduction of this angle below 15 degrees signifies hindfoot parallelism, a diagnostic hallmark of congenital talipes equinovarus (clubfoot).

#### **Alternate Names:**

Talocalcaneal Angle; AP Kite Angle

### **Klatskin Tumor**

A cancer arising where the right and left bile ducts join near the liver. It can block bile flow and cause jaundice, itching, pale stools, or weight loss.

### **Klebsiella**

A genus of Gram-negative bacteria that can cause pneumonia, urinary-tract infection, bloodstream infection, and other healthcare-associated infections.

### **Klebsiella Oxytoca**

*Klebsiella oxytoca* is a bacterium that can cause urinary, bloodstream, wound, and hospital-acquired infections and can sometimes cause antibiotic-associated hemorrhagic colitis.

### **Klebsiella Pneumonia**

Pneumonia caused by *Klebsiella* bacteria. It can cause fever, cough, chest pain, breathlessness, and sometimes thick or blood-stained sputum; severe infection may destroy lung tissue and is more likely in people with serious illness or weakened defenses.

### **Klebsiella Pneumoniae**

*Klebsiella pneumoniae* is a bacterium that can cause pneumonia, urinary infection, bloodstream infection, and liver abscess; some strains are highly antibiotic-resistant.

### **Klebsiella Rhinoscleromatis**

A gram-negative bacillus causing chronic granulomatous infection of the nasal passages and upper airway, leading to rhinoscleroma with progressive nasal obstruction and tissue destruction.

### **Klebsiella Singaporensis**

*Klebsiella singaporensis* is a rare environmental bacterium that has occasionally been recovered from human clinical specimens; its significance depends on the specimen and symptoms.

### **Klebsiella Variicola**

*Klebsiella variicola* is a bacterium closely related to *K. pneumoniae* that can cause human infections, including bloodstream and urinary infections.

---

### Kleihauer-Betke Test

A quantitative acid-elution blood test used to detect and calculate the volume of fetomaternal hemorrhage; exposure to an acid buffer elutes adult hemoglobin (HbA) from maternal erythrocytes while acid-resistant fetal hemoglobin (HbF) inside fetal red blood cells remains intact and stains dark pink, allowing calculation of necessary Rh immune globulin (RhoGAM) dosing.

#### Alternate Names:

Acid Elution Test; Fetomaternal Hemorrhage Assay

### Kleine-Levin Syndrome

A rare disorder causing repeated episodes of excessive sleep, often with confusion, reduced activity, and changes in behavior or appetite. Episodes last days to weeks and are separated by periods of more typical function.

### Kleptomania

A mental-health disorder involving recurrent inability to resist impulses to steal items that are not needed for personal use or monetary value. The acts are typically preceded by rising tension and followed by relief or gratification, and cause distress or impairment.

#### Alternate Names:

Compulsive Stealing Pathological Stealing

### Klinefelter's Syndrome

Alternate name; see *Klinefelter Syndrome*.

### Klinefelter Syndrome

A genetic condition in males caused by one or more extra X chromosomes, most often the 47,XXY pattern. It can cause small testes, low testosterone, infertility, tall stature, less facial and body hair, breast enlargement, weak bones, and learning or language difficulties. A chromosome test identifies the condition.

#### Alternate Names:

47,XXY Syndrome; XXY Syndrome; Klinefelter-Reifenstein-Albright Syndrome

### Klippel-Feil Syndrome

A condition present from birth in which two or more neck bones are joined together. It may cause a short neck, low hairline, limited neck movement, hearing loss, or spinal and kidney problems.

### Klippel-Trenaunay Syndrome

A rare congenital vascular disorder characterized by a classic triad of features: capillary malformations (port-wine stains), soft tissue and bone enlargement (hypertrophy, usually affecting one leg or arm), and slow-flow venous and lymphatic malformations (such as persistent embryonic veins, prominent varicose veins, and lymphedema). It is typically caused by non-inherited (somatic) mutations in the *PIK3CA* gene. Complications include limb length discrepancy, chronic pain, recurrent skin infections (cellulitis), and blood clots (deep vein thrombosis and pulmonary embolism). Management is multidisciplinary and includes compression garments, physiotherapy, vascular procedures, and orthopedic care.

#### Alternate Names:

KTS; Klippel-Trénaunay Syndrome; Angio-Osteohypertrophy Syndrome; KTW Syndrome; KTWS

### Klippel-Trenaunay-Weber Syndrome

A congenital vascular and limb-overgrowth disorder with capillary malformation, venous or lymphatic abnormalities, and enlargement of a limb or its digits. It can cause pain, bleeding, ulcers, clotting, or pulmonary embolism and requires vascular assessment.

### Klüver-Bucy Syndrome

A profound neurobehavioral syndrome resulting from bilateral damage or surgical ablation of the anterior temporal lobes, specifically including the amygdaloid nuclei and hippocampus. Clinical hallmarks include visual agnosia (psychic blindness), compulsive oral exploration of objects (hyperorality), hypermetamorphosis, marked passivity or docility, altered dietary preferences, and inappropriate hypersexuality.

#### Alternate Names:

---

Kluver-Bucy Syndrome; Bilateral Amygdala Ablation Syndrome

### **Klumpke Palsy**

An injury to the lower trunk of the brachial plexus involving spinal nerve roots C8 and T1, typically resulting from upward traction on an abducted arm during birth or trauma, causing paralysis of intrinsic hand muscles resulting in a 'claw hand' deformity and ipsilateral Horner syndrome (if sympathetic T1 root fibers are disrupted).

#### **Alternate Names:**

Klumpke-Dejerine Palsy; Lower Trunk Brachial Plexus Injury

### **Klumpke Paralysis**

A nerve injury affecting the lower part of the network that supplies the arm and hand. It can cause severe hand weakness, numbness along the inner forearm and hand, and occasionally drooping of the eyelid and a smaller pupil on one side.

### **Klumpke's Paralysis**

Lower brachial-plexus palsy involving mainly the C8 and T1 nerve roots, causing hand weakness and sensory loss; Horner syndrome may accompany severe cases.

### **Kluyvera**

A genus of gram-negative bacteria in the Enterobacteriales group; species can rarely cause opportunistic human infections.

### **Kluyvera Ascorbata**

A gram-negative Kluyvera species found in the environment and occasionally associated with urinary, bloodstream, or other opportunistic infections.

### **Kluyvera Cryocrescens**

A gram-negative environmental bacterium that rarely causes opportunistic human infection.

### **Kluyvera Georgiana**

A rarely encountered Gram-negative Kluyvera species that may appear in clinical samples. It can

occasionally cause opportunistic infection, especially in people with serious illness, weakened defenses, or invasive medical devices.

### **Kluyvera Intermedia**

A Gram-negative bacterium occasionally found in human samples and rarely causing opportunistic urinary, wound, bloodstream, or other infections.

### **Knee**

The synovial joint between the femur, tibia, and patella, stabilized by the cruciate and collateral ligaments and menisci; it permits flexion and extension and bears body weight.

### **Knee Arthroplasty**

Surgery that replaces damaged parts of the knee joint with artificial components. It is usually performed for severe pain and loss of function from arthritis or major joint damage.

### **Knee Arthroscopy**

Knee arthroscopy is a minimally invasive procedure in which a camera and small instruments are inserted into the knee to diagnose or treat selected joint problems.

### **Knee Bursitis**

Inflammation of a fluid-filled sac around the knee, causing local swelling, tenderness, and pain with kneeling or movement. Repeated pressure, injury, arthritis, gout, or infection can contribute; fever or redness may occur with infection.

#### **Alternate Names:**

Prepatellar Bursitis; Housemaid's Knee; Pes Anserine Bursitis

### **Kneecap**

A flat, rounded bone at the front of the knee, embedded in the tendon of the thigh muscles. It protects the joint and improves the leverage used to straighten the knee; injury or arthritis can cause pain and restricted movement.

---

## Kneecap Dislocation

An injury in which the patella slips completely out of the groove at the end of the thighbone, usually toward the outside. It causes sudden pain, deformity, swelling, and difficulty walking and needs prompt assessment.

## Knee-Chest Position

A position in which the patient rests on the knees and chest, with the hips flexed; it may be used for spinal or anorectal procedures.

## Knee CT Scan

A detailed X-ray scan of the knee made by a computer. It is useful for complex fractures, bone alignment, arthritis, infection, masses, and selected soft-tissue problems, but it exposes the person to more radiation than a plain X-ray.

## Knee Dislocation

Complete anatomical disruption and loss of articulation between the distal femur and proximal tibia, representing a severe orthopaedic limb-threatening emergency frequently associated with multi-ligament rupture and high incidence of popliteal artery injury and peroneal nerve palsy.

## Knee Disorders

Problems affecting the knee joint, including injuries to ligaments or menisci, arthritis, bursitis, tendon disorders, infection, and fractures. They may cause pain, swelling, stiffness, instability, locking, or difficulty bearing weight.

## Knee Effusion

The abnormal accumulation of excess synovial fluid, pus, or blood within the joint capsule of the knee, clinically detected by the ballottement of the patella or bulge sign, warranting diagnostic arthrocentesis to differentiate inflammatory, septic, or hemarthrotic causes.

## Knee Hyperextension

Bending the knee backward beyond its normal range. It can strain or tear ligaments and may cause pain, swelling, instability, or difficulty bearing weight.

## Knee Injury And Meniscus Tears

A meniscus tear is damage to the cartilage cushion between the thighbone and shinbone, often after twisting or with degeneration. Pain, swelling, catching, or locking may occur; treatment ranges from rehabilitation to surgery according to tear, symptoms, and stability.

## Knee-Jerk

The automatic kick of the lower leg when the tendon below the kneecap is tapped. It tests part of the nerve pathway from the thigh and lower spinal cord and can help identify nerve or spinal problems.

## Knee Jerk Reflex

A monosynaptic deep tendon stretch reflex elicited by percussing the patellar ligament below the patella with a reflex hammer, stretching muscle spindles within the quadriceps femoris and provoking reflex knee extension mediated through the femoral nerve and spinal segments L2, L3, and L4.

### Alternate Names:

Patellar Reflex; Quadriceps Reflex

## Knee Joint

A synovial joint formed mainly by the femur, tibia, and patella. It permits flexion and extension with limited rotation and is stabilized by ligaments, menisci, muscles, and the joint capsule.

## Knee Joint Replacement

Surgery replacing damaged surfaces of the knee with artificial parts to relieve severe pain and improve movement. It is commonly used for advanced arthritis; infection, blood clots, stiffness, loosening, persistent pain, or repeat surgery are possible.

## Knee Meniscus

A pair of crescent-shaped fibrocartilaginous pads (medial and lateral menisci) interposed between the femoral condyles and the tibial plateau, functioning to distribute compressive axial loads, absorb shock, lubricate the knee joint, and deepen the tibial articular surfaces.



---

## **Knee Microfracture Surgery**

An operation that makes small holes in the bone beneath a damaged area of knee cartilage so marrow cells can form repair tissue. It may improve symptoms in selected small defects, but the new tissue is not identical to normal cartilage and results vary.

## **Knee MRI Scan**

Magnetic resonance imaging of the knee using a magnetic field and radio waves to evaluate cartilage, menisci, ligaments, tendons, bone marrow, muscles, and joint fluid.

## **Knee Pain**

Pain felt in or around the knee, commonly caused by injury, osteoarthritis, bursitis, tendinitis, or inflammatory disease. Swelling, instability, inability to bear weight, or fever may also occur.

### **Alternate Names:**

Gonalgia; Knee Arthralgia

## **Knee Prosthesis**

An artificial joint component used to replace damaged surfaces of the knee, usually including femoral and tibial components and sometimes a patellar component.

## **Knee Replacement**

An operation that replaces part or all of a damaged knee joint with an artificial joint to reduce pain and improve movement. Recovery includes wound care, exercises, and rehabilitation.

## **Kniest Dysplasia**

A rare inherited disorder of bone and cartilage growth that causes short stature, enlarged joints, unusual facial features, and progressive joint problems. Some people also develop serious eye, hearing, or breathing complications.

## **Knife-Cut Ulcers From Crohn'S Disease**

Deep, narrow ulcers in the intestinal lining that look like cuts and can occur in Crohn disease. They may cause bleeding, narrowing of the bowel, abnormal connections, pain, or blockage and are identified

during intestinal examination.

## **Knock-Knee**

Genu valgum, a leg alignment in which the knees angle inward and may touch while the ankles remain apart.

## **Knock Knees**

Medial angulation of the knees causing the knees to touch while the ankles remain apart, also called genu valgum. It is common during childhood development but may be pathologic when severe, asymmetric, persistent, or associated with pain or growth disturbance.

### **Alternate Names:**

Genu Valgum

## **Knotless Anchor**

A small surgical anchor securing sutures in bone without requiring tied knots. Orthopedic surgeons use it to repair tendons, ligaments, or other soft tissues, with risks including loosening, infection, or tissue failure.

## **Knuckle**

A projecting finger joint, usually a metacarpophalangeal joint; it may be affected by trauma, inflammation, or arthritis.

## **Knuckle Pad**

A firm, usually painless thickening of skin and tissue over a finger joint. It is commonly related to repeated pressure or a fibrous tissue change and is generally harmless.

## **Kocher Clamp**

A surgical instrument with teeth used to hold firm tissue or clamp blood vessels. Its strong grip is useful during operations, but careless use can crush or injure delicate tissue.

## **Kocher-Debre-Semelaigne Syndrome**

A rare pediatric clinical syndrome associated with chronic, untreated congenital or juvenile hypothyroidism, characterized by generalized muscular pseu-

---

dohypertrophy ('infant Hercules' phenotype), muscle stiffness, sluggish deep tendon reflexes, and myxedema.

**Alternate Names:**

Hypothyroid Muscular Pseudohypertrophy; Debré-Sémélaigne Syndrome

**Kocher-Debré-SéMéLaigne Syndrome**

A rare condition in infants or young children with severe, untreated underactive thyroid disease. Muscles may look unusually large but are weak and slow, with delayed growth and development, puffy facial features, and other signs of hypothyroidism. Thyroid-hormone treatment usually improves the muscle changes and weakness.

**Alternate Names:**

Cretinism with Muscular Hypertrophy; Infantile Hypothyroid Myopathy

**Kocher Incision**

An open surgical subcostal oblique abdominal incision made parallel to and approximately 2 to 3 cm below the right costal margin, traditionally utilized for open cholecystectomy and biliary tract surgical access.

**Kocher Maneuver**

A surgical technique in which the peritoneum along the lateral border of the duodenum is sharply incised, mobilizing the duodenum and head of the pancreas medially and anteriorly to gain access to retroperitoneal structures including the inferior vena cava, aorta, and posterior pancreas.

**Alternate Names:**

Duodenal Mobilization; Kocherization

**Koch's Bacillus**

The bacterium that causes tuberculosis, an infection most often affecting the lungs but capable of involving many organs.

**Alternate Names:**

Tubercle Bacillus; *M. tuberculosis*

**Koebnerisation**

Development of lesions of an existing skin disease at

sites of trauma or injury, classically seen in psoriasis, vitiligo, and lichen planus.

**Koebner Phenomenon**

Development of lesions of a skin disease, classically psoriasis or vitiligo, at sites of injury or other trauma.

**Kohler Disease**

A rare, self-limiting childhood osteochondrosis characterized by avascular necrosis of the tarsal navicular bone, occurring predominantly in boys aged 3 to 7 years, presenting with antalgic limp, medial midfoot pain, and localized dorsal swelling.

**Alternate Names:**

Kohler Bone Disease

**KOH Preparation**

A test in which a skin, hair, or nail sample is treated with potassium hydroxide and viewed under a microscope to look for fungi. A negative result does not always rule out infection if the sample is small or treatment has already begun.

**Koilocyte**

A cell with changes often seen after infection with human papillomavirus (HPV). Under a microscope, it has an enlarged or irregular nucleus and a clear space around the nucleus. The finding supports HPV-related change but is not enough by itself to determine the diagnosis.

**Koilonychia**

Thin, brittle nails that become concave and curve upward at the edges, giving them a spoon-like shape. It is often linked to iron deficiency but can also result from poor nutrition, repeated injury, inherited conditions, or other illness; new or widespread changes need evaluation.

**Kombucha**

A fermented tea made with a culture of bacteria and yeast. It may contain sugar, acids, caffeine, and a small amount of alcohol; contaminated or excessive products can cause illness, especially in vulnerable people.

---

## Konno Procedure

Open-heart surgery to widen a severely narrowed path through which blood leaves the left ventricle. The surgeon enlarges the area below and around the aortic valve, often replacing or repairing the valve at the same time. It is used for complex or recurrent obstruction and carries the usual risks of major heart surgery.

### Alternate Names:

Konno-Rastan Procedure; Aortoventriculoplasty

## Koplik Spots

Small white or bluish spots on a red background inside the cheek, usually opposite the back teeth. They are characteristic of measles and often appear shortly before the skin rash.

## Koplik's Spots

Small white or bluish-gray spots on the inner cheek, often with a red base, that are characteristic of measles and may appear shortly before the skin rash.

## Korotkoff Sounds

Sounds heard through a stethoscope over an artery as a blood-pressure cuff is deflated; their appearance and disappearance are used to estimate systolic and diastolic pressure.

## Korsakoff's Syndrome

A chronic amnesic disorder, commonly associated with severe thiamine deficiency and alcohol-use disorder, causing impaired new learning, memory loss, and sometimes confabulation.

## Korsakoff Syndrome

A chronic amnesic disorder caused most often by severe thiamine deficiency associated with alcohol-use disorder. It causes impaired formation of new memories, loss of past memories, and sometimes confabulation.

## Kostmann Syndrome

An autosomal recessive form of severe congenital neutropenia caused by mutations in the HAX1 or

ELANE genes, characterized by early arrest of granulopoiesis at the promyelocyte stage in the bone marrow, profound absolute neutropenia (less than 200/uL), and life-threatening bacterial infections starting in early infancy.

### Alternate Names:

Severe Congenital Neutropenia Type 1; Infantile Genetic Agranulocytosis

## Koumpounophobia

An intense fear or strong distress related to buttons. It is a specific phobia when it causes significant anxiety or interferes with daily life, and it can be treated with psychological therapies such as exposure-based therapy.

## Kounis Syndrome

A heart attack or temporary narrowing of a heart artery triggered by a severe allergic reaction. It may cause chest pain together with rash, swelling, wheezing, or low blood pressure.

## KP

Abbreviation; see *Keratosis Pilaris*.

## Krabbe Disease

A rare inherited disorder in which missing enzymes damage myelin, the protective covering of nerves. Early disease can cause irritability, stiffness, seizures, loss of skills, and progressive problems with movement, vision, or development.

## Krabbe's Disease

A rare inherited disorder in which missing enzymes cause myelin loss, progressively damaging nerves and producing weakness, stiffness, vision problems, or developmental decline.

## KRAS Gene

A gene that helps control cell growth and division. Certain changes can keep its growth signal switched on, contributing to cancers and affecting which targeted treatments are likely to work.

---

## Kraurosis

Older term; see *Lichen Sclerosus* or the specific vulvar condition.

## Kraurosis Vulvae

Advanced progressive atrophy, shrinkage, and sclerosis of the female external genitalia occurring in chronic untreated vulvar lichen sclerosus, characterized by resorption of the labia minora, narrowing of the vaginal introitus, and thin parchment-like epidermal fragility.

### Alternate Names:

Vulvar Lichen Sclerosus; Atrophic Vulvitis

## Krebs Cycle

A series of chemical reactions in mitochondria that oxidizes acetyl-CoA and generates energy-rich molecules used for cellular energy production.

### Alternate Names:

Tricarboxylic Acid Cycle; TCA Cycle

## Kringles

Small folded parts of certain proteins that help them attach to other molecules. They are found in proteins involved in blood clot breakdown and in some growth signals.

## Krukenberg Spindle

A vertical, brownish line of pigment on the inner surface of the cornea. It may occur with pigment dispersion in the eye and can be associated with changes in eye pressure.

### Alternate Names:

Endothelial Pigment Spindle; Pigment Dispersion Corneal Spindle

## Krukenberg's Tumor

Alternate spelling; see *Krukenberg Tumor*.

## Krukenberg Tumor

An ovarian tumor caused by cancer that has spread from another organ, most often the stomach or digestive tract. It may cause abdominal swelling, pain, menstrual changes, or digestive symptoms.

## Kt/V

A number used to estimate how much urea a dialysis treatment removes. It combines the dialyzer's cleaning ability and treatment time with the person's body-water volume; clinicians use it with symptoms, fluid balance, laboratory results, and the dialysis schedule.

## Kugelberg-Welander Disease

A milder form of inherited spinal muscular atrophy, also called type 3 spinal muscular atrophy. It usually begins after 18 months of age and causes slowly worsening weakness in muscles close to the body, especially the hips and thighs. Children may have a waddling walk or trouble climbing stairs. Many walk independently, and life expectancy is often near normal with current care.

### Alternate Names:

Spinal Muscular Atrophy Type 3 (SMA 3); Juvenile Spinal Muscular Atrophy

## Kulchitsky Cells

Specialized hormone-producing cells found mainly in the digestive and respiratory systems. They release chemical signals, including serotonin, that help regulate movement, secretion, and other organ functions.

### Alternate Names:

EC Cells; Argentaffin Cells

## Kupffer Cells

Immune cells lining the small blood channels in the liver. They remove old blood cells, particles, and some germs from blood coming from the digestive organs.

### Alternate Names:

Stellate Macrophages of the Liver

## Kupffer's Cell

A Kupffer cell is a resident macrophage lining the liver sinusoids; it removes microbes, damaged blood cells, and particles from portal blood and contributes to hepatic inflammation and iron handling.

---

## Kuru

Kuru is a rare fatal prion disease historically associated with ritual cannibalism, causing progressive ataxia, tremor, and neurologic decline.

## Kussmaul Breathing

Deep, rapid, labored breathing that helps remove extra carbon dioxide when the blood is dangerously acidic. It can occur in diabetic ketoacidosis, severe kidney failure, or other illnesses.

## Kussmaul Respiration

Alternate name; see *Kussmaul Breathing*.

## Kussmaul Sign

A rise or failure to fall in the pressure of the large neck veins during breathing in. It suggests that the right side of the heart cannot fill normally and may occur with severe heart failure, constrictive pericarditis, or other conditions affecting right-heart filling.

## Kveim-Siltzbach Test

An old skin test once used to support the diagnosis of sarcoidosis. Material from sarcoid tissue was injected under the skin and the area was later biopsied. The test is rarely used now because of safety, availability, and accuracy concerns.

### Alternate Names:

Kveim Test; Sarcoidosis Skin Test

## Kwashiorkor

Severe malnutrition in which too little protein and other nutrients cause swelling, poor growth, muscle loss, skin or hair changes, and weaker immunity. It can occur in children or adults.

## Kyasanur Forest Disease

An acute tick-borne viral hemorrhagic fever caused by Kyasanur Forest disease virus, a flavivirus transmitted to humans and non-human primates primarily by *Haemaphysalis* ticks (*Haemaphysalis spinigera*) in the Western Ghats region of South Asia. Following an incubation period of 3 to 8 days, it presents with sudden onset of high fever, frontal

headache, conjunctival injection, severe prostration, myalgia, and hemorrhagic manifestations (such as epistaxis, hematemesis, or melena), frequently followed by a secondary biphasic stage marked by meningoencephalitis and tremors.

### Alternate Names:

KFD; Monkey Fever; Kyasanur Forest Disease Virus Infection

## Kynurenine

A substance made when the body breaks down the amino acid tryptophan. It is involved in chemical signaling in the brain and immune system and may change during infection, inflammation, or some diseases.

## Kyphoplasty

A minimally invasive procedure that uses a balloon to create space in a collapsed vertebral body and then injects bone cement to stabilize a painful compression fracture.

## Kyphoscoliosis

A combination of sideways curvature and excessive forward rounding of the spine. Severe deformity can reduce chest movement and lung capacity, causing breathlessness or, over time, breathing failure and high blood pressure in the lung arteries.

## Kyphosis

An exaggerated forward curve of the upper spine, sometimes producing a rounded back. Causes include posture, osteoporosis and spinal fractures, growth disorders, arthritis, or inflammation. It may cause pain, weakness, or worsening curvature.

### Alternate Names:

Round Back

## Kyrlle Disease

A perforating dermatosis characterized by large, intensely itchy, keratotic papules that expel abnormal dermal material through the epidermis. It is strongly associated with diabetes and chronic kidney disease and may improve when associated illness is controlled.

---

## Kytococcus

A genus of gram-positive cocci found on skin and in the environment that can rarely cause opportunistic infection.

## Kytococcus Sedentarius

A skin-associated gram-positive coccus that produces enzymes linked to pitted keratolysis and can rarely cause invasive infection.

## Kava

A plant-derived product traditionally used in the Pacific and marketed for anxiety or relaxation. It can cause sedation, drug interactions, and serious liver injury and should not be treated as risk-free because it is herbal.

## KC

Common abbreviation for keratoconus, a progressive thinning and cone-shaped bulging of the cornea; see *Keratoconus*.

## Kegel Exercises

Alternate plural form; see *Kegel Exercise*.

## Keratosis Seborrheic

Alternate word order; see *Seborrheic Keratosis*.

## Knee Replacement Surgery

Surgery that replaces part or all of a damaged knee joint with an artificial joint to reduce pain and improve function; see *Knee Replacement*.

## Kneecap Instability

Recurrent or persistent abnormal movement of the patella within or out of the femoral groove, causing pain, giving way, or dislocation; assessment may include examination and imaging.

## Kneecap Tracking Disorder

Abnormal alignment or movement of the patella during knee bending and straightening, sometimes causing front-of-knee pain, grinding, or instability.

## Kratom

A plant product containing mitragynine and related compounds that can have stimulant or opioid-like effects. It can cause dependence, withdrawal, sedation, seizures, liver injury, and dangerous interactions, especially with other sedatives or opioids.

## KTW Syndrome

Abbreviation; see *Klippel-Trenaunay-Weber Syndrome*.



# L

## LAAO

Alternate name; see *Left Atrial Appendage Occlusion*.

## Labbe Vein

The inferior anastomotic vein of the cerebral cortex that traverses the surface of the temporal lobe, establishing an anastomotic venous connection between the superficial middle cerebral vein (Sylvian vein) and the transverse dural venous sinus.

### Alternate Names:

Inferior Anastomotic Vein; Vein of Labbé

## Labetalol

A medicine that lowers blood pressure by slowing the heart and relaxing blood vessels. It is also used for severe high blood pressure during pregnancy and may cause a slow heartbeat, low blood pressure, or breathing problems in people with asthma.

## Labia

The folds of skin surrounding the entrance to the vagina and urethra, consisting mainly of the labia majora and labia minora.

## Labial Sounds

Speech sounds made with one or both lips, such as the sounds in “p,” “b,” and “m.” Difficulty making them may occur with problems affecting the lips, mouth, nerves, or speech muscles.

## Labia Majora

The paired outer folds of skin of the vulva that help protect the urethral and vaginal openings and contain connective tissue, glands, and fat.

## Labia Minora

The paired inner folds of tissue of the vulva, lying within the labia majora and surrounding the vestibule and its openings.

## Labiaplasty

Surgical reshaping or reduction of the labia, usually the labia minora, performed for functional symptoms, reconstruction, or selected cosmetic indications. Risks include pain, bleeding, infection, scarring, and altered sensation.

## Labile

Easily or rapidly changing, such as blood pressure, blood sugar, mood, or symptoms. Labile measurements may reflect illness, medicines, stress, hormones, or changing treatment and often require repeated observation.

## Labile Hypertension

Blood pressure that changes substantially over short periods. Stress, pain, medicines, illness, and measurement conditions can contribute, so repeated properly performed readings help confirm the pattern.

## Lability

A tendency to change easily or unpredictably, such as rapidly shifting emotions or blood pressure. Its medical importance depends on the symptom, severity, and underlying cause.

## Labium

A lip-like anatomical structure; in obstetrics and gynecology, usually one of the labia of the vulva.

---

## Labium Majus

One of the paired outer folds of the vulva containing skin, connective tissue, glands, vessels, and nerves and protecting the more internal genital structures.

## Labium Minus

One of the paired inner folds of the vulva bordering the vestibule and helping protect the urethral and vaginal openings.

## Labor

The process in which regular contractions open the cervix and lead to delivery of the baby and placenta. It includes the stages of cervical opening, birth of the baby, and delivery of the placenta.

## Laboratory

A facility or department where clinical specimens are examined using microbiologic, hematologic, chemical, molecular, or other diagnostic methods.

## Laboratory Medicine

The medical specialty that uses laboratory tests on blood, tissue, and other specimens for diagnosis, screening, monitoring, and treatment.

## Laboratory Procedure

A method used to collect, prepare, examine, or measure a patient sample such as blood, urine, tissue, or a swab. Results can help diagnose disease or guide treatment and must be interpreted with the clinical findings.

## Laboratory Test

An examination of a body fluid, tissue, or other specimen to obtain information for diagnosis, monitoring, screening, or treatment.

## Labor Complication

A maternal or fetal problem occurring during labor or delivery, such as abnormal labor progress, hemorrhage, infection, fetal distress, or malpresentation, requiring assessment and management based on the cause.

## Labor Dystocia

Labor that is unusually slow or difficult because of weak contractions, the baby's position or size, or the shape of the mother's pelvis. Treatment depends on the cause and may include medicines or an operation.

## Labor Pain

Pain caused by uterine contractions, cervical dilation, and stretching of pelvic tissues during labor; its intensity and management vary with labor stage and individual factors.

## Labor: Signs And Symptoms

Labor is the process of regular uterine contractions that opens the cervix and leads to birth. Possible signs include increasingly regular contractions, rupture of the membranes, a blood-stained mucus discharge, pelvic pressure, and back pain. Heavy bleeding, reduced fetal movement, severe pain, or other complications may occur.

## Labour

The process of childbirth, consisting of regular uterine contractions that cause progressive cervical effacement and dilation, leading to delivery of the baby and placenta.

## Labra

The plural of labrum: tough rings of tissue around certain joint sockets, especially the shoulder and hip. They deepen the socket, improve stability, and help distribute pressure during movement.

## Labral Tear

A tear in the fibrocartilaginous rim of a joint socket, commonly the shoulder or hip, causing pain, clicking, instability, or reduced range of motion.

## Labrum

A ring of tough, flexible tissue around the socket of a joint. It deepens and stabilizes the shoulder or hip socket.

---

## **Labyrinth**

The inner-ear system that contains the cochlea for hearing and the balance organs that sense head movement and position. Disease or injury can cause dizziness, unsteadiness, ringing, or hearing loss.

## **Labyrinth Disorder**

A problem affecting inner-ear hearing or balance structures, potentially causing spinning sensations, nausea, unsteadiness, hearing changes, or abnormal eye movements.

## **Labyrinthine Membrane**

The delicate, fluid-containing system inside the inner ear that contains the organs for hearing and balance. Injury or abnormal fluid pressure can cause dizziness, hearing loss, or ringing in the ear.

## **Labyrinthitis**

Inflammation of the inner ear that can cause sudden spinning dizziness, imbalance, nausea, ringing in the ear, and sometimes hearing loss.

## **Laceration**

A torn wound caused by blunt or irregular force. It may involve only the skin or extend into fat, muscle, tendons, nerves, or blood vessels; deep, gaping, dirty, or heavily bleeding wounds need medical care.

## **Laceration Suture and Staple Care**

Home-care instructions and wound hygiene protocols following wound closure with sutures or staples. Involves keeping the incision clean and dry for the initial 24–48 hours, daily inspection for signs of wound infection (erythema, purulent exudate, localized warmth), gentle cleansing, and scheduled removal by a clinician.

### **Alternate Names:**

Suture Home Care; Staple Wound Management; Laceration Closure Care

## **Lachman Sign**

The main test for a torn knee ligament (ACL); with the knee slightly bent, the doctor pulls the shin forward to check if the joint is loose.

## **Lachman Test**

A knee examination in which the lower leg is gently pulled forward while the knee is slightly bent. Too much movement may suggest an injury to the anterior cruciate ligament.

## **Lachrymal**

Relating to tears or the structures that produce and drain them; also spelled lacrimal.

## **Lacosamide**

An antiseizure medicine used mainly for focal seizures. It may cause dizziness, double vision, poor coordination, or changes in the heart's electrical rhythm.

## **Lacquer Poisoning**

Harm from swallowing or breathing lacquer or its solvents. It may irritate the mouth and lungs, cause dizziness or drowsiness, and cause severe lung injury if liquid enters the lungs.

## **Lacrimal**

Relating to tears or the lacrimal glands and drainage passages.

## **Lacrimal Apparatus**

The glands and passages that make, spread, and drain tears. It includes the tear gland, small openings near the inner eyelids, the tear sac, and the duct that drains tears into the nose.

## **Lacrimal Disease**

A disorder of the tear-producing or tear-draining system, including dry eye, dacryocystitis, obstruction, inflammation, or abnormal tearing.

## **Lacrimal Duct**

A drainage channel that carries tears from the lacrimal sac into the nasal cavity. Blockage can cause watering of the eye, recurrent infection, swelling near the inner corner, or discharge.

## **Lacrimal Gland**

One of two glands above the outer part of the eye

---

that make the watery part of tears. The tears spread across the eye and then drain through passages near the inner corner.

### **Lacrimal Gland Tumor**

A benign or malignant growth arising in the tear-producing lacrimal gland, often causing swelling, pain, or displacement of the eye.

### **Lacrimal Sac**

A small sac at the inner corner of the eye that collects tears before they pass through the lacrimal duct into the nose.

### **Lacrimation**

Production and release of tears by the lacrimal glands; excessive tearing may result from irritation, allergy, infection, or blocked tear drainage.

### **La Crosse Encephalitis**

A mosquito-borne viral encephalitis caused by La Crosse virus, usually affecting children and causing fever, headache, seizures, or altered consciousness.

### **Lactam**

A ring-shaped chemical structure found in some medicines. Beta-lactam antibiotics use this structure to weaken bacterial cell walls, but allergies and bacterial resistance can limit their use.

### **Lactase**

An enzyme in the small intestine that breaks lactose, the natural sugar in milk, into absorbable sugars. Low lactase can cause bloating, cramps, gas, diarrhea, or discomfort after dairy foods.

### **Lactase Deficiency**

Alternate name; see *Lactose Intolerance*.

### **Lactate**

A product of cellular metabolism measured in blood. An increased level may occur with strenuous exercise, impaired oxygen delivery, infection, medicines, liver disease, or other conditions and must be interpreted clinically.

### **Lactate Dehydrogenase**

An enzyme found in many body tissues. Its blood level may rise when cells are damaged, but the test does not identify the cause by itself.

### **Lactate Dehydrogenase Test**

A blood test measuring lactate dehydrogenase, an enzyme released when cells are injured; increased activity is nonspecific and may occur with hemolysis, liver disease, muscle injury, infection, or malignancy.

### **Lactation**

The production and release of breast milk after childbirth. Early milk, called colostrum, is followed by mature milk; hormones help make and release the milk.

#### **Alternate Names:**

Milk Secretion; Galactopoiesis

### **Lactational Amenorrhea**

The temporary absence of menstrual periods during breastfeeding because ovulation is suppressed. Breastfeeding prevents pregnancy reliably only when specific conditions are met, so other contraception may be needed.

### **Lactation Disorder**

A problem with making, releasing, or transferring breast milk after childbirth. Examples include low supply, excess milk, painful engorgement, mastitis, or difficulty with the baby's latch; assessment considers both parent and infant.

### **Lactation Mastitis**

Alternate name; see *Mastitis*.

### **Lactation Nutrition**

Eating and drinking enough during breastfeeding to support the parent's health and milk production. Most people need a varied diet and regular fluids; supplements may be needed for a diagnosed deficiency or specific dietary pattern.

### **Lacteals**

Tiny lymph vessels in the lining of the small intes-

---

tine that absorb most digested fats and fat-soluble vitamins. They carry these nutrients through the lymph system before they enter the bloodstream.

### **Lactic Acid**

An organic acid produced during normal metabolism, especially when pyruvate is converted to lactate. Blood lactate may rise for many reasons, including impaired oxygen delivery, strenuous exercise, or medicines.

### **Lactic Acidosis**

Metabolic acidosis associated with increased lactate production or reduced lactate clearance. Causes include impaired tissue oxygen delivery, medicines, liver disease, seizures, and other disorders.

### **Lactic Acid Test**

A blood test measuring lactate, which accumulates when production exceeds clearance; it helps assess tissue hypoperfusion, sepsis, severe exercise, metabolic disease, or other causes of lactic acidosis.

### **Lactiferous Duct**

One of 15 to 20 excretory ducts in the female mammary gland that drains milk from glandular lobules toward the nipple, expanding into a lactiferous sinus beneath the areola before opening onto the nipple surface.

#### **Alternate Names:**

Galactophore; Mammary Duct

### **Lactobacillus**

A genus of bacteria that produces lactic acid and normally forms part of the healthy vaginal and intestinal microbiota; some species are used as probiotics.

### **Lactoferrin**

Lactoferrin is an iron-binding glycoprotein in milk and mucosal secretions that limits microbial iron access and contributes to innate immune defense.

### **Lactose**

The principal sugar in milk, composed of glucose

and galactose. Lactase in the small intestine digests it.

### **Lactose Fermentation**

The ability of bacteria to break down lactose and produce acid or gas. It is used in the laboratory to help identify different types of bacteria.

### **Lactose Intolerance**

A condition in which the body cannot fully digest lactose, the sugar in milk and dairy products, because it does not make enough lactase. It may cause bloating, gas, abdominal pain, nausea, or diarrhea after consuming lactose.

### **Lactose Tolerance Test**

A test for difficulty digesting lactose, the sugar in milk. After lactose is consumed, blood glucose or breath hydrogen is measured; symptoms and results help assess lactose malabsorption but do not diagnose every cause of digestive discomfort.

### **Lactose Tolerance Tests**

Tests that evaluate digestion and absorption of lactose by measuring blood glucose response after lactose ingestion or detecting hydrogen in exhaled breath; they may support evaluation of lactose malabsorption.

### **Lactulose**

A medicine that softens stool by drawing water into the bowel. It is also used to lower harmful ammonia levels in some people with severe liver disease. Gas and diarrhea can occur.

### **Lacuna**

A small space or cavity in tissue, such as a space containing an osteocyte in bone.

### **Lacunar Infarct**

A small stroke in a deep part of the brain caused by blockage of one of the small arteries supplying that area. It can affect movement, feeling, speech, or coordination, depending on the location.

#### **Alternate Names:**

---

Lacunar Stroke; Subcortical Small Vessel Infarction

### **Lacunar Stroke**

A small ischemic stroke caused by blockage of a tiny artery deep in the brain, often producing weakness, numbness, clumsiness, or speech problems.

### **Lacunar Syndromes (Small-Vessel Strokes)**

Clinical patterns caused by small deep cerebral infarcts in penetrating-artery territories. Recognized patterns include pure motor hemiparesis, pure sensory stroke, sensorimotor stroke, ataxic hemiparesis, and dysarthria-clumsy-hand syndrome.

### **Ladd Bands**

Abnormal bands of tissue present at birth that can cross and squeeze the first part of the small intestine when the bowel is in an abnormal position. They may cause green vomiting and blockage in an infant and are released during the Ladd procedure.

#### **Alternate Names:**

Ladd's Peritoneal Bands; Ladd Fibrous Bands

### **Ladd Procedure**

An operation for a child whose intestines developed in an abnormal position and may twist. The surgeon untwists the intestine, releases bands causing blockage, and broadens its attachment to reduce future twisting.

#### **Alternate Names:**

Ladd Operation; Ladd's Procedure

### **Laetrile**

A semisynthetic amygdalin-related compound promoted as an alternative cancer treatment; credible evidence has not established anticancer benefit, and cyanide poisoning is a serious risk.

### **Lafora Disease**

A rare inherited epilepsy that usually begins during adolescence and progressively worsens. Seizures, brief muscle jerks, visual symptoms, memory problems, and loss of physical or thinking abilities may occur.

### **Lagophthalmos**

Inability to close the eyelids completely. Exposed cornea can become dry, irritated, scratched, infected, or damaged, especially during sleep, so lubricating treatment and protection may be needed.

### **Lambda**

A landmark on the back of the skull where two skull seams meet. It is used to describe head shape and the position of nearby structures, especially in newborn examinations and imaging.

### **Lambert-Eaton Myasthenic Syndrome**

A rare condition in which the immune system interferes with nerve signals to muscles. It causes weakness, especially in the hips and shoulders, reduced reflexes, and sometimes dry mouth. It can be associated with small-cell lung cancer.

### **Lamellar**

Made of thin layers or plates. The word may describe the organized layers of mature bone, the skin barrier, or structures in the eye and nervous system. Its exact meaning depends on the body structure being described.

### **Lamellar Body**

A small storage structure inside certain cells that contains layered fats. In lung cells, it stores and releases surfactant, the substance that helps keep the tiny air sacs open.

### **Lamellar Bone**

Mature bone made of organized layers of mineralized tissue. It forms most adult compact and spongy bone and provides strength while being continually remodeled.

### **Lamellar Ichthyosis**

An inherited skin disorder present from birth that causes dry skin covered with large, dark, plate-like scales over much of the body, including the arms and legs. It may reduce sweating and cause overheating, and can affect the hair, nails, eyes, or palms and soles.



---

## Lamin

A structural protein forming part of the nuclear lamina, which supports the inner surface of the cell nucleus.

## Lamina

A flat piece of bone that forms part of the back of a vertebra. The two laminae help complete the bony arch around the spinal cord and can be removed during some spine operations.

## Lamina Cribrosa

A fenestrated, mesh-like collagenous sheet of the sclera through which unmyelinated axons of retinal ganglion cells exit the eye to form the optic nerve; the primary site of axonal injury and excavation in glaucoma.

## Lamina Propria

A thin layer of supporting tissue beneath the inner lining of organs such as the intestine, airways, and bladder. It contains small blood vessels, immune cells, glands, and connective tissue.

## Laminectomy

An operation that removes part of a spinal bone to make more room for the spinal cord or nerves. It may be used for spinal narrowing, a tumor, or selected disk problems.

## Laminin

A protein in the supporting framework around cells that helps cells attach to underlying tissue. It contributes to tissue structure, development, repair, and the filtering barrier in some organs.

## Lamivudine

Lamivudine is a nucleoside reverse-transcriptase inhibitor used with other antiretrovirals for HIV and as an antiviral for chronic hepatitis B; resistance and rare lactic acidosis or hepatic complications require monitoring.

## Lamotrigine

Lamotrigine is an antiseizure medicine that inhibits

voltage-gated sodium channels and is also used for bipolar-maintenance treatment; rapid dose escalation increases the risk of serious rash including Stevens–Johnson syndrome.

## Lancet

A small, sterile, usually single-use device with a sharp point used to puncture the skin and obtain a capillary-blood sample. Lancets are commonly used for fingertip glucose testing and other point-of-care blood tests.

### Alternate Names:

Blood Lancet; Fingerstick Lancet

## Landolt Ring

A ring-shaped symbol with a gap used to test visual sharpness. The person identifies which direction the gap faces; smaller symbols indicate better visual acuity.

## Langerhans' Cell

An immune cell in the skin that captures foreign substances and helps activate T-cell responses.

## Langerhans Cell Histiocytosis

A rare disorder in which abnormal immune cells build up in one or more organs, commonly the bones, skin, lungs, lymph nodes, or pituitary gland. Symptoms and treatment depend on which organs are affected.

## Langerhans-Cell Histiocytosis

A rare disorder in which abnormal immune cells collect in one or more organs, especially bone, skin, lungs, lymph nodes, or the pituitary gland. Symptoms depend on the organs involved and may include bone pain, rash, cough, or hormone problems.

## Langerhans Cells

Immune cells in the outer skin that capture foreign substances and help start immune responses. They connect skin defenses with nearby lymph tissue and can contribute to allergic or inflammatory reactions.

---

## Langer Lines

Natural lines of skin tension that reflect the predominant orientation of collagen fibers in the dermis and may influence surgical scar direction.

## Language Barrier

Difficulty communicating because a patient and healthcare professional do not share a language well enough for safe understanding. A trained interpreter helps explain symptoms, consent, instructions, and risks.

## Language Development

The gradual development of understanding, sounds, words, grammar, and communication during childhood. Delays may be related to hearing loss, developmental conditions, neurologic disease, or limited language exposure.

## Language Disorder

A developmental or acquired impairment in understanding, producing, or using spoken, written, or signed language that affects communication and daily function.

## Language Therapy

Treatment that improves understanding, expression, reading, writing, or other language skills after developmental or acquired impairment.

## Lanolin

Lanolin is a waxy substance derived from sheep wool used as an emollient and in topical products; allergy or irritation can occur.

## Lanolin Poisoning

Swallowing a small amount of lanolin, a wool-derived moisturizer, usually causes little more than stomach upset. Large amounts or products with added medicines can cause more serious poisoning.

## Lansoprazole

A medicine that strongly reduces stomach acid. It is used for heartburn, ulcers, and some *H. pylori* infections; long-term use may affect mineral or vitamin

levels and infection risk.

## Lanthanum

A chemical element whose carbonate salt can bind dietary phosphate and lower phosphate absorption in people with advanced kidney disease.

## Lanugo

Fine, soft hair that grows all over the body of a fetus and is typically shed before birth.

## Lanugo Hair

Lanugo: the fine, soft, usually unpigmented hair covering a fetus and sometimes present on a premature newborn; it is normally shed before or shortly after birth.

## Laparoscope

A slender instrument with a light and camera used to view the abdominal or pelvic cavity through a small incision during minimally invasive surgery.

## Laparoscopic Cholecystectomy

Surgical removal of the gallbladder through small abdominal incisions using a laparoscope and instruments, commonly performed for symptomatic gallstones or gallbladder inflammation.

## Laparoscopic Colectomy

Laparoscopic colectomy is removal of part or all of the colon through small incisions using a camera and specialized instruments.

## Laparoscopic Gallbladder Removal

Surgery to remove the gallbladder through several small abdominal openings using a camera and instruments. It commonly treats symptomatic gallstones or gallbladder inflammation; bleeding, infection, bile leakage, and injury to bile ducts are possible.

## Laparoscopic Gastric Banding

A bariatric operation that places an adjustable band around the upper stomach to create a small pouch and restrict food passage; it is less commonly performed than other bariatric procedures.

---

## Laparoscopic Hysterectomy

Removal of the uterus using laparoscopic instruments through small abdominal incisions. The cervix, fallopian tubes, and ovaries may or may not also be removed; the approach and extent depend on the indication, anatomy, and surgical plan.

## Laparoscopic Nephrectomy

A minimally invasive surgical procedure in which a diseased or donor kidney is dissected and extracted using laparoscopy or robotic assistance, categorized as radical (en bloc removal with Gerota fascia and adrenal gland for renal cell carcinoma) or partial (nephron-sparing surgery).

## Laparoscopic Surgery

Surgery performed through small cuts using a camera and long instruments. It often causes less pain and faster recovery than open surgery, but still has risks such as bleeding, infection, and injury to nearby organs.

## Laparoscopy

Keyhole surgery that uses a small camera and instruments through small cuts in the abdomen or pelvis. It can be used to look for a problem, take a sample, or treat conditions such as appendicitis, gallstones, endometriosis, or an ectopic pregnancy. Recovery is often easier than after open surgery, but risks and recovery time depend on the procedure.

### Alternate Names:

Keyhole Surgery; Minimally Invasive Abdominal Surgery; Diagnostic Laparoscopy; Laparoscopic Surgery

## Laparotomy

An operation that uses a larger cut in the abdomen to reach the abdominal or pelvic organs.

## Lapatinib

Lapatinib is a targeted cancer medicine that blocks HER2 and epidermal-growth-factor receptors and is used for selected advanced breast cancers.

## Lapatinib Ditosylate

Lapatinib ditosylate is the salt form of lapatinib, a targeted medicine used for selected HER2-positive breast cancers.

## Laplace Law

A biophysical principle stating that the wall tension ( $T$ ) in a hollow sphere or cylinder is directly proportional to the transmural pressure ( $P$ ) times the radius ( $r$ ) divided by wall thickness (or  $2 \times$  thickness for a sphere), explaining ventricular wall stress in cardiac dilation and alveolar stability with surfactant.

## Large Airway Obstruction

A blockage in the windpipe or large breathing tubes. It can cause noisy breathing, shortness of breath, and difficulty clearing mucus and may be caused by narrowing, a tumor, or weakness of the airway wall.

## Large Bowel Obstruction

A blockage in the colon, commonly caused by cancer, twisting of the bowel, scar tissue, or diverticular disease. It can cause increasing swelling, cramping, vomiting, and inability to pass stool or gas.

## Large Bowel Resection

Surgical removal of a segment of the colon or rectum, followed by reconnection of the bowel or creation of a stoma when necessary to treat cancer, obstruction, ischemia, inflammation, or other disease.

## Large Colon

The large colon is the main portion of the large intestine that absorbs water, stores fecal material, and moves contents toward the rectum.

## Large For Gestational Age

A fetus or newborn larger than expected for the number of weeks of pregnancy, usually above the 90th percentile. Maternal diabetes, family size, and excess pregnancy weight gain can contribute and may increase birth complications.

---

## Large Intestine

The final part of the digestive tract, extending from the small intestine to the rectum. It absorbs water and helps form and move stool.

## Lesser Toe Deformities

Deformities in which one or more toes bend abnormally at the middle or end joints. They may cause pain, rubbing, corns, calluses, or difficulty wearing shoes.

### Alternate Names:

Claw Toe; Hammer Toe; Hammertoe; Mallet Toe

## Lupus Arthritis

Joint pain or inflammation associated with systemic lupus erythematosus. It commonly affects multiple joints and is considered alongside other lupus manifestations and possible overlapping disease.

## Luteinized Follicle Cyst

A usually harmless ovarian cyst that develops during pregnancy as hormone-sensitive ovarian tissue enlarges. It may be found by chance or as a pelvic mass and often settles after pregnancy.

## Laron Syndrome

A rare inherited growth disorder in which the body cannot respond properly to growth hormone. It causes severe short stature, characteristic facial features, and very low levels of the growth signal needed for normal bone development.

## Larva Currens From Strongyloides

A rapidly moving, itchy skin rash caused by *Strongyloides* larvae moving through the skin. It often appears around the buttocks or trunk and may indicate ongoing intestinal infection requiring antiparasitic treatment.

## Larva Migrans

A condition caused by migrating larval parasites in human tissues, including cutaneous larva migrans in the skin and visceral larva migrans in internal organs.

## Laryngeal

Relating to the larynx, the organ in the throat containing the vocal cords and helping protect the airway.

## Laryngeal Botulinum Toxin Injection

Targeted electromyography- or laryngoscopy-guided injection of botulinum neurotoxin into specific intrinsic laryngeal muscles (such as the thyroarytenoid or posterior cricoarytenoid) to reduce involuntary spasms in focal laryngeal dystonias, predominantly adductor or abductor spasmodic dysphonia.

### Alternate Names:

Laryngeal Botulinum Toxin Injection; Laryngeal Botox; Spasmodic Dysphonia Injection

## Laryngeal Cancer

Cancer of the larynx, or voice box. Tobacco, alcohol, and some HPV infections increase risk. Persistent hoarseness, painful or difficult swallowing, noisy breathing, or a neck lump may occur.

## Laryngeal Candidiasis

A yeast infection of the voice box, usually causing hoarseness, throat pain, or difficulty swallowing. It is more likely with weakened immunity or inhaled steroid use and may require examination of the larynx and antifungal treatment.

## Laryngeal Carcinoma

Laryngeal carcinoma is cancer arising in the voice box and may cause hoarseness, swallowing difficulty, pain, or breathing problems.

## Laryngeal Cartilage

Laryngeal cartilage forms the structural framework of the voice box, maintains the airway, and supports vocal-fold movement.

## Laryngeal Cleft

A laryngeal cleft is a congenital opening between the airway and esophagus that can allow aspiration and cause feeding or breathing problems.

---

## Laryngeal Edema

Swelling of the tissues of the larynx that can narrow the upper airway and cause hoarseness, stridor, or respiratory distress. It may result from allergy, infection, trauma, or airway instrumentation.

## Laryngeal Fistula

An abnormal passage between the voice box and nearby tissue or the swallowing tract. It can allow food or saliva to enter the airway and cause coughing, repeated chest infections, or breathing problems.

## Laryngeal Hemorrhage

Bleeding within or around the larynx, which may result from trauma, instrumentation, vocal strain, vascular lesions, infection, or tumor and can threaten the airway.

## Laryngeal Masks

Laryngeal masks are supraglottic airway devices placed above the vocal cords to maintain ventilation during anesthesia or emergencies.

## Laryngeal Mucositis

Inflammation and open sores in the lining of the voice box, often after radiation, chemotherapy, infection, or irritation. It can cause hoarseness, throat pain, painful swallowing, or breathing difficulty.

## Laryngeal Nerve

A branch of the vagus nerve supplying the larynx; injury can cause hoarseness, swallowing difficulty, or breathing problems.

## Laryngeal Nerve Damage

Injury to the recurrent or superior laryngeal nerves causing hoarseness, vocal-cord weakness, swallowing difficulty, or airway compromise. Common causes include thyroid or neck surgery, malignancy, trauma, and compression along the nerve pathway.

## Laryngeal Obstruction

Partial or complete blockage of the laryngeal airway caused by edema, foreign body, tumor, trauma,

infection, vocal-cord dysfunction, or other structural disease.

## Laryngeal Papillomatosis

Benign papillomatous growths caused by HPV in the larynx or respiratory tract. They may cause hoarseness, stridor, or airway obstruction.

### Alternate Names:

Recurrent Respiratory Papillomatosis

## Laryngeal Prosthesis

A device used to help restore speech after removal of the larynx, commonly through a one-way valve placed between the trachea and esophagus.

## Laryngeal Stridor

A harsh, high-pitched noise caused by turbulent airflow through a narrowed upper airway, often most noticeable during inspiration.

## Laryngeal Web

A laryngeal web is a thin membrane across part of the airway, usually congenital or acquired after injury, causing hoarseness or airway narrowing.

## Laryngectomy

A person who has undergone laryngectomy, surgical removal of all or part of the larynx; total laryngectomy separates the airway from the mouth and nose.

## Laryngectomy

Surgery to remove part or all of the larynx, usually for cancer or severe structural disease. After total removal, breathing occurs permanently through an opening in the neck; partial surgery may preserve some voice and swallowing.

## Laryngectomy Tube

A tube placed in the neck opening after larynx surgery to help keep the airway open and allow suctioning or ventilation when needed. Mucus blockage, bleeding, infection, and tissue overgrowth can occur.

---

## Laryngitis

Inflammation of the voice box that causes hoarseness or temporary voice loss. Common causes include viral infection, voice overuse, smoke or other irritation, and reflux.

## Laryngocele

An air-filled pouch formed when a small space beside the vocal cords becomes enlarged. It may cause hoarseness, a neck swelling that changes with pressure, coughing, or blockage of the airway.

## Laryngomalacia

A birth condition in which soft tissue above an infant's vocal cords partly collapses during breathing in, causing noisy breathing. It often improves with growth, but feeding trouble, pauses in breathing, or poor growth may occur.

## Laryngopharyngeal Reflux

Backflow of stomach contents into the throat and voice box. It may cause hoarseness, frequent throat clearing, cough, a lump-like feeling, excess mucus, or throat tightness, often without typical heartburn.

## Laryngopharynx

The lower part of the throat behind the voice box, leading to the esophagus. It helps direct swallowed food away from the airway and is a site where some throat cancers can develop.

## Laryngoscope

An instrument used to see the throat and vocal cords, especially while placing a breathing tube. Common blades are curved Macintosh and straight Miller types; use can injure teeth, lips, or throat.

## Laryngoscopy

An examination of the voice box and nearby throat using a mirror or a thin camera passed through the mouth or nose. It helps investigate hoarseness, swallowing problems, airway narrowing, bleeding, or suspected tumors.

## Laryngoscopy And Nasolaryngoscopy

An examination of the voice box and nearby airway using a small rigid or flexible camera passed through the mouth or nose. It helps investigate hoarseness, swallowing difficulty, noisy breathing, bleeding, or a suspected growth.

## Laryngospasm

Sudden tightening of the vocal cords that partly or completely blocks breathing. It can be triggered by saliva, blood, reflux, airway procedures, or anesthesia and may cause noisy breathing, choking, or inability to breathe.

## Laryngostenosis

Narrowing of the voice box, or larynx, that can restrict breathing. Causes include scarring after intubation, injury, inflammation, tumors, and congenital narrowing; noisy breathing or breathlessness may occur.

## Laryngotracheobronchitis

Alternate name; see *Croup*.

## Larynx

The voice box in the front of the neck between the throat and windpipe. It contains the vocal cords, produces sound, helps keep food and liquid out of the lungs, and regulates airflow during breathing.

## Larynx Cancer

Laryngeal cancer is a malignant tumor of the voice box and may cause persistent hoarseness, swallowing pain, neck mass, ear pain, cough, or breathing difficulty. Examination and biopsy are used for diagnosis; treatment depends on site and stage.

### Alternate Names:

Laryngeal Carcinoma; Cancer of the Voice Box

## Larynx Disease

Any disorder affecting the larynx, including inflammation, infection, structural lesions, vocal-cord dysfunction, neurologic disease, or cancer.



---

## Larynx Muscle

A muscle of the voice box that moves the vocal folds and helps protect the airway and produce sound.

## Larynx Neoplasm

An abnormal growth in the voice box. It may be noncancerous or cancerous and can cause hoarseness, swallowing difficulty, pain, breathing problems, or a neck lump.

## Lasegue Sign

A physical examination test for lumbosacral nerve root irritation (radiculopathy / sciatica, particularly L4, L5, or S1); with the patient supine, passive straight leg raising of the symptomatic leg provokes sharp radicular pain radiating down the posterior leg below the knee between 30 and 70 degrees of hip flexion.

### Alternate Names:

Straight Leg Raise Test; SLR Test

## Laser

A device that produces a concentrated beam of light. Depending on its wavelength and power, it can measure tissue, cut or seal tissue, remove abnormal growths, or treat selected eye and skin conditions.

## Laser Ablation

The use of focused laser energy to heat, destroy, or seal targeted tissue. The technique is used in selected vascular, ophthalmic, dermatologic, and other procedures.

## Laser-Assisted In Situ Keratomileusis

A refractive surgical procedure that reshapes the cornea using a laser to correct short-sightedness, long-sightedness, or astigmatism, thereby reducing dependence on glasses or contact lenses.

### Alternate Names:

LASIK

## Laser Assisted Uvula Palatoplasty

A procedure using laser energy to reshape or remove selected soft-palate and uvula tissue, intended mainly to reduce snoring in carefully selected pa-

tients.

### Alternate Names:

LAUP

## Laser Fiber

A thin optical fiber that carries laser energy into the body during treatment. It can break urinary stones, destroy selected tumors, seal blood vessels, or treat other targeted tissues.

## Laser Photocoagulation

A laser treatment that seals or destroys leaking or abnormal blood vessels, especially in the retina, helping limit bleeding, swelling, or damaging new vessel growth.

## Laser Surgery

Laser surgery uses a focused beam of light to cut, remove, seal, or reshape tissue in selected procedures.

## Laser Surgery For The Skin

Use of concentrated light to remove, vaporize, seal, or reshape selected skin tissue. It may treat scars, abnormal blood vessels, unwanted hair, or some lesions; redness, pigment change, burns, scarring, and infection are possible.

## Laser Therapy

Laser therapy uses controlled light energy to treat selected skin, eye, vascular, dental, or other medical conditions.

## Laser Therapy For Cancer

Use of an intense, focused beam of light to destroy or shrink selected cancer tissue or control bleeding and blockage. It is useful only for certain tumors and sites; treatment may be combined with surgery, medicines, or radiation.

## Laser Trabeculoplasty

A laser treatment for some forms of open-angle glaucoma that improves drainage of fluid from the eye and lowers eye pressure. The effect may be temporary, and continued examinations or other treatment may still be needed.

---

## **LASIK**

A laser operation that reshapes the clear front surface of the eye to correct short-sightedness, long-sightedness, or astigmatism. It can reduce dependence on glasses, but dry eye, glare, infection, under-correction, and other vision problems are possible.

## **LASIK Eye Surgery**

A laser operation that reshapes the cornea to correct short-sightedness, long-sightedness, or astigmatism and reduce dependence on glasses. Dry eyes, glare, infection, under- or over-correction, and rarely serious vision loss can occur.

## **Lassa Fever**

A viral illness that can cause fever, weakness, vomiting, bleeding, and damage to several organs. It is usually spread by contact with urine or droppings of infected rodents in parts of West Africa and sometimes by contact with body fluids.

## **Lassa Virus**

An arenavirus that causes Lassa fever, a potentially severe hemorrhagic illness acquired mainly through contact with infected rodent excreta or body fluids.

## **Lassitude**

A state of marked fatigue, weakness, or lack of energy that may occur with systemic illness, anemia, endocrine disease, infection, medication effects, or psychological conditions.

## **Last Menstrual Period**

The first day of the most recent normal menstrual bleeding. Clinicians use it to estimate how many weeks a pregnancy has lasted and the expected delivery date, but the estimate is less reliable with irregular cycles or uncertain dates.

## **Latanoprost**

Latanoprost is a prostaglandin F<sub>2</sub>-alpha analogue eye drop that increases uveoscleral outflow and lowers intraocular pressure in glaucoma or ocular hypertension; iris pigmentation and eyelash growth may occur.

## **Latarjet Procedure**

Shoulder-stabilization surgery that moves part of the coracoid bone, with attached tissue, to the front edge of the shoulder socket. It helps prevent repeated dislocation when bone loss or failed previous treatment makes other repairs unsuitable.

## **Late Decelerations**

A repeated, gradual slowing of a baby's heart rate that starts after a uterine contraction begins and recovers after the contraction ends. It can indicate reduced oxygen transfer through the placenta during labor.

## **Late Effects**

Health problems appearing months or years after cancer treatment, including heart, lung, hormone, fertility, nerve, bone, or second-cancer complications.

## **Late Gadolinium Enhancement**

A heart-MRI finding in which contrast dye remains longer in areas of heart muscle that are scarred, injured, or inflamed. Its pattern helps doctors estimate the cause and extent of heart damage.

### **Alternate Names:**

LGE; Delayed Contrast Enhancement; Myocardial Scar Imaging

## **Latent**

Present but inactive, hidden, or not currently producing symptoms, as in latent tuberculosis infection or a latent virus.

## **Latent Autoimmune Diabetes In Adults**

Diabetes caused by the immune system gradually damaging insulin-producing cells in an adult. It may first resemble type 2 diabetes, but insulin production usually declines over time and insulin treatment may eventually be needed.

### **Alternate Names:**

Type 1.5 Diabetes; Late-Onset Autoimmune Diabetes of Adulthood

---

### **Latent Infection**

Persistent infection in which an infectious agent remains in a host in a nonreplicating or intermittently replicating state without active clinical disease. Reactivation can occur when immune control is impaired, as in latent tuberculosis.

### **Latent Phase Of Labor**

The early part of the first stage of labor, beginning with regular contractions and gradual cervical change before active labor. The transition to active labor is assessed clinically, and the duration varies widely between individuals.

### **Latent Syphilis**

Latent syphilis is syphilis infection without current symptoms, identified by testing and classified by duration to guide treatment and follow-up.

### **Latent Tuberculosis**

Infection with *Mycobacterium tuberculosis* in which the bacteria remain inactive and cause no symptoms but may later become active.

### **Latent Tuberculosis Infection**

An immune response to *Mycobacterium tuberculosis* without clinical or microbiologic evidence of active disease. It is usually asymptomatic and not contagious, but it can reactivate; treatment decisions depend on age, immune status, exposure risk, test results, and drug interactions.

### **Late-Onset Hypogonadism**

A condition in which an older man has symptoms of low testosterone and repeatedly low early-morning blood levels. Symptoms may include reduced sexual desire, erection problems, tiredness, reduced muscle strength, increased body fat, weak bones, or low mood. Other illnesses and medicines should also be assessed before treatment.

#### **Alternate Names:**

LOH; Age-Related Testosterone Deficiency; Testosterone Deficiency Syndrome

### **Late Preterm Birth**

Births occurring from 34 weeks 0 days through 36 weeks 6 days of gestation. These infants have greater risks than term infants of respiratory, temperature, glucose, jaundice, feeding, and readmission problems.

#### **Alternate Names:**

34-36 Weeks

### **Lateral**

Toward or located at the side, away from the body's midline. For example, the ears are lateral to the nose; the opposite term is medial.

### **Lateral Canthotomy And Cantholysis**

An emergency operation that opens the outer corner of the eyelids and releases tight supporting tissue around the eye. It lowers dangerous pressure after severe eye injury or bleeding and may protect vision.

### **Lateral Collateral Ligament Injury**

A stretch or tear of the strong ligament on the outer side of the knee, often caused by a blow to the inner knee or a force pushing the lower leg inward. It causes outer-knee pain and may make the knee unstable.

### **Lateral Cuneiform**

One of three small bones in the middle of the foot. It lies between the other cuneiform bones and the cuboid, joins the third toe bone, and helps support the foot's arch.

### **Lateral Cutaneous Nerve of the Thigh**

A sensory nerve arising from the L2 and L3 ventral rami of the lumbar plexus that passes under the lateral end of the inguinal ligament to provide sensation to the anterolateral thigh; entrapment here causes meralgia paresthetica.

#### **Alternate Names:**

Lateral Femoral Cutaneous Nerve; LFCN

### **Lateral Decubitus Position**

A side-lying position in which the patient rests on the nonoperative side to provide access to the chest,

---

hip, or flank.

### **Lateral Elbow Tendinopathy**

Alternate name; see *Tennis Elbow*.

### **Lateral Epicondylitis**

Pain and tendon damage at the outer elbow, usually from repeated gripping or wrist movement rather than tennis alone. Pain and tenderness worsen when lifting, gripping, or extending the wrist.

**Alternate Names:**

Lateral Epicondylitis; Lateral Elbow Tendinopathy

### **Lateral Epicondylitis**

Alternate name; see *Tennis Elbow*.

### **Lateral Geniculate Nucleus**

A laminated thalamic sensory relay nucleus that receives visual information directly from retinal ganglion cell axons via the optic tract and projects through the optic radiations (geniculocalcarine tract) to the primary visual cortex (Brodmann area 17).

**Alternate Names:**

LGN; Lateral Geniculate Body

### **Lateral Medullary Syndrome**

A stroke in the outer part of the lower brainstem. It may cause dizziness, poor coordination, trouble swallowing or speaking, loss of feeling on one side of the face, and loss of pain or temperature sensation on the opposite side of the body.

**Alternate Names:**

Wallenberg Syndrome; PICA Syndrome

### **Lateral Meniscus**

The lateral meniscus is a crescent-shaped cartilage pad on the outer side of the knee that distributes load and stabilizes movement.

### **Lateral Pontine Syndrome**

A stroke in the outer part of the bridge-like section of the brainstem. It may cause facial weakness, hearing loss, dizziness, poor coordination, and loss of pain or temperature sensation on the opposite

side of the body.

**Alternate Names:**

Marie-Foix Syndrome; AICA Infarction Syndrome

### **Lateral Rotation**

Turning a limb or body part away from the body's midline. Rotating the hip so the knee and foot point outward is one example; limited or painful movement may indicate injury.

### **Lateral Sclerosis**

Damage affecting nerve pathways along the side of the spinal cord that carry movement signals. It may cause increasing stiffness and weakness, as in some motor-neuron diseases; the exact diagnosis requires the associated clinical findings.

### **Lateral Traction**

Lateral traction is a pulling force applied sideways to a body part to improve alignment, reduce a dislocation, or relieve pressure.

### **Lateral Ventricle**

One of two fluid-filled spaces inside the brain that produce and contain cerebrospinal fluid, helping protect and support the brain.

### **Lateral X-Ray**

An X-ray image taken from the side, providing a lateral view of the body part being examined.

### **Latex Agglutination Test**

A laboratory test in which tiny coated particles clump together when they meet a matching substance. The reaction can help detect selected germs, antibodies, or proteins in a patient sample.

### **Latex Allergy**

An immune-mediated hypersensitivity reaction to natural rubber latex. Contact may cause dermatitis, urticaria, rhinitis, asthma, or anaphylaxis, especially in healthcare and occupational settings.

**Alternate Names:**

Natural Rubber Latex Allergy; NRL Allergy

---

## Lathyrism

A neurologic disorder caused by prolonged or excessive ingestion of certain *Lathyrus* legumes, especially grass pea, characterized primarily by progressive spastic weakness of the legs.

## Latissimus Dorsi

A broad back muscle that extends and internally rotates the arm and assists movements such as climbing.

## Latissimus Dorsi Muscle

A broad muscle covering much of the back that pulls the arm down, backward, and toward the body. It is used in climbing, swimming, and rowing and may also be used to reconstruct tissue during surgery.

## Lattice Degeneration

A peripheral retinal degeneration characterized by elongated areas of retinal thinning, pigment changes, and abnormal vessels. It can increase the risk of retinal tears or detachment in some people.

## Lavage

Washing a body space, organ, or wound with sterile fluid. It may be used to remove blood, infected material, or harmful substances, or to collect a sample; the method and risks depend on the part of the body treated.

## Lavage Fluid

Liquid used to wash a body space, wound, or organ during a medical procedure, or collected afterward for laboratory analysis. Its contents can help detect infection, bleeding, cancer cells, inflammation, or abnormal proteins.

## Lawton IADL Scale

A tool measuring ability to manage complex daily tasks such as telephone use, shopping, cooking, housekeeping, transportation, medicines, and finances.

## Laxative Overdose

Taking too much laxative can cause severe diarrhea,

vomiting, dehydration, and abnormal body salts. Some products can disturb heart rhythms or damage the bowel, especially after large exposures or in children.

## Laxatives For Constipation

Laxatives relieve constipation through fiber and water retention, stool softening, osmotic effects, or increased bowel contractions. The suitable type depends on the cause, duration, age, medicines, kidney function, and risk of blockage; persistent bleeding, vomiting, or severe pain may indicate another condition.

### Alternate Names:

Aperients; Cathartics; Purgatives

## Lazy Eye

Reduced vision caused by abnormal visual development during childhood, usually in one eye but sometimes in both. It may result from eye misalignment, unequal focusing, or an obstruction such as a cataract; the medical term is amblyopia.

### Alternate Names:

Amblyopia; Functional Amblyopia

## LCIS

Abbreviation; see *Lobular Carcinoma In Situ*.

## LDH Isoenzyme Blood Test

A blood test separating lactate dehydrogenase isoenzymes to help suggest the tissue source of an elevated total LDH, although more specific tests are preferred for most conditions.

## LDL

Low-density lipoprotein, a blood particle that carries cholesterol to tissues. Too much LDL can contribute to fatty buildup in artery walls and increase the risk of heart attack and stroke; risk-based treatment may lower it.

## LDL Test

A blood test measuring low-density lipoprotein cholesterol, a major atherogenic lipoprotein used with the lipid profile to estimate cardiovascular risk

---

and guide lipid-lowering treatment.

### **L-Dopa**

A medicine that the brain converts into dopamine. It is usually combined with carbidopa or benserazide to treat Parkinson disease.

### **Lead**

A toxic heavy metal that can harm the nervous system, blood-forming system, kidneys, cardiovascular system, and reproductive system. Exposure may occur through contaminated paint dust or soil, drinking water, workplaces, hobbies, imported products, or other sources; children and fetuses are especially vulnerable to neurodevelopmental effects.

### **Lead And Tap Water**

Drinking water can become contaminated with lead when it contacts lead-containing service lines, plumbing, fixtures, or solder. Risk reduction may require testing, use of an appropriately certified filter, flushing or other interim measures, and repair or replacement of the source according to local public-health guidance; boiling water does not remove lead.

### **Leadless Pacemaker**

A small pacemaker placed directly inside a heart chamber without a conventional chest pocket or transvenous lead. It provides pacing for selected slow or unreliable heart rhythms; device selection depends on the person's cardiac anatomy and pacing needs.

### **Lead Pipe Rigidity**

Constant stiffness that gives the same resistance throughout a joint's movement. It may occur in severe movement disorders, advanced Parkinson disease, or neuroleptic malignant syndrome.

#### **Alternate Names:**

Plastic Rigidity; Extrapyramidal Lead-Pipe Hyper-tonia

### **Lead-Pipe Rigidity**

Smooth, continuous muscle stiffness where an arm or leg resists movement evenly in every direction, like bending a lead pipe.

### **Lead Poisoning**

Harm caused by exposure to lead, which can damage the brain, nerves, blood, kidneys, and reproductive system. Children may develop learning, behavior, growth, or developmental problems. Sources include old paint and dust, contaminated soil or water, workplace or hobby materials, and some imported products. Diagnosis uses blood lead testing; treatment starts by stopping exposure, with chelation for selected severe cases.

### **Lead-Time Bias**

The false appearance that screening improves survival because it finds a disease earlier, even though the person dies at the same time as without earlier diagnosis.

### **Lead Toxicity and Nutrition**

Nutritional strategies and dietary interventions aimed at minimizing intestinal absorption of ingested lead, particularly in young children. Adequate dietary intake of iron, calcium, zinc, and vitamin C competitively inhibits divalent metal transporter 1 (DMT1) uptake of lead.

#### **Alternate Names:**

Nutritional Lead Protection; Dietary Prevention of Lead Absorption

### **Learned Insomnia**

Ongoing difficulty sleeping maintained by worry, sleep-related habits, and learned associations that make the bed or bedtime trigger alertness instead of sleep.

### **Learning Difficulties**

Persistent problems acquiring or using academic skills such as reading, writing, or mathematics, not explained solely by low intelligence or inadequate education.

### **Learning Disabilities**

Persistent difficulties acquiring or using skills such as reading, writing, mathematics, language, attention, or coordination despite appropriate opportunity and instruction; assessment considers development, education, and other neurologic or sensory factors.



---

## Learning Disorder

A neurodevelopmental disorder causing persistent difficulty acquiring or using academic skills such as reading, writing, or mathematics despite appropriate instruction.

## Leber Congenital Amaurosis

A rare inherited disorder of the retina that causes severe vision loss from birth or early infancy. Babies may have wandering eye movements, sensitivity to light, or unusual responses of the pupils; genetic testing can help confirm the cause.

## Leber Hereditary Optic Neuropathy

An inherited disorder of the energy-producing parts of cells that damages the optic nerves. It usually causes painless, rapidly worsening loss of central vision in one eye followed by the other, often in young adults. The altered mitochondrial DNA is passed through the mother; genetic testing can confirm the diagnosis.

### Alternate Names:

Lhon

## Leber's Hereditary Optic Neuropathy

An inherited mitochondrial disorder that causes painless loss of central vision, usually first in one eye and then the other. It mainly affects young men, and genetic counseling and specialist eye care may help.

## Lecithin-Sphingomyelin Ratio

A laboratory ratio measured on amniotic fluid obtained via amniocentesis to evaluate fetal pulmonary maturity and the risk of neonatal respiratory distress syndrome. Lecithin (dipalmitoylphosphatidylcholine) concentrations rise sharply relative to constant sphingomyelin levels after 35 weeks of gestation; an L/S ratio of 2.0 or greater indicates adequate pulmonary surfactant production.

### Alternate Names:

L/S Ratio; Amniotic Fluid Lecithin-Sphingomyelin Ratio

## Leech

A blood-feeding annelid used in selected reconstruc-

tive procedures to relieve venous congestion; leech bites can cause bleeding and infection.

## Leech Therapy

Carefully controlled use of medicinal leeches to remove congested blood from selected tissue, especially after reconstructive surgery. Their saliva also prevents clotting. Bleeding, infection, anemia, and allergic reactions are possible, so clinical monitoring is essential.

## Leflunomide

Leflunomide is a disease-modifying antirheumatic drug that inhibits dihydroorotate dehydrogenase and lymphocyte proliferation; it treats rheumatoid or psoriatic arthritis but can cause hepatotoxicity, cytopenias, hypertension, and fetal harm.

## Le Fort Fractures

A classical orthopedic and maxillofacial classification of traumatic midface fractures based on René Le Fort's lines of structural facial weakness. Subdivided into Le Fort I (horizontal maxillary fracture separating the dental arch from the upper midface), Le Fort II (pyramidal fracture extending through the nasal bones, ethmoid, and maxillary sinus walls), and Le Fort III (complete craniofacial disjunction separating the facial skeleton from the cranial base).

### Alternate Names:

Le Fort Midface Fractures; Maxillofacial Le Fort System

## Left Anterior Descending Artery

A major branch of the left coronary artery that travels in the anterior interventricular groove and supplies the anterior left ventricle, much of the interventricular septum, and the apex. Its occlusion can cause extensive anterior myocardial infarction.

## Left Anterior Descending Coronary Artery

A major heart artery supplying the front wall, septum, and much of the left ventricle; blockage can cause a serious heart attack.

### Alternate Names:

LAD Artery

---

### Left Anterior Fascicular Block

A cardiac intraventricular conduction defect caused by delay or block in the thin anterior fascicle of the left bundle branch, manifesting on electrocardiogram as marked left axis deviation (between -45 and -90 degrees), small Q waves in leads I and aVL, and small R waves in leads II, III, and aVF.

#### Alternate Names:

LAFB; Left Anterior Hemiblock

### Left Atrial Appendage Occlusion

A procedure that seals a small pouch of the heart where blood clots often form in people with atrial fibrillation. A closure device is placed through a catheter, usually from a vein in the groin. It may reduce stroke risk when long-term blood-thinning medicine is unsuitable, but short-term medication and follow-up are still needed.

#### Alternate Names:

LAAO; Left Atrial Appendage Closure; Watchman Procedure

### Left Atrial Myxoma

A usually benign primary cardiac tumor, commonly arising from the interatrial septum in the left atrium. It may obstruct mitral inflow or cause embolic and constitutional symptoms such as fever, weight loss, or fatigue.

### Left Atrium

The upper-left chamber of the heart. It receives oxygen-rich blood from the lungs and passes it through the mitral valve into the left ventricle.

### Left Bundle Branch

Part of the heart's electrical wiring that carries signals through the left lower chamber and helps it contract. Damage or blockage can delay the heart-beat signal and appear on an electrocardiogram.

### Left Bundle Branch Block

Delay or interruption of electrical signals through the left bundle branch. It changes the ECG pattern and may reflect heart disease, although some people have no symptoms.

#### Alternate Names:

LBBB

### Left Circumflex Coronary Artery

A heart artery curving around the left side and back, supplying parts of the left ventricle and sometimes the lower heart wall.

#### Alternate Names:

Circumflex Artery

### Left Heart Catheterization

A procedure that passes a thin tube into the left side of the heart or nearby arteries to measure pressures, assess valves, or look for blocked coronary arteries.

### Left Heart Ventricular Angiography

An imaging study that injects contrast into the left ventricle to assess ventricular contraction, wall motion, ejection fraction, and selected structural abnormalities.

### Left Hypochondriac Region

The upper-left area of the abdomen, beneath the lower ribs. It overlies part of the stomach, spleen, pancreas, and colon; pain or swelling there may arise from digestive, splenic, or nearby chest disease.

### Left Iliac Region

The lower-left area of the abdomen, overlying part of the large intestine and urinary tract. Pain there may occur with diverticular disease, bowel inflammation, urinary problems, or other pelvic conditions.

### Left Main Coronary Artery

The short vessel arising from the left aortic sinus and dividing principally into the left anterior descending and circumflex arteries. Significant left-main disease jeopardizes a large proportion of left-ventricular myocardium.

### Left-Sided Superior Vena Cava

A birth-related variation in which a large vein on the left side of the chest drains blood through an unusual route to the heart. It usually causes no symptoms but matters when placing central lines

---

or performing heart surgery.

### **Left Testicular Torsion**

Acute rotation of the left testis on its spermatic cord, obstructing blood flow and causing ischemia. It is a urological emergency presenting with sudden severe left scrotal pain, nausea, vomiting, and a high-riding, horizontally oriented testis with absent cremasteric reflex.

### **Left Ventricle**

The lower-left chamber of the heart. It receives oxygen-rich blood from the left atrium and pumps it through the aortic valve into the aorta and throughout the body.

### **Left Ventricular Assist Device**

A pump placed in or near the heart to help the left side pump blood in advanced heart failure. A cable connects the pump to a controller and batteries outside the body. It may support someone while waiting for a transplant or provide long-term support when a transplant is not suitable. It can cause bleeding, infection, clots, or stroke and needs close follow-up.

#### **Alternate Names:**

LVAD; Left Ventricular Assist System; Mechanical Circulatory Support; Ventricular Assist Device

### **Left-Ventricular Assist Device**

An implanted mechanical pump that helps the heart's main pumping chamber send blood around the body. It may support people with severe heart failure while they await a transplant or as long-term treatment; infection, bleeding, and blood clots are important risks.

### **Left Ventricular Hypertrophy**

Thickening of the muscle wall of the heart's main pumping chamber. It is often caused by long-term high blood pressure or a narrowed aortic valve and can make the heart stiff, leading to breathlessness, rhythm problems, or heart failure.

### **Left Ventricular Non-Compaction**

A genetic cardiomyopathy resulting from the ar-

rest of normal embryonic myocardial compaction, characterized by prominent ventricular trabeculations and deep intertrabecular recesses ("spongy myocardium"), predisposing to heart failure, systemic thromboembolism, and ventricular arrhythmias.

#### **Alternate Names:**

LVNC; Spongy Myocardium

### **Left Ventricular Noncompaction**

A rare heart-muscle disorder in which the main pumping chamber has an unusually spongy inner muscle layer. It may cause no symptoms or lead to heart failure, abnormal rhythms, blood clots, or stroke.

### **Leg**

The lower limb, often meaning the region between the knee and ankle, although it may also refer broadly to the entire lower extremity.

### **Leg CT Scan**

A series of detailed X-ray pictures of the leg combined by a computer. It can show complex fractures, bone alignment, infection, tumors, blood-vessel problems, and selected soft-tissue injuries.

### **Leg Fracture**

Alternate name; see *Broken Leg*.

### **Legg-Calve-Perthes Disease**

A childhood hip disorder in which the blood supply to the top of the thighbone is temporarily reduced. The bone can weaken and change shape, causing a limp, hip or knee pain, stiffness, and limited movement.

### **Legg-Calvé-Perthes Disease**

A childhood hip disorder in which reduced blood flow temporarily damages the rounded top of the thighbone. It can cause a limp, hip or knee pain, stiffness, and later deformity as the bone heals.

### **Legionella**

Legionella is a group of water-associated bacteria that can cause Legionnaires' disease, a severe pneu-

---

monia, or Pontiac fever, a milder illness.

### **Legionella Longbeachae**

*Legionella longbeachae* is a *Legionella* species associated particularly with exposure to contaminated potting mix and capable of causing pneumonia.

### **Legionella Pneumonia**

Pneumonia caused by *Legionella* bacteria, usually after inhaling contaminated water droplets. It can cause high fever, cough, breathlessness, diarrhea, confusion, and low oxygen.

### **Legionella Pneumophila**

A bacterium found in water systems that can cause severe pneumonia after a person inhales contaminated water droplets. It may also cause a milder fever illness; older age, smoking, and weakened immunity increase risk.

### **Legionellosis**

An infection caused by *Legionella* bacteria, ranging from a mild flu-like illness to Legionnaires' disease with pneumonia.

### **Legionnaire Disease And Pontiac Fever**

Legionnaires disease is a severe pneumonia caused by *Legionella*, while Pontiac fever is a milder flu-like illness from the same bacteria. Exposure to contaminated water systems is typical; pneumonia, confusion, low oxygen, or high fever may occur in the more serious illness.

#### **Alternate Names:**

Legionellosis; *Legionella* Infection

### **Legionnaires' Disease**

A severe form of pneumonia caused by *Legionella pneumophila*, a Gram-negative aerobic bacillus that is found naturally in freshwater environments (lakes, streams) and artificially in water systems (air conditioning cooling towers, hot tubs, decorative fountains, plumbing systems, showers, humidifiers).

### **Legionnaires Disease**

A severe pneumonia caused by *Legionella*, usually

acquired by inhaling contaminated water aerosols or aspiration. Fever, cough, breathlessness, diarrhea, confusion, and low sodium may occur; laboratory testing and prompt active treatment are important because routine beta-lactams do not cover *Legionella* reliably.

### **Leg Lengthening And Shortening**

Operations that gradually increase or decrease the length of a leg bone to correct a major difference between the legs or a deformity. Recovery is prolonged and may involve an external or internal support device and rehabilitation.

### **Leg MRI Scan**

Magnetic resonance imaging of the leg using a magnetic field and radio waves to evaluate muscles, tendons, ligaments, nerves, marrow, joints, and soft-tissue masses.

### **Leg Or Foot Amputation**

Surgical removal of part or all of a leg or foot, usually because of severe injury, infection, poor blood flow, or cancer. Rehabilitation supports mobility afterward.

### **Leg Pain**

Pain in the lower limb caused by muscle or tendon injury, joint disease, nerve problems, vascular disease, infection, or other conditions.

### **Leg Pain With Cramping**

Painful tightening or spasms in a leg muscle; causes include exertion, dehydration, medicines, nerve or muscle disorders, and reduced arterial blood flow causing intermittent claudication.

### **Leg Swelling**

Enlargement or puffiness of one or both legs caused by fluid retention, venous disease, injury, infection, inflammation, or a blood clot.

### **Leg Ulcer**

An open sore on the lower leg that heals slowly or not at all. Causes include poor vein or artery

---

circulation, diabetes-related nerve damage, pressure, and inflammation. Venous ulcers often occur near the ankle and may be accompanied by leg swelling or varicose veins; the cause should be assessed because treatment depends on it.

### **Leigh Disease**

A progressive mitochondrial disorder of early life characterized by symmetric lesions of the brainstem and basal ganglia, developmental regression, movement abnormalities, and lactic acidosis.

### **Leigh's Syndrome**

A rare progressive neurologic disorder, usually beginning in infancy or childhood, caused by problems with cellular energy production. It can damage the brainstem and cause developmental regression, movement problems, seizures, or breathing difficulties.

### **Leigh Syndrome**

A rare inherited disorder of cellular energy production, usually beginning in infancy or childhood. It can damage the brainstem and deep brain areas, causing loss of developmental skills, movement problems, seizures, and breathing difficulties.

#### **Alternate Names:**

Subacute Necrotizing Encephalomyelopathy; Mitochondrial Leigh Disease

### **Leiomyoma**

A usually noncancerous tumor made of smooth muscle. It most often develops in the uterus, where it is called a fibroid, but it can also occur in the skin, digestive tract, or other organs.

### **Leiomyomatosis**

Leiomyomatosis is development of multiple benign smooth-muscle tumors or proliferations, which may affect the uterus, skin, lungs, or other sites.

### **Leiomyomatosis Peritonealis Disseminata**

A rare benign condition with multiple smooth-muscle nodules scattered over the peritoneum, often associated with pregnancy or hormonal stimulation and sometimes mimicking peritoneal carcinomato-

sis.

### **Leiomyosarcoma**

A malignant tumor of smooth muscle, most often arising in the uterus, retroperitoneum, or gastrointestinal tract.

### **Leiomyosarcoma Of The Vulva**

A rare malignant smooth-muscle tumor of the vulva, usually presenting as a solitary enlarging mass. Histologic atypia, mitotic activity, necrosis, and invasion help determine malignant behavior.

### **Leishmania**

A group of single-celled parasites transmitted mainly by infected sandflies. Different species cause skin, mucosal, or visceral leishmaniasis, ranging from localized ulcers to life-threatening illness.

### **Leishmania Donovanii**

A parasite spread by infected sandflies that causes visceral leishmaniasis. It can infect organs such as the spleen, liver, and bone marrow, causing prolonged fever, weight loss, enlarged organs, and reduced blood-cell counts.

### **Leishmania Infection**

A parasitic infection caused by *Leishmania* protozoa and transmitted by infected sandflies; it can cause cutaneous, mucocutaneous, or visceral leishmaniasis.

### **Leishmania Major**

A parasite commonly causing cutaneous leishmaniasis in parts of Africa, Asia, and the Middle East. Sandfly transmission produces one or more skin ulcers that may heal slowly and leave scars.

### **Leishmania Mexicana**

A parasite that causes mainly cutaneous leishmaniasis in the Americas. Infection after a sandfly bite may produce a persistent skin nodule or ulcer and, less commonly, involve mucous membranes.

---

### **Leishmaniasis**

An infection caused by *Leishmania* parasites transmitted by sandflies. It may produce skin ulcers, damage to the nose or mouth, or a life-threatening illness affecting organs such as the spleen, liver, and bone marrow.

### **Leishmania Tropica**

A sandfly-transmitted parasite that commonly causes localized cutaneous leishmaniasis. It produces chronic skin sores that may heal over months and leave permanent scars.

### **Lemaitre's Sign**

Pain or discomfort felt in the urethra or penis during a rectal examination. It has traditionally been associated with inflammation near the prostate or back part of the urinary passage but is not specific for one diagnosis.

### **Lemierre Syndrome**

A potentially severe oropharyngeal infection complicated by septic thrombophlebitis of the internal jugular vein and frequently metastatic infection, classically caused by *Fusobacterium necrophorum*.

### **Lenalidomide**

Lenalidomide is an immunomodulatory anticancer drug used for multiple myeloma and selected myelodysplastic or lymphoma disorders; myelosuppression, thrombosis, infection, and severe teratogenicity are important risks.

### **Lenegre Disease**

An idiopathic, sclerodegenerative process characterized by slowly progressive fibrocalcific degeneration and loss of conduction fibers within the His-Purkinje system, leading to bundle branch blocks, bifascicular block, and spontaneous complete atrioventricular block in middle-aged individuals.

#### **Alternate Names:**

Lenègre-Lev Disease; Primary Conduction System Sclerosis

### **Lengthening Of Limb**

A surgical or orthopedic process that gradually increases the length of a bone, often using controlled distraction after osteotomy.

### **Length-Time Bias**

The tendency of screening to preferentially detect slower-growing or less aggressive disease because it remains detectable for a longer preclinical period. It can exaggerate the apparent benefit of screening.

### **Lennox-Gastaut Syndrome**

A severe childhood-onset developmental and epileptic encephalopathy with multiple seizure types, characteristic slow spike-and-wave EEG patterns, cognitive impairment, and often treatment-resistant seizures.

### **Lenograstim**

A recombinant granulocyte colony-stimulating factor that increases neutrophil production and is used to prevent or treat some forms of neutropenia.

### **Lens**

A clear, flexible structure behind the iris and pupil that bends light onto the retina. Muscles change its shape for near or distant focus; clouding causes cataract, while displacement can blur vision or raise eye pressure.

### **Lens Coloboma**

A lens coloboma is a congenital segmental defect or notch in the eye lens caused by incomplete development.

### **Lens Disease**

A disorder of the eye lens, including cataract, dislocation, inflammation-related opacity, or developmental abnormality, that may impair focusing or vision.

### **Lens Dislocation**

Displacement of the crystalline lens from its normal position because of trauma, zonular weakness, surgery, or connective-tissue disease; it can cause



---

blurred vision, glaucoma, or inflammation.

### **Lensometer**

An instrument measuring the prescription, power, and optical alignment of eyeglass lenses. It helps verify new glasses, assess existing lenses, and check whether corrective strength matches the prescribed values.

#### **Alternate Names:**

Lensmeter; Focimeter

### **Lens Prescription**

Written instructions specifying the lens power, correction, and other features needed for eyeglasses or contact lenses in each eye.

### **Lens Subluxation**

Partial displacement of the crystalline lens while some zonular attachments remain, potentially causing blurred or fluctuating vision, refractive change, or glaucoma.

### **Lentiform Nucleus**

A group of deep brain structures involved in controlling movement. It includes the putamen and globus pallidus and works with other brain regions to help start and regulate actions.

### **Lentigines**

Alternate plural name; see *Lentigo*.

### **Lentigo**

A lentigo is a sharply defined flat brown skin spot caused by increased melanocytes or pigment and may be benign or, rarely, atypical.

### **Lentigo Maligna**

A melanoma in situ that occurs on chronically sun-damaged skin, most commonly on the face of elderly individuals. It presents as a large, irregularly pigmented, slow-growing macule with variegated shades of brown and black. The radial growth phase may persist for years before invasion occurs.

### **Lentigo Maligna Melanoma**

An invasive melanoma arising in a lentigo maligna on chronically sun-damaged skin, usually of the head or neck; it presents as an enlarging irregular pigmented patch or plaque.

### **LEOPARD Syndrome**

A rare inherited disorder that can cause many dark skin spots, abnormal heart rhythms, widely spaced eyes, narrowing of the pulmonary valve, genital differences, slow growth, and hearing loss. The combination and severity vary between people.

### **Leopold Maneuvers**

A set of gentle abdominal examinations used to estimate the baby's position and which part is presenting before birth. The findings help guide care but may be limited by body size, fluid level, or the baby's movement and are confirmed with other assessment when needed.

### **Lepidoptera**

The insect order containing butterflies and moths; contact with some caterpillars can cause skin, eye, or systemic reactions.

### **Lepirudin**

Lepirudin is a direct thrombin inhibitor formerly used as an anticoagulant in heparin-induced thrombocytopenia; bleeding and kidney clearance are important considerations.

### **Lepromatous Leprosy**

A form of leprosy with a high number of bacteria and usually widespread skin involvement. It can cause numerous poorly defined patches, nodules, or diffuse thickening, including on the face, and may affect the nose, eyes, and peripheral nerves. Sensation may be preserved early, but untreated nerve damage can lead to numbness, weakness, and disability.

### **Lepromin Skin Test**

A skin test using harmless material from the leprosy bacterium to show how strongly the immune system reacts. It may help classify the type of leprosy but

---

cannot by itself diagnose active infection.

### **Leprosy**

A long-term infection caused mainly by bacteria that affect the skin and cooler nerves in the arms, legs, and face. It can cause pale or reddish patches with reduced feeling, thickened nerves, weakness, and eye or nose problems. A skin sample or other tests may confirm it. Treatment uses several antibiotics, commonly rifampicin, dapsone, and clofazimine, to prevent lasting nerve damage and disability.

### **Leptin**

A hormone produced mainly by adipose tissue that signals stored-energy status to the brain and influences appetite and energy balance.

### **Leptin Receptor**

A cell-surface protein that receives signals from leptin, a hormone made mainly by fat tissue. It helps the brain regulate appetite, energy use, growth, and reproductive function in response to stored body energy.

### **Leptomeningeal**

Relating to the pia mater and arachnoid mater, the two thin inner layers covering the brain and spinal cord; cancer or infection can spread in these layers.

### **Leptomeningeal Metastases**

Spread of cancer cells to the pia and arachnoid membranes or cerebrospinal fluid. It may cause headache, nausea, cranial-nerve findings, radicular pain, weakness, sensory change, or bladder dysfunction; MRI and cerebrospinal-fluid testing support diagnosis.

### **Leptomeningeal Metastasis**

Spread of cancer to the thin membranes and fluid surrounding the brain and spinal cord. It can cause headache, vomiting, weakness, confusion, seizures, or changes in vision and is treated according to the cancer type and extent.

### **Leptomeningeal Sarcoma**

Leptomeningeal sarcoma is a rare malignant tumor involving the membranes and fluid spaces around the brain or spinal cord.

### **Leptomeninges**

The arachnoid mater and pia mater, the two delicate inner membranes covering the brain and spinal cord.

### **Leptomeningitis**

Inflammation or infection of the leptomeninges, the arachnoid and pia mater covering the brain and spinal cord.

### **Leptospira**

A genus of spiral-shaped bacteria that includes the causes of leptospirosis, usually acquired through water or soil contaminated with infected animal urine.

### **Leptospirosis**

A bacterial infection caused by spiral-shaped *Leptospira* bacteria, usually spread when cut skin or eyes touch water, mud, or soil contaminated with the urine of infected animals, especially rats and farm animals. Symptoms often begin suddenly with fever, severe headache, intense calf and lower back muscle pain, and characteristic red eyes without discharge (conjunctival suffusion). A severe form, known as Weil's disease, can develop rapidly, causing yellow skin and eyes (jaundice), kidney failure, severe bleeding, and lung hemorrhage. Diagnosis is made through blood tests, antibody tests, or PCR. Mild cases are treated with oral antibiotics like doxycycline, while severe cases require intravenous penicillin or ceftriaxone in a hospital.

#### **Alternate Names:**

Canefield Fever; Canicola Fever; Seven-Day Fever; Weil's Disease

#### **Alternate Names:**

Weil's Disease; Leptospira Infection; Mud Fever; Canicola Fever

### **Leriche Syndrome**

Aortoiliac occlusive disease causing reduced blood flow to the pelvis and lower limbs. The classic pre-

---

sentation includes buttock or thigh claudication, diminished femoral pulses, and erectile dysfunction, although symptoms vary with the extent of collateral circulation.

### **Leri-Weill Dyschondrosteosis**

A rare autosomal dominant pseudoautosomal skeletal dysplasia caused by haploinsufficiency of the short stature homeobox (SHOX) gene on the X and Y chromosomes, characterized by disproportionate mesomelic short stature, shortened forearms and lower legs, and bilateral Madelung wrist deformity.

#### **Alternate Names:**

Leri-Weill Syndrome; SHOX Deficiency Mesomelic Dysplasia

### **Lesch-Nyhan Syndrome**

An inherited disorder of purine metabolism caused by severe HPRT enzyme deficiency, leading to high uric acid, neurologic problems, and characteristic self-injurious behavior.

### **Lesion**

Any localized abnormal change in tissue caused by disease, injury, inflammation, infection, or a tumor. A lesion may be visible, felt, or found only on imaging or microscopic examination.

### **Lesion Identification**

The process of locating and describing an abnormal area in the body using examination, imaging, endoscopy, laboratory tests, or tissue sampling.

### **Lesser Curvature**

The shorter, inward curve along the upper and inner edge of the stomach. It is attached to a fold of tissue leading to the liver and contains blood vessels that supply the stomach.

### **Lesser Omentum**

A fold of tissue that connects the liver to the lesser curvature of the stomach and the first part of the small intestine. It contains blood vessels, ducts, nerves, and lymph vessels.

### **Lesser Peritoneal Cavity**

The peritoneal recess behind the stomach and lesser omentum, also called the lesser sac or omental bursa. It communicates with the greater peritoneal cavity through the epiploic foramen.

#### **Alternate Names:**

Lesser Sac; Omental Bursa

### **Lesser Trochanter**

A small bony projection on the inner upper thigh-bone where the iliopsoas muscle attaches. It helps bend the hip; injury may occur from a sudden muscle pull in adolescents or, in adults, from weakened bone.

### **Let-Down Reflex**

Release of breast milk when oxytocin makes the small muscles around milk-producing glands contract. Suckling, breast stimulation, or seeing or hearing a baby can trigger it, although stress may delay it.

### **Lethal**

Able to cause death. In medicine, the word may describe a disease, injury, dose of a substance, or genetic change that prevents survival.

### **Lethal-Means Counseling**

A preventive intervention that reduces access to firearms, medications, toxic substances, or other methods of self-harm during periods of elevated risk. It is conducted collaboratively and is an important component of suicide prevention.

### **Lethal Triad**

A dangerous combination of low body temperature, excess acid in the blood, and poor blood clotting. It can develop after severe injury or major bleeding, and each problem can worsen the others.

### **Lethargy**

Lethargy is reduced energy, alertness, or responsiveness; it may result from sleep loss, infection, metabolic disease, medicines, depression, or neurologic illness. Sudden marked lethargy can occur

---

with serious illness.

### **Letrozole**

Letrozole is an aromatase inhibitor that lowers oestrogen production and is used mainly for hormone-receptor-positive breast cancer and ovulation induction; hot flushes, arthralgia, bone loss, and fetal harm are important considerations.

### **Letterer-Siwe Disease**

An acute, disseminated, aggressive multisystem form of Langerhans cell histiocytosis occurring in infants and toddlers under 2 years of age, characterized by cutaneous seborrheic and purpuric eruptions, osteolytic bone lesions, hepatosplenomegaly, lymphadenopathy, and pancytopenia.

#### **Alternate Names:**

Acute Disseminated LCH; Multisystem Langerhans Histiocytosis

### **Leucocyte**

Another spelling of leukocyte, meaning a white blood cell. Leukocytes help defend against infection and other threats; major groups include neutrophils, lymphocytes, monocytes, eosinophils, and basophils.

### **Leucocytosis**

A higher-than-usual number of white blood cells in the blood. It may occur with infection, inflammation, physical stress, medicines, smoking, or blood and bone-marrow disorders, so the type of cell and the person's symptoms help find the cause.

### **Leucoderma**

Loss or marked reduction of skin pigmentation, often used as a descriptive term for depigmented patches.

### **Leucomalacia**

Softening or loss of tissue after injury, poor blood flow, infection, or inflammation. In newborns, the term often refers to damage of the brain's white matter near the fluid spaces, which may affect movement or development.

### **Leucopenia**

A lower-than-usual number of white blood cells. It may result from medicines, infections, immune disorders, nutritional deficiency, or bone-marrow disease and can increase the risk of infection, especially when neutrophils are low.

### **Leucoplakia**

A persistent white patch on a moist body lining, often inside the mouth, that cannot be wiped away. It may result from irritation and sometimes contains changes that can precede cancer.

### **Leucopoiesis**

The production and maturation of white blood cells, mainly in the bone marrow. It is regulated by growth factors and increases or changes during infection, inflammation, and some blood disorders.

### **Leucovorin**

Leucovorin is a folate medicine used to reduce the toxicity of methotrexate, enhance some chemotherapy regimens, and treat certain folate-related problems.

### **Leucovorin Calcium**

Leucovorin calcium is the calcium salt of leucovorin, used as rescue therapy after methotrexate and in selected colorectal-cancer treatments.

### **Leukaemia**

Alternate spelling; see *Leukemia*.

### **Leukapheresis**

A blood-filtering procedure that removes excess white blood cells and returns the remaining blood. It may rapidly reduce very high counts causing symptoms in leukemia, but it is temporary and does not replace treatment of the underlying disease.

### **Leukemia**

Cancer that starts in blood-forming tissue, usually the bone marrow, and produces large numbers of abnormal blood cells. These cells can crowd out healthy white blood cells, red blood cells, and

---

platelets, making it harder to fight infection, carry oxygen, and stop bleeding. This may cause tiredness, repeated infections, easy bruising, or bleeding. Leukemia usually does not form a solid tumor. It is grouped as acute or chronic according to how quickly it progresses and as lymphoid or myeloid according to the type of blood-forming cell involved.

**Alternate Names:**

Hematologic Malignancy; Blood Cancer

**Leukemia Cutis**

A condition in which leukemia cells collect in the skin, causing firm red, purple, or brown bumps, nodules, or patches. It often accompanies acute myeloid leukemia and reflects the underlying blood cancer.

**Leukemia Genetics**

Changes in genes or chromosomes in leukemia cells help identify the leukemia type, estimate its likely course, and choose treatment. The meaning of each change depends on the person's specific leukemia.

**Leukemic Infiltration Of The Gingiva**

Swelling of the gums caused by leukemia cells collecting in gum tissue. The gums may become red, soft, enlarged, and prone to bleeding, especially in some acute myeloid leukemias.

**Leukemoid Reaction**

A marked reactive increase in the white-cell count, usually with prominent neutrophilia and immature granulocytes, caused by severe infection, inflammation, malignancy, or tissue injury. It resembles leukemia but is not a clonal hematologic malignancy.

**Leukoaraiosis**

White-matter changes seen on brain scans, often related to aging and long-term small-vessel disease such as high blood pressure. Extensive changes may be associated with slower thinking, walking problems, or stroke risk.

**Leukocoria**

A white or pale reflection appearing in the pupil instead of the normal red reflection. It may result

from cataract, retinal disease, infection, or a tumor such as retinoblastoma and is especially important in infants and children.

**Leukocyte**

A white blood cell that helps defend the body against infection and other threats. Main types include neutrophils, lymphocytes, monocytes, eosinophils, and basophils; unusually high or low numbers can reflect infection, inflammation, medicines, or bone-marrow disease.

**Leukocyte Adhesion Deficiency**

A rare inherited immune disorder in which white blood cells cannot attach to blood-vessel walls and reach infected tissue normally. It can cause delayed umbilical-cord separation, repeated skin infections with little pus, poor wound healing, gum disease, and dangerously severe infection in infants.

**Leukocyte Alkaline Phosphatase**

A measure of an enzyme in certain white blood cells. It was formerly used to help distinguish a reactive response to infection from chronic myeloid leukemia, but newer genetic and blood tests are usually more informative.

**Leukocyte Disorder**

A disorder involving the number, function, or development of white blood cells, including leukopenia, leukocytosis, immune deficiency, and malignant leukocyte proliferation.

**Leukocyte Esterase Urine Test**

A urine dipstick test detecting leukocyte esterase released by white blood cells; a positive result supports pyuria but does not by itself diagnose urinary tract infection.

**Leukocytes**

White blood cells that defend the body against infections, abnormal cells, and other threats. Major types include neutrophils, lymphocytes, monocytes, eosinophils, and basophils, each with different immune functions.

---

## Leukocyte Transfusion

Transfusion of donated white blood cells in rare, carefully selected situations involving life-threatening infection and severe inability to produce neutrophils. Benefits are uncertain and risks include fever, lung injury, immune reactions, and infection transmission.

## Leukocytoclastic Vasculitis

A pattern of inflammation in small blood vessels, usually identified by examining a skin biopsy. It commonly causes raised purple or red spots that can be felt, especially on the legs. Medicines, infections, and immune diseases may trigger it; evaluation checks whether inflammation is limited to the skin or also affects internal organs.

### Alternate Names:

Cutaneous Small-Vessel Vasculitis

## Leukocytosis

A higher-than-usual number of white blood cells. It may occur with infection, inflammation, stress, smoking, medicines, or blood and bone-marrow disorders; the specific cell type and symptoms help identify the cause.

## Leukocytosis in Pregnancy

A mild rise in white blood cells can be normal during pregnancy, labor, or stress. Marked or persistent elevation with fever, pain, urinary or respiratory symptoms, or abnormal differential counts may indicate infection, inflammation, medication effect, or another disorder.

## Leukoderma

Loss of skin pigment producing sharply defined pale or white patches, usually because melanocytes have been destroyed or have stopped functioning.

### Alternate Names:

Hypopigmentation; Achromoderma

## Leukodystrophy

A group of inherited disorders that damage myelin, the protective covering around nerves in the brain and spinal cord. They can cause progressive problems with movement, thinking, vision, hearing, or

other nervous-system functions.

## Leukoencephalopathy

Leukoencephalopathy is disease of the brain's white matter caused by inherited, infectious, toxic, vascular, metabolic, inflammatory, or other processes.

## Leukonychia

White areas or bands in one or more nails. Causes include minor injury to the nail, inherited traits, infection, medicines, or illness; the pattern and whether it grows out with the nail help determine the cause.

## Leukonychia Striata

Longitudinal white bands running along the nail plate, usually caused by changes in the nail matrix. They may be inherited or follow repeated minor trauma, skin disease, or some medicines; persistent or widespread changes can be linked to other illness.

## Leukonychia Totalis

Uniform white discoloration involving most or all of one or more nail plates. It is often inherited but can occasionally be associated with illness or medication; persistent changes affecting many nails can occur with other conditions.

## Leukopenia

An abnormally low white-blood-cell count, which may increase susceptibility to infection depending on the cells affected and the severity of the reduction.

## Leukoplakia

A persistent white patch or plaque in the mouth that cannot be rubbed off and cannot be characterized as another condition. It is a clinical description rather than a specific tissue diagnosis; some lesions contain precancerous changes or may progress to cancer, so assessment and often biopsy are needed.

### Alternate Names:

Smokers Keratosis (leukoplakia)

## Leukopoiesis

The process by which the bone marrow makes and develops white blood cells. Production increases



---

or changes during infection, inflammation, immune disorders, and some blood diseases.

### **Leukorrhea**

Whitish vaginal discharge that may be normal, especially during hormonal changes, or may result from infection or inflammation. Foul odor, itching, pain, bleeding, or a new change may suggest an underlying condition.

### **Leukostasis**

A medical emergency in which very high numbers of abnormal leukocytes impair microvascular blood flow, potentially causing respiratory distress, neurologic symptoms, organ ischemia, or hemorrhage.

### **Leukotriene**

A leukotriene is an inflammatory lipid mediator made from arachidonic acid that contributes to bronchoconstriction, mucus, vascular leak, and immune-cell recruitment.

### **Leukotriene Blockers**

Medicines that reduce the effects or production of leukotrienes, substances involved in inflammation and airway narrowing. They help prevent some asthma or allergy symptoms but do not usually relieve a sudden asthma attack.

#### **Alternate Names:**

Leukotriene Antagonists

### **Leukotriene Modifiers**

Medicines that reduce the production or effects of leukotrienes, inflammatory chemicals that narrow airways and cause allergy symptoms. They help prevent some asthma or allergy problems but do not usually relieve a sudden asthma attack.

### **Leukotriene Receptor**

A cell-surface protein activated by leukotrienes, inflammatory chemicals that can narrow airways, increase mucus, and promote swelling. Blocking these receptors can help prevent selected asthma and allergy symptoms.

### **Leukotrienes**

Chemical signals released during inflammation that can narrow airways, increase mucus, and cause swelling. They contribute to asthma, allergies, and some other inflammatory conditions.

### **Leuprolide**

Leuprolide is a gonadotropin-releasing hormone agonist used to suppress sex-hormone production in conditions such as prostate cancer, endometriosis, and fibroids.

### **Leuprolide Acetate**

Leuprolide acetate is the salt form of leuprolide, used to suppress ovarian or testicular hormone production for selected medical conditions.

### **Levalbuterol**

An inhaled short-acting bronchodilator that stimulates beta-2 receptors and relaxes airway muscle. It relieves wheezing and shortness of breath but may cause tremor, nervousness, headache, or a fast heartbeat.

### **Levamisole**

Levamisole is an antiparasitic and immunomodulating drug used selectively; serious low white-cell counts and vasculitic reactions can occur.

### **Levator Ani**

A group of pelvic-floor muscles that supports the bladder, bowel, and reproductive organs and helps control urination and bowel movements. Weakness or injury may contribute to pelvic organ prolapse or incontinence.

### **Levator Ani Muscle**

A broad, symmetrical muscular sheet forming the major constituent of the pelvic diaphragm, composed of the puborectalis, pubococcygeus, and iliococcygeus muscles, which supports the pelvic viscera and maintains urinary and fecal continence.

### **Levator Ani Syndrome**

A chronic functional anorectal pain disorder charac-

---

terized by episodic or continuous dull aching, fullness, or pressure sensation in the rectum and perineum that is aggravated by prolonged sitting, associated with tenderness and spasm of the levator ani and puborectalis muscle on digital rectal palpation.

**Alternate Names:**

Chronic Proctalgia; Levator Spasm Syndrome

**Lev Disease**

An acquired degenerative disorder of the cardiac conduction system characterized by progressive idiopathic sclerosis and fibrocalcification of the fibrous cardiac skeleton, atrioventricular node, and proximal bundle branches, leading to high-grade atrioventricular block and syncope in elderly patients.

**Alternate Names:**

Lenegre-Lev Disease; Idiopathic Conduction System Sclerosis

**Levetiracetam**

Levetiracetam is a broad-spectrum antiseizure medicine that binds synaptic vesicle protein SV2A; it treats focal and generalized seizures and may cause somnolence, dizziness, or behavioral and mood changes.

**Levine Sign**

A physical examination finding in which a patient instinctively places a clenched fist over the sternum when describing ischemic chest pain, serving as a classic nonverbal clinical indicator of angina pectoris or acute myocardial infarction.

**Alternate Names:**

Levine's Sign; Clenched Fist Chest Pain Sign

**Levobunolol**

Levobunolol is a beta-blocker eye drop used to lower pressure inside the eye in glaucoma or ocular hypertension.

**Levocardia**

Levocardia is a normal left-sided position of the heart; it may occur with normal organ arrangement or with other congenital laterality abnormalities.

**Levocarnitine**

Levocarnitine is the biologically active form of carnitine used to treat selected carnitine deficiencies and help transport fatty acids into mitochondria.

**Levodopa**

Levodopa is a dopamine precursor that crosses the blood-brain barrier and is converted to dopamine, usually with carbidopa or benserazide, to treat Parkinson disease; dyskinesias, hallucinations, nausea, and motor fluctuations can develop.

**Levofloxacin**

Levofloxacin is a fluoroquinolone antibiotic that inhibits bacterial DNA gyrase and topoisomerase IV; tendon injury, peripheral neuropathy, QT prolongation, dysglycaemia, and central-nervous-system effects are important class risks.

**Levoleucovorin**

Levoleucovorin is the active isomer of leucovorin, used as methotrexate rescue and with selected chemotherapy regimens.

**Levoleucovorin Calcium**

Levoleucovorin calcium is the calcium salt of levoleucovorin, used to reduce methotrexate toxicity and support selected cancer treatments.

**Levomepromazine**

A phenothiazine medicine with antipsychotic, sedative, antiemetic, and analgesic properties, used in selected palliative-care and psychiatric settings.

**Levonorgestrel**

A progestin hormone used in birth-control pills, hormonal intrauterine devices, and emergency contraception to prevent or delay ovulation and alter cervical mucus.

**Levorphanol**

Levorphanol is a potent opioid analgesic that can relieve severe pain but also cause sedation, dependence, and respiratory depression.

---

### Levothyroxine

A synthetic form of thyroxine (T4) used as thyroid hormone replacement in hypothyroidism and in selected other thyroid conditions; dose is individualized and monitored with laboratory tests.

### Levothyroxine Sodium

A manufactured form of thyroid hormone used to replace hormone missing in hypothyroidism. The dose is adjusted using blood tests, and it is usually taken consistently on an empty stomach.

#### Alternate Names:

Levothyroxine

### Levulose

An older name for fructose, a simple sugar found in fruit, honey, and many sweeteners.

### Lewy Bodies

Abnormal deposits of the protein alpha-synuclein inside nerve cells. They are seen in Parkinson disease and dementia with Lewy bodies and are associated with movement, thinking, sleep, and visual-perception problems.

### Lewy Body

A Lewy body is an abnormal neuronal inclusion containing alpha-synuclein, characteristic of Parkinson disease and Lewy-body dementia.

### Lewy Body Dementia

A progressive brain disorder caused by abnormal alpha-synuclein protein deposits called Lewy bodies. It causes dementia with changing attention or alertness, recurrent visual hallucinations, Parkinson-like movement problems, sleep disorders such as acting out dreams, and sometimes automatic body-function problems such as constipation or dizziness on standing. Some people are very sensitive to antipsychotic medicines, so treatment must be individualized.

#### Alternate Names:

DLB; Dementia with Lewy Bodies

### Lewy Body Disease

A progressive brain disorder caused by abnormal protein deposits in nerve cells. It may cause changing attention, visual hallucinations, movement problems, stiffness, sleep-related behaviors, and gradual difficulty with thinking and daily activities.

### Leydig Cell

A cell between the testicular tubules that produces testosterone, mainly in response to luteinizing hormone.

### Leydig Cells

Cells found between the sperm-making tubes in the testes. In response to luteinising hormone, they make testosterone, which supports male sexual development, sperm production, sex drive, and muscle and bone health.

### Leydig Cell Testicular Tumor

A usually rare sex-cord stromal tumor arising from testosterone-producing Leydig cells in the testis; it may cause a testicular mass or hormone-related signs.

### Leydig Cell Tumor

A testicular sex cord-stromal tumor that secretes testosterone or estrogen, causing precocious puberty or gynecomastia; most are benign and treated by orchidectomy.

### Lyme Arthritis

Joint inflammation caused by Lyme disease, often with swelling in a large joint such as the knee. It can appear weeks or months after infection and is evaluated and treated as part of Lyme disease.

### Löfgren Syndrome

A sudden form of sarcoidosis causing swollen lymph nodes near the lungs, tender red lumps on the shins, and painful swollen ankles. It often improves over time; involvement of other organs can occur.

### LGE

Alternate name; see *Late Gadolinium Enhancement*.

---

## LH

Luteinising hormone, a hormone released by the pituitary gland. In people with ovaries it helps trigger ovulation and supports hormone production; in people with testes it stimulates testosterone production. Blood tests can help assess reproductive or pituitary problems.

### Alternate Names:

Luteinizing Hormone

## Lhermitte-Duclos Disease

A rare, usually slow-growing abnormal area in the cerebellum, the part of the brain that helps control balance and movement. It may cause headache, dizziness, poor coordination, or seizures and can be associated with PTEN-related conditions, but it is not by itself proof of one diagnosis.

### Alternate Names:

Dysplastic Gangliocytoma of the Cerebellum; Cerebellar Hamartoma

## Lhermitte Sign

A sudden electric shock-like sensation that travels down the neck and back into the arms or legs when bending the head forward, often seen in multiple sclerosis.

## Lhermitte's Sign

Alternate spelling; see *Lhermitte Sign*.

## LH Response To GnRH Blood Test

A hormone test that measures luteinizing hormone before and after a medicine that stimulates its release. It helps assess delayed or early puberty and problems involving the brain, pituitary gland, ovaries, or testes.

## Libido

A person's sexual desire or interest. It can change with age, relationships, mood, stress, sleep, illness, pregnancy, menopause, hormone levels, and medicines; a change is important mainly when it causes distress or affects a person's wellbeing.

## Libido And Sexual Performance

Sexual desire and sexual function can be influenced by hormones, blood flow, nerves, medicines, illness, mood, stress, sleep, and relationships. A persistent or distressing change deserves an assessment for treatable causes.

## Libman-Sacks Endocarditis

A sterile, nonbacterial endocardial valvulopathy with verrucous vegetations, most strongly associated with systemic lupus erythematosus and antiphospholipid syndrome. It usually affects the mitral or aortic valve and may cause regurgitation or systemic emboli; infective endocarditis must be excluded when vegetations or fever are present.

### Alternate Names:

Marantic Endocarditis; Verrucous Endocarditis; Nonbacterial Verrucous Endocarditis

## Lice

Small insects that live on hair or clothing and feed on blood. Head, body, and pubic lice cause itching and may leave eggs attached to hair. They spread mainly through close contact and are not a sign of poor hygiene.

## Lice Infestation

Alternate term; see *Lice*.

## Lice Infestations

Alternate plural form; see *Lice*.

## Licensed Clinical Social Worker

A social worker licensed to provide clinical assessment and psychotherapy or other mental-health services within the scope of local law and training.

## Lichen

A descriptive term for a firm, flat-topped, raised skin lesion or for a skin pattern resembling lichen planus. It describes appearance rather than a single disease or cause.

## Lichen Amyloidosis

A localized cutaneous amyloidosis causing intensely

---

itchy, rough, hyperkeratotic papules, commonly on the shins or upper back. The deposited amyloid usually derives from damaged keratinocytes and remains confined to the skin.

### **Lichenification**

Thick, leathery skin with more visible skin lines caused by repeated scratching or rubbing. It often develops in long-lasting eczema, allergies, or other itchy skin conditions.

### **Lichenified**

Lichenified skin is thickened and leathery from chronic rubbing or scratching, commonly occurring in longstanding eczema or itch disorders.

### **Lichen Myxedematosus**

A group of rare disorders in which mucin accumulates in the skin, producing firm papules, plaques, or diffuse thickening. The generalized form may be associated with monoclonal gammopathy and internal-organ disease, whereas localized forms may remain confined to the skin.

### **Lichen Nitidus**

A benign inflammatory skin disorder causing numerous tiny, shiny, flat-topped, skin-colored or pink papules, commonly on the trunk, arms, genital area, or hands. It is often asymptomatic and may resolve spontaneously, although itch or extensive disease may need treatment.

### **Lichenoid Drug Eruption**

A lichenoid inflammatory skin reaction caused by a medicine, often resembling lichen planus but with a broader distribution or different clinical and histologic features; improvement may follow withdrawal of the culprit drug.

### **Lichen Planopilaris**

A lymphocytic inflammatory scalp disorder that attacks hair follicles and can cause scarring alopecia, perifollicular scale, redness, and permanent hair loss.

### **Lichen Planus**

A noncontagious, immune-mediated inflammatory condition that can affect the skin, mouth, genitals, nails, scalp, or hair follicles. On the skin it often causes itchy, shiny, flat-topped bumps that may look red, purple, brown, or gray, especially on the wrists, ankles, or lower back. In the mouth it may cause a lacy white pattern, soreness, or ulcers; genital disease may cause soreness or erosions; nail disease can cause ridges, thinning, splitting, or loss; and scalp disease can cause scarring hair loss.

#### **Alternate Names:**

Lichen Ruber Planus; Cutaneous Lichen Planus

### **Lichen Ruber Planus**

Alternate name; see *Lichen Planus*.

### **Lichen Sclerosus**

A long-term inflammatory skin condition that usually affects the genital or anal area, causing pale, thin, itchy, easily injured skin. Treatment relieves symptoms and lowers scarring; persistent sores need review because cancer risk is slightly increased.

#### **Alternate Names:**

White Spot Disease

### **Lichen Simplex Chronicus**

A thickened, leathery plaque caused by chronic scratching and rubbing (itch-scratch cycle). It commonly affects the neck, ankles, lower legs, and genital area. It is a form of neurodermatitis associated with stress and anxiety.

### **Lichen Spinulosus**

A benign follicular eruption of grouped, rough, spiny papules, often forming round or oval plaques on the trunk, buttocks, or limbs. It commonly begins in childhood or adolescence and may fade spontaneously.

### **Lichen Striatus**

A usually harmless skin condition causing small, flat, pink, brown, or pale bumps in a line along a limb. It often affects children and usually clears without treatment.

---

## Licorice Poisoning

Excess glycyrrhizin from licorice inhibits 11-beta-hydroxysteroid dehydrogenase type 2 and produces apparent mineralocorticoid excess. Hypertension, hypokalemia, metabolic alkalosis, edema, weakness, and dysrhythmias may occur; stop the source and correct potassium and blood pressure under medical supervision.

## Liddle Syndrome

A rare inherited cause of early high blood pressure in which the kidneys retain too much sodium and lose too much potassium. It may begin in childhood or young adulthood and usually requires specialist treatment.

## Liddle Syndrome Treatment

Treatment of Liddle syndrome lowers blood pressure and the body's excessive sodium retention, usually with a medicine that blocks the overactive kidney channel. Potassium and kidney function are monitored, and relatives may need testing.

## Lidocaine

Lidocaine is a local anaesthetic that blocks voltage-gated sodium channels and, intravenously, can treat selected ventricular arrhythmias; excessive exposure may cause seizures, hypotension, arrhythmia, or cardiac arrest.

## LieberküHn's Crypts

Tubular glands in the intestinal lining between villi that contain stem, absorptive, and secretory cells. They renew the lining, produce intestinal substances, and help maintain the gut's protective barrier.

## Life Expectancy

The average number of additional years a person of a specified age is expected to live under current mortality conditions.

## Lifetime Risk

The probability that a person will develop a specified disease or experience an outcome during the remainder of their life or over a defined lifetime period.

## Li-Fraumeni Syndrome

An inherited condition that greatly increases the chance of developing several cancers, often at a young age. Cancers may include breast cancer, sarcoma, brain tumors, adrenal cancer, or leukemia; specialized screening is usually advised.

## Ligament

A dense, fibrous connective tissue band that connects bone to bone, providing passive joint stability and guiding joint motion. Ligaments are composed predominantly of type I collagen fibers (parallel orientation, crimped pattern) and elastin fibers (10-20% in some ligaments), arranged in bundles.

## Ligamentum Flavum

A series of paired, thick elastic ligaments connecting the laminae of adjacent vertebrae from the axis to the sacrum, whose high elastin content imparts a characteristic yellow color and helps preserve the upright posture while preventing excessive spinal flexion.

## Ligamentum Teres Hepatis

The obliterated fibrous remnant of the embryonic left umbilical vein that carried oxygenated blood from the placenta to the fetus, running within the free inferior border of the falciform ligament from the umbilicus to the left branch of the portal vein.

### Alternate Names:

Round Ligament of the Liver

## Ligand

Any molecule that binds specifically to a biological receptor, macromolecule, or active site. In pharmacology and physiology, ligands include hormones, neurotransmitters, drugs, and toxins that bind receptors to initiate (agonist), block (antagonist), or modulate signaling.

## Ligase

An enzyme that joins two molecules, often using energy from ATP. Ligases help repair DNA, copy genetic material, and build or maintain other important molecules.



---

## Ligation

The surgical process of tying off a blood vessel or duct using a suture or clip to prevent bleeding or block fluid flow.

## Ligature

Any cord, wire, rope, fabric, or flexible object used to constrict the neck, bind limbs, suspend the body, or inflict injury. The resulting pale groove-shaped abrasion may be horizontal with petechiae in homicidal strangulation, or oblique and upward in hanging.

## Light

Visible electromagnetic radiation; in medicine it is used for examination, phototherapy, imaging, and selected treatments.

## Light Chain-Associated Kidney Disease

Kidney injury caused by abnormal immunoglobulin light chains produced by a plasma-cell or B-cell clone. Light chains may cause cast nephropathy, amyloid, or other deposits, leading to proteinuria, acute kidney injury, or chronic kidney disease.

## Light-Chain Deposition Disease

A monoclonal plasma-cell disorder in which non-amyloid immunoglobulin light chains deposit along kidney and sometimes other organ structures. It commonly causes proteinuria, high blood pressure, declining kidney function, or nephrotic syndrome.

## Light Coagulation

Light coagulation uses intense focused light, usually a laser, to heat and seal tissue or blood vessels.

## Light Cycle

A light cycle is the recurring pattern of light and darkness that helps regulate circadian rhythms and biological activity.

## Lightening

In late pregnancy, descent of the fetus into the pelvis before labor, which may reduce upper-abdominal pressure but increase pelvic pressure and urinary frequency.

## Lighter Fluid Poisoning

Harm from swallowing or breathing lighter fluid, a hydrocarbon. The greatest danger is liquid entering the lungs, causing chemical pneumonia, cough, or breathing difficulty.

## Lightheadedness

A feeling of near-fainting, floating, or unsteadiness without necessarily perceiving that the surroundings are spinning; causes include dehydration, low blood pressure, anemia, medications, and heart or neurologic conditions.

## Light Microscope

An instrument that uses visible light and lenses to magnify cells, tissues, microorganisms, or other small structures. It is widely used in clinical laboratories, pathology, microbiology, and research.

## Light Microscopy

Examination of cells, tissues, microorganisms, or other specimens using visible light and a microscope. Different stains and preparation methods reveal structure, infection, inflammation, blood cells, or abnormal growth.

## Light-Near Dissociation

A pupil abnormality in which the pupils constrict when focusing on a nearby object but respond poorly or not at all to light. It can occur with certain nerve disorders, infections, diabetes, or midbrain disease.

## Lightning Injuries

Injuries caused by direct or indirect lightning exposure, potentially affecting the heart, nervous system, skin, hearing, muscles, and lungs. Cardiac arrest, neurologic deficits, burns, trauma from falls, and temporary skin branching may occur.

## Lightning Pains

Brief, stabbing, electric-shock-like pains caused by irritated or damaged nerves. They classically occur in tabes dorsalis but may also accompany shingles, nerve compression, diabetes, or other neurological disorders.

---

## Light Reflex

Automatic narrowing of the pupil when light enters the eye. It tests the retina, optic nerve, brain pathways, and nerve that controls the pupil; an unequal or absent response may indicate disease.

## Light Scattering

Deflection of light when it meets particles, cells, molecules, or irregular structures. Measuring the scattered light can help estimate particle size or concentration in laboratory tests and can affect the appearance of tissues on imaging.

## Light's Criteria

A set of laboratory comparisons used to decide whether fluid around the lung is caused by inflammation or disease in the chest rather than by pressure or fluid imbalance elsewhere. Protein and enzyme levels in the fluid are compared with blood levels; the result helps guide further testing and treatment.

## Light Therapy

Treatment that uses selected wavelengths or intensity of light for conditions such as some skin disorders, seasonal depression, or neonatal jaundice.

## Lignocaine

The international nonproprietary name for lidocaine, a local anesthetic and antiarrhythmic drug.

## Likelihood Ratio

A measure of how strongly a test result supports or argues against a disease. A value above one supports disease, a value below one argues against it, and larger changes are more informative.

## Lilliputian Hallucination

A visual hallucination of unusually small people, animals, or objects; it can occur with neurologic, psychiatric, toxic, infectious, or sensory conditions.

## Lily Of The Valley Poisoning

Poisoning from eating lily-of-the-valley plant parts, which contain substances that can disrupt the heart-beat. Nausea, vomiting, dizziness, slow or irregular

pulse, confusion, or fainting may occur.

## Limb

An arm or leg, including its bones, joints, muscles, tendons, nerves, blood vessels, and skin. Limb problems may affect movement, sensation, strength, circulation, or function.

## Limb Bud

A limb bud is an embryonic outgrowth that develops into an arm or leg through coordinated signaling, growth, and tissue differentiation.

## Limb Development

Limb development is embryonic formation and growth of the bones, joints, muscles, nerves, vessels, and skin of the arms and legs.

## Limb Fracture

A break in a bone of an upper or lower limb, assessed according to location, displacement, open or closed status, joint involvement, and associated neurovascular injury.

## Limb Girdle

A limb girdle is the skeletal connection of a limb to the trunk, such as the shoulder or pelvic girdle.

## Limb-Girdle Muscular Dystrophies

Limb-girdle muscular dystrophies are inherited muscle diseases causing progressive weakness predominantly around the hips and shoulders, with variable cardiac and respiratory involvement.

## Limbic Encephalitis

Inflammation of the limbic system, often presenting subacutely with memory impairment, seizures, psychiatric or behavioral change, and sleep disturbance. It may be autoimmune, paraneoplastic, infectious, or of unknown cause.

## Limbic Lobe

A curved region on the inner surface of the brain involved in memory, emotion, motivation, and some responses controlled by the nervous system.

---

## **Limbic System**

A group of connected brain structures involved in memory, emotion, motivation, learning, and some automatic body responses, such as changes in heart rate during stress.

## **Limb Injury**

Trauma affecting an arm or leg, including injury to bone, joint, muscle, tendon, ligament, nerve, blood vessel, or skin.

## **Limb-Length Discrepancy**

A difference in the lengths of the two legs or, less commonly, the two arms. It may result from growth differences, injury, infection, surgery, or a developmental condition and can affect posture, walking, and joint loading.

## **Limb Pain**

Pain arising from an arm or leg, with causes including trauma, nerve disease, vascular insufficiency, infection, inflammation, and musculoskeletal disorders.

## **Limb Plethysmography**

A noninvasive test measuring changes in limb volume or blood flow, used to evaluate arterial or venous obstruction and sometimes to assess vascular responses.

## **Limb Prosthesis**

A limb prosthesis is an artificial device replacing part or all of an arm or leg to restore mobility, function, appearance, or independence.

## **Limb Reduction Defects**

Birth defects in which part or all of an arm or leg does not form normally. The missing or shortened part may involve a bone, joint, hand, foot, or the whole limb.

## **Limbs**

The arms and legs, including their bones, joints, muscles, nerves, blood vessels, and skin. Limb problems may affect strength, sensation, movement, cir-

ulation, balance, daily activities, or independence.

## **Limbus**

The border where the clear cornea meets the white outer coat of the eye. It contains cells that help renew the corneal surface and may be affected by injury, inflammation, or disease.

### **Alternate Names:**

Corneoscleral Limbus

## **Limbus Corneae**

The corneal limbus is the circular border where the transparent cornea meets the white sclera and conjunctiva, containing stem cells important for corneal repair.

## **Limited Range Of Motion**

Limited range of motion is reduced ability to move a joint through its normal extent because of pain, stiffness, weakness, swelling, or structural change.

## **Linagliptin**

Linagliptin is a DPP-4 inhibitor used with diet and exercise to improve blood glucose control in type 2 diabetes.

## **Lincomycin**

Lincomycin is a lincosamide antibiotic used for selected serious bacterial infections; it can cause antibiotic-associated diarrhea and colitis.

## **Lindane**

Lindane is an organochlorine insecticide formerly used for lice or scabies; neurotoxicity and environmental persistence limit its use.

## **Lindsay's Nails**

A distinctive nail abnormality in which the proximal forty to fifty percent of the nail bed is white and opaque while the distal fifty to sixty percent is pink, red, or brownish. It is strongly associated with chronic kidney disease and uraemia.

### **Alternate Names:**

Half-and-Half Nails; Lindsay Nails

---

## Linea

A line or linear anatomical marking, often used in names such as linea alba or linea nigra.

### Linea Alba

A dense, fibrous median collagenous band formed by the interlacing decussating aponeuroses of the external oblique, internal oblique, and transversus abdominis muscles, extending vertically from the xiphoid process to the pubic symphysis.

### Linea Aspera

Prominent longitudinal posterior femoral shaft ridge from converging muscle attachment lines. Insertion for adductor magnus, vastus medialis/lateralis, and short head of biceps femoris. Thickest femoral cortex, resisting bending forces. Diverges distally into supracondylar lines.

### Linea Nigra

A dark brown or black vertical line appearing on the abdomen during pregnancy, extending from the pubic symphysis to the umbilicus or beyond. Caused by hormonal stimulation of melanocytes, resulting in hyperpigmentation. More prominent in darker-skinned women.

## Linear

Arranged in a straight or nearly straight line, describing the shape or distribution of a skin lesion or eruption.

### Linear Accelerator

A machine accelerating electrons to produce high-energy X-rays or electron beams that precisely deliver radiation treatment to selected tumors.

### Linear Energy Transfer

The amount of energy that radiation deposits as it travels through tissue. Radiation with higher transfer deposits energy more densely and may cause more biological damage for the same total dose.

### Linear Fracture

A narrow break that runs along the length of a bone

without moving the pieces far apart or breaking the bone into many fragments. Treatment depends on the bone, the injury's stability, and whether nearby nerves, blood vessels, or organs are affected.

### Linear IgA Bullous Dermatitis

An autoimmune blistering disease caused by linear deposition of IgA at the basement membrane zone. It produces tense blisters, sometimes in annular or "string of pearls" patterns, and may be idiopathic or drug-induced.

### Linear Nevus Sebaceous Syndrome

A rare disorder involving a linear sebaceous nevus, a skin lesion caused by abnormal skin glands, together with possible eye, brain, or skeletal abnormalities. It requires evaluation based on the organs involved.

### Lined Canister

A container with a protective inner lining used to collect or store a medical substance, specimen, waste, or device component. The lining helps prevent leakage, contamination, chemical reaction, or damage to the container; its exact use depends on the equipment.

### Lines of Blaschko

Non-random developmental cutaneous pathways representing the routes of epidermal cell migration and clonal proliferation during embryogenesis, which become apparent in X-linked mosaic and cutaneous mosaic disorders (such as incontinentia pigmenti and linear epidermal nevi).

#### Alternate Names:

Blaschko Lines; Blaschko's Tracks

### Lines of Zahn

Microscopic and gross alternating pale and dark laminations seen in arterial and intracardiac thrombi formed in rapid pulsatile blood flow, composed of pale layers of platelets and fibrin mesh alternating with dark layers of trapped red blood cells, distinguishing antemortem thrombi from postmortem clots.

#### Alternate Names:

Zahn Lines; Laminations of Zahn

---

## Linezolid

Linezolid is an oxazolidinone antibiotic that blocks formation of the bacterial 70S initiation complex and treats resistant Gram-positive infections; thrombocytopenia, neuropathy, lactic acidosis, and serotonin interactions require caution.

## Lingua

The anatomical Latin term for the tongue, a muscular organ used for chewing, swallowing, speaking, and tasting. The front two-thirds lies in the mouth and the back third lies in the throat; related terms include sublingual, meaning under the tongue.

## Lingual

Relating to the tongue or the tongue side of a body structure. Lingual muscles, nerves, blood vessels, and surfaces help with taste, speech, swallowing, and movement of food.

## Lingual Artery

A branch of the external carotid artery supplying the tongue, floor of mouth, and sublingual structures. It arises anterior to the hypoglossal nerve and courses deep to the hyoglossus muscle. Lingual artery aneurysm or thrombosis is rare but can cause life-threatening hemorrhage or tongue ischemia.

## Lingual Thyroid

A lingual thyroid is thyroid tissue at the base of the tongue caused by failure of normal thyroid descent and may obstruct swallowing or breathing or cause hypothyroidism.

## Lingual Tonsil

The lingual tonsil is lymphoid tissue at the base of the tongue that contributes to immune surveillance of the throat.

## Lingual Tonsils

Collections of lymphoid tissue at the base of the tongue.

## Lingua Villosa Nigra

A benign condition characterized by elongation and

dark pigmentation of the filiform papillae of the tongue, often associated with tobacco use, poor oral hygiene, or antimicrobial therapy.

## Liniment

A liquid or semiliquid preparation rubbed onto the skin, usually to provide temporary relief of muscle or joint discomfort. Some contain irritating or warming substances and should not be used on broken skin, near the eyes, or under tight coverings unless directed.

## Linitis Plastica

A form of stomach cancer that spreads widely through the stomach wall, making it thick and stiff. It may cause early fullness, weight loss, nausea, vomiting, or vague upper-abdominal discomfort.

## Linoleic Acid

An essential omega-6 polyunsaturated fatty acid obtained from food. The body uses it to make cell membranes and signaling molecules.

## Linolenic Acid

A polyunsaturated fatty acid. Alpha-linolenic acid is an essential omega-3 nutrient from plant foods, while other forms occur in animal tissues and have different roles.

## Liothyronine Sodium

A manufactured form of triiodothyronine, an active thyroid hormone, used in selected thyroid disorders. It acts quickly and can cause fast heartbeat, tremor, anxiety, bone loss, or heart strain if excessive.

## Lip

The lips are mobile folds of skin and mucosa containing muscle, blood vessels, and sensory nerves; they form the oral seal and assist speech, facial expression, feeding, and swallowing.

## Lipase

An enzyme that breaks down fats into smaller molecules. Blood lipase testing is commonly used when pancreatitis is suspected, although an abnor-

---

mal result is not specific in every case.

### **Lipase Test**

A lipase test measures a pancreatic digestive enzyme in blood and supports evaluation of acute pancreatitis and selected pancreatic or gastrointestinal disorders.

### **Lip Cancer**

Cancer developing in the tissue of the upper or lower lip. It may appear as a persistent sore, lump, crust, bleeding area, numbness, or color change.

### **Lip Disease**

A disorder of the lips, such as infection, inflammation, ulceration, trauma, pigmentation change, benign tumor, or malignancy.

### **Lipedema**

A long-term disorder causing an unusual, usually symmetrical buildup of painful fat in the legs and sometimes arms. The feet and hands are usually spared, while easy bruising, tenderness, and swelling may occur.

### **Lipemia Retinalis**

A creamy pink-white appearance of the blood vessels at the back of the eye caused by extremely high blood triglycerides. It signals severe fat buildup in the blood and a high risk of sudden pancreatitis.

### **Lipemic Index**

A laboratory estimate of how cloudy a blood sample is because of excess fats. Marked cloudiness can interfere with some test results and may prompt review of the sample or repeat testing.

### **Lipid**

A fat or fat-like substance, including triglycerides, cholesterol, phospholipids, and fat-soluble vitamins. Lipids store energy, form cell membranes, help make hormones, and help the body absorb vitamins A, D, E, and K.

### **Lipid A**

The part of a substance on the outer surface of some bacteria that triggers a strong immune reaction. Large amounts can contribute to widespread inflammation, shock, and abnormal clotting.

### **Lipid Bilayer**

The double layer of fat-like molecules forming the basic structure of cell membranes. It separates the cell from its surroundings and controls movement of substances.

### **Lipid Disorder**

An abnormal level or handling of fats in the blood or tissues, such as high LDL cholesterol, high triglycerides, or unusually low protective HDL cholesterol.

### **Lipid Emulsion Therapy**

Emergency intravenous treatment for severe poisoning from a local anesthetic or selected other fat-soluble substances. The fat may draw some drug away from the heart and brain and provide energy to the heart; it is given under medical supervision.

### **Lipid Metabolism**

The processes by which the body digests, transports, stores, makes, and breaks down fats for energy, cell membranes, and hormone production.

### **Lipidosis**

A disorder involving abnormal accumulation or metabolism of lipids in tissues, often due to inherited enzyme defects.

### **Lipid Panel**

A routine blood test measuring circulating blood fats to assess heart and blood vessel disease risk. It measures four key markers: total cholesterol, LDL cholesterol (often termed "bad" cholesterol because it contributes to artery-clogging plaque), HDL cholesterol ("good" cholesterol that helps clear excess fat from arteries), and triglycerides (stored fat used for energy). It evaluates risk for heart attacks and strokes, helps guide diet and lifestyle changes, and monitors response to cholesterol-lowering medicines such as statins.



---

**Alternate Names:**

Lipid Profile; Cholesterol Panel; Coronary Risk Panel; Lipid Profile Test

**Lipid Profile**

Alternate name; see *Lipid Panel*.

**Lipid Profile Test**

Alternate name; see *Lipid Panel*.

**Lipids**

A group of mostly water-insoluble substances that includes fats, oils, fatty acids, phospholipids, and cholesterol. Lipids store energy, form cell membranes, help make hormones, and help the body absorb vitamins A, D, E, and K.

**Lipid Transport**

Movement of fats through blood and tissues by lipoproteins, carrier proteins, or other mechanisms for energy use, storage, membrane formation, and hormone production.

**Lip Infection**

Infection of the lip caused by bacteria, viruses, fungi, or an infected wound. It may cause pain, redness, swelling, blisters, cracks, pus, or fever; rapidly spreading swelling or breathing difficulty may occur in a serious infection.

**Lipitor**

Lipitor is a brand name for atorvastatin, a statin medicine that lowers low-density lipoprotein cholesterol and reduces the risk of heart attack and stroke. Muscle symptoms and liver-related effects are possible.

**Lip Moisturizer Poisoning**

Swallowing a small amount of ordinary lip moisturizer usually causes mild stomach upset or loose stools. Products containing camphor, salicylates, or other medicines can cause serious poisoning and require poison-control advice.

**Lip Neoplasm**

A lip neoplasm is an abnormal growth in the lip that may be benign or malignant; persistent sores or lumps require assessment.

**Lipoatrophy**

Loss of the layer of fat beneath the skin, causing dents or hollow areas. It can follow repeated injections, inflammation, autoimmune disease, HIV treatment, or inherited disorders and may alter medicine absorption at injection sites.

**Lipoblastoma**

A usually benign tumor of immature adipose tissue occurring mainly in infants and young children; it may be localized or diffuse and can recur if incompletely removed.

**Lipodermatosclerosis**

A chronic fibrosing inflammatory panniculitis of the subcutaneous adipose tissue of the lower extremity secondary to severe chronic venous insufficiency, producing induration, hyperpigmentation, and the classic "inverted champagne bottle" leg appearance.

**Alternate Names:**

Sclerosing Panniculitis

**Lipodystrophy**

Abnormal loss or distribution of body fat. It may be inherited or caused by medicines, metabolic disease, or repeated injections and can lead to insulin resistance, diabetes, high triglycerides, or changes in appearance.

**Lipodystrophy Syndromes**

Rare disorders in which body fat is absent from some areas, present from birth or lost later, and may collect excessively elsewhere. They can cause severe insulin resistance, diabetes, fatty liver, and very high triglycerides.

**Lipofuscin**

Lipofuscin is a yellow-brown, non-degradable pigment made of oxidized lipid and protein residues that accumulates in long-lived cells with ageing and

---

in some degenerative diseases.

### **Lipogranulomatosis**

Inflammation in which fat accumulates and triggers organized collections of immune cells called granulomas. It may affect skin, soft tissue, or internal organs and can result from injury, leakage of oily material, or metabolic disease.

### **Lipohypertrophy**

A firm or lumpy buildup of fat under the skin, often at repeated insulin-injection sites. Injecting into these areas can change how insulin is absorbed.

### **Lipoid Nephrosis**

An older name for a kidney disorder, most often in children, in which the kidney filters leak large amounts of protein into the urine. It can cause swelling around the eyes, legs, or abdomen.

### **Lipoidosis**

Abnormal buildup of fats within cells or tissues, usually because a metabolic pathway cannot process or transport them normally. It may be inherited or acquired and can damage organs such as the liver, nerves, or kidneys.

### **Lipoid Pneumonia**

Lung inflammation caused by oil or fat entering the airways or building up in lung tissue. It may follow inhalation or swallowing of oil-based products and can cause cough, breathlessness, fever, or chest changes on imaging.

### **Lipolysis**

Breakdown of stored body fat into fatty acids and glycerol so they can be used for energy. Fasting and stress hormones increase this process, while insulin generally reduces it.

### **Lipoma**

A common benign tumor made of mature fat cells. It is usually soft, movable, painless, and slow-growing beneath the skin of the trunk, neck, or limbs; a deep, firm, painful, or rapidly enlarging fatty lump needs

assessment to distinguish it from other tumors.

### **Lipomas**

Plural of lipoma; see *Lipoma*.

### **Lipomastia**

Alternate name; see *Pseudogynecomastia*.

### **Lipomatosis**

A condition in which mature fat tissue grows in multiple or broad areas rather than forming one separate lump. It may be symmetrical, involve the trunk or limbs, and be difficult to remove when it extends between nearby structures.

### **Lipomatous Neoplasm**

A growth made partly or mainly of fat cells. It may be noncancerous, like a lipoma, or cancerous, like a liposarcoma; a deep, firm, painful, or rapidly enlarging lump needs specialist assessment.

### **Lipophilicity**

Lipophilicity is the tendency of a substance to dissolve in fats or oils rather than water, influencing drug absorption and distribution.

### **Lipopolysaccharide**

Lipopolysaccharide is the outer-membrane endotoxin of Gram-negative bacteria, consisting of lipid A, core polysaccharide, and O antigen; circulating lipid A can trigger cytokine release, fever, shock, and disseminated inflammation.

### **Lipoprotein**

A particle made of fat and protein that carries cholesterol and other fats through the blood. Different types transport dietary fat, triglycerides, or cholesterol; high LDL levels can contribute to artery disease.

### **Lipoprotein A**

A cholesterol-carrying particle similar to LDL whose level is largely inherited. A high level can increase the long-term risk of heart attack, stroke, and narrowing of the aortic valve.

---

**Alternate Names:**

Lp(a)

**Lipoprotein(a)**

An independent, genetically determined causal risk factor for atherosclerotic cardiovascular disease and calcific aortic valve stenosis, consisting of an LDL-like particle covalently linked by a disulfide bond to a highly polymorphic, plasminogen-like glycoprotein called apolipoprotein(a).

**Alternate Names:**

Lp(a)

**Lipoprotein Analysis**

A blood test measuring cholesterol, triglycerides, and particles such as LDL and HDL to estimate cardiovascular risk and guide treatment.

**Alternate Names:**

Lipid Panel; Lipid Profile

**Lipoprotein Lipase**

An enzyme on capillary walls that breaks down triglycerides in chylomicrons and very-low-density lipoproteins so tissues can use or store the fatty acids. Its activity is regulated by nutritional and hormonal signals.

**Lipoprotein Metabolism**

The body's production, transport, modification, and removal of particles that carry cholesterol and other fats through the bloodstream.

**Liposarcoma**

A malignant tumor of fat cells, usually arising in the deep soft tissues of the thigh or retroperitoneum; well-differentiated, myxoid, and pleomorphic subtypes are recognized.

**Liposarcoma Of The Breast**

A rare malignant mesenchymal tumor with adipocytic differentiation arising in breast soft tissue. Diagnosis requires histopathologic and molecular evaluation, and treatment is generally wide excision with sarcoma-focused care.

**Liposomal Amphotericin B**

A liposomal formulation of amphotericin B used in the treatment of severe systemic fungal infections, with reduced nephrotoxicity compared to conventional amphotericin B. The liposomal formulation encapsulates amphotericin B within lipid vesicles, which alters the drug's biodistribution and reduces renal tubular damage.

**Liposome**

A liposome is a tiny vesicle made of a lipid bilayer that can carry medicines or other substances into or around the body.

**Liposuction**

A surgical procedure that uses a cannula and suction to remove localized deposits of subcutaneous fat for body contouring. It is not a treatment for obesity or a substitute for diet and exercise. Risks may include bleeding, infection, contour irregularity, fluid accumulation or shifts, sensory changes, and anesthesia-related complications.

**Alternate Names:**

Lipoplasty

**Lips**

The paired movable folds surrounding the opening of the mouth, involved in speech, facial expression, eating, and oral sensation.

**Lipschütz Bodies**

Older term; see *Acute Genital Ulcers*.

**Lip Swelling**

Enlargement of one or both lips caused by local injury, infection, inflammation, dental disease, medication, or angioedema. Sudden swelling with tongue or throat swelling, breathing difficulty, wheezing, faintness, or widespread hives may indicate a severe allergic reaction.

**Alternate Names:**

Cheilitis; Lip Edema

**Liquefaction**

Conversion of a solid or partly solid material into

---

liquid, such as tissue breakdown during liquefactive necrosis or melting of a substance.

### **Liquefactive Necrosis**

A pattern of tissue death in which enzymes digest dead cells until the area becomes soft or liquid. It commonly occurs in brain injury and some infections and may leave a fluid-filled cavity.

### **Liquid Bandage for Lacerations**

Application of topical cyanoacrylate tissue adhesives to seal minor, clean, low-tension superficial lacerations and abrasions. Provides a waterproof antimicrobial barrier that sloughs off naturally as epidermal re-epithelialization occurs over 5–10 days.

#### **Alternate Names:**

Tissue Adhesive Closure; Topical Skin Glue; Cyanoacrylate Liquid Bandage

### **Liquid Biopsy**

A non-invasive laboratory technique that analyzes tumor-derived biomarkers, including circulating tumor cells, cell-free circulating tumor DNA (ctDNA), and exosomes isolated from peripheral blood or other bodily fluids for cancer detection and genomic profiling.

#### **Alternate Names:**

Liquid Biopsy

### **Liquid Chromatography**

A laboratory method that separates dissolved substances as a liquid carries them through a material that interacts differently with each substance. It is used to identify or measure medicines, hormones, metabolites, proteins, and toxins.

### **Liquid Diet**

A diet consisting partly or entirely of liquids; clear-liquid and full-liquid diets differ in allowed foods and may be prescribed temporarily for procedures, illness, or swallowing or digestive problems.

### **Liquid Incense**

A scented liquid product that may contain solvents, oils, or other chemicals. Swallowing or inhaling it

can cause irritation, drowsiness, vomiting, seizures, or lung injury; the product container is needed for poison-control assessment.

### **Liquid Nitrogen**

Liquid nitrogen is extremely cold nitrogen used to freeze tissue in cryotherapy; contact can cause frost-bite and its gas can displace oxygen.

### **Liquor**

A liquid; in clinical usage it may refer to cerebrospinal fluid or a medicinal preparation, depending on context.

### **LI-RADS**

A standardized system for reporting liver-imaging findings in people at increased risk of liver cancer, especially those with cirrhosis. Its categories estimate how likely a finding is to be cancer and help guide follow-up or biopsy.

### **Liraglutide**

Liraglutide is a GLP-1 receptor agonist used to improve glucose control in type 2 diabetes and, at some doses, support weight management.

### **Lisch Nodule**

A Lisch nodule is a harmless pigmented hamartoma of the iris, commonly associated with neurofibromatosis type 1.

### **Lisch Nodules**

Benign, dome-shaped melanocytic hamartomas on the iris, commonly seen in neurofibromatosis type 1. They usually do not affect vision and support the diagnosis when interpreted with other clinical findings.

### **Lisfranc Fracture**

A severe midfoot injury characterized by fracture, subluxation, or dislocation of the tarsometatarsal joints, classically involving disruption of the strong Lisfranc ligament connecting the medial cuneiform to the base of the second metatarsal bone.

#### **Alternate Names:**

---

Lisfranc Joint Complex Injury; Tarsometatarsal Fracture-Dislocation

### **Lisfranc Injury**

A spectrum of traumatic fracture-dislocations involving the tarsometatarsal articulation of the mid-foot, typically disrupting the critical Lisfranc ligament extending between the medial cuneiform and the base of the second metatarsal.

### **Lisinopril**

Lisinopril is an angiotensin-converting-enzyme inhibitor used for hypertension, heart failure, and kidney protection in some diabetes; cough, hyperkalemia, renal-function changes, angioedema, and fetal toxicity are important risks.

### **Lissencephaly**

A birth condition in which the brain's surface has few or no normal folds because nerve cells did not move to their usual positions during development. It can cause seizures, severe developmental impairment, feeding problems, and movement difficulties.

### **Listeria**

Listeria is a genus of bacteria; *L. monocytogenes* can cause foodborne illness, bloodstream infection, and meningitis, especially in pregnancy, older adults, and people with weakened immunity.

#### **Alternate Names:**

Listeria Species; Listerial Infection

### **Listeria Infection**

An infection caused by Listeria bacteria, usually acquired from contaminated food. It may cause fever, muscle aches, diarrhea, or vomiting; infection can be severe in pregnancy, newborns, older adults, and people with weakened immune systems.

### **Listeria Monocytogenes**

A gram-positive, facultatively anaerobic rod that causes listeriosis, a foodborne infection. The organism is transmitted through contaminated food, particularly deli meats, soft cheeses, and ready-to-eat foods. It can grow at refrigeration temperatures.

### **Listeriosis**

Infection with *Listeria monocytogenes*, usually acquired from contaminated food. It may cause fever and diarrhea or severe disease such as bloodstream infection or meningitis, especially in pregnancy, newborns, older adults, and people with weakened immunity.

### **Listeriosis In Pregnancy**

A foodborne infection that may cause only fever, tiredness, or muscle aches in the pregnant person but can spread to the fetus. It can cause miscarriage, stillbirth, premature birth, or severe infection in the newborn.

### **Lithiasis**

The formation or presence of stones (calculi) in an organ or duct. The term is often named for the site, such as nephrolithiasis in the kidney, ureterolithiasis in the ureter, or cholelithiasis in the gallbladder.

### **Lithium**

Lithium is a mood stabilizer used mainly for bipolar disorder and selected augmentation treatment; its narrow therapeutic range requires monitoring for tremor, thyroid or renal dysfunction, dehydration, drug interactions, and toxicity.

### **Lithium Carbonate**

A mood-stabilizing medicine used mainly to treat bipolar disorder and reduce the risk of manic or depressive episodes; serum levels, kidney function, thyroid function, and drug interactions require monitoring.

### **Lithium Nephropathy**

Kidney injury related to long-term or toxic lithium exposure, most often causing impaired urine concentration and excessive urination, and sometimes chronic tubulointerstitial scarring or hyperparathyroidism. Monitoring lithium levels, kidney function, and calcium helps reduce risk.

### **Lithium Toxicity**

Poisoning caused by too much lithium in the body, often after dehydration, illness, a dose change, or

---

a medicine interaction. Nausea, diarrhea, worsening tremor, weakness, confusion, poor coordination, seizures, or reduced consciousness may occur.

### **Lithotomy**

A surgical procedure for removing a stone, or the position with the hips and knees flexed and legs supported.

### **Lithotomy Position**

A supine position in which the hips and knees are flexed and the legs are raised and supported to provide access to the pelvis and perineum.

### **Lithotripsy**

A medical procedure that breaks hard mineral stones located in the kidneys, ureters, bladder, or gallbladder into tiny, sand-like fragments so they can pass out of the body naturally in the urine or be removed. The most common form is Extracorporeal Shock Wave Lithotripsy (ESWL), a completely non-invasive technique that generates high-energy acoustic shock waves outside the body and focuses them through the skin and soft tissues onto the stone under X-ray or ultrasound guidance, shattering it into dust without cuts or incisions. When stones are very dense, stuck low in the ureter, or unsuitable for ESWL, intracorporeal or laser lithotripsy is used, where a thin endoscope is threaded up the urinary tract and a fine holmium laser fiber fragments the stone under direct camera visualization. Lithotripsy relieves the severe cramping flank pain (renal colic) and prevents kidney obstruction and recurrent urinary tract infections caused by large stones.

#### **Alternate Names:**

Extracorporeal Shock Wave Lithotripsy; ESWL; Laser Lithotripsy; Stone Fragmentation

### **Lithotripsy Probe**

A device that fragments a stone using mechanical, ultrasonic, laser, or shock-wave energy during a stone-removal procedure.

### **Lithotripter**

A device that breaks urinary, bile-duct, or other

stones into smaller pieces using shock waves, laser energy, or another treatment method.

### **Lithotripter Probe**

A specialized probe delivering mechanical, electrical, or laser energy to break urinary or bile-duct stones into fragments that can be removed.

### **Litmus**

A dye that changes color with acidity or alkalinity and is used as a simple pH indicator.

### **Little Area**

The highly vascular region of the anterior nasal septum where several arteries meet and where most anterior nosebleeds originate.

### **Little's Disease**

An older name for spastic cerebral palsy, a condition caused by early brain injury or abnormal development that produces stiff muscles and movement difficulties, mainly in the legs. It is not progressive, although needs change over time.

### **Live Attenuated Vaccines**

Vaccines containing weakened forms of viruses or bacteria. They usually produce a strong, long-lasting immune response without causing illness in healthy people, but some are unsuitable for people with severely weakened immune systems or during certain pregnancies.

#### **Alternate Names:**

Live Vaccines

### **Live Birth**

A live birth is delivery of an infant showing signs of life such as breathing, heartbeat, or movement, regardless of gestational age.

### **Livedoid Vasculopathy**

A chronic occlusive disorder of small dermal vessels causing painful purpuric lesions, ulcers, and porcelain-white stellate scars, usually on the lower legs and ankles.



---

## Livedo Racemosa

A persistent, irregular, branching net-like pattern of reddish-blue skin caused by disease of small or medium blood vessels. It may occur with autoimmune, clotting, inflammatory, or other systemic disorders and needs evaluation.

## Livedo Reticularis

A net-like, violaceous discoloration of the skin caused by altered blood flow in small cutaneous vessels. It may be temporary and cold-induced or associated with vascular, autoimmune, infectious, or clotting disorders.

## Liver

The largest internal organ, beneath the right ribs. It processes nutrients, makes bile to help digest fats, stores energy, helps regulate blood chemicals, and breaks down medicines and harmful substances.

## Liver Abscess

A pocket of pus in the liver, usually caused by bacteria or an amoeba. It may cause fever, chills, or pain in the upper-right abdomen. Treatment depends on the cause and size and may include antimicrobial medicines and drainage.

## Liver Anatomy and Physiology

Alternate name; see *Liver*.

## Liver Biopsy

Percutaneous, transjugular, or laparoscopic sampling of liver tissue for histologic evaluation when noninvasive tests do not answer the clinical question. It may stage fibrosis or characterize inflammation, fatty liver disease, infiltrative disease, or a mass; complications include pain and bleeding.

## Liver Cancer

A malignant tumor involving the liver. Primary liver cancers begin in liver cells or bile-duct cells, with hepatocellular carcinoma the most common primary type in adults. Cancer that begins elsewhere and spreads to the liver is metastatic liver cancer and is more common than primary liver cancer. Risk factors for primary liver cancer include cirrhosis,

chronic viral hepatitis, alcohol-related liver injury, fatty-liver inflammation, and some toxins.

### Alternate Names:

Hepatic Malignancy

## Liver Circulation

The liver receives oxygen-rich blood from the hepatic artery and nutrient-rich blood from the portal vein. Blood passes through liver tissue and leaves through the hepatic veins to the heart.

## Liver Cirrhosis

Alternate name; see *Cirrhosis*.

## Liver Damage

Liver damage is injury or loss of liver-cell function caused by infection, alcohol, medicines, toxins, fatty disease, autoimmune disease, or other processes.

## Liver Disease

Any condition damaging or impairing liver structure or function. Causes include infections, alcohol, medicines, toxins, fat buildup, immune disease, inherited disorders, blocked bile flow, and cancer.

## Liver Disease and Pregnancy

Liver disorders may begin during pregnancy or predate it and can affect maternal health, fetal growth, medication choices, and timing of delivery. Abnormal liver tests, jaundice, severe itching, right-upper-abdominal pain, vomiting, or high blood pressure require prompt assessment.

## Liver Disease Diet

An individualized eating plan for a person with liver disease. It may address adequate energy and protein, sodium or fluid intake, alcohol avoidance, and nutrient deficiencies according to the condition.

## Liver Diseases

Disorders of the liver, including hepatitis, fatty liver, cirrhosis, tumors, and biliary disease, that can impair metabolism and detoxification.

---

## Liver Disorder

Any disease affecting the liver, including fatty liver, viral hepatitis, autoimmune disease, scarring, blockage of bile flow, or cancer. Symptoms may be absent or include jaundice, fatigue, pain, swelling, or abnormal blood tests.

## Liver Dysfunction

Liver dysfunction is reduced ability of the liver to perform tasks such as processing nutrients, making proteins, clearing toxins, and producing bile.

## Liver Elastography

A noninvasive ultrasound- or MRI-based test that estimates liver stiffness. It helps assess scarring (fibrosis) and cirrhosis, but stiffness can also rise from inflammation, congestion, or other conditions and must be interpreted with other findings.

## Liver Enzymes

Blood measurements including ALT, AST, alkaline phosphatase, and GGT that can suggest liver or bile-duct injury. Abnormal results require interpretation with symptoms, medicines, imaging, and other tests.

## Liver Failure

A life-threatening loss of the liver's ability to make proteins, process nutrients, remove harmful substances, and regulate blood clotting. It can cause jaundice, bleeding, confusion, swelling, and multiple-organ failure.

## Liver Fibrosis

Excessive scar formation in the liver after ongoing injury or inflammation. It can progress to cirrhosis, reduce liver function, and result from viral hepatitis, alcohol, fatty liver, immune disease, or other causes.

## Liver Fluke

A parasitic flatworm that infects the liver or bile ducts, including *Clonorchis*, *Opisthorchis*, and *Fasciola* species.

## Liver Function

Alternate name; see *Liver Function Tests*.

## Liver Function Tests

Blood tests used to assess liver injury, bile-flow abnormalities, and liver synthetic function. They commonly include aminotransferases, alkaline phosphatase, bilirubin, albumin, and clotting tests; results must be interpreted together with the clinical context.

## Liver Health

The ability of the liver to perform its many jobs, including processing nutrients and medicines, making important blood proteins, producing bile, storing energy, and helping remove harmful substances. Alcohol, infections, obesity, medicines, and other diseases can damage the liver.

## Liver Infection

Infection of liver tissue, such as a viral hepatitis infection or a bacterial liver abscess. Symptoms may include fever, tiredness, nausea, right-upper-abdominal pain, jaundice, or confusion and require medical evaluation.

## Liver Lobule

A small repeating unit of liver tissue in which blood from branches of the portal vein and hepatic artery flows past liver cells toward a central vein, while bile flows in the opposite direction.

## Liver Mass

An unusual area found in the liver, often on an ultrasound, CT scan, or MRI. It may be a harmless cyst or growth, a primary liver cancer, or cancer that spread from elsewhere; further testing may be needed.

## Liver Metastases

Liver metastases are cancer deposits that spread to the liver from another primary tumor and may cause no symptoms or produce pain, weight loss, or liver dysfunction.

## Secondaries In The Liver

Informal name for cancer that has spread to the liver from another part of the body; see *Liver Metastases*.

---

## Liver Neoplasm

A liver neoplasm is an abnormal growth in the liver that may be benign or malignant and may arise from liver cells, bile ducts, or metastasis from another cancer.

## Liver Pain

Discomfort in the upper-right abdomen that may arise from the liver, gallbladder, bile ducts, stomach, muscles, or other organs. Persistent pain, jaundice, fever, vomiting, or confusion may indicate an underlying illness.

## Liver Regeneration

The liver's ability to restore lost tissue by growing and reorganizing remaining cells after injury or partial surgical removal.

## Liver Scan

A nuclear medicine or imaging examination of the liver used to evaluate its size, shape, tissue distribution, focal lesions, blood flow, or selected functional abnormalities.

## Liver Shunt

An abnormal or surgically created channel that diverts blood around the liver, such as a portosystemic shunt; it can reduce detoxification and contribute to hepatic encephalopathy.

## Liver Spots

Alternate name; see *Age Spots*.

## Liver Transplant

Surgical replacement of a diseased liver with a healthy liver or part of a liver from a donor.

## Liver Transplantation

Surgery to replace a severely diseased or failing liver with a healthy donor liver or part of one. It may be used for advanced liver disease, sudden liver failure, or selected liver cancers; lifelong medicines help prevent rejection.

## Living Donor

Individual who donates an organ (kidney, liver segment, or other tissue) while alive for transplantation into a recipient.

## Living Will

A type of advance directive that records a person's preferences for future medical treatment if the person cannot communicate or decide.

## Livor Mortis

Post-mortem purplish-red discoloration of dependent body parts resulting from gravitational settling of blood in uncompressed capillaries and small veins following cessation of circulation.

### Alternate Names:

Hypostasis; Postmortem Staining

## Lizard Bite

A puncture or tissue injury caused by a lizard; care includes cleaning and assessing for infection, tetanus, venom exposure, and significant tissue damage according to the species and circumstances.

## LLETZ

A procedure that uses a thin electrically heated wire loop to remove abnormal tissue from the cervix. The tissue is examined in the laboratory and the procedure can treat precancerous cervical changes.

### Alternate Names:

LEEP (loop Electrosurgical Excision Procedure)

## Lloyd-Davies Position

A modified lithotomy position in which the patient lies supine with lower-profile leg support (less hip flexion) combined with a Trendelenburg tilt. It is commonly used for pelvic and colorectal surgery, allowing simultaneous abdominal and perineal surgical access.

## Loading Dose

The first, sometimes larger, dose of a medicine given to reach a helpful amount in the body quickly. It is followed by regular maintenance doses and is selected according to the medicine and the patient's

---

condition.

### **Loa Loa**

The African eye worm, a filarial nematode transmitted by Chrysops deer flies, endemic to the rainforests of West and Central Africa. Adult worms migrate subcutaneously, causing Calabar swellings, and may cross the subconjunctiva of the eye, causing visible worm migration. Microfilariae are detected in daytime peripheral blood.

### **Lobaplatin**

Lobaplatin is a platinum-based antineoplastic drug used in some countries and studied for its ability to damage cancer-cell DNA.

### **Lobar Hemorrhage**

Bleeding into one of the large regions, or lobes, of the brain. It can result from long-term high blood pressure, fragile blood vessels, blood-thinning medicines, injury, or other conditions and may cause sudden headache, weakness, confusion, seizures, or loss of consciousness.

### **Lobar Pneumonia**

Pneumonia affecting a large section or an entire lobe of the lung. It can cause fever, cough, chest pain, breathlessness, and low oxygen and may be caused by bacteria, viruses, or other organisms.

### **Lobe**

A lobe is a distinct subdivision of an organ or structure, such as a lung, liver, brain, thyroid, or gland.

### **Lobectomy**

Surgical removal of a lobe of an organ. Most commonly lung lobectomy (removal of one lobe of the lung, performed for lung cancer, bronchiectasis, abscess, or benign tumor) or temporal lobectomy (removal of the anterior temporal lobe for drug-resistant temporal lobe epilepsy).

### **Lobotomy**

A historical brain operation that cut or damaged nerve connections in the front part of the brain. It

was once used for severe mental illness but caused serious changes in personality and ability and is no longer an accepted treatment.

### **Lobular Carcinoma**

Lobular carcinoma is a cancer arising from cells of a breast lobule, the gland that produces milk; it may be invasive or confined to the lobule.

### **Lobular Carcinoma In Situ**

Abnormal cells confined to the breast lobules that increase the risk of developing invasive breast cancer.

### **Lobule**

A lobule is a small subdivision of an organ or tissue, such as a liver lobule, lung lobule, or breast lobule.

### **Lobules**

Small sections that make up an organ. In the breast, lobules are groups of glands that can make milk; in other organs, the meaning depends on the organ's structure and function.

### **Local**

Limited to a particular body site or region rather than involving the whole body. A local treatment or symptom affects the treated or affected area, although nearby or wider effects can sometimes occur.

### **Local Anesthesia**

Temporary loss of sensation, especially pain, in a small, specific body area caused by a local anesthetic applied to the surface or injected nearby. The person usually remains conscious, although sedation may be added; excessive dosing or accidental injection into a blood vessel can cause seizures, heart problems, or other toxicity.

#### **Alternate Names:**

Local Anaesthesia

### **Local Anesthetic Systemic Toxicity**

A dangerous reaction when a local numbing medicine enters the bloodstream in excessive

---

amounts. Early signs include tingling around the mouth, ringing in the ears, dizziness, or agitation; seizures and heart problems can follow.

### **Local Anesthetic Toxicity**

Harm from too much local anesthetic in the bloodstream. Early warning signs include ringing in the ears, numbness around the mouth, dizziness, or agitation; seizures, low blood pressure, abnormal rhythms, or cardiac arrest can follow and require emergency treatment.

### **Local Invasion**

Growth of a tumor into nearby tissues rather than only at its original site. It is a feature of malignant behavior and differs from distant metastasis.

### **Localised**

Confined to a limited area or a small number of adjacent body sites rather than widespread.

### **Localization**

Determining the body site responsible for a symptom, sign, lesion, or function. Accurate localization helps guide examination, imaging, diagnosis, treatment, surgery, rehabilitation, and assessment of nerve or organ damage.

### **Localized**

Limited to one particular area, organ, or body site without evidence of distant spread. A localized disease may still be serious, but treatment can sometimes focus on the affected area.

#### **Alternate Names:**

Localized Disease

### **Localized Edema**

Swelling confined to a particular body region because of fluid accumulation, commonly caused by venous or lymphatic obstruction, inflammation, trauma, allergy, or local infection.

### **Localized Lipodystrophy**

Loss of subcutaneous fat in a limited body area, sometimes after injections, trauma, inflammation,

pressure, or panniculitis. It may be cosmetic or reflect an underlying local or systemic disorder.

### **Localized Scleroderma**

An autoimmune skin disorder causing localized thickening and hardening of the skin, with or without involvement of deeper tissues; it differs from systemic sclerosis.

### **Local Therapy**

Local therapy treats a disease at or near the affected area rather than throughout the body, for example with a cream, injection, surgery, or focused radiation.

### **Lochia**

The normal vaginal discharge after childbirth. It is usually red and bloody at first, then becomes pink or brown, and finally yellowish or white over the following weeks. Heavy bleeding, large clots, fever, or foul odor needs assessment.

### **Locked-In Syndrome**

A condition in which a person remains conscious and able to think but is almost completely paralyzed and cannot speak. Some people can communicate by blinking or moving their eyes.

#### **Alternate Names:**

Pseudocoma; De-Efferented State

### **Lockjaw**

A serious neurologic illness caused by tetanospasmin produced by *Clostridium tetani*. It causes painful muscle stiffness and spasms, often beginning in the jaw.

#### **Alternate Names:**

Trismus; *Clostridium Tetani* Infection

### **Locomotion**

Movement of an organism or person from one place to another using coordinated muscles, bones, nerves, and balance systems.

### **Locomotive System**

The musculoskeletal and nervous systems working

---

together to produce posture, balance, and movement from place to place.

### **Locomotor Ataxia**

A late manifestation of untreated tertiary syphilis (neurosyphilis) caused by infection with *Treponema pallidum*, involving progressive degeneration of the dorsal columns (fasciculus gracilis and cuneatus) and dorsal roots of the spinal cord.

#### **Alternate Names:**

Syphilitic Myelopathy; Posterior Sclerosis

### **Locoregional**

Affecting a defined local area and nearby tissues or lymph nodes, without necessarily involving the whole body.

### **Locus**

The specific location of a gene or other DNA sequence on a chromosome. It is described using the chromosome number and a coded position on its short or long arm.

### **Locus Ceruleus**

A small brainstem region containing nerve cells that release norepinephrine. It helps regulate alertness, attention, stress responses, sleep, and some automatic body functions.

#### **Alternate Names:**

Locus Coeruleus

### **Locus Coeruleus**

The locus coeruleus is a brainstem nucleus whose noradrenaline-producing neurons influence arousal, attention, stress, sleep, and pain.

### **Locus Minoris Resistentiae**

Latin for a site of lesser resistance, meaning a body area considered more vulnerable to injury, infection, or disease because of local weakness or prior damage.

### **Lod Score**

A statistical measure used in genetic linkage analysis to compare how likely it is that two genetic markers are inherited together rather than independently.

Higher positive scores provide stronger evidence of linkage in the studied family.

### **Loeffler Endocarditis**

Eosinophil-mediated endomyocardial disease, usually associated with hypereosinophilic syndrome, in which eosinophil infiltration and degranulation cause myocardial or endocardial injury, mural thrombus, and later endocardial fibrosis. It can produce restrictive physiology, valvular regurgitation, emboli, and heart failure; the disease may progress through necrotic, thrombotic, and fibrotic stages.

#### **Alternate Names:**

Eosinophilic Endomyocardial Fibrosis; Fibroplastic Parietal Endocarditis

### **Loeffler Syndrome**

A benign, self-limiting pulmonary syndrome characterized by transient migratory pulmonary infiltrates, peripheral blood eosinophilia, dry cough, and low-grade fever, classically triggered by the transpulmonary larval migration of helminth parasites such as *Ascaris lumbricoides*.

#### **Alternate Names:**

Simple Pulmonary Eosinophilia; Loeffler Pneumonia

### **Loeys-Dietz Syndrome**

Loeys-Dietz syndrome is an inherited connective-tissue disorder associated with arterial aneurysm or dissection, distinctive skeletal and craniofacial features, allergic disease, and variable organ involvement. Lifelong vascular imaging and coordinated specialist care are important.

#### **Alternate Names:**

LDS; TGF-Beta Receptor Vasculopathy

### **Log Roll Test**

A physical examination manoeuvre used to assess hip joint pathology. With the patient supine and the lower extremity in full extension, the examiner passively rolls the thigh internally and externally; pain elicited during rotation indicates intra-articular hip pathology, such as femoral neck fracture, labral tear, or synovitis.



---

## Loiasis

Loiasis is infection with the filarial worm *Loa loa*, transmitted by deerflies and sometimes causing transient eye-worm migration and localized swelling.

## Loin

The area of the lower back between the ribs and pelvis, including the side of the body called the flank. Pain there may come from muscles, the spine, kidneys, or nearby organs.

## Lomotil Overdose

Taking too much diphenoxylate-atropine can cause extreme sleepiness, confusion, dry mouth, fast heart-beat, slowed breathing, coma, or delayed worsening, especially in children.

## Lomustine

Lomustine is an oral chemotherapy medicine used for selected brain tumors and other cancers; delayed low blood counts are an important risk.

## Lonafarnib

A farnesyltransferase-inhibiting medicine used for selected progeroid syndromes, in which abnormal protein processing contributes to premature aging features. It can cause gastrointestinal symptoms, liver abnormalities, and drug interactions.

## Long-Acting Beta-2 Agonists

Inhaled medicines that keep the airways relaxed for many hours and reduce breathing symptoms in asthma or chronic obstructive pulmonary disease. For asthma, they are used with an inhaled corticosteroid, not alone.

## Long-Acting Reversible Contraception

Extended-release contraceptive modalities that provide highly effective pregnancy prevention over months to years without requiring daily patient adherence, including subdermal progestin implants, copper and levonorgestrel intrauterine devices (IUDs), and injectable depot medroxyprogesterone acetate.

**Alternate Names:**

LARC; Long-Acting Contraception; Extended-Release Contraception

## Long Arm Of A Chromosome

The longer segment of a chromosome, designated q, extending from the centromere toward the chromosome's end.

## Long Bones

Bones that are longer than they are wide, such as the femur, tibia, humerus, and radius. Their shafts provide support and leverage for movement, while their ends form or help form joints; bone marrow inside them produces blood cells.

## Long COVID

Ongoing or new health problems that begin or continue after COVID-19. Symptoms may include tiredness, worsening after activity, difficulty thinking, breathlessness, changes in smell or taste, sleep problems, or effects on several organs.

## Longitudinal Axis

An imaginary line running along the length of a body or body part, from one end to the other. Rotation around this line is called longitudinal rotation.

## Longitudinal Section

A cut or image made along the long axis of an organ, limb, vessel, or other structure.

## Longitudinal Studies

Longitudinal studies follow participants or measurements over time to examine change, incidence, exposures, outcomes, and temporal relationships.

## Longitudinal Vaginal Septum

A wall of tissue that divides the vagina lengthwise, present from birth. It may cause painful sex, difficulty inserting tampons, blocked menstrual flow, or no symptoms. It can occur with an unusually shaped uterus and may be removed surgically when it causes problems.

---

## Long QT Syndrome

A heart rhythm disorder in which the heart takes too long to reset between beats. It may be inherited or caused by medicines or abnormal body salts and can lead to fainting, dangerous rhythms, or sudden death.

## Long-Sightedness

Hyperopia, a refractive error in which distant objects may be clearer than near objects and accommodation is required for focusing.

## Long-Term Care

Ongoing help with daily activities, nursing, medical needs, or supervision for a person who cannot live fully independently. Care may be provided at home, in assisted living, or in a nursing facility and should match changing needs.

## Long-Term Care Facility

A residential or care setting providing ongoing assistance, nursing, rehabilitation, or personal care to people who cannot live independently or need sustained support.

## Long-Term Complications Of Diabetes

Chronic complications of diabetes caused mainly by vascular and metabolic injury, including retinopathy, kidney disease, neuropathy, cardiovascular disease, and foot disease.

## Long-Term Effect

A health effect that develops or persists after prolonged exposure, treatment, or disease. It may be beneficial, harmful, or uncertain.

## Long-Term Memory

Information retained for a long period, from hours or days to a lifetime. It includes explicit memories of facts and events and implicit skills or habits.

## Long-Term Potentiation

Long-term potentiation is a lasting increase in the strength of communication between nerve cells after repeated stimulation and is important in learning

and memory.

## Long-Term Rehabilitation

Ongoing rehabilitation for chronic disease, lasting disability, or persistent injury. Goals include maintaining function, preventing complications, adapting activities, and supporting independence and quality of life.

## Loop Diuretic

A strong water tablet that makes the kidneys remove extra salt and water in urine. It is used for fluid buildup, heart failure, some kidney problems, and high blood pressure; dehydration and low body salts can occur.

## Loop Diuretics

A group of strong water tablets that make the kidneys remove salt and water quickly. They treat fluid buildup and some heart, kidney, or liver disorders; excessive use can cause dehydration, low salts, low blood pressure, or kidney problems.

## Loop Electrosurgical Excision Procedure

A procedure that uses a thin heated wire loop to remove abnormal tissue from the cervix. It both treats precancerous cervical changes and provides tissue for laboratory examination. Bleeding, infection, narrowing of the cervix, and a small increased risk of later premature birth can occur.

### Alternate Names:

LEEP; Large Loop Excision of the Transformation Zone

## Loop Of Henle

A U-shaped part of each kidney's tiny filtering tube. It helps the kidney conserve water and control the concentration of urine by allowing water and salts to move differently along its two limbs.

## Loose Bodies

Free fragments of cartilage, bone, or other tissue within a joint or body cavity that may cause pain, catching, or locking.

---

## Loose Body

A free-floating intra-articular fragment of bone, calcified hyaline cartilage, or fibrous tissue within a synovial joint space ('joint mouse'), arising from osteochondritis dissecans, osteoarthritis, or trauma, capable of causing abrupt painful mechanical joint locking.

### Alternate Names:

Joint Mouse; Intra-Articular Loose Fragment

## Loperamide

Loperamide is a medicine that slows intestinal movement and reduces diarrhea. It may be unsafe in bloody or fever-related diarrhea, and excessive doses can cause severe abnormal heart rhythms.

## Loperamide Hydrochloride

Loperamide hydrochloride is an antidiarrheal medicine that slows movement through the intestine; excessive doses can cause dangerous heart rhythms.

## Lopinavir

Lopinavir is an HIV protease inhibitor used with ritonavir in antiretroviral treatment; drug interactions are common.

## Loratadine

An antihistamine used to relieve sneezing, runny nose, itchy or watery eyes, and hives. It usually causes less drowsiness than older antihistamines, but sleepiness, headache, or dry mouth can still occur.

## Lorazepam

Lorazepam is a benzodiazepine used short term for severe anxiety, seizures, or sedation; it can cause drowsiness, dependence, and breathing suppression.

## Lordosis

The inward curve of the neck or lower back. A curve that is unusually pronounced may be related to posture, pregnancy, muscle imbalance, spinal changes, or another condition and may cause back discomfort.

## Losartan

Losartan is an angiotensin-receptor blocker used for high blood pressure, some kidney disease, and heart failure; it can raise potassium levels.

## Losartan Potassium

Losartan potassium is the salt form of losartan, an angiotensin-receptor blocker used to lower blood pressure and protect selected patients with kidney or heart disease.

## Loss Of Appetite

Reduced desire to eat. It may result from illness, medicines, mood disorders, pain, or other causes.

## Loss Of Consciousness

A temporary loss of awareness and responsiveness. It may result from fainting, seizure, head injury, low blood sugar, intoxication, or other serious conditions and requires assessment when unexplained, prolonged, or associated with injury.

## Loss of Bladder Control

Alternate name; see *Urinary Incontinence*.

## Loss-Of-Function Mutation

A genetic change that reduces or abolishes the normal activity or production of a gene product; its clinical effect depends on the gene, variant, and inheritance pattern.

## Loss-Of-Function Variant

A genetic variant that reduces or abolishes the normal activity of the encoded gene product. Disease risk depends on the gene, inheritance pattern, residual function, and whether one or both copies are affected.

## Loss Of Heterozygosity

Loss of heterozygosity is loss of one of two different versions of a gene in a cell, a change that can contribute to cancer when a tumor-suppressor gene is affected.

---

### Loss of Smell

Partial or complete inability to detect odors, commonly caused by nasal obstruction, infection, inflammation, head injury, neurologic disease, or certain medicines.

### Loteprednol Etabonate

Loteprednol etabonate is a corticosteroid eye medicine used for eye inflammation; prolonged use can raise eye pressure and increase infection risk.

### Lou Gehrig Disease

Amyotrophic lateral sclerosis, a progressive neurodegenerative disease affecting upper and lower motor neurons and causing weakness, muscle wasting, and eventual respiratory impairment.

### Lou Gehrig's Disease

Alternate name; see *Amyotrophic Lateral Sclerosis*.

### Louse

Singular form; see *Lice*.

### Lovastatin

Lovastatin is a statin medicine that lowers LDL cholesterol and reduces cardiovascular risk; muscle injury and liver effects are uncommon but important risks.

### Low Back Pain

Pain or discomfort in the lower back, commonly caused by muscle or ligament strain, disc disease, arthritis, or spinal nerve irritation. Weakness, numbness, new bladder or bowel problems, fever, major trauma, or unexplained weight loss are warning signs requiring prompt medical assessment.

#### Alternate Names:

Lumbago; LBP; Lower Back Strain

### Low Birth Weight

A birth weight below 2,500 grams (about 5 pounds 8 ounces), whether caused by being born early, slower growth before birth, or both. Lower weights carry greater risks and need careful newborn monitoring.

### Low Blood Oxygen

An abnormally low oxygen level in arterial blood, usually related to lung disease, impaired breathing, circulation problems, or reduced oxygen availability.

### Low Blood Potassium

Low blood potassium, or hypokalemia, can cause weakness, cramps, constipation, abnormal heart rhythms, or paralysis and may result from losses, medicines, or hormonal disease.

### Low Blood Pressure

A systolic blood pressure below 90 mmHg or diastolic blood pressure below 60 mmHg, resulting in inadequate end-organ perfusion. Etiologies include hypovolemia, septic shock, cardiogenic impairment, adrenal insufficiency, orthostatic autonomic failure, and antihypertensive overdosage.

#### Alternate Names:

Low Blood Pressure; Hypotension; Orthostatic Hypotension

### Low Blood Pressure Causes

Alternate name; see *Hypotension*.

### Low Blood Sodium

Low blood sodium, or hyponatremia, reflects excess water relative to sodium and can cause headache, confusion, seizures, or coma when severe or rapidly developing.

### Low Blood Sugar

Low blood sugar, or hypoglycemia, occurs when circulating glucose is insufficient for normal brain and body function, causing sweating, tremor, confusion, seizures, or loss of consciousness.

### Low Body Temperature

Alternate name; see *Hypothermia*.

### Low-Calorie Diet

An eating plan that provides less energy than a person usually uses, often to support weight loss or a medical goal. It should still provide adequate protein, vitamins, minerals, fluids, and fiber.

---

**Alternate Names:**

Calorie-Restricted Diet

**Low-Density Lipoprotein**

A particle that carries cholesterol through the blood. Too much LDL cholesterol can collect in artery walls, forming fatty deposits that narrow arteries and increase the risk of heart attack and stroke.

**Low-Dose CT Scan**

A CT scan performed with less radiation than a standard scan. It is used for purposes such as lung-cancer screening or follow-up of lung nodules, but abnormal findings may require additional tests.

**Lower Airway Obstruction**

Narrowing or blockage of the breathing tubes inside the lungs. Asthma, chronic lung disease, infection, swelling, or a foreign object may cause wheezing, cough, breathlessness, or reduced airflow.

**Lower Esophageal Ring**

A thin ring of tissue near the junction of the esophagus and stomach, often called a Schatzki ring. It can narrow the esophagus and cause intermittent difficulty swallowing solid food or food becoming stuck.

**Lower Esophageal Sphincter**

A ring of muscle where the esophagus joins the stomach. It opens during swallowing and normally closes afterward to limit backward flow of stomach contents and acid.

**Alternate Names:**

LES

**Lower Extremity**

The lower extremity is the hip, thigh, knee, leg, ankle, and foot, including their bones, joints, muscles, nerves, vessels, and skin.

**Lower Extremity Amputation Dressing Care**

Systematic wound care, sterile dressing changes, and residual limb inspection following lower extrem-

ity amputation. Aims to promote healing, manage exudate, monitor for surgical site infection or flap ischemia, and prepare the residual limb with compression wrapping for prosthetic fitting.

**Alternate Names:**

Stump Dressing Change; Residual Limb Care; Amputation Wound Management

**Lower Extremity Arterial Bypass**

A vascular surgical bypass operation (such as femoropopliteal or femorotibial bypass) utilizing an autologous saphenous vein or synthetic graft to detour arterial blood flow around a hemodynamically significant occlusion in the peripheral arteries of the leg.

**Alternate Names:**

Fem-Pop Bypass; Leg Arterial Bypass; Peripheral Vascular Bypass Graft

**Lower Extremity Peripheral Artery Disease**

Atherosclerotic stenosis or occlusion of the arterial vascular tree supplying the lower extremities, leading to intermittent claudication, rest pain, non-healing ischemic ulcerations, and critical limb-threatening ischemia.

**Alternate Names:**

Lower Extremity PAD; Leg Arteriosclerosis Obliterans; Peripheral Vascular Disease of Legs

**Lower Extremity Venogram**

Contrast imaging of the veins of a leg, used selectively to evaluate suspected deep-vein thrombosis or venous obstruction when noninvasive imaging is inadequate.

**Lower Extremity Venous Doppler**

An ultrasound test of the leg veins that shows blood flow and whether the veins compress normally. It is commonly used to look for a deep-vein blood clot.

**Lower Gastrointestinal Bleeding**

Bleeding that originates in the lower digestive tract, including the small intestine, colon, rectum, or anus. It most often appears as bright red or dark maroon blood passed with bowel movements, or as hidden

---

blood leading to anemia; common causes include diverticular bleeding, hemorrhoids, polyps, colon cancer, and colitis. Heavy bleeding, dizziness, or fainting may occur.

**Alternate Names:**

Lower GI bleeding; LGIB; Lower gastrointestinal hemorrhage; Hematochezia.

**Lower Leg**

The part of the limb between the knee and ankle, containing the tibia, fibula, muscles, nerves, and blood vessels.

**Lower Motor Neuron Syndrome**

A pattern of weakness caused by damage to nerves that carry movement signals from the spinal cord or brainstem to muscles. It may cause muscle wasting, twitching, low muscle tone, and reduced reflexes.

**Lower Segment Cesarean Section**

Alternate name; see *Cesarean Section*.

**Alternate Names:**

LSCS; Lower Segment Cæsarian Section

**Lower Urinary Tract Symptoms**

Problems with storing or passing urine, including urgency, frequency, nighttime urination, weak stream, straining, leakage, or a feeling of incomplete emptying. Causes include infection, prostate enlargement, bladder disorders, nerve disease, medicines, and blockage.

**Lowe's Syndrome**

A rare inherited disorder, also called oculocerebrorenal syndrome, that mainly affects the eyes, brain, and kidneys. Features may include congenital cataracts, developmental difficulties, low muscle tone, and kidney-related loss of minerals.

**Low Eye Tension**

Alternate name; see *Ocular Hypotony*.

**Low-Fiber Diet**

An eating plan that limits foods such as whole grains, nuts, seeds, raw vegetables, and fruit skins. It may

temporarily reduce stool volume for selected bowel problems or medical procedures.

**Low-Flow Low-Gradient Aortic Stenosis**

A severe narrowing of the aortic valve in which less blood moves through the valve and the pressure difference is lower than expected. Echocardiography and other tests help confirm severity.

**Alternate Names:**

LF-LG Aortic Stenosis; Low-Gradient Severe Aortic Stenosis

**Low-Flow Priapism**

Alternate name; see *Priapism*.

**Low FODMAP Diet**

A temporary eating plan that limits certain poorly absorbed carbohydrates that can cause gas, bloating, pain, or diarrhea in some people with irritable bowel syndrome. Foods are gradually reintroduced with dietitian guidance to identify personal triggers.

**Low Grade**

Describes a disease or tumor that appears less aggressive or more slowly developing than higher-grade forms. The exact meaning and outlook depend on the specific disease and test used.

**Low-Grade Appendiceal Mucinous Neoplasm**

A mucin-producing appendiceal neoplasm with low-grade cytologic atypia that may dilate the appendix and rupture, disseminating mucin into the peritoneum as pseudomyxoma peritonei.

**Low-Grade Astrocytoma**

A usually slow-growing glioma composed of astrocytic cells, often arising in the brain or spinal cord. Symptoms depend on location and may include seizures, headache, weakness, or visual change; molecular classification and grade guide prognosis and care.

**Low-Grade Central Osteosarcoma**

A rare low-grade malignant osteogenic tumor aris-



---

ing within the medullary cavity of bone. It may resemble fibrous dysplasia or a benign fibro-osseous lesion and requires complete oncologic excision.

### **Low-Grade Endometrial Stromal Sarcoma**

A malignant uterine stromal tumor resembling proliferative-phase endometrial stroma, characterized by infiltrative myometrial or vascular growth and often driven by characteristic gene fusions.

### **Low HDL Cholesterol (Hypoalphalipoproteinemia)**

A low concentration of high-density lipoprotein cholesterol, associated with metabolic disease, obesity, smoking, genetic disorders, or other causes. Cardiovascular risk is assessed from the entire lipid profile and clinical context rather than HDL alone.

### **Low Hemoglobin Count**

A hemoglobin level below the expected range, often indicating anemia from blood loss, reduced production, nutritional deficiency, inflammation, or increased destruction of red blood cells.

### **Low LDL Cholesterol (Hypobetalipoproteinemia)**

An unusually low concentration of low-density lipoprotein cholesterol, often related to inherited apolipoprotein B disorders, other genetic traits, malabsorption, illness, or lipid-lowering treatment. The cause and associated liver or nutritional findings guide evaluation.

### **Low Molecular Weight Heparin**

A blood-thinning medicine used to prevent or treat blood clots. It is usually injected under the skin and can cause bleeding, bruising, or, rarely, a serious fall in platelets.

### **Low-Molecular-Weight Heparin**

A group of blood-thinning medicines, including enoxaparin and dalteparin, made from heparin. They reduce clot formation and usually have a more predictable effect than standard heparin, but bleeding and low platelets can occur.

**Alternate Names:**

LMWH

### **Low Nasal Bridge**

A low nasal bridge is reduced height of the nasal root and may be a normal familial feature or part of a congenital, genetic, or developmental syndrome.

### **Low Neutrophil Count**

Alternate name; see *Neutropenia*.

### **Lown-Ganong-Levine Syndrome**

A cardiac pre-excitation syndrome mediated by an accessory atriofascicular bypass tract (James fibers) that bypasses the normal slow conduction of the upper AV node, characterized on ECG by a short PR interval (less than 120 ms) with normal narrow QRS duration and recurrent paroxysmal supraventricular tachycardias.

**Alternate Names:**

LGL Syndrome; Accelerated AV Conduction Syndrome

### **Low-Phosphorus Diet**

A dietary pattern limiting phosphorus intake for selected patients with chronic kidney disease or persistent hyperphosphatemia. It emphasizes avoidance of phosphate additives while maintaining adequate protein and overall nutritional status.

### **Low Platelet Count**

Alternate name; see *Thrombocytopenia*.

### **Low Potassium**

An abnormally low potassium level in the blood, which can cause weakness, cramps, abnormal heart rhythms, or paralysis and may result from losses, medicines, or hormonal disorders.

### **Low-Potassium Diet**

A dietary pattern limiting potassium intake when clinically indicated, usually for hyperkalemia or advanced kidney disease. It must be individualized because unnecessary restriction can reduce diet quality and cause inadequate intake of fruits and vegetables.

---

### Low-Residue Diet

A short-term diet limiting dietary fiber and foods that leave substantial undigested material in the bowel. It may be used in selected gastrointestinal conditions or around certain procedures but is not a universal treatment for inflammatory bowel disease.

### Low-Salt Diet

A low-salt diet limits sodium intake and may help control blood pressure or fluid retention in conditions such as heart failure and kidney disease.

### Low-Set Ear

An ear positioned lower on the side of the head than expected relative to facial landmarks; it may be an isolated variation or a feature of a genetic or developmental condition.

### Low-Set Ears And Pinna Abnormalities

Differences in the position, size, shape, or folds of the external ear, including ears positioned lower than usual. They may occur alone or with a congenital syndrome, so hearing and other physical findings may need assessment.

### Low Sperm Count

A semen condition in which sperm concentration is below the expected reference range. It may reduce the chance of pregnancy, but fertility also depends on sperm movement, shape, ejaculation, and the partner's health.

### Low T

Alternate name; see *Low Testosterone*.

### Low-Tension Glaucoma

Alternate name; see *Normal-Tension Glaucoma*.

### Low Testosterone

A condition in which a person has symptoms of testosterone deficiency and repeatedly low morning testosterone levels. Causes may involve the testes, ovaries, pituitary gland, medicines, obesity, or another illness and need medical evaluation.

**Alternate Names:**

Male Hypogonadism; Testosterone Deficiency

### Low Vision

Visual impairment that cannot be fully corrected with spectacles, contact lenses, medication, or surgery, impairing daily activities. Managed with low-vision aids (magnifiers, telescopes, adaptive devices), lighting, and occupational therapy.

### Low White Blood Cell Count

A reduced number of white blood cells, which can increase susceptibility to infection and may result from medicines, infection, autoimmune disease, nutritional deficiency, or bone marrow disorders.

### Low White Cell Count in Cancer

A reduced number of infection-fighting blood cells in a person with cancer or receiving cancer treatment. The risk of serious infection depends on which cells are low and how severely they are reduced.

### Lozenges

Solid medicinal preparations that dissolve slowly in the mouth, usually to provide local treatment for the throat or oral cavity.

### LRINEC Score

A blood-test score that estimates the chance of a serious soft-tissue infection such as necrotizing fasciitis. It uses inflammation, blood count, sodium, kidney, and blood-sugar results. The score cannot by itself confirm or exclude the infection; severe pain, rapid swelling, or serious illness may occur.

**Alternate Names:**

Laboratory Risk Indicator for Necrotizing Fasciitis; LRINEC

### LSD

Lysergic acid diethylamide is a potent hallucinogen that can cause altered perception, visual hallucinations, autonomic effects, and unpredictable changes in mood and behavior.

### LSD Substance Use

LSD is a potent hallucinogen that alters perception,

---

mood, and thought and can cause panic, accidents, persistent perceptual symptoms, or psychosis in susceptible people.

### **LSD Toxicity**

Hallucinogen poisoning causing perceptual distortion, anxiety, panic, agitation, hallucinations, mydriasis, tachycardia, hypertension, and impaired judgment. Severe agitation, hyperthermia, seizures, trauma, or persistent psychosis can occur; use a calm environment, supportive care, and benzodiazepines when clinically indicated.

### **L-theanine**

An amino acid found mainly in tea leaves. It may influence relaxation and alertness, but supplements can interact with medicines and are not proven treatments for disease.

### **Lubiprostone**

Lubiprostone activates intestinal chloride channels and is used for selected forms of chronic constipation and irritable bowel syndrome with constipation.

### **Lubricant**

A substance that reduces friction between surfaces; medical lubricants are used on skin, mucosa, instruments, or devices and should be selected for tissue compatibility and intended use.

### **Lucilia**

A genus of blowflies; some species can cause wound myiasis, while sterile larvae have occasionally been used in maggot debridement.

### **Ludwig Angina**

A rapidly spreading infection of the floor of the mouth and tissues under the jaw, usually arising from an infected lower molar. Tongue elevation, difficulty swallowing, drooling, and airway obstruction can develop rapidly.

### **Ludwig's Angina**

A rapidly progressive, life-threatening cellulitis of the submandibular, sublingual, and submental

spaces, typically originating from an odontogenic infection of the lower second or third molars.

### **Lugano Classification**

A system used to describe how far Hodgkin or non-Hodgkin lymphoma has spread and how it responds to treatment. It combines examination, blood tests, and imaging, often PET with CT. Stages range from disease in one area to widespread disease; the results help guide treatment and follow-up.

### **Lugol's Iodine**

An aqueous solution of iodine and potassium iodide used as an antiseptic, in selected thyroid treatments, and in some laboratory procedures.

### **Lugol's Solution**

Lugol's solution is a mixture of iodine and potassium iodide used in selected laboratory, diagnostic, and medical applications; it can irritate tissue and affect thyroid function.

### **Lumbago**

A non-specific term for low back pain, typically referring to acute or chronic pain in the lumbar region (between the lower ribs and the gluteal folds). Lumbago is a symptom, not a diagnosis.

### **Lumbar**

Relating to the lower back, especially the region or five vertebrae between the thoracic spine and sacrum.

### **Lumbar Disc Herniation**

Protrusion or extrusion of lumbar disc material through the annulus fibrosus, which may irritate or compress a nerve root and cause back or leg pain. Symptoms, examination findings, and imaging determine its significance and treatment.

### **Lumbar Disk Replacement**

Surgical replacement of a worn or herniated intervertebral disc in the lower back with an artificial prosthetic disc to relieve nerve pain while preserving spinal flexibility.

---

**Alternate Names:**

Artificial Disc Replacement; Total Disc Arthroplasty; Lumbar Arthroplasty

**Lumbar Hernia**

A rare type of abdominal wall hernia in which abdominal or retroperitoneal contents (such as fat, intestine, or kidney) protrude through an anatomical defect in the posterolateral lumbar wall. These defects typically occur in the superior lumbar triangle (Grynfeltt-Lesshaft triangle, the most common site) or the inferior lumbar triangle (Petit's triangle). They may be congenital or acquired following flank surgery (such as kidney surgery), trauma, or severe muscular strain. Symptoms include a soft, reducible swelling in the lower back or flank that enlarges with coughing; surgical repair using mesh is typically indicated to prevent incarceration or bowel obstruction.

**Alternate Names:**

Grynfeltt-Lesshaft Hernia; Petit's Hernia; Flank Hernia; Dorsal Hernia

**Lumbar Lordosis**

The normal anteriorly convex physiological curvature of the lumbar spine. Pathologic hyperlordosis (swayback) or hypolordosis (flatback) may develop secondary to muscular imbalances, obesity, pregnancy, spondylolisthesis, or neuromuscular disorders.

**Alternate Names:**

Lumbar Inward Curvature; Hyperlordosis; Swayback

**Lumbar Lordosis Lower Spine**

Lumbar lordosis is the normal inward curve of the lower spine; excessive or reduced curvature may be associated with posture, muscle imbalance, hip disease, vertebral deformity, or pain. Examination and imaging are guided by symptoms rather than appearance alone.

**Lumbar MRI Scan**

Magnetic resonance imaging of the lumbar spine used to evaluate vertebrae, discs, spinal canal, nerve roots, marrow, and soft tissues in conditions such

as radiculopathy or spinal stenosis.

**Lumbar Plexus**

The lumbar plexus is a network of nerves from the upper lumbar spinal roots that supplies the lower abdomen, pelvis, and parts of the thigh and leg.

**Lumbar Puncture**

A procedure that uses a needle in the lower back to collect cerebrospinal fluid, the liquid around the brain and spinal cord. The fluid can be tested for infection, bleeding, inflammation, or cancer; pressure can also be measured, and some medicines can be given. Clinicians check bleeding risk and possible increased pressure in the skull first. Headache, back pain, bleeding, and infection are possible complications.

**Alternate Names:**

LP; Spinal Tap

**Lumbar Puncture Opening Pressure**

The hydrostatic cerebrospinal fluid pressure measured using a vertical manometer with the patient in the relaxed lateral decubitus position during lumbar puncture, normally ranging between 10 and 20 cm H<sub>2</sub>O in adults; elevated in idiopathic intracranial hypertension and meningitis.

**Lumbar Radiculopathy**

Irritation or pressure on a nerve leaving the lower spine, causing pain that travels into the buttock or leg. It may also cause numbness, tingling, or weakness and is often called sciatica.

**Lumbar Region**

The lumbar region is the lower part of the back between the ribs and pelvis, containing the five lumbar vertebrae.

**Lumbar Spinal Stenosis**

Alternate name; see *Lumbar stenosis*.

**Lumbar Spine**

The lower part of the backbone between the ribs and pelvis, usually made of five vertebrae. It bears much

---

of the body's weight and contains nerves supplying the legs; discs, joints, or nerves there may cause lower-back pain or sciatica.

**Alternate Names:**

L-Spine; Lower Vertebral Column

**Lumbar Spine CT Scan**

Computed tomography of the lumbar spine that provides detailed images of vertebral bone and alignment and is particularly useful for detecting fractures or other bony abnormalities.

**Lumbar Spondylosis**

Alternate name; see *Spondylosis*.

**Lumbar Stenosis**

Narrowing of the spinal canal or nerve openings in the lower back. It can squeeze spinal nerves and cause back pain, leg pain, numbness, weakness, or symptoms that worsen with walking.

**Alternate Names:**

LSS; Lumbar Spinal Stenosis; Neurogenic Claudication

**Lumbar Vertebrae**

The five large bones of the lower spine between the ribs and pelvis. They bear much of the body's weight, allow bending, protect spinal nerves, and can be affected by fractures, arthritis, or disc problems.

**Lumbosacral Radiculopathy**

Pain, sensory change, or weakness caused by irritation or compression of a nerve root in the lower lumbar or sacral spine. Symptoms often radiate from the back or buttock into the leg in a dermatomal pattern.

**Lumbosacral Region**

The lumbosacral region is the junction between the lower lumbar spine and the sacrum at the back of the pelvis.

**Lumbosacral Spine CT**

Computed tomography of the lower spine and sacrum using x-rays and computer reconstruction

to evaluate fractures, bone lesions, alignment, and selected causes of nerve compression.

**Lumbosacral Spine X-Ray**

Plain radiographic imaging of the lumbar spine and sacrum used to evaluate alignment, fractures, degenerative changes, deformity, and some bone lesions.

**Lumbricals**

Small hand or foot muscles arising from tendons of the flexor digitorum profundus or flexor digitorum longus and assisting flexion-extension coordination.

**Lumen**

The inner space of a hollow organ, vessel, or tube.

**Luminescence**

Luminescence is emission of light from a substance through a process other than high-temperature incandescence, such as fluorescence or chemiluminescence.

**Luminescence Type**

Luminescence type identifies the mechanism by which a material emits light, such as fluorescence, phosphorescence, bioluminescence, or chemiluminescence.

**Lumpectomy**

Breast-conserving surgery that removes a cancer or other abnormal area with a small margin of nearby tissue. Radiation or additional treatment may be recommended afterward, depending on the diagnosis and risk of recurrence.

**Lump In The Abdomen**

An abdominal lump may be a hernia, enlarged organ, cyst, stool, inflammation, or tumor; pain, vomiting, fever, growth, or a pulsatile mass requires prompt assessment.

**Lumpy Jaw**

A chronic granulomatous infection of the jaw, usually cervicofacial actinomycosis, causing swelling and sinus formation.

---

## Lunate

A small crescent-shaped bone in the center of the wrist. It helps the wrist move and can be injured by falls; loss of its blood supply can cause Kienböck disease and bone collapse.

## Lunate Bone

A crescent-shaped carpal bone situated in the center of the proximal carpal row between the scaphoid and triquetrum, articulating with the distal radius; vulnerable to avascular necrosis (Kienbock disease) and anterior dislocation into the carpal tunnel.

## Lung

One of the two breathing organs in the chest. The lungs take oxygen into the blood and remove carbon dioxide. The right lung has three sections, and the left has two to leave room for the heart.

## Lung Abscess

A pocket of infection and pus that forms inside damaged lung tissue, often after inhaling stomach contents or because of bacteria. It may cause fever, cough, foul-smelling sputum, chest pain, and breathlessness.

## Lung Anatomy

Alternate name; see *Lungs*.

## Lung Cancer

Cancer that begins in the lungs or airways. The main types are small-cell and non-small-cell cancer. Smoking is the leading risk factor, but radon, asbestos, air pollution, inherited risk, and other exposures can also contribute. Early disease may cause no symptoms.

### Alternate Names:

Bronchogenic Carcinoma; Pulmonary Neoplasm

## Lung Capacity

The amount of air the lungs can hold or move, measured with breathing tests such as spirometry and lung-volume testing. Results help assess conditions including asthma, chronic obstructive pulmonary disease, and restrictive lung disease.

## Lung Collapse

An informal phrase for loss of lung expansion. It may refer to atelectasis, in which part of the lung loses air, or pneumothorax, in which air around the lung compresses it. Use the more specific term when the cause is known.

## Lung Compliance

How easily the lungs stretch when a person breathes in. Stiff lungs require more effort, while overly flexible lungs may not spring back well, as can occur in emphysema.

## Lung Cyst

A lung cyst is a round air- or fluid-containing space in lung tissue, caused by developmental disease, infection, emphysema, inflammation, or another disorder.

## Lung Development

Lung development is embryonic and postnatal formation and maturation of airways, alveoli, blood vessels, and gas-exchange capacity.

## Lung Diffusion Testing

A breathing test that measures how well oxygen-like gas moves from the lung air sacs into the blood. It helps assess diseases such as emphysema, lung scarring, or disorders of the lung blood vessels.

## Lung Disease

Lung disease includes disorders affecting the breathing tubes, air sacs, lung tissue, covering, blood vessels, or chest wall. These conditions may cause cough, breathlessness, chest discomfort, wheezing, or low oxygen.

## Lung Diseases

Conditions affecting the airways, alveoli, pleura, or pulmonary circulation, including infection, asthma, COPD, fibrosis, and cancer.

## Lung Fibroma

A usually noncancerous growth made of fibrous tissue in or near the lung. It often causes no symptoms



---

and is found on imaging, but its appearance, growth, and location may require further imaging or biopsy.

### **Lung Fluke**

A parasitic flatworm, usually of the *Paragonimus* group, acquired by eating undercooked freshwater crustaceans. Infection may cause chronic cough, chest pain, blood in sputum, or lung lesions and can rarely affect the brain.

### **Lung Function Tests**

Physiologic tests that measure aspects of breathing, including airflow, lung volumes, and gas transfer, to assess respiratory function.

### **Lung Herniation**

Protrusion of lung tissue through a weak or torn area of the chest wall, usually after chest trauma or surgery. It may appear as a bulge that enlarges with coughing and can cause pain or breathing difficulty.

### **Lung Injury**

Damage to lung tissue or the pulmonary circulation caused by trauma, infection, inhaled substances, inflammation, aspiration, oxygen toxicity, or other disease.

### **Lung Lavage**

A procedure in which doctors wash parts of the airways or lungs with sterile fluid and then remove it. It may clear certain material from the lungs or collect a sample for testing.

### **Lung Metastases**

Lung metastases are cancer deposits that have spread to the lungs from another primary tumor and may cause cough, breathlessness, chest pain, or no symptoms.

### **Lung Needle Biopsy**

CT- or ultrasound-guided percutaneous sampling of a lung or pleural lesion to evaluate selected masses, infection, or diffuse disease. Pneumothorax and pulmonary hemorrhage are important complications.

### **Lung Neoplasm**

An abnormal growth in the lung or airways that may be noncancerous or cancerous. It may cause cough, blood in sputum, chest pain, breathlessness, or no symptoms; imaging and sometimes tissue sampling determine its nature.

### **Lung Nodule**

A rounded, usually well-defined opacity in the lung measuring up to 3 cm on imaging. It may be benign, infectious, inflammatory, or an early manifestation of lung cancer and often requires follow-up assessment.

### **Lung PET Scan**

Positron-emission tomography, often combined with computed tomography, using a radiotracer to assess metabolic activity in lung nodules, cancer, infection, or inflammation.

### **Lung Plethysmography**

A breathing test performed in a sealed room to measure how much air the lungs hold, including air that cannot be breathed out, and how much resistance the airways provide.

### **Lung Problems And Volcanic Smog**

Volcanic smog contains gases and fine particles that irritate eyes and airways and can worsen asthma or lung disease; limiting exposure and following public-health advice reduce harm.

### **Lung Resection**

Surgery to remove a portion or all of a lung, usually to treat cancer or another localized disease. The amount removed may range from a small wedge or segment to a lobe or an entire lung, depending on the disease and the person's lung function.

### **Lung Protective Ventilation**

A ventilator approach that uses gentle breath sizes and carefully controlled pressures to reduce further lung injury. It is especially important in acute respiratory distress and requires adjustment to oxygen levels, blood gases, and the person's condition.

---

## Lungs

The paired respiratory organs in the chest that exchange oxygen and carbon dioxide between air and blood; they also contribute to acid–base regulation.

### Lungs Design And Purpose

The lungs exchange oxygen and carbon dioxide through branching airways and microscopic alveoli surrounded by capillaries. The airways filter, warm, and humidify air, while respiratory muscles drive ventilation; disease can impair airflow, diffusion, or blood flow.

#### Alternate Names:

Pulmonary Anatomy and Function; Respiratory Mechanics

### Lung Surgery

An operation on the lungs or nearby structures to remove or sample tissue, repair injury, or improve breathing. The procedure depends on the disease and may involve lung resection; bleeding, infection, air leakage, and breathing problems are possible.

### Lung Transplant

Surgery to replace one or both severely diseased lungs with donor lungs. It may be considered for advanced lung failure from conditions such as emphysema, lung scarring, cystic fibrosis, or pulmonary hypertension and requires lifelong immune-suppressing medicines.

### Lung Transplantation

Surgical replacement of a diseased lung with a healthy lung from a donor. Indications include IPF, COPD, CF, PAH, and bronchiectasis. Options include single lung transplant, double lung transplant, or heart-lung transplant. Post-transplant immunosuppression is required lifelong. Complications include rejection and infection.

### Lung Volume Reduction Surgery

An operation that removes the most damaged, over-inflated parts of the lungs in carefully selected people with severe emphysema. This gives healthier lung tissue and the diaphragm more room to work, which may improve breathing and activity. It has

important risks, including air leakage, infection, and respiratory failure.

#### Alternate Names:

LVRs; Reduction Pneumoplasty; Bilateral Lung Volume Reduction

## Lunula

The pale crescent-shaped area at the base of a fingernail or toenail. It is the visible front part of the nail-forming region, although its size and visibility vary normally.

## Lupus

A group of autoimmune diseases in which the immune system attacks the body's own tissues. Systemic lupus erythematosus is one form and may affect the skin, joints, kidneys, blood cells, nervous system, and other organs.

### Lupus Anticoagulant

An antiphospholipid autoantibody directed against phospholipid-protein complexes that paradoxically prolongs in vitro phospholipid-dependent coagulation tests (such as the dilute Russell viper venom time or aPTT) while promoting a potent in vivo hypercoagulable state with arterial and venous thrombosis.

#### Alternate Names:

LA; LAC

### Lupus Band Test

A skin-biopsy test that looks for immune proteins along the junction between the skin's layers. The pattern may support lupus, but it must be interpreted with symptoms and other test results.

#### Alternate Names:

LBT; Dermoepidermal Junction Immunofluorescence

### Lupus Erythematosus

A long-term autoimmune disease in which the immune system attacks the body's own tissues. It can cause tiredness, joint pain, rashes, and inflammation or damage in organs such as the kidneys, lungs, heart, brain, or blood vessels.

---

### **Lupus Miliaris Disseminatus Faciei**

A chronic granulomatous eruption of numerous red-brown or skin-colored papules, usually around the eyelids and central face. It may heal with small depressed scars; diagnosis often requires biopsy to distinguish it from acne, rosacea, sarcoidosis, and other granulomatous disorders.

### **Lupus Nephritis**

Kidney inflammation caused by systemic lupus erythematosus. It can cause blood or protein in the urine, high blood pressure, and reduced kidney function. Classification is based largely on kidney-biopsy findings, and treatment depends on the class, severity, kidney function, and individual risks.

### **Lupus Panniculitis**

A chronic inflammatory form of cutaneous lupus affecting subcutaneous fat, producing firm deep nodules that may heal with lipoatrophy, scarring, or overlying discoid changes.

### **Lupus Pernio**

Persistent red-purple or blue-purple swelling and discoloration of the nose, cheeks, lips, ears, or fingers, usually associated with sarcoidosis. It often indicates longer-lasting disease and may cause scarring or cosmetic changes.

### **Lupus Vulgaris**

A long-lasting form of tuberculosis affecting the skin. It usually causes slowly enlarging reddish-brown patches or plaques that may scar and is treated with a combination of tuberculosis medicines.

### **Luteal Cell**

A cell in the ovarian corpus luteum that produces progesterone after ovulation and supports the uterine lining.

### **Luteal Phase Defect**

A term used when the part of the menstrual cycle after ovulation may not provide enough progesterone or enough time for the uterine lining to support pregnancy. The diagnosis is uncertain and cannot be determined from cycle length alone.

### **Alternate Names:**

Luteal Phase Deficiency; LPD

### **Lutein**

A yellow plant pigment found in leafy green vegetables and concentrated in the retina. Along with related pigments, it helps filter blue light and may protect eye tissues from oxidative damage.

### **Luteinization**

Luteinization is the transformation of an ovarian follicle after ovulation into a corpus luteum that produces progesterone.

### **Luteinizing Hormone**

A hormone released by the pituitary gland. In people with ovaries, a mid-cycle rise triggers ovulation and supports hormone production; in people with testes, it stimulates testosterone production. Blood testing helps assess puberty, fertility, and pituitary or reproductive disorders.

### **Luteinizing Hormone Blood Test**

A blood test measuring luteinizing hormone, which stimulates ovulation and ovarian hormone production in females and testosterone production in males; it helps evaluate puberty, fertility, and pituitary or gonadal disorders.

### **Luteinizing Hormone Deficiency**

Inadequate luteinizing-hormone secretion from hypothalamic or pituitary disease, causing reduced gonadal steroid production, irregular or absent ovulation or menstruation, infertility, or delayed puberty.

### **Luteinizing Hormone-Releasing Hormone**

An older name for gonadotropin-releasing hormone (GnRH), a hypothalamic hormone released in pulses that stimulates the anterior pituitary to secrete luteinizing hormone and follicle-stimulating hormone.

### **Lutembacher Syndrome**

A rare congenital heart defect complex defined by

---

the combination of an atrial septal defect (usually secundum type) and acquired or congenital mitral valve stenosis. The left-to-right atrial shunt partially decompresses the left atrium, reducing pulmonary venous congestion but markedly augmenting right ventricular volume overload and pulmonary arterial blood flow.

**Alternate Names:**

Lutembacher Disease

**Luteolysis**

Luteolysis is the breakdown of the corpus luteum, causing progesterone levels to fall when pregnancy has not occurred.

**Luteoma**

A luteoma is a usually benign ovarian mass of hormone-producing lutein cells, sometimes associated with pregnancy and androgen excess.

**Luteoma Of Pregnancy**

A benign, pregnancy-associated ovarian proliferation of luteinized stromal cells driven by hormonal stimulation. It may cause maternal or fetal androgen excess and usually regresses after delivery.

**Luteotropic Hormone**

A hormone that supports the corpus luteum; the term traditionally refers mainly to prolactin, although usage varies by species and source.

**Lutetium-177**

A radioactive form of lutetium that releases therapeutic radiation and a small amount of detectable radiation. Attached to a targeting molecule, it can deliver treatment to selected neuroendocrine or prostate cancer cells while limiting exposure to other tissues.

**Luxation**

Complete dislocation of a joint, where the bone ends lose normal contact. It can stretch or tear nearby ligaments, nerves, and blood vessels, causing severe pain, deformity, swelling, and inability to move.

**LVAD**

Left-ventricular assist device, an implanted pump helping the heart's main chamber circulate blood. It may support people with severe heart failure while awaiting transplant or as long-term treatment.

**Alternate Names:**

Left-Ventricular Assist Device

**LVNC**

Alternate name; see *Left Ventricular Noncompaction*.

**LVRS**

Alternate name; see *Lung Volume Reduction Surgery*.

**Lyase**

An enzyme that breaks or joins certain chemical bonds without adding water or using oxidation as the main step. Lyases participate in many processes that make or use energy in cells.

**Lyme Disease**

A bacterial infection spread by infected ticks; black-legged ticks transmit it in the United States. Early symptoms may include an expanding rash, fever, tiredness, headache, and muscle or joint aches; untreated infection can later affect nerves, joints, or the heart.

**Alternate Names:**

Lyme Borreliosis; *Borrelia burgdorferi* Infection

**Lyme Disease Blood Test**

A blood test that looks for immune proteins made after infection with Lyme disease bacteria. Results depend on symptoms, tick exposure, and timing; early testing can be negative and a positive result does not always prove active infection.

**Lyme Neuroborreliosis**

Neurological manifestations of dissemination of the spirochete *Borrelia burgdorferi* during early disseminated or late Lyme disease. The classic early disseminated presentation is Bannwarth syndrome, marked by painful radiculoneuritis, cranial neuropathy.

---

thy (frequently bilateral facial nerve palsy), and lymphocytic meningitis with intrathecal antibody production.

**Alternate Names:**

Neurological Lyme Disease; Bannwarth Syndrome

## **Lymph**

A clear fluid collected from body tissues and carried through lymph vessels. It transports immune cells and excess fluid to lymph nodes and eventually returns it to the bloodstream.

## **Lymphadenectomy**

Surgical removal of lymph nodes, either sentinel or regional, for staging or therapeutic purposes; examples include axillary, inguinal, cervical, and retroperitoneal lymph node dissection.

## **Lymphadenitis**

Inflammation or infection of one or more lymph nodes, causing swelling and sometimes tenderness, warmth, or pus. Causes include viruses, bacteria, cat-scratch disease, tuberculosis, and other disorders.

## **Lymphadenopathy**

Enlargement of one or more lymph nodes. It may be localized or widespread and can result from infection, inflammation, immune disease, medicines, or cancer. Persistent, hard, or unexplained nodes can have an important underlying cause.

## **Lymphangiectasia**

Dilatation of lymphatic vessels that can cause clear or blood-tinged vesicles, swelling, or leakage of lymph through the skin or mucosa. It may be primary or follow surgery, radiation, inflammation, infection, or obstruction of lymphatic drainage.

## **Lymphangiectasis**

Abnormal widening of lymph vessels. It may be present from birth or develop later and can cause swelling, leakage of lymph, loss of protein through the intestine, or fluid around the lungs.

## **Lymphangiogenesis**

Formation of new lymph vessels from existing vessels. This can help repair tissue and drain fluid, but it may also contribute to chronic swelling, inflammation, or the spread of cancer.

## **Lymphangiogram**

A lymphangiogram uses contrast imaging to show lymphatic vessels and nodes, helping evaluate obstruction, leakage, malformation, or spread of disease.

## **Lymphangiography**

An imaging test that uses contrast dye to show lymphatic vessels and their flow. It can help find blocked vessels, leaks, or abnormal connections, although newer scans are often preferred.

## **Lymphangioliomyomatosis**

A rare lung disease in which unusual cells grow in the lungs and lymphatic vessels, causing many small cysts. It mainly affects women and can lead to breathlessness, collapsed lung, bleeding, or kidney tumors.

## **Lymphangioma**

A noncancerous birth defect in which lymph vessels form abnormally and become enlarged. It usually appears in infancy or childhood as a soft swelling, often in the neck, armpit, or chest.

## **Lymphangiomatosis**

A rare disorder in which abnormal lymph vessels grow in one or more organs. It may cause swelling, fluid buildup, bleeding, bone damage, or breathing problems, depending on the tissues involved.

## **Lymphangiomyoma**

A rare, usually noncancerous growth made of smooth-muscle-like cells around lymph vessels. It is most often associated with lymphangioliomyomatosis and may contribute to cysts or blockage of lymph flow.

---

### **Lymphangiosarcoma**

A rare, fast-growing cancer arising from the lining of lymph vessels. It may develop in tissue with long-standing lymph swelling and can appear as a bruise-like, purple, or ulcerated skin lesion.

### **Lymphangitis**

Inflammation and infection of lymphatic vessels, usually spreading from an infected skin wound or cellulitis; tender red streaks, fever, and enlarged regional lymph nodes may occur.

### **Lymphatic Basin**

The group of lymph vessels and lymph nodes that drains a particular body area. Doctors use this drainage pattern to examine nearby nodes and assess possible spread of infection or cancer.

### **Lymphatic Capillary**

The smallest type of lymphatic vessel. It begins in body tissues, collects extra tissue fluid, and carries it into larger lymphatic vessels.

### **Lymphatic Circulation**

The movement of lymph from body tissues through lymphatic vessels and nodes back into the bloodstream.

### **Lymphatic Disease**

Any disorder affecting lymph vessels, lymph nodes, or related organs. It may cause swelling, repeated infection, abnormal fluid collections, or cancer and may result from birth defects, injury, infection, inflammation, or tumor growth.

### **Lymphatic Drainage**

The movement of lymph away from a body area through lymphatic vessels and lymph nodes.

### **Lymphatic Filariasis**

A mosquito-borne infection caused by thread-like worms that live in lymph vessels. Repeated infection can block lymph flow and cause long-term swelling of the legs, arms, breasts, or genitals.

### **Lymphatic Malformation**

A birth defect in which lymphatic vessels form abnormally. It may cause soft, fluid-filled spaces in the skin or deeper tissues.

### **Lymphatic Mapping**

Imaging or tracer-guided identification of lymphatic vessels and lymph nodes that drain a body area. It helps plan surgery, locate a sentinel node, investigate lymphedema, or assess possible cancer spread.

### **Lymphatic Obstruction**

Blocked or damaged lymph vessels or nodes that prevent normal fluid drainage. The affected area may develop persistent swelling called lymphedema.

### **Lymphatic System**

A network of lymph, lymphatic vessels, lymph nodes, and lymphoid organs and tissues, including the spleen, thymus, tonsils, and bone marrow. It returns excess tissue fluid to the bloodstream, transports absorbed dietary fats and fat-soluble nutrients, and supports immune-cell development, surveillance, and responses to infection.

### **Lymphatic Tissue**

Lymphatic tissue contains immune cells and structures that filter lymph, trap antigens, and coordinate immune defense.

### **Lymphatic Trunk**

A large lymphatic vessel formed by the joining of smaller lymphatic vessels. It drains a body region and empties into a lymphatic duct.

### **Lymphatic Valve**

A flap inside a lymphatic vessel that helps keep lymph moving in one direction.

### **Lymphatic Vessel**

A lymphatic vessel transports lymph from tissues through lymph nodes and eventually returns fluid to the bloodstream.



---

## Lymphedema

Chronic swelling caused by a buildup of protein-rich lymph fluid when lymphatic vessels or lymph nodes cannot drain it normally. It most often affects an arm or leg, but may also involve the trunk, breast, face, or genitals. The affected area may feel heavy, tight, or uncomfortable and may develop inflammation, thickened skin, reduced movement, or repeated skin infections. Primary lymphedema results from abnormal lymphatic development or inherited changes; secondary lymphedema follows damage or blockage from causes such as surgery, radiation, cancer, infection, or injury.

## Lymph Leakage

Lymph leakage is escape of lymphatic fluid after injury, surgery, or obstruction and may cause swelling, fluid collections, or chylous drainage.

## Lymph Node

A lymph node is a small immune organ that filters lymph and helps activate immune responses to infection, inflammation, or abnormal cells.

## Lymph Node Biopsy

Removal of part or all of a lymph node for microscopic and sometimes microbiologic or molecular examination to evaluate malignancy, infection, or inflammatory disease.

## Lymph Node Culture

Laboratory culture of lymph-node tissue or aspirate to identify bacteria, fungi, or mycobacteria causing lymphadenitis; culture is interpreted with histology and other tests.

## Lymph Node Dissection

Surgery that removes one or more groups of lymph nodes to check for cancer or treat cancer spread. Removed tissue is examined in a laboratory, and the operation can cause swelling or numbness.

## Lymph Node / Gland

A small immune organ along a lymph vessel that filters lymph and helps the body recognize infection, inflammation, or abnormal cells. Nodes are found

in groups in the neck, armpits, chest, abdomen, and groin.

## Lymph Nodes

Small immune organs filtering lymph fluid and helping detect infections or abnormal cells. They may enlarge or become tender with infection, inflammation, immune disease, or cancer spread.

### Alternate Names:

Lymph Glands

## Lymph Node Structure

A lymph node is a small, bean-shaped immune organ with a capsule, spaces that carry lymph, and areas rich in immune cells. It filters lymph and may enlarge during infection, inflammation, or cancer spread.

## Lymphoblast

An immature cell that normally develops into a type of infection-fighting white blood cell. Large numbers of abnormal lymphoblasts can build up in the blood or bone marrow in acute lymphoblastic leukemia.

## Lymphoblast Count

A lymphoblast count measures immature lymphoid cells in blood or bone marrow and may help evaluate or monitor acute lymphoblastic leukemia.

## Lymphoblastic Lymphoma

An aggressive lymphoma of immature B- or T-cell precursors, closely related to acute lymphoblastic leukemia. It often presents with lymph-node, mediastinal, bone, skin, or marrow disease; immunophenotyping and marrow assessment help classify the extent.

## Lymphocele

A lymphocele is a collection of lymphatic fluid caused by disruption or blockage of lymph vessels, often after surgery or transplantation.

## Lymphocyte

A type of white blood cell that recognizes infections

---

and abnormal cells and helps coordinate immune defense. The main types are B cells, which make antibodies, and T cells, which direct or carry out cellular immune responses.

### **Lymphocyte Activation**

The process by which a lymphocyte recognizes a target and begins responding. Activated lymphocytes multiply and help coordinate antibody or cellular immune defense.

### **Lymphocyte Count**

A lymphocyte count measures lymphocytes in the blood and can help assess infection, immune deficiency, inflammation, or blood disorders.

### **Lymphocyte Proliferation**

Rapid multiplication of lymphocytes after immune stimulation. It helps build a response to infection, but abnormal or uncontrolled proliferation can occur in lymphoid cancers.

### **Lymphocytic**

Relating to lymphocytes or containing an increased proportion of these white blood cells. A lymphocytic pattern may occur with viral infection, chronic inflammation, autoimmune disease, or certain cancers.

### **Lymphocytic Choriomeningitis**

A rodent-borne viral infection caused by lymphocytic choriomeningitis virus (LCMV), an arenavirus naturally carried and excreted by the common house mouse (*Mus musculus*). Human infection acquired through inhalation or ingestion of contaminated dust or food typically follows a biphasic course, presenting with an initial flu-like illness followed in severe cases by aseptic meningitis, meningoencephalitis, or congenital infection causing hydrocephalus and chorioretinitis if acquired during pregnancy.

#### **Alternate Names:**

LCMV Infection; Lymphocytic Meningitis

### **Lymphocytic Colitis**

A form of microscopic colitis causing long-lasting, watery diarrhea. The colon may look normal during

examination, but a biopsy shows increased immune cells in its lining; medicines, autoimmune disease, or other facts may contribute.

#### **Alternate Names:**

Microscopic Colitis; IEL-Predominant Colitis

### **Lymphocytic Interstitial Pneumonia**

A rare inflammatory lung disease in which immune cells collect in the tissue around the air sacs. It may cause cough, breathlessness, and abnormal scans and is associated with autoimmune disease or HIV.

### **Lymphocytic Leukemia**

Lymphocytic leukemia is a blood cancer in which abnormal lymphoid cells multiply in the bone marrow, blood, lymph nodes, or other tissues.

### **Lymphocytic Mastitis**

A noncancerous breast lump caused by inflammation and toughening of breast tissue. It is associated with diabetes or autoimmune disease and can look like breast cancer on examination or scans, so testing may be needed.

### **Lymphocytoma Cutis**

Lymphocytoma cutis is a benign skin lymphoid proliferation presenting as red or violaceous plaques or nodules and sometimes associated with infection or immune stimulation.

### **Lymphocytopenia**

An abnormally low number of lymphocytes, a type of white blood cell that helps fight infection. It may be temporary after illness or stress, or result from medicines, immune disorders, infections, poor marrow function, or inherited conditions. Severity and duration determine the risk of infection and need for evaluation.

### **Lymphocytosis**

An abnormally increased number or proportion of lymphocytes in the blood, often associated with viral infection, some bacterial infections, or lymphoproliferative disorders.

---

## Lymphoedema

Alternate spelling; see *Lymphedema*.

## Lymphofollicular Hyperplasia

Lymphofollicular hyperplasia is enlargement or increased formation of lymphoid follicles in response to immune stimulation, infection, inflammation, or other triggers.

## Lymphogranuloma Venereum

A sexually transmitted infection caused by certain types of *Chlamydia*. It may begin with a small painless sore, followed by painful groin swelling or inflammation of the rectum, sometimes with bleeding or discharge.

## Lymphoid

Resembling or relating to lymphatic tissue, lymphocytes, or organs involved in lymphocyte development and immune function.

## Lymphoid Hyperplasia

Lymphoid hyperplasia is benign increase in lymphoid tissue caused by immune stimulation, infection, inflammation, or sometimes a reactive response near a tumor.

## Lymphoid Organs

Organs that contain lymphoid tissue and support the development, storage, or activity of immune cells. They include the bone marrow, thymus, lymph nodes, spleen, and tonsils.

## Lymphoid Tissue

Tissue containing lymphocytes and supporting immune responses, found in lymph nodes, spleen, thymus, tonsils, and mucosal sites.

## Lymphokine

Older term; see *Cytokine*.

## Lymphoma

A cancer of lymphocytes, a type of white blood cell. Hodgkin and non-Hodgkin lymphoma include many subtypes that may grow slowly or quickly and

can cause swollen lymph nodes, fever, night sweats, weight loss, or organ symptoms.

## Lymphoma Overview

Alternate name; see *Lymphoma*.

## Lymphomatoid Granulomatosis

A rare Epstein-Barr-virus-associated lymphoproliferative disorder that commonly involves the lungs and may also affect the skin, central nervous system, or kidneys. It can produce nodules, cough, breathlessness, or systemic symptoms and requires tissue diagnosis and specialist hematology care.

## Lymphomatoid Papulosis

Lymphomatoid papulosis is a chronic, recurring skin disorder with cancer-like immune cells. The bumps often heal on their own, but some people later develop lymphoma.

## Lymphopenia

Lymphopenia is an abnormally low number of lymphocytes, caused by infection, medicines, immune disease, malnutrition, marrow disorders, or other conditions.

## Lymphoplasmacytic Lymphoma

Alternate name; see *Waldenström Macroglobulinemia*.

## Lymphopoiesis

The production and development of lymphocytes, a group of white blood cells that help fight infection and abnormal cells. These cells develop mainly in the bone marrow and thymus.

## Lymphoproliferative Disorder

A condition in which lymphocytes multiply excessively or abnormally. It may be temporary after infection or immune stimulation, or may represent a cancer such as leukemia or lymphoma.

## Lymphoscintigraphy

A scan that follows a small radioactive tracer through lymph vessels. It can map drainage to the

---

first lymph node draining a tumor or show impaired lymph flow in lymphedema and other disorders.

### **Lymph System**

The lymph system includes lymph, lymphatic vessels, lymph nodes, spleen, thymus, and related tissues that return fluid and support immune defense.

### **Lynch Syndrome**

An inherited condition that greatly increases the risk of colorectal, womb, ovarian, stomach, urinary-tract, and several other cancers. Regular screening can detect some cancers early, and relatives may need genetic counseling or testing.

### **Lysergic Acid Diethylamide**

A potent hallucinogenic drug (psychedelic) classified as a Schedule 1 controlled substance (high abuse potential, no accepted medical use). LSD acts as a partial agonist at serotonin 5-HT<sub>2A</sub> receptors in the prefrontal cortex, leading to altered sensory perception, mood, and cognition.

### **Lysine**

Lysine is an essential amino acid needed for protein production, growth, and tissue repair; it must be obtained from food.

### **Lysis**

The breakdown, dissolution, or destruction of cells or tissue; examples include hemolysis, the breakdown of red blood cells, and thrombolysis, the dissolution of a blood clot.

### **Lysogenic Conversion**

A change in a bacterium's characteristics after a virus that infects bacteria inserts its genetic material into the bacterial chromosome and provides new genes, such as genes that increase disease-causing ability.

### **Lysogeny**

Lysogeny is a state in which a bacteriophage's genetic material is maintained inside a bacterial cell and replicated with the host genome.

### **Lysophagy**

A form of selective autophagy that removes damaged lysosomes, the cell compartments that digest waste. It helps maintain cell health; failure of lysophagy can allow damaged lysosomes to accumulate and promote inflammation or cell injury.

### **Lysophosphatidylcholine**

A fat-like molecule found in cell membranes and blood. It helps transport fats and can influence inflammation and cell communication; abnormal levels may occur in metabolic or inflammatory diseases.

### **Lysophospholipids**

Lipids related to phospholipids but containing one fewer fatty-acid chain. They are parts of cell membranes and can act as signaling molecules, influencing inflammation, blood vessels, immune responses, and cell movement.

### **Lysosomal Storage Disorders**

A group of inherited diseases in which missing or faulty lysosomal enzymes cause substances to build up inside cells. Accumulation can damage organs, bones, nerves, or other tissues; treatment depends on the specific disorder.

### **Lysosome**

A small compartment inside a cell containing substances that break down waste, worn-out cell parts, and material brought into the cell. Lysosome disorders can cause harmful material to accumulate in tissues.

### **Lysozyme**

A natural enzyme that helps defend against bacteria by breaking down part of their protective outer wall. It is found in tears, saliva, breast milk, and some immune cells.

### **Lysozyme Test**

A lysozyme test measures the enzyme lysozyme in a sample and may support evaluation of selected blood, inflammatory, or infectious disorders.

---

## Lyssavirus

A group of viruses that includes rabies virus and related viruses. They mainly spread through the bite or saliva of an infected mammal and can cause a severe brain infection. Prompt wound cleaning and exposure-appropriate vaccination can prevent disease before symptoms begin.

## Lytic

Causing or involving lysis, or the breakdown of cells, tissue, or another biological structure; a lytic bone lesion is an area of bone destruction.

## Lytic Lesion

A lytic lesion is an area of bone destruction caused by processes such as cancer, infection, or some metabolic bone disorders.

## Lab Results

Measurements from blood, urine, tissue, or other specimens used with the history and examination to evaluate health, diagnose disease, monitor treatment, or assess risk; an abnormal result is not a diagnosis by itself.

## Labhart-Willi Syndrome

Alternate name; see *Prader-Willi Syndrome*.

## Labor and Delivery

The process of uterine contractions, cervical dilation, birth of the baby, delivery of the placenta, and immediate postpartum care.

## Labor and Delivery Complications

Problems arising during labor, birth, or the immediate postpartum period, including abnormal fetal position, prolonged labor, excessive bleeding, infection, hypertensive emergencies, or fetal distress.

## Labyrinthine Positional Nystagmus

Involuntary rhythmic eye movements triggered by head position or movement because of abnormal vestibular signaling; evaluation distinguishes peripheral inner-ear causes from central neurologic causes.

## LAM

Abbreviation for lymphangioleiomyomatosis, a rare cystic lung disease; see *Lymphangioleiomyomatosis*.

## Laxatives

Medicines or preparations that help relieve constipation by adding bulk, softening stool, drawing water into the bowel, or stimulating bowel movement. The choice depends on the cause and duration of constipation and the person's health.

## Lazy Eye (Amblyopia)

Alternate name; see *Lazy Eye* and *Amblyopia*.

## LCH

Abbreviation for Langerhans cell histiocytosis; see *Langerhans Cell Histiocytosis*.

## LDL Cholesterol

Cholesterol carried in low-density lipoprotein particles; high levels contribute to atherosclerosis and cardiovascular risk. See *LDL*.

## LGMD

Abbreviation for limb-girdle muscular dystrophy; see *Limb-Girdle Muscular Dystrophies*.

## LGS

Abbreviation for Lennox-Gastaut syndrome; see *Lennox-Gastaut Syndrome*.

## Lichen Annularis

Alternate name for granuloma annulare, a benign inflammatory ring-shaped skin eruption; see *Granuloma Annulare*.

## Lichen Planus Sclerosus Atrophicus

Historical or combined name; see *Lichen Sclerosus*.

## Lichen Psoriasis

A descriptive or overlapping term for psoriasis with lichenified, thickened, scaly plaques; the diagnosis should distinguish psoriasis from lichen planus or chronic eczema.

---

## **Lichen Urticatus**

Alternate name for papular urticaria, an itchy hypersensitivity reaction to insect bites; see *Papular Urticaria*.

## **Life Support**

Medical treatments that temporarily replace or support vital body functions, such as mechanical ventilation, dialysis, circulatory support, or artificial nutrition and hydration, when illness prevents adequate function.

## **Lingua Fissurata**

Alternate name for fissured tongue, in which grooves form on the tongue surface; see *Fissured Tongue*.

## **Lingua Nigra**

Alternate name for black hairy tongue, a usually harmless dark or furry coating caused by elongation and discoloration of tongue papillae; see *Black Hairy Tongue*.

## **Linguofacial Dyskinesia**

Involuntary repetitive movements of the tongue, lips, jaw, or face, often related to medication exposure or neurologic disease; evaluation depends on the pattern and cause.

## **Liver Cancer Treatment**

Treatment of liver cancer may include surgery, ablation, embolization, radiation, immunotherapy, targeted therapy, chemotherapy, or transplantation according to tumor type, stage, liver function, and overall health.

## **Localized Epidermolysis Bullosa**

An inherited or acquired blistering disorder limited mainly to particular areas of skin or mucosa; friction can cause blisters, erosions, pain, and infection, and the subtype determines management.

## **Low Back Pain Deciding About MRI**

The decision to obtain lumbar MRI depends on red flags, neurologic findings, duration, treatment response, and whether imaging would change man-

agement; routine early imaging is not useful for uncomplicated low back pain.

## **Low Back Pain Exercises**

Carefully selected movement and strengthening exercises may improve function in nonspecific low back pain; the program should be adapted to symptoms and stopped for progressive weakness, numbness, or other warning signs.

## **Low Potassium Syndrome**

Alternate name for hypokalemia, an abnormally low blood potassium level that can cause weakness, cramps, paralysis, or dangerous heart-rhythm changes; see *Hypokalemia*.

## **Low-Dose Aspirin Therapy**

Regular small-dose aspirin used for selected cardiovascular or obstetric indications. It can cause bleeding and is not appropriate for everyone; primary-prevention use should be based on individualized medical advice.

## **Lumbar Pain**

Pain arising in the lower back, often from muscles, joints, discs, nerves, or other spinal or nonspinal causes; see *Low Back Pain*.

## **Lung Carcinoid Tumors**

Neuroendocrine tumors arising in the lung, ranging from typical slow-growing tumors to atypical tumors with greater malignant potential; see *Carcinoid Tumor*.

## **Lung Disease/Respiratory Problems**

Disorders affecting the airways, lung tissue, pleura, pulmonary blood vessels, or breathing control that may cause cough, shortness of breath, wheezing, chest discomfort, or low oxygen; see *Lung Disease*.

## **Lung Nocardiosis**

Lung infection caused by *Nocardia* bacteria, usually in people with weakened immunity. It can cause cough, fever, chest pain, or nodules and may spread to the brain or other organs; see *Nocardiosis*.



---

## **Lymph Node Cancer**

Cancer involving lymph nodes, either lymphoma arising in lymphatic tissue or cancer spread from another organ; diagnosis requires examination of tissue and staging.

## **Lymphangioleiomyomatosis**

Alternate spelling; see *Lymphangioleiomyomatosis*.



# M

## MacConkey Agar

A laboratory growth medium used mainly to grow and distinguish certain intestinal bacteria. It helps show whether bacteria can break down lactose, a sugar.

## Macerate

To soften and break down tissue after prolonged contact with fluid. The term may describe skin, wounds, or changes in a fetus that has died before delivery.

## Maceration

Softening and breakdown of tissue caused by prolonged exposure to moisture or fluid; it may occur in skin, wounds, or a fetus after death.

## Macewen Sign

An unusually hollow sound heard when part of the skull is tapped. It may occur with enlarged fluid spaces in the brain or rarely a brain abscess, but it cannot diagnose either condition by itself.

## Machupo Virus

An arenavirus that causes Bolivian hemorrhagic fever, a potentially fatal illness characterized by fever, pain, and bleeding; it is endemic in parts of Bolivia.

## Macroamylasemia

Macroamylasemia is a benign condition in which amylase binds to immunoglobulin and remains in blood, causing high serum amylase with low urinary clearance.

## Macroangiopathy

A disease of the body's larger blood vessels, often caused by fatty buildup or a blood clot. It can reduce blood flow to the heart, brain, legs, or other organs.

## Macroautophagy

A cell-cleaning process in which a cell encloses damaged parts or unwanted proteins, breaks them down, and reuses some of their materials. It helps maintain cell health during stress and normal turnover.

## Macrocephalus

Having an abnormally large head, usually defined by head circumference above the expected range for age and sex.

## Macrocephaly

An unusually large head for a person's age and sex. It may run in a family or be linked to fluid buildup, increased brain growth, or another condition.

## Macrocyte

An abnormally large red blood cell, commonly seen in macrocytosis and megaloblastic or other anemias.

## Macrocytic

Describing a cell, especially a red blood cell, that is abnormally large; macrocytic anemia features enlarged red blood cells.

## Macrocytic Anaemia

Alternate British spelling; see *Macrocytic Anemia*.

---

## Macrocytic Anemia

Anemia in which red blood cells are larger than usual. Common causes include vitamin B12 or folate deficiency, alcohol use, liver disease, underactive thyroid, some bone-marrow disorders, and recovery from blood loss. It may cause tiredness, pale skin, shortness of breath, or tingling.

### Alternate Names:

Macrocytosis; Large-Cell Anemia; High-MCV Anemia

## Macrocytosis

An abnormal enlargement of mature circulating red blood cells defined by a mean corpuscular volume (MCV) exceeding 100 fL, commonly resulting from vitamin B12 or folate deficiency (megaloblastic anemia), alcoholism, chronic liver disease, reticulocytosis, or myelodysplastic syndrome.

### Alternate Names:

Macrocytic Erythrocytes; Elevated MCV

## Macroductyly

Fingers or toes that are unusually large because their bones and soft tissues overgrow. It is usually present from birth, may affect one digit or several, and can interfere with movement or footwear.

## Macrodonia

Teeth that are larger than expected. It may affect one tooth or many teeth and can cause crowding, bite problems, or difficulty cleaning; treatment depends on the cause and symptoms.

## Macrogonitosomia

Unusual enlargement of the external genitals caused by excessive hormone activity before birth or during childhood. Possible causes include disorders of the adrenal glands, testes, ovaries, or other hormone systems.

## Macroglobulin

A large protein in the blood, often made of several protein units joined together. Immunoglobulin M is a major example; abnormal amounts can occur in some immune or blood disorders.

## Macroglossia

Macroglossia is an abnormally enlarged tongue that can affect feeding, speech, airway, or dental development and may be congenital or acquired.

## Macrognaia

Abnormal enlargement of the jaw, which may be associated with pituitary gigantism, tumors, or other disorders; also called a prognathic mandible.

## Macrolide

A class of antibiotics that stops bacteria from making proteins. Examples include azithromycin, clarithromycin, and erythromycin; side effects and medicine interactions vary.

## Macrolides

Antibiotics that stop bacteria from making proteins and treat selected infections. Common examples include azithromycin, clarithromycin, and erythromycin.

## Macronutrients

Nutrients needed in large amounts, mainly carbohydrates, proteins, and fats. They provide energy, build and repair tissues, support hormones and cell membranes, and help maintain normal body functions.

## Macrophage

A large immune cell that surrounds and removes germs, dead cells, and debris. It also helps coordinate inflammation and tissue repair.

## Macrophage Activation

The process by which macrophages, a type of immune cell, respond to infection or tissue injury. They may destroy germs and damaged cells, alert other immune cells, and release substances that promote inflammation or repair.

## Macrophage Activation Syndrome

A dangerous complication of some rheumatic diseases in which immune cells become overactive and cause severe inflammation, fever, low blood counts,

---

and organ problems.

### **Macrophage Count**

The number or percentage of macrophages found in a tissue, body fluid, or laboratory sample. The result may help assess infection, inflammation, tissue repair, or a blood disorder, but its meaning depends on the sample and other findings.

### **Macropsia**

A visual disturbance in which objects appear larger than they really are. It may occur with migraine, seizures, retinal problems, medicines, or brain disorders and can temporarily affect depth and distance judgment.

### **Macroscopic**

Large enough to be seen without a microscope. A macroscopic examination records a specimen's visible size, shape, color, surface, and other gross features before microscopic testing.

### **Macroscopic Hematuria**

Visible blood in the urine, also called gross hematuria; it may arise from the kidneys, ureters, bladder, prostate, or urethra and requires evaluation for causes such as infection, stones, trauma, or malignancy.

### **Macrosomia**

Macrosomia means unusually large fetal or newborn size, often associated with diabetes, prolonged pregnancy, parental size, or other factors.

### **Macrostomia**

An abnormally wide mouth or oral opening, usually a congenital facial cleft caused by incomplete fusion of the embryonic facial processes.

### **Macula**

The central part of the retina responsible for sharp, detailed, and color vision. Damage from aging, diabetes, inflammation, or other disease can cause blurred, distorted, or missing central vision.

### **Macula Densa**

A specialized plaque of densely packed columnar epithelial cells located in the wall of the distal convoluted tubule adjacent to the afferent and efferent arterioles of its originating glomerulus; senses luminal sodium chloride concentrations and modulates renin secretion via tubuloglomerular feedback.

### **Macula Lutea**

The small yellowish area in the center of the retina responsible for sharp vision. Its center, called the fovea, provides the clearest vision.

### **Macular**

Relating to a macule, a flat, circumscribed change in skin color without a change in texture or elevation. Macules may be caused by pigment, inflammation, vascular change, or other processes and are distinguished from raised lesions such as papules.

### **Macular Amyloidosis**

A localized cutaneous amyloidosis characterized by mottled or rippled brown-gray patches, usually on the upper back. It is often associated with chronic friction or rubbing and is distinct from systemic amyloidosis.

### **Macular Cyst**

A cystic defect in the macula, the central retinal area responsible for sharp vision, often associated with vitreous traction and potentially causing distortion or loss of central vision.

### **Macular Degeneration**

Damage to the macula, the central part of the retina responsible for sharp, straight-ahead vision. The term most often refers to age-related macular degeneration. In the dry form, the macula becomes thinner and may develop deposits called drusen; advanced dry disease can involve loss of light-sensitive retinal cells and usually progresses gradually. In the wet form, abnormal blood vessels grow beneath the retina and leak blood or fluid, often causing faster loss or distortion of central vision. Straight lines may look wavy, and blank or blurry areas may develop. Peripheral vision is usually preserved, but

---

reading, recognizing faces, and other detailed tasks may become difficult.

### Macular Drusen

Small yellow deposits beneath the retina in the macula, the area responsible for sharp central vision. A few small deposits may occur with aging, while many large deposits can signal increased risk of macular degeneration.

### Macular Edema

Swelling of the macula caused by fluid leaking from retinal blood vessels. It can blur or distort central vision and may result from diabetes, vein blockage, inflammation, surgery, or other eye disease.

### Macular Hole

A small opening in the center of the retina that can cause a missing or distorted area in central vision. Some cases need surgery.

### Macular Pucker

A thin layer of scar-like tissue on the retina that can wrinkle the macula and cause distorted or blurred central vision.

### Macular Skin Lesion

Relating to a macule, a small, flat, circumscribed change in skin color that is neither raised nor depressed.

### Macular Sparing

Retention of central visual acuity and central visual field representation in the presence of an otherwise complete homonymous hemianopia, classically resulting from occlusion of the posterior cerebral artery sparing the occipital pole due to collateral blood flow from the middle cerebral artery or bilateral cortical macular representation.

### Macule

A flat, clearly defined change in skin color that cannot be felt as a raised or depressed area. Macules may be lighter, darker, red, or brown and can have many causes.

### Macules

Small, flat, circumscribed changes in skin color that are neither raised nor depressed; they are distinguished from raised papules and fluid-filled vesicles.

### Maculopapular Lesion

A skin lesion combining flat areas of altered color (macules) with small raised solid lesions (papules); it can occur in infections, drug eruptions, inflammatory diseases, and other conditions.

### Madarosis

Loss or absence of eyelashes, eyebrows, or other hair, caused by local disease, inflammation, trauma, or systemic conditions.

### Mad Cow Disease

Alternate name; see *Bovine Spongiform Encephalopathy*.

### Maddrey Discriminant Function

A validated mathematical prognostic scoring index used in hepatology to assess the severity of acute alcoholic hepatitis and predict short-term mortality. Calculated as  $4.6 \times (\text{patient PT} - \text{control PT in seconds}) + \text{serum total bilirubin in mg/dL}$ ; a score of 32 or greater defines severe acute alcoholic hepatitis carrying an approximate 30- to 50-percent 30-day mortality, identifying patients who benefit significantly from corticosteroid therapy.

#### Alternate Names:

MDF; Maddrey DF Score

### Madelung Deformity

A developmental or post-traumatic wrist deformity resulting from premature physal arrest of the medial and volar aspects of the distal radius, leading to shortening and volar bowing of the radius with prominent dorsal subluxation of the ulnar head.

#### Alternate Names:

Madelung Wrist; Radial Dyschondrosteosis Deformity



---

## Madura Foot

A chronic localized infection of the skin and deeper tissues of the foot, caused by fungi or certain bacteria and characterized by swelling, draining sinuses, and granules.

### Alternate Names:

Mycetoma Pedis; Maduromycosis

## Maduramycosis

Mycetoma, a chronic subcutaneous infection caused by true fungi or filamentous bacteria and characterized by swelling, sinuses, and granules.

## Madurella

Madurella is a genus of fungi that can cause mycetoma, a chronic infection of skin and deeper tissue after implantation through the skin.

## Madurella Mycetomatis

A fungus that commonly causes eumycetoma, a chronic infection producing nodules, draining sinuses, and grains, usually after traumatic implantation in the foot or limb.

## Maffucci Syndrome

A rare condition causing multiple benign cartilage growths and abnormal blood-vessel growths, usually beginning in childhood. It can deform bones, cause fractures, and increase the risk of certain cancers.

## MAG3 Renal Scan

A kidney scan that uses a small radioactive tracer to show how well each kidney works and how urine drains. It can help detect blockage or delayed drainage.

## Magenblase Syndrome

Stomach discomfort and bloating caused by swallowing too much air, often while eating or drinking. The swallowed air may cause burping, fullness, or visible swelling and can be worsened by chewing gum, fizzy drinks, eating quickly, or anxiety.

## Magnesium

A mineral needed for healthy nerves, muscles, bones,

heart rhythm, and many body processes. Too little can cause weakness, tremor, muscle cramps, or abnormal heart rhythms.

## Magnesium Blood Test

A blood test measuring serum magnesium, an electrolyte involved in neuromuscular function, cardiac rhythm, and metabolism; it helps evaluate deficiency, excess, and related electrolyte disorders.

## Magnesium Carbonate

A magnesium salt used in some antacid or laxative preparations and as a magnesium supplement; excessive intake may cause toxicity, especially in renal failure.

## Magnesium Chloride

A magnesium salt used in some oral supplements and medical preparations to provide magnesium or, in concentrated solutions, to help correct magnesium deficiency; dosing depends on the elemental magnesium content.

## Magnesium Deficiency

An abnormally low body magnesium level, often caused by poor intake, gastrointestinal loss, renal wasting, or medications. It may cause weakness, tremor, muscle cramps, seizures, cardiac arrhythmias, hypokalemia, and hypocalcemia.

## Magnesium Hydroxide

An osmotic laxative and antacid that neutralizes gastric acid and draws water into the bowel.

## Magnesium In Diet

Dietary magnesium comes from foods such as nuts, seeds, legumes, whole grains, and leafy vegetables and supports nerve, muscle, bone, and heart function.

## Magnesium Oxide

An inorganic magnesium compound used in some supplements and antacids; it can also have a laxative effect, and excessive use may cause hypermagnesemia, especially with kidney impairment.

---

## **Magnesium Sulfate**

A magnesium salt used intravenously or intramuscularly to treat selected magnesium deficiencies and conditions such as eclampsia, and used orally as an osmotic laxative; it is also known as Epsom salt in its hydrated form.

## **Magnesium Sulfate For Neuroprotection**

Magnesium given through a vein before some very early preterm births to lower the baby's risk of brain injury. The timing and dose are guided by obstetric and newborn-care specialists.

## **Magnesium Sulphate**

A magnesium salt used medically for severe magnesium deficiency, prevention or treatment of eclamptic seizures, and selected arrhythmias or asthma indications.

## **Magnetic Resonance Angiography**

MRI techniques used to depict arteries or veins, with or without contrast. It may evaluate stenosis, aneurysm, dissection, thrombosis, and vascular malformation without ionizing radiation.

## **Magnetic Resonance Cholangiopancreatography**

A special MRI scan that shows the bile ducts, gallbladder, and pancreatic duct without inserting a camera into them. It can detect stones, narrowing, inflammation, leaks, and some blockages without using X-rays.

### **Alternate Names:**

MRCP; MR Cholangiopancreatography

## **Magnetic Resonance Imaging**

A scan that uses a strong magnetic field and radio waves to make detailed pictures inside the body without X-rays. It is especially useful for the brain, spinal cord, joints, muscles, and many organs; some scans require injected contrast.

## **Magnetic Resonance Safety**

Precautions used during an MRI scan to prevent injury from the strong magnet, radio waves, heating,

loud noise, and metal objects. Staff must check implants, metal fragments, medicines, pregnancy, and other risks before scanning.

## **Magnetic Resonance Spectroscopy**

A special MRI test that measures chemicals in tissue rather than only making a picture of its shape. It may help study brain disorders or provide extra information about a tumor, but results must be interpreted with ordinary imaging and other tests.

## **Magnetic Resonance Urography**

An MRI scan of the kidneys, ureters, and bladder. It shows the urinary tract without X-rays and may be performed with or without contrast material.

## **Magnetic Resonance Venography**

An MRI test that produces images of veins, with or without injected contrast. It can assess suspected clots or narrowing in veins of the brain, abdomen, pelvis, or limbs and does not use X-rays.

### **Alternate Names:**

MRV; MR Venography

## **Magnetic Seed**

A tiny magnetic marker placed in a breast lesion or lymph node to help a surgeon find it. A special detector locates the marker during surgery, providing an alternative to some wire or radioactive-marker methods.

## **Magnetic Therapy**

Treatment that uses a magnetic field for a health purpose. Ordinary static magnets have not been shown to treat disease, while specialized magnetic stimulation devices have established uses for some neurologic or psychiatric conditions.

## **Magnetocardiography**

A test that records the very weak magnetic fields produced by the heart's electrical activity using sensors outside the body. It may help study heart rhythm or electrical conduction, usually in specialist or research settings.

---

## Magnetoencephalography

A noninvasive technique that records the weak magnetic fields produced by neuronal electrical activity, using sensors outside the head to localize and study brain function and some epileptic discharges.

## Magnusiomyces Capitatus

An opportunistic yeast formerly known as *Geotrichum capitatum* or *Saprochaete capitata*; it can cause invasive infections, especially in people with severe immunosuppression or hematologic malignancy.

## Mahvash Disease

A rare inherited disorder in which the body's cells do not respond normally to glucagon, a hormone that helps raise blood sugar. It can enlarge the pancreas, increase glucagon levels, and disturb blood-sugar control.

## Main-En-Griffe

A claw-like hand position usually caused by damage to the ulnar nerve, which controls several hand muscles. The ring and little fingers may bend at the smaller joints, weaken the grip, and become difficult to straighten.

## Maintenance

Ongoing care or treatment intended to preserve health, prevent relapse or deterioration, or sustain a therapeutic result; maintenance therapy is continued after initial control of a condition.

## Maintenance Dose

The regular amount of a medicine needed to keep its effect at the desired level. The dose depends on how quickly the body removes the medicine and how much the body absorbs.

## Maintenance Of Wakefulness Test

A sleep test that measures how long a person can remain awake in a quiet setting during the day.

## Maintenance Therapy

Treatment continued after a disease has improved

or been controlled to maintain that benefit, prevent return or worsening, or delay progression. The treatment may be a medicine, procedure, or other care, and its duration depends on the condition, response, risks, and treatment goals.

## Maisonneuve Fracture

A spiral fracture of the proximal third of the fibula associated with disruption of the distal tibiofibular syndesmosis, interosseous membrane, and either an avulsion fracture of the medial malleolus or rupture of the deep deltoid ligament of the ankle.

## Maize

Corn, a cereal grain used as food and feed; contaminated maize can expose people to fungal toxins such as aflatoxin.

## Majeed Syndrome

A rare autoinflammatory disorder caused by LPIN2 variants, characterized by chronic recurrent multifocal osteomyelitis, congenital dyserythropoietic anemia, and inflammatory skin disease.

## Majewski Syndrome

A rare inherited skeletal disorder involving abnormal bone growth and distinctive craniofacial or limb findings; the exact features depend on the genetic subtype.

## Majocchi Granuloma

A deep follicular infection caused by dermatophyte fungi, producing inflamed papules, pustules, or nodules and sometimes abscesses. It often follows shaving or topical steroid use and requires systemic antifungal treatment rather than topical therapy alone.

## Major

Larger, greater, or more significant than another related structure or condition; for example, the teres major muscle is larger than the teres minor muscle.

## Major Depression

Alternate name; see *Major Depressive Disorder*.

---

## Major Depression With Psychotic Features

A major depressive episode accompanied by delusions or hallucinations, usually mood-congruent, requiring prompt psychiatric assessment and treatment.

## Major Depressive Disorder

A mood disorder involving one or more major depressive episodes. During an episode, five or more symptoms occur during the same two-week period, including either depressed mood or loss of interest or pleasure (anhedonia). Other symptoms may include changes in sleep, appetite, energy, concentration, movement, guilt, worthlessness, or thoughts of death. The symptoms cause clinically significant distress or impairment.

### Alternate Names:

Major Depression; Clinical Depression

## Major Depressive Episode

A period of at least two weeks with depressed mood or loss of interest or pleasure plus additional emotional, cognitive, physical, or behavioral symptoms that cause significant distress or impairment. It may occur in major depressive disorder, bipolar disorder, or other conditions and must be distinguished from symptoms caused by substances or another medical condition.

## Major Histocompatibility Complex

A group of genes and cell-surface proteins that display pieces of germs or abnormal cells to the immune system. In humans, these proteins are called HLA.

## Major Histocompatibility Complex Molecule

A cell-surface protein that displays small pieces of germs or abnormal cells to immune cells. This helps the immune system recognize what belongs to the body and what may need an immune response.

### Alternate Names:

MHC Molecule

## Major Trauma

A serious injury that may threaten life, limb, or

organ function. It can involve multiple body regions or major damage to the brain, spine, chest, abdomen, pelvis, or blood vessels.

## Malabsorption

A condition in which the body cannot properly digest or absorb nutrients from food. It may cause chronic diarrhea, greasy stools, weight loss, anemia, or vitamin and mineral deficiencies.

## Malabsorption Syndrome

Impaired absorption of nutrients from the small intestine, causing diarrhea, weight loss, anemia, vitamin deficiencies, or other nutritional complications.

## Malacia

Abnormal softening or weakening of tissue. It may affect cartilage, bone, or brain tissue after faulty development, injury, infection, reduced blood supply, or another disease; symptoms depend on the tissue involved.

## Malacoplakia

A rare long-term inflammation in which immune cells collect mineral deposits. It most often affects the bladder or digestive tract and can form a mass that looks like cancer.

## Maladaptation

An adjustment to a situation or environment that does not work well and harms health or daily functioning. The term may describe physical, psychological, or behavioral responses to illness or stress.

## Maladaptive Stress Response

A lasting or overly strong reaction to stress that interferes with health, daily life, or relationships. It may include constant worry, poor sleep, irritability, avoidance, or physical symptoms and can improve with support and appropriate treatment.

## Malaise

Malaise is a general feeling of illness, discomfort, or lack of well-being that commonly accompanies infection, inflammation, chronic disease, medication

---

effects, or other systemic conditions.

### Malakoplakia

A rare long-term inflammation in which immune cells collect mineral deposits. It most often affects the bladder or digestive tract and can form a mass that resembles cancer.

### Malar

Relating to the cheekbone or cheek area. A malar rash is a red or purple rash across the cheeks and bridge of the nose, often seen in lupus. A malar fracture is a break in the cheekbone and may cause swelling, numbness, or a changed facial shape.

### Malaria

A serious infection caused by *Plasmodium* parasites and spread by infected *Anopheles* mosquitoes. The parasites first multiply in the liver and then enter red blood cells, causing fever, chills, sweating, headache, and weakness. Severe malaria can cause seizures, coma, severe anemia, kidney failure, or breathing problems. Some species can remain in the liver and cause later recurrences.

#### Alternate Names:

Plasmodium Infection; Paludism; Ague

### Malar Rash

A red or violaceous rash across the cheeks and bridge of the nose, often sparing the folds beside the nose. It is classically associated with systemic lupus erythematosus but can occur with other conditions; appearance alone does not establish a diagnosis.

### Malassezia

Malassezia is a yeast normally found on skin that can contribute to dandruff, seborrheic dermatitis, pityriasis versicolor, and some folliculitis.

### Malassezia Furfur

Malassezia furfur is a yeast that normally lives on skin but can multiply excessively. It is linked with dandruff, oily skin inflammation, and pityriasis versicolor, which causes lighter or darker patches.

### Malassezia (Pityrosporum) Folliculitis

An overgrowth of Malassezia yeast in hair follicles causing many small, itchy, fairly uniform papules and pustules, usually on the chest, back, shoulders, or forehead. Heat, sweating, antibiotics, oils, and immune suppression can contribute.

### Malassimilation

Failure to properly digest food or absorb its nutrients. It may cause diarrhea, weight loss, bloating, anemia, or vitamin shortages and can result from pancreatic disease, bowel disorders, surgery, infection, or food intolerance.

### Malate Dehydrogenase

An enzyme in energy-producing pathways that changes malate into oxaloacetate. It helps cells process nutrients and is involved in transferring chemical energy within mitochondria and other cellular compartments.

### Malathion

An organophosphate insecticide that inhibits acetylcholinesterase; topical formulations have also been used to treat head lice, but exposure or ingestion can cause cholinergic toxicity.

### Malathion Poisoning

Poisoning caused by the insecticide malathion. It can produce sweating, excess saliva, vomiting, diarrhea, wheezing, slow heartbeat, muscle twitching, weakness, seizures, or breathing failure.

### MALDI-TOF Mass Spectrometry

A laboratory method that identifies bacteria or fungi by comparing their protein pattern with a reference database. It can identify an organism quickly, but separate testing is needed to determine antibiotic sensitivity.

### Male

Describing a sex associated with reproductive anatomy that produces or is organized around small gametes (sperm); chromosomes, hormones, anatomy, and gender identity do not always fit a simple binary pattern.



---

## Male Breast Cancer

Cancer developing in breast tissue in a man. It is uncommon and may cause a new lump, nipple discharge, skin change, or nipple retraction.

### Alternate Names:

Male Mammary Carcinoma; Breast Carcinoma in Men

## Male Breast Pathology

Disorders affecting male breast tissue, including gynecomastia, benign epithelial lesions, inflammatory conditions, and carcinoma. Evaluation of a persistent mass or nipple change requires clinical examination and appropriate imaging, with biopsy when indicated.

## Male Chlamydial Infection

A sexually transmitted infection in men caused by *Chlamydia trachomatis* (serovars D–K), most frequently manifesting as non-gonococcal urethritis, epididymo-orchitis, or proctitis, characterized by dysuria, mucoid urethral discharge, and testicular tenderness.

### Alternate Names:

Male Chlamydial Urethritis; Chlamydia in Men; Male Chlamydia

## Male Condom

A thin sheath placed over the penis before sexual contact to reduce the risk of pregnancy and transmission of many sexually transmitted infections by acting as a physical barrier.

## Male Genitalia

The external and internal reproductive organs of a male, including the penis, scrotum, testes, epididymides, vas deferens, seminal vesicles, prostate, and associated ducts and glands.

## Male Genital Sores

Ulcerations, blisters, or skin lesions on the penis, scrotum, or perianal region. Major causes include sexually transmitted infections (such as genital herpes, syphilis, and chancroid), infectious balanitis, inflammatory skin diseases, fixed drug eruptions, trauma, or penile neoplasia.

### Alternate Names:

Male Genital Ulcers; Penile Lesions; Penile Ulcers

### Alternate Names:

Male Genital Ulcers; Penile Ulcerations

## Male Groin Pain

Pain in the groin or adjacent lower abdomen, hip, or testicle in a male; causes include muscle strain, hernia, infection, urinary stones, or referred pain.

## Male Hypoactive Sexual Desire Disorder

Persistently reduced sexual thoughts or desire in a man that causes personal distress or relationship difficulty. It differs from erection problems, although both can occur together; medicines, illness, mood, and relationship factors may contribute.

## Male Hypogonadism

Reduced production of testosterone, impaired sperm production, or both due to disease of the testes or the hypothalamic-pituitary system. Possible features include low libido, erectile difficulty, infertility, reduced muscle mass, fatigue, and loss of bone density.

### Alternate Names:

Low Testosterone; Testosterone Deficiency

## Male Infertility

Reduced ability to achieve a pregnancy because of factors affecting sperm production, sperm transport, ejaculation, sexual function, or other reproductive processes.

### Alternate Names:

Male Subfertility; Male Factor Infertility

## Male Menopause

Alternate informal term; see *Male Hypogonadism*.

## Male Pattern Baldness

Androgenetic alopecia, a patterned progressive loss of scalp hair caused by genetic susceptibility and androgen effects on hair follicles.

## Male Pelvis

The lower part of the abdomen between the hip



---

bones in a male; compared with the female pelvis, it is generally narrower, taller, and more robust, with a narrower pubic arch and sacrum.

### **Male Phenotype**

The observable reproductive anatomy, secondary sex characteristics, and other traits associated with male sex development, produced by interacting genetic, hormonal, and developmental factors and showing natural variation.

### **Male Reproductive System**

The organs and tissues that produce sperm and male sex hormones and deliver sperm, including the testes, ducts, accessory glands, penis, and scrotum.

### **Male Self-Catheterization**

Intermittent insertion and removal of a catheter through the male urethra to drain the bladder when neurologic, obstructive, or other conditions prevent adequate emptying.

### **Male Urethritis**

Inflammation of the male urethra, commonly caused by sexually transmitted infection but also by irritation or other infection. Dysuria, urethral discharge, itching, or meatal irritation may occur.

### **Malformation**

An abnormal structural development present from birth, caused by disturbed formation of an organ or body part.

### **Malignancy**

Cancerous disease in which abnormal cells grow uncontrollably and may invade nearby tissue or spread to distant organs. Its likely behavior and treatment depend on the cancer type, extent, grade, and important molecular findings.

### **Malignancy-Associated Hypercoagulability**

A cancer-related tendency toward excessive clot formation caused by tumor-associated coagulation activation, inflammation, platelet interaction, treat-

ment, immobility, or vascular injury. See *Cancer-Associated Thrombosis*.

### **Malignant**

Cancerous, with the ability to invade nearby tissues or spread to distant parts of the body. Malignant cells grow abnormally and may damage organs, but behavior varies among cancer types and stages.

### **Malignant Acanthosis Nigricans**

A paraneoplastic syndrome in which velvety, hyperpigmented, verrucous plaques develop rapidly in flexural and mucosal areas as a marker of an underlying internal malignancy, most commonly gastric adenocarcinoma. Unlike benign acanthosis nigricans, the malignant form is extensive, abrupt in onset, and may involve the oral mucosa.

### **Malignant Ameloblastoma**

A rare cancer arising from ameloblastoma-forming tissue, usually in the jaw, with potential to invade locally or spread to distant organs.

### **Malignant Ascites**

Buildup of fluid in the abdomen caused by cancer involving the abdominal lining or nearby organs. It can cause swelling, discomfort, early fullness, nausea, or breathlessness and may require drainage and cancer-directed treatment.

### **Malignant Bone Neoplasm**

A cancer that begins in bone or nearby supporting tissue. It may cause persistent bone pain, swelling, or a fracture; scans and a biopsy identify the type and extent before treatment.

### **Malignant Fibrous Histiocytoma**

Alternate name; see *Undifferentiated Pleomorphic Sarcoma*.

### **Malignant Giant Cell Tumor Of Tendon Sheath**

A rare locally aggressive malignant tumor resembling a giant cell tumor of tendon sheath, characterized by destructive growth and potential for metas-

---

tasis; diagnosis requires expert soft-tissue pathology review.

### **Malignant Glioma**

An aggressive primary tumor arising from glial or glial-like cells of the central nervous system; high-grade examples include anaplastic astrocytoma and glioblastoma.

### **Malignant Hypertension**

A medical emergency in which very high blood pressure is causing new injury to organs such as the brain, heart, kidneys, or eyes. It can cause headache, confusion, chest pain, breathlessness, vision changes, seizures, or kidney failure.

### **Malignant Hyperthermia**

A rare, life-threatening reaction to certain anesthetic medicines that causes rapidly rising body temperature, severe muscle stiffness, fast heart rate, and dangerous chemical changes. It affects people with certain inherited susceptibilities.

### **Malignant Mastocytosis**

An older name for a severe disorder in which abnormal mast cells multiply and collect in organs. It can damage the bone marrow, liver, spleen, or other tissues and may cause flushing, faintness, bleeding, or organ failure.

### **Malignant Melanoma**

A cancer that begins in melanocytes, the cells that make skin pigment. It may appear as a changing dark or unusual mole and can spread to other tissues, especially when diagnosed at a later stage.

### **Malignant Mesothelioma**

A cancer of the thin lining around the lungs, abdomen, or heart. It is strongly linked to asbestos exposure and may cause chest or abdominal pain, fluid buildup, cough, or breathlessness.

### **Malignant Myoepithelioma**

A rare cancer arising from supportive contractile cells in a gland, most often a salivary gland. It

can grow into nearby tissue and spread elsewhere, causing a slowly enlarging or painful mass.

### **Malignant Neoplasm**

A cancerous growth in which abnormal cells multiply without control, invade nearby tissues, and may spread to distant organs through the blood or lymphatic system.

#### **Alternate Names:**

Cancer; Malignancy

### **Malignant Otitis Externa**

An invasive infection of the external ear canal and skull base, usually caused by *Pseudomonas aeruginosa* in older adults with diabetes or immunocompromise; it can cause severe pain and cranial-nerve complications.

### **Malignant Peripheral Nerve Sheath Tumor**

An uncommon soft-tissue cancer arising from or near peripheral nerves. It may cause a growing painful mass, numbness, or weakness and is treated with specialist surgery, radiation, or medicines when appropriate.

### **Malignant Pleural Effusion**

Buildup of fluid around the lung because cancer involves the lung lining or has spread there. It may cause breathlessness, chest discomfort, or cough; drainage relieves symptoms, while cancer treatment may reduce recurrence.

### **Malignant Pustule**

An older name for the skin form of anthrax, caused by *Bacillus anthracis* and beginning as an itchy papule that develops into a painless black eschar.

### **Malignant Syphilis**

A severe form of secondary syphilis that causes fever and widespread crusted, ulcerated, or tissue-damaging skin lesions. It is more common with weakened immunity and requires prompt specialist treatment for syphilis.

---

### **Malignant Teratoma Of The Mediastinum**

A malignant germ-cell tumor containing tissues from more than one embryonic layer and arising in the mediastinum; evaluation uses imaging, tumor markers, and tissue diagnosis.

### **Malignant Thymoma**

An older term for an invasive thymic epithelial tumor; modern usage distinguishes invasive thymoma from thymic carcinoma, which is a separate, more aggressive thymic malignancy.

### **Malignant Tumor**

A cancerous growth capable of invading nearby tissue or spreading to distant sites. Its behavior and treatment depend on its type, grade, stage, and molecular features.

### **Malignant Vulvar Lesions**

Cancerous or potentially invasive lesions of the vulva, most commonly squamous-cell carcinoma but also melanoma, adenocarcinoma, basal-cell carcinoma, or sarcoma. Persistent, ulcerated, pigmented, bleeding, painful, or changing lesions require examination and biopsy.

### **Malingering**

Deliberately pretending to have, or exaggerating, symptoms to obtain an outside benefit such as money, medicines, shelter, or avoiding an obligation. It is not itself a mental illness and should not be diagnosed without reliable evidence.

### **Mallampati Classification**

A grading system used before anesthesia to estimate how difficult it may be to place a breathing tube. The clinician looks inside the open mouth and records which parts of the throat can be seen; it is only one part of the airway assessment.

### **Malleolus**

A bony prominence at the ankle formed by the lower end of the tibia medially or fibula laterally.

### **Mallet**

A surgical hammer used with tools such as osteotomes to cut or shape bone. It delivers controlled force during operations, helping surgeons work on bone while limiting unnecessary injury to nearby tissues.

### **Mallet Finger**

An injury to the extensor tendon at the distal finger joint, causing inability to actively straighten the fingertip. It may result from a forced flexion injury and sometimes includes an avulsion fracture.

### **Mallet Toe**

A flexion deformity of the distal interphalangeal joint, causing the end of a lesser toe to bend downward. It may be flexible or rigid and can cause pain, rubbing, corns, or difficulty wearing shoes; it differs anatomically from hammertoe and claw toe.

### **Mallet Toe and Hammertoe**

Alternate name; see *Hammertoe*.

### **Malleus**

The largest of the three small bones in the middle ear, also called the hammer. It is attached to the eardrum and passes sound vibrations to the incus and then the inner ear.

**Common name:** Hammer

### **Mallory Bodies**

Irregular, ropey, eosinophilic cytoplasmic inclusions within degenerating hepatocytes composed of tangled, ubiquitinated cytokeratin intermediate filaments (cytokeratins 8 and 18), characteristically observed in alcoholic hepatitis, non-alcoholic steatohepatitis (MASH), and Wilson disease.

**Alternate Names:**

Mallory-Denk Bodies; Mallory Hyaline

### **Mallory Body**

A clump of damaged proteins inside a liver cell, seen most often in alcohol-related liver disease but also in other liver conditions. It is a laboratory finding, not a diagnosis by itself.

---

## Mallory-Weiss Syndrome

Non-transmural, longitudinal mucosal tears localized to the gastroesophageal junction or gastric cardia, typically provoked by severe, forceful retching, vomiting, or coughing, presenting clinically as acute upper gastrointestinal hemorrhage and hematemesis.

### Alternate Names:

Mallory-Weiss Tear; Gastroesophageal Laceration

## Mallory-Weiss Tear

A superficial tear in the lining of the lower esophagus or upper stomach, usually caused by forceful vomiting or retching. It commonly causes vomiting of blood. Many tears stop bleeding on their own within a few days, but upper endoscopy may be needed for diagnosis and treatment.

## Malnutrition

Poor health caused by too little, too much, or an imbalance of nutrients. It can impair growth, immunity, wound healing, muscle function, and organ function and may involve undernutrition, nutrient deficiencies, or excess weight.

## Malocclusion

A mismatch in the position of the upper and lower teeth or jaws that affects how the teeth meet. It may cause crowding, an overbite, an underbite, difficulty chewing, speech problems, jaw discomfort, or trouble cleaning the teeth.

## Malocclusion Of Teeth

Misalignment of the upper and lower teeth or jaws that affects the bite, chewing, speech, appearance, or oral hygiene.

## Malondialdehyde

A chemical formed when fats are damaged by oxidation. Laboratory researchers may measure it as an indirect sign of oxidative stress, but the result alone does not diagnose a disease.

## Malpighian Bodies

Renal glomeruli within Bowman's capsules, where

blood filtration begins; the term may also refer historically to lymphoid nodules in the spleen.

## Malposition

An abnormal position of an organ, body part, implanted device, or fetus; in obstetrics, fetal malposition refers to an abnormal orientation of the presenting part during labor.

## Malposition of the Uterus

A uterus tilted or oriented differently from the usual position, such as retroversion or retroflexion. It is often a normal variant, but pain, pressure, infertility, urinary symptoms, or a mass may indicate an underlying condition requiring evaluation.

## Malpresentation

A fetal presentation other than the usual vertex presentation, including breech, face, brow, shoulder, or compound presentation. It may complicate labor and influence delivery planning.

## Malrotation

Abnormal rotation or fixation of an organ during development, most often referring to abnormal rotation of the intestine and its risk of volvulus.

## MALS

Abbreviation; see *Median Arcuate Ligament Syndrome*.

## Malta Fever

Brucellosis, a zoonotic infection caused by *Brucella* species and characterized by intermittent fever, sweats, fatigue, and joint or systemic symptoms.

## Maltase

An intestinal enzyme that breaks maltose into two glucose molecules.

## MALT Lymphoma

Alternate name; see *MALT Lymphoma*.

## Maltodextrin

A mixture of short glucose polymers produced by

---

partial hydrolysis of starch, used as a rapidly digestible carbohydrate and as a thickener, filler, or carrier in foods and medicines.

### **Maltose**

A reducing disaccharide composed of two glucose molecules joined by an alpha-1,4 glycosidic bond, produced during starch digestion and hydrolyzed by maltase.

### **Malt-Worker's Lung**

An occupational hypersensitivity pneumonitis caused by inhalation of organic dusts associated with malting or grain processing.

### **Mammaglobin A**

A secretory protein expressed predominantly in mammary tissue and used as an immunohistochemical marker that can help identify breast origin in some metastatic carcinomas.

### **Mammaplasty**

Surgery to change the size or shape of the breast or to rebuild it after breast removal. It may enlarge, reduce, lift, or reconstruct the breast; possible risks include bleeding, infection, altered sensation, scarring, and problems with healing.

### **Mammary**

Relating to the breast or milk-producing gland. Mammary glands develop during puberty and pregnancy and produce milk after childbirth under hormone control. The term may describe normal breast tissue or diseases such as cysts, infection, benign growths, or breast cancer.

### **Mammary Duct Ectasia**

A noncancerous inflammation in which milk ducts beneath the nipple become widened and filled with thick material. It may cause sticky green, brown, or creamy discharge, nipple tenderness, nipple pulling inward, or a lump that can look like breast cancer on examination or imaging.

#### **Alternate Names:**

Duct Ectasia; Plasma Cell Mastitis; Periductal Mastitis; Comedomastitis

### **Mammary Epithelium**

The epithelial lining of the mammary ducts and secretory alveoli; these cells help produce, secrete, and transport milk and can give rise to many breast carcinomas.

### **Mammary Gland**

The milk-producing gland in the breast. It contains small milk-making lobules connected to ducts that carry milk to the nipple, and its growth and activity are controlled by hormones.

### **Mammary Glands**

Paired glands of the anterior chest that produce milk after childbirth. Each breast contains lobules, ducts, connective tissue, and fat and is influenced by hormonal and reproductive changes.

#### **Alternate Names:**

Breasts; Lactiferous Glands

### **Mammary Implant**

An artificial device placed in the breast for reconstruction or enlargement. Implants may contain saline or silicone gel and may be placed above or below the chest muscle. Possible complications include infection, rupture, hard scar tissue, changes in sensation, and rarely implant-associated lymphoma.

### **Mammary Paget Disease**

An intraepidermal spread of malignant glandular cells into the nipple epidermis, usually associated with underlying breast carcinoma and presenting as persistent eczematous or crusted nipple change.

### **Mammillary Bodies**

A pair of small, spherical hypothalamic nuclei situated at the posterior termination of the anterior columns of the fornix on the ventral surface of the diencephalon, forming an essential component of the Papez circuit for memory consolidation; severely damaged in Wernicke encephalopathy.

#### **Alternate Names:**

Corpora Mamillaria

---

## **Mammillary Body**

One of a pair of rounded nuclei on the underside of the hypothalamus that participate in memory circuits connecting the hippocampus with the thalamus; injury can contribute to memory impairment.

## **Mammogram**

A mammogram is a low-dose x-ray examination of breast tissue used for screening and evaluation of masses, asymmetry, calcifications, or other abnormalities.

## **Mammographic Calcifications**

Mineral deposits seen in breast tissue on a mammogram. Large deposits are usually noncancerous, while the shape and arrangement of very small deposits can be linked to early breast cancer or other changes and may lead to further imaging or biopsy.

### **Alternate Names:**

Breast Microcalcifications; Mammographic Microcalcifications; Breast Calcifications

## **Mammography**

X-ray imaging of the breast used to screen for or evaluate breast abnormalities. Screening intervals depend on age, risk, symptoms, and the guideline being followed. Standard views include the cranio-caudal and mediolateral-oblique projections.

## **Mammoplasty**

Plastic or reconstructive breast surgery that changes breast size, shape, or contour. It may reduce or enlarge the breasts, lift them, rebuild them after cancer surgery, or correct differences present from birth.

## **Management of Third-Stage Labor**

Care from birth of the baby until delivery of the placenta, including monitoring, uterotonic medication when indicated, controlled cord traction by trained clinicians, and assessment for retained placenta or postpartum hemorrhage.

## **Mandatory Reporting**

A legal or regulatory duty to report specified find-

ings or circumstances—such as suspected child or elder abuse, certain communicable diseases, or some injuries—to designated authorities; requirements vary by jurisdiction.

## **Mandible**

The large movable bone forming the lower face and holding the lower teeth. It supports chewing and speech and contains nerves and blood vessels that supply the teeth, gums, and lip.

### **Alternate Names:**

Inferior Maxilla; Jawbone

## **Mandibular**

Relating to the mandible, the lower jawbone. It holds the lower teeth and moves at the jaw joints. A mandibular fracture can cause pain, swelling, numbness, loose teeth, or a changed bite.

## **Mandibular Advancement Device**

An oral appliance that holds the lower jaw or tongue forward during sleep to enlarge or stabilize the upper airway, usually for selected patients with obstructive sleep apnea or snoring.

## **Mandibular Condyle**

The rounded superior end of each mandibular ramus that articulates with the temporal bone at the temporomandibular joint and permits jaw movement.

## **Mandibular Fracture**

A break in the lower jaw bone typically caused by trauma, presenting with pain, swelling, malocclusion, and difficulty opening the mouth; may require surgical fixation.

## **Mandibular Prosthesis**

An artificial implant or device used to replace, reconstruct, stabilize, or restore part of the mandible after trauma, tumor removal, congenital deformity, or other loss of jaw structure.

## **Mandibulofacial Dysostosis**

A congenital craniofacial disorder, classically Treacher Collins syndrome, involving underdevelop-



---

ment of the mandible and cheekbones with variable abnormalities of the ears, eyes, and palate.

### **Manganese**

An essential trace element required in small amounts. Excessive exposure, especially by inhalation, can injure the nervous system and cause manganism, a movement disorder resembling parkinsonism.

### **Mania**

A distinct period of abnormally elevated, expansive, or irritable mood and increased energy or activity, with associated symptoms such as reduced need for sleep, rapid speech, racing thoughts, inflated self-esteem, or risky behavior. It causes marked impairment, requires hospitalization, or includes psychotic features; the episode usually lasts at least one week when hospitalization is not required.

### **Manic**

Relating to mania, a pathological episode of abnormally elevated, expansive, or irritable mood with increased energy or activity and associated changes in thought, speech, sleep, or judgment.

### **Manic-Depression**

Older term; see *Bipolar Disorder*.

### **Manic-Depressive Illness**

Alternate name; see *Bipolar Disorder*.

### **Manic Disorder**

Alternate name; see *Bipolar Disorder*.

### **Manic Episode**

A distinct period of abnormally elevated, expansive, or irritable mood and increased energy or activity, lasting at least seven days unless hospitalization is required sooner or psychotic features occur. It includes associated changes such as reduced need for sleep, rapid speech, racing thoughts, inflated self-esteem, or risky behavior and causes marked impairment.

### **Manic Mood**

An abnormally elevated, expansive, or persistently irritable mood accompanied by increased energy or activity, reduced need for sleep, rapid speech, racing thoughts, inflated self-esteem, or risky behavior; it is characteristic of a manic episode.

### **Manifest**

Clearly apparent or expressed; in clinical usage, a disease may manifest through particular signs or symptoms.

### **Manifestation**

A sign, symptom, or other observable expression of a disease or health condition. Manifestations may be noticed by the person, found during examination, or detected through laboratory tests or imaging.

### **Manipulation**

Deliberate physical movement of a body part or joint, often performed therapeutically or diagnostically by a trained clinician.

### **Manipulation Procedure**

Deliberate movement of a body part by a trained clinician to restore motion, alignment, or function. It may be performed with or without anesthesia; soreness, bleeding, nerve injury, fracture, or worsening symptoms are possible depending on the procedure.

### **Mannan**

A polysaccharide composed primarily of mannose units, found in the cell walls or capsules of many fungi and in some plants; fungal mannans can be recognized by the immune system and used in laboratory testing.

### **Mannheimia**

A genus of gram-negative bacteria that mainly infects livestock, especially the respiratory tract of cattle and sheep.

### **Mannitol**

A medicine that pulls water into the urine. It is

---

given by injection in selected emergencies to reduce swelling or pressure around the brain or inside the eye and in some kidney-related situations; it can worsen dehydration or heart failure.

### **Mannose**

A six-carbon monosaccharide related to glucose that forms part of glycoproteins and glycolipids and is important in protein glycosylation and cell recognition.

### **Mannose-Binding Lectin Deficiency**

Low levels or reduced activity of a blood protein that helps the immune system recognize and attack germs. Many people have no symptoms, but some are more prone to repeated infections, especially if other immune defenses are weak.

### **Mannosidosis**

An inherited lysosomal storage disorder caused by deficient alpha-mannosidase activity, leading to accumulation of mannose-rich oligosaccharides and variable intellectual, skeletal, immune, hearing, and neurologic abnormalities.

### **Manometer**

An instrument that measures pressure in a liquid or gas. In medicine, it may measure pressure in blood vessels, the lungs, the bladder, or another body space.

### **Manometry**

A test that measures pressure and muscle coordination inside an organ or passage, such as the esophagus, stomach, rectum, or urinary tract. It helps identify movement and sphincter disorders.

### **Mansonella Perstans**

A filarial worm that causes mansonellosis, usually acquired through bites of infected midges and often producing mild or no symptoms.

### **Mansonella Streptocerca**

A filarial worm transmitted by biting midges that lives mainly in the skin and can cause itchy skin

disease or small nodules.

### **Mansonelliasis**

Infection with filarial nematodes of the genus *Mansonella*, transmitted by biting midges or blackflies; many infections are asymptomatic, but some cause itching, skin lesions, fever, or eosinophilia.

### **Mantle Cell Lymphoma**

A usually fast-growing cancer of B lymphocytes that often enlarges lymph nodes and the spleen and may involve the bone marrow, blood, or digestive tract. A characteristic chromosome change helps confirm the diagnosis and guide treatment.

### **Mantle Field**

A historical radiation-treatment field encompassing lymph-node regions in the neck, mediastinum, and upper chest, formerly used for Hodgkin lymphoma; modern treatment generally uses more limited involved-site fields.

### **Mantoux Test**

A skin test for immune sensitization to tuberculosis. A small amount of purified protein derivative is injected into the forearm, and the raised area is measured 48 to 72 hours later; a positive result does not by itself prove active disease.

### **Manual**

Performed by hand rather than automatically, or a written guide explaining procedures and instructions. In healthcare, manual techniques may assess, move, treat, or support body structures safely.

### **Manual Removal Of Placenta**

A procedure in which a clinician removes a placenta that has not separated or cannot be delivered after birth. The timing and need depend on bleeding, maternal condition, and local clinical protocols.

### **Manual Vacuum Aspiration**

A procedure that uses a handheld syringe and a thin tube to remove tissue from the uterus. It may treat early pregnancy loss or end a pregnancy,

---

depending on local law and clinical circumstances, and is performed by trained clinicians.

**Alternate Names:**

MVA; Mini-Suction Curettage

**Manubrium**

The broad, upper quadrangular segment of the sternum articulating superiorly with the clavicles (sternoclavicular joints) and first costal cartilages, and joining inferiorly with the sternal body at the sternal angle of Louis (level of T4-T5 intervertebral disc).

**Alternate Names:**

Manubrium Sterni

**Manus**

The hand, including the wrist and fingers, in anatomical terminology.

**MAOIs**

A class of medicines used mainly for depression and some other conditions. They slow the breakdown of certain brain chemical messengers and can have important interactions with foods and other medicines.

**Map-Dot-Fingerprint Type Corneal Dystrophy**

A usually bilateral corneal epithelial basement membrane dystrophy producing map-like, dot-like, or fingerprint-like patterns and sometimes recurrent corneal erosions or visual blurring.

**Maple Syrup Urine Disease**

An inherited disorder in which the body cannot properly break down three amino acids from protein. Harmful substances build up, causing sweet-smelling urine, poor feeding, sleepiness, abnormal movements, seizures, or brain injury without treatment.

**Mapping**

Charting the location of genes or other identifiable regions on chromosomes; the term may also describe charting functional or anatomical regions in other

medical contexts.

**Marantic**

Relating to a sterile, noninfective process associated with wasting; marantic endocarditis refers to nonbacterial thrombotic endocarditis.

**Marasmus**

Severe undernutrition caused by prolonged lack of calories and protein. It leads to marked loss of body fat and muscle, poor growth in children, weakness, and increased vulnerability to infection.

**Maraviroc**

A CCR5 antagonist (entry inhibitor) used in combination antiretroviral therapy for HIV-1 infection in patients with CCR5-tropic virus. Maraviroc binds to the human CCR5 coreceptor on CD4+ T cells, preventing HIV-1 gp120 binding and viral entry.

**Marble Bone Disease**

Osteopetrosis, an inherited disorder of defective bone resorption causing abnormally dense but fragile bones and possible marrow failure.

**Marburg Virus**

A filovirus that causes Marburg virus disease, a severe viral hemorrhagic fever in humans and some nonhuman primates. The virus is associated with Egyptian rousette fruit bats and can first spread to people after contact with infected bats. It then spreads between people through direct contact with infected blood, other body fluids, or materials contaminated with them, not through ordinary airborne spread. The disease may begin with high fever, severe headache, tiredness, and muscle aches, followed by vomiting, diarrhea, rash, bleeding, shock, or organ failure.

**Marburgvirus**

A group of filoviruses that includes Marburg virus and Ravn virus, the viruses that cause Marburg virus disease. The disease is a severe viral hemorrhagic fever that can spread from fruit bats to people and between people through direct contact with infected body fluids.

---

### Marcus Gunn Jaw-Winking Phenomenon

An uncommon congenital synkinetic neuro-ophthalmological anomaly characterized by unilateral upper eyelid retraction or elevation occurring synchronously with movements of the lower jaw (such as chewing, sucking, or lateral jaw excursion). Caused by aberrant congenital innervation of the ipsilateral levator palpebrae superioris muscle (cranial nerve III) by motor branches of the trigeminal nerve (cranial nerve V<sub>3</sub>) supplying the pterygoid muscles.

#### Alternate Names:

Jaw-Winking Syndrome; Marcus Gunn Ptosis

### Marcus Gunn Pupil

A relative afferent pupillary defect (RAPD) detected during the swinging-flashlight test, wherein the affected pupil displays paradoxical dilation (or reduced constriction) when the light source is shifted from the unaffected eye to the abnormal eye, indicating unilateral or asymmetric optic nerve or retinal pathology.

#### Alternate Names:

Relative Afferent Pupillary Defect; RAPD

### Marden-Walker Syndrome

A rare inherited disorder characterized by a distinctive facial appearance, low muscle tone, limited joint movement, poor growth, and developmental delay. Care is supportive and involves several specialties.

### Marek's Disease

A contagious herpesvirus disease of chickens that can cause nerve paralysis, tumors, weight loss, or death.

### Marfanoid Habitus

A body habitus characterized by tall stature, long limbs, long fingers (arachnodactyly), and joint laxity. Seen in Marfan syndrome, but also in other connective tissue disorders and homocystinuria. Differentiated from Marfan syndrome by absence of aortic root dilation and lens subluxation.

### Marfan's Syndrome

Alternate spelling; see *Marfan Syndrome*.

### Marfan Syndrome

An autosomal dominant connective-tissue disorder usually caused by a disease-causing variant in the *FBN1* gene, which encodes fibrillin-1. It may affect the aortic root and heart valves, eyes, skeleton, joints, lungs, and dura. Features can include aortic-root enlargement, lens dislocation, tall or long-limbed stature, long fingers, scoliosis, chest-wall differences, and joint laxity; the severity varies among individuals.

### Martinus Beijerinck

A Dutch microbiologist who helped establish the concept of viruses as infectious agents that reproduce only in living cells and contributed to the development of enrichment culture methods.

### Marginal Ulcer

An ulcer occurring at or near a gastrojejunal anastomosis, often after gastric bypass or other gastric surgery, where acid, ischemia, smoking, nonsteroidal anti-inflammatory drugs, or other factors may contribute.

### Marginal Zone Lymphoma

A slow-growing non-Hodgkin lymphoma that begins in immune cells in the marginal zone of lymphoid tissue.

#### Alternate Names:

MZL

### Maribavir

Maribavir is an antiviral medicine used to treat certain cytomegalovirus infections that are resistant or refractory to other treatment after transplantation.

### Marie-Strumpell Disease

Ankylosing spondylitis, a chronic inflammatory disease primarily affecting the sacroiliac joints and spine that can cause back pain, stiffness, and progressive spinal fusion.

### Marijuana

Cannabis preparations containing cannabinoids such as THC; medical effects and risks depend on dose,

---

formulation, route, and patient factors.

**Alternate Names:**

Cannabis; Weed; Pot

**Marijuana/Cannabis**

Cannabis is a plant containing active chemicals such as THC and cannabidiol. THC can affect memory, coordination, mood, and judgment; cannabis may also interact with medicines and can cause dependence.

**Marijuana Intoxication**

Temporary changes after using cannabis, including impaired attention, memory, coordination, judgment, or perception. Anxiety, panic, drowsiness, vomiting, or psychosis may occur. Severe symptoms or accidental ingestion by a child can be dangerous.

**Marijuana Substance Use**

Marijuana or cannabis can impair attention, memory, coordination, and driving and may cause dependence, anxiety, psychosis, or respiratory harm when smoked.

**Marine Animal Stings Or Bites**

Injuries caused by marine animals through biting, stinging, spines, or venom; effects range from local pain and swelling to allergic, neurologic, or systemic toxicity.

**Marine Toxin**

A poisonous substance produced or accumulated by marine organisms, such as some fish, shellfish, algae, or bacteria, that can cause neurologic, gastrointestinal, or other poisoning.

**Marital Disharmony**

Persistent conflict or distress in a marriage or partnership that may contribute to psychological symptoms or require relationship-focused support.

**Marital Status**

A demographic or social-status variable describing whether a person is unmarried, married or partnered, separated, divorced, widowed, or in another

legally or culturally recognized relationship; it may be relevant to social and health assessments.

**Marital Therapy**

Counseling for couples that addresses communication, conflict, emotional distress, intimacy, and relationship problems. It may help partners improve interaction and make informed decisions about their relationship.

**Marjolin Ulcer**

A rare, aggressive cutaneous squamous cell carcinoma arising within longstanding chronic non-healing wounds, burn scars (cicatrices), chronic venous stasis ulcers, or chronic osteomyelitis sinus tracts.

**Alternate Names:**

Scar Carcinoma; Burn Scar Squamous Cell Carcinoma

**Marked**

Pronounced, conspicuous, or substantially greater than usual; in a clinical report, the exact meaning depends on the finding and the comparison standard.

**Marker**

An identifiable biological feature used to track a gene, cell, disease, or process; a genetic marker is a detectable DNA region inherited with a nearby gene, while a blood or tumor marker is a measurable substance associated with a condition.

**Maroteaux-Lamy Syndrome**

A rare inherited disorder in which certain complex sugars build up in the body. It can cause short stature, stiff joints, enlarged liver or spleen, cloudy corneas, heart-valve disease, breathing problems, and coarse facial features. Intelligence is often normal. Treatment may include enzyme replacement, supportive care, or stem-cell transplantation.

**Alternate Names:**

Mucopolysaccharidosis Type Vi; MPS VI; Arylsulfatase B Deficiency

---

## **Marrow**

The soft tissue inside bones, including red marrow that produces blood cells and yellow marrow rich in fat.

## **Marrow Hyperplasia**

Increased activity and numbers of blood-forming cells in the bone marrow. It may occur when the body is replacing blood lost or destroyed, or when tissues are not receiving enough oxygen.

## **Marshall Syndrome**

A rare periodic-fever syndrome, usually associated with PFAPA features, including recurrent fever, aphthous ulcers, pharyngitis, and cervical adenitis, often beginning in childhood.

## **Marsupialization**

A surgical technique that opens a cyst or abscess and sutures its edges to the surrounding skin or mucosa to permit continuing drainage.

## **Martin-Bell Syndrome**

An older name for fragile X syndrome, an inherited neurodevelopmental disorder caused by CGG-repeat expansion in the FMR1 gene.

## **MASA Syndrome**

An X-linked neurodevelopmental disorder characterized by intellectual disability, absent or limited speech, shuffling gait, spasticity, and adducted or clasped thumbs; it may also include short stature and lumbar lordosis. The name is an older acronym and terminology should be used sensitively.

## **Masculinization**

Development of male-typical physical characteristics, caused by normal puberty, androgen excess, androgen treatment, or certain disorders.

## **MASH**

Abbreviation; see *Metabolic Dysfunction-Associated Steatohepatitis*.

## **Mask**

A device placed over the face to deliver oxygen, anesthesia, ventilation, or protection; the meaning depends on clinical context.

## **Masked Data**

Data altered, obscured, or replaced so that identifying values or sensitive details are hidden while selected analytical properties may be retained; masking is used to protect privacy and can be reversible or irreversible.

## **Masklike Face**

An expressionless face with little spontaneous animation, also called masklike facies; it may occur in Parkinson disease, myotonic dystrophy, and other disorders.

## **MASLD**

Abbreviation; see *Metabolic Dysfunction-Associated Steatotic Liver Disease*.

## **Masochism**

Sexual or nonsexual arousal or gratification associated with receiving pain, humiliation, or degradation; distress or impairment determines clinical significance.

## **Mason Fracture**

A clinical and radiographic classification system for fractures of the radial head: Type I is non-displaced; Type II is a displaced partial-head marginal fracture; Type III is a severely comminuted fracture involving the entire head; and Type IV is a radial head fracture associated with elbow dislocation.

### **Alternate Names:**

Radial Head Fracture; Mason Classification of Radial Head Fracture

## **Masoprocycl**

A synthetic derivative of nordihydroguaiaretic acid that inhibits lipoxygenase pathways and has been investigated for topical treatment of actinic keratoses and for antineoplastic activity.



---

## Mass

A discrete area of abnormal tissue that is larger or more substantial than a nodule. A mass may be benign or malignant and can occur in many organs.

## Massage

Manual manipulation of soft tissues using techniques such as stroking, kneading, or pressure, used for relaxation or symptom relief; it is not a substitute for diagnosis or treatment of an underlying disease.

## Massage Therapy

Massage therapy uses hands-on manipulation of soft tissues to promote relaxation or help manage selected pain and movement problems.

## Mass Casualty Incident

An event causing more injured or ill people than nearby medical services can immediately manage. Health teams use triage to prioritize urgent treatment, transport, and limited supplies.

## Masseter

A powerful muscle of chewing that elevates the mandible and lies over the lateral surface of the mandibular ramus.

## Masseter Muscle

A strong chewing muscle running from the cheekbone to the lower jaw. It raises the jaw to close the mouth and may become painful or enlarged with teeth grinding.

## Massive Hemoptysis

Life-threatening coughing of blood that threatens the airway, breathing, or circulation; a fixed volume threshold is not required. It requires urgent airway assessment, stabilization, and definitive treatment of the bleeding source.

## Massive Ovarian Edema

Marked stromal edema causing enlargement of one or both ovaries, usually due to partial torsion or hormonal stimulation. It can mimic an ovarian neoplasm and may cause pain or virilization.

## Massive Pulmonary Embolism

A high-risk, life-threatening pulmonary embolism characterized by extensive clot obstruction causing acute right ventricular strain and hemodynamic collapse (systolic blood pressure < 90 mmHg or obstructive shock). See *Massive Pulmonary Embolus*.

## Massive Pulmonary Embolus

A severe blockage of the main blood vessels to the lungs by a blood clot. It can suddenly strain or stop the right side of the heart and cause very low blood pressure, severe breathlessness, fainting, bluish skin, or cardiac arrest. Treatment may include clot-dissolving medicine or removal of the clot.

### Alternate Names:

Massive PE; High-Risk Pulmonary Embolism; Fulminant Pulmonary Embolism

## Massive Transfusion Protocol

A coordinated protocol for life-threatening hemorrhage that provides rapid, balanced delivery of red cells, plasma, and platelets with monitoring and correction of fibrinogen, calcium, temperature, and acid-base abnormalities.

## Mass Screening

Testing a large group of people who do not yet have symptoms to find disease or risk early. A useful screening program has a suitable test, an effective follow-up pathway, and benefits that outweigh harms.

## Mass Spectrometer

A laboratory instrument that identifies chemicals by giving their particles an electrical charge and separating them according to size and charge. It can measure medicines, toxins, proteins, and other substances.

## Mass Spectrometry

An analytical method that identifies and measures molecules by their mass-to-charge ratios; coupled with chromatography, it is widely used in clinical chemistry, toxicology, pharmacology, and proteomics.

---

## **Mastalgia**

Breast pain or tenderness, which may be cyclical with menstruation or noncyclical. Causes include hormonal changes, cysts, infection, trauma, and musculoskeletal pain.

## **Mast Cell**

An immune cell found in tissues that releases substances involved in allergy, inflammation, and defense against some germs. Its products can cause itching, swelling, mucus production, and changes in blood pressure.

## **Mast Cell Activation Syndrome**

A condition involving recurrent symptoms from inappropriate release of mast-cell mediators, such as flushing, itching, hives, abdominal symptoms, wheezing, or low blood pressure. Diagnosis requires compatible episodes, objective evidence of mediator release, and response to appropriate treatment; many similar conditions must be excluded.

## **Mast Cell Sarcoma**

An exceptionally rare, aggressive neoplasm composed of immature or atypical mast cells forming a destructive mass in tissue or bone. Unlike systemic mastocytosis, it is characterized by a localized sarcomatous tumor and may progress to disseminated mast-cell disease or acute leukemia. Clinical manifestations depend on the involved site and may include pain, swelling, organ dysfunction, or mediator-related symptoms such as flushing and hypotension. Diagnosis requires histopathology, mast-cell immunophenotyping, and systemic staging; management generally involves specialist-directed intensive antineoplastic therapy.

## **Mast-Cell Sarcoma**

An extremely rare cancer made of abnormal mast cells, immune cells involved in allergic and inflammatory reactions.

## **Mastectomy**

Surgery to remove breast tissue from one or both breasts, usually to treat or prevent selected breast cancers. Most mastectomies remove nearly all breast

tissue; whether the nipple, areola, skin, or nearby lymph nodes are removed depends on the type. A lumpectomy removes a tumor or abnormal area while leaving most of the breast in place.

## **Masticate**

To chew food, using the teeth, tongue, and muscles of mastication to mechanically break it down and mix it with saliva before swallowing.

## **Mastication**

Chewing food with the teeth and moving it with the tongue before swallowing. It breaks food into smaller pieces and mixes it with saliva, making it easier to swallow and begin digesting.

## **Masticatory Muscle**

One of the muscles that moves the mandible during chewing, chiefly the masseter, temporalis, medial pterygoid, and lateral pterygoid muscles.

## **Mastitis**

Inflammation of breast tissue, often occurring during lactation but also possible outside lactation. It may cause localized pain, warmth, swelling, and fever; infection and inflammatory causes must be distinguished clinically.

### **Alternate Names:**

Puerperal Mastitis; Breast Inflammation

## **Mastocytoma**

A localized collection or tumor of mast cells, usually presenting in the skin as a solitary cutaneous mastocytoma; rubbing the lesion can cause itching, redness, or swelling from mediator release.

## **Mastocytosis**

A disorder in which too many mast cells, immune cells involved in allergic reactions, collect in the skin or internal organs. It may cause itchy spots, flushing, abdominal symptoms, wheezing, or low blood pressure.

## **Mastodynia**

Pain or tenderness in one or both breasts, also

---

called mastalgia. It may follow the menstrual cycle or result from injury, infection, medicines, cysts, or other breast conditions; a new persistent lump, skin change, or bloody discharge may indicate another breast problem.

### **Mastoid**

Relating to the mastoid part of the temporal bone behind the ear. This area contains air spaces connected with the middle ear and can become infected, causing pain, swelling, or drainage.

### **Mastoid Antrum**

An air-containing cavity in the mastoid portion of the temporal bone that communicates with the middle-ear attic.

### **Mastoid Bone**

The portion of the temporal bone located behind the ear.

### **Mastoidectomy**

Surgery to remove infected or diseased air cells in the bone behind the ear. It may treat chronic middle-ear infection, cholesteatoma, mastoiditis, or complications that have spread beyond the ear.

### **Mastoiditis**

Mastoiditis is infection and inflammation of the mastoid air cells behind the ear, usually following middle-ear infection and potentially causing serious spread.

### **Mastoid Process**

The conical bony projection of the temporal bone behind the ear, containing air cells and providing attachment for several neck muscles; infection of its air cells is mastoiditis.

### **Mastopexy**

Breast-lift surgery that removes extra skin and reshapes breast tissue to raise the breasts and nipples. It can improve shape but does not necessarily reduce breast size; scars, altered sensation, asymmetry, and changes in breastfeeding ability are possible risks.

### **Alternate Names:**

Breast Lift

### **Masturbation**

Self-stimulation of the genitals for sexual pleasure. It is a common human behavior; it is generally not harmful unless it causes injury, distress, or interferes with daily life.

### **Alternate Names:**

Autoeroticism; Solitary Sex

### **Materia Medica**

An older term for the body of medicinal substances and information about their properties and uses.

### **Maternal**

Maternal means relating to a mother or the pregnant person, especially in connection with pregnancy, childbirth, or postpartum health.

### **Maternal Age**

Maternal age is the age of the pregnant person and can influence fertility, pregnancy risks, screening choices, and obstetric care.

### **Maternal Expulsive Efforts**

The pregnant person's efforts to push the baby down and out during the second stage of labor. Pushing may follow the body's urge or be guided by the birth team, depending on the situation and chosen pain relief.

### **Alternate Names:**

Bearing Down; Second Stage Expulsive Efforts

### **Maternal Health**

The physical and mental health of a person during pregnancy, childbirth, and the postpartum period. Care includes prevention, antenatal monitoring, safe delivery, and treatment of complications.

### **Maternal Mortality**

Maternal mortality is death during pregnancy or within the defined postpartum period from a pregnancy-related cause, measured as a population health indicator.

---

### **Maternal Mortality Rate**

The number of maternal deaths related to pregnancy per 100,000 live births in a specified population and period.

### **Maternal Obesity**

Obesity present before or during pregnancy. It can increase the risk of high blood pressure, diabetes, cesarean birth, and health problems for the pregnant person or baby.

### **Maternal Physiologic Changes In Pregnancy**

Normal body changes during pregnancy that support the growing fetus and prepare for birth. They affect blood volume, heart function, breathing, digestion, kidneys, hormones, breasts, skin, and the musculoskeletal system.

### **Maternal Sepsis**

Organ dysfunction caused by a dysregulated maternal response to infection during pregnancy, childbirth, or the postpartum period. It may arise from uterine, urinary, respiratory, wound, or other infection and requires prompt cultures, antimicrobial therapy, source control, and hemodynamic support.

### **Maternal Serum Alpha-Fetoprotein**

A blood test during pregnancy, usually at 15–20 weeks, that measures a protein made by the fetus. An unusually high or low result may be linked to an incorrect pregnancy date, more than one fetus, or certain birth defects or chromosome conditions. It is a screening test, not a diagnosis, and abnormal results need follow-up.

### **Maternal Vaccine**

A vaccine given during pregnancy to protect the pregnant person and, through transferred antibodies, provide early protection to the fetus or newborn; recommended products and timing depend on current guidance.

### **Maternity**

The state or service relating to pregnancy, childbirth, and the care of the mother and newborn.

### **Maternity Services**

Healthcare for pregnancy, labor and birth, the postpartum period, and the newborn. Services may include prenatal checks, midwifery or obstetric care, delivery support, breastfeeding help, and emergency care.

### **Materteral**

Relating to a maternal aunt or to the genetic relationship between an aunt and her niece or nephew; synonymous with materterine.

### **Mathematics Disorder**

A learning disorder involving persistent difficulty understanding numbers, calculation, quantity, or mathematical reasoning despite appropriate instruction.

### **Mating**

Sexual pairing or reproduction between organisms that permits exchange or transfer of gametes and may lead to fertilization.

### **Matrix Band**

A thin dental strip placed around a prepared tooth to temporarily replace a missing wall and contain restorative material while a filling or crown buildup is formed.

### **Matrix Metalloproteinase**

An enzyme that cuts proteins in the extracellular matrix surrounding cells. It helps with normal tissue repair and remodeling, but excessive or poorly controlled activity can promote inflammation, tissue destruction, or cancer spread.

### **Maturation**

The process through which cells, tissues, organs, or a person develop and become fully functional. Normal maturation may involve changes in size, structure, ability, hormone activity, and response to the environment.

### **Maturation Disorders Of Erythropoiesis**

Anemia categories caused by ineffective development

---

or maturation of red-cell precursors in the bone marrow. They include disorders of hemoglobin synthesis, such as iron deficiency and thalassemia, and disorders of nuclear maturation, such as vitamin B12 or folate deficiency and myelodysplasia.

### **Mature Onset Diabetes**

Older term; see *Type 2 Diabetes Mellitus*.

### **Mature Teratoma**

A germ-cell tumor composed of well-differentiated tissues from one or more embryonic germ layers; many are benign, but behavior depends on the site, age, and tumor type, and some can undergo malignant transformation.

### **Maxilla**

The paired bones forming the upper jaw and supporting the upper teeth. They also form parts of the cheeks, nose, eye sockets, hard palate, and the air-filled sinuses behind the cheeks.

### **Maxillary**

Relating to the maxilla, the upper jawbone. It supports the upper teeth, forms part of the eye socket and roof of the mouth, and contains the maxillary sinuses. Problems may include tooth infection, sinusitis, or facial fractures.

### **Maxillary Sinus**

The maxillary sinus is a paired, air-filled cavity in the maxilla beneath each orbit. It drains into the nasal cavity and may become inflamed or infected in sinusitis.

### **Maxillary Sinusitis**

Inflammation or infection of the air-filled spaces behind the cheekbones. It may cause cheek or tooth pain, blocked nose, thick nasal discharge, reduced smell, cough, or fever and can follow a cold, allergy, or dental infection.

### **Maxillofacial**

Relating to the jaw, face, mouth, and associated bones and soft tissues. Maxillofacial care may in-

clude treatment of injuries, congenital differences, infections, and tumors.

### **Maxillofacial Prosthesis**

An artificial replacement for part of the face, jaw, mouth, nose, eye area, or ear after injury, cancer surgery, or a birth difference. Examples include devices that close a palate opening or replace an eye, nose, or ear.

### **Maximal Breathing Capacity**

The greatest volume of air that can be moved in and out of the lungs in a specified period during maximal effort; it is related to but distinct from standard spirometric measures.

### **Maximum**

The maximum is the greatest value in a set of measurements or the highest level reached by a variable, such as a recorded temperature, blood pressure, or drug concentration.

### **Maximum Plasma Concentration**

The highest measured plasma concentration after a dose.  $C_{max}$  and the time to reach it are used to describe absorption rate, peak-related toxicity, and comparative pharmacokinetics.

**Alternate Names:**

$C_{max}$

### **Maximum Thickness**

Maximum thickness is the greatest measured thickness of a structure or specimen. It may be reported in imaging, pathology, or assessment of a lesion, organ, or tissue layer.

### **Maximum Tolerated Dose**

The highest dose of a drug that produces acceptable, manageable toxicity without causing unacceptable adverse effects, determined through dose-escalation studies, most commonly in Phase I clinical trials. The MTD identifies the dose range for subsequent efficacy studies.

### **Mayer-Rokitansky-Küster-Hauser**

---

## Syndrome

A condition present from birth in which the uterus and much of the vagina do not develop, although the ovaries and usual female puberty changes develop normally. It often first becomes noticeable when menstruation does not begin. Kidney, hearing, or spine abnormalities may also occur. Treatment and fertility options depend on the individual's anatomy and goals.

### Alternate Names:

MRKH Syndrome; Müllerian Agenesis; Vaginal Agenesis

## Mayer-Rokitansky-Kuster-Hauser Syndrome

A congenital disorder of female reproductive tract development (Müllerian agenesis) characterized by congenital absence or severe hypoplasia of the uterus and upper two-thirds of the vagina in an individual with a 46,XX karyotype, normal functional ovaries, and normal secondary sexual characteristics, presenting as primary amenorrhea.

### Alternate Names:

MRKH Syndrome; Müllerian Agenesis

## May-Hegglin Anomaly

An inherited MYH9-related disorder with macrothrombocytopenia, giant platelets, and characteristic leukocyte inclusions; hearing, kidney, or cataract abnormalities may occur.

## Mayo Scissors

Strong surgical scissors designed to cut fascia, muscle, sutures, and other tough materials. Their broad blades provide force and control, and they may be straight or curved for different surgical tasks.

## Maytansine

Maytansine is a potent antimetabolic compound that binds tubulin and inhibits microtubule assembly. Its derivatives are used as cytotoxic payloads in some antibody-drug conjugates.

## May-Thurner Syndrome

Compression of the left pelvic vein by the right pelvic artery, which can slow blood flow and increase

the risk of a clot or swelling in the left leg.

## Maze Procedure

A heart operation for atrial fibrillation. The surgeon creates carefully placed lines of scar tissue in the upper heart chambers to block abnormal electrical signals and help restore a regular rhythm. It may be performed with other heart surgery; recurrence, bleeding, stroke, or the need for a pacemaker are possible.

### Alternate Names:

Cox-Maze Procedure; Surgical Ablation for Atrial Fibrillation; Maze Operation

## Mazindol

Mazindol is a sympathomimetic appetite suppressant that affects central catecholamine signaling. It has been used for short-term obesity treatment in some countries and may cause stimulant-related adverse effects.

## MB ChB

Abbreviation for Bachelor of Medicine and Bachelor of Surgery, a medical degree awarded in many countries.

## MCAD Deficiency

An inherited disorder in which the body cannot properly break down certain medium-chain fats for energy.

## MCA Peak Systolic Velocity

An ultrasound measurement of blood speed in a major artery of the fetus's brain. An unusually high value may suggest fetal anemia and can help guide monitoring or further testing during pregnancy.

## McArdle Disease

A rare inherited muscle disorder in which muscles cannot release usable energy from stored sugar during exercise. Activity can cause early muscle pain, cramps, exhaustion, or dangerous muscle breakdown with dark urine. Some people improve after slowing down briefly and restarting gently, called the second-wind phenomenon. Avoiding sudden strenuous exercise helps prevent attacks.



---

**Alternate Names:**

GSD V; Myophosphorylase Deficiency; Glycogen Storage Disease Type V

**McBurney Point**

A specific anatomical surface point situated one-third of the distance along an imaginary line drawn from the right anterior superior iliac spine (ASIS) to the umbilicus, corresponding to the anatomical base of the vermiform appendix and the site of maximal tenderness in acute appendicitis.

**Alternate Names:**

McBurney's Point; Appendiceal Surface Landmark

**McBurney Sign**

Tenderness when pressing on the lower right side of the belly about halfway between the belly button and the hip bone, which is a key sign of an infected appendix (appendicitis).

**McBurney's Point**

A location in the lower-right abdomen where tenderness may occur when the appendix is inflamed. Pain there can support a diagnosis of appendicitis but cannot confirm it by itself.

**Mcc Gene**

A gene that helps regulate cell growth and is considered a tumour-suppressor gene. Loss or alteration of its function has been found in some colorectal and other cancers, but the finding alone does not diagnose cancer.

**McCune-Albright Syndrome**

A condition caused by a change affecting hormone signals in some body cells. It may cause irregular skin coloring, fragile or unevenly growing bones, and early puberty or other hormone problems.

**McDonald Cerclage**

A common operation that places a strong stitch around the cervix through the vagina. It may help keep the cervix closed when there is a high risk of very early birth; timing and suitability depend on the pregnancy history and examination.

**MCH**

The average amount of hemoglobin in one red blood cell, reported in a complete blood count. Hemoglobin carries oxygen, and the result helps classify some types of anemia.

**MCHC**

The average concentration of hemoglobin inside red blood cells, reported in a complete blood count. It helps clinicians assess and classify anemia together with the other blood-count results.

**MCL Injury**

Abbreviation; see *Medial Collateral Ligament Injury*.

**MC Lymphoma**

Abbreviation; see *Mantle Cell Lymphoma*.

**McMurray Test**

A knee examination in which the leg is flexed, rotated, and extended to place stress on the menisci. Pain or a palpable click may suggest a meniscal lesion but is not diagnostic alone.

**McRoberts Maneuver**

A positioning maneuver for managing shoulder dystocia where the mother's legs are hyperflexed onto her abdomen, with thighs against the abdomen. This flattens the sacrum, increases the anteroposterior diameter of the pelvis, and rotates the symphysis pubis superiorly. Often combined with suprapubic pressure.

**MCRPC**

Abbreviation; see *Metastatic Castration-Resistant Prostate Cancer*.

**MCV**

Mean corpuscular volume: the calculated average volume of a red blood cell, derived from hematocrit and red-cell count and reported as part of a complete blood count.

---

## MDMA

3,4-Methylenedioxymethamphetamine, a psychoactive stimulant and hallucinogen commonly known as ecstasy or molly; it can produce serious cardiovascular, neurologic, and temperature-regulation effects.

## MDMA Toxicity

Serotonergic and sympathomimetic toxicity from MDMA, with agitation, hyperthermia, diaphoresis, tachycardia, hypertension, jaw clenching, seizures, rhabdomyolysis, and potentially severe hyponatremia from excess water intake. Treatment includes rapid cooling, benzodiazepines, electrolyte assessment, and careful management of complications.

## MDS

Abbreviation; see *Myelodysplastic Neoplasms*.

## Mead Acid

An omega-9 fatty acid made when the body lacks enough essential omega-6 and omega-3 fatty acids. Increased levels can support laboratory assessment of essential-fatty-acid deficiency, although results must be interpreted with nutrition and illness history.

## Meadow Saffron

*Colchicum autumnale*, a plant containing colchicine; ingestion can cause severe gastrointestinal illness, multiorgan toxicity, and death.

## Mean

The mean is the arithmetic average, calculated by adding all values and dividing by the number of values. It is commonly used to summarize clinical measurements but can be distorted by extreme values.

## Mean Arterial Pressure

An estimate of the average pressure pushing blood through the body's organs during each heartbeat. Clinicians use it with other findings to judge whether organs are receiving enough blood.

## Mean Cell Volume

The average volume of a red blood cell, calculated from the hematocrit and red-cell count and reported as MCV in a complete blood count.

## Mean Corpuscular Hemoglobin Concentration

The average concentration of oxygen-carrying hemoglobin inside red blood cells. It is part of a complete blood count and helps classify the cause of anemia.

## Mean Corpuscular Volume

The average size of red blood cells, reported in a complete blood count. Small cells or large cells can point toward different causes of anemia and must be interpreted with the other blood results.

## Mean Diameter

Mean diameter is the average diameter of a structure or lesion, usually calculated from one or more measured dimensions. The method used should be specified when measurements are clinically compared.

## Mean Thickness

Mean thickness is the average thickness calculated from multiple measurements of a tissue, structure, or specimen. It is used in imaging, pathology, and physiologic measurements.

## Measles

A highly contagious viral infection causing fever, cough, runny nose, red eyes, and a rash that spreads from the face downward. It can cause pneumonia, brain inflammation, or other serious complications.

## Measles Virus

A negative-sense RNA virus that causes measles, a highly contagious respiratory illness. The virus is transmitted by respiratory droplets and is one of the most infectious pathogens known. Symptoms include fever, cough, coryza, conjunctivitis, and rash.

---

### Measurable Residual Disease

Cancer cells that remain after treatment and can be detected with a sensitive, validated test. The amount may be too small to see on scans or by routine examination, but it can help estimate relapse risk and guide further treatment.

### Measurement

Measurement is the quantitative determination of a physical, biologic, or clinical characteristic using a defined method and unit. Interpretation depends on accuracy, precision, and the clinical context.

### Meatal

Relating to a meatus, a natural opening or passage in the body. It commonly refers to the opening of the urethra, the ear canal, or one of the passages inside the nose. Meatal narrowing can obstruct urine or air flow.

### Meatal Stenosis

Narrowing of the urethral meatus, which may cause a deflected or narrow urinary stream, spraying, dysuria, or difficulty emptying the bladder.

### Meatus

A natural opening or passage into or out of a body structure. Examples include the opening of the urethra, the ear canal, and the small passages inside the nose.

### Meatuses

Natural openings or passages in the body. In the nose, the meatuses are channels beneath the nasal folds that help air move through the nose and allow drainage from the sinuses and tear ducts.

### Mebendazole

A medicine used to treat several intestinal worm infections. It disrupts structures the worms need to absorb nutrients and survive; treatment choices and dosing depend on the parasite, age, pregnancy, and local guidance.

### Mebrofenin Scan

A nuclear-medicine scan that follows a tracer through the liver and bile ducts. It can assess gallbladder function, bile blockage, leaks, or selected causes of jaundice and abdominal pain.

### Mecamylamine

Mecamylamine is an orally active ganglionic nicotinic-receptor blocker that reduces sympathetic nerve transmission and lowers blood pressure. It has been used for severe hypertension and may cause postural hypotension, dizziness, constipation, and dry mouth.

### Mecarzole

Mecarzole is a synonym for carbendazim, a benzimidazole fungicide that interferes with fungal microtubule formation. It is an agricultural chemical rather than a routine human medicine and may be toxic if exposure occurs.

### ME/CFS

Abbreviation; see *Myalgic Encephalomyelitis*.

### Mechanical Circulatory Support

Machines or pumps that help the heart move blood when it cannot pump enough on its own. They may be used temporarily during severe illness or longer term while awaiting, or instead of, a heart transplant.

### Mechanical Insufflation-Exsufflation

A cough-assist device that gives a breath in followed quickly by suction-like pressure out. It helps people with weak breathing muscles clear mucus, especially in neuromuscular disease or spinal-cord injury.

### Mechanical Ventilation

Breathing support from a machine that pushes air and oxygen into the lungs and helps remove carbon dioxide. It may be delivered through a mask or a breathing tube when illness or injury prevents adequate breathing.

---

## Mechanical Ventilator

A mechanical ventilator is a machine that supports or replaces spontaneous breathing by delivering controlled or assisted breaths, usually through a mask or an endotracheal tube. Settings are adjusted to oxygenation, ventilation, mechanics, and patient needs.

## Mechanism Of Death

The physiologic or biochemical derangement that causes death (the final failure) as opposed to the underlying disease or injury (cause of death) and the circumstances (manner). Examples: ventricular fibrillation, cerebral hemorrhage, sepsis, respiratory failure, exsanguination, cardiac tamponade.

## Mechanism Of Toxicity

The way a substance causes injury or illness in the body. It may interfere with an enzyme or receptor, damage cell membranes or DNA, disrupt energy production, or trigger an abnormal immune response. Understanding the mechanism helps guide prevention and treatment.

## Mechanoreceptor

A sensory receptor that detects physical forces such as touch, pressure, stretch, vibration, sound, or movement. It converts mechanical stimulation into electrical signals carried by nerves to the brain or spinal cord.

## Mechlorethamine

Mechlorethamine is an alkylating antineoplastic agent that cross-links DNA and inhibits cell division. Topical gel is used for cutaneous T-cell lymphoma; systemic use is limited by toxicities such as myelosuppression and tissue injury.

## Meckel Diverticulectomy

Surgical removal of a Meckel diverticulum, usually performed when it causes bleeding, inflammation, obstruction, perforation, or is found incidentally in a high-risk setting.

## Meckel Diverticulum

A small pouch present from birth on the lower small

intestine, caused by a leftover connection from early development. It often causes no symptoms, but it can bleed, become inflamed, or cause blockage or twisting of the bowel. A Meckel scan may help find it when bleeding is suspected; treatment depends on symptoms and complications.

### Alternate Names:

Meckel's Diverticulum; Vitelline Duct Remnant; Omphalomesenteric Diverticulum; Congenital Ileal Diverticulum

## Meckel-Gruber Syndrome

A severe inherited birth disorder that causes major abnormalities of the brain, kidneys, and limbs. Typical findings include a sac of brain tissue at the back of the skull, enlarged cystic kidneys, and extra fingers or toes. Poor lung development is common, and affected pregnancies usually do not survive to birth.

### Alternate Names:

Meckel Syndrome; Dysencephalia Splanchnocystica

## Meckel's Diverticulum

Alternate name; see *Meckel Diverticulum*.

## Meclizine

Meclizine is a first-generation antihistamine with anticholinergic and vestibular-suppressant effects. It is used to prevent or treat motion sickness and vertigo and may cause drowsiness, dry mouth, and blurred vision.

## Meclofenamate Overdose

Taking too much meclufenamate, a nonsteroidal anti-inflammatory drug, may cause nausea, vomiting, stomach pain or bleeding, drowsiness, kidney injury, seizures, or metabolic acidosis. Urgent medical or poison-control advice is needed after a suspected overdose.

## Meconium

The thick, dark first stool of a newborn, composed of intestinal secretions, shed cells, mucus, and swallowed amniotic material.

---

## Meconium Aspiration

Breathing meconium, a newborn's first stool, into the airways before or around birth. It can block airflow and inflame the lungs, causing breathing difficulty and low oxygen.

## Meconium Aspiration Syndrome

Breathing problems in a newborn after meconium, the first stool, enters the airways before or during birth. It can block airways and inflame the lungs, causing rapid breathing, low oxygen, and need for respiratory support.

## Meconium Ileus

A blockage of a newborn's small intestine by unusually thick first stool. It is strongly associated with cystic fibrosis and may cause a swollen abdomen, green vomiting, and failure to pass stool soon after birth.

## Meconium Peritonitis

Inflammation of the tissue lining a fetus's abdomen after a hole in the intestine allows first stool to escape before birth. It can cause abdominal fluid or calcium deposits and may be linked to bowel blockage or cystic fibrosis.

## Meconium-Stained Amniotic Fluid

Green or yellow amniotic fluid caused by passage of fetal stool before birth. It may occur with a mature or distressed fetus and increases the risk of breathing problems after delivery; newborns are assessed and supported as needed.

## MECP2 Duplication Syndrome

A rare X-linked genetic disorder caused by extra copies or increased dosage of the MECP2 gene. It often causes developmental delay, low muscle tone, seizures, recurrent infections, and breathing or feeding problems, especially in boys.

## Medial

Medial means toward or situated nearer the body's midline; it contrasts with lateral and is used to describe anatomical position, movement, or imaging findings.

## Medial Collateral Ligament Injury

A stretch or tear of the strong ligament on the inner side of the knee, often caused by a blow to the outer knee or a twisting injury. It causes inner-knee pain and swelling; severe tears may make the knee unstable.

## Medial Cuneiform

The largest of three small bones in the inner middle part of the foot. It joins the navicular bone and the first toe bone, helps support the arch, and forms part of the joint that can be injured in a Lisfranc injury.

## Medial Epicondylitis

Alternate medical name; see *Golfer's Elbow*.

## Medial Gastrocnemius Strain

Overstretching or tearing of the medial head of the gastrocnemius muscle, often during sudden acceleration or push-off. It causes abrupt calf pain, tenderness, swelling, and pain with walking or ankle movement.

## Medial Geniculate Nucleus

A specialized thalamic relay nucleus that receives ascending auditory inputs from the inferior colliculus via the brachium of the inferior colliculus and projects auditory radiations to the primary auditory cortex in the superior temporal gyrus (Brodmann areas 41 and 42).

### Alternate Names:

MGN; Medial Geniculate Body

## Medial Lemniscus

A prominent ascending sensory white matter tract in the brainstem formed by the decussating internal arcuate fibers from the contralateral gracile and cuneate nuclei of the medulla, conveying conscious proprioception, vibration, and discriminative fine touch to the ventral posterolateral (VPL) nucleus of the thalamus.

## Medial Medullary Syndrome

A brainstem stroke syndrome (Dejerine syndrome)

---

caused by infarction of the anteromedial medulla oblongata (typically due to occlusion of the anterior spinal artery or vertebral artery), characterized by ipsilateral hypoglossal nerve (CN XII) palsy (tongue deviates toward lesion), contralateral hemiparesis sparing the face, and contralateral loss of vibration and proprioception.

**Alternate Names:**

Dejerine Anterior Bulbar Syndrome; Inferior Alternating Hemiplegia

**Medial Meniscus**

The medial meniscus is a crescent-shaped fibrocartilage pad between the medial femoral condyle and medial tibial plateau. It distributes load and contributes to knee stability; tears may cause joint-line pain, swelling, or locking.

**Medial Rotation**

Turning a limb toward the body's midline. For example, turning the thigh so the knee and foot point inward is medial rotation; the movement is assessed at joints such as the shoulder and hip.

**Medial Tibial Stress Syndrome**

Alternate name; see *Shin Splints*.

**Median**

Located in or relating to the middle plane or midpoint of a structure. In medicine, it describes anatomy, measurements, statistical values, nerve positions, or locations between two sides.

**Median Arcuate Ligament Syndrome**

Compression of the celiac artery and nearby nerves by the median arcuate ligament, potentially causing post-meal abdominal pain and weight loss.

**Alternate Names:**

Dunbar Syndrome; MALS

**Median Nerve**

A major nerve running from the neck through the arm and forearm to the hand. It helps bend the wrist and fingers, moves the thumb, and provides feeling to the thumb, index, middle, and part of

the ring finger; compression can cause carpal-tunnel symptoms.

**Median Nerve Compression**

Alternate name; see *Carpal Tunnel Syndrome*.

**Median Neuropathy**

Dysfunction of the median nerve caused by compression, trauma, inflammation, or another lesion. Numbness, tingling, pain, or weakness may affect the thumb, index, middle, and radial half of the ring finger or related muscles.

**Median Plane**

A sagittal plane that passes directly through the midline of the body, dividing it into equal right and left halves.

**Median Rhomboid Glossitis**

A smooth, red, diamond-shaped depapillated area on the dorsum of the tongue caused by chronic *Candida* infection, usually asymptomatic and managed with antifungal therapy.

**Median Survival**

The time by which half of the people in a study have experienced the specified event, such as death. It is a summary measure of survival in a defined population.

**Mediastinal Cyst**

A mediastinal cyst is a fluid- or tissue-filled cystic lesion in the mediastinum. Common types include bronchogenic, thymic, and pericardial cysts; symptoms depend on size and compression, and imaging guides evaluation.

**Mediastinal Disease**

A disorder affecting structures in the mediastinum, including the heart, great vessels, thymus, lymph nodes, nerves, and tumors or cysts within the central chest.

**Mediastinal Emphysema**

Mediastinal emphysema, or pneumomediastinum,



---

is the presence of free air in the mediastinum. It can follow alveolar rupture, trauma, surgery, or esophageal injury and may cause chest pain, neck crepitus, or respiratory compromise.

### **Mediastinal Hemorrhage**

Bleeding into the mediastinum, commonly caused by trauma, vascular rupture, surgery, or invasive procedures, and potentially causing airway or cardiovascular compression.

### **Mediastinal Lymphadenopathy**

Enlargement of lymph nodes in the mediastinum. It may occur with infection, inflammation, lymphoma, or cancer that has spread from another organ.

### **Mediastinal Lymph Nodes**

Lymph nodes in the central chest that filter lymph from the lungs, heart, esophagus, and nearby tissues.

### **Mediastinal Lymphoma**

Lymphoma involving lymph nodes or other lymphoid tissue in the mediastinum. It may cause cough, chest pressure, breathlessness, superior vena cava obstruction, or constitutional symptoms; CT, PET, biopsy, and immunophenotyping establish the type and extent.

### **Mediastinal Mass**

An abnormal growth or collection of tissue in the mediastinum, the central compartment of the chest. Causes include tumors, enlarged lymph nodes, cysts, and vascular abnormalities.

### **Mediastinal Neoplasm**

A mediastinal neoplasm is an abnormal growth in the central space of the chest between the lungs; it may arise from lymph nodes, thymus, nerves, or other tissues.

### **Mediastinal Pleura**

The mediastinal pleura is the portion of parietal pleura lining the lateral surfaces of the mediastinum. It separates the mediastinal contents from the pleural cavity and reflects around the lung root.

### **Mediastinal Shift**

Mediastinal shift is displacement of the mediastinal structures away from their usual position. It may result from increased pressure on one side, such as a tension pneumothorax, or loss of volume on the other, such as lung collapse.

### **Mediastinal Tumor**

A mediastinal tumor is a mass in the central chest compartment between the lungs and may be benign or malignant.

### **Mediastinitis**

Mediastinitis is inflammation or infection of the central chest compartment, often after esophageal rupture, surgery, trauma, or spread of infection.

### **Mediastinoscopes**

Mediastinoscopes are slender endoscopic instruments used during mediastinoscopy to inspect the mediastinum and obtain tissue samples, especially from mediastinal lymph nodes. The procedure is performed through a small incision near the sternum.

### **Mediastinoscopy**

A surgical procedure that accesses the mediastinum through a small incision near the sternum to inspect or biopsy mediastinal lymph nodes or masses. It is one option for staging lung cancer when less-invasive sampling is nondiagnostic or unsuitable.

### **Mediastinoscopy With Biopsy**

A procedure using an instrument inserted through a small incision near the sternum to inspect the mediastinum and obtain lymph-node or mass samples for diagnosis and staging.

### **Mediastinotomy**

A surgical opening into the mediastinum to examine or reach structures in the central chest.

### **Mediastinum**

The central part of the chest between the lungs. It contains the heart, major blood vessels, windpipe,

---

food pipe, thymus, nerves, and lymph nodes.

### **Mediators Of Inflammation**

Molecules that start, strengthen, limit, or coordinate inflammation. They include histamine, prostaglandins, cytokines, and other signals that change blood vessels, attract immune cells, cause symptoms, and support healing.

### **Medical Abbreviations**

Alternate name; see *Common medical abbreviations*.

### **Medical Cannabis And Cannabidiol**

Cannabis and cannabidiol are substances from the cannabis plant used in some medical settings. Their effects and risks vary by product and dose; they may cause sleepiness, impaired judgment, interactions, or dependence and do not treat every condition.

### **Medical Condition**

A state affecting health or body function, whether temporary, long-lasting, symptomatic, or silent. A medical condition may result from disease, injury, inherited traits, infection, or normal biological changes.

### **Medical Device**

An instrument, implant, machine, software program, or other product used to diagnose, prevent, monitor, or treat disease or injury. Examples include pace-makers, syringes, imaging equipment, and blood-glucose meters.

### **Medical Entomology**

The study of insects and related arthropods that cause disease or transmit disease-causing organisms to humans, including their biology, ecology, physiology, and genetics.

### **Medical Ethics**

Moral principles that guide healthcare decisions, including respect for a person's choices, helping rather than harming, fairness, privacy, and honest communication. Decisions consider the patient's wishes, likely benefits and harms, professional duties,

and relevant law.

### **Medical Examiner**

A physician who investigates sudden, unexpected, violent, or suspicious deaths. The examiner may order an autopsy, determine the medical cause of death, and provide information for the official death certificate; exact duties vary by location.

### **Medical History**

A medical history is a structured account of a person's current concerns, past illnesses and operations, medications, allergies, family history, social history, and relevant exposures. It helps guide examination, diagnosis, and treatment.

### **Medical Illness**

The subjective experience of being unwell, which may or may not correspond to an identifiable disease.

### **Medical Jargon**

Specialized words and phrases used by healthcare professionals. If they are not explained, patients may misunderstand their condition, test results, or treatment, so clinicians should use plain language whenever possible.

### **Medical Jurisprudence**

The branch of law (and its study) that deals with the rights and duties of physicians and the legal aspects of medical practice, distinct from but overlapping with forensic medicine.

### **Medical Malpractice**

Legal tort occurring when a healthcare provider fails to exercise the standard of reasonable care, skill, or diligence expected of a prudent practitioner of similar standing, resulting in injury, harm, or death to a patient.

#### **Alternate Names:**

Clinical Malpractice; Medical Malpractice

### **Medical Marijuana**

Cannabis used under medical supervision to relieve

---

symptoms or support treatment of selected conditions. Products contain cannabinoids such as THC or cannabidiol; effects, interactions, impairment, and dependence risk vary by product and dose.

### **Medical Nutrition Therapy**

Personalized food and nutrition care provided by a qualified dietitian to manage a health problem. It may support people with diabetes, kidney disease, digestive disorders, cancer, obesity, or malnutrition.

### **Medical Officer**

A physician appointed to provide clinical care or medical oversight for an institution, service, or organization.

### **Medical Physicist**

A specialist who applies physics to medical imaging, radiation treatment, and radiation safety. Medical physicists check equipment, calculate radiation doses, and help ensure that scans and cancer treatments are accurate and as safe as possible.

### **Medical Physics**

The application of physics to medicine, including diagnostic imaging, radiation therapy, radiation safety, and physiological measurement.

### **Medical Pin**

A medical pin is a slender metal fixation device inserted into bone or across a joint to stabilize a fracture or support orthopedic treatment. Pins may be temporary or permanent and require monitoring for infection and displacement.

### **Medical Practitioner**

A licensed physician qualified to diagnose, prevent, and treat illness.

### **Medical Privacy**

Protection of information that can identify a patient and control over its sharing. It includes keeping health records secure, limiting access to appropriate people, and obtaining permission when disclosure is not otherwise allowed.

### **Medical Procedure**

A planned clinical action or intervention performed by healthcare professionals to diagnose, measure, monitor, treat, or prevent a health condition, or to improve bodily function and structure. Medical procedures range in complexity from non-invasive evaluations (such as an electrocardiogram, ultrasound, or blood pressure check) to minimally invasive tests (such as drawing blood, inserting a catheter, performing an endoscopy, or taking a needle biopsy) and major invasive surgeries (such as tumor resections, joint replacements, or organ transplants). Procedures are classified by their primary clinical purpose: diagnostic (identifying the underlying cause of disease), therapeutic or curative (treating an illness, correcting a defect, or removing damaged tissue), palliative (relieving symptoms, obstruction, or pain to improve quality of life), or preventive/prophylactic (such as screening colonoscopies to remove precancerous polyps). Depending on the level of invasiveness and tissue involvement, procedures may be performed awake, with local numbing, under procedural sedation, or under general anesthesia, requiring strict sterile technique to reduce risks of bleeding, infection, and tissue injury.

#### **Alternate Names:**

Clinical Procedure; Healthcare Procedure; Medical Intervention; Surgical Procedure

### **Medical Records**

Written or electronic information about a person's health, including symptoms, examinations, test results, diagnoses, medicines, procedures, and follow-up. Accurate records help healthcare professionals coordinate safe care and track changes over time.

### **Medical Symbol**

A commonly recognized sign representing medicine, especially the Rod of Asclepius: a staff with one serpent. The winged, two-serpent caduceus is often used incorrectly as a medical symbol.

### **Medical Tattooing**

Medical tattooing (permanent makeup, micropigmentation) covers depigmented skin (vitiligo, vitiligo camouflage), restores areolas after mastectomy,

---

and masks scars. Trichopigmentation simulates hair on bald scalp. Medical tattooing uses specialized pigments and techniques.

### **Medical Tests**

Diagnostic procedures used to screen for, confirm, or monitor disease through analysis of biologic specimens, imaging, physiologic measurements, or genetic material.

### **Medicinal**

Relating to a substance or treatment used for the prevention, diagnosis, relief, or treatment of disease.

### **Medicine**

A substance or treatment used to prevent, diagnose, relieve, or treat disease. Medicines may be prescription, over-the-counter, biological, or traditional, and all can have benefits, risks, and interactions.

### **Medicines For Back Pain**

Medicines for back pain may include selected anti-inflammatory drugs, acetaminophen, muscle relaxants, or other treatments depending on the cause and patient risks.

### **Medicines For Osteoporosis**

Osteoporosis medicines reduce bone loss or increase bone formation and may include bisphosphonates, denosumab, or anabolic treatments chosen according to fracture risk.

### **Medicines For Sleep**

Medicines for sleep may help short-term insomnia but can cause dependence, falls, confusion, or next-day impairment; treatment should also address the cause of poor sleep.

### **Medicolegal**

Pertaining to the application of medical knowledge to legal questions and the intersection of medicine and the law. Covers forensic pathology, toxicology, psychiatry, and clinical issues: informed consent, malpractice, medical records, disability, personal injury, occupational disease, and public health law.

### **Meditation**

Meditation is a practice of intentionally directing attention, often toward the breath, a sensation, or an object, while observing thoughts without judgment. It may promote relaxation and stress management but does not replace indicated medical treatment.

### **Mediterranean Anemia**

Alternate name; see *Thalassemia*.

### **Mediterranean Diet**

An eating pattern rich in vegetables, fruits, beans, whole grains, nuts, olive oil, and fish, with less red and processed meat and added sugar. It can support cardiovascular health as part of an overall balanced diet.

### **Mediterranean Fever**

An older name for brucellosis, a bacterial infection that can cause prolonged fever, sweats, fatigue, and joint or muscle pain.

### **Medium**

A substance or condition through which something occurs; in microbiology, a culture medium supports the growth of microorganisms.

### **MEDLARS**

Medical Literature Analysis and Retrieval System, an electronic bibliographic system developed by the U.S. National Library of Medicine to index and retrieve biomedical literature; it preceded MEDLINE.

### **Medroxyprogesterone**

Medroxyprogesterone is a progestogen used for contraception, abnormal uterine bleeding, endometriosis, and hormone therapy; depot injections may cause irregular bleeding, weight change, delayed fertility return, and reduced bone density.

### **Medroxyprogesterone Acetate**

Medroxyprogesterone acetate is a synthetic progestin used in hormonal contraception, treatment of endometriosis or abnormal uterine bleeding, and hormone therapy. It suppresses ovulation and produces

---

progestational effects; formulation-specific risks and contraindications apply.

### **Medulla**

The inner or central part of an organ, such as the renal medulla or adrenal medulla.

### **Medulla Oblongata**

The lower part of the brainstem, joined to the spinal cord. It helps control breathing, heart rate, blood pressure, swallowing, coughing, vomiting, and other automatic functions.

### **Medullary**

Relating to the medulla or inner portion of an organ.

### **Medullary Carcinoma**

A cancer whose cells grow in broad groups and are surrounded by many immune cells. The term is used most often for cancers of the breast or thyroid, and its behavior depends on the organ and tumor findings.

### **Medullary Cavity**

The hollow space inside the shaft of a long bone. It contains bone marrow and blood vessels.

### **Medullary Sponge Kidney**

A condition present from birth in which tiny urine-collecting tubes inside the kidneys are widened. Urine minerals may form stones or deposits, causing blood in the urine, repeated infections, pain, or no symptoms.

### **Medullary Thyroid Carcinoma**

A cancer of hormone-producing thyroid cells that make calcitonin. It may occur alone or as part of an inherited syndrome; blood tests, imaging, and a biopsy help confirm it.

### **Medulloblastoma**

A fast-growing cancer that usually begins in the cerebellum, the part of the brain involved in balance and coordination. It mainly affects children and may cause headache, vomiting, unsteady walking,

or other neurologic problems.

### **Medulloepithelioma**

A medulloepithelioma is a rare embryonal tumor arising most often from the nonpigmented ciliary epithelium of the eye in children. It may cause visual symptoms, neovascular glaucoma, or a visible mass and is treated according to its extent and pathology.

### **Mees' Lines**

True leukonychia characterized by single or multiple paired, non-blanching transverse white bands across the nail plate that migrate distally as the nail grows. Classically associated with acute arsenic or thallium poisoning, Hodgkin lymphoma, and severe systemic illness.

#### **Alternate Names:**

Aldrich-Mees Lines; Mees Lines

### **Mefenamic Acid**

A nonsteroidal anti-inflammatory medicine used for short-term pain, including menstrual pain. It can irritate or bleed from the stomach, affect kidney function, and increase cardiovascular risk, especially with prolonged use or in people with risk factors.

### **Mefloquine**

Mefloquine is an antimalarial drug used for prevention and treatment of susceptible malaria. It has a long half-life but can cause neuropsychiatric, vestibular, and cardiac adverse effects, so patient-specific contraindications and warnings are important.

### **Megabecquerel**

A megabecquerel (MBq) is a unit of radioactivity equal to one million nuclear disintegrations per second. It is used to state the activity of radiopharmaceuticals and other radioactive materials.

### **Megacalycosis**

Megacalycosis is congenital, nonobstructive dilatation of the renal calyces, usually with otherwise preserved renal architecture and function. It is often incidental but must be distinguished from obstructive hydronephrosis.

---

## Megacolon

Abnormal enlargement and dilation of the colon, which may be congenital, acquired, toxic, or inflammatory.

## Megakaryocyte

A very large cell in the bone marrow that makes platelets, the small blood parts that help stop bleeding. It grows long extensions into nearby blood vessels, where pieces break off and enter the bloodstream as platelets.

## Megakaryocytopoiesis

Megakaryocytopoiesis is the formation and maturation of megakaryocytes in the bone marrow, followed by platelet production. It is regulated in large part by thrombopoietin and other hematopoietic signals.

## Megaloblast

An unusually large, immature cell in bone marrow that should develop into a red blood cell. It usually reflects faulty DNA production, commonly caused by vitamin B12 or folate deficiency.

## Megaloblastic Anaemia

Macrocytic anemia caused by defective DNA synthesis, most often from vitamin B12 or folate deficiency.

## Megaloblastic Anemia

Anemia caused by impaired DNA synthesis, usually related to vitamin B12 or folate deficiency, with abnormally large red blood cells.

## Megalocephaly

Abnormally large head size, which may be familial, developmental, metabolic, or associated with brain disease.

## Megamonas Funiformis

Megamonas funiformis is an anaerobic bacterium of the genus Megamonas, commonly associated with the intestinal microbiota. Its detection in a clinical specimen requires interpretation alongside symptoms, specimen quality, and other microbiologic findings.

## Megamonas Hypermegale

Megamonas hypermegale is an anaerobic bacterium in the genus Megamonas, primarily associated with the gastrointestinal microbiota. Its presence in a specimen is interpreted in relation to the site, clinical symptoms, and other microbiologic results.

## Megasphaera

Megasphaera is a genus of anaerobic bacteria found in the oral cavity and gastrointestinal and genital microbiota. Species identification may be relevant in microbiome studies or when recovered from a clinical specimen.

## Megaureter

An abnormally widened ureter, the tube that carries urine from the kidney to the bladder. It may result from a blockage, urine flowing backward, or abnormal ureter development and can cause infection, kidney swelling, or reduced kidney function.

## Megestrol

Megestrol is a synthetic progestin used in some settings to treat hormone-responsive cancers and, in certain formulations, anorexia or severe weight loss. It can cause glucocorticoid-like effects, thromboembolic risk, and adrenal suppression.

## Megestrol Acetate

Megestrol acetate is the acetate salt of the progestin megestrol. Depending on formulation and indication, it is used for advanced endometrial or breast cancer and for appetite stimulation in selected patients with severe weight loss.

## Meglitinides

Short-acting diabetes medicines that encourage the pancreas to release insulin around mealtimes. They lower blood sugar but can cause low blood sugar and weight gain, so doses are usually linked to meals and skipped when a meal is skipped.

## Megrim

Older term; see *Migraine*.



---

## Meibomian Gland Dysfunction

A chronic, diffuse abnormality of the modified sebaceous tarsal glands of the eyelids, characterized by terminal duct obstruction and qualitative or quantitative alterations in lipid secretion (meibum), representing the leading cause of evaporative dry eye syndrome.

### Alternate Names:

MGD; Posterior Blepharitis

## Meibomian Glands

Sebaceous glands in the eyelids that produce the oily layer of the tear film and help prevent tear evaporation.

## Meibomianitis

Meibomianitis is inflammation or blockage of oil-producing eyelid glands, causing irritated eyelid margins, dry eye, crusting, or recurrent styes.

## Meige Syndrome

A movement disorder causing repeated involuntary eyelid squeezing and muscle spasms around the mouth, jaw, or tongue. Symptoms may worsen with stress, speaking, chewing, or touch.

## Meigs Syndrome

The benign clinical triad of an ovarian fibroma (or other benign solid pelvic ovarian tumor), ascites, and pleural effusion (typically right-sided), which resolves completely and permanently following surgical resection of the ovarian neoplasm.

## Meigs' Syndrome

A rare combination of a usually benign ovarian fibroma, fluid in the abdomen, and fluid around the lungs. The fluid often improves after the ovarian tumor is removed, but cancer and other causes must be excluded.

## Meiosis

A specialized cell division that produces haploid eggs or sperm from a diploid precursor cell and creates genetic variation through recombination and chromosome assortment.

## Meiotic Recombination

Meiotic recombination is exchange of DNA between homologous chromosomes during meiosis, usually through crossing over. It creates new combinations of alleles and contributes to genetic variation.

## Meissner Corpuscle

An encapsulated, rapidly adapting mechanoreceptor located in the dermal papillae of glabrous (hairless) skin (fingertips, palms, soles, lips), sensitive to low-frequency vibrations (30 to 50 Hz), light touch, and dynamic skin flutter.

### Alternate Names:

Tactile Corpuscle

## Meissner Plexus

A network of unmyelinated autonomic nerve fibers and parasympathetic postganglionic neuronal cell bodies located within the submucosa of the gastrointestinal tract, primarily regulating local glandular secretions, mucosal absorption, and submucosal blood flow.

### Alternate Names:

Submucosal Plexus

## Meissner's Corpuscles

Small touch sensors in hairless skin, especially the fingertips, palms, and soles. They help the nervous system detect gentle contact, texture, and movement across the skin.

## MEK Inhibitor

A targeted medicine that blocks MEK proteins in the MAPK signaling pathway, which can slow growth of some cancers with pathway changes. Side effects vary and may include rash, diarrhea, heart effects, and eye problems.

## Melancholia

An older term for severe depression, often describing pervasive sadness, loss of pleasure, psychomotor change, and biological symptoms.

## Melancholic Features

A pattern of severe depression marked by almost

---

complete loss of pleasure, very low mood, major changes in movement or sleep, poor appetite or weight loss, excessive guilt, and symptoms that are often worse in the morning.

### Melanin

Melanin is a pigment produced by melanocytes that gives skin, hair, and eyes their colour and absorbs ultraviolet radiation; its amount and distribution influence pigmentation disorders and melanoma biology.

### Melanoblast

A precursor cell that develops into a melanocyte and produces pigment after migrating into the skin and other tissues.

### Melanocyte

A melanocyte is a pigment-producing cell derived from the neural crest. It makes melanin in specialized organelles called melanosomes and transfers the pigment to neighboring keratinocytes, helping protect skin from ultraviolet radiation.

### Melanocytes

Pigment-producing cells in the skin and hair follicles. They make melanin, the pigment that gives skin and hair much of their color and helps protect skin cells from ultraviolet radiation.

### Melanocytic Nevus

Common moles formed by collections of pigment-producing melanocytes. Most are harmless, but a mole that changes in size, shape, color, or symptoms should be assessed for possible melanoma.

### Melanocytic Nevus

A melanocytic nevus is a usually benign, localized proliferation of melanocytes that appears as a macule, papule, or nodule. Changes in size, color, border, or symptoms may require assessment for atypia or melanoma.

### Melanocytic CNS Tumor

A tumor showing melanocytic differentiation within

the CNS or meninges, ranging from benign melanocytoma to malignant melanoma. Evaluation includes imaging, histology, and molecular assessment to determine behavior.

### Melanoderma

Increased pigmentation or darkening of the skin caused by increased melanin production or an increased number of melanocytes.

### Melanoma

Melanoma is a malignant tumor of melanocytes, most often arising in skin but also in eye or mucosal sites; early detection matters because it can invade and metastasize.

#### Alternate Names:

Cutaneous Melanoma; Malignant Melanoma

### Melanoma Of The Eye

A cancer arising from pigment-producing cells in the eye, most often the uveal tract. It may cause blurred vision, flashes, a growing dark spot, or no symptoms at first and requires specialist eye assessment.

### Melanoma Skin Cancer

Alternate name; see *Melanoma*.

### Melanoma Surveillance

Ongoing examination of the skin and, when appropriate, the eyes or other previously affected sites after melanoma or in people at increased risk. It may include self-checks, clinician examination, dermoscopy, photographs, and evaluation of new or changing lesions.

### Melanoma Treatment

Treatment depends on the tumor's depth, spread, molecular findings, overall health, and previous treatment. Localized melanoma is usually removed surgically; selected cases need lymph-node assessment or additional medicine. Disease that has spread may require immune-based or targeted treatment, surgery, radiation, or combinations.

---

## Melanonychia

Brown, gray, or black pigmentation of a nail plate caused by increased melanin or pigment-producing cells. It may affect several nails from medicines or systemic disease, but a new or changing single-nail band requires assessment for nail-unit melanoma.

## Melanophore

A melanophore is a pigment-containing cell, especially one in lower vertebrates, whose melanin granules can disperse or aggregate to alter visible coloration. In mammals, melanocytes are the principal melanin-producing cells.

## Melanism

Abnormal accumulation or increased production of melanin in skin, mucosa, an organ, or other tissue.

## Melanism Coli

Melanism coli is dark pigmentation of the colon lining caused by lipofuscin accumulation, classically associated with prolonged stimulant-laxative use. It is usually benign and may fade after stopping the trigger, but the underlying constipation and medication use should be reviewed.

## Melanosome

A melanosome is a specialized organelle in a melanocyte in which melanin is synthesized, stored, and transported. Melanosomes are transferred to keratinocytes and influence skin, hair, and eye pigmentation.

## Melasma

An acquired, chronic pigmentary disorder characterized by symmetric, hyperpigmented, light-to-dark brown macules and patches on sun-exposed areas of the face (malar, forehead, upper lip, chin). It primarily affects women of childbearing age and individuals with darker skin types (Fitzpatrick types III-V).

### Alternate Names:

Facial Melanism; Mask of Pregnancy

## MELAS Syndrome

A mitochondrial disorder involving encephalopathy, lactic acidosis, and stroke-like episodes, often beginning before adulthood and caused by pathogenic mitochondrial DNA variants.

## Melatonin

A hormone produced mainly by the pineal gland that helps coordinate sleep and wake timing. Darkness usually increases its release, while light suppresses it; supplements can cause drowsiness and medicine interactions.

## Melatonin Receptor

A cell-surface receptor activated by the hormone melatonin. The main types, MT1 and MT2, help regulate the body's sleep-wake clock and seasonal rhythms and are targets of some sleep medicines.

## Melatonin Receptor Agonists

Medicines that act like melatonin at receptors involved in the body's sleep-wake clock. They may help some people fall asleep, but choice of treatment depends on the cause of insomnia, other medicines, and health conditions.

## MELD

Model for End-Stage Liver Disease, a laboratory-based score used to estimate severity of advanced liver disease and help prioritize adults for liver transplantation. Traditional versions use bilirubin, creatinine, and INR; current implementations may include additional variables.

## MELD-Na Score

An updated mathematical scoring model derived from the Model for End-Stage Liver Disease (MELD), incorporating serum sodium alongside serum creatinine, total bilirubin, and international normalized ratio (INR). It improves 90-day mortality prediction in patients with decompensated cirrhosis and severe hyponatremia, utilized internationally as the primary objective scoring metric for donor liver transplant allocation.

### Alternate Names:

MELD-Sodium Score; Sodium-Adjusted MELD

---

## MELD Score

A medical score estimating the severity and short-term risk of death from advanced liver disease. It uses laboratory results and helps guide urgency for liver transplantation; the calculation may vary as official versions are updated.

## Melena

Black, sticky, often foul-smelling stool caused by digested blood, usually from bleeding in the stomach or upper intestine. It can signal serious bleeding and requires prompt medical evaluation.

## Melibiose

A disaccharide composed of galactose and glucose, produced by hydrolysis of raffinose and metabolized by some microorganisms.

## Melio

An older shortened form related to melioidosis, the infection caused by *Burkholderia pseudomallei*.

## Melioidosis

An infectious disease caused by *Burkholderia pseudomallei*, usually acquired from contaminated soil or water through inoculation or inhalation. It may cause pneumonia, skin infection, abscesses, sepsis, or chronic disease and is more severe in people with diabetes or other risk factors.

## Melitensis

A species designation, especially *Brucella melitensis*, a bacterium that causes brucellosis and is associated with goats and sheep.

## Melitten

Melitten, more commonly spelled melittin, is the principal peptide toxin in honeybee venom. It disrupts cell membranes and contributes to pain, inflammation, hemolysis, and other effects of envenomation.

## Melkersson-Rosenthal Syndrome

A rare neuromucocutaneous disorder characterized by the classic clinical triad of recurrent orofacial

edema (especially granulomatous cheilitis of the lips), relapsing peripheral facial nerve (CN VII) palsy, and a fissured tongue (lingua plicata).

### Alternate Names:

Cheilitis Granulomatosa Triad; Fissured Tongue and Facial Palsy Syndrome

## Melorheostosis

A rare sclerosing bone dysplasia producing irregular, flowing cortical hyperostosis on imaging. It may cause pain, stiffness, limb deformity, restricted movement, contractures, or soft-tissue fibrosis and can affect one or several contiguous bones.

## Meloxicam

Meloxicam is a nonsteroidal anti-inflammatory drug that inhibits cyclooxygenase enzymes and reduces prostaglandin synthesis. It is used for pain and inflammation, but can cause gastrointestinal, renal, cardiovascular, and hypersensitivity adverse effects.

## Melphalan

A chemotherapy medicine that damages cancer cells by interfering with their DNA. It is used for selected blood cancers and other cancers, but can also suppress bone marrow and increase infection or bleeding risk.

## Memantine

An NMDA receptor antagonist (uncompetitive, low-to-moderate affinity) used for moderate to severe Alzheimer disease, often added to an acetylcholinesterase inhibitor. Blocks excessive glutamate-mediated excitotoxicity while preserving normal synaptic transmission. Once or twice daily oral.

## Memantine Hydrochloride

Memantine hydrochloride is an NMDA-receptor antagonist used to help manage moderate-to-severe Alzheimer disease. It reduces pathologic glutamate-mediated excitotoxicity; common adverse effects include dizziness, headache, constipation, and confusion.

## Membrane

A thin sheet of tissue or material that surrounds

---

a cell, organ, or body cavity. Membranes can protect structures, separate spaces, allow selected substances to pass, and produce or absorb fluids.

### **Membrane Channel**

A membrane channel is a protein-forming pore across a cell membrane that permits selected ions or molecules to pass down an electrochemical or concentration gradient. Many channels are gated by voltage, ligands, or mechanical forces.

### **Membrane Fusion**

Membrane fusion is merging of two lipid bilayers into one continuous membrane. It is required for processes such as synaptic vesicle release, enveloped-virus entry, fertilization, and intracellular vesicle trafficking.

### **Membrane Lipids**

Fat-like molecules that form the main structure of cell membranes. They create a flexible barrier, help cells communicate, and provide attachment sites for proteins; abnormal lipid composition can affect signaling and cell function.

### **Membrane Microdomain**

A membrane microdomain is a small, specialized region of a cell membrane with a distinct composition of lipids and proteins. It can organize receptors, signaling molecules, transporters, and cytoskeletal interactions.

### **Membrane Part**

A membrane part is a structural or molecular component of a biological membrane, such as a lipid, protein, carbohydrate, channel, or receptor. These components determine membrane integrity, transport, signaling, and cell recognition.

### **Membrane Potential**

The electrical potential difference across a cell membrane, approximately -70 mV in most neurons (resting membrane potential). Established by the unequal distribution of ions (primarily K<sup>+</sup>, Na<sup>+</sup>, Cl<sup>-</sup>) across the membrane and selective permeability.

### **Membrane Protein**

A protein attached to or embedded in a cell membrane. It may transport substances, receive signals, act as an enzyme, join cells, anchor the cytoskeleton, or help the immune system identify cells.

### **Membrane Raft**

A membrane raft is a cholesterol- and sphingolipid-enriched membrane microdomain that helps cluster selected proteins and signaling molecules. Raft organization is dynamic and may change during receptor signaling or membrane trafficking.

### **Membranoproliferative Glomerulonephritis**

A pattern of kidney-filter damage that can cause blood and protein in the urine, swelling, and reduced kidney function. Causes include immune disorders, infections, and some blood cancers.

### **Membranous**

Relating to, consisting of, or resembling a membrane. In medicine, the term may describe a thin tissue layer or a disease pattern involving such a layer.

### **Membranous Labyrinth**

A continuous system of communicating, endolymph-filled epithelial sacs and ducts lodged within the bony labyrinth of the inner ear, comprising the cochlear duct (hearing) and the vestibular apparatus (utricle, saccule, and three semicircular ducts for balance and equilibrium).

### **Membranous Nephropathy**

A kidney disease in which the filtering units are injured by immune proteins deposited along their delicate filter. It often causes large amounts of protein in the urine and swelling and may be linked to autoimmune disease, infection, medicines, or cancer.

### **Memory**

The ability to learn, store, and recall information. Memory depends on several connected areas of the brain and can be affected by sleep problems, stress, medicines, injury, illness, aging, or brain disorders.

---

## Memory Cell

A long-lasting B cell or T cell formed after an infection or vaccination. If the same germ is encountered again, memory cells respond faster and more strongly, helping prevent illness or reduce its severity.

## Memory Disorder

Impairment of encoding, storage, retrieval, or recognition of information that exceeds normal variation and may result from neurodegeneration, injury, vascular disease, medication, or other causes.

## Memory Impairment

Memory impairment is reduced ability to learn, store, retrieve, or recognize information. It may be transient or persistent and can result from normal aging, medications, sleep problems, neurologic disease, psychiatric illness, or other medical conditions.

## Memory Loss

Reduced ability to recall past information, form new memories, or both. Causes include normal aging, medicines, head injury, seizures, stroke, infection, substance use, neurocognitive disorders, and psychological conditions; sudden memory loss can occur with serious illness.

## Memory Span

The number of items, such as words or digits, that a person can briefly retain and recall, usually in the original order; it is a measure of short-term or working memory.

## Memory T Cells And B Cells

Long-lived immune cells that remember earlier exposures and respond faster when they meet the same target again. They support lasting protection after infection or vaccination, though protection may fade.

### Alternate Names:

Memory Lymphocytes

## MEN 2

Abbreviation; see *Multiple Endocrine Neoplasia Type 2*.

## Menarche

The first menstrual period, marking the onset of menstruation. Its timing varies with genetics, nutrition, health, and pubertal development; markedly early or delayed menarche may require assessment.

## Mendelian Inheritance

Inheritance of traits determined by single genes and transmitted according to dominant, recessive, or sex-linked patterns.

## Mendelism

The principles of inheritance based on Gregor Mendel's work, including segregation of alleles and independent assortment of genes; Mendelian traits may follow autosomal or X-linked dominant or recessive patterns.

## Mendelson Syndrome

A chemical pneumonitis caused by inhalation of acidic stomach contents, classically during anesthesia or childbirth. It can cause cough, breathing difficulty, and impaired oxygenation.

## Menetrier Disease

A rare stomach disorder in which the lining develops unusually large folds and makes excess mucus. Protein may leak from the stomach, lowering the protein level in the blood and causing swelling; stomach acid is often reduced. It usually affects the upper part of the stomach.

### Alternate Names:

Hypertrophic Protein-Losing Gastropathy; Giant Hypertrophic Gastritis

## Meniere Disease

An inner-ear disorder causing repeated attacks of spinning dizziness, usually with fluctuating hearing loss, ringing in the ear, and a feeling of pressure or fullness. Attacks often last from 20 minutes to several hours.



---

**Alternate Names:**

Endolymphatic Hydrops; Meniere's Syndrome

**Meniere's Disease**

An inner-ear disorder causing episodes of vertigo, fluctuating hearing loss, tinnitus, and a feeling of fullness in one ear. Treatment aims to reduce attacks and protect hearing.

**Meningeal Carcinomatosis**

Spread of cancer cells to the thin coverings and fluid around the brain and spinal cord. It may cause headache, nausea, weakness, seizures, vision changes, or problems with movement and needs specialist evaluation.

**Meningeal Leukemia**

Meningeal leukemia is leukemia involving the membranes or fluid around the brain and spinal cord, potentially causing headache, weakness, or cranial-nerve problems.

**Meningeal Lymphoma**

Meningeal lymphoma is lymphoma affecting the membranes or fluid spaces around the brain and spinal cord.

**Meningeal Neoplasm**

A meningeal neoplasm is an abnormal growth arising in or near the membranes surrounding the brain and spinal cord.

**Meninges**

Three protective layers surrounding the brain and spinal cord. Fluid between two of the layers cushions the nervous system; bleeding or infection in these layers can be dangerous.

**Meningioma**

A usually slow-growing tumor arising from cells associated with the meninges. Most are benign, but some are atypical or malignant and can cause symptoms by compressing the brain, spinal cord, or cranial nerves.

**Meningioma Pathology**

A usually extra-axial tumor showing meningotheelial cells in syncytial sheets, whorls, or other patterns, sometimes with psammoma bodies. Histologic grade reflects mitotic activity, invasion, and other atypical or anaplastic features.

**Meningism**

Meningism is a syndrome of neck stiffness, headache, and sometimes photophobia or vomiting that resembles meningeal irritation without necessarily indicating meningitis. Serious causes, including infection or subarachnoid hemorrhage, require prompt evaluation.

**Meningismus**

Neck stiffness and other signs of irritation around the brain and spinal cord without proven inflammation of those coverings. It can occur with fever, severe headache, dehydration, bleeding, or other illnesses and needs evaluation.

**Meningitis**

Inflammation of the meninges around the brain and spinal cord, caused by bacteria, viruses, fungi, parasites, medicines, or immune disease. Fever, headache, neck stiffness, photophobia, vomiting, altered mental status, or seizures may occur, but the classic triad is often incomplete. Suspected bacterial meningitis is an emergency requiring blood tests and cerebrospinal-fluid evaluation when safe.

**Meningocele**

A birth defect in which the protective coverings around the brain or spinal cord bulge through an opening in the skull or spine, forming a fluid-filled sac. It may cause nerve problems and needs specialist assessment.

**Meningocele Repair**

Surgery to close an opening in the spine and return the protective coverings of the spinal cord to their proper position. It aims to prevent injury or infection; nerve, bladder, bowel, movement, or fluid problems may remain or develop.

---

## Meningococcal Disease

A serious infection caused by *Neisseria meningitidis* that can produce meningitis or rapidly spreading bloodstream infection. Fever, severe headache, stiff neck, confusion, vomiting, or a purple rash may occur.

## Meningococcal Infection

An infection caused by the bacterium *Neisseria meningitidis*. It may remain mild or cause life-threatening meningitis or bloodstream infection, with fever, headache, stiff neck, vomiting, confusion, or a purple rash developing quickly.

## Meningococcal Meningitis

Acute bacterial meningitis caused by *Neisseria meningitidis*, often accompanied by bloodstream infection; it is a medical emergency requiring immediate antibiotics and public-health measures.

## Meningococcemia

Meningococcemia is bloodstream infection with *Neisseria meningitidis*, causing fever, rash, shock, clotting failure, and potentially fatal sepsis.

## Meningococcus

*Neisseria meningitidis*, a bacterium that can cause meningitis or rapidly spreading bloodstream infection. It spreads through respiratory secretions and may cause fever, headache, rash, shock, or death.

## Meningoencephalitis

Inflammation of both the meninges surrounding the brain and the brain tissue itself, usually caused by infection or autoimmune disease. Fever, headache, neck stiffness, confusion, seizures, or reduced consciousness requires urgent medical assessment.

## Meningomyelocele

A birth defect in which the protective coverings and spinal-cord tissue bulge through an opening in the spine. It can cause weakness or paralysis of the legs and bladder or bowel problems.

## Meniscal Allograft Transplantation

Surgery placing a donated meniscus into a knee after substantial loss of the person's own meniscus. It may reduce pain in carefully selected younger patients, but healing is variable and infection, stiffness, graft failure, or later arthritis can occur.

## Meniscal Injuries

Tears or other damage to the knee menisci, which are crescent-shaped pieces of cartilage that help distribute load and stabilize the knee. Symptoms may include joint-line pain, swelling, clicking, or restricted motion. Diagnosis is based on examination and, when needed, imaging.

### Alternate Names:

Meniscal Injury

## Meniscal Tear

A split in one of the two rubbery cartilage pads that cushion the knee. It may follow a twisting injury or gradual wear and can cause pain along the joint line, swelling, clicking, catching, locking, or a feeling that the knee gives way.

## Meniscectomy

An operation that removes part or all of a damaged meniscus, the rubbery pad that cushions the knee. Surgeons usually remove only the torn portion when possible, because preserving meniscus helps protect the joint from later wear.

## Meniscus

A crescent-shaped fibrocartilage structure in the knee, or a curved fluid surface in a container.

## Meniscus Tear

Alternate name; see *Torn Meniscus*.

## Menkes Disease

A rare inherited disorder in which the body cannot transport copper properly. Low copper in the brain and other tissues can cause poor growth, developmental decline, seizures, weak muscles, and sparse, brittle, or unusually twisted hair.

---

### Menkes Kinky Hair Syndrome

An X-linked disorder of copper transport caused by ATP7A dysfunction. It produces progressive neurodegeneration, seizures, developmental delay, hypopigmented or brittle kinky hair, connective-tissue abnormalities, and failure to thrive.

### Menogaril

Menogaril is a semisynthetic anthracycline derivative of nogalamycin. It intercalates into DNA and inhibits topoisomerase II, thereby impairing DNA replication and repair; it was investigated as an antineoplastic agent.

### Menometrorrhagia

Excessive uterine bleeding occurring both at expected menstrual times and at irregular intervals; the older term corresponds broadly to heavy and irregular menstrual bleeding and warrants evaluation for underlying causes.

### Menopausal

Relating to menopause or the years around the permanent end of menstrual periods. Hormone changes may cause hot flashes, sleep problems, mood changes, vaginal dryness, and increased bone-loss risk.

### Menopausal Hormone Therapy

Treatment with estrogen, with progestogen when needed to protect the uterus, for troublesome hot flashes, night sweats, or vaginal symptoms during menopause. It may affect clot, stroke, breast-cancer, and other risks, so treatment is individualized.

### Menopausal Hot Flashes

Alternate name; see *Hot Flashes*.

### Menopause

The permanent end of menstrual periods because of loss of ovarian activity, usually diagnosed after 12 consecutive months without menstruation. Hormonal changes may cause hot flashes, night sweats, vaginal symptoms, sleep problems, or other symptoms.

### Menopause and Mood Disorders

Mood symptoms can occur during the menopausal transition because of hormonal fluctuation, sleep disruption, vasomotor symptoms, life stress, or an independent depressive or anxiety disorder. Persistent, severe, or suicidal symptoms require mental-health assessment; treatment is individualized.

### Menopause And Perimenopause

Perimenopause is the transition before menopause, when menstrual periods and hormone levels become irregular. Menopause is reached after 12 consecutive months without a period when no other cause explains it.

### Menopause Status

Menopause status describes whether a person is premenopausal, perimenopausal, or postmenopausal. Natural menopause is clinically defined after 12 consecutive months without menstruation when no other cause explains the amenorrhea.

### Menopause Syndrome

A nontechnical term for the group of symptoms that may occur during the menopausal transition, such as hot flashes, night sweats, sleep problems, mood changes, vaginal dryness, and menstrual irregularity. Menopause itself is a normal life stage, not a disease.

### Menorrhagia

Older term; see *Heavy Menstrual Bleeding*.

### Menotropin

Menotropin is a urinary-derived preparation containing follicle-stimulating and luteinizing hormone activity, used to stimulate follicular development or spermatogenesis; ovarian hyperstimulation and multiple pregnancy are important risks.

### Menses Duration

Menses duration is the number of days of menstrual bleeding in a cycle, counted from the first day of bleeding until bleeding stops. Persistently prolonged, heavy, or irregular bleeding warrants clinical assessment.

---

## Menstrual

Relating to menstruation or the menstrual cycle, including the monthly shedding of the uterine lining and the bleeding that follows. Cycle length and bleeding vary, but very heavy, painful, irregular, or unexpected bleeding may need evaluation.

## Menstrual Cramps

Cramping pain in the lower abdomen during or around menstruation, medically called dysmenorrhea. Pain may radiate to the back or thighs and may occur with nausea, diarrhea, headache, or fatigue. Severe, worsening, or newly developed pain requires evaluation for causes such as endometriosis or fibroids.

### Alternate Names:

Dysmenorrhea; Period Pain; Painful Menstruation

## Menstrual Cycle

The recurring hormonal and uterine changes that prepare the body for possible pregnancy. Cycle length and bleeding pattern vary among individuals and across life stages.

## Menstrual Disorders

Conditions involving abnormal menstrual timing, regularity, duration, amount of bleeding, or pain. They include absent or infrequent periods, irregular cycles, unusually heavy or prolonged bleeding, painful periods, and bleeding between periods or after sex. Causes include pregnancy, hormonal or ovulatory disorders, fibroids, endometriosis, thyroid disease, medicines, structural abnormalities, and perimenopause; persistent, heavy, or unexpected bleeding requires assessment.

## Menstruation

The periodic shedding of the endometrial lining as part of the menstrual cycle when pregnancy does not occur. The timing, duration, and amount of bleeding vary with age, hormonal state, medicines, and health conditions.

## Menstruation Disturbance

Alternate name; see *Menstrual Disorders*.

## Mental

Relating to thinking, memory, emotion, behavior, or other functions of the mind. It does not by itself mean that a condition is psychiatric.

## Mental Age

An estimate of cognitive functioning expressed as the age level at which a standardized test score is typical; it is not the same as chronological age.

## Mental Disorder

Alternate singular form; see *Mental Disorders*.

## Mental Disorders

Health conditions involving a clinically significant disturbance in cognition, emotion regulation, or behavior that causes distress, impairment, or both. Examples include mood, anxiety, psychotic, neurodevelopmental, trauma-related, personality, and substance use disorders. A mental-health concern or symptom does not necessarily constitute a mental disorder.

## Mental Disorders In Children

Mental disorders in children involve persistent, developmentally atypical disturbances in mood, thinking, behavior, or functioning that cause clinically significant distress or impairment. Symptoms must be interpreted in relation to the child's developmental stage and distinguished from normal variation, temporary reactions, and age-appropriate behavior.

## Mental Fatigue

Mental fatigue is a subjective feeling of reduced mental energy, concentration, or cognitive endurance after sustained effort or illness. Persistent or severe symptoms may accompany sleep disorders, depression, neurologic disease, medication effects, or other medical conditions.

## Mental Health

Mental health encompasses emotional, psychological, and social well-being, affecting how individuals think, feel, behave, and handle stress, relationships, and decision-making.

---

## Mental Health Crisis

A period of severe emotional, behavioral, or psychological distress in which a person's usual ability to cope or maintain safety is overwhelmed. It may involve suicidal thoughts, severe agitation, psychosis, substance intoxication, or inability to care for basic needs; it may or may not meet criteria for a psychiatric emergency.

## Mental Illness

A broad, commonly used term for mental disorders and related conditions affecting mood, thinking, behavior, perception, or daily functioning. See *Mental Disorders*.

## Mental Illness In Children

Broad, commonly used term; see *Mental Disorders In Children*.

## Mental Process

A mental process is a cognitive or affective activity of the mind, such as perception, attention, memory, reasoning, language, emotion, or decision-making.

## Mental Recall

Mental recall is retrieval of previously learned or stored information from memory. It may be assessed by asking a person to remember facts, events, words, or experiences after a delay.

## Mental Status Examination

A structured part of a clinical interview that assesses a person's appearance, behavior, speech, mood, thinking, perceptions, attention, memory, awareness, judgment, and understanding of their condition.

## Mental Status Testing

A structured assessment of appearance, behavior, alertness, orientation, attention, memory, language, thought, perception, mood, insight, and judgment.

## Mentation

The process of thinking, perceiving, reasoning, and other mental activity.

## Menthol

A mint-derived or synthetic compound producing a cooling sensation and used in some topical and respiratory preparations.

## Menthol Poisoning

Poisoning after swallowing, inhaling, or having substantial skin exposure to menthol. It may cause nausea, abdominal pain, dizziness, drowsiness, agitation, seizures, coma, or breathing problems.

## Mepacrine

An older antimalarial and antiprotozoal medicine, also called quinacrine, now used only in selected indications.

## Meperidine

An opioid analgesic, also called pethidine, whose use is limited by adverse effects and an active neurotoxic metabolite.

## Meperidine Hydrochloride

Meperidine hydrochloride is the hydrochloride salt of the opioid analgesic meperidine. It produces analgesia through opioid-receptor activity and carries risks of sedation, respiratory depression, seizures, dependence, and serotonin toxicity with interacting drugs.

## Meperidine Hydrochloride Overdose

Taking too much meperidine hydrochloride, an opioid, can cause extreme sleepiness, pinpoint pupils, slow or stopped breathing, low blood pressure, seizures, or coma. Delayed toxicity and serotonin toxicity are possible, so emergency assessment is required.

## Mephenesin

An older centrally acting muscle relaxant with sedative effects. It is rarely used now and can impair alertness, coordination, and breathing when combined with other sedatives.

## Mephentermine

A stimulant medicine formerly used in selected situ-

---

ations to raise dangerously low blood pressure. It increases activity of certain adrenaline-like receptors and can cause fast heartbeat, anxiety, or abnormal heart rhythms.

### **Mephenytoin**

Mephenytoin is a hydantoin anticonvulsant formerly used to treat seizures. It is metabolized partly to ethotoin and has limited modern use because of potentially serious adverse effects, including hypersensitivity and blood dyscrasias.

### **Mepivacaine**

A local anesthetic medicine that temporarily blocks pain signals from nerves. It is injected for numbing skin, dental areas, individual nerves, or larger body regions during procedures.

### **Meprobamate**

An older medicine that reduces anxiety and causes sleepiness. It can cause dependence, dangerous breathing and nervous-system depression, and overdose, especially with alcohol or other sedatives, so it is rarely used today.

### **Meprobamate Overdose**

Taking too much meprobamate can cause severe drowsiness, confusion, poor coordination, low blood pressure, slowed breathing, coma, or death. Alcohol and other sedatives increase the danger.

### **Mepyramine**

An antihistamine that blocks H1 receptors and may cause sedation; it is also called pyrilamine.

### **Meralgia Paresthetica**

Burning, tingling, or numbness over the outer thigh caused by compression of the lateral femoral cutaneous nerve.

#### **Alternate Names:**

Burning Thigh Pain; Bernhardt-Roth Syndrome

### **Merbromin Poisoning**

Poisoning from merbromin, an older mercury-containing antiseptic. Swallowing or repeated ex-

posure may injure the mouth, stomach, kidneys, or nervous system; the amount and route determine risk and require poison-control or medical assessment.

### **Mercaptopurine**

A thiopurine antimetabolite used in selected leukemias and inflammatory disorders, requiring monitoring for marrow and liver toxicity.

### **Mercurialism**

Long-term poisoning caused by mercury exposure. It may damage the nervous system, kidneys, digestive tract, mouth, or mental health, with effects depending on the mercury form, dose, and exposure duration.

### **Mercuric Chloride Poisoning**

Severe poisoning from mercuric chloride, a corrosive inorganic mercury salt. It can burn the mouth and digestive tract and cause vomiting, bloody diarrhea, shock, kidney failure, or neurologic injury; emergency treatment is required.

### **Mercuric Oxide Poisoning**

Poisoning from mercuric oxide, an inorganic mercury compound that can irritate or burn tissues and damage the kidneys and nervous system. Swallowing or inhalation can cause severe poisoning.

### **Mercury**

A metal used in some industrial and laboratory products. Its chemical symbol is Hg; mmHg denotes millimeters of mercury, a unit of pressure. Exposure can harm the nervous system, kidneys, lungs, or digestive tract, so it must be handled and disposed of safely.

#### **Alternate Names:**

Hg

### **Mercury Poisoning**

Illness caused by exposure to elemental, inorganic, or organic mercury. Effects depend on the form and dose and may include nerve, kidney, lung, or digestive injury; urgent advice is needed after significant exposure.



---

## Meridian

A reference line or plane; in traditional acupuncture it denotes a proposed pathway for qi, while in anatomy it may describe a longitudinal line.

## Merkel Cell Carcinoma

An aggressive neuroendocrine skin cancer arising from Merkel cells. It presents as a rapidly growing, painless, red-violet nodule on sun-exposed skin of elderly patients. Approximately 80 percent are associated with Merkel cell polyomavirus. Risk factors include immunosuppression, UV exposure, and fair skin.

### Alternate Names:

MCC; Neuroendocrine Carcinoma of the Skin

## Merkel Cells

Specialized touch-sensing cells in the deepest layer of the skin that help detect light touch and sustained pressure.

## Mermaid Syndrome

An extremely rare congenital anomaly in which the legs are fused to varying degrees, also called sirenomelia.

## Meropenem

Meropenem is a broad-spectrum carbapenem antibiotic that inhibits bacterial cell-wall synthesis and treats severe susceptible infections, including meningitis; allergy, gastrointestinal effects, seizures, and resistance are possible.

## Meropenem-Vaborbactam

An intravenous antibiotic combination used for some serious infections caused by bacteria resistant to many other antibiotics. The choice depends on laboratory susceptibility testing, infection site, kidney function, and local guidance.

## MERRF Syndrome

A rare inherited disorder affecting how cells produce energy. It commonly causes muscle jerks, seizures, balance problems, and muscle weakness, and may also affect hearing, vision, or other organs.

## MERS-CoV

Middle East respiratory syndrome coronavirus, the virus that causes MERS. Infection can cause fever and respiratory illness ranging from mild disease to severe pneumonia, especially in people with other health conditions.

## Merthiolate Poisoning

Poisoning from merthiolate, commonly referring to the mercury-containing antiseptic thimerosal. Small skin exposures usually cause local irritation, but swallowing or repeated exposure may cause corrosive, kidney, or neurologic injury and needs advice.

## Merycism

Rumination, the repeated regurgitation and rechewing of food after eating.

## Mesalamine

Mesalamine, or 5-aminosalicylic acid, is an anti-inflammatory drug used to induce and maintain remission in ulcerative colitis and some other inflammatory bowel conditions. It acts mainly topically in the intestinal mucosa and can rarely affect renal function.

## Mesangial Cell

Specialized contractile pericyte-like cells located in the central stalk (mesangium) of the renal glomerulus between capillary loops, responsible for phagocytosing macromolecules, synthesizing mesangial matrix, and contracting in response to angiotensin II to regulate glomerular capillary filtration surface area.

## Mesangiolytic

Mesangiolytic is dissolution or disruption of the mesangial tissue within a renal glomerulus. It may occur in severe endothelial injury and can be associated with microvascular lesions and glomerular capillary aneurysmal changes.

## Mesangium

The mesangium is the supportive tissue between glomerular capillary loops, composed of mesangial cells and extracellular matrix. Mesangial cells pro-

---

vide structural support, regulate capillary flow, and clear certain substances; abnormal expansion occurs in several glomerular diseases.

### **Mescaline**

A naturally occurring psychedelic substance found in certain cacti that alters perception and may cause hallucinations and autonomic effects.

### **Mesencephalon**

The mesencephalon, or midbrain, is the part of the brainstem between the pons and diencephalon. It contains pathways and nuclei involved in eye movements, motor control, arousal, and auditory and visual reflexes.

### **Mesenchymal Cell**

A connective-tissue precursor that can develop into bone, cartilage, fat, muscle, or tendon cells.

### **Mesenchymal Chondrosarcoma**

Mesenchymal chondrosarcoma is a rare aggressive cancer containing cartilage-forming tissue, often arising in bone or soft tissue.

### **Mesenchymal CNS Tumor**

A rare tumor arising from connective or supporting tissue in or around the brain or spinal cord rather than from nerve cells. Its behavior is determined by examination of tissue, imaging, and sometimes genetic testing.

### **Mesenchyme**

Embryonic connective tissue that gives rise to much of the skeleton, connective tissue, blood vessels, and smooth muscle.

### **Mesenchymoma**

A mesenchymoma is a rare tumor containing more than one type of mesenchymal tissue, sometimes including cartilage, bone, fat, or vascular elements. Behavior depends on the site, age, histology, and presence of malignant components.

### **Mesenteric**

Relating to the mesentery, a fold of tissue supporting the intestines. It contains blood vessels, nerves, lymph vessels, and fat that help supply, drain, anchor, and protect the intestines.

### **Mesenteric Angiography**

Contrast imaging of the arteries supplying the intestines, used to evaluate mesenteric ischemia, bleeding, aneurysm, stenosis, occlusion, or vascular anatomy.

### **Mesenteric Artery**

A mesenteric artery supplies blood to portions of the intestine and associated abdominal organs. The superior and inferior mesenteric arteries are the major vessels; obstruction or embolism can cause acute mesenteric ischemia.

### **Mesenteric Artery Aneurysm**

An abnormal bulge in an artery supplying the intestines. It may cause no symptoms, but rupture can cause severe internal bleeding, and clotting or reduced blood flow can injure the bowel.

### **Mesenteric Artery Ischemia**

Inadequate blood flow through the mesenteric arteries, usually caused by embolism, thrombosis, or severe low-flow states. Acute disease presents with severe abdominal pain out of proportion to examination and can progress to bowel infarction.

### **Mesenteric Cyst**

A rare, usually noncancerous, fluid-filled sac in the tissue that supports the intestines. Small cysts may cause no symptoms, while larger ones can cause swelling, pain, infection, twisting, or bowel blockage. Imaging helps locate the cyst, and treatment depends on its size, symptoms, and cause.

#### **Alternate Names:**

Mesenteric Lymphangioma; Chylolymphatic Cyst; Enteric Mesenteric Cyst

### **Mesenteric Ischemia**

Reduced blood flow to the intestines, usually from

---

a blocked or narrowed blood vessel. Sudden blockage may cause severe abdominal pain, often with little tenderness at first, nausea, vomiting, diarrhea, or bloody stool, and is a medical emergency. Long-term narrowing may cause cramping pain after meals, fear of eating, and weight loss. Severe or untreated disease can lead to bowel death, infection, or shock.

### **Mesenteric Lymphadenitis**

Inflammation of lymph nodes in the tissue supporting the intestines, often after a viral or bacterial infection. It can cause right-sided abdominal pain, fever, nausea, and tenderness and may resemble appendicitis.

**Alternate Names:**  
Mesenteric Adenitis

### **Mesenteric Lymph Nodes**

Lymph nodes in the tissue that supports the intestines. They filter lymph from the bowel and help the immune system respond to infection or other abnormal changes.

### **Mesenteric Veins**

Mesenteric veins drain blood from the small and large intestines and carry it mainly through the superior and inferior mesenteric veins to the portal venous system. Thrombosis or impaired flow can cause intestinal congestion and ischemia.

### **Mesenteric Venous Thrombosis**

A blood clot in a mesenteric vein that impairs intestinal venous drainage and can cause abdominal pain, bowel edema, ischemia, infarction, or gastrointestinal bleeding.

### **Mesentery**

A fold of the abdominal lining that attaches the intestines to the back wall of the abdomen. It carries blood vessels, nerves, and lymph vessels to the bowel and helps hold the intestines in place.

### **Mesh**

A sheet-like implant made from synthetic or biological material that reinforces weakened tissue. It

may be used in selected hernia or pelvic-floor operations; possible problems include pain, infection, scarring, movement of the implant, or injury to nearby organs.

### **Mesial**

Located toward the body's midline; in dentistry, located toward the front or midline of a tooth. The term helps describe tooth surfaces, positions, cavities, gum disease, and dental restorations.

### **Mesial Surface**

The mesial surface is the surface of a tooth or other structure oriented toward the midline. In dentistry, mesial surfaces contact adjacent teeth and can be involved in caries, periodontal disease, or restoration margins.

### **Mesna**

A protective medicine given with certain chemotherapy drugs to reduce irritation and bleeding of the bladder. It binds harmful breakdown products in the urine and does not treat the cancer itself.

### **Mesoderm**

The middle germ layer in an embryo, which develops into muscles, bones, cartilage, blood vessels, and the kidneys.

### **Mesometrium**

The portion of the broad ligament of the uterus that extends from the lateral uterus toward the pelvic wall and contains associated vessels, nerves, and connective tissue.

### **Mesonephric**

Mesonephric means relating to the mesonephros, a temporary embryonic kidney, or to structures derived from its ducts and tubules. Mesonephric remnants can give rise to uncommon benign or malignant lesions of the genital tract.

### **Mesonephric Adenocarcinoma**

A rare HPV-independent cervical adenocarcinoma arising from mesonephric remnants, usually present-

---

ing as a cervical mass or abnormal bleeding and requiring histopathologic diagnosis.

### **Mesonephric Tumor**

A mesonephric tumor is a rare tumor arising from remnants of embryonic mesonephric ducts, usually in the female reproductive tract.

### **Mesothelial Neoplasm**

A mesothelial neoplasm is a growth arising from the lining of body cavities, such as the pleura, peritoneum, or pericardium.

### **Mesothelin**

Mesothelin is a cell-surface glycoprotein expressed mainly by mesothelial cells and overexpressed in several cancers, including mesothelioma and some ovarian and pancreatic cancers. It is being studied as a biomarker and therapeutic target.

### **Mesothelioma**

A cancer arising from the thin lining of body cavities, most often the pleura around the lungs. It is strongly associated with asbestos exposure and may cause chest pain, breathlessness, or fluid around the lung.

### **Mesothelium**

The mesothelium is a single layer of specialized cells lining serous body cavities and covering their organs, including the pleura, peritoneum, and pericardium. It produces lubricating fluid and participates in inflammation and repair.

### **Messenger RNA**

A temporary copy of genetic instructions made from DNA. It carries those instructions to cell structures that build proteins and is then broken down; some vaccines use messenger RNA to teach the immune system to recognize a target.

### **Met**

Abbreviation for methionine, an essential sulfur-containing amino acid; it is also represented by the one-letter code M and the codons AUG.

### **Metabolic**

Relating to metabolism, the chemical processes by which cells produce and use energy and materials.

### **Metabolic Acidosis**

A primary fall in plasma bicarbonate causing acidemia unless respiratory compensation or another process changes the pH. It is classified by anion gap and may result from lactic or ketoacidosis, kidney failure, toxins, or bicarbonate loss from the gastrointestinal tract or kidneys; assessment includes pH, bicarbonate, respiratory compensation, electrolytes, and the cause.

### **Metabolic Alkalosis**

A primary increase in plasma bicarbonate causing alkalemia unless respiratory compensation or another process changes the pH. Common causes include vomiting, gastric suction, diuretics, mineralocorticoid excess, and alkali administration; hypokalemia and chloride depletion are frequent, and urine chloride helps distinguish chloride-responsive from chloride-resistant forms.

### **Metabolic Bone Disease**

A group of disorders in which abnormal handling of calcium, phosphate, vitamin D, parathyroid hormone, or other factors weakens or changes bone. Examples include osteomalacia, rickets, and some disorders caused by kidney or endocrine disease.

### **Metabolic Bone Disease of Prematurity**

Reduced bone mineral density and defective postnatal skeletal mineralization in very-low-birth-weight or extremely preterm infants. Results from inadequate accretion of calcium, phosphorus, and vitamin D typically transferred during the third trimester, exacerbated by prolonged total parenteral nutrition or diuretic therapy.

#### **Alternate Names:**

Osteopenia of Prematurity; Neonatal Rickets; Metabolic Bone Disease in Preterm Infants

### **Metabolic Disease**

A disorder that changes how the body processes or uses substances such as sugars, fats, proteins, min-

---

erals, or hormones. It may be inherited or acquired and can affect energy, growth, blood chemistry, or organ function.

### **Metabolic Disorders**

Alternate plural form; see *Metabolic Disease*.

### **Metabolic Dysfunction-Associated Steatohepatitis**

Alternate name; see *Fatty Liver Disease*.

### **Metabolic Dysfunction-Associated Steatotic Liver Disease**

Alternate name; see *Fatty Liver Disease*.

### **Metabolic Equivalent of Task**

A physiological measure expressing the energy cost of physical activities, defined as the ratio of metabolic rate during activity to resting metabolic rate, standardized as the consumption of 3.5 mL of oxygen per kilogram of body weight per minute (1 MET).

**Alternate Names:**

MET

### **Metabolic Equivalents**

Units that compare the energy used during an activity with the energy used while resting. They provide a rough estimate of exercise intensity, but actual energy use varies between people.

**Alternate Names:**

METs

### **Metabolic Myopathies**

Muscle disorders caused by impaired energy metabolism, often producing exercise intolerance, cramps, weakness, muscle pain, or episodes of rhabdomyolysis.

### **Metabolic Neuropathies**

Nerve damage caused by a metabolic or systemic disorder, such as diabetes, kidney failure, vitamin deficiency, or an inherited metabolic disease. Symptoms commonly include numbness, burning pain, tingling, weakness, or problems with balance.

### **Metabolic Rate**

The speed at which the body uses energy to keep organs working and support activity. It changes with body size, age, muscle amount, hormones, illness, temperature, food intake, and exercise.

### **Metabolic Response**

A metabolic response is a change in energy use, substrate metabolism, or biochemical pathways in response to conditions such as fasting, exercise, infection, trauma, or medication. The response may be adaptive or contribute to illness when excessive or prolonged.

### **Metabolic Syndrome**

A combination of increased waist size, high blood pressure, high blood sugar, high triglycerides, or low HDL cholesterol. Together these findings increase the risk of heart disease, stroke, and type 2 diabetes.

**Alternate Names:**

Insulin Resistance Syndrome; Syndrome X

### **Metabolic Tumor Volume**

The estimated amount of tumor tissue that shows increased activity on a PET scan. It can help measure disease burden and treatment response, especially in lymphoma, but inflammation and scan technique can affect the result.

### **Metabolism**

The chemical processes cells use to turn food and medicines into energy and building materials, and to break down substances the body no longer needs.

### **Metabolite**

A substance made, changed, or used when the body processes a medicine, nutrient, or naturally occurring molecule. Measuring metabolites in blood or urine can show body function, disease, or drug effects.

### **Metacarpal**

Relating to one of the five bones forming the palm between the wrist and fingers. Metacarpal fractures can cause pain, swelling, deformity, reduced grip,

---

and difficulty moving the hand.

### **Metacarpal Bone**

A metacarpal bone is one of the five long bones of the hand between the wrist bones and the proximal phalanges. Each has a base, shaft, and head; fractures can affect alignment, grip, and joint function.

### **Metacarpals**

The five bones forming the palm, numbered from the thumb to the little finger. Each connects the wrist to a finger and helps with grip and hand movement; breaks can cause pain, swelling, deformity, or reduced function.

### **Metacarpophalangeal Joint**

A knuckle joint connecting a palm bone to a finger or thumb bone. It allows the digit to bend, straighten, and move from side to side and can be injured by impact or twisting.

### **Metacarpophalangeal Joint Dislocation**

Displacement of a finger or thumb's metacarpophalangeal joint, usually after trauma or forced hyperextension. Pain, deformity, swelling, and restricted movement occur; associated fractures, tendon injury, or neurovascular damage must be assessed.

### **Metacarpus**

The part of the hand between the wrist and fingers, containing the five metacarpal bones.

### **Metachromatic Leukodystrophy**

A rare inherited disorder in which harmful fatty substances build up in nerve cells and damage the protective covering of nerves. It causes worsening movement, thinking, speech, or vision problems, often beginning in childhood.

### **Metagonimus Yokogawai**

Metagonimus yokogawai is a small intestinal fluke acquired by eating raw or inadequately cooked freshwater fish. Infection can cause abdominal symptoms and, rarely, ectopic migration of eggs or adult worms.

### **Metal Artifact**

Distortion or loss of detail on an X-ray, CT, or MRI image caused by metal such as an implant, clip, or bullet. It can hide nearby findings, so special imaging methods or another test may be needed.

### **Metal Fume Fever**

A short-lived influenza-like illness caused by inhalation of fumes from heated metals, especially zinc; symptoms include fever, chills, headache, muscle aches, and cough.

### **Metalloendopeptidase**

A metalloendopeptidase is an endopeptidase whose catalytic activity depends on a bound metal ion, commonly zinc. It cleaves peptide bonds within proteins and includes enzymes involved in extracellular matrix remodeling and signaling.

### **Metalloporphyrins**

Metalloporphyrins are porphyrin molecules containing a centrally coordinated metal ion. Heme, which contains iron, is a biologically important example and forms the functional group of hemoglobin, myoglobin, and many enzymes.

### **Metalloprotease**

A metalloprotease is a proteolytic enzyme that requires a metal ion, commonly zinc, for catalytic activity. Metalloproteases regulate extracellular matrix turnover, cell signaling, inflammation, and tissue remodeling; dysregulation contributes to disease.

### **Metalloprotein**

A protein that contains a metal ion or binds one tightly as part of its structure or function. Iron, zinc, copper, and magnesium in metalloproteins support oxygen transport, enzyme activity, electron transfer, or regulation.

### **Metallothionein**

Metallothionein is a small, cysteine-rich protein that binds metals such as zinc, copper, and cadmium. It helps regulate essential metals and may protect cells from metal toxicity and oxidative stress.



---

## Metanephrine

Metanephrine is a metabolite of epinephrine formed mainly by catechol-O-methyltransferase. Plasma or urinary metanephrine measurements are used in evaluation for catecholamine-secreting tumors such as pheochromocytoma and paraganglioma.

## Metaphase

The stage of cell division in which chromosomes align at the cell's equatorial plane.

## Metaphase Plate

The metaphase plate is the plane at the center of a dividing cell where chromosomes align during metaphase. It is a geometric arrangement rather than a discrete anatomical structure.

## Metaphyseal Corner Fractures

Small corner or bucket-handle breaks near the growth plates of an infant's long bones. They are highly concerning for forceful injury, especially when multiple or at different healing stages, but must be interpreted with the full medical and imaging assessment.

## Metaphyseal Widening

Metaphyseal widening is enlargement of the growth-plate or metaphyseal region of a developing long bone. It may be seen in disorders such as rickets, skeletal dysplasia, infection, or healing injury and must be interpreted with imaging and clinical findings.

## Metaphysis

The wider part of a growing long bone between its shaft and end. In children it lies next to the growth plate, where new bone forms and the bone lengthens.

## Metaplasia

A usually reversible change in which one mature cell type is replaced by another that better tolerates ongoing irritation or stress. Some persistent forms may increase the risk of further abnormal changes.

## Metaplastic Carcinoma

A cancer whose cells have changed to resemble a different type of tissue. The term is used especially for an uncommon, often aggressive breast cancer; diagnosis requires examination of a tissue sample.

## Metapneumovirus

A respiratory virus that can cause cough, fever, bronchiolitis, or pneumonia, especially in infants, older adults, and people with chronic disease.

## Metastasectomy

Surgery to remove one or more cancer deposits that have spread from the original tumor. It may help selected people when the deposits are limited and removable, but benefit depends on the cancer type, location, and other treatment options.

## Metastases To The Breast

Secondary malignant deposits in breast tissue from a nonmammary primary tumor, most often melanoma, lymphoma, or carcinoma from another organ. Imaging and biopsy are needed because the treatment differs from primary breast carcinoma.

## Metastasis

The spread of cancer cells from the original tumor to a separate part of the body, where they form new tumors. Spread can occur through the blood, lymphatic system, or body cavities.

## Metastasis-Free Survival

The length of time after treatment before cancer spreads to a distant part of the body or the person dies. Researchers use it to compare treatments and estimate how well they delay spread.

## Metastasis Of Colorectal Cancer

Spread of cancer from the colon or rectum to another part of the body, most often the liver, lungs, or abdominal lining. Symptoms and treatment depend on the sites involved, number of tumors, and tumor test results.

---

### **Metastatic Adenocarcinoma**

A gland-forming cancer that has spread from where it began to distant organs such as the liver, lungs, bones, or brain. Identifying the original site is important because treatment depends on its origin and spread.

### **Metastatic Brain Tumor**

A metastatic brain tumor is cancer that has spread to the brain from a primary tumor elsewhere in the body.

### **Metastatic Breast Cancer**

Breast cancer that has spread from the breast to distant organs or tissues, such as bone, liver, lung, or brain. It is treatable, but usually considered advanced and not curable.

### **Metastatic Calcification**

Buildup of calcium salts in otherwise healthy tissues because the blood calcium level is too high. It can affect the kidneys, lungs, stomach, or blood vessels and may damage their function.

### **Metastatic Cancer**

Cancer that has spread from its original site to distant organs or tissues. The spread may occur through the blood or lymphatic system and can form new tumors. Metastatic cancer is often called stage 4 cancer, although staging depends on the cancer type. The new tumors retain the characteristics of the original cancer; for example, breast cancer that spreads to the lung remains metastatic breast cancer, not lung cancer.

#### **Alternate Names:**

Stage 4 Cancer; Advanced Metastatic Cancer; Secondary Cancer

### **Metastatic Cancer With Unknown Primary Site**

Cancer proven to have spread but for which the original primary tumor cannot be identified after appropriate evaluation. Pathology, immunohistochemistry, imaging, and molecular testing may suggest the origin and guide site-directed or empiric treatment.

### **Metastatic Castration-Resistant Prostate Cancer**

Prostate cancer that has spread and continues growing despite treatment that lowers male hormones. Care may include hormone-targeting medicines, chemotherapy, radiation, immune treatment, or medicines for bone protection.

### **Metastatic Colon Cancer**

Colon cancer that has spread beyond the bowel, most often to the liver, lungs, or abdominal lining. Symptoms and treatment depend on where it has spread, the tumor's genetic findings, and the person's overall health.

### **Metastatic Colorectal Cancer**

Alternate name; see *Colorectal Cancer*.

### **Metastatic Inguinal Lymphadenopathy**

Enlargement of lymph nodes in the groin because cancer cells have spread to them, often from the leg, genital area, lower abdomen, or anus. Examination and imaging help identify the original cancer and its extent.

### **Metastatic Lung Cancer**

Lung cancer that has spread beyond the lungs to organs such as the brain, bones, liver, or adrenal glands. Treatment depends on the cancer type, molecular findings, areas affected, and overall health.

### **Metastatic Pleural Tumor**

A cancer deposit in the thin lining around the lung that has spread from another tumor, often from the lung, breast, or digestive tract. It may cause chest pain, breathlessness, or fluid around the lung.

### **Metastin**

Metastin is an older name for kisspeptin, a peptide derived from the KISS1 gene. It stimulates hypothalamic gonadotropin-releasing hormone secretion and helps regulate puberty, reproductive function, and tumor-cell signaling.

---

## Metatarsal

Relating to one of the five long bones in the middle and front of the foot. Metatarsal bones connect the ankle bones to the toe bones and may be injured by falls, twisting, or repeated stress.

## Metatarsal Bone

A metatarsal bone is one of the five long bones of the forefoot between the tarsal bones and proximal phalanges. Metatarsal fractures and alignment abnormalities can affect weight bearing and gait.

## Metatarsalgia

Pain in the front or ball of the foot, often arising from pressure or irritation around the metatarsal heads.

## Metatarsals

The five bones forming the front half of the foot between the ankle and toes. They support body weight and help with walking; overuse or injury can cause pain or fractures.

## Metatarsal Valgus

Metatarsal valgus is outward deviation of one or more metatarsal bones relative to the foot's normal axis. It may contribute to forefoot deformity and altered load distribution and is assessed clinically and radiographically.

## Metatarsophalangeal Joint

A joint between a metatarsal bone of the foot and the first bone of a toe. It helps the toes bend and supports walking.

## Metatarsus

The part of the foot between the tarsal bones and toes, containing the five metatarsal bones.

## Metatarsus Adductus

A foot deformity in which the forefoot is angled inward relative to the hindfoot, often recognized in infancy and ranging from flexible to rigid.

## Metatarsus Varus

Metatarsus varus is inward deviation of the forefoot caused by medial angulation of the metatarsal bones. It is often recognized in infancy and may be flexible or rigid; management depends on severity and persistence.

## Metencephalon

The metencephalon is the embryonic brain region that develops into the pons and cerebellum. These structures participate in motor coordination, balance, cranial-nerve functions, and relay of signals between the brain and spinal cord.

## Meteorism

Abdominal distension caused by excessive gas in the gastrointestinal tract.

## Meter

A metric unit of length equal to 100 centimeters; medical usage may also refer to a measuring device.

## Metered-Dose Inhaler

A pressurized inhaler that releases a measured amount of medicine as a spray. Correct timing and breathing are needed for the medicine to reach the lungs; a spacer can make use easier and improve delivery.

## Metestrus

The stage of the reproductive cycle after ovulation in many mammals, when the corpus luteum develops and progesterone rises. The exact timing and biological features differ among species.

## Metformin

A commonly used medicine for type 2 diabetes that lowers the amount of sugar made by the liver and helps the body use insulin better. Nausea or diarrhea are common; serious illness or severe kidney problems increase rare toxicity risk.

## Metformin Hydrochloride

Metformin hydrochloride is the hydrochloride salt of metformin, an oral biguanide used chiefly for

---

type 2 diabetes. It lowers blood glucose without directly stimulating pancreatic insulin release and is available in immediate- and extended-release formulations.

### **Metformin In Prediabetes**

Metformin may reduce progression from prediabetes to type 2 diabetes, especially in people at high risk. Its use depends on age, body size, prior gestational diabetes, kidney function, other conditions, and patient preference; lifestyle intervention remains important.

### **Methacholine Challenge Test**

A breathing test used when asthma is suspected but routine breathing tests are normal. The person inhales increasing amounts of methacholine, which can narrow sensitive airways; breathing is measured after each dose under medical supervision.

### **Methacholine Chloride**

Methacholine chloride is a muscarinic cholinergic agonist used in bronchoprovocation testing. In susceptible people it causes airway narrowing, allowing assessment for bronchial hyperresponsiveness under controlled medical supervision.

### **Methadone**

A long-acting opioid medicine used in selected pain cases and treatment of opioid-use disorder. It reduces withdrawal and cravings but can cause dangerous breathing suppression, sedation, dependence, or heart-rhythm changes.

### **Methadone Hydrochloride**

Methadone hydrochloride is the hydrochloride salt of methadone. It provides long-lasting opioid-receptor activity for selected analgesic and opioid-use-disorder indications, with risks of respiratory depression, dependence, overdose, and cardiac arrhythmia.

### **Methadone Overdose**

Taking too much methadone can cause profound sleepiness, pinpoint pupils, slow or stopped breathing, coma, and death. Effects may be delayed or

prolonged, and abnormal heart rhythms can occur.

### **Methaemoglobinaemia**

A condition in which hemoglobin is changed so it cannot carry or release oxygen normally. It may cause blue or gray skin, headache, breathlessness, confusion, or severe oxygen shortage after exposure to certain medicines or chemicals.

### **Methamidophos**

Methamidophos is a highly toxic organophosphorus insecticide that inhibits acetylcholinesterase. Exposure can cause cholinergic poisoning with sweating, salivation, vomiting, bronchospasm, bradycardia, muscle weakness, and potentially respiratory failure.

### **Methamphetamine**

Methamphetamine is a potent central nervous system stimulant that increases synaptic catecholamines. It has limited medical use in some settings but is commonly misused; toxicity and chronic use can cause cardiovascular complications, psychosis, dependence, and neurocognitive harm.

### **Methamphetamine Overdose**

Taking too much methamphetamine can cause agitation, dangerously high temperature, fast heartbeat, high blood pressure, chest pain, seizures, stroke, abnormal rhythms, or severe confusion. It is a life-threatening poisoning.

### **Methamphetamines**

Powerful stimulant drugs that increase activity in the brain and body. Repeated or high-dose use can cause dependence, severe heart problems, overheating, sleeplessness, paranoia, hallucinations, seizures, stroke, or death.

### **Methamphetamine Toxicity**

Severe stimulant toxicity causing agitation, paranoia or psychosis, mydriasis, diaphoresis, tachycardia, hypertension, hyperthermia, seizures, myocardial ischemia, stroke, or rhabdomyolysis. Benzodiazepines, active cooling, ECG monitoring, and treatment of hyperthermia, seizures, ischemia, and complications

---

are central.

### **Methane**

Methane is a colorless, odorless, highly flammable gas produced by anaerobic decomposition and normal intestinal microbial activity. It can displace oxygen in enclosed spaces and cause asphyxiation; it is not itself a direct systemic poison at ordinary exposure levels.

### **Methane Sulfonate**

Methane sulfonate, or mesylate, is the anion or ester derived from methanesulfonic acid. Mesylate salts are commonly used to improve the formulation of medicines; some methanesulfonate esters are potent alkylating agents and toxic impurities.

### **Methanobrevibacter Smithii**

A methane-producing archaeon commonly found in the human intestine. It consumes hydrogen made by other gut microorganisms and may influence intestinal fermentation and gas production.

### **Methanol**

A poisonous alcohol whose breakdown products can cause severe acid buildup, blindness, brain injury, or death. Serious poisoning can develop even before symptoms appear.

### **Methanol Poisoning**

Poisoning by methanol, a type of alcohol found in some industrial products. Its breakdown can cause severe acid buildup, blindness, brain injury, or death, sometimes before symptoms appear.

### **Methanol Test**

A blood test that measures methanol, a poisonous alcohol. It helps confirm toxic exposure, but treatment may begin before results because methanol can cause severe metabolic acidosis and delayed blindness or brain injury.

### **Methazolamide**

Methazolamide is a carbonic anhydrase inhibitor used mainly to lower intraocular pressure in glau-

coma. It reduces aqueous humor production and can cause metabolic acidosis, electrolyte abnormalities, kidney stones, or sulfonamide-related reactions.

### **Methemoglobin**

Hemoglobin in which iron has been oxidized so it cannot carry oxygen normally. High levels cause methemoglobinemia, with cyanosis, headache, breathlessness, confusion, or severe tissue oxygen deficiency.

### **Methemoglobinemia**

Methemoglobinemia occurs when hemoglobin iron is oxidized and cannot carry oxygen normally, causing cyanosis, headache, breathlessness, or severe tissue hypoxia.

### **Methenamine**

Methenamine is a urinary antiseptic that decomposes in acidic urine to release formaldehyde, which inhibits bacterial growth. It is used for prevention of recurrent urinary tract infection in selected patients, not for treatment of active upper-tract infection.

### **Methicillin-Resistant Staphylococcus Aureus**

A strain of *Staphylococcus aureus* that is resistant to several common antibiotics. It can cause skin, wound, lung, blood, bone, or other infections, and treatment is chosen using the infection site and laboratory testing.

### **Methicillin-Resistant Staphylococcus Aureus Infection**

Alternate name; see *MRSA Infection*.

### **Methidathion**

Methidathion is an organophosphorus insecticide and acaricide that inhibits acetylcholinesterase. Exposure can produce a cholinergic toxidrome with secretions, bronchospasm, bradycardia, gastrointestinal symptoms, weakness, and respiratory failure.

### **Methimazole**

A medicine that reduces production of thyroid hor-

---

mones and is used mainly for hyperthyroidism and Graves disease. It can rarely cause severe low white-cell counts or liver injury; fever or sore throat during treatment needs prompt medical advice.

### **Methiocarb**

Methiocarb is a carbamate pesticide that inhibits acetylcholinesterase reversibly. Poisoning causes excessive muscarinic and nicotinic cholinergic activity and may require urgent supportive care and antidotal treatment.

### **Methionine**

An essential amino acid containing sulfur that the body uses to build proteins and make other substances. People must obtain it from food, and it can be converted into homocysteine and related compounds.

### **Methionine Enkephalin**

Methionine enkephalin is an endogenous opioid pentapeptide that binds opioid receptors and participates in modulation of pain, stress, reward, and neuroendocrine function. It is one of the products of proenkephalin processing.

### **Methocarbamol**

Methocarbamol is a centrally acting skeletal-muscle relaxant used with rest and physical measures for acute painful musculoskeletal conditions. It can cause drowsiness, dizziness, impaired coordination, and additive central nervous system depression.

### **Methohexital**

Methohexital is an ultra-short-acting barbiturate anesthetic used for induction of general anesthesia and, in selected settings, procedural sedation. It depresses central nervous system activity and may cause respiratory or cardiovascular depression.

### **Methomyl**

Methomyl is a carbamate insecticide that reversibly inhibits acetylcholinesterase. Poisoning produces cholinergic symptoms such as salivation, sweating, vomiting, bronchospasm, bradycardia, weakness, and respiratory compromise.

### **Methotrexate**

An antimetabolite that inhibits folate-dependent enzymes and is used in cancer and selected autoimmune or inflammatory disorders.

### **Methotrexate In Dermatology**

An immunomodulatory antimetabolite used in selected inflammatory and autoimmune skin diseases, including psoriasis, eczema, and some connective-tissue disorders. Treatment requires attention to liver disease, blood counts, infection risk, drug interactions, and strict pregnancy precautions.

### **Methotrimeprazine**

Methotrimeprazine, also called levomepromazine, is a phenothiazine with antipsychotic, sedative, antiemetic, and analgesic properties. It is used in some countries for psychiatric illness and palliative-care symptoms and can cause sedation, hypotension, and anticholinergic effects.

### **Methoxamine**

Methoxamine is a predominantly alpha-adrenergic agonist that increases vascular tone and blood pressure. It has been used as a vasopressor for hypotension but is not a routine first-line agent in many current practice settings.

### **Methoxsalen**

Methoxsalen is a psoralen that intercalates into DNA and, after exposure to UVA, forms cross-links that suppress abnormal lymphocyte or skin-cell activity. It is used with UVA phototherapy for selected skin diseases; photosensitivity and skin-cancer risk require precautions.

### **Methoxyflurane**

A vapor medicine formerly used to cause general anesthesia and, in some settings, provide short-term pain relief. High or repeated exposure can damage the kidneys, so use requires careful medical supervision.

### **Methyclothiazide**

Methyclothiazide is a thiazide diuretic that inhibits sodium and chloride reabsorption in the distal con-



---

volute tubule. It is used for hypertension or edema and may cause hypokalemia, hyponatremia, hyperuricemia, and glucose intolerance.

### **Methyl Alcohol**

Another name for methanol, a toxic alcohol that can cause severe acidosis and blindness.

### **Methylation**

Attachment of a small chemical group to DNA, proteins, or other molecules. On DNA, this can change how strongly a gene is used and helps regulate normal development, cell function, and some diseases.

### **Methyl Bromide**

Methyl bromide, or bromomethane, is a volatile fumigant and alkylating chemical. Inhalation can cause severe neurologic, respiratory, renal, and cardiovascular toxicity, including delayed seizures and pulmonary edema.

### **Methylcellulose**

A plant-fiber product used as a bulk-forming laxative and as a thickener in medicines and foods. It absorbs water and adds bulk to stool; it can be unsafe when a bowel blockage is present.

### **Methyl Chloride**

Methyl chloride, or chloromethane, is a volatile industrial gas formerly used in refrigerants and other applications. High-level inhalation exposure can depress the central nervous system and cause headache, ataxia, seizures, or cardiac effects.

### **Methyldopa**

A blood-pressure medicine that acts in the brain to reduce nerve signals that tighten blood vessels. It is used in selected cases, including pregnancy, and may cause sleepiness, dizziness, or liver problems.

### **Methylene Blue**

A medicine used mainly for serious methemoglobinemia, a condition in which blood cannot carry oxygen normally. It may be unsafe in some people with an inherited red-cell enzyme deficiency and can interact

with certain antidepressants.

### **Methylene Blue Test**

A test or procedure using methylene blue dye to trace fluid flow, identify a structure, or detect a leak or abnormal connection. The purpose depends on the body site; allergy and blood-related complications are uncommon but possible.

### **Methylene Chloride**

Methylene chloride, or dichloromethane, is a volatile industrial solvent. Inhalation can cause central nervous system depression and metabolism to carbon monoxide can produce tissue hypoxia; high exposure may be fatal.

### **Methylenedioxymethamphetamine**

3,4-Methylenedioxymethamphetamine, abbreviated MDMA and commonly called ecstasy or molly, is a psychoactive drug with stimulant and hallucinogenic effects that increases serotonergic signaling and can cause serious toxicity.

### **Methylergometrine**

An ergot alkaloid that increases uterine contraction and may be used to control postpartum hemorrhage when appropriate.

### **Methyl Isocyanate**

Methyl isocyanate is a highly volatile and reactive industrial chemical. Inhalation or eye and skin exposure can cause severe irritation, pulmonary edema, and delayed respiratory injury; it was the toxic agent in the Bhopal disaster.

### **Methylmalonic Acid Blood Test**

A blood test measuring methylmalonic acid to help detect vitamin B12 deficiency, especially when the B12 result is borderline. Kidney impairment can raise the result and must be considered during interpretation.

### **Methylmalonic Acidemia**

An inherited metabolic disorder in which the body cannot properly process certain fats and proteins,

---

causing methylmalonic acid to build up. Episodes may cause vomiting, dehydration, acidosis, low blood sugar, seizures, or developmental and kidney problems.

### **Methylmalonic Acid Level**

The amount of methylmalonic acid measured in blood or urine. A high level can support vitamin B12 deficiency, especially when the B12 result is unclear, but kidney disease can also raise it.

### **Methylmalonic Aciduria**

A disorder in which methylmalonic acid builds up in blood and urine, usually because of an inherited enzyme or vitamin B12-processing problem. It may cause vomiting, poor feeding, acidosis, developmental problems, kidney disease, or neurologic injury.

### **Methylmercury Compounds**

Methylmercury compounds are organic mercury compounds that can accumulate in aquatic food chains and cross the blood-brain and placental barriers. Exposure primarily injures the nervous system and can impair fetal neurodevelopment.

### **Methylmercury Poisoning**

Poisoning caused by methylmercury, usually through contaminated food. It primarily injures the nervous system and may cause numbness, poor coordination, vision or hearing changes, and impaired fetal brain development.

### **Methylmethacrylate**

Methyl methacrylate is a volatile monomer used to make acrylic plastics and dental or orthopedic bone cement. Vapors may irritate the eyes and airways, and liquid exposure can cause dermatitis or sensitization.

### **Methylmorphine**

Another name for codeine, an opioid medicine used for selected pain or cough indications. It can cause drowsiness, constipation, dependence, and dangerous breathing suppression, especially with alcohol or other sedatives.

### **Methylnaltrexone**

Methylnaltrexone is a peripherally acting mu-opioid-receptor antagonist used to treat opioid-induced constipation when laxatives are insufficient. It generally preserves central analgesia but can cause abdominal pain, diarrhea, and rarely gastrointestinal perforation.

### **Methylnaltrexone Bromide**

Methylnaltrexone bromide is the bromide salt of methylnaltrexone, a peripherally restricted opioid antagonist. It reverses opioid effects in the gastrointestinal tract and is used for opioid-induced constipation under indication-specific precautions.

### **Methyl Parathion**

Methyl parathion is an organophosphorus insecticide that inhibits acetylcholinesterase. Exposure can produce severe cholinergic poisoning with secretions, bronchospasm, bradycardia, weakness, seizures, and respiratory failure.

### **Methylphenidate**

A stimulant medicine used mainly to treat attention-deficit/hyperactivity disorder and narcolepsy, a condition causing excessive sleepiness. It affects brain chemicals involved in attention and alertness; reduced appetite, sleep problems, increased blood pressure, and misuse are possible risks.

### **Methylprednisolone**

A synthetic glucocorticoid used to suppress inflammation and immune activity.

### **Methyl Salicylate Overdose**

Taking too much methyl salicylate, found in some oils and pain-relief products, can cause salicylate poisoning with vomiting, ringing in the ears, rapid breathing, confusion, seizures, acidosis, or coma. Emergency assessment is needed.

### **Methyltestosterone**

A synthetic androgen related to testosterone. It can affect sexual development and other androgen-sensitive tissues but may cause liver injury, adverse cholesterol changes, infertility, or virilization and is

---

used only in selected medical situations.

### **Metoclopramide**

A medicine that helps the stomach empty and relieves nausea and vomiting. It is used for selected digestive problems and treatment-related nausea; prolonged or high-dose use can cause involuntary movements and other nervous-system effects.

### **Metolazone**

A water-removing medicine used for high blood pressure and swelling. It helps the kidneys remove salt and water and is often combined with another diuretic, but can disturb sodium, potassium, kidney function, or hydration.

### **Metolcarb**

Metolcarb is a carbamate insecticide that can inhibit acetylcholinesterase. Exposure may cause cholinergic poisoning with excessive secretions, gastrointestinal symptoms, bronchospasm, bradycardia, weakness, or respiratory compromise.

### **Metopic Ridge**

A metopic ridge is a palpable or visible line along the forehead from fusion of the metopic suture; it may be normal or associated with trigonocephaly.

### **Metoprolol**

Metoprolol is a relatively beta-1-selective blocker used for hypertension, angina, heart failure, rate control, and some post-infarction settings; bradycardia, hypotension, fatigue, and bronchospasm can occur.

### **Metric System**

The metric system is a decimal system of measurement based on units such as the meter, kilogram, and liter. Modern medicine generally uses the International System of Units, with conventional units retained for some measurements.

### **Metritis**

Inflammation or infection of the uterus, often after childbirth, miscarriage, abortion, or a uterine

procedure. It may cause fever, lower abdominal pain, unusual discharge, bleeding, or tenderness and usually needs treatment.

### **Metronidazole**

An antibiotic and antiparasitic medicine used for certain infections caused by bacteria that grow without oxygen and by some parasites. It can cause nausea or a metallic taste; nerve problems may occur with prolonged use.

### **Metronidazole-Associated Encephalopathy**

A rare complication of prolonged or high-dose metronidazole therapy, causing cerebellar dysfunction (ataxia, dysarthria, nystagmus) and sometimes confusion or seizures. MRI may show T2-hyperintense lesions in the cerebellar dentate nuclei, corpus callosum splenium, and brainstem. Symptoms typically resolve with drug discontinuation.

### **Metronomic Chemotherapy**

Cancer treatment given in frequent, usually smaller doses over a long period instead of occasional high doses. The goal is to slow cancer growth and sometimes its blood supply; use depends on the cancer and treatment plan.

### **Metrorrhagia**

Uterine bleeding that occurs at unexpected times, including between menstrual periods. Causes include pregnancy, hormonal changes, fibroids, polyps, infection, medicines, and cancer. New, heavy, prolonged, or postmenopausal bleeding can have an underlying medical cause.

### **METs**

Metabolic equivalents used to estimate activity intensity or energy use. One MET is approximately the energy used while resting; higher values indicate more demanding activity and help describe exercise capacity.

#### **Alternate Names:**

Metabolic Equivalents

---

## Metirapone

A medicine that blocks one step in the production of cortisol, a hormone made by the adrenal glands. It is used for selected hormone tests or treatment under specialist supervision because levels can change quickly.

## Metzenbaum Scissors

Fine, lightweight surgical scissors designed to separate and cut delicate soft tissue. Their relatively narrow blades are useful for careful dissection, but they are not intended for cutting hard material or heavy sutures.

## Mevalonate Kinase Deficiency

An inherited autoinflammatory disorder caused by changes in the MVK gene. It can cause repeated fever attacks, swollen lymph nodes, abdominal pain, rash, joint pain, or mouth ulcers; the milder form is often called hyperimmunoglobulin D syndrome.

## Mevinphos

Mevinphos is a highly toxic organophosphorus insecticide that inhibits acetylcholinesterase. Exposure can rapidly cause cholinergic crisis, including bronchorrhea, bronchospasm, muscle weakness, seizures, and respiratory failure.

## Mexacarbate

Mexacarbate is a carbamate insecticide that interferes with acetylcholinesterase. Toxic exposure may produce muscarinic and nicotinic cholinergic symptoms and requires prompt decontamination and medical evaluation.

## Mexiletine

Mexiletine is an oral class IB antiarrhythmic and sodium-channel blocker used for selected ventricular arrhythmias and sometimes neuropathic pain. It can cause gastrointestinal and neurologic effects and may worsen or provoke arrhythmias in some patients.

## Meyerozyma Guilliermondii

*Meyerozyma guilliermondii* is a yeast formerly classified as *Candida guilliermondii*. It is usually an

environmental or colonizing organism but can cause opportunistic bloodstream or other infections, particularly in immunocompromised patients.

## MF

Abbreviation; see *Mycosis Fungoides*.

## MFS

Abbreviation; see *Myxofibrosarcoma*.

## MG

Abbreviation; see *Myasthenia Gravis*.

## MGUS

See *Monoclonal Gammopathy of Undetermined Significance*.

## MHC Molecule

A protein on the surface of a cell that displays small pieces of proteins to T cells. This helps the immune system recognize infected or abnormal cells and distinguish the body's own cells from foreign material.

### Alternate Names:

Major Histocompatibility Complex Molecule

## Mianserin

An antidepressant medicine used in some countries to treat depression. It changes signaling by several brain chemicals and can cause sleepiness, increased appetite, weight gain, dizziness, or other medicine-related effects.

## MIBG Scan

A nuclear medicine scan that uses a small amount of radioactive MIBG to find certain tumors of nerve-related tissue. It is used especially for pheochromocytoma, paraganglioma, and neuroblastoma; medicines and thyroid protection may be needed before the scan.

## MIBG Scintiscan

A nuclear medicine scan using radiolabeled metaiodobenzylguanidine to image tissues derived from the sympathetic nervous system, including se-

---

lected pheochromocytomas, paragangliomas, and neuroblastomas.

### **Mica**

A group of sheet silicate minerals; inhalation of mica dust can cause occupational lung disease.

### **Micafungin**

An echinocandin antifungal agent used in the treatment and prophylaxis of invasive candidiasis and esophageal candidiasis. Micafungin inhibits beta-(1,3)-D-glucan synthase, disrupting fungal cell wall synthesis and leading to osmotic instability and cell death.

### **Micelle**

A tiny cluster of molecules that helps carry fats and fat-soluble substances through intestinal fluid for absorption.

### **Michaelis-Menten Kinetics**

A mathematical model describing the rate of enzyme-catalyzed reactions by relating reaction velocity ( $V$ ) to substrate concentration ( $[S]$ ), defined by the maximal velocity ( $V_{max}$ ) and the Michaelis constant ( $K_m$ , the substrate concentration at half-maximal velocity).

### **Miconazole**

An antifungal medicine used in creams, gels, or other preparations for infections of the skin, mouth, or vagina. It stops fungi from making an essential part of their outer covering; irritation and medicine interactions are possible, especially with some oral products.

### **Microabscess**

A very small pocket of pus containing immune cells, damaged tissue, and sometimes germs. It may occur in the skin or an organ because of infection or inflammation and is usually a finding rather than a diagnosis.

### **Microalbuminuria**

A small but abnormal amount of albumin, a blood

protein, in the urine. It can be an early sign of kidney damage, especially in diabetes or high blood pressure, but temporary illness or exercise can also raise it and repeat testing may be needed.

### **Microalbuminuria Test**

A urine test measuring small amounts of albumin, usually by an albumin-to-creatinine ratio, to detect early kidney damage, especially in diabetes or hypertension.

### **Microaneurysm**

A very small balloon-like bulge in a blood vessel wall. In the retina, microaneurysms are an early sign of diabetic eye disease and may leak fluid, causing swelling and blurred vision. Their number and associated retinal changes help show how diabetes is affecting the eyes.

### **Microangiopathic Hemolysis**

Destruction of red blood cells as they pass through damaged or narrowed small blood vessels. Blood tests may show fragmented cells, anemia, and other signs of red-cell breakdown; causes include severe high blood pressure, pregnancy complications, infection, and clotting disorders.

### **Microangiopathic Hemolytic Anemia**

An anemia caused by mechanical fragmentation of red blood cells as they pass through damaged or narrowed small blood vessels, producing schistocytes and often elevated hemolysis markers.

### **Microangiopathy**

Disease or damage affecting the body's smallest blood vessels. Thickened or injured vessel walls can reduce blood flow or allow leakage and may damage organs, especially in diabetes, high blood pressure, or clotting disorders.

### **Microarray**

A laboratory test using a small chip covered with DNA probes. It can measure activity of many genes or find selected missing or extra DNA segments and is used mainly for specialized genetic or research testing.

---

## Microarray Analysis

Microarray analysis measures the abundance or hybridization of many nucleic-acid sequences simultaneously using an array of immobilized probes. It can assess gene expression, copy-number changes, or selected genetic variants.

## Microbacterium

A genus of gram-positive bacteria found in soil and water that can rarely cause bloodstream or other opportunistic infections.

## Microbe

A microorganism, such as a bacterium, fungus, protozoan, or virus.

### Alternate Names:

Microorganism

## Microbial Biofilms

Microbial biofilms are organized communities of microorganisms attached to a surface and enclosed in a self-produced extracellular matrix. Biofilms can form on tissues and medical devices and often show increased tolerance to antimicrobials and host defenses.

## Microbicide

A microbicide is a chemical or biological agent that kills microorganisms or inactivates them. The term may refer to disinfectants, topical agents, or investigational products intended to reduce sexual transmission of infection.

## Microbiology

The study of microorganisms, including bacteria, viruses, fungi, and parasites, and how they interact with people, animals, plants, and the environment. In medicine, it supports the diagnosis, prevention, and treatment of infectious diseases.

## Microbiome

The community of microorganisms and their genetic material living in a body site, especially the intestine. These organisms can influence digestion, immunity, and health, although the effects differ by person and

condition.

## Microcalcification

A very small calcium deposit in tissue. On a mammogram, its shape and arrangement may reflect normal change, injury, infection, or early breast cancer, so suspicious findings may require further imaging or biopsy.

## Microcephaly

Microcephaly is head size substantially smaller than expected for age and sex, reflecting genetic, developmental, infectious, toxic, or acquired brain causes.

## Microcirculation

The movement of blood through the body's smallest blood vessels. In these vessels, oxygen and nutrients pass to tissues and waste products pass into the blood; poor microcirculation can slow healing and damage organs.

## Microcolon

Microcolon is an abnormally small-caliber colon, often identified on contrast imaging. It may result from reduced fetal intestinal use or obstruction and is associated with conditions such as small-left-colon syndrome, meconium ileus, or distal intestinal obstruction.

## Microcornea

Microcornea is an abnormally small cornea, usually defined in adults by a horizontal diameter below about 10 mm. It may be isolated or associated with microphthalmia, glaucoma, cataract, or genetic syndromes.

## Microcurie

A microcurie ( $\mu\text{Ci}$ ) is a unit of radioactivity equal to one-millionth of a curie, or approximately 37 kilobecquerels. It may be used to express the activity of radioactive tracers or sources.

## Microcystic Adnexal Carcinoma

A rare, slow-growing but locally aggressive malignant adnexal tumor, usually on the head or neck,



---

with follicular and ductal differentiation; it may extend deeply and requires complete excision.

### **Microcytic Anemia**

Anemia in which red blood cells are smaller than usual. Common causes include iron deficiency, thalassemia, long-term inflammation, lead exposure, and some bone-marrow disorders. Blood counts and iron tests help find the cause; symptoms may include tiredness, weakness, pale skin, or shortness of breath.

#### **Alternate Names:**

Microcytic Hypochromic Anemia; Small-Cell Anemia; Low-MCV Anemia

### **Microdermabrasion**

A cosmetic treatment that gently removes the outermost layer of skin with fine crystals or a rough-tipped device. It may improve the look of mild uneven texture, dullness, or some marks, but can cause redness, irritation, infection, or temporary darkening.

### **Microdialysis**

A testing method that uses a very small tube to collect chemicals from the fluid around cells. It can measure local levels of medicines or natural substances, including in the brain or beneath the skin.

### **Microdiscectomy**

A spinal operation that removes part of a herniated disc pressing on a nerve, usually in the lower back. It may reduce pain, numbness, or weakness caused by nerve pressure. Possible complications include infection, bleeding, nerve injury, or a return of the disc problem.

#### **Alternate Names:**

Lumbar Microdiscectomy; Microdiskectomy; Minimally Invasive Discectomy

### **Microdissection**

Microdissection is separation or isolation of very small tissue regions or structures under microscopic or highly magnified visualization. It is used in surgery, pathology, and molecular research to obtain

selected cells or tissue.

### **Microdissection Testicular Sperm Extraction**

A surgical procedure that uses a microscope to search the testicle for small areas making sperm. It may help some men with very low or absent sperm production obtain sperm for assisted reproduction. The procedure can cause pain, bleeding, infection, or reduced testicular function, and sperm may not be found.

#### **Alternate Names:**

Micro-TESE; Microsurgical TESE

### **Microdontia**

Abnormally small teeth, which may affect one or several teeth and can be inherited or developmental.

### **Microfibril**

A very small thread-like structure that supports tissues. In connective tissue, microfibrils help organize elastic fibers and provide strength; in plants, they are made mainly of cellulose.

### **Microfilament**

A microfilament is a thin cytoskeletal filament composed mainly of actin, about 7 nm in diameter. Microfilaments support cell shape, motility, contraction, cytokinesis, and intracellular transport.

### **Microfilament Proteins**

Proteins that make, organize, move, or regulate actin microfilaments. They help cells maintain shape, migrate, divide, contract, and transport materials; examples include actin, myosin, and actin-binding proteins.

### **Microfilaria**

The tiny, immature form of a filarial worm that may circulate in blood or live in the skin. Mosquitoes, flies, or other biting insects can pick them up and spread the infection to another person.

### **Microglandular Adenosis**

Microglandular adenosis is an uncommon benign

---

proliferation of small glands in the breast, often associated with hormonal influences such as pregnancy or progestin use. It can mimic invasive carcinoma and is diagnosed histologically.

### **Microglia**

Microglia are resident immune cells of the central nervous system. They survey neural tissue, remove debris, and respond to injury or infection; persistent activation can contribute to neuroinflammation.

### **Microgliosis**

Microgliosis is proliferation or increased activation of microglia in the central nervous system, usually in response to injury, infection, degeneration, or other inflammation. It is a histopathologic finding rather than a specific disease.

### **Micrognathia**

Micrognathia is an unusually small lower jaw that may affect feeding, teeth, facial development, or airway breathing.

### **Micrognathism**

Micrognathism is abnormal smallness of the jaw, usually referring to the mandible. It may be congenital or acquired and can affect feeding, dentition, airway patency, speech, or facial development.

### **Micrograms**

A unit of mass equal to one millionth of a gram, written as mcg. It is commonly used for small medicine doses and nutrient amounts; confusing micrograms with milligrams can cause a serious dosing error.

### **Micrographia**

Abnormally small handwriting, often becoming progressively smaller within a sentence. It commonly occurs in Parkinson disease and other movement disorders because of reduced movement size and control.

### **Microinjection**

Microinjection is delivery of a very small volume of

fluid into a cell, tissue, or other microscopic target using a fine needle or micropipette. It is used in assisted reproduction, experimental biology, and some therapeutic procedures.

### **Micrometastasis**

A small group of cancer cells that has spread from the original tumor but is too small to detect on routine scans or examination. It may be found in tissue samples and can affect staging and treatment decisions.

### **Micrometer**

An instrument or unit used to measure very small distances; one micrometer is one-millionth of a meter.

### **Micromolar**

A concentration unit equal to one millionth of a mole per liter, written as  $\mu\text{mol}/L$ .

### **Micromole**

A micromole ( $\mu\text{mol}$ ) is an amount of substance equal to one-millionth of a mole, or approximately  $6.022 \times 10^{17}$  entities. It is commonly used for small quantities and concentrations of biochemical substances.

### **Micron**

A unit measuring one-millionth of a meter, commonly used to describe the size of cells, bacteria, and particles.

### **Microneme**

A microneme is a small secretory organelle in many apicomplexan parasites. It releases proteins that support host-cell attachment, motility, and invasion during the parasite life cycle.

### **Micronization**

Micronization is a process that reduces particles to micrometer-scale dimensions, often to increase surface area and improve dissolution or absorption of a drug. The resulting particle size and formulation still determine clinical performance.

---

### **Micronized**

Processed into very small particles to improve dispersion or absorption.

### **Micronutrient**

A micronutrient is a vitamin or mineral required in small amounts for normal growth, metabolism, and physiologic function. Deficiency or excess can cause disease, and requirements vary with age, pregnancy, illness, and other factors.

### **Micronutrients**

Vitamins and minerals required in small amounts for enzymatic, hormonal, structural, hematologic, neurologic, and immune functions. They include fat-soluble vitamins (A, D, E, and K), water-soluble vitamins, and essential minerals.

### **Microorchidism**

Abnormally small testes, which may result from impaired development, genetic syndromes, or testicular atrophy after injury or infection.

### **Microorganism**

A microscopic living organism, such as a bacterium, fungus, protozoan, or alga; viruses are acellular infectious agents.

### **Microorganism Genetics**

The study of genetic material, inheritance, variation, and gene activity in microorganisms. It helps explain how microbes grow, cause disease, develop drug resistance, and exchange useful or harmful traits.

### **Micropenis**

An unusually small penis caused by inadequate androgen effect or other developmental conditions, defined clinically by a stretched penile length more than 2.5 standard deviations below the mean for age and population. Evaluation may identify endocrine or genetic causes.

### **Microphallus**

An abnormally small penis, also called micropenis,

usually defined relative to age and developmental standards; it may reflect hormonal or developmental disorders.

### **Microphthalmia**

A birth defect in which one or both eyes are smaller than usual. It may occur with other eye or genetic conditions and can reduce vision. Treatment depends on the cause and may include glasses, treatment for poor vision, or surgery.

### **Microphthalmos**

A birth condition in which one or both eyeballs are unusually small. Vision may be reduced, and other eye or genetic problems may occur, so affected children need an eye examination and ongoing specialist care.

### **Microplastics**

Very small plastic particles, generally less than five millimeters across, found in the environment and sometimes in food, water, and air. Human health effects are still being studied.

### **Micropsia**

A change in vision in which objects appear smaller or farther away than they really are. It can occur with migraine, retinal disease, brain disorders, medicines, or other problems and should be assessed if it is new, persistent, or associated with neurological symptoms.

### **MicroRNA**

A very small, non-coding RNA that helps control which genes are active, mainly by reducing the production of specific proteins. MicroRNAs are being studied as possible disease markers and treatment targets.

### **Microsatellite Instability**

A tumor pattern in which short repeated DNA sequences gain or lose units because the cell's DNA repair system is faulty. A high level can occur in inherited or sporadic cancers and may help predict benefit from some immune treatments.

---

## Microscope

An instrument that magnifies structures too small to see clearly with unaided eyes. Light, electron, and fluorescence microscopes help examine cells, tissues, blood, microbes, and other samples.

## Microscopic

Visible only with a microscope or relating to microscopic examination.

## Microscopic Anatomy

The study of cells, tissues, and fine structural details that require magnification for examination; it includes cytology and histology.

## Microscopic Colitis

Long-lasting inflammation of the colon that usually causes frequent, watery, non-bloody diarrhea while the colon looks normal during routine colonoscopy. Tiny tissue samples are needed for diagnosis; the main forms are collagenous and lymphocytic colitis.

### Alternate Names:

Collagenous Colitis; Lymphocytic Colitis

## Microscopic Examination

Examination of cells or tissue with a microscope to look for normal structure, infection, inflammation, or cancer. The sample may be stained and assessed together with its visible appearance, imaging, and other laboratory results.

## Microscopic Hematuria

Alternate name; see *Hematuria*.

## Microscopic Polyangiitis

An autoimmune disease that inflames and damages small blood vessels. It can affect the kidneys, lungs, skin, nerves, or other organs and may cause blood in the urine, breathing problems, rash, weakness, or bleeding.

## Microscopy

Microscopy is examination of cells, tissues, organisms, or materials with a microscope. Light, fluo-

rescence, electron, and other microscopy methods provide different information about structure and composition.

## Microsecond

A unit of time equal to one-millionth of a second.

## Microspectrophotometry

Microspectrophotometry measures the absorption or emission spectrum of a microscopic region or individual cell. It can quantify pigments, nucleic acids, proteins, or other chromophores in small biological samples.

## Microsphere

A microsphere is a small spherical particle, often made of polymer, glass, or another material, used in drug delivery, imaging, embolization, or laboratory assays. Its size, composition, and surface properties determine its biologic behavior.

## Microspherophakia

Microspherophakia is an abnormally small, spherical crystalline lens. It can occur in inherited disorders such as Weill–Marchesani syndrome and may cause lens displacement, pupillary block, angle-closure glaucoma, or reduced vision.

## Microsporidia

A group of microscopic spore-forming organisms related to fungi. They can cause prolonged diarrhea, eye disease, muscle disease, or widespread infection, especially in people with weakened immune systems.

## Microsporidiosis

An infection caused by microsporidia, microscopic organisms related to fungi that produce resistant spores. It most often affects the intestine and may cause long-lasting diarrhea and weight loss. It can also affect the eyes, muscles, kidneys, brain, or several organs, especially in people with weakened immune systems.

## Microsporum

Microsporum is a genus of dermatophyte fungi that

---

can cause ringworm of the skin, scalp, or hair.

### **Microsporium Audouinii**

*Microsporium audouinii* is a dermatophyte that mainly infects the scalp and can cause tinea capitis, especially in children.

### **Microsporium Distortum**

*Microsporium distortum* is a rare dermatophyte related to *M. canis* that can cause skin or scalp infection.

### **Microsporium Ferrugineum**

A fungus that can cause ringworm of the scalp or skin, particularly in children. Laboratory examination can help confirm the infection and distinguish it from other causes of rash or hair loss.

### **Microsporium Gypseum**

*Microsporium gypseum* is a soil-associated dermatophyte that can cause ringworm of the skin or scalp.

### **Microstomia**

Microstomia is abnormal narrowing or smallness of the mouth opening. It may be congenital or result from scarring, burns, surgery, trauma, or connective-tissue disease and can impair eating, speech, oral hygiene, and dental treatment.

### **Micro-Surgery**

Surgery performed with magnification and fine instruments to repair very small structures such as nerves and vessels.

### **Microtechnology**

Microtechnology is the design and manufacture of devices or structures with micrometer-scale features. Medical applications include microfluidic diagnostics, sensors, drug-delivery systems, and minimally invasive instruments.

### **Microtia**

A birth defect in which the outer ear is smaller than usual or does not form fully. The ear canal may also be narrow or absent, which can reduce hearing.

### **Microtome**

An instrument that cuts extremely thin slices of tissue for examination under a microscope. The slices are mounted on slides, stained, and studied to identify normal structures or disease.

### **Microtubule**

A tiny hollow protein tube inside a cell. Microtubules help maintain cell shape, move materials within the cell, support cell division, and form part of cilia and other moving structures.

### **Microtubule Bundle**

A microtubule bundle is an organized group of parallel or otherwise coordinated microtubules within a cell. Bundles provide structural support and guide transport or movement, including in axons, cilia, and dividing cells.

### **Microvascular**

Relating to the body's smallest blood vessels, including tiny arteries, capillaries, and veins. These vessels deliver oxygen and nutrients, remove waste, and control blood flow through tissues.

### **Microvascular Disease**

Disease affecting the body's smallest blood vessels, which can reduce blood flow and oxygen delivery to organs or tissues. Diabetes, high blood pressure, smoking, inflammation, and aging can damage these vessels.

### **Microvasculature**

The network of the body's smallest blood vessels, including tiny arteries, capillaries, and veins. It delivers oxygen and nutrients, removes waste, controls tissue blood flow, and helps regulate fluid movement between blood and tissues.

### **Microvilli**

Tiny finger-like projections on the surface of cells. In the small intestine, they increase the area available for absorbing nutrients.

---

### Microvillus

A microvillus is a microscopic, actin-supported projection of a cell membrane that increases surface area. Microvilli are abundant on intestinal epithelial cells, where they support nutrient absorption, and on renal tubular cells.

### Microvolt

A microvolt ( $\mu\text{V}$ ) is one-millionth of a volt. It is used to express small bioelectric signals, such as those recorded by electroencephalography, electromyography, electrocardiography, or evoked-potential testing.

### Microwave Ablation

A procedure that uses microwave energy to heat and destroy selected tissue, usually under imaging guidance.

### Microwave Therapy

Treatment that uses microwave energy to heat or destroy selected tissue. It has specialized uses, such as treating some tumors or controlling bleeding, and the risks depend on the body area and device used.

### Micturition

The process of passing urine. The bladder squeezes while the muscles at its outlet relax, under coordinated control from the brain and nerves.

### Midaxillary Line

An imaginary vertical line on the side of the body passing through the apex of the axilla (armpit).

### Midazolam

Midazolam is a short-acting benzodiazepine used for procedural sedation, anesthesia, and treatment of acute seizures in appropriate settings. It enhances GABA-A receptor activity and can cause profound sedation and respiratory depression, especially with opioids or other depressants.

### Midazolam Hydrochloride

Midazolam hydrochloride is the hydrochloride salt of the short-acting benzodiazepine midazolam. It

is used for procedural sedation, anesthesia, and selected seizure emergencies and can cause dose-related sedation, hypotension, and respiratory depression.

### Midbrain

The upper part of the brainstem, between the lower brainstem and the main brain. It helps control eye movements, movement, alertness, and automatic responses to sights and sounds.

### Midclavicular Line

An imaginary vertical line on the front of the body passing through the midpoint of the clavicle (collarbone).

### Middle Ear

An air-filled space between the eardrum and inner ear that contains the malleus, incus, and stapes. These ossicles transmit sound vibrations to the inner ear, and the auditory tube helps equalize middle-ear pressure.

### Middle Ear Implant

An implanted hearing device that electrically or mechanically stimulates middle-ear structures to improve sound transmission in selected hearing loss. It differs from a cochlear implant, which directly stimulates the inner-ear auditory nerve pathway.

### Middle Ear Infection

Infection or inflammation in the space behind the eardrum. It often follows a cold and may cause ear pain, fever, temporary hearing loss, fluid behind the eardrum, or irritability in children.

### Middle East Respiratory Syndrome

A serious respiratory infection caused by a coronavirus, usually linked to close contact with infected camels or people. It may cause fever, cough, pneumonia, breathing failure, or kidney and other organ problems.

#### Alternate Names:

MERS; Camel Flu



---

### **Middle Insomnia**

Middle insomnia is repeated or prolonged awakening during the middle portion of the sleep period, often with difficulty returning to sleep. It may occur with stress, mood disorders, sleep apnea, pain, substances, medications, or other sleep disorders.

### **Middle Lobe Syndrome**

Chronic or recurrent atelectasis and infection of the right middle lobe due to bronchial obstruction. The long, narrow right middle lobe bronchus is susceptible to compression by lymph nodes, tumors, or mucous plugging. It may be caused by bronchiectasis, malignancy, or granulomatous disease. Treatment depends on the underlying cause.

### **Middle Mediastinum**

The central part of the lower mediastinum that contains the heart, pericardium, roots of the great vessels, and phrenic nerves.

### **Mid-Dorsal**

Situated in the middle region of the back or on the dorsal side of a structure.

### **Mid-Forearm**

The middle part of the forearm, between the elbow and wrist. Measurements or injections described as mid-forearm use this region as a reference point.

### **Mid-Gut**

The embryonic middle portion of the primitive gut, which develops into much of the small intestine and proximal colon.

### **Midgut Volvulus**

A life-threatening twisting of the intestine that can cut off its blood supply, usually because the intestine formed abnormally before birth. Babies may suddenly develop green vomiting, a swollen or painful abdomen, blood in the stool, or extreme illness. It is an emergency requiring immediate imaging and surgery to untwist and protect the bowel.

#### **Alternate Names:**

Volvulus Neonatorum; Midgut Torsion

### **Mid-Leg**

The central part of the lower limb between the knee and ankle. It contains the shin bones, muscles, nerves, blood vessels, and soft tissues that support walking and ankle movement.

### **Mid-Line**

The imaginary or anatomical line dividing a structure or body into right and left portions.

### **Midomafetamine**

Midomafetamine is another name for 3,4-methylenedioxymethamphetamine, a psychoactive amphetamine derivative. It is being studied in controlled psychotherapy-assisted research but can cause hyperthermia, hyponatremia, cardiovascular toxicity, serotonin toxicity, and misuse.

### **Midostaurin**

Midostaurin is a multikinase inhibitor used with chemotherapy for newly diagnosed acute myeloid leukemia with a susceptible FLT3 mutation and for advanced systemic mastocytosis. It can cause nausea, cytopenias, infection, and other treatment-related toxicities.

### **Midriff**

Alternate informal term; see *Abdomen*.

### **Mid-Stream Urine Specimen**

A urine sample collected after initial urine is voided, reducing contamination from the urethral opening and used commonly for culture.

### **Mid-Systolic Click**

A sharp heart sound heard during the middle or later part of the heartbeat, often associated with mitral-valve prolapse. It may be followed by a murmur if the valve allows blood to leak backward.

### **Midtrimester**

The middle part of pregnancy, usually covering about weeks 14 through 27. During this period the fetus grows rapidly, routine anatomy screening is often performed, and pregnancy symptoms and

---

risks vary from person to person.

### Mid-Upper-Arm Circumference

The circumference of the upper arm measured at the midpoint between the acromion and olecranon processes. It is a practical anthropometric measure of muscle and fat stores, particularly useful when body weight or height cannot be measured reliably.

### Midwife

A trained and regulated professional who provides care during pregnancy, labor, birth, and the postnatal period.

### Midwifery

The professional practice of supporting and managing normal pregnancy, childbirth, postpartum recovery, and newborn care.

### Mifepristone

Mifepristone is a progesterone-receptor antagonist and, at higher doses, a glucocorticoid-receptor antagonist. It is used with misoprostol for medication abortion under applicable protocols and for selected cases of hyperglycemia associated with Cushing syndrome.

### Migraine

A recurring neurological disorder that causes attacks of moderate to severe, often throbbing or pulsing head pain. The pain may occur on one or both sides of the head and may be accompanied by nausea, vomiting, or increased sensitivity to light, sound, or smells. Some people have an aura, such as flashing lights, blind spots, tingling, numbness, or temporary speech problems, which usually develops gradually and resolves completely. Migraine attacks may last from several hours to a few days, and some occur with little or no headache.

#### Alternate Names:

Migraine Headache; Hemicrania

### Migraine And Stroke Symptoms

Migraine aura usually develops gradually and resolves, whereas stroke symptoms often begin suddenly and cause persistent focal deficits such as

weakness, speech trouble, facial droop, or vision loss. New or atypical neurologic symptoms require emergency stroke assessment.

### Migraine With Aura

A primary headache disorder characterized by recurrent attacks of fully reversible visual, sensory, or speech symptoms that develop gradually over at least five minutes and last no more than sixty minutes, followed by headache with the characteristics of migraine.

### Migraine Without Aura

A primary headache disorder characterized by recurrent attacks of moderate to severe, pulsating, unilateral headache lasting four to seventy-two hours, associated with nausea, vomiting, photophobia, and phonophobia.

### Migrainous Neuralgia

Older term; see *Cluster Headache*.

### Migrating Motor Complex

Repeating waves of muscle activity that move through the stomach and small intestine between meals. They help clear leftover food, secretions, and some bacteria before the next meal.

### Migration

Movement of cells, organisms, substances, or a body structure from one place to another. Normal examples include immune-cell movement and fetal development; abnormal migration can contribute to infection, cancer spread, or blood clots.

### Migratory Thrombophlebitis

Recurrent or shifting venous thrombosis associated with an underlying malignancy; an alternate term for the thrombotic manifestation of Trousseau syndrome.

### Milan System for Salivary Cytopathology

A six-level system for reporting results from a needle sample of a salivary-gland lump. Results range from an insufficient sample or noncancerous finding to

---

a finding suspicious for or showing cancer. The system helps estimate cancer risk and guide further care.

### **Mild**

Mild describes relatively low intensity, severity, or extent of a symptom, sign, disease, or abnormality. The meaning depends on the condition and the clinical scale used and does not by itself indicate that no treatment or evaluation is needed.

### **Mild Cognitive Impairment**

A noticeable decline in memory or another thinking skill that is greater than expected for age but does not substantially prevent everyday activities. It may remain stable, improve, or progress to dementia depending on the cause.

### **Mildew Remover Poisoning**

Poisoning from a mildew-removal product, often containing bleach, acids, or other corrosive chemicals. It can burn the skin, eyes, mouth, and digestive tract or release dangerous fumes; leave the area and seek urgent advice.

### **Mile**

A mile is a unit of distance equal to 1,609.344 meters. It may be used in descriptions of walking or running distance, although scientific and clinical measurements generally use SI units.

### **Milia**

Milia are tiny, firm, white or yellow bumps caused by small cysts containing keratin, a protein in the skin. They are common in newborns and may appear on the face in adults; they are harmless and usually go away on their own.

### **Miliaria**

A sweat-retention disorder caused by blockage of eccrine ducts, producing small itchy or prickly papules.

### **Miliaria Crystallina**

A harmless heat rash made of tiny clear blisters when sweat becomes trapped just beneath the skin's

surface. It can occur in hot, humid conditions or during fever and usually settles when the skin is cooled and kept dry.

### **Miliaria Profunda**

Alternate name; see *Heat Rash*.

### **Miliaria Rubra**

An itchy prickly heat rash caused by sweat-duct blockage in the epidermis, producing small red papules or vesicles.

### **Miliary**

Describing many tiny spots or lesions that resemble millet seeds. The term is often used for tuberculosis or another disease spread widely through an organ or the bloodstream.

### **Miliary TB**

Tuberculosis that has spread through the bloodstream, producing many tiny areas of infection in the lungs and sometimes the liver, brain, bone marrow, or other organs. It can cause fever, weight loss, night sweats, cough, or organ-specific symptoms.

### **Miliary Tuberculosis**

Tuberculosis that spreads through the bloodstream and forms many tiny areas of infection. It may affect the lungs, liver, brain, bone marrow, or other organs and can cause prolonged fever, weight loss, night sweats, cough, or organ-specific symptoms.

### **Milieu Therapy**

Milieu therapy uses the structured social environment of a treatment setting as part of care, especially in some psychiatric and rehabilitation programs.

### **Milium**

A small, firm, white or yellow bump filled with keratin, usually appearing on the face. It is harmless, does not spread by infection, and may disappear on its own or be removed safely by a clinician.

### **Milk-Alkali Syndrome**

A condition caused by taking too much calcium

---

and absorbable alkali, leading to high blood calcium, excessive blood alkalinity, and kidney problems. Severe cases can cause confusion, vomiting, dehydration, or abnormal heart rhythms.

### Milker's Node

A localized nodular skin infection caused by parapoxvirus acquired from infected cattle, also called orf-like milker's nodule.

### Milkman Syndrome

Multiple small weak areas or incomplete breaks in bones, also called Looser zones. They can occur in severe osteomalacia or rickets and may cause bone pain, weakness, or fractures.

#### Alternate Names:

Looser-Milkman Syndrome; Pseudofracture Syndrome

### Milk Of Magnesia

A liquid suspension of magnesium hydroxide used to neutralize stomach acid and relieve constipation. It draws water into the bowel and can cause diarrhea, especially when taken too often or in kidney disease.

### Milky Urine

Urine with a cloudy or white appearance caused by chyle, pus, crystals, fat, or other substances; evaluation identifies the cause.

### Millard-Gubler Syndrome

A ventral pontine stroke syndrome caused by an ischemic or hemorrhagic lesion in the lower pontine tegmentum, characterized by ipsilateral abducens nerve (CN VI) palsy, ipsilateral peripheral facial nerve (CN VII) palsy, and contralateral hemiparesis of the upper and lower extremities.

#### Alternate Names:

Ventral Pontine Syndrome; Facial Abducens Hemiplegia Syndrome

### Miller Fisher Syndrome

Alternate name; see *Fisher Syndrome*.

### Miller's Lung

Hypersensitivity pneumonitis caused by inhaling material associated with the granary weevil (wheat weevil, *Sitophilus granarius*), especially in people working with contaminated grain or flour.

### Millicurie

A millicurie (mCi) is one-thousandth of a curie, equal to approximately 37 megabecquerels. It is used to express the activity of radioactive sources and some radiopharmaceutical doses.

### Milliequivalent

A unit used to measure the amount and electrical combining power of charged substances, such as sodium, potassium, calcium, or chloride, in medicines and laboratory results.

### Milligram

A metric unit of mass equal to one-thousandth of a gram. Its abbreviation is mg, and it is commonly used for medicine doses and small quantities of substances.

### Milligram-Percent

An older concentration unit meaning milligrams of a substance in 100 milliliters of fluid. Its number is the same as milligrams per deciliter, written mg/dL, which is preferred in current medical reporting.

### Milligray

A unit of absorbed ionizing-radiation dose equal to one-thousandth of a gray, or 0.001 joule per kilogram; its abbreviation is mGy.

### Milliliter

A unit of volume equal to one-thousandth of a liter and one cubic centimeter. It is commonly used to measure liquid medicines, blood, urine, laboratory samples, and other small amounts of fluid.

### Millimeter

A unit of length equal to one-thousandth of a meter.

---

### Millimolar

A concentration unit equal to one thousandth of a mole per liter, written as mmol/L.

### Millimole

A unit measuring the amount of a chemical substance equal to one-thousandth of a mole. Laboratory tests use millimoles to report substances such as glucose, sodium, potassium, and other electrolytes.

### Million

A number equal to 1,000,000; laboratory counts may use it as a numerical multiplier.

### Milliosmole

A milliosmole (mOsm) is one-thousandth of an osmole, a unit reflecting the number of osmotically active particles in a solution. Osmolality and osmolarity are commonly reported in mOsm/kg or mOsm/L.

### Millipede Toxin

Irritating defensive chemicals released by some millipedes. Contact can cause burning, redness, blistering, or dark staining of skin and severe eye irritation; rinse exposed areas with water and seek care for eye symptoms.

### Millisecond

A millisecond (ms) is one-thousandth of a second. It is used to measure brief physiologic events, reaction times, nerve conduction intervals, and timing of electrical or imaging signals.

### Milliunit

A milliunit (mU) is one-thousandth of a unit of biologic or enzymatic activity. Because a unit is defined according to the substance or assay, the clinical meaning of a milliunit depends on the specific test or medication.

### Millivolt

A unit of electrical voltage equal to one-thousandth of a volt. Small voltages are measured in millivolts in tests of the heart, brain, nerves, muscles, and

other electrical activity in the body.

### Milrinone

A medicine given by vein for short-term support in severe heart failure or after heart surgery. It helps the heart squeeze more strongly and widens blood vessels, but can lower blood pressure or trigger abnormal heart rhythms.

### Milroy Disease

An inherited primary lymphedema caused most often by pathogenic variants affecting VEGFR3 signaling. It typically presents with chronic swelling of the lower limbs from birth or early infancy and may be associated with papillomatosis, hydrocele, or recurrent skin infection.

### Milroy's Disease

An inherited form of primary lymphedema, usually present from birth or early life and caused commonly by lymphatic-development abnormalities.

### Mimesis

Imitation or mimicry; in medicine, the term can refer to simulation of an organic disease by psychological symptoms or to one organic disease resembling another.

### Mimicry

Resemblance of one organism, structure, or clinical pattern to another; in medicine, it may describe a disorder that imitates another disease.

### Minamata Disease

A nervous-system disorder caused by poisoning with methylmercury, often after eating contaminated seafood. It can cause numbness, poor coordination, vision or hearing problems, muscle weakness, speech difficulty, and developmental harm in children.

### Mind

The collection of mental processes and capacities involved in consciousness, thought, perception, memory, emotion, judgment, and behavior.

---

## Mindful Eating

Paying attention to hunger and fullness, eating slowly, and noticing food's taste and texture. It may improve awareness of eating habits and help some people reduce distracted eating or overeating.

## Mineral

An inorganic element or compound required for body structure, fluid balance, enzyme activity, or other physiological functions.

## Mineralocorticoid

A steroid hormone, mainly aldosterone, that helps the kidneys retain sodium and water while removing potassium. Too much can raise blood pressure and lower potassium; too little can cause dehydration and weakness.

## Mineralocorticoid Receptor

A protein inside cells that responds mainly to aldosterone, a hormone made by the adrenal glands. It helps the kidneys retain salt and water and remove potassium, thereby affecting blood pressure and body-fluid balance.

## Mineralocorticoids

Steroid hormones, chiefly aldosterone, that promote sodium retention and potassium and hydrogen secretion in the kidney.

## Mineral Oil

Mineral oil is a purified mixture of liquid hydrocarbons derived from petroleum. It is used in some topical products, cosmetics, and laxatives; aspiration of liquid mineral oil can cause lipoid pneumonia.

## Mineral Oil Overdose

Swallowing a large amount of mineral oil is dangerous mainly because it can enter the lungs during vomiting or reflux and cause lipoid pneumonia.

## Minerals

Naturally occurring elements needed by the body for bones, nerves, muscles, blood formation, fluid balance, and other functions. They include calcium,

iron, magnesium, potassium, sodium, and zinc; both deficiency and excess can be harmful.

## Mineral Spirits Poisoning

Poisoning from mineral spirits, petroleum-based solvents used in paints and cleaning products. Swallowing or inhalation may cause dizziness, drowsiness, chemical pneumonitis, or breathing failure, especially after liquid enters the lungs.

## Miner's Asthma

Asthma caused or worsened by repeated inhalation of dust, fumes, or other substances during mining work. It can cause wheezing, cough, chest tightness, and breathlessness that improve after leaving the exposure.

## Miner's Nystagmus

An older term for visual disturbance and nystagmus reported in underground miners, associated with poor lighting and occupational conditions.

## Minimal Change Disease

A kidney disorder, especially common in children, in which the kidney filters look nearly normal on routine examination but leak large amounts of protein. It can cause swelling around the eyes, legs, or abdomen and is often treatable.

## Minimal-Change Disease

A glomerular disorder in which the kidney looks nearly normal on routine microscopy but podocyte changes cause heavy protein loss in the urine. It is a common cause of nephrotic syndrome in children and often responds to corticosteroids.

## Minimally Invasive Aortic Valve Surgery

Aortic valve surgery performed through a smaller incision than a full median sternotomy, such as a mini-thoracotomy or partial sternotomy. Transcatheter aortic valve replacement (TAVR) is a separate procedure.

### Alternate Names:

Minimally Invasive Aortic Valve Replacement; MI-AVR



---

### **Minimally Invasive Esophagectomy**

Surgical resection of part or all of the esophagus performed using thoracoscopy and laparoscopy or robotic assistance through small keyhole incisions, reducing postoperative recovery time.

#### **Alternate Names:**

MIE; Thoracoscopic Esophagectomy; Robotic Esophagectomy

### **Minimally Invasive Gastrointestinal Surgery**

Gastrointestinal operations performed through small incisions or natural orifices using laparoscopic, robotic, or endoscopic techniques, often reducing recovery time while retaining procedure-specific risks.

### **Minimally Invasive Heart Bypass Surgery**

Coronary artery bypass graft (CABG) surgery performed through a small thoracotomy incision between the ribs, frequently on a beating heart without cardiopulmonary bypass, avoiding a traditional median sternotomy.

#### **Alternate Names:**

MIDCAB; Minimally Invasive Direct Coronary Artery Bypass; Less Invasive CABG

### **Minimally Invasive Hip Replacement**

Hip-replacement surgery performed through a smaller or muscle-sparing approach to reduce tissue disruption. It still replaces damaged joint surfaces and has risks such as infection, blood clots, dislocation, fracture, nerve injury, or implant loosening.

### **Minimally Invasive Mitral Valve Surgery**

Surgical mitral valve repair or replacement performed through a right mini-thoracotomy or video-assisted thoracoscopic approach, avoiding a traditional median sternotomy. Results in reduced surgical trauma, less postoperative pain, and faster recovery.

#### **Alternate Names:**

MIMVS; Mini-Thoracotomy Mitral Repair; Port-Access Mitral Surgery

### **Minimally Invasive Prostate Resection**

Endoscopic or robotic transurethral procedures (such as holmium laser enucleation of the prostate [HoLEP], transurethral resection of the prostate [TURP], or robotic simple prostatectomy) performed to remove obstructive adenomatous tissue in benign prostatic hyperplasia.

#### **Alternate Names:**

HoLEP; Endoscopic Prostatectomy; Minimally Invasive BPH Resection

### **Minimally Invasive Spine Surgery**

Spine surgery performed through small incisions using specialized instruments, often reducing muscle disruption and recovery time. It still carries risks such as nerve injury, infection, bleeding, and persistent symptoms.

### **Minimally Invasive Surgery**

Surgery performed through small incisions or natural openings using specialized instruments, usually reducing tissue disruption and recovery time.

### **Minimal Residual Disease**

A small number of cancer cells that remain after treatment and cannot be seen on routine examination or ordinary scans. Highly sensitive blood, bone-marrow, tissue, or molecular tests may detect them and estimate relapse risk.

### **Mini-Mental State Examination**

A brief questionnaire that checks orientation, attention, memory, language, and simple thinking skills. The score can be affected by education, language, hearing, vision, culture, or delirium and cannot diagnose dementia by itself.

### **Minimization**

A psychological or communication process of downplaying the importance or severity of symptoms, events, or problems.

### **Minimum Alveolar Concentration**

A standard measure of how much inhaled anesthetic is needed to prevent movement in half of people

---

during a specified surgical stimulus. It helps compare anesthetic strength and guide dosing during anesthesia.

### **Minimum Bactericidal Concentration**

The lowest concentration of an antimicrobial that kills a defined proportion of the original bacterial inoculum, commonly demonstrated by subculture without growth. MBC is distinct from MIC and is not routinely required for every infection.

### **Minimum Effect Concentration**

The lowest concentration of a drug in the blood or at its site of action that produces a defined measurable effect.

### **Minimum Inhibitory Concentration**

The smallest amount of an antibiotic or other antimicrobial that stops a particular germ from visibly growing in a laboratory test. It helps guide treatment but must be interpreted with the infection and the patient's condition.

### **Minimum Lethal Concentration**

The lowest concentration of a substance associated with death in a defined test population or experimental setting.

### **Minimum Lethal Dose**

The smallest dose of a substance reported to cause death under specified conditions; it varies by species, route, and individual.

### **Minisatellite Repeat**

A minisatellite repeat is a tandemly repeated DNA sequence with repeat units commonly ranging from about 10 to 100 base pairs. Variation in repeat number can serve as a genetic marker and, in some loci, influence gene regulation or disease risk.

### **Mini-Stroke**

A lay term usually referring to a transient ischemic attack, in which focal neurologic symptoms resolve without persistent infarction.

### **MINOCA**

A heart attack occurring without a major blockage in the coronary arteries. It is an initial diagnosis; further tests are needed to find causes such as a small clot, artery spasm, artery-wall tear, or inflammation.

### **Minocycline**

An antibiotic used for some bacterial infections and for inflammatory acne. It can cause nausea, dizziness, skin or tooth discoloration, sun sensitivity, and rarely liver or immune problems; the course should be taken as prescribed.

### **Minor**

Smaller, lesser, or less significant than a related structure or condition; for example, the teres minor muscle is smaller than the teres major muscle.

### **Minority Health**

The health status and healthcare experiences of populations affected by social, economic, cultural, or structural disadvantage. Differences in exposure, income, environment, discrimination, and access to care can contribute to health disparities.

### **Minor Response**

A minor response is a limited improvement after treatment that does not meet the predefined criteria for a partial or complete response. The exact threshold depends on the disease, study protocol, and outcome being measured.

### **Minoxidil**

A vasodilator used topically for pattern hair loss and orally in selected cases of resistant hypertension.

### **Minute Ventilation**

The total amount of air breathed in one minute, calculated from breath size multiplied by breathing rate. It may rise with exercise or illness and fall with sedation, muscle weakness, airway disease, or poor breathing effort.

---

## Miosis

Unusually small pupils or normal pupil narrowing in response to light. It may be caused by medicines or toxins, eye disease, or problems affecting nerves in the brain or eye.

## Miosis Disorder

Abnormally small pupils or an abnormal pupillary constriction response, caused by medications, neurologic disease, ocular disease, or exposure to certain toxins.

## Miotic

A medicine or substance that makes the pupil smaller. Miotics may be used in glaucoma or during some eye procedures and can affect vision in dim light.

### Alternate Names:

Miotic Agent; Pupillary Constrictor

## Miracidium

The ciliated larval stage of a trematode that hatches from an egg and infects a suitable snail host.

## Mirage Syndrome

A rare genetic disorder, usually related to SAMD9 variants, that can cause growth restriction, adrenal insufficiency, recurrent infections, and distinctive developmental findings.

## MIRD Schema

A method used to estimate how much radiation reaches organs and tissues after a radioactive medicine is given. It helps specialists plan treatment and assess safety for radiopharmaceutical procedures.

## Mirizzi Syndrome

A gallstone stuck near the gallbladder outlet presses on a nearby bile duct and blocks the flow of bile. It can cause jaundice, abdominal pain, fever, or infection and may require an endoscopic or surgical procedure.

## Mirror Therapy

A rehabilitation exercise in which a mirror makes movement of the stronger limb appear to come from the affected limb. It may help some people with phantom-limb pain or movement recovery after stroke.

## Mirror-Writing

Writing in which letters or words are reversed as if seen in a mirror; it may occur transiently in childhood or with neurologic disease.

## Mirtazapine

Mirtazapine is a noradrenergic and specific serotonergic antidepressant used for major depressive disorder. It commonly causes sedation and increased appetite or weight and can rarely cause neutropenia or other serious adverse effects.

## Misalignment

Misalignment is abnormal positioning or orientation of two or more structures relative to their intended relationship. It may affect bones, joints, teeth, eyes, implants, or medical devices and can impair function or healing.

## Miscarriage

Spontaneous loss of a pregnancy before fetal viability, commonly caused by chromosomal abnormalities but requiring clinical assessment.

### Alternate Names:

Spontaneous Abortion; Early Pregnancy Loss

## Misdiagnosis

Misdiagnosis is assignment of an incorrect diagnosis or failure to recognize the correct condition. It can lead to inappropriate treatment, delayed care, avoidable harm, or unnecessary testing and may result from atypical presentation, incomplete information, or system factors.

## Mismatch Repair

A system that checks newly copied DNA and corrects small errors. When it fails, genetic changes can accumulate and increase the risk of cancers such

---

as colorectal and womb cancer.

### Mismatch Repair Deficiency

Loss of DNA mismatch-repair function that produces microsatellite instability and a high tumor mutational burden. Seen in Lynch syndrome and some sporadic colorectal, endometrial, and other cancers, and associated with responsiveness to immune checkpoint inhibitors.

### MISME

Abbreviation; see *Neurofibromatosis Type 2*.

### Misophonia

Misophonia is an intense emotional or physiologic reaction to specific repetitive sounds, such as chewing or tapping. It is not simple annoyance; behavioral strategies, sound management, and therapy can reduce impairment, while assessment considers anxiety, obsessive-compulsive symptoms, and sensory sensitivity.

### Misoprostol

A synthetic prostaglandin analogue used for selected gynecologic and obstetric indications, including cervical ripening, labor induction, and medical management of some pregnancies. It can cause uterine contractions, bleeding, gastrointestinal effects, or uterine tachysystole; use depends on the clinical setting and gestational age.

#### Alternate Names:

Prostaglandin E1 Analogue

### Missed Dose

A missed dose is a scheduled dose of a medicine that was not taken or administered. The appropriate response depends on the drug and timing; doubling a dose can be dangerous, so the product instructions or a clinician or pharmacist should guide the next dose.

### Missed Miscarriage

A nonviable intrauterine pregnancy that has not been expelled and may produce few or no symptoms of miscarriage. Diagnosis requires appropriate ultrasound or other clinical evaluation before man-

agement is selected.

#### Alternate Names:

Missed Abortion; Silent Miscarriage; Early Pregnancy Failure

### Missense Mutation

A missense mutation is a DNA sequence change that substitutes one amino acid for another in a protein. Its effect may be harmful, beneficial, or neutral depending on the affected residue, protein function, and genetic context.

### Missense Variant

A single-nucleotide substitution that changes one amino acid in a protein. Its clinical effect ranges from benign to pathogenic and must be assessed using molecular, functional, population, and clinical evidence.

### Mistletoe Poisoning

Poisoning after eating mistletoe leaves or berries. It may cause nausea, vomiting, diarrhea, abdominal pain, dizziness, low blood pressure, slow or irregular heartbeat, seizures, or confusion; children and symptomatic people need urgent assessment.

### Mite

A tiny arachnid; some mites cause dermatitis or transmit disease, and house-dust mites can trigger allergy.

### Mites

Tiny arthropods related to ticks and spiders. Some live harmlessly on skin, while others cause allergy, scabies, or irritation; treatment depends on the species and condition.

### Mithramycin

An older antineoplastic antibiotic, also called plinamycin, whose use is limited by substantial toxicity.

### Mitobronitol

Mitobronitol is an alkylating antineoplastic compound formerly investigated or used in selected hematologic malignancies. Its cytotoxic effects can

---

damage DNA and cause myelosuppression and other treatment-related toxicities.

### **Mitochondria**

Small structures inside cells that make much of the energy cells need. They also help control calcium, cell survival, and several metabolic processes; mitochondrial disorders can affect organs that need high energy, such as the brain, muscles, heart, and eyes.

### **Mitochondrial Cristae**

Folds of the inner membrane of mitochondria, the energy-producing parts of cells. The folds provide space for the machinery that turns nutrients into usable cellular energy.

### **Mitochondrial Diabetes**

Diabetes caused by a change in mitochondrial DNA, which is usually passed through the mother. It may occur with hearing loss, muscle problems, or other mitochondrial symptoms; insulin production is often reduced and genetic testing may confirm the cause.

### **Mitochondrial Disease**

A group of genetic disorders caused by impaired mitochondrial energy production. Symptoms vary and may affect the brain, muscles, nerves, heart, eyes, or multiple organs; onset and severity can differ widely even within one family.

### **Mitochondrial DNA**

Small circular DNA molecules located primarily in mitochondria that encode some components of oxidative phosphorylation and are inherited predominantly through the maternal line.

### **Mitochondrial DNA Depletion Syndrome MDS MDDS**

A group of inherited disorders in which some tissues contain too little mitochondrial DNA to make energy normally. Depending on the form, the brain, muscles, liver, or several organs may be affected, often beginning in infancy or childhood.

### **Mitochondrial Fission**

Mitochondrial fission is division of one mitochondrion into two or more mitochondria. It helps distribute mitochondria, remove damaged portions through mitophagy, and respond to cell-cycle or metabolic signals; excessive fission may contribute to disease.

### **Mitochondrial Fusion**

Mitochondrial fusion is joining of mitochondria into a connected network. It permits exchange of mitochondrial contents and helps maintain bioenergetic function and genome integrity; fusion and fission are normally balanced dynamically.

### **Mitochondrial Inheritance**

Inheritance of mitochondrial DNA, which is transmitted predominantly through the egg and therefore usually from the mother. Examples include Leber hereditary optic neuropathy and MELAS. Heteroplasmy means that mutant and normal mitochondrial DNA coexist in a cell.

### **Mitochondrial Matrix**

The mitochondrial matrix is the protein-rich compartment enclosed by the inner mitochondrial membrane. It contains mitochondrial DNA, ribosomes, and enzymes for the citric-acid cycle, fatty-acid oxidation, and other metabolic processes.

### **Mitochondrial Membrane**

The mitochondrial membrane system consists of an outer membrane and a highly folded inner membrane separated by an intermembrane space. It controls transport and houses the respiratory chain and ATP-producing machinery.

### **Mitochondrial Myopathy**

A muscle disorder caused by faulty mitochondria, the cell structures that make energy. It may cause weakness, exercise intolerance, drooping eyelids, hearing loss, seizures, or problems in other organs.

### **Mitochondrial Ribosome**

A mitochondrial ribosome is a ribosome within mitochondria that translates proteins encoded by mito-

---

chondrial DNA. Its structure differs from cytosolic ribosomes and defects in mitochondrial translation can cause energy-production disorders.

### **Mitochondrial Swelling**

Enlargement of mitochondria caused by disruption of ion and water balance, often indicating cellular injury, impaired oxidative phosphorylation, or permeability-transition changes.

### **Mitochondrion**

A mitochondrion is a membrane-bound organelle that generates much of a cell's ATP through oxidative phosphorylation. It also participates in apoptosis, calcium handling, metabolism, and production or regulation of reactive oxygen species.

### **Mitogen**

A mitogen is a substance that stimulates a cell, especially a lymphocyte, to enter the cell cycle and proliferate. Mitogens include growth factors, lectins, and other signals that activate proliferative pathways.

### **Mitolactol**

Mitolactol is an alkylating antineoplastic compound formerly used or investigated in cancer treatment. It damages DNA and can cause cytotoxic effects such as myelosuppression and gastrointestinal toxicity.

### **Mitomycin**

Mitomycin is a chemotherapy medicine that damages DNA and is used for selected cancers; it can suppress bone marrow and injure the kidneys or bladder.

### **Mitomycins**

Mitomycins are antitumor antibiotics that become activated in cells and cross-link DNA, inhibiting replication and transcription. Mitomycin C is the best-known member and is used in selected cancers and ophthalmic or surgical applications.

### **Mitophagy**

The process by which a cell identifies and breaks

down damaged or unnecessary mitochondria, the cell's energy-producing structures. This helps maintain healthy cells; abnormal mitophagy may contribute to some diseases.

### **Mitosis**

Cell division that produces two genetically similar daughter cells with the same chromosome number as the parent cell.

### **Mitosporic Fungi**

Fungi that reproduce by asexual spores; the term is mainly used for fungi whose sexual stage is unknown or not observed.

### **Mitotane**

Mitotane is an adrenolytic drug used for selected adrenocortical carcinomas and to suppress excess adrenal steroid production. It can cause gastrointestinal and neurologic effects, adrenal insufficiency, and numerous drug interactions.

### **Mitotic Anaphase**

Mitotic anaphase is the stage of mitosis in which sister chromatids separate and move toward opposite poles of the cell. Accurate spindle attachment and chromosome segregation are required for formation of genetically similar daughter cells.

### **Mitotic Figure**

A cell seen under a microscope while it is dividing. The number of dividing cells in a tissue sample can help estimate how quickly a tumor or other tissue is growing.

### **Mitotic Metaphase**

Mitotic metaphase is the stage of mitosis in which duplicated chromosomes align at the cell equator with sister kinetochores attached to opposite spindle poles. A checkpoint helps prevent premature chromosome separation.

### **Mitotic Prophase**

Mitotic prophase is the early stage of mitosis in which chromatin condenses into visible chro-



---

mosomes, the nucleolus disappears, and the mitotic spindle begins to form. In late prophase or prometaphase, the nuclear envelope breaks down.

### **Mitotic Recombination**

Mitotic recombination is exchange or rearrangement of DNA between homologous chromosomes during or around mitotic cell division. It can repair DNA or generate loss of heterozygosity and is relevant to tumor evolution and genetic disease.

### **Mitotic Telophase**

Mitotic telophase is the stage after chromosome separation when chromosomes reach opposite poles, decondense, and become enclosed in newly forming nuclear envelopes. Cytokinesis usually follows or overlaps with telophase.

### **Mitoxantrone**

A chemotherapy and immunosuppressive medicine that interferes with DNA and is used for selected cancers and some forms of multiple sclerosis.

### **MitraClip**

A transcatheter device that creates a double orifice in the mitral valve by clipping the anterior and posterior leaflets together, reducing mitral regurgitation. It is indicated for patients with significant (3+ or 4+) mitral regurgitation who are at high risk for surgical mitral valve repair.

### **Mitral**

Relating to the heart's mitral valve, which lies between the left upper and lower chambers. The valve opens for blood flow forward and closes to prevent backward leakage during each heartbeat.

### **Mitral Annular Calcification**

Chronic calcium and fibrous deposition in the ring supporting the mitral valve. It is more common with older age and chronic kidney disease and can restrict leaflet motion, causing or worsening mitral regurgitation or degenerative mitral stenosis; echocardiography and sometimes CT assess its extent and consequences.

### **Mitral Atresia**

A serious heart defect present from birth in which the opening of the mitral valve is absent or completely closed. Blood cannot flow normally into the heart's main pumping chamber, so newborns need urgent specialist care.

### **Mitral Facies**

A distinctive physical appearance characterized by a purplish or violaceous malar flush and patchy cyanosis across the cheeks, bridge of the nose, and lips. It is caused by chronic pulmonary hypertension, systemic vasoconstriction, and reduced cardiac output in severe mitral stenosis.

### **Mitral Insufficiency**

Failure of the mitral valve to close completely, allowing blood to leak back into the left atrium during systole.

### **Mitral Regurgitation**

Leakage through the mitral valve during heart contraction (systole), allowing blood to flow backward from the left ventricle into the left atrium. It may result from abnormal valve leaflets or supporting structures, infection, or heart damage, and can cause breathlessness, fatigue, palpitations, or heart failure.

### **Mitral Stenosis**

Narrowing of the mitral valve opening, which restricts blood flow from the left atrium to the left ventricle. Causes include rheumatic valve disease and age-related calcium buildup around the valve. Increased pressure in the left atrium may lead to breathlessness, atrial fibrillation, pulmonary hypertension, or right-sided heart failure.

### **Mitral Stenosis Murmur**

A low rumbling heart sound heard when blood passes through a narrowed mitral valve. It may be accompanied by a sharp opening sound and is best assessed with a stethoscope and heart imaging.

### **Mitral Valve**

The two-flap valve between the heart's left upper and lower chambers. It opens to let blood pass

---

forward and closes to prevent backward flow; it may become narrowed, leaky, infected, or weakened.

### **Mitral Valve Insufficiency**

Leakage of the mitral valve that allows blood to flow backward into the left atrium when the heart contracts. It may cause breathlessness, fatigue, palpitations, swelling, or heart failure.

### **Mitral Valve Prolapse**

Displacement of one or both mitral-valve leaflets backward into the left atrium during ventricular systole, often because of myxomatous or other degenerative valve changes. It may be harmless, but can cause mitral regurgitation, palpitations, arrhythmias, or rarely infective endocarditis or sudden cardiac death.

#### **Alternate Names:**

MVP; Barlow Syndrome; Click-Murmur Syndrome; Floppy Mitral Valve

### **Mitral Valve Regurgitation**

Backward leakage of blood through the mitral valve when the left ventricle contracts, which can cause breathlessness, fatigue, atrial enlargement, or heart failure.

#### **Alternate Names:**

MR; Mitral Incompetence; Mitral Insufficiency

### **Mitral-Valve Repair**

A surgical or transcatheter procedure to restore mitral valve competence without replacement, typically for regurgitation from prolapse, chordal rupture, or ischemic dysfunction. Repair preserves native valve tissue, maintains left ventricular function, and avoids lifelong anticoagulation required by mechanical prostheses.

### **Mitral Valve Stenosis**

Narrowing of the mitral valve that restricts blood flow from the left atrium to the left ventricle.

#### **Alternate Names:**

MS; Mitral Stenosis

### **Mitral Valvotomy**

A procedure that opens a narrowed mitral valve so blood can move more easily from the upper to the lower left heart chamber. It may use a balloon catheter or surgery and is suitable only for selected valve anatomy.

### **Mitral Valvuloplasty**

Mitral valvuloplasty is a procedure that widens a narrowed mitral valve, often with a balloon catheter or surgical repair.

### **Mitsuokella**

Mitsuokella is a genus of anaerobic bacteria found mainly in the mouth and digestive tract; human infection is uncommon.

### **Mittelschmerz**

Recurrent one-sided lower-abdominal or pelvic pain occurring around ovulation. It is usually brief or lasts up to a day or two, but severe, persistent, or atypical pain requires assessment for other causes.

### **Mixed Aortic Valve Disease**

Narrowing of the aortic valve (aortic stenosis) and backward leakage through it (aortic regurgitation) occurring together. The two problems affect blood flow and the heart at the same time, so their combined severity is assessed rather than judging each in isolation.

### **Mixed Connective Tissue Disease**

An autoimmune illness with features of lupus, scleroderma, and inflammatory muscle disease. It may cause color changes in fingers with cold, swollen hands, joint pain, muscle weakness, swallowing problems, or inflammation around organs.

#### **Alternate Names:**

MCTD; Sharp Syndrome

### **Mixed Connective Tissue Disease-Associated Arthritis**

Joint pain or inflammation occurring with mixed connective tissue disease, an autoimmune condition with features of lupus, systemic sclerosis, and in-

---

inflammatory muscle disease.

### **Mixed Hearing Loss**

Hearing loss caused by both a problem carrying sound through the outer or middle ear and a problem in the inner ear or hearing nerve. The best treatment depends on the causes found.

### **Mixed-Lineage Leukemia**

An acute leukemia in which the malignant blasts show defining features of more than one hematopoietic lineage, commonly both myeloid and lymphoid; it is also called mixed-phenotype acute leukemia.

### **Mixed Mitral Valve Disease**

Narrowing of the mitral valve (mitral stenosis) and backward leakage through it (mitral regurgitation) occurring together. The two problems affect blood flow and pressure in the heart at the same time, so their combined severity is assessed rather than judging each in isolation.

### **Mixed Oedema**

Edema caused by more than one mechanism or affecting more than one tissue compartment; interpretation depends on clinical context.

### **Mixed Urinary Incontinence**

Urine leakage caused by both a sudden strong urge to urinate and pressure from coughing, sneezing, laughing, or exercise. Treatment may include bladder training, pelvic-floor exercises, medicines, or procedures.

### **Miyoshi Myopathy**

A rare inherited muscle disease that usually begins in adolescence or early adulthood with progressive weakness of the calf muscles. People may have difficulty standing on tiptoe or climbing stairs; weakness later spreads to the thighs and hips. Blood tests often show high muscle-enzyme levels. Physical therapy helps maintain movement and function.

#### **Alternate Names:**

Miyoshi Distal Myopathy; Distal Dysferlinopathy

### **MME**

Membrane metalloendopeptidase, also called neprilysin, CD10, neutral endopeptidase, or CALLA; it is a cell-surface enzyme and marker expressed in several tissues and in some acute lymphoblastic leukemias.

### **MéNièRe'S Disease**

An inner-ear disorder causing repeated attacks of spinning dizziness, changing hearing loss, ringing, and pressure or fullness in one ear. Symptoms may last minutes to hours and can affect balance and daily life.

#### **Alternate Names:**

Meniere Disease; Endolymphatic Hydrops

### **MéNéTrier'S Disease**

A rare stomach disorder in which the stomach lining becomes unusually thick and loses protein into the digestive tract. It may cause upper-abdominal pain, nausea, vomiting, weight loss, swelling, or low blood protein.

### **Mobile**

Capable of moving or being moved; a mass or structure described as mobile is not fixed to adjacent tissue.

### **Mobile Genetic Elements**

Genetic elements that can move within and between genomes. These include plasmids, transposons, and integrons. Mobile genetic elements can carry antibiotic resistance genes and virulence factors. They facilitate horizontal gene transfer and contribute to the rapid evolution of bacteria.

### **Mobile Health**

The use of phones, tablets, wearable devices, or similar portable technology to support health care. Uses include reminders, remote monitoring, education, communication, symptom tracking, and some diagnostic tools; privacy, accuracy, accessibility, and clinical oversight remain important.

#### **Alternate Names:**

mHealth; Digital Health

---

## Mobile Health Unit

A mobile health unit is a vehicle or transportable facility equipped to provide health services at changing locations. It may deliver screening, vaccination, primary care, diagnostic testing, emergency support, or outreach to underserved communities.

## Mobility

The ability to move or be moved, including functional movement of a person or range of motion of a joint.

## Mobilization

The process of restoring or encouraging movement, such as early patient activity, joint movement, or surgical release of tissue.

## Mobiluncus

A genus of curved anaerobic bacteria associated with bacterial vaginosis and occasionally isolated from other clinical specimens.

## Mobiluncus Curtisi

A curved bacterium that grows without oxygen and may be present with bacterial vaginosis. Its detection can support assessment, but symptoms, examination, and other test results are needed to interpret vaginal findings.

## Mobiluncus Mulieris

A curved bacterium that grows without oxygen and may be associated with bacterial vaginosis. Finding it can support a diagnosis, but symptoms, examination, and other laboratory results are also important.

## Mobitz Type I

A form of second-degree atrioventricular (AV) block characterized on electrocardiography by progressive PR interval prolongation in successive beats until a single P wave fails to conduct to the ventricles (dropped QRS), after which the cycle repeats (Wenckebach periodicity), typically localized within the AV node.

**Alternate Names:**

Wenckebach Block; Second-Degree AV Block Mobitz I

## Mobitz Type I Block

A form of second-degree atrioventricular block characterized electrocardiographically by progressive lengthening of the PR interval in consecutive cardiac cycles until an atrial P wave fails completely to conduct to the ventricles (dropped QRS complex). The cycle then resets with a shorter PR interval. Typically localized to physiological or pathological fatigue within the atrioventricular node, often benign and reversible.

**Alternate Names:**

Wenckebach Block; Mobitz I AV Block

## Mobitz Type II

A high-grade form of second-degree atrioventricular (AV) block characterized on electrocardiography by intermittent, non-conducted P waves (dropped QRS complexes) occurring without prior PR interval prolongation, typically localized in the infranodal His-Purkinje system and carrying a high risk of progression to complete heart block.

**Alternate Names:**

Second-Degree AV Block Mobitz II; Infranodal AV Block

## Mobitz Type II Block

A heart rhythm problem in which some signals from the upper chambers fail to reach the lower chambers without the warning lengthening of the PR interval. It can suddenly worsen into complete heart block and usually needs urgent specialist assessment; a pacemaker is often recommended when the cause is not reversible.

**Alternate Names:**

Mobitz II AV Block; Infranodal Second-Degree AV Block

## Mobius Syndrome

A rare condition present from birth that affects certain facial and eye-moving nerves. It can cause facial weakness or paralysis, trouble closing the eyes, limited outward eye movement, and feeding or speech difficulties.

---

## Modafinil

A prescription medicine that promotes wakefulness in narcolepsy, shift-work sleep disorder, and persistent sleepiness from obstructive sleep apnea when airway treatment is adequate. It can cause headache, anxiety, insomnia, serious rash, or dependence and is not a substitute for sleep or airway treatment.

## Modality

A particular method used for treatment, diagnosis, imaging, or sensory stimulation. Examples include surgery, medicines, ultrasound, MRI, physiotherapy, and forms of counseling; the best modality depends on the health problem and the person's needs.

## Mode

The most frequent value in a dataset or a particular manner or method of operation.

## Model

A simplified representation used to describe, explain, predict, or study a biological or clinical process.

## Moderate To Severe Disabilities

Moderate to severe disability describes substantial limitations in physical, cognitive, sensory, communication, or adaptive functioning that require ongoing support. Needs vary widely and may include medical care, rehabilitation, assistive technology, education, and assistance with daily activities.

## Moderation

Moderation is reduction or control of the intensity, amount, or severity of a process, symptom, behavior, or treatment effect. In clinical research, a moderator is a factor that changes the strength or direction of an association.

## Moderator

A factor that changes the strength or direction of a relationship or treatment effect; it may also mean a person who guides a discussion.

## Modes Of Inheritance

Patterns by which genetic traits pass through fami-

lies, including autosomal dominant, autosomal recessive, X-linked, mitochondrial, and other patterns.

## Modified Biophysical Profile

A pregnancy assessment that combines a fetal heart-rate test with measurement of the amniotic-fluid volume. A nonreactive heart-rate result or low fluid level may suggest that further assessment or delivery is needed, depending on gestational age and the clinical situation.

### Alternate Names:

Modified BPP; NST with Amniotic Fluid Index

## Modulator

A substance or factor that alters the activity of a biological pathway, receptor, gene, or physiological process.

## Modus Operandi

The characteristic method by which a person performs an action; it is chiefly a forensic or legal term.

## Modus Vivendi

A practical arrangement or way of living that allows people or groups with differences to coexist.

## Moebius Syndrome

A congenital disorder characterized mainly by facial paralysis and inability to abduct the eyes, caused by developmental cranial-nerve abnormalities.

## Moellerella

A rare group of bacteria that can occasionally cause urinary, bloodstream, or other infection, mainly in people who are seriously ill or have weakened defenses. Laboratory identification is needed to determine whether a specimen represents true infection.

## Moellerella Wisconsensis

A rare type of bacterium that can occasionally cause urinary, bloodstream, wound, or other infections, especially in people with serious illness or weakened immunity. Laboratory testing identifies it and guides antibiotic choice.

---

### Moeller's Glossitis

An older term for a smooth, painful, inflamed tongue associated especially with nutritional deficiency, including iron, folate, or vitamin B12 deficiency.

### MOGAD

Abbreviation; see *MOG Antibody Disease*.

### MOG Antibody-Associated Disorder

Alternate name; see *MOG Antibody Disease*.

### Mogibacterium

A genus of anaerobic bacteria found in the mouth and intestine and occasionally associated with dental or other infection.

### Mohs Micrographic Surgery

A skin-cancer operation that removes the visible tumor a thin layer at a time and examines each layer under a microscope until no cancer remains. It preserves as much healthy tissue as possible; bleeding, infection, scarring, or recurrence can occur.

### Mohs Surgery

A micrographically controlled technique in which skin cancer is excised in thin layers and examined margin-by-margin, achieving high cure rates for basal cell and squamous cell carcinoma while sparing normal tissue.

### Moisture

Water or other liquid present on a surface or in tissue; excessive moisture can cause maceration and skin breakdown.

### Molar

Relating to a back tooth used for chewing, or to a chemical concentration measured in moles per liter.

### Molar Pregnancy

An abnormal pregnancy in which placental tissue grows excessively instead of developing normally. In a complete mole no embryo forms; in a partial mole abnormal fetal tissue may be present. It can cause bleeding, severe nausea, or unusually high

pregnancy-hormone levels and requires specialist follow-up.

#### Alternate Names:

Hydatid Mole; Gestational Trophoblastic Disease

### Molasses

A viscous syrup produced during sugar refining; it is rich in sugar and may contain variable amounts of minerals.

### Mold

A multicellular fungus that grows as branching filaments; molds can cause allergy, food spoilage, or opportunistic infection.

### Mold Allergy

An allergic response to mold spores that may cause nasal, eye, skin, or lower-respiratory symptoms.

#### Alternate Names:

Fungal Hypersensitivity; Mold-Induced Rhinitis

### Mold Exposure

Mold exposure can irritate the eyes, skin, and airways and may worsen asthma or allergic rhinitis; invasive infection is uncommon and mainly affects severely immunocompromised people. Reducing moisture and remediating contaminated areas are more useful than unproven detoxification.

### Mole

A common usually harmless skin growth made of pigment-producing cells. It may be flat or raised and brown, black, pink, or skin-colored. A new or changing mole, or one unlike the others, should be checked for skin cancer.

### Molecular

Relating to molecules or processes occurring at the molecular level.

### Molecular Chaperone

A molecular chaperone is a protein that assists other proteins with folding, assembly, transport, or refolding without becoming part of the final structure. Chaperones can also help prevent or resolve protein



---

aggregation.

### **Molecular Diagnosis**

Molecular diagnosis uses analysis of DNA, RNA, proteins, or other molecular markers to identify a disease, pathogen, inherited variant, tumor subtype, or treatment-relevant characteristic. Results require appropriate assay validation and clinical interpretation.

### **Molecular Dynamics**

A computer method that predicts how atoms and molecules move and interact over time. Researchers use it to study protein shape, drug binding, chemical reactions, and other biological processes; its results depend on the model and assumptions used.

### **Molecular Imaging**

Imaging that shows biological molecules, cell activity, or biochemical processes in the body, often using a targeted radioactive or fluorescent tracer. It can help characterize tumors, inflammation, infection, or organ function.

### **Molecular Pathology**

The study of disease-related changes in genes, gene activity, or proteins within cells and tissues. It can help identify a disease, classify a tumor, estimate likely outcome, detect inherited risk, or select a treatment.

### **Molecular Pathology Of Colorectal Cancer**

The assessment of molecular alterations that classify colorectal tumors and guide prognosis or therapy, including mismatch-repair deficiency or microsatellite instability, RAS mutations, BRAF mutation, HER2 amplification, and selected gene fusions.

### **Molecular Probe**

A molecular probe is a labeled or otherwise detectable molecule designed to bind a particular nucleic-acid sequence, protein, metabolite, or other target. Probes are used in imaging, assays, microscopy, and molecular diagnostics.

### **Molecular Response**

A molecular response is a measurable change in a molecular marker, such as a pathogen load, oncogenic transcript, mutation burden, or biomarker, after disease progression or treatment. Its definition and clinical significance depend on the disease and assay.

### **Molecular Testing**

Analysis of DNA, RNA, or other molecular features to identify variants, fusions, expression patterns, or infectious agents for diagnosis, prognosis, or treatment selection.

### **Molecular Weight**

The mass of one molecule, usually expressed in atomic mass units; for a substance used in chemistry, its numerical value corresponds to molar mass in grams per mole. It helps calculate concentrations and doses.

### **Molecule**

A group of chemically bonded atoms forming the smallest unit of a substance that retains its chemical properties.

### **Moles**

Moles are common collections of pigment-producing melanocytes and are usually benign. A lesion that changes in size, shape, color, symmetry, or symptoms, or differs markedly from others, should be examined for melanoma.

### **Molimina**

The symptoms and physical changes that occur before menstruation, including breast tenderness, bloating, or mood changes.

### **Mollusc**

A soft-bodied invertebrate; the term is also used in names of mollusc-borne or mollusc-associated parasitic diseases.

### **Mollusca**

The animal group containing snails, slugs, clams,

---

and related species. Some mollusks can cause illness through venom, contaminated food, or parasites they carry; medical risk depends on the species and exposure.

### **Molluscipoxvirus**

A group of poxviruses that includes the virus causing molluscum contagiosum. Infection produces small, firm, usually painless skin bumps with a central dimple and spreads through close skin contact or shared items.

### **Molluscum Contagiosum**

A benign superficial skin infection caused by a molluscum poxvirus. It produces small, pearly, firm papules with a central depression, spreads through close skin contact, sexual contact, or shared objects, and usually resolves spontaneously; lesions may be numerous, large, or persistent in people with weakened immunity.

### **Molten Metal Fumes**

Fumes produced during welding, smelting, or casting that can cause metal-fume fever, airway irritation, or metal toxicity.

### **Molteno Implant**

A non-valved glaucoma drainage device (aqueous shunt) surgically implanted to reduce elevated intraocular pressure (IOP) in refractory and complicated glaucomas. It consists of a flexible silicone tube inserted into the anterior chamber (or pars plana) that shunts aqueous humor to one or two thin episcleral polypropylene plates sutured to the sclera beneath Tenon's capsule in the equatorial region, where fluid passively diffuses through a fibrous capsule into the orbital vascular and lymphatic circulation. Because the device is non-valved and offers no intrinsic resistance to flow, the tube is temporarily occluded at surgery (using an absorbable ligature or intraluminal ripcord suture) during the first 4–6 weeks until fibrous bleb encapsulation develops, thereby preventing early postoperative hypotony, anterior chamber collapse, and choroidal detachment. It is indicated when conventional filtration surgery (trabeculectomy) has failed or has

a high probability of failure, including neovascular glaucoma, uveitic glaucoma, and post-penetrating keratoplasty glaucoma.

#### **Alternate Names:**

Molteno Glaucoma Drainage Device; Molteno Aqueous Shunt; Molteno Seton

### **Molybdenum**

An essential trace element required in small amounts for enzymes; excessive exposure can affect copper balance.

### **Molybdenum-99**

A radioactive form of molybdenum that changes into technetium-99m. It is used in generator systems to produce technetium-99m, a radioactive tracer used for many medical scans.

### **Molybdenum Cofactor Deficiency**

An inherited disorder in which cells cannot make or use the molybdenum cofactor required by enzymes such as sulfite oxidase and xanthine dehydrogenase. Severe neonatal disease may cause feeding difficulty, seizures, abnormal muscle tone, encephalopathy, and progressive brain injury.

### **Molybdosis**

Illness caused by excessive molybdenum exposure, which may disturb copper metabolism and cause systemic effects.

### **Mometasone**

Mometasone is a potent synthetic glucocorticoid used topically or by inhalation for inflammatory skin or airway disease. It reduces local inflammation but may cause skin atrophy, oral candidiasis, adrenal effects, or other corticosteroid adverse effects.

### **Mometasone Furoate**

Mometasone furoate is the furoate ester of the corticosteroid mometasone. Depending on formulation, it is used for inflammatory skin disease, allergic rhinitis, or asthma; local and systemic corticosteroid precautions apply.

---

## Momordica

A genus of plants including bitter melon; preparations may affect blood glucose and can interact with medicines.

## Monascus

Monascus is a genus of fungi used in some fermented foods; some species produce pigments, while contamination can produce harmful mycotoxins.

## Monckeberg Medial Calcific Sclerosis

A benign, non-inflammatory form of arteriosclerosis characterized by dystrophic ring-like calcification localized within the tunica media of medium and small muscular arteries without luminal obstruction or intimal ulceration, producing palpable 'lead-pipe' rigidity of peripheral vessels.

### Alternate Names:

Mönckeberg Arteriosclerosis; Medial Calcific Sclerosis

## Mondor Disease

A benign, self-limited superficial thrombophlebitis affecting the subcutaneous veins of the anterior chest wall and breast (lateral thoracic, thoracoepigastric, or superior epigastric veins), presenting as a palpable, painful, cord-like indurated subcutaneous band under tension.

## Mondor'S Disease

Superficial thrombophlebitis of the anterior chest wall veins, typically the lateral thoracic, thoracoepigastric, or superior epigastric veins. Presents as a tender, cord-like subcutaneous structure with skin retraction. Usually self-limited; associated with breast surgery, trauma, or hypercoagulable states.

## Mongolian Spot

A Mongolian spot, now preferably called congenital dermal melanocytosis, is a flat blue-gray patch caused by melanocytes in the dermis, usually present at birth. It is benign and often fades during childhood but should be documented to distinguish it from bruising.

## Monilethrix

Monilethrix is an inherited hair-shaft disorder in which hair has regularly spaced narrow segments and breaks easily, producing a beaded appearance. It commonly causes short, brittle scalp hair and may involve follicles elsewhere.

## Monitored Anesthesia Care

Anesthesia care in which a trained professional gives medicines for comfort or pain relief and continuously watches breathing, heart rate, blood pressure, and oxygen. The level of sedation can be adjusted and emergency support is available.

## Monitoring Device

A monitoring device measures and displays or records a physiologic variable, such as heart rhythm, oxygen saturation, blood pressure, glucose, temperature, or respiratory rate. Accuracy, calibration, and clinical context determine how results should be interpreted.

## Monitoring Test

A test repeated over time to follow a disease, treatment response, medicine safety, or a possible complication. The result is interpreted with symptoms and earlier measurements.

## Monkeypox

Former name; see *Mpox*.

## Monkeypox (Mpox)

Alternate name; see *Mpox*.

## Mono

Alternate name; see *Infectious Mononucleosis*.

## Monoamine Oxidase Inhibitor (MAOI) Toxicity

Poisoning or interaction involving an MAOI that may cause severe hypertension, tachycardia, agitation, hyperthermia, seizures, or serotonin syndrome with clonus and hyperreflexia. Stop serotonergic or sympathomimetic drugs, provide benzodiazepines and supportive care, and manage rapidly chang-

---

ing blood pressure and temperature in an intensive setting.

### **Monoamine Oxidase Inhibitors**

Alternate form; see *Monoamine Oxidase Inhibitor*.

### **Monoarticular Joint Pain**

Pain affecting one joint, caused by trauma, infection, crystal arthritis, osteoarthritis, or inflammatory arthritis. Acute pain with swelling, warmth, fever, or inability to bear weight can occur with septic arthritis.

### **Monobactams**

A class of antibiotics used mainly for certain infections caused by oxygen-using Gram-negative bacteria. Aztreonam is the main example; laboratory testing and allergy history help guide its use.

### **Monoblast**

An early bone marrow cell that develops into a monocyte, a type of infection-fighting white blood cell.

### **Monoblast Count**

A monoblast count is the number or proportion of monoblasts, immature precursors of monocytes, in a blood or bone-marrow sample. Increased blasts may support evaluation for acute myeloid leukemia or another marrow disorder and require expert morphologic or immunophenotypic interpretation.

### **Monochorionic Twin Pregnancy**

A twin pregnancy in which identical twins share a single placenta. Because the babies share placental blood vessels, the pregnancy carries unique risks such as twin-to-twin transfusion syndrome and unequal growth, requiring frequent ultrasound monitoring throughout pregnancy. Twins may have separate inner fluid sacs (diamniotic) or rarely share a single sac (monoamniotic).

#### **Alternate Names:**

Monochorionic twins; MCDA twins; MCMA twins.

### **Monoclonal**

Made by, or arising from, one original cell and therefore very similar to that cell. A monoclonal antibody is designed to recognize one specific target in the body or in a laboratory test.

### **Monoclonal Antibodies**

Antibodies produced from a single B-cell clone and directed against one antigenic target; they are used in diagnosis and treatment.

### **Monoclonal Antibody**

A laboratory-made immune protein designed to recognize one specific target. Monoclonal antibodies are used in tests and treatments for cancer, inflammatory disease, infection, and other conditions.

### **Monoclonal Gammopathy Of Undetermined Significance**

A condition in which abnormal plasma cells make a small amount of one unusual antibody. It usually causes no symptoms or organ damage but can rarely progress to multiple myeloma or another blood cancer, so monitoring is recommended.

### **Monoclonal Gammopathy of Undetermined Significance (MGUS)**

An asymptomatic clonal plasma-cell or B-cell disorder with a small monoclonal immunoglobulin, limited marrow involvement, and no myeloma-defining organ damage. It can progress to a related malignancy and requires monitoring.

### **Monocrotaphos**

Monocrotaphos is a highly toxic organophosphorus insecticide that inhibits acetylcholinesterase. Exposure can cause acute cholinergic poisoning with secretions, bronchospasm, vomiting, bradycardia, weakness, seizures, and respiratory failure.

### **Monocular**

Involving or affecting one eye, or occurring with vision from one eye. Monocular vision can reduce depth perception and may result from eye disease, injury, or loss of vision in the other eye.

---

## Monocyte

A type of white blood cell that travels in the blood and can enter tissues, where it becomes a macrophage or another immune cell. It helps remove germs and damaged material and coordinates inflammation and repair.

## Monocytes

Large white blood cells that circulate in the blood and can enter tissues, where they become macrophages or some dendritic cells. They help remove debris and coordinate inflammation and immune defense.

## Monocytopenia

An abnormally low number of monocytes, a type of white blood cell, in the blood. It may occur with bone-marrow disorders, medicines, infections, or severe illness and is interpreted with the full blood count.

## Monocytosis

Monocytosis is an increased absolute number of circulating monocytes. It may be reactive to infection or inflammation or associated with a hematologic neoplasm; persistent or marked monocytosis requires appropriate evaluation.

## Monogenic

Describing a condition or trait caused mainly by a change in one gene. The change may be inherited from a parent or arise for the first time in the affected person.

## Monogenic Obesity

Severe or early-onset obesity caused mainly by a disease-causing change in one gene. The affected gene may alter appetite or the body's control of energy use.

## Monograph

A monograph is a detailed, authoritative document devoted to one subject. In health care it may describe a drug's identity, quality, pharmacology, indications, dosing, safety, or regulatory standards.

## Monoiodotyrosine

Monoiodotyrosine is an iodinated tyrosine residue formed during thyroid-hormone synthesis. Coupling of iodinated tyrosine residues within thyroglobulin contributes to production of thyroxine and triiodothyronine.

## Monokines

Monokines are cytokines produced chiefly by monocytes and macrophages, including mediators such as tumor necrosis factor and interleukin-1. They regulate inflammation, immune responses, fever, and communication among cells.

## Monomer

A monomer is a relatively small molecule that can chemically bond with other monomers to form a polymer. Biological examples include amino acids and nucleotides; some industrial monomers can irritate tissues or be toxic.

## Monomethylhydrazine

Monomethylhydrazine is a volatile, corrosive, highly toxic chemical used as a rocket propellant. Exposure can injure the eyes, skin, lungs, liver, and nervous system and may cause seizures or systemic poisoning.

## Mononegavirales

An order of viruses with nonsegmented, negative-sense RNA genomes, including rabies, measles, Ebola, and respiratory syncytial viruses.

## Mononeuritis Multiplex

A painful, asymmetrical, asynchronous sensory and motor peripheral neuropathy affecting at least two separate, non-contiguous named peripheral nerves, classically provoked by systemic necrotizing vasculitides (such as polyarteritis nodosa, microscopic polyangiitis, or cryoglobulinemia).

### Alternate Names:

Multifocal Neuropathy

## Mononeuropathy

Mononeuropathy is dysfunction or injury of a sin-

---

gle peripheral nerve. Common causes include compression, trauma, ischemia, inflammation, and metabolic disease; symptoms follow the nerve's distribution and may include pain, numbness, weakness, or autonomic changes.

### Mononuclear Leukocyte

A white blood cell with one main, unsegmented nucleus. Monocytes and lymphocytes are examples; they help remove damaged material, recognize germs, and coordinate immune responses.

### Mononucleosis

Common shortened term; see *Infectious mononucleosis*.

### Mononucleosis Spot Test

A rapid blood test detecting heterophile antibodies associated with infectious mononucleosis; it may be negative early in illness or in young children, so other testing may be needed.

### Monoplegia

Complete paralysis of one arm or one leg. It can result from a stroke, brain or spinal-cord disease, or severe damage to a nerve. The cause is found through a neurological examination and, when needed, brain or spine imaging and nerve tests.

### Monopodium

A single foot or one-sided foot structure; the term is uncommon and may be used descriptively in developmental anatomy.

### Monorchidism

The presence of only one testis, either because the other failed to develop, descended, or was removed.

### Monosaccharide

The simplest form of sugar, which cannot be split into a smaller carbohydrate. Glucose and fructose are examples; cells use these sugars mainly for energy.

### Monosodium Glutamate

MSG, the sodium salt of glutamic acid, used as a flavor enhancer because it contributes umami taste; it occurs naturally in foods such as tomatoes and aged cheese.

### Monosodium Urate

Needle-shaped crystals formed from uric acid that can collect in joints and other tissues when the blood uric-acid level is high. They cause the inflammation of gout and may be identified in joint fluid with a microscope.

### Monosomy

A genetic condition in which a cell has one copy of a chromosome instead of the usual two. Effects depend on the chromosome and whether all cells or only some cells are affected.

### Monospot Test

A rapid blood test that detects heterophile antibodies associated with infectious mononucleosis. It may be negative early in illness or in young children and does not directly detect Epstein-Barr virus; specific antibody testing may be needed.

#### Alternate Names:

Heterophile Antibody Test; Paul-Bunnell Test

### Monotest

A rapid blood test that looks for heterophile antibodies produced during some cases of infectious mononucleosis. A negative result does not always exclude infection, especially early in illness or in young children.

### Monounsaturated Fat

A fat whose fatty acids contain one carbon-carbon double bond. It occurs in foods such as olive oil, avocados, nuts, and some seeds.

### Monozygotic Twin

A monozygotic twin develops when one fertilized egg divides into two embryos. The twins share nearly identical nuclear genomes but can differ through developmental, epigenetic, mitochondrial, and envi-



---

ronmental factors.

### **Mons Veneris**

The rounded fatty prominence over the pubic bone in front of the external female genitalia.

### **Monteggia Fracture**

A traumatic fracture of the proximal third of the ulnar shaft accompanied by dislocation or subluxation of the radial head at the proximal radioulnar joint (PRUJ), classified into four types by Bado.

#### **Alternate Names:**

Monteggia Fracture-Dislocation; Proximal Ulnar Fracture with Radial Dislocation

### **Monteggia Fracture-Dislocation**

A traumatic forearm injury consisting of a fracture of the proximal third of the ulna associated with anterior, posterior, or lateral dislocation of the radial head at the proximal radioulnar joint.

### **Montelukast**

Montelukast is a leukotriene CysLT<sub>1</sub>-receptor antagonist used for asthma prevention and allergic rhinitis; it is not a rescue bronchodilator, and neuropsychiatric adverse effects require attention.

### **Montelukast Sodium**

Montelukast sodium is the sodium salt of montelukast, a leukotriene-receptor antagonist used for asthma prevention and allergic rhinitis. It is not a rescue bronchodilator and carries an important warning for possible neuropsychiatric adverse effects.

### **Montevideo Units**

A measurement of the strength of uterine contractions over 10 minutes during labor. It is calculated from pressure readings obtained by a catheter inside the uterus and helps clinicians assess whether contractions are strong enough to support labor progress.

### **Montgomery Tubercles**

Small oil-producing glands in the darker skin around

the nipple. They often become more noticeable during pregnancy and breastfeeding and release oil that helps protect and lubricate the nipple; they are usually normal.

### **Montreal Cognitive Assessment**

A brief screening test of memory, attention, language, problem-solving, visual-spatial skills, and orientation. It can identify possible thinking problems but does not by itself diagnose dementia; education, language, sensory impairment, and acute illness affect the score.

### **Mood**

A lasting emotional state experienced by a person and noticed by others. Mood may be described as depressed, anxious, irritable, elevated, or stable and can change with illness, medicines, stress, or life events.

### **Mood Disorder**

Alternate name; see *Mood Disorders*.

### **Mood Disorders**

A group of mental disorders characterized primarily by clinically significant depression, mania, hypomania, or combinations of these mood episodes. The group includes depressive disorders, bipolar disorders, and cyclothymic disorder.

### **Mood Stabilizers**

Medicines used mainly to treat or prevent episodes of mania and depression in bipolar disorder. Examples include lithium, valproate, carbamazepine, and lamotrigine; each has different benefits, interactions, monitoring needs, and risks during pregnancy.

### **Mood Swing**

A mood swing is a noticeable change in emotional state or affect over time. Occasional changes can be normal, while frequent, severe, or impairing swings may occur with mood disorders, substance use, medical illness, medications, or sleep disruption.

---

## Morale

Morale is the confidence, motivation, and sense of shared purpose within a person or group. It can affect teamwork and occupational well-being but is not itself a specific medical diagnosis.

## Moraxella

Moraxella is a genus of bacteria that can colonize the respiratory tract and cause ear, sinus, eye, or respiratory infections.

## Moraxella Atlantae

Moraxella atlantae is a rare Moraxella species; its recovery from a clinical sample should be interpreted with the site and evidence of infection.

## Moraxellaceae

A family of bacteria that includes Moraxella and related organisms. Some normally live on the skin or in the airways, while others can cause ear, sinus, lung, wound, or bloodstream infections.

## Moraxella Lincolnii

Moraxella lincolnii is a rare respiratory-tract bacterium; clinical importance depends on the specimen and evidence of infection.

## Moraxella Oblonga

Moraxella oblonga is a rare Moraxella species found in respiratory or oral specimens; significance depends on clinical evidence of infection.

## Moraxella Osloensis

An uncommon bacterium that can occasionally cause bloodstream, respiratory, wound, or other opportunistic infections, usually in people with serious illness or weakened immunity.

## Morbidity

Illness or disability in a person or group of people. It can also mean how many people have a disease or how much illness or disability a disease causes.

## Morbilli

The medical or Latin name for measles, an acute

contagious viral infection with fever and a characteristic rash.

## Morbilli Form

Resembling measles in appearance, especially a widespread rash made of flat and slightly raised spots. This description does not prove that a person has measles because medicines, infections, and other illnesses can cause similar rashes.

## Morbillivirus

A genus of paramyxoviruses that includes measles virus and related animal viruses causing fever, respiratory illness, rash, or neurologic disease.

## Morbus

A Latin word meaning disease. It appears in some older or formal disease names, but the word itself does not describe the cause, symptoms, or treatment of the condition.

## Morcellation

The division of tissue into smaller fragments, often during minimally invasive surgery; it may spread an unrecognized cancer and requires appropriate case selection.

## Mordant

A substance that helps fix a dye to tissue or a material, used in some histological staining methods.

## Morgagni Hernia

A congenital diaphragmatic hernia occurring through the anterior retrosternal foramen of Morgagni, resulting from incomplete fusion of parts of the diaphragm. Abdominal contents may move into the chest and cause respiratory or gastrointestinal symptoms.

## Morganella

Morganella is a genus of gram-negative bacteria that can cause urinary, wound, bloodstream, and other opportunistic infections.

---

## Morganella Morganii

A gram-negative bacterium that can cause urinary, wound, bloodstream, or postoperative infections, especially in hospitalized or medically vulnerable people. Antibiotic choice requires susceptibility testing because resistance is common.

## Morgellons Disease

Morgellons is a controversial condition involving distressing skin sensations and the belief that fibers or materials emerge from the skin. Symptoms are real to the person and require respectful evaluation for dermatologic, neurologic, psychiatric, toxic, or infectious causes without reinforcing an unproven explanation.

## Moribund

Critically ill and close to death, with little or no chance of recovery from the underlying condition. A moribund person may be unconscious, have very low blood pressure, cold or mottled skin, and irregular or failing breathing. Care focuses on urgent treatment when possible and comfort when death is inevitable.

## Morning Sickness

Nausea, with or without vomiting, during pregnancy. It often begins early and improves by the middle of pregnancy, but severe or persistent vomiting can cause dehydration, weight loss, and nutrient problems and may need medical treatment.

## Moro Reflex

A normal newborn startle response that causes the baby to spread out the arms, open the hands, and then draw the arms inward. It usually fades by about 3 to 6 months; an absent or unequal response may indicate a nerve or developmental problem.

## Morphea

A localized sclerosing skin disorder that produces areas of thickened, hardened, or discolored skin and differs from systemic sclerosis by usually lacking internal-organ involvement.

## Morpheus

The Greek and Roman mythological god of dreams;

the name is the source of *morphine*, an opioid named for its association with sleep and dreams.

## Morphinans

Morphinans are a group of chemical compounds with a fused-ring structure related to morphine. The class includes opioid analgesics and other compounds with varying receptor activity, potency, and clinical effects.

## Morphine

Morphine is a potent opioid analgesic used for selected moderate-to-severe pain and palliative-care indications. It activates opioid receptors and can cause respiratory depression, sedation, constipation, nausea, tolerance, dependence, and overdose.

## Morphine Overdose

Taking too much morphine can cause extreme sleepiness, pinpoint pupils, slow or stopped breathing, low blood pressure, coma, and death. Naloxone can sometimes reverse opioid-related breathing suppression.

## Morphine Sulfate

Morphine sulfate is a salt formulation of the opioid analgesic morphine. It is available in immediate- and extended-release preparations for selected pain indications and carries risks of respiratory depression, misuse, dependence, and fatal overdose.

## Morphogenesis

The biological process by which cells and tissues change shape, move, and organize to form organs and body structures during development or tissue repair.

## Morphologic Response

A morphologic response is a treatment-related change in the visible or microscopic structure of a lesion, tumor, tissue, or cell population. It may include shrinkage, necrosis, differentiation, or altered cellular features and is interpreted using disease-specific criteria.

---

## Morphology

The study or description of the form, shape, and structural features of organisms, organs, tissues, cells, or other biological structures.

## Morquio Syndrome

A rare inherited disorder in which complex sugars build up and interfere mainly with bone and cartilage growth. It can cause short stature, an unusually shaped chest and spine, loose joints, cloudy corneas, and pressure on the spinal cord in the neck. Intelligence is usually unaffected. Treatment may include enzyme replacement, surgery, and careful monitoring of breathing and the spine.

### Alternate Names:

Mucopolysaccharidosis Type Iv; MPS IV

## Mortality

Death, or the number of deaths in a particular group during a stated period. A mortality measure may describe all causes of death or deaths from a specific disease and population.

## Mortality Rate

The number of deaths in a specified population and time period, often reported for a particular disease or cause. It helps compare the impact of illness between groups when the populations are defined appropriately.

## Morton Neuroma

Thickening and irritation of a nerve in the ball of the foot, usually between the third and fourth toes. It can cause burning pain, numbness, or the feeling of a pebble under the foot, often worsened by tight shoes or walking.

### Alternate Names:

Interdigital Neuroma; Morton's Metatarsalgia; Planar Interdigital Neuroma

## Morton's Metatarsalgia

Pain in the forefoot beneath the metatarsal heads, caused by pressure or irritation and distinct from Morton neuroma.

## Morton's Neuroma

Thickening and irritation of tissue around a nerve, usually between the third and fourth toes. It causes burning forefoot pain or a feeling of a pebble in the shoe and may improve with wider footwear.

## Morton's Toe

A foot shape in which the second toe is longer than the big toe, often because the first metatarsal is relatively short. It is usually a normal variation but may contribute to pressure or forefoot pain.

## Morula

A morula is an early embryo consisting of a compact ball of blastomeres formed after several cleavage divisions. It develops into a blastocyst before implantation in the uterine lining.

## Morvan's Disease

A rare autoimmune neurologic syndrome with severe insomnia, hallucinations, confusion, pain, and peripheral nerve hyperexcitability.

## Mosaic

Describing a person or tissue composed of two or more genetically different cell populations derived from one zygote, often because of a post-zygotic mutation or chromosomal event; mosaicism can alter the severity and distribution of a genetic disorder.

## Mosaicism

The presence of groups of cells in one person that have different genetic material, even though all the cells came from the same fertilized egg. It may or may not cause disease, and its effects depend on which cells are affected.

## Mosquito

A small blood-feeding insect whose bites may transmit malaria, dengue, West Nile virus, and other infections.

## Mosquito-Borne

Spread by mosquitoes, usually describing an infection or parasite carried from one host to another by

---

a mosquito bite. Examples include malaria, dengue, West Nile virus, yellow fever, and some forms of encephalitis.

### **Mosquito Clamp**

A small hemostatic clamp used to compress small blood vessels.

### **Motilin**

A hormone made mainly in the small intestine that helps trigger waves of movement through the stomach and small intestine, especially between meals. Its activity contributes to the pattern that clears leftover food and secretions from the upper digestive tract.

### **Motility**

The ability of an organ to move its contents or itself. Intestinal motility moves food and waste forward, while abnormal motility can cause pain, constipation, diarrhea, vomiting, or swallowing problems.

### **Motion Analysis**

A detailed study of how a person moves during walking, reaching, or other activities. Clinicians use measurements and video to find movement problems, understand their causes, plan treatment, and track improvement.

### **Motion Artifact**

Image distortion or blurring caused by movement of the patient, organs, or equipment during image acquisition.

### **Motion Correction**

Processing or acquisition methods that reduce image blurring and errors caused by patient, breathing, or organ motion. Examples include image registration, tracking, gating, and navigator methods.

### **Motion Perception**

Motion perception is the ability of the visual, vestibular, and proprioceptive systems to detect and interpret movement of the body or surrounding objects. Disturbance may cause vertigo, oscillopsia,

imbalance, or visual-motion sensitivity.

### **Motion Sickness**

Nausea, dizziness, sweating, or vomiting caused by sensory mismatch during travel or other motion. It can occur in cars, boats, aircraft, amusement rides, or virtual-reality environments and usually improves when motion stops. Severe or persistent symptoms require assessment for other causes.

### **Motivation**

Motivation is the set of processes that initiate, direct, and sustain goal-directed behavior. Reduced or excessive motivation may occur with mood disorders, neurologic disease, substance use, medication effects, or other conditions.

### **Motivational Interviewing**

A collaborative counseling approach that helps a person explore mixed feelings and strengthen their own reasons for changing a health behavior. The clinician listens without judgment, asks open questions, reflects the person's views, and supports informed choice.

### **Motor**

Relating to movement or to the nerves and brain pathways that control movement. A motor problem may cause weakness, stiffness, poor coordination, tremor, or loss of normal control.

### **Motor Aphasia**

Difficulty speaking or writing caused by damage to language areas of the brain, often after a stroke. A person may know what they want to say but struggle to find words, form sentences, or produce speech.

### **Motor Cortex**

Areas at the front of the brain that plan and control voluntary movement. Each side mainly controls the opposite side of the body. Injury can cause weakness, loss of fine movement, or difficulty planning purposeful actions.

---

## Motor Delay

Motor delay is slower-than-expected acquisition of gross- or fine-motor skills in a child. It may reflect neuromuscular, developmental, genetic, sensory, environmental, or systemic causes and warrants age-appropriate developmental assessment.

## Motor End-Plate

The specialized area where a nerve signal reaches a skeletal muscle fiber. Chemical signals released there start muscle contraction, and damage can cause weakness, fatigue, or abnormal muscle function.

## Motor Endplate

The specialized, highly folded post-synaptic sarcolemmal region of a skeletal muscle fiber directly underlying an axon terminal at the neuromuscular junction, densely packed with nicotinic acetylcholine receptors and acetylcholinesterase enzymes.

### Alternate Names:

Neuromuscular Junction Postsynaptic Membrane

## Motor Incoordination

Impaired coordination of voluntary movement, often caused by disease affecting the cerebellum or other parts of the nervous system.

### Alternate Names:

Ataxia

## Motor Learning

A relatively lasting change in the ability to perform a motor skill resulting from practice, experience, feedback, and adaptation. Rehabilitation uses task-specific repetition and meaningful practice to promote motor learning.

## Motor Neuron

A nerve cell that carries signals from the brain or spinal cord to skeletal muscle, allowing movement. Damage can cause weakness, muscle wasting, stiffness, cramps, or loss of voluntary control.

### Alternate Names:

Motor Nerve Cell; Motoneuron

## Motor Neuron Disease

Alternate spelling; see *Motor Neurone Disease*.

## Motor Neuron Disorders Pathology

Pathologic changes affecting upper or lower motor neurons, which may include neuronal loss, chromatolysis, gliosis, corticospinal-tract degeneration, muscle denervation atrophy, and variable involvement of brainstem or spinal motor nuclei.

## Motor Neurone Disease

A group of progressive disorders affecting upper and lower motor neurons, including amyotrophic lateral sclerosis. They cause worsening weakness, muscle wasting, stiffness, cramps, and difficulty with movement; speech, swallowing, and breathing may become affected as the disease advances. Cognitive or behavioral changes occur in some forms.

## Motor Neurones

Nerve cells that carry signals from the brain or spinal cord to skeletal muscles, producing movement. Injury or disease can cause weakness, wasting, stiffness, cramps, poor coordination, or loss of voluntary control.

## Motor Stereotypy

A repeated, predictable movement such as hand flapping, rocking, or body waving. It may occur during normal development or with autism, intellectual disability, neurologic disease, stress, or other conditions.

## Motor Tic

A motor tic is a sudden, brief, repetitive movement such as blinking, facial grimacing, or shoulder shrugging. Tics are often temporarily suppressible and may occur in transient tic disorder, persistent tic disorder, or Tourette syndrome.

## Motor Unit

One movement-controlling nerve cell and all the muscle fibers it activates. Small units allow precise movements, while large units produce more force. Damage to motor units can cause weakness, wasting, cramps, or abnormal results on nerve and muscle



---

tests.

### **Motor Vehicle Accident**

A collision or other road-traffic event causing injury or property damage. Medical evaluation may be needed for visible or delayed injuries, including head, neck, chest, abdominal, or musculoskeletal trauma.

### **Mottled Enamel**

Patchy, discolored, or structurally altered tooth enamel, classically caused by excessive fluoride during tooth development.

### **Mottled Teeth**

Teeth with irregular white, brown, or pitted enamel, often due to dental fluorosis or developmental enamel defects.

### **Moulage**

A model, cast, or realistic reproduction of a body part or lesion used for teaching, simulation, or forensic documentation.

### **Mould**

British spelling of mold, a multicellular fungus that grows as branching filaments.

### **Mountain Sickness**

Illness caused by rapid ascent to high altitude, ranging from acute mountain sickness to life-threatening pulmonary or cerebral edema.

### **Mouth**

The oral cavity that serves as the entry point of the digestive tract, used for eating, breathing, and speaking.

### **Mouth Cancer**

Cancer arising in the lips, gums, tongue, cheeks, floor of the mouth, palate, or other oral tissues.

#### **Alternate Names:**

Oral Cancer; Oral Cavity Carcinoma; Oral Squamous Cell Carcinoma

### **Mouth Diseases**

Disorders affecting the oral cavity, including infections, ulcers, inflammatory disease, dental disease, mucosal lesions, and tumors.

### **Mouth Neoplasm**

A mouth neoplasm is an abnormal growth in the oral cavity that may be benign or malignant; persistent sores, lumps, or patches need assessment.

### **Mouth Sores**

Painful ulcers, blisters, raw areas, or inflamed patches inside or around the mouth. Trauma, infection, immune disease, medicines, nutritional deficiency, and cancer can cause them; sores lasting more than two weeks need assessment.

### **Mouth-To-Mouth Resuscitation**

Rescue breathing in which air is delivered from a rescuer to a person who is not breathing normally, usually as part of CPR.

### **Mouth Ulcer**

Open sores affecting the oral mucosa, commonly caused by aphthous ulcers, trauma, infection, irritation, or systemic inflammatory disease. Most resolve within two weeks; persistent, recurrent, unusually large, or indurated ulcers require evaluation.

#### **Alternate Names:**

Oral Ulcer; Aphthous Ulcer; Canker Sore

### **Mouthwash**

A liquid rinsed around the mouth and then expelled to freshen breath, deliver an active ingredient, or reduce selected oral microorganisms.

### **Mouthwash Overdose**

Swallowing too much mouthwash may cause alcohol poisoning, vomiting, drowsiness, low blood sugar, breathing problems, or coma; products containing fluoride or other ingredients can cause additional toxicity.

### **Movement**

Movement is a change in position of the body or

---

a body part produced by coordinated activity of the nervous system, muscles, and joints. Abnormal movement may be excessive, reduced, involuntary, poorly coordinated, or restricted.

### **Movement Disorder**

A neurologic condition causing excessive, reduced, or otherwise abnormal movement. Hyperkinetic disorders include tremor, dystonia, chorea, myoclonus, tics, and dyskinesia; hypokinetic disorders include bradykinesia and parkinsonism.

### **Movement Disorders**

Alternate name; see *Movement Disorders*.

### **Movement Incoordination**

Impairment in the ability to execute smooth, coordinated voluntary movements (ataxia), resulting from lesions or dysfunction within the cerebellum, spinocerebellar tracts, vestibular apparatus, or proprioceptive pathways. Manifests as dysmetria, intention tremor, and gait instability.

#### **Alternate Names:**

Ataxic Incoordination; Motor Incoordination; Dysdiadochokinesia

### **Moxalactam**

Moxalactam is a beta-lactam antibiotic of the oxacephem group with activity against many gram-negative bacteria and some gram-positive organisms. It is rarely used today and may cause allergic reactions, diarrhea, or bleeding related to impaired coagulation.

### **Moxibustion**

A traditional treatment that warms selected skin areas with burning mugwort or another heated material. Evidence of benefit varies by condition, and burns, smoke exposure, allergic reactions, or infection can occur.

### **Moxifloxacin**

An antibiotic used for selected bacterial infections. It interferes with bacterial reproduction but can cause serious side effects, including tendon injury, nerve symptoms, blood-sugar changes, or dangerous

heart rhythms, so use is selective.

### **Moyamoya Disease**

A rare condition in which major arteries supplying the brain gradually narrow or close. Small fragile blood vessels form to compensate, but the condition can cause transient attacks, strokes, headaches, seizures, or thinking problems.

### **Moynihan's Hump**

An anatomical variation in which the right hepatic artery forms an abnormally tortuous, caterpillar-like loop that extends close to the gallbladder neck and cystic duct within Calot's triangle. During gallbladder removal (cholecystectomy), this loop can be mistaken for the smaller cystic artery; accidental clipping or injury can cause severe arterial bleeding or damage blood flow to the right lobe of the liver.

#### **Alternate Names:**

Moynihan Caterpillar Turn; Caterpillar Turn of the Right Hepatic Artery; Hepatic Artery Tortuosity

### **MPH**

Master of Public Health, a graduate degree focused on population health, disease prevention, health policy, epidemiology, and related public-health practice.

### **MPO**

Myeloperoxidase, an enzyme in neutrophil and monocyte granules that uses hydrogen peroxide and halide ions to generate antimicrobial oxidants; MPO antibodies are also used in evaluating some autoimmune vasculitides.

### **Mpox**

A viral disease caused by monkeypox virus, an orthopoxvirus. It spreads mainly through close contact with lesions, body fluids, respiratory particles, or contaminated objects and may cause fever, swollen lymph nodes, fatigue, and painful lesions on the skin or mucosal surfaces; severe disease is more likely in children, pregnancy, and immunocompromise.

#### **Alternate Names:**

Simian Pox; Monkeypox Virus Infection

---

## **MPTP**

A toxic chemical that can permanently damage movement-controlling nerve cells and produce symptoms resembling Parkinson disease. It is mainly important in research and is not a routine medical treatment.

## **MRA**

Magnetic resonance angiography, a noninvasive MRI-based technique that produces images of blood vessels and can help evaluate aneurysms, stenoses, malformations, and atherosclerosis.

## **MR Angiography**

An MRI scan that creates pictures of blood vessels without using X-rays. It can help assess narrowed or blocked arteries, aneurysms, dissections, clots, and abnormal blood-vessel connections.

## **MRCP**

Magnetic resonance cholangiopancreatography, an MRI technique that noninvasively images the bile ducts, gallbladder, and pancreatic duct without requiring endoscopic catheterization.

## **MRD**

Minimal residual disease: a small number of malignant cells remaining after treatment, detectable by sensitive methods such as flow cytometry or PCR even when routine tests show remission.

## **MR Enterography**

A specialized MRI scan of the small intestine after drinking contrast liquid. It can show inflammation, narrowing, fistulas, abscesses, bleeding, or other complications of Crohn disease and does not use X-rays.

## **MRI**

Magnetic resonance imaging uses a strong magnetic field, radiofrequency energy, and computer processing to produce detailed images of body tissues without ionizing radiation. Contrast material may be used when clinically appropriate.

## **MRI And Low Back Pain**

MRI may be used for low back pain when serious disease is suspected, neurologic deficits are present, or symptoms persist despite appropriate care; many uncomplicated cases do not need early MRI.

## **MRI Brain**

An MRI scan used to examine the brain for tumors, stroke, bleeding, infection, inflammation, multiple sclerosis, or degenerative disease. Different settings highlight different tissues and abnormalities; contrast may be used when needed.

## **MRI Contrast**

A contrast agent administered during magnetic resonance imaging to improve visibility of selected tissues or abnormalities.

## **MRI Joint**

Evaluation of musculoskeletal structures: cartilage, ligaments, tendons, menisci, labrum. Knee: ACL, meniscus tears. Shoulder: rotator cuff, labrum. Hip: labral tears, FAI.

## **MRI Sequences**

Different image settings used during an MRI scan to highlight anatomy, fluid, fat, bleeding, restricted water movement, or inflammation. Radiologists combine several settings because each shows different types of abnormality.

## **MRI-Targeted Fusion Prostate Biopsy**

A prostate biopsy guided by combining a previous MRI with real-time ultrasound. The images help the clinician direct needles toward areas that look suspicious for cancer, sometimes along with routine samples from other areas. The procedure can cause blood in the urine or semen, infection, pain, or temporary difficulty passing urine.

### **Alternate Names:**

Fusion Biopsy; Mri-Ultrasound Fusion Biopsy

## **Mrna Precursor**

An initial RNA transcript that is processed by capping, splicing, and polyadenylation to form mature

---

messenger RNA.

### MR Neurography

Magnetic resonance imaging optimized to show peripheral nerves and abnormalities such as swelling, compression, or injury.

### MRSA

A strain of *Staphylococcus aureus* resistant to several common antibiotics. It may live harmlessly on the skin or cause skin, wound, lung, bone, joint, or bloodstream infection; treatment is guided by culture results.

### MRSA Decolonization

Measures intended to reduce MRSA carriage on the skin or in the nose. Protocols may include topical agents and antiseptic bathing, but the indication, regimen, and duration depend on the person's risk, infection history, setting, and local guidance.

### MRSA Infection

Methicillin-resistant *Staphylococcus aureus* is a strain of staphylococcus resistant to several beta-lactam antibiotics. It can colonize without illness or cause skin, wound, lung, bone, joint, or bloodstream infection; culture-guided treatment and hygiene reduce spread.

#### Alternate Names:

Methicillin-Resistant *Staphylococcus Aureus* Infection

### MS Alternative Therapies

Alternate name; see *Alternative Therapy for Multiple Sclerosis*.

### MSG Symptom Complex

A group of symptoms such as headache, flushing, or palpitations attributed by some people to monosodium glutamate; evidence for a consistent syndrome is limited and symptoms may have other causes.

### MST

Multiple subpial transection, a neurosurgical pro-

cedure in which shallow cuts are made in the cerebral cortex to interrupt seizure pathways when the epileptogenic region cannot safely be removed.

### MTOR

A protein inside cells that helps control growth, division, energy use, and response to nutrients. Medicines that block mTOR, such as rapamycin-related drugs, are used in some cancers, transplant treatments, and other selected conditions.

### Mucha-Habermann Disease

An acute, severe, self-limiting variant of pityriasis lichenoides (PLEVA) characterized by the sudden eruption of widespread erythematous macules that rapidly evolve into necrotic, papulovesicular, and hemorrhagic crusted lesions that heal leaving small varioliform (pox-like) scars.

#### Alternate Names:

PLEVA; Pityriasis Lichenoides et Varioliformis Acuta

### Mu-Chain Disease

A rare B-cell disorder in which abnormal immunoglobulin heavy chains are produced, sometimes causing anemia, enlarged lymph nodes, or other features of lymphoproliferative disease.

### Mucin

Mucin is a high-molecular-weight glycoprotein that forms mucus and helps lubricate and protect epithelial surfaces. Abnormal mucin production or accumulation occurs in some respiratory, gastrointestinal, and neoplastic diseases.

### Mucinoses

A group of disorders in which excessive mucin, a gel-like substance in connective tissue, builds up in the skin or other tissues. Causes may be primary, metabolic, autoimmune, or associated with another disease.

### Mucinous

Mucinous means containing, producing, or resembling mucin. In pathology, the term describes lesions or tumors with extracellular or intracellular mucin

---

and must be interpreted with the organ, architecture, and other histologic features.

### **Mucinous Adenocarcinoma**

Mucinous adenocarcinoma is a gland-forming cancer that produces abundant mucus and can arise in organs such as the colon, pancreas, lung, or breast.

### **Mucinous Borderline Ovarian Tumor**

An ovarian mucinous epithelial tumor with cellular proliferation and atypia but without destructive stromal invasion; it may be large and unilateral.

### **Mucinous Cystadenoma**

A mucinous cystadenoma is a usually benign cystic epithelial tumor that produces mucin, commonly arising in the ovary or pancreas. Large lesions may cause pain or pressure, and pathology is needed to distinguish benign from borderline or malignant mucinous tumors.

### **Mucinous Cystadenoma of Pancreas**

Alternate name; see *Pancreatic Cysts*.

### **Mucinous Cyst Of The Vulva**

A benign mucus-filled cyst of the vulva, arising from obstruction or metaplastic mucinous epithelium in a vulvar gland or duct; persistent or atypical masses require examination and possible biopsy.

### **Mucinous Ovarian Carcinoma**

A malignant ovarian epithelial tumor composed of invasive mucin-producing cells; metastatic gastrointestinal or pancreatobiliary carcinoma must be excluded.

### **Muckle-Wells Syndrome**

An inherited cryopyrin-associated autoinflammatory syndrome causing recurrent fever, rash, joint or muscle pain, and progressive sensorineural hearing loss. Anti-inflammatory treatment can reduce inflammation and prevent complications.

### **Mucocele**

A soft, mucus-filled swelling caused by leakage or

blockage of a small salivary gland, most often on the inner lower lip after minor injury. It may come and go, but a persistent or recurrent swelling should be examined and may need removal.

### **Mucocele and Ranula**

Mucus-filled lesions caused by leakage or blockage of a salivary gland duct. A mucocele usually affects the lower lip, while a ranula arises in the floor of the mouth; persistent, recurrent, or large lesions may need dental or oral-surgery treatment.

### **Mucociliary Escalator**

A critical pulmonary innate defense mechanism consisting of ciliated pseudo-stratified columnar epithelial cells and goblet-cell-derived mucus lining the conducting tracheobronchial tree, whose coordinated rhythmic upward ciliary beating sweeps inhaled particulate matter and pathogens out of the respiratory tract toward the pharynx.

### **Mucocutaneous Leishmaniasis**

A chronic, mutilating metastatic form of leishmaniasis caused by New World species of the *Leishmania* subgenus *Viannia* (principally *Leishmania braziliensis*), which disseminates from a primary cutaneous ulcer to mucosal tissues through hematogenous or lymphatic routes. It causes progressive, destructive granulomatous ulceration and cartilaginous necrosis of the nasal septum, palate, pharynx, and larynx, clinically presenting with nasal obstruction, epistaxis, septal perforation, and severe facial disfigurement.

#### **Alternate Names:**

Espundia; American Mucocutaneous Leishmaniasis; Nasopharyngeal Leishmaniasis

### **Mucocutaneous Lymph Node Syndrome**

Alternate name; see *Kawasaki Disease*.

### **Mucoepidermoid Carcinoma**

Mucoepidermoid carcinoma is a cancer containing mucus-producing and squamous-like cells, most often arising in salivary glands or the lungs.

---

## Mucoid

Resembling mucus or having a thick, slippery, mucus-like appearance. The word describes a substance or finding and does not by itself identify its cause.

## Mucolipidoses

A group of inherited lysosomal storage disorders in which cells cannot properly process certain lipids and carbohydrate-containing substances. Material accumulates in tissues and may cause developmental delay, skeletal changes, vision problems, and organ disease.

## Mucolytics

Medicines that make thick mucus thinner and easier to clear from the airways. They may help some people cough up mucus, but treatment depends on the cause of congestion and the person's overall condition.

## Mucolytic Therapy

Treatment that makes thick mucus easier to move and cough up. Different medicines work by breaking mucus proteins, removing extra DNA, or drawing water into airway secretions; the choice depends on the disease and the person's ability to clear mucus.

## Mucopolysaccharides

Long chains of sugar molecules that form part of connective tissue and help support joints, skin, cartilage, and other structures. Inherited problems breaking them down can cause storage diseases.

## Mucopolysaccharidoses

A group of inherited lysosomal storage diseases in which glycosaminoglycans accumulate because one of several degradation enzymes is deficient. The disorders can affect the skeleton, joints, airways, heart, eyes, abdominal organs, and nervous system, with severity varying by type.

## Mucopolysaccharidosis

The singular name for any of a group of inherited lysosomal storage disorders in which glycosaminoglycans accumulate because of deficient enzyme ac-

tivity. Types differ in their genetic cause, age of onset, and effects on the skeleton, organs, airways, eyes, heart, and nervous system.

### Alternate Names:

Gargoylism (deprecated historical term)

## Mucopolysaccharidosis Type I

An inherited lysosomal storage disorder caused by deficient alpha-L-iduronidase, usually because of disease-causing variants in the *IDUA* gene. It ranges from severe Hurler syndrome to milder forms and may affect the skeleton, airways, heart, eyes, abdominal organs, and nervous system.

### Alternate Names:

Hurler Syndrome; Hurler-Scheie Syndrome; Scheie Syndrome

## Mucopolysaccharidosis Type II

Mucopolysaccharidosis type II, or Hunter syndrome, is an X-linked iduronate-2-sulfatase deficiency causing accumulation of glycosaminoglycans and multi-system disease.

## Mucopolysaccharidosis Type III

Mucopolysaccharidosis type III, or Sanfilippo syndrome, is an inherited disorder of heparan-sulfate breakdown causing progressive neurodevelopmental and behavioral decline.

## Mucopolysaccharidosis Type IV

Mucopolysaccharidosis type IV, or Morquio syndrome, is an inherited disorder of keratan-sulfate breakdown causing skeletal and connective-tissue abnormalities with variable neurologic involvement.

## Mucopolysaccharidosis VII

Mucopolysaccharidosis VII, or Sly syndrome, is a lysosomal storage disorder caused by deficient beta-glucuronidase activity. Accumulation of glycosaminoglycans can cause skeletal, connective-tissue, visceral, developmental, and sometimes neurologic abnormalities.

## Mucopus

A mixture of mucus and pus produced by inflamma-



---

tion or infection. It may appear in the nose, sinuses, airways, bowel, cervix, or other mucus-lined areas and should be interpreted with symptoms.

### **Mucor**

Mucor is a genus of molds; related Mucorales fungi can cause mucormycosis, a serious invasive infection mainly in people with severe immune or metabolic problems.

### **Mucorales**

Mucorales is an order of molds that includes fungi capable of causing mucormycosis, a rapidly invasive infection of the sinuses, lungs, skin, or other tissues.

### **Mucormycosis**

A rare but rapidly destructive infection caused by certain molds. It most often affects people with severely weakened immunity or uncontrolled diabetes and can damage the sinuses, eyes, lungs, skin, brain, or several organs.

### **Mucor Ramosissimus**

A type of mold found in the environment that rarely causes infection in people. When infection occurs, it is usually in someone with severely weakened immune defenses and requires identification of the fungus and specialist treatment.

### **Mucosa**

Mucosa is the moist epithelial lining of body passages such as the mouth, nose, airways, gastrointestinal tract, and reproductive or urinary tracts.

### **Mucosa-Associated Lymphoid Tissue**

Collections of immune tissue in the moist linings of the digestive, breathing, and urinary or reproductive tracts. They help detect and respond to substances and germs entering through these surfaces.

### **Mucosa-Associated Lymphoid Tissue Lymphoma**

Alternate name; see *MALT Lymphoma*.

### **Mucosal**

Relating to a mucous membrane, the moist lining of areas such as the mouth, nose, airways, digestive tract, and genital or urinary tract. Mucosa helps protect tissues, produce mucus, absorb substances in some organs, and support local immune defense.

### **Mucosal Candidiasis**

Candida yeast infection of a mucous membrane, commonly producing white plaques, soreness, redness, or burning in the mouth or genital tract. Antibiotics, diabetes, inhaled steroids, immune suppression, and moisture can increase risk.

### **Mucosal Immunity**

Mucosal immunity is immune defense at surfaces lining the respiratory, gastrointestinal, genitourinary, and other tracts. It includes epithelial barriers, mucus, secretory IgA, antimicrobial molecules, innate cells, and locally regulated adaptive responses.

### **Mucosal Infection**

An infection involving a moist epithelial lining such as the mouth, eye, nose, airway, gastrointestinal tract, or genital tract.

### **Mucosal Irritation**

Inflammation or injury of moist linings in the eyes, nose, throat, mouth, stomach, or intestines. It may cause burning, pain, redness, swelling, or sores after chemicals, medicines, infection, or friction.

### **Mucosal Melanoma**

A melanoma arising from melanocytes in mucosal surfaces such as the sinonasal tract, oral cavity, anorectal region, or genital tract; it is uncommon and often diagnosed at an advanced stage.

### **Mucosal Warts**

Warts caused by human papillomavirus on moist lining tissues such as the mouth, genitals, or anus. They may be painless or cause itching, bleeding, or discomfort; some HPV types increase cancer risk and need medical assessment.

---

## Mucositis

Painful inflammation and sores in the moist lining of the mouth or digestive tract, often after chemotherapy or radiation. It can make eating and swallowing difficult and may increase the risk of infection.

## Mucous

Relating to mucus, the viscous fluid produced by mucous membranes and glands that lubricates surfaces and helps trap particles and microorganisms.

## Mucous Cysts

Small fluid-filled lumps near the end joint of a finger, often associated with osteoarthritis. They may cause tenderness, nail changes, or leakage and should not be punctured without medical advice.

### Alternate Names:

Digital Mucous Cyst; Myxoid Cyst

## Mucous Membrane

A moist tissue lining body cavities that open to the outside, including the nose, mouth, and lungs.

### Alternate Names:

Mucosa

## Mucous Membrane (Cicatricial) Pemphigoid (MMP)

A group of autoimmune subepidermal blistering disorders affecting mucous membranes, especially the mouth, eyes, nose, and genitals. Repeated inflammation can scar and threaten vision, airway function, or other organ function, so specialist diagnosis and treatment are important.

## Mucous Membrane Pemphigoid

A chronic autoimmune subepidermal blistering disease affecting mucous membranes, especially the eyes and mouth; scarring can impair vision or function.

## Mucus

Mucus is a viscoelastic secretion containing water, mucin glycoproteins, salts, antimicrobial factors, and cellular material. It lubricates and protects epithelial surfaces and traps particles and microor-

ganisms for clearance.

## Mucus Plugging

Accumulation of thick, tenacious mucus within airways causing obstruction. It is common in asthma, CF, bronchiectasis, and chronic bronchitis. It impairs gas exchange, promotes infection, and causes atelectasis. Treatment includes airway clearance techniques, mucolytics, and bronchodilators.

## Muehrcke Lines

Paired, white, non-palpable, transverse parallel bands spanning the width of the nail bed, separated by normal pink nail bed, which do not migrate with nail growth and blanch with digital pressure (apparent leukonychia), characteristic of severe hypoalbuminemia (serum albumin less than 2 g/dL).

### Alternate Names:

Muehrcke's Lines; Paired White Bands of Hypoalbuminemia

## Muehrcke Lines of the Fingernails

Paired transverse white bands in the nail bed that remain fixed as the nail grows. They may occur with low blood protein, severe illness, chemotherapy, or other systemic stress and usually improve when the underlying problem resolves.

## Muehrcke's Nails

Paired transverse white bands in the nail beds, usually associated with hypoalbuminemia or other systemic illness. The bands remain fixed as the nail grows because they arise from the nail bed rather than the nail plate.

## Muenke Syndrome

A craniosynostosis syndrome caused by a pathogenic FGFR3 variant. Premature fusion of one or more skull sutures may produce an abnormal head shape and can be accompanied by developmental, hearing, limb, or neurological findings.

## MUGA Scan

A heart scan that uses a small radioactive tracer and records the heart's pumping with each heartbeat. It measures how much blood the main chambers

---

pump and can assess wall movement.

**Alternate Names:**

Radionuclide Angiography; Radionuclide Ventriculography

**Muir-Torre Syndrome**

An autosomal dominant cancer predisposition syndrome (a variant of Lynch syndrome) caused by germline mutations in DNA mismatch repair genes (MSH2, MLH1), characterized by the combination of at least one sebaceous gland neoplasm (sebaceous adenoma, sebaceoma) or keratoacanthomas with internal visceral malignancies (especially colorectal and genitourinary carcinomas).

**Alternate Names:**

MTS; Sebaceous Neoplasm and Visceral Cancer Syndrome

**Mulberry Molars**

Permanent back teeth with many small, rounded, irregular cusps. They are classically linked with congenital syphilis but can also result from other developmental problems affecting tooth formation before birth.

**Mulibrey Nanism**

Mulibrey nanism is a rare inherited multisystem disorder, usually caused by biallelic variants in *TRIM37*. It may cause severe growth restriction, characteristic facial features, pericardial or hepatic disease, insulin resistance, and other organ involvement.

**Muller Cells**

Supporting glial cells of the retina that help nourish retinal nerve cells and maintain the retina's structure and chemical environment.

**Mullerian Agenesis**

A congenital developmental anomaly in 46,XX females resulting from embryonic failure of the paramesonephric (Müllerian) ducts to develop, causing congenital absence of the upper two-thirds of the vagina, cervix, and uterus, with normal functional ovaries and secondary sexual characteristics.

**Alternate Names:**

Mayer-Rokitansky-Kuster-Hauser Syndrome; MRKH Syndrome

**Mullerian Duct Anomalies**

Congenital differences in development of the uterus, cervix, fallopian tubes, or upper vagina. Examples include septate, bicornuate, didelphys, unicornuate, and agenesis patterns; they may cause menstrual obstruction, infertility, pregnancy loss, or preterm birth.

**Multicenter Study**

A multicenter study is a clinical or research investigation conducted at two or more sites under a coordinated protocol. It can improve enrollment and generalizability but requires harmonized methods, quality control, and site-specific oversight.

**Multicentric**

Having more than one center or focus; in oncology, describing separate primary tumor foci, often also called multifocal or polycentric.

**Multicentric Reticulohistiocytosis**

A rare systemic non-Langerhans cell histiocytosis characterized by papulonodular skin or mucosal lesions and often destructive inflammatory arthritis. It may involve other organs and can be associated with an underlying malignancy, so evaluation is multidisciplinary.

**Multicystic Dysplastic Kidney**

A birth defect in which one kidney develops as a cluster of cysts and has little or no working tissue. The other kidney often provides most kidney function but may also need monitoring for related urinary problems.

**Multidetector Computed Tomography**

A CT scan that uses many detectors to collect multiple thin image slices at the same time. It produces detailed pictures quickly and is useful for injuries, blood vessels, organs, and some virtual examinations, but uses X-rays.

**Multidisciplinary Disaster Victim**

---

## Identification

The coordinated process of identifying victims of mass-fatality incidents (natural disasters, transport crashes, terrorist attacks) using multiple disciplines: fingerprinting, dental records, DNA profiling, forensic anthropology/radiology, and personal effects, with ante-mortem data collection and comparison.

## Multidisciplinary Team

Healthcare professionals from different fields who work together on one person's care. Depending on the problem, the team may include doctors, nurses, therapists, pharmacists, psychologists, dietitians, and social workers.

## Multidrug-Resistant TB

Tuberculosis caused by bacteria resistant to at least the two most important first-line medicines, isoniazid and rifampin. It needs specialized medicines, susceptibility testing, and close monitoring because treatment is harder and may last many months.

## Multidrug-Resistant Tuberculosis

Tuberculosis caused by bacteria resistant to at least isoniazid and rifampicin, two important tuberculosis medicines. It requires specialist testing and a longer combination treatment chosen according to drug-susceptibility results.

## Multienzyme Complexes

Groups of enzymes organized together so that several steps in a chemical pathway occur efficiently. This arrangement can speed reactions, limit loss of intermediates, and coordinate metabolism inside cells.

## Multifactorial

Caused or influenced by several factors rather than one single cause. These may include genes, age, lifestyle, infections, medicines, environment, and other health conditions, with each factor adding to the overall risk.

## Multifocal

Arising from or involving two or more distinct foci or sites, such as separate tumor foci or vascular

lesions.

## Multifocal Atrial Tachycardia

An atrial tachyarrhythmia with at least three distinct P-wave morphologies, irregular rhythm, and a rate above 100 beats per minute. It is strongly associated with severe pulmonary disease, hypoxia, and metabolic stress.

## Multifocal Motor Neuropathy

A rare, chronic, acquired autoimmune peripheral neuropathy characterized by slowly progressive, asymmetric, distal lower motor neuron weakness and muscle atrophy predominantly affecting the upper extremities without sensory loss. Characterized electrophysiologically by persistent multifocal motor conduction blocks outside usual entrapment sites, with high circulating titers of IgM anti-GM1 ganglioside antibodies in over 50 percent of patients; responds selectively to intravenous immunoglobulin.

### Alternate Names:

MMN; Multifocal Motor Neuropathy with Conduction Block

## Multigravida

A person who has been pregnant two or more times, including pregnancies that did not continue.

## Multi-Infarct Dementia

A decline in memory and thinking caused by several strokes that damage different parts of the brain. Problems may include slowed thinking, poor planning, mood changes, walking difficulty, or memory loss; the term is now often included under vascular cognitive impairment.

### Alternate Names:

Vascular Dementia; Vascular Cognitive Impairment

## Multilocular

Having several separate spaces or compartments. The word may describe a cyst, tumor, or other lesion seen on imaging; its significance depends on the organ involved, the appearance of the walls, and the person's symptoms.

---

### Multi-Minicore Disease

A rare inherited muscle disorder in which muscle fibers contain multiple small areas of damage. It can cause low muscle tone, weakness, spinal curvature, and breathing problems.

#### Alternate Names:

MmD; Minicore Myopathy; Multicore Disease

### Multimodal Analgesia

Use of analgesic techniques with different mechanisms to improve pain control while reducing reliance on any single drug, especially opioids. Components may include acetaminophen, nonsteroidal anti-inflammatory drugs, regional anesthesia, local anesthetics, and selected adjuvants.

### Multimodal Imaging

Use of two or more imaging methods, such as CT, MRI, ultrasound, or nuclear imaging, to examine the same disease or body region. Combining methods can provide complementary structural, functional, or molecular information.

### Multimorbidity

The coexistence of two or more chronic health conditions in the same person, without requiring one to be considered the primary disease. It complicates treatment, increases polypharmacy risk, and requires coordinated patient-centered care.

### Multinodular Goiter

An enlarged thyroid containing multiple nodules. Thyroid function may be normal, underactive, or overactive; evaluation commonly includes thyroid-function tests and ultrasound, with further assessment of suspicious nodules or compressive symptoms.

### Multinucleated Hepatocyte

A multinucleated hepatocyte is a liver cell containing two or more nuclei, usually from nuclear division without complete cytokinesis. It may be a normal or reactive finding, but its significance depends on the number, morphology, and associated liver abnormalities.

### Multipara

A person who has had two or more pregnancies that reached the stage of potential fetal viability, regardless of whether the pregnancies resulted in live births.

### Multiparametric MRI Of The Prostate

A detailed prostate MRI that combines several types of images to show prostate structure, water movement, and blood flow. It helps identify suspicious areas and guide biopsy or treatment, but an abnormal scan does not by itself prove cancer.

### Multiparous

Having given birth two or more times to a fetus or fetuses beyond the stage of viability.

### Multiple Bacterial Abscesses

Several pockets of pus caused by bacterial infection in one or more organs. They may develop when bacteria spread through the blood or extend from nearby tissue and can cause fever, pain, or organ problems; imaging helps locate them.

### Multiple Biliary Hamartomas

Multiple small, harmless birth-related changes in the liver's bile ducts that look like tiny cysts on scans. They usually cause no symptoms and are often found by chance, but imaging helps distinguish them from cancer or other liver disease.

### Multiple Endocrine Neoplasia

A group of inherited, usually autosomal dominant cancer-predisposition syndromes causing tumors or overactivity in two or more hormone-making glands during a person's lifetime. Types include MEN1, MEN2, and related subtypes with different gene associations and tumor risks.

### Multiple Endocrine Neoplasia I

MEN type 1 is an inherited tumor syndrome involving parathyroid, pituitary, and pancreatic or duodenal neuroendocrine tumors, often caused by MEN1 variants.



---

## Multiple Endocrine Neoplasia II

MEN type 2 is an inherited syndrome with medullary thyroid cancer, pheochromocytoma, and sometimes parathyroid disease, commonly involving RET variants.

## Multiple Endocrine Neoplasia Type 1

An inherited condition that causes tumors in several hormone-making organs, especially the parathyroid glands, pituitary gland, and pancreas or nearby intestine. It is usually caused by a change in the MEN1 gene and requires lifelong monitoring.

## Multiple Endocrine Neoplasia Type 2

An inherited condition that greatly increases the risk of medullary thyroid cancer and may cause adrenal or parathyroid tumors. Type 2B can also cause nerve-related mouth bumps and distinctive body features; lifelong specialist monitoring is needed.

## Multiple Endocrine Neoplasia Type 2A

An inherited condition that greatly increases the risk of medullary thyroid cancer and can also cause adrenal tumors and overactive parathyroid glands. Regular genetic and specialist monitoring helps detect these problems early.

### Alternate Names:

MEN 2A; MEN2A; Sipple Syndrome

## Multiple Endocrine Neoplasia Type 2B

An inherited RET-associated syndrome characterized by medullary thyroid carcinoma, pheochromocytoma, mucosal neuromas, and a marfanoid habitus. The thyroid-cancer risk is high and requires early specialist genetic and surgical assessment.

## Multiple Endocrine Neoplasia Type 2 (MEN2)

An inherited cancer-predisposition syndrome, usually caused by a pathogenic RET variant, featuring medullary thyroid carcinoma and, in some people, pheochromocytoma and parathyroid disease. Genetic counseling, biochemical testing, and preventive thyroid surgery may be needed.

## Multiple Endocrine Neoplasia Type 4

A rare, autosomal dominant inherited tumor syndrome caused by germline mutations in the *CDKN1B* gene, which encodes the p27<sup>Kip1</sup> tumor suppressor protein. It presents with features overlapping with MEN1, most commonly characterized by primary hyperparathyroidism (parathyroid adenomas causing high blood calcium) and anterior pituitary adenomas (such as prolactinomas or growth hormone-secreting tumors). Patients may also develop pancreatic neuroendocrine tumors, adrenal adenomas, and reproductive organ tumors. It is typically diagnosed in patients with MEN1-like symptoms who test negative for mutations in the *MEN1* gene.

### Alternate Names:

MEN 4; MEN4; Multiple Endocrine Neoplasia Type IV; CDKN1B-Related MEN

## Multiple Exostoses

Multiple exostoses, or hereditary multiple osteochondromas, is an inherited disorder causing multiple cartilage-capped bony projections near growth plates. Lesions can deform bones, restrict motion, compress nerves or vessels, and rarely undergo malignant transformation.

## Multiple Fibrofolliculomas

Numerous harmless small skin-colored growths arising from hair follicles, usually on the face and trunk. They can be a sign of Birt-Hogg-Dubé syndrome, which also increases the risk of kidney tumors and collapsed lung.

## Multiple Inherited Schwannomas

Alternate name; see *Neurofibromatosis Type 2*.

## Multiple Mononeuropathy

Damage to several separate nerves in different parts of the body, often developing unevenly. It can cause patchy pain, numbness, weakness, or muscle wasting and may result from blood-vessel inflammation, diabetes, infection, or another systemic disease.

## Multiple Myeloma

A cancer of plasma cells that usually develops in



---

the bone marrow and often produces a monoclonal immunoglobulin. It can cause bone lesions, anemia, kidney dysfunction, high calcium, infection, or other organ damage.

**Alternate Names:**

Plasma Cell Myeloma; Kahler Disease; Myelomatosis

### **Multiple Myeloma: Deadly To Chronic**

Multiple myeloma is a plasma-cell cancer that produces a monoclonal protein and can cause bone lesions, anemia, kidney injury, high calcium, infections, and fatigue. It is often treatable as a chronic illness, though relapse and complications remain possible.

### **Multiple Organ Dysfunction Syndrome**

Abnormal function of two or more organs during severe illness, such as sepsis, shock, trauma, or pancreatitis. Organ function may worsen over time and requires close monitoring and treatment of the underlying cause.

### **Multiple Organ Failure**

A life-threatening condition in which two or more organs stop working properly, usually during severe infection, shock, injury, or widespread inflammation. Intensive care supports the organs while doctors treat the underlying cause.

### **Multiple Pregnancy**

A pregnancy involving two or more fetuses. Twins may develop from two separate eggs or from one fertilized egg that divides; triplets and higher-order pregnancies are less common. Multiple pregnancies have higher risks of premature birth, growth problems, and pregnancy complications and require closer monitoring.

### **Multiple Sclerosis**

A chronic immune-mediated disease of the brain, spinal cord, and optic nerves. The immune system mistakenly attacks myelin, the protective covering around nerve fibers, and may also damage the nerve fibers themselves. This creates areas of inflammation and scarring, called lesions, that slow or block

nerve signals. Symptoms vary and may include vision changes, numbness, weakness, stiffness, poor coordination, balance problems, fatigue, thinking difficulties, or bladder and bowel problems. Some people have attacks followed by partial or complete recovery, while others develop steadily worsening symptoms; mixed patterns also occur.

**Alternate Names:**

MS; Disseminated Sclerosis

### **Multiple Sclerosis MS And Pregnancy**

Multiple sclerosis usually does not prevent pregnancy, but symptoms, disease-modifying medicines, relapse risk, delivery planning, and postpartum care require coordination. Some treatments must be stopped or changed, and relapse risk can shift after delivery.

### **Multiple Sleep Latency Test**

A daytime sleep study measuring how quickly a person falls asleep during several scheduled nap opportunities. It helps evaluate excessive daytime sleepiness and narcolepsy after nighttime sleep has been assessed.

### **Multiple System Atrophy**

A progressive brain disorder affecting movement and automatic body functions. It may cause stiffness and slowness, poor balance, fainting from low blood pressure, bladder problems, speech changes, or difficulty swallowing.

**Alternate Names:**

Shy-Drager Syndrome; MSA

### **Multiple System Atrophy Cerebellar Type**

An adult-onset neurodegenerative synucleinopathy (MSA-C) characterized primarily by progressive cerebellar ataxia (gait instability, limb dysmetria, dysarthria, oculomotor abnormalities) accompanied by severe autonomic failure and variable pyramidal signs.

**Alternate Names:**

MSA-C; Cerebellar Multiple System Atrophy; Olivopontocerebellar Atrophy Variant

---

## Multiple System Atrophy Parkinsonian Type

A progressive neurodegenerative disorder (MSA-P) characterized predominantly by parkinsonian features (rigidity, akinesia, postural instability) that respond poorly to levodopa, combined with early severe autonomic failure (orthostatic hypotension, urinary incontinence) and variable pyramidal dysfunction.

### Alternate Names:

MSA-P; Parkinsonian Multiple System Atrophy; Striatonigral Degeneration Variant

## Multiple System Failure

Failure of two or more organ systems, also called multiple-organ failure or multiple-organ dysfunction syndrome; it may occur during severe illness such as sepsis, shock, or major trauma.

## Multiple Vitamin Overdose

Taking too many multivitamins can cause poisoning, especially from iron, vitamin A, or vitamin D. Symptoms may include vomiting, abdominal pain, confusion, liver injury, kidney injury, or dangerous calcium levels; urgent advice is needed.

## Multiplex PCR Panel

A laboratory test that looks for genetic material from several germs in one sample. Different panels test respiratory, intestinal, blood, or other specimens; a detected germ may represent infection, harmless carriage, or material left after an infection.

### Alternate Names:

Multiplex Polymerase Chain Reaction; Syndromic Molecular Panel

## Multiplicity

Multiplicity is the state of having more than one occurrence, copy, lesion, tumor, or pregnancy component. Its interpretation depends on context, such as multiple gestation, multifocal disease, or copy number in genetics.

## Multipolar Neuron

A nerve cell with one long output branch and several receiving branches. This common type of neuron

includes many cells that control movement or pass signals within the brain and spinal cord.

## Multisystem Inflammatory Syndrome In Children

A serious delayed inflammatory illness occurring in some children after infection with SARS-CoV-2, involving fever and two or more organ systems and potentially causing shock or cardiac dysfunction.

## Multivesicular Body

A multivesicular body is an endosomal compartment containing many small intraluminal vesicles. It can fuse with the plasma membrane to release exosomes or with a lysosome to degrade its cargo.

## Mumps

A contagious viral infection caused by a paramyxovirus and spread through respiratory droplets, saliva, and direct contact. It typically begins with fever, headache, and fatigue, followed by tender swelling of one or both parotid salivary glands (parotitis) below and in front of the ears, making chewing and swallowing painful. Complications are more common in adolescents and adults and include orchitis (painful testicular inflammation that can cause atrophy and reduced fertility), viral meningitis, pancreatitis, and permanent sensorineural hearing loss. Treatment is supportive with rest, fluids, and pain relief. It is highly preventable with the standard two-dose Measles-Mumps-Rubella (MMR) vaccine.

### Alternate Names:

Epidemic Parotitis; Mumps Parotitis; Paramyxoviral Parotitis

## Mumps Orchitis

Acute inflammation of the testis occurring in approximately 20 to 30 percent of post-pubertal males infected with the mumps paramyxovirus, presenting with sudden severe testicular pain, unilateral or bilateral scrotal edema, high fever, and risking subsequent testicular atrophy and impaired spermatogenesis.

---

## Mumps Parotitis

Alternate name; see *Mumps*.

## Mumps Virus

The virus that causes mumps, a contagious infection spread through saliva and respiratory droplets. It commonly causes fever and painful swelling near the jaw, and can sometimes inflame the brain, pancreas, or reproductive organs; vaccination reduces risk.

## Munchausen Syndrome

Alternate name; see *Factitious Disorder*.

## Mupirocin

Mupirocin is a topical antibiotic that inhibits bacterial isoleucyl-tRNA synthetase. It is used for selected skin infections and nasal eradication of certain staphylococci; resistance can develop with inappropriate or prolonged use.

## Mural

Relating to a wall, especially the wall of a body cavity, organ, or blood vessel.

## Mural Thrombosis

Formation of a blood clot along the wall of a heart chamber or blood vessel rather than freely within the lumen.

## Mural Thrombus

A mural thrombus is a blood clot that forms on the wall of a heart chamber or large blood vessel, typically in areas of damaged endothelium, stasis, or turbulent flow.

## Muramidase

Muramidase, or lysozyme, is an enzyme that cleaves the peptidoglycan backbone of bacterial cell walls. It is present in secretions and innate immune cells and contributes to defense against susceptible bacteria.

## Murine Typhus

An acute endemic rickettsial illness caused by *Rickettsia typhi* and transmitted to humans primarily through the feces of infected rodent fleas (*Xenop-*

*sylla cheopis*). It typically presents with high prolonged fever, severe frontal headache, myalgias, arthralgias, and a maculopapular rash that typically spares the face, palms, and soles.

### Alternate Names:

Endemic Typhus; Flea-Borne Typhus

## Murmur

An extra sound heard during a heartbeat, usually caused by fast or turbulent blood flow through the heart or nearby blood vessels. Some murmurs are harmless, while others may be caused by heart valve or structural problems.

## Murmur Classification

Heart murmurs are classified by timing, grade, shape, location, radiation, and pitch. Timing: systolic (between S1 and S2), diastolic (between S2 and S1), or continuous. Grade: I (barely audible) to VI (heard without stethoscope). Shape: crescendo, decrescendo, or plateau.

## Murphy Sign

A sudden catch or stop in breathing when a doctor presses under the right ribcage as the patient takes a deep breath in, indicating an inflamed gallbladder.

## Murphy's Sign

Inspiratory arrest caused by pain when the right upper quadrant is palpated, traditionally suggesting acute cholecystitis.

## Musca

A fly genus including the housefly; some species can spread germs by carrying them on their bodies.

## Muscae Volitantes

Moving spots, threads, or cobweb-like shapes seen in the visual field, usually caused by shadows from vitreous opacities.

## Muscarine

Muscarine is a mushroom-derived agonist of muscarinic acetylcholine receptors. Ingestion of muscarine-containing mushrooms can cause a cholin-

---

ergic syndrome with salivation, sweating, tearing, abdominal cramps, vomiting, diarrhea, bronchospasm, and bradycardia.

### **Muscarinic Antagonist**

A muscarinic antagonist blocks acetylcholine at muscarinic receptors in smooth muscle, glands, the heart, and the central nervous system. Examples include atropine, ipratropium, and oxybutynin; effects can include dry mouth, blurred vision, urinary retention, tachycardia, and confusion.

#### **Alternate Names:**

Antimuscarinic

### **Muscidae**

Muscidae is a family of flies that includes the housefly. Some species can mechanically carry pathogens to food or wounds, and their larvae may occasionally cause myiasis.

### **Muscimol**

Muscimol is a psychoactive compound found in some *Amanita* mushrooms and an agonist at GABA-A receptors. Ingestion can cause altered perception, sedation, agitation, incoordination, or seizures and may require toxicologic evaluation.

### **Muscle**

Body tissue that shortens to create force and movement. Skeletal muscle moves the bones and supports posture, heart muscle pumps blood, and smooth muscle moves substances through organs such as the bowel and blood vessels.

### **Muscle Aches**

Pain or soreness in one or more muscles. Common causes include overuse, injury, infection, medicines, and conditions affecting the whole body.

### **Muscle Atrophy**

Loss or wasting of muscle tissue caused by disuse, nerve injury, inflammation, malnutrition, or systemic disease. It may lead to weakness and reduced function; treatment focuses on the cause, nutrition, and appropriately supervised rehabilitation.

### **Muscle Belly**

The thick, fleshy middle part of a muscle between its attached ends. It contains most of the muscle fibers that shorten to produce force.

### **Muscle Biopsy**

Removal of a small sample of skeletal muscle for microscopic, histochemical, immunologic, or molecular examination in suspected myopathy, inflammation, infection, or infiltrative disease.

### **Muscle Cell**

A cell specialized to contract; skeletal, cardiac, and smooth muscle cells perform different movement and organ functions.

### **Muscle Contraction**

The tightening of a muscle, allowing it to produce force or movement. Muscle fibers use calcium and energy from ATP to make their protein filaments slide against each other.

### **Muscle-Contraction Headache**

A pressure-like headache caused by prolonged tightening or sensitivity of muscles in the scalp, neck, or jaw. Stress, poor posture, fatigue, or jaw clenching may contribute; frequent, severe, or changing headaches need medical assessment.

#### **Alternate Names:**

Tension-Type Headache; Stress Headache

### **Muscle Contracture**

Persistent shortening or tightening of a muscle or nearby tissues that limits joint movement and does not fully relax with stretching. It may follow scarring, prolonged spasticity, immobility, injury, or, more rarely, a muscle-energy disorder.

#### **Alternate Names:**

Skeletal Muscle Contracture; Metabolic Contracture

### **Muscle Cramp**

A muscle cramp is a sudden, involuntary, painful contraction of a muscle. It may follow exertion, dehydration, electrolyte disturbance, pregnancy, medication use, nerve disease, or vascular disease; recur-

---

rent cramps need assessment of the cause.

### **Muscle Cramps**

Alternate plural form; see *Muscle Cramp*.

#### **Alternate Names:**

Charley Horse; Involuntary Muscle Spasm

### **Muscle Development**

Muscle development is the formation, growth, differentiation, and maturation of muscle tissue. It depends on genetic programs, neural input, hormones, nutrition, mechanical loading, and repair processes.

### **Muscle Disorder**

A disease affecting muscle structure or function, causing weakness, pain, cramps, stiffness, wasting, or abnormal movement. Causes include inherited conditions, inflammation, nerve disease, medicines, infection, and metabolic problems.

### **Muscle Fascicle**

A bundle of muscle fibers surrounded by connective tissue. Several fascicles join to form a muscle.

### **Muscle Fatigue**

A reduced ability of a muscle to produce force after prolonged or repeated activity. It may result from exertion, poor sleep, illness, medicines, nerve problems, or muscle disease, and improves when the cause is treated.

### **Muscle Fiber**

A muscle fiber is an individual muscle cell containing contractile myofibrils. Skeletal muscle fibers vary in size and metabolic properties, while cardiac and smooth muscle cells have distinct structures and functions.

### **Muscle Fibers**

Individual muscle cells containing contractile proteins. Skeletal muscle fibers vary in size and properties, including endurance-oriented slow fibers and powerful, faster-fatiguing fibers.

#### **Alternate Names:**

Muscle Cells; Myocytes

### **Muscle Function Loss**

Reduced ability of a muscle or muscle group to generate movement or force, caused by muscle disease, nerve injury, neuromuscular-junction disorders, pain, or disuse.

### **Muscle Hypertrophy**

Muscle hypertrophy is enlargement of individual muscle fibers, usually in response to resistance, hormonal stimulation, or compensatory workload. It can be physiologic or associated with disease, depending on the muscle and cause.

### **Muscle Insertion**

The attachment of a muscle to the body part that usually moves when the muscle contracts. The insertion is often farther from the body's center than the muscle's origin.

### **Muscle-Invasive Bladder Cancer**

Bladder cancer that has grown into the bladder muscle layer. It has a greater risk of spreading than cancer limited to the bladder lining and is classified by examination of bladder tissue.

### **Muscle Neoplasm**

A muscle neoplasm is an abnormal growth in skeletal or smooth muscle or nearby soft tissue and may be benign or malignant.

### **Muscle Origin**

The attachment of a muscle to the body part that usually stays more fixed when the muscle contracts. It is often closer to the body's center than the muscle's insertion.

### **Muscle Pain**

Pain, soreness, or aching in one or more muscles caused by exertion, injury, infection, inflammation, medicines, or an underlying medical condition. Persistent, severe, or weakness-associated pain needs assessment.

#### **Alternate Names:**

---

Myalgia; Muscle Ache; Myodynia

### **Muscle Protein**

A protein found in muscle that provides structure, produces contraction, stores or transports substances, or regulates muscle activity. Important examples include actin, myosin, troponin, dystrophin, and myoglobin.

### **Muscle Relaxant**

A medicine that reduces skeletal-muscle spasms or tightness. Some act in the brain or spinal cord and may cause sleepiness or weakness. Neuromuscular-blocking medicines are a distinct group that temporarily paralyzes muscles and requires breathing support and close monitoring.

### **Muscle Relaxants**

See *Muscle Relaxant*.

### **Muscle Relaxation**

The return of a muscle to a less tense state after contraction, when calcium levels inside the muscle cell fall and the contractile proteins stop generating force.

### **Muscle Rigidity**

Muscle rigidity is abnormal resistance to passive movement caused by sustained muscle contraction or altered tone. It occurs in conditions such as Parkinson disease, dystonia, meningitis, neuroleptic malignant syndrome, and tetanus.

### **Muscle Spasm**

A sudden, involuntary tightening of a muscle that may cause pain, stiffness, or an abnormal position. Spasms can follow overuse, dehydration, mineral changes, nerve problems, medicines, pregnancy, or injury.

### **Muscle Spasms**

Muscle spasms are sudden involuntary contractions caused by exertion, dehydration, electrolyte disturbance, nerve disease, medicines, pregnancy, or musculoskeletal problems. Recurrent, widespread,

painful, or weakness-associated spasms need evaluation for an underlying cause.

### **Muscle Spindle**

A small sensor inside skeletal muscle that detects how much the muscle is stretched and how quickly it changes length. It helps maintain posture, muscle tone, balance, and coordinated movement.

### **Muscle Strain Treatment**

Care for a stretched or torn muscle, usually involving temporary activity reduction, gradual movement, rehabilitation, and pain relief chosen safely for the person. Severe swelling, weakness, deformity, inability to use the limb, or persistent pain needs assessment.

### **Muscle Strength**

Muscle strength is the maximal force a muscle or muscle group can generate under specified conditions. It is assessed by functional or resisted testing and may be reduced by muscle, nerve, neuromuscular-junction, central, or systemic disorders.

### **Muscle-Tendon Unit**

The connected working unit made of a muscle and its tendon. It produces force and transfers that force to a bone or other structure.

### **Muscle Tissue**

Specialized tissue that contracts to produce movement, maintain posture, move substances through organs, and generate heat. Skeletal, heart, and smooth muscle have different structures and control systems for their jobs.

### **Muscle Tone**

The slight, continuous activity present in resting muscles that gives them some resistance to being stretched. Abnormally high or low muscle tone can occur with disorders of the brain, spinal cord, nerves, or muscles.



---

## **Muscle Tonus**

The slight, continuous activity present in resting muscles, which gives them resistance to being stretched. Too much or too little tone can occur with disorders of the brain, spinal cord, nerves, or muscles.

## **Muscle Twitching**

Brief involuntary contractions of a small group of muscle fibers, often called fasciculations; persistent or widespread twitching may require evaluation.

## **Muscle Weakness**

Reduced ability to generate force with one or more muscles. Causes include nerve or muscle disease, pain, inactivity, medicines, metabolic disorders, and aging.

## **Muscular**

Relating to muscle or made of muscle tissue. It may describe strength, body build, pain, weakness, or a structure containing muscle.

## **Muscular Dystrophy**

A group of more than 30 inherited muscle diseases that cause progressive weakness and breakdown of skeletal muscles. Different gene changes affect proteins that strengthen and protect muscle fibers, so the age of onset, muscles affected, and severity vary by type. Weakness may cause difficulty walking or doing everyday activities. Some forms also affect the heart, breathing muscles, swallowing, bones, brain, or other organs. Muscular dystrophy is not contagious and is not caused by ordinary injury or physical activity.

### **Alternate Names:**

MD; Inherited Myopathy

## **Muscular Endurance**

The ability of a muscle or muscle group to keep working during repeated or prolonged effort. It depends on muscle strength, energy supply, blood flow, nerve control, conditioning, and overall health.

## **Muscularis**

A layer of smooth muscle in the wall of a hollow organ, especially the digestive tract. Its coordinated contractions move contents through the organ and can also change the diameter or pressure of the passage.

### **Alternate Names:**

Muscular Layer; Tunica Muscularis

## **Muscular System**

The body system made up mainly of skeletal, smooth, and heart muscle. It produces movement, maintains posture, moves substances through organs, pumps blood, and generates heat.

## **Musculocutaneous Nerve**

A nerve from the neck and upper chest that controls several muscles used to bend the elbow and turn the forearm. It also carries sensation from part of the outer forearm; injury can cause weakness or numbness there.

## **Musculoskeletal**

Relating to muscles, bones, joints, tendons, ligaments, and related tissues.

## **Musculoskeletal Deformity**

A musculoskeletal deformity is an abnormal shape, alignment, or structure of bone, joint, muscle, or related tissue. It may be congenital, developmental, traumatic, degenerative, neuromuscular, or caused by disease.

## **Musculoskeletal Disease**

A disorder affecting bones, joints, muscles, tendons, ligaments, cartilage, or other supporting tissues, causing pain or limited movement.

## **Musculoskeletal Injury**

Trauma to muscles, bones, joints, tendons, ligaments, cartilage, or related soft tissues, ranging from strains and sprains to fractures and dislocations.

---

## Musculoskeletal Pain

Pain arising from muscles, bones, joints, ligaments, tendons, or related soft tissues. Common causes include injury, overuse, arthritis, and inflammation; persistent pain, weakness, swelling, or neurologic symptoms should be assessed.

## Musculoskeletal System

The musculoskeletal system consists of bones, joints, cartilage, ligaments, tendons, muscles, and associated connective tissues. It provides support, protects organs, enables movement, and contributes to mineral storage and blood-cell production.

## Musculoskeletal Tumor

An abnormal growth arising in bone, muscle, cartilage, connective tissue, or related structures. It may be benign or malignant and can cause a lump, pain, swelling, reduced function, or fracture.

## Musculus

The Latin word for muscle, used in formal anatomical names. For example, “musculus deltoideus” means the deltoid muscle. Abbreviations may be used in anatomy texts to identify one or several muscles.

## MUSE

A medicated urethral system that delivers a pellet into the urethra to treat erectile dysfunction.

### Alternate Names:

Medicated Urethral System for Erection

## Mushroom

The visible fruiting body of a fungus. Edible mushrooms can provide nutrients, but wild mushrooms may be poisonous or confused with safe species; uncertain mushrooms should not be eaten.

## Mushroom Poisoning

Illness caused by eating toxic fungi, with effects ranging from gastrointestinal upset to liver failure, neurologic toxicity, or death.

## Mushroom Toxicity

Poisoning from an edible-looking or otherwise toxic mushroom, with syndromes ranging from early gastrointestinal illness or hallucinations to delayed hepatic or renal failure. The species, amount, and time to symptoms are important; contact a poison center, preserve a specimen or photograph, and provide syndrome-specific supportive treatment.

## Music Therapy

The planned use of music by a trained therapist to support health and wellbeing. It may help with pain, stress, communication, movement, memory, emotional expression, or coping during illness and rehabilitation.

## MuSK-Positive Myasthenia Gravis

A form of myasthenia gravis in which antibodies interfere with nerve-to-muscle communication. It often causes weakness of the face, mouth, throat, neck, or breathing muscles and requires specialist care.

### Alternate Names:

MuSK Myasthenia Gravis; MuSK-Ab-Positive MG

## Mustard Compounds

Mustard compounds are chemical agents containing reactive sulfur or nitrogen mustard groups that can alkylate DNA and other molecules. Some are chemotherapy drugs; others are highly toxic chemical-warfare agents.

## Mustard Gas

Mustard gas, or sulfur mustard, is a vesicant chemical-warfare agent that damages skin, eyes, and respiratory tissues and can cause blistering, airway injury, and delayed complications. It is not a true gas at ordinary temperatures and has no specific widely available antidote.

## Mutagen

A chemical, physical agent, or biological factor capable of causing a genetic mutation. Mutagenicity does not by itself prove that an agent causes cancer.

---

## Mutant

A mutant is a cell, organism, or genetic sequence carrying a mutation relative to a reference or parental form. The change may have no effect, alter a trait, or cause disease, depending on its location and functional consequence.

## Mutant Proteins

Proteins whose amino-acid sequence differs because of a genetic mutation. The change may have no effect or may alter folding, stability, location, or function and contribute to inherited disease, cancer, or drug resistance.

## Mutate

To mutate means to undergo a change in genetic sequence. Mutations arise spontaneously or from replication errors, DNA damage, or mutagen exposure and may be inherited, acquired in somatic cells, harmful, neutral, or beneficial.

## Mutation

A heritable or acquired alteration in the DNA sequence. Variants may be single-nucleotide substitutions, insertions, deletions, copy-number changes, or larger rearrangements and may be benign, pathogenic, or of uncertain significance.

## Mutation Abnormality

A mutation abnormality is an atypical change in DNA sequence or chromosome structure identified relative to a reference. Its clinical significance must be classified using evidence, such as pathogenic, likely pathogenic, uncertain significance, likely benign, or benign.

## Mutation Rate

The frequency at which new DNA changes arise in a cell, organism, or generation. The rate differs between genes and species and can be influenced by DNA-copying errors, repair systems, radiation, chemicals, or other factors.

## Mute

Describing a person who does not speak, whether because of a physical, neurologic, developmental, or

psychological reason or by choice; current clinical language should describe the specific communication limitation or preference.

## Mutism

Absence or refusal of speech despite adequate ability to speak, occurring in conditions such as catatonia, selective mutism, or neurologic disease.

## Myalgia

Muscle pain or aching. It may affect one area after injury or exercise, or occur throughout the body with infection, inflammation, medicine effects, thyroid disease, or other conditions. Severe pain with weakness or dark urine needs prompt medical assessment because muscle breakdown may be occurring.

### Alternate Names:

Myodynia; Muscle Pain

## Myalgic Encephalomyelitis

A complex illness involving substantial activity-limiting fatigue, post-exertional worsening, unrefreshing sleep, and often cognitive or orthostatic symptoms.

### Alternate Names:

ME/CFS; SEID; Systemic Exertional Intolerance Disease

## Myalgic Encephalomyelitis/Chronic Fatigue Syndrome

A complex illness characterized by substantial reduction in activity, post-exertional malaise, unrefreshing sleep, and cognitive or autonomic symptoms. Management is individualized and focuses on pacing, symptom relief, and treatment of coexisting conditions.

## Myasthenia

Muscle weakness that often becomes more pronounced with activity and lessens after rest. The term may refer generally to this pattern of weakness or to specific disorders such as myasthenia gravis.

## Myasthenia Crisis

A life-threatening complication of myasthenia gravis

---

characterized by severe exacerbation of respiratory muscle weakness producing impending acute ventilatory failure, often triggered by systemic infection, surgery, or medication withdrawal, requiring endotracheal intubation and treatment with plasma-pheresis or intravenous immunoglobulin.

### **Myasthenia Gravis**

A chronic autoimmune disorder of the neuromuscular junction, where nerves communicate with skeletal muscles. Antibodies most often target acetylcholine receptors and sometimes proteins such as muscle-specific kinase (MuSK) or LRP4, disrupting signals that cause muscles to contract. Weakness typically worsens with repeated use and improves with rest. It may affect the eyelids and eye movements, facial expression, chewing, speech, swallowing, neck, arms, legs, or breathing muscles. Sensation is usually preserved; thymus abnormalities, including thymoma, may also occur.

#### **Alternate Names:**

MG

### **Myasthenia Gravis and Pregnancy**

Pregnancy and the postpartum period can alter autoimmune neuromuscular weakness, while some medicines and anesthesia may worsen respiratory or swallowing function. Neurology and obstetric planning addresses medication safety, delivery, neonatal weakness, and crisis risk.

### **Myasthenic Crisis**

A life-threatening worsening of myasthenia gravis that weakens the muscles needed for breathing or swallowing. Infection, surgery, stress, pregnancy, or certain medicines may trigger it. Severe breathlessness, choking, inability to swallow, or rapidly worsening weakness may occur.

#### **Alternate Names:**

Myasthenia Gravis Crisis; Acute Myasthenic Exacerbation

### **Myasthenic Syndrome**

A disorder of neuromuscular transmission causing fluctuating weakness; it may refer to myasthenia gravis or a congenital or Lambert-Eaton syndrome

depending on context.

### **Mycelium**

The network of fine branching threads that makes up the growing part of a fungus. It spreads through soil, food, or tissue and can produce the structures that form fungal spores.

### **Mycetoma**

A chronic localized infection of skin and subcutaneous tissue caused by fungi or filamentous bacteria, producing swelling, sinuses, and grains.

### **Mycobacteriaceae**

A family of slow-growing or rapidly growing, acid-fast bacteria that includes *Mycobacterium* species causing tuberculosis, leprosy, and nontuberculous infections.

### **Mycobacterial Culture**

A laboratory test that tries to grow *Mycobacterium* bacteria from a sample such as sputum, tissue, or fluid. It can identify tuberculosis or other mycobacteria and may show which medicines will work, but results can take weeks.

### **Mycobacterial Infection**

Infection caused by bacteria in the genus *Mycobacterium*. It includes tuberculosis, leprosy, and nontuberculous mycobacterial disease; symptoms and treatment depend on the species and affected organs.

### **Mycobacterium**

A genus of acid-fast bacteria that includes the causes of tuberculosis and leprosy and several nontuberculous infections.

### **Mycobacterium Abscessus**

A rapidly growing nontuberculous mycobacterium that can cause pulmonary, skin, and soft-tissue infections and is often difficult to treat because of multidrug resistance.

---

### **Mycobacterium Avium**

A member of the *Mycobacterium avium* complex that can cause lung disease, lymph-node infection, or disseminated infection, especially in people with weakened immunity.

### **Mycobacterium Avium Complex**

A group of nontuberculous mycobacteria that can cause chronic pulmonary disease or disseminated infection, particularly in people with impaired immunity. Treatment depends on the species, site, severity, susceptibility, and patient factors.

### **Mycobacterium Bovis**

A tuberculosis-complex bacterium that mainly infects cattle but can cause tuberculosis in humans through unpasteurized dairy or animal exposure.

### **Mycobacterium Chelonae**

A rapidly growing nontuberculous mycobacterium that can cause skin, soft-tissue, bone, eye, or disseminated infection.

### **Mycobacterium Fortuitum**

A rapidly growing nontuberculous mycobacterium that can cause wound, catheter, bone, lung, or other opportunistic infections.

### **Mycobacterium Gordona**

A usually nonpathogenic water-associated nontuberculous mycobacterium that can occasionally cause disease in people with weakened immunity.

### **Mycobacterium Haemophilum**

A nontuberculous mycobacterium that can cause skin, joint, and bone infections, especially in people with impaired immunity.

### **Mycobacterium Infection**

Infection caused by a *Mycobacterium* species, including tuberculosis, leprosy, and nontuberculous mycobacterial disease.

### **Mycobacterium Kansaii**

A nontuberculous mycobacterium that commonly

causes chronic lung disease resembling tuberculosis and can occasionally cause disseminated infection.

### **Mycobacterium Leprae**

The bacterium that causes leprosy, a long-lasting infection affecting the skin and nerves. It usually spreads only after prolonged close contact, develops slowly, and can cause numbness, weakness, or skin lesions without treatment.

### **Mycobacterium Marinum**

A nontuberculous mycobacterium associated with fish tanks and aquatic exposure that can cause chronic skin and soft-tissue infection.

### **Mycobacterium Scrofulaceum**

A nontuberculous mycobacterium associated especially with cervical lymph-node infection in children and occasional other disease.

### **Mycobacterium Szulgai**

A rare nontuberculous bacterium that can cause long-lasting lung disease or other infections, especially in people with underlying lung problems or weakened immunity. Diagnosis requires specialist laboratory testing and clinical assessment.

### **Mycobacterium Tuberculosis**

The bacterium that causes tuberculosis. It spreads through the air and most often affects the lungs, but can involve other organs. Latent infection has no symptoms and does not spread; active disease may cause cough, fever, night sweats, or weight loss.

### **Mycobacterium Ulcerans**

A mycobacterium that causes Buruli ulcer, a chronic destructive skin infection associated with the toxin mycolactone.

### **Mycology**

The branch of microbiology that studies fungi, including yeasts, moulds, and mushrooms. Medical mycology focuses on fungal pathogens that cause disease in humans. Fungal infections (mycoses) are classified as superficial, cutaneous, subcutaneous,

---

or systemic.

### **Mycophenolate Mofetil**

Mycophenolate mofetil is a prodrug of mycophenolic acid that inhibits inosine monophosphate dehydrogenase and lymphocyte proliferation. It is used as an immunosuppressant, especially after organ transplantation, and can cause infection, cytopenias, gastrointestinal effects, and serious fetal harm.

### **Mycophenolic Acid**

The active immune-suppressing substance formed from mycophenolate medicines. It blocks an enzyme needed by multiplying lymphocytes and helps prevent organ-transplant rejection, but can cause infections, low blood counts, diarrhea, and fetal harm.

### **Mycoplasma**

A genus of very small bacteria lacking a cell wall, including organisms that cause respiratory and genitourinary infections.

### **Mycoplasma Genitalium**

A sexually transmitted bacterium that can cause urethritis, cervicitis, pelvic inflammatory disease, and, less commonly, proctitis. Many infections are asymptomatic. Diagnosis is usually by a nucleic acid amplification test, and macrolide resistance is common, so treatment should follow current susceptibility-informed guidance.

### **Mycoplasma Hominis**

A wall-less bacterium associated with the genital tract that can cause pelvic, postpartum, urinary, wound, or joint infection.

### **Mycoplasmal Pneumonia**

Pneumonia caused by *Mycoplasma* species, most often *Mycoplasma pneumoniae*, usually producing a gradual cough and fever.

### **Mycoplasma Pneumonia**

Atypical pneumonia most common in ages 5–35. Dry cough, low-grade fever, headache. Patchy in-

filtrates on CXR disproportionate to symptoms (“walking pneumonia”).

### **Mycoplasma Pneumoniae**

A bacterium without a cell wall that can cause bronchitis, sore throat, or pneumonia, especially in school-age children and young adults. Illness often develops gradually with cough, fever, headache, or tiredness; complications outside the lungs are uncommon but possible.

### **Mycoplasma Pneumoniae-Associated Mucositis**

Inflammatory lesions of the oral, ocular, or genital mucosa occurring as an extrapulmonary manifestation of *Mycoplasma pneumoniae* infection, often presenting with target-like skin lesions.

### **Mycoplasmataceae**

A family of very small bacteria that lack a cell wall and includes *Mycoplasma* and related genera.

### **Mycoses**

Plural of mycosis; see *Mycosis*.

### **Mycosis**

An infection caused by a fungus, including a yeast or mold. Mycoses may be superficial, cutaneous, subcutaneous, or systemic, depending on the tissues involved; the site and organism determine symptoms and treatment.

### **Mycosis Fungoides**

The most common cutaneous T-cell lymphoma, presenting with progressive patches, plaques, and tumors on the skin with an indolent course. Staging and therapy depend on skin involvement and extracutaneous spread.

#### **Alternate Names:**

MF; Cutaneous T-Cell Lymphoma

### **Mycosis Fungoides Lymphoma**

Mycosis fungoides is a type of cutaneous T-cell lymphoma that usually begins with persistent patches or plaques on the skin.



---

### **Mycotic Aneurysm**

An infected arterial wall or aneurysm, usually caused by bacteria rather than fungi. It may develop from bloodstream infection or an infected embolus and can rupture, causing life-threatening hemorrhage.

### **Mycotoxin**

A poisonous substance made by some fungi that can contaminate food, crops, or indoor environments. Exposure may damage the liver, kidneys, nervous system, immune system, or other organs depending on the toxin and dose.

### **Mydriasis**

Dilation of the pupil, produced mainly by sympathetic activity through the dilator pupillae muscle or by reduced parasympathetic activity. Causes include darkness, medicines, toxins, neurologic disease, and ocular injury.

### **Mydriatic Agents**

A medicine that widens the pupil. Some, such as tropicamide or atropine, also temporarily prevent focusing; others, such as phenylephrine, mainly widen the pupil. These medicines are used to examine the inside of the eye and sometimes to ease pain and prevent scarring during eye inflammation. They can trigger a dangerous rise in eye pressure in people with narrow drainage angles.

### **Myelin**

Myelin is an insulating sheath around nerve fibers that speeds electrical conduction; it is produced by oligodendrocytes in the CNS and Schwann cells in the PNS.

### **Myelination**

Myelination is formation of a myelin sheath around an axon by oligodendrocytes in the central nervous system or Schwann cells in the peripheral nervous system. It increases the speed and efficiency of nerve conduction.

### **Myelin Sheath**

The insulating covering around many nerve fibers

that allows electrical signals to travel quickly. It is made by different support cells in the brain, spinal cord, and peripheral nerves and can be damaged in disorders such as multiple sclerosis.

### **Myelitis**

Myelitis is inflammation of the spinal cord or, less commonly, other neural tissue described by the term. Transverse myelitis can cause weakness, sensory changes, and bladder or bowel dysfunction and may result from infection, immune disease, reduced blood flow, or pressure on the spinal cord.

### **Myeloablative Agonist**

A nonstandard expression that most likely means *myeloablative agent*, a drug or treatment that severely or completely destroys blood-forming cells in the bone marrow. Myeloablative therapy may be used before a stem-cell transplant and causes profound, temporary low blood counts; “agonist” is not the usual term in this context.

### **Myeloablative Chemotherapy**

Myeloablative chemotherapy destroys or severely suppresses bone-marrow blood-cell production and is often used before a stem-cell transplant.

### **Myeloblast**

An immature precursor of granulocytes in the bone marrow; excess myeloblasts are characteristic of acute myeloid leukemia.

### **Myelocyte**

A developing granulocyte precursor in bone marrow that is more mature than a myeloblast and normally absent or rare in peripheral blood.

### **Myelocytic Leukaemia**

A blood cancer arising from cells that normally develop into several blood-cell types. It includes acute and chronic myeloid forms, which can cause anemia, infections, bleeding, fatigue, or an enlarged spleen.

---

## Myelodysplastic Neoplasms

A group of clonal bone-marrow disorders in which blood cells develop abnormally and are made ineffectively. They can cause low red-cell, white-cell, or platelet counts, leading to anemia, infection, or bleeding; some types can progress to acute myeloid leukemia. They are also widely known as myelodysplastic syndromes (MDS).

## Myelodysplastic Or Myeloproliferative Neoplasms

Myelodysplastic and myeloproliferative neoplasms are clonal bone-marrow disorders causing abnormal blood-cell production. They may produce anemia, infection, bleeding, high counts, enlarged spleen, thrombosis, or progression to acute leukemia; classification and mutation testing guide care.

## Myelodysplastic Syndrome

See *Myelodysplastic Neoplasms*.

## Myelodysplastic Syndromes

See *Myelodysplastic Neoplasms*.

## Myeloencephalitis

Myeloencephalitis is inflammation involving the spinal cord and brain, caused by infection, immune disease, toxins, or other neurologic processes.

## Myelofibrosis

A group of bone-marrow disorders in which fibrous tissue disrupts normal blood-cell production. It includes primary myelofibrosis and secondary forms caused by other blood or marrow diseases.

## Myelogenous

Relating to blood-forming cells that develop from the bone marrow's myeloid cell line.

### Alternate Names:

Myeloid

## Myelogram

An imaging study of the spinal canal performed after contrast material is introduced around the spinal cord.

## Myelography

Contrast injected into subarachnoid space for spinal imaging. Used when MRI contraindicated. CT myelography more common now.

## Myeloid

Relating to bone marrow or to the non-lymphoid blood-cell lineages derived from hematopoietic stem cells.

## Myeloid Cell

A blood-forming cell from the myeloid lineage, which produces red cells, platelets, and several white blood-cell types.

## Myeloid Leukemia

Myeloid leukemia is a blood cancer arising from abnormal myeloid precursor cells in the bone marrow and blood.

## Myeloid Sarcoma

Myeloid sarcoma is a mass of immature myeloid cells occurring outside the bone marrow, often associated with acute myeloid leukemia.

## Myelolipoma

A usually benign tumor composed of mature adipose tissue and blood-forming tissue, most often arising in the adrenal gland and often found incidentally.

## Myeloma

A cancer of plasma cells, usually called multiple myeloma when it affects many bone-marrow sites. It can cause bone pain, anemia, kidney damage, high blood calcium, infections, or abnormal proteins in blood and urine.

## Myeloma Cast Nephropathy

Kidney damage caused by abnormal proteins from multiple myeloma blocking and injuring the kidney's filtering tubes. It can cause sudden kidney failure and is treated by addressing the myeloma and supporting kidney function.

---

## Myelomalacia

Softening of the spinal cord, usually due to ischemia, trauma, hemorrhage, or other cord injury.

## Myelomeningocele

A birth defect in which the spinal cord and its protective coverings bulge through an opening in the spine. It can cause weakness or paralysis, loss of sensation, bladder or bowel problems, and fluid buildup around the brain.

## Myelomonocyte

A bone-marrow precursor that can develop toward either the monocyte or granulocyte lineages. An immature bone marrow cell that can develop into either monocytes or certain infection-fighting white blood cells.

## Myelopathy

Spinal-cord dysfunction caused by compression, ischemia, inflammation, infection, trauma, degeneration, or tumor. It may produce bilateral weakness or upper-motor-neuron signs, a sensory level, gait disturbance, pain, and bowel or bladder dysfunction; MRI and urgent evaluation are important when compression or rapid progression is suspected.

## Myeloperoxidase

An enzyme in neutrophil and monocyte granules that uses hydrogen peroxide and chloride to generate antimicrobial oxidants; antibodies against it occur in some autoimmune diseases.

## Myelopoiesis

The production and development of myeloid blood cells, including granulocytes, monocytes, and some precursor cells, in the bone marrow.

## Myeloproliferative Disorders

Clonal bone-marrow disorders causing overproduction of one or more blood-cell lineages, including polycythemia vera, essential thrombocythemia, and myelofibrosis.

## Myeloproliferative Neoplasm

A group of clonal blood and bone-marrow disorders in which the marrow makes too many cells of one or more blood-cell types. Examples include polycythemia vera, essential thrombocythemia, and myelofibrosis; the specific disorder determines its effects and risks.

### Alternate Names:

Myeloproliferative Neoplasms; MPN; MPNs

## Myelosuppression

Reduced bone marrow production of red cells, white cells, and platelets, causing anemia, neutropenia, and thrombocytopenia; a common dose-limiting toxicity of chemotherapy and radiation that increases infection and bleeding risk.

## Myelosuppressive Therapy

Myelosuppressive therapy reduces bone-marrow production of blood cells, sometimes intentionally during cancer treatment but with risk of anemia, infection, and bleeding.

## Myenteric Plexus

A network of enteric nerves between the circular and longitudinal muscle layers of the gastrointestinal tract that coordinates intestinal motility.

## Myhre Syndrome

A rare inherited connective-tissue disorder caused by variants in SMAD4 and characterized by short stature, stiff joints, distinctive facial features, hearing loss, and cardiovascular abnormalities.

## Myiasis

Infestation of living tissue by larvae (maggots) of various fly species, which feed on host tissue and can cause pain, secondary infection, and tissue destruction. It may be cutaneous, involving the skin and subcutaneous tissue, or affect other body sites including the nose, ears, and wounds.

## Myoblast

An early muscle cell that develops and joins with others to form skeletal muscle fibers.

---

## Myoblastoma

A rare tumor made of cells resembling developing muscle cells. The term has been used for more than one tumor type, so diagnosis requires modern microscopic and molecular classification.

### Alternate Names:

Granular Cell Tumor; Abrikossoff Tumor

## Myocardial

Relating to the myocardium, the muscular wall of the heart.

## Myocardial Biopsy

Removal of a small sample of heart muscle, usually by catheter, for microscopic or molecular examination of suspected myocarditis, rejection, infiltrative disease, or other cardiomyopathy.

## Myocardial Bridging

Myocardial bridging occurs when a segment of a coronary artery tunnels through heart muscle and may be compressed during contraction.

## Myocardial Contusion

A bruise of the heart muscle caused by blunt chest trauma, which may produce chest pain, arrhythmias, or impaired heart function.

## Myocardial Infarction

Death of part of the heart muscle because its blood supply is suddenly reduced or blocked, usually by a blood clot on an artery plaque. It may cause chest pressure or pain, breathlessness, sweating, nausea, or no clear symptoms. Diagnosis uses symptoms, ECG findings, and heart-damage blood tests. STEMI and NSTEMI describe different ECG patterns and do not simply mean complete versus partial blockage. Emergency treatment aims to restore blood flow and prevent further clots.

### Alternate Names:

Heart Attack; MI; Acute Myocardial Infarction; STEMI; NSTEMI

## Myocardial Ischemia

Reduced blood flow and oxygen supply to heart

muscle, usually from narrowed coronary arteries, potentially causing chest pain or heart damage.

### Alternate Names:

Cardiac Ischemia; Myocardial Hypoxia

## Myocardial Perfusion Imaging

A heart scan that compares blood flow to the heart muscle at rest and during exercise or medicine-induced stress. It can show areas with reduced blood supply or old heart-muscle damage.

## Myocardial Reperfusion

Restoration of blood flow to heart muscle after ischemia, which can limit injury but may also cause reperfusion-related tissue damage.

## Myocardial Rupture

A tear in the heart wall or other heart structure, usually a severe complication of myocardial infarction.

## Myocardial Viability

The ability of heart muscle weakened by poor blood flow to recover if blood supply is restored. Imaging tests can show whether the tissue is still alive and may help guide decisions about restoring blood flow.

## Myocarditis

Inflammation of the heart muscle, often after an infection or because of an immune reaction. It can cause chest pain, breathlessness, abnormal heart rhythms, reduced pumping, or sudden severe illness; severity ranges from mild inflammation to heart failure or shock.

## Myocarditis Diagnosis

Doctors assess suspected heart-muscle inflammation using symptoms, examination, an electrocardiogram, blood tests, heart imaging, and sometimes MRI or biopsy. No single test proves every case, so results are interpreted together and other causes are excluded.

## Myocarditis Pathology

Inflammatory infiltrates with associated cardiomy-

---

ocyte injury or necrosis that is not explained by ischemia. The pattern may be lymphocytic, eosinophilic, giant-cell, granulomatous, or mixed depending on the cause.

### **Myocarditis Treatment**

Treatment depends on the cause and severity. Care may include rest from strenuous exercise, medicines for heart failure or abnormal rhythms, treatment of infection or autoimmune inflammation when appropriate, and hospital monitoring for severe cases.

### **Myocardium**

The thick muscular middle layer of the heart wall that contracts to pump blood throughout the body.

### **Myoclonic Epilepsy**

An epilepsy syndrome in which seizures include brief, shock-like involuntary muscle jerks; the specific syndrome determines associated seizure types, age of onset, and prognosis.

### **Myoclonic Seizure**

A generalized seizure characterized by brief, shock-like, involuntary single or multiple muscle contractions without loss of consciousness. Commonly affects the upper extremities, resulting in sudden dropping or throwing of objects, and is a hallmark feature of juvenile myoclonic epilepsy.

### **Myoclonus**

A sudden, brief, involuntary muscle jerk or group of jerks.

### **Myocyte**

A differentiated muscle cell specialized for contraction. Skeletal myocytes are long, multinucleated fibers; cardiac myocytes are branching cells joined by intercalated discs; and smooth-muscle myocytes are spindle-shaped cells without organized sarcomeres. Myocyte injury or death contributes to myopathy, rhabdomyolysis, and myocardial infarction.

#### **Alternate Names:**

Muscle Cell

### **Myodegeneration**

Degeneration or breakdown of muscle fibers, caused by inherited, inflammatory, ischemic, toxic, or other disorders.

### **Myoepithelial Carcinoma Pathology**

An infiltrative malignant neoplasm composed predominantly of myoepithelial cells, which may be clear, epithelioid, spindle, or plasmacytoid and show variable immunoreactivity for myoepithelial markers. It most often arises in salivary tissue or soft tissue.

### **Myoepithelial Cell**

A contractile cell around certain glands that helps push secretions from breast or salivary tissue.

### **Myoepithelioma**

A myoepithelioma is a tumor composed of myoepithelial cells and can be benign or malignant, commonly arising in salivary or other glands.

### **Myoepithelioma Of Soft Tissue**

A rare growth in soft tissue made of myoepithelial cells, which have features of both gland cells and muscle-like support cells. It may be harmless or cancerous; imaging and tissue examination determine treatment.

### **Myofascial Pain Syndrome**

A regional pain disorder involving sensitive trigger points in muscle and fascia, with pain that may be referred to other areas and often worsens with pressure or movement.

### **Myofibril**

A contractile thread-like structure inside a muscle fiber, composed of repeating sarcomeres containing actin and myosin.

### **Myofibrillar Myopathy**

Myofibrillar myopathy is a group of inherited muscle disorders with abnormal protein aggregates and progressive skeletal-muscle weakness, sometimes involving heart or breathing muscles.

---

## Myofibrils

Contractile structures inside muscle fibers made of repeating sarcomeres. Long contractile structures inside muscle fibers, made of repeating sarcomeres containing actin and myosin. Their shortening produces muscle contraction.

## Myofibroblastoma

A benign mesenchymal tumor composed of myofibroblastic cells in collagenous or myxoid stroma, most often arising in the mammary or inguinal region but potentially occurring at other soft-tissue sites.

## Myofibroblastoma Of The Breast

A rare benign stromal tumor composed of spindle-shaped myofibroblastic cells, usually occurring in older adults. It is typically a well-circumscribed painless mass and is diagnosed by histology with immunohistochemistry.

## Myofibroblasts

Cells that combine features of fibroblasts and smooth muscle cells and help contract wounds and produce scar tissue.

## Myofibromatosis

Myofibromatosis is a benign proliferation of myofibroblastic tumors, often in infants or children and involving skin, muscle, bone, or internal organs.

## Myofilament

A contractile protein filament within a muscle fiber; actin and myosin myofilaments interact through cross-bridges to produce muscle contraction.

## Myofilaments

Thin actin and thick myosin protein strands inside muscle cells. They slide past one another using energy from ATP, shortening the muscle and producing contraction.

### Alternate Names:

Contractile Filaments; Actin and Myosin Filaments

## Myogenin

A muscle-specific transcription factor required for skeletal-muscle differentiation and used as an immunohistochemical marker in diagnosing some soft-tissue tumors.

## Myoglobin

An oxygen-binding protein stored in skeletal and heart muscle. When muscle is injured, myoglobin can enter the blood and urine; large amounts may contribute to kidney damage and are interpreted with other tests.

## Myoglobin Blood Test

A blood test measuring myoglobin released from injured skeletal or cardiac muscle; it rises early after muscle injury but is nonspecific and is not preferred over troponin for myocardial infarction.

## Myoglobinuria

A condition in which muscle protein enters urine, often after severe muscle injury or breakdown.

## Myoglobin Urine Test

A urine test detecting myoglobin released during muscle breakdown, used in evaluating rhabdomyolysis; substantial myoglobinuria can contribute to acute kidney injury.

## Myokymia

Involuntary, fine, continuous rippling contractions of muscle fibers, often seen around the eye but also associated with nerve hyperexcitability or neurologic disease.

## Myoma

A usually benign tumor made of muscle tissue. The term often refers to a leiomyoma, especially a uterine fibroid, but myomas can arise from smooth or skeletal muscle in other sites.

## Myomalacia

Softening and breakdown of muscle tissue, sometimes affecting heart muscle after serious injury or poor blood supply.



---

## Myomas

Plural of myoma; see *Myoma*.

## Myomectomy

Surgical removal of uterine fibroids while leaving the uterus in place. It may improve bleeding, pain, or pressure symptoms and can preserve the possibility of pregnancy, although risks and recurrence remain.

## Myometrium

The thick muscle layer forming most of the uterus. It contracts during menstruation and labor, responds to hormones, and may develop conditions such as fibroids, adenomyosis, or abnormal thickening.

## Myopathic Facies

A characteristic expressionless, mask-like facial appearance marked by drooping eyelids (ptosis), prominent facial muscle wasting, an inability to bury the eyelashes, and a transverse, tented upper lip (tapir mouth). It occurs due to weakness of the facial musculature in conditions such as facioscapulohumeral dystrophy and myotonic dystrophy.

## Myopathy

A disease that directly affects muscles and causes weakness, fatigue, cramps, or pain. Causes include inherited conditions, inflammation, medicines, hormone disorders, infections, toxins, and problems with muscle energy production.

## Myopericytoma

A myopericytoma is a usually benign perivascular tumor composed of cells around small blood vessels, often occurring in skin or soft tissue.

## Myopia

A refractive error in which light focuses in front of the retina, usually because the eyeball is too long or the eye focuses too strongly. Distant objects appear blurred while near objects are usually clearer. It can be corrected with glasses, contact lenses, or selected refractive procedures.

### Alternate Names:

Short-Sightedness; Nearsightedness

## Myosin

A protein that uses energy to pull on actin, another muscle protein, and produce movement or force. It is important for contraction of skeletal, heart, and smooth muscle and also helps movement inside many other cells.

## Myosin ATPase

The activity by which the muscle protein myosin breaks down ATP, the cell's energy source. This releases energy that lets myosin pull on actin and produce muscle contraction.

## Myositis

Inflammation of muscle tissue caused by autoimmune disease, infection, medicines, or injury. It may produce weakness, pain, swelling, or fatigue; evaluation often includes examination, blood tests, imaging, and sometimes muscle testing.

## Myositis Ossificans

A noncancerous condition in which abnormal bone forms inside a muscle or soft tissue, most often following a severe blunt injury, deep bruise, or surgery, commonly in the thigh or upper arm. It causes a firm, painful lump and joint stiffness and may improve as the abnormal bone matures.

### Alternate Names:

Traumatic myositis ossificans; Heterotopic ossification of muscle; Myositis ossificans.

## Myotonia

Delayed relaxation of a muscle after voluntary contraction or stimulation, often caused by an inherited or acquired muscle-channel disorder.

## Myotonia Congenita

An inherited skeletal-muscle channelopathy causing delayed relaxation after voluntary contraction. Stiffness often improves with repeated movement and may be accompanied by muscle hypertrophy; onset and severity vary between genetic forms.

## Myotonia Fluctuans

A rare inherited muscle-channel disorder causing

---

stiffness that varies from day to day or after activity. It is usually mild and does not cause lasting weakness; genetic testing may identify a change in the *SCN4A* gene.

**Alternate Names:**

Potassium-Aggravated Myotonia Fluctuans

**Myotonia Permanens**

A rare inherited muscle-channel disorder causing continuous, severe muscle stiffness, including at rest. It may begin in infancy and affect the face, throat, or breathing muscles; it is usually related to a change in the *SCN4A* gene and requires specialist care.

**Alternate Names:**

Severe Sodium Channel Myotonia; Continuous Myotonia

**Myotonic Disorder**

A neuromuscular disorder characterized by delayed relaxation after muscle contraction, often with weakness, stiffness, or progressive muscle wasting.

**Myotonic Dystrophy**

An inherited disorder causing delayed muscle relaxation, progressive weakness, and wasting. It can also affect the heart, breathing, swallowing, eyes, hormones, or thinking; symptoms and severity vary between types and people.

**Myotonic Dystrophy Type 2**

An inherited multisystem disorder caused by CCTG repeat expansion in CNBP, producing myotonia, proximal or axial weakness, muscle pain, cataracts, and variable endocrine or cardiac features.

**Myringitis**

Inflammation of the eardrum that may cause ear pain, blisters, drainage, or temporary hearing changes.

**Myringotomy**

A small opening made in the eardrum to drain fluid or relieve pressure in the middle ear. A ventilation tube may be inserted to keep the opening open temporarily. Possible problems include infection,

scarring, persistent drainage, or a hole that does not close.

**Myringotomy With Tympanostomy Tube**

Incision of the tympanic membrane with placement of a ventilation tube to equalize middle-ear pressure and permit drainage. It is used selectively for persistent middle-ear effusion, recurrent acute otitis media, or associated hearing and developmental problems.

**Myristica Oil Poisoning**

Poisoning from concentrated nutmeg oil, which can cause nausea, vomiting, agitation, fast heartbeat, dizziness, hallucinations, confusion, seizures, or coma. Suspected ingestion needs prompt poison-control or emergency assessment.

**Myxedema**

Severe hypothyroidism causing accumulation of mucin in tissues, with swelling, slowed body functions, and other symptoms.

**Myxedema Coma**

A life-threatening medical emergency representing the decompensated, end-stage state of severe, long-standing, untreated or undertreated hypothyroidism. Typically precipitated by an acute physiological stressor (such as sepsis, cold exposure, myocardial infarction, stroke, or sedatives), it is clinically defined by a classic triad of profound hypothermia (often core temperature < 35°C), altered mental status (ranging from lethargy and confusion to stupor and coma), and multisystem organ slowing. Characteristic complications include severe sinus bradycardia, hypotension, hypoventilation leading to hypercapnic respiratory acidosis, dilutional hyponatremia, and diffuse non-pitting periorbital and peripheral myxedema.

**Alternate Names:**

Myxoedema Coma; Decompensated Hypothyroidism; Myxedematous Coma

**Myxofibroma**

A myxofibroma is a benign tumor containing fibrous and gelatinous myxoid tissue, often arising in the

---

jaw or other connective tissue.

### **Myxoid Cyst**

A myxoid cyst is a gelatinous fluid-filled lesion near a joint or tendon, commonly on a finger near the distal joint and nail.

### **Myxoma**

A usually noncancerous tumor, most often in the upper left chamber of the heart. It may obstruct blood flow or release clots, causing breathlessness, fainting, fever, or stroke-like symptoms; heart ultrasound is commonly used for diagnosis.

### **Myxopapillary Ependymoma**

Myxopapillary ependymoma is a usually slow-growing spinal tumor arising from ependymal cells, commonly in the conus or filum terminale.

### **Myxosarcoma**

A rare cancer of soft tissue containing abundant jelly-like material. It can grow into nearby tissue or spread to distant organs; diagnosis requires examination of a tissue sample.

### **MZL**

Abbreviation; see *Marginal Zone Lymphoma*.

### **Male Sexual Problems**

Sexual symptoms in men, including erectile dysfunction, reduced desire, ejaculation disorders, infertility, or penile curvature; causes may be vascular, hormonal, neurologic, medication-related, psychological, or mixed.

### **Malignant Appendical Tumor**

Malignant tumor arising in the appendix; see *Appendiceal Cancer*.

### **Malignant Carcinoid Syndrome**

Severe carcinoid syndrome caused by hormone-releasing neuroendocrine cancer, potentially producing flushing, diarrhea, bronchospasm, right-sided heart-valve disease, and hemodynamic complications; see *Carcinoid Syndrome*.

### **Malignant Large Bowel Tumor**

Cancer arising in the colon or rectum; see *Colorectal Cancer*.

### **Mallet Toes**

Alternate plural form; see *Mallet Toe*.

### **Malocclusion and Orthodontics**

Misalignment of the teeth or jaws and the dental treatment used to improve bite, spacing, function, or facial balance; see *Malocclusion*.

### **Mammary Paget's Disease**

A rare breast cancer involving the nipple and areola, often associated with underlying ductal carcinoma; see *Paget Disease Of The Nipple*.

### **Mandibular Ameloblastoma**

An ameloblastoma arising in the lower jaw, usually a slow-growing locally invasive odontogenic tumor; see *Ameloblastoma*.

### **Manic Depression in Children and Teens**

Older name for bipolar disorder beginning in childhood or adolescence; episodes of mania, hypomania, or depression require specialist assessment.

### **Manic-Depressive Disorder**

Older name for bipolar disorder, characterized by episodes of mania or hypomania and episodes of depression; see *Bipolar Disorder*.

### **Metabolic Dysfunction-Associated Steatohepatitis (MASH)**

Alternate name; see *Metabolic Dysfunction-Associated Steatohepatitis*.

### **Metastatic Melanoma**

Melanoma that has spread from the skin or another primary site to lymph nodes or distant organs; treatment depends on molecular findings, extent, symptoms, and overall health.

### **Micropolygyria with Muscular Dystrophy**

A rare congenital syndrome combining abnormal

---

folding of the cerebral cortex with muscular dystrophy, causing developmental, seizure, movement, and muscle weakness findings.

### **Migraine Headaches**

Alternate name; see *Migraine*.

### **Milk Allergy**

An immune reaction to proteins in cow's milk, often beginning in infancy. It may cause hives, vomiting, wheezing, eczema, diarrhea, or anaphylaxis; it differs from lactose intolerance, which is difficulty digesting milk sugar.

#### **Alternate Names:**

Cow's Milk Allergy; Cow's Milk Protein Allergy; CMPA

### **Milk Intolerance**

Difficulty tolerating milk, often because of lactose malabsorption, causing bloating, cramps, gas, or diarrhea; it is not the same as milk-protein allergy.

### **Mini-Stroke (TIA)**

Alternate name for a transient ischemic attack, a temporary episode of focal neurologic dysfunction caused by reduced brain, spinal-cord, or retinal blood flow without lasting infarction; urgent evaluation is required because stroke risk is increased.

### **Missed Period**

Absence of an expected menstrual period. Pregnancy is common, but stress, weight change, endocrine disease, medications, perimenopause, and other conditions can also contribute.

### **Mitchell Disease**

Alternate name for erythromelalgia, a disorder causing attacks of burning pain, warmth, and redness in the extremities; see *Erythromelalgia*.

### **Mixed Connective Tissue Disease (MCTD)**

Alternate name; see *Mixed Connective Tissue Disease*.

### **Mixed Hepatic Porphyrria**

An acute hepatic porphyria in which inherited impairment of heme synthesis can cause episodic abdominal and neurologic symptoms and, in some people, photosensitive skin disease.

### **Monomodal Visual Amnesia**

A rare visual-memory disorder in which a person has marked difficulty recognizing or recalling visual information while other memory modalities may be relatively preserved; neurologic evaluation is needed to identify the cause.

### **Morgagni-Turner-Albright Syndrome**

Alternate name for Turner syndrome, a chromosomal condition involving complete or partial loss of one X chromosome and characteristic growth, ovarian, heart, kidney, and hearing findings; see *Turner Syndrome*.

### **Morvan Disease – Syringomyelia**

Historical or imprecise index label combining Morvan syndrome and syringomyelia, which are distinct neurologic disorders; see *Morvan's Disease* and *Syringomyelia*.

### **Mountain Fever**

A feverish illness associated with high-altitude exposure or, in some contexts, a tick-borne infection; the exposure history and clinical findings determine the specific diagnosis.

### **Mouth (Oral) Cancer**

Alternate name; see *Mouth Cancer*.

### **MPS**

Abbreviation for mucopolysaccharidosis, a group of inherited lysosomal storage disorders; see *Mucopolysaccharidosis*.

### **MPS Disorder**

Alternate name; see *Mucopolysaccharidosis*.

### **MPS Disorder I**

Alternate name; see *Mucopolysaccharidosis Type I*.

---

## **MPS Disorder II**

Alternate name; see *Mucopolysaccharidosis Type II*.

## **MPS I**

Abbreviation; see *Mucopolysaccharidosis Type I*.

## **MPS II**

Abbreviation; see *Mucopolysaccharidosis Type II*.

## **MS (Multiple Sclerosis)**

Alternate name; see *Multiple Sclerosis*.

## **MSA**

Abbreviation for multiple system atrophy, a progressive neurodegenerative disorder affecting movement, autonomic function, and coordination; see *Multiple System Atrophy*.

## **MSL**

Abbreviation for multiple symmetric lipomatosis, a disorder with symmetric deposits of fatty tissue around the neck, shoulders, or trunk; see *Multiple Symmetric Lipomatosis*.

## **Mud Fever**

A historical name used for certain febrile illnesses associated with muddy or wet environments; the exact disease depends on the clinical and exposure context.

## **Multiple Endocrine Deficiency Syndrome**

A syndrome involving deficiency of more than one endocrine hormone, caused by autoimmune, genetic, pituitary, hypothalamic, or other disease; evaluation identifies which glands are affected.

## **Multiple Lentigines Syndrome**

An inherited disorder causing numerous lentigines with variable heart, endocrine, eye, hearing, or developmental abnormalities; it may overlap with Noonan-spectrum syndromes.

## **Multiple Peripheral Neuritis**

Inflammation or injury affecting multiple peripheral nerves, causing pain, numbness, weakness, or

autonomic symptoms; see *Peripheral Neuritis*.

## **Multiple Polyposis of the Colon**

Development of many colorectal polyps, which may be inherited or acquired and may increase colorectal cancer risk; see *Polyposis*.

## **Multiple Pregnancy: Twins or More**

Alternate name; see *Multiple Pregnancy*.

## **Multiple Sclerosis (MS)**

Alternate name; see *Multiple Sclerosis*.

## **Multiple Sulfatase Deficiency**

A rare inherited lysosomal-storage disorder caused by deficient activation of multiple sulfatase enzymes, producing progressive neurologic, skeletal, skin, and organ disease.

## **Multiple Symmetric Lipomatosis**

A disorder characterized by symmetric, usually nonencapsulated fatty deposits around the neck, shoulders, upper trunk, or proximal limbs; see *Multiple Symmetric Lipomatosis*.

## **Munchausen Syndrome by Proxy**

Older name for factitious disorder imposed on another, in which a caregiver falsifies, exaggerates, or induces illness in another person; see *Factitious Disorder Imposed on Another*.

## **Murk Jansen Metaphyseal Chondrodysplasia**

Alternate name for Jansen metaphyseal chondrodysplasia; see *Jansen Metaphyseal Chondrodysplasia*.

## **Muscular Dystrophies**

Alternate plural form; see *Muscular Dystrophy*.

## **Myelodysplastic Syndrome (MDS)**

Alternate name; see *Myelodysplastic Syndrome*.

## **Manual Therapy or Bodywork**

Hands-on treatment involving mobilization, manipulation, massage, or soft-tissue techniques to address

---

pain, stiffness, or movement limitations; risks and suitability depend on the condition and technique.

### **Melanoma/Skin Cancer**

Alternate name; see *Melanoma*.

### **Mono (Infectious Mononucleosis)**

Alternate name; see *Infectious Mononucleosis*.



# N

## **Nabothian Cyst**

A small, harmless mucus-filled lump on the cervix caused by a blocked cervical gland. It is common and usually causes no symptoms or need for treatment; an unusual or bleeding lesion still needs examination.

## **Nabumetone**

A nonsteroidal anti-inflammatory medicine used for pain and inflammation in selected joint or muscle conditions. It can irritate the stomach, affect kidneys, raise cardiovascular risk, or cause allergic reactions, especially with prolonged use.

## **N-Acetylcysteine**

An antidote used mainly for acetaminophen overdose. It helps protect the liver and may still help when treatment starts late or liver injury has already begun.

## **N-Acetylglutamate Synthetase Deficiency**

An inherited urea-cycle disorder in which deficient N-acetylglutamate synthetase fails to activate carbamoyl phosphate synthetase I. Ammonia can accumulate and cause vomiting, sleepiness, confusion, seizures, coma, or developmental injury.

## **Nadir**

The lowest level reached by a measurement or symptom. After chemotherapy, it often means the period when blood-cell counts are lowest and the risk of infection or bleeding is greatest.

## **Naegle Rule**

A method for estimating a baby's due date by adding

seven days and about nine months to the first day of the last menstrual period. It is less reliable with irregular cycles or uncertain dates.

## **Naegeli-Franceschetti-Jadassohn Syndrome**

A rare inherited ectodermal disorder causing reticulate skin pigmentation, reduced sweating, abnormal teeth or nails, and sometimes loss of fingerprints. Heat intolerance and dental problems may be important; genetic and multidisciplinary care is useful.

## **Naegleria Fowleri**

A free-living microscopic amoeba found mainly in warm freshwater lakes, rivers, and hot springs. If water containing it enters the nose, the amoeba can travel to the brain and cause primary amebic meningoencephalitis (PAM), a rare, rapidly worsening brain infection that is nearly always fatal. Early symptoms may include headache, fever, nausea, or vomiting, followed by stiff neck, confusion, seizures, or coma. Infection does not occur from swallowing the water and does not spread from person to person.

## **Naegleria Infection**

A rare infection caused when warm freshwater containing *Naegleria fowleri* enters the nose and the amoeba reaches the brain. It causes primary amebic meningoencephalitis (PAM), which rapidly inflames and destroys brain tissue. Symptoms may include severe headache, fever, nausea, vomiting, stiff neck, confusion, seizures, or coma. It is not acquired by swallowing contaminated water and does not spread between people.

---

## Naevus

The British spelling of nevus, a usually benign localized skin or mucosal lesion such as a mole.

## Naevus Flammeus

A congenital vascular capillary malformation (port-wine stain) presenting as a flat, sharply demarcated, pink-to-violaceous cutaneous patch that does not regress spontaneously, classically involving the ophthalmic (V1) distribution of the trigeminal nerve in Sturge-Weber syndrome.

### Alternate Names:

Port-Wine Stain; Capillary Vascular Malformation

## Naffziger Sign

A clinical physical sign for spinal nerve root compression or spinal space-occupying lesions; manual bilateral compression of the internal jugular veins causes transient elevation of intraspinal venous and CSF pressure, reproducing or worsening radiating radicular pain in the affected dermatome.

### Alternate Names:

Naffziger's Sign; Jugular Compression Radiculopathy Test

## Nail

A hard protective plate on the end of each finger and toe. Nails protect the tips, help with fine tasks, and may show changes caused by injury, infection, skin disease, or illness elsewhere.

## Nail Abnormalities

Changes in nail color, shape, thickness, texture, or attachment caused by trauma, infection, inflammatory disease, systemic illness, medication, or congenital conditions.

## Nail Bed Injury

Trauma to the tissue beneath a nail that may cause bleeding under the nail, laceration, nail separation, fracture of the fingertip, or permanent nail deformity. Assessment is important when the nail is badly disrupted or the fingertip is crushed.

## Nail Changes

Alterations in nail color, shape, texture, or attachment that may result from local injury, skin disease, infection, medicines, nutritional deficiency, or systemic illness. Their significance depends on the pattern and other clinical findings.

## Nail Clipping For Fungal Testing

Laboratory examination of clipped nail material by microscopy, culture, or another validated test for fungal infection. Testing can help distinguish nail fungus from injury, psoriasis, or other nail disorders, especially before oral antifungal treatment.

## Nail Clubbing

Enlargement and rounding of the ends of the fingers or toes with more curved nails. It may run in families or signal disease of the lungs, heart, digestive system, or another organ, especially when it develops recently.

## Nail-Fold Capillaroscopy

A test that uses magnification to examine the tiny blood vessels near the nails. It can help distinguish primary Raynaud phenomenon from Raynaud phenomenon linked to conditions such as systemic sclerosis, dermatomyositis, or mixed connective-tissue disease.

### Alternate Names:

Periungual Capillaroscopy; NFC Examination

## Nail Fungus

A fungal infection of a fingernail or toenail causing thickening, discoloration, brittleness, distortion, or separation from the nail bed.

### Alternate Names:

Toenail Fungus

## Nail Furrows

Nail furrows are grooves or depressions running across or along a nail; transverse Beau lines can follow systemic illness or trauma, while longitudinal ridges are often age-related but may reflect local or systemic disease.

---

## Nail Infection

Infection of the nail or surrounding tissue, commonly caused by fungi or bacteria. It may cause thickening, discoloration, separation of the nail, pain, swelling, or drainage; diabetes and poor circulation increase the risk of complications.

## Nail Injuries

Trauma to the nail plate, nail bed, nail folds, or fingertip that may cause bruising, laceration, avulsion, fracture, or later nail-growth abnormalities.

## Nail Patella Syndrome

Usually an autosomal-dominant disorder caused by a pathogenic variant in *LMX1B*, characterized by abnormal or absent nails, small or absent kneecaps, and elbow or pelvic abnormalities such as iliac horns. Glaucoma and kidney disease with proteinuria may also occur.

## Nail-Patella Syndrome

A rare inherited condition affecting the nails, kneecaps, elbows, and pelvis. Some people also develop kidney disease or glaucoma. The kneecaps may be small, absent, or displaced, and lifelong monitoring may be needed.

## Nail Pits

Small dents in the surface of a nail. They are common in psoriasis but may also occur with hair-loss disorders, eczema, lichen planus, or other conditions affecting nail growth.

## Nail Pitting

Punctate, cupuliform depressions in the dorsal nail plate resulting from parakeratotic defective keratinization within the proximal nail matrix, characteristically seen in psoriasis, alopecia areata, and reactive arthritis.

## Nail Psoriasis

Psoriasis affecting the nails can cause small pits, separation from the nail bed, yellow or red-brown patches, thickening, crumbling, or ridges. It may occur with skin psoriasis or psoriatic arthritis and can be severe or newly developed.

## Nails

Hard plates of keratin covering the ends of fingers and toes. Changes in color, shape, thickness, or attachment can result from injury, infection, skin disease, nutritional problems, or other illness.

## Nail Surgery

A procedure on the nail plate, nail bed, nail fold, or matrix to treat ingrown nails, tumors, infection, trauma, or painful deformity. Risks include bleeding, infection, scarring, altered nail growth, and recurrence.

## Nalbuphine

An opioid pain medicine with mixed effects, used for moderate to severe pain under medical supervision.

## Naloxone

A medicine that blocks opioids and can quickly restore breathing during an opioid overdose. It comes as a nasal spray or injection. Emergency help is still needed because its effect may wear off before the opioid does.

## Naltrexone

A longer-acting medicine that blocks opioid effects and reduces alcohol-related cravings in some people. Opioid use during treatment can trigger withdrawal.

## Nanocolloid

Very small particles carrying a radioactive tracer. After injection, they can show how lymph drains and help locate the first lymph node receiving drainage from a tumor.

## Nanomedicine

The use of very small materials or devices in medicine to help diagnose, prevent, monitor, or treat disease. Examples include nanoparticle drug carriers, imaging agents, biosensors, and some vaccine or cancer treatments; their effects and safety depend on the material, size, dose, and how the body handles them.

---

## Nanoparticle

A particle with dimensions typically measured in nanometres. In medicine, nanoparticles may carry drugs, improve imaging, or support diagnostic tests; their safety depends on composition, size, dose, route, and how the body processes them.

## Nanoscience

The study of materials and structures at approximately the nanometre scale, where surface and physical properties may differ from those of larger materials. It supports research in drug delivery, imaging, diagnostics, and biomaterials.

## Nanotechnology

The design and use of materials, devices, or systems at the nanometre scale. Medical applications include targeted drug delivery, imaging agents, biosensors, tissue engineering, and diagnostic tests; safety depends on the specific material and exposure.

## Napelline

A poisonous plant chemical related to aconite alkaloids. Exposure can disturb nerve signals and heart rhythm, causing tingling, weakness, vomiting, dangerous blood-pressure changes, or life-threatening cardiac poisoning.

## Naphazoline

A topical alpha-adrenergic agonist used as a nasal decongestant or ocular vasoconstrictor; prolonged use can worsen congestion.

## Naphthalene Poisoning

Harm from mothballs or other products containing naphthalene. It can destroy red blood cells, causing weakness, jaundice, dark urine, kidney injury, seizures, or coma.

## Naproxen

A nonsteroidal anti-inflammatory drug used for pain, fever, and inflammation.

## Naproxen Sodium Overdose

Taking too much naproxen can cause stomach

pain, vomiting, bleeding, drowsiness, kidney injury, seizures, or breathing problems. Large or intentional overdoses can be life-threatening.

## Narcissism

A pattern of excessive self-focus, need for admiration, or sense of superiority. Mild traits are common, while persistent traits that impair relationships or functioning may occur in narcissistic personality disorder.

## Narcissistic Personality Disorder

A long-lasting pattern of an unusually strong need for admiration, a sense of special importance, and difficulty recognizing other people's feelings. It causes problems in relationships or daily life and is diagnosed by a trained mental-health professional after a full assessment.

### Alternate Names:

NPD; Megalomania; Pathological Narcissism

## Narcolepsy

A chronic disorder that affects the brain's control of sleep and wakefulness. It causes overwhelming daytime sleepiness and sudden, irresistible sleep episodes, even after enough nighttime sleep; nighttime sleep may also be broken. The boundary between sleep and wakefulness is less clear, so some features of rapid-eye-movement (REM) sleep may occur while awake or when falling asleep or waking. Symptoms may include cataplexy, a brief loss of muscle control triggered by strong emotions; sleep paralysis; and vivid dream-like experiences. Type 1 narcolepsy includes cataplexy and is often associated with low hypocretin (orexin), a brain chemical that helps maintain wakefulness; type 2 usually does not include cataplexy.

## NARP Syndrome

A rare inherited mitochondrial disorder that can cause nerve damage, muscle weakness, loss of coordination, seizures, and vision problems. Symptoms and severity vary widely.

### Alternate Names:

NARP; Neuropathy, Ataxia, and Retinitis Pigmentosa

---

### Narrow Angle Glaucoma

A condition in which the drainage area inside the eye is unusually narrow, raising the risk of a sudden rise in eye pressure. Severe eye pain, blurred vision, halos, nausea, or vomiting may occur during an acute attack.

### Narrowband UVB Phototherapy

A treatment that uses carefully measured ultraviolet light for conditions such as psoriasis, vitiligo, and eczema. Treatment requires eye protection and monitoring of the skin.

### Narrow-Complex Tachycardia

A fast heart rhythm that usually begins in the upper chambers of the heart. It produces narrow-looking beats on an ECG and may cause palpitations, dizziness, chest discomfort, breathlessness, or fainting.

### Narrow Pulse Pressure

A pulse pressure (systolic minus diastolic blood pressure) of less than 30 mmHg or less than twenty-five percent of systolic pressure. It reflects reduced stroke volume or marked peripheral vasoconstriction, commonly seen in severe aortic stenosis, cardiac tamponade, hypovolaemic shock, and advanced heart failure.

### Narrow Therapeutic Index

A medicine for which the helpful dose is close to the harmful dose. It requires careful dosing and sometimes blood tests to avoid toxicity.

### Nasal

Relating to the nose or the space inside it. The nose warms, moistens, and filters incoming air and helps with smell and speech. Nasal disease may cause blockage, bleeding, discharge, reduced smell, or difficulty breathing.

### Nasal Bleeding

Alternate name; see *Epistaxis*.

### Nasal Cannula

A soft tube with two small prongs that rest inside

the nostrils to deliver extra oxygen. It is used when a person needs mild to moderate breathing support, and the oxygen flow is adjusted while breathing is monitored.

### Nasal Cavity

The paired space inside the nose, extending from the nostrils to the nasopharynx. It warms, humidifies, and filters inspired air and contains the receptors responsible for smell.

### Nasal Congestion

A blocked or stuffy feeling in the nose caused by swelling of the nasal lining, extra mucus, or a structural problem. It occurs with colds, allergies, sinus disease, irritants, and other conditions.

### Nasal Congestion And Rhinorrhea

Nasal obstruction or stuffiness together with a runny nose, commonly caused by viral infection, allergic or nonallergic rhinitis, sinus disease, irritants, or structural blockage. Persistent unilateral symptoms, recurrent bleeding, fever, facial swelling, or breathing difficulty requires evaluation.

### Nasal Discharge

Alternate name; see *Rhinorrhea*.

### Nasal Corticosteroid Sprays

Nasal corticosteroid sprays reduce mucosal inflammation and congestion in allergic rhinitis, nasal polyps, and some chronic sinus disorders; correct direction away from the septum limits irritation and nosebleeds.

### Nasal Disorders

Conditions affecting the nose or nasal passages, including infection, allergy, trauma, obstruction, polyps, and tumors.

### Nasal Endoscopy

An examination in which a thin camera is used to look inside the nose and upper throat. It can help find swelling, polyps, a foreign object, bleeding, or a growth.

---

## Nasal Flaring

Widening and outward movement of the nostrils during breathing. It is a visible sign of increased work of breathing, particularly in infants and children, and may accompany airway obstruction, bronchiolitis, pneumonia, asthma, or severe systemic illness.

## Nasal Foreign Bodies

Objects lodged in a nasal passage, most often in children. Unilateral obstruction, foul-smelling discharge, bleeding, pain, or difficulty breathing may occur; button batteries and paired magnets require urgent removal.

## Nasal Fracture

A break in one or more bones or cartilaginous structures of the nose, usually caused by facial trauma. Pain, swelling, bruising, deformity, bleeding, and nasal obstruction may occur; a septal hematoma requires prompt treatment.

## Nasal Implant

A medical device placed inside or near the nose to support weakened tissue, change nasal shape, or improve airflow. The implant may be temporary or permanent, and risks include infection, movement, scarring, or breathing problems.

## Nasal Mucosal Biopsy

Removal of a small piece of tissue from the nose for laboratory examination. It may help investigate persistent inflammation, infection, unusual deposits, or a growth; soreness, bleeding, and infection are possible complications.

## Nasal Obstruction

Blocked airflow through one or both nasal passages. Causes include congestion, allergy, infection, a deviated septum, polyps, trauma, or a mass. Persistent one-sided obstruction or bleeding requires medical evaluation.

## Nasal Passage

One of the air channels inside the nose. Its lining warms, moistens, and filters inhaled air before it reaches the throat; swelling, mucus, polyps, or a

structural blockage can make breathing difficult.

## Nasal Polyps

Soft, usually noncancerous growths in the lining of the nose or sinuses. They can cause a blocked nose, runny nose, and reduced sense of smell.

### Alternate Names:

Sinonasal Polyps; Nasal Polyposis

## Nasal Septal Hematoma

A collection of blood inside the wall dividing the nostrils, usually after a nose injury. It can block breathing and damage the nose if it is not drained promptly.

## Nasal Septum

The wall that divides the nasal cavity into right and left sides. It consists mainly of cartilage in front and bone behind and helps support the nose and direct airflow.

## Nasal Septum Perforation

A hole through the cartilage or bone separating the nasal passages, which may cause crusting, nosebleeds, whistling, or obstruction and can result from trauma, surgery, intranasal drugs, or inflammatory disease.

## Nasal Spray

A liquid or mist delivered into the nose. Different sprays contain saline, corticosteroids, antihistamines, or decongestants; correct use and duration depend on the product.

## Nasal Stent

A small device placed inside the nose to keep an airway open or support healing after surgery. It may cause crusting, blockage, infection, bleeding, or pressure injury.

## Nasal Vestibulitis

Inflammation or infection of the nasal vestibule, commonly caused by *Staphylococcus* bacteria and producing tenderness, crusting, redness, or small pustules near the nostrils.



---

### Nasogastric Feeding Tube

A flexible tube passed through the nose into the stomach to deliver enteral nutrition or medicines, or to decompress or aspirate the stomach.

### Nasogastric Tube

A tube passed through the nose into the stomach to remove contents or deliver nutrition, fluids, or medicines when clinically indicated.

### Nasojejunal Feeding Tube

An enteral feeding catheter passed through the nose, pharynx, stomach, and pylorus into the proximal jejunum under endoscopic or fluoroscopic guidance, indicated for patients requiring enteral nutrition who cannot tolerate gastric feeding due to severe gastroparesis, recurrent aspiration, or acute pancreatitis.

#### Alternate Names:

NJ Tube; Post-Pyloric Feeding Tube

### Nasolacrimal Duct

A membranous canal enclosed within a bony canal that conveys tears from the lacrimal sac downward into the inferior meatus of the nasal cavity beneath the inferior nasal concha, guarded by the mucosal valve of Hasner.

#### Alternate Names:

Tear Duct

### Nasopharyngeal Airway

A soft tube inserted through the nose to maintain a passage between the nares and pharynx. It may be useful in a spontaneously breathing patient with partial upper-airway obstruction but is relatively contraindicated in suspected basal skull fracture or significant nasal trauma.

### Nasopharyngeal Cancer

Cancer arising in the upper throat behind the nose, most often a carcinoma associated with Epstein-Barr virus in endemic regions. Nasal obstruction, nosebleeds, hearing loss, neck nodes, or cranial-nerve symptoms may occur; endoscopy, biopsy, and imaging establish extent.

### Nasopharyngeal Carcinoma

A cancer arising in the upper throat behind the nose. It may cause a blocked or bleeding nose, hearing problems, a neck lump, headache, or swallowing trouble; risk varies with genetics, infections, and geographic background.

#### Alternate Names:

NPC; Nasopharyngeal Cancer

### Nasopharyngeal Culture

A laboratory test in which a sample from behind the nose is placed in a growth medium to look for selected bacteria or fungi. A negative result does not exclude every infection, and other tests may be more sensitive.

### Nasopharynx

The upper part of the throat behind the nose and above the soft palate. It carries air, contains the openings of the tubes that connect to the middle ears, and includes nearby lymph tissue that helps defend against infection.

### Natal Cleft

The groove between the buttocks, also called the intergluteal or sacral cleft.

### Natal Teeth

Teeth present at birth, most often involving the lower central incisors. They may be loose, interfere with feeding, or injure the infant's tongue and require dental assessment.

### Natriuresis

Removal of sodium from the body in urine. Water usually leaves with the sodium, which can reduce fluid volume and blood pressure; the process may be increased by natural hormones or water-removing medicines.

### Natriuretic

Causing the kidneys to remove more sodium in urine. Water usually leaves with the sodium, which can lower blood volume and pressure. The term describes certain body hormones and some medicines;

---

related blood tests can help assess heart failure.

### **Natural Immunity**

Protection that develops after a person's immune system responds to an infection, or temporary protection passed to a newborn through the placenta or breast milk. It may be incomplete or fade and is not always safer than vaccination.

### **Natural Killer Cell**

A type of white blood cell that can recognize and destroy some virus-infected or abnormal cells without previous exposure to that specific target. It helps provide early immune defense and releases signals that guide other immune cells.

#### **Alternate Names:**

NK Cells; Large Granular Lymphocytes

### **Natural Killer Cells**

Alternate plural form; see *Natural Killer Cell*.

### **Nausea**

An unpleasant feeling that you may vomit. It can occur with stomach or intestinal illness, pregnancy, medicines, migraine, inner-ear problems, metabolic disorders, or serious abdominal or neurologic disease. Persistent vomiting, severe pain, dehydration, blood, or confusion may signal a serious condition.

### **Nausea And Vomiting**

Nausea is the urge to vomit; vomiting is forceful emptying of stomach contents. Causes include infection, medicines, pregnancy, migraine, obstruction, and other illnesses.

### **Nausea And Vomiting – Adults**

Nausea and vomiting in adults can result from infection, stomach or bowel disease, medicines, pregnancy, migraine, metabolic illness, neurologic disease, or blockage. Severe pain, blood, confusion, dehydration, chest symptoms, or persistent symptoms may indicate a serious condition.

### **Nausea And Vomiting During Early**

### **Pregnancy**

Nausea and vomiting are common in early pregnancy, but persistent symptoms with inability to keep fluids down, weight loss, dehydration, abdominal pain, fever, blood, or altered mental status may indicate hyperemesis gravidarum or another condition and require medical evaluation.

### **Navel**

The navel, or umbilicus, is the scar left after the umbilical cord separates and marks the former connection between fetus and placenta.

### **Navicular**

A boat-shaped bone in the middle of the foot that helps support the arch. Injury to it can cause foot pain and may heal slowly because part of the bone has a limited blood supply.

### **Navicular Fracture**

A break in the navicular bone of the foot, caused by acute trauma or repetitive loading. Midfoot pain, swelling, and tenderness over the navicular may occur, and some fractures are difficult to detect without advanced imaging.

### **NCS**

Usually an abbreviation for nerve-conduction study, a test that measures how quickly and strongly electrical signals travel through nerves. It can help assess nerve damage, pressure on a nerve, or some causes of numbness and weakness.

### **NDM-1**

An enzyme made by some bacteria that makes them resistant to many important antibiotics. Infections caused by these bacteria can be difficult to treat and require laboratory testing and careful infection control.

### **Near-Drowning**

Survival for at least 24 hours following suffocation or submersion in a liquid medium, characterized pathologically by alveolar surfactant washout, pulmonary edema, ventilation-perfusion mismatch, acute respiratory distress syndrome, and hypoxic-ischemic

---

encephalopathy.

**Alternate Names:**

Non-Fatal Drowning; Submersion Injury

**Nearsightedness**

A common focusing problem in which nearby objects are clear but distant objects look blurry because light focuses in front of the retina. Glasses, contact lenses, or selected procedures can improve vision.

**Alternate Names:**

Myopia; Short-Sightedness

**Nebulizer**

A device that converts liquid medicine into an aerosol for inhalation through a mouthpiece or mask. Common types include jet nebulizers, ultrasonic nebulizers, and vibrating-mesh nebulizers; they differ in how the aerosol is generated and delivered.

**Alternate Names:**

Nebuliser; Atomizer

**NEC**

Abbreviation; see *Neuroendocrine Carcinoma*.

**Necator**

A group of hookworms that live in the small intestine. They enter through the skin, mature in the gut, and feed on blood, potentially causing abdominal symptoms, iron-deficiency anemia, weakness, and poor growth.

**Necator Americanus**

A hookworm species that infects humans through skin penetration by larvae and causes intestinal blood loss.

**Neck**

The neck connects the head to the trunk and contains the cervical spine, airway, esophagus, thyroid, vessels, nerves, muscles, and lymph nodes.

**Neck And Back Pain**

Pain in the neck, upper back, lower back, or surrounding tissues, commonly caused by strain, age-related changes, disc problems, or trauma. Weak-

ness, numbness, walking difficulty, bowel or bladder changes, fever, cancer history, or major trauma may indicate a serious cause.

**Neck Cancer**

Cancer arising in tissues of the neck, such as the throat, thyroid, salivary glands, lymph nodes, or other structures. Symptoms may include a persistent lump, pain, swallowing trouble, voice change, or unexplained weight loss.

**Neck Dissection**

Surgical removal of cervical lymph nodes and sometimes surrounding tissues to treat or stage head and neck cancer.

**Neck Lump**

A localized swelling or mass in the neck caused by enlarged lymph nodes, thyroid or salivary disease, cyst, infection, or tumor; persistent or concerning masses require evaluation.

**Neck Mass**

A lump or swelling in the neck caused by an enlarged lymph node, congenital cyst, thyroid or salivary-gland lesion, infection, or tumor. A persistent, enlarging, firm, fixed, or unexplained mass—especially in an adult—can have an important underlying cause.

**Neck Pain**

Pain or stiffness in the neck, commonly caused by muscle strain, poor posture, injury, arthritis, degenerative disc disease, or nerve compression. It may be localized or spread to the shoulders, arms, or head.

**Neck Stiffness**

Reduced ability to move the neck comfortably because of muscle tightness, inflammation, injury, or irritation of tissues around the spine or meninges. Severe stiffness with fever, headache, confusion, rash, or neurologic symptoms requires urgent assessment.

**Alternate Names:**

Cervicalgia; Cervical Pain; Neck Stiffness

---

## Neck Pain And Dizziness

Neck pain with dizziness may reflect muscle strain, migraine, inner-ear disease, medication, blood-pressure change, or less commonly vascular or neurologic disease. Sudden severe symptoms, weakness, speech or vision change, fainting, or inability to walk may occur with a serious condition.

## Neck Sprain

Alternate name; see *Whiplash*.

## Neck Strain

Alternate name; see *Whiplash*.

## Neck X-Ray

Plain radiographic imaging of the cervical spine or neck soft tissues used to evaluate alignment, fractures, degenerative change, foreign bodies, or selected airway abnormalities.

## Necrobiosis Lipoidica

A chronic inflammatory skin disorder characterized by yellow-brown, atrophic plaques, usually on the shins. It is associated with diabetes but can occur without diabetes.

## Necrobiosis Lipoidica Diabeticorum

A chronic inflammatory skin disorder producing yellow-brown atrophic plaques with a raised border, most often on the shins and associated with diabetes, although it can occur without diabetes.

## Necrolytic Acral Erythema

A rare rash with dark, thickened, or blistered patches, usually on the feet and ankles. It is often linked to long-term hepatitis C but can also occur with low zinc or other metabolic problems. Testing is needed to find the cause.

### Alternate Names:

NAE; Hepatitis C-Associated Necrolytic Erythema

## Necrolytic Migratory Erythema

A recurrent, erosive, erythematous eruption that often affects periorificial and intertriginous skin. It is classically associated with glucagonoma but may

have other causes, so evaluation should consider nutritional, hepatic, pancreatic, and metabolic disease.

## Necrosis

Necrosis is death of cells or tissue within a living body caused by ischemia, infection, toxins, trauma, inflammation, or other injury.

## Necrotizing Anterior Scleritis

A severe inflammation of the white outer coat of the eye that causes tissue damage and can threaten vision. It usually causes intense eye pain, redness, and light sensitivity.

## Necrotizing Enterocolitis

A serious illness in which inflammation damages the intestine, mainly in premature or medically fragile newborns. A swollen abdomen, feeding problems, green vomit, bloody stools, or unusual sleepiness may occur.

## Necrotizing Fasciitis

A life-threatening infection that rapidly destroys tissue beneath the skin. Early skin changes may be mild despite severe pain and deeper damage. Treatment usually includes surgery and antibiotics.

## Necrotizing Granuloma

A granulomatous inflammatory lesion containing central tissue necrosis, commonly associated with infections such as tuberculosis or fungal disease but requiring clinical and histologic correlation.

## Necrotizing Infections

Severe infections that cause tissue death through germs, toxins, blocked blood flow, or intense inflammation. Examples include necrotizing fasciitis, necrotizing pneumonia, and necrotizing enterocolitis. Rapidly worsening pain, swelling, skin changes, severe illness, or organ problems may occur.

## Necrotizing Pneumonia

Severe pneumonia in which infection causes necrosis and cavitation of lung tissue, often accompanied by

---

sepsis, respiratory failure, or abscess formation.

### **Necrotizing Sialometaplasia**

A benign, self-limiting, ischemic inflammatory lesion of the salivary glands (most commonly involving the minor salivary glands of the hard palate) that clinically and histologically mimics mucoepidermoid carcinoma or squamous cell carcinoma.

#### **Alternate Names:**

Ischemic Sialadenitis; Benign Palatal Sialometaplasia

### **Necrotizing Soft Tissue Infection**

A rapidly progressive infection of the tissue beneath the skin or in muscle that causes severe pain, swelling, severe illness, and tissue death. Treatment may include surgery and antibiotics.

### **Necrotizing Vasculitis**

Inflammation of blood-vessel walls causing fibrinoid necrosis and potential vessel narrowing, thrombosis, hemorrhage, ischemia, or organ injury.

### **Needle**

A hollow, sharp tube used to inject medicine or remove blood, fluid, or tissue. Needles come in different lengths and widths; the appropriate size depends on the procedure and body site.

### **Needle Biopsy**

Removal of cells, fluid, or a tissue sample through a needle for cytologic or histopathologic examination. Fine-needle aspiration primarily collects cells, whereas core biopsy collects tissue architecture.

### **Needle Decompression**

An emergency procedure used to release trapped air from around a collapsed lung when a tension pneumothorax is causing severe breathing or circulation problems. A trained clinician places a needle or catheter through the chest, then usually inserts a chest tube for ongoing treatment. It is a temporary lifesaving measure, not a complete repair.

#### **Alternate Names:**

Needle Thoracostomy; Tension Pneumothorax De-

compression; Emergency Chest Decompression

### **Needle Driver**

A surgical instrument with gripping jaws and handles used to hold and guide a suture needle through tissue. Its design helps control the needle while protecting the operator's fingers.

#### **Alternate Names:**

Needle Holder Suture Needle Holder

### **Needle Holder**

Alternate name; see *Surgical Needle Holder*.

### **Needlestick Injury**

A puncture from a needle or sharp object that may expose someone to blood or body fluids. It can create a risk of infection, including HIV or hepatitis, and may lead to testing or preventive treatment.

### **Neer Sign**

A physical examination finding used to evaluate for subacromial impingement syndrome. The examiner stabilizes the patient's scapula while passively elevating the internally rotated arm in full forward flexion; reproduction of anterior shoulder pain indicates impingement of the rotator cuff tendons against the coracoacromial arch.

#### **Alternate Names:**

Neer Impingement Test

### **Neer Test**

A clinical diagnostic physical examination maneuver for subacromial shoulder impingement; the examiner stabilizes the patient's scapula to prevent upward rotation and passively performs forced forward elevation and internal rotation of the fully extended arm, driving the greater tuberosity against the anteroinferior acromion to reproduce anterior shoulder pain.

#### **Alternate Names:**

Neer Impingement Sign; Neer Shoulder Test

### **Negative Feedback**

A control process in which a change triggers a response that reduces or reverses the original change.

---

It helps keep body functions, such as temperature, blood sugar, and hormone levels, within a useful range.

### **Negative Predictive Value**

The chance that a person with a negative test result truly does not have the condition. It depends on test accuracy and how common or likely the condition was before testing.

### **Negative Pressure Pulmonary Edema**

A form of non-cardiogenic pulmonary edema provoked by vigorous, forceful inspiratory efforts against an acutely obstructed upper airway (such as post-extubation laryngospasm), generating extreme negative intrathoracic pressures that increase venous return and cause alveolar capillary stress failure.

#### **Alternate Names:**

NPPE; Post-Obstructive Pulmonary Edema

### **Negative Pressure Ventilation**

Breathing support that uses pressure around the chest to help the lungs expand, instead of pushing air directly through the airway. It is used rarely, including for some neuromuscular conditions.

### **Negative-Strand RNA Virus**

A virus whose genetic material cannot be read directly to make proteins. After entering a cell, it must first make a readable copy of its RNA before producing new virus particles.

### **Negative Symptoms**

A reduction or loss of usual emotional expression, speech, motivation, pleasure, or interest in other people. They can occur in schizophrenia and some other conditions and may interfere with work, relationships, and daily activities.

### **Neglect**

Failure to provide a child with necessary food, shelter, supervision, education, emotional care, or medical care, causing harm or risk of harm. The term can also describe ignoring one side of space after brain injury; the intended meaning depends on context.

### **Neglected Tropical Disease**

An infectious or parasitic disease that mainly affects populations in tropical or subtropical regions with limited resources. Examples include schistosomiasis, leprosy, dengue, and lymphatic filariasis.

### **Neglect Syndrome**

A neurologic problem in which a person fails to notice or respond to one side of the body or surrounding space, usually after injury to the opposite side of the brain. It is more than ordinary inattention.

### **Neisseria Gonorrhoeae**

A gram-negative diplococcus that causes gonorrhea, a sexually transmitted infection. The organism has developed resistance to many antibiotics including fluoroquinolones and macrolides.

### **Neisseria Infection**

Infection caused by a pathogenic *Neisseria* species. The most important human infections are gonorrhea caused by *N. gonorrhoeae* and meningococcal disease caused by *N. meningitidis*; illness may affect the genital tract, throat, rectum, joints, bloodstream, or meninges, and treatment depends on the site and antibiotic susceptibility.

#### **Alternate Names:**

*Neisseria* Infections

### **Neisseria Meningitidis**

A gram-negative diplococcus that causes meningitis and meningococcemia. The organism has a polysaccharide capsule that is the target of conjugate vaccines. Serogroups A, B, C, W, X, and Y cause most disease. Meningitis presents with fever, headache, neck stiffness, and altered mental status.

### **Nelson Syndrome**

Enlargement and hypersecretion of an adrenocorticotrophic-hormone-producing pituitary tumour after bilateral adrenalectomy, usually for Cushing disease. It may cause rapid skin hyperpigmentation, headache, visual-field loss, and other effects of an expanding sellar mass.



---

### **Nemaline Myopathy**

A group of inherited muscle diseases that cause weakness because muscle fibers do not develop or work normally. Severe forms begin in infancy and may affect movement, feeding, or breathing; milder forms begin later. The severity varies widely, and care may include breathing support, nutrition, physical therapy, and genetic counseling.

#### **Alternate Names:**

Rod Body Myopathy; Nemaline Rod Myopathy

### **Neoadjuvant Therapy**

Treatment given before the main treatment, such as surgery or radiotherapy, to shrink a tumor or make it easier to treat. It may also show how the tumor responds and help doctors plan the next treatment.

### **Neocortex**

The outer part of the cerebral hemispheres involved in conscious perception, language, planning, memory, and voluntary movement.

### **Neoehrlichiosis**

An infection caused by the bacterium *Candidatus Neoehrlichia mikurensis*, usually transmitted by ticks. It can cause fever and blood-vessel inflammation, especially in people with weakened immunity.

### **Neologisms**

Newly created words, phrases, or idiosyncratic combinations of word fragments that have meaning only to the speaker and are incomprehensible to others. They are characteristic features of formal thought disorder in schizophrenia and fluent (Wernicke) aphasia.

### **Neonatal**

Relating to a newborn, especially during the first 28 days after birth. This period includes important changes in breathing, feeding, temperature control, immunity, circulation, and adjustment to life outside the womb.

### **Neonatal Abstinence Syndrome**

Withdrawal symptoms in a newborn after exposure

to opioids or certain other medicines or substances during pregnancy. Symptoms may include trembling, irritability, poor feeding, vomiting, diarrhea, sweating, or breathing problems.

### **Neonatal Cephalohematoma**

A collection of blood beneath the outer covering of a newborn's skull, usually caused by birth trauma. It is limited by the skull's seams and generally resolves over time, but it can be associated with injury or jaundice.

### **Neonatal Conjunctivitis**

Inflammation of the eye's outer lining during the first month of life, caused by chemical irritation, bacteria, or viruses acquired during delivery. Pus-like discharge, eyelid swelling, and corneal involvement may occur.

### **Neonatal Cystic Fibrosis Screening Test**

A newborn screening test, usually measuring immunoreactive trypsinogen with reflex DNA or sweat testing, used to identify infants at increased risk for cystic fibrosis.

### **Neonatal Diabetes Mellitus**

Diabetes first diagnosed during the first six months of life. It may be temporary or lifelong and is usually caused by a genetic change affecting insulin production; specialist care is needed.

### **Neonatal Galactorrhea**

A temporary milk-like breast secretion in a newborn caused by exposure to maternal hormones. It usually resolves without treatment, and pressure on the breasts can cause irritation or infection.

#### **Alternate Names:**

Neonatal Milk Secretion; Newborn Galactorrhea

### **Neonatal Herpes Simplex**

Herpes simplex infection in a baby during the first 28 days of life, usually acquired around delivery. It may affect the skin, eyes, mouth, brain, or several organs and can resemble sepsis. Treatment uses antiviral medicine.

---

### Neonatal Herpes Simplex Virus

A severe neonatal infection caused by herpes simplex virus, typically acquired during passage through the birth canal, presenting within the first month of life as skin, eye, and mouth disease, central nervous system disease, or disseminated disease with multiorgan involvement.

### Neonatal Hyperbilirubinemia

An excess of bilirubin in a newborn's blood, causing yellow skin or eyes. Mild cases are common, but rapidly rising or very high levels can damage the brain; treatment may include frequent feeding, phototherapy, or rarely an exchange transfusion.

### Neonatal Encephalopathy

A syndrome of disturbed brain function in a newborn, ranging from abnormal alertness or muscle tone to seizures or coma. Causes include hypoxic-ischemic injury, infection, stroke, metabolic disorders, and genetic conditions; it is broader than hypoxic-ischemic encephalopathy.

#### Alternate Names:

Newborn Encephalopathy

### Neonatal Hyperglycemia

An abnormally elevated blood glucose level in infants and neonates (typically defined as blood glucose > 145 mg/dL or > 8.0 mmol/L). Common causes include extreme prematurity, intravenous dextrose overinfusion, sepsis, severe physiological stress, corticosteroid therapy, or transient neonatal diabetes mellitus.

#### Alternate Names:

Infantile Hyperglycemia; High Blood Glucose in Infants; Newborn Hyperglycemia

### Neonatal Hypertension

Sustained blood pressure elevation in infants and neonates exceeding age- and birthweight-adjusted norms. Frequently secondary to umbilical artery catheter-related renal artery thrombosis, congenital renal anomalies, bronchopulmonary dysplasia, or congenital heart defects.

#### Alternate Names:

Infant Hypertension; Neonatal High Blood Pressure;

Infantile Hypertension

### Neonatal Hypocalcemia

Low serum calcium in a newborn, which may cause jitteriness, poor feeding, hypotonia, apnea, or seizures. Causes include prematurity, maternal diabetes, birth asphyxia, and disorders of parathyroid function.

### Neonatal Hypoglycemia

Blood sugar that is too low in a newborn. It may cause trembling, sleepiness, poor feeding, breathing pauses, or seizures and is more likely with prematurity, unusual birth size, or maternal diabetes; prompt monitoring and treatment are needed.

### Neonatal Hypothermia

A newborn's body temperature below 36.5°C. It can result from exposure, prematurity, infection, or illness and may worsen breathing, blood sugar, and circulation problems. Prompt warming and assessment are needed; skin-to-skin contact and a warm, dry environment help prevent it.

### Neonatal Hypothyroidism

Insufficient thyroid hormone production in a newborn, which can impair brain development and growth if untreated; newborn screening enables early diagnosis and treatment.

### Neonatal Hypoxic-Ischemic Encephalopathy

Brain injury in a newborn caused by too little oxygen or blood flow around the time of birth. It may cause poor alertness, abnormal muscle tone, seizures, breathing difficulty, or injury to other organs. Selected babies benefit from carefully controlled cooling started soon after birth.

### Neonatal Infections

Infections occurring during the first weeks of a newborn's life. They may begin before or soon after birth from exposure during pregnancy or delivery, or later from the environment or medical devices. Warning signs include poor feeding, temperature change, breathing difficulty, unusual sleepiness, irri-

---

tability, or color change.

### Neonatal Jaundice

Yellow discoloration of a newborn's skin or eyes caused by elevated bilirubin. It is common but may require treatment or investigation when severe, early, or prolonged.

#### Alternate Names:

Neonatal Hyperbilirubinemia; Icterus Neonatorum

### Neonatal Lupus

A condition caused by antibodies passed from a pregnant person to a baby. It may cause a temporary rash, low blood counts, liver problems, or a slow heart rhythm. Most effects improve as the antibodies leave the baby's blood, but heart block can be serious and may require a pacemaker.

#### Alternate Names:

Neonatal Lupus Erythematosus; NLE

### Neonatal Mechanical Ventilation

Invasive or non-invasive mechanical positive-pressure ventilation specifically tailored for neonates and infants with respiratory failure, severe respiratory distress syndrome (RDS), bronchopulmonary dysplasia, or apnea of prematurity. Emphasizes lung-protective strategies such as high-frequency oscillatory ventilation and synchronized intermittent mandatory ventilation.

#### Alternate Names:

Pediatric Mechanical Ventilation; Neonatal Ventilation Support; Infant Mechanical Ventilation

### Neonatal Neutropenia

An absolute neutrophil count (ANC) below 1,000/ $\mu$ L to 1,500/ $\mu$ L in infants, or below 500/ $\mu$ L in severe cases. Etiologies include neonatal sepsis, maternal preeclampsia, alloimmune or autoimmune neutropenia, twin-to-twin transfusion, bone marrow failure syndromes (such as Kostmann syndrome), or drug exposure.

#### Alternate Names:

Infant Neutropenia; Neonatal Low Neutrophil Count; Pediatric Neutropenia

### Neonatal Opioid Withdrawal Syndrome

Withdrawal symptoms in a newborn exposed to opioid medicines or drugs during pregnancy. The baby may be hard to settle, tremble, feed poorly, vomit, have loose stools, sweat, or breathe rapidly. Quiet surroundings, swaddling, skin-to-skin contact, and feeding support are used first; some babies need hospital medicines and monitoring.

#### Alternate Names:

NOWS; Opioid-Induced NAS

### Neonatal Peripheral Arterial Line

A fine indwelling cannula placed in a peripheral artery (such as the radial, posterior tibial, or dorsalis pedis artery) of an infant for continuous invasive hemodynamic arterial blood pressure monitoring and frequent arterial blood gas sampling in neonatal intensive care.

#### Alternate Names:

Infant Arterial Line; Pediatric A-Line; Neonatal Arterial Cannulation

### Neonatal Peripheral IV Line

A short peripheral venous catheter inserted into a superficial vein in the hand, foot, forearm, or scalp of an infant for short-term hydration, medication delivery, or blood transfusions.

#### Alternate Names:

Infant PIV; Pediatric Peripheral Cannula; Neonatal IV Cannulation

### Neonatal PICC Line

A small-caliber (1.0–2.0 Fr) central venous catheter inserted percutaneously into a peripheral vein of an infant or neonate with the tip positioned in the superior or inferior vena cava, providing reliable vascular access for total parenteral nutrition and hyperosmolar medications.

#### Alternate Names:

Neonatal Percutaneous Central Line; Infant PICC; Pediatric Percutaneous Central Catheter

### Neonatal Pneumothorax

Pathologic accumulation of extrapulmonary air within the pleural space in a newborn or infant, which may occur spontaneously or secondary to

---

meconium aspiration, respiratory distress syndrome, or mechanical barotrauma.

**Alternate Names:**

Infant Pneumothorax; Neonatal Air Leak Syndrome; Newborn Pleural Air Collection

**Neonatal Polycythemia**

An abnormally high red-cell concentration in a newborn, often defined by a venous hematocrit of at least 65 percent. It may increase blood viscosity and cause respiratory, neurologic, or metabolic symptoms.

**Neonatal Respiratory Distress Syndrome**

Breathing difficulty in a premature newborn caused mainly by insufficient pulmonary surfactant, leading to stiff lungs and impaired oxygen exchange.

**Neonatal Resuscitation**

Emergency steps used to help a newborn who is not breathing well or has a slow heart rate after birth. They may include warmth, airway support, assisted breathing, and chest compressions.

**Neonatal Seizures**

Seizures occurring in a newborn, usually during the first 28 days after birth. They may be subtle or detectable only with electroencephalography (EEG). Causes include oxygen-related brain injury, stroke or bleeding, infection, low blood sugar or calcium, and genetic or metabolic conditions; suspected seizures need urgent medical assessment.

**Neonatal Sepsis**

A serious infection causing illness throughout a newborn's body during the first 28 days of life. It may begin before or after birth and can cause poor feeding, temperature change, breathing difficulty, sleepiness, or color change.

**Neonatal Skin Conditions**

Skin changes common in newborns, including temporary rashes, tiny white spots, heat rash, and small yellow bumps. Most are harmless, but blistering, widespread redness, fever, poor feeding, or a rapidly worsening rash may signal illness.

**Neonatal Testicular Torsion**

Twisting of the spermatic cord around a testicle during the newborn period, reducing or stopping its blood supply. It may appear as a firm, enlarged, discolored, or painless scrotal swelling.

**Neonatal Thyrotoxicosis**

Excess thyroid activity in a newborn, usually caused by thyroid-stimulating antibodies passed from the mother. It may cause a fast heartbeat, irritability, poor feeding, poor weight gain, sweating, or breathing problems and usually needs temporary treatment and monitoring.

**Neonatal Transition**

The newborn's adjustment from receiving oxygen through the placenta to breathing with the lungs after birth. The lungs fill with air, blood flow changes, and the baby must begin controlling temperature, blood pressure, and feeding.

**Neonate**

A neonate is a newborn infant during the first 28 completed days after birth, a period requiring attention to feeding, temperature, jaundice, infection, breathing, and transition to extrauterine life.

**Neonatologist**

A pediatrician who specializes in the medical care of newborn infants, particularly premature and critically ill neonates.

**Neoplasia**

Abnormal, uncontrolled growth of cells that forms a new mass or tissue. It may be benign and remain localized, or malignant and invade nearby tissues or spread elsewhere.

**Neoplasm**

A new, abnormal growth of cells that forms a mass or tissue. A neoplasm may be benign and stay localized, or malignant and invade nearby tissues or spread to distant organs.

---

## Neoplastic

Relating to an abnormal new growth of cells. A neoplastic growth may be noncancerous and localized or cancerous and able to invade nearby tissue or spread.

## Neostigmine

An acetylcholinesterase inhibitor used to reverse selected nondepolarizing neuromuscular blockade and to treat some disorders of neuromuscular transmission or bowel and bladder motility.

## Nephoptosis

Abnormal downward movement or mobility of a kidney, also called floating kidney.

## Nephrectomy

Surgery to remove part or all of a kidney. It may treat kidney cancer, severe kidney damage, infection, or a kidney that no longer works.

## Nephric

Relating to the kidney. In embryology, nephric structures are involved in development of the urinary system.

## Nephritic Syndrome

A pattern of kidney inflammation that causes blood in the urine, reduced urine output, swelling, and high blood pressure. Protein may also leak into the urine. It can reduce kidney function and requires evaluation to find and treat the cause.

## Nephritis

Inflammation of kidney tissue that can cause blood or protein in the urine, swelling, high blood pressure, reduced urine, or loss of kidney function. Infection, medicines, and immune diseases are possible causes.

## Nephroblastoma

Another name for Wilms tumor, the most common kidney cancer in children. It may appear as a painless abdominal lump, abdominal pain, blood in the urine, fever, or high blood pressure and needs specialist treatment.

## Nephrocalcinosis

Buildup of calcium deposits inside kidney tissue. It may result from high calcium levels, hormone or kidney disorders, inherited conditions, dehydration, or medicines and can lead to stones, blood in the urine, or reduced kidney function.

## Nephrogenic Diabetes Insipidus

A condition in which the kidneys do not respond properly to vasopressin, the hormone that helps conserve water. It causes large amounts of dilute urine and intense thirst; inherited changes, kidney disease, electrolyte problems, or medicines may cause it.

## Nephrogenic Fibrosing Dermopathy

An older name for nephrogenic systemic fibrosis, a rare condition causing thick, tight, hardened skin and sometimes stiff joints in people with severe kidney disease, usually after exposure to certain gadolinium contrast agents.

## Nephrogenic Systemic Fibrosis

A rare disorder in which skin and connective tissue become thick and hardened, sometimes limiting movement. It is linked mainly to certain gadolinium contrast exposures in people with severe kidney impairment and may affect internal tissues.

## Nephrolithiasis

Formation of stones in the kidneys or urinary tract. Stones can cause severe side or back pain, blood in the urine, nausea, or blockage; dehydration, inherited conditions, medicines, and diet can contribute.

## Nephrologist

A doctor who specializes in kidney disease. Nephrologists assess reduced kidney function, high blood pressure related to kidney problems, kidney inflammation, and mineral or salt disturbances and may manage dialysis or transplant care.

## Nephrology

The medical specialty concerned with kidney disease, fluid and electrolyte disorders, hypertension, dialysis, and kidney transplantation.



---

## Nephrolysis

Surgical freeing of a kidney from surrounding adhesions or tissue.

## Nephron

The microscopic working unit of a kidney. Each nephron filters blood, returns needed water and substances to the body, and removes waste to make urine; each kidney contains about one million.

## Nephron Function

Each kidney contains about one million nephrons, its microscopic filtering units. A nephron filters blood, removes wastes into urine, and returns needed water and substances to the bloodstream while helping control salts, acid, and blood pressure.

## Nephronophthisis

A rare inherited kidney disease that gradually reduces the kidneys' ability to concentrate urine. It may cause excessive thirst and urination, anemia, growth problems, and eventually kidney failure.

## Nephropathy

Disease or damage of the kidneys. It may reduce waste removal, disturb fluid and salt balance, cause protein or blood in the urine, or eventually lead to kidney failure.

## Nephroptosis

Abnormal downward movement of a kidney when a person stands, sometimes causing flank discomfort, urinary symptoms, or intermittent obstruction.

### Alternate Names:

Wandering Kidney; Ren Mobilis; Hypermobility Kidney

## Nephrosclerosis

Hardening and scarring of kidney tissue, commonly caused by long-standing high blood pressure or blood-vessel disease. Progressive damage can reduce filtering ability, cause protein in urine, and lead to chronic kidney disease.

## Nephrostomy

A procedure that creates a drainage path from the kidney through the skin when urine cannot pass normally to the bladder. A temporary tube usually carries urine into a bag while the blockage is treated.

## Nephrostomy Tube

A drainage tube passed through the skin into the kidney to carry urine into a bag when the ureter is blocked. Blockage may be caused by a stone, tumor, or narrowing. Infection with blockage can require urgent drainage; the tube may be temporary or long-term.

## Nephrotic Range Proteinuria

Very large protein loss in the urine, usually more than 3.5 grams per day in an adult. It often indicates serious damage to the kidney filters and can cause swelling, low blood protein, high cholesterol, blood clots, or infection.

## Nephrotic Syndrome

A group of findings caused by kidney-filter damage: a large amount of protein in the urine, low albumin in the blood, swelling, and often high blood cholesterol. It may cause frothy urine and swelling around the eyes, legs, or feet, and can increase the risk of blood clots or infection. Causes include diseases of the kidney filters and illnesses such as diabetes or lupus.

### Alternate Names:

Nephrosis; Protein-losing nephropathy; Glomerular nephrosis.

## Nephrotoxic

Describing a substance or condition that can damage the kidneys or reduce their function. Examples include some medicines, poisons, infections, and severe dehydration; risk depends on dose, duration, and kidney health.

## Nephrotoxicity

Kidney damage caused by a medicine, poison, contrast material, or other substance. It may reduce urine output or raise blood creatinine; risk is lowered by avoiding triggers when possible and adjusting



---

doses for kidney function.

### Nernst Equation

A fundamental biophysical formula that calculates the equilibrium electrical potential across a semipermeable cell membrane for a specific single ion species based on its intracellular and extracellular concentration gradient at physiological body temperature.

### Nerve

A bundle of nerve fibers that carries messages between the brain or spinal cord and the rest of the body. Nerves control movement and carry sensations such as touch, pain, and temperature.

### Nerve Biopsy

Removal of a small piece of a nerve for laboratory examination when serious nerve disease is suspected. It may help identify inflammation, infection, abnormal deposits, or a tumor, but can leave permanent numbness or weakness near the biopsy site.

### Nerve Block

An injection of numbing medicine near a nerve or group of nerves to temporarily stop pain signals. It may help diagnose the source of pain or provide pain relief during or after a procedure; weakness or other temporary effects can occur.

### Nerve Compression

Alternate name; see *Pinched Nerve*.

### Nerve Conduction Studies

Tests that stimulate peripheral nerves with small electrical pulses and record how quickly and strongly signals travel. They help identify nerve damage, compression, loss of the nerve's insulating covering, or loss of nerve fibers.

### Nerve Conduction Study

A test that gives small electrical pulses to a nerve and measures how quickly and strongly the signal travels. It helps identify nerve damage, pressure on a nerve, or loss of its protective covering.

### Nerve Conduction Velocity

The speed at which an electrical signal travels along a peripheral nerve. A nerve-conduction test measures this speed and the signal strength to help identify nerve damage, pressure on a nerve, or damage to its insulating covering.

### Nerve Deafness

Hearing loss caused by damage to the cochlea, auditory nerve, or central auditory pathways rather than blockage of sound transmission through the outer or middle ear.

#### Alternate Names:

Sensorineural Hearing Loss; SNHL

### Nerve Endings

The tiny ends of nerves that detect sensations such as touch, pain, heat, or pressure, or release signals to muscles and glands. Injury or irritation can cause pain, numbness, tingling, or abnormal sweating.

### Nerve Fiber

The long, thread-like part of a nerve cell that carries signals. Most nerve fibers are axons, sometimes covered by a fatty insulating layer called myelin.

### Nerve Fiber Types

A physiological classification of peripheral nerve axons based on diameter and conduction velocity: group A (thickly myelinated motor and sensory fibers: alpha, beta, gamma, delta), group B (thinly myelinated preganglionic autonomic fibers), and group C (unmyelinated, slow postganglionic and pain/temperature C-fibers).

#### Alternate Names:

Erlanger-Gasser Classification

### Nerve Impulse

An electrical signal that travels along a nerve cell or nerve fiber to carry information.

### Nerve Injury

Damage to a nerve that can cause pain, numbness, tingling, weakness, altered reflexes, or changes in sweating, blood pressure, digestion, or bladder func-

---

tion. Recovery depends on the nerve, severity, cause, and treatment.

### **Nerve Plexus**

A network where nerves join and branch to supply a body region. The brachial plexus, for example, supplies much of the arm.

### **Nerve Root**

The beginning part of a spinal nerve where it connects with the spinal cord. Sensory and motor roots carry signals into and out of the cord.

### **Nerve Sheath Myxoma**

A benign myxoid peripheral-nerve-sheath tumor, usually presenting as a slow-growing papule or nodule on the fingers, hands, or upper extremities.

### **Nerve Sparing**

A surgical approach that tries to preserve nearby nerves needed for erection, sensation, or other functions. It may improve recovery after prostate or other pelvic surgery, but success depends on disease and nerve involvement.

### **Nerve Tract**

A group of nerve fibers traveling together inside the brain or spinal cord. Fibers in a tract carry signals between specific regions, so damage can cause predictable movement, sensation, or organ-control problems.

### **Nervous Colon Syndrome**

An older name for irritable bowel syndrome, a disorder causing recurrent abdominal pain with diarrhea, constipation, or both without a structural explanation. Symptoms may improve with individualized diet, medicines, and stress management.

### **Nervous System**

The brain and spinal cord (central nervous system), peripheral nerves and ganglia (peripheral nervous system), and supporting cells that detect and process information. It coordinates sensation, movement, automatic body functions, cognition, emotion,

learning, and behavior.

### **Nervous Tissue**

Tissue made of nerve cells and supporting cells that detects changes, processes information, and sends signals through the body. It forms the brain, spinal cord, and nerves and helps control movement, sensation, and organ function.

### **Nesidioblastosis**

A rare diffuse or focal hyperfunctional hypertrophy and hyperplasia of pancreatic beta cells budding from the pancreatic ductal epithelium, causing endogenous persistent hyperinsulinemic hypoglycemia in neonates (congenital hyperinsulinism) and adults (post-gastric bypass hyperinsulinemic hypoglycemia).

#### **Alternate Names:**

Congenital Hyperinsulinism; Pancreatic Beta-Cell Hyperplasia

### **NET**

Abbreviation; see *Neuroendocrine Tumors*.

### **Netherton Syndrome**

A severe autosomal recessive genodermatosis caused by mutations in the SPINK5 gene encoding the serine protease inhibitor LEKTI, characterized by the classic triad of congenital ichthyosiform erythroderma (ichthyosis linearis circumflexa), trichorrhexis invaginata ('bamboo hair'), and severe atopic diathesis.

#### **Alternate Names:**

SPINK5 Deficiency; Ichthyosis Linearis Circumflexa Syndrome

### **Nettle Rash**

An itchy rash made of raised patches that come and go, often within hours. It results from substances released in the skin and may be triggered by infection, foods, medicines, pressure, heat, or no clear cause.

### **Neural**

Relating to nerves, nerve cells, or the nervous sys-

---

tem. In pregnancy, the neural tube develops into the brain and spinal cord; failure of it to close can cause birth defects such as spina bifida or anencephaly.

### Neural Crest

A temporary embryonic multipotent cellular population located along the borders of the closing neural folds that undergoes epithelial-to-mesenchymal transition and migrates extensively to form peripheral sensory ganglia, autonomic ganglia, Schwann cells, melanocytes, craniofacial bones, and adrenal chromaffin cells.

### Neuralgia

Neuralgia is sharp, burning, or electric pain along the distribution of a nerve, caused by irritation, compression, infection, injury, or other disease.

### Neuralgia Amyotrophica

An acute, idiopathic autoimmune inflammatory brachial plexopathy (Parsonage-Turner syndrome) characterized by the sudden onset of severe, unprovoked, sharp unilateral shoulder and periscapular pain, followed days to weeks later by patchy flaccid muscle weakness and rapid wasting in the distribution of the upper brachial plexus.

#### Alternate Names:

Parsonage-Turner Syndrome; Brachial Radiculitis

### Neural Pathway

A connected route of nerve cells and nerve fibers that carries signals between different parts of the nervous system.

### Neural Tube Defects

Birth defects caused when the early structure that becomes the brain and spinal cord does not close completely during the first month of pregnancy. They include anencephaly, encephalocele, and spina bifida and vary from fatal to mild.

### Neuraminidase Inhibitors

Antiviral medicines that block a surface protein used by influenza viruses to release new virus particles from infected cells. They work best when started early and are used according to the influenza type,

illness severity, and risk of complications.

### Neurapraxia

The mildest form of peripheral nerve injury in the Seddon classification, characterized by transient conduction block caused by focal demyelination or localized ischemia without structural disruption of the underlying axon or its connective tissue sheaths; complete recovery occurs spontaneously within days to weeks.

### Neurilemma

The outer layer of a Schwann cell that surrounds and supports a nerve fiber outside the brain and spinal cord.

### Neurilemmoma

Alternate spelling; see *Schwannoma*.

### Neurinomatosis

Alternate name; see *Schwannomatosis*.

### Neuritic Plaques

Abnormal beta-amyloid deposits surrounded by damaged nerve branches, commonly found in Alzheimer disease. They are one pathological feature of the disease and occur with other changes, including abnormal tau protein and nerve-cell loss.

#### Alternate Names:

Senile Plaques; Amyloid Plaques

### Neuritis

Inflammation or irritation of a nerve, often causing pain, altered sensation, weakness, or other changes in nerve function.

### Neuroacanthocytosis

A group of rare inherited disorders causing movement or psychiatric symptoms together with abnormal spiny red blood cells called acanthocytes. Neurologic assessment and genetic evaluation may be needed.

### Neuroanatomy

The study of the structure and organization of the

---

nervous system, including the brain, spinal cord, peripheral nerves, and their connections.

### **Neuroblastoma**

A childhood cancer arising from immature nerve cells, commonly in or near the adrenal glands or spine. It may cause an abdominal mass, pain, weakness, or other symptoms and can range from mild to aggressive.

#### **Alternate Names:**

Sympathicoblastoma; Neural Crest Tumor

### **Neurocardiogenic Syncope**

Alternate name; see *Vasovagal Syncope*.

### **Neurocognitive Disorder**

A decline from a person's previous level of cognitive ability, such as memory, attention, language, executive function, or judgment, that develops after normal earlier development. Mild neurocognitive disorder causes modest decline while independence is largely preserved; major neurocognitive disorder causes substantial decline that interferes with independent daily life and is often called dementia. Causes include brain disease, stroke, injury, infection, substance use, or medicines. Delirium is distinguished by its sudden onset and fluctuating attention and awareness.

### **Neurocognitive Disorders**

Alternate plural form; see *Neurocognitive Disorder*.

### **Neurocutaneous Syndrome**

A disorder affecting both the nervous system and the skin, often because of an inherited genetic change. Examples include neurofibromatosis and tuberous sclerosis.

### **Neurocutaneous Syndromes Pathology**

Pathologic features of inherited disorders affecting the skin, nervous system, eyes, and sometimes other organs. Findings vary by syndrome and may include hamartomas, vascular malformations, neural tumors, or characteristic pigmentary lesions.

### **Neurocysticercosis**

Infection of the brain or spinal cord by the larval form of the pork tapeworm. It can cause seizures, headaches, confusion, or movement problems and is acquired by swallowing tapeworm eggs from contaminated food or hands.

### **Neurocytoma**

A usually slow-growing tumor made of nerve-cell-like cells, most often found in the fluid-filled spaces inside the brain. It may block fluid flow and cause headache, vomiting, vision changes, or balance problems; imaging and tissue testing guide care.

### **Neurodegeneration**

Progressive damage and loss of nerve cells, causing worsening problems with movement, memory, thinking, sensation, or other nervous-system functions.

### **Neurodegeneration With Brain Iron Accumulation**

A group of inherited disorders in which excess iron builds up in certain brain regions and gradually damages nerve cells. Symptoms may include abnormal movements, muscle stiffness, seizures, speech or thinking problems, and vision changes.

### **Neurodegenerative Disease**

A disease in which nerve cells progressively lose function or die. Examples include Alzheimer disease, Parkinson disease, and Huntington disease.

### **Neurodegenerative Disorder**

Another term for a condition involving progressive damage or loss of nerve cells, which can affect movement, memory, behavior, sensation, or autonomic function.

### **Neurodermatitis**

A chronic itchy skin condition in which repeated scratching or rubbing makes a patch thick, leathery, and irritated. Stress, dry skin, or an initial rash may start the itch-scratch cycle.

#### **Alternate Names:**

Lichen Simplex Chronicus; Circumscribed Neuro-

---

dermatitis

### Neurodevelopmental Disorders

A group of conditions that begin during the developmental period and affect cognition, communication, behavior, motor function, learning, or adaptive skills. Examples include attention-deficit/hyperactivity disorder, autism spectrum disorder, intellectual disability, communication disorders, specific learning disorder, motor disorders, and tic disorders.

### Neuroendocrine

Relating to the connection between the nervous system and hormone system, or to cells that receive nerve signals and can release hormones. Neuroendocrine cells are found in many organs and can sometimes form tumors.

### Neuroendocrine Carcinoma

A cancer arising from cells with nerve-like and hormone-related features. It can begin in the lung or digestive system, often grows quickly, and may release hormones that cause additional symptoms.

### Neuroendocrine Carcinoma of the Skin

Alternate name; see *Merkel Cell Carcinoma*.

### Neuroendocrine Tumor

A growth arising from cells that receive nerve signals and can release hormones. It may begin in organs such as the intestine, pancreas, lung, or appendix and may grow slowly or rapidly; some tumors cause hormone-related symptoms.

### Neuroendocrine Tumors

Growths arising from hormone-producing cells that also respond to nerve signals. They can occur in many organs, may release excess hormones, and range from slow-growing tumors to aggressive cancers.

#### Alternate Names:

NETs; Neuroendocrine Neoplasms; Carcinoid Tumors

### Neuroendocrine Tumors Grading

Classification of a neuroendocrine tumor by how quickly its cells divide, measured by the Ki-67 index and mitotic count. Lower grades usually grow more slowly; higher grades grow faster and may behave more aggressively. The grade helps guide treatment and prognosis.

### Neuroendocrinology

The medical field that studies how the brain and nerves control hormones and how hormones affect the nervous system. It includes conditions involving the pituitary, adrenal glands, thyroid, reproduction, growth, and body-fluid balance.

### Neurofibrillary Tangles

Twisted clumps of an abnormal protein called tau inside nerve cells. They damage the cells and are a characteristic finding in Alzheimer disease and some other disorders that cause progressive memory or thinking problems.

#### Alternate Names:

Paired Helical Filaments; Tau Tangles

### Neurofibroma

A benign peripheral nerve sheath tumor. Solitary neurofibromas are common and present as soft, flesh-colored papules or nodules. Multiple neurofibromas are characteristic of neurofibromatosis type 1 (NF1).

### Neurofibromatosis

A group of inherited conditions in which tumors or other changes develop in tissues containing nerves. Type 1 commonly causes café-au-lait spots, soft nerve tumors, bone changes, and learning difficulties; type 2 mainly affects hearing and balance nerves and can cause other nervous-system tumors.

### Neurofibromatosis-1

An inherited condition that can cause light-brown skin patches, freckles in the armpits or groin, soft growths along nerves, bone changes, vision problems, learning difficulties, or blood-vessel complications. Findings vary widely between people.

---

## Neurofibromatosis 2

An inherited condition in which tumors can grow on nerves, especially the nerves for hearing and balance. It may cause hearing loss, ringing in the ears, balance problems, cataracts, or other brain and spinal-cord tumors.

## Neurofibromatosis Type 1

An autosomal dominant inherited condition caused by a disease-causing change in the *NF1* gene. It may cause light-brown skin patches, freckles in skin folds, soft growths along nerves, bone problems, vision complications, learning difficulties, or blood-vessel problems.

### Alternate Names:

Neurofibromatosis Type I NF1 Peripheral Neurofibromatosis Von Recklinghausen's Disease Von Recklinghausen's Neurofibromatosis Von Recklinghausen's Phakomatosis

## Neurofibromatosis Type 2

An autosomal dominant genetic disorder, usually caused by a disease-causing change in the *NF2* gene, characterized by tumors involving nerves, especially bilateral vestibular schwannomas, with possible hearing loss, balance problems, and other nerve tumors.

### Alternate Names:

NF2; Bilateral Acoustic Neurofibromatosis; MISME Syndrome

## Neurofibromatosis Type I

Alternate spelling; see *Neurofibromatosis Type 1*.

## Neurofibromatosis Type II

Alternate spelling; see *Neurofibromatosis Type 2*.

## Neurofibrosarcoma

Alternate name; see *Malignant Peripheral Nerve Sheath Tumor*.

## Neurogenesis

The formation of new nerve cells. It happens mainly before birth but can continue in limited areas of the adult brain.

## Neuroimmunology

The study of interactions between the nervous system and the immune system. It includes immune-related diseases of the brain, spinal cord, nerves, and muscles and the effects of inflammation on neural function.

## Neurogenic

Caused by, arising from, or related to the nervous system. For example, neurogenic bladder results from nerve damage that disrupts bladder storage or emptying. The specific symptoms and treatment depend on which nerves are affected.

## Neurogenic Bladder

Bladder problems caused by damage to the nerves that control storing or passing urine. A person may have urgency, leakage, difficulty emptying, or urine retention; repeated infection and high bladder pressure can damage the kidneys.

## Neurogenic Claudication

A characteristic symptom complex of lumbar spinal canal stenosis characterized by bilateral dull aching, numbness, cramping, and heaviness in the buttocks, thighs, and calves provoked by walking or standing (lumbar extension) and relieved promptly by lumbar flexion (sitting or leaning forward over a shopping cart).

### Alternate Names:

Pseudoclaudication; Spinal Stenosis Claudication

## Neurogenic Megacolon

Severe widening of the large intestine caused by damage to the nerves that control bowel movement. It can cause long-term constipation, abdominal swelling, and stool blockage and may lead to dangerous bowel inflammation or rupture.

## Neurogenic Orthostatic Hypotension

A drop in blood pressure after standing because nerve damage prevents the body from tightening its blood vessels and maintaining pressure. It may cause light-headedness, blurred vision, weakness, fainting, or falls.



---

## Neurogenic Pulmonary Edema

Rapid-onset, non-cardiogenic pulmonary edema developing acutely after severe central nervous system insults (such as massive subarachnoid hemorrhage, head trauma, or status epilepticus), provoked by an intense sympathetic surge that induces sudden pulmonary vasoconstriction and capillary leak.

### Alternate Names:

NPE

## Neurogenic Shock

A dangerous fall in blood pressure caused by sudden loss of nerve control over blood-vessel tone, usually after a serious spinal-cord injury. The heart rate may be unusually slow and the skin warm.

## Neuroglia

Supporting cells of the nervous system that nourish, protect, insulate, and maintain nerve cells. Different types help control the nerve environment, form protective coverings, remove waste, and respond to injury or infection.

## Neurography

An MRI method that produces detailed pictures of nerves outside the brain and spinal cord. It can help assess trapped, injured, inflamed, or enlarged nerves and may help find some nerve tumors.

## Neurohypophysis

The back part of the pituitary gland that stores and releases oxytocin and vasopressin. These hormones help control childbirth, milk release, water balance, and urine concentration.

## Neurolathyrism

A toxic neurologic disorder caused by excessive consumption of *Lathyrus sativus*, producing progressive spastic weakness of the legs.

## Neuroleptic

An older term for an antipsychotic medicine, used to treat conditions such as schizophrenia, severe mania, or psychosis. The term remains in names such as neuroleptic malignant syndrome; these

medicines may cause sleepiness, movement problems, or metabolic effects.

## Neuroleptic Agents

Antipsychotic medicines used for conditions such as schizophrenia, bipolar disorder, severe agitation, or nausea, depending on the drug. They can cause movement, metabolic, heart-rhythm, and sedation-related effects.

### Alternate Names:

Antipsychotics

## Neuroleptic Agent Toxicity

Overdose or adverse effects from an antipsychotic or other dopamine-blocking agent, causing sedation, hypotension, anticholinergic effects, seizures, QT prolongation, or other dysrhythmias. Neuroleptic malignant syndrome is a distinct delayed reaction with hyperthermia and rigidity; ECG, airway, temperature, and neurologic monitoring guide supportive treatment.

## Neuroleptic Malignant Syndrome

A rare, life-threatening reaction to some antipsychotic or other dopamine-blocking medicines. It causes very high temperature, severe muscle stiffness, confusion, sweating, unstable blood pressure, and a fast heart rate.

## Neurological

Related to the brain, spinal cord, or nerves. Neurological symptoms may include weakness, numbness, seizures, headache, dizziness, confusion, or problems with movement, speech, vision, or memory.

## Neurological Disease

Any disease affecting the brain, spinal cord, peripheral nerves, muscles, or their connections. Symptoms may include weakness, numbness, seizures, headache, movement changes, or problems with thinking.

## Neurological Examination

A clinical assessment of mental status, cranial nerves, motor function, reflexes, coordination, sensation, and gait. It helps localize nervous-system

---

dysfunction and determine the need for further testing.

### **Neurological Rehabilitation**

Treatment and training that help a person regain movement, communication, thinking, or independence after stroke, brain or spinal-cord injury, Parkinson disease, multiple sclerosis, or another nervous-system disorder.

### **Neurologic Deficit**

Loss or impairment of a neurologic function, such as strength, sensation, coordination, vision, language, or consciousness, caused by dysfunction of the nervous system.

### **Neurologic Disorders**

A group of conditions affecting the brain, spinal cord, peripheral nerves, neuromuscular junction, or muscles. Examples include stroke, epilepsy, migraine, multiple sclerosis, Parkinson disease, Alzheimer disease, peripheral neuropathy, meningitis, neuromuscular disorders, and movement disorders.

### **Neurologic Localization**

Using the pattern of weakness, numbness, reflex changes, coordination problems, vision, speech, or thinking changes to identify where the nervous system is affected. This helps clinicians choose likely causes and appropriate tests.

### **Neurologist**

A medical doctor who diagnoses and treats disorders of the brain, spinal cord, nerves, and muscles. Neurologists assess symptoms such as weakness, numbness, seizures, tremor, headaches, memory change, and movement or balance problems.

### **Neurology**

The medical specialty concerned with diseases of the brain, spinal cord, peripheral nerves, and muscles.

### **Neurology PET**

PET imaging used in selected neurologic disorders to assess brain metabolism or receptor activity, such

as in some dementias, epilepsy evaluations, and movement disorders.

### **Neurolysis**

A procedure that frees a nerve from scar tissue or pressure, or deliberately interrupts selected nerve fibers to relieve pain. The purpose and possible effects depend on the nerve treated and the technique used.

### **Neuroma**

A benign growth or reactive thickening involving nerve tissue, often producing pain or altered sensation.

### **Neuromodulation**

Changing nerve activity with electrical, magnetic, chemical, or other signals. Treatments include spinal cord stimulation, deep brain stimulation, and some noninvasive brain-stimulation methods.

### **Neuromuscular**

Relating to nerves, muscles, or the place where a nerve tells a muscle to contract. Neuromuscular diseases can cause weakness, tiredness with activity, muscle wasting, cramps, or problems with breathing or swallowing.

### **Neuromuscular Blockade Reversal**

The pharmacological restoration of normal neuromuscular transmission following surgical paralysis, achieved by administering an acetylcholinesterase inhibitor paired with an anticholinergic (neostigmine and glycopyrrolate) or a selective encapsulating reversal agent (sugammadex for rocuronium/vecuronium).

### **Neuromuscular Blocking Agents**

Medicines used during anesthesia to temporarily relax skeletal muscles, making insertion of a breathing tube and some operations easier. They stop muscle movement but do not provide pain relief or unconsciousness and require breathing support and monitoring.

---

## Neuromuscular Disorders

A group of conditions affecting motor nerves, neuromuscular junctions, or skeletal muscles. Examples include myasthenia gravis, Lambert-Eaton syndrome, muscular dystrophy, spinal muscular atrophy, motor neuron disease, inflammatory myositis, myopathies, and peripheral neuropathies.

## Neuromuscular Hamartoma

A rare benign developmental lesion in which mature skeletal muscle and nerve fascicles are intermixed, typically involving a major peripheral nerve and causing enlargement or neuropathic symptoms.

## Neuromuscular Junction

The small connection where a movement-controlling nerve communicates with a skeletal muscle. The nerve releases a chemical messenger there, causing the muscle to contract.

### Alternate Names:

NMJ; Motor End-Plate

## Neuromyelitis Optica

An autoimmune disorder that mainly causes inflammation of the optic nerves and spinal cord. It can cause vision loss and weakness or altered sensation and is treated with immune-directed therapy.

## Neuromyelitis Optica Spectrum Disorder

An immune-system disorder that repeatedly inflames the optic nerves, spinal cord, or parts of the brain. Attacks may cause vision loss, limb weakness, numbness, vomiting, or bladder problems and can lead to lasting disability without treatment.

### Alternate Names:

Neuromyelitis Optica NMO NMOSD Devic Disease Devic's Disease

## Neuromyotonia

A rare disorder in which nerves repeatedly stimulate muscles, causing ongoing stiffness, twitching, cramps, delayed muscle relaxation, and sometimes excessive sweating. It may be inherited or caused by an immune reaction. An electrical nerve and muscle test helps diagnosis; treatment aims to calm nerve activity and address any underlying cause.

### Alternate Names:

Isaacs Syndrome; Continuous Muscle Fiber Activity Syndrome

## Neuron

A nerve cell that receives, processes, and sends messages using electrical signals and chemical messengers. It has a cell body, branches that receive information, and usually a long fiber that carries messages to other cells.

## Neuronal Ceroid Lipofuscinoses

A group of inherited disorders in which waste material builds up inside nerve cells. They usually cause worsening seizures, vision loss, movement problems, and difficulty thinking or learning, with symptoms and age of onset varying by type.

## Neuronal Migration Disorder

A developmental brain disorder in which nerve cells do not move to their normal positions before birth. It can cause abnormal brain structure, seizures, developmental delay, or learning problems.

## Neurone

The British spelling of neuron, an excitable nerve cell that receives and transmits signals.

## Neuronopathy

A disorder caused mainly by damage to nerve-cell bodies, often in sensory or motor nerve groups. It may cause numbness, imbalance, weakness, or loss of reflexes.

## Neurons

Specialized nerve cells that receive, process, and transmit electrical or chemical signals. They communicate with one another and with muscles or glands.

## Neuron-Specific Enolase

A protein that can be released by nerve cells and some neuroendocrine tumors. A raised blood level may support evaluation or monitoring in selected situations but is not specific enough to diagnose a

---

disease alone.

### **Neuron-Specific Enolase Tumor Marker**

Neuron-specific enolase, a protein that may be increased in some neuroendocrine cancers or after nerve-cell injury. It can support monitoring in selected situations but is not specific enough to diagnose cancer alone.

### **Neuro-Ophthalmology**

The medical specialty that evaluates vision problems caused by the brain, optic nerves, or eye-movement pathways. It assesses symptoms such as unexplained vision loss, double vision, abnormal pupils, or visual-field changes.

### **Neuropathic**

Caused by disease or dysfunction of nerves; neuropathic skin symptoms may include burning, electric-shock pain, altered sensation, numbness, or itch.

### **Neuropathic Pain**

Pain caused by disease or injury of the nerves that carry sensation, the spinal cord, or the brain. It is often burning, shooting, electric, or accompanied by numbness and may follow diabetes, shingles, injury, or nerve compression.

### **Neuropathic Ulcer**

A wound caused or worsened by loss of protective feeling, most often in the foot. A person may not notice pressure or injury, allowing the wound to deepen; diabetes, poor circulation, and infection can delay healing and increase the risk of amputation.

### **Neuropathy**

Disease or damage affecting nerves outside the brain and spinal cord. It may cause burning pain, numbness, tingling, weakness, balance problems, or altered sweating and can result from diabetes, medicines, toxins, poor nutrition, compression, or inherited disease.

### **Neuropeptide**

A peptide released by neurons that acts as a signal-

ing molecule in the nervous or endocrine system.

### **Neuropeptides**

Short chains of amino acids released by nerve cells as chemical signals. They can influence pain, appetite, stress, mood, sleep, blood pressure, inflammation, hormone release, and communication between nerve cells.

### **Neuropharmacology**

The study of how medicines and other substances affect the brain, spinal cord, and nerves. It examines their actions, benefits, side effects, interactions, and uses in treating neurological or psychiatric conditions.

### **Neurophysiology**

The study of how the nervous system functions, including electrical signaling, sensory processing, movement, reflexes, and communication between nerve cells. Clinical neurophysiology uses tests such as EEG and nerve-conduction studies to assess nervous-system function.

### **Neurophysiotherapy**

Physical therapy focused on movement and function problems caused by disorders of the brain, spinal cord, or nerves. Treatment may address strength, balance, walking, coordination, and daily activities.

### **Neuroplasticity**

The brain's ability to change its connections and activity in response to learning, experience, or injury. This process supports recovery and skill development, especially with repeated practice and rehabilitation, but recovery varies with the injury and the person.

### **Neuroplegic**

Causing paralysis or severe loss of nerve-controlled movement. The effect may result from a neurologic disease, spinal injury, toxin, or medicine and can impair breathing or other vital functions when major nerve pathways are affected.

---

## Neuropsychiatry

The clinical field linking neurologic and psychiatric disorders, including cognitive and behavioral consequences of brain disease.

## Neuropsychopharmacology

The study of how medicines and other substances alter brain function and behavior, especially in neurological and psychiatric disorders. It links drug actions on neural systems with therapeutic effects and adverse effects.

## Neuroradiology

The use of imaging to diagnose disorders of the brain, spine, head and neck, and peripheral nerves.

## Neuroretinitis

Inflammation of the optic nerve and nearby retina that can cause blurred or reduced vision, blind spots, and pain with eye movement. Infections and immune disorders are possible causes and require specialist evaluation.

## Neurosarcoidosis

Sarcoidosis affecting the brain, spinal cord, coverings around them, or peripheral nerves. It may cause headaches, facial weakness, vision problems, seizures, weakness, numbness, or hormone disturbances and usually needs specialist treatment.

## Neuroscience

The scientific study of the nervous system, including its structure, function, development, diseases, and treatment.

## Neurosciences

Neurosciences are the scientific disciplines studying the nervous system, including its structure, function, development, disease, and treatment.

## Neurosurgeon

A surgeon trained to diagnose and operate on disorders of the brain, spinal cord, spine, nerves, and nearby supporting structures. Neurosurgeons may treat tumors, injuries, bleeding, pressure, pain, or

blood-vessel problems.

## Neurosurgery

Surgery for disorders of the brain, spinal cord, spine, peripheral nerves, or supporting structures. Procedures may remove tumors, relieve pressure, repair injuries, stop bleeding, treat blood vessels, or improve severe pain.

## Neurosyphilis

Neurosyphilis is infection of the nervous system by *Treponema pallidum* and can cause meningitis, stroke, sensory loss, cognitive change, or tabes dorsalis.

## Neurotoxicity

Injury or dysfunction of the nervous system caused by a medicine, drug, toxin, heavy metal, infection, or other harmful exposure.

## Neurotransmission

The process by which nerve cells communicate with one another or with muscles and glands using electrical signals and chemical messengers. It involves release of a neurotransmitter, movement across a synapse, receptor binding, and termination or removal of the signal.

## Neurotransmitter

A chemical messenger released by a nerve cell. It crosses a tiny gap to affect another nerve cell or a muscle or gland cell, either increasing or reducing its activity.

## Neurotransmitter Receptors

Proteins on or inside cells that receive chemical messages released by nerve cells. When a message attaches to a receptor, it can increase or decrease activity, affecting movement, mood, pain, heart rate, or other body functions.

## Neurotrophic

Describing a substance or effect that supports the growth, survival, connection, or repair of nerve cells. Neurotrophic signals help nerves develop and main-

---

tain function and are studied in nerve injury and degenerative disease.

### **Neurotrophic Keratopathy**

Corneal damage caused by reduced sensation from injury or disease of the trigeminal nerve. It can lead to poor healing, recurrent epithelial defects, ulceration, and vision loss, sometimes with little pain.

### **Neurotropic**

Having a preference for nervous tissue or mainly affecting the brain, spinal cord, or nerves. The term is often used for infections, medicines, or other substances that act in these tissues.

### **Neurovascular Examination**

A check of movement, feeling, pulses, skin color, temperature, and blood return in an injured arm or leg. Repeated checks can detect worsening nerve damage, reduced blood flow, or dangerous swelling and pressure.

### **Neutral Alignment**

A posture in which the body is arranged in a straight, balanced line while preserving normal spinal curves.

### **Neutral Endopeptidase**

A membrane-bound zinc metallopeptidase expressed in vascular endothelial and renal proximal tubular cells that degrades circulating natriuretic peptides (ANP, BNP, CNP), bradykinin, and substance P; pharmacologically inhibited by sacubitril in heart failure therapy.

#### **Alternate Names:**

Neprilysin; NEP; CD10

### **Neutralization**

Removal or reduction of an effect by counteraction, such as an acid-base reaction or antibody-antigen binding.

### **Neutralizing Antibodies**

Antibodies that bind a pathogen or its toxin and block its ability to enter cells, attach to tissues, or

cause its usual biologic effects.

### **Neutropenia**

An abnormally low number of neutrophils, white blood cells that help fight bacterial and fungal infections. The lower the count, the greater the infection risk; fever during severe neutropenia can signal a serious infection.

#### **Alternate Names:**

Neutrocytopenia; Low Absolute Neutrophil Count

### **Neutropenic Enterocolitis**

A life-threatening, necrotizing transmural inflammation of the cecum and ascending colon occurring in severely neutropenic oncology patients receiving intensive myelosuppressive chemotherapy, presenting with fever, right lower quadrant abdominal pain, and bowel wall thickening on CT.

#### **Alternate Names:**

Typhlitis

### **Neutropenic Fever**

A medical emergency in oncology patients defined as a single oral temperature of 38.3 degrees Celsius (101 F) or sustained temperature of 38.0 degrees Celsius for one hour in the setting of an absolute neutrophil count (ANC) below 500 cells/microliter, requiring immediate empiric broad-spectrum antipseudomonal antibiotic administration.

#### **Alternate Names:**

Febrile Neutropenia

### **Neutrophil**

The most common infection-fighting white blood cell in the bloodstream. It quickly enters injured tissue, surrounds and destroys many bacteria and fungi, and helps start inflammation; unusually low levels increase infection risk.

### **Neutrophil Extracellular Traps**

Webs of extracellular decondensed chromatin fibers decorated with antimicrobial granule proteins (myeloperoxidase, elastase, histones) extruded by activated neutrophils during a specialized form of cell death (NETosis) to trap and kill extracellular bacterial pathogens; implicated in thrombosis and



---

autoimmune disease.

**Alternate Names:**  
NETs

### Neutrophilic Dermatositis

A group of inflammatory skin diseases in which infection-fighting white blood cells collect in the skin without causing a typical skin infection. Examples include Sweet syndrome and pyoderma gangrenosum; the rash may be painful, swollen, or associated with fever.

### Neutrophilic Eccrine Hidradenitis

An uncommon skin inflammation in which infection-fighting white cells collect around sweat glands. It often occurs during chemotherapy and causes painful red or purple bumps or patches, usually on the trunk or limbs.

### Neutrophil Recruitment

The movement of neutrophils from the bloodstream into injured or infected tissue. They first slow and attach to the blood-vessel lining, then pass through it and follow chemical signals to the affected area.

### Neutrophils

The most abundant type of white blood cell. Neutrophils rapidly migrate to sites of infection or injury and ingest microorganisms and cellular debris.

### Nevi of Ota and Ito

Benign dermal melanocytic nevi causing blue-gray pigmentation around the eye and forehead in nevus of Ota or over the shoulder and upper back in nevus of Ito. They may persist for life; ocular examination and monitoring of changing lesions are appropriate.

### Nevoid Basal Cell Carcinoma Syndrome

Gorlin syndrome, an inherited disorder increasing the risk of basal-cell cancers, jaw cysts, and other developmental abnormalities.

### Nevus

A usually harmless, localized change or growth in the skin or mucous membrane, often called a mole.

It may contain pigment-producing cells or involve blood vessels or other skin structures; a changing lesion needs assessment.

### Nevus Anemicus

A congenital vascular anomaly presenting as a localized, well-circumscribed, pale, hypopigmented macule or patch that remains pale after rubbing or temperature changes due to localized hypersensitivity of cutaneous alpha-adrenergic receptors producing constant arteriolar vasoconstriction.

**Alternate Names:**

Anemic Nevus; Pharmacologic Vasoconstrictive Nevus

### Nevus Araneus (Spider Nevus)

A small central red vessel with radiating branches that blanch with pressure and refill from the center. It is common in children, pregnancy, and liver disease; numerous or new lesions may prompt assessment for an underlying condition.

### Nevus Comedonicus

A congenital hamartoma of the pilosebaceous unit presenting as a localized cluster of dilated follicular ostia filled with keratinous plugs, resembling closely grouped open comedones. It typically appears at birth or in early childhood on the face, neck, or trunk, and may be associated with other developmental anomalies.

### Nevus Comedonicus Syndrome

A rare disorder in which multiple comedone-like skin lesions occur with abnormalities of the eyes, bones, nervous system, or other organs. It is usually present from birth or early childhood.

### Nevus of Ito

A benign dermal melanocytic hamartoma characterized by mottled, patchy slate-gray, blue-brown, or hyperpigmented macules distributed along the sensory branches of the posterior supraclavicular and lateral cutaneous brachial nerves (involving the supraclavicular, acromial, and scapular regions).

**Alternate Names:**

Omohyoid Nevus; Dermal Melanocytosis of Shoul-

---

der

### **Nevus of Ota**

A congenital or acquired dermal melanocytosis presenting as mottled blue-gray or brownish-black pigmentation in the distribution of the ophthalmic (V1) and maxillary (V2) branches of the trigeminal nerve, commonly involving the ipsilateral sclera, conjunctiva, and facial skin.

#### **Alternate Names:**

Oculodermal Melanocytosis; Nevus Fusco-Caeruleus Ophthalmomaxillaris

### **Nevus Sebaceus**

A congenital or early-onset hamartoma of sebaceous and other adnexal structures, usually a yellow-orange hairless plaque on the scalp or face; secondary tumors may rarely arise within it.

### **Newborn**

A baby during the first 28 days after birth, when special newborn health monitoring and care are important.

### **Newborn Circumcision**

Surgical removal of the foreskin from the penis, typically performed within the first few days of life. Medical benefits include reduced risk of urinary tract infections, sexually transmitted infections, and penile cancer. Complications include bleeding, infection, and meatal stenosis.

### **Newborn Head Molding**

Temporary shaping or overlapping of a newborn's skull bones during passage through the birth canal; it usually resolves as the skull returns to its normal contour.

### **Newborn Hearing Screen**

A quick test, usually before discharge from the birth facility, that checks whether a newborn responds normally to sound. A result needing repeat testing does not prove hearing loss; timely follow-up is important when the screen is not passed.

### **Newborn Infant Development**

The normal physical, neuromotor, sensory, and social milestone progression of a healthy full-term infant during the first month of life (neonatal period), including primitive reflexes (Moro, rooting, suckling, palmar grasp), visual tracking, auditory responsiveness, and weight recovery.

#### **Alternate Names:**

Neonatal Development; Infant Milestone Progression; Newborn Neuromotor Development

### **Newborn Jaundice**

Newborn jaundice is yellow discoloration from elevated bilirubin; physiologic patterns are common, but early, severe, rapidly rising, or conjugated jaundice requires evaluation for serious disease.

### **Newborn Metabolic Screen**

A dried blood spot test (heel stick) collected 24-48 hours after birth to screen for congenital disorders before symptoms develop.

### **Newborn Screening**

Testing shortly after birth for selected treatable genetic, metabolic, hormone, hearing, and heart conditions before symptoms appear. A small blood sample is usually taken from the heel; the conditions screened and timing vary by country or region.

### **Newborn Screening Panel**

A standardized public health laboratory testing program performed on dried blood spots collected via neonatal heel prick within 24 to 48 hours of life, screening for early, treatable inborn errors of metabolism, hemoglobinopathies, and endocrine disorders (such as phenylketonuria, congenital hypothyroidism, cystic fibrosis, and sickle cell disease).

### **Newborn Screening Tests**

Tests performed shortly after birth to detect selected treatable genetic, endocrine, metabolic, hematologic, hearing, or critical congenital heart conditions before symptoms develop.

---

### New Daily Persistent Headache

A headache that begins on a clearly remembered day and becomes continuous within 24 hours, persisting for at least three months. It may resemble migraine or tension-type headache; a new persistent headache requires medical evaluation to exclude secondary causes such as infection, pressure disorders, or vascular disease.

### New York Heart Association Functional Classification

A standardized clinical grading scale assessing the severity of functional symptomatic limitation in patients with heart failure based on physical exertion, ranging from Class I (no symptoms with ordinary physical activity) to Class IV (symptoms present at rest).

#### Alternate Names:

NYHA Functional Classification; NYHA Class

### Nexium

Brand name; see *Esomeprazole*.

### Next-Generation Sequencing

A laboratory method that reads many DNA sections at the same time. It can examine selected genes, nearly all protein-coding genes, the whole genome, or tumor DNA; results may miss some changes and require expert interpretation.

### Nexus

A connection or central point linking structures, processes, or ideas. In medical writing, its meaning depends on the context and may describe an anatomic connection, a group of related causes, or a link between two findings.

### NF1

Abbreviation; see *Neurofibromatosis Type 1*.

### NF2

Abbreviation; see *Neurofibromatosis Type 2*.

### NF2-Related Schwannomatosis

A genetic tumor-predisposition disorder, usually in-

volving pathogenic variants in NF2, characterized by schwannomas, especially bilateral vestibular schwannomas, with possible hearing loss, tinnitus, balance problems, meningiomas, or other nerve tumors.

### Niacin

Vitamin B3, needed to help cells release energy and perform many chemical reactions; it forms the coenzymes NAD<sup>+</sup> and NADP<sup>+</sup> that carry electrons in metabolism. Severe deficiency causes pellagra, with skin changes, digestive problems, confusion, weakness, and potentially life-threatening complications.

#### Alternate Names:

Vitamin B3; Nicotinic Acid

### Niacinamide

The amide form of vitamin B3, used as a nutritional supplement and in selected dermatologic preparations. Unlike niacin, it usually does not cause flushing at nutritional doses.

### Niacin Deficiency

A lack of vitamin B3, usually from a diet poor in niacin and the amino acid tryptophan. When severe it causes pellagra, with dermatitis, diarrhea, and neuropsychiatric disturbance, and it may complicate alcohol use disorder or conditions that impair absorption. Niacin replacement and treatment of the underlying cause correct the deficiency.

### Niacin For Cholesterol

Niacin lowers triglycerides and can raise HDL cholesterol, but its routine cardiovascular benefit is limited when added to effective statin therapy; flushing, liver injury, hyperglycaemia, and hyperuricaemia limit high-dose use.

### Niacin Vitamin B3

Alternate name; see *Niacin*.

### Nialamide

An older monoamine-oxidase-inhibitor antidepressant that is no longer commonly used.

---

## Nicardipine

A calcium-channel-blocking medicine that relaxes blood vessels and lowers blood pressure. It is used for selected urgent or ongoing blood-pressure problems and may cause headache, flushing, dizziness, fast heartbeat, or swelling.

## Nickel Allergy

An allergic skin reaction caused by contact with nickel-containing materials.

### Alternate Names:

Nickel Hypersensitivity; Nickel Contact Dermatitis

## Nickel-Chromium

A metal mixture used in some dental crowns and medical devices. It is strong and less costly than some alternatives, but people allergic to nickel may develop skin or mouth reactions after contact.

## Niclosamide

An antiparasitic medicine used in some countries to treat selected intestinal tapeworm infections. It acts mainly in the gut and does not reliably treat larval tapeworm infections in tissues, such as neurocysticercosis.

## Nicolau Syndrome

A rare, acute iatrogenic cutaneous adverse reaction characterized by severe localized pain, livedoid erythema, and rapid ischemic skin necrosis (embolia cutis medicamentosa) occurring at the site of intramuscular or intra-articular drug injections, caused by accidental intra-arterial or peri-arterial drug deposition and microvascular thrombosis.

### Alternate Names:

Embolia Cutis Medicamentosa; Livedo-like Dermatitis

## Nicotinamide

Alternate name; see *Niacinamide*.

## Nicotinamide Adenine Dinucleotide

A molecule present in all living cells that helps transfer energy from food and supports enzymes involved in cell repair and communication. It exists

in forms commonly written as NAD<sup>+</sup> and NADH.

## Nicotine

An addictive stimulant found in tobacco and some vaping products that acts on neuronal nicotinic receptors.

## Nicotine Addiction

Alternate name; see *Nicotine Dependence*.

## Nicotine And Tobacco

Nicotine poisoning can follow swallowing tobacco or liquid nicotine, breathing concentrated vapor, or skin contact. It may cause nausea, vomiting, sweating, dizziness, fast heartbeat, seizures, or breathing failure.

## Nicotine Dependence

A pattern of compulsive nicotine use with cravings, loss of control, tolerance, and withdrawal symptoms when use is reduced or stopped.

### Alternate Names:

Tobacco Use Disorder; Nicotine Addiction

## Nicotine Poisoning

Nicotine poisoning causes early nausea, vomiting, salivation, dizziness, tachycardia, and agitation followed by weakness, bradycardia, seizures, or respiratory failure in severe exposure.

## Nicotine Replacement Therapy

Nicotine medicines help reduce withdrawal when stopping tobacco. Using too much or combining products improperly can cause nausea, vomiting, dizziness, sweating, fast heartbeat, confusion, or seizures; keep them away from children.

## Nicotinic

Relating to nicotine or to nicotinic acetylcholine receptors. These receptors respond to the chemical messenger acetylcholine and help control nerve signals, skeletal-muscle movement, attention, and some automatic body functions.

---

## Nicotinic Acetylcholine Receptor

A pentameric ligand-gated non-selective cation channel responsive to acetylcholine and nicotine, functioning to generate rapid excitatory postsynaptic potentials at the neuromuscular junction (muscle type) and throughout autonomic ganglia and the central nervous system (neuronal type).

### Alternate Names:

nAChR

## Nicotinic Acid

Alternate name; see *Niacin*.

## NICU Staff

NICU staff are the healthcare professionals who monitor and treat newborns in a neonatal intensive care unit and support their families.

## Nidation

Implantation of a fertilized egg into the lining of the uterus early in pregnancy. The term is largely synonymous with implantation.

## Nidus

A starting point or central focus from which a process develops. In medicine, it may mean the core of a kidney stone, the focus of an infection, or the central part of an abnormal blood-vessel network.

## Niemann-Pick Disease

A group of rare inherited lysosomal storage disorders in which lipids accumulate in cells. The group includes acid sphingomyelinase deficiency, formerly called Niemann-Pick disease types A and B, and Niemann-Pick disease type C, caused by impaired intracellular cholesterol trafficking. Features vary by type and may include enlargement of the liver or spleen, lung disease, difficulty coordinating movement, and progressive neurologic impairment.

## Nifedipine

A medicine that relaxes blood vessels to lower blood pressure and relieve some types of chest pain. It may cause flushing, headache, dizziness, ankle swelling, or a fast heartbeat; the dose depends on the condi-

tion and formulation.

## Night Blindness

Difficulty seeing in dim light or darkness. Causes include vitamin-A deficiency, inherited retinal disorders, cataracts, glaucoma, and some medicines. New or worsening night blindness needs an eye examination because some causes can be treated.

## Night-Blindness

Difficulty seeing in dim light, called nyctalopia; causes include vitamin A deficiency, retinal disease, and inherited disorders.

## Night Eating Syndrome

A recurring pattern of eating much of the day's food in the evening or after waking at night, often with little morning appetite and sleep problems. It is a related eating pattern that may occur with depression, obesity, or other eating disorders and is not automatically a formal eating-disorder diagnosis.

## Night Leg Cramps

Sudden, painful involuntary contractions of leg muscles occurring during sleep or rest, often affecting the calf or foot.

## Nightmare Disorder

Repeated frightening dreams that cause distress, wake a person, or interfere with sleep and daytime activities. It may be linked with stress, trauma, medicines, mental health conditions, or other sleep disorders.

## Nightmares

Vivid frightening dreams that cause distress or wake a person, sometimes linked to stress, illness, or medicines.

## Night Sweats

Episodes of excessive sweating during sleep that soak clothing or bedding. They may occur with infection, hormonal changes, cancer, medication effects, or other systemic illness; persistent unexplained night sweats warrant assessment.

---

**Alternate Names:**

Sleep Hyperhidrosis; Nocturnal Diaphoresis

**Night-Sweats**

Episodes of excessive sweating during sleep; causes include infection, malignancy, hormonal change, medicines, and environmental heat.

**Night Sweats in Men**

Alternate name; see *Night Sweats*.

**Night Terrors**

Episodes of abrupt partial awakening from deep non-REM sleep with intense fear, screaming, autonomic arousal, and limited recall afterward, occurring most often in children.

**Night Terrors In Children**

Night terrors are episodes of sudden fear, screaming, autonomic arousal, and apparent confusion during non-REM sleep; children usually remain difficult to awaken and have little memory afterward.

**NIH Stroke Scale**

The National Institutes of Health Stroke Scale is a standardized, validated clinical assessment tool used to quantify the severity of acute ischemic stroke.

**Nijmegen Breakage Syndrome**

A rare inherited disorder that affects DNA repair and the immune system. It may cause a small head, short stature, developmental or learning difficulties, frequent infections, and a high risk of lymphoma or other cancers. People with it are unusually sensitive to radiation, so genetic and specialist care is important.

**Alternate Names:**

NBS; Ataxia-Telangiectasia Variant 1

**Nikethamide**

An old medicine once used to stimulate breathing. It is no longer a routine treatment because its benefit is limited and excessive doses can cause agitation, shaking, seizures, or other dangerous effects.

**Nikolsky Sign**

Peeling or separation of the top skin layer when nearby skin is gently rubbed. It can occur in serious blistering disorders, but the sign is not specific and must be interpreted with the rash and other findings.

**Nikolsky's Sign**

Shearing or peeling of the superficial epidermis when apparently normal skin beside a blister is rubbed; it may occur in pemphigus and other blistering disorders.

**Nipah Virus**

A zoonotic virus in the genus *Henipavirus*, with fruit bats as its natural host. It can spread from infected bats or other animals to people, through contaminated food such as raw date-palm sap, and between people through close contact. Infection ranges from no symptoms to fever, headache, vomiting, and respiratory illness such as cough or difficulty breathing. Severe disease may cause inflammation of the brain (encephalitis), confusion, seizures, coma, or death. There is currently no licensed vaccine or specific approved treatment for Nipah virus disease; management relies on supportive care.

**Nipah Virus Infection**

A zoonotic illness caused by Nipah virus. Fruit bats are the natural host, and infection can pass to people through infected animals, contaminated food such as raw date-palm sap, or close contact with an infected person. Some infections cause no symptoms; illness may otherwise cause fever, headache, muscle pain, vomiting, cough, or difficulty breathing and can progress to brain inflammation (encephalitis), confusion, seizures, coma, or death. Early intensive supportive care may improve survival, but there is currently no licensed vaccine or specific approved treatment.

**Nipple**

The projection at the centre of the breast through which milk is released from the mammary ducts; it contains smooth muscle and sensory nerve endings.



---

## Nipple-Areola Complex

The nipple together with the darker skin around it, called the areola. It contains ducts, smooth muscle, glands, and sensory nerves.

## Nipple Discharge

Fluid leaking from one or both nipples when not pregnant or breastfeeding. Most nipple discharge is harmless, coming from multiple milk ducts in both breasts only when squeezed, and is usually caused by normal hormone changes or medications. Discharge that leaks on its own from a single duct in one breast, especially if clear or bloody, is less common. The most frequent cause is a small, harmless growth inside a milk duct (a papilloma), while breast cancer accounts for about 5% to 15% of these cases. Doctors identify the cause using a physical examination, imaging such as ultrasound or mammography, and duct tests.

## Nipple Inversion Or Retraction

Inward turning or pulling-in of the nipple. It may be longstanding and benign, but new one-sided inversion or retraction can occur with duct disease or breast cancer.

## Nipple Retraction

Alternate name; see *Nipple Inversion Or Retraction*.

## Niridazole

An older medicine once used against schistosomiasis, a parasitic infection. Its use is now limited because of serious possible side effects, including seizures, liver injury, and damage to blood-cell production.

## Nissen Fundoplication

A surgical antireflux procedure in which the gastric fundus is completely wrapped 360 degrees around the lower esophagus to reinforce the lower esophageal sphincter (LES) and repair a hiatal hernia, used for refractory gastroesophageal reflux disease (GERD).

### Alternate Names:

360-Degree Complete Fundoplication; Total Antireflux Fundoplication

## Nit

The egg of a louse, usually attached firmly to a hair shaft or clothing fiber. Nits may be confused with dandruff, but they do not brush away easily. Finding live lice confirms an active infestation; treatment and fine combing remove lice and eggs.

## Nitabuch Layer

A normal, dense, fibrinoid layer formed at the junction between the invading syncytiotrophoblast and the decidua basalis (stratum compactum) in normal pregnancy, which acts as a physiological anatomical boundary limiting deep placental invasion (congenitally deficient in placenta accreta).

### Alternate Names:

Stria of Nitabuch; Decidual Fibrinoid Boundary

## Nitrates

Medicines that widen blood vessels by releasing nitric oxide. They can relieve angina and lower the heart's workload but may cause headache, flushing, dizziness, or dangerously low blood pressure with certain medicines.

## Nitrate Tolerance

Reduced effect of nitrate medicines, such as nitroglycerin, after continuous use. It may involve changes in blood-vessel signaling and the body's counter-response to the medicine.

### Alternate Names:

Nitroglycerin Tachyphylaxis; Nitrate Resistance

## Nitrazepam

A benzodiazepine medicine that slows brain activity and can cause sleepiness, calmness, and seizure prevention. It can cause dependence, falls, memory problems, and dangerously slow breathing, especially with alcohol or opioids.

## Nitric Acid

A strong corrosive chemical that can burn skin, eyes, mouth, throat, and digestive organs. Breathing its fumes can injure the lungs, and swallowed or inhaled exposure can cause severe poisoning.

---

## Nitric Acid Poisoning

Exposure to nitric acid can severely burn the skin, eyes, mouth, throat, and stomach and may damage the lungs if fumes are inhaled.

## Nitric Oxide

A short-lived signaling molecule made by many cells. It relaxes blood-vessel muscle, helping regulate blood flow and pressure, and also participates in nerve signaling and immune defense. Too much or too little production may contribute to disease.

## Nitric Oxide Synthase

An enzyme family that catalyzes the synthesis of the gaseous signaling molecule nitric oxide (NO) and L-citrulline from L-arginine, oxygen, and NADPH, comprising constitutive calcium-dependent isoforms (endothelial eNOS, neuronal nNOS) and a calcium-independent inducible isoform (iNOS).

**Alternate Names:**

NOS

## Nitrofurantoin

An antibacterial medicine used mainly for bladder infections caused by susceptible bacteria. It concentrates in urine but is not suitable for many kidney infections; nausea, lung reactions, liver injury, or nerve problems are uncommon but important risks.

## Nitrogen

An essential element in amino acids, proteins, nucleic acids, and other biological molecules; atmospheric nitrogen is largely inert to humans.

## Nitrogen Balance

The difference between dietary nitrogen intake, estimated mainly from protein intake, and nitrogen loss, primarily in urine. Positive balance supports growth or tissue repair; negative balance occurs with inadequate intake, inflammation, trauma, or catabolic disease.

## Nitrogenous Base

A nitrogen-containing component of nucleic acids; adenine, guanine, cytosine, thymine, and uracil pair

in DNA or RNA to encode genetic information.

## Nitroglycerin Overdose

Too much nitroglycerin can cause severe headache, flushing, dizziness, fainting, low blood pressure, fast heartbeat, or breathing difficulty.

## Nitromors

A trade name for paint or varnish stripper formulations; exposure can cause chemical burns or toxic inhalation.

## Nitrosoureas

A group of chemotherapy medicines that can cross the blood-brain barrier and damage DNA; examples include carmustine and lomustine.

## Nitrous Oxide

An inhaled anesthetic gas with analgesic and anesthetic-sparing properties. It has rapid onset and offset but relatively low potency, expands closed gas spaces, and may contribute to diffusion hypoxia or postoperative nausea.

## Nitrous Oxide For Labor

A 50:50 mixture of nitrous oxide and oxygen (Entonox) inhaled by the mother during labor for analgesia. Self-administered via a mask or mouthpiece; the woman controls her own dosing, taking a breath 30 seconds before a contraction.

## Nizatidine

A medicine that reduces stomach-acid production by blocking histamine signals in stomach cells. It may relieve ulcers or reflux symptoms, although availability varies and treatment should consider kidney function and medicine interactions.

## NMR

NMR, or nuclear magnetic resonance, measures how atomic nuclei respond to a magnetic field and is used in chemical analysis and imaging.

## Nocardia Asteroides Infection

An infection caused by *Nocardia asteroides*, a soil-

---

associated bacterium. It most often affects the lungs, skin, or brain and is more likely when immunity is weakened; treatment requires species identification and prolonged antibiotics.

### **Nocardia Infection**

An infection caused by *Nocardia* bacteria, usually affecting the lungs, skin, or brain, especially in people with weakened immunity.

### **Nocardiosis**

An infection caused by *Nocardia* bacteria found in soil and water. It most often affects the lungs, skin, or brain, especially in people with weakened immunity, and may cause cough, fever, skin sores, confusion, or seizures.

### **Nocebo**

A harmful or unpleasant symptom or treatment effect caused or amplified by negative expectations rather than by the treatment's specific pharmacologic action.

### **Nociception**

The nervous system's detection and processing of potentially harmful heat, pressure, or chemicals. It can lead to pain, but pain also depends on how the brain interprets these signals.

### **Nociceptor**

A sensory nerve ending that detects potentially damaging pressure, heat, cold, or chemicals. It sends warning signals through nerves to the spinal cord and brain, where they can be experienced as pain.

### **Nociceptors**

Sensory nerve endings that detect potentially damaging pressure, heat, cold, or chemicals. They send warning signals through the nervous system, although pain also depends on how the brain interprets those signals.

### **Nocturia**

Waking during the night one or more times to urinate and then returning to sleep. It may result

from evening fluids, medicines, bladder or prostate problems, diabetes, heart disease, or sleep disorders.

### **Nocturnal**

Occurring at night or during sleep. Examples include bedwetting, nighttime cough, breathing pauses during sleep, nighttime low oxygen, or breathlessness that wakes a person. The cause depends on the symptom and requires assessment when persistent or severe.

### **Nocturnal Emission**

Involuntary ejaculation during sleep, sometimes called a wet dream. It is a common and normal event during male puberty and adolescence.

### **Nocturnal Enuresis**

Repeated involuntary urination during sleep in a child who is old enough to have developed nighttime bladder control. It may run in families; pain, daytime wetting, constipation, or sudden onset needs evaluation.

#### **Alternate Names:**

Nocturnal Incontinence; Sleep Enuresis

### **Nocturnal Enuresis Entry**

Alternate name; see *Bed-Wetting*.

### **Nocturnal Hypoventilation**

A reduction in minute ventilation occurring specifically during sleep, resulting in nighttime hypercapnia and hypoxemia, commonly encountered in neuromuscular diseases (such as ALS or muscular dystrophies), severe kyphoscoliosis, and obesity hypoventilation syndrome.

### **Nocturnal Incontinence**

Alternate name; see *Incontinence*.

### **Nocturnal Lagophthalmos**

Incomplete closure of the eyelids during sleep, which may cause dryness, irritation, and exposure injury to the cornea.

---

## Nocturnal Polyuria

Production of an unusually large share of the day's urine during nighttime sleep. It can contribute to repeated nighttime urination and may be related to fluid shifts, sleep disorders, medicines, or other medical conditions.

## Nodal Rhythm

A heartbeat rhythm that starts near the atrioventricular node instead of the heart's usual pacemaker. It may occur when the normal pacemaker is slow or blocked and can cause a slow pulse, dizziness, or no symptoms.

## Node

A localized swelling, structure, or point of connection; examples include lymph nodes and electrical conduction nodes.

## Node of Ranvier

Periodic, unmyelinated gaps or interruptions occurring at regular intervals along the length of a myelinated nerve axon where high concentrations of voltage-gated sodium channels enable rapid saltatory nerve impulse conduction.

### Alternate Names:

Ranvier Node; Myelin Sheath Gap

## Nodular

Made up of or containing one or more rounded lumps called nodules. The word may describe a skin lesion, thyroid swelling, lung finding, or other tissue. Its importance depends on the nodule's size, appearance, location, growth, and associated symptoms.

## Nodular Fasciitis

A rapidly growing benign self-limited fibroblastic and myofibroblastic proliferation, usually presenting as a painful or tender superficial mass. It can mimic sarcoma clinically and microscopically.

## Nodular Localized Cutaneous Amyloidosis

A localized cutaneous amyloidosis presenting as one

or more firm, waxy, skin-colored or yellow-brown nodules. It is usually confined to the skin, but examination may be needed to distinguish it from systemic amyloidosis or other nodular disorders.

## Nodular Lymphoid Hyperplasia

Enlargement of small areas of immune tissue, most often in the small intestine. It is usually harmless but may be associated with reduced antibody production or intestinal infection; its significance depends on symptoms and the test findings.

## Nodular Melanoma

A type of skin cancer that grows downward early and often appears as a rapidly enlarging firm bump. It may be black, brown, pink, or skin-colored, so any quickly changing or bleeding lump needs prompt assessment.

## Nodule

A small, rounded area of abnormal tissue that may be solid or fluid-filled and may occur in the skin, lung, thyroid, or another organ. Its meaning depends on location, size, appearance, growth, and symptoms; some nodules are harmless, while others need testing or follow-up.

## Noise-Induced Hearing Loss

Noise-induced hearing loss is permanent injury to inner-ear hair cells from repeated or intense sound exposure, often with tinnitus or difficulty understanding speech. Limiting exposure, increasing distance, taking quiet breaks, and using well-fitted hearing protection reduce risk.

## Noise-Related Hearing Loss

Alternate name; see *Hearing Loss*.

## Noma

Noma is a rapidly progressive gangrenous infection of the mouth and face, usually in severely malnourished or immunocompromised children. It destroys oral and facial tissue and is treated with antibiotics, nutritional support, and sometimes surgery.

---

## Nomenclature

Nomenclature is the organized system used to name and classify diseases, organisms, chemicals, anatomy, medicines, or other entities.

## Nomogram

A graphical tool that estimates a quantity or risk from related measurements. The Rumack–Matthew nomogram is used in selected cases of acute acetaminophen ingestion.

## Nonaccidental Injury

An injury caused deliberately or by neglect rather than by an accident. In children, suspicious patterns may include injuries inconsistent with the history, injuries at different stages of healing, or injuries in protected areas.

## Non-Accidental Trauma

An injury caused deliberately or through abuse rather than by an accident. Clinicians consider it when the explanation does not fit the injuries, when injuries are repeated or occur at different healing stages, or when a child cannot explain what happened; suspected cases require safeguarding assessment.

## Non-Adherence

Failure to follow an agreed treatment plan, which may result from access barriers, side effects, beliefs, misunderstanding, or other circumstances.

## Non-Alcoholic Steatohepatitis

An aggressive, progressive form of metabolic dysfunction-associated steatotic liver disease (MASLD), now called metabolic dysfunction-associated steatohepatitis (MASH), characterized by liver fat with inflammation and cell injury, with or without scarring. It can progress to cirrhosis and liver cancer.

### Alternate Names:

NASH; Metabolic Dysfunction-Associated Steatohepatitis (MASH)

## Nonalcoholic Steatohepatitis Pathology

Steatotic liver disease with hepatocyte ballooning, lobular inflammation, and variable perisinusoidal or portal fibrosis, usually in the absence of a history sufficient to explain the injury by alcohol alone. The modern term MASH is also used.

## Non-Allergic Rhinitis

A chronic nasal syndrome characterized by sneezing, rhinorrhea, nasal congestion, and post-nasal drip without clinical or laboratory evidence of IgE-mediated allergic sensitization, frequently triggered by weather changes, cold dry air, chemical odors, or foods (vasomotor rhinitis).

### Alternate Names:

Vasomotor Rhinitis; NAR

## Nonallergic Rhinitis

Nasal congestion, sneezing, or runny nose caused by irritants or other triggers rather than an allergic immune response. Triggers may include changes in temperature or humidity, smoke, odors, medicines, or hormonal changes.

### Alternate Names:

Vasomotor Rhinitis; Idiopathic Rhinitis

## Nonallergic Rhinopathy

Nonallergic rhinopathy causes congestion, runny nose, or sneezing without an allergic immune reaction, often triggered by odors, smoke, weather changes, or irritants.

## Non-Articular Rheumatism

An older term for painful disorders of muscles, tendons, bursae, or other tissues outside the joints.

## Nonarticular Rheumatism/Regional Pain Syndrome

A broad clinical category of pain arising outside the joints, including disorders of muscles, tendons, bursae, fascia, nerves, or other regional soft tissues. The cause may be overuse, injury, inflammation, nerve dysfunction, or altered pain processing.



---

### Non-Asthmatic Eosinophilic Bronchitis

A cause of long-lasting cough in which certain white blood cells inflame the airways without the variable narrowing seen in asthma. Breathing tests may be normal, but an inhaled corticosteroid can reduce the cough when this diagnosis is confirmed.

### Nonbacterial Prostatitis

Chronic pelvic pain syndrome (CP/CPPS) characterized by persistent pelvic, perineal, or genital pain and urinary symptoms for at least 3 months without evidence of active urinary tract bacterial infection on standard cultures.

#### Alternate Names:

Chronic Pelvic Pain Syndrome in Men; Abacterial Prostatitis; Category III Prostatitis

### Nonbacterial Thrombotic Endocarditis

Small, sterile blood clots that form on heart valves without infection. It is often linked to cancer, lupus, or another condition that makes blood clot easily. The clots can break off and cause stroke or blockage elsewhere, so treatment focuses on the cause and preventing further clots.

#### Alternate Names:

NBTE; Marantic endocarditis; Libman-Sacks-like nonbacterial endocarditis

### Noncandidal Fungal Infections of the Mouth

Fungal infections of the oral cavity caused by organisms other than *Candida*, including selected dermatophytes, molds, and endemic fungi. They may produce plaques, ulcers, swelling, or deep tissue disease and often require culture, biopsy, or systemic treatment.

### Non-Celiac Gluten Intolerance

Alternate name; see *Non-Celiac Gluten Sensitivity*.

### Non-Celiac Gluten Sensitivity

Symptoms attributed to gluten or wheat that improve after removal, despite testing not showing celiac disease or wheat allergy. Diagnosis requires careful medical evaluation because other foods and conditions can cause similar symptoms.

### Non-Communicable Disease

A chronic, non-transmissible medical condition that is not caused by acute infectious agents and typically progresses slowly over a prolonged duration. Arising from a multifactorial interplay of genetic, physiological, environmental, and behavioral factors, the four primary groups responsible for the vast majority of global mortality and disability are cardiovascular diseases (such as coronary artery disease, stroke, and hypertension), cancers, chronic respiratory diseases (such as COPD and asthma), and diabetes mellitus. Management and public-health prevention focus on early screening, long-term pharmacological control, and modifying shared behavioral risk factors, including tobacco use, physical inactivity, unhealthy diet, and harmful alcohol consumption.

#### Alternate Names:

NCD; Non-Communicable Diseases; Noncommunicable Disease; Chronic Non-Communicable Disease

### Non-Coding RNA (ncRNA)

An RNA molecule that is not translated into a protein. Non-coding RNAs can regulate gene activity, RNA processing, protein production, and chromosome function; abnormal regulation is involved in some cancers and other diseases.

### Non Communicating Hydrocephalus

A buildup of cerebrospinal fluid in the brain because a blockage prevents it from moving between the brain's fluid-filled spaces and reaching the area where it is absorbed. It can increase pressure on the brain and cause headache, vomiting, sleepiness, vision problems, or difficulty walking.

### Noncompetitive Antagonist

A medicine that reduces another medicine or natural chemical's effect by attaching to its receptor or signaling pathway in a way that extra amounts of the activating substance cannot fully overcome.

### Noncompetitive Inhibition

A form of enzyme inhibition in which a substance binds away from the enzyme's working site and reduces its activity. Because it does not compete directly with the usual substrate, adding more sub-



---

strate does not fully overcome the inhibition.

### **Non-Compliant Angioplasty Balloon**

A small balloon on a catheter that keeps its shape when inflated at high pressure to widen a narrowed blood vessel. It is used during selected vascular procedures and can rarely cause bleeding, vessel injury, or blockage.

#### **Alternate Names:**

High-Pressure Angioplasty Balloon; Non-Compliant Balloon

### **Non-Contact Tonometry**

A quick estimate of pressure inside the eye using a brief puff of air against the clear front surface of the eye. It does not touch the eye and can screen for glaucoma, but unusual results may need confirmation.

### **Non-Convulsive Status Epilepticus**

A prolonged state of continuous electrographic seizure activity lasting 30 minutes or longer characterized by altered mental status, confusion, coma, or subtle motor automatisms (such as eyelid fluttering) in the complete absence of generalized tonic-clonic motor convulsions, diagnosed via continuous EEG.

#### **Alternate Names:**

NCSE

### **Nonconvulsive Status Epilepticus**

Ongoing seizure activity without prominent rhythmic limb jerking, often presenting as prolonged confusion, impaired awareness, or subtle twitching. An electroencephalogram can confirm it, and treatment is urgent.

### **Nondisjunction**

Failure of chromosome copies to separate properly when a cell divides. This can produce cells with too many or too few chromosomes and may cause conditions such as Down syndrome, Turner syndrome, or some miscarriages.

### **Non-Fatal Drowning**

Survival after breathing is impaired by submersion

or immersion in liquid. A person may still develop lung injury, brain injury, or delayed breathing problems, even after apparently recovering.

### **Non-Functioning Pituitary Adenoma**

A usually noncancerous pituitary tumor that does not release enough active hormones to cause a hormone syndrome. As it enlarges, it may cause headache, reduced vision, or loss of normal pituitary function.

### **Nongenital Warts**

Common benign skin growths caused by cutaneous human papillomaviruses, usually appearing as rough papules on the hands, fingers, elbows, or knees; plantar warts affect the soles. They spread by direct contact or self-inoculation, may persist for months or years, and are distinct from anogenital warts.

### **Non-Hodgkin Lymphoma**

A group of cancers that begin in immune cells called lymphocytes. Some grow slowly and others quickly; they may cause swollen lymph nodes, fever, night sweats, weight loss, or symptoms in an affected organ.

#### **Alternate Names:**

NHL; B-Cell or T-Cell Lymphoma

### **Non-Hodgkin Lymphoma In Children**

A group of childhood cancers arising from immune cells called lymphocytes. Some grow quickly and can cause swollen nodes, fever, abdominal symptoms, or breathing problems; the exact type and spread determine treatment.

### **Non-Hodgkin's Lymphoma**

Alternate spelling; see *Non-Hodgkin lymphoma*.

### **Non-Hodgkins Lymphomas**

Alternate spelling and plural form; see *Non-Hodgkin lymphoma*.

### **Non-Infectious Cystitis**

Inflammation of the bladder without infection by bacteria or another microorganism. It may cause

---

pain, urgency, and frequent urination after irritants, radiation, medicines, chemical exposure, or other bladder conditions.

### **Noninfectious Cystitis**

A heterogeneous group of bladder inflammatory conditions occurring in the absence of active bacterial infection, encompassing radiation cystitis, chemotherapy-induced cystitis (e.g., cyclophosphamide), chemical cystitis, and interstitial cystitis.

#### **Alternate Names:**

Abacterial Cystitis; Sterile Cystitis; Chemical/Radiation Cystitis

### **Noninvasive**

Not involving a surgical cut, a procedure entering the body, or a device placed inside it. Examples include measuring blood pressure with a cuff, external imaging, and breathing support through a mask. Risks depend on the method used.

### **Non-Invasive Positive Pressure Ventilation**

The delivery of mechanical ventilatory support without an invasive endotracheal tube or tracheostomy, utilizing a tight-fitting facial or nasal mask to provide continuous positive airway pressure (CPAP) or bilevel positive airway pressure (BiPAP) in acute hypercapnic respiratory failure or cardiogenic pulmonary edema.

#### **Alternate Names:**

NIPPV; NPPV

### **Non-Invasive Prenatal Testing**

A blood test during pregnancy that estimates the chance of selected chromosome conditions, especially trisomy 21, 18, and 13, by analyzing small DNA fragments from the placenta. It is screening, not diagnosis; an abnormal result needs confirmatory testing.

### **Non-Invasive Procedure**

An examination or treatment that does not cut or pierce the skin or place an instrument inside the body. Examples include blood-pressure checks, ECGs, ultrasound, MRI, X-rays, and physical ther-

apy. These procedures are generally safer than invasive procedures, but each can still have limits or risks.

#### **Alternate Names:**

Noninvasive Procedure; Non-Invasive Medical Procedure; Noninvasive Test; Non-Invasive Technique

### **Non-Invasive Ventilation**

Breathing support delivered through a face or nose mask instead of a tube placed in the windpipe. Continuous or two-level air pressure can improve breathing in selected people with respiratory failure, sleep apnea, or fluid in the lungs.

### **Nonischemic Priapism**

Alternate name; see *Priapism*.

### **Nonketotic Hyperglycemic Hyperosmolar Syndrome**

A life-threatening diabetic emergency in which very high blood sugar causes profound dehydration and concentrated blood, usually without major ketone buildup. It can cause extreme thirst, weakness, confusion, seizures, coma, and organ failure.

### **Non-Ketotic Hyperosmolar State**

A life-threatening acute metabolic emergency occurring predominantly in type 2 diabetes mellitus, characterized by extreme hyperglycemia (glucose greater than 600 mg/dL), profound hyperosmolality (plasma effective osmolality greater than 320 mOsm/kg), and severe dehydration in the absence of significant ketoacidosis.

#### **Alternate Names:**

Hyperosmolar Hyperglycemic State; HHS

### **Nonmelanoma Skin Cancer**

A group of skin cancers that arise from the cells of the epidermis, excluding melanoma. The two most common types are basal cell carcinoma (BCC) and squamous cell carcinoma (SCC), which together account for the vast majority of all skin cancers diagnosed globally.

---

### **Non-Muscle-Invasive Bladder Cancer**

Bladder cancer limited to the inner lining or connective tissue beneath it, without invasion of the bladder muscle. It can recur and may progress to muscle-invasive disease.

### **Non-Obstructive Azoospermia**

Complete absence of sperm in semen because the testicles make too few or no sperm, rather than because a tube is blocked. Causes include genetic conditions, undescended testes, infections, cancer treatment, or an unknown problem. Testing may include hormone tests and genetic studies; some people can use surgically retrieved sperm for assisted reproduction.

#### **Alternate Names:**

NOA; Secretory Azoospermia; Testicular Failure Azoospermia

### **Nonpathogenic**

Nonpathogenic means not ordinarily capable of causing disease in a given host or setting; an organism can still cause opportunistic infection when immunity or barriers are impaired.

### **Non-Pitting Edema**

Swelling of subcutaneous tissue that does not leave a persistent indentation when firm digital pressure is applied, usually due to local accumulation of protein-rich fluid, mucopolysaccharides, or chronic cellular infiltration. Common causes include lymphedema, pretibial myxedema in Graves disease, and severe lipedema.

### **Non-Polio Enterovirus Infection**

Alternate name; see *Enterovirus*.

### **Non-Proliferative Diabetic Retinopathy**

An early form of diabetes-related damage to the retina, the light-sensitive tissue at the back of the eye. Small blood vessels may leak or swell, causing blurred vision, but abnormal new vessels have not formed.

### **Nonproliferative Retinopathy**

Retinal disease in which small blood vessels leak or become damaged without abnormal new vessels growing. Diabetes is a common cause; progression can impair vision, especially when the macula is affected.

#### **Alternate Names:**

Background Retinopathy; NPDR

### **Non-Radiographic Axial Spondyloarthritis**

Inflammatory arthritis affecting the spine or joints between the spine and pelvis without definite changes on a plain X-ray. It may cause persistent back pain and morning stiffness, and may occur with eye inflammation, psoriasis, or inflammatory bowel disease.

### **Non-Rapid Eye Movement Sleep**

Sleep that includes lighter and deeper stages without rapid eye movements. Deep stages support physical restoration, while lighter stages help the body transition into and out of sleep.

### **Non-Rebreather Mask**

A high-flow oxygen delivery device featuring an attached reservoir bag and one-way directional valves that prevent entrainment of room air and rebreathing of exhaled gas, capable of delivering inspired oxygen fractions (FiO<sub>2</sub>) between 80 and 95 percent at flow rates of 10 to 15 L/min.

#### **Alternate Names:**

NRB Mask

### **Non-REM Sleep**

Sleep stages without rapid eye movements, ranging from light sleep to deep slow-wave sleep. These stages support physical restoration, memory processing, and the normal transition between waking and sleep.

#### **Alternate Names:**

NREM sleep

### **Non-Responder**

A person or disease that shows little or no expected

---

response to a treatment, vaccine, or test stimulus.

### **Nonselective Beta-Blockers**

Medicines that block two main types of adrenaline receptor. They slow the heart and lower blood pressure but can also narrow the airways, so people with asthma or certain lung diseases may need a different medicine.

### **Nonseminomatous Germ Cell Tumor**

A cancer arising from reproductive cells, most often in the testicle. It may grow and spread quickly; blood tests, scans, and tissue examination guide staging and treatment.

#### **Alternate Names:**

NSGCT; Nonseminoma

### **Non-Seminomatous Germ Cell Tumor**

A major clinicopathological category of testicular and extragonadal germ cell neoplasms comprising embryonal carcinoma, yolk sac tumor, choriocarcinoma, and teratoma (or mixed variants). Characterized by aggressive clinical behavior with early hematogenous and retroperitoneal lymphatic dissemination, producing characteristic serum tumor markers (alpha-fetoprotein and human chorionic gonadotropin); treated with radical orchiectomy followed by platinum-based combination chemotherapy.

#### **Alternate Names:**

NSGCT; Nonseminoma

### **Nonseminomatous Testicular Tumors**

Testicular germ-cell tumors containing or resembling embryonal carcinoma, yolk-sac tumor, choriocarcinoma, teratoma, or mixed components. They may produce elevated AFP or hCG and require staging with tumor markers, imaging, and examination of the contralateral testis.

### **Nonsense Variant**

A change in DNA that tells a cell to stop making a protein too early. The resulting shortened or missing protein may cause disease, but the effect depends on the gene, the location of the change, and how much normal protein remains.

### **Non-Small Cell Lung Cancer**

The most common group of lung cancers, including adenocarcinoma and squamous-cell cancer. It may start in different parts of the lung and can spread; treatment depends on its stage, cell type, gene changes, and the person's health.

### **Non-Small Cell Lung Carcinoma**

A broad histological grouping of primary epithelial malignancies of the lung comprising approximately 85 percent of all lung cancers, including adenocarcinoma, squamous cell carcinoma, and large cell carcinoma, distinguished by clinical behavior and targeted therapy sensitivity from small cell lung cancer.

#### **Alternate Names:**

NSCLC

### **Nonspecific Interstitial Pneumonia**

A pattern of lung inflammation or scarring that develops relatively evenly over time. It is often linked to connective-tissue disease, medicines, or an unknown cause and can cause breathlessness and a dry cough.

### **Nonspecific Interstitial Pneumonitis**

A pattern of lung inflammation and scarring that develops fairly evenly through the lung tissue. It may cause a dry cough and breathlessness and is often linked to autoimmune disease, medicines, or no identified cause.

### **Non-Specific Urethritis**

Inflammation of the tube that carries urine out of the body when a specific cause has not yet been identified. It may cause burning urination or discharge and is often linked to sexually transmitted infections, so testing and treatment of partners may be needed.

### **Non-ST Elevation Myocardial Infarction**

An acute coronary syndrome characterized by ischemic myocardial necrosis with detectable elevated cardiac biomarkers (cardiac troponin I or T) in the absence of acute persistent ST-segment elevation on ECG (presenting instead with ST-segment depres-

---

sion, T-wave inversion, or normal tracing).

**Alternate Names:**

NSTEMI; Non-ST-Segment Elevation Infarct

**Non-ST-Elevation Myocardial Infarction**

A heart attack in which blood tests show heart-muscle injury but the electrocardiogram does not show persistent ST-segment elevation. It is usually caused by reduced or blocked coronary blood flow and may require urgent assessment and treatment.

**Nonsteroidal Anti-inflammatory Drug (NSAID) Toxicity**

Overdose causing nausea, vomiting, abdominal pain, gastrointestinal bleeding, tinnitus, drowsiness, acute kidney injury, metabolic acidosis, seizures, or coma in severe cases. Assess renal function, acid-base status, and coingestants; treatment is mainly supportive, with activated charcoal considered soon after a potentially serious ingestion.

**Nonsteroidal Anti-Inflammatory Drugs**

A group of medicines that reduce pain, fever, and inflammation by lowering the body's production of prostaglandins. They can cause stomach bleeding, kidney injury, fluid retention, heart problems, or allergic reactions, especially at high doses or with long use.

**Nonsteroidal Anti-Inflammatory Drugs And Ulcers**

Nonsteroidal anti-inflammatory drugs can reduce stomach-protective prostaglandins and cause gastritis, ulcers, and gastrointestinal bleeding. Risk rises with age, prior ulcer, anticoagulants, steroids, kidney disease, and high doses; safer alternatives or acid protection may be needed.

**Alternate Names:**

NSAID-Induced Peptic Ulcer; NSAID Gastropathy

**Non-Stress Test**

A test during pregnancy that monitors the baby's heart rate while the baby moves. A healthy response usually includes temporary heart-rate increases; an abnormal result may reflect sleep, medicines, or a problem requiring further assessment.

**Non-ST-Segment Elevation Myocardial Infarction**

An acute coronary syndrome characterized by evidence of myocardial necrosis (elevated cardiac troponin biomarkers) in the absence of acute ST-segment elevation on electrocardiogram, typically reflecting subendocardial ischemia caused by a non-occlusive mural thrombus.

**Alternate Names:**

NSTEMI

**Nonsyndromic**

Nonsyndromic means occurring without the additional characteristic features of a recognized syndrome; it may still be genetic, congenital, or clinically significant.

**Nontropical Sprue**

Alternate name; see *Celiac Disease*.

**Nontuberculous Mycobacteria**

Mycobacteria other than the tuberculosis complex and *M. leprae*. They are found in soil and water and can cause lung, skin, lymph-node, or disseminated infections, especially in people with underlying illness.

**Nontuberculous Mycobacterial Infection**

Infection caused by *Mycobacterium* species other than *M. tuberculosis*, often affecting the lungs but sometimes the lymph nodes, skin, or other organs. Disease is more likely in people with structural lung disease or impaired immunity.

**Nontuberculous Mycobacterial Lung Disease**

Chronic lung infection caused by environmental *Mycobacterium* species other than *M. tuberculosis*.

**Nontuberculous Mycobacterial (NTM) Diseases**

Infections caused by environmental mycobacteria other than the tuberculosis complex and *M. leprae*. They may affect the lungs, skin, lymph nodes, or multiple organs, particularly in people with struc-



---

tural lung disease or impaired immunity. It may produce cough, fatigue, weight loss, breathlessness, or coughing blood, especially in people with structural lung disease. Diagnosis requires compatible imaging and repeated microbiologic testing; treatment is prolonged and individualized.

### **Nonulcer Dyspepsia**

Alternate name; see *Functional Dyspepsia*.

### **Nonulcer Stomach Pain**

Alternate name; see *Functional Dyspepsia*.

### **Nonunion And Malunion**

Nonunion means a broken bone fails to heal after an expected period, leaving pain or instability. Malunion means it heals in a poor position, causing deformity, altered movement, or strain on nearby joints. Both may require further treatment.

### **Nonunion Fracture**

A broken bone that has failed to heal after the expected period and shows no further progress toward joining. It may cause ongoing pain, movement at the fracture, weakness, or deformity and may require surgery or other treatment.

### **Noonan Syndrome**

An inherited condition that may cause distinctive facial features, short stature, heart defects, delayed development, easy bruising, or swelling from lymph problems. Its features vary, and regular heart, growth, hearing, and developmental checks are important.

### **Noonan Syndrome With Multiple Lentigines**

A genetic disorder also called LEOPARD syndrome, characterized by multiple lentigines, distinctive facial features, heart disease, short stature, and sometimes deafness or genital abnormalities.

### **Noradrenaline**

A chemical messenger acting as both a hormone and neurotransmitter. It helps regulate alertness,

attention, stress responses, heart activity, and blood-vessel tightening, which can raise blood pressure.

#### **Alternate Names:**

Norepinephrine

### **No-Reflow Phenomenon**

Failure of blood to reach small vessels and damaged tissue even after a larger blocked artery has been reopened. It can occur after a heart attack or stroke and may limit recovery despite successful treatment of the main blockage.

### **Norepinephrine**

A hormone and nerve messenger involved in alertness, stress responses, heart function, and blood-vessel tightening. It can increase blood pressure and is also used in hospitals to support blood pressure during some forms of shock.

#### **Alternate Names:**

Noradrenaline

### **Norethisterone**

A manufactured progesterone-like medicine used for contraception and selected menstrual problems or endometriosis. It can change bleeding patterns and may cause nausea, headache, mood changes, breast tenderness, or other hormone-related effects.

### **Norgestrel**

A synthetic progestogen used in some hormonal contraceptives. It helps prevent pregnancy by suppressing ovulation and changing cervical mucus and the uterine lining.

### **Norketamine**

Norketamine is an active metabolite of ketamine with NMDA-receptor antagonist activity; it contributes to ketamine's analgesic and psychoactive effects and may persist after the parent drug is cleared.

### **Normal**

Within an expected range or consistent with usual anatomy, physiology, or function for a specified population.



---

## Normal and Abnormal Puerperium

The puerperium is the weeks after birth while the uterus and other body systems recover. Normal findings include lochia, uterine involution, breast changes, fatigue, and emotional fluctuation; heavy bleeding, fever, severe pain, headache, breathlessness, or profound mood change may signal a complication.

## Normal Flora

The bacteria, fungi, and other microorganisms that normally live on the skin and in the mouth, intestine, and urinary or genital tract without causing illness. They can help block harmful germs, support immunity, and contribute to digestion and vitamin production.

## Normal Growth And Development

Normal growth and development describes expected physical growth, movement, language, learning, and social progress for a person's age and stage of life.

## Normalization

A mathematical adjustment that makes measurements comparable by correcting for known differences in scale, background, equipment response, or other measurement conditions.

## Normal Labor and Delivery

Spontaneous labor with progressive cervical dilation, descent and birth of the fetus, delivery of the placenta, and stable maternal and newborn adaptation. Monitoring remains important because labor can become prolonged or complicated by fetal distress, malposition, bleeding, or infection.

## Normal Labour

Spontaneous labour at term with regular uterine contractions, progressive cervical dilatation, and delivery through the birth canal without major complication.

## Normal Pressure Hydrocephalus

A buildup of cerebrospinal fluid that enlarges spaces inside the brain and may cause a slow, unsteady walk, reduced thinking, and loss of bladder control.

Some people improve with specialist evaluation and drainage treatment.

## Normal-Pressure Hydrocephalus

A disorder in which cerebrospinal fluid accumulates and causes gait difficulty, cognitive decline, and urinary problems despite near-normal measured pressure.

## Normal Range

A normal range is the interval of results commonly found in a reference population; a result outside it does not by itself prove disease.

## Normal Saline

A sterile approximately 0.9% sodium-chloride solution used for fluid replacement, irrigation, and dilution of medicines.

## Normal-Tension Glaucoma

A type of open-angle glaucoma in which the optic nerve is damaged and vision may be lost even though pressure inside the eye remains within the normal range.

## Normoblast

A nucleated immature red-cell precursor in bone marrow that normally loses its nucleus before entering circulation.

## Normocalcemic Hyperparathyroidism

Condition of elevated parathyroid hormone with persistently normal serum calcium and normal vitamin D levels. Exclusion of secondary causes of elevated PTH including vitamin D deficiency, renal insufficiency, and medications is required. Represents an early or mild form of primary hyperparathyroidism. May progress to overt hypercalcemia.

## Normocytic Anaemia

Alternate British spelling; see *Normocytic Anemia*.

## Normocytic Anemia

A form of anemia in which circulating red blood cells are of normal average size (mean corpuscular

---

volume [MCV] between 80 and 100 fL), but total red blood cell count and hemoglobin concentration are decreased. Diagnostic evaluation relies on the reticulocyte count to differentiate between: (1) hyperproliferative causes with elevated reticulocytes, such as acute blood loss (hemorrhage) or premature red cell destruction (hemolytic anemia), and (2) hypoproliferative causes with low reticulocytes, including anemia of chronic disease, chronic kidney disease (erythropoietin deficiency), aplastic anemia, and bone marrow infiltration. Symptoms include fatigue, weakness, dizziness, and pallor.

**Alternate Names:**

Normocytic Normochromic Anemia; Normal-MCV Anemia

**Normotensive**

Having blood pressure within the expected range for the individual and measurement setting. Having blood pressure within the expected range for a person, based on the measurement setting.

**Norovirus**

A highly contagious virus that causes acute gastroenteritis, or inflammation of the stomach and intestines. Symptoms usually begin 12–48 hours after exposure and include sudden nausea, vomiting, watery diarrhea, stomach cramps, and sometimes fever, headache, or body aches. It spreads mainly when tiny particles of stool or vomit enter the mouth, through infected people, contaminated food or water, surfaces, or particles released during vomiting. Most illness improves within 1–3 days, but severe fluid loss can cause dehydration, especially in infants, older adults, and people with other illnesses.

**Norovirus Infection**

Acute gastroenteritis caused by norovirus. Symptoms usually begin 12–48 hours after exposure and include nausea, vomiting, watery diarrhea, abdominal cramps, and sometimes fever, headache, or body aches. Infection spreads easily through infected people, contaminated food or water, surfaces, or particles released during vomiting. Most cases improve within 1–3 days, but fluid loss may cause

dehydration, especially in infants, older adults, and people with other illnesses.

**Norrie Disease**

A rare inherited eye disorder in which the retina develops abnormally, usually causing severe vision loss or blindness from birth or early infancy. Some people also develop hearing loss, developmental differences, or problems in the eye's blood vessels.

**Nortriptyline**

A medicine used to treat depression and sometimes long-lasting nerve pain. It changes the levels of certain chemical messengers in the nervous system and may cause dry mouth, constipation, sleepiness, dizziness, heart-rhythm problems, or overdose toxicity.

**Norwegian Scabies**

A severe form of scabies in which thick crusts contain very large numbers of mites. It is extremely contagious and occurs especially in people with weakened immunity or difficulty sensing or scratching the skin, requiring prompt treatment and contact precautions.

**Norwood Procedure**

The first operation for many newborns with hypoplastic left heart syndrome and related heart defects. It rebuilds the main artery so the right ventricle can send blood to the body, creates a wide opening between the upper heart chambers, and provides controlled blood flow to the lungs through a shunt or tube. It is usually performed within the first days of life and is followed by further staged operations.

**Alternate Names:**

Stage 1 Norwood Reconstruction; Norwood Operation

**Nose**

The external structure and internal passages used for breathing and smell. The nose warms, moistens, and filters air and contains bone, cartilage, skin, mucus, blood vessels, and smell-sensing tissue.

---

## Nosebleed

Singular form; see *Nosebleeds*.

## Nosebleeds

Bleeding from blood vessels inside the nose, commonly caused by dryness, irritation, injury, infection, allergies, medicines, or high blood pressure. Episodes may be brief or recurrent and can vary in amount.

## Nose Fracture

A break in one or more nasal bones caused by trauma, potentially causing pain, swelling, deformity, bleeding, or nasal obstruction.

## Nosocomial

Acquired during healthcare, especially in a hospital, rather than being present or incubating when the person entered care. A nosocomial infection may be caused by bacteria, viruses, fungi, or other organisms and can involve drug-resistant organisms.

## Nosocomial Infection

A hospital-acquired infection developing during or after hospitalization that was not present or incubating at admission, classically defined by onset beyond 48 hours. Common sites are bloodstream, urinary tract, lungs, and surgical wounds; causative organisms are frequently multidrug-resistant, prevented by hand hygiene and stewardship.

## Nosocomial Norovirus Infection

Healthcare-associated norovirus gastroenteritis characterized by acute, explosive onset of watery diarrhea, severe non-bloody emesis, and abdominal cramps. Outbreak management necessitates strict patient isolation, contact precautions, rigorous hand hygiene with soap and water (as alcohol-based rubs are less effective against non-enveloped virions), and hypochlorite-based surface disinfection.

### Alternate Names:

Hospital-Acquired Norovirus; Healthcare Norovirus Outbreak; Hospital Winter Vomiting Disease

## Nosocomial Vancomycin-Resistant

## Enterococci

Enterococcal bacteria that resist vancomycin and can spread in healthcare settings. They may cause urinary, wound, bloodstream, or other infections, particularly in people who are seriously ill or have devices such as catheters; treatment requires susceptibility testing.

## Nostril

One of the two openings at the front of the nose. Each nostril leads into the nasal passage, where hairs and mucus help trap particles before air reaches the deeper airways.

## Notalgia Paresthetica

A localized neuropathic itch or burning sensation, usually between or just below the shoulder blades, caused by irritation or entrapment of thoracic sensory nerves. Scratching may produce a persistent darkened patch; treatment focuses on reducing itch and addressing contributing spine or nerve disease when present.

## Notencephalocele

A rare congenital neural-tube defect involving protrusion of brain tissue or meninges through a skull defect; terminology varies by source.

## Notifiable Disease

An infection or other condition that healthcare professionals or laboratories must report to public-health authorities under local law. Reporting helps detect outbreaks, arrange control measures, protect contacts, and plan services; the exact list and reporting rules vary by location.

## Notochord

A flexible embryonic rod that guides axial development and largely disappears as the vertebral column forms.

## Novel Biomarkers of Kidney Function

Emerging blood, urine, or tissue measurements intended to detect kidney injury or altered filtration earlier or more specifically than serum creatinine alone. Examples include selected tubular-injury pro-

---

teins, but clinical usefulness depends on validation, timing, and the patient's context.

## **NOWS**

Alternate name; see *Neonatal Opioid Withdrawal Syndrome*.

## **Noxious**

Harmful, poisonous, or capable of causing tissue injury or pain. Noxious stimuli activate protective sensory pathways such as pain and withdrawal reflexes.

## **NREM Sleep**

The part of sleep without rapid eye movements, ranging from light sleep to deep slow-wave sleep. It alternates with REM sleep in repeated cycles and supports physical restoration, learning, and memory.

## **NSAID**

A nonsteroidal anti-inflammatory drug used to reduce pain, fever, and inflammation. Examples include ibuprofen and naproxen; these medicines can irritate the stomach, cause bleeding or kidney problems, and increase heart risks in some people.

### **Alternate Names:**

Nonsteroidal anti-inflammatory drug

## **NSAID-Induced Ulcers**

Alternate name; see *Peptic Ulcer Disease*.

## **NSTEMI**

A heart attack in which a blocked or narrowed coronary artery injures heart muscle but does not produce the typical persistent ST-segment elevation on an ECG. It may cause chest pressure, breathlessness, sweating, nausea, or faintness.

### **Alternate Names:**

Non-ST-Elevation Myocardial Infarction

## **NTD**

NTD commonly means neural-tube defect, a congenital brain or spinal-cord malformation associated with inadequate folate and other factors.

## **Nuchal**

Relating to the back of the neck. Nuchal rigidity means painful or limited neck bending and can occur with meningitis or bleeding around the brain. Nuchal translucency is a first-trimester ultrasound measurement used to estimate certain pregnancy risks.

## **Nuchal Cord**

Wrapping of the umbilical cord around the fetal neck, occurring once or more times. It is common and often harmless, but monitoring during labor is needed if fetal heart-rate abnormalities occur.

## **Nuchal Rigidity**

Stiffness or pain that makes it difficult to bend the neck forward. It may occur with meningitis, bleeding around the brain, severe muscle problems, or other illnesses; sudden stiffness with fever or severe headache may signal a serious condition.

## **Nuchal Translucency Screening**

An ultrasound measurement of the clear space at the back of a fetus's neck, usually performed early in pregnancy. A larger-than-expected measurement can indicate increased risk of chromosome or heart problems but is not a diagnosis.

## **Nuchal Translucency Test**

An early-pregnancy ultrasound measurement of the clear space at the back of the fetus's neck. It estimates the chance of certain chromosome or structural problems but cannot diagnose them; abnormal results need follow-up testing.

## **Nuclear**

Relating to the nucleus of a cell, the center of an atom, or medical tests and treatments that use radioactivity. The intended meaning depends on the surrounding medical context.

## **Nuclear Factor Kappa B**

A ubiquitous inducible master transcription factor complex that coordinates immune and inflammatory responses, cell survival, and cytokine transcription; normally held inactive in the cytoplasm by I-kappa-

---

B inhibitor proteins until activated by inflammatory cytokines, bacterial LPS, or viral infection.

**Alternate Names:**

NF-kappaB; NF-kB

## **Nuclear Medicine**

A medical specialty that uses small amounts of radioactive substances to diagnose or treat disease. Scans show body function or chemical activity, while treatments deliver radiation to selected tissues.

## **Nuclear Medicine Imaging**

Imaging that uses a radioactive tracer and a specialized detector to show organ function, blood flow, chemical activity, or disease distribution. Common examples include PET, SPECT, bone scans, thyroid scans, and radionuclide cardiac imaging.

## **Nuclear Pore Complex**

A large, multiprotein cylindrical channel spanning the double-membrane nuclear envelope, composed of multiple nucleoporin proteins that regulate and mediate the selective bidirectional transport of macromolecules (RNA, ribonucleoproteins, proteins) between the nucleus and the cytoplasm.

**Alternate Names:**

NPC

## **Nuclear Sclerosis**

A common age-related cataract in which the center of the eye's lens gradually becomes yellow and hard. It can slowly blur vision, reduce contrast, or alter color perception and is treated when it interferes with daily activities.

## **Nuclear Stress Test**

A test that compares blood flow to heart muscle at rest and during exercise or medicine-induced stress using a radioactive tracer. It can identify areas of reduced perfusion that suggest coronary artery disease.

**Alternate Names:**

Nuclear medicine tests

## **Nuclear Ventriculography**

A heart scan that follows a small radioactive tracer through the bloodstream to measure how much blood the heart pumps with each beat and whether the heart walls move normally.

## **Nucleic Acid**

DNA or RNA, the molecules that store, copy, and use genetic instructions. DNA usually stores the long-term information, while RNA helps read those instructions and make proteins or perform other jobs in cells.

## **Nucleic Acid Hybridization**

A laboratory method in which matching DNA or RNA strands bind together. This allows a test to find, identify, or measure a particular genetic sequence from a person, tumor, or infectious organism.

## **Nucleolus**

A dense region within the cell nucleus where ribosomal RNA is produced and ribosome subunits begin assembly.

## **Nucleoside**

A small molecule made of a nitrogen-containing base joined to a sugar. Adding one or more phosphate groups makes a nucleotide, the building block used to make DNA or RNA and to carry energy in cells.

## **Nucleoside Analogs**

Medicines that resemble the building blocks used to make DNA or RNA. Some viruses use them by mistake, which interrupts viral copying; the medicine chosen depends on the virus, illness, kidney function, and other factors.

## **Nucleotide**

A small molecule made of a sugar, a nitrogen-containing base, and one or more phosphate groups. Nucleotides build DNA and RNA and also carry energy used by cells.

---

## Nucleus

The membrane-bound cell organelle containing most genetic material, or a defined cluster of neurons in the central nervous system.

## Nucleus Accumbens

A group of nerve cells deep in the brain that helps process reward, motivation, learning, and reinforcement. Its activity is involved in normal behavior and in disorders such as addiction and some psychiatric illnesses.

## Nucleus Ambiguus

A longitudinal column of branchial motor lower motor neurons located within the medullary reticular formation that provides motor innervation to the skeletal striated muscles of the soft palate, pharynx, and larynx via cranial nerves IX (glossopharyngeal), X (vagus), and cranial XI.

## Nucleus Pulposus

The soft, gel-like center of a spinal disk, surrounded by a tough outer ring. It helps spread pressure between vertebrae, but a tear may allow it to bulge or press on nerves.

## Nulligravida

An obstetric term for a person who has never been pregnant. It is different from nullipara, which means never having given birth beyond the stage of pregnancy used in a particular medical record system.

## Nullipara

A person who has never carried a pregnancy beyond the stage of fetal viability; the adjective is nulliparous.

## Null Mutation

A genetic change that prevents a gene from producing a functional protein or RNA. The effect depends on the gene, the other copy, and whether the change occurs in body cells or reproductive cells.

## Numb

Numb means lacking normal sensation or emotional

responsiveness; physical numbness can result from nerve, brain, circulation, metabolic, or medication-related problems.

## Number Needed To Harm

The estimated number of people who must receive a treatment, rather than the comparison treatment, for one extra person to experience a specified harmful outcome during a stated period. A smaller number means the harm is more frequent.

## Number Needed To Treat

The number of people who must receive an intervention for one additional person to experience the specified beneficial outcome compared with the control over a defined time. It is the reciprocal of absolute risk reduction.

## Numbness

Numbness is reduced or absent sensation, often from nerve compression, neuropathy, stroke, spinal disease, circulation problems, or anesthetic effects.

## Numbness And Tingling

Unusual reduced sensation or prickling, often caused by nerve compression, neuropathy, spinal disease, vitamin deficiency, circulation problems, or other neurologic disorders. Sudden one-sided symptoms, weakness, speech trouble, or facial drooping require emergency assessment.

### Alternate Names:

Paresthesia; Pins and Needles

## Numbness In Hands

Reduced or absent sensation in one or both hands, commonly caused by nerve compression, neuropathy, injury, circulation problems, or systemic disease.

## Nummular

Coin-shaped or round. The term is commonly used for nummular eczema, which causes itchy, coin-shaped patches of inflamed skin.

## Nummular Dermatitis

An inflammatory skin condition causing itchy, coin-



---

shaped patches, often on the arms or legs. Dry skin, irritants, and skin injury may contribute; treatment commonly includes moisturizers and prescribed anti-inflammatory medicines.

### **Nummular Eczema**

An inflammatory skin condition causing intensely itchy, coin-shaped red patches, sometimes with blisters, crusting, or scaling, usually on the arms or legs. Dry skin, irritation, or injury can trigger it and similar rashes may need examination.

### **Nursemaid's Elbow**

Nursemaid's elbow is subluxation of the radial head after sudden traction on a young child's extended or pronated arm; the child avoids using the arm, and gentle clinician reduction usually restores movement.

### **Nurse Practitioner**

A nurse practitioner is an advanced-practice nurse who can assess patients, diagnose and treat many conditions, prescribe medicines where authorized, and provide preventive care.

### **Nursery**

A place or service for the care of infants, including specialized newborn care in a hospital.

### **Nursing**

The professional practice of caring for people, promoting health, preventing illness, and supporting recovery or comfort.

### **Nursing Home**

A residential facility that provides ongoing nursing care, assistance with daily activities, and medical or rehabilitation services for people who cannot live independently.

### **Nursing Home Acquired Pneumonia**

Pneumonia in a nursing-home or long-term-care resident, often associated with frailty, swallowing impairment, immobility, chronic disease, or viral exposure. Evaluation should consider severity, as-

piration, goals of care, and recent healthcare or antibiotic exposure rather than assuming resistant infection.

### **Nut Allergies**

Nut allergy is an immune-mediated reaction to tree nuts or peanuts that can cause hives, gastrointestinal or respiratory symptoms, or life-threatening anaphylaxis.

### **Nut-Allergy**

An immune-mediated reaction to one or more nuts, potentially causing urticaria, anaphylaxis, or gastrointestinal and respiratory symptoms.

### **Nutation**

A nodding or rocking movement; in anatomy it commonly describes movement of the sacrum relative to the pelvis.

### **Nutcracker Esophagus**

An older name for a swallowing disorder in which the esophagus contracts with unusually high pressure. It can cause intermittent chest pain or difficulty swallowing; current testing and clinical assessment distinguish it from reflux and other causes.

### **Nutcracker Syndrome**

Compression of the vein draining the left kidney, usually between two nearby arteries. It may cause blood in the urine, pelvic or flank pain, or enlarged veins; mild cases may be monitored, while persistent severe symptoms may need treatment.

### **Nutrient**

A substance the body needs for energy, growth, repair, or normal function. Nutrients include protein, fats, carbohydrates, vitamins, minerals, and water; too little or too much of some nutrients can cause illness.

### **Nutrigenomics**

The study of how food and nutrients interact with genes and how genetic differences influence responses to diet. It is used in research on metabolism

---

and disease risk; many proposed personalized-nutrition applications remain under study.

## **Nutrition**

How the body takes in, digests, absorbs, and uses food and its nutrients for health.

## **Nutritional Anemia**

Anemia caused by inadequate intake, absorption, or use of nutrients needed for red-cell production, especially iron, vitamin B12, or folate.

## **Nutritional Assessment**

A structured evaluation of food intake, body measurements, laboratory results, symptoms, and health conditions. It identifies nutritional problems and guides a safe, individualized eating plan or nutrition support.

## **Nutritional Deficiency**

A lack of a vitamin, mineral, protein, energy, or other nutrient needed for normal body function. It may result from inadequate intake, poor absorption, increased needs, or disease and can impair growth, blood formation, immunity, nerves, or other organs.

## **Nutritional Dermatoses**

Skin changes caused by too little or too much of a nutrient. For example, vitamin C deficiency can cause easy bruising and bleeding gums, while niacin deficiency can cause a sun-sensitive rash, diarrhea, and confusion.

## **Nutritional Disorders**

Health problems caused by too little, too much, or poor use of nutrients. Examples include undernutrition, obesity, vitamin or mineral shortages, and problems caused by the body being unable to digest or absorb food properly.

## **Nutritional Screening**

A brief check for risk of malnutrition using weight change, food intake, illness, and other factors. A concerning result leads to a fuller assessment and an individualized nutrition plan.

## **Nutrition-Focused Physical Examination**

A structured examination for physical findings associated with malnutrition, including loss of subcutaneous fat, muscle wasting, edema, ascites, reduced functional status, and micronutrient-related signs. It complements dietary history, anthropometry, laboratory data, and disease assessment.

## **Nutrition In Pregnancy**

Eating enough varied food and obtaining appropriate vitamins and minerals to support the pregnant person and growing fetus. Important nutrients include folate, iron, calcium, vitamin D, iodine, and protein; individual needs and supplements should be reviewed with a prenatal clinician.

## **Nutrition Risk Screening 2002**

A hospital screening questionnaire that considers recent food intake, weight change, illness severity, and age to identify adults at risk of malnutrition. A high score prompts a fuller assessment; it is not itself a diagnosis.

## **Nutrition Support**

Use of oral nutrition supplements, tube feeding, or intravenous nutrition when a person cannot meet nutritional needs by ordinary eating. The method depends on the digestive tract, illness, nutrition needs, and risks of treatment.

## **Nutritive**

Providing nourishment that the body can use for energy, growth, repair, or normal function. Nutritive substances include protein, fat, carbohydrates, vitamins, minerals, and water.

## **Nux Vomica**

The seed of *Strychnos nux-vomica*, containing strychnine and brucine; ingestion can cause severe convulsant poisoning.

## **Nyctalopia**

Impaired vision in dim light or darkness, also called night blindness.

---

## NYHA Functional Classification

A four-class system describing how heart failure symptoms limit activity: class I causes no limitation, class II limits ordinary activity, class III limits less-than-ordinary activity, and class IV causes symptoms even at rest.

## Nystagmus

Involuntary rhythmic movement of the eyes. It may occur with inner-ear disease, neurologic disorders, reduced vision, medicines, or normal early development; new nystagmus with dizziness, weakness, or vision change may signal an underlying disorder.

## Nystagmus Retractorius

An unusual eye movement in which the eyes pull backward and may move toward each other when a person tries to look upward. It is a characteristic sign of a problem in the upper part of the brain, but other findings are needed to identify the cause.

### Alternate Names:

Retraction-Convergence Nystagmus; Koerber-Salus-Elschnig Sign

## Nail-Biting

Repetitive biting of the fingernails or surrounding skin, often associated with stress, habit, or a body-focused repetitive behavior. It can damage the nail, skin, teeth, or jaw and may introduce infection.

## Narcoleptic Syndrome

Alternate name for narcolepsy, a neurologic sleep disorder causing excessive daytime sleepiness and, in some people, cataplexy, sleep paralysis, or vivid hallucinations; see *Narcolepsy*.

## Nasal Septal Perforation Repair

A procedure to close or reconstruct a hole in the nasal septum, using local tissue, grafts, or other techniques when symptoms or progressive damage justify repair.

## Naturopathic Medicine

A system of complementary or alternative health practices that may use diet, supplements, herbs,

and other treatments. Products can interact with medicines or cause harm, and naturopathic care should not replace proven treatment for serious disease.

## Navicular Fracture of the Wrist

A fracture of the wrist navicular, more commonly called the scaphoid. It may cause pain in the anatomic snuffbox after a fall and can heal poorly because part of the bone has limited blood supply.

## NDI

Abbreviation for nephrogenic diabetes insipidus, in which the kidneys do not respond adequately to vasopressin and produce excessive dilute urine; see *Nephrogenic Diabetes Insipidus*.

## Nearsightedness (Myopia)

Alternate name; see *Nearsightedness*.

## Necrotizing Fasciitis of the Perineum

Necrotizing soft-tissue infection of the perineum or genital region, commonly called Fournier gangrene. Severe pain, swelling, skin discoloration, fever, or rapid progression requires emergency surgery and antibiotics.

## Needle Aspiration Biopsy

A biopsy that uses a thin needle to remove cells or fluid from a lump, organ, or lesion for laboratory examination. Results are interpreted with imaging and clinical findings and may require a larger biopsy.

## Neonatal Listeriosis

Listeria infection acquired before birth or during delivery that can cause sepsis, pneumonia, meningitis, or other serious illness in a newborn; see *Listeriosis*.

## Nephrocarcinoma

A carcinoma arising in the kidney, usually referring to renal cell carcinoma; see *Renal Cell Carcinoma*.

## Nephropathy Diabetic

Alternate name; see *Diabetic Nephropathy*.

---

## Nicotine Withdrawal

Symptoms occurring after reducing or stopping nicotine, such as irritability, craving, anxiety, low mood, restlessness, poor concentration, increased appetite, and sleep disturbance. Symptoms usually improve with time and can be treated with behavioral and medication support.

## Night Eating

A pattern of eating after dinner or waking to eat during the night, sometimes with reduced morning appetite; persistent distress or impairment may indicate night eating syndrome or another eating disorder.

## Nitroglycerin for Chest Pain

Nitroglycerin may relieve angina caused by reduced coronary blood flow, but new, severe, persistent, or unexplained chest pain requires emergency evaluation rather than repeated self-treatment.

## Noninfectious Uveitis

Inflammation inside the eye not caused by an active infection, often related to autoimmune or inflammatory disease. It can threaten vision and requires prompt ophthalmic assessment.

## Non-Scarring Epidermolysis Bullosa

An epidermolysis bullosa subtype in which genetic fragility causes blisters or erosions without primary scarring; severity and complications depend on the subtype.

## Nonthrombocytopenic Idiopathic Purpura

Purpura or easy bruising with a normal platelet count and no identified cause; the evaluation considers vessel fragility, medicines, coagulation disorders, nutritional deficiency, and systemic disease.

## Nontraditional Medicine

Treatments outside conventional evidence-based medical practice, including some complementary, alternative, or traditional systems. Safety and effectiveness vary, and interactions or delayed diagnosis are possible.

## Normal Menstrual Cycle

The recurring hormonal and uterine changes that usually produce menstrual bleeding at a predictable interval; cycle length, flow, and symptoms vary, and persistent changes require evaluation.

## Noroviruses

A group of highly contagious viruses that cause acute gastroenteritis with vomiting, diarrhea, cramps, and dehydration; see *Norovirus*.

## Norrbottnian Gaucher Disease

A rare northern-European genetic form of Gaucher disease, usually a non-neuronopathic type caused by inherited glucocerebrosidase deficiency; see *Gaucher Disease*.

## NPS

Abbreviation for nail-patella syndrome, an inherited disorder affecting nails, kneecaps, elbows, pelvis, eyes, and sometimes kidneys; see *Nail Patella Syndrome*.

## NS

Abbreviation commonly used for Noonan syndrome, a genetic condition with characteristic facial features, short stature, congenital heart disease, bleeding tendency, or developmental differences; see *Noonan Syndrome*.

## Nutritional Rickets

Defective bone mineralization in a growing child caused by inadequate vitamin D, calcium, phosphate, or absorption, leading to bone pain, growth problems, or deformity; see *Rickets*.

## Nystagmus Benign Paroxysmal Positional

Positional nystagmus caused by movement of displaced inner-ear crystals, usually as part of benign paroxysmal positional vertigo; see *Benign Paroxysmal Positional Vertigo*.

# O

## OAB

Abbreviation; see *Overactive Bladder*.

## Ober Test

An orthopedic examination maneuver used to detect tightness or contracture of the tensor fasciae latae and iliotibial (IT) band; with the patient in a lateral decubitus position, the examiner abducts and extends the upper leg at the hip and releases it; failure of the thigh to adduct smoothly toward the examination table indicates a positive test.

### Alternate Names:

Ober's Test; Iliotibial Band Tightness Test

## Obesity

A chronic disease involving abnormal or excessive accumulation of body fat that may impair health. It is linked with a higher risk of conditions such as type 2 diabetes, high blood pressure, heart disease, stroke, sleep apnea, osteoarthritis, and some cancers. Many factors can contribute, including genes, metabolism, hormones, medicines, eating patterns, activity, environment, and social conditions; it is not caused by one factor or by lack of willpower alone. Body mass index (BMI) is commonly used to classify obesity in adults, but it does not measure body fat directly and is interpreted with other information.

## Obesity Hypoventilation Syndrome

A breathing disorder in which excess body weight contributes to too little breathing and a buildup of carbon dioxide while awake. It often occurs with sleep apnea and may cause sleepiness, headaches, breathlessness, low oxygen, and heart strain.

## Obex Region

A small area at the lower end of the brain's fourth fluid-filled space, where it narrows into the central canal of the spinal cord. It lies near centers controlling breathing, swallowing, and vomiting.

## OB/GYN

Abbreviation for obstetrics and gynecology, the medical specialties that care for pregnancy, childbirth, the period after birth, and the reproductive system.

## Objective Response Rate

The percentage of people in a cancer study whose tumors shrink by a predefined amount or disappear during treatment. It does not show how long the response lasts or whether treatment improves survival.

## Objects Or Insects In Ear

An object or insect in the ear canal can cause pain, hearing loss, bleeding, discharge, or a sensation of movement. Deep, sharp, chemical, or eardrum-near objects can damage the ear and may need professional removal.

### Alternate Names:

Aural Foreign Body; Foreign Body in Ear

## Oblique Fracture

A break in a bone that runs diagonally across it.

## Oblique Plane

A slice or view through the body or an organ at an angle other than straight across. Medical imaging can use oblique planes to show structures that are difficult to see in standard horizontal, vertical, or

---

front-to-back views.

### **Obliteration**

Complete closing, destruction, or disappearance of a space, passage, or hollow channel in the body. It may result from scarring, inflammation, growth, injury, or treatment and can prevent normal movement of air, blood, fluid, or other contents.

### **Observation**

Planned monitoring of a person, symptom, disease, or test result without immediate treatment or a procedure. Follow-up timing and signs that require reassessment should be defined.

### **Obsession**

A recurrent, intrusive, unwanted thought, image, or urge that causes distress and is difficult to control.

### **Obsessions**

Repeated, unwanted thoughts, images, or urges that enter the mind despite attempts to stop them and cause distress. Common examples involve contamination, harm, mistakes, symmetry, or taboo subjects; having an obsession does not mean a person will act on it.

### **Obsessive-Compulsive And Related Disorders**

A group of mental disorders involving recurrent intrusive thoughts, repetitive behaviors or mental acts, preoccupation with perceived appearance flaws, difficulty discarding possessions, or repetitive body-focused behaviors. Examples include obsessive-compulsive disorder, body dysmorphic disorder, hoarding disorder, trichotillomania, and excoriation disorder.

### **Obsessive Compulsive Disorder**

Alternate spelling; see *Obsessive-Compulsive Disorder*.

### **Obsessive-Compulsive Disorder**

A mental-health disorder involving unwanted, repeated thoughts, images, or urges and repetitive

behaviors or mental acts performed to reduce distress or prevent a feared event. The symptoms are time-consuming or cause clinically significant distress or impairment.

### **Obsessive-Compulsive Personality Disorder**

A long-lasting pattern of excessive concern with order, rules, perfection, and control. It can make a person inflexible, slow to complete tasks, or unable to delegate and can interfere with relationships or work.

### **Obstetrical Forceps**

A surgical instrument used in selected vaginal births to guide the baby's head when delivery needs assistance. Safe use requires the correct position, an experienced clinician, and assessment of risks and alternatives.

### **Obstetric Anal Sphincter Injury**

A severe tear during childbirth that extends into the muscles controlling the anus. It may cause pain, difficulty controlling gas or stool, or later incontinence and needs prompt examination, repair, and follow-up.

### **Obstetric Cholestasis**

Alternate name; see *Intrahepatic Cholestasis of Pregnancy*.

### **Obstetric Epidural Anesthesia**

A regional neuraxial analgesic technique in which a local anesthetic and opioid are continuously infused through a catheter into the lumbar epidural space to provide pain relief during labor and delivery.

#### **Alternate Names:**

Epidural Analgesia in Labor; Labor Epidural; Obstetric Epidural

### **Obstetric Fistula**

A hole that forms between the birth canal and the bladder or rectum after prolonged, obstructed labour without timely medical care. It causes continuous leakage of urine or stool and may lead to skin infections, kidney problems, and social isolation.



---

Most cases can be prevented with timely maternity care, and many can be repaired with surgery.

### **Obstetric Forceps**

Paired curved instruments placed around the fetal head during selected assisted vaginal births. They can help complete delivery when the baby or pregnant person needs delivery sooner and conditions are suitable.

#### **Alternate Names:**

Delivery Forceps; Instrumental Extraction Tongs

### **Obstetrician**

A medical doctor who cares for pregnant people during pregnancy, labor, birth, and the postpartum period. Obstetricians manage routine and high-risk pregnancies, perform deliveries, and treat complications such as high blood pressure, diabetes, bleeding, or preterm labor.

### **Obstetric Nursing**

Obstetric nursing provides care during pregnancy, labor, birth, and the postpartum period, including monitoring, education, support, and recognition of complications.

### **Obstetrics**

The branch of medicine concerned with care during pregnancy, labor, birth, and the weeks after birth, including routine care and treatment of complications affecting the pregnant person or baby.

### **Obstetric Ultrasound**

An ultrasound scan during pregnancy used to check the baby's age, growth, anatomy, position, heart-beat, and sometimes the placenta and fluid.

### **Obstipation**

Severe constipation with inability to pass stool or gas. It can occur with a blockage in the intestine and may cause abdominal swelling, pain, vomiting, or serious illness.

### **Obstructed Infected Urinary System**

A blocked urinary tract with an infection behind

the blockage. It is a medical emergency because infection can spread quickly and the blockage may need urgent relief.

### **Obstructed Labour**

Failure of the fetus to pass through the birth canal despite adequate uterine contractions, commonly because of disproportion, abnormal presentation, or an obstructing lesion.

### **Obstruction**

A blockage that prevents normal movement through a body passage, such as an airway, intestine, bile duct, blood vessel, or urinary tract. Causes include swelling, stones, scars, tumors, clots, and foreign objects.

### **Obstructive Azoospermia**

A cause of male infertility in which no sperm enters the semen because a reproductive tube is blocked or missing, although the testes make sperm normally. Causes include a missing vas deferens, vasectomy, infection-related scarring, blocked ejaculatory ducts, or surgical injury. Testicle size and hormone levels are often normal. Treatment may reopen the tube or collect sperm for use with in-vitro fertilization.

#### **Alternate Names:**

OA; Excretory Azoospermia

### **Obstructive Hydrocephalus**

A buildup of cerebrospinal fluid caused by a blockage in its normal flow through the brain's ventricles.

### **Obstructive Jaundice**

Yellowing of the skin or eyes caused by blocked bile flow, usually from a blockage in the bile ducts. Causes include gallstones, scarring, or a tumor. It may cause dark urine, pale stools, itching, or upper-abdominal pain. Fever or chills with jaundice may signal infection of the bile ducts and require urgent medical assessment.

### **Obstructive Nephropathy**

Kidney damage caused by a blockage that prevents urine from draining normally. Stones, tumors, scarring, an enlarged prostate, or birth-related problems

---

can cause it; untreated blockage may lead to infection, permanent loss of kidney function, or kidney failure.

### **Obstructive Parotitis**

Inflammation of a saliva gland near the ear because its drainage tube is blocked, often by a stone. Painful swelling usually worsens during meals, and trapped saliva can lead to repeated infection and tenderness.

### **Obstructive Shock**

A life-threatening form of shock caused by a blockage that prevents the heart from filling or pumping blood properly. Causes include tension pneumothorax, cardiac tamponade, massive pulmonary embolism, and severe constrictive pericarditis. Signs may include very low blood pressure, fast heartbeat, cold clammy skin, and swollen neck veins. Treatment removes or bypasses the blockage.

#### **Alternate Names:**

Extracardiac Obstructive Shock

### **Obstructive Sleep Apnea**

Repeated narrowing or closing of the upper airway during sleep, causing pauses in breathing, snoring, fragmented sleep, daytime tiredness, and sometimes reduced oxygen levels.

#### **Alternate Names:**

OSA; Obstructive Sleep Apnea Syndrome

### **Obstructive Sleep Apnea Syndrome**

Alternate name; see *Obstructive sleep apnea*.

### **Obstructive Uropathy**

A blockage anywhere between the kidneys and the urethra that slows or stops urine flow. Stones, scarring, an enlarged prostate, tumors, or pressure from nearby tissue may cause it and can damage the kidneys if not relieved.

### **Obtunded**

Having reduced alertness and slowed responses but still responding to a loud voice or touch. It can result from medicines, illness, metabolic problems,

poisoning, or brain disease.

### **Obturator**

Relating to a structure that closes an opening. In the pelvis, the obturator nerve helps control muscles that draw the thigh inward and carries feeling from part of the inner thigh; the obturator artery supplies nearby muscles.

### **Obturator Canal**

A narrow fibro-osseous tunnel in the superior aspect of the obturator foramen bounded by the obturator groove of the pubic bone and the upper margin of the obturator membrane, transmitting the obturator nerve, artery, and vein from the pelvis into the medial thigh compartment.

### **Obturator Foramen**

A large anatomical aperture in the pelvic bone bounded by the ischium and pubis, almost completely closed in life by the fibrous obturator membrane except at its anterosuperior margin (the obturator canal).

### **Obturator Hernia**

A rare hernia in which part of the intestine pushes through an opening in the pelvic bone. It can block the bowel and may cause pain in the inner thigh.

### **Obturator Nerve**

A nerve from the lumbar plexus that supplies muscles responsible for bringing the thigh inward and carries sensation from part of the inner thigh.

### **Obturator Sign**

A physical examination finding in acute appendicitis or pelvic peritonitis; with the patient supine, passive internal rotation of the flexed right hip stretches the obturator internus muscle against the inflamed pelvic appendix, provoking acute right lower quadrant or pelvic pain.

#### **Alternate Names:**

Cope's Obturator Test; Obturator Internus Sign

---

### **Obtuse Marginal Artery**

A branch of the circumflex coronary artery that runs along the outer border of the heart and supplies part of the left ventricle. Its size and number vary between people; blockage can cause a heart attack.

### **Occipital**

Relating to the back of the head, the occipital bone, or the occipital lobe. These structures protect and support the brain and help process visual information and head or neck movement.

### **Occipital Bone**

The bone at the back and base of the skull. It surrounds the opening where the spinal cord joins the brain and connects with the first neck vertebra.

### **Occipital Diastasis**

Widening or separation of a joint between bones at the back of a newborn's skull. It may follow difficult birth or abnormal development, and imaging assesses its extent and associated injury.

### **Occipital Hematoma**

A collection of blood in the scalp or tissues over the back of the head, often after trauma. A rapidly enlarging swelling or associated neurologic symptoms may indicate a serious injury.

### **Occipital Lobe**

The part of the brain at the back of the head that receives and interprets visual signals. Injury or disease may cause missing areas of vision, difficulty recognizing objects, or visual hallucinations.

### **Occipital Neuralgia**

Brief attacks of sharp, stabbing, or electric-shock pain caused by irritation of nerves at the back of the head. Pain may spread from the upper neck across the scalp and can be associated with tenderness or sensitivity to touch.

### **Occiput**

The back part of the skull and the back of the head. It protects the rear of the brain and includes the

occipital bone, which joins the skull base and neck.

### **Occiput Posterior Position**

A position during labor in which the back of the baby's head faces the pregnant person's back. It can cause back pain or slow descent, although many babies rotate to a better position during labor.

### **Occiput Transverse Position**

A fetal head position in which the back of the baby's head faces the pregnant person's side. It often rotates during labor; if it persists, descent may be slow and assisted delivery or cesarean birth may be needed.

### **Occlude**

To block or close a passage, vessel, airway, duct, or opening. An occlusion may be caused by a clot, stone, swelling, scar, tumor, foreign object, or deliberate medical treatment.

### **Occlusal Adjustment**

Selective reshaping of tooth surfaces to improve how the upper and lower teeth meet and reduce abnormal contact or bite interference.

### **Occlusal Guard**

A removable device worn over the teeth to protect them from grinding, clenching, or injury. A night guard may reduce tooth wear during sleep, while a sports mouthguard protects against blows; the device should fit properly and does not treat every cause of jaw pain.

### **Occlusal Splint**

A removable dental appliance that changes or protects tooth contact and may reduce effects of grinding, jaw-joint strain, or selected bite problems.

### **Occlusion**

Complete blockage of a blood vessel or body passage. It may be caused by a blood clot, plaque, pressure, swelling, or an intentional medical procedure; the effects depend on the location and can include loss of blood flow, pain, or organ damage.

---

### **Occlusive Vascular Disease**

Disease in which narrowing or blockage of a blood vessel reduces or stops blood flow to tissues.

### **Occult**

Hidden or not readily visible or detectable on routine examination. An occult disease or bleeding may be found only with imaging, microscopy, or laboratory testing.

### **Occult Blood**

Blood that is not visible to the naked eye and is detected by laboratory testing. In stool, it may be detected by a chemical guaiac test or an immunochemical test; the method and specimen determine the collection instructions and possible dietary or medicine effects. A positive stool test indicates blood is present but does not establish its cause.

### **Occult Blood Screen**

Alternate name; see *Fecal Occult Blood Test*.

### **Occult Gastrointestinal Bleeding**

Slow or intermittent bleeding somewhere in the digestive tract that cannot be seen in the stool. It may cause iron-deficiency anemia, tiredness, or a positive stool blood test and can have many possible sources.

### **Occult Primary Cancer**

Alternate name; see *Carcinoma of Unknown Primary*.

### **Occupational Asthma**

Asthma caused or worsened by substances at work, such as flour dust, latex, chemicals, or fumes. Symptoms may improve on days away from work and worsen during exposure; careful exposure history and breathing tests help confirm it.

### **Occupational COPD**

Chronic obstructive pulmonary disease caused or worsened by repeated workplace exposure to dust, fumes, chemicals, smoke, or other inhaled hazards. It causes lasting airflow limitation, cough, phlegm,

and breathlessness; smoking and other risks may also contribute.

### **Occupational Dermatoses**

Skin diseases caused or worsened by workplace conditions, such as repeated wet work, chemicals, gloves, dust, plants, or friction. Common problems include irritant or allergic contact dermatitis, often affecting the hands; reducing exposure and protecting the skin help prevent recurrence.

### **Occupational Disease**

An illness caused or significantly worsened by workplace exposures or conditions. Examples include asthma from dust, hearing loss from noise, skin disease from chemicals, and lung disease from harmful particles.

### **Occupational Exposure**

Contact with a potentially harmful substance, energy source, or condition during work. Examples include radiation, chemicals, dust, noise, heat, infection, and repetitive strain. Risk depends on the hazard, amount, duration, and protective controls; monitoring and prevention are part of occupational health.

### **Occupational Exposure Limit**

The highest recommended or legally permitted workplace exposure to a hazardous substance over a specified time. The limit varies by substance, country, route of exposure, and whether it applies to a brief peak or an average work shift.

### **Occupational Health**

The medical field focused on preventing work-related illness, injury, and harmful exposure while supporting workers' health.

### **Occupational Hearing Loss**

Hearing damage caused by repeated or prolonged workplace noise or substances that harm the ear. It is usually permanent and may cause difficulty understanding speech, especially in background noise; reducing exposure and using effective hearing protection help prevent it.

---

## Occupational Lung Disease

Lung disease caused or worsened by dusts, fumes, gases, vapors, fibers, or other exposures at work. Examples include occupational asthma, pneumoconiosis, hypersensitivity pneumonitis, and asbestos-related disease.

## Occupational Rehabilitation

Assessment, treatment, and workplace support that help a person return to work or remain employed after illness, injury, or disability. It may include graded activity, skills training, equipment, job changes, and coordination with employers and clinicians.

## Occupational Safety And Health

Rules and practices that protect workers from injuries, illness, dangerous exposures, and unsafe conditions. Measures include training, protective equipment, ventilation, safe procedures, health monitoring, and reporting or controlling workplace hazards.

## Occupational Therapist

A healthcare professional who helps people perform daily activities through rehabilitation, adaptive techniques, environmental changes, and assistive equipment.

## Occupational Therapy

Rehabilitation that helps a person perform daily activities, work, school, and other meaningful tasks after illness, injury, or disability. It may use exercises, practice, adaptive equipment, changes to the environment, and strategies for movement, thinking, or sensory problems.

## Occupational Therapy Assessment

An evaluation of how illness, injury, disability, or development affects daily activities, school, work, and self-care. The therapist identifies strengths and barriers and recommends exercises, skills practice, equipment, or environmental changes.

## OCD

Obsessive-compulsive disorder, a condition involving recurrent unwanted thoughts or urges and/or repet-

itive behaviors or mental acts that cause distress or interfere with daily life.

### Alternate Names:

Obsessive-Compulsive Disorder

## Ochre

A yellow-brown color or earth pigment. In medicine, ochre discoloration can describe yellow-brown skin or tissue changes caused by pigment, blood breakdown, chronic congestion, or other conditions.

## Ochronosis

Blue-black or brown discoloration of cartilage, tendons, skin, or the whites of the eyes caused by buildup of a chemical called homogentisic acid. It is most often linked to the inherited disorder alcaptonuria and may cause arthritis.

## OCP

OCP commonly means oral contraceptive pill, a medicine containing estrogen, progestin, or both to prevent pregnancy and sometimes treat menstrual disorders.

## Octreoscan

Nuclear-medicine imaging using indium-111 octreotide, which binds to somatostatin receptors. It was used to locate some neuroendocrine tumors and their spread; newer receptor-targeted PET scans are often preferred.

## Octreotide

Octreotide is a somatostatin analogue that suppresses growth hormone, several gastrointestinal hormones, and intestinal secretion; it is used for acromegaly, certain neuroendocrine tumors, and selected secretory diarrhoeas.

## Ocular

Relating to the eye or vision. The term may describe an eye structure, examination, medicine, injury, infection, or disease affecting sight. Ocular emergencies such as sudden severe pain or sudden loss of vision need immediate eye care.



---

## Ocular Albinism

Ocular albinism is an inherited disorder with reduced pigment primarily in the eyes, causing decreased visual acuity, nystagmus, photophobia, and abnormal retinal development.

## Ocular Bobbing

A repeated downward movement of both eyes followed by a slower return to the normal position. It can signal a serious problem in the lower brain.

### Alternate Names:

Fast Downward Ocular Movement; Pontine Ocular Bobbing

## Ocular Burns and Chemical Injuries

Injuries to the eye from chemicals, heat, radiation, or other agents. Damage may involve the conjunctiva, cornea, limbus, eyelids, or deeper structures and can threaten vision.

## Ocular Cicatricial Pemphigoid

A chronic, progressive autoimmune blistering disease mediated by autoantibodies targeting basement membrane components in mucosal tissues, presenting with recurrent conjunctival inflammation, subepithelial fibrosis, symblepharon formation, trichiasis, and progressive blinding corneal opacity.

### Alternate Names:

Mucous Membrane Pemphigoid; Cicatricial Conjunctivitis

## Ocular Fixation

The ability to hold the eyes steadily on a target; abnormal fixation can impair visual acuity, tracking, or alignment.

## Ocular Flutter

An involuntary ocular motility disorder characterized by rapid, conjugate, horizontal saccadic oscillations without an intersaccadic pause around the point of visual fixation, occurring in paraneoplastic syndromes, viral encephalitis, and cerebellar dysfunction.

### Alternate Names:

Opsoclonus Variant; Horizontal Saccadic Flutter

## Ocular Foreign Body

A foreign particle, such as dust, metal, wood, or glass, lodged on the cornea, conjunctiva, or within the eye globe, causing pain, redness, tearing, and foreign body sensation.

### Alternate Names:

Foreign Body in Eye; Corneal Foreign Body; Intraocular Foreign Body

## Ocular Histoplasmosis Syndrome

An eye condition linked to an earlier histoplasmosis infection. It can cause scars in the retina and, in some people, abnormal blood vessels that damage central vision.

## Ocular Hypertension

Higher-than-normal pressure inside the eye without current signs of glaucoma damage. It increases the future risk of glaucoma and needs regular eye checks.

## Ocular Hypotony

An abnormally low intraocular pressure, typically defined as five to six millimetres of mercury or lower. It can result from decreased aqueous humour production by the ciliary body or excessive outflow from surgical filtering blebs, cyclodialysis clefts, or wound leaks, potentially causing hypotony maculopathy, choriorretinal folds, or choroidal detachment.

### Alternate Names:

Hypotony, Ocular; Ocular Hypotension; Low Eye Tension

## Ocular Ischemic Syndrome

Eye findings caused by severely reduced blood flow through the arteries supplying the eye, usually from advanced carotid artery disease. It may cause gradual or sudden vision loss, eye pain, retinal changes, or abnormal new blood vessels.

## Ocular Larva Migrans

Eye infection caused by migrating larvae of *Toxocara* roundworms, usually affecting one eye. It can cause inflammation inside the eye, a white pupil, reduced vision, crossed eyes, or retinal damage. Diagnosis uses the eye findings, exposure history, and sometimes blood tests.



---

### Ocular Melanoma

A cancer that begins in pigment-producing cells of the eye.

### Ocular Migraine

A migraine-related visual disturbance, often involving flashing lights, zigzag lines, or a temporary blind spot. It may occur with or without headache.

### Ocular Motility

The ability of the eyes to move together in a coordinated way. Abnormal eye movements or poor alignment can cause double vision, eyestrain, dizziness, or reduced vision and may result from muscle, nerve, brain, or eye disease.

### Ocular Myasthenia Gravis

A form of myasthenia gravis in which weakness affects only the muscles that move the eyes and lift the eyelids. It causes drooping eyelids or double vision that worsens with use and improves with rest. Symptoms may later spread to swallowing, speech, breathing, or limb muscles, so follow-up is important.

#### Alternate Names:

OMG; Ocular MG

### Ocular Myiasis

Infestation of the eye or nearby tissues by fly larvae. It can cause pain, irritation, tearing, and vision problems.

### Ocular Orbit

The bony cavity around the eye. It contains the eyeball, eye-movement muscles, nerves, blood vessels, glands, and protective tissues.

### Ocular Pain

Pain in or around the eye caused by surface irritation, infection, inflammation, trauma, glaucoma, or orbital disease. Severe pain with vision loss, light sensitivity, nausea, or injury requires urgent ophthalmic assessment.

### Ocular Rosacea

Rosacea affecting the eyelids or surface of the eyes, causing burning, redness, a gritty feeling, dryness, or light sensitivity. Persistent inflammation can affect the cornea and may impair the eye surface.

### Ocular Surface Disease

A group of conditions affecting the eye's surface, eyelids, or tears, including dry eye and eyelid inflammation. Symptoms may include burning, redness, irritation, and blurred or changing vision.

### Ocular Toxoplasmosis

A retinal infection or inflammation caused by *Toxoplasma gondii*, which can produce blurred vision, floaters, pain, and recurrent retinal scars.

### Oculocardiac Reflex

A slowing of the heartbeat or similar body response caused by pressure or pulling on the eye or tissues around it.

### Oculocephalic Reflex

A brainstem reflex sometimes tested in an unconscious person by carefully moving the head. In people who cannot focus their eyes, the eyes normally move opposite to the head movement. The response may suggest that some brainstem pathways are working, but the test has safety limits and must be interpreted with the full examination.

### Oculocerebrorenal Syndrome

A rare inherited disorder, also called Zellweger spectrum disorder, that affects the brain, eyes, kidneys, liver, and other organs. Severe forms begin in newborns and can cause weak muscles, seizures, feeding problems, and major developmental problems.

#### Alternate Names:

Lowe Syndrome; OCRL

### Oculocutaneous Albinism

Alternate name; see *Albinism*.

### Oculogyric Crisis

A prolonged, involuntary upward turning of the eyes,

---

often caused by a medicine or neurologic disorder.

### **Oculomotor Nerve**

The third cranial nerve, controlling most eye movements, eyelid lifting, and pupil narrowing. Damage can cause double vision, a drooping eyelid, an outward-turning eye, or an abnormally enlarged pupil.

### **Oculomotor Nerve Palsy**

Weakness or paralysis of muscles supplied by the third cranial nerve, causing eyelid drooping, double vision, abnormal eye position, and sometimes a dilated pupil.

### **Oculomotor Nucleus**

A group of nerve cells in the midbrain that controls most eye movements, lifts the upper eyelid, and narrows the pupil through the third cranial nerve.

### **Oculopharyngeal Muscular Dystrophy**

An inherited muscle disease that usually begins in middle age and gradually weakens the eyelids and throat muscles. Drooping eyelids, difficulty swallowing, and later weakness of the hips and shoulders are typical. A gene test can confirm the diagnosis; swallowing assessment, exercises, and surgery may help prevent complications.

**Alternate Names:**

OPMD

### **Oculoplastic Procedures**

Operations on the eyelids, tear ducts, eye socket, or nearby facial tissues to restore function, repair injury, remove disease, or change appearance. Risks include bleeding, infection, scarring, dry eyes, and rarely injury affecting vision.

### **Oculosympathetic Palsy**

Alternate name; see *Horner's Syndrome*.

### **Oculovestibular Reflex**

A brainstem reflex evaluated by cold caloric stimulation in which ice-cold water irrigated into the external auditory canal of a comatose patient with

intact brainstem function produces slow conjugate tonic deviation of both eyes toward the irrigated side (mediated by cranial nerves III, VI, and VIII).

**Alternate Names:**

Cold Caloric Test; Caloric Vestibulo-Ocular Reflex

### **ODD**

Abbreviation; see *Oppositional Defiant Disorder*.

### **Odontalgia**

Pain coming from a tooth or the tissues around it. Common causes include decay, a cracked tooth, infection, gum disease, or injury; swelling, fever, difficulty swallowing, or severe pain needs prompt dental or medical care.

### **Odontoblast**

A cell along the inner part of a tooth that makes dentin, the hard layer beneath enamel. Odontoblasts produce dentin while teeth develop and can make limited repairs after injury.

### **Odontoclast**

A cell that breaks down the hard tissues of a tooth. This process helps baby teeth be shed naturally, but unwanted tooth resorption can weaken a tooth and may follow injury, inflammation, pressure, or other disease.

### **Odontodysplasia**

Odontodysplasia is abnormal development of tooth enamel and dentin, producing thin, weak, discolored teeth that may erupt late or be easily damaged.

### **Odontogenesis**

The process by which teeth form and develop before and after birth. Problems during development can cause missing teeth, weak enamel, abnormal dentin, or changes in tooth shape and eruption.

### **Odontogenic Cutaneous Fistula**

An abnormal tunnel from an infected tooth to the skin, usually on the chin, cheek, or jaw. It may appear as a small draining opening; treating the tooth infection usually closes the tunnel.

---

## Odontogenic Cyst

A fluid-filled sac arising from tissues involved in tooth development, usually within the jaw. It may be painless or cause swelling, tooth displacement, infection, or bone damage.

## Odontogenic Fibroma

An odontogenic fibroma is a rare benign tumor arising from tooth-forming connective tissue, usually presenting as a jaw lesion.

## Odontogenic Infection

An infection arising from a tooth or the tissues around it, usually caused by dental caries, pulp infection, gum disease, or trauma. It may form an abscess and spread into the jaw, face, neck, or bloodstream.

## Odontogenic Keratocyst

A cyst that develops in the jaw from tooth-forming tissue. It may grow into bone without causing symptoms, can recur after removal, and is sometimes associated with an inherited skin-cancer syndrome; imaging and tissue examination guide treatment.

## Odontogenic Keratocyst Pathology

An odontogenic cyst typically lined by a thin, uniform parakeratinized stratified squamous epithelium with a palisaded basal layer. Satellite cysts and a tendency to recur may be present.

## Odontogenic Neoplasm

An odontogenic neoplasm is a growth arising from tissues involved in tooth formation, usually in the jaw; it may be benign or malignant.

## Odontogenic Tumors and Cysts

Alternate name; see *Jaw Tumors and Cysts*.

## Odontoid Process

A peg-like projection from the second neck vertebra that fits into the first neck vertebra and allows the head to turn from side to side.

## Odontoma

A noncancerous overgrowth of tooth-forming tissue, usually found in the jaw and often discovered when a permanent tooth fails to erupt. It may look like several small teeth or a solid calcified mass and is usually removed if it blocks eruption.

## Odontopathy

Any disease or abnormality affecting a tooth or the tissues supporting it. Examples include tooth decay, gum disease, infection, abnormal tooth development, wear, and injury. Untreated dental infection can spread to surrounding tissues and occasionally to the bloodstream.

## Odynophagia

Pain when swallowing, often felt as burning, squeezing, or a sticking sensation in the throat or chest. Causes include infection, reflux, ulcers, medicines, injury, and other swallowing-tube disorders.

## Oedema

British spelling of edema: abnormal fluid accumulation in tissues that causes swelling. Causes include injury, infection, venous disease, heart or kidney failure, medicines, and lymphatic blockage.

## Oesophageal

Relating to the oesophagus, the muscular tube carrying swallowed material from the pharynx to the stomach.

## Oesophageal Atresia

A birth defect in which the swallowing tube ends blindly instead of connecting normally to the stomach. A baby may drool, cough, choke, or turn blue during feeds; urgent specialist care and surgery are required, especially when an abnormal connection to the airway is also present.

## Oesophageal Ulcer

An open sore in the swallowing tube between the throat and stomach. Reflux, infection, medicines, corrosive injury, inflammation, or cancer may cause it, leading to painful swallowing, chest pain, bleeding, or difficulty swallowing.

---

## Oesophageal Varices

Enlarged fragile veins in the wall of the swallowing tube, usually caused by high pressure in the liver's circulation. They may cause no symptoms until they rupture, leading to vomiting blood or black stools and a life-threatening emergency.

## Oesophagitis

Alternate spelling; see *Esophagitis*.

## Oesophagoscopy

Examination of the food pipe using a thin viewing tube passed through the mouth. It can investigate swallowing difficulty, pain, bleeding, or abnormal tissue, and can remove a foreign body, take samples, stop bleeding, or widen a narrowing. Sedation or anesthesia is usually used.

## Oesophagus

The muscular tube that transports swallowed food and liquid from the pharynx to the stomach.

## Oestradiol

British spelling of estradiol, the main naturally occurring estrogen during the reproductive years. It supports reproductive development, the menstrual cycle, bone health, and other body functions.

## Oestriol

A form of estrogen made in large amounts during pregnancy by the placenta using substances from the fetus and pregnant person. Its level can be measured in some prenatal screening tests, but it is not interpreted alone to diagnose a problem.

## Oestrogen

A group of hormones important for reproductive development, menstrual cycles, pregnancy, bone strength, and other body functions. They are made mainly by the ovaries before menopause and by other tissues, including the placenta during pregnancy.

## Oestrogen Replacement Therapy

Treatment with estrogen, sometimes combined with

a progestogen, to relieve symptoms caused by low estrogen such as hot flushes and vaginal dryness. Benefits and risks depend on age, medical history, uterus status, dose, and route, so treatment is individualized.

## Oestrogens

A group of steroid hormones involved in reproductive development, menstrual cycling, pregnancy, bone, and other physiological functions.

## Oestrone

A form of estrogen made by the ovaries and by conversion of other hormones in body tissues. It becomes the main estrogen after menopause and affects reproductive tissues, bones, blood vessels, and metabolism.

## Ofatumumab

A monoclonal antibody against CD20 used for selected blood cancers and relapsing multiple sclerosis; it can increase infection risk and reduce B cells.

## Off-Label

Use of an approved medicine for an indication, dose, route, or population not specified in its regulatory approval.

## Off-Label Use

Use of an approved medicine for a condition, dose, route, or age group not specifically included in its official approval. A clinician may choose it when evidence and expected benefits support its use.

## Offspring

The young or descendants produced by one organism or by a pair. In medicine and genetics, offspring may be studied to understand inherited traits, pregnancy outcomes, developmental conditions, and disease risk.

## Ofloxacin

A fluoroquinolone antibiotic used for selected susceptible bacterial infections. It can cause serious tendon, nerve, heart-rhythm, and other adverse ef-

---

fects.

### **Ogilvie'S Syndrome**

Another spelling of Ogilvie syndrome: severe widening of the large intestine without a physical blockage, often after surgery or during serious illness. It needs medical monitoring because pressure can reduce blood flow or cause a bowel perforation.

### **Ogilvie Syndrome**

A dangerous widening of the large intestine without a physical blockage, usually in a seriously ill or recently operated patient. Abdominal swelling, pain, nausea, or failure to pass stool may occur, and the bowel can perforate if untreated.

### **Ohm**

The standard unit of electrical resistance, equal to one volt per ampere. In medical equipment, resistance affects how electrical current flows through devices, sensors, wires, and body tissues.

### **OHSS**

Abbreviation; see *Ovarian Hyperstimulation Syndrome*.

### **Oidium**

A fragment or spore-like fungal element formed by fragmentation of a hypha; the term is also used in older fungal classifications.

### **Oil**

A hydrophobic liquid used as a nutrient, solvent, emollient, or medicinal preparation; safety depends on the specific oil.

### **Oily Hair**

Hair that appears greasy because the scalp produces excess oil or because of product buildup. It may occur with infrequent washing, hormonal changes, or some skin conditions and is not poisoning.

### **Oily Skin**

Skin that produces more oil than usual, often causing shine, enlarged-looking pores, or acne. Gentle

cleansing and non-comedogenic products may help; persistent acne or irritation may need medical assessment.

### **Ointment**

A thick, semisolid preparation applied to the skin, inside the nose, or other body surfaces. Its oily base helps protect the area and keep it moist; some ointments also contain medicines that treat infection, inflammation, pain, or other conditions.

### **Ointment Bases**

Semisolid materials that carry medicines applied to the skin or moist body surfaces. They affect how easily a product spreads, how much moisture it retains, how much medicine enters tissue, and how long it remains stable.

### **Okadaic Acid**

A toxin made by certain marine organisms that can accumulate in shellfish. Eating contaminated shellfish may cause diarrhoea and vomiting; in laboratories it is used to study enzymes that remove phosphate groups from proteins.

### **Okihiro Syndrome**

A rare inherited condition caused by changes in the SALL4 gene. It may combine abnormalities of the thumb or forearm with difficulty moving the eyes, and may also affect hearing, the heart, kidneys, or spine.

### **Olanzapine**

Olanzapine is an atypical antipsychotic that blocks dopamine and serotonin receptors and treats schizophrenia and bipolar disorder; weight gain, diabetes, dyslipidaemia, sedation, and movement disorders are important risks.

### **Olaparib**

Olaparib is a PARP inhibitor used for selected cancers with BRCA or other homologous-recombination repair defects; anaemia, nausea, fatigue, myelosuppression, and rare myelodysplastic syndrome or acute myeloid leukaemia may occur.

---

## Old Age

Later life; health assessment should consider function, comorbidity, cognition, and social circumstances rather than age alone.

## Oleander Poisoning

Poisoning from eating oleander, which contains toxins that affect the heart. Nausea, vomiting, dizziness, confusion, slow or irregular heartbeat, fainting, or seizures may occur.

## Olecranon

The prominent bony process at the proximal end of the ulna forming the point of the elbow.

## Olecranon Bursitis

Inflammation of the subcutaneous synovial bursa overlying the olecranon process of the ulna, presenting with fluctuant, localized swelling over the posterior elbow, provoked by repetitive trauma ("student's elbow"), gout, or bacterial infection (septic bursitis).

### Alternate Names:

Student Elbow; Miner Elbow

## Olecranon Fossa

A deep triangular depression on the posterior surface of the distal humerus superior to the trochlea that accommodates the curved olecranon process of the ulna during full elbow extension.

## Oleic Acid

A monounsaturated fat found in olive oil, nuts, seeds, avocados, and many other foods. Replacing some saturated fats with it may support healthier blood cholesterol levels as part of a balanced diet.

## Oleum

A very corrosive mixture of sulfur trioxide and sulfuric acid, used industrially and dangerous to skin, eyes, and lungs.

## Olfaction

The sense of smell. Odor molecules stimulate special cells in the upper part of the nose, which send signals

to the brain to identify and perceive odors.

## Olfactory

Relating to the sense of smell or the nerves and brain pathways that process odors. Problems in this system can cause reduced smell, loss of smell, distorted smells, or unusual odor sensations.

## Olfactory Bulb

The olfactory bulb is a forebrain structure that receives signals from nasal smell receptors and begins processing odor information.

## Olfactory Cortex

The olfactory cortex is a group of brain regions that interprets smell signals and links them with memory, emotion, and behavior.

## Olfactory Gyrus

The olfactory gyrus is a cortical region associated with processing olfactory information and includes areas near the medial temporal and frontal lobes.

## Olfactory Learning

Olfactory learning is acquisition or modification of behavior or memory through exposure to odors and their associations.

## Olfactory Mucosa

The olfactory mucosa is specialized nasal lining containing receptor neurons, supporting cells, and glands that detect odor molecules.

## Olfactory Nerve

The first cranial nerve, which carries smell signals from specialized cells high in the nose to the brain. Injury or disease can reduce or eliminate the sense of smell or distort odors.

## Olfactory Neuroblastoma

A rare malignant tumor arising in the upper nasal cavity near the olfactory nerves, often causing nasal blockage, nosebleeds, or reduced smell.



---

## Olfactory Tract

The olfactory tract is a bundle of nerve fibers carrying smell information from the olfactory bulb to cerebral regions involved in odor perception.

## Olfactory Tubercle

The olfactory tubercle is a ventral forebrain region involved in odor processing, reward, motivation, and integration with limbic circuits.

## Oligoasthenoteratozoospermia

A semen abnormality in which sperm numbers are low, movement is reduced, and fewer sperm have a typical shape. It can reduce fertility and may result from testicular disease, hormone problems, heat, medicines, smoking, or an unknown cause. Repeat semen testing and evaluation help guide treatment or assisted reproduction.

### Alternate Names:

OAT Syndrome; OAT

## Oligoclonal Bands

Groups of immune proteins found in blood or the fluid surrounding the brain and spinal cord. Bands found only in that fluid suggest immune activity inside the nervous system, but they can occur in multiple sclerosis, infections, and other inflammatory conditions.

## Oligocythaemia

An unusually low number of blood cells, especially red blood cells. The uncommon term may describe anemia or another low blood-cell count, so the specific cell type and cause must be identified with a blood test.

## Oligodendrocyte

A supporting cell in the central nervous system that forms myelin around axons. Myelin helps nerve signals travel quickly; oligodendrocytes can be damaged in disorders such as multiple sclerosis.

## Oligodendroglia

Oligodendroglia are glial cells of the central nervous system that produce myelin around axons and

support nerve conduction.

## Oligodendroglial Neoplasm

A brain tumor made of cells that resemble oligodendrocytes, the cells that form the insulating coating around nerve fibers. Diagnosis uses tissue examination and genetic testing, and the tumor's behavior depends on its grade and molecular features.

## Oligodendroglioma

A brain tumor defined by characteristic changes in the IDH gene and loss of parts of chromosomes 1 and 19. It usually begins in the cerebral hemispheres and may cause seizures, headache, or other neurologic symptoms; grade affects its behavior and treatment.

## Oligodontia

A condition present from birth in which many permanent teeth fail to develop, usually more than six excluding wisdom teeth. It may affect chewing, speech, appearance, and jaw development, requiring dental replacement or orthodontic care.

## Oligogenic Inheritance

A pattern in which changes in a small number of genes work together to influence a disease or characteristic. The combination and strength of the changes can make symptoms differ among affected people.

## Oligohydramnios

Too little amniotic fluid around the baby during pregnancy. It may result from leaking membranes, placental problems, dehydration, a pregnancy that continues beyond the due date, or problems affecting the baby's kidneys.

## Oligomenorrhea

Oligomenorrhea is infrequent or widely spaced menstrual bleeding, often from anovulation, polycystic ovary syndrome, endocrine disease, undernutrition, excessive exercise, stress, or perimenopause.

## Oligomenorrhoea

Menstrual periods that occur less often than ex-

---

pected, usually with cycles longer than about 35 days. Causes include pregnancy, polycystic ovary syndrome, thyroid disease, stress, low body weight, intense exercise, and some medicines; evaluation depends on age and symptoms.

### **Oligometastasis**

Cancer that has spread to only a small number of separate sites, often no more than five. In selected people, surgery, focused radiation, or another local treatment may control these deposits for a long time.

### **Oligonucleotide**

A short piece of DNA or RNA made in the laboratory or found in cells. It can be used to detect genetic material, block or change gene messages, or help read and study genes.

### **Oligopeptide**

A short chain of amino acids, usually containing only a few to about 20 building blocks. Oligopeptides can act as hormones, signaling molecules, antimicrobial substances, or parts of larger proteins.

### **Oligosaccharide**

A short chain of sugar molecules joined together. Oligosaccharides form parts of cell-surface molecules, affect how cells recognize one another, and occur naturally in foods where some are used by gut bacteria.

### **Oligospermia**

A lower-than-expected sperm concentration in semen. It can reduce fertility and may result from hormone problems, infections, medicines, heat, toxins, varicoceles, genetic conditions, or problems with sperm production.

### **Oliguria**

Abnormally low urine production. It may result from dehydration, low blood flow, blockage, kidney disease, or medicines and can be an early sign of sudden kidney injury.

### **Olivary**

Relating to the olive-shaped inferior olivary nucleus of the medulla or to an olive-like structure.

### **Olivary Nucleus**

The olivary nucleus is a brainstem structure that sends timing and motor-learning signals to the cerebellum through climbing fibers.

### **Olive Pontocerebellar Atrophy**

A group of progressive brain disorders that damage areas controlling balance and coordination. They may cause worsening unsteadiness, clumsy movements, slurred speech, stiffness, or other neurologic problems.

### **Olivopontocerebellar Atrophy**

Olivopontocerebellar atrophy is a degenerative condition affecting the inferior olives, pons, and cerebellum, causing progressive ataxia and other neurologic dysfunction.

### **Ollier Disease**

A rare, non-hereditary skeletal dysplasia characterized by multiple benign cartilaginous enchondromas developing predominantly in the metaphyses and diaphyses of long bones with marked asymmetrical skeletal involvement and significant risk of malignant chondrosarcoma transformation.

#### **Alternate Names:**

Multiple Enchondromatosis; Dyschondroplasia

### **Ollier's Disease**

A nonhereditary skeletal disorder with multiple enchondromas, causing deformity, limb-length discrepancy, and an increased risk of malignancy.

### **Olmesartan Medoxomil**

An oral prodrug converted to olmesartan, an angiotensin II receptor blocker used to treat high blood pressure.

### **Olopatadine Hydrochloride**

Olopatadine hydrochloride is an antihistamine used in eye drops or nasal spray to treat allergic conjunc-

---

tivitis or rhinitis.

## **OLP**

Abbreviation; see *Oral Lichen Planus*.

## **Olsenella**

Olsenella is a genus of anaerobic bacteria found mainly in the mouth and digestive tract; human infection is uncommon.

## **Olsenella Uli**

An anaerobic gram-positive bacterium found in the mouth and associated with periodontal disease; invasive infection is rare.

## **Omacetaxine Mepesuccinate**

Omacetaxine mepesuccinate is a protein-synthesis inhibitor used for selected chronic myeloid leukemia resistant or intolerant to tyrosine-kinase inhibitors.

## **Omalizumab**

Omalizumab is a monoclonal antibody that binds circulating IgE and reduces allergic inflammation; it is used for selected allergic asthma, chronic urticaria, nasal polyps, and food allergy, with rare anaphylaxis.

## **Omasum**

The omasum is the third compartment of a ruminant stomach, absorbing water and small particles before contents reach the abomasum.

## **Omega-3**

A family of polyunsaturated fats found in fish, flaxseed, walnuts, and some supplements, supporting heart and brain health.

## **Omega-3 Fatty Acid**

A type of polyunsaturated fat found in fish, seafood, flaxseed, chia, walnuts, and some oils. Omega-3s support cell membranes and cardiovascular function; supplements can interact with medicines and are not a substitute for treatment.

## **Omega-3 Fatty Acids**

A family of fats found in oily fish, seafood, some seeds, nuts, and plant oils. EPA and DHA support cell and eye function, while ALA is a plant form; supplements are not appropriate for everyone and can interact with medicines.

## **Omega-6**

A family of essential polyunsaturated fats, found in vegetable oils, nuts, seeds, and many prepared foods. The body uses them for cell membranes and signaling. Linoleic acid is the main dietary form; replacing saturated fats with unsaturated fats generally supports heart health.

## **Omega-6 Fatty Acids**

Essential polyunsaturated fats found in plant oils, nuts, seeds, and many prepared foods. The body needs them for cell membranes and signaling, but overall health depends on balanced intake with other fats.

### **Alternate Names:**

n-6 Fatty Acids; Omega-6 Polyunsaturated Fatty Acids

## **Omega-N-Methylarginine**

Omega-N-methylarginine is a methylated arginine compound that can inhibit nitric-oxide synthase and affect vascular signaling.

## **Omega Sign**

A characteristic facial expression observed in patients with severe melancholic depression, produced by tonic contraction of the corrugator supercilii and procerus muscles forming wrinkles and furrows on the forehead resembling the Greek letter omega.

### **Alternate Names:**

Veraguth's Fold; Depressive Facial Omega

## **Omenn Syndrome**

A severe inherited immune deficiency that usually begins in infancy. Babies may develop widespread red, peeling skin, swollen lymph nodes, enlarged liver or spleen, diarrhea, poor growth, and repeated serious infections.

---

## Omental Caking

The radiographic (CT) and macroscopic surgical appearance of a thickened, diffuse, infiltrative nodular mass replacing the greater omentum, resulting from extensive peritoneal carcinomatosis (most commonly originating from ovarian, gastric, pancreatic, or colonic adenocarcinomas).

### Alternate Names:

Peritoneal Carcinomatosis; Metastatic Omental Thickening

## Omentectomy

Removal of part or all of the greater omentum, commonly performed during staging and cytoreduction for ovarian and other intra-abdominal cancers.

## Omentum

A fold of fatty tissue inside the abdomen that hangs from the stomach and partly covers the intestines. It contains blood vessels and immune cells and can help limit the spread of infection or inflammation.

## Omeprazole

Omeprazole is a proton-pump inhibitor that irreversibly suppresses gastric H<sup>+</sup>/K<sup>+</sup>-ATPase; it treats reflux and peptic-ulcer disease and supports *Helicobacter pylori* eradication, while prolonged use may affect magnesium, B12, bone, and infection risk.

## Ommaya Reservoir

A surgically implanted reservoir connected to a catheter in a brain ventricle, used to sample cerebrospinal fluid or deliver selected medicines.

## Omphalitis

Infection of a newborn's umbilical stump, usually during the first weeks of life. Redness spreading onto the skin, swelling, pus, bad smell, fever, poor feeding, or unusual sleepiness may occur, and infection can spread rapidly.

## Omphalocele

A birth defect in which abdominal organs protrude through an opening at the navel into a protective sac.

Other chromosome or organ abnormalities may occur, so newborn assessment and imaging are needed.

## Omphalocele Repair

Treatment to place an infant's abdominal organs back inside the abdomen and close the opening in the abdominal wall. Repair may be immediate or staged according to the size of the defect and the baby's breathing and circulation.

## Onchocerca Volvulus

A parasitic roundworm spread by repeated bites of infected blackflies that breed near fast-flowing rivers. Adult worms live in lumps under the skin and release young worms that can migrate through the skin and eyes.

## Onchocerciasis

A parasitic infection caused by *Onchocerca volvulus* and spread by infected blackflies. It can cause intensely itchy skin, nodules, skin changes, and eye inflammation that may lead to permanent visual impairment or blindness.

## Oncocytic Neoplasm

A growth made of cells containing unusually many energy-producing structures, giving them a granular appearance under a microscope. It may be noncancerous or cancerous, depending on the organ and the tumor's other features.

## Oncocytoma

A usually noncancerous tumor made of cells containing many energy-producing structures. It most often develops in the kidney but can occur in other organs; testing is needed to distinguish it from cancer.

## Oncogene

A gene that can make cells grow or survive without normal control when it is changed, overactive, or present in too many copies. Oncogene changes can contribute to the development and progression of cancer.

---

## **Oncogenic**

Capable of causing or promoting cancer. An oncogenic change, virus, gene, or chemical can increase abnormal cell growth or reduce normal control of cell division.

## **Oncogenicity**

The ability of a substance, infection, genetic change, or other factor to contribute to the development of cancer.

## **Oncogenic Osteomalacia**

A rare acquired paraneoplastic metabolic bone disorder caused by small, benign mesenchymal tumors that oversecrete fibroblast growth factor 23 (FGF23), which induces renal phosphate wasting, severe hypophosphatemia, suppressed 1,25-dihydroxyvitamin D production, and defective osteoid mineralization.

### **Alternate Names:**

Tumor-Induced Osteomalacia; TIO

## **Oncogenic Virus**

A virus capable of contributing to cancer by altering cell growth, suppressing tumor defenses, or integrating or maintaining oncogenic genetic material.

## **Oncologic Emergency**

A cancer-related problem that can quickly threaten life, organ function, or the nervous system and needs urgent treatment. Examples include spinal-cord pressure, blocked major chest veins, dangerously high calcium, tumor breakdown, airway blockage, and infection during low white-cell counts.

## **Oncologist**

A doctor who specializes in cancer care. Medical oncologists use medicines, radiation oncologists use radiation, and surgical oncologists operate; many people receive care from more than one type of oncologist.

## **Oncology**

The medical field concerned with preventing, finding, and treating cancer and supporting people after

treatment. Oncologists may use surgery, radiation, medicines, or combinations of treatments and work with other professionals to provide symptom and emotional support.

## **Oncology Nursing**

Oncology nursing provides cancer-related assessment, treatment support, symptom control, education, coordination, survivorship, and end-of-life care.

## **Oncology Nutrition**

Nutritional care for people with cancer, addressing intake, symptoms, treatment effects, weight, muscle mass, and personal goals.

### **Alternate Names:**

Cancer Nutrition Support; Oncology Nutrition

## **Oncology PET**

A positron-emission tomography scan, often combined with CT, that uses a small radioactive tracer to show active cancer in the body. It may help assess spread, treatment response, or recurrence, but inflammation can also appear active.

## **Oncology Rehabilitation**

Healthcare that helps people prevent or manage cancer-related problems before, during, or after treatment. It may address fatigue, pain, weakness, nerve symptoms, swelling, movement, nutrition, work, daily activities, and emotional wellbeing.

## **Oncolytic Virotherapy**

Oncolytic virotherapy uses modified viruses that preferentially infect and destroy cancer cells and may stimulate an immune response against the tumor.

## **Oncolytic Virus**

A virus selected or modified to infect and destroy cancer cells more readily than normal cells. It may also stimulate the immune system to attack the tumor; use is limited to selected cancers and requires specialist treatment.

---

## Oncolytic Viruses

Viruses selected or engineered to infect and destroy cancer cells while stimulating an immune response against the tumor.

## Oncoprotein

A protein that promotes abnormal cell growth or survival and can contribute to cancer. It may result from a cancer-causing gene, a mutation, a chromosomal change, or infection with certain viruses.

## Oncosis

A form of cell injury in which cells swell and may die, usually because severe lack of oxygen or another injury disrupts energy production and the movement of salt and water across the cell membrane.

## Oncotic Pressure

The pulling force created by proteins in the blood, mainly albumin, that helps keep fluid inside blood vessels. When blood protein is low, fluid may leak into tissues and cause swelling.

## Oncotoxic

Harmful to cancer cells or relating to substances that damage tumors. An oncotoxic treatment may also affect healthy cells, so its benefits, side effects, dose, and safety require careful medical monitoring.

## Ondansetron

Ondansetron is a selective 5-HT<sub>3</sub>-receptor antagonist used to prevent nausea and vomiting after chemotherapy, radiotherapy, surgery, or other triggers; constipation, headache, and QT prolongation can occur.

## Ondansetron Hydrochloride

The hydrochloride salt of ondansetron, a serotonin 5-HT<sub>3</sub> receptor blocker used to prevent nausea and vomiting.

## Ondine Curse

A life-threatening autonomic disorder of respiratory control (congenital central hypoventilation syndrome) caused by mutations in the PHOX2B gene,

characterized by absent or severely blunted autonomic ventilatory responses to hypercapnia and hypoxemia, resulting in severe sleep-induced respiratory failure.

### Alternate Names:

Congenital Central Hypoventilation Syndrome; CCHS

## Ondine's Curse

A rare disorder in which automatic breathing becomes inadequate, especially during sleep, because brain signals fail to drive breathing normally. It can cause dangerously low oxygen and may require assisted ventilation.

### Alternate Names:

CCHS; Congenital Central Hypoventilation Syndrome

## One-Hour Glucose Challenge Test

A screening test for diabetes during pregnancy, usually performed at 24–28 weeks. After drinking a measured glucose solution, blood sugar is checked one hour later. A high result does not diagnose diabetes but usually leads to a longer confirmatory test.

## Oneirophrenia

A dreamlike state with confusion or impaired reality testing; the term is uncommon and does not represent a specific modern psychiatric diagnosis.

## Onset

Onset is the time or manner in which a symptom, disease, or physiologic change begins, such as sudden, gradual, acute, or insidious onset.

## Onychia

Inflammation or infection of the nail or nail-producing tissue. It can cause redness, swelling, pain, pus, or changes in nail growth and may result from bacteria, fungi, injury, or skin disease. Treatment depends on the cause.

## Onychocryptosis

Onychocryptosis is an ingrown nail, usually at the



---

great toe, in which the nail edge penetrates surrounding skin and causes pain or inflammation.

### **Onychogryphosis**

Severe thickening and overgrowth of a nail, which becomes curved and may resemble a claw. It is more common in older adults and can follow injury, fungal infection, poor circulation, or difficulty caring for the feet; it may cause pain in shoes.

### **Onycholysis**

Separation of the nail plate from the underlying nail bed, beginning distally or laterally. Causes include trauma, psoriasis, fungal infection, thyroid disease, medications, and chemical exposure; the underlying cause should be identified.

### **Onychomadesis**

Onychomadesis is proximal separation and shedding of a nail caused by temporary arrest of nail-matrix growth, often after infection, systemic illness, trauma, or medication exposure.

### **Onychomycosis**

A fungal infection of a nail, more common in toenails than fingernails. The nail may become thick, yellow or white, brittle, crumbly, or separated from the skin. A sample may be tested before prolonged antifungal treatment because other nail diseases look similar.

### **Oocyte**

An immature egg cell in an ovary. It develops before birth, remains paused until a menstrual cycle, and may mature and be released for possible fertilization.

### **Oocyte Cryopreservation**

Freezing unfertilized eggs for possible use in future pregnancy. Hormone medicines stimulate the ovaries, and the eggs are collected through the vagina with a needle before being rapidly frozen. Eggs may be stored before cancer treatment or for other fertility-preservation reasons, but freezing does not guarantee a future pregnancy.

#### **Alternate Names:**

Egg Freezing; Oocyte Vitrification

### **Oocyte Retrieval**

A procedure used in assisted reproduction to collect mature eggs from the ovaries. After hormone stimulation, a clinician guides a thin needle through the vagina using ultrasound and gently draws fluid containing the eggs into a container for laboratory fertilization.

#### **Alternate Names:**

OPU; Egg Retrieval; Follicular Aspiration

### **Oogenesis**

The formation and maturation of egg cells, or ova, in the ovaries. It begins before birth and continues through the reproductive years, with changes during each menstrual cycle.

### **Oogonium**

An oogonium is an early female germ cell that proliferates during embryonic development and gives rise to primary oocytes.

### **Oophorectomy**

Surgery to remove one or both ovaries. It may treat cancer, a cyst, infection, torsion, or reduce inherited cancer risk; removing both ovaries before natural menopause causes an immediate fall in ovarian hormones.

### **Oophoritis**

Inflammation of an ovary, usually associated with pelvic infection or, less commonly, autoimmune disease. It may cause lower abdominal or pelvic pain, fever, abnormal discharge, or menstrual changes; severe pain requires assessment to exclude ovarian torsion or abscess.

### **Ooplasm**

Ooplasm is the cytoplasm of an egg cell, containing organelles, nutrients, and molecular factors that support fertilization and early embryo development.

### **Opacity**

Clouding of a normally clear part of the eye, such as the cornea, lens, or fluid inside the eye. It can blur vision and may result from injury, infection,

---

inflammation, swelling, or age-related change.

### **Open Abdominal Aortic Aneurysm Repair**

Open surgery to repair a widened, weakened area of the abdominal aorta. The surgeon reaches the aorta through the abdomen and replaces the damaged section with a strong artificial tube.

#### **Alternate Names:**

Open AAA Repair; Open Abdominal Aortic Aneurysm Repair; Surgical AAA Repair

### **Open Aortic Valve Surgery**

Conventional surgical repair or replacement of a diseased aortic valve performed through a full median sternotomy with cardiopulmonary bypass and cardioplegic arrest.

#### **Alternate Names:**

Surgical Aortic Valve Replacement; SAVR; Open Aortic Valve Replacement

### **Open-Angle Glaucoma**

A type of glaucoma in which the eye's drainage angle remains open, but fluid drains too slowly. Pressure may rise and damage the optic nerve, causing gradual, permanent vision loss, often without early symptoms.

### **Open Bite**

An open bite is a malocclusion in which upper and lower teeth do not meet in part of the mouth when the jaws are closed.

### **Open-Book Fracture**

A severe pelvic injury in which the front or back pelvic joints separate, making the pelvis widen like an open book. It usually follows major trauma and can cause life-threatening internal bleeding or organ injury.

### **Open Esophagectomy**

Conventional surgical resection of the esophagus performed through larger open thoracic and abdominal incisions (such as an Ivor Lewis or McKeown approach) for advanced esophageal tumors.

#### **Alternate Names:**

Open Esophageal Resection; Transthoracic Esophagectomy

### **Open Fracture**

A fracture associated with a wound that communicates between the fracture site and the external environment. It carries increased risks of infection and tissue damage.

#### **Alternate Names:**

Open Fracture

### **Open Fractures**

Broken bones associated with a wound that connects the fracture to the outside air. They have a high risk of infection and tissue damage; treatment includes cleaning, antibiotics, and specialist care.

### **Open Gallbladder Removal**

Surgery to remove the gallbladder through one larger abdominal incision. It may be needed when keyhole surgery is unsafe or cannot be completed; bleeding, infection, bile leakage, and injury to nearby ducts or organs are possible.

### **Open-Heart Surgery**

Surgery in which the chest is opened to repair or treat the heart or its major blood vessels. Some operations use a heart–lung machine while the heart is temporarily stopped; risks depend on the operation and the person's health.

### **Opening Snap**

A sharp sound heard soon after the heart starts to relax, caused by a stiff mitral valve opening. It suggests mitral valve narrowing and must be interpreted with the rest of the heart examination.

#### **Alternate Names:**

Mitral Opening Snap; OS Sound

### **Open Lung Biopsy**

An open lung biopsy removes a small piece of lung through a surgical chest incision for microscopic diagnosis when less invasive tests are insufficient.

---

## Open Mitral Valve Surgery

Conventional cardiac surgical repair or mechanical/bioprosthetic replacement of the mitral valve performed via a median sternotomy with cardiopulmonary bypass. Indicated for severe mitral regurgitation or mitral stenosis when percutaneous or minimally invasive approaches are not feasible.

### Alternate Names:

Open Mitral Repair; Open Mitral Valve Replacement; Sternotomy Mitral Surgery

## Open Pleural Biopsy

An open pleural biopsy removes a sample of the lining around the lung through surgery to investigate cancer, infection, or unexplained pleural disease.

## Open Prostatectomy

Surgery through an abdominal incision to remove the part of an enlarged prostate that blocks urine flow. It can improve severe urinary obstruction but may cause bleeding, infection, urine leakage, or changes in sexual and urinary function.

## Open Reduction And Internal Fixation

Surgery to expose a broken bone, put the pieces back into position, and hold them with plates, screws, wires, or rods. It is used when a cast or other nonsurgical treatment cannot provide safe alignment or stability.

### Alternate Names:

ORIF

## Open Surgery

Surgery performed through a relatively large incision that allows the surgeon to see and reach the affected area directly. It is used when a larger working space is needed, although recovery may be longer than after some keyhole procedures.

## Open Surgical Wound Care

Care of a surgical wound left partly or completely open to drain or heal gradually, usually with dressings and scheduled assessment for infection, bleeding, and healthy tissue growth.

## Operable

Able to be treated with surgery because the disease can be safely reached or removed.

## Operant Conditioning

Learning in which the likelihood of a behavior changes because of its consequences, such as reinforcement or punishment.

## Operating Room

An operating room is a controlled sterile or semisterile clinical area equipped for surgical procedures, anesthesia, monitoring, and recovery support.

## Operating Tables

Adjustable tables designed to support and position patients during surgery or other procedures while allowing access and maintaining safety.

## Operation

A planned surgical procedure used to diagnose, repair, remove, replace, or improve part of the body. It may be performed through an incision or with instruments inserted through small openings, depending on the condition and goal.

## Operative Dentistry

The branch of dentistry that repairs teeth damaged by decay, fractures, wear, or developmental defects. Treatment may include fillings, bonding, reshaping, and other procedures that restore appearance, strength, and function.

## Operative Vaginal Delivery

Vaginal birth assisted with a vacuum cup or forceps when the baby's position is suitable and birth needs to be completed promptly. Reasons may include prolonged pushing or concern about the baby's condition; tears, bleeding, or injury to the baby can occur.

## Operator

A person who performs a procedure or operates equipment; in healthcare, training and authorization are essential.

---

## **Operculum**

A lid, covering, or flap-like structure; in anatomy it may cover an opening or cortical region.

## **Ophthalmia**

Inflammation of the eye, especially the conjunctiva or other surface tissues. It may cause redness, discharge, pain, light sensitivity, or blurred vision and can result from infection, allergy, injury, or inflammation.

## **Ophthalmia Neonatorum**

Inflammation or infection of a newborn baby's eye during the first weeks of life. It may be caused by bacteria passed during birth, other infections, or chemical irritation; redness, swelling, or pus may occur and can threaten the cornea and vision.

## **Ophthalmic**

Relating to the eye. An ophthalmic medicine, drop, ointment, or device is designed for use on or around the eye and must be used as directed to prevent injury or infection.

## **Ophthalmic Emergency**

An eye problem that needs immediate assessment to prevent permanent vision loss or serious injury. Examples include sudden vision loss, retinal detachment, severe eye infection, chemical exposure, a penetrating injury, or rapidly painful high eye pressure.

## **Ophthalmic Solution**

A sterile liquid medicine made for application to the eye. It must remain clean and should not be shared or used after contamination because contaminated drops can cause serious eye infections.

## **Ophthalmitis**

Inflammation of the eye caused by infection, immune disease, injury, or surgery. It may cause redness, pain, light sensitivity, discharge, or blurred vision and needs prompt assessment when severe.

## **Ophthalmologist**

A medical doctor trained to diagnose and treat eye disease and vision problems. Ophthalmologists can prescribe medicines and glasses, perform eye procedures and surgery, and manage conditions affecting the eye, optic nerve, and visual system.

## **Ophthalmology**

The medical specialty concerned with the eyes, vision, and the parts of the nervous system used for seeing. It includes eye examinations, treatment of eye disease, vision correction, and procedures or surgery when needed.

## **Ophthalmomyiasis**

Infestation of the eye by fly larvae. Larvae may remain on the eye surface or enter deeper tissues, causing intense irritation, pain, inflammation, or vision loss; prompt examination is needed for safe removal and treatment.

## **Ophthalmoplegia**

Weakness or paralysis of one or more muscles that move the eyes, causing limited eye movement, double vision, or an abnormal eye position. Causes include nerve disease, stroke, diabetes, thyroid disease, injury, and muscle disorders.

## **Ophthalmoscope**

A lighted instrument used to examine the back and internal structures of the eye, including the retina, macula, optic nerve, and retinal blood vessels. Direct ophthalmoscopes provide a magnified view through the pupil, while indirect ophthalmoscopes use a separate light and lens to provide a wider view. Findings may relate to diabetes, high blood pressure, glaucoma, infection, inflammation, or other eye and nervous-system disorders.

## **Ophthalmoscopy**

Examination of the inside of the eye with a lighted instrument. It allows a clinician to inspect the retina, optic nerve, and blood vessels for changes caused by diabetes, high blood pressure, glaucoma, inflammation, or other disease.

---

## Opiate

A drug derived directly from chemicals in the opium poppy, such as morphine or codeine, or a closely related semisynthetic form. Opiates can relieve pain but may cause sleepiness, constipation, dependence, and dangerously slow breathing.

## Opiate Alkaloid

An opiate alkaloid is a naturally occurring alkaloid from the opium poppy, such as morphine or codeine, with opioid analgesic and dependence potential.

## Opiate And Opioid Withdrawal

Opioid withdrawal is a predictable syndrome after reducing or stopping opioid exposure, producing anxiety, sweating, muscle aches, diarrhea, abdominal cramps, nausea, and autonomic symptoms.

## Opiate Assay

A laboratory test that detects or measures opiates or their breakdown products in urine, blood, saliva, or another specimen. Results may support clinical care or toxicology testing, but some synthetic opioids require separate tests.

## Opiate Overdose

A life-threatening amount of an opioid medicine or drug that slows brain and breathing activity. Extreme sleepiness, pinpoint pupils, slow or absent breathing, blue lips, or inability to wake may occur. Naloxone can sometimes restore breathing.

## Opiate Overdose Triad

The common combination of extreme sleepiness or unconsciousness, very slow or stopped breathing, and very small pupils after an opioid overdose. Not every person has all three signs. Naloxone may restore breathing.

## Opioid

A natural or synthetic substance that acts on opioid receptors to relieve pain and may cause sedation or dependence.

## Opioid Agonist Treatment

Treatment for opioid use disorder using a prescribed medicine, usually methadone or buprenorphine, that activates opioid receptors in a controlled way. It reduces withdrawal and cravings, lowers illicit opioid use and overdose risk, and works best with ongoing medical support.

## Opioid Analgesic Medications

A class of potent analgesic agents that bind to central and peripheral  $\mu$ -opioid receptors to modulate nociceptive transmission. Used for moderate-to-severe acute pain and cancer-related pain; carries significant risks of respiratory depression, sedation, physical dependence, tolerance, and constipation.

### Alternate Names:

Opioid Analgesics; Narcotic Analgesics; Opiate Pain Relievers

### Alternate Names:

Opioid Analgesics; Narcotic Analgesics

## Opioid Intoxication

Opioid intoxication can cause euphoria, drowsiness, slowed breathing, pinpoint pupils, impaired coordination, coma, and death.

## Opioid Overdose

A medical emergency characterized by the classic triad of respiratory depression, miosis (pinpoint pupils), and decreased level of consciousness. It can progress to respiratory arrest, cardiovascular collapse, and death.

## Opioid Poisoning

Alternate name; see *Opioid Toxicity*.

## Opioid Receptor

A cell-surface protein activated or blocked by opioids and naturally produced endorphins. Mu, kappa, and delta receptors influence pain, breathing, mood, bowel movement, and reward; excessive mu-receptor activation can suppress breathing.

## Opioid Reversal

Emergency treatment of opioid toxicity with nalox-

---

one, a competitive opioid-receptor antagonist. It is titrated to restore adequate breathing, with repeated doses or monitoring sometimes needed because its effect may wear off before the opioid.

## Opioids

Medicines and drugs that act on opioid receptors to relieve pain. They can cause sleepiness, constipation, dependence, and life-threatening breathing suppression, especially with alcohol or sedatives; overdose may be reversed with naloxone.

## Opioid Toxicity

Toxic effects of opioid agonists, classically respiratory depression, central nervous system depression, and miosis. Management prioritizes ventilation and airway support; naloxone is titrated to restore adequate breathing while recognizing the risk of recurrent toxicity and withdrawal.

## Opioid Toxidrome

A poisoning pattern caused by opioid excess, characterized by respiratory depression, reduced consciousness, and small pupils; naloxone and ventilatory support may be required.

## Opioid Use Disorder

A problematic pattern of opioid use that causes clinically significant impairment or distress. It is diagnosed when at least two of 11 symptoms occur within 12 months, such as inability to cut down, craving, hazardous use, or continued use despite harm. It may involve prescribed or illicit opioids; tolerance or withdrawal alone does not establish the disorder when opioids are taken solely under appropriate medical supervision. Effective treatment includes medication, such as buprenorphine, methadone, or naltrexone, with counseling or other recovery support as needed.

**Alternate Names:**

OD

## Opioid Withdrawal

A characteristic group of symptoms occurring after abrupt reduction or cessation of prolonged opioid use, including anxiety, muscle aches, sweating, ab-

dominal cramps, diarrhea, nausea, and insomnia.

## Opisthorchis Felineus

A liver fluke acquired by eating raw or undercooked freshwater fish; chronic infection can cause bile-duct inflammation and increase cholangiocarcinoma risk.

## Opisthotonos

A severe, abnormal posture characterized by extreme hyperextension and rigid backward arching of the head, neck, and spine due to sustained, violent spasm of the axial extensor muscles. In severe cases, the back bridges off the bed so that the body rests entirely on the back of the head and the heels. It results from severe neurological injury or loss of normal central nervous system motor inhibition (such as disinhibition of spinal motor neurons). Hallmark causes include generalized tetanus (due to tetanospasmin blocking inhibitory GABA and glycine release), strychnine poisoning, severe bacterial meningitis, kernicterus (acute bilirubin encephalopathy in newborns), severe brainstem injury, rabies, and acute neuroleptic-induced dystonic reactions. Management requires immediate airway protection, muscle relaxation (such as high-dose benzodiazepines or neuromuscular blockade), and targeted treatment of the underlying cause.

## Opisthotonus

Alternate spelling; see *Opisthotonos*.

**Alternate Names:**

Opisthotonus; Opisthotonic Posturing; Arc de Cercle; Tetanic Posturing

## Opium

The dried latex of *Papaver somniferum* containing opioid alkaloids; it can cause dependence and respiratory depression.

## Oppenheim Sign

A pathological neurological reflex elicited by firmly stroking down the anterior medial aspect of the tibia with the examiner's thumb and knuckles, provoking extension (dorsiflexion) of the great toe and fanning of the other toes in corticospinal upper motor neuron dysfunction.



---

**Alternate Names:**

Oppenheim Reflex; Oppenheim's Sign

**Opponens**

A muscle that moves a body part toward an opposing surface, especially the thumb across the palm.

**Opportunistic Infection**

An infection that causes illness mainly when the immune system is weakened or a normal protective barrier is damaged. Causes may include cancer treatment, HIV, transplantation, severe illness, or prolonged use of immune-suppressing medicines.

**Opportunistic Pathogen**

An infectious microorganism (bacterium, fungus, virus, or parasite) that rarely causes disease in an immunocompetent host with intact natural defense mechanisms, but causes severe, life-threatening infection when host immune defenses are compromised (such as *Pneumocystis jirovecii* in advanced HIV infection).

**Opposition**

Movement of the thumb across the palm so it can touch the fingertips. It is important for grasping and depends on several small movements at the joint where the thumb meets the wrist.

**Oppositional Defiant Disorder**

A childhood or adolescent disorder involving a persistent pattern, lasting at least six months, of angry or irritable mood, argumentative or defiant behavior, or vindictiveness toward an authority figure or other person. The pattern causes clinically significant distress or impairment and is not limited to conflicts with siblings.

**Opsoclonus**

Rapid, involuntary, back-and-forth eye movements in many directions without pauses. It may occur with muscle jerks and poor coordination, sometimes because of an immune reaction, infection, medicine, or an underlying tumor, and needs medical evaluation.

**Opsoclonus-Myoclonus-Ataxia Syndrome**

A rare, devastating neuro-inflammatory condition characterized by the sudden onset of chaotic, involuntary, conjugate, multidirectional saccadic eye movements without intersaccadic intervals (opsoclonus), combined with widespread myoclonic muscle jerking of the limbs and trunk, cerebellar ataxia, and severe encephalopathy. In pediatric patients, it occurs in approximately 50 percent of cases as a paraneoplastic manifestation of an occult neuroblastoma.

**Alternate Names:**

Dancing Eye-Dancing Feet Syndrome; Kinsbourne Syndrome; OMS

**Opsoclonus-Myoclonus Syndrome**

A rare neurologic disorder causing rapid, involuntary eye movements, muscle jerks, poor coordination, and sometimes behavioral or cognitive changes. In children it may be associated with neuroblastoma or an immune reaction.

**Opsonin**

An immune protein that coats a germ and marks it for destruction. Antibodies and complement proteins can act as opsonins by helping infection-fighting cells recognize and engulf the germ.

**Opsonization**

The coating of a germ or other particle with immune proteins so that infection-fighting cells can recognize, capture, and destroy it more easily. Antibodies and complement proteins commonly perform this function.

**Optical Coherence Tomography**

A painless scan that uses reflected light to create detailed cross-sectional pictures of the retina and other eye structures. It helps detect and monitor glaucoma, macular disease, swelling, and other changes without touching the eye.

**Optical Coherence Tomography Angiography**

A scan that uses optical coherence tomography to show blood flow in small vessels of the retina and

---

the layer beneath it, without injecting a dye.

### **Optical Illusion**

A visual perception in which an object, image, or movement appears different from its physical properties. Illusions are common, but new or persistent visual distortions may occur with eye, brain, medicine, or migraine-related disorders.

### **Optical Rotation**

The turning of plane-polarized light as it passes through a substance whose molecules have a particular three-dimensional arrangement. Measuring it can help identify or assess the concentration of some chemicals and medicines.

### **Optic Atrophy**

Damage and loss of optic-nerve fibers, causing permanent reduction in vision, color vision, or part of the visual field. Causes include glaucoma, poor blood supply, inflammation, pressure, injury, toxins, and inherited disorders; treatment targets the cause when possible.

### **Optic Chiasm**

An X-shaped junction beneath the brain where some fibers from the optic nerves cross. Damage can cause characteristic loss of parts of the visual field, depending on the location and cause.

### **Optic Chiasma**

The crossing point beneath the brain where some fibers from the two optic nerves switch sides. Damage there can cause loss of the outer parts of both visual fields.

### **Optic Disc**

The small area at the back of the eye where the optic nerve and blood vessels pass through the retina. It has no light-sensing cells, so it creates the eye's normal blind spot.

### **Optic Disc Drusen**

Calcified, hyaline, extracellular proteinaceous globular deposits located within the pre-laminar optic

nerve head, which exhibit autofluorescence on fundus imaging and produce elevation of the optic disc margin that can mimic true papilledema (pseudopapilledema).

#### **Alternate Names:**

Optic Nerve Head Drusen; Pseudopapilledema Drusen

### **Optic Disc Edema**

Swelling where the optic nerve enters the eye. It may result from raised pressure around the brain, inflammation, poor blood supply, or other eye or nerve disease. New swelling with headache or vision change requires urgent assessment.

### **Optic Disc Swelling**

Enlargement or elevation of the optic nerve head where it enters the eye. Causes range from raised pressure inside the skull to inflammation, ischemia, or local eye disease; sudden vision changes require urgent assessment.

### **Optic Disk**

The point where the optic nerve leaves the retina; it has no photoreceptors and forms the blind spot.

### **Optic Glioma**

A usually slow-growing tumor of the optic nerve or the visual pathway leading to the brain, most often in children. It may cause reduced vision, loss of part of the visual field, a bulging eye, or abnormal eye movements and is associated with neurofibromatosis type 1.

#### **Alternate Names:**

Optic Pathway Glioma; Optic Nerve Glioma

### **Optician**

A technician who fits and dispenses eyeglasses or contact lenses.

### **Optic Nerve**

The nerve that carries visual signals from each eye to the brain. Damage can cause reduced vision, loss of color or part of the visual field, and sometimes pain with eye movement.

---

### Optic Nerve Atrophy

Permanent damage and loss of nerve fibers carrying vision from the eye to the brain. It can reduce sharpness, color vision, or part of the visual field and may follow glaucoma, inflammation, poor blood supply, pressure, injury, or toxins.

### Optic Nerve Disorder

A disease affecting the optic nerve, which carries visual signals from the eye to the brain; it can cause reduced vision, color loss, or visual-field defects.

### Optic Nerve Glioma

Alternate name; see *Optic Glioma*.

### Optic Neuritis

Inflammation or demyelination of the optic nerve, often causing subacute vision loss, impaired color vision, and pain with eye movement. It may be associated with multiple sclerosis or other inflammatory, infectious, or immune-mediated disorders.

### Optic Pathway Glioma

Alternate name; see *Optic Glioma*.

### Optic Tract

A bundle of nerve fibers carrying visual information from the optic chiasma to brain areas that process sight and control visual reflexes. Damage causes loss of a characteristic part of the visual field.

### Optokinetic Nystagmus

Rhythmic eye movements produced when a person views a large moving visual scene, consisting of a slow tracking phase followed by a rapid resetting phase.

### Optometrist

A trained eye-care professional who examines vision and many ocular conditions and prescribes optical correction within local scope of practice.

### Optometry

The healthcare profession focused on examining vision and the eyes, prescribing glasses or contact

lenses, detecting eye disease, and providing treatment or referral within the professional scope allowed by local law.

### Oral

Relating to the mouth, or taken through the mouth. Oral medicines are swallowed and absorbed through the digestive tract unless otherwise specified.

### Oral Administration

Taking a medicine by mouth so it is absorbed through the digestive tract; absorption can be affected by food, vomiting, interactions, or gastrointestinal disease.

### Oral Antidiabetic Agents

Alternate name; see *Antidiabetic Agents*, *Oral*.

### Oral Appliance Therapy

Treatment using a removable dental appliance during sleep, commonly to advance the mandible or retain the tongue and reduce upper-airway collapse in selected obstructive sleep apnea.

### Oral Burn

Thermal, chemical, or electrical injury to the oral mucosal lining, palate, tongue, or lips, resulting in acute pain, erythema, epithelial desquamation, ulceration, and potential upper airway compromise.

#### Alternate Names:

Oral Burn; Oral Cavity Burn; Mouth Burn; Thermal Palatal Burn

### Oral Cancer

Cancer of the lips or tissues inside the mouth, including the tongue, gums, cheeks, floor of the mouth, and palate. Tobacco, alcohol, and areca nut or betel quid increase risk. A persistent sore, lump, red or white patch, pain, or numbness needs dental or medical assessment; HPV is more strongly linked to cancers of the oropharynx.

### Oral Candidiasis

A yeast infection of the mouth, usually caused by *Candida*. It may produce wipeable white patches,

---

soreness, burning, altered taste, or cracks at the mouth corners. Antibiotics, steroid inhalers, dentures, diabetes, dry mouth, and weak immunity increase risk.

### **Oral Cavity**

The space inside the mouth, extending from the lips to the throat and bounded by the cheeks, palate, tongue, and floor of the mouth. It receives food, begins digestion, and helps with speech, taste, and breathing.

### **Oral Cavity Cancer**

Cancer arising in the lips, tongue, gums, inner cheeks, floor of the mouth, or palate. Tobacco, alcohol, betel quid, and some HPV infections increase risk; persistent sores, lumps, patches, pain, or numbness need examination and often biopsy.

### **Oral Cholecystogram**

An older X-ray test in which swallowed contrast collects in the gallbladder, allowing gallstones and gallbladder function to be assessed. Ultrasound and other imaging methods have largely replaced it.

### **Oral Contraceptive**

A birth-control pill containing progestin alone or progestin combined with estrogen. It mainly prevents ovulation and thickens cervical mucus; it does not protect against sexually transmitted infections.

### **Oral Contraceptive Pill**

An oral hormonal contraceptive, usually a combined estrogen–progestin pill or a progestin-only pill, that prevents ovulation and/or changes cervical mucus and the uterine lining.

### **Oral Cutaneous Fistulas**

Abnormal channels connecting the mouth or an oral structure with the skin surface, often caused by dental infection, salivary-gland disease, trauma, surgery, or inflammatory disease. Persistent drainage or swelling requires dental, oral-surgery, or medical evaluation.

### **Oral-Facial-Digital Syndrome**

A group of genetic disorders affecting the mouth, face, fingers, and toes. Features vary and may include oral abnormalities, unusual facial development, shortened or joined digits, and neurologic or kidney problems.

### **Oral Fibromas and Fibromatoses**

Benign fibrous overgrowths in the mouth or jaw region. A solitary oral fibroma is often a firm response to chronic irritation, whereas fibromatoses can infiltrate nearby tissue and recur; persistent lesions may need biopsy.

### **Oral Glucose Tolerance Test**

A test in which a person drinks a measured sugar solution and has blood sugar checked afterward. It helps assess diabetes, prediabetes, or gestational diabetes when other tests do not give a clear answer.

### **Oral Hairy Leukoplakia**

A benign, painless, non-malignant mucocutaneous lesion occurring on the lateral borders of the tongue in immunocompromised individuals (predominantly HIV-positive patients), caused by opportunistic Epstein-Barr virus (EBV) replication, presenting as white, corrugated, non-scrapable vertical folds.

#### **Alternate Names:**

OHL

### **Oral Health**

Oral health is the condition of the teeth, gums, mouth, jaw, and related functions such as eating, speaking, and swallowing.

### **Oral Hemangiomas**

Benign blood-vessel lesions of the lips, tongue, gums, or other oral tissues that may appear red, blue, or purple and bleed after trauma. Evaluation distinguishes them from vascular malformations and other tumors; treatment depends on symptoms and risk.

### **Oral Hemorrhage**

Bleeding from the mouth, gums, tongue, throat, or nearby tissues, caused by trauma, dental disease,

---

infection, medicines, or a bleeding disorder.

### **Oral Herpes**

Herpes simplex infection involving the mouth or lips, most often caused by HSV-1. It may cause no symptoms; when symptoms occur, painful blisters or ulcers may develop. Recurrent cold sores often begin with tingling or burning as the virus reactivates from nerve cells.

#### **Alternate Names:**

Herpes Simplex Labialis; Cold Sores; Fever Blisters; Oral HSV-1 Infection

### **Oral Human Papillomavirus Infection**

Oral human papillomavirus infection affects the mouth or throat; many infections clear without symptoms, while persistent high-risk types can contribute to some cancers.

### **Oral Hygiene**

Practices maintaining oral health: brushing twice daily with fluoride toothpaste, flossing daily, mouthwash (optional), regular dental visits. Proper brushing technique: soft brush, 45-degree angle to gum-line, gentle circular motions. Interdental cleaning essential for plaque removal.

### **Oral Hypoglycemics Overdose**

Taking too much diabetes medicine by mouth can cause dangerously low blood glucose, especially with insulin-releasing medicines. Sweating, shaking, confusion, seizures, or unconsciousness require immediate glucose treatment and emergency advice.

### **Oral Leukoplakia**

A persistent white patch in the mouth that cannot be rubbed off and is not explained by another condition. It is often linked to tobacco, alcohol, or betel-nut use and can sometimes develop into mouth cancer, so a persistent patch needs dental or medical assessment.

### **Oral Leukoplakia and Erythroplakia Pathology**

Histopathologic evaluation of persistent white or red oral mucosal lesions for hyperkeratosis, epithelial

dysplasia, carcinoma in situ, or invasive squamous cell carcinoma. The clinical appearance alone does not determine the diagnosis.

### **Oral Lichen Planus**

A long-lasting inflammatory condition causing lace-like white lines, red patches, or painful mouth sores, often on the cheeks or gums. It may fluctuate over time; persistent painful or changing areas need dental review to exclude other disease.

#### **Alternate Names:**

OLP; Mucosal Lichen Planus

### **Oral Medications for Type 2 Diabetes**

Oral medicines for type 2 diabetes lower glucose through effects on insulin, liver glucose production, kidney reabsorption, or intestinal absorption; choice depends on health and treatment goals.

### **Oral Moniliasis**

Oral moniliasis is oral candidiasis, a *Candida* yeast infection causing white plaques, soreness, burning, or altered taste.

### **Oral Mucosa**

The moist lining of the mouth, including the inner lips and cheeks, gums, tongue, and palate. It protects deeper tissues, provides sensation, stays lubricated, and can develop ulcers, infections, inflammation, or tumors.

### **Oral Mucous Cyst**

An oral mucous cyst is a mucus-filled lesion caused by salivary-gland duct injury or blockage, often appearing on the lower lip or floor of the mouth.

### **Oral Nutritional Supplement**

A calorie-, protein-, or micronutrient-containing product consumed by mouth to supplement, not automatically replace, usual food intake. The product and dose should match the patient's needs, swallowing ability, disease restrictions, and tolerance.

### **Oral Pathology**

The study and diagnosis of diseases affecting the

---

mouth, teeth, jaws, and related tissues. These include tooth decay, infections, cysts, inflammation, developmental changes, and tumors. Examination, imaging, laboratory tests, or a tissue sample may be needed to identify the cause.

### **Oral Potentially Malignant Disorders**

Conditions affecting the mouth lining that carry an increased risk of developing into oral cancer. Examples include leukoplakia, erythroplakia, and some long-lasting inflammatory or inherited disorders; appearance alone cannot determine the risk.

### **Oral Pyogenic Granuloma**

A rapidly growing, friable vascular overgrowth of the gums or other oral mucosa, often related to irritation, trauma, or pregnancy. It bleeds easily and may recur unless the lesion and its local cause are treated.

### **Oral Radiology**

The use and interpretation of X-rays and other scans of the mouth, teeth, jaws, face, and nearby structures. These images help find decay, infection, fractures, tumors, bone disease, and problems related to treatment planning.

### **Oral Rehydration Salts**

A balanced mixture of glucose and electrolytes dissolved in clean water to replace fluid and salts lost through diarrhea or vomiting.

### **Oral Rehydration Therapy**

Replacement of water and electrolytes by mouth using a balanced glucose-and-salt solution, especially to prevent or treat dehydration from diarrhoea or vomiting.

**Alternate Names:**

ORT

### **Oral Route**

The administration of a medicine through the mouth for swallowing and absorption through the gastrointestinal tract. Clear instructions should spell out “by mouth” to reduce medication-route errors.

**Alternate Names:**

PO; Per Os; By Mouth

### **Oral Sex Practices**

Oral stimulation of the genitals for sexual pleasure. Fellatio: oral stimulation of the penis. Cunnilingus: oral stimulation of the vulva and clitoris. Analingus (rimming): oral stimulation of the anus.

**Alternate Names:**

Oral-Genital Stimulation; Fellatio and Cunnilingus

### **Oral Spray**

A liquid medicine or preparation sprayed into the mouth or throat for local treatment or absorption through oral tissues.

### **Oral Squamous Cell Carcinoma**

The most common type of mouth cancer, arising from the cells lining the lips, tongue, gums, cheeks, or palate. A sore, lump, red or white patch, bleeding, pain, or numbness that persists needs examination and often a biopsy.

### **Oral Submucous Fibrosis**

A chronic, progressive scarring disorder of the mouth lining, strongly associated with chewing areca nut or betel quid. Fibrous bands can make the mouth stiff and reduce how widely it opens; burning, difficulty eating spicy food, or trouble swallowing may occur. It is a potentially malignant condition and needs dental or specialist assessment.

**Alternate Names:**

OSMF

### **Oral Surgery**

Operations on the mouth, teeth, jaws, or nearby facial tissues. Procedures include tooth removal, dental implants, tissue sampling, cyst removal, repair of injury, and correction of jaw problems.

### **Oral Thrush**

A fungal infection of the mouth, usually caused by *Candida*, producing creamy white patches, soreness, or altered taste.

**Alternate Names:**



---

Oral Candidiasis; Oropharyngeal Candidiasis

### Oral Toxicity

Harm to the mouth, throat, or digestive system caused by a swallowed poison, medicine, treatment, or excessive dose. It may cause burning, sores, nausea, vomiting, abdominal pain, or organ injury and may require urgent poison-control or medical advice.

### Oral Yeast Infection

Alternate name; see *Oral thrush*.

### Orange Peel Skin

Skin with small dimples resembling an orange peel because swelling stretches the skin around hair follicles. It may result from blocked lymph drainage, especially in breast disease, and a new breast change needs assessment.

### Orbicularis

A circular muscle surrounding an opening, such as orbicularis oculi around the eye or orbicularis oris around the mouth.

### Orbit

The bony socket that contains and protects the eye, its muscles, nerves, blood vessels, and supporting tissues. Problems in the orbit can cause pain, swelling, double vision, bulging, or impaired eye movement.

### Orbital

Relating to the orbit, the bony cavity containing the eye, or to structures within it. Orbital disease may affect the eye, muscles, nerves, or surrounding tissues.

### Orbital Apex Syndrome

A clinical syndrome resulting from compressive, neoplastic, or inflammatory lesions located at the orbital apex involving both the superior orbital fissure and the optic canal, characterized by combined palsies of cranial nerves III, IV, V1, and VI (complete ophthalmoplegia, ptosis, and forehead sensory loss) associated with optic nerve (CN II) atrophy

and vision loss.

#### Alternate Names:

Apex Orbitae Syndrome; Combined Superior Orbital Fissure and Optic Canal Syndrome

### Orbital Blowout Fracture

A break in one of the thin bones forming the eye socket, usually after blunt trauma. It can cause swelling, double vision, numbness, or restricted eye movement and may need urgent evaluation.

### Orbital Cavity

The bony space in the skull that contains and protects the eye, its muscles, nerves, blood vessels, and supporting tissues.

### Orbital Cellulitis

A serious infection in the tissues around and behind the eye. It can cause eyelid swelling, fever, pain with eye movement, double vision, or reduced vision and can spread to the brain.

### Orbital Contusion

A bruise of the tissues around the eye caused by blunt trauma, often producing swelling, bruising, and tenderness.

### Orbital Floor Fracture

A traumatic 'blowout' fracture of the thin bony orbital floor (maxillary bone) caused by sudden blunt impact to the globe and orbit, presenting with enophthalmos, periorbital ecchymosis, infraorbital nerve (V2) paresthesias, and vertical diplopia due to mechanical entrapment of the inferior rectus muscle.

#### Alternate Names:

Blowout Fracture of the Orbit; Orbital Floor Blowout

### Orbital Fracture

A break in one or more bones forming the eye socket, usually caused by facial trauma and potentially associated with double vision, eye-movement restriction, or injury to the eye.

---

## Orbital Fractures

Breaks in the bones surrounding the eye, which may cause swelling, double vision, numbness, restricted eye movement, or injury to the eye itself.

## Orbital Irradiation

Radiation treatment directed at tissues within the eye socket, sometimes used for severe thyroid eye disease.

## Orbital Metastasis

Cancer that has spread to the bony socket around the eye from another organ, commonly the breast, lung, or prostate. It may cause bulging of the eye, pain, double vision, restricted movement, or reduced sight.

## Orbital Myositis

Inflammation of one or more muscles that move the eye. It can cause pain, eyelid swelling, double vision, and restricted eye movement; treatment depends on the cause and may require urgent specialist assessment.

## Orbital Neoplasm

An abnormal growth in the eye socket that may arise from the eye, optic nerve, muscles, glands, bone, or other tissues. It can cause bulging, pain, double vision, or vision loss and may be benign or cancerous.

## Orbital Pseudotumor

Painful inflammation of tissues around the eye that is not caused by infection or a true tumor. It may cause eyelid swelling, bulging of the eye, double vision, and painful or limited eye movement.

### Alternate Names:

Idiopathic Orbital Inflammatory Disease; IOID

## Orbital Tumors

Benign or malignant growths arising within or around the eye socket. They may cause a visible mass, proptosis, double vision, pain, impaired eye movements, or vision loss; examination and orbital imaging guide biopsy and treatment.

## Orbital Varix

An enlarged, flexible vein in the eye socket. The eye may bulge more when a person bends, coughs, or strains and then return to normal when upright. It can rarely bleed, clot, or press on the optic nerve; imaging helps confirm the diagnosis.

### Alternate Names:

Orbital Venous Malformation; Intermittent Proptosis; Orbital Phlebectasia; Orbital Venous Anomaly

## Orbit CT Scan

A CT scan of the eye socket that uses X-rays and computer images to show the bones, eye muscles, nerves, blood vessels, and surrounding tissues. It can help assess injury, infection, bleeding, or a growth.

## Orchialgia

Pain in one or both testicles. Sudden severe pain, swelling, nausea, or a high-riding testicle may indicate twisting of the testicle. Longer-lasting pain can result from infection, injury, varicocele, nerve pain, or other causes.

## Orchidectomy

Surgical removal of one or both testes, also called orchiectomy.

## Orchiectomy

Surgical removal of one or both testicles. Radical inguinal orchiectomy is diagnostic and therapeutic for testicular cancer; bilateral orchiectomy is a hormonal therapy for prostate cancer.

## Orchitis

Orchitis is inflammation of one or both testes, often from infection such as mumps or bacteria, causing pain, swelling, and sometimes impaired fertility.

## Order Coleoptera

The insect order containing beetles, characterized by hardened forewings that protect the flying wings; some species affect health as pests or vectors.

---

## Order Spirochaetales

A biological order of spiral-shaped bacteria that includes groups causing syphilis, Lyme disease, relapsing fever, and leptospirosis. Identification of the species determines medical significance and treatment.

## Orexin

Orexin, or hypocretin, is a hypothalamic neuropeptide that promotes wakefulness, arousal, appetite, and energy regulation; loss of orexin signaling is strongly associated with narcolepsy type 1.

## Orexin Receptor Antagonists

Medications that block orexin (hypocretin) receptors to promote sleep. Suvorexant and lemborexant are approved for insomnia. They reduce wakefulness by blocking the arousal system. They have lower abuse potential than benzodiazepines and do not cause respiratory depression.

## Orf

A zoonotic skin infection caused by orf virus, a parapoxvirus of sheep and goats. Humans acquire it through contact with infected animals or contaminated fomites; a painful or tender nodular, ulcerating lesion usually develops on the hand or finger and often resolves without treatment.

## Organ

A structure composed of two or more different types of tissues that has a recognizable shape and performs specific functions.

## Organelle

A specialized part inside a cell that performs a particular job. Examples include the nucleus, which stores genetic instructions, and mitochondria, which release energy; organelles work together to keep the cell alive.

## Organ Failure

Severe loss of an organ's ability to perform essential functions, such as filtering blood, exchanging oxygen, or pumping blood.

## Organic

Relating to living matter or carbon-containing chemistry; in psychiatry, an older term for symptoms caused by a physical brain disorder.

## Organic Chemical

A carbon-containing chemical compound, usually also containing hydrogen and sometimes oxygen, nitrogen, sulfur, or other elements. Organic chemicals include medicines, fats, sugars, proteins, solvents, and many toxins.

## Organic Disease

An illness caused by a physical change or measurable disturbance in the body, such as infection, inflammation, tissue damage, or abnormal organ function. The term is older and is less useful when symptoms have not yet been explained.

## Organic Impurity

An organic impurity is an unintended carbon-containing chemical in a medicine, material, or specimen that may affect safety, quality, or measurement.

## Organization Name

The formal name of a group, institution, company, or agency; in healthcare records it should identify the responsible organization accurately.

## Organ of Corti

The specialized sensory receptor organ for auditory transduction situated upon the basilar membrane within the cochlear duct of the inner ear, containing inner hair cells (the primary auditory transducers) and outer hair cells (electromotile cochlear amplifiers) innervated by the cochlear nerve.

### Alternate Names:

Spiral Organ

## Organogenesis

The process during embryonic development in which groups of cells form and organize into organs. Disruption can lead to congenital abnormalities, while the process is also studied in tissue engineering and regenerative medicine.

---

## Organophosphate

A class of chemicals that can inhibit acetylcholinesterase. Some are pesticides or nerve agents and can cause cholinergic poisoning.

## Organophosphate Poisoning

A serious poisoning caused by pesticides or nerve agents that excessively stimulate nerves and muscles. It may cause sweating, drooling, vomiting, diarrhea, pinpoint pupils, muscle twitching, wheezing, weakness, seizures, or breathing failure.

## Organophosphate Toxicity

Acetylcholinesterase inhibition causing a cholinergic toxidrome: miosis, sweating, salivation, lacrimation, vomiting, diarrhea, bronchorrhea, bronchospasm, bradycardia, fasciculations, weakness, seizures, and respiratory failure. Immediate decontamination, airway support, atropine, and early pralidoxime are key treatments; delayed neuromuscular weakness can occur.

## Organosilicon Compounds

Organosilicon compounds contain silicon bonded to carbon and are used in materials, medicines, devices, and industry; biological effects depend on the specific compound.

## Organ System

A group of organs that work together to perform an important body function, such as breathing or digestion.

## Organ Transplant

Surgical transfer of an organ from a donor to a recipient to replace failing organ function.

## Organ Transplantation

Replacement of a failing organ with a healthy organ from a donor. Transplants can restore function but require careful matching, lifelong monitoring, and medicines to prevent rejection; infection, rejection, and medicine-related harm remain possible.

## Organ Transplants

The surgical transfer of an organ from a donor to a recipient whose organ function is severely impaired. Solid-organ transplantation may involve the kidney, liver, heart, lung, pancreas, or intestine and requires donor assessment, recipient selection, and ongoing follow-up.

## Orgasm

The peak of sexual pleasure, usually accompanied by involuntary muscle and body responses that vary among individuals.

## Orgasmic Dysfunction

Persistent or recurrent difficulty reaching orgasm, markedly reduced orgasmic intensity, or absent orgasm that causes clinically significant distress. Medical, psychological, medication-related, hormonal, and relationship factors may contribute.

## Orgasmic Dysfunction in Women

Alternate name; see *Orgasmic Dysfunction*.

## Orientation

Awareness of personal identity, place, time, and situation. Clinicians assess orientation as part of mental-status and neurologic examinations.

## Orifice

An opening leading into or out of a body structure, organ, or passage, allowing materials to enter or leave.

## Orlistat

A medicine used with a reduced-calorie diet to help with weight loss or weight control. It stops some dietary fat from being absorbed, so oily stools, gas, and urgent bowel movements can occur; it can also reduce absorption of vitamins and some medicines.

## Ornithine

A nonessential amino acid and intermediate in the urea cycle that helps convert toxic ammonia into urea.

---

### Ornithine Decarboxylase

An enzyme that converts ornithine to putrescine, an early step in polyamine synthesis involved in cell growth.

### Ornithine Transcarbamylase Deficiency

OTC deficiency is an inherited urea-cycle disorder causing ammonia accumulation, vomiting, confusion, seizures, or coma, especially during metabolic stress.

### Orogastric Tube

A flexible tube passed through the mouth into the stomach. It may deliver nutrition or medicines, remove stomach contents, or decompress the stomach when nasal placement is unsuitable.

### Orolabial Herpes Simplex

Alternate name; see *Oral Herpes*.

### Oropharyngeal

Oropharyngeal means relating to the mouth and the part of the throat behind it, including structures involved in breathing, swallowing, and speech.

### Oropharyngeal Airway

A rigid device placed through the mouth to keep the tongue from blocking the throat in an unconscious person. It must not be used in someone with a gag reflex and does not prevent vomit from entering the lungs.

### Oropharyngeal Cancer

Alternate name; see *Oropharyngeal Carcinoma*.

### Oropharyngeal Carcinoma

Cancer arising in the part of the throat behind the mouth, including the tonsils, back of the tongue, soft palate, or throat wall. Tobacco, alcohol, and some HPV infections increase risk; assessment usually includes examination, tissue sampling, and scans.

### Oropharyngeal Dysphagia

Difficulty moving food or liquid from the mouth through the throat. It may cause coughing, choking,

a wet voice, food coming through the nose, or food entering the airway and needs assessment when persistent.

### Oropharyngeal Neoplasm

An oropharyngeal neoplasm is an abnormal growth in the throat behind the mouth, including the tonsils and base of the tongue; it may be benign or malignant.

### Oropharynx

The part of the throat behind the mouth, including the tonsils and back of the tongue. It helps move food and liquid toward the esophagus and carries air toward the voice box.

### Oropharynx Lesion Biopsy

Sampling of an oropharyngeal lesion, usually through the mouth under local or procedural anesthesia, for histopathologic evaluation of infection, dysplasia, or cancer. Persistent or suspicious lesions require specialist assessment.

### Oropouche Fever

An emerging arboviral disease caused by the Oropouche virus, an orthobunyavirus transmitted to humans primarily by biting midges (*Culicoides paraensis*) and mosquitoes in South and Central America. It manifests as an acute febrile syndrome characterized by abrupt high fever, severe retro-orbital headache, chills, photophobia, arthralgias, and myalgias, with frequent symptom recrudescence within 1 to 2 weeks and occasional progression to aseptic meningitis.

#### Alternate Names:

Oropouche Virus Disease; Oropouche Fever Arbovirus Infection

### Orotic Aciduria

An unusually high amount of orotic acid in urine. It may result from a rare inherited disorder affecting blood-cell and DNA building blocks, causing anemia and poor growth, or from a urea-cycle disorder with high ammonia.

---

## Orphan Disease

A rare disease affecting relatively few people; definitions and eligibility for orphan-drug programs vary by jurisdiction.

## Orphenadrine

A centrally acting medicine with anticholinergic properties used in some countries for painful muscle spasm; it can cause drowsiness, dry mouth, and confusion.

## Orthodontic

Relating to the diagnosis, prevention, or treatment of problems with tooth and jaw position. Orthodontic treatment may use braces, aligners, retainers, or other appliances.

## Orthodontic Appliance

A device used to move teeth, guide jaw growth, maintain alignment, or correct a bite problem.

## Orthodontic Bracket

A bonded or banded attachment that transmits forces from an orthodontic archwire to a tooth. Bracket position and wire sequence guide controlled tooth movement.

## Orthodontic Retainer

A removable or fixed appliance used after active orthodontic treatment to maintain tooth position while periodontal and alveolar tissues remodel. Retention is often needed long term because relapse can occur.

## Orthodontics

The dental specialty that corrects teeth or jaws that do not align properly. Braces, clear aligners, or other appliances can improve biting, chewing, appearance, and cleaning; retainers are usually needed afterward because teeth can shift back.

## Orthodontic Treatment

Treatment that uses braces, wires, aligners, or other appliances to move teeth and guide jaw development. It can improve bite, spacing, and cleaning

access; treatment requires assessment and long-term retention.

## Orthodontic Wires

Metal or other wires used with braces to apply controlled forces that move teeth or help maintain their position.

## Orthodontist

A dentist specially trained to diagnose and treat teeth or jaws that do not align properly. Orthodontists use braces, clear aligners, retainers, and sometimes surgery to improve biting, chewing, appearance, and oral health.

## Orthodromic AVRT

The most common form of atrioventricular reentrant tachycardia in Wolff-Parkinson-White syndrome, in which the electrical cardiac impulse conducts antegradely down the normal atrioventricular node and His-Purkinje system (yielding a narrow QRS complex) and retrogradely up the accessory pathway (bundle of Kent) back to the atria.

## Orthognathic Surgery

Corrective surgery to reposition the upper jaw, lower jaw, or both when bone alignment causes problems with biting, chewing, speaking, breathing, or facial development. It is often combined with orthodontic treatment.

## Orthohepadnavirus

A group of viruses that includes hepatitis B virus and related viruses infecting mammals. These viruses carry genetic material as DNA and may cause long-lasting liver infection, inflammation, scarring, or cancer.

## Orthokeratosis

Thickening of the outer skin layer with fully mature skin cells. It can be a normal protective response, such as on the soles, or part of conditions causing thick, scaly, or rough skin.



---

## **Orthomyxoviridae**

A family of viruses that includes influenza A, B, and C. They mainly infect the airways and can cause flu, outbreaks, and sometimes severe pneumonia or other complications.

## **Orthopaedics**

The British spelling of orthopedics, the medical and surgical specialty concerned with the musculoskeletal system.

## **Orthopaedic Surgery**

Surgery for disorders and injuries involving bones, joints, muscles, ligaments, tendons, and related tissues. Procedures may repair fractures, replace joints, correct deformity, stabilize the spine, or relieve pressure on nerves.

## **Orthopedic Biologics**

Biological materials used to help bones, tendons, or other tissues heal. They include a person's own bone, donated bone, processed tissue, and synthetic substitutes; benefits and risks vary by the injury and procedure.

## **Orthopedic Implant**

A device placed in or on the body to repair, support, replace, or stabilize a bone or joint. Examples include plates, screws, rods, joint replacements, and artificial ligaments. Possible complications include infection, loosening, breakage, wear, or bone damage.

## **Orthopedic Infections**

Infections involving bones, joints, muscles, or orthopedic implants. Examples include osteomyelitis, septic arthritis, prosthetic-joint infection, and some diabetic foot infections.

## **Orthopedic Rehabilitation**

Treatment that restores movement, strength, daily function, and independence after injury or surgery involving bones, joints, muscles, ligaments, or tendons. It may include exercises, activity training, pain management, and assistive equipment.

## **Orthopedics**

The branch of surgery concerned with the musculoskeletal system, including bones, joints, ligaments, tendons, muscles, and nerves.

## **Orthopedic Screws**

Metal or other screws used to hold broken bones or surgical implants in position while bone heals. Their shape and size are chosen for the bone and purpose. Complications can include loosening, breakage, infection, or injury to nearby structures.

## **Orthopedic Services**

Orthopedic services diagnose and treat problems of bones, joints, muscles, ligaments, tendons, and related movement systems.

## **Orthopedist**

A physician who diagnoses and treats problems of bones, joints, muscles, ligaments, tendons, and related tissues. Treatment may include rehabilitation, medicines, injections, or surgery.

### **Alternate Names:**

Orthopedic Surgeon; Orthopedist Surgeon

## **Orthopnea**

Shortness of breath that begins or worsens when lying flat and improves when sitting or standing. It often occurs with heart or lung disease, but other causes are possible; new or worsening symptoms need medical assessment.

## **Orthopneic Position**

Sitting upright, often leaning forward with the arms supported, to make breathing easier. A person who suddenly needs this position or cannot lie flat because of breathlessness should seek medical assessment.

## **Orthopnoea**

Shortness of breath that starts or worsens when lying flat and improves when sitting or standing. It often occurs with heart or lung disease; new or worsening difficulty lying down because of breathlessness needs medical assessment.

---

## **Orthopoxvirus**

A genus of poxviruses that includes variola, mpox, vaccinia, and cowpox viruses. Infection may cause fever and skin lesions; severity and transmission depend on the specific virus.

## **Orthoptera**

The insect order containing grasshoppers, crickets, katydids, and locusts; some species cause bites, allergies, or crop-related health effects.

## **Orthoptic**

Relating to assessment and treatment of binocular vision, eye alignment, and disorders such as strabismus.

## **Orthoptist**

An eye-care professional who assesses and treats problems with eye movement, eye alignment, and the use of both eyes together. Orthoptists commonly help manage strabismus, amblyopia, double vision, and some vision problems in children and adults.

## **Orthoreovirus**

A genus of nonenveloped, segmented double-stranded RNA viruses; human infection is often mild or asymptomatic but can occasionally cause disease.

## **Orthorexia**

A term used for an excessive or rigid preoccupation with eating foods considered healthy. It is not a universally recognized formal psychiatric diagnosis but may cause nutritional or social impairment.

## **Orthorexia Nervosa**

A term used for an unhealthy preoccupation with eating foods considered pure or healthy, leading to severe restriction, distress, or impaired daily life. It is not yet a formal standalone diagnosis in major classification systems, and professional assessment is important.

## **Orthoscopic**

Orthoscopic means providing a straight or undis-

torted view, as in an optical instrument or examination.

## **Orthosis**

A brace, splint, shoe insert, or similar device worn outside the body to support, align, protect, or improve movement of a body part. Unlike a prosthesis, it does not replace a missing body part.

## **Orthostatic Hypotension**

A fall in blood pressure after standing, often causing dizziness, blurred vision, weakness, or fainting. It may result from dehydration, medicines, autonomic nerve problems, or prolonged bed rest. Repeated symptoms or falls need medical assessment.

## **Orthostatic Hypotension and Autonomic Failure**

Disorders in which autonomic nerves or their central pathways cannot maintain blood pressure and other automatic body functions. They may cause orthostatic hypotension, faintness, urinary or bowel dysfunction, sweating abnormalities, sexual dysfunction, or impaired heart-rate responses.

## **Orthostatic Intolerance**

Dizziness, fast heartbeat, weakness, blurred vision, fatigue, or faintness that develops or worsens on standing because the body does not maintain blood flow to the brain adequately. Causes include dehydration, medicines, nerve disorders, and some chronic conditions.

## **Orthotic**

Orthotic means relating to a brace, splint, insert, or device that supports, aligns, protects, or improves function of a body part.

## **Orthotic Devices**

External supports that hold, align, protect, or assist a body part, such as a brace, splint, shoe insert, or spinal support. They can reduce pain and improve movement, but poor fit may cause pressure, numbness, skin injury, or restricted circulation.

---

## Orthotic Training

Instruction and practice for safely using a brace or other support. It includes putting it on and removing it, checking skin, cleaning it, walking with it, and recognizing problems needing review.

## Ortolani Maneuver

A hip examination in infants used to look for developmental hip dislocation. The clinician gently moves the bent leg outward and feels for a shift of the hip back into place. A clunk can support the finding, but the examination and ultrasound or X-ray are considered together.

### Alternate Names:

Ortolani's Sign; Hip Reduction Maneuver

## Ortolani Sign

A reassuring click or clunk felt when a doctor gently spreads an infant's hips outward, showing that a dislocated hip has slid back into its socket.

## Ortolani's Sign

A palpable or audible clunk produced when a dislocated but reducible infant hip is gently abducted and reduced; it is a screening sign of developmental dysplasia of the hip.

## Ortolani's Test

A hip examination maneuver in infants that detects a reducible dislocation by producing a palpable clunk as the femoral head enters the acetabulum.

## Os

A bone or bony structure; in Latin anatomical terminology it means bone, while in obstetrics it means an opening.

## Osborn Wave

A notch or extra wave on an electrocardiogram, classically seen when the body is dangerously cold. It can also occur in other conditions and must be interpreted with the person's temperature and other findings.

### Alternate Names:

J Wave; Hypothermic Wave

## Oscillation

Repeated movement or variation around a central position or value.

## Oscillator

A device or system that produces repeated movement or an electrical signal at regular intervals.

## Oscillograph

An instrument or computer display that records a changing signal over time. It can show electrical, sound, pressure, or movement waves and has been used in medical tests to study signals from the heart, brain, muscles, or breathing.

## Oscillometry

A method that measures pressure-related vibrations, commonly used by automated blood-pressure cuffs and some vascular devices. The result estimates blood pressure or blood-flow characteristics but may be inaccurate with movement or irregular rhythms.

## Oscillopsia

The disturbing illusion that the visual environment is moving or bouncing, often caused by vestibular or eye-movement dysfunction.

## Oscitation

Yawning, a reflex involving opening the mouth and deep inhalation that is associated with sleepiness, fatigue, or other physiologic states.

## Oseltamivir

Oseltamivir is an antiviral medicine used to treat or prevent influenza, with the greatest benefit when started early.

## Oseltamivir Phosphate

Oseltamivir phosphate is the salt form of oseltamivir, an antiviral medicine used for treatment or prevention of influenza.

## OSFED

A recognized eating disorder in which symptoms cause significant distress or interfere with health or

---

daily life but do not meet every criterion for another named eating disorder. It still requires assessment and treatment.

### Osgood-Schlatter Disease

Painful irritation of the growing area where the kneecap tendon attaches to the shinbone. It commonly affects active children and teenagers and causes tenderness or swelling below the knee.

### Osler Nodes

Tender red or purple lumps on the pads of the fingers or toes. They can occur with infective endocarditis and some other conditions; they are an important clue but do not prove the diagnosis, which requires a full medical assessment.

### Osler Triad

A historical clinical triad (Austrian syndrome) characterized by the simultaneous co-occurrence of pneumococcal pneumonia, pneumococcal endocarditis (classically involving the aortic valve), and pneumococcal meningitis, carrying high mortality in patients with asplenia or chronic alcohol dependence.

### Osler-Weber-Rendu Disease

An autosomal dominant vascular dysplasia (hereditary hemorrhagic telangiectasia) caused by mutations in the ENG or ACVRL1 genes, characterized clinically by recurrent epistaxis, widespread mucocutaneous telangiectasias, and visceral arteriovenous malformations (AVMs) in the lungs, brain, liver, and gastrointestinal tract.

#### Alternate Names:

Hereditary Hemorrhagic Telangiectasia; HHT

### Osmidrosis

Strong, unpleasant body odor caused when skin bacteria break down sweat and skin oils. It commonly affects the armpits or feet and may improve with washing, dry clothing, antiperspirants, treatment of skin infection, or assessment of an underlying condition.

### Osmium

A dense metallic element; some osmium compounds

are highly toxic and can damage the eyes and respiratory tract.

### Osmolal Gap

The difference between measured serum osmolality (via freezing point depression osmometry) and calculated serum osmolality ( $2 \times [\text{Na}^+] + [\text{Glucose}]/18 + [\text{BUN}]/2.8$ ), where a gap exceeding 10 mOsm/kg indicates the presence of unmeasured exogenous osmotically active substances (such as toxic alcohols: methanol, ethylene glycol, or isopropanol).

#### Alternate Names:

Osmolar Gap

### Osmolality

A measure of how many dissolved particles are present in a kilogram of fluid. Blood osmolality depends mainly on sodium, glucose, and urea and helps determine how water moves between the bloodstream and cells.

### Osmolality Blood Test

A blood osmolality test measures the concentration of dissolved particles in blood and helps assess hydration, electrolyte disorders, and certain poisonings.

### Osmolality Urine Test

A urine osmolality test measures how concentrated the urine is and helps evaluate the kidneys' ability to regulate water balance.

### Osmolar

Describing a liquid according to the number of dissolved particles in each liter. This property affects water movement between body compartments and helps clinicians assess blood chemistry, hydration, kidney function, and intravenous fluids.

### Osmolarity

The concentration of osmotically active particles per liter of solution.

### Osmole

A unit representing the amount of a substance that

---

contributes one mole of osmotically active particles.

### **Osmoreceptor**

A sensory cell or neural receptor that detects changes in osmotic concentration, especially in the hypothalamus.

### **Osmoregulation**

Control of water and dissolved-solute balance in the body or an organism, maintaining appropriate cell volume and internal concentration.

### **Osmosis**

Movement of water through a membrane toward the side with more dissolved particles. This process helps determine how water is distributed between blood, cells, and surrounding tissues.

### **Osmotic Demyelination Syndrome**

Serious brain injury that can occur when long-standing low blood sodium is corrected too quickly. Symptoms may appear days later and include confusion, difficulty speaking or swallowing, weakness, movement problems, or reduced consciousness.

### **Osmotic Diarrhea**

Diarrhea caused by poorly absorbed solutes retaining water in the intestinal lumen. It typically improves with fasting and may result from carbohydrate malabsorption, laxatives, or other osmotically active substances.

### **Osmotic Diuresis**

Increased urine production caused by poorly reabsorbed solutes in the kidney tubules, such as glucose during marked hyperglycemia.

### **Osmotic Fragility Test**

A test that measures the susceptibility of red blood cells to hemolysis when placed in progressively more dilute saline solutions. Increased fragility supports hereditary spherocytosis, although the result may also be affected by other hemolytic disorders and recent transfusion.

### **Osmotic Pressure**

The force that draws water across a membrane toward the side containing more dissolved particles. It helps control how water moves between blood, cells, and surrounding tissues.

### **Os Odontoideum**

Os odontoideum is a separate ossicle near the second cervical vertebra's odontoid process that can cause atlantoaxial instability or spinal-cord risk.

### **Osseointegration**

The firm attachment of living bone to the surface of an implant, such as a dental implant or artificial joint. Good stability, healthy bone, healing, and infection control support this process.

### **Osseous**

Relating to or composed of bone. For example, an osseous lesion is a bone lesion.

### **Osseous Metaplasia**

Osseous metaplasia is formation of mature bone in tissue where bone is not normally present, sometimes after injury, inflammation, or within a lesion.

### **Ossicle**

A small bone, especially one of the three tiny middle-ear bones. The malleus, incus, and stapes transmit eardrum vibrations to the inner ear, allowing sound to be heard.

### **Ossicles**

Three smallest body bones forming the ossicular chain in the middle ear: malleus (hammer, attached to tympanic membrane), incus (anvil), stapes (stirrup, footplate in oval window).

### **Ossicular Discontinuity**

Partial or complete separation of the middle-ear ossicles, usually caused by trauma, chronic ear disease, or surgery. It produces conductive hearing loss and may be evaluated with otoscopy, audiometry, and imaging.

**Alternate Names:**

---

Ossicular Chain Discontinuity; Ossicular Disruption

### **Ossicular Fusion**

Abnormal joining or fixation of the middle-ear ossicles (malleus, incus, or stapes), which can restrict sound transmission and cause conductive hearing loss. Evaluation and treatment depend on the underlying congenital, inflammatory, traumatic, or postoperative cause.

### **Ossicular Prosthesis**

An artificial replacement or support for one or more middle-ear ossicles, used during ear surgery to restore sound transmission in selected conductive hearing loss.

### **Ossification**

The formation of bone. It occurs during growth and fracture healing and can also happen abnormally in soft tissues after severe injury, burns, or nervous-system damage.

### **Ossifying Fibroma**

An ossifying fibroma is a usually benign fibro-osseous tumor that replaces normal bone with fibrous tissue and mineralized material, often in the jaw.

### **Osteitis**

Inflammation of bone caused by infection, injury, inflammatory disease, or another process. It may cause localized pain, swelling, warmth, reduced movement, or fever and sometimes requires imaging, antibiotics, or surgery.

### **Osteitis Deformans**

Osteitis deformans is Paget disease of bone, in which abnormal remodeling produces enlarged, weakened, or deformed bones.

### **Osteitis Fibrosa**

Weakening of bone caused by prolonged high levels of parathyroid hormone. In severe cases, bone may be replaced by fibrous tissue and develop painful weak spots.

### **Osteitis Fibrosa Cystica**

A rare, severe bone complication of long-standing hyperparathyroidism. Excess parathyroid hormone causes bone to break down, become weak, and sometimes form cyst-like areas called brown tumors. It may cause bone pain, deformity, or fractures.

#### **Alternate Names:**

Von Recklinghausen Disease Of Bone

### **Osteitis Pubis**

A usually noninfectious inflammation or stress-related disorder of the pubic symphysis and nearby tissues, causing pain in the groin or lower abdomen. It may follow athletic overuse, pelvic surgery, pregnancy, or childbirth.

### **Osteoarthritis**

A joint disease in which cartilage and other tissues gradually break down or change. It can cause pain, stiffness, swelling, bone changes, and restricted movement. It usually develops over time but may occur earlier after joint injury, abnormal joint structure, or previous joint disease.

#### **Alternate Names:**

Degenerative Joint Disease; DJD; Hypertrophic Arthritis

### **Osteoarthropathy**

Osteoarthropathy is disease affecting bones and joints, often describing degenerative, hypertrophic, metabolic, or systemic-associated changes.

### **Osteoarthrosis**

Another name for osteoarthritis, a joint disorder in which cartilage and nearby bone gradually change, causing pain, stiffness, and reduced movement.

### **Osteoblast**

A cell that builds new bone. It makes the soft framework of bone and helps add calcium and phosphate to harden it. Some osteoblasts become osteocytes, which maintain mature bone.

### **Osteoblastoma**

A rare benign bone-forming tumor, usually in the



---

spine or long bones, that can cause persistent pain and sometimes neurologic symptoms.

### **Osteoblasts**

Cells that build new bone. They produce the material that forms bone and help control its hardening with calcium and phosphate. Some osteoblasts become osteocytes, which help maintain mature bone.

### **Osteocalcin**

A protein made mainly by osteoblasts that participates in bone mineralization and serves as a marker of bone formation.

### **Osteocalcin Assay**

A blood test measuring osteocalcin, a protein made by bone-forming cells. It can provide information about bone formation and turnover, but results are affected by age, kidney function, medicines, and the laboratory method.

### **Osteochondritis**

Inflammation or injury involving bone and its overlying cartilage, often near a joint. The term includes conditions such as osteochondritis dissecans, in which a fragment may loosen.

### **Osteochondritis Dissecans**

An area of damaged bone and cartilage beneath a joint surface that may loosen or separate. It most often affects the knee in adolescents and young adults and can cause activity-related pain, swelling, catching, or locking. Treatment depends on lesion stability, age, and symptoms.

### **Osteochondroma**

A usually noncancerous cartilage-covered growth projecting from the surface of a bone, often near a growth plate. It may be single or occur as multiple inherited growths and can cause pain, deformity, or pressure on nearby nerves or vessels.

### **Osteochondromatosis**

Osteochondromatosis is formation of multiple cartilage-capped bony outgrowths, usually around

growth plates, that can cause deformity, pain, or nerve and vessel compression.

### **Osteochondrosis**

A general name for several disorders affecting developing bone and nearby cartilage, often near a growth area or joint. The specific condition depends on the bone and age involved; it may cause pain, swelling, stiffness, or limited movement.

### **Osteoclasis**

Deliberate surgical breaking or manipulation of a bone to correct a deformity or restore alignment. It requires stabilization while the bone heals and may cause bleeding, infection, or nerve injury.

### **Osteoclast**

A specialized bone cell that breaks down old bone during normal renewal and repair. Osteoclast activity helps regulate bone strength and blood calcium; excessive activity can contribute to osteoporosis and other diseases causing bone loss.

### **Osteoclastoma**

See *Giant Cell Tumor Of Bone*.

### **Osteoclasts**

Cells that break down old bone. This normal process helps bones grow, repair themselves, and release calcium into the blood. Too much osteoclast activity causes bone loss, as in osteoporosis, Paget disease, and some cancers that spread to bone. Medicines such as bisphosphonates and denosumab can slow osteoclast activity and reduce bone weakening.

### **Osteocyte**

A mature bone cell embedded in the hard bone it helped form. Osteocytes sense strain and chemical changes and send signals that help control bone strength, repair, and removal of old bone.

### **Osteocytes**

Mature bone cells trapped inside the hard bone they helped form. They sense strain and changes in the bone and send signals that help control bone repair,

---

strengthening, and breakdown.

### **Osteodystrophy**

Abnormal bone growth, strength, or renewal caused by a body-wide or metabolic disorder. It may result from kidney disease, mineral imbalance, hormone problems, or inherited conditions and can cause pain or fractures.

### **Osteogenesis**

The formation and development of bone. It occurs in the growing skeleton and during fracture repair. Abnormal osteogenesis can contribute to birth disorders, poor healing, or fragile bones.

### **Osteogenesis Imperfecta**

A group of genetic disorders, also called brittle bone disease, in which bones break easily, sometimes after mild injury or for no clear reason. The disorders are often caused by gene changes that reduce or alter type I collagen, a protein that gives strength to bone and other connective tissues. Osteogenesis imperfecta is present from birth and can range from mild to severe; severe forms may cause fractures before birth. Other features may include short height, loose joints, blue or gray whites of the eyes, hearing loss, weak or discolored teeth, weak muscles, or a curved spine.

### **Osteogenic Sarcoma**

A cancerous bone tumor whose cells make immature bone tissue. It often develops near the ends of growing long bones and may cause persistent pain, swelling, reduced movement, or a fracture.

### **Osteoid**

The newly formed, unmineralized organic framework of bone, produced mainly by osteoblasts. Minerals are later deposited in it to make mature hard bone.

### **Osteoid Osteoma**

A small noncancerous bone tumor, usually in an adolescent or young adult. It often causes persistent pain that is worse at night and may improve with anti-inflammatory medicine; scans help locate it and guide treatment.

### **Osteology**

The study of bones, including their structure, development, function, injuries, and diseases. It supports the assessment and treatment of fractures, bone deformities, weak bones, infections, and bone tumors.

### **Osteolysis**

Osteolysis is progressive destruction or resorption of bone caused by tumors, inflammation, prosthesis reactions, infection, metabolic disease, or other processes.

### **Osteolytic**

Causing or characterized by breakdown and loss of bone tissue. Osteolytic lesions can weaken bones and occur with cancers, infections, inflammation, metabolic disorders, or excessive activity of bone-removing cells.

### **Osteoma**

A benign, slow-growing tumor made of bone. It often develops in the skull or facial bones and may cause no symptoms unless it blocks a passage or presses on nearby structures.

### **Osteoma Cutis**

Osteoma cutis is formation of bone within the skin or subcutaneous tissue, occurring as a primary disorder or secondary to acne, trauma, or another condition.

### **Osteomalacia**

Osteomalacia is inadequate mineralization of adult bone, commonly from vitamin-D deficiency or phosphate disorders, causing bone pain, weakness, and fractures.

### **Osteomyelitis**

Infection and inflammation of bone, usually caused by bacteria and sometimes fungi or other organisms. It can cause pain, fever, swelling, reduced movement, bone damage, or a persistent draining wound.

### **Osteomyelitis In Children**

Childhood osteomyelitis is infection of bone, often caused by bacteria entering through blood or

---

trauma, producing pain, fever, swelling, and impaired movement.

### **Osteon**

A tube-shaped structural unit of hard, compact bone arranged around a small channel containing blood vessels and nerves. Osteons help give bone strength while allowing it to receive nutrients.

### **Osteonecrosis**

Death of bone tissue when its blood supply is seriously reduced or interrupted. It may cause pain and, near a joint, collapse of the bone and joint damage. Causes include injury, some blood disorders, long-term steroid use, heavy alcohol use, and sometimes no known cause.

#### **Alternate Names:**

Avascular Necrosis; Aseptic Necrosis; Ischemic Bone Necrosis; AVN

### **Osteonecrosis Of The Jaw**

Death of jawbone tissue, often associated with antiresorptive or antiangiogenic medications, radiation, infection, or trauma, and characterized by exposed bone or a nonhealing oral wound.

### **Osteonectin**

An extracellular-matrix protein involved in collagen organization, mineral binding, and tissue remodeling; it is also called SPARC.

### **Osteons**

Cylindrical structural units of compact bone arranged around central canals.

#### **Alternate Names:**

Haversian Systems

### **Osteopath**

A physician trained in medicine and surgery who earns a Doctor of Osteopathic Medicine degree.

#### **Alternate Names:**

Doctor of Osteopathic Medicine; DO

### **Osteopathy**

A term used differently in different countries. It

may refer to physician-led osteopathic medicine or to manual treatment focused on the muscles and joints; the practitioner's training and legal scope should be checked locally.

### **Osteopenia**

Bone density that is below the usual range but does not meet the bone-density threshold for osteoporosis. In some adults, a DXA scan describes osteopenia with a T-score between -1.0 and -2.5; this range is not used to classify everyone. Fracture risk also depends on age, previous fractures, falls, medicines, and other health factors.

### **Osteopetrosis**

A rare inherited disorder in which old bone is not removed normally. Bones become unusually dense but brittle and may fracture; crowding of the bone marrow and nerves can cause anemia, infections, vision loss, or other problems.

### **Osteophyte**

A bony outgrowth forming at the edge of a joint, commonly with osteoarthritis. It may cause stiffness, pain, reduced movement, or pressure on nearby nerves, although some osteophytes cause no symptoms.

### **Osteoplasty**

Surgery to repair, reshape, or reconstruct bone to restore structure, movement, or function. The technique depends on the bone problem and may be combined with fixation, grafting, or joint surgery.

### **Osteopoikilosis**

Osteopoikilosis is an inherited benign bone condition with multiple small dense spots on imaging, usually asymptomatic and discovered incidentally.

### **Osteopontin**

A protein involved in bone remodeling, immune responses, inflammation, and attachment between cells and tissues. Its level may change in several diseases, but a measurement is not by itself a specific diagnosis.

---

## **Osteoporosis**

A bone disease in which bone is broken down faster than it is replaced, or its structure and strength change. The bones become less dense and more likely to break, sometimes after a minor fall or other low-impact force. Osteoporosis often causes no symptoms until a fracture occurs. Fractures commonly affect the hip, spine, and wrist; weakened bones in the spine may collapse and cause back pain or loss of height. Osteoporosis can affect anyone, but it is more common in females, especially after menopause, when lower estrogen levels can speed bone loss. It can also affect males. Risk is influenced by age, family history, hormones, inadequate calcium or vitamin D, some medicines, smoking, alcohol, and other health conditions.

## **Osteoporosis Fracture Risk Assessment**

An evaluation of future fracture risk using bone density, age, previous fractures, falls, medicines, family history, and other health factors. The result helps determine whether lifestyle measures, monitoring, or bone-strengthening treatment is appropriate.

## **Osteoporotic Fracture**

A fracture resulting from weakened, porous bone after a low-energy event that would not usually break healthy bone.

## **Osteoradionecrosis**

Death of bone within an area treated with radiation, most often the jaw after head-and-neck cancer treatment. The bone may become exposed or fail to heal after injury or dental work and can become infected.

## **Osteosarcoma**

An aggressive cancer that begins in bone-forming cells and produces abnormal immature bone. It often develops near the ends of long bones in children and young adults and may cause persistent pain, swelling, or a fracture.

## **Osteosclerosis**

Abnormally increased bone density seen on imaging or examination. It may result from inherited disease,

altered bone renewal, chronic stress, inflammation, medicines, or cancer; dense bone can still be weak or painful.

## **Osteosynthesis**

Surgical repair and stabilization of a broken bone using implants such as plates, screws, rods, nails, or wires. The implants hold the pieces in alignment while the bone heals.

## **Osteotome**

A sharp surgical instrument used to cut or shape bone.

## **Osteotomy**

A surgical procedure involving cutting and repositioning of bone to correct deformity, realign joint surfaces, alter weight-bearing forces, or change limb length.

## **Osteotomy Of The Knee**

Surgery that cuts and reshapes a knee-region bone to correct alignment and redistribute weight across the joint.

## **Ostium**

A natural opening into or out of an organ, cavity, blood vessel, or duct. It allows air, fluid, or other contents to pass between connected body spaces.

## **Ostomy**

A surgically created opening that connects part of the bowel or urinary tract to the skin so stool or urine can leave the body. A colostomy or ileostomy passes stool; a urostomy passes urine. A pouch usually collects the output and protects the surrounding skin.

## **Ostomy Appliance**

A pouching system that collects stool or urine from a surgically created opening in the abdomen. Colostomy and ileostomy appliances collect stool; urostomy appliances collect urine. The skin around the opening needs regular protection and monitoring.

---

### **Ostraceous Psoriasis**

A rare severe form of psoriasis in which plaques become very thick, hard, and layered, resembling an oyster shell. It can be difficult to treat and may require specialist care and medicines affecting the whole immune system.

### **Otalgia**

The medical term for earache or ear pain. It is divided into two main types: primary and referred. Primary ear pain originates directly inside the ear, most often caused by middle ear infections, swimmer's ear (infection of the outer ear canal), buildup of earwax, pressure changes, or eardrum injury. Referred ear pain comes from nearby body parts that share the same sensory nerves with the ear, including tooth decay, jaw joint problems (temporomandibular joint issues), throat infections, or neck strain. Doctors find the cause by inspecting the ear canal and eardrum with a lighted scope (otoscope) and checking the teeth, mouth, and jaw when the ear looks healthy.

#### **Alternate Names:**

Earache; Ear Pain

### **Othello Syndrome**

A delusional belief that a partner is unfaithful despite insufficient evidence; it may occur with psychiatric, neurologic, or substance-related disorders.

### **Other Pain Conditions**

Pain conditions that do not fit neatly into a single category, including nociplastic pain, fibromyalgia, chronic tension-type headache, temporomandibular pain, chronic pelvic pain, and complex regional pain syndrome. Diagnosis and treatment depend on the specific condition.

### **Other Race**

Other race is a demographic category for a person who does not select or fit the available listed race categories; classification varies by system and should not be treated as a biological diagnosis.

### **Otic Barotrauma**

Injury to the ear caused by failure to equalize pres-

sure during flying, diving, or other pressure changes.

### **Otic Capsule**

The dense part of the skull bone that surrounds and protects the inner ear, including the hearing and balance organs. Disease or abnormal remodeling in this bone can affect hearing or balance.

### **Otic Ointment**

A semisolid medicine formulated for application to the outer ear or ear canal; it should be used only as directed, especially when the eardrum may be perforated.

### **Otitis**

Inflammation of the ear, which may be infectious or noninfectious. Otitis externa affects the ear canal, otitis media affects the middle ear, and inner-ear inflammation may cause vertigo or hearing loss. Symptoms can include pain, fever, blockage, discharge, or reduced hearing and vary with the location and cause.

### **Otitis Externa**

Inflammation or infection of the outer ear canal, commonly causing ear pain, itching, tenderness, swelling, or discharge. Keeping water and objects out of the ear can help while it heals.

#### **Alternate Names:**

Swimmer's Ear; Tropical Ear Swimmer's Ear; External Otitis

### **Otitis Media**

Inflammation or infection in the middle ear, often after a cold or other upper-airway infection. It may cause ear pain, fever, reduced hearing, irritability, or fluid behind the eardrum.

### **Otitis Media With Effusion**

Fluid behind the eardrum without the severe pain, fever, or pus of an acute ear infection. It may cause blocked hearing or speech and learning problems in children and often clears, but persistent fluid needs hearing assessment.

---

## Otoacoustic Emissions

Very soft sounds naturally produced by a healthy inner ear that can be measured with a small probe placed in the ear canal. The test is quick and painless and is commonly used to screen newborn hearing and assess inner-ear function.

### Alternate Names:

OAE; Cochlear Emissions

## Otocephaly

A severe congenital craniofacial malformation characterized by absent or markedly underdeveloped mandible, ventral displacement or fusion of the ears, and associated facial and airway abnormalities. It is usually lethal because of profound impairment of breathing and feeding.

## Otolaryngologist

An otolaryngologist is a physician specializing in disorders of the ear, nose, throat, head, neck, voice, swallowing, and related surgery.

## Otolaryngology

The medical specialty treating disorders of the ears, nose, throat, head, and neck. Specialists manage hearing, balance, breathing, swallowing, voice, sinus, facial, and related surgical problems.

## Otolith

A tiny calcium-containing crystal in the inner ear that helps sense gravity and head movement. If it moves into the wrong balance canal, it can cause brief spinning attacks known as positional vertigo.

## Otolithiasis

The displacement of microscopic calcium carbonate crystals (otoconia) from the macula of the utricle into one or more fluid-filled semicircular canals of the inner ear (most frequently the posterior semicircular canal). Free-floating otolith debris deflects the cupula during rapid head position changes, causing transient, paroxysmal, rotational vertigo and torsional nystagmus characteristic of benign paroxysmal positional vertigo (BPPV).

### Alternate Names:

Canalithiasis; Cupulolithiasis; Otoconial Displace-

ment

## Otoliths

Tiny calcium-carbonate crystals in inner-ear organs that sense gravity and straight-line movement. If they move into the wrong balance canal, they can cause brief spinning attacks called positional vertigo.

## Otology

The study and medical care of the ear and hearing.

## Otomycosis

A superficial fungal infection of the external auditory canal, most commonly caused by *Aspergillus niger*, *Aspergillus fumigatus*, or *Candida albicans*, presenting with intense aural pruritus, otalgia, conductive hearing loss, and white or black cotton-like fungal debris filling the external ear canal.

### Alternate Names:

Fungal Otitis Externa; Mycotic Otitis

## Otoplasty

Surgery to alter the shape, position, or proportion of the external ear, commonly to correct prominent or congenital ear differences.

## Otorhinolaryngology

The medical specialty concerned with diseases of the ear, nose, throat, head, and neck; also called otolaryngology.

### Alternate Names:

Otolaryngology; ENT; Head and Neck Surgery

## Otorrhea

Fluid, pus, blood, or other discharge flowing from the ear. Causes include infection, eardrum perforation, injury, or a growth; persistent, bloody, foul-smelling, or clear discharge needs medical assessment. Clear watery discharge after a head injury may be cerebrospinal fluid and requires emergency assessment.

## Otorrhoea

Alternate spelling; see *Otorrhea*.



---

## Otosalpingitis

Inflammation, mucosal edema, and catarrhal obstruction of the Eustachian tube connecting the middle ear to the nasopharynx, predisposing to negative middle ear pressure, tympanic membrane retraction, and serous otitis media.

### Alternate Names:

Eustachian Salpingitis; Tubotympanitis

## Otosclerosis

An abnormal hardening and fixation of middle-ear bones that can gradually reduce hearing by blocking sound transmission.

## Otoscope

A lighted instrument used to examine the ear canal and eardrum. It can help identify earwax, infection, a foreign object, fluid, swelling, or a perforated eardrum. Examination findings are interpreted with symptoms and, when needed, hearing tests.

## Otoscopy

Looking into the ear canal and at the eardrum with a lighted instrument called an otoscope. It can help find earwax, infection, a foreign object, a hole in the eardrum, fluid, or other causes of ear symptoms.

## Ototoxicity

Damage to the inner ear from a medicine, chemical, or other toxic exposure. It may cause hearing loss, ringing in the ears, dizziness, or imbalance and can be temporary or permanent.

### Alternate Names:

Ear Toxicity; Vestibulotoxicity; Cochleotoxicity

## Ottawa Ankle Rules

A highly sensitive clinical decision-making algorithm used in acute emergency settings to determine whether radiography is required following acute ankle or midfoot trauma based on localized bony tenderness (posterior edge of distal malleoli, navicular bone, or base of fifth metatarsal) and inability to bear weight.

## Ounce

An ounce is a unit of mass or fluid volume whose exact conversion depends on the measurement system; medication instructions should use unambiguous units.

## Outcome

The result or final effect of a disease, treatment, exposure, or healthcare intervention. Outcomes may include recovery, symptom control, complications, disability, quality of life, or death and can be measured in different ways.

## Outcome Measurement

The use of defined measures to assess a patient's symptoms, function, quality of life, or other results before and after care.

## Outer Ear

The visible auricle and the external auditory canal, which collect and conduct sound toward the eardrum.

## Outer Ear Infection

Alternate name; see *Swimmer's Ear*.

## Outpatient

A person who receives medical evaluation or treatment without being admitted overnight to a hospital or other inpatient facility.

## Outpatient Rehabilitation

Rehabilitation provided without hospital admission to improve function after illness, injury, surgery, or disability. The setting, frequency, and duration are individualized.

## Output

The amount of a substance produced or eliminated by the body or by an organ over a specified time, such as urine output or cardiac output.

## Ova

Plural of ovum; eggs or egg cells, and in parasitology, eggs of worms or other parasites.

---

## Ovalocyte Count

An ovalocyte count estimates the number or proportion of oval-shaped red blood cells on a blood smear and may help evaluate anemia.

## Ovarian

Relating to an ovary. The ovaries release eggs and produce hormones. Ovarian problems include cysts, inflammation, reduced function, twisting that cuts off blood supply, and tumors; sudden severe pelvic pain may indicate twisting and needs emergency care.

## Ovarian (Adnexal) Torsion

Twisting of an ovary, often together with its fallopian tube, around its blood supply. Sudden unilateral pelvic pain, nausea, or vomiting may occur; it is a surgical emergency because prolonged torsion can cause ovarian necrosis.

## Ovarian Cancer

Cancer that begins in an ovary. Most are epithelial cancers; other types can begin in egg-producing or hormone-producing cells. Fallopian-tube and primary peritoneal cancers are related and are often grouped with ovarian cancer because they can behave and be treated similarly, but they begin in different tissues. Ovarian cancer may cause persistent bloating, pelvic or abdominal pain, or early fullness, though early disease may have few symptoms.

### Alternate Names:

Malignant Ovarian Neoplasm; Ovarian Malignancy

## Ovarian Carcinoid

Alternate form; see *Ovarian Neuroendocrine Tumor*.

## Ovarian Cyst

Alternate singular form; see *Ovarian Cysts*.

## Ovarian Cyst Rupture

Breakage of an ovarian cyst that may cause sudden pelvic pain and internal bleeding or, if the cyst is infected, fever and inflammation. Assessment depends on vital signs, pregnancy testing, ultrasound, blood

loss, and the severity and persistence of symptoms.

## Ovarian Cysts

Fluid-filled or partly solid sacs in or on an ovary. Most are harmless, cause no symptoms, and go away without treatment. Some cause pelvic pain, pressure, bloating, or menstrual changes. A cyst may rarely burst or twist the ovary, causing sudden severe pain and requiring urgent assessment. Ultrasound is commonly used for evaluation.

### Alternate Names:

Ovarian Fluid-Filled Sacs

## Ovarian Dysgerminomas

Malignant ovarian germ-cell tumors that occur most often in adolescents and young adults. They may present as a rapidly enlarging pelvic mass or pain; treatment commonly involves fertility-sparing surgery with staging and, when indicated, chemotherapy.

## Ovarian Failure

Reduced ovarian activity causing irregular or absent periods, difficulty becoming pregnant, and symptoms of low estrogen such as hot flashes or vaginal dryness. When it occurs before age 40, it is usually called primary ovarian insufficiency and needs evaluation.

## Ovarian Fibromata

Usually noncancerous tumors made of fibrous tissue that develop in an ovary. They may cause no symptoms or produce pelvic pressure, pain, menstrual changes, or fluid in the abdomen when large.

## Ovarian Follicle

A small fluid-filled structure in an ovary that contains an immature egg and supporting cells. During each menstrual cycle, one follicle usually matures and releases an egg during ovulation.

## Ovarian Gonadoblastoma

A rare tumor containing cells that normally develop into eggs and cells that support sex-hormone production. It often occurs in an atypically developed gonad, may contain Y-chromosome material, and

---

can produce hormones.

### Ovarian Hemorrhage

Bleeding into or around an ovary, often from a ruptured cyst or blood vessel. It may cause sudden pelvic pain, weakness, dizziness, or internal blood loss and can require urgent assessment or surgery.

### Ovarian Hyperstimulation Syndrome

A complication of fertility treatment in which the ovaries enlarge and fluid moves into the abdomen or other tissues. It may cause bloating, pain, vomiting, reduced urine, breathlessness, blood clots, or kidney problems and can be severe.

**Alternate Names:**

OHSS

### Ovarian Infection

Infection involving an ovary, often as part of pelvic inflammatory disease or an abscess, causing pelvic pain, fever, and abnormal discharge.

### Ovarian Insufficiency

Reduced ovarian hormone production or follicular function, which may cause irregular or absent menstruation, infertility, hot flashes, vaginal symptoms, and reduced bone density. The cause, age at onset, pregnancy goals, and associated autoimmune or genetic findings guide evaluation.

### Ovarian Metastasis

Cancer that has spread to an ovary from another organ, commonly the digestive tract, breast, or reproductive organs. It may resemble a primary ovarian cancer, so tissue testing is important because treatment depends on the original site.

### Ovarian Neuroendocrine Tumor

A rare ovarian tumor arising from cells with neuroendocrine features. It may be a primary ovarian tumor or part of a tumor arising elsewhere, and its behavior ranges from well-differentiated to aggressive disease.

### Ovarian Overproduction Of Androgens

Ovarian androgen overproduction is excessive production of male-pattern hormones by the ovaries, which may cause acne, excess facial hair, irregular periods, or infertility.

### Ovarian Pregnancy

A rare ectopic pregnancy that implants in an ovary rather than inside the uterus or fallopian tube. It may cause one-sided pelvic pain or internal bleeding and usually needs urgent ultrasound assessment and treatment.

### Ovarian Reserve Testing

Clinical and laboratory assessment of a woman's remaining primordial follicle pool and reproductive potential, measured primarily using early follicular phase serum anti-Müllerian hormone (AMH) concentration, day-3 follicle-stimulating hormone (FSH) with estradiol, and transvaginal ultrasound antral follicle count (AFC).

### Ovarian Rupture

Ovarian rupture is tearing of an ovary or ovarian cyst, which may cause sudden pelvic pain and internal bleeding and sometimes requires urgent treatment.

### Ovarian Teratoma

A tumor of an ovary made from reproductive cells that can form different kinds of tissue, such as skin, hair, or bone. Mature teratomas are usually non-cancerous; immature teratomas can be cancerous.

### Ovarian Thecoma

Alternate form; see *Thecoma*.

### Ovarian Torsion

Twisting of an ovary, often together with its fallopian tube, that can cut off blood flow. It causes sudden one-sided pelvic pain, often with nausea or vomiting, and is a gynecologic emergency that is usually treated with surgery.

---

## Ovaries

The paired female gonads that produce oocytes and secrete sex hormones, mainly estrogen and progesterone. They are located on either side of the uterus and participate in the menstrual cycle and reproduction.

### Alternate Names:

Female Gonads; Oophoron

## Ovariectomy

Older term; see *Oophorectomy*.

## Ovary

A reproductive organ that releases eggs and produces hormones such as estrogen and progesterone. The ovaries help control menstrual cycles, ovulation, pregnancy preparation, and many body changes during reproductive life.

## Ovary Disease

Any disorder affecting an ovary, including cysts, infection, inflammation, tumors, hormonal dysfunction, torsion, or bleeding.

## Ovary Neoplasm

An abnormal growth in an ovary that may be non-cancerous, have uncertain behavior, or be cancerous. Symptoms can include bloating, pelvic pain, pressure, or abnormal bleeding, but some growths are found incidentally on imaging.

## Overactive Bladder

A pattern of a sudden, difficult-to-delay need to urinate, usually with frequent urination and waking at night, with or without leakage. Causes include bladder irritation, nerve problems, obstruction, medicines, and sometimes no identifiable cause.

### Alternate Names:

OAB; Urgency Frequency Syndrome

## Overactive Bladder Syndrome

A symptom complex characterized by urinary urgency, with or without urgency urinary incontinence, usually accompanied by daytime frequency and nocturia in the complete absence of a proven urinary

tract infection or other obvious local pathological condition.

### Alternate Names:

OAB

## Overall Survival

The length of time from diagnosis, treatment, or study entry until death from any cause. Doctors and researchers use it to describe how long people live and to compare treatments. It differs from time without worsening disease because it includes survival regardless of whether the illness has progressed.

## Overbite

Vertical overlap of the upper front teeth over the lower front teeth; excessive overbite can damage gums, teeth, or jaw function.

## Overdenture

A removable denture supported by dental implants or remaining natural teeth. This support can improve stability, chewing, speech, and comfort, while attachments may help preserve jawbone and reduce denture movement.

## Overdose

An overdose is exposure to a medicine, drug, or substance in an amount capable of causing toxicity, with effects determined by dose, formulation, route, timing, and patient factors.

## Overflow Incontinence

Involuntary leakage of urine occurring when the intravesical pressure exceeds the outlet resistance in a chronically overdistended, poorly contracting bladder, caused by mechanical bladder outlet obstruction (e.g., benign prostatic hyperplasia) or an acontractile detrusor muscle in diabetic or neurogenic bladder.

### Alternate Names:

Paradoxical Incontinence; Overdistension Incontinence

---

### Overflow Urinary Incontinence

Urine leakage caused by a bladder that remains too full because it does not empty properly. Weak flow, difficulty starting, frequent small amounts, dribbling, or constant leakage may occur; blockage or nerve disease can be responsible.

### Overgrowth Syndrome

A group of conditions present from birth in which the whole body, one side, or particular tissues grow more than expected. Some forms also cause developmental differences or increase the risk of certain tumors.

### Over-Hydration

Excess accumulation of body water, which may cause edema, dilutional hyponatremia, or heart and lung complications.

### Overlap Myositis

Muscle inflammation occurring together with another autoimmune connective-tissue disease, such as lupus, systemic sclerosis, mixed connective-tissue disease, or Sjögren syndrome. It may cause muscle weakness, skin changes, joint symptoms, swallowing problems, or lung disease; treatment is tailored to the affected organs.

### Overlapping Autoimmune Syndrome

A rheumatological or hepatological disorder in which an individual patient simultaneously meets full clinical, serological, and histological diagnostic criteria for two or more distinct established autoimmune diseases (such as autoimmune hepatitis overlapping with primary biliary cholangitis).

**Alternate Names:**  
Overlap Syndrome

### Overlay Denture

A removable denture that fits over remaining teeth, roots, or implants to improve support, chewing, and retention.

### Over-Nutrition

Excessive intake or accumulation of energy or nu-

trients that can contribute to obesity, metabolic disease, or toxicity.

### Overriding Aorta

A birth-related heart abnormality in which the main artery leaving the heart sits over a hole between the lower chambers and receives blood from both. It commonly occurs as part of tetralogy of Fallot.

### Oversedate

To give excessive sedation, causing unusually deep sleep, slowed responses, impaired breathing, or difficulty maintaining the airway.

### Over-The-Counter

A medicine available without a prescription for self-treatment of specified common problems. It can still cause harm, interact with prescribed medicines, or be unsafe with certain illnesses, so the label and dose should be followed carefully.

### Overt Proteinuria

Abnormally high protein loss in the urine that is detectable by routine testing and may indicate kidney disease.

### Overuse Syndrome

Alternate name; see *Repetitive Strain Injury*.

### Overweight

Body weight above the usual range for height. In adults, a Body Mass Index of 25 to less than 30 kg/m<sup>2</sup> is commonly classified as overweight, but it does not directly measure body fat.

### Oviduct

A tube carrying an ovum from the ovary toward the uterus; in humans it is the uterine or fallopian tube.

### Ovomucin

A glycoprotein in egg white that contributes to its gel-like viscosity and helps stabilize the egg-white structure.

### Ovotesticular Disorder of Sexual

---

## Development

A difference of sex development in which a person has both ovarian and testicular tissue, in the same or different gonads. Evaluation is multidisciplinary and considers anatomy, hormones, chromosomes, genetics, fertility, wellbeing, and the person's preferences.

## Ovotestis

A gonad containing both ovarian-type and testicular tissue. It is an uncommon difference of sex development and may produce ovarian hormones, testicular hormones, eggs, sperm, or none of these; evaluation is individualized.

## Ovulation

Release of a mature egg from an ovarian follicle, usually around the middle of the menstrual cycle.

## Ovulation Home Test

An ovulation home test detects the rise in urinary luteinizing hormone that usually occurs shortly before ovulation.

## Ovulation Induction

Pharmacological stimulation of the ovaries to promote the development, maturation, and release of one or more mature fertilizable oocytes in women suffering from anovulation or oligo-ovulation (such as in polycystic ovary syndrome), utilizing agents such as letrozole, clomiphene citrate, or exogenous gonadotropins.

## Ovum

A mature female gamete; in humans, the cell released at ovulation is usually a secondary oocyte, which completes its final meiotic division when a sperm enters. The resulting genetic material combines with the sperm's to form a zygote, the first cell of a developing embryo.

## Oxacillin

A penicillinase-resistant penicillin used mainly to test for methicillin resistance in staphylococci; it is not commonly used as the preferred treatment for most infections.

## Oxalate

A salt or ester of oxalic acid that can form kidney stones and binds calcium.

## Oxalate Nephropathy

Renal tubular injury and interstitial fibrosis resulting from the intrarenal precipitation and deposition of insoluble calcium oxalate crystals within renal tubular lumens, seen in enteric hyperoxaluria (due to fat malabsorption, Crohn disease, or bariatric surgery) and ethylene glycol intoxication.

## Oxalates

Oxalates are salts or esters of oxalic acid that can bind calcium and form calcium-oxalate kidney stones; excess dietary absorption or inherited hyperoxaluria can cause renal injury.

## Oxalic Acid

A dicarboxylic acid whose oxalate salts can contribute to calcium-oxalate kidney stones; concentrated exposure is toxic.

## Oxalic Acid Poisoning

Swallowing oxalic acid can burn the digestive tract and bind calcium, causing muscle cramps, abnormal heart rhythms, seizures, kidney injury, or shock. It is a medical emergency.

## Oxaliplatin

Oxaliplatin is a platinum chemotherapy drug that forms DNA cross-links and is used especially in colorectal cancer regimens; acute cold-triggered neuropathy and cumulative sensory neuropathy are characteristic toxicities.

## Oxaloacetate

A small molecule used in the cell's energy-making cycle. It combines with another molecule to begin a step that produces energy and can also help make glucose and the amino acid aspartate.

## Oxandrolone

An anabolic steroid used in selected conditions involving severe weight loss or catabolic stress; it can



---

affect the liver, lipids, and hormone function.

### **Oxaprozin**

A nonsteroidal anti-inflammatory drug used to relieve pain and inflammation in conditions such as osteoarthritis and rheumatoid arthritis.

### **Oxazepam**

A benzodiazepine medicine used short-term for anxiety or alcohol withdrawal, causing sleepiness and possible dependence.

### **Oxazepam Overdose**

Taking too much oxazepam, a sedative medicine, can cause severe sleepiness, confusion, poor coordination, slowed breathing, coma, or death, especially with alcohol or other sedatives.

### **Oxazole**

Oxazole is a five-membered chemical ring containing oxygen and nitrogen that is used as a building block in medicinal and other chemistry.

### **Oxcarbazepine**

Oxcarbazepine is an antiseizure medicine that blocks voltage-gated sodium channels and treats focal seizures; dizziness, ataxia, rash, and clinically important hyponatraemia may occur.

### **Oxidant**

A substance that can damage cells by taking electrons from other molecules. Oxidants are produced normally during metabolism and inflammation, but excessive amounts can cause oxidative stress and tissue injury.

#### **Alternate Names:**

Oxidizing Agent; Electron Acceptor

### **Oxidants**

Substances or reactive molecules that accept electrons or promote oxidation; excessive oxidant activity can damage proteins, lipids, and DNA.

### **Oxidation**

A chemical process that changes a substance

through loss of electrons or reaction with oxygen.

### **Oxidation-Reduction**

A chemical reaction involving transfer of electrons, in which one substance is oxidized and another is reduced.

### **Oxidative Phosphorylation**

A process mainly occurring at the inner mitochondrial membrane in which electron transfer drives a proton gradient that powers ATP synthase to produce ATP. It is linked to, but distinct from, the earlier pathways that generate the electron carriers.

### **Oxidative Stress**

A harmful imbalance between reactive oxygen-containing molecules and the body's protective antioxidants. Excessive oxidative stress can injure cell membranes, proteins, and DNA and may contribute to inflammation, aging, or disease, although the term alone does not identify a specific illness.

### **Oxidoreductase**

An enzyme that transfers electrons or hydrogen between molecules during a chemical reaction. These enzymes are important in energy production, drug breakdown, and many other cell processes.

### **Oxime**

A type of chemical compound. Some oximes, such as pralidoxime, are used as emergency antidotes because they can restore an important nerve enzyme after organophosphate pesticide poisoning.

### **Oximeter**

A device that estimates blood oxygen level, usually by shining light through a finger, toe, or earlobe. A pulse oximeter also displays pulse rate, but readings can be affected by movement, poor circulation, or skin factors.

### **Oximetry**

Measurement of oxygen levels in blood or tissue. A pulse oximeter estimates blood oxygen through a skin sensor, while arterial testing measures oxy-

---

gen more directly; results can be affected by poor circulation or motion.

### **Oxter**

Oxter is an anatomical term for the armpit or axilla.

### **Oxybuprocaine**

A local anaesthetic eye drop that briefly numbs the surface of the eye. It may be used for examinations or short procedures. Repeated use can delay healing and damage the cornea.

### **Oxycephaly**

A cone-shaped skull caused by premature fusion of cranial sutures, also called acrocephaly.

### **Oxycodone**

Oxycodone is an opioid analgesic used for moderate-to-severe pain; sedation, constipation, respiratory depression, tolerance, dependence, and overdose risk increase with alcohol or other sedatives.

### **Oxycodone Hydrochloride**

The hydrochloride salt of oxycodone, an opioid analgesic used for moderate-to-severe pain; it can cause sedation, constipation, dependence, and fatal respiratory depression in overdose.

### **Oxygen**

A gas essential for aerobic metabolism and tissue energy production.

### **Oxygenases**

Enzymes that incorporate oxygen from molecular oxygen into a substrate; monooxygenases add one oxygen atom and dioxygenases add both.

### **Oxygenated Blood**

Blood that has picked up oxygen in the lungs and carries it through arteries to body tissues. It is usually bright red, while blood returning to the lungs contains less oxygen and more carbon dioxide.

### **Oxygenation**

The process of supplying oxygen to blood and tis-

sues, or the degree to which hemoglobin and tissues are adequately oxygenated.

### **Oxygenator**

A machine that adds oxygen to blood and removes carbon dioxide when the lungs cannot do this adequately. It is used during heart-lung bypass surgery and some forms of extracorporeal life support.

### **Oxygen Consumption**

The amount of oxygen the body uses in a given time. It reflects the body's energy needs and rises with exercise, fever, or illness; it can be measured during exercise testing to assess heart and lung function.

### **Oxygen Content**

The amount of oxygen carried in blood, mainly bound to hemoglobin with a small dissolved component.

### **Oxygen Delivery**

The amount of oxygen transported from the lungs to body tissues each minute. It depends mainly on cardiac output, the amount of hemoglobin, and how much oxygen the hemoglobin carries. Inadequate delivery can injure organs when the body cannot compensate.

### **Oxygen Desaturation Index**

The average number of oxygen desaturation events per hour of sleep, usually reported from a sleep study as the ODI; interpretation depends on the recording method and clinical context.

### **Oxygen Dissociation Curve**

A sigmoidal physiological graph relating the percentage saturation of hemoglobin with oxygen ( $S_aO_2$ ) to the partial pressure of oxygen ( $P_aO_2$ ), whose position is modulated by temperature, arterial pH, carbon dioxide tension, and 2,3-bisphosphoglycerate concentration.

#### **Alternate Names:**

Oxygen-Hemoglobin Dissociation Curve; Oxyhemoglobin Dissociation Curve

---

### **Oxygen Enhancement Ratio**

A measure of how much oxygen increases the effect of radiation on cells. It compares the radiation dose needed with little oxygen to the dose needed when oxygen is normal to produce the same amount of cell injury.

### **Oxygen-Hemoglobin Dissociation Curve**

A graph showing how strongly hemoglobin holds oxygen at different oxygen levels. Temperature, acidity, and carbon dioxide change this binding and therefore affect how readily oxygen is released to tissues.

### **Oxygen Mask**

A mask that delivers supplemental oxygen through the nose and mouth. The type and flow rate are chosen according to oxygen needs; oxygen should be used as prescribed and monitored.

### **Oxygen Pulse**

The amount of oxygen used by the body with each heartbeat, estimated during exercise as oxygen uptake divided by heart rate.

### **Oxygen Safety**

Practices that reduce fire, explosion, pressure, and treatment risks during oxygen use, including keeping oxygen away from flames, smoking, oils, and unsafe equipment.

### **Oxygen Saturation**

The percentage of hemoglobin in the blood carrying oxygen. A finger or ear sensor estimates it without taking blood, while an arterial blood test measures it more directly; the healthy target varies with the person's condition.

### **Oxygen Tent**

A canopy or enclosure used to deliver oxygen around a patient's head; modern oxygen masks, nasal cannulas, or ventilatory devices are usually more precise.

### **Oxygen Therapy**

Giving extra oxygen through a tube, mask, or

breathing machine when the blood oxygen level is too low or is expected to fall. The amount is prescribed and monitored because too much oxygen can also be harmful; oxygen must be kept away from flames.

### **Oxygen Therapy In Infants**

Carefully measured extra oxygen given to a baby whose blood oxygen is too low or whose lungs are not exchanging gases adequately. Clinicians monitor oxygen closely because both too little and too much can injure developing organs.

### **Oxygen Toxicity**

Tissue injury caused by exposure to oxygen at excessively high partial pressures or concentrations, potentially affecting the lungs, eyes, or central nervous system.

### **Oxygen Transport**

The movement of oxygen from the lungs to tissues, mainly bound to hemoglobin and partly dissolved in plasma, with delivery affected by blood flow and hemoglobin concentration.

### **Oxyhaemoglobin**

Hemoglobin with oxygen attached to it. It forms mainly in the lungs and carries oxygen through the bloodstream to body tissues.

### **Oxyhemoglobin**

Hemoglobin bound to oxygen, formed mainly in the lungs and carrying oxygen through arterial blood to tissues.

### **Oxylipins**

Oxylipins are oxygenated fatty-acid derivatives that act as local mediators of inflammation, vascular tone, pain, and other physiologic responses.

### **Oxymetazoline**

A topical alpha-adrenergic agonist used as a nasal decongestant or ocular vasoconstrictor; prolonged nasal use can cause rebound congestion.

---

## Oxymetholone

Oxymetholone is an anabolic steroid used in selected severe anemias; it can cause liver injury, fluid retention, and hormonal side effects.

## Oxymorphone

A potent opioid analgesic used for severe pain; it can cause sedation, constipation, dependence, and life-threatening respiratory depression.

## Oxyphil Cell

A cell with many mitochondria, giving it a distinctive granular appearance under a microscope. Oxyphil cells occur in tissues such as the parathyroid and thyroid, where their significance depends on the organ and the surrounding findings.

## Oxyphil Cell Carcinoma

Alternate name; see *Hürthle Cell Carcinoma*.

## Oxyphilic Adenoma

An oxyphilic adenoma is a usually benign glandular tumor composed of cells rich in mitochondria, such as an oncocyctic parathyroid or salivary-gland tumor.

## Oxytocic

Causing the womb to contract. Oxytocic medicines, such as oxytocin, may start or strengthen labor or help control heavy bleeding after birth; they require monitoring because overly strong or frequent contractions can harm the pregnant person or baby.

## Oxytocics

Medicines that stimulate uterine contractions, used to start or strengthen labor and to prevent or treat heavy bleeding after birth. They require monitoring because overly strong or frequent contractions can harm the pregnant person or baby.

## Oxytocin

A hormone made in the brain and released by the pituitary gland. It helps the uterus contract during labor and causes milk to flow during breastfeeding; a manufactured form is used under monitoring to induce labor or control bleeding after birth.

## Oxytocin Infusion

A controlled drip of synthetic oxytocin into a vein to start or strengthen labor or help control heavy bleeding after birth. It requires continuous assessment because overly strong or frequent contractions can harm the pregnant person or baby.

## Oxyuriasis

Infection with pinworms, especially *Enterobius vermicularis*. The worms commonly cause itching around the anus at night, disturbed sleep, and sometimes abdominal discomfort; treatment usually includes medicine and household hygiene.

## Ozaena

A chronic form of atrophic rhinitis in which the inside of the nose becomes thin and widened, with dry crusts, a foul smell, and a blocked sensation despite an open passage. Medical assessment is needed to exclude infection and other causes.

## Ozena

Ozena is a chronic form of atrophic rhinitis with widened nasal passages, crusting, foul odor, and impaired mucosal function.

## Ozone

Ozone is a reactive form of oxygen that can irritate and injure the airways and lungs when inhaled, despite useful atmospheric and disinfecting roles.

## Oral Allergy Syndrome

An itchy or tingling reaction of the mouth, lips, tongue, or throat after eating certain raw fruits, vegetables, or nuts, usually in people with pollen allergy. Cooking often prevents the reaction, but swelling, breathing trouble, vomiting, or widespread symptoms require urgent assessment because a more serious allergic reaction is possible.

## Opioid Dependence

A state in which the body adapts to repeated opioid exposure and withdrawal symptoms occur when the medicine or drug is reduced or stopped. Physical dependence can occur during appropriate pain treatment and is different from opioid use disorder,

---

which involves problematic use and loss of control.

### **Opioid Addiction**

Alternate name; see *Opioid Use Disorder*.

### **Optic Nerve Swelling**

Swelling of the optic nerve head where the nerve enters the eye. It may be caused by raised pressure inside the skull, inflammation, reduced blood supply, infection, or other eye or nerve disease; new swelling with headache or vision change needs prompt evaluation.





# P

## **P2Y12 Inhibitors**

A subclass of oral antiplatelet medications—including clopidogrel, ticagrelor, and prasugrel—that selectively block the platelet P2Y12 adenosine diphosphate receptor, preventing ADP-mediated platelet activation and arterial thrombosis.

### **Alternate Names:**

P2Y12 Receptor Antagonists; ADP Receptor Antagonists

## **P53 Tumor Suppressor**

A protein that helps prevent damaged cells from dividing or surviving. Changes in the TP53 gene can allow abnormal cells to multiply and contribute to many cancers, but a gene change alone does not determine a person's outcome.

## **Pacemaker**

A small implanted device that sends electrical signals to the heart when its natural rhythm is too slow or unreliable. It may treat fainting or symptoms from slow heartbeats and certain conduction blocks; regular checks monitor the battery and wires.

### **Alternate Names:**

Artificial Pacemaker; Cardiac Pacemaker

## **Pacemaker Implantation**

A procedure to place a pacemaker under the skin of the chest and connect it to the heart with one or more wires, or place a leadless device inside the heart. The device supports an abnormally slow or unreliable rhythm; the approach depends on the rhythm problem and the person's anatomy.

## **Pacemaker Complications**

Problems after pacemaker placement can include bleeding, infection, a collapsed lung, heart or blood-vessel injury, movement or breakage of a lead, blood clots, battery failure, or an abnormal heartbeat. New chest pain, fainting, fever, or worsening breathlessness may signal a serious problem.

## **Pacemaker Malfunction**

Failure of a pacemaker system to sense, pace, capture, or communicate as intended. Causes include lead displacement or fracture, battery depletion, programming problems, electromagnetic interference, and myocardial or metabolic factors.

## **Pacemaker-Mediated Tachycardia**

A rapid reentrant rhythm involving a dual-chamber pacemaker and the patient's cardiac conduction system. It can cause palpitations, dizziness, or reduced device effectiveness and is managed by device reprogramming or other therapy.

## **Pacemakers and Implantable Cardioverter-Defibrillators**

Implantable devices that provide bradycardia pacing, resynchronization, or detection and treatment of dangerous ventricular arrhythmias. Assessment includes indications, programming, leads, battery status, complications, and follow-up.

## **Pacemaker Syndrome**

An adverse hemodynamic and symptomatic consequence of ventricular pacing (typically VVI mode) resulting from loss of physiological atrioventricular synchrony, retrograde ventriculoatrial conduction,

---

and cannon A waves, producing fatigue, syncope, dizziness, and neck pulsations.

### **Pachyderma**

Abnormal thickening and coarsening of the skin, which may occur with chronic inflammation, lymphedema, endocrine disease, or pachydermoperiostosis.

### **Pachydermoperiostosis**

The primary form of hypertrophic osteoarthropathy, usually inherited and characterized by thickened oily or furrowed facial skin, enlargement of the fingers and toes, clubbing, and painful periosteal new bone formation. Severity varies and specialist evaluation is needed.

### **Pachygyria**

A birth-related brain-development abnormality in which the brain's outer surface has fewer, broader folds than usual. It can cause developmental delay, learning difficulties, seizures, movement problems, or other neurologic effects of varying severity.

### **Pachymeningitis**

Inflammation and thickening of the tough outer covering of the brain and spinal cord. It may cause headache, nerve problems, or pressure on the brain and can result from infection, immune disease, or other inflammation.

### **Pachymetry**

Measurement of the thickness of the cornea, the clear front surface of the eye. It is used to assess corneal disease and helps place eye-pressure readings in context.

### **Pachyonychia Congenita**

An inherited skin disorder causing very thick, abnormal nails and painful thick skin on the palms and soles. Some people also develop blisters, cysts, excess sweating, or white patches in the mouth.

### **Pacing**

Planning activity, rest, and effort so symptoms do

not flare while endurance gradually improves. It may help with fatigue, chronic pain, heart or lung disease, and recovery after infection.

### **Pacinian Corpuscle**

A specialized nerve ending deep in the skin and connective tissue that detects deep pressure and vibration. It helps the nervous system sense contact, movement, and the handling of objects.

### **Paclitaxel**

Paclitaxel is a taxane chemotherapy drug that stabilizes microtubules and prevents cell division; it treats several cancers and can cause myelosuppression, neuropathy, alopecia, myalgia, and hypersensitivity reactions.

### **Paclitaxel Liposome**

A liposomal formulation of paclitaxel designed to alter drug distribution and delivery; paclitaxel disrupts microtubules and is used as chemotherapy in selected cancers.

### **PACS**

A computer system used by healthcare services to store, retrieve, view, and share medical images such as X-rays, CT scans, MRI scans, and ultrasound studies.

### **PAES**

Alternate name; see *Popliteal artery entrapment*.

### **Page Kidney**

High blood pressure caused when blood or another mass presses on a kidney from the outside. The pressure activates the kidney's blood-pressure system and may follow injury, bleeding, or treatment with blood thinners.

### **Paget Disease Of Bone**

A long-term disorder in which bone is broken down and rebuilt too quickly and unevenly. Affected bones may become enlarged, weak, painful, or bent and can fracture or press on nearby nerves; many people have no symptoms.

---

**Alternate Names:**

Paget's Disease of Bone; Osteitis Deformans

**Paget Disease Of The Anus**

A rare cancer in the skin around the anus that may look like persistent eczema, redness, itching, soreness, or an erosion. A biopsy is needed, and doctors usually check for cancer in nearby or internal organs.

**Paget Disease Of The Nipple**

A rare form of breast cancer affecting the skin of the nipple and areola. It can cause persistent one-sided redness, scaling, crusting, itching, discharge, or an ulcer. An underlying ductal cancer is present in many cases. Diagnosis uses breast imaging and a skin biopsy; treatment depends on the underlying cancer.

**Alternate Names:**

Mammary Paget's Disease; Paget's Disease of the Breast; Nipple Paget Disease

**Paget Disease Of The Vulva**

A rare cancer in the skin of the vulva that may cause a persistent itchy, red, scaly, sore, or moist patch. A biopsy is needed, and assessment may look for another cancer beneath or elsewhere.

**Pagetoid Reticulosis**

A rare localized variant of cutaneous T-cell lymphoma with solitary slowly enlarging scaly plaques containing atypical pagetoid lymphocytes.

**Paget-Schroetter Syndrome**

An effort-induced deep vein thrombosis (effort thrombosis) of the axillary-subclavian venous system, typically occurring in young, athletic individuals following repetitive, strenuous upper-extremity overhead exertion (e.g., throwing or weightlifting), caused by extrinsic compression of the vein at the thoracic outlet.

**Alternate Names:**

Effort Thrombosis of Subclavian Vein; Upper Extremity DVT

**Paget's Disease**

An ambiguous term that may refer to Paget disease of bone or Paget disease of the breast. Paget disease of bone is a chronic disorder of abnormal bone breakdown and regrowth that can enlarge, weaken, or deform bone; mammary Paget disease is a malignancy involving the nipple and areola, often associated with underlying breast cancer.

**Paget's Disease of the Nipple**

A rare breast cancer involving the skin of the nipple and areola. It may cause redness, scaling, itching, discharge, or a lump.

**Paget's Disease of the Vulva**

A rare cancer or precancerous condition affecting the skin of the vulva. It may cause persistent itching, burning, redness, or a scaly patch and needs biopsy assessment.

**Pain**

An unpleasant physical and emotional experience linked to actual or possible tissue injury. Pain can be short-lived or persistent and is influenced by nerves, inflammation, health, emotions, past experiences, and social circumstances.

**Pain Assessment Scales**

Tools that help a person describe pain intensity or its effect on daily life. They may use numbers, words, faces, or observation of behavior when self-report is not possible. A score tracks change but does not identify the cause or fully measure suffering.

**Painful Bladder Syndrome**

Alternate name; see *Interstitial cystitis*.

**Painful Menstrual Periods**

Painful menstrual periods, or dysmenorrhea, are cramps during menstruation caused by uterine contractions or conditions such as endometriosis or fibroids.

**Painful Swallowing**

Painful swallowing, or odynophagia, is discomfort

---

when swallowing and may result from infection, inflammation, injury, reflux, or a blockage.

### **Painful Intercourse**

Pain in the genital or pelvic area before, during, or after sexual activity; medically called dyspareunia. Causes include dryness, infection, inflammation, pelvic-floor spasm, endometriosis, injury, and other gynecologic or urologic conditions.

### **Pain Management**

Care that identifies the cause of pain and reduces pain while improving function and quality of life. It may combine medicines, physical treatment, psychological support, procedures, and treatment of the underlying condition, with risks reviewed regularly.

### **Painsomnia**

Sleep difficulty or disturbed sleep caused or worsened by pain. It can create a cycle in which poor sleep increases pain sensitivity and fatigue; management addresses the pain, sleep habits, medicines, mood, and underlying condition.

### **Pain Medicine**

The medical specialty concerned with assessing and treating acute, chronic, and complex pain through medication, procedures, rehabilitation, and multidisciplinary care.

#### **Alternate Names:**

Pain Management; Algology

### **Pain Perception**

The process by which the nervous system notices possible or actual tissue injury and the brain interprets it as pain. Attention, emotions, past experiences, inflammation, and medicines can change how pain feels.

### **Pain Threshold**

The minimum intensity of a stimulus that a person perceives as painful; it varies with tissue, context, attention, inflammation, medications, and neurologic state.

### **Pain Unit**

A clinical service specializing in assessment and treatment of acute, chronic, cancer, neuropathic, or procedural pain using medicines, procedures, rehabilitation, and psychological care.

### **PA/IVS**

Alternate name; see *Pulmonary atresia with intact ventricular septum*.

### **Palatal Muscle**

Muscle of the soft palate that elevates or tenses the palate and helps separate the nasal and oral cavities during swallowing and speech.

### **Palatal Myoclonus**

Repeated, involuntary twitching of the soft palate, sometimes causing a clicking sound in the ear or difficulty speaking and swallowing. It may follow a brainstem injury or occur without an identified cause.

### **Palate**

The roof of the mouth, consisting of a bony hard palate and muscular soft palate.

### **Palate Neoplasm**

An abnormal growth in the roof of the mouth. It may be noncancerous or cancerous and can cause a lump, pain, bleeding, loose teeth, difficulty swallowing, or speech changes.

### **Palatine**

Relating to the palate, the roof of the mouth, or the palatine bones. Palatine structures include the tonsils, nerves, and blood vessels of the mouth; problems may include tonsillitis, cleft palate, or infection near a tooth.

### **Palatine Tonsil**

One of two masses of lymphoid tissue at the sides of the throat that help detect inhaled or swallowed pathogens.

---

### Palatorrhaphy

Palatorrhaphy is surgical repair of a cleft or defect of the palate to restore separation between the oral and nasal cavities and improve feeding and speech.

### Palatoschisis

A congenital cleft or failure of fusion of the palate, commonly called cleft palate, which may affect feeding, speech, hearing, and dental development.

### Paleness

Unusual lightening of the skin or mucous membranes, which may result from reduced blood flow, anemia, shock, cold, or normal variation.

### Paleo Diet

An eating pattern based on foods thought to resemble those available to early humans, such as meat, fish, vegetables, fruits, nuts, and seeds. It excludes or limits grains, legumes, and dairy and may not provide balanced nutrition for everyone.

### Pale or Clay-Colored Stools

Light-colored stool caused by reduced bile pigment reaching the intestine, potentially indicating bile-duct obstruction, gallbladder disease, liver disease, or certain transient conditions.

### Palindromic Rheumatism

Repeated attacks of joint pain and swelling that begin suddenly and resolve completely within hours or days, leaving no lasting joint damage between attacks. Some people later develop rheumatoid arthritis, especially when specific antibodies are present.

### Paliperidone

An antipsychotic medicine used to treat schizophrenia and schizoaffective disorder. It changes the action of certain brain chemicals and is available as tablets or long-acting injections; possible effects include sleepiness, movement problems, weight gain, hormone changes, and heart-rhythm changes.

### Palivizumab

A monoclonal antibody against respiratory syncytial virus fusion protein, used prophylactically in selected high-risk infants and young children during RSV season.

### Palla Sign

A chest radiographic sign of acute pulmonary embolism consisting of an enlarged, dilated descending right pulmonary artery (diameter greater than 16 mm on PA projection) terminating abruptly due to a large impacted thromboembolus and downstream vasoconstriction.

#### Alternate Names:

Palla's Sign; Descending Pulmonary Artery Sign

### Pallesthesia

The ability to feel vibration, usually tested with a vibrating tuning fork on the fingers or toes. Reduced vibration sense may result from nerve disease, spinal-cord problems, poor circulation, or other neurologic conditions.

### Palliative Care

Medical care for serious illness that relieves pain and other symptoms and supports daily life, emotional health, and family needs. It can begin at any stage and can be provided alongside treatment intended to control or cure disease.

### Palliative Radiation

Radiation therapy given to relieve symptoms caused by cancer, such as pain, bleeding, obstruction, or pressure from a tumor. It is usually intended to improve comfort or function rather than cure the cancer and may be delivered in a short treatment course.

### Palliative Rehabilitation

Rehabilitation for a person with serious or life-limiting illness that focuses on comfort, independence, meaningful activities, and quality of life. It may address weakness, breathlessness, pain, fatigue, movement, communication, swallowing, or daily tasks.

---

## Palliative Surgery

Surgery intended to relieve symptoms, prevent complications, or improve function when the underlying disease may not be curable. It can ease blockage, bleeding, pain, breathing difficulty, or problems caused by advanced cancer.

## Palliative Treatment

Treatment intended to relieve symptoms and improve comfort or quality of life rather than cure the underlying disease.

## Pallidotomy

Brain surgery that makes a carefully placed small injury in an area involved in movement control. It may reduce severe Parkinson disease symptoms such as involuntary movements when medicines no longer provide adequate control.

## Pallister-Hall Syndrome

A rare inherited condition that may cause an extra finger or toe, a noncancerous growth near the brain's hormone-control center, and abnormalities of the anus, kidneys, airway, or pituitary hormones.

## Pallor

Unusual paleness of skin or mucous membranes, often due to reduced blood flow, anemia, or other illness.

## Palm

The inner surface of the hand between the wrist and fingers. It contains thick skin, creases, muscles, nerves, and blood vessels that help with grip, touch, movement, and protection.

## Palmar

Relating to the palm or palmar surface of the hand, including the palmar sides of the fingers.

## Palmar Erythema

A persistent, symmetrical reddening of the thenar and hypothenar eminences of the palms with sparing of the central palm, often accompanied by warmth. It results from localized microvascular dilatation

driven by elevated circulating estrogen levels or altered hepatic metabolism in chronic liver disease, cirrhosis, pregnancy, and thyrotoxicosis.

## Palmar Erythema Of Pregnancy

Redness of both palms that can occur during pregnancy, often because of normal blood-vessel and hormone changes. It is usually harmless and fades after birth, but new itching, yellowing, pain, or other symptoms need assessment.

## Palmar Grasp Reflex

A primitive reflex in which an infant flexes the fingers around an object placed in the palm. It is normally present at birth and gradually diminishes as voluntary grasp develops; asymmetry or persistence may warrant neurologic assessment.

## Palmar Hyperhidrosis

Alternate name; see *Hyperhidrosis*.

## Palmar Reflex

The palmar reflex is an infant response in which stroking the palm causes finger flexion; persistence or asymmetry beyond the expected age may indicate neurologic dysfunction.

## PALM-COEIN Classification

A system clinicians use to organize the causes of abnormal bleeding from the uterus. PALM covers structural causes such as polyps, adenomyosis, fibroids, and cancer; COEIN covers clotting, ovulation, lining, medicine-related, and other causes. It helps guide evaluation and treatment.

### Alternate Names:

FIGO AUB Classification; PALM-COEIN System

## Palmitate

Palmitate is the salt or ester form of palmitic acid and participates in fatty-acid metabolism, lipid storage, and protein modification.

## Palmitic Acid

A 16-carbon saturated fatty acid common in dietary and stored fats; excess saturated-fat intake can con-



---

tribute to adverse lipid profiles, although the body also synthesizes it normally.

### **Palmoplantar Hyperkeratosis**

Excessive thickening of the skin on the palms and soles, inherited or acquired, causing calluses, fissures, pain, impaired walking, and sometimes infection.

### **Palmoplantar Keratoderma (Keratosis Palmaris et Plantaris)**

Excessive thickening of the skin on the palms and soles. It may be inherited or acquired and can cause fissures, pain, impaired walking, sweating problems, or secondary infection.

### **Palonosetron Hydrochloride**

A long-acting 5-HT<sub>3</sub> serotonin-receptor antagonist used to prevent nausea and vomiting caused by chemotherapy, surgery, or selected other treatments.

### **Palpable**

Palpable means detectable by touch during physical examination, such as a mass, enlarged organ, lymph node, pulse, tenderness, or abnormal surface.

### **Palpable Purpura**

Raised, indurated, non-blanching purpuric papules or plaques resulting from transmural inflammation and fibrinoid necrosis of post-capillary venules with red cell extravasation, serving as the clinical hallmark of cutaneous small-vessel leukocytoclastic vasculitis and IgA vasculitis (Henoch-Schönlein purpura).

### **Palpate**

To examine part of the body by feeling it carefully with the hands. A healthcare professional may assess size, shape, tenderness, warmth, movement, swelling, firmness, pulses, or unusual lumps.

### **Palpation**

Examination by touching the body to assess tenderness, temperature, texture, masses, pulses, organ size, and movement.

### **Palpebra**

The anatomical term for an eyelid. Eyelids protect the eye, spread tears over the cornea, and limit light entry; disorders include drooping, inward or outward turning, inflammation, styes, and blocked oil glands.

### **Palpebral Conjunctiva**

The thin, moist membrane lining the inner surfaces of the eyelids. It joins the membrane covering the front of the eye and helps protect, lubricate, and move tears across the eye's surface.

### **Palpebral Slant**

The direction and inclination of the axis between the inner and outer canthi of the eyes. An upward (upslanting) or downward (downslanting) palpebral slant may represent normal familial variation or serve as a dysmorphic feature in genetic syndromes such as Down syndrome or Noonan syndrome.

#### **Alternate Names:**

Palpebral Fissure Slant; Upslanting Palpebral Fissures; Downslanting Palpebral Fissures

### **Palpitation**

Awareness of the heart beating fast, forcefully, irregularly, or as if it skipped a beat. Causes include stress, stimulants, anemia, thyroid disease, medicines, and abnormal rhythms.

#### **Alternate Names:**

Cardiac Palpitations; Palpitations

### **PALS**

Alternate name; see *Pediatric Advanced Life Support*.

### **Palsy**

Weakness or loss of movement in a body part, often affecting muscles supplied by a particular nerve. It may be temporary or lasting and can result from nerve injury, infection, stroke, inflammation, or a problem present from birth.

### **Palsy Of The Long Thoracic Nerve**

Weakness of the nerve that activates a muscle hold-

---

ing the shoulder blade against the ribs. It can cause a winged shoulder blade, difficulty raising the arm, and pain after injury, infection, surgery, or pressure on the nerve.

### **Pamidronic Acid**

An intravenous bisphosphonate that inhibits osteoclast-mediated bone resorption and is used for selected hypercalcemia, bone complications of cancer, and Paget disease.

### **Pampiniform Plexus**

A network of veins in the spermatic cord that drains the testis and helps cool blood entering the testicular artery.

### **Panacinar Emphysema**

A form of emphysema in which air sacs and small airways are damaged throughout affected areas of the lung. It can cause breathlessness and reduced lung function and is strongly associated with severe alpha-1 antitrypsin deficiency.

#### **Alternate Names:**

Panlobular Emphysema; Lower Lobe Emphysema

### **Pancoast Syndrome**

Symptoms caused by a cancer at the top of the lung growing into nearby nerves or other tissues. It may cause severe shoulder or arm pain, hand weakness, numbness, and drooping of one eyelid with a smaller pupil.

### **Pancoast Tumor**

A lung cancer at the top of the lung that can grow into nearby nerves, blood vessels, or the spine. It may cause severe shoulder or arm pain, hand weakness, or drooping of one eyelid.

### **Pancolitis**

Inflammation affecting the entire large intestine. It may result from ulcerative colitis, infection, reduced blood flow, or another inflammatory condition and can cause diarrhea, abdominal pain, fever, or bleeding.

### **Pancreas**

An organ behind the stomach that makes digestive juices and hormones. Its digestive enzymes enter the small intestine, while insulin and other hormones help control blood sugar and digestion.

### **Pancreas Divisum**

A birth-related difference in which two drainage tubes of the pancreas do not join normally. Most people have no symptoms, but some develop repeated pancreatitis because pancreatic fluid does not drain freely.

#### **Alternate Names:**

Isolated Dorsal Pancreatic Duct

### **Pancreas Enzyme**

An enzyme produced by the pancreas, such as amylase, lipase, or proteases, that helps digest carbohydrate, fat, or protein in the small intestine.

### **Pancreas Infection**

Infection involving the pancreas or a pancreatic collection, usually occurring with pancreatitis, surgery, trauma, or spread from nearby infection.

### **Pancreas Inflammation**

Alternate name; see *Pancreatitis*.

### **Pancreas Neoplasm**

An abnormal growth in the pancreas that may be noncancerous, precancerous, or cancerous. Its effects depend on its location and may include pain, jaundice, digestive problems, or diabetes.

### **Pancreas Transplant**

A procedure that replaces a diseased pancreas with a healthy donor pancreas, usually to restore insulin production in selected people with severe diabetes.

### **Pancreas Transplantation**

Alternate name; see *Pancreas Transplant*.

### **Pancreatectomy**

Surgery to remove part or all of the pancreas. It may treat a pancreatic tumor, severe injury, or another

---

serious disease; removal of pancreatic tissue can affect digestion and blood-sugar control.

### **Pancreatic Abscess**

A pocket of pus in or near the pancreas, usually after severe or infected pancreatitis. It may cause continuing fever, worsening abdominal pain, vomiting, or sepsis and often needs drainage and antibiotics.

### **Pancreatic Adenocarcinoma**

The most common cancer of the pancreas, arising from cells in its digestive-juice ducts. It may cause jaundice, abdominal or back pain, weight loss, or no early symptoms; smoking, chronic pancreatitis, diabetes, obesity, and inherited risks can contribute.

### **Pancreatic Cancer**

Cancer that begins in the pancreas, most often in cells lining its digestive-juice ducts. It may cause jaundice, abdominal or back pain, weight loss, or digestive problems, but early cancer may cause no symptoms; treatment depends on type and spread.

#### **Alternate Names:**

Pancreatic Carcinoma; Pancreatic Adenocarcinoma; Cancer of the Pancreas

### **Pancreatic Carcinoma**

Pancreatic carcinoma is cancer arising in the pancreas, most often from duct cells, and may cause jaundice, pain, weight loss, or digestive problems.

### **Pancreatic Cysts**

Fluid-filled sacs in or around the pancreas. Some follow inflammation or injury, while others are growths that can become cancerous; imaging and sometimes fluid or tissue testing guide follow-up or treatment.

#### **Alternate Names:**

Pancreatic Pseudocysts; Mucinous Cystic Neoplasms; Serous Cystadenomas

### **Pancreatic Diseases**

Disorders affecting the pancreas's digestive or hormone-producing functions, including inflammation, cysts, tumors, duct blockage, and diabetes-

related damage. They may cause abdominal pain, vomiting, weight loss, jaundice, abnormal blood sugar, or poor digestion.

### **Pancreatic Duct**

The main tube carrying digestive fluid from the pancreas into the first part of the small intestine. It often joins the bile duct before opening into the intestine; blockage can cause pain, jaundice, or pancreatitis.

### **Pancreatic Ductal Adenocarcinoma**

The most common exocrine pancreatic malignancy, arising from ductal epithelium and often presenting late with painless jaundice, weight loss, pain, or pancreatitis. Diagnosis and staging rely on imaging, tissue sampling when needed, and multidisciplinary assessment.

### **Pancreatic Elastase**

A stool test that estimates how well the pancreas makes digestive enzymes. A low result supports pancreatic insufficiency, which can cause greasy stools, weight loss, and vitamin deficiency, but watery stool and other factors can affect interpretation.

### **Pancreatic Enzyme**

A digestive substance made by the pancreas and released into the small intestine. Amylase breaks down starch, lipase breaks down fat, and proteases break down protein; bicarbonate helps neutralize stomach acid.

### **Pancreatic Exocrine Insufficiency**

Alternate name; see *Exocrine Pancreatic Insufficiency*.

### **Pancreatic Fistula**

A pancreatic fistula is an abnormal passage through which pancreatic juice leaks from the pancreas, often after trauma or surgery, causing fluid collections, ascites, or skin drainage.

### **Pancreatic Hormones**

Hormones produced by pancreatic islet cells, includ-

---

ing insulin, glucagon, somatostatin, and pancreatic polypeptide, which regulate glucose, digestion, and energy metabolism.

### **Pancreatic Inflammatory Cysts**

Alternate name; see *Pancreatic Cysts*.

### **Pancreatic Insufficiency**

Failure of the pancreas to produce or deliver enough digestive enzymes or other secretions for normal digestion and absorption. It may cause greasy stools, diarrhea, bloating, weight loss, or vitamin deficiency; exocrine insufficiency specifically refers to inadequate digestive enzymes.

### **Pancreatic IPMN**

Alternate name; see *Pancreatic Cysts*.

### **Pancreatic IPMN Entry**

Alternate name; see *Pancreatic Cysts*.

### **Pancreatic Islet**

A pancreatic islet is a cluster of endocrine cells that releases hormones such as insulin, glucagon, and somatostatin to regulate blood glucose and digestion.

### **Pancreatic Juice**

Pancreatic juice is an alkaline digestive secretion containing bicarbonate and enzymes such as amylase, lipase, and proteases, delivered into the duodenum.

### **Pancreatic Mass**

A lump or abnormal area seen in the pancreas on an imaging test. It may be a cyst, inflammation, scar, or tumor; further scans, blood tests, or a tissue sample may be needed to determine its cause.

### **Pancreatic Necrosis**

Death of pancreatic or nearby fatty tissue, usually during severe acute pancreatitis. The dead tissue may remain sterile or become infected, causing fever, worsening pain, sepsis, and organ failure.

**Alternate Names:**

Necrotizing Pancreatitis; Acute Necrotizing Pancreatitis

### **Pancreatic Neuroendocrine Tumor**

A tumor arising from hormone-related cells in the pancreas. It may release excess hormones or cause no hormone symptoms and can grow slowly or aggressively; its behavior depends on size, spread, cell features, and growth rate.

### **Pancreatic Neuroendocrine Tumors**

Tumors arising from hormone-making cells in the pancreas. Some release excess hormones, such as insulin or gastrin, while others do not. Their behavior depends on size, spread, cell appearance, growth rate, and hormone production.

**Alternate Names:**

pNETs; Islet Cell Tumors; Pancreatic Endocrine Neoplasms

### **Pancreatic Non-Inflammatory Cysts**

Alternate name; see *Pancreatic Cysts*.

### **Pancreaticobiliary Cytology**

Cytologic evaluation of pancreatic and biliary lesions, usually from fine-needle aspiration, brushings, or bile-duct sampling. It is used to assess neoplasia, inflammation, and obstructive lesions alongside imaging and histology.

### **Pancreaticoduodenectomy**

Pancreaticoduodenectomy, or the Whipple procedure, removes the head of the pancreas and nearby parts of the digestive tract to treat selected tumors and other serious disease.

### **Pancreatic Pseudocyst**

A walled-off collection of pancreatic fluid that usually develops after acute or chronic pancreatitis. It has no true cell lining and may cause abdominal pain, vomiting, early fullness, infection, bleeding, or blockage of nearby organs.

### **Pancreatic Serous Cystadenoma**

Alternate name; see *Pancreatic cysts*.

---

## Pancreatic Surgery

An operation on the pancreas to remove diseased tissue, drain a collection, repair an injury, or treat cancer. The type of surgery depends on the problem and its location.

## Pancreatic Trauma

Injury to the pancreas caused by blunt or penetrating trauma. Assessment may require CT, laboratory testing, and observation or intervention according to the injury.

## Pancreatitis

Inflammation of the pancreas that may be acute or chronic. Causes include gallstones, alcohol, medicines, metabolic disorders, and other conditions. Acute disease may cause severe upper-abdominal pain, vomiting, fluid loss, infection, tissue death, or organ failure; chronic disease can permanently scar the pancreas and cause poor digestion, weight loss, or diabetes.

### Alternate Names:

Pancreatic Inflammation; Acute or Chronic Pancreatitis

## Pancreatoblastoma

A rare cancer of the pancreas that occurs mainly in children. It may cause an abdominal lump, pain, vomiting, weight loss, or hormone-related symptoms and requires specialist imaging, biopsy, staging, and treatment.

## Pancreatography

Imaging of the tubes that carry digestive juices from the pancreas, usually after contrast dye is introduced. It can show narrowing, blockage, leaks, stones, or other changes and may be done during ERCP or with MRI; ERCP also carries a risk of pancreatitis.

## Pancreazymine

An older name for cholecystokinin, a hormone that stimulates pancreatic enzyme secretion and gallbladder contraction.

## Pancrelipase

Pancrelipase is a mixture of pancreatic digestive enzymes used as replacement therapy for exocrine pancreatic insufficiency, including that caused by cystic fibrosis or pancreatic disease.

## Pancuronium

A long-acting medicine that temporarily prevents skeletal muscles from contracting during anesthesia or assisted breathing. It causes paralysis but does not relieve pain or cause unconsciousness and must be given with reliable airway and breathing support.

## Pancytopenia

A reduction in red blood cells, white blood cells, and platelets at the same time. It can cause tiredness, infections, and bleeding and may result from medicines, infection, immune disease, or a problem in the bone marrow.

## PANDAS

A proposed condition in which obsessive-compulsive symptoms or movement tics begin suddenly or worsen after a streptococcal infection in a child. The diagnosis remains debated, so clinicians must also assess common neurologic, psychiatric, and infectious causes.

## Pandemic

An epidemic that spreads across multiple countries or continents and affects many people. It may occur when an infectious agent spreads widely among populations with limited previous immunity.

## Panendoscopy

Examination of several connected areas of the throat and upper swallowing and breathing passages with a viewing instrument, often including the throat, voice box, and swallowing tube. Tissue samples may be taken during the procedure.

## Paneth Cell

A cell at the base of a small-intestinal gland that releases substances helping control bacteria and other microbes in the gut. It supports the intestinal lining's normal defense system.

---

### **Panhypopituitarism**

Failure of the pituitary gland to make most or all of its important hormones. It can cause tiredness, low blood pressure, infertility, growth or thyroid problems, and inability to respond normally to illness; lifelong hormone replacement may be needed.

### **Panic**

Panic is sudden intense fear or distress with autonomic symptoms such as racing heart, sweating, trembling, breathlessness, or a sense of impending danger.

### **Panic Attack**

A sudden surge of intense fear or discomfort that usually reaches peak intensity within minutes. Symptoms may include a racing or pounding heart, sweating, trembling, breathlessness, chest pain, nausea, dizziness, tingling, a sense of losing control, or fear of dying. A first or atypical episode—especially with chest pain or breathing difficulty—should not be assumed to be panic until urgent medical causes are considered.

### **Panic Attacks**

Plural of panic attack; see *Panic Attack*.

### **Panic Attacks And Panic Disorder**

Panic attacks are sudden episodes of intense fear; panic disorder involves recurrent attacks and persistent concern about having more.

### **Panic Disorder**

Repeated unexpected panic attacks followed by ongoing worry about more attacks or major changes in behavior to avoid them. Attacks peak within minutes and may cause a racing heart, breathlessness, trembling, dizziness, or fear of dying.

### **Panlobular Emphysema**

A type of permanent lung damage affecting the whole small air-sac unit, often more severe in the lower lungs. It is strongly associated with alpha-1 antitrypsin deficiency and can cause breathlessness, cough, and reduced exercise ability.

### **Panmyelopathy**

A term for widespread disease affecting the bone marrow and sometimes the spinal cord. The exact meaning depends on context and may involve reduced production of several blood-cell types together with neurologic problems.

### **Panniculectomy**

Surgical removal of a hanging apron of excess lower-abdominal skin and fat, usually after substantial weight loss; it is distinct from a full abdominoplasty.

### **Panniculitis**

Inflammation of the layer of fat beneath the skin, usually causing tender red or purple lumps, often on the legs. Causes include infection, immune disease, medicines, injury, and pancreatic disease; treatment depends on the cause.

### **Panniculus**

A panniculus is a fold or layer of subcutaneous fatty tissue, especially the lower abdominal apron that may overhang the pubic region.

### **Pannus**

An abnormal layer of inflamed tissue that grows over a joint or the clear front surface of the eye. It can damage cartilage or reduce vision and may occur with rheumatoid arthritis, trachoma, or other chronic inflammation.

### **Panophthalmitis**

Severe infection and inflammation involving the entire eye and its surrounding tissues. It can cause intense pain, redness, swelling, discharge, and major vision loss.

### **Panoramic Radiograph**

An extraoral radiograph displaying both jaws, teeth, temporomandibular regions, and surrounding structures in one image. It is useful for surveying dentition, development, impaction, and gross osseous pathology but has less fine detail than intraoral radiographs.



---

### **Panretinal Photocoagulation**

A laser treatment that places many small burns in the outer retina to reduce the abnormal blood-vessel growth caused by severe retinal disease, especially diabetic retinopathy. It can preserve central vision but may reduce side or night vision.

### **Pantothenic Acid**

Vitamin B5, needed to help cells make energy and produce fatty acids and certain hormones.

### **Pantothenic Acid And Biotin**

Two B vitamins involved in energy metabolism, fatty-acid synthesis, and other enzyme reactions; deficiency is uncommon but can occur with severe malnutrition, malabsorption, or specific inherited disorders.

### **Panuveitis**

Inflammation affecting all the main internal layers around the eye, including the iris, the gel inside the eye, and the tissues near the retina. It can cause pain, redness, light sensitivity, floaters, and vision loss.

### **Papain**

A protein-breaking enzyme found in papaya latex. It is used in some foods and products, but evidence for many medicinal claims is limited; it can irritate tissues or cause allergic reactions.

### **Papillary Fibroelastoma**

A usually noncancerous, small growth attached to a heart valve. It often causes no symptoms but can release a clot and lead to a stroke or other blockage, especially when it is mobile; treatment depends on its size, location, and risk.

### **Papillary Muscle**

Muscles inside the lower chambers of the heart that connect by thin cords to the heart valves. They tighten when the heart contracts to keep the valves from turning inside out and prevent blood from flowing backward.

### **Papillary Thyroid Carcinoma**

The most common type of thyroid cancer. It often grows slowly and may spread to nearby neck lymph nodes; treatment and outlook depend on the tumor's size, spread, and other features.

### **Papillectomy**

Surgical removal of a papilla or papillary growth, such as a tumor near the bile and pancreatic ducts.

### **Papilledema**

Swelling of the area where the optic nerve enters the eye because pressure inside the skull is raised. It can threaten vision and may result from bleeding, a mass, infection, or other serious illness.

### **Papilloedema**

Swelling of the optic discs caused by raised intracranial pressure.

### **Papilloma**

A usually noncancerous growth from a surface lining, often appearing as a wart, skin tag, or small finger-like projection. Some are linked to human papillomavirus; the site, appearance, and tissue examination determine whether further treatment is needed.

### **Papillomatosis**

A condition in which multiple papillomas, or wart-like overgrowths of epithelial tissue, develop. Causes and risks vary by body site, virus, inherited condition, or other disease.

### **Papillomatosis Cutis Lymphostatica**

Thick, rough, wart-like skin caused by long-standing swelling from poor lymph or vein drainage, usually in the legs. The skin may become cobblestoned and prone to infection; controlling the swelling and protecting the skin are important.

### **Papillomavirus**

A papillomavirus, especially human papillomavirus, which can cause warts and some anogenital, oropharyngeal, and other cancers.

---

## Papilloma With Atypia / DCIS

### Pathology

Evaluation of an intraductal papillary breast lesion for fibrovascular cores, epithelial atypia, and ductal carcinoma in situ. The key distinction is between benign papilloma, atypical proliferation, in situ carcinoma, and invasive carcinoma.

### Pap Smear

A screening test in which cells are collected from the cervix and examined for early changes that could become cervical cancer or for cancer itself. It may be done alone or with a test for high-risk human papillomavirus (HPV).

### Pap Test

A screening test for cervical cancer and precancerous changes in which cells are collected from the cervix and examined under a microscope. It may also reveal inflammation or some infections, but an abnormal result does not by itself diagnose cancer.

### Papular

A papule: a small, solid, elevated skin lesion <1 cm in diameter (palpable), of various colors and etiologies.

### Papular Urticaria

A hypersensitivity reaction to insect bites, especially from fleas, bedbugs, or mosquitoes, causing recurrent intensely itchy papules or small vesicles, often clustered on exposed areas in children. Diagnosis is clinical.

### Papule

A solid, elevated, circumscribed primary skin lesion measuring up to one centimetre in diameter, containing no visible fluid. Papules may be dome-shaped, flat-topped, umbilicated, or verrucous, and arise from localized cellular infiltration, epidermal hyperplasia, metabolic deposition, or dermal edema in conditions such as acne vulgaris, lichen planus, molluscum contagiosum, viral warts, and psoriasis.

#### Alternate Names:

Papules; Primary Skin Lesion

### Papules

Alternate name; see *Papule*.

### PAPVR

Alternate name; see *Partial anomalous pulmonary venous return*.

### Paracentesis

A procedure in which a clinician uses a needle to remove fluid from the abdomen. The fluid may be tested to find its cause or removed to relieve pressure and discomfort.

### Paracetamol

Also called acetaminophen, paracetamol reduces pain and fever. Excessive doses can cause serious liver injury, especially with alcohol or other products containing paracetamol.

### Paracetamol Overdose

Taking too much paracetamol, also called acetaminophen, can seriously damage the liver even before symptoms appear. The risk depends on the amount, timing, repeated doses, and health of the person; an antidote can reduce the risk of liver failure when given promptly.

### Paracetamol Toxicity

Liver injury caused by excessive acetaminophen (paracetamol) exposure. Severe poisoning may cause nausea, abdominal pain, liver failure, or death and requires urgent assessment.

### Paracusia

A hearing disturbance in which sounds are perceived unusually, inaccurately, or differently from their actual qualities. It may occur with hearing loss, inner-ear disease, neurological conditions, medicines, or mental health disorders.

### Paradichlorobenzene Poisoning

Harm from mothballs or other products containing paradichlorobenzene. It may cause headache, nausea, dizziness, liver injury, or destruction of red blood cells; significant exposure needs poison-

---

control advice.

### **Paraesophageal Hernia**

A hiatal hernia in which part of the stomach moves through the diaphragm beside the esophagus. It may cause obstruction or strangulation even when reflux is absent.

### **Paraesthesia**

An unusual skin sensation such as tingling, pins and needles, burning, prickling, or numbness. It can result from pressure on a nerve, nerve disease, poor circulation, medicines, vitamin deficiency, or other causes.

### **Paraffin**

A mixture of hydrocarbons used in laboratory tissue processing, topical preparations, and other applications; some forms are flammable.

### **Paraffin Poisoning**

Harm from swallowing or breathing paraffin wax or liquid paraffin. Small swallowed amounts may cause stomach upset, but liquid entering the lungs can cause chemical pneumonia.

### **Paraganglioma**

A rare tumor arising from specialized nerve-related cells outside the adrenal glands. Some release hormones that cause high blood pressure, sweating, headaches, or a fast heartbeat; inherited risk may require genetic assessment.

### **Paragonimiasis**

A parasitic infection caused by lung flukes, usually acquired by eating raw or undercooked freshwater crab or crayfish. It commonly affects the lungs and may cause a long-lasting cough, chest pain, fever, or blood-stained sputum.

### **Paragraphia**

A language problem in which a person writes an unintended word, letter, or symbol instead of the one intended. It may occur with damage to language areas of the brain, such as after a stroke, and is

assessed with other speaking, reading, and writing abilities.

### **Parainfectious**

Describing a problem that occurs during or soon after an infection, often because the immune system reacts to the infection rather than because germs directly invade the affected tissue.

### **Parainfluenza**

Parainfluenza viruses are respiratory viruses that commonly cause colds, croup, bronchitis, or pneumonia, especially in children; illness ranges from mild to severe airway disease.

### **Paraldehyde**

An older sedative and anticonvulsant medicine with limited current use.

### **Paralysis**

Loss of voluntary muscle movement in part or all of the body due to nervous-system or muscle disease.

### **Paralysis Agitans**

An older name for Parkinson disease, characterized by tremor, rigidity, and bradykinesia.

### **Paralytic Ileus**

Failure of intestinal peristalsis without a mechanical blockage, causing abdominal distension, pain, vomiting, and absent or reduced bowel sounds.

### **Paramedical**

Relating to healthcare professionals or services that support medical practice, such as physiotherapy, radiography, or emergency medical care.

### **Parametritis**

Inflammation or infection of the tissue beside the uterus, often after pelvic infection, childbirth, miscarriage, or surgery. It may cause pelvic pain, fever, tenderness, abnormal discharge, and requires treatment based on its cause.

---

## Paramnesia

A memory disturbance involving false, distorted, or inaccurately familiar recollections.

## Paramyotonia Congenita

A rare inherited muscle disorder in which muscles become stiff and slow to relax, especially after repeated movement or exposure to cold. The face, eyelids, neck, and hands are often affected; temporary weakness may follow an attack. Avoiding cold and strenuous triggers helps, and medicines may reduce symptoms.

### Alternate Names:

Eulenburg Disease; Paramyotonia Congenita of Eulenburg

## Paranasal Sinuses

Air-filled spaces in the bones around the nose that connect with the nasal passages. Their lining produces mucus, which drains into the nose; inflammation or blockage can cause facial pressure, congestion, and pain.

## Paraneoplastic Autonomic Neuropathy

Autoimmune injury to autonomic nerves associated with cancer, causing problems such as abnormal blood-pressure control, sweating, heart-rate variability, gastrointestinal motility, bladder function, or sexual function. It may accompany broader paraneoplastic neurologic syndromes.

## Paraneoplastic Cerebellar Degeneration

A subacute, debilitating autoimmune paraneoplastic neurological disorder characterized by progressive pancerebellar ataxia, truncal instability, dysarthria, nystagmus, and oscillopsia. Mediated by cytotoxic T cells and cross-reactive circulating onconeural antibodies (such as anti-Yo in ovarian/breast cancer, anti-Hu in small cell lung cancer, or anti-Tr in Hodgkin lymphoma) causing extensive Purkinje cell loss.

### Alternate Names:

PCD; Subacute Cerebellar Degeneration

## Paraneoplastic Disorder

A health problem caused indirectly by cancer rather

than by the tumor invading or spreading to the affected organ. Tumors may release hormones or trigger immune reactions that affect the nerves, skin, blood, joints, or other distant organs.

## Paraneoplastic Encephalomyelitis

An autoimmune inflammatory disorder of the brain, spinal cord, or peripheral nervous system triggered by an underlying cancer rather than direct tumor invasion. It may cause multifocal neurologic symptoms and requires evaluation for neural antibodies and an occult malignancy.

## Paraneoplastic Endocrine Syndromes

Hormonal disorders caused by a cancer or by the body's response to it rather than by direct tumor growth in an endocrine gland. Examples include inappropriate antidiuretic hormone secretion, ectopic adrenocorticotrophic hormone production, and hypercalcemia caused by tumor-related factors.

## Paraneoplastic Neurological Disorder

Alternate name; see *Paraneoplastic syndromes of the nervous system*.

## Paraneoplastic Neurological Syndrome

Alternate name; see *Paraneoplastic syndromes of the nervous system*.

## Paraneoplastic Neurologic Syndrome

A nervous-system problem caused by the immune response to a cancer rather than by the tumor pressing on the nerves. It may cause weakness, loss of balance, memory problems, seizures, or abnormal sensation and can occur before the cancer is found.

## Paraneoplastic Neurologic Syndromes

Rare disorders in which an immune response to cancer also attacks the nervous system. Symptoms may include weakness, imbalance, seizures, memory change, or sensory problems and can precede cancer diagnosis.

## Paraneoplastic Pemphigus

A rare, often severe autoimmune blistering disease

---

associated with an underlying tumor, especially a lymphoid cancer. It causes painful mouth sores and skin lesions and requires specialist treatment.

### **Paraneoplastic Syndrome**

A group of symptoms caused by substances or immune reactions from a cancer, rather than by the tumor pressing on or spreading to the affected area. It may cause hormone, nerve, skin, blood, or joint problems; treating the cancer may improve them.

### **Paraneoplastic Syndromes Of The Nervous System**

Nervous-system problems caused by an immune reaction to a cancer rather than by the cancer growing into the nerves or brain. They may cause weakness, poor balance, seizures, memory change, or abnormal sensation.

#### **Alternate Names:**

Paraneoplastic Neurological Syndromes; PNS

### **Paranoia**

A thought process characterized by excessive or irrational suspiciousness and distrust of others, often manifesting as delusions of persecution, reference, or grandeur.

### **Paranoid**

Characterized by persistent or excessive suspicion and distrust, often involving beliefs that others intend harm. Severe or fixed beliefs may occur in a psychotic disorder and need professional assessment.

### **Paranoid Ideation**

Suspicious or persecutory thoughts that other people intend harm, deceive, monitor, or plot against the person. The significance depends on the degree of conviction, distress, functional effect, and associated symptoms.

### **Paranoid Personality Disorder**

A longstanding pattern of distrust and suspiciousness in which other people's motives are interpreted as threatening or deceitful.

### **Paranoid Psychosis**

An older term for psychosis dominated by delusions of persecution, reference, or grandeur.

### **Paraparesis**

Weakness in both legs caused by disease affecting the spinal cord, brain, nerves, or muscles.

### **Paraphasia**

An unintended substitution, omission, or distortion of a sound or word during speech. It commonly occurs with aphasia after brain injury or stroke and may affect naming, conversation, reading, or writing.

### **Paraphilia**

A persistent and intense sexual interest involving atypical objects, situations, or activities. A paraphilia by itself is not a mental disorder. It becomes a paraphilic disorder when it causes significant personal distress or problems in daily life, apart from distress caused only by social disapproval, or when acting on it involves harm, risk of harm, or people who do not or cannot consent.

### **Paraphilic Disorder**

Alternate name; see *Paraphilic Disorders*.

### **Paraphilic Disorders**

The group of mental disorders in which a paraphilia causes significant personal distress or problems in daily life, apart from distress caused only by social disapproval, or involves harm, risk of harm, or people who do not or cannot consent. The group includes voyeuristic, exhibitionistic, frotteuristic, fetishistic, sexual masochism, sexual sadism, pedophilic, and transvestic disorders, as well as other specified and unspecified paraphilic disorders. A paraphilia without these consequences is not, by itself, a paraphilic disorder.

### **Paraphimosis**

Paraphimosis occurs when a retracted foreskin cannot be returned over the glans, causing painful swelling and possible loss of blood flow. Treatment involves promptly returning the foreskin to its

---

normal position.

### **Paraplegia**

Paralysis of both legs and usually part of the lower trunk. Paralysis of both legs and often part of the trunk, usually from spinal-cord or neurologic disease.

### **Parapneumonic Effusion**

A collection of fluid between the lung and chest wall that develops with pneumonia. Small uncomplicated collections may improve with antibiotics, while infected or loculated fluid can cause persistent fever, chest pain, or breathlessness and may require drainage.

### **Parapneumonic Pleural Effusion**

Pleural fluid accumulation associated with bacterial pneumonia. It may be uncomplicated or progress to a complicated effusion or empyema, requiring imaging, pleural-fluid analysis, antibiotics, and sometimes drainage.

### **Parapneumonic Pleural Effusions and Empyema Thoracis**

Fluid that accumulates beside a pneumonia is a parapneumonic effusion; infected, purulent, or loculated fluid is empyema. Assessment uses imaging and pleural-fluid analysis, and complicated infection often requires antibiotics plus drainage.

### **Parapsoriasis**

A group of chronic scaly patches or thin plaques that resemble psoriasis but have different causes and risks. Some forms remain stable, while large-plaque disease can rarely evolve into cutaneous T-cell lymphoma and needs follow-up.

### **Paraquat**

A highly toxic herbicide. Ingestion can cause injury to the mouth and gastrointestinal tract, kidney failure, and delayed lung injury; severe poisoning has a high risk of death.

### **Paraquat Poisoning**

A medical emergency caused by swallowing or significant exposure to the herbicide paraquat. It can burn the mouth and digestive tract and later cause lung, kidney, or liver failure.

### **Paraseptal Emphysema**

A form of emphysema affecting air sacs near the outer surface of the lung. It may cause few symptoms but can form fragile air pockets that rupture and produce a collapsed lung, especially in tall young adults or people who smoke.

### **Parasite**

An organism that lives on or inside another organism, called the host, and obtains food or shelter from it. Parasites may cause disease in the skin, intestine, blood, or organs.

### **Parasitemia**

The presence and amount of parasites in the bloodstream. It is especially used for malaria and some other blood infections; a higher amount may indicate more severe disease, but interpretation depends on the parasite and the person's symptoms.

### **Parasitic**

Parasitic means relating to an organism that lives in or on a host and obtains nutrients at the host's expense, sometimes causing disease.

### **Parasitic Infection**

An illness caused by a parasite—an organism that lives on or inside another living creature (the host) and survives by taking nutrients at the host's expense. Human parasites belong to three main groups: single-celled protozoa (such as *Plasmodium* causing malaria, *Entamoeba* causing amebic dysentery, and *Giardia* causing giardiasis, which can multiply rapidly inside the body), helminths or parasitic worms (such as roundworms, tapeworms, pinworms, hookworms, and flukes, which are multicellular and can live in tissues for years), and ectoparasites (such as scabies mites, lice, fleas, and ticks that feed on or burrow into the skin). Parasites enter the body through contaminated water or unwashed food, un-



---

dercooked meat, insect bites, walking barefoot on contaminated soil, or direct physical contact. Symptoms depend on the parasite and can range from watery or bloody diarrhea, stomach cramps, and nausea, to prolonged fevers, extreme fatigue, severe anemia, skin rashes, and chronic organ enlargement or damage. Parasitic infections are diagnosed using stool tests, blood smears, or tissue exams, and are treated with targeted antiparasitic medicines (such as antiprotozoals or worm-clearing anthelmintics). Access to clean drinking water, proper sanitation, thorough cooking, and insect protection are key to prevention.

**Alternate Names:**

Parasitic Infections; Parasitic Disease; Parasitosis; Parasitic Infestation

**Parasitic Worm**

A worm that lives in or on a host and uses the host for nourishment. Human infections may affect the intestine, blood, lymph system, lungs, liver, skin, or other tissues and can cause anemia, pain, or organ damage.

**Parasitology**

The branch of microbiology that studies parasites, including protozoa, helminths, and ectoparasites. Parasitic infections are a major global health problem, particularly in tropical and subtropical regions. Protozoan parasites include *Plasmodium* (malaria), *Trypanosoma* (sleeping sickness, Chagas disease), and *Leishmania* (leishmaniasis).

**Parasitosis**

Infestation or infection by a parasite; the term may describe a confirmed parasitic disease or, in delusional parasitosis, a fixed false belief of infestation without evidence.

**Parasomnia**

A parasomnia is an undesirable movement, behavior, emotion, or experience occurring during sleep or transitions between sleep and wakefulness.

**Parasomnias**

Unwanted movements, behaviors, emotions, or expe-

riences that occur during sleep or at the transition between sleep and wakefulness.

**Parasternal Heave**

A palpable lifting or thrusting sensation felt over the left parasternal border during systole, examined using the heel of the examiner's hand. It is a sign of right ventricular hypertrophy or right ventricular volume overload, commonly seen in severe pulmonary hypertension or congenital heart defects.

**Parasuicide**

Older term; see *Suicide Attempt* or *Intentional Self-Harm*.

**Parasympathetic Nervous System**

The part of the automatic nervous system that supports resting functions such as digestion, saliva, urination, and slowing the heartbeat. It also helps the body conserve energy and recover after activity.

**Parathyroid**

One of the usually four small glands behind the thyroid gland. They release parathyroid hormone, which helps keep calcium and phosphate levels balanced in the blood and bones.

**Parathyroid Adenoma**

A parathyroid adenoma is a usually benign tumor of a parathyroid gland that can secrete excess parathyroid hormone and cause primary hyperparathyroidism with high calcium.

**Parathyroid Cancer**

Parathyroid cancer is a rare malignant tumor of the parathyroid gland that often causes excessive parathyroid hormone and high blood calcium.

**Parathyroid Carcinoma**

A rare malignant tumor of the parathyroid gland that commonly causes excessive parathyroid-hormone secretion, hypercalcemia, kidney stones, bone disease, or related symptoms.

---

## Parathyroid Crisis

A life-threatening rise in blood calcium caused by severe overactivity of the parathyroid glands. It may cause extreme thirst, vomiting, constipation, confusion, weakness, abnormal heart rhythm, kidney injury, or coma.

## Parathyroidectomy

Surgery to remove one or more overactive parathyroid glands. It may treat primary or kidney-related hyperparathyroidism when hormone excess causes high calcium, kidney stones, bone disease, or other complications; calcium levels must be monitored afterward.

## Parathyroid Gland

A small gland behind the thyroid that releases parathyroid hormone to regulate calcium and phosphate.

## Parathyroid Gland Removal

Surgery to remove one or more overactive parathyroid glands, usually for persistent high blood calcium. It can improve symptoms and protect bones or kidneys; low calcium, bleeding, voice-nerve injury, or persistent disease may occur.

## Parathyroid Glands

Usually four small endocrine glands behind the thyroid that secrete parathyroid hormone to regulate blood calcium and phosphate levels.

## Parathyroid Hormone

A hormone that regulates calcium and phosphate levels in the blood and supports bone remodeling.

## Parathyroid Hormone Blood Test

A blood test measuring parathyroid hormone, interpreted with calcium, phosphate, vitamin D, and kidney function to evaluate parathyroid and calcium disorders.

## Parathyroid Hormone Physiology

When blood calcium falls, the parathyroid glands release more parathyroid hormone. The hormone

acts on bone and kidneys and increases active vitamin D, helping raise calcium and maintain the balance needed for nerves, muscles, and bones.

## Parathyroid Hormone-Related Protein Blood Test

A blood test measuring parathyroid hormone-related protein, used mainly to evaluate humoral hypercalcemia of malignancy and selected disorders of calcium regulation.

## Parathyroid Hyperplasia

Enlargement of one or more small glands behind the thyroid, causing them to release too much parathyroid hormone. This can raise blood calcium and may occur with kidney disease or inherited multiple-gland disorders.

## Parathyroid Surgery

Operations on the parathyroid glands, usually performed for selected cases of primary or secondary hyperparathyroidism. The decision depends on symptoms, calcium levels, kidney involvement, bone disease, overall health, and current clinical guidance.

## Paratubal Cyst

A usually benign fluid-filled cyst adjacent to the fallopian tube, arising from paratubal or embryologic remnants; most are asymptomatic but large cysts may cause pain or torsion.

## Paratyphoid Fever

An infection caused by *Salmonella Paratyphi* bacteria, usually spread through contaminated food or water. It can cause prolonged fever, headache, weakness, abdominal symptoms, and sometimes serious complications without antibiotic treatment.

## Parechovirus

A group of viruses that commonly cause mild respiratory or gastrointestinal illness but can cause severe sepsis-like illness, meningitis, or encephalitis in newborns and young infants.

---

## Parenchyma

The working tissue of an organ, distinguished from its supporting framework, blood vessels, and connective tissue. For example, lung parenchyma performs gas exchange, while liver parenchyma processes nutrients and harmful substances.

## Parenchymal

Parenchymal means relating to the functional cells of an organ, distinguished from its supporting connective tissue, vessels, and framework.

## Parenteral

Parenteral means delivered by a route other than the gastrointestinal tract, commonly by injection or infusion such as intravenous, intramuscular, or subcutaneous administration.

## Parenteral Nutrition

Nutrition given through a vein when the digestive tract cannot safely absorb enough nutrients. The solution can contain water, glucose, amino acids, fats, vitamins, and minerals. It can cause bloodstream infection, liver problems, or abnormal blood sugar and salts, so it requires monitoring.

## Parenting

The care, protection, guidance, and support provided to a child. Healthy parenting includes meeting basic needs, setting safe limits, nurturing development, and seeking help when concerns arise.

## Paresis

Partial weakness or incomplete paralysis of voluntary muscles. It may affect one limb, one side of the body, both legs, or all four limbs, depending on the nerves, brain, or spinal cord involved.

## Paresthesia

An unusual sensation such as tingling, prickling, burning, pins and needles, or numbness. Brief symptoms often follow pressure on a nerve, while persistent or spreading symptoms need assessment for nerve or circulation problems.

## Paresthesia (Pins and Needles)

Alternate name; see *Paresthesia*.

## Paricalcitol

Paricalcitol is a vitamin-D receptor activator used to reduce secondary hyperparathyroidism in selected chronic kidney disease, with monitoring for calcium and phosphate changes.

## Parietal

Relating to a wall of a body cavity or to the parietal bone or lobe of the skull and brain.

## Parietal Bone

One of two broad bones forming much of the skull's sides and roof. They protect the brain, meet along the top midline, and connect with several neighboring skull bones.

## Parietal Cell Antibodies

Circulating autoimmune antibodies directed against the gastric parietal cell H<sup>+</sup>/K<sup>+</sup>-ATPase proton pump, detected in over 90 percent of patients with autoimmune atrophic gastritis and pernicious anemia.

### Alternate Names:

PCA; Anti-Parietal Cell Antibodies

## Parietal Lobe

A part of the brain that processes touch, pain, temperature, body position, and spatial awareness. Damage may cause numbness, difficulty recognizing objects by touch, poor coordination, or trouble understanding where parts of the body are.

## Parietal Pericardium

The outer serous layer of the pericardium that lines the inner surface of the fibrous sac surrounding the heart.

## Parietal Pleura

The pleural layer lining the chest wall, diaphragm, and mediastinum; it is sensitive to pain when inflamed.

---

### Parinaud Oculoglandular Syndrome

Conjunctivitis with nearby lymph-node swelling, usually caused by infections such as Bartonella, tularemia, or certain viruses.

### Parinaud Syndrome

A disorder caused by dysfunction of the dorsal mid-brain, characterized by impaired upward gaze, abnormal pupil responses, and convergence-retraction eye movements. Causes include tumors, stroke, multiple sclerosis, and hydrocephalus.

### Paris System For Reporting Urinary Cytology

A standardized framework for urinary cytology that emphasizes detection of high-grade urothelial carcinoma and uses defined categories to improve diagnostic consistency and clinical communication.

### Parity

Parity is the number of pregnancies that reached a viable gestational age, regardless of whether the infants were born alive or stillborn; it differs from gravidity.

### Park-Bench Position

A modified lateral position in which the upper body is partly rotated toward prone; it is used for selected posterior brain and neck procedures.

### Parkinson Disease

Alternate spelling; see *Parkinson's Disease*.

### Parkinsonian Gait

A characteristic abnormal gait seen in Parkinson disease, characterized by universal stooped posture, diminished reciprocal arm swing, shuffling short steps (*marche a petits pas*), difficulty initiating movement (freezing), and involuntary acceleration forward (festination).

#### Alternate Names:

Festinating Gait

### Parkinsonism

A group of movement problems involving slowness,

stiffness, shaking, or balance difficulty. Parkinson disease is one cause, but medicines, other brain disorders, and several less common conditions can produce similar symptoms.

### Parkinson's Disease

A progressive disorder of the nervous system that mainly affects movement. It develops when nerve cells in a part of the brain called the substantia nigra become damaged or die. These cells normally make dopamine, a chemical messenger that helps produce smooth, purposeful movement. Reduced dopamine can cause tremor, slowness of movement, muscle stiffness, balance or walking problems, a softer voice, and smaller handwriting. Non-movement symptoms may include constipation, loss of smell, tiredness, sleep problems, depression, or changes in thinking. Symptoms and their rate of progression vary from person to person.

#### Alternate Names:

Idiopathic Parkinsonism; Paralysis Agitans; PD

### P Arm Of A Chromosome

The shorter arm of a chromosome, labeled p, opposite the longer q arm. A genetic report may use this label to show where a gene change or missing or extra DNA segment is located.

### Paromomycin

Paromomycin is an aminoglycoside antibiotic used mainly in the intestine for selected parasitic infections and sometimes for other infections under specialist care.

### Paronychia

Infection or inflammation of the skin around a fingernail or toenail, causing pain, redness, swelling, and sometimes pus. It may follow nail biting, injury, a hangnail, or repeated water exposure.

### Parosteal Osteosarcoma

Parosteal osteosarcoma is a usually lower-grade bone cancer that grows on the outer surface of a bone, often near the back of the thigh.

---

## Parotidectomy

Surgery to remove part or all of a parotid salivary gland, usually because of a tumor. Possible complications include facial-nerve weakness, bleeding, infection, saliva leakage, numbness, or sweating while eating.

## Parotid Gland

The largest major saliva-producing gland, located in front of and below each ear. It releases saliva into the mouth near the upper back teeth and can become swollen from infection, blockage, inflammation, or a tumor.

## Parotid Neoplasm

An abnormal growth in a parotid salivary gland in front of or below the ear. Most are noncancerous, but some are cancerous; a painless lump, facial weakness, pain, or rapid growth may occur.

## Parotid Tumor

An abnormal growth in a parotid salivary gland. Most parotid tumors are noncancerous, but some are malignant and may cause a lump, pain, or facial-nerve weakness.

## Parotid Tumors

Abnormal growths arising in the parotid salivary glands, located in front of and below the ears.

## Parotitis

Inflammation or infection of one or both parotid salivary glands, which lie in front of and below the ears. It can cause painful swelling, fever, dry mouth, or pus near the upper back teeth. Causes include viruses such as mumps, bacteria, blocked ducts, dehydration, and autoimmune disease.

## Paroxetine Hydrochloride

Paroxetine hydrochloride is an SSRI antidepressant used for depression and several anxiety-related disorders; it can cause sexual side effects and withdrawal symptoms if stopped abruptly.

## Paroxysm

A sudden, often brief attack or burst of symptoms or body activity that may recur. Examples include an episode of pain, coughing, rapid heartbeat, seizure, or involuntary movement; the cause depends on the symptom.

## Paroxysmal Atrial Fibrillation

A clinical subtype of atrial fibrillation in which recurrent, uncoordinated atrial tachyarrhythmias terminate spontaneously or with pharmacological/electrical intervention within seven days of onset (usually within 48 hours).

### Alternate Names:

PAF

## Paroxysmal Atrial Tachycardia

Sudden episodes of rapid, regular atrial rhythm that start and stop abruptly, often producing palpitations, lightheadedness, chest discomfort, or shortness of breath.

## Paroxysmal Cold Hemoglobinuria

A rare immune disorder in which cold temperatures trigger destruction of red blood cells when the blood warms again. It can cause sudden anemia, fever, back pain, jaundice, or dark urine, often after an infection.

## Paroxysmal Hemicrania

A rare headache disorder causing brief, repeated attacks of severe one-sided head or facial pain.

## Paroxysmal Nocturnal Dyspnea

Sudden breathlessness that wakes a person after they have been asleep for a while, often improving after sitting or standing. It commonly occurs with heart failure but may have other heart or lung causes.

## Paroxysmal Nocturnal Haemoglobinuria

A rare acquired blood disorder in which abnormal blood-forming cells make red blood cells vulnerable to immune destruction. It can cause anemia, dark urine, blood clots, low blood counts, kidney injury,

---

or bone-marrow failure.

### **Paroxysmal Nocturnal Hemoglobinuria**

A rare acquired blood disorder in which red cells are destroyed by the immune system. It can cause dark urine, anemia, blood clots in unusual locations, low blood counts, and kidney or other organ problems.

### **Paroxysmal Supraventricular Tachycardia**

Repeated episodes of a very fast, usually regular heartbeat that start and stop suddenly and begin in the upper chambers or connecting electrical tissues of the heart. It may cause pounding heartbeat, dizziness, chest discomfort, breathlessness, or fainting.

#### **Alternate Names:**

PSVT; Paroxysmal Atrial Tachycardia

### **Pars**

A part or portion of an anatomical structure, used in names such as pars interarticularis.

### **Pars Compacta**

A dopamine-producing part of the substantia nigra, a movement-control area deep in the brain. Loss of these nerve cells contributes to the slowness, stiffness, and tremor of Parkinson disease.

### **Parsonage-Turner Syndrome**

An acute neuro-inflammatory disorder affecting the brachial plexus, typically presenting with the abrupt onset of excruciating, unremitting, burning shoulder and periscapular pain, followed days to weeks later by patchy multifocal muscle weakness, flaccid paresis, and rapid denervation atrophy of the shoulder girdle and arm muscles. Often triggered by prior viral infection, vaccination, trauma, or surgical stress.

#### **Alternate Names:**

Neuralgic Amyotrophy; Acute Brachial Radiculitis; Brachial Plexitis

### **Pars Planitis**

A form of inflammation inside the eye, near the edge of the retina. It may cause floaters, blurred vision,

and swelling or other retinal complications and is treated according to its severity and cause.

### **Partial**

Partial means incomplete or involving only a portion, such as partial loss, partial response, partial thickness, or partial seizure; the specific meaning depends on context.

### **Partial Agonist**

A medicine that attaches to a cell receptor and activates it, but produces a smaller effect than a full activator even when all available receptors are occupied. It can reduce the effect of a stronger activator at the same receptor.

### **Partial Anomalous Pulmonary Venous Connection**

Alternate name; see *Partial anomalous pulmonary venous return*.

### **Partial Breast Brachytherapy**

Radiation treatment placed inside or next to the area where a breast cancer was removed, usually after breast-conserving surgery. It treats selected early cancers over a shorter course but may cause skin changes, breast firmness, infection, or delayed healing.

### **Partial Cystectomy**

Surgery to remove part of the urinary bladder, usually for a tumor or localized disease while leaving the rest of the bladder in place. The remaining bladder may need monitoring, and urinary or surgical complications can occur.

### **Partial Denture**

A removable replacement for some missing teeth, supported by remaining teeth, gums, or both. It may use metal or plastic and can improve chewing, speech, appearance, and tooth stability.

### **Partial Hysterectomy**

A common but imprecise term usually referring to supracervical or subtotal hysterectomy, in which the



---

uterine body is removed while the cervix remains. The fallopian tubes and ovaries may or may not be removed separately.

### **Partial Knee Replacement**

Surgery replacing only the damaged compartment of the knee with an artificial surface while preserving healthy bone, ligaments, and cartilage. It may relieve pain from localized arthritis; infection, clots, stiffness, loosening, or later arthritis elsewhere can occur.

### **Partial Nephrectomy**

Partial nephrectomy removes a diseased part of a kidney while preserving the remaining healthy tissue.

### **Partial Remission**

A reduction in signs, symptoms, or measurable disease burden that does not meet criteria for complete remission; the definition depends on the disease and response criteria used.

### **Partial Response**

A measurable improvement after treatment in which a tumor or other disease becomes smaller or less active but does not disappear completely. The exact amount of improvement required is defined by the cancer type, test method, or study criteria.

### **Partial Seizure**

A seizure beginning in a localized region of one cerebral hemisphere, now called a focal seizure, with symptoms determined by the involved network and awareness either preserved or impaired.

### **Partial Thromboplastin Time**

A blood test measuring how long a clot takes to form through several clotting pathways. It helps investigate unusual bleeding and monitor unfractionated heparin; abnormal results may reflect a clotting-factor problem, medicine effect, or another illness.

### **Partial Volume Artifact**

A scan distortion that occurs when one small image area contains different tissues. Their signals are blended, which can blur borders or hide a small finding; thinner slices or higher-resolution images may reduce the effect.

### **Partial-Volume Effect**

A scan limitation in which a small structure and the surrounding tissue are measured together, making the structure appear blurred or its activity seem too high or low. Thinner slices or correction methods may reduce the effect.

### **Participation**

A person's involvement in life situations and activities, a central concept in rehabilitation and the International Classification of Functioning that extends beyond impairment alone.

### **Particle Size**

Particle size is the diameter or size distribution of particles in a powder, suspension, aerosol, or formulation, influencing dissolution, absorption, stability, and delivery.

### **Particulate Matter**

Small solid or liquid particles suspended in air. Fine particles can penetrate the lungs and increase cardiovascular and respiratory risk.

### **Partogram**

A graphic record of cervical change, fetal descent, contractions, and maternal or fetal observations during labor. It can support communication and recognition of abnormal progress, but findings must be interpreted with current labor guidance and the overall clinical situation.

#### **Alternate Names:**

Labor Progress Chart; Partographic Curve

### **Parturition**

The process of childbirth, including cervical ripening, labor contractions, delivery of the fetus, and expulsion of the placenta.

---

## Paruresis

Difficulty urinating in the presence of other people or in public places, often called shy bladder syndrome. It is related to anxiety and can improve with behavioral therapy and gradual exposure.

## Parvimonas Micra

An anaerobic gram-positive bacterium found in the mouth and gastrointestinal tract that can cause dental, bone, joint, or bloodstream infections.

## Parvoviridae

A family of very small DNA viruses; human parvovirus B19 can cause fifth disease, anemia, or fetal complications.

## Parvovirus

A small DNA virus; parvovirus B19 is the main human pathogen and can cause rash, joint symptoms, or red-cell aplasia.

## Parvovirus B19

A human virus spread mainly through respiratory droplets. It commonly causes a mild rash illness in children, but can cause joint pain or severe anemia and may harm a fetus during pregnancy.

## Parvovirus B19 Infection

An infection that may cause fever, a slapped-cheek rash, joint pain, or temporary reduction in red-blood-cell production. It can be serious in people with certain blood disorders or weakened immunity and can affect pregnancy after significant exposure.

## Parvovirus Infection

An infection caused by parvovirus B19 that commonly causes mild fever and rash but may be serious in some people.

### Alternate Names:

Fifth Disease; Erythema Infectiosum; Slapped Cheek Disease

## Passive Exercise

Movement of a person's joint or limb by another person, a machine, or gravity without active muscle

effort from the patient. It can maintain flexibility and reduce stiffness when illness or weakness limits movement, but forceful movement can injure tissues.

## Passive Heat Therapy

Use of external heat, such as warm packs, baths, or saunas, to relax muscles and temporarily reduce pain or stiffness. Excessive heat can cause burns, dehydration, or fainting.

## Passive Immunity

Protection produced by transferred antibodies rather than by the person's own immune response, such as maternal antibodies or immunoglobulin treatment.

## Passive Immunization

Protection produced by giving preformed antibodies, such as immunoglobulin or antivenom, which acts quickly but is temporary because the recipient does not generate durable immune memory.

## Passive Smoking

Breathing smoke from another person's tobacco or nicotine product. Secondhand smoke increases the risk of respiratory disease, heart disease, stroke, and lung cancer and is especially harmful to children and during pregnancy.

## Pasteurella

A genus of bacteria carried by animals that can cause wound infections after bites and other human disease.

## Pasteurella Aerogenes

*Pasteurella aerogenes* is a rare *Pasteurella* species; its recovery from a clinical sample should be interpreted with the source and evidence of infection.

## Pasteurella Canis

*Pasteurella canis* is a bacterium found in dogs and cats that can cause wound or soft-tissue infection after an animal bite.

---

## Pasteurellaceae

Pasteurellaceae is a family of bacteria that includes *Pasteurella* and related animal-associated organisms capable of causing bite-wound and respiratory infections.

## Pasteurella Multocida

A bacterium commonly carried by animals that can cause wound infections after bites or scratches and, less often, respiratory or bloodstream infection.

## Pasteurellosis

Infection caused by *Pasteurella* bacteria, commonly introduced through animal bites or scratches and causing cellulitis, abscesses, or occasionally invasive disease.

## Pasteurization

A heat treatment reducing microbial load in liquids by heating below boiling point to inactivate vegetative pathogens and spoilage organisms while preserving quality.

## Pastia Lines

Darker red lines that appear in skin creases during some rashes, especially scarlet fever. They support the diagnosis when other signs fit but are not specific by themselves.

### Alternate Names:

Pastia's Sign; Pastia Lines of Scarlet Fever

## Patau's Syndrome

A congenital disorder caused by an extra copy of chromosome 13. It can cause severe developmental problems, brain and heart defects, cleft lip or palate, and other organ abnormalities.

## Patau Syndrome

A genetic condition caused by an extra copy of chromosome 13. It can cause severe problems involving brain development, growth, the eyes, heart, kidneys, hands or feet, and the lip or palate. Many affected pregnancies end before birth, and surviving infants need complex specialist care.

## Patch

A flat, circumscribed area of altered skin color larger than approximately 1 cm. It is the larger counterpart of a macule and may be caused by pigmentary, vascular, inflammatory, or neoplastic disease.

## Patches

Patches are flat, distinct areas of altered skin color larger than a macule; their appearance, distribution, and associated symptoms help identify the cause.

## Patch Graft

A piece of natural or synthetic material sewn onto a blood vessel to widen or repair it during patch angioplasty. Materials may include a patient's tissue, bovine pericardium, or synthetic PTFE.

## Patch Test

A skin test used to identify delayed allergic contact reactions to substances applied under adhesive patches.

## Patch Testing

Patch testing identifies allergens causing allergic contact dermatitis. The standard series includes 35 allergens. Additional custom trays include occupational and personal allergens. Patches are applied to the upper back for 48 hours and removed. Reading at 48 hours (initial) and 72 to 96 hours (delayed) identifies positive reactions.

## Patchy Skin Color

Areas of skin that are lighter, darker, redder, or otherwise different in color from the surrounding skin. Causes include vitiligo, birthmarks, inflammation, infection, sun exposure, injury, and changes in blood flow; new or changing patches should be assessed.

## Patella

The largest sesamoid bone in the human body, embedded within the quadriceps tendon at the front of the knee. It improves the mechanical efficiency of knee extension, protects the front of the joint, and distributes forces across the femoral trochlea.

---

**Alternate Names:**

Rotula; Knee Cap

**Patellar Apprehension Test**

A knee examination used to look for kneecap instability. The examiner gently moves the kneecap outward while the knee is slightly bent. Fear, pain, or a muscle tightening response may support instability, but the test is interpreted with the history and other findings.

**Alternate Names:**

Fairbank Apprehension Test

**Patellar Bulge Test**

A physical examination manoeuvre used to detect small intra-articular effusions in the knee joint. The examiner sweeps the medial aspect of the knee upward to displace fluid, then sweeps downward on the suprapatellar and lateral aspects; a visible fluid bulge or wave returning to the medial hollow indicates a small effusion.

**Alternate Names:**

Brush Test; Stroke Test; Sweep Test

**Patellar Dislocation**

Complete displacement of the kneecap from the femoral groove, usually laterally, causing acute pain, deformity, swelling, and difficulty extending the knee.

**Patellar Reflex**

A monosynaptic deep tendon stretch reflex elicited by sharply tapping the patellar tendon beneath the patella with a reflex hammer, producing involuntary quadriceps contraction and knee extension; tests the integrity of the L3-L4 spinal nerve roots and femoral nerve.

**Alternate Names:**

Knee-Jerk Reflex

**Patellar Tap Test**

A clinical examination technique used to detect moderate to large knee joint effusions. The examiner compresses the suprapatellar pouch with one hand and pushes the patella downward against the femoral condyles with the other; a floating sensation

followed by a distinct tap or click against the femur confirms fluid accumulation.

**Alternate Names:**

Patellar Ballottement Test

**Patellar Tendinitis**

Alternate name; see *Patellar tendinitis*.

**Patellar Tendinopathy**

An overuse mechanical tendinopathy of the patellar tendon near its origin at the inferior pole of the patella, common in jumping athletes, presenting with localized anterior knee pain exacerbated by jumping, squatting, and descending stairs.

**Alternate Names:**

Jumper Knee

**Patellofemoral Instability**

A tendency of the patella to move abnormally or dislocate from the femoral trochlea. Risk factors include abnormal trochlear shape, a high-riding patella, ligament insufficiency, altered limb alignment, and previous dislocation.

**Patellofemoral Osteoarthritis**

Degenerative disease affecting the cartilage and bone surfaces between the kneecap and the femoral groove. It commonly causes pain at the front of the knee with stairs, squatting, kneeling, or prolonged sitting.

**Patellofemoral Pain Syndrome**

Pain around or behind the kneecap, often worsened by running, stairs, squatting, or sitting with the knee bent. It commonly results from repeated loading, muscle weakness, movement changes, or kneecap tracking problems.

**Alternate Names:**

Runner's Knee; Chondromalacia Patellae

**Patellofemoral Syndrome**

Pain around or behind the kneecap related to movement and loading of the patellofemoral joint, often without a major structural injury.

---

## Patent

Patent means open, unobstructed, or functioning, as in a patent airway, vessel, duct, or congenital fetal channel.

## Patent Ductus Arteriosus

A blood-vessel connection between the heart's main lung artery and the aorta that remains open after birth instead of closing. It can send extra blood to the lungs and strain the heart; effects range from none to heart failure.

## Patent Foramen Ovale

A persistent flap-like opening between the atria that normally permits fetal blood flow and usually closes after birth. It is often an incidental finding but may allow a clot to pass from the venous to the arterial circulation in selected circumstances.

## Patent Urachus

A patent urachus is persistence of the fetal channel between the bladder and umbilicus after birth, allowing urine to drain from the umbilicus and predisposing to infection.

## Patent Urachus Repair

Surgery to close an abnormal open connection between the bladder and the navel that remains from fetal development. It may stop urine leaking from the umbilicus and prevent infection; treatment depends on the child's anatomy and health.

## Paternal Depression

Depression occurring in a father during pregnancy or after a child is born. Symptoms may include low mood, irritability, withdrawal, sleep or appetite changes, and loss of interest; treatment can help both parent and family.

## Paternal Postnatal Depression

Depression in a father during the first year after a baby's birth. It may be linked to stress, sleep loss, previous depression, or a partner's depression and deserves professional assessment.

## Paternal Uniparental Disomy 14

A rare genetic condition resulting from inheritance of both copies of chromosome 14 from the father, leading to dysregulated expression of imprinted genes on chromosome 14. It presents with severe thoracic hypoplasia, pulmonary hypoplasia, and characteristic facial features. It is a lethal condition in most cases.

### Alternate Names:

Kagami-Ogata Syndrome; UPD(14)pat

## Paternity Testing

A DNA test that assesses whether a person is the biological parent of a child. Samples are usually collected from the child and possible parent, and the result reports either exclusion or a calculated probability of biological relationship.

## Pathogen

An organism or infectious agent that can enter the body and cause disease. Examples include viruses, bacteria, fungi, and parasites.

## Pathogenesis

The way a disease begins and develops, including its causes, changes in cells or tissues, and progression over time. Understanding pathogenesis helps explain symptoms, complications, and possible treatment targets.

## Pathogenic

Pathogenic means capable of causing disease in a susceptible host; pathogenicity depends on the organism or factor, dose, route, and host defenses.

## Pathogenicity

The capacity of a microorganism to cause disease in a susceptible host. Pathogenicity is a qualitative property, whereas virulence describes the relative degree of disease-causing capacity among strains or organisms.

## Pathogenic Variant

A change in DNA with strong evidence that it causes or contributes to a disease. Its importance depends

---

on the gene, the person's symptoms and family history, and whether one or both copies of the gene are affected.

### **Pathognomonic**

Describes a finding considered highly specific for a particular disease. Truly pathognomonic findings are uncommon; most findings must still be interpreted with the history, examination, and other test results.

### **Pathologic**

Pathologic means related to disease or an abnormal structural or functional change, rather than normal anatomy or physiology.

### **Pathological Fracture**

See *Pathologic Fracture*.

### **Pathological Laughter And Crying**

Alternate name; see *Pseudobulbar affect*.

### **Pathological Stealing**

Alternate name; see *Kleptomania*.

### **Pathologic Dilatation**

Abnormal widening of a hollow organ, duct, blood vessel, or other structure beyond its expected size because of obstruction, weakness, pressure, or disease.

### **Pathologic Fracture**

A break in bone weakened by a disease, such as osteoporosis, cancer, infection, or another bone disorder. It can happen after a minor injury or sometimes during ordinary activity; treatment addresses both the fracture and its cause.

### **Pathologic Hypertrophy**

Enlargement of an organ because its cells become larger, usually after long-term extra workload or hormone stimulation. It can help at first but may eventually impair function, as in an overworked heart.

### **Pathologic Nipple Discharge**

Nipple fluid that occurs without squeezing, comes from one breast or one duct, or is bloody or clear. It may have a benign cause, but it needs clinical assessment and often imaging to decide whether further testing is needed.

### **Pathologic Ossification**

Abnormal formation of bone in soft tissue or in a location where bone should not develop, as in heterotopic ossification after trauma, neurologic injury, or surgery.

### **Pathologic Process**

A pathologic process is an abnormal biological change that produces disease, such as infection, inflammation, ischemia, degeneration, abnormal growth, or immune injury.

### **Pathologic Response**

A pathologic response is an abnormal reaction of cells, tissues, or organs to injury, infection, immune activation, toxins, or other stressors that contributes to disease.

### **Pathologist**

A pathologist is a physician who diagnoses disease by examining tissues, cells, blood, or body fluids and may guide laboratory testing, cancer classification, and treatment decisions.

### **Pathology**

Study of disease. Includes general pathology (cell injury, inflammation, neoplasia) and systemic pathology (organ-specific diseases).

### **Pathology Artifact**

A change introduced while collecting, preserving, cutting, staining, or imaging a sample rather than by disease. Recognizing these false changes prevents an incorrect diagnosis.

### **Pathophysiology**

The study of the functional changes produced by disease or injury and the mechanisms through which



---

they cause signs and symptoms.

### **Patient**

A person receiving medical care, advice, examination, or treatment. Good care includes listening to the patient's concerns, explaining choices clearly, respecting informed decisions, protecting privacy, and reducing avoidable harm.

### **Patient Advocacy**

Acting on behalf of a patient to ensure that the patient's needs, rights, and preferences are addressed in healthcare.

### **Patient Autonomy**

A patient's right and capacity to make informed voluntary decisions about healthcare after receiving understandable information about options, benefits, risks, and alternatives.

### **Patient-Centered Care**

Healthcare planned around a person's needs, values, preferences, and goals. It includes clear communication, shared decisions, respect for cultural and personal choices, coordination between professionals, and support for the person and family.

### **Patient-Centered Outcome**

An outcome that reflects what matters to the patient, including symptoms, function, quality of life, treatment burden, and ability to achieve personal goals, rather than relying only on laboratory or imaging measures.

### **Patient-Controlled Analgesia**

A method in which a patient self-administers prescribed doses of analgesic medication using a programmable pump with preset safety limits. The medication, dose, and lockout interval are individualized according to the patient's condition and monitoring needs.

### **Patient Education**

The process of explaining a health condition, treatment, medicines, exercises, equipment, warning

signs, and prevention strategies in a way the patient can understand and use.

### **Patient Observation**

Systematic monitoring of a patient's symptoms, vital signs, examination findings, behavior, and response to treatment to detect deterioration or guide management.

### **Patient Positioning**

Placing and supporting a patient in a position that permits safe care, accurate examination or imaging, and adequate comfort.

### **Patient Preparation**

The instructions and assessments completed before a test or procedure, such as fasting, medicine review, hydration, consent, or skin preparation when required.

### **Patient-Reported Outcome Measure**

A validated questionnaire completed directly by a patient to assess symptoms, function, health status, or quality of life without clinician interpretation of the response.

### **Patient Rights**

Ethical and legal protections for people receiving care, including informed consent, privacy, access to information, respectful treatment, participation in decisions, and refusal of treatment where lawful.

### **Patient Safety**

Preventing avoidable harm during healthcare. It includes accurate identification, safe prescribing and medicine use, infection prevention, clear communication, fall prevention, and checking procedures and results.

### **Patlak-Rutland Analysis**

A graph used with some nuclear-medicine scans to estimate how quickly a tracked substance moves from the blood into tissue and stays there. It can help measure tissue activity, including activity in some tumors.

---

### Patrick Test

A physical examination maneuver (FABER test: Flexion, ABduction, External Rotation) used to evaluate sacroiliac joint dysfunction or hip joint pathology; with the patient supine, the foot of the tested leg is placed on the opposite knee in a 'figure-4' position while the examiner gently presses downward on the tested knee; anterior groin pain indicates hip pathology, whereas posterior pain localizes to the sacroiliac joint.

#### Alternate Names:

FABER Test; Patrick's FABER Maneuver

### Pattern Recognition Receptors

Proteins on or inside immune cells that detect common features of germs or damaged tissue. Their activation helps start the body's early immune response.

### Pattern-Recognition Receptors

Sensors on or inside immune cells that detect common features of germs or damaged cells. Their activation starts early immune responses, including inflammation and signals that help coordinate protection and tissue repair.

### Pauci-Immune Glomerulonephritis

A necrotizing and crescentic glomerulonephritis characterized on renal biopsy by focal glomerular capillary necrosis and cellular crescent formation with minimal or absent deposition of immunoglobulins or complement on immunofluorescence, strongly associated with antineutrophil cytoplasmic antibodies (ANCA-associated vasculitis).

#### Alternate Names:

ANCA-Associated Glomerulonephritis; Pauci-Immune Crescentic GN

### Paxillin

Paxillin is a focal-adhesion adaptor protein that links integrin signaling to the actin cytoskeleton and helps regulate cell adhesion, migration, and mechanical sensing.

### PA X-Ray

A posteroanterior X-ray is taken with the beam

traveling from the back toward the front, commonly for a chest image. This position reduces enlargement of the heart's outline compared with some other views.

### Pazopanib

An oral tyrosine-kinase inhibitor targeting vascular endothelial growth factor and other receptors, used for selected advanced renal-cell carcinoma and soft-tissue sarcoma.

### PBA

Alternate name; see *Pseudobulbar affect*.

### PBG Urine Test

A urine test measuring porphobilinogen, used urgently to evaluate suspected acute hepatic porphyria during compatible neurovisceral or abdominal attacks.

### PCBs

PCBs, or polychlorinated biphenyls, are persistent industrial chemicals that accumulate in tissues and can affect the nervous, endocrine, immune, and reproductive systems.

### PCL Injuries

Sprains or tears of the posterior cruciate ligament, a strong ligament inside the knee. They may follow a blow to the shin or forceful bending and can cause pain, swelling, stiffness, or a feeling that the knee gives way.

### PCO

PCO may mean polycystic ovaries or posterior capsular opacification; the intended condition depends on the clinical context.

### PCO Disease

Polycystic ovary syndrome, a hormonal and metabolic disorder involving irregular ovulation, androgen excess, and sometimes polycystic-appearing ovaries.

---

## PCOS

A common hormone disorder that can cause irregular ovulation or periods, higher levels of androgen hormones, acne, or excess facial hair. Some people have ovaries with many small follicles, but that finding is not required for diagnosis.

## PCP

Alternate name; see *Pneumocystis Jirovecii Pneumonia*.

## PCP and Ketamine Toxicity

Dissociative-drug toxicity causing agitation, hallucinations, impaired coordination, nystagmus, analgesia, tachycardia, hypertension, hyperthermia, seizures, or coma. PCP often produces severe behavioral disturbance, whereas ketamine can cause profound sedation; reduce stimulation, protect the airway, use benzodiazepines when needed, and treat hyperthermia or complications.

## PDA

Alternate name; see *Patent ductus arteriosus*.

## PDE5 Inhibitors

Medicines that improve blood flow into the penis by relaxing blood vessels and are mainly used for erectile dysfunction. Combining them with nitrate medicines can cause a dangerous fall in blood pressure.

## Peak Expiratory Flow Rate

The fastest speed at which a person can blow air out after taking a full breath. A handheld meter measures it in litres per minute; lower or changing values can help monitor airway narrowing, especially in asthma.

## Peak Flow Meter

A handheld device that measures peak expiratory flow rate, the fastest speed of air expelled after a full breath, usually reported in liters per minute. It provides a simple measure of large-airway flow but does not measure the full range of airflow and lung volumes assessed by spirometry.

## Peak Plasma Concentration

The highest measured concentration of a medicine in the blood after a dose. It is often called C<sub>max</sub> and is influenced by the dose, absorption speed, formulation, and route of administration.

## PE Anticoagulation

Anticoagulant treatment for pulmonary embolism to prevent clot extension and recurrence. Options include direct oral anticoagulants, heparins, and vitamin K antagonists; the choice and duration depend on bleeding risk, pregnancy, kidney function, cancer, provoking factors, and other clinical features.

## Peanut Allergy

An immune reaction to peanut proteins that can cause symptoms ranging from hives to life-threatening anaphylaxis.

### Alternate Names:

Peanut Hypersensitivity; Peanut Anaphylaxis

## Peanut Oil

Peanut oil is oil extracted from peanuts; refined oil usually contains little allergenic protein, whereas unrefined products may trigger reactions in susceptible people.

## Pearly Penile Papules

Small, benign, dome-shaped papules arranged circumferentially around the glans penis, representing normal anatomical variants requiring no treatment.

## Peau D'Orange

Peau d'orange is dimpled skin resembling an orange peel, caused by edema tethered around hair follicles; it can occur with breast cancer, infection, or lymphatic obstruction.

## Pectinate Line

A boundary inside the anal canal where the lining, nerve supply, blood vessels, and lymph drainage change. Conditions above and below this line may cause different symptoms and may be treated differently.

---

## **Pectoral**

Relating to the chest muscles or the front of the chest. The pectoral muscles move the arm and help stabilize the shoulder. Pain in this area may come from muscle strain, the ribs, breast tissue, or organs inside the chest.

## **Pectoral Girdle**

The pectoral girdle consists of the clavicles and scapulae, attaching the upper limbs to the trunk and providing broad surfaces for shoulder and arm movement.

## **Pectoralis**

Relating to the chest muscles, especially pectoralis major and pectoralis minor. These muscles help move the arm, support the shoulder, and connect the chest wall with the upper limb.

## **Pectoralis Major Muscle**

The large chest muscle that moves the arm across the body, turns it inward, and helps with pushing. It attaches to the collarbone, breastbone, ribs, and upper arm.

## **Pectoralis Minor Muscle**

A small chest muscle beneath the larger pectoralis major. It helps draw the shoulder blade forward and downward, hold it against the ribs, and assist forceful breathing; tightness can contribute to rounded shoulders.

## **Pectoralis Muscles**

The pectoralis major and minor muscles of the chest, which move and stabilize the shoulder and assist with movements such as pushing, adduction, and breathing.

## **Pectoriloquy**

Abnormally clear transmission of spoken sounds through the chest wall, often suggesting lung consolidation or another change in lung tissue.

## **Pectus Carinatum**

A chest-wall shape present from childhood in which

the breastbone pushes outward, sometimes called pigeon chest. It may cause no health problems, but some people have chest discomfort, reduced exercise tolerance, or emotional distress.

## **Pectus Excavatum**

A chest-wall shape present from childhood in which the breastbone is pushed inward, sometimes called funnel chest. It may cause no symptoms, but severe cases can affect exercise, breathing, posture, or emotional wellbeing.

### **Alternate Names:**

Sunken Chest; Funnel Chest

## **Pectus Excavatum Repair**

Surgical correction of a sunken sternum and adjacent costal cartilage, performed when chest-wall deformity causes significant symptoms, physiologic impairment, or psychosocial impact.

## **Pediatric Advanced Life Support**

Emergency training and guidance for recognizing and treating life-threatening breathing, circulation, or heart problems in infants and children. It covers resuscitation, airway support, shock, abnormal rhythms, and care after recovery.

### **Alternate Names:**

PALS; Pediatric Resuscitation Protocol

## **Pediatric AIDS**

Pediatric AIDS is advanced HIV infection in a child, with severe immune suppression or AIDS-defining illness; antiretroviral therapy and prevention of opportunistic infection are essential.

## **Pediatric Anesthesia**

Anesthesia for infants and children uses age- and weight-appropriate medicines, equipment, and monitoring. Children's airways and breathing reserves differ from adults, so they can develop low oxygen levels or changes in blood pressure quickly and need specialized care.

## **Pediatric Antireflux Surgery**

Surgical reconstruction of the gastroesophageal junc-

---

tion—most commonly via laparoscopic Nissen fundoplication—in infants and children with medically refractory gastroesophageal reflux disease, recurrent aspiration, or severe failure to thrive.

**Alternate Names:**

Pediatric Fundoplication; Pediatric Anti-Reflux Surgery; Nissen Fundoplication in Children

**Pediatric Apnea**

A pause or cessation of breathing in an infant or child. Causes include prematurity, infection, airway obstruction, seizures, metabolic disease, medication exposure, and central or obstructive sleep disorders.

**Pediatric Appendicitis**

Acute inflammation of the vermiform appendix, usually caused by luminal obstruction. Children may have evolving periumbilical-to-right-lower-quadrant pain, anorexia, vomiting, fever, and guarding, but presentation can be atypical in young children; timely surgical and antimicrobial assessment reduces perforation risk.

**Pediatric Assessment Triangle**

A rapid first look at a sick child that does not require touching the child. Clinicians observe appearance, effort of breathing, and skin color or circulation. The findings show how urgently help is needed and guide the next examination and treatment.

**Alternate Names:**

PAT; Pediatric Visual Triage Triangle

**Pediatric Brain Tumor**

A primary neoplasm originating within the parenchyma of the brain or surrounding meninges and cranial nerves during childhood or adolescence. Common histologic types include pilocytic astrocytoma, medulloblastoma, ependymoma, and diffuse intrinsic pontine glioma.

**Alternate Names:**

Pediatric Brain Tumor; Childhood Brain Tumor; Pediatric CNS Neoplasm

**Pediatric Brain Tumors**

Tumors that develop in a child's brain or nearby tissues. They may be noncancerous or cancerous

and can cause headache, vomiting, seizures, vision changes, weakness, or changes in behavior; treatment depends on the type, location, size, and age of the child.

**Pediatric Cancer**

Cancer occurring in infants, children, or adolescents. Common types include leukemia, brain tumors, lymphoma, neuroblastoma, and Wilms tumor; prognosis varies by type and stage.

**Pediatric Cardiopulmonary Resuscitation**

Basic life support resuscitation algorithm for pediatric patients aged 1 year to the onset of puberty, employing compressions to a depth of approximately 2 inches (5 cm) using a one- or two-handed technique at a 30:2 or 15:2 compression-to-ventilation ratio.

**Alternate Names:**

Pediatric CPR; Child Basic Life Support; BLS Pediatric

**Pediatric Cataract**

Clouding of the natural lens in an infant or child. It may be present at birth or develop after birth and can interfere with visual development, causing amblyopia or reduced vision.

**Pediatric Central Venous Line**

A small-caliber central vascular access catheter specifically designed and placed in neonates and pediatric patients for total parenteral nutrition, critical drug infusions, hemodialysis, or advanced hemodynamic monitoring.

**Alternate Names:**

Neonatal Central Line; Pediatric CVC; Infant Venous Catheter

**Pediatric Crutch Transfers**

Physical therapy instruction for pediatric patients utilizing axillary crutches to safely transition between seated and standing positions by grasping both crutches on the injured side and supporting weight on the unaffected extremity.

**Alternate Names:**

Crutch Chair Transfers in Children; Crutch Trans-

---

fers

### **Pediatric Crutch Gait Training and Fitting**

Clinical fitting and rehabilitation gait training for children utilizing axillary or forearm crutches, ensuring proper axillary clearance (2 – 3 finger-breadths) and handgrip alignment to prevent compressive crutch palsy.

#### **Alternate Names:**

Pediatric Crutch Fitting; Pediatric Gait Training; Axillary Crutch Training in Children; Crutch Gait Training; Crutch Fitting

### **Pediatric Crutch Stair Training**

Physical therapy gait training for pediatric crutch users navigating stairs safely (advancing the unaffected extremity first when ascending, and advancing crutches with the affected extremity when descending).

#### **Alternate Names:**

Stair Ambulation with Crutches; Crutch Stair Training

### **Pediatric Dehydration**

A deficit of body water and electrolytes in a child, commonly caused by vomiting, diarrhea, fever, poor intake, or excessive losses. Signs include reduced urination, dry mucosa, lethargy, tachycardia, and altered perfusion.

### **Pediatric Dentistry**

Dental care for infants, children, and adolescents, including prevention and treatment of tooth decay, care of developing teeth, growth-related problems, behavior support, and adaptations for children with special healthcare needs.

### **Pediatric Dermatology**

The medical specialty concerned with skin, hair, and nail disorders in infants, children, and adolescents. It includes congenital lesions, inflammatory diseases, infections, vascular lesions, and tumors.

### **Pediatric Dermatology Conditions**

Skin conditions in children may include birthmarks, eczema, infections, vascular growths, pigment changes, and inherited disorders. Most are manageable, but a rapidly changing, painful, bleeding, or infected lesion needs medical assessment.

### **Pediatric Diabetic Ketoacidosis**

A life-threatening complication of diabetes in which very high blood sugar and a lack of insulin make the blood acidic. A child may urinate and drink excessively, vomit, have abdominal pain, deep breathing, fruity-smelling breath, sleepiness, or coma.

### **Pediatric Early Warning Score**

A hospital scoring system that combines a child's behavior, breathing, heart rate, circulation, and need for oxygen to detect worsening illness early. A rising score prompts closer observation or urgent review; it supports, but does not replace, clinical judgment.

#### **Alternate Names:**

PEWS; Pediatric Early Warning System

### **Pediatric Echocardiogram**

A noninvasive ultrasound examination of the heart in infants and children used to evaluate congenital structural heart defects, valve function, and cardiac blood flow.

#### **Alternate Names:**

Pediatric Echo; Congenital Echocardiography; Pediatric Cardiac Ultrasound

### **Pediatric Eczema**

Atopic dermatitis in children commonly begins in infancy with facial and extensor involvement; flexural disease predominates in older children. Pruritus, xerosis, sleep disturbance, and scratching are characteristic.

### **Pediatric Feeding Tube**

A small, flexible plastic tube inserted through the nose (nasogastric/nasojejunal) or directly into the stomach (gastrostomy) to deliver breast milk, formula, or medications to infants unable to feed adequately by mouth.



---

**Alternate Names:**

Infant Feeding Tube; Pediatric NG Tube; Infant Enteral Tube

**Pediatric Fever**

An elevated body temperature in a child, usually caused by infection but sometimes by inflammation, medicines, or environmental heat. Evaluation depends on age, appearance, duration, and associated findings.

**Pediatric Foreign Body Ingestion**

Swallowing of a nonfood object by a child, with risk determined by the object, location, size, symptoms, and time since ingestion. Batteries, magnets, sharp objects, and obstructing objects require urgent evaluation.

**Pediatric Gastroesophageal Reflux Disease**

Reflux that causes troublesome symptoms or complications in a child, such as feeding difficulty, pain, or poor growth. Care may include feeding adjustments or, in selected cases, thickened feeds, dietary changes, or medicine under clinical guidance. Infants should sleep flat on their backs; reflux is not a reason for side, stomach, or inclined sleep positioning.

**Alternate Names:**

Pediatric GERD; Childhood Acid Reflux; Pediatric Gastroesophageal Reflux

**Alternate Names:**

Pediatric GERD; Infant Acid Reflux

**Pediatric Glaucoma**

Glaucoma occurring in an infant or child, often involving abnormal development or blockage of the eye's drainage system. Increased eye pressure can enlarge or damage the eye and impair vision.

**Pediatric Growth Hormone Deficiency**

Inadequate secretion of growth hormone by the anterior pituitary gland in infants and children, leading to deceleration of growth velocity, proportionate short stature, delayed bone age, and abnormal body composition. Causes include pituitary hypoplasia,

genetic defects, midline cranial defects, intracranial tumors, radiation therapy, or idiopathic hypopituitarism.

**Alternate Names:**

Pediatric GH Deficiency; Childhood Hypopituitarism; Growth Hormone Deficiency in Children

**Alternate Names:**

Pediatric GH Deficiency; Pituitary Dwarfism

**Pediatric Health**

The physical, mental, emotional, and social well-being of infants, children, and adolescents. Care includes prevention, vaccinations, growth and development checks, treatment of illness, and support for the child and family.

**Pediatric Heart Surgery**

An operation to repair or improve a child's heart or major blood vessels, often for a congenital heart defect. Procedures may close abnormal openings, improve blood flow, or replace a valve; recovery and risks depend on the defect and operation.

**Pediatric Hepatitis A**

An acute viral infection of the liver caused by the hepatitis A virus (HAV) in children, transmitted predominantly through the fecal-oral route via contaminated food or water. While often self-limiting, supportive hydration is critical and routine pediatric vaccination provides durable protection.

**Alternate Names:**

Childhood Hepatitis A; Pediatric HAV Infection; Acute Viral Hepatitis A in Children

**Pediatric Hepatitis B**

A viral liver infection caused by the hepatitis B virus (HBV) in pediatric patients, most commonly acquired perinatally from an infected mother or during early childhood. Chronic carrier states are highest when acquired in infancy; universal infant vaccination and post-exposure immunoprophylaxis prevent transmission.

**Alternate Names:**

Childhood Hepatitis B; Perinatal HBV Infection; Chronic Pediatric Hepatitis B

---

## Pediatric Hepatitis C

Infection with the hepatitis C virus (HCV) in infants and children, primarily acquired vertically through perinatal maternal-fetal transmission or through parenteral exposure. Modern direct-acting antiviral (DAA) regimens provide high cure rates in pediatric cohorts.

### Alternate Names:

Childhood Hepatitis C; Pediatric HCV Infection; Perinatal Hepatitis C

## Pediatric Hypercholesterolemia

Elevated circulating total cholesterol or low-density lipoprotein (LDL) levels in children and adolescents. Causes include familial hypercholesterolemia, obesity, insulin resistance, endocrine disorders, or high-saturated-fat diets.

### Alternate Names:

Childhood Hyperlipidemia; Pediatric Dyslipidemia; Familial Hypercholesterolemia in Children

## Pediatric Hypertension

Blood pressure in children and adolescents persistently at or above the 95th percentile for age, sex, and height. Secondary etiologies (such as renal parenchymal disease or aortic coarctation) are common, especially in young children.

### Alternate Names:

Childhood Hypertension; Pediatric High Blood Pressure; Adolescent Hypertension

## Pediatrician

A medical doctor who specializes in the health and medical care of children from birth through adolescence (typically up to age 16-18).

### Alternate Names:

Paediatrician

## Pediatric Limp

An abnormal walking pattern or refusal to walk in a child. Causes range from minor injury and transient synovitis to infection, fracture, inflammatory disease, neuromuscular disease, or malignancy; fever, severe pain, swelling, or inability to bear weight requires prompt assessment.

## Pediatric Meningitis

Inflammation or infection of the coverings of a child's brain and spinal cord. Fever, headache, stiff neck, vomiting, light sensitivity, confusion, seizures, or a rash may occur; babies may be irritable, feed poorly, or have a bulging soft spot. It can progress quickly and is life-threatening.

## Pediatric Midline Catheters

A peripheral vascular access device (typically 3 to 8 cm in length) inserted into a peripheral vein in the upper extremity of infants and children, with the catheter tip terminating distal to the axillary line. Utilized for intermediate-duration intravenous fluid therapy and medication administration lasting 1 to 4 weeks.

### Alternate Names:

Pediatric Midline Venous Access; Infant Midline IV; Neonatal Midline Catheter

## Pediatric Myocarditis

Acute or chronic inflammation of the myocardium in infants and children, most commonly triggered by viral infections (enteroviruses, adenovirus, parvovirus B19, SARS-CoV-2) or post-infectious autoimmune responses. Manifests as tachypnea, poor feeding, tachycardia, cardiomegaly, arrhythmias, or acute congestive heart failure.

### Alternate Names:

Childhood Myocarditis; Pediatric Myocardial Inflammation; Acute Pediatric Carditis

## Pediatric Nasal Congestion

A stuffy or runny nose in a child is usually caused by a viral infection or allergy; breathing difficulty, dehydration, prolonged fever, or symptoms in an infant need prompt assessment.

## Pediatric Nursing

Pediatric nursing provides age-appropriate care to infants, children, and adolescents, accounting for growth, development, family involvement, communication, safety, and medication differences.

## Pediatric Obesity

Excess body fat in a child or adolescent, assessed by

---

comparing weight and height with other children of the same age and sex. It can affect breathing, joints, blood sugar, sleep, and emotional wellbeing; assessment should be supportive and non-stigmatizing.

### **Pediatric Obstructive Sleep Apnea**

Repeated narrowing or blockage of a child's throat during sleep, causing pauses in breathing, snoring, restless sleep, daytime behavior or attention problems, and sometimes poor growth. Enlarged tonsils or adenoids are common causes.

### **Pediatric Ophthalmology**

The subspecialty caring for eye and vision disorders in infants, children, and adolescents, including strabismus, amblyopia, congenital abnormalities, refractive error, and childhood eye disease.

### **Pediatric Orthopedics**

The branch of orthopedics focused on bone, joint, muscle, and movement problems in infants, children, and adolescents. It includes developmental conditions, limb differences, fractures, sports injuries, spine deformities, and growth-related disorders.

### **Pediatric Pancreatitis**

Acute or chronic inflammatory disease of the pancreas in infants, children, and adolescents. Common pediatric etiologies include abdominal blunt trauma, viral infections (mumps, Coxsackie, EBV), medications, systemic illness, gallstones, and genetic mutations (*PRSS1*, *SPINK1*, *CFTR*).

#### **Alternate Names:**

Childhood Pancreatitis; Pediatric Pancreatic Inflammation; Acute Pediatric Pancreatitis

### **Pediatric Pneumonia**

Infection or inflammation of the lung tissue in an infant, child, or adolescent. It may cause cough, fever, fast or difficult breathing, chest pain, or low oxygen and can be caused by bacteria, viruses, fungi, or aspiration.

### **Pediatric Poisoning**

Toxic exposure in an infant, child, or adolescent, most often from medicines, household products,

plants, gases, or recreational substances. Risk depends on the agent, dose per kilogram, route, formulation, age, and time since exposure; symptoms may include vomiting, altered consciousness, respiratory depression, seizures, or organ injury. Obtain poison-center guidance and do not induce vomiting unless specifically directed.

### **Pediatric Ptosis**

Congenital or acquired drooping of the upper eyelid in infants and children, most commonly caused by developmental dysgenesis of the levator palpebrae superioris muscle. Timely surgical correction is indicated to prevent deprivation amblyopia.

#### **Alternate Names:**

Congenital Ptosis; Pediatric Blepharoptosis; Childhood Drooping Eyelid

### **Pediatric Pump Gastrostomy Feeding**

Continuous or intermittent enteral nutrition delivered to pediatric patients through a gastrostomy tube using an electronic feeding pump. Provides precise, controlled flow rates and volumes for children who cannot tolerate bolus feeding.

#### **Alternate Names:**

Pediatric Enteral Pump Feeding; Pediatric G-Tube Pump Feeding

#### **Alternate Names:**

Pediatric G-Tube Pump Feeding; Continuous Enteral Infusion

### **Pediatric Rash Evaluation**

Systematic clinical assessment of acute cutaneous eruptions in infants and young children under two years of age. Etiologies encompass benign viral exanthems (roseola infantum, enteroviral hand-foot-mouth disease), common inflammatory dermatoses (atopic dermatitis, seborrheic dermatitis, diaper dermatitis), and life-threatening dermatologic emergencies (meningococcemia, Kawasaki disease).

#### **Alternate Names:**

Infant Rash Assessment; Toddler Dermatitis; Childhood Exanthem Evaluation

### **Pediatric Rehabilitation**

Therapy that helps children with conditions present

---

from birth, injuries, illness, or developmental disabilities gain skills and independence. Care considers growth, family goals, school participation, communication, movement, learning, and daily activities.

## **Pediatrics**

The branch of medicine concerned with the health and medical care of infants, children, and adolescents from birth to age 18 (or up to age 16 in the UK).

**Alternate Names:**

Paediatrics

## **Pediatric Sepsis**

Sepsis occurring in an infant, child, or adolescent, in which the body's response to infection causes organ dysfunction. Signs vary with age and may include abnormal temperature, breathing difficulty, unusual sleepiness, poor feeding, or reduced urine.

## **Pediatric Shock**

A dangerous condition in which a child's organs do not receive enough blood flow or oxygen. Causes include severe fluid loss, infection, heart failure, or a blockage. Early signs may include a fast heartbeat, pale or cold skin, unusual sleepiness, confusion, or little urine.

## **Pediatric Skin Tumors**

Skin growths in children include harmless birth-related or blood-vessel growths, cysts, and, rarely, cancer. A lump that grows quickly, bleeds, ulcerates, changes color, or causes pain should be assessed by a clinician.

## **Pediatric Sleep Apnea**

Pediatric sleep apnea is repeated interruption or reduction of breathing during sleep, commonly from enlarged tonsils or adenoids, causing snoring, restless sleep, behavior changes, or impaired growth.

## **Pediatric Total Parenteral Nutrition**

Intravenous nutrition for infants who cannot receive enough nutrients through the gastrointestinal tract; it requires careful monitoring of glucose, electrolytes, liver function, infection, and catheter complications.

## **Pediatric Toxicology**

The medical study and treatment of poisoning or harmful exposure in children. Common exposures include medicines, iron, household chemicals, and plants. Symptoms vary with the substance and may include vomiting, sleepiness, breathing problems, seizures, or burns; urgent poison-center or emergency advice may be needed.

## **Pediatric Volvulus**

Twisting of a loop of intestine around its mesentery, causing obstruction and potentially loss of blood supply. In children it is often related to intestinal malrotation and is a surgical emergency when acute.

## **Pedicle**

A pedicle is a stalk or attachment containing vessels, nerves, or other supporting structures, such as the base of a polyp, flap, or vertebral arch.

## **Pedicle Skin Graft**

A piece of skin moved to cover a wound while remaining attached to its original blood supply through a tissue bridge.

## **Pediculosis**

Infestation with lice, small insects that live on the scalp, body, or pubic hair. It causes itching and spreads mainly through close contact or shared personal items.

## **Pediculosis and Pthiriasis (Lice Infestation)**

Infestation of scalp, body, or pubic hair by lice. It causes itching, visible lice or attached eggs, and sometimes excoriations or secondary infection; treatment requires an effective pediculicide plus measures to limit reinfestation.

## **Pediculosis Capitis**

Infestation of the scalp by head lice, causing itching and attached eggs or lice in the hair.

## **Pediculus Humanus Capitis**

A blood-sucking parasite commonly known as the

---

louse (plural, lice) that can cause an itchy scalp; infestations are highly contagious and especially common in school-age children.

### **Pedigree**

A family diagram that shows relationships and who has a particular trait or disease. Doctors and genetic counselors use it to look for inheritance patterns and estimate the chance that relatives may be affected.

### **Pedodontics**

The branch of dentistry concerned with infants, children, and adolescents.

### **Pedophilia**

A continuing sexual interest in children who have not reached puberty. The term describes an interest and is not synonymous with pedophilic disorder; sexual contact with a child is abuse and cannot be consensual.

### **Pedophilic Disorder**

A mental-health disorder involving recurrent, intense sexual fantasies, urges, or behaviors toward children who have not reached puberty. Diagnostic criteria include clinically significant distress or impairment, or acting on the urges, together with required age and duration criteria.

### **Pedunculate**

Pedunculate describes a structure, lesion, or polyp attached by a narrow stalk rather than a broad base.

### **Pedunculated**

Attached to the skin or another surface by a narrow stalk, rather than having a broad base.

### **Peeling Skin**

Loss or shedding of the outer layer of skin, which may result from dryness, sunburn, irritation, allergy, infection, inflammation, or a systemic illness.

### **PEEP**

Pressure kept in the airways at the end of breathing

out, usually by a ventilator or breathing machine. It helps keep the lung air sacs open and improve oxygen levels, but too much pressure can injure the lungs or lower blood flow.

### **PEF**

PEF, or peak expiratory flow, is the fastest airflow produced during a forceful exhalation and is used chiefly to monitor variable airway obstruction such as asthma.

### **PEG Tube**

A flexible feeding tube placed through the abdominal wall into the stomach with camera guidance. It provides nutrition, fluids, or medicines when swallowing is unsafe or inadequate, and requires regular skin care.

### **Pel-Ebstein Fever**

A cyclical, recurring fever pattern observed in a subset of patients with Hodgkin lymphoma, characterized by regular episodes of high fever lasting 1 to 2 weeks alternating with afebrile intervals of similar duration.

#### **Alternate Names:**

Pel-Ebstein Pattern; Cyclical Lymphoma Fever

### **Peliosis Hepatis**

A usually uncommon condition in which multiple blood-filled spaces form within the liver. It may be linked to medicines, hormones, toxins, infections, cancer, or other diseases. It often causes no symptoms but can rarely cause liver failure or life-threatening bleeding if a space ruptures.

### **Pelizaeus-Merzbacher Disease**

A rare inherited disorder in which the brain does not make myelin, the protective covering around nerve fibers, normally. It usually begins in infancy with poor muscle tone, unusual eye movements, and delayed development, followed by stiffness, balance problems, and increasing movement difficulty. There is no cure; care supports movement, feeding, breathing, and communication.

#### **Alternate Names:**

PMD; PLP1-Related Hypomyelinating Leukodys-



---

trophy

### **Pellagra**

A disease caused by too little niacin, leading to skin changes, diarrhea, and problems with thinking.

### **Pellet**

A pellet is a small compacted mass of medication, material, or tissue; drug pellets may be formulated for implantation or controlled release.

### **Pelvic**

Relating to the pelvis, the bony ring and associated organs, muscles, vessels, and nerves in the lower trunk.

### **Pelvic Binder**

A circumferential device applied around the pelvis to reduce motion and hemorrhage from suspected unstable pelvic-ring injury. It should be centered over the greater trochanters and used with definitive imaging and hemorrhage-control management.

### **Pelvic Bone**

The ring of bones formed by the paired hip bones, sacrum, and coccyx that supports the trunk, protects pelvic organs, and transmits weight to the lower limbs.

#### **Alternate Names:**

Innominate Bone; Coxal Bone

### **Pelvic Cavity**

The space inside the lower part of the body enclosed by the pelvic bones. It contains the bladder, rectum, reproductive organs, blood vessels, nerves, and supporting muscles.

### **Pelvic Congestion**

Enlargement or pooling of blood in the veins of the pelvis, which may cause pelvic pressure, discomfort, or pain.

### **Pelvic Congestion Syndrome**

Chronic pelvic pain associated with enlarged or inefficient pelvic veins, often worsening with prolonged

standing, before menstruation, or after intercourse. Diagnosis requires evaluation for other causes of pelvic pain and may involve pelvic imaging; treatment is individualized by gynecology or vascular specialists.

### **Pelvic CT Scan**

Computed tomography of the pelvis using x-rays and computer reconstruction to evaluate bones, organs, soft tissues, masses, trauma, infection, and bleeding.

### **Pelvic Diaphragm**

The group of muscles and supporting tissue forming the floor of the pelvis. It supports the bladder, bowel, and reproductive organs and helps control urine and stool.

### **Pelvic Examination**

Clinical inspection and palpation of the external genitalia, vagina, cervix, uterus, and adnexa to evaluate bleeding, pain, discharge, lesions, pregnancy-related findings, or infection.

### **Pelvic Exenteration**

Very extensive surgery removing some pelvic organs and nearby tissues, usually for selected cancers that have returned or grown locally when less extensive treatment cannot control them. Recovery is prolonged and reconstruction may be needed.

### **Pelvic Floor**

A group of muscles and connective tissues supporting the pelvic organs and controlling urinary and fecal continence.

### **Pelvic Floor Dysfunction**

Weakness, overactivity, poor coordination, or injury of the muscles and connective tissues supporting the bladder, bowel, and reproductive organs. Symptoms may include urinary or fecal problems, pelvic pain, constipation, pain with sex, or pelvic organ prolapse; assessment may involve pelvic-health clinicians.



---

## Pelvic Floor Health And Disorders

The pelvic floor is a group of muscles and supporting tissues that hold the bladder, bowel, and reproductive organs in place and help control urine, stool, and sexual function. Weakness, tightness, injury, or poor coordination can cause pain, leakage, constipation, or prolapse.

### Pelvic Floor Muscles

Muscles at the bottom of the pelvis that support the bladder, bowel, and reproductive organs and help control urine, stool, and sexual function.

### Pelvic Floor Muscle Training Exercises

Pelvic floor muscle exercises strengthen the muscles supporting the bladder, bowel, and reproductive organs and may improve selected forms of urinary or fecal leakage.

### Pelvic Fracture

A break in one or more bones of the pelvic ring, often caused by major trauma but sometimes occurring in weakened bone. The injury may be stable or unstable and can be associated with bleeding or damage to nearby organs, nerves, or blood vessels.

### Pelvic Girdle

The ring of bones connecting the lower limbs to the trunk, consisting of the hip bones, sacrum, and associated joints.

### Pelvic Girdle Pain In Pregnancy

Pain arising from the pelvic joints and surrounding tissues during pregnancy, often affecting the pubic symphysis, sacroiliac joints, hips, or lower back. It may impair walking, turning in bed, or climbing stairs.

### Pelvic Infection

An infection involving pelvic organs or tissues, including pelvic inflammatory disease, urinary infection, abscess, or postoperative infection.

### Pelvic Inflammatory Disease

Infection and inflammation of the upper reproduc-

tive tract, including the uterus, fallopian tubes, and nearby tissues. It often follows an ascending infection: chlamydia and gonorrhea are important causes, but other genital-tract bacteria can also be involved. Symptoms vary and may include lower-abdominal pain, unusual discharge or bleeding, pain during sex, or tenderness on examination; fever may occur, but some cases cause mild symptoms or none.

### Pelvic Inflammatory Disease (PID)

Alternate name; see *Pelvic Inflammatory Disease*.

### Pelvic Inlet

The upper opening of the bony birth canal, formed by the lower spine, hip bones, and front pelvic joint. Its shape and size help determine how a baby can pass through the pelvis during childbirth.

### Pelvic Kidney

A kidney that remains in the pelvis rather than ascending to its usual lumbar position during development; it may be asymptomatic or associated with abnormal drainage, stones, infection, or vascular anatomy.

### Pelvic Laparoscopy

A minimally invasive procedure using a camera inserted through small abdominal incisions to inspect and treat pelvic organs and structures.

### Pelvic Lymphadenectomy

Pelvic lymphadenectomy is surgical removal of lymph nodes in the pelvis to check for or treat spread of cancer.

### Pelvic Mass

A lump or abnormal enlargement in the pelvis arising from reproductive, urinary, gastrointestinal, lymphatic, or other tissues.

### Pelvic Neoplasm

A pelvic neoplasm is an abnormal growth in the pelvis, arising from organs, bones, soft tissues, or lymph nodes; it may be benign or malignant.

---

## Pelvic Organ Prolapse

Dropping or bulging of the bladder, uterus, rectum, or nearby tissue into or toward the vagina because supporting muscles and tissues have weakened. It may cause pressure, a vaginal bulge, urinary or bowel problems, or discomfort during sex.

## Pelvic Organ Prolapse Quantification System

A standard examination used to measure how far the vagina, uterus, bladder, or rectum has moved downward. Measurements are taken while bearing down and describe the severity of prolapse to help plan treatment.

### Alternate Names:

POP-Q System; POP-Q Classification

## Pelvic Outlet

The lower opening of the bony pelvis and birth canal, bordered by the pubic bone, sitting bones, and tailbone. Its shape and size affect passage through the pelvis during childbirth.

## Pelvic Pain

Pain in the lower abdomen, pelvis, or area between the genitals and anus. Causes include reproductive, urinary, bowel, muscle, nerve, and other conditions. Sudden severe pain, pregnancy, fever, fainting, vomiting, or heavy bleeding needs urgent assessment.

## Pelvic Pain During Early Pregnancy

Pain in the pelvis or lower abdomen during early pregnancy, caused by normal uterine changes, miscarriage, ectopic pregnancy, ovarian disease, infection, or other conditions. Severe or one-sided pain, shoulder pain, fainting, fever, or vaginal bleeding requires emergency assessment.

## Pelvic Ring Fracture

A high-energy traumatic skeletal disruption of the pelvic ring structure (involving the sacrum, sacroiliac joints, and pubic rami), classified by Young-Burgess mechanism (lateral compression, antero-posterior compression, vertical shear); frequently complicated by life-threatening retroperitoneal venous plexus hemorrhage.

## Pelvic Spleen

A spleen located unusually low in the abdomen or pelvis because its supporting tissues are loose or absent. It may feel like a pelvic lump and can twist, lose its blood supply, or rupture, causing sudden pain and requiring urgent care.

## Pelvic Viscus

An organ located in the pelvic cavity, such as the bladder, rectum, uterus, ovaries, prostate, or related structures.

## Pelvimeter

An instrument used to estimate the size and shape of the pelvis, especially during obstetric assessment. Pelvic measurements are only one part of judging whether vaginal birth is likely and are interpreted with the clinical situation.

## Pelvimetry

Measurement of the size and shape of the pregnant person's pelvis, usually for selected childbirth concerns. Results are considered with the baby's position, size, labor progress, and overall clinical situation.

## Pelvis

The bony ring and associated soft tissues at the base of the trunk, connecting the spine with the lower limbs.

## Pelvis MRI Scan

Magnetic resonance imaging of the pelvis used to evaluate organs, muscles, bones, joints, nerves, blood vessels, and soft-tissue disease without ionizing radiation.

## Pelvis X-Ray

Plain radiographic imaging of the pelvic bones and hips used to evaluate fractures, alignment, arthritis, deformity, and selected bone lesions.

## Pemberton Sign

Facial redness or bluish discoloration, neck-vein swelling, or breathing difficulty that appears when

---

both arms are raised. It may suggest that an enlarged thyroid or another chest mass is pressing on major veins or the airway.

### **Pemberton'S Sign**

Facial redness or bluish discoloration, neck-vein swelling, or breathing difficulty that appears when both arms are raised. It may suggest that an enlarged thyroid or another chest mass is pressing on major veins or the airway.

### **Pemetrexed**

Pemetrexed is a folate-antimetabolite chemotherapy drug that inhibits several enzymes involved in nucleotide synthesis, used especially for mesothelioma and nonsquamous non-small-cell lung cancer; folic acid and vitamin B12 reduce toxicity.

### **Pemetrexed Disodium**

The disodium salt of pemetrexed, an antifolate chemotherapy medicine used with other treatment for selected non-small-cell lung cancer and mesothelioma.

### **Pemoline**

Pemoline is a stimulant formerly used for attention-deficit disorders; serious liver toxicity led to withdrawal or severe restriction in many countries.

### **Pemphigoid**

A group of immune-system diseases that cause firm, often itchy blisters when antibodies attack the deeper layers of skin or mucous membranes. The type, location, age, and associated illness determine treatment.

### **Pemphigoid Gestationis**

An autoimmune blistering disease of pregnancy characterised by intensely pruritic urticarial plaques and tense vesicles or bullae, typically arising in the second or third trimester around the umbilicus and spreading to the trunk and extremities.

### **Pemphigus**

A group of autoimmune diseases causing fragile

blisters when antibodies attack connections between skin cells.

### **Pemphigus Erythematosus**

A localized superficial pemphigus variant with fragile blisters, erosions, and crusted plaques, often in seborrheic areas such as the face, scalp, and upper chest. Autoantibodies disrupt epidermal cell adhesion; diagnosis generally requires biopsy and immunofluorescence.

### **Pemphigus Foliaceus**

An autoimmune blistering skin disease caused by antibodies against proteins that hold skin cells together. It causes superficial fragile blisters and crusted sores but usually does not affect mucous membranes.

#### **Alternate Names:**

Brazilian Pemphigus Foliaceus; Endemic Pemphigus Foliaceus; Fogo Selvagem; South American Pemphigus

### **Pemphigus Vulgaris**

A rare autoimmune blistering disease that often begins with painful sores in the mouth and can also cause fragile blisters on the skin or other moist linings. The blisters break easily, leaving raw areas that can lead to infection, fluid loss, or difficulty eating and drinking.

### **Penbutolol**

Penbutolol is a beta-blocker formerly used for high blood pressure; it can slow the heart rate and worsen asthma in susceptible people.

### **Pencil Eraser Swallowing**

Swallowing a small pencil eraser usually causes little poisoning risk, but it may lodge in the airway or digestive tract. Choking, breathing difficulty, persistent vomiting, or severe pain requires emergency care.

### **Pencil Swallowing**

Swallowing a pencil or part of one can injure the mouth, throat, or digestive tract and may block the airway. Do not force food or drink; breathing

---

trouble, bleeding, or severe pain needs emergency care.

### **Pendred Syndrome**

An autosomal recessive genetic disorder caused by biallelic mutations in the SLC26A4 gene encoding the anion exchanger pendrin, characterized by congenital bilateral sensorineural hearing loss, temporal bone malformations (enlarged vestibular aqueduct), and euthyroid or hypothyroid multinodular goiter.

#### **Alternate Names:**

Pendred's Syndrome; Goiter and Sensorineural Deafness Syndrome

### **Pendular**

Moving or swinging freely like a pendulum; pendular nystagmus has a similar speed in both directions.

### **Pendular Nystagmus**

Involuntary, rhythmic, sinusoidal oscillations of the eyes in which the movements in both directions occur at equal velocity without distinct fast or slow phases. It most commonly arises from congenital visual deprivation, optic nerve abnormalities, or demyelinating brainstem diseases.

### **Pendulous**

Pendulous means hanging loosely from a supporting structure, as in a pedunculated lesion, drooping tissue, or an organ suspended by a stalk.

### **Penetrance**

The proportion of people with a particular gene change who develop the related trait or condition. Penetrance may be complete, incomplete, or influenced by other factors.

### **Penetrating Abdominal Trauma**

Injury to the abdomen caused by a projectile, sharp object, or other penetrating force. It may damage abdominal organs or blood vessels and requires urgent assessment for internal bleeding or organ injury; management depends on the mechanism, examination, stability, and imaging findings.

### **Penetrating Wounds**

Injuries caused when an object enters the skin and damages underlying tissue. They may injure blood vessels, nerves, organs, or bones and require assessment for bleeding, contamination, retained objects, and infection.

### **Penicillin**

A family of antibiotics that kill bacteria by stopping them from making cell walls. Different types treat different bacterial infections; they do not work against viruses. Allergy, resistance, and kidney problems can affect the choice of medicine.

### **Penicillin Allergy**

An immune reaction to penicillin antibiotics that may cause rash, swelling, breathing difficulty, or anaphylaxis.

#### **Alternate Names:**

Penicillin Hypersensitivity; Beta-Lactam Allergy

### **Penicillin G**

Penicillin G is an injectable penicillin antibiotic used for infections such as syphilis and selected streptococcal or other susceptible bacterial infections.

### **Penicillin V**

Penicillin V is an oral penicillin antibiotic used for selected susceptible infections, including some throat and dental infections.

### **Penicillium**

A genus of molds that includes species used to make medicines and foods and species that can cause allergy or opportunistic infection.

### **Penile**

Penile means relating to the penis, the external male organ involved in urination and sexual reproduction.

### **Penile Adhesions**

Abnormal attachment of penile skin to the glans, often after circumcision or during childhood. They may cause irritation or conceal part of the glans and are assessed and treated according to severity.

---

## Penile Augmentation

A cosmetic or reconstructive procedure intended to increase penile length, girth, or both. Techniques and results vary, and complications may include scarring, deformity, infection, altered sensation, or erectile problems.

## Penile Cancer

An uncommon cancer of the penis, usually arising from surface skin cells. It may appear as a persistent sore, lump, bleeding area, or skin change; HPV, smoking, foreskin problems, and chronic inflammation can increase risk.

## Penile Disease

A disorder affecting the penis, including infection, inflammation, erectile or structural disease, trauma, skin disease, or malignancy.

## Penile Edema

Swelling of the penis caused by inflammation, infection, trauma, allergic reaction, impaired lymphatic drainage, or a constricting ring or retracted foreskin. Sudden painful swelling with an irreducible foreskin is an emergency.

## Penile Erection

Stiffening and enlargement of the penis when blood fills erectile tissue and is temporarily held there. It can occur with sexual or other stimulation; persistent difficulty getting or keeping an erection may be erectile dysfunction.

## Penile Fracture

A tear in the tough covering of an erect penis, usually after forceful bending. A popping sound, immediate loss of erection, swelling, and bruising may occur; urgent surgery can prevent lasting problems.

## Penile Hygiene

Routine cleaning and drying of the penis and foreskin, with gentle retraction only when it moves freely and replacement afterward. Good hygiene helps reduce irritation and infection without harsh soaps or forceful retraction.

## Penile Implant

A surgically implanted device used to produce penile rigidity in selected people with erectile dysfunction that has not responded to other treatments.

## Penile Infection

Infection of the penis or foreskin caused by bacteria, fungi, viruses, or parasites, producing pain, redness, discharge, sores, or swelling.

## Penile Intraepithelial Neoplasia

A precancerous change in the surface cells of the penis. It may appear as a persistent red, white, scaly, or wart-like patch on the glans or foreskin and may be linked to HPV or chronic inflammation. Examination, biopsy, treatment, and follow-up reduce the risk of invasive cancer.

### Alternate Names:

PeIN; Erythroplasia of Queyrat; Bowen Disease of the Penis

## Penile Intraepithelial Neoplasia (PeIN or PIN)

Alternate name; see *Penile Intraepithelial Neoplasia*.

## Penile Leukoplakia

A persistent white patch on penile skin or mucosa that may reflect irritation, inflammation, or a precancerous disorder and requires assessment.

## Penile Lichen Sclerosus

A chronic inflammatory and sclerosing disorder of the penile skin and foreskin, causing pale tight skin, itching, fissures, pain, or narrowing. It can scar the foreskin or urethral opening and requires treatment and surveillance for complications.

## Penile Prosthesis

An implanted device used to produce an erection when erectile dysfunction has not improved with other treatments. It may be inflatable or bendable. Surgery can cause infection, bleeding, pain, device failure, or erosion into nearby tissue.

---

## Penile Torsion

Rotation of the penis around its longitudinal axis, usually a congenital condition in which the glans or urethral opening is rotated. Symptomatic or marked torsion may require urologic evaluation and surgical correction.

## Penile Trauma

Injury to the penis caused by blunt force, bending, penetration, crushing, burns, bites, or medical instruments. It may damage the skin, erectile tissue, blood vessels, or urethra. A penile fracture is a tear in the covering of the erectile tissue, usually after forceful bending of an erect penis. It may cause pain, swelling, bruising, bleeding, or difficulty passing urine.

## Penis

An external genital organ that contains the outlet for urine and can deliver sperm during sexual activity. It becomes firm when filled with blood during an erection and contains sensitive tissue, skin, blood vessels, and nerves.

## Penis Carcinoma

Penis carcinoma is cancer of the penis, most often arising from skin or squamous cells and sometimes linked to HPV infection.

## Penis Pain

Penis pain may result from infection, injury, inflammation, urinary problems, or sexual conditions; severe pain, swelling, discharge, or inability to urinate requires prompt assessment.

## Penoscrotal Transposition

A congenital difference in which the penis and scrotum are positioned abnormally relative to each other, often associated with hypospadias or other genital and urinary anomalies.

## Pentad Of TTP

The five findings once considered typical of thrombotic thrombocytopenic purpura: low platelets, destruction of red blood cells, neurologic symptoms, kidney problems, and fever. Most patients do not

have all five, so urgent blood testing is essential when TTP is suspected.

## Pentalogy of Cantrell

A rare congenital thoracoabdominal malformation syndrome characterized by the constellation of five midline defects: midline supraumbilical abdominal wall defect (omphalocele), lower sternal cleft defect, anterior diaphragmatic defect, diaphragmatic pericardial defect, and congenital intracardiac anomalies (such as ectopia cordis and VSD).

### Alternate Names:

Thoracoabdominal Syndrome; Cantrell Pentalogy

## Pentamidine

An antimicrobial medicine used for certain parasitic infections and *Pneumocystis pneumonia*, with potentially serious side effects.

## Pentamidine Isethionate

An antiprotozoal and antimicrobial drug used in selected cases of *Pneumocystis pneumonia*, leishmaniasis, trypanosomiasis, and other infections; toxicity includes hypoglycemia, kidney injury, and arrhythmia.

## Pentatrichomonas Hominis

*Pentatrichomonas hominis* is a flagellated intestinal protozoan generally regarded as nonpathogenic; finding it in stool usually indicates exposure to fecal contamination.

## Pentazocine

An opioid pain medicine with mixed agonist and antagonist effects, used for some moderate or severe pain. It can cause drowsiness, dependence, and slowed breathing and may trigger withdrawal in a person dependent on other opioids.

## Pentazocine Overdose

Taking too much pentazocine, an opioid pain medicine, can cause extreme sleepiness, confusion, slow breathing, pinpoint pupils, seizures, or coma. Call emergency services immediately and give naloxone when available and appropriate.



---

### Pentetic Acid

A chelating compound that binds metal ions and is used in selected radiopharmaceuticals, contrast agents, or laboratory applications.

### Pentobarbital Overdose

Taking too much pentobarbital, a sedative, can severely slow brain activity and breathing, causing low blood pressure, coma, or death. It requires immediate emergency treatment, especially with alcohol or other sedatives.

### Pentosan Polysulfate

A semi-synthetic sulfated polysaccharide used for selected bladder pain syndrome and historically as an anticoagulant; prolonged use can cause pigmentary maculopathy and bleeding.

### Pentose

A pentose is a five-carbon sugar, including ribose in RNA and deoxyribose in DNA, and participates in nucleotide and carbohydrate metabolism.

### Pentose Phosphate Pathway

A cell pathway that uses glucose to make NADPH, which protects cells from chemical damage, and ribose, which helps build DNA and RNA. It is especially important in the liver and red blood cells.

### Pentoxifylline

A methylxanthine derivative that improves red-cell flexibility and is used in some countries for peripheral vascular disease; benefit varies by condition.

### Penumbra

Brain tissue surrounding an area of infarction that is functionally impaired but potentially salvageable if blood flow is restored promptly.

### Peplomycin

Peplomycin is a bleomycin-related chemotherapy medicine used in some countries for selected cancers; lung toxicity is an important risk.

### Peppermint

*Mentha piperita*, used as a flavoring and in some preparations for dyspeptic symptoms; concentrated oil can be hazardous.

### Peppermint Oil Overdose

Swallowing a large amount of peppermint oil can cause nausea, vomiting, abdominal pain, dizziness, allergic reactions, seizures, or lung injury if aspirated. Contact poison control promptly and do not induce vomiting.

### Pepsin

A stomach enzyme that begins breaking proteins into smaller fragments. It is released as inactive pepsinogen and activated by stomach acid, helping digestion before food enters the small intestine.

### Pepsin A

Pepsin A is a stomach protease activated by acid that begins digestion of dietary proteins into smaller peptides.

### Pepsinogen

An inactive precursor released by gastric chief cells and converted to pepsin in acidic conditions.

### Peptic Esophagitis

Inflammation and injury of the esophageal lining caused mainly by repeated gastroesophageal reflux of acid or bile, producing heartburn, chest discomfort, swallowing pain, or strictures.

### Peptic Stricture

A circumferential, fibrotic luminal narrowing of the distal esophagus resulting from chronic, untreated acid peptic injury and collagen deposition in severe gastroesophageal reflux disease (GERD), presenting with progressive solid food dysphagia.

#### Alternate Names:

Esophageal Peptic Stricture

### Peptic Ulcer

An open sore in the lining of the stomach or duodenum, most commonly caused by *Helicobacter pylori*

---

infection or NSAID use.

**Alternate Names:**

Gastric or Duodenal Ulcer; Peptic Ulcer Disease; PUD

**Peptic Ulcer Bleeding**

Haemorrhage caused by erosion of an artery or other blood vessel beneath a gastric or duodenal peptic ulcer. It is a common cause of upper gastrointestinal bleeding and may present with haematemesis, coffee-ground vomiting, melaena, or haemodynamic instability.

**Peptic Ulcer Cross-Reference**

Alternate name; see *Peptic ulcer*.

**Peptic Ulcer Disease**

One or more open sores in the lining of the stomach or first part of the small intestine. Common causes include *Helicobacter pylori* infection and anti-inflammatory medicines; ulcers may cause burning pain, bleeding, or a dangerous perforation.

**Peptidase**

An enzyme that breaks peptide bonds in proteins or peptides.

**Peptide**

A short chain of amino acids linked by peptide bonds. Peptides can form parts of proteins or act independently as hormones, neurotransmitters, immune signals, or medicines.

**Peptide Biosynthesis**

The process by which cells make peptides, short chains of amino acids, by linking amino acids according to genetic instructions. Some peptides act as hormones, neurotransmitters, or signals between cells.

**Peptide Bond**

The chemical link joining two amino acids. Repeated peptide bonds form short peptides and long proteins, which cells use for structure, chemical reactions, communication, transport, and many other

functions.

**Peptide Hormone**

A hormone made of amino acids that binds a cell-surface receptor and regulates processes such as metabolism, growth, reproduction, stress, and fluid balance.

**Peptide Metabolism**

The body's production, breakdown, and use of peptides. These processes regulate signaling molecules, hormones, digestion, and protein turnover.

**Peptide Transport**

The movement of peptides into, out of, or within cells and tissues through transport proteins, vesicles, or other cellular mechanisms. It helps control peptide signaling, digestion, and removal from the body.

**Peptide Yy**

Peptide YY (PYY) is a hormone released mainly by cells in the intestine after eating. It helps reduce appetite and slow movement of food through the digestive tract and is being studied in relation to body weight and metabolism.

**Peptidoglycan**

A strong mesh-like material forming much of the bacterial cell wall. Some antibiotics kill bacteria by preventing them from making or repairing this protective layer.

**Peptococcaceae**

A family of anaerobic gram-positive cocci found in the human microbiota and sometimes associated with infection.

**Peptococcus**

A genus of anaerobic gram-positive cocci that can contribute to mixed infections, especially in damaged or poorly oxygenated tissue.

**Peptococcus Niger**

An anaerobic gram-positive coccus found in the

---

human microbiota and rarely associated with mixed infection.

### **Peptoniphilus**

A genus of anaerobic gram-positive cocci found on skin and mucosal surfaces that can cause opportunistic infection.

### **Peptoniphilus Hareii**

An anaerobic gram-positive coccus occasionally recovered from wounds, abscesses, or other clinical specimens. An anaerobic Gram-positive bacterium sometimes found in wounds and abscesses, where it may contribute to mixed infections.

### **Peptoniphilus Indolicus**

An anaerobic gram-positive coccus found in the genital and other microbiota that can occasionally contribute to mixed infection.

### **Peptostreptococcus**

A group of anaerobic gram-positive cocci associated with dental, skin, abdominal, pelvic, and other mixed infections.

### **Peracetic Acid**

A strong oxidizing disinfectant and sterilant used for equipment and surfaces; concentrated exposure irritates or burns skin, eyes, and airways.

### **Peracute**

Peracute describes disease or a clinical event that develops extremely rapidly, often over minutes to hours, with little time for adaptation or diagnosis.

### **Percentile**

A percentile indicates the percentage of a reference population with a value below a measurement, such as height, weight, or test score.

### **Perception**

The brain's process of organizing and interpreting sensory information into an experience of objects, events, body sensations, or the environment.

### **Perceptual**

Relating to the process of detecting and interpreting sensory information.

### **Perceptual Disorder**

A clinically significant disturbance in the way sensory information is interpreted, including altered visual, auditory, tactile, or body perception.

### **Perceptual Distortion**

An inaccurate or altered interpretation of a real sensory stimulus, which may occur with neurologic disease, psychiatric illness, drugs, migraine, seizures, or sensory impairment.

### **Perceptual Masking**

Reduction in the ability to detect or identify one sensory stimulus because another stimulus overlaps it in time, space, or processing.

### **PERC Rule**

A checklist used by emergency clinicians for people already judged very unlikely to have a blood clot in the lungs. If all criteria are reassuring, testing may be avoided; it is not suitable for home use or for people with a strong suspicion of clot.

#### **Alternate Names:**

Pulmonary Embolism Rule-out Criteria; PERC Score

### **Percussion**

Physical examination in which the body surface is tapped and the resulting sound or vibration is interpreted to estimate underlying air, fluid, or solid tissue.

### **Percussion Myotonia**

A muscle that stays contracted briefly after it is tapped. This may occur in myotonic dystrophy, myotonia congenita, and some other muscle disorders; the finding is interpreted with the rest of the examination.

### **Percussion Note**

The sound heard when the chest is gently tapped

---

during examination. A normal lung sounds hollow; a dull sound may suggest fluid or infection, while an unusually hollow sound may suggest excess air or a collapsed lung.

### **Percutaneous**

Performed through the skin, usually using a needle, catheter, wire, or other instrument. This approach can diagnose or treat disease while avoiding a larger open operation, but bleeding and infection remain possible.

### **Percutaneous Biopsy**

Removal of tissue through the skin using a needle or other instrument, often guided by ultrasound, CT, or another imaging method.

### **Percutaneous Closure**

Closing a heart or blood-vessel opening using a device delivered through a catheter inserted through the skin. It can avoid open surgery in selected patients but has procedure-related risks.

### **Percutaneous Coronary Intervention**

A procedure that guides a thin tube through a blood vessel to a narrowed heart artery, then widens it with a balloon and often supports it with a stent. It can restore blood flow during a heart attack or treat selected severe coronary disease.

#### **Alternate Names:**

PCI; Coronary Angioplasty With Stent Placement

### **Percutaneous Discectomy**

Percutaneous discectomy removes or decompresses part of a herniated spinal disc through a small needle or incision.

### **Percutaneous Diskectomy**

A minimally invasive procedure removing part of a herniated spinal disk through a small skin opening to relieve nerve pressure.

### **Percutaneous Drainage**

Placement of a needle or catheter through the skin to drain a fluid collection or obstructed body space,

often with imaging guidance.

### **Percutaneous Electrical Nerve Stimulation**

Electrical stimulation delivered through the skin to reduce pain signals.

#### **Alternate Names:**

PENS

### **Percutaneous Endoscopic Gastrostomy**

Placement of a feeding tube through the abdominal wall into the stomach using a camera passed through the mouth. It may provide long-term nutrition, fluids, or medicines when swallowing is unsafe or inadequate.

### **Percutaneous Kidney Procedures**

Procedures performed through a small needle or tube passed through the skin into or near the kidney. They may obtain a biopsy, drain infection or urine, or remove stones; bleeding, infection, urine leakage, or injury to nearby organs are possible.

### **Percutaneous Nephrolithotomy**

A procedure that removes a large or complex kidney stone through a small opening in the back. A narrow instrument reaches the kidney, breaks or removes the stone, and may leave a temporary drainage tube; bleeding, infection, and injury are possible.

### **Percutaneous Nephrolithotripsy**

Percutaneous nephrolithotripsy removes or fragments kidney stones through a tract made through the skin into the kidney, usually for larger or complex stones.

### **Percutaneous Nephrostomy**

A procedure that places a drainage tube through the skin of the back into the kidney. Urine drains into a bag when the normal path to the bladder is blocked or needs to be bypassed. It may be used for stones, tumors, infection, or a narrowed urine passage and requires follow-up care.

#### **Alternate Names:**

PCN; Nephrostomy Tube Placement; Percutaneous

---

Renal Drainage

### **Percutaneous Transhepatic Cholangiogram**

A percutaneous transhepatic cholangiogram uses a needle through the liver to inject contrast into bile ducts for imaging and may guide drainage or stent placement.

### **Percutaneous Transhepatic Cholangiography**

PTC is imaging of the bile ducts after contrast is injected through a needle passed through the skin and liver; drainage or stenting may follow.

### **Percutaneous Transluminal Coronary Angioplasty**

Balloon-based treatment that widens a narrowed coronary artery, often followed by stent placement. A catheter procedure using balloon inflation to widen a narrowed coronary artery, sometimes followed by stent placement.

#### **Alternate Names:**

PTCA; Percutaneous Coronary Intervention

### **Percutaneous Umbilical Blood Sampling**

A procedure that uses ultrasound to guide a needle into the umbilical cord and collect a fetal blood sample during pregnancy. It may investigate anemia, infection, or genetic problems and carries risks such as bleeding, infection, and pregnancy loss.

### **Perforant Pathway**

A group of nerve fibers carrying information from a brain area involved in memory to the hippocampus. It helps the brain form and retrieve memories and can be affected in some dementias and seizure disorders.

### **Perforated Bowel**

A perforated bowel is a hole through the intestinal wall allowing contents to leak into the abdomen, causing peritonitis and potentially life-threatening sepsis.

### **Perforated Eardrum**

Alternate name; see *Ruptured eardrum (perforated eardrum)*.

### **Perforating Folliculitis**

A rare follicular disorder in which itchy keratotic papules develop around follicles and abnormal keratin is expelled through the epidermis. It is associated with chronic kidney disease, diabetes, and other systemic illness and may need treatment of both the skin and underlying disease.

### **Perforation**

A hole or complete tear through the wall of an organ or tissue. It can release contents into surrounding spaces, cause severe infection or bleeding, and often requires urgent medical treatment.

### **Perforation Of A Viscus**

A hole through the wall of a hollow abdominal organ, such as the stomach, small intestine, or colon. Contents can leak into the abdomen and cause severe infection; ulcers, inflammation, poor blood flow, and injury are possible causes.

### **Performance**

How effectively a person, organ, treatment, test, or system carries out a defined task. Examples include exercise performance, heart performance, test performance, and response to treatment.

### **Performance Anxiety**

Anxiety about sexual performance that can interfere with sexual function.

### **Performance Status**

A rating of how well a person can move, work, and perform daily activities while ill. Clinicians use it to estimate prognosis, decide whether some treatments are safe, and track change over time, especially during cancer care.

### **Perfusion**

The flow of blood through the small vessels of a tissue or organ. Good perfusion supplies oxygen

---

and nutrients and removes waste; poor perfusion can cause pain, weakness, tissue injury, or organ failure.

### **Perfusion Defect**

An area receiving unusually little blood flow, identified by an imaging or function test. It may reflect a blocked artery, narrowed vessel, poor heart output, inflammation, or damaged tissue.

### **Perfusion Imaging**

Imaging that shows how blood flows through an organ or tissue. It can help assess problems such as stroke, reduced heart blood flow, or tumor blood supply.

### **Perfusion PET**

Positron-emission tomography that measures tissue blood flow using a suitable radioactive tracer. Cardiac perfusion PET can show reduced blood flow, viability, and sometimes the effects of treatment.

### **Perianal**

Located around the anus; the term describes the skin, glands, pain, inflammation, abscesses, fistulas, and other findings in that region.

### **Perianal Abscess**

A localized collection of pus near the anus caused by infected anal glands or surrounding tissue, producing severe throbbing pain, swelling, fever, and sometimes a fistula.

### **Perianal Glands**

Small glands near the anus that contribute secretions to the local tissue; obstruction or infection can contribute to perianal abscesses and fistulas.

### **Perianal Streptococcal Cellulitis**

A bacterial skin infection around the anus, usually caused by group A *Streptococcus*. It may cause bright-red skin, pain, itching, discharge, or pain during bowel movements.

### **Periapical Abscess**

An acute or chronic localized collection of pus at the apex of a tooth's root, arising from pulpal infection and necrosis (usually due to untreated dental caries or trauma).

### **Periapical Granuloma**

A chronic inflammatory lesion at the apex of a tooth, usually resulting from pulpal necrosis and persistent apical infection. It may be asymptomatic and is diagnosed through clinical, radiographic, and sometimes histopathologic assessment.

### **Periappendiceal**

Periappendiceal means located around the appendix, as in inflammation, fluid, an abscess, or fat stranding near the appendiceal tissue.

### **Periarteritis Nodosa**

An older name for polyarteritis nodosa, a necrotizing vasculitis of medium-sized arteries that can cause skin, nerve, muscle, kidney, and abdominal ischemic symptoms without typical glomerulonephritis.

### **Periarticular**

Periarticular means located around a joint, including nearby tendons, bursae, ligaments, capsule, and soft tissues rather than the joint surface itself.

### **Pericardial**

Relating to the pericardium, the thin two-layered sac surrounding the heart. A small amount of fluid between its layers allows the heart to beat smoothly, while excess fluid or inflammation can interfere with heart function.

### **Pericardial Cavity**

The thin fluid-containing space between the layers of the sac surrounding the heart. A small amount of fluid lets the heart move smoothly; excess fluid or blood can press on the heart and impair its pumping.

### **Pericardial Effusion**

Abnormal accumulation of fluid in the sac around the heart. Small or slowly developing effusions may



---

cause no symptoms, while rapid or large accumulation can impair heart filling and cause cardiac tamponade; significance depends on the amount, rate of accumulation, and cause.

### **Pericardial Effusion Etiology**

Fluid around the heart may result from infection, cancer, kidney failure, low thyroid activity, autoimmune disease, heart injury, surgery, radiation, trauma, or medicines. Tests and the person's history help identify the cause.

### **Pericardial Fluid Analysis**

Laboratory examination of fluid removed from the pericardial space, usually during pericardiocentesis. Cell counts, chemistry, cytology, cultures, and targeted tests help investigate infection, cancer, tuberculosis, inflammation, or bleeding; results must be interpreted with the clinical context.

### **Pericardial Fluid Culture**

A microbiologic culture of fluid from the pericardial space to identify bacteria, fungi, or other organisms in suspected infectious pericarditis.

### **Pericardial Fluid Gram Stain**

A microscopic stain of pericardial fluid that can rapidly suggest bacterial organisms, although a negative result does not exclude infection.

### **Pericardial Friction Rub**

A scratchy or grating sound heard when the inflamed layers around the heart rub together. It can support a diagnosis of acute pericarditis, especially when heard along the left side of the breastbone, but it may come and go and is not present in every case.

**Alternate Names:**

Pericardial Rub; Triphasic Pericardial Sound

### **Pericardial Inflammation**

Alternate name; see *Pericarditis*.

### **Pericardial Knock**

An extra early heart sound caused when a stiff, scarred sac around the heart suddenly stops the

heart from filling. It may suggest constrictive pericarditis and must be interpreted with other examination and test findings.

### **Pericardial Knock Sound**

An early diastolic, high-pitched, sharp, knocking extra heart sound heard on auscultation in patients with chronic constrictive pericarditis. Occurs 0.08 to 0.12 seconds after aortic valve closure ( $S_2$ ), earlier than a standard third heart sound ( $S_3$ ), caused by the sudden abrupt cessation of ventricular filling when the expanding ventricular myocardium collides with the rigid, non-compliant, calcified pericardium.

**Alternate Names:**

Pericardial Knock; Constrictive Knock Sound

### **Pericardial Mesothelioma**

A rare malignant tumor arising from the mesothelial lining around the heart, which can cause effusion, constriction, or impaired cardiac filling.

### **Pericardial Metastases**

Cancer that has spread to the sac around the heart from another part of the body. It may cause inflammation, fluid or blood around the heart, breathlessness, chest discomfort, or impaired heart filling.

### **Pericardial Rub**

A high-pitched, superficial, grating or scratching heart sound produced by the friction between inflamed parietal and visceral pericardial layers. It characteristically has up to three components per cardiac cycle (atrial systole, ventricular systole, and early ventricular diastole) and is best heard at the lower left sternal border with the patient leaning forward in expiration.

**Alternate Names:**

Pericardial Friction Rub

### **Pericardial Stripping**

Surgery to remove part or most of the thickened or scarred sac around the heart. It may relieve constrictive pericarditis, in which the stiff sac prevents the heart from filling normally; bleeding, rhythm problems, and other major surgical risks are possible.

---

## Pericardial Tamponade

A life-threatening buildup of fluid, blood, or pus in the sac around the heart. The pressure prevents the heart from filling properly and can cause low blood pressure, breathlessness, fainting, shock, or cardiac arrest. Bedside ultrasound can support the diagnosis, but the examination and cause are also important. Emergency drainage with a needle or surgery may be needed.

### Alternate Names:

Cardiac Tamponade; Pericardial Compression; Hemopericardium with Tamponade

## Pericardial Wall

The layered tissue forming the protective sac around the heart. Its inner surfaces produce a small amount of lubricating fluid that lets the heart beat with little friction.

## Pericardial Window

An operation that creates an opening in the sac around the heart so extra fluid can drain into a nearby space and be absorbed. It may prevent repeated fluid buildup and allow a tissue sample to be taken.

### Alternate Names:

Subxiphoid Pericardiostomy; Pleuropericardial Window; Pericardiostomy

## Pericardiectomy

Surgery to remove part or most of the sac around the heart when thickening or scarring prevents the heart from filling normally. It is major surgery and may improve constrictive pericarditis but carries bleeding and other serious risks.

## Pericardiocentesis

A needle or catheter is used to remove fluid or blood from the sac around the heart, usually to relieve dangerous pressure or investigate an effusion. Imaging guidance improves safety, but bleeding, abnormal rhythm, organ injury, or infection can occur.

## Pericardiostomy

A surgical opening made in the pericardium to drain

fluid or prevent recurrent accumulation around the heart.

## Pericarditis

Inflammation of the sac around the heart. It often causes sharp chest pain that worsens with deep breathing or lying down and improves when sitting up and leaning forward. It may be caused by infections, autoimmune diseases, heart injury, kidney disease, cancer, or some medicines. Severe cases can cause fluid buildup around the heart.

### Alternate Names:

Pericardial Sac Inflammation; Acute Pericarditis

## Pericardium

A two-layered protective sac surrounding the heart and beginnings of major blood vessels, reducing friction during each heartbeat.

## Perichondritis

Perichondritis is inflammation or infection of tissue surrounding cartilage, often affecting the outer ear after trauma or piercing and potentially damaging cartilage.

## Perichondritis Of The Ear

Infection or inflammation of the auricular perichondrium, often after trauma, piercing, or surgery. *Pseudomonas aeruginosa* is an important cause; delayed treatment can produce cartilage necrosis and permanent deformity.

## Perichondrium

A tough layer of tissue surrounding most cartilage. It supplies nutrients, contains cells that help cartilage grow and repair, and can become inflamed or infected, especially around the outer ear.

## Pericoronitis

Inflammation of the soft tissue around the crown of a partially erupted tooth, most often a mandibular third molar. Food retention and bacterial accumulation may cause pain, swelling, trismus, fever, or spread into deeper spaces.

---

## Pericytes

Pericytes are contractile support cells wrapped around small blood vessels that help regulate capillary stability, blood flow, repair, and the blood-brain barrier.

## Periductal Mastitis

A chronic inflammatory condition of the subareolar breast ducts primarily seen in young women and strongly associated with cigarette smoking. Toxins in tobacco smoke damage the ductal lining, causing squamous metaplasia and keratin plugging of the lactiferous ducts. The blocked duct dilates, ruptures, and incites chronic inflammation and secondary infection, presenting as a painful periareolar mass, nipple retraction, or recurrent subareolar abscesses that frequently drain and form a chronic fistula (Zuska's disease). Treatment includes smoking cessation, targeted antibiotics, abscess drainage, or surgical duct excision.

### Alternate Names:

Zuska's Disease; Subareolar Abscess; Squamous Metaplasia of Lactiferous Ducts; Periareolar Breast Abscess

## Perifolliculitis Capitis Abscedens et Suffodiens

A severe chronic scalp folliculitis, also called dissecting cellulitis, in which deep nodules, abscesses, and interconnected tunnels cause purulent drainage and permanent scarring alopecia. It is part of the follicular-occlusion spectrum and requires specialist treatment.

## Peri-Implantitis

Inflammation around a dental implant with progressive loss of the supporting bone. It may cause bleeding, swelling, pus, pain, or implant looseness and requires dental assessment.

## Perilymph

Fluid in the bony labyrinth of the inner ear that surrounds the membranous labyrinth and helps transmit sound and balance signals.

## Perilymphatic Fistula

An abnormal leak between the fluid spaces of the inner ear and the middle ear, often after injury, pressure changes, or surgery. It may cause dizziness, hearing loss, ringing, or symptoms triggered by sound or pressure.

## Perimenopausal

Perimenopausal describes the years of hormonal and menstrual transition before and around menopause, when cycles, bleeding, hot flashes, sleep, and fertility may change.

## Perimenopause

The transition time in a woman's life that begins when ovaries produce less estrogen and menstruation becomes less frequent, and ends when the ovaries no longer produce eggs and menstruation stops.

## Perimetry

A test that measures the full field of vision, including central and side vision. It helps detect and monitor problems such as glaucoma, optic-nerve disease, retinal disease, and disorders affecting the visual pathways in the brain.

## Perimysium

A layer of connective tissue that surrounds groups of muscle fibers, forming bundles called fascicles. It carries small blood vessels and nerves within the muscle.

## Perinatal

Perinatal refers to the period around birth, conventionally including late pregnancy and the first days or weeks after delivery depending on the definition used.

## Perinatal Asphyxia

Too little oxygen or inadequate breathing before, during, or soon after birth. Severe cases can injure the brain and other organs; rapid newborn assessment, breathing support, and treatment of the cause are essential.

---

## Perinatal Death

Death of a fetus or newborn during the late pregnancy, birth, or early newborn period; the exact time range varies by definition.

## Perinatal Psychiatry

The subspecialty focused on psychiatric disorders during pregnancy and the postpartum period. Common conditions include perinatal depression, perinatal anxiety, perinatal OCD, and postpartum psychosis. Medication selection must balance maternal mental health with fetal or neonatal exposure risks.

## Perinatal Transmission

Transmission of an infection or condition from mother to infant during pregnancy, labor, delivery, or breastfeeding, as with HIV, hepatitis B, syphilis, or certain genetic and congenital disorders.

## Perinatologist

A perinatologist is an obstetrician specializing in high-risk pregnancy, fetal medicine, prenatal diagnosis, and complications affecting the pregnant person or fetus.

## Perindopril

Perindopril is an ACE inhibitor used for high blood pressure and cardiovascular risk reduction; it can cause cough, high potassium, or rarely angioedema.

## Perineal

Relating to the area between the external genitals and anus, which can be affected by injury or infection.

## Perineal Hemostat

A clamp used during procedures to compress a blood vessel or hold tissue and control bleeding.

### Alternate Names:

Perineal Hemostatic Clamp; Pelvic Bleeding Forceps

## Perineal Laceration

Tearing of the perineal tissues during vaginal delivery. The injury is graded by depth: it may involve skin, perineal muscle, the anal sphincter, or the

rectal mucosa.

## Perineal Laceration Classification

A system grading tears during childbirth by depth: first degree affects skin, second degree also affects muscle, third degree involves the anal-control muscle, and fourth degree extends into the rectum. Accurate examination and repair reduce later pain and bowel-control problems.

## Perineal Needle Holder

A surgical instrument used to hold and guide a suture needle while repairing tissues of the perineum. Its locking handle allows controlled needle placement and helps protect the surgeon's hands.

### Alternate Names:

Perineorrhaphy Needle Holder; Perineal Needle Holder

## Perineal Protection

Measures used during the pushing stage of childbirth to reduce severe tearing of the tissue between the vagina and anus. They may include controlled delivery of the baby's head, support of the perineum, a warm compress, and a selective approach to episiotomy.

## Perineal Suture

A thread or strand of material used to approximate tissues (bring wound edges together) and hold them in place until healing occurs. See Suture.

### Alternate Names:

Perineal Suture; Perineorrhaphy Suture

## Perineal Tear

A split in the tissue between the vagina and anus during vaginal birth. First-degree tears affect skin, second-degree tears reach muscle, third-degree tears involve the anal-control muscle, and fourth-degree tears extend into the rectum; severe tears need careful repair and follow-up.

## Perineum

The area between the external genitalia and the anus. It contains skin, muscles, nerves, and connec-

---

tive tissue and may be affected by injury, infection, childbirth, or surgery.

### **Perineural**

Perineural means around a nerve; the term may describe normal connective tissue, inflammation, tumor spread, or imaging findings surrounding a nerve.

### **Perineuronal Satellitosis**

A microscopic pattern in which supporting brain cells cluster around a nerve cell. It can occur in some brain tumors, inflammation, or reactive changes and has meaning only with the surrounding tissue findings.

### **Perinuclear Space**

The narrow space between the two membranes surrounding a cell nucleus. It connects with part of the cell's internal membrane network and helps form the nuclear envelope.

### **Periodic Abstinence**

Periodic abstinence is avoiding intercourse during estimated fertile days to reduce pregnancy likelihood; cycle timing is imperfect and does not prevent infections.

### **Periodic Acid**

A chemical used to stain tissue samples. In the periodic-acid-Schiff test, it helps highlight sugars, mucus, certain fungi, and supporting membranes so they can be seen under a microscope.

### **Periodic Breathing**

Repeated cycles of shallow or absent breaths followed by deeper breaths. It can occur normally for short periods in some infants or during sleep, but persistent episodes may reflect a brain, heart, or lung disorder.

### **Periodic Limb Movement Disorder**

Repeated involuntary limb movements during sleep that may disrupt rest.

**Alternate Names:**

PLMD; Periodic Leg Movements in Sleep; Nocturnal Myoclonus

### **Periodic Limb Movement Sleep Disorder**

Alternate name; see *Periodic limb movement disorder*.

### **Periodic Paralysis Syndrome**

Periodic paralysis syndromes are rare inherited or acquired disorders causing episodic muscle weakness associated with abnormal potassium handling or thyroid disease. Attacks may follow rest after exercise, meals, or stress; ECG, potassium testing, and genetic or endocrine evaluation guide care.

### **Periodontal Abscess**

A localized accumulation of pus within periodontal tissues, commonly associated with a periodontal pocket, foreign material, or impaired drainage. It may cause pain, swelling, suppuration, tooth mobility, and systemic symptoms and requires urgent dental assessment.

### **Periodontal Cyst**

A periodontal cyst is a fluid-filled epithelial-lined lesion near the tooth-supporting tissues, with effects ranging from incidental discovery to local bone destruction.

### **Periodontal Disease**

Infection or inflammation of the tissues supporting teeth, including gums and jawbone. It can cause bleeding, swelling, bad breath, loose teeth, bone loss, and tooth loss, but treatment can slow progression.

### **Periodontal Dressing**

A protective material placed over the gums after selected dental or periodontal procedures. It can protect the surgical area while it heals, reduce irritation, and support comfort; it should be kept clean and reviewed if it becomes loose, painful, foul-smelling, or causes swelling.

### **Periodontal Flap Surgery**

Dental surgery in which the gum is gently lifted so

---

a clinician can clean infection and deposits below the gumline and, when needed, reshape accessible bone before replacing the gum.

### **Periodontal Graft**

Tissue placed around a tooth to cover an exposed root, thicken the gum, or repair gum loss. The tissue may come from the person's mouth, a donor, or a processed substitute.

#### **Alternate Names:**

Periodontal Soft Tissue Graft; Gingival Graft

### **Periodontal Ligament**

Strong tissue fibers that attach a tooth's root to the surrounding jawbone. They cushion chewing forces, help the body sense pressure on the tooth, and can become inflamed or damaged in gum disease or grinding.

### **Periodontal Pocket**

An abnormally deep space between a tooth and the surrounding gum caused by loss of attachment. Plaque can collect there, leading to bleeding, infection, bone loss, loose teeth, and eventual tooth loss.

### **Periodontal Probing**

A dentist gently measures the space between the gum and tooth with a marked instrument. Deeper spaces, bleeding, and loss of support can indicate gum disease; measurements help assess severity and monitor treatment.

### **Periodontics**

The dental specialty focused on preventing, diagnosing, and treating gum disease and tissues supporting the teeth.

### **Periodontist**

A periodontist is a dentist specializing in prevention, diagnosis, and treatment of gum disease and supporting tissues, including periodontal surgery and implants.

### **Periodontitis**

A chronic inflammatory disease caused by dysbiotic dental plaque that destroys the periodontal ligament and alveolar bone supporting the teeth. It can cause bleeding gums, periodontal pockets, tooth mobility, recession, bad breath, and tooth loss; severity and progression vary and require dental evaluation.

#### **Alternate Names:**

Pyorrhea; Periodontal Disease; Periodontal Gum Disease; Advanced Gum Disease

### **Periodontium**

The tissues that hold and support a tooth, including gums, ligament, root covering, and jawbone. Disease of these tissues can cause bleeding, gum recession, loose teeth, bone loss, and tooth loss.

### **Periodontium Disease**

Disease of the tissues supporting the teeth, including gingiva, periodontal ligament, cementum, and alveolar bone; periodontitis can cause attachment loss and tooth loss.

### **Perionychium**

The skin and soft tissue surrounding a fingernail or toenail, including the nail folds and nearby cuticle; infection or inflammation causes paronychia.

### **Perioperative**

Perioperative means relating to the period before, during, and after an operation, including preparation, anesthesia, recovery, and complication prevention.

### **Perioral Dermatitis**

A papulopustular eruption around the mouth, sparing the vermilion border. It is more common in young women and associated with topical corticosteroid use. Other triggers include fluoridated toothpaste, cosmetics, and inhaler steroids.

### **Periorbital**

Located around the eye socket or eyelids. Periorbital swelling may result from injury, allergy, infection, or fluid retention.



---

### **Periorbital Cellulitis**

A bacterial infection of the eyelid and tissues around the eye that causes redness, swelling, and tenderness. It must be distinguished from orbital cellulitis, which can threaten vision and spread more deeply.

### **Periorbital Edema**

Swelling of the tissues around the eye, caused by allergy, infection, trauma, venous or lymphatic congestion, kidney disease, thyroid eye disease, or other systemic conditions.

### **Periorbital Infection**

Infection of the tissues around the eye; it can cause swelling and redness and must be distinguished from orbital cellulitis, which threatens vision.

### **Periosteal Osteosarcoma**

A cancerous bone tumor that develops beneath the thin tissue covering a bone, usually along the shaft of a long bone. It often makes cartilage and causes pain or swelling, requiring specialist cancer treatment.

### **Periosteal Reaction**

The non-specific radiographic and histological response of the periosteum to cortical bone irritation, injury, infection, or underlying malignancy, manifesting as solid, laminated ("onion-skin"), spicular ("sunburst"), or Codman triangle bone formation.

### **Periosteum**

A blood-vessel-rich, pain-sensitive tissue layer covering the outer surface of bone, except where joint cartilage covers it. It supplies nutrients, supports repair, and provides attachment for tendons and ligaments.

### **Periostitis**

Inflammation of the periosteum, the pain-sensitive tissue covering the outer surface of bone. It may cause localized bone pain, tenderness, or swelling and can result from repeated stress, infection, injury, or another disease.

### **Peripancreatic Malignancy**

Cancer involving the tissues around the pancreas, either by spreading from the pancreas or from another organ. It may cause abdominal or back pain, weight loss, jaundice, or newly developed diabetes and usually needs detailed imaging and specialist care.

### **Peripartum**

Relating to the period around childbirth, including late pregnancy, labor, delivery, and the early weeks after birth. Clinicians may use a more specific time range for a particular condition or study.

### **Peripartum Cardiomyopathy**

A usually dilated cardiomyopathy that develops toward the end of pregnancy or during the postpartum period and may cause systolic heart failure. Features can include breathlessness, difficulty lying flat, swelling, fatigue, or chest discomfort.

#### **Alternate Names:**

PPCM; Pregnancy-Associated Cardiomyopathy

### **Peripartum Cardiomyopathy Prognosis**

Recovery of ventricular function varies widely. Prognosis is influenced by the initial ejection fraction, ventricular size, symptoms, treatment, complications, and follow-up; persistent dysfunction requires ongoing cardiac care.

### **Peripartum (Postpartum) Cardiomyopathy (PPCM)**

A form of heart-muscle failure developing toward the end of pregnancy or in the months after delivery. It may cause breathlessness, orthopnea, swelling, chest discomfort, palpitations, or fatigue and requires urgent cardiac evaluation and pregnancy-compatible treatment planning.

### **Peripheral**

Peripheral means away from the center or central nervous system; in clinical use it may describe nerves, vessels, lesions, circulation, or symptoms outside the brain, spinal cord, or central structures.

---

## Peripheral Angioplasty

A percutaneous vascular intervention that dilates stenosed or occluded peripheral arteries—primarily in the lower extremities, renal arteries, or carotid vessels—using balloon dilation and optional stent placement.

### Alternate Names:

PTA; Peripheral Arterial Angioplasty

## Peripheral Artery Disease

Narrowing of arteries supplying the limbs, usually caused by atherosclerosis. It may cause exertional leg pain relieved by rest, reduced pulses, nonhealing wounds, or chronic limb-threatening ischemia, a severe reduction in blood flow associated with persistent pain or tissue loss. Also called peripheral arterial disease.

### Alternate Names:

PAD; Peripheral Arterial Occlusive Disease; PAOD

## Peripheral Blood

Peripheral blood is blood obtained from a vein or capillary rather than directly from an organ or marrow, used for counts, smears, chemistry, and other testing.

## Peripheral Blood Smear

Microscopic examination of stained blood spread on a slide. It shows the size, shape, and appearance of red cells, white cells, and platelets and can help investigate anemia, infection, and blood disorders.

## Peripheral Blood Stem Cells

Immature blood-forming cells collected from the bloodstream after medicines encourage them to leave the bone marrow. They can be returned to the same person or given to another person to rebuild blood production after intensive treatment.

## Peripheral Cyanosis

A bluish or gray-blue discoloration of the fingers, toes, hands, or feet caused by reduced peripheral blood flow or increased oxygen extraction in tissues. The lips and tongue are usually not affected, distinguishing it from central cyanosis.

## Peripheral Edema

Swelling due to accumulation of fluid in the interstitial space of the limbs, most commonly the lower extremities.

## Peripheral Ischemia

Inadequate arterial blood flow to an arm, leg, digit, or other peripheral tissue, causing pain, pallor, coolness, weak pulses, numbness, ulcers, or tissue loss.

## Peripherally Inserted Central Catheter

A long, flexible catheter inserted into the basilic or cephalic vein and advanced so the tip sits in the superior vena cava. Inserted at the bedside, often by a specialized PICC nurse, with ultrasound and ECG or X-ray confirmation. Indications: long-term IV antibiotics (>2 weeks), TPN, chemotherapy, and frequent blood draws.

## Peripheral Nerve

A nerve outside the brain and spinal cord that carries motor, sensory, and autonomic signals between the central nervous system and the rest of the body.

## Peripheral Nerve Blocks

Injections of local anesthetic near a nerve or group of nerves to numb a specific area and reduce pain during or after a procedure. Ultrasound guidance may improve accuracy and safety.

## Peripheral Nerve Injuries

Damage to nerves outside the brain and spinal cord caused by trauma, compression, stretching, disease, or toxins. It may cause pain, numbness, weakness, or loss of function; recovery depends on the severity and location of the injury.

## Peripheral Nerve Tumors

Growths arising from or around nerves outside the brain and spinal cord; they may be benign or cancerous.

## Peripheral Nervous System

All nerves outside the brain and spinal cord. It carries messages between the central nervous system

---

and the skin, muscles, organs, and glands, controlling movement, sensation, and automatic functions such as heartbeat and digestion.

### **Peripheral Neuritis**

Alternate name; see *Peripheral Neuropathy*.

### **Peripheral Neurofibromatosis**

Alternate name; see *Neurofibromatosis type 1*.

### **Peripheral Neuropathy**

Damage or dysfunction of nerves outside the brain and spinal cord. It may cause burning pain, tingling, numbness, weakness, or problems with sweating and other automatic body functions, often beginning in the feet or hands.

#### **Alternate Names:**

Peripheral Nerve Damage; Polyneuropathy; Neuritis

### **Peripheral Oedema**

Alternate spelling of Peripheral Edema; swelling caused by excess interstitial fluid, usually in the limbs.

### **Peripheral Vascular Disease**

A broad term for disease of blood vessels outside the heart and brain. It is often used to mean peripheral artery disease, in which narrowed leg arteries cause pain with walking, coldness, wounds that heal slowly, or tissue loss; the vessel involved should be specified because venous disorders are different.

#### **Alternate Names:**

PVD; Peripheral Angiopathy

### **Peripheral Vision**

The parts of the visual field seen away from the direct line of sight. Loss can result from eye disease, nerve damage, brain injury, or glaucoma and may increase falls or collisions.

#### **Alternate Names:**

Side Vision; Visual Field

### **Periplasm**

The periplasm is the space between the inner and outer membranes of Gram-negative bacteria, con-

taining enzymes, transport proteins, and structural components.

### **Periprosthetic Joint Infection**

Infection involving an artificial joint and nearby tissues. It may cause persistent pain, swelling, drainage, fever, or loosening. Diagnosis uses examination, blood tests, joint-fluid testing, cultures, and imaging; treatment may require antibiotics and surgery.

### **Perirenal Abscess**

A perirenal abscess is a collection of pus around the kidney, usually from spread of a urinary or kidney infection, causing fever, flank pain, and possible sepsis.

### **Perirenal Extramedullary Hematopoiesis**

Production of blood cells in tissue around the kidney instead of mainly in the bone marrow. It is an unusual response to severe long-term anemia or bone-marrow disease and may appear as masses on imaging.

### **Perirenal Hematoma**

A collection of blood around the kidney, usually caused by trauma, a bleeding lesion, a procedure, or anticoagulation.

### **Peristalsis**

Peristalsis is coordinated, wave-like contraction of smooth muscle that propels food, fluid, or other contents through the gastrointestinal tract and some ducts.

### **Peritoneal**

Peritoneal means relating to the peritoneum, the thin membrane lining the abdominal cavity and covering many abdominal organs.

### **Peritoneal Cancer**

Cancer arising from or involving the peritoneum, the membrane lining the abdominal cavity and covering many abdominal organs. It may be primary or metastatic and can cause abdominal swelling, pain,

---

bowel symptoms, or ascites; imaging, fluid analysis, and tissue diagnosis help determine the origin.

### **Peritoneal Carcinomatosis**

Cancer that has spread widely across the lining of the abdomen, often from the bowel, stomach, ovary, or another organ. It may cause abdominal swelling, pain, fluid buildup, weight loss, or bowel blockage; scans and fluid or tissue tests guide treatment.

### **Peritoneal Cavity**

The narrow space inside the abdominal lining around the abdominal and pelvic organs. A small amount of fluid normally lubricates it; blood, pus, excess fluid, or cancer cells can collect there during disease.

### **Peritoneal Dialysis**

A treatment for kidney failure that uses the lining of the abdomen to remove waste and extra water. Clean dialysis fluid enters through a permanent tube and is later drained; it can be done at home but may cause infection, fluid imbalance, or catheter problems.

### **Peritoneal Dialysis-Associated Peritonitis**

Infection or inflammation of the abdominal lining during peritoneal dialysis, usually related to contamination of the dialysis catheter or fluid. It may cause cloudy drainage, abdominal pain, fever, or nausea.

### **Peritoneal Dialysis Catheter**

A soft tube placed through the abdominal wall into the abdominal cavity for dialysis. Dialysis fluid enters and leaves through it to remove waste and extra water; infection, blockage, leakage, or displacement can occur.

### **Peritoneal Diseases**

Disorders affecting the thin lining of the abdomen, including infection, inflammation, cancer, bleeding, scar tissue, or abnormal fluid buildup. Symptoms depend on the cause and may include abdominal pain, swelling, fever, or vomiting.

### **Peritoneal Fluid**

The small amount of slippery fluid normally present inside the abdominal cavity. Excess fluid, blood, pus, or cancer cells can collect there and may cause swelling, pain, infection, or other signs of abdominal disease.

### **Peritoneal Fluid Analysis**

Laboratory testing of fluid collected from the abdominal cavity. Cell counts, protein, albumin, cultures, and other tests help determine whether the fluid is caused by liver disease, infection, cancer, bleeding, or another problem.

### **Peritoneal Fluid Culture**

Culture of ascitic fluid to detect microorganisms, particularly in suspected spontaneous bacterial peritonitis or secondary peritoneal infection.

### **Peritoneal Infection**

Infection of the lining of the abdominal cavity, often called peritonitis. Causes include a ruptured appendix or bowel, abdominal surgery, pelvic infection, or contamination during dialysis. Severe abdominal pain, fever, vomiting, or a rigid abdomen requires urgent assessment.

### **Peritoneal Lavage**

Washing or sampling the abdominal cavity with sterile fluid. It was once used more often to detect internal bleeding or intestinal contents after injury; imaging and other tests now commonly provide this information.

### **Peritoneal Necrosis**

Death of peritoneal tissue caused by ischemia, severe infection, inflammation, trauma, or tumor, potentially leading to peritonitis, adhesions, sepsis, or organ failure.

### **Peritoneal Sac**

The space formed by the abdominal lining around the abdominal and pelvic organs. It normally contains only a small amount of lubricating fluid, but blood, pus, excess fluid, or cancer can collect there.

---

## Peritoneum

A thin, slippery lining that covers the inside of the abdomen and many abdominal organs. It allows organs to move smoothly; inflammation, infection, bleeding, fluid, or cancer in this lining can cause severe abdominal symptoms.

## Peritoneum Neoplasm

An abnormal growth in the lining of the abdominal cavity. It may begin there or spread from cancer in another organ and can cause abdominal pain, swelling, fluid buildup, or bowel blockage.

## Peritonitis

Inflammation or infection of the peritoneum, the lining inside the abdomen. It may result from a hole in the stomach or bowel, appendicitis, abdominal injury, surgery, or infected abdominal fluid. It can cause severe abdominal pain, tenderness, a swollen or hard abdomen, fever, nausea, vomiting, or inability to pass stool or gas. It is a medical emergency because it can cause sepsis or shock.

## Peritonitis In PD

Infection of the abdominal lining in a person receiving peritoneal dialysis. Cloudy drainage, abdominal pain, fever, nausea, or feeling unwell needs prompt testing of the drainage and antibiotics; severe or repeated infection may require catheter removal.

## Peritonsillar Abscess

A collection of pus beside a tonsil, usually after a throat infection. It can cause severe one-sided throat pain, fever, muffled speech, drooling, difficulty opening the mouth, or trouble breathing and requires urgent treatment and drainage.

## Periungual Fibroma

A noncancerous, firm growth arising beside or beneath a fingernail or toenail. It may distort the nail and can occur alone or as one of several skin findings in tuberous sclerosis complex.

## Periurethral

Located around the urethra, the tube that carries urine out of the body. The term may describe glands,

infection, an abscess, a urethral diverticulum, or tissue and treatments used for urinary leakage.

## Periventricular Leukomalacia

Injury to the brain's developing nerve fibers near its fluid-filled spaces, most often in premature infants. It may later affect movement, muscle tone, vision, learning, or development and is associated with cerebral palsy.

## Perivitelline Space

The perivitelline space is the narrow region between an egg's plasma membrane and surrounding vitelline or zona material, important in fertilization and early development.

## Perleche

Inflammation, soreness, and cracking at one or both corners of the mouth. Saliva, ill-fitting dentures, yeast or bacterial infection, skin irritation, and iron or vitamin shortages can contribute; treatment depends on the cause and keeping the area dry.

## Permanent Teeth

The adult set of teeth that replaces the deciduous teeth.

## Permease

A permease is a membrane transport protein that facilitates movement of a specific molecule or ion across a cell membrane.

## Permethrin

A skin-applied medicine that kills scabies mites and head lice. It is usually applied as directed and washed off later; itching may continue briefly after treatment even when parasites are gone.

## Permissive Hypercapnia

A ventilator approach that allows carbon dioxide in the blood to remain above normal so that gentler pressures and smaller breaths can protect injured lungs. It is used selectively and requires close monitoring because excessive acidity or carbon dioxide can be harmful.

---

## **Pernicious Anaemia**

A form of vitamin B12 deficiency caused by immune damage to stomach cells needed for B12 absorption. It can cause anemia, tiredness, sore tongue, numbness, balance problems, and nerve damage.

## **Pernicious Anemia**

Anemia caused by vitamin B12 deficiency when the immune system damages stomach cells that make a substance needed for B12 absorption. It may cause tiredness, sore tongue, numbness, balance problems, or permanent nerve damage if untreated.

### **Alternate Names:**

Biermer Disease; Addisonian Anemia

## **Pernio**

Alternate name; see *Chilblains*.

## **Peroneal**

Relating to the fibula or the muscles and nerve on the outer side of the lower leg.

## **Peroneal Muscular Atrophy**

Alternate name; see *Charcot-Marie-Tooth disease*.

## **Peroneal Nerve**

The common fibular nerve, a branch of the sciatic nerve that winds around the fibular neck and supplies sensation and muscles responsible for ankle dorsiflexion and eversion.

## **Peroneal Nerve Palsy**

A peripheral lower motor neuropathy caused by trauma or compression of the common fibular (peroneal) nerve as it courses superficially around the neck of the fibula, presenting with foot drop (loss of ankle dorsiflexion and eversion) and sensory numbness over the anterolateral lower leg and dorsum of the foot.

### **Alternate Names:**

Common Fibular Nerve Palsy; Foot Drop Palsy

## **Peroneal Tendon Syndromes**

Disorders of the peroneal tendons, including tendinopathy, subluxation, and tears, usually caus-

ing pain or swelling behind the outer ankle and sometimes ankle instability. They may follow overuse, an inversion injury, or structural abnormalities.

## **Peroxidase**

An enzyme that uses hydrogen peroxide or another peroxide to oxidize a substrate; thyroid peroxidase is essential for thyroid-hormone synthesis.

## **Peroxisomal Disorder**

An inherited condition in which small cell structures that process certain fats do not form or work properly. It may affect the brain, nerves, liver, eyes, hearing, bones, or development.

## **Peroxisome**

A small structure inside a cell that breaks down certain fats and harmful substances. It also helps control hydrogen peroxide and makes substances needed for cell membranes and nerve function.

## **Peroxisome Proliferation**

An increase in the number or size of peroxisomes, small cell structures that process certain fats and harmful chemicals. It can be triggered by some medicines, chemicals, hormones, or metabolic conditions and may alter cell function.

## **Perphenazine**

A first-generation (typical) phenothiazine antipsychotic used for schizophrenia and, in some settings, severe nausea or vomiting. Adverse effects can include sedation, orthostatic hypotension, extrapyramidal symptoms, tardive dyskinesia, and neuroleptic malignant syndrome.

## **Perseveration**

Involuntary repetition of a word, idea, movement, or response after the appropriate stimulus has changed, associated with frontal or basal-ganglia disease, aphasia, delirium, or psychiatric illness.

## **Persistent Asthma**

Asthma symptoms or airway narrowing that occur regularly rather than only occasionally and usually



---

require ongoing controller treatment. Severity is judged by symptoms, night waking, reliever use, lung function, and activity limits; treatment is adjusted over time.

### **Persistent Cloaca**

Persistent cloaca is a congenital malformation in which the rectum, vagina, and urinary tract join into one channel, requiring specialist evaluation and reconstructive care.

### **Persistent Depressive Disorder**

A long-lasting depressive disorder with depressed mood or irritability for at least two years in adults or one year in children and adolescents. Other symptoms may include low energy, low self-esteem, sleep or appetite changes, poor concentration, or hopelessness.

### **Persistent Fetal Vasculature**

Persistence after birth of fetal blood vessels that normally regress before or soon after birth. It usually affects one eye and may cause a small eye, cataract, abnormal pupil reflection, retinal changes, or reduced vision.

### **Persistent Generalized Lymphadenopathy**

The chronic enlargement of lymph nodes measuring greater than one centimetre in diameter across two or more extra-inguinal anatomical sites persisting for at least three months, serving as a recognized clinical manifestation of chronic human immunodeficiency virus (HIV) infection.

**Alternate Names:**

PGL

### **Persistent Pulmonary Hypertension Of The Newborn**

Failure of the normal fall in pulmonary vascular resistance after birth, causing right-to-left shunting and hypoxaemia. Management requires urgent neonatal respiratory and circulatory support.

### **Persistent Pupillary Membranes**

Strands left from a temporary membrane that covered the developing eye before birth. They may

cross the pupil and are usually harmless, but dense strands can interfere with vision and occasionally need treatment.

### **Persistent Vegetative State**

A disorder of consciousness lasting more than one month following severe acute brain injury, characterized by complete wakefulness without awareness (spontaneous eye opening, sleep-wake cycles, and intact brainstem autonomic reflexes) in the total absence of purposeful, voluntary behavioral responses to sensory stimuli.

**Alternate Names:**

PVS; Unresponsive Wakefulness Syndrome

### **Personal Emergency Response System**

A wearable or nearby device that lets a person summon help during an emergency.

**Alternate Names:**

PERS; Medical Alert System

### **Personality**

Personality is the relatively enduring pattern of thoughts, emotions, motivations, and behaviors through which a person experiences and responds to the world.

### **Personality Change**

Personality change is a new or marked alteration in behavior, emotional responses, judgment, or interpersonal style, which may reflect neurologic disease, psychiatric illness, substances, or medical conditions.

### **Personality Disorder**

Alternate name; see *Personality Disorders*.

### **Personality Disorders**

A group of mental disorders involving enduring, inflexible patterns of thought, emotion, behavior, and relationships that differ markedly from the person's cultural context and cause distress or impaired functioning. Traditional groupings include Cluster A (paranoid, schizoid, and schizotypal), Cluster B (antisocial, borderline, histrionic, and narcissistic),

---

and Cluster C (avoidant, dependent, and obsessive-compulsive personality disorders).

### Personality Test

A psychological assessment using structured questions or stimuli to describe enduring traits, patterns, or possible personality pathology; results require validated instruments and clinical interpretation.

### Personalized Medicine

Healthcare that uses a person's genes, biological measurements, environment, and lifestyle to guide prevention or treatment. It can improve treatment selection but does not guarantee a response.

### Personal Protective Equipment

Clothing or equipment that protects a person from infection, chemicals, blood, or other hazards. It may include gloves, gowns, masks, respirators, goggles, or face shields; the correct items depend on the hazard and must be put on, removed, and disposed of safely.

### Person With Disability

A person with disability has a long-term physical, sensory, intellectual, or mental-health impairment that may interact with barriers to limit participation; care should be accessible and person-centered.

### Perspiration

Sweat released by skin glands. Eccrine glands mainly help cool the body, while apocrine glands contribute to odor after their secretions are changed by skin bacteria; unusually heavy sweating is called hyperhidrosis.

### Pertussis

Alternate name; see *Whooping Cough*.

### Pertussis Toxin

A toxin produced by *Bordetella pertussis* that disrupts cell signaling and contributes to the cough and systemic effects of whooping cough.

### Pertuzumab

A monoclonal antibody targeting the HER2 receptor and blocking its dimerization, used with other medicines for selected HER2-positive cancers.

### Pes

Pes means foot; it is also used in terms describing foot shape or deformity, such as pes planus or pes cavus.

### Pes Anserinus

The combined tendon insertion of the sartorius, gracilis, and semitendinosus muscles on the antero-medial proximal tibia; inflammation there is pes-anserine bursitis.

### Pes Cavus

A high-arched foot deformity characterized by an abnormally elevated medial longitudinal arch that does not flatten with weight-bearing, frequently associated with neuromuscular disorders including Charcot-Marie-Tooth disease, Friedreich ataxia, and spinal dysraphism.

#### Alternate Names:

High-Arched Foot; Talipes Cavus

### Pes Planus

Alternate name; see *Flatfoot* (*Pes Planus*).

### Pessaries

Devices placed inside the vagina to support organs that have descended, such as the bladder, uterus, or rectum. Ring, cube, and other designs suit different shapes and needs; regular removal and follow-up help prevent sores, discharge, or infection.

### Pessary

A device placed inside the vagina to support pelvic organs, treat prolapse, or sometimes prevent leakage of urine.

### Pessimism

A tendency to expect negative outcomes or interpret situations unfavorably.

---

## **Pesticide**

A substance used to control insects, weeds, fungi, or other pests; exposure can cause poisoning.

## **Pesticide Poisoning**

Toxic exposure to an insecticide, herbicide, fungicide, rodenticide, or related agricultural chemical. Clinical findings and treatment depend on the active ingredient: cholinergic toxidromes, corrosive injury, methemoglobinemia, hepatic or renal injury, seizures, shock, and coagulopathy are possible. Identify the product, decontaminate safely, and use agent-specific toxicology guidance.

## **Pesticides**

Chemicals used to kill insects, weeds, fungi, rodents, or other pests. Exposure can irritate the skin, eyes, lungs, or digestive tract; some pesticides can cause sweating, vomiting, weakness, seizures, or breathing failure.

## **Pesticides On Fruits And Vegetables**

Small pesticide residues on properly washed produce usually do not cause acute poisoning. Swallowing concentrated pesticide or developing vomiting, sweating, breathing difficulty, weakness, or confusion requires immediate poison-control or emergency advice.

## **Pestilence**

Pestilence is an old term for a severe, widespread, often fatal infectious disease outbreak; modern usage usually specifies the pathogen and epidemiologic event.

## **Pet Allergy**

An allergic reaction to proteins in an animal's skin flakes, saliva, or urine, commonly causing rhinitis, conjunctivitis, or asthma.

### **Alternate Names:**

Animal Dander Allergy; Pet Dander Hypersensitivity

## **PET/CT**

A scan that combines PET, which shows active body

processes, with CT, which shows detailed anatomy. It helps locate disease and assess selected cancers, heart conditions, and brain disorders; it uses a radioactive tracer and X-rays.

## **Petechia**

Alternate name; see *Petechiae*.

## **Petechiae**

Pinpoint, flat, non-blanching red, purple, or brown macules measuring less than two to three millimetres in diameter, caused by minor capillary hemorrhage and extravasation of erythrocytes into the skin or mucous membranes. Major causes include thrombocytopenia (such as immune thrombocytopenia, leukemia, or aplastic anemia), platelet dysfunction, life-threatening infections (meningococcemia, infective endocarditis, viral hemorrhagic fevers), vasculitis, and acute mechanical elevation of venous pressure from violent coughing or vomiting.

### **Alternate Names:**

Petechia; Pinpoint Hemorrhages

## **Petit Mal**

Older term; see *Absence Seizure*.

## **Petit Mal Seizure**

Alternate name; see *Absence seizure*.

## **PET/MRI**

A scan that combines information about tissue activity with detailed MRI images of soft tissues. It is useful for selected brain, cancer, and heart problems and does not use X-rays, but takes longer than many other imaging tests.

## **Petrissage**

A massage technique involving kneading, squeezing, or lifting soft tissues. It should be adapted or avoided when there is acute injury, infection, clot risk, or fragile skin.

## **Petrolatum**

A semisolid purified mixture of hydrocarbons used on skin as a protective ointment and moisture bar-

---

rier. It reduces water loss, shields minor irritation, and supports healing of dry, cracked, or chafed skin.

### **Petroleum Jelly Overdose**

Swallowing a small amount of petroleum jelly usually causes little harm, but larger amounts can cause diarrhea or vomiting. If it enters the lungs, it may cause inflammation and breathing problems; contact poison control after a significant exposure.

### **Petrosal Sinus**

Paired dural venous sinuses draining the cavernous sinus: the superior petrosal sinus courses along the crest of the petrous temporal bone into the transverse sinus, while the inferior petrosal sinus travels in the petro-occipital fissure through the anterior jugular foramen into the internal jugular vein.

### **Petrositis**

Inflammation or infection of the petrous portion of the temporal bone, usually arising from complicated otitis media and potentially affecting nearby cranial nerves.

### **Petrous**

Describing dense, hard bone, especially the part of the temporal bone that houses inner-ear structures.

### **PET Scan**

Nuclear medicine imaging using radioactive tracers (most commonly FDG, a glucose analog) to detect metabolically active tissues.

#### **Alternate Names:**

PET Imaging; PET-CT

### **PET Scan For Breast Cancer**

Positron-emission tomography, usually combined with computed tomography, used selectively to assess metabolically active breast cancer, staging, recurrence, or treatment response.

### **PET Scanner**

A machine that detects radiation from a tracer inside the body and creates images showing how actively tissues use substances such as glucose. It is used to

assess selected cancers and other disorders.

### **PET Tracers**

Radioactive substances used during a PET scan to highlight particular body processes. Different tracers show glucose use, bone activity, or specific cell receptors; the appropriate tracer depends on the clinical question.

### **Peutz-Jeghers's Syndrome**

An inherited condition causing dark patches around the mouth and many growths in the stomach or intestines. The growths may bleed, cause blockage, or increase the risk of digestive and other cancers.

### **Peutz-Jeghers Syndrome**

An inherited condition causing dark spots on the lips, mouth, fingers, or skin and multiple growths in the digestive tract. These growths can bleed or block the bowel and increase the risk of several cancers.

#### **Alternate Names:**

PJS

### **PEWS**

Alternate name; see *Pediatric Early Warning Score*.

### **Pexophagy**

A cell-cleaning process that removes damaged or unneeded peroxisomes, the small structures that process certain fats and harmful chemicals. The cell breaks them down and reuses some of their materials.

### **Peyer Patches**

Clusters of immune tissue in the wall of the lower small intestine. They sample material from the gut and help the immune system recognize and respond to germs while tolerating normal food and bacteria.

### **Peyer's Patches**

Organized lymphoid tissue in the mucosa and submucosa of the ileum. These aggregates sample intestinal contents and contribute to immune surveillance of the gastrointestinal tract.

---

**Alternate Names:**

Ileal Lymphoid Nodules; Aggregated Lymphoid Nodules

**Peyronie Disease**

Scar tissue forming inside the penis, causing a new bend or indentation that may make erections painful or intercourse difficult. Symptoms may stabilize or worsen; treatment ranges from observation and injections to surgery in selected severe cases.

**Peyronie's Disease**

A condition in which scar tissue forms in the penis and causes a curved or painful erection, shortening, or difficulty with sexual activity. Treatment depends on severity and symptoms.

**PFAPA Syndrome**

A recurrent fever syndrome, usually beginning in childhood, characterized by periodic fever with some combination of mouth ulcers, sore throat, and swollen neck lymph nodes; it is commonly abbreviated PFAPA.

**Pfeiffer Syndrome**

An autosomal dominant craniosynostosis syndrome caused by FGFR1 or FGFR2 mutations.

**Alternate Names:**

Classic Type Pfeiffer Syndrome Features: craniosynostosis (bilateral coronal), midface hypoplasia, proptosis, hypertelorism, broad thumbs and great toes (medial deviation), and variable syndactyly.

**Alternate Names:**

Acrocephalosyndactyly Type V

**P-Glycoprotein**

A transport protein in cell membranes that pumps many medicines and other substances out of cells. It affects how medicines are absorbed and removed and can cause interactions when its activity is blocked or increased.

**pH**

A measure of how acidic or alkaline a solution is, usually on a scale from 0 to 14. Body fluids have

tightly controlled pH ranges, and major changes can be dangerous.

**Phacoemulsification**

The standard modern microsurgical technique for cataract extraction in which an ultrasonic handpiece vibrating at high frequencies (28–40 kHz) emulsifies (fragments and liquefies) the cataractous crystalline lens nucleus and cortex, which are simultaneously aspirated while balanced salt solution irrigates the anterior chamber to maintain intraocular pressure and cool the tip. Performed through a micro-incision (typically 1.8–2.8 mm) via a continuous curvilinear capsulorhexis (CCC), it preserves the crystalline lens capsule (capsular bag) to allow immediate in-the-bag implantation of a foldable artificial intraocular lens (IOL). Compared with traditional extracapsular cataract extraction (ECCE), phacoemulsification offers self-sealing sutureless incisions, minimal surgically induced astigmatism, rapid visual recovery, and lower complication rates. Key surgical complications include posterior capsule rupture (PCR), vitreous loss, dropped nucleus, and corneal endothelial decompensation.

**Alternate Names:**

Phaco; Ultrasonic Cataract Extraction; Small-Incision Cataract Surgery; Phacoemulsification with IOL Implantation

**Phacomatosis**

A broad group of congenital hereditary neurocutaneous syndromes characterized by widespread hamartomatous developmental lesions simultaneously involving structures of ectodermal origin (central nervous system, retina, and skin), including neurofibromatosis, tuberous sclerosis, and von Hippel-Lindau disease.

**Alternate Names:**

Neurocutaneous Syndrome

**Phagocyte**

An immune cell that surrounds, swallows, and breaks down germs, dead cells, or foreign particles. Phagocytes help prevent infection, remove damaged material, and alert other immune cells to danger.

---

## Phagocytes

Immune cells that surround, swallow, and break down germs, dead cells, and foreign particles. They help prevent infection, clear damaged material, and signal other immune cells during inflammation and tissue repair.

## Phagocytic Vesicle

A small membrane-bound compartment formed inside a cell after it surrounds and swallows a germ, dead cell, or particle. It later joins a digestive compartment so the material can be broken down.

## Phagocytosis

The process by which a cell surrounds, takes in, and breaks down a particle or microorganism. Immune cells use it to remove germs, dead cells, and harmful debris from tissues.

## Phagolysosome

A compartment formed when an immune cell's swallowed-particle compartment joins a digestive compartment. Enzymes and other chemicals inside it destroy germs and break down cellular debris.

## Phagophore

A temporary membrane that grows around damaged or unnecessary material inside a cell during recycling. It closes around the material so the cell can break it down and reuse some of its parts.

## Phalangeal Fractures

Breaks in the bones of the fingers or toes, usually caused by crushing, direct impact, or twisting. Pain, swelling, bruising, deformity, and difficulty moving the digit may occur; alignment and nail-bed injury should be assessed.

## Phalanges

The bones of the fingers and toes. The thumb and big toe each have two, while the other digits usually have three. Fractures, arthritis, infection, or nerve problems in these bones can cause pain, swelling, stiffness, or difficulty using the hand or foot.

## Phalanx

One of the bones of a finger or toe. Each phalanx has a base, shaft, and end; the end bone supports the fingertip or toenail. The thumb and big toe have two phalanges, while other digits usually have three.

## Phalanx Unspecified

A phalanx is one of the bones of a finger or toe; “unspecified” indicates that the particular digit or proximal, middle, or distal segment is not identified.

## Phalen Maneuver

A diagnostic physical examination provocative test for carpal tunnel syndrome; the patient rests both elbows on a flat surface and maintains full passive wrist flexion (at 90 degrees) by pressing the dorsal surfaces of the hands together for 60 seconds; reproduction of paresthesias or pain in the median nerve distribution indicates a positive test.

### Alternate Names:

Phalen's Test; Wrist Flexion Test

## Phalen Sign

Tingling or numbness in the fingers after pressing the backs of the hands together with wrists bent down for one minute, used to check for carpal tunnel syndrome.

## Phalen Test

A test in which the wrists are bent to see whether numbness or tingling develops. Reproduced symptoms may support carpal tunnel syndrome, but the result must be interpreted with examination and other tests.

## Phantom

A test object made to resemble human tissue or organs so medical imaging equipment can be checked and calibrated. Different phantoms test image quality, measurements, or specific body regions.

## Phantom Limb

The feeling that an amputated body part is still present. Sensations may include position, move-



---

ment, itching, pressure, or pain and often change over time after amputation.

**Alternate Names:**

Phantom Limb Sensation; Phantom Limb Pain

**Phantom Limb Pain**

Painful sensations perceived in a limb that has been amputated, caused by persistent or reorganized activity in peripheral nerves and central sensory pathways. It may feel burning, shooting, cramping, or electric.

**Phantom Pain**

Pain perceived in a body part that has been amputated, caused by changes in peripheral nerves and central pain-processing pathways.

**Phantom Pregnancy**

A false belief or physical state in which a person has pregnancy-like symptoms without an ongoing pregnancy. Symptoms can include abdominal enlargement, breast changes, nausea, or missed periods and require medical evaluation.

**Phantosmia**

Perception of a smell when no corresponding odor is present. It may occur with sinonasal disease, migraine, seizures involving olfactory networks, head injury, neurodegenerative disease, infection, or other neurological conditions.

**Pharmaceutical Adjuvant**

A pharmaceutical adjuvant is an inactive or auxiliary ingredient added to improve a medicine's stability, absorption, preservation, delivery, or therapeutic effect.

**Pharmaceutical Excipient**

A pharmaceutical excipient is a non-active ingredient in a dosage form, such as a filler, binder, coating, solvent, preservative, or disintegrant.

**Pharmaceutical Solutions**

Liquid preparations in which one or more active pharmaceutical ingredients are dissolved in a suit-

able vehicle for administration, with concentration, sterility, stability, and route determining safe use.

**Pharmacist**

A medicine expert who prepares and dispenses drugs, checks safety, and explains correct use and possible side effects.

**Pharmacodynamic Drug Interaction**

An interaction in which two medicines have additive, opposing, or otherwise changed effects on the same body system, even when their blood concentrations are not altered.

**Pharmacodynamics**

The study of the biochemical and physiological effects of drugs on the body and their mechanisms of action, describing what the drug does to the body.

**Pharmacoepidemiology**

The study of how medicines are used and affect health in large populations. It helps assess effectiveness, safety, adverse effects, and patterns of prescribing.

**Pharmacogenetics**

Pharmacogenetics studies how inherited genetic variation alters a person's response to medicines, including efficacy, metabolism, and adverse-effect risk, to support individualized prescribing.

**Pharmacogenetic Testing**

Testing for genetic variants that influence a person's response, metabolism, efficacy, or toxicity of particular medicines and may support individualized prescribing.

**Pharmacogenomics**

The study of how genetic variations influence individual responses to drugs, including drug efficacy, toxicity, and metabolism. Genetic polymorphisms in drug-metabolizing enzymes, transporters, and receptors can lead to significant inter-individual variability in drug response.

---

### Pharmacogenomic Testing

Testing that examines genetic variants affecting drug metabolism, transport, targets, or adverse reactions to help select or dose certain medicines.

### Pharmacokinetic Drug Interaction

An interaction in which one medicine changes another medicine's absorption, distribution, metabolism, or elimination, altering its concentration or duration of action.

### Pharmacokinetics

The study of what the body does to a medicine, including its absorption, distribution, metabolism, and elimination. It helps explain drug concentrations over time and supports dose selection.

### Pharmacological Stress Test

A heart stress test using medicine to imitate exercise when a person cannot exercise adequately. Imaging or heart recordings show blood flow and function, while possible effects include chest discomfort, breathlessness, or abnormal rhythm.

### Pharmacologic Substance

A pharmacologic substance is a chemical that produces a biological effect by interacting with cells, receptors, enzymes, transporters, or other targets.

### Pharmacologist

A pharmacologist studies how medicines and chemicals act in living systems, including mechanisms, metabolism, toxicity, and therapeutic effects.

### Pharmacology

The study of medicines and other drugs, including how they work, how the body handles them, their uses, side effects, and interactions. This knowledge helps guide safe choice, dosing, testing, and monitoring of treatment.

### Pharmacopoeia

A pharmacopoeia is an official reference specifying standards for identity, purity, strength, testing, and preparation of medicines.

### Pharmacoproteomics

The study of how a person's protein patterns influence medicine response, drug effects, or toxicity. It is an emerging part of precision medicine.

### Pharmacotherapy

Pharmacotherapy is treatment with medicines chosen to prevent, relieve, or control a disease or symptom.

### Pharmacovigilance

The science and activities relating to the detection, assessment, understanding, and prevention of adverse drug reactions and other drug-related problems.

### Pharmacy

The healthcare field concerned with preparing, checking, supplying, and explaining medicines. Pharmacists help choose safe doses, identify interactions and allergies, support treatment decisions, and teach people how to use medicines correctly.

### Pharmacy Facility

A pharmacy facility is an authorized location where medicines may be stored, prepared, dispensed, monitored, and supplied under applicable professional and safety standards.

### Phytotherapy

A treatment approach using medicinal plants or plant-derived preparations. Safety and effectiveness vary by product and condition; interactions, contamination, incorrect dosing, and delayed effective care are possible.

### Pharyngeal

Relating to the pharynx, or throat. The pharynx connects the nose and mouth with the voice box, food pipe, and breathing passages. Infection, swelling, narrowing, or a tumor there may cause sore throat, swallowing difficulty, noisy breathing, or voice changes.

---

## Pharyngeal Carcinoma

Pharyngeal carcinoma is cancer arising in the throat and may cause persistent sore throat, swallowing difficulty, hoarseness, or a neck mass.

## Pharyngeal Disease

A disorder affecting the throat, including infection, inflammation, blockage, injury, muscle or nerve problems, or a tumor. Symptoms may include sore throat, swallowing difficulty, choking, voice changes, or breathing problems.

## Pharyngeal Hemorrhage

Bleeding from the pharynx caused by trauma, surgery, infection, vascular lesions, tumor, or severe mucosal disease and potentially threatening the airway.

## Pharyngeal Mucositis

Inflammation and ulceration of the throat lining, commonly caused by radiation, chemotherapy, infection, trauma, or medications and producing pain, difficulty swallowing, and reduced intake.

## Pharyngeal Stenosis

Narrowing of the pharyngeal passage caused by scarring, congenital abnormality, tumor, inflammation, or surgery, which may impair swallowing or breathing.

## Pharyngeal Tonsil

Adenoid tissue in the roof of the throat behind the nose that helps immune defense, especially in children.

## Pharyngectomy

Surgery to remove part or all of the throat, usually to treat cancer. Depending on the extent, reconstruction or a temporary feeding tube may be needed, and treatment can affect swallowing, speech, breathing, or appearance.

## Pharyngitis

Acute inflammation of the mucous membranes of the pharynx, commonly producing sore throat,

odynophagia, pharyngeal erythema, and cervical lymphadenopathy. Caused primarily by respiratory viruses or Group A *Streptococcus*.

### Alternate Names:

Sore Throat; Acute Pharyngitis; Pharyngeal Inflammation

## Pharyngolaryngeal Pain

Pain involving the pharynx and larynx, commonly associated with infection, inflammation, irritation, trauma, reflux, or malignancy.

## Pharyngomaxillary Space Abscess

A pharyngomaxillary space abscess is a deep infection containing pus near the pharynx and maxilla that can cause severe pain, swelling, difficulty swallowing, and airway risk.

## Pharynx

The muscular passage behind the nose and mouth that carries air toward the voice box and food toward the esophagus. It helps with breathing, swallowing, and speech.

## Pharynx Necrosis

Death of tissue in the throat caused by severe infection, ischemia, radiation, caustic injury, or tumor, with risk of deep infection, bleeding, airway compromise, and sepsis.

## Pharynx Neoplasm

A pharynx neoplasm is an abnormal growth in the throat that may be benign or malignant.

## Phase-Contrast Microscopy

A microscope technique that makes transparent cells and other specimens easier to see by converting small differences in light passing through them into visible contrast. It is useful for examining living, unstained cells.

## Phase II Metabolism

A stage of drug breakdown in which the body attaches another chemical group to a medicine or its earlier breakdown product. This usually makes the

---

substance easier to remove in urine or bile.

### Phase I Metabolism

An early step in the body's breakdown of many medicines. Enzymes, mainly in the liver, chemically change a medicine by adding or exposing a reactive part; this may inactivate it, activate it, or prepare it for further breakdown.

### Phe

Phe is the standard abbreviation for phenylalanine, an essential amino acid incorporated into proteins and metabolized through a pathway disrupted in phenylketonuria.

### Phelan-McDermid Syndrome

A genetic disorder that commonly causes low muscle tone, delayed development, intellectual disability, very limited speech, and autism-related behaviors. Some people have reduced sensitivity to pain, seizures, feeding problems, or kidney and heart abnormalities. It usually results from loss of genetic material near the end of chromosome 22 or a change in the *SHANK3* gene.

#### Alternate Names:

22q13.3 Deletion Syndrome

### Phenacetin

Phenacetin is a former analgesic and fever-reducing drug withdrawn or restricted because long-term use can cause kidney damage and increase cancer risk.

### Phenazopyridine

Phenazopyridine is a urinary analgesic that temporarily relieves dysuria without treating its cause; it can turn urine orange and prolonged or excessive use may cause haemolysis or methemoglobinemia.

### Phencyclidine

A dissociative drug that can cause altered perception, agitation, hallucinations, impaired coordination, or dangerous behavior.

### Phencyclidine Overdose

A dangerous amount of the dissociative drug phen-

cyclidine can cause agitation, hallucinations, high body temperature, high blood pressure, seizures, loss of coordination, coma, or injury. Severe or unusual behavior requires emergency care.

### Phencyclidine Substance Use

Phencyclidine is a dissociative drug that can cause agitation, hallucinations, loss of coordination, high blood pressure, seizures, dangerous behavior, or coma.

### Phenelzine

An antidepressant that blocks monoamine oxidase, an enzyme breaking down several brain chemicals. It has important interactions with medicines and tyramine-rich foods and can cause dangerous high blood pressure if precautions are ignored.

### Phenethylamines

Phenethylamines are a chemical family including neurotransmitters, medicines, and psychoactive substances that can affect adrenergic, dopaminergic, or serotonergic signaling.

### Phenformin

Phenformin is a former biguanide diabetes medicine withdrawn because it can cause life-threatening lactic acidosis.

### Phenindione

An older oral anticoagulant that interferes with vitamin-K-dependent clotting factors.

### Pheniramine Overdose

Taking too much pheniramine, an antihistamine, can cause extreme sleepiness or agitation, dry mouth, large pupils, fast heartbeat, seizures, abnormal rhythms, or coma. Immediate poison-control or emergency advice is needed.

### Phenobarbital

Phenobarbital is a long-acting barbiturate anti-seizure medicine that enhances GABA-A receptor inhibition; sedation, respiratory depression, cognitive effects, dependence, and enzyme-inducing drug

---

interactions are important risks.

### **Phenobarbital Overdose**

Taking too much phenobarbital can cause severe drowsiness, poor coordination, low blood pressure, slowed breathing, coma, or death. Emergency treatment is required, especially if alcohol or another sedative was also taken.

### **Phenobarbitone**

The British name for phenobarbital, a barbiturate anticonvulsant and sedative.

### **Phenol**

A corrosive aromatic compound used in chemical and laboratory applications and in some medical preparations.

### **Phenols**

A group of aromatic chemicals that can burn skin and mucous membranes and cause systemic poisoning if absorbed or swallowed.

### **Phenomenon**

An observable event, sign, or pattern in the body. Examples include fingers changing color in Raynaud phenomenon, early-morning blood-sugar rises in the dawn phenomenon, and upward eye movement when the eyelids close in Bell phenomenon.

### **Phenothiazine**

Phenothiazine is a chemical class that includes several antipsychotic and antiemetic medicines, with possible sedation, movement, and cardiac adverse effects.

### **Phenothiazine Overdose**

Taking too much of a phenothiazine medicine can cause sleepiness, confusion, low blood pressure, seizures, abnormal heart rhythms, or severe muscle stiffness and fever. It requires immediate medical assessment.

### **Phenothiazines**

A class of medicines that includes antipsychotics

and antiemetics. They can cause sedation, low blood pressure, movement disorders, heart-rhythm changes, and other adverse effects.

### **Phenotype**

The observable features of a person, such as appearance, growth, body functions, and behavior. These features result from the interaction between inherited genetic information and the environment.

### **Phenoxybenzamine**

A long-acting alpha-blocker used mainly before surgery for a catecholamine-secreting tumor such as pheochromocytoma. It can cause dizziness, low blood pressure, rapid heartbeat, and nasal congestion.

### **Phenoxymethylpenicillin**

Penicillin V, an oral penicillin used for selected susceptible infections.

### **Phenprocoumon**

Phenprocoumon is a long-acting vitamin-K antagonist anticoagulant that reduces clotting-factor production and requires coagulation monitoring.

### **Phermine**

A stimulant medicine that reduces appetite and may be used short term with diet and activity changes for weight management. It can raise heart rate and blood pressure and may cause insomnia, anxiety, dependence, or dangerous interactions with some medicines.

### **Phenthoate**

Phenthoate is an organophosphate insecticide that can inhibit acetylcholinesterase and cause cholinergic poisoning after exposure.

### **Phentolamine**

Phentolamine is a short-acting alpha-adrenergic blocker used for catecholamine-related hypertension and to limit tissue injury after certain vasoconstrictor leaks.

---

### Phenylacetate

A salt or ester of phenylacetic acid used in some metabolic and experimental medical applications; it can bind nitrogen-containing metabolites.

### Phenylalanine

An essential amino acid incorporated into proteins and converted to tyrosine; impaired metabolism causes phenylketonuria and requires dietary or medical management.

### Phenylbutazone

An older anti-inflammatory pain medicine whose use is limited by serious toxicity. It can cause stomach bleeding, kidney or liver injury, fluid retention, blood-cell problems, allergic reactions, and other dangerous effects.

### Phenylbutyrate

A nitrogen-scavenging medicine converted to phenylacetate, which combines with glutamine and facilitates nitrogen excretion in selected urea-cycle disorders.

### Phenylephrine

A medicine that narrows blood vessels. It may reduce a blocked nose, raise dangerously low blood pressure in hospital care, or widen the pupil in an eye examination; it can worsen high blood pressure, heart problems, or urinary difficulty in some people.

### Phenylketonuria

An autosomal recessive metabolic disorder, usually caused by disease-causing variants in the *PAH* gene that reduce phenylalanine hydroxylase activity. Phenylalanine accumulates in the blood and tissues, while tyrosine availability may decrease. Untreated or poorly controlled disease can affect brain development and may be associated with intellectual disability, seizures, eczema, behavioral or psychiatric symptoms, and a musty odor. Severity ranges from classic to milder forms, and the disorder is commonly identified through newborn screening.

#### Alternate Names:

PKU; Phenylalanine Hydroxylase Deficiency

### Phenylpropanolamine

A sympathomimetic decongestant formerly used in some products; it can raise blood pressure and has been restricted because of safety concerns.

### Phenytoin

A medicine used to prevent or control some seizures. It reduces repeated electrical activity in brain cells; levels in the blood may need monitoring because interactions and side effects can include dizziness, poor coordination, gum overgrowth, skin reactions, or serious toxicity.

### Phenytoin Overdose

Too much phenytoin can cause nausea, severe dizziness, double vision, involuntary eye movements, poor coordination, confusion, abnormal heart rhythms, or coma. Significant or intentional overdoses can be life-threatening.

### Phenytoin Sodium

The sodium salt of phenytoin, an anticonvulsant used for selected focal and generalized tonic-clonic seizures; toxicity can cause nystagmus, ataxia, arrhythmia, or severe skin reactions.

### Phenytoin Toxicity

Dose-related toxicity causing horizontal nystagmus, ataxia, dysarthria, lethargy, confusion, or coma; severe overdose can cause hypotension, dysrhythmias, or seizures, particularly after rapid intravenous administration. Serial serum levels, ECG, and neurologic assessment guide supportive care; there is no routine specific antidote.

### Pheochromocytoma

A rare tumor, usually in an adrenal gland, that may release excess adrenaline-like hormones. It can cause episodes of high blood pressure, headache, sweating, a pounding heartbeat, shakiness, or paleness.

#### Alternate Names:

Chromaffin Cell Tumor; Adrenal Medullary Tumor; Pheo



---

## **Pheresis**

A procedure that removes a selected part of blood, such as plasma, platelets, or white blood cells, and returns the remaining blood. It may treat disease or collect a needed blood component from a donor.

## **Pheromone**

A pheromone is a chemical signal released by an organism that alters behavior or physiology in another member of the same species.

## **Phialophora**

A genus of darkly pigmented fungi that can cause chromoblastomycosis or other subcutaneous and opportunistic infections.

## **Phialophora Verrucosa**

A dark-pigmented fungus that commonly causes chromoblastomycosis, a long-lasting infection of skin and tissue beneath it. Infection usually follows implantation through injured skin and produces slowly enlarging warty or raised lesions.

## **Philadelphia Chromosome**

A rearrangement of genetic material between chromosomes 9 and 22 that creates the BCR::ABL1 gene. It is found in most cases of chronic myeloid leukemia and some acute leukemias and helps guide treatment.

## **Philodendron Poisoning**

Chewing philodendron leaves releases irritating crystals that can cause immediate burning and swelling of the mouth, tongue, or throat. Breathing or swallowing difficulty may occur.

## **Phimosis**

Tightness of the foreskin that prevents it from being pulled back over the head of the penis. It can be normal in young children, but pain, repeated infection, scarring, or difficulty passing urine may indicate a problem.

## **Phimosis and Paraphimosis**

Alternate name; see *Penis Disorders*.

## **Phlebitis**

Inflammation of a vein causing pain, warmth, redness, or tenderness along its course. It may follow irritation from an intravenous line or occur with a clot in a surface vein; sudden limb swelling or breathlessness may indicate a more serious clot.

## **Phlebitis And Thrombophlebitis**

Phlebitis is inflammation of a vein, while thrombophlebitis includes a clot within the inflamed vein. Superficial disease causes localized pain and redness; deep-vein thrombosis can embolize to the lungs and is suspected with limb swelling, pain, or breathlessness.

## **Phlebography**

An imaging test in which contrast dye is injected into a vein to show the veins on X-ray. It can detect a blood clot or an abnormal vein but is now often replaced by ultrasound.

## **Phlebotomy**

Taking blood from a vein, usually with a needle, for testing, donation, or treatment. The procedure may cause brief pain, bruising, bleeding, fainting, or rarely infection; trained staff use safe collection and labeling practices.

### **Alternate Names:**

Venipuncture

## **Phlebovirus**

A genus of segmented RNA viruses, many transmitted by sandflies or mosquitoes; some cause fever, encephalitis, or hemorrhagic disease.

## **Phlegm**

Thick mucus brought up by coughing from the lower airways. Its amount and appearance can change with infection, asthma, smoking, chronic lung disease, or other conditions.

## **Phlegmasia Alba Dolens**

Painful, marked swelling and pallor of a limb caused by extensive deep-vein thrombosis, usually involving major venous outflow. It can threaten the limb and

---

circulation.

**Alternate Names:**

Milk Leg; Femoral Thrombophlebitis

**Phlegmasia Cerulea Dolens**

Phlegmasia cerulea dolens is a severe, extensive deep-vein thrombosis that obstructs venous outflow, causing painful swollen blue discoloration and risking limb ischemia or pulmonary embolism.

**PH Meter**

An instrument that measures acidity or alkalinity as pH in a solution or specimen; clinical applications include gastric, urinary, laboratory, and environmental measurements.

**PH Monitoring**

Measurement of acidity, usually in the esophagus, over time with a small sensor. It can help detect and measure acid reflux when symptoms or endoscopy do not give a clear answer.

**Phobia**

An excessive, persistent fear of a specific object or situation that leads to avoidance or distress.

**Phobias**

Persistent, excessive fear of a specific object or situation that leads to avoidance or significant distress.

**Phobic Disorder**

A disorder characterized by clinically significant fear and avoidance of a particular object or situation.

**Phocomelia**

A rare congenital limb-development defect in which hands or feet are attached close to the trunk because of absent or severely shortened long bones.

**Phonation**

Production of voiced sound when exhaled air makes the vocal folds vibrate in the larynx; disorders cause hoarseness, weak voice, or loss of voice.

**Phonocardiogram**

A graphic recording of heart sounds and murmurs, traditionally made with a microphone over the chest. It can document timing and duration but is usually interpreted with examination and echocardiography.

**Phonocardiography**

Recording heart sounds and murmurs with a microphone, producing a tracing that can help evaluate heart valve problems.

**Phonological Disorder**

A communication disorder involving persistent difficulty producing or organizing speech sounds beyond what is expected for age and language development.

**Phonophobia**

Unusual sensitivity to sound or distress caused by ordinary sounds. It often occurs with migraine but may also accompany concussion, infection, anxiety, or other conditions; new severe sound sensitivity with headache or neurological symptoms needs assessment.

**Phorate**

Phorate is a highly toxic organophosphate pesticide whose exposure can produce excessive cholinergic activity and medical emergency.

**Phorbols**

Phorbols are diterpene compounds related to phorbol esters that can activate cellular protein kinase C signaling.

**Phorometer**

An instrument used to measure ocular alignment and the amount of prism correction needed for binocular vision disorders.

**Phosgene**

A highly toxic industrial gas that reacts with moist respiratory tissues and can cause delayed pulmonary edema, cough, chest tightness, and respiratory failure after exposure.

---

## Phosgene Oxime

A highly corrosive chemical warfare agent that causes immediate severe pain, skin and eye injury, airway damage, and tissue necrosis on contact.

## Phosmet

Phosmet is an organophosphate insecticide that can inhibit acetylcholinesterase and cause cholinergic toxicity after substantial exposure.

## Phosphamidon

A poisonous organophosphate insecticide. Exposure can cause sweating, excess saliva, vomiting, diarrhea, small pupils, muscle twitching, breathing difficulty, seizures, or weakness because it disrupts nerve signals; suspected poisoning needs urgent medical help.

## Phosphate

An inorganic ion and component of nucleotides, phospholipids, proteins, and bone mineral. It participates in ATP energy transfer, phosphorylation-dependent signaling, intracellular buffering, and acid excretion by the kidney.

## Phosphate Binders

Medicines taken with meals that bind phosphate in food inside the intestine, reducing its absorption. They are used mainly in advanced kidney disease when blood phosphate is high. The choice depends on calcium levels, vascular calcification, other illnesses, and treatment response.

## Phosphate Homeostasis

The body keeps blood phosphate within a useful range through the coordinated actions of the parathyroid glands, vitamin D, kidneys, intestines, and bones. Kidney disease can disrupt this balance.

## Phosphatidic Acid

A membrane phospholipid and signaling intermediate that serves as a precursor for other glycerolipids and influences membrane curvature, trafficking, and cell growth.

## Phosphatidylcholine

A major membrane phospholipid and component of pulmonary surfactant, lipoproteins, and bile; it also participates in cell signaling.

## Phosphatidylethanolamines

Membrane phospholipids containing ethanolamine that help determine membrane structure, fusion, and cell signaling. A group of cell-membrane fats that help maintain membrane shape, join membranes, and support cell signaling.

## Phosphatidylinositol

A membrane phospholipid that serves as a precursor for signaling molecules and helps anchor proteins to cell membranes.

## Phosphatidylserine

A membrane phospholipid normally concentrated on the inner cell surface that becomes exposed during apoptosis and helps signal cell removal.

## Phosphatidylserines

Molecules of phosphatidylserine, a membrane phospholipid involved in apoptosis, coagulation, membrane structure, and cell signaling.

## Phosphaturic Mesenchymal Tumor

A rare tumor that releases a substance causing the kidneys to lose too much phosphate. Low phosphate can soften bones and cause bone pain, weakness, or fractures; removing the tumor often corrects the problem.

## Phosphine

A highly toxic gas used in some industrial or pest-control settings that can cause severe lung, heart, liver, kidney, and neurologic injury.

## Phosphodiesterase-5 Inhibitors

Medicines that block an enzyme called PDE-5, increasing a chemical signal that relaxes blood vessels. They are used for erectile dysfunction and some forms of high blood pressure in the lung; nitrates must not be combined with them.

---

### Phosphofructokinase-1

A key enzyme that controls how cells break down sugar to produce energy, especially when energy demand increases.

### Phospholipase C Pathway

A system cells use to turn some hormone or nerve signals at their surface into internal actions. It can change calcium levels and control contraction, secretion, growth, or other cell responses.

### Phospholipid

A type of lipid with a water-attracting phosphate-containing head and water-repelling fatty-acid tails. Phospholipids form the basic double-layered structure of cell membranes and also help with cell signaling and transport.

### Phosphorus

An essential element in bone, nucleic acids, phospholipids, and ATP.

### Phosphorus Blood Test

A blood test measuring inorganic phosphate, interpreted with calcium, parathyroid hormone, vitamin D, and kidney function to evaluate mineral and renal disorders.

### Phosphorus In Diet

Dietary phosphorus comes from protein-rich foods and many processed foods; people with kidney disease may need individualized limits.

### Phosphorylation

Addition of a phosphate group to a molecule, often regulating protein activity or energy transfer.

### Photoallergy

An immune-mediated skin reaction that occurs when ultraviolet light activates a sensitizing substance on or in the skin, causing an eczematous rash.

### Photocoagulation

Treatment that uses intense light, usually a laser, to

heat and seal targeted tissue. In the eye, it can close leaking or abnormal blood vessels and limit damage from conditions such as diabetic retinopathy.

### Photodermatosis

A skin disorder caused or aggravated by exposure to ultraviolet or visible light, including polymorphous light eruption, phototoxic reactions, and photoallergic reactions.

### Photodynamic Therapy

A treatment that combines a photosensitizing agent with light to generate reactive oxygen species and destroy cancer cells, used for actinic keratosis, superficial skin cancers, and selected visceral tumors.

### Photodynamic Therapy For Cancer

Treatment in which a light-sensitive medicine collects in selected abnormal cells and is activated by a specific light, producing substances that destroy nearby cancer cells. It is used for certain accessible cancers and may cause pain, swelling, or temporary light sensitivity.

### Photoelectric Effect

An interaction in which tissue absorbs an X-ray and releases an electron. It contributes to how much radiation is absorbed and helps create contrast between tissues on an X-ray image.

### Photographic Fixative Poisoning

Harm from swallowing or inhaling chemicals used to process photographs. Effects depend on the product but may include irritation, vomiting, breathing problems, or chemical burns; keep the container and contact poison control.

### Photomultiplier Tube

A sensitive device that converts tiny flashes of light into electrical signals. In some medical scanners, groups of these devices help detect and locate radiation emitted by a tracer.

### Photopeak

The part of a radiation-measurement graph show-

---

ing photons that deposited their full energy in the detector. Clinicians use it to select the energy range for interpreting some nuclear medicine scans.

### **Photopheresis**

A treatment in which some blood cells are removed, mixed with a light-sensitive medicine, exposed to ultraviolet light, and returned to the body. It is used for selected immune, blood, and skin disorders under specialist care.

### **Photophobia**

Unusual discomfort or pain caused by light, sometimes leading a person to avoid bright surroundings. It may occur with migraine, corneal injury, eye inflammation, infection, concussion, or meningitis. Severe eye pain, vision loss, fever, stiff neck, or a sudden severe headache needs urgent care.

### **Photoplethysmography**

A noninvasive optical method that detects blood-volume changes in tissue. It is used in devices such as pulse oximeters and some heart-rate monitors.

### **Photoreceptors**

Specialized cells in the retina that detect light. Rods support dim-light and peripheral vision, while cones provide color and sharp central vision; damage can reduce sight or cause blind areas.

### **Photorefractive Keratectomy**

Laser surgery that removes a very thin surface layer of the cornea and reshapes the layer beneath it. It can reduce short-sightedness, long-sightedness, or astigmatism, but recovery is slower than some procedures and pain, haze, infection, or imperfect vision can occur.

### **Photosensitivity**

An unusually strong reaction of the skin to sunlight, causing redness, burning, itching, blistering, or a rash on exposed areas. It may result from a disease, inherited tendency, medicine, or chemical.

### **Photosensitization**

Increased sensitivity to ultraviolet or visible light caused by a medication, chemical, or disease, producing an exaggerated skin reaction.

### **Photosensitizing Drugs**

Alternate name; see *Photosensitivity to drugs*.

### **Phototherapy**

Treatment using carefully controlled light for conditions such as newborn jaundice, psoriasis, eczema, or seasonal depression. The wavelength, dose, duration, and eye or skin protection depend on the condition treated.

### **Phototoxic Dermatitis**

A sunburn-like skin injury caused when a medicine or chemical makes the skin react strongly to ultraviolet light. It affects exposed areas and may cause redness, pain, swelling, blisters, or later darkening.

### **Phototoxicity**

Skin injury occurring when a chemical makes skin unusually sensitive to ultraviolet light. It resembles severe sunburn and may cause redness, pain, blistering, or later darkening after medicine or chemical exposure.

### **Phrenic**

Relating to the diaphragm or the phrenic nerves, which arise from the neck and carry the main motor supply to the diaphragm for breathing.

### **Phrenic Nerve**

The nerve arising mainly from C3-C5 that supplies the diaphragm and carries sensory fibers from nearby structures.

### **Phrenic Nerve Palsy**

Paresis or complete paralysis of the phrenic nerve (arising from cervical roots C3, C4, C5), leading to denervation, elevated position, and paradoxical movement of the ipsilateral hemidiaphragm during inspiration. Common etiologies include mediastinal malignancy (bronchogenic carcinoma), thoracic

---

surgical trauma (cardiac or pulmonary resections), cervical spine trauma, and severe neuralgic amyotrophy.

**Alternate Names:**

Diaphragmatic Hemiparalysis; Phrenic Nerve Injury

**PHT**

Alternate name; see *Portal hypertension*.

**Phthiriasis**

Infestation with lice, especially pubic lice, causing itching and sometimes visible lice or eggs attached to hair. Treatment usually includes an approved lice-killing product, washing or sealing recently used clothing and bedding, and evaluation or treatment of close or sexual contacts.

**Phyllodes Tumor**

A rare breast tumor that can be benign, borderline, or cancerous. It often appears as a painless lump that grows quickly. Treatment usually involves removing the tumor with a safe margin; the operation depends on the tumor's size, type, and location.

**Alternate Names:**

Phyllodes Tumor of the Breast; Cystosarcoma Phyllodes; Periductal Fibroadenoma

**Phyllodes Tumors**

Alternate name; see *Phyllodes Tumor*.

**Physiatrist**

A doctor who specializes in rehabilitation and in improving movement, function, independence, and quality of life after illness, injury, or disability.

**Physical Activity**

Any movement produced by skeletal muscles that increases energy expenditure, including occupational, household, transportation, and recreational movement. Regular activity supports cardiovascular, metabolic, musculoskeletal, and mental health.

**Physical Child Abuse**

Physical child abuse is intentional physical injury or harmful force inflicted on a child, requiring im-

mediate safety assessment, medical care, and safeguarding action.

**Physical Dependence**

A state in which the body adapts to a substance or medication and develops withdrawal symptoms when it is reduced or stopped.

**Physical Exam Frequency**

How often a person should receive a physical examination. The interval depends on age, symptoms, chronic conditions, medicines, pregnancy, and preventive-care recommendations rather than one schedule for everyone.

**Physical Examination**

A systematic assessment of the body using inspection, palpation, percussion, auscultation, and other examination methods.

**Physical Half-Life**

The time needed for half the radioactive atoms in a substance to change through radioactive decay. It describes the substance itself and does not include how quickly the body removes it.

**Physical Medicine**

The medical specialty focused on function, rehabilitation, pain, and restoration of independence after illness or injury.

**Physical Therapy**

Alternate name; see *Physiotherapy*.

**Physical Therapy Assessment**

A detailed evaluation of movement, strength, flexibility, sensation, balance, walking, pain, and daily activities. The findings guide treatment goals, exercise choices, equipment needs, safety advice, and progress monitoring.

**Physician Assistant**

A nationally certified and state-licensed medical clinician who practices medicine collaboratively with or under the supervision of physicians. PAs obtain



---

medical histories, perform physical examinations, order and interpret diagnostic studies, formulate medical diagnoses, prescribe pharmacotherapy, assist in surgery, and perform clinical procedures across primary care and surgical specialties.

### Physicians' Desk Reference

A reference source containing prescribing information, indications, contraindications, dosing, warnings, and adverse effects for medicines and selected products.

### Physiologic

Physiologic means relating to normal body function or a process occurring within the expected biological range.

### Physiological Dead Space

The part of a breath that does not exchange oxygen and carbon dioxide with blood. It includes the air passages and lung areas that receive little or no blood flow.

### Physiologic Jaundice Of Newborn

A common temporary yellowing of a newborn's skin and eyes that begins after the first day of life. It results from normal newborn red-cell breakdown and immature liver processing of bilirubin; jaundice that starts earlier or becomes severe needs urgent assessment.

### Physiologic Jaundice Of The Newborn

Physiologic newborn jaundice is a temporary bilirubin rise after birth caused by immature bilirubin processing, but early or severe jaundice needs urgent evaluation.

### Physiology

The study of how healthy body parts, organs, and living organisms normally work together. It examines processes such as breathing, circulation, digestion, movement, hormone action, nerve signaling, and maintaining internal balance.

### Physiotherapy

A healthcare profession dedicated to restoring, maintaining, and maximizing movement, physical strength, and functional independence in people affected by injury, illness, surgery, or disability. Treatment is tailored to individual needs and utilizes therapeutic exercise (strengthening, stretching, and aerobic conditioning), hands-on manual therapy (joint mobilization and soft-tissue release), gait and balance retraining, and physical modalities (heat, cold, ultrasound, and electrical stimulation). Specialized branches include orthopedic rehabilitation (e.g., post-fracture and joint replacement recovery), neurological rehabilitation (e.g., stroke and Parkinson disease recovery), cardiopulmonary physiotherapy (e.g., chest clearance and cardiac rehab), and pediatric and geriatric care.

#### Alternate Names:

Physical Therapy; PT; Movement Rehabilitation Therapy

### Physostigmine

A reversible acetylcholinesterase inhibitor that may reverse severe central anticholinergic delirium in carefully selected patients after assessment for contraindications and cardiac conduction abnormalities.

### Phytomenadione

A form of vitamin K needed by the liver to make several blood-clotting proteins. It is used to prevent or treat bleeding caused by vitamin K deficiency or by medicines that block vitamin K.

### Phytophotodermatitis

A phototoxic reaction caused when psoralens or related furocoumarins from plants such as limes, celery, or wild parsnip contact the skin and are followed by ultraviolet-A exposure. Burning, sharply patterned redness, streaks, or blisters may be followed by prolonged postinflammatory hyperpigmentation; it is not an immune-mediated sun allergy.

### Pia Mater

The thin, delicate inner covering that closely follows the surface of the brain and spinal cord. Together

---

with the other coverings and fluid around the nervous system, it helps protect these organs.

### **Pica**

Repeated eating of things that have no useful nutritional value, such as soil, paper, ice, or hair, when this is not expected for a person's age or culture. It may cause poisoning, infection, blockage, or poor nutrition and can be linked with iron deficiency or mental-health conditions.

### **PICC Line Dressing Care**

A sterile clinical procedure for replacing the transparent semipermeable dressing and securement device over a peripherally inserted central catheter insertion site to prevent catheter-related bloodstream infections (CRBSI).

#### **Alternate Names:**

PICC Dressing Change; Central Line Dressing Care; PICC Sterile Maintenance

### **PICC Line Flushing**

The routine pulsatile push-pause administration of sterile normal saline or heparin lock solution through a PICC lumen to clear intraluminal blood, maintain catheter patency, and prevent thrombotic occlusion.

#### **Alternate Names:**

PICC Flush Protocol; Central Catheter Heparinization; PICC Patency Flushing

### **PICC Line Insertion**

Ultrasound-guided percutaneous cannulation of a deep peripheral vein (such as the basilic, cephalic, or brachial vein) in the upper arm, advancing the radiopaque catheter tip under fluoroscopic or electrocardiographic guidance to the cavoatrial junction.

#### **Alternate Names:**

PICC Placement; Peripherally Inserted Central Catheterization; PICC Cannulation

### **Pick Disease**

A distinct subtype of frontotemporal lobar degeneration (FTLD-tau) characterized pathologically by circumscribed knife-edge frontotemporal cortical lobar atrophy and intracellular round, silver-staining, tau-positive inclusions (Pick bodies), presenting

with progressive behavioral disinhibition, personality change, and language impairment.

#### **Alternate Names:**

Frontotemporal Dementia with Pick Bodies; FTLD-Tau

### **Pick's Disease**

An older term often referring to frontotemporal dementia, a group of disorders causing progressive changes in behavior, personality, language, or executive function. The term is now used less consistently.

### **Pickwickian Syndrome**

An older name for obesity hypoventilation syndrome, involving obesity, daytime hypercapnia, and sleep-disordered breathing.

### **PID**

Alternate name; see *Pelvic inflammatory disease*.

### **Piebaldism**

A rare condition present from birth in which some areas of skin and hair lack pigment. It usually causes stable, sharply outlined white patches and often a white section of hair at the front; it is commonly inherited through a change in the KIT gene.

### **Piedra**

A superficial fungal infection of hair shafts producing hard or soft nodules attached to the hair. Black piedra is caused by *Piedraia hortae*, while white piedra is caused by *Trichosporon* species; treatment may include hair trimming and antifungal therapy.

### **Pierre Robin Sequence**

A condition present at birth in which a small lower jaw allows the tongue to fall backward and may block breathing. Many affected infants also have a U-shaped opening in the roof of the mouth and feeding difficulty.

### **Pierson Syndrome**

A rare inherited condition causing severe kidney protein loss from birth and unusually small pupils. It can lead to swelling, kidney failure, poor vision,

---

and developmental problems.

### **Piezogenic Pedal Papules**

Small, soft skin-colored papules caused by fat herniating through connective tissue at the edges of the heels when standing. They are common and usually harmless, but may be painful with prolonged standing or in people with connective-tissue disorders.

### **Pigeon Breast**

A congenital structural deformity of the anterior chest wall characterized by an abnormal anterior protrusion of the sternum and adjacent costal cartilages, giving the chest a prominent, pouter-pigeon appearance.

#### **Alternate Names:**

Pectus Carinatum

### **Pigment**

A substance that gives skin, hair, eyes, tissue, or other material its color. Pigments may be made by the body, received from outside, or altered by disease, injury, medicines, or aging.

### **Pigmentary Glaucoma**

A secondary open-angle glaucoma caused by dispersion of iris pigment, which accumulates in the trabecular meshwork and impairs aqueous outflow. It may cause raised intraocular pressure and progressive optic-nerve damage.

### **Pigmentary Mosaicism**

Uneven skin coloring caused by groups of skin cells carrying different genetic information. It may produce lighter or darker lines, swirls, or patches; most cases affect only the skin, but associated developmental problems should be assessed when present.

### **Pigmentation**

Coloration of skin, hair, eyes, or other tissue produced mainly by melanin, carotenoids, blood, or exogenous material; abnormal changes may reflect inflammation, endocrine disease, drugs, or tumors.

### **Pigmentation Disorder**

A condition causing abnormal increase, decrease, or uneven distribution of pigment in the skin, mucosa, hair, or eye.

### **Pigment Cirrhosis**

Liver cirrhosis associated with abnormal pigment or iron deposition, as may occur in hereditary hemochromatosis.

### **Pigment Dispersion Syndrome**

An ocular condition characterized by the mechanical shedding of pigment granules from the posterior surface of the iris (due to rubbing against lens zonules), which deposit on the corneal endothelium (Krukenberg spindle) and obstruct the trabecular meshwork, predisposing to pigmentary glaucoma.

### **Pigmented Birthmarks**

Congenital cutaneous lesions characterized by localized melanin accumulation or melanocytic hyperplasia present at birth or emerging during early infancy, encompassing congenital melanocytic nevi, café-au-lait macules, dermal melanocytosis (Mongolian spots), and lentigines.

#### **Alternate Names:**

Congenital Melanocytic Nevi; Pigmented Nevi; Birthmarks, Pigmented

### **Pigmented Purpuric Dermatitis**

A group of chronic harmless eruptions caused by leakage of small amounts of blood from superficial vessels. They produce reddish-brown or copper-colored speckles and patches, commonly on the lower legs, and usually do not cause systemic illness.

### **Pilar Cyst**

A usually harmless, firm lump beneath the skin that forms from a hair follicle, most often on the scalp. It may be inherited, become inflamed or painful, and can usually be removed if it causes symptoms or repeated problems.

### **Piles**

Enlarged or swollen blood vessels and supporting

---

tissue in the anus or lower rectum. Internal piles may cause painless bright-red bleeding or tissue protruding during bowel movements; external piles may cause pain, itching, or a tender lump.

### **Piles (Haemorrhoids)**

Alternate name; see *Piles*.

### **Pill**

A solid medicine designed to be taken by mouth. It may contain one or more active ingredients and may be made to release them immediately or slowly; pills should be taken as directed and not crushed unless the instructions allow it.

#### **Alternate Names:**

Tablet

### **Pilocarpine**

A medicine that activates muscarinic receptors, causing pupil constriction and increased secretions. It may lower eye pressure in glaucoma or relieve dry mouth, but can cause sweating, blurred vision, headache, or breathing problems.

### **Pilocytic Astrocytoma**

A usually slow-growing brain tumor arising from support cells, most often in children. It commonly affects the cerebellum or visual pathway and may cause headache, vomiting, seizures, balance problems, or vision changes.

### **Piloerection**

Elevation of hair shafts producing gooseflesh, caused by contraction of arrector pili muscles under sympathetic stimulation from cold, fear, pain, fever, or certain drugs.

### **Pilomatrix Carcinoma**

A rare skin cancer that develops from cells involved in hair formation. It usually appears as a firm lump on the head, neck, or arm and can grow into nearby tissue or spread elsewhere.

### **Pilomatrixoma**

Pilomatrixoma is a usually benign skin tumor de-

rived from hair-matrix cells, presenting as a firm subcutaneous nodule, often on the head or neck.

### **Pilomatrixoma (Pilomatricoma)**

A usually benign, firm skin tumor arising from hair-matrix cells, most often on the head, neck, or upper limbs of children and young adults. It may be mistaken for a cyst; excision confirms the diagnosis and is usually curative.

### **Pilonidal Cyst**

A cyst or sinus in the cleft between the buttocks, usually containing hair and skin debris. It may become infected, painful, swollen, and drain pus.

#### **Alternate Names:**

Pilonidal Sinus; Sacrococcygeal Fistula

### **Pilonidal Dimple**

Alternate name; see *Sacral dimple*.

### **Pilonidal Disease**

A condition in which hair and skin debris collect in a small pit or tunnel in the cleft between the buttocks. It can cause repeated swelling, pain, drainage, or an abscess.

### **Pilonidal Fistula**

An abnormal tunnel from a pilonidal sinus to the skin, usually in the natal cleft, that can drain pus or recur after abscess formation.

### **Pilonidal Sinus Disease**

A chronic sinus or cyst in the natal cleft, usually containing hair and debris. It may cause recurrent pain, swelling, drainage, or abscess formation.

### **Pilus**

A tiny hair-like structure on some bacteria. It may help bacteria attach to human cells, move, or transfer genetic material to other bacteria, which can contribute to infection or antibiotic resistance.

### **Pimozide**

An antipsychotic medicine used for selected disorders involving severe tics or abnormal thoughts. It

---

can cause movement problems, sleepiness, and dangerous heart-rhythm changes, so dosing and heart monitoring require careful medical supervision.

### **Pimple**

A common lay term for a small inflamed or blocked skin lesion, often an acne papule or pustule.

### **Pin Care**

Pin care keeps the skin around external fixation or traction pins clean and monitored for redness, swelling, drainage, loosening, or infection.

### **Pinched Nerve**

Compression or irritation of a nerve that can cause pain, tingling, numbness, or weakness along the nerve's distribution.

#### **Alternate Names:**

Compressed Nerve; Nerve Compression; Radiculopathy

### **Pinch Strength**

Pinch strength is the force produced when the thumb presses against one or more fingers and can help assess hand, nerve, muscle, and functional impairment.

### **Pindolol**

Pindolol is a beta-blocker used for selected cardiovascular conditions; it can slow the heart rate and worsen asthma in susceptible people.

### **Pineal Cyst**

A fluid-filled sac in the small gland deep in the brain that helps regulate sleep-related signals. Most are found incidentally and cause no harm, but a large cyst can rarely cause headache, eye problems, or fluid buildup.

### **Pineal Germinoma**

A pineal germinoma is a malignant germ-cell tumor in the pineal region, often causing headache, eye-movement abnormalities, or hydrocephalus and usually responsive to radiotherapy and chemotherapy.

### **Pineal Gland**

A small hormone-producing gland deep in the brain that releases melatonin. Melatonin helps coordinate the body's day-night clock and influences sleep timing, seasonal rhythms, and related biological functions.

### **Pineal Gland Tumor**

A growth in or near the pineal gland. It may block the normal flow of brain fluid and cause headache, nausea, vomiting, sleepiness, or problems moving the eyes; symptoms and treatment depend on its type and size.

### **Pinealoma**

A primary neoplasm arising within the pineal gland region, predominantly including germ cell tumors (germinomas, non-germinomatous germ cell tumors) and pineal parenchymal tumors (pineocytoma, pineoblastoma); commonly presents with obstructive hydrocephalus and Parinaud dorsal midbrain syndrome.

### **Pineal Parenchymal Tumor**

A tumor arising from cells of the pineal gland, ranging from slow-growing to aggressive. It may block the flow of brain fluid and cause headache, vomiting, sleep changes, or abnormal eye movements.

### **Pineoblastoma**

An aggressive cancer in the pineal region of the brain, occurring mainly in children. It may block brain-fluid flow and cause headache, vomiting, sleepiness, or eye-movement problems and requires specialist treatment.

### **Pineocytoma**

Pineocytoma is a usually slow-growing tumor of pineal cells that may obstruct cerebrospinal-fluid flow and cause raised intracranial pressure.

### **Pine Oil Poisoning**

Swallowing or inhaling concentrated pine oil can irritate the mouth and stomach and may cause drowsiness, seizures, or lung injury if it enters the lungs.

---

## Pinguecula

A pinguecula is a benign yellowish raised growth on the conjunctiva near the cornea, often associated with ultraviolet light, wind, and dust; it may irritate the eye.

## Pinhole Meatus

An unusually tiny external opening, most often the opening of the urethra. It can make urine flow weak or spraying, cause discomfort, or make catheter placement difficult.

## Pink Eye

A common name for conjunctivitis, inflammation of the eye covering causing redness, irritation, watering, or discharge.

### Alternate Names:

Conjunctival Inflammation; Acute Conjunctivitis

## Pink Puffer

An older, nonformal term for a person with emphysema who has severe breathlessness, a thin build, and pursed-lip breathing. The term is not a separate diagnosis and is less useful than describing the person's COPD and test results.

## Pinna

The visible external part of the ear that collects sound.

## Pinocytosis

A process in which a cell takes in small drops of surrounding fluid and dissolved substances. The cell membrane folds inward to form tiny bubbles that carry the material inside for use, transport, or breakdown.

## Pins and Needles

A common name for paresthesia, an abnormal tingling, prickling, or numb sensation. Brief episodes often follow pressure on a nerve; persistent symptoms need evaluation.

## Pint

A pint is a unit of volume; its exact amount varies

by measurement system, so the local standard must be specified when measuring fluids or medicines.

## Pinta

A long-lasting skin infection caused by *Treponema carateum* bacteria. It spreads through close skin contact and produces changing stages of discolored, scaly, or thickened patches, usually without damage to internal organs.

## Pinworm

A small intestinal worm that commonly causes intense itching around the anus at night. Infection spreads when microscopic eggs are swallowed or transferred from contaminated hands or surfaces; treatment usually includes medicine, hygiene, and sometimes treating household contacts.

## Pinworm Infection

Pinworm infection is an intestinal infestation by *Enterobius vermicularis*, causing nighttime anal itching, disturbed sleep, and sometimes abdominal or genital irritation. Treatment of the person and close household contacts, hygiene, and repeat dosing reduce reinfection.

### Alternate Names:

Enterobiasis; Oxyuriasis; Threadworm Infection

## Pinworm Infestation

Alternate name; see *Pinworm Infection*.

## Pinworms

Pinworms are intestinal nematodes, *Enterobius vermicularis*, that cause nighttime anal itching; treatment, hand hygiene, and laundering help prevent household reinfection.

## Pinworm Test

A test for *Enterobius vermicularis*, usually the early-morning adhesive-tape test applied to the perianal skin before bathing or defecation.

## Pioglitazone

A medicine for type 2 diabetes that helps muscle and fat cells respond better to insulin, lowering



---

blood sugar. It can cause weight gain and swelling and may worsen heart failure, so it is unsuitable for some people.

### **Piperacillin**

Piperacillin is an extended-spectrum penicillin antibiotic often combined with tazobactam to treat serious infections including *Pseudomonas*; allergy, diarrhoea, cytopenias, electrolyte effects, and renal dosing are important.

### **Piperazine**

An older antiparasitic medicine that paralyzes certain intestinal worms, especially roundworms and pinworms. Its use depends on local guidance and the particular infection.

### **Piperonyl Butoxide**

A pesticide synergist that inhibits insect detoxification enzymes and increases the activity of certain insecticides; it can irritate skin, eyes, or airways.

### **Piperonyl Butoxide With Pyrethrins Poisoning**

Exposure to these insecticide ingredients can irritate the skin, eyes, or airways and may cause coughing, nausea, vomiting, dizziness, or wheezing. Severe breathing symptoms or swallowed product require urgent poison-control or emergency advice.

### **Pipotiazine**

A long-acting phenothiazine antipsychotic given by injection in selected settings. It can cause movement disorders, drowsiness, and other adverse effects and requires clinical monitoring.

### **Pipradrol**

An older central nervous system stimulant with limited current use.

### **Piracetam**

A nootropic compound used in some countries for selected neurologic indications; evidence and approval vary.

### **PI-RADS**

A standard scoring system for prostate MRI findings. It estimates how likely an area is to contain an important prostate cancer and helps guide decisions about follow-up or biopsy; it does not diagnose cancer by itself.

### **Pirfenidone**

Pirfenidone is an antifibrotic medicine used to slow decline in idiopathic pulmonary fibrosis; nausea, photosensitivity, fatigue, and liver-enzyme elevation may occur and require monitoring.

### **Piriformis Syndrome**

Pain or altered sensation in the buttock or down the leg when the piriformis muscle irritates or presses on the sciatic nerve. Other causes, especially a spinal nerve problem, must be considered before treatment.

#### **Alternate Names:**

Piriformis Muscle Spasm; Deep Gluteal Syndrome

### **Pirimicarb**

Pirimicarb is a selective carbamate insecticide targeting aphids; exposure can inhibit acetylcholinesterase and cause cholinergic symptoms.

### **Piromidic Acid**

An older quinolone-like antibacterial investigated or used for urinary and gastrointestinal infections; current use depends on local availability, susceptibility, and safety guidance.

### **Piroxicam**

A long-acting nonsteroidal anti-inflammatory drug used for pain and inflammation; gastrointestinal bleeding, kidney injury, cardiovascular events, and hypersensitivity are important risks.

### **Piroxicam Overdose**

Taking too much piroxicam, an anti-inflammatory pain medicine, can cause stomach pain, vomiting, bleeding, kidney injury, drowsiness, seizures, or breathing problems. Large or intentional overdoses can be life-threatening.

---

## Pisa Syndrome

A sustained involuntary sideways flexion of the trunk, often associated with medicines, Parkinson disease, or other neurologic disorders.

## Pisiform

A small, pea-shaped wrist bone embedded in a tendon on the little-finger side of the palm. It helps form the passageway for the ulnar nerve and artery and may hurt after injury or with arthritis.

## Pisiform Bone

A small pea-shaped carpal bone embedded in the tendon of flexor carpi ulnaris on the ulnar side of the wrist; fracture or arthritis can cause localized wrist pain.

## Pits

Small depressions or holes in a surface. In medicine, pits may occur in nails, skin, teeth, or other tissues and can result from normal anatomy, injury, infection, inflammation, or inherited conditions.

## Pitted Keratolysis

A superficial, non-inflammatory bacterial infection of the stratum corneum of the plantar foot (most commonly caused by *Kytococcus sedentarius* or *Corynebacterium* species), presenting with punched-out superficial crateriform pits, hyperhidrosis, and severe sulfurous malodor (bromhidrosis).

## Pitting Edema

Swelling in which pressure applied to the skin leaves a temporary indentation, usually because of excess interstitial fluid.

## Pituicytoma

A rare, usually slow-growing tumor near the pituitary gland. It may cause headache, vision problems, or hormone disturbances by pressing on nearby structures and is usually assessed with brain imaging and tissue examination.

## Pituitary

The pituitary is an endocrine gland at the base of

the brain that releases hormones controlling growth, thyroid and adrenal function, reproduction, lactation, and water balance.

## Pituitary Adenoma

A usually noncancerous growth arising from the front part of the pituitary gland. It may release too much of a hormone or press on nearby tissue, causing headache, vision changes, or hormone problems.

## Pituitary Apoplexy

Sudden bleeding or loss of blood supply inside a pituitary growth, causing severe headache, vision changes, vomiting, eye-movement problems, or confusion. It is an emergency because hormone failure can cause life-threatening low blood pressure.

## Pituitary Carcinoma

Pituitary carcinoma is a rare malignant pituitary tumor defined by spread to distant or separate sites, often producing excess hormones.

## Pituitary Disease

A disorder of the pituitary gland that alters hormone secretion or causes mass effects, including adenomas, inflammation, hemorrhage, genetic disease, or hypopituitarism.

## Pituitary Dwarfism

Pituitary dwarfism is severe short stature caused by deficient growth hormone or broader pituitary dysfunction, with evaluation focused on endocrine and genetic causes.

## Pituitary Gland

An endocrine gland at the base of the brain that regulates growth, thyroid, adrenal, reproductive, lactation, and water-balance hormones.

## Pituitary Hormone

A hormone secreted by the anterior or posterior pituitary that regulates growth, thyroid and adrenal function, reproduction, lactation, water balance, or other endocrine organs.

---

## Pituitary Insufficiency

Alternate name; see *Hypopituitarism*.

## Pituitary Macroadenoma

A pituitary adenoma measuring at least 10 mm. It may compress the optic chiasm and cause visual-field loss or headache, or alter pituitary hormone production; MRI, visual-field testing, and endocrine assessment are central to evaluation.

## Pituitary Microadenoma

A pituitary adenoma smaller than 10 mm. It may be hormonally inactive and found incidentally or may produce prolactin, ACTH, growth hormone, or other hormones; evaluation depends on imaging and endocrine findings.

## Pituitary Neoplasm

A pituitary neoplasm is an abnormal growth in or near the pituitary gland and may affect hormones, vision, or nearby brain structures.

## Pituitary Neuroendocrine Tumor

Alternate name; see *Pituitary tumors and adenomas*.

## Pituitary Tumor

A pituitary tumor is a growth in the pituitary gland that may be noncancerous but can cause hormone excess, hormone deficiency, or visual problems.

## Pituitary Tumors Pathology

Pathologic assessment of pituitary neuroendocrine tumors, including cell lineage, hormone production, reticulin architecture, cytologic features, proliferative activity, and invasion. Immunohistochemistry helps classify the tumor.

## Pityriasis

A group of skin conditions marked by fine, flaky scaling. Examples include pityriasis rosea and pityriasis versicolor, which have different causes and may produce itching or lighter, darker, or pink patches.

## Pityriasis Alba

Pityriasis alba is a common benign skin condition causing faint, dry, lighter patches, especially on children's faces, often more noticeable after tanning.

## Pityriasis Lichenoides

An uncommon inflammatory skin disorder producing recurrent small scaly papules, with forms ranging from mild chronic lesions to an acute ulceronecrotic eruption.

## Pityriasis Rosea

An acute, self-limited eruption often beginning with a herald patch followed by oval scaly lesions distributed along skin-cleavage lines. It usually resolves within several weeks; treatment is symptomatic, and atypical, severe, recurrent, or persistent disease should be reassessed.

### Alternate Names:

Pityriasis Rosea of Gibert; Christmas Tree Rash

## Pityriasis Rubra Pilaris

A group of rare inflammatory skin disorders that cause well-defined, reddish-orange scaly patches and rough bumps around hair follicles. Patches may spread widely while leaving islands of normal skin. The palms and soles may become thick, yellowish, dry, and cracked. The course varies and may last for months or years.

## Pityriasis Versicolor

Alternate name; see *Tinea versicolor*.

## Pivampicillin

An orally absorbed prodrug of ampicillin used for selected bacterial infections; its use is limited in some settings by adverse effects and availability.

## Pivot Joint

A joint in which one bone rotates around another bone. The joints between the first two neck bones and parts of the forearm are examples.

## Pixantrone

Pixantrone is an anthracenedione antineoplastic

---

drug developed for selected aggressive lymphomas, with potential marrow, infection, and cardiac risks.

## **PJI**

Alternate name; see *Periprosthetic joint infection*.

## **PJS**

Alternate name; see *Peutz-Jeghers syndrome*.

## **PKD**

Alternate name; see *Polycystic kidney disease*.

## **Placebo**

An inactive substance, treatment, or procedure designed to resemble an active treatment. In a clinical trial, it is given to a control group so researchers can compare outcomes and estimate the specific effect of the treatment being studied.

### **Alternate Names:**

Sham Treatment; Inactive Control

## **Placebo Control**

A placebo control is an inactive treatment designed to resemble the intervention in a clinical trial, helping separate treatment effects from expectation and natural change.

## **Placebo Effect**

A real change in symptoms or wellbeing that can occur after a person receives an inactive treatment, partly because of expectations, attention, and the treatment setting. It does not prove that the inactive treatment changed the disease itself.

## **Placenta**

The temporary organ connecting mother and fetus, allowing exchange of oxygen, nutrients, wastes, and hormones.

## **Placenta Abruptio**

Placental abruptio is premature separation of the placenta from the uterine wall, causing bleeding, abdominal pain, uterine tenderness, fetal compromise, or maternal shock.

## **Placenta Accreta**

Placenta accreta spectrum occurs when placental tissue abnormally attaches to or invades the uterine wall, creating major risk of hemorrhage during attempted delivery or removal.

## **Placenta Accreta Spectrum**

Abnormal adherence or invasion of placental villi into the uterine wall, ranging from accreta to increta and percreta; it can cause severe postpartum hemorrhage and requires planned obstetric management.

## **Placenta Disorder**

A disorder of placental implantation, structure, perfusion, or separation, including previa, accreta spectrum, abruptio, infarction, or placental insufficiency.

## **Placenta Increta**

Abnormal placental invasion into the myometrium without reaching the serosa, increasing the risk of massive hemorrhage at delivery and often requiring cesarean hysterectomy.

## **Placental Abnormalities**

Disorders of placental location, attachment, structure, or separation that may affect pregnancy or cause bleeding.

## **Placental Abruptio**

Premature detachment of a normally implanted placenta from the uterine myometrium prior to delivery of the fetus, classically presenting with painful dark vaginal bleeding, uterine hypertonicity, and fetal distress.

### **Alternate Names:**

Abruptio Placentae; Premature Placental Separation; Accidental Antepartum Hemorrhage

## **Placental Accreta Spectrum**

Abnormal placental attachment to the myometrium, ranging from accreta (adherent to decidua basalis) to increta (invading myometrium) to percreta (invading through serosa). Major risk factor is prior

---

caesarean section with anterior placenta previa.

### **Placental Hormone**

A hormone produced by placental tissue, such as hCG, progesterone, estrogens, or human placental lactogen, that supports pregnancy, fetal growth, and maternal metabolic adaptation.

### **Placental Infarction**

A region of placental tissue damaged by inadequate maternal blood flow and coagulation, which may be incidental or, when extensive, associated with fetal growth restriction or pregnancy complications.

### **Placental Insufficiency**

Placental insufficiency is inadequate placental delivery of oxygen and nutrients, which can cause fetal growth restriction, abnormal Doppler findings, reduced fluid, or fetal distress.

### **Placental Lactogen**

Placental lactogen is a hormone made by the placenta that alters maternal metabolism and supports breast development and fetal nutrient availability during pregnancy.

### **Placental Mesenchymal Dysplasia**

A rare abnormal development of the placenta in which supporting tissue becomes enlarged and cyst-like. It can resemble a molar pregnancy on scans and may be associated with poor fetal growth, fetal abnormalities, or pregnancy complications.

### **Placental Separation Signs**

Clinical signs indicating the placenta has separated from the uterine wall and is ready for delivery. Classic signs: lengthening of the umbilical cord (cord lengthens as the placenta descends), a sudden gush of blood, and uterine elevation and firming (the uterus rises and becomes globular as the placenta moves into the lower segment).

### **Placental Site Nodule**

A benign, usually incidental remnant of intermediate trophoblast at a prior implantation site, com-

posed of small nests of chorionic-type or extravillous trophoblast.

### **Placenta Percreta**

Placenta percreta is the most invasive form of placenta accreta spectrum, with villi growing through the uterine wall and sometimes into nearby organs such as the bladder.

### **Placenta Praevia**

A condition in which the placenta lies over or near the opening of the cervix. It can cause painless bleeding later in pregnancy and may require monitoring, activity advice, or cesarean delivery.

### **Placenta Previa**

Implantation of the placenta over or near the internal cervical opening, which can cause painless bleeding during the second half of pregnancy.

### **Placentation**

The developmental process by which the placenta forms and attaches to the uterus, establishes maternal–fetal circulation, and develops the exchange and endocrine functions needed for pregnancy.

### **Plagiocephaly**

Asymmetric flattening of an infant's skull. The common positional form develops from repeated pressure on one area; rare skull-shape changes from early fusion of a cranial suture are called synostotic plagiocephaly and need medical assessment.

#### **Alternate Names:**

Flat Head Syndrome; Deformational Plagiocephaly

### **Plague**

A serious infection caused by *Yersinia pestis*, transmitted mainly by infected fleas or respiratory droplets in pneumonic disease.

### **Plain Radiography**

An imaging method that uses X-rays to create a two-dimensional projection of internal structures. It is used to assess bones, the chest, abdomen, teeth,

---

and other selected body regions.

**Alternate Names:**

Roentgenography; Projection Radiography

**Planar**

Flat or arranged in one plane. In medical imaging, the term describes a two-dimensional view or projection, unlike three-dimensional imaging that shows depth and allows reconstruction from multiple directions.

**Plane Joint**

A joint with mostly flat surfaces that slide over one another. Small movements occur in several directions, as in some wrist and foot joints.

**Plantar**

Relating to the sole or underside of the foot, including the plantar surfaces of the toes.

**Plantar Fasciitis**

Pain from irritation and small injuries in the strong band of tissue supporting the sole of the foot. Heel pain is often worst with the first steps after waking or resting and may worsen after prolonged standing.

**Plantar Fibromatosis**

A benign overgrowth of fibrous tissue in the plantar fascia that forms firm nodules in the sole of the foot. It may be painless or cause pressure pain while walking; treatment depends on symptoms and can include footwear changes, injections, or surgery.

**Plantar Flexion**

Downward movement of the foot at the ankle, pointing the toes away from the shin. It is produced mainly by the calf muscles and is used in standing on tiptoe, walking, running, and jumping.

**Plantar Hyperhidrosis**

Alternate name; see *Hyperhidrosis*.

**Plantaris Muscle**

A small posterior-leg muscle with a long tendon that assists weakly with knee flexion and plantar

flexion; it can be strained during sudden athletic movement.

**Plantar Nerve**

A nerve supplying sensation to the sole of the foot and motor branches to intrinsic foot muscles; compression can cause burning pain, numbness, or altered toe movement.

**Plantar Neuroma**

Alternate name; see *Morton neuroma*.

**Plantar Reflex**

Movement of the toes after the sole of the foot is gently stroked. In most adults the toes bend downward; upward movement of the big toe may indicate a problem in brain or spinal nerve pathways controlling movement.

**Plantar Response**

Movement of the toes and foot after the sole is stimulated. The pattern helps clinicians assess pathways between the brain, spinal cord, and leg; an unusual upward movement of the big toe in an adult may indicate neurologic disease.

**Plantar Wart**

A wart on the sole of the foot caused by human papillomavirus. Pressure from standing and walking can push it inward and make it painful; small black dots may represent tiny clotted blood vessels.

**Alternate Names:**

Verruca; Foot Wart

**Plant Fertilizer Poisoning**

Fertilizer exposure may irritate the mouth, skin, eyes, stomach, or lungs. Some products can dangerously alter blood oxygen or body salts; keep the container and contact poison control after swallowing or significant exposure.

**Plant Poisoning**

Illness caused by eating, touching, or inhaling a toxic plant or plant product; effects range from local irritation to organ failure.



---

## **Plant Thorn Arthritis**

Plant thorn arthritis is persistent joint inflammation after a thorn puncture introduces plant material, sometimes requiring removal of foreign material and treatment of infection or inflammation.

## **Plant Thorn Synovitis**

Inflammation of a joint after penetration by a plant thorn or splinter, sometimes involving an infection introduced by the injury. Symptoms may develop gradually as persistent joint pain, swelling, and limited movement; evaluation may require removal of retained material and targeted treatment.

### **Alternate Names:**

Thorn-Induced Arthritis; Foreign Body Synovitis

## **Plaque**

A localized buildup or raised area of material. In the skin, a plaque is a broad, raised lesion often formed by joined papules; in an artery, plaque is a fatty deposit in the vessel wall that can narrow the artery.

## **Plaque And Tartar On Teeth**

Dental plaque is a bacterial biofilm that forms on teeth, while tartar is hardened mineralized plaque; both contribute to cavities, gingivitis, and periodontitis.

## **Plaque Assay**

A laboratory test that estimates the number of infectious virus particles in a sample. The sample is placed on living cells, and each infectious particle may produce a visible area of damaged cells that can be counted.

## **Plaque Psoriasis**

The most common form of psoriasis, producing well-demarcated raised red plaques with silvery scale, commonly on elbows, knees, scalp, and lower back and sometimes associated with arthritis.

## **Plasma**

The liquid part of blood carrying cells, proteins, nutrients, hormones, wastes, and other dissolved

substances through the body.

## **Plasma Amino Acids**

A blood test that measures amino acids in plasma. It can help evaluate inherited metabolic disorders, poor nutrition, liver disease, or other conditions affecting amino-acid processing.

## **Plasmablastic Lymphoma**

Plasmablastic lymphoma is an aggressive B-cell lymphoma often associated with immune deficiency, HIV infection, or immunosuppression.

## **Plasma Cell**

An immune cell that develops from a B cell and produces antibodies. Antibodies help recognize and control germs and other foreign substances.

## **Plasma Cell Mastitis**

A chronic benign inflammatory breast lesion with prominent plasma-cell infiltration, often involving ducts beneath the nipple and potentially causing a mass, nipple retraction, or discharge. It can clinically mimic carcinoma.

## **Plasma Cell Myeloma**

A cancer of antibody-producing plasma cells, usually in the bone marrow. It can produce an abnormal antibody and damage bones, kidneys, blood-cell production, or calcium balance, causing bone pain, fractures, anemia, infections, or kidney problems.

## **Plasma Cell Neoplasm**

A clonal disorder of antibody-producing plasma cells, ranging from localized plasmacytoma to multiple myeloma with marrow, bone, kidney, calcium, or blood abnormalities.

## **Plasma Cells**

Immune cells that make antibodies. They develop from B cells after the body recognizes an infection or other foreign substance. Abnormal plasma-cell growth can cause diseases such as multiple myeloma and may produce excessive abnormal antibodies.

---

## Plasma Concentration

The amount of a substance present in a defined volume of plasma at a given time. It is used to interpret exposure, toxicity, therapeutic drug monitoring, and pharmacokinetic relationships.

## Plasmacytoma

A solitary mass of neoplastic plasma cells, which may occur in bone (solitary plasmacytoma of bone) or soft tissue (extramedullary plasmacytoma). It can progress to multiple myeloma.

## Plasma Expanders

Fluids given through a vein to increase circulating blood volume. Crystalloid solutions are commonly used, while selected colloids have limited roles; treatment depends on the cause of fluid loss and patient condition.

## Plasma Kallikrein

A blood enzyme that helps produce bradykinin, a substance that widens blood vessels and increases leakage from them. It contributes to inflammation and swelling and is involved in some inherited swelling disorders.

## Plasmalogen

A special type of fat found in cell membranes, especially in the brain, nerves, and heart. The body makes it partly in structures called peroxisomes; very low levels occur in some inherited metabolic disorders.

## Plasma Membrane

The selectively permeable lipid bilayer surrounding a cell, containing proteins that control transport, signaling, adhesion, recognition, and communication with the extracellular environment.

## Plasma Osmolarity

The concentration of dissolved particles in the liquid part of blood. It mainly reflects water balance and substances such as sodium, glucose, and urea; abnormal values can indicate dehydration, excess water, or metabolic disease.

## Plasmapheresis

A specialized medical procedure that filters and separates the pale yellow liquid portion of blood (the plasma) from blood cells. In therapeutic plasma exchange, whole blood is drawn through an intravenous catheter into a machine that spins or filters it, separating out the patient's plasma—which often carries disease-causing autoantibodies, abnormal proteins, or inflammatory toxins—and discarding it. The patient's red blood cells, white blood cells, and platelets are then mixed with replacement fluids (such as donor plasma or sterile albumin solution) and safely returned to the bloodstream. Plasmapheresis is an established, life-saving treatment for severe autoimmune and neurological disorders where the immune system attacks the body's own tissues, including Guillain-Barré syndrome, myasthenia gravis crisis, thrombotic thrombocytopenic purpura (TTP), chronic inflammatory demyelinating polyneuropathy (CIDP), and Goodpasture syndrome. It is also used in blood donation centers to harvest plasma from healthy donors.

### Alternate Names:

Therapeutic Plasma Exchange; TPE; Plasma Exchange; Apheresis

## Plasma Substitutes

Intravenous fluids intended to expand plasma volume, including crystalloids and colloid solutions; choice depends on the clinical indication and risks.

## Plasma / Urine Catecholamine Assay

Blood or urine tests measuring catecholamines or their breakdown products. They may help investigate pheochromocytoma or paraganglioma, but medicines, stress, posture, diet, and collection conditions can affect results.

### Alternate Names:

Fractionated Catecholamines; Free Metanephrines

## Plasmid

A plasmid is a small usually circular DNA molecule separate from a bacterial chromosome that can replicate independently and sometimes carries antibiotic-resistance genes.

---

## Plasmids

Small DNA rings found separately from the main chromosome in many bacteria. They can copy themselves and may carry genes that help bacteria survive, including genes for antibiotic resistance or disease-causing traits.

## Plasmin

An enzyme that breaks down fibrin, the protein mesh holding a blood clot together. It is formed from plasminogen and helps remove clots after a damaged blood vessel has been repaired.

## Plasminogen

An inactive blood protein that can be changed into plasmin, an enzyme that breaks down the protein framework of blood clots. The body activates it when a clot needs to be removed.

## Plasminogen Activator

An enzyme or medicine that changes plasminogen into plasmin, which breaks down the protein mesh of blood clots. These medicines can dissolve dangerous clots but may cause serious bleeding.

## Plasminogen Inactivators

Substances or medicines that reduce the conversion of plasminogen into plasmin, thereby slowing breakdown of blood clots. They may be used or studied when limiting bleeding is important.

## Plasmodium

A group of parasites spread through bites from infected *Anopheles* mosquitoes and responsible for malaria. The parasites first infect the liver, then red blood cells, causing fever and potentially severe organ complications.

## Plasmodium Vivax

A parasite spread by infected *Anopheles* mosquitoes that causes malaria. It infects red blood cells and can remain dormant in the liver, allowing illness to return months or years after the first infection.

## Plasmodium Vivax Malaria

Malaria caused by the parasite *Plasmodium vivax* and spread by infected mosquitoes. Fever attacks may recur about every two days, and dormant liver forms can cause later relapses; treatment must clear both blood and liver stages when safe.

### Alternate Names:

Vivax Malaria; Benign Tertian Malaria

## Plaster Of Paris

A powder that hardens when mixed with water. In healthcare it is used to make casts and splints that hold an injured bone or joint still while it heals; swelling, numbness, or increasing pain under a cast needs urgent review.

## Plastic and Reconstructive Surgery

Surgery that restores form or function after congenital difference, injury, disease, or removal of tissue.

## Plastic Bronchitis

A rare condition producing large, branching mucous casts that form in the airways, causing life-threatening airway obstruction, often associated with congenital heart disease or post-fontan circulation.

## Plastic Casting Resin Poisoning

Exposure to uncured casting resin or its solvents can irritate or burn the skin, eyes, lungs, and digestive tract. Swallowing or breathing fumes may cause vomiting, dizziness, or lung injury and requires poison-control advice.

## Plastic Resin Hardener Poisoning

Resin hardeners may contain corrosive or irritating chemicals that burn the skin, eyes, mouth, throat, or stomach. Fumes can injure the lungs; rinse exposed areas and seek urgent poison-control advice after significant exposure.

## Plastic Surgery

Surgery that repairs, rebuilds, or changes body tissues to improve physical function, appearance, or both.

---

## Platelet

A small piece of a blood cell that helps stop bleeding. Platelets have no nucleus, gather at damaged blood vessels, and stick together to form an early blood clot. They are made in the bone marrow from larger cells called megakaryocytes.

## Platelet Aggregation

The clumping of platelets at an injured blood vessel to form an early plug that helps stop bleeding.

## Platelet Aggregation Test

A laboratory test measuring platelet clumping in response to agonists such as ADP, collagen, or epinephrine, used to evaluate platelet function disorders and medication effects.

## Platelet Antibodies Blood Test

A blood test detecting antibodies directed against platelet antigens, used selectively in suspected immune thrombocytopenia, neonatal alloimmune thrombocytopenia, or platelet-transfusion refractoriness.

## Platelet Count

The number of platelets in a given volume of blood, used to assess bleeding or clotting risk.

## Platelet Function Assay

An automated in vitro laboratory test measuring primary platelet-related hemostasis under high shear stress conditions, determining the closure time required for platelets in citrated whole blood to occlude an aperture coated with collagen/epinephrine or collagen/ADP.

### Alternate Names:

PFA-100

## Platelet Function Disorders

Conditions in which platelets do not adhere, activate, or aggregate normally, causing impaired primary hemostasis. They may be inherited or acquired and can cause easy bruising, nosebleeds, heavy menstrual bleeding, or bleeding after procedures.

## Platelet Plug Formation

The first step in stopping bleeding after a blood vessel is injured. Platelets attach to the damaged area, become active, and clump together to form a temporary plug.

### Alternate Names:

Primary Hemostasis; Platelet Aggregation

## Platelet-Rich Plasma

A preparation made by concentrating platelets from a person's own blood and injecting it into a selected tissue. It releases growth factors and is used in some musculoskeletal, wound, and hair-loss treatments, but evidence and benefit vary by condition and technique.

## Platelets

Small cell fragments made in the bone marrow that circulate in the blood and help stop bleeding. They attach to damaged blood vessels, become activated, and join together to form an initial plug.

### Alternate Names:

Blood Platelets; Coagulation Fragments

## Platybasia

An unusually flattened base of the skull. It may cause no symptoms, but in some people it narrows the space around the lower brain or spinal cord and contributes to headache, imbalance, weakness, or other nerve problems.

## Platypnea

Dyspnea that occurs in the upright position and is relieved by lying down. It is the opposite of orthopnea. It is rare and may be associated with hepatopulmonary syndrome, intracardiac shunting, or pulmonary arteriovenous malformations.

## Platypnea-Orthodeoxia Syndrome

A clinical syndrome characterized by dyspnea (platypnea) and arterial oxygen desaturation (orthodeoxia) that are precipitated or worsened when assuming an upright seated or standing posture and relieved in the recumbent (supine) position, typically caused by a posture-dependent intracardiac right-to-left shunt through a patent foramen ovale

---

or pulmonary AVMs.

**Alternate Names:**

Orthodeoxia; Postural Shunt Hypoxemia

**Platysma**

A thin superficial muscle of the neck that tenses the skin and assists lower-jaw and lower-lip movement.

**Pleiotropic Cytokine**

An immune-signaling protein that affects several types of cells or produces more than one effect, such as changing inflammation, blood-cell production, immune activity, or tissue repair.

**Pleiotropy**

The ability of one gene change to influence several traits or body systems. For example, one inherited disorder may affect the brain, skin, blood, and other organs at the same time.

**Pleomorphic**

Pleomorphic means having multiple forms or marked variation in cell or tissue appearance; in pathology, it may be a feature of malignancy but is not diagnostic alone.

**Pleomorphic Adenoma**

A usually noncancerous tumor of a salivary gland, most often the parotid gland near the ear. It usually grows slowly as a painless lump, but long-standing tumors can rarely become cancerous.

**Pleomorphic Adenoma Of The Vulva**

An exceptionally rare benign mixed epithelial and mesenchymal tumor of the vulva, resembling pleomorphic adenoma of salivary tissue and requiring histopathologic confirmation.

**Pleomorphic Adenoma Pathology**

A usually benign salivary-gland neoplasm containing variable epithelial and myoepithelial elements in a chondromyxoid, myxoid, or cartilaginous-appearing stroma. Incomplete excision and long duration increase recurrence or malignant-transformation risk.

**Pleomorphic Dermal Sarcoma**

An uncommon high-grade cutaneous sarcoma, usually on chronically sun-damaged skin of older adults, composed of markedly pleomorphic cells and capable of local recurrence or metastasis.

**Pleomorphic Xanthoastrocytoma**

A circumscribed astrocytic tumor, often arising superficially in the cerebral hemispheres of younger patients and associated with seizures. Most are low grade, but anaplastic transformation can occur.

**Pleomorphism**

Variation in the size and shape of cells or their nuclei within a tissue. Marked pleomorphism is a feature of many high-grade malignancies but can also occur in some reactive or degenerative conditions.

**Plethoric**

Plethoric means unusually full, congested, or ruddy, often describing the appearance associated with increased blood volume or red-cell mass.

**Plethysmography**

A test that measures changes in the size of a body part or the amount of blood or air it contains. It can assess circulation in the limbs, lung volume, or changes in blood flow during breathing or exercise.

**Pleura**

A thin, two-layered lining surrounding each lung and covering the inside of the chest wall. A small amount of fluid between its layers allows the lungs to move smoothly during breathing.

**Pleural**

Relating to the pleura, the membranes surrounding the lungs and lining the inside of the chest.

**Pleural Cavity**

The narrow space between the two slippery layers surrounding each lung. A small amount of fluid normally allows smooth breathing, while extra air, blood, pus, or fluid can restrict lung expansion.

---

## **Pleural Decortication**

An operation that removes a thick scar-like layer from the lung's covering so a trapped lung can expand again. It may be used for long-standing infected fluid, blood, or inflammation around the lung when other treatment is insufficient. Risks include bleeding, infection, air leakage, and breathing complications.

### **Alternate Names:**

Decortication of the Lung; Pulmonary Decortication; Thoracoscopic Decortication

## **Pleural Effusion**

An abnormal buildup of fluid in the space between a lung and the chest wall. It may result from heart failure, infection, inflammation, cancer, or injury, and can cause shortness of breath, cough, or chest pain.

## **Pleural Fluid Analysis**

Laboratory testing of fluid collected from around the lung. Cell counts, protein, acidity, glucose, cultures, and other tests help determine whether the fluid is caused by infection, cancer, heart failure, inflammation, or another problem.

## **Pleural Fluid Culture**

Culture of pleural fluid to detect bacteria, fungi, or mycobacteria in suspected pleural-space infection; specimens should be collected appropriately, often in blood-culture bottles.

## **Pleural Fluid Glucose**

Pleural-fluid glucose may be reduced in empyema, complicated parapneumonic effusion, rheumatoid pleuritis, tuberculosis, and malignancy. It is interpreted with pH, cell count, protein, LDH, microbiology, imaging, and the clinical setting.

## **Pleural Fluid Gram Stain**

A pleural-fluid Gram stain looks for bacteria in fluid around the lung and may support evaluation of empyema or bacterial pleural infection.

## **Pleural Fluid PH**

Pleural-fluid pH helps assess infection and other causes of effusion. A markedly low pH supports a complicated parapneumonic effusion in the appropriate clinical setting, but drainage decisions also depend on imaging, fluid appearance, microbiology, glucose, and the patient's condition.

## **Pleural Fluid Smear**

Microscopic examination of fluid removed from the space around the lung. It may show inflammatory cells, bacteria, fungi, cancer cells, or other material and is interpreted with fluid chemistry, cultures, imaging, and symptoms.

## **Pleural Friction Rub**

A scratchy or grating sound heard through a stethoscope when inflamed layers around the lung rub together during breathing. It may occur with infection, a blood clot in the lung, or other causes of pleural inflammation.

## **Pleural Mesothelioma**

A cancer arising from the pleura, the thin lining around the lungs and inner chest wall. It is strongly associated with asbestos exposure and may cause chest pain, breathlessness, or fluid around the lung.

## **Pleural Needle Biopsy**

A pleural needle biopsy removes a small sample of the pleural lining through a needle for evaluation of cancer, infection, or inflammation.

## **Pleural Neoplasm**

A pleural neoplasm is an abnormal growth in the lining around the lung, such as mesothelioma or metastatic cancer.

## **Pleural Perforation**

A breach in the pleural lining allowing air or fluid to enter the pleural space, potentially causing pneumothorax or hydropneumothorax, often from trauma, procedures, or underlying lung disease.



---

## Pleural Plaque

A usually harmless area of thickened or calcified tissue on the lining around the lung, most often caused by past asbestos exposure. It is not the same as asbestos-related lung scarring.

## Pleural Rub

A rough, creaking or grating auscultatory sound heard during both inspiration and expiration, caused by the friction between roughened, inflamed visceral and parietal pleura. It is heard in pleuritis, pulmonary infarction, or pneumonia and ceases when the patient holds their breath.

### Alternate Names:

Pleural Friction Rub

## Pleural Sac

The potential space and enclosing pleural membranes around a lung, normally containing a small amount of lubricating fluid.

## Pleural Space

The narrow potential space between the lung's inner covering and the chest wall's lining. It normally contains a small amount of lubricating fluid; air, blood, or excess fluid can impair breathing.

## Pleural Tuberculosis

Tuberculous infection or inflammation of the pleura, often producing a lymphocyte-rich pleural effusion, chest pain, fever, and breathlessness and requiring microbiologic or tissue evaluation.

### Alternate Names:

Tuberculous Pleural Effusion; Tubercular Pleural Effusion; Pleural Effusion, Tubercular; Tuberculous Pleurisy

## Pleurectomy

Surgical removal of part or all of the pleura, used in the treatment of malignant pleural mesothelioma and recurrent malignant pleural effusion.

## Pleurisy

Inflammation of the thin layers around the lungs, often causing sharp chest pain that worsens with

deep breathing or coughing. Causes include viral infection, pneumonia, a blood clot, autoimmune disease, or injury; associated features vary with the underlying cause.

### Alternate Names:

Pleuritic Chest Pain; Pleural Inflammation; Pleuritis

## Pleuritic Pain

Sharp chest pain that worsens with breathing or coughing, often caused by inflammation of the pleura or nearby tissues.

## Pleuritis

Alternate name; see *Pleurisy*.

## Pleurodesis

A procedure that makes the two layers around a lung stick together so air or fluid cannot collect repeatedly. It may use a chemical or surgery and can cause pain, fever, infection, or breathing complications.

## Pleurodynia

Severe, repeated pain in the chest or upper abdomen caused by inflammation of breathing muscles, often during infection with coxsackievirus. Pain may worsen with breathing or movement and usually resolves as infection improves.

## Pleuropulmonary Blastoma

A rare aggressive childhood cancer arising in the lung or pleura, usually associated with DICER1-related tumor predisposition.

## Plexiform Fibrohistiocytic Tumor

A rare low-grade malignant fibroblastic and histiocytic tumor, usually arising in the dermis or subcutis of the upper extremities of young people; local recurrence is common and metastasis is uncommon.

## Plexiform Neurofibroma

A plexiform neurofibroma is a diffuse nerve-sheath tumor, strongly associated with neurofibromatosis type 1, that can grow along multiple nerve branches

---

and sometimes become malignant.

**Alternate Names:**

PN PNF

**Plexus**

A network formed by intersecting nerves, blood vessels, lymphatic vessels, or other structures. Examples include the brachial nerve plexus and the celiac vascular plexus.

**Pliability**

Pliability is the ability of a material or tissue to bend, stretch, or deform without breaking; it is relevant to skin, vessels, lungs, and medical devices.

**Plica**

A fold of tissue, especially a fold in the lining of a joint. Plicae are often normal, but an irritated plica around the knee can cause pain, clicking, swelling, or a catching feeling with movement.

**Plitidepsin**

Plitidepsin is an antineoplastic compound that targets eukaryotic elongation factor 1A and has been studied for selected cancers and viral infections.

**Ploidies**

Ploidy describes the number of complete chromosome sets in a cell, such as haploid, diploid, or polyploid; abnormal ploidy can characterize tumors or developmental disorders.

**PLS**

Alternate name; see *Primary lateral sclerosis*.

**Plummer-Vinson Syndrome**

A condition involving iron-deficiency anemia, difficulty swallowing, and thin webs in the upper swallowing tube. Treating the iron deficiency and stretching the web can improve swallowing and may reduce complications.

**Pluralibacter Gergoviae**

A gram-negative environmental bacterium that can contaminate medicines or cosmetics and rarely cause

opportunistic infection.

**Pluripotential**

Able to develop into many, but not every, type of cell in the body. Pluripotent stem cells can produce cells from the three early tissue groups and are studied for development, disease, and regenerative treatment.

**Plutonium**

A radioactive heavy metal whose inhalation or ingestion can cause internal radiation exposure and toxicity.

**Pm**

An abbreviation that may mean afternoon, post-mortem, or picometer; meaning depends on clinical context.

**PMBCL**

Alternate name; see *Primary mediastinal large B-cell lymphoma*.

**PMBL**

Alternate name; see *Primary mediastinal large B-cell lymphoma*.

**PMDD**

Premenstrual dysphoric disorder is a severe cyclic mood disorder occurring before menstruation with irritability, depression, anxiety, and functional impairment that improve after menses.

**PML**

PML commonly means progressive multifocal leukoencephalopathy, a JC-virus demyelinating brain infection during severe immune suppression, but it may have other meanings.

**PN**

Alternate name; see *Plexiform neurofibroma*.

**Pneumatic**

Relating to air or gas under pressure. Medical examples include an air-pressure ear examination, in-

---

flatable compression sleeves used to reduce clot risk, and an inflatable tourniquet used during some operations.

### **Pneumatic Otoscopy**

Examination of the eardrum with a lighted instrument while gently changing the air pressure in the ear canal. Reduced eardrum movement may suggest fluid behind it or poor middle-ear ventilation.

### **Pneumatic Retinopexy**

A procedure for selected retinal detachments in which an intravitreal gas bubble is injected and laser or cryotherapy seals the retinal break while positioning helps the bubble support the repair.

### **Pneumatic Tourniquet**

An inflatable cuff used to temporarily reduce blood flow to a limb during surgery or selected procedures.

### **Pneumatocele**

A thin-walled, air-filled space in the lung that often develops after infection, inflammation, or injury. Many resolve naturally, but large or infected spaces can impair breathing or require drainage or surgery.

### **Pneumatosis**

Gas trapped within the wall of an organ or other tissue, most often the bowel. It may be harmless, but when caused by poor blood flow, infection, or a tear, it can signal a life-threatening abdominal emergency.

### **Pneumatosis Cystoides Coli**

Multiple gas-filled cysts within the wall of the large intestine. They may be harmless and found by chance, or may signal bowel blockage, poor blood flow, infection, or another serious illness when pain or other warning signs occur.

### **Pneumatosis Cystoides Intestinalis**

Gas-filled cysts within the bowel wall. They may be harmless and found incidentally, but can also result from poor blood flow, infection, blockage, lung disease, or medicines. Severe pain, fever, or

shock may indicate bowel injury.

### **Pneumatosis Intestinalis**

Gas within the wall of the intestine, seen as small bubbles or lines on an imaging test. It may be harmless, but pain, fever, shock, or signs of poor blood flow may indicate bowel injury.

### **Pneumocephalus**

Air or another gas inside the skull, usually after a head injury, brain surgery, or a skull fracture near the sinuses. A large or trapped collection can press on the brain and cause headache, vomiting, confusion, or reduced alertness.

### **Pneumococcal Infection**

Infection caused by *Streptococcus pneumoniae*, ranging from ear or sinus infection to pneumonia, meningitis, or bloodstream infection.

### **Pneumococcal Infections (*Streptococcus pneumoniae*)**

Infections caused by *Streptococcus pneumoniae*, ranging from otitis and sinusitis to pneumonia, bloodstream infection, and meningitis. Risk is increased by young or older age, absent spleen, chronic disease, smoking, or immune compromise; vaccination and prompt treatment reduce complications.

### **Pneumococcal Meningitis**

A dangerous infection of the brain and spinal-cord coverings caused by *Streptococcus pneumoniae*. It can cause fever, headache, stiff neck, confusion, seizures, hearing loss, or death and requires immediate treatment.

### **Pneumococcal Pneumonia**

An acute bacterial infection of the lung parenchyma caused by *Streptococcus pneumoniae*, the single most common cause of community-acquired pneumonia globally. It classically presents with sudden high fever, shaking chills (rigors), pleuritic chest pain, productive cough with rust-colored sputum, and lobar consolidation on chest imaging.

#### **Alternate Names:**

*Streptococcus Pneumoniae* Pneumonia; Pneumo-

---

coccal Lung Infection

### **Pneumococcal Sepsis**

Sepsis caused by *Streptococcus pneumoniae*, often arising from pneumonia, meningitis, or another invasive pneumococcal infection.

### **Pneumoconiosis**

A group of work-related lung diseases caused by breathing mineral or industrial dust over time. Dust can scar lung tissue and cause cough, breathlessness, reduced oxygen, and progressive breathing disability.

### **Pneumocystis**

A genus of fungi that includes *Pneumocystis jirovecii*, which can cause pneumonia in people with weakened cellular immunity.

### **Pneumocystis Infection**

Infection by *Pneumocystis jirovecii*, most often causing pneumonia with dry cough, fever, and progressive shortness of breath in immunocompromised people.

### **Pneumocystis Jirovecii Pneumonia**

Opportunistic pneumonia caused by *Pneumocystis jirovecii*, especially in people with advanced HIV infection or other cellular immunodeficiency. It causes progressive exertional dyspnea, dry cough, fever, and hypoxemia.

### **Pneumocystis Pneumonia**

Opportunistic pneumonia caused by *Pneumocystis jirovecii*, usually in people with impaired cell-mediated immunity. It may cause progressive exertional dyspnea, dry cough, fever, and hypoxemia with diffuse ground-glass changes. Diagnosis and treatment require clinical, microbiologic, and radiologic assessment; trimethoprim-sulfamethoxazole is commonly used.

### **Pneumocyte**

A cell lining the air sacs of the lungs. Type I cells allow oxygen and carbon dioxide to pass between

air and blood, while type II cells make surfactant, which helps keep the air sacs open.

### **Pneumolysis**

Surgical separation or freeing of lung tissue or pleural adhesions.

### **Pneumomediastinum**

Air trapped in the central space of the chest between the lungs. It may follow severe coughing, an injury, a medical procedure, or a tear in the airway or swallowing tube and can cause chest pain or breathlessness.

### **Pneumonectomy**

Surgery to remove an entire lung. It is usually performed for selected lung cancers or severe one-sided lung disease when removing only part of the lung is not suitable.

### **Pneumonia**

An infection of one or both lungs that inflames the air sacs. The air sacs may fill with fluid or pus, making breathing harder. Pneumonia is commonly caused by bacteria or viruses and less often by fungi or parasites. Symptoms may include cough, fever or chills, chest pain, and shortness of breath.

### **Pneumonia in Immunocompromised Patients**

Lung infection in a person with impaired immune defenses from disease, medicines, transplantation, cancer, or other causes. The organism spectrum is broad, symptoms may be muted, and evaluation often requires early imaging, microbiologic testing, and tailored antimicrobial or supportive treatment.

### **Pneumonitis**

Inflammation of lung tissue, often caused by a medicine, radiation, an immune reaction, inhaled irritants, or material entering the lungs rather than by infection. It may cause dry cough and breathlessness.

---

## **Pneumopericardium**

Air around the heart, usually after a chest injury, surgery, a medical procedure, or a tear in the esophagus. A large or trapped collection can prevent the heart from filling and pumping properly and requires urgent assessment.

## **Pneumoperitoneum**

Air or gas inside the abdominal cavity around the organs. It may be expected after abdominal surgery, but unexpected gas can indicate a torn intestine and requires urgent assessment, especially with pain or illness.

## **Pneumoplasty**

Surgery to repair or reshape part of the lung or an airway. It may be used to remove a damaged area, improve airflow, or restore the shape of tissue after injury or disease.

## **Pneumotachometer**

An instrument that measures the speed and amount of air moving during breathing. It can help assess airflow limitation and is used in lung-function testing, often together with other measurements.

## **Pneumothorax**

Air in the space between the lung and chest wall (pleural space), which can partly or completely collapse the lung. It may happen spontaneously, after an injury, or after a medical procedure and can cause sudden chest pain or breathlessness; severe symptoms need emergency care.

## **Pneumovirinae**

Pneumovirinae is a subfamily of viruses that includes respiratory pathogens such as respiratory syncytial virus and human metapneumovirus.

## **Pneumovirus**

A group or genus of respiratory viruses; human pneumovirus terminology often refers to human metapneumovirus or related paramyxoviruses.

## **PNF**

Alternate name; see *Plexiform neurofibroma*.

## **Pocket Erosion**

Loss of tissue at the margin or base of a periodontal or surgical pocket, potentially exposing deeper structures and promoting infection, bleeding, or impaired healing.

## **Podagra**

A sudden gout attack in the big-toe joint, causing intense pain, redness, warmth, and swelling.

## **Podiatrist**

A healthcare professional specializing in diseases and injuries of the feet, ankles, and related lower limbs. Care may include examination, wound treatment, medicines, footwear advice, procedures, and prevention of complications.

## **Podiatry**

The healthcare field focused on the feet, ankles, and related lower-limb conditions. It includes prevention, diagnosis, treatment, wound care, mobility support, and management of problems linked with diabetes or circulation.

## **Podocyte**

A specialized cell covering the tiny blood vessels that filter blood in the kidney. Its delicate extensions help keep proteins in the bloodstream; damage can cause protein to leak into urine.

## **Podofilox**

Podofilox is a topical antimitotic medicine used to treat selected external anogenital warts caused by human papillomavirus; irritation and correct application are important.

## **Podophyllin**

Podophyllin is a plant-derived resin with antimitotic activity formerly used topically for warts; it can be severely toxic if absorbed and is generally replaced by safer products.

---

## Podophyllum

A plant genus containing podophyllotoxin, used in some topical wart treatments but toxic if misused.

## Podosome

A temporary attachment structure on the surface of some cells, including immune cells and bone-removing cells. It helps the cell grip tissue and break down nearby material while moving or carrying out its function.

## POEMS Syndrome

A rare disorder linked to abnormal plasma cells that affects several body systems. It may cause nerve damage, enlarged organs, hormone problems, skin changes, fluid buildup, eye swelling, and unusual bone changes.

### Alternate Names:

Crow-Fukase Syndrome

## Poikilocytosis

The presence of red blood cells with unusually varied shapes on a blood smear. It may occur with anemia, inherited blood disorders, or damage to red blood cells.

## Poikiloderma

A skin pattern combining mottled pigmentation, visible small vessels, and epidermal thinning, seen in connective-tissue disease, chronic sun damage, inflammation, and some cutaneous lymphomas.

## Poikilotherm

An organism whose body temperature changes with surroundings, unlike humans, who normally maintain a relatively stable temperature.

## Poinsettia Plant Exposure

Chewing poinsettia usually causes mild mouth irritation, nausea, or stomach upset, not severe poisoning. Rinse the mouth and contact poison control if symptoms are severe, a large amount was swallowed, or another product was involved.

## Point

A specific location, spot, or value; in medicine it may refer to a tender point, pressure point, or measurement point.

## Point Mutation

A change affecting one DNA building block. Depending on its location, it may have no effect, change the protein made by a gene, stop protein production early, or contribute to an inherited condition or cancer.

## Point-Of-Care Testing

A test performed near the patient, such as in a clinic, ambulance, home, or hospital ward, instead of a central laboratory. Results are available quickly, but trained users, quality checks, and clinical interpretation remain essential.

## Point-Of-Care Ultrasound

Ultrasound performed and interpreted at the bedside by the treating clinician to answer a focused question or guide a procedure. It can assess the heart, lungs, abdominal fluid, blood vessels, and bladder, but does not replace a complete imaging study when one is needed.

## Poison

A substance that can cause injury, illness, or death when enough enters the body. Harm depends on the substance, amount, route, timing, and person's health.

## Poison-Center Consultation

Specialist toxicology advice used to identify hazards, interpret exposure and laboratory data, select antidotes, guide observation or disposition, and coordinate public-health response. Consultation is appropriate for significant, uncertain, mixed, pediatric, or unusual exposures.

## Poison Control

A specialist service that gives immediate advice after exposure to a medicine, chemical, plant, or other possible poison. They assess the substance and amount and advise on first aid, observation,



---

emergency care, or further treatment.

### Poison Control Emergency Services

Specialized regional 24-hour telephone triage and emergency clinical toxicological consultation services staffed by toxicologists and specialist nurses to guide immediate management of toxic ingestions, overdoses, bites, and chemical exposures.

#### Alternate Names:

Poison Helpline; Poison Control Hotline; Toxicological Emergency Consultation

### Poisoning

Harmful effects caused by swallowing, inhaling, injecting, or absorbing a toxic substance. The effects depend on the substance, amount, route, timing, and person's health; inducing vomiting can worsen some poisonings.

### Poisoning First Aid

Basic actions used after exposure to a toxic substance, such as leaving contaminated air and removing chemicals from skin or eyes. The correct response depends on the substance and route of exposure; inducing vomiting or giving food can worsen some poisonings.

### Poison Ivy Rash

An itchy allergic contact dermatitis caused by exposure to urushiol, an oil found in poison ivy and related plants. The rash commonly appears as red streaks or patches with swelling, small bumps, or blisters; the pattern and severity vary with the amount and route of exposure.

### Pokeweed Poisoning

Eating pokeweed, especially its roots or unripe parts, can cause burning in the mouth, vomiting, diarrhea, weakness, low blood pressure, or seizures. Significant ingestion requires immediate poison-control or emergency advice.

### Poland Syndrome

A congenital condition characterized by absence or underdevelopment of part of the pectoral muscles on one side, sometimes accompanied by abnormalities

of the hand on the same side.

### Polar

Relating to a pole or direction, or describing a molecule with slightly different electrical charges at its ends. In biology and chemistry, polarity affects how substances dissolve, move across cell membranes, and interact with other molecules.

### Polarity

The arrangement or direction of positive and negative electrical, chemical, or biological properties. Cell polarity helps organize tissue structure and directs processes such as movement, division, absorption, and secretion.

### Polarography

Polarography is an electrochemical analytical method that measures current while voltage changes at an electrode to identify or quantify reducible substances.

### Pol Gene

The HIV gene encoding enzymes required for viral replication, including protease, reverse transcriptase, and integrase; antiretroviral drugs target these products.

### Polidocanol

Polidocanol is a detergent sclerosant injected to close selected varicose veins or vascular lesions by damaging the vessel lining and promoting fibrosis.

### Polio

An infection caused by poliovirus, also called poliomyelitis. It spreads mainly when virus from an infected person's stool enters the mouth, and it can also spread through respiratory secretions. Most infected people have no symptoms or have a short, mild illness with fever, sore throat, tiredness, nausea, headache, or stomach pain. Fewer than 1% of infections cause the virus to affect the brain or spinal cord, leading to meningitis, muscle weakness, or acute flaccid paralysis, which is often uneven between the limbs and may become permanent. Paralysis of the muscles used for breathing can be fatal.

---

Vaccination prevents polio.

### **Poliomyelitis**

Alternate name; see *Polio*.

### **Poliosis**

A localized patch of white or depigmented hair caused by reduced melanin in follicles, occurring in vitiligo, alopecia areata, inflammatory disease, genetic syndromes, or after injury.

### **Poliovirus**

An enterovirus that causes poliomyelitis. It is transmitted mainly by the fecal-oral route; most infections are asymptomatic, but a small proportion cause meningitis or acute flaccid paralysis by damaging motor neurons in the spinal cord. Diagnosis uses molecular testing, usually reverse-transcription PCR of stool or other appropriate specimens.

### **Pollakisuria**

Pollakisuria is unusually frequent urination, often in small amounts, caused by irritation, infection, stones, bladder disorders, anxiety, or other urinary conditions.

### **Pollakiuria**

Frequent urination, often in small amounts. It may occur with urinary infection, irritation, diabetes, pregnancy, medicines, or anxiety; in some children it is a temporary benign pattern, but pain, fever, blood, thirst, or weight loss needs assessment.

### **Pollen**

Tiny plant grains released for reproduction that can trigger allergies, asthma symptoms, or eye inflammation.

### **Pollen-Food Allergy Syndrome**

An allergic reaction to certain raw fruits, vegetables, or nuts in people sensitized to related pollen proteins. It usually causes itching or tingling of the mouth and throat and is also called oral allergy syndrome.

### **Pollex**

The anatomical term for the thumb, the first and most mobile digit of the hand. It has two bones and can touch the fingertips, allowing precise grasping; injury or arthritis can cause pain and loss of hand function.

### **Pollution**

Contamination of air, water, soil, or surroundings by substances that may harm health or ecosystems.

### **Poly A**

A sequence of adenine nucleotides added to the end of many messenger RNA molecules, helping regulate stability, transport, and translation.

### **Polyadenylation**

Addition of a polyadenosine tail to the 3' end of many eukaryotic messenger RNAs, promoting stability, nuclear export, and efficient translation.

### **Polyamine**

A small positively charged molecule, such as putrescine, spermidine, or spermine, required for cell growth and nucleic-acid function.

### **Polyamine Catabolism**

The enzyme-controlled breakdown of polyamines, natural substances involved in cell growth and function. This process produces chemicals that can affect cell signals, oxidative stress, inflammation, development, and cancer biology.

### **Polyarteritis**

Inflammation of several arteries, which can reduce blood flow and damage organs. Causes and effects vary; polyarteritis nodosa is a rare form that mainly affects medium-sized arteries and may cause fever, muscle pain, skin changes, nerve problems, or abdominal pain.

### **Polyarteritis Nodosa**

A rare disease in which inflammation damages medium-sized arteries. It can reduce blood flow to the skin, nerves, kidneys, muscles, or intestines,

---

causing fever, weight loss, nerve pain or weakness, skin changes, high blood pressure, or abdominal pain.

**Alternate Names:**

PAN; Kussmaul-Maier Disease

**Polyarthrititis**

Inflammation involving multiple joints, commonly defined as five or more. It is a clinical pattern rather than a single diagnosis and may result from autoimmune, infectious, crystal-related, or other inflammatory disorders.

**Polyarticular Joint Pain**

Pain affecting multiple joints, caused by systemic inflammatory disease, infection, osteoarthritis, crystal disease, medication, or other conditions. The pattern, duration, swelling, and associated symptoms guide evaluation.

**Polybrominated Biphenyl**

A group of brominated industrial chemicals that can persist in the environment and cause toxic effects after exposure.

**Polychlorinated Biphenyl**

A group of persistent industrial chemicals that can accumulate in body fat and are associated with cancer, endocrine, immune, and neurologic effects.

**Polychondritis**

A relapsing immune-related disease that inflames and damages cartilage. It may affect the ears, nose, joints, voice box, windpipe, or inner ear, causing pain, deformity, breathing problems, hearing loss, or balance symptoms.

**Polychromasia**

The presence of unusually bluish, immature red blood cells on a blood smear. It usually means bone marrow is releasing cells quickly, often because of blood loss, red-cell destruction, or treatment response.

**Polychromatophilia**

Polychromatophilia is bluish-gray staining of red blood cells on a blood smear, reflecting increased circulating immature reticulocytes from stimulated red-cell production.

**Polycystic Kidney Disease**

An inherited disorder in which many fluid-filled cysts grow in the kidneys, which may enlarge and gradually lose function. The main forms are autosomal dominant polycystic kidney disease, usually recognized in adulthood, and autosomal recessive polycystic kidney disease, which can affect infants or children. Complications may include high blood pressure, pain, blood in the urine, kidney failure, or cysts in the liver.

**Alternate Names:**

PKD

**Polycystic Liver Disease**

An inherited condition in which many fluid-filled cysts develop in the liver. The cysts may enlarge the liver and cause abdominal fullness or pain; liver function is often preserved, but kidney cysts may also occur.

**Polycystic Ovarian Syndrome**

A hormonal and metabolic disorder involving ovulatory dysfunction and androgen excess, often causing irregular periods, acne, excess hair, infertility, or metabolic risk.

**Polycystic Ovary**

Polycystic ovaries contain many small follicles and may occur as a normal variant or with polycystic ovary syndrome. PCOS additionally involves ovulatory dysfunction, androgen excess, irregular periods, infertility, acne, or metabolic risk and requires clinical—not ultrasound alone—diagnosis.

**Alternate Names:**

Polycystic Ovarian Morphology; PCOM

**Polycystic Ovary Syndrome**

A common hormonal and metabolic condition that affects ovulation and the levels or effects of certain hormones. It may cause irregular or absent periods,

---

reduced or absent ovulation, acne, extra facial or body hair, scalp hair thinning, or difficulty becoming pregnant. It may also cause higher levels or effects of androgens, hormones found in everyone but usually present at higher levels in males. Some people have many small follicles in the ovaries, but this finding is not required for the condition. Insulin resistance, weight gain, darkened thickened skin, type 2 diabetes, and other metabolic problems may also occur. In adults, the diagnosis is based on at least two of three features—ovulation problems, androgen excess, or polycystic-appearing ovaries—after other causes have been considered.

### **Polycythaemia**

An increased red-cell mass or concentration, which may be primary or secondary to another condition.

### **Polycythemia**

An abnormally high red-cell concentration or total red-cell amount. It may result from a bone-marrow disorder, low oxygen, dehydration, smoking, medicines, or another illness and can increase the risk of blood clots.

### **Polycythemia Vera**

A long-term bone-marrow cancer that produces too many red blood cells and sometimes too many white cells and platelets. The thicker blood can cause headaches, itching, clots, bleeding, or an enlarged spleen.

#### **Alternate Names:**

Primary Polycythemia; Polycythaemia Vera

### **Polydactyly**

A birth condition in which a person has one or more extra fingers or toes. The extra digit may be on the thumb or big-toe side, the little-finger or little-toe side, or between the usual digits. It may occur alone or as part of a genetic syndrome. Treatment depends on its size and function and may involve reconstructive surgery.

### **Polydipsia**

Excessive thirst that leads to drinking unusually large amounts of fluid. Common causes in-

clude uncontrolled diabetes, water-balance disorders, medicines, and some mental health conditions.

### **Polyene**

A molecule containing multiple carbon-carbon double bonds; polyene medicines such as amphotericin B act against fungal cell membranes.

### **Polyethylene**

A plastic polymer used in packaging and medical devices; implanted polyethylene components may wear over time and produce foreign-body reactions.

### **Polyethylene Glycol**

A polymer used in medicines, especially laxatives that draw water into stool to ease constipation.

### **Polygenic**

Influenced by variants in many genes, often together with environmental factors. Height, blood pressure, and risk for many common diseases are polygenic traits.

### **Polygenic Disease**

A disease influenced by small effects from many genes together with environment and lifestyle. Examples include common diabetes, high blood pressure, heart disease, and some cancers; risk is inherited, not certainty.

### **Polygenic Hypercholesterolemia**

Elevated low-density lipoprotein cholesterol caused by the combined effect of multiple genetic variants and environmental factors rather than one dominant pathogenic variant. Levels and cardiovascular risk vary, and secondary causes should also be assessed.

### **Polygenic Inheritance**

Inheritance in which many genes, each with a small effect, together influence a trait or disease risk. Height, skin color, blood pressure, diabetes risk, and heart-disease risk are examples; environment also contributes.

---

### **Polygenic Obesity**

Obesity influenced by many small genetic differences together with eating patterns, activity, sleep, medicines, and the surrounding environment.

### **Polygenic Risk Score**

An estimate of a person's inherited tendency toward a disease, calculated from many small genetic differences. It changes risk assessment but does not predict certainty and must be considered with family history, environment, and other health factors.

### **Polyglandular Autoimmune Syndrome**

An autoimmune disorder in which two or more endocrine glands are damaged, causing combinations of thyroid, adrenal, parathyroid, pancreatic, or gonadal dysfunction.

### **Polyglandular Autoimmune Syndrome Type 1**

An inherited autoimmune syndrome, usually beginning in childhood, involving chronic mucocutaneous candidiasis and autoimmune damage to the parathyroid and adrenal glands. Other organs may be affected, and the exact combination varies.

### **Polyglandular Autoimmune Syndrome Type 2**

An autoimmune syndrome, usually beginning in adulthood, characterized by autoimmune adrenal insufficiency together with autoimmune thyroid disease and/or type 1 diabetes. Other autoimmune disorders may coexist.

### **Polygraphy**

Polygraphy is the simultaneous recording of several physiologic signals, such as breathing, heart rate, oxygen level, and movement, often during sleep testing.

### **Polyhydramnios**

An unusually large amount of amniotic fluid during pregnancy. It may cause abdominal discomfort, breathlessness, or swelling and can increase the risks of early birth, abnormal fetal position, cord problems, or heavy bleeding after delivery.

### **Polyketide Synthase**

An enzyme system that builds complex natural products from repeated carbon units, producing many antibiotics, toxins, and other biologically active molecules.

### **Polymastia**

The presence of extra breast tissue, sometimes with an additional nipple, along the embryonic milk line. Accessory breast tissue can enlarge or develop the same benign or malignant conditions as normal breast tissue.

### **Polymenorrhea**

Menstrual periods that occur unusually often, commonly at intervals shorter than about 21 days. Causes include hormonal changes, thyroid disease, fibroids, bleeding disorders, medicines, and the menopausal transition; persistent frequent bleeding needs evaluation.

### **Polymenorrhoea**

British spelling of polymenorrhea: menstrual periods occurring unusually often, commonly at intervals shorter than about 21 days.

### **Polymer**

A large molecule made of repeating smaller units called monomers; polymers are widely used in biology, medicines, and medical materials.

### **Polymerase Chain Reaction**

A laboratory method that makes millions of copies of a selected piece of DNA so it can be detected and studied. It is used to identify infections, inherited conditions, cancer-related gene changes, and other genetic material. The test must be chosen and interpreted for the specific question.

### **Polymerization**

Chemical joining of many small monomers into a larger polymer, a process important in biomolecules, plastics, dental materials, drug delivery, and laboratory techniques.

---

## Polymicrobial Infections

Infections caused by two or more kinds of germs at the same time. They commonly affect contaminated wounds, the abdomen, or the lungs and may require treatment directed at several different organisms.

## Polymicrogyria

A brain-development abnormality in which the outer brain surface has too many small, unusually formed folds. It may cause seizures, developmental delay, movement problems, or learning difficulties, depending on the areas involved.

## Polymorphic

Having more than one form or appearance. The term may describe a rash, crystal, cell, gene variant, heart rhythm, or tumor. Its medical importance depends on the structure being described and the other clinical findings.

## Polymorphic Eruption of Pregnancy

An itchy inflammatory eruption of pregnancy, usually beginning in abdominal stretch marks late in pregnancy and spreading to the trunk or limbs while sparing the umbilicus. It is uncomfortable but generally does not harm the fetus and usually resolves after delivery.

## Polymorphic Light Eruption

An itchy rash triggered by sunlight, often appearing in spring or after the first strong sun exposure. It may improve as the skin adapts, and sun protection reduces recurrences.

## Polymorphism

The presence of two or more inherited genetic forms or visible traits within a population.

## Polymorphonuclear Leukocyte

A white blood cell with a multilobed nucleus, especially a neutrophil, eosinophil, or basophil, involved in innate immune defense and inflammation.

## Polymorphous

Having lesions or clinical features that differ in

shape, size, or stage and therefore show multiple forms.

## Polymorphous Light Eruption

An acquired, usually seasonal photodermatosis in which ultraviolet exposure causes an itchy eruption of papules, plaques, vesicles, or other morphologies in susceptible people. It often appears hours to days after sun exposure, especially in spring or early summer, and may improve as the skin becomes repeatedly exposed.

### Alternate Names:

PMLE; Sun Poisoning; Polymorphic Light Eruption

## Polymyalgia Rheumatica

An inflammatory disorder causing aching and morning stiffness around the shoulders and hips in older adults.

### Alternate Names:

PMR; Senile Arteritic Gout

## Polymyositis

An uncommon immune-related disease causing gradual, usually symmetrical weakness in muscles close to the body, especially the shoulders, hips, and thighs. It can make climbing stairs, lifting, or swallowing difficult and may raise muscle-enzyme levels.

### Alternate Names:

PM; Idiopathic Inflammatory Myopathy

## Polymyxin

A group of antibiotics that disrupt bacterial cell membranes, used mainly for selected gram-negative infections.

## Polymyxin B

An antibiotic active against selected gram-negative bacteria; kidney and nerve toxicity limit its use to difficult infections.

## Polyneuritis

Inflammation or injury affecting multiple peripheral nerves, causing symmetric pain, numbness, weakness, or autonomic dysfunction; causes include immune disease, infection, toxins, and deficiency.



---

## **Polyneuropathy**

A generalized disorder affecting multiple peripheral nerves, often causing symmetric numbness, tingling, burning pain, weakness, or impaired reflexes in the limbs.

## **Polynucleotide**

A chain of linked nucleotides forming DNA, RNA, or a related laboratory molecule. Its sequence stores or carries genetic information and determines how cells make proteins or regulate biological processes.

## **Polyoma Virus**

A polyomavirus; some human polyomaviruses are widespread and cause serious disease mainly in immunocompromised people.

## **Polyomavirus Hominis 2**

A human polyomavirus designation associated with a virus found in people; clinically important disease is uncommon and mainly associated with immune suppression.

## **Polyp**

A growth projecting from a mucous membrane; it may be benign, precancerous, or rarely cancerous.

## **Polyp Biopsy**

A polyp biopsy removes part or all of a polyp for microscopic examination to determine whether it is benign, precancerous, or cancerous.

## **Polypeptide**

A chain of amino acids joined by peptide bonds. One or more folded polypeptide chains make up many proteins.

## **Polyphagia**

Unusually strong or persistent hunger that may lead to eating much more than usual. Causes include poorly controlled diabetes, an overactive thyroid, certain medicines, inherited conditions, and some mental-health disorders.

## **Polypharmacology**

The study or use of a medicine that affects several targets in the body. This can be helpful when one disease involves multiple pathways, but it may also increase side effects and medicine interactions.

## **Polypharmacy**

Use of several medicines at the same time, often five or more. Some combinations are necessary, but taking many medicines increases the risk of interactions, side effects, confusion, missed doses, and prescribing errors, especially in older adults.

## **Polyposis**

A condition in which many polyps develop, especially in the bowel or other digestive organs. Polyposis may be inherited or acquired, and some types increase the risk of colorectal or other cancers.

## **Polypropylene**

A synthetic plastic polymer used in medical supplies, sutures, meshes, and containers; implanted materials may rarely cause infection or foreign-body reactions.

## **Polyp Surveillance Colonoscopy**

A follow-up colon examination after polyps have been found or removed. The timing depends on the number, size, type, and microscopic features of the polyps and on the person's overall risk.

## **Polyradiculitis**

Inflammation or other disease affecting several spinal nerve roots. It may cause weakness, numbness, pain, and reduced reflexes; causes include infections, autoimmune disorders, and toxins.

## **Polyradiculopathy**

Disease affecting multiple spinal nerve roots, producing pain, weakness, sensory loss, or reflex changes in a distribution determined by the involved roots.

## **Polysaccharide**

A carbohydrate made of many linked simple sugar units. Examples include starch, glycogen, cellulose,

---

and some bacterial capsules.

### **Polysome**

A group of ribosomes attached to one messenger RNA molecule at the same time. Together they make many copies of the protein instructions, helping cells produce proteins efficiently.

### **Polysomnography**

An overnight sleep study that records brain activity, eye movements, muscle activity, heart rhythm, air-flow, breathing effort, and blood oxygen. It helps identify sleep-related breathing problems, unusual movements, seizures, and other sleep disorders.

### **Polysplenia**

A congenital condition with multiple small spleens and variable abnormalities of organ arrangement, heart, blood vessels, or intestinal rotation.

### **Polystyrene**

A synthetic plastic polymer used in containers and laboratory materials; burning it can release hazardous fumes.

### **Polytene Chromosome**

An enlarged chromosome formed by repeated DNA replication without cell division, displaying visible bands and commonly used to study gene activity in insect tissues.

### **Polythelia**

The presence of one or more additional nipples, usually along the embryonic milk line. It is often harmless, although accessory tissue can rarely have the same disorders as normal breast tissue.

### **Polythiazide**

A thiazide diuretic formerly used to treat high blood pressure and fluid retention by increasing renal salt and water excretion.

### **Polytrauma**

Serious injuries affecting two or more body regions or organ systems, usually caused by a major ac-

cident, fall, blast, or violence. The injuries may include bleeding, fractures, brain injury, chest injury, or damage to internal organs.

### **Polyunsaturated Fat**

A type of fat usually liquid at room temperature, found in fish, vegetable oils, nuts, and seeds. Some are essential because the body cannot make them.

### **Polyurethane**

A family of synthetic polymers used in medical tubing, implants, dressings, and other products; tissue response depends on formulation and site.

### **Polyuria**

Producing unusually large amounts of urine, commonly more than about three liters daily in an adult. Causes include diabetes, excess fluids, medicines, kidney problems, hormone disorders, and some inherited conditions.

### **Polyvalent**

Effective against or involving multiple types, strains, antigens, or targets.

### **Polyvinyl**

A family of polymers used in plastics and medical materials; properties vary according to attached chemical groups.

### **Pomalidomide**

An immunomodulatory medicine used with other treatments for selected multiple myeloma and related blood cancers; it can cause severe birth defects and blood abnormalities.

### **Pompe Disease**

A rare inherited disorder in which a missing enzyme causes stored sugar to build up inside muscle cells. Babies may develop severe weakness, an enlarged heart, and breathing or feeding problems; later-onset disease mainly weakens the limbs and breathing muscles. Enzyme replacement and supportive care can slow or improve the disease.

**Alternate Names:**

---

Glycogen Storage Disease Type II; Acid Maltase Deficiency; GSD II

### **Pompholyx**

A form of dyshidrotic eczema causing intensely itchy small blisters on the hands or feet.

### **Pompholyx Eczema**

Pompholyx, or dyshidrotic eczema, causes intensely itchy small blisters on the palms, sides of fingers, or soles and may leave painful peeling or cracks.

### **Pons**

The middle part of the brainstem, containing pathways and nuclei involved in movement, sensation, breathing, and cranial-nerve function.

### **Pontiac Fever**

A mild, self-limited flu-like illness caused by *Legionella* bacteria without the pneumonia typical of Legionnaires' disease.

### **Pontine**

Relating to the pons, a part of the brainstem between the midbrain and medulla. The pons helps carry movement and feeling signals and contains centers involved in eye movement, facial movement, hearing, balance, and breathing. A stroke or bleed there can cause weakness, loss of sensation, trouble speaking or swallowing, coma, or a locked-in state. Rapid correction of very low sodium can also damage it.

### **Pontine Myelinolysis**

A severe non-inflammatory demyelinating neurological disorder of the central pons caused by rapid, overly aggressive intravenous correction of chronic hyponatremia, presenting with spastic quadriparesis, pseudobulbar palsy, dysarthria, dysphagia, and the "locked-in" syndrome.

#### **Alternate Names:**

Central Pontine Myelinolysis; Osmotic Demyelination Syndrome; ODS

### **Poor Circulation**

Reduced or abnormal blood flow, usually referring to arterial insufficiency or venous disease, causing coolness, color change, pain, swelling, ulcers, or delayed wound healing.

### **Poor Color Vision**

Alternate name; see *Color blindness*.

### **Popcorn Lung Bronchiolitis Obliterans**

Bronchiolitis obliterans is irreversible scarring and narrowing of small airways that can follow toxic inhalation, infection, transplantation, or autoimmune disease. It causes persistent cough and breathlessness with obstructive lung testing; exposure avoidance and specialist care are important.

### **Popliteal**

Relating to the hollow behind the knee. This area contains the popliteal artery, vein, and nerves and can be affected by an artery aneurysm, a blood clot, or a Baker cyst.

### **Popliteal Aneurysm**

A weakened, bulging section of the artery behind the knee. It may form blood clots, block blood flow, or send clots into the lower leg, causing pain, coldness, numbness, or tissue damage. Larger aneurysms may need repair.

### **Popliteal Artery**

The main artery behind the knee. It continues from the thigh artery and divides into arteries supplying the lower leg and foot; blockage or injury can cause pain, coldness, numbness, or loss of circulation.

### **Popliteal Artery Aneurysm**

An abnormal enlargement of the artery behind the knee that may cause limb ischemia, clotting, or embolization.

### **Popliteal Artery Entrapment Syndrome**

An uncommon vascular cause of exertional lower extremity claudication in young, athletic individuals without cardiovascular risk factors, resulting from

---

an anomalous anatomical relationship between the popliteal artery and the medial head of the gastrocnemius muscle that compresses the artery during plantarflexion.

### **Popliteal Cyst**

A fluid-filled swelling behind the knee, also called a Baker cyst, caused by communication between the joint and a bursa and often associated with arthritis, meniscal injury, or inflammation.

### **Popliteal Fossa**

Diamond-shaped space behind the knee, bounded superolaterally by the biceps femoris, superomedially by the semimembranosus and semitendinosus, and inferiorly by the two heads of the gastrocnemius. It contains the popliteal artery and vein, tibial nerve, and common fibular nerve; the popliteal pulse is palpated with the knee slightly flexed.

### **Popliteal Vein**

A deep vein behind the knee formed by the union of the anterior and posterior tibial veins. It continues as the femoral vein and can be affected by deep-vein thrombosis.

### **Population Attributable Fraction**

The estimated proportion of disease in a whole population that could be prevented if a specific harmful exposure were removed, assuming the exposure truly causes the disease and other conditions stayed the same.

### **Population Attributable Risk**

The estimated amount of disease in a population linked to a specific exposure. It helps show the potential public-health benefit of reducing that exposure, but it depends on accurate data and a causal relationship.

### **Population Genetics**

The study of how gene frequencies and genetic variation change within and between populations through inheritance, mutation, selection, migration, and genetic drift.

### **Population Group**

A defined set of people sharing a characteristic such as age, location, exposure, ancestry, occupation, or health condition for epidemiologic or clinical analysis.

### **Porcelain Gallbladder**

Hard calcium deposits in the wall of the gallbladder, usually after long-standing inflammation or gallstones. It may cause no symptoms but can be associated with gallbladder disease; imaging findings guide discussion of monitoring or removal.

### **Porcine**

Porcine means derived from or relating to pigs; porcine tissues, enzymes, and biologic products may be used in medicine after appropriate processing and screening.

### **Pore**

A pore is a small opening or channel in a surface, membrane, tissue, or material that permits passage of fluid, gas, particles, or cellular contents.

### **Porencephaly**

A fluid-filled cavity within the brain caused by abnormal development or damage before or after birth. Effects vary with its size and location and may include seizures, weakness, movement problems, or developmental delay.

### **Porfimer Sodium**

A photosensitizing medicine used with light therapy to destroy selected abnormal tissues, including some tumors; it causes prolonged light sensitivity.

### **Porin**

A porin is a channel-forming protein in an outer membrane, especially of Gram-negative bacteria, allowing passage of selected small molecules.

### **Pork Tapeworm**

A parasite acquired by eating undercooked pork; its larvae can invade tissues and cause cysticercosis.

---

### Porokeratosis

A group of skin conditions that cause one or more ring-shaped, scaly patches with a raised border. Some forms can rarely develop into skin cancer, so changing, bleeding, or persistent lesions need dermatology assessment.

### Poroma

A usually noncancerous tumor arising from a sweat gland. It often appears as a flesh-colored, pink, or red lump on the palms, soles, or other skin and may be removed if painful, bleeding, or uncertain in diagnosis.

### Porphyria

A group of mostly inherited disorders of heme biosynthesis in which reduced activity of a pathway enzyme causes porphyrins or their precursors to accumulate. Acute hepatic porphyrias can cause episodic abdominal, neurologic, or psychiatric symptoms, while cutaneous porphyrias can cause photosensitivity and skin injury; some forms have both patterns. Porphyria cutanea tarda is usually acquired rather than inherited.

### Porphyria Cutanea Tarda

The most common porphyria, caused by reduced activity of hepatic uroporphyrinogen decarboxylase (UROD), leading to accumulation of porphyrins in the liver and skin. It causes fragile, blistering skin on sun-exposed areas. Associated factors include hepatitis C, excess iron, alcohol use, estrogen exposure, and certain genetic UROD variants.

### Porphyrias

Alternate name; see *Porphyria*.

### Porphyrin

A ring-shaped compound involved in heme and other biological molecules; abnormal accumulation occurs in porphyrias.

### Porphyrins Blood Test

A blood porphyrin test measures porphyrins or related compounds and helps evaluate suspected porphyria and disorders of haem production.

### Porphyrins Urine Test

A urine porphyrin test measures porphyrins or their precursors and helps investigate suspected porphyria or abnormal haem metabolism.

### Porphyromonas

A genus of anaerobic gram-negative bacteria found especially in the mouth and associated with periodontal and other infections.

### Porphyromonas Bennonis

An anaerobic oral bacterium rarely isolated from human specimens and occasionally associated with mixed infection.

### Porphyromonas Catoniae

An anaerobic gram-negative bacterium found in the mouth and occasionally associated with periodontal or mixed infection.

### Porphyromonas Endodontalis

A mouth bacterium that grows without oxygen and is associated with infected tooth pulp and root canals. It may contribute to dental infection, but diagnosis depends on symptoms, examination, imaging, and laboratory findings.

### Porphyromonas Levii

A *Porphyromonas* species found mainly in animals and rarely associated with human infection. An animal-associated bacterium rarely linked to human infection, usually in wounds or other opportunistic settings.

### Porphyromonas Somerae

A bacterium that grows without oxygen and is occasionally found in human wounds, abscesses, or other samples. When it causes infection, treatment depends on laboratory identification, infection location, and antibiotic susceptibility.

### Portacaval Shunting

A surgical or image-guided connection that diverts blood around the liver to lower dangerously high pressure in the portal veins. It may reduce bleed-

---

ing risk but can worsen confusion caused by liver disease.

### **Portal Circulation**

The flow of blood from the digestive organs and spleen through the portal vein to the liver before it returns to the general circulation.

### **Portal Hypertension**

Abnormally high pressure in the veins carrying blood from the digestive organs to the liver, most often because of cirrhosis. It can enlarge veins in the esophagus or stomach, cause abdominal fluid buildup, enlarge the spleen, or lead to bleeding.

**Alternate Names:**

PHT

### **Portal Hypertensive Colopathy**

Changes in the colon caused by high pressure in veins around the liver, usually from advanced liver disease. Enlarged fragile blood vessels may cause visible or hidden bleeding and anemia.

### **Portal Hypertensive Gastropathy**

Changes in the stomach lining caused by high pressure in the veins around the liver. The lining may look mottled and bleed slowly or suddenly, causing anemia or vomiting of blood in severe cases.

### **Portal System**

A network of veins that carries blood from the stomach, intestines, spleen, pancreas, and gallbladder to the liver. The liver processes nutrients and many absorbed substances before the blood returns to the general circulation.

### **Portal Vein**

The large vein that carries blood from the digestive organs, spleen, pancreas, and gallbladder to the liver. A clot or increased pressure in this vein can cause enlarged abdominal veins, bleeding, or fluid buildup.

### **Portal Vein Obstruction**

Partial or complete blockage of blood flow through

the portal vein or its branches, caused by thrombosis, tumor, compression, inflammation, or scarring. It may produce portal hypertension, enlarged veins, abdominal fluid, intestinal injury, or gastrointestinal bleeding.

### **Portal Vein Thrombosis**

A blood clot that partly or completely blocks the portal vein or one of its branches. It may occur with cirrhosis, cancer, abdominal infection or inflammation, surgery, or a clotting disorder. It can cause portal hypertension, enlarged veins, spleen enlargement, intestinal injury, or gastrointestinal bleeding.

### **Portion Of Cytosol**

A region of the cell's aqueous internal fluid outside membrane-bound organelles, containing ions, metabolites, proteins, and enzymes that support cellular reactions.

### **Portion Size**

The amount of food or drink served or eaten at one time, distinct from a standardized serving size and important for estimating energy, carbohydrate, sodium, and nutrient intake.

### **Portopulmonary Syndrome**

High blood pressure in the arteries of the lungs occurring with portal hypertension or serious liver disease. It can cause breathlessness, tiredness, chest discomfort, or fainting and affects decisions about liver treatment or transplantation.

### **Port-Wine Stain**

A permanent birthmark caused by widened small blood vessels, appearing as flat pink, red, or purple skin.

### **Position**

The location or arrangement of a body part, organ, fetus, or patient. Position can affect examination findings, movement, comfort, circulation, breathing, childbirth, medical procedures, and interpretation of imaging.



---

## **Positional Asphyxia**

Fatal or potentially fatal impairment of breathing caused by body position that restricts chest movement, airway patency, or venous return. Risk is increased by restraint, intoxication, obesity, and inability to reposition; findings are nonspecific and require scene and medical correlation.

## **Positive**

Positive means that a test, finding, result, or response is present or above a defined threshold; interpretation depends on the test and clinical setting.

## **Positive Airway Pressure Treatment**

Treatment that uses pressurized air delivered through a mask or similar device to keep the airway open during sleep or breathing difficulty. It is commonly used for obstructive sleep apnea; mask discomfort, dry airways, leaks, or skin irritation may occur.

## **Positive End-Expiratory Pressure**

Pressure kept in the airways at the end of breathing out, usually by a ventilator. It helps keep the lung's air sacs open and improve oxygen levels, but excessive pressure can injure the lungs or lower blood pressure.

## **Positive Feedback**

A body-control process in which a response strengthens the original change, making the process grow until a specific event is completed. Examples include contractions during labor and blood clotting.

## **Positive Finding**

A positive finding is an observed symptom, examination sign, image, laboratory result, or test response that supports the presence of a specified condition or feature.

## **Positive Predictive Value**

The chance that a person with a positive test result truly has the condition being tested for. It depends on test accuracy and how common the condition is among the people tested.

## **Positive Predictive Value In Surveillance**

The proportion of reported signals or alerts that represent true events meeting the surveillance case definition. It is influenced by the specificity of the signal and the underlying prevalence of the condition.

## **Positive Pressure Ventilation**

Breathing support that pushes air or oxygen into the lungs under pressure. It may be delivered through a mask or a breathing tube and can support or replace a person's own breathing during respiratory failure.

## **Positive Psychology**

The study of wellbeing, strengths, resilience, relationships, and factors that help people function and thrive. Its practices may support coping and quality of life but do not replace treatment for illness.

## **Positive-Strand RNA Virus**

A virus whose RNA can be read directly by an infected cell as instructions for making viral proteins. This allows the virus to begin producing new parts soon after entering the cell.

## **Positive Symptoms**

Symptoms reflecting an excess or distortion of normal mental functions, such as hallucinations, delusions, disorganized speech, or grossly disorganized behavior. They are contrasted with negative symptoms, which reflect loss or reduction of function.

## **Positron**

A particle with the same mass as an electron but a positive charge. Some medical tracers release positrons; when one meets an electron, energy is produced and detected during a PET scan.

## **Positron Camera**

A positron camera detects paired gamma rays from positron annihilation and is used in positron-emission tomography to map tracer distribution.

## **Positron Emission Mammography**

PET imaging of breast. Higher sensitivity than

---

mammography for dense breasts. Limited availability. A breast imaging test combining positron emission scanning with mammography to show tissue activity and structure.

### **Positron Emission Tomography**

An imaging test that uses a small amount of radioactive tracer to show tissue activity, such as glucose use, blood flow, or specific proteins. It is often combined with CT or MRI to locate abnormal areas more precisely.

### **Post-Acute Rehabilitation**

Time-limited interdisciplinary care after an acute illness, injury, or hospitalization to restore mobility, self-care, cognition, communication, and safe community function. Intensity and setting depend on medical stability and rehabilitation potential.

### **Post-Adoptive Depression**

Depression that begins after adopting a child. Stress, sleep loss, adjustment demands, past trauma, or unmet expectations may contribute; persistent sadness, loss of interest, or thoughts of self-harm need prompt mental-health assessment.

### **Postanesthesia Care Unit**

The postanesthesia care unit is a monitored recovery area where patients are observed during the immediate postoperative period. Admission assessment includes vital signs, level of consciousness, pain assessment, and surgical site evaluation.

### **Post-Cardiac Arrest Management**

Care after return of spontaneous circulation that supports breathing and circulation, identifies and treats the cause of the arrest, manages temperature and seizures when indicated, and assesses neurologic recovery over time.

### **Post-Cardiac Injury Syndrome**

Alternate name; see *Dressler syndrome*.

### **Post-Chemotherapy Cognitive**

### **Impairment**

Alternate name; see *Chemo brain*.

### **Postcholecystectomy Syndrome**

Persistence or recurrence of symptoms after cholecystectomy. Causes include retained common bile duct stones, sphincter of Oddi dysfunction, bile salt malabsorption, irritable bowel syndrome, and other gastrointestinal conditions. Evaluation includes liver function tests, MRCP, and endoscopic ultrasound.

### **Post Cibum**

A medication instruction meaning that a medicine should be taken after meals. Clear instructions should spell this out to prevent confusion with other uses of the abbreviation P.C.

#### **Alternate Names:**

PC; After Meals

### **Postcoital Bleeding**

Bleeding from the vagina after sexual intercourse. Causes include infection, dryness, injury, polyps, pregnancy-related conditions, or cancer.

#### **Alternate Names:**

Contact Bleeding; Postcoital Vaginal Bleeding

### **Post-Concussional Syndrome**

Persistent physical, cognitive, emotional, or sleep symptoms after a concussion or mild traumatic brain injury.

### **Post-Concussion Syndrome**

Ongoing symptoms after a concussion, such as headache, dizziness, tiredness, poor concentration, sleep problems, or mood changes. Symptoms usually improve gradually; evaluation and a stepwise return to activity help guide recovery.

### **Post-Dural Puncture Headache**

A headache that begins after a spinal tap or accidental puncture during an epidural. It is usually worse when sitting or standing and improves when lying down because spinal fluid has leaked; neck stiffness, nausea, ringing ears, or light sensitivity may occur.

---

## Posterior

Located behind or toward the back of the body or a body structure. The term is used to describe anatomical direction, position, movement, imaging findings, and areas affected by disease or injury.

## Posterior Chamber

The eye space between the iris and lens, filled with clear fluid that nourishes tissues and helps maintain pressure.

## Posterior Cord Syndrome

An incomplete spinal cord injury syndrome resulting from selective damage to the posterior columns (fasciculus gracilis and cuneatus), caused by posterior spinal artery infarction, tabes dorsalis, or subacute combined degeneration, characterized by loss of proprioception, vibration sense, and two-point discrimination below the lesion with preserved motor strength and pain/temperature sensation.

### Alternate Names:

Dorsal Column Syndrome; Posterior Column Myelopathy

## Posterior Cruciate

Usually referring to the posterior cruciate ligament, a strong knee ligament that limits backward movement of the shinbone. Injury can cause pain, swelling, and knee instability.

## Posterior Cruciate Ligament Injury

A stretch or tear of the ligament deep inside the knee that prevents the shinbone from moving too far backward. It may follow a blow to the shin, a fall on a bent knee, or severe twisting and can cause pain, swelling, and instability.

## Posterior Cruciate Ligament Tear

A partial or complete tear of the strong ligament deep inside the knee that limits backward movement of the shinbone. It often follows a forceful blow to the shin or a fall on a bent knee and may cause pain, swelling, and knee instability. Treatment depends on severity and other injuries.

## Posterior Descending Artery

A heart artery that runs along the back groove between the lower chambers and supplies part of the back and lower heart muscle. It usually branches from the right coronary artery but may arise from another heart artery.

## Posterior Drawer Test

A knee examination in which the lower leg is gently pushed backward while the knee is bent. Excessive backward movement may indicate injury to the ligament that stabilizes the back of the knee.

## Posterior Epistaxis

Nasal hemorrhage originating from arteries situated in the posterior nasal cavity or nasopharynx (most commonly branches of the sphenopalatine artery or Woodruff plexus), typically severe, profuse, draining down the oropharynx, and requiring posterior nasal packing or arterial embolization.

## Posterior Fossa Tumor

A growth in the lower back part of the skull, where the cerebellum and brainstem are located. It may cause headache, vomiting, unsteady movement, eye problems, or changes in breathing by pressing on nearby tissue or blocking brain fluid.

## Posterior Heel Bursitis

Inflammation of a bursa near the back of the heel, causing pain and swelling.

### Alternate Names:

Retrocalcaneal Bursitis; Achilles Bursitis

## Posterior Keyhole Foraminotomy

An operation through a small opening in the back that enlarges the passage where a spinal nerve leaves the spine. It relieves pressure from causes such as a disk bulge or arthritis and may reduce radiating pain or weakness.

## Posterior Mediastinum

The back part of the lower mediastinum, in front of the thoracic spine. It contains the esophagus, descending aorta, azygos veins, thoracic duct, and

---

nerves.

### Posterior Myocardial Infarction

A heart attack affecting the back wall of the heart, usually because a coronary artery is blocked. It may be missed on a standard ECG, so additional ECG leads or imaging may be needed when symptoms suggest it.

### Posterior Pituitary

The back part of the pituitary gland, which stores and releases hormones made in the brain. Vasopressin helps control body water and blood pressure, while oxytocin supports uterine contractions and release of breast milk.

### Posterior Prolapse

Alternate name; see *Posterior vaginal prolapse (rectocele)*.

### Posterior Reversible Encephalopathy Syndrome

Brain swelling caused by disturbed blood-flow regulation, often linked to very high blood pressure, pregnancy-related hypertension, kidney disease, severe infection, or immune-suppressing medicines. Headache, seizures, confusion, or vision loss require emergency treatment because prompt care can prevent lasting injury.

### Posterior Rhinorrhea

Posterior rhinorrhea is mucus draining from the nose into the throat, commonly causing throat clearing or cough from allergy, infection, irritation, or sinus disease.

### Posterior Subcapsular Cataract

A cloudy area at the back of the eye's lens, just beneath its outer capsule. Because it often lies in the path of light, it can cause glare, blurred vision, and difficulty reading, especially in bright light.

### Posterior Synechiae

Abnormal fibrous adhesions between the posterior surface of the iris and the anterior lens capsule, com-

monly formed during episodes of acute or chronic iritis and causing an irregular, non-reactive pupil.

### Posterior Tibial Artery

A major artery running down the back of the lower leg and behind the inner ankle. It supplies the lower leg and sole of the foot; its pulse is checked when assessing blood flow to the foot.

### Posterior Urethral Valves

Abnormal tissue folds present in the urethra of some male babies that block urine leaving the bladder. They can cause a weak urine stream, swollen kidneys, urinary infection, or kidney damage and require specialist treatment.

### Posterior Uveitis

Inflammation at the back of the eye involving the retina, the layer beneath it, or both. It may cause floaters, blurred or missing vision, or eye pain and can result from infection, immune disease, or another inflammatory condition.

### Posterior Vaginal Prolapse

Alternate name; see *Rectocele*.

### Posterior Vitreous Detachment

Separation of the jelly inside the eye from the retina, usually as an age-related change. It may cause new flashes or floaters; urgent eye examination is needed because a retinal tear or detachment can occur at the same time.

### Posteroanterior

Posteroanterior means directed from back to front; a PA chest radiograph uses this projection to reduce magnification of the heart.

### Post-Exposure Prophylaxis

Medicine or vaccination given after a possible exposure to prevent infection from developing. It is used for selected exposures to HIV, hepatitis B, rabies, and other infections; the correct treatment depends on the pathogen and must begin promptly.

---

### **Post-Finasteride Syndrome**

A controversial set of symptoms, chiefly sexual, that reportedly persist or begin after stopping treatment with finasteride or other 5-alpha-reductase inhibitors used for hair loss or prostate symptoms. Reported features include reduced libido, erectile or ejaculatory difficulties, decreased genital sensation, depression, anxiety, fatigue, and difficulty thinking. Its cause is uncertain and its existence is disputed; evaluation excludes hormonal, psychiatric, and other medical causes, and management is supportive.

### **Postganglionic Fiber**

An autonomic nerve fiber extending from a ganglion to its target organ after the synapse, carrying sympathetic or parasympathetic signals to smooth muscle, glands, or the heart.

### **Postherpetic Neuralgia**

Persistent nerve pain in an area affected by shingles after the rash has healed.

#### **Alternate Names:**

PHN; Shingles-Related Nerve Pain

### **Posthitis**

Inflammation of the foreskin, causing redness, swelling, itching, pain, discharge, or difficulty retracting it; infection, irritation, diabetes, and poor hygiene are possible contributors.

### **Post-Hoc**

Performed or analyzed after an event; in statistics, a post-hoc analysis follows the planned primary analysis.

### **Posthumous**

Occurring, discovered, or published after a person's death. A posthumous diagnosis may be based on records, stored samples, or autopsy findings.

### **Post-Infectious Cough**

A chronic cough persisting 3–8 weeks after a URI, caused by transient bronchial hyperresponsiveness and airway inflammation. It is self-limited and

usually resolves without treatment. Inhaled corticosteroids, antitussives, and time are the mainstays of management.

### **Postinflammatory Hyperpigmentation**

Darkening of skin after inflammation or injury, caused by increased melanin production or its deposition in the epidermis or dermis.

### **Post-Intensive Care Syndrome**

New or worsened physical, thinking, emotional, or mental-health problems that persist after critical illness or an intensive-care stay. It may include muscle weakness, poor memory or concentration, anxiety, depression, sleep problems, or post-traumatic stress.

### **Post-Kala-Azar Dermal Leishmaniasis**

A skin condition caused by *Leishmania donovani* that can develop after visceral leishmaniasis (kala-azar), often after apparent recovery. It occurs mainly in South Asia and East Africa and may cause pale or red patches, raised bumps, or nodules. Parasites can remain in the skin and be transmitted by sand flies in areas where the disease occurs.

#### **Alternate Names:**

PKDL; Post-Kala-Azar Dermatitis

### **Postmenopausal**

Postmenopausal describes the period after menopause, when ovarian estrogen production and menstrual periods have permanently ceased.

### **Postmenopausal Bleeding**

Vaginal bleeding after menopause. Even light or one-time bleeding needs medical evaluation because causes include tissue thinning, polyps, hormone effects, and cancer.

### **Postmenopausal Osteoporosis**

Bone loss caused by lower estrogen levels associated with menopause. Sometimes called type I osteoporosis.

### **Postmenopause**

The stage of life after menopause, when menstrual

---

periods have permanently stopped because the ovaries no longer release eggs regularly and produce much less estrogen.

### **Post-Mortem**

Occurring after death or relating to examination of a body afterward. A post-mortem examination can help determine the cause of death, identify disease, and provide information for families or legal investigations.

### **Postmortem Artifact**

A change that happens to a body after death and can look like disease or injury. Causes include decomposition, medical resuscitation, embalming, insects, animals, or movement of blood after death; examiners must distinguish it from changes before death.

### **Postmortem Changes**

Changes that occur after death, including cooling, livor mortis, rigor mortis, drying, autolysis, and putrefaction. Their timing varies with temperature, environment, body condition, and other factors and cannot alone establish an exact time of death.

### **Postmortem Interval**

The estimated time between death and the examination or discovery of a body. Investigators consider body temperature, stiffness, pooling of blood, chemical changes, environment, and insect activity, but the estimate is not exact.

### **Postmortem Lividity**

Purplish-red discoloration of dependent body parts following death, caused by gravitational pooling of blood in uncompressed capillaries and venules. Starts within 30 min–2 hours, becomes fixed after 6–12 hours.

#### **Alternate Names:**

Cadaveric Lividity; Postmortem Suggillation

### **Postmortem Toxicology**

Testing blood, tissues, or other samples after death for medicines, alcohol, poisons, or their breakdown products. Results must be interpreted with the

medical history, dose, sample site, decomposition, and possible movement of substances after death.

### **Postmortem Vitreous Analysis**

Laboratory examination of vitreous humor for electrolytes, glucose, ketones, drugs, alcohols, and other analytes. Vitreous potassium may provide a rough postmortem interval estimate, but results require cautious interpretation.

### **Post-Myocardial Infarction Pericarditis**

Inflammation of the sac around the heart that occurs after a heart attack. It may begin soon after the heart attack or develop later as part of a delayed immune reaction.

### **Postmyocardial Infarction Syndrome**

Alternate name; see *Dressler syndrome*.

### **Post-Nasal Drip**

Alternate name; see *Chronic Rhinitis*.

### **Postnatal**

Relating to the period after birth, especially the newborn and infant period. Postnatal care includes checking breathing, feeding, temperature, weight, jaundice, development, and signs of infection or other illness.

### **Postnatal Anxiety**

Excessive anxiety occurring after childbirth. It may involve persistent worry, panic, intrusive thoughts, restlessness, or physical symptoms and can be treated with therapy, support, and sometimes medicine.

### **Postnatal Care**

Health care for the parent and baby after birth. It includes checking bleeding, blood pressure, wounds, pain, feeding, mood, newborn growth, and warning signs, while providing guidance about recovery, contraception, vaccination, and follow-up.

### **Postnatal Depression**

Alternate name; see *Post-Natal Depression*.



---

## Post-Natal Depression

Depression beginning during pregnancy or after childbirth, with persistent low mood or loss of interest and possible effects on sleep, bonding, and daily functioning.

## Post-Obstructive Pneumonia

Lung infection developing beyond a blocked airway. A tumor, inhaled object, thick mucus, or narrowing can prevent drainage and allow infection; antibiotics may not cure it until the blockage is found and relieved.

## Postop

Postop is an abbreviation for postoperative, meaning relating to the period after a surgical procedure.

## Postoperative

Postoperative means occurring or provided after surgery, including recovery, wound care, pain control, rehabilitation, and complication monitoring.

## Post-Operative Atelectasis

Partial collapse of air sacs in the lung after surgery. Anesthesia, pain, shallow breathing, mucus, and lying still can contribute, causing shortness of breath or low oxygen. Pain control, movement, and breathing exercises may help prevent or treat it.

## Postoperative Care

Care after surgery that monitors breathing, circulation, pain, fluids, the wound, movement, eating, and possible complications. It may include early walking, exercises, medicines, and prevention of blood clots.

## Postoperative Cognitive Dysfunction

Problems with memory, attention, thinking, or planning after surgery. They may be temporary or persistent, especially after major operations; assessment compares the person with their usual abilities and checks for delirium, medicines, infection, or other causes.

## Postoperative Complications

Problems that develop after an operation, including infection, bleeding, blood clots, breathing problems, organ dysfunction, wound separation, or medication reactions. Risk depends on the procedure and the person's health; severe pain, fever, breathlessness, or heavy bleeding needs urgent care.

## Postoperative Corneal Edema

Swelling of the cornea after eye surgery, caused by endothelial dysfunction, inflammation, elevated intraocular pressure, or surgical trauma. It can blur vision and may resolve or persist depending on the injury.

## Postoperative Delirium

A sudden change in attention and thinking after surgery. A person may become confused, restless, unusually sleepy, or see things that are not there; symptoms may fluctuate and require prompt assessment for medical causes.

## Postoperative Hemorrhage

Bleeding after surgery that may occur at the operative site or internally and can cause shock, anemia, or the need for urgent intervention.

## Postoperative Ileus

Temporary slowing or stopping of bowel movement after surgery, especially abdominal surgery. It can cause bloating, nausea, vomiting, pain, and inability to pass gas or stool; medicines, electrolyte problems, and the operation may contribute.

## Postoperative Infection

An infection developing after surgery, involving the incision, an organ space, implanted material, or the bloodstream.

## Postoperative Nausea And Vomiting

Nausea, retching, or vomiting occurring after surgery. Risk is influenced by factors such as previous motion sickness or postoperative nausea, anesthetic technique, opioid use, and patient characteristics. Prevention and treatment are tailored to the individual risk and procedure.

---

## Postoperative Period

The postoperative period is the time after an operation when recovery, pain control, wound care, mobility, nutrition, complications, and follow-up are monitored.

## Postoperative Pulmonary Complication

A breathing problem developing after surgery, such as partial lung collapse, pneumonia, aspiration, worsening asthma or COPD, or respiratory failure. Risk is reduced by suitable pain control, movement, breathing exercises, and careful monitoring.

## Post-Orgasmic Illness Syndrome

A rare condition in which flu-like symptoms, extreme tiredness, headache, nasal or eye symptoms, and difficulty thinking begin soon after orgasm and last for days. The cause is uncertain; repeated episodes need medical assessment to exclude other conditions.

## Postpartum

Relating to the period after childbirth. Recovery includes changes in bleeding, pain, breasts, hormones, sleep, mood, and body function; important complications include heavy bleeding, infection, high blood pressure, blood clots, depression, and psychosis.

### Alternate Names:

Puerperium; Postnatal Period

## Postpartum Anxiety

Excessive, persistent, or distressing anxiety during pregnancy or after childbirth that interferes with sleep, daily function, bonding, or self-care. It may occur with panic, intrusive worries, or depression and deserves clinical assessment; thoughts of self-harm or harm to the baby require emergency help.

## Postpartum Blues

A common, short-lived period of tearfulness, mood changes, irritability, or anxiety beginning soon after childbirth and usually improving within two weeks. Support, rest, and practical help are important; persistent, severe, psychotic, or suicidal symptoms require prompt assessment for a postpartum mental-health disorder.

## Postpartum Care

Clinical care during the weeks after childbirth that assesses bleeding, uterine involution, blood pressure, infection, wound healing, breastfeeding, mental health, contraception, and recovery.

## Postpartum Contraception

Contraceptive counseling and care after childbirth, taking account of breastfeeding, venous-thromboembolism risk, medical conditions, fertility intentions, and the timing of method initiation.

## Postpartum Depression

Depression that begins during pregnancy or after childbirth and lasts beyond ordinary short-term mood changes. It may cause persistent sadness, loss of interest, guilt, anxiety, sleep or appetite changes, difficulty bonding, or thoughts of self-harm and needs treatment.

### Alternate Names:

PPD; Postnatal Depression

## Postpartum Endometritis

Infection of the inner lining of the uterus after childbirth, more common after cesarean birth. It may cause fever, lower-abdominal pain, uterine tenderness, fast heartbeat, or foul-smelling discharge and needs prompt treatment.

## Post-Partum Haemorrhage

Excessive bleeding after childbirth, most commonly caused by failure of the uterus to contract, retained placental tissue, genital-tract trauma, or clotting problems.

## Postpartum Hemorrhage

Excessive bleeding after childbirth, usually caused by poor uterine contraction, retained placental tissue, genital-tract trauma, or a clotting disorder. It can cause shock and requires urgent obstetric assessment and treatment.

## Postpartum Infection

An infection after childbirth, commonly involving the uterus, surgical wound, bladder, breasts, or

---

bloodstream. Fever, worsening abdominal or pelvic pain, foul-smelling discharge, wound redness, breast redness, or feeling very unwell requires prompt medical assessment.

### **Postpartum Infections**

Infections occurring after birth, including endometritis, wound infection, urinary infection, mastitis, and sepsis. Fever, worsening pelvic or wound pain, foul discharge, heavy bleeding, rapid heart rate, confusion, or weakness require prompt assessment.

### **Postpartum Preeclampsia**

New-onset hypertension with pre-eclampsia features after childbirth, sometimes occurring several days or weeks postpartum. Severe headache, visual symptoms, chest pain, shortness of breath, or seizures require urgent assessment.

### **Postpartum Psychosis**

A rare but severe psychiatric emergency that can begin suddenly during the days or weeks after childbirth. Symptoms may include hallucinations, delusions, paranoia, severe confusion, mania, depression, disorganized behavior, or rapidly changing moods. It can worsen quickly and may threaten the safety of the mother or baby, so immediate medical assessment is required.

#### **Alternate Names:**

Puerperal Psychosis; Postnatal Psychosis

### **Postpartum Thyroiditis**

Inflammation of the thyroid during the first year after childbirth, caused by an immune reaction. It may first cause anxiety, sweating, and a fast heart-beat, followed by tiredness, cold intolerance, or low thyroid function; recovery varies.

### **Postpartum Urinary Retention**

Inability to empty the bladder adequately after childbirth, sometimes associated with epidural anesthesia, perineal trauma, swelling, or prolonged labor. Prompt assessment prevents bladder overdistension and injury.

### **Postpericardiotomy Syndrome**

An immune-mediated post-cardiac-injury syndrome that develops days to weeks after heart surgery. It may cause fever, pleuritic or positional chest pain, shortness of breath, and fluid around the heart or lungs; symptoms can recur.

### **Post-Pneumonectomy Syndrome**

A rare problem after removal of one lung in which the remaining chest structures shift and press on the airway or major blood vessels. It may cause breathlessness, noisy breathing, repeated infections, or difficulty swallowing.

### **Post-Polio Syndrome**

Progressive muscle weakness, fatigue, and pain that begin many years after recovery from acute poliomyelitis. It typically affects muscles previously weakened by polio, and sometimes regions thought to have been spared, and is thought to result from slow decline of overworked surviving motor neurons. There is no specific diagnostic test, so other causes of new weakness are excluded first; management is supportive and includes adjusted exercise, energy conservation, and symptom control.

### **Post-Prandial**

Occurring after a meal. Post-prandial blood glucose helps assess diabetes, while post-prandial low blood pressure or chest pain may indicate a circulation or heart problem and needs assessment.

### **Postprandial Period**

The postprandial period is the interval after eating when digestion and nutrient absorption occur and glucose, insulin, and other metabolic responses change.

### **Postrenal AKI**

Sudden loss of kidney function caused by blocked urine flow. Prostate enlargement, stones, tumors, scarring, or bladder-nerve problems may be responsible; kidney damage becomes more likely when the blockage is severe or lasts a long time.

---

### Post-Splenectomy Complications

Post-splenectomy complications include infection by encapsulated bacteria, blood clots, bleeding, and changes in blood-cell counts; vaccination and follow-up reduce risk.

### Poststreptococcal Glomerulonephritis

Alternate spelling; see *Post-Streptococcal Glomerulonephritis*.

### Post-Streptococcal Glomerulonephritis

Kidney inflammation that can develop after a group A streptococcal throat or skin infection. It may cause dark urine, swelling, reduced urine, high blood pressure, tiredness, or kidney failure and needs medical evaluation.

### Poststreptococcal Pustulosis

A rare immune-related outbreak of pus-filled spots after a streptococcal throat infection. The spots usually appear on reddened skin of the trunk or limbs and should be assessed to confirm the cause and exclude infection or a medicine reaction.

### Post-Surgical Pain

Pain after an operation caused by tissue injury and inflammation. Its severity depends on the procedure and the person; uncontrolled pain can limit breathing, movement, sleep, and recovery and should be discussed with the care team.

### Post-Surgical Rehabilitation

A planned program to restore movement, strength, independence, and safe return to daily activities after surgery. It may include wound care, pain control, exercises, mobility training, and advice tailored to the operation and the person's recovery.

### Postsynaptic Membrane

The receiving surface of a nerve or muscle cell where a neighboring nerve releases a chemical signal. Receptors in this surface detect the signal and help start or change an electrical response.

### Postsynaptic Potential

A graded, local change in the electrical membrane potential of a postsynaptic neuron or muscle cell generated by the binding of neurotransmitters to ligand-gated receptors, classified as depolarizing excitatory postsynaptic potentials (EPSP) or hyperpolarizing inhibitory postsynaptic potentials (IPSP).

**Alternate Names:**

PSP

### Post-Term Infant

An infant born at or after 42 weeks of gestation, beyond the usual term-pregnancy period; risks depend on placental function, fetal condition, and circumstances of delivery.

### Post-Term Pregnancy

Pregnancy lasting 42 completed weeks or longer. The placenta may provide less support, the baby may become unusually large, and amniotic fluid may decrease. Clinicians monitor parent and baby and discuss delivery timing to reduce complications.

### Postterm Pregnancy

A pregnancy continuing to 42 weeks or longer. Risks may include reduced placental function, low amniotic fluid, fetal overgrowth, meconium passage, stillbirth, and difficult delivery, so fetal surveillance and delivery planning are considered.

### Post-Test Probability

The estimated chance that a person has a disease after a test result is known. It depends on the original likelihood of disease and how accurately the test distinguishes people with and without the condition.

### Post-Thrombotic Syndrome

Chronic pain, heaviness, edema, skin change, and sometimes ulceration developing after deep-vein thrombosis because of persistent venous obstruction and valvular damage. Compression, exercise, skin care, and selected interventions may reduce disability.

---

### Post-Transfusion Purpura

A rare immune reaction that causes a severe fall in platelets about 5–10 days after a blood transfusion. It can produce pinpoint skin bleeding, bruising, nosebleeds, or dangerous internal bleeding and requires urgent treatment.

### Post-Traumatic Headache

A headache beginning or continuing after a head or neck injury. It may resemble migraine or tension headache and can occur with dizziness, poor concentration, sleep problems, nausea, or light sensitivity.

### Post-Traumatic Stress Disorder

A mental-health disorder that may follow actual or threatened death, serious injury, sexual violence, or another traumatic event. It causes ongoing unwanted memories, nightmares, avoidance, emotional changes, and heightened alertness that interfere with daily life.

### Post-Traumatic Stress Disorder (PTSD)

Alternate name; see *Post-Traumatic Stress Disorder*.

### Postural

Postural means relating to body position or posture, including symptoms or vital signs that change when a person sits, stands, or lies down.

### Postural Drainage

Using specific body positions so gravity helps mucus move from smaller airways into larger ones, where it can be coughed out. A respiratory clinician selects positions based on the affected lung area and the person's condition.

### Postural Hypotension

A drop in blood pressure after standing that may cause dizziness, blurred vision, weakness, or fainting. It can result from dehydration, medicines, nerve disorders, blood loss, or heart problems and increases fall risk.

### Postural Orthostatic Tachycardia

### Syndrome

A disorder of the autonomic nervous system, which helps control heart rate and blood flow. When a person stands, the heart rate rises unusually quickly while blood pressure does not fall substantially, causing ongoing symptoms of poor tolerance of upright posture. In adults, the heart rate typically rises by at least 30 beats per minute within 10 minutes of standing; the increase is at least 40 beats per minute in adolescents. Symptoms may include dizziness, fainting or near-fainting, racing heartbeat, tiredness, weakness, headache, blurred vision, nausea, trouble thinking, or poor ability to exercise.

### Postural Tremor

An involuntary, rhythmic oscillation that appears when an affected body part is actively maintained against gravity, such as holding the arms outstretched. It is the characteristic tremor of essential tremor and physiologic tremor exaggerated by thyrotoxicosis, stress, or medications.

### Posture

The alignment and position of the body while resting or moving. Posture depends on bones, joints, muscles, nerves, vision, and balance and can affect comfort, breathing, movement, and injury risk.

### Post-Vasectomy Pain Syndrome

Persistent or recurring testicular or scrotal pain lasting at least three months after vasectomy. The pain may result from nerve irritation, inflammation, or pressure in the reproductive tract.

#### Alternate Names:

PVPS; Chronic Post-Vasectomy Pain

### Post-Void Residual

The amount of urine remaining in the bladder after urination, measured by ultrasound or catheterization. A persistently high residual may indicate incomplete emptying, but its significance depends on the volume, timing, symptoms, and clinical context.

### Post-Void Residual Volume

The amount of urine remaining in the bladder immediately after urination, measured by ultrasound



---

or catheterization. An elevated residual suggests incomplete emptying but must be interpreted with timing, bladder volume, symptoms, and repeated measurements.

### **Potassium**

The chemical symbol for potassium, a major intracellular electrolyte needed for nerve signaling, muscle contraction, and cardiac electrical activity.

### **Potassium Acetate**

A potassium salt used to replace potassium or provide acetate in selected medical and laboratory applications; dosing requires monitoring of kidney function and serum potassium.

### **Potassium Binders**

Medicines taken by mouth that trap potassium in the intestine so it leaves the body in stool. They help lower persistently high blood potassium in selected people, especially with kidney disease, but require monitoring for constipation, swelling, and other electrolyte changes.

### **Potassium Carbonate Poisoning**

Swallowing potassium carbonate can irritate or burn the mouth, throat, and stomach and may raise blood potassium dangerously. Vomiting, weakness, abnormal heartbeat, or breathing difficulty requires emergency care.

### **Potassium Channel**

A membrane protein that allows potassium ions to cross cells and helps control electrical excitability, heart rhythm, and hormone release.

### **Potassium Chloride**

A salt used to prevent or treat low blood potassium. It may be taken by mouth or given into a vein; too much can cause dangerous heart-rhythm problems, especially when kidney function is reduced.

### **Potassium Citrate**

A medicine that makes urine less acidic and increases urinary citrate, helping prevent some kidney

stones. It can dangerously raise blood potassium in people with kidney disease or those taking certain medicines.

### **Potassium Cyanide**

A highly poisonous chemical that prevents cells from using oxygen. Swallowing or inhaling it can rapidly cause headache, confusion, seizures, collapse, heart failure, and death and requires emergency treatment.

### **Potassium Deficiency**

An abnormally low blood potassium level, which can cause weakness, cramps, constipation, abnormal heart rhythms, or paralysis.

### **Potassium Dichromate**

A toxic, corrosive chemical that can burn the skin, eyes, mouth, and digestive tract. Swallowing or inhaling it may damage the kidneys, liver, blood, or lungs and requires urgent poison-control or emergency care.

### **Potassium Hydroxide Poisoning**

A severe chemical burn caused by swallowing or contacting potassium hydroxide, a strong alkali. It can damage the mouth, throat, stomach, skin, eyes, and lungs; rinse exposed areas and seek emergency help immediately.

### **Potassium In Diet**

Dietary potassium comes from fruits, vegetables, beans, dairy, and other foods; intake may need adjustment in kidney disease or with medicines that change potassium levels.

### **Potassium Iodide**

Potassium iodide supplies iodide and is used in selected thyroid conditions, radiation emergencies, and some expectorant preparations; excess can disturb thyroid function.

### **Potassium Ion**

The positively charged form of potassium, the principal intracellular cation that helps establish resting



---

membrane potential, nerve transmission, muscle contraction, and cardiac rhythm.

### **Potassium Permanganate**

A chemical sometimes used as a very dilute skin antiseptic or drying solution. Concentrated material can burn tissue, stain skin, and poison if swallowed, so it must be prepared and used carefully.

### **Potassium-Sparing Diuretic**

A diuretic that promotes sodium excretion while reducing potassium loss, such as spironolactone or amiloride.

### **Potassium-Sparing Diuretics**

Water tablets that help remove salt and water while limiting potassium loss. They are used for selected blood-pressure, heart, liver, or hormone conditions, but can raise potassium dangerously, especially in kidney disease or with certain medicines.

### **Potassium Test**

A potassium test measures potassium in blood, usually to assess heart, muscle, kidney, or medication-related electrolyte problems.

### **Potassium Urine Test**

A urine potassium test measures potassium excretion and helps determine whether abnormal blood potassium is related to kidney losses or other causes.

### **Potbellies And Toddlers**

A rounded abdomen in a toddler is often normal body shape or posture, but persistent distension with pain, vomiting, poor growth, constipation, or a mass needs medical assessment.

### **Potency**

The amount of a drug required to produce a specified effect. A more potent drug produces that effect at a lower dose, but potency does not indicate the maximum effect.

### **Potential**

Potential describes a capacity, possibility, or stored

energy that may become clinically relevant depending on circumstances and additional findings.

### **Potential Acuity Meter**

An eye-testing device that estimates the vision a person might achieve after treatment of an eye problem such as a cataract. Results can be limited by retinal, nerve, or other disease.

### **Potocki-Lupski Syndrome**

A genetic condition caused by an extra copy of a small section of chromosome 17. It may cause low muscle tone, feeding and growth problems, developmental or learning difficulties, autism-related features, behavior differences, and congenital heart defects.

#### **Alternate Names:**

PTLS; Duplication 17p11.2 Syndrome

### **Pott Disease**

Tuberculosis infection of the bones of the spine. It can cause persistent back pain, collapse or bending of the spine, an abscess, and pressure on the spinal cord leading to weakness or paralysis.

### **Potter Sequence**

A cascade of neonatal physical anomalies resulting directly from severe oligohydramnios in utero (frequently caused by bilateral renal agenesis or posterior urethral valves), comprising pulmonary hypoplasia, compressed dysmorphic facial features (Potter facies), and extremity limb positioning deformities.

### **Potter Syndrome**

A pattern of birth abnormalities caused by very low amniotic fluid, often because both fetal kidneys are absent or cannot make urine. It may cause underdeveloped lungs, distinctive facial features, and limb changes and is usually life-threatening.

### **Pott Puffy Tumor**

A subperiosteal abscess of the frontal bone associated with underlying frontal bone osteomyelitis, arising as a severe suppurative complication of untreated frontal sinusitis or direct cranial trauma,

---

presenting with fluctuant forehead swelling and headache.

**Alternate Names:**

Frontal Subperiosteal Abscess; Frontal Bone Osteomyelitis

**Pott's Disease**

Tuberculosis of the spine, causing vertebral destruction, back pain, deformity, or neurologic compromise.

**Pouch**

A sac-like body structure or a surgically created reservoir. Examples include a cheek pouch, a pouch in the digestive tract, and an ileal pouch made from small intestine.

**Pouch of Douglas**

A space in the pelvis between the uterus and rectum. In a person without a uterus, the comparable space lies between the bladder and rectum; fluid or blood can collect there.

**Pound**

A unit measuring weight or force, with meaning determined by context. In healthcare, pounds may describe body weight, pressure, equipment loads, or medicine instructions and should be distinguished from metric units.

**Povidone**

Povidone is a water-soluble polymer used in medicines, eye products, and other preparations; it is also part of povidone-iodine antiseptic.

**Povidone-Iodine**

Povidone-iodine is an iodine-containing antiseptic used on skin or tissue before selected procedures; it can irritate skin and affect the thyroid with extensive use.

**Powassan Encephalitis**

Inflammation of the brain caused by Powassan virus, a tick-borne flavivirus. It can cause fever, headache, vomiting, confusion, seizures, or weakness and may

require hospitalization.

**Powassan Virus**

A tick-borne virus that can cause encephalitis or meningitis. Severe infection may cause headache, fever, confusion, seizures, or weakness; there is no specific antiviral treatment.

**Powassan Virus Disease**

Powassan virus disease is a rare tick-borne infection that can cause fever, headache, vomiting, confusion, seizures, meningitis, or encephalitis. Treatment is supportive, and prevention relies on avoiding ticks and removing them promptly.

**Powder Diffraction**

An analytical method that identifies crystalline substances by measuring how an X-ray or other beam is diffracted by powdered material.

**Power**

The rate at which work is performed; in exercise, it combines force and movement speed.

**Power Training**

Exercise designed to improve the ability to produce force rapidly.

**Alternate Names:**

Explosive Strength Training; Ballistic Training

**Pox**

A disease producing vesicles, pustules, or pitted scars; the term is used in names such as smallpox and cowpox.

**Poxviridae**

A family of large DNA viruses that includes orthopoxviruses, molluscipoxvirus, and parapoxviruses; some cause significant human disease.

**Poxviridae Infection**

An infection caused by a poxvirus, ranging from localized skin lesions to severe systemic disease depending on the virus and host.

---

## Poxvirus

A family of large DNA viruses that includes variola, vaccinia, mpox, and molluscum contagiosum viruses.

## Ppar Alpha

A protein inside cells that helps control how the body uses fats. Fibrate medicines activate it to lower some blood fats, especially triglycerides.

## Ppar Delta

A protein inside cells that helps regulate fat use, energy balance, inflammation, and the development of some tissues. It is mainly studied in laboratory and drug research.

## PPD

PPD may mean purified protein derivative used in tuberculosis skin testing or post-partum depression; context determines the intended meaning.

## PPD Skin Test

A PPD skin test places a small amount of tuberculin under the skin to assess immune evidence of tuberculosis exposure; interpretation depends on risk and the size of the reaction.

## Practitioner

A person qualified and authorized to practice a healthcare profession.

## Prader-Labhart-Willi Syndrome

Alternate name; see *Prader-Willi syndrome*.

## Prader-Willi Syndrome

A genetic disorder that often causes weak muscle tone and feeding difficulty in infancy, followed later by unusually strong hunger, obesity, short stature, learning difficulties, hormone problems, and behavioral changes.

### Alternate Names:

Prader-Labhart-Willi Syndrome

## Pragmatic

Focused on what works in ordinary real-world con-

ditions. A pragmatic clinical trial studies how an intervention performs in usual healthcare rather than under highly controlled conditions.

## Pragmatic Trial

A pragmatic trial evaluates an intervention under usual real-world care conditions to estimate how well it works in routine practice.

## Pralatrexate

A folate-blocking chemotherapy medicine used for some relapsed or treatment-resistant peripheral T-cell lymphomas. It can cause mouth sores, low blood counts, infection, fatigue, nausea, and other serious treatment-related effects.

## Pralidoxime

An oxime that can reactivate inhibited acetylcholinesterase when administered early after organophosphate poisoning, particularly for nicotinic weakness and respiratory-muscle dysfunction. It is used with atropine and supportive care according to exposure and local guidance.

## Pramipexole

A medicine that stimulates dopamine receptors and is used mainly for Parkinson disease or restless legs syndrome. It can cause nausea, sleepiness, dizziness, sudden sleep, hallucinations, or impulse-control problems.

## Prandial

Prandial means relating to a meal, as in preprandial before eating or postprandial after eating.

## Prasugrel

An antiplatelet medicine that prevents blood cells from clumping. It is used in selected people with acute coronary syndrome receiving coronary stents, but it increases bleeding risk and is unsuitable for some patients.

## Pravastatin

A statin that lowers LDL cholesterol by inhibiting HMG-CoA reductase.

---

### **Pravastatin Sodium**

A statin medicine that lowers harmful LDL cholesterol by reducing cholesterol production in the liver. It helps reduce heart attack and stroke risk alongside diet, exercise, and other risk-factor treatment.

### **Praziquantel**

A medicine used to treat several infections caused by parasitic worms, including schistosomiasis, tapeworms, and flukes. It damages the worms so the body can remove them; dizziness, sleepiness, nausea, or abdominal discomfort may occur.

### **Prealbumin**

Another name for transthyretin, a protein made mainly by the liver that carries thyroid hormone and vitamin A in the blood. Its blood level changes with inflammation, illness, kidney disease, and nutrition, so it is not a nutrition measure by itself.

### **Preauricular Sinuses**

Small congenital pits or channels near the front of the ear caused by incomplete fusion during development. They may remain harmless or become infected, swollen, painful, or draining; recurrent infection may require surgical excision.

### **Prebiotics**

Food components, usually nondigestible fibers, that feed selected helpful bacteria in the large intestine. Sources include garlic, onions, leeks, asparagus, bananas, and whole grains; increasing them gradually may reduce gas.

### **Prebiotics And Probiotics**

Prebiotics are food components that feed certain helpful gut bacteria. Probiotics are live microorganisms found in some foods or supplements. Their effects depend on the specific product and health condition; they are not interchangeable treatments.

### **Pre-Cancerous**

Describing abnormal cells or tissue that are not cancer but have a higher-than-usual chance of becoming cancer. The risk varies by the tissue and the severity of change; monitoring or removal may

reduce the risk.

### **Precancerous Condition**

A precancerous condition contains abnormal changes that may progress to cancer but are not cancer themselves.

### **Precancerous Polyp**

A precancerous polyp is a growth with cellular changes that can become cancerous over time if not removed or monitored appropriately.

### **Precapillary Sphincter**

A ring of muscle at the entrance to some capillaries that helps control how much blood flows into a capillary network.

### **Precaution**

An action taken in advance to lower the chance of harm, infection, or error. Examples include hand-washing, protective equipment, checking a medicine, using a seat belt, and avoiding a known trigger.

### **Precipitate Labor**

An abnormally rapid labor and childbirth characterized by the expulsion of the fetus within less than three hours from the onset of regular, painful uterine contractions. It results from unusually intense and frequent uterine contractions (uterine tachysystole or hypertonicity) overcoming low maternal soft-tissue resistance in the birth canal, most commonly occurring in multiparous women or following excessive labor-inducing medications (such as oxytocin). Although brief, it carries significant maternal risks, including severe cervical and perineal lacerations, uterine rupture, amniotic fluid embolism, and acute postpartum hemorrhage from uterine exhaustion (atony). Major fetal and neonatal hazards include fetal distress and hypoxia from restricted placental blood flow during unrelenting contractions, intracranial trauma or hemorrhage from rapid skull compression and decompression, and risk of unassisted delivery (such as cord tearing or fall injury).

#### **Alternate Names:**

Precipitate Labour; Precipitous Labor; Precipitous

---

Labour; Tumultuous Labor; Rapid Delivery

### **Precipitating Factor**

A precipitating factor is an event or exposure that triggers the onset or worsening of symptoms or disease in a susceptible person.

### **Precision Medicine**

An approach that matches prevention or treatment to biological differences between people, often using genetic, molecular, or clinical information. It is related to but not identical to personalized medicine.

### **Precision Oncology**

An approach to cancer care that uses information about a tumor's genes, proteins, or other features to help select prevention, testing, or treatment. It aims to match treatment to the biology of the cancer rather than to its location alone.

### **Preclinical**

Before human testing, or describing disease changes that occur before symptoms become noticeable or cause recognized illness.

### **Precocious Puberty**

Development of puberty-related body changes earlier than expected. It may result from early brain signals, hormone-producing disorders, medicines, or other conditions and requires evaluation to protect growth and emotional wellbeing.

#### **Alternate Names:**

Early Puberty Premature Puberty

### **Preconception Care**

Health care before pregnancy to improve the chance of a healthy pregnancy. It reviews medical conditions, medicines, vaccinations, genetic risks, nutrition, folic acid, substance exposure, and lifestyle, and helps plan changes before conception.

### **Preconception Counseling**

Healthcare advice before pregnancy to improve the health of the pregnant person and future baby. It reviews medicines, vaccines, chronic diseases, nutri-

tion, genetic risks, harmful exposures, and recommended supplements.

### **Precordial Catch Syndrome**

Brief, sharp, localized chest pain near the heart that worsens with breathing and usually occurs in otherwise healthy children or adolescents.

### **Precordium**

The area of the front chest wall over the heart and nearby large blood vessels. A clinician examines this area by looking, feeling for the heartbeat or unusual movement, and listening for heart sounds and murmurs.

### **Precursor**

A substance that the body can change into another substance with a useful biological effect. For example, some vitamins, hormones, and medicines begin as precursors and are changed by enzymes in particular tissues.

### **Prediabetes**

Blood glucose higher than normal but below the diagnostic threshold for diabetes, usually identified by A1C, fasting glucose, or an oral glucose-tolerance test; it increases future diabetes and cardiovascular risk.

#### **Alternate Names:**

Borderline Diabetes; Impaired Glucose Tolerance; Impaired Fasting Glucose

### **Prediabetes Syndrome**

Blood glucose levels above normal but below the diabetes threshold, associated with increased future risk of type 2 diabetes and cardiovascular disease.

### **Predictive Test**

A test that estimates whether a person or disease is likely to respond to a particular treatment.

### **Predispose**

To predispose is to increase a person's susceptibility or likelihood of developing a disease or response when additional triggers occur.

---

### **Prednimustine**

Prednimustine is a former antineoplastic drug combining a corticosteroid-related group with an alkylating component and causing immunosuppression and marrow toxicity.

### **Prednisolone**

Prednisolone is a corticosteroid used to reduce inflammation and immune activity; prolonged or high-dose use can cause infection, bone loss, and other effects.

### **Prednisone**

Prednisone is a corticosteroid used for inflammatory, allergic, and autoimmune conditions; it should often be tapered after prolonged use.

### **Preeclampsia**

A disorder that can develop during pregnancy, usually after 20 weeks, or during the weeks after childbirth. It involves new high blood pressure together with protein in the urine or signs of organ injury. Possible findings include a low platelet count, abnormal kidney or liver function, severe headache, vision changes, upper-abdominal pain, or fluid in the lungs. During pregnancy, it can affect the placenta and fetal growth. It may progress to eclampsia, which is preeclampsia with seizures.

#### **Alternate Names:**

Pregnancy-Related Hypertension

### **Pre-eclampsia**

Alternate spelling; see *Preeclampsia*.

### **Preeclampsia and Eclampsia**

Alternate name; see *Pregnancy: Preeclampsia and Eclampsia*.

### **Preeclampsia With Severe Features**

A severe form of preeclampsia involving very high blood pressure or damage to organs such as the brain, liver, kidneys, lungs, or blood-clotting system. Severe headache, vision changes, breathing difficulty, or upper-abdominal pain requires emergency care.

### **Preexcitation Syndrome**

A heart-rhythm condition in which an extra electrical pathway activates the lower heart chambers too early. It can cause episodes of a very fast heartbeat, dizziness, or fainting and may require specialist assessment.

### **Pre-Existing Diabetes And Pregnancy**

Pre-existing type 1 or type 2 diabetes during pregnancy increases risks to parent and fetus, so glucose control, medicines, fetal growth, blood pressure, and eye and kidney health need coordinated care.

### **Prefrontal**

Relating to the anterior part of the frontal lobe involved in planning, judgment, working memory, and behavior.

### **Prefrontal Cortex**

The prefrontal cortex is the front part of the frontal lobe involved in planning, judgment, working memory, impulse control, and social behavior.

### **Pregabalin**

Pregabalin is used for certain nerve pain and seizures and may cause dizziness, sleepiness, swelling, or dependence.

### **Preganglionic Fiber**

A nerve fiber that carries automatic body signals from the brain or spinal cord to a relay station before they reach an organ. These signals help control the heart, blood vessels, glands, digestion, and bladder.

### **Pregnancy**

The period during which an embryo or fetus develops in the uterus, usually lasting about 40 weeks from the first day of the last menstrual period until birth.

### **Pregnancy And Herpes**

Genital herpes during pregnancy can affect delivery planning and rarely infect the newborn; antiviral treatment and examination of lesions near labor help reduce transmission risk.



---

## Pregnancy And Travel

Travel during pregnancy depends on gestational age, health, destination infections, transport, and access to care; prevention includes vaccines as appropriate, food and water safety, movement, and medical planning.

## Pregnancy and Urolithiasis

Kidney or ureteral stones during pregnancy can cause colicky flank pain, hematuria, nausea, obstruction, infection, or preterm contractions. Ultrasound is generally preferred for initial imaging, and treatment balances symptom control, drainage, infection management, and fetal safety.

## Pregnancy And Work

Work during pregnancy is usually possible but may require adjustments for heavy lifting, prolonged standing, heat, chemicals, radiation, infection, night work, or other occupational hazards.

## Pregnancy at Age 35 or Older

Pregnancy in a person who will be 35 years or older at the estimated time of delivery. It is associated with increased risks of some chromosomal conditions and pregnancy complications; individual risk depends on overall health and pregnancy history.

### Alternate Names:

Advanced Maternal Age

## Pregnancy: Bleeding During The First Trimester

First-trimester bleeding may occur with implantation, cervical changes, miscarriage, ectopic pregnancy, or other causes. Assessment uses symptoms, examination, serial pregnancy hormone testing, and ultrasound; heavy bleeding, severe pain, faintness, or shoulder pain is urgent.

## Pregnancy Complication

A maternal, fetal, placental, or labor-related problem that disrupts an otherwise expected pregnancy, such as preeclampsia, hemorrhage, diabetes, growth restriction, or preterm labor.

## Pregnancy Diagnosis

Confirmation of pregnancy using a urine or blood test for human chorionic gonadotropin, followed by ultrasound when needed to establish location, viability, and gestational age. A positive test with pain or bleeding requires assessment for ectopic pregnancy.

## Pregnancy-Induced Hypertension

New-onset hypertension during pregnancy without the proteinuria or organ dysfunction that defines preeclampsia. Diagnosis and treatment depend on repeated blood-pressure measurements, gestational age, symptoms, and evidence of maternal or fetal complications.

## Pregnancy Nutrition

Food and supplements during pregnancy that support the pregnant person and fetal growth. Needs vary, but a varied diet usually includes enough protein, folate, iron, iodine, calcium, vitamin D, and other nutrients; individual advice is important.

## Pregnancy Of Unknown Location

A positive pregnancy test without a confirmed intrauterine or ectopic pregnancy on ultrasound. Serial human chorionic gonadotropin measurements and repeat imaging help determine whether the pregnancy is early, failing, or ectopic.

## Pregnancy Outcome

Pregnancy outcome describes what happens to a pregnancy, such as live birth, miscarriage, stillbirth, preterm birth, or other maternal or fetal result.

## Pregnancy: Placenta Previa

Placenta previa occurs when the placenta lies over or near the cervical opening and can cause painless bright-red bleeding, especially later in pregnancy. Digital vaginal examination is avoided until location is known; recurrent bleeding requires obstetric planning and sometimes cesarean delivery.

## Pregnancy: Preeclampsia And Eclampsia

Preeclampsia is new hypertension during pregnancy with proteinuria or organ dysfunction; eclampsia

---

is the occurrence of seizures. Severe headache, visual symptoms, upper-abdominal pain, shortness of breath, or seizure requires emergency obstetric treatment.

**Alternate Names:**

Toxemia of Pregnancy; Hypertensive Disorders of Pregnancy

**Pregnancy-Related Hypertension**

Alternate name; see *Preeclampsia*.

**Pregnancy Status**

Pregnancy status indicates whether a person is pregnant, not pregnant, or of unknown status and can affect medication, imaging, and treatment decisions.

**Pregnancy Test**

A urine or blood test that detects human chorionic gonadotropin, a hormone made soon after an embryo attaches to the uterus. Testing too early can be falsely negative; an unexpected result may need repeat testing or medical assessment.

**Pregnancy Trauma**

Physical injury during pregnancy, including falls, motor-vehicle crashes, assault, or abdominal impact. Maternal stabilization comes first, followed by assessment for placental abruption, bleeding, uterine injury, fetal compromise, and preterm labor.

**Pregnancy Trimester**

One of three approximate stages of pregnancy: first through 13 weeks, second from 14 through 27 weeks, and third from 28 weeks until birth. The stages help organize fetal development, screening, and care.

**Pregnane**

Pregnane is a 21-carbon steroid hydrocarbon that forms the structural backbone of progesterone and several corticosteroid and progestin compounds.

**Pregnanolone**

Pregnanolone is a neuroactive steroid metabolite that can modulate GABA-A receptors and influence neuronal excitability, mood, and sedation.

**Pregnant**

Pregnant means carrying a developing embryo or fetus in the uterus after conception; pregnancy status affects medicines, imaging, and clinical decisions.

**Pregnenolone**

Pregnenolone is a steroid hormone precursor made from cholesterol that can be converted into progesterone, corticosteroids, and sex hormones.

**Prehn Sign**

A test in which lifting the painful testicle is said to relieve pain in epididymitis but not testicular torsion. This finding is unreliable and cannot safely rule out torsion. Sudden severe testicle pain, swelling, nausea, or a high-riding testicle requires urgent assessment, usually with Doppler ultrasound and immediate surgery when torsion cannot be excluded.

**Alternate Names:**

Prehn's Sign

**Prehypertension**

Older term; see *Elevated Blood Pressure*.

**Alternate Names:**

High-Normal Blood Pressure; Elevated Blood Pressure

**Preimplantation Genetic Diagnosis**

Genetic testing of embryos created through in-vitro fertilization before transfer to the uterus, used to identify selected inherited disorders or chromosomal abnormalities. It requires genetic counseling and does not guarantee implantation, a healthy pregnancy, or absence of all disease.

**Preimplantation Genetic Testing**

Genetic testing of cells from an embryo created through in-vitro fertilization before possible transfer to the uterus, used in selected situations to identify specific inherited disorders or chromosome-number abnormalities.

**Preload**

The amount of blood filling the heart's lower chambers and stretching their muscle just before each

---

heartbeat. It mainly depends on blood returning to the heart and influences how strongly the heart can pump.

### **Preload Reduction**

A decrease in the amount of blood returning to the heart, which lowers the heart's filling pressure and workload. It may occur with fluid loss or medicines and can help some forms of heart failure.

### **Premalignant**

Describing cells or tissue changes that have a higher chance of becoming cancer but have not grown into nearby tissues. Not every premalignant change becomes cancer; follow-up or removal depends on its type, location, size, and risk.

### **Premalignant Lesions of the Endometrium**

Abnormal endometrial gland growth, usually endometrial hyperplasia with atypia, that can precede endometrial carcinoma. Diagnosis requires sampling of the uterine lining; treatment depends on fertility goals, cancer risk, and whether invasion is present.

### **Premature**

Occurring earlier than expected or before the usual developmental or pregnancy-related time. A premature birth occurs before thirty-seven completed weeks and may increase risks involving breathing, feeding, temperature, and development.

### **Premature Adrenarche**

Early appearance of pubic or axillary hair, body odor, or mild acne from adrenal androgen production without true central puberty; evaluation excludes virilizing disorders.

### **Premature Atrial Contraction**

An early heartbeat that starts in the upper chambers of the heart rather than at the usual pacemaker. It may feel like a skipped or extra beat and is often harmless, but frequent episodes or fainting need assessment.

### **Premature Baby**

An infant born before 37 completed weeks of pregnancy, with risks related to immature lungs, brain, feeding, temperature control, immunity, and other organ systems.

### **Premature Birth**

Birth before 37 completed weeks of pregnancy; the earlier the birth, the greater the risks of respiratory, neurological, and developmental complications.

### **Premature Closure**

Closing of a body opening, growth plate, or developmental connection earlier than expected. The result depends on the site and may include blockage, abnormal body shape, or restricted growth; treatment depends on the cause.

### **Premature Ejaculation**

A persistent or recurrent pattern of ejaculation that occurs sooner than desired, with limited control and associated personal distress. Assessment considers the person's typical experience and sexual context.

#### **Alternate Names:**

Rapid Ejaculation; PE

### **Premature Infant**

An infant born before 37 completed weeks of pregnancy. Earlier birth increases the risk of breathing, feeding, temperature-control, infection, and developmental problems.

### **Premature Labor**

Regular uterine contractions with cervical change before 37 completed weeks of pregnancy, which can lead to preterm birth and requires assessment of membranes, infection, and fetal well-being.

### **Premature Menopause**

Loss of ovarian function and menstruation before age 40, now often termed primary ovarian insufficiency.

### **Premature Ovarian Failure**

Older name; see *Premature Ovarian Insufficiency*.

---

### Premature Ovarian Insufficiency

Reduced or intermittent ovarian function before age 40, causing irregular or absent periods, reduced fertility, and sometimes hot flashes, vaginal dryness, or low bone density. Causes include genetic conditions, autoimmune disease, chemotherapy, radiation, surgery, or no identifiable cause.

### Premature Puberty

Alternate name; see *Precocious puberty*.

### Premature Rupture Of Membranes

Breaking of the amniotic sac before labor begins. If it happens before 37 weeks, it is called preterm rupture; a gush or continuing leak of fluid needs prompt obstetric assessment because infection and early birth may follow.

### Premature Thelarche

Isolated breast development in a young child without other signs of puberty, often transient but assessed for growth acceleration, estrogen exposure, or true precocious puberty.

### Premature Ventricular Contraction

An early heartbeat that starts in a ventricle instead of following the heart's usual rhythm. It may feel like a skipped or extra beat and is often harmless, but frequent PVCs may occur with fainting, chest pain, or breathlessness.

**Alternate Names:**

PVC

### Premature Ventricular Contractions

Ectopic beats originating from the ventricles, appearing as wide, bizarre QRS complexes that occur earlier than expected. PVCs are common in the general population and are usually benign. Frequent PVCs (>10% of total beats) may cause left ventricular dysfunction.

**Alternate Names:**

PVCs; Premature Ventricular Complexes; Ventricular Extrasystoles

### Prematurity

Birth before 37 completed weeks of pregnancy. Prematurity can affect breathing, feeding, temperature control, immunity, brain development, and growth, especially when birth occurs very early.

### Premenstrual

Occurring during the days before menstruation; the term commonly refers to cyclical symptoms such as mood change, breast tenderness, bloating, headache, or fatigue.

### Premenstrual Breast Changes

Premenstrual breast changes include cyclic fullness, tenderness, swelling, or lumpiness caused by hormonal variation; a persistent one-sided mass, skin change, or bloody discharge needs evaluation.

### Premenstrual Dysphoric Disorder

A severe form of premenstrual symptoms that repeatedly causes depression, irritability, anxiety, mood swings, or loss of control along with physical symptoms and major disruption of work, relationships, or daily life before menstruation.

### Premenstrual Syndrome

A recurring group of physical and emotional symptoms in the days or weeks before menstruation that improves when bleeding begins. Symptoms may include irritability, mood changes, bloating, breast tenderness, headache, tiredness, or appetite changes.

### Pre-MiRNA

A pre-miRNA is an early hairpin-shaped microRNA transcript processed in the nucleus before becoming a mature regulator of gene expression.

### Premolar Tooth

One of the permanent teeth between the canine teeth and molars. Premolars usually have two raised points and help tear and crush food; decay, crowding, or infection may require dental treatment.

### Prenatal

Prenatal means occurring during pregnancy before

---

birth, including care, testing, development, counseling, and treatment of the pregnant person or fetus.

### **Prenatal Care**

Regular health care during pregnancy to monitor the pregnant person and developing baby. Visits may include blood-pressure and urine checks, blood tests, ultrasound, screening, vaccination, nutrition and medicine review, and advice about warning signs.

### **Prenatal Depression**

Depression during pregnancy. Symptoms may include persistent sadness, loss of interest, guilt, hopelessness, sleep or appetite changes, and thoughts of self-harm; prompt professional support is important.

### **Prenatal Development**

Prenatal development is growth and differentiation of the embryo and fetus from conception to birth, influenced by genes, nutrition, maternal health, and exposures.

### **Prenatal Diagnosis**

Testing during pregnancy to identify a possible genetic, chromosome, or structural condition in the fetus. It may use ultrasound, blood-based screening, or diagnostic tests such as chorionic-villus sampling and amniocentesis, depending on the clinical question and stage of pregnancy.

### **Prenatal Exposure**

Contact of a fetus with a medicine, substance, infection, or environmental factor during pregnancy. Effects depend on the agent, dose, timing, and fetal susceptibility.

### **Prenatal Genetic Counseling**

A consultation that reviews the risk of inherited or chromosomal conditions, available screening and diagnostic tests, possible results, and reproductive options before or during pregnancy.

### **Prenatal Testing**

Screening and diagnostic tests performed during pregnancy to assess fetal health and development or identify genetic, chromosomal, and structural conditions. Tests may include ultrasound, maternal blood tests, cell-free DNA screening, chorionic-villus sampling, and amniocentesis.

### **Prenatal Vitamins**

Supplements designed to provide nutrients needed during pregnancy, commonly including folic acid, iron, iodine, and vitamin D. The appropriate product and dose depend on diet, medical conditions, laboratory results, and pregnancy needs; excessive doses can be harmful.

### **Prenylation**

A cell process that attaches a fat-like group to a protein, helping direct where the protein works and how it interacts with other proteins. Abnormal prenylation can contribute to some inherited disorders and cancers.

### **Preop**

Preop is an abbreviation for preoperative, meaning relating to preparation and assessment before surgery.

### **Preoperative Assessment**

A medical review before surgery that checks illnesses, medicines, allergies, previous anesthesia problems, heart and lung function, and the airway. Tests are ordered when indicated, and correctable risks are addressed before anesthesia and the operation.

### **Preoperative Care**

Assessment and preparation before an operation. It may include reviewing illnesses and medicines, checking allergies, performing needed tests, explaining anesthesia and recovery, planning fasting and medicines, and reducing risks such as infection or blood clots.

### **Preoperative Evaluation**

Medical assessment before surgery to identify and reduce risks. It reviews health history, medicines,

---

allergies, physical findings, needed tests, anesthesia concerns, and preparations that improve safety during and after surgery.

### **Preoptic Area**

The preoptic area is a hypothalamic region involved in temperature regulation, sleep, reproductive behavior, and hormone control.

### **Prepubertal**

Prepubertal means before the physical and hormonal changes of puberty have begun or progressed to reproductive maturity.

### **Prepuce**

A fold of skin covering and protecting the end of an external sex organ. The penile prepuce is the foreskin over the head of the penis; the clitoral prepuce is the hood over the clitoris.

### **Prerenal Azotemia**

A rise in kidney waste products because the kidneys receive too little effective blood flow, without initial damage to the kidney tissue itself. Dehydration, blood loss, severe infection, heart failure, or liver disease may cause it and can lead to kidney injury if untreated.

### **Presbyacusis**

Alternate spelling; see *Presbycusis*.

### **Presbycusis**

Bilateral, usually symmetric age-related sensorineural hearing loss, initially affecting high frequencies and speech discrimination in noise. Management includes hearing rehabilitation, hearing aids, communication strategies, and cochlear implantation in selected severe cases.

### **Presbyopia**

The gradual loss of the eye's ability to focus on nearby objects with age. It commonly causes difficulty reading small print or doing close work and is corrected with reading glasses, contact lenses, or other prescribed lenses.

### **Presbyphonia**

Age-related changes in the voice caused by thinning or weakening of the vocal folds and changes in the voice box. It may produce a weak, breathy, rough, or quieter voice.

### **Preschooler Development**

Preschooler development includes rapid gains in language, motor skills, thinking, self-care, play, and social-emotional regulation; milestones vary, but loss of skills warrants assessment.

### **Preschooler Test Or Procedure Preparation**

Preparing a preschooler for a test or procedure uses simple explanations, play, comfort objects, and caregiver support while following any fasting or medicine instructions.

### **Prescription**

A written or electronic instruction from an authorized healthcare prescriber for a medicine, device, or treatment. It states what is needed and how to use it, allowing a pharmacy or service to provide care safely.

### **Prescription Drug Misuse**

Use of prescription medicines in a way not directed by a clinician, including taking someone else's medicine, taking extra doses, or using them for intoxication; dependence, overdose, and interactions may result.

### **Prescription Medications**

Pharmaceutical drugs that require authorization from a licensed healthcare provider before they can be dispensed to a patient. These medications are regulated because they carry risks that necessitate professional oversight, including potential side effects, drug interactions, and the need for precise dosing.

### **Young-Onset Dementia**

Dementia beginning before the usual older-adult age range. Modern assessment specifies the underlying disease and documents the age at onset.



---

**Alternate Names:**

Presenile Dementia (deprecated historical term)

**Presenilin**

Presenilin is a component of the gamma-secretase complex that processes membrane proteins; inherited variants can cause familial early-onset Alzheimer disease.

**Presentation**

The part of a fetus that is directed toward the birth canal, or the symptoms and signs with which a patient seeks care.

**Preseptal Cellulitis**

An infection of the eyelid and tissues around the eye, without infection inside the eye socket. It causes redness, swelling, pain, and fever; pain with eye movement, reduced vision, or a bulging eye may indicate deeper orbital infection and needs emergency assessment.

**Alternate Names:**

Periorbital Cellulitis

**Preservation**

Maintenance of tissue, cells, organs, or function by preventing deterioration.

**Preservative**

A substance added to food, medicine, or cosmetics to prevent microbial growth or chemical breakdown.

**Pressor**

Raising blood pressure, usually by tightening blood vessels or increasing heart output. Pressor medicines may treat dangerously low blood pressure but require monitoring because they can reduce circulation to some tissues.

**Pressoreceptor**

A pressure-sensitive receptor, especially a baroreceptor in the carotid sinus or aortic arch, that detects arterial stretch and helps regulate heart rate and blood pressure.

**Pressure**

Force applied over a particular area. Medical pressure measurements include blood pressure, pressure inside the skull, eye pressure, and airway pressure, and abnormal values can damage organs or impair function.

**Pressure Atrophy**

Thinning or wasting of tissue caused by prolonged pressure. Pressure can compress small blood vessels, reduce oxygen delivery, and gradually damage the affected tissue; the result depends on its location and duration.

**Pressure-Flow Study**

A bladder test that measures pressure while the bladder fills and while a person urinates, together with urine flow. It helps identify blockage or weak bladder muscle when difficulty emptying cannot be explained by symptoms alone.

**Pressure Injury**

Localized damage to the skin and/or underlying soft tissue caused by prolonged pressure, shear, or friction, most commonly over bony prominences.

**Pressure Sore**

Alternate name; see Bedsores (pressure ulcers). A localized injury to skin and underlying tissue caused by prolonged pressure or shear, also called a pressure ulcer.

**Pressure Support Ventilation**

A ventilator mode in which the machine gives extra pressure with each breath the person starts, making breathing easier and reducing the work of the breathing muscles. It is often used while reducing ventilator support.

**Pressure Ulcer**

Localized damage to the skin and underlying soft tissue resulting from sustained, unrelieved mechanical pressure, often compounded by shear forces and friction over bony prominences (most commonly the sacrum, ischial tuberosities, greater trochanters, and heels) in immobilized or debilitated individu-

---

als. Compressive occlusion of microvascular capillary beds produces localized ischemia, hypoxia, and tissue necrosis. Clinical severity is classified from Stage I (non-blanchable erythema of intact skin) through Stage IV (full-thickness skin and tissue loss with exposed fascia, muscle, tendon, or bone), along with unstageable wounds and deep tissue pressure injuries.

**Alternate Names:**

Pressure Sore; Decubitus Ulcer; Bedsore; Pressure Injury

**Pressure Ulcer Prevention**

Measures that reduce prolonged pressure and tissue injury, including repositioning, support surfaces, skin inspection, moisture control, nutrition, and mobility.

**Pressure Urticaria**

A form of chronic inducible urticaria in which sustained pressure causes localized swelling, erythema, and pain, usually after a delay of several hours. Lesions may last 8 to 72 hours and commonly affect the hands, feet, buttocks, or areas compressed by clothing or sitting.

**Pre-Synaptic**

Located before a synapse, especially on the nerve ending that releases a chemical messenger. Presynaptic activity controls how much signal reaches the next nerve cell, muscle, or gland.

**Presynaptic Terminals**

Presynaptic terminals are the ends of neurons that release neurotransmitters into synapses in response to electrical activity.

**Presyncope**

A feeling that fainting is about to occur without actually losing consciousness. It may cause lightheadedness, weakness, sweating, nausea, or blurred vision and can result from low blood pressure, dehydration, abnormal heart rhythm, medicines, or other illness.

**Preterm Birth**

Birth before 37 completed weeks of gestation. It may follow spontaneous labor or rupture of the membranes, or may be medically indicated because of maternal or fetal complications.

**Preterm Infant**

A baby born before 37 completed weeks of pregnancy. Earlier birth increases the chance of breathing, feeding, temperature-control, infection, and developmental problems; care depends on the baby's maturity, weight, and health.

**Preterm Labor**

Regular uterine contractions causing cervical change before 37 weeks gestation. Risk factors include prior preterm delivery, multiple gestation, infection, and cervical insufficiency. Diagnosis requires regular contractions with documented cervical change.

**Preterm Labor and Birth**

Regular uterine contractions causing cervical change before 37 completed weeks, potentially leading to preterm birth. Management depends on gestational age, membranes, infection, fetal status, and cause and may include corticosteroids, magnesium sulfate, antibiotics, or transfer to specialist care.

**Preterm Premature Rupture Of Membranes**

Breaking of the amniotic sac before labor begins and before 37 completed weeks of pregnancy. It may cause a gush or continuing leak of fluid and increases the risks of infection, early birth, and problems from too little amniotic fluid.

**Pretest Probability**

The estimated probability that a patient has a disease before the result of a diagnostic test is known. It is based on prevalence, symptoms, examination findings, risk factors, and prior test results and should guide test selection and interpretation.

**Pretibial Myxedema**

Thickened, firm skin over the shins caused by an immune reaction associated with Graves disease.

---

The skin may look swollen, waxy, or dimpled and usually does not leave a lasting indentation when pressed.

### **Pretibial Myxoedema**

Thickened, waxy swelling of the skin over the shins, most often associated with Graves disease.

### **Prevalence**

The proportion of people in a defined population who have a disease or characteristic at a particular time or during a stated period. It includes both newly diagnosed and previously existing cases.

### **Preventive Dentistry**

Dental care that lowers the risk of tooth decay, gum disease, and tooth loss. It includes brushing with fluoride toothpaste, cleaning between teeth, protective sealants when appropriate, healthy eating advice, and regular dental examinations.

### **Preventive Health Care**

Preventive health care aims to prevent disease or detect it early through vaccination, screening, healthy behaviors, and routine medical care.

### **Preventive Medicine**

The medical field focused on preventing illness and promoting health through individual care, screening, vaccination, and community programs.

### **Prevertebral Air**

Air seen in the soft tissues in front of the spine on a neck X-ray. It can indicate injury or a hole in the swallowing or breathing passages, a deep infection, or recent surgery and may require urgent evaluation.

### **Priapism**

Prolonged, painful erection lasting longer than 4 hours, occurring independently of sexual desire or stimulation. Medical emergency categorized into two types: ischemic (low-flow, veno-occlusive) and non-ischemic (high-flow, arterial).

#### **Alternate Names:**

Persistent Penile Erection; Low-Flow or High-Flow

Priapism

### **Prickly Heat**

Miliaria, an itchy rash caused by blockage of sweat ducts.

### **Prick Test**

A skin test in which small amounts of suspected allergens are introduced into the skin.

#### **Alternate Names:**

Skin-Prick Test; Percutaneous Allergy Test

### **Primary**

First in time, origin, importance, or site; in disease classification it often means arising independently rather than secondary to another cause.

### **Primary Adrenal Insufficiency**

Alternate name; see *Addison's disease*.

### **Primary Adrenal Lymphoma**

A rare lymphoma arising primarily in the adrenal gland, most commonly diffuse large B-cell lymphoma. It often presents bilaterally and may cause adrenal insufficiency.

### **Primary Aldosteronism**

Excess production of aldosterone by the adrenal glands, usually from an adrenal nodule or enlargement of both glands. It increases sodium retention and may cause hypertension, low potassium, or suppressed renin.

### **Primary Alveolar Hypoventilation**

A rare disorder of inadequate automatic breathing, often related to PHOX2B or other autonomic-control abnormalities, causing elevated carbon dioxide especially during sleep.

### **Primary Amebic Meningoencephalitis**

A fulminant, rapidly fatal central nervous system infection caused by the free-living thermophilic amoeba *Naegleria fowleri*, contracted when warm contaminated freshwater is inhaled into the nasal cavity during swimming, diving, or nasal irrigation.

---

The amoebae penetrate the olfactory neuroepithelium and cribriform plate to reach the brain, causing acute hemorrhagic necrotizing meningoencephalitis characterized by severe frontal headache, high fever, meningismus, altered olfaction, seizures, and rapid coma, carrying a mortality rate exceeding 97%.

**Alternate Names:**

PAM; Naegleriasis; Primary Amoebic Meningoencephalitis; Fatal Amebic Meningoencephalitis

**Primary Amenorrhea**

Failure to establish menarche by age 15 in the presence of normal secondary sexual development, or by age 13 in the absence of secondary sexual characteristics. Causes include gonadal dysgenesis, müllerian agenesis, hypothalamic-pituitary disorders, or anatomical outflow tract obstruction.

**Alternate Names:**

Primary Amenorrhea

**Primary Amyloidosis**

Primary amyloidosis, usually AL amyloidosis, results from abnormal plasma-cell proteins depositing as amyloid in organs such as the heart, kidneys, nerves, or gastrointestinal tract.

**Primary and Secondary Fibrinolysis**

The physiologic enzymatic breakdown of fibrin blood clots by plasmin. Primary fibrinolysis occurs without preceding clotting, whereas secondary fibrinolysis is a reactive response to widespread clot formation, as seen in disseminated intravascular coagulation (DIC).

**Alternate Names:**

Hyperfibrinolysis; Clot Dissolution; Fibrinolytic System

**Primary And Secondary Hyperaldosteronism**

Excess aldosterone production, from an adrenal source in primary disease or from increased renin stimulation in secondary disease, causing hypertension, sodium retention, and sometimes low potassium.

**Primary Angioplasty**

Angioplasty used as the initial treatment to reopen an artery during an acute heart attack.

**Primary Biliary Cholangitis**

A chronic autoimmune liver disease in which small intrahepatic bile ducts are progressively damaged. It can cause cholestasis, itching, fatigue, and eventually cirrhosis; antimitochondrial antibodies often support the diagnosis.

**Primary Biliary Cirrhosis**

Alternate name; see *Primary biliary cholangitis*.

**Primary Biliary Cirrhosis Treatment**

Alternate name; see *PBC*.

**Primary Care**

The usual first place people go for healthcare. It includes prevention, check-ups, treatment of common illnesses, care of long-term conditions, basic mental-health support, and referral or coordination with specialists when needed.

**Primary Ciliary Dyskinesia**

An inherited disorder in which tiny moving hairs lining the airways do not clear mucus properly. It can cause repeated sinus and lung infections, permanent airway widening, hearing problems, infertility, and organs arranged differently from usual.

**Primary Ciliary Dyskinesia (Kartagener Syndrome)**

An inherited disorder of motile cilia that impairs clearance of airway secretions. It causes chronic wet cough, recurrent sinus and ear disease, bronchiectasis, infertility in some patients, and sometimes situs inversus; long-term airway care and infection prevention are important.

**Primary Vulvar Clear Cell Adenocarcinoma**

A rare cancer of the vulva made up of clear-looking cells. Diagnosis requires examination of a tissue sample and tests that distinguish it from cancer

---

that has spread from the kidney or another organ.

### **Primary CNS Lymphoma**

An aggressive lymphoma confined to the brain, spinal cord, meninges, cerebrospinal fluid, or eyes at diagnosis. It often causes focal neurologic deficits, cognitive change, seizures, or raised intracranial pressure and requires specialist imaging, biopsy, and systemic evaluation.

### **Primary Cutaneous B-Cell Lymphoma**

A type of lymphoma that starts in the skin and has not spread elsewhere when diagnosed. It may cause one or more painless lumps or plaques; behavior and treatment depend on the subtype.

### **Primary Endpoint**

The main outcome chosen in advance to judge whether a treatment or other intervention works in a clinical trial. It may be a symptom, laboratory result, complication, quality-of-life measure, or survival.

### **Primary Hyperaldosteronism**

Excess production of the hormone aldosterone by the adrenal glands, usually from a small gland growth or enlargement of both glands. It can cause difficult-to-control high blood pressure, low potassium, muscle weakness, or abnormal heart rhythms.

### **Primary Hyperoxaluria**

A group of inherited disorders in which the liver makes too much oxalate. This can cause repeated kidney stones, calcium deposits, kidney damage, and eventually failure of the kidneys.

### **Primary Hyperparathyroidism**

An endocrine disorder caused by autonomous, excessive secretion of parathyroid hormone (PTH) from one or more parathyroid glands, leading to hypercalcemia and hypophosphatemia. It is most commonly caused by a single benign parathyroid adenoma ( 85% of cases), four-gland parathyroid hyperplasia ( 15%), or rarely parathyroid carcinoma (<1%), and may occur sporadically or as part of MEN1, MEN2A, or MEN4 syndromes. Many patients are

asymptomatic, detected on routine blood testing, but symptomatic disease classically manifests as “stones, bones, abdominal groans, and psychiatric moans” (nephrolithiasis, osteoporosis or osteitis fibrosa cystica, peptic ulcers and pancreatitis, and depression or lethargy). Diagnosis is established by elevated serum calcium alongside inappropriately elevated intact PTH. Definitive curative treatment is surgical parathyroidectomy guided by preoperative localization imaging (such as Tc-99m sestamibi scanning or 4D-CT).

#### **Alternate Names:**

PHPT; Primary Hyperparathyroid Disease; Parathyroid Hyperactivity

### **Primary Hypersomnia**

Excessive daytime sleepiness that is not explained by too little sleep, another sleep disorder, a medicine, substance use, or another illness. It can cause unintended sleep, long sleep, and difficulty becoming fully alert after waking.

### **Primary Hypertension**

Persistently high blood pressure without one identifiable underlying disease. It results from a combination of genetic, environmental, vascular, and lifestyle factors and can often be controlled with lifestyle changes and medicines.

### **Primary Hypertrophic Osteoarthropathy**

A rare inherited condition causing enlarged, rounded fingertips, thickened facial or scalp skin, and new bone growth along the limbs. It may cause joint pain, stiffness, swelling, and excessive sweating.

### **Primary Hypoparathyroidism**

Inadequate parathyroid hormone production causing low calcium and high phosphate, with possible tingling, cramps, tetany, seizures, cataracts, or chronic bone and dental effects.

### **Primary Hypothyroidism**

Reduced thyroid-gland production of thyroxine and triiodothyronine, usually causing high TSH and symptoms such as fatigue, cold intolerance, constipation, dry skin, and slowed heart rate.



---

### Primary Immune Response

The body's first targeted immune response to a new germ or other foreign substance. It develops over days, produces antibodies and immune cells, and creates memory that allows a faster response next time.

### Primary Immunodeficiency

A group of usually inherited disorders in which part of the immune system is missing or does not work properly. They can cause repeated or unusually severe infections and may also cause autoimmune disease or inflammation.

### Primary Immunodeficiency Disease

An inherited or genetic disorder of immune development or function that causes unusual, recurrent, severe, or persistent infections and sometimes autoimmunity.

### Primary Immunodeficiency Disease PIDD

Primary immunodeficiency diseases are inherited disorders of immune development or function that cause unusual, severe, recurrent, or persistent infections and sometimes autoimmunity or inflammation. Evaluation may include blood counts, immunoglobulins, functional tests, and genetic testing.

#### Alternate Names:

PID; Inborn Errors of Immunity; IEI

### Primary Lactase Deficiency

Reduced lactase enzyme activity developing with age, causing lactose malabsorption and possible bloating, abdominal pain, gas, or diarrhea after dairy consumption.

### Primary Lateral Sclerosis

A rare progressive nerve disorder causing increasing muscle stiffness and weakness, usually beginning in the legs. Speech, swallowing, walking, and emotional control may become affected over time.

### Primary Lymphoid Organs

The bone marrow and thymus, where lymphocytes are made and mature.

### Primary Lymphoma Of The Brain

Primary central-nervous-system lymphoma is a lymphoma arising in the brain, spinal cord, eyes, or surrounding fluid without systemic disease at diagnosis.

### Primary Metabolism

The essential chemical activity inside cells that produces energy, builds needed substances, breaks down waste, and keeps cells alive and functioning.

### Primary Microrna

The first RNA copy made from a microRNA gene. It is processed inside and outside the cell nucleus into a shorter molecule that helps control which proteins cells make.

### Primary Myelofibrosis

A long-term bone-marrow disorder in which scar tissue replaces healthy marrow. Blood-cell production may move to the spleen or liver, causing anemia, tiredness, easy bleeding or infection, bone pain, and an enlarged spleen.

#### Alternate Names:

Agnogenic Myeloid Metaplasia

### Primary Open-Angle Glaucoma

A slowly progressive disease that damages the optic nerve, usually without pain or early symptoms. Vision is gradually lost from the sides inward; lowering pressure inside the eye can slow or prevent further damage.

### Primary Osteoporosis

Weakening and thinning of bone caused mainly by aging, loss of estrogen after menopause, or age-related changes, rather than by another disease or medicine. It increases the risk of fractures, especially in the hip, spine, and wrist.

### Primary Ovarian Insufficiency

Loss of normal ovarian activity before age 40, characterized by irregular or absent menses, reduced estrogen, and intermittently or persistently elevated gonadotropins. Causes include genetic, autoimmune,



---

iatrogenic, toxic, infectious, and unknown factors.

### Primary Polycythemia

Alternate name; see *Polycythemia Vera*.

### Primary Prevention

Actions taken before a disease begins to prevent it, such as vaccination, avoiding tobacco, reducing harmful exposures, using sun protection, and controlling health risks. It differs from screening, which looks for disease that may already be present.

### Primary Progressive Aphasia

A neurodegenerative disorder causing gradual worsening of language abilities, including speaking, naming, comprehension, reading, or writing.

### Primary Progressive Multiple Sclerosis

A form of multiple sclerosis that gradually worsens from the beginning, without clear attacks and recoveries. It commonly affects walking, strength, balance, bladder control, and thinking; medicines and rehabilitation may slow disability.

**Alternate Names:**

PPMS; Primary Progressive MS

### Primary Sclerosing Cholangitis

A long-term disease in which inflammation and scarring narrow the bile tubes inside and outside the liver. It can cause itching, tiredness, jaundice, repeated infections, and liver damage and is often associated with inflammatory bowel disease.

### Primary Systemic Amyloidosis

A form of AL amyloidosis in which abnormal immunoglobulin light chains made by a plasma-cell clone deposit in organs such as the heart, kidneys, nerves, liver, or gastrointestinal tract. Symptoms depend on the organs involved and require hematology evaluation.

### Primary Tuberculosis

The first infection with the bacterium that causes tuberculosis. It often begins in the lungs and nearby lymph nodes; the immune system may contain it

without symptoms, or it may progress to active disease.

### Primary Tumor

The original tumor, or site where a cancer began, from which cancer cells may spread to form secondary or metastatic tumors. A cancer may be described as having an unknown primary when the original site cannot be identified.

### Primidone

Primidone is an antiseizure medicine used for epilepsy and sometimes essential tremor; it can cause sleepiness, dizziness, and drug interactions.

### Primigravida

A person who is pregnant for the first time. First pregnancies may have a longer labor and require monitoring for conditions such as high blood pressure, but the term itself does not indicate a problem.

### Primipara

A person who has given birth once after the fetus could survive outside the womb, regardless of outcome.

### Primitive

Relating to an early developmental stage or a basic, less specialized form. In medicine, primitive cells may be immature and capable of developing into several more specialized cell types.

### Primitive Node

The primitive node is an embryonic organizer at the cranial end of the primitive streak that helps establish body axes and germ-layer formation.

### Primitive Streak

The primitive streak is an embryonic structure through which cells migrate during gastrulation to form the three germ layers and establish body organization.

### Primordial

Present at the earliest stage of development or ex-

---

isting in an original, basic form. Examples include primordial germ cells and primordial follicles.

### Primordial Prevention

Preventing the emergence of disease risk factors by addressing social, economic, behavioural, and environmental conditions early. Examples: promoting healthy diets and physical activity in children, reducing tobacco exposure, and creating safe environments for exercise.

### PR Interval

The ECG interval from the start of atrial depolarization to the start of ventricular depolarization, reflecting atrioventricular conduction.

### Prinzmetal Angina

An atypical form of angina pectoris (vasospastic angina) caused by transient, focal epicardial coronary artery vasospasm rather than fixed atherosclerotic plaque obstruction, presenting with episodic ischemic chest pain occurring at rest (characteristically between midnight and early morning) accompanied by transient ST-segment elevation on ECG.

#### Alternate Names:

Vasospastic Angina; Variant Angina

### Prion

An infectious protein folded into an abnormal shape that can make other proteins fold incorrectly. Prions cause rare, serious brain diseases that worsen over time; they are not ordinary bacteria or viruses and currently have no cure.

### Prion Disease

A group of rare, rapidly progressive, and fatal brain disorders caused by prions—infectious agents made entirely of misfolded protein that contain no genetic material (no DNA or RNA). Normally, healthy prion proteins exist in brain cells, but when a protein misfolds into an abnormal shape, it triggers a chain reaction that forces neighboring healthy proteins to misfold into the same defective shape. These clumped proteins destroy brain tissue and leave microscopic holes, giving the brain a sponge-like appearance

(transmissible spongiform encephalopathies). Prion diseases develop in three ways: sporadic (occurring spontaneously in late adulthood without a known trigger, accounting for approximately 85% of cases such as sporadic Creutzfeldt–Jakob disease), genetic or inherited (caused by mutations in the *PRNP* gene passed through families, such as familial Creutzfeldt–Jakob disease, fatal familial insomnia, and Gerstmann–Sträussler–Scheinker disease), or acquired (contracted through exposure to infected tissue, such as eating beef from cattle with bovine spongiform encephalopathy or “mad cow disease”, contaminated surgical instruments, or ritual cannibalism as in kuru). Although the incubation period can last years or decades without symptoms, once illness begins it causes rapid memory loss, progressive dementia, severe unsteadiness and balance loss (ataxia), sudden muscle jerks (myoclonus), and loss of speech and movement. Prions are extraordinarily resistant to standard heat and chemical sterilization. There is currently no cure, vaccine, or disease-halting treatment, and the conditions are universally fatal, with medical care centered on comfort and palliative support.

#### Alternate Names:

Prion Diseases; Transmissible Spongiform Encephalopathies; TSE; Prion Disorders

### Prism

A clear optical device that bends light. In eye care, prisms can measure or correct eye misalignment, reduce double vision, and be built into glasses or used during examinations.

### Privacy

Privacy is a person’s right and expectation that personal information, communications, and bodily matters are protected from unauthorized access or disclosure.

### Privacy Breach

A privacy breach is unauthorized access, use, disclosure, loss, or exposure of personal or health information.

---

### Private Mutation

A genetic variant confined to a family, individual, tumor, or small population rather than commonly seen in the general population; its significance requires genetic and clinical interpretation.

### PRK

Photorefractive keratectomy is laser eye surgery that reshapes the cornea to reduce short-sightedness, long-sightedness, or astigmatism. The surface layer is removed and grows back during healing; pain, haze, infection, or imperfect vision can occur.

### Probe

A blunt instrument used to examine an opening, cavity, wound, tooth, or narrow passage. Different probes can measure depth, check whether a passage is open, guide a procedure, or trace a sinus or fistula.

### Probenecid

A uricosuric medicine used to prevent gout attacks in selected patients and to alter renal excretion of some drugs.

### Probiotic

A live microorganism that may provide a health benefit when taken in sufficient amounts. Benefits depend on the specific strain and condition; a probiotic is not automatically effective or safe for every person.

### Probiotics

Probiotics are live microorganisms that may provide a health benefit when consumed in adequate amounts, although effects are strain-specific and vary by condition.

### Probucol

Probucol is a lipid-lowering drug that reduces cholesterol but may prolong the QT interval and is not widely used in current practice.

### PROC

PROC is the gene symbol for protein C, an anticoagulant protein that limits clot formation; inherited

deficiency increases thrombosis risk.

### Procainamide

An antiarrhythmic medicine that can slow abnormal electrical conduction in the heart. It is used in selected rhythm disorders and requires monitoring because it can affect blood pressure, the heart rhythm, and immune or blood systems.

### Procaine

An ester local anesthetic that temporarily blocks nerve signals and reduces pain. It is now used mainly in limited or historical settings because newer anesthetics are commonly preferred and allergic reactions can occur.

### Procalcitonin

A biomarker that may rise during some bacterial infections and systemic inflammation. It can support assessment of infection and antibiotic decisions, but it cannot by itself diagnose or exclude bacterial infection or determine sepsis severity.

### Procarbazine

A chemotherapy medicine used in some lymphomas and brain tumors. It damages the DNA of rapidly growing cells and can cause nausea, low blood counts, infertility, or other side effects; alcohol and many medicines may interact with it.

### Procarbazine Hydrochloride

A salt form of procarbazine, a chemotherapy medicine that damages cancer cells. It is used for selected lymphomas and brain tumors and can cause low blood counts, nausea, infertility, or serious medicine and food interactions.

### Procedural Memory

Memory for how to perform learned skills, such as riding a bicycle, typing, or playing an instrument. It usually works without consciously recalling each step and differs from memory for facts or events.

### Procedural Sedation

Use of medicines to reduce anxiety, discomfort, and

---

awareness during a medical procedure while maintaining breathing and protective reflexes. Continuous monitoring is required because sedation can unexpectedly become deeper and slow breathing.

### **Procedure**

A planned medical or surgical action performed to diagnose, treat, prevent, monitor, or relieve a health problem. It may involve an examination, sample collection, medicine delivery, device placement, or operation, with risks depending on the method and person.

#### **Alternate Names:**

Medical Procedure; Clinical Procedure; Intervention

### **Procentriole**

An immature structure that forms next to an existing centriole as a cell prepares to divide. It later matures into a centriole, which helps organize the fibers that separate chromosomes.

### **Procerus Muscle**

A small facial muscle over the bridge of the nose that draws the medial eyebrows downward and produces horizontal wrinkles between the eyebrows.

### **Process**

A process is an ordered series of biological, physical, chemical, or operational changes that produces a result.

### **Processing Bodies**

Processing bodies are cytoplasmic RNA-protein granules where messenger RNAs may be stored, decapped, degraded, or regulated.

### **Prochlorperazine**

A phenothiazine medicine used mainly for severe nausea or vomiting and, in selected cases, psychotic disorders. It can cause drowsiness, low blood pressure, and movement disorders.

### **Prochlorperazine Overdose**

Taking too much prochlorperazine can cause severe sleepiness, confusion, low blood pressure, abnormal

heart rhythms, seizures, or muscle stiffness with fever. It requires immediate emergency or poison-control advice.

### **Procidentia**

The most severe form of uterine prolapse, in which the uterus and cervix descend through and outside the vaginal opening. It may cause pressure, bleeding, sores, difficulty urinating, or bowel problems and requires gynecologic assessment.

#### **Alternate Names:**

Complete Uterovaginal Prolapse; Third-Degree Uterine Prolapse

### **Procollagen**

An unfinished form of collagen made inside cells. After chemical changes and removal of its end sections outside the cell, it joins with other molecules to form strong collagen fibers in skin, bone, tendons, and other tissues.

### **Proctalgia**

Pain in or around the rectum or anal canal. Causes include fissures, hemorrhoids, abscesses, inflammation, injury, and functional muscle pain; sudden severe pain or pain with fever, bleeding, or a lump needs medical assessment.

### **Proctalgia Fugax**

A benign functional anorectal disorder characterized by sudden, severe, fleeting, cramp-like spasms of intense pain localized to the rectum and anus lasting from seconds to several minutes (maximum 30 minutes) and resolving completely, occurring predominantly at night due to levator ani and anal sphincter spasm.

#### **Alternate Names:**

Fugacious Rectal Pain; Nocturnal Rectal Spasm

### **Proctitis**

Inflammation of the rectum. It can cause a constant urge to pass stool, rectal pain, bleeding, mucus or pus, and pelvic discomfort. Causes include inflammatory bowel disease, sexually transmitted infections, radiation treatment, and diversion of stool through an opening in the abdomen. Diagnosis may

---

require an examination, stool or swab tests, and a small tissue sample. Treatment depends on the cause.

**Alternate Names:**

Rectal Inflammation

**Proctocolitis**

Inflammation of the rectum and nearby lower colon. It may cause bleeding, mucus, diarrhea, urgency, pain, or difficulty passing stool and can result from infection, inflammatory bowel disease, reduced blood flow, radiation, or other causes.

**Proctology**

The part of medicine concerned with diseases of the anus and rectum. Care may include assessment and treatment of hemorrhoids, fissures, fistulas, abscesses, bowel-control problems, infections, inflammation, and cancers.

**Proctoscope**

An instrument used to inspect the anal canal and rectum.

**Proctoscopy**

Examination of the anal canal and rectum with a short viewing instrument. It can help investigate bleeding, pain, inflammation, ulcers, polyps, or a lump and may allow a small tissue sample to be taken.

**Proctostomy**

An operation that creates an opening from the rectum through the abdominal wall or nearby skin so stool can leave the body. It may be temporary or permanent and requires care of the opening and surrounding skin.

**Prodromal**

Describing early, nonspecific signs or symptoms that occur before the characteristic clinical features of a disease become apparent.

**Prodrome**

An early symptom or group of symptoms that ap-

pears before a disease episode or major event. Examples include tiredness before migraine, tingling before a seizure, or fever before an infection becomes obvious.

**Prodrug**

A medicine given in a form that the body changes into its active form. This design may improve absorption, delivery to a target tissue, duration of action, or tolerability.

**Productive Cough**

A cough that brings up mucus or phlegm from the airways. It can occur with infections, asthma, chronic lung disease, or other conditions and does not by itself identify the cause.

**Profunda**

A deep artery or vessel, commonly referring to the profunda femoris artery in the thigh. This artery supplies muscles and nearby tissues and can be important in bleeding, circulation, or vascular procedures.

**Progenitor**

A partly developed cell that can multiply and mature into one or a limited range of specialized cell types. Progenitor cells help replace or repair tissues but have less developmental potential than stem cells.

**Progeny**

Biological offspring or descendants. In genetics, the term usually means children or later descendants whose traits may be studied to understand how characteristics or disorders are inherited. In cell biology, it can mean daughter cells produced by one parent cell.

**Progeria**

A rare genetic disorder in which features resembling accelerated aging begin during childhood. The usual form is Hutchinson–Gilford progeria syndrome, caused by a disease-causing variant in the *LMNA* gene that produces abnormal lamin A. Most cases result from a new variant rather than inheri-

---

tance from an affected parent. Children are often healthy-looking at birth and later develop slowed growth, loss of body fat, hair loss, thin aged-looking skin, joint stiffness, and characteristic facial features. Accelerated atherosclerosis can cause early cardiovascular disease, while intellectual development is usually preserved.

**Alternate Names:**

Hutchinson–Gilford Progeria Syndrome Hutchinson–Gilford Syndrome HGPS

**Progestational**

Relating to progesterone-like effects or preparation of the uterus for pregnancy. Progestational medicines can alter menstrual bleeding, prevent ovulation, support the uterine lining, or protect it from excessive estrogen stimulation.

**Progesterone**

A steroid hormone produced mainly by the corpus luteum and placenta that supports the endometrium and pregnancy.

**Progesterone Antagonist**

A progesterone antagonist blocks progesterone receptors and may be used for selected reproductive, gynecologic, or other medical purposes.

**Progesterone Receptor**

A protein inside or on a cell that binds progesterone. Some breast cancers have these receptors, which helps describe the cancer's biology.

**Progesterone Supplementation**

Use of progesterone during pregnancy in selected people at risk of preterm birth, especially when ultrasound shows a short cervix. Vaginal progesterone may be considered in that situation; routine use after a previous preterm birth without a short cervix is not recommended, and 17-hydroxyprogesterone caproate is no longer FDA-approved for this purpose.

**Progesterone Test**

A blood test that measures the level of progesterone, a hormone involved in ovulation and pregnancy.

It may help assess ovulation, infertility, abnormal uterine bleeding, fertility treatment, or certain pregnancy problems. Results depend on the timing of the menstrual cycle and whether the person is pregnant.

**Alternate Names:**

Serum Progesterone Progesterone Blood Test

**Progestin**

A progestin is a synthetic progesterone-like hormone used in contraception, hormone therapy, and treatment of selected gynecologic conditions.

**Progestogen**

A natural or synthetic hormone with effects like progesterone. Progestogens help regulate the menstrual cycle and pregnancy and are used in some contraceptives and hormone treatments.

**Progestogen-Only Pill**

Oral contraceptive tablets formulated solely with a progestin without an estrogen component. They prevent conception primarily by thickening cervical mucus to impede sperm penetration, altering endometrial receptivity, and in certain formulations inhibiting the mid-cycle luteinizing hormone surge.

**Alternate Names:**

Progestin-Only Pill; POP; Mini-Pill

**Proglottid**

One segment of a tapeworm. Each mature segment contains reproductive organs and may release eggs in stool.

**Prognathism**

Forward projection of the upper or lower jaw relative to the facial skeleton, affecting bite, appearance, chewing, speech, and sometimes airway function.

**Prognosis**

The expected course and outcome of a disease or condition.

**Prognostic Factor**

A prognostic factor is a clinical, pathologic, or bio-



---

logic feature associated with the likely course, outcome, or survival of a disease.

### **Prognostic Test**

A test that provides information about a disease's likely course or outcome. Results may estimate risks, guide treatment choices, or monitor response, but they cannot predict an individual future with certainty.

### **Programmed Cell Death**

A controlled process by which cells die as part of normal development, tissue maintenance, or disease. Apoptosis is the best-known form; other regulated forms include necroptosis and pyroptosis.

### **Progression**

Worsening or advancement of a disease, such as increasing tumor size, new lesions, or spread to another site.

### **Progression-Free Survival**

The length of time after a defined starting point, such as diagnosis or treatment, during which a person remains alive without the cancer worsening. It is a group measure used in research and cannot predict exactly what will happen to one person.

### **Progressive**

Increasing, worsening, or spreading over time, or moving forward in a particular direction. A progressive disease may cause gradually increasing symptoms, but the word alone does not explain the cause, speed, or expected outcome.

### **Progressive Disease**

A disease that is worsening, spreading, or becoming more active over time. In cancer, progression may mean that tumors have grown, new tumors have appeared, or existing disease has clearly worsened on examination or imaging.

### **Progressive Lenticular Degeneration**

An older name for Wilson disease, an inherited disorder causing copper buildup. Excess copper can

damage the liver, brain, eyes, kidneys, and other organs unless treatment lowers copper levels.

### **Progressive Multifocal Leukoencephalopathy**

A serious brain infection caused by reactivation of JC virus in people with severely weakened immune systems. It damages the brain's nerve insulation and may cause worsening weakness, clumsiness, vision or speech problems, confusion, or seizures.

### **Progressive Muscular Atrophy**

A motor-neuron disorder predominantly affecting lower motor neurons and causing progressive muscle wasting and weakness.

### **Progressive Supranuclear Palsy**

A progressive brain disorder that causes balance problems, frequent falls, stiffness, slowed movement, speech or swallowing difficulty, and trouble moving the eyes, especially looking down. It resembles Parkinson disease but has different features and treatment needs.

#### **Alternate Names:**

Steele-Richardson-Olszewski Syndrome Supranuclear Palsy

### **Proinflammatory**

Describing a substance, cell, or immune response that starts or increases inflammation. Such activity can help fight infection or repair injury, but excessive or prolonged inflammation can damage healthy tissue.

### **Proinsulin**

An unfinished hormone made by pancreatic beta cells before it is split into insulin and C-peptide. Measuring proinsulin can provide information about insulin production and certain disorders of glucose regulation.

### **Projectile Vomiting**

Forceful vomiting expelled with unusual intensity, which may occur with gastric outlet obstruction, raised intracranial pressure, infection, migraine, or other causes; severity and associated signs guide

---

evaluation.

### **Projection**

A view or image made by directing X-rays or other imaging energy through the body toward a detector. Different projections show body structures from different angles and can help locate an abnormality.

### **Prokaryote**

A prokaryote is a cell or organism without a membrane-bound nucleus, including bacteria and archaea.

### **Prokinetic**

A medicine that promotes gastrointestinal motility and movement of contents through the digestive tract; use depends on the underlying disorder and safety profile.

### **Prolactin**

A hormone released mainly by the pituitary gland that supports breast development and milk production. Abnormal levels can affect periods, fertility, sexual function, pregnancy, breastfeeding, and sometimes vision because of pituitary growths.

### **Prolactin Blood Test**

A prolactin blood test measures the hormone prolactin and helps evaluate abnormal milk production, menstrual or fertility problems, and some pituitary disorders.

### **Prolactin Deficiency**

Inadequate production of prolactin by the pituitary, most clearly recognized by failure to produce breast milk after delivery. It may occur with broader pituitary disease, surgery, hemorrhage, or injury and is evaluated with other pituitary hormones.

### **Prolactinoma**

A usually benign pituitary tumor that produces too much prolactin. It may cause milk production, menstrual or fertility problems, low sex hormones, headache, or vision loss and is often treated with dopamine-agonist medicines.

### **Alternate Names:**

Prolactin-Secreting Adenoma; Lactotroph Adenoma

### **Prolactinoma Treatment**

Treatment may include a dopamine agonist, surgery, radiation, or observation, depending on tumor size, symptoms, hormone levels, treatment response, and patient preferences. Dopamine agonists often lower prolactin levels and reduce tumor size.

### **Prolactin Physiology**

Prolactin is a hormone made by the pituitary gland that supports breast development and milk production. Levels normally rise during pregnancy, breastfeeding, sleep, stress, and breast stimulation.

### **Prolactin Receptor**

A cell-surface receptor activated by prolactin to regulate mammary development, milk production, reproduction, immune effects, and other tissue responses.

### **Prolapse**

Downward displacement or protrusion of an organ or tissue from its normal position, often through a supporting structure or body opening.

### **Prolapsed Bladder**

Alternate name; see *Anterior Vaginal Prolapse*.

### **Prolapsed Cord**

A prolapsed umbilical cord descends beside or ahead of the fetus after membrane rupture, risking cord compression and reduced fetal oxygen; it is an obstetric emergency.

### **Prolapsed Disk**

A spinal disk that bulges or ruptures and may compress a nearby nerve; also called a herniated disk.

### **Alternate Names:**

Herniated Nucleus Pulposus; Slipped Disc; Herniated Disc

### **Prolapsed Uterus**

Alternate name; see *Uterine prolapse*.

---

### **Prolapse Of Orbital Fat**

Bulging of the eye's normal cushioning fat through a weak spot in the thin tissue around the eye. It usually appears as a soft, painless lump in an upper or lower eyelid, often with aging, but other lumps may look similar.

### **Proliferating Pilar Tumor**

A usually low-grade follicular tumor arising from a trichilemmal cyst, most often on the scalp. It may be large, ulcerated, recurrent, or locally aggressive, so complete excision and histologic assessment are important.

### **Proliferating Trichilemmal Tumor**

A usually low-grade follicular tumor arising from a trichilemmal cyst, most often on the scalp; large or recurrent lesions may show locally aggressive or malignant behavior.

### **Proliferation**

Increase in the number of cells or organisms through growth and division. It may be normal, part of healing, or abnormal as in uncontrolled tumor growth.

### **Proliferative**

Characterized by increased cell multiplication or new tissue growth. The meaning depends on context: some proliferative changes are normal or healing, while others are precancerous or cancerous.

### **Proliferative Diabetic Retinopathy**

An advanced form of diabetic eye disease in which poor blood supply causes fragile new blood vessels to grow on the retina or nearby structures. They can bleed, scar, detach the retina, and threaten vision; specialist treatment can reduce these risks.

### **Proliferative Fasciitis**

A benign, reactive fibroblastic proliferation resembling nodular fasciitis but containing distinctive ganglion-like cells; it usually grows quickly and resolves or stabilizes after local treatment.

### **Proliferative Myositis**

A benign reactive proliferation within skeletal muscle that may present as a rapidly enlarging mass and mimic sarcoma. Imaging and biopsy are used to establish the diagnosis.

### **Proliferative Retinopathy**

Advanced retinal disease in which abnormal new blood vessels grow and may bleed or impair vision.

### **Proline**

An amino acid used to build proteins. Its ring-shaped side chain helps give collagen and other proteins their structure.

### **Prolonged Capillary Refill Time**

A delay of greater than two to three seconds for normal skin color to return after moderate digital blanching pressure is applied to a distal nail bed or pulp and released. It is a bedside indicator of reduced peripheral microvascular perfusion, shock, dehydration, or hypothermia.

### **Prolonged Deceleration**

A temporary slowing of the fetal heart rate lasting at least two minutes but less than ten minutes. It may result from reduced blood flow, excessive contractions, cord problems, or placental separation and requires prompt assessment during labor.

### **Prolonged Second Stage**

The pushing stage of labor lasting longer than expected. The safe duration depends on whether it is a first birth, use of epidural anesthesia, progress, and the health of the parent and fetus; the team reassesses before deciding on further care.

### **Polymphocyte Count**

The number or proportion of polymphocytes, immature-appearing lymphoid cells, in blood or marrow; an increased count may occur in selected lymphoid leukemias and requires morphology and immunophenotyping.

---

### **Prolymphocytic Leukemia**

Prolymphocytic leukemia is a rare aggressive leukemia with large abnormal lymphocytes in the blood, bone marrow, spleen, or lymph nodes.

### **Promazine**

A first-generation phenothiazine antipsychotic with sedative properties; it is now used infrequently in many settings and availability varies by country. It can cause drowsiness, orthostatic hypotension, anticholinergic effects, extrapyramidal symptoms, and other serious antipsychotic adverse effects.

### **Promecarb**

Promecarb is a carbamate insecticide that inhibits acetylcholinesterase and can cause cholinergic poisoning after substantial exposure.

### **Prometaphase**

Prometaphase is the stage of mitosis after nuclear-envelope breakdown when spindle microtubules attach to chromosome kinetochores.

### **Promethazine**

Promethazine is an antihistamine used for allergy symptoms, nausea, motion sickness, and sometimes sedation; it can cause marked sleepiness and breathing problems in young children.

### **Promethazine Overdose**

Too much promethazine can cause extreme drowsiness, confusion, agitation, fast heartbeat, seizures, breathing problems, or coma. Children are particularly vulnerable; significant or intentional overdoses need immediate emergency advice.

### **Prominent Iris Collarette**

A visible thickening or ridge of the iris at the junction of the pupillary and ciliary zones, representing the site where the embryonic pupillary membrane was attached. It is typically a normal anatomical variant but can be a clinical sign in conditions such as Axenfeld-Rieger syndrome or iridocorneal endothelial syndrome.

### **Promonocyte Count**

The number or proportion of promonocytes, immature monocyte precursors, in blood or marrow; increased cells may support evaluation of acute or chronic myeloid leukemia.

### **Promontory**

A part of the body that projects outward like a small ridge or bump. Examples include the sacral promontory in the pelvis and the cochlear promontory in the inner ear; the exact meaning depends on the structure named.

### **Promoter**

A region of DNA near a gene where proteins attach to start copying the gene into RNA. Promoters help determine when a gene is active and how much of its protein is made.

### **Promoters**

DNA control regions that help start gene activity when regulatory proteins attach. They influence how much messenger RNA and protein a gene makes and are important in development, normal function, and disease.

### **Promyelocytes**

Immature cells in the bone-marrow pathway that normally develops into several types of white blood cells. Large numbers in blood or marrow may occur in leukemia and need examination with other blood and cell tests.

### **Pronation**

Turning the forearm so the palm faces down when the elbow is bent, or backward when the arm hangs at the side. During this movement, the radius rotates across the ulna.

### **Pronator**

A muscle that turns a limb or part of it into a pronated position. In the forearm, pronator muscles rotate the palm downward and help stabilize the elbow during movement.

---

### **Pronator Drift**

A subtle test for nerve or brain weakness where holding the arms straight out with palms up and eyes closed causes one hand to turn inward and slowly drift down.

### **Prone**

Prone means lying face downward with the front of the body against a surface; positioning affects breathing, pressure injury, and procedures.

### **Pronephric**

Relating to the earliest temporary kidney system formed in an embryo. In humans it does not provide effective kidney function and is replaced during development by later kidney structures.

### **Pronephros**

The first temporary kidney structure formed in a human embryo. It is not functional as a kidney and largely disappears as the permanent kidneys develop.

### **Prone Position**

A position in which the patient lies face down, commonly providing access to the back, spine, buttocks, or posterior legs.

### **Prone Positioning**

Placing a person face down to improve breathing, especially during severe acute respiratory distress syndrome. It can improve oxygen levels by helping more of the lung take part in breathing, but requires careful monitoring of the airway, skin, eyes, and pressure points.

### **Propafenone**

A heart-rhythm medicine that slows electrical signals through the heart. It is used for selected irregular rhythms, including episodes of atrial fibrillation, but can worsen or cause dangerous rhythms and is unsuitable for some people with structural heart disease.

### **Propane**

Propane is a flammable hydrocarbon gas used as fuel; inhalation or displacement of oxygen can cause suffocation, and liquid contact can cause cold burns.

### **Propane Poisoning**

Breathing concentrated propane can displace oxygen and cause dizziness, headache, confusion, unconsciousness, abnormal heart rhythms, or suffocation. Leave the area if safe, avoid flames, and call emergency services for symptoms or significant exposure.

### **Propanol**

Propanol is an alcohol solvent with two structural isomers; ingestion or substantial exposure can depress the nervous system and irritate tissues.

### **Propanolol**

A nonselective beta-adrenergic blocker used for hypertension, arrhythmias, angina, tremor, migraine prevention, and other indications.

### **Propantheline**

An older medicine that reduces certain nerve signals to relax the gut and decrease sweating. It can cause dry mouth, blurred vision, constipation, difficulty urinating, fast heartbeat, or confusion and is rarely used today.

### **Propellant**

A propellant is a gas or volatile substance that expels a product from a container or delivers medicine in an inhaler.

### **Properdin**

A blood protein that strengthens one part of the body's early immune defense. It helps mark and damage certain germs; too little properdin can increase susceptibility to serious bacterial infections.

### **Properdin Deficiency**

A rare inherited immune-system disorder in which a complement protein is missing or low. It increases the risk of severe infections, especially meningococcal disease, and is evaluated with immune tests and

---

a history of recurrent infections.

### **Propetamphos**

Propetamphos is an organophosphate insecticide that can inhibit acetylcholinesterase and cause cholinergic toxicity after exposure.

### **Prophage**

A prophage is bacteriophage genetic material integrated into a bacterial chromosome or maintained in a dormant state inside a bacterium.

### **Prophase**

Prophase is the early stage of cell division when chromosomes condense, the spindle begins forming, and the nucleolus disappears.

### **Prophenofos**

Prophenofos is an organophosphate pesticide whose toxicity results mainly from acetylcholinesterase inhibition and excess cholinergic activity.

### **Prophylactic**

Prophylactic means intended to prevent disease, infection, injury, or another unwanted outcome before it occurs.

### **Prophylactic Mastectomy**

Prophylactic mastectomy removes one or both breasts to greatly reduce cancer risk in selected people at very high genetic or clinical risk.

### **Prophylaxis**

Medicine or other preventive care used to lower risk of disease, infection, or recurrence before it occurs.

### **Propionates**

Salts or esters of propionic acid used in food preservation, metabolism, and some medicines; clinical effects depend on the specific compound and exposure.

### **Propionibacterium**

A former bacterial genus name now largely replaced by *Cutibacterium*; its members are common skin

organisms and can cause acne or device-related infection.

### **Propionic Acidemia**

A rare inherited disorder in which the body cannot properly break down certain proteins and fats. Harmful acids can build up, especially during illness or fasting, causing vomiting, extreme sleepiness, poor feeding, seizures, or coma. Newborn screening can detect it; treatment uses a controlled diet, avoids fasting, and manages attacks urgently.

#### **Alternate Names:**

Propionic Aciduria (PA); Propionyl-CoA Carboxylase Deficiency

### **Propofol**

A short-acting medicine given through a vein to start or maintain anesthesia and provide procedural sedation. It acts quickly but can lower blood pressure, slow breathing, or cause dangerous reactions without monitoring.

### **Propofol Infusion Syndrome**

A rare, potentially fatal illness after prolonged or high-dose propofol infusion. It may cause severe blood acidity, muscle breakdown, high potassium, kidney failure, heart failure, or dangerous heart rhythms. Stopping propofol and urgent intensive care are required when it is suspected.

### **Propoxur**

Propoxur is a carbamate insecticide that inhibits acetylcholinesterase and may cause sweating, salivation, vomiting, breathing difficulty, or weakness after exposure.

### **Propoxycaine**

A local anesthetic formerly used in some topical or dental preparations; toxicity can affect the nervous system and heart.

### **Propoxyphene**

Propoxyphene is a former opioid analgesic withdrawn in many countries because of overdose toxicity, respiratory depression, and dangerous cardiac arrhythmias.



---

## Propranolol

A beta-blocking medicine that slows heart rate and reduces the body's response to adrenaline. It treats selected heart conditions, tremor, migraine prevention, anxiety symptoms, and other problems, but can worsen asthma or low blood sugar.

## Proprioception

Awareness of body position and movement without looking, based on signals from muscles, tendons, joints, skin, and balance organs. Reduced proprioception can cause clumsiness, poor coordination, falls, or difficulty walking.

## Proprioceptor

A sensory receptor that detects position, movement, or muscle stretch.

## Proptosis

The abnormal forward protrusion or displacement of one or both eyeballs from the orbit. Major causes include thyroid eye disease (Graves ophthalmopathy), orbital cellulitis, orbital tumors, idiopathic orbital inflammation, and vascular malformations such as carotid-cavernous fistulas.

### Alternate Names:

Exophthalmos; Ocular Proptosis; Proptosis of Eye; Bulging Eyes

## Proptosis of Eye

Alternate name; see *Proptosis*.

## Propulsion

Movement that drives the body or another object forward. In some neurologic disorders, impaired forward movement can cause short, shuffling steps or a tendency to lean and fall forward.

## Propyl Alcohol

Propyl alcohols are flammable solvents; ingestion or substantial exposure can irritate tissues and depress the central nervous system, with toxicity depending on the isomer and dose.

## Propylene Glycol

Propylene glycol is a solvent and humectant used in medicines, foods, and cosmetics; large exposures can cause metabolic or kidney problems.

## Propylthiouracil

Propylthiouracil is an antithyroid drug that reduces thyroid-hormone synthesis and conversion of T4 to T3; liver injury and fetal risks require careful use.

## Pro Re Nata

A medication or treatment instruction meaning that it should be used only when a specified symptom or clinical circumstance occurs, within clearly prescribed dose and frequency limits.

### Alternate Names:

PRN; As Needed

## Prosencephalon

The front part of the early developing brain. It later forms the cerebral hemispheres, thalamus, hypothalamus, and related structures that control thought, sensation, movement, hormones, and other functions.

## Prosopagnosia

Difficulty or inability to recognize familiar faces despite adequate vision and general intelligence. It may be developmental or result from brain injury or disease.

## Prosoplasia

Development of a more specialized cell function or structure within a tissue. The change may occur during normal development or as an adaptation, and its medical meaning depends on the tissue and surrounding disease process.

## Prosoposchisis

A rare congenital facial cleft or fissure caused by abnormal fusion of facial processes during embryonic development, potentially affecting the nose, mouth, eyes, and feeding or breathing.

---

## Prostacyclin Analogs

Medications that mimic prostacyclin (PGI<sub>2</sub>), causing vasodilation and inhibiting platelet aggregation in the pulmonary vasculature. They include epoprostenol (continuous IV), treprostinil (subcutaneous, IV, inhaled, oral), and iloprost (inhaled). They are used for severe PAH.

## Prostaglandin

A locally acting lipid mediator derived from arachidonic acid that influences pain, inflammation, fever, vascular tone, platelet function, gastric protection, kidney function, and uterine contraction.

## Prostaglandin Analogs

Eye-drop medicines that lower pressure inside the eye by helping fluid drain. They treat glaucoma or ocular hypertension and may cause eye redness, eyelash growth, or darkening of the iris or skin around the eye.

## Prostaglandin Degradation

Enzymatic inactivation and removal of prostaglandins, which limits the duration and intensity of their local effects on vessels, inflammation, smooth muscle, kidneys, stomach, and reproductive tissues.

## Prostaglandin E<sub>2</sub>

A prostaglandin that mediates inflammation, pain, fever, vasodilation, gastric protection, cervical ripening, and uterine contraction and is used in selected obstetric and cardiovascular treatments.

## Prostaglandin F<sub>2</sub>-Alpha

A prostaglandin that promotes smooth-muscle contraction and luteolysis; analogues are used in glaucoma, postpartum hemorrhage, labor-related care, and veterinary medicine.

## Prostaglandin Receptor

A protein on or inside a cell that recognizes a prostaglandin and starts signals affecting blood-vessel width, inflammation, pain, smooth-muscle activity, platelets, or reproduction.

## Prostaglandins

Locally acting lipid mediators involved in pain, inflammation, vascular tone, gastric protection, and reproductive physiology.

## Prostaglandins E

The E series of prostaglandins, lipid mediators that influence inflammation, vascular tone, gastric protection, kidney blood flow, cervical ripening, and uterine contraction.

## Prostaglandins F

The F series of prostaglandins, lipid mediators involved in smooth-muscle contraction, luteal function, vascular responses, inflammation, and labor-related uterine activity.

## Prostate

A gland below the bladder surrounding part of the urine tube. It adds fluid to semen and may enlarge, become inflamed, or develop cancer, causing urinary, sexual, or pelvic symptoms.

## Prostate Adenocarcinoma

The most common type of prostate cancer, arising from gland cells in the prostate. It often grows slowly but can spread, and its growth is usually stimulated by male sex hormones; grade and stage help determine treatment.

## Prostate Biopsy

Removal of small samples of prostate tissue with a needle to look for cancer under a microscope. The needle may pass through the skin between the scrotum and anus or through the rectum; bleeding, infection, and temporary difficulty urinating are possible risks.

## Prostate Brachytherapy

Radiation treatment in which small radioactive sources are placed inside or next to the prostate to destroy cancer cells. It is used for selected cancers and may cause temporary urinary or bowel symptoms and later erectile problems.

---

## Prostate Cancer

Cancer that forms in the prostate gland. It may grow slowly or behave aggressively and may cause no symptoms early on. Risk increases with age, family history, inherited gene changes, and certain racial or ethnic backgrounds. Prostate-specific antigen testing and sometimes a digital rectal examination may help evaluate risk, but diagnosis requires a prostate biopsy. Tumor grade and stage help guide treatment.

## Prostate Cancer Screening

Prostate cancer screening may use a PSA blood test with or without a digital rectal examination after discussing possible benefits, false positives, and harms.

## Prostate Cancer Staging

Determining how far prostate cancer has grown or spread. Clinicians combine examination, PSA level, tumor grade, biopsy findings, and imaging to decide whether it is confined to the prostate, nearby, or in distant organs.

## Prostate Cancer Treatment

Care for cancer of the prostate, chosen according to cancer grade, stage, growth rate, age, and health. Options may include active surveillance, surgery, radiation, hormone-blocking treatment, or medicines for advanced disease.

## Prostate Carcinoma

Prostate carcinoma is cancer arising from the prostate gland and may grow slowly or behave aggressively.

## Prostatectomy

Surgery to remove part or all of the prostate, commonly for prostate cancer or severe urinary obstruction. Possible effects include urinary leakage, erectile dysfunction, bleeding, and infertility.

## Prostate Disease

Any disorder affecting the prostate, including benign enlargement, inflammation, infection, or cancer. It may cause difficulty urinating, weak flow, urgency,

pelvic pain, blood in urine, or no symptoms initially.

## Prostate Enlargement

Alternate name; see *Benign prostatic hyperplasia*.

## Prostate Gland

The prostate, a gland below the bladder that surrounds the urethra and contributes secretions to semen.

## Prostate Gland Enlargement

Alternate name; see *Benign prostatic hyperplasia*.

## Prostate Infection

Infection or inflammation of the prostate, usually causing pelvic or urinary symptoms and sometimes fever or bloodstream infection.

## Prostate Neoplasm

An abnormal mass of cells arising in the prostate. It may be noncancerous, such as a glandular growth, or cancerous; symptoms and treatment depend on its type, size, and spread.

## Prostate-Specific Antigen

A protein made mainly by the prostate and measured in blood. A high or rising level can occur with prostate cancer, but also with prostate enlargement, infection, urine retention, ejaculation, or recent procedures. The result is considered with age, examination, trends, and other tests rather than used alone.

## Prostate-Specific Antigen Blood Test

A PSA blood test measures prostate-specific antigen; an elevated result may reflect enlargement, inflammation, infection, or cancer and is not diagnostic by itself.

## Prostatic Fluid

Prostatic fluid is a mildly alkaline secretion that forms part of semen and contains substances supporting sperm function and survival.

---

## Prostatic Obstruction

Blockage of urine flow caused by enlargement, inflammation, scarring, stones, or a tumor of the prostate or nearby structures.

## Prostatic Urethral Stent

A small device placed in the prostatic urethra to help maintain urine flow when the prostate obstructs it.

## Prostatism

Urinary symptoms caused by blockage or poor function at the bladder outlet, often related to an enlarged prostate. Symptoms include weak flow, straining, urgency, frequent urination, nighttime urination, or incomplete emptying.

## Prostatitis

Inflammation of the prostate gland. It may be caused by a bacterial infection or by inflammation involving the prostate, pelvic muscles, or nerves without a proven infection. Symptoms may include pain in the pelvis, area between the scrotum and anus, penis, scrotum, lower abdomen, or lower back; painful or frequent urination; difficulty emptying the bladder; painful ejaculation; and, in acute bacterial prostatitis, fever and chills. Recognized types include acute bacterial prostatitis, chronic bacterial prostatitis, chronic prostatitis/chronic pelvic pain syndrome, and asymptomatic inflammatory prostatitis, which causes no symptoms and is found during evaluation for another problem.

## Prostatodynia

Prostatodynia is an older term for pelvic or prostate-region pain without proven bacterial infection, now often considered within chronic pelvic pain syndrome.

## Prosthesis

An artificial device replacing or supporting a missing body part or function. Prostheses may replace limbs, joints, teeth, eyes, or heart valves and require fitting, care, adjustment, and monitoring.

## Prosthetic

Relating to an artificial device that replaces or supports a missing or damaged body part. Examples include artificial limbs, joints, heart valves, teeth, and blood-vessel grafts; complications can include wear, loosening, skin problems, or infection.

## Prosthetic Graft

A prosthetic graft is an artificial tube or patch used to replace or bypass a damaged blood vessel or other tubular structure; infection and thrombosis are important risks.

## Prosthetic Heart Valve

An artificial valve that replaces a diseased heart valve and keeps blood moving in the correct direction. Mechanical valves are durable but usually require lifelong anticoagulation; tissue valves generally avoid lifelong anticoagulation unless another indication exists but can degenerate over time.

## Prosthetic Limb

An artificial limb used to replace some or all of a missing arm or leg. It may improve movement, balance, or daily activities and is fitted to the person's remaining limb, strength, goals, and comfort. Training helps with safe use and skin care.

## Prosthetics

The design, fitting, and use of artificial replacements for missing body parts, such as arms, legs, teeth, or eyes. Successful use depends on comfort, strength, balance, skin health, goals, and rehabilitation.

## Prosthetic Training

Rehabilitation teaching a person to use an artificial limb safely and effectively. Training includes skin and residual-limb care, putting it on, walking, balance, stairs, transfers, daily activities, and community mobility.

## Prosthetic Valve Endocarditis

A severe endocardial infection affecting an artificial mechanical or bioprosthetic heart valve, categorized as early (within one year of implantation, predominantly caused by *Staphylococcus epidermidis* or

---

*Staphylococcus aureus*) or late (more than one year post-surgery, mimicking native valve endocarditis).

**Alternate Names:**

PVE

**Prosthodontics**

The dental specialty that replaces missing teeth and restores chewing, speech, and appearance. Treatment may use crowns, bridges, implants, dentures, or facial prostheses and is planned around oral health and remaining tissues.

**Protamine**

A medicine that binds to and reverses the blood-thinning effect of heparin. It is given carefully because it can cause low blood pressure, allergic reactions, slow heart rate, or dangerous clotting.

**Protanopia**

Protanopia is complete absence of red-cone function, causing a form of red-green color-vision deficiency and altered perception of red and related colors.

**Protean**

Highly variable in form, appearance, or behavior; in medicine it may describe a disease with changing or diverse manifestations.

**Proteaphagy**

The process by which a cell uses its recycling system to remove whole proteasomes, the structures that normally break down unwanted proteins. It helps adjust protein disposal during stress or changes in nutrition.

**Protease**

An enzyme that breaks proteins into smaller pieces by cutting the chemical links between their amino acids. Proteases help digest food and control many processes inside and outside cells.

**Protease Inhibitors**

Antiviral medicines that block an enzyme viruses need to cut newly made proteins into working parts. They are used for selected viral infections and may

interact with many other medicines.

**Proteasome**

A barrel-shaped protein machine inside cells that breaks down unwanted, damaged, or short-lived proteins. This controls cell growth and signaling and prevents harmful protein buildup.

**Proteasome Binding**

The attachment of a medicine or other molecule to a cell's protein-destroying machine. This can increase, reduce, or otherwise change the breakdown of proteins and may alter cell survival or function.

**Proteasome Inhibitor**

A medicine that blocks the proteasome, a cell structure that breaks down unwanted proteins. These drugs are used especially in multiple myeloma and some blood cancers.

**Protective**

Serving to prevent or reduce injury, infection, or other harm.

**Protective Agent**

A substance, device, or treatment used to reduce harm to the body. Examples include sunscreen protecting skin from ultraviolet light, barrier creams protecting skin, and medicines that protect the stomach or bones during selected treatments. The appropriate agent depends on the hazard and the person's health.

**Protegrin**

A small antimicrobial peptide, usually derived from pig immune cells, that can disrupt microbial membranes.

**Protein**

A large molecule made of one or more chains of amino acids folded into a specific three-dimensional shape. Proteins provide structure and perform functions such as catalyzing reactions, transporting substances, sending signals, regulating cells, and defending against infection; their amino-acid sequence

---

helps determine their shape and function.

### **Proteinaceous Plug**

A blockage made primarily of protein-rich material, mucus, cellular debris, or secretions that can obstruct an airway, duct, catheter, or other passage.

### **Protein Acetylation**

Addition of an acetyl group to a protein, often changing its activity, localization, stability, or interactions; histone acetylation is a major regulator of gene expression.

### **Protein Binding**

The temporary attachment of a medicine to proteins in the blood. The unattached portion can usually leave the bloodstream, reach tissues, and produce effects; changes in protein levels or other medicines can alter the amount available.

### **Protein Biosynthesis**

The way cells make proteins: they copy information from DNA into messenger RNA, then use ribosomes to join amino acids in the instructed order. The new chain folds and may be modified before it works.

### **Protein C**

A natural blood-clotting regulator that helps prevent excessive clotting. Low levels may be inherited or acquired and increase clot risk; starting warfarin in severe deficiency can rarely cause skin injury.

### **Protein C Blood Test**

A protein C blood test measures a natural anticoagulant and may help evaluate unexplained or recurrent blood clots or suspected inherited deficiency.

### **Protein C Deficiency**

A condition in which too little or poorly functioning protein C reduces the body's natural control of blood clotting. It may be inherited or acquired and increases the risk of deep-vein thrombosis and pulmonary embolism.

### **Protein Deficiency**

Inadequate protein intake, digestion, absorption, or use, causing loss of muscle, impaired immunity and wound healing, edema, or growth failure when severe or prolonged.

### **Protein Denaturation**

Loss of a protein's normal three-dimensional structure from heat, chemicals, pH, or mechanical stress, often impairing its biological function without breaking its peptide backbone.

### **Protein Dephosphorylation**

Removal of a phosphate group from a protein by a phosphatase, often reversing or modifying signaling, enzyme activity, localization, or cell-cycle control.

### **Protein Electrophoresis**

A laboratory test that separates proteins in serum or urine according to their electrical properties, helping detect abnormal protein patterns.

### **Protein-Energy Malnutrition**

Severe lack of calories, protein, or both. It can cause weight loss, muscle wasting, weakness, swelling, poor growth, weak immunity, and slow healing; nutrition must often be restored gradually and carefully.

### **Protein Engineering**

Design or modification of proteins through genetic, biochemical, or computational methods to alter activity, stability, specificity, expression, or therapeutic usefulness.

### **Protein Expression**

The process by which a cell uses genetic instructions to make a protein, then folds, changes, and transports that protein so it can perform its function.

### **Protein Folding**

The process by which a newly made protein bends into the three-dimensional shape needed for its job. Incorrect folding can stop the protein from working or cause harmful protein deposits in cells.



---

## Protein Glycosylation

The attachment of sugar chains to proteins. This helps proteins fold, remain stable, reach the correct part of a cell, and work properly; inherited problems with this process can affect many organs.

## Protein Import

Movement of a newly made protein into a cell structure or across a cell membrane. Address labels on the protein and transport systems guide it to the correct location, where it can fold and work properly.

## Protein In Diet

Dietary protein supplies amino acids needed for muscle, enzymes, hormones, immunity, and tissue repair; requirements vary with age, health, and kidney function.

## Protein Isoform

One of several related protein forms produced from different genes, alternative splicing, or post-translational processing, with potentially different locations or functions.

## Protein Kinase

An enzyme that adds a small chemical group to a protein and changes how it works. Protein kinases help control cell signals, metabolism, growth, division, and other functions.

## Protein-Losing Enteropathy

Excessive loss of plasma proteins through the gastrointestinal tract, causing low albumin, swelling, diarrhea, or nutritional deficiency.

## Protein Malnutrition

Poor health caused by inadequate protein or energy intake, leading to weight loss, muscle wasting, weak immunity, or impaired healing.

## Protein Metabolism

The synthesis, folding, modification, breakdown, and recycling of proteins and amino acids in cells and tissues.

## Protein Methylation

Addition of small chemical groups called methyl groups to selected parts of a protein. This can change the protein's activity or location and can alter how cells switch genes on or off.

## Protein Overexpression

Production of an abnormally large amount of a protein, caused by gene amplification, transcriptional activation, altered degradation, or other mechanisms and potentially driving disease or serving as a biomarker.

## Protein Phosphorylation

Attachment of a small phosphate group to a protein by a cell enzyme. This acts like a reversible switch that can change the protein's activity, location, or interactions and helps control cell signals and division.

## Protein Prenylation

Attachment of a fat-like chemical group to a protein, helping position it in a cell membrane. This placement allows some proteins to send signals or help move materials within the cell.

## Protein Purification

Separation of a desired protein from cells or a mixture using properties such as charge, size, solubility, affinity, or binding, while preserving its structure and activity.

## Protein Refolding

Return of a denatured or newly synthesized protein to its functional three-dimensional structure, sometimes requiring chaperones or controlled laboratory conditions.

## Protein S

A natural blood protein that helps activated protein C limit clotting. Too little or abnormal protein S can increase the risk of clots in deep veins or the lungs.

---

### **Protein S Blood Test**

A protein S blood test measures a natural anticoagulant and may help evaluate suspected inherited or acquired tendency to form blood clots.

### **Protein S Deficiency**

A condition in which the blood has too little or poorly functioning protein S, reducing natural control of clotting. It may be inherited or acquired and can increase the risk of deep-vein thrombosis or pulmonary embolism.

### **Protein Secretion**

Release of a protein from a cell or into a particular cell compartment. Cells use this process to send hormones, digestive enzymes, antibodies, and other proteins to where they are needed.

### **Protein Sequence**

The precise order of amino acids joined together to make a protein. This order helps determine the protein's three-dimensional shape, location, interactions, and biological job.

### **Protein Splicing**

A process in which a newly made protein removes an internal section and joins the remaining sections together. The finished protein may then fold into the form needed for its normal function.

### **Protein Subunit**

One protein component of a larger protein assembly. Several subunits may join to create the final structure and allow the assembly to perform its biological function.

### **Protein Synthesis**

The process by which a cell uses messenger RNA instructions to join amino acids into a protein. The resulting protein folds into a working shape and may be changed or moved before performing its function.

### **Protein Translocation**

Movement of a newly synthesized or mature protein across a membrane or to a different cellular compart-

ment, directed by targeting signals and transport machinery.

### **Protein Transport**

Cellular movement of proteins between compartments, across membranes, or outside the cell, ensuring that each protein reaches its functional location.

### **Protein Unfolding**

Loss of a protein's native three-dimensional structure, caused by heat, chemicals, mutations, or cellular stress, which can impair function or promote aggregation.

### **Proteinuria**

An unusually high amount of protein in urine. It can result from kidney disease, diabetes, high blood pressure, infection, exercise, fever, or pregnancy and may require repeat testing to find the cause.

### **Proteobacteria**

A large group of bacteria containing many gram-negative species, including *Escherichia*, *Salmonella*, *Pseudomonas*, and *Vibrio*. Some are harmless or helpful, while others cause intestinal, urinary, lung, wound, or bloodstream infections.

### **Proteogenomics**

Integration of genomic, transcriptomic, and proteomic data to identify proteins, discover coding variants, characterize tumors or microbes, and connect DNA changes with protein function.

### **Proteoglycan**

A large molecule made of a protein with many attached sugar chains. It helps connective tissues and cartilage hold water, resist pressure, maintain structure, and send signals between cells.

### **Proteolipid Protein**

A protein containing a substantial fat component, especially important in the myelin sheath around nerve fibers.

---

## Proteolipids

Proteins tightly associated with lipid molecules or lipid membranes, important in myelin, mitochondrial membranes, ion transport, and other membrane structures.

## Proteolysis

Enzymatic breakdown of proteins into smaller peptides or amino acids. It is needed for digestion, cell regulation, tissue remodeling, and removal of damaged proteins.

## Proteolytic

Relating to the breakdown of proteins by cutting the links between their amino acids. Proteolytic enzymes help digest food, activate clotting and immune proteins, remove damaged proteins, and control cell death; medicines can block some of these enzymes.

## Proteome

The complete set of proteins produced by a cell, tissue, organism, or biological condition at a particular time.

## Proteomic Profile

A proteomic profile is a pattern of proteins measured in a biological sample to study disease, treatment response, or biological processes.

## Proteomics

The large-scale study of proteins, including their abundance, structure, modifications, interactions, and functions. Large-scale study of proteins, including their amounts, structures, changes, interactions, and roles in cells or disease.

## Proteostasis

The balance of protein production, folding, movement, and breakdown that keeps cells functioning. Failure of proteostasis can contribute to aging, neurodegeneration, and other diseases.

## Proteus

A group of Gram-negative bacteria that can cause

urinary, wound, bloodstream, or other infections. *Proteus* urinary infections may produce urease, raising urine pH and encouraging certain kidney stones.

## *Proteus* Hauseri

A gram-negative bacterium in the *Proteus* group that is rarely isolated from human clinical specimens.

## *Proteus* Mirabilis

A gram-negative bacterium that commonly causes urinary-tract infection and can produce urease-related kidney or bladder stones.

## *Proteus* Penneri

A gram-negative *Proteus* species that can rarely cause urinary, wound, bloodstream, or other opportunistic infections.

## Proteus Syndrome

A rare mosaic overgrowth disorder, usually caused by AKT1 variants, producing asymmetric enlargement of bones, skin, or other tissues and an increased risk of tumors.

## *Proteus* Vulgaris

A gram-negative bacterium that can cause urinary, wound, bloodstream, and other infections and often produces urease.

## Prothionamide

A thioamide antimycobacterial drug used as part of selected multidrug regimens for drug-resistant tuberculosis; gastrointestinal, hepatic, neurologic, and thyroid adverse effects require monitoring.

## Prothrombin

A blood protein made with the help of vitamin K. When a blood vessel is injured, it is changed into thrombin, which helps form a stable blood clot.

## Prothrombin Deficiency

A rare bleeding disorder caused by too little or poorly functioning prothrombin, a protein needed for blood clotting. It can cause easy bruising, nose-

---

bleeds, heavy menstrual bleeding, or prolonged bleeding after injury or surgery.

### **Prothrombin Time**

A blood test measuring how quickly a sample forms a clot. It is often reported as the INR and helps assess clotting disorders, liver disease, vitamin K deficiency, and the effect of warfarin.

#### **Alternate Names:**

PT

### **Proton Pump Inhibitors**

Proton pump inhibitors suppress gastric acid secretion and treat reflux, ulcers, and other acid-related disorders; prolonged or inappropriate use should be periodically reviewed.

#### **Alternate Names:**

PPIs; Proton Pump Inhibitor Medicines

### **Proton Therapy**

Radiation treatment using high-energy protons to damage cancer cells. The beam can deliver much of its dose at a selected depth, potentially reducing radiation to nearby healthy tissue; suitability depends on the tumor and treatment plan.

### **Proto-Oncogene**

A normal gene that helps control cell growth, division, or survival. If it becomes abnormally active because of a genetic change or extra copies, it can act as an oncogene and contribute to cancer.

### **Protozoa**

A traditional group of single-celled organisms with nuclei. Some cause human disease, including malaria, giardiasis, and amoebiasis; modern biological classification places them in several different groups.

### **Protozoal Infection**

Infection caused by a disease-producing protozoan parasite. Protozoal infections may affect the intestine, blood, brain, eyes, skin, or genital tract and include malaria, amoebiasis, giardiasis, leishmaniasis, toxoplasmosis, trichomoniasis, and trypanosomiasis;

transmission and treatment depend on the organism and site of disease.

#### **Alternate Names:**

Protozoan Disease; Protozoiasis

### **Protractile**

Able to move forward or outward from its usual position. The tongue and lower jaw are examples of body structures that can be actively protruded.

### **Protractor**

An instrument or muscle that moves a body part forward or outward. In anatomy, a protractor muscle may move the jaw, shoulder blade, or another structure away from its usual position.

### **Protrusion**

A part that projects outward or a disc or tissue displaced beyond its normal boundary.

### **Protuberance**

A protuberance is a structure or area that projects outward from a surface, such as a normal bone prominence, swelling, or mass.

### **Provisional Diagnosis**

A temporary working diagnosis made from available evidence before all test results are known, and subject to change.

### **Provisional Tic Disorder**

A disorder with motor or vocal tics lasting less than one year, beginning in childhood and not better explained by another condition.

### **Provocation**

An action or stimulus that brings on a symptom, response, or test finding. Clinicians may use controlled provocation during examination, but it should be stopped if it causes dangerous pain or other harm.

### **Proximal**

Proximal means nearer to the point of attachment or origin; in a limb, the shoulder or hip end is proximal to the hand or foot.

---

### Proximal Deep Vein Thrombosis

A blood clot in a major deep vein of the leg at or above the knee, such as the popliteal, femoral, or iliac vein. It has a higher risk of traveling to the lungs than a clot limited to the calf. It may cause one-sided swelling, warmth, redness, or pain, but some people have no symptoms. Ultrasound and the clinical assessment help establish the diagnosis, and treatment usually includes anticoagulation when safe.

#### Alternate Names:

Proximal DVT; Above-Knee Deep Vein Thrombosis; Proximal Lower Extremity DVT

### Proximal DVT

Alternate name; see *Proximal Deep Vein Thrombosis*.

### Proximal Humerus Fractures

Breaks near the upper end of the upper-arm bone, close to the shoulder. They may cause pain, swelling, bruising, and difficulty moving the arm; treatment depends on displacement, joint involvement, bone quality, nerve or blood-vessel injury, and the person's needs.

### Proximal Radioulnar Joint

The joint near the elbow where the two forearm bones rotate around each other. It allows the palm to turn up or down; injury or dislocation can limit rotation and cause pain.

### Proximal Renal Tubular Acidosis

Proximal renal tubular acidosis occurs when the kidney cannot reclaim enough bicarbonate, causing excess acid in the blood and often low potassium.

### Proximal Tubule

The first part of the kidney's microscopic filtering tube after the filtering unit. It returns much of the filtered water, salt, bicarbonate, sugar, amino acids, and phosphate to the blood.

### Proximal Tubule Reabsorption

The kidney's return of much of the filtered water

and useful substances, including salt, bicarbonate, glucose, amino acids, and phosphate, from the first tubule back into the blood.

### Prune Belly Syndrome

A condition present at birth, usually in boys, involving weak or absent abdominal muscles, undescended testicles, and abnormalities of the urinary tract. Kidney problems, repeated infections, breathing difficulty, and impaired growth may occur.

### Prurigo

A group of intensely itchy skin disorders with papules or nodules caused by inflammation and scratching.

### Prurigo Nodularis

A long-lasting skin disorder with intensely itchy, firm bumps called nodules. Repeated scratching or rubbing causes and maintains an itch-scratch cycle; treatment aims to reduce itching, heal the skin, and prevent new nodules.

### Pruritic

Itchy or causing an urge to scratch; pruritus can result from skin inflammation, allergy, infection, systemic disease, medications, or neuropathic mechanisms.

### Pruritic Eruption

An itchy skin eruption that may result from allergic, inflammatory, infectious, drug-related, or autoimmune disease; diagnosis depends on its morphology, distribution, timing, and associated findings.

### Pruritis

The medical term for itching, an unpleasant urge to scratch. Causes include dry skin, allergy, inflammation, infection, medicines, nerve disorders, and internal disease.

### Pruritis Ani

Alternate name; see *Anal itching*.

---

## Pruritus

An unpleasant itchy feeling that creates the desire to scratch. It may result from dry skin, allergy, infection, medicines, liver or kidney disease, nerve problems, or other conditions; severe or widespread itching needs assessment.

## Pruritus and Systemic Disease

Generalized or persistent itching can be a manifestation of kidney or liver disease, iron deficiency, thyroid disease, hematologic disease, diabetes, infection, or malignancy. Examination and targeted tests are guided by the history and associated findings.

## Pruritus Ani

Persistent itching around the anus with multiple potential causes including hemorrhoids, fissures, infections such as pinworms and candidiasis, dermatologic conditions including psoriasis and lichen sclerosus, and idiopathic causes.

## Pruritus In CKD

Persistent itching associated with advanced chronic kidney disease, often worse in people receiving dialysis. It may occur without a visible rash and can disturb sleep and quality of life; treatment is individualized after other causes are assessed.

## Prussian Blue

A blue pigment used in laboratory staining and as an antidote for selected radioactive cesium or thallium poisoning.

## PSA

Prostate-specific antigen, a protein made mainly by prostate cells and measured in blood. A raised level may occur with enlargement, inflammation, infection, or cancer and does not diagnose cancer alone.

### Alternate Names:

Prostate-Specific Antigen; Total PSA

## Psammoma

A tumor or lesion containing rounded, layered calcium deposits called psammoma bodies. These de-

posits can occur in several tumors and are identified by microscopic examination rather than symptoms alone.

## PSA Test

A laboratory blood test that measures prostate-specific antigen (PSA), a protein made mainly by prostate cells. An elevated level may occur with prostate cancer, benign prostate enlargement, or infection or inflammation of the prostate, so the result does not diagnose cancer by itself and must be interpreted with the clinical context and other findings.

## PSA Velocity

The rate at which prostate-specific antigen concentration changes over time. It may provide additional information in prostate evaluation but should not be used alone to diagnose cancer.

## Pseudarthrosis

Failure of a broken bone to heal, leaving persistent movement or a false joint at the fracture site. It can cause pain, weakness, deformity, and loss of function and may require surgery.

## Pseudoaneurysm

A weak area or leak in an artery in which blood collects outside the normal artery but remains connected to it. It may follow injury, surgery, infection, or a catheter procedure and can enlarge, burst, or send clots downstream.

## Pseudoangiomatous Stromal Hyperplasia

A benign breast stromal proliferation forming slit-like spaces within dense collagen that may mimic vascular channels. It can be incidental or present as a palpable or imaging-detected mass.

## Pseudobulbar Affect

Sudden, difficult-to-control episodes of laughing or crying that do not match how a person feels or the situation. It can occur after stroke or with other brain and nerve diseases and is different from depression.

### Alternate Names:



---

PBA; Emotional Incontinence; Pathological Laughter and Crying

### **Pseudocholinesterase Deficiency**

Reduced activity of butyrylcholinesterase, inherited or acquired, causing prolonged paralysis after medicines such as succinylcholine or mivacurium. People may need extended breathing support after anesthesia; enzyme or genetic testing can help confirm the condition.

### **Pseudoclubbing**

Nail-tip broadening or flattening that resembles clubbing but does not have the characteristic change in the angle between the nail and surrounding skin. It may occur with some bone, joint, skin, or inherited conditions and needs assessment of the underlying cause.

### **Pseudo-Cushing Syndrome**

A reversible state with clinical or biochemical features resembling cortisol excess without autonomous Cushing disease. Severe alcohol use, depression, obesity, sleep disorders, or acute illness may contribute, and repeat or specialized testing may be needed.

### **Pseudocyesis**

A condition in which a person has pregnancy-like symptoms, such as missed periods, abdominal enlargement, nausea, or breast changes, without an actual pregnancy. A pregnancy test and examination distinguish it from pregnancy and identify other possible causes.

### **Pseudodementia**

Memory and thinking complaints caused by depression or another potentially reversible problem rather than progressive brain-cell loss. Careful assessment is needed because depression and true dementia can occur together or resemble each other.

### **Pseudoephedrine**

An oral decongestant that shrinks swollen blood vessels in the nose and improves blockage from colds or sinus problems. It may raise blood pressure, cause sleeplessness or palpitations, and interact with some

medicines.

### **Pseudofolliculitis of the Beard**

An inflammatory reaction caused by shaved or trimmed hairs curling back into the skin, producing itchy or tender papules and pustules in the beard area. Avoiding close shaving, changing shaving technique, and treating inflammation can reduce recurrence and scarring.

### **Pseudogout**

A joint disease caused by calcium pyrophosphate crystals collecting in cartilage. It can cause sudden pain, swelling, warmth, and stiffness, often in a knee or wrist, and may also produce longer-term joint symptoms.

#### **Alternate Names:**

Calcium Pyrophosphate Dihydrate Deposition Disease; CPPD

### **Pseudogout Flare**

A sudden attack of joint pain and swelling caused by calcium pyrophosphate crystals. The knee or wrist is often affected; the joint may be red and warm, and fluid testing can confirm the diagnosis.

### **Pseudogynecomastia**

Bilateral enlargement of male breasts due to excessive subareolar and chest wall adipose tissue accumulation without true glandular or ductal tissue proliferation, commonly observed in obesity and distinguishable from true gynecomastia by the absence of a firm, distinct subareolar glandular disc on palpation.

#### **Alternate Names:**

Lipomastia; Adipomastia

### **Pseudohypertrophic Muscular Dystrophy**

An older name for Duchenne or related muscular dystrophy with enlarged-appearing muscles caused by fat and connective tissue.

### **Pseudohypoglycemia**

A falsely low glucose measurement that does not reflect the blood glucose available to tissues. It may

---

result from poor peripheral circulation, shock, severe edema, or delayed processing of a blood sample and should be checked with a reliable venous or alternative-site measurement.

### **Pseudohypoparathyroidism**

A group of inherited disorders in which the body does not respond properly to parathyroid hormone. This can cause low calcium, high phosphate, muscle cramps, seizures, and sometimes short fingers or other physical features.

### **Pseudo-Kaposi Sarcoma (Acroangiodermatitis)**

A benign vascular eruption, usually on the lower legs or feet, that can resemble Kaposi sarcoma. It is often associated with chronic venous insufficiency, arteriovenous malformations, or dialysis access and may cause violaceous plaques or ulcers.

### **Pseudomelanosis Duodeni**

A usually harmless dark discoloration of the first part of the small intestine seen during endoscopy. It results from pigment-containing material in the lining and may be associated with chronic illness, medicines, or mineral deposits.

### **Pseudomembranous Colitis**

Severe inflammation of the colon, usually caused by overgrowth of *Clostridioides difficile* after antibiotics. It can cause frequent watery diarrhea, fever, abdominal pain, dehydration, and serious complications.

#### **Alternate Names:**

*Clostridioides Difficile* Colitis; Antibiotic-Associated Colitis

### **Pseudomonas**

A group of Gram-negative bacteria that can cause opportunistic infections, especially in damaged tissues or people with weakened immunity.

### **Pseudomonas Aeruginosa**

A gram-negative bacterium that can cause pneumonia, bloodstream, urinary, wound, and ear infections, especially in hospitalized or immunocompromised

people; it often has antibiotic resistance.

### **Pseudomonas Aeruginosa Infection**

Infection caused by *Pseudomonas aeruginosa*, an opportunistic gram-negative bacterium that can affect the lungs, wounds, urinary tract, blood, or other sites.

### **Pseudomonas Aeruginosa Resistance**

The ability of *Pseudomonas aeruginosa* to survive medicines that would normally kill it. Resistance may be natural or develop through genetic changes, so laboratory testing is needed to choose an effective antibiotic.

### **Pseudomonas Folliculitis**

An infection of hair follicles by *Pseudomonas aeruginosa*, often after exposure to contaminated hot tubs or pools. It causes itchy red papules and pustules under swimwear and usually resolves, although severe or recurrent disease may need medical treatment.

### **Pseudomyxoma Peritonei**

A rare condition in which jelly-like tumor material gradually fills the abdomen, usually after a mucus-producing appendix tumor ruptures. It can cause increasing abdominal size, pain, bowel blockage, or poor nutrition and requires specialist treatment.

### **Pseudophakia**

Having an artificial lens inside the eye after the natural lens has been removed, usually during cataract surgery. The replacement lens focuses light, but later clouding of its supporting capsule or other eye problems can still affect vision.

### **Pseudoporphyria**

A blistering and skin-fragility disorder that resembles porphyria cutanea tarda but occurs without the characteristic porphyrin excess. It may be triggered by kidney failure, dialysis, medicines, or ultraviolet exposure and usually affects the backs of the hands.

### **Pseudopseudohypoparathyroidism**

An inherited condition caused by paternally trans-

---

mitted mutations in the *GNAS* gene, characterized by the somatic phenotype of Albright hereditary osteodystrophy (short stature, round face, brachydactyly, subcutaneous ossifications) in the presence of completely normal calcium, phosphorus, and parathyroid hormone levels due to intact maternal allele expression in renal tubules.

**Alternate Names:**

PPHP

**Pseudorabies**

Aujeszky disease, a herpesvirus infection mainly affecting pigs and other animals. People are rarely infected, but exposed humans may develop serious nervous-system disease; it is different from rabies.

**Pseudothrombocytosis Due To Red Cell Fragments**

A falsely high platelet count caused when automated equipment mistakes very small broken red-cell fragments for platelets. A blood-smear examination can distinguish this error from a genuinely high platelet count.

**Pseudotumor Cerebri**

An older name for idiopathic intracranial hypertension, a condition in which pressure around the brain rises without a tumor. It can cause daily headache, brief dimming of vision, whooshing in the ears, and swelling of the optic nerves that may threaten sight.

**Pseudoxanthoma Elasticum**

A rare inherited disorder in which elastic tissue becomes hardened and damaged. It can cause yellowish or loose skin, bleeding or vision loss from eye blood vessels, and narrowing or bleeding in arteries.

**Alternate Names:**

PXE; Gronblad-Strandberg Syndrome

**Psittacosis**

An infection caused by *Chlamydia psittaci*, usually acquired by breathing dust from infected birds or their droppings. It commonly causes fever, headache, cough, and pneumonia and requires appropriate antibiotic treatment.

**Psittacosis (Parrot Fever)**

A systemic infection caused by *Chlamydia psittaci*, usually acquired from infected birds or their dried secretions. It commonly causes atypical pneumonia with fever, headache, cough, and sometimes hepatitis or neurologic disease; exposure history is diagnostically important.

**PSMA**

A protein found in large amounts on many prostate cancer cells and in smaller amounts in some normal tissues. Special tracers can use it to show prostate cancer on scans or deliver radiation to cancer cells.

**Psoas**

A large muscle running from the lower spine to the thigh bone. It helps bend the hip, lift the leg, maintain upright posture, and stabilize the spine during movement.

**Psoas Abscess**

A collection of pus in or around the psoas muscle, caused by infection spreading through the blood or from nearby tissues. It may cause back or hip pain, fever, difficulty walking, or few symptoms and usually requires imaging and targeted treatment.

**Psoas Sign**

An abdominal physical examination sign elicited by passive hyperextension of the right hip with the patient in the left lateral decubitus position, or active right hip flexion against resistance; exacerbation of right lower quadrant abdominal pain indicates retrocecal appendicitis inflaming the underlying psoas muscle.

**Psoriasis**

A long-term immune-mediated disease that causes inflammation and faster replacement of skin cells, leading to raised, thick, dry, scaly patches. Patches may be red, purple, brown, darker, or lighter than nearby skin and often affect the scalp, elbows, knees, trunk, palms, or soles. Nails may become pitted, ridged, thick, or separated from the nail bed. Symptoms may flare and then improve. Psoriasis is not contagious and may occur with psoriatic arthritis,

---

which affects joints and tendon attachments.

### **Psoriatic Arthritis**

An inflammatory arthritis associated with psoriasis of the skin or nails. It may affect the spine, peripheral joints, digits, or tendon insertions, and diagnosis uses the clinical pattern together with appropriate laboratory and imaging findings.

#### **Alternate Names:**

PsA; Arthropathia Psoriatica

### **Psoriatic Plaque**

The characteristic cutaneous lesion of plaque psoriasis (psoriasis vulgaris), consisting of a sharply demarcated, erythematous, indurated plaque covered with adherent, silvery-white micaceous scales, typically distributed symmetrically over extensor surfaces (elbows, knees, scalp, lumbosacral region).

### **PSP**

PSP commonly means progressive supranuclear palsy, a neurodegenerative disorder causing early postural instability, vertical gaze impairment, rigidity, and cognitive change.

### **Psychiatric**

Relating to the assessment, diagnosis, treatment, or prevention of disorders of mood, thought, perception, behavior, or emotional functioning.

### **Psychiatric Emergency**

A mental-health condition involving an immediate and serious risk to life, safety, basic functioning, or the safety of others. Examples include imminent suicide risk, severe psychosis, dangerous agitation, catatonia, or severe substance intoxication or withdrawal. It is a subset of mental-health crises distinguished by the immediacy and severity of risk.

### **Psychiatrist**

A medical doctor who assesses, diagnoses, and treats mental-health conditions. A psychiatrist may provide talking therapy, prescribe medicines, coordinate other treatments, and assess how mental or physical illness affects safety and daily life.

### **Psychiatry**

The medical specialty concerned with mental health and the diagnosis and treatment of mental disorders.

### **Psychoacoustics**

The study of how people perceive and interpret sound, including loudness, pitch, masking, localization, speech, and hearing thresholds.

### **Psychoactive**

Affecting mood, thought, perception, behavior, alertness, or other mental processes. Psychoactive substances include prescribed medicines, alcohol, and recreational drugs.

### **Psychoactive Substance**

A substance that alters mood, perception, cognition, behavior, or consciousness.

### **Psychoanalysis**

A psychological theory and talking treatment emphasizing unconscious thoughts, feelings, memories, and conflicts. Traditional treatment explores patterns in relationships and behavior, while modern approaches vary in methods, goals, and evidence.

### **Psychodermatology**

The area of care that considers links between skin conditions and emotional health. Stress can worsen problems such as eczema, psoriasis, acne, or hives, while visible skin disease can cause anxiety or low mood; both aspects may need treatment.

### **Psychodidae**

A family of small hairy flies that includes sand flies; some species transmit pathogens such as *Leishmania* and therefore have public-health importance.

### **Psychodrama**

A therapy using guided acting to explore emotions, relationships, and behavior in a supportive setting.

### **Psychodynamic Therapy**

Talking therapy that explores how past experiences, relationships, feelings, and patterns outside a per-

---

son's immediate awareness may influence current distress and behavior. The therapist and patient use the therapeutic relationship to develop insight and healthier ways of coping.

### **Psychogenic**

Caused or strongly influenced by psychological factors such as stress, trauma, or emotional conflict. The symptoms are real and not deliberately produced, and physical causes should still be considered.

### **Psychogenic Erectile Dysfunction**

Difficulty getting or keeping an erection mainly related to stress, anxiety, depression, relationship concerns, or other emotional factors. Physical causes can coexist, so medical assessment should consider health, medicines, and psychological wellbeing.

### **Psychogenic Non-Epileptic Seizures**

Episodes that resemble epileptic seizures but are not caused by the abnormal electrical discharges of epilepsy. They are real and involuntary and may involve shaking, stiffening, unresponsiveness, or altered movement; video-EEG can help establish the diagnosis.

### **Psycholinguistics**

The study of how the brain and mind produce, understand, acquire, and remember language, including speech, reading, grammar, and meaning.

### **Psychological**

Relating to thoughts, feelings, behavior, or the way these affect health. Psychological factors can influence pain, sleep, coping, treatment choices, and recovery, but describing a factor as psychological does not mean symptoms are imagined.

### **Psychological Adjustment**

The process of adapting thoughts, emotions, and behavior to stress, illness, disability, loss, developmental change, or other life circumstances.

### **Psychological Dependence**

A pattern in which a person feels a strong emotional or mental need for a substance or behavior, often with craving, preoccupation, or distress when it is unavailable.

### **Psychological Health**

Alternate name; see *Mental Health*.

### **Psychological Homeostasis**

The ongoing adjustment of thoughts, emotions, and behavior that helps a person maintain psychological stability while responding to stress or change. Severe disruption can contribute to anxiety, depression, or impaired daily functioning.

### **Psychological Support**

Emotional and practical help for people coping with illness, disability, pain, or difficult treatment. It may include counseling, coping skills, support groups, and assessment or treatment of anxiety, depression, and adjustment problems.

### **Psychological Transfer**

The process in which feelings, expectations, or relationship patterns from earlier experiences become directed toward someone in the present, especially a therapist. Recognizing this pattern can help understand distress and relationships during therapy.

### **Psychological Trauma**

Emotional or psychological effects that may follow an event or ongoing situation involving actual or threatened death, serious injury, violence, abuse, or another overwhelming experience. Responses vary and may include fear, intrusive memories, avoidance, emotional numbness, sleep problems, or changes in mood and behavior.

### **Psychology**

The scientific study of behavior, emotion, learning, and mental processes. Psychological care may include assessment, counseling, and behavioral therapies.



---

## **Psychometrics**

The design and use of tests that measure mental abilities, feelings, behavior, or personality. It also examines whether a test gives consistent results and truly measures the quality it claims to measure.

## **Psychomotor**

Relating to the connection between mental activity and physical movement, including the speed, coordination, and purposeful control of actions.

## **Psychomotor Agitation**

Restless, excessive, or purposeless movement associated with inner tension, anxiety, mania, delirium, depression, substance effects, or medication reactions.

## **Psychomotor Disorder**

A condition affecting the speed, starting, coordination, or expression of movement and mental activity. A person may move unusually slowly or quickly, seem restless, or have difficulty organizing purposeful actions.

## **Psychomotor Performance**

The ability to coordinate mental processing with physical movement, assessed through tasks involving speed, precision, reaction time, balance, or sequencing.

## **Psychomotor Retardation**

A noticeable slowing of mental processes, speech output, and physical movements, manifested by decreased spontaneous gestures, prolonged latency in answering questions, sluggish gait, and reduced facial expressiveness. It is a core feature of severe major depressive disorder and melancholia.

## **Psychoneuroimmunology**

The study of how thoughts, emotions, the brain, hormones, and the immune system influence one another in health and illness. It examines links involving stress, inflammation, infection, and recovery.

## **Psycho-Oncology**

The clinical field addressing psychological, social, behavioral, and communication needs of people with cancer and their families during diagnosis, treatment, survivorship, and end-of-life care.

## **Psychopathic**

Describing severe, persistent antisocial traits such as limited empathy, deceitfulness, and disregard for others; it is not a precise diagnosis.

## **Psychopathology**

The study and description of abnormal mental processes, symptoms, behaviors, and mechanisms underlying psychiatric disorders. The study or description of abnormal mental states and behavior.

## **Psychopharmacology**

The study of how medicines affect mood, behavior, thinking, sleep, and other mental functions in people.

## **Psychophysics**

The study of how physical stimuli, such as light, sound, pressure, or heat, are experienced by a person. It measures detection thresholds, ability to distinguish stimuli, and perceived strength.

## **Psychophysiology**

The study of links between psychological processes and measurable bodily functions such as heart rate, skin conductance, muscle activity, and brain signals.

## **Psychosexual**

Relating to how psychological factors interact with sexual development, feelings, behavior, relationships, or function. Mental health, past experiences, body image, culture, medicines, and physical conditions can all influence sexuality.

## **Psychosexual Development**

Development of sexual identity, interest, relationships, and behavior across the lifespan as influenced by biology, cognition, attachment, culture, and experience.



---

## Psychosis

A state in which a person loses contact with reality. It may involve strongly held false beliefs, hearing or seeing things others do not, confused speech, or markedly disorganized behavior and can result from mental, medical, neurologic, or substance-related causes.

## Psychosocial Rehabilitation

Support that helps a person with mental illness build skills, relationships, confidence, and independence. It may include daily-living training, education, work support, peer support, therapy, and help taking part in family and community life.

## Psychosocial Support

Practical, emotional, and social help for people facing illness or difficult treatment. It may include counseling, support groups, help with work or money worries, family support, and care for anxiety, depression, fear, loneliness, or grief.

## Psychosomatic

Relating to physical symptoms influenced by psychological processes or stress.

## Psychosomatic Disorder

An older broad term for physical symptoms influenced by psychological and social factors. The symptoms are real; modern assessment considers both body and mind rather than assuming symptoms are imagined.

## Psychosomatic Medicine

Medical care that considers how body processes, thoughts, emotions, relationships, and social conditions interact in illness. Physical symptoms are real, and assessment looks for both medical and psychological factors without assuming symptoms are imagined.

## Psychosurgery

Surgery intended to alter brain circuits for severe, treatment-resistant psychiatric illness; it is rarely used and highly specialized.

## Psychotherapy

Structured treatment using therapeutic communication and psychological techniques to reduce distress, change unhelpful patterns, and improve coping or relationships. Common approaches include cognitive behavioral and interpersonal therapies.

## Psychotic

Describing psychosis, a syndrome involving impaired reality testing. It may include delusions, hallucinations, disorganized thinking or speech, or markedly disorganized behavior. Psychosis can occur with schizophrenia-spectrum disorders, severe mood disorders, substances or medicines, delirium, neurologic disease, or other medical conditions.

## Psychotic Disorder

Alternate name; see *Psychotic Disorders*.

## Psychotic Disorders

A group of mental disorders characterized by impaired reality testing, such as delusions, hallucinations, disorganized thought or speech, or markedly disorganized behavior. The group includes schizophrenia-spectrum disorders and other conditions in which psychosis is a defining feature; psychotic symptoms can also occur with mood disorders, substances, medicines, neurologic disease, or other medical conditions.

## PTCA

Percutaneous transluminal coronary angioplasty, a procedure using a small balloon to open a narrowed heart artery. A stent is often placed afterward to help keep the artery open and restore blood flow.

### Alternate Names:

Percutaneous Transluminal Coronary Angioplasty; Coronary Angioplasty

## PTCL

Alternate name; see *Peripheral T-cell lymphoma*.

## PTE

Alternate name; see *Pulmonary Thromboendarterectomy*.

---

## Pteridine

A nitrogen-containing chemical structure found in folate vitamins, natural pigments, and other active body compounds. Pteridine-containing molecules participate in cell growth, chemical reactions, color production, and inherited metabolic disorders.

## Pterygium

A raised, often triangular growth of the clear covering of the eye that extends onto the cornea. Long-term sun, wind, dust, or dry-eye exposure may contribute; it can cause irritation, redness, blurred vision, or astigmatism.

## Pthirus Pubis

The insect commonly called the pubic louse. It lives mainly in coarse genital hair and causes intense itching and visible lice or eggs; it spreads mainly through close body or sexual contact.

## PTH-Related Peptide

A hormone-like substance sometimes made in excess by cancer cells. It acts partly like parathyroid hormone and can raise blood calcium, causing thirst, constipation, weakness, confusion, dehydration, or kidney problems.

## Ptilosis

An abnormal hair pattern or hair disorder, including excess growth, unusual direction, or loss. The term is nonspecific, so the body site, appearance, timing, and underlying cause must be assessed.

## PTLD

Post-transplant lymphoproliferative disorder, an abnormal overgrowth of immune cells after organ or stem-cell transplantation. It ranges from a reversible response to lymphoma and may cause fever, swollen lymph nodes, organ problems, or unexplained weight loss.

## Ptoxis

Drooping of an eyelid or another body structure. Eyelid ptosis may result from muscle, nerve, tendon, aging, injury, or a neurologic disorder.

## PTSD

Alternate name; see *Post-Traumatic Stress Disorder*.

## Ptyalin

An older name for salivary amylase, the enzyme that begins starch digestion in the mouth.

## Ptyalism

Excessive saliva or drooling. It may result from increased saliva production or difficulty swallowing, and can occur with pregnancy, mouth irritation, reflux, medicines, or disorders affecting the brain, nerves, or muscles.

## Pubertal Failure

Puberty that does not begin or stops progressing as expected. Causes may involve the brain's hormone centers, pituitary gland, testes or ovaries, chromosomes, chronic illness, inadequate nutrition, excessive exercise, or other factors.

## Puberty

The stage when hormone changes cause the body to mature sexually and become capable of reproduction. It includes growth, development of breasts or testes and genitals, body hair, voice changes, and menstrual periods or sperm production.

## Puberty In Boys

Puberty in boys is the hormone-driven development of testes, penis, body hair, voice, muscle, and fertility, usually progressing through predictable physical stages.

## Puberty In Girls

Puberty in girls is the hormone-driven development of breasts, body hair, growth, menstrual cycles, and reproductive maturity, usually progressing through predictable stages.

## Pubes

The area in front of the pelvis over the pubic bone, or the coarse hair that grows there during puberty. Changes in pubic hair can help assess sexual devel-

---

opment, but must be interpreted with other physical findings.

### **Pubic**

Relating to the pubis, the front part of the pelvis, or the area covered by hair above the genitals.

### **Pubic Lice**

Pubic lice are parasitic insects infesting coarse body hair, causing intense itching and visible nits; treatment and assessment of close or sexual contacts are needed.

### **Pubic Symphysis**

The joint at the front of the pelvis where the two pubic bones meet. A strong pad of cartilage joins the bones and allows very little movement.

### **Pubis**

The front part of the pelvic bone, meeting its opposite side at the pubic symphysis.

### **Public Assistance**

Government or community support providing eligible people with food, housing, income, healthcare, disability, or other essential services.

### **Public Exposure**

Radiation received by other people from a patient who has undergone nuclear medicine treatment. The amount depends on the radioactive substance and treatment; the medical team gives specific instructions to limit exposure when needed.

### **Public Health**

The science and practice of protecting and improving the health of communities through education, policy-making, and research for disease and injury prevention.

### **Public Health Emergency**

A serious event threatening the health of many people, such as a pandemic, disaster, or deliberate biological attack.

### **Puborectalis Muscle**

A loop-shaped pelvic muscle that surrounds the lower rectum and helps control bowel movements. At rest it pulls the rectum forward to make a bend that helps hold stool; relaxing it helps stool pass.

### **Pudendal**

Relating to the external genitalia, anus, or perineum. The pudendal nerve carries sensation from these areas and helps control several muscles involved in continence and sexual function.

### **Pudendal Arteries**

Arteries supplying blood to the external genital and perineal region.

### **Pudendal Block**

Injection of local anesthetic near the pudendal nerve to numb the lower vagina, vulva, anus, and perineum. It may reduce pain during late labor, assisted delivery, or repair of a birth-related tear.

### **Pudendal Nerve**

The nerve carrying sensation and motor signals to much of the perineum and pelvic floor.

### **Pudendal Nerve Block**

A regional obstetric anesthesia technique in which local anesthetic is infiltrated transvaginally near the ischial spine and sacrospinous ligament to block the pudendal nerve (S2-S4), providing sensory anesthesia to the perineum, vulva, and lower vagina for forceps delivery or episiotomy repair.

### **Puerperal**

Relating to the period after childbirth, usually the first six weeks. During this time the body recovers from pregnancy and birth, the womb becomes smaller, bleeding gradually settles, and problems such as infection, heavy bleeding, or blood clots may occur.

### **Puerperal Disorder**

A physical or mental health disorder occurring during the postpartum period, including infection,

---

bleeding, hypertension, depression, or psychosis.

### **Puerperal Fever**

Fever occurring after childbirth, often caused by infection of the uterus, birth canal, urinary tract, breast, wound, or bloodstream. It requires prompt assessment because untreated infection can spread and become life-threatening.

### **Puerperal Infection**

An infection after childbirth, commonly involving the uterus, surgical wound, urinary tract, breast, or bloodstream, often causing fever, pain, or foul discharge.

### **Puerperal Psychosis**

A rare, severe mental-health emergency that begins soon after childbirth. Rapid mood changes, confusion, false beliefs, hallucinations, or disorganized behavior can develop quickly and may threaten the safety of the parent or baby, requiring immediate assessment.

### **Puerperal Sepsis**

A serious infection of the reproductive tract during labor or the six weeks after childbirth. Fever, pelvic pain, foul-smelling discharge, weakness, or fast breathing require urgent medical assessment and antibiotic treatment.

#### **Alternate Names:**

Postpartum Sepsis; Puerperal Septicemia

### **Puerperium**

The period after childbirth, usually about six weeks, when the body recovers from pregnancy and birth. The uterus shrinks, bleeding settles, milk production develops, and health problems may still arise.

### **Puffy Eyes**

Alternate name; see *Bags under eyes*.

### **Pugilistic Stance**

A flexed posture of the arms, hands, and legs sometimes seen in bodies exposed to intense heat. It results from heat-related muscle and tendon con-

traction, not from purposeful movement, and does not by itself show when or how death occurred.

### **Pulling A Patient Up In Bed**

A safe repositioning technique using adequate staff, friction-reducing equipment, and coordinated movement to move a patient toward the head of the bed while preventing falls, shear, and staff injury.

### **Pull Test**

A standardized clinical test used to evaluate postural stability and balance recovery. The examiner stands behind the patient and delivers a brisk, backward tug to the patient's shoulders; an inability to recover balance within one or two steps indicates impaired postural reflexes in Parkinson's disease or atypical parkinsonism.

### **Pulmonary**

Relating to the lungs or the blood vessels and airways that support breathing. Pulmonary disease may involve infection, asthma, chronic obstructive lung disease, high pressure in lung vessels, blood clots, fluid, scarring, or cancer.

### **Pulmonary Acinus**

The smallest part of the lung that exchanges gases. It begins after a small airway and contains respiratory bronchioles, alveolar ducts, and alveoli where oxygen enters blood and carbon dioxide leaves it.

### **Pulmonary Actinomycosis**

Pulmonary actinomycosis is a chronic bacterial infection of the lungs caused by *Actinomyces* and may form mass-like lesions or spread across tissue planes.

### **Pulmonary Alveolar Hemorrhage**

Bleeding into the lung's air sacs. It may cause coughing up blood, breathlessness, low oxygen, anemia, and widespread lung shadows and can result from immune disease, infection, medicines, injury, or a clotting problem.

### **Pulmonary Alveolar Microlithiasis**

A rare inherited lung disease in which tiny mineral

---

deposits build up inside the air sacs. It may cause no symptoms at first, but later can lead to cough, breathlessness, low oxygen, and progressive lung scarring.

### **Pulmonary Alveolar Proteinosis**

A rare lung disease in which a fatty material builds up inside the air sacs and interferes with oxygen exchange. It can cause gradually worsening breathlessness, cough, tiredness, and low oxygen and may be immune-related or secondary to another illness.

### **Pulmonary Angiography**

Contrast imaging of the pulmonary arteries, usually by catheter. It can diagnose pulmonary vascular abnormalities but is now used less often for pulmonary embolism because CT pulmonary angiography is less invasive.

### **Pulmonary Arterial Hypertension**

Group 1 pulmonary hypertension caused by increased resistance in the small pulmonary arteries, which become narrowed, remodeled, or obstructed. It may be idiopathic, heritable, drug- or toxin-induced, or associated with connective-tissue disease, HIV, portal hypertension, congenital heart disease, or schistosomiasis. Symptoms include exertional breathlessness, fatigue, chest discomfort, dizziness, fainting, or swelling; diagnosis requires right-heart catheterization.

### **Pulmonary Arteries**

The two large blood vessels carrying oxygen-poor blood from the right side of the heart to the lungs. In the lungs, the blood releases carbon dioxide and picks up oxygen.

#### **Alternate Names:**

Main Pulmonary Trunk; Right and Left Pulmonary Arteries

### **Pulmonary Arteriovenous Fistula**

An abnormal direct connection between an artery and vein in the lung that lets blood bypass normal oxygen exchange. It can cause low oxygen, breathlessness, blue skin, stroke-causing clots, or rarely a brain abscess.

### **Pulmonary Arteriovenous Malformation**

An abnormal direct connection between an artery and vein in the lung. Blood bypasses normal oxygen exchange, which can cause low oxygen, breathlessness, coughing blood, stroke, or a brain abscess; some cases are linked to an inherited blood-vessel disorder.

### **Pulmonary Artery**

The large blood vessel carrying oxygen-poor blood from the right side of the heart to the lungs. There, blood releases carbon dioxide, receives oxygen, and returns through pulmonary veins.

### **Pulmonary Artery Catheter**

A thin tube passed through the right side of the heart into a lung artery to measure pressures, blood flow, and heart performance. It is used selectively in severe heart or lung disease and critical illness because insertion has risks.

### **Pulmonary-Artery Dissection**

A rare, life-threatening tear in the wall of an artery carrying blood from the heart to the lungs. It may cause sudden chest pain, breathlessness, low blood pressure, or collapse and requires emergency specialist care.

### **Pulmonary Artery Pulsatility Index**

A measurement calculated during right-heart catheterization from pulmonary-artery pressure and right-atrial pressure. A low result may indicate weak right-ventricle pumping and a higher risk of right-sided heart failure in seriously ill patients.

#### **Alternate Names:**

PAPi; PAP Index

### **Pulmonary Artery Wedge Pressure**

An indirect hemodynamic measure of left atrial pressure and left ventricular end-diastolic filling pressure, obtained by inflating a balloon at the tip of a pulmonary artery catheter (Swan-Ganz) until it occludes a small pulmonary arterial branch.

#### **Alternate Names:**

PAWP; PCWP; Pulmonary Capillary Wedge Pressure

---

## **Pulmonary Aspergilloma**

A pulmonary aspergilloma is a ball of *Aspergillus* fungus and debris growing in a pre-existing lung cavity, sometimes causing coughing of blood.

## **Pulmonary Atresia**

A heart defect present from birth in which the valve or passage from the right ventricle to the pulmonary artery does not open normally. Blood cannot reach the lungs through the usual route, so newborns may become blue or severely unwell and need urgent specialist care.

### **Alternate Names:**

Pulmonic Valve Atresia; PA-IVS; PA-VSD

## **Pulmonary Blastoma**

A rare malignant lung tumor composed of primitive epithelial and mesenchymal tissue, usually treated with specialist oncology care.

## **Pulmonary Capillary Wedge Pressure**

A pressure measured with a special catheter in a small lung artery to estimate pressure on the left side of the heart. It can help assess heart failure or fluid overload in critically ill patients, but breathing support and other conditions affect its meaning.

## **Pulmonary Circuit**

The circulation in which blood travels from the right ventricle to the lungs for gas exchange and returns to the left atrium.

## **Pulmonary Circulation**

The blood-flow circuit from the right ventricle through the pulmonary arteries and lung capillaries to the left atrium, where blood releases carbon dioxide and gains oxygen.

## **Pulmonary Compliance**

How easily the lungs expand when pressure changes during breathing. Low compliance means the lungs are stiff and harder to inflate; high compliance can mean they expand easily but may not recoil effectively.

## **Pulmonary Contusion**

Bruising of the lung after a blow or crush injury to the chest. Bleeding and swelling inside the lung can reduce oxygen exchange and worsen during the first day or two; treatment supports breathing and addresses associated injuries.

## **Pulmonary Disease and Pregnancy**

Lung disease can worsen during pregnancy or be affected by pregnancy-related changes in oxygen demand and ventilation. Asthma, pneumonia, pulmonary embolism, and chronic lung disease require prompt assessment when breathlessness is severe, new, or associated with chest pain or low oxygen.

## **Pulmonary Edema**

A buildup of fluid in the lungs that can make breathing difficult. It is often caused by heart failure but may also result from kidney problems, lung injury, medicines, or high altitude. Sudden severe breathing difficulty is an emergency.

### **Alternate Names:**

Fluid in the Lungs; Pulmonary Congestion; Alveolar Flooding

## **Pulmonary Embolism**

A blockage of a pulmonary artery, usually caused by a blood clot that traveled from a deep vein. It may cause sudden breathlessness, chest pain that worsens with breathing, rapid heartbeat, coughing blood, dizziness, or fainting. A large embolism can severely strain the heart or cause cardiac arrest and requires emergency assessment.

### **Alternate Names:**

PE; Pulmonary Thromboembolism; Blood Clot in Lungs

## **Pulmonary Embolism Treatment**

Treatment for a blood clot in the lung usually uses anticoagulant medicine to stop growth and prevent new clots. Severe or unstable cases may need clot-dissolving medicine or a procedure; choice and duration depend on the person's risks and cause.

## **Pulmonary Embolus**

A blockage of a pulmonary artery, usually by a



---

blood clot that traveled from a deep vein, causing sudden breathlessness, chest pain, low oxygen, or collapse.

### **Pulmonary Emphysema**

Alternate name; see *Emphysema*.

### **Pulmonary Eosinophilia**

A group of disorders in which eosinophils accumulate in lung tissue or air spaces, causing infiltrates, cough, wheezing, breathlessness, or low oxygen. Causes include parasites, medicines, asthma, allergic disease, and eosinophilic lung syndromes; treatment targets the cause and inflammation.

### **Pulmonary Fibrosis**

Scarring and stiffening of lung tissue, often as part of an interstitial lung disease. The scarring can make it harder for the lungs to expand and for oxygen to pass into the blood. It may cause worsening shortness of breath, a dry cough, and tiredness. Causes include some autoimmune diseases, medicines, radiation, and inhaled dusts or chemicals; sometimes no cause is found. Idiopathic pulmonary fibrosis is a specific type with no known cause.

### **Pulmonary Fistula**

An abnormal passage between the lung or an airway and another structure, such as the space around the lung, skin, or swallowing tube. It can allow air or infected fluid to travel where it should not.

### **Pulmonary Function Tests**

Breathing tests that measure how much air the lungs hold, how quickly air can be moved, and how well oxygen passes into the blood. They help assess asthma, chronic obstructive lung disease, scarring, and other breathing problems. Results depend on effort, technique, age, and body size.

### **Pulmonary Hemorrhage**

Bleeding into the airways or lung tissue, which may cause coughing blood, low oxygen, anemia, or respiratory failure.

### **Pulmonary Hemosiderosis**

A condition in which repeated bleeding into the lung's air sacs leaves iron deposits behind. It may cause cough, breathlessness, coughing blood, anemia, and low oxygen and can result from immune disease or other causes.

### **Pulmonary Hypertension**

Abnormally high blood pressure in the blood vessels that carry blood from the right side of the heart to the lungs. Narrowed, blocked, or damaged vessels make the right side of the heart work harder to pump blood and may eventually weaken it. Pulmonary hypertension may occur on its own or result from left-sided heart disease, lung disease or low oxygen, long-term blood clots, or other conditions affecting the blood vessels. Symptoms may include breathlessness, tiredness, chest discomfort, dizziness, fainting, or swelling from heart strain.

#### **Alternate Names:**

PH; Pulmonary Arterial Hypertension; High Blood Pressure in Lungs

### **Pulmonary Hypertension Classification**

A clinical grouping based on cause: disease of the pulmonary arteries, left-sided heart disease, lung disease or low oxygen, chronic blood clots, or several other mechanisms. Identifying the group guides testing and treatment because medicines useful for one group may be harmful in another.

### **Pulmonary Hypoplasia**

Underdevelopment of one or both lungs, resulting in too little lung tissue for normal breathing. It may occur with congenital abnormalities, low amniotic fluid, or compression before birth.

### **Pulmonary Infarction**

Death of a small area of lung tissue after its blood supply is blocked, usually by a pulmonary embolism. It may cause sharp pain with breathing, coughing up blood, breathlessness, or a shadow on imaging.

### **Pulmonary Infection**

An infection of the lungs or lower respiratory tract caused by bacteria, viruses, fungi, or other organ-

---

isms.

### **Pulmonary Langerhans Cell Histiocytosis**

A rare lung disease strongly linked to cigarette smoking, in which unusual immune cells damage small airways and form lung cysts. It can cause cough, breathlessness, chest pain, or a collapsed lung. Stopping smoking is essential and may stabilize the disease; specialist treatment is needed if it progresses.

#### **Alternate Names:**

PLCH; Histiocytosis X of the Lung; Pulmonary Eosinophilic Granuloma

### **Pulmonary Lobes**

The anatomical divisions of the lungs separated by interlobar fissures. The right lung has superior, middle, and inferior lobes; the left lung has superior and inferior lobes and a lingula.

### **Pulmonary Mass**

A space-occupying lesion in or near the lung, which may represent infection, inflammation, benign growth, or cancer and requires appropriate imaging assessment.

### **Pulmonary Nocardiosis**

A lung infection caused by *Nocardia* bacteria, usually affecting people with weakened immunity and sometimes spreading to the brain or skin.

### **Pulmonary Nodule**

A rounded opacity in the lung, usually smaller than 3 cm, caused by scar, infection, inflammation, or tumor and assessed according to size and risk factors.

### **Pulmonary Oedema**

Fluid collecting in lung tissue or air sacs, making breathing difficult and reducing oxygen transfer. It often results from heart failure but can also follow infection, toxins, kidney disease, or severe injury.

### **Pulmonary Reactance**

A measure of how the stretch and movement of the lungs and chest wall affect airflow during breathing. It is obtained during specialized tests and can

help describe stiffness or altered movement of the respiratory system.

### **Pulmonary Regurgitation (Pulmonic Regurgitation)**

Backward flow from the pulmonary artery into the right ventricle during diastole because the pulmonary valve does not close completely. Mild regurgitation may be asymptomatic, whereas severe or long-standing regurgitation can enlarge or weaken the right ventricle and cause exercise intolerance, arrhythmias, or right-sided heart failure.

### **Pulmonary Rehabilitation**

A supervised, multidisciplinary program combining exercise training, education, behavioral support, and self-management for chronic respiratory disease. It improves exercise capacity, dyspnea, functional status, and quality of life.

### **Pulmonary-Renal Syndrome**

A group of diseases that cause inflammation or bleeding in the lungs together with inflammation or injury in the kidneys. It is often associated with small-vessel vasculitis or another autoimmune disease.

### **Pulmonary Sequestration**

A birth-related area of lung tissue that does not connect normally to the airways and receives blood from an unusual artery. It may cause repeated lung infections, cough, or breathing problems, although some cases cause no symptoms.

### **Pulmonary Stenosis**

Narrowing that obstructs blood leaving the right ventricle on its way to the lungs, usually at the pulmonary valve. It is often congenital and increases right-ventricular pressure; severe disease may cause a murmur, fatigue, breathlessness, fainting, or right-heart failure.

### **Pulmonary Surfactant**

A slippery mixture made by cells lining the lung's air sacs that reduces surface tension and helps keep the sacs open. Too little in premature babies can

---

cause serious breathing difficulty.

### **Pulmonary Thromboendarterectomy**

Major surgery to remove old clots and scar tissue from the arteries of the lungs. It may improve or cure long-term clot-related high blood pressure in the lungs in selected people, but it carries serious surgical risks.

#### **Alternate Names:**

PTE; Pulmonary Endarterectomy; PEA

### **Pulmonary Trunk**

The large vessel leaving the right ventricle that divides into the right and left pulmonary arteries.

### **Pulmonary Tuberculosis**

Tuberculosis infection involving the lungs, caused by *Mycobacterium tuberculosis*. It may cause prolonged cough, fever, night sweats, weight loss, or coughing blood and can spread through the air.

### **Pulmonary Valve**

A one-way valve between the right lower heart chamber and the artery leading to the lungs. It opens to let blood leave the heart and closes to prevent backward flow; narrowing or leakage can strain the right side of the heart.

### **Pulmonary Valve Insufficiency**

Leakage of blood backward through the pulmonary valve into the right lower heart chamber while the heart relaxes. Mild leakage may cause no symptoms, but severe disease can enlarge the heart and reduce exercise ability.

### **Pulmonary Valve Regurgitation**

Leakage through the pulmonary valve that allows blood to flow back into the right lower heart chamber as the heart relaxes. Mild leakage may cause no symptoms; severe leakage can enlarge and weaken the right side of the heart.

#### **Alternate Names:**

Pulmonic Regurgitation; Pulmonary Incompetence; PR

### **Pulmonary Valve Stenosis**

Alternate name; see *Pulmonary valve stenosis*.

### **Pulmonary Vascular Resistance**

The opposition to blood flow through the lung blood vessels. High resistance makes the right side of the heart pump harder and can lead to high pressure in the lungs, right-heart enlargement, and heart failure.

### **Pulmonary Vasodilator Therapy**

Treatment with medicines that widen blood vessels in the lungs and lower pressure on the right side of the heart. It is used for selected types of pulmonary hypertension after specialist testing because it may be harmful in other types.

### **Pulmonary Vein**

The four veins (two from each lung) returning oxygenated blood from the lungs to the left atrium, a unique exception in the venous system (carrying oxygenated blood). Superior and inferior pulmonary veins enter the left atrium posteriorly.

### **Pulmonary Vein Isolation**

A catheter procedure for selected people with atrial fibrillation. The clinician creates small areas of scar around the veins entering the upper heart chamber to block abnormal electrical signals. It can reduce episodes but may not cure them; bleeding, stroke, narrowing of a vein, or other complications are possible.

#### **Alternate Names:**

PVI; Antral Pulmonary Vein Isolation; AFib Catheter Ablation

### **Pulmonary Veins**

Usually four veins that carry oxygen-rich blood from the lungs to the heart's left upper chamber. They are unusual veins because they carry oxygen-rich rather than oxygen-poor blood.

### **Pulmonary Veno-Occlusive Disease**

A rare progressive disease in which small veins in the lungs become narrowed or blocked. Pressure

---

then rises in the lung circulation, causing breathlessness and low oxygen; some medicines for pulmonary hypertension can trigger life-threatening fluid in the lungs.

### **Pulmonary Ventilation**

The movement of air into and out of the lungs. It brings oxygen into the air sacs and removes carbon dioxide; inadequate ventilation can cause low oxygen or a buildup of carbon dioxide.

### **Pulmonary Ventilation/Perfusion Scan**

An imaging test comparing air reaching different lung regions with blood flow through those regions. A mismatch may suggest a blood clot in the lung, while other patterns can indicate lung or blood-vessel disease.

### **Pulmonary Wedge Pressure**

A pressure measured through a special catheter in a small lung artery to estimate the pressure in the left side of the heart. It helps assess fluid overload or heart failure in seriously ill patients.

### **Pulmonic Valve Stenosis**

Narrowing of the valve between the right ventricle and pulmonary artery, usually present from birth, that creates right-ventricular outflow obstruction. Echocardiography assesses severity; selected symptomatic or severe cases can be treated with catheter balloon valvuloplasty or surgery.

### **Pulmonologist**

A pulmonologist is a physician specializing in diseases of the lungs and breathing system, including asthma, COPD, infections, and sleep-related breathing disorders.

### **Pulmonology**

The medical specialty concerned with the lungs and breathing system. Pulmonologists diagnose and treat asthma, chronic lung disease, infections, sleep-related breathing disorders, and other respiratory conditions.

### **Pulp**

The soft living tissue in the center of a tooth, containing nerves and blood vessels. It can become inflamed or infected after decay, cracks, or injury and may then require root-canal treatment or removal.

### **Pulp Canal**

The pulp canal is the passage within a tooth root containing dental pulp, nerves, and blood vessels; infection may require root-canal treatment.

### **Pulp Disease**

A problem affecting the living tissue inside a tooth, including inflammation, infection, loss of blood supply, or tissue death. It may cause toothache or sensitivity and can follow decay, a crack, a filling, or injury.

### **Pulpectomy**

Complete removal of infected or damaged pulp from inside a tooth, usually a primary tooth. The space is cleaned and filled with a material that can safely resorb as the adult tooth develops.

### **Pulpitis**

Inflammation of the dental pulp, usually caused by caries, trauma, cracks, or restorative treatment. Reversible pulpitis causes brief pain provoked by cold or sweet stimuli; irreversible pulpitis causes spontaneous or lingering pain and may progress to pulp necrosis, requiring endodontic treatment or extraction.

### **Pulposus**

Relating to the nucleus pulposus, the gelatinous central portion of an intervertebral disk. Relating to the soft, jelly-like center of an intervertebral disk, which helps cushion spinal movement.

### **Pulpotomy**

Removal of inflamed soft tissue from the crown of a baby tooth while keeping the root tissue alive. It treats suitable decay or injury and preserves the tooth until it naturally falls out.

---

## **Pulp Stone**

A small area of hardened mineral inside the soft center of a tooth. It often causes no symptoms and is found on dental imaging, but it may complicate root-canal treatment if it blocks the canal.

## **Pulsatile Proptosis**

An abnormal forward protrusion of the eyeball characterized by visible or palpable pulsations synchronous with the arterial pulse, classically accompanied by an audible orbital bruit. It is a cardinal sign of a direct carotid-cavernous fistula or congenital sphenoid wing dysplasia in neurofibromatosis type 1.

## **Pulsatile Tinnitus**

A rhythmical acoustic sound perceived in one or both ears that beats synchronously with the patient's heartbeat, frequently signifying an underlying vascular anomaly such as dural arteriovenous fistula, internal carotid artery dissection, carotid stenosis, glomus jugulare tumor, or idiopathic intracranial hypertension.

## **Pulse**

A pulse is the rhythmic expansion of an artery produced by each heartbeat; its rate, rhythm, strength, and symmetry provide clues about circulation and cardiac function.

## **Pulse Examination**

Checking arterial pulses for speed, regularity, strength, and equality on both sides. It helps assess blood flow, heart rhythm, circulation to limbs, and possible blockage or reduced blood supply.

## **Pulse Irregular**

An irregular pulse has uneven timing or varying beats and may reflect premature beats, atrial fibrillation, other arrhythmias, or measurement variation.

## **Pulseless Disease**

Takayasu arteritis, an inflammation of large arteries that can narrow or block blood flow. It may cause weak pulses, unequal blood pressure, limb pain,

dizziness, vision problems, or organ damage.

## **Pulseless Electrical Activity**

Cardiac arrest in which an organized electrical rhythm appears on the ECG but there is no palpable pulse or effective mechanical cardiac output. It is a non-shockable rhythm requiring high-quality CPR, epinephrine, and rapid evaluation for reversible causes, including the Hs and Ts.

## **Pulse Oximeter**

A small device, often placed on a fingertip, that estimates blood oxygen and pulse rate using light. Movement, cold fingers, poor circulation, nail products, and device limitations can make the reading inaccurate.

## **Pulse Oximetry**

A painless method of estimating blood oxygen with a light-based sensor, usually on a finger, toe, or ear. The result must be interpreted with symptoms and circulation because it can be inaccurate in some situations.

## **Pulse Pressure**

The systolic blood pressure minus the diastolic blood pressure. A widened or narrowed value may provide clues about circulation but must be interpreted with the full blood-pressure reading and clinical context.

## **Pulse Rate**

Pulse rate is the number of palpable arterial beats per minute and usually approximates heart rate, although some arrhythmias produce a pulse deficit.

## **Pulsus**

A term used in medicine to describe the pulse and named patterns of arterial beats. The pattern may provide clues about heart rhythm, pumping strength, blood-vessel resistance, breathing, or circulation.

## **Pulsus Alternans**

A pulse rhythm where strong and weak beats alternate back and forth, showing that the main heart



---

pump is weak and struggling.

### **Pulsus Bisferiens**

An abnormal peripheral arterial pulse waveform characterized by two distinct systolic peaks per cardiac cycle separated by a midsystolic dip, classic for severe aortic regurgitation, combined aortic stenosis and regurgitation, or hypertrophic obstructive cardiomyopathy (HOCM).

#### **Alternate Names:**

Biphasic Pulse; Bisferious Pulse

### **Pulsus Paradoxus**

An unusually large drop in blood pressure during breathing in, usually more than 10 millimeters of mercury. It can occur when the heart is compressed by fluid, or during severe asthma, chronic lung disease, or other heart conditions.

### **Pulsus Parvus et Tardus**

A weak, slow-rising pulse in the neck. It may occur with severe narrowing of the aortic valve, but it is not specific and must be assessed with the heart examination and tests.

#### **Alternate Names:**

Anacrotic Pulse; Slow-Rising Arterial Pulse

### **Pulvinar**

A region at the back of the thalamus, a deep brain structure. It helps direct visual attention, combine sensory information, and communicate with areas of the cerebral cortex.

### **Pumice**

A porous volcanic stone used as an abrasive; excessive dust inhalation can irritate the lungs.

### **Pump Failure**

Severe weakening of the heart's pumping ability, causing inadequate blood flow to organs.

#### **Alternate Names:**

Cardiogenic Shock; Acute Ventricular Failure

### **Pump-Oxygenator**

The part of a heart-lung machine that moves blood

and adds oxygen while removing carbon dioxide during some heart operations. It temporarily replaces the pumping and gas-exchange work of the heart and lungs. Use can be associated with bleeding, inflammation, clots, or organ complications.

### **Punch Biopsy**

A dermatological procedure in which a circular, hollow blade instrument is rotated through the skin into the subcutaneous fat to obtain a full-thickness cylindrical specimen of skin tissue for histopathological examination.

#### **Alternate Names:**

Punch Biopsy

### **Punctal Occlusion**

Partial or complete blocking of the small tear-drain openings near the inner eyelids, often with plugs or a procedure, to keep tears on the eye surface. It may help selected people with dry eye; watering, irritation, or infection can occur.

### **Punctate**

Made up of, or marked by, small spots, dots, or points. The term describes appearance and does not by itself identify a diagnosis.

### **Punctate Palmoplantar Keratoderma**

An inherited condition causing many small, hard areas of thickened skin on the palms and soles. It often begins in adolescence or adulthood and can make walking or using the hands painful; rarely, a skin cancer develops.

### **Puncture**

A small hole or wound made by a pointed object, needle, bite, or sharp instrument. A puncture may be intentional for an injection or sample, or accidental and may carry infection, bleeding, or deeper injury risks.

### **Puncture Biopsy**

Alternate name; see *needle biopsy*: removal of cells or tissue through a percutaneous puncture for microscopic, molecular, or microbiologic examination; imaging may guide the target and route.



---

## Puncture Wound

A wound produced when a pointed object penetrates the skin and underlying tissue, creating a small surface opening with potentially deep tissue damage and infection risk.

## Pungent

Describes a strong, sharp smell or taste that may irritate the nose, mouth, eyes, or other moist linings.

## Punicic Acid

A conjugated polyunsaturated fatty acid found mainly in pomegranate seed oil and studied for lipid and inflammatory effects; clinical supplement benefits remain uncertain.

## Punishment

Punishment is a consequence intended to reduce a behavior; in healthcare, coercive or harmful punishment is inappropriate and behavior support should prioritize safety and dignity.

## Pupil

The central opening in the colored iris through which light enters the eye. Muscles change its size to control incoming light, and unequal or poorly reactive pupils may indicate eye or nerve disease.

## Pupillary

Relating to the pupil, the opening in the iris that controls how much light enters the eye. The pupil normally becomes smaller in bright light and larger in dim light; unequal size or poor reaction can indicate an eye, nerve, or brain problem.

## Pupillary Light Reflex

The normal narrowing of the pupil when light enters the eye. It tests the eye, optic nerve, brain pathways, and nerves that control the pupil.

## Pure Red Cell Aplasia

A rare disorder in which the bone marrow stops making red blood cells while other blood cells may remain normal. It causes severe anemia, tiredness, breathlessness, and pale skin and may be inherited

or acquired from infection, immune disease, cancer, or medicines.

### Alternate Names:

PRCA; Erythroblastopenia

## Pure Tone Audiometry

A hearing test in which a person signals whenever they hear tones of different pitches and loudness. It measures the quietest sounds detectable through air and bone and helps identify the type and degree of hearing loss.

## Pure-Tone Audiometry

A hearing test that measures the softest tones a person can hear at selected frequencies through air and bone conduction. The results are plotted on an audiogram and help distinguish conductive from sensorineural hearing loss.

### Alternate Names:

Pure-Tone Threshold Audiometry; Pure-Tone Hearing Test

## Purgative

A strong laxative that causes rapid emptying of the bowel. It may be used for bowel preparation before some procedures, but excessive use can cause diarrhea, dehydration, and dangerous changes in body salts.

## Purge

To empty the bowel, stomach, or body of a substance; the term may also mean forceful vomiting or diarrhea.

## Purging

Deliberate behaviors intended to compensate for eating, such as self-induced vomiting or misuse of laxatives, diuretics, or excessive exercise.

## Purging Disorder

Repeated self-induced vomiting or misuse of laxatives, diuretics, or other methods to influence weight or body shape without recurrent binge eating. It can cause dangerous salt disturbances, tooth damage, dehydration, heart-rhythm problems, and tears

---

in the food pipe.

### **Purine**

A nitrogen-containing building block of DNA, RNA, and energy-carrying molecules. The body breaks down purines into uric acid, which can accumulate and contribute to gout or kidney stones.

### **Purine Nucleosides**

Molecules made from a purine base, such as adenine or guanine, joined to a sugar. Examples include adenosine and guanosine, which help form genetic material and participate in cell energy and signaling.

### **Purine Nucleotides**

Molecules containing adenine or guanine joined to a sugar and phosphate groups. They build DNA and RNA and include ATP and GTP, which store energy and help transmit signals inside cells.

### **Purines**

Nitrogen-containing building blocks, mainly adenine and guanine, used to make DNA, RNA, and energy-carrying molecules. The body breaks them down into uric acid, which can form crystals and contribute to gout or kidney stones.

### **Purinoceptor**

A protein on a cell's surface that detects substances such as ATP, ADP, or adenosine outside the cell. It helps regulate cell communication, inflammation, blood-vessel activity, and other body responses.

### **Purkinje Cell**

A major nerve cell in the cerebellum, the brain region that coordinates movement and balance. It sends signals that help fine-tune posture, coordination, and learned movements.

### **Purkinje Fibers**

Special electrical fibers spread through the lower chambers of the heart. They rapidly carry each heartbeat's signal so the ventricles contract together and pump blood efficiently.

### **Purkinje Fibres**

Specialized heart-muscle fibers that rapidly spread electrical signals through the ventricles. Their coordinated activity helps the lower chambers contract together and pump blood efficiently to the lungs and body.

### **Purpura**

Non-blanching red, purple, or violaceous macules, papules, or patches produced by the extravasation of erythrocytes into the dermis or submucosa. It is classified by lesion size into petechiae (less than three millimetres), purpura (three to ten millimetres), and ecchymoses (greater than one centimetre), and divided clinically into palpable purpura (indicating small-vessel vasculitis) and non-palpable purpura (caused by thrombocytopenia, platelet dysfunction, vascular fragility, or coagulation disorders).

### **Purpura Fulminans**

A rapidly worsening, life-threatening condition in which widespread blood clots and clotting-system failure cause purple skin patches, bleeding, and tissue death. It can follow severe infection, loss of natural anticoagulants, or another serious illness.

### **Purulent**

Containing or producing pus, a thick fluid made of immune cells, dead tissue, and often germs. Purulent drainage can occur in an abscess or infected wound, but its appearance alone does not identify the organism or the treatment needed.

### **Purulent Pericarditis**

A serious bacterial infection of the sac around the heart, causing pus and fluid to collect there. Fever, chest pain, breathlessness, and pressure on the heart may occur; urgent drainage and antibiotic treatment are needed.

### **Pus**

A thick fluid made of dead cells, germs, immune cells, and damaged tissue. It commonly collects during infection and may appear in wounds, abscesses, infected organs, or other inflamed body spaces.

---

### **Push Enteroscopy**

An examination of the upper small intestine using a long flexible camera passed through the mouth. It allows clinicians to see the lining, take samples, stop bleeding, or treat selected lesions.

### **Pustular Psoriasis**

A form of psoriasis characterized by sterile pus-filled pustules on inflamed skin. It may be localized to the palms and soles or generalized with fever and systemic illness; generalized disease can be an emergency requiring urgent specialist care.

### **Pustule**

A small raised skin lesion filled with pus. Pustules can occur with acne, follicle infection, impetigo, psoriasis, or a medicine reaction; their cause is determined from their location, appearance, symptoms, and timing.

### **Pustules**

Pustules are small raised skin lesions filled with pus, occurring in acne, folliculitis, infections, inflammatory disorders, or drug reactions.

### **Pustulosis**

A skin disorder characterized by sterile or infectious pustules, which may be localized or widespread and associated with inflammation, fever, psoriasis, drug reactions, or infection.

### **Putamen**

A deep brain structure that helps select and smooth purposeful movements and supports learning of repeated movement habits. Damage to it can cause abnormal movements, weakness, or changes in behavior.

### **Putrefaction**

Decomposition of body tissues after death caused mainly by bacteria and natural enzymes. It can produce color changes, swollen tissues from gas, visible veins, a strong odor, and gradual breakdown into liquid material; the rate depends on the environment.

### **Putrescine**

A chemical with a strong unpleasant smell made when proteins break down. It is found in decaying material and is also produced in small amounts during normal cell processes; its presence alone does not diagnose disease.

### **PUVA**

A light treatment that combines a light-sensitizing medicine called psoralen with ultraviolet-A exposure. It can treat selected skin diseases but may cause burns, eye injury, premature skin aging, and increased skin-cancer risk.

### **PVC**

A PVC, or premature ventricular contraction, is an early heartbeat arising in a ventricle; occasional PVCs are common, while frequent or symptomatic beats need evaluation.

### **PVD**

Alternate name; see *Peripheral artery disease*.

### **PVI**

Alternate name; see *Pulmonary Vein Isolation*.

### **PVPS**

Alternate name; see *Post-vasectomy pain syndrome*.

### **P Wave**

The first small wave on an ECG, showing the electrical activation of the heart's upper chambers. Its shape and timing can help identify abnormal heart rhythms, enlargement of a chamber, or problems with electrical conduction.

### **Pyarthrosis**

Infection of a joint with pus in the joint space, usually bacterial septic arthritis, causing acute pain, swelling, restricted movement, fever, and risk of cartilage destruction.

### **Pycnodysostosis**

A rare inherited bone disorder in which bones are unusually dense but fragile. It can cause short

---

stature, frequent fractures, short fingers, delayed closure of skull bones, crowded teeth, and problems with the jaw.

### **Pyelitis**

Inflammation of the renal pelvis, the funnel-shaped part of the kidney that drains urine. It is often associated with urinary infection and may cause flank pain, fever, or painful urination.

### **Pyelo**

Pyelo is an informal shortening usually referring to pyelonephritis, an infection of the kidney and renal pelvis that can cause fever and flank pain.

### **Pyelogram**

An image of the kidney drainage area and ureters after contrast material is given. It can show blockage, stones, narrowing, leaks, or structural changes in the urinary collecting system.

### **Pyelography**

Imaging of the renal pelvis and ureters using contrast, by X-ray or another imaging method. It can show obstruction, stones, leaks, strictures, or structural abnormalities.

### **Pyelolithotomy**

Surgical removal of a kidney stone through an incision into the renal pelvis. Less invasive stone treatments are often preferred when suitable.

### **Pyelonephritis**

Infection and inflammation of the kidney and its urine-collecting area, usually caused by bacteria traveling upward from the bladder. Symptoms may include fever, chills, side pain, nausea, and painful urination.

### **Pyelotomy**

An incision into the renal pelvis, usually made to remove an obstruction or kidney stone.

### **Pylephlebitis**

An infected blood clot in the large vein carrying

blood from the abdominal organs to the liver. It may follow appendicitis, diverticulitis, or another abdominal infection and can cause fever, abdominal pain, jaundice, and life-threatening sepsis.

### **Pyloric Antrum**

The pyloric antrum is the lower stomach region that grinds food and regulates its passage toward the pyloric canal and duodenum.

### **Pyloric Canal**

The narrow passage between the pyloric antrum and the pyloric sphincter. It carries stomach contents toward the duodenum.

### **Pyloric Sphincter**

A ring of muscle at the outlet of the stomach. It opens in a controlled way to let partly digested food enter the first part of the small intestine and helps prevent the intestine's contents from flowing backward.

#### **Alternate Names:**

Pyloric Valve; Gastric Pylorus

### **Pyloric Stenosis**

Narrowing of the outlet between the stomach and small intestine. In infants, the usual form is thickening of the outlet muscle, causing progressively forceful, non-green vomiting. In adults, narrowing is usually acquired, for example from ulcer scarring or a tumor. It can prevent food from leaving the stomach and cause dehydration.

### **Pyloric Stenosis In Infants**

Infantile pyloric stenosis is narrowing of the stomach outlet that causes forceful, non-bilious vomiting in young infants and usually requires surgical treatment.

### **Pyloroplasty**

Surgery to widen the pylorus, the outlet of the stomach, to improve passage of stomach contents. It may be used with treatment for obstruction or selected gastric-motility disorders.

---

## **Pylorus**

The distal part of the stomach leading into the duodenum.

## **Pyoderma**

A bacterial infection of the skin that can produce pus-filled spots, crusts, open sores, or deeper inflammation. Staphylococcus and Streptococcus are common causes; the infection may spread and require medical treatment, especially with fever, pain, or rapidly worsening redness.

## **Pyoderma Gangrenosum**

A rare inflammatory skin disease that causes very painful ulcers, often with a dark purple edge. It may occur with bowel inflammation, arthritis, or blood disorders; injury can make it worse, so treatment differs from ordinary wound infection.

## **Pyogenesis**

The formation of pus, a thick fluid made mainly of immune cells, dead tissue, and germs. It commonly occurs during bacterial infection or severe inflammation.

## **Pyogenic**

Producing pus or associated with a pus-forming infection. Pyogenic infections commonly cause redness, warmth, swelling, pain, and a collection of pus, although the cause and treatment vary.

## **Pyogenic Granuloma**

A common benign vascular proliferation (lobular capillary hemangioma) of the skin or mucous membranes that arises rapidly following minor trauma or during pregnancy ('pregnancy tumor' / granuloma gravidarum), presenting as an erythematous, smooth or lobulated, pedunculated or sessile vascular papule prone to profuse bleeding.

### **Alternate Names:**

Lobular Capillary Hemangioma; Granuloma Gravidarum

## **Pyogenic Granuloma (Lobular Capillary**

## **Hemangioma)**

A rapidly growing, benign vascular nodule of skin or mucosa that often follows minor trauma or occurs during pregnancy. It is bright red, friable, and prone to bleeding; treatment may include curettage, cautery, excision, or other local therapy.

## **Pyogenic Liver Abscess**

A pyogenic liver abscess is a pus-filled infection in the liver, usually from biliary, abdominal, or blood-stream spread, causing fever, pain, and possible sepsis.

## **Pyometra**

A collection of pus inside the uterus, usually caused by infection combined with blockage of drainage through the cervix. It may cause lower abdominal pain, fever, tenderness, or foul-smelling discharge; in older adults it can be linked to cervical narrowing or a tumor and needs prompt evaluation.

## **Pyonephrosis**

Pus trapped in the kidney's urine-collecting system, usually because infection occurs behind a blockage. It is a medical emergency requiring urgent drainage, antibiotics, and treatment of the obstruction.

## **Pyosalpinx**

A fallopian tube filled with pus, usually because of pelvic infection. It may cause lower abdominal pain, fever, unusual discharge, infertility, or an abscess and generally requires prompt treatment.

## **Pyostomatitis Vegetans**

A rare inflammatory mouth condition causing many small yellowish pus-filled spots on red, swollen tissue, especially the inner cheeks and gums. It is strongly associated with inflammatory bowel disease and may appear before bowel symptoms.

## **Pyramid**

A structure shaped like a pyramid. In the brain, the medullary pyramids contain nerve fibers that carry movement commands from the brain to the spinal cord; other pyramids include pyramid-shaped areas in the kidney.

---

## Pyramidal Cell

A triangular-shaped excitatory neuron with a prominent apical dendrite, found especially in the cerebral cortex and hippocampus and involved in long-range brain signaling.

## Pyramidal Tract

The pyramidal tract is a major motor pathway from the cerebral cortex to the brainstem and spinal cord that controls voluntary movement.

## Pyramidal Tracts

Nerve pathways from brain movement areas down the brainstem and spinal cord, carrying signals controlling voluntary movement.

## Pyrazinamide

An antibiotic used with other medicines during the initial treatment of tuberculosis. It helps shorten treatment but can injure the liver or raise uric acid, so monitoring and prescribed dosing are important.

## Pyrazines

A family of chemical compounds found naturally in some foods and used in flavors, medicines, and industry. Their effects on health depend on the specific compound, amount, and route of exposure.

## Pyrazole

A small five-membered chemical structure used to build medicines, pesticides, and other compounds. The word names a chemical family and does not identify one particular treatment or effect.

## Pyrazoles

Pyrazoles are a family of chemical compounds containing a pyrazole ring, including some medicines and pesticides.

## Pyrenochaeta

A group of fungi found mainly in soil and plants that rarely infect people. When infection occurs, it usually affects tissue beneath injured skin or people with weakened immunity.

## Pyrethrins

Natural insect-killing chemicals derived from chrysanthemum flowers and used in some pest-control products. Exposure can irritate skin and airways and, in large amounts, may cause vomiting, tremors, seizures, or breathing problems.

## Pyrethrum

The dried flower heads of certain *Chrysanthemum* species, used as a natural source of pyrethrins, insecticidal compounds that affect insect sodium channels. Pyrethrum and pyrethrins differ from synthetic pyrethroids; exposure may irritate the skin or airways and can cause toxicity with substantial exposure.

## Pyridine

A nitrogen-containing ring-shaped chemical structure found in many medicines, vitamins, and biologically active substances. Pyridine groups can change how compounds dissolve, react, travel through the body, and produce medical effects.

## Pyridostigmine

A medicine that increases acetylcholine, a chemical helping nerves activate muscles. It mainly treats myasthenia gravis and may cause stomach cramps, diarrhea, sweating, increased saliva, muscle twitching, or slow heartbeat.

## Pyridostigmine Bromide

Pyridostigmine bromide increases communication between nerves and muscles and is used mainly for myasthenia gravis; it may cause abdominal cramps, diarrhea, or excess secretions.

## Pyridoxal Phosphate

The active form of vitamin B6 used by many enzymes. It helps the body process amino acids, make certain brain chemicals and red blood cells, and maintain normal nerve function.

## Pyridoxine

One form of vitamin B6, converted in the body to pyridoxal phosphate, a coenzyme for amino-acid metabolism. It is needed for processing food, mak-



---

ing red blood cells, and supporting nerves and the immune system; deficiency can cause anemia or nerve problems, while excessive supplements can damage nerves.

**Alternate Names:**

Vitamin B6

**Pyridoxine Deficiency**

Deficiency of vitamin B6, which is needed for amino-acid metabolism, neurotransmitter synthesis, hemoglobin production, and normal nerve function. It may cause dermatitis, glossitis, anemia, seizures, neuropathy, or immune dysfunction.

**Pyrimethamine**

A parasite-fighting medicine that blocks folate use in certain protozoa. It is used for selected infections, often with another medicine, and can suppress bone marrow or cause serious skin reactions.

**Pyrimidine**

A six-membered nitrogen-containing ring that forms the bases cytosine, thymine, and uracil in nucleic acids.

**Pyrimidine Analog**

A medicine or chemical resembling a pyrimidine nucleotide that can interfere with DNA or RNA synthesis, often for cancer or antiviral treatment.

**Pyrimidine Dimer**

A DNA lesion formed when adjacent pyrimidine bases bond abnormally after ultraviolet exposure, potentially causing mutations if unrepaired.

**Pyrimidine Nucleosides**

Nucleosides containing cytosine, thymine, or uracil bases linked to a sugar, serving as components or precursors of DNA and RNA.

**Pyrimidine Nucleotides**

Genetic and energy-related building blocks containing pyrimidine bases, including cytidine, uridine, and thymidine. Cells use them to make DNA, RNA, cell membranes, and other substances needed for

growth and repair.

**Pyrimidines**

Nitrogenous bases containing a single six-membered heterocyclic ring, serving as structural building blocks for nucleic acids. The primary pyrimidines are cytosine (C), thymine (T, present in DNA), and uracil (U, present in RNA).

**Pyrogen**

A pyrogen is a substance that induces fever, commonly through immune signaling; bacterial endotoxin is an important example in contaminated medicines or fluids.

**Pyroglyphidae**

Pyroglyphidae is a mite family that includes house-dust mites whose allergens can trigger allergic rhinitis, asthma, or atopic dermatitis.

**Pyromania**

A mental disorder involving recurrent deliberate fire-setting, mounting tension or arousal before the act, and pleasure, gratification, or relief afterward, without a primary external motive such as financial gain, revenge, concealment of a crime, or political purpose. The behavior is not better explained by conduct disorder, mania, psychosis, or substance use.

**Pyrophosphatase**

An enzyme that hydrolyzes inorganic pyrophosphate and helps drive many biosynthetic reactions forward. An enzyme that breaks down pyrophosphate, helping many chemical reactions needed to build cellular materials.

**Pyropoikilocytosis**

A rare inherited disorder that makes red blood cells unusually small, irregular, and fragile. The cells break down easily, causing hemolytic anemia with tiredness, jaundice, pale skin, or an enlarged spleen, often beginning in infancy or childhood.

---

## Pyroptosis

A form of controlled cell death that causes inflammation. The cell swells, breaks open, and releases signals that alert the immune system, often in response to infection or severe cell stress.

## Pyrosequencing

A laboratory method for reading selected DNA sequences by detecting light produced as DNA building blocks are added. It can identify some genetic changes or microbes but is not suitable for every sequencing need.

## Pyrexia

Alternate medical name; see *Fever*.

## Pyrexia Of Unknown Origin In A Hemodialysis Patient

Descriptive phrase for unexplained fever in a person receiving hemodialysis, not a separate disease. Infection of the dialysis access or bloodstream is an important possibility, but other causes must also be considered; see *Fever Of Unknown Origin*.

## Pyrosis

Alternate medical name; see *Heartburn*.

## Pyrroles

A group of five-membered nitrogen-containing ring compounds found in heme, bile pigments, medicines, and other biologically active molecules.

## Pyrrolidonecarboxylic Acid

A cyclic amino-acid derivative, also called pyroglutamic acid, found in proteins and metabolic pathways; abnormal accumulation may occur in selected metabolic or medication-related conditions.

## Pyrrolnitrin

Pyrrolnitrin is an antimicrobial compound produced by some bacteria and studied for antifungal activity and biological control.

## Pyruvate

A small molecule produced when cells break down

glucose. Cells can use it to make more energy, convert it into other substances, or produce lactate when oxygen or energy-processing pathways are limited.

## Pyruvate Carboxylase Deficiency

An inherited disorder caused by deficient pyruvate carboxylase, an enzyme needed for gluconeogenesis and other metabolic pathways. It can cause lactic acidosis, low blood glucose, poor growth, seizures, abnormal movements, developmental impairment, or liver failure.

## Pyruvate Decarboxylase

Pyruvate decarboxylase is an enzyme that converts pyruvate to acetaldehyde and carbon dioxide during fermentation in some organisms.

## Pyruvate Dehydrogenase

An enzyme complex that changes pyruvate, made when cells break down sugar, into acetyl-CoA for energy production. Inherited deficiency can cause excess lactic acid, poor growth, developmental problems, or neurologic disease.

## Pyruvate Dehydrogenase Complex Deficiency (PDCD)

An inherited disorder in which the pyruvate dehydrogenase complex cannot efficiently convert pyruvate into acetyl-CoA. It may cause lactic acidosis, poor growth, developmental impairment, seizures, low muscle tone, ataxia, or other neurologic disease.

## Pyruvate Kinase

An enzyme that helps red blood cells make energy from glucose. If it is inheritedly deficient, red cells may break down too early, causing chronic hemolytic anemia, jaundice, gallstones, tiredness, or an enlarged spleen.

## Pyruvate Kinase Blood Test

A pyruvate kinase blood test measures the enzyme in red blood cells and may help evaluate pyruvate kinase deficiency, an inherited cause of hemolytic anemia.

---

### Pyruvate Kinase Deficiency

A rare inherited condition in which red blood cells cannot make enough energy and break down early. It causes anemia of varying severity, jaundice, tiredness, gallstones, an enlarged spleen, and sometimes excess iron in the body.

### Pyuria

Presence of white blood cells or pus in urine, often indicating urinary-tract inflammation or infection.

### PYY

Peptide YY is a hormone released by the intestine after eating. It helps signal fullness and slows movement of food through the digestive tract; abnormal signaling is being studied in obesity and other metabolic disorders.

### Panic And Anxiety Disorders

Alternate name; see *Anxiety Disorders*.

### Parkinson's Disease Center

Alternate name; see *Parkinson's Disease*.

### Paroxysmal Nocturnal Hemoglobinuria (PNH)

Alternate name; see *Paroxysmal Nocturnal Hemoglobinuria*.

### Pediatric ADHD

Attention-deficit/hyperactivity disorder in children or adolescents, involving developmentally inappropriate inattention, hyperactivity, and/or impulsivity that persist across settings and impair functioning.

### Physical Allergy Urticaria

Hives triggered by a physical stimulus such as pressure, cold, heat, sunlight, vibration, water, or exercise. The trigger causes mast-cell activation and itchy raised welts; severe breathing or circulation symptoms require emergency care.

### PID (Pelvic Inflammatory Disease)

Alternate name; see *Pelvic Inflammatory Disease*.

### Pimples (Acne)

Alternate name; see *Acne*.

### PMS

Abbreviation; see *Premenstrual Syndrome*.

### Porcelain Inlay

A tooth-colored ceramic restoration bonded into a prepared cavity within a tooth. It can repair selected areas of decay or damage while preserving more tooth structure than a full crown.

### Postcoital Test

A fertility test that examines cervical mucus after intercourse for sperm presence and movement. It is rarely used now because results are variable and better-established fertility evaluations are available.

### Posttraumatic Stress Disorder (PTSD)

Alternate name; see *Post-Traumatic Stress Disorder*.

### Posttraumatic Stress Syndrome

Older or alternate name for *Post-Traumatic Stress Disorder*.

### Preeclampsia And High Blood Pressure

Alternate name; see *Preeclampsia*.

### Pregnancy With Asthma

Asthma during pregnancy may change as hormone, breathing, and immune conditions change. Poorly controlled asthma can reduce oxygen for the pregnant person and fetus, so prescribed controller treatment and regular obstetric and respiratory follow-up are important.

### Premenstrual Syndrome (PMS)

Alternate name; see *Premenstrual Syndrome*.

### Prevention Of Liver Cancer

Measures that lower the risk of liver cancer, especially preventing or treating chronic hepatitis B or C, limiting alcohol, avoiding tobacco, maintaining a healthy weight, and managing metabolic liver dis-

---

ease. People at high risk may need regular surveillance.

### **Paroxysmal Positional Nystagmus**

Involuntary, rhythmic eye movements triggered by a change in head position. It may accompany benign paroxysmal positional vertigo or arise from another vestibular or neurologic disorder; persistent, atypical, or neurologic-associated nystagmus needs evaluation.

### **Pfeiffer Syndrome Type I**

The milder form of Pfeiffer syndrome, a genetic craniosynostosis disorder usually associated with broad or deviated thumbs and great toes and premature fusion of some skull sutures. Severity and developmental effects vary.

### **Phakomatosis TS**

Alternate name; see *Tuberous Sclerosis Complex*.

### **Pharmacogenic Myopathy**

Muscle weakness or muscle injury caused by a medicine or drug. The presentation can include pain, cramps, elevated creatine kinase, or rhabdomyolysis, and evaluation considers the timing, dose, interactions, and other causes.

### **Phenylketonuria (PKU)**

Alternate name; see *Phenylketonuria*.

### **PPS (Post-Polio Syndrome)**

Alternate name; see *Post-Polio Syndrome*.

### **Prader Willi Syndrome**

Alternate spelling; see *Prader-Willi Syndrome*.

### **Prealbumin Tyr-77 Amyloidosis**

A rare hereditary transthyretin amyloidosis caused by a Tyr77 transthyretin variant. Misfolded protein can deposit in nerves, the heart, or other organs and may cause neuropathy, cardiomyopathy, or gastrointestinal symptoms.

### **Early-Onset Dementia**

Dementia beginning unusually early in life. Modern assessment identifies the specific cause, such as Alzheimer disease, frontotemporal dementia, vascular disease, or an inherited disorder, while also documenting the age at onset.

#### **Alternate Names:**

Premature Senility Syndrome (deprecated historical term)

### **Early-Onset Tremor**

Tremor beginning before older adulthood. The cause should be assessed for essential tremor, medication effects, dystonia, Parkinson disease, metabolic disease, or another neurologic disorder.

#### **Alternate Names:**

Presenile Tremor Syndrome (deprecated historical term)

### **Pressure Sores**

Alternate name; see *Pressure Sore*.

### **Primary Biliary Cirrhosis (PBC)**

Alternate name; see *Primary Biliary Cholangitis*.

### **Primary Cutaneous Amyloidosis**

Deposition of amyloid proteins limited mainly to the skin, causing itchy rippled, lichen-like, or darker patches. It differs from systemic amyloidosis, although assessment may be needed when symptoms or findings suggest disease beyond the skin.

### **Pudendal Neuralgia**

Pain caused by irritation or entrapment of the pudendal nerve, often felt in the perineum and worsened by sitting. It may include burning, tingling, altered sensation, or sexual and urinary symptoms and requires assessment for other pelvic causes.

### **Pulmonary Hypertension Due to Another Condition**

Pulmonary hypertension associated with another condition, such as left-heart disease, lung disease or hypoxia, chronic blood clots, or systemic disease. Classification by cause guides evaluation and

---

treatment.

### **Pulmonary Lymphangiomyomatosis**

Alternate name; see *Lymphangiomyomatosis*.

### **Pump Lung**

An older occupational term for byssinosis, a work-related airway disease caused by inhalation of cotton, flax, or hemp dust. Symptoms may include chest tightness, cough, wheezing, and breathlessness that worsen during or after workplace exposure.

### **Pure Cutaneous Histiocytosis**

A rare disorder in which abnormal histiocytes accumulate in the skin without recognized internal-organ involvement. It can produce papules, nodules, or plaques and must be distinguished from systemic histiocytic disease.

### **Purpura Hemorrhagica ITP**

Alternate name; see *Immune Thrombocytopenia*.

### **PVD (Peripheral Vascular Disease)**

Alternate name; see *Peripheral Vascular Disease*.

### **Polycystic Ovary Syndrome (PCOS)**

Alternate name; see *Polycystic Ovary Syndrome*.

### **PPH**

Abbreviation; see *Postpartum Hemorrhage*.

### **Prostate Enlargement/BPH**

Alternate name for benign prostatic hyperplasia, noncancerous enlargement of the prostate that can obstruct urine flow and cause weak stream, urgency, frequency, nighttime urination, or incomplete emptying.

### **Physical Abuse**

Deliberate use of physical force that causes or risks injury, pain, disability, or death. In clinical care, suspected abuse requires attention to immediate safety, documentation, and reporting duties under local law.

### **PI**

Abbreviation for primary immunodeficiency, a group of inherited or acquired disorders in which immune defenses are absent, reduced, or dysfunctional, leading to recurrent or unusual infections and sometimes autoimmune disease.

### **Pick's Disease**

A historical name for a subtype of frontotemporal dementia characterized by progressive changes in behavior, personality, language, or executive function and, on pathology, characteristic Pick bodies in affected neurons.

### **Prader-Labhart-Willi Fancone Syndrome**

Historical or variant naming for Prader-Willi syndrome, a chromosome 15 imprinting disorder causing infantile hypotonia and feeding difficulty followed by hyperphagia, obesity risk, hypogonadism, developmental differences, and short stature.

### **Pulmonary Lymphangiomyomatosis**

Alternate spelling for lymphangiomyomatosis, a rare cystic lung disease that mainly affects women and can cause progressive breathlessness, recurrent pneumothorax, chylous effusions, and abnormal growth of smooth-muscle-like cells.





# Q

## Q Angle

The quadriceps angle formed by the intersection of a line drawn from the anterior superior iliac spine (ASIS) to the central patella and a line from the central patella to the tibial tubercle (normally 10 to 14 degrees in males, 15 to 17 degrees in females); an increased Q angle accentuates lateral patellar displacement and predisposes to patellofemoral pain syndrome.

### Alternate Names:

Quadriceps Angle; Patellofemoral Tracking Angle

## Q-Angle Measurement

An orthopedic clinical measurement of the lateral pull of the quadriceps femoris muscle relative to the patellofemoral joint, defined as the angle formed between a line drawn from the anterior superior iliac spine to the center of the patella and a second line extending from the patellar center to the tibial tuberosity. An abnormally elevated Q-angle (greater than 20 degrees) predisposes to patellofemoral pain syndrome, lateral patellar subluxation, and chondromalacia patellae.

### Alternate Names:

Quadriceps Angle; Patellar Q-Angle

## Q Arm Of A Chromosome

The longer arm of a chromosome, identified by the letter q. A genetic report may use this label to show where a gene change or missing, extra, or rearranged DNA segment is located.

## Q-Banding

A laboratory staining method that creates bright and dim bands on chromosomes when viewed under

ultraviolet light. The pattern helps show chromosome structure and some rearrangements.

## Q Fever

An infection caused by the bacterium *Coxiella burnetii*, usually caught by breathing dust contaminated by infected cattle, sheep, goats, or their birth products. It often causes sudden fever, severe headache, tiredness, and muscle aches; some people develop pneumonia, hepatitis, or long-term heart-valve infection.

## QRS Axis

The general direction of the heart's electrical signal as the lower chambers contract, measured on an electrocardiogram. A shifted direction can occur with changes in the heart muscle, conduction system, or strain on the heart.

## QRS Complex

The group of waves on an electrocardiogram showing the electrical signal that makes the heart's lower chambers contract. An unusually wide or shaped complex may indicate a conduction problem, abnormal rhythm, or electrolyte disturbance.

## QRS Duration

The time taken for the electrical signal to spread through the heart's lower chambers, measured on an electrocardiogram. A longer-than-usual time can occur with a conduction problem or an abnormal rhythm.

## Q-Switch Laser

A specialized dermatologic and ophthalmologic laser

---

modality that delivers ultra-short pulses (nanosecond domain) of high peak energy, used for selective photothermolysis of cutaneous pigmented lesions, tattoo pigment clearance, and YAG posterior capsulotomy.

**Alternate Names:**

Quality-Switched Laser; Nanosecond Pulsed Laser

**QTc Dispersion**

The difference between the longest and shortest corrected QT intervals on an electrocardiogram. It describes variation in the heart's recovery time between beats, but its meaning is limited and depends on the full ECG and clinical setting.

**Alternate Names:**

Interlead QT Dispersion; Spatial Repolarization Heterogeneity

**QT Interval**

The time on an electrocardiogram representing electrical activation and recovery of the heart's lower chambers. Because heart rate changes this time, clinicians often use a corrected value when assessing rhythm risk.

**QT Prolongation**

An unusually long QT interval on an electrocardiogram. It may result from an inherited condition, a medicine, or abnormal body salts and can increase the risk of a rapid, dangerous rhythm that may cause fainting or cardiac arrest.

**Quadrant**

One of four sections used to describe a location, especially in the abdomen. The four abdominal quadrants help clinicians record where pain, tenderness, a lump, or another finding is located.

**Quadrantanopia**

Loss of vision in one quarter of the normal visual field, often from damage to the brain pathways carrying sight. The missing area may occur on the same side in both eyes and can affect reading, driving, walking, or safety.

**Quadrantectomy**

Surgery to remove one section of an organ, most often part of the breast containing a tumor plus a margin of healthy tissue. The need for lymph-node surgery or further treatment depends on the tumor and test results.

**Quadrate Lobe**

A small area on the underside of the liver between the gallbladder and the main vessels and ducts entering the liver. It is part of the left functional section of the liver.

**Alternate Names:**

Lobus Quadratus Hepatis; Couinaud Segment IVb

**Quadratus Femoris**

A small muscle deep in the buttock that turns the thigh outward and helps draw it toward the body.

**Quadratus Lumborum**

A muscle along the back of the abdomen that bends the body to one side and helps steady the lowest rib during breathing.

**Quadratus Plantae**

A deep muscle in the sole of the foot that helps the long toe-flexing muscle pull the toes straight downward.

**Alternate Names:**

Flexor Accessorius

**Quadriceps**

The four muscles on the front of the thigh that extend the knee; the rectus femoris also assists hip flexion. The quadriceps tendon connects them to the patella and contributes to the extensor mechanism.

**Quadriceps Active Test**

A knee examination used to look for injury to the posterior cruciate ligament. The person tightens the thigh muscle while the knee is bent, and forward movement of a previously sagging shinbone may support the finding. The test does not confirm a tear by itself.

**Alternate Names:**

---

Active Quadriceps Drawer Test; PCL Active Contraction Sign

### Quadriceps Contusion

A blunt injury to the muscles at the front of the thigh, often called a charley horse. It can cause torn muscle fibers, bleeding within the muscle, pain, swelling, and limited knee movement.

#### Alternate Names:

Charley Horse; Intramuscular Thigh Hematoma

### Quadriceps Femoris

The group of four muscles at the front of the thigh. It straightens the knee for standing, walking, climbing, and kicking; one part also helps bend the hip.

### Quadriceps Femoris Muscle

The four-muscle group at the front of the thigh that straightens the knee. It is essential for standing, walking, climbing, and rising from a chair; injury can cause pain and difficulty extending the leg.

### Quadriceps Injury

Damage to the quadriceps muscle or tendon caused by sudden force, overuse, or direct trauma. It may cause front-of-thigh pain, bruising, weakness, or difficulty extending the knee. Severe pain, a visible or felt gap, or inability to bear weight may occur with a major tear.

### Quadriceps Tendinitis

Overuse-related inflammation or degeneration of the quadriceps tendon, causing pain above the kneecap that worsens with jumping, running, or resisted knee extension.

### Quadriceps Tendon Rupture

Complete traumatic disruption of the confluent tendon of the quadriceps femoris muscle at its insertion into the superior pole of the patella, typically occurring following eccentric loading on a flexed knee in patients over 40 years of age, presenting with a palpable suprapatellar gap and inability to perform an active straight leg raise.

#### Alternate Names:

Suprapatellar Tendon Rupture; Extensor Mechanism Disruption

### Quadriceps Tendon Ruptures

A tear of the tendon connecting the quadriceps muscle to the patella, causing weakness or inability to extend the knee. Complete tears often require surgical repair; treatment depends on the extent of injury and the person's functional needs.

### Quadrigeminal Cistern

A subarachnoid cistern located dorsal to the midbrain tectum (corpora quadrigemina / superior and inferior colliculi), ventral to the splenium of the corpus callosum and tentorium cerebelli, containing the vein of Galen, the basal veins of Rosenthal, and the posterior cerebral arteries.

#### Alternate Names:

Cisterna Quadrigeminalis; Cisterna Ambiens Dorsalis

### Quadrigeminal Plate

The dorsal roof (tectum) of the midbrain bearing four rounded sensory eminences: the paired superior colliculi (visual reflexes and saccadic eye movements) and inferior colliculi (auditory relay processing).

#### Alternate Names:

Tectal Plate; Lamina Tecti

### Quadrilateral Space Syndrome

A neurovascular compression syndrome caused by entrapment of the axillary nerve and posterior circumflex humeral artery within the quadrilateral space (bounded by teres minor, teres major, long head of triceps, and humerus), presenting with posterior shoulder ache, paresthesias over the lateral shoulder, and deltoid weakness.

#### Alternate Names:

Quadrangular Space Syndrome; Axillary Nerve Compression Syndrome

### Quadriparesis

Weakness affecting both arms and both legs. It may result from disease or injury to the neck spinal cord, brain, nerves, or muscles and can impair movement, breathing, and bladder or bowel control.

---

## Quadriplegia

Paralysis affecting both arms and both legs, usually with weakness of the trunk, because of damage to the neck portion of the spinal cord or another major nervous-system injury. It can impair breathing and bladder or bowel control.

## Quadruple Screen Test

A second-trimester blood test that combines four pregnancy-related markers to estimate the chance of selected chromosome conditions and neural-tube defects. It does not diagnose a condition; abnormal results need further assessment.

## Quadruplet Pregnancy

A pregnancy involving four fetuses. It has increased risks of preterm birth, restricted fetal growth, high blood pressure, metabolic problems, and illness in the pregnant person.

## Quad Screen

A second-trimester blood screening test that estimates the chance of certain chromosome conditions and neural-tube defects in the fetus. It estimates risk rather than making a diagnosis; an abnormal result may require ultrasound or diagnostic testing.

### Alternate Names:

Quadruple Marker Test; AFP-Plus; Second-Trimester Screen

## Qualitative HCG Blood Test

A diagnostic serological laboratory test that detects the presence or absence of human chorionic gonadotropin (hCG) in maternal blood. Primarily utilized for early confirmation of pregnancy.

### Alternate Names:

Qualitative Serum Pregnancy Test; Blood hCG Screen; Qualitative Beta-hCG

## Quality-Adjusted Life Year

A standardized composite health economic metric that combines both the quantity (length) and quality (health-related utility weight from 0 for death to 1 for perfect health) of life generated by a medical intervention, widely used in cost-effectiveness analysis and health technology assessment.

### Alternate Names:

QALY

## Quality Of Life

A person's experience of physical health, emotional well-being, relationships, independence, and ability to do meaningful activities. Illness and treatment may affect these areas differently for each person.

## Quantitative Bence-Jones Protein Test

A urine test measuring abnormal light-chain proteins made by some plasma-cell cancers. It can help diagnose or monitor disorders such as multiple myeloma, but results are interpreted with blood tests and other findings.

## Quantitative Computed Tomography

A CT scan that measures bone density while showing bone structure. It can help assess osteoporosis, unusual bone loss, or fracture risk when standard bone-density testing is unsuitable or more detail is needed.

## Quantitative HCG Blood Test

A blood test that measures the amount of human chorionic gonadotropin (hCG). It can help assess early pregnancy, possible ectopic pregnancy or miscarriage, and some pregnancy-related or germ-cell tumors.

### Alternate Names:

Quantitative Beta-hCG; Serial hCG Test; Serum hCG Level

## Quantitative Immunoglobulins

A blood test measuring the main classes of antibodies. Low levels may suggest an immune deficiency, while high or uneven levels may occur with infection, inflammation, liver disease, or some blood cancers; results need clinical interpretation.

## Quantitative Nephelometry Test

A laboratory test that estimates the amount of selected proteins by measuring how much light they scatter. It can measure antibodies, complement proteins, and other proteins involved in immune, kidney, or blood disorders.

---

### Quantitative PET

Analysis of a PET scan that estimates how much radioactive tracer is taken up by tissue. Standardized measurements can help compare disease activity or treatment response, but results are affected by the tracer, timing, blood sugar, and scanning method.

### Quantitative Sensory Testing

A test that measures how strongly a person senses vibration, temperature, or pain. It can help assess nerve damage affecting small or large nerve fibers.

#### Alternate Names:

QST; Psychophysical Sensory Threshold Assay

### Quarantine

Temporary separation or movement restrictions for people exposed to an infectious disease while monitoring whether they become ill. It differs from isolation, which separates people who are already infected.

### Quartan Fever

A fever pattern in which attacks recur about every 72 hours, classically seen with the malaria parasite *Plasmodium malariae*. Fever timing alone cannot diagnose malaria; testing is needed in anyone with possible exposure and fever.

### Queckenstedt Test

A classical diagnostic maneuver performed during lumbar puncture; bilateral manual compression of the internal jugular veins causes a prompt rise in CSF pressure on spinal manometry in normal individuals; failure of CSF pressure to rise indicates a complete mechanical blockage in the spinal subarachnoid space.

#### Alternate Names:

Queckenstedt's Sign; Jugular Venous Spinal Manometry Test

### Queensland Tick Typhus

An Australian spotted fever group rickettsial infection caused by the obligate intracellular bacterium *Rickettsia australis*, transmitted to humans by the bite of the paralysis tick *Ixodes holocyclus*. Clinically characterized by fever, headache, regional

lymphadenopathy, a localized necrotic inoculation eschar at the tick bite site, and a maculopapular or vesicular rash; responds rapidly to doxycycline.

#### Alternate Names:

QTT; Australian Tick Typhus

### Quellung Reaction

A classic microbiological microscopic diagnostic reaction in which specific polyvalent or monovalent anticapsular antibodies bind to the polysaccharide capsule of encapsulated bacteria (such as *Streptococcus pneumoniae*, *Neisseria meningitidis*, or *Haemophilus influenzae*), increasing its refractive index and causing it to appear sharply demarcated and swollen.

#### Alternate Names:

Neufeld Quellung Phenomenon; Capsular Swelling Test

### Quetiapine Fumarate

A form of quetiapine, a medicine used for schizophrenia, bipolar disorder, and selected depression treatment. It can cause sleepiness, dizziness on standing, weight or blood-sugar changes, and movement problems.

### Quickening

The first time a pregnant person feels fetal movement, often as a flutter or tapping sensation during the second trimester. Timing varies with the pregnancy, body shape, and position of the placenta.

### Quick Relievers

Fast-acting asthma medicines used when symptoms suddenly worsen. They quickly relax tightened airway muscles and can ease wheezing, coughing, chest tightness, and breathlessness, but they do not replace regular treatment that controls airway inflammation.

#### Alternate Names:

Rescue Inhalers; Fast-Acting Bronchodilators

### Quick Sequential Organ Failure Assessment

A bedside score used with suspected infection to identify people at increased risk of serious illness.

---

It considers changes in alertness, fast breathing, and low blood pressure. It supports, but does not replace, a full assessment for sepsis.

**Alternate Names:**

qSOFA; Quick SOFA Score

**Quick Test**

The historical and foundational one-stage prothrombin time (PT) coagulation test introduced by Armand Quick in 1935, which measures the extrinsic and common pathways of coagulation (factors II, V, VII, X, and fibrinogen) following addition of tissue thromboplastin and calcium.

**Alternate Names:**

Quick Prothrombin Time; One-Stage Prothrombin Assay

**Quinalphos**

A poisonous insecticide that disrupts nerve-signal control. Significant exposure can cause sweating, excess saliva, vomiting, muscle twitching, weakness, seizures, breathing failure, or death.

**Quinapril**

Quinapril is an angiotensin-converting-enzyme inhibitor used for hypertension and heart failure; cough, hyperkalaemia, renal-function changes, angioedema, and fetal toxicity are important risks.

**Quincke Edema**

An acute, episodic, well-demarcated swelling of the deep dermis, subcutaneous tissues, or submucosal tissues (angioedema), classically involving the lips, eyelids, tongue, larynx, or gastrointestinal tract, mediated by histamine release or excessive bradykinin generation.

**Alternate Names:**

Angioneurotic Edema; Acute Angioedema

**Quincke Needle**

A traditional spinal needle featuring a sharp, medium-length cutting bevel with a matching stylet, utilized for diagnostic lumbar puncture and spinal anesthesia, historically associated with a higher incidence of post-dural puncture headache compared to atraumatic pencil-point needles.

**Quincke Pulse**

A physical clinical sign of severe aortic regurgitation characterized by visible, rhythmic, alternating subungual capillary flushing and blanching occurring synchronously with the cardiac cycle when gentle pressure is applied to the distal edge of a fingernail.

**Alternate Names:**

Quincke's Sign; Subungual Capillary Pulsation

**Quincke Pulse Sign**

An eponymous physical cardiovascular sign characterized by alternating subungual erythema and blanching of the nail bed capillary plexus, synchronized with ventricular systole. Elicited by applying gentle compressive pressure to the tip of a fingernail; it represents a manifestation of the hyperdynamic wide pulse pressure and rapid runoff characteristic of severe chronic aortic regurgitation.

**Alternate Names:**

Quincke Capillary Pulsation; Quincke Sign

**Quincke's Disease**

Sudden swelling of deeper layers of the skin or mucosa, commonly affecting the lips, face, tongue, throat, or bowel; airway involvement is an emergency.

**Alternate Names:**

Angioedema; Giant Urticaria

**Quincke's Edema**

Older term; see *Angioedema*.

**Quincke's Pulse**

Visible flushing and paling of the nail bed or skin with each heartbeat. It may occur when the difference between the upper and lower blood-pressure numbers is unusually wide, especially with severe aortic-valve leakage, but it is not diagnostic alone.

**Quincke's Sign**

Alternating redness and paling of the nail bed or skin with each heartbeat, usually caused by a wide pulse pressure. It can be seen in severe aortic-valve leakage but is not diagnostic by itself.



---

## Quinidine

A heart-rhythm medicine that changes electrical conduction in the heart. It is used selectively because it can cause dangerous abnormal rhythms, low blood pressure, low platelets, diarrhea, or other serious side effects.

## Quinidine Syncope

Sudden syncope or cardiac arrest occurring in patients treated with quinidine or related class Ia antiarrhythmic agents, caused by marked prolongation of the QT interval triggering self-terminating or degenerating episodes of polymorphic ventricular tachycardia (torsades de pointes).

### Alternate Names:

Antiarrhythmic Drug-Induced Syncope; Quinidine Torsades Syncope

## Quinine

A medicine derived from cinchona bark that can treat selected malaria infections. It is used less often because safer options are available and it can cause low blood sugar, ringing in the ears, blood-cell problems, or abnormal heart rhythms.

## Quinine Amblyopia

Acute toxic retinal and optic nerve damage caused by quinine toxicity (cinchonism), presenting with sudden bilateral visual acuity loss, marked concentric visual field constriction, retinal arteriolar attenuation, and optic disc pallor.

### Alternate Names:

Quinine Retinopathy; Cinchona Retinal Toxicity

## Quinolone

A class of antibiotics that stop bacteria from copying and repairing their DNA. Some treat serious infections, but side effects can include tendon injury, nerve symptoms, abnormal heart rhythms, or severe diarrhea, so use depends on the infection and risks.

## Quinolone Antibiotics

A major synthetic bactericidal antibiotic class (fluoroquinolones, such as ciprofloxacin, levofloxacin, and moxifloxacin) that selectively inhibits bacterial type II topoisomerases (DNA gyrase and topoisomerase

IV), thereby arresting bacterial DNA replication and transcription.

### Alternate Names:

Fluoroquinolones

## Quinolone-Induced Tendinopathy

An adverse musculoskeletal drug reaction associated with fluoroquinolone antimicrobial therapy (such as ciprofloxacin or levofloxacin), characterized by painful tendinitis and an elevated risk of tendon rupture (predominantly the Achilles tendon), especially in patients receiving concurrent corticosteroids or with renal impairment.

### Alternate Names:

Fluoroquinolone Tendonitis; Fluoroquinolone-Associated Achilles Rupture

## Quinsy

An abscess beside a tonsil, usually after a throat infection. It can cause severe one-sided throat pain, fever, muffled speech, drooling, difficulty opening the mouth, or breathing difficulty. Treatment usually includes drainage and antibiotics.

## Quinsy Peritonsillar Abscess

A suppurative localized collection of purulent exudate in the potential space between the palatine tonsil capsule and the superior pharyngeal constrictor muscle, typically arising as a complication of acute bacterial tonsillopharyngitis (frequently involving *Streptococcus pyogenes* and oral anaerobes). Characterized by severe unilateral sore throat, odynophagia, hot potato muffled voice, trismus, and medial displacement of the tonsil with contralateral uvular deviation.

### Alternate Names:

Quinsy; Peritonsillar Abscess; PTA

## Quinupristin-Dalfopristin

An intravenous antibiotic combination used rarely for selected serious infections caused by resistant gram-positive bacteria. It can cause muscle or joint pain, liver problems, infusion reactions, and medicine interactions.

---

## Quit Smoking

Alternate name; see *Smoking Cessation*.

## Quorum Sensing

A density-dependent bacterial intercellular communication mechanism mediated by the release and detection of small diffusible autoinducer signaling molecules, coordinating group behaviors such as biofilm formation, bioluminescence, and virulence factor expression once a threshold bacterial population density is achieved.

## Quotidian Fever

A fever pattern in which fever rises or attacks occur about every 24 hours. It can occur with malaria and other infections, but the timing of fever alone does not identify the cause.

## Quotidian Malaria

A recurring pattern of malarial paroxysms characterized by daily febrile spikes occurring at 24-hour intervals, classic for infection caused by *Plasmodium knowlesi* or resulting from overlapping broods of *Plasmodium vivax* or *Plasmodium falciparum*.

### Alternate Names:

24-Hour Malarial Paroxysm; Daily Malarial Fever

## Q Wave

A downward movement that may begin the electrical pattern of a heartbeat on an electrocardiogram. It can be normal in some recording positions or suggest previous heart-muscle injury, depending on its size and location.

## Q Waves

Early downward waves on an electrocardiogram. Some patterns may suggest a previous heart attack or another heart problem, but Q waves must be interpreted with the other ECG findings and the person's history.

### Alternate Names:

Pathologic Q Wave; Myocardial Infarction Q Wave

# R

## Rabeprazole

A medicine that greatly reduces acid made by the stomach. It treats reflux and ulcers and may be combined with antibiotics for *Helicobacter pylori* infection; possible effects include headache, diarrhea, and rarely low magnesium or bowel infection.

## Rabies

A viral infection of the brain and nervous system caused by rabies virus, a virus that infects mammals. It spreads mainly when saliva from an infected animal enters a bite, scratch, broken skin, or the eyes, mouth, or another mucous membrane. After a variable symptom-free period, early symptoms may include fever, weakness, headache, or pain, itching, or unusual sensations near the exposure site. Later symptoms may include confusion, agitation, hallucinations, fear of water, difficulty swallowing, paralysis, coma, and death. Once symptoms begin, rabies is nearly always fatal, but vaccination before exposure or post-exposure treatment can prevent disease before symptoms start.

## Rabies Vaccination

Immunization used before or after exposure to reduce the risk of rabies. Post-exposure management also includes wound care and, when indicated, rabies immunoglobulin.

## Rabies Virus

The virus that causes rabies, a usually fatal viral infection of the central nervous system transmitted most often through the saliva of an infected animal.

## Rachitic Rosary

A physical examination finding in pediatric rickets (vitamin D deficiency) characterized by prominent, beadlike, symmetrical nodular expansion of the costochondral junctions along the anterolateral rib cage resulting from disordered osteoid mineralization and chondrocyte hypertrophy.

### Alternate Names:

Beaded Ribs; Costochondral Rosary

## Radial

Relating to the radius bone or the thumb side of the forearm and hand. The word can also describe structures arranged like spokes spreading from a central point.

## Radial Artery

An artery that runs along the thumb side of the forearm to the wrist and hand. Its pulse can be felt at the wrist, and it supplies blood to the forearm and parts of the hand.

## Radial Artery Thrombosis

A blood clot blocking the artery on the thumb side of the forearm, sometimes after an arterial catheter or injury. It may cause a painful, pale, cold, numb, or weak hand and can threaten hand circulation.

## Radial Club Hand

A birth-related difference in which the thumb-side forearm bone is small or absent and the hand bends toward the thumb side. The thumb may also be underdeveloped, and treatment depends on movement, function, and associated conditions.

---

### **Radial Head Subluxation**

A common elbow injury in young children, often called nursemaid's elbow, caused when a sudden pull partly slips a forearm bone out of its supporting band. The child may stop using the arm but usually has little swelling. A trained clinician can put the bone back in place.

#### **Alternate Names:**

Nursemaid's Elbow; Pulled Elbow; Annular Ligament Displacement

### **Radial Nerve**

A major nerve of the arm that helps straighten the elbow, wrist, and fingers and carries feeling from part of the back of the hand. Injury can cause wrist drop, finger weakness, or numbness.

### **Radial Nerve Dysfunction**

Impaired function of the radial nerve causing weakness of wrist or finger extension and sensory changes over part of the posterior arm or hand; causes include compression, trauma, and systemic neuropathy.

### **Radial Nerve Palsy**

Dysfunction of the radial nerve causing weakness of wrist and finger extension, often producing wrist drop, with sensory loss over part of the back of the hand in some cases.

### **Radial Tunnel Syndrome**

Irritation or pressure on the radial nerve near the elbow, causing aching outer-forearm pain and sometimes weakness. Symptoms may worsen with repeated wrist or forearm movement and require evaluation to exclude other causes.

### **Radial Vein**

Veins running alongside the radial artery on the thumb side of the forearm. They drain blood from the forearm and join other deep veins near the elbow.

### **Radiation**

Energy traveling as particles or electromagnetic waves, including X-rays, gamma rays, ultraviolet

light, and radio waves. Depending on its type and amount, radiation can diagnose disease, treat cancer, or damage living tissue.

### **Radiation Biology**

The study of how radiation affects cells, tissues, organisms, and populations. It examines injury, repair, mutations, cancer risk, treatment effects, and protective measures after exposure to ionizing or nonionizing radiation.

### **Radiation Contamination**

Radioactive material present on or inside a person, object, surface, or body site when it should not be. It can spread or continue exposing tissue and may require isolation, monitoring, removal of the material, and specialist advice.

### **Radiation Dermatitis Skin Care**

Care of skin affected by radiation treatment. The skin may become red, dry, sore, peeling, or moist, and care focuses on protecting the skin and treating irritation or infection when present.

#### **Alternate Names:**

Radiation Skin Care; Radiation Dermatitis Management; Post-Radiotherapy Skin Care

### **Radiation Dosage**

The amount of radiation given during an imaging test or radiation treatment. The dose is planned to obtain the needed benefit while limiting unnecessary exposure.

### **Radiation Dose**

The amount of ionizing radiation absorbed or received by a person or tissue. Dose varies with the examination, equipment, protocol, and patient size; imaging should be justified and optimized to obtain the needed information with the lowest reasonable exposure.

### **Radiation Emergency**

An event involving actual or suspected harmful radiation exposure or contamination that requires urgent protective and medical measures.

---

### **Radiation Enteritis**

Inflammation or injury of the small or large intestine after radiation treatment. Early symptoms may include diarrhea, cramps, nausea, or urgency; months or years later, scarring can cause narrowing, poor absorption, bleeding, or blockage.

### **Radiation Exposure**

Contact with ionizing or non-ionizing radiation from medical imaging, treatment, sunlight, workplaces, or other sources. The possible harm depends mainly on the type, dose, body area, and duration of exposure.

### **Radiation Fibrosis**

Long-term thickening and stiffening of tissue caused by radiation injury, sometimes appearing months or years after treatment. It can affect skin, lungs, bowel, or other organs and may cause pain, reduced movement, or shortness of breath.

### **Radiation Field**

The area and distribution of radiation delivered during imaging or radiation treatment. In therapy, its size, shape, direction, and dose are planned to cover the target while limiting exposure to nearby healthy tissue.

### **Radiation-Induced Angiosarcoma**

A rare, fast-growing cancer of blood-vessel lining cells that develops years after radiation in previously treated tissue. It may appear as a bruise, purple patch, swelling, or lump and requires prompt specialist assessment.

### **Radiation Leukemia**

Radiation leukemia is leukemia that develops after significant exposure to ionizing radiation; it is a late treatment- or exposure-related complication.

### **Radiation Monitoring**

Measurement of radiation levels or individual exposure using devices such as personal dosimeters and area monitors.

### **Radiation Necrosis**

Permanent death of tissue caused by radiation injury, often developing months or years after treatment. In the brain it may cause headache, seizures, swelling, weakness, or speech and vision problems and can resemble returning cancer on scans.

### **Radiation Oncologist**

A doctor who plans and oversees radiation treatment for cancer and selected noncancerous conditions. They choose treatment areas and doses, coordinate care, monitor side effects, and adjust therapy during treatment.

### **Radiation Oncology**

A medical specialty that uses ionizing radiation to treat cancer. Treatment may aim to destroy or control tumors, prevent cancer from returning, or relieve symptoms.

### **Radiation Physics**

The study of how radiation is produced, travels, interacts with tissues, and is measured. It supports medical imaging, radiation treatment, dose calculation, and protection from unnecessary exposure.

### **Radiation Pneumonitis**

Inflammation of lung tissue occurring weeks or months after radiation to the chest. It may cause dry cough, fever, breathlessness, or low oxygen and can later leave permanent scarring; new breathing symptoms need medical assessment.

### **Radiation Poisoning**

Illness caused by substantial exposure to ionizing radiation, potentially producing skin injury, bone-marrow suppression, gastrointestinal symptoms, or organ failure.

### **Radiation Pregnancy Screening**

Assessment of pregnancy status before procedures involving ionizing radiation or radioactive materials when pregnancy is possible. The decision to proceed requires clinical justification, risk optimization, and consideration of gestational age and fetal dose.

---

### **Radiation Proctitis**

Inflammation and injury of the rectum after pelvic radiation therapy, causing rectal pain, urgency, diarrhea, mucus, or bleeding.

### **Radiation Protection**

The practice of keeping radiation exposure as low as reasonably achievable while obtaining the required diagnostic or therapeutic result. It relies on justification, optimization, time reduction, distance, shielding, and appropriate contamination control.

### **Radiation Recall**

Inflammation that returns in an area previously treated with radiation after a later trigger, often a medicine. The skin or lining may become red, sore, swollen, peel, or blister, and the treating team should review the suspected trigger.

### **Radiation-Recall Dermatitis**

An inflammatory skin reaction that occurs in a previously irradiated area after exposure to a triggering substance, often months or years later. The area may become red, swollen, painful, or blistered and requires clinical evaluation.

### **Radiation Safety**

Measures that limit unnecessary exposure to ionizing radiation while allowing needed imaging or treatment. They include using the shortest practical exposure time, keeping as much distance as possible, using shielding, preventing contamination, and following pregnancy-related precautions when relevant.

### **Radiation Sickness**

Acute radiation syndrome caused by high-dose whole-body or substantial partial-body exposure to ionizing radiation. It may produce nausea, vomiting, marrow failure, infection, bleeding, gastrointestinal injury, neurovascular collapse, and death depending on dose.

### **Radiation Therapy**

Treatment that uses carefully aimed ionizing radiation to damage cancer cells so they stop dividing or

die. It may be given from outside the body or from a source placed near or inside a tumor. Side effects depend on the body area and dose and may include tiredness or skin changes.

### **Radiation Tolerance**

The amount of radiation that a tissue or organ can receive before injury becomes likely. It varies with the tissue, dose, treatment schedule, and previous radiation.

### **Radical Cystectomy**

Surgery to remove the urinary bladder, usually for muscle-invasive or high-risk bladder cancer. Because urine can no longer leave through the bladder, another route for urine drainage is created.

### **Radical Cystoprostatectomy**

Radical cystoprostatectomy removes the urinary bladder, prostate, and nearby structures, usually to treat invasive bladder cancer in men.

### **Radical Hysterectomy**

An operation that removes the uterus and cervix together with surrounding parametrial tissues and the upper vagina, usually with pelvic lymph-node assessment or removal. It is used mainly for selected invasive cervical cancers and is more extensive than a simple or total hysterectomy.

### **Radical Lymphadenectomy**

Extensive surgery to remove lymph nodes and nearby tissue to look for or treat cancer spread. It may provide important staging information but can cause pain, nerve or vessel injury, and long-term swelling from lymph-fluid blockage.

### **Radical Mastectomy**

Radical mastectomy removes the breast, chest muscles, and many underarm lymph nodes; it is now rarely used because less extensive cancer surgery is often effective.

### **Radical Nephrectomy**

Radical nephrectomy removes an entire kidney and



---

sometimes nearby tissues, usually to treat a kidney tumor.

### **Radical Prostatectomy**

Surgery to remove the whole prostate gland and usually the nearby seminal vesicles, most often to treat prostate cancer that has not spread widely. The urinary passage is rejoined to the bladder; possible effects include urine leakage, erection problems, bleeding, and infection.

### **Radicular Cyst**

An inflammatory odontogenic cyst arising at the apex of a nonvital tooth, usually from chronic pulpal infection and apical periodontitis.

### **Radiculitis**

Irritation or inflammation of a nerve root where it leaves the spine. It can cause pain spreading along an arm or leg, tingling, numbness, weakness, or changed reflexes because of a disk problem, narrowing, infection, or other disease.

### **Radiculopathy**

Dysfunction of a spinal nerve root, usually from disc herniation, narrowing, or inflammation. It may cause shooting pain, tingling, numbness, weakness, or altered reflexes along the affected nerve pathway.

### **Radioactive Iodine**

A radioactive form of iodine taken up by thyroid cells. It is used to image the thyroid or destroy overactive thyroid tissue and selected thyroid cancer cells; radiation-safety precautions are needed after some treatments.

### **Radioactive Iodine Ablation**

Treatment using radioactive iodine to destroy remaining thyroid tissue or selected thyroid cancer cells after surgery. Thyroid cells absorb the iodine; the decision depends on cancer type, spread, recurrence risk, and the person's overall health.

### **Radioactive Iodine Uptake**

A test measuring the percentage of administered

radioactive iodine taken up by the thyroid over time, used to evaluate hyperthyroidism and selected thyroid disorders.

### **Radioactive Iodine Uptake Test**

A nuclear medicine test that measures how much radioactive iodine the thyroid gland absorbs. It helps evaluate hyperthyroidism and some thyroid disorders.

### **Radioactivity**

Energy released when unstable atoms change into more stable forms. The energy may be particles or rays and can be used in medical imaging or treatment, but enough exposure can damage living tissue.

### **Radioallergosorbent Test**

An in vitro radioimmunoassay that detects and quantifies allergen-specific IgE antibodies in a patient's blood serum, used to identify allergic sensitization to environmental aeroallergens, foods, medications, or venoms without risking in vivo anaphylaxis.

#### **Alternate Names:**

RAST

### **Radiobiology**

The study of how radiation affects living cells and tissues, including DNA damage, cell death, repair, mutations, treatment effects, and the development of cancer.

### **Radiocarpal Joint**

The main wrist joint between the forearm bone on the thumb side and several wrist bones. It allows the hand to bend forward, backward, and from side to side and may be injured by falls or arthritis.

### **Radiochemical Purity**

The proportion of a radioactive medicine present in the intended chemical form. Unwanted forms can make a scan less accurate, send radiation to the wrong tissues, or reduce treatment effectiveness.

---

## Radiochemistry

The chemistry of radioactive substances and compounds. In medicine it includes making and labeling radiopharmaceuticals, studying radioactive decay, separating products, and checking radiochemical purity and safety.

## Radiocontrast

A substance given during an imaging test to make selected tissues, blood vessels, or body spaces easier to see. Depending on the agent, allergic reactions, kidney injury, or leakage-related damage may occur.

## Radiodermatitis

Skin injury caused by ionizing radiation, such as radiation treatment. It may cause redness, dryness, itching, hair loss, peeling, sores, or long-term skin changes; severity depends on the dose, treatment area, and repeated exposure.

## Radioembolisation

Alternate name; see *Radioembolization*.

## Radioembolization

A treatment that delivers tiny radioactive beads through an artery to selected liver tumors. The beads reduce the tumor's blood supply and release radiation nearby; suitability depends on liver function, tumor extent, and blood flow.

## Radio-Femoral Delay

A clinically palpable lag or latency in the arrival of the femoral pulse compared to the simultaneous radial pulse, often accompanied by reduced femoral pulse amplitude. It is a classic clinical sign of coarctation of the aorta and extensive aortoiliac occlusive disease.

## Radiofrequency

A type of electromagnetic energy, between some radio waves and microwaves, used in communication and in selected medical tests and treatments. Medical uses include heating tissue or delivering energy during procedures.

**Alternate Names:**

Radiofrequency Energy; RF Radiation

## Radiofrequency Ablation

A procedure that uses high-frequency electrical energy to heat and destroy selected tissue, often under imaging guidance.

## Radiofrequency Ablation For Chronic Pain

A procedure that applies radiofrequency energy to selected sensory nerves or other targets to interrupt pain signaling for a limited period in carefully selected chronic pain conditions.

## Radiograph

An image made with X-rays. Dense structures such as bone appear light, while air appears dark and soft tissues appear in shades of gray. Radiographs are used to assess bones, the chest, teeth, and selected abdominal or foreign-body problems; they use a small amount of radiation.

**Alternate Names:**

X-Ray; Roentgenogram; Radiographic Image

## Radiographer

A trained professional who performs diagnostic imaging examinations or delivers radiotherapy, depending on specialty and jurisdiction.

## Radiography

Creation of images using X-rays. It is useful for bones, chest disease, dental structures, and many other conditions; radiation exposure is kept as low as reasonably achievable.

## Radioimmunoassay

A laboratory test that uses an antibody and a small radioactive label to measure a hormone, medicine, or protein in a sample. It can be very sensitive but is often replaced by newer nonradioactive tests.

## Radioimmunoguided Surgery

Surgery in which a radioactive tracer attached to an antibody helps the surgeon locate tissue carrying a particular marker, often tumor tissue. Special

---

equipment detects the tracer during the operation.

### **Radioimmunotherapy**

Treatment that attaches a radioactive substance to an antibody so the antibody carries radiation toward cells with a particular marker, often cancer cells. Radiation can still affect nearby normal tissue and cause side effects.

### **Radioiodine**

A radioactive form of iodine used in nuclear medicine to image or treat thyroid tissue. It is taken up by thyroid cells; precautions are needed to limit radiation exposure to others.

### **Radioiodine Therapy**

Treatment using radioactive iodine, which is taken up mainly by thyroid cells to destroy overactive thyroid tissue or selected thyroid cancer cells. It requires individualized dosing and temporary radiation-safety precautions; pregnancy and breast-feeding are contraindications.

### **Radioisotope**

An unstable isotope that emits radiation as it decays; radioisotopes are used in medical imaging, diagnosis, and selected treatments.

#### **Alternate Names:**

Radioactive Isotope; Radiotracer

### **Radioligand Therapy**

A cancer treatment in which a radioactive particle is attached to a molecule that binds a target on cancer cells. The molecule delivers radiation to cells carrying that target; use depends on the cancer, target expression, organ function, and treatment-specific safety monitoring.

### **Radiologic Health**

The safe use of medical imaging and radiation-based treatment to protect patients, workers, and the public from unnecessary ionizing radiation while preserving clinical benefit. Safety measures include appropriate justification, shielding, and dose optimization.

### **Radiologic-Pathologic Correlation**

Comparing scan findings with tissue examined under a microscope to determine whether the sample explains the abnormal area. If they do not match, another sample or further testing may be needed.

### **Radiologic Response**

The change seen on medical scans after treatment, such as shrinkage, disappearance, stability, growth, or new lesions. Standard measurement rules help classify the response, but scans do not always show the full biological effect.

### **Radiologic Technologist**

A trained health professional who performs selected medical-imaging examinations and helps protect patients from unnecessary radiation exposure.

### **Radiologist**

A doctor trained to interpret medical images and sometimes perform image-guided procedures. Radiologists evaluate X-rays, ultrasound, CT, MRI, and other scans to diagnose disease, guide treatment, or obtain tissue samples.

### **Radiology**

The medical field that uses images to find and monitor disease and guide procedures. Radiologists interpret X-rays, ultrasound, CT, MRI, and other scans and may perform image-guided biopsies, drainages, injections, or treatments.

### **Radiology-Patient Identification**

Verification of the correct patient, examination, body site, indication, pregnancy status when relevant, allergies, renal function, and prior imaging before image acquisition or intervention. It is a core diagnostic and procedural safety step.

### **Radiology Report**

A written explanation of an imaging study. It describes why the study was done, the important findings, and their meaning; urgent or unexpected findings should be communicated promptly to the treating team.

---

## **Radiolucent**

Describing a material that lets more X-rays pass through and therefore appears darker on an X-ray or CT image. Air is highly radiolucent; some stones, foreign bodies, and abnormal tissues can also have this appearance.

## **Radiomimetic**

Radiomimetic describes a chemical or treatment that produces cellular or DNA damage resembling the effects of ionizing radiation.

## **Radionuclide**

An unstable form of an atom that releases radiation as it changes into a more stable form. In medicine, radionuclides can be attached to substances for imaging or used to deliver treatment to selected tissues.

## **Radionuclide Angiography**

An imaging test in which a small amount of radioactive tracer is used to follow blood flow through the heart or blood vessels. A camera detects the tracer and produces information about circulation or heart pumping.

## **Radionuclide Cystogram**

A nuclear medicine examination that tracks radio-labeled urine to detect vesicoureteral reflux or evaluate bladder emptying.

## **Radionuclide Generator**

A device that produces a short-lived radioactive substance from a longer-lasting radioactive substance. It supplies certain tracers for nuclear scans or treatment when needed and must be operated under strict safety and quality controls.

## **Radionuclide Heart Scan**

A nuclear medicine study that uses a radioactive tracer to evaluate myocardial blood flow, heart muscle viability, or ventricular function.

## **Radionuclide Imaging**

A scan that uses a small amount of radioactive tracer to show the structure or activity of an organ.

A camera detects the tracer, helping evaluate blood flow, bone, thyroid, kidney, heart, or cancer-related changes.

## **Radionuclide Imaging Or Scan**

A scan that detects radiation from a small amount of radioactive tracer given to a person. The tracer collects in selected organs or tissues, allowing clinicians to examine blood flow, structure, or function and identify some diseases.

### **Alternate Names:**

Scintigraphy; Nuclear Medicine Scan; Gamma Scan

## **Radionuclides**

Radioactive forms of atoms used in nuclear medicine. Attached to a suitable substance, they can show how an organ works on a scan or deliver radiation to selected diseased tissue during treatment.

## **Radionuclide Therapy**

Treatment that delivers radiation from inside the body using a radioactive substance directed toward selected tissue. It can treat some thyroid diseases, cancers, or painful bone lesions. The medicine, dose, precautions, and side effects depend on the target and the person's kidney, blood, and overall health.

## **Radionuclidic Purity**

The proportion of radioactivity in a preparation that comes from the intended radioactive atom rather than unwanted radioactive contaminants. High purity helps ensure the expected scan or treatment and limits unnecessary radiation to other tissues.

## **Radio-Opaque**

Blocking X-rays and appearing relatively white on a radiograph; radiopaque is the modern spelling.

## **Radiopaque**

Describing a material that attenuates X-rays strongly and therefore appears relatively white on a radiograph or CT image. Bone, metal, and some contrast agents are radiopaque.

---

## **Radiopharmaceutical**

A medicine containing a small amount of radioactive material. It can show how an organ or tissue works on a scan or deliver radiation to selected diseased cells; its distribution determines the information or treatment it provides.

## **Radiopharmaceutical Extravasation**

Leakage of a radioactive medicine from the vein into nearby tissue during an injection. It can reduce the accuracy of a scan and, with treatment doses, injure local tissue; staff assess and document the event.

## **Radiopharmaceutical Labeling Kit**

A prepared set of sterile ingredients used to attach a radioactive atom to a substance that travels to a particular organ or follows a body process. The finished product is tested before it is given for imaging.

## **Radiopharmaceutical Quality Control**

Testing performed to verify identity, radionuclidic purity, radiochemical purity, pH, sterility, endotoxin limits, and other release specifications before clinical administration. Requirements depend on the preparation and jurisdiction.

## **Radiopharmacy**

The preparation, testing, storage, and dispensing of medicines containing radioactive materials. Radiopharmacy requires accurate activity measurement, cleanliness, sterility checks, quality control, safe handling, and protection of workers and patients.

## **Radioprotective Agent**

A medicine or substance intended to reduce injury to healthy tissue from ionizing radiation. It may limit chemical damage or support cell repair, but its benefit and risks depend on the type, dose, and treatment setting.

## **Radio-Radial Delay**

A palpable asymmetry or latency between the right and left radial arterial pulses when examined simultaneously. It indicates proximal arterial obstruction affecting one subclavian artery, as seen in aortic

dissection, subclavian steal syndrome, or thoracic aortic aneurysm.

## **Radiosensitization**

Making cells more vulnerable to radiation damage. Some medicines or treatments are combined with radiotherapy to increase damage to cancer cells, while treatment planning aims to limit harm to nearby healthy tissue.

## **Radiosurgery**

Radiosurgery delivers highly focused radiation to a small target, usually in the brain or body, without making a conventional surgical incision.

## **Radiotherapy**

Treatment using high-energy radiation to damage and control abnormal cells, especially cancer. It may be given from outside or inside the body and can cause temporary or lasting effects in nearby healthy tissue.

## **Radiotherapy Bolus**

A layer of material placed on the skin during selected radiation treatments so more radiation reaches the skin surface. It is positioned carefully to deliver the planned dose to the target area.

## **Radiotherapy Dosage**

Radiotherapy dosage is the amount of ionizing radiation prescribed for a treatment target, usually divided into carefully planned fractions.

## **Radiotracer**

A small amount of radioactive material attached to a substance that travels through a specific body process. A scanner detects where it goes, helping show blood flow, organ function, bone activity, or selected disease-related changes.

## **Radioulnar Joint**

A joint between the radius and ulna bones of the forearm. It allows the hand and forearm to turn palm-up and palm-down.

---

## Radium

A naturally radioactive chemical element that releases penetrating radiation as it breaks down. Internal exposure can damage bone and other tissues and increase cancer risk, so handling requires strict safety controls.

## Radium-223

A radioactive medicine that behaves like calcium and collects in areas of increased bone activity. It is used for selected prostate cancers that have spread to bone and may cause low blood counts or other side effects.

## Radius

The forearm bone on the thumb side. It joins the elbow and wrist and rotates around the ulna, allowing the palm to turn up or down.

## Radius Fracture

A break in the radius, one of the two bones of the forearm, often occurring near the wrist after a fall onto an outstretched hand.

## Radon

A naturally occurring radioactive gas that can accumulate indoors and increase lung-cancer risk when inhaled over time.

## Ragged Red Fibers

An appearance seen in a muscle biopsy when abnormal mitochondria build up around muscle fibers. It supports the diagnosis of some mitochondrial muscle diseases but is not specific to one condition. The biopsy result must be interpreted with symptoms, blood tests, genetic testing, and other findings.

### Alternate Names:

RRF; Ragged-Red Muscle Fibers

## Raised Hemidiaphragm

A raised hemidiaphragm is elevation of one side of the diaphragm, caused by abdominal pressure, lung volume loss, phrenic-nerve dysfunction, or diaphragmatic disease.

## Rale

A rale, or crackle, is an abnormal discontinuous breath sound produced by fluid, collapsed airways, or opening of small airways during breathing.

## Rales

Discontinuous, adventitious non-musical respiratory lung sounds heard on auscultation, generated by the sudden explosive reopening of collapsed small airways and alveoli containing fluid or exudate; categorized into fine crackles (high-pitched, dry) and coarse crackles (low-pitched, wet).

### Alternate Names:

Crackles; Crepitations

## Raloxifene

A medicine that acts like estrogen in bones but blocks some estrogen effects elsewhere. It may reduce spine fractures and selected breast-cancer risk after menopause, but can increase blood-clot risk.

## Raloxifene Hydrochloride

Raloxifene hydrochloride is a selective estrogen-receptor modulator used to prevent or treat osteoporosis and reduce breast-cancer risk in selected postmenopausal women.

## Raltegravir Potassium

Raltegravir potassium is an HIV integrase inhibitor used with other antiretroviral medicines to suppress HIV replication.

## Raltitrexed

A chemotherapy medicine used for some colorectal cancers. It blocks a cell process needed to make DNA, slowing the growth of cancer cells; possible effects include diarrhea, mouth sores, low blood counts, tiredness, and liver problems.

## Ramipril

Ramipril is an angiotensin-converting-enzyme inhibitor used for hypertension, heart failure, and cardiovascular risk reduction; cough, hyperkalemia, renal-function changes, angioedema, and fetal toxicity are important risks.



---

## Rampant Caries

Severe tooth decay affecting many teeth or surfaces at once. It may be linked with frequent sugar, dry mouth, vomiting, poor oral hygiene, limited dental care, medicines, or other health problems.

## Ramsay Hunt Syndrome

Reactivation of the shingles virus near the facial nerve, causing severe ear pain, a blistering rash in or around the ear or mouth, and weakness on one side of the face. Hearing loss or dizziness may occur; early treatment can improve recovery.

## Ramsay Hunt Syndrome Type I

A rare neurological disorder characterized by progressive myoclonus epilepsy, intentional cerebellar tremor, progressive ataxia, and mild cognitive decline. Historically termed dyssynergia cerebellaris myoclonica; distinct from Ramsay Hunt syndrome type II (herpes zoster oticus).

### Alternate Names:

Dyssynergia Cerebellaris Myoclonica; Dentatorubral Tremor Syndrome

## Ramsay Hunt Syndrome Type II

Reactivation of latent varicella-zoster virus within the geniculate ganglion of the facial nerve (cranial nerve VII), presenting with the clinical triad of acute ipsilateral peripheral facial paralysis, otalgia, and a painful vesicular eruption on the external auditory canal, auricle, or soft palate.

### Alternate Names:

Herpes Zoster Oticus

## Ramucirumab

A monoclonal antibody that blocks vascular endothelial growth factor receptor 2 and is used in selected advanced cancers, including gastric, colorectal, lung, and liver cancers.

## Ramus

A ramus is a branch-like extension of a bone, nerve, artery, or other structure, such as the mandibular ramus.

## Random Allocation

Assignment of participants to study groups by chance to reduce selection bias and balance known and unknown factors.

## Randomised Controlled Trial

A study in which participants are assigned by chance to interventions or controls and outcomes are compared.

## Randomization Validation

Checking that a clinical trial's assignment process truly places participants into study groups by chance and is implemented as planned. This helps reduce selection bias and protects the reliability of comparisons between treatments.

## Randomized Controlled Trial

An experimental study in which participants are assigned by chance to different interventions or a control condition and followed for specified outcomes.

## Range Of Motion

The amount a joint can move in different directions. It may be measured by a healthcare professional or by the person moving alone; pain, swelling, stiffness, weakness, injury, or a shortened muscle can reduce it.

## Ranibizumab

An intravitreal monoclonal antibody fragment that inhibits vascular endothelial growth factor and treats neovascular macular degeneration, diabetic macular edema, retinal vein occlusion, and related retinal disease.

## Ranke Complex

The late, healed, fibrocalcified sequela of primary pulmonary tuberculosis, consisting of a calcified subpleural Ghon parenchymal focus and calcified ipsilateral draining hilar or mediastinal lymph nodes visible on chest radiography.

### Alternate Names:

Calcified Primary Tuberculous Complex; Healed

---

Ghon Complex

### Ranolazine

Ranolazine is an antianginal drug that inhibits the late sodium current in cardiac cells and reduces chronic angina without substantially lowering heart rate or blood pressure; QT prolongation and drug interactions require caution.

### Ranson Criteria

An older scoring system that estimates the severity and risk of death in acute pancreatitis using clinical and blood-test findings at admission and again after 48 hours. It is less commonly used now because newer scores and organ-failure assessment may be more practical.

### RANTES

RANTES, now called CCL5, is a chemokine that attracts immune cells and participates in inflammation, antiviral defense, and immune signaling.

### Ranula

A mucocele (mucus extravasation cyst) of the sublingual salivary gland, presenting as a blue-translucent, fluctuant swelling in the floor of the mouth, typically lateral to the lingual frenulum. A ranula results from trauma or obstruction of a sublingual gland duct, causing mucus accumulation in the sublingual space.

### Raoultella

Raoultella is a genus of gram-negative bacteria that can cause opportunistic urinary, respiratory, wound, or bloodstream infections.

### Rape

Nonconsensual sexual penetration or sexual violence; it is a serious crime and a medical, safeguarding, and trauma-care concern.

### Raphe

A seam-like line or ridge where structures join, such as the median raphe or raphe nuclei of the brainstem.

### Raphe Nuclei

The raphe nuclei are brainstem groups of neurons that produce serotonin and influence mood, pain, sleep, arousal, and autonomic functions.

### Rapid Eye Movement Sleep

A sleep stage with rapid eye movements, vivid dreaming, and active brain patterns. Most skeletal muscles are temporarily relaxed during it, while breathing and heart rate can become more variable.

#### Alternate Names:

REM Sleep; Paradoxical Sleep

### Rapid Eye Movement Sleep Behavior Disorder

Alternate name; see *REM sleep behavior disorder*.

### Rapid Gastric Emptying

Abnormally rapid movement of food from the stomach into the small intestine, sometimes called dumping syndrome.

### Rapid Heartbeat

Alternate name; see *Tachycardia*.

### Rapidly Progressive Glomerulonephritis

Severe inflammation of the kidney's filtering units that causes kidney function to worsen over days or weeks. It may cause blood or protein in urine, swelling, high blood pressure, or reduced urine.

### Rapid Plasma Reagin Test

A blood test that detects antibodies associated with syphilis. It is useful for screening and monitoring treatment, but false-positive and false-negative results occur, so reactive results need a confirmatory treponemal test.

### Rapid Sequence Induction

A method of giving anesthetic and muscle-relaxing medicines to place a breathing tube quickly while reducing the chance of vomit entering the lungs. It is used when urgent airway control is needed and requires preparation for difficult intubation.

---

## Rapid Sequence Intubation

An emergency airway management technique involving the near-simultaneous administration of a rapid-acting sedative induction agent (such as etomidate or propofol) and a neuromuscular blocking paralytic agent (succinylcholine or rocuronium) without intervening bag-mask ventilation, designed to minimize aspiration risk in an unfasted patient.

### Alternate Names:

RSI

## Rapid-Sequence Intubation

An emergency method for placing a breathing tube quickly after giving medicines that induce unconsciousness and temporarily relax the muscles. It is used when airway protection is urgent, with oxygen preparation, backup equipment, blood-pressure monitoring, and confirmation that the tube is correctly positioned.

## Rapid Shallow Breathing

Breathing that is faster than normal but moves only small amounts of air each time. It may occur with lung disease, pain, anxiety, metabolic illness, or tiring breathing muscles and can be a warning sign of respiratory failure.

## Rapid Strep Test

A test performed on a throat swab to detect Group A Streptococcus. It provides a rapid result, but a negative result may require confirmatory testing depending on the patient's age, symptoms, and local guidance.

## Rapp-Hodgkin Syndrome

A TP63-related ectodermal dysplasia characterized by abnormalities of the hair, teeth, nails, skin, and sweat glands, often with cleft lip or palate. It is also called Rapp-Hodgkin syndrome and requires individualized genetic and specialist care.

### Alternate Names:

Rapp-Hodgkin Type Ectodermal Dysplasia

## Rare Disease

A condition affecting relatively few people in a defined population; many rare diseases are genetic,

require specialized diagnosis, and may have limited evidence or treatment options.

## Rarefaction

A reduction in density or pressure, such as the low-pressure part of a sound wave. In medicine, understanding rarefaction helps explain ultrasound imaging, sound transmission, and pressure changes in tissues.

## Rasburicase

A recombinant urate-oxidase enzyme that converts uric acid to the more soluble allantoin. It is used in selected patients at risk of or experiencing tumor-lysis-related hyperuricemia. It can cause serious hypersensitivity or hemolysis in people with glucose-6-phosphate dehydrogenase deficiency and requires appropriate screening and monitoring.

## Rash

A visible change in skin color, texture, or surface, such as spots, bumps, blisters, or scaling. Causes include infection, allergy, medicines, inflammation, and other disease; rapid or severe rashes need assessment.

### Alternate Names:

Exanthem; Skin Eruption; Dermatitis

## Rasmussen Encephalitis

A rare, progressive, chronic immune-mediated neuro-inflammatory disorder occurring predominantly in children, characterized by unilateral hemispheric inflammation, drug-resistant focal motor seizures (frequently *epilepsia partialis continua*), progressive contralateral hemiparesis, cognitive deterioration, and progressive unilateral cortical atrophy on neuroimaging.

### Alternate Names:

Chronic Focal Encephalitis; Rasmussen Syndrome

## Rasp

A roughened surgical instrument used to file, shape, or smooth bone or other hard tissue.

---

## Ras Peptide

A small protein fragment related to Ras proteins, which help control cell growth and division. Changes in Ras signaling can contribute to cancer.

## RAST

An older name for a blood test that measures IgE antibodies directed against specific allergens. Modern laboratories usually use newer specific-IgE immunoassays; results show sensitization and must be interpreted with symptoms and exposure history.

### Alternate Names:

Radioallergosorbent Test; Specific IgE Test

## Rastelli Procedure

Open-heart surgery for certain complex birth defects involving the heart's major arteries, a hole between the lower chambers, and a narrowed route to the lungs. A patch directs blood from the left ventricle to the aorta, and a tube with a valve connects the right ventricle to the lung arteries. The tube may need later replacement.

### Alternate Names:

Rastelli Operation; Rastelli Repair

## Rat Bite Fever

A systemic bacterial infection caused by *Streptobacillus moniliformis* or *Spirillum minus*, transmitted through rat bites or contact with rat-contaminated food or water. It presents with relapsing fever, rash, polyarthritis, and lymphadenopathy. The characteristic rash is petechial or maculopapular on the extremities.

## Rat-Bite Fever

Alternate name; see *Rat Bite Fever*.

## Rathke Cleft Cyst

A benign, non-neoplastic, mucin-filled sellar or suprasellar epithelial cyst derived from remnants of the embryonic Rathke pouch, which may compress the optic chiasm (causing bitemporal hemianopia) or normal pituitary gland (producing hypopituitarism).

### Alternate Names:

RCC; Pituitary Cleft Cyst

## Rathke Pouch

An embryonic ectodermal diverticulum arising from the roof of the stomodeum (primitive oral cavity) that ascends toward the developing brain to form the anterior pituitary gland (adenohypophysis: pars distalis, pars intermedia, pars tuberalis); incomplete obliteration gives rise to craniopharyngiomas or Rathke cleft cysts.

## Ratio

A ratio compares two quantities by division, such as a measurement of risk, concentration, or body proportions.

## Rationalization

A psychological defense in which a person creates plausible explanations for behavior or feelings while avoiding the underlying motive or conflict.

## Rat Lungworm Disease

Infection caused by the rat lungworm parasite *Angiostrongylus cantonensis*, usually after eating contaminated raw or undercooked produce or animals. It can cause eosinophilic meningitis with headache, neck stiffness, or neurologic symptoms.

## RATS

Alternate name; see *Robotic-Assisted Thoracoscopic Surgery*.

## Ravuconazole

Ravuconazole is a triazole antifungal medicine studied and used in some settings for invasive fungal infections.

## Raw Egg

Raw eggs may contain *Salmonella* and are safest when properly refrigerated, handled hygienically, and cooked unless pasteurized products are used.

## Raynaud Phenomenon

Episodes in which small blood vessels in the fingers or toes suddenly narrow after cold or stress.

---

The digits may turn white, blue, then red and become numb or painful; severe or new symptoms may indicate another disease.

### **Raynaud's Disease**

The primary form of Raynaud phenomenon, in which small blood vessels in the fingers or toes narrow episodically without an associated underlying disease. Cold or emotional stress may trigger attacks that cause numbness, tingling, or pain and a characteristic change in color from white to blue and then red as blood flow returns.

### **Raynaud's Phenomenon**

Episodic constriction of small arteries in the fingers or toes, causing color changes, numbness, or pain after cold exposure or emotional stress.

#### **Alternate Names:**

Raynaud Disease; Raynaud Syndrome; Vasospastic Attack

### **Raynaud's Syndrome**

Episodes in which fingers or toes change color and become cold, numb, or painful after cold exposure or stress because small blood vessels narrow. It may occur alone or with autoimmune disease.

### **Raynaud Syndrome**

Episodes in which small blood vessels in the fingers or toes narrow after cold or stress, causing numbness, pain, and color changes that may progress from white to blue to red. It may occur alone or result from another disease; severe attacks can damage tissue.

### **Razor Bumps**

Inflamed bumps caused by hairs curling back into the skin after shaving or other hair removal. They are more common with coarse or curly hair and may cause itching, pain, dark marks, or scarring; avoiding very close shaving and using proper shaving technique can reduce recurrence.

### **RBC Count**

A measurement of the number of red blood cells in a volume of blood, interpreted with hemoglobin,

hematocrit, and red-cell indices to evaluate anemia or erythrocytosis.

### **RBC Indices**

Measurements describing the size, hemoglobin content, and variation of red blood cells. They help classify anemia and interpret a complete blood count alongside hemoglobin and other results.

### **RBC Nuclear Scan**

An imaging test using a small amount of the person's blood labeled with a radioactive tracer. It can help locate ongoing bleeding in the digestive tract or assess blood flow through the heart.

### **RBC Urine Test**

A urine examination detecting or counting red blood cells, used to evaluate hematuria from urinary-tract infection, stones, glomerular disease, tumors, trauma, or other causes.

### **RDA**

Recommended dietary allowance, the daily nutrient intake sufficient for nearly all healthy people in a defined group.

#### **Alternate Names:**

Recommended Dietary Allowance; Recommended Daily Allowance

### **Reabsorption**

The transport of a substance from a body cavity or tubular fluid back into tissue or the bloodstream, such as renal tubular recovery of water and filtered solutes.

### **Reaction**

A response of the body or mind to a stimulus, illness, medicine, treatment, or event. Examples include an allergic reaction, an immune reaction, a treatment response, and a psychological reaction; the cause and severity determine its importance.

### **Reactivation Tuberculosis**

Active tuberculosis developing when a previously contained infection becomes active again, often

---

years later. It commonly affects the upper lungs and may cause a long-lasting cough, coughing blood, fever, night sweats, weight loss, and spread to others through the air.

### **Reactive Arthritis**

Painful inflammation of joints that develops during or after certain intestinal or genital infections. It may also cause eye inflammation, skin changes, or urinary symptoms, and the triggering infection may already have cleared.

#### **Alternate Names:**

Reiter Syndrome; Post-Infectious Arthritis

### **Reactive Attachment Disorder**

A rare childhood condition in which a child persistently avoids or barely responds to comfort from caregivers. It is linked to severe neglect or unstable care early in life and may involve sadness, irritability, limited trust, and difficulty forming relationships.

### **Reactive Gastropathy Pathology**

Gastric mucosal injury characterized by foveolar hyperplasia, edema, smooth-muscle fibers in the lamina propria, vascular congestion, and relatively little active inflammation. Chemical injury, bile reflux, and medicines are common associations.

### **Reactive Hypoglycemia**

A fall in blood sugar occurring within a few hours after eating. It may cause shakiness, sweating, hunger, weakness, dizziness, or confusion; repeated episodes need assessment to confirm low sugar and identify the cause.

### **Reactive Oxygen Species**

Unstable oxygen-containing molecules produced during normal cell activity, infection, or inflammation. Small amounts can help cells signal and defend against germs, but excess amounts can damage cell membranes, proteins, and DNA.

### **Reactive Perforating Collagenosis**

A skin condition in which damaged collagen from deeper skin is pushed through the surface. It causes small, firm, very itchy bumps with a central plug,

often on the arms, legs, or hands, and may be associated with diabetes or kidney disease.

### **Reality Testing**

The ability to recognize whether perceptions, thoughts, and beliefs match shared reality. Clinicians assess it during a mental-status examination; impaired reality testing can occur with psychosis, severe mood episodes, delirium, or some brain disorders.

### **Rebound Effect**

Worsening or return of symptoms after stopping or reducing a treatment that had been suppressing them.

### **Rebound Headache**

Alternate name; see *Medication overuse headaches*.

### **Rebound Insomnia**

Return or worsening of insomnia after stopping a sleep medicine.

### **Rebound Tenderness**

A sharp increase in stomach pain when pressure on the belly is suddenly released, which signals that the inner lining of the abdomen is inflamed.

### **Recalcitrant**

Resistant to treatment, control, healing, or change; the term often describes a persistent disease or infection.

### **Receiver Operating Characteristic**

A graph showing how a diagnostic test's ability to detect disease changes as its cutoff is changed. It compares true-positive and false-positive results; the area under the curve summarizes overall discrimination, not clinical usefulness by itself.

### **Recent Memory**

Recent memory is the ability to recall information or events from the near past and can be impaired by aging, delirium, brain disease, depression, or medicines.



---

## Receptor

A protein on or inside a cell that recognizes a specific signal, such as a hormone, nerve chemical, or medicine, and starts or changes a cell response.

## Receptor Binding

The attachment of a signaling molecule, medicine, or other ligand to a receptor on or inside a cell. Binding can activate, block, or change the receptor and its downstream cellular response.

## Receptor Desensitization

A reduced cell response after a receptor is stimulated continuously or repeatedly. The change can develop quickly and helps prevent excessive signaling, but it may also reduce the effect of a medicine or natural body signal.

## Receptor Downregulation

A decrease in the number or sensitivity of cell receptors after a signal stimulates them for a long time. The cell may remove receptors from its surface or break them down, becoming less responsive to the signal or medicine.

## Receptors

Proteins or structures on or inside cells that recognize specific signals and start responses such as growth, movement, or hormone action.

## Receptor Tyrosine Kinases

Cell-surface proteins that receive signals from hormones and growth factors. When switched on, they activate internal pathways controlling growth, survival, movement, and repair; abnormal activity can contribute to cancer and other disease.

## Receptor Upregulation

An increase in the number or sensitivity of cell receptors after prolonged blocking of a signal or reduced stimulation. This adjustment can make a cell respond more strongly when the signal or medicine is later present.

## Recession Of Medial Rectus Muscle

An eye operation that moves the medial rectus muscle backward to weaken its pull and correct certain inward eye deviations.

## Recessive

Describing a gene change whose effect usually appears only when a person has two altered copies, one inherited from each parent. A person with one copy may carry it without having the condition.

## RECIST

Standard rules for judging how a solid tumor changes on scans during treatment. Using measured lesions, they classify disease as completely gone, partly reduced, stable, or worsened; the categories support research and clinical decisions but do not replace overall assessment.

## Recognizing Medical Emergencies

Recognizing a medical emergency includes identifying severe breathing difficulty, chest pain, stroke signs, major bleeding, seizure, collapse, poisoning, or altered consciousness and calling emergency services.

## Recognizing Teen Depression

Recognizing depression in adolescents involves looking for persistent low mood or irritability, loss of interest, sleep or appetite changes, poor concentration, hopelessness, or self-harm thoughts.

## Recombinant

Produced by combining genetic material from different sources or by using engineered DNA in a cell or organism.

## Recombinant DNA

DNA made by joining genetic material from different sources. It is used in biotechnology to produce medicines such as insulin and growth hormone, develop some vaccines, and create vehicles for gene therapy.

---

## Recombinant Protein

A protein produced by putting its gene into a laboratory cell and growing the cell so it makes the protein. This method supplies some hormones, vaccines, antibodies, enzymes, and research materials.

## Recombinant Tissue Plasminogen Activator

A laboratory-made form of tissue plasminogen activator that helps convert plasminogen to plasmin and dissolve blood clots. It is used in selected emergencies, such as some ischemic strokes, heart attacks, or pulmonary emboli, when benefits outweigh bleeding risk.

### Alternate Names:

rtPA; Alteplase; Recombinant tPA

## Recombinase

An enzyme that cuts, joins, or rearranges DNA segments. It helps cells exchange genetic material, repair some DNA damage, and assemble the gene parts needed to make a varied immune response.

## Recombinational Repair

A way cells repair a serious DNA break by using a matching, undamaged DNA copy as a guide. It restores genetic information accurately; inherited problems with this repair can increase cancer risk and treatment sensitivity.

## Recommended Daily Allowance

The Recommended Dietary Allowance is the average daily nutrient intake sufficient for nearly all healthy people in a specified age and sex group.

## Recommended Dietary Allowance

The average daily intake expected to meet the nutrient needs of nearly all healthy people in a defined group.

### Alternate Names:

RDA

## Record

A health record is a documented account of a person's history, findings, tests, diagnoses, treatments,

decisions, and outcomes.

## Recovery Position

A side-lying position used for an unconscious but normally breathing person to help keep the airway open and reduce aspiration risk.

## Recreation Therapy

Recreation therapy uses adapted games, arts, exercise, and leisure activities to support physical, emotional, cognitive, or social functioning.

## Recrudescence

Recurrence of symptoms or infection after apparent improvement, often from persistence of the original organism or disease process.

## Rectal

Rectal means relating to the rectum, the final section of the large intestine before the anal canal.

## Rectal Biopsy

Removal of a small piece of tissue from the rectum for examination under a microscope. It can help investigate missing nerve cells, inflammation, infection, abnormal growth, or other rectal disease.

## Rectal Bleeding

Passage of bright-red or dark blood from the rectum, caused by hemorrhoids, fissures, polyps, inflammatory bowel disease, diverticular disease, or cancer. Persistent, heavy, or unexplained bleeding requires evaluation.

### Alternate Names:

Hematochezia; Lower Gastrointestinal Bleeding

## Rectal Cancer

Cancer developing in the rectum, the final part of the large intestine. Symptoms may include bleeding, altered bowel habits, pain, or weight loss; treatment may combine surgery, radiation, chemotherapy, immunotherapy, or targeted therapy.

### Alternate Names:

Rectal Carcinoma; Adenocarcinoma of the Rectum

---

## Rectal Carcinoma

Rectal carcinoma is cancer arising in the rectum and may cause bleeding, altered bowel habits, pain, or anemia.

## Rectal Culture

Laboratory culture of a rectal specimen to detect selected bacterial pathogens or colonization, depending on the clinical indication.

## Rectal Disease

A disorder affecting the rectum, including inflammation, infection, bleeding, obstruction, prolapse, functional disease, or neoplasm.

## Rectal Fistula

A rectal fistula is an abnormal tract connecting the rectum with skin or another organ, often causing drainage, infection, or passage of stool through an abnormal route.

## Rectal Hemorrhage

Bleeding from the rectum, which may result from hemorrhoids, fissure, inflammation, ischemia, polyps, cancer, trauma, or other gastrointestinal disease.

## Rectal Inflammation

Alternate name; see *Proctitis*.

## Rectal Itching

Alternate name; see *Anal itching*.

## Rectal Mucositis

Rectal mucositis is inflammation and injury of the rectal lining, often after radiation, chemotherapy, infection, inflammatory disease, or trauma.

## Rectal Necrosis

Rectal necrosis is death of rectal tissue from inadequate blood flow, severe inflammation, infection, pressure, or another destructive process.

## Rectal Neoplasm

An abnormal growth in the rectum that may be

noncancerous, precancerous, or cancerous. It can cause bleeding, altered bowel habits, pain, anemia, or no symptoms and may require examination and biopsy.

## Rectal Obstruction

Impaired passage through the rectum caused by fecal impaction, stricture, tumor, foreign body, inflammation, or functional outlet disorder.

## Rectal Perforation

A tear or hole through the rectal wall that can release bowel contents and cause peritonitis, sepsis, or severe bleeding.

## Rectal Polyposis

Rectal polyposis is the presence of multiple polyps in the rectum, which may be sporadic or part of an inherited syndrome and can increase cancer risk.

## Rectal Polyps

Alternate name; see *Colon Polyps*.

## Rectal Prolapse

The rectum slipping downward and partly or completely coming through the anus. It may cause a visible bulge, bleeding, mucus, discomfort, or difficulty controlling or passing stool and can result from weakened pelvic support, straining, or nerve problems.

## Rectal Prolapse Repair

A surgical or procedural treatment that restores a prolapsed rectum to its normal position and secures or removes affected tissue according to the prolapse and patient factors.

## Rectal Route

The rectal route administers a medicine or fluid through the anus into the rectum, where it may act locally or be absorbed systemically.

## Rectal Stenosis

Abnormal narrowing of the rectal lumen caused by inflammation, scarring, surgery, radiation, ischemia,

---

trauma, or neoplasm.

### Rectal Tenesmus

Rectal tenesmus is a persistent or urgent feeling of needing to pass stool despite an empty or nearly empty rectum, caused by inflammation, infection, mass, or motility disorder.

### Rectal Ulcer

Alternate name; see *Solitary rectal ulcer syndrome*.

### Rectocele

Bulging of the rectum into the back wall of the vagina because supporting tissues have weakened. It may cause pelvic pressure, difficulty emptying the bowel, or a vaginal bulge.

### Rectosigmoid Junction

The point where the sigmoid colon joins the rectum.

### Rectosigmoid Neoplasm

A rectosigmoid neoplasm is an abnormal growth where the sigmoid colon joins the rectum and may be benign or malignant.

### Rectouterine Pouch

The rectouterine pouch, or pouch of Douglas, is the peritoneal recess between the uterus and rectum where fluid or disease may collect.

### Rectovaginal Fistula

Abnormal connection between the rectum and vagina causing passage of stool or gas through the vagina. Causes include obstetric injury particularly third and fourth degree perineal tears, Crohn disease, radiation therapy, and surgical complications. Presents with vaginal discharge of stool, flatus, or recurrent vaginal infections.

### Rectum

The final straight section of the large intestine, approximately 12-15 cm long, extending from the sigmoid colon to the anal canal. The rectum stores feces before defecation.

### Rectus Abdominis

The rectus abdominis is a paired abdominal muscle that flexes the trunk, stabilizes the pelvis, and helps compress abdominal contents.

### Rectus Abdominis Muscle

The paired vertical muscles at the front of the abdomen, commonly called the “six-pack.” They bend the trunk forward, stabilize the pelvis, and help increase pressure inside the abdomen; stretching can cause separation during pregnancy or after major weight change.

### Rectus Abdominis Sheath

A fibrous aponeurotic envelope formed by the abdominal wall muscles and surrounding the rectus abdominis muscle. It contains the superior and inferior epigastric vessels and related nerves.

### Rectus Sheath Hematoma

A collection of blood within the sheath surrounding the rectus abdominis muscle, often caused by muscle strain, coughing, trauma, anticoagulation, or vessel injury. It may cause localized abdominal pain and a tender mass.

### Recur

To recur means to return or happen again, as symptoms, disease, infection, or a tumor after a period of improvement or treatment.

### Recurrence

The return of a disease after it improved or seemed to disappear. Cancer may return at the original site, nearby lymph nodes, or another part of the body; recurrence does not always mean that treatment is no longer possible.

### Recurrence-Free Survival

The length of time after treatment during which a disease does not return. In cancer studies, it usually ends when cancer is found again or when a person dies, depending on the study rules. It measures control of disease, not a guarantee that treatment has cured it.

---

## Recurrence Risk

The chance that a disease or condition will return after treatment or occur again in a future pregnancy or family member. The estimate depends on the condition, previous history, test results, inheritance pattern, and other risk factors.

## Recurrent

Happening again after a period of improvement, remission, or apparent absence. A recurrent condition may return at the original site or elsewhere; examples include repeated urinary infections, repeated seizures, or cancer returning after treatment.

## Recurrent Breast Cancer

Breast cancer that returns after a period when tests found no detectable disease. It may recur near the original site or elsewhere and is assessed with examination, imaging, biopsy, and other tests.

## Recurrent or Metastatic Breast Cancer

Breast cancer that returns after treatment or has spread from the breast to distant organs or tissues. Assessment and treatment depend on the site of disease, hormone-receptor and HER2 status, prior therapy, symptoms, and the person's goals.

## Recurrent Caries

Tooth decay that develops beneath or beside an existing filling, crown, or other restoration. It may result from leaking edges, plaque, frequent sugar, or dry mouth and can cause sensitivity, pain, or further tooth damage.

## Recurrent Disease

Return of a disease after a period of improvement, remission, or apparent absence, either at the original site or elsewhere.

## Recurrent Early Pregnancy Loss

Repeated miscarriages occurring early in pregnancy. Evaluation may consider uterine anatomy, chromosomal factors, antiphospholipid syndrome, endocrine or metabolic disease, parental genetics, and other causes, although no cause is found in many cases.

## Recurrent Fractures

Repeated bone fractures, suggesting persistent trauma, osteoporosis, bone fragility, metabolic disease, tumor, or an inherited skeletal disorder.

## Recurrent Laryngeal Nerve

A branch of the vagus nerve supplying most intrinsic muscles of the larynx; injury can cause hoarseness.

## Recurrent Laryngeal Nerve Palsy

A peripheral cranial neuropathy resulting from surgical injury (e.g., during thyroidectomy), trauma, or mediastinal tumor compression affecting the recurrent laryngeal branch of the vagus nerve (CN X), causing ipsilateral vocal cord immobility, hoarseness, and loss of airway protective reflexes.

### Alternate Names:

RLN Palsy; Vocal Cord Paralysis

## Recurrent Oral Ulcers

Alternate name; see *Canker sore*.

## Recurrent Pregnancy Loss

Two or more miscarriages. Possible causes include chromosome changes, an unusual shape of the uterus, hormone or thyroid problems, antiphospholipid syndrome, or other medical conditions, although no cause is found in many people. Evaluation is individualized and may identify treatment that improves the chance of a future healthy pregnancy.

### Alternate Names:

Recurrent Miscarriage; Habitual Abortion; RPL

## Recurrent Prostate Cancer

Alternate name; see *Prostate cancer recurrence*.

## Recurrent Respiratory Papillomatosis

A disease in which wart-like growths caused by human papillomavirus develop in the larynx or other breathing passages. The growths can cause hoarseness, noisy breathing, airway narrowing, or repeated recurrence.

---

## Recurrent Urinary Tract Infection

Repeated urinary tract infections separated by periods without infection. They may affect the bladder or kidneys and can be related to bacteria that persist, incomplete bladder emptying, stones, urinary blockage, or other urinary-tract problems.

## Red Birthmarks

Red birthmarks are vascular skin marks present at or soon after birth, ranging from harmless capillary marks to lesions requiring monitoring or specialist treatment.

## Red Blood Cell

A blood cell that carries oxygen from the lungs to the body's tissues. It contains hemoglobin, has a flattened, indented shape, and lacks a nucleus, which helps it bend through small blood vessels. Red blood cells also carry some carbon dioxide back to the lungs and help keep the blood's acid level balanced. They are made mainly in the bone marrow and normally live for about 120 days.

## Red Blood Cell Casts

Tube-shaped groups of red blood cells seen in urine. They form inside the kidney's small tubes and strongly suggest bleeding or inflammation in the kidney's filtering units, such as glomerulonephritis or vasculitis.

### Alternate Names:

RBC Casts

## Red Blood Cell Labeling

A nuclear-medicine technique in which a person's red blood cells are marked with a small amount of radioactive material. Scans can then show ongoing digestive-tract bleeding or measure blood flow and heart function.

## Red Bone Marrow

Bone marrow that makes red blood cells, white blood cells, and platelets. It is found mainly in certain bones and in the spongy parts of many bones.

## Red Cell Distribution Width

A standard automated complete blood count parameter that measures the degree of variation in red blood cell volume (anisocytosis); an elevated RDW aids in distinguishing iron deficiency anemia (high RDW) from uncomplicated thalassemia trait (normal RDW).

### Alternate Names:

RDW; Erythrocyte Anisocytosis Index

## Red Cell Indices

Standardized automated complete blood count parameters derived from hemoglobin, hematocrit, and erythrocyte count that quantify red blood cell physical properties: mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC).

## Red Eye

Redness of the white part of the eye caused by irritation, dryness, allergy, infection, injury, inflammation, or increased surface blood flow.

## Red Flag

A symptom, sign, or test result suggesting a potentially serious condition that requires urgent assessment or escalation of care. Red flags are context-specific and should not be used as a substitute for clinical judgment.

## Red Hepatization

The second pathological stage of acute lobar pneumonia (occurring 2 to 4 days after infection), in which the affected lung lobe becomes consolidated, heavy, and firm ('liver-like') due to massive filling of alveolar spaces with packed erythrocytes, neutrophils, and fibrin strands.

### Alternate Names:

Second Stage of Lobar Pneumonia; Consolidation Stage

## Red Infarction

Tissue death caused by loss of blood flow in which blood also leaks into the damaged area, making it dark red. It is more likely in organs with a second blood supply or when a vein is blocked.



---

**Alternate Names:**  
Hemorrhagic Infarction

### Red Man Syndrome

An adverse, non-IgE-mediated pseudoallergic hypersensitivity reaction caused by rapid intravenous infusion of vancomycin, in which direct degranulation of mast cells and basophils triggers massive histamine release, producing flushing, erythema, and pruritus of the face, neck, and upper torso.

**Alternate Names:**  
Vancomycin Flushing Reaction

### Red-Man Syndrome

An acute, non-immunologic (anaphylactoid) rate-dependent adverse drug reaction caused by rapid intravenous infusion of vancomycin, mediated by direct mast cell and basophil degranulation with histamine release, presenting with erythematous flushing and pruritus of the face, neck, and upper torso.

**Alternate Names:**  
Vancomycin Flushing Syndrome; Histaminoid Infusion Reaction

### Red Nucleus

A group of nerve cells in the midbrain that helps coordinate movement through connections with the cerebellum and spinal cord. Damage to nearby pathways can cause weakness, tremor, or abnormal movement.

### Reductase Deficiency

A disorder or laboratory finding caused by inadequate activity of a reductase enzyme, impairing a specific metabolic or hormone pathway.

### Reduction

Restoration of a displaced fracture or joint to better alignment, or decrease in amount, size, or activity.

### Reduction Division

The first division of meiosis, when a reproductive cell's chromosome number is reduced by half. It produces cells with one chromosome from each pair

and helps create genetic variation in eggs and sperm.

### Reduction Mammoplasty

Surgery that removes breast tissue, fat, and excess skin to reduce breast size and relieve symptoms such as neck, shoulder, or back discomfort; it may also reshape the breasts.

### Redundant Mitral Valve

Excess leaflet tissue of the mitral valve, most commonly the posterior leaflet, which may prolapse into the left atrium during systole. Often associated with myxomatous degeneration and mitral valve prolapse. May be asymptomatic or cause mild mitral regurgitation detected incidentally on echocardiography.

### Reduviidae

A family of insects including assassin bugs; some species transmit pathogens such as the parasite causing Chagas disease.

### Reed-Sternberg Cells

Large abnormal immune cells derived from B lymphocytes and commonly found in classic Hodgkin lymphoma. Their appearance in a tissue sample helps pathologists confirm that diagnosis.

### Reexpansion Pulmonary Edema

A sudden buildup of fluid in a lung after it rapidly expands from a collapsed state, often after drainage of a large or long-standing pneumothorax. It can cause severe breathlessness and low oxygen and needs urgent treatment.

### Refeeding Syndrome

Dangerous changes in body salts and fluid balance when food is restarted after severe or prolonged undernutrition. Low phosphate, potassium, or magnesium can cause weakness, swelling, abnormal heart rhythms, breathing failure, or death and requires careful medical monitoring.

### Reference Interval

The range of results usually found in a selected healthy reference group, often covering the middle

---

95 percent. A result outside it does not by itself prove disease and must be interpreted with symptoms, history, and the laboratory's own method.

### **Referral**

A request for assessment or treatment by another healthcare professional or service. Referrals help patients access specialized expertise, tests, procedures, rehabilitation, or coordinated care for a particular concern.

### **Referred Otalgia**

Ear pain felt in an anatomically normal ear caused by sensory nerve sharing between the ear and distant head and neck structures via cranial nerves V3 (auriculotemporal nerve), VII (facial), IX (glossopharyngeal - Jacobson nerve), X (vagus - Arnold nerve), and upper cervical nerves C2-C3; mandates careful examination of the pharynx and larynx for occult malignancy.

### **Referred Pain**

Pain felt in an area away from the tissue causing it because nearby or connected nerves share pathways to the brain. Examples include shoulder pain from irritation below the diaphragm or arm and jaw pain during a heart attack.

### **Reflex**

A rapid, automatic response to a stimulus, such as pulling the hand away from heat. Reflex testing helps assess pathways involving the nerves, spinal cord, and brain.

### **Reflex Action**

A rapid, automatic response to a stimulus that occurs through a nerve pathway called a reflex arc. Reflexes protect the body, maintain posture, and provide information about nerve and spinal-cord function.

### **Reflex Anoxic Seizure**

A brief fainting episode that can look like a seizure after sudden pain, fright, or emotional stress. The heart slows briefly, the person becomes pale and unresponsive, and jerking may occur before rapid

recovery.

### **Reflex Arc**

The nerve pathway that produces a rapid, automatic response. It includes a sensor, a sensory nerve, a processing center in the spinal cord or brain, a motor nerve, and the muscle or gland that responds.

### **Reflex Epilepsy**

Reflex epilepsy is epilepsy in which specific stimuli such as flashing light, sound, reading, or touch reliably trigger seizures.

### **Reflexes**

Automatic, rapid responses to stimuli carried through nerve pathways called reflex arcs. Testing them helps assess sensory nerves, motor nerves, the spinal cord, and parts of the brain.

### **Reflux**

The backward flow of a body fluid, especially stomach contents into the esophagus or urine from the bladder toward the kidneys. Symptoms and risks depend on the type, frequency, and underlying cause.

### **Reflux Esophagitis Pathology**

Mucosal injury associated with gastroesophageal reflux, often showing basal-zone hyperplasia, elongation of lamina-propria papillae, intercellular edema, and variable eosinophils or neutrophils.

### **Reflux Laryngitis**

Irritation or inflammation of the larynx attributed to reflux of stomach contents into the throat or upper airway. Hoarseness, throat clearing, cough, globus sensation, or throat discomfort may occur, although other causes of these symptoms should also be assessed.

### **Reflux Nephropathy**

Renal scarring and chronic kidney damage caused by vesicoureteral reflux, recurrent urinary infection, or both. It may lead to hypertension, proteinuria, impaired growth, and chronic kidney disease.

---

## Refraction

The bending of light as it passes through the cornea, aqueous humour, lens, and vitreous humour. In an eye examination, refraction determines the optical power needed to focus a clear image on the retina; refractive power is measured in dioptres.

## Refractive Disorder

A vision problem caused by the eye not focusing light correctly on the retina. Nearsightedness, farsightedness, astigmatism, and presbyopia are common refractive disorders and may be corrected with glasses, lenses, or surgery.

## Refractive Error

A refractive error occurs when the eye does not focus light correctly on the retina, causing nearsightedness, farsightedness, astigmatism, or presbyopia.

## Refractive Errors And Astigmatism

Refractive errors are vision disorders caused by the eye's inability to properly focus light onto the retina, resulting in blurred vision.

## Refractive Surgery

Eye surgery that changes the shape of the cornea or another part of the eye to reduce short-sightedness, long-sightedness, or astigmatism. It may reduce dependence on glasses but can cause dry eyes, glare, infection, or imperfect vision.

## Refractometer

A refractometer measures how much light bends through a substance and is used for fluid concentration, optical properties, or eye refraction.

## Refractometry

Measurement of the eye's refractive power to determine the optical correction needed for myopia, hyperopia, astigmatism, or presbyopia.

## Refractory

Resistant to a treatment or not responding adequately to it. The term may refer to cancer or other diseases.

## Refractory Anemia

An older term for anemia that does not improve with usual treatment. It may refer to a myelodysplastic syndrome or another cause, so evaluation looks for blood-cell disorders, bleeding, nutritional problems, kidney disease, and other conditions.

## Refractory Cancer

Refractory cancer is cancer that does not respond adequately to a treatment or begins progressing during treatment.

## Refractory Disease

Disease that does not improve adequately with appropriate standard treatment or that continues to worsen despite therapy. The term does not mean that every possible treatment has failed.

## Refractory Period

The brief recovery time after a nerve or muscle cell sends an electrical signal. During this time it cannot, or can only partly, send another signal, which helps control the speed and direction of nerve and heart activity.

## Refrigerant Poisoning

Poisoning from inhaling or contacting a refrigerant gas or liquid. Exposure can cause dizziness, abnormal heart rhythms, suffocation, frostbite, chemical burns, or lung injury.

## Refsum Disease

A rare inherited disorder in which a fatty substance builds up in the body. It may cause night blindness, loss of smell, nerve damage, poor coordination, hearing loss, and scaly skin.

## Regadenoson Pharmacological Stress

A medicine used during a heart scan to widen the heart's arteries when a person cannot exercise adequately. It can cause flushing, headache, chest discomfort, shortness of breath, or abnormal heart rhythms.

---

## Regeneration

Growth or replacement of damaged or lost cells and tissues, helping restore structure or function after injury.

## Regimen

A planned schedule for medicines, treatments, diet, exercise, or other healthcare steps. It states what to do, how much or how often, and for how long, and may be changed according to response or side effects.

## Regional

Relating to a particular body area, geographic area, or localized distribution rather than the whole body.

## Regional Anesthesia

Numbing a larger body area by injecting local anesthetic near the nerves supplying it. The person is usually awake, although sedation may be added; the block can provide numbness during surgery and pain relief afterward.

### Alternate Names:

Conduction Anesthesia; Nerve Block Anesthesia

## Regional Chemotherapy

Treatment that delivers anticancer medicine directly to a body region or organ to produce a high local concentration while reducing, but not eliminating, exposure elsewhere. It is suitable only for selected cancers and carries procedure-related and medicine-related risks.

## Regional Enteritis

An older name for Crohn disease, a chronic inflammatory bowel disease that can affect any part of the digestive tract.

## Regional Ileitis

An older name for Crohn disease affecting mainly the terminal ileum. It causes long-term inflammation that may lead to abdominal pain, diarrhea, weight loss, narrowing, fistulas, or nutritional problems.

## Regulation Of Health Professions

Legal and professional systems that govern health-care workers' education, licensing, permitted duties, conduct, and accountability. Regulation aims to protect patients, maintain competence, support ethical care, and address unsafe practice.

## Regulatory Approval

Authorization by a competent regulatory authority for a medicine, device, or procedure to be marketed or used for specified purposes after review of evidence.

## Regulatory Authority

A regulatory authority is a government or official body that sets and enforces standards for medicines, devices, facilities, professionals, or public health.

## Regulatory Gene

A gene that controls when, where, or how strongly another gene is used. Changes in regulatory genes or their control regions can alter development, cell growth, metabolism, or disease risk.

## Regulatory T Cells

Immune cells that limit excessive immune reactions and help prevent attacks against the body's own tissues. They develop mainly in the thymus or from other T cells in tissues. Problems with their function can contribute to autoimmune disease.

## Regurgitation

Backward flow of blood through a heart valve that does not close fully. Mild leakage may cause no symptoms; more severe leakage can enlarge or weaken the heart and cause breathlessness, tiredness, or swelling.

## Rehab

A structured rehabilitation program that helps restore function, independence, participation, or quality of life after illness or injury.

## Rehabilitation

A coordinated process that helps a person restore or

---

adapt physical, cognitive, psychological, and social function after illness, injury, surgery, or disability.

### **Rehabilitation Clinic**

An outpatient facility where a team helps people regain or adapt movement, communication, thinking, daily activities, or independence after illness, injury, surgery, or disability. Services may include physical, occupational, speech, or other rehabilitation therapies.

### **Rehabilitation Engineering**

The use of engineering and technology to improve independence after illness or disability. It includes artificial limbs, braces, wheelchairs, communication aids, home adaptations, and devices that help control the environment.

### **Rehabilitation Motivation**

A person's willingness and ability to take part in rehabilitation and work toward functional goals. Pain, mood, fatigue, cognition, support, confidence, and the expected benefit can all influence participation.

### **Rehabilitation Potential**

The expected capacity to improve function, independence, participation, or quality of life through rehabilitation. It depends on diagnosis, prognosis, cognition, motivation, comorbidity, social support, and environmental resources.

### **Rehabilitation Unit**

A hospital unit for people who need coordinated medical care and intensive therapy after a serious illness or injury, such as stroke, brain or spinal-cord injury, or major surgery. The team works to restore safe movement, daily skills, communication, and independence.

### **Rehydration**

Replacement of water and electrolytes lost through dehydration, commonly using oral fluids or intravenous treatment when necessary.

### **Reid Index**

A microscope measurement comparing the thickness of mucus-making glands with the wall of a large airway. A higher value may be seen in chronic bronchitis, but it is not used alone to diagnose the condition.

#### **Alternate Names:**

Bronchial Gland-to-Wall Ratio; Reid Histopathologic Index

### **Reifenstein Syndrome**

An inherited disorder of androgen action, usually involving partial androgen insensitivity in an individual with a 46,XY chromosome pattern.

### **Reiki**

Reiki is a complementary practice involving light touch or hands near the body. Evidence that it treats disease is limited.

### **Reinfection**

A new episode of infection by the same or a related pathogen after the original infection has resolved. Reinfection is distinct from relapse, in which the original infection persists or returns from an incompletely eradicated focus.

### **Reinke Edema**

Chronic polypoid degeneration and diffuse, gelatinous mucoid edema within the superficial lamina propria (Reinke's space) of both true vocal cords, strongly associated with chronic cigarette smoking and vocal strain, presenting with progressive hoarseness and a deepened voice.

#### **Alternate Names:**

Polypoid Corditis; Diffuse Vocal Fold Edema

### **Reinnervation**

Restoration of nerve supply to a muscle or tissue after nerve repair, regeneration, or collateral nerve growth.

### **Rejection**

An immune response in which the body recognizes transplanted tissue or an organ as foreign and at-

---

tacks it. Rejection can damage or destroy the transplant, may be sudden or gradual, and is monitored and treated with immune-suppressing medicines.

**Alternate Names:**

Allograft Rejection; Graft Rejection

**Relapse**

Return of a disease or its signs and symptoms after a period of improvement or remission. The term is used for cancer, infections, mental-health and substance-use disorders, and other conditions; its precise meaning depends on the disease.

**Relapse-Free Survival**

The time from completion of treatment to recurrence of the cancer or death from any cause.

**Relapsing Fever**

An acute febrile illness caused by *Borrelia* species, characterized by recurrent episodes of fever separated by afebrile intervals.

**Relapsing Polychondritis**

A rare immune disease in which episodes of inflammation damage cartilage. It commonly affects the ears, nose, eyes, joints, or windpipe and may cause pain, swelling, hearing or breathing problems; airway involvement can be life-threatening.

**Relapsing-Remitting Multiple Sclerosis**

A form of multiple sclerosis in which neurological symptoms appear in attacks and then improve partly or completely for periods of time. Symptoms may include vision changes, numbness, weakness, balance problems, or fatigue. Treatment can reduce relapses and new nerve damage, although the course varies between people.

**Alternate Names:**

RRMS; Relapsing-Remitting MS

**Relative Afferent Pupillary Defect**

Unequal response of the pupils because one eye or optic nerve sends less light information to the brain than the other. During a swinging-light test, both pupils may appear to dilate when the light moves to

the affected eye; it can indicate retinal or optic-nerve disease.

**Alternate Names:**

RAPD; Marcus Gunn Pupil

**Relative Biological Effectiveness**

A comparison of how strongly two types of radiation produce the same biological effect at the same absorbed dose. It depends on the radiation type, dose, tissue, and biological outcome.

**Relative Bradycardia**

A physical finding in which the heart rate fails to increase appropriately in response to fever (sphygmothermic dissociation), where pulse rises by less than 10 beats per minute per degree Celsius of fever, characteristic of *Salmonella* typhoid fever, Legionnaires' disease, and yellow fever (Faget sign).

**Alternate Names:**

Sphygmothermic Dissociation; Inappropriate Bradycardia in Fever

**Relative Risk**

A comparison of the chance of an outcome in an exposed group with its chance in an unexposed group. A value above 1 suggests higher risk, a value below 1 suggests lower risk, and 1 suggests no difference.

**Relaxation**

A reduction in muscle tension, physiologic arousal, or psychological stress; techniques may include breathing, biofeedback, or progressive muscle relaxation.

**Relaxation Response**

The body's calming changes after activities such as slow breathing, meditation, or progressive muscle relaxation. Heart rate, breathing, muscle tension, and stress-related alertness may decrease; these practices can support health but do not replace needed medical treatment.

**Relaxation Techniques For Stress**

Relaxation techniques for stress include slow breath-



---

ing, progressive muscle relaxation, meditation, guided imagery, and other methods that reduce physical arousal.

### **Relaxation Therapy**

Techniques such as breathing, muscle relaxation, or imagery used to reduce stress, arousal, or symptom burden.

### **Relaxin**

Relaxin is a reproductive hormone that helps remodel connective tissue and may contribute to cervical, pelvic, and vascular changes during pregnancy.

### **Religion**

A system of spiritual beliefs and practices that may influence health decisions, coping, diet, treatment preferences, and end-of-life care.

### **Relocation Test**

A shoulder examination often performed after an apprehension test. The examiner gently supports the front of the shoulder while the arm is in a vulnerable position. Relief of fear or pain may support shoulder instability, but the result must be interpreted with the rest of the examination.

#### **Alternate Names:**

Fowler Test; Jobe Relocation Test

### **REM**

Rapid eye movement, a sleep stage associated with vivid dreams and intense brain activity.

#### **Alternate Names:**

Rapid Eye Movement; REM Sleep

### **REM Behavior Disorder**

A sleep disorder where a person physically acts out dreams because normal muscle relaxation during REM sleep is lost.

#### **Alternate Names:**

RBD; REM Sleep Behavior Disorder

### **Remedy**

A treatment or measure intended to relieve a symptom or illness; safety and effectiveness depend on

the specific remedy and condition.

### **Remifentanyl**

A very short-acting opioid pain medicine given by infusion during anesthesia or intensive procedures. Its effects wear off quickly after the infusion stops, but it can slow breathing, lower blood pressure, and cause opioid-related complications.

### **Remission**

A period when the signs and symptoms of a disease decrease or disappear. Complete remission means no evidence of the disease can be found with available tests; partial remission means it has improved but remains detectable.

### **Remittent Fever**

Fever that rises and falls during the day but remains above normal temperature, rather than returning completely to normal.

### **Remodeling**

The body's ongoing process of removing old tissue and building new tissue to adapt to use, injury, growth, or disease. Bone remodeling responds to stress, hormones, nutrition, and medicines throughout life.

### **Remote Surgery**

Surgery performed when the surgeon is physically separated from the patient and controls robotic instruments through a telecommunications system. It may also include remote surgical assistance or guidance, with a qualified local team available to manage the patient and respond to emergencies.

#### **Alternate Names:**

Telesurgery Telepresence Surgery

### **Removal Technique**

The method used to take away a foreign object, diseased tissue, stone, clot, or other material from the body. The appropriate method depends on its location, size, cause, and risk of bleeding, infection, or injury to nearby structures.

---

## REM Rebound

An increase in rapid-eye-movement sleep after sleep loss, REM deprivation, or stopping a medicine that suppresses REM sleep. It may cause vivid dreams, nightmares, or temporary changes in sleep patterns.

## REM Sleep

A sleep stage with rapid eye movements, active brain activity, and temporary relaxation of most skeletal muscles. Vivid dreams are common, and REM sleep supports memory, learning, emotional processing, and healthy development.

## REM Sleep Behavior Disorder

A sleep disorder in which the normal muscle paralysis of REM sleep is reduced or absent, allowing a person to move or vocalize while dreaming.

### Alternate Names:

Rapid Eye Movement Sleep Behavior Disorder

## Renal

Renal means relating to the kidneys, which filter blood, regulate fluid and electrolytes, produce hormones, and excrete urine.

## Renal Abscess

A localized collection of pus within or around the kidney, usually caused by infection. Diagnosis and treatment may require imaging, antibiotics, and drainage.

## Renal Adenoma

A renal adenoma is a usually benign kidney tumor, often discovered incidentally; its size, imaging features, and growth determine whether follow-up is needed.

## Renal Agenesis

A condition present at birth in which one or both kidneys fail to form. One absent kidney may cause no symptoms if the other works well, but absence of both kidneys is usually fatal before or shortly after birth.

## Renal and Ureteral Injury

Mechanical, penetrating, blunt, or iatrogenic trauma affecting the kidneys or ureters. Clinical manifestations include gross or microscopic hematuria, flank pain, retroperitoneal hemorrhage, urinoma formation, or obstructive hydronephrosis requiring contrast-enhanced CT staging and potential surgical or endoscopic stenting.

### Alternate Names:

Upper Urinary Tract Trauma; Renal Laceration; Ureteral Disruption

## Renal And Urological Diseases

Renal and urological diseases affect the kidneys, urinary tract, bladder, urethra, or male reproductive urinary organs.

## Renal Anemia

Anemia associated with chronic kidney disease. Diseased kidneys make less erythropoietin, a hormone that stimulates red-cell production, and iron deficiency or inflammation may add to the problem, causing tiredness and breathlessness.

## Renal Arteriography

Renal arteriography uses contrast X-rays to visualize kidney arteries and may identify narrowing, bleeding, aneurysm, or vascular anatomy during an intervention.

## Renal Artery

One of the two main arteries carrying blood from the abdominal aorta to the kidneys. Each divides into smaller branches inside the kidney so blood reaches the filtering tissue.

## Renal Artery Aneurysm

A focal enlargement of a renal artery, often asymptomatic but capable of causing hypertension, thromboembolism, or rupture. Management depends on size, symptoms, anatomy, pregnancy, and growth, and may involve surveillance, endovascular treatment, or surgery.

---

## Renal Artery Disease

Alternate name; see *Renal artery stenosis*.

## Renal Artery Doppler

Doppler ultrasound that measures blood flow in the renal arteries. It may help assess narrowing or blockage that can contribute to renovascular hypertension.

## Renal Artery Embolism

Sudden blockage of an artery supplying the kidney, often by a clot from the heart in atrial fibrillation or another heart condition. It may cause abrupt side pain, blood in urine, nausea, or kidney injury.

## Renal Artery Stenosis

Narrowing of an artery supplying a kidney. It can reduce kidney blood flow and cause difficult-to-control high blood pressure or worsening kidney function. Causes include fatty buildup and fibromuscular dysplasia.

### Alternate Names:

RAS; Renal Artery Narrowing; Renovascular Disease

## Renal Biopsy

Alternate name; see *Kidney biopsy*. It is used selectively for unexplained acute kidney injury, nephrotic syndrome, glomerular hematuria, or transplant dysfunction; bleeding is the main complication.

## Renal Calculi

Alternate medical term; see *Nephrolithiasis*.

## Renal Calculus

A kidney stone, a hard mass of mineral crystals formed in the urinary tract. It may pass without symptoms or cause severe wave-like pain, blood in the urine, blockage, vomiting, or urinary infection.

## Renal Cancer

Alternate name; see *Kidney cancer*.

## Renal Carcinoma

Renal carcinoma is cancer arising from kidney tissue,

most commonly renal cell carcinoma. Cancer arising from kidney tissue, most often from cells lining the kidney's small filtering tubes.

## Renal Cell Carcinoma

A cancer beginning in the kidney's small filtering tubes, most often the clear-cell type. It may cause blood in urine, pain, weight loss, fever, or no symptoms until found on imaging.

## Renal Cell Carcinoma Pathology

Pathologic classification of renal cell carcinoma by growth pattern, cytology, immunophenotype, necrosis, sarcomatoid or rhabdoid change, and invasion. Clear cell, papillary, chromophobe, and other recognized subtypes have different features and behavior.

## Renal Changes In Pregnancy

Pregnancy increases renal blood flow and glomerular filtration. These changes lower usual serum creatinine values and may increase urinary loss of glucose and amino acids.

## Renal Circulation

Blood flow through the kidneys, which supports filtration, oxygen delivery, hormone production, and regulation of fluid and electrolytes.

## Renal Clearance

A measure of how quickly the kidneys remove a substance from the blood. It is usually calculated from blood and urine levels and helps assess kidney function or adjust some medicine doses.

## Renal Colic

Severe, wave-like pain caused when a kidney stone or other blockage stretches the urine tube. Pain usually starts in the side and may move toward the groin, with nausea, vomiting, or blood in urine; fever with blockage is an emergency.

## Renal Corpuscle

The first part of a kidney filtering unit, made of a tiny ball of blood vessels inside a cup-shaped capsule. Water and small dissolved substances pass out of

---

the blood here to begin urine formation.

### Renal Cortex

The outer layer of the kidney. It contains many of the structures that filter blood and process the first fluid that will become urine, while extensions of the cortex separate the inner kidney pyramids.

### Renal Cryptococcosis

Infection of the kidney by *Cryptococcus* species, usually as part of disseminated cryptococcosis in an immunocompromised person.

### Renal Cyst

Alternate name; see *Kidney Cysts*.

### Renal Dialysis

Alternate term for dialysis: kidney replacement therapy using hemodialysis or peritoneal dialysis to remove excess fluid and dissolved waste products when kidney function is inadequate.

### Renal Diet

An eating plan adjusted for kidney disease. Depending on kidney function and treatment, it may limit sodium, potassium, phosphorus, or fluid while providing enough calories and protein; needs differ between people and during dialysis.

### Renal Disease

Any condition affecting the kidneys, including infection, inflammation, stones, obstruction, inherited disease, or chronic loss of filtering function. Kidney disease can alter fluid balance, blood pressure, and body chemistry.

### Renal Diseases

Disorders affecting kidney tissue or function, including glomerular, tubular, vascular, infectious, obstructive, and neoplastic disease.

### Renal Dose Adjustment

Modification of a medication dose, dosing interval, or both to account for reduced kidney clearance. The adjustment should use an appropriate estimate

of renal function, drug-specific pharmacokinetics, treatment indication, and therapeutic monitoring when available.

### Renal Dysplasia

Abnormal development of kidney tissue before birth. It may affect one or both kidneys and can cause reduced kidney function, high blood pressure, urine reflux, or other urinary-tract problems.

### Renal Dysplasia And Cystic Disease

A group of developmental or inherited kidney disorders in which the kidneys have abnormal tissue, cysts, or both. They may affect one or both kidneys and can reduce kidney function or cause high blood pressure.

### Renal Failure

Severe loss of kidney function, causing waste, extra fluid, and abnormal body salts to build up. It may develop suddenly and sometimes improve, or progress gradually as chronic kidney disease; dialysis or transplant may be needed.

### Renal Function

The kidneys' ability to filter blood, remove waste, regulate fluids and electrolytes, maintain acid-base balance, and produce or activate hormones.

### Renal Glomerulus

A tiny ball of blood vessels inside each kidney filtering unit. Water and small dissolved substances pass from the blood through its walls to begin urine formation, while blood cells and most proteins remain in the bloodstream.

### Renal Hamartoma

A renal hamartoma is a benign disorganized overgrowth of mature tissues in the kidney, sometimes used to describe an angiomyolipoma-like lesion.

### Renal Hypertension

High blood pressure caused by kidney disease or narrowing of the renal arteries. High blood pressure caused by kidney disease or narrowed renal arteries,

---

often requiring treatment of both causes.

### **Renal Hypertrophy**

Renal hypertrophy is enlargement of kidney tissue, often compensating for loss or reduced function of the other kidney but sometimes caused by disease.

### **Renal Hypoperfusion**

Reduced blood flow to the kidneys, causing decreased glomerular filtration and potentially acute kidney injury. Causes include dehydration, blood loss, shock, heart failure, renal artery disease, and severe hypotension. Management requires treating the cause and restoring adequate circulation while avoiding further kidney injury.

### **Renal Infarction**

Death of part of the kidney caused by a blocked blood vessel. It may cause sudden flank or abdominal pain, nausea, blood in the urine, or reduced kidney function.

### **Renal Insufficiency**

Renal insufficiency is reduced kidney function that impairs waste removal, fluid and electrolyte balance, acid handling, or hormone production.

### **Renal Leukemia Infiltration**

Spread of leukemia cells into kidney tissue. It may enlarge both kidneys and reduce kidney function, sometimes causing sudden kidney injury, blood or protein in urine, or high blood pressure; treatment focuses on the underlying leukemia.

### **Renal Lithiasis**

Alternate name; see *Kidney stones*.

### **Renal Manifestations of Sickle Cell Disease**

Kidney complications of sickling disorders, including excessive urination, impaired concentration, hematuria, proteinuria, papillary necrosis, acute kidney injury, and progressive chronic kidney disease. Blood pressure and urine abnormalities require ongoing monitoring.

### **Renal Mass**

An abnormal area seen in or on a kidney during imaging. It may be a harmless cyst, infection, scar, benign growth, or cancer; its appearance, size, growth, and sometimes biopsy determine the next step.

### **Renal Medulla**

The inner part of the kidney containing tubes and blood vessels that help concentrate urine. It receives the fluid processed by the outer kidney and directs the finished urine toward the kidney's central collecting area.

### **Renal Necrosis**

Renal necrosis is death of kidney tissue caused by severe ischemia, toxins, infection, obstruction, or other injury and can cause acute kidney failure.

### **Renal Oncocytoma**

A renal oncocytoma is a usually benign kidney tumor arising from oncocytic cells; imaging may not always distinguish it reliably from renal cancer.

### **Renal Osteodystrophy**

The bone changes that are part of chronic kidney disease mineral and bone disorder. Changes in calcium, phosphate, vitamin D, parathyroid hormone, and bone remodeling can affect bone strength and growth; the pattern and treatment vary with the person and kidney disease.

### **Renal Papillary Necrosis**

Renal papillary necrosis is death of the kidney papillae from ischemia, infection, obstruction, diabetes, sickle-cell disease, or analgesic injury and may cause blood or obstruction.

### **Renal Parenchyma**

The working tissue of the kidney, containing the tiny filtering units and supporting structures that remove waste from blood, balance body fluids, and produce urine.

---

## Renal Pelvis

The funnel-shaped expansion of the ureter within the kidney, formed by the union of major calyces. It collects urine from the renal papillae and directs it into the ureter.

## Renal Pelvis Or Ureter Cancer

Cancer of the renal pelvis or ureter usually arises from urothelial cells lining the urinary tract and may cause blood in the urine.

## Renal Perfusion Scintiscan

A nuclear medicine study using a radiotracer to evaluate renal blood flow and relative perfusion of the kidneys.

## Renal Pyramids

Cone-shaped structures in the renal medulla that contain tubules and collecting ducts and drain urine toward the renal papillae.

## Renal Reabsorption

The return of water and useful substances, such as salt, sugar, and bicarbonate, from kidney tubes into the blood. This prevents their unnecessary loss and helps control body fluids and chemistry.

## Renal Replacement Therapy

Treatment that takes over some functions of severely damaged kidneys, including removing waste and extra fluid and correcting dangerous chemical imbalances. Options include hemodialysis, peritoneal dialysis, continuous treatment in intensive care, or kidney transplantation.

## Renal Replacement Therapy Indications

Reasons to consider urgent dialysis include dangerously high potassium, severe acid buildup, certain poisonings, fluid overload causing breathing difficulty, or symptoms from waste buildup. The decision also considers kidney recovery, overall health, and the person's goals.

## Renal Salt-Wasting Syndrome

Excessive loss of sodium and other salts through

the kidneys, causing low blood sodium, dehydration, low blood pressure, or muscle symptoms. Causes include inherited tubular disorders, medicines, kidney disease, and hormonal problems.

## Renal Sarcoma

Renal sarcoma is a rare malignant tumor arising from connective or supporting tissues of the kidney.

## Renal Scan

A nuclear medicine test using a small radioactive tracer to assess kidney blood flow, drainage, or separate function. The tracer and scan method are chosen according to the clinical question.

## Renal Scintigraphy

Nuclear imaging that follows a radioactive tracer through the kidneys. It can compare kidney function, assess blood flow and urine drainage, detect obstruction, or monitor a transplanted kidney.

## Renal Stone

A renal stone is a solid crystal aggregate formed in the kidney that can pass through the urinary tract or obstruct urine flow.

## Renal Transitional Cell Carcinoma

A urothelial carcinoma arising in the renal pelvis or calyces rather than the kidney's filtering tissue. It may cause blood in the urine, flank pain, or obstruction and is evaluated with imaging, urine testing, endoscopy, and tissue diagnosis.

## Renal Transplantation

Surgical placement of a functioning donor kidney into a person with advanced kidney failure. It can restore kidney function but requires suitability assessment and long-term immunosuppression.

### Alternate Names:

Kidney Transplant; Kidney Transplantation

## Renal Trauma

Kidney injury caused by a blow, fall, crash, sports injury, or penetrating wound. Assessment may include urine testing and CT imaging; treatment ranges



---

from observation to procedures or surgery depending on severity.

### Renal Tubular Acidosis

A group of disorders in which kidney tubules do not remove enough acid or return enough bicarbonate to the blood. This makes the blood too acidic and may cause weakness, kidney stones, bone disease, or high potassium.

#### Renal Tubular Acidosis Type 1

A metabolic acidosis caused by an impaired ability of distal intercalated renal tubular cells to secrete hydrogen ions into the collecting duct lumen. Characterized by normal anion gap metabolic acidosis, inappropriately high urinary pH (greater than 5.5 in the setting of systemic acidemia), hypokalemia, hypocitraturia, and recurrent nephrolithiasis or nephrocalcinosis.

##### Alternate Names:

Distal RTA; Type 1 RTA

#### Renal Tubular Acidosis Type 2

A normal anion gap hyperchloremic metabolic acidosis caused by defective proximal tubular reabsorption of filtered bicarbonate, resulting in bicarbonate wasting into the urine. Urinary pH is typically elevated initially but drops below 5.5 when systemic acidosis becomes severe; frequently associated with generalized proximal tubular dysfunction (Fanconi syndrome) including glucosuria, aminoaciduria, phosphaturia, and osteomalacia.

##### Alternate Names:

Proximal RTA; Type 2 RTA

#### Renal Tubular Acidosis Type 4

A normal anion gap metabolic acidosis characterized by persistent hyperkalemia caused by absolute aldosterone deficiency or renal tubular resistance to aldosterone (hypoaldosteronism). Impaired distal sodium reabsorption and potassium/hydrogen excretion reduce renal ammonium generation, yielding an acidic urinary pH (typically less than 5.5); common in diabetic nephropathy and chronic tubulointerstitial nephritis.

##### Alternate Names:

Hyperkalemic RTA; Hypoaldosteronemic RTA

### Renal Tubular Disorders

Diseases affecting the kidney tubules, which reabsorb water and useful substances and remove selected wastes into urine. They can disturb acid-base balance, electrolytes, urine concentration, or blood pressure.

#### Renal Tubule

A tiny nephron tube that reabsorbs needed water and substances and removes others into urine.

#### Renal Tubule Acidosis

A group of disorders in which the kidney tubules do not remove enough acid or conserve enough bicarbonate. The blood becomes too acidic, and complications may include low potassium, kidney stones, bone disease, weakness, or poor growth.

#### Renal Ultrasound

Noninvasive imaging used to assess kidney size, structure, blood flow, and urinary obstruction. It is particularly useful when obstruction, cystic disease, or structural abnormality is suspected and is often used in the evaluation of acute or chronic kidney disease.

#### Renal Vascular Disease

Alternate name; see *Renal artery stenosis*.

#### Renal Vein

One of two large veins that carry blood away from the kidneys to the body's main abdominal vein. A clot in a renal vein can impair drainage and damage kidney function.

#### Renal Vein Thrombosis

Renal vein thrombosis is a clot in a kidney vein that can cause flank pain, blood or protein in urine, kidney dysfunction, swelling, or pulmonary embolism.

#### Renal Venogram

Contrast imaging of the renal veins used selectively to evaluate venous anatomy, thrombosis, obstruction.

---

tion, or vascular abnormalities.

### **Renaut Body**

A Renaut body is a specialized cluster of cells in the peripheral nerve endoneurium, associated with pressure sensitivity and sometimes nerve pathology.

### **Renin**

An enzyme from kidney cells that starts a hormone chain controlling blood pressure, salt balance, and fluid levels.

### **Renin-Angiotensin-Aldosterone System**

A hormone system that raises blood pressure and preserves body fluid when circulation or kidney blood flow falls. It narrows blood vessels and signals the adrenal glands to retain salt and water; overactivity contributes to hypertension and heart failure.

### **Renin-Angiotensin-Aldosterone System Physiology**

When blood pressure or kidney blood flow falls, the kidneys release renin. A chain of chemical signals then narrows blood vessels and prompts the adrenal glands to retain salt and water, raising blood pressure and restoring circulation.

### **Renin-Angiotensin System**

A hormone and enzyme system that regulates blood pressure, fluid balance, and vascular tone. Renin initiates formation of angiotensin II, which promotes vasoconstriction and aldosterone release; excessive activation contributes to hypertension and heart failure.

### **Renin Blood Test**

A blood test measuring renin concentration or activity, interpreted with aldosterone, posture, sodium status, medications, and blood pressure to evaluate hypertension or mineralocorticoid disorders.

### **Rennin**

An older spelling that may refer to renin or to rennet enzyme; in humans, renin regulates blood pressure

and fluid balance.

### **Renography**

A kidney scan that follows a small amount of radioactive tracer as it enters and leaves the kidneys. It can compare kidney function and help identify reduced blood flow or blocked urine drainage.

### **Renovascular Disease**

Disease of the blood vessels supplying or draining the kidneys. It includes narrowing, blockage, aneurysm, or other vessel abnormalities and may reduce kidney blood flow or contribute to high blood pressure.

### **Renovascular Hypertension**

High blood pressure caused by reduced blood flow through one or both kidney arteries, usually from narrowing. It may be suspected when hypertension begins suddenly or is difficult to control; treatment addresses blood pressure and, in selected cases, the narrowed artery.

### **Renshaw Cell**

A nerve cell in the spinal cord that helps limit and regulate the activity of motor neurons.

### **Reoviridae Infection**

An infection caused by a reovirus; illness varies by virus and host and is often mild or asymptomatic in humans.

### **Rep**

An abbreviation most often meaning one repetition of an exercise or movement. In medical records it can also mean representative or another phrase, so the full wording should be used when the meaning is important.

### **Repair Of Webbed Fingers Or Toes**

Surgery to separate joined fingers or toes and reconstruct the skin between them. It may improve movement, growth, function, or appearance; wound problems, scarring, and recurrence of joining are possible.

---

## Repeat Surgery

Repeat surgery is a second or subsequent operation on the same or related area to treat persistent disease, recurrence, complications, or failure of prior treatment.

## Reperfusion

Restoration of blood flow to tissue after it has been reduced or interrupted. Reperfusion can save injured tissue but may also cause additional inflammation or oxidative injury.

## Reperfusion Injury

Damage that occurs when blood flow returns to tissue after a period of inadequate flow. The returning oxygen and inflammation can worsen injury, causing swelling, bleeding, or loss of function in organs such as the heart, brain, kidneys, or limbs.

## Reperfusion Therapy

Treatment that restores blood flow to tissue deprived of oxygen, such as clot-dissolving medicine, angioplasty, stenting, or surgery. Benefit depends on the tissue, duration of blockage, and risk of bleeding.

## Repetitive Motion Disorders

Conditions caused or worsened by repeated movement, force, vibration, or awkward posture. They may affect muscles, tendons, nerves, or joints and can cause pain, numbness, weakness, or reduced function.

## Repetitive Nerve Stimulation

A test that gives a nerve several small electrical impulses while recording the muscle's response. A changing response may support a disorder in communication between nerves and muscles, such as myasthenia gravis or Lambert-Eaton syndrome. The result is interpreted with the examination and other tests.

### Alternate Names:

RNS; Repetitive Stimulation Test

## Repetitive Strain Injury

Pain or dysfunction caused by repeated forceful

movements, sustained positions, or overuse of muscles, tendons, nerves, or other tissues.

## Replantation Of An Avulsed Tooth

Putting a completely knocked-out permanent tooth back into its socket after an injury. The tooth should be handled by its crown, kept moist, and seen urgently; later splinting, root treatment, and follow-up may be needed to prevent loss.

## Replantation Of Digits

Microsurgical replacement of a completely amputated finger or toe, reconnecting bone, tendons, nerves, arteries, and veins when feasible.

## Replication Initiation

The time and process when copying of DNA or another genetic molecule begins. Specific proteins recognize a starting region, separate the strands, and recruit enzymes that build matching new genetic material.

## Replicon

A unit of genetic material that can replicate from one origin, such as a chromosome segment, plasmid, or viral genome.

## Repolarization

The return of the membrane potential toward the resting value after depolarization. In neurons and skeletal/cardiac muscle, repolarization is primarily caused by the efflux of potassium through voltage-gated potassium channels and inactivation of sodium channels.

## Reportable Diseases

Reportable diseases are illnesses that must be notified to public-health authorities under applicable laws to support surveillance and outbreak control.

## Reporting Standards

Agreed rules for recording and communicating clinical or imaging findings. Structured reports improve completeness and consistency and can make important results, follow-up needs, and comparisons easier

---

for clinicians to understand.

### **Repressed Memory Therapy**

A controversial approach aimed at recovering allegedly repressed memories; suggestive techniques can create false memories and require caution.

### **Repression**

The reduction or stopping of gene activity, cell activity, behavior, or a body process. In genetics, regulatory proteins can silence genes; in psychology, the term may describe unconsciously keeping distressing thoughts outside awareness.

### **Reproduction**

The biological process by which living organisms produce new offspring, either sexually or without combining reproductive cells. In humans, it involves reproductive organs, hormones, eggs, sperm, pregnancy, and childbirth.

### **Reproductive Cells**

Cells used for reproduction, mainly sperm in males and eggs in females. Each normally carries half the usual number of chromosomes.

### **Reproductive History**

Reproductive history records pregnancies, births, miscarriages, abortions, menstrual function, fertility, contraception, and related conditions relevant to health assessment.

### **Reproductive Hormones**

Hormones that regulate sexual development, reproductive organs, ovulation, menstruation, sperm production, pregnancy, and lactation. Examples include estrogen, progesterone, testosterone, luteinizing hormone, and follicle-stimulating hormone.

### **Reproductive Organs**

The organs involved in reproduction. Female: ovaries (produce eggs and hormones), fallopian tubes (transport eggs, site of fertilization), uterus (nurtures the developing fetus), cervix (lower part of uterus, dilates during labor), vagina (passage for

sperm and baby), and vulva (external genitalia).

### **Reproductive System**

The organs and tissues that produce gametes and reproductive hormones and support sexual development and reproduction. Depending on the anatomy present, it may enable sperm or egg production, fertilization, pregnancy, childbirth, and lactation.

### **Reproductive Technique**

A reproductive technique is a medical or laboratory method used to assist conception, preserve fertility, diagnose reproductive disease, or manage pregnancy.

### **Reproductive Toxicity**

Harmful effects of a substance on fertility, pregnancy, the developing baby, birth outcomes, or later growth. Risks depend on the substance, dose, timing, route, and individual health and require careful exposure assessment.

### **Reproductive Years**

The part of life when a person may be able to become pregnant or produce sperm. Fertility varies widely and is affected by age, health, reproductive organs, hormones, and medicines; it should not be assumed from age alone.

### **Reprogram**

To reprogram a cell means to alter its gene-expression state so it acquires a different developmental potential or function.

### **Resectable**

Describing a tumor or lesion that can be removed completely by surgery with acceptable risk and an appropriate margin, based on its extent and relationship to vital structures.

### **Resection**

Surgical removal of part or all of an organ, tissue, or abnormal growth. The removed material may be examined to confirm diagnosis, determine treatment, or reduce disease while preserving useful function.

**Alternate Names:**

---

Surgical Excision; Surgical Removal

### **Resectoscope**

An endoscopic surgical instrument with a viewing system and working loop used to remove or cauterize tissue, especially during prostate or bladder procedures.

### **Reserve Volume**

The additional amount of air that can be inhaled or exhaled beyond a normal tidal breath.

### **Reservoir**

A human, animal, or environmental habitat in which an infectious agent normally lives, multiplies, and persists. A reservoir may serve as the source from which the agent is transmitted to a susceptible host.

### **Reservoir Of Infection**

A reservoir of infection is a person, animal, environment, or material in which an infectious organism normally lives and from which transmission can occur.

### **Resident**

A resident is a physician or other trainee receiving supervised specialty education after initial professional qualification; the term can also mean a person living in a place.

### **Residential Mobility**

Residential mobility is movement between homes or communities and can affect continuity of care, social support, exposure, and health access.

### **Residual**

Residual means remaining after a process, treatment, measurement, or event, such as residual urine, disease, risk, or volume.

### **Residual Limb**

The portion of a limb remaining after amputation. Care includes edema control, skin inspection, shaping, strengthening, desensitization, contracture prevention, and prosthetic preparation.

### **Residual Limb Pain**

Pain in the remaining portion of a limb after amputation.

#### **Alternate Names:**

Stump Pain

### **Residual Neoplasm**

A residual neoplasm is tumor tissue that remains after treatment intended to remove or destroy the original tumor.

### **Residual Risk**

Residual risk is the chance of harm that remains after preventive measures, treatment, screening, or controls have been applied.

### **Residual Tumor**

A residual tumor is cancer or other tumor tissue that remains after surgery, radiation, chemotherapy, or another treatment.

### **Residual Volume**

The amount of air remaining in the lungs after breathing out as forcefully as possible. It keeps small airways open and cannot be measured accurately by ordinary spirometry alone.

### **Resistance**

The ability of a microorganism to survive a medicine that previously inhibited or killed it. Resistance can make infections harder to treat and may spread between organisms through genetic material.

#### **Alternate Names:**

Antimicrobial Resistance; Drug Resistance

### **Resistance Training**

Exercise in which muscles work against weights, bands, body weight, or other resistance. Regular training can improve strength, endurance, balance, bone health, function, and independence when progressed safely.

### **Resistant Hypertension**

High blood pressure remaining above goal despite appropriate use of three or more medicines, includ-

---

ing a water pill.

**Alternate Names:**

Refractory Hypertension; Treatment-Resistant Hypertension

**Resistant Starch**

Starch that resists digestion in the small intestine and is fermented by gut bacteria in the colon.

**Resistin**

A signaling protein associated with adipose tissue and inflammation; its human clinical significance remains under study.

**Resolution**

The ability of an imaging system to distinguish separate structures, or the improvement or disappearance of a disease or symptom.

**Resonance**

Amplification or characteristic vibration produced when a system is driven at a suitable frequency; used in sound and MRI contexts.

**Resorcinol**

A phenolic compound used in selected topical dermatologic preparations; concentrated exposure can irritate or poison.

**Resorption**

The body's breakdown and removal of tissue or material, especially old bone. The released components may be absorbed, reused, or removed from the body.

**Resorption Atelectasis**

Atelectasis caused by complete obstruction of an airway, leading to absorption of trapped air by the pulmonary capillaries. Common causes include mucus plugging, foreign body, tumor, and post-surgical shallow breathing. The affected area appears as increased density on CXR with volume loss.

**Respiration**

The processes of breathing, gas exchange, and cellular energy production involving oxygen and carbon

dioxide.

**Respiratory**

Respiratory means relating to breathing, the airways, lungs, gas exchange, or the organs and muscles that move air.

**Respiratory Acidosis**

A condition in which breathing does not remove enough carbon dioxide, making the blood too acidic. It can result from lung disease, blocked airways, weak breathing muscles, brain disorders, or medicines that slow breathing.

**Respiratory Alkalosis**

A condition in which breathing removes too much carbon dioxide, making the blood less acidic. It can occur with anxiety, pain, fever, low oxygen, pregnancy, severe infection, or other conditions that cause rapid breathing.

**Respiratory Arrest**

Stopping of effective breathing. It is an immediate emergency that may require airway support, rescue breathing or ventilation, and treatment of the cause.

**Respiratory Burst**

A rapid release of chemically reactive oxygen molecules by certain immune cells after they engulf germs. This helps kill the germs inside the cell, although excessive activity can also damage nearby tissue.

**Respiratory Changes In Pregnancy**

Pregnancy changes breathing as the growing uterus raises the diaphragm and hormones increase the drive to breathe. Breaths may become deeper, overall ventilation increases, and the amount of air left after a normal breath out decreases.

**Respiratory Compliance**

The physical measure of the distensibility and elastic stretchability of the lungs and chest wall, expressed mathematically as the change in lung volume per unit change in distending pressure ( $\Delta V/\Delta P$ ); re-



---

duced in acute respiratory distress syndrome, pulmonary edema, and pulmonary fibrosis.

**Alternate Names:**

Lung Compliance

**Respiratory Depression**

Breathing that is abnormally slow or shallow, causing inadequate ventilation and possible carbon-dioxide retention or low oxygen. Medicines, brain disorders, and metabolic illness can cause it.

**Respiratory Disease**

Any disease affecting the airways, lungs, breathing muscles, or gas exchange. Examples include asthma, chronic obstructive pulmonary disease, pneumonia, tuberculosis, and lung cancer.

**Respiratory Disorder**

A disease affecting the airways, lungs, pleura, respiratory muscles, or control of breathing and causing impaired ventilation, oxygenation, or gas exchange.

**Respiratory Distress**

Difficult or labored breathing caused by problems such as asthma, infection, a blood clot, a collapsed lung, an allergic reaction, or heart failure. Fast breathing, struggling to speak, blue lips, or severe exhaustion may occur.

**Respiratory Distress Syndrome**

A serious breathing problem in premature newborns whose lungs do not make enough surfactant, a substance that keeps the tiny air sacs open. Babies may breathe rapidly, grunt, or need oxygen and breathing support. Antenatal steroids, noninvasive breathing support, and surfactant treatment improve survival.

**Respiratory Drive**

The brain's signal to the breathing muscles that controls how often and how deeply a person breathes. It responds mainly to carbon-dioxide and oxygen levels and can be reduced by brain disease, severe illness, or sedating medicines.

**Respiratory Failure**

A life-threatening inability of the lungs to provide enough oxygen or remove enough carbon dioxide. It may cause severe breathlessness, confusion, blue or gray lips, extreme sleepiness, or collapse. Causes include lung disease, infection, heart problems, weak breathing muscles, brain injury, and drug overdose.

**Respiratory Failure Types 1 and 2**

The clinical classification of acute respiratory failure: Type 1 (hypoxemic,  $P_aO_2 < 60$  mmHg with normal or low  $P_aCO_2$ ) resulting from ventilation-perfusion mismatch or shunt (e.g., pneumonia, pulmonary edema); and Type 2 (hypercapnic,  $P_aCO_2 > 45 - 50$  mmHg) resulting from alveolar hypoventilation (e.g., COPD, neuromuscular weakness, sedative overdose).

**Respiratory Gating**

A radiation-treatment method that tracks breathing and delivers the beam only during a selected part of the breathing cycle. It helps account for movement of tumors in the chest or upper abdomen and may reduce radiation to nearby healthy tissue.

**Respiratory Health**

The proper functioning of the lungs and airways, enabling efficient gas exchange and oxygen delivery to the body. Maintaining respiratory health involves avoiding tobacco smoke, minimizing exposure to air pollutants and occupational hazards, and receiving vaccinations against respiratory infections like influenza and pneumococcal disease.

**Respiratory Illness**

A short-term or long-term illness affecting breathing or the respiratory tract. Symptoms may include cough, fever, congestion, wheezing, shortness of breath, or chest discomfort, depending on the cause.

**Respiratory Infection**

An infection of the upper or lower respiratory tract caused by viruses, bacteria, fungi, or other organisms.

---

## Respiratory Infections

Illnesses caused by pathogens including viruses, bacteria, and fungi that affect the airways and lungs.

## Respiratory Insufficiency

Inadequate breathing or gas exchange that causes low oxygen, high carbon dioxide, or both. It can result from lung or airway disease, weak breathing muscles, or problems in the brain's breathing control and may progress to respiratory failure.

## Respiratory Muscle

A muscle used for breathing, especially the diaphragm and muscles between the ribs. These muscles expand and contract the chest to move air into and out of the lungs.

## Respiratory Neoplasm

A respiratory neoplasm is an abnormal growth in the airways or lungs and may be benign or malignant.

## Respiratory Paralysis

Respiratory paralysis is failure of the muscles needed for breathing because of nerve, muscle, brainstem, toxin, or spinal-cord dysfunction and is an emergency.

## Respiratory Physiology

The study of how breathing works, including movement of air, exchange of oxygen and carbon dioxide in the lungs, transport of gases in blood, and control of breathing.

## Respiratory Protection

Measures and equipment used to reduce inhalation of harmful dusts, fumes, gases, aerosols, or infectious particles.

## Respiratory Quotient

The ratio of carbon dioxide produced to oxygen used by the body's cells. The value changes according to whether the body is mainly using carbohydrate, fat, or protein and helps estimate metabolism and energy needs.

## Respiratory Rate

The number of breaths taken per minute. The expected rate varies with age, activity, temperature, illness, and other factors.

## Respiratory Rehabilitation

A program of exercise, education, breathing techniques, and self-management that improves function and quality of life in people with chronic respiratory disease.

## Respiratory Syncytial Virus

A respiratory virus that commonly causes bronchiolitis and pneumonia in infants and young children. Infection is usually mild in adults but may be severe in infants, older adults, and people with chronic heart or lung disease.

## Respiratory Syncytial Virus Infection

A common contagious respiratory illness caused by respiratory syncytial virus (RSV) that usually causes cold-like symptoms. It can also cause bronchiolitis or pneumonia, especially in infants and young children. Severe illness is more likely in older adults and people with certain heart, lung, or immune-system conditions. People can get RSV more than once.

### Alternate Names:

RSV; Human Orthopneumovirus

## Respiratory Syncytial Virus Bronchiolitis

An acute lower respiratory tract infection occurring predominantly in infants under two years of age caused by respiratory syncytial virus (RSV), characterized by bronchiolar mucosal inflammation, edema, and epithelial sloughing leading to wheezing, tachypnea, cough, and intercostal retractions.

## Respiratory System

The nose, mouth, pharynx, larynx, trachea, bronchi, bronchioles, lungs, pleura, and breathing muscles, including the diaphragm and chest-wall muscles. It moves air into and out of the lungs and exchanges oxygen and carbon dioxide between the alveoli and blood.

---

## Respiratory Therapy

Respiratory therapy includes treatments and breathing support that improve ventilation, oxygenation, airway clearance, or lung function.

## Respiratory Tract Infections

Infections affecting any part of the respiratory tract, classified by anatomical location into upper respiratory tract infections (URTIs) and lower respiratory tract infections (LRTIs).

## Respiratory Transport

Respiratory transport is movement of oxygen and carbon dioxide between lungs, blood, and tissues, largely through hemoglobin and dissolved gas.

## Respiratory Viruses

Viruses that cause respiratory tract infections. Major respiratory viruses include influenza, respiratory syncytial virus (RSV), parainfluenza virus, adenovirus, and human metapneumovirus. SARS-CoV-2 causes COVID-19. Symptoms range from mild upper respiratory illness to severe pneumonia.

## Respiratory Viruses And Face Masks

Face masks can reduce spread of respiratory viruses by blocking droplets and some aerosols, especially in crowded or healthcare settings; fit and local guidance matter.

## Respirologist

A doctor specializing in diseases of the lungs and breathing system. They assess breathing problems, interpret tests, manage chronic and acute illness, and may evaluate people for lung transplantation.

## Respirovirus

A genus of paramyxoviruses that includes human parainfluenza viruses, which can cause croup and other respiratory infections.

## Respite Care

Temporary care for a person who needs support so that a usual family or unpaid caregiver can rest, attend appointments, or manage other responsibilities.

ties. It may be provided at home, in a facility, or through short-term services.

## Response Assessment

Evaluation of how a disease responds to treatment. Clinicians may use symptoms, examination, blood tests, scans, tumor measurements, or activity levels to decide whether treatment is helping, failing, or needs adjustment.

## Response Evaluation Criteria

Standardized rules for judging whether a disease, especially a tumor, has responded to treatment, remained stable, or progressed. RECIST mainly uses tumor measurements on imaging; other systems may use metabolic or blood-based findings.

## Responsible Drinking

Responsible drinking means limiting alcohol according to health guidance, avoiding driving or hazardous activities after drinking, and avoiding alcohol when it is unsafe or contraindicated.

## Rest

Rest is reduced physical or mental activity used to recover, though excessive inactivity can impair strength and function in some conditions.

## Restenosis

Re-narrowing of a blood vessel after it has been opened by angioplasty or another procedure.

## Resting Energy Expenditure

The energy the body uses while resting to keep essential functions going, such as breathing, circulation, temperature control, and cell activity. It is affected by body size, age, muscle, illness, hormones, and recent food or activity.

### Alternate Names:

REE; Resting Metabolic Rate; Basal Metabolic Rate

## Resting Membrane Potential

The small electrical difference across a nerve or muscle cell when it is not sending a signal. Unequal amounts of charged particles inside and outside the

---

cell create it and allow the cell to respond quickly to stimulation.

### Resting Phase

The resting phase is a period of relative inactivity, such as G0 in the cell cycle or a physiologic interval between active events.

### Resting Tremor

An involuntary, rhythmic oscillation (typically four to six hertz) occurring in a body part that is fully supported against gravity and relaxed, which diminishes or temporarily ceases with voluntary movement. It typically presents as a pill-rolling hand tremor in Parkinson's disease and other parkinsonian syndromes.

### Restless Anal Syndrome

An uncommon term for uncomfortable sensations around the anus with an urge to move, touch, or relieve the area. The cause and best treatment are uncertain, so other anal and bowel conditions should be assessed.

### Restless Legs

Alternate name; see *Restless Legs Syndrome*.

### Restless Legs Syndrome

A neurological and sleep-related condition causing an irresistible urge to move the legs, often with crawling, pulling, aching, or tingling sensations. Symptoms begin or worsen during rest, improve temporarily with movement, and are worse in the evening or at night; they may disrupt sleep and daytime function. Iron deficiency, kidney disease, pregnancy, some medicines, and genetic factors may contribute.

#### Alternate Names:

RLS; Willis-Ekbom Disease

### Restless Leg Syndrome

Alternate name; see *Restless Legs Syndrome*.

### Restlessness

Restlessness is an uncomfortable need to move or

inability to remain still, caused by anxiety, akathisia, restless-legs syndrome, pain, stimulants, withdrawal, or medical illness.

### Restorative Dentistry

Specialty of restoring damaged teeth. Procedures: fillings, inlays, onlays, crowns, bridges, implants. Goals: function, aesthetics, durability.

### Restraint

A device, medicine, or other measure used to limit a person's movement. Restraints can cause injury, distress, falls, and loss of independence, so trained staff should use the least restrictive option only when necessary and monitor closely.

### Restraint-Related Harm

Physical or psychological injury associated with using a device, medication, or environmental restriction to limit a person's movement. Older adults are at risk of falls, pressure injury, delirium, aspiration, and functional decline; the least restrictive alternatives should be prioritized.

### Restriction Enzyme

An enzyme made by bacteria that cuts DNA at particular short sequences. Scientists use these enzymes to analyze, join, or modify DNA in laboratory research and medical testing.

### Restrictive Cardiomyopathy

A disease in which stiff heart muscle cannot relax and fill normally. It may cause breathlessness, tiredness, swelling, or abnormal rhythms and can result from conditions such as amyloidosis, sarcoidosis, iron overload, or scarring.

### Restrictive Lung Disease

A pattern of reduced total lung capacity caused by stiff lung tissue, chest-wall limitation, respiratory-muscle weakness, obesity, or other factors. It may cause exertional breathlessness and low oxygen; lung volumes, imaging, muscle strength, and the underlying cause guide management.

---

## Resuscitation

Emergency measures used to restore or support breathing and circulation, including opening the airway, rescue breathing, chest compressions, defibrillation, medicines, and treatment of the cause.

### Alternate Names:

Cardiopulmonary Resuscitation; CPR; Clinical Revival

## Resuscitative Thoracotomy

Emergency chest surgery used for selected trauma patients who are in cardiac arrest or close to it, especially after a penetrating chest injury. It may relieve blood around the heart, control bleeding, or allow direct heart compressions. It is a last-resort procedure with a high risk of death and complications.

### Alternate Names:

Emergency Department Thoracotomy; EDT; Clamshell Thoracotomy; Trauma Thoracotomy

## Retained Placenta

Failure of the placenta to come out after childbirth, usually when it remains in the uterus for about 30 minutes or longer or bleeding is excessive. It can cause severe bleeding or infection.

## Retained Products Of Conception

Pregnancy tissue that remains in the uterus after childbirth, miscarriage, or abortion. It may cause continuing or heavy bleeding, cramping, fever, or infection and may be treated with medicine or a procedure to remove it.

## Retainer

A dental appliance worn after orthodontic treatment to help keep teeth in their corrected positions.

## Retching

Repeated involuntary effort to vomit without necessarily producing stomach contents.

## Retention Of Urine

Inability to empty the bladder completely or at all despite making urine. It may cause painful full-

ness, leaking, or kidney damage and can result from blockage, weak bladder muscles, medicines, or nerve disease.

## Retention Time

The time a substance takes to pass through a laboratory separation system from injection until it is detected. Comparing it with a tested reference can help identify a substance, but timing alone is not definitive proof.

## Rete Testis

A network of small spaces inside the testis that collects sperm from the seminiferous tubules and passes them to the small ducts leading to the epididymis.

## Reticular

Net-like or arranged in a network pattern, describing the appearance or distribution of a skin lesion, rash, or vascular pattern.

## Reticular Activating System

A network of brain structures that helps maintain wakefulness, alertness, attention, and transitions between sleep and waking. Damage or severe disturbance can cause unusual sleepiness, reduced awareness, or coma.

## Reticular Dermis

The deeper layer of the dermis containing dense connective tissue, collagen, elastin, blood vessels, nerves, glands, and hair follicles.

## Reticular Erythematous Mucinosi

A rare primary cutaneous mucinosis presenting with erythematous, reticulated macules and plaques with dermal mucin accumulation on the mid-chest and upper back in middle-aged individuals, exacerbated by ultraviolet light exposure.

### Alternate Names:

REM Syndrome; Plaque-like Cutaneous Mucinosi

## Reticular Formation

A network of nerve cells in the brainstem that helps control alertness, sleep, attention, muscle tone, pos-

---

ture, breathing, heart function, and responses to pain.

### **Reticulin**

A network-forming type of collagen that supports bone marrow, liver, lymph nodes, and other organs.

### **Reticulocyte**

A young red blood cell recently released from bone marrow. The number of reticulocytes shows how actively the marrow is replacing red cells and helps distinguish reduced production from blood loss or red-cell breakdown.

### **Reticulocyte Count**

The number or percentage of immature red blood cells in blood. It reflects recent bone-marrow red-cell production and helps assess anemia, especially when interpreted with the hemoglobin and clinical context.

### **Reticulocyte Production Index**

A calculated hematological parameter that corrects the raw reticulocyte percentage for both the patient's degree of anemia (hematocrit) and the premature release of shift-reticulocytes from the bone marrow into circulating blood. Calculated as raw reticulocyte percent multiplied by (patient hematocrit / normal hematocrit) divided by maturation factor in days; an RPI of 2.0 or greater indicates an adequate marrow erythropoietic response to hemolysis or acute hemorrhage, whereas an RPI below 2.0 signifies an inadequate hypoproliferative marrow response.

#### **Alternate Names:**

RPI; Corrected Reticulocyte Count

### **Reticulocytes**

Immature red blood cells recently released from bone marrow; their proportion helps assess red-cell production.

### **Reticulocytopenia**

An abnormally low reticulocyte count indicating reduced effective red-cell production, which may occur with marrow failure, nutrient deficiency, in-

flammation, or kidney disease.

### **Reticulocytosis**

Reticulocytosis is an increased number of immature red blood cells in blood, usually indicating stimulated marrow production after blood loss or hemolysis.

### **Reticulo-Endothelial System**

An older name for the body's network of macrophages and related immune cells in the blood, liver, spleen, lymph nodes, and tissues. These cells remove germs, particles, and worn-out blood cells.

### **Reticuloendothelial System**

A diffuse anatomical network of specialized phagocytic cells (primarily monocytes and tissue macrophages) situated in reticular connective tissue frameworks of the spleen, liver (Kupffer cells), lymph nodes, and bone marrow, responsible for filtering particulate foreign matter, microbial pathogens, and senescent erythrocytes.

#### **Alternate Names:**

Mononuclear Phagocyte System; MPS

### **Reticulophagy**

A cell-cleaning process that removes worn or damaged parts of the endoplasmic reticulum, the cell's internal membrane network. The material is broken down and recycled, helping cells control quality and respond to stress.

### **Reticulum**

A network-like structure; in anatomy or cell biology it may refer to a tissue network, endoplasmic reticulum, or ruminant stomach compartment.

### **Retina**

The light-sensitive neural tissue lining the back of the eye that converts light into nerve signals.

### **Retina Disorder**

A disease affecting the retina, including vascular, degenerative, inflammatory, infectious, traumatic, inherited, or neoplastic conditions that can impair



---

vision.

### **Retina Hemorrhage**

Bleeding within or beneath the retina caused by vascular disease, diabetes, hypertension, trauma, retinal vein occlusion, inflammation, or other disorders.

### **Retinal Artery Occlusion**

Blockage of a retinal artery causing sudden, usually painless loss of vision. It may indicate a wider blood-vessel or clotting disorder.

### **Retinal Cone**

A light-sensing cell in the retina that provides color vision and sharp detail, especially in daylight. Damage to cone cells can reduce color perception and central vision.

### **Retinal Damage**

Injury or disease affecting the retina, potentially causing blurred vision, field loss, flashes, floaters, or permanent visual impairment.

### **Retinal Degeneration**

Progressive damage or loss of light-sensitive cells in the retina at the back of the eye. It may cause blurred or reduced central vision, night blindness, or loss of side vision and can result from inherited disease, aging, or other injury.

### **Retinaldehyde**

A form of vitamin A used in the eye's light-sensing cycle. Light changes it inside photoreceptor cells, helping start the nerve signals that allow vision.

### **Retinal Detachment**

A separation of the retina, the light-sensitive tissue at the back of the eye, from the layer beneath it. It may follow a retinal tear, scar tissue, or fluid buildup. Sudden floaters, flashes, a curtain-like shadow, or reduced vision may occur, and permanent sight loss is possible.

### **Retinal Detachment Repair**

Surgical or pneumatic treatment to close retinal breaks and reattach the retina. Techniques include vitrectomy, scleral buckling, and pneumatic retinopexy, selected according to the type and location of detachment.

#### **Alternate Names:**

Retinal Detachment Surgery

### **Retinal Diseases**

Disorders that damage the retina, the light-sensitive tissue at the back of the eye. They may cause flashes, floaters, distorted vision, blind spots, night-vision problems, or gradual or sudden sight loss.

### **Retinal Disorders**

Diseases affecting the light-sensing layer at the back of the eye. Causes include blood-vessel problems, inherited conditions, aging, inflammation, infection, injury, and cancer, potentially causing blurred vision or blindness.

### **Retinal Dysplasia**

Abnormal development and arrangement of the retina, the light-sensing layer at the back of the eye. It may impair vision and can occur with conditions present from birth, genetic disorders, or problems affecting development before birth.

### **Retinal Dystrophy**

Gradual loss or damage of retinal cells, often caused by an inherited condition. It may begin with night blindness, difficulty seeing in dim light, or loss of side vision and can progress to serious sight impairment.

### **Retinal Fold**

A retinal fold is a crease or ridge in the retina caused by traction, developmental abnormality, detachment, swelling, or another ocular process.

### **Retinal Fundus**

The inside back surface of the eye viewed during funduscopy, including the retina, optic nerve, macula, and retinal blood vessels.

---

## Retinal Ganglion

A nerve cell in the retina that collects signals from other retinal cells and sends them through the optic nerve to the brain. It is essential for carrying visual information out of the eye.

## Retinal Hemorrhages

Bleeding in the light-sensitive tissue at the back of the eye. It may result from diabetes, high blood pressure, injury, bleeding disorders, or increased pressure in the skull; in infants and children, the pattern must be assessed carefully for its cause.

## Retinal Implant

An implanted visual prosthesis that detects or processes light and electrically stimulates remaining retinal or visual-pathway cells in selected severe retinal disorders. It provides limited visual information and is distinct from an intraocular lens.

## Retinal Ischemia

Too little blood and oxygen reaching the retina, often because a retinal artery is blocked or blood-vessel disease is severe. It may cause sudden, painless sight loss and is an emergency because prolonged injury can become permanent.

## Retinal Laser Photocoagulation

An ophthalmic procedure utilizing targeted laser energy to induce localized thermal denaturation of retinal tissue. Widely used to seal microvascular leakages in diabetic macular edema, ablate ischemic peripheral retina in proliferative diabetic retinopathy, or secure retinal tears to prevent rhegmatogenous retinal detachment.

### Alternate Names:

Laser Photocoagulation; Retinal Laser Therapy; Panretinal Photocoagulation

## Retinal Layers

The organized layers of nerve cells, supporting cells, and blood vessels that make up the retina. Together they detect light, begin visual processing, and send signals through the optic nerve to the brain.

## Retinal Macula

The macula is the central region of the retina responsible for detailed straight-ahead vision, reading, and color discrimination.

## Retinal Neoplasm

A retinal neoplasm is an abnormal growth arising in or near the retina, such as retinoblastoma or a metastatic tumor.

## Retinal Perforation

A full-thickness hole or tear in the retina. Fluid can pass through the opening and separate the retina from the tissue beneath it, causing flashes, floaters, or loss of vision.

## Retinal Pigment

Retinal pigment refers to melanin-containing pigment in the retinal pigment epithelium, which supports photoreceptors and absorbs excess light.

## Retinal Pigment Epithelium

A layer of cells beneath the retina that supports the light-sensing cells, absorbs extra light, and helps remove worn-out cell material.

## Retinal Prosthesis

An implanted device designed to provide limited visual signals when the retina is severely damaged. A camera and electronic system stimulate remaining visual pathways; it may provide light or simple pattern perception, not normal sight.

## Retinal Tear

A break through the retina, often caused when the jelly inside the eye pulls on it. Fluid can pass through the tear and detach the retina; new flashes, floaters, or a curtain-like shadow may occur. Treatment may include laser or freezing treatment.

## Retinal Vasculitis

Inflammation of blood vessels in the retina. Leaking or blocked vessels can cause blurred vision, bleeding, swelling, or permanent sight loss; it may be linked to infection, autoimmune disease, or inflammation

---

elsewhere in the body.

### **Retinal Vein Occlusion**

A blockage in a vein that drains the retina. It can cause sudden or gradual blurred vision from bleeding and swelling, especially in the macula, and may lead to abnormal blood-vessel growth or permanent sight loss.

### **Retinitis**

Inflammation or infection of the retina, the light-sensitive tissue at the back of the eye. It may cause blurred vision, floaters, flashes, or loss of sight and needs prompt eye assessment.

### **Retinitis Pigmentosa**

A group of inherited retinal diseases in which the retina's light-sensing cells gradually stop working. Inheritance varies by gene and may be autosomal dominant, autosomal recessive, X-linked, or mitochondrial. Early symptoms often include poor night vision and loss of side vision, followed by narrowing of the visual field and possible severe sight loss.

### **Retinoblastoma**

The most common eye cancer in infants and young children, starting in the retina at the back of the eye. It is caused by changes in the *RB1* gene. It can be inherited (often affecting both eyes) or occur by chance (usually affecting only one eye). The most common early signs are leukocoria (an abnormal white glow in the pupil, often noticed in flash photographs instead of the normal red reflex) and strabismus (crossed or misaligned eyes). Diagnosis is made through a specialized dilated eye examination and imaging scans (ultrasound or MRI) to check tumor size and optic nerve involvement. Treatment focuses on saving the child's life and preserving vision using chemotherapy (given into the bloodstream or directly into the eye's blood vessels), laser therapy, freezing (cryotherapy), or surgical removal of the eye (enucleation) in advanced cases.

#### **Alternate Names:**

Childhood Eye Cancer; Retinal Glioma; Malignant Retinoblastoma

### **Retinoblastoma Protein**

A protein made from the *RB1* gene that helps prevent cells from dividing when they should not. Loss of its protective function can contribute to retinoblastoma and other cancers.

### **Retinoic Acid**

An active form of vitamin A that helps control how cells develop and which genes they use. Related medicines are used for selected skin conditions and some cancers but can cause irritation or serious pregnancy risks.

### **Retinoid**

A vitamin-A-related substance used in selected skin treatments and other medical care. Retinoids can improve acne or abnormal cell growth but may irritate skin, interact with medicines, and harm an unborn baby.

### **Retinoids**

Vitamin-A-related compounds used in selected dermatologic, ophthalmic, and oncologic treatments.

### **Retinol**

A form of vitamin A needed for vision, healthy skin and body linings, growth, and immune function. It comes from animal foods and some supplements; too much can be harmful, especially during pregnancy.

### **Retinopathy**

Damage or disease of the retina, the light-sensing layer at the back of the eye. Diabetes, high blood pressure, prematurity, blocked vessels, aging, inflammation, and inherited conditions can cause it and may lead to sight loss.

### **Retinopathy Of Prematurity**

Abnormal growth of blood vessels in the retina of a premature or very low-birth-weight infant. Mild disease may resolve, but severe disease can cause retinal detachment and blindness, so eligible infants need scheduled eye examinations.

---

### Retinoschisis

Splitting between layers of the retina. It may result from aging or an inherited condition and can cause reduced sharpness or loss of part of the visual field; it differs from retinal detachment but may look similar on examination.

### Retinoscope

An instrument that shines light into the eye to help determine refractive error. A lighted instrument used to estimate focusing error by observing how light reflects from the retina.

### Retinoscopy

An eye examination that uses a retinoscope to estimate refractive error by observing how light reflects from the retina.

### Retirement

Retirement is withdrawal from regular employment or a career; changes in income, activity, social contact, and health coverage can affect wellbeing.

### Retracted Nipple

Alternate name; see *Nipple Inversion Or Retraction*.

### Retractile Testicle

Alternate name; see *Retractile testicle*.

### Retractile Testis

A testis that moves between the scrotum and groin because of a strong reflex but can be gently brought into the scrotum. It should be monitored because it may later become undescended.

### Retractor

A surgical instrument used to hold tissue or an organ aside so the area can be seen and reached during an examination or operation. Some retractors are held by hand; others stay in place by themselves.

### Retrobulbar Hemorrhage

Bleeding behind the eyeball within the orbit, often after trauma or surgery, that can rapidly increase orbital pressure and threaten vision.

### Retrobulbar Neuritis

Inflammation of the optic nerve behind the eyeball, causing reduced or blurred vision and pain, especially when the eye moves. It may be associated with multiple sclerosis or another immune condition and needs prompt assessment.

### Retroesophageal

Retroesophageal means located behind the esophagus, such as a mass, lymph node, vessel, or collection in the posterior mediastinum.

### Retrognathia

Retrognathia is backward positioning of the lower jaw relative to the upper jaw and may affect facial structure, bite, airway, or sleep breathing.

### Retrograde

Moving backward or against the normal direction, such as retrograde amnesia or retrograde blood flow.

### Retrograde Amnesia

Loss of memory for events that occurred before an injury, illness, or other triggering event, with variable preservation of more remote memories.

### Retrograde Cystography

Retrograde cystography uses contrast placed through the urethra to outline the bladder and detect leaks, injury, or structural abnormalities.

### Retrograde Degeneration

Breakdown of a nerve fiber that spreads back toward the nerve cell body after the fiber is injured. It can reduce movement, sensation, or organ function depending on which nerve is affected and how severe the injury is.

### Retrograde Ejaculation

A condition in which semen enters the bladder instead of leaving through the penis during orgasm. It may cause little or no semen to appear and can result from diabetes, surgery, nerve problems, or medicines. It may affect fertility.

---

### **Retrograde Intrarenal Surgery**

RIRS is a minimally invasive procedure using a flexible scope passed through the urinary tract to inspect and fragment or remove kidney stones.

### **Retrograde Urethrogram**

An X-ray study that outlines the urethra after contrast material is introduced through its opening. It can show narrowing, tears, blockage, or abnormal connections, especially after suspected urethral injury.

### **Retroperitoneal Fibrosis**

Formation of abnormal scar-like tissue in the space behind the abdominal organs. It often surrounds the tubes draining urine from the kidneys, causing back or abdominal pain, urine blockage, and reduced kidney function; medicines, immune disease, or cancer can cause it.

### **Retroperitoneal Hematoma**

A collection of extravasated blood within the retroperitoneal anatomical space, commonly provoked by pelvic fractures, penetrating or blunt abdominal trauma, rupture of an abdominal aortic aneurysm, or femoral vascular catheterization; classified into zones I (central), II (flank), and III (pelvic).

### **Retroperitoneal Hemorrhage**

Bleeding in the space behind the abdominal lining. It may follow injury, blood-thinning treatment, a procedure, or rupture of a blood vessel and can cause back or abdominal pain, weakness, low blood pressure, and life-threatening blood loss.

### **Retroperitoneal Inflammation**

Retroperitoneal inflammation is inflammatory disease in the space behind the peritoneum, caused by infection, pancreatitis, urinary disease, trauma, medicines, or fibrosis.

### **Retroperitoneal Neoplasm**

A retroperitoneal neoplasm is an abnormal growth in the tissue space behind the abdominal lining and may be benign or malignant.

### **Retroperitoneal Space**

The retroperitoneal space is the area behind the peritoneal lining containing the kidneys, adrenal glands, pancreas, parts of the bowel, vessels, nerves, and lymphatics.

### **Retroperitoneum**

The space behind the abdominal lining. It contains the kidneys, adrenal glands, ureters, major blood vessels, and parts of the bowel and pancreas; bleeding, infection, tumors, or inflammation there can cause back or abdominal symptoms.

### **Retropharyngeal Abscess**

A collection of pus in the tissues behind the throat, usually after infection or injury. Fever, severe throat or neck pain, difficulty swallowing, drooling, noisy breathing, or a stiff neck can occur. The abscess can threaten the airway.

### **Retropharyngeal Space**

A potential fascial space in the neck located between the buccopharyngeal fascia of the pharynx anteriorly and the alar layer of the prevertebral fascia posteriorly, extending from the skull base down into the superior mediastinum (providing an anatomical conduit for the spread of retropharyngeal abscesses into the mediastinum).

### **Retrospective Studies**

Retrospective studies examine existing records or past exposures and outcomes; they are efficient but vulnerable to missing data, bias, and confounding.

### **Retrospective Study**

A study that examines existing records or past exposures to investigate disease causes, treatments, or outcomes. It can be efficient, but missing information and differences between groups may limit reliable conclusions.

### **Retrosternal Thyroid Surgery**

An operation to remove part or all of a thyroid gland that extends behind the breastbone. It may relieve pressure on the airway or swallowing tube and requires planning around nearby nerves, blood

---

vessels, and the parathyroid glands.

### **Retrotracheal**

Retrotracheal means located behind the trachea, such as a mass, lymph node, vessel, or collection in the posterior mediastinum.

### **Retrotransposition**

The copying and insertion of a genetic sequence into a new place in the genome using an RNA copy as an intermediate. New insertions can be harmless or disrupt a gene and cause disease.

### **Retroversion**

Backward tilting or turning of an organ, especially the uterus toward the spine. A backward-tilted uterus is often normal, but in some people it is linked with pain, endometriosis, or difficult examinations.

### **Retroversion Of The Uterus**

A uterine position in which the uterus tilts backward toward the spine; it is often a normal variant but can be associated with pain or endometriosis.

### **Retroviridae**

A family of RNA viruses that make a DNA copy of their genetic material and insert it into infected cells. HIV is a member; these viruses can cause persistent infection, immune disease, or cancer.

### **Retroviridae Infection**

An infection caused by a retrovirus, which may persist through genomic integration and cause immune, neurologic, or malignant disease.

### **Retrovirus**

A type of virus that uses RNA as its genetic material instead of DNA. It uses an enzyme called reverse transcriptase to make a DNA copy of its RNA. This copy can become part of the DNA of an infected cell, allowing the virus to make more copies. HIV is an example of a retrovirus.

### **Rett Syndrome**

A rare genetic disorder that affects brain develop-

ment and occurs almost exclusively in girls. Development may appear typical for the first 6 to 18 months, followed by loss of previously learned language, purposeful hand use, coordination, or movement. Repeated hand movements such as hand-wringing or clapping are common. Slowed head growth, seizures, breathing changes, feeding problems, sleep problems, and curvature of the spine (scoliosis) may occur. Most cases are caused by a new change in the *MECP2* gene; rare males with related gene changes can also be affected.

### **Return Of Spontaneous Circulation**

Restoration of a sustained heartbeat and blood flow after cardiac arrest, shown by a pulse, blood pressure, or other signs of circulation. Intensive post-arrest care then supports breathing, blood pressure, brain function, temperature, and treatment of the cause.

### **Reuptake**

The removal of a released nerve chemical from the space between nerve cells by returning it to the releasing cell or taking it into a nearby cell. This limits and ends the signal.

### **Revalidation**

Formal reassessment that a healthcare professional remains fit, competent, and up to date to practice.

### **Revascularisation**

Restoration of blood flow to an organ or limb by a procedure such as angioplasty, stenting, or surgical bypass.

### **Revascularization**

Restoration of blood flow to an area by bypass surgery, balloon treatment, stent placement, or another procedure. It can relieve symptoms, preserve tissue, and reduce damage caused by a blocked or narrowed vessel.

### **Revascularization Surgery**

Revascularization surgery restores blood flow around a blocked or narrowed vessel, such as in coronary artery bypass or limb bypass surgery.



---

### Reverberation Artifact

An ultrasound image error caused by repeated reflections, sometimes reduced by changing the angle or using specialized imaging.

### Reversal Learning

Reversal learning is the ability to change a learned response when the previously rewarded or correct choice is no longer so.

### Reverse Total Shoulder Replacement

Shoulder-replacement surgery that places the rounded joint surface on the shoulder blade and the socket on the upper-arm implant. This changes shoulder mechanics and can help when severe arthritis occurs with an irreparable rotator cuff.

### Reverse Transcriptase

An enzyme that synthesizes DNA using RNA as a template.

### Reverse Transcriptase Inhibitor

An antiviral medicine that blocks reverse transcriptase, used in combination treatment for HIV or selected infections.

### Reverse Transcriptase Inhibitors

A major class of antiretroviral medications utilized in the management of HIV-1 infection and chronic hepatitis B, categorized into nucleoside/nucleotide reverse transcriptase inhibitors (NRTIs, such as emtricitabine and tenofovir) and non-nucleoside reverse transcriptase inhibitors (NNRTIs, such as efavirenz and doravirine).

#### Alternate Names:

RTIs

### Reverse Transcription

The production of DNA using RNA as a template, carried out by reverse transcriptase. Some viruses use this process to copy their genetic material, and laboratories use it to study RNA.

### Reverse Trendelenburg Position

A supine position in which the head and upper body

are elevated higher than the feet. The head-up tilt uses gravity to shift abdominal organs downward, improving surgical visualization during head, neck, upper gastrointestinal, and laparoscopic procedures while facilitating ventilation.

#### Alternate Names:

Head-Up Tilt Position; Anti-Trendelenburg Position

### Reversible Cell Injury

Temporary cell damage that can resolve when the harmful cause is removed. Cells may swell, store excess fat, or produce less energy, but their membranes and genetic material remain sufficiently intact to recover.

### Revision Arthroplasty

Surgical replacement, exchange, or repair of a failed or painful joint prosthesis. Indications include infection, aseptic loosening, instability, periprosthetic fracture, wear, and recurrent dislocation.

### Revision Total Hip Arthroplasty

Surgery to remove and replace some or all parts of a previous hip replacement that have loosened, worn out, become infected, dislocated repeatedly, or failed. It is more complex than first-time replacement and may require bone reconstruction.

### Revision Total Knee Arthroplasty

Surgery to remove and replace parts of a previous knee replacement because of loosening, wear, infection, instability, stiffness, or persistent pain. It may require specialized implants or bone grafting and has higher risks than the first operation.

### Reye-Johnson Syndrome

Alternate name; see *Reye Syndrome*.

### Reye's Syndrome

A rare, potentially fatal illness causing acute brain dysfunction and liver injury, associated particularly with viral illness and aspirin exposure in children.

### Reye Syndrome

A rare, serious illness causing sudden brain swelling

---

and liver dysfunction, usually in a child or teenager recovering from a viral infection. Repeated vomiting, confusion, unusual sleepiness, seizures, or loss of consciousness may occur; aspirin exposure is associated with increased risk.

**Alternate Names:**

Reye-Johnson Syndrome; Fatty Liver with Encephalopathy

**Reynolds Pentad**

A combination of fever or chills, pain in the upper right abdomen, jaundice, low blood pressure, and confusion. Together these signs suggest severe infection and blockage of the bile ducts, which requires emergency treatment.

**Rezafungin**

A long-acting echinocandin antifungal medicine that inhibits fungal cell-wall synthesis. It is used for selected invasive candidiasis infections; its long duration permits an intermittent dosing schedule under specialist supervision.

**R-Factor**

A laboratory measure comparing the amount of an antibiotic needed to stop a bacterium with the expected effective amount. A higher value suggests reduced susceptibility and may help guide antibiotic choice.

**Rgs Proteins**

Regulators of G-protein signaling are proteins that speed the switch-off of signals from G-protein-coupled receptors. They help control responses to hormones, neurotransmitters, and other signals and may influence disease when altered.

**Rhabdoid Tumor**

A rare, fast-growing cancer that usually affects infants and young children. It may begin in the kidney, brain, or other tissues; changes in genes that control cell growth, especially SMARCB1, are common.

**Rhabdomyolysis**

Rhabdomyolysis is rapid breakdown of skeletal muscle that releases muscle proteins and salts into

the bloodstream. Causes include crush injury, extreme exertion, seizures, heat illness, medicines, and metabolic disease. It may cause muscle pain, weakness, dark urine, kidney injury, abnormal heart rhythms, or dangerous pressure in a limb. A high creatine kinase level supports the diagnosis.

**Rhabdomyolysis And AKI**

Severe muscle breakdown releases muscle proteins and salts, potentially causing acute kidney injury, abnormal body salts, and abnormal heart rhythms. Findings may include muscle pain, weakness, dark urine, and a high creatine kinase level.

**Rhabdomyoma**

A usually noncancerous tumor made of heart-muscle cells, seen mainly in infants and children. It is strongly associated with tuberous sclerosis and may shrink on its own, but large or multiple tumors can block blood flow or disturb the heart rhythm.

**Rhabdomyosarcoma**

Rhabdomyosarcoma is a malignant tumor showing skeletal-muscle differentiation, occurring most often in children but also in adults.

**Rhabdomyosarcoma Pathology**

A malignant mesenchymal tumor showing skeletal-muscle differentiation. Histologic patterns include embryonal, alveolar, spindle-cell, and pleomorphic forms, with primitive cells, rhabdomyoblasts, and variable myogenin or MyoD1 expression.

**Rhegmatogenous Retinal Detachment**

The most common type of retinal detachment, caused by a hole or tear that lets the eye's internal fluid pass beneath the retina. It can cause flashes, floaters, a curtain-like shadow, or sight loss.

**Rhesus Factor**

The Rh blood-group antigen, especially the D antigen, whose presence or absence is important in transfusion and pregnancy.

---

## Rhesus Incompatibility

Rhesus incompatibility occurs when an RhD-negative pregnant person forms antibodies against RhD-positive fetal red cells, risking haemolytic disease in a later or affected pregnancy; anti-D immunoglobulin prevents sensitization.

## Rhesus System

The Rh blood-group system, a group of inherited red-cell antigens including D, C, c, E, and e. RhD status is commonly reported as positive or negative and is important in transfusion compatibility and prevention of hemolytic disease of the fetus and newborn.

## Rheumatic Chorea

Involuntary, irregular, fidgety movements caused by an immune reaction after group A streptococcal infection. It is a feature of acute rheumatic fever and may affect the face, arms, legs, speech, or handwriting.

## Rheumatic Disease

A broad group of disorders affecting joints, muscles, bones, connective tissue, or immune function, including arthritis and lupus.

## Rheumatic Fever

An immune-mediated inflammatory illness that can develop after a group A streptococcal infection, usually strep throat. It may affect the heart, joints, skin, or brain. Heart inflammation can cause lasting valve damage (rheumatic heart disease). Diagnosis uses clinical criteria and evidence of a preceding streptococcal infection.

## Rheumatic Heart Disease

Permanent damage to heart valves after rheumatic fever, an immune reaction following group A streptococcal throat infection. It most often affects the mitral valve and can cause breathlessness, tiredness, swelling, abnormal rhythms, or heart failure.

## Rheumatism

Pain, stiffness, or discomfort in joints and surrounding soft tissues.

## Rheumatoid Arthritis

A long-term autoimmune disease in which the immune system mistakenly attacks the lining of the joints. This causes inflammation, pain, swelling, warmth, and stiffness, often worse after waking or resting. It commonly affects the same joints on both sides of the body, especially the hands, wrists, and feet. Over time, the inflammation can damage cartilage and bone and reduce joint function. It may also cause tiredness, low-grade fever, or loss of appetite and can affect the eyes, lungs, heart, blood vessels, skin, nerves, or blood cells.

## Rheumatoid Arthritis and Pregnancy

Rheumatoid arthritis often changes during pregnancy and may flare after delivery. Care balances disease control, joint function, medication safety, contraception or fertility goals, and fetal or newborn considerations, with coordinated rheumatology and obstetric review.

## Rheumatoid Arthritis Cause Gastrointestinal Issues

Rheumatoid arthritis can affect the gastrointestinal system indirectly through inflammation, associated autoimmune disease, reduced mobility, or medicines such as nonsteroidal anti-inflammatory drugs. Abdominal pain, bleeding, persistent diarrhea, or weight loss needs evaluation rather than attribution to arthritis alone.

## Rheumatoid Arthritis Inflammation Of The Brain

Direct brain inflammation is uncommon in rheumatoid arthritis, but vasculitis, medication effects, infection, stroke risk, or another autoimmune process can cause neurologic symptoms. New confusion, seizure, weakness, severe headache, or vision change requires urgent assessment.

## Rheumatoid Factor

An antibody that mistakenly recognizes part of another antibody. A blood test may support the diagnosis of rheumatoid arthritis, but it can also be positive in other diseases or in healthy people.

---

## Rheumatoid Lung Disease

Lung involvement associated with rheumatoid arthritis, including interstitial lung disease, pleural disease, nodules, or airway abnormalities.

## Rheumatoid Nodule

A rheumatoid nodule is a firm inflammatory lump under the skin or near a joint in rheumatoid arthritis, often over pressure points.

## Rheumatoid Nodules

Firm inflammatory lumps under the skin or near joints in people with rheumatoid arthritis, often over pressure points. They may cause no symptoms but can become tender or be affected by pressure.

## Rheumatoid Pneumoconiosis

Rheumatoid pneumoconiosis is lung disease involving rheumatoid arthritis and dust-related lung nodules, historically called Caplan syndrome.

## Rheumatoid Synovitis

Long-lasting immune inflammation of the tissue lining a joint in rheumatoid arthritis. It causes pain and swelling and can gradually damage cartilage, bone, and joint movement.

## Rheumatoid Vasculitis

Inflammation of small or medium-sized blood vessels associated with rheumatoid arthritis. It can reduce blood flow and affect the skin, nerves, eyes, or internal organs.

## Rheumatologist

A medical doctor who diagnoses and treats diseases affecting joints, muscles, bones, and connective tissues, especially autoimmune and inflammatory conditions such as rheumatoid arthritis, lupus, gout, and vasculitis.

## Rheumatology

The medical specialty concerned with inflammatory, autoimmune, and musculoskeletal disorders.

## Rh Factor

A system of antigens on the surface of red blood cells, especially the D antigen. A person is Rh-positive when the D antigen is present and Rh-negative when it is absent. Rh incompatibility can affect pregnancy and transfusion safety.

### Alternate Names:

Rh D Antigen; Rhesus Antigen

## Rh Incompatibility

An immune reaction that can occur when an Rh-negative pregnant person carries an Rh-positive fetus and forms antibodies against fetal red blood cells. Anti-D immune globulin can prevent sensitization.

## Rhinencephalon

An older term for brain regions involved mainly in smell and closely connected with emotion and memory. Modern anatomy usually names the individual structures instead of using this broad term.

## Rhinitis

Inflammation or irritation of the lining of the nose. It can cause a blocked or runny nose, sneezing, itching, and reduced smell and may result from allergy, infection, smoke, strong smells, medicines, or other triggers.

## Rhinitis Medicamentosa

A condition of severe rebound nasal mucosal congestion and edema caused by prolonged, repetitive use of topical alpha-adrenergic nasal decongestant sprays (such as oxymetazoline) beyond 3 to 5 days.

### Alternate Names:

Rebound Nasal Congestion; Topical Decongestant Rhinitis

## Rhinocephaly

A very rare birth defect involving severe abnormal development of the face and forebrain, with a nose-like structure that may lie unusually high or between the eyes. It is usually associated with major neurologic and developmental abnormalities.

---

## Rhinocerebral Mucormycosis

A rapidly progressive, fulminant, angioinvasive fungal infection caused by molds of the order Mucorales (such as *Rhizopus* and *Mucor*), occurring primarily in patients with diabetic ketoacidosis, severe neutropenia, or hematologic malignancies. Fungal hyphae invade blood vessels causing extensive thrombosis, tissue infarction, and black necrotic eschars of the nasal mucosa and hard palate, rapidly extending into the orbit and cavernous sinus.

### Alternate Names:

Invasive Rhinocerebral Zygomycosis

## Rhinophyma

Progressive thickening and enlargement of the nose caused by long-standing changes often associated with rosacea. It may produce a rough, irregular appearance and can sometimes interfere with breathing or require surgical treatment.

## Rhinoplasty

Surgery to reshape, repair, or reconstruct the nose for breathing, injury, birth differences, or appearance. Recovery includes swelling and bruising, and results depend on anatomy, technique, healing, and patient goals.

## Rhinorrhea

Excessive watery or mucoid discharge from the nose, commonly caused by infection, allergy, irritants, or changes in temperature.

## Rhinorrhoea

British spelling of rhinorrhea, meaning drainage of watery or mucus-like fluid from the nose. Causes include allergy, infection, irritation, and rarely a cerebrospinal-fluid leak.

## Rhinoscleroma

A chronic, slowly progressive granulomatous infection of the nose and upper airway caused by *Klebsiella rhinoscleromatis*. It may progress from foul-smelling discharge and obstruction to scarring, deformity, and airway narrowing and usually needs prolonged antibiotics.

## Rhinoscopy

Examination of the nasal cavity and nasopharynx using an instrument or endoscope to assess the mucosa, septum, turbinates, masses, bleeding, or obstruction.

## Rhinosporidiosis

A chronic localized granulomatous disease of the mucous membranes caused by *Rhinosporidium seeberi*, an aquatic protistan parasite in the clade Mesomycetozoea, endemic primarily to rural wetland communities in India and Sri Lanka. It classically manifests in the anterior nasal cavity or nasopharynx as painless, friable, highly vascular, sessile or pedunculated polypoid lesions with a characteristic "strawberry-like" appearance dotted with visible white spherules (sporangia), causing nasal obstruction, epistaxis, and foreign-body sensation.

### Alternate Names:

Nasal Rhinosporidiosis; *Rhinosporidium Seeberi* Infection

## Rhinovirus

A genus of non-enveloped, positive-sense RNA viruses in the Picornaviridae family. Rhinoviruses are a common cause of the common cold and spread mainly through respiratory secretions and contaminated hands or surfaces.

## Rhinovirus A

A species group of rhinoviruses that commonly causes the common cold and can worsen asthma or trigger wheezing.

## Rhinovirus C

A rhinovirus group associated with common-cold symptoms and, in some children, more severe wheezing or lower-respiratory illness.

## Rhinoviruses

A group of viruses that commonly cause colds. They spread through droplets, hands, and contaminated surfaces and may worsen asthma or cause more serious breathing illness in vulnerable people.

---

## **Rh Isoimmunization**

Maternal alloantibody production against fetal Rh D-positive red blood cells in an Rh D-negative mother, occurring when fetal RBCs enter the maternal circulation (fetomaternal hemorrhage).

## **Rhizomucor**

A genus of molds that can cause mucormycosis, especially in people with diabetes, neutropenia, or severe immune suppression.

## **Rhizomucor Pusillus**

A mold that can cause invasive mucormycosis involving the lungs, sinuses, skin, or other organs in severely immunocompromised people.

## **Rhizopus**

A genus of molds that includes important causes of mucormycosis, a rapidly invasive infection of sinuses, lungs, skin, or other tissues.

## **Rhizopus Oryzae**

A mold, also called *Rhizopus arrhizus*, that is a common cause of invasive mucormycosis.

## **Rhizotomy**

A procedure that cuts or heats selected nerve roots to reduce severe pain, muscle stiffness, or abnormal reflexes. It is used only in carefully selected cases because it can cause permanent numbness, weakness, or other nerve problems.

## **Rhodium**

A chemical element used in industrial and dental materials; its compounds can irritate tissues and may cause allergic reactions.

## **Rhodococcus**

A genus of gram-positive bacteria that includes *Rhodococcus equi*, an opportunistic pathogen causing pneumonia mainly in immunocompromised people.

## **Rhodococcus Equi**

A gram-positive bacterium that can cause cavitory

pneumonia and disseminated infection, especially in people with advanced immune suppression.

## **Rhodopsin**

A light-sensitive protein in rod cells of the retina that enables vision in dim light.

## **Rhodospirillaceae**

A family of metabolically diverse bacteria found in soil and water; some species carry out photosynthesis or nitrogen fixation.

## **Rhodospirillum**

A genus of spiral-shaped aquatic bacteria capable of photosynthesis or nitrogen fixation; human infection is rare.

## **RhoGAM**

A brand name for Rh immune globulin, which contains anti-D antibodies. It is given to eligible RhD-negative pregnant people after specified pregnancy events and around delivery to reduce formation of antibodies that could harm an RhD-positive fetus or newborn.

## **Rhombencephalon**

The hindbrain region of an early embryo. It develops into the pons, medulla, cerebellum, and related structures that help control breathing, balance, movement, swallowing, and other essential functions.

## **Rhonchi**

Low-pitched breathing sounds caused by air moving through narrowed or mucus-filled larger airways, often changing after coughing.

## **Rhonchus**

An abnormal sound heard through a stethoscope during breathing, caused by air moving through fluid, mucus, narrowed airways, or abnormal lung tissue.

### **Alternate Names:**

Crackles; Crepitations; Rales



---

## Rhubarb Leaves Poisoning

Poisoning after eating large amounts of rhubarb leaves, which contain oxalates and other irritants. It may cause burning, vomiting, abdominal pain, low calcium, kidney injury, weakness, or abnormal heart rhythms.

## Rhythm Method

A fertility-awareness method that estimates fertile days from menstrual-cycle timing to avoid or achieve pregnancy.

## Rib

One of 12 paired, curved bones forming the thoracic cage, protecting the heart, lungs, and great vessels. True ribs (1–7) attach directly to the sternum via costal cartilage. False ribs (8–10) attach indirectly to the sternum via the costal cartilage of rib 7. Floating ribs (11–12) have no anterior attachment.

### Alternate Names:

Costal Bone; Costa

## Rib Angle

The point where a rib bends most sharply as it curves from the spine toward the front of the chest.

## Ribavirin

An antiviral medicine used in selected viral infections, usually in combination and under specialist supervision.

## Rib Cage

The rib cage consists of ribs, sternum, thoracic vertebrae, muscles, and connective tissues that protect chest organs and assist breathing.

## Ribcage Pain

Ribcage pain may arise from muscles, ribs, joints, lungs, heart, or other organs; sudden severe pain or breathing difficulty may indicate a serious condition.

## Rib Fracture

A break in one or more ribs, usually causing localized chest pain worsened by breathing, coughing, or movement and sometimes associated with injury to

underlying organs.

## Rib Head

The back end of a rib that joins with the bodies of one or two thoracic vertebrae.

## Rib Neck

The narrowed part of a rib between its head and tubercle.

## Riboflavin

A water-soluble vitamin that forms the coenzymes FAD and FMN. These cofactors participate in oxidation-reduction reactions, including electron transfer, fatty-acid oxidation, and energy metabolism.

### Alternate Names:

Lactoflavin; Vitamin G

## Riboflavin Deficiency

Deficiency of vitamin B2, which can cause inflammation or fissuring of the mouth and lips, sore throat, dermatitis, anemia, and other mucosal or systemic findings.

## Ribonuclease

An enzyme that cuts RNA, a molecule involved in reading genetic instructions, into smaller pieces. Ribonucleases help control how long RNA lasts, recycle cell material, and protect cells from some unwanted genetic material.

## Ribonuclease H

An enzyme that removes the RNA strand when RNA and DNA are joined together. Cells use it during DNA copying, and some viruses depend on related activity when making DNA from their RNA.

## Ribonuclease III

An enzyme that cuts double-stranded RNA into smaller pieces. The products can help control gene activity, make proteins, or process genetic material in cells and some microorganisms.

---

## Ribonucleic Acid

RNA is a nucleic acid that carries genetic instructions, helps make proteins, and regulates cell activity; some viruses use RNA as their genetic material.

## Ribonucleoprotein

A structure made of RNA joined to one or more proteins. It helps make, process, transport, or control RNA; antibodies against certain ribonucleoproteins can occur in autoimmune diseases.

## Ribonucleoside

A molecule made of a ribose sugar joined to one of the bases used in RNA. Examples include adenosine, guanosine, cytidine, and uridine; adding phosphate groups produces RNA building blocks.

## Ribonucleotide Reductase

An enzyme that changes RNA building blocks into the related building blocks needed to make DNA. Cells require it to copy and repair DNA, so it is important for growth and is a target of some cancer medicines.

## Ribonucleotides

Nucleotides containing ribose sugar and RNA bases, serving as RNA building blocks and cellular energy or signaling molecules.

## Ribophagy

A cell-cleaning process that breaks down some ribosomes, the structures that make proteins, so their parts can be recycled. It helps cells adjust protein production when nutrients or growth conditions change.

## Ribose

A five-carbon sugar used to build RNA, ATP, and other molecules involved in storing genetic information and transferring energy. Cells make ribose from other nutrients and use it in many normal chemical processes.

## Ribosomal Protein

A protein that combines with ribosomal RNA to

form a ribosome, the cell structure that builds proteins. Inherited changes in some ribosomal proteins can cause bone-marrow failure or developmental syndromes.

## Ribosomal RNA

RNA that forms the main structural and working part of ribosomes, the cell structures that make proteins. It helps position messenger RNA and joins amino acids together; its genes are also used in some laboratory studies to identify organisms.

## Ribosome

A cellular structure made of RNA and proteins that reads messenger RNA and links amino acids together to make proteins.

## Ribosome Subunit

One of the two RNA-and-protein parts of a ribosome. The small and large parts join around messenger RNA and work together to assemble amino acids into a protein.

## Rib Tubercle

A small projection near the back of a rib that joins with the transverse process of a thoracic vertebra.

## Richter's Syndrome

Transformation of chronic lymphocytic leukemia or small lymphocytic lymphoma into an aggressive lymphoma, most often diffuse large B-cell lymphoma.

## Richter Transformation

The rapid clinico-pathological transformation of chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL) into an aggressive, high-grade diffuse large B-cell lymphoma (DLBCL), presenting with rapidly enlarging lymphadenopathy, systemic B symptoms, and elevated serum LDH.

### Alternate Names:

Richter's Syndrome; High-Grade Transformation of CLL

## Ricin

Ricin is a highly toxic plant protein that blocks

---

protein synthesis; ingestion, inhalation, or injection can cause severe organ injury and death.

### **Rickets**

A condition in children whose growing bones do not harden normally, usually because of too little vitamin D, calcium, or phosphate. It can cause weak or bowed bones, bone pain, and problems with growth.

#### **Alternate Names:**

Nutritional Rickets; Hypocalcemic Rickets

### **Rickettsia**

A group of small intracellular bacteria transmitted mainly by arthropods and causing diseases such as typhus and spotted fevers.

### **Rickettsiaceae**

A family of bacteria that live inside cells and includes several organisms spread by ticks, fleas, lice, or mites. They can cause fever, rash, blood-vessel inflammation, and serious organ complications.

### **Rickettsia Felis**

A flea-associated bacterium that can cause a spotted-fever-like illness in people. Infection may produce fever, headache, rash, fatigue, or muscle aches and is assessed with exposure history and laboratory testing.

### **Rickettsia Infections**

Rickettsial infections are illnesses caused by intracellular bacteria spread mainly by ticks, fleas, lice, or mites and often causing fever and rash.

### **Rickettsia Japonica**

A bacterium spread by ticks and associated with Japanese spotted fever. Infection may cause fever, headache, rash, and a dark mark where the tick bit; treatment uses suitable antibiotics.

### **Rickettsialpox**

An infection spread by house mites and caused by *Rickettsia akari*. It usually begins with a small dark sore at the bite site, followed by fever and a

blister-like rash.

### **Rickettsia Prowazekii**

A bacterium that causes epidemic typhus and is mainly spread through infected body-lice feces. Disease can cause high fever, headache, rash, confusion, and serious organ complications without prompt antibiotic treatment.

### **Rickettsia Rickettsii**

A bacterium spread by infected ticks that causes Rocky Mountain spotted fever. Fever, severe headache, rash, muscle aches, or confusion may develop; prompt antibiotic treatment is important because the illness can rapidly become life-threatening.

### **Rickettsia Typhi**

A bacterium spread mainly by infected fleas and responsible for murine typhus. The illness commonly causes fever, headache, chills, and sometimes a rash; prompt antibiotic treatment usually works, but untreated infection can become severe.

### **Riddoch Phenomenon**

A form of visual statuagnosia in which a patient with homonymous hemianopia or cortical blindness from an occipital lobe lesion is able to perceive moving visual targets within the blind field while remaining unable to visualize stationary, static stimuli.

#### **Alternate Names:**

Statuagnosia; Visual Motion Perception Preservation

### **Ridged Cranial Sutures**

Ridged cranial sutures are raised skull seams in an infant and may reflect normal molding or premature suture fusion, which requires assessment for craniosynostosis.

### **Ridged Nail**

A ridged nail has longitudinal or transverse grooves in its surface, which may result from aging, trauma, inflammation, nutritional problems, or systemic disease.

---

## Ridge Preservation

A dental procedure performed after a tooth is removed to reduce loss of the jawbone that held the tooth. Material is placed in the empty socket, sometimes with a covering, to help maintain shape for healing or a future implant.

## Riedel's Thyroiditis

A rare form of thyroid inflammation in which tough scar-like tissue makes the thyroid hard and may spread into nearby structures. It can cause a neck lump, difficulty swallowing or breathing, or reduced thyroid function.

## Riedel Thyroiditis

A rare, chronic fibroinflammatory disorder of the thyroid gland (an organ manifestation of IgG4-related disease) characterized by dense, invasive collagenous fibrous tissue replacing normal thyroid tissue and extending into adjacent strap muscles and trachea, presenting as a stony-hard, fixed ('woody') goiter.

### Alternate Names:

Invasive Fibrous Thyroiditis; Woody Thyroiditis

## Rieger Syndrome

A genetic condition affecting development of the front of the eye and often the teeth. It may cause glaucoma, unusual teeth, or changes around the navel and can result from changes in genes such as PITX2 or FOXC1.

## Riehl Melanosis (Pigmented Contact Dermatitis)

An acquired gray-brown to slate-colored facial or neck hyperpigmentation associated with contact or photocontact exposure to fragrances, cosmetics, dyes, or other chemicals. Itching or redness may precede pigmentation; identifying and avoiding the trigger, sometimes with patch or photopatch testing, helps guide management.

## Rifabutin

A rifamycin antibiotic used mainly for selected mycobacterial infections and tuberculosis regimens; it has important drug interactions and can discolor

body fluids.

## Rifampicin

A rifamycin antibiotic used in tuberculosis and selected other bacterial infections, usually in combination to prevent resistance.

## Rifampin

Rifampin is a rifamycin antibiotic that inhibits bacterial DNA-dependent RNA polymerase and is a key component of tuberculosis treatment; it causes orange body fluids and has many drug interactions and hepatotoxicity risk.

## Rifampin Toxicity

Adverse pharmacological effects of the first-line rifamycin antibiotic rifampin, including benign orange-red discoloration of body secretions (urine, tears, sweat), drug-induced hepatotoxicity, an immune-mediated flu-like syndrome with intermittent dosing, and potent induction of hepatic cytochrome P450 enzymes (CYP3A4).

## Rifamycin

Rifamycin is an antibiotic that inhibits bacterial RNA polymerase; related rifamycin medicines are used for tuberculosis and other infections.

## Rifaximin

Rifaximin is a minimally absorbed rifamycin antibiotic used for selected intestinal infections, prevention of recurrent hepatic encephalopathy, and some diarrhoeal syndromes; indication and resistance patterns guide use.

## Rift Valley Fever

A mosquito-borne viral disease affecting animals and humans, usually causing fever but sometimes severe eye, brain, or hemorrhagic disease.

## Right Atrium

The upper-right chamber of the heart. It receives oxygen-poor blood from the body and passes it through the tricuspid valve into the right ventricle, which sends it to the lungs.

---

### Right Bundle Branch

A group of heart fibers that carries the electrical signal to the right lower chamber so it contracts. If conduction is delayed or blocked, the ECG shows right bundle-branch block; this may be harmless or reflect heart or lung disease.

### Right Bundle Branch Block

Delay or interruption of electrical signals through the right bundle branch. It changes the ECG pattern and may occur without symptoms or with heart or lung disease.

**Alternate Names:**

RBBB

### Right Coronary Artery

A blood vessel that branches from the aorta and supplies much of the right side and lower part of the heart. A blockage can cause a heart attack or abnormal heart rhythm.

### Right Coronary Artery Occlusion

A blockage of the artery supplying much of the right and lower heart. It can cause a heart attack, slow or abnormal rhythms, low blood pressure, or failure of the right side of the heart.

### Right Heart Failure

Failure of the right side of the heart to pump blood effectively to the lungs. It may result from left-sided heart failure, high pressure in the lung arteries, a heart attack, lung clots, heart-muscle disease, or a stiff sac around the heart. It can cause swollen legs, neck veins, an enlarged liver, abdominal fluid, and weight gain.

### Right Heart Ventricular Angiography

An imaging test in which contrast dye and X-rays show the right lower heart chamber and how it pumps blood. It is used selectively to assess structure, movement, or blood flow during heart evaluation.

### Right Hypochondriac Region

The upper-right area of the abdomen, beneath the

lower ribs. It contains much of the liver, the gallbladder, and part of the intestine. Pain there may arise from these organs.

### Right Iliac Region

The lower-right area of the abdomen, containing the appendix, beginning of the large intestine, and part of the small intestine. Pain there may occur with appendicitis, bowel disease, urinary problems, or a hernia and must be interpreted with other findings.

### Righting Reflex

The righting reflex is an automatic response that restores head and body orientation when balance or position changes.

### Right Lower Quadrant Abdominal Pain

Alternate name; see *Appendicitis*.

### Right Lymphatic Duct

A short vessel that returns lymph from the right arm, right side of the head and neck, and right upper chest to the bloodstream near the right side of the neck. Damage or blockage can cause swelling in these areas.

### Right Sixth Cranial Nerve Palsy

Weakness of the nerve that moves the right eye outward. It can cause double vision, usually worse when looking to the right or at a distance, and may result from diabetes, high pressure in the skull, inflammation, stroke, or other disease.

### Right Ureterovesical Calculus

A stone lodged at the junction of the right ureter and bladder, causing flank pain, hematuria, and urinary urgency or frequency. May require endoscopic or lithotripsy intervention.

### Right Ventricle

The lower-right chamber of the heart. It receives oxygen-poor blood from the right atrium and pumps it through the pulmonary valve into the lung arteries to receive oxygen.

---

### Right Ventricular Strain

An electrocardiographic or echocardiographic pattern reflecting acute or chronic right ventricular pressure and volume overload, manifesting on ECG with T-wave inversions in leads V1 through V4, S1Q3T3 pattern, right bundle branch block, or rightward axis deviation (characteristic of acute massive pulmonary embolism).

### Rigid And Flexible Sigmoidoscopy

A sigmoidoscopy examines the rectum and lower colon with a flexible or rigid instrument. It can evaluate bleeding, pain, altered bowel habits, inflammation, polyps, or tumors; preparation, reach, comfort, and biopsy capability differ by technique.

### Rigidity

Abnormally increased resistance when a relaxed limb is moved by another person. It can occur with Parkinson disease, brain or spinal injury, severe infection, certain medicines, or other movement disorders.

### Rigler Sign

A classic abdominal radiographic sign of massive pneumoperitoneum (free intra-abdominal air) on plain supine radiographs, where both the inner (mucosal) and outer (serosal) surfaces of the bowel wall are clearly outlined by gas.

#### Alternate Names:

Double Wall Sign; Bas-Relief Sign

### Rigler Triad

Three imaging findings that strongly suggest gallstone ileus: blockage of the small intestine, air in the bile ducts, and a gallstone in the intestine. Not every person has all three findings.

### Rigor

Marked stiffness or rigidity; rigor mortis is post-mortem muscle stiffening, while rigor may also mean a severe chill.

### Rigor Mortis

Stiffening of the body's muscles after death. It

develops as the muscles lose the energy needed to relax and later fades as decomposition advances; its timing is affected by temperature, activity, illness, and the environment.

### Rilpivirine

An HIV medicine that blocks an enzyme the virus needs to copy itself. It is taken with other antiretroviral medicines, and missed doses or drug interactions can allow resistance to develop.

### Riluzole

A medicine used to modestly slow disease progression in amyotrophic lateral sclerosis and to treat selected forms of spinal muscular atrophy.

### Rimmed Vacuoles

Small empty-looking spaces with a visible border seen inside muscle cells on biopsy. They indicate that muscle cells are breaking down or recycling parts of themselves and can occur in conditions such as inclusion-body myositis and GNE myopathy. The finding must be interpreted with symptoms and other tests.

#### Alternate Names:

Autophagic Rimmed Vacuoles

### Ring Block

Local anesthetic injected around the base of a finger or toe to numb the entire digit.

### Ring Chromosome

A chromosome whose two ends have joined to form a ring, often with loss or disruption of genetic material. It may cause developmental or other health problems, depending on the chromosome and changes involved.

### Ringling In The Ear

Alternate name; see *Tinnitus*.

### Ringling In The Ears

Alternate name; see *Tinnitus*.

#### Alternate Names:

Tinnitus; Aurium Acusticus



---

## Ring Vaccination

A public-health strategy in which close contacts and contacts of contacts around an identified case are vaccinated to create a protective ring and limit transmission.

## Ringworm

A contagious fungal infection of the skin, scalp, hair, or nails. It commonly causes an itchy, scaly rash with a ring-like edge; scalp infection may cause broken hairs or hair loss. The appearance and complications vary by the affected site.

### Alternate Names:

Tinea Corporis; Dermatophytosis; Tinea

## Ringworm Of The Body

A contagious fungal infection of body skin causing an itchy, scaly, ring-shaped rash. It spreads through contact with infected people, animals, or objects and usually responds to antifungal treatment.

## Ringworm Of The Foot

Alternate name; see *Athlete's foot*.

## Ringworm Of The Scalp

A fungal infection of the scalp and hair that can cause scaling, broken hairs, and hair loss.

## Rinne and Weber Tests

Complementary bedside tuning fork examinations (typically using a 512 Hz fork) to evaluate auditory hearing loss: Rinne compares air conduction to bone conduction in each ear, while Weber tests lateralization of sound when the vibrating fork is placed on the midline forehead.

## Rinnes Test

Rinne's test compares air and bone conduction with a tuning fork to help assess conductive hearing loss.

## Rinne Test

The Rinne test is a screening test that compares air conduction and bone conduction hearing using a vibrating tuning fork (typically 512 Hz) to help differentiate between conductive and sensorineural

hearing loss.

## RIPE Therapy

Standard 4-drug regimen for active TB: Rifampin, Isoniazid, Pyrazinamide, Ethambutol. Intensive phase 2 months, continuation phase 4 months (total 6 months). Pyridoxine prevents isoniazid-induced neuropathy.

## Rippling Muscle Disease

A rare, benign myopathy characterized by mechanically-induced muscle contractions that spread across the muscle as a visible wave (rippling), typically triggered by stretch, percussion, or movement.

## Risedronate Sodium

The sodium salt of risedronate, a bisphosphonate used to prevent or treat osteoporosis and other disorders of excessive bone resorption.

## Risk

Risk is the probability of an unwanted event or outcome, considered together with its severity and the circumstances that influence it.

## Risk Adjustment

Risk adjustment accounts for differences in baseline patient characteristics so outcomes or payments can be compared more fairly across groups.

## Risk Factor

A characteristic, exposure, or behavior linked with a higher chance of disease or another outcome. It may be changeable, such as smoking, or unchangeable, such as age or inherited traits.

## Risk Management

Finding possible sources of harm, judging how likely and serious they are, and taking steps to reduce them. In healthcare, it includes safe systems, reporting problems, learning from errors, and checking whether safety measures work.

---

## Risk Register

A documented list of identified risks, their likelihood, impact, controls, owners, and review status.

## Risks Of Hip And Knee Replacement

Possible problems after joint replacement include infection, blood clots, bleeding, nerve or vessel injury, dislocation, stiffness, loosening, wear, persistent pain, and need for repeat surgery. Risk varies with the operation and the person's health.

## Risks Of Tobacco

Tobacco use increases risks of cancer, heart and blood-vessel disease, stroke, lung disease, pregnancy complications, addiction, and harm from second-hand smoke.

## Risks Of Underage Drinking

Alcohol use by minors can impair brain development, judgment, learning, safety, and mental health and increases risks of injury, dependence, and unsafe sexual or substance-related behavior.

## Risk-Taking

Risk-taking is behavior involving deliberate or inadvertent exposure to possible harm, influenced by perception, reward, development, mental health, and circumstances.

## Risky Drinking

Alcohol use that increases the chance of health, safety, work, relationship, or social problems. Risk depends on amount, frequency, pattern, personal health, medicines, pregnancy, age, and circumstances such as driving.

### Alternate Names:

Hazardous Drinking; At-Risk Alcohol Consumption

## Risperidone

Risperidone is an atypical antipsychotic that blocks dopamine and serotonin receptors and treats schizophrenia, bipolar disorder, and selected severe behavioral symptoms; hyperprolactinaemia, metabolic effects, movement disorders, and sedation may occur.

## Ristocetin-Induced Platelet Aggregation

A blood test that checks how platelets clump when exposed to ristocetin, a substance that requires von Willebrand factor and a platelet surface protein. It helps investigate von Willebrand disease and some platelet disorders.

## Risus Sardonicus

A fixed, strained grin caused by painful facial-muscle spasms. It is classically associated with tetanus or strychnine poisoning and reflects severe muscle overactivity.

## Ritonavir

Ritonavir is an HIV protease inhibitor mainly used at low doses to boost levels of other antiviral medicines by slowing their breakdown.

## Ritual

A ritual is a repeated, structured behavior or ceremony; repetitive rituals may also occur as compulsions in obsessive-compulsive disorder.

## Rituximab

Rituximab is a monoclonal antibody against CD20 that depletes B cells and is used for B-cell malignancies and selected autoimmune diseases; infusion reactions, infections, hepatitis-B reactivation, and rare PML are important risks.

## Rivaroxaban

Rivaroxaban is an oral direct factor Xa inhibitor used to prevent or treat venous thromboembolism and reduce stroke risk in selected atrial fibrillation; bleeding and renal-function-dependent dosing are important considerations.

## Rivastigmine

A medicine that increases acetylcholine signaling and may temporarily improve thinking or daily function in Alzheimer disease or Parkinson disease dementia. Common effects include nausea, vomiting, diarrhea, appetite loss, dizziness, and slow heart-beat.

---

## River Blindness

Onchocerciasis, a parasitic infection caused by *Onchocerca volvulus* and transmitted by blackflies. It can cause severe itching, skin disease, and visual impairment.

## RâLe

Older term; see *Crackles*.

## RLS

Restless legs syndrome, an urge to move the legs with uncomfortable sensations, especially at night.

### Alternate Names:

Restless Legs Syndrome; Willis-Ekbom Disease

## RMDs

Abbreviation; see *Repetitive Motion Disorders*.

## RNA

Ribonucleic acid, a molecule that helps cells read genetic instructions and make proteins. Some RNA carries messages from DNA, some helps build proteins, and some controls cell activity or forms part of viruses.

## RNA Binding

The attachment of a protein or other molecule to RNA, affecting RNA stability, transport, folding, processing, or translation.

## RNA Biosynthesis

The cellular production of RNA from a DNA template or, for some viruses, from an RNA template.

## RNA Degradation

The enzyme-controlled breakdown of RNA molecules. It controls how long genetic messages last, removes damaged or unnecessary messages, recycles their building blocks, and helps regulate protein production and cell behavior.

## RNA Folding

The process by which an RNA strand forms secondary and three-dimensional structures that influence its function.

## RNA Helicase

An enzyme that uses energy to separate, reshape, or move RNA structures and RNA-protein complexes. It helps cells copy, process, transport, and destroy RNA and is important for some viruses.

## RNA Interference

A gene-silencing process in which small RNAs guide proteins to degrade or block specific messenger RNA molecules.

## RNase P

An RNA-protein enzyme complex that processes precursor transfer RNA by removing its extra 5'-prime sequence.

## RNA Splicing

Removal of introns and joining of exons from a precursor RNA to produce mature messenger RNA or other functional RNA.

## RNA Transport

Movement of RNA within a cell or between cells, often mediated by transport proteins and required for translation or signaling.

## RNA Virus

A virus whose genetic material is RNA; its replication often involves an RNA-dependent polymerase and may have a high mutation rate.

## Road Rage

Alternate name; see *Intermittent explosive disorder*.

## Robinow Syndrome

A rare inherited skeletal-development disorder with distinctive facial features, short stature, vertebral or rib abnormalities, and genital differences.

## Robot-Assisted Rehabilitation

Robotic devices for repetitive task training. Upper extremity, lower extremity, gait training. Objective measurement, high repetition.

---

### Robotic-Assisted Thoracoscopic Surgery

Chest surgery performed through several small openings using instruments controlled by a surgeon from a console. The camera provides a magnified three-dimensional view and the instruments can move precisely. It may reduce the size of the incisions, but bleeding, infection, air leakage, and conversion to open surgery remain possible.

#### Alternate Names:

RATS; Robotic Thoracic Surgery; Robotic Lobectomy

### Robotic Hysterectomy

Hysterectomy performed through minimally invasive abdominal access using a surgeon-controlled robotic platform. The extent of uterine, cervical, tubal, and ovarian removal depends on the indication and operative plan.

### Robotic Surgery

A surgical technique in which a surgeon controls a computer-assisted system that moves a camera and small instruments. It is usually performed through small incisions and may allow precise movements, but the system does not operate independently.

#### Alternate Names:

Robot-Assisted Surgery; Computer-Assisted Surgery; RAS

### Rocker-Bottom Foot

A congenital foot deformity characterized by a prominent, downward-convex plantar surface with a dorsiflexed midfoot and rigid equinovarus hind-foot (prominent calcaneus), classically caused by a congenital vertical talus and strongly associated with trisomy 18 (Edwards syndrome) and trisomy 13 (Patau syndrome).

### Rocky Mountain Spotted Fever

A severe, potentially fatal tick-borne rickettsial infection caused by *Rickettsia rickettsii* transmitted by *Dermacentor* ticks, presenting with high fever, severe headache, myalgias, and a petechial maculopapular rash that characteristically begins on the wrists and ankles and spreads centripetally to the trunk, palms, and soles.

#### Alternate Names:

RMSF

### Rod

A rod is a long cylindrical structure; in the retina, a rod photoreceptor supports dim-light and peripheral vision.

### Rodenticide Toxicity

A heterogeneous group of poisonings whose presentation depends on the active ingredient. Anticoagulant products cause delayed coagulopathy and bleeding; bromethalin causes neurologic toxicity, cholecalciferol causes hypercalcemia and kidney injury, and metal phosphides can cause shock and organ failure. Identify the product and provide agent-specific treatment.

### Rodent Ulcer

An older name for basal cell carcinoma, a common locally invasive skin cancer.

### Rod Photoreceptor

A light-sensitive cell in the retina that supports vision in dim light and the peripheral field. Rods do not provide color vision; their loss can cause night blindness and narrowing of the visual field.

### Rods

Retinal photoreceptor cells that provide vision in dim light and contribute to peripheral vision but do not detect color.

### Rods And Cones

Two types of light-sensing cells in the retina. Rods help with dim-light and side vision, while cones provide sharp and color vision in brighter light.

### Roentgen Rays

Alternate name; see *X-Ray*.

#### Alternate Names:

Roentgen Radiation; Electromagnetic X-Radiation

### Rofecoxib

Rofecoxib is a selective COX-2 anti-inflammatory

---

drug withdrawn in many countries because of increased cardiovascular risk.

### **Romano-Ward Syndrome**

The most common congenital long QT syndrome, inherited in an autosomal dominant pattern without associated neurosensory deafness (due to mutations in cardiac potassium or sodium channel genes *KCNQ1*, *KCNH2*, or *SCN5A*), predisposing to syncope, polymorphous ventricular tachycardia (torsades de pointes), and sudden cardiac death.

### **Romanus Lesion**

An early inflammatory radiographic finding in ankylosing spondylitis characterized by enthesitis with focal erosion and surrounding reactive sclerosis at the anterior superior and inferior corners of vertebral bodies ('shiny corners'), preceding syndesmophyte formation.

#### **Alternate Names:**

Shiny Corner Sign; Spondylitis Anterior Enthesitis

### **Rombergism**

Unsteadiness or falling when standing with feet together and eyes closed, indicating impaired proprioception or vestibular function.

### **Romberg Sign**

Loss of balance when standing with the feet together and the eyes closed, suggesting impaired proprioception or vestibular function.

### **Romberg's Sign**

Increased unsteadiness when a person stands with the feet together and closes the eyes, suggesting impaired proprioception or vestibular function.

### **Romberg Test**

The Romberg test is a clinical neurological test used to assess the integrity of the dorsal column-medial lemniscus pathway (proprioception) and the cerebellum in maintaining balance. The test is performed by asking the patient to stand with feet together and eyes open, and then to close the eyes.

### **Romidepsin**

A histone deacetylase inhibitor used for selected T-cell lymphomas; it can cause cytopenias, infection risk, and ECG-related effects.

### **Romiplostim**

A thrombopoietin-receptor agonist that stimulates platelet production and is used mainly for chronic immune thrombocytopenia.

### **Rongeur**

A strong, jawed surgical instrument used to bite and remove small pieces of bone or other hard tissue.

### **Ronnel**

Ronnel is an organophosphate insecticide that can inhibit acetylcholinesterase and cause cholinergic toxicity after exposure.

### **Root Canal**

The channel inside a tooth root containing the pulp and its nerves and blood vessels.

### **Root Canal Therapy**

Dental treatment in which infected or inflamed pulp is removed from a tooth, the canal is cleaned and sealed, and the tooth is restored.

### **Root Canal Treatment**

Alternate term; see *Root Canal Therapy*.

### **Root Caries**

Tooth decay affecting exposed root surfaces after gum recession or loss of protective covering. It is more common with aging, dry mouth, or limited cleaning and may be slowed with fluoride and dental care.

### **Root Filling**

Dental treatment sealing a cleaned tooth root canal to prevent bacteria entering again and causing reinfection.

### **Root Fracture**

A crack or break in the root of a tooth. A hori-

---

zontal fracture may sometimes be stabilized and monitored, while a vertical fracture often allows bacteria to enter and may require removal of the tooth. Treatment depends on the fracture's location and extent.

### **Rooting Reflex**

A primitive reflex present from birth to 3-4 months, triggered by stroking the infant's cheek or corner of the mouth. The infant turns toward the stimulus and opens the mouth, searching for the nipple. Facilitates breastfeeding. Absence may indicate neurologic impairment. Disappears as voluntary feeding develops.

### **Ropinirole**

A medicine that stimulates dopamine receptors and is used for Parkinson disease and restless legs syndrome. It may cause nausea, dizziness, sleepiness, low blood pressure, hallucinations, swelling, or impulse-control problems such as gambling.

### **Ropivacaine**

A long-acting local anesthetic used for regional nerve blocks and epidural anesthesia; excessive blood levels can cause neurologic or cardiac toxicity.

### **Ropivacaine Hydrochloride**

The hydrochloride salt of ropivacaine, a long-acting local anesthetic used for regional and epidural anesthesia.

### **Rorschach Test**

A projective psychological test using inkblots; its diagnostic validity is limited and interpretation requires specialist training.

### **Rosacea**

A chronic inflammatory skin disorder that most often affects the central face. It may cause recurrent flushing, persistent redness, visible small blood vessels, burning or stinging, and acne-like bumps or pustules without being acne. Eye involvement may cause irritation or inflammation, and long-standing disease may produce thickening of the skin, especially of the nose (rhinophyma). Rosacea is not

contagious, and its severity and features vary over time.

### **Rosai-Dorfman Disease**

A rare, benign, non-Langerhans cell histiocytic proliferative disorder (sinus histiocytosis with massive lymphadenopathy) characterized clinically by massive painless bilateral cervical lymphadenopathy and fever, and pathologically by sinusoidal infiltration of S100-positive histiocytes demonstrating emperipolesis.

#### **Alternate Names:**

Sinus Histiocytosis with Massive Lymphadenopathy; SHML

### **Rose Bengal**

Rose bengal is a dye used in eye-surface testing and laboratory or diagnostic procedures; it highlights damaged or poorly protected conjunctival cells.

### **Roseola**

Roseola infantum is a common viral illness, usually caused by human herpesvirus 6, with several days of high fever followed by a pink trunk rash as the fever resolves; febrile seizures may occur.

#### **Alternate Names:**

Exanthema Subitum; Sixth Disease; Roseola Infantum

### **Roseola Infantum**

A common childhood illness, usually caused by human herpesvirus 6 and sometimes HHV-7. Several days of high fever are followed by a pink or red trunk rash as the fever resolves; febrile seizures may occur, and illness is usually self-limited.

### **Roseolovirus**

A genus of herpesviruses that includes human herpesvirus 6 and 7, commonly associated with roseola and sometimes serious disease after reactivation.

### **Rosiglitazone**

A medicine for type 2 diabetes that helps the body respond better to insulin and lower blood sugar. It can cause weight gain, swelling, fractures, or



---

worsening heart failure and is used selectively.

### **Rosiglitazone Maleate**

The maleate salt of rosiglitazone, a thiazolidinedione that improves insulin sensitivity in type 2 diabetes but may cause weight gain, edema, fractures, and heart-failure worsening.

### **Ross Procedure**

An operation for severe aortic-valve disease, especially in some children and young adults. The patient's own pulmonary valve is moved to the aortic position, and a donor valve or tube replaces it. The moved valve can grow in children and usually avoids lifelong blood-thinning medicine, but either valve may later narrow or leak.

#### **Alternate Names:**

Pulmonary Autograft Procedure; Ross Operation

### **Ross River Virus**

A mosquito-borne alphavirus that causes Ross River fever, with fever, rash, and often prolonged joint pain or swelling.

### **Rostral**

Positioned toward the nose or front of the head. In descriptions of the brain, it usually means toward the forehead or front rather than toward the back of the head.

### **Rosuvastatin Calcium**

Rosuvastatin calcium is a statin medicine that lowers LDL cholesterol and reduces the risk of cardiovascular events.

### **Rotation**

Movement around an axis; in anatomy it describes turning a body part inward, outward, or around its longitudinal axis.

### **Rotator Cuff**

The group of four shoulder muscles and tendons that stabilize the glenohumeral joint and rotate the arm.

#### **Alternate Names:**

Musculotendinous Cuff; SITS Muscles

### **Rotator Cuff Arthropathy**

Shoulder-joint damage that develops after long-standing, irreparable tears of the rotator-cuff tendons. It can cause pain, weakness, grinding, and inability to raise the arm; severe cases may need specialized shoulder replacement.

### **Rotator Cuff Exercises**

Rotator cuff exercises strengthen and stretch shoulder muscles and are selected according to the injury, pain, range of motion, and rehabilitation plan.

### **Rotator Cuff Injury**

Damage to one or more shoulder tendons or muscles that stabilize and move the arm. It may cause pain, weakness, night discomfort, or limited movement after overuse, aging, or sudden injury.

### **Rotator Cuff Problems**

Rotator cuff problems include tendinopathy, bursitis, and tears that cause shoulder pain, weakness, painful motion, and difficulty with overhead activity.

### **Rotator Cuff Repair**

Surgery to reattach a torn shoulder tendon to the upper-arm bone, usually with small instruments or an open approach. Recovery requires structured rehabilitation and may take months; stiffness, weakness, recurrent tearing, or infection can occur.

### **Rotator Cuff Tear**

A partial or complete tear of one or more tendons stabilizing the shoulder, causing pain, weakness, and difficulty raising or rotating the arm.

### **Rotator Cuff Tendinitis**

Alternate name; see *Rotator cuff syndrome*.

### **Rotavirus**

A very contagious virus that commonly causes vomiting and watery diarrhea in infants and young children. It can cause dangerous dehydration; vaccination greatly reduces severe illness, and testing is not

---

always needed.

### Rotavirus Antigen Test

A rotavirus antigen test detects rotavirus proteins in stool and can help diagnose viral gastroenteritis, especially in infants and young children.

### Rotaviruses

A group of viruses causing vomiting and watery diarrhea, especially in young children; vaccination lowers severe disease risk.

### Rotavirus Infection

A highly contagious viral gastroenteritis causing vomiting and watery diarrhea, especially in infants and young children; vaccination prevents most severe disease.

### Rotavirus Vaccine

A live-attenuated oral viral vaccine administered to young infants in a 2- or 3-dose series to prevent severe dehydrating gastroenteritis and diarrhea caused by rotavirus infection, carrying a slight, transient risk of intussusception.

### Rotenone

Rotenone is a naturally occurring pesticide that inhibits mitochondrial complex I and can be toxic to people and animals after exposure.

### Rothmund-Thomson Syndrome

A rare inherited condition that causes a mottled, red, and thin skin rash beginning in infancy, short stature, sparse hair, skeletal or dental problems, and sometimes cataracts. It also increases the risk of certain cancers.

### Roth Spots

Small areas of bleeding in the retina with pale centers, seen during an eye examination. They may occur with infection of the heart valves, severe anemia, leukemia, diabetes, or other illnesses and are not specific to one diagnosis.

### Roth's Spots

Small retinal haemorrhages with pale centres, associated with conditions such as infective endocarditis, severe anaemia, leukaemia, and diabetes.

### Rotor Syndrome

A benign, autosomal recessive inherited hyperbilirubinemia caused by mutations in *SLCO1B1* and *SLCO1B3* genes, resulting in defective hepatic reuptake and storage of conjugated bilirubin, presenting with lifelong, asymptomatic conjugated hyperbilirubinemia with normal liver histology and elevated urinary coproporphyrin.

#### Alternate Names:

Rotor's Syndrome; Familial Conjugated Hyperbilirubinemia

### Rotterdam Criteria

A commonly used approach for diagnosing polycystic ovary syndrome in adults. After other causes are excluded, diagnosis generally requires at least two of: irregular or absent ovulation, signs or blood-test evidence of high androgen levels, or polycystic-appearing ovaries on ultrasound. Ultrasound alone does not establish the diagnosis.

### Rotter Nodes

Small interpectoral lymph nodes situated between the pectoralis major and pectoralis minor muscles that provide an alternate pathway for breast lymphatic drainage directly to Level II and III axillary lymph nodes.

#### Alternate Names:

Interpectoral Lymph Nodes; Rotter's Nodes

### Rouleau

A stack or column of red blood cells, commonly seen in blood with increased plasma proteins.

### Rouleaux

A pattern where red blood cells stack like coins, often because blood plasma contains extra proteins.

### Rouleaux Formation

Linear, coin-like stacks or rolls of red blood cells

---

visualized on a peripheral blood smear, occurring when elevated plasma concentrations of asymmetric acute-phase proteins (fibrinogen or monoclonal immunoglobulins) neutralize the normal negative repulsive zeta potential of erythrocytes, characteristically seen in multiple myeloma and severe inflammatory states.

### Round

Round means circular or approximately spherical; in anatomy or pathology it describes shape and must be interpreted with the structure being described.

### Round Back

Alternate name; see *Kyphosis*.

### Round Ligament

A round ligament is a cord-like supporting structure, including the uterine ligaments that pass toward the groin and help maintain uterine position.

### Round Ligament of the Uterus

A fibrous and muscular cord originating from the superior-lateral aspect of the uterine cornu anterior to the fallopian tube, traversing through the inguinal canal to terminate in the subcutaneous tissue of the labium majus, functioning to hold the uterus in an anteverted and anteflexed position.

### Round Ligament Pain

Sharp or aching pain in the lower abdomen or groin caused by stretching of the round ligaments as the uterus enlarges during pregnancy. Persistent, severe, or bleeding-associated pain requires medical evaluation.

### Round Pneumonia

A lung infection that appears as a rounded shadow on a chest X-ray rather than the usual patchy pattern. It occurs especially in children and can resemble a tumor, so follow-up imaging may be needed after treatment.

### Roundworm

A parasitic worm, often referring to *Ascaris lumbricoides*,

acquired by swallowing eggs from contaminated soil, food, or water. It may cause abdominal symptoms, poor nutrition, or intestinal blockage when many worms are present.

### Roundworms

Nematode parasites that may infect intestines or tissues, causing disease depending on their species and number.

### Roussy-Levy Syndrome

An autosomal dominant hereditary motor and sensory neuropathy (a phenotypic variant of Charcot-Marie-Tooth disease type 1A) characterized by early-onset sensory ataxia, prominent postural hand tremor, distal muscle weakness and wasting, pes cavus, and absent deep tendon reflexes.

#### Alternate Names:

Hereditary Areflexic Dystasia; CMT-Tremor Variant

### Route Of Administration

The way a medicine enters the body, such as by mouth, under the tongue, injection, inhalation, skin application, or through a vein. The route affects how quickly and how much of the medicine reaches the bloodstream.

### Roux-en-Y Gastric Bypass

A bariatric and metabolic surgical procedure in which a small (15 to 30 mL) upper gastric pouch is created and anastomosed to a Roux limb of proximal jejunum (gastrojejunostomy), bypassing the remaining stomach, duodenum, and proximal jejunum, connected to a downstream biliopancreatic limb (jejunojejunostomy).

#### Alternate Names:

RYGB

### Roux Stasis Syndrome

A chronic functional disorder of gastric and small bowel motility developing in up to 30 percent of patients following a Roux-en-Y gastrojejunostomy or gastrectomy. Characterized by delayed gastric emptying, severe postprandial fullness, nausea, vomiting of non-bilious food, and chronic epigastric pain, re-

---

sulting from the loss of normal duodenal pacemaking coordination in the transected Roux jejunal limb.

**Alternate Names:**

Roux-en-Y Stasis Syndrome; Chronic Roux Limb Stasis

**Rovsing Sign**

Pain in the lower right abdomen that occurs when the lower left abdomen is pressed. This examination finding can support suspicion of appendicitis, but it is not diagnostic by itself and must be interpreted with the symptoms and other tests.

**Roxithromycin**

A macrolide antibiotic used in some countries for selected bacterial infections. It stops bacteria making proteins, but resistance, medicine interactions, stomach upset, liver problems, and heart-rhythm changes can occur.

**RPR Test**

A blood screening test for syphilis that detects antibodies produced during tissue injury from the infection. A reactive result does not prove active syphilis and requires a second, more specific test and clinical assessment.

**RS3PE Syndrome**

Remitting seronegative symmetrical synovitis with pitting edema, an inflammatory syndrome causing sudden swelling of the hands or feet, joint pain, and stiffness, usually in older adults. It may occur with another rheumatic disease, infection, or cancer.

**Alternate Names:**

Remitting Seronegative Symmetrical Synovitis with Pitting Edema

**RSI**

Repetitive strain injury, a group of pain and dysfunction syndromes associated with repeated movement or sustained posture.

**RSV**

A common respiratory virus that can cause mild cold symptoms or serious bronchiolitis and pneumonia,

especially in infants and older adults.

**RSV Antibody Test**

An RSV antibody test detects antibodies to respiratory syncytial virus and may help assess past exposure or immune response, but is not usually the main test for acute infection.

**RSV Infection**

Alternate name; see *Respiratory Syncytial Virus Infection*.

**RT-PCR**

A laboratory test that converts RNA into DNA and then makes many copies of a selected genetic sequence so it can be detected. It can identify RNA viruses and measure selected genes in clinical samples.

**Rubber**

Rubber is an elastic material used in gloves, devices, and equipment; natural-rubber latex can cause irritant or allergic reactions.

**Rubber Band Ligation**

The most common and effective non-surgical outpatient procedure performed to treat symptomatic internal hemorrhoids. Through a small lighted viewing scope (an anoscope) inserted into the anal canal, a clinician uses a suction or mechanical ligator instrument to grasp the redundant hemorrhoidal tissue above the pain-sensitive dentate line and snap a tiny, tight elastic rubber band firmly around its base. The tight band strangles the hemorrhoid's blood supply; within 7 to 10 days, the ischemic hemorrhoidal tissue withers, dies, and painlessly detaches, sloughing off and passing out unnoticed during a bowel movement. The remaining small scar fixes the surrounding mucosa to the underlying tissue, preventing further prolapse or bleeding without requiring surgical incisions, general anesthesia, or hospital recovery.

**Alternate Names:**

Hemorrhoidal Banding; RBL; Band Ligation of Hemorrhoids; Elastic Band Ligation

---

## Rubber Cement Poisoning

Poisoning from swallowing or inhaling rubber cement, which contains volatile solvents. It may cause dizziness, drowsiness, confusion, breathing problems, or chemical pneumonitis if it enters the lungs.

## Rubber Dams

Sheets used in dentistry to isolate teeth from saliva and the rest of the mouth during procedures.

## Rubella

A contagious viral infection causing mild fever, swollen lymph nodes, and a fine rash. Infection during pregnancy can cause miscarriage, fetal growth problems, hearing loss, eye disease, congenital heart disease, or other congenital rubella syndrome; vaccination prevents most cases.

### Alternate Names:

German Measles; Three-Day Measles

## Rubella Virus

The virus that causes rubella, a contagious illness with fever, swollen glands, and rash. Infection during pregnancy can seriously harm the developing baby, causing hearing loss, eye disease, heart problems, or other birth defects.

## Rubeola

Measles, a highly contagious viral illness caused by measles virus and characterized by fever, cough, conjunctivitis, and a widespread maculopapular rash.

### Alternate Names:

Measles; Hard Measles; Morbilli

## Rubeosis Iridis

Growth of abnormal new blood vessels on the colored part of the eye. It usually results from poor blood flow in the retina, often caused by diabetes or a blocked retinal vein, and can lead to painful glaucoma and vision loss.

## Rubidium

A chemical element whose radioactive isotopes can be used in medical imaging; nonmedical rubidium compounds can affect potassium-related physiology.

## Rubin Maneuver

An emergency childbirth technique for shoulder dystocia, when a baby's shoulder is stuck behind the birthing person's pelvic bone. A clinician applies pressure and rotates the baby's shoulder to reduce its width and help complete delivery safely.

## Rubinstein-Taybi Syndrome

A genetic developmental disorder associated with characteristic facial features, short stature, broad thumbs and toes, and intellectual disability. It is commonly caused by pathogenic variants in CREBBP or EP300.

## Rubivirus

A genus of viruses that includes rubella virus, which causes rubella and congenital rubella syndrome.

## Rubredoxins

Small proteins containing iron that help move electrons in some bacteria and other organisms. They take part in chemical reactions that release energy and protect cells from harmful oxygen-related chemicals.

## Rubricyte Count

A measurement of rubricytes, an older name for immature red blood cells, in blood or bone marrow. It may help assess how actively the marrow is making red cells, but modern tests usually report reticulocytes or other marrow findings.

## Rule Of Nines

A quick method for estimating the percentage of body surface burned by assigning simple proportions to body regions. The estimate helps guide fluids and emergency care, but children and unusual body shapes require more accurate methods.

## Rule Out

Rule out means evaluate a possibility and determine whether it is present; it does not mean the condition has already been excluded.

---

## Rumination

Repeated regurgitation and rechewing of food after eating, without the usual vomiting process. It is a behavior or symptom that may occur in rumination disorder or with other gastrointestinal, developmental, or behavioral conditions.

## Rumination Disorder

A feeding or eating disorder involving repeated, usually effortless return of recently eaten food to the mouth, followed by rechewing, reswallowing, or spitting. It differs from ordinary vomiting and causes clinically significant distress, impairment, or nutritional or medical consequences.

## Rumination Syndrome

Alternate name; see *Rumination Disorder*.

## Ruminococcus Gnavus

An intestinal bacterium involved in mucin and carbohydrate metabolism and studied in relation to gut inflammation.

## Rumpel-Leede Sign

Small pinpoint bleeding spots that appear below a blood-pressure cuff after it has briefly been inflated. The sign may suggest fragile small blood vessels or a low platelet count, but it is not specific and is rarely used alone today.

### Alternate Names:

Tourniquet Test; Capillary Fragility Test

## Rumpel-Leede Test

A clinical diagnostic test evaluating capillary wall fragility and thrombocytopenia in which a sphygmomanometer cuff is inflated around the arm midway between systolic and diastolic pressure for five minutes; the appearance of more than 10 to 20 petechiae in a designated circular area distal to the cuff indicates increased capillary fragility (as seen in dengue fever, thrombocytopenia, or scurvy).

### Alternate Names:

Capillary Fragility Test; Tourniquet Test

## Runner's Knee

Alternate name; see *Patellofemoral pain syndrome*.

## Runny Nose

Excess fluid or mucus draining from the nose, commonly caused by infection, allergy, irritants, or sinus disease.

## Rupture

A tear or complete break in an organ, vessel, muscle, or other tissue, often causing bleeding or loss of function.

## Ruptured Abdominal Aortic Aneurysm

A burst in a weakened section of the main artery in the abdomen, causing severe internal bleeding. Sudden abdominal or back pain, fainting, sweating, or collapse may occur.

## Ruptured Disk

A spinal disk whose outer layer has torn, allowing inner material to bulge outward.

### Alternate Names:

Herniated Disc; Slipped Disc; Herniated Nucleus Pulposus

## Ruptured Eardrum

A ruptured eardrum is a tear in the tympanic membrane caused by infection, pressure, trauma, or foreign objects and may cause pain, drainage, hearing loss, or dizziness.

## Ruptured Perforated Eardrum

A perforated eardrum is a tear in the tympanic membrane from infection, pressure change, trauma, or an inserted object. Pain, drainage, hearing loss, or ringing may occur; keeping the ear dry and avoiding unprescribed drops helps while healing or repair is assessed.

## Russell-Silver Syndrome

A condition present from birth that causes poor growth before and after birth, a relatively large head, feeding difficulty, uneven growth of the limbs or body, and characteristic facial features. It is



---

usually related to changes in gene regulation.

### **Russell Viper Venom Time**

A blood-clotting test used as part of testing for lupus anticoagulants. The venom activates clotting in the test tube; a prolonged result that shortens when extra phospholipid is added supports the presence of a lupus anticoagulant but does not establish the diagnosis by itself.

#### **Alternate Names:**

dRVVT

### **Rx**

A common abbreviation for prescription or prescribed treatment. It comes from the Latin word meaning “take” and is used on medication orders and in medical writing.

### **Ryle’s Tube**

A soft tube passed through the nose into the stomach to remove stomach contents or give fluids, nutrition, or medicines. Position must be checked before use because a misplaced tube can enter the airway and cause serious harm.

### **Raynaud Disease**

Alternate name; see *Raynaud’s Disease*.

### **Ringworm (Tinea)**

Alternate name; see *Ringworm*.

### **Respiratory Syncytial Virus (RSV)**

Alternate name; see *Respiratory Syncytial Virus*.

### **Restless Legs Syndrome (RLS)**

Alternate name; see *Restless Legs Syndrome*.

### **Rheumatoid Arthritis (RA)**

Alternate name; see *Rheumatoid Arthritis*.

### **Willis-Ekbom Disease**

Another name for restless legs syndrome, a neurologic sleep-related movement disorder causing an urge to move the legs and uncomfortable sensations that worsen at rest and at night.



# S

## S3 Gallop Pathology

An extra heart sound heard just after the normal second heart sound as the lower chambers fill rapidly. It can be normal in children and pregnancy but may suggest excess fluid or heart failure in an older adult.

## S4 Gallop Pathology

An extra heart sound heard just before the normal first heart sound when an upper chamber pushes blood into a stiff lower chamber. It may occur with long-standing high blood pressure, a narrowed aortic valve, or thickened heart muscle.

## SAA

Usually an abbreviation for serum amyloid A, a blood protein that rises during inflammation. Persistently high levels can contribute to abnormal protein deposits in organs during some long-lasting inflammatory diseases.

## Sabin Vaccine

The oral polio vaccine containing live attenuated poliovirus strains. It produces strong intestinal immunity but, rarely, vaccine-derived poliovirus can occur, so vaccine policy depends on local public-health guidance.

## Sabre Tibia

A skeletal physical deformity characterized by prominent anterior bowing and cortical thickening of the tibia, classically resulting from congenital syphilis (chronic periostitis) or Paget disease of bone.

### Alternate Names:

Saber Shin; Anterior Tibial Bowing

## Saccade

A quick, coordinated eye movement that shifts the eyes from one target to another. Abnormal saccades can occur with disorders affecting the eye muscles, balance system, or brain and nerves.

## Saccular Aneurysm

A rounded pouch bulging from one side of an artery because that part of the vessel wall is weak. It may enlarge, form clots, or burst; risk depends on its location, size, shape, and growth.

## Sacculle

A small, fluid-filled membranous sac situated within the vestibule of the inner ear, containing a vertically oriented macula with hair cells and otoliths that detect linear vertical acceleration and gravity.

## Sachet Poisoning

Illness caused by swallowing, inhaling, or getting the contents of a sachet on the skin or in the eyes. Effects depend on the substance and amount; some sachets contain medicines or dangerous chemicals.

## Sacral

Sacral means relating to the sacrum, the triangular bone at the base of the spine that connects with the pelvis.

## Sacral Agenesis

A rare congenital neural tube and caudal developmental defect characterized by complete or partial absence of the sacrum and lower spine, strongly associated with poorly controlled maternal diabetes mellitus.

---

**Alternate Names:**

Caudal Regression Syndrome

**Sacral Dimple**

A small indentation in the skin over the lower spine that is present at birth. Most are harmless, but a deep, large, high, hairy, discolored, or draining dimple may need imaging for an underlying spinal abnormality.

**Alternate Names:**

Pilonidal Dimple; Coccygeal Dimple

**Sacral Fistula**

An abnormal channel near the lower spine that connects the skin with deeper tissue or another body space. It may cause persistent drainage, pain, repeated infection, or a skin opening and can result from birth differences, injury, or pilonidal disease.

**Sacral Nerves**

The spinal nerves arising from the sacral spinal cord and supplying the pelvis, pelvic floor, bladder, bowel, and lower limbs.

**Sacral Nerve Stimulation**

Treatment that sends mild electrical pulses to nerves near the lower spine to improve bladder or bowel control. It may help selected people with persistent urinary or fecal leakage, retention, or constipation after other treatments have not worked.

**Sacral Nerve Stimulator**

An implanted device that sends mild electrical signals to nerves near the lower spine to improve bladder or bowel control in selected people. It may help incontinence or some emptying problems when other treatments have not worked.

**Sacral Plexus**

A network of spinal nerves in the pelvis that supplies the lower limbs, pelvis, and parts of the bladder, bowel, and reproductive organs.

**Sacral Region**

The lower back and posterior pelvic region overlying

the sacrum, a fused triangular bone that connects the lumbar spine with the pelvis.

**Sacral Vertebra**

One of the fused vertebrae forming the sacrum, which supports the spine, joins the pelvic bones, and transmits trunk weight to the lower limbs.

**Sacral Vertebrae**

Five spinal bones that fuse to form the sacrum at the back of the pelvis. They transfer weight from the spine to the hips and contain openings through which sacral nerves pass.

**Sacrococcygeal**

Sacrococcygeal means relating to the sacrum and coccyx at the lower end of the vertebral column.

**Sacrococcygeal Teratoma**

A tumor present from birth at the base of the spine and made of several tissue types. It may grow outside or inside the pelvis; a large fetal tumor can strain the heart and increase amniotic fluid.

**Sacrocolpopexy**

An operation that supports a prolapsed top of the vagina or uterus by attaching it to the strong tissue over the sacrum, usually with surgical mesh. It may be performed through open or small incisions; bleeding, infection, mesh problems, bowel or bladder injury, and recurrent prolapse are possible.

**Sacroiliac Joint**

The joint connecting the triangular sacrum at the base of the spine to each hip bone. It transfers weight between the spine and legs and may become painful or inflamed after injury, pregnancy, arthritis, or infection.

**Sacroiliac Joint Dysfunction**

Alternate name; see *Sacroiliac joint pain*.

**Sacroiliac Joint Injury**

Sprain, strain, or traumatic disruption of the ligaments and tissues supporting the sacroiliac joint. It

---

can cause localized low-back or buttock pain that worsens with transfers, stairs, prolonged standing, or asymmetric loading.

### **Sacroiliac Joint Pain**

Sacroiliac-joint pain arises near the junction of the lower spine and pelvis and may follow injury, pregnancy, arthritis, leg-length or gait changes, or inflammation. Pain can radiate to the buttock or thigh; examination distinguishes it from hip or spine disease.

#### **Alternate Names:**

SI Joint Dysfunction; Sacroiliitis

### **Sacroiliitis**

Inflammation of one or both joints connecting the lower spine to the pelvis. It can cause pain in the lower back, buttock, or back of the thigh and may result from inflammatory arthritis, infection, injury, or wear.

### **Sacrospinous Ligament Fixation**

An operation through the vagina that attaches the top of the vagina or cervix to a strong pelvic ligament to treat prolapse. It avoids an abdominal incision but may cause buttock pain, bleeding, infection, nerve injury, or recurrent prolapse.

#### **Alternate Names:**

Sacrospinous Colpopexy; Sacrospinous Suspension

### **Sacrum**

A large triangular bone at the back of the pelvis formed by five fused spinal bones. It connects the spine to the hip bones, transfers body weight to the legs, and contains openings for nerves supplying the pelvis and lower limbs.

### **SAD**

An abbreviation for seasonal affective disorder, a recurring pattern of depression or mood change linked to a particular season, often winter and reduced daylight. Treatment may include therapy, medicines, or supervised light therapy.

#### **Alternate Names:**

Seasonal Depression; Winter Depression

### **Saddle Anesthesia**

Loss of cutaneous touch and pain sensation over the perineum, buttocks, anus, and medial upper thighs corresponding to spinal cord nerve roots S3 through S5 (the anatomical region that would contact a riding saddle), serving as a hallmark red-flag clinical sign of cauda equina syndrome.

#### **Alternate Names:**

Perineal Sensory Loss; S3-S5 Dermatomal Anesthesia

### **Saddle Joint**

A joint in which each bone has one rounded and one hollow surface, fitting together like a saddle. It allows movement in two main directions, as at the base of the thumb.

### **Saddle Nose Deformity**

A physical collapse and depression of the cartilaginous nasal dorsum resulting from destruction or ischemic necrosis of the nasal septal cartilage, seen in granulomatosis with polyangiitis (GPA), relapsing polychondritis, congenital syphilis, chronic intranasal cocaine use, and nasal trauma.

#### **Alternate Names:**

Pug Nose; Depressed Nasal Dorsum

### **Saddle Pulmonary Embolism**

A large blood clot lodged where the main artery from the heart divides toward both lungs. It can severely block blood flow, causing sudden breathlessness, chest pain, fainting, low blood pressure, or cardiac arrest.

#### **Alternate Names:**

Saddle Embolus; Bifurcation Pulmonary Embolism

### **S-Adenosylmethionine**

A substance made naturally in the body that helps chemical reactions involving cell membranes, genetic material, and brain chemicals. It is also sold as a supplement, but its benefits and interactions vary by condition.

### **Sadism**

Sexual arousal or gratification from inflicting pain,

---

humiliation, or suffering; clinical significance depends on distress, impairment, or nonconsent.

### **Sad Mood**

Sad mood is a feeling of unhappiness or low mood that may be a normal response or a symptom of depression, grief, illness, or another condition.

### **Safe Sex**

Sexual practices that reduce the risk of sexually transmitted infections and unintended pregnancy, including condoms or other barriers, testing, vaccination, and informed consent.

### **Safety Of Drugs**

The process of balancing a medicine's expected benefits against side effects, interactions, and reasons it may be unsafe. Safe use includes checking the person, dose, allergies, other medicines, monitoring needs, and instructions.

### **Safety Plan**

A collaboratively written, practical plan for responding to escalating suicidal thoughts or emotional crisis. It identifies warning signs, internal coping strategies, supportive contacts, professional resources, and methods to restrict access to lethal means.

### **Safety Study**

A safety study evaluates whether a medicine, device, procedure, or treatment causes harmful effects and identifies risks before or during clinical use.

### **Saffron**

Saffron is a spice made from the dried stigmas of *Crocus sativus*. It is used in food and studied as a possible supplement, but it is not a substitute for proven medical treatment.

### **Safrole**

Safrole is a natural aromatic compound found in some plants and oils; high or prolonged exposure is toxic and has carcinogenic concern.

### **Sagittal**

Oriented in a vertical direction that divides the body or a structure into right and left parts. A midsagittal plane divides the parts equally; a parasagittal plane does not.

### **Sagittal Axis**

An imaginary line running from front to back through the body or a body part. Side-to-side movements, such as moving a limb away from or toward the body, occur around this axis.

### **Sagittal Plane**

An imaginary vertical plane dividing the body or an organ into right and left portions. A midsagittal plane divides them equally; a parasagittal plane divides them unequally.

### **Sagittal Sinus Thrombosis**

A form of cerebral venous sinus thrombosis involving the superior sagittal sinus, presenting with headache, papilledema, seizures, bilateral motor deficits, and the "empty delta sign" on contrast neuroimaging.

#### **Alternate Names:**

Superior Sagittal Sinus Thrombosis; SSST

### **Salaam Attacks**

Brief repeated bending spasms or sudden head drops in infants. They may be seizures and can affect development if they continue.

### **Salbutamol**

A quick-relief inhaled medicine that relaxes tightened airway muscles in asthma and chronic obstructive lung disease. It eases wheezing and breathlessness but may cause shaking, a fast heartbeat, or palpitations and does not replace treatment that controls airway inflammation.

#### **Alternate Names:**

Proventil; Ventolin

### **Salicylate**

A group of chemicals that includes aspirin and can reduce pain, fever, or inflammation. Excessive amounts may cause poisoning with vomiting,



---

ringing ears, fast breathing, confusion, dehydration, or dangerous blood chemistry changes.

### Salicylate Toxicity

Poisoning caused by too much aspirin or another salicylate. It may cause vomiting, ringing in the ears, fever, sweating, fast or deep breathing, dehydration, confusion, seizures, or coma.

### Salicylic Acid

A medicine applied to the skin to soften and remove excess surface skin. It is used for acne, psoriasis, warts, corns, and similar thickened or blocked skin; irritation is common and large exposures can be toxic.

### Saline

A sterile solution of sodium chloride in water. Different concentrations are used for intravenous fluids, wound or eye irrigation, inhalation, laboratory work, or dilution of medicines.

#### Alternate Names:

Normal Saline; Physiological Saline

### Saline Infusion Sonohysterography

An ultrasound test in which sterile salt water is placed inside the uterus to outline its cavity. It can show polyps, fibroids, scar tissue, or an unusually shaped cavity and may cause temporary cramping or light bleeding.

### Saline Nasal Washes

Rinsing the nasal passages with sterile, distilled, or appropriately treated saline to loosen mucus and reduce nasal irritants; unsafe water can cause serious infection.

### Saliva

Fluid made by the salivary glands that moistens food, begins digestion, protects the mouth and teeth, and helps swallowing and taste. Too little saliva can cause dry mouth, mouth soreness, and tooth decay.

### Salivary Duct Stones

Calcified deposits obstructing a salivary duct, most often the submandibular duct, causing recurrent painful swelling of a gland during meals and sometimes infection.

### Salivary Gland

A gland that makes saliva and releases it into the mouth. The three major paired glands are the parotid, submandibular, and sublingual glands; many smaller glands lie in the mouth lining.

### Salivary Gland Biopsy

Removal of salivary-gland tissue or a needle sample for microscopic examination of suspected autoimmune disease, inflammation, infection, or neoplasm.

### Salivary Gland Cancer

Cancer arising in a salivary gland, most often the parotid near the ear. It may cause a persistent lump, pain, numbness, dry mouth, or facial weakness; imaging and biopsy identify the type and guide treatment.

### Salivary Gland Infection

Infection of a salivary gland, usually bacterial and often involving the parotid or submandibular gland. Painful swelling, fever, dry mouth, and purulent drainage from the duct may occur.

### Salivary Gland Infections

Infections of a salivary gland or its duct, commonly bacterial or viral, causing painful swelling, tenderness, fever, dry mouth, or purulent drainage.

### Salivary Gland Neoplasm

An abnormal growth in a salivary gland that may be noncancerous or cancerous. It can cause a lump, pain, facial weakness, dry mouth, or swallowing problems and may require imaging and biopsy.

### Salivary Glands

Glands that make saliva, including the parotid, submandibular, and sublingual glands. Saliva moistens food, begins digestion, protects teeth, helps swal-

---

lowing, and supports taste and mouth health.

### Salivary Gland Scintigraphy

An imaging test that measures salivary-gland function after a small radioactive tracer is given. It may help assess dryness from Sjögren syndrome, blocked ducts, inflammation, radiation injury, or other gland problems.

### Salivary Gland Stones

Stones formed from mineral deposits in a salivary gland or its drainage tube. They can block saliva flow and cause swelling or pain, especially during meals; the medical term is sialolithiasis.

### Salivary Gland Tumors

Tumors arising in the salivary glands. Many are benign, especially in the parotid gland; pleomorphic adenoma is a common benign type. Malignant types include mucoepidermoid, adenoid cystic, and acinic cell carcinoma.

#### Alternate Names:

Salivary Neoplasms; Parotid Tumors

### Salivation

Production and release of saliva by the salivary glands. Saliva moistens food, begins digestion, protects teeth, and helps swallowing; excessive or reduced salivation can have many causes.

### Salk Vaccine

The inactivated poliovirus vaccine, given by injection to protect against poliomyelitis. It contains killed virus and cannot cause polio; it differs from the older oral live-attenuated vaccine.

### Salmonella

A genus of Gram-negative, facultatively anaerobic, rod-shaped bacteria in the Enterobacteriaceae family. It includes non-typhoidal strains that cause gastroenteritis and typhoidal strains that can cause enteric fever.

### Salmonella Choleraesuis

A Salmonella serotype adapted to pigs that can

cause severe, invasive infection in humans, including bloodstream infection and pneumonia.

### Salmonella Enterica

A major Salmonella species causing human disease. It can produce diarrhea and fever, enteric fever, or bloodstream infection after contaminated food, water, animals, or infected people spread the bacteria.

### Salmonella Enterocolitis

Salmonella enterocolitis is intestinal infection causing diarrhea, fever, abdominal cramps, and sometimes bacteremia after contaminated food, water, or animal exposure.

### Salmonella Food Poisoning

Alternate name; see *Salmonellosis*.

### Salmonella Infection

An infection caused by Salmonella bacteria, usually acquired through contaminated food, water, animals, or hands. It commonly causes diarrhea, fever, cramps, and vomiting but can spread into blood or other organs.

### Salmonella Infections

Infections caused by Salmonella bacteria, ranging from gastroenteritis to enteric fever or invasive bloodstream disease.

### Salmonella Typhi

The bacterium that causes typhoid fever, usually acquired through contaminated food or water and capable of spreading through the bloodstream.

### Salmonella Typhimurium

A common Salmonella serotype that usually causes foodborne gastroenteritis but can cause invasive disease in vulnerable people.

### Salmonella Typhisuis

A Salmonella serotype mainly associated with swine; human infection is uncommon but may cause enteric or invasive disease.

---

## Salmonella Vaccines

Vaccines used mainly to prevent typhoid fever, including typhoid conjugate, Vi polysaccharide, and oral live-attenuated vaccines; routine vaccination does not prevent all Salmonella gastroenteritis.

## Salmonellosis

Infection with Salmonella bacteria (non-typhoidal), typically acquired from contaminated food (eggs, poultry, meat). Presents with acute gastroenteritis: diarrhea, fever, and abdominal cramps within 6-72 hours of ingestion. Self-limited in immunocompetent patients. Bacteremia can occur, especially in immunocompromised.

### Alternate Names:

Salmonella Enteritis; Salmonella Gastroenteritis

## Salmon Patches

Flat pink or red capillary birthmarks, commonly on the eyelids, forehead, or nape; many fade during infancy.

## Salpingectomy

Surgical removal of a fallopian tube. It may be used to treat a tubal ectopic pregnancy, hydrosalpinx, or other tubal disease. The approach and whether the tube is removed depend on the clinical condition, fertility plans, and the state of the opposite tube.

## Salpingitis

Inflammation or infection of one or both fallopian tubes, usually as part of pelvic inflammatory disease spreading upward from the vagina or cervix. It may cause lower-abdominal pain, fever, unusual discharge, infertility, or pregnancy outside the uterus.

## Salpingitis Isthmica Nodosa

Small thickened areas and side pockets in the narrow part of a fallopian tube, often after inflammation. They can narrow the tube, making pregnancy harder to achieve and increasing the risk of an ectopic pregnancy.

### Alternate Names:

SIN

## Salpingography

Contrast imaging of the fallopian tubes, usually performed as part of hysterosalpingography to assess whether the tubes are open and to examine the uterine cavity.

## Salpingo-Oophorectomy

Surgery to remove one or both fallopian tubes and ovaries. It may treat cancer, an ovarian or tubal problem, or reduce cancer risk in people with certain inherited gene changes. Removing both ovaries causes permanent loss of ovarian hormone production.

## Salpingo-OöPhorectomy

Surgical removal of a fallopian tube and ovary, also called salpingo-oophorectomy.

## Salpingostomy

An operation that opens a fallopian tube to remove an ectopic pregnancy while leaving the tube in place. It may preserve fertility in selected cases, but persistent pregnancy tissue, bleeding, scarring, or another ectopic pregnancy can occur.

### Alternate Names:

Tubal Incision; Ectopic Tubostomy

## Salt

An ionic chemical compound; dietary sodium chloride is essential in small amounts but excess intake can increase blood pressure.

## Saltatory Conduction

The rapid movement of an electrical signal along a nerve fiber covered by myelin. The signal travels between small uncovered gaps in the covering, allowing fast communication with other nerves and muscles.

## Salter-Harris Fracture

A break involving the growth plate of a child or adolescent's bone. The fracture pattern helps predict the risk of disturbed growth; pain, swelling, and inability to use the limb require prompt assessment and follow-up.

---

## Salter-Harris Fractures

A pediatric orthopaedic classification system categorizing fractures involving the cartilaginous growth plate (physis): Type I (straight across physis), Type II (above physis into metaphysis), Type III (lower through epiphysis), Type IV (through metaphysis, physis, and epiphysis), and Type V (rammed/crushed physis).

## Salvage Therapy

Treatment used when a disease, especially cancer, has not improved with earlier treatment or has returned. The choice depends on the disease, previous treatments, overall health, and treatment goals.

## Sample Fixation

Preserving a tissue or cell sample soon after collection so it does not decay or change shape. The preserved sample can then be processed, stained, and examined under a microscope for diagnosis.

## Samter Triad

A clinical symptom complex (aspirin-exacerbated respiratory disease / AERD) characterized by the triad of bronchial asthma, chronic rhinosinusitis with recurrent nasal polyposis, and acute severe bronchospasm or naso-ocular reactions following the ingestion of aspirin or nonsteroidal anti-inflammatory drugs (NSAIDs).

### Alternate Names:

Aspirin-Exacerbated Respiratory Disease; AERD

## Sandfly Fever

An acute viral illness transmitted by phlebotomine sandflies, causing fever, headache, muscle pain, and sometimes rash.

## Sandhoff Disease

A rare inherited disorder in which fatty substances build up in nerve cells. It usually begins in infancy or childhood with loss of skills, weakness, seizures, movement problems, and sometimes an enlarged liver or spleen.

## Sandifer Syndrome

A pediatric paroxysmal dystonic movement disorder occurring in infants with severe gastroesophageal reflux disease (GERD) or hiatal hernia, characterized by episodes of spasmodic torticollis, lateral head tilting, and arching of the back (opisthotonos) during or shortly after feeding to alleviate heartburn.

### Alternate Names:

Sandifer's Syndrome; Gastroesophageal Reflux Dys-tonia

## Sandpaper Tooth

A tooth with a rough, grainy enamel surface that can hold plaque and feel uneven to the tongue. It may result from weak enamel, early mineral loss, fluorosis, or wear and should be checked by a dental professional.

## Sanfilippo Syndrome

A rare inherited disorder in which the body cannot break down part of a complex sugar. The buildup mainly affects the brain and causes progressive developmental and behavioral changes.

### Alternate Names:

Mucopolysaccharidosis Type III; MPS III

## Sanguineous

Containing or resembling fresh blood. The word often describes bright-red drainage from a wound; a large or rapidly increasing amount after surgery may indicate active bleeding.

## Sanitation

Public-health measures that safely manage human waste, water, food, hygiene, and environmental contamination to prevent infection and disease transmission.

## SA Node

A small group of specialized cells in the upper right heart chamber that normally starts each heartbeat. It produces regular electrical signals that spread through the heart and coordinate its rhythm.

---

## Sano Shunt

A temporary tube connecting the lower-right heart chamber to the lung artery during the first operation for some babies with severe left-sided heart defects. It sends blood to the lungs while the heart is reconstructed; blockage or abnormal flow can be dangerous.

### Alternate Names:

RV-PA Conduit; Norwood-Sano Modification

## Santorini Duct

An accessory pancreatic duct that drains the anterosuperior head of the pancreas directly into the duodenum via the minor duodenal papilla (located approximately 2 cm proximal to the major duodenal papilla of Vater).

### Alternate Names:

Accessory Pancreatic Duct; Duct of Santorini

## Saphenous

Relating to the superficial veins of the lower limb, especially the great or small saphenous vein.

## Saphenous Nerve

The largest cutaneous sensory terminal branch of the femoral nerve, descending through the adductor canal and piercing the deep fascia at the knee to travel alongside the great saphenous vein, supplying sensation to the medial lower leg, ankle, and medial foot arch.

## Saphenous Vein

A superficial vein of the lower limb that drains the foot and leg. The great and small saphenous veins are clinically important in venous disease and may be used as graft material in selected vascular or cardiac bypass procedures.

## SAPHO Syndrome

A rare inflammatory condition combining bone or joint pain with acne or pus-filled skin lesions. It often affects the breastbone, collarbone, spine, or jaw and can resemble infection or cancer, so diagnosis and treatment require specialist assessment.

## Sapovirus

A virus that can cause vomiting and watery diarrhea, especially in children. It spreads through contaminated food, water, surfaces, or close contact.

## Sapovirus Infection

An acute enteric infection caused by sapoviruses, positive-sense single-stranded RNA viruses in the family Caliciviridae that infect humans across all age groups. Clinical manifestations include acute watery diarrhea, nausea, vomiting, abdominal cramping, and low-grade fever, often responsible for outbreaks of foodborne or waterborne gastroenteritis in daycares, schools, and hospitals that resolve with supportive rehydration within 48 to 72 hours.

### Alternate Names:

Sapovirus Gastroenteritis; Human Sapovirus Infection

## Sapropterin

A synthetic form of tetrahydrobiopterin used for some people with phenylketonuria and certain tetrahydrobiopterin deficiencies; response and diet requirements vary.

## Saquinavir

An HIV protease inhibitor used with other antiretroviral medicines; drug interactions, gastrointestinal effects, and cardiovascular concerns limit its use.

## Saquinavir Mesylate

The mesylate salt form of saquinavir, an HIV medicine that blocks viral protein processing. It is used with other antiretroviral drugs, but interactions, liver problems, and dangerous heart-rhythm changes require careful monitoring.

## Sarcocystis Hominis

A protozoan parasite acquired by eating undercooked beef; people may develop intestinal symptoms, while cattle serve as an intermediate host.

## Sarcocystis Suihominis

A protozoan parasite acquired by eating undercooked pork that can cause intestinal sarcocystosis;

---

pigs serve as an intermediate host.

### **Sarcocystosis**

Parasitic infection by *Sarcocystis* species, acquired from undercooked meat or contaminated food and sometimes causing muscle pain, fever, diarrhea, or no symptoms.

### **Sarcoglycan Complex**

A group of proteins that helps strengthen the outer covering of muscle cells during movement. Inherited problems with these proteins can cause progressive muscle weakness and sometimes heart-muscle disease.

### **Sarcoglycanopathy**

A group of inherited muscle diseases caused by changes in sarcoglycan genes. They usually cause progressive weakness of the hips and shoulders and may affect the heart.

#### **Alternate Names:**

Sarcoglycan-Deficient Limb-Girdle Muscular Dystrophy; LGMD R3-R6

### **Sarcoid Arthropathy**

Joint inflammation associated with sarcoidosis. It may cause joint pain or swelling, often at the ankles during acute disease, and can occur with other sarcoidosis findings.

### **Sarcoidosis**

An inflammatory condition in which small clumps of immune cells, called granulomas, form in organs. The cause is unknown. It most often affects the lungs and lymph nodes in the chest but can also affect the skin, eyes, heart, nervous system, or other organs. Some people have no symptoms; others have symptoms that depend on which organs are affected.

#### **Alternate Names:**

Besnier-Boeck-Schaumann Disease; Boeck Sarcoid

### **Sarcoidosis Skin Manifestations**

Skin involvement in sarcoidosis can cause firm bumps, raised or discolored patches, scars that

change, or lupus pernio on the nose or face. The appearance varies and may require a skin examination or biopsy.

### **Sarcolemma**

The thin outer covering of a muscle cell. It carries electrical signals that start contraction and helps maintain the cell's structure and internal chemical balance.

### **Sarcoma**

A malignant tumor arising from connective or supporting tissues, including bone, cartilage, fat, muscle, and blood vessels. Examples include osteosarcoma, liposarcoma, and rhabdomyosarcoma.

### **Sarcomere**

The smallest working unit that makes skeletal and heart muscle contract. It contains protein threads that slide past one another, shortening the muscle and producing force.

### **Sarcopenia**

Loss of muscle strength and muscle mass that reduces physical function, often with aging but also with inactivity, poor nutrition, or chronic disease. It can cause weakness, slower walking, falls, and difficulty with daily activities.

### **Sarcopenic Obesity**

Excess body fat together with low muscle mass, strength, or physical ability. It can increase the risk of falls, disability, metabolic disease, and complications after surgery.

### **Sarcoplasm**

The material inside a muscle cell surrounding its specialized structures. It contains energy-producing parts, stored fuel, oxygen-holding myoglobin, and the protein machinery that produces contraction.

### **Sarcoplasmic Reticulum**

A network inside muscle cells that stores calcium. It releases calcium to start contraction and takes calcium back up so the muscle can relax.



---

## Sarcoptes

The mite genus containing *Sarcoptes scabiei*, the cause of scabies.

## Sarcoptes Scabiei

The microscopic mite that causes scabies. It burrows into the outer layer of skin and lays eggs, producing intense itching and a rash or thin lines. It spreads mainly through prolonged close skin contact.

## Sarcoptic Mange

Alternate name; see *Scabies*.

## Sargramostim

A recombinant granulocyte-macrophage colony-stimulating factor used to promote white-cell recovery after selected cancer treatments or transplantation; bone pain and fever may occur.

## Sarnat Staging

A three-level examination used to describe the severity of brain dysfunction in a newborn after reduced oxygen or blood flow. Mild disease causes increased alertness, moderate disease causes poor responsiveness and seizures, and severe disease causes coma and weak or absent reflexes; the stage helps guide urgent treatment and prognosis.

### Alternate Names:

Sarnat and Sarnat Grading; Sarnat Score; HIE Staging

## SARS

Severe acute respiratory syndrome, a serious lung infection caused by the SARS coronavirus. It can cause fever, cough, breathing difficulty, and pneumonia and spreads mainly through close contact with respiratory droplets; suspected cases require public-health and medical guidance.

## SARS-CoV-2

The coronavirus that causes COVID-19. It spreads mainly through particles released when an infected person breathes, talks, coughs, or sneezes and can cause anything from no symptoms to severe lung or other organ disease.

## SARS Virus

The coronavirus that causes severe acute respiratory syndrome, a serious respiratory infection that can spread through respiratory droplets and close contact.

## Sartorius Muscle

The longest muscle in the human body, a slender strap-like muscle of the anterior thigh originating from the anterior superior iliac spine (ASIS) and inserting into the pes anserinus on the proximal medial tibia; acts to flex, abduct, and laterally rotate the hip, and flex the knee ("tailor's muscle").

## Sassafras Oil Overdose

Poisoning from swallowing or otherwise being exposed to too much sassafras oil. It may cause nausea, vomiting, seizures, breathing problems, liver injury, or other serious effects.

## Satellite DNA

Repeated stretches of DNA found mainly in tightly packed parts of chromosomes. They help organize chromosome structure and can vary between people, usually without providing instructions for making proteins.

## Satellite Virus

A small virus-like infectious agent that can multiply only when a separate helper virus is present. It uses the helper's machinery and may change how severe or persistent the combined infection becomes.

## Satiation

The feeling of fullness that terminates a meal, produced by gastrointestinal stretch, nutrients, hormones, and brain signals regulating food intake.

## Satiety

The feeling of fullness that reduces the desire to eat after a meal. It is influenced by stomach stretching, food composition, hormones, and brain signals; persistent early fullness can indicate illness.

---

## Saturated Fat

A dietary fat whose fatty-acid chains contain no carbon-carbon double bonds. It is found in foods such as fatty meat, butter, full-fat dairy, coconut oil, and palm oil; replacing some saturated fat with unsaturated fat can help lower LDL cholesterol and cardiovascular risk.

## Saturation

A state in which available binding, transport, or processing capacity is nearly full. Once capacity is reached, extra medicine or substance may remain in the blood or produce a disproportionately larger effect.

## Sausage Digit

Diffuse swelling of an entire finger or toe, often caused by inflammation in joints, tendons, and surrounding tissues.

### Alternate Names:

Dactylitis

## Saxitoxin

A powerful poison made by some marine algae that can build up in shellfish. Eating contaminated seafood may cause tingling, weakness, trouble breathing, or paralysis.

## SBBO

Alternate name; see *Small intestinal bacterial overgrowth*.

## Scab

A crust of dried blood, serum, pus, or tissue fluid over a healing skin injury.

## Scabies

An infestation caused by tiny mites that burrow into the skin, causing intense itching that is often worse at night and a rash or thin burrows. It spreads mainly through prolonged close skin-to-skin contact.

### Alternate Names:

Sarcoptic Acariasis; Seven-Year Itch

## SCAD

Alternate name; see *Spontaneous coronary artery dissection*.

## Scaffold

A three-dimensional framework used in tissue engineering to hold cells in place while they grow and organize into new tissue. It may be made from natural or synthetic material and may gradually dissolve.

## Scaffold Seeding

Placing living cells onto or inside a three-dimensional framework so they can attach, multiply, and begin forming tissue in laboratory or regenerative-medicine research.

## Scalded Mouth Syndrome

Alternate name; see *Burning mouth syndrome*.

## Scalded Skin Syndrome

Alternate name; see *Scalded-Skin Syndrome*.

## Scalded-Skin Syndrome

A serious infection, usually in young children, in which toxins from *Staphylococcus* bacteria make the outer skin layer separate. The skin becomes painful, red, and easily blistered or peeled; prompt treatment is needed to prevent dehydration and infection.

## Scalds

Burns caused by hot liquids or steam. They may damage only the surface skin or extend into deeper tissue; severity depends on depth, size, location, and age. Large, deep, facial, genital, hand, or circumferential burns need urgent specialist assessment, and unusual patterns in children require safeguarding evaluation.

## Scalene Muscles

A group of three paired cervical muscles (anterior, middle, and posterior scalenes) originating from cervical transverse processes and inserting onto the first and second ribs; the interscalene triangle be-

---

tween the anterior and middle scalenes transmits the brachial plexus roots and subclavian artery (site of thoracic outlet syndrome).

### **Scales**

Scales are flakes or plates of shed skin on the surface, occurring with dryness, eczema, psoriasis, fungal infection, or other dermatologic conditions.

### **Scaling**

Scaling is flaking or shedding of the outer skin layer and may occur with dryness, eczema, psoriasis, fungal infection, or other skin disease.

### **Scaling And Root Planing**

A deep dental cleaning that removes hardened plaque and bacteria above and below the gumline and smooths tooth roots. It reduces gum inflammation and may slow periodontal disease, but severe damage may need additional treatment.

### **Scalp**

The skin and underlying tissues covering the skull. It contains hair follicles, glands, blood vessels, nerves, and connective tissue and can be affected by injury, infection, or skin disease.

### **Scalp and Body Ringworm**

Alternate name; see *Ringworm*.

### **Scalpel**

A sharp surgical knife used to make precise incisions or cuts in tissue. It may be a reusable handle with a detachable blade or a disposable single-piece instrument. Blade designs and numbers differ in size and shape; for example, a No. 10 blade is commonly used for larger incisions, a No. 11 for puncture or stab incisions, and a No. 15 for shorter precise cuts.

#### **Alternate Names:**

Surgical Knife; Lancette

### **Scalpel Handle**

The reusable handle that holds a detachable scalpel blade. Different handle and blade sizes are chosen for the type and precision of the incision.

### **Scalp Psoriasis**

Psoriasis affecting the scalp, causing sharply demarcated red plaques covered by silvery scale that may extend beyond the hairline. It can itch, crack, or bleed and is not contagious. Treatment may include medicated shampoos, topical corticosteroids, vitamin-D analogues, or systemic therapy for extensive disease.

#### **Alternate Names:**

Sebopsoriasis of Scalp; Scalp Plaque Psoriasis

### **Scalp Ringworm**

A contagious fungal infection of the scalp, also called tinea capitis. It can cause scaling, itching, broken hairs, bald patches, or swollen tender areas and usually needs prescription antifungal medicine taken by mouth to prevent persistent hair loss or spread.

### **Scalp Tenderness**

Pain when the scalp is touched or brushed. It can occur with giant cell arteritis, migraine, skin problems, nerve pain, or other conditions, so it is not diagnostic by itself.

#### **Alternate Names:**

Temporal Arteritis Scalp Allodynia; Cranial Hyperesthesia

### **Scan**

An imaging examination that creates pictures of body structures or functions using methods such as ultrasound, X-rays, CT, MRI, or radioactive tracers. The method and interpretation depend on the clinical question.

### **Scan Gallium**

A nuclear-medicine scan using radiolabeled gallium to identify areas of inflammation, infection, or some tumors; uptake is interpreted with anatomy and clinical context.

### **Scanning**

Systematic examination of a body region or specimen using imaging, microscopy, or another detection method to locate, characterize, or monitor abnormalities.

---

## Scanning Electron Microscopy

A method that scans a specimen's surface with an electron beam to produce a highly magnified, three-dimensional-looking image of its shape and surface details.

## Scanning Speech

Speech that is unusually slow, broken into separate syllables, and spoken with uneven rhythm or emphasis. It can occur when disease affects the cerebellum or other parts of the nervous system.

## Scanning Techniques

Methods used to create images or measurements of the body or a sample. Examples include ultrasound, X-ray, CT, MRI, and nuclear-medicine scans; the method is chosen according to the clinical question.

## Scaphoid

A small wrist bone on the thumb side, commonly injured when someone falls onto an outstretched hand. Pain in the hollow just behind the thumb may indicate a fracture, which sometimes heals poorly because part of the bone has limited blood flow.

## Scaphoid Abdomen

A scaphoid abdomen is an unusually hollowed abdominal contour, classically seen with severe diaphragmatic hernia in a newborn but also possible with marked wasting.

## Scaphoid Bone

The largest proximal-row carpal bone on the thumb side of the wrist; its waist is vulnerable to fracture and avascular necrosis because of its blood supply.

## Scaphoid Fractures

Breaks in the small wrist bone near the base of the thumb, often after falling onto an outstretched hand. Pain may be felt in the hollow beside the thumb; some fractures heal poorly because part of the bone has a limited blood supply.

## Scapholunate Advanced Collapse

The most common predictable pattern of wrist

osteoarthritis, resulting from untreated chronic scapholunate ligament dissociation or chronic scaphoid nonunion. Pathological kinematics cause rotatory subluxation of the scaphoid with progressive articular cartilage wear progressing through three distinct stages: radial styloid-scaphoid impingement (stage I), complete scaphoid facet arthritis (stage II), and capitolunate joint arthritis (stage III), while the radiolunate joint is characteristically spared.

### Alternate Names:

SLAC Wrist; Scapholunate Collapse Arthropathy

## Scaphycephaly

A long, narrow head shape caused when the major seam running from front to back of the skull closes too early. It is a form of craniosynostosis and may require specialist assessment to protect brain and skull development.

## Scapula

The shoulder blade, a flat triangular bone at the back of the chest. It connects the upper arm to the collarbone and provides attachment for muscles that move and stabilize the shoulder.

## Scapular Winging

Prominence of the medial border of the scapula lifting away from the posterior thoracic wall during forward arm elevation or pushing against a wall, classically caused by paralysis of the serratus anterior muscle due to long thoracic nerve injury.

## Scapulothoracic Joint

The functional gliding connection between the shoulder blade and the back of the chest wall. It is not a true joint but works with nearby joints to position and move the arm.

## Scar

Fibrous tissue formed during healing after injury, inflammation, or surgery.

## Scarlatina

Scarlet fever, an illness caused by toxin-producing group A *Streptococcus* and characterized by sore

---

throat, fever, and a fine rough-textured rash.

**Alternate Names:**

Scarlet Fever; Second Disease

**Scarlet Fever**

An illness caused by toxin-producing group A *Streptococcus*. It usually causes sore throat, fever, and a widespread fine, rough-textured rash; a red tongue and pale skin around the mouth may also occur.

**Scar Revision**

A procedure that improves the appearance, contour, flexibility, or symptoms of a scar through excision, rearrangement, grafting, laser, injection, or other selected techniques.

**Scarring Alopecia**

Permanent hair loss caused when inflammation or injury destroys hair follicles and replaces them with scar tissue. Early scalp redness, scaling, itching, pain, or spreading hair loss needs dermatologic assessment because lost follicles cannot regrow hair.

**Scars**

A scar is fibrous tissue formed during healing after injury, surgery, inflammation, or burns. Scars may be flat, depressed, raised, or contractured and can itch, hurt, restrict movement, or affect appearance; treatment depends on maturity and type.

**Scatter Correction**

A medical-imaging method that estimates and reduces errors caused when detected radiation changes direction after passing through tissue. It can improve the accuracy of a nuclear-medicine image or measured activity.

**Scatter Events**

Radiation particles that change direction after interacting with tissue or another material. In medical imaging they can place signals in the wrong location or reduce contrast, so scanners use design and correction methods to limit their effect.

**Scavenger Receptor**

A protein on or inside a cell that recognizes and helps remove altered fats, germs, cell debris, or dying cells. It supports early immune defense but can also help fatty material build up in artery walls.

**Scavenging System**

Equipment that collects and removes waste anesthetic gases from an operating room. It protects staff from repeated exposure and may use a vacuum system or passive venting to carry gases safely away.

**Scedosporium**

A genus of environmental molds that can cause difficult-to-treat lung, sinus, skin, bone, or disseminated infections, especially in immunocompromised people.

**Scedosporium Apiospermum**

A mold that can cause localized or disseminated infection, including mycetoma, pneumonia, sinusitis, and brain abscesses; treatment often requires specialist antifungal care.

**Scedosporium Prolifican**

An environmental mold, now commonly classified as *Lomentospora prolificans*, that causes severe disseminated infection and is often resistant to antifungal drugs.

**Schaalia Radingae**

A normally slow-growing anaerobic bacterium of the human mouth and skin that can rarely cause abscesses or other invasive infections.

**Schaalia Turicensis**

A filamentous anaerobic bacterium that may be part of normal flora but can cause actinomycosis-like abscesses and other opportunistic infections.

**Schatzki Ring**

A thin ring of tissue narrowing the lower esophagus, often near a hiatus hernia. It may cause occasional difficulty swallowing solid food or a piece of food becoming stuck; treatment can widen the narrowed

---

area.

**Alternate Names:**

Lower Esophageal Mucosal Ring; B Ring

**Scheuermann Disease**

A juvenile osteochondrosis of the spine in which vertebral end plates develop irregularities and anterior wedging during skeletal growth. It commonly produces a rigid thoracic kyphosis, although the thoracolumbar or lumbar spine may also be involved. Symptoms can include back pain, fatigue, limited spinal motion, and postural deformity that becomes more apparent during adolescence. Diagnosis is based on clinical examination and characteristic radiographs; management depends on deformity severity, skeletal maturity, pain, and progression.

**Scheuermann's Disease**

A growth disorder beginning in adolescence that causes several spinal bones to become wedge-shaped. This can produce a lasting rounded upper back, stiffness, pain, and sometimes breathing or activity limitations.

**Schiff Base**

A chemical compound formed when a primary amine reacts with an aldehyde or ketone; Schiff bases occur in biochemistry and drug chemistry.

**Schiller Test**

An examination of the cervix in which iodine solution is applied to highlight areas of abnormal tissue. Areas that do not stain may need further examination or biopsy; the test alone cannot diagnose cancer.

**Schilling Test**

An older test that measured how well the body absorbed vitamin B12, sometimes used to investigate pernicious anemia or intestinal disease. It is rarely performed now because blood tests and antibody testing have largely replaced it.

**Schirmer Test**

A test in which standardized filter paper strips measure tear production over a set time, commonly used

to evaluate aqueous-deficient dry eye.

**Schistocyte Count**

Measurement of fragmented red blood cells on a blood smear, supporting evaluation of microangiopathic hemolysis, mechanical valve damage, disseminated intravascular coagulation, or severe burns.

**Schistocytes**

Broken, irregular, or helmet-shaped red blood cells seen on a blood-smear test. They form when red cells are damaged in abnormal small vessels, by artificial heart valves, or in other illnesses and may signal serious red-cell destruction.

**Schistomicide**

A medicine that kills parasitic blood flukes responsible for schistosomiasis. Treatment choice depends on the parasite species and the organs affected, and medical advice is needed because infection may cause lasting damage.

**Schistosoma**

A group of parasitic worms that live in blood vessels. Their young forms can enter through skin exposed to contaminated fresh water and later cause intestinal, urinary, liver, or other disease.

**Schistosoma Haematobium**

A parasitic blood fluke that can cause urinary schistosomiasis, including hematuria and chronic injury of the urinary tract.

**Schistosoma Intercalatum**

A parasitic blood fluke that causes intestinal schistosomiasis, mainly in parts of Central and West Africa, after freshwater exposure.

**Schistosomiasis**

An infection caused by parasitic worms called *Schistosoma*. The worms live in freshwater snails, and their larvae can penetrate the skin when a person swims, wades, or works in contaminated freshwater. The infection may cause an itchy rash, fever, abdominal pain, diarrhea, blood in the stool or urine, and



---

painful urination. Long-term infection can damage the intestine, liver, urinary tract, or other organs.

### **Schistosomiasis (Bilharzia)**

Alternate name; see *Schistosomiasis*.

#### **Alternate Names:**

Snail Fever; Katayama Fever

### **Schizencephaly**

A rare condition present from birth in which a fluid-filled cleft extends through part of the brain. It results from abnormal early brain development and may cause seizures, weakness, movement problems, or delayed development, depending on the cleft's size and location.

### **Schizoaffective Disorder**

A mental-health disorder in which a person has symptoms of schizophrenia, such as hallucinations or disorganized thinking, together with major depression or mania. Psychotic symptoms also occur for a period without a major mood episode.

### **Schizogony**

A way some single-celled parasites reproduce without mating. One parasite cell grows and divides internally to produce many new cells, which are then released to infect other cells.

### **Schizoid Personality Disorder**

A long-standing pattern of limited interest in close relationships and restricted emotional expression. It differs from schizophrenia because hallucinations and fixed delusions are not central features.

#### **Alternate Names:**

Schizoid Disorder; Asocial Personality

### **Schizophrenia**

A serious, long-term mental disorder that can affect a person's thoughts, perceptions, emotions, behavior, and ability to function. Symptoms are often grouped as psychotic, negative, and cognitive. Psychotic symptoms include hallucinations, such as hearing or seeing things that others do not, delusions, which are firmly held beliefs not supported

by shared evidence, and disorganized thought or speech. Negative symptoms may include reduced emotional expression, motivation, pleasure, or social interest. Cognitive symptoms may affect attention, memory, planning, and decision-making. Symptoms and their severity vary between people and may come and go or persist over time.

### **Schizotypal Personality Disorder**

A long-term pattern of discomfort with close relationships, unusual beliefs or perceptions, unusual speech or behavior, and difficulty interpreting social situations. It is different from schizophrenia because persistent hallucinations and major loss of reality testing are not typical.

#### **Alternate Names:**

Schizotypal Disorder; STPD

### **Schlemm Canal**

A circular, endothelium-lined vessel situated in the iridocorneal angle of the anterior chamber, which drains aqueous humor from the trabecular meshwork into the collector channels and episcleral venous system.

#### **Alternate Names:**

Canal of Schlemm; Scleral Venous Sinus

### **Schlemm's Canal**

A circular drainage channel at the edge of the cornea that carries fluid from the front of the eye into blood vessels. Poor drainage can raise pressure inside the eye and contribute to glaucoma.

#### **Alternate Names:**

Canal of Schlemm

### **Schmorl Node**

An intravertebral disc herniation characterized by the protrusion and displacement of nucleus pulposus cartilage through a defect in the cartilaginous vertebral endplate into the trabecular bone marrow of an adjacent vertebral body.

#### **Alternate Names:**

Schmorl's Node; Intravertebral Herniation

---

## Schnitzler Syndrome

A rare inflammatory disease causing a long-lasting hives-like rash, repeated fever, bone or joint pain, and raised inflammation tests. It is usually associated with an abnormal IgM blood protein and requires specialist evaluation.

## School-Age Children Development

School-age development includes advances in growth, coordination, language, learning, independence, social relationships, and emotional regulation, with broad normal variation.

## School-Based Rehabilitation

Therapy and other support services provided in a school setting to help a child participate in education and daily school activities.

## School Sores

Alternate name; see *Impetigo*.

## Schwann Cell

A supporting cell of the peripheral nervous system. It wraps around many nerve fibers to form their insulating myelin covering and helps maintain or repair peripheral nerves.

## Schwann Cells

Glial cells of the peripheral nervous system that form myelin around many axons and support the repair and maintenance of peripheral nerves.

## Schwannoma

A usually noncancerous, slow-growing tumor arising from cells that support nerves. It may cause a lump, pain, numbness, or weakness; a tumor involving the hearing and balance nerve can cause hearing loss, ringing, or dizziness.

### Alternate Names:

Neurilemmoma

## Schwannomatosis

A condition in which multiple schwannomas, usually slow-growing tumors of the covering of nerves, develop. The main problem is often long-lasting

pain, although numbness, weakness, or other nerve symptoms can occur depending on tumor location.

### Alternate Names:

Neurinomatosis

## SCI

Alternate name; see *Spinal cord injury*.

## Sciatica

Pain, tingling, numbness, or weakness that travels from the lower back or buttock down the leg, usually because a nerve root is irritated by a disk problem or narrowing of the spine. Progressive weakness, numbness around the inner thighs or genitals, or new loss of bladder or bowel control may indicate severe nerve compression.

### Alternate Names:

Lumbar Radiculopathy; Sciatic Neuralgia

## Sciatic Nerve

The body's largest nerve, running from the lower spine through the buttock and down the back of the leg. It supplies movement and sensation to much of the leg and foot.

## Sciatic Neuritis

Alternate name; see *Sciatica*.

## Scimitar Syndrome

A birth defect in which some veins from the right lung drain to the wrong large vein instead of directly to the heart. The right lung may be smaller, and the condition can cause repeated chest infections, breathing problems, high pressure in the lung arteries, or heart failure.

## Scintigraphy

Nuclear-medicine imaging in which a small amount of radioactive tracer is given and a camera records where it collects. The pattern can show the function, blood flow, or activity of organs such as bone, thyroid, kidneys, or heart.

## Scintillation

The brief release of light by a material after it ab-

---

sorbs radiation. Detectors measure these flashes to estimate radiation and create images in some medical scans.

### Scintillation Crystal

A material in some medical detectors that changes incoming radiation into tiny flashes of light. The flashes are measured to create images or estimate the amount of radioactive tracer present.

### Scintillations

Flashes, sparkles, or zigzag lines seen in the field of vision. They often occur during a migraine aura but may also result from an eye or brain problem; sudden new flashes with floaters or a curtain-like shadow need urgent eye assessment.

### Scirrhus

An older term for a hard cancer containing a large amount of scar-like tissue. It was used especially for some breast or stomach cancers and describes texture rather than a separate cancer type.

### Scissors

Instruments with two blades used to cut tissue, sutures, dressings, or other material. Surgical types vary in length, blade shape, and sharpness; examples include Mayo and Metzenbaum scissors.

### Sclera

The sclera is the tough white outer coat of the eye that protects the globe and provides attachment for eye muscles.

### Scleral Buckling

A retinal-detachment repair procedure in which a silicone band or buckle is placed around the outside of the eye to support the retinal break and reduce traction.

### Scleral Icterus

A yellow pigmentation of the ocular conjunctiva and sclera caused by the accumulation of bilirubin in tissues with high elastin content. It becomes clinically detectable when serum total bilirubin levels

exceed 2.5 to 3.0 mg/dL (43 to 51 micromol/L) and serves as an early physical sign of jaundice and liver or biliary disease.

### Scleral Inflammation

Alternate name; see *Scleritis*.

### Scleredema

A sclerosing skin disorder causing thick, hard, often waxy skin, classically on the neck, upper back, and shoulders. It may follow infection, occur with diabetes or a monoclonal gammopathy, or rarely be associated with other systemic disease.

### Scleredema Diabeticorum

Scleredema diabeticorum is thickening and hardening of the skin on the upper back and neck, usually associated with long-standing diabetes and obesity.

### Scleritis

A severe, painful inflammation of the sclera (the tough, white outer wall of the eye). It is characterized by a deep, boring eye ache that often radiates to the forehead or temple and worsens at night, accompanied by intense redness with a bluish or purple hue. Unlike mild surface eye redness (episcleritis), scleritis affects the deeper protective coat of the eye and is frequently linked to autoimmune conditions such as rheumatoid arthritis or lupus. Severe or untreated inflammation can cause thinning of the eye wall and impair vision.

#### Alternate Names:

Deep Scleral Inflammation; Anterior Scleritis; Posterior Scleritis; Scleral Inflammation

### Sclerodactyly

A localized, progressive thickening, tightening, and induration of the skin of the fingers and toes, often resulting in tapered, shiny digits, loss of normal skin folds, and flexion contractures. It is a hallmark cutaneous feature of systemic sclerosis (scleroderma) and CREST syndrome.

### Scleroderma

A group of autoimmune diseases that cause inflammation and too much collagen to build up, making

---

skin and sometimes deeper tissues hard, thick, and tight. Localized scleroderma, also called morphea or linear scleroderma, mainly affects the skin and tissues just beneath it. Systemic scleroderma, also called systemic sclerosis, can affect the skin, small blood vessels, and internal organs such as the esophagus, lungs, kidneys, and heart. The cause is not fully known, and symptoms vary with the type and organs involved.

**Alternate Names:**

Systemic Sclerosis; SSc

### **Scleroderma Renal Crisis**

A life-threatening complication of systemic sclerosis (predominantly diffuse cutaneous subtype), characterized by the abrupt onset of severe accelerated hypertension, acute oliguric or anuric renal failure, microangiopathic hemolytic anemia, and thrombocytopenia. Driven by endothelial injury and dramatic renin-angiotensin-aldosterone axis hyperactivation; treated urgently with angiotensin-converting enzyme (ACE) inhibitors.

**Alternate Names:**

SRC; Systemic Sclerosis Renal Crisis

### **Scleroma**

A long-lasting infection, usually of the nose and upper airway, caused by *Klebsiella rhinoscleromatis*. It can produce scar tissue, blocked breathing, nasal discharge, voice changes, or facial deformity.

### **Scleromyxedema**

A rare long-term skin disorder in which many small, firm, waxy bumps and thickened skin develop, especially on the face, neck, and hands. It may also affect muscles, joints, swallowing, lungs, or other organs.

### **Sclerosing Cholangitis**

Inflammation and scarring that narrow or block the tubes carrying bile from the liver. It may cause itching, jaundice, abdominal pain, repeated infections, and liver damage; the cause guides treatment and follow-up.

### **Sclerosing Mesenteritis**

A rare inflammatory condition causing thickening and scarring in the tissue that supports the intestines. It may cause abdominal pain, nausea, weight loss, or bowel blockage, although some people have no symptoms.

### **Sclerosing Peritonitis**

Thickening and scarring of the lining inside the abdomen, sometimes after long-term peritoneal dialysis. The scar tissue can restrict the intestines and cause pain, vomiting, poor nutrition, or bowel blockage.

### **Sclerosing Sialadenitis**

Long-term inflammation that makes a salivary gland swollen and hard because scar-like tissue forms inside it. It may be related to an immune disorder and can cause dry mouth, pain, or repeated swelling.

### **Sclerosis**

Abnormal hardening or scarring of tissue, often from excess connective tissue, chronic inflammation, degeneration, or neurologic plaques; the affected organ determines the specific disease.

### **Sclerosteosis**

A rare inherited condition in which bones, especially the skull, become abnormally thick. It may cause a broad or tall build, joined fingers, headaches, hearing loss, facial weakness, or pressure on nerves inside the skull.

### **Sclerotherapy**

A treatment in which an irritating or closing agent is injected into a blood vessel or body cavity to make it scar and seal, commonly used for varicose veins and selected vascular lesions.

### **Sclerotomy**

A sclerotomy is a surgical opening made through the sclera, the white outer layer of the eye, often during vitreoretinal surgery.

---

## Scoliosis

An abnormal sideways curve of the spine that may look like the letter “C” or “S.” The bones of the spine may also rotate, changing the shape of the back or rib cage. In many children and teenagers, the cause is unknown; this is called idiopathic scoliosis. Other causes include differences present at birth, nerve or muscle conditions, spinal injury, or age-related changes. Mild scoliosis may cause no symptoms, while larger curves may cause uneven shoulders or hips, a prominent shoulder blade, back pain, reduced movement, or, rarely, breathing problems.

### Alternate Names:

Spinal Curvature

## Scombroid Poisoning

A foodborne intoxication caused by the ingestion of improperly refrigerated dark-flesh scombroid fish (such as tuna, mackerel, bonito, and mahi-mahi). Bacterial contamination by species possessing histidine decarboxylase converts endogenous muscular histidine into high levels of heat-stable histamine; manifests within minutes with a sharp, peppery taste, intense facial flushing, throbbing headache, palpitations, urticaria, and abdominal cramps.

### Alternate Names:

Histamine Fish Poisoning; Scombrototoxicity

## Scopolamine

Scopolamine is a muscarinic antagonist used mainly to prevent motion sickness and postoperative nausea; dry mouth, blurred vision, urinary retention, confusion, and other anticholinergic effects may occur.

## Scopolamine Hydrobromide

The hydrobromide salt of scopolamine, an anticholinergic medicine used mainly to prevent motion sickness and postoperative nausea.

## Scopulariopsis

Scopulariopsis is a mold genus that rarely causes opportunistic infection, especially in people with immune suppression or implanted devices.

## Scorbutic

Relating to scurvy caused by severe vitamin C deficiency. It can cause easy bruising, bleeding gums, poor wound healing, anemia, bone or joint pain, and weakness; treatment with vitamin C and an adequate diet usually leads to recovery.

## Scorpion Fish Sting

A painful injury from the venomous spines of a scorpion fish, causing severe local pain and swelling and sometimes whole-body symptoms. Spines or venom may remain in the injury.

## Scorpions

Venomous arachnids whose stings can cause local pain and, depending on the species, sweating, muscle spasms, breathing problems, or other serious symptoms.

## Scorpion Sting

A venomous injury from a scorpion, usually causing immediate pain, burning, redness, or tingling. Some species can cause muscle twitching, abnormal eye movements, breathing difficulty, or dangerous autonomic symptoms.

### Alternate Names:

Scorpion Envenomation; Scorpion Bite

## Scorpion Stings

Alternate name; see *Scorpion sting*.

## Scotoma

A blind or blurred spot within an otherwise usable field of vision. It may result from retinal disease, optic-nerve damage, migraine, stroke, or another eye or brain disorder.

## Scotty Dog Sign

The normal radiological configuration of the posterior vertebral elements on oblique lumbar spine radiographs resembling a Scottish terrier dog; a radiolucent line crossing the neck of the dog indicates a defect or fracture in the pars interarticularis (spondylolysis).

### Alternate Names:

---

Scottish Terrier Sign; Pars Interarticularis Collar Sign

### **Scrape**

A scrape is a superficial abrasion in which friction removes part of the skin surface; cleaning, protection, and monitoring for infection support healing.

### **Screening**

Testing people without symptoms to identify a disease or an increased risk of disease. The appropriate test, age range, interval, and follow-up depend on the condition, individual risk, and current guidance; screening benefits must be balanced against harms.

### **Screening And Diagnosis For HIV**

Testing for human immunodeficiency virus using an antigen-antibody assay or nucleic-acid test, followed by confirmatory testing and assessment of immune status when positive.

### **Screening Procedure**

A test or examination offered to people without symptoms to detect disease or risk early enough for effective prevention or treatment; eligibility and intervals depend on evidence and risk.

### **Screening Program**

An organized public-health service that systematically invites a defined population to undergo testing for an asymptomatic condition, with quality-assured follow-up, diagnosis, treatment, and evaluation of benefits and harms.

### **Screening Study**

A planned investigation of an asymptomatic population to identify early disease, precursor lesions, or risk factors using a validated test and follow-up pathway.

### **Screening Test**

A test applied to people without symptoms or a known diagnosis to identify those at increased risk of a condition. A screening result usually requires a diagnostic or confirmatory evaluation before treat-

ment is initiated.

### **Screen Time And Children**

Time children spend using televisions, computers, phones, tablets, or game devices; excessive or poorly timed use can affect sleep, activity, learning, mood, and family routines.

### **Screws**

Threaded metal devices used to hold broken bones together or attach plates and other implants securely to bone.

### **Script**

A script is a written set of instructions or a prescription; in healthcare, a prescription specifies the medicine, dose, route, timing, and duration.

### **Scrivener's Palsy**

An older term for task-specific hand dystonia or cramp occurring during writing.

### **Scrotal Calcinosis**

Firm, usually painless calcium deposits in the skin of the scrotum that appear as one or more whitish or yellowish lumps. They may become tender, ulcerate, or release a chalky material; removal is considered if they cause symptoms.

### **Scrotal Infection**

Infection of the scrotal skin or underlying structures, which may cause redness, swelling, warmth, pain, or drainage. Rapidly worsening pain, fever, discoloration, or severe swelling is an emergency because serious soft-tissue infection must be excluded.

### **Scrotal Masses**

Lumps or swellings in the scrotum, arising from the skin, fluid around a testicle, veins, infection, hernia, or a testicular tumor. A new, firm, painless, or rapidly growing lump needs prompt examination and ultrasound.

### **Scrotal Pain**

Pain in the scrotum caused by infection, trauma,



---

hernia, fluid collections, varicocele, testicular torsion, or other conditions affecting the testes and surrounding structures.

### **Scrotal Septum**

A tissue wall that divides the scrotum into right and left compartments.

### **Scrotal Swelling**

Enlargement of the scrotum caused by fluid, inflammation, infection, hernia, varicocele, hydrocele, tumor, trauma, or testicular torsion; sudden pain is an emergency.

### **Scrotal Ultrasound**

Ultrasound imaging of the scrotum and testes used to evaluate pain, swelling, masses, torsion, infection, trauma, and blood flow.

### **Scrotum**

The scrotum is the external skin-and-muscle sac containing the testes and epididymides; its muscles help regulate testicular temperature, and swelling or pain may indicate infection, torsion, hernia, or tumor.

### **Scrub In**

To scrub in is to perform surgical hand antisepsis and put on sterile clothing and gloves before participating in a sterile procedure.

### **Scrubs**

Scrubs are protective work garments commonly worn by healthcare staff; they reduce clothing contamination but do not replace hand hygiene or other precautions.

### **Scrub Typhus**

An infection spread by tiny mites called chiggers. It can cause high fever, headache, swollen lymph nodes, rash, and sometimes a dark sore at the bite site. Severe illness can affect the lungs, kidneys, heart, or brain and needs prompt treatment.

#### **Alternate Names:**

Tsutsugamushi Disease; Mite-Borne Typhus

### **Scrub Up**

To scrub up means to clean and antiseptically prepare the hands and forearms before a sterile procedure.

### **Scurf**

Fine flakes or scales shed from the skin, especially the scalp. It may occur with dry skin, dandruff, eczema, psoriasis, infection, or irritation and may be accompanied by itching or redness.

### **Scurvy**

A disease caused by severe vitamin C deficiency. It can cause bleeding or swollen gums, easy bruising, small skin bleeds, poor wound healing, anemia, bone or joint pain, and fatigue.

### **Seabather's Eruption**

An itchy papular or vesicular eruption caused by contact with stinging larvae in seawater, often beneath swimwear. Symptoms may begin after leaving the water and usually resolve over days, though severe reactions may need medical treatment.

### **Sea-Blue Histiocyte**

An immune cell that appears blue-green under a microscope because it contains stored fats or cell-waste pigment. It may be seen in some storage, blood, or metabolic disorders, but the finding alone does not identify a disease.

### **Seafood Poisoning**

Clinical toxic syndromes resulting from the ingestion of seafood containing marine toxins or accumulated bio-toxins, including ciguatera fish poisoning, scombroid (histamine) toxicity, paralytic shellfish poisoning, and tetrodotoxin poisoning.

#### **Alternate Names:**

Marine Toxin Poisoning; Shellfish Toxicity; Ichthyosarcotoxism

### **Sea-Sickness**

Motion sickness caused by conflicting signals from the inner ear, eyes, and body during travel on a boat or ship. It may cause nausea, vomiting, dizziness,

---

sweating, headache, and pallor.

### Seasonal Affective Disorder

Depression that repeatedly begins during a particular season, most often autumn or winter, and improves at another time of year. Symptoms may include low mood, loss of interest, tiredness, oversleeping, increased appetite, and social withdrawal.

### Seasonal Affective Disorder (SAD)

Alternate name; see *Seasonal Affective Disorder*.

### Seasonal Affective Disorder Syndrome

Alternate name; see *Seasonal Affective Disorder*.

### Seasonal Allergies

An allergic reaction that occurs at specific times of the year when outdoor plants release tiny pollen grains into the air. In spring, tree pollen (such as oak, birch, and cedar) is the most common trigger; in summer, grass pollen causes symptoms; and in late summer and autumn, weed pollen (especially ragweed) and outdoor mold spores are common culprits. When a sensitized person inhales pollen, the immune system mistakenly identifies it as a threat, causing the release of histamine and other chemicals that inflame the nasal passages, sinuses, and eyes. Common symptoms include frequent sneezing, a congested or runny nose with clear drainage, an itchy throat or roof of the mouth, and red, itchy, watery eyes (allergic conjunctivitis). Treatment includes tracking pollen counts, keeping windows closed during high-pollen days, using daily corticosteroid nasal sprays, taking oral antihistamines to control itching and sneezing, and rinsing nasal passages with saline.

#### Alternate Names:

Hay Fever; Seasonal Allergic Rhinitis; Pollinosis; Seasonal Allergy

### Sebaceous Adenoma

A sebaceous adenoma is a usually benign tumor of sebaceous glands that can appear as a yellowish skin nodule and may be associated with Muir-Torre syndrome.

### Sebaceous Carcinoma

An uncommon cancer arising from sebaceous, or oil-producing, glands. It often appears as a firm or yellowish eyelid or skin nodule and may require biopsy and specialist treatment.

### Sebaceous Cyst

A common but imprecise term for an epidermoid cyst, a keratin-filled cyst beneath the skin.

### Sebaceous Epithelioma

A benign sebaceous neoplasm, often presenting as a yellowish papule or nodule. Multiple or unusual lesions may warrant assessment for a DNA mismatch-repair syndrome such as Muir-Torre syndrome.

### Sebaceous Gland

A small gland in the skin that secretes sebum (oil) into hair follicles to lubricate and waterproof the skin and hair.

### Sebaceous Gland Carcinoma

Sebaceous gland carcinoma is an uncommon skin cancer arising from oil glands, often appearing as a firm yellowish eyelid or skin lump.

### Sebaceous Glands

Skin glands that produce sebum, an oily substance usually released into hair follicles. They are most abundant on the face and scalp and contribute to acne when blocked or inflamed.

### Sebaceous Hyperplasia

A benign enlargement of sebaceous glands, presenting as small yellowish papules with central umbilication, usually on the face of adults.

### Seborrhea

Seborrhea is excessive or altered oil production that contributes to greasy scale and inflammation, commonly in seborrheic dermatitis of the scalp, face, or chest. Gentle cleansing, medicated shampoos, and topical anti-inflammatory or antifungal treatment may help.

---

## Seborrheic Dermatitis

A common long-term inflammation of oily skin, especially the scalp, eyebrows, sides of the nose, chest, and skin folds. It causes red or pink patches with greasy white or yellow scale; dandruff is a mild scalp form.

### Alternate Names:

Seborrhea; Dandruff; Cradle Cap

## Seborrheic Keratoses

Common, benign, epidermal skin tumors presenting as waxy, stuck-on, pigmented plaques with a rough, verrucous surface, typically appearing on the trunk and face of middle-aged and older adults. They are the most common benign skin neoplasm and are caused by keratinocyte proliferation.

## Seborrheic Keratosis

A benign epidermal tumor appearing as stuck-on, waxy, hyperpigmented plaques with a verrucous surface. They are extremely common in older adults and increase in number with age. Seborrheic keratoses are not premalignant.

### Alternate Names:

Basal Cell Papilloma; Senile Wart

## Seborrhoea

A chronic inflammatory skin condition affecting sebum-rich areas: scalp, face (nasolabial folds, eyebrows), chest, and intertriginous areas. See Seborrheic Dermatitis.

## Seborrhoeic

Relating to increased or altered activity of sebaceous glands or to areas rich in these glands, such as the scalp, face, chest, and upper back.

## Seborrhoeic Keratosis

A common harmless skin growth that may be brown, black, or light-colored with a waxy, rough, or stuck-on appearance. It is not contagious or cancerous, but changing lesions should be checked.

## Sebum

An oily substance made by skin glands that lu-

bricates and protects the skin and hair. Excess production can contribute to oily skin, acne, and dandruff, while changes may accompany some skin disorders.

## Seckel Syndrome

A rare inherited condition causing severe growth restriction before and after birth, a very small head, characteristic facial features, and often developmental or intellectual disability. Bone, blood, and other birth differences may also occur.

## Secobarbital Overdose

Poisoning from taking too much secobarbital, a sedative medicine. It can cause extreme sleepiness, confusion, slowed breathing, coma, or death.

## Secondary Adrenal Insufficiency

Reduced production of adrenal hormones because the pituitary or brain does not provide enough stimulation, often after long-term steroid treatment is stopped suddenly. It can cause fatigue, weakness, nausea, low blood pressure, and low blood sugar; skin darkening and salt loss are less typical than in primary adrenal failure.

## Secondary Amenorrhea

Absence of menstruation for at least three months in someone with previously regular cycles or six months with previously irregular cycles, caused by pregnancy, endocrine, ovarian, uterine, or systemic factors.

## Secondary Attack Rate

The proportion of susceptible people who become infected after contact with an initial case during a defined period. It helps estimate how readily an infection spreads in households, schools, healthcare settings, or other close-contact groups.

## Secondary Brain Cancer

Alternate name; see *Brain metastases*.

## Secondary Cancer

A secondary cancer is a tumor that has spread from

---

its original site to another part of the body. It is also called a metastasis or metastatic cancer.

### **Secondary Hyperparathyroidism**

Overactivity of the parathyroid glands in response to low calcium, abnormal phosphate, vitamin-D deficiency, or kidney disease. It can weaken bones, cause bone pain, and contribute to calcium deposits in blood vessels.

### **Secondary Hypertension**

High blood pressure caused by an identifiable medical condition, medicine, or substance, rather than primary (essential) hypertension. Causes include kidney disease or narrowed kidney arteries, hormone disorders, sleep apnea, pregnancy-related conditions such as preeclampsia, and certain medicines or drugs, including steroids, stimulants, decongestants, and some pain relievers. Treating the cause may improve blood pressure.

### **Secondary Hypothyroidism**

Low thyroid-hormone production because the pituitary or nearby brain area does not provide enough thyroid-stimulating signal. It may cause tiredness, feeling cold, constipation, dry skin, slow heart rate, or menstrual changes.

### **Secondary Immune Response**

A faster and stronger defense when the body meets a germ or foreign substance again. Memory immune cells from the first exposure respond quickly and usually produce antibodies and other defenses more effectively.

### **Secondary Immunodeficiency**

Reduced immune function caused by an acquired condition or exposure rather than a primary genetic disorder. Causes include HIV infection, cancer, malnutrition, medicines, and chronic disease; the manifestations depend on which immune functions are impaired.

### **Secondary Infections**

Infections that develop during or after another disease, injury, or treatment because of tissue damage,

altered defenses, or immunosuppression.

### **Secondary Lymphoid Organs**

Organs where mature lymphocytes meet foreign substances and begin immune responses. They include lymph nodes, the spleen, tonsils, and other lymphoid tissues.

### **Secondary Osteoporosis**

Bone loss caused by another illness, medication, or prolonged immobility.

### **Secondary Osteosarcoma**

An osteogenic sarcoma arising in previously abnormal or treated bone, such as bone affected by Paget disease, prior radiation, or chronic bone lesions; prognosis and treatment depend on stage and resectability.

### **Secondary Parkinsonism**

Secondary parkinsonism is slowed movement, rigidity, tremor, or postural instability caused by another condition, medicine, toxin, vascular injury, or structural brain disease.

### **Secondary Peritonitis**

Diffuse or localized acute bacterial or chemical inflammation of the peritoneal cavity resulting from direct breach of an intra-abdominal hollow viscus, such as a perforated peptic ulcer, ruptured appendix, ischemic bowel perforation, or anastomotic leak.

#### **Alternate Names:**

Perforated Viscus Peritonitis; Acute Secondary Peritoneal Inflammation; Surgical Peritonitis

### **Secondary Polycythemia**

An absolute increase in red blood cell mass driven by elevated circulating erythropoietin levels, either as an appropriate physiological response to chronic systemic hypoxia (chronic cyanotic lung/heart disease, high altitude, smoking) or pathologically from erythropoietin-secreting tumors (renal cell carcinoma, cerebellar hemangioblastoma, hepatocellular carcinoma).

---

## Secondary Postpartum Hemorrhage

Heavy or abnormal bleeding occurring more than 24 hours and up to 12 weeks after childbirth. Causes include retained placental tissue, infection, poor healing where the placenta was attached, or a bleeding disorder. Heavy bleeding, dizziness, fainting, fever, or severe pain may occur.

### Alternate Names:

Late Postpartum Hemorrhage; Delayed PPH

## Secondary Prevention

Finding and treating a disease early to stop it worsening or causing complications. Examples include screening for some cancers, checking blood pressure, testing for early diabetes, and treating an abnormality before it produces serious symptoms.

## Secondary Progressive Multiple Sclerosis

A phase of multiple sclerosis in which disability gradually worsens over time after an earlier relapsing pattern. Relapses may still occur, but progressive walking, strength, vision, bladder, thinking, or other problems can develop even between attacks.

### Alternate Names:

SPMS; Secondary-Progressive MS

## Secondary Sexual Characteristics

Physical changes that develop during puberty and are not directly part of the reproductive organs, such as breast development, facial or body hair, voice changes, and changes in body shape.

## Secondary Shock

Circulatory failure occurring as a consequence of another condition, such as infection, trauma, bleeding, heart failure, allergy, or endocrine crisis, with inadequate tissue perfusion.

## Secondary Syphilis

The stage of syphilis that occurs weeks or months after the first sore when the bacteria spread through the body. It may cause a widespread rash, including the palms and soles, swollen glands, fever, tiredness, mouth patches, or moist lesions and requires treatment.

## Secondary Systemic Amyloidosis

Secondary systemic amyloidosis results when chronic inflammation or infection drives production of serum amyloid A, which deposits in organs such as the kidneys, liver, and spleen.

## Secondary Tumor

A tumor formed when cancer cells spread from the original site to another part of the body.

## Second-Degree AV Block

A heart-rhythm problem in which some electrical signals from the upper chambers fail to reach the lower chambers. The heartbeat may be slow or irregular; certain patterns can progress to complete heart block and need urgent assessment.

## Second-Degree Burn

A burn that damages the outer skin and part of the layer beneath it. The area is usually painful, red, moist, and blistered; deeper burns may look pale and heal slowly. Large, deep, or sensitive-area burns need medical assessment.

## Second-Hand Smoke

Smoke inhaled from burning tobacco products or exhaled by a person who is smoking. It contains toxic substances and increases health risks in exposed people.

## Second-Look Surgery

Second-look surgery is a planned or follow-up operation performed to inspect healing, remove remaining disease, assess treatment response, or manage complications.

## Second Malignancy

A new, separate cancer that develops after an earlier cancer. The risk may be influenced by inherited factors, previous radiation or chemotherapy, age, and other exposures; it is different from recurrence of the first cancer.

## Second Messengers

Small molecules inside cells that pass along mes-



---

sages received from hormones, nerve chemicals, or other outside signals. They help control contraction, secretion, metabolism, gene activity, and cell growth.

### **Second Messenger Systems**

Internal signaling systems that carry a message from a hormone, nerve chemical, or growth factor at the cell surface to processes inside the cell. They can change contraction, secretion, metabolism, gene activity, or cell growth.

### **Second Opinion**

An independent medical assessment by another qualified clinician to help confirm or clarify a diagnosis, prognosis, or treatment plan.

### **Second Stage of Labor**

The part of childbirth that begins when the cervix is fully open and ends when the baby is born. It includes descent of the baby and pushing; its length varies with previous births, pain relief, the baby's position, and the health of parent and baby.

### **Second Trimester**

The middle part of pregnancy, from 14 through 27 completed weeks. The fetus grows rapidly, movement is often felt, and ultrasound and blood screening may assess development and certain conditions; the timing and tests vary by pregnancy and local care.

### **Secretin**

A hormone released by the first part of the small intestine when acidic stomach contents enter it. Secretin prompts the pancreas and bile ducts to release alkaline fluid, helping neutralize acid and support digestion.

### **Secretin Stimulation Test**

A test in which secretin is given and changes in blood or digestive fluid are measured. It may assess pancreatic function or investigate a rare tumor that releases excess gastrin and stomach acid.

### **Secretion**

The release of a substance from cells or glands into the blood, a body passage, or outside the body. Examples include saliva, hormones, digestive juices, mucus, and substances removed into urine or bile.

### **Secretory Diarrhea**

Watery diarrhea caused when the intestine releases too much water and salt or absorbs too little. It may continue even without eating and can result from infection, medicines, bile-acid problems, hormone-producing tumors, or inflammation.

### **Section**

A defined part or subdivision of an organ, document, study, or procedure; in anatomy and imaging, the term identifies a specific slice or region examined.

### **Sedation**

The use of medicines to make a person relaxed, drowsy, or less aware during a procedure while maintaining appropriate monitoring and breathing.

### **Sedation Dentistry**

The use of medicine during dental care to reduce fear, discomfort, or awareness. It may involve inhaled gas, tablets, or medicines through a vein; deeper sedation can affect breathing and requires trained staff and continuous monitoring.

### **Sedation Levels**

The depth of medicine-induced drowsiness, ranging from minimal relaxation with normal responses to deep sedation with limited response and possible breathing or airway support. Sedation can deepen unexpectedly, so appropriate monitoring is required.

### **Sedative**

A sedative is a medicine or substance that reduces alertness and may cause relaxation or sleepiness. It can impair coordination and breathing, especially when combined with alcohol or opioids.

### **Sedative-Hypnotic**

A medicine or substance that slows brain activ-



---

ity to reduce anxiety or cause sleep. Examples include benzodiazepines, barbiturates, and some sleep medicines. Too much can cause confusion, coma, or dangerously slow breathing, especially with alcohol or opioids.

### **Sedative, Hypnotic, or Anxiolytic Use Disorder**

A pattern of problematic use of sedative, hypnotic, or anxiolytic medicines causing impairment, distress, tolerance, withdrawal, or inability to control use.

### **Sedative-Hypnotic Toxidrome**

A pattern of poisoning caused by substances that slow brain activity. It may cause extreme sleepiness, slurred speech, poor coordination, slow breathing, coma, or loss of airway protection, especially when combined with alcohol or opioids.

### **Sedatives**

Medicines or substances that slow brain activity and may produce calmness, sleepiness, or reduced anxiety. Some are also used to induce sleep, but not every sedative is a sleep medicine. They can impair coordination and dangerously slow breathing, especially with alcohol or opioids.

#### **Alternate Names:**

Sedatives and Hypnotics

### **Sedimentation Rate**

The erythrocyte sedimentation rate, a nonspecific blood test measuring how quickly red cells settle in a tube and sometimes supporting assessment of inflammation when interpreted with other findings.

### **Sed Rate**

A blood test measuring how quickly red blood cells settle, used as a nonspecific marker of inflammation.

#### **Alternate Names:**

Erythrocyte Sedimentation Rate; ESR

### **Segmental Innervation**

The organization of nerve supply according to spinal cord segments. Each segment contributes to characteristic dermatomes, myotomes, and reflexes, so

changes in sensation, strength, or reflexes can help localize a neurologic lesion.

### **Segmental Pressures**

Blood-pressure readings taken at several points on the arms and legs to compare circulation. A large difference between levels can help locate a narrowed or blocked artery in peripheral artery disease.

### **Segmentation**

Repeated squeezing of short sections of the intestine that mixes food with digestive fluids and brings it against the intestinal lining for absorption. Unlike peristalsis, it mainly mixes contents rather than pushing them forward.

### **Segmentectomy**

Surgery removing one or more defined sections of a lung rather than an entire lobe. It may treat a small, localized lung cancer or another focal problem while preserving more lung tissue; bleeding, air leakage, infection, and reduced breathing capacity are possible.

#### **Alternate Names:**

Sublobar Resection; Anatomical Segmentectomy; Pulmonary Segmentectomy

### **SEID**

Alternate name; see *Myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)*.

### **Seidel Test**

An eye examination using fluorescein dye to look for leaking fluid from a full-thickness injury to the cornea or eye wall. Streaming or dilution of the dye suggests a leak, but a negative test does not rule out a serious eye injury.

#### **Alternate Names:**

Seidel's Test; Aqueous Humor Leakage Test

### **Seizure**

A temporary episode caused by abnormal electrical activity in the brain. It may cause shaking, stiffening, unusual sensations, staring, confusion, or loss of awareness and can be triggered by fever, illness,

---

injury, medicines, or epilepsy.

### Seizure Disorder

Alternate name; see *Epilepsy*.

### Seizure Disorders

Conditions characterized by recurrent unprovoked seizures or an enduring predisposition to them, classified by seizure type and cause and treated according to epilepsy syndrome and risk.

### Seizure Disorders in Pregnancy

Epilepsy and new-onset seizures during pregnancy require assessment of seizure type, medicine exposure, adherence, sleep, metabolic causes, and possible eclampsia. Management balances seizure control with fetal safety and includes coordinated neurology and obstetric care.

### Seizures

Seizures are transient episodes of abnormal excessive brain electrical activity causing changes in awareness, movement, sensation, behavior, or autonomic function.

### Selective Antibody Deficiency With Normal Immunoglobulins

An immune problem in which the total amount of antibodies is normal but the body does not make enough protective antibodies against some germs after vaccination. Repeated ear, sinus, or chest infections may occur.

### Selective Deficiency Of IgA

A usually inherited immune problem in which the antibody IgA is very low or absent while other antibodies are near normal. Some people have repeated sinus, chest, or bowel infections, allergies, or autoimmune disease; others have no symptoms.

### Selective Estrogen Receptor Modulator

A medicine that acts like estrogen in some tissues and blocks estrogen effects in others. Different medicines in this group are used for conditions such as osteoporosis, breast-cancer risk reduction,

or breast cancer.

### Selective Estrogen Receptor Modulators

Medicines that act like estrogen in some tissues and block estrogen effects in others, depending on the organ.

#### Alternate Names:

SERMs

### Selective IgA Deficiency

An immune condition with very low or absent immunoglobulin A, an antibody protecting mucous-lined surfaces. Some people have repeated respiratory or intestinal infections, allergies, autoimmune disease, or no symptoms.

### Selective Mutism

A childhood anxiety disorder in which a person speaks comfortably in some settings but repeatedly cannot speak in others, such as school or unfamiliar social situations. It lasts at least one month and can interfere with learning, relationships, or daily life.

### Selective Serotonin Reuptake Inhibitors

Antidepressants that increase serotonin signaling by slowing its reabsorption by nerve cells. They treat depression and several anxiety disorders, but effects take time and may include nausea, sexual problems, sleep changes, or withdrawal symptoms.

#### Alternate Names:

SSRIs

### Selective Serotonin Reuptake Inhibitor (SSRI)

Alternate name; see *Selective Serotonin Reuptake Inhibitors*.

### Selective Serotonin-Reuptake Inhibitors

Antidepressants that increase the availability of serotonin, a chemical used for communication between nerve cells. They treat depression and anxiety disorders; nausea, sleep changes, sexual problems, and withdrawal symptoms can occur.

---

### Selective Serotonin Reuptake Inhibitor Toxicity

SSRI overdose is often mild, causing nausea, vomiting, tremor, drowsiness, or agitation, but interactions can produce serotonin syndrome with clonus, hyperreflexia, hyperthermia, and autonomic instability. Citalopram and escitalopram can prolong the QT interval; stop serotonergic drugs, obtain ECG monitoring, and provide supportive care and benzodiazepines when indicated.

### Selective Tubal Occlusion Procedure

Selective tubal occlusion blocks a fallopian tube to prevent sperm and egg from meeting and is a permanent contraceptive procedure.

### Selegiline

A medicine that blocks monoamine oxidase B, used for Parkinson disease and sometimes depression; interactions require caution.

### Selenium

A mineral needed for antioxidant enzymes, thyroid function, reproduction, and immune activity. Deficiency is uncommon in many regions, while excessive supplements can cause hair or nail changes, stomach upset, nerve problems, or toxicity.

### Selenium-75

A radioactive form of selenium used mainly as a calibration or testing source in imaging and measurement equipment. It releases radiation as it decays and has a half-life of about 120 days.

### Selenium Deficiency

Low levels of selenium, a trace element needed for antioxidant enzymes and thyroid hormone activity. It is uncommon in most regions and usually subclinical, but severe deficiency contributes to Keshan disease, affecting heart muscle, and Kashin-Beck disease in selenium-poor areas. Assessment and replacement are guided by clinical and laboratory findings.

### Selenium In Diet

Selenium in diet provides a trace mineral needed for

antioxidant enzymes, thyroid hormone metabolism, and reproduction. Excess supplements can cause toxicity, so intake should remain within recommended limits.

### Selenium Sulphide

A skin-applied medicine that reduces certain fungi and excess scaling. It is used for dandruff and seborrheic dermatitis and may cause irritation, dryness, discoloration, or an unpleasant smell if misused.

### Self-Diagnosing Body Dysmorphic Disorder

Alternate name; see *BDD*.

### Self-Diagnosis

Self-diagnosis is identifying a health problem without a professional assessment. It can be inaccurate, so concerning or persistent symptoms should be evaluated by a clinician.

### Self-Efficacy

A person's confidence in their ability to manage a health condition, follow a treatment plan, and handle symptoms. It can influence participation in care and health outcomes.

### Self-Harm

Deliberate injury or poisoning of oneself. It may be related to overwhelming emotions, mental illness, trauma, or suicidal thoughts, and can include cutting, burning, hitting, or taking too much medicine. Every episode deserves compassionate safety assessment.

### Self Harming

Alternate name; see *Self-injury/cutting*.

### Self-Help Group

A peer group in which members support one another while changing a harmful behavior or coping with a problem.

#### Alternate Names:

Support Group; Peer Support Group

---

## Self-Injury/Cutting

Deliberate injury to one's own body, often used to cope with overwhelming emotions and requiring compassionate assessment of safety and mental health.

### Alternate Names:

Nonsuicidal Self-Injury; NSSI; Self-Harm

## Self-Management

Actions a person takes to monitor and manage a health condition, including treatment use, symptom monitoring, and lifestyle measures.

## Self-Poisoning

Intentional ingestion or exposure to a potentially harmful substance, often requiring urgent medical and mental-health assessment.

## Self-Retaining Retraction

Holding tissue away from an operative field with a retractor that remains in position without continuous manual support.

### Alternate Names:

Self-Retaining Retraction; Mechanical Retraction

## Self-Testing For COVID-19

Testing a sample collected at home, usually from the nose, to check for COVID-19 infection. A negative result does not always rule out infection, especially early in illness.

## Sella Turcica

A saddle-shaped hollow in a bone at the base of the skull that holds the pituitary gland. The pituitary controls many hormones, so changes in this area can affect growth, reproduction, metabolism, and the body's response to stress.

## Semaglutide

Semaglutide is a glucagon-like peptide-1 receptor agonist used for selected patients with type 2 diabetes or chronic weight management. It slows gastric emptying and reduces appetite and glucose levels; nausea, gallbladder disease, pancreatitis risk, and contraindications require review.

## Semen

The fluid released during ejaculation, containing sperm from the testes mixed with fluids from the seminal vesicles, prostate, and other glands. It carries and supports sperm; changes in amount, color, blood, or pain may need assessment.

## Semen Analysis

A laboratory examination measuring semen volume and sperm number, movement, shape, and vitality. It is a common first test when assessing fertility; results vary, so an abnormal result is often repeated and interpreted with the couple's history.

## Semen DNA Analysis

A test that examines genetic material inside sperm, usually for selected infertility or repeated-pregnancy-loss evaluations. It may detect increased DNA damage, but its usefulness and treatment implications depend on the clinical situation.

## Semicircular Canals

Three curved, fluid-filled tubes in each inner ear that sense head rotation and help maintain balance. Problems affecting them can cause spinning dizziness, nausea, or difficulty keeping the eyes steady when the head moves.

## Semi-Fowler Position

A semiupright position in which the head and upper body are elevated about 30 to 45 degrees.

## Semilunar Cartilages

The two crescent-shaped pads of tough cartilage inside the knee, commonly called the menisci. They cushion the joint, spread weight, and improve stability; a tear can cause pain, swelling, catching, or locking.

## Semilunar Valves

The aortic and pulmonary valves, which open during ventricular contraction and close during ventricular relaxation to prevent backflow into the ventricles.

---

## Seminal Fluid Analysis

A laboratory test of semen that measures the amount of fluid and the number, movement, and shape of sperm. It helps assess fertility and may check whether sperm remain after a vasectomy.

### Alternate Names:

Semen Analysis; Spermiogram

## Seminal Vesicle

One of a paired set of male accessory reproductive glands behind the urinary bladder. Its fructose-rich secretion contributes substantially to semen and joins the vas deferens through the ejaculatory duct.

## Seminal Vesicles

Paired glands behind the bladder that produce much of the fluid in semen. Their secretions contain nutrients and other substances that support sperm movement and help form the ejaculate.

## Seminiferous Tubules

Tiny, coiled tubes inside the testes where sperm are made. Supporting cells nourish developing sperm and help create the environment needed for sperm production.

## Seminoma

A cancerous tumor arising from sperm-forming cells in the testis. It often grows slowly and is highly treatable with surgery, radiation, chemotherapy, or careful monitoring depending on its stage.

## Semont Maneuver

A rapid, side-to-side liberatory repositioning maneuver used in the clinical management of posterior semicircular canal benign paroxysmal positional vertigo (BPPV), designed to dislodge cupulolithiasis or canalithiasis debris into the utricle.

### Alternate Names:

Semont Liberatory Maneuver; Liberatory Maneuver for BPPV

## Senescence

Changes and gradual loss of function associated with aging. In cells, senescence is a lasting state in

which a cell stops dividing but remains active; an accumulation of such cells may contribute to aging and disease.

### Alternate Names:

Biological aging

## Senile Lentigines

Flat, sharply defined brown or tan spots caused by chronic ultraviolet exposure, commonly appearing on sun-exposed skin with age.

## Senile Purpura

Alternate name; see *Actinic Purpura*.

## Senna

A plant-derived stimulant laxative containing sennosides, used short term for constipation and sometimes to empty the bowel before a procedure. It increases intestinal activity and may cause abdominal cramps or diarrhea; prolonged or excessive use can cause dependence or fluid and electrolyte disturbances.

## Sensate Focus Techniques

Structured, gradual touch exercises used in sex therapy to reduce performance pressure, improve body awareness, and rebuild comfortable intimacy. Partners agree on boundaries and initially avoid goals such as intercourse or orgasm.

### Alternate Names:

Sensate Focus; Masters and Johnson Therapy

## Sensation

Awareness of changes such as touch, pain, temperature, sound, movement, or body position. Special nerve endings detect these changes and send messages to the brain, where they are experienced and interpreted.

## Sense

A sense is a capacity to detect and interpret stimuli such as touch, temperature, pain, sound, light, smell, taste, position, or internal body signals.

---

## Sensitisation

A process in which repeated exposure makes the body's immune, nerve, or behavioral response stronger or easier to trigger. Sensitization can contribute to allergy, chronic pain, and some learned responses.

## Sensitivity

In testing, the proportion of people who truly have a condition and receive a positive result. In other settings, it means how strongly a person, cell, or instrument responds to a stimulus.

## Sensitivity Analysis

Sensitivity analysis tests how much a study's conclusion changes when assumptions, measurements, or input values are varied.

## Sensitized

Having antibodies against human-cell markers called HLA. This can happen after pregnancy, transfusion, or transplantation and may make it harder to find a suitable organ donor or increase the risk of transplant rejection.

## Sensorimotor Polyneuropathy

Damage to many nerves causing both altered sensation and weakness, usually starting in the feet or hands. Numbness, burning pain, poor balance, reduced reflexes, and muscle wasting may occur; diabetes is one possible cause.

## Sensorineural Deafness

Sensorineural deafness is hearing loss from dysfunction of the inner ear, auditory nerve, or central pathways rather than blockage of sound conduction.

## Sensorineural Hearing Loss

Hearing loss caused by damage to the inner ear or the nerve carrying sound to the brain. It may cause difficulty understanding speech, especially in noise, and can result from aging, loud noise, infection, inherited conditions, or medicines.

## Sensorium

A person's level of consciousness, awareness, attention, and understanding of surroundings. Changes may include confusion, drowsiness, agitation, disorientation, or reduced responsiveness and can signal illness, medicines, intoxication, or brain injury.

## Sensory

Relating to sensation or to nerves and pathways that carry information such as touch, pain, temperature, position, hearing, or vision to the brain.

## Sensory Cortex

Areas of the brain that receive and interpret information about touch, pain, temperature, pressure, and body position. Damage can cause numbness, altered sensation, or difficulty recognizing where a body part is.

## Sensory Deprivation

A major reduction in normal input from sight, sound, touch, movement, or other senses. Prolonged deprivation can affect mood, thinking, sleep, orientation, and perception, especially in isolated or vulnerable people.

## Sensory Level

A clear line on the body below which feeling is reduced or changed. It can suggest a problem in the spinal cord, especially when accompanied by weakness, altered reflexes, or bladder and bowel problems, and needs prompt assessment.

## Sensory Nerves

Nerves that carry information such as touch, pain, temperature, and position from the body to the spinal cord and brain. Damage may cause numbness, tingling, burning pain, or reduced awareness of injury.

## Sensory Neuron

A nerve cell that carries information from receptors toward the brain and spinal cord. It helps detect touch, pain, temperature, pressure, body position, sound, light, smell, and taste.



---

## Sensory Processing Disorder

Difficulty noticing, interpreting, or responding to sound, touch, movement, light, or body position. It may affect behavior, learning, coordination, or daily activities and can occur with other developmental conditions; assessment and support are individualized.

## Sensory Receptors

Specialized cells or nerve endings that detect stimuli such as touch, temperature, pain, light, sound, or chemical changes.

## Sentinel Event

An unexpected healthcare event that causes death or serious physical or psychological harm, or creates a major risk of such harm. It requires prompt investigation and system changes to reduce the chance of recurrence.

## Sentinel Injury

An unexpected injury in an infant who cannot yet move independently, such as bruising, a mouth injury, or a fracture. It may be an early sign of serious harm and requires careful examination, explanation, safeguarding assessment, and documentation.

## Sentinel Loop

A localized, persistently gas-filled, dilated segment of small intestine visible on plain abdominal radiographs, reflecting focal paralytic ileus produced by adjacent acute intraperitoneal inflammation (most commonly acute pancreatitis or acute appendicitis).

### Alternate Names:

Sentinel Loop Sign; Localized Adynamic Ileus

## Sentinel Lymph Node

The first lymph node or group of nodes most likely to receive fluid draining from a tumor. Removing and examining them can show whether cancer has begun to spread and help determine its stage.

## Sentinel Lymph-Node Biopsy

Surgical removal and pathologic examination of the first draining lymph node or nodes from a primary

tumor. It is used for selected melanomas, breast cancers, and other tumors to assess regional metastatic risk and guide staging.

## Sentinel Lymph Node Biopsy For Melanoma

Sentinel lymph-node biopsy identifies and removes the first draining lymph node or nodes from a melanoma site to detect microscopic spread. The result helps stage disease and guide further treatment; it does not remove all regional nodes routinely.

## Sentinel Node

A sentinel node is the first lymph node or group of nodes to which cancer cells are most likely to spread from a primary tumor. A sentinel lymph-node biopsy helps assess spread.

## Sentinel Node Biopsy

Alternate name; see *sentinel lymph-node biopsy: removal and examination of the first draining node or nodes to which a tumor is likely to spread, used to assess regional metastasis*.

## Sentinel Surveillance

Ongoing collection of health information from selected clinics, laboratories, or communities to track disease trends. It can provide early warning of outbreaks but does not count every case in the population.

## Separated Cranial Sutures

Separated cranial sutures in an infant can reflect normal growth or increased intracranial pressure, hydrocephalus, prematurity, or other illness and require clinical assessment.

## Separation Anxiety

Separation anxiety is developmentally excessive fear or distress when apart from an attachment figure, causing avoidance, physical complaints, sleep problems, or impairment. Treatment may include gradual exposure, family-based support, cognitive behavioral therapy, and selected medication.

---

## Separation Anxiety Disorder

Excessive and persistent fear about being apart from a parent, partner, or other attachment figure. It may cause repeated reassurance-seeking, refusal to leave home, nightmares, physical symptoms, or difficulty attending school or work.

## Separation Anxiety In Children

Developmentally excessive distress when separated from an attachment figure, with worry, refusal, physical symptoms, nightmares, or school avoidance that impairs daily life.

## Separation Of The Iris

A tear that pulls the colored part of the eye away from its normal attachment, usually after a blunt injury. It may cause double vision, glare, an irregular pupil, bleeding, or increased eye pressure and needs eye-specialist assessment.

## Sepsis

Life-threatening organ dysfunction caused by a dysregulated host response to infection. Findings may include altered mental status, hypotension, fast breathing or heart rate, low urine output, elevated lactate, low oxygenation, or other organ injury; the infection source and organ dysfunction must be assessed urgently.

### Alternate Names:

Systemic Inflammatory Response Syndrome

## Sepsis Syndrome

A range of severe responses to infection, from infection-related organ dysfunction to septic shock and failure of several organs. The condition can worsen quickly and may damage several organs.

## Septal Defect

An abnormal opening in the septum between the heart's chambers that permits blood to pass between the right and left sides of the heart.

## Septal Hematoma

A collection of blood inside the wall dividing the two sides of the nose, usually after an injury. It

causes blockage and painful swelling and can destroy cartilage and deform the nose if untreated.

## Septate Uterus

A congenital difference in which a band of tissue partly or completely divides the inside of the uterus. Many people have no symptoms, but it can increase miscarriage or some pregnancy risks; imaging confirms it, and selected cases are treated by removing the dividing tissue.

### Alternate Names:

Uterine Septum

## Septic

Relating to a serious infection or to harmful germs in a normally sterile body site. For example, septic arthritis is an infected joint, while septic shock is life-threatening organ and circulation failure caused by an overwhelming infection.

## Septic Abortion

A miscarriage or induced pregnancy loss complicated by infection of the uterus or surrounding tissues. Fever, chills, worsening abdominal pain, foul discharge, fast heartbeat, or faintness may occur; it is an emergency requiring antibiotics, supportive care, and removal of infected retained tissue when necessary.

### Alternate Names:

Infected Abortion; Postabortal Sepsis

## Septicaemia

Older term; see *Bacteremia* or *Sepsis*, depending on the clinical findings.

## Septic Arthritis

Infection of a joint, usually bacterial, causing acute pain, swelling, restricted movement, and systemic illness.

### Alternate Names:

Infectious Arthritis; Pyogenic Arthritis; Bacterial Arthritis; Non-Gonococcal Bacterial Arthritis

## Septic Embolism

A piece of infected material travels through the

---

bloodstream and blocks a smaller blood vessel, spreading infection to another organ. It often starts with an infected heart valve and can cause abscesses or tissue damage.

### Septicemia

Older term; see *Bacteremia* or *Sepsis*, depending on the clinical findings.

### Septic Pulmonary Infarction

Death of part of the lung after infected clots or other infected material block small lung arteries. It may cause fever, chest pain worse with breathing, cough, coughing blood, and lung abscesses.

### Septic Shock

A life-threatening complication of sepsis in which infection causes dangerously low blood pressure and poor blood flow to organs, even after initial fluids. Treatment may include antibiotics, fluids, and medicines to support circulation.

### Septic Shock Criteria

Clinical diagnostic criteria defined by consensus Sepsis-3 guidelines: persistent sepsis-induced hypotension requiring vasopressor therapy to maintain a mean arterial pressure of at least 65 mmHg, accompanied by a serum lactate level exceeding 2 mmol/L despite adequate fluid resuscitation.

### Septic Thrombophlebitis

Infected thrombosis within a vein, causing local inflammation and possible bloodstream infection or septic emboli. It may complicate intravenous devices, pelvic infection, skin infection, or deep soft-tissue infection.

### Septo-Optic Dysplasia

A condition present from birth involving underdevelopment of one or both optic nerves, parts of the brain's midline, and sometimes the pituitary gland. It may cause poor vision, involuntary eye movements, seizures, developmental differences, or hormone deficiencies.

### Septoplasty

Septoplasty is surgery to straighten or reshape the nasal septum to improve airflow or address structural obstruction.

### Septum

A wall or partition separating two spaces or tissues, such as the nasal septum or the septum between heart chambers.

### Sequela

A lasting consequence or condition that remains after an earlier disease, injury, infection, or treatment has ended, such as paralysis after stroke or scarring after infection.

### Sequelae

Lasting effects or complications that remain after an illness or injury. Examples include weakness after a stroke, breathing problems after severe infection, or pain after an injury; sequelae may be physical, mental, emotional, or social.

### Sequential Therapy

Treatment in which different medicines or interventions are given one after another rather than at the same time. The sequence may improve safety, effectiveness, or tolerance.

### Sequester

To sequester means to isolate or set apart; in medicine, sequestered tissue or material is separated from its normal circulation or surroundings.

### Sequestrum

A piece of dead bone that has separated from healthy bone, usually because of a long-lasting bone infection. It can prevent healing, cause drainage or pain, and may need surgery after infection is treated.

### Ser

Ser is the standard abbreviation for serine, an amino acid incorporated into proteins and involved in metabolism and cell signaling.

---

## Serine

Serine is a nonessential amino acid used in protein synthesis, one-carbon metabolism, membrane formation, and signaling.

## Serious Adverse Event

An unwanted medical event that results in death, is life-threatening, requires or prolongs hospitalization, causes major disability, or causes a birth defect. It may occur during treatment without being caused by the treatment.

## Serious Health Problems Symptoms

Alternate name; see *Symptoms of serious disease*.

## SERMs

Selective estrogen receptor modulators that have estrogen-like effects in some tissues and blocking effects in others.

### Alternate Names:

Selective Estrogen Receptor Modulators

## Seroconversion

Development of detectable antibodies in blood after infection, vaccination, or another immune-triggering event, showing a changed immune status.

## Serodiagnostic

Relating to diagnosis using antibodies, antigens, or other markers measured in blood serum. These tests can show infection, immunity, inflammation, or other conditions but must be interpreted with timing and clinical findings.

## Serologic Test

A blood test that detects antibodies, antigens, or other immune markers to help diagnose infection, assess immunity, or monitor response to treatment.

## Serology

The laboratory evaluation of serum for antibodies, antigens, or other immune markers. Results may indicate current infection, previous exposure, vaccination, autoimmune reactivity, or passive antibody transfer and must be interpreted with timing and

clinical context.

## Serology For Brucellosis

Blood testing for antibodies to *Brucella* species, interpreted with exposure history and symptoms because cross-reactions and early false-negative results can occur.

## Seroma

A pocket of clear body fluid that forms near a surgical incision or injured tissue. It may cause swelling, pressure, or discomfort and often resolves, though large or painful collections may need drainage.

## Seronegative Arthritis

Inflammatory arthritis in which common blood tests for rheumatoid factor and anti-CCP antibodies are negative. It includes several diseases with different patterns, such as spine, skin, bowel, or eye involvement, so symptoms and examination guide diagnosis.

## Serosanguineous

Serosanguineous describes thin wound or body fluid containing both clear serum and a small amount of blood, often appearing pink or pale red.

## Serositis

Inflammation of the smooth lining around the lungs, heart, or abdominal organs. It can cause sharp pain that worsens with breathing or movement and may cause fluid to collect around the affected organ.

## Serotonin

A chemical messenger used by nerve cells and other tissues. It helps regulate mood, sleep, appetite, bowel function, and pain; too much activity from medicines can cause serotonin syndrome.

### Alternate Names:

5-Hydroxytryptamine; 5-HT

## Serotonin Blood Test

A blood test measuring serotonin, used selectively in evaluating suspected serotonin-producing neuroendocrine tumors; many medicines and specimen

---

factors affect results.

### **Serotonin-Norepinephrine Reuptake Inhibitors**

Antidepressants that increase the activity of serotonin and norepinephrine, chemical messengers between nerve cells. They treat depression, some anxiety disorders, and certain nerve or long-lasting pain conditions; nausea, sweating, sleep changes, and withdrawal symptoms can occur.

### **Serotonin Syndrome**

A potentially life-threatening toxic reaction caused by excessive serotonin activity, usually after combining, increasing, or overdosing on serotonergic medicines. It often begins within minutes to hours and may cause agitation, sweating, diarrhea, tremor, overactive reflexes or clonus, muscle stiffness, fever, fast heart rate, blood-pressure changes, or confusion; severe cases require urgent medical care.

### **Serotype**

A distinct version of a microorganism or antigen identified by how antibodies react to it.

### **Serous**

Thin, watery, and usually clear, resembling blood serum. Serous fluid can occur normally or collect in a wound, body cavity, or blister.

### **Serous Borderline Ovarian Tumor**

An ovarian tumor whose cells look abnormal but have not invaded nearby supporting tissue. It may stay in the ovary or form deposits on its surface; treatment and follow-up depend on its extent and microscopic features.

### **Serous Cystadenoma of Pancreas**

Alternate name; see *Pancreatic Cysts*.

### **Serous Fluid Cytology**

Examination of cells in fluid collected from around the lungs, heart, or abdominal organs. It can look for cancer cells, infection, or inflammation; additional tests may be needed when the result is un-

clear.

### **Serous Membranes**

Thin, smooth linings that cover organs and reduce friction inside closed spaces, including the linings around the lungs, heart, and abdominal organs.

### **Serous Tubal Intraepithelial Carcinoma**

An early, noninvasive cancerous change in cells at the end of a fallopian tube near the ovary. It is thought to be the starting point for many high-grade cancers of the ovary, tube, or abdominal lining.

### **Serpentine Supravenous Hyperpigmentation**

A pattern of dark, wavy skin discoloration that follows the paths of surface veins, usually on the upper chest or neck. It can occur with vein inflammation, poor blood flow, some medicines, or cancer and requires assessment of the underlying cause.

### **Serrated Pathway**

One route by which some colon cancers develop from flat or saw-toothed polyps. These polyps may be easy to miss during colonoscopy and can acquire cell changes that increase cancer risk, so removal and follow-up are important.

### **Serrated Polyposis Syndrome**

A disorder characterized by multiple serrated lesions or polyps in the colon and an increased risk of colorectal cancer, requiring colonoscopic surveillance and removal of significant lesions.

### **Serration**

A row of small tooth-like projections or notches along an edge, used to describe anatomy, cells, instruments, or imaging appearances.

### **Serratus Anterior Muscle**

A muscle along the side of the chest that holds the shoulder blade against the ribs and helps raise the arm. Injury to its nerve can make the shoulder blade stick out and weaken pushing or overhead movements.

---

## Sertoli Cell

A support cell inside the seminiferous tubules that helps sperm develop and forms part of the blood-testis barrier.

## Sertoli Cells

Somatic "nurse" cells spanning the full thickness of the seminiferous tubule epithelium that physically support and nourish developing germ cells, form the blood-testis barrier via tight junctions, and secrete androgen-binding protein, inhibin B, and anti-Müllerian hormone.

### Alternate Names:

Sustentacular Cells of the Testis

## Sertoli-Cell Tumor

A tumor arising from Sertoli cells, which support sperm development in the testicle. It is often benign but may produce estrogen or other hormones and cause breast enlargement, infertility, or a testicular mass.

## Sertoli-Leydig Cell Tumor

A rare ovarian tumor that may produce male hormones, causing increased facial hair, acne, a deeper voice, or irregular periods. Treatment depends on the tumor's stage and features.

## Sertoli-Leydig Tumor

Alternate form; see *Sertoli-Leydig Cell Tumor*.

## Serum

The clear liquid left after blood has clotted. It contains water, salts, hormones, antibodies, nutrients, and waste products but lacks fibrinogen and most clotting proteins, making it useful for laboratory tests.

## Serum Albumin Test

A quantitative clinical blood assay measuring serum albumin levels to assess hepatic protein synthetic capacity, nutritional status, and oncotic pressure, as well as renal protein loss in nephrotic syndrome.

### Alternate Names:

Serum Albumin Test; Blood Albumin Test; Albumin

Level; Serum Albumin Assay

## Serum Chloride Test

A clinical laboratory measurement of chloride ( $\text{Cl}^-$ ) concentration in serum or plasma, essential for calculating the serum anion gap and evaluating acid-base balance, hydration status, and renal tubular function.

### Alternate Names:

Serum Chloride; Blood Chloride Test; Serum Electrolytes Chloride

## Serum Concentration

The amount of a substance measured in the liquid part of clotted blood. Results may reflect when the sample was taken, how the substance is absorbed and removed, and whether the level is expected to help or harm.

## Serum Free Hemoglobin Test

A blood test measuring hemoglobin not contained within red cells, used to support evaluation of intravascular hemolysis, transfusion reactions, or mechanical red-cell destruction.

## Serum Glutamic Oxaloacetic Transaminase

SGOT, now called AST, is an enzyme released by injured liver, heart, skeletal muscle, or other cells; its level must be interpreted with other findings.

## Serum Glutamic Pyruvic Transaminase

SGPT, now called ALT, is an enzyme concentrated in liver cells and commonly measured as a marker of hepatocellular injury.

## Serum Herpes Simplex Antibodies

A blood test for antibodies against herpes simplex virus types 1 or 2. IgG usually indicates previous infection, while IgM is less reliable; results do not show the infection site or prove that a current lesion is caused by herpes.

## Serum Iron Test

A blood test measuring circulating iron bound



---

mainly to transferrin; it is interpreted with ferritin, transferrin saturation, and blood counts rather than used alone to diagnose iron deficiency.

### **Serum Osmolal Gap**

The difference between the measured concentration of dissolved particles in blood and the calculated value. A high gap may suggest an unmeasured substance, including a toxic alcohol, but other illnesses and laboratory factors can also affect it.

### **Serum Phenylalanine Screening**

A blood test measuring phenylalanine, used in newborn screening and follow-up of phenylketonuria or other disorders of phenylalanine metabolism.

### **Serum Progesterone**

A blood test measuring progesterone, a hormone made mainly after ovulation and during pregnancy. The result can help assess ovulation, ovarian function, fertility treatment, or early pregnancy, but interpretation depends strongly on timing and pregnancy status.

### **Serum Protein Electrophoresis**

A blood test that separates proteins in serum into groups. It can show abnormal antibody production or low protein levels and is often followed by immunofixation when a plasma-cell disorder is suspected.

### **Serum Sickness**

A delayed immune reaction, usually one to three weeks after exposure to certain medicines, injected proteins, or antiserum. It can cause fever, rash or hives, painful swollen joints, enlarged lymph nodes, and sometimes organ problems.

### **Serum Therapy**

Prevention or treatment using antibody-containing plasma, immunoglobulin, antitoxin, or antivenom. It provides immediate but usually temporary protection against a specific infection or toxin; allergic reactions and other infusion-related problems are possible.

### **Serum Uric Acid**

A blood test measuring uric acid, a purine-breakdown product. The result helps assess gout, kidney stones, kidney function, or effects of some cancer treatments.

### **Serving Size**

Serving size is the standardized amount used to describe the nutrients or calories in food and may differ from the amount a person actually eats.

### **Sesamoid Bones**

Small bones embedded within or beside tendons, where they protect the tendon and improve leverage. Examples include the kneecap and the small bones beneath the big toe.

### **Sesamoiditis**

Painful inflammation of tissues around a sesamoid bone, most often beneath the big toe. Pain usually worsens with walking, running, dancing, or pressure on the front of the foot.

### **Sesquipedalian**

Sesquipedalian means characterized by very long words; clear healthcare communication favors language that patients can understand.

### **Sessile**

Attached directly by a broad base rather than a narrow stalk. A sessile polyp can be harder to remove completely and may require careful examination because its shape can hide abnormal tissue.

### **Set**

In exercise, a group of consecutive repetitions performed before a rest period. In medicine, “set” can also mean a prepared group of instruments, tests, or equipment.

### **Sever Disease**

A painful irritation of the growth area at the back of the heel in growing children, often during sports. Heel pain worsens with running or jumping and usually improves with activity adjustment, heel support,

---

calf stretching, and time.

### Severe Acute Respiratory Syndrome

A serious contagious respiratory illness caused by the SARS-associated coronavirus, SARS-CoV. It usually begins with fever and may cause chills, muscle aches, headache, diarrhea, dry cough, and breathing difficulty. Most affected people develop pneumonia, which can progress to low blood oxygen, respiratory failure, and death. The name usually refers to the disease identified in the 2002–2004 outbreak; SARS-CoV-2 causes COVID-19, a different disease.

### Severe Brain Injury Coma

Alternate name; see *Coma*.

### Severe Combined Immunodeficiency

A group of inherited conditions in which important immune cells do not develop or work properly. Affected babies can develop severe, repeated infections early in life and need urgent specialist care, protective measures, and immune-restoring treatment.

### Severe Combined Immunodeficiency (SCID)

Alternate name; see *Severe Combined Immunodeficiency*.

### Severe Dengue

The modern World Health Organization (WHO) classification term for the critical, life-threatening form of dengue virus infection, typically developing around the time of fever defervescence (days 3 to 7). It is characterized by one or more of three major clinical manifestations: severe plasma leakage resulting in hypovolemic shock (dengue shock syndrome) or respiratory distress from fluid accumulation, severe bleeding (such as massive gastrointestinal hemorrhage), or severe organ impairment including acute liver failure, encephalitis, or myocarditis.

#### Alternate Names:

Dengue Shock Syndrome; Severe Dengue Virus Infection; Critical Dengue

### Severe Obesity

Obesity with a Body Mass Index of 40 kg/m<sup>2</sup> or higher, or 35 kg/m<sup>2</sup> or higher together with a serious obesity-related health problem. Body Mass Index does not fully measure health.

#### Alternate Names:

Class III Obesity; Morbid Obesity; Extreme Obesity

### Sevoflurane

An inhaled anesthetic gas used to start and maintain general anesthesia. It acts in the brain to produce unconsciousness and reduces awareness of pain; it can lower breathing and blood pressure and may trigger malignant hyperthermia in susceptible people.

### Sex Change

A nontechnical term for gender transition or medical gender-affirming care; respectful, current terminology should be used.

### Sex Chromosomes

The X and Y chromosomes, which contribute to sex development and carry many other genes.

### Sex Cord-Stromal Ovarian Tumor

Pathologic evaluation of ovarian tumors derived from granulosa, theca, Sertoli, Leydig, or related stromal cells. Patterns, hormone production, nuclear features, and immunohistochemistry help distinguish the tumor type and malignant potential.

### Sex Headaches

Headaches brought on by sexual activity, including a gradually increasing ache or a sudden severe headache at orgasm. A first or thunderclap headache requires urgent assessment to exclude bleeding or other dangerous causes.

### Sex Hormones

Hormones involved in sexual development, reproduction, menstrual cycles, pregnancy, and related body functions. Major groups include estrogens, progesterone, and androgens, which also affect bones, muscles, mood, and tissues.

---

### Sex-Linked Dominant

A pattern of inheritance in which a disease-causing gene variant lies on a sex chromosome and one altered copy can cause the condition. The chance of passing it on differs for mothers and fathers and for children of different sexes.

### Sex-Linked Inheritance

Inheritance of a trait caused by a gene on a sex chromosome, commonly the X chromosome.

### Sex-Linked Recessive

A pattern in which a disease-causing gene variant, usually on the X chromosome, commonly causes disease in males with one altered copy. Females usually need two altered copies, while those with one may be carriers or have milder symptoms.

### Sex Therapy

Structured psychological or behavioral therapy for sexual concerns or dysfunction.

### Sexual Abuse

Nonconsensual sexual contact, exploitation, or behavior; it is a safeguarding and health emergency when immediate danger or injury exists.

### Sexual Addiction

Legacy label; see *Compulsive Sexual Behavior Disorder*.

### Sexual Assault

Any sexual contact or sexual act without freely given consent, including unwanted touching, penetration, coercion, or sexual activity when a person cannot consent. It can cause injury, infection, pregnancy, and psychological trauma and may require medical and safeguarding support.

#### Alternate Names:

Nonconsensual Sexual Contact; Sexual Violence

### Sexual Assault Prevention

Sexual-assault prevention includes consent education, recognizing coercion, safer environments, bystander intervention, and access to confidential sup-

port; responsibility always lies with the person who commits the assault.

### Sexual Dysfunction

A group of conditions involving persistent or recurrent difficulty with sexual desire, arousal, erection, ejaculation, orgasm, or pain that causes clinically significant distress. Causes may be physical, psychological, medication-related, hormonal, or related to life circumstances.

### Sexual Dysfunction Disorders

Alternate name; see *Sexual Dysfunction*.

### Sexual Health

Sexual health includes physical, emotional, mental, and social well-being related to sexuality. It includes contraception, fertility, consent, relationships, and prevention of sexually transmitted infections.

### Sexual Intercourse

Sexual activity involving penetration, most commonly penile-vaginal, but sometimes penile-anal or other forms depending on the people involved. It may be for pleasure, intimacy, or reproduction and requires freely given consent.

#### Alternate Names:

Copulation; Coition; Sexual Union

### Sexually Transmitted Disease

A sexually transmitted disease is an infection spread mainly through sexual contact. The term sexually transmitted infection is also used because some infections cause no symptoms.

### Sexually Transmitted Disease (STD)

Alternate name; see *Sexually Transmitted Disease*.

### Sexually Transmitted Diseases

Alternate name; see *Sexually Transmitted Infections*.

### Sexually Transmitted Diseases in Pregnancy

Alternate name; see *Sexually transmitted infections*.

---

## Sexually Transmitted Diseases in Women

Alternate name; see *Sexually transmitted infections*.

## Sexually Transmitted Disease Treatments

Alternate name; see *Sexually Transmitted Infections*.

## Sexually Transmitted Infection

Alternate name; see *Sexually Transmitted Infections*.

## Sexually Transmitted Infections

Infections caused by bacteria, viruses, fungi, or parasites that spread mainly through sexual contact, including vaginal, anal, or oral sex and genital skin-to-skin contact. Some can also spread through blood exposure or from a pregnant person to a baby. Many cause no symptoms, especially early in the infection.

### Alternate Names:

STIs; Venereal Diseases; STDs

## Sexual Masochism Disorder

A mental-health diagnosis involving repeated sexual arousal from being humiliated, beaten, bound, or otherwise made to suffer, when the pattern causes significant distress or disrupts daily life, or involves a person who has not freely consented.

## Sexual Obsession

Alternate name; see *Compulsive sexual behavior*.

## Sexual Rehabilitation

Assessment and treatment to help restore or adapt sexual function after illness, injury, surgery, disability, or medicine effects. It may include education, counseling, pelvic-floor therapy, devices, or medicines.

## Sexual Sadism

A recurrent sexual interest in another person's physical or psychological suffering or humiliation. The interest alone is not a mental disorder; acts without consent are abuse.

## Sexual Sadism Disorder

A mental-health diagnosis involving repeated sexual arousal from another person's physical or emotional suffering, when it causes clinically significant distress or impairment or involves acting with someone who has not freely consented.

## Sexual Violence

Sexual violence is any sexual act, attempt, contact, or coercion without freely given consent and can cause physical injury, infection, pregnancy, trauma, and long-term health effects.

## Sezary Disease

Alternate name; see *Sezary syndrome*.

## Sezary Syndrome

An uncommon form of cutaneous T-cell lymphoma involving widespread red skin and abnormal lymphocytes in the blood.

### Alternate Names:

Sezary Disease; Mycosis Fungoides Variant

## Sgarbossa Criteria

A set of electrocardiogram findings used to look for a heart attack when an existing conduction problem or pacemaker makes the tracing difficult to read. The findings can support, but cannot prove, an acute blocked heart artery.

### Alternate Names:

Modified Sgarbossa Criteria; Smith-Modified Sgarbossa Rule

## Shadowing Artifact

An ultrasound artifact in which a strongly attenuating or reflecting structure produces a dark area behind it because little sound reaches the tissues beyond. Stones, bone, and gas commonly cause shadowing.

## Shaken Baby Syndrome

Older term; see *Abusive Head Trauma*.

## Shaken Impact Syndrome

A term for abusive head trauma involving shaking

---

with or without impact, especially in infants.

### Shampoo Ingestion

Accidental swallowing of a small amount of shampoo usually causes mouth or stomach irritation; larger or concentrated exposures may cause vomiting or breathing problems and warrant poison-control advice.

### Shared Decision-Making

A collaborative process in which clinicians and patients combine best available evidence with patient values, preferences, goals, and circumstances to choose among reasonable options.

### Shark Attack

A shark attack can cause severe lacerations, bleeding, tissue loss, drowning, and shock; emergency rescue, hemorrhage control, and trauma care are critical.

### Sharp-Force Defense Wounds

Cuts or other injuries on the hands or forearms that may occur when a person attempts to protect against an attack. The distribution and morphology of defensive injuries may help reconstruct an assault but are not independently proof of a particular event.

#### Alternate Names:

Defensive Wounds; Defense Injuries

### Sharp Force Trauma

An injury caused by a sharp or pointed object, such as a knife, broken glass, or needle. It may produce a cut or a deeper puncture and can damage blood vessels, nerves, organs, or bones; deep or heavily bleeding wounds need emergency care.

### Sharps Container

A strong, leak-resistant container for used needles, lancets, blades, and other sharp medical items. It helps prevent injury and infection and should be clearly labeled, kept upright, and closed or replaced before it becomes too full.

### Sharps Injury

A percutaneous or mucocutaneous injury caused by a needle or other sharp instrument that may expose a person to blood or body fluids. Immediate wound care, incident reporting, risk assessment, and pathogen-specific post-exposure management are required.

### Shave Biopsy

Removal of a raised or superficial skin lesion by shaving off a thin layer with a sterile blade. The tissue is examined under a microscope; bleeding, infection, scarring, or an inadequate sample can occur.

### Sheehan's Syndrome

Loss of pituitary hormones caused by damage to the pituitary gland after severe bleeding or dangerously low blood pressure during or after childbirth. It may cause inability to breastfeed, extreme tiredness, low blood pressure, menstrual changes, or other hormone deficiencies.

### Sheehan Syndrome

Permanent injury to the pituitary gland caused by severe bleeding or very low blood pressure during or after childbirth. Because the gland makes several hormones, symptoms may include inability to breastfeed, low blood pressure, fatigue, and menstrual or fertility problems.

#### Alternate Names:

Simmonds Disease; Postpartum Hypopituitarism

### Shellac Poisoning

Poisoning from swallowing or inhaling shellac, a coating product that may contain alcohol or other solvents. It can cause irritation, vomiting, dizziness, drowsiness, or lung injury if it enters the lungs.

### Shellfish Allergy

An immune reaction to proteins in crustaceans or mollusks that can cause hives, digestive or respiratory symptoms, or anaphylaxis.

#### Alternate Names:

Crustacean Hypersensitivity; Shellfish Hypersensitivity



---

## Shellfish Poisoning

Illness caused by toxins accumulated in contaminated shellfish. Depending on the toxin, symptoms may include vomiting, diarrhea, tingling, weakness, confusion, breathing difficulty, or paralysis and require urgent medical advice.

## Shielding

Material placed between a person and a source of radiation to reduce exposure. The material depends on the radiation: lead or tungsten is used for many X-rays and gamma rays, while plastic or acrylic can be useful for beta radiation.

## Shifting Dullness

A change in the sound made when tapping the belly as a person turns onto their side, showing that extra fluid is shifting with gravity inside the stomach cavity.

## Shift Work Sleep Disorder

A circadian rhythm sleep-wake disorder caused by a recurring work schedule that overlaps the usual sleep period, producing insomnia and/or excessive sleepiness, reduced sleep, and impaired functioning. Symptoms persist in association with the schedule, typically for at least 3 months.

### Alternate Names:

SWSD; Shift-Work Sleep Disorder

## Shiga Toxin-Producing *Escherichia coli* Infection

Infection caused by Shiga toxin-producing strains of *Escherichia coli*. It commonly causes diarrhea and abdominal cramps and may cause bloody diarrhea. A small number of people, especially children, develop hemolytic uremic syndrome, which can damage the kidneys.

### Alternate Names:

STEC; Shiga Toxin-Producing *E. coli* Infection; Verocytotoxin-Producing *E. coli* Infection

## Shigella

A genus of bacteria causing shigellosis, an acute diarrheal illness that may include fever and bloody stools.

## Shigellosis

Shigellosis is intestinal infection caused by *Shigella* bacteria, producing diarrhea that may be bloody, fever, cramps, and dehydration.

## Shilajit

Shilajit is a mineral-rich, resin-like substance used in some traditional medicines. Product purity and safety vary, and contamination with heavy metals is possible.

## Shin Bone Fever

An older nonspecific term for painful inflammation or infection involving the shin or nearby tissues.

## Shingles

Herpes zoster, reactivation of varicella-zoster virus in sensory nerves causing a painful one-sided blistering rash. Pain, burning, or tingling may begin before the rash. Eye or ear involvement can threaten vision or hearing, and persistent nerve pain after the rash heals is called postherpetic neuralgia; prompt assessment is important for facial or eye-area rashes.

## Shingles And Pregnancy

Shingles is reactivation of varicella-zoster virus and usually causes a painful one-sided blistering rash. It is not the same as primary chickenpox exposure, but pregnancy requires prompt clinical advice about treatment, pain, and contact with people at risk.

## Shin Splints

A common term for medial tibial stress syndrome, an overuse injury causing pain and tenderness along the inner border of the shinbone. It is associated with repetitive impact or a sudden increase in activity and involves irritation of tissues around the bone. It is distinct from a stress fracture and acute compartment syndrome, which are separate causes of lower-leg pain.

### Alternate Names:

Medial Tibial Stress Syndrome

## Shin Spot

A shin spot is a nonspecific mark or lesion on the



---

lower leg; its significance depends on appearance, duration, symptoms, injury, and associated disease.

### **Shirodkar Cerclage**

An operation that places a strong stitch high around the cervix to help prevent it opening too early during pregnancy. It is used in selected people at risk of late miscarriage or premature birth and can cause bleeding, infection, or injury to nearby organs.

### **Shisha**

Shisha is tobacco or another smoking mixture used in a water pipe. The water does not remove the main toxic substances, and smoking shisha can expose users to nicotine, carbon monoxide, and cancer-causing chemicals.

### **Shivering**

Repeated, involuntary muscle contractions that generate heat, usually in response to cold or fever. Severe or unexplained shivering may occur with infection, medicine reactions, or other illness and should be assessed with the other symptoms.

### **Shock**

A life-threatening failure of circulation in which organs do not receive enough blood and oxygen. It can result from severe blood or fluid loss, heart failure, overwhelming infection, a severe allergic reaction, or an obstruction to blood flow. Signs may include confusion, faintness, cold or mottled skin, rapid breathing, little urine, or very low blood pressure and require emergency treatment.

### **Shock and Pregnancy**

Life-threatening inadequate tissue perfusion during pregnancy or after birth, caused by hemorrhage, sepsis, cardiogenic disease, embolism, anaphylaxis, or other processes. Rapid maternal resuscitation, hemorrhage or infection control, and fetal assessment are essential.

### **Shock Lung**

Older term; see *Acute Respiratory Distress Syndrome*.

### **Shock Wave Lithotripsy**

A procedure that focuses repeated sound-pressure waves from outside the body onto a kidney or ureter stone to break it into smaller pieces that can pass in urine. Blood in the urine, pain, blockage, infection, or incomplete stone removal can occur.

### **Shone Complex**

A group of congenital left-sided obstructive heart lesions, classically including supraventricular mitral ring, parachute mitral valve, subaortic stenosis, and coarctation of the aorta. The combination and severity vary and may lead to heart failure or systemic outflow obstruction.

### **Short Arm Of A Chromosome**

The shorter section of a chromosome, labeled p from the French word for “small.” Missing, extra, or rearranged DNA in this section can alter genes and cause disease.

### **Short Bones**

Small, roughly cube-shaped bones that provide stability and limited movement. The wrist and ankle contain most of the body’s short bones.

### **Short Bowel Syndrome**

A condition in which the small intestine is shortened or damaged and cannot absorb enough nutrients from food. It most often occurs after surgery to remove part of the small intestine and may cause diarrhea, weight loss, and nutritional deficiencies.

### **Short-Chain Fatty Acids**

Small fatty substances made when bacteria in the large intestine break down dietary fiber. They help nourish the bowel lining and may influence inflammation and metabolism; their effects vary with diet and the person’s gut bacteria.

### **Shortness Of Breath**

Alternate name; see *Dyspnea*.

### **Short Philtrum**

A short philtrum is reduced distance between the

---

nose and upper lip and may be an isolated feature or part of a genetic, developmental, or prenatal-exposure syndrome.

### **Short QT Syndrome**

A rare inherited heart-rhythm disorder in which the heart's electrical recovery time is unusually short. It can cause atrial or ventricular arrhythmias, fainting, or sudden cardiac death; diagnosis requires specialist ECG and family-history assessment, and treatment may include an implantable defibrillator.

**Alternate Names:**

SQTS

### **Short-Sight**

Myopia, a refractive error in which distant objects are focused in front of the retina.

### **Short Stature**

Height substantially below the expected range for age and sex, requiring assessment of growth pattern and cause.

### **Short-Term Memory**

The ability to hold and use a small amount of information for seconds or minutes. It helps with following instructions, doing mental tasks, and remembering a recent conversation before information is stored more permanently or forgotten.

**Alternate Names:**

Working Memory; Immediate Memory

### **Shotty**

Shotty describes small, firm, discrete lymph nodes, often reactive after infection; persistent, enlarging, hard, or generalized nodes require evaluation.

### **Shoulder**

The area where the upper limb joins the body, including the shoulder blade, collarbone, and upper-arm bone. Its main joint allows a wide range of arm movement but is less stable and can be injured or dislocated.

### **Shoulder Arthroscopy**

A minimally invasive procedure using a camera and instruments through small incisions to diagnose and treat selected disorders of the shoulder joint and surrounding tissues.

### **Shoulder Bursitis**

Shoulder bursitis is inflammation of a fluid-filled sac around the shoulder, often associated with rotator-cuff disease or repetitive overhead activity. Pain with elevation and lying on the affected side is common; activity modification, therapy, medicines, injection, or treatment of the underlying cause may help.

**Alternate Names:**

Subacromial Bursitis; Subdeltoid Bursitis

### **Shoulder CT Scan**

Computed tomography of the shoulder used to evaluate fractures, bone alignment, arthritis, tumors, infection, and complex bony anatomy.

### **Shoulder Dislocation**

Complete displacement of the upper arm bone from the shoulder socket, usually caused by a fall, collision, or forceful twisting. It causes severe pain, deformity, and inability to move the arm; urgent reduction and assessment for associated fractures or nerve and blood-vessel injury are required.

### **Shoulder Dystocia**

A childbirth emergency in which the baby's head is born but a shoulder becomes stuck behind the mother's pelvic bone. It requires immediate maneuvers because delay can cause lack of oxygen, fractures, or nerve injury in the baby and heavy bleeding in the mother.

### **Shoulder Impingement**

Pain associated with movement of the shoulder, often involving the rotator-cuff tendons or subacromial tissues. Symptoms include pain with overhead activity and sometimes weakness; assessment and treatment are individualized and may include exercise-based rehabilitation.

---

## Shoulder Impingement Syndrome

Mechanical compression and irritation of the rotator cuff tendons (principally the supraspinatus) and subacromial bursa beneath the coracoacromial arch during shoulder abduction and forward elevation, presenting with anterolateral shoulder pain and a painful arc between 60 and 120 degrees.

### Alternate Names:

Subacromial Impingement Syndrome; Neer Impingement

## Shoulder Instability

Excessive or recurrent movement of the shoulder joint caused by injury, laxity, or altered joint anatomy. Associated lesions may include a Bankart tear, Hill-Sachs lesion, or injury to the joint capsule or ligaments.

## Shoulder MRI Scan

Magnetic resonance imaging of the shoulder used to evaluate rotator-cuff tendons, labrum, cartilage, ligaments, muscles, bone marrow, and joint fluid.

## Shoulder Pain

Pain in or around the shoulder caused by injury, tendon or bursa disorders, arthritis, nerve disease, or referred pain. Sudden weakness, deformity, chest symptoms, fever, or pain after major trauma requires prompt evaluation.

## Shoulder Replacement

Surgery replacing damaged surfaces of the shoulder with artificial parts to relieve severe pain and improve movement. The implant may replace the joint in the usual or reverse configuration; infection, stiffness, dislocation, loosening, or nerve injury can occur.

## Shulman Syndrome

Alternate name; see *Eosinophilic fasciitis*.

## Shunt

A passage that diverts blood, cerebrospinal fluid, urine, or another fluid from one channel to another.

## Shwachman Syndrome

An inherited multisystem disorder usually involving exocrine pancreatic insufficiency, bone-marrow dysfunction, and skeletal abnormalities. Children may have steatorrhea, poor growth, recurrent infections, neutropenia, anemia, or increased risk of myeloid malignancy.

## Shy-Drager Syndrome

Alternate name; see *Multiple system atrophy*.

## SIADH

A condition in which the body releases too much antidiuretic hormone, causing it to retain water and dilute the blood sodium level. It may cause nausea, headache, confusion, seizures, or coma and requires treatment of the cause and sodium abnormality.

## Sialadenitis

Inflammation or infection of a salivary gland, causing painful swelling near the cheek or under the jaw, often worse during meals. Causes include bacteria, viruses, dehydration, or a blocked duct; persistent fever or pus needs medical care.

## Sialagogues

Substances or medicines that increase saliva production. They may help some people with dry mouth, but treatment also considers hydration, medicines, oral care, and the cause of reduced saliva.

## Sialogram

An imaging study of a salivary gland and its ducts after contrast is introduced through the duct opening, used to assess obstruction, stones, strictures, or chronic inflammation.

## Sialography

An imaging procedure that uses contrast material to show salivary glands and their ducts. It can help identify blockage, stones, narrowing, chronic inflammation, or tumors, although newer imaging is often preferred.

---

### Sialidosis (Mucopolidosis Type I)

A rare inherited lysosomal storage disorder caused by deficient alpha-neuraminidase, usually because of *NEU1* variants. It may cause coarse facial features, skeletal or eye abnormalities, myoclonus, ataxia, developmental impairment, or vision problems, with severity varying by type.

### Sialolithiasis

Formation of a stone in a salivary gland or its drainage tube, most often beneath the jaw. It can cause repeated swelling and pain during meals and may lead to blockage or infection.

### Sialorrhea

Excess saliva or difficulty controlling and swallowing it, causing drooling. It commonly occurs with neurologic disease, swallowing difficulty, dental problems, or some medicines; treatment addresses the cause and may include swallowing therapy or selected medicines.

### Sialorrhoea

Excess saliva that pools in the mouth or dribbles out as drooling. It is often caused by difficulty controlling or swallowing saliva rather than making too much. It may occur with cerebral palsy, Parkinson disease, motor-neuron disease, or other conditions and can cause skin irritation or choking. Swallowing therapy and other treatments may help.

### Sialuria

A rare inherited disorder in which cells make and release too much sialic acid, a sugar used on cell surfaces. It may cause developmental delay, enlarged liver and spleen, distinctive facial features, and variable learning difficulties.

### Sibling

A person's brother or sister; family history from siblings can inform genetic and disease risk assessment.

### SIBO

Alternate name; see *Small intestinal bacterial overgrowth*.

### Sicca Syndrome

Persistent dryness of the eyes and mouth caused by reduced tears and saliva. It may result from Sjögren syndrome, medicines, dehydration, radiation, or other illnesses and can lead to eye irritation, mouth sores, tooth decay, or difficulty swallowing.

### Sickle Cell

An inherited group of blood disorders caused by abnormal hemoglobin. Red blood cells may become rigid and curved, causing anemia, severe pain, blocked blood flow, and damage to organs.

### Sickle-Cell Anaemia

See *Sickle Cell Anemia*.

### Sickle Cell Anemia

An inherited blood disorder in which abnormal haemoglobin makes red blood cells stiff, sticky, and crescent-shaped. These cells break down easily and can block small blood vessels, causing anaemia, severe pain, stroke, lung problems, jaundice, and serious infections.

### Sickle Cell Crisis

A sudden complication of sickle-cell disease, usually caused by sickled red cells blocking blood flow and causing severe pain. Other crises can damage the lungs, reduce blood-cell production, or trap blood in the spleen; urgent care may be needed.

### Sickle Cell Dactylitis

Painful swelling of the hands or feet caused by blockage of small blood vessels during a sickle-cell episode, especially in infants and young children. It may be an early sign of sickle-cell disease and requires appropriate clinical evaluation.

### Sickle Cell Disease

A group of autosomal recessive inherited hemoglobin disorders involving hemoglobin S, which can make red blood cells rigid and prone to sickling. Sickling and red-cell breakdown can cause anemia, episodes of blocked blood flow, and damage to organs; forms include hemoglobin SS, hemoglobin SC, and hemoglobin Sbeta-thalassemia.

---

### Sickle Cell Test

A laboratory test detecting hemoglobin S or assessing sickling, used in screening and diagnosis of sickle-cell trait or sickle-cell disease; definitive testing identifies hemoglobin variants.

### Sickle Cell Trait

An inherited state in which a person has one gene for hemoglobin S and one usual hemoglobin gene. Most people have no symptoms and do not develop sickle cell disease, but rare problems can occur with severe dehydration, low oxygen, or extreme exertion.

### Sick Sinus Syndrome

A problem with the heart's natural pacemaker that causes the heartbeat to become too slow, pause, or alternate between fast and slow. It may cause dizziness, fainting, tiredness, palpitations, or reduced exercise ability.

#### Alternate Names:

Bradycardia-Tachycardia Syndrome

### Side Effect

An unintended effect that occurs while a medicine is used at a normal dose. It may be mild, such as nausea or drowsiness, or serious, such as bleeding, severe allergy, organ injury, or an abnormal heart rhythm.

### Side Effects

Unintended effects that occur while a medicine or treatment is used. They range from mild symptoms to serious harm; the risk depends on the treatment, dose, duration, other medicines, and the person's health.

### Side-Effects

Unintended effects that occur while a medicine or treatment is used. They range from mild symptoms to serious harm; the risk depends on the treatment, dose, duration, other medicines, and the person's health.

### Sideroblastic Anemia

Anemia caused by faulty hemoglobin production.

Iron builds up inside developing red blood cells instead of being used properly; causes include inherited disorders, alcohol, some medicines or toxins, vitamin B6 deficiency, and bone-marrow disease.

### Siderosis

Accumulation of iron in body tissues, or lung disease from breathing iron-containing dust. It may cause no symptoms, but long-term dust exposure can produce cough, breathing difficulty, and permanent lung damage.

### SIDS

Sudden infant death syndrome, the sudden unexplained death of an infant younger than one year after thorough investigation.

### Sievert

A unit used to estimate how much biological harm ionizing radiation could cause. It accounts for the type of radiation and the sensitivity of different tissues and is used when discussing radiation risk.

#### Alternate Names:

Equivalent Radiation Dose Unit; Sv Unit

### Sight

The ability to see, involving the eyes and visual nervous system.

### Sigmoid

Sigmoid means S-shaped; the sigmoid colon is the curved section of large intestine between the descending colon and rectum.

### Sigmoid Colon

The curved, S-shaped section of the large intestine between the descending colon and rectum. It stores stool before bowel movements and is a common site for diverticular disease, twisting, and colorectal cancer.

### Sigmoidoscopy

Endoscopic examination of the rectum and lower colon using a flexible or rigid sigmoidoscope. It can identify inflammation, bleeding, polyps, or tumors

---

and may allow biopsy or treatment.

### **Sigmoid Volvulus**

Twisting of the S-shaped lower colon around the tissue that supports it, blocking the bowel. It can cause severe abdominal swelling, pain, vomiting, and inability to pass stool or gas and needs urgent treatment because the blood supply may be cut off.

### **Sign**

An observable or measurable finding that may show disease or a change in health. Examples include a rash, fever, swelling, abnormal heartbeat, or high blood pressure; a sign differs from a symptom, which is felt and reported by the person.

### **Signal Peptide**

A short part of a newly made protein that acts like an address label, directing the protein to the cell membrane, outside the cell, or a particular internal compartment. It is often removed after delivery.

### **Signal-To-Noise Ratio**

A comparison between the useful information in a test or image and the unwanted background variation. A higher ratio usually produces a clearer image or more reliable measurement.

### **Signal Transduction**

The chain of events by which a hormone, nerve chemical, or medicine attaches to a cell and produces a response inside it. The response can change cell activity, movement, growth, survival, or gene use.

### **Signature**

A signature is a person's identifying mark or electronic confirmation of approval, consent, authorship, or verification.

### **Signature Strengths**

Personal character strengths that a person values and naturally uses.

### **Signet Ring Cell Carcinoma**

A cancer whose cells contain mucus that pushes

the cell nucleus to one side, giving a ring-like appearance under a microscope. It most often begins in the stomach or colon and may grow widely or aggressively.

### **Sign Language**

Sign language is a complete visual language using hand shape, movement, position, facial expression, and body movement rather than spoken sound.

### **Sign of Leser-Trelat**

The sudden appearance or rapid increase of numerous seborrheic keratoses, sometimes reported as a paraneoplastic sign. Most seborrheic keratoses are harmless and common, so the pattern is not diagnostic by itself and should prompt clinical context rather than automatic cancer conclusions.

### **Signs Of An Asthma Attack**

An asthma attack may cause worsening wheeze, cough, chest tightness, breathlessness, rapid breathing, difficulty speaking, or reduced peak flow; severe symptoms need emergency care.

### **Sildenafil Citrate**

The citrate salt of sildenafil, a phosphodiesterase-5 inhibitor used for erectile dysfunction and pulmonary arterial hypertension.

### **Silent Heart Attack**

A heart attack that causes no recognized pain or typical warning signs. It may be discovered later through an electrocardiogram or imaging and still damages heart muscle and increases future risks.

#### **Alternate Names:**

Unrecognized Myocardial Infarction; Silent MI

### **Silent Ischemia**

Reduced blood flow to heart muscle without typical chest pain or other recognized symptoms. It may still damage the heart and is detected through testing rather than symptoms alone.

#### **Alternate Names:**

Asymptomatic Myocardial Ischemia



---

### Silent Mutation

A DNA change that usually does not alter the amino acid in the protein made from the gene. It may still affect how the gene is copied, how long its RNA lasts, or how much protein is produced.

### Silent Myocardial Infarction

An acute ischemic myocardial infarction that occurs in the absence of classic chest pain or associated typical symptoms, recognized retrospectively on routine electrocardiogram (pathological Q waves) or echocardiography; prevalent in elderly individuals, diabetics (due to autonomic neuropathy), and women.

#### Alternate Names:

Painless MI; Silent Ischemia

### Silent Sinus Syndrome

Silent sinus syndrome is painless inward collapse of the maxillary sinus and floor of the orbit, often causing facial asymmetry, enophthalmos, or a sunken eye. Imaging confirms the diagnosis, and endoscopic treatment restores sinus ventilation when indicated.

### Silent Thyroiditis

Silent thyroiditis is painless autoimmune inflammation of the thyroid that may briefly release excess hormone before causing temporary hypothyroidism and eventual recovery.

### Silicate

A salt or ester containing silicon and oxygen, found in minerals and industrial materials; inhalation of some silicate dusts can irritate the lungs or contribute to pneumoconiosis.

### Silicon Dioxide

A mineral found in soil, rock, dust, and industrial materials. Breathing crystalline forms over time can scar the lungs, cause silicosis, and increase the risk of lung cancer.

### Silicone

A family of synthetic silicon-containing polymers used in medical devices, implants, sealants, and

consumer products; medical risks depend on the product and exposure.

### Silicone Elastomer

A flexible, rubber-like silicone material used in implants, tubing, dressings, and other devices; complications may include infection, leakage, allergy, or device failure.

### Silicone Implants

Medical implants containing silicone gel or elastomer used for breast reconstruction or augmentation; complications may include rupture, capsular contracture, infection, malposition, or need for revision.

### Silicone Mastitis

A granulomatous inflammatory reaction to silicone released from a breast implant or injected silicone. It may cause nodules, pain, lymphadenopathy, or imaging findings that mimic malignancy.

### Silicone Oil Hyperoleon

A complication after eye surgery in which silicone oil breaks into tiny droplets and moves into the front of the eye. It can raise eye pressure, irritate the eye, damage the cornea, and sometimes requires removal or washing of the oil.

### Silicones

Synthetic polymers used in medical devices, implants, dressings, and other products; adverse effects depend on formulation and site.

### Silicosis

A permanent lung-scarring disease caused by breathing crystalline silica dust. It can cause cough, breathlessness, reduced lung function, tuberculosis risk, and progressive respiratory disability after workplace or environmental exposure.

### Silk

Silk is a protein fiber used in textiles and some medical sutures or biomaterials; reactions and tissue response depend on the product and processing.

---

### Silk Road Disease

Alternate name; see *Behcet syndrome*.

### Siltuximab

A monoclonal antibody that binds interleukin-6 and is used to treat adults with multicentric Castleman disease who are negative for HIV and human herpesvirus 8.

### Silver

A metal used in some dressings, creams, and medical devices for antimicrobial activity; excessive exposure can cause permanent blue-gray skin discoloration called argyria.

### Silver Diamine Fluoride

A liquid applied to a tooth to slow or stop active tooth decay. It usually turns the treated decayed area black and may be useful when drilling or immediate restorative treatment is difficult.

### Silver Nitrate

A caustic antiseptic used topically to cauterize tissue, control minor bleeding, or treat some wounds; it can burn healthy skin and permanently stain tissue.

### Silver Poisoning

Silver poisoning, or argyria, follows excessive silver exposure and can cause permanent blue-gray skin discoloration and, rarely, organ or neurologic effects.

### Silver-Russell Syndrome

A growth disorder usually present from birth, marked by poor growth before and after birth, a small or asymmetrical body, and characteristic facial features. Most cases are linked to changes in gene activity rather than a change in the gene itself.

### Silver Sulfadiazine

A topical antimicrobial cream formerly used widely for burns; it may delay healing in some wounds and is not appropriate for everyone, including some newborns and people with sulfonamide allergy.

### Silymarin

A mixture of flavonolignans from milk thistle marketed as a dietary supplement; evidence for treating liver disease is limited and product quality varies.

### Simeprevir

An older hepatitis C virus NS3/4A protease inhibitor formerly used with other antivirals; it is no longer a usual first-line treatment because better direct-acting regimens are available.

### Simethicone

Simethicone is a non-absorbed antifoaming agent that coalesces gas bubbles in the gastrointestinal tract and is used for symptomatic relief of bloating; it does not treat the underlying cause of pain.

### Simian Crease

A simian crease is a single transverse crease across the palm, which may be a normal variant or occur with some chromosomal or developmental conditions.

### Simmonds Disease

An older term for severe pituitary-gland failure. It can cause extreme tiredness, weight loss, low blood pressure, loss of menstrual periods or sexual function, and deficiencies of several hormones.

### Simmonds Test

A calf-squeeze test for a possible torn Achilles tendon. If squeezing the calf does not make the foot point downward, a tear is suspected, but examination and sometimes imaging are needed.

#### Alternate Names:

Thompson Test; Thompson-Simmonds Calf Squeeze Test

### Simmonds-Thompson Test

Another name for the calf-squeeze test used when an Achilles tendon tear is suspected. A missing downward movement of the foot supports the diagnosis but does not replace a full assessment.

#### Alternate Names:

Thompson Test; Simmonds Test

---

### Simple Febrile Seizure

A brief, generalized seizure caused by fever in a child, usually between six months and five years old. It lasts less than 15 minutes and does not recur during the same illness. The child should be assessed to find the cause of fever, but most children recover fully and do not develop epilepsy.

#### Alternate Names:

Benign Febrile Seizure; Typical Febrile Seizure

### Simple Goiter

A simple goiter is noncancerous enlargement of the thyroid without obvious thyroid-hormone excess or deficiency, caused by iodine imbalance, stimulation, or other factors.

### Simple Mastectomy

Simple mastectomy removes the breast tissue and nipple while leaving most chest muscles and usually the underarm lymph nodes.

### Simple Prostatectomy

Surgical removal of the enlarged central portion of the prostate to relieve bladder-outlet obstruction from benign prostatic hyperplasia; it does not remove the entire prostate.

### Simple Pulmonary Eosinophilia

A usually mild lung reaction in which eosinophils, a type of white blood cell, collect temporarily in the lungs. It may cause cough, fever, or wheezing and can follow medicines, parasites, or other exposures. Symptoms and chest shadows often clear on their own, but the cause should be assessed.

### Simplexvirus

A genus of herpesviruses that includes herpes simplex virus types 1 and 2, which cause cold sores, genital herpes, and other infections.

### Simpson Test

A bedside fatigability test used to assess neuromuscular transmission in suspected ocular myasthenia gravis. The patient is asked to maintain upward gaze for one minute; progressive worsening of ptosis

or the development of diplopia constitutes a positive test result.

#### Alternate Names:

Sustained Upward Gaze Test

### Simuliidae

The family of black flies; bites can cause itchy skin reactions, and some species transmit *Onchocerca volvulus*, the parasite that causes river blindness.

### Simultanagnosia

An inability to see more than one object, or the whole visual scene, at the same time even though basic eyesight may be adequate. It usually results from damage to brain areas that combine visual information.

### Simvastatin

A statin medicine that lowers LDL, often called bad cholesterol, by reducing cholesterol production in the liver. It helps lower the risk of heart attack and stroke; muscle pain, liver problems, and medicine interactions are possible.

### Sincalide

A synthetic form of cholecystokinin used during gallbladder imaging to stimulate contraction and measure bile flow; it may cause abdominal pain, nausea, or flushing.

### Sindbis Virus

A mosquito-borne virus that can cause fever, rash, and joint pain. It occurs mainly in parts of Africa, Europe, Asia, and Australia.

### Sinding-Larsen-Johansson Syndrome

A painful overuse injury at the lower tip of the kneecap in growing children and adolescents. Pain usually worsens with running or jumping and improves as growth finishes and the irritated area heals.

### Sinecatechins

A topical green-tea catechin ointment used for external genital and perianal warts caused by human

---

papillomavirus; it can cause local irritation and should not be used internally.

### **Singer's Nodule**

A small, usually noncancerous thickening on a vocal fold caused by repeated voice strain. It can cause lasting hoarseness, a rough voice, or reduced vocal range; voice therapy and reducing strain are often helpful.

### **Singing**

Singing is producing musical sounds with the voice. It can support communication and social well-being, while excessive or improper voice use may cause vocal strain.

### **Single-Dose Toxicity**

Harm caused by one exposure to a substance or medicine, assessed according to dose, route, timing, and the exposed person's characteristics.

### **Single-Fiber Electromyography**

A specialized electrical test that records signals from individual muscle fibers. It measures small variations in the timing of nerve-to-muscle communication and can detect disorders such as myasthenia gravis, even when routine nerve tests are normal. A fine needle is used, and the test may be uncomfortable.

#### **Alternate Names:**

SFEMG; Single-Fiber EMG

### **Single Nucleotide Polymorphism**

A common inherited difference in one DNA building block. Most have little or no effect, but some are linked with disease risk, ancestry, or how a person responds to a medicine.

### **Single Palmar Crease**

A single transverse palmar crease is one crease crossing the palm instead of the usual two; it can be a normal variant or a feature of genetic syndromes.

### **Single-Photon Absorptiometry**

An older scan that used a small amount of radia-

tion to estimate bone density, usually in an arm or other peripheral bone. More accurate bone-density methods have largely replaced it.

#### **Alternate Names:**

SPA

### **Single Photon Emission Computed Tomography**

A nuclear medicine scan that uses a radioactive tracer and a rotating camera to create three-dimensional images of blood flow or organ function. It can assess the heart, bones, brain, kidneys, or other organs.

### **Single-Photon Emission Computed Tomography**

A nuclear-medicine scan that uses a radioactive tracer and a rotating camera to create three-dimensional images of blood flow or organ activity. It can assess the heart, brain, bones, kidneys, and other organs.

### **Single-Photon Emission Computed Tomography/Computed Tomography**

A combined scan that uses a radioactive tracer to show organ function and CT to show detailed body structure. The two sets of images are matched to help locate abnormal areas.

### **Singlet Oxygen**

A highly reactive form of oxygen involved in oxidative reactions, immune defenses, photodynamic therapy, and tissue injury from oxidative stress.

### **Single Ventricle**

A group of heart defects present at birth in which one lower chamber does most of the work of pumping blood to both the body and lungs. Babies usually need specialist care and staged operations.

### **Singultus**

Medical term for hiccups; see *Hiccups*.

### **Sin Nombre Virus Infection**

Alternate name; see *Hantavirus pulmonary syn-*

---

*drome.*

### Sinoatrial Block

A cardiac conduction disturbance in which electrical impulses generated by the sinoatrial node are delayed or blocked from exiting the node to depolarize the surrounding atrial myocardium, manifesting on electrocardiogram as intermittent dropped P waves and pauses in sinus rhythm.

#### Alternate Names:

SA Exit Block

### Sinoatrial Node

The heart's natural pacemaker in the right atrium that normally initiates each heartbeat.

#### Alternate Names:

Cardiac Pacemaker; SA Node

### Sinuatrial Node

The sinoatrial node, a group of specialized cells in the right atrium that normally initiates the heartbeat and sets the cardiac rhythm.

### Sinus

A sinus is a cavity, channel, or tract; in anatomy it may mean a paranasal air space, venous channel, or abnormal passage from a diseased area to the surface.

### Sinus Arrest

Transient failure of the sinoatrial node pacemaker cells to spontaneously generate an electrical impulse, producing a pause on electrocardiogram that is not an exact mathematical multiple of the baseline PP interval, lasting from seconds to several seconds before an escape pacemaker emerges.

#### Alternate Names:

Sinus Pause

### Sinus Arrhythmia

Sinus arrhythmia is normal variation in sinus-heart rate, commonly speeding during inhalation and slowing during exhalation, especially in young people.

### Sinus Bradycardia

Sinus bradycardia is a sinus rhythm slower than the expected rate; it can be normal during sleep or athletic conditioning but may also reflect drugs or disease.

### Sinus CT Scan

Computed tomography of the paranasal sinuses used to evaluate chronic or complicated sinusitis, anatomic obstruction, trauma, inflammatory disease, or tumors.

### Sinuses

Air-filled cavities within the skull bones lined by respiratory mucosa, connected to the nasal cavity via ostia.

### Sinus Headache

Pain attributed to a sinus headache is often migraine, but sinus inflammation can cause facial pressure, congestion, reduced smell, and pain that worsens with bending. Fever, purulent discharge, persistent symptoms, or eye or neurologic signs guide evaluation.

### Sinus Histiocytosis

An increase in cleanup immune cells within the open spaces of a lymph node. It is often a response to nearby inflammation, infection, a tumor, or tissue breakdown and is interpreted with the rest of the biopsy.

### Sinusitis

Inflammation of the lining of one or more paranasal sinuses. It may follow a viral respiratory infection or be associated with allergy, nasal inflammation, or other causes. Common features include nasal obstruction, discharge, facial pressure or pain, headache, and reduced smell. Acute and chronic forms differ in duration and pattern.

### Sinusitis And Sinus Problems

Alternate name; see *Sinusitis*.

---

## Sinus Lift

A dental procedure that adds bone beneath the lining of the maxillary sinus in the upper jaw. It may create enough support for a dental implant when the jawbone has become too thin.

## Sinus MRI Scan

Magnetic resonance imaging of the paranasal sinuses and surrounding soft tissues used to evaluate tumors, invasive infection, orbital or intracranial extension, and selected inflammatory disease.

## Sinus Node

The heart's natural pacemaker, located in the right atrium.

### Alternate Names:

Sinoatrial Node; SA Node

## Sinusoid

A small, wide blood vessel with unusually open walls, found especially in the liver, spleen, and bone marrow. Its structure allows blood to exchange cells, proteins, nutrients, and waste with nearby tissue.

## Sinusoidal Fetal Heart Rate Pattern

A rare, smooth, regular wave pattern on an unborn baby's heart-rate tracing with little normal variation. It can indicate severe anemia or low oxygen and requires immediate assessment and monitoring.

## Sinusoidal Obstruction Syndrome

Injury and blockage of the small blood vessels in the liver, most often after high-dose chemotherapy given before a stem-cell transplant. It can cause painful liver enlargement, jaundice, fluid buildup, swelling of the abdomen, and rapid weight gain.

### Alternate Names:

Hepatic Veno-Occlusive Disease; VOD/SOS

## Sinusoids

Specialized, low-pressure microvascular blood vessels featuring wide, irregular lumens, fenestrated or discontinuous endothelial cells, and an incomplete or absent basement membrane, found in the liver, spleen, and bone marrow to facilitate unrestricted

transcellular exchange of macromolecules and blood cells.

## Sinus Pain

Pressure or pain around the forehead, cheeks, nose, or eyes caused by sinus inflammation, infection, migraine, or other conditions. Severe headache, eye swelling, vision change, high fever, or neurologic symptoms requires urgent assessment.

## Sinus Rhomboidalis

A diamond-shaped depression forming the floor of the fourth brain ventricle, near structures controlling movement and vital functions.

## Sinus Rhythm

A heart rhythm started by electrical impulses from the sinoatrial node, the natural pacemaker. It is usually regular, although sinus rhythm can be too fast, too slow, or affected by illness.

## Sinus Tachycardia

A heart rhythm originating in the sinoatrial node with a rate faster than normal, commonly occurring as a physiologic response to exercise, fever, pain, dehydration, or stress.

## Sinus X-Ray

Plain radiographic imaging of the paranasal sinuses; it has limited sensitivity and is often replaced by CT when detailed evaluation is needed.

## Sipuleucel-T

A personalized cancer immunotherapy made from a patient's immune cells and used for selected advanced prostate cancers. The cells are trained to recognize a prostate-cancer target before being returned to the patient.

## Sirenomelia

A rare congenital malformation in which the lower limbs are partially or completely fused. It is usually associated with severe abnormalities of the genitourinary, gastrointestinal, vascular, and spinal systems and is often fatal in the neonatal period.



---

## **Sirolimus**

Sirolimus is an mTOR inhibitor used for immunosuppression after transplantation and selected vascular or lymphatic disorders; mouth ulcers, hyperlipidaemia, cytopenias, infection, impaired wound healing, and pneumonitis can occur.

## **Sirtuins**

A family of enzymes that change other proteins and help regulate how cells use energy, respond to stress, and switch genes on or off. Their possible roles in aging and disease are still being studied.

## **Sisomicin**

An aminoglycoside antibiotic related to gentamicin that has activity against some aerobic gram-negative bacteria; use is limited by kidney and hearing toxicity.

## **Sister Mary Joseph Nodule**

A metastatic tumor deposit in or around the umbilicus, usually from an intra-abdominal or pelvic malignancy.

## **Sister Mary Joseph's Nodule**

A firm, hard nodule in the umbilicus resulting from metastatic malignancy, most commonly from intra-abdominal or pelvic cancers including gastric, ovarian, and colorectal carcinoma. It indicates advanced, disseminated disease with a poor prognosis. Diagnosis is by fine needle aspiration biopsy.

## **Sitagliptin Phosphate**

Sitagliptin phosphate is a DPP-4 inhibitor that prolongs incretin activity and improves glucose-dependent insulin secretion in type 2 diabetes; dose adjustment in kidney disease and rare pancreatitis or severe joint pain are important considerations.

## **Sitosterolemia**

A rare inherited disorder in which the body absorbs and stores too much plant sterol, a substance found in plant foods. The buildup can cause early artery disease, skin lumps, anemia, easy bleeding, or low platelets.

## **Sitting Position**

An upright or semiupright position in which the patient's torso is elevated and the hips and knees are flexed; it is used for selected shoulder and neurosurgical procedures.

## **Sit-to-Stand Postural Drift**

A tendency to lose balance or move away from a stable position while standing up from a chair. It may reflect weakness, poor coordination, dizziness, or impaired balance and should be interpreted with the full examination.

### **Alternate Names:**

Postural Instability on Rising; Chair Rise Imbalance

## **Situational Syncope**

A reflex faint triggered by a specific situation such as coughing, urination, defecation, swallowing, laughter, or prolonged standing, caused by transient reduced cerebral perfusion.

## **Situs Ambiguus**

A birth difference in which internal organs are arranged irregularly rather than in the usual left-right pattern. It is often associated with congenital heart defects and an absent, small, or abnormally functioning spleen.

## **Situs Inversus**

A congenital condition in which the thoracic and abdominal organs are reversed or mirrored from their normal positions. Complete situs inversus (situs inversus totalis) involves all organs.

## **Situs Inversus Totalis**

A congenital condition in which the major visceral organs are mirrored from their normal positions, with the liver on the left, heart on the right, and other viscera similarly reversed.

## **Sitz Bath**

Immersion of the perineal and anal region in warm or prescribed water for symptom relief in conditions such as hemorrhoids, anal fissures, or postpartum discomfort.

---

## SI Units

The International System of Units used for standardized measurements, including meter, kilogram, second, mole, and pascal.

## Six Fingers Or Toes

Six fingers or toes is polydactyly, a congenital extra-digit difference that may occur alone or as part of a genetic syndrome.

## Six-Minute Walk Test

A simple test measuring how far a person can walk in six minutes on a flat path. Clinicians record distance, symptoms, heart rate, and oxygen level to assess exercise ability and monitor heart or lung disease. Rest breaks are allowed, and the result is interpreted with other findings.

### Alternate Names:

6MWT; Six-Minute Walk Distance

## Sixth Disease

Alternate name; see *Roseola*.

## Sixth Nerve Palsy

Dysfunction of the abducens nerve (cranial nerve VI) producing weakness or paralysis of the ipsilateral lateral rectus muscle, characterized by convergent strabismus (esotropia) and horizontal binocular diplopia worsening on gaze toward the paretic side; frequently caused by microvascular ischemia or elevated intracranial pressure.

### Alternate Names:

Abducens Nerve Palsy

## Size Perception

The brain's judgment of how large or small an object or body part is. Vision, touch, distance, and context all contribute; eye or brain disorders can make objects appear unusually large or small.

## Sjögren Disease

The current preferred name for Sjögren syndrome, an autoimmune disease that commonly causes dry eyes and dry mouth and may affect other organs.

## Sjögren Disease-Associated Arthritis

Joint pain or inflammation associated with Sjögren disease. Joint pain is more common than objective joint inflammation, so symptoms are interpreted alongside examination and other findings.

## SjöGren's Disease

Alternate name; see *Sjögren disease*.

## SjöGren's Syndrome

Older name for Sjögren disease.

### Alternate Names:

SjöGren's Disease

## SjöGren Syndrome

Older spelling and name for Sjögren disease.

## Sjogren-Larsson Syndrome

A rare autosomal recessive neurocutaneous disorder caused by mutations in the ALDH3A2 gene encoding fatty aldehyde dehydrogenase (FALDH), characterized by the clinical triad of congenital ichthyosis, spastic diplegia or tetraplegia, and severe intellectual disability with retinal glistening white dots.

### Alternate Names:

SLS; Fatty Aldehyde Dehydrogenase Deficiency

## Sjogren's Syndrome

Older name for Sjögren disease, an autoimmune disease that commonly causes dry eyes and dry mouth.

## Sjogren Syndrome

Older name for Sjögren disease, an autoimmune disease that commonly causes dry eyes and dry mouth.

## SJS

Alternate name; see *Stevens-Johnson syndrome*.

## Skeletal Development

Skeletal development is formation, growth, remodeling, and maturation of bones and cartilage from embryonic life through adulthood.

---

## **Skeletal Dysplasia**

An inherited disorder of bone or cartilage development that may cause short or uneven growth, unusually shaped limbs, spine problems, or other differences in the skeleton. Effects vary widely between conditions.

## **Skeletal Limb Abnormalities**

Skeletal limb abnormalities are differences in the formation, length, alignment, or structure of an arm or leg caused by genetic, developmental, vascular, injury, or other factors.

## **Skeletal Muscle**

Striated muscle attached to bones and generally under voluntary control.

## **Skeletal Muscles**

Muscles attached to bones by tendons that produce movement, maintain posture, and generate heat. They are usually under voluntary control but also support automatic actions such as breathing.

## **Skeletal Muscle Tissue**

Voluntary striated muscle attached to bones or other structures, composed of contractile fibers that generate movement, posture, heat, and joint stability.

## **Skeletal Myocytes**

Skeletal myocytes are striated muscle cells responsible for voluntary movement, posture, heat production, and stabilization of joints.

## **Skeletal Survey**

A set of X-rays of most of a child's bones, used especially when multiple injuries or a medical condition affecting bone development is suspected. It can show fractures, bone abnormalities, and signs of healing and must be interpreted with the full clinical assessment.

## **Skeletal System**

The skeletal system consists of bones, cartilage, joints, and supporting ligaments that provide structure, protect organs, enable movement, and store

minerals.

## **Skeletal X-Ray**

A skeletal X-ray series uses radiographs of selected bones and joints to evaluate fractures, alignment, bone density patterns, arthritis, infection, tumors, and developmental abnormalities.

## **Skeleton**

The body's framework of bones and cartilage that supports movement and protects organs. It also stores minerals, contains marrow that makes blood cells, and provides attachment points for muscles and ligaments.

## **Skeletonization**

Final stage of postmortem decomposition where soft tissues, organs, and skin have completely decayed or been scavenged, leaving only clean osseous skeletal remains.

## **Skene Gland Cyst**

A fluid-filled sac that forms when a Skene gland, also called a paraurethral gland, becomes blocked. These glands are located beside the opening of the urethra. Small cysts may cause no symptoms. Larger or infected cysts may cause pain, painful urination, difficulty passing urine, pain during sex, recurrent urinary infections, or an abscess. Diagnosis is usually made by pelvic examination. Symptomatic cysts may need surgical removal or marsupialization.

## **Skilled Nursing Facility**

A facility where people receive 24-hour nursing care and rehabilitation after illness, injury, surgery, or a hospital stay. Care may include wound care, medicines, injections, therapy, and help with daily activities until the person can return home or move to another setting.

## **Skilled Nursing Or Rehabilitation Facilities**

Facilities that provide 24-hour nursing care, rehabilitation, medication management, and assistance with daily activities after illness, injury, surgery, or functional decline.

---

## Skin

The body's outer covering, consisting mainly of the epidermis and dermis with appendages such as hair, nails, and glands. It protects against injury, infection, ultraviolet light, and water loss; detects touch, pressure, pain, and temperature; helps regulate body temperature; and contributes to immune defense and vitamin D production.

### Skin Abscess

A painful, swollen pocket of pus under the skin, usually caused by a bacterial infection. It may need drainage and antibiotics.

### Skin Absorption

Movement of a substance through the skin into underlying tissues or the bloodstream. Absorption depends on the substance, skin condition, contact time, and exposed area.

### Skin Aging

Natural aging and long-term sun exposure gradually thin and dry the skin, reduce elasticity, slow healing, and cause wrinkles or dark spots. Limiting ultraviolet exposure, using protective clothing, and applying broad-spectrum sunscreen can reduce further damage.

### Skin Aging And Appearance

Skin aging is a complex biological process influenced by both intrinsic factors, such as genetics and natural aging, and extrinsic factors, particularly ultraviolet radiation exposure.

### Skin Allergies And Contact Dermatitis

Inflammatory skin reactions triggered by exposure to allergens or irritants that come into direct contact with the skin.

### Skin Allergy Testing

Tests in which small amounts of suspected allergens are placed on or just under the skin and the reaction is observed. They can show sensitization to foods, medicines, venoms, or environmental allergens but must be interpreted with symptoms and exposure history.

## Skin And Aging

Skin aging involves intrinsic and extrinsic changes. Intrinsic aging causes thinning, wrinkling, and fragility. Extrinsic (photo)aging causes wrinkles, telangiectasias, lentigines, and actinic keratoses. Collagen loss, elastin degradation, and decreased cell turnover contribute to aging skin.

## Skin And Environment

Environmental factors significantly affect skin health. Pollution causes oxidative stress, inflammation, and accelerated aging. Particulate matter penetrates skin, generating free radicals. Blue light from screens causes hyperpigmentation, particularly in darker skin tones. Indoor heating and air conditioning cause dryness.

## Skin Appendages

Structures that grow from or lie within the skin, including hair follicles, oil glands, and sweat glands. They help lubricate and cool the body and can be affected by acne, infection, inflammation, or inherited disorders.

## Skin Barrier

Alternate term; see *Skin Barrier Function*.

## Skin Barrier Function

The outer skin barrier limits water loss and helps keep out irritants, allergens, and germs. Its tightly packed surface cells and natural oils maintain protection; injury or inflammation can make skin dry, cracked, and more vulnerable.

## Skin Biopsy

Removal of a small skin sample for examination under a microscope. It can help diagnose skin cancer, infection, inflammatory disease, blistering disease, or an unexplained rash; the method depends on the lesion.

## Skin Biopsy Methods

The sampling method is chosen according to the lesion's appearance and depth. A punch takes a round full-thickness sample, a shave removes a superficial layer, and an excision removes the whole

---

lesion when complete removal is appropriate.

### **Skin Biopsy Techniques**

Skin biopsy is selected according to the suspected diagnosis and lesion depth. Punch biopsy samples full-thickness skin; shave biopsy samples superficial lesions; excisional biopsy removes an entire small lesion; and incisional biopsy samples part of a large lesion.

### **Skin Blushing/Flushing**

Temporary reddening or warming of the skin caused by widening of small blood vessels. It may result from heat, emotion, exercise, fever, medicines, or hormone changes.

### **Skin Cancer**

Cancer that begins in cells of the skin. Common types include basal cell carcinoma, squamous cell carcinoma, and melanoma; less common types arise from other skin structures. Risk is influenced by ultraviolet exposure, immune suppression, genetic susceptibility, and certain preexisting lesions. A skin cancer may appear as a changing, bleeding, growing, scaly, or nonhealing spot.

#### **Alternate Names:**

Cutaneous Malignancy; Non-Melanoma Skin Cancer

### **Skin Cancer Epidemiology**

Skin cancers are common and include basal-cell carcinoma, squamous-cell carcinoma, and melanoma. Incidence and mortality vary by tumor type, geography, skin phenotype, ultraviolet exposure, immune status, and access to diagnosis and treatment.

### **Skin Cancer Screening**

Examination of the skin for new or changing spots, especially in people at increased risk. Clinicians may use a full-body examination, a magnifying instrument, or a biopsy; routine screening recommendations vary by individual risk.

### **Skin Cancer Staging**

Skin cancer staging guides treatment and prognosis. Melanoma: Breslow thickness (most important

prognostic factor), ulceration, mitotic rate, sentinel lymph node status (AJCC 8th edition).

### **Skin Cancer Surgery**

Surgery to remove skin cancer with a margin of apparently healthy tissue. The method depends on the cancer type, size, location, and spread and may include standard excision or Mohs surgery, which checks the edges during removal.

### **Skin Carcinoma**

Cancer arising from skin cells, most commonly basal-cell or squamous-cell carcinoma. It may appear as a changing sore, lump, scaly patch, or wound that does not heal and needs examination.

### **Skin Care**

Measures that protect and maintain the skin, including gentle cleansing, moisturization, sun protection, wound care, avoidance of irritants, and treatment of identified disease.

### **Skin Care And Incontinence**

Skin protection for people with urine or stool leakage, including gentle cleansing, thorough drying, absorbent products, barrier creams, regular changing, and treatment of the underlying cause.

### **Skin Cells**

Skin cells form the layers of the skin and include keratinocytes, melanocytes, immune cells, and sensory cells. They provide protection, pigmentation, sensation, and repair.

### **Skin Color**

The visible color of skin, determined mainly by melanin, blood flow, oxygenation, and other pigments.

### **Skin Culture**

Laboratory testing of a swab, scraping, fluid, or tissue sample from the skin to identify bacteria, fungi, or other organisms causing infection.

---

## Skin Cyst

A skin cyst is a closed sac under or within the skin, often filled with fluid, keratin, or other material. Many are harmless, but a painful, infected, or changing cyst should be assessed.

## Skin Diseases

Conditions affecting skin, hair, nails, glands, or subcutaneous tissue, including infection, inflammation, neoplasia, and genetic disorders.

## Skin Erosion

A shallow loss of the epidermis that leaves a moist or crusted surface but does not extend deeply into the dermis, often caused by friction, scratching, blister rupture, or inflammation.

## Skin Examination

Systematic inspection and palpation of the skin, hair, nails, and mucous membranes to assess color, lesions, texture, distribution, vascular signs, infection, and cancer warning features.

## Skin Findings In Newborns

Common skin changes in newborns, such as milia, newborn rash, peeling, birthmarks, or bluish hands and feet. Most are harmless, but fever, blisters, widespread redness, or a sick appearance needs prompt assessment.

## Skin Flaps

Pieces of skin and underlying tissue moved to cover a wound while keeping or reconnecting their blood supply. Flaps may come from nearby tissue or another part of the body and are used after injury, cancer removal, or reconstructive surgery.

## Skin Graft

Skin taken from one area of the body and placed over a wound, burn, ulcer, or surgical defect. A graft must obtain blood from the wound surface to survive and may fail, scar, change color, or require further surgery.

### Alternate Names:

Skin Grafting

## Skin-Grafting

Surgery that moves skin from a donor area to cover a wound, burn, ulcer, or tissue defect. The graft needs blood supply from the wound bed and may fail, scar, or change color.

## Skin Immune System

The cells and protective substances in skin that recognize germs, remove damaged cells, and control inflammation. It works with the skin barrier to prevent infection and support healing after injury.

## Skin Induration

Unusual hardness or firmness of the skin or tissue beneath it. Causes include scar formation, inflammation, swelling, infection, an infiltrating disease, or a growth; examination determines the cause.

## Skin Infection

Infection of the skin or underlying soft tissue, commonly caused by bacteria, fungi, or viruses. Redness, warmth, swelling, pain, drainage, or fever may occur; rapidly spreading redness, severe pain, blisters, or systemic illness requires prompt medical assessment.

## Skin Integrity

The condition of skin and underlying tissue when it remains intact and protected from injury. Prolonged pressure, friction, moisture, poor blood flow, and illness can cause breakdown, especially over bony areas.

## Skin Irritation

Inflammation or discomfort caused by direct contact with an irritant, often producing redness, burning, itching, dryness, or scaling.

## Skin Lesion

An abnormal area of skin that may result from infection, inflammation, injury, allergy, or a benign or malignant growth.

## Skin Lesion Aspiration

Removal of fluid or cells from a skin lump with a



---

needle for testing or drainage.

### **Skin Lesion Biopsy**

Removal of part or all of a skin lesion for microscopic examination to diagnose infection, inflammatory disease, benign tumor, or skin cancer.

### **Skin Lesion KOH Exam**

Microscopic examination of skin scrapings treated with potassium hydroxide to detect fungal hyphae or yeast in suspected superficial fungal infection.

### **Skin Lesion Of Blastomycosis**

A blastomycosis skin lesion results from dissemination of *Blastomyces* fungal infection and may appear as verrucous or ulcerated lesions with raised borders.

### **Skin Lesion Removal**

Removal of an unwanted, suspicious, or diseased area of skin by shaving, cutting, freezing, scraping, or another method. Tissue may be sent for laboratory examination; bleeding, infection, scarring, and incomplete removal are possible.

### **Skin Lumps**

Skin lumps are localized bumps or masses caused by cysts, lipomas, inflamed follicles, infections, benign tumors, or less commonly cancer; persistence or change warrants assessment.

### **Skin Mastocytoma**

A localized collection of mast cells in the skin, usually a yellow-brown or reddish plaque or nodule in a child that may swell, itch, or redden when rubbed.

### **Skin Microbiome**

The community of bacteria, fungi, viruses, and other microorganisms normally living on the skin. They interact with the skin and immune system; changes in this community may contribute to some skin diseases.

### **Skin Mycosis**

A fungal infection of the skin, hair, or nails, also called cutaneous mycosis or dermatomycosis. It may

be superficial or extend deeper into the skin and can cause scaling, itching, inflammation, discoloration, or lesions; diagnosis depends on the appearance and, when needed, microscopy or culture.

### **Skin Necrosis**

Death of skin tissue from ischemia, infection, pressure, toxins, trauma, vasculitis, or severe inflammation, producing discoloration, blisters, ulcers, or black eschar.

### **Skin Neoplasm**

A skin neoplasm is an abnormal growth in the skin that may be benign, precancerous, or malignant.

### **Skin Nodules**

Firm, raised lumps in or under the skin. They may result from inflammation, infection, injury, cysts, or benign or malignant growths.

### **Skin Of Color**

A broad term for people with more deeply pigmented skin. Some conditions may appear differently, and dark marks or raised scars may develop more readily after inflammation or injury; skin cancer can occur in every skin tone.

### **Skin Or Nail Culture**

Laboratory culture of skin, hair, or nail material to identify bacteria or fungi when infection is suspected; results require clinical correlation.

### **Skin Pigmentation**

Skin color is produced mainly by melanin made by pigment-producing cells. Sun exposure, hormones, inflammation, medicines, and illness can make areas darker or lighter; new or changing discoloration may need examination.

### **Skin Resurfacing**

Procedures that improve skin texture, scars, wrinkles, or uneven color by removing or changing surface skin layers. Methods include lasers, chemical peels, and abrasion; risks include infection, pigment changes, and scarring.

---

## Skin Scraping

Removal of a small amount of surface skin scale or material for laboratory examination, commonly to look for fungi, mites, or other causes of a rash.

## Skin Self-Exam

A skin self-exam is a regular check of the skin for new or changing moles, sores, lumps, or patches that may need medical assessment.

## Skin Tag

A small, harmless (benign) growth of soft skin that hangs from a narrow stalk (peduncle). Commonly appearing in body folds and friction areas—such as the neck, armpits, groin, eyelids, and under the breasts—they consist of loose collagen fibers and small blood vessels. While medically harmless, they are frequently associated with skin friction, aging, pregnancy, obesity, or insulin resistance, and can be easily removed if irritated or for appearance.

### Alternate Names:

Acrochordon; Fibroepithelial Polyp; Cutaneous Papilloma

## Skin Test

Alternate name; see *diagnostic or immunologic skin tests, including prick, patch, and tuberculin testing*.

## Skin Turgor

The skin's ability to return to its normal position after being gently pinched. Reduced turgor may occur with dehydration, although it is less reliable in older adults.

## Skin Writing

A skin condition in which light scratching or pressure causes temporary raised red lines or welts.

## Skipped Heartbeats

Alternate name; see *Palpitation*.

## Skull

The bony framework of the head that protects the brain and supports the face, eyes, ears, and jaw.

## Skull Fracture

A break in one or more skull bones caused by trauma. It may occur with bleeding inside the skull, brain injury, or leakage of clear fluid from the nose or ear and requires medical assessment.

## Skull X-Ray

A skull X-ray uses low-dose radiation to show skull bones and may help assess selected fractures, foreign bodies, or bone abnormalities.

## Slanted Ear

An ear whose upper edge slopes unusually compared with typical ear orientation; it may be isolated or part of a genetic condition.

## SLAP Lesion

A tear in the upper rim of the shoulder socket, sometimes where the biceps tendon attaches. It may follow a fall, lifting injury, or repeated overhead activity and can cause pain, clicking, weakness, or instability.

## Slapped Cheek Disease

Alternate name; see *Parvovirus infection*.

## Slapped Cheek Syndrome

Erythema infectiosum, a parvovirus B19 infection causing a slapped-cheek rash and sometimes joint or hematologic complications.

## SLAP Tears

Tears of the superior glenoid labrum extending from anterior to posterior, sometimes involving the biceps tendon attachment. They may cause pain, clicking, or instability, especially with overhead activity.

## SLE

Systemic lupus erythematosus, a chronic autoimmune disease that can affect skin, joints, kidneys, blood, nervous system, and other organs.

## Sleep

A naturally recurring state in which awareness and responses to the surroundings decrease while the

---

brain and body follow organized patterns. Sleep supports learning, memory, emotional health, physical recovery, immune function, and metabolism.

**Alternate Names:**

Somnolence; Slumber

**Sleep Disorders in Children**

Children and teens need age-appropriate sleep, and disorders may involve insomnia, circadian delay, sleep apnea, parasomnias, restless legs, or insufficient opportunity. Snoring, pauses in breathing, unusual daytime sleepiness, mood change, or poor school performance warrants evaluation.

**Sleep Apnea**

Repeated pauses or reductions in breathing during sleep. It can result from the airway closing, the brain failing to signal breathing, or both, and may cause loud snoring, gasping, poor sleep, daytime sleepiness, and low oxygen.

**Sleep Apnoea**

Alternate name; see *Sleep Apnea*.

**Sleep Architecture**

The pattern and order of sleep stages during a night or other sleep period. It includes lighter sleep, deep sleep, and dream-related REM sleep; illness, medicines, age, stress, and sleep disorders can change how much time is spent in each stage.

**Sleep Bruxism**

Repeated grinding or clenching of the teeth during sleep, which may cause tooth wear, jaw pain, or headaches.

**Sleep Debt**

The accumulated difference between the amount of sleep a person needs and the amount obtained over time; it may impair alertness, mood, cognition, and performance.

**Sleep Diary**

A daily record of sleep and wake times, awakenings, naps, medicines, caffeine, and related factors used

to assess sleep patterns and treatment response.

**Sleep Disorder**

Alternate name; see *Sleep Disorders*.

**Sleep Disorders**

A group of conditions affecting sleep quality, timing, duration, breathing, movement, or behavior. Major groups include insomnia, sleep-related breathing disorders, central disorders of hypersomnolence, circadian rhythm sleep-wake disorders, parasomnias, and sleep-related movement disorders. Examples include sleep apnea, narcolepsy, restless legs syndrome, and sleep paralysis.

**Sleep Disorders In Older Adults**

Sleep problems in later life, including insomnia, sleep apnea, restless legs, and circadian changes; medication effects, pain, depression, and medical illness should be assessed.

**Sleep Efficiency**

The percentage of time in bed that a person is actually asleep. It is calculated by dividing total sleep time by time in bed; a low value may occur with insomnia, repeated awakenings, or other sleep disorders.

**Sleep Fragmentation**

Repeated brief awakenings or partial arousals that interrupt otherwise continuous sleep. It can reduce restorative sleep and occur with sleep apnea, pain, noise, movement disorders, medicines, or illness.

**Sleep Hygiene**

Habits and surroundings that support regular, restorative sleep. Helpful measures include a consistent schedule, a dark and quiet bedroom, relaxing before bed, and limiting caffeine, alcohol, nicotine, and stimulating screen use near bedtime.

**Sleeping Sickness**

An infection caused by *Trypanosoma* parasites and spread by tsetse flies in sub-Saharan Africa. It may begin with fever and swollen nodes, then progress

---

to sleep disturbance, confusion, and other nervous-system problems.

### **Sleeplessness**

Alternate lay term; see *Insomnia Disorder*.

### **Sleep Onset Latency**

The time required to fall asleep after attempting to sleep; prolonged latency may occur with insomnia, anxiety, circadian misalignment, stimulants, or another sleep disorder.

### **Sleep Paralysis**

Temporary inability to move or speak when falling asleep or waking, while awareness is partly or fully preserved. Episodes usually last seconds to minutes and may include frightening hallucinations. Recurrent episodes can be associated with sleep deprivation, irregular sleep, narcolepsy, or other sleep disorders.

### **Sleep Related Breathing Disorders**

Sleep-related breathing disorders cause abnormal ventilation or oxygenation during sleep, including obstructive and central sleep apnea and sleep-related hypoventilation. Snoring, witnessed pauses, gasping, morning headache, or daytime sleepiness may prompt sleep testing and treatment.

#### **Alternate Names:**

Sleep-Disordered Breathing; SDB

### **Sleep-Related Hypoventilation**

Breathing that is too shallow during sleep to remove enough carbon dioxide or maintain normal oxygen. It can result from obesity, lung or muscle disease, chest-wall problems, brain disorders, or sedating medicines.

### **Sleep-Related Hypoxemia**

Abnormally low blood oxygen during sleep that may occur with sleep-disordered breathing, lung disease, neuromuscular weakness, hypoventilation, or other medical conditions.

### **Sleep Spindles**

Short bursts of regular brain activity that occur mainly during stage 2 non-REM sleep. They are seen on a sleep recording and are thought to help protect sleep and process newly learned information.

### **Sleep Studies**

Overnight tests that record brain activity, eye movements, heart rate, breathing, blood oxygen, body position, and limb movements. They help identify sleep apnea, unusual movements, seizures during sleep, and other sleep disorders.

#### **Alternate Names:**

Sleep Diagnostic Study; Nocturnal Somnography

### **Sleepwalking**

Recurrent walking or other complex behavior during sleep, usually with little memory afterward. It is more common in children and may be triggered by sleep deprivation, stress, fever, medicines, or other sleep disorders; injury prevention is important.

### **Sleepwalking And Children**

Sleepwalking in children is usually benign and often resolves with age; regular sleep, hazard prevention, and evaluation of frequent, violent, injurious, or atypical episodes are important.

### **Sleepwalking Disorder**

A sleep disorder involving repeated episodes of walking or other complex behavior during sleep, usually with little memory afterward, that cause injury, distress, sleep disruption, or impaired daytime functioning.

### **Sleeve Gastrectomy**

An operation that removes most of the stomach, leaving a narrow sleeve-shaped section. It limits how much food the stomach can hold and changes appetite hormones; bleeding, leakage, reflux, and nutritional problems can occur.

### **Sleeve Lobectomy**

An operation that removes a lung lobe and the nearby affected part of the main breathing tube,

---

then reconnects the remaining airway. It can remove a centrally located tumor while preserving more healthy lung than removing the whole lung. Possible complications include air leakage, narrowing of the airway, infection, and bleeding.

**Alternate Names:**

Bronchial Sleeve Resection; Sleeve Resection of the Lung

**Sliding Filament**

The process by which muscles contract: two protein fibers slide past each other, shortening the muscle's working units. Calcium and energy from ATP allow the fibers to attach, pull, release, and repeat the cycle.

**Sliding Hiatal Hernia**

A sliding hiatal hernia occurs when the stomach and gastroesophageal junction move upward through the diaphragm, often contributing to reflux.

**Sling**

A supportive device or surgically placed strip used to support an injured limb or urinary structures.

**Slipped Capital Femoral Epiphysis**

A hip problem in growing children or teenagers in which the upper end of the thigh bone slips through its growth area. It may cause hip, groin, thigh, or knee pain and a limp and needs urgent assessment to prevent further slipping.

**Slipped Disk**

A spinal disk whose tough outer layer tears or weakens, allowing its inner material to bulge and irritate nearby nerves. It may cause back or neck pain, numbness, weakness, or sciatica.

**Alternate Names:**

Herniated Disc; Intervertebral Disc Prolapse

**Slipped Vertebra**

Forward movement of one spinal bone compared with the bone below it, also called spondylolisthesis. It may cause back pain, stiffness, hamstring tightness, or pressure on a nerve, although some people

have no symptoms.

**Alternate Names:**

Spondylolisthesis

**Slipping Rib Syndrome**

Pain caused by excessive movement of the lower ribs, usually from loosened cartilage. The pain may be sharp and can sometimes be reproduced by pressing or moving the rib.

**Slit Lamp**

An eye-examination microscope that uses magnification and a narrow beam of light to inspect the eyelids, cornea, lens, and other eye structures. It helps detect infection, injury, cataract, and glaucoma-related changes.

**Alternate Names:**

Biomicroscope

**Slit-Lamp Biomicroscope**

Alternate name; see *Slit Lamp*.

**Slit Lamp Biomicroscopy**

A detailed eye examination using a bright, narrow beam of light and a microscope to view the front of the eye. Additional lenses can help examine the retina and other structures farther back.

**Slit-Lamp Exam**

Alternate name; see *Slit Lamp Biomicroscopy*.

**Slit Lamp Examination**

A fundamental ophthalmological diagnostic procedure utilizing a specialized high-intensity light source focused through a slit aperture paired with a binocular biomicroscope to examine the anterior segment of the eye (cornea, anterior chamber, iris, crystalline lens) and vitreous under high magnification.

**Alternate Names:**

Biomicroscopy

**Slit-Lamp Examination**

A diagnostic ophthalmologic procedure utilizing a binocular biomicroscope paired with an adjustable,

---

high-intensity slit-beam light source, allowing detailed optical sectioning and high-magnification stereoscopic examination of the eyelids, cornea, anterior chamber, iris, lens, and anterior vitreous.

**Alternate Names:**

Slit-Lamp Biomicroscopy; Anterior Segment Biomicroscopy

**SLL**

Alternate name; see *Small lymphocytic lymphoma*.

**Slough**

Soft, dead, or detached tissue covering or separating from a wound, ulcer, or inflamed surface. It can delay healing and may need gentle removal after blood flow, infection, and the wound's depth are assessed.

**Slowed or Arrested Breathing**

Pathologic depression of the respiratory rate (bradypnea) or complete cessation of ventilatory air flow (apnea), resulting from central nervous system depression, drug toxicity (such as opioid overdose), traumatic brain injury, upper airway obstruction, or terminal cardiopulmonary arrest requiring emergent airway management and resuscitation.

**Alternate Names:**

Bradypnea; Apnea; Respiratory Arrest; Hypopnea

**Slow Heartbeat**

Alternate name; see *Bradycardia*.

**Slow Heart Rate**

A heart rate slower than expected, often below 60 beats per minute in an adult. It may be normal in some people but can cause dizziness, fainting, tiredness, or shortness of breath when it reduces blood flow.

**Slow-Twitch Fiber**

A type of muscle fiber suited to steady activity over a long time. It uses oxygen efficiently and resists tiredness, helping with posture, walking, and endurance activities rather than short bursts of maximum force.

**Alternate Names:**

Type I Muscle Fiber; Red Muscle Fiber

**Slow-Wave Sleep**

The deepest stage of non-REM sleep, marked by slow brain waves and reduced response to sounds or touch. It supports physical recovery, memory, and hormone release and is often reduced by ageing, illness, alcohol, or sleep disruption.

**Alternate Names:**

Deep Sleep; Delta Sleep; Stage N3 Sleep

**Small Airway Disease**

Narrowing, inflammation, or blockage of the smallest breathing tubes in the lungs. It may cause cough, wheezing, breathlessness, or trapped air and can occur with asthma, COPD, infection, or other lung disease.

**Small Bowel Adenocarcinoma**

A malignant gland-forming epithelial tumor of the small intestine. It is uncommon and may cause occult bleeding, obstruction, abdominal pain, or weight loss; risk is increased in selected inherited and inflammatory disorders.

**Small Bowel Bacterial Overgrowth**

Excess bacteria in the small intestine, where fewer bacteria are normally present. It can cause bloating, abdominal discomfort, diarrhea, weight loss, and poor absorption of nutrients; some cases involve methane-producing organisms instead.

**Small Bowel Bleeding**

Bleeding from the small intestine. It may cause black or maroon stool, hidden blood loss, or iron-deficiency anemia and may require capsule endoscopy, enteroscopy, imaging, or other tests to find the source.

**Small Bowel Follow-Through**

A series of X-rays taken after swallowing contrast, usually barium, to follow its movement through the small intestine. It can show narrowing, obstruction, inflammation, or abnormal transit.



---

### Small Bowel Obstruction

Partial or complete blockage of the small intestine. Common causes include adhesions, hernias, tumors, and inflammatory narrowing. Symptoms may include abdominal pain, vomiting, distension, and inability to pass stool or gas.

### Small Bowel Prolapse

Alternate name; see *Small bowel prolapse (enterocele)*.

### Small Bowel Resection

Surgical removal of a diseased segment of the small intestine followed by reconnection or stoma formation when needed.

### Small Bowel Series

A series of X-ray images of the small intestine after swallowing contrast material, usually barium. It can show narrowing, blockage, abnormal movement, inflammation, tumors, or changes from Crohn disease.

### Small Bowel Tissue Smear/Biopsy

Examination of cells or a tissue sample from the small intestine to investigate poor nutrient absorption, celiac disease, inflammation, infection, or a tumor.

### Small-Bowel Transplantation

Surgery replacing a severely diseased or nonfunctioning small intestine with donor bowel. It may help people unable to receive enough nutrition through the gut but requires lifelong rejection-preventing medicines and monitoring.

### Small-Bowel Volvulus

Twisting of the small intestine that can block its contents and cut off its blood supply. It may cause severe abdominal pain, vomiting, swelling, and inability to pass stool or gas.

### Small Calorie

A small calorie is a unit of heat energy; food labels usually use a kilocalorie, written Calorie with a

capital C, equal to 1,000 small calories.

### Small Cell Carcinoma

A fast-growing cancer made of small cells, most often beginning in the lung and strongly linked to smoking. It may shrink at first with treatment but often returns or spreads early, including to the brain.

### Small-Cell Carcinoma

A fast-growing cancer, most often starting in the lung and strongly linked with smoking. It may shrink quickly with chemotherapy or radiation but often returns or spreads early, including to the brain.

### Small Cell Lung Cancer

An aggressive lung cancer strongly associated with smoking and early spread. It often responds initially to chemotherapy or radiation, but recurrence is common and treatment depends on the stage and overall health.

### Small Cell Lung Carcinoma

An aggressive, high-grade neuroendocrine malignancy of the lung strongly linked to heavy cigarette smoking, characterized histologically by small sheets of cells with scant cytoplasm, granular nuclear chromatin ("salt-and-pepper"), absence of nucleoli, high mitotic rates, and frequent paraneoplastic syndromes (SIADH, Lambert-Eaton).

#### Alternate Names:

SCLC

### Small Eye

An eye that is smaller than usual because it did not develop fully before birth. Vision may be reduced, especially when other eye structures are also affected, and assessment can identify associated conditions.

### Small Fiber Neuropathy

Damage to tiny nerves that carry pain, temperature, and automatic body signals. It can cause burning feet, tingling, poor temperature sensation, abnormal sweating, digestive symptoms, or heart-rate changes; diabetes is one possible cause.

---

### Small For Gestational Age

Birth weight or size below the expected range for gestational age, caused by constitutional small size, placental insufficiency, infection, genetic disease, or maternal factors.

### Small Incision Lenticule Extraction Surgery

Small incision lenticule extraction, or SMILE, is a laser eye procedure that removes a small piece of corneal tissue through a tiny incision to correct some cases of nearsightedness and astigmatism.

### Small Intestinal Bacterial Overgrowth

An abnormal increase or change in bacteria within the small intestine, often causing bloating, abdominal pain, nausea, diarrhea, fullness, weight loss, or nutrient deficiency. Causes include impaired intestinal movement, structural problems, surgery, and some systemic diseases; persistent symptoms require medical evaluation.

### Small-Intestinal Cancer

Malignant tumors arising in the small intestine, classified mainly as adenocarcinoma, neuroendocrine tumors, lymphomas, or sarcomas. They are far less common than colorectal cancer and often present late with vague abdominal pain, bleeding, obstruction, or weight loss; risk is increased in some inherited syndromes, celiac disease, and Crohn disease. Diagnosis uses endoscopy, capsule endoscopy, or imaging, and treatment depends on the tumor type and stage.

### Small Intestinal Fungal Overgrowth

A proposed condition in which excess fungi are found in the small intestine and may be associated with bloating, nausea, diarrhea, or abdominal discomfort. Symptoms are nonspecific, testing is not standardized, and other causes should be assessed before attributing symptoms to it.

### Small Intestinal Ischemia And Infarction

Inadequate blood supply to the small intestine that can progress to irreversible tissue death, perforation, sepsis, and shock.

### Small Intestinal Mucosa

The inner lining of the small intestine. Its folds, villi, and microvilli greatly increase the area available for digestion and nutrient absorption.

### Small Intestine

The long, coiled part of the digestive tract between the stomach and large intestine. It receives bile and pancreatic juices, completes digestion, and absorbs most nutrients and much of the body's water.

### Small Intestine Aspirate And Culture

Collection and culture of fluid aspirated from the small intestine to evaluate suspected bacterial overgrowth or selected infections; the procedure is invasive and results require clinical correlation.

### Small Intestine Cancer

A malignant tumor arising in the small intestine. Types include adenocarcinoma, neuroendocrine tumor, lymphoma, and sarcoma; symptoms and treatment depend on the tumor type and extent.

### Small Intestine Prolapse

Alternate name; see *Small bowel prolapse (enterocele)*.

### Smallpox

A severe contagious disease caused by variola virus, producing fever and rash; global eradication was certified in 1980.

### Small Round Blue Cell Tumor

A tumor whose cells look small, round, and blue under a microscope. This appearance occurs in several cancers, especially in children, so additional tissue tests are needed to identify the exact type.

### Small Saphenous Vein

A superficial vein running up the back of the calf and usually joining the popliteal vein behind the knee. It helps return blood from the foot and calf to the heart.

---

## Small Vessel Disease

Disease affecting small blood vessels, which can impair blood flow and contribute to brain, heart, kidney, or other organ problems.

### Alternate Names:

Coronary Microvascular Disease; Microangiopathy

## Small-Vessel Vasculitis

A group of diseases that inflame small blood vessels and can affect the skin or internal organs. Skin involvement may cause raised purple spots, blisters, or ulcers; other symptoms depend on the organs involved. Leukocytoclastic vasculitis is a common biopsy pattern in skin-limited small-vessel vasculitis, but the terms are not interchangeable in every case.

## SMART

A way to set goals that are specific, measurable, achievable, relevant, and time-limited. SMART goals state exactly what is wanted, how progress will be checked, and when the goal should be reached.

### Alternate Names:

SMART Criteria; Specific Measurable Goals

## Smashed Fingers

A smashed finger may injure skin, nail, bone, tendon, or nerve; deformity, severe pain, numbness, impaired movement, or blood under the nail may require urgent evaluation.

## Smear Of Duodenal Fluid Aspirate

A smear of duodenal fluid aspirate examines fluid from the first part of the small intestine for selected parasites or other microscopic findings.

## Smegma

A collection of shed cells, oils, and moisture around the genital folds, which may promote irritation if hygiene is poor.

## Smell

The sense of detecting airborne chemicals through receptors in the nasal lining and processing the signals in the brain. Reduced or lost smell can follow infection, nasal disease, head injury, medicines, or

neurologic disease.

## SMILE

A laser eye operation that corrects short-sightedness and some short-sighted astigmatism. The laser creates a thin piece of tissue inside the cornea, which is removed through a small opening without making a larger corneal flap.

## Smile Design

Planning smile based on facial features, gum display, tooth proportions. Digital smile design software available.

## Smith Fracture

An extra-articular fracture of the distal radius with volar (palmar) displacement and angulation of the distal fracture fragment ('reverse Colles fracture'), typically caused by a direct fall onto a flexed wrist.

### Alternate Names:

Reverse Colles Fracture; Volar Distal Radius Fracture

## Smith-Lemli-Opitz Syndrome

A rare inherited disorder affecting cholesterol production before birth. It may cause growth and developmental problems, distinctive facial features, heart or limb abnormalities, feeding difficulty, and genital differences.

## Smith-Magenis Syndrome

A genetic neurodevelopmental disorder, usually involving deletion of 17p11.2 or an RAI1 variant. It causes developmental delay, intellectual disability, distinctive facial features, sleep disturbance, behavioral dysregulation, and variable congenital abnormalities.

## Smoke Evacuator

A device that captures and filters smoke produced during surgery by electrical, laser, or ultrasonic instruments. Removing the plume reduces staff and patient exposure to particles, irritating gases, and potentially infectious material.

---

## Smoke Inhalation

Injury from breathing smoke, hot gases, or combustion products, potentially causing airway burns, poisoning, and lung injury.

## Smoke Inhalation Injury

Pulmonary injury caused by inhalation of toxic products of combustion.

## Smokeless Tobacco

Smokeless tobacco delivers nicotine and causes dependence, oral lesions, gum recession, tooth loss, cardiovascular risk, and cancers of the mouth, throat, and pancreas. Quitting reduces harm, and counseling plus approved medicines improve cessation success.

### Alternate Names:

Chewing Tobacco; Snuff; Dip

## Smoker Keratosis

Alternate name; see *Leukoplakia*.

## Smoker Lung

Alternate name; see *Chronic obstructive pulmonary disease*.

## Smoking

Inhalation of smoke from burning tobacco or other substances, associated with addiction, cancer, cardiovascular disease, and lung disease.

## Smoking And Asthma

Smoking irritates and inflames airways, worsens asthma control, increases attacks, and reduces treatment response; stopping smoking and avoiding secondhand smoke improve outcomes.

## Smoking And COPD

Smoking is the major preventable cause of COPD and accelerates lung-function loss; quitting reduces progression and exacerbations even after COPD is established.

## Smoking and Pregnancy

Smoking during pregnancy exposes the fetus to nicot-

tine and other harmful chemicals and increases risks such as poor growth, preterm birth, placental problems, and stillbirth. Stopping at any time is beneficial.

## Smoking And Surgery

Smoking before or after an operation increases the risk of breathing problems, wound infection, poor healing, blood clots, heart complications, and failure of repaired tissue. Stopping as early as possible improves recovery, even if surgery is soon.

## Smoking Cessation

The process of stopping tobacco or nicotine use. It may involve counseling, behavioral support, medicines, or nicotine replacement.

## Smoking Cessation Medications

Medicines that reduce nicotine withdrawal and cravings, including nicotine replacement, bupropion, and varenicline, used with behavioral support and individualized for health risks and preferences.

## Smoldering Multiple Myeloma

An asymptomatic plasma-cell disorder with increased clonal plasma cells or monoclonal protein but no myeloma-defining organ damage; it requires risk-based monitoring for progression to active multiple myeloma.

### Alternate Names:

Asymptomatic Myeloma; Indolent Myeloma; SMM

## Smooth Muscle

Muscle that works automatically in the walls of hollow organs, blood vessels, and airways. It controls processes such as bowel movement, bladder emptying, blood-vessel width, airway narrowing, and childbirth contractions.

## Smothering

Mechanical obstruction of the external nose and mouth that prevents effective ventilation. It may be caused by a covering, compression, or an obstructing soft material; findings are often nonspecific and must be interpreted with the scene, history, and complete examination.

---

## Snake Bite

Snake bites can cause puncture wounds, tissue injury, bleeding, paralysis, or venom-related effects on clotting, nerves, muscles, kidneys, or the heart. Venomous bites can become life-threatening, and cutting, sucking, icing, or applying a tourniquet can worsen injury.

### Alternate Names:

Snake Envenomation; Ophidism

## Snake-Bite

A wound from a snake that may cause local injury, envenomation, infection, or systemic toxicity.

## Snake Bites

Wounds caused by a snake's teeth; venomous bites may cause pain, swelling, bleeding, tissue damage, paralysis, abnormal clotting, or shock. Cutting, sucking, icing, or applying a tourniquet can worsen injury.

## Snapping Finger

Trigger finger, painful catching or locking of a finger caused by stenosing inflammation around a flexor tendon.

## Snapping Hip Syndrome

A clicking or snapping sensation in the hip, sometimes with pain, caused by a tendon moving over bone or by a problem inside the joint. Activity adjustment and exercises often help; persistent pain may need imaging or other treatment.

## Sneddon Syndrome

A rare disorder in which small arteries become blocked, causing a net-like purplish skin pattern and repeated strokes or stroke-like events. Headache, high blood pressure, memory problems, or heart-valve changes may also occur.

## Sneezing

A forceful reflex expulsion of air through the nose and mouth, often triggered by nasal irritation.

## Snellen Chart

A chart of letters or symbols used to measure visual acuity at a specified distance.

## Snoring

Noisy breathing during sleep caused by vibration of relaxed or narrowed tissues in the throat. It can be harmless, but loud frequent snoring with pauses in breathing, choking, morning headaches, or daytime sleepiness may indicate sleep apnoea.

## Snoring And Sleep Apnea

Snoring is the sound produced by vibrating soft tissues in the upper airway during sleep, often resulting from partial airway obstruction. Sleep apnea is a more serious condition characterized by repeated episodes of complete or partial airway collapse during sleep, leading to intermittent hypoxia, frequent arousal, and fragmented sleep.

## Snort

A snort is a forceful nasal or respiratory sound; it may occur with congestion, sleep-disordered breathing, irritation, or inhalation of a substance.

## Snot

Alternate informal term; see *Mucus*.

## Snow Blindness

A painful temporary burn of the eye's clear front surface caused by intense ultraviolet light, often reflected from snow, ice, water, or sand. Symptoms may include gritty pain, tearing, redness, light sensitivity, and blurred vision.

## SNRIs

A class of antidepressants that increase the activity of serotonin and norepinephrine, chemical messengers between nerve cells. They treat depression, some anxiety disorders, and certain nerve or long-lasting pain conditions.

## Snuff

Snuff is finely ground tobacco inhaled through the nose or placed in the mouth and can cause nicotine

---

dependence, oral disease, and cardiovascular harm.

### **Snuffles**

A runny or congested nose; in newborns, persistent snuffles can be a sign of congenital syphilis.

### **Social Anxiety Disorder**

Intense, persistent fear of being judged, embarrassed, or rejected in social or performance situations. It can lead to avoidance, physical anxiety symptoms, and difficulty with school, work, relationships, or daily activities.

### **Social History**

Information about a person's living situation, relationships, occupation, education, tobacco or substance use, diet, safety, and other social factors relevant to health and care.

### **Social Isolation**

Limited contact or interaction with other people. Persistent isolation may adversely affect mental and physical health.

### **Social Media Addiction**

Social media addiction is a pattern of difficult-to-control use that causes distress or interferes with sleep, work, school, relationships, or daily activities. It is not a universally defined formal diagnosis.

### **Social Phobia**

Alternate name; see *Social anxiety disorder (social phobia)*.

### **Social Worker**

A professional who helps patients and families manage practical, emotional, and social problems related to illness. They may arrange community resources, support discharge planning, assist with benefits, and provide counseling or referrals.

### **Sociophobia**

Older term; see *Social Anxiety Disorder*.

## **Sodium**

A mineral and body salt helping control fluid balance, nerves, muscles, and blood pressure; abnormal levels can be harmful.

### **Sodium Bicarbonate Therapy**

Treatment using sodium bicarbonate to make overly acidic blood less acidic in selected conditions, such as some cases of kidney disease. It can increase sodium and fluid load and requires monitoring.

### **Sodium Bisulfate Poisoning**

Poisoning from sodium bisulfate, an acidic chemical used in some cleaners and pool products. It can burn the skin, eyes, mouth, and digestive tract; fumes may irritate breathing passages.

### **Sodium Blood Test**

A blood test measuring serum sodium, a major extracellular electrolyte involved in water balance and nerve and muscle function; results are interpreted with fluid status and other electrolytes.

### **Sodium Carbonate Poisoning**

Poisoning from sodium carbonate, an alkaline chemical found in some cleaning products. It may irritate or burn the eyes, skin, mouth, and digestive tract; vomiting can worsen injury after swallowing it.

### **Sodium Chloride**

The salt sodium chloride; sterile solutions are used for fluid replacement, irrigation, and drug dilution.

### **Sodium Cromoglycate**

A mast-cell stabilizer used in selected allergy and asthma treatments; also called cromolyn.

### **Sodium Diatrizoate**

A water-soluble iodinated contrast agent used for selected radiographic examinations. It can cause allergic-like reactions, kidney stress, or aspiration complications, so its use depends on the procedure and patient risks.



---

## Sodium Hydroxide Poisoning

Severe poisoning from sodium hydroxide, a strong alkali in some drain and oven cleaners. It can rapidly burn the mouth, throat, esophagus, stomach, skin, and eyes; emergency assessment is required and vomiting must not be induced.

## Sodium Hypochlorite

The active chemical in many bleach and disinfectant products. It kills many germs when used correctly, but concentrated liquid can burn skin, eyes, and the digestive tract; mixing it with acids or ammonia can release dangerous gas.

## Sodium Hypochlorite Poisoning

Poisoning from bleach containing sodium hypochlorite. Swallowing may irritate or burn the mouth and digestive tract; mixing bleach with acids or ammonia can release dangerous gases causing cough, chest pain, or breathing failure.

## Sodium In Diet

Dietary sodium helps regulate fluid balance and nerve and muscle function, but excess intake can raise blood pressure; processed foods are a major source and intake should follow individual health advice.

## Sodium Polystyrene Sulfonate

A resin that binds potassium in the intestine and has been used to lower high blood potassium. It can cause constipation and serious intestinal injury, so its use requires careful medical assessment.

## Sodium-Potassium Pump

A ubiquitous, electrogenic P-type ATPase enzyme complex in eukaryotic plasma membranes that hydrolyzes one molecule of ATP to export three sodium ions from the cell in exchange for importing two potassium ions, maintaining intracellular ion gradients, resting membrane potential, and cellular osmotic volume.

### Alternate Names:

Na<sup>+</sup>/K<sup>+</sup>-ATPase

## Sodium Urine Test

A urine test measuring sodium excretion, used with blood tests and clinical findings to assess volume status, kidney handling of sodium, and causes of hyponatremia or hypertension.

## Sodium Valproate

An anticonvulsant and mood stabilizer; it carries important pregnancy and reproductive risks and requires specialist prescribing.

## Soft Markers For Aneuploidy

Ultrasound findings during pregnancy that are common in healthy fetuses but occur somewhat more often when there is an extra or missing chromosome. They do not diagnose a chromosome condition and are interpreted with other screening results.

## Soft Tissue Augmentation

Restoration or enhancement of volume and contour in skin or underlying soft tissue using fillers, fat transfer, implants, or other reconstructive techniques.

## Soft-Tissue Calcinosis

Buildup of calcium deposits in skin, muscles, tendons, or other soft tissues. The deposits may form hard lumps, cause pain or limited movement, break through the skin, or occur because of kidney disease, abnormal calcium levels, inflammation, or tissue damage.

## Soft Tissue Injury

Damage to muscles, tendons, ligaments, fascia, skin, or other non-bony tissues, often caused by trauma or overuse.

## Soft Tissue Rheumatism

An older broad term for pain or inflammation affecting muscles, tendons, ligaments, or bursae. Modern diagnosis usually names the specific condition, such as tendinitis, bursitis, or a muscle strain.

## Soft Tissue Sarcoma

A group of malignant tumors arising in muscle, fat,

---

fibrous tissue, blood vessels, or other connective tissues. Treatment depends on the subtype, location, stage, and relationship to nearby structures.

**Alternate Names:**

Soft-Tissue Malignancy; Musculoskeletal Sarcoma

**Soft Tissue Tumors**

Growths arising in muscle, fat, fibrous tissue, blood vessels, nerves, or other supporting tissues. Many are harmless, but some are cancerous; a growing, deep, firm, painful, or persistent lump needs assessment.

**Soil Contamination**

The presence of harmful substances in soil, such as heavy metals, pesticides, fuel, or industrial waste. People may be exposed through food, water, dust, or skin; testing, removal, covering, or treatment of the soil can reduce risk.

**Solanine Plant Poisoning**

Toxic ingestion of glycoalkaloids (primarily solanine and chaconine) present in high concentrations in green potato tubers, potato sprouts, and green nightshade foliage, causing acute gastrointestinal and neurological symptoms.

**Alternate Names:**

Potato Glycoalkaloid Toxicity; Green Potato Poisoning; Solanine Intoxication

**Solar Elastosis**

Solar elastosis is degeneration and thickening of elastic fibers in skin after long-term ultraviolet exposure. It contributes to leathery, wrinkled, yellowish sun-damaged skin.

**Solar Keratosis**

Alternate name; see *Actinic Keratosis*.

**Solar Lentigines**

Alternate name; see *Age Spots*.

**Solar Plexus**

A network of involuntary nerves in the upper abdomen, centered around the celiac plexus. It helps

regulate abdominal organs, and irritation or injury may contribute to upper-abdominal pain or altered organ function.

**Solar Purpura**

Alternate name; see *Actinic Purpura*.

**Solar Urticaria**

A rare allergic-type reaction to sunlight or other light that causes itchy red raised patches within minutes on exposed skin. The rash usually improves after leaving the light, but severe widespread symptoms need medical care.

**Solder Poisoning**

Poisoning from swallowing solder or inhaling fumes or dust during heating. Products may contain lead, tin, antimony, or other metals and can cause stomach symptoms, anemia, nerve injury, or kidney damage; exposure needs evaluation.

**Soleus Muscle**

A deep calf muscle that joins the Achilles tendon and helps point the foot downward. It is important for standing, walking, and running; injury or overuse can cause calf pain and weakness.

**Solid Pseudopapillary Pancreatic Tumor**

Alternate name; see *Pancreatic Cysts*.

**Solitary Fibrous Tumor**

A solitary fibrous tumor is a usually slow-growing growth of connective tissue that can arise in the pleura, meninges, or other sites and may rarely behave aggressively.

**Solitary Kidney**

A solitary kidney means a person has one functioning kidney, because the other kidney is absent, removed, or not working. Many people remain healthy but may need blood-pressure and kidney-function monitoring.

**Solitary Pulmonary Nodule**

A single small spot in the lung surrounded by oth-

---

erwise normal lung tissue. It may be a scar, infection, harmless growth, or early cancer; its size, appearance, and change over time determine whether follow-up or biopsy is needed.

### **Solution**

A homogeneous mixture in which one or more substances are dissolved in a solvent; medical solutions include saline and antiseptics.

### **Solvent Abuse**

Harmful inhalation or ingestion of volatile solvents, which can cause neurologic, cardiac, liver, kidney, or respiratory injury.

### **Solvent Exposure**

Contact with organic solvents by inhalation, ingestion, or skin absorption. Acute exposure may cause irritation or central nervous system effects; repeated exposure may damage organs.

### **Somatic**

Relating to the body, especially the body wall and musculoskeletal system, or to physical symptoms.

### **Somatic Mutation**

A genetic change acquired by a body cell during life rather than inherited from a parent. It is passed to descendants of that cell and may contribute to cancer, but it usually is not passed to children.

### **Somatic Nervous System**

The part of the peripheral nervous system that carries sensory information from the skin, muscles, and joints to the central nervous system and controls voluntary movement of skeletal muscles. Its functions also include reflexes that may occur without conscious control.

### **Somatic Symptom And Related Disorders**

A group of mental disorders involving distressing physical symptoms, excessive health-related thoughts or behaviors, illness fears, functional neurologic symptoms, or intentionally produced illness. Examples include somatic symptom disorder, illness

anxiety disorder, functional neurologic disorder, and factitious disorder. Malingering is separate because it involves intentional symptom production for an external benefit.

### **Somatic Symptom Disorder**

A mental disorder in which one or more physical symptoms cause significant distress or disruption of daily life, together with excessive thoughts, feelings, or behaviors related to the symptoms. The symptoms are real and may occur with or without an identified medical condition; their severity is not determined solely by whether a medical cause is found.

### **Somatic Variant**

A change in DNA acquired by a body cell after conception rather than inherited from a parent. It usually cannot be passed to children but may affect only some tissues or contribute to cancer.

### **Somatization**

The experience or expression of psychological distress through physical symptoms such as pain, fatigue, or stomach problems. The symptoms are real and not intentional; assessment should consider both physical and psychological causes.

### **Somatostatin**

A hormone made in the brain, pancreas, and digestive tract that slows growth hormone, insulin, glucagon, and digestive-hormone release. It also reduces movement and secretion within parts of the digestive system.

### **Somatostatin Analog**

A medicine that imitates somatostatin, a hormone that slows the release of several hormones and digestive secretions. It is used for selected hormone-producing tumors, acromegaly, severe diarrhea, and some bleeding conditions.

### **Somatostatinoma**

A rare hormone-producing tumor usually arising in the pancreas or small intestine. It may cause diabetes, diarrhea or greasy stools, gallstones, weight

---

loss, or no symptoms; treatment depends on whether it has spread and may include surgery or cancer medicines.

### **Somatotropic Pituitary Adenoma**

A usually noncancerous pituitary tumor that makes too much growth hormone. In adults it causes acromegaly, with enlargement of the hands, feet, jaw, or other tissues; in children it can cause excessive height before growth ends.

### **Somatotropin**

Growth hormone, produced by the pituitary gland. It supports childhood growth and influences protein, fat, and carbohydrate metabolism; excess or deficiency can cause characteristic growth and metabolic disorders.

### **Somatotype**

A descriptive body-build classification; it is not a disease diagnosis.

### **Somnambulism**

Sleepwalking or performing other activities while only partly awake, usually during deep sleep. Episodes are more common in children and may be triggered by lack of sleep, stress, illness, or medicines; safety measures are important, especially if wandering or injury occurs.

### **Somniloquy**

Talking, laughing, or making sounds during sleep without knowing it. It is usually harmless, but frequent or disruptive episodes may occur with sleepwalking, nightmares, breathing problems during sleep, medicines, or other sleep disorders.

#### **Alternate Names:**

Nocturnal Vocalization; Sleep Utterance; Parasomnia Vocalization

### **Somnolence**

Unusual sleepiness or a tendency to fall asleep when a person should be awake. It may result from too little sleep, medicines, alcohol or drugs, sleep disorders, metabolic illness, or neurologic disease.

### **Somnolent**

Somnolent means unusually sleepy or drowsy, which may result from sleep deprivation, medicines, intoxication, metabolic illness, or neurologic disease.

### **Somogyi Effect**

A rebound hyperglycemia occurring in the early morning in response to an unrecognized, nocturnal episode of hypoglycemia (typically occurring between 2:00 AM and 3:00 AM) in patients with diabetes mellitus treated with insulin. Hypoglycemia triggers a surge of counterregulatory stress hormones (epinephrine, glucagon, cortisol, and growth hormone) stimulating hepatic gluconeogenesis and glycogenolysis; distinguished from the dawn phenomenon by checking 3:00 AM blood glucose levels.

#### **Alternate Names:**

Somogyi Phenomenon; Rebound Hyperglycemia

### **Somogyi Phenomenon**

A proposed pattern of morning hyperglycemia following unrecognized overnight hypoglycemia, with counterregulatory hormone release. Continuous or overnight glucose measurements are needed to distinguish it from the more common dawn phenomenon before changing therapy.

### **Sonographer**

A healthcare professional trained to obtain diagnostic ultrasound images. They position the patient, operate the scanner, record appropriate views and measurements, and document findings for interpretation by a qualified clinician.

### **Sonohysterography**

Sonohysterography is an ultrasound examination in which sterile fluid is placed in the uterus to outline the uterine cavity and assess problems such as polyps or fibroids.

### **Soporifics**

Medicines or substances that cause sleepiness or promote sleep. They may impair alertness, coordination, and breathing, especially when combined with alcohol or other sedatives.

---

## Sorbitol

A sugar alcohol used as a sweetener and in some medicines. Larger amounts draw water into the bowel and act as a laxative, but may cause gas, cramps, diarrhea, or dehydration.

## Sore

Painful or tender, or describing an open area on skin or a moist body lining. A sore may result from injury, infection, inflammation, pressure, allergy, or another medical condition.

## Sore Throat

Pain, scratchiness, or irritation in the throat. Common causes include viral infections, bacterial infections, allergies, dry air, smoke, voice strain, and acid reflux; persistent or severe symptoms, drooling, inability to swallow, a muffled voice, neck swelling, or breathing difficulty need prompt medical assessment.

## Sotalol

A medicine with beta-blocking and class III antiarrhythmic effects, used for selected cardiac rhythm disorders.

## Sotos Syndrome

A genetic disorder that causes unusually rapid growth in childhood, a large head, distinctive facial features, and delayed development or learning difficulties. Coordination, behavior, seizures, heart problems, or kidney problems may also occur. Growth usually slows with age. A change in the *NSD1* gene causes most cases.

### Alternate Names:

Cerebral Gigantism; NSD1-Related Overgrowth Syndrome

## Sound

Mechanical energy transmitted as waves and perceived by hearing; clinical sound assessment includes speech and respiratory or cardiac sounds.

## Source Control

A procedure or intervention that eliminates, drains,

debrides, or otherwise controls an anatomic focus of infection or contamination. Examples include drainage of an abscess, removal of infected tissue or device, and repair of a perforation.

## Soy

Soy is a legume providing protein and isoflavones; it is nutritious for many people but can cause allergy and may interact with some medicines or thyroid treatment timing.

## Soya

Soybean and products made from it; soy is a food source and can cause allergy in susceptible people.

## Soybeans

Soybeans are legumes that provide protein, fiber, and compounds called isoflavones. They are used in foods such as tofu and soy milk; allergy is possible.

## Soy Protein Intolerance

Adverse gastrointestinal reaction to soy proteins, most often seen in infants and young children. It may cause vomiting, diarrhoea, abdominal pain, blood or mucus in the stool, feeding difficulty, or poor growth.

## Space Medicine

The medical specialty concerned with human health and performance during aviation and space travel. It addresses altered gravity, radiation, isolation, motion sickness, sleep, emergencies, exercise, and returning safely to Earth.

## Spacer

A chamber attached to a metered-dose inhaler that holds the released medicine briefly, making it easier to breathe into the lungs and reducing medicine left in the mouth or throat.

### Alternate Names:

Valved Holding Chamber; Aerosol Spacer

## Spanish Fly

A historical name for cantharidin-containing blister beetles; ingestion is poisonous and it is not a safe

---

aphrodisiac.

### **Spasm**

An involuntary, often painful tightening of one muscle or a group of muscles. Spasms may result from fatigue, dehydration, nerve disease, injury, medicines, inflammation, or problems affecting the brain or spinal cord.

### **Spasmodic Dysphonia**

Spasmodic dysphonia is a focal laryngeal dystonia causing involuntary voice-box spasms and strained, breathy, or interrupted speech.

### **Spasmodic Torticollis**

A form of neck dystonia with involuntary muscle contractions causing twisting, tilting, or abnormal neck posture.

### **Spasmus Nutans**

An uncommon early-childhood syndrome of rapid shimmering eye movements, head nodding, and torticollis that usually resolves but requires evaluation to exclude neurologic or ocular disease.

### **Spastic**

Describes muscles that are unusually tight and resist being stretched, especially when moved quickly. It may cause stiffness, jerking, abnormal posture, and difficulty controlling movement after nervous-system injury.

### **Spastic Colon**

Alternate name; see *Irritable bowel syndrome*.

### **Spastic Diplegia**

A clinical subtype of cerebral palsy historically associated with prematurity and periventricular leukomalacia, characterized by symmetrical spastic hypertonia affecting the lower extremities significantly more than the upper extremities, producing a characteristic "scissor gait" with hip adduction and internal rotation.

### **Spastic Gait**

An abnormal walking pattern characterized by stiff, dragging legs, internal rotation of the foot, plantar flexion, and a semicircular circumduction movement of the affected lower limb during the swing phase. It is caused by unilateral or bilateral corticospinal tract (upper motor neuron) damage, such as in cerebral palsy, stroke, or multiple sclerosis.

### **Spasticity**

Velocity-dependent increase in muscle tone caused by an upper motor-neuron lesion, producing stiffness, spasms, abnormal reflexes, and restricted movement.

### **Spatial Resolution**

The ability of an imaging system to show two nearby small objects as separate. Better resolution shows finer detail, but the result depends on the scanner, movement, body part, scan settings, and the type of imaging used.

### **Speaking with a Tracheostomy Tube**

Speech with a tracheostomy tube may be supported by cuff deflation, a speaking valve, or other methods when airflow and airway protection are adequate; specialist assessment is required.

### **Special Care Baby Unit**

A ward within a hospital that provides specialist care for newborn babies who need additional support but not intensive care.

### **Special Stain**

A histochemical stain that highlights a particular tissue component, organism, or deposited substance not adequately demonstrated by routine hematoxylin and eosin. Examples include periodic acid-Schiff, Ziehl-Neelsen, Congo red, and Gomori methenamine silver stains.

### **Specific Absorbed Fraction**

The portion of energy released by radiation in one body area that is absorbed by a particular organ or tissue. It helps calculate the radiation dose during nuclear-medicine treatment or testing.



---

### Specific Activity

The amount of radioactivity in a given amount of material. It matters when preparing radioactive medicines because enough activity is needed for imaging or treatment while limiting the amount of substance and unnecessary exposure.

### Specific Gravity

A comparison of urine density with water that gives an estimate of how concentrated the urine is. It may rise with dehydration, sugar, or protein in urine and fall when the kidneys cannot concentrate urine or when excess fluid is present.

### Specificity

A test's ability to correctly identify people who do not have the condition being tested.

### Specific Learning Disorder

A developmental condition causing lasting difficulty learning or using reading, writing, or mathematics skills despite appropriate teaching. The difficulty interferes with school, work, or daily activities and is not explained only by poor instruction, vision, hearing, or intellectual disability.

### Specific Phobia

A disorder involving marked, persistent fear or anxiety about a particular object or situation. The fear is excessive relative to the actual danger, leads to avoidance or intense distress, and interferes with daily functioning.

### Specific Reading Disability

Alternate name; see *Dyslexia*.

### Specimen Adequacy

The degree to which a collected clinical specimen is sufficient in quantity, correctly identified, properly transported, and representative of the site being tested. Inadequacy can cause invalid, misleading, or falsely negative results.

### Specimen Container

A clean, suitable container used to collect, identify,

protect, and transport a patient sample for laboratory testing. The container depends on the sample and test; it should be labeled correctly, closed securely, and kept free from contamination.

### Specimen Transport Medium

A special liquid or gel that keeps germs or cells suitable for testing while a patient sample travels to the laboratory. The medium, temperature, container, and delivery time depend on the sample and suspected infection.

### Speckled Lentiginous Nevus (Nevus Spilus)

A flat tan or brown patch containing multiple darker macules or papules. It is usually benign, but changing, enlarging, or atypical components should be examined because melanoma can rarely arise within a melanocytic nevus.

### Spectral Analysis

Examination of the different energy levels within radiation or another signal. In medical imaging, it can separate useful information from background scatter and help identify materials or improve image interpretation.

### SPECT Scan

A nuclear-medicine scan that uses a small radioactive tracer and a rotating camera to create three-dimensional pictures of blood flow or organ function. It can examine the heart, brain, bones, kidneys, or other organs.

#### Alternate Names:

SPECT Imaging; Single-Photon Emission Tomography

### Speculum

An instrument used to gently separate or hold open the walls of a body passage so that the area can be visualized, examined, sampled, or treated. Common types include vaginal, nasal, ear, and rectal specula; design and size depend on the body site and intended procedure.

---

## Speech

The production of spoken language using breathing, voice, and movements of the tongue, lips, palate, and other structures. Hearing, brain function, teeth, illness, and dental appliances can affect speech.

## Speech And Language Delay

Slower-than-expected development of speech sounds, vocabulary, understanding, or use of language in a child. Causes may include hearing loss, developmental disorders, autism, neurologic disease, or other medical and environmental factors.

## Speech Audiometry

Hearing testing that measures detection or recognition of speech at different loudness levels. It commonly includes speech reception threshold and word-recognition testing, complementing pure-tone audiometry.

### Alternate Names:

Speech Discrimination Testing; Speech Audiometric Evaluation

## Speech Delay

Speech or language development that is slower than expected for a child's age. Causes may include hearing loss, developmental differences, autism, neurologic disease, or limited language access; loss of previously learned skills needs prompt assessment.

## Speech Discrimination

A hearing test that measures how well a person understands spoken words at a comfortable loudness. The result helps assess hearing function beyond the ability to detect pure tones.

## Speech Disorders

Conditions that affect making speech sounds, speaking smoothly, using the voice, or controlling the movements needed for speech. They may follow hearing loss, developmental differences, brain injury, stroke, nerve disease, or other conditions and can often be assessed by a speech therapist.

## Speech Impairment In Adults

Difficulty producing or understanding speech caused by aphasia, dysarthria, apraxia, hearing loss, neurologic disease, structural problems, or cognitive disorder; sudden onset requires stroke assessment.

## Speech-Language Pathologist

A clinician who evaluates and treats disorders of speech, language, voice, communication, cognition, and swallowing across childhood, adulthood, and older age.

## Speech-Language Pathology Assessment

An evaluation of speech, language, voice, communication, swallowing, and sometimes related cognitive skills. It combines interview, clinical observation, and standardized or functional tests to guide therapy.

## Speech Therapy

Assessment and treatment to help with speaking, understanding language, voice, communication, or swallowing. A speech and language therapist may use exercises, practice, communication aids, and advice for the person and family.

## Sperm

Mature male gametes produced in the testes. Each normally carries 23 chromosomes and has a head containing genetic material, a midpiece that supplies energy, and a tail that helps it move; semen contains sperm plus fluids from reproductive glands.

## Spermarche

The onset of sperm production and the first ejaculation, marking an important stage of male puberty.

## Spermatic

Relating to sperm or structures that produce, transport, or support sperm. Spermatic structures include the testes, ducts, blood vessels, nerves, and tissues involved in male reproductive function.

## Spermatic Cord

A cord-like structure carrying the vas deferens, tes-

---

ticular blood vessels, nerves, lymph vessels, and supporting tissue. It extends through the groin to the testis and may be affected by hernia, twisting, or infection.

### **Spermatocele**

A usually harmless, fluid-filled cyst beside or within the epididymis, the coiled tube behind the testicle. It often causes no symptoms but may feel like a lump and should be checked if newly noticed or painful.

### **Spermatogenesis**

The production and maturation of sperm in the seminiferous tubules of the testes. It depends on germ cells, Sertoli cells, hormones, and suitable testicular temperature.

### **Spermatogonium**

An early male reproductive cell in the testis that divides and develops into primary spermatocytes, beginning sperm production.

### **Spermatorrhea**

Involuntary leakage or discharge of semen, sometimes occurring during sleep or without orgasm. Occasional nocturnal emissions can be normal; persistent discharge, pain, blood, or urinary symptoms needs assessment.

### **Spermatozoa**

Mature male reproductive cells, also called sperm. Each normally carries half the usual chromosome number and has a head, middle section, and tail that helps it move toward an egg.

### **Spermatozoon**

A single mature male reproductive cell, or sperm cell. The singular form of spermatozoa: one mature male reproductive cell capable of carrying genetic material to an egg.

### **Sperm Cell**

Alternate name; see *Sperm*.

### **Spermicide**

A contraceptive substance that kills or immobilizes sperm, often used with condoms or other barrier methods.

### **Sperm Retrieval**

A procedure that obtains sperm from semen, the epididymis, or testicular tissue for fertility treatment or laboratory testing. The method depends on whether sperm are present in the semen and on the cause of infertility.

#### **Alternate Names:**

Sperm Extraction; TESE; PESA; Surgical Sperm Recovery

### **S Phase**

The stage of the cell cycle when a cell copies all of its DNA before dividing. This duplication allows each new cell to receive a complete set of genetic instructions.

### **Sphenoid**

Relating to the sphenoid bone, a central bone at the base of the skull. It helps form the eye sockets and skull base and contains the hollow spaces called the sphenoid sinuses.

### **Sphenoidal Sinus**

A paired air-filled sinus within the sphenoid bone at the base of the skull. It lies near important nerves and blood vessels, so inflammation or surgery in this area requires careful assessment.

### **Sphenoid Bone**

A central bone at the base of the skull that helps form the eye sockets and skull base. It contains the sphenoid sinuses and a hollow that holds the pituitary gland.

### **Sphenoid Sinusitis**

Inflammation or infection of the sphenoid sinus, which lies deep behind the nasal cavity. It may cause deep headache, facial pain, or visual symptoms and can rarely spread to nearby nerves or the brain; persistent or severe symptoms require

---

medical assessment.

### Spherocytes

Red blood cells that are unusually round and dense instead of having their normal flattened shape. They may occur when the cell membrane is damaged or removed, especially in hereditary spherocytosis or immune red-cell destruction.

### Sphincter

A ring-shaped muscle that opens and closes a body passage. Sphincters help control urine, stool, food, bile, and other fluids; examples include the muscles around the anus, bladder outlet, and stomach outlet.

### Sphincter Of Oddi

A muscular valve controlling the release of bile and pancreatic juice into the first part of the small intestine. It opens during digestion and abnormal blockage or tightening can cause pain or pancreatitis.

### Sphygmograph

An instrument or sensor that records the shape and timing of the pulse in an artery. The tracing can provide information about circulation and arterial stiffness.

### Sphygmomanometer

A device used to measure arterial blood pressure with an inflatable cuff and pressure-measuring system. Manual versions commonly use an aneroid or mercury manometer with a stethoscope, while electronic versions use sensors to analyze pressure oscillations and display the reading.

#### Alternate Names:

Blood Pressure Gauge

### Spider Angioma

Alternate name; see *Spider Naevi*.

### Spider Bite

An injury caused by a spider, usually producing a small painful or itchy area. Some species can cause worsening tissue damage, severe muscle cramps,

sweating, vomiting, breathing problems, or other serious effects; rapidly worsening symptoms need emergency care.

### Spider Bites

Alternate name; see *Black widow and brown recluse bites*.

### Spider Naevi

Vascular cutaneous lesions characterized by a central, pulsating red arteriole from which fine, radiating telangiectatic capillaries branch outward like spider legs. Digital pressure blanches the lesion, and upon release it refills rapidly from the center outwards; multiple spider naevi in the superior vena cava distribution (face, neck, upper chest) are classic physical signs of hyperestrogenic states in chronic liver disease, cirrhosis, and pregnancy.

#### Alternate Names:

Spider Angioma; Spider Naevus; Spider Telangiectasia; Nevus Araneus; Vascular Spider

### Spider Naevus

Alternate name; see *Spider Naevi*.

### Spider Telangiectasia

Alternate name; see *Spider Naevi*.

### Spigelian Hernia

A ventral abdominal wall hernia protruding through a defect in the Spigelian aponeurosis along the semilunar line (between the lateral border of the rectus abdominis muscle and the transversus abdominis aponeurosis), most frequently located below the arcuate line.

#### Alternate Names:

Lateral Ventral Hernia; Spigelian Aponeurosis Hernia

### Spina Bifida

A birth defect in which the neural tube and the bones protecting the spinal cord do not close completely during early development. It can range from mild to severe and may cause weakness, loss of feeling, or problems controlling the bladder or bowel.

---

## Spina Bifida Meningocele

A form of spina bifida in which the protective coverings around the spinal cord bulge through an opening in the backbone and form a fluid-filled sac. The spinal cord is usually not inside the sac, so weakness or loss of bladder and bowel control may be less severe.

## Spina Bifida Myelomeningocele

The most severe common form of spina bifida, in which the spinal cord and its coverings bulge through an opening in the backbone. It can cause weakness or paralysis, loss of feeling, and difficulty controlling the bladder or bowel below the affected area.

## Spina Bifida Occulta

A usually mild form of spina bifida in which the vertebral arch does not fully close without herniation of the spinal cord or meninges. It is often asymptomatic but may be associated with a skin dimple, hair tuft, or other finding over the lower spine.

## Spinal Analgesia

Pain relief produced by injecting numbing medicine, sometimes with an opioid, into the fluid around the lower spinal nerves. It may cause temporary leg weakness, itching, difficulty urinating, low blood pressure, headache, or slowed breathing.

## Spinal And Bulbar Muscular Atrophy

An X-linked disorder caused by a CAG-repeat expansion in the androgen-receptor gene. It usually affects adult men and gradually damages motor neurons, causing muscle weakness and wasting, cramps, twitching, difficulty speaking or swallowing, and sometimes breast enlargement or reduced fertility. Treatment supports movement, swallowing, breathing, and daily activities.

### Alternate Names:

X-Linked Spinal and Bulbar Muscular Atrophy; Kennedy Disease; Kennedy's Disease; SBMA

### Alternate Names:

Kennedy Disease; SBMA

## Spinal And Epidural Anesthesia

Spinal and epidural anesthesia deliver local anesthetic near spinal nerves to numb the lower body; spinal is a single injection, while epidural uses a catheter for adjustable dosing.

## Spinal Anesthesia

An injection of numbing medicine into the fluid around the lower spinal nerves to temporarily block pain and movement in the lower body. Possible effects include low blood pressure, difficulty urinating, headache, infection, bleeding, or rarely excessive numbness and breathing problems.

### Alternate Names:

Subarachnoid Block; Spinal Block

## Spinal Anesthesia For Cesarean Delivery

A regional anesthetic injected into the fluid around the lower spinal nerves to numb the lower body during a cesarean birth. The mother remains awake, while blood pressure, breathing, and the baby's condition are monitored.

## Spinal Angiography

X-ray imaging of blood vessels supplying the spinal cord after contrast is injected. It is used to locate vascular malformations, fistulas, aneurysms, or sources of bleeding and may guide treatment.

## Spinal Artery

An artery supplying the spinal cord and its coverings, including the anterior spinal artery and paired posterior spinal arteries; blockage can cause characteristic neurologic deficits.

## Spinal Block

A spinal block is injection of local anesthetic into cerebrospinal fluid in the lower back, producing temporary numbness and weakness for surgery or pain control.

## Spinal Canal

The spinal canal is the bony passage through the vertebral column containing the spinal cord, its coverings, nerve roots, and blood vessels.

---

## Spinal Cavity

The posterior body cavity formed by the vertebral column that contains and protects the spinal cord.

## Spinal Column

The vertebral column, consisting of vertebrae, discs, ligaments, and joints that support and protect the spinal cord.

## Spinal Cord

The cylindrical nervous tissue within the vertebral canal that carries signals between brain and body and mediates reflexes.

## Spinal Cord Abscess

A spinal cord or epidural abscess is a collection of infection near the spinal cord that can compress neural tissue and cause pain, fever, weakness, paralysis, or bladder problems.

## Spinal Cord and Spine Disorders

Diseases or injuries affecting the spinal bones, discs, ligaments, spinal cord, or nerve roots. They may cause pain, weakness, numbness, balance problems, deformity, or changes in bladder and bowel function.

## Spinal Cord Compression

Pressure on the spinal cord or its nerve roots, often from a tumor, infection, bleeding, fracture, or slipped disc. It can cause severe back pain, weakness, numbness, walking difficulty, or loss of bladder or bowel control.

## Spinal Cord Disease

A disorder affecting the spinal cord, including compression, inflammation, vascular injury, infection, tumor, degeneration, or trauma, and potentially causing weakness, sensory loss, or autonomic dysfunction.

## Spinal Cord Infarction

Injury to the spinal cord caused by interruption of its blood supply, often involving the anterior spinal artery or aortic circulation. It typically produces sudden back or limb pain followed by weakness,

sensory loss, and bladder or bowel dysfunction.

## Spinal Cord Injuries

Traumatic damage to the spinal cord that can cause temporary or permanent motor, sensory, autonomic, respiratory, or bowel and bladder dysfunction. Severity depends on the level and completeness of injury.

## Spinal Cord Injury

Damage to the spinal cord that can cause weakness, paralysis, altered sensation, or loss of bladder and bowel control. Severity is often graded from complete to incomplete using a standardized neurological examination.

### Alternate Names:

SCI

## Spinal Cord Injury: Treatments And Rehabilitation

Spinal-cord injury can cause temporary or permanent weakness, sensory loss, autonomic dysfunction, and bowel or bladder problems. Acute care protects the spine and treats complications; rehabilitation addresses mobility, skin, breathing, function, equipment, sexuality, and community participation.

## Spinal Cord Neoplasms

Tumors arising in the spinal cord, its coverings, or nearby tissues. They may be noncancerous or cancerous and can cause pain, weakness, numbness, walking problems, or bladder and bowel changes by pressing on nerves.

## Spinal Cord Stimulation

An implanted treatment that delivers electrical impulses to the epidural space to alter pain signaling, used for selected chronic neuropathic or ischemic pain after careful assessment.

## Spinal Cord Trauma

Spinal cord trauma is injury to the cord from force such as fracture, dislocation, or penetrating damage, potentially causing weakness, sensory loss, autonomic dysfunction, or paralysis.



---

## Spinal Curvature

Alternate name; see *Scoliosis*.

## Spinal Disease

Any disorder affecting the vertebrae, discs, joints, ligaments, spinal cord, or nerve roots. It may cause back or neck pain, stiffness, weakness, numbness, walking difficulty, or changes in bladder or bowel control.

## Spinal Dysraphism

A group of birth defects caused by incomplete development or closure of the spine and spinal cord before birth. Some forms have an open sac on the back, while others are hidden and may later cause weakness, pain, or bladder and bowel problems.

## Spinal Epidural Abscess

A collection of pus between the covering of the spinal cord and the surrounding bone, usually caused by infection. Severe back pain, fever, weakness, numbness, difficulty walking, or loss of bladder or bowel control may occur.

## Spinal Fluid Leak

Alternate name; see *CSF leak* (*Cerebrospinal fluid leak*).

## Spinal Fracture

A fracture of one or more vertebrae caused by trauma, osteoporosis, tumor, or other bone-weakening disease.

## Spinal Fusion

Surgery that joins two or more spinal bones so they heal into one stable segment. Bone and often screws, rods, or cages are used; the operation may reduce movement and pain from selected conditions but can cause infection, nerve injury, or problems at nearby levels.

## Spinal Ganglion

A small group of nerve-cell bodies beside the spinal cord. It contains sensory nerve cells that carry information about touch, pain, temperature, and

body position from the skin, muscles, and joints.

## Spinal Injury

Damage to the spinal cord, vertebrae, discs, ligaments, or nearby nerves caused by trauma or another disease. It may cause pain, weakness, numbness, paralysis, or bladder and bowel problems and can be life-threatening when severe.

## Spinal Metastasis

Spread of a cancer to the vertebrae, spinal canal, or nearby epidural tissues. Back pain, weakness, numbness, gait change, or bowel or bladder dysfunction may signal spinal cord or nerve compression and requires urgent assessment.

## Spinal Muscular Atrophy

Usually an autosomal recessive inherited disorder causing progressive loss of motor neurons and muscle weakness, often beginning in infancy or childhood but sometimes appearing in adulthood. Severity varies; breathing, swallowing, and mobility may be affected. Early diagnosis allows genetic counseling and disease-modifying care.

## Spinal Neoplasms

Tumors arising in the spinal cord, its coverings, vertebrae, or nearby tissues. They may be non-cancerous or cancerous and can cause back pain, weakness, numbness, walking difficulty, or bladder and bowel changes when they press on nerves.

## Spinal Nerve

One of 31 pairs of nerves leaving the spinal cord through openings between the spinal bones. Each carries messages for sensation, movement, and automatic functions between the body and spinal cord.

## Spinal Precautions

Measures used after possible spinal injury to limit movement until the spine and spinal cord have been assessed. They may include careful positioning, support of the neck, and safe transfer; the exact approach depends on the injury and examination.

---

## Spinal Reflex

A fast, automatic response that is processed through the spinal cord. It can occur before the brain becomes aware of the stimulus.

## Spinal Shock

A temporary loss of reflexes, muscle tone, and automatic functions below a spinal-cord injury. The affected limbs may initially be limp, and blood-pressure, bladder, bowel, or sexual problems can occur while the injury is assessed and treated.

## Spinal Stenosis

Narrowing of the passage containing the spinal cord or nerve roots. It may cause back pain, numbness, weakness, or leg pain with walking that improves when sitting or leaning forward.

### Alternate Names:

Vertebral Canal Stenosis; Lumbar Spinal Narrowing

## Spinal Stenosis Lumbar

Alternate name; see *Lumbar stenosis*.

## Spinal Tap

Alternate name; see *Lumbar puncture*.

## Spinal Tumor

An abnormal growth in the spinal bones, spinal cord, its coverings, or nearby tissues. It may be noncancerous or cancerous; symptoms depend on its location and may include pain, weakness, numbness, or loss of bladder control.

## Spinal Tumors

Alternate name; see *Spinal neoplasms*.

## Spindle-Cell Sarcoma

A group of cancers made of long, narrow cells under the microscope. They can arise from muscle, fibrous tissue, or nerve coverings; their behavior and treatment vary by the exact tumor type and location.

## Spindle Pole

One end of the temporary structure formed during

cell division. Fibers extending from each pole attach to duplicated chromosomes and pull the copies into the two new cells.

## Spindle Pole Body

A structure in fungal cells that organizes the fibers used to separate duplicated chromosomes during cell division. It performs a role similar to the centrosome in animal cells.

## Spine

The vertebral column, consisting of stacked vertebrae that support the body and protect the spinal cord.

## Spine Disorders

Conditions affecting the bones, disks, joints, ligaments, nerves, or spinal cord. They may cause back or neck pain, stiffness, weakness, numbness, balance problems, or changes in bladder and bowel control.

## Spine Injury

Damage to the bones, discs, ligaments, nerves, or spinal cord. It may cause back pain, weakness, numbness, loss of movement, or bladder and bowel problems and can require emergency care.

## Spinocerebellar Ataxia

Alternate name; see *Spinocerebellar Ataxias*.

## Spinocerebellar Ataxias

A group of rare inherited disorders that progressively damage the cerebellum and sometimes other parts of the nervous system. The cerebellum helps control balance, coordination, and precise movement. These disorders may cause unsteady walking, poor hand coordination, slurred or unclear speech, shaking, abnormal eye movements, muscle stiffness or weakness, and difficulty swallowing. The age of onset, symptoms, and rate of progression vary widely. Many types are inherited in an autosomal dominant pattern, but the inheritance pattern differs among types.

---

## **Spinothalamic Tract**

The primary ascending somatic sensory pathway conveying pain, thermal sensation, and crude non-discriminative touch from peripheral free nerve endings through the spinal cord (decussating within 1 to 2 segments in the anterior white commissure) to the ventral posterolateral (VPL) nucleus of the thalamus.

### **Alternate Names:**

Anterolateral System; ALS

## **Spinous Process**

The backward-pointing bony projection of a vertebra that can be felt along the middle of the back. Muscles and ligaments attach to it and help support and move the spine.

## **Spiradenocarcinoma**

A rare malignant sweat-gland tumor that may arise in a long-standing spiradenoma, presenting as a rapidly enlarging, painful, or ulcerated nodule.

## **Spiradenoma**

A usually benign painful sweat-gland tumor presenting as a dermal nodule; multiple spiradenomas may occur in Brooke-Spiegler syndrome.

## **Spiral Organ**

The organ of Corti, a sensory structure in the cochlea containing hair cells that convert sound vibrations into nerve signals.

## **Spiramycin**

A macrolide antibiotic used for selected bacterial infections and sometimes during pregnancy for toxoplasmosis. Choice depends on the infection, local guidance, pregnancy stage, allergies, resistance, and other treatment needs.

## **Spirillum Minus**

*Spirillum minus* is a spiral Gram-negative bacterium that causes rat-bite fever, characterized by fever, rash, and migratory joint pain after rodent exposure; culture is difficult and diagnosis may be clinical or molecular.

## **Spirit**

A volatile liquid preparation or alcohol; the meaning depends on the medical, pharmaceutical, or general context.

## **Spiritual Pain**

Distress involving meaning, purpose, beliefs, identity, or connection, often addressed in palliative and supportive care.

## **Spirochaetaceae**

A family of spiral-shaped bacteria that includes medically important spirochetes such as *Borrelia*, *Treponema*, and *Leptospira*.

## **Spirochaetales Infection**

Infection caused by spiral-shaped bacteria in the order Spirochaetales, including syphilis, Lyme disease, relapsing fever, and leptospirosis.

### **Alternate Names:**

Spirochaetal Infections; Spirochetal Infections

## **Spirochaete**

Another spelling of spirochete, a thin, flexible, spiral-shaped bacterium. Important genera include *Treponema*, *Borrelia*, and *Leptospira*.

## **Spirochete**

A thin, spiral-shaped bacterium that moves with a corkscrew-like motion. Important examples cause syphilis, Lyme disease, relapsing fever, and leptospirosis; diagnosis depends on the organism and may use blood tests or other samples.

## **Spirogram**

A graph recording how much air a person breathes and how quickly it moves during a lung-function test. It helps identify patterns suggesting airway narrowing, restricted lung expansion, or poor effort.

## **Spirometer**

An instrument that measures airflow and the volume of air moved during specified breathing maneuvers, especially forced expiration. It does not by itself measure all lung volumes; residual volume and total

---

lung capacity require additional pulmonary-function testing.

### **Spirometry**

A breathing test that measures how much air a person can blow out and how quickly. It helps identify narrowing of the airways, as in asthma or chronic obstructive pulmonary disease, and can monitor treatment. The person must follow instructions and blow forcefully several times; results are interpreted with age, height, and symptoms.

### **Spironolactone**

An aldosterone antagonist and potassium-sparing diuretic used for heart failure, hypertension, hyperaldosteronism, and other indications.

### **Spirulina**

Spirulina is a type of cyanobacterium sold as a dietary supplement. Products may provide nutrients, but contamination and medication interactions are possible, and it is not a reliable substitute for a balanced diet.

### **Spit**

Saliva or other fluid expelled from the mouth. Increased saliva or repeated spitting may occur with mouth irritation, nausea, reflux, difficulty swallowing, medication effects, or habit.

### **Spitting**

Forceful expulsion of saliva or other material from the mouth. New or excessive spitting may occur with oral disease, reflux, nausea, difficulty swallowing, medication effects, or habit.

### **Spitz Nevus**

A benign melanocytic nevus, often a pink or pigmented papule in children or young adults, that may resemble melanoma clinically or histologically.

### **Spitz Nevus Pathology**

A usually benign melanocytic proliferation composed of epithelioid or spindle cells, often symmetrical and well circumscribed, with maturation,

Kamino bodies, and characteristic epidermal changes. Some lesions overlap morphologically with melanoma.

### **Splanchnic**

Relating to internal organs, especially abdominal organs and their blood vessels or nerves. Splanchnic circulation supplies the digestive organs, while splanchnic nerves carry involuntary signals that regulate their function.

### **Splanchnic Nerve**

A nerve carrying automatic signals between the spinal cord and internal organs, especially organs in the chest and abdomen. It helps regulate digestion, blood-vessel tone, and other functions that do not require conscious control.

### **Spleen**

A lymphoid organ in the upper-left abdomen. Its red pulp filters blood and removes aged or abnormal blood cells; its white pulp supports immune responses. It also stores platelets and helps clear blood-borne microorganisms.

### **Spleen Development**

The formation and maturation of the spleen before and after birth. The spleen gradually develops its blood-filtering and immune functions, including removal of aged blood cells and responses to germs in the bloodstream.

### **Spleen Diseases**

Disorders affecting splenic structure or function, including splenomegaly, infarction, infection, cysts, abscess, rupture, vascular disease, and hematologic or infiltrative disorders.

### **Spleen Disorder**

A disease affecting the spleen, such as enlargement, infection, infarction, rupture, inflammation, cyst, or tumor, with possible effects on blood cells and immunity.

---

## Spleen Hemangioma

A spleen hemangioma is a usually benign blood-vessel tumor of the spleen, often incidental but occasionally associated with enlargement, anemia, or rupture risk.

## Spleen Hilum

The indented area on the inner surface of the spleen where its blood vessels, nerves, and lymph vessels enter or leave. The splenic vein carries blood from the spleen toward the liver through the portal circulation.

## Spleen Removal

Surgery to remove the spleen, which may be needed for injury, an enlarged or overactive spleen, blood disease, or a tumor. Without a spleen, the risk of severe infection increases, so vaccination and medical advice about fever are important.

## Splenectomy

Surgical removal of the spleen, performed for selected traumatic, hematologic, malignant, vascular, or other disorders. It increases the risk of severe infection with encapsulated bacteria and requires preventive planning.

## Splenic Abscess

A localized collection of pus in the spleen, usually from bloodstream infection, trauma, infarction, or spread from nearby disease and potentially causing fever, left-upper-quadrant pain, and sepsis.

## Splenic Artery Aneurysm

An abnormal bulge in the artery that supplies the spleen. It often causes no symptoms, but rupture can cause life-threatening internal bleeding; risk is of particular concern during pregnancy.

## Splenic Cyst

A fluid-filled lesion within the spleen, classified as true cysts with an epithelial lining, including congenital, epidermoid, and dermoid cysts, or pseudocysts without a lining, typically resulting from trauma, infection, or pancreatitis. Most are asymptomatic and discovered incidentally on imaging.

## Splenic Flexure

The bend of the colon where the transverse colon becomes the descending colon, near the spleen.

## Splenic Flexure Syndrome

Pain, fullness, or bloating in the upper-left abdomen caused by gas collecting at the bend where the transverse colon meets the descending colon. Symptoms may improve after passing gas or stool.

## Splenic Infarction

Death of part of the spleen because its blood supply is blocked. It may cause sudden severe pain in the upper-left abdomen, sometimes spreading to the left shoulder, with nausea or fever. Causes include blood clots, heart or blood disorders, infections, inflammation of blood vessels, or injury. CT or ultrasound is commonly used for diagnosis, and complications may include an abscess or rupture.

## Splenic Infection

Infection of the spleen, often an abscess caused by bacteria spreading through the bloodstream or from nearby tissue. It may cause fever, chills, left-upper-abdominal pain, and severe illness and usually needs urgent medical treatment.

## Splenic Neoplasm

An abnormal growth in the spleen. It may be non-cancerous, cancerous, or cancer that has spread from another organ; symptoms and treatment depend on its type, size, and effect on nearby tissues or blood cells.

## Splenic Rupture

A tear of the spleen, usually after blunt abdominal trauma but sometimes spontaneous in an enlarged or diseased spleen. It can cause life-threatening internal bleeding; diagnosis and treatment depend on hemodynamic stability and imaging, with urgent surgery or embolization sometimes required.

## Splenomegaly

Abnormal enlargement of the spleen. It may result from infection, liver disease, blood disorders, cancer, or other conditions and may cause no symptoms.

---

Some people feel fullness or pain in the upper-left abdomen. An enlarged spleen can trap blood cells and is more easily injured or ruptured.

### **Spliceosome**

The spliceosome is a nuclear RNA-protein complex that removes introns from pre-messenger RNA and joins exons to form mature RNA.

### **Splice-Site Variant**

A variant affecting the sequences required for accurate removal of introns and joining of exons during RNA processing. It may cause abnormal transcripts, exon skipping, intron retention, or reduced functional protein.

### **Splint**

A device that supports and limits movement of an injured or painful body part. Splints may be temporary or definitive and should not be so tight that they impair circulation or sensation.

### **Splinter**

A splinter is a small piece of wood, glass, metal, or another material embedded in the skin. It should be removed carefully; medical care may be needed if it is deep, infected, or near an eye or joint.

### **Splinter Haemorrhages**

Small, narrow lines of bleeding beneath a nail. They often follow minor injury but can also occur with heart-valve infection, blood-vessel inflammation, psoriasis, or other illnesses and need context for interpretation.

### **Splinter Hemorrhage**

A splinter hemorrhage is a thin line of bleeding under a fingernail or toenail. It may follow injury, but multiple or unexplained lesions can occur with medical conditions and deserve evaluation.

### **Splinter Hemorrhages**

Thin, lengthwise lines of bleeding beneath a nail. Minor injury is common, but they can also occur with heart-valve infection, blood-vessel inflammation,

psoriasis, or other illnesses; their meaning depends on the overall clinical picture.

### **Splinter Removal**

Taking a piece of wood, metal, glass, or another material out of the skin. Small superficial splinters may be removed with clean tweezers, while deep, painful, infected, or eye- or nail-area splinters need clinical care.

### **Splints**

Devices that support, immobilize, align, or protect an injured body part, reducing pain and allowing healing.

### **Split Brain**

A state after surgical or pathological disruption of the corpus callosum in which information transfer between cerebral hemispheres is reduced, producing characteristic disconnection findings.

### **Split-Thickness Skin Graft**

A graft containing the epidermis and part of the dermis, used to cover wounds or burns and allowing the donor site to heal from remaining skin structures.

### **Split-Thickness Skin Grafts**

Grafts containing the epidermis and part of the dermis, used to cover burns, wounds, ulcers, or surgical defects. They take more readily than full-thickness grafts but may contract, differ in color or texture, and leave a donor-site wound.

### **Splitting**

A psychological defense pattern in which a person or situation is viewed as entirely good or entirely bad, with difficulty integrating positive and negative qualities. It can occur in several mental-health conditions and is not diagnostic by itself.

### **Spondees**

A two-syllable word with equal emphasis on both parts, used during speech-hearing tests. Examples help measure the quietest loudness at which a person can recognize spoken words accurately.



---

## Spondylarthritis

A family of inflammatory rheumatic diseases affecting the spine, sacroiliac joints, entheses, peripheral joints, skin, eyes, or bowel, including ankylosing spondylitis and psoriatic arthritis.

## Spondylitis

Inflammation of one or more vertebrae or parts of the spine. Causes include infection, autoimmune disease, and inflammatory arthritis; pain, stiffness, fever, or neurologic symptoms may occur.

## Spondyloarthritis

A group of inflammatory joint diseases that often affect the spine and joints between the spine and pelvis. They may also cause tendon pain, swollen fingers or toes, eye inflammation, psoriasis, or bowel inflammation.

## Spondylolisthesis

Slipping of one spinal bone forward or backward compared with the bone below it. It may cause back pain, stiffness, leg pain, numbness, or weakness, although some people have no symptoms.

### Alternate Names:

Lumbosacral Spondylolisthesis

## Spondylolysis

A small stress crack or defect in part of a spinal bone, usually in the lower back. It may cause back pain in adolescents or athletes and can sometimes allow one spinal bone to slip on another.

### Alternate Names:

Lumbosacral Spondylolysis

## Spondylosis

Wear-related changes in the bones, discs, and small joints of the spine. It becomes more common with age and may cause stiffness or pain, but many people have no symptoms; nerve pressure can cause numbness or weakness.

## Sponge

A sponge is a porous material used to absorb fluid, clean or dress wounds, or deliver a product; retained

surgical sponges are a preventable complication.

## Spongiform Encephalopathy

A serious progressive brain disease in which brain tissue develops tiny sponge-like spaces. Prion diseases are important examples and can cause worsening memory, movement, coordination, behavior, and eventually loss of independence.

## Spongy Bone

Porous trabecular bone found inside many bones, especially near joints.

### Alternate Names:

Cancellous Bone; Trabecular Bone

## Spontaneous Abortion

Unintentional loss of a pregnancy before fetal viability. Causes include chromosomal abnormalities, uterine or cervical problems, infection, and maternal medical conditions.

### Alternate Names:

Early Pregnancy Loss; Spontaneous Pregnancy Loss

## Spontaneous Bacterial Peritonitis

An infection of ascitic fluid without an identifiable abdominal source, usually occurring in people with cirrhosis and ascites. It may cause fever, abdominal pain, tenderness, or worsening illness. Diagnosis is made by testing ascitic fluid, usually showing at least 250 neutrophils per mm<sup>3</sup>.

## Spontaneous Bacterial Peritonitis Diagnostic Criteria

The diagnostic consensus criteria for spontaneous bacterial peritonitis (SBP), defined as an acute bacterial infection of ascitic fluid in patients with advanced cirrhosis in the absence of a surgically treatable intra-abdominal source. Established on diagnostic paracentesis when the ascitic fluid absolute neutrophil count (ANC) equals or exceeds 250 cells per cubic millimeter ( $0.25 \times 10^9/L$ ).

### Alternate Names:

SBP Diagnostic Criteria; Ascitic Neutrophil Cutoff

---

### Spontaneous Breathing Trial

A monitored period during which a person on a breathing machine does most of the breathing with little or no machine assistance. It helps clinicians judge whether the person may breathe safely after the tube is removed.

### Spontaneous Coronary Artery Dissection

A tear within the wall of an artery supplying the heart, which can narrow or block blood flow and cause a heart attack. It often affects women and may be associated with pregnancy, severe stress, or fibromuscular dysplasia. Treatment depends on blood flow and stability; medicines or a stent may be used, and emergency care is needed for chest pain.

**Alternate Names:**

SCAD

### Spontaneous Miscarriage

Pregnancy loss occurring without an intentional medical or surgical procedure.

### Spontaneous Perforation

Unexpected rupture of a hollow organ without a clear external injury, allowing contents to escape and potentially causing peritonitis, mediastinitis, sepsis, or hemorrhage.

### Spontaneous Pneumothorax

Air leaking into the space around a lung without a direct injury. It may occur in an otherwise healthy person or because of lung disease and can cause sudden chest pain or breathlessness.

### Spontaneous Vaginal Delivery

Birth through the vagina without forceps, a vacuum device, or cesarean surgery. It usually occurs during the second stage of labor as contractions and pushing move the baby through the birth canal.

**Alternate Names:**

Spontaneous Vaginal Delivery; SVD; Normal Vaginal Delivery

### Sporadic

Occurring irregularly or as an isolated case rather than in a predictable pattern or family cluster.

### Sporothrix

A genus of environmental fungi that can cause sporotrichosis, usually after inoculation through the skin and sometimes through inhalation or dissemination.

### Sporothrix Schenckii

A group of fungi that causes sporotrichosis, a typically nodular skin and lymphatic infection acquired from plants, soil, or animal scratches.

### Sporotrichosis

Sporotrichosis is a fungal infection caused by *Sporothrix* species, often introduced through plant matter and producing nodules along lymphatic channels.

### Sporozoa

An older group name for spore-forming protozoan parasites, including *Plasmodium*.

### Sporozoites

Moving, infective forms produced by certain parasites. For example, mosquitoes inject *Plasmodium* sporozoites into people, after which they travel to the liver and begin the malaria infection cycle.

### Sports Cream Overdose

Excessive use or swallowing of a muscle or joint pain cream, often containing menthol, methyl salicylate, or capsaicin. It may cause skin burns, salicylate poisoning, vomiting, ringing ears, confusion, or seizures.

### Sportsman's Groin

Long-lasting groin or lower-abdominal pain related to repeated sports activity, especially kicking, sprinting, twisting, or quick changes of direction. It is often called sports hernia, although there may be no true hernia. Examination and sometimes imaging help identify the cause; treatment may include rest,

---

guided exercises, and, in selected persistent cases, surgery.

**Alternate Names:**

Athletic Pubalgia; Sports Hernia; Gilmore's Groin; Inguinal Disruption; Core Muscle Injury

**Sports Medicine**

Medical care focused on preventing, diagnosing, and treating exercise- and sports-related injuries and conditions. It also supports safe training, rehabilitation, performance, and return to activity.

**Sports Physical**

A preparticipation health evaluation that reviews medical history, symptoms, medications, family risks, and examination findings to identify conditions needing evaluation before sports.

**Alternate Names:**

Sports Physicals

**Sports Rehabilitation**

Treatment and training that help a person recover safe movement, strength, fitness, and confidence after a sports injury. It may include exercises, hands-on care, education, gradual return to activity, and steps to reduce the chance of another injury.

**Spots Before The Eyes**

Visual floaters, flashes, or scotomata perceived in the visual field; sudden onset requires urgent assessment.

**Spots In Front Of The Eyes**

Floaters or visual spots caused by shadows from vitreous opacities, inflammation, bleeding, retinal tears, migraine, or other eye or neurologic conditions; sudden flashes or a curtain-like loss are urgent.

**Spotted Fever**

A group of infections usually spread by ticks and caused by *Rickettsia* bacteria. They can produce fever, headache, rash, muscle aches, and serious illness requiring prompt diagnosis and antibiotic treatment.

**Spousal Abuse**

Spousal abuse is physical, sexual, emotional, financial, or coercive harm by an intimate partner; immediate safety planning and confidential support may be needed.

**Spouse**

A spouse is a person's legally or socially recognized partner; involving a spouse in healthcare requires the patient's consent unless law provides otherwise.

**Sprain**

An injury in which a ligament that supports a joint is stretched or torn. It can cause pain, swelling, bruising, and looseness; severity ranges from minor stretching to a complete tear and joint instability.

**Sprained Ankle**

Stretching or tearing of one or more ankle ligaments, usually after rolling the ankle inward. It may cause pain, swelling, bruising, and instability.

**Sprains**

Injuries in which ligaments around a joint are stretched or torn. They can cause pain, swelling, bruising, and joint looseness; severity ranges from minor stretching to complete tearing.

**Spray Drying**

Spray drying converts a liquid mixture into a dry powder by atomizing it into a heated gas stream, often used in pharmaceutical and food manufacturing.

**Sprengel Deformity**

A birth difference in which one shoulder blade sits higher than usual because it did not move to its normal position during development. It may cause uneven shoulders, limited arm movement, or neck and back discomfort.

**Spring Fever Henoch-Schonlein**

Alternate name; see *Henoch-Schonlein purpura*.

---

## Sprue

A condition causing poor absorption of nutrients from the small intestine. The term may describe celiac disease or tropical sprue, which have different causes and require appropriate testing and treatment.

## Spur Cell

Alternate name; see *Acanthocyte*.

## Spur-Cell Hemolytic Anemia

A severe anemia in which abnormal, spiky red blood cells are rapidly removed from the blood, usually because of advanced liver disease. It can cause jaundice, weakness, and worsening anemia.

## Spurling Test

A clinical provocative test for cervical radiculopathy; with the patient seated, the neck is extended and laterally rotated toward the affected side, and gentle downward axial pressure is applied to the vertex of the skull; reproduction of sharp radiating dermatomal pain down the arm indicates nerve root compression.

### Alternate Names:

Spurling's Maneuver; Cervical Foraminal Compression Test

## Sputum

Mucus or other material brought up by coughing from the airways or lungs. Its amount, color, smell, and blood content may provide clues, but sputum alone cannot identify the cause; laboratory testing may be needed.

## Sputum Culture

A laboratory test that grows germs from mucus coughed up from the lungs or collected by suction. It can help identify bacterial, fungal, or mycobacterial infection and guide treatment, but a poor sample can give misleading results.

## Sputum Direct Fluorescent Antibody Test

A laboratory test that uses glowing antibody mark-

ers to look for selected germs in mucus from the lungs. It can provide a rapid result, but culture or molecular testing may also be needed.

## Sputum Fungal Smear

Examination of coughed-up lung mucus under a microscope to look for fungi. A culture, blood test, or molecular test may be needed to confirm the specific infection.

## Sputum Gram Stain

A quick microscope test that colors a sputum sample to show some bacteria and inflammatory cells and to judge whether the sample is suitable for culture.

## Sputum Stain For Mycobacteria

A microscope test that looks for tuberculosis-causing and related bacteria in mucus from the lungs. A positive result supports infection, but a negative result does not exclude it; culture or molecular testing may also be needed.

## Squamous Cell

A flat cell that forms part of the covering or lining of the skin and many body passages. These cells protect against friction and infection; changes in them can cause inflammation, precancer, or squamous cell cancer in organs such as the skin, mouth, cervix, lungs, or esophagus.

## Squamous Cell Carcinoma

A cancer of flat cells that line the skin or certain body passages. On the skin it may appear as a rough, scaly, crusted, wart-like, or nonhealing sore and can invade nearby tissue or spread if untreated.

## Squamous Cell Carcinoma in Situ

Squamous cell carcinoma in situ, also called Bowen disease, is an early skin cancer limited to the outer skin layer. It can become invasive if untreated.

## Squamous Cell Carcinoma Of The Lung

A type of non-small-cell lung cancer that usually begins in the lining of a larger airway. It is strongly linked to smoking but can occur in nonsmokers.

---

Symptoms may include a persistent cough, coughing blood, chest pain, breathlessness, or repeated chest infections; treatment depends on stage and tumor features.

### **Squamous Cell Carcinoma Of The Skin**

A skin cancer arising from squamous cells, often appearing as a scaly, crusted, wart-like, or ulcerated growth.

#### **Alternate Names:**

Cutaneous Squamous Cell Carcinoma; cSCC

### **Squamous Cells**

Flat epithelial cells with scale-like appearance that line surfaces such as skin, mouth, cervix, and esophagus; abnormal numbers or forms can reflect irritation, infection, dysplasia, or cancer.

### **Squamous Cell Skin Cancer**

Squamous cell skin cancer is a malignant keratinocyte tumor often arising on sun-exposed skin and capable of local invasion or metastasis, especially when high risk.

### **Square Foot**

A square foot is a unit of area equal to a square measuring one foot on each side, used to describe surfaces, rooms, wounds, or treatment areas.

### **Square Inch**

A square inch is a unit of area equal to a square measuring one inch on each side, used for small surfaces or wound measurements.

### **Squint**

Misalignment of the eyes so they do not point in the same direction. It may be constant or intermittent and can cause double vision, poor depth perception, or reduced vision in one eye.

### **SRT**

Stereotactic radiotherapy, a precise form of external radiation treatment that uses three-dimensional imaging and careful positioning to target a small tumor or other abnormal area. It is often given

in two to five sessions to limit exposure to nearby tissue.

### **SSRI**

An SSRI is a selective serotonin-reuptake inhibitor antidepressant used for depression and several anxiety-related disorders; effects and withdrawal symptoms require monitoring.

### **SSRIs**

Selective serotonin reuptake inhibitors, a group of medicines that increase serotonin signaling in the brain. They are used for depression and several anxiety-related disorders; effects and side effects develop over time, not immediately.

### **Stable Angina**

Repeated, predictable chest pressure or discomfort brought on by activity or stress because the heart muscle receives too little blood. It usually improves with rest or prescribed medicine; new, severe, prolonged, or rest pain may be a heart attack and needs emergency care.

#### **Alternate Names:**

Chronic Stable Angina; Angina Pectoris

### **Stable Coronary Artery Disease**

Coronary artery narrowing causing a consistent, predictable pattern of angina.

#### **Alternate Names:**

Stable Ischemic Heart Disease; Chronic Coronary Syndrome

### **Stable Disease**

An oncology response category in which a cancer has neither shrunk enough to meet criteria for partial response nor grown enough to meet criteria for progressive disease. It describes change over a specified assessment interval and does not mean that the cancer is cured or inactive.

### **Stabs**

Brief, sharp pains or penetrating injuries described as stabbing. Causes vary with location and may include nerve irritation, muscle strain, infection,

---

inflammation, injury, or serious disease requiring urgent assessment.

### Stab Wound

A deep puncture caused by a pointed or sharp object. The wound may damage blood vessels, nerves, organs, or bone even when the skin opening looks small; heavy bleeding, severe pain, or a wound to the chest or abdomen requires emergency care.

### Stafne Bone Defect

A Stafne bone defect is a usually harmless, developmental indentation in the lower jaw caused by nearby salivary-gland tissue. It is often found incidentally on imaging.

### Stage 4 Prostate Cancer

Alternate name; see *Metastatic (stage 4) prostate cancer*.

### Stage I

A cancer stage that generally indicates limited disease, although the exact definition and prognosis depend on the cancer type and staging system.

### Stage II

A cancer stage that usually indicates a larger or more locally advanced tumor than stage I, without distant spread. The exact tumor size, lymph-node findings, prognosis, and treatment vary by cancer type and staging system.

### Stage III

A cancer stage that often indicates more extensive local or regional disease, with the precise criteria depending on the cancer type.

### Stage IV

A cancer stage that generally indicates spread to distant organs or sites. Prognosis and treatment vary by cancer type and individual factors.

### Stages Of Change

A model describing common steps people may pass through when changing a behavior. These in-

clude not yet considering change, thinking about it, preparing, taking action, and maintaining the change; people may move backward or forward between stages.

### Alternate Names:

Transtheoretical Model; Prochaska Model

### Stages of Labor

The three sequential stages of childbirth: the first stage extends from the onset of labor to complete cervical dilation and includes latent and active phases; the second stage extends from complete dilation to birth of the baby; and the third stage extends from birth to delivery of the placenta.

### Staghorn Calculi

Large branching kidney stones that fill part or all of the kidney's urine-collecting system. They are often linked with infection, can damage kidney tissue, and usually require specialist treatment to remove them.

### Staghorn Calculus

A branching kidney stone that fills part or all of the renal pelvis and calyces, often associated with infection.

### Staging

Determining how far a cancer has grown or spread by assessing the original tumor, nearby lymph nodes, and distant organs. The stage helps estimate outlook and choose treatment.

### Staining

A laboratory method that uses colored dyes to make cells, tissue structures, or microorganisms easier to see under a microscope. Different stains highlight different features and can help guide diagnosis, although results may need culture or other tests for confirmation.

### Stainless Steel

A corrosion-resistant alloy containing iron and chromium, used for instruments, implants, orthodontic appliances, and some dental crowns. Its suitability depends on the grade, strength, and tissue exposure.



---

## Stammering

Alternate name for stuttering, a speech-fluency difficulty involving repeated sounds or words, prolonged sounds, or blocks that interrupt speaking. It is common in childhood and may continue into adulthood; speech-language therapy can help.

## Standard Deviation

A number that shows how widely measurements differ from their average. A small standard deviation means values are close together, while a large one means they are more spread out; it does not by itself show whether results are good or bad.

## Standard Eye Exam

A comprehensive eye examination assessing visual acuity, refraction, pupils, eye movements, external structures, anterior eye, intraocular pressure, and ocular fundus as indicated.

## Standardized Extract

An herbal extract adjusted to contain a measured amount of selected ingredient, improving consistency but not guaranteeing safety or effectiveness.

## Standardized Uptake Value

A number used on a PET scan to estimate how much radioactive tracer a tissue has taken up, adjusted for the injected amount and body size. It can help compare areas or follow change, but it does not by itself prove cancer or another disease.

## Standard Of Care

The conduct, skill, diligence, and judgment a reasonably prudent practitioner of the same specialty would exercise under similar circumstances; the benchmark for medical negligence. It is established through guidelines, peer practice, and expert testimony rather than fixed statute.

## Standard Precautions

Basic infection-control measures used for every patient, because blood, body fluids, nonintact skin, and mucous membranes may carry germs. They include hand hygiene, appropriate protective equipment, safe injections, sharps safety, and respiratory

etiquette.

## Stannosis

A usually mild form of pneumoconiosis caused by long-term inhalation of tin oxide dust. It may produce visible lung changes on imaging, often without major breathing impairment.

## Stanozolol

A synthetic anabolic androgenic steroid with limited medical use and potential liver, lipid, and cardiovascular toxicity.

## Stapedectomy

Surgical removal or replacement of the stapes, usually to treat otosclerosis and conductive hearing loss.

## Stapedius

A small middle-ear muscle that helps limit movement of the stapes and may protect the inner ear from loud sounds.

## Stapes

The smallest middle-ear bone, passing sound vibrations from the incus to the inner ear for hearing.

## Staph Infection

Alternate informal name; see *Staphylococcal Infection*.

## Staph Infections In The Hospital

Infections caused by *Staphylococcus* bacteria that begin during or after healthcare treatment, or require hospital care. They may involve wounds, blood, lungs, bones, or devices and can be difficult to treat when the bacteria resist antibiotics.

## Staphylococcal Food Poisoning

An acute foodborne intoxication caused by the ingestion of preformed heat-stable enterotoxins produced by *Staphylococcus aureus* in improperly refrigerated or handled foods (such as dairy products, pastries, processed meats, and salads). Characterized by an abrupt onset of severe nausea, spasmodic abdominal

---

cramping, and violent vomiting within 1 to 6 hours of consumption, usually resolving spontaneously within 24 to 48 hours.

**Alternate Names:**

Staphylococcal Enterotoxigenesis; Staph Food Poisoning

**Staphylococcal Infection**

An infection caused by *Staphylococcus aureus* or another *Staphylococcus* species. It may affect the skin, bone, joints, lungs, heart valves, blood, or other tissues. Some strains resist multiple antibiotics. Diagnosis and antimicrobial treatment depend on the site, severity, and susceptibility of the organism.

**Alternate Names:**

Staph Infection; Staphylococcal Disease

**Staphylococcal Meningitis**

Staphylococcal meningitis is infection and inflammation of the meninges caused by *Staphylococcus* bacteria, often after surgery, trauma, bloodstream infection, or an invasive device.

**Staphylococcal Pneumonia**

Pneumonia caused by *Staphylococcus* bacteria, sometimes after influenza or another viral infection. It may cause fever, cough, chest pain, breathlessness, lung abscesses, or infected fluid around the lung and needs medical treatment.

**Staphylococcal Scalded Skin**

Alternate name; see *Staphylococcal Scalded Skin Syndrome*.

**Staphylococcal Scalded Skin Syndrome**

A toxin-mediated skin disorder caused by *Staphylococcus aureus*, usually affecting infants or young children. It produces widespread redness, tenderness, fragile superficial blisters, and peeling that can resemble a burn and requires prompt treatment.

**Staphylococcus**

A genus of gram-positive cocci that includes *S. aureus* and common skin colonizers and pathogens.

**Staphylococcus Aureus**

A bacterium that commonly lives on the skin or in the nose but can cause infections of the skin, blood, lungs, bones, joints, or heart valves. Some strains are resistant to several antibiotics.

**Stapling**

Closing or joining tissue with metal or absorbable surgical staples. Staples may be used for skin, blood vessels, or parts of the digestive tract and must be selected and removed or absorbed appropriately.

**Starch Poisoning**

Illness after swallowing a starch product, most often causing choking, blockage, or stomach upset rather than systemic poisoning. Inhaled powder can irritate the airways; breathing difficulty or persistent vomiting requires medical care.

**Stargardt Disease**

An inherited eye disorder that gradually damages the macula, the central part of the retina. It usually begins in childhood or young adulthood and causes loss of detailed central vision; side vision is often preserved.

**Starling Equation**

A mathematical physical equation describing the transcapillary fluid filtration rate ( $J_v$ ) as a function of the hydraulic conductance and the balance between hydrostatic pressure gradients (capillary minus interstitial) and oncotic pressure gradients (capillary minus interstitial) across microvascular endothelia.

**Starling Forces**

The balance of pressures that pushes water out of small blood vessels or pulls it back in. Changes can cause swelling in tissues, dehydration of tissues, or fluid buildup around organs.

**Starvation**

Severe lack of food or usable energy over time, causing weight and muscle loss, weakness, changes in body chemistry, reduced immunity, and eventually organ failure. Restarting nutrition after prolonged

---

starvation must be done carefully because dangerous shifts can occur.

### **Stasis**

Slowing or stopping of normal movement of blood, lymph, intestinal contents, or other body fluids.

### **Stasis Dermatitis**

Inflammation of the lower-leg skin caused by chronic venous hypertension and edema. It may cause itching, redness, scaling, brown discoloration, weeping, or ulcers around the ankles. Management focuses on treating venous disease, compression when safe, skin care, and evaluation for infection or arterial insufficiency.

### **Stasis Dermatitis And Ulcers**

Stasis dermatitis and ulcers result from chronic venous hypertension in the lower legs, causing swelling, brown skin change, inflammation, itching, and slow-healing wounds.

### **Stasis Ulcer**

A stasis ulcer is a chronic lower-leg wound caused by venous hypertension and poor return of blood, commonly near the ankle and associated with edema and skin changes.

### **STAT**

A medical instruction meaning immediately or with urgent priority, such as a stat test, medicine, or consultation.

### **Statin**

See *Statins*.

### **Statin Drugs**

See *Statins*.

### **Statin-Induced Myopathy**

Muscle pain, weakness, or tenderness caused by a statin cholesterol-lowering medicine. Symptoms may occur without muscle damage or, rarely, with severe muscle breakdown that can injure the kidneys. A clinician may check the medicine, symptoms, mus-

cle enzymes, and other causes before changing treatment.

### **Alternate Names:**

Statin-Associated Muscle Symptoms (SAMS); Statin Myotoxicity

### **Statins**

A class of medicines that lowers LDL cholesterol by reducing cholesterol production in the liver. They reduce the risk of heart attack and stroke in appropriate people; muscle symptoms, liver problems, and medicine interactions should be reviewed.

### **Alternate Names:**

HMG-CoA Reductase Inhibitors; Lipid-Lowering Agents

### **Statins In Diabetes**

Cholesterol-lowering treatment for people with diabetes who may benefit from reduced risk of heart attack and stroke. The medicine and dose depend on age, cardiovascular risk, kidney function, other medicines, and possible side effects.

### **Station**

A measure of how far the baby's presenting part, usually the head, has descended through the pelvis during labor. It is compared with the mother's pelvic bones and helps clinicians follow progress toward birth.

### **Status Asthmaticus**

A severe asthma attack that does not improve enough with the first treatments. Breathing may become dangerously difficult, with exhaustion, inability to speak normally, or low oxygen, and emergency hospital treatment is required.

### **Status Epilepticus**

A seizure lasting long enough or recurring often enough that recovery does not occur between seizures, creating a persistent seizure state and neurologic emergency. Operational treatment thresholds are about 5 minutes for generalized tonic-clonic seizures, 10 minutes for focal seizures, and 10 to 15 minutes for absence seizures; status may be convulsive or nonconvulsive.

---

## Status Migrainosus

A severe migraine attack that persists for more than 72 hours and may require urgent medical treatment, especially when dehydration, vomiting, or medication overuse is present.

## Stay Away From Asthma Triggers

Avoiding asthma triggers such as smoke, allergens, infections, fumes, and cold air can reduce symptoms and attacks, alongside prescribed controller and reliever treatment.

## STDs

Alternate name; see *Sexually transmitted diseases (STDs)*.

## Steady State

The point during regular medicine use when the amount entering the body over time is about equal to the amount leaving it. Blood levels then become relatively stable; reaching this point usually takes several doses and depends on the medicine's half-life.

## Steal Syndrome

Reduced blood flow to a hand, foot, or other area because a bypass graft or dialysis fistula diverts too much blood. It may cause pain, coolness, pale or blue skin, numbness, or weakness and requires vascular assessment.

## Steatocystoma Multiplex

A disorder producing multiple sebum-filled cysts arising from the pilosebaceous units, commonly on the trunk, chest, arms, or face. Lesions are usually skin-colored and painless but may become inflamed; treatment depends on symptoms and extent.

## Steatoma

A fatty or sebaceous mass; the term may refer to a lipoma or, in the ear, a cholesteatoma depending on context.

## Steatorrhea

The passage of excess fat in the stool, causing greasy, pale, foul-smelling, floating stools—a hallmark of

fat malabsorption.

## Steatorrhoea

Excess fat in stool, causing pale, bulky, greasy, foul-smelling bowel movements. It suggests poor fat digestion or absorption from pancreatic, liver, bile-duct, intestinal, or other digestive disease.

## Steatotic Liver Disease (SLD)

A group of liver disorders characterized by excess fat accumulation in liver cells. Causes include metabolic dysfunction, alcohol, medicines, and other conditions; inflammation, fibrosis, cirrhosis, or liver cancer may develop in some people.

## Steele-Richardson-Olszewski Syndrome

Alternate name; see *Progressive supranuclear palsy*.

## Stein-Leventhal Syndrome

Alternate name; see *Polyendocrine metabolic ovarian syndrome*.

## ST-Elevation Myocardial Infarction

A heart attack associated with characteristic ST-segment elevation on an electrocardiogram, usually indicating sudden complete blockage of a coronary artery. It causes acute injury to heart muscle and requires rapid emergency evaluation and restoration of blood flow.

## Stem Cell

An unspecialized cell with two defining abilities: self-renewal, or making more stem cells, and differentiation, or developing into specialized cells. The range of possible cell types depends on the stem cell's potency; adult stem cells are often tissue-restricted, while embryonic and induced pluripotent stem cells can form many cell types.

## Stem Cell Donation

The collection of healthy blood-forming stem cells from a donor for transplantation into another person. Cells may be collected from the bloodstream by apheresis after medication increases their number in the blood, or directly from bone marrow under

---

anesthesia. Donor and recipient compatibility, especially human leukocyte antigen matching, helps improve transplant success.

### **Stem Cells**

Cells that can self-renew and produce specialized cell types within a range determined by their potency. They help maintain or repair some tissues, and blood-forming stem-cell transplantation is an established clinical use; many other stem-cell applications remain investigational.

### **Stem Cell Transplantation**

Hematopoietic stem cell transplantation replaces diseased or ablated bone marrow with healthy donor or autologous stem cells.

### **STEMI**

A type of heart attack caused by sudden blockage of a heart artery, usually producing characteristic changes on an electrocardiogram. It is an emergency because quickly restoring blood flow can limit permanent heart damage.

#### **Alternate Names:**

ST-Elevation Myocardial Infarction

### **Stenosing Tenosynovitis**

Alternate name; see *Trigger finger*.

### **Stenosis**

Abnormal narrowing of a passage, blood vessel, heart valve, spinal canal, or other opening. Narrowing may restrict flow or compress nearby structures.

### **Stenosis of the LAD**

Narrowing of a major artery supplying the front of the heart. Reduced blood flow may cause chest pain during exertion or a heart attack if the narrowing becomes severe or the artery suddenly blocks.

### **Stenotic**

Stenotic means narrowed, as in a stenotic heart valve, blood vessel, airway, intestine, or other passage.

### **Stenotrophomonas Maltophilia Resistance**

The ability of *Stenotrophomonas maltophilia* bacteria to survive several antibiotics. Treatment depends on the infected body site, illness severity, and laboratory testing that identifies which medicines are likely to work.

### **Stensen Duct**

The main excretory duct of the parotid gland that traverses the masseter muscle and opens into the oral cavity through a small papilla on the buccal mucosa opposite the second maxillary molar.

#### **Alternate Names:**

Parotid Duct; Stensen's Duct

### **Stent**

A tube or scaffold placed inside a vessel or other passage to keep it open.

### **Stent Graft**

A fabric-covered stent placed inside a blood vessel to reinforce the vessel wall or exclude an aneurysm from blood flow.

### **Stenting**

Placement of a small mesh tube to keep a narrowed or blocked passage open. Vascular stents restore blood flow, while drug-coated types release medicine to reduce repeated narrowing.

### **Stent Placement**

Insertion of a mesh tube or other support to keep a narrowed or blocked passage open. Stents may be used in arteries, bile ducts, ureters, airways, or the esophagus; bleeding, blockage, infection, movement, or repeated narrowing can occur.

### **Stent Restenosis**

Re-narrowing of a blood vessel within or adjacent to a previously placed stent, usually caused by tissue growth, recurrent atherosclerosis, or thrombosis.

### **Stepping Reflex**

A primitive reflex present from birth to 2 months,

---

elicited by holding the infant upright with feet touching a flat surface. The infant makes stepping-like movements as if walking. Disappears by 2 months as voluntary motor control develops. Reappears around 9-10 months as voluntary walking emerges.

### **Stereognosis**

The ability to recognize a familiar object by feeling it without looking. It requires intact skin sensation, nerve pathways, and areas of the brain that interpret touch; loss may occur after nerve or brain injury.

### **Stereotactic**

Stereotactic describes a technique using three-dimensional coordinates to target a precise location in the brain or another body structure.

### **Stereotactic Biopsy**

Image-guided biopsy using three-dimensional coordinates to target a lesion that may be difficult to localize by direct palpation. It is used in the breast and, with specialized systems, the brain, lung, and other organs.

### **Stereotactic Body Radiation Therapy**

A focused form of external radiation that delivers a high dose to a small body tumor in a few treatments. It may treat selected lung, liver, kidney, pancreatic, or spinal tumors while limiting exposure to nearby tissue.

### **Stereotactic Brain Biopsy**

A precision, minimally invasive neurosurgical procedure used to safely obtain small tissue samples from deep, delicate, or high-risk areas of the brain for microscopic examination. Using high-resolution pre-operative MRI or CT scans and a specialized three-dimensional stereotactic head frame or frameless optical infrared navigation system, a neurosurgeon maps exact mathematical coordinates to the suspected tumor or infection. Through a tiny circular burr hole (approximately 3 to 5 mm) in the skull, the surgeon gently passes a specialized hollow biopsy needle straight to the target lesion without disrupting surrounding healthy brain tissue. The extracted tissue specimens are examined by a neu-

ropathologist to determine the exact tumor type, grade, genetic markers, or underlying infection, allowing doctors to design targeted chemotherapy, radiation, or medical therapy.

#### **Alternate Names:**

Stereotactic Needle Biopsy; Brain Biopsy; Frameless Stereotactic Biopsy

### **Stereotactic Breast Biopsy**

A minimally invasive percutaneous core-needle biopsy procedure utilizing digital mammographic triangulation from angled X-ray views to precisely target and sample nonpalpable microcalcifications or architectural distortions.

#### **Alternate Names:**

Stereotactic Breast Biopsy; Stereotactic Core Needle Biopsy; Mammographically Guided Breast Biopsy

### **Stereotactic Radiosurgery**

Precise radiation treatment delivering one or a few high doses to a small target, often in the brain, without open surgery.

### **Stereotaxis**

A technique using a three-dimensional coordinate system to target deep structures, especially in brain procedures.

### **Stereotypic Movement Disorder**

Repeated, apparently purposeless movements such as hand flapping, rocking, or head banging that interfere with activities or cause injury. It is more common in childhood and developmental conditions and requires assessment for other causes.

### **Sterile Field**

A designated area maintained free of viable microorganisms during an invasive procedure. Only sterile equipment and materials may contact the field or the sterile portions of instruments.

### **Sterile Site**

A body area or sample that normally contains no living microorganisms, such as blood, spinal fluid, joint fluid, or tissue from a closed space. Finding a



---

germ there is usually important, although accidental contamination must be considered.

### **Sterile Technique**

Methods used to prevent introduction of microorganisms into a sterile field, wound, tissue, device, or body site through hand hygiene, barriers, aseptic handling, and appropriate preparation.

### **Sterilisation**

Destruction or removal of all forms of microbial life, or permanent contraception depending on context.

### **Sterility**

Inability to reproduce, or complete absence of living microorganisms from an object or preparation, depending on context. In healthcare, sterile equipment, implants, and medicines must remain free from contamination to prevent infection.

### **Sterilization**

A process that destroys or removes all forms of microbial life, including spores, from equipment, medicines, or other materials. In another context, sterilization means a permanent method intended to prevent pregnancy.

### **Sternal Angle**

The ridge where the manubrium joins the body of the sternum. It marks the level where the second ribs attach and helps locate structures inside the chest.

### **Sternal Exploration Or Closure**

Surgical reopening or repair of the breastbone after median sternotomy, performed to treat infection, bleeding, instability, dehiscence, or other postoperative chest complications.

### **Sternal Fracture**

A break in the breastbone, most often caused by direct chest impact or compression. It may cause localized pain and tenderness and can accompany cardiac, pulmonary, or mediastinal injury.

### **Sternal Line**

An imaginary vertical line along the outer edge of the breastbone. It is used as a surface landmark when describing the location of chest structures or physical findings.

### **Sternoclavicular Joint**

A saddle-type synovial joint between the clavicle and manubrium of the sternum. It is the only bony connection between the upper limb and axial skeleton and permits movement of the shoulder girdle.

### **Sternocleidomastoid Muscle**

A prominent muscle on each side of the neck that turns and bends the head and helps stabilize the neck. It can also assist forceful breathing; strain or spasm may cause neck pain and limited movement.

### **Sternocostal Joint**

A joint where a rib or its cartilage meets the breastbone. These joints help form the front of the chest wall.

### **Sternotomy**

Surgical division of the sternum, usually along the midline, to provide access to the heart or mediastinal structures.

### **Sternum**

The breastbone forming the front of the central chest wall.

### **Sternutation**

Sneezing, a reflex forceful expulsion of air through the nose and mouth triggered by irritation of nasal or upper-airway sensory receptors.

### **Steroid Myopathy**

Muscle weakness caused by prolonged exposure to corticosteroid medicines or excessive cortisol. It usually develops gradually and affects the hips, thighs, shoulders, and upper arms, making it hard to rise from a chair or climb stairs. Weakness may improve when the cause is treated, but steroid doses must be changed only with medical guidance.

---

**Alternate Names:**

Corticosteroid-Induced Myopathy; Glucocorticoid Myopathy

**Steroids**

A broad family of compounds with a characteristic four-ring structure, including body hormones, corticosteroids, and sex hormones. Their effects depend on the type; prolonged corticosteroid use can suppress immunity and bone health.

**Alternate Names:**

Corticosteroids; Glucocorticoids

**Stertor**

A low-pitched snoring, gurgling, or rattling sound caused by partly blocked air passages in the nose or throat. It may occur during sleep or illness; noisy breathing with severe sleepiness, blue lips, or breathing difficulty is an emergency.

**Stethoscope**

An instrument used for auscultation, the examination of sounds produced within the body. It usually has a chest piece with a diaphragm and sometimes a bell, flexible tubing, and earpieces; electronic models convert sounds into amplified or recorded signals. It is used to examine heart, lung, bowel, and vascular sounds.

**Stevens-Johnson Syndrome**

A rare, life-threatening reaction usually caused by a medicine or infection. It causes painful widespread skin and mucous-membrane damage, blistering, and peeling and requires emergency hospital treatment.

**Alternate Names:**

SJS

**Stewart-Treves Syndrome**

An angiosarcoma arising in long-standing lymphedema, classically after mastectomy but also after congenital lymphedema, radiation, infection, or other lymphatic damage. New bruised, purple, nodular, or ulcerated areas in chronically swollen skin need urgent biopsy.

**STI**

Alternate name; see *Sexually Transmitted Infections*.

**Stickler Syndrome**

An inherited connective-tissue disorder that may cause severe nearsightedness, retinal detachment, hearing loss, early arthritis, and differences in the jaw or palate. Regular eye and hearing checks are important.

**Stiff Baby Syndrome**

A nonspecific description of unusual stiffness or increased muscle tone in an infant, not a single diagnosis. Causes include brain, nerve, muscle, genetic, or metabolic disorders; persistent stiffness, poor feeding, seizures, or breathing problems need urgent assessment.

**Stiffness**

Difficulty or discomfort when moving a body part. It may result from joint disease, muscle problems, nerve disorders, inflammation, injury, prolonged rest, or aging and can limit everyday activities.

**Stiff-Person Syndrome**

A rare immune-related nervous-system disorder causing persistent muscle stiffness and sudden painful spasms. Noise, touch, movement, or emotional stress may trigger spasms, leading to falls and difficulty walking. Blood tests may find related antibodies. Treatment can reduce stiffness and immune activity, but symptoms often require long-term care.

**Alternate Names:**

SPS; Stiff-Man Syndrome; Moersch-Woltman Syndrome

**Stigma**

A mark, characteristic, or social discrediting label; in medicine it may refer to a sign or to harmful social attitudes.

**Stilboestrol**

Historical name; see *Diethylstilbestrol*.

---

## Stilet

A thin, pointed instrument placed inside a cannula or tube to guide insertion, clear a blockage, or provide firmness. It must be used carefully because it can injure tissue or blood vessels.

## Still Disease-Associated Arthritis

Joint inflammation occurring in Still disease. In children, the related diagnosis is systemic juvenile idiopathic arthritis; in adults it is adult-onset Still disease.

## Stillbirth

The death of a fetus before or during delivery after the gestational-age or birth-weight threshold used by a country or health system. In the United States, it usually refers to loss at or after 20 weeks of pregnancy; the World Health Organization commonly uses 28 weeks for international comparisons. Causes may include problems with the placenta, fetal growth, infection, genetic or developmental conditions, or maternal illness, but some stillbirths remain unexplained.

## Still Disease

An inflammatory disease that can affect the whole body. In children it is called systemic juvenile idiopathic arthritis; in adults it is called adult-onset Still disease. It can cause fever, joint pain or swelling, and a rash.

### Alternate Names:

Systemic Juvenile Idiopathic Arthritis; Adult-Onset Still Disease; AOSD

## Still's Disease

An inflammatory illness that can affect children or adults. It commonly causes repeated high fevers, a fleeting rash, joint pain or swelling, and marked inflammation; it can affect other organs and usually requires specialist treatment.

## Still's Murmur

A common harmless heart murmur in children, usually heard between ages 2 and 7. It has a soft, musical sound, becomes louder with fever or exercise, and usually disappears as the child grows; the

heart is otherwise normal.

### Alternate Names:

Innocent Vibratory Murmur; Normal Pediatric Murmur; Twanging-String Murmur

## Stimulant

A substance that increases activity in the brain or other body systems. Stimulants may improve alertness and attention, but excessive or inappropriate use can cause dependence, anxiety, fast heartbeat, high blood pressure, or seizures.

## Stimulants

Substances that increase activity in the brain or the body's fight-or-flight system. They may improve alertness and attention but can also cause insomnia, anxiety, dependence, fast heartbeat, high blood pressure, or seizures.

## Stimulant Use Disorder

A pattern of using stimulant drugs that causes significant problems or loss of control despite harm. Stimulants include cocaine and amphetamine-type drugs; stopping may cause tiredness, low mood, increased appetite, or excessive sleep.

## Stimulus

A stimulus is a physical, chemical, sensory, or social change that evokes a response from a cell, tissue, organism, or person.

## Stingray

A sea-animal injury caused by a stingray spine, usually producing severe pain, swelling, and a puncture wound. Medical care may be needed to clean the wound, remove retained spines, and treat whole-body reactions.

## Stings

Injuries caused when an insect, animal, or plant injects venom or irritating material. They may cause pain, swelling, itching, allergic reactions, tissue damage, or serious poisoning depending on the source.

---

## Stippling

Patterned dermal abrasions and punctate hemorrhages surrounding a firearm entrance wound, caused by unburned and partially burned gunpowder grains striking the skin at close range (15–75 cm).

### Alternate Names:

Gunpowder Stippling; Powder Tattooing

## Stitch

A suture used to close tissue, or a sharp transient pain, often in the side during exercise.

## St John's Wort

A herbal product made from *Hypericum perforatum* and sometimes used for depressive symptoms. It can cause side effects and reduce the effectiveness of many medicines, including some contraceptives, anticoagulants, transplant medicines, and HIV treatments.

## Stochastic Effects

Possible effects of radiation, especially cancer or inherited genetic changes, whose chance increases with dose. The severity cannot be predicted from the dose, and even low exposure may carry some risk.

## Stoddard Solvent Poisoning

Poisoning from Stoddard solvent, a petroleum-based cleaning solvent. Inhalation or swallowing can cause headache, dizziness, drowsiness, skin irritation, or aspiration-related lung injury; do not induce vomiting and seek advice.

## Stokes-Adams Syndrome

Sudden fainting caused by a temporary dangerous slowing or stopping of the heartbeat, often from heart block or another abnormal rhythm. It requires urgent heart assessment because injury or cardiac arrest can occur.

## Stoma

A natural opening or a surgically created opening connecting an organ to the skin or another organ.

## Stomach

A muscular digestive organ between the oesophagus and small intestine that stores and begins digestion of food.

## Stomach Cancer

Cancer that begins in the stomach, most often in gland-forming cells lining the stomach. Risk factors include *Helicobacter pylori* infection, smoking, chronic inflammation, certain foods, and inherited conditions; symptoms and treatment depend on type and stage.

### Alternate Names:

Gastric Carcinoma; Gastric Neoplasm

## Stomach Cramps

Intermittent gripping or spasmodic upper-abdominal pain caused by gas, infection, indigestion, ulcer disease, gallbladder or pancreatic disease, obstruction, or other causes.

## Stomach Diseases

Disorders affecting the stomach, including inflammation, ulcers, movement problems, blockage, bleeding, and tumors. Symptoms may include pain, nausea, vomiting, early fullness, weight loss, indigestion, or blood in vomit or stool.

## Stomach Flu

A common informal term for viral gastroenteritis, an infection of the stomach and intestines caused by viruses rather than influenza viruses. It may cause watery diarrhea, vomiting, abdominal cramps, fever, and dehydration; some infections cause few or no symptoms.

## Stomach Pain

Stomach pain is discomfort in the upper abdomen. Causes range from indigestion and infection to ulcers, gallbladder disease, or pancreatitis; severe or persistent pain needs medical assessment.

## Stomach Paralysis

Stomach paralysis, or gastroparesis, is delayed emptying of stomach contents without obstruction, caus-

---

ing early fullness, nausea, vomiting, and bloating.

### **Stomach Tube**

A tube passed through the mouth or nose into the stomach for drainage, feeding, sampling, or treatment.

### **Stomach Ulcer**

See *Gastric Ulcer*.

### **Stomach Washout**

A procedure that rinses and removes stomach contents through a tube. It is rarely used for selected serious poisonings because it can cause choking, lung injury, bleeding, or other complications.

### **Stoma Complications**

Problems involving an opening made from the intestine or urinary tract to the skin. They include skin irritation, bleeding, retraction, prolapse, narrowing, blockage, high output, infection, and a parastomal hernia; sudden severe pain, color change, or no output needs urgent care.

### **Stomatitis**

Inflammation, soreness, redness, or ulcers of the lining of the mouth. Causes include infection, irritation, nutritional deficiency, medicines, and cancer treatment; severe pain, dehydration, fever, or inability to eat or drink needs medical care.

### **Stomatodynia**

Alternate name; see *Burning mouth syndrome*.

### **Stone**

A hard lump formed inside an organ or tube, commonly from mineral salts or other body substances. Stones may block flow, cause pain or infection, and occur in kidneys, gallbladder, or ducts.

### **Stonefish Sting**

A puncture injury from the venomous spines of a stonefish, causing severe pain, swelling, bleeding, and sometimes weakness, breathing problems, or shock. Hot-water immersion may reduce pain, but

severe poisoning can affect breathing and circulation.

### **Stool**

Fecal material expelled from the bowel, composed of water, undigested residue, bacteria, cells, mucus, and bile pigments; color, form, frequency, and blood can provide clinical clues.

### **Stool C Difficile Toxin**

A stool test for harmful substances made by *Clostridioides difficile* bacteria. A positive result supports active infection when a person has compatible diarrhea, but bacteria can be present without causing illness.

### **Stool Examination**

Laboratory analysis of a stool sample for blood, infection, parasites, inflammation, malabsorption, or other digestive abnormalities.

#### **Alternate Names:**

Fecal Analysis; Stool Test; Coprology

### **Stool Gram Stain**

Microscopic examination of a stool specimen after Gram staining; it may provide limited information about bacterial flora or inflammatory cells but is not a general test for infectious diarrhea.

### **Stool Guaiac Test**

A chemical fecal occult blood test that detects the peroxidase activity of heme in stool. Dietary factors, medicines, and specimen collection can affect results. It is a screening or detection test rather than a diagnosis; a positive result requires clinical follow-up and usually evaluation for a gastrointestinal source.

### **Stool Holding**

Alternate name; see *Encopresis*.

### **Stool Occult Blood Test**

Alternate name; see *Fecal occult blood test*.

### **Stool Ova And Parasites Exam**

Microscopic examination of stool for parasite eggs,

---

cysts, larvae, or other forms, sometimes requiring multiple specimens to improve detection.

### **Stools**

Feces, the material expelled from the digestive tract. Changes in color, consistency, frequency, or blood can provide clues about digestive health but are not diagnostic alone.

### **Stools – Floating**

Feces that remain on the water surface, often from trapped gas or diet but sometimes from excess fat in malabsorption; persistent greasy, pale, foul-smelling stool needs evaluation.

### **STOPP/START**

A checklist used to review medicines in older adults. STOPP identifies medicines that may be inappropriate or harmful, while START identifies useful treatments that may have been missed; clinical judgment is still required.

### **Storage Diseases**

A group of inherited metabolic disorders characterized by accumulation of substrates within cells due to specific enzyme deficiencies. Lysosomal storage diseases result from defects in lysosomal hydrolytic enzymes, leading to accumulation of undigested macromolecules within lysosomes.

### **Stork Bite**

A stork bite is a common harmless pink birthmark, usually a nevus simplex on the forehead, eyelids, nose, or neck, that often fades during childhood.

### **Strabismus**

A condition in which the eyes are not aligned and do not point in the same direction. One eye may turn inward, outward, upward, or downward. It may be constant or come and go. Strabismus can cause double vision in adults and may lead to amblyopia, or reduced vision in one eye, when it begins during childhood. Common types include esotropia, exotropia, hypertropia, and hypotropia.

### **Straddle Injury**

A direct blunt compressive trauma to the perineum occurring when a patient falls astride a hard, rigid object (such as a bicycle crossbar, fence, or beam), frequently causing crushing rupture of the anterior bulbous urethra beneath the pubic symphysis.

### **Straight Leg Raise Test**

A provocative physical examination test for lumbosacral nerve root irritation (L4-S1, typically due to herniated intervertebral disc), in which the patient lies supine and the examiner passively elevates the extended lower extremity; reproduction of electric shooting radicular pain down the leg between 30 and 70 degrees indicates a positive result.

#### **Alternate Names:**

Lasegue Sign; SLR Test

### **Strain**

An injury from overstretching or tearing a muscle or tendon, or a force applied to tissue.

### **Strain Imaging**

Ultrasound or magnetic-resonance imaging that measures how much tissue deforms during movement, especially how heart muscle shortens or thickens with each heartbeat. It can reveal early weakness even when the total amount of blood pumped appears normal.

### **Strains**

A strain is overstretching or tearing of a muscle or tendon, causing pain, tenderness, swelling, weakness, or reduced function after force or overuse.

### **Strangulated Hernia**

A hernia in which the blood supply to the trapped tissue is cut off, causing ischaemia and possible tissue death; it is a surgical emergency.

### **Strangury**

Painful, urgent, or difficult urination in which only small amounts of urine are passed. It may result from inflammation or irritation of the bladder or urethra, infection, stones, or obstruction.



---

## Strangulation

Compression of a body structure, especially the neck or a hernia, that compromises blood flow or airway.

## Strapping

Application of adhesive tape or bandage to support, protect, or restrict movement of a body part.

## Stratum Corneum

The outermost skin layer, made mainly of flattened dead cells filled with keratin. It forms a protective barrier that limits water loss and helps block chemicals, germs, and physical injury.

## Strawberry Hemangioma

Alternate name; see *Hemangioma*.

## Strawberry Tongue

A tongue that looks bright red and bumpy like a strawberry, often seen in childhood illnesses like scarlet fever or Kawasaki disease.

## Strength

The ability of a muscle or group of muscles to produce force. Healthcare professionals may test strength by comparing movements against resistance; reduced strength can result from pain, poor effort, muscle disease, nerve problems, or brain injury.

## Strength Testing

Assessment of how much force a muscle or muscle group can produce, using examination or equipment. It helps identify weakness and monitor recovery, progression, or response to treatment.

## Strength Training

Exercise against resistance to improve muscle force, endurance, power, bone health, and function. The program should be progressed safely according to ability and medical needs.

### Alternate Names:

Resistance Training; Weight Training

## Strep Throat

A bacterial infection of the throat and tonsils caused by group A Streptococcus, also called Streptococcus pyogenes. It commonly causes a sudden sore throat, pain when swallowing, fever, and swollen or red tonsils, sometimes with white patches.

## Streptococcal Cellulitis With Lymphangitis

A streptococcal skin infection with inflammation spreading along nearby lymphatic vessels. Redness, warmth, pain, swelling, fever, and red streaks may occur and require prompt antibiotic treatment.

## Streptococcal Infection

An infection caused by bacteria of the Streptococcus group. Depending on the species and site, it may cause strep throat, scarlet fever, impetigo, cellulitis, pneumonia, heart-valve infection, bloodstream infection, or severe tissue infection.

## Streptococcal Infections

Infections caused by Streptococcus bacteria, which may affect the throat, skin, lungs, heart valves, blood, or other tissues. Diagnosis and antibiotic choice depend on the site and species; rapidly worsening illness, severe pain, or breathing difficulty is urgent.

## Streptococcal Pharyngitis

Alternate name; see *Strep Throat*.

## Streptococcal Screen

A streptococcal screen looks for group A Streptococcus, usually from a throat swab, to help diagnose strep throat.

## Streptococcal Toxic Shock Syndrome

A fulminant, life-threatening invasive infection caused by Streptococcus pyogenes (group A Streptococcus) producing streptococcal pyrogenic exotoxins that act as superantigens. It is characterized by the sudden onset of hypotension, shock, and multiorgan failure, frequently in association with severe invasive soft-tissue infections such as necrotizing fasciitis; it carries an exceptionally high mortality rate.

---

**Alternate Names:**

STSS; Streptococcal TSS; Toxic Shock-Like Syndrome

**Streptococcus**

A group of round bacteria that can cause throat infections, pneumonia, skin infections, heart-valve infection, or serious bloodstream illness. Different species spread and respond to antibiotics differently, so testing may guide treatment.

**Streptococcus Agalactiae**

Group B Streptococcus, a bacterium that can live in the bowel or genital tract without causing symptoms. It can pass to a baby during birth and cause serious newborn infection, so pregnancy screening and preventive antibiotics are used when indicated.

**Streptococcus pneumoniae**

Streptococcus pneumoniae is a bacterium that can cause pneumonia, meningitis, ear infection, sinusitis, and bloodstream infection. Vaccination helps prevent serious disease.

**Streptococcus Pyogenes**

Group A Streptococcus, a bacterium that causes strep throat, scarlet fever, impetigo, cellulitis, and sometimes severe invasive infection. Untreated infection can occasionally lead to rheumatic fever or kidney inflammation.

**Streptokinase**

A thrombolytic protein that activates plasminogen and dissolves clots; use is limited by bleeding risk and alternatives.

**Streptomycin**

An aminoglycoside antibiotic used in selected tuberculosis and other infections.

**Stress**

A physical and emotional response to perceived demands or threats. It can change heart rate, breathing, muscle tension, attention, sleep, and digestion; persistent or overwhelming stress may affect health.

**Stress Cardiomyopathy**

A temporary weakening of the heart muscle that often follows severe emotional or physical stress. It can cause chest pain, breathlessness, or fainting and may resemble a heart attack, even when no heart artery is blocked.

**Stress Echocardiography**

An ultrasound scan of the heart performed while the heart works harder through exercise or medicine. It looks for changes in heart-muscle movement that may suggest reduced blood flow.

**Stress Fracture**

A small crack in a bone caused by repeated activity or loading. Pain usually develops gradually in one spot and worsens with activity; continuing to stress the bone can make the crack larger or complete.

**Stress Fractures**

Small cracks in bone caused by repeated loading that exceeds the bone's ability to repair itself. They commonly affect the feet or legs and cause localized pain that worsens with activity.

**Stress Incontinence**

Involuntary leakage of urine during activities that increase intra-abdominal pressure, such as coughing, sneezing, lifting, or exercise.

**Stress Injuries**

Tissue injuries caused by repeated loading or overuse without sufficient recovery. Examples include stress fractures, tendinopathy, and muscle strain.

**Stressors**

Events, conditions, or circumstances that trigger a physical or emotional stress response. Examples include illness, pain, sleep loss, trauma, work demands, and relationship difficulties.

**Stress Related**

Describes symptoms or health problems that are triggered or worsened by physical or emotional stress. Stress can affect sleep, pain, digestion, mood,

---

heart rate, and blood pressure, but symptoms still require assessment for other causes.

### **Stress Response**

Body changes that prepare a person to respond to a perceived threat or challenge. Hormones and nerve signals alter heart rate, breathing, blood flow, attention, digestion, immunity, and energy use.

#### **Alternate Names:**

Fight-or-Flight Response; General Adaptation Syndrome

### **Stress Shielding**

Loss or weakening of bone next to an implant because the implant carries more force than the bone normally would. Over time, this can reduce bone density and affect how securely the implant stays in place.

### **Stress Tests**

Diagnostic tests that assess organ function under controlled physical, pharmacologic, or physiologic stress, commonly evaluating the heart for inducible ischemia or abnormal rhythm.

### **Stress Ulcer Prophylaxis**

Measures used to reduce the risk of stomach or duodenal bleeding in selected critically ill people. The need for treatment depends on bleeding risk, other medicines, and the person's overall condition.

### **Stress Urinary Incontinence**

Leakage of urine when coughing, sneezing, laughing, lifting, exercising, or otherwise increasing pressure inside the abdomen. It commonly results from weakness of the pelvic-floor muscles or the bladder outlet and can often be treated.

### **Stretch Mark**

A stretch mark is a linear scar caused by stretching and changes in the deeper skin, often during pregnancy, growth, or rapid weight change. It is harmless and usually fades over time.

### **Stretch Marks**

Linear bands of atrophic skin, also called striae, caused by stretching and hormonal or corticosteroid effects; they commonly occur during pregnancy, growth, weight change, or steroid exposure.

### **Stretch Reflex**

An automatic tightening of a muscle when it is suddenly stretched, such as the lower-leg movement after tapping the knee tendon. It helps maintain posture and provides information about nerve and spinal-cord function.

### **Stria**

A line, streak, or band in tissue, such as a stretch mark. Striae may result from skin stretching, growth, hormone changes, medicines, or underlying disease and can fade over time.

#### **Alternate Names:**

Striae; Stretch Lines

### **Striae**

Linear bands or streaks in the skin, commonly called stretch marks. They develop when deeper skin layers stretch or change rapidly during growth, pregnancy, weight change, muscle gain, or hormone treatment.

### **Striae Atrophicae**

Linear bands of thin, wrinkled skin caused by stretching and disruption of dermal connective tissue; commonly called stretch marks.

#### **Alternate Names:**

Stretch Marks; Striae Distensae

### **Striae Distensae (Stretch Marks)**

Linear bands caused by stretching and structural change in the dermis during growth, pregnancy, rapid weight change, muscle gain, or corticosteroid exposure. They may begin red or purple and later become pale and atrophic.

### **Striae Gravidarum**

Stretch marks that develop during pregnancy, commonly on the abdomen, breasts, thighs, or buttocks.

---

They may begin as red or purple lines and gradually fade to lighter bands.

### **Stricture**

An abnormal narrowing of a passage in the body, such as the esophagus, intestine, bile duct, or urethra. Scarring, inflammation, injury, surgery, or cancer may cause blockage and symptoms related to the affected organ.

### **Stridor**

A harsh, high-pitched breathing sound caused by air squeezing through a narrowed windpipe or voice box. It is heard loudest over the neck and usually happens while breathing in. This makes it different from wheezing, which comes from inside the lungs when breathing out. In babies and young children, common causes include croup, an object stuck in the throat, or floppy airway cartilage. In adults, causes include severe allergic swelling, vocal-cord problems, infections, or growths in the neck. Doctors find the cause by checking oxygen levels, taking neck X-rays, or looking into the throat with a tiny camera.

### **String Sign**

A very narrow, thread-like passage seen on an imaging test of the digestive tract. It may occur with narrowing from Crohn disease or with narrowing at the stomach outlet in an infant, but it is not specific to one condition.

### **String Test For Intestinal Parasites**

A string test collects a sample from the upper digestive tract on a swallowed capsule and string to look for selected intestinal parasites.

### **Stripping of Membranes**

A procedure performed during a cervical exam where the examiner inserts a finger through the internal os and sweeps the membranes circumferentially, separating them from the lower uterine segment. Releases endogenous prostaglandins, promoting cervical ripening and potentially initiating labor.

#### **Alternate Names:**

Membrane Sweeping; Cervical Sweep

### **Stroke**

Brain injury caused by interrupted blood flow to part of the brain (ischemic stroke) or bleeding in or around the brain (hemorrhagic stroke). It may cause sudden facial drooping, arm or leg weakness or numbness, speech trouble, vision loss, severe headache, dizziness, or confusion. The effects vary according to the brain area involved and the extent of injury.

### **Stroke Index**

A measure of the amount of blood pumped by the heart with each beat, adjusted for body-surface area. It helps assess heart function and is interpreted with other examination, imaging, and hemodynamic findings.

### **Stroke Rehabilitation**

Rehabilitation after a stroke helps improve movement, speech, swallowing, thinking, mood, and daily activities while reducing complications and preventing another stroke.

### **Stroke Risk Factors**

Stroke risk factors include high blood pressure, smoking, diabetes, high cholesterol, atrial fibrillation, prior stroke, and certain blood or vascular disorders.

### **Stroke Volume**

The volume of blood ejected by the left ventricle with each heartbeat, normally 60-100 mL. It is determined by three factors: preload, afterload, and myocardial contractility. Stroke volume equals end-diastolic volume minus end-systolic volume. Reduced stroke volume occurs in heart failure, hypovolemia, and cardiogenic shock.

### **Stroma**

The supporting tissue of an organ that surrounds and holds its working cells. It may contain connective tissue, blood vessels, nerves, and immune cells; the exact structure and role vary between organs.

### **Strongyloides Stercoralis**

A soil-transmitted roundworm that can enter

---

through skin and live in the human intestine. Infection may cause rash, abdominal symptoms, or no symptoms, but severe hyperinfection can occur when immunity is weakened.

### **Strongyloidiasis**

An intestinal worm infection that can spread throughout the body and become life-threatening when immunity is weakened.

### **Structured Reporting**

Reporting that uses a standard format, headings, and required fields to organize findings. It can improve completeness, consistency, and communication in imaging, pathology, and other clinical reports, but does not replace professional judgment.

### **Struma Ovarii**

A specialized mature ovarian teratoma composed predominantly of thyroid tissue; it may be hormonally active or develop malignant thyroid-type transformation.

### **Struvite Stones**

Kidney or urinary-tract stones formed during infection with certain bacteria. They can grow into large branching stones that fill the kidney and may damage it or block urine flow.

### **Strychnine**

A poison causing severe painful muscle spasms and convulsions while awareness may remain; emergency treatment is required.

### **ST Segment**

The part of an electrocardiogram between the electrical activation and recovery signals of the heart's lower chambers. Abnormal elevation or depression can occur with a heart attack, inflammation, or other conditions and must be interpreted with symptoms and the full tracing.

### **ST-Segment Elevation**

Elevation of the ECG ST segment above the isoelectric baseline, reflecting abnormal ventricular repolarization.

It may indicate acute coronary occlusion but can also occur in pericarditis, early repolarization, ventricular aneurysm, or other conditions; interpretation requires clinical context and serial testing.

### **Stucco Keratosis**

Multiple small, rough, whitish or gray-brown keratotic papules, usually on the lower legs and sometimes the arms or trunk of older adults. They are benign and may be treated for cosmetic or symptomatic reasons.

### **Student's Elbow**

Olecranon bursitis, an inflammation of the bursa over the point of the elbow, often related to repeated pressure or trauma.

### **Stuffy Nose**

Blocked or difficult nasal breathing caused by swollen tissue, mucus, allergy, infection, irritants, or structural narrowing.

### **Stuffy Or Runny Nose – Adult**

A stuffy or runny nose in an adult commonly results from viral infection, allergy, irritants, or sinus disease; persistent, severe, bloody, or one-sided symptoms need review.

### **Stump Pain**

Alternate name; see *Residual limb pain*.

### **Stunted Growth**

Stunted growth is height substantially below expected levels for age, often due to long-term undernutrition, chronic illness, infection, or other factors. Children with poor growth should be evaluated.

### **Stupor**

A state of reduced consciousness in which the patient is unresponsive to most environmental stimuli but can be temporarily aroused by vigorous, repeated stimulation (painful stimuli, loud shouting).

---

## Sturge-Weber Syndrome

A condition present from birth involving abnormal blood vessels in the skin and sometimes on the surface of the brain. It may cause a port-wine birthmark, seizures, weakness, headaches, developmental problems, or glaucoma, depending on which areas are affected.

## Stutter

Stuttering is a speech-fluency disorder involving repetitions, prolongations, or blocks in speech. Speech-language therapy can improve communication and confidence.

## Stuttering

A speech-flow problem involving repeated sounds or words, stretched sounds, pauses, or blocks when speaking. It is not caused by low intelligence; support may include speech therapy, communication strategies, and help with anxiety or teasing.

## St Vitus's Dance

An older name for Sydenham chorea, involuntary movements following group A streptococcal infection.

## Sty

Alternate name; see *Stye*.

## Stye

A tender, localized infection or inflammation of an eyelid gland, usually producing a painful red lump near the eyelid margin. It is also called a hordeolum.

## Styptics

Substances that help stop minor surface bleeding by tightening tissue, narrowing small blood vessels, or supporting clot formation. They are used for limited bleeding and do not replace emergency care for severe blood loss.

## Subacute

Developing or lasting longer than an acute condition but not as long as a chronic one. The exact time range varies with the disease.

## Subacute Bacterial Endocarditis

Older term for subacute infective endocarditis, a slowly developing infection of the heart's inner lining, usually involving a valve. It is most often caused by bacteria, but other organisms can also cause it. Symptoms may include fever, tiredness, weight loss, a new murmur, or complications from infected material travelling in the blood; see *Infective Endocarditis*.

## Subacute Combined Degeneration

Damage to parts of the spinal cord, usually from vitamin B12 deficiency. It can cause numbness, weakness, poor balance, and difficulty walking.

## Subacute Combined Degeneration Of The Cord

Damage to the spinal cord, usually caused by vitamin B12 deficiency, affecting nerve pathways for vibration, position sense, and movement. It may cause tingling, weakness, poor balance, or difficulty walking and can become permanent without treatment.

## Subacute Cutaneous Lupus Erythematosus (SCLE)

A photosensitive form of cutaneous lupus causing annular or papulosquamous red scaly lesions on sun-exposed skin. It may be associated with anti-Ro/SSA antibodies, medicines, or systemic lupus features and needs sun protection and medical evaluation.

## Subacute Rehabilitation

Rehabilitation provided after the early phase of illness or injury for people who still need nursing care and therapy but no longer need intensive hospital treatment. It may occur in a specialized facility or outpatient setting.

## Subacute Sclerosing Panencephalitis

A rare, progressive brain disease caused by measles virus remaining in the brain years after the original infection. It can cause behavior or learning changes, jerking movements, seizures, loss of coordination, and severe disability; prevention depends on measles



---

vaccination.

### **Subacute Thyroiditis**

Temporary inflammation of the thyroid, often after a viral illness, causing painful neck swelling, fever, and tiredness. Thyroid hormone levels may first be high and later low before returning to normal; treatment relieves pain and other symptoms.

### **Subaortic Stenosis**

Narrowing just below the aortic valve that obstructs blood leaving the heart. It may cause a heart murmur, chest pain, fainting, breathlessness, or damage to the aortic valve and sometimes requires surgery.

### **Subarachnoid**

Situated beneath the arachnoid, one of the thin coverings around the brain and spinal cord. The subarachnoid area contains protective fluid and blood vessels; bleeding there can cause a sudden severe headache and is an emergency.

### **Subarachnoid Cisterns**

Anatomical enlargements and pools of the subarachnoid space formed where the arachnoid membrane bridges over deep depressions between the brain and skull, filled with cerebrospinal fluid, including the cisterna magna, interpeduncular cistern, ambient cistern, and chiasmatic cistern.

### **Subarachnoid Haemorrhage**

Bleeding into the space around the brain containing cerebrospinal fluid, often causing sudden severe headache.

### **Subarachnoid Hemorrhage**

Bleeding into the space around the brain, often from a ruptured blood-vessel bulge. It typically causes a sudden, extremely severe headache and may cause vomiting, neck stiffness, confusion, weakness, or loss of consciousness.

### **Subarachnoid Space**

The space between two thin coverings of the brain and spinal cord, called the arachnoid and pia mater.

It contains cerebrospinal fluid and blood vessels and helps cushion the nervous system.

### **Subareolar Abscess**

A pus-filled infection beneath the areola, often associated with periductal inflammation and smoking, causing a painful lump, redness, drainage, or recurrent fistula.

### **Subchorionic Hemorrhage**

Bleeding between the chorion and the uterine wall caused by partial separation of the gestational sac membranes. It may produce vaginal bleeding in early or mid-pregnancy and is assessed according to bleeding, gestational age, and ultrasound findings.

### **Subclavian**

Relating to structures beneath the clavicle, especially the subclavian arteries and veins that supply or drain the upper limbs.

### **Subclavian Artery**

A major artery beneath the collarbone that supplies the arm and contributes blood flow to the neck, chest, and brain. Narrowing or blockage can cause arm pain, weakness, or symptoms from reduced blood flow to the brain.

### **Subclavian Steal Syndrome**

A condition in which narrowing of an artery supplying the arm redirects blood away from the back of the brain, especially during arm exercise. It may cause arm pain or weakness, dizziness, fainting, blurred vision, or imbalance. Treatment controls vascular risk factors and may restore blood flow when symptoms are significant.

### **Subclavian Vein**

The continuation of the axillary vein beyond the lateral border of the first rib. It joins the internal jugular vein behind the sternoclavicular joint to form the brachiocephalic vein.

### **Subclavian Vein Thrombosis**

A blood clot in the large vein beneath the collarbone.

---

It may be related to a central venous catheter or pacemaker lead, or to repeated strenuous arm activity and compression at the thoracic outlet (Paget-Schroetter syndrome). It can cause arm or neck swelling, pain, bluish discoloration, and pulmonary embolism.

### Subclinical

Present without clear symptoms or signs that a person or clinician can notice. A subclinical problem may be found through a blood test, scan, screening examination, or other test and may or may not later cause symptoms.

### Subclinical Disease

Disease that produces no obvious symptoms or signs but can be detected by screening, examination, or laboratory testing. It may remain mild or later become clinically apparent, depending on the condition.

### Subclinical Hypothyroidism

An early biochemical state of mild thyroid failure defined by an elevated serum thyroid-stimulating hormone (TSH) concentration in the presence of strictly normal circulating free thyroxine (free  $T_4$ ) and free triiodothyronine (free  $T_3$ ) levels. It represents a compensatory pituitary response that maintains euthyroid peripheral hormone concentrations. Although often asymptomatic and discovered incidentally on routine blood tests, patients may exhibit subtle symptoms (mild fatigue, cold intolerance, constipation), atherogenic dyslipidemia (elevated LDL cholesterol), and a heightened risk of progression to overt primary hypothyroidism, particularly when antithyroid peroxidase (anti-TPO) autoantibodies are present.

#### Alternate Names:

Mild Thyroid Failure; Subclinical Thyroid Failure; Compensated Hypothyroidism

### Subclinical Infection

An infection that produces no noticeable symptoms or signs but may still be detectable by testing and may sometimes be transmissible.

### Subconjunctival Hemorrhage

A small blood vessel broken beneath the clear surface of the eye, causing a sharply demarcated red patch that usually resolves without treatment.

#### Alternate Names:

Hyposphagma; Eye Capillary Bleed

### Subconscious

Mental processes outside immediate awareness; it is a psychological term rather than a specific diagnosis.

### Subcorneal Pustular Dermatitis

A rare recurring skin disorder causing shallow, non-infectious pus-filled blisters just beneath the outer skin layer, usually on the trunk or skin folds. It may resemble psoriasis and can be associated with blood or immune disorders.

### Subcu

Subcu is an abbreviation for subcutaneous, meaning beneath the skin; clear clinical instructions should spell out the route.

### Subculture

The transfer of microorganisms from one culture to fresh culture medium, used to maintain pure cultures, isolate colonies, increase viability, prepare standardized inocula, and perform identification/susceptibility testing. On solid media, subculture aims to obtain isolated single colonies from a mixed or stock culture.

### Subcutaneous

Subcutaneous means beneath the skin; injections by this route deliver medicine into the fatty tissue between skin and muscle.

### Subcutaneous Emphysema

Air trapped beneath the skin, often after chest injury, surgery, a punctured lung, or a breathing procedure. It may cause swelling and a crackling feeling; treatment addresses the air leak, and breathing difficulty or rapid spread requires urgent care.

### Subcutaneous Implantable

---

### Cardioverter-Defibrillator

A device placed under the skin that detects and treats dangerous fast heart rhythms with an electrical shock. Its wires run outside the heart and blood vessels; infection, inappropriate shocks, movement, or device failure are possible.

### Subcutaneous Injections

Administration of a medicine into the fatty tissue beneath the skin, commonly used for insulin, anti-coagulants, and selected biologic medicines.

### Subcutaneous Layer

The tissue layer beneath the dermis, made mainly of fat and connective tissue. It cushions deeper structures, insulates the body, stores energy, and contains larger blood vessels and nerves.

### Subcutaneous Tissue

The layer beneath the skin that connects it to deeper structures. It contains fat, connective tissue, larger blood vessels, and nerves and provides insulation, cushioning, and energy storage.

### Subdural

Located beneath the dura, one of the protective coverings around the brain and spinal cord. A subdural hematoma is bleeding in this space, often after head injury, and may cause headache, vomiting, confusion, weakness, seizures, or loss of consciousness. Significant bleeding needs urgent assessment.

### Subdural Effusion

Accumulation of usually clear fluid in the subdural space, often after trauma, surgery, infection, or cerebrospinal-fluid disturbance and sometimes causing headache or pressure effects.

### Subdural Haematoma

A collection of blood between protective coverings of the brain, usually after small veins tear during a head injury. It may develop suddenly or gradually and cause headache, confusion, drowsiness, weakness, seizures, or loss of consciousness.

### Subdural Hematoma

A collection of blood between two protective coverings of the brain, usually after small veins tear during a head injury. It can cause headache, confusion, weakness, seizures, or sleepiness and may require urgent treatment.

### Subdural Hemorrhage

Bleeding between two protective coverings of the brain, usually after tearing of small veins during a head injury. It may cause sudden or gradually worsening headache, confusion, drowsiness, weakness, seizures, or loss of consciousness and is diagnosed with brain imaging.

### Subendocardial Myocytes

Heart-muscle cells in the inner layer of the myocardium, which are especially vulnerable to reduced blood flow.

### Subependymal Giant Cell Astrocytoma

A usually slow-growing brain tumor that develops near a passage where fluid leaves the brain's cavities. It is strongly associated with tuberous sclerosis and can block fluid flow, causing headache, vomiting, or enlarged brain cavities.

### Subependymoma

A rare, usually benign slow-growing ependymal tumor, commonly arising in the ventricular system of adults. Symptoms result from obstruction of cerebrospinal-fluid pathways or incidental detection.

### Subfertility

Alternate name; see *Infertility*.

### Subgaleal Hematoma

A dangerous collection of blood within the loose areolar connective tissue space between the epicranial aponeurosis (galea aponeurotica) and the cranial periosteum, capable of crossing suture lines and expanding massively across the entire cranial vault, causing hypovolemic shock in neonates following vacuum-assisted delivery.

---

### **Subgaleal Hemorrhage**

A potentially life-threatening accumulation of blood in the space beneath the scalp's epicranial aponeurosis. It can expand rapidly in a newborn, particularly after vacuum-assisted delivery, and requires urgent recognition and treatment.

### **Subinvolution**

Slower-than-expected return of an organ to its usual size, especially the uterus after childbirth. It may cause prolonged or heavy bleeding, pelvic pain, or an enlarged tender uterus and needs assessment for retained tissue, infection, or other causes.

### **Subjacent**

An anatomical term describing a structure, tissue, or lesion located beneath another structure. It indicates relative position and does not necessarily mean that the two structures are touching; the relationship may be described on physical examination, imaging, or during surgery.

### **Subjective**

Based on a person's own experience or report rather than a directly measured finding. Pain, nausea, fatigue, dizziness, and mood are subjective symptoms, while temperature and blood pressure are objective measurements.

### **Sublimation**

A psychological defense process in which unacceptable impulses are redirected into socially acceptable activity; in chemistry it is solid-to-gas transition.

### **Sublingual**

Sublingual means beneath the tongue; medicines given this way can be absorbed through oral mucosa and may bypass much first-pass liver metabolism.

### **Sublingual Epidermoid Cyst**

A benign cyst lined by stratified squamous epithelium beneath the tongue mucosa, presenting as a slow-growing midline swelling that may require surgical excision if symptomatic.

### **Sublingual Gland**

One of the paired major salivary glands, located beneath the tongue. It produces mainly mucous saliva that enters the mouth through several small ducts and helps lubricate the oral cavity.

### **Subluxation**

Partial displacement of a joint in which some contact between the joint surfaces remains. It may cause pain, swelling, instability, limited movement, nerve injury, or repeated episodes after injury or disease.

### **Submandibular Gland**

One of the major salivary glands, located beneath the lower jaw. It makes much of the saliva present when a person is not eating and releases it into the floor of the mouth through a small duct.

### **Submucosa**

The supportive layer beneath the moist lining of organs such as the intestine, airway, and bladder. It contains blood vessels, nerves, glands, and connective tissue and helps nourish and anchor the lining.

### **Subphrenic Abscess**

A collection of pus beneath the diaphragm, usually caused by infection after abdominal surgery, a burst abdominal organ, or spread from nearby tissue. It may cause fever, upper-abdominal or shoulder pain, breathing discomfort, and sepsis and usually needs drainage and antibiotics.

### **Subq**

Subq is an abbreviation for subcutaneous, referring to tissue beneath the skin and the route used for some injections.

### **Subscapular Artery**

A branch of the axillary artery that supplies the shoulder blade region, chest wall, and nearby muscles. It contributes to the network of vessels supporting the shoulder and upper limb.

---

## Substance

A substance is a material with defined or variable chemical composition; medical effects depend on identity, dose, route, duration, and host factors.

## Substance Abuse

Legacy, nonspecific label; see *Substance Use Disorder*.

## Substance Abuse Disorder

Legacy label; see *Substance Use Disorder*.

## Substance Abuse Psychiatry

An older term for the clinical specialty concerned with substance use disorders and co-occurring mental-health conditions. Current terminology generally uses *Addiction Psychiatry* or *Addiction Medicine*.

## Substance Dependence

A state in which repeated substance use produces physiologic adaptation, so that reducing or stopping the substance may cause withdrawal. It is distinct from, but may occur with, substance-use disorder.

## Substance Intoxication

A temporary change in thinking, behavior, or body function caused by recent use of alcohol, a drug, or another substance. Effects depend on the substance, amount, timing, combinations, and the person's health.

## Substance Use

The consumption of alcohol, tobacco or nicotine, cannabis, prescription or nonprescription drugs, inhalants, or other psychoactive substances. Use may be occasional, prescribed, nonmedical, or associated with a substance use disorder; the term alone does not imply harmful use.

## Substance Use Disorder

A clinically significant pattern of alcohol, tobacco or nicotine, cannabis, prescription or nonprescription drug, inhalant, or other psychoactive-substance use that causes impaired control, social or occupational

impairment, risky use, or clinically significant distress. Features may include craving, unsuccessful efforts to reduce or stop use, substantial time spent obtaining or using the substance, failure to meet major responsibilities, continued use despite harm, tolerance, or withdrawal; tolerance and withdrawal are not required.

### Alternate Names:

Drug Addiction; Chemical Dependency; Substance Abuse Disorder

## Substance Use In Pregnancy

Use of alcohol, tobacco or nicotine, cannabis, opioids, cocaine, or other substances during pregnancy. Effects depend on the substance, dose, timing, route, and other factors and may include maternal complications, impaired fetal growth, preterm birth, or neonatal withdrawal.

## Substance Withdrawal

Physical or mental symptoms that occur after reducing or stopping heavy or prolonged substance use. Symptoms range from anxiety and nausea to seizures, severe confusion, dangerous blood-pressure changes, or breathing problems, depending on the substance.

## Substantia Nigra

A crescent-shaped basal ganglia midbrain structure containing dopaminergic neurons in the dorsal pars compacta (which project to the striatum and degenerate in Parkinson disease) and GABAergic output neurons in the ventral pars reticulata.

## Substernal Goiter

An enlarged thyroid gland that extends from the neck into the upper chest. It may press on the windpipe or esophagus, causing shortness of breath, cough, or difficulty swallowing. Imaging helps assess its size and position.

## Substrate

A substance on which an enzyme acts or a material used by a biochemical pathway.

---

## Subtalar Joint

The joint between the talus and heel bone beneath the ankle. It helps the foot tilt inward and outward.

## Subungual Exostosis

A benign osteocartilaginous outgrowth arising beneath the nail, usually of a toe or finger. It causes a painful firm subungual mass and nail deformity and is treated by excision when symptomatic.

## Succenturiat Lobe

An accessory lobe of the placenta, connected to the main placenta by blood vessels that traverse the membranes. If undetected after delivery, the succenturiat lobe may be retained in the uterus, causing postpartum hemorrhage or infection. Diagnosis: careful inspection of the placenta and membranes after delivery.

## Succenturiate Placenta

A succenturiate placenta has one or more accessory lobes separate from the main placenta, which can increase risk of retained tissue or fetal-vessel injury.

## Succinate Dehydrogenase

An enzyme inside mitochondria that helps cells make energy. It changes one small molecule into another while passing electrons through the energy-producing system; inherited or tumor-related changes can have clinical importance.

## Succinylcholine

A very fast, short-acting medicine that temporarily paralyzes skeletal muscles to help place a breathing tube or perform emergency procedures. It can dangerously raise potassium, stop breathing longer than expected, or trigger malignant hyperthermia in susceptible people.

## Succumb

To succumb means to die from or be overcome by a disease, injury, exposure, or other serious process.

## Succussion Splash

A splashing sound auscultated over the epigastrium

or thorax while gently shaking the patient's torso. In the abdomen, a splash heard more than three to four hours after a meal indicates delayed gastric emptying or gastric outlet obstruction; in the chest, it indicates hydropneumothorax.

### Alternate Names:

Hippocratic Succussion Splash

## Sucking Blister

A blister or worn area of skin in a newborn, usually on a hand, finger, forearm, or lip, caused by repeated sucking before or during feeding. It is usually harmless and heals on its own.

## Sucking Reflex

A primitive reflex present from birth, elicited by placing an object in the infant's mouth. The infant rhythmically sucks on the object. Coordinated sucking and swallowing typically develop by 34-36 weeks gestational age. Premature infants may lack coordinated suck-swallow, requiring gavage feeding.

## Sucrose

Common table sugar, a disaccharide made of glucose and fructose. Digestive enzymes split it into these simpler sugars for absorption.

## Sucrose Lysis Test

An older blood test once used to look for paroxysmal nocturnal hemoglobinuria, a disorder in which red blood cells are unusually vulnerable to destruction. Flow-cytometry testing has largely replaced it because it is more specific and reliable.

## Suction

Removal of fluid, secretions, blood, or other material using negative pressure. It may be used in airway care, surgery, wound care, or drainage.

## Suction Device

Equipment that creates negative pressure to remove fluid, blood, mucus, or other material. It usually includes a pump, tubing, collection container, and suction catheter or tip and is used in airway care, surgery, or wound drainage.



---

### Suction Tip

The end attachment of a suction device used to remove blood, fluid, secretions, or debris. Common types include the Yankauer tip for oropharyngeal suction and surgical suction tips such as the Frink tip.

### Sudden Cardiac Death

Unexpected death from a cardiac cause, usually occurring rapidly after symptom onset or without witnessed symptoms. Causes include coronary disease, cardiomyopathy, inherited arrhythmia syndromes, and bradyarrhythmia; prevention depends on the underlying risk.

#### Alternate Names:

SCD; Massive Fatal Arrhythmia

### Sudden Infant Death Syndrome

The unexplained sudden death of an infant younger than one year after a complete investigation, including medical history, death-scene review, and autopsy. Safe sleep on the back on a firm, clear surface and avoiding smoke reduce risk, but do not prevent every case.

### Sudden Infant Death Syndrome (SIDS)

Alternate name; see *Sudden Infant Death Syndrome*.

#### Alternate Names:

SIDS

### Sudden Natural Death Pathology

Postmortem evaluation of an unexpected death from natural disease, commonly involving coronary artery disease, myocardial infarction, cardiomyopathy, myocarditis, pulmonary embolism, epilepsy, or intracranial disease. The cause may remain indeterminate despite complete examination.

### Sudden Sensorineural Hearing Loss

A rapid loss of hearing caused by a problem in the inner ear or hearing nerve, often noticed on waking or over a few days. It may be accompanied by ringing or dizziness and needs urgent assessment because early treatment can improve recovery in some cases.

### Sudden Unexpected Death in Epilepsy

An unexpected death in a person with epilepsy that is not caused by drowning, injury, prolonged seizure, or another known condition. It is uncommon; controlling seizures, taking prescribed treatment, and discussing nighttime safety may reduce risk.

### Sudden Unexpected Death In Infancy

The sudden, unexpected death of an infant that requires investigation of the medical history, circumstances, death scene, and sometimes autopsy. The cause may be identified, or the death may meet the definition of sudden infant death syndrome.

### Sudden Unexpected Infant Death

The sudden and unexpected death of an infant younger than 1 year, whether or not the cause is known before investigation. Sudden infant death syndrome is the unexplained subtype after complete scene, history, and autopsy evaluation.

### Sudden Visual Loss

Rapid loss or marked reduction of vision in one or both eyes caused by retinal, optic-nerve, vascular, inflammatory, traumatic, or neurologic disease. It requires urgent localization and evaluation.

### Sudeck Atrophy

An acute post-traumatic or post-surgical manifestation of complex regional pain syndrome type I (CRPS-I) characterized by severe burning neuropathic pain, autonomic vasomotor and sudomotor dysfunction, localized edema, and rapid patchy radiographic osteopenia of the affected limb.

#### Alternate Names:

Sudeck's Dystrophy; Post-Traumatic Osteoporosis

### Sudek's Atrophy

An older name for complex regional pain syndrome with pain, swelling, autonomic change, and regional bone loss.

### Sudorifics

Medicines or other treatments that increase sweating. They have limited specific medical uses; ex-

---

cessive sweating caused by a treatment can lead to dehydration, overheating, or abnormal body salts.

### **Suffocation**

Inability to breathe enough because the airway is blocked, oxygen is unavailable, the chest is compressed, or another problem prevents normal breathing. It quickly causes unconsciousness and death without immediate rescue.

### **Sugammadex**

A medicine used during anesthesia to quickly reverse muscle paralysis caused by selected muscle-relaxing medicines. It binds those medicines so the body can remove them; allergic reactions and changes in heart rate are possible.

### **Sugar Substitutes**

Sugar substitutes provide sweetness with little or no sugar; types differ in energy, digestion, after-taste, and effects on glucose and gastrointestinal symptoms.

### **Sugar-Water Hemolysis Test**

An older laboratory test assessing red-cell resistance to hypotonic sugar solution, historically used in evaluating paroxysmal nocturnal hemoglobinuria and now largely replaced by flow-cytometric testing.

### **Suicidal Ideation**

Thoughts about dying or killing oneself, with or without a plan or intention to act. Any current plan, intent, access to a method, recent attempt, or inability to stay safe is an emergency requiring immediate support and professional help.

### **Suicidal Intent**

The degree to which a person expects or wants to die as a result of suicidal thoughts or behavior. It is assessed with the person's own account, planning, preparation, access to means, and other clinical information.

### **Suicidality**

The presence or risk of suicidal thoughts, intent,

plans, or behavior. The term describes a clinical domain rather than a diagnosis and requires assessment of immediacy, access to means, past behavior, protective factors, and safety.

### **Suicidal Plan**

A stated or considered method, timing, place, or other preparation for ending one's own life. A plan is an important risk indicator and requires immediate professional safety assessment.

### **Suicide**

Death caused by an intentional action to end one's life. Suicidal thoughts, plans, or behavior are urgent medical concerns; immediate support, safety measures, and professional assessment can help protect the person.

### **Suicide And Suicidal Behavior**

Thoughts, plans, attempts, or acts intended to cause one's own death; any current intent, plan, access to means, or inability to stay safe requires immediate emergency support.

### **Suicide Attempt**

An act in which a person intentionally tries to end their own life but does not die. A suicide attempt can cause serious injury even when the person did not expect to die and requires prompt medical and mental-health assessment.

### **Suicide Risk Assessment**

A clinical evaluation of suicidal thoughts, intent, plan, access to means, past attempts, self-harm, psychiatric and medical illness, substance use, protective factors, and immediate safety. It informs but does not replace ongoing clinical judgment and safety planning.

### **Sulcus**

A groove or shallow furrow on an organ or body surface, especially the brain or heart. Brain sulci separate raised areas and increase the cerebral cortex's folded surface area.

---

## Sulfa Drugs

A group of medicines, especially sulfonamide antibiotics, that block a chemical pathway bacteria need to grow. They treat selected infections but can cause allergic reactions and other side effects; the term may also refer to nonantibiotic sulfonamides.

## Sulfamethoxazole

A sulfonamide antibiotic used commonly with trimethoprim in selected infections.

## Sulfasalazine

An anti-inflammatory medicine used for inflammatory bowel disease and rheumatoid or other inflammatory arthritis. It changes immune activity and can cause nausea, rash, low blood counts, liver problems, or kidney problems.

## Sulfatide

A fatty substance containing sulfate found mainly in myelin, the insulating covering of nerve fibers. Abnormal buildup or deficiency can disrupt nerve function and occurs in some inherited metabolic disorders.

## Sulfite Oxidase Deficiency

A rare inherited metabolic disorder in which sulfite cannot be adequately converted to sulfate. Toxic sulfite and related compounds can cause neonatal seizures, feeding difficulty, abnormal muscle tone, developmental impairment, and progressive brain injury.

## Sulfite Sensitivity

Sulfite sensitivity is an adverse reaction to sulfite preservatives, sometimes causing wheeze, flushing, headache, or other symptoms, especially in people with asthma.

## Sulfonamide

A medicine containing a sulfonamide chemical group. Some are antibiotics that stop bacteria making folate, which they need to grow; others reduce inflammation or increase urine flow. Allergic reactions and other side effects are possible.

## Sulfonamides

Sulfonamides are drugs containing a sulfonamide group, including antibacterial agents that inhibit bacterial folate synthesis; allergic reactions, severe skin reactions, cytopenias, renal effects, and crystaluria are possible.

## Sulfonylurea

An oral diabetes medicine that makes the pancreas release more insulin. It can lower blood sugar but may cause dangerously low blood sugar, hunger, or weight gain.

## Sulfonylureas

Oral medicines for type 2 diabetes that stimulate the pancreas to release more insulin. They can lower blood glucose but may cause low blood sugar, hunger, or weight gain.

## Sulfur Colloid Scan

A nuclear-medicine scan in which a small radioactive tracer is used to show activity in the liver, spleen, bone marrow, or lymph nodes. It can help assess how these organs work or identify abnormal tissue distribution.

## Sulfuric Acid Poisoning

Severe poisoning from sulfuric acid, a highly corrosive chemical. It can cause deep burns of the skin, eyes, mouth, throat, and digestive tract and dangerous fumes; emergency care is required and vomiting must not be induced.

## Sulindac Overdose

Taking too much sulindac, a nonsteroidal anti-inflammatory drug, may cause nausea, vomiting, stomach bleeding, drowsiness, kidney injury, liver injury, seizures, or metabolic acidosis. A suspected overdose needs prompt medical or poison-control advice.

## Sulphonamides

A group of medicines, including some antibiotics, that block a chemical pathway bacteria need to grow. They are used for selected infections and other conditions; allergic reactions, interactions, and

---

blood or kidney problems can occur.

### **Sulphonylureas**

Oral medicines that stimulate pancreatic insulin release to lower blood glucose in type 2 diabetes.

### **Sumatriptan**

A migraine medicine that narrows certain blood vessels and changes nerve signaling to stop acute migraine or cluster headaches.

### **Summer Diarrhoea**

An older term for seasonal diarrheal illness, often referring to infectious diarrhea in children.

### **Sunburn**

An acute inflammatory injury of the skin caused by excessive ultraviolet radiation, resulting in redness, pain, warmth, and sometimes blistering or peeling.

### **Sundowning**

Late-day or evening worsening of confusion, agitation, wandering, or behavioral symptoms in some people with dementia or delirium, influenced by fatigue, light, routines, and unmet needs.

### **Sunflower Syndrome**

A rare epilepsy in which a person repeatedly turns toward bright light and waves a hand near the eyes, often triggering absence seizures. It usually begins in childhood and requires neurologic assessment and seizure treatment.

### **Sunken Chest**

Alternate name; see *Pectus excavatum*.

### **Sunken Fontanelle**

A noticeable inward depression of an infant's soft spot, which is a classic physical sign of moderate to severe dehydration or significant malnutrition requiring prompt fluid replacement.

#### **Alternate Names:**

Depressed Fontanelle; Sunken Soft Spot; Dehydration Sign in Infants

### **Sun Protection**

Measures that reduce exposure to ultraviolet radiation, including shade, protective clothing, sunglasses, and broad-spectrum sunscreen. They help prevent sunburn, premature skin aging, eye damage, and ultraviolet-related skin cancers.

#### **Alternate Names:**

UV Protection; Solar Defense

### **Sunscreens**

Topical products that absorb, scatter, or reflect ultraviolet radiation to reduce sunburn and cumulative ultraviolet skin damage. They may be chemical, mineral, or combined formulations and are commonly described by their broad-spectrum protection and sun-protection factor (SPF).

### **Sunstroke**

A life-threatening form of heat illness in which body temperature rises dangerously high and the nervous system is impaired.

### **Superantigen**

A substance made by some bacteria that activates many immune cells at once. This can cause an excessive release of inflammatory chemicals and contribute to illnesses such as toxic shock syndrome or some forms of scarlet fever.

### **Superficial**

Near the surface of the body or an organ, rather than deep inside it. The term may describe tissues, veins, burns, wounds, or infections that mainly affect the skin and tissue just beneath it.

### **Superficial Fascia**

The soft tissue layer just beneath the skin. It contains fat, small blood vessels, nerves, and connective tissue and helps insulate the body and allow the skin to move.

#### **Alternate Names:**

Subcutaneous Tissue; Hypodermis; Camper and Scarpa Fascia

---

### **Superficial Spreading Melanoma**

The most common cutaneous melanoma subtype, characterized initially by radial growth across the epidermis and an irregularly pigmented patch or plaque before invasion develops.

### **Superficial Thrombophlebitis**

Inflammation and a blood clot in a vein just beneath the skin, causing a tender, firm, red cord. It often improves with treatment, but a spreading clot or symptoms of deep-vein thrombosis require medical assessment.

### **Superimposed Preeclampsia**

Preeclampsia developing in a person who already has chronic hypertension. It may present with new or worsening proteinuria, thrombocytopenia, abnormal liver or kidney tests, neurologic symptoms, or other features of maternal organ dysfunction.

### **Superinfection**

A new infection that develops during or after treatment of a previous infection, often involving an antimicrobial-resistant organism or an organism not susceptible to the original therapy, such as *Clostridioides difficile* after antibiotic exposure.

### **Superior**

Above or toward the head in the body, or higher than another structure. For example, the neck is superior to the chest. The opposite direction is inferior, meaning below or toward the feet.

### **Superior Mediastinum**

The upper part of the mediastinum, above the level of the sternal angle. It contains the thymus, major blood vessels, trachea, esophagus, nerves, and lymph nodes.

### **Superior Mesenteric Artery**

A major artery branching from the abdominal aorta. It supplies blood to much of the small intestine and to parts of the large intestine, including the right colon and nearby transverse colon.

### **Superior Mesenteric Artery Syndrome**

Pressure on part of the small intestine where it passes between two major arteries, causing blockage. It may follow major weight loss and cause fullness, upper-abdominal pain, nausea, and vomiting; treatment restores nutrition and relieves the obstruction.

### **Superior Semicircular Canal Dehiscence**

A gap or very thin area in the bone covering one balance canal in the inner ear. Loud sounds or pressure may cause dizziness, imbalance, hearing changes, or hearing one's own voice unusually loudly; diagnosis uses symptoms, hearing tests, and imaging.

### **Superior Vena Cava**

A large vein in the chest that carries oxygen-poor blood from the head, neck, arms, and upper chest to the right side of the heart.

### **Superior Vena Caval Obstruction**

Narrowing or blockage of the superior vena cava, the large vein that returns blood from the head, neck, arms, and upper chest to the heart. Causes include a chest tumor, a blood clot, or pressure from a nearby structure. It may cause superior vena cava syndrome if blood return is sufficiently impaired.

#### **Alternate Names:**

SVCO; Superior Vena Cava Obstruction

### **Superior Vena Cava Syndrome**

A group of symptoms and signs caused by impaired blood flow through the superior vena cava, usually because it is compressed or blocked. It may cause swelling of the face, neck, or arms, visible neck veins, cough, or shortness of breath; severe or rapidly worsening symptoms need urgent assessment.

### **Supernumerary Digit**

An extra finger or toe present from birth, ranging from a small skin tag to a well-formed digit with bone and joints. It may occur alone or with a genetic syndrome and is evaluated for function, circulation, and associated anomalies.

---

## Supernumerary Nipple

An additional nipple or areolar structure along the embryonic milk line, usually harmless and sometimes mistaken for a mole. It may darken or enlarge with hormonal changes and generally needs treatment only if symptomatic or diagnostically uncertain.

## Supernumerary Nipples

Supernumerary nipples are extra nipples along the embryonic milk line, usually harmless but occasionally associated with underlying breast or urinary-tract developmental anomalies.

## Supernumerary Rib

An extra rib beyond the typical twelve pairs, most commonly a cervical rib above the first thoracic rib, which may compress the brachial plexus or subclavian vessel causing thoracic outlet syndrome.

## Supernumerary Teeth

Teeth that develop in addition to the usual number. They may remain hidden or appear in the mouth and can crowd, displace, or prevent nearby teeth from erupting; a common type develops between the upper front teeth.

## Supervised Exercise Therapy

A structured, monitored exercise program used to improve walking ability and reduce claudication symptoms in people with peripheral artery disease. The exercise plan is adjusted to the person's symptoms and functional capacity.

## Supination

Rotation of the forearm so that the palm faces anteriorly in anatomical position, or upward when the elbow is flexed. The radius and ulna become parallel.

## Supine

Lying flat on the back with the face and front of the body facing upward. This position is commonly used for examinations, imaging, and surgery. Staying in this position for a long time can increase the risk of pressure injuries; late in pregnancy, it may

also reduce blood return to the heart.

## Supine Hypotensive Syndrome

A drop in blood pressure that can occur when a pregnant person lies flat on the back, because the enlarged uterus reduces blood return to the heart. Dizziness, nausea, sweating, or faintness may improve by turning onto the side.

## Supine Position

Lying on the back with the face and front of the body upward. It is commonly used for examination, imaging, and surgery, but prolonged use can cause pressure injury and late pregnancy may reduce blood return to the heart.

## Supplemental Oxygen Therapy

Administration of oxygen at a concentration above that of room air to treat or prevent clinically important hypoxemia. Delivery method and oxygen target depend on the illness, baseline oxygenation, risk of carbon-dioxide retention, and monitoring results; excessive oxygen can also cause harm.

## Supplementation

Use of a vitamin, mineral, protein, fiber, or other nutrient preparation to correct a deficiency or meet a clinically identified increased requirement. The indication, dose, interactions, product quality, and risk of toxicity should be assessed before use.

## Supplements

Dietary supplements contain substances such as vitamins, minerals, herbs, or amino acids. They can have side effects and interactions, so patients should discuss their use with a healthcare professional.

## Supportive Care

Treatment that maintains essential body functions, relieves symptoms, prevents complications, and supports recovery while the underlying cause is treated.

## Suppository

A solid preparation inserted into the rectum, vagina, or urethra where it dissolves or melts.



---

## Suppression

Reduction or inhibition of a function, response, symptom, or process.

## Suppuration

The formation or discharge of pus, a thick fluid made from immune cells, germs, and damaged tissue. It usually occurs with infection and may form an abscess that needs drainage and treatment.

## Supracervical Hysterectomy

Hysterectomy in which the uterine body is removed while the cervix is left in place; it is also called subtotal or partial hysterectomy. The fallopian tubes and ovaries may be managed separately.

## Supracondylar Humerus Fracture

A break in the upper part of the elbow end of the arm bone, common in children after a fall on an outstretched hand. Swelling, pain, deformity, or inability to move the elbow may occur. Doctors must check circulation and nerves because severe fractures can damage an artery or cause dangerous swelling; some need urgent surgery.

### Alternate Names:

Gartland Supracondylar Fracture; Pediatric Elbow Fracture

## Supraglottic Airway Device

A breathing device placed above the vocal cords to help deliver oxygen and ventilation during anesthesia or emergencies. It can be inserted more quickly than a breathing tube but does not protect against aspiration as reliably.

## Supranuclear Ophthalmoplegia

Weakness or loss of control of eye movements caused by damage to brain pathways above the eye-movement nerves. It may make it difficult to look in certain directions.

## Supranuclear Palsy

Alternate name; see *Progressive supranuclear palsy*.

## Suprapubic

Located above the pubic bone in the lower abdomen. The term may describe pain or examination in this area, or a procedure such as inserting a catheter through the lower abdominal wall directly into the bladder when a tube cannot be passed through the urethra.

## Suprapubic Catheter

A tube placed through the lower abdominal wall directly into the bladder to drain urine. It may be used when urine cannot drain safely through the urethra or when longer-term bladder drainage is needed.

## Suprapubic Catheter Care

Care of a drainage tube that enters the bladder through the lower abdomen, including cleaning the site, keeping the tube unobstructed, securing it, and monitoring urine. Fever, increasing pain, leakage, bleeding, or no urine flow needs medical advice.

## Suprarenal Glands

Two small glands above the kidneys. Their outer layer makes hormones that regulate salt, blood pressure, stress, and metabolism; their inner layer makes adrenaline and related hormones.

## Suprascapular Nerve

A nerve from the brachial plexus supplying shoulder muscles that help lift the arm and rotate it outward.

## Suprascapular Neuropathy

Dysfunction of the suprascapular nerve from compression, traction, trauma, or an adjacent lesion. It may cause deep shoulder pain, weakness of abduction or external rotation, and wasting of the supraspinatus or infraspinatus muscles.

## Supraspinatus Muscle

The most frequently injured muscle of the rotator cuff, originating from the supraspinous fossa of the scapula and passing beneath the acromion to insert onto the superior facet of the greater tubercle of the humerus, responsible for initiating the first 15 degrees of shoulder abduction.

---

## Suprasternal Notch

The shallow dip at the upper border of the manubrium, between the inner ends of the collarbones.

## Supraventricular Tachycardia

A fast heart rhythm beginning above the ventricles, often starting and stopping suddenly, causing palpitations, dizziness, or weakness.

**Alternate Names:**

SVT

## Sural Nerve

A sensory nerve formed mainly from branches of the tibial and common fibular nerves. It supplies sensation to the posterolateral lower leg and lateral border of the foot.

## Suramin

An antiparasitic medicine used mainly for selected early forms of African trypanosomiasis. It requires specialist use because it can cause serious kidney, nerve, allergic, and other adverse effects.

## Surfactant

A substance that lowers surface tension. In the lungs, it coats the tiny air sacs and helps keep them open when a person breathes out; babies born very early may have too little and need treatment.

## Surgeon

A physician trained to perform operations and invasive procedures, and to evaluate and care for patients before and after surgery.

## Surgery

Treatment in which a clinician operates on tissues or organs to diagnose, remove, repair, replace, or improve a body structure. Risks and recovery depend on the procedure and the patient's health.

## Surgery For Pancreatic Cancer

An operation to remove a pancreatic tumor and sometimes nearby organs or lymph nodes when the cancer can be safely removed. It is major surgery

and may cause bleeding, infection, digestive problems, diabetes, or delayed stomach emptying.

## Surgery For Pilonidal Cyst

An operation to drain or remove a recurrent or complicated tunnel or cyst near the cleft of the buttocks. Wound care and keeping the area clean are important; infection, slow healing, and recurrence can occur.

## Surgical Abortion

A procedure that removes a pregnancy from the uterus to end a pregnancy or treat an early pregnancy loss. Common methods include vacuum aspiration and dilation and evacuation; the method depends on how far the pregnancy has progressed.

**Alternate Names:**

Vacuum Aspiration; Suction Curettage; Induced Abortion

## Surgical Drain

A tube or device placed during or after surgery to remove blood, pus, bile, urine, or other fluid. The amount and appearance of drainage are monitored; infection, bleeding, blockage, and tissue injury are possible.

## Surgical Drill

A powered instrument used to cut, shape, or make holes in bone during surgery. It requires careful control to avoid heat injury, damage to nearby tissues, infection, or unintended bone weakening.

## Surgical Excision

Cutting out a lesion, tumor, organ, or other unwanted tissue using an operation. The removed material may be examined under a microscope; bleeding, infection, scarring, damage to nearby structures, and incomplete removal are possible.

## Surgical Fixation

Stabilization of a bone, joint, or other structure with devices such as plates, screws, rods, wires, or external frames.

---

## **Surgical Forceps**

Handheld instruments used during surgery to hold, grasp, move, or manipulate tissue, sutures, dressings, or other materials. Their shape and pressure are selected to limit tissue injury.

### **Alternate Names:**

Surgical Grasping Forceps; Tissue Forceps

## **Surgical Hemostasis**

Control of bleeding during an operation using pressure, ligation, sutures, clips, electrosurgery, topical agents, vessel sealing, or other techniques. Effective hemostasis requires identification of the bleeding source and correction of coagulopathy when present.

## **Surgical Hemostat**

A clamping surgical instrument used to control bleeding by compressing a blood vessel or tissue.

### **Alternate Names:**

Surgical Hemostatic Forceps; Mosquito Clamp

## **Surgical Instruments**

Tools used during operations, including scalpels, scissors, forceps, clamps, retractors, needle holders, and devices controlling bleeding. Each instrument is designed for a specific task and must be cleaned, sterilized, and handled safely.

## **Surgical Margin**

The outer edge of tissue removed during an operation. A laboratory examination checks whether disease reaches that edge: a clear margin suggests no disease was seen there, while an involved margin may mean more treatment is needed.

## **Surgical Repair**

An operation that restores or rebuilds damaged tissue, an organ, or a body function. It may close wounds, reconnect structures, stabilize bones, repair blood vessels, or correct defects present from birth.

## **Surgical Retractor**

A surgical instrument used to hold tissues apart, giving doctors a clearer and safer view of the operating area.

### **Alternate Names:**

Tissue Retractor; Wound Retractor

## **Surgical Risk Stratification**

Estimation of the likelihood of perioperative complications using patient factors, urgency, procedure risk, functional capacity, and disease-specific assessment. It informs optimization, monitoring level, shared decisions, and postoperative planning but cannot eliminate uncertainty.

## **Surgical Safety Checklist**

A structured checklist used at defined points before anesthesia, before incision, and before leaving the operating room to verify patient identity, procedure, site, consent, equipment, specimens, and team communication.

## **Surgical Scissors**

Surgical instruments designed to cut tissue, sutures, dressings, or other materials. Different shapes and blade types are selected for delicate, deep, curved, or heavy-duty work.

### **Alternate Names:**

Surgical Scissors; Operating Shears

## **Surgical Site Infection**

Infection involving the incision, organ, or operative space within the surveillance period after surgery. It may be superficial incisional, deep incisional, or organ-space and requires appropriate culture, antimicrobial therapy, drainage, debridement, or reoperation when indicated.

## **Surgical Stapler**

A device that places metal or absorbable staples to close skin, join tissue, or seal a hollow organ during an operation. Incorrect placement can cause bleeding, leakage, narrowing, or injury and must be checked before completion.

## **Surgical Time-Out**

A deliberate team pause immediately before incision to verify patient identity, procedure, site, consent, allergies, antibiotics, imaging, implants, equipment, and anticipated critical events. It is a core surgical-

---

safety intervention.

### **Surgical Wound Classification**

Categorization of an operative wound as clean, clean-contaminated, contaminated, or dirty/infected according to the degree of microbial exposure and existing infection. The class helps estimate surgical-site infection risk and guide prophylaxis.

### **Surgical Wound Infection Treatment**

Management of infection in or around a surgical wound, including assessment of depth and severity, drainage or debridement when needed, wound care, and targeted antimicrobial therapy when indicated.

### **Surrogacy**

Surrogacy is an arrangement in which a person carries a pregnancy for intended parent or parents; medical, legal, ethical, and psychosocial requirements vary by jurisdiction.

### **Surrogate**

A substitute or proxy. In medical research, a surrogate outcome is a measurable change, such as a blood-test result, used to estimate a patient-important outcome such as fewer symptoms or longer survival.

### **Surrogate Decision-Maker**

A person authorized to make healthcare decisions for someone who lacks decision-making capacity, guided by the person's known values and preferences.

### **Surveillance**

The planned collection and review of health information to detect changes in disease, follow a known condition, or identify problems early. It may involve routine reports, active case finding, selected monitoring sites, or regular follow-up of an individual.

### **Surveillance Imaging**

Planned repeat imaging used to monitor disease, treatment response, or possible recurrence. The timing and type of scan depend on the condition, previous treatment, symptoms, risks, and expected

usefulness of results.

### **Survival Rate**

The percentage of people alive after a specified time from diagnosis or treatment, commonly measured at five years.

### **Survivorship**

The period of living during and after cancer treatment, together with the care provided during that time. It may include checking for return of cancer, treating late effects, supporting emotional health, improving quality of life, and managing other illnesses.

### **Survivorship Care Plan**

A written summary of cancer treatment, recommended follow-up, possible late effects, screening, medicines, and contacts for ongoing care. It helps the person and healthcare team coordinate care after treatment.

### **Susac Syndrome**

A rare disease in which inflammation and blockage of tiny blood vessels can affect the brain, eyes, and inner ears. It may cause confusion or headache, vision loss, and hearing loss; all features may not appear at once.

### **Susceptibility**

The likelihood of being affected by a disease, exposure, medicine, or microorganism. Susceptibility can reflect genetics, age, immunity, health conditions, previous exposure, dose, or the organism's drug resistance.

### **Susceptibility Testing**

Laboratory testing that shows which antibiotics are likely to stop a person's bacterial infection. The result helps clinicians choose treatment, especially when resistance is possible.

### **Susceptibility-Weighted Imaging**

An MRI technique that is especially sensitive to small veins, blood products, iron, and some calcium

---

deposits in tissue, particularly in the brain.

### Suspected Cause

A condition, exposure, or factor believed to explain a symptom or disease but not yet proven. Further history, examination, testing, or observation may support or reject the suspected cause.

### Suspension

A liquid medicine in which fine drug particles are spread through the liquid rather than dissolved. The container may need shaking so each dose contains the intended amount; settling, inaccurate measuring, and contamination can cause problems.

### Sustained Virologic Response

No detectable hepatitis C virus in the blood 12 or more weeks after antiviral treatment. This is considered a cure of that infection, but it does not protect against catching hepatitis C again.

### Suture

A stitch or material used to close or join tissue.

### Suture Anchor

A small implant that secures a tendon, ligament, or other soft tissue to bone. It may be made from metal, polymer, or bioabsorbable material and can fail through pullout, breakage, migration, or inflammatory reaction.

### SUV

Standardized uptake value, a number estimating how much radioactive tracer is concentrated in tissue during a PET scan. Clinicians compare it with surrounding tissue and clinical findings to assess tumors or inflammation.

### SVC Obstruction

Blockage or compression of the superior vena cava, the large vein returning blood from the head, neck, arms, and upper chest to the heart. It may cause facial or arm swelling, prominent neck veins, cough, headache, or breathing difficulty.

### SVC Syndrome

Blockage or compression of the superior vena cava, the large vein returning blood from the upper body to the heart. It may cause swelling of the face, neck, or arms, visible neck veins, cough, or breathing difficulty.

### SVT

Alternate name; see *Supraventricular tachycardia*.

### Swab

A small absorbent stick used to collect a sample from the skin, throat, nose, wound, or another body site. The sample can be tested for infection or other conditions, and correct collection affects the result.

### Swallowing

The movement of food, liquid, or saliva from the mouth through the throat and esophagus into the stomach. Problems may cause coughing, choking, food sticking, or material entering the airway.

#### Alternate Names:

Deglutition; Ingestion

### Swallowing Assessment

An evaluation of how safely and effectively a person moves food, liquid, or saliva from the mouth to the stomach. It may include observation, X-ray or camera testing, and recommendations about exercises, posture, or food and drink textures.

### Swallowing Chalk

Ingestion of chalk, which is usually low in toxicity in small amounts but can cause choking, constipation, or gastrointestinal upset; repeated eating may indicate pica.

### Swallowing Difficulty

Dysphagia, or impaired movement of food or liquid from the mouth to the stomach, caused by oropharyngeal, esophageal, neuromuscular, structural, or functional disorders.

### Swallowing Problems

Difficulty moving food, liquid, or saliva safely from

---

the mouth to the stomach; dysphagia can cause choking, aspiration, weight loss, or dehydration and may need evaluation.

### **Swallowing Soap**

Ingestion of soap, which commonly causes mouth or stomach irritation, nausea, or diarrhea; concentrated or large exposures can injure tissue or affect breathing.

### **Swallowing Sunscreen**

Ingestion of sunscreen, usually causing mild gastrointestinal upset but potentially causing choking or aspiration depending on amount and product; breathing difficulty or persistent vomiting requires urgent help.

### **Swallowing Syncope**

Fainting triggered by swallowing, usually through an abnormal reflex that slows the heart or lowers blood pressure; recurrent episodes require evaluation for rhythm and structural heart disease.

### **Swallowing Therapy**

Assessment and treatment to make swallowing safer and more effective. A speech-language therapist may teach exercises, positioning, food-texture changes, and strategies to reduce aspiration into the airway.

### **Swallow Study**

An X-ray test that records a person swallowing food or liquid mixed with barium. It shows how the mouth and throat work, whether material enters the airway, and whether changes could make swallowing safer.

### **Swan-Ganz Catheter**

A thin tube passed through the right side of the heart into a lung artery to measure pressures and estimate how much blood the heart pumps. It is used selectively in seriously ill patients because complications are possible.

### **Swan-Ganz Catheterization**

A procedure in which a thin catheter is passed through the right side of the heart into a lung artery to measure heart pressures and blood flow. It is used selectively in critically ill patients because bleeding, infection, rhythm problems, or vessel injury can occur.

### **Swan Neck Deformity**

A characteristic joint deformity of the hand in which there is hyperextension of the proximal interphalangeal (PIP) joint accompanied by compensatory flexion of the distal interphalangeal (DIP) joint. It is classically seen in advanced rheumatoid arthritis, systemic lupus erythematosus, and chronic tendon imbalance.

### **Swan-Neck Deformity**

A finger shape problem where the middle joint bends backward and the fingertip curls down, commonly caused by rheumatoid arthritis.

### **Sweat**

Fluid produced by sweat glands, mainly to cool the body as it evaporates. Sweating can also increase with exercise, fever, stress, hormones, or some medicines.

### **Sweat Chloride Test**

A test that measures salt levels in sweat to help diagnose cystic fibrosis. A high result supports the diagnosis, but it is interpreted with symptoms and may be repeated or followed by genetic testing because other factors can affect results.

### **Sweat Electrolytes Test**

A sweat electrolytes test measures salts, especially chloride, in sweat and is commonly used to help diagnose cystic fibrosis.

### **Sweat Gland**

A skin gland that produces sweat, which contributes mainly to temperature regulation and skin surface function.



---

## Sweat Glands

Eccrine and apocrine glands that produce sweat or related secretions.

## Sweat Gland Tumor

A growth arising from cells of a sweat gland. Most are benign skin lumps, but some are cancerous and may grow, ulcerate, bleed, or spread. Examination and biopsy determine the type; complete removal is often used, with wider treatment and follow-up for cancerous tumors.

## Sweating

Production of sweat by eccrine and apocrine glands for heat regulation and other responses; excessive, absent, or asymmetric sweating can indicate medication, autonomic, endocrine, or neurologic disease.

## Sweating And Body Odor

Excess perspiration or noticeable odor produced when skin bacteria break down sweat; it may be influenced by hygiene, diet, medicines, hormones, or disease.

### Alternate Names:

Diaphoresis and Bromhidrosis; Perspiration and Osmidrosis

## Sweetened Beverages

Sweetened beverages contain added sugars and can increase calorie intake, dental caries, obesity, and diabetes risk when consumed frequently.

## Sweet's Syndrome

An inflammatory skin disorder causing sudden fever and painful red or purple raised patches or nodules. It may follow infection, result from a medicine, or occur with inflammatory disease or cancer, especially blood cancers.

## Sweet Syndrome

An inflammatory skin disorder characterized by sudden fever and tender red or violaceous plaques or nodules containing many neutrophils. It may be associated with infection, medicines, inflammatory disease, or cancer.

## Swelling

Abnormal enlargement caused by fluid accumulation, inflammation, bleeding, infection, tumor, or tissue injury; location, onset, pain, and associated symptoms identify the likely cause.

## Swimmer's Ear

Swimmer's ear, also called otitis externa, is inflammation or infection of the skin lining the outer ear canal. It often develops when moisture remains in the canal, allowing bacteria or fungi to grow. Scratching, cotton swabs, or other irritation can also damage the canal's protective skin. Symptoms include ear pain, itching, redness, swelling, drainage, and pain when the outer ear is touched.

### Alternate Names:

Otitis Externa; Tropical Ear

## Swimmer's Itch

An itchy rash caused by an allergic reaction to microscopic parasites released by infected snails in freshwater or saltwater.

### Alternate Names:

Cercarial Dermatitis; Schistosome Dermatitis

## Swimming Pool Granuloma

A slowly developing skin infection caused by a water-dwelling bacterium after exposure to an aquarium, swimming pool, or natural water. It can cause persistent bumps or sores, often spreading along the arm or hand, and needs specific antibiotic treatment.

## Swine Flu

Alternate informal term; see *Influenza A Virus*.

## Swollen Ankles And Swollen Feet

Enlargement of the ankles or feet caused by injury, venous insufficiency, lymphatic disease, medication, pregnancy, or systemic fluid retention from heart, kidney, or liver disease. Sudden one-sided swelling with pain, redness, warmth, or breathlessness may indicate thrombosis or infection and requires urgent assessment.

### Alternate Names:

Peripheral Dependent Edema; Pedal Edema

---

## Swollen Glands

Lymph nodes that are larger than usual, often because the immune system is responding to infection or inflammation. They may be tender, but hard, fixed, enlarging, or unexplained nodes require medical assessment.

### Alternate Names:

Lymphadenomegaly; Adenopathy

## Swollen Gums

Pathologic enlargement, edema, or hypertrophy of the gingival tissue. Etiologies include plaque-induced gingivitis, periodontal disease, acute oral infections, hormonal fluctuations during pregnancy, nutritional deficiencies (such as scurvy), leukemia, or drug-induced gingival hyperplasia from medications like calcium channel blockers, phenytoin, and cyclosporine.

### Alternate Names:

Gingival Swelling; Gingival Hyperplasia; Gingival Enlargement; Enlarged Gums

### Alternate Names:

Gingival Hyperplasia; Gingival Enlargement

## Swollen Lymph Glands

Enlarged lymph nodes, usually caused by infection or inflammation and less often by cancer. They commonly occur in the neck, armpits, or groin; persistent, hard, fixed, or unexplained swelling needs medical assessment.

### Alternate Names:

Lymphadenopathy; Lymph Node Enlargement

## Swollen Lymph Nodes

Swollen lymph nodes are enlarged immune glands, commonly from infection but sometimes from inflammation, medication, or cancer; persistent, hard, or generalized nodes need evaluation.

### Alternate Names:

Lymphadenopathy; Adenopathy; Bubo

## Swollen Lymph Nodes Entry

Alternate name; see *Swollen Lymph Nodes*.

## Swollen Tongue

Enlargement of the tongue caused by injury, infection, allergy, inflammation, bleeding, or another condition; sudden breathing difficulty is an emergency.

## Swyer Syndrome

A form of pure 46,XY gonadal dysgenesis resulting from mutations in the *SRY* gene or related sex-determining pathways, presenting as a phenotypic female with normal external female genitalia, streak gonads, absent puberty, and primary amenorrhea, carrying a high risk of gonadoblastoma in the streak gonads.

### Alternate Names:

46,XY Pure Gonadal Dysgenesis

## Sycosis Barbae

Long-lasting inflammation or infection of hair follicles in the beard area, usually involving bacteria and ingrown hairs. It can cause painful bumps, pus, crusting, scarring, and hair loss and needs appropriate treatment.

## Sydenham Chorea

A delayed neurologic complication of group A streptococcal infection causing involuntary jerky movements, hypotonia, emotional lability, and coordination problems, usually in children.

## Sydenham's Chorea

A delayed immune complication of group A streptococcal infection, usually in children. It causes involuntary, irregular movements and may cause clumsiness, muscle weakness, emotional changes, or difficulty with speech and daily tasks.

## Symblepharon

Scar tissue that joins the inside of the eyelid to the eye's surface, usually after a severe burn, inflammation, or eye disease. It can restrict eye movement, cause dryness or pain, and may need specialist treatment.

---

## Sympathectomy

Surgical interruption or destruction of sympathetic nerves to reduce abnormal vasoconstriction, sweating, or pain in selected conditions.

## Sympathetic

Relating to the part of the autonomic nervous system that prepares the body for activity or stress by changing heart rate, blood-vessel tone, sweating, breathing, and digestion. In everyday language, it can also mean showing concern for another person.

## Sympathetic Ganglion

A small group of nerve cells outside the brain and spinal cord that relays automatic signals to blood vessels, sweat glands, and internal organs. These signals help control heart rate, blood pressure, digestion, and body responses to stress.

## Sympathetic Nervous System

The autonomic division that prepares the body for activity or stress by increasing heart rate and altering vascular, airway, and digestive function.

## Sympatholytic

A medicine or treatment that reduces adrenaline-like nerve activity. It may lower blood pressure, slow the heart, or reduce excessive sweating, but the effects and risks depend on the specific medicine and require appropriate prescribing.

## Sympathomimetic Drugs

Medicines that imitate the body's adrenaline-like nerve activity, often by stimulating alpha or beta receptors. They may open airways, raise blood pressure, or reduce congestion but can cause fast heartbeat, anxiety, or dangerous effects.

## Sympathomimetic Syndrome

Alternate name; see *Sympathomimetic Toxidrome*.

## Sympathomimetic Toxicity

Excess catecholamine activity causing agitation, mydriasis, diaphoresis, tachycardia, hypertension, hyperthermia, chest pain, seizures, stroke, or rhab-

domyolysis. Unlike anticholinergic toxicity, sweating is usually prominent; benzodiazepines, active cooling, ECG monitoring, and treatment of ischemia, hypertension, and complications are central.

## Sympathomimetic Toxidrome

A poisoning pattern caused by excessive adrenergic stimulation, producing agitation, sweating, dilated pupils, rapid heart rate, high blood pressure, elevated temperature, tremor, or seizures.

## Symphysis

A joint where two bones are joined by a pad of tough cartilage. The pubic symphysis connects the front pelvic bones and normally allows slight movement, especially during pregnancy and childbirth.

## Symphysis-Fundal Height

The distance from the top of the pubic bone to the top of the uterus, measured through the abdomen during pregnancy. From about 24 to 36 weeks, the measurement often roughly matches the number of weeks of pregnancy. A larger or smaller result may lead to an ultrasound to check fetal growth, fluid, or the pregnancy dates.

### Alternate Names:

Fundal Height Measurement; SFH

## Symphysis Pubis

The fibrocartilaginous joint connecting the left and right pubic bones at the front of the pelvis.

## Symptom

A change or problem felt or reported by a patient that may suggest illness, such as pain, nausea, or breathlessness.

## Symptomatic

Having symptoms caused by a disease or condition; symptomatic treatment relieves those symptoms without necessarily treating the underlying cause.

## Symptomatic Treatment

Care aimed at relieving symptoms such as pain, fever, nausea, itching, or cough without directly

---

removing the underlying cause. It may be used alone or alongside treatment for the disease itself.

### Synaesthesia

Synaesthesia is a neurological trait in which stimulation of one sense or mental process automatically produces an additional sensory experience, such as seeing colors when hearing sounds.

### Synapse

A connection where one nerve cell communicates with another nerve cell, muscle, or gland. Signals cross the small gap using chemicals or electrical changes to control movement, sensation, thought, and body functions.

### Synapse Assembly

The developmental process by which neurons form organized connections with other neurons or target cells, using cell-adhesion molecules and signaling proteins.

### Synapsins

A family of neuronal proteins that help organize synaptic vesicles and regulate the release of neurotransmitters.

### Synaptic Cleft

The very small gap between a presynaptic neuron and its target cell across which neurotransmitters travel.

### Synaptic Membrane

The specialized membrane at a neuronal junction, including the neurotransmitter-releasing presynaptic membrane and receptor-containing postsynaptic membrane.

### Synaptic Plasticity

The ability of connections between nerve cells to become stronger, weaker, or structurally different in response to activity. It supports learning, memory, adaptation, and some changes in chronic pain.

### Synaptic Transmission

Communication between nerve cells or between a nerve cell and another target across a synapse. Electrical signals trigger release of chemical messengers, which bind receptors and change the receiving cell's activity.

### Synaptic Vesicle

A small membrane-bound sac in a nerve ending that stores neurotransmitters and releases them into the synaptic cleft when the neuron is activated.

### Synaptonemal Complex

A temporary protein structure that pairs matching chromosomes during egg or sperm formation. It helps chromosomes exchange genetic material accurately before cells divide and is important for normal reproductive development.

### Synaptosome

A sealed nerve-terminal structure produced when brain tissue is mechanically disrupted; it is used to study neurotransmitter release and synaptic function.

### Synaptotagmin

A calcium-sensing protein on nerve-cell storage sacs that helps trigger rapid release of chemical nerve signals. It is essential for communication between nerve cells and between nerves and muscles.

### Synarthrosis

A joint with little or no movement, such as a skull suture or tooth anchored in its socket, in which bones are joined by fibrous tissue or cartilage.

### Synchondroses

Cartilage-based joints in which bones are joined by hyaline cartilage and allow little or no movement. Growth plates and the joint between the first rib and the breastbone are examples.

### Synchronized Cardioversion

Delivery of an electrical shock timed with the heart's electrical cycle to restore a more organized rhythm.

---

It treats selected fast rhythms; sedation, clot prevention, monitoring, and energy settings depend on the rhythm and patient.

**Alternate Names:**

Direct Current Cardioversion; Electrical Cardioversion; R-Wave Synchronized Shock

**Syncope**

A brief loss of consciousness and muscle control caused by a temporary reduction in blood flow to the brain, followed by spontaneous recovery. Common causes include reflex fainting, a sudden blood-pressure drop, dehydration, and heart disease; prolonged confusion may suggest another cause.

**Alternate Names:**

Passing Out; Temporary Loss of Consciousness; Swoon

**Syndactyly**

Congenital joining or webbing of two or more fingers or toes. It may occur alone or with a genetic syndrome and sometimes requires reconstructive surgery.

**Syndecan**

A group of proteins on the cell surface that help cells attach to their surroundings and receive signals. They contribute to tissue development, healing, and responses to injury.

**Syndesmophyte**

A thin, bony growth that forms along the edges of the vertebrae, often in ankylosing spondylitis, and may bridge adjacent vertebrae.

**Syndesmoses**

Fibrous joints in which two bones are held together by strong ligaments or a sheet of connective tissue. They allow little movement; the joint between the lower leg bones near the ankle is one example.

**Syndrome**

A group of symptoms, signs, or test findings that commonly occur together. A syndrome may have several possible causes and does not always name

one specific disease.

**Syndrome Of Inappropriate ADH Secretion**

Excessive antidiuretic hormone activity causing water retention and low blood sodium despite no major swelling or dehydration. It may result from lung disease, medicines, cancer, or brain disorders and can cause confusion or seizures.

**Syndrome of Inappropriate Antidiuresis (SIAD)**

Excessive vasopressin activity causes retention of free water, inappropriately concentrated urine, and usually low serum sodium despite no appropriate physiologic stimulus. Causes include medicines, lung or brain disease, cancer, and postoperative stress; severe hyponatremia can cause seizures or coma.

**Syndrome Of Inappropriate Antidiuretic Hormone**

Excessive release or effect of antidiuretic hormone causing impaired free-water excretion, low serum sodium, and inappropriately concentrated urine in the absence of an appropriate physiologic stimulus.

**Syndrome Of Inappropriate Antidiuretic Hormone Secretion**

Excessive activity of the water-retaining hormone vasopressin causes the body to retain water and lowers the blood sodium level. Causes include medicines, lung disease, cancer, and brain disorders; severe cases can cause confusion or seizures.

**Alternate Names:**

SIADH; Schwartz-Bartter Syndrome

**Syndrome X**

Alternate name; see *Metabolic syndrome*.

**Synechia**

An abnormal band of scar-like tissue joining surfaces that should remain separate. In the eye, it may attach the iris to the lens or cornea and interfere with fluid drainage or vision.

---

## Synechiaie

Abnormal bands of scar-like tissue joining surfaces that should remain separate. They may occur in the eye, uterus, nose, or other sites and can restrict movement, block passages, or impair function.

## Synergist

A muscle, drug, or factor that works together with another to produce or enhance an effect.

## Synesthesia

Synesthesia is a stable perceptual phenomenon in which stimulation of one sensory pathway automatically produces an experience in another, such as seeing colors with sounds.

## Synostoses

Fusions between two bones in which bone tissue joins them together. They may be present from birth or develop later and can limit growth or movement depending on their location.

## Synostosis

Fusion of two bones by bone tissue, present from birth or developing later. It may be harmless, but fusion in the skull, forearm, or other important area can restrict growth or movement.

## Synovectomy

An operation to remove part or all of an inflamed lining inside a joint. It may reduce pain and swelling in selected cases of arthritis or repeated joint inflammation and can be done through small cuts or open surgery.

## Synovial Biopsy

A synovial biopsy removes a small sample of the joint lining to investigate infection, inflammatory arthritis, crystal disease, or an unusual joint lesion.

## Synovial Bursa

A small, fluid-filled sac near a joint that reduces rubbing between moving tissues such as tendons, muscles, skin, and bone. Irritation or inflammation of a bursa causes bursitis, pain, and swelling.

## Synovial Chondromatosis

A usually noncancerous condition in which the joint lining forms many small cartilage nodules that may break loose. It can cause pain, swelling, catching, and limited movement, often in the knee, hip, or elbow.

## Synovial Cyst

A synovial cyst is a fluid-filled sac arising from or near a joint or tendon sheath, sometimes compressing nearby nerves and causing pain or weakness.

## Synovial Fluid

A slippery fluid inside a freely movable joint that reduces friction and helps nourish cartilage. Changes in its amount or contents can occur with injury, infection, bleeding, crystals, or arthritis.

## Synovial Fluid Analysis

Testing fluid removed from a swollen or painful joint. The laboratory examines its appearance, cells, crystals, and germs to help distinguish injury, arthritis, gout, bleeding, and joint infection; a very hot painful joint needs urgent assessment.

## Synovial Joint

A freely movable joint in which the articulating bones are separated by a fluid-filled joint cavity enclosed by a capsule.

## Synovial Joints

Freely movable joints in which the connecting bone surfaces are separated by a small fluid-filled space and enclosed by a joint capsule. Examples include the knee, hip, shoulder, and elbow.

## Synovial Membrane

The thin lining inside a freely moving joint. It makes synovial fluid, which reduces rubbing and helps nourish the smooth joint surfaces; inflammation of this lining causes swelling and pain.

## Synovial Reaction To Implanted Prostheses

Inflammation or tissue change around an artificial



---

joint caused by wear particles, loosening, corrosion, infection, or an immune reaction. Pain, swelling, or reduced function after replacement needs evaluation to identify the cause.

### Synovial Sarcoma

A cancer of soft tissue that often develops near a joint in an arm or leg, especially in adolescents and young adults. Diagnosis requires tissue testing; treatment may include surgery, radiation, or chemotherapy.

### Synoviocytes

Synoviocytes are cells lining the joint synovium that produce lubricating fluid and inflammatory mediators and help maintain or damage joint tissue.

### Synovitis

Inflammation of the lining inside a joint, causing swelling, warmth, pain, or reduced movement. It may follow injury or occur with arthritis, gout, autoimmune disease, or infection; a hot, very painful joint needs urgent assessment.

### Synovium

Tissue lining a joint or tendon sheath that makes lubricating fluid and helps support smooth movement.

### Synovography

An imaging test in which contrast is placed inside a joint to outline its capsule, lining, or possible tear. It is used selectively; MRI, ultrasound, or other scans are often preferred for joint problems.

### Synthetic Androgen

A synthetic androgen is a manufactured compound that activates androgen receptors and may mimic testosterone; therapeutic and anabolic use can suppress gonadal function and cause acne, mood effects, infertility, liver injury, or cardiovascular harm.

### Synthetic Biology

Synthetic biology combines biology and engineering to design or modify biological systems. It has po-

tential medical uses but also requires careful safety and ethical oversight.

### Syphilis

A sexually transmitted infection caused by *Treponema pallidum* that progresses through primary, secondary, latent, and tertiary stages and can affect the nervous system, eyes, heart, or fetus.

### Syphilis Serology

Blood tests used to screen for syphilis and monitor response to treatment. A reactive result usually needs confirmation with a different test and interpretation alongside symptoms, examination, and treatment history.

#### Alternate Names:

Non-Treponemal Antibodies; RPR and VDRL Tests

### Syphilitic Alopecia

Non-scarring hair loss occurring as a manifestation of secondary syphilis, presenting with diffuse thinning or patchy, moth-eaten alopecia of the scalp, eyebrows, or beard. It is thought to result from direct spirochaetal invasion of hair follicles during the spirochaetemic phase.

### Syphilitic Aseptic Meningitis

Syphilitic aseptic meningitis is inflammation of the meninges caused by *Treponema pallidum*, producing headache, neck stiffness, cranial-nerve symptoms, or other neurologic findings.

### Syphilitic Gumma

A destructive lump or area of inflammation caused by long-standing untreated syphilis. It can affect the skin, bones, liver, brain, or other organs and may form a firm mass or ulcer; appropriate antibiotic treatment can prevent further damage.

### Syringe

A medical device with a hollow barrel, movable plunger, and tip that may connect to a needle, catheter, or other fitting. It is used to draw up, measure, inject, or remove fluids. Common types include disposable, oral, insulin, tuberculin, and safety syringes.

---

**Alternate Names:**

Medical Syringe

**Syringe Drivers**

Small pumps that push medicine from a syringe continuously or at programmed times. They provide controlled delivery when precise dosing is needed, including pain relief, symptom control, or medicines given under the skin.

**Syringocystadenoma Papilliferum**

A usually harmless sweat-gland skin growth, often appearing as a small bump or plaque on the scalp or face. It may develop within a birthmark and can resemble other lesions, so diagnosis may require examination or biopsy.

**Syringoma**

A syringoma is a benign tumor of sweat-duct cells, usually appearing as small skin-colored or yellowish papules around the eyelids or elsewhere.

**Syringomyelia**

A fluid-filled cavity that develops inside the spinal cord. It can slowly damage the cord and cause pain, weakness, loss of pain or temperature sensation, stiffness, or problems with bladder, bowel, or sexual function.

**Syrinx**

A fluid-filled cavity within the spinal cord, usually related to disturbed movement of the fluid around the brain and spinal cord. It can damage nerve tissue and cause pain, weakness, stiffness, or loss of sensation; a syrinx in the cord is called syringomyelia.

**Syrup**

A syrup is a viscous liquid preparation, often sweetened, used as a medicine vehicle or oral dosage form; concentration and ingredients determine safe dosing.

**Systematic Desensitization**

A behavioral therapy technique that pairs gradual, planned exposure to a feared situation with relaxation or other coping skills. It is used mainly for

phobias and anxiety and is adapted to the person's response.

**Systemic**

Affecting the whole body or relating to an entire system rather than a localized site.

**Systemic Antifungals**

Antifungal medicines taken orally or given intravenously for infections that are extensive, invasive, refractory, or unsuitable for topical treatment. Choice and duration depend on the organism, site, severity, drug interactions, organ function, and local resistance patterns.

**Systemic Chemotherapy**

Systemic chemotherapy uses medicines that travel through the bloodstream to kill or slow cancer cells throughout the body.

**Systemic Circuit**

The circulation that carries oxygenated blood from the left ventricle through the body and returns deoxygenated blood to the right atrium.

**Systemic Dermatoses**

Skin manifestations that arise as secondary signs of internal medical diseases, such as endocrine disorders, autoimmune conditions, internal malignancies, or nutritional deficiencies.

**Alternate Names:**

Cutaneous Manifestations of Systemic Disease; Systemic Skin Signs

**Systemic Disease**

A disease that affects the whole body or several organs rather than one local area. Examples include autoimmune disease, widespread infection, hormonal disorders, and some cancers.

**Systemic Effects**

Effects involving the whole body rather than one local area. A medicine, infection, toxin, or disease may produce systemic cardiac, neurologic, kidney, liver, blood, or other effects.

---

## Systemic Exertional Intolerance Disease

Alternate name; see *Myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)*.

## Systemic Inflammatory Response Syndrome

A widespread inflammatory reaction that may follow infection, injury, burns, pancreatitis, or surgery. It can cause abnormal temperature, heart rate, breathing rate, or white blood-cell count and may progress to organ failure.

## Systemic Juvenile Idiopathic Arthritis

A childhood inflammatory disease causing repeated high fevers, a temporary pink rash, and swollen or painful joints. It can also enlarge lymph nodes, liver, or spleen and may affect other organs, requiring specialist monitoring and treatment.

## Systemic Lupus

Systemic lupus erythematosus is an autoimmune disease that can affect skin, joints, kidneys, blood, nervous system, lungs, and heart. Symptoms fluctuate, and treatment is tailored to organ involvement with sun protection, monitoring, and immunomodulatory medicines.

### Alternate Names:

Systemic Lupus Erythematosus; SLE

## Systemic Lupus Erythematosus

A long-term autoimmune disease in which the immune system mistakenly attacks the body's own tissues and causes inflammation. It can affect the skin, joints, kidneys, heart, lungs, brain, and blood cells. Symptoms vary widely and may include fatigue, fever, joint pain or swelling, rashes, hair loss, mouth sores, or swelling. The disease often has periods of increased activity called flares and periods of fewer symptoms called remission; inflammation can cause organ damage in some people.

## Systemic Lupus Erythematosus (SLE) and Pregnancy

SLE can affect fertility, kidneys, blood counts, blood vessels, placenta, and fetal growth, while pregnancy can trigger disease activity. Preconception plan-

ning, compatible medicines, antibody testing, blood-pressure and kidney monitoring, and maternal-fetal surveillance are important.

## Systemic Mastocytosis

A disorder in which too many mast cells build up in the skin, bone marrow, digestive tract, or other organs. The cells can release chemicals causing flushing, itching, faintness, stomach symptoms, bone problems, or severe allergic reactions.

## Systemic Scleroderma

An autoimmune disease that causes tight, thickened skin and may scar blood vessels and internal organs. The esophagus, lungs, kidneys, heart, fingers, and other areas can be affected, so care depends on the organs involved.

## Systemic Sclerosis

An autoimmune disease in which inflammation, blood-vessel changes, and excess scar-like tissue can affect the skin and internal organs. The skin may become tight and hard; the digestive tract, lungs, kidneys, and heart may also be affected. Raynaud phenomenon, in which fingers or toes change color in response to cold or stress, is common.

## Systemic Sclerosis-Associated Arthritis

Joint pain or inflammation occurring in systemic sclerosis. Stiffness, swelling, or reduced movement may occur, and symptoms can overlap with other connective-tissue diseases.

## Systemic Therapy

Treatment that travels through the bloodstream or otherwise affects the whole body. Examples include chemotherapy, hormone treatment, targeted treatment, and immunotherapy; it differs from local treatment such as surgery or focused radiation.

## Systemic Vascular Resistance

The opposition to blood flow in the body's arteries, controlled mainly by how narrow or wide the small arteries are. It helps determine blood pressure and the work required of the heart.

---

## Systemic Vasculitis

Inflammation of blood vessels that may involve large, medium, or small vessels and impair blood flow or cause tissue injury. Examples include ANCA-associated vasculitis and IgA vasculitis.

### Alternate Names:

Systemic Vasculitis; Generalized Vasculitis

## Systemic Venous

Systemic venous refers to veins returning deoxygenated blood from the body to the right side of the heart, excluding the pulmonary circulation.

## Systole

The part of each heartbeat when a heart chamber contracts and pushes blood forward. Ventricular systole sends blood to the lungs and body, while arterial pressure rises and the heart's valves coordinate flow.

## Systolic

Systolic means relating to contraction of the heart, especially the pressure in arteries when the ventricles eject blood.

## Systolic Anterior Motion

Movement of the mitral valve toward the path where blood leaves the heart during each heartbeat. It can partly block blood flow and cause the valve to leak, especially in hypertrophic cardiomyopathy. Symptoms may include breathlessness, chest pain, dizziness, or fainting; treatment depends on the cause and severity.

### Alternate Names:

SAM; SAM of the Mitral Valve

## Systolic Murmur

An abnormal heart sound auscultated during ventricular systole (between the first and second heart sounds, S1 and S2). It may be midsystolic/ejection (as in aortic or pulmonary stenosis) or holosystolic/pansystolic (as in mitral or tricuspid regurgitation, or ventricular septal defect).

## Systolic Murmurs

Extra heart sounds heard while the heart is contracting. They may result from fast blood flow, a narrowed heart valve, or another structural problem; their significance depends on the sound and the person's symptoms.

## Systolic Pressure

The maximum arterial pressure during ventricular contraction, recorded as the upper number of a blood-pressure reading.

## Sézary Syndrome

An uncommon form of skin lymphoma in which abnormal immune cells are found in the skin and blood. It usually causes widespread red, scaly, itchy skin and enlarged lymph nodes and requires specialist care.

## Sacroccygeal Chordoma

A malignant tumor arising from remnants of the embryonic notochord at the base of the spine. It may cause tailbone pain, a mass, bowel or bladder symptoms, or nerve problems and is assessed with imaging and biopsy.

## SAD (Seasonal Affective Disorder)

Alternate name; see *Seasonal Affective Disorder*.

## SAM-e

Abbreviation; see *S-Adenosylmethionine*.

## Sarcoid Of Boeck

Historical name for sarcoidosis, an inflammatory disease in which clusters of immune cells form in organs such as the lungs, lymph nodes, skin, or eyes.

## SARS (Severe Acute Respiratory Syndrome)

Alternate name; see *Severe Acute Respiratory Syndrome*.

## Scarring Bullosa

A group of blistering disorders in which repeated inflammation and blistering heal with scarring, po-

---

tentially affecting skin, mucous membranes, eyes, mouth, or the airway. Diagnosis and treatment depend on the specific disorder.

### **Schereshevskii-Turner Syndrome**

Alternate name; see *Turner Syndrome*.

### **Secondary Psychosis**

Psychotic symptoms caused or contributed to by a medical illness, medicine, substance, or neurologic disorder rather than a primary psychotic disorder. New hallucinations, delusions, or confusion require urgent assessment for reversible causes.

### **Severe Hearing Loss**

Hearing loss that substantially limits access to speech or environmental sound. It may be conductive, sensorineural, or mixed and may require hearing aids, cochlear implantation, communication support, or treatment of an underlying cause.

### **Skin Rash**

A visible change in the skin, such as redness, scaling, bumps, blisters, or discoloration. Causes include infection, allergy, inflammation, medicines, heat, and systemic disease; rapid spread, skin pain, blistering, fever, or mucosal involvement needs prompt care.

### **Skin Tags**

Soft, usually harmless skin growths attached by a narrow stalk, often in skin folds. They can catch on clothing or become irritated; a changing, bleeding, painful, or atypical lesion should be examined before removal.

### **Sleep Epilepsy**

Epilepsy in which seizures occur predominantly during sleep or transitions between sleep and wakefulness. Events may be subtle or involve movements, vocalizations, confusion, or injury and require neurologic evaluation.

### **Slipped Disc**

A common lay term for a herniated or prolapsed intervertebral disk, in which disk material extends

beyond its normal boundary and may irritate a nearby nerve, causing back or neck pain, numbness, or weakness.

### **Small Lymphocytic Lymphoma**

A usually slow-growing B-cell non-Hodgkin lymphoma that is closely related to chronic lymphocytic leukemia. It commonly causes enlarged lymph nodes and is diagnosed with tissue examination and immunophenotyping.

### **Spastic Dysphonia**

A voice disorder caused by involuntary spasms of the muscles used for speaking, also called spasmodic dysphonia. The voice may sound strained, broken, breathy, or weak; assessment distinguishes it from structural and neurologic causes.

### **Spermatocele (Epididymal Cyst)**

A usually benign, fluid-filled cyst arising in the epididymis near the testicle. It may cause a painless scrotal lump or heaviness and is distinguished from tumors, hydrocele, infection, and other causes by examination and ultrasound.

### **Spherocytic Anemia**

Anemia caused by fragile, sphere-shaped red blood cells that are removed early by the spleen. It is often inherited as hereditary spherocytosis and may cause jaundice, gallstones, fatigue, or an enlarged spleen.

### **Spinal Arachnoiditis**

Chronic inflammation and scarring of the arachnoid membrane around spinal nerves. It may cause persistent burning pain, tingling, weakness, or bowel and bladder problems after infection, surgery, trauma, bleeding, or chemical irritation.

### **Spinal Chordoma**

A malignant notochord-derived tumor arising along the spine, often in the sacrum. It can produce progressive pain, a mass, weakness, numbness, or bowel and bladder dysfunction and requires specialist imaging and treatment.



---

### Spinal Meningitis

Inflammation or infection of the membranes surrounding the spinal cord, often occurring with meningitis involving the brain. Fever, severe headache, neck or back stiffness, confusion, rash, or neurologic changes require emergency evaluation.

### Spinal Muscular Atrophy (SMA)

Alternate name; see *Spinal Muscular Atrophy*.

### Splenomegaly With Rheumatoid Arthritis

Enlargement of the spleen occurring with rheumatoid arthritis, classically in Felty syndrome and sometimes with low white blood-cell counts. Infection risk and other causes of spleen enlargement require assessment.

### Stevens Johnson Syndrome

Alternate spelling; see *Stevens-Johnson Syndrome*.

### Subcortical Dementia

Cognitive decline associated predominantly with disease of structures beneath the cerebral cortex, often causing slowed thinking, executive dysfunction, mood change, gait problems, and impaired attention. Causes include small-vessel disease and some neurodegenerative disorders.

### Subcortical Ischemic Vascular Disease

Chronic small-vessel disease causing reduced blood flow and tissue injury in deep brain structures. It may contribute to slowed thinking, gait difficulty, mood symptoms, urinary problems, and vascular dementia.

### Subendocardial Sclerosis

Fibrous thickening of the inner lining of the heart, often involving the ventricles or valves. Its significance depends on the cause and may include effects on filling, conduction, or cardiac function.

### Sucrose Intolerance

Inability to digest or absorb sucrose adequately, often because of congenital or acquired sucrase-

isomaltase deficiency. Eating sucrose may cause bloating, cramps, gas, and diarrhea; diagnosis and dietary management should avoid unnecessary restriction.

### Suppurative Arthritis

A joint infection producing pus and inflammation, usually caused by bacteria. It can rapidly destroy cartilage and cause fever, severe joint pain, swelling, and limited movement, requiring urgent drainage and antimicrobial treatment.

### Symmastia

An abnormal connection or loss of the normal separation between the breasts across the midline of the chest, sometimes occurring after breast surgery or reconstruction. Management depends on symptoms, anatomy, and cause.

### Symptomatic Porphyria

Porphyria causing clinical attacks or ongoing symptoms from abnormal heme production. Manifestations may include severe abdominal pain, neurologic or psychiatric symptoms, blistering or sun-sensitive skin, or liver complications depending on the type.

### Spider Veins

Small visible red, blue, or purple superficial veins, usually on the legs or face. They are generally harmless but may be associated with venous insufficiency; sudden pain, swelling, skin ulceration, or one-sided change requires assessment.

### Sphingomyelinase Deficiency

Deficiency of acid sphingomyelinase caused by variants in the *SMPD1* gene. It causes Niemann-Pick disease types A, A/B, or B, with lipid accumulation affecting the spleen, liver, lungs, brain, or other organs.

### Spielmeyer-Vogt Disease

An older name for neuronal ceroid lipofuscinosis, a group of inherited lysosomal disorders causing progressive loss of vision, seizures, movement problems, and cognitive decline.



---

### **Spielmeyer-Vogt-Batten Syndrome**

Alternate historical name for juvenile neuronal ceroid lipofuscinosis, commonly called Batten disease. It usually begins in childhood with vision loss followed by seizures, developmental regression, movement problems, and dementia.

### **Synergistic Necrotizing Fasciitis**

A rapidly progressive necrotizing soft-tissue infection caused by mixed aerobic and anaerobic organisms acting together. Severe pain, swelling, skin discoloration, fever, toxicity, or rapid progression requires immediate surgery, antibiotics, and critical care.

### **Systemic Aleukemic Reticuloendotheliosis**

A historical name for a leukemic reticuloendothelial disorder, most often referring to the aleukemic form of hairy-cell leukemia. It may cause low blood counts, an enlarged spleen, infections, fatigue, or bleeding and is diagnosed by blood, marrow, and immunophenotypic studies.



# T

## **T3**

Triiodothyronine, the more active of the main thyroid hormones. It helps control how quickly the body uses energy and supports normal growth, development, heart activity, and body temperature.

## **T3RU Test**

An older blood test that indirectly estimated how much thyroid hormone could bind to blood proteins. It was once used with other thyroid tests but has largely been replaced by direct free-thyroxine measurements.

## **T3 Test**

A blood test measuring triiodothyronine, used with thyroid-stimulating hormone and thyroxine to evaluate thyroid function, particularly suspected hyperthyroidism.

## **T4**

Thyroxine, the main hormone released by the thyroid gland. Much of it is changed into the more active T3 in body tissues. T4 helps control energy use, growth, development, and body temperature.

## **Tabes Dorsalis**

A late nervous-system complication of untreated syphilis that damages sensory nerves and parts of the spinal cord. It can cause severe brief pains, numbness, poor balance, loss of reflexes, and bladder or sexual problems.

## **Tablet**

A solid medicine made to be swallowed, chewed, dissolved in the mouth, or used in another specified

way. Some tablets release medicine immediately, while others are coated or designed to release it slowly.

## **Tablet Splitting**

Dividing a tablet into smaller parts. Extended-release, enteric-coated, and some specialized tablets may release medicine incorrectly or become unsafe when split.

## **TAB Vaccine**

An older vaccine against typhoid fever and paratyphoid A and B. It has largely been replaced by newer typhoid vaccines and does not protect against every cause of fever or every type of paratyphoid infection.

## **Tacalcitol**

A vitamin-D-like medicine applied to skin affected by psoriasis or selected scaling disorders. It helps regulate skin-cell growth and inflammation but can irritate skin or raise calcium if overused.

## **Tache Noire**

A dark scab or area of dead tissue at the site of a tick bite. It can occur with some spotted-fever infections and is interpreted with fever, rash, travel, and exposure history.

## **Tachyarrhythmia**

An abnormally fast heart rhythm, which may arise from the atria, ventricles, or the heart's conduction system. Symptoms can include palpitations, dizziness, chest discomfort, shortness of breath, or fainting; fainting or severe symptoms require urgent

---

assessment.

### **Tachycardia**

A heart rate faster than expected for a person's age and situation. It may be a normal response to exercise, stress, fever, or dehydration, or may result from a heart or other medical problem.

#### **Alternate Names:**

Rapid Heartbeat; Fast Heart Rate

### **Tachycardia-Bradycardia Syndrome**

A variant manifestation of sick sinus syndrome characterized by alternating paroxysmal supraventricular tachyarrhythmias (commonly atrial fibrillation or atrial flutter) and periods of severe sinus bradycardia or prolonged sinus pauses upon arrhythmia termination.

#### **Alternate Names:**

Tachy-Brady Syndrome

### **Tachycardia-Induced Cardiomyopathy**

Weakening and dilation of the heart muscle caused by a persistent or frequently recurring rapid heart rhythm, often improving when the arrhythmia is controlled.

### **Tachykinin**

A family of small signaling proteins, including substance P, that help regulate pain, inflammation, smooth-muscle activity, and communication between nerve cells.

### **Tachyphrenia**

Tachyphrenia is rapid flow of thoughts or accelerated mental activity, often described in association with mania, stimulants, or other psychiatric states.

### **Tachyphylaxis**

A rapid loss of a medicine's effect after repeated doses over a short time. It differs from tolerance, which usually develops more slowly; changing treatment may restore the effect under medical supervision.

### **Tachypnea**

Abnormally rapid breathing. It may occur with exercise or anxiety, but can also signal fever, low oxygen, pneumonia, blood clot in the lung, heart failure, severe infection, pain, or acid buildup.

### **Tachypnoea**

Breathing faster than expected for a person's age and situation. It may result from exercise, fever, pain, anxiety, lung disease, low oxygen, heart failure, severe infection, or acid buildup and must be interpreted with other findings.

### **Tachysystole**

More than five uterine contractions in 10 minutes, averaged over 30 minutes, during labor. It can reduce the baby's oxygen supply, especially when the fetal heart rate is abnormal, and may follow medicines used to start or strengthen labor.

#### **Alternate Names:**

Uterine Tachysystole; Excessive Uterine Activity

### **TACO**

Transfusion-associated circulatory overload is breathing difficulty caused by too much fluid entering the circulation during or soon after a blood transfusion. It can cause lung fluid, high blood pressure, and swelling, especially in people with heart or kidney disease or those receiving rapid or large transfusions.

### **TAC Regimen**

A chemotherapy combination of docetaxel, doxorubicin, and cyclophosphamide, used for selected breast cancers; it can cause infection risk, nausea, hair loss, heart effects, and other toxicities.

### **Tacrolimus**

A medicine that suppresses immune activity. It is used to prevent rejection after an organ transplant and, in different preparations, to treat eczema and some other inflammatory skin conditions; infection risk and other side effects require monitoring.

---

## Tactile

Relating to the sense of touch, including feeling pressure, texture, vibration, temperature, or pain. Touch signals travel through nerves to the brain; reduced or abnormal touch sensation may result from nerve, spinal-cord, brain, or skin disease.

## Tactile Agnosia

Inability to recognize a familiar object by touch even though basic sensation is intact. It usually results from a problem in areas of the brain that interpret sensory information.

## Tactile Fremitus

Vibrations felt on the chest when a person speaks. Changes in these vibrations can help clinicians identify lung problems such as pneumonia, fluid around the lung, a collapsed lung, or reduced air movement.

## Tactile Sense

The ability to perceive touch, pressure, vibration, texture, and related mechanical stimuli through receptors in the skin and deeper tissues.

## Tadalafil

Tadalafil is a phosphodiesterase-5 inhibitor used for erectile dysfunction, benign prostatic hyperplasia, and pulmonary hypertension; headache, flushing, hypotension, priapism, and dangerous nitrate interactions are important risks.

## Taenia

A group of tapeworms that includes *Taenia solium*, the pork tapeworm, and *Taenia saginata*, the beef tapeworm. Adult worms can live in the intestine; *T. solium* eggs can also cause cysts in body tissues.

## Taenia Saginata

*Taenia saginata* is the beef tapeworm acquired from undercooked infected beef; intestinal infection may cause abdominal symptoms or passage of motile segments but does not usually cause human cysticercosis.

## Taeniasis

Intestinal infection caused by adult *Taenia* tapeworms. It is usually acquired by eating undercooked beef or pork containing larval cysts. It may cause abdominal discomfort, nausea, diarrhea, or passage of worm segments. Swallowing *Taenia solium* eggs causes cysticercosis in tissues, which is a different infection.

## Taenia Solium

A pork tapeworm that infects people who eat undercooked pork or swallow its eggs. Adult worms live in the intestine, while eggs can cause cysticercosis, including dangerous infection of the brain.

## Tailbone Trauma

Injury to the coccyx, the small bone at the bottom of the spine, usually after a fall or prolonged pressure. It causes pain when sitting, standing, or passing stool. Most injuries are bruises and improve over time; persistent or severe pain may indicate a fracture or another problem.

## Tail Of The Epididymis

The lower part of the epididymis where mature sperm are stored before entering the vas deferens.

## Takayasu Arteritis

An immune-mediated large-vessel vasculitis affecting the aorta and its major branches. Inflammation can narrow, block, or weaken these arteries, causing limb pain with use, weak or unequal pulses, blood-pressure differences, bruits, fever, fatigue, vision or neurological symptoms, or organ ischemia.

## Takayasu Disease

A large-vessel vasculitis that primarily affects the aorta and its major branches. It may cause limb claudication, weak or unequal pulses, blood-pressure differences, bruits, fever, fatigue, or arterial narrowing and aneurysms. Imaging and inflammatory assessment guide immunosuppressive and vascular treatment.

## Takayasu'S Arteritis

Inflammation of the aorta and its major branches

---

that can narrow or weaken the arteries. It may cause weak pulses, unequal blood pressure, arm or leg pain with use, fever, fatigue, or high blood pressure.

### **Takotsubo Cardiomyopathy**

A temporary weakening of the heart muscle, often after intense emotional or physical stress. It can resemble a heart attack with chest pain and ECG changes, but the heart arteries are usually not blocked; most people recover with appropriate care.

### **Takotsubo Syndrome**

A temporary stress-related weakening of the heart muscle, also called stress cardiomyopathy or broken-heart syndrome. It can resemble an acute heart attack with chest pain, breathlessness, and ECG changes despite no obstructing coronary artery lesion; complications can include heart failure, arrhythmia, or shock.

#### **Alternate Names:**

Stress Cardiomyopathy; Broken-Heart Syndrome

### **Talactoferrin Alfa**

Talactoferrin alfa is a recombinant lactoferrin-related immunomodulatory protein studied as a treatment or adjunct in cancer and infectious disease.

### **Talaporfin Sodium**

Talaporfin sodium is a photosensitizer used or studied in photodynamic therapy, where light activation produces local cytotoxic effects.

### **Talar Dome Lesion**

An osteochondral defect or fracture involving the articular hyaline cartilage and subchondral bone of the superior articular surface (dome) of the talus, typically provoked by traumatic ankle inversion sprains, causing chronic deep ankle pain and mechanical catching.

#### **Alternate Names:**

Osteochondritis Dissecans of the Talus; OCD of the Talus

### **Talar Tilt Test**

An orthopedic stress examination maneuver used to assess the structural integrity of the calcaneofibular ligament (CFL) of the ankle; with the patient seated and knee flexed, the examiner inverts the talus within the ankle mortise; talar tilt exceeding 10 to 15 degrees or 5 degrees greater than the contralateral uninjured side indicates a complete CFL tear.

#### **Alternate Names:**

Inversion Stress Test of Ankle; CFL Integrity Test

### **Talc**

A soft mineral; inhalation of talc dust can cause talcosis, and contaminated injected talc can injure vessels and lungs.

### **Talcosis**

Lung damage caused by breathing talc dust or, rarely, injecting talc-containing crushed medicines. It can cause scarring, cough, and shortness of breath; severity depends on the amount and type of exposure.

### **Talcum Powder Poisoning**

Harmful effects from swallowing or inhaling talcum powder. Swallowing usually causes mild stomach upset, but inhaling it can injure the lungs and make breathing difficult.

### **Talipes**

A deformity of the foot and ankle present at birth or acquired later. It may affect position, flexibility, walking, and shoe fit, and treatment depends on severity, cause, age, and function.

### **Talipes Calcaneovalgus**

A foot deformity in which the heel is turned outward and the ankle is dorsiflexed so the foot points upward; flexible newborn cases often improve, while rigid cases need evaluation.

### **Talipes Equinovarus**

Clubfoot, a birth condition in which the foot turns inward and downward and is difficult to move into



---

a normal position. Early treatment usually uses gentle stretching, repeated casts, and a brace to support walking development.

### **Tall Stature**

Tall stature means height substantially above the expected range for age and sex and may be familial, endocrine, genetic, or associated with another condition.

### **Talofibular Ligament Injury**

Sprain or tear of one or more ligaments connecting the fibula to the talus, especially the anterior talofibular ligament on the outer ankle. It commonly follows an inversion injury and may cause pain, swelling, bruising, or instability.

### **Talus**

The ankle bone between the shinbones and the heel and foot bones. It transfers body weight through the ankle and has a limited blood supply, so some fractures can damage the bone and cause delayed healing or bone death.

### **Tamiflu**

Tamiflu is the brand name for oseltamivir, an antiviral medicine used to treat or help prevent influenza. It works best when started early and does not replace flu vaccination.

### **Tamoxifen**

A medicine that blocks or changes oestrogen effects in breast tissue. It treats and helps prevent some breast cancers, but can increase the risk of blood clots and, rarely, cancer of the uterine lining.

### **Tamoxifen Citrate**

The citrate salt of tamoxifen, a selective estrogen-receptor modulator used to treat or prevent some breast cancers; blood clots and endometrial cancer are important uncommon risks.

### **Tamoxifen Toxicity**

Adverse pharmacological effects of the selective estrogen receptor modulator tamoxifen used in breast

cancer treatment, including hot flashes, increased risk of deep vein thrombosis and pulmonary embolism, and endometrial hyperplasia/carcinoma via agonistic estrogenic effects on the uterus.

### **Tampon**

A material inserted into the vagina to absorb menstrual blood; prolonged use can rarely lead to toxic shock syndrome.

### **Tamponade**

Pressure on an organ caused by accumulated fluid, blood, or gas. Cardiac tamponade occurs when fluid or blood around the heart prevents it from filling and pumping properly.

### **Tamsulosin Hydrochloride**

The hydrochloride salt of tamsulosin, an alpha-1 blocker that relaxes prostate and bladder-neck muscle to improve urinary symptoms from benign prostatic enlargement.

### **Tangier Disease**

A rare inherited disorder causing extremely low HDL cholesterol and buildup of cholesterol in tissues. It may cause enlarged yellow-orange tonsils, enlarged liver or spleen, nerve symptoms, or clouding of the cornea; cardiovascular risk varies and specialist testing is needed.

### **Tanner Staging**

A five-stage scale describing physical changes during puberty, including breast or genital development and pubic hair. It helps clinicians assess whether development is absent, typical, early, delayed, or progressing unevenly.

### **Tantalum**

A corrosion-resistant metal used in some surgical implants and medical devices; exposure risk depends on the form, dose, and route.

### **Tapentadol**

A prescription pain medicine with opioid-like and nerve-signal effects, used for selected moderate or

---

severe pain. It can cause sleepiness, constipation, dependence, seizures, and dangerously slow breathing, especially with alcohol or other sedatives.

### Tapeworm

A parasitic flatworm with a segmented body that may live in the intestine or form larval cysts in tissues. People acquire different species by eating infected meat or fish or by ingesting eggs from contaminated sources. Symptoms and complications depend on the species and site.

### Tapeworm Infection

Infection caused by parasitic tapeworms. *Taenia* species may live as adult worms in the intestine, while some tapeworm larvae can form cysts in tissues.

### Tapotement

A massage technique using rhythmic tapping, patting, or percussion. It can worsen injury or bleeding in some situations.

### Tapping

Repeated light striking or percussion used during an examination, treatment, or test. It may help assess reflexes, pain, body sounds, or fluid and can also be used in chest physiotherapy or other treatments.

### TAPSE

Alternate name; see *Tricuspid Annular Plane Systolic Excursion*.

### TAPVR

Alternate name; see *Total anomalous pulmonary venous return*.

### Tarantula Spider Bite

A bite from a tarantula that usually causes local pain, redness, and swelling. Some species also release irritating hairs that can affect the skin or eyes.

### Tardieu Spots

Tiny pinpoint areas of bleeding under the skin or on internal surfaces. They can form when small blood

vessels rupture during severe pressure or oxygen deprivation, but their presence alone cannot establish the cause or timing of death.

### Alternate Names:

Subpleural Ecchymoses; Petechial Hemorrhages

### Tardive Dyskinesia

Repeated, involuntary movements that may develop after long-term use of some antipsychotic medicines. They often affect the mouth, tongue, face, arms, or legs and may continue after the medicine is reduced or stopped.

### Tardive Dystonia

A persistent or recurrent movement disorder caused by prolonged exposure to dopamine-receptor-blocking medicines. Sustained twisting postures or repetitive muscle contractions may affect the neck, trunk, face, or limbs.

### Target Cell

A cell that responds to a particular hormone, medicine, toxin, antigen, or other chemical signal because it has the needed receptor or other binding site.

### Target Cells

Abnormal erythrocytes (codocytes) characterized on peripheral blood smear by a central dense area of hemoglobin surrounded by a clear ring and an outer perimeter of hemoglobin (resembling a bullseye target), resulting from an excess of membrane surface area relative to intracellular volume, characteristic of hemoglobinopathies (HbC, HbS), thalassemia, post-splenectomy states, and obstructive liver disease.

### Alternate Names:

Codocytes; Bullseye Erythrocytes

### Target-Controlled Infusion

A computer-controlled pump that adjusts the rate of an intravenous medicine to reach and maintain a planned concentration in the blood or at its site of action. It is used mainly during anesthesia and requires close monitoring.

---

## Targeted Radiation Therapy

Radiation directed precisely at a tumor or other abnormal tissue to damage its DNA while limiting exposure to nearby healthy tissue. It may be delivered from outside the body or through an implanted source.

## Targeted Temperature Management

Careful control of body temperature after cardiac arrest or another critical illness, usually in an unconscious patient. It aims to limit further brain injury and requires continuous monitoring while the person is cooled or kept from developing fever.

## Targeted Therapies For Cancer

Cancer medicines designed to act on specific features of cancer cells or their surroundings. They may be more selective than traditional chemotherapy, but can still cause important side effects.

## Targeted Therapy

Treatment that acts on a specific change or feature helping cancer cells grow, survive, or spread. Testing the tumor may identify a target and predict benefit. These treatments can be more selective than traditional chemotherapy but may still cause serious side effects and resistance.

## Target Organ

An organ especially affected by a disease, hormone, toxin, medicine, or other exposure. Effects may occur because the organ has specific receptors, receives the substance, or is especially vulnerable to damage.

## Target Sign

A classic cross-sectional ultrasonographic and CT finding of intussusception characterized by concentric alternating hyperechoic and hypoechoic rings formed by the invaginating bowel (intussusciens and intussusceptum) with mesenteric fat and fluid ('donut sign').

### Alternate Names:

Donut Sign of Intussusception; Bullseye Sign

## Target Tissue

Tissue that responds to a hormone, medicine, toxin, or other signal because its cells have the appropriate receptor or molecular target. The response may change growth, activity, metabolism, or survival.

## Tarlov Cyst

A fluid-filled sac around a nerve near the lower spine. Many cause no symptoms and are found by chance, but a larger cyst can cause lower-back or buttock pain, numbness, weakness, or bladder, bowel, or sexual problems.

## Tarlov Cysts

Tarlov cysts are fluid-filled sacs around the nerve roots, most often in the sacral spine. Many cause no symptoms, but larger or symptomatic cysts may cause pain, numbness, or bladder problems.

## Tar Remover Poisoning

Poisoning from swallowing, inhaling, or skin contact with tar-removing chemicals. Effects vary by product and may include irritation, dizziness, vomiting, or breathing problems.

## Tars

A thick black substance from organic material; tobacco tar contains chemicals that can damage lungs and cause cancer.

## Tarsal

Relating to the tarsus, the group of bones forming the ankle and hindfoot. Tarsal bones help support body weight and allow foot movement.

## Tarsal Bones

The seven bones of the hindfoot and midfoot that connect the ankle to the metatarsals and form the arches and load-bearing framework of the foot.

### Alternate Names:

Tarsus; Ankle Bones

## Tarsal Coalition

An abnormal connection between two foot bones, usually present from birth. It can make the foot stiff

---

or flat and cause pain, repeated ankle sprains, or difficulty walking, often becoming noticeable during adolescence.

### **Tarsal Joints**

The articulations among the tarsal bones and between the tarsals and metatarsals, providing stability and controlled movement for walking and adapting the foot to surfaces.

### **Tarsal Tunnel Syndrome**

Pressure on the tibial nerve as it passes through a narrow space near the inner ankle. It can cause burning pain, tingling, or numbness in the sole and may result from injury, swelling, diabetes, or a mass. Treatment depends on the cause.

### **Tarsorrhaphy**

A procedure that partially or completely closes the eyelids to protect the cornea or reduce exposure in conditions such as severe dry eye or facial nerve palsy.

### **Tar Staining**

A yellowish-brown discoloration observed on the fingers, fingernails, or facial hair caused by prolonged cutaneous deposition of tar and nicotine particulates from cigarette smoking. It serves as an objective physical sign of chronic heavy tobacco consumption.

### **Tarsus**

The group of seven bones forming the ankle and the back and middle parts of the foot. These bones connect the lower-leg bones to the metatarsals and help support body weight, absorb forces, and adapt the foot during walking.

### **TAR Syndrome**

Thrombocytopenia-absent-radius syndrome, an inherited condition causing low platelet counts and absent or severely shortened radius bones in the forearms. It may also cause limb differences, heart defects, kidney problems, or milk intolerance.

### **Tartar**

A hard deposit of mineralized dental plaque, also called calculus.

### **Tarui Disease**

A rare inherited muscle-energy disorder caused by deficiency of an enzyme needed to break down sugar for energy. Exercise can trigger early fatigue, cramps, muscle breakdown, and dark urine; some people also have chronic anemia or high uric acid.

#### **Alternate Names:**

Glycogen Storage Disease Type VII (GSD VII);  
Muscle Phosphofructokinase Deficiency

### **Tassinari Syndrome**

An older term for a childhood epilepsy disorder with frequent abnormal brain activity during sleep, seizures, and possible effects on learning or development. Modern clinicians usually use a more specific diagnosis based on the child's seizures and sleep recording.

### **Taste**

The sensation produced by the interaction of chemical substances with taste receptor cells (TRCs) on the tongue, palate, pharynx, epiglottis, and upper esophagus. Taste is one of the five primary senses (gustation).

### **Taste Bud**

A taste bud is a cluster of specialized sensory cells in papillae of the tongue and some oral tissues that detects sweet, sour, salty, bitter, and umami stimuli and transmits signals through cranial nerves.

### **Taste Disorder**

Abnormal reduction, loss, distortion, or unpleasant alteration of taste, caused by infection, medications, oral disease, nerve injury, neurologic disease, or other factors.

### **Taste Receptor**

A sensor on taste-bud cells that detects chemicals in food and helps produce the sensations of sweet, salty, sour, bitter, and savory (umami). Signals

---

then travel through nerves to the brain.

### Tattoos

Tattoos are permanent designs made by placing pigment into the skin. Risks include infection, allergic reactions, scarring, and difficulty removing the pigment.

### Taurine

Taurine is a sulfur-containing compound involved in normal cell, nerve, muscle, and bile-acid function. It is present in foods and added to some energy drinks; high caffeine intake in such drinks can be harmful.

### Taurodontism

A tooth-development difference with an enlarged pulp chamber and shortened roots, usually affecting molars. It may occur alone or with genetic conditions and can complicate dental treatment or root-canal procedures.

### Taussig-Bing Anomaly

A rare birth defect in which both major arteries leave the right lower heart chamber and a hole lies between the lower chambers below the pulmonary artery. It can cause severe low oxygen and heart failure in infancy and usually requires early specialist surgery.

#### Alternate Names:

Taussig-Bing Syndrome; DORV with Subpulmonic VSD

### TAVR

Alternate name; see *Transcatheter Aortic Valve Replacement*.

### Taxanes

Chemotherapy medicines, including paclitaxel and docetaxel, that disrupt cell division by stabilizing tiny internal structures. They treat several cancers but may cause low blood counts, nerve symptoms, hair loss, allergic reactions, or fluid retention.

### Taxotere

Taxotere is the brand name for docetaxel, a chemotherapy medicine that interferes with cell division. It is used for several cancers and can cause low blood counts, infection risk, hair loss, and other side effects.

### Tay-Sachs Disease

A rare autosomal recessive lysosomal storage disorder usually caused by disease-causing variants in the *HEXA* gene. Reduced or absent beta-hexosaminidase A activity prevents normal breakdown of GM2 ganglioside, which accumulates mainly in neurons. The infantile form causes progressive loss of previously acquired skills, weakness, an exaggerated startle response, seizures, and vision changes; juvenile and late-onset forms may progress more slowly and have variable movement, speech, psychiatric, or cognitive features.

#### Alternate Names:

GM2 Gangliosidosis Type 1 Hexosaminidase A Deficiency HEXA Deficiency

### Tazarotene

A topical retinoid used for acne, plaque psoriasis, and some sun-damaged skin. It can irritate skin and increases sun sensitivity; pregnancy precautions are important.

### TB

Alternate name; see *Tuberculosis*.

### TBG Blood Test

A blood test measuring thyroxine-binding globulin, the principal thyroid-hormone transport protein; it helps interpret abnormal total thyroid-hormone levels when binding-protein changes are suspected.

### TBI

Alternate name; see *Traumatic brain injury*.

### TB Test

A tuberculosis test looks for infection with *Mycobacterium tuberculosis* using a skin test, blood test, or examination of respiratory specimens.

---

## **TCA Cycle**

A series of chemical reactions inside cells that breaks down products of carbohydrates, fats, and proteins. It produces carbon dioxide and energy-carrying molecules used to make the cell's energy supply.

## **T Cell**

A type of white blood cell that matures in the thymus and helps protect the body from infections and abnormal cells. Some T cells kill infected cells, while others coordinate or limit immune responses.

## **T-Cell**

A type of white blood cell that helps direct immune responses or destroys infected and abnormal cells. T cells mature in the thymus and include several groups with different jobs.

## **T-Cell Count**

A laboratory measurement of T lymphocytes in blood, often including CD4 cells, used to assess immune function and monitor conditions such as HIV infection or immunosuppression.

## **T-Cell Lymphoma**

A cancer of T lymphocytes, a type of white blood cell. It includes several different diseases that may affect lymph nodes, skin, blood, or organs and can cause swollen nodes, fever, weight loss, rash, or other symptoms.

## **T-Cell Lymphomas**

Cancers that arise from T cells, a type of white blood cell. Different forms may affect lymph nodes, skin, blood, or internal organs and can cause swollen nodes, fever, weight loss, rash, infections, or other symptoms.

## **T-Cell Receptor**

A protein on a T cell that recognizes a particular piece of a germ or abnormal cell when another cell displays it. This recognition helps start or direct a targeted immune response.

## **T-Cells**

White blood cells that mature in the thymus and direct immune responses against infections and abnormal cells. Different types kill affected cells, coordinate defenses, remember threats, or reduce excessive inflammation.

### **Alternate Names:**

Thymus-Derived Lymphocytes; T Cells

## **Td**

Td is a vaccine protecting against tetanus and diphtheria, generally used as a booster in adolescents and adults according to local schedules.

## **Tdap Vaccination In Pregnancy**

Administration of tetanus, diphtheria, and acellular pertussis (Tdap) vaccine during each pregnancy, optimally at 27-36 weeks. The goal: boost maternal antibodies to protect the newborn from pertussis (whooping cough) before the infant can be vaccinated at 2 months.

## **Td Immunization**

Vaccination against tetanus and reduced-dose diphtheria, used for routine boosters, catch-up immunization, and wound-related protection according to age and vaccination history.

## **Tear**

A split or rupture in tissue, such as a muscle, tendon, ligament, meniscus, skin, or organ, caused by trauma, overuse, degeneration, or pressure.

## **Teardrop Cells**

Red blood cells shaped like teardrops on a blood smear. They may occur when the bone marrow is scarred or crowded by another disease, but they are not specific to one condition.

### **Alternate Names:**

Dacrocytes; Tear-Drop Poikilocytes

## **Tear Film Break-Up Time**

A quantitative clinical ophthalmologic test measuring tear film stability and evaporative dry eye; after instilling topical fluorescein, the interval between a



---

complete blink and the appearance of the first dry black spot in the pre-corneal tear film under cobalt blue light is timed (normal greater than 10 seconds; less than 10 seconds indicates tear film instability).

**Alternate Names:**

TBUT; Tear Breakup Assay

**Tearing**

Excessive production or inadequate drainage of tears, commonly caused by irritation, allergy, dry-eye reflex tearing, infection, eyelid malposition, or lacrimal-duct obstruction.

**Tears**

Fluid made mainly by the lacrimal glands that lubricates, nourishes, cleans, and protects the eye's surface. Tears contain water, salts, oils, mucus, and immune substances that help maintain clear, comfortable vision.

**Technetium-99**

A radioactive form of technetium that changes slowly by radioactive decay and is related to technetium-99m, the more widely used imaging isotope. Technetium isotopes are used in nuclear medicine and research.

**Technetium-99M**

A short-lived radioactive form of technetium used in many nuclear-medicine scans. It is attached to different substances so doctors can examine bones, the heart, kidneys, brain, or other organs; the radiation is detected by a special camera.

**Technique**

The method, equipment, and sequence of steps used for a medical examination, procedure, or imaging study. Correct technique improves safety, accuracy, and consistency of results.

**Tectum**

The dorsal roof plate of the midbrain located posterior to the cerebral aqueduct, composed of the paired superior colliculi (mediating visual orientation reflexes and saccades) and inferior colliculi (relaying auditory pathways).

**Alternate Names:**

Tectum Mesencephali; Quadrigeminal Plate

**TEE**

TEE, or transesophageal echocardiography, uses an ultrasound probe in the esophagus to obtain detailed images of heart valves, chambers, clots, and nearby vessels.

**Teenage Pregnancy**

Pregnancy in a person aged 19 years or younger. Compared with adult pregnancy, it may carry higher risks of anemia, high blood pressure, preterm birth, and low birth weight, although outcomes improve with timely prenatal care and support.

**Teenagers And Drugs**

Drug use during adolescence can affect brain development, school and relationships, safety, and mental health; early, nonjudgmental screening and support can reduce harm.

**Teenagers And Sleep**

Teenagers commonly need about eight to ten hours of sleep, but schedules, screens, caffeine, mood, school demands, and sleep disorders can reduce sleep quality.

**Teen Depression**

Depression in an adolescent, causing persistent sadness, irritability, loss of interest, low energy, sleep or appetite changes, hopelessness, or thoughts of death. Symptoms last long enough to impair school, relationships, or daily life and need assessment.

**Alternate Names:**

Adolescent Depression; Major Depressive Disorder in Adolescents

**TEER**

Alternate name; see *Transcatheter Edge-to-Edge Repair*.

**Teeth**

Hard, mineralized structures in the mouth used for cutting, tearing, and grinding food. They are held

---

in the jaw by surrounding bone and gums and can be affected by decay, infection, injury, or wear.

### Teeth Grinding

Alternate name; see *Teeth grinding (bruxism)*.

### Teeth-Grinding

Alternate name; see *Bruxism*. Repetitive clenching or grinding of teeth, often during sleep and capable of causing tooth wear and jaw pain.

### Teething

Teething is eruption of an infant's primary teeth through the gums and may cause gum discomfort, drooling, and chewing; high fever or severe illness suggests another cause.

### Teichopsia

A visual disturbance involving shimmering zigzags, flashing patterns, or fortress-like shapes. It commonly occurs as a migraine aura but can have other causes, especially when new, persistent, or associated with vision loss.

### Teicoplanin

A glycopeptide antibiotic used for selected serious infections caused by susceptible Gram-positive bacteria, including some resistant staphylococci. Kidney function and blood levels may need monitoring.

### Telangiectasia

A visible, permanently dilated small blood vessel near the skin or mucosal surface, occurring alone or with conditions such as rosacea, liver disease, or inherited vascular syndromes.

### Telangiectasis

Widening of small blood vessels near the skin or a moist body lining, producing visible red or purple lines or patches. Causes include sun exposure, rosacea, hormones, liver disease, and inherited conditions.

### Teledermatology

Teledermatology is the use of video, photographs,

or other digital communication to assess and manage skin conditions remotely. Some problems still require an in-person examination or biopsy.

### Telegraphic Speech

A style of verbal expression in which speech consists predominantly of content words (nouns and verbs) with omission of grammatical morphemes, prepositions, conjunctions, and auxiliary verbs, resembling a telegram. It is characteristic of expressive (Broca) non-fluent aphasia.

### Telehealth

Delivery of healthcare through secure electronic communication or remote monitoring, including video visits, telephone care, messaging, and transmission of clinical data.

### Telemedicine

Delivery of medical care at a distance using information and communication technology. It may include video or telephone consultations, secure messages, remote diagnosis, treatment, and review of clinical information.

### Teleradiology

The remote review of medical images, such as X-rays, CT scans, or MRI scans, by a qualified radiologist who is not at the imaging site. It can provide urgent or specialist interpretation when local expertise is unavailable and must protect patient information.

### Tele-Rehabilitation

Rehabilitation provided remotely through video visits, phone calls, apps, or monitoring devices. It may include assessment, home exercises, education, and progress checks when in-person care is difficult to access.

### Telogen and Anagen Effluvium

Diffuse, usually nonscarring hair shedding caused by premature entry of follicles into the resting telogen phase or abrupt injury to actively growing anagen follicles. Telogen shedding commonly follows illness or stress after a delay, whereas anagen shedding can begin within days to weeks of chemotherapy,

---

radiation, or toxin exposure.

### **Telogen Effluvium**

A common, non-scarring form of diffuse hair shedding that occurs when more hairs than usual enter the resting (telogen) phase. Shedding often begins 2–4 months after illness, fever, surgery, childbirth, major stress, rapid weight loss, nutritional deficiency, or certain medicines. It is usually temporary, with regrowth after the trigger is corrected, although recovery may take several months.

### **Telomere**

A protective DNA region at the end of each chromosome. It helps keep genetic material stable during cell division and usually becomes shorter as cells divide and age.

### **Temazepam**

A sedative medicine sometimes used for short-term insomnia. It can cause drowsiness, poor coordination, memory problems, dependence, and dangerously slow breathing, especially when combined with alcohol, opioids, or other sedatives.

### **Temperature**

A measurement of body heat. The result varies with the measurement site, time, activity, age, and illness.

### **Temperature Measurement**

Measurement of body or environmental temperature using a thermometer or sensor at a specified site, interpreted with method, age, timing, and clinical context.

### **Temple**

The region of the skull at the side of the head between the forehead and ear.

### **Temple Syndrome**

A genomic-imprinting disorder involving chromosome 14q32 that causes prenatal and postnatal growth restriction, hypotonia, developmental delay, feeding difficulty, short stature, and abnormal

pubertal development.

### **Temporal**

Relating either to time or to the temple area on the side of the skull, including the temporal lobe.

### **Temporal Arteritis**

Inflammation of arteries, especially those at the temples, usually in adults over 50. It may cause a new headache, scalp tenderness, jaw pain while chewing, fever, or vision problems and can threaten sight.

### **Temporal Arteritis Pathology**

Granulomatous inflammation of medium and large arteries, often with disruption of the internal elastic lamina, intimal hyperplasia, medial inflammation, and giant cells. Biopsy findings may be segmental and can persist after treatment.

### **Temporal Artery**

An artery running along the side of the scalp, supplying nearby tissues; inflammation can cause temporal arteritis.

### **Temporal Bones**

The two skull bones on the sides and base of the head. They contain the ear structures and passages for important nerves and blood vessels and help protect the brain.

### **Temporal Lobe**

One of the main areas on each side of the brain, near the ears. It helps process sounds, understand language, form memories, and process emotions.

### **Temporal Lobe Epilepsy**

Epilepsy in which seizures begin in the part of the brain near the ears that helps process memory, emotion, and sound. Episodes may cause unusual feelings, déjà vu, staring, loss of awareness, or repetitive movements.

### **Temporal Lobe Seizure**

A seizure that begins in the temporal lobe of the

---

brain and may cause altered awareness, unusual sensations, or repetitive movements.

**Alternate Names:**

Focal Impaired Awareness Seizure; Psychomotor Seizure

**Temporalis**

A broad muscle on each side of the head, over the temple. It helps close the jaw for chewing; its back fibers also pull the lower jaw backward.

**Alternate Names:**

Temporal Muscle

**Temporary Loss Of Consciousness**

A brief period of unresponsiveness with loss of posture, followed by recovery. Fainting is common, but seizures, abnormal heart rhythms, low blood sugar, and other illnesses can also cause it; unexplained or repeated episodes need assessment.

**Temporary Unconsciousness**

A brief episode of loss of awareness and responsiveness, often caused by syncope but also possible with seizure, concussion, hypoglycemia, or other conditions requiring assessment.

**Temporomandibular Disorders**

Alternate name; see *TMJ disorders*.

**Temporomandibular Joint**

The jaw joint in front of each ear, where the lower jaw meets the skull. It allows the mouth to open, close, and move side to side; problems can cause pain, clicking, stiffness, or limited movement.

**Temporomandibular Joint Disorder**

Pain or impaired movement involving the jaw joint or chewing muscles. It may cause jaw pain, clicking, stiffness, headache, or difficulty opening the mouth. Clenching, grinding, injury, arthritis, and muscle tension can contribute; treatment usually begins with simple pain relief and jaw-rest measures.

**Temporomandibular Joint Dysfunction**

A musculoskeletal disorder affecting the temporo-

mandibular joint, articular disc, and masticatory muscles, characterized by preauricular facial pain, clicking or crepitus on jaw movement, and restricted mandibular opening.

**Alternate Names:**

TMD; TMJ Syndrome

**Temporomandibular Joint Replacement**

Surgery that removes a severely damaged jaw joint and replaces it with an artificial joint. It may be considered for ankylosis, advanced joint destruction, or persistent severe symptoms after other treatments have failed.

**Temporomandibular Joint Syndrome**

Pain or dysfunction involving the jaw joint and chewing muscles, often causing jaw pain, clicking, restricted opening, headache, or tooth-related symptoms. Causes include clenching, trauma, arthritis, and stress-related muscle tension. Persistent locking, severe pain, or inability to eat requires dental or medical evaluation.

**TEN**

Alternate name; see *Toxic epidermal necrolysis*.

**Tenascin**

A group of proteins in the tissue surrounding cells that help organize tissue structure and influence growth, healing, and inflammation. Levels may change in wounds, developing tissues, and some tumors.

**Tender**

Painful or sensitive when touched or pressed, a finding that may indicate inflammation, infection, trauma, ischemia, or another underlying process.

**Tenderness**

Pain or discomfort caused by pressing or touching a body part. It may indicate inflammation, injury, infection, or another local or systemic problem.

**Tendinitis**

Inflammation of a tendon, often associated with

---

repetitive loading, overuse, or injury. It can cause localized tenderness, stiffness, swelling, or pain with movement. Tendinitis is distinct from tendinopathy, a broader term that includes tendon pain and structural change without necessarily involving active inflammation.

**Alternate Names:**

Tendonitis; Tendinopathy

**Tendinopathy**

Pain and abnormal structure or function of a tendon, often related to repeated loading, aging, or injury. It can cause localized tenderness, stiffness, weakness, or pain with movement and does not necessarily involve active inflammation.

**Tendinosis**

Long-lasting wear and disorganization of tendon tissue, usually from repeated loading or an injury rather than sudden inflammation. It can cause pain and reduced strength; treatment may include activity changes and rehabilitation.

**Tendon**

A strong band of tissue that connects a muscle to a bone and transfers the muscle's pull to move a joint. Tendons have a limited blood supply, so some injuries heal slowly.

**Tendon Injury**

Damage to a tendon ranging from microscopic overuse injury to partial or complete rupture, causing localized pain, weakness, swelling, or loss of function.

**Tendonitis**

Alternate spelling; see *Tendinitis*.

**Tendon Repair**

Surgery to reconnect a torn tendon to itself or to bone so that movement can be restored. It is followed by protected movement and rehabilitation; re-tearing, stiffness, weakness, infection, and nerve or vessel injury can occur.

**Tendon Rupture**

A partial or complete tear of a tendon, usually caused by sudden force, chronic tendon disease, or injury. It can cause sudden pain, swelling, weakness, and loss of the movement normally produced by that tendon.

**Tendon Sheath**

A thin lubricated covering around some tendons that reduces friction as the tendon moves; inflammation is called tenosynovitis.

**Tendon Transfer**

Surgical relocation of a functioning tendon to restore movement or balance when the original muscle-tendon unit is absent, paralyzed, or irreparable. Successful transfer requires an adequate donor, appropriate excursion, and a stable recipient joint.

**Tendon Xanthoma**

Firm, often painless cholesterol deposits in tendons, especially the Achilles tendons and tendons of the fingers. They can occur with inherited high cholesterol and other lipid disorders and should lead to a cholesterol assessment.

**Alternate Names:**

Tendinous Xanthoma; Achilles Tendon Xanthoma

**Tenecteplase**

A clot-dissolving medicine that helps restore blood flow during selected ischemic strokes and heart attacks. It is given through a vein, usually as one dose, but can cause serious bleeding, including bleeding in the brain.

**Tenericutes**

A former classification group for bacteria that lack a rigid cell wall, including Mycoplasma. The name is mainly used in older biological classifications; it is not a diagnosis or a routine clinical test result.

**Tenesmus**

A distressing sensation of needing to empty the bowel or bladder despite little output, commonly caused by rectal inflammation, infection, tumor,

---

constipation, or urinary irritation.

### Teniposide

Teniposide is a podophyllotoxin-derived antineoplastic drug that inhibits topoisomerase II and is used in selected leukemia or lymphoma regimens.

### Tennis Elbow

Alternate name; see *Lateral Epicondylitis*.

### Tennis Elbow Surgery

An operation to remove or repair damaged tendon tissue at the outer elbow when persistent tennis-elbow pain has not improved with nonsurgical care. Recovery requires rehabilitation, and pain, stiffness, nerve injury, or continued symptoms are possible.

### Tenodesis

Surgical fixation of a tendon to bone or another tendon to restore function, stabilize a joint, or treat tendon imbalance.

### Tenofovir

An antiviral medicine used with other drugs to treat HIV and, in some forms, to treat or prevent hepatitis B. It stops viruses copying their genetic material; kidney and bone effects require appropriate monitoring.

### Tenosynovitis

Tenosynovitis is inflammation of the sheath surrounding a tendon, producing pain, swelling, and restricted gliding, commonly in the hand or wrist.

### Tenotomy

Surgery that cuts or releases a tendon to reduce tightness, correct a deformity, relieve pressure, or change movement. It may be performed through a small opening or a larger incision; weakness, bleeding, infection, nerve injury, or recurrence can occur.

### Tenovaginitis

Inflammation of a tendon and its surrounding sheath, also called tenosynovitis.

### Tensile Strength

The ability of a tissue or material to resist being pulled apart; it is important in wound healing and the integrity of sutures, tendons, and other connective tissues.

### Tension

A pulling force or state of being stretched; in medicine it may describe tissue tension, blood-vessel wall tension, muscle tightness, or psychological strain depending on context.

### Tension Headache

A common headache causing mild or moderate pressure or tightness, often on both sides or like a band around the head. It usually lacks vomiting and may be linked with stress, poor sleep, muscle tension, or prolonged screen or neck strain.

### Tension Pneumothorax

A life-threatening buildup of air between the lung and chest wall that presses on the lung and heart. It can cause severe breathing difficulty, low blood pressure, a fast heartbeat, one-sided absent breath sounds, and collapse. Treatment rapidly releases the trapped air and usually includes a chest tube.

#### Alternate Names:

Tension Air Leak; Pleural Tension

### Tension-Type Headache

A usually bilateral, pressing or tightening headache of mild-to-moderate intensity without prominent nausea, often associated with stress, muscle tenderness, or poor sleep.

### Tentorium

A strong fold of the brain's protective covering separating the cerebellum from the back parts of the cerebrum. It supports brain structures and contains an opening through which the brainstem passes.

### Tentorium Cerebelli

A broad, crescent-shaped dural fold that arches over the posterior cranial fossa, physically separating the overlying occipital lobes of the cerebral hemispheres



---

from the underlying cerebellum; its anterior free edge forms the tentorial notch (incisura).

### **Teprotumumab**

A monoclonal antibody targeting the insulin-like growth factor 1 receptor and used to treat thyroid eye disease in selected patients.

### **Teratocarcinoma**

A teratocarcinoma is a malignant germ-cell tumor containing tissues from more than one embryonic layer, often arising in the testis or ovary.

### **Teratogen**

A medicine, chemical, infection, radiation exposure, or other influence that can harm an embryo or fetus and cause birth defects, growth problems, abnormal development, or pregnancy loss. Risk depends on the exposure, amount, timing, and individual factors.

### **Teratogenesis**

The development of a birth defect, impaired growth, abnormal function, or pregnancy loss after harmful exposure during pregnancy. Risk depends on the substance or infection, amount, timing, route, and genetic factors; exposure does not automatically mean that harm will occur.

### **Teratogenic**

Capable of causing congenital malformations, growth impairment, functional deficits, or pregnancy loss when an embryo or fetus is exposed during a vulnerable developmental period.

### **Teratogenic Drugs**

Medicines that can harm an embryo or fetus and cause birth defects, poor growth, abnormal development, or pregnancy loss. The risk depends on the medicine, dose, timing, and stage of pregnancy.

### **Teratogenicity**

The ability of a medicine, chemical, infection, or other exposure to harm an embryo or fetus. Risk depends on the substance, amount, timing, and in-

dividual factors; exposure does not guarantee harm, but should be discussed promptly with a healthcare professional.

### **Teratology and Drug Use During Pregnancy**

Teratology is the study of how genetic, environmental, infectious, and medicine exposures can affect embryonic or fetal development. Risk depends on the substance, dose, timing, route, and maternal factors; medication decisions should use reliable pregnancy-specific guidance rather than abrupt self-discontinuation.

### **Teratoma**

A tumor made of several types of tissue, such as skin, muscle, bone, or nerve tissue, because it develops from cells capable of forming different body tissues. It may be noncancerous or cancerous and can occur in the ovary, testicle, or other sites.

### **Teratozoospermia**

Teratozoospermia is a high proportion of abnormally shaped sperm and may contribute to reduced fertility, though semen results must be interpreted as a whole.

### **Terazosin Hydrochloride**

The hydrochloride salt of terazosin, an alpha-1 blocker used for urinary symptoms from benign prostatic enlargement and sometimes for hypertension; dizziness and first-dose fainting can occur.

### **Terbinafine**

An antifungal medicine that damages the outer covering of fungi, stopping their growth or killing them. It is used on the skin or taken by mouth for selected fungal infections; liver problems and taste changes are possible.

### **Terbufos**

Terbufos is a highly toxic organophosphate pesticide that inhibits acetylcholinesterase and can cause severe cholinergic poisoning.

---

### Terbutaline

A beta-2 agonist used as a bronchodilator; it has limited and specialist use as a tocolytic.

### Teres Major Muscle

A thick muscle of the posterior shoulder originating from the dorsal inferior angle of the scapula and inserting onto the medial lip of the intertubercular sulcus of the humerus; innervated by the lower subscapular nerve, it acts to adduct, medially rotate, and extend the arm.

### Terfenadine

An older antihistamine withdrawn or restricted in many places because interactions and excessive blood levels can cause dangerous heart-rhythm abnormalities.

### Teriparatide For Osteoporosis

An injectable medicine related to parathyroid hormone that stimulates new bone formation. It is used for selected people with severe osteoporosis or high fracture risk, usually for a limited period, and is followed by treatment that helps maintain the bone gained.

### Terlipressin

A medicine similar to vasopressin that tightens certain blood vessels. It is used in selected hospital treatments for life-threatening bleeding from enlarged esophageal veins or kidney failure related to advanced liver disease; reduced blood flow and breathing problems are important risks.

### Term

Describing a pregnancy or birth at 37 or more completed weeks. Birth before 37 weeks is called preterm; birth after the expected period may be called post-term.

### Term Birth

Birth at or after 37 completed weeks of pregnancy. Early term is 37 to 38 weeks, full term is 39 to 40 weeks, and late or post-term describes later births; gestational age helps guide newborn care.

### Terminal Care

Care focused on comfort, symptom relief, dignity, and support for a person with an advanced illness who is approaching the end of life.

### Terminal Delirium

Delirium occurring near the end of life, often caused by advanced organ failure, medication effects, metabolic disturbance, infection, or reduced cerebral perfusion. Management prioritizes comfort, reversible contributors when consistent with goals, and support for the patient and family.

### Terminal Duct Lobular Unit

The smallest working unit of the breast, made of a small duct and the lobules connected to it. Many common breast changes and cancers begin in this area.

### Terminal Ileitis

Inflammation of the distal ileum, commonly associated with Crohn disease but also caused by infection, ischemia, medication, or other inflammatory conditions and producing pain, diarrhea, or bleeding.

### Terminal Illness

An advanced disease that is expected to lead to death, although the timing is uncertain and treatment may sometimes slow it. Care may focus on comfort, symptom relief, personal goals, and support for the patient and family.

### Terminal Insomnia

Waking too early and being unable to return to sleep, a pattern that can occur in depression, circadian-rhythm disorders, aging, or other insomnia conditions.

### Terry's Lines

Alternate name; see *Terry's Nails*.

### Terry's Nails

An abnormal nail appearance in which the proximal eighty percent or more of the nail bed is opaque and chalky-white (apparent leukonychia) with ob-

---

scuration of the lunula, bordered by a narrow distal band of pink or reddish-brown vascular nail bed. It is classically associated with liver cirrhosis, chronic congestive heart failure, type 2 diabetes mellitus, and systemic sclerosis, as well as normal aging.

**Alternate Names:**

Terry Nails; Terry's Lines; Terry Lines

**Terson Syndrome**

The occurrence of intraocular hemorrhage (subhyaloid, preretinal, or vitreous hemorrhage) in association with acute aneurysmal subarachnoid hemorrhage or severe traumatic brain injury, caused by sudden severe intracranial pressure elevation transmitted through the optic nerve sheath.

**Alternate Names:**

Terson's Syndrome; Subarachnoid Intraocular Hemorrhage

**Tertian Fever**

A fever that returns about every 48 hours. This pattern can occur with some malaria infections, especially those caused by *Plasmodium vivax* or *P. falciparum*; the pattern alone does not confirm malaria.

**Tertiary Hyperparathyroidism**

Persistent overproduction of parathyroid hormone after long-standing kidney disease or another cause of parathyroid overactivity. It can raise the blood calcium level and damage bones or kidneys.

**Tertiary Prevention**

Care that reduces complications, disability, or suffering from an established disease. Examples include rehabilitation, long-term disease management, preventing relapses, controlling symptoms, restoring function, and providing palliative care.

**Testes**

The paired male reproductive organs in the scrotum. They produce sperm and testosterone and may be affected by infection, injury, torsion, tumors, or disorders of development.

**Testicle**

One of the male reproductive glands in the scrotum. It produces sperm and testosterone; sudden severe testicular pain is an emergency because it may indicate torsion.

**Testicle Lump**

A localized mass or enlargement in or around a testis caused by cyst, infection, inflammation, torsion, trauma, or tumor; a new intratesticular mass requires prompt evaluation.

**Testicle Pain**

Alternate name; see *Testicular Pain*.

**Testicles**

The paired male reproductive glands in the scrotum that produce sperm and testosterone; pain, enlargement, a new lump, or sudden change requires examination.

**Testicular Appendage Torsion**

Twisting of a small, nonessential remnant attached to the testicle or epididymis. It commonly causes localized scrotal pain and swelling in children and may produce a small tender blue spot under the scrotal skin.

**Testicular Artery**

An artery that supplies blood to a testis and travels in the spermatic cord.

**Testicular Atrophy**

Shrinkage of a testicle with loss of some functioning tissue. Causes include an undescended testicle, infection, twisting, injury, varicocele, hormone problems, aging, or other damage; it may reduce sperm or testosterone production.

**Testicular Biopsy**

Removal of a small sample of testicular tissue for microscopic examination, used selectively to evaluate infertility, atypical lesions, or suspected testicular disease.

---

## Testicular Cancer

Cancer arising in a testicle, most often in younger adult men. It commonly appears as a painless lump, enlargement, heaviness, or ache. Ultrasound and blood tests help assess a suspected tumor; treatment is highly effective for many people, especially when found early.

## Testicular Cancer In Children

Cancer arising in a child's testicle. It may appear as a painless lump, swelling, or enlargement; the tumor types and behavior differ with age.

### Alternate Names:

Pediatric Testicular Neoplasm; Pediatric Testicular Tumor

## Testicular Choriocarcinoma

A rare, highly aggressive nonseminomatous germ-cell tumor containing trophoblastic cells. It often produces markedly elevated hCG and may spread early through the bloodstream, including to the lungs or brain; urgent specialist evaluation is required.

## Testicular Descent

The movement of the testes from the abdomen into the scrotum before or soon after birth.

## Testicular Diseases

Conditions affecting the testis, including twisting, infection, injury, tumors, blood-vessel problems, and abnormal development. Symptoms may include pain, swelling, a lump, fertility problems, or changes in hormone production.

## Testicular Disorders

Conditions affecting the testes or surrounding structures, including torsion, infection, inflammation, trauma, tumors, and hormonal disorders. Symptoms may include pain, swelling, a mass, infertility, or changes in sexual development and require appropriate evaluation.

### Alternate Names:

Testicular Pathology; Scrotal Disorders

## Testicular Failure

Testicular failure occurs when the testes do not produce enough testosterone, sperm, or both. Causes include genetic conditions, injury, infection, medicines, and damage from chemotherapy or radiation.

## Testicular Function

The two main jobs of the testes: making sperm and producing the sex hormone testosterone.

## Testicular Gonadoblastoma

A rare tumor made from developing reproductive cells and supporting cells in an abnormally developed or undescended testicle. It is often associated with Y-chromosome material and may increase the risk of another germ-cell cancer.

## Testicular Hydrocele

A buildup of clear fluid around a testicle, causing painless swelling of the scrotum. It may be present from birth or develop after injury, infection, inflammation, or another condition; examination helps exclude other causes of swelling.

## Testicular Infection

Alternate name; see *Orchitis*.

## Testicular Interstitium

The tissue between the seminiferous tubules. It contains Leydig cells, blood vessels, lymph vessels, and connective tissue.

## Testicular Lobule

One of the small sections inside a testis that contains seminiferous tubules.

## Testicular Mass

A lump or enlargement within or around a testicle caused by tumor, infection, inflammation, cyst, hematoma, or torsion. A solid mass within the testicle can be linked to cancer or another serious condition.

---

### Testicular Microlithiasis

Multiple tiny calcifications within the testicular tissue seen on ultrasound. The finding alone does not establish cancer, and follow-up depends on other risk factors and clinical findings.

### Testicular Pain

Pain in one or both testes. Causes include torsion, infection, trauma, inflammation, hernia, and tumors; pain may also arise from nearby structures.

### Testicular Self-Exam

A testicular self-exam is a gentle check for new lumps, swelling, heaviness, or changes in size or shape; a new abnormality should be medically assessed.

### Testicular Seminoma

A malignant germ-cell tumor of the testis, usually affecting younger adults. It commonly presents as a painless testicular enlargement or mass; diagnosis uses ultrasound, orchiectomy pathology, tumor markers, and staging imaging.

### Testicular Torsion

A medical emergency in which the cord supplying a testicle twists and cuts off its blood supply. It usually causes sudden, severe one-sided scrotal pain, swelling, nausea, or vomiting. Treatment usually involves immediate surgery.

#### Alternate Names:

Torsion of Spermatic Cord; Testis Twisting

### Testicular Torsion Emergency

Rotation of the spermatic cord that compromises testicular blood flow, causing sudden unilateral scrotal pain, nausea, a high-riding testis, and an abnormal cremasteric reflex. Immediate surgical exploration and detorsion are required; imaging must not delay treatment when suspicion is high.

### Testicular Torsion Overview

Alternate name; see *Testicular torsion*.

### Testicular Torsion Repair

Urgent surgery to untwist a spermatic cord and restore testicular blood flow, followed by fixation of the affected and usually opposite testis to prevent recurrence.

### Testicular Trauma

Injury to a testicle caused by a blow, crush injury, penetrating wound, or other force. It may cause pain, swelling, bruising, bleeding, or rupture of the tough covering around the testicle.

### Testicular Tumor

Alternate name; see *Testicular cancer*.

### Testicular Vein

A vein that drains blood from a testis through the spermatic cord. The right and left sides drain into different larger veins.

### Testis

One of the two male reproductive glands in the scrotum. It produces sperm and testosterone and is surrounded by protective tissue. Sudden pain, swelling, or a new lump needs prompt medical assessment.

#### Alternate Names:

Male Gonad; Orchis

### Testis Disorder

A condition affecting a testicle, including torsion, infection, tumor, trauma, infertility-related damage, hormonal dysfunction, or abnormal development.

### Testis Neoplasm

A tumor arising in testicular tissue, most commonly a germ-cell tumor in younger adults; painless enlargement, heaviness, or a solid mass requires prompt ultrasound and evaluation.

### Test Meal

A meal with known ingredients and amounts given before measuring a body response, such as stomach emptying, acid production, blood sugar, or hormone release. The person follows specific preparation and

---

timing instructions.

### **Testosterone**

A steroid sex hormone produced mainly by the testes and, in smaller amounts, by the ovaries and adrenal glands. It supports reproductive development, muscle and bone maintenance, red-blood-cell production, libido, and other physiologic functions; levels vary with age, sex, time of day, illness, and medicines.

### **Testosterone Deficiency**

Alternate name; see *Male hypogonadism*.

### **Testosterone Propionate**

A short-acting injectable testosterone ester used in selected hormone-replacement or androgen-deficiency settings; risks include erythrocytosis, infertility, acne, mood effects, and prostate-related monitoring needs.

### **Tests For H Pylori**

Tests for *Helicobacter pylori* infection, including urea breath testing, stool antigen testing, biopsy-based testing, or selected blood tests; choice depends on treatment and clinical context.

### **Test Strips**

Narrow strips containing chemicals that react with a sample to measure or detect a substance. Common examples test urine for blood or infection, or test blood for glucose or ketones. Results can be affected by storage, timing, contamination, and incorrect use.

### **Tetanus**

A serious infection caused by bacteria entering a wound and producing a toxin that causes severe muscle stiffness and painful spasms. It can cause lockjaw, breathing problems, and death.

#### **Alternate Names:**

Lockjaw

### **Tetanus Toxin**

A toxin made by the bacterium *Clostridium tetani*.

It blocks calming signals in the nervous system, causing painful muscle stiffness and spasms, including lockjaw. Vaccination and proper wound care reduce the risk of tetanus.

### **Tetany**

Muscle cramps, spasms, or twitching caused by increased nerve and muscle excitability, most often from low calcium or magnesium. Tingling around the mouth or in the hands may occur; severe spasms or breathing difficulty may occur in serious cases.

### **Tethered Cord Syndrome**

A condition in which the spinal cord is abnormally attached and stretched inside the spinal canal. It may cause back or leg pain, weakness, numbness, foot changes, or bladder and bowel problems, sometimes worsening during growth.

### **Tetrabenazine**

A medicine that reduces storage and release of certain brain chemicals involved in movement. It treats selected involuntary movement disorders but may cause sleepiness, depression, restlessness, stiffness, swallowing problems, or abnormal movements.

### **Tetracaine**

Tetracaine is a potent ester local anaesthetic used topically or for spinal anaesthesia; excessive dose can cause seizures, cardiovascular toxicity, and methemoglobinemia.

### **Tetrachloroethylene**

Tetrachloroethylene, or perchloroethylene, is an industrial solvent used in dry cleaning; substantial exposure can affect the nervous system, liver, and kidneys.

### **Tetrachlorvinphos**

An organophosphate insecticide that can block acetylcholinesterase. Significant exposure may cause sweating, vomiting, pinpoint pupils, muscle twitching, slow breathing, seizures, or coma.



---

## **Tetracycline**

An antibiotic that stops bacteria making proteins needed for growth. It treats selected infections but can cause stomach upset, sensitivity to sunlight, and permanent tooth discolouration if used during tooth development.

## **Tetracyclines**

A group of antibiotics that stop bacteria making proteins needed for growth. Examples include tetracycline, doxycycline, and minocycline. Side effects can include stomach upset and increased sensitivity to sunlight.

## **Tetracycline Stain**

Gray, yellow, or brown tooth discolouration caused by tetracycline-family antibiotics taken while teeth are developing. The colour is within the tooth and may require professional whitening, veneers, or crowns for cosmetic improvement.

## **Tetrahydrozoline Poisoning**

Poisoning from swallowing eye drops or nasal drops containing tetrahydrozoline. It can cause extreme sleepiness, slow heart rate, low blood pressure, and breathing problems, especially in children.

## **Tetralogy Of Fallot**

A heart defect present from birth with four features: a hole between the lower chambers, narrowing of the route from the right ventricle to the lungs (usually pulmonary stenosis), an aorta positioned over the hole, and thickening of the right ventricle. Reduced blood flow to the lungs can lower the oxygen level in the blood.

## **Tetraplegia**

Paralysis or severe weakness affecting both arms and both legs, usually because the spinal cord in the neck has been injured or damaged by disease. Trunk movement, sensation, breathing, and bladder or bowel control may also be affected.

## **Tet Spell**

A sudden episode of severe bluish coloring and low oxygen in a child with unrepaired tetralogy of Fallot.

Crying, feeding, or exertion may trigger it; the child may breathe rapidly, become limp, or faint. It is an emergency requiring calming, positioning, oxygen, and urgent specialist care.

### **Alternate Names:**

Hypercyanotic Spell; Paroxysmal Dyspneic Attack; Blue Spell

## **TGA**

Alternate name; see *Transient global amnesia*.

## **Th1/Th2/Th17/Tfh/Treg Balance**

A description of how several types of helper immune cells work together. They help fight infections, support antibody production, promote inflammation, or limit excessive immune activity; disruption may contribute to allergy or autoimmune disease.

## **Thalamic Pain Syndrome**

A central neuropathic pain syndrome developing weeks to months after a cerebrovascular infarct involving the ventral posterolateral (VPL) nucleus of the thalamus (posterior cerebral artery occlusion), characterized by burning, allodynic, and severe intractable contralateral hemibody pain.

### **Alternate Names:**

Dejerine-Roussy Syndrome

## **Thalamus**

A pair of structures deep within the brain that relay most sensory information, such as touch, vision, and hearing, to the outer brain. They also help regulate movement, alertness, and memory.

## **Thalassaemia**

Alternate spelling; see *Thalassemia*.

## **Thalassemia**

A group of inherited blood disorders in which the body makes too little of one part of hemoglobin, the oxygen-carrying protein in red blood cells. This causes anemia, ranging from mild to severe, and may require regular treatment.

### **Alternate Names:**

Cooley's Anemia Mediterranean Anemia

---

## Thalassemia Major

A severe inherited blood disorder in which the body makes too little hemoglobin. It causes serious anemia beginning in childhood and may require regular blood transfusions, with monitoring for iron buildup and organ damage.

## Thalidomide

A medicine that changes immune activity and is used for selected cancers and inflammatory conditions. It can cause severe birth defects, nerve damage, blood clots, sleepiness, and other harms, requiring strict safety controls.

## Thallium

Thallium is a toxic heavy metal that can cause gastrointestinal illness followed by painful nerve damage, hair loss, skin changes, kidney or liver injury, seizures, and heart or brain problems.

### Alternate Names:

Thallium Toxicity; Thallotoxicosis

## Thallium-201

A radioactive substance sometimes used in heart-blood-flow scans. Heart muscle takes it up in relation to its blood supply, and a special camera shows areas receiving less blood.

## Thallium Poisoning

Alternate name; see *Thallium*.

## Thallium Scan

A nuclear-medicine test using thallium-201 to show blood flow to heart muscle or selected tumors. It is now less common because other tracers are often preferred, but may still have specific uses.

## Thallium Toxicity

Heavy-metal poisoning that may begin with gastrointestinal symptoms and progress to painful sensorimotor neuropathy, encephalopathy, alopecia, and nail changes such as Mees lines. Blood or urine testing supports diagnosis; severe poisoning requires toxicology consultation and may be treated with Prussian blue plus supportive care.

## Thanatophoric Dysplasia

A severe inherited disorder of bone growth causing extremely short limbs, a narrow chest, a large head, and abnormal skull and spine development. The small chest can prevent adequate breathing after birth and is often life-threatening.

## Theca

A connective-tissue sheath, especially the theca surrounding an ovarian follicle.

## Theca Lutein Cyst

A usually temporary ovarian cyst caused by strong stimulation from pregnancy-related hormones or fertility treatment. It is often present in both ovaries and may cause abdominal pressure or pain; it commonly shrinks when hormone levels return to normal.

## Thecoma

A usually benign ovarian sex-cord stromal tumor composed of theca-like cells, often producing estrogen and presenting with abnormal uterine bleeding or an incidental adnexal mass.

## Thelarche

The onset of breast development, usually the first visible sign of female puberty. It commonly begins between ages 8 and 13, with substantial normal variation.

## Thenar Eminence

The fleshy mound at the base of the thumb, formed by muscles that move the thumb. Damage to the supplying nerve or muscles can weaken thumb opposition and make grasping difficult.

## Theophylline

A medicine that can relax airway muscles and improve breathing in selected lung diseases. Its safe and harmful doses are close together, so blood levels, medicine interactions, heart rhythm, and side effects need monitoring.

---

## Theophylline Toxicity

Toxicity causing persistent nausea and vomiting, tremor, agitation, tachycardia, hypokalemia, hyperglycemia, seizures, or life-threatening atrial or ventricular dysrhythmias. Measure serial serum levels and electrolytes; benzodiazepines treat seizures, repeated activated charcoal may reduce absorption, and hemodialysis is considered for severe poisoning.

## Theranostic Pair

Two related medical substances designed to find and treat the same target in the body. One helps show where the target is on an image; the other delivers treatment to cells carrying that target.

## Theranostics

A medical approach that uses one targeted substance to find abnormal cells and a related substance to treat them. In some cancers, the treatment substance carries radiation directly to cells with the target.

## Therapeutic Abortion

Intentional termination of a pregnancy because of medical, fetal, or other legally recognized indications, according to applicable law and clinical guidance.

## Therapeutic Alliance

The working relationship between a patient and a therapist. It includes trust, agreement about treatment goals, and cooperation on the steps of therapy, and can influence how well treatment works.

## Therapeutic Drug Monitoring

The clinical practice of measuring specific drug concentrations in blood at timed intervals (peak and trough levels) to individualize dosages, ensuring efficacy while preventing toxicity for drugs with narrow therapeutic windows (e.g., aminoglycosides, vancomycin, digoxin, lithium, theophylline, phenytoin).

**Alternate Names:**

TDM; Serum Drug Level Monitoring

## Therapeutic Equivalence

Two medicine products are therapeutically equivalent when they contain the same active ingredient, have the same strength and dosage form, meet quality standards, and are expected to have the same clinical effect when used as directed.

## Therapeutic Exercise

Planned physical activity used to improve strength, flexibility, balance, movement, endurance, or daily function and to reduce pain or disability. A clinician selects and gradually adjusts exercises according to the person's condition, abilities, symptoms, and goals.

## Therapeutic Hypothermia

Deliberate, closely monitored cooling of the body to reduce tissue damage in selected emergencies, such as after some cardiac arrests or in severe overheating-related illness. Cooling is used with treatment of the underlying cause and can cause abnormal heart rhythms, shivering, or electrolyte changes.

## Therapeutic Index

A measure of how close a medicine's effective dose is to a dose that causes harm. A narrow index means small dose changes may be risky and require careful monitoring.

## Therapeutics

The part of medicine concerned with treating disease and improving health. It includes medicines, procedures, surgery, rehabilitation, and other treatments used to prevent, control, or cure illness.

## Therapeutic Window

The range of medicine exposure that is high enough to work but low enough to avoid unacceptable harm. Medicines with a narrow window require careful dosing and sometimes blood-level monitoring.

## Therapy

Treatment intended to prevent, relieve, or cure disease or to improve function and quality of life.

---

## Thermal Injury

Tissue damage caused by heat. Initial care includes stopping the exposure, removing loose constricting material, cooling briefly with clean water, covering the injury, and assessing depth and extent. Larger or deeper burns require specialized evaluation, fluid management, pain control, wound care, and tetanus assessment.

### Alternate Names:

Heat Injury; Scald or Thermal Trauma

## Thermal Therapy

Treatment using heat or cold to influence pain, swelling, muscle tension, or tissue temperature. Effects and safety depend on the condition and method used.

## Thermogenesis

The production of heat by the body. It comes mainly from normal metabolism, muscle activity such as shivering, digestion, and specialized fat tissue that releases stored energy as heat.

## Thermography

Recording the pattern of heat released from the body surface with a special camera. It may show temperature differences, but these are not specific for one disease and usually require confirmation with standard examination or tests.

## Thermoluminescent Dosimeter

A small radiation-monitoring device that records accumulated exposure. When the device is heated, its detector releases light in proportion to the radiation it absorbed, allowing the exposure to be measured.

## Thermometer Scales

Numbering systems used to report temperature, such as Celsius, Fahrenheit, and Kelvin. The scale should be identified when a temperature is communicated.

## Thermoneutral Zone

The range of surrounding temperatures in which the body can maintain its normal temperature with-

out shivering or sweating. The range varies with clothing, activity, age, health, and humidity.

## Thermoplasty

A procedure for selected adults with severe asthma that uses controlled heat delivered through a bronchoscope to reduce extra muscle in the airway walls. It may reduce severe attacks but does not cure asthma and has procedural risks.

## Thermoreceptor

A nerve ending that senses warmth or cold in the skin and other tissues. Signals travel through the spinal cord to the brain, helping a person recognize temperature and helping the body control heat through sweating, shivering, and changes in skin blood flow. Damage to these nerves can reduce temperature sensation and increase the risk of burns or cold injury.

## Thermoregulation

The body's control of internal temperature by balancing heat production and heat loss. It uses sweating, shivering, changes in skin blood flow, behavior, and brain signals to keep temperature within a safe range.

## Thiabendazole

An antiparasitic medicine used for selected worm infections and, historically, some fungal infections. It can cause nausea, dizziness, liver problems, or nervous-system effects, so safer alternatives are often preferred.

## Thiamin

Alternate name; see *Thiamine*.

## Thiamine

Vitamin B1, a water-soluble vitamin required for carbohydrate metabolism and nerve function. Severe deficiency can cause beriberi or Wernicke encephalopathy.

### Alternate Names:

Vitamin B1; Aneurin

---

### Thiamine Deficiency

A lack of vitamin B1, needed to release energy from carbohydrates and for normal nerve and heart function. It results from prolonged poor intake or absorption, heavy alcohol use, or increased requirements and causes beriberi, with neuropathy, heart dysfunction, or swelling, and Wernicke encephalopathy when the brain is affected. Prompt thiamine replacement treats or prevents these consequences.

### Thiazide Diuretics

Water tablets that make the kidneys remove more salt and water in urine. They are used mainly for high blood pressure and some fluid-retention disorders and can lower sodium or potassium levels.

### Thiazide Overdose

Taking too much of a thiazide diuretic can cause dehydration, low blood pressure, abnormal sodium or potassium levels, confusion, or abnormal heart rhythms.

### Thiazides

Water tablets that make the kidneys remove more salt and water in urine. They are used mainly for high blood pressure and some fluid-retention disorders and can lower sodium or potassium levels.

### Thiazolidinedione Drugs

A group of diabetes medicines that make body tissues more sensitive to insulin by activating a protein called PPAR-gamma. Pioglitazone is the main example; possible effects include weight gain, swelling, bone fractures, and worsening heart failure.

### Thiazolidinediones

A group of diabetes medicines that make the body more sensitive to insulin, helping lower blood glucose. They can cause weight gain and swelling and may worsen heart failure or increase fracture risk.

### Thigh

The part of the lower limb between the hip and knee. It contains the femur, large muscles, major blood vessels, and nerves that help move and support the leg.

### Thimerosal

A preservative containing ethylmercury that is used in some multidose vaccines and medical products. Ethylmercury leaves the body faster than methylmercury; its safety depends on the product, amount, and applicable regulations.

### ThinPrep Pap Test

A cervical screening test in which cells collected from the cervix are placed in a liquid vial and examined for precancerous changes, cancer, or sometimes HPV-related abnormalities.

#### Alternate Names:

Liquid-Based Pap Test; Liquid-Based Cytology

### Thiopental Sodium

A very short-acting anesthetic given through a vein to start general anesthesia or, in selected critical situations, reduce brain activity and pressure. It can dangerously slow breathing, lower blood pressure, or cause allergic reactions.

### Thiopentone Sodium

A very short-acting anesthetic medicine given into a vein to start general anesthesia in selected settings. It can suppress breathing, lower blood pressure, and cause allergic reactions, so close monitoring is required.

### Thioridazine

An older antipsychotic medicine used only in selected situations because it can cause severe sleepiness, movement problems, and dangerous changes in heart rhythm, especially at high doses or with interacting medicines.

### Thioridazine Overdose

Poisoning from taking too much thioridazine, an antipsychotic medicine. It can cause severe sleepiness, low blood pressure, seizures, and dangerous heart rhythms.

### Thiotepa

A chemotherapy medicine that damages DNA and stops cells from multiplying. It is used for selected

---

cancers and sometimes before a stem-cell transplant; side effects may include low blood counts, infection, nausea, mouth sores, and organ injury.

### **Third-Degree Atrioventricular Block**

A serious heart-rhythm problem in which signals from the upper chambers do not reach the lower chambers. The chambers beat independently, often causing a very slow pulse, fainting, or heart failure. A pacemaker is often used.

#### **Alternate Names:**

Complete Heart Block; Complete AV Block

### **Third-Degree AV Block**

A severe heart-rhythm problem in which electrical signals from the upper heart chambers do not reach the lower chambers. The chambers beat independently, often causing a very slow pulse, dizziness, fainting, or collapse.

### **Third-Degree Burn**

A full-thickness burn that destroys the epidermis and dermis and may extend into subcutaneous tissue. The skin may appear white, brown, or charred and sensation may be reduced. Assessment includes burn depth, extent, inhalation injury, and associated trauma; deep or extensive burns require specialist treatment and may need grafting.

### **Third-Degree Perineal Tear**

A severe tear during vaginal birth that extends from the vaginal opening into the muscles controlling the anus. It needs prompt examination and careful repair, followed by bowel-softening treatment and follow-up; pain, infection, difficulty controlling stool, or a fistula can result.

#### **Alternate Names:**

Third-Degree Obstetric Tear; OASIS

### **Third Heart Sound**

A quiet extra sound heard right after the normal heartbeat, sounding like a gallop; it can be normal in young athletes but often signals heart strain in older adults.

### **Third Stage Of Labor**

The part of childbirth between the baby's birth and delivery of the placenta. The placenta normally separates and is delivered within minutes; heavy bleeding or a placenta that does not deliver may occur with complications.

### **Third Trimester**

The period of pregnancy from 28 weeks until birth. The fetus grows rapidly and continues developing its lungs, brain, and other organs. Prenatal visits monitor blood pressure, growth, movement, and symptoms; tests and screening are scheduled according to local guidance and individual risk.

### **Thirst**

The conscious desire to drink, regulated mainly by body-fluid concentration and circulating blood volume. Excessive thirst may occur with dehydration, diabetes, medicines, kidney problems, or other conditions and may need evaluation.

### **Thomas Test**

A physical examination manoeuvre used to assess for hip flexion contractures and tightness of the iliopsoas muscle. With the patient supine, one hip is fully flexed to flatten the lumbar lordosis; if the contralateral extended thigh lifts off the examination table, a fixed flexion deformity of that hip is present.

### **Thompson Test**

A physical examination test (Simmonds-Thompson test) used to evaluate the integrity of the Achilles tendon; with the patient prone and feet hanging off the examination table, the examiner firmly squeezes the middle third of the calf musculature; absence of involuntary plantarflexion of the foot indicates a complete rupture of the Achilles tendon.

#### **Alternate Names:**

Simmonds Test; Calf Squeeze Test

### **Thomsen Disease**

An inherited muscle disorder causing stiffness because muscles relax slowly after they contract. Symptoms often begin in childhood, affect the



---

hands, legs, eyelids, or jaw, and improve after repeated movement. Strength and life expectancy are usually preserved, although stiffness can interfere with activity.

**Alternate Names:**

Autosomal Dominant Myotonia Congenita; Myotonia Congenita (Thomsen Type)

**Thoracentesis**

A medical procedure in which a thin needle or catheter is gently inserted through the back of the chest wall into the pleural space (the narrow gap between the lungs and the inner chest wall) to withdraw accumulated excess fluid (a pleural effusion). Thoracentesis is performed for two key reasons: diagnostic (collecting a small sample of fluid for laboratory analysis to find the underlying cause, such as pneumonia, heart failure, kidney disease, tuberculosis, or cancer) and therapeutic (removing a large volume of trapped fluid, typically 1 to 1.5 liters, to relieve pressure on a collapsed lung and rapidly ease shortness of breath and chest tightness). Often guided by real-time ultrasound to identify the safest insertion spot, the procedure is carried out under local anesthesia while the patient sits upright, carrying low risks when properly monitored.

**Alternate Names:**

Pleural Tap; Pleurocentesis; Pleural Fluid Aspiration; Thoracocentesis

**Thoracic**

Thoracic means relating to the chest, the thoracic spine, or structures within the thorax such as the heart and lungs.

**Thoracic Aortic Aneurysm**

A weakened, widened area of the aorta within the chest. It may cause no symptoms, but can cause chest or back pain or press on nearby structures; dissection or rupture can cause sudden severe pain, collapse, and life-threatening bleeding. Risk is increased by hypertension, inherited disorders, a bicuspid valve, or smoking.

**Thoracic Cage**

The bony and cartilaginous framework formed by

the thoracic vertebrae, ribs, costal cartilages, and sternum. It protects thoracic organs and assists respiratory movements.

**Thoracic Cavity**

The space inside the chest, enclosed by the ribs and breathing muscles. It contains the lungs, heart, esophagus, trachea, major blood vessels, and other structures protected by the chest wall.

**Thoracic Duct**

The body's largest lymph vessel. It collects excess fluid and immune cells from most of the body and returns them to the bloodstream near the neck; injury can cause lymph fluid to leak into the chest.

**Thoracic Endometriosis**

Endometriosis involving the chest, usually around the lung. Symptoms such as chest pain, breathlessness, or coughing may recur around menstruation and can rarely result in air or blood around the lung.

**Thoracic Inlet**

The upper opening of the chest, surrounded by the first thoracic vertebra, first ribs, and upper sternum. Major nerves, blood vessels, the airway, and the food pipe pass through it.

**Thoracic Outlet**

The lower opening of the chest, bordered by the lower ribs and diaphragm. It allows structures such as the aorta, food pipe, and large veins to pass between the chest and abdomen.

**Thoracic Outlet Syndrome**

Pressure on nerves or blood vessels as they pass from the neck into the arm. It may cause neck, shoulder, or arm pain, tingling, weakness, swelling, color change, or a cold hand. Treatment depends on the cause and may include physical therapy or surgery.

**Alternate Names:**

TOS; Neurovascular Compression of Thoracic Outlet

---

## Thoracic Skeleton

The bony and cartilaginous framework of the chest, made up of the thoracic vertebrae, ribs, sternum, and costal cartilages.

## Thoracic Spine

The middle part of the spine made up of the 12 thoracic vertebrae. It supports the ribs and protects the spinal cord in the chest region.

## Thoracic Spine CT Scan

Computed tomography of the thoracic spine used to evaluate vertebral fractures, alignment, bone lesions, spinal canal narrowing, and other bony abnormalities.

## Thoracic Spine X-Ray

Plain radiographic imaging of the thoracic spine used to assess vertebral alignment, fractures, curvature, degenerative change, and selected bone lesions.

## Thoracic Surgeon

A surgeon who performs operations on organs and structures in the chest, such as the lungs, esophagus, mediastinum, and sometimes the heart.

## Thoracic Vertebrae

The twelve vertebrae of the chest region, each articulating with ribs and protecting the thoracic spinal cord while supporting the trunk and rib cage.

## Thoracolumbar Spine Trauma

Injury to the middle or lower back. Doctors assess the bones, supporting ligaments, spinal cord, and nerves to determine whether the spine is stable and whether treatment needs a brace, monitoring, or surgery.

## Thoracoscopy

A minimally invasive procedure using a camera inserted through the chest wall to examine or treat the pleura, lungs, or mediastinum.

## Thoracotomy

A surgical incision into the chest wall to access the

thoracic cavity (pleural space, lungs, heart, great vessels, esophagus, diaphragm, mediastinum).

## Thorax

The thorax is the chest region between the neck and abdomen, enclosed by ribs and containing the heart, lungs, major vessels, and mediastinal structures.

## Thought Blocking

An abrupt interruption in the train of thought before it is completed, as if the thought was removed or pushed out of the patient's mind. The patient may pause mid-sentence, appear confused, and report that the thought was "taken out" or "blocked." It is a formal thought disorder seen in schizophrenia.

## Thought Disorders

Conditions in which thinking is unusually disorganized, interrupted, fixed on false beliefs, or difficult to express. They can occur in psychosis, mood disorders, brain disease, substance effects, or other conditions and require assessment of the underlying cause.

## Threadworm

A small intestinal worm, also called a pinworm, or the infection it causes. The female worm lays eggs around the anus, commonly causing intense itching at night; infection spreads when eggs are swallowed.

## Threadworm Infection

An intestinal parasitic infection caused by *Enterobius vermicularis*, also called pinworm. Infection spreads when eggs are swallowed, often after contact with contaminated hands, bedding, food, or surfaces.

## Threadworms

Threadworms, also called pinworms, are small intestinal worms that commonly cause itching around the anus at night. Treatment usually includes an anthelmintic medicine and hygiene measures for household contacts.

---

### Threatened Miscarriage

Vaginal bleeding during early pregnancy with a closed cervix and no passage of pregnancy tissue; the pregnancy may remain viable. Evaluation commonly includes clinical assessment, serial human chorionic gonadotropin testing when indicated, and ultrasound.

#### Alternate Names:

Threatened Abortion; Early Pregnancy Bleeding

### Three-Day Measles

Alternate name; see *Rubella*.

### Three-Dimensional Ultrasound

Ultrasound that combines many sound-wave measurements to create a three-dimensional image. It is used in obstetrics, gynecology, and other settings to show anatomy from several angles.

### Three Swab Test

A diagnostic bedside test used to investigate continuous urine leakage from the vagina after pelvic surgery or childbirth. Three cotton swabs are placed inside the vagina and a colored dye (such as methylene blue) is put into the bladder; blue staining on the upper swabs indicates a bladder fistula (vesicovaginal fistula), whereas unstained urine indicates a leak from the ureter (ureterovaginal fistula).

#### Alternate Names:

Moir three-swab test; Three-swab test; Vaginal swab dye test.

### Threonine

An essential amino acid used to build proteins and other body molecules. The body cannot make enough of it, so it must come from food.

### Threshold

The lowest amount of a stimulus or measurement needed to produce a detectable response or effect.

### Thrill

A palpable vibration over the chest or a blood vessel caused by turbulent blood flow.

### Throat

The passage behind the mouth and nose that connects them with the voice box and esophagus. It carries air toward the lungs and food toward the stomach and contains the tonsils and other protective tissues.

### Throat Cancer

Cancer arising in the pharynx or nearby structures of the throat. Persistent hoarseness, difficulty or pain with swallowing, a neck lump, unexplained weight loss, or persistent sore throat requires medical evaluation, especially in people who use tobacco or alcohol.

#### Alternate Names:

Pharyngeal Cancer; Laryngeal Carcinoma

### Throat Or Larynx Cancer

Throat or larynx cancer is malignant growth in the pharynx or voice box and may cause persistent hoarseness, swallowing difficulty, pain, or a neck lump.

### Throat Swab Culture

Culture of a throat specimen to detect bacterial pathogens, especially group A Streptococcus, when evaluating pharyngitis.

#### Alternate Names:

Throat Culture; Pharyngeal Swab Culture

### Throbbing

A pulsating or beating quality of pain or another sensation. It can occur with migraine, inflammation, injury, infection, or increased awareness of a pulse.

### Throckmorton Sign

A radiographic observation (John Thomas sign) in which the shadow of the male penis on an AP pelvis radiograph points toward the side of a unilateral hip or pelvic fracture.

#### Alternate Names:

John Thomas Sign; Penile Orientation Sign

### Thrombectomy

A procedure to remove a blood clot from a blood

---

vessel and restore blood flow. A catheter may remove the clot from a blocked brain artery during some ischemic strokes; surgery is used in selected other situations.

### Thrombin

A blood-clotting substance that changes fibrinogen into fibrin, the protein threads that form a clot, and also activates platelets. The body makes thrombin after a blood vessel is injured; too much or too little clotting can cause disease.

### Thrombin Time

A coagulation assay measuring the time required for plasma to form a fibrin clot after the addition of exogenous bovine or human thrombin, evaluating the final common pathway conversion of fibrinogen to fibrin, prolonged by heparin, dabigatran, hypofibrinogenemia, or dysfibrinogenemia.

#### Alternate Names:

TT; Thrombin Clotting Time

### Thromboangiitis Obliterans

An inflammatory disease of small and medium blood vessels, strongly associated with tobacco use. It can cause pain, color changes, ulcers, or gangrene in the hands or feet. Complete tobacco cessation is essential; further treatment depends on ischemia, tissue loss, and other complications.

#### Alternate Names:

Buerger's Disease; Presenile Gangrene

### Thrombocyte

A small blood-cell fragment made in the bone marrow. Platelets stick to damaged blood vessels, become activated, and join together to form the first plug that stops bleeding. Too few platelets or poorly working platelets can cause easy bruising, tiny skin spots, nosebleeds, or other bleeding. Too many or overly active platelets can increase the risk of unwanted clots.

### Thrombocytes

Another name for platelets: small blood-cell fragments made in the bone marrow that form plugs to stop bleeding. Too few can cause bruising or

bleeding, while too many or overly active platelets can increase the risk of unwanted clots.

### Thrombocythemia

Alternate name; see *Essential Thrombocythemia*.

### Thrombocytopathy

A problem with how platelets work, even when the platelet count is normal or nearly normal. It may cause easy bruising, nosebleeds, heavy menstrual bleeding, or prolonged bleeding after an injury or procedure.

### Thrombocytopenia

A low number of platelets, the blood cells that help stop bleeding. It may result from reduced production, increased destruction, trapping in an enlarged spleen, or dilution and can cause bruising, bleeding, or no symptoms.

#### Alternate Names:

Low Platelet Count

### Thrombocytopenia in Pregnancy

A low platelet count during pregnancy or after delivery caused by gestational thrombocytopenia, immune thrombocytopenia, hypertensive disease, HELLP syndrome, infection, medicines, or another disorder. The cause and trend guide bleeding risk, treatment, anesthesia planning, and fetal evaluation.

### Thrombocytopenic

Characterized by too few circulating platelets. It may cause easy bruising, nosebleeds, heavy menstrual bleeding, or other bleeding, although some people have no symptoms.

### Thrombocytosis

A higher-than-normal number of platelets in the blood. It may be a temporary response to infection, inflammation, bleeding, or iron deficiency, or result from a bone-marrow disorder. It can increase bleeding or clotting risk depending on the cause.

---

## Thromboelastography

A blood test that shows how quickly a clot forms, how strong it becomes, and how it dissolves. It can help guide blood-product treatment during severe bleeding, major surgery, or critical illness.

## Thromboembolism

Obstruction of a blood vessel by a clot that formed elsewhere and traveled through the circulation.

## Thromboembolism in Pregnancy

A blood clot in a vein or artery that may travel to the lungs or another organ during pregnancy or postpartum. Pregnancy increases venous thromboembolism risk; new unilateral swelling, chest pain, sudden breathlessness, coughing blood, or collapse requires emergency assessment.

## Thrombolysis

The dissolution of a blood clot using a medicine such as a fibrinolytic agent. It is used only in selected conditions because it can cause serious bleeding.

## Thrombolysis For PE

Emergency treatment using a clot-dissolving medicine for a life-threatening lung clot causing low blood pressure or shock. It can restore blood flow quickly but carries a serious risk of bleeding.

## Thrombolysis For Stroke

Emergency treatment for selected strokes caused by a blood clot. A clot-dissolving medicine is given through a vein within a limited time after symptoms begin, after brain imaging excludes bleeding; serious bleeding is a possible complication.

## Thrombolytic Agents

Medicines that dissolve an established blood clot by activating the body's clot-breaking system. They are reserved for selected emergencies because they can cause serious bleeding, including bleeding in the brain.

### Alternate Names:

Thrombolytics; Clot-Dissolving Medicines

## Thrombolytic Therapy

Treatment with a medicine that breaks down a blood clot and may restore blood flow in selected emergencies such as certain strokes, heart attacks, or lung clots. The main danger is serious bleeding, including bleeding in the brain.

## Thrombophilia

A tendency for blood to form clots too easily. It may be inherited or develop with cancer, pregnancy, some medicines, inflammation, or other conditions and can increase the risk of clots in veins.

## Thrombophilias in Pregnancy

Inherited or acquired tendencies toward abnormal blood clotting that can increase maternal venous thromboembolism risk and may be associated with placental complications. Testing and anticoagulation decisions depend on personal and family history, the specific thrombophilia, and pregnancy risk.

## Thrombophilia Testing

Blood tests that look for inherited or acquired reasons a person may form blood clots too easily. Testing is considered selectively after a clot or with a strong family history because results may not change treatment.

## Thrombophlebitis

Inflammation of a vein associated with a blood clot. Superficial thrombophlebitis may cause a tender, warm, red, firm cord along a vein just beneath the skin. Thrombosis in a deep vein is deep-vein thrombosis and can break away and cause a life-threatening pulmonary embolism; sudden limb swelling, chest pain, or breathlessness requires urgent assessment.

## Thrombopoiesis

The process by which the bone marrow makes platelets. Platelets develop from large marrow cells and enter the bloodstream, where they help stop bleeding.

## Thrombosed Intracranial Aneurysm

A widened brain artery containing a blood clot. The

---

clot may partly block the aneurysm but can also press on nearby brain structures, causing headache, vision or nerve problems, or seizures.

### **Thrombosis**

Formation of a blood clot inside a blood vessel or the heart. The clot can block blood flow and damage tissue, causing problems such as deep-vein thrombosis, stroke, heart attack, or loss of blood supply to a limb.

### **Thrombosis Of The Inferior Vena Cava**

A blood clot in the large vein that returns blood from the legs and abdomen to the heart. It may cause swelling in both legs or abdominal pain and can extend from a deep-vein clot.

### **Thrombotic Microangiopathy**

A group of disorders in which tiny clots form in small blood vessels, damaging red blood cells and lowering the platelet count. The clots can injure the kidneys, brain, heart, or other organs. Causes include immune disorders, infections, medicines, pregnancy, and severe illness; urgent specialist treatment is often needed.

### **Thrombotic Thrombocytopenic Purpura**

A rare, life-threatening disorder in which small-vessel clots destroy red blood cells and use up platelets. It is usually caused by severe deficiency of ADAMTS13, a protein that helps regulate clotting. It requires urgent hospital treatment.

#### **Alternate Names:**

TTP

### **Thrombotic Thrombocytopenic Purpura Pentad**

The classic, historical clinical pentad characterizing thrombotic thrombocytopenic purpura (TTP), comprising severe microangiopathic hemolytic anemia (with prominent schistocytes), severe thrombocytopenia, fluctuating neurological deficits, acute renal insufficiency, and fever. Pathophysiologically caused by severe deficiency of the von Willebrand factor-cleaving metalloprotease ADAMTS13.

#### **Alternate Names:**

TTP Clinical Pentad; Moschcowitz Pentad

### **Thrombotic Vasculopathy**

Damage to blood vessels caused by clots forming within them. It can reduce blood flow and injure skin or organs and may occur with autoimmune disease, severe infection, cancer, or disorders that cause widespread clotting.

### **Thromboxane**

A chemical made mainly by activated platelets that makes platelets stick together and narrows blood vessels. These actions help stop bleeding but can also contribute to harmful clot formation.

### **Thrombus**

A blood clot that forms inside a blood vessel or the heart and remains attached where it formed. It may partly or completely block blood flow, or break loose and travel elsewhere as an embolus.

### **Thrombus Organization**

The gradual healing of a blood clot as it is replaced by scar tissue and sometimes channels that allow blood to pass through again. This process can make an old clot firm and attached to the vessel wall.

### **Thrush**

A Candida infection, usually of the mouth or throat, causing creamy white plaques, soreness, altered taste, or discomfort with eating or swallowing. Vaginal candidiasis is a separate condition. Risk increases with antibiotics, inhaled corticosteroids, diabetes, or immune suppression.

#### **Alternate Names:**

Oral Thrush; Pseudomembranous Candidiasis

### **Thrush in Children**

Thrush is Candida overgrowth in the mouth, producing white plaques, soreness, or feeding difficulty; Candida can also affect skin folds or diaper areas. Persistent, recurrent, severe, or immune-associated infection needs pediatric evaluation and appropriate antifungal treatment.



---

### Thrush In Newborns

Oral *Candida* infection in an infant, causing white plaques on the tongue or mouth lining that may interfere with feeding and can be passed between infant and breastfeeding parent.

### Thumb Arthritis

Degeneration or inflammation of a thumb joint, especially the joint at the base of the thumb, causing pain, stiffness, and reduced grip.

#### Alternate Names:

Basal Joint Arthritis; CMC Joint Arthritis

### Thumb Carpometacarpal Arthritis

Arthritis of the carpometacarpal joint at the base of the thumb, often due to osteoarthritis. It can cause pain with pinching or gripping, stiffness, and reduced hand function.

### Thumb CMC Arthritis

Arthritis of the joint at the base of the thumb, where the thumb meets the wrist. It causes pain with gripping or pinching, stiffness, swelling, and reduced hand strength; advanced disease may change the joint's shape.

### Thumb-Sucking

A common childhood habit of sucking the thumb; persistent habit can affect dental and facial development.

### Thymic Carcinoma

A malignant epithelial tumor of the thymus that is generally more aggressive and biologically distinct from thymoma. It may invade nearby structures or spread to other sites.

### Thymic Hyperplasia

Thymic hyperplasia is enlargement of the thymus from increased lymphoid tissue or rebound growth and may be incidental. It can be associated with autoimmune disease, infection, endocrine conditions, or stress; imaging and clinical context distinguish it from a mass.

### Thymic Neoplasm

Alternate name; see *Thymoma*.

### Thymine

Thymine is one of the four main bases in DNA. It pairs with adenine through hydrogen bonds and helps store genetic information.

### Thymocyte

An immature T lymphocyte developing in the thymus, where it undergoes selection before becoming a mature T cell.

### Thymoma

A tumor arising from the thymus, an immune-system organ behind the breastbone. It may cause chest pressure or no symptoms and is associated with conditions such as myasthenia gravis; behavior depends on whether it grows beyond the thymus.

#### Alternate Names:

Thymic Epithelial Tumor; Thymic Neoplasm

### Thymus

An immune-system organ behind the breastbone where immature T cells mature and learn not to attack the body's own tissues. It is most active in childhood and becomes smaller after puberty.

### Thymus Gland

An immune-system organ behind the breastbone where immature T lymphocytes learn to recognize harmful foreign material without attacking the body's own tissues. It is most active during childhood and gradually becomes smaller after puberty.

### Thyrocalcitonin

Another name for calcitonin, a hormone made by thyroid C cells that lowers blood calcium mainly by reducing bone resorption.

### Thyroglobulin Surveillance

Blood testing for thyroglobulin after treatment of some thyroid cancers. A rising or unexpectedly detectable level may suggest remaining or returning thyroid tissue, but results must be interpreted with

---

the treatment history and antibody testing.

### **Thyroglossal Duct Cyst**

A fluid-filled lump in the middle of the neck caused by tissue left behind as the thyroid developed before birth. It often moves when swallowing or sticking out the tongue and is usually assessed with ultrasound.

### **Thyroid**

An endocrine gland in the front of the neck that produces thyroid hormones, which regulate metabolism, growth, and development.

### **Thyroid Cancer**

A malignant tumor of the thyroid. Major types include papillary, follicular, medullary, and anaplastic carcinoma, which differ in behavior and treatment.

#### **Alternate Names:**

Thyroid Carcinoma; Malignant Thyroid Neoplasm

### **Thyroid Cancer Staging**

The process of determining how far thyroid cancer has grown or spread. Doctors consider the tumor's size, nearby lymph nodes, distant spread, cancer type, age, and other features to guide treatment and estimate outlook.

### **Thyroid Cartilage**

The largest cartilage of the larynx, formed by two plates that meet in front and create the laryngeal prominence. It protects the vocal folds and provides attachment for muscles and ligaments.

### **Thyroid Diseases**

Conditions affecting the thyroid gland, including underactivity, overactivity, inflammation, enlargement, nodules, and cancer. They may change energy, weight, temperature tolerance, heart rate, bowel habits, mood, or neck appearance and are assessed with examination and blood tests.

### **Thyroid Disorders**

Conditions affecting the thyroid gland, which makes hormones that regulate energy use, temperature,

heart rate, growth, and development. Disorders may cause an underactive or overactive gland, inflammation, enlargement, nodules, or cancer.

### **Thyroidectomy**

Surgery to remove part or all of the thyroid gland. It may treat thyroid cancer, an overactive thyroid, or a large goiter; possible complications include bleeding, voice-nerve injury, and low calcium.

### **Thyroid Eye Disease**

Alternate name; see *Dysthyroid Eye Disease*.

### **Thyroid Fine-Needle Aspiration**

A procedure that uses a thin needle, usually guided by ultrasound, to remove cells from a thyroid lump. A laboratory examines the cells to help determine whether the lump is benign or needs further treatment.

### **Thyroid Follicular Carcinoma**

A cancer arising from thyroid follicle cells and the second most common differentiated thyroid cancer. It tends to spread through blood vessels to the lungs or bones; diagnosis requires showing invasion of the capsule or blood vessels.

### **Thyroid Function Tests**

Blood tests, mainly thyroid-stimulating hormone and free T4, used to assess how well the thyroid is working. Results help identify an underactive or overactive thyroid but must be interpreted with symptoms and medicines.

### **Thyroid Gland**

A butterfly-shaped gland at the front of the neck. It makes hormones that help control energy use, temperature, heart rate, growth, and development.

### **Thyroid Gland Enlargement**

Alternate name; see *Goiter*.

### **Thyroid Gland Removal**

Surgery to remove part or all of the thyroid gland for cancer, overactivity, a large goiter, or a sus-

---

picious nodule. Possible effects include bleeding, voice changes from nerve injury, low calcium, and the need for lifelong thyroid hormone.

### Thyroid Hormone

The hormones T3 and T4, made by the thyroid gland, that help control energy use, temperature, heart rate, growth, and brain development. Too little can slow body functions, while too much can make them overactive.

### Thyroid Hormone Resistance

A rare disorder in which body tissues respond incompletely to thyroid hormones, usually because of an inherited change affecting a thyroid-hormone receptor. Blood levels of thyroid hormones are often high while TSH is not fully suppressed, and symptoms vary between people.

### Thyroid Hormones

Thyroxine (T4) and triiodothyronine (T3), hormones that regulate metabolic rate, growth, brain development, heat production, and cardiovascular and gastrointestinal activity.

### Thyroid Hormone Toxicity

Clinical toxicity caused by excessive thyroid hormone activity, often from excessive replacement or other exogenous exposure. Symptoms may include palpitations, tremor, anxiety, heat intolerance, diarrhea, weight loss, arrhythmia, or severe thyrotoxicosis.

### Thyroid Drugs

Medicines that treat thyroid disorders. Thyroid hormone replacement, usually levothyroxine, treats an underactive thyroid; antithyroid medicines such as methimazole or propylthiouracil reduce hormone production in an overactive thyroid. The choice and monitoring depend on the condition and patient factors.

#### Alternate Names:

Thyroid Medicines; Thyroid Medication

### Thyroiditis

Inflammation of the thyroid gland. It may cause

neck pain, fever, or temporary overactivity followed by underactivity; some forms are painless and autoimmune. Features vary according to the type and duration.

### Thyroiditis Classification

Thyroid inflammation is classified by its cause and pattern. Hashimoto disease is usually long-term and autoimmune; subacute thyroiditis is often painful and follows an infection; other forms may occur after childbirth, medicines, or injury.

### Thyroid Lymphoma

A lymphoma arising in the thyroid, often in the setting of chronic autoimmune thyroiditis. It may present as a rapidly enlarging neck mass with swallowing or breathing difficulty; ultrasound, tissue biopsy, and staging distinguish it from thyroid carcinoma.

### Thyroid Nodule

Alternate name; see *Thyroid Nodules*.

### Thyroid Nodule Evaluation

Assessment of a lump in the thyroid using examination, thyroid blood tests, and ultrasound. Ultrasound features help estimate cancer risk and decide whether repeat imaging or a needle sample is needed.

### Thyroid Nodules

Discrete growths within the thyroid gland that may be solid, cystic, benign, or malignant. Most are benign. They may be associated with rapid growth, hoarseness, swallowing difficulty, or breathing problems, and assessment may include thyroid-function testing, ultrasound, or selected needle biopsy.

#### Alternate Names:

Colloid Nodule; Thyroid Lump; Adenomatous Goiter

### Thyroid Papillary Carcinoma

The most common type of thyroid cancer, arising from cells that make thyroid hormone. It often grows slowly and can spread to nearby lymph nodes; outlook depends on its size, spread, and other tumor features.

---

### Thyroid Peroxidase Antibody

An immune protein directed against thyroid peroxidase, an enzyme needed to make thyroid hormones. A high level supports autoimmune thyroid disease, especially Hashimoto disease, but does not by itself show how well the thyroid is working.

### Thyroid Peroxidase Test

A blood test measuring antibodies against thyroid peroxidase, an enzyme needed to make thyroid hormones. Elevated levels support autoimmune thyroid disease but do not by themselves show how well the thyroid is working.

### Thyroid Preparation Overdose

Taking too much thyroid hormone medicine can cause a fast or irregular heartbeat, shaking, sweating, anxiety, vomiting, fever, or, in severe cases, heart failure or a dangerous thyroid crisis.

### Thyroid Scan

A nuclear-medicine test that shows how actively the thyroid or a thyroid lump takes up a small radioactive tracer. An overactive lump is rarely cancerous; a less-active lump needs assessment with other tests.

### Thyroid Scintigraphy

Another name for a thyroid scan. It uses a small radioactive tracer to show thyroid activity, lumps, or thyroid tissue in an unusual location; results are interpreted with blood tests and ultrasound.

### Thyroid Stimulating Hormone

A pituitary hormone measured in the blood as the primary screening test for thyroid gland disorders. It operates on an inverse feedback loop: when the thyroid is underactive (hypothyroidism), the pituitary releases extra TSH to prompt more hormone production, resulting in a high TSH level. Conversely, when the thyroid produces too much hormone (hyperthyroidism), pituitary TSH output drops, resulting in a low TSH level. It is commonly tested alongside free thyroxine (free T4) to evaluate fatigue, unexplained weight changes, heart rate fluctuations, and to monitor thyroid medication dosing.

### Alternate Names:

TSH; Thyrotropin; Thyroid Function Screen; TSH Blood Test; Thyroid-Stimulating Hormone

### Thyroid-Stimulating Hormone

Alternate name; see *Thyroid Stimulating Hormone*.

### Thyroid Storm

A life-threatening, sudden worsening of an overactive thyroid. It can cause very high fever, a fast or irregular heartbeat, severe agitation or confusion, vomiting, heart failure, and organ failure, often after infection, surgery, or injury.

### Thyroid Storm Diagnostic Criteria

Standardized quantitative clinical scoring criteria (Burch-Wartofsky Point Scale) assessing the likelihood of decompensated, life-threatening thyrotoxicosis based on the severity of thermoregulatory dysfunction, central nervous system effects, cardiovascular tachycardia/arrhythmias, congestive heart failure, and gastrointestinal-hepatic impairment.

### Alternate Names:

Burch-Wartofsky Score

### Thyroid Storm Following Thyroidectomy

Acute, life-threatening decompensation from thyroid hormone excess after thyroid surgery, usually in a person with inadequately controlled thyrotoxicosis. Fever, marked tachycardia, agitation or delirium, heart failure, vomiting, and shock may occur.

### Thyroid Surgery

Operations that remove part or all of the thyroid gland. They may be performed for selected cancers, enlarging or compressive nodules, goiter, hyperthyroidism, or diagnostic uncertainty; the procedure depends on the disease and its extent.

### Thyroid Ultrasound

An imaging test using sound waves to examine the thyroid, thyroid lumps, cysts, and nearby lymph nodes. It can identify features that need monitoring or a needle sample but cannot by itself confirm cancer.

---

### Thyrotoxic Myopathy

Muscle weakness caused by an overactive thyroid, usually affecting the hips, thighs, shoulders, and upper arms. Weakness may occur without pain and can improve when thyroid hormone levels return to normal; severe weakness or swallowing or breathing difficulty needs urgent care.

#### Alternate Names:

Hyperthyroid Myopathy

### Thyrotoxicosis

A condition caused by too much thyroid hormone activity in the body, whatever the source. It may cause weight loss, fast heartbeat, tremor, sweating, anxiety, diarrhea, heat intolerance, or severe illness.

### Thyrotoxic Periodic Paralysis

Repeated brief attacks of severe muscle weakness caused by an overactive thyroid, often with low blood potassium. Episodes commonly affect the hips and shoulders and may follow rest after exercise or a high-carbohydrate meal.

### Thyrotroph Adenoma

A usually noncancerous pituitary tumor arising from cells that make thyroid-stimulating hormone. It may produce too much hormone, causing an overactive thyroid, headaches, or vision problems if it presses on nearby structures.

### Thyrotrophin-Releasing Hormone

A hormone made in the hypothalamus, a part of the brain, that tells the pituitary gland to release thyroid-stimulating hormone and prolactin. It helps link brain signals with thyroid activity and some reproductive functions.

### Thyrotrophin-Stimulating Hormone

Another name for thyroid-stimulating hormone, a pituitary hormone that tells the thyroid gland to make and release thyroid hormones. Blood levels help assess whether the thyroid is underactive or overactive.

### Thyrotropin

Thyroid-stimulating hormone, a pituitary hormone that stimulates thyroid growth and thyroid-hormone production; it is also called TSH.

### Thyrotropin Alfa

A recombinant form of thyroid-stimulating hormone used to stimulate residual thyroid tissue or differentiated thyroid cancer cells during selected diagnostic or treatment protocols.

### Thyroxine

The thyroid hormone T4, which contains four iodine atoms. The thyroid releases it into the blood, and much of it is converted to the more active hormone triiodothyronine (T3).

#### Alternate Names:

T4; Tetraiodothyronine

### Thyroxine-Binding Globulin

A blood protein that carries most thyroid hormones through the bloodstream. Changes in its amount can alter measured total hormone levels without necessarily changing the amount of active, unbound hormone.

### TIA

A brief episode of stroke-like symptoms caused by temporary reduced blood flow to part of the brain, without lasting brain injury on standard assessment. It is an emergency warning of possible stroke.

#### Alternate Names:

Transient Ischemic Attack; Mini-Stroke

### Tibia

The larger, weight-bearing bone of the lower leg, extending from the knee to the ankle. It forms part of both joints and is commonly called the shinbone.

### Tibial Artery

An artery of the lower leg, chiefly the anterior and posterior tibial arteries, supplying the leg, ankle, and foot and contributing to pedal pulses.

---

### **Tibialcrural Artery**

A term describing the arteries and branching vessels near the lower leg and ankle, especially connections between the tibial and fibular arteries. These vessels supply blood to the ankle and foot.

### **Tibial Fracture**

A break in the tibia, the major weight-bearing bone of the lower leg, often caused by high-energy trauma, twisting injury, or repetitive stress.

### **Tibialis Anterior Muscle**

A muscle along the front of the shin that lifts the front of the foot and helps turn it inward. Weakness can cause foot drop and make walking difficult.

### **Tibialis Posterior Muscle**

A deep muscle behind the shin that points the foot downward, turns it inward, and helps support the foot's arch. Damage or weakness may contribute to pain or a gradually flattened foot.

### **Tibial Nerve**

A large nerve running from the back of the knee down the back of the leg and behind the inner ankle. It controls several leg and foot muscles and carries sensation from parts of the foot.

### **Tibial Shaft Fractures**

Breaks in the middle portion of the shinbone, often caused by a high-energy injury but sometimes by a twist or lower-force event. Pain, swelling, deformity, or inability to bear weight are common. Treatment depends on alignment, skin and muscle damage, and whether the fracture is open.

### **Tibia Vara**

Tibia vara is inward angulation of the upper shinbone causing bowing of the leg, classically associated with Blount disease or other growth disorders.

### **Tibiofibular Joint**

A joint between the tibia and fibula bones of the lower leg. The upper joint allows small movements, while the lower connection helps stabilize the ankle.

### **Tic**

A sudden, rapid, recurrent, nonrhythmic movement or sound that a person may feel an urge to make. Tics can be motor or vocal and may worsen with stress.

### **Ticarcillin**

A penicillin antibiotic with activity against many gram-negative bacteria, including *Pseudomonas*; it is usually given with clavulanate and can cause allergy or electrolyte disturbances.

### **Tic Disorder**

A neurologic condition causing repeated, sudden movements or sounds that are difficult to control, such as blinking, shrugging, throat clearing, or grunting. Tics may be briefly suppressed, often worsen with stress, and vary in severity.

### **Tic Douloureux**

A chronic pain condition characterized by recurrent, sudden, severe, electric shock-like pain in the distribution of the trigeminal nerve (typically V2/V3). See Trigeminal Neuralgia.

### **Tick**

A small blood-feeding arachnid that can attach to people or animals and transmit infections such as Lyme disease, rickettsial diseases, and babesiosis.

### **Tick Bite**

A bite from a tick, which may cause local irritation and can sometimes spread infections such as Lyme disease, ehrlichiosis, or Rocky Mountain spotted fever. Remove the tick promptly and watch for fever or a spreading rash.

### **Tick Bites**

Alternate name; see *Tick-borne diseases*.

### **Tick-borne Diseases**

Tick-borne diseases are infections or other illnesses transmitted by ticks, including Lyme disease, babesiosis, ehrlichiosis, and some viral diseases. Removing ticks promptly can reduce risk.



---

### Tick-Borne Encephalitis

A viral infection transmitted mainly by ticks that can cause fever, meningitis, encephalitis, or lasting neurological complications.

### Tickborne Relapsing Fever

An infection caused by certain *Borrelia* bacteria and spread by soft ticks. It causes repeated episodes of fever, headache, muscle aches, and joint pain separated by periods of feeling better.

### Tick Paralysis

Rapidly worsening weakness caused by poison released by an attached tick. Weakness often starts in the legs and moves upward; breathing can become difficult, but symptoms usually improve after prompt tick removal.

### Tick Removal

Removing an attached tick from the skin. The tick is usually grasped close to the skin and pulled steadily; burning, crushing, or covering it with chemicals can increase exposure to fluids.

### Ticks

Small arachnids that attach to skin and may transmit infections or cause tick paralysis.

#### Alternate Names:

Tick Infestation; Ixodid Bites

### Tick Typhus

A rickettsial infection transmitted by ticks, usually causing fever, headache, rash, and sometimes severe systemic illness.

### Ticlopidine

An antiplatelet medicine that reduces the blood's ability to form clots. It was used to prevent stroke or heart-related clots in selected people but is now less common because it can cause serious blood-count and bleeding problems.

### Tics

Brief, repeated movements or sounds that are difficult to control, such as blinking, shrugging, throat

clearing, or grunting. They may be held back for a short time and often worsen with stress; persistent or impairing tics need assessment.

### Tidal Volume

The amount of air breathed in or out with one normal breath. It varies with body size, activity, illness, and breathing support and can be measured during lung-function testing.

### Tidy Wound

A clean, sharply cut wound with little contamination, crushing, or dead tissue and clearly defined healthy edges. After examination and cleaning, it may often be closed immediately, depending on its depth, location, and age.

### Tietze's Syndrome

Alternate spelling; see *Tietze Syndrome*.

### Tietze Syndrome

A related but distinct chest-wall disorder causing painful, visible swelling at one or a few upper costochondral or sternocostal joints. It should not be treated as simply another name for costochondritis.

### Tigecycline

An antibiotic used in selected serious infections when other treatments are unsuitable. It stops bacteria making proteins and commonly causes nausea and vomiting; it is not appropriate for every type of infection.

### Tight Junction

A seal between neighboring lining cells that controls what can pass between them. It helps keep separate the cell surfaces facing the body cavity and the underlying tissue.

### Tilorone

Tilorone is a synthetic antiviral and immunomodulating compound used or studied in some countries; evidence and indications vary by setting.

---

### **Tilted Disc Syndrome**

A congenital appearance in which the optic disc is angled or rotated, often with an unusual shape of nearby retina and blood vessels. Vision may be normal, but some people have visual-field changes or other eye conditions.

### **Tilt Table Test**

A supervised test for unexplained fainting. A person lies secured on a table that is tilted upright while blood pressure, heart rate, and symptoms are recorded to look for fainting caused by abnormal blood-pressure or heart-rate responses.

### **Tilt Table Testing**

A supervised test in which a patient is moved from a lying to an upright or head-up tilt position while blood pressure, heart rate, and symptoms are monitored. It may help evaluate reflex syncope, orthostatic hypotension, or selected autonomic disorders when the diagnosis remains uncertain; it does not replace assessment for cardiac or neurologic causes.

### **Timed Up-And-Go Test**

A functional mobility test measuring the time required to stand from a chair, walk a short distance, turn, return, and sit. Prolonged performance suggests impaired mobility and increased fall risk but must be interpreted with gait, vision, cognition, and environment.

### **Time In Range**

Percentage of time that continuous glucose monitoring readings remain within the target glucose range of seventy to one hundred eighty milligrams per deciliter for most adults with diabetes. General target is greater than seventy percent corresponding to approximately seventeen hours per day.

### **Time Of Death**

The actual or estimated time when a person died. After death, investigators may use body temperature, stiffening, settling of blood, decomposition, insect activity, and chemical changes to estimate it, but the estimate has limits.

### **Time-Of-Flight PET**

A PET scanning method that uses the arrival times of detected radiation to better locate where it came from. This can improve image quality and help show disease more clearly.

### **Timing Of Breastfeeding**

The timing and frequency of breastfeeds, usually guided by early hunger cues and effective milk transfer rather than a rigid clock schedule, with special considerations for newborns and preterm infants.

### **TIMI Risk Score**

A clinical score estimating short-term risk in people with unstable angina or a non-ST-elevation heart attack. It uses age, risk factors, known artery disease, recent symptoms, ECG changes, and heart-injury blood tests to help guide treatment; it supports but does not replace clinical judgment.

#### **Alternate Names:**

TIMI Score; TIMI Risk Score for UA/NSTEMI; TIMI Score for STEMI

### **Timolol**

Timolol is a non-selective beta blocker used in glaucoma eye drops and for selected cardiovascular conditions; topical ocular use can still cause bradycardia, hypotension, bronchospasm, or fatigue.

### **Timolol Maleate**

The maleate salt of timolol, a beta blocker used in glaucoma and selected cardiovascular conditions.

### **Timothy Syndrome**

A rare inherited disorder affecting the heart and several body systems. It can cause a dangerously prolonged heart rhythm, heart defects, webbed fingers or toes, and developmental or immune problems.

### **Tinea**

A fungal infection of the skin, hair, or nails caused by fungi that feed on a protein called keratin. Common forms include ringworm, athlete's foot, and jock itch.

---

### **Tinea Barbae**

A fungal infection of the beard or moustache area. It can cause an itchy, scaly rash, inflamed hair follicles, pus-filled spots, or painful lumps and may spread from infected animals or people.

### **Tinea Capitis**

A fungal infection of the scalp and hair, most common in children. It may cause scaling, itching, broken or falling hair, swollen tender areas, or pus-filled sores and usually needs prescription treatment.

### **Tinea Corporis**

A fungal infection of the body skin, often causing an itchy, scaly, ring-shaped rash with a clearer center. It spreads through contact with infected people, animals, or contaminated objects.

### **Tinea Cruris**

Alternate name; see *Jock itch*.

### **Tinea Faciei**

A fungal infection of the face that can cause a red, scaly, ring-shaped or uneven rash. It may resemble eczema or rosacea and can spread from infected people, animals, or another infected body area.

### **Tinea Favosa**

A long-lasting fungal infection of the scalp or skin that causes yellow, cup-shaped crusts around hair follicles. It can lead to scarring and permanent hair loss if untreated and is now uncommon in many regions.

### **Tinea Imbricata**

A long-lasting fungal skin infection found mainly in parts of the Pacific and Southeast Asia. It causes overlapping, scaly rings that form a distinctive wood-grain pattern on the body.

### **Tinea Incognito**

A fungal skin infection whose appearance has been changed by using a steroid cream or other immune-suppressing treatment. It may become less scaly but more widespread or unusual, so testing may be

needed before targeted treatment.

### **Tinea Infection**

A fungal infection affecting skin, hair, or nails. It may cause a ring-shaped scaly rash, peeling or cracking between the toes, an itchy groin rash, scalp scaling, or thickened discolored nails.

#### **Alternate Names:**

Ringworm Infection; Dermatophyte Mycosis

### **Tinea Manuum**

A dermatophyte fungal infection of one or both hands. It may cause dry scaling, redness, itching, thickened skin, or blisters and often occurs with fungal infection of the feet or nails.

### **Tinea Nigra**

A usually harmless fungal infection causing a painless brown or black patch, most often on the palm or sole. It may resemble a mole or skin cancer and can be confirmed with a skin examination or sample.

### **Tinea Pedis**

A fungal infection of the feet, also called athlete's foot. It may cause itching, peeling or cracking between the toes, dry scaling on the soles, blisters, or sores and can spread to the nails.

### **Tinea Unguium**

A fungal infection of the nail plate, also called onychomycosis, causing thickening, discoloration, brittleness, or separation of the nail.

### **Tinea Versicolor**

A common yeast-related skin condition causing lighter or darker patches with fine scaling, usually on the chest, back, or upper arms. The patches become more noticeable after surrounding skin tans.

#### **Alternate Names:**

Pityriasis Versicolor

### **Tinel Sign**

Tingling or an electric sensation felt along a nerve when the nerve is gently tapped. It may suggest nerve irritation or recovery after injury, but it is not

---

diagnostic by itself and must be interpreted with the examination and other tests.

### **Tinidazole**

Tinidazole is a nitroimidazole antimicrobial effective against anaerobic bacteria and protozoa such as *Giardia* and *Trichomonas*; metallic taste, gastrointestinal effects, neuropathy, and an alcohol interaction may occur.

### **Tinnitus**

Tinnitus is perception of sound without an external source, often related to hearing loss, noise exposure, ear disease, medications, vascular causes, or neurologic factors.

#### **Alternate Names:**

Ringing in Ears; Tinnitus Aurium

### **Tinzaparin**

A low-molecular-weight heparin anticoagulant used for prevention or treatment of venous thromboembolism; bleeding and heparin-induced thrombocytopenia are important risks.

### **Tinzaparin Sodium**

The sodium salt of tinzaparin, a low-molecular-weight heparin used to prevent or treat venous blood clots; bleeding and heparin-induced thrombocytopenia are important risks.

### **Tiopronin**

A thiol-containing medicine used in selected cases of cystinuria to increase urinary cystine solubility; rash, proteinuria, and blood abnormalities may occur.

### **Tiotropium Bromide**

A long-acting inhaled medicine that relaxes airway muscles and keeps the airways open. It is used regularly to control COPD and sometimes asthma; it does not provide immediate relief during a sudden attack.

### **Tip Lesion Variant FSGS**

A distinct histological and clinical subtype of focal segmental glomerulosclerosis (FSGS) characterized

by the presence of at least one segmental lesion involving the tubular pole (origin of the proximal tubule) of the glomerular tuft, characterized by foam cells, podocyte hypertrophy, and endocapillary cellularity. Clinically associated with abrupt-onset full nephrotic syndrome, well-preserved baseline renal function, and favorable response to systemic corticosteroid therapy.

#### **Alternate Names:**

Glomerular Tip Lesion FSGS; Tip Variant FSGS

### **TIPS**

A procedure that creates a channel inside the liver between veins carrying blood into and out of it. The channel lowers high pressure in the portal vein and may treat repeated variceal bleeding or difficult-to-control abdominal fluid.

### **TI-RADS**

An ultrasound scoring system for thyroid lumps. It uses features such as the lump's composition, shape, edges, and calcifications to estimate cancer risk and help decide whether follow-up or needle sampling is needed.

### **Tirbanibulin**

A topical medicine used in short courses to treat actinic keratoses on the face or scalp; local redness, scaling, or irritation may occur.

### **Tired**

Feeling a lack of energy or physical or mental stamina, which may result from sleep loss, exertion, infection, anemia, endocrine disease, medication, depression, or other illness.

### **Tiredness**

Fatigue or reduced energy and endurance that can be physical, mental, or both; persistent unexplained fatigue warrants assessment for sleep, mood, systemic, medication, and metabolic causes.

### **Tirofiban**

An intravenous medicine that temporarily prevents platelets from sticking together. It is used in selected urgent heart conditions or coronary procedures to

---

reduce clotting, but can cause serious bleeding and requires close monitoring.

### **Tissue**

A group of cells and the supporting material around them that work together for a particular function. Examples include muscle, nerve, bone, connective, and lining tissues.

### **Tissue Adhesions**

Bands of fibrous scar tissue that abnormally connect surfaces or organs, often after surgery, inflammation, infection, or bleeding and potentially causing pain, infertility, or obstruction.

### **Tissue Banks**

Facilities that recover, screen, process, preserve, store, and distribute donated human tissue for transplantation, research, or reconstructive treatment under safety and traceability standards.

### **Tissue Congestion**

Accumulation of blood or fluid within tissue because of impaired venous drainage, inflammation, obstruction, or increased vascular flow, producing swelling, redness, or impaired function.

### **Tissue Distribution**

The movement of a medicine or other substance from the bloodstream into body tissues and organs. How much reaches each tissue depends on blood flow, binding, body composition, and the substance's properties.

### **Tissue Elastance**

How strongly a tissue or organ resists being stretched. In the lungs and chest wall, high elastance means more pressure is needed to expand them, which can make breathing harder.

### **Tissue Engineering**

The use of cells, supportive materials called scaffolds, and biologically active substances to create or repair tissues that restore, maintain, or improve body function.

### **Tissue Expander**

A temporary balloon placed beneath the skin and gradually filled with fluid to stretch the skin. The extra skin can later be used for breast, scalp, burn, or other reconstruction.

### **Tissue Expansion**

Gradual stretching of skin or soft tissue with an implanted or external expander so additional tissue can be created for reconstruction; it usually requires staged treatment and monitoring for infection, pain, or device problems.

### **Tissue Kallikrein**

An enzyme made in several tissues that releases chemical messengers affecting local blood flow, blood-vessel leakiness, pain, and inflammation.

### **Tissue Membrane**

A thin layer of tissue that covers a surface, surrounds an organ, or separates spaces in the body. Its structure and function depend on the type of tissue forming it.

### **Tissue Plasminogen Activator**

A natural blood protein that starts the process of breaking down clots. Manufactured forms are used in selected emergencies, such as some strokes, heart attacks, and lung clots, but can cause serious bleeding.

### **Tissue Plasminogen Activator-Associated Angioedema**

Sudden swelling of the face, lips, tongue, or throat after treatment with a clot-dissolving medicine. Throat swelling can block breathing.

### **Tissue Transglutaminase**

An enzyme found throughout the body. In celiac disease, the immune system may make antibodies against it, so a blood test for these antibodies is commonly used during evaluation.

### **Tissue Typing**

Testing tissue antigens, especially HLA, to assess

---

compatibility for transplantation.

### **Titanium**

A strong, lightweight, corrosion-resistant metal used in implants, prostheses, dental devices, and surgical instruments. Bone can form a stable connection with some titanium implant surfaces.

### **Titer**

A laboratory measure showing how much antibody or another detectable substance is present in a sample. It is often reported as the greatest dilution that still produces a measurable reaction; meaning depends on the test and timing.

### **Titration**

Gradually changing a medicine dose to find the amount that gives the desired effect with acceptable side effects, or measuring a substance by reacting it with a solution of known strength.

### **Titre**

A measure of antibody or another substance's concentration, determined by repeatedly diluting a sample until detection stops. Titres can help assess infection, immunity, or response, but interpretation depends on the test and timing.

### **Titubation**

A regular, trembling or nodding movement of the head or body. It can occur when the part of the brain that helps control balance and coordination is diseased.

### **Tizanidine**

A medicine that reduces nerve signals causing muscle tightness and spasms. It is used for selected painful spasms or spasticity but can cause sleepiness, dizziness, low blood pressure, dry mouth, and liver problems.

#### **Alternate Names:**

Zanaflex; Central Alpha-2 Agonist Muscle Relaxant

### **T-Lymphocyte**

A white blood cell that matures in the thymus and

directs cell-based immune responses. Different T cells kill infected cells, coordinate immunity, remember previous threats, or reduce excessive immune activity.

### **Tmax**

The time after a dose when a medicine reaches its highest measured concentration in blood or plasma.

### **TMD**

Alternate name; see *TMJ disorders*.

### **TMJ Disorders**

Temporomandibular joint disorders affect the jaw joints or chewing muscles and may cause jaw pain, clicking, stiffness, headaches, or limited movement.

#### **Alternate Names:**

Temporomandibular Joint Disorders; TMD

### **TNBC**

Alternate name; see *Triple-negative breast cancer*.

### **TNF Alpha**

A chemical messenger made by immune cells that promotes inflammation. It helps fight infection but excessive or prolonged activity can contribute to autoimmune disease, fever, tissue damage, and some cancers.

### **TNM Classification**

A cancer-staging system in which T describes the size or local extent of the primary tumor, N describes involvement of regional lymph nodes, and M describes distant metastasis. The combined findings estimate disease extent, guide treatment, compare outcomes, and inform prognosis; criteria vary by cancer type and staging edition.

### **TNM Classification Of Cancer Stages**

Alternate name; see *TNM Classification*.

### **TNM Staging**

Alternate name; see *TNM Classification*.



---

## **Tobacco**

A plant product containing nicotine and many toxic combustion or processing chemicals. Its use causes dependence and increases the risk of cancer, cardiovascular disease, and lung disease.

## **Tobacco Use**

Tobacco use exposes the body to nicotine and toxic combustion or aerosol chemicals, causing addiction and increased risks of cancer, vascular disease, and lung disease.

## **Tobramycin**

An aminoglycoside antibiotic used for selected serious bacterial infections, including some Gram-negative infections. It can damage the kidneys or hearing and is used with dosing and laboratory monitoring when appropriate.

## **Tocolysis**

Use of medicine to slow contractions for a short time when preterm birth may occur. The delay may allow treatment to help the baby's lungs mature or transfer to a hospital with newborn specialists; it does not usually prevent birth permanently.

## **Tocolytic Agents**

Tocolytic agents suppress uterine contractions temporarily to delay preterm birth, often allowing time for corticosteroids or transfer to specialist care.

## **Tocolytics**

Medicines that temporarily slow contractions before term. They may provide time for corticosteroids, magnesium when indicated, or transfer to specialist care, but their use depends on pregnancy duration, the cause of labor, and risks to the pregnant person or fetus.

## **Tocolytic Therapy**

Medicines used for a short time to slow contractions during threatened preterm birth. The delay may allow time for treatments that improve newborn outcomes or for transfer to a hospital with appropriate neonatal care; they do not usually stop preterm birth permanently.

## **Tocopherol**

A form of vitamin E, a fat-soluble vitamin that helps protect cell membranes. Deficiency is uncommon but may cause nerve problems, muscle weakness, or abnormal red blood cells; excessive supplements can also be harmful.

## **Toddler Feeding Patterns and Diet**

Nutritional and feeding guidance for young children from 6 months to 2 years of age, transitioning from pureed foods to family meals with diverse nutrient-rich textures, iron-rich foods, and healthy mealtime routines.

### **Alternate Names:**

Toddler Nutrition; Weaning and Complementary Feeding; Child Diet 6-24 Months

## **Toddler's Fracture**

A small spiral crack in the lower shinbone of a young child, usually after a minor twist. The child may suddenly limp or refuse to walk, even though there is little swelling or deformity. The first X-ray may look normal; a repeat X-ray after about two weeks may show healing. Treatment commonly uses a supportive boot or cast, based on examination and imaging.

### **Alternate Names:**

Cast-Tibia Fracture; Childhood Spiral Tibial Fracture

## **Toddler Test Or Procedure Preparation**

Preparing a toddler for a test or procedure uses simple explanations, play, distraction, comfort objects, caregiver support, and any required fasting or medicine instructions.

## **Todd Paralysis**

A transient, focal neurological deficit (most commonly hemiparesis, but may include aphasia or neglect) occurring immediately following a focal or secondarily generalized epileptic seizure, typically resolving completely within 15 minutes to 36 hours.

### **Alternate Names:**

Todd's Paresis; Postictal Hemiparesis

---

## Todd Paresis

A transient, focal neurological deficit (most commonly hemiparesis, monoparesis, hemihypesthesia, or aphasia) that follows a focal motor or secondarily generalized epileptic seizure. Symptoms typically resolve spontaneously within minutes to 36 hours; it is hypothesized to arise from localized neuronal metabolic exhaustion, transient lactic acidosis, or sustained Todd post-ictal inhibition.

### Alternate Names:

Todd Paralysis; Post-Ictal Paresis

## Toe-Brachial Index

The ratio between blood pressure measured at a toe and in the arm. It helps assess circulation in the legs, especially when stiff ankle arteries make ankle-pressure testing unreliable.

## Toe Joint

Any joint connecting the bones of a toe. Toe joints include the metatarsophalangeal and interphalangeal joints; they permit toe movement and may be affected by arthritis, gout, injury, or deformity.

### Alternate Names:

Metatarsophalangeal Joint; Interphalangeal Joint of Toe; MTP Joint

## Toenail

A toenail is the hard keratin plate covering the end of a toe and protecting its underlying tissue; injury, fungus, and ingrowth can cause disease.

## Toenail Fungus

Alternate name; see *Nail fungus*.

## Toe Pressure

The blood pressure measured in a toe, useful for assessing circulation in the foot. It can remain reliable when ankle arteries are too stiff to compress, as may occur with diabetes or kidney disease.

## Toes

The five digits at the end of each foot. The big toe has two bones and helps with balance and push-off; the other toes usually have three bones each. Injury,

arthritis, tight shoes, or nerve and muscle problems can cause pain or deformity.

## Togaviridae

A family of viruses that includes some mosquito-borne viruses, such as chikungunya and eastern equine encephalitis virus, which can cause fever, rash, joint pain, or brain inflammation.

## Togaviridae Infections

Infections caused by viruses in the Togaviridae family. Symptoms vary by virus and may include fever, rash, joint pain, or encephalitis.

## Tokophobia

Tokophobia is an intense fear of pregnancy or childbirth that can cause substantial distress or avoidance. Support may include psychological therapy, birth planning, and specialist maternity care.

## Tolazamide

An older sulfonylurea diabetes medicine that increases insulin release; hypoglycemia and weight gain can occur, especially in older adults or people with kidney disease.

## Tolcapone

A medicine used with levodopa for selected people with Parkinson disease to prolong levodopa's effect. It can cause diarrhea, involuntary movements, confusion, and potentially fatal liver injury, so liver monitoring is required.

## Tolerable Upper Intake Level

The highest average daily intake of a nutrient unlikely to cause harm in almost all healthy people. It includes food, drinks, and supplements and is not a recommended amount to consume.

### Alternate Names:

UL

## Tolerance

A decreased response to a drug after repeated administration, requiring dose escalation to achieve the same effect. Unlike tachyphylaxis, tolerance

---

develops over weeks to months of chronic use.

### **Toll-Like Receptor**

A protein on certain immune cells that recognizes common features of germs or damaged tissue. Activating these receptors helps start inflammation and other defenses against infection.

### **Toll-Like Receptors**

Receptors of the innate immune system that recognize characteristic molecules from microbes or damaged tissue. Their activation starts inflammatory and antimicrobial responses.

**Alternate Names:**

TLRs

### **Tolmetin**

A nonsteroidal anti-inflammatory drug used for pain and arthritis; gastrointestinal bleeding, kidney injury, and cardiovascular events are important risks.

### **Tolmetin Overdose**

Taking too much tolmetin, a nonsteroidal anti-inflammatory medicine, can cause stomach bleeding, vomiting, kidney injury, drowsiness, seizures, or other serious effects.

### **Tolnaftate**

A topical antifungal medicine used for some dermatophyte infections such as athlete's foot and ringworm; it is not effective for every type of fungal infection.

### **Tolosa-Hunt Syndrome**

A rare neuro-ophthalmic disorder characterized by severe, unremitting, painful unilateral periorbital/retro-orbital pain accompanied by ipsilateral ophthalmoplegia (palsies of cranial nerves III, IV, and VI) and sensory loss in the V1 distribution, caused by idiopathic granulomatous inflammation within the cavernous sinus or superior orbital fissure, exquisitely responsive to systemic corticosteroid therapy.

**Alternate Names:**

Painful Ophthalmoplegia; Cavernous Sinus Granu-

lomatous Syndrome

### **Tolterodine Tartrate**

Tolterodine tartrate is an antimuscarinic medicine used for overactive bladder and urinary urgency; dry mouth, constipation, and urinary retention can occur.

### **Toluene**

Toluene is a volatile solvent that can irritate tissues and, with substantial inhalation or ingestion, injure the nervous system, liver, kidneys, or heart.

### **Toluene And Xylene Poisoning**

Poisoning from solvents containing toluene or xylene, often through swallowing or inhalation. It can cause dizziness, confusion, abnormal heart rhythms, breathing problems, or organ injury.

### **Toluene Toxicity**

Solvent poisoning from inhalation or ingestion that can cause CNS depression, arrhythmias, renal tubular acidosis, hypokalemia, weakness, and altered mental status. Check electrolytes, acid-base status, renal function, and ECG; remove the exposure and provide supportive care while correcting life-threatening potassium or acid-base abnormalities.

### **Tolvaptan**

A vasopressin V2-receptor antagonist used in selected patients with hyponatremia or autosomal dominant polycystic kidney disease; thirst, rapid sodium correction, and liver injury require attention.

### **Tomogram**

A tomogram is an image of a selected cross-sectional layer of the body or another object, produced by tomography such as CT or MRI.

### **Tomography**

Imaging that produces cross-sectional views, or slices, through the body. Computed tomography uses X-rays, while other forms use ultrasound, magnetic fields, or radioactive tracers.

---

## Tomosynthesis

An imaging test that takes several low-dose X-ray pictures from different angles and combines them into thin layers. It is commonly used to examine breast tissue in three dimensions.

## Tomotherapy

Tomotherapy combines CT imaging with rotating, precisely shaped radiation beams to deliver radiation treatment while adjusting for the target and nearby tissues.

## Tongue

A muscular organ in the mouth that helps move food, form speech sounds, swallow, and detect taste. Swelling, sores, weakness, numbness, or restricted movement may result from injury, infection, inflammation, nerve disease, or cancer.

## Tongue Biopsy

A tongue biopsy removes a small tissue sample from a tongue lesion for microscopic examination of infection, inflammation, precancer, or cancer.

## Tongue Carcinoma

Tongue carcinoma is cancer of the tongue, usually squamous-cell cancer, and may cause a persistent sore, lump, pain, or difficulty speaking or swallowing.

## Tongue Disease

Any disorder affecting the tongue, such as infection, inflammation, injury, nerve disease, nutritional deficiency, or cancer. Symptoms may include pain, sores, swelling, altered taste, or difficulty speaking or swallowing.

## Tongue Neoplasm

An abnormal growth in the tongue that may be non-cancerous, precancerous, or cancerous. A persistent lump, ulcer, pain, bleeding, numbness, or difficulty speaking or swallowing needs examination and may require biopsy.

## Tongue Problems

Changes in the tongue, such as pain, swelling, sores, color change, numbness, or difficulty moving or tasting. Causes include injury, infection, inflammation, nutritional deficiency, and other disease.

## Tongue-Tie

A congenital condition characterized by an abnormally short, thick, or tight lingual frenulum that tethers the bottom of the tongue tip to the floor of the mouth, restricting tongue mobility. In infants, it can cause latching difficulty and painful breastfeeding; in older children, it may affect speech articulation, eating, or dental hygiene. Severe restriction may be treated with a simple surgical release (frenotomy).

### Alternate Names:

Ankyloglossia; Lingual Frenulum Tethering; Short Lingual Frenulum; Tongue Tie

## Tonic-Clonic Seizure

A seizure causing loss of awareness, first with stiffening of the body and then with repeated jerking movements. Sleepiness, confusion, headache, or muscle soreness commonly follows.

## Tonic Neck Reflex

A normal early-infant reflex, sometimes called the fencing posture. When the infant's head turns to one side, the arm on that side straightens and the opposite arm bends. It usually fades by about six months; marked asymmetry may suggest a neurologic problem.

### Alternate Names:

Asymmetric Tonic Neck Reflex; ATNR; Fencing Reflex

## Tonic Pupil

A pupil that reacts poorly to light but constricts slowly when focusing on something nearby. It often results from injury to nerves controlling the pupil and should be assessed with other eye and neurologic findings.

## Tonics

Preparations traditionally claimed to improve

---

strength, energy, appetite, or general health. Their ingredients, safety, and evidence vary widely, and some may interact with medicines or contain harmful amounts of active substances.

### **Tonic Seizures**

Seizures causing sustained stiffening of muscles, usually for seconds to minutes. They may affect one area or the whole body, cause falls or breathing difficulty, and require individualized seizure evaluation.

### **Tonometer**

An instrument that measures pressure inside the eye. Eye-pressure readings help detect glaucoma or monitor treatment, but they must be interpreted with the optic nerve, visual field, cornea, and other findings.

### **Tonometry**

Measurement of pressure inside the eye, usually to assess for glaucoma or monitor its treatment. Methods include applanation, noncontact air-puff, rebound, and electronic devices; results must be interpreted with corneal thickness and the clinical context.

### **Tonsil**

A small mass of immune tissue at the back of the throat. Tonsils help detect germs entering through the mouth and nose but can become enlarged or infected, causing sore throat, difficulty swallowing, or breathing problems.

#### **Alternate Names:**

Palatine Tonsil; Tonsilla

### **Tonsillar Cancer**

Cancer arising in a tonsil, most often from the surface lining cells. It may cause a persistent sore throat, pain felt in the ear, trouble swallowing, a tonsil that is larger on one side, or a lump in the neck.

### **Tonsillar Neoplasm**

A tonsillar neoplasm is an abnormal growth in a tonsil that may be benign or malignant and can cause a persistent sore throat, swallowing difficulty,

or neck swelling.

### **Tonsillectomy**

Surgery to remove the tonsils, performed for selected repeated infections, obstructive sleep apnea, or other tonsil disease after the expected benefits and risks are assessed.

### **Tonsillitis**

Inflammation or infection of the tonsils, usually caused by a virus and sometimes by group A streptococcal bacteria. It can cause sore throat, fever, swollen neck glands, painful swallowing, and white patches.

### **Tonsil Stones**

Small, hard lumps formed when food particles, mucus, dead cells, and bacteria become trapped and calcify in the folds of the tonsils. They are often harmless but may cause bad breath, throat irritation, a feeling that something is stuck in the throat, recurrent inflammation, or visible white or yellow deposits.

### **Tonus**

The normal resting tension or partial contraction of a muscle or tissue. Abnormally low or high tone can occur with nerve, muscle, brain, spinal-cord, or developmental disorders.

### **Tooth**

A hard mineralized structure anchored in the jaw, used for biting, tearing, chewing, and grinding food.

### **Tooth Abrasion**

Loss of tooth substance caused by external mechanical friction, such as aggressive brushing, abrasive products, or habits, producing grooves, notches, or sensitivity.

### **Tooth Abscess**

A tooth abscess is a localized collection of pus from bacterial infection of the tooth or surrounding tissues, causing pain, swelling, and possible spreading infection.

---

## **Toothache**

Pain arising from a tooth or nearby gum, ligament, bone, or other supporting tissue. Common causes include decay, infection, cracks, trauma, and gum disease.

## **Toothaches**

Alternate name; see *Toothache*.

## **Tooth Anatomy**

A tooth has a hard outer covering called enamel, a supporting layer called dentin, and a living center called the pulp that contains nerves and blood vessels. The root anchors it in the jaw through surrounding tissues.

## **Tooth Ankylosis**

Fusion of a tooth root to alveolar bone through loss of the periodontal ligament, preventing normal eruption and making the tooth appear infraoccluded as neighboring teeth grow.

## **Tooth Attrition**

Wear of tooth surfaces caused by tooth-to-tooth contact, commonly from chewing, bruxism, or malocclusion, producing flattened cusps and possible sensitivity.

## **Tooth Avulsion**

Complete displacement of a tooth from its socket, usually caused by an injury. The tooth, socket, periodontal ligament, and surrounding tissues may all be damaged; permanent teeth are more commonly affected.

## **Tooth Brushing**

Tooth brushing removes plaque and food from teeth and gums. Brushing twice daily with fluoride toothpaste helps prevent tooth decay and gum disease.

## **Tooth Calcification**

Deposition of mineral that forms and hardens enamel, dentin, cementum, and other tooth structures during development; abnormal timing or amount can affect tooth strength and appearance.

## **Tooth Component**

Tooth components include enamel, dentin, cementum, pulp, and supporting tissues, each contributing to tooth strength, sensation, nutrition, and attachment.

## **Tooth Crowding**

Tooth crowding occurs when teeth lack enough space to align normally, often from tooth size, jaw size, eruption pattern, or retained primary teeth.

## **Tooth Crown**

The tooth crown is the visible portion above the gum, covered mainly by hard enamel and containing underlying dentin and pulp.

## **Tooth Decay**

Gradual destruction of tooth structure caused when mouth bacteria make acids from sugars. It can cause sensitivity, pain, holes, infection, and tooth loss, but fluoride, cleaning, diet, and dental care help prevent progression.

### **Alternate Names:**

Dental Caries; Cavities

## **Tooth Discoloration**

Abnormal tooth color can result from plaque, decay, trauma, fluorosis, medicines, enamel defects, or staining, and the cause determines treatment.

## **Tooth Disease**

Any disorder affecting a tooth, including decay, infection, cracks, developmental defects, or wear. It may cause pain, sensitivity, discoloration, swelling, or difficulty chewing and should be assessed by a dental professional.

## **Tooth Enamel**

Tooth enamel is the highly mineralized outer covering of the crown that protects dentin and pulp from wear, acid, and bacteria.

## **Tooth Erosion**

Tooth erosion is the chemical loss of tooth enamel caused by acids, not by bacteria. Acidic foods,



---

drinks, reflux, and repeated vomiting can contribute to it.

### **Tooth Eruption**

The process by which a developing tooth moves through the gum into the mouth. Baby teeth usually appear from about 6 months to 3 years, and permanent teeth from about 6 years onward; timing varies and delays may need assessment.

### **Tooth Extraction**

Removal of a tooth from its socket because of severe decay, infection, periodontal disease, fracture, impaction, orthodontic planning, or another clinical indication.

### **Tooth Fracture**

A crack or break in a tooth caused by injury, biting hard material, decay, grinding, or weakened enamel. It may expose the inner nerve, causing pain, sensitivity, infection, or loss of the tooth.

### **Tooth Infection**

Infection of the tooth, its pulp, or surrounding tissues, commonly caused by untreated decay, cracks, or gum disease. It may cause toothache, sensitivity, swelling, fever, or drainage. Facial swelling, difficulty swallowing or breathing, or spreading infection requires emergency assessment.

### **Tooth Loss**

Tooth loss is absence of a tooth due to decay, gum disease, trauma, developmental absence, or extraction and can affect chewing, speech, and oral health.

### **Tooth Luxation**

Injury that loosens or moves a tooth without completely knocking it out. The tooth may be pushed inward, outward, or sideways and needs prompt dental assessment to protect the tooth and surrounding tissues.

### **Toothpaste Overdose**

Swallowing too much toothpaste can cause stomach upset, vomiting, or diarrhea. Large amounts of flu-

oride toothpaste can cause serious salt and calcium disturbances, especially in children.

### **Tooth Replantation**

Replacement of an accidentally displaced or avulsed tooth into its socket, followed by stabilization and dental follow-up.

### **Tooth Resorption**

Breakdown of part of a tooth, usually starting inside the tooth or on its root. It may follow injury, infection, orthodontic treatment, or occur without a known cause and can weaken or destroy the tooth.

### **Tooth Root**

The tooth root anchors a tooth in its socket and contains dentin, cementum, pulp, nerves, and blood vessels beneath the gum.

### **Tooth Sensibility Testing**

Dental tests using cold, heat, electricity, or touch to see whether a tooth's nerve responds. The results do not directly measure blood flow inside the tooth and must be combined with examination and dental imaging.

### **Tooth Sensitivity**

Brief pain caused by cold, heat, sweet or acidic foods, brushing, or touch. It often occurs when protective enamel or gum tissue is worn away and the inner dentin is exposed; persistent sensitivity needs dental assessment.

### **Tophaceous Gout**

Advanced, long-lasting gout in which firm lumps of uric-acid crystals build up in and around joints. These lumps, called tophi, may develop on the toes, fingers, ears, or elbows and damage nearby tissue.

### **Tophus**

A firm lump made of uric-acid crystals that collects in or around a joint or in soft tissue after long-standing gout. Tophi can cause pain, inflammation, skin breakdown, deformity, and damage to bone or cartilage.

---

## Topographic Spinal Cord Anatomy

The organization of the spinal cord into segments, gray matter, white-matter tracts, roots, enlargements, and vascular territories. This anatomy explains patterns of motor, sensory, reflex, and autonomic loss after cord injury or disease.

## Topical

Applied directly to skin or a moist lining, usually to act at the treatment site with limited whole-body exposure.

## Topical Antibiotics

Antibiotics applied to the skin to treat selected superficial bacterial infections. The choice and duration depend on the organism, site, extent of infection, resistance patterns, and local guidance. Unnecessary or prolonged use can cause irritation and antimicrobial resistance.

## Topical Antifungals

Medicines applied to the skin to treat fungal infections such as athlete's foot, ringworm, or jock itch. The product must match the fungus and body site; irritation or treatment failure can occur.

## Topical Calcineurin Inhibitors

Non-steroid creams or ointments that reduce immune inflammation in eczema, especially on the face, eyelids, or skin folds. Burning or itching may occur briefly after application.

## Topical Corticosteroid Potency

The strength of a steroid cream, ointment, lotion, or gel used on the skin. Mild products are used for sensitive areas or children, while stronger products may be needed briefly for thick or severe inflammation. Choice depends on the diagnosis, body site, age, duration, and risk of skin thinning or other effects.

## Topical Retinoids

Vitamin-A-related medicines applied to the skin for acne and some sun-related skin changes. They help prevent blocked pores and increase skin-cell turnover but may cause dryness, irritation, and increased sun

sensitivity; pregnancy precautions vary by product.

## Topiramate

An antiseizure medicine also used to prevent migraine. It may cause tingling, slowed thinking, weight loss, kidney stones, or acid imbalance and can harm a developing fetus; pregnancy planning and medicine changes require medical advice.

## Topoisomerase

An enzyme that temporarily cuts and rejoins DNA to control its twisting during replication, transcription, and chromosome separation.

## Topoisomerase II

An enzyme that cuts both DNA strands, passes another DNA segment through the break, and rejoins the strands; it is targeted by some anticancer drugs.

## Topoisomerase Inhibitors

Drugs that interfere with DNA topoisomerases; some are antibiotics and others are anticancer medicines, with toxicity depending on the drug and target enzyme.

## Topotecan

A topoisomerase-I inhibitor used for selected ovarian, small-cell lung, and cervical cancers; bone-marrow suppression is a major toxicity.

## Topotecan Hydrochloride

A salt form of topotecan, a chemotherapy medicine that interferes with DNA copying. It treats selected cancers but can cause dangerously low blood counts, infection, bleeding, diarrhea, nausea, or fatigue.

## TORCH Panel

A group of blood tests used when pregnancy or newborn findings suggest infection acquired before birth. TORCH traditionally refers to Toxoplasma, other infections, rubella, cytomegalovirus, and herpes; individual tests and antibody results need careful clinical interpretation.

---

## **TORCH Screen**

Testing for selected congenital infections—classically Toxoplasma, Other infections, Rubella, Cytomegalovirus, and Herpes—when fetal or newborn findings suggest an in-utero infection; broad panels require careful interpretation.

## **Toremifene**

A selective estrogen-receptor modulator used for metastatic breast cancer in postmenopausal women; hot flashes and, rarely, dangerous QT prolongation may occur.

## **Torn ACL**

A torn anterior cruciate ligament is a knee injury commonly caused by sudden pivoting, landing, or contact, producing a pop, swelling, instability, and difficulty changing direction. Rehabilitation, bracing, or reconstruction is selected according to activity, associated injury, and goals.

## **Torn Meniscus**

A torn meniscus is a split in the knee's shock-absorbing cartilage caused by twisting or degeneration. Pain, swelling, catching, or locking may occur; physical therapy or arthroscopic treatment depends on tear pattern, symptoms, age, and associated disease.

### **Alternate Names:**

Meniscal Tear; Torn Semilunar Cartilage

## **Torpor**

A state of greatly reduced activity, alertness, or responsiveness. In a patient, marked or new torpor may signal a serious illness, medicine effect, or metabolic problem.

## **Torsade De Pointes**

A dangerous, fast heart rhythm in which the lower chambers beat in a twisting pattern. It is often linked with a prolonged electrical recovery time, abnormal body salts, certain medicines, or inherited heart conditions and can cause collapse or cardiac arrest.

## **Torsades De Pointes**

A dangerous, fast heart rhythm arising from the lower chambers and often associated with a prolonged QT interval. It may cause dizziness, fainting, or cardiac arrest and requires urgent treatment and correction of the cause.

## **Torsemide**

A water tablet that helps the kidneys remove salt and water. It treats swelling caused by heart, kidney, or liver disease but can cause dehydration, low blood pressure, or abnormal body-salt levels.

## **Torsion**

Twisting of a body structure around its supporting axis, which may reduce or stop blood flow. Torsion of the testicle, ovary, or bowel can destroy tissue and often requires emergency surgery.

## **Torsion Dystonia**

A movement disorder in which repeated or sustained muscle contractions twist part of the body into abnormal postures. It may affect one area or many areas and can be inherited or develop after illness, injury, or medicine exposure.

## **Torsion of Appendix Testis**

Twisting of a small leftover tissue tag near the top of a testicle, usually in a child. It can cause local scrotal pain and a small blue spot, but sudden scrotal pain must be assessed urgently to rule out twisting of the testicle itself.

### **Alternate Names:**

Hydatid of Morgagni Torsion; Blue Dot Sign

## **Torsion of Testicular Appendage**

Twisting and infarction of a vestigial appendage attached to the testis or epididymis, usually in children. Localized upper-pole pain and a tender blue-dot sign may occur, but testicular torsion must be excluded.

## **Torsion Of The Appendix Testis**

Twisting of a tiny leftover structure attached to the upper part of the testicle. It is common in

---

boys before puberty and causes sudden, localized scrotal pain and tenderness; a small blue spot may be visible through the skin. The testicle usually still has normal blood flow. Rest, support, and pain medicine usually allow it to heal, but urgent assessment is needed to exclude testicular torsion.

**Alternate Names:**

Hydatid of Morgagni Torsion; Appendix Testis Torsion

**Torticollis**

A condition in which the neck muscles pull the head into a tilted or rotated position. It may be present from birth or result from pain, injury, muscle spasm, nerve disease, or another condition.

**Alternate Names:**

Cervical Dystonia; Loxia

**Torulaspora**

A genus of yeasts used in food and fermentation; human infection is rare and usually opportunistic.

**Torus Fracture**

A stable incomplete break in a child's bone caused by compression, often in the wrist after a fall. The bone bulges rather than breaking all the way through; it usually heals well with a splint or cast and follow-up as advised.

**Alternate Names:**

Buckle Fracture; Metaphyseal Buckle Fracture; Torus Fracture of Radius

**Torus Mandibularis**

A harmless, usually painless bony lump on the tongue side of the lower jaw, commonly near the premolar teeth. It usually needs no treatment unless it is repeatedly injured or interferes with dentures.

**Torus Palatinus**

A harmless, usually painless bony lump along the middle of the roof of the mouth. It grows slowly and usually needs no treatment unless it becomes irritated or interferes with eating or dentures.

**Total Abdominal Colectomy**

Surgery to remove the entire colon through an abdominal incision or keyhole approach. The small intestine may be joined to the rectum or brought through the abdominal wall as an ileostomy, depending on the disease and safety.

**Total Anomalous Pulmonary Venous Connection**

Alternate name; see *Total anomalous pulmonary venous return*.

**Total Anomalous Pulmonary Venous Return**

A birth defect in which the veins returning oxygen-rich blood from the lungs connect to the wrong side of the heart. Babies may have blue skin, rapid breathing, poor feeding, or heart failure. Severe obstruction is an emergency, and surgery is needed to reroute the veins.

**Total Artificial Heart**

A mechanical device that replaces the pumping function of both ventricles and is used in selected patients with severe biventricular heart failure, usually as a bridge to transplantation. It requires specialized center-based management and carries risks including bleeding, infection, thrombosis, and device malfunction.

**Total-Body Skin Examination**

A systematic examination of the skin, scalp, nails, and visible mucous membranes from head to toe to look for new, changing, or suspicious lesions. It may include dermoscopy and is especially useful for people at increased risk of skin cancer.

**Total Body Surface Area**

The percentage of the skin affected by a burn or another widespread process. Clinicians estimate it using the person's palm or body-area charts to judge severity and guide treatment.

**Total Breech Extraction**

A delivery procedure in which a clinician brings out a breech baby by the feet. It is mainly used for a

---

second twin in selected circumstances, not routinely for a single full-term breech baby because of injury risks.

### **Total Cholesterol**

Total cholesterol is the combined cholesterol carried in major blood lipoproteins; cardiovascular risk is assessed with HDL, LDL, triglycerides, and other factors.

### **Total Daily Energy Expenditure**

The total energy the body uses in one day, including energy needed at rest, digestion, movement, and physical activity. It varies with body size, age, muscle mass, health, and activity level.

### **Total Haemolytic Complement Test**

A blood test that measures whether a group of immune proteins can work together to destroy cells. An abnormal result may suggest immune deficiency, autoimmune disease, infection, inflammation, or a laboratory problem.

### **Total Hip Arthroplasty**

Surgery that replaces a damaged hip joint with artificial parts. It is used for severe arthritis, fractures, or other joint damage when pain and loss of movement continue despite other treatment.

### **Total Hysterectomy**

Removal of the uterus and cervix. The fallopian tubes and ovaries are not necessarily removed and should be specified separately, such as salpingectomy or oophorectomy.

### **Total Intravenous Anesthesia**

General anesthesia given entirely through medicines injected into a vein rather than inhaled gases. It produces controlled unconsciousness during surgery, while trained anesthesia staff continuously monitor breathing, blood pressure, and other vital functions.

### **Total Iron-Binding Capacity**

A clinical laboratory blood test measuring the maximum concentration of iron that can be bound by

circulating serum transferrin molecules, elevated in iron deficiency anemia and decreased in anemia of chronic disease, hemochromatosis, and malnutrition.

#### **Alternate Names:**

TIBC

### **Total Iron Binding Capacity**

A blood-test estimate of how much iron the protein transferrin can carry. It is interpreted with blood iron, ferritin, blood counts, and other findings to assess iron deficiency, inflammation, or iron overload; it does not measure stored iron directly.

### **Total Knee Arthroplasty**

Surgery that replaces damaged surfaces of the knee with artificial parts to relieve severe pain and improve movement. It may be recommended for advanced arthritis or major joint damage after other measures are inadequate; rehabilitation and monitoring help restore function and detect complications.

### **Total Knee Replacement**

Surgery that replaces damaged surfaces of the knee with artificial parts to relieve severe pain and improve function. Recovery includes exercises and gradual activity; infection, blood clots, stiffness, loosening, persistent pain, or repeat surgery may occur.

#### **Alternate Names:**

Total Knee Arthroplasty

### **Total Laryngectomy**

Total laryngectomy removes the entire voice box and creates a permanent opening in the neck for breathing, usually to treat advanced laryngeal cancer.

### **Total Lesion Glycolysis**

A PET-scan measurement combining the amount of active tumor tissue with how much radioactive sugar-like tracer it absorbs. It estimates the tumor's overall activity and may help monitor treatment, but does not determine outlook alone.

---

### Total Lung Capacity

The greatest amount of air the lungs can hold after a full breath in. Lung-volume testing measures it; the amount may be too high when lungs are overinflated or too low when the lungs or chest cannot expand normally.

### Total Pancreatectomy

Total pancreatectomy removes the entire pancreas and causes permanent loss of insulin and digestive-enzyme production, requiring lifelong replacement treatment.

### Total Parenteral Nutrition

Nutrition given through a vein when the digestive system cannot absorb enough food. The solution can provide water, sugar, protein, fat, vitamins, and minerals, but requires monitoring for infection, blood-sugar changes, and liver or electrolyte problems.

### Total Proctocolectomy And Ileal-Anal Pouch

Total proctocolectomy with an ileal-anal pouch removes the colon and rectum and creates a pouch from the ileum that is connected to the anus.

### Total Proctocolectomy With Ileostomy

Total proctocolectomy with ileostomy removes the colon and rectum and diverts the end of the small intestine through an abdominal stoma.

### Total Protein

The combined amount of albumin and other proteins in the blood. A blood test may help assess nutrition, liver or kidney disease, inflammation, or abnormal antibody production.

### Total Sleep Time

The total amount of time spent asleep during a defined sleep period or sleep study, excluding periods of wakefulness.

### Total Thyroidectomy

The complete surgical resection of both lobes and

the isthmus of the thyroid gland, indicated for aggressive thyroid malignancies, compressive multinodular goiters, or medically refractory Graves disease, carrying specific risks of recurrent laryngeal nerve injury and permanent hypoparathyroidism.

### Touch

The sensory perception of contact, pressure, texture, vibration, or movement across the skin. Signals travel through sensory nerves to the spinal cord and brain.

### Tourette Disorder

A condition beginning in childhood or adolescence that causes repeated movements and sounds, called tics, for at least one year. Tics may include blinking, shrugging, throat clearing, or words and often change over time.

### Tourette's

Alternate name; see *Tourette syndrome*.

### Tourette's Syndrome

A neurodevelopmental disorder involving multiple motor tics and at least one vocal tic persisting for more than one year.

### Tourette Syndrome

A neurodevelopmental disorder characterized by multiple motor tics and at least one vocal tic occurring over at least one year. Tics are sudden, rapid, recurrent, stereotyped movements or vocalizations. They are often preceded by premonitory urges. Tics typically worsen in adolescence and may improve in adulthood.

### Alternate Names:

Tourette's Disorder; Gilles de la Tourette Syndrome

### Tourniquet

A band or pneumatic device placed around a limb to temporarily occlude or greatly reduce arterial blood flow. Emergency tourniquets are designed for control of severe limb bleeding, while pneumatic surgical tourniquets create a relatively bloodless operative field. Prolonged inflation can cause nerve, muscle, or skin injury.



---

### **Tourniquet Test**

A bedside test that looks for small skin spots after brief pressure from a blood-pressure cuff. It may show increased capillary fragility or low platelets, but it is not specific and cannot diagnose dengue or another disease by itself.

#### **Alternate Names:**

Rumpel-Leede Test; Hess Test; Capillary Fragility Test

### **Towel Clamp**

A surgical clamp used to secure drapes or hold materials in place during an operation. Its sharp points must be applied carefully because they can puncture skin or damage underlying tissue.

### **Townes Syndrome**

A rare inherited birth disorder that commonly causes blockage or absence of the anal opening and unusual thumbs. Kidney, heart, hearing, foot, or genital abnormalities may also occur. The specific problems vary, so treatment is planned around each child's findings.

### **Toxaphene**

A persistent organochlorine insecticide mixture that can accumulate in soil, water, and body fat. Significant exposure may overstimulate the nervous system and cause tremors, seizures, liver injury, or other toxic effects.

### **Toxbase**

A clinical poison-information service that helps healthcare professionals assess chemical or medicine exposures. It provides evidence-based advice about expected symptoms, necessary tests, treatment, observation, and urgency for particular substances.

### **Toxic**

Poisonous or capable of harming living cells, tissues, organs, or the whole body. Toxicity depends on the substance, dose, route, duration, and individual susceptibility.

### **Toxic Adenoma**

A single, non-cancerous (benign) thyroid lump that produces too much thyroid hormone on its own, independent of the body's normal hormonal controls. This causes an overactive thyroid (hyperthyroidism) with symptoms such as a rapid or irregular heart-beat, unexplained weight loss, shaky hands, sweating, anxiety, and heat intolerance, but without the bulging eye signs seen in Graves' disease. Blood tests show very low pituitary thyroid-stimulating hormone (TSH) and high thyroid hormone levels, while a radioactive thyroid scan characteristically shows a single bright "hot nodule" that absorbs tracer while the rest of the thyroid remains quiet. Definitive treatment involves radioactive iodine therapy to destroy the overactive lump or surgery to remove the affected half of the thyroid gland (lobectomy).

#### **Alternate Names:**

Toxic Thyroid Adenoma; Autonomous Thyroid Nodule; Plummer Disease; Hot Thyroid Nodule

### **Toxic Anterior Segment Syndrome**

A sterile, noninfectious inflammatory reaction in the anterior eye after surgery, caused by toxic substances or contaminants introduced during the procedure. Severe corneal edema, anterior chamber inflammation, and elevated pressure may occur.

### **Toxic Dose**

The amount of a substance that causes harmful effects. It varies with the substance, route, duration of exposure, age, health, and individual susceptibility.

### **Toxic Effect**

Harmful change caused by a chemical, medicine, poison, radiation, or other exposure. The effect may involve one or more organs and depends on the dose, route, duration, metabolism, and individual susceptibility.

### **Toxic Encephalopathy**

Brain dysfunction caused by a poison, chemical, or medicine. It may cause confusion, unusual behavior, memory or movement problems, seizures, or reduced alertness and needs assessment for the exposure and

---

its effects.

### Toxic Epidermal Necrolysis

A rare, life-threatening reaction usually to a medicine in which widespread painful skin and moist body linings blister and peel. It can cause fluid loss, infection, and organ problems and requires emergency specialist care.

**Alternate Names:**

TEN

### Toxicity

The ability of a substance, treatment, or exposure to cause harmful effects in a living organism. Severity depends on the dose, route, duration, metabolism, target organs, and individual susceptibility; effects may be temporary or permanent.

### Toxicity Grade

A standardized rating of how severe a harmful effect is, often used for adverse effects of medicines or cancer treatment. Grades generally range from mild to life-threatening and help guide monitoring and management.

### Toxic Megacolon

A life-threatening widening of the large intestine during severe bowel inflammation or infection. It may cause severe abdominal swelling and pain, fever, rapid heart rate, dehydration, or shock and requires urgent hospital treatment.

### Toxic Metabolic Encephalopathy

Temporary or lasting brain dysfunction caused by a poison, medicine, organ failure, infection, or abnormal body chemistry. It may cause confusion, unusual sleepiness, agitation, seizures, or coma and requires treatment of the underlying problem.

### Toxic Neuropathy

Peripheral-nerve dysfunction caused by a toxic exposure, medicine, metabolic toxin, or substance-use-related injury. Symptoms may include burning pain, numbness, tingling, weakness, or impaired balance.

### Toxic Nodular Goiter

An enlarged thyroid containing one or more nodules that make too much thyroid hormone. It can cause weight loss, sweating, shaking, fast or irregular heartbeat, anxiety, or heat intolerance and is assessed with blood tests and imaging.

### Toxicodendron Dermatitis

An intensely pruritic, erythematous, papulovesicular, and linear contact dermatitis resulting from cutaneous exposure to urushiol resin found in poison ivy, poison oak, or poison sumac.

**Alternate Names:**

Urushiol Dermatitis; Rhus Dermatitis; Poison Ivy Rash

### Toxicodendron Poisoning

Urushiol-induced allergic contact dermatitis from poison ivy, poison oak, or poison sumac. An intensely pruritic, often linear vesicular eruption appears after a delayed interval and can spread through contaminated clothing, tools, or animal fur; severe facial, genital, widespread, or inhalational reactions need medical treatment.

### Toxicodendron Species

Plant species belonging to the genus *Toxicodendron* that contain the lipophilic antigenic resin urushiol, which triggers a type IV cell-mediated allergic contact dermatitis upon epidermal contact.

**Alternate Names:**

Toxicodendron Plants; Urushiol-Containing Plants; Poison Ivy Oak Sumac Species; Poison Ivy; Poison Oak; Poison Sumac

### Toxicokinetic Description

An account of how a harmful substance enters the body, is absorbed, distributed, changed by metabolism, and eliminated. These processes determine the concentration and duration of exposure at target organs.

### Toxicology

The study of poisons and harmful effects of chemicals, medicines, and other substances. It includes how exposure happens, how the body is affected,

---

how poisoning is recognized, how it can be prevented, and how it is treated.

### **Toxicology Screen**

A test of urine, blood, or another sample for medicines, drugs, alcohol, or poisons. Results must be interpreted with the person's symptoms, timing, and prescribed medicines.

### **Toxicology Specimen Preservation**

Collection, labeling, storage, and transport of biological specimens to preserve analytes and prevent contamination or degradation. Appropriate specimens may include peripheral blood, urine, vitreous humor, bile, liver, hair, or gastric contents depending on the suspected exposure and timing.

### **Toxic Shock Syndrome**

A rare, life-threatening illness caused by toxins made by certain bacteria. It can cause sudden fever, a widespread rash, low blood pressure, vomiting or diarrhea, and failure of organs such as the kidneys or heart.

### **Toxic Synovitis**

Temporary inflammation of the lining of the hip joint, usually in a child after a viral illness. It can cause hip pain and limping and must be distinguished from a serious joint infection.

### **Toxidrome**

A recognizable group of symptoms and examination findings suggesting poisoning by a particular type of substance. Examples include opioid, stimulant, sedative, and cholinergic patterns; the pattern helps guide urgent treatment but does not identify the exact poison by itself.

### **Toxin**

A poisonous substance made by a living organism, such as a bacterium, plant, animal, or fungus. Toxins can damage specific tissues or disrupt nerve, muscle, or cellular functions.

#### **Alternate Names:**

Biological Poison; Endotoxin or Exotoxin

### **Toxocara**

A genus of parasitic roundworms that commonly infect dogs and cats. People acquire infection by swallowing eggs from contaminated soil or food; migrating larvae can cause fever, enlarged organs, or eye or nervous-system disease.

### **Toxocara Canis**

A dog roundworm whose eggs can infect people. Larvae migrate through tissues and may cause toxocariasis, with fever, cough, enlarged liver, eye inflammation, or rarely nervous-system involvement.

### **Toxocara Cati**

A cat roundworm whose eggs can infect people. The larvae may migrate through organs and cause toxocariasis, including fever, cough, enlarged liver, eye inflammation, or rarely neurologic disease.

### **Toxocariasis**

Infection caused by dog or cat roundworm larvae after a person swallows eggs from contaminated soil, food, or hands. The larvae can migrate through the body and cause fever, cough, enlarged liver, eye inflammation, or rarely brain disease.

### **Toxoid**

A bacterial toxin that has been chemically or otherwise inactivated so it cannot cause its usual harm but can still stimulate protective immunity. Toxoids are used in vaccines against diseases such as tetanus and diphtheria.

### **Toxoid Vaccines**

Vaccines made from bacterial toxins that have been detoxified so they cannot cause their usual harm but can train the immune system to neutralize the toxin. Examples protect against tetanus and diphtheria.

### **Toxoplasma**

A genus of intracellular parasites; *Toxoplasma gondii* is the species that infects people. Infection is often mild, but it can seriously affect a fetus or people with weakened immunity, especially the brain and eyes.

---

### **Toxoplasma Blood Test**

A blood test detecting antibodies or genetic material related to *Toxoplasma gondii* infection, interpreted with symptoms, immune status, pregnancy, and timing of exposure.

### **Toxoplasma Chorioretinitis**

Alternate name; see *Ocular Toxoplasmosis*.

### **Toxoplasma Gondii**

A parasite that causes toxoplasmosis. Cats are its main host and can spread infective forms in their stool; people may also acquire it from undercooked meat, contaminated soil, or contaminated food and water.

### **Toxoplasma Meningoencephalitis**

Alternate name; see *Toxoplasmosis*.

### **Toxoplasmosis**

An infection caused by the single-celled parasite *Toxoplasma gondii*. People may acquire it by eating undercooked contaminated meat or shellfish, swallowing the parasite from cat feces or contaminated soil, or consuming contaminated food or water. Most people with a healthy immune system have no symptoms; others may have swollen lymph nodes, fever, tiredness, or muscle aches. Infection can cause serious disease of the eyes or brain in people with weakened immunity, may reactivate after immune suppression, and can pass from a newly infected pregnant person to the fetus. Person-to-person spread is otherwise uncommon but can occur through an infected organ or blood transfusion.

#### **Alternate Names:**

Toxoplasma Gondii Infection; Ocular Toxoplasmosis

### **Toxoplasmosis Pathology**

Tissue injury caused by *Toxoplasma gondii*, often showing necrotizing inflammation, tachyzoites, tissue cysts containing bradyzoites, and variable lymphohistiocytic or mixed inflammatory infiltrates.

### **TPA**

Tissue plasminogen activator, a natural clot-

dissolving protein or a laboratory-made medicine based on it. In selected emergencies it helps break down clots, but it can cause serious bleeding.

#### **Alternate Names:**

Tissue Plasminogen Activator; Fibrinolytic Enzyme

### **TPMT Polymorphism**

An inherited difference in the thiopurine methyltransferase enzyme, which breaks down some immune-suppressing medicines. Low enzyme activity can cause dangerously low blood counts unless dosing is reduced or another treatment is chosen.

### **TPN**

TPN, or total parenteral nutrition, supplies nutrients intravenously when the gastrointestinal tract cannot be used adequately.

### **Trabectedin**

A chemotherapy medicine that attaches to DNA and disrupts cancer-cell growth and repair. It is used for selected advanced soft-tissue sarcomas and ovarian cancers and can cause low blood counts, liver injury, nausea, or muscle damage.

### **Trabecula**

A small beam, strut, or partition of tissue, especially within spongy bone. Trabeculae form a supportive internal network that helps bones bear forces while keeping them relatively light.

### **Trabecular Bone**

Light, honeycomb-like bone found inside many bones and near their ends and joints. It supports the body, contains bone marrow, and changes quickly when weight, hormones, medicines, or diseases such as osteoporosis affect bone strength.

#### **Alternate Names:**

Spongy Bone; Cancellous Bone

### **Trabecular Meshwork**

A sieve-like tissue near the front of the eye through which the clear fluid called aqueous humor drains. If the drainage pathway becomes blocked, pressure inside the eye may rise and damage the optic nerve

---

in glaucoma.

### **Trabeculectomy**

An operation for glaucoma that creates a new drainage path for fluid inside the eye. Lowering eye pressure can protect the optic nerve, but bleeding, infection, scarring, or pressure that becomes too low can occur.

### **Trabeculoplasty**

Laser treatment for some forms of glaucoma that improves drainage of the clear fluid inside the eye. Better drainage can lower eye pressure and reduce the risk of further optic-nerve damage.

### **Trace Elements**

Minerals needed in very small amounts for normal enzyme function, blood formation, thyroid activity, and other body processes.

### **Trace Mineral**

A mineral needed in very small amounts for health. Examples include iron, zinc, copper, iodine, and selenium; too little or too much can impair blood formation, thyroid function, immunity, growth, or other body processes.

### **Tracer**

A detectable substance used to follow a biological, chemical, or physical process. Medical tracers can show blood flow, organ function, metabolism, or disease location through imaging or laboratory testing.

### **Trachea**

The windpipe connecting the larynx to the right and left main bronchi. Cartilage rings help keep it open so air can move to and from the lungs.

### **Tracheal Agenesis With Bronchoesophageal Fistula**

A very rare birth defect in which the windpipe is absent or severely underdeveloped and an abnormal passage connects the breathing system with the food pipe. Newborns have extreme breathing difficulty and need immediate specialist care.

### **Tracheal Foreign Body**

An object stuck in the windpipe. It can cause choking, noisy breathing, severe breathing difficulty, or complete blockage and is an emergency requiring immediate removal by trained clinicians.

### **Tracheal Rupture**

A tear in the windpipe caused by severe injury, surgery, a medical instrument, or pressure inside the airway. It can let air leak into the chest and cause breathing difficulty, infection, or life-threatening complications.

### **Tracheal Stenosis**

Narrowing of the windpipe, often after prolonged breathing-tube use, tracheostomy, injury, or inflammation. It may cause worsening breathlessness, noisy breathing, or wheezing and can require airway procedures or surgery.

### **Tracheal Tug**

A downward, rhythmic traction or pulling sensation on the cricoid cartilage and trachea synchronized with cardiac systole, felt by placing the examiner's fingers under the cricoid cartilage while the patient extends the neck. It is a physical sign of an aneurysm of the aortic arch downwardly displacing the left main bronchus.

#### **Alternate Names:**

Oliver Sign; Cardarelli Sign

### **Tracheitis**

Inflammation or infection of the windpipe. Mild cases may occur with a viral illness, but bacterial infection can rapidly block a child's airway and cause high fever, a harsh cough, noisy breathing, and severe illness. It requires emergency airway and antibiotic treatment.

### **Trachelectomy**

Trachelectomy is surgical removal of the cervix, sometimes with nearby tissue, used in selected cervical cancers or other conditions.



---

## Tracheobronchial Injury

Traumatic rupture or laceration of the cervical trachea or major bronchi, usually occurring within 2 cm of the carina following severe blunt thoracic deceleration trauma, presenting with massive subcutaneous emphysema, pneumomediastinum, and a continuous large air leak through a thoracostomy tube.

### Alternate Names:

TBI

## Tracheobronchomalacia

Weakness of the windpipe or larger airways that allows them to collapse during breathing. It may cause noisy breathing, chronic cough, repeated infections, or difficulty clearing mucus and may be present from birth or develop later.

## Tracheobronchomegaly

Unusual widening of the windpipe and the main airways, usually because their supporting tissue is weak. It can cause a weak cough, breathlessness, repeated chest infections, and difficulty clearing mucus.

## Tracheoesophageal Fistula

An abnormal opening between the windpipe and food pipe, usually present from birth. It can let milk enter the lungs, causing coughing, choking, blue coloring, repeated chest infections, or poor growth, and usually needs surgery.

## Tracheoesophageal Fistula And Esophageal Atresia Repair

Surgery in a newborn to close an abnormal connection between the windpipe and food pipe and join separated parts of the food pipe when possible. The operation is urgent; breathing problems, leakage, narrowing, reflux, and feeding difficulties may persist.

## Tracheomalacia

Weakness of the windpipe wall that allows it to narrow or collapse during breathing out. It may cause noisy breathing, cough, breathlessness, repeated infections, or difficulty clearing mucus and may be

present from birth or develop later.

## Tracheostomy

A surgically created artificial opening (stoma) on the front of the neck leading directly into the trachea (windpipe), through which a curved plastic or metal breathing tube (a tracheostomy tube) is placed. It allows air to enter the lungs directly, bypassing severe upper airway obstructions (such as from throat swelling, facial trauma, vocal cord paralysis, or tumors), facilitates the suctioning of deep mucus, and provides a secure route for long-term mechanical ventilation when patients cannot breathe independently. A tracheostomy may be temporary (removed once the underlying condition resolves) or permanent, and requires diligent daily care and suctioning to keep the tube patent and prevent airway obstruction or infection.

### Alternate Names:

Trach; Surgical Stoma of the Trachea; Tracheostomy Stoma

## Tracheostomy Care

Care of a breathing tube and neck opening, including cleaning the skin and tube, suctioning mucus when needed, checking airflow, and keeping emergency equipment nearby. Blockage, bleeding, infection, or accidental tube displacement requires prompt attention.

## Tracheostomy Tube

A tube inserted through a tracheostomy opening to maintain access to the airway and permit breathing, suction, or mechanical ventilation.

## Tracheotomy

A surgical procedure in which an incision (cut) is made through the skin of the neck into the windpipe (trachea) to create a direct airway. It is performed in planned or emergency situations to overcome acute upper-airway obstruction (from foreign bodies, severe infections, or head and neck trauma) or to facilitate prolonged breathing machine support. While “tracheotomy” refers to the surgical act of making the incision into the trachea, the resulting opening and airway passage created by the opera-



---

tion is called a tracheostomy.

**Alternate Names:**

Tracheal Incision; Surgical Airway Procedure

**Trachoma**

A chronic eye infection caused by certain types of *Chlamydia trachomatis*. It spreads through contact with infected eye or nose discharge, contaminated items, or, in some areas, flies. Repeated infections can scar the inside of the eyelid, turn the eyelid inward, and cause the eyelashes to rub against the eye. This can scar the cornea and lead to permanent visual impairment or blindness.

**Trachyonychia**

Rough, thin, ridged nails with many tiny pits, sometimes called twenty-nail dystrophy when most or all nails are affected. It may occur on its own or with psoriasis, alopecia areata, eczema, or lichen planus.

**Traction**

A controlled pulling force used to align or immobilize a fracture, reduce a dislocation, or relieve pressure on tissues. It may be applied manually, with weights, or through a medical device.

**Traction Alopecia**

Mechanical hair loss resulting from sustained, repetitive tension pulling on hair follicles over prolonged periods (from tight braids, ponytails, hair weaves, or extensions), initially causing reversible perifollicular erythema before progressing to irreversible follicular scarring and permanent patchy alopecia.

**Tractional Retinal Detachment**

Separation of the light-sensing retina from the back of the eye when bands of scar tissue pull it away. It commonly occurs in advanced diabetic eye disease and can cause vision loss.

**Tractotomy**

Surgery that cuts a selected nerve pathway in the brain or spinal cord, usually to reduce severe pain or another difficult neurologic symptom. It can cause permanent changes in sensation or movement.

**Traditional Medicine**

Traditional medicine includes health practices, remedies, and healing systems developed within cultural traditions. Products and practices vary in evidence and safety, so they should not replace effective care without professional advice.

**Tra Gene**

A gene involved in sex determination and sexual differentiation in some organisms, especially insects; its function and sequence vary by species.

**Trained Immunity**

A lasting change in the body's first-line immune response after an earlier infection or exposure. The later response may be stronger or altered, although this process does not provide the same specific protection as ordinary immune memory.

**Train-Of-Four Monitoring**

A test used during anesthesia that gives four brief electrical signals to a nerve and measures the muscle responses. It shows how much a muscle-relaxing medicine remains active and whether muscle strength has recovered.

**Trajectory**

The path a projectile follows in flight and, in wound ballistics, the course it traverses through tissue from entrance to exit or rest. Internal trajectory is reconstructed from entrance and exit defects, the organ track, recovered fragments, and clothing damage; angles disclose shooter position and victim posture.

**TRALI**

A rare, life-threatening lung reaction that develops during or within several hours after a blood transfusion. It causes sudden breathing difficulty and low blood oxygen from fluid leaking into the lungs, without being caused by excess fluid in the circulation.

**Tram-Track Appearance**

A double-line pattern seen under a microscope in some kidney biopsies. It reflects thickening or splitting of the kidney's filtering membranes and can

---

occur in certain types of glomerulonephritis.

### **Trance**

An altered state of awareness with unusual attention, responsiveness, or sense of surroundings. It may occur during hypnosis, seizures, intoxication, sleep-related events, or psychological states and requires context for interpretation.

### **Tranexamic Acid**

A medicine that helps control bleeding by preventing the body from breaking down blood clots too quickly. It is used for selected heavy bleeding, including some surgical, injury-related, menstrual, and postpartum bleeding.

### **Transabdominal Cerclage**

Placement of a strong band around the upper cervix through abdominal surgery, usually before pregnancy or early in pregnancy, to reduce the risk of premature opening in severe cervical weakness. It requires cesarean birth and carries surgical, bleeding, infection, and pregnancy risks.

#### **Alternate Names:**

TAC; Cervico-Isthmic Cerclage

### **Transabdominal Pelvic Ultrasound**

A non-invasive diagnostic ultrasound examination of the pelvic organs (uterus, endometrium, ovaries, fallopian tubes, and urinary bladder) performed transabdominally through a full acoustic bladder window.

#### **Alternate Names:**

Transabdominal Pelvic Sonography; TA Pelvic Ultrasound; Transabdominal Pelvic Scan

### **Transabdominal Scan**

An ultrasound scan performed by moving the probe over the abdomen. It can examine pelvic organs and, during pregnancy, assess the baby's growth and anatomy, the placenta, and the amount of fluid around the baby.

### **Transaminitis**

Higher-than-normal levels of liver-related enzymes

in the blood. This may indicate liver-cell injury, but the result does not identify the cause; medicines, infection, alcohol, fatty liver, and muscle injury can contribute.

### **Transarterial Chemoembolization**

A treatment in which a catheter delivers chemotherapy and blocks blood flow to a liver tumor. It can slow or shrink selected tumors that cannot be removed surgically, but may cause pain, fever, liver injury, or other complications.

### **Transbronchial Biopsy**

Removal of a small lung-tissue sample using a camera passed through the airways. It can investigate infection, inflammation, cancer, or rejection after transplant; bleeding and a collapsed lung are possible complications.

### **Transcatheter Aortic Valve Replacement**

Replacement of a narrowed aortic heart valve using a folded artificial valve delivered through a catheter, usually from an artery in the groin. It avoids open-heart surgery for selected people, but bleeding, stroke, blood-vessel injury, rhythm problems, leakage, or need for a pacemaker can occur.

#### **Alternate Names:**

TAVR; TAVI; Transcatheter Aortic Valve Implantation

### **Transcatheter Edge-To-Edge Repair**

A catheter procedure that joins parts of the mitral-valve leaflets to reduce leakage. It may help selected people with severe symptoms who are not good candidates for open surgery; bleeding, infection, stroke, valve narrowing, residual leakage, or device problems are possible.

#### **Alternate Names:**

TEER; MitraClip Procedure; Percutaneous Mitral Valve Repair

### **Transcatheter Pulmonary Valve**

Replacement of a diseased pulmonary valve with a prosthetic valve delivered through a catheter. It is used in selected congenital or previously operated right-ventricular outflow tracts.

---

### **Transcervical Foley Catheter For Cervical Ripening**

A soft balloon catheter placed through the cervix and inflated to help soften and open the cervix before labor induction. It is a mechanical method and may cause cramping, bleeding, infection, or accidental rupture of the membranes.

### **Transcervical Resection Of Endometrium**

A camera-guided procedure through the cervix that removes or destroys the uterine lining to reduce heavy menstrual bleeding. It is not suitable for future pregnancy and requires careful evaluation beforehand.

### **Transcranial Doppler**

An ultrasound test that measures blood-flow speed in arteries inside the skull. It can help detect narrowed vessels, spasm after brain bleeding, or increased stroke risk in selected conditions.

### **Transcranial Doppler Scanning**

An ultrasound test measuring blood-flow speed and direction in major arteries inside the skull. It can help assess narrowing, blockage, vasospasm, emboli, or circulation after certain brain injuries.

#### **Alternate Names:**

Transcranial Doppler; TCD

### **Transcranial Doppler Ultrasound**

An ultrasound test that measures blood-flow speed and direction in major arteries inside the skull. It can help assess stroke risk, vessel narrowing, bleeding-related spasm, and blood flow changes.

### **Transcranial Magnetic Stimulation**

A noninvasive treatment or brain-function test that uses magnetic pulses near the scalp to induce electrical activity in selected brain regions. Repetitive stimulation is used for some depression and other conditions; effects and risks depend on the protocol and person's health.

### **Transcription**

The process by which a cell copies information from

DNA into RNA. The RNA copy can direct protein production or help control cell activity.

### **Transcription Factor**

A protein that attaches to DNA and increases or decreases the copying of selected genes into RNA. Transcription factors help control cell growth, development, specialization, and responses to hormones or injury.

### **Transcutaneous Electrical Nerve Stimulation**

TENS: low-voltage electrical pulses delivered through electrodes on the skin to reduce pain for some people. It does not treat every cause of pain and should be used according to clinical guidance.

#### **Alternate Names:**

TENS

### **Transdermal**

Delivering a medicine through the skin, usually with a patch or gel, so it enters the bloodstream gradually. The medicine, dose, skin condition, heat, and application site affect how much is absorbed.

### **Transesophageal Echocardiography**

An ultrasound scan of the heart made with a probe passed into the esophagus. It gives close views of the heart, valves, and nearby blood vessels.

### **Trans Fat**

An unsaturated fat containing at least one trans-configured double bond. Industrial trans fats, formerly common in partially hydrogenated oils, can raise LDL cholesterol, lower HDL cholesterol, and increase cardiovascular risk; small amounts also occur naturally in some meat and dairy foods.

### **Trans Fatty Acid**

An unsaturated fat with a trans shape, often formed during industrial processing and associated with higher heart disease risk.

#### **Alternate Names:**

Trans Fat; Hydrogenated Fat

---

## Transferase

An enzyme that moves a chemical group from one molecule to another. Transferases help cells make and break down substances, and some transferase levels in blood help identify tissue injury or disease.

## Transference

Feelings or expectations that a patient developed in an earlier important relationship and now directs toward a therapist or another healthcare professional. Discussing this pattern can help explain relationship difficulties during therapy.

## Transferrin

A blood protein made mainly by the liver that carries iron to tissues. Its level and the amount of iron it carries help assess iron deficiency, inflammation, and iron overload when interpreted with other blood tests.

## Transferrin Saturation

The percentage of transferrin's iron-binding sites occupied by iron. It is calculated from iron and total iron-binding capacity and helps evaluate iron deficiency and iron overload.

### Alternate Names:

TSAT

## Transfer Training

Teaching a person how to move safely between places, such as from a bed to a chair or toilet. Training may include correct body position, assistance from another person, a transfer board, or lifting equipment.

## Transforming Growth Factor-Beta

A multifunctional cytokine superfamily that orchestrates wound healing, extracellular matrix synthesis, collagen deposition, tissue fibrosis, cell cycle arrest, and immunosuppression by inhibiting epithelial and lymphocyte proliferation while stimulating myofibroblasts.

### Alternate Names:

TGF-beta

## Transfusion

Delivery of donated blood or blood components into a person's bloodstream. It can replace lost red cells, platelets, or clotting factors but carries risks including reactions, infection, fluid overload, and lung injury.

## Transfusion Of Neonate

Administration of a blood product to a newborn for clinically significant anemia, bleeding, or another specific indication. The decision, product characteristics, volume, and rate depend on the infant's symptoms, laboratory results, gestational age, and local transfusion protocol.

## Transfusion Reaction

A harmful response to donated blood or blood components. It may cause fever, chills, rash, breathing difficulty, pain, low blood pressure, or destruction of red cells. The transfusion must be stopped and the person assessed urgently if a reaction is suspected.

## Transfusion-Related Acute Lung Injury

A life-threatening complication of blood product transfusion developing within six hours of transfusion, caused by donor anti-HLA or anti-neutrophil antibodies activating recipient neutrophils in the pulmonary microvasculature, producing non-cardiogenic pulmonary edema and acute hypoxemic respiratory failure.

### Alternate Names:

TRALI

## Transient Acantholytic Dermatitis (Grover Disease)

An itchy eruption of small red or skin-colored papules and vesicles caused by temporary loss of cohesion between epidermal cells. It commonly affects the trunk of older adults and may flare with heat, sweating, or dry skin.

## Transient Familial Hyperbilirubinemia

An inherited tendency for bilirubin, a yellow waste product, to rise temporarily in the blood and cause episodes of jaundice. It usually reflects slower processing by the liver without major liver injury and

---

may worsen during illness, fasting, or stress.

### **Transient Global Amnesia**

A sudden temporary inability to form new memories, often with repeated questions and uncertainty about recent events. The person remains awake and knows who they are; symptoms usually resolve within 24 hours, but urgent assessment is needed to exclude stroke or seizure.

#### **Alternate Names:**

TGA; Acute Amnestic Syndrome

### **Transient Hypogammaglobulinemia Of Infancy**

A temporary delay in an infant's production of antibodies, the proteins that help fight infection. Some infants have repeated ear, sinus, or chest infections, but antibody levels usually improve as the immune system matures.

#### **Alternate Names:**

THI; Transient Infantile Hypogammaglobulinemia

### **Transient Ischaemic Attack**

A brief episode of stroke-like symptoms caused by temporarily reduced blood flow to the brain, eye, or spinal cord without lasting tissue damage. Weakness, numbness, speech trouble, or vision loss needs emergency assessment because stroke may follow.

#### **Alternate Names:**

TIA; Mini-Stroke

### **Transient Ischaemic Attacks Or Episodes**

Short episodes of weakness, numbness, speech trouble, vision loss, or other neurological symptoms caused by temporarily reduced blood flow to the brain. Symptoms resolve without lasting brain injury, but a TIA is an emergency warning of possible stroke.

### **Transient Ischemic Attack**

A temporary episode of stroke-like symptoms caused by reduced blood flow to the brain, eye, or spinal cord without lasting tissue damage. It can be an early warning of a possible stroke even when symptoms resolve.

### **Transient Ischemic Attack Cross-Reference**

Alternate name; see *Transient ischemic attack*.

### **Transient Neonatal Pustular Melanosis**

A harmless newborn eruption in which superficial pustules rupture and leave small pigmented macules with a collarette of scale. It is more common in infants with darker skin and usually resolves without treatment.

### **Transient Pain**

Pain that is brief and resolves rather than continuing or recurring persistently. Even brief pain may need assessment if it is severe, recurrent, or associated with warning symptoms.

### **Transient Synovitis**

Temporary inflammation of the hip lining, usually in children ages 3 to 8 and often after a viral illness. It causes hip, groin, thigh, or knee pain and a limp, but usually little fever or illness. Rest and pain medicine generally help. Doctors must distinguish it from a dangerous joint infection, especially when fever, severe pain, or refusal to walk is present.

#### **Alternate Names:**

Toxic Synovitis; Irritable Hip; Transient Synovitis of the Hip

### **Transient Tachypnea Of The Newborn**

Temporary rapid breathing after birth, usually caused by delayed clearance of fetal lung fluid. It is more common after caesarean birth and typically improves within one to three days with supportive care.

### **Transillumination**

Transillumination shines light through a body part or structure to reveal fluid, cavities, or other differences in tissue density.

### **Transitional Care**

Coordinated healthcare when a person moves between settings or levels of care, such as from hospital to home or from children's to adult services. It in-



---

cludes sharing information, arranging medicines and follow-up, and preventing avoidable complications.

### **Transitional Epithelium**

A stretchable protective lining in the kidneys' drainage areas, ureters, bladder, and part of the urethra. It allows these organs to expand as urine collects without allowing urine to leak into surrounding tissue.

#### **Alternate Names:**

Urothelium; Transitional Urothelial Lining

### **Transjugular Intrahepatic Portosystemic Shunt**

A channel created through the liver between veins that bring blood to it and veins that carry blood away. It lowers high pressure caused by advanced liver disease and can reduce repeated bleeding or severe abdominal fluid buildup; confusion and shunt blockage are possible.

#### **Alternate Names:**

TIPS; Transjugular Portosystemic Shunt

### **Translation**

The cellular process of making a protein from instructions carried by messenger RNA. A ribosome reads the message in order, while transfer molecules bring the matching amino acids. The process starts, builds the chain, and stops at a signal; the finished chain folds into a working protein.

### **Translation Initiation**

The first stage of protein synthesis, when a ribosome recognizes messenger RNA, selects the start codon, and assembles the initiator transfer RNA and ribosomal subunits.

### **Translocation**

Movement from one place to another. In genetics, it means that a piece of one chromosome moves to another chromosome or location, which can alter genes and sometimes cause cancer or inherited disease.

### **Transmembrane Transport**

Movement of salts, nutrients, medicines, or other

substances across a cell's outer membrane. Channels and carrier proteins may allow passage directly, while pumps use energy to move substances against a concentration difference.

### **Transmission**

Passage of an infection, signal, genetic trait, force, or substance from one source to another; in infectious disease, route, dose, and host susceptibility determine risk.

### **Transmural Infarction**

A heart attack that damages the full thickness of part of the heart muscle because blood flow has been blocked for a prolonged period. It can weaken the heart and cause dangerous complications.

### **Transplacental**

Passing from the pregnant person's blood across the placenta to the fetus, or in the opposite direction. Antibodies, medicines, chemicals, and some infections can cross the placenta, with effects that depend on the substance and timing.

### **Transplant**

Transfer of cells, tissue, or an organ from one site or person to another to replace damaged tissue or restore function. Donor matching, rejection prevention, infection risk, and long-term follow-up depend on the type of transplant.

### **Transplantation**

Transfer of an organ, tissue, or cells from a donor to a recipient to replace or restore failing function. Care includes donor matching, surgery, prevention of rejection, infection monitoring, and long-term follow-up.

#### **Alternate Names:**

Organ Transplant; Allotransplantation

### **Transplantation Conditioning**

Treatment given before some stem-cell transplants to suppress the recipient's bone marrow and immune system and make room for donated cells. It may use chemotherapy, radiation, or both and can cause infection, bleeding, infertility, or organ injury.



---

## Transplant Immunology

The study of how the immune system reacts to donated organs, tissues, or cells. Blood-group and tissue matching, antibody testing, and immunosuppressive medicines help reduce rejection, but they also increase infection risk and require regular monitoring of the transplant and medicine effects.

## Transplant Rejection

An immune reaction in which the body attacks a donated organ or tissue. It may occur soon after transplantation or later and can reduce organ function; regular monitoring and medicines help prevent or treat it.

## Transplant Surgery

An operation that replaces a failing organ or tissue with a donated one. It requires careful donor and recipient assessment, medicines to reduce rejection, infection prevention, and long-term follow-up.

## Transport

Movement of a substance, cell, signal, or person from one place to another. In healthcare, the term may describe movement through the body, across a cell membrane, or safe transfer between care settings.

## Transport Protein

A protein that carries a substance through body fluids or across a cell boundary. It helps control how nutrients, medicines, salts, and waste products are absorbed, distributed, and used.

## Transport Vesicle

A transport vesicle is a membrane-bound carrier that moves proteins, lipids, or other cargo between cellular compartments.

## Transposition Of Great Vessels

Alternate name; see *Transposition of the great arteries*.

## Transposition Of The Great Arteries

A heart defect present at birth in which the two main arteries are connected to the wrong lower chambers.

Blood returning to the body may not receive oxygen; newborns usually need urgent treatment and surgery.

### Alternate Names:

Complete TGA; D-TGA; Transposition of Great Vessels

## Transposition of the Great Vessels

A cyanotic congenital heart defect resulting from embryological failure of the aorticopulmonary septum to spiral, such that the aorta arises directly from the right ventricle and the pulmonary artery originates from the left ventricle, creating two parallel, independent circulations requiring a mixing communication for survival.

### Alternate Names:

TGA; D-Transposition of the Great Arteries; d-TGA

## Transposons

Movable pieces of genetic material that can shift between locations in a cell's DNA. In bacteria, they may carry antibiotic-resistance genes and help resistance spread between bacteria.

## Transrectal Ultrasonography

Ultrasound imaging of the prostate and nearby structures using a probe placed in the rectum. It can guide biopsies and assess size or abnormalities, but may cause discomfort, bleeding, or infection.

### Alternate Names:

TRUS

## Transsphenoidal Hypophysectomy

A minimally invasive skull-base neurosurgical procedure performed to remove pituitary gland tumors (pituitary adenomas) or craniopharyngiomas located at the base of the brain. Instead of cutting open the skull or leaving visible facial scars, the surgical team passes high-definition endoscopes and micro-instruments directly through the patient's nostrils, across the nasal cavity, and through the sphenoid sinus bone into the sella turcica—the bony saddle cradling the pituitary gland. By operating from below, neurosurgeons and otolaryngologists can decompress the optic chiasm to restore threatened vi-

---

sion, normalize excessive hormone production (such as in Cushing disease or acromegaly), and remove tumors with faster recovery times, minimal brain retraction, and reduced complication rates compared to traditional open craniotomies.

**Alternate Names:**

Transsphenoidal Surgery; Endoscopic Transsphenoidal Pituitary Resection; Pituitary Adenoma Surgery; TSS

**Transthoracic Echocardiogram**

A non-invasive point-of-care or formal diagnostic ultrasound procedure utilizing a transducer placed on the anterior chest wall to evaluate cardiac chamber dimensions, ventricular systolic and diastolic function, valvular morphology, and hemodynamics via Doppler imaging.

**Alternate Names:**

TTE

**Transthoracic Echocardiography**

An ultrasound scan of the heart made with a probe placed on the chest. It shows the heart chambers, valves, pumping movement, and blood flow.

**Transthyretin Amyloidosis**

A disease in which abnormal transthyretin protein builds up in organs, especially the heart or nerves. It may be inherited or develop with age without an inherited gene change.

**Alternate Names:**

ATTR Amyloidosis; ATTR-CM; Amyloid Transthyretin Disease

**Transtracheal**

Transtracheal means passing through or relating to the trachea, such as a catheter, oxygen delivery route, or diagnostic procedure.

**Transtympanic**

Transtympanic means passing through the tympanic membrane, such as medicine delivered into the middle ear or a procedure crossing the eardrum.

**Transudate**

A thin fluid with little protein that leaks from blood vessels because of pressure or blood-protein changes rather than inflammation. It may collect around organs, such as the lungs, heart, or abdomen.

**Transudation**

Movement of relatively watery, low-protein fluid out of blood vessels when pressure or blood-protein levels change. This process can cause some types of swelling or fluid collection around organs.

**Transudative Pleural Effusion**

A buildup of relatively thin fluid around a lung caused by changes in blood pressure or protein balance, most often heart failure. Treatment focuses on the underlying condition and drainage may be needed if breathing is affected.

**Transurethral**

Performed through the urethra, the tube that carries urine out of the body. Instruments passed through it can examine the urinary tract, remove tissue, treat blockage, or place a device.

**Transurethral Incision Of The Prostate**

An operation making small cuts in the prostate through an instrument passed in the urethra. It can improve urine flow in selected people with prostate enlargement while preserving most prostate tissue.

**Alternate Names:**

TUIP

**Transurethral Microwave Thermotherapy**

A treatment for some people with an enlarged prostate in which a catheter uses microwave heat to shrink or destroy prostate tissue blocking urine flow. It is less commonly used now and can cause pain, bleeding, infection, or temporary urinary problems.

**Alternate Names:**

TUMT

**Transurethral Needle Ablation**

A treatment using needles passed through the urethra to deliver radiofrequency heat into enlarged

---

prostate tissue. The heat destroys selected tissue and may improve urine flow, but relief develops gradually and complications can occur.

**Alternate Names:**

TUNA

### **Transurethral Resection**

Removal of tissue through an instrument passed along the urethra, commonly for prostate or bladder disease.

### **Transurethral Resection of Bladder Tumor**

A procedure in which a urologist passes a camera and small cutting tool through the urethra to remove a visible bladder tumor and collect tissue for testing. The results show the type and depth of the cancer and guide further treatment. Some early tumors are treated during the same procedure, but follow-up treatment may still be needed.

**Alternate Names:**

TURBT; Transurethral Bladder Tumor Resection; Bladder Tumor Resection

### **Transurethral Resection Of The Prostate**

An endoscopic operation that removes obstructing prostate tissue to improve urine flow in selected people with benign prostatic enlargement. Alternatives and indications depend on symptoms, prostate size, complications, and patient preference.

**Alternate Names:**

TURP; Transurethral Prostate Resection

### **Transvaginal**

Performed through the vagina, commonly describing an ultrasound examination or a procedure involving the uterus, cervix, ovaries, or other pelvic structures.

### **Transvaginal Ultrasound**

Transvaginal ultrasound uses a covered probe in the vagina to produce images of the uterus, ovaries, cervix, pregnancy, or nearby pelvic structures.

### **Transverse**

Running across or perpendicular to the long axis or

midline.

### **Transverse Colon**

The section of the large intestine that crosses the upper abdomen from the right side to the left. It absorbs water and moves partly digested food toward the descending colon.

### **Transverse Lie**

A position in which the baby lies sideways across the uterus rather than head-down or bottom-down. A shoulder or arm may be closest to the birth canal; if the position persists near labor, vaginal birth is usually not possible and cesarean birth is recommended.

### **Transverse Myelitis**

Inflammation across part of the spinal cord that can develop over hours or days and cause weakness, numbness, pain, or problems controlling the bladder or bowel. It needs prompt evaluation because treatment may limit lasting disability.

### **Transverse Plane**

An imaginary horizontal slice that divides the body or an organ into upper and lower parts. Medical scans often use this view to show structures at a particular level.

### **Transverse Vaginal Septum**

A congenital band of tissue crossing the vagina and partly or completely blocking it. A complete blockage can cause absent periods and monthly pelvic pain after puberty; imaging confirms the condition, and surgery may create an open vaginal passage.

### **Transversus Abdominis Muscle**

The deepest broad muscle of the abdominal wall. It tightens the abdomen, supports the spine, and helps with coughing, breathing out, lifting, and other movements that increase pressure inside the abdomen.

### **Transvestic Disorder**

A mental-health diagnosis involving repeated sexual

---

arousal from cross-dressing that causes significant distress, interferes with life, causes harm, or involves another person without consent. Cross-dressing alone is not a disorder and is different from gender identity.

### **Tranylecypromine**

An antidepressant that increases certain brain chemicals by blocking their breakdown. It can interact dangerously with many medicines and foods high in tyramine, so careful prescribing and a medicine-free waiting period before or after some treatments are required.

### **Trapezium Bone**

A small wrist bone at the base of the thumb. It forms the thumb's highly mobile joint with the hand and can be affected by fractures or arthritis that causes pain during gripping and pinching.

### **Trapezius Muscle**

A large triangular muscle across the upper back and neck. It helps lift, lower, and draw the shoulder blades backward and supports movement and stability of the neck and shoulders.

### **Trapezoid**

A small wedge-shaped wrist bone between the trapezium and capitate. It joins mainly with the index-finger metacarpal and helps stabilize the thumb-side of the wrist; fractures are uncommon.

### **Trapezoid Bone**

A small distal-row carpal bone between the trapezium and capitate that articulates mainly with the index metacarpal and helps stabilize the radial side of the hand.

### **Trapped Lung**

A condition in which a lung cannot fully expand because a fibrous layer or another restriction prevents normal movement. It may follow long-standing pleural inflammation, infection, bleeding, or cancer and can cause persistent pleural fluid or breathlessness.

### **TRAP Sequence**

A rare complication of an identical-twin pregnancy in which one twin has a severely malformed heart or no heart and receives blood from the other twin through shared placental vessels. The healthy twin may develop heart strain and needs specialist monitoring.

### **Trastuzumab**

Trastuzumab is a monoclonal antibody targeting HER2 used for HER2-positive breast, gastric, and other cancers; infusion reactions and potentially reversible cardiomyopathy require cardiac monitoring.

### **Trauma**

Physical injury caused by an outside force, or the mental response to a deeply distressing event. Physical trauma may damage organs, bones, or blood vessels; psychological trauma can affect emotions, sleep, and daily life.

### **Trauma- And Stressor-Related Disorders**

A group of mental disorders involving exposure to a traumatic or stressful event or condition and subsequent emotional, behavioral, or physiologic symptoms. Examples include acute stress disorder, post-traumatic stress disorder, adjustment disorder, reactive attachment disorder, and disinhibited social engagement disorder.

### **Trauma Center**

A hospital organized to provide rapid, coordinated care for seriously injured people. Trauma centers differ in available specialists, equipment, surgery, intensive care, and ability to treat complex injuries.

### **Trauma Surgery**

Emergency surgery and other immediate care for seriously injured people. The team first checks and protects breathing, the neck and spine, circulation, brain function, and the whole body, while controlling bleeding and repairing damaged organs.

### **Traumatic Amputation**

Loss of a body part caused by an injury. It can cause life-threatening bleeding, shock, infection, and

---

lasting disability; emergency care controls bleeding, treats damaged tissue, preserves the separated part when possible, and begins rehabilitation and psychological support.

### **Traumatic Bleb**

A fragile, fluid-filled area that forms after glaucoma drainage surgery and is damaged by an injury. It may leak or become infected and can cause dangerously low eye pressure or vision problems.

### **Traumatic Brain Injury**

Traumatic Brain Injury (TBI) is a change in brain function or damage to the brain caused by an external force. It may result from a bump, blow, or jolt to the head or body, a fall, blast, or an object penetrating the skull. Effects range from brief confusion or headache to memory problems, seizures, loss of consciousness, permanent disability, or death. Later swelling, bleeding, low oxygen, or other complications can cause additional brain damage.

**Alternate Names:**

TBI

### **Traumatic Events And Children**

Traumatic events can affect children's safety, behavior, sleep, learning, and emotions; supportive caregivers, stable routines, and professional assessment help identify persistent trauma responses.

### **Traumatic Grief**

Alternate name; see *Complicated grief*.

### **Traumatic Hemorrhagic Shock**

Shock caused by severe blood loss after an injury. Reduced circulating blood volume limits oxygen delivery to organs and may cause fast heartbeat, pale or clammy skin, confusion, low blood pressure, and organ failure.

### **Traumatic Injury**

Damage to the body caused by an external force, such as a fall, collision, blow, cut, burn, or penetrating object. Injury may affect skin, bones, organs, blood vessels, nerves, or the brain.

### **Traumatic Bladder and Urethral Injury**

Damage to the bladder or urethra caused by an accident, surgery, or penetrating injury. Blood in the urine, inability to urinate, pelvic pain, or shock requires urgent evaluation.

### **Traumatic Shock**

A life-threatening failure of blood flow after serious injury, often because of major bleeding, but sometimes because of heart damage or another cause. It can cause pale clammy skin, confusion, rapid breathing, and collapse and requires immediate treatment.

### **Traumatology**

The medical field concerned with physical injuries and their consequences. It includes emergency assessment, control of bleeding, repair of damaged tissues, prevention of complications, surgery, and rehabilitation.

### **Traumeel S**

Traumeel S is a multi-ingredient complementary product marketed for pain and inflammation; evidence, formulation, and safety depend on the product and it should not replace needed care.

### **Travelers' Diarrhea**

Acute diarrhea occurring during or shortly after travel, usually caused by contaminated food or water and commonly involving bacterial, viral, or parasitic infection.

### **Travel Sickness**

Nausea, dizziness, sweating, or vomiting caused by conflicting signals from the eyes and balance organs during travel or other motion. Looking toward the horizon, fresh air, gradual exposure, and suitable preventive medicines may help; severe or persistent symptoms need assessment.

### **Travel Vaccinations**

Travel vaccinations are vaccines recommended or required for protection against infections encountered in particular countries or regions. Recommendations depend on the destination, activities, health conditions, and previous vaccinations.

---

### **Travoprost**

Travoprost is a prostaglandin analogue eye drop that increases uveoscleral outflow and lowers intraocular pressure in glaucoma or ocular hypertension; iris pigmentation and eyelash changes may occur.

### **Trazodone**

A prescription medicine used mainly for depression and sometimes for sleep problems. It can cause drowsiness, dizziness, low blood pressure, falls, and rare prolonged painful erection; dangerous reactions are more likely with certain medicines or substances.

### **Trazodone Overdose**

Poisoning from taking too much trazodone. It can cause severe sleepiness, vomiting, low blood pressure, seizures, abnormal heart rhythms, or a prolonged painful erection.

### **Trb Gene**

A gene involved in biological processes that vary by species; its precise identity and function must be interpreted from the organism and gene nomenclature used.

### **Trd Gene**

The gene encoding the delta chain of the T-cell receptor in humans, contributing to gamma-delta T-cell antigen recognition.

### **Treacher Collins Syndrome**

An inherited condition affecting development of the face and skull. It may cause small cheekbones and jaw, downward-slanting eyes, malformed ears, and hearing loss; breathing or feeding may also be difficult. Intelligence is usually normal. Care may include airway support, hearing treatment, and reconstructive surgery.

### **Treacle**

A thick sugar syrup with little current medical use. Older medical writing sometimes used the word for a sweetened mixture or supposed antidote; it is not a modern treatment or diagnosis.

### **Treatment-Resistant Depression**

Depression that does not improve adequately after appropriate trials of standard treatments. It requires reassessment of diagnosis, adherence, comorbidity, and treatment options.

### **Treatment-Resistant Schizophrenia**

Schizophrenia that continues to cause important symptoms or disability despite adequate trials of at least two appropriate antipsychotic medicines. The term requires careful review of the diagnosis, treatment adherence, substance use, medical causes, and the adequacy of previous treatment.

### **Treat-To-Target Strategy**

A treatment approach in which care is checked and adjusted regularly toward a clear goal, such as no disease activity, controlled blood pressure, or a safe blood-glucose range. The goal and changes are agreed with the patient and based on ongoing monitoring.

### **Trematoda**

A class of parasitic flatworms called flukes; human infections include schistosomiasis and liver or lung fluke disease.

### **Tremelimumab**

A monoclonal antibody that blocks CTLA-4 and enhances T-cell activity; it is used with durvalumab for selected unresectable hepatocellular carcinoma and other cancers, with immune-related adverse effects possible.

### **Tremor**

An involuntary, rhythmic shaking of a body part. It may occur at rest, while holding a position, or during movement and can result from medicines, caffeine, anxiety, essential tremor, Parkinson disease, or other neurologic or metabolic conditions.

### **Trench Fever**

A louse-borne infection caused by Bartonella quintana, causing recurring fever, headache, weakness, and bone pain.



---

### **Trench Foot**

An injury caused by keeping feet cold and wet for a long time. It can cause numbness, swelling, pain, skin color changes, blisters, infection, and tissue death without gradual warming and medical care.

### **Trench Mouth**

Trench mouth is acute necrotizing ulcerative gingivitis causing painful bleeding gums, foul breath, and ulceration, often associated with plaque, smoking, stress, or poor nutrition.

### **Trendelenberg Position**

A position in which the patient lies supine with the feet higher than the head; clinical uses and risks depend on context.

### **Trendelenburg**

Trendelenburg position places the body supine with the feet higher than the head; its use is procedure-specific and may affect breathing, pressure, and circulation.

### **Trendelenburg Gait**

An abnormal gait pattern characterized by a downward pelvic drop toward the unsupported side during the swing phase, or a lateral trunk lurch over the weight-bearing side to compensate for hip abductor weakness. It occurs due to weakness or paralysis of the gluteus medius and minimus muscles or superior gluteal nerve lesion.

### **Trendelenburg Position**

A position lying on the back with the head and upper body lower than the feet. It is used for selected procedures but can worsen breathing, pressure in the head, or blood pressure in some patients.

### **Trendelenburg Sign**

A physical clinical sign indicating weakness or paralysis of the hip abductor muscles (gluteus medius and minimus) innervated by the superior gluteal nerve; when the patient stands on one leg (the affected stance side), the unsupported pelvis tilts downward toward the opposite non-weight-bearing swing side.

### **Alternate Names:**

Trendelenburg Hip Abductor Sign; Positive Trendelenburg Test

### **Trendelenburg Test**

A clinical test used to evaluate the integrity and strength of the hip abductor mechanism (gluteus medius and minimus). The patient stands on one leg; a drop of the pelvis on the contralateral unsupported side indicates abductor weakness or hip instability on the stance side.

### **Treosulfan**

An alkylating chemotherapy drug used in conditioning before allogeneic hematopoietic stem-cell transplantation; infection, mucositis, infertility, and organ toxicities may occur.

### **Trephine**

A circular cutting instrument used to remove a disk of tissue or bone, such as during a bone-marrow biopsy or corneal procedure.

### **Treponema**

A group of spiral-shaped bacteria, including *Treponema pallidum*, which causes syphilis. Different species can live in people or the environment, and identifying the species helps determine whether disease and treatment are present.

### **Treponemal Infections**

Infections caused by spiral-shaped *Treponema* bacteria. They include syphilis and related diseases and may affect the skin, nerves, heart, eyes, or other organs; diagnosis usually combines clinical findings with blood or lesion tests.

### **Treponema Pallidum**

The spiral-shaped bacterium that causes syphilis. It spreads mainly through contact with an infectious sore during sex and can cross the placenta to an unborn baby. Untreated infection can later affect the skin, nerves, eyes, heart, or brain.

---

## Treponema Pallidum Infection

Infection with the bacterium *Treponema pallidum*, which causes syphilis and can affect the skin, nervous system, cardiovascular system, and other organs.

## Trepopnea

Breathlessness that occurs or becomes worse when lying on one side but not the other. It can result from disease affecting one lung, fluid around a lung, or pressure from a nearby mass and should be assessed with other symptoms.

## Tretinoin

A vitamin-A-related medicine applied to the skin for acne and sun-related skin changes, or taken by mouth for a type of acute leukemia. Skin irritation is common; oral treatment can cause serious fluid buildup, fever, and breathing problems.

## TRG Gene

A gene encoding the gamma chain of the T-cell receptor, contributing to gamma-delta T-cell development and antigen recognition.

## Triage

Sorting patients by urgency and resource needs so those at greatest risk receive timely care.

## Trial Of Labor After Cesarean

An attempt to give birth vaginally after a previous cesarean delivery. Whether it is appropriate depends on the prior uterine incision, obstetric history, current pregnancy, facility resources, and the person's preferences. Counseling includes the possibility of successful VBAC, repeat cesarean delivery, and uterine rupture.

## Triamcinolone

A corticosteroid medicine used on skin, by injection, or in other forms to reduce inflammation and itching.

## Triamcinolone Acetonide

A synthetic corticosteroid used topically, by injection, or in other preparations to reduce inflammation;

prolonged or extensive use can cause local and systemic steroid effects.

## Triamterene

A water tablet that helps the kidneys remove salt and water while conserving potassium. It is used for high blood pressure or swelling, but can cause dangerously high potassium, kidney problems, or kidney stones.

## Triangle of Doom

An essential laparoscopic anatomical region in the pelvis bounded by the vas deferens medially, the spermatic (testicular) vessels laterally, and the peritoneal reflection inferiorly, containing the external iliac artery and vein and deep circumflex iliac vein, which must be avoided during laparoscopic inguinal hernia mesh fixation to prevent catastrophic vascular injury.

### Alternate Names:

Laparoscopic Vascular Danger Triangle; Inguinal Triangle of Doom

## Triangle of Pain

A laparoscopic anatomical region in the groin bounded by the spermatic vessels medially, the iliopectic tract superolaterally, and the peritoneal reflection inferiorly, containing the femoral branch of the genitofemoral nerve, the lateral femoral cutaneous nerve, and the femoral nerve, where placement of tacks or staples must be avoided during hernia repair to prevent chronic neuropathic groin pain.

### Alternate Names:

Laparoscopic Neural Danger Triangle; Inguinal Triangle of Pain

## Triazenes

A group of chemical compounds that includes some chemotherapy medicines. Their effects and risks depend on the specific drug; possible problems include nausea, low blood counts, infection, and damage to other organs.

## Triaziquone

An older alkylating anticancer drug that cross-links DNA; its clinical use is limited by toxicity and avail-

---

ability.

### **Triazolam**

Triazolam is a short-acting benzodiazepine hypnotic for short-term insomnia; amnesia, falls, dependence, respiratory depression, and dangerous interactions with opioids or other sedatives can occur.

### **Triazole**

A five-membered nitrogen-containing chemical ring found in several antifungal medicines and other drugs; clinical effects depend on the specific compound.

### **Triazophos**

An organophosphate insecticide that blocks acetylcholinesterase, allowing nerve signals to remain active. Significant exposure can cause sweating, vomiting, diarrhea, small pupils, breathing difficulty, weakness, seizures, or life-threatening poisoning.

### **Triceps**

A muscle with three heads, especially the triceps brachii at the back of the upper arm. It straightens the elbow and helps stabilize the shoulder; injury or nerve problems can weaken pushing and lifting.

### **Triceps Brachii Muscle**

The large muscle at the back of the upper arm. It straightens the elbow and helps move and stabilize the shoulder; injury or nerve damage can weaken pushing and lifting.

### **Triceps Reflex**

A monosynaptic deep tendon stretch reflex elicited by tapping the triceps brachii tendon immediately superior to the olecranon process with a reflex hammer, stretching muscle spindles and provoking reflex forearm extension mediated through the radial nerve (evaluating spinal cord segments C7 and C8).

#### **Alternate Names:**

Triceps Tendon Reflex; C7 Reflex

### **Trichiasis**

Inward misdirection of eyelashes so they rub against

the eye.

### **Trichilemmal Carcinoma**

A rare malignant follicular tumor showing outer-root-sheath differentiation, usually occurring on sun-damaged skin of older adults as a papule or nodule; complete excision is generally required.

### **Trichilemmal Cyst (Pilar Cyst)**

A usually benign keratin-filled cyst arising from the outer root sheath of a hair follicle, most often on the scalp. It is typically a firm, mobile lump and may become inflamed, rupture, or rarely develop a proliferating pilar tumor.

### **Trichilemmoma**

A usually harmless tumor arising from a hair follicle, most often appearing as a small skin-colored bump on the face. Multiple growths can be associated with an inherited condition called Cowden syndrome.

### **Trichinellosis**

Trichinellosis is an infection caused by *Trichinella* worms, usually after eating undercooked infected meat. It may cause diarrhea followed by fever, muscle pain, and swelling around the eyes.

### **Trichiniasis**

An infection caused by *Trichinella* larvae, acquired from undercooked meat.

### **Trichinosis**

Infection with *Trichinella* worms, usually acquired by eating undercooked infected pork or wild game. It may cause diarrhea, fever, muscle pain, and swelling around the eyes.

### **Trichobezoar**

A mass of swallowed hair in the digestive tract, usually the stomach, which may cause pain, vomiting, or blockage.

### **Trichoblastoma**

A rare benign follicular tumor of the scalp or face, usually presenting as a solitary dermal or subcuta-

---

neous nodule; it can resemble basal-cell carcinoma and is diagnosed histologically.

### **Trichoepithelioma**

A rare benign follicular adnexal tumor that usually presents as a skin-coloured papule or nodule on the face or scalp; it can resemble basal-cell carcinoma and may require histopathologic confirmation.

### **Trichomonas**

A flagellated protozoan, *Trichomonas vaginalis*, that infects the urogenital tract and causes trichomoniasis. It is transmitted mainly through sexual contact and may cause discharge, irritation, or no symptoms.

### **Trichomonas Vaginalis**

A flagellated protozoan that causes trichomoniasis, a sexually transmitted infection.

### **Trichomoniasis**

A sexually transmitted infection caused by the parasite *Trichomonas vaginalis*. It may cause genital irritation and discharge or no symptoms; testing and treatment of partners help prevent reinfection.

### **Trichophyton**

A group of fungi that grow in keratin and cause ringworm infections of skin, hair, or nails. They spread through contact with infected people, animals, or objects and are treated with antifungal medicines.

### **Trichorrhexis Nodosa**

A hair-shaft defect in which weakened points split into brush-like ends, caused by inherited disease, trauma, chemicals, heat, or nutritional or metabolic disorders.

### **Trichostasis Spinulosa**

A follicular disorder in which many fine vellus hairs become retained in a dilated follicle, producing dark, rough, spiny papules often on the back, chest, or face. It is benign and may improve with topical keratolytics, retinoids, or hair-removal methods.

### **Trichothiodystrophy Syndrome**

A group of inherited disorders characterized by brittle sulfur-deficient hair with a distinctive banded appearance, often accompanied by photosensitivity, ichthyosis, developmental delay, short stature, or recurrent infection.

### **Trichotillomania**

Recurrent pulling out of one's hair, causing hair loss, distress, or impairment, together with repeated attempts to reduce or stop the behavior.

### **Trichuriasis**

Intestinal infection caused by the whipworm *Trichuris trichiura*, usually acquired by swallowing eggs from contaminated soil or food. Heavy infection can cause abdominal pain, diarrhea, anemia, or rectal prolapse.

### **Tricuspid Annular Plane Systolic Excursion**

A measurement made during an echocardiogram that shows how far part of the tricuspid valve moves as the right ventricle contracts. A low value may indicate reduced right-heart pumping and is interpreted with the complete scan and clinical findings.

#### **Alternate Names:**

TAPSE; Tricuspid Annular Excursion; RV Longitudinal Systolic Function

### **Tricuspid Atresia**

A congenital absence or complete closure of the tricuspid valve, blocking normal flow from the right atrium to the right ventricle. Survival depends on an atrial communication and usually a ventricular septal defect or patent ductus to allow blood to reach the lungs; newborns typically have cyanosis and need urgent specialist care and staged treatment.

### **Tricuspid Incompetence**

Incomplete closure of the tricuspid heart valve, allowing blood to leak backward into the right upper chamber. Severe leakage can cause swelling, liver congestion, irregular heartbeat, fatigue, and breathlessness.

---

## Tricuspid Regurgitation

Backward flow through the tricuspid valve from the right ventricle to the right atrium during systole. It may result from primary valve disease or annular enlargement caused by right-ventricular or right-atrial dilation, pulmonary hypertension, or left-sided heart disease; severe regurgitation can cause fatigue, edema, ascites, liver congestion, or right-sided heart failure.

## Tricuspid Stenosis

Narrowing of the tricuspid valve opening, making it harder for blood to move from the right upper chamber to the right lower chamber. It can cause fatigue, swelling, enlarged liver, or neck-vein fullness.

## Tricuspid Valve

The heart valve between the right upper and right lower chambers. It opens to let blood move forward and closes to prevent backflow; leakage or narrowing can strain the heart.

## Tricuspid Valve Disease

Alternate name; see *Tricuspid regurgitation* and *Tricuspid stenosis*.

## Tricyclic Antidepressant Drugs

A group of older antidepressants that increase certain chemical signals in the brain. They can also cause dry mouth, constipation, sleepiness, dangerous heart rhythms, and serious poisoning in overdose.

## Tricyclic Antidepressants

Antidepressants that block reuptake of serotonin and norepinephrine, with additional anticholinergic, antihistaminic, and alpha-adrenergic blocking effects. Examples include amitriptyline, nortriptyline, imipramine, and desipramine. They are effective for depression, neuropathic pain, and migraine prophylaxis.

### Alternate Names:

Tricyclics; First-Generation Antidepressants

## Tricyclic Antidepressant Toxicity

Rapid anticholinergic and sodium-channel toxicity causing dry skin, mydriasis, delirium, seizures, hypotension, QRS widening, ventricular dysrhythmias, and coma. Obtain an ECG early; airway support, intravenous sodium bicarbonate for QRS widening or ventricular toxicity, and benzodiazepines for seizures are key interventions.

## Trifluoperazine

A phenothiazine antipsychotic used for selected psychotic disorders. It can cause drowsiness, movement disorders, and other serious effects, so treatment requires medical supervision.

## Trigeminal Nerve

The fifth cranial nerve, carrying sensation from the face and controlling muscles used for chewing.

## Trigeminal Neuralgia

A chronic pain disorder causing sudden, repeated bursts of severe electric-shock-like, stabbing, or burning pain on one side of the face along one or more branches of the trigeminal nerve, the fifth cranial nerve. Attacks usually last from a fraction of a second to about two minutes and may occur in clusters, with pain-free periods between them. Light touch, brushing the teeth, eating, drinking, speaking, or a breeze may trigger an attack. The usual cause is pressure on the nerve near the brainstem, but multiple sclerosis, a tumor, or another nerve disorder can sometimes be responsible.

### Alternate Names:

Tic Douloureux; Fothergill Disease

## Trigger

A trigger is an event, exposure, stimulus, or condition that initiates or worsens a symptom, disease episode, behavior, or physiologic response.

## Trigger Finger

A condition in which a finger tendon catches as it moves through a tight covering. It can cause pain, clicking, stiffness, or locking and may improve with rest, injections, splinting, or surgery.

### Alternate Names:

---

Stenosing Tenosynovitis; Snapping Digit

### Trigger Finger Release

A procedure that opens the tight tunnel around a finger tendon so the finger can move more freely. It may be offered when rest, a splint, or an injection has not helped. Movement often improves, but recovery and results vary and infection, nerve injury, pain, or continued symptoms can occur.

#### Alternate Names:

A1 Pulley Release; Trigger Thumb Release; Stenosing Tenosynovitis Surgery; Percutaneous Trigger Finger Release

### Trigger Thumb

A condition in which the thumb tendon catches as it moves through a tight covering. It may cause pain, clicking, stiffness, or locking and can often be treated with rest, splinting, an injection, or surgery.

### Triglyceride

A lipid composed of glycerol and three fatty acids, transported in blood and stored in adipose tissue.

### Triglyceride Level

A triglyceride level measures a type of fat in the blood; persistently high levels can increase cardiovascular risk and, when very high, cause pancreatitis.

### Triglycerides

Triglycerides are blood fats made of glycerol and three fatty acids; elevated levels contribute to cardiovascular risk and, when very high, pancreatitis risk.

### Trigone

A triangular area, especially the smooth region at the base of the urinary bladder between the ureter openings and urine outlet. It helps direct urine flow and can be affected by inflammation or tumors.

### Triiodothyronine

The thyroid hormone T3, which contains three iodine atoms and has stronger biological activity than T4. It helps regulate metabolism, heart rate, tem-

perature, and growth.

### Trimester

One of the three approximately three-month periods of pregnancy. Each trimester has typical developmental milestones and screening or care needs.

### Trimethoprim

An antibiotic that blocks bacterial folate production and is often combined with sulfamethoxazole. It treats selected infections but can cause rash, blood-cell changes, high potassium, kidney effects, or medicine interactions.

### Trimethoprim-Sulfamethoxazole

An antibiotic combination that blocks bacteria from making folate, which they need to grow. It treats selected urinary, lung, skin, and other infections but can cause allergy, blood-count changes, high potassium, or kidney problems.

### Trimethylaminuria

A metabolic disorder in which impaired oxidation of trimethylamine causes a characteristic fish-like body odor in sweat, urine, and breath. It may be inherited or acquired and can cause substantial psychosocial distress despite limited physical illness.

### Trimipramine

A sedating tricyclic antidepressant used for depression in some countries. Its anticholinergic and cardiovascular effects can cause dry mouth, constipation, blurred vision, drowsiness, orthostatic hypotension, or dangerous toxicity in overdose; interactions and suicide risk require clinical monitoring.

### Trinitrin

Another name for glyceryl trinitrate or nitroglycerin, a nitrate vasodilator.

### Trinitrophenol

Picric acid, a corrosive and potentially explosive chemical once used in medicine, laboratories, and industry.



---

### Trinucleotide Repeat

A short DNA sequence of three building blocks repeated many times. Extra copies can interfere with genes and cause inherited disorders such as Huntington disease, fragile X syndrome, myotonic dystrophy, or Friedreich ataxia.

### Trip

A trip is a journey or an accidental stumble; in drug-related language it can also mean an episode of altered perception after a hallucinogen.

### Tripe Palms

Diffuse, velvety, thickened hyperkeratosis of the palmar skin with a cerebriform, corrugated appearance resembling the lining of a cow's stomach (tripe), which is strongly associated with underlying malignancy, most commonly gastric adenocarcinoma. It may occur alone or in conjunction with acanthosis nigricans.

### Triple-Negative Breast Cancer

Breast cancer whose cells lack three common growth-related markers: estrogen receptors, progesterone receptors, and HER2. It does not respond to treatments aimed at those markers, so treatment may include surgery, chemotherapy, immunotherapy, or other options based on the cancer's features.

### Triplets

Three offspring born from the same pregnancy or three infants born in a multiple birth.

### Triple X Syndrome

A sex-chromosome variation in which a female has an extra X chromosome. Many affected people have typical sexual development and fertility, while some have tall stature, speech or learning difficulties, coordination problems, or anxiety; severity varies widely.

#### Alternate Names:

Trisomy X; 47,XXX Syndrome

### Triploidy Syndrome

A chromosome condition in which cells have three

complete sets of chromosomes instead of two. It usually causes severe growth restriction and multiple birth defects and often prevents long-term survival; rare mosaic forms may be less severe.

### Triptan

A medicine for stopping a migraine attack. It changes signals in pain pathways and narrows certain blood vessels. It is not suitable for everyone, especially people with some heart or blood-vessel conditions.

### Triptans

A group of medicines used to stop a migraine attack. They act on serotonin receptors and pain pathways; they are not suitable for some people with heart or blood-vessel disease and can cause chest or pressure sensations.

### Triquetrum

One of the small bones on the little-finger side of the wrist. It helps stabilize the wrist and can be injured by a fall or direct blow, causing pain and swelling.

### Trismus

Difficulty opening the mouth because the jaw muscles are tight or painful. Causes include dental infection, injury, medicines, muscle disorders, and tetanus; severe or rapidly worsening restriction can interfere with eating or breathing and needs urgent assessment.

### Trisodium Phosphate Poisoning

Poisoning from swallowing or contacting trisodium phosphate, a strong alkaline cleaner. It can burn the mouth, throat, eyes, and skin and may cause vomiting or breathing problems.

### Trisomy

A chromosome condition in which cells have an extra copy of a whole chromosome or chromosome segment, giving three copies instead of the usual two. It may affect all cells or only some cells (mosaicism); effects depend on the chromosome or segment involved and may include developmental differences,

---

birth defects, or pregnancy loss.

### Trisomy 13

Patau syndrome, a chromosomal disorder caused by an extra chromosome 13 and associated with severe developmental, brain, facial, heart, kidney, and limb abnormalities.

#### Alternate Names:

D Trisomy Syndrome; Patau Syndrome

### Trisomy 18

Edwards syndrome, a chromosomal disorder caused by an extra chromosome 18 and associated with growth restriction, characteristic craniofacial and limb findings, heart disease, and severe developmental impairment.

### Trisomy 21

Down syndrome, a genetic condition caused by an extra copy of chromosome 21. It may affect learning, growth, muscle tone, the heart, hearing, vision, or other organs; the features and level of support needed vary widely.

### Trisomy 21 Syndrome

Alternate name; see *Down syndrome*.

### Trisomy X

Alternate name; see *Triple X syndrome*.

### Trocar

A pointed instrument placed inside a hollow tube to enter a body cavity or tissue. It is used in minimally invasive procedures to create an opening for cameras, instruments, or drainage tubes.

### Trochanter

A bony projection on the thigh bone, especially the larger outer or smaller inner projection. Muscles and tendons attach there, and irritation around the greater trochanter can cause outer-hip pain.

### Trochanteric Bursitis

Alternate name; see *Greater trochanteric pain syndrome*.

### Troche

A troche is a small medicated lozenge that dissolves slowly in the mouth or throat for local or systemic drug delivery.

### Trochlear Nerve

Cranial nerve IV, which supplies the superior oblique eye muscle. Injury can cause vertical or tilted double vision, often worse when looking down or inward.

### Trochlear Nerve Palsy

A cranial neuropathy involving the fourth cranial nerve (CN IV), causing paralysis of the superior oblique muscle, presenting with vertical and torsional diplopia (characteristically worse on downgaze and reading), an upward and inward hypertropia of the affected eye, and a compensatory head tilt toward the opposite contralateral shoulder (Bielschowsky head tilt test).

#### Alternate Names:

Fourth Nerve Palsy; CN IV Palsy

### Trophic

Relating to the nutrition, growth, development, or maintenance of tissue. Trophic changes may occur when nerve supply or blood flow is impaired.

### Trophoblast

The outer cell layer of an early embryo that attaches to the uterus and helps form the placenta. It supports implantation, exchanges nutrients, and produces hormones needed to maintain early pregnancy.

### Trophoderm

The outer layer of cells in a very early embryo before it attaches to the uterus. It helps the embryo implant and develops into tissues that form much of the placenta.

### Trophozoite

The active stage of a single-celled parasite. In this stage it feeds, grows, and often multiplies; it differs from the more resistant form that some parasites

---

use to survive outside the body.

### **Tropical Disease**

A tropical disease is an infection or other health condition more common in tropical and subtropical regions, often because of local climate, vectors, or living conditions. Travel history helps guide diagnosis.

### **Tropical Diseases**

Illnesses more common in warm tropical or subtropical regions, including malaria, dengue, schistosomiasis, and other neglected infections.

### **Tropical Medicine**

The medical specialty concerned with diseases common in tropical and subtropical regions and illnesses related to travel. It includes infections, parasites, insect-borne diseases, environmental hazards, prevention, diagnosis, and treatment.

### **Tropical Sprue**

A disorder causing long-lasting diarrhea and poor absorption of nutrients, reported mainly in tropical and subtropical areas. It can lead to weight loss and deficiencies of folate, iron, and vitamin B12. Testing helps exclude infection and other causes; treatment usually includes antibiotics and replacement of missing nutrients.

### **Tropical Ulcer**

A chronic ulcer occurring in tropical regions, often associated with infection, trauma, or poor wound healing.

### **Troponin**

A protein released into the blood when heart muscle is injured. A raised level can support a heart-attack diagnosis, but it may also result from other heart, lung, kidney, or severe systemic illnesses.

### **Troponin I**

An inhibitory subunit of the troponin regulatory complex of cardiac striated muscle; cardiac troponin I (cTnI) possesses an amino acid sequence unique

to myocardium, serving as an exquisitely sensitive and specific clinical blood biomarker for acute myocardial injury and infarction.

#### **Alternate Names:**

cTnI; Cardiac Troponin I

### **Troponin Test**

A troponin test measures heart-muscle proteins in the blood and helps assess possible heart injury, especially during suspected heart attack.

### **Trouble Swallowing**

Difficulty moving food or liquid from the mouth to the stomach. It may cause coughing, choking, pain, or a feeling that food is stuck in the throat or chest.

### **Trousseau Sign**

A hand cramp where the fingers stiffen and the thumb pulls inward when a blood pressure cuff is pumped up on the arm, showing low calcium in the blood.

### **Trousseau Sign Of Latent Tetany**

A hand spasm that may occur when blood flow to the arm is briefly reduced with a blood-pressure cuff. It can suggest low blood calcium, but it is not specific and must be interpreted with symptoms and blood tests.

#### **Alternate Names:**

Trousseau Sign; Carpopedal Spasm Sign

### **Trousseau Sign of Malignancy**

Spontaneous, migratory superficial thrombophlebitis characterized by recurrent tender, erythematous venous cords appearing sequentially in different superficial veins of the extremities and trunk, strongly associated with underlying visceral mucinous adenocarcinomas, particularly of the pancreas.

#### **Alternate Names:**

Trousseau Syndrome; Migratory Thrombophlebitis

### **Trousseau Syndrome**

Repeated or shifting blood clots in veins associated with active cancer, especially some cancers of the

---

pancreas, stomach, or lung. The cancer can make the blood clot too easily; it differs from the hand spasm called Trousseau sign.

## Trp

Trp is the standard abbreviation for tryptophan, an essential amino acid used in protein synthesis and serotonin and niacin pathways.

## Ovotesticular Disorder of Sex Development

A difference of sex development in which both ovarian and testicular tissue are present, either in one ovotestis or in separate gonads. Chromosomal patterns and external anatomy vary. The deprecated historical term “true hermaphroditism” may appear in older records, but current medical writing should use ovotesticular disorder of sex development or ovotesticular DSD.

### Alternate Names:

Ovotesticular DSD; True Hermaphroditism (deprecated historical term)

## True Ribs

The first seven pairs of ribs, each attached directly to the breastbone by its own flexible cartilage. They protect the heart and lungs and move with the chest during breathing.

## Truncation

A shortened or incomplete form caused by removal of part of a sequence, structure, chromosome, protein, or process; the effect depends on what was lost and where.

## Truncus Arteriosus

A birth defect in which one large blood vessel leaves the heart instead of separate vessels for the body and lungs. A large hole between the lower chambers usually allows oxygen-poor and oxygen-rich blood to mix, causing blue skin and heart failure. Early surgery is usually required.

## Trunk

The central part of the body excluding the head and limbs, or a major nerve or vessel trunk.

## Trypanosoma

A genus of protozoan parasites causing African and American trypanosomiasis.

## Trypanosoma Brucei

A protozoan parasite transmitted by tsetse flies that causes African trypanosomiasis, or sleeping sickness. The human-infecting subspecies differ in geographic distribution and typical disease course; infection can progress from fever and lymph-node enlargement to neurologic involvement.

## Trypanosomiasis

Disease caused by Trypanosoma parasites, including African sleeping sickness and Chagas disease. Infected insects spread them, and illness may affect blood, lymph nodes, the nervous system, heart, or digestive organs.

## Trypsin

A digestive enzyme made by the pancreas and activated in the small intestine. It breaks proteins in food into smaller parts and activates other pancreatic digestive enzymes.

## Trypsin And Chymotrypsin In Stool

Stool trypsin and chymotrypsin testing measures digestive enzymes and may help assess pancreatic enzyme production, although other tests are often preferred.

## Trypsinogen Test

A trypsinogen test measures a pancreatic enzyme precursor and may help evaluate pancreatic function or newborn screening for cystic fibrosis.

## Tryptophan

An amino acid the body cannot make and must obtain from food. It is used to make proteins and can be changed into serotonin, melatonin, and niacin, which help regulate mood, sleep, and other body functions.

## Tsetse Fly

A blood-feeding fly found in parts of sub-Saharan

---

Africa. Its bite can spread *Trypanosoma* parasites, which cause African sleeping sickness; infection may cause fever, headaches, behavior changes, sleep disturbance, and serious nervous-system disease.

## **TSH**

Thyroid-stimulating hormone, made by the pituitary gland to tell the thyroid to produce its hormones. A blood TSH measurement is commonly the first test for many thyroid problems.

## **TSH-Receptor Antibody**

An antibody that attaches to the thyroid's control receptor. Some antibodies overstimulate the thyroid and cause Graves disease, while others block its activity; testing can help diagnose thyroid disease and assess pregnancy-related risk.

## **TSH Test**

A TSH test measures thyroid-stimulating hormone in the blood and is a common initial test of thyroid function.

## **TSI Test**

A blood test that detects antibodies which overstimulate the thyroid. It can support the diagnosis or monitoring of Graves disease and may help assess risk to a fetus during pregnancy.

## **Tsutsugamushi Fever**

Scrub typhus, a mite-borne infection caused by *Orientia tsutsugamushi*. It may cause fever, headache, rash, an eschar, and organ complications and is treated with an appropriate antibiotic.

## **Tubal Ligation**

A permanent birth-control procedure that closes, cuts, or removes the fallopian tubes so sperm and egg cannot normally meet. It does not protect against sexually transmitted infections, and pregnancy after the procedure, though uncommon, can be ectopic.

## **Tubal Ligation Reversal**

Microsurgical reconnection of previously divided or

blocked fallopian-tube segments to restore a possible path for sperm and egg; success depends on remaining tube length, age, scarring, and original method.

## **Tubal Pregnancy**

An ectopic pregnancy that implants in a fallopian tube instead of the uterus. It can rupture and cause internal bleeding; one-sided pain, vaginal bleeding, dizziness, or fainting may occur.

## **Tubercle**

A small rounded projection or nodule; it may refer to bone anatomy or a tuberculous lesion.

## **Tuberculide**

A skin eruption caused by an immune reaction to tuberculosis elsewhere in the body. The skin lesions usually contain no bacteria and may improve when the underlying tuberculosis is treated.

## **Tuberculin**

Protein material from *Mycobacterium tuberculosis* used in a skin test for immune sensitization. A positive result suggests exposure or vaccination but does not by itself prove active tuberculosis or show when exposure occurred.

## **Tuberculin Skin Test**

A small amount of tuberculosis protein is injected into the skin. The raised, firm area is measured 2 to 3 days later; the result depends on its size and the person's risk factors and does not by itself distinguish latent from active tuberculosis.

## **Tuberculin Test**

A skin test that checks whether the immune system has reacted to tuberculosis proteins. A positive result suggests infection or previous vaccination and does not by itself prove active disease.

## **Tuberculoid Leprosy**

A form of leprosy in which the immune system keeps the number of bacteria low. It usually causes a few pale or reddish skin patches with reduced feeling

---

and can injure nearby nerves, causing weakness or numbness.

### **Tuberculosis**

An infection caused by *Mycobacterium tuberculosis* bacteria. It usually affects the lungs but can affect other parts of the body. TB infection may remain inactive, causing no symptoms and not spreading to others, or develop into active TB disease. Active TB in the lungs or throat can spread through the air. Symptoms may include a lasting cough, fever, night sweats, and weight loss.

**Alternate Names:**  
TB

### **Tuberculosis In Children**

Tuberculosis infection in a child, usually acquired by breathing air containing the bacteria from an infectious person. It may be silent or cause cough, fever, poor growth, night sweats, or swollen lymph nodes; testing and treatment depend on exposure and findings.

### **Tuberculosis (TB)**

See *Tuberculosis*.

### **Tuberculous Meningitis**

A life-threatening infection of the coverings of the brain caused by tuberculosis bacteria. It may cause persistent headache, fever, vomiting, neck stiffness, confusion, seizures, or weakness.

### **Tuberculous Pleural Effusion**

See *Pleural Tuberculosis*.

### **Tuberous Sclerosis**

A rare genetic disorder, also called tuberous sclerosis complex, that causes growths in the brain and other organs, including the skin, kidneys, heart, and lungs. It may cause seizures, developmental or behavioral problems, and skin changes; the severity varies widely.

### **Tuberous Sclerosis Complex**

An autosomal dominant genetic condition, usually

caused by a disease-causing change in the *TSC1* or *TSC2* gene, in which noncancerous growths develop in several organs, especially the brain, skin, kidneys, heart, and lungs. It may cause seizures, developmental or learning difficulties, skin changes, kidney problems, or breathing disease. Lifelong monitoring and treatment are tailored to the organs affected.

### **Tubes**

Hollow structures that carry air, fluid, food, or medical instruments. Examples include breathing tubes, fallopian tubes, drainage tubes, and blood vessels; the specific structure must be named for the term to be clinically precise.

### **Tubography**

Contrast imaging of the fallopian tubes, usually as part of hysterosalpingography, to see whether they are open. It may help assess infertility but can cause temporary cramping, bleeding, infection, or rarely injury.

### **Tubo-Ovarian Abscess**

A pocket of pus involving a fallopian tube and ovary, usually caused by a pelvic infection. It can cause pelvic pain, fever, and severe illness and may need antibiotics, drainage, or surgery.

### **Tubular Adenoma**

A benign glandular polyp, most often in the colon, that can develop dysplastic changes and is considered a precursor lesion for some colorectal cancers.

### **Tubular Maximum**

The maximum transport rate across the renal tubular epithelium for a filtered solute (such as glucose, phosphate, or amino acids), representing the point of carrier protein saturation beyond which any additional filtered load spills directly into the urine.

**Alternate Names:**

T<sub>m</sub>; Transport Maximum

### **Tubular Reabsorption**

The process by which kidney tubules return useful water and dissolved substances from filtered fluid to the blood. This prevents the loss of most water,



---

salt, glucose, and other substances in urine.

### **Tubular Secretion**

The movement of selected medicines, waste products, and extra ions from the blood into kidney tubules. This helps the kidneys remove substances from the body and regulate fluid and salt balance.

### **Tubule**

A small tube in the body. In the kidney, tubules change filtered fluid by taking useful water and substances back into the blood and leaving waste to form urine; other tubules carry fluids in different organs.

### **Tubuloglomerular Feedback**

A kidney control system that adjusts blood flow and filtration according to how much salt reaches part of a kidney tubule. It helps keep filtration and fluid balance reasonably stable.

### **Tubulointerstitial Diseases of the Kidney**

Disorders that primarily injure renal tubules and the surrounding interstitial tissue rather than the glomeruli. They may cause impaired urine concentration, electrolyte or acid-base abnormalities, sterile pyuria, mild proteinuria, and acute or chronic loss of kidney function.

### **Tubulointerstitial Nephritis**

A pattern of renal parenchymal inflammation predominantly centered on the renal interstitial compartment and tubules rather than the glomeruli, classically triggered by allergic adverse reactions to medications (beta-lactams, NSAIDs, PPIs), autoimmune diseases, or direct infections.

#### **Alternate Names:**

TIN; Interstitial Nephritis

### **Tubulointerstitial Nephropathy**

Kidney damage involving the small tubes that process urine and the tissue around them. Medicines, infections, toxins, autoimmune disease, and inherited conditions can cause it and may reduce kidney function.

### **Tularaemia**

A zoonotic infection caused by *Francisella tularensis*, acquired from animals, ticks, contaminated water, or aerosols.

### **Tularemia**

Tularemia is infection with *Francisella tularensis* acquired from ticks, animals, contaminated water, or aerosols and causing ulcer-glandular, pulmonary, or other disease.

#### **Alternate Names:**

Deerfly Fever; Rabbit Fever; *Francisella Tularensis* Infection

### **Tularemia Blood Test**

A tularemia blood test looks for antibodies or other evidence of infection with *Francisella tularensis*; antibodies may not appear early in illness.

### **Tulle Gras**

A paraffin-impregnated gauze dressing used to cover wounds and burns.

### **Tullio Phenomenon**

Dizziness, eye movements, or balance problems triggered by loud sounds. It can occur with inner-ear conditions such as superior semicircular canal dehiscence or a perilymphatic fistula, but testing is needed to find the cause.

#### **Alternate Names:**

Sound-Induced Vertigo; Acoustic Vestibular Phenomenon

### **Tumescence**

Swelling or enlargement caused by increased blood flow, fluid, or tissue growth. The term commonly describes erection but can also refer to temporary or abnormal enlargement in other body tissues.

### **Tumor**

An abnormal mass formed by cells growing more than they should or surviving when they should not. A tumor may be noncancerous and remain localized, or cancerous and invade nearby tissue or spread.

---

## **Tumor Board**

A meeting in which doctors from different specialties review a person's cancer diagnosis and test results together. They discuss the possible treatment options and coordinate a care plan.

## **Tumor Burden**

The total amount of cancer in the body, estimated from the number, size, and spread of tumor deposits. It can help doctors assess disease extent and treatment response but does not alone determine a person's outlook.

## **Tumor Differentiation**

How closely tumor cells resemble the normal cells from which they arose. Well-differentiated tumors look and behave more like normal tissue, while poorly differentiated tumors often behave more aggressively.

## **Tumor Embolism**

A blockage caused when cancer cells or a piece of a tumor travels through the bloodstream and lodges in another vessel, most often in the lungs. It can impair blood flow and may cause breathing or other organ problems.

## **Tumor Grade**

A description of how abnormal cancer cells look under a microscope compared with normal cells. Lower-grade tumors often grow more slowly, while higher-grade tumors may grow or spread more quickly; grade is different from cancer stage.

## **Tumor Grading**

The process of examining cancer cells under a microscope and assigning a grade according to how different they look from normal cells. Lower-grade tumors often grow more slowly, while higher-grade tumors may behave more aggressively; grade is different from stage.

## **Tumor Heterogeneity**

Differences between cancer cells in the same tumor or between the original tumor and its spread. These differences can affect how quickly cancer grows and

why some cells respond to treatment while others resist it.

## **Tumor-Infiltrating Lymphocyte Therapy**

A cellular cancer treatment using immune cells removed from a tumor, selected or expanded in the laboratory, and returned to the person to attack cancer cells. It is used for selected advanced cancers and requires specialized treatment because serious immune and treatment-related complications can occur.

## **Tumor Lysis Syndrome**

A medical emergency caused by rapid cancer-cell breakdown, releasing substances that disturb blood chemistry. It can damage kidneys, trigger seizures or abnormal heart rhythms, and requires prevention, monitoring, and urgent treatment.

## **Tumor Marker**

A substance found in blood, urine, tissue, or another body fluid that may be produced by cancer or by the body's response to it. Tumor markers can help monitor treatment or recurrence and sometimes guide diagnosis, but most are not specific enough to prove cancer alone.

## **Tumor Mutational Burden**

The number of gene changes found in a sample of cancer. A high number may help predict response to some immunotherapy, but the result depends on the test, cancer type, and other tumor features.

## **Tumor Necrosis Factor**

An immune-system messenger that promotes inflammation, fever, and activation of other immune cells. Helpful during infection, it can also contribute to autoimmune disease, tissue damage, and wasting when produced excessively.

## **Tumor Necrosis Factor Receptor-Associated Periodic Syndrome**

A rare inherited autoinflammatory disorder caused by variants in the TNFRSF1A gene, producing recurrent prolonged fever, muscle pain, abdominal pain, rash, and inflammation; it is also called

---

TRAPS.

### **Tumor Staging**

The process of determining how large a cancer is and whether it has spread to nearby lymph nodes or distant organs. The results help guide treatment and estimate outlook.

### **Tumor Suppressor**

A gene or protein that helps prevent abnormal cell growth by slowing cell division, repairing DNA damage, or removing damaged cells. Loss of this protection can contribute to cancer.

### **Tumor Suppressor Gene**

A gene that helps prevent uncontrolled cell growth by slowing division, repairing DNA damage, or directing abnormal cells to die. If both copies or the gene's function are lost or disabled, cells may grow abnormally and cancer risk can increase.

### **Tumor-Suppressor Gene**

A gene that helps prevent uncontrolled cell growth by slowing cell division, repairing damaged DNA, or removing abnormal cells. Loss of its protective function can increase the chance of cancer.

### **Tumour**

An abnormal mass caused by uncontrolled or inappropriate cell growth; it may be noncancerous or cancerous.

### **Tungiasis**

A skin infestation caused by the sand flea *Tunga* penetrans, producing an itchy or painful lesion, usually on the feet.

### **Tunica Adventitia**

The tough outer layer of a blood vessel. It contains supportive tissue, nerves, and small vessels and helps anchor the vessel to nearby structures.

### **Tunica Albuginea**

A tough layer of connective tissue surrounding the testis. It helps protect the testicular tissue and

sends inward partitions that divide it into lobules.

### **Tunica Media**

The middle layer of a blood-vessel wall, made mainly of muscle and elastic tissue. It can tighten or relax to change vessel width and help control blood pressure and blood flow.

### **Tunica Vaginalis**

A thin, slippery covering around most of the testis, formed from the abdominal lining during testicular descent. It allows the testis to move smoothly inside the scrotum.

### **Tunnel Vision**

Loss of peripheral visual field leaving a narrowed central field, caused by glaucoma, retinal disease, optic-nerve or brain lesions, severe hypoxia, or other conditions.

### **Turbinates**

Curved structures inside the nose that warm, moisten, and filter breathed air. They contain bone covered by soft tissue and can swell during allergies, infection, or irritation and make nasal breathing difficult.

### **Turbinates Hypertrophy**

Long-lasting enlargement of the tissues inside the nose, often from allergies, irritation, sinus disease, or hormonal changes. It can block nasal airflow and cause a stuffy nose, mouth breathing, snoring, or reduced smell.

### **Turbinate Surgery**

An operation that reduces or reshapes swollen tissues along the side walls of the nose to improve airflow. It may help persistent nasal blockage when other treatment fails; bleeding, crusting, dryness, scarring, or ongoing obstruction are possible.

### **Turcot Syndrome**

A rare inherited cancer-predisposition pattern involving colon polyps or colon cancer together with a brain tumor. It can result from mismatch-repair

---

gene changes or the APC gene change seen in familial adenomatous polyposis, so genetic counseling and screening are important.

**Alternate Names:**

Brain Tumor-Polyposis Syndrome

**Turgor**

The firmness and springiness of body tissue caused partly by its fluid content. Skin that remains tented after being gently pinched may suggest dehydration, but this sign is not reliable by itself.

**Turner Hypoplasia**

A defect in the enamel of a permanent tooth caused by infection or injury involving the baby tooth above it. The affected tooth may have a discolored, rough, or weakened area.

**Turner's Syndrome**

A chromosome condition affecting females in which one X chromosome is missing or altered. It commonly causes short stature and reduced ovarian function, with delayed or absent puberty and infertility. Heart, kidney, hearing, and thyroid problems may also occur, so lifelong medical follow-up is important.

**Turner Syndrome**

A chromosome condition affecting females in which one X chromosome is missing or altered. It commonly causes short stature and reduced ovarian function, with delayed or absent puberty and infertility. Heart, kidney, hearing, and thyroid problems may also occur, so lifelong medical follow-up is important.

**Turning Patients Over In Bed**

Repositioning a patient at planned intervals to protect skin, improve comfort and breathing, prevent pressure injuries and contractures, and reduce complications of immobility.

**Turpentine Oil Poisoning**

Poisoning from swallowing or inhaling turpentine oil. It can irritate or damage the lungs, stomach, and nervous system and may cause chemical pneumonia.

**Tussis**

Cough; a forceful reflex expiration that helps clear secretions, irritants, or foreign material from the airways.

**TVS**

TVS commonly means transvaginal sonography, an ultrasound examination using a probe in the vagina to assess pelvic organs and early pregnancy.

**T Wave**

The part of an ECG that shows the heart's lower chambers resetting electrically after each beat. Changes in its height or direction can occur with reduced blood flow, abnormal body salts, heart strain, or other conditions.

**T-Wave Alternans**

T-wave alternans is a beat-to-beat change in the size or shape of the heart's T wave and may indicate increased risk of dangerous ventricular rhythms.

**Twin**

A twin is one of two offspring from the same pregnancy, either identical from one fertilized egg or fraternal from two eggs and two sperm.

**Twin Placenta**

The number and arrangement of placentas and the sacs surrounding twins. Knowing whether twins share a placenta or a sac helps estimate risks and determines how closely the pregnancy is monitored.

**Twin Pregnancy**

A pregnancy involving two fetuses. Clinical risk depends on chorionicity and amnionicity, with increased risks including preterm birth, fetal growth problems, and hypertensive disorders.

**Twins**

Two offspring from the same pregnancy, either monozygotic or dizygotic.

**Twin-To-Twin Transfusion Syndrome**

A serious complication of an identical-twin preg-

---

nancy in which shared placental blood vessels send too much blood to one twin and too little to the other. It requires urgent specialist monitoring and treatment.

### **Twin-Twin Transfusion Syndrome**

A complication of an identical-twin pregnancy sharing one placenta, in which blood flows unevenly between the twins. One twin may receive too little blood and fluid, while the other receives too much; both need urgent specialist monitoring and treatment.

### **Twitch**

A brief involuntary muscle contraction or movement, which may be benign, medication-related, stress-related, or a sign of nerve, muscle, or brain disease.

### **Twitching**

Repeated involuntary muscle contractions or fasciculations, commonly caused by fatigue, stimulants, electrolyte disturbance, nerve irritation, medicines, or neurologic disease.

### **Tympanic**

Tympanic means relating to the eardrum or middle ear, or resembling a drum in shape or sound.

### **Tympanic Bulla**

The tympanic bulla is a hollow or expanded bony chamber surrounding the middle ear, especially prominent in many mammals.

### **Tympanic Cavity**

The air-filled chamber of the middle ear. It contains the eardrum and three tiny bones that transmit sound vibrations toward the inner ear, while a passage to the throat helps equalize air pressure.

### **Tympanic Hyperectasis**

Abnormal bulging or stretching of the eardrum, usually related to long-lasting pressure changes in the middle ear. It may cause blocked-ear sensations or reduced hearing and requires assessment for the underlying cause.

### **Tympanic Membrane**

The thin eardrum separating the outer ear canal from the middle ear. Sound makes it vibrate, and those vibrations move the middle-ear bones; infection, injury, or pressure can tear it.

### **Tympanic Membrane Perforation**

A tear or hole in the eardrum caused by infection, trauma, pressure changes, or instrumentation, potentially producing pain, drainage, or conductive hearing loss.

### **Tympanites**

Abdominal distension caused by excessive gas in the stomach or intestines, producing a drum-like sound on percussion and sometimes pain, belching, or altered bowel movements.

### **Tympanometer**

An instrument that changes pressure in the ear canal and measures movement of the tympanic membrane and middle-ear system. It is used to perform tympanometry.

### **Tympanometry**

An objective test that varies air pressure in the ear canal while measuring movement of the tympanic membrane and middle-ear pressure. It helps assess middle-ear fluid, eustachian-tube dysfunction, perforation, and ossicular or tympanic-membrane abnormalities.

### **Tympanoplasty**

Surgical repair or reconstruction of the tympanic membrane and, when necessary, the middle-ear ossicles to close a perforation or improve hearing.

### **Tympanosclerosis**

Scarring and calcification of the tympanic membrane or middle-ear lining, often following recurrent infection or previous inflammation. It may be asymptomatic or cause conductive hearing loss when the ossicles are affected.

#### **Alternate Names:**

Myringosclerosis; Tympanic Membrane Calcifica-

---

tion

### **Tympanostomy Tube Insertion**

The most commonly performed pediatric surgical procedure worldwide, in which a tiny cylindrical ventilation tube (grommet) is placed across the tympanic membrane (eardrum). An ear, nose, and throat (ENT) specialist makes a microscopic incision in the eardrum (myringotomy), suctions out accumulated thick, sticky fluid from the middle ear cavity, and positions the tiny tube across the incision. The tube acts as an artificial pressure-equalizing valve, providing continuous airflow ventilation to the middle ear and allowing trapped fluid to drain freely, effectively compensating for an immature or malfunctioning Eustachian tube. It is primarily indicated for persistent middle ear fluid (otitis media with effusion or “glue ear”) causing conductive hearing impairment and speech delays, or for frequent recurrent acute ear infections. Tubes typically stay in place for 6 to 18 months before naturally extruding on their own as the eardrum heals.

#### **Alternate Names:**

Ear Tube Insertion; Grommet Insertion; Myringotomy with Tube Insertion; Tympanostomy

### **Tympanostomy Tubes**

Small tubes placed through the eardrum to let air into the middle ear and help fluid drain. They may be recommended for repeated ear infections or persistent fluid causing hearing problems; drainage, blockage, scarring, or early tube loss can occur.

### **Tympanum**

The middle-ear cavity or, in some contexts, the tympanic membrane.

### **Tympany**

Tympany is a drumlike sound on percussion, often produced by gas in a hollow organ such as the stomach or intestine.

### **Tyndall Effect**

The optical scattering of light by suspended inflammatory cells and proteinaceous exudate (‘aqueous

flare’) in the anterior chamber of the eye viewed with a narrow slit-lamp biomicroscope beam against a dark background, serving as a cardinal physical sign of active anterior uveitis (iritis / iridocyclitis).

#### **Alternate Names:**

Aqueous Flare; Anterior Chamber Tyndall Phenomenon

### **Type 1 Diabetes**

A long-term disease in which the immune system destroys the pancreas cells that make insulin. Without insulin treatment, blood sugar rises dangerously and can lead to diabetic ketoacidosis.

#### **Alternate Names:**

T1D; Insulin-Dependent Diabetes Mellitus; IDDM

### **Type 1 Diabetes In Children**

Type 1 diabetes diagnosed during childhood, causing high blood sugar and requiring insulin, monitoring, nutrition planning, and education.

### **Type 1 Diabetes Mellitus**

Type 1 diabetes is an autoimmune disease in which the immune system attacks and destroys the insulin-producing beta cells of the pancreas, resulting in absolute insulin deficiency.

### **Type 2 Diabetes**

A long-term disease in which the body does not use insulin effectively and gradually makes less of it. High blood sugar can damage the eyes, kidneys, nerves, heart, and blood vessels.

#### **Alternate Names:**

T2D; Non-Insulin-Dependent Diabetes Mellitus; NIDDM

### **Type 2 Diabetes In Children**

Type 2 diabetes beginning during childhood or adolescence, often involving insulin resistance and requiring lifestyle and medical management.

### **Type 2 Diabetes Meal Planning**

Nutritional guidance focusing on consistent carbohydrate intake, portion control, dietary fiber, lean proteins, and healthy fats to maintain target blood



---

glucose and body weight.

**Alternate Names:**

Diabetic Meal Planning; Nutritional Therapy for Type 2 Diabetes; Diabetic Diet Plan

**Type 2 Diabetes Medications**

Alternate name; see *Oral Antidiabetic Agents*.

**Type 2 Diabetes Mellitus**

Type 2 diabetes is a chronic metabolic disorder characterized by insulin resistance and progressive pancreatic beta-cell dysfunction, leading to hyperglycemia.

**Type 3B Diabetes**

Type 3b diabetes is a form of diabetes caused by disease or removal of the pancreas, which can impair both insulin and digestive-enzyme production.

**Type Ia Glycogen Storage Disease**

Von Gierke disease, caused by glucose-6-phosphatase deficiency. It typically causes severe fasting hypoglycemia, lactic acidosis, hyperlipidemia, hyperuricemia, hepatomegaly, and impaired growth.

**Type Ib Glycogen Storage Disease**

A glycogen-storage disorder caused by impaired glucose-6-phosphate transport in the endoplasmic reticulum. It can cause fasting hypoglycemia, lactic acidosis, hepatomegaly, neutropenia, recurrent infections, and inflammatory bowel disease.

**Type I Hypersensitivity**

A rapid allergic reaction in which the immune system releases substances such as histamine after exposure to an allergen. It can cause hay fever, asthma symptoms, hives, food allergy, or life-threatening anaphylaxis.

**Type II Hypersensitivity**

An immune reaction in which antibodies attach to a person's cells or tissues and cause injury or destruction. It contributes to some blood-cell disorders, medicine reactions, and autoimmune diseases.

**Type III GSD (Forbes-Cori)**

An inherited glycogen-storage disorder caused by debranching-enzyme deficiency. It can cause fasting hypoglycemia, hepatomegaly, muscle weakness, elevated liver enzymes, and sometimes cardiomyopathy, with severity varying by subtype.

**Type III Hypersensitivity**

An immune reaction in which clusters of antibodies and foreign or abnormal proteins settle in tissues and trigger inflammation. It can damage the skin, joints, blood vessels, kidneys, or other organs.

**Type III Hypersensitivity Reaction**

Tissue injury caused when antigen-antibody clusters deposit in tissues and activate inflammation. This reaction can damage blood vessels, joints, kidneys, skin, or other organs, as in serum sickness or lupus.

**Type II Muscle Fiber**

A fast-contracting muscle fiber that can produce high force quickly but usually tires sooner than a type I fiber.

**Type I Muscle Fiber**

A slow-contracting muscle fiber that uses oxygen efficiently and can keep working for a long time. It is important for posture and endurance.

**Type IV GSD (Andersen)**

An inherited disorder caused by branching-enzyme deficiency, leading to abnormal glycogen accumulation. Progressive liver disease, cirrhosis, growth problems, muscle disease, or neuromuscular involvement may occur.

**Type IV Hypersensitivity**

A delayed immune reaction driven mainly by T cells rather than antibodies. It usually develops hours to days after exposure and can cause contact dermatitis, some medicine reactions, or inflammation in an organ.

**Types Of Ileostomy**

Types of ileostomy include end, loop, and continent

---

procedures, which divert stool through an opening in the abdominal wall when the colon or rectum cannot be used normally.

### **Type V Glycogen Storage Disease**

An inherited muscle disorder, also called McArdle disease, in which muscle cells cannot properly break down glycogen for energy. Exercise can cause muscle pain, cramps, and sometimes dark urine from muscle injury.

### **Type VI Glycogen Storage Disease**

Hers disease, caused by hepatic glycogen-phosphorylase deficiency. It usually causes mild fasting hypoglycemia, hepatomegaly, growth delay, and elevated liver enzymes, with symptoms often improving with age.

### **Type VII Glycogen Storage Disease**

Tarui disease, caused by phosphofructokinase deficiency in muscle and red blood cells. Exercise intolerance, muscle cramps, rhabdomyolysis, and hemolytic anemia may occur, sometimes with a second-wind phenomenon.

### **Typhlitis**

A serious infection and inflammation of the first part of the large intestine, usually in someone with very low white-cell counts after cancer treatment. It can cause right-sided abdominal pain, fever, diarrhea, and bowel damage and needs urgent care.

#### **Alternate Names:**

Neutropenic Enterocolitis; Ileocecal Syndrome; Cecitis

### **Typhoid**

Alternate name; see *Typhoid Fever*.

### **Typhoid Fever**

A severe systemic bacterial infection caused by *Salmonella* Typhi, transmitted through contaminated food or water. It causes a gradually rising step-ladder fever, headache, stomach pain, constipation or “pea-soup” diarrhea, a relatively slow heart rate, and faint pink spots on the abdomen (rose spots). Untreated infection can lead to ulcera-

tion of the small intestine with severe bleeding or bowel perforation in the third week. Diagnosis is made by blood or bone marrow culture, and treatment requires targeted antibiotics like ceftriaxone or azithromycin.

#### **Alternate Names:**

Enteric Fever; Salmonella Typhi Infection; Typhoid

### **Typhus**

A group of bacterial infections spread mainly by lice, fleas, or mites. They can cause fever, headache, rash, and serious illness affecting several organs; the type and treatment depend on the bacteria and insect involved.

### **Typhus Fever**

A group of bacterial infections spread by lice, fleas, or mites. They commonly cause sudden fever, severe headache, weakness, and sometimes rash or organ injury; the exact illness determines testing and antibiotic treatment.

### **Typical Antipsychotics**

Older antipsychotic medicines used for conditions such as psychosis, mania, or severe agitation. They can reduce hallucinations and delusions but may cause stiffness, tremor, involuntary movements, sleepiness, or other serious effects.

### **Tyramine**

A naturally occurring amine found in some aged or fermented foods. In people taking monoamine oxidase inhibitors, excessive tyramine can cause a dangerous rise in blood pressure.

### **Tyrosine**

An amino acid used to build proteins and make thyroid hormones, melanin, and adrenaline-like chemicals. The body usually makes enough from phenylalanine, but inherited metabolic disorders can disrupt this pathway.

#### **Alternate Names:**

L-Tyrosine; 4-Hydroxyphenylalanine; Tyr

---

## Tyrosine Kinase Inhibitors

A major class of targeted antineoplastic pharmacological agents that competitively inhibit the adenosine triphosphate-binding catalytic pocket of tyrosine kinase enzymes, blocking downstream oncogenic proliferative signaling cascades (e.g., imatinib for BCR-ABL, osimertinib for EGFR).

### Alternate Names:

TKIs

## Tyrosinemia

An inherited disorder in which the body cannot break down the amino acid tyrosine normally. Harmful substances can build up and damage the liver, kidneys, nerves, or eyes; severity depends on the type and treatment.

## Tzanck Smear

A microscope test of cells scraped from a blister or sore. It may suggest a herpes infection or pemphigus, but it cannot reliably identify the cause and is often replaced by more specific laboratory tests.

## Tardive Oral Dyskinesia

Involuntary repetitive movements of the mouth, tongue, jaw, or face that develop after prolonged exposure to dopamine-blocking medicines. It may persist after the medicine is stopped and needs medication review and specialist assessment.

## Tarsal Cyst

A fluid-filled or other cystic lesion in the ankle or foot region, which may cause a lump, pain, pressure, or nerve symptoms. Examination and imaging determine whether it is a ganglion, synovial cyst, or another lesion.

## TB (Tuberculosis)

Alternate name; see *Tuberculosis*.

## TBM

Abbreviation; see *Tuberculous Meningitis*.

## TCOF1

A gene in which disease-causing variants can pro-

duce Treacher Collins syndrome, a craniofacial developmental disorder. Genetic counseling and assessment of breathing, hearing, vision, feeding, and speech may be needed.

## TCS

Abbreviation; see *Treacher Collins Syndrome*.

## TENS

Abbreviation; see *Transcutaneous Electrical Nerve Stimulation*.

## TMJ Disorder

Alternate name; see *Temporomandibular Joint Disorder*.

## Tonic Pupil Syndrome

A pupil that is enlarged or irregular and reacts poorly to light but constricts slowly with near focus, usually because of damage to postganglionic parasympathetic nerves. New pupil asymmetry, headache, drooping eyelid, or neurologic symptoms requires urgent assessment.

## TORCH Syndrome

A group of congenital infections caused by Toxoplasma, Other infections, Rubella, Cytomegalovirus, and Herpes simplex, which may cause growth restriction, rash, jaundice, eye disease, hearing loss, brain abnormalities, or other newborn problems.

## Torn Rotator Cuff

A partial or complete tear of one or more rotator-cuff tendons, causing shoulder pain, weakness, night symptoms, or difficulty lifting or rotating the arm.

## Transient Acantholytic Dermatitis

A recurrent itchy papular rash caused by temporary loss of cohesion between skin cells. It often affects the trunk of older adults and may be triggered by heat, sweating, sunlight, medicines, or illness; it is also called transient acantholytic dermatosis or Grover disease.

---

### **Transient Ischemic Attack (TIA)**

Alternate name; see *Transient Ischemic Attack*.

### **Transthyretin Familial Amyloid Polyneuropathy**

An inherited transthyretin amyloidosis in which abnormal protein deposits damage peripheral and autonomic nerves. It may cause progressive numbness, pain, weakness, blood-pressure or heart-rate changes, gastrointestinal symptoms, and sometimes heart disease.

### **Transthyretin Methionine-30 Amyloidosis (Type I)**

A hereditary transthyretin amyloidosis caused by a methionine-at-position-30 transthyretin variant, classically producing progressive peripheral and autonomic neuropathy with variable cardiac, kidney, or eye involvement.

### **Traumatic Brain Injury (Concussion)**

A head injury that disrupts brain function temporarily or permanently. Concussion is a mild traumatic brain injury and may cause headache, dizziness, confusion, memory problems, or sensitivity to light; worsening headache, repeated vomiting, seizure, weakness, or reduced alertness is an emergency.

### **Traveler's Diarrhea**

Diarrhea during or shortly after travel, usually caused by contaminated food or water and commonly by bacteria, viruses, or parasites. Oral rehydration is important; blood, high fever, severe dehydration, or persistent symptoms require medical evaluation.

### **Treacher Collins-Franceschetti Syndrome 1**

A craniofacial developmental disorder, usually caused by a variant affecting TCOF1, that may cause underdeveloped cheekbones and jaw, malformed ears, hearing loss, cleft palate, and airway or feeding problems.

### **Trisomy 13 Syndrome**

Alternate name; see *Patau Syndrome*.

### **Trisomy 18 Syndrome**

Alternate name; see *Edwards Syndrome*.

### **Tuberose Sclerosis**

Historical or variant name for tuberous sclerosis complex, a genetic disorder causing benign tumors in multiple organs, including the brain, skin, kidneys, heart, lungs, or eyes.

### **Tuberous Sclerosis-1**

A form of tuberous sclerosis complex associated with pathogenic variants in the *TSC1* gene. Features may include seizures, developmental or behavioral differences, skin findings, and benign tumors in several organs.

### **Turner-Varny Syndrome**

Alternate name; see *Turner Syndrome*.

### **Twin Twin Transfusion Syndrome**

Alternate name; see *Twin-to-Twin Transfusion Syndrome*.

### **Temporary Blindness**

A transient loss of vision in one or both eyes. It can result from retinal or brain ischemia, migraine, seizures, changes in blood pressure, eye disease, or other causes; sudden new loss of vision requires emergency assessment.

# U

## UA (Urinalysis)

UA commonly means urinalysis, laboratory examination of urine for blood, protein, cells, glucose, infection, and other findings.

## Ubiquitin-Proteasome System

The primary non-lysosomal intracellular protein degradation pathway in eukaryotic cells, in which damaged, misfolded, or regulatory proteins are tagged with polyubiquitin chains by ubiquitin ligases (E1, E2, E3) and subsequently recognized and degraded by the 26S proteasome complex.

### Alternate Names:

UPS; Ubiquitination Degradation Pathway

## UBT (Urea Breath Test)

UBT commonly means urea breath test, which detects active *Helicobacter pylori* infection by measuring labeled carbon dioxide after ingested urea.

## UDP-Glucuronosyltransferase

A group of enzymes, found mainly in the liver, that helps the body process and remove bilirubin, medicines, hormones, and other substances. Inherited differences in these enzymes can cause some types of jaundice or change how medicines work.

## UDP-Glucuronosyltransferase Deficiency

Reduced activity of a liver enzyme that prepares bilirubin for removal from the body. Inherited deficiency can cause intermittent mild jaundice or, in severe forms, dangerous bilirubin buildup.

## Uhl Anomaly

A rare heart defect present from birth in which

much of the muscle of the right lower chamber is missing. The chamber may pump poorly, causing blue coloring, abnormal rhythms, or heart failure in infancy or childhood.

## Uhthoff Phenomenon

A transient, reversible worsening of pre-existing neurological symptoms or visual blurring occurring in patients with multiple sclerosis when body temperature is elevated by physical exercise, hot baths, or fever, resulting from heat-induced conduction block in chronically demyelinated axons.

### Alternate Names:

Uhthoff Sign

## Ulcer

A crater-like open sore caused by loss of the surface layers of skin or a mucous membrane, sometimes extending into deeper tissue. Causes include poor blood flow, pressure, infection, nerve damage, inflammation, injury, and cancer; the appearance and cause determine treatment.

## Ulceration

The formation of an open sore caused by loss of surface tissue, usually involving skin or a mucous membrane and sometimes extending into deeper tissue.

## Ulcerative Colitis

A long-term disease causing inflammation and open sores in the inner lining of the rectum and colon. It usually starts in the rectum and spreads continuously through part of the colon. It may cause repeated diarrhea, blood or mucus in the stool, ur-

---

gent bowel movements, abdominal cramps, tiredness, weight loss, or a feeling that the bowel has not emptied. Symptoms often flare and then improve. Severe disease may cause heavy bleeding, anemia, dehydration, or a hole in the bowel; long-standing extensive disease increases the risk of colorectal cancer.

**Alternate Names:**

UC; Idiopathic Proctocolitis

### Ulcerative Colitis Pathology

Chronic, usually continuous mucosal inflammation beginning in the rectum and extending proximally. Findings include crypt architectural distortion, cryptitis, crypt abscesses, mucosal ulceration, and inflammatory polyps in active or longstanding disease.

### Ulcerative Keratitis

An ulcerative infection or inflammatory defect of the cornea that can cause pain, redness, photophobia, and impaired vision and may threaten sight if untreated.

### Ulcerative Proctitis

Inflammation and ulceration limited to the lining of the rectum, often causing rectal bleeding, urgency, tenesmus, mucus, and diarrhea.

### Ulcer Healing Drugs

Medicines used to promote healing of gastrointestinal ulcers, usually by suppressing acid secretion or protecting the mucosa; treatment also addresses causes such as *H. pylori* infection or NSAID use.

### Ulcers

Plural of ulcer; see *Ulcer*.

### Ulerythema

A group of follicular keratinization disorders causing rough plugged follicles, redness, and sometimes atrophic scarring, most often on the cheeks, eyebrows, or upper arms. The term may refer to variants such as ulerythema ophryogenes and requires clinical distinction from eczema or acne.

### Ullrich Congenital Muscular Dystrophy

An inherited muscle disease caused by problems with collagen VI, a protein that supports muscle and connective tissue. It begins in infancy with weakness, low muscle tone, tight joints near the body, unusually loose joints in the hands and feet, and later breathing problems.

**Alternate Names:**

UCMD; Ullrich Scleroatonic Muscular Dystrophy

### Ulna

The forearm bone on the side of the little finger. Its upper end forms most of the hinge of the elbow, and its lower end helps form the wrist. Together with the radius, it supports rotation of the forearm and movement of the hand.

### Ulnar

Relating to the ulna or to the side of the forearm and hand nearest the little finger. Pressure on the ulnar nerve in this area can cause tingling or numbness in the little and ring fingers.

### Ulnar Artery

A major artery running along the little-finger side of the forearm and into the hand. It supplies blood to the forearm and hand and helps form the hand's arterial arches.

### Ulnar Claw Hand

A distinctive resting hand deformity (Duchenne claw hand) resulting from a distal ulnar nerve injury at the wrist, characterized by hyperextension of the metacarpophalangeal (MCP) joints and hyperflexion of the proximal and distal interphalangeal (PIP/DIP) joints of the fourth and fifth digits due to loss of the medial lumbricals and interossei.

**Alternate Names:**

Duchenne Claw Hand; Main en Griffe

### Ulnar Collateral Ligament

A thick triangular ligament on the medial side of the elbow joint (consisting of anterior, posterior, and transverse bands) that stabilizes against valgus stress, commonly torn in overhead throwing athletes ('Tommy John' injury).



---

**Alternate Names:**

UCL of the Elbow; Medial Collateral Ligament of Elbow

**Ulnar Deviation**

A characteristic musculoskeletal deformity of the hand in advanced chronic rheumatoid arthritis, characterized by the abnormal lateral (ulnar-directed) subluxation and drifting of the fingers at the metacarpophalangeal joints resulting from extensor tendon stretching and volar plate destruction.

**Alternate Names:**

Ulnar Drift

**Ulnar Drift**

A classic progressive hand deformity seen in chronic rheumatoid arthritis characterized by abnormal ulnar deviation of the second through fifth fingers at the metacarpophalangeal (MCP) joints, resulting from chronic synovitis, stretching of collateral ligaments, and extensor tendon subluxation.

**Alternate Names:**

Ulnar Deviation of Digits; Rheumatoid Ulnar Drift

**Ulnar Hypoplasia**

Underdevelopment or partial absence of the ulna, the forearm bone on the little-finger side. Present from birth in most cases, it may shorten or bend the forearm, limit elbow or wrist movement, and occur with finger or other body differences.

**Ulnar Nerve**

A nerve running from the neck through the inner arm and behind the elbow to the hand. It controls some hand and forearm muscles and carries feeling from the little finger and part of the ring finger.

**Ulnar Nerve Dysfunction**

Impairment of the ulnar nerve, commonly caused by compression at the elbow (cubital tunnel) or wrist (Guyon canal). Symptoms include numbness or tingling in the little and ring fingers, weakened grip, and hand-muscle wasting; severe lesions may produce an ulnar claw.

**Ulnar Nerve Entrapment**

Compression or irritation of the ulnar nerve, commonly at the elbow or wrist, causing numbness or tingling in the ring and little fingers and sometimes weakness of the hand.

**Ulnar Neuropathy**

A peripheral compression neuropathy of the ulnar nerve occurring most commonly at the elbow within the cubital tunnel (cubital tunnel syndrome) or at the wrist within Guyon canal, presenting with numbness and paresthesias in the fifth and medial half of the fourth digit, intrinsic hand weakness, and diminished grip strength.

**Alternate Names:**

Ulnar Nerve Entrapment; Cubital Tunnel Neuropathy

**Ulnar Neuropathy at Guyon Canal**

Entrapment or compressive neuropathy of the ulnar nerve as it traverses the fibro-osseous canal (pisohamate tunnel) at the wrist between the pisiform and the hook of the hamate. Subdivided into zone I (proximal trunk lesion with combined motor and sensory deficits), zone II (deep motor branch lesion with pure motor clawing and intrinsic muscle weakness sparing hypothenar sensation), and zone III (superficial sensory branch lesion with pure sensory paresthesias of the fifth and medial fourth digits).

**Alternate Names:**

Guyon Canal Syndrome; Handlebar Palsy

**Ulnar Paradox**

The apparently unexpected finding that damage to the ulnar nerve at the wrist can cause more clawing of the fingers than damage higher in the arm. The difference occurs because some finger-bending muscles remain active after a wrist injury.

**Ulnar-Sided Wrist Pain**

Alternate name; see *Ulnar wrist pain*.

**Ulnar Tunnel Syndrome**

An entrapment neuropathy of the ulnar nerve as it passes through Guyon canal (the ulnar tunnel at the wrist between the pisiform bone and hook

---

of the hamate), causing sensory loss in the medial ring and little fingers and motor weakness of the hypothenar and intrinsic hand muscles.

**Alternate Names:**

Guyon Canal Syndrome; Handlebar Palsy

**Ulnar Vein**

Veins on the little-finger side of the forearm that carry blood away from the forearm and hand. They join other deep veins near the elbow to continue toward the heart.

**Ulnar Wrist Pain**

Pain on the side of the wrist nearest the little finger, often involving the ulna or nearby tendons and ligaments.

**Alternate Names:**

Ulnar-Sided Wrist Pain

**Ultrafiltration**

A process in which pressure pushes water and small dissolved substances through a very fine filter while larger particles remain behind. The kidneys use it to begin forming urine, and dialysis uses it to remove excess fluid.

**Ultra-processed food**

Ultra-processed food is an industrially formulated product made mostly from refined ingredients and additives, often with little intact whole food. Frequent high intake is associated with poorer health outcomes, although effects vary by the food and overall diet.

**Ultrasonic Therapy**

Treatment using high-frequency sound energy to heat, move, break up, or otherwise affect tissue. It is used in selected physical-therapy procedures and some operations; benefits and risks depend on the device, body site, and condition.

**Ultrasonography**

A scan using high-frequency sound waves to create images of organs, tissues, vessels, or a fetus without radiation.

**Ultrasound**

An imaging test that uses high-frequency sound waves to show organs, blood vessels, tissues, or a developing fetus without ionizing radiation. Different examinations can assess the abdomen, pelvis, kidneys, thyroid, heart, blood flow, or testicles.

**Ultrasound Biomicroscopy**

Very high-frequency ultrasound that creates detailed images of the front part of the eye, including the cornea, iris, lens, and drainage angle when an ordinary eye examination is not enough.

**Ultrasound Contrast**

Tiny gas-filled bubbles given into a vein to improve ultrasound images of blood flow and selected organs. They are usually cleared through the lungs but can rarely cause allergic or cardiopulmonary reactions.

**Ultrasound-Guided Biopsy**

An image-guided interventional procedure in which real-time ultrasonography is utilized to continuously visualize and steer a biopsy needle (core needle or fine-needle aspiration) into a target lesion or mass (such as in the thyroid, breast, liver, or lymph nodes) for histological diagnosis.

**Ultrasound-Guided Breast Biopsy**

A percutaneous core-needle or vacuum-assisted tissue sampling procedure performed under continuous real-time sonographic guidance to biopsy solid masses, architectural abnormalities, or complex cystic breast lesions.

**Alternate Names:**

Ultrasound-Guided Breast Biopsy; Sonographically Guided Core Biopsy; Ultrasound Breast Biopsy

**Ultrasound Therapy**

Use of sound waves at therapeutic intensity to heat or mechanically affect tissue. It is used in selected rehabilitation settings, but evidence and suitability vary by condition and it is not appropriate over every body area.

---

## Ultrasound Transducer

The probe placed against the skin or inside a body opening during an ultrasound scan. It sends and receives sound waves to create images or guide a procedure; different probes are used for different body areas.

## Ultraviolet Rays

Invisible electromagnetic radiation with wavelengths shorter than visible violet light. Excess exposure can cause sunburn, photoaging, eye injury, and skin cancers; controlled ultraviolet exposure has selected therapeutic uses.

## Ultraviolet Therapy

Controlled exposure to ultraviolet light used for selected skin disorders such as psoriasis, eczema, and vitiligo; treatment requires eye and skin protection and monitoring for burns and long-term skin cancer risk.

## Umbilical

Relating to the umbilicus or navel, including structures, lesions, discharge, or abnormalities located at or connected with it.

## Umbilical Artery Catheter

A specialized vascular catheter inserted into one of the two umbilical arteries in critically ill neonates, advancing into the descending thoracic aorta (high placement, T6-T8) or infrarenal abdominal aorta (low placement, L3-L5) for continuous arterial blood pressure monitoring and arterial blood gas sampling.

### Alternate Names:

UAC

## Umbilical Artery Doppler

An ultrasound measurement of blood flow through the umbilical cord. It helps assess how well the placenta is supplying the fetus, especially when fetal growth is reduced or pregnancy complications affect placental function.

## Umbilical Catheters

Umbilical catheters provide vascular access through

a newborn's umbilical vessels for fluids, medicines, monitoring, or blood sampling while complications are monitored.

## Umbilical Cord

The connection between a fetus and the placenta. It contains blood vessels that carry oxygen, nutrients, and waste between them and is protected by a soft, cushioning tissue.

## Umbilical Cord Blood Gas

A test of blood taken from the umbilical cord after birth to assess the baby's oxygen supply and acid-base balance around delivery. Arterial and venous results provide different information and must be interpreted with the baby's condition, gestational age, and delivery circumstances.

## Umbilical Cord Complications

Problems involving the umbilical cord, including prolapse, compression, knots, abnormal insertion, short or long cord, and vascular abnormalities. They may reduce fetal oxygenation or complicate delivery and require fetal monitoring and obstetric management.

## Umbilical Cord Prolapse

The umbilical cord slips through the cervix ahead of or alongside the baby, usually after the waters break. The cord can be squeezed and reduce the baby's oxygen supply, making this an obstetric emergency.

## Umbilical Granuloma

A small moist pink or red lump of healing tissue at a baby's navel after the cord falls off. It may ooze clear fluid but should not cause spreading redness, fever, or marked tenderness; a clinician can treat it and exclude infection or another abnormal connection.

### Alternate Names:

Granuloma Umbilicale; Neonatal Umbilical Granuloma

## Umbilical Hernia

A soft bulge caused when tissue pushes through the opening at the navel. Many infant hernias close as the child grows, while adult hernias may enlarge or become trapped and painful and sometimes need

---

surgery.

### **Umbilical Region**

The central part of the abdomen around the navel. Pain, swelling, a hernia, or a lump in this area may arise from the bowel, abdominal wall, blood vessels, or other organs and is assessed with examination and, when needed, imaging.

### **Umbilical Vein**

The umbilical vein carries oxygen- and nutrient-rich blood from the placenta to the fetus during pregnancy.

### **Umbilication**

Formation of a central hollow or navel-like depression in a skin lesion or other body structure. It can occur in some infections, inflammatory lesions, tumors, or normal anatomical features and must be interpreted with the lesion's appearance.

### **Umbilicus**

The navel, a scar on the front of the abdomen left after the umbilical cord separates at birth. It may be affected by a hernia, infection, discharge, or other skin changes.

#### **Alternate Names:**

Belly Button; Omphalos

### **Uncal Herniation**

A life-threatening subtype of transtentorial herniation in which expanding supratentorial mass effect pushes the medial temporal lobe (uncus) downward over the free edge of the tentorium cerebelli, compressing the ipsilateral oculomotor nerve (CN III, causing fixed dilated 'blown' pupil), the ipsilateral cerebral peduncle (contralateral hemiparesis), and the posterior cerebral artery (occipital infarction).

#### **Alternate Names:**

Lateral Transtentorial Herniation; Uncinate Herniation

### **Uncal Transtentorial Herniation**

A life-threatening intracranial herniation syndrome in which an expanding supratentorial mass displaces

the uncus of the medial temporal lobe medially and downward across the tentorial incisura, compressing the ipsilateral oculomotor nerve (cranial nerve III; producing an unreactive dilated pupil) and the ipsilateral cerebral peduncle (producing contralateral hemiparesis).

### **Unconjugated Hyperbilirubinemia**

An elevation of circulating indirect (water-insoluble) bilirubin in blood plasma exceeding physiological limits, resulting from excessive bilirubin overproduction (hemolysis), impaired hepatic uptake (heart failure, portosystemic shunts), or defective glucuronosyltransferase conjugation (Gilbert syndrome, Crigler-Najjar syndrome, neonatal jaundice).

### **Unconscious**

Not awake and not responsive to ordinary external stimuli. Unconsciousness may result from syncope, seizure, trauma, intoxication, metabolic disease, stroke, or other emergencies.

### **Unconsciousness**

Loss of awareness and normal response to surroundings, ranging from brief fainting to prolonged coma. It can result from reduced brain blood flow, injury, seizures, low blood sugar, drugs, or serious illness.

### **Uncoupling Protein**

A family of mitochondrial inner membrane transport proteins (such as UCP1 / thermogenin) that dissipate the proton electrochemical gradient across the inner mitochondrial membrane as heat without ATP synthesis, playing a vital role in non-shivering thermogenesis in brown adipose tissue.

#### **Alternate Names:**

Thermogenin; UCP1

### **Uncover Test**

An ophthalmologic test performed following the cover test to detect latent strabismus (phoria). As the occluder is removed from an eye, the examiner observes whether that eye rapidly refixates from a deviated position back to alignment with the target.

---

## Underventilation

Breathing that is too slow or shallow to remove enough carbon dioxide. It may also lower oxygen and can result from sedatives, weak breathing muscles, severe obesity, lung disease, or brain problems.

## Underweight

Body weight that is lower than expected for a person's height, age, and build. It may be healthy for some people but can also reflect poor nutrition, illness, or unintentional weight loss and is interpreted with the whole clinical picture.

## Undescended Testicle

See *Cryptorchidism*.

## Undescended Testicle Repair

An operation that moves a testicle into the scrotum and fixes it there. It is usually performed in childhood to support fertility, allow examination, and reduce risks related to twisting, injury, and cancer.

## Undescended Testis

See *Cryptorchidism*.

## Undifferentiated

Not showing the recognizable features of a specialized cell or tissue. An undifferentiated tumor has cells that look very unlike normal tissue, so its origin and behavior may be difficult to determine.

## Undifferentiated Arthritis

Inflammatory arthritis with joint symptoms and signs that do not yet fit one specific named arthritis. The pattern may become clearer over time or remain an undifferentiated condition.

## Undifferentiated Cancer

Cancer made of cells that look very unlike normal cells and do not clearly show the tissue where they began. Special laboratory tests and imaging may be needed to identify its type and guide treatment.

## Undifferentiated Carcinoma

A high-grade carcinoma whose cells lack recogniz-

able features of a specific tissue or tumor subtype; diagnosis and treatment require pathology, immunostaining, and staging.

## Undifferentiated Connective Tissue Disease

An autoimmune connective-tissue disease with symptoms or laboratory findings suggesting systemic inflammation but not enough features for a specific diagnosis such as lupus or systemic sclerosis.

## Undifferentiated Connective-Tissue Disease

An autoimmune connective-tissue disorder with symptoms, examination findings, or laboratory abnormalities suggesting systemic autoimmunity but not meeting criteria for a defined disease such as lupus or systemic sclerosis. Features and long-term course vary.

## Undifferentiated Pleomorphic Sarcoma

An uncommon, often fast-growing cancer of soft tissue or bone whose cells vary greatly in shape and do not resemble a specific normal tissue. Treatment depends on its site, size, and spread.

### Alternate Names:

UPS; Malignant Fibrous Histiocytoma; MFH

## Undifferentiated Pleomorphic Sarcoma Of The Breast

A rare high-grade breast sarcoma formerly called malignant fibrous histiocytoma. It is a diagnosis of exclusion after other specific sarcomas and metastatic carcinoma have been ruled out histologically.

## Undifferentiated Spondyloarthritis

Inflammatory arthritis with features of the spondyloarthritis group but insufficient or atypical findings for classification as ankylosing spondylitis, psoriatic arthritis, reactive arthritis, enteropathic arthritis, or another defined subtype. Symptoms may involve the spine, sacroiliac joints, peripheral joints, entheses, skin, eyes, or bowel.

## Undulant Fever

An infection caused by *Brucella* bacteria, usually



---

acquired from infected animals or unpasteurized dairy products. It can cause repeated or fluctuating fever, sweats, tiredness, joint pain, and serious organ complications.

### **Unerupted Tooth**

A tooth that has not emerged into the oral cavity at the expected time. It may be normally developing, delayed, impacted, or prevented from erupting by obstruction or abnormal position.

### **Ungual**

Relating to a fingernail or toenail. The term may describe a finding beneath, around, or within the nail, such as a bruise, fungal infection, wart, or pigmented lesion.

### **Unicameral Bone Cyst**

A noncancerous, fluid-filled space in a bone, usually in the upper arm or thigh of a child or teenager. It may be found after a weakened bone breaks. An imaging sign called a fallen-leaf sign can support the diagnosis, but it is not specific by itself.

#### **Alternate Names:**

Simple Bone Cyst; Solitary Bone Cyst

### **Unicornuate Uterus**

A congenital condition in which only one side of the uterus develops fully, usually with one fallopian tube. Some people have no symptoms, but it can increase risks of infertility, pregnancy outside the uterus, miscarriage, or premature birth and may require specialist pregnancy care.

### **Unilateral**

Affecting, occurring on, or involving one side of the body or one paired organ, such as unilateral weakness or unilateral eye disease.

### **Unilateral Gynecomastia**

Enlargement of breast glandular tissue affecting one breast in a male patient. A new, firm, fixed, painful, or nipple-associated mass can have causes other than gynecomastia.

### **Unilateral Microphthalmos**

A congenital condition in which one eye is abnormally small. It may be associated with other ocular malformations and can cause substantial visual impairment.

### **Unilateral Spatial Neglect**

A common neurobehavioral syndrome following acute brain injury (most frequently a right-sided stroke involving the right inferior parietal lobule or temporoparietal junction) in which the patient fails to attend to, respond to, or orient toward sensory stimuli presented on the contralesional (typically left) hemispace, despite intact primary visual and sensory pathways.

#### **Alternate Names:**

Hemispatial Neglect; Left Visual Neglect; Spatial Inattention

### **Uninodular Goiter**

An enlarged thyroid containing one main lump, or nodule. Blood tests and ultrasound help assess thyroid function and the chance of cancer; some nodules need follow-up or a needle sample.

### **Unintentional Weight Loss**

Unintentional weight loss is reduction in body weight without planned dietary or activity change and may result from malignancy, endocrine disease, infection, gastrointestinal disease, medication, or mood disorder.

### **Uniparental Disomy**

A chromosome difference in which both copies of a chromosome, or part of one, come from the same parent. It can cause disease by disrupting genes that depend on parental origin or by giving a child two copies of a harmful gene variant.

### **Universal Donor**

An individual with blood group O Rh-negative (O-neg), whose red blood cells lack A, B, and Rh(D) surface antigens, making their packed red blood cells safe for emergency transfusion into recipients of any ABO and Rh blood group without immediate hemolytic transfusion reactions.



---

**Alternate Names:**

Type O Negative Donor; Universal Erythrocyte Donor

**Universal Recipient**

An individual with blood group AB Rh-positive (AB-pos), whose plasma lacks anti-A, anti-B, and anti-Rh(D) isohemagglutinins, allowing them to safely receive packed red blood cells of any ABO/Rh blood group.

**Alternate Names:**

Type AB Positive Recipient; Universal Blood Recipient

**Unloader Knee Braces**

Specialized orthotic knee braces designed to redistribute mechanical weight and load away from an osteoarthritic or damaged compartment of the knee joint (such as the medial or lateral tibiofemoral compartment) to the healthier compartment, reducing pain and improving mobility.

**Alternate Names:**

Unloader Knee Bracing; Offloader Braces; Osteoarthritis Knee Brace

**Unopposed Estrogen**

Estrogen treatment given without a progestogen to someone with a uterus. Long-term use can overstimulate the uterine lining and increase cancer risk, so protection is usually recommended when appropriate.

**Unresectable**

Describing a tumor or lesion that cannot be completely removed safely by surgery because of its location, extent, invasion of vital structures, or the patient's operative risk.

**Unresponsive Coma**

A state of profound unconsciousness in which a person cannot be awakened or follow commands. Causes include brain injury, stroke, seizures, poisoning, infection, and metabolic failure. It is a medical emergency.

**Unsaturated Fat**

A dietary fat whose fatty-acid chains contain one or more carbon-carbon double bonds. Monounsaturated and polyunsaturated fats are common in vegetable oils, nuts, seeds, and fish; replacing saturated fat with unsaturated fat generally supports healthier LDL cholesterol levels.

**Unstable Angina**

Chest pain caused by reduced blood flow to the heart that is new, worsening, lasts longer, or occurs at rest, without clear evidence of heart-muscle death. It is an emergency warning of possible heart attack.

**Unsteadiness**

Impaired balance or gait stability, which may result from vestibular disease, neurologic disorders, weakness, medication effects, visual impairment, or musculoskeletal problems.

**Untidy Wound**

An irregular or contaminated wound with crushing, devitalized tissue, tissue loss, foreign material, or associated injury to vessels, nerves, tendons, or bone. It has increased infection risk and may require exploration, irrigation, debridement, and delayed or staged closure.

**Upper Airway Biopsy**

Removal of tissue from the nasal cavity, pharynx, larynx, or related upper airway for microscopic evaluation of suspected infection, inflammation, infiltrative disease, or tumor.

**Upper Airway Cough Syndrome**

A common clinical etiology of chronic cough provoked by post-nasal drip and mechanical or chemical stimulation of upper airway sensory cough receptors, secondary to allergic rhinitis, non-allergic rhinitis, or chronic rhinosinusitis.

**Alternate Names:**

UACS; Post-Nasal Drip Syndrome

**Upper Airway Obstruction**

Narrowing or blockage of the breathing passage in

---

the nose, throat, or voice box. It can cause noisy breathing, breathlessness, voice changes, or blue lips and can become life-threatening when severe or rapidly worsening.

### **Upper Airway Resistance Syndrome**

Repeated extra effort to breathe through a partly narrowed upper airway during sleep. It can cause brief repeated awakenings, unrefreshing sleep, morning headache, and daytime sleepiness even without clearly positive standard sleep-apnea testing.

**Alternate Names:**

UARS

### **Upper Arm**

The portion of the upper limb between the shoulder and elbow, containing the humerus and muscles, nerves, and vessels that serve the arm.

### **Upper Endoscopy**

A flexible camera examination of the food pipe, stomach, and first part of the small intestine. It can identify inflammation, ulcers, bleeding, narrowing, or tumors and can allow tissue samples or treatment.

### **Upper Esophageal Sphincter**

A ring of muscle at the top of the esophagus that opens during swallowing and coordinates food passage.

**Alternate Names:**

UES

### **Upper Extremity**

The anatomic region extending from the shoulder through the arm, forearm, wrist, and hand. Disorders may involve its bones, joints, muscles, nerves, vessels, or skin.

### **Upper Extremity Deep Vein Thrombosis**

A thrombosis occurring in the deep venous system of the arm or thoracic inlet (axillary, subclavian, or brachiocephalic veins), categorized as primary (effort-induced venous thrombosis or Paget-Schroetter syndrome) or secondary (most commonly related to indwelling central venous catheters or pacemaker

leads).

**Alternate Names:**

UEDVT

### **Upper Extremity Venous Doppler**

An ultrasound examination of the arm and related veins that assesses compressibility and blood flow, commonly to evaluate suspected venous thrombosis.

### **Upper Gastrointestinal Bleeding**

Bleeding from the esophagus, stomach, or first part of the small intestine. It may cause vomiting of red or coffee-ground blood, black stools, dizziness, weakness, or fainting. Common causes include ulcers, inflammation, tears, and enlarged veins; significant bleeding can be life-threatening.

### **Upper GI And Small Bowel Series**

A fluoroscopic x-ray examination in which swallowed contrast outlines the esophagus, stomach, duodenum, and small intestine to evaluate structure and transit.

### **Upper GI Series**

An X-ray examination of the food pipe, stomach, and first part of the small intestine after swallowing contrast material. It can show narrowing, blockage, ulcers, abnormal movement, or other structural problems.

### **Upper Leg**

The portion of the lower limb between the hip and knee, containing the femur and the thigh muscles, nerves, and blood vessels.

### **Upper Limb Disorders**

Conditions affecting the shoulder, arm, elbow, forearm, wrist, or hand, including musculoskeletal, nerve, vascular, and work-related disorders.

### **Upper Motor Neuron**

A nerve cell in the brain or spinal cord that controls nerve cells supplying muscles. Damage to these pathways can cause weakness, stiffness, exaggerated reflexes, or difficulty controlling movement.

---

## Upper Motor Neuron Lesion

Damage or interruption of the corticospinal (pyramidal) or corticobulbar motor pathways within the cerebral cortex, internal capsule, brainstem, or spinal cord, characterized clinically by muscle spasticity ('clasp-knife' hypertonia), hyperreflexia, clonus, extensor plantar response (positive Babinski sign), and pronator drift without significant muscle atrophy.

### Alternate Names:

Pyramidal Tract Lesion; UMN Syndrome

## Upper Motor Neuron Syndrome

A group of findings caused by damage to movement pathways in the brain or spinal cord. It may cause weakness, stiffness, muscle spasms, exaggerated reflexes, or reduced control of movement.

## Upper Respiratory Infection

An infection of the nose, sinuses, pharynx, or larynx, most often caused by a virus and producing symptoms such as congestion, sore throat, cough, or runny nose.

## Upper Respiratory Tract Infection

An acute, typically self-limiting viral or bacterial infection of the mucosal structures of the upper aerodigestive tract above the larynx, encompassing the common cold, rhinitis, pharyngitis, tonsillitis, and acute rhinosinusitis.

### Alternate Names:

URTI; Upper Respiratory Infection; URI

## Upper-Tract Urothelial Carcinoma

A cancer arising from the urothelial lining of the renal pelvis or ureter. It may cause blood in the urine, flank pain, or blockage and is distinct from urothelial cancer arising in the bladder.

## Upright Labor Positions

Maternal positions during labor, such as walking, standing, squatting, kneeling, or sitting upright. They may improve comfort and mobility and can be used when safe and acceptable to the laboring person and clinical team.

## Upward Gaze

Elevation of both eyes to direct the visual axis upward. It depends on coordinated activity of the extraocular muscles and their brainstem control pathways.

## Urachal Abnormalities

Birth-related problems caused by incomplete closure of the urachus, the temporary fetal connection between the bladder and navel. They include a urachal cyst, sinus, or open connection and may cause infection, navel discharge, abdominal pain, or urine leakage.

## Urachal Carcinoma

A rare cancer arising from a leftover connection between the bladder and the navel, usually near the top of the bladder. It may cause blood or mucus in the urine, pelvic pain, or urinary changes.

## Urachal Cyst

A fluid-filled sac left from the temporary fetal connection between the bladder and navel. It may cause no symptoms, but infection can produce lower-abdominal pain, fever, navel discharge, or urinary symptoms.

## Urachal Fistula

A congenital anomaly resulting from complete failure of obliteration of the embryonic urachus (allantoic canal), creating a persistent patent fibrous lumen connecting the bladder dome directly to the umbilicus, presenting as continuous or intermittent urine drainage from the umbilicus.

### Alternate Names:

Patent Urachus; Persistent Urachal Canal

## Urachal Mucinous Cystic Tumor

A rare tumor arising from the urachal remnant between the bladder dome and umbilicus. It may be benign or malignant, with mucinous histology, and requires surgical excision.

## Urachus

An embryologic connection between the fetal blad-

---

der and umbilicus that normally becomes the median umbilical ligament after birth. Persistent urachal remnants can form cysts, sinuses, or other complications.

### **Uracil-DNA Glycosylase**

A DNA repair enzyme that initiates the base excision repair (BER) pathway by specifically recognizing and hydrolyzing the N-glycosidic bond of mutagenic uracil residues formed by spontaneous deamination of cytosine in DNA, generating an abasic (AP) site for downstream repair.

**Alternate Names:**

UDG

### **Uraemia**

A serious illness caused by severe kidney failure and buildup of waste products in the blood. It may cause nausea, itching, tiredness, confusion, bleeding, or inflammation around the heart.

### **Uraemic Foetor**

A distinctive, fishy, ammoniacal, or urine-like breath odor caused by the breakdown of elevated salivary urea into ammonia by oral bacterial urease in patients with advanced chronic kidney disease or acute renal failure.

**Alternate Names:**

Uremic Fetor; Ammoniacal Breath

### **Uraemic Syndrome**

The collection of symptoms and biochemical abnormalities caused by severe accumulation of uraemic waste products in advanced kidney failure.

### **Uranium**

A naturally occurring radioactive heavy metal. Internal exposure can damage the kidneys, while inhaled or retained radioactive material can also expose tissues to ionizing radiation.

### **Urate**

A form of uric acid, a waste product made when the body breaks down certain substances in food and cells. Too much urate in the blood can form

crystals in joints and cause gout.

### **Urate-Lowering Therapy**

Long-term treatment that lowers urate in the blood, helping dissolve gout crystals and prevent further attacks and joint damage. The medicine is chosen according to kidney function, other illnesses, and treatment goals.

### **Urate Nephropathy**

Acute kidney injury caused by the intratubular precipitation of needle-shaped monosodium urate and uric acid crystals within the renal collecting ducts and distal tubules, typically occurring in the setting of acute tumor lysis syndrome following cytotoxic chemotherapy for high-turnover hematological malignancies. Marked by an elevated urinary uric acid-to-creatinine ratio greater than 1.0.

**Alternate Names:**

Acute Uric Acid Nephropathy; Crystal-Induced Urate Nephropathy

### **Urates**

Salts or ions of uric acid. Monosodium urate crystals can accumulate in joints in gout, and urate crystals may contribute to urinary stones.

### **Urea**

The principal nitrogen-containing waste product of protein metabolism, formed in the liver and excreted mainly by the kidneys.

### **Urea Breath Test**

A noninvasive test for active *Helicobacter pylori* infection: the person drinks labeled urea, and labeled carbon dioxide in exhaled breath indicates bacterial urease activity.

### **Urea Cycle**

A series of chemical reactions in the liver that changes toxic ammonia, produced when the body breaks down protein, into urea. The kidneys then remove urea in urine.

---

## Urea Cycle Disorder

An inherited problem in the liver's process for removing ammonia. Ammonia can build up in the blood and cause vomiting, poor feeding, sleepiness, confusion, seizures, or brain injury, especially during illness or fasting.

## Urea Nitrogen Urine Test

A urine test measuring urea nitrogen excretion, interpreted with blood urea nitrogen, creatinine, diet, and urine volume to assess protein metabolism and renal handling.

## Ureaplasma

A genus of very small bacteria without a cell wall that can colonize the genital or urinary tract and is sometimes associated with urethritis or pregnancy complications.

## Ureaplasma Urealyticum

A cell-wall-free bacterium that can colonize the genital or urinary tract and is sometimes associated with nongonococcal urethritis, pregnancy complications, or infection in vulnerable newborns.

## Uremia

A clinical syndrome caused by advanced kidney failure and accumulation of waste products and other biologically active substances. Symptoms may include nausea, itching, fatigue, confusion, bleeding, or pericarditis; laboratory values and symptoms must be interpreted together.

## Uremic Bleeding

A qualitative hemorrhagic diathesis occurring in patients with severe acute kidney injury or end-stage renal disease, resulting from uremic toxin-induced platelet dysfunction (impaired GPIIb/IIIa activation and defective von Willebrand factor interaction), characteristically treated with desmopressin (DDAVP), cryoprecipitate, or dialysis.

## Uremic Encephalopathy

Brain dysfunction caused by the buildup of waste products in severe kidney failure. It may cause confusion, extreme sleepiness, muscle twitching,

seizures, or coma. It is a medical emergency and is often treated by addressing the kidney failure, sometimes with dialysis.

## Uremic Fetor

A characteristic fishy, ammoniacal, or urine-like odor on the breath of patients with severe end-stage renal disease (ESRD), produced by the bacterial breakdown of elevated concentrations of salivary urea into ammonia by oral urease-producing microflora.

### Alternate Names:

Ammoniacal Breath; Uremic Breath Odor

## Uremic Frost

Rare white, powdery crystals of urea that appear on the skin when severe untreated kidney failure causes very high waste levels in the blood. It indicates advanced kidney failure.

## Uremic Pericarditis

Inflammation of the sac around the heart caused by severe kidney failure and buildup of waste products. It may cause sharp chest pain, a scratchy heart sound, or fluid around the heart; fluid can sometimes compress the heart and impair its pumping.

## Uremic Platelet Dysfunction

Reduced ability of platelets to stick together and stop bleeding because of advanced kidney failure. It can cause easy bruising, nosebleeds, heavy menstrual bleeding, or bleeding from the stomach or bowel and may improve when the kidney failure is treated.

## Uremic Pruritus

A refractory, distressing cutaneous itch affecting patients with advanced chronic kidney disease and end-stage renal disease on hemodialysis or peritoneal dialysis, related to immune dysregulation, microinflammation, peripheral neuropathy, and opioid receptor imbalance.

### Alternate Names:

Chronic Kidney Disease-Associated Pruritus; CKD-aP

---

### Uremic Stomatitis

Inflammation of the mouth caused by severe kidney failure and buildup of waste products. It may cause a bad taste, mouth soreness, white patches, ulcers, or bleeding gums and tongue.

### Ureter

One of two muscular tubes that carry urine from a kidney to the bladder. A stone or narrowing can block a ureter and cause severe side pain, swelling of the kidney, infection, or kidney damage.

### Ureteral Calculi

Stones that form in or pass through a ureter, the tube carrying urine from a kidney to the bladder.

### Ureteral Catheters

Thin tubes passed into a ureter to drain urine, inject contrast, measure pressure, or guide a procedure. Possible complications include infection, bleeding, a tear, movement of the tube, or blockage.

### Ureteral Injury During Gynecologic Surgery

Damage to a ureter during pelvic surgery, including transection, ligation, thermal injury, devascularization, or obstruction. It may cause flank pain, urine leakage, hydronephrosis, fever, or delayed kidney injury and requires prompt imaging and urologic management.

### Ureteral Obstruction

A blockage that prevents urine from flowing from a kidney to the bladder. It may cause side pain, infection, kidney swelling, or reduced kidney function. Common causes include stones, scar tissue, tumors, and pressure from nearby tissues.

### Ureteral Orifice

One of two slit-like mucosal openings situated at the posterolateral angles of the trigone of the urinary bladder through which urine from the ureters enters the bladder lumen, guarded by an oblique intramural muscular tunnel that acts as a functional physiological valve to prevent vesicoureteral reflux.

### Ureteral Perforation

A tear or hole in the ureter, sometimes caused by a stone, trauma, surgery, or an endoscopic procedure. Urine may leak into surrounding tissues, causing pain, infection, or inflammation; prompt imaging and urologic assessment are required.

### Ureteral Polyp

Usually a benign fibroepithelial growth arising from the ureter. It may cause hematuria, flank pain, or obstruction and must be distinguished from urothelial malignancy when evaluated.

### Ureteral Reimplantation

Surgery that moves the lower end of a ureter to a new opening in the bladder. It may be used to correct urine flowing backward, blockage, injury, or an abnormal ureter entry.

#### Alternate Names:

Ureteroneocystostomy; Ureteral Neoimplantation

### Ureteral Retrograde Brush Biopsy

A small brush passed through a thin camera into the ureter to collect cells from its lining. A laboratory examines the cells for cancer or other abnormal changes.

### Ureteral Stenosis

Narrowing of the ureter caused by scar tissue, prior surgery, radiation, stones, inflammation, or congenital disease, impairing urine drainage and possibly causing hydronephrosis or recurrent infection.

### Ureteral Stent

A thin tube placed inside the ureter to keep urine flowing from the kidney to the bladder when a stone, narrowing, tumor, or swelling blocks the passage. It can cause urinary urgency, discomfort, blood in the urine, infection, or mineral buildup and is usually temporary.

### Ureteral Stricture

Narrowing of the ureter caused by fibrosis, surgery, instrumentation, radiation, infection, malignancy, or congenital disease. It can produce upstream



---

hydronephrosis, pain, infection, and declining renal function.

### **Ureter Duplex**

A birth variation in which one kidney has two urine tubes instead of one. It may cause no problems but can lead to urine flowing backward, blockage, repeated infections, or poor drainage.

### **Ureter Fissus**

A birth difference in which a ureter is divided lengthwise or partly duplicated. It may cause no symptoms, but abnormal urine flow can lead to urine reflux, blockage, repeated infection, or poor kidney drainage.

### **Ureteric Bud**

An embryonic epithelial outgrowth arising from the posteromedial wall of the mesonephric (Wolffian) duct that invades the metanephric blastema, branching repeatedly to form the ureter, renal pelvis, major and minor calyces, and collecting duct system.

#### **Alternate Names:**

Metanephric Diverticulum

### **Ureteric Calculi**

Kidney stones that have moved into a ureter, the tube carrying urine from a kidney to the bladder. They can cause severe wave-like side pain, blood in the urine, nausea, blockage, or infection.

### **Ureteric Colic**

Severe, paroxysmal, agonizing flank and abdominal pain radiating anteriorly and inferiorly toward the ipsilateral groin, testicle, or labium majus, caused by acute smooth muscle spasm and proximal hydronephrotic distension from an obstructing urinary calculus (nephrolithiasis) in the ureter.

#### **Alternate Names:**

Renal Colic; Ureteral Calculus Pain

### **Ureteric Perforation**

A tear or hole in the ureter that can allow urine to leak into surrounding tissue, usually after instrumentation, surgery, trauma, or a stone; it needs

prompt urologic assessment.

### **Ureteritis**

Inflammation of the ureter, usually related to infection, obstruction, stone passage, instrumentation, or adjacent inflammatory disease. It may cause flank or abdominal pain, urinary symptoms, hematuria, and fever.

### **Ureter Obstruction**

Blockage of urine flow from a kidney to the bladder, commonly caused by a stone, stricture, tumor, or external compression and potentially leading to hydronephrosis, infection, or kidney injury.

### **Ureterocele**

A balloon-like dilation of the lower ureter where it enters the bladder, usually congenital; it can obstruct urine flow and cause infection, hydronephrosis, reflux, stones, or impaired renal function.

### **Ureterolithiasis**

A kidney stone located in the ureter, the tube from the kidney to the bladder. It can block urine and cause severe wave-like side pain, blood in the urine, nausea, infection, or kidney damage.

### **Ureteropelvic Junction Obstruction**

A blockage where the kidney's urine-collecting area joins the ureter. It may be present from birth or develop later and can cause kidney swelling, side pain, infections, stones, or loss of kidney function.

### **Ureteroscope**

A thin camera instrument passed through the urinary passage to examine the ureter and kidney. It can help remove stones, take tissue samples, or place a stent.

### **Ureteroscopes**

Small endoscopes used to inspect the ureter and renal collecting system and to perform procedures such as stone removal, biopsy, or stent placement.

---

## Ureteroscopy

Ureteroscopy uses a small endoscope passed through the urinary tract to inspect, biopsy, or treat ureteral or kidney stones and other lesions.

## Ureterostomy Procedure

Surgery that creates an opening from a ureter to the skin or another urinary conduit to divert urine when normal drainage to the bladder is impossible or must be bypassed.

## Urethra

The tube that carries urine from the bladder out of the body. In people with male reproductive anatomy it also carries semen; the shorter urethra in people with female reproductive anatomy makes bladder infections more likely.

## Urethra Disorder

A condition affecting the urethra, such as infection, inflammation, stricture, trauma, congenital abnormality, or tumor, that may cause pain, discharge, weak stream, or retention.

## Urethral Cancer

Cancer arising from the urethra, usually from its lining cells. It may cause blood in the urine, a lump, discharge, pain, or difficulty passing urine.

## Urethral Carcinoma

A rare cancer arising from the urethra that may cause bleeding, a lump, discharge, pain, or difficult urination; diagnosis requires examination, imaging, and biopsy.

## Urethral Caruncle

A benign, fleshy, vascular, exophytic polypoid outgrowth of the posterior margin of the external urethral meatus occurring almost exclusively in postmenopausal women due to chronic hypoestrogenism and local urethral prolapse, presenting with spotting, dysuria, or a tender reddish meatal nodule.

### Alternate Names:

Benign Meatal Vascular Polyp; Estrogen-Deficient Urethral Lesion

## Urethral Catheter

A tube inserted through the urethra into the bladder to drain urine, obtain a specimen, instill medication, or monitor output. Urinary infection and urethral trauma are important risks.

## Urethral Diverticulum

A pocket beside the urethra that connects with the urine tube, often developing after infection or blockage of nearby glands. It may cause burning urination, urine dribbling after voiding, repeated infections, pain during sex, or a vaginal-wall lump.

## Urethral Fistula

An abnormal channel between the urethra and another epithelial surface, often the vagina or skin. It can cause continuous urine leakage, recurrent infection, and local irritation.

## Urethral Hypermobility

Excessive movement of the urethra when abdominal pressure rises, contributing to urine leakage during coughing, lifting, or exercise.

## Urethral Infection

Infection involving the urethra, commonly caused by bacteria, including sexually transmitted organisms. Symptoms may include burning urination, discharge, frequency, or pelvic discomfort; testing and appropriate treatment help prevent complications and transmission.

## Urethral Injury

Damage to the urethra caused by trauma, a pelvic fracture, a penetrating wound, or a medical instrument. It may cause blood at the urethral opening, difficulty passing urine, bruising, or urine leakage into nearby tissues.

## Urethral Meatus

The external opening of the urethra through which urine leaves the body. Its position may be abnormal in conditions such as hypospadias or epispadias.

---

## Urethral Obstruction

Partial or complete blockage of urine leaving the bladder, often from stricture, stone, prostate disease, tumor, or a congenital valve. Acute retention can damage the bladder or kidneys.

## Urethral Pressure Profile

A test using a small catheter to measure pressure along the urethra and assess how well it closes. It may help investigate certain types of urine leakage or difficulty emptying the bladder.

## Urethral Prolapse

A circumferential eversion and protrusion of the distal urethral mucosa through the external urethral meatus, presenting as a reddish, doughnut-shaped or rosette-like congestion at the vulva, typically occurring in prepubertal Black girls or frail postmenopausal women.

### Alternate Names:

Urethral Mucosal Prolapse; Meatal Eversion

## Urethral Sling

A strip of synthetic material or the person's own tissue placed beneath the urethra to support it and reduce urine leakage during coughing, lifting, or exercise. Possible complications include urgency, difficulty urinating, infection, or erosion.

## Urethral Sphincter

Muscles around the urine tube that help keep urine from leaking and relax when a person urinates. Nerve or muscle damage can weaken control and cause leakage or difficulty emptying the bladder.

## Urethral Stenosis

Scar-related narrowing of the urethral channel that restricts urinary flow and can cause spraying, weak stream, recurrent infection, retention, or bladder damage.

## Urethral Stent

A small tube placed inside the urethra to help keep a narrowed or blocked segment open and allow urine to pass. Migration, blockage, infection, pain, or

tissue injury can occur.

## Urethral Stricture

A narrowing of the urethra, the tube that carries urine out of the body, usually caused by scar tissue. The narrowing can slow or block urine flow and may cause a weak or spraying stream, straining, pain with urination, dribbling, or incomplete bladder emptying. It can lead to repeated urinary infections, urinary retention, and, if severe or prolonged, damage to the bladder or kidneys. Causes include injury, infection, surgery, radiation, or previous use of a catheter or other instrument; sometimes no cause is found.

## Urethral Syndrome

Urinary frequency, urgency, dysuria, or discomfort without a clear bacterial urinary-tract infection; causes may include irritation, pelvic-floor dysfunction, or inflammation.

## Urethral Trauma

Injury to the urethra caused by a pelvic fracture, blunt or penetrating trauma, or medical instrumentation. It may cause bleeding, difficulty passing urine, bruising, or leakage of urine into nearby tissues.

## Urethral Discharge

Fluid or pus coming from the urethral opening. It may result from urethritis, including sexually transmitted infection, but can also follow irritation or injury and warrants appropriate medical assessment and testing.

## Urethritis

Inflammation of the urethra, often caused by infection, including sexually transmitted infections, but sometimes by irritation or injury. It can cause burning urination, itching, pelvic discomfort, or discharge.

## Urethrocele

Bulging or downward displacement of the urethra, usually into the front wall of the vagina. It may cause pressure, urinary leakage, difficulty emptying

---

the bladder, discomfort, or no symptoms.

### Urethrography

An X-ray test in which contrast liquid outlines the urethra, the tube carrying urine out of the body. It can show narrowing, leaks, injury, abnormal connections, or birth differences.

### Urethroplasty

Surgery that rebuilds or widens the urine tube, most often to treat a narrowing called a stricture. It may use nearby tissue or a graft and aims to improve urine flow.

### Urethroscopy

Examination of the urethra with a thin camera passed through its opening. It can identify narrowing, injury, inflammation, stones, tumors, or other abnormalities and may allow treatment during the same procedure.

### Urge Incontinence

Involuntary leakage of urine preceded immediately by a sudden, intense, compelling desire to void that is difficult to defer, resulting from involuntary detrusor muscle contractions (detrusor overactivity / overactive bladder) during the bladder filling phase.

#### Alternate Names:

Detrusor Overactivity Incontinence; Urgency Urinary Incontinence

### Urgency

A sudden, difficult-to-delay need to urinate. It may result from a urinary infection, bladder irritation, overactive bladder, a stone, nerve disease, prostate enlargement, or pelvic-floor problems.

### Urge Urinary Incontinence

Involuntary leakage of urine that follows a sudden, difficult-to-delay need to urinate. It may occur with overactive bladder, urinary infection, bladder irritation, nerve disease, or other urinary-tract problems.

### Uric Acid

A waste product formed when the body breaks down

purines. High blood levels can cause gout or kidney stones, although many people with high levels have no symptoms.

### Uric Acid Nephrolithiasis

The formation of pure uric acid urinary stones resulting from chronic acidic urine (urine pH less than 5.5), hyperuricosuria, and low urine volume, which appear radiolucent on plain abdominal radiographs but are readily visualized as hyperdense calculi on non-contrast CT.

#### Alternate Names:

Uric Acid Calculi; Radiolucent Kidney Stones

### Uric Acid Nephropathy

Kidney injury caused when too much uric acid forms crystals in the kidney's small tubes. It can occur during rapid breakdown of cancer cells or severe uric-acid excess.

### Uric Acid Stones

Kidney stones composed primarily of uric acid. They form more readily in acidic urine and are associated with hyperuricemia, gout, high purine intake, or chronic diarrhea. Some can dissolve when urine is alkalinized.

### Uric Acid Urine Test

A urine test measuring uric acid excretion, used in evaluation of gout, kidney stones, uric-acid overproduction, and selected metabolic disorders.

### Uricosuric Drug

A medicine that increases renal excretion of uric acid, such as probenecid, used in selected patients with gout.

### Uridine Triacetate

An oral prodrug that supplies uridine and is used as an emergency antidote for severe or life-threatening toxicity from fluorouracil or related fluoropyrimidines, under specialist direction.

### Urinal

A portable container used to collect urine from

---

someone who cannot easily reach a toilet. It should be emptied, cleaned, and stored safely to reduce spills, odor, and infection risk.

## Urinalysis

Laboratory examination of urine for its appearance, acidity, blood, cells, protein, sugar, ketones, bacteria, and other substances. It can provide clues to infection, kidney disease, bleeding, diabetes, dehydration, and other illness.

## Urinary

Relating to urine or to the organs and processes involved in making, storing, and passing urine, including the kidneys, ureters, bladder, and urethra.

## Urinary Anion Gap

A calculation using urine sodium, potassium, and chloride levels. It can help doctors assess how the kidneys are handling acid during some types of blood acidosis. The result is only one part of the assessment and cannot by itself identify the cause.

### Alternate Names:

UAG; Urine Net Charge

## Urinary Bladder

A hollow muscular organ in the lower abdomen that stores urine until it is released through the urethra.

## Urinary Bladder Disorder

A disease affecting bladder storage or emptying, including infection, overactivity, pain syndrome, stones, tumor, neurologic dysfunction, or outlet obstruction.

## Urinary Bladder Squamous Cell Carcinoma Pathology

An invasive malignant squamous neoplasm of the bladder showing keratinization, intercellular bridges, or both. It may be associated with chronic irritation, infection, stones, or schistosomiasis and is assessed for depth of invasion.

## Urinary Calculi

Stones formed from crystallized minerals in the urinary tract. They may cause colicky pain, hematuria, obstruction, infection, or kidney injury depending on their size and location.

## Urinary Casts

Small tube-shaped particles formed inside kidney tubules and passed in urine. Their appearance can provide clues about kidney inflammation, bleeding, infection, or damage.

## Urinary Catheter

A hollow, flexible tube inserted through the urethra into the bladder to drain urine; it may be used intermittently or left in place.

## Urinary Catheterization

Insertion of a tube into the bladder to drain urine, measure output, obtain a specimen, or deliver treatment; infection, trauma, and blockage are possible complications.

## Urinary Catheters

Urinary catheters drain the bladder when a person cannot empty it normally or accurate urine measurement is needed, with risks including infection, trauma, blockage, and discomfort.

## Urinary Diversion

Surgical or catheter-based rerouting of urine when the normal passage from the kidneys to the urethra cannot be used, often to a stoma or bowel reservoir.

## Urinary Fistula

An abnormal connection between part of the urinary tract and another organ or the skin, allowing urine to leak outside its normal pathway. Causes include surgery, childbirth injury, inflammation, and cancer.

## Urinary Frequency

The need to urinate more often than usual, which may result from infection, bladder irritation, increased urine production, or other urinary disorders.

---

## Urinary Hesitation

Difficulty starting or maintaining urination, which may result from prostate enlargement, urethral narrowing, medications, nerve disorders, pain, or impaired bladder contraction.

## Urinary Incontinence

Unwanted leakage of urine. It may happen with coughing or exercise, a sudden strong urge, an overfull bladder, difficulty reaching the toilet, or nerve and muscle problems. More than one pattern may occur in the same person.

### Alternate Names:

Involuntary Urination; Loss of Bladder Control; Enuresis

## Urinary Lithiasis

The formation or presence of stones anywhere in the urinary tract, including the kidneys, ureters, bladder, or urethra.

## Urinary Retention

Inability to empty the bladder fully. It may be sudden and painful or develop gradually with a weak stream, frequent small amounts of urine, or leakage from an overfull bladder. Blockage, medicines, nerve problems, or weak bladder muscles can cause it.

## Urinary Retraining

Scheduled toilet visits and bladder-control techniques used to reduce urgency, frequent urination, or leakage. A clinician may combine it with pelvic-floor exercises, fluid changes, or treatment of the underlying cause.

## Urinary Stasis

Slowing or stopping of normal urine flow, which can promote urinary infection, stone formation, or obstruction-related kidney injury.

## Urinary System

The kidneys, ureters, bladder, and urethra. Together they filter blood, make and transport urine, store and remove it, and help regulate fluid volume, electrolytes, acid-base balance, blood pressure, and

hormone-related functions such as red-cell production.

## Urinary Tract

The kidneys, ureters, bladder, and urethra. Together these organs make urine, carry it to the bladder, store it, and remove it from the body.

## Urinary Tract Disorder

Any disease of the kidneys, ureters, bladder, or urethra, including infection, stones, obstruction, congenital abnormalities, cancer, and functional or neurologic disorders.

## Urinary Tract Imaging

Imaging tests used to examine the kidneys, ureters, bladder, and urethra. Methods include ultrasound, X-ray studies, computed tomography, magnetic resonance imaging, and nuclear scans, chosen according to the suspected problem.

## Urinary Tract Infection

Infection of the urinary tract, classified as lower infection such as cystitis or urethritis, or upper infection such as pyelonephritis. Symptoms vary; dysuria, frequency, urgency, fever, flank pain, or nausea may occur. *Escherichia coli* is a common cause, but organisms vary by setting and patient factors.

## Urinary Tract Infection In Adults

A urinary infection in an adult involving the bladder, urethra, kidneys, or another part of the urinary tract. Symptoms vary by site and may include painful urination, frequency, urgency, blood in the urine, fever, or flank pain.

## Urinary Tract Infection In Children

A urinary infection in a child involving the bladder, kidneys, ureters, or urethra. Symptoms may include fever, painful or frequent urination, new wetting, abdominal or back pain, vomiting, or unusual irritability.

### Alternate Names:

Pediatric UTI; Childhood Cystitis



---

## Urinary Tract Infection In Pregnancy

A urinary infection or asymptomatic bacterial growth occurring during pregnancy. Pregnancy increases the risk of upper urinary infection, so screening and treatment follow pregnancy-specific clinical guidance.

## Urinary Tract Infections in Pregnancy

Bacterial infection of the urethra, bladder, or kidneys during pregnancy. Even asymptomatic bacteriuria can progress to pyelonephritis, preterm birth, or sepsis, so screening, culture-directed pregnancy-safe treatment, and follow-up are important.

## Urinary Tract Neoplasm

A benign or malignant growth in the kidneys, ureters, bladder, or urethra; evaluation depends on its site, symptoms, tissue type, and extent of spread.

## Urinary Tract Obstruction

An anatomical or functional impediment to the normal antegrade flow of urine along the urinary tract from the renal calyces to the urethral meatus, predisposing to hydronephrosis, postrenal acute kidney injury, urinary stasis, and urosepsis.

## Urinary Urgency

A sudden, difficult-to-delay need to urinate. It may occur with urinary infection, bladder irritation, overactive bladder, stones, neurologic disease, or other conditions. Urgency with fever, flank pain, blood in the urine, or inability to urinate requires medical assessment.

## Urination

The process of releasing urine from the bladder through the urethra and out of the body.

### Alternate Names:

Micturition; Voiding

## Urination Disorder

Abnormal storage or emptying of urine causing frequency, urgency, dysuria, incontinence, hesitancy, weak stream, retention, or excessive or reduced urine

output.

## Urine

The liquid waste produced by the kidneys, containing water and dissolved substances eliminated from the blood.

## Urine 24-Hour Volume

Twenty-four-hour urine volume is the total urine produced in a day and helps assess hydration, kidney function, endocrine disease, or abnormal urine production.

## Urine Albuminuria

An abnormal amount of the blood protein albumin in urine. Persistent albumin loss can be an early sign of kidney disease and also increases the risk of worsening kidney function and cardiovascular disease.

## Urine Alkalinization

Treatment that makes urine less acidic, usually with a carefully monitored medicine, to help remove certain poisons such as salicylate. It can disturb body salts or fluid balance and is used only for selected cases.

## Urine Amylase Test

A urine test that measures the amount of amylase passed from the body. It may help assess pancreatitis, salivary-gland problems, or unusually high amylase levels in the blood.

### Alternate Names:

Urine Amylase; Urinary Amylase Test; Amylase Urine Test

## Urine Analysis

Laboratory testing of urine for blood, cells, protein, sugar, microbes, and other clues to illness. It can help assess urinary infection, kidney disease, diabetes, dehydration, and other conditions.

## Urine Bilirubin Test

A clinical urinalysis that detects conjugated (water-soluble) bilirubin in urine. A positive result indi-

---

cates hepatobiliary disease, hepatocellular injury, or biliary tract obstruction; unconjugated bilirubin is bound to albumin and does not filter into urine.

**Alternate Names:**

Urine Bilirubin; Urinary Bilirubin Test

**Urine Calcium Test**

A diagnostic urinalysis quantifying urinary calcium excretion, typically measured in a 24-hour urine collection. It is utilized in the evaluation and metabolic workup of nephrolithiasis, hyperparathyroidism, osteomalacia, and idiopathic hypercalciuria.

**Alternate Names:**

Urinary Calcium; 24-Hour Urine Calcium; Urine Calcium Excretion

**Urine Catecholamine Test**

A diagnostic biochemical assay measuring 24-hour urinary excretion of epinephrine, norepinephrine, dopamine, and their metanephrine metabolites to diagnose neuroendocrine tumors such as pheochromocytoma, paraganglioma, and neuroblastoma.

**Alternate Names:**

24-Hour Urine Catecholamines; Urinary Catecholamines; Fractionated Urinary Catecholamines

**Urine Chemistry**

Measurement of chemical constituents in urine, such as glucose, protein, electrolytes, ketones, blood, and pH, to evaluate kidney function and systemic disease.

**Urine Chloride Test**

A diagnostic urinalysis quantifying urinary chloride excretion, crucial in differentiating chloride-responsive (e.g., vomiting, diuretic use) from chloride-resistant (e.g., hyperaldosteronism) metabolic alkalosis.

**Alternate Names:**

Urinary Chloride; Urine Chloride Excretion; Spot Urine Chloride

**Urine Concentration Test**

A test of the kidney's ability to concentrate urine, assessed by urine osmolality or specific gravity after

controlled fluid conditions when appropriate.

**Urine Conductivity**

Urine conductivity measures how readily urine carries electrical current and reflects its dissolved electrolyte and solute concentration.

**Urine Culture**

A laboratory test that grows microorganisms from a urine specimen to identify urinary-tract infection and guide antimicrobial selection. Results may report the organism, quantity of growth, and antimicrobial susceptibility. Proper collection is important because contamination can produce misleading growth; results must be interpreted with symptoms and urinalysis.

**Urine Cytology**

Microscopic examination of cells shed into urine, mainly used to help detect high-grade urothelial carcinoma and some other abnormalities. A negative result does not exclude all urinary-tract cancers.

**Urine Dipstick**

A rapid, point-of-care biochemical test consisting of a plastic strip with reagent pads that undergo colorimetric changes upon chemical reaction with urine, screening for leukocytes, nitrite, urobilinogen, protein, pH, blood, specific gravity, ketones, bilirubin, and glucose.

**Urine Drainage Bags**

Urine drainage bags collect urine from a catheter or urinary diversion and require correct positioning, emptying, hygiene, and monitoring for blockage or infection.

**Urine Electrolytes Test**

A diagnostic urinalysis measuring the concentrations of sodium, potassium, and chloride in urine to evaluate acute kidney injury, dehydration, electrolyte imbalances, and acid-base disturbances.

**Alternate Names:**

Urinary Electrolytes; Spot Urine Electrolytes; 24-Hour Urine Electrolytes

---

## Urine Formation

The process by which the kidneys make urine: they filter blood, return needed water and substances to the body, and add selected wastes and extra ions to the fluid that leaves as urine.

## Urine HCG Test

A urine test that detects human chorionic gonadotropin, a hormone produced during pregnancy. Testing too early or using diluted urine can give a false-negative result; a positive result confirms pregnancy but does not show its location or health.

### Alternate Names:

Urine hCG Test; Gravidex Test

## Urine Hemoglobin

Hemoglobin detected in urine, which may reflect intravascular hemolysis or contamination with intact red blood cells. Interpretation requires comparison with urine microscopy and clinical findings.

## Urine Immunoelectrophoresis

A specialized clinical laboratory test that separates urinary proteins by agarose gel electrophoresis followed by immunodiffusion with monospecific antibodies. Primarily utilized to identify and characterize monoclonal light chains (Bence Jones proteins) in plasma cell dyscrasias such as multiple myeloma.

### Alternate Names:

Urine IEP; Urinary Protein Immunoelectrophoresis; Urine Immunofixation Electrophoresis

## Urine Immunofixation

An immunochemical laboratory technique combining high-resolution electrophoresis with specific antibody precipitation to identify and type monoclonal free light chains (kappa or lambda) and intact immunoglobulins excreted in urine.

### Alternate Names:

Urine IFE; Urinary Immunofixation Electrophoresis; Bence Jones Protein IFE

## Urine Osmolality

A measure of how concentrated urine is. It helps assess whether the kidneys can conserve or remove

water and must be interpreted with fluid intake, medicines, kidney function, and blood tests.

## Urine PH

A measurement of urine acidity or alkalinity. It varies with diet, metabolism, medications, infection, and kidney function and can influence the formation of some urinary stones.

## Urine PH Test

A test measuring urine acidity or alkalinity, useful in evaluating acid-base handling, urinary infection, kidney stones, and some metabolic conditions.

## Urine Protein Dipstick Test

A screening test using a reagent strip to detect mainly albumin in urine; positive results require confirmation and quantitative assessment when clinically indicated.

## Urine Protein Electrophoresis Test

A laboratory test that separates urine proteins by electrical charge and size to help detect abnormal proteins, including monoclonal light chains associated with plasma-cell disorders.

## Urine Retention

Alternate name; see *Urinary Retention*.

## Urine Sediment

The cells, crystals, casts, and other particles seen under a microscope after urine is concentrated. Their appearance can provide clues to bleeding, infection, stones, or kidney inflammation.

## Urine Sodium

The amount of sodium in urine. It can help assess fluid balance and kidney handling of salt, but results are strongly affected by diet, medicines, hydration, and the timing of the sample.

## Urine Specific Gravity Test

A test comparing the density of urine with the density of water. It gives a rough estimate of how concentrated the urine is and can help assess hydra-

---

tion and the kidneys' ability to conserve water.

### **Urine Toxicology**

Laboratory analysis of urine for medicines, drugs, or their metabolites. Results require interpretation because detection does not always establish impairment or the timing of exposure.

### **Urinoma**

An encapsulated or non-encapsulated extravasation of urine into the retroperitoneal space or perirenal tissues, resulting from traumatic rupture or iatrogenic instrumentation injury to the renal pelvis, calyces, or ureter.

#### **Alternate Names:**

Urine Pseudocyst

### **Urinometer**

An instrument that measures how dense urine is compared with water. The result gives an estimate of urine concentration and may help assess hydration or the kidney's ability to conserve water.

### **URI (Upper Respiratory Infection)**

URI usually means upper respiratory infection, commonly a viral infection of the nose, throat, or upper airways causing congestion, cough, and sore throat.

### **Urobilinogen**

A colourless product formed in the intestine from bilirubin and partly reabsorbed and excreted in urine; abnormal levels may provide clues to liver disease or haemolysis.

### **Urodynamics**

A group of tests that measures how the bladder stores urine, how pressure changes as it fills, and how the urethra and bladder release urine. It helps investigate leakage, urgency, weak stream, or difficulty emptying when simpler tests are inconclusive.

### **Urodynamic Testing**

A group of tests that measures how the bladder and urethra store and release urine. It can help explain leakage, urgency, difficulty emptying, or repeated

retention when simpler tests do not identify the cause.

### **Uroflow**

A test that measures how quickly and steadily urine leaves the body during urination. The result can suggest blockage, weak bladder muscles, or other emptying problems, but it is interpreted with symptoms and sometimes additional tests.

### **Uroflowmetry**

A noninvasive test measuring how fast and steadily urine leaves the body. A low or abnormal flow may reflect blockage, weak bladder contraction, or an incomplete test and must be interpreted with symptoms and other findings.

### **Urogenital**

Relating to the urinary and reproductive systems or structures involving both. The term may describe organs, infections, cancers, birth conditions, injuries, symptoms, tests, or treatments affecting these closely connected body systems.

### **Urogenital Fistula**

An abnormal connection between the urinary and genital tracts, such as a vesicovaginal or ureterovaginal fistula. It commonly causes continuous urinary leakage and requires specialist evaluation.

### **Urogenital Sinus**

A birth-related abnormality in which the urethra and vagina join into a common channel before reaching the outside of the body. It may occur with differences in the external genitalia, urinary tract, or reproductive organs.

### **Urogenital System**

The related urinary and reproductive organs and their associated ducts, vessels, and supporting tissues. The systems share developmental origins and anatomic relationships.

### **Urogenital Tract**

The combined urinary and reproductive organs and

---

passages. These include kidneys, urine tubes, bladder, urethra, reproductive organs, and connected structures that develop partly from related tissues before birth.

### **Urography**

Imaging of the urinary tract, traditionally using X-rays after contrast dye is given. It can show the kidneys, urine tubes, bladder, narrowing, stones, leaks, blockage, or other structural problems.

### **Urokinase**

A plasminogen activator produced by the body and also used as a thrombolytic medicine to help dissolve blood clots.

### **Urolithiasis**

Formation of stones in the urinary tract, including the kidneys, ureters, bladder, or urethra. Stone composition varies by population and may include calcium oxalate, calcium phosphate, uric acid, struvite, or cystine.

### **Urologic Diagnostic Yield**

The proportion of urinary-system tests or procedures that provide a clear, useful diagnosis. A test may have a low yield when the sample is too small, the disease is difficult to detect, or the result does not change care.

### **Urologist**

A doctor who specializes in the urinary system and, in many practices, the male reproductive system. Urologists diagnose and treat kidney stones, infections, urine leakage, prostate problems, cancers, infertility, and conditions requiring procedures or surgery.

### **Urology**

The medical specialty that diagnoses and treats diseases of the kidneys, urine tubes, bladder, and urethra, as well as the male reproductive organs. Treatment may include medicines, procedures, or surgery.

### **Uropathy**

Disease or damage affecting the kidneys or urinary tract. Blockage, infection, stones, urine flowing backward, medicines, and other conditions can cause it and may reduce kidney function.

### **Urosepsis**

A life-threatening whole-body response to a severe urinary infection. It may start in the bladder or kidney and cause fever or low temperature, confusion, low blood pressure, reduced urine, organ failure, or shock.

### **Urostomy Leak**

Escape of urine beneath or around a urostomy appliance or from a surgical connection. Persistent leakage can irritate the skin and may signal poor appliance fit, obstruction, infection, or a surgical complication.

### **Urostomy Procedure**

Surgery that diverts urine through a stoma into an external pouch or conduit, commonly after bladder removal or when the lower urinary tract cannot safely transport urine.

### **Urostomy Prolapse**

Protrusion of a segment of bowel used as a urostomy stoma beyond its usual level. It may cause bleeding, swelling, appliance problems, or impaired urine drainage and should be assessed clinically.

### **Urostomy Stenosis**

Narrowing of a urinary stoma or conduit that impedes drainage, causing reduced urine output, leakage, infection, or kidney-pressure changes and sometimes requiring revision.

### **Urothelial Carcinoma**

A cancer arising from the stretchable lining of the urinary tract, most often the bladder but also the kidney's drainage area, ureters, or urethra. Blood in the urine is common, and treatment depends on location, depth, grade, and spread.

---

## Upper Tract Urothelial Carcinoma

Tumors arising from the urothelial lining of the upper urinary tract. Blood in the urine, flank pain, obstruction, or incidental imaging findings may occur; urine cytology, imaging, ureteroscopy, and tissue assessment guide diagnosis and treatment.

## Urothelium

A specialized stretchable lining of the renal pelvis, ureters, bladder, and part of the urethra. It forms a protective barrier between urine and underlying tissues and allows the bladder to expand.

## Urotrauma

Injury to the urinary or genital organs, including the kidneys, ureters, bladder, urethra, or external genitalia. It may result from blunt force, penetrating wounds, fractures, surgery, or medical instruments.

## Ursodeoxycholic Acid

A hydrophilic bile acid used for primary biliary cholangitis and selected cholestatic liver diseases; it may also dissolve certain small cholesterol gallstones when bile flow is preserved.

## Urticaria

A skin reaction causing transient, itchy raised welts that change location and usually resolve within 24 hours. It may occur with angioedema, which is deeper swelling often affecting the lips, eyelids, tongue, or extremities.

## Urticarial Vasculitis

Inflammation of small blood vessels causing hive-like patches that usually last longer than 24 hours and may burn, bruise, or leave dark marks. It can be limited to the skin or linked with joint, kidney, lung, or other disease.

## Urticaria Multiforme

A short-lived allergic-type rash in children that causes widespread, ring-shaped or target-like raised patches, often with swelling of the hands, feet, or face. It may follow an infection or medicine and usually fades without scarring.

## Urticaria Pigmentosa

A skin condition in which too many mast cells collect in the skin, causing tan or brown spots or bumps that swell and itch when rubbed. Flushing or other body-wide symptoms can occur.

## Usher Syndrome

An inherited condition causing hearing loss together with gradually worsening vision from retinal disease. The age and severity vary by type; hearing, vision, balance, communication, and genetic support may all be needed.

## Ustekinumab

A biologic medicine that blocks immune signals involved in inflammation. It is used for psoriasis, psoriatic arthritis, Crohn disease, ulcerative colitis, and other approved conditions; infections and allergic reactions are important risks.

## Usual Interstitial Pneumonia

A pattern of permanent scarring in the lungs, often seen in idiopathic pulmonary fibrosis. Scarring is usually greatest near the outer and lower parts of the lungs and can form small spaces called honeycombing. It causes increasing breathlessness and cough and is assessed with lung tests and CT imaging.

## Usual Interstitial Pneumonia Pathology

A patchy, temporally heterogeneous interstitial fibrosing pattern with subpleural or paraseptal fibrosis, fibroblastic foci, architectural distortion, and honeycomb change. Clinical and radiologic correlation is required.

## Ut Dict.

An abbreviation of the Latin phrase “ut dictum,” meaning “as stated” or “as directed.” In medical notes it refers the reader to information already given or to an instruction previously specified.

## Uterine Adnexae

The structures beside and connected to the uterus, mainly the fallopian tubes, ovaries, and supporting tissues. Problems in these structures can cause pelvic pain, abnormal bleeding, infertility, or an



---

ectopic pregnancy.

### **Uterine Artery Embolization**

A minimally invasive procedure that blocks blood flow to uterine fibroids so they shrink. It can reduce heavy bleeding and pelvic pressure, but may cause pain, infection, or effects on fertility and future pregnancy.

### **Uterine Atony**

Failure of the uterus to contract effectively after childbirth, causing heavy postpartum bleeding. It is the most common cause of postpartum hemorrhage and may be managed with medicines, uterine massage, procedures, or surgery, depending on severity and response.

#### **Alternate Names:**

Postpartum Uterine Atony

### **Uterine Balloon Tamponade**

A treatment for severe bleeding after childbirth when medicines do not control bleeding. A soft balloon is placed inside the uterus and filled with sterile fluid to press against the uterine wall and slow or stop blood loss while other treatment is arranged.

### **Uterine Bleeding**

Bleeding from the uterus outside the expected menstrual pattern or in an abnormal amount. Causes include pregnancy complications, fibroids, polyps, hormonal changes, medicines, infection, and cancer; heavy or postmenopausal bleeding can have an underlying medical cause.

### **Uterine Cancer**

Cancer arising in the uterus, most commonly from the endometrium. Abnormal vaginal bleeding, bleeding after menopause, pelvic pain, or unusual discharge requires evaluation; early diagnosis often permits effective treatment.

### **Uterine Cavity**

The hollow space inside the uterus, lined by the endometrium. A fertilized egg normally implants in this lining, which thickens during the menstrual

cycle and supports pregnancy; polyps, fibroids, scar tissue, or abnormal shape may affect it.

### **Uterine Compression Suture**

A surgical hemostatic technique (exemplified by the B-Lynch, Hayman, or Cho suture) in which heavy absorbable sutures are placed through the uterine wall to mechanically compress the anterior and posterior walls of an atonic uterus, controlling refractory postpartum hemorrhage while preserving future fertility.

#### **Alternate Names:**

B-Lynch Suture; Uterine Bracing Suture

### **Uterine Contraction**

Tightening of the uterine muscle. Contractions normally occur during menstruation and labor, but painful or unusually frequent contractions can occur with pregnancy complications, dehydration, infection, or medicines and may require assessment.

### **Uterine Diseases**

Conditions affecting the womb, including fibroids, adenomyosis, endometriosis, infection, unusual bleeding, prolapse, and cancer. Symptoms may include pain, heavy periods, infertility, pressure, discharge, or bleeding after menopause.

### **Uterine Fibroid**

A benign tumor arising from the smooth muscle of the uterus. Fibroids may be submucosal, intramural, or subserosal and may cause heavy menstrual bleeding, pelvic pressure, pain, urinary symptoms, or reproductive problems.

#### **Alternate Names:**

Uterine Myoma; Fibromyoma

### **Uterine Fibroid Embolization**

A minimally invasive procedure in which particles block blood flow to uterine fibroids, causing them to shrink. It can reduce heavy bleeding and pressure symptoms but may cause pain, infection, or effects on fertility.

---

## Uterine Fibroids

Benign tumors of uterine smooth muscle, also called leiomyomas. They may cause heavy menstrual bleeding, pelvic pressure, pain, infertility, or no symptoms.

### Alternate Names:

Uterine Leiomyomata; Uterine Myomas; Fibromyomas

## Uterine Fistula

A uterine fistula is an abnormal passage connecting the uterus with another organ or the skin, often after surgery, childbirth injury, infection, or cancer.

## Uterine Fornix

One of the shallow spaces at the top of the vagina surrounding the cervix. A clinician can examine these spaces during a pelvic examination, and selected procedures may use them as an access route.

## Uterine Gland

A small gland in the lining of the uterus that produces fluid and helps support the early embryo. Its activity changes during the menstrual cycle and pregnancy.

## Uterine Growths

Uterine growths may include fibroids, polyps, adenomyosis, or rarely cancer. They can cause heavy or irregular bleeding, pelvic pressure, pain, infertility, or no symptoms; examination and imaging determine the type and treatment.

## Uterine Hemorrhage

Heavy or abnormal bleeding from the uterus, occurring during pregnancy, after childbirth, or at other times. Causes include pregnancy complications, fibroids, hormonal disorders, and malignancy; severe bleeding with dizziness or weakness is an emergency.

## Uterine Inversion

A rare, life-threatening complication of childbirth in which the uterus turns partly or completely inside out and may protrude into or beyond the vagina. It can cause severe bleeding, shock, and intense pain.

Treatment restores the uterus and controls blood loss.

## Uterine Leiomyoma

A benign tumor of uterine smooth muscle that may cause heavy menstrual bleeding, pelvic pressure, pain, or no symptoms.

### Alternate Names:

Fibroid

## Uterine Leiomyosarcoma

A rare malignant tumor of uterine smooth muscle that may cause abnormal bleeding, pelvic pain, or an enlarging mass; diagnosis is based on tissue examination and staging.

## Uterine Monitoring

Recording how often, how long, and how strongly the uterus contracts. During labor it helps assess progress, response to induction medicines, and excessive contractions that could affect the baby.

## Uterine Myomectomy

Surgical removal of uterine fibroids while preserving the uterus; it may be performed through hysteroscopy, laparoscopy, or open surgery depending on fibroid size and location.

## Uterine Obstruction

Blockage that prevents normal passage through the uterus or cervix, which may cause menstrual outflow obstruction, infertility, pregnancy complications, or retained blood or tissue.

## Uterine Perforation

A hole through the wall of the uterus, most often caused by an instrument used during a procedure such as hysteroscopy or removal of uterine tissue. It can cause internal bleeding or injury to nearby organs and may need repair.

## Uterine Polyps

Growths arising from the inner lining of the uterus that may cause irregular bleeding, heavy periods, or fertility problems.

---

**Alternate Names:**

Endometrial Polyps; Intrauterine Polyps

**Uterine Prolapse**

Descent of the uterus toward or through the vaginal opening because pelvic-floor muscles and supporting tissues have weakened. It may cause pelvic pressure, a bulge, backache, urinary or bowel problems, or discomfort during sex; childbirth and aging increase risk.

**Alternate Names:**

Descensus Uteri; Pelvic Organ Prolapse; Uterine Herniation

**Uterine Rupture**

A complete tear through the wall of the uterus, usually during pregnancy or labor. It can cause severe internal bleeding and reduce the baby's oxygen supply; previous cesarean surgery is an important risk factor.

**Uterine Rupture in Pregnancy**

A tear through the uterine wall, most often during labor in a scarred uterus but sometimes in an unscarred uterus. Sudden pain, bleeding, fetal heart-rate abnormality, loss of station, or maternal shock may occur; emergency surgery is required.

**Uterine Sarcoma**

Uterine sarcoma is a rare malignant tumor arising from uterine muscle, connective tissue, or related cells and may cause bleeding, pain, or an enlarging mass.

**Uterine Tachysystole**

More than five contractions in 10 minutes, averaged over 30 minutes during labor. Very frequent contractions can reduce the baby's oxygen supply, especially after labor medicines, and may require prompt treatment adjustment.

**Uterine Tube**

A paired muscular tube that transports the oocyte from the ovary toward the uterus and is the usual site of fertilization.

**Uterine Tumors**

Alternate name; see *Uterine fibroids*.

**Utero-Ovarian Ligament**

A band of tissue connecting each ovary to the uterus. It helps hold the ovary in position within the pelvis and lies near the fallopian tube and other pelvic supporting tissues.

**Alternate Names:**

Ovarian Ligament; Proper Ovarian Ligament

**Uteroplacental Insufficiency**

Inadequate maternal blood flow and nutrient delivery across the intervillous spaces of the placenta to the developing fetus, resulting from defective spiral artery remodeling, maternal vascular disease, or placental infarction; the primary cause of fetal growth restriction and intrapartum late decelerations.

**Alternate Names:**

Placental Insufficiency

**Uterosacral Ligament**

One of two strong bands of tissue supporting the uterus and top of the vagina by attaching them toward the sacrum at the back of the pelvis. Weakness can contribute to pelvic-organ prolapse; endometriosis can also affect or cause pain in this area.

**Uterus**

A hollow muscular organ in the pelvis between the bladder and rectum. Its inner lining thickens to prepare for pregnancy and is shed during menstruation. A fertilized egg normally implants in this lining, and a fetus develops in the uterus during pregnancy.

**Common name:** Womb

**Uterus Didelphys**

A congenital condition in which two separate uteruses develop, often with two cervixes and a dividing wall in the vagina. Many people can become pregnant, but miscarriage, premature birth, abnormal fetal position, or other pregnancy complications are more common.

**Alternate Names:**

Double Uterus; Didelphic Uterus

---

## Uterus Disorder

A condition affecting the uterus, including fibroids, adenomyosis, endometriosis, infection, abnormal bleeding, congenital anomalies, prolapse, and cancer.

## Uterus Neoplasm

A benign or malignant growth arising in the uterus, including fibroids, endometrial cancer, and sarcomas; diagnosis depends on examination, imaging, and tissue sampling.

## UTI

Alternate name; see *Urinary Tract Infection*.

## Utricle

A small balance organ in the inner ear that senses head position and movement in a straight line, such as forward, backward, or side-to-side motion.

## Uvea

The middle layer of the eye, made up of the iris, ciliary body, and choroid. It controls the amount of light entering the eye, helps focus vision, and supplies blood to eye tissues.

## Uveal Disease

A disorder of the iris, ciliary body, or choroid, including uveitis, tumors, vascular disease, and inflammation that may cause eye pain, photophobia, floaters, or visual loss.

## Uveal Melanoma

The most common primary cancer inside the adult eye. It begins in pigment-producing cells of the iris, ciliary body, or choroid and may cause visual changes or be found during an eye examination; it can spread to other organs.

## Uveal Neoplasm

A tumor arising in the uveal tract of the eye—the iris, ciliary body, or choroid; malignant melanoma may threaten vision and spread to distant organs.

## Uveitic Glaucoma

Secondary open-angle or closed-angle glaucoma caused by active intraocular inflammation (anterior uveitis); elevated intraocular pressure results from inflammatory cell and debris trabecular meshwork obstruction (trabeculitis), posterior synechiae causing pupillary block (iris bombé), or topical corticosteroid therapy.

### Alternate Names:

Inflammatory Glaucoma; Secondary Uveitic Ocular Hypertension

## Uveitis

Inflammation of the uveal tract—the iris, ciliary body, and choroid—and sometimes nearby eye tissues. It may result from infection, autoimmune or other inflammatory disease, trauma, or an unknown cause. Symptoms can include eye pain, redness, light sensitivity, floaters, blurred vision, or reduced vision; prompt eye assessment is important because untreated inflammation can damage vision.

### Alternate Names:

Intraocular Inflammation; Iritis; Cyclitis

## Uveoparotid Fever

A form of sarcoidosis causing fever, swelling of both parotid glands near the ears, and inflammation in the front of the eyes. It may also cause facial-nerve weakness; treatment depends on the organs affected and severity.

## UV Light

Ultraviolet radiation, an invisible part of sunlight or artificial light that can affect the skin and eyes and may cause sunburn, aging, or skin cancer.

## Uvula

A small, soft flap hanging from the back of the roof of the mouth. It helps with swallowing and speech; swelling can cause gagging, pain, snoring, or difficulty breathing.

## Uvulitis

Inflammation and swelling of the uvula, the small tissue hanging at the back of the throat. Infection, allergy, injury, dehydration, or irritation may

---

cause it; severe swelling or breathing difficulty can threaten the airway.

### **Uvulopalatopharyngoplasty**

An operation that removes or reshapes parts of the soft palate, the hanging uvula, and the throat to widen the breathing passage. It may help selected people with obstructive sleep apnea, but can cause pain, swallowing changes, bleeding, or recurrence.

### **U Wave**

A small wave that may appear after the T wave on an electrocardiogram. A prominent U wave can occur with low potassium, a slow heart rate, or some medicines and must be interpreted with the complete tracing and clinical findings.

### **Underactive Thyroid**

Alternate name; see *Hypothyroidism*.

### **Urinary Tract Infection (UTI)**

Alternate name; see *Urinary Tract Infection*.

### **Ureteral Stones**

Stones lodged in a ureter, the tube carrying urine from a kidney to the bladder. They may cause severe wave-like flank or groin pain, blood in the urine, nausea, blockage, or infection and sometimes need urgent treatment.

### **Uterine Polyp**

A growth arising from the inner lining of the uterus. Many are benign, but they can cause irregular or heavy bleeding, bleeding after menopause, or infertility and may require imaging, sampling, or removal.

### **Universal Precautions**

Infection-control practices that treat blood and certain body fluids as potentially infectious, regardless of a person's diagnosed condition. They include hand hygiene, appropriate gloves and protective barriers, safe sharps handling, and measures to prevent exposure to bloodborne pathogens.

### **Urostomy**

A surgically created opening that diverts urine from the urinary tract through the abdominal wall into an external pouch or appliance. It may be formed with a segment of intestine after bladder removal or when normal urinary drainage is not possible.

### **Ultrasound Scan**

Alternate name; see *Ultrasound*.

### **Ulcerative Colitis (UC)**

Alternate name; see *Ulcerative Colitis*.





# V

## **Vaccination**

Giving a vaccine to train the immune system to recognize a specific infection. Vaccination lowers the risk of illness, severe complications, and sometimes transmission; the recommended schedule depends on age, health, and local guidance.

## **Vaccination Schedule**

A time-based plan for administering vaccines according to age, previous doses, medical conditions, exposure risks, and public-health guidance. Schedules vary by country and are updated as recommendations change.

## **Vaccinations/Immunizations During Pregnancy**

Vaccines given during pregnancy are selected according to safety, disease risk, season, exposure, and local recommendations. Inactivated vaccines such as influenza and Tdap are commonly used when indicated, while some live vaccines are deferred until after pregnancy.

## **Vaccine**

A vaccine trains the immune system to recognize a specific infection or toxin and respond more effectively. Vaccination lowers the risk of severe disease and helps protect communities.

## **Vaccine Allergy**

An allergic reaction to a vaccine or one of its ingredients. Reactions may be local and mild or, rarely, involve hives, swelling, breathing difficulty, or anaphylaxis; suspected reactions are assessed to identify the responsible ingredient and guide future

vaccination.

## **Vaccine-Associated Paralytic Poliomyelitis**

An extremely rare form of polio paralysis caused by a weakened poliovirus used in the oral polio vaccine changing back to a harmful form.

### **Alternate Names:**

VAPP

## **Vaccines**

Preparations that train the immune system to recognize a particular germ or toxin before exposure. They reduce the risk of infection and severe illness and may also reduce complications and spread.

## **Vaccines and Immunizations**

Vaccines train the immune system to recognize specific pathogens or toxins and reduce the risk of infection, severe disease, complications, and transmission.

## **Vaccinia**

A virus used to make smallpox vaccines. Rare accidental infection can cause a sore at the contact site or more widespread illness, especially in people with weakened immunity.

## **Vaccinia Virus**

Vaccinia virus is a poxvirus used to make smallpox vaccines and studied in some experimental cancer treatments; accidental infection is uncommon but possible.

---

## **VACTERL Association**

A pattern of birth defects affecting several of the spine, anus, heart, food pipe, windpipe, kidneys, or limbs. There is no single test; the pattern of differences is used to identify it and rule out other conditions.

## **Vacuum-Assisted Delivery**

A vaginal birth assisted by a suction cup placed on the baby's scalp when labor is not progressing or the baby needs to be born promptly. It is used only when the cervix is fully open and the baby's position and head size are suitable. Possible complications include scalp injury, bleeding, or tears.

## **Vacuum Curettage**

A procedure that removes tissue from the uterus using suction, commonly for uterine evacuation or endometrial sampling. Risks include bleeding, infection, retained tissue, and rarely uterine injury.

## **Vacuum Delivery**

Assisted vaginal birth using a suction cup placed on the baby's scalp. It may be used when labor is not progressing or the baby needs to be born promptly and can cause scalp injury, bleeding, or tears.

## **Vacuum Erection Devices**

Devices with a cylinder and pump that draw blood into the penis to produce an erection. A flexible ring may hold the erection during sex; bruising, numbness, or discomfort can occur, especially if the ring is left on too long.

### **Alternate Names:**

Vacuum Erection Pump; Penis Pump

## **Vacuum Extractor**

A cup-shaped device applied to the fetal scalp to assist vaginal delivery when specific maternal or fetal indications are present.

## **Vagal Maneuver**

A physical action that affects the vagus nerve and can slow the heart's electrical activity. Examples include bearing down and, in clinical settings, cer-

tain pressure or cold-stimulation methods. These maneuvers may help assess or stop some fast heart rhythms.

### **Alternate Names:**

Vagal Stimulation Technique; Parasympathetic Provocative Maneuver

## **Vagal Nerve Stimulation**

A treatment that uses an implanted device to send small electrical signals to the vagus nerve in the neck. It may be used for selected cases of epilepsy or depression that have not improved with other treatments.

### **Alternate Names:**

VNS

## **Vagal Paraganglioma**

A usually slow-growing tumor arising from nerve-related cells near the vagus nerve in the neck. It may cause a neck mass, hoarseness, swallowing difficulty, or nerve symptoms and can be identified with imaging and specialist assessment.

## **Vagina**

A muscular, elastic canal connecting the cervix to the vaginal opening. It allows menstrual blood to leave the body, receives a penis or other objects during vaginal sex, and serves as the birth canal.

## **Vagina Carcinoma**

Cancer arising from the vagina. It may cause unusual bleeding, discharge, pain, or a visible sore, although early cancer may cause no symptoms; examination and tissue testing are needed for diagnosis.

## **Vaginal Agenesis**

A condition in which the vagina is absent or has not fully formed before birth. It may occur with differences in the uterus, kidneys, hearing, or spine; care is individualized and may include counseling, gradual treatment, or surgery when wanted.

## **Vaginal Atresia**

Absence or complete blockage of the vaginal canal

---

from birth. It may cause absent menstrual flow, repeated monthly pelvic pain, or difficulty with penetration.

### **Vaginal Atrophy**

Thinning, dryness, and irritation of vaginal tissues, usually caused by lower estrogen after menopause or other hormone changes. It can cause burning, itching, pain during sex, bleeding, or urinary symptoms and may improve with moisturizers, lubricants, or prescribed hormone treatment.

#### **Alternate Names:**

Atrophic Vaginitis; Vulvovaginal Atrophy; Genitourinary Syndrome of Menopause; GSM; Vaginal Dryness

### **Vaginal Bleeding**

Unexpected, unusually heavy, or prolonged bleeding from the vagina, including bleeding between periods, after sexual intercourse, during pregnancy, or after menopause. Causes include hormonal changes, infection, pregnancy-related conditions, structural lesions, medications, and cancer.

### **Vaginal Bleeding Between Periods**

Bleeding between menstrual periods may result from contraception, infection, pregnancy, polyps, fibroids, hormonal changes, or cancer. Persistent or heavy bleeding can have a serious cause.

### **Vaginal Bleeding During Early Pregnancy**

Bleeding from the vagina during the first part of pregnancy can result from cervical changes, miscarriage, ectopic pregnancy, infection, or other conditions. A small amount may occur without a serious cause, but bleeding with heavy loss, severe or one-sided pain, shoulder pain, dizziness, or fainting can indicate a serious problem.

### **Vaginal Bleeding During Late Pregnancy**

Bleeding from the vagina later in pregnancy may be caused by placenta previa, placental separation, labor, cervical disease, or other complications. Significant bleeding can be a serious pregnancy complication.

### **Vaginal Bleeding In Early Pregnancy**

Alternate name; see *Vaginal Bleeding During Early Pregnancy*.

### **Vaginal Bleeding In Late Pregnancy**

Alternate name; see *Vaginal Bleeding During Late Pregnancy*.

### **Vaginal Bleeding In Pregnancy**

Broad index label for bleeding during pregnancy; see *Vaginal Bleeding During Early Pregnancy* and *Vaginal Bleeding During Late Pregnancy*. Heavy bleeding or bleeding with pain, dizziness, fainting, or reduced fetal movement can indicate a serious pregnancy complication.

### **Vaginal Cancer**

A rare cancer of the vagina, most often arising from its surface lining. It may cause bleeding, discharge, pain, or a lump; treatment depends on its type, size, location, and spread.

#### **Alternate Names:**

Vaginal Carcinoma; Malignant Vaginal Neoplasm

### **Vaginal Candidiasis**

Alternate name; see *Yeast infection (vaginal)*.

### **Vaginal Cuff**

The upper end of the vagina that is closed with stitches after the uterus is removed during a hysterectomy.

### **Vaginal Cuff Dehiscence**

An opening of the stitched upper end of the vagina after hysterectomy. It may cause pelvic pain, bleeding, discharge, or tissue bulging through the opening.

### **Vaginal Cysts**

Vaginal cysts are fluid-filled or tissue-lined sacs in or beside the vaginal wall, commonly benign but sometimes causing pain, pressure, infection, or obstruction.

---

## Vaginal Dilator

A smooth device used to gently stretch or keep open the vaginal canal when it has become narrow or short because of scarring, surgery, radiation, or a difference present from birth.

## Vaginal Discharge

Vaginal discharge may be normal or may indicate infection, inflammation, hormonal change, or another condition. Normal discharge is usually clear to whitish and odourless and varies with the menstrual cycle, pregnancy, sexual arousal, and some contraceptives.

## Vaginal Dryness

Reduced vaginal lubrication, commonly associated with menopause, breastfeeding, hormonal changes, medications, or irritation. It may cause discomfort, itching, painful intercourse, or minor bleeding.

## Vaginal Fistula

An abnormal passage between the vagina and another organ, commonly the bladder or rectum. It may cause continuous urine or stool leakage, infection, skin irritation, and emotional distress.

## Vaginal Foreign Body

An object retained in the vagina, most often a tampon or other inserted material. It may cause discharge, odor, bleeding, pain, or infection.

## Vaginal Fornix

One of the recesses surrounding the cervix where the vagina meets the cervix. The posterior fornix is adjacent to the peritoneal cul-de-sac and is relevant in pelvic examination and some procedures.

## Vaginal Fusion And Duplication

Birth-related differences in which the vaginal canal is partly joined, divided, or duplicated. They may occur with abnormalities of the uterus, cervix, urinary tract, or external genitalia and may cause obstruction, pain, menstrual problems, or difficulty with intercourse.

## Vaginal Health

Vaginal health includes normal vaginal tissues, fluids, microorganisms, and comfort. Persistent odor, itching, pain, bleeding, or unusual discharge can indicate infection or another condition.

## Vaginal Hematoma

A collection of blood within the vaginal wall or surrounding tissues, usually caused by childbirth, trauma, or surgery and potentially producing severe pain, swelling, or blood loss.

## Vaginal Hemorrhage

Heavy or clinically significant bleeding from the vagina. Causes include childbirth, pregnancy complications, trauma, infection, structural disease, hormonal disorders, and gynecologic cancer; severe or pregnancy-related bleeding can be life-threatening.

## Vaginal Hypoplasia

Vaginal hypoplasia is underdevelopment of the vagina, usually present from birth. It may occur with other reproductive or urinary-tract differences and can affect menstruation, sexual function, or fertility.

## Vaginal Hysterectomy

Surgical removal of the uterus through the vagina. It may be used for selected benign uterine conditions and usually avoids an abdominal incision, although suitability depends on anatomy and disease.

## Vaginal Inflammation

Alternate name; see *Vaginitis*.

## Vaginal Intercourse

The insertion of the penis into the vagina for sexual pleasure or reproduction. The most common form of sexual intercourse. Vaginal intercourse can lead to pregnancy if sperm are deposited in the vagina.

## Vaginal Intraepithelial Neoplasia

Abnormal, precancerous cells in the surface lining of the vagina, often related to human papillomavirus. The changes vary from mild to severe and may

---

require monitoring, treatment, or removal.

### **Vaginal Introitus**

The entrance to the vagina. It is part of the external genital area and may be examined when assessing pain, infection, injury, scarring, or pelvic-floor problems.

### **Vaginal Itching**

Itching or irritation of the vulva or vagina caused by candidiasis, bacterial vaginosis, sexually transmitted infections, dermatitis, hormonal changes, or other conditions. It may occur with discharge, odor, soreness, or burning.

### **Vaginal Laceration**

A tear in the vaginal tissue, often caused by childbirth, sexual activity, trauma, or a medical procedure. It may cause pain, swelling, or bleeding.

### **Vaginal Lubrication**

Natural moisture made by the vaginal tissues and nearby glands. It usually increases with sexual arousal and may decrease with hormonal changes, some medicines, illness, or irritation.

### **Vaginal Membrane**

A thin fold of tissue associated with the vagina. The term may refer to the hymen at the vaginal opening or to an abnormal membrane that blocks the vaginal canal.

### **Vaginal Microbiome**

The community of microorganisms living in the vagina, commonly dominated by *Lactobacillus* species that help maintain an acidic environment. Changes in this community may contribute to bacterial vaginosis, candidiasis, or other symptoms.

### **Vaginal Mucosa**

The moist inner lining of the vagina. It contains folds that allow the vaginal canal to stretch and helps protect the tissue.

### **Vaginal Odor**

An unusual or unpleasant odor from vaginal discharge that may result from bacterial vaginosis, infection, retained material, hormonal changes, or other conditions.

### **Vaginal Opening**

The external entrance to the vagina. It is part of the vulvar vestibule and may be affected by infection, inflammation, trauma, scarring, or congenital variation.

### **Vaginal Pessary**

A removable device placed in the vagina to support the vaginal walls or nearby pelvic organs, usually for pelvic-organ prolapse or some forms of urine leakage.

### **Vaginal pH**

A measure of how acidic the vagina is. The vaginal environment is normally mildly acidic, and the pH can change with hormones, semen, blood, medicines, or infection.

### **Vaginal Prolapse**

Downward bulging of a vaginal wall because the tissues supporting the vagina have weakened. The bulge may involve the bladder, rectum, or the top of the vagina after hysterectomy.

### **Vaginal Rugae**

Soft folds in the inner lining of the vagina that allow it to stretch and return toward its usual shape.

### **Vaginal Seeding**

Placing vaginal fluid on a baby born by cesarean section in an attempt to expose the baby to the parent's vaginal microorganisms. Clear health benefits have not been established, and harmful infections may be transmitted.

### **Vaginal Speculum**

An instrument inserted into the vagina and gently opened to separate the vaginal walls and visualize the vagina and cervix. It may be used for pelvic

---

examination, cervical cytology such as a Pap test, swab collection, assessment of bleeding or discharge, and selected procedures. Size and design are chosen according to anatomy, comfort, and the clinical purpose.

### **Vaginal Stenosis**

Narrowing or shortening of the vaginal canal, which may result from surgery, radiation, scarring, inflammation, trauma, or congenital development. It can cause pain, difficult examination, or difficulty with penetration.

### **Vaginal Swab**

A sample of fluid or cells collected from the vagina for laboratory testing, often to look for infection or other changes.

### **Vaginal Trauma**

Injury to the vaginal tissues caused by childbirth, sexual activity, foreign objects, procedures, or blunt or penetrating trauma. It may cause bleeding, pain, swelling, or urinary symptoms; severe bleeding or deep injury can be serious.

### **Vaginal Ultrasound Probe**

A specialized endocavitary ultrasound transducer used through the vagina to produce images of the uterus, cervix, ovaries, nearby pelvic structures, or an early pregnancy. It sends and receives sound waves and does not use ionizing radiation.

#### **Alternate Names:**

Endovaginal Transducer Transvaginal Ultrasound Transducer

### **Vaginal Vault Prolapse**

Descent of the top of the vagina into or through the vaginal opening, usually after removal of the uterus. It may cause pelvic pressure, a bulge, backache, or difficulty emptying the bladder or bowel; pelvic-floor therapy, a support device, or surgery may help.

#### **Alternate Names:**

Post-Hysterectomy Vault Prolapse; Apical Prolapse

### **Vaginal Vestibule**

The space of the vulva between the labia minora that contains the external openings of the urethra and vagina and the ducts of the greater and lesser vestibular glands.

### **Vaginal Wall**

The layers of muscle, connective tissue, and moist lining that form the walls of the vaginal canal.

### **Vaginal Yeast Infection**

Alternate name; see *Vulvovaginal Candidiasis*.

### **Vagina Neoplasm**

An abnormal growth in the vagina that may be benign, precancerous, or cancerous. It can cause bleeding, discharge, pain, or a lump, although early growths may cause no symptoms.

### **Vaginectomy**

Surgical removal of part or all of the vagina, usually to treat selected cancers or severe disease; effects may include changes in sexual function, fertility, urination, and bowel function.

### **Vaginismus**

Alternate name; see *Genito-Pelvic Pain/Penetration Disorder*.

### **Vaginitis**

Inflammation or irritation of the vagina that may cause discharge, itching, burning, odor, pain, or discomfort with urination or sex. Causes include infections, hormones, chemicals, allergies, and tissue dryness.

### **Vaginoplasty**

Vaginoplasty is surgery to create, repair, or reshape the vagina. It may be performed for congenital differences, injury, cancer treatment, pelvic-floor problems, or gender-affirming care.

### **Vaginospasm**

Alternate name; see *Genito-Pelvic Pain/Penetration Disorder*.



---

## Vagotomy

Surgery that cuts selected branches of the vagus nerve to reduce signals that stimulate stomach acid and movement. It was once used for difficult stomach ulcers but is now uncommon because medicines and less invasive treatments are available.

## Vagus Nerve

The tenth cranial nerve, carrying signals between the brain and the heart, lungs, throat, voice box, and digestive tract. It helps regulate heart rate, breathing, swallowing, voice, and digestion.

## Vagus Nerve Stimulation

A treatment that sends regular, mild electrical pulses to the vagus nerve through an implanted device in the chest or, in some settings, a handheld device. It is used for selected epilepsy, treatment-resistant depression, and other conditions; benefits may take time and side effects can include voice change, cough, or throat discomfort.

## Valacyclovir

An antiviral medicine that the body changes into acyclovir to stop herpes viruses from multiplying. It treats cold sores, genital herpes, and shingles; the dose may need adjustment in kidney disease, and excessive levels can affect the nervous system.

## Valganciclovir

An antiviral medicine used mainly to prevent or treat cytomegalovirus infection after transplantation. The body changes it into ganciclovir, which blocks viral multiplication; it can lower blood counts, injure the kidneys, affect fertility, and harm a fetus.

## Valgus

A limb or joint deformity in which the part farther from the body bends outward, away from the body's center line. Examples include a bunion and knock-knees.

## Valgus Stress Test

A physical examination manoeuvre used to assess the structural integrity of medial collateral ligaments, most commonly at the knee or elbow. A

medial-directed force is applied to the joint while stabilizing the limb; excessive joint opening, laxity, or pain indicates a medial collateral ligament sprain or rupture.

## Valinemia

A rare inherited condition in which the body cannot process the amino acid valine normally. High valine levels may cause feeding problems, low muscle tone, developmental delay, or other nervous-system symptoms.

## Valley Fever

The common name for coccidioidomycosis, a fungal infection caused by *Coccidioides* species and acquired by inhaling airborne spores from soil. It usually causes a mild respiratory illness but can disseminate to skin, bone, joints, or meninges, especially in people at higher risk.

## Valproate

An antiseizure medicine also used for selected mood disorders. It can affect the liver, pancreas, blood cells, weight, and fetal development, so age, pregnancy possibility, other illnesses, and medicines must be considered.

### Alternate Names:

Valproic Acid; Divalproex Sodium

## Valproate Toxicity

Overdose or adverse exposure causing CNS depression, vomiting, hyperammonemic encephalopathy, hepatotoxicity, pancreatitis, thrombocytopenia, metabolic acidosis, or cerebral edema. Measure serum valproate, glucose, ammonia, liver tests, and acid-base status; L-carnitine may be considered and severe poisoning may require hemodialysis.

## Valproic Acid

A medicine used for certain seizures, bipolar disorder, and migraine prevention. It can cause liver injury, pancreatitis, birth defects, and developmental harm during pregnancy.

## Valproic Acid Toxicity

Acute overdose or adverse metabolic toxicity of the

---

broad-spectrum antiepileptic agent valproate, characterized clinically by central nervous system depression, high anion gap metabolic acidosis, hyperammonemia (due to urea cycle inhibition) with encephalopathy, and drug-induced microvesicular hepatic necrosis; treated with L-carnitine.

### Valrubicin

A chemotherapy medicine placed directly into the bladder for selected bladder cancers that have not responded adequately to other treatment. It can cause bladder irritation, blood in the urine, infection, and, rarely, effects throughout the body.

### VRE Infection

An infection caused by vancomycin-resistant enterococci, bacteria that commonly live in the intestine but can cause urinary-tract, wound, bloodstream, heart-valve, or other infections. Treatment depends on the infection site and susceptibility testing because usual antibiotics may not work.

### Visceral Pain

Pain arising from internal organs, often felt as pressure, cramping, aching, or poorly localized discomfort rather than sharp surface pain. It may result from stretching, inflammation, reduced blood flow, obstruction, or injury of an organ.

### Valsalva

Straining while trying to breathe out against a closed throat, like bearing down during a bowel movement. It changes pressure in the chest and can affect heart rate and blood pressure, so it is not safe for everyone.

### Valsalva Maneuver

Alternate name; see *Valsalva manoeuvre*.

### Valsalva Retinopathy

Preretinal (subhyaloid or sub-internal limiting membrane) hemorrhage occurring within the macula following a sudden, dramatic rise in intrathoracic or intra-abdominal pressure (such as from forceful coughing, vomiting, or heavy lifting) that transmits back through the valveless venous system into

retinal capillaries.

### Valsalva's Manoeuvre

Forceful straining while trying to breathe out against a closed airway, like bearing down during a bowel movement. It changes pressure in the chest and can affect heart rate and blood pressure; it is used in selected tests but may be unsafe for some people.

#### Alternate Names:

Valsalva Maneuver

### Valsartan

Valsartan is an angiotensin-II receptor blocker used for hypertension, heart failure, and selected cardiovascular indications; hyperkalaemia, renal-function changes, hypotension, and fetal toxicity are important risks.

### Valves

Structures that permit fluid or blood to move predominantly in one direction and prevent backward flow; cardiac valves and venous valves are important examples.

### Valvotomy

Surgical or catheter-based division of narrowed valve tissue to improve blood flow, most often performed for selected stenotic heart valves.

### Valvulae Conniventes

Large, permanent transverse circular folds of the mucous membrane and submucosa that project into the lumen of the small intestine (most prominent and numerous in the jejunum), which do not disappear with intestinal distension and act to triple the mucosal surface area for absorption.

#### Alternate Names:

Plicae Circulares; Kerckring Folds

### Valvular Disease

Alternate term; see *Heart valve disease*.

### Valvular Endocarditis

Alternate name; see *Endocarditis*.

---

## Valvular Heart Disease

A broad spectrum of structural and functional disorders affecting one or more of the four cardiac valves (aortic, mitral, tricuspid, pulmonic), classified hemodynamically as valvular stenosis (narrowing and outflow obstruction) or valvular regurgitation (insufficiency and backward flow).

### Alternate Names:

VHD

## Valvular Insufficiency

Incomplete closure of a heart valve, allowing blood to leak backward when the heart contracts or relaxes. It can affect the mitral, aortic, tricuspid, or pulmonary valve and may eventually strain the heart.

## Valvuloplasty

A procedure used to widen or repair a narrowed or abnormal heart valve, by balloon catheter or open surgery depending on the valve and condition.

## Valvulotomy

A procedure that cuts open a narrowed heart valve to improve blood flow. It may be performed with a catheter or surgery; the valve can leak afterward and narrowing may recur.

## Vancomycin

An antibiotic used for selected serious bacterial infections and, by mouth, for certain infections of the large intestine. It can cause infusion reactions, kidney injury, hearing problems, or low blood counts and requires medical monitoring.

## Vancomycin Flushing Reaction

A reaction that can occur when vancomycin is given too quickly into a vein. It may cause flushing, warmth, itching, and redness of the face, neck, and upper body.

### Alternate Names:

Red Man Syndrome; Red Neck Syndrome

## Vancomycin-Resistant Enterococcus

Enterococcus bacteria that are resistant to van-

comycin, usually because of altered cell-wall precursors. VRE can cause healthcare-associated urinary, bloodstream, wound, or other infections and may require susceptibility-guided treatment.

## Van der Woude Syndrome

An autosomal dominant congenital disorder caused by mutations in the IRF6 gene, characterized by the combination of paramedian lower-lip pits (fistulae), cleft lip, and cleft palate, representing the most frequent form of syndromic orofacial clefting.

### Alternate Names:

VWS; Lip-Pit Syndrome

## Vandetanib

A targeted cancer medicine used for selected medullary thyroid cancers. It blocks signals that help cancer cells grow and blood vessels form; diarrhea, high blood pressure, skin effects, and dangerous heart-rhythm changes require monitoring.

## Vaping

Vaping uses an electronic device to heat a liquid and produce an aerosol for inhalation. Aerosols may contain nicotine and other substances that can harm the lungs and cause dependence.

## Vaporiser

A device that converts a liquid medicine or volatile substance into a vapor for inhalation or controlled delivery, such as in anesthesia.

## Variable Decelerations

Brief drops in a baby's heart rate that vary in timing and shape, often caused by pressure on the umbilical cord during labor. Occasional drops may be harmless, but frequent or prolonged drops require assessment of the baby's oxygen supply.

## Variant Angina Pectoris

Chest pain caused by a temporary tightening of an artery supplying the heart, often occurring at rest or at night. Episodes may be severe and can disturb the heart rhythm; sudden or prolonged chest pain needs emergency care.

---

**Alternate Names:**

Prinzmetal Angina; Vasospastic Angina

**Variant CJD**

Alternate name; see *Creutzfeldt-Jakob disease*.

**Variant Classification**

The process of estimating whether a genetic change is harmless, uncertain, or likely to cause disease. Specialists use laboratory evidence, family history, population data, and the person's clinical findings.

**Varibaculum Cambriense**

A bacterial species in the genus *Varibaculum* reported from human clinical specimens. Its role may be opportunistic and requires identification with the clinical context and susceptibility data.

**Variceal Band Ligation**

A procedure using a flexible camera to place small bands around enlarged veins in the food pipe. The bands stop blood flow through the veins and are used to control or prevent bleeding.

**Alternate Names:**

Endoscopic Variceal Ligation; EVL

**Variceal Hemorrhage**

Severe bleeding from enlarged, fragile veins in the food pipe or stomach, usually caused by high pressure in the liver's blood vessels. It can cause vomiting blood, black stools, shock, and death without urgent treatment.

**Varicella**

A primary infection with varicella-zoster virus that causes fever and a generalized itchy rash. See Chickenpox.

**Alternate Names:**

Chickenpox; Varicella-Zoster Virus Infection

**Varicella-Zoster Virus**

The virus that causes chickenpox and shingles. After chickenpox, it can remain inactive in nerve tissue and reactivate years later as a painful, blistering rash called shingles.

**Varicellovirus**

Varicellovirus is a genus of herpesviruses that includes varicella-zoster virus, the cause of chickenpox and shingles.

**Varices**

Abnormally dilated and tortuous veins, commonly occurring in the legs or in the oesophagus and stomach as a result of portal hypertension.

**Varicocele**

Enlargement of veins in the scrotum, often on the left. It may cause aching, a soft "bag of worms" feeling, testicular growth differences, or reduced fertility, but many cause no symptoms.

**Varicocele Embolization**

A procedure that uses a thin tube to place coils or another material in enlarged scrotal veins. This blocks abnormal backward blood flow and reduces pressure in the veins.

**Varicose Ulcer**

An open sore near the ankle caused by long-term poor drainage through the leg veins. It may be surrounded by swelling or skin discoloration and often needs compression and treatment of the underlying vein problem.

**Varicose Veins**

Enlarged, twisted superficial veins, usually in the legs, caused by weak or damaged vein walls or valves that allow blood to pool and flow backward. They may cause visible bulging veins, aching, heaviness, itching, cramps, swelling, skin discoloration, bleeding, superficial inflammation or clotting, and difficult-to-heal venous ulcers.

**Varicose Veins and Venous Insufficiency**

Enlarged superficial veins and inadequate return of blood from the legs because venous valves or the vein wall do not work effectively. Aching, heaviness, swelling, skin discoloration, dermatitis, and ulcers may occur.

---

## Variegate Porphyria

An inherited disorder of heme production that can cause attacks of abdominal pain, vomiting, nerve symptoms, confusion, or dark urine and can also cause fragile, blistering skin after sunlight exposure.

## Variola

A highly contagious and deadly disease caused by variola virus, eradicated globally by 1980 through vaccination. See Smallpox.

## Varix

An abnormally dilated, tortuous vein; multiple varices commonly occur in portal hypertension or chronic venous disease.

## Varus

A limb or joint deformity in which the part farther from the body bends inward, toward the body's center line. Examples include bowlegs and some inward-turning foot deformities.

## Varus Stress Test

A clinical examination manoeuvre used to evaluate the lateral collateral ligaments of a joint, particularly the knee or elbow. A lateral-directed force is applied across the joint line while stabilizing the limb; increased laxity, gapping, or localized pain signifies injury to the lateral collateral ligament complex.

## Vas

A duct or vessel; in reproductive anatomy, the vas deferens transports sperm from the epididymis toward the ejaculatory duct.

## Vasa Deferentia

The paired muscular ducts that transport mature sperm from the epididymides toward the ejaculatory ducts during ejaculation.

### Alternate Names:

Ductus Deferentes; Vas Deferens

## Vasa Efferentia

Small tubes that carry sperm from the testicle's

collecting channels to the epididymis, where sperm mature. They also remove some fluid and help move sperm onward.

## Vasa Previa

A pregnancy complication in which fetal blood vessels run across or near the cervix without the usual protection of the placenta or umbilical cord. If the membranes break, the vessels can tear and cause rapid fetal blood loss, so specialist planning is essential.

## Vasa Recta

Long, hairpin-shaped blood vessels in the inner part of the kidney. They run beside the loops of Henle and help the kidneys concentrate urine.

## Vasa Vasorum

Small blood vessels in the walls of large arteries and veins. They supply oxygen and nutrients to the outer parts of these vessel walls.

## Vascular

Relating to blood vessels or, sometimes, lymphatic vessels. Vascular conditions include narrowed or blocked arteries, aneurysms, blood clots, abnormal vessel connections, and problems with veins; the term may also describe tests, surgery, or medicines involving these vessels.

## Vascular Access

A route into a blood vessel for medicines, fluids, blood tests, monitoring, or dialysis. It may be a short-term catheter, a surgically joined artery and vein, or a synthetic graft, depending on the purpose and expected duration.

## Vascular Access For Hemodialysis

A route that allows blood to leave the body and return during dialysis. It may be a joined artery and vein, a synthetic graft, or a large vein catheter; each has different risks of infection, clotting, and failure.

---

### Vascular Balloon Catheter

An inflatable balloon mounted at the distal tip of a flexible vascular catheter used to dilate stenotic or occluded blood vessels during angioplasty, deliver and expand endovascular stents, or temporarily occlude vascular flow.

#### Alternate Names:

Angioplasty Balloon; Vascular Balloon Dilator; Balloon Catheter

### Vascular Bed

The network of blood vessels supplying a particular organ or tissue. Its vessels widen or narrow to adjust blood flow according to the tissue's needs.

### Vascular Bypass

Surgery that uses a vein or artificial tube to carry blood around a blocked part of an artery and restore blood flow.

#### Alternate Names:

Surgical Bypass; Bypass Grafting; Arterial Bypass Graft

### Vascular Claudication

Alternate name; see *Claudication*.

### Vascular Clip

A small device placed during a procedure to close or control a blood vessel. It may stop bleeding, prevent blood flow to a selected area, or secure a vessel permanently.

#### Alternate Names:

Hemostatic Clip; Surgical Vessel Clip; Ligation Clip

### Vascular Compliance

The ability of a blood vessel to stretch when pressure or blood volume increases. More compliant vessels can hold more blood with a smaller rise in pressure.

#### Alternate Names:

Angiopathic Complications; Vascular Adverse Events

### Vascular Complications

Problems involving blood vessels after disease, injury, or a procedure, such as bleeding, hematoma,

thrombosis, embolism, vessel injury, or reduced blood flow to a limb. Symptoms and urgency depend on the vessel and tissue affected.

### Vascular Dementia

Problems with thinking, memory, mood, or walking caused by strokes or long-term damage to the brain's small blood vessels. Symptoms may appear suddenly or gradually, and treatment focuses on preventing further vessel damage and supporting daily function.

#### Alternate Names:

VaD; Multi-Infarct Dementia; Vascular Cognitive Impairment

### Vascular Disease

Vascular disease affects arteries, veins, or lymphatic vessels and includes atherosclerosis, aneurysm, peripheral artery disease, venous thrombosis, varicose disease, and inflammatory disorders. Symptoms and treatment depend on the vessel and may involve risk-factor control, medicines, procedures, or surgery.

### Vascular Ehlers-Danlos Syndrome

A rare inherited connective-tissue disorder that makes arteries and hollow organs unusually fragile. It may cause thin translucent skin, easy bruising, characteristic facial features, or serious spontaneous blood-vessel or bowel rupture; specialist care and genetic counseling are important.

#### Alternate Names:

vEDS; Ehlers-Danlos Syndrome Type IV

### Vascular Embolic Particles

Small materials placed in a blood vessel during a procedure to block blood flow. They may be used to control bleeding or reduce blood flow to a tumor, fibroid, or other abnormal area.

#### Alternate Names:

Vascular Embolization Particle; PVA Particles

### Vascular Embolization

Intentional blockage of a blood vessel with material delivered through a catheter. It may stop bleeding, reduce blood flow to a tumor, or close an aneurysm



---

or abnormal vessel connection.

**Alternate Names:**

Catheter Embolization; Therapeutic Embolization

**Vascular Embolization Coil**

A catheter-delivered wire device used to block selected blood vessels, often to treat aneurysms or bleeding.

**Alternate Names:**

Endovascular Coil

**Vascular Embolization Plug**

A vascular occlusion device used to block a selected vessel under image guidance. A device placed in a blood vessel to block selected blood flow during image-guided treatment.

**Alternate Names:**

Vascular Occlusion Plug; Embolization Plug; Amplatzer Plug

**Vascular Endothelial Growth Factor**

A protein that helps new blood vessels form and makes blood vessels more leaky. Some medicines block it to treat certain eye diseases and cancers.

**Alternate Names:**

VEGF; Vascular Permeability Factor

**Vascular Endothelial Growth Factor Inhibitor**

A medicine that blocks vascular endothelial growth factor, a protein that helps blood vessels grow. These medicines are used for some cancers and eye diseases involving abnormal new blood vessels.

**Alternate Names:**

Anti-VEGF Agent

**Vascular Endothelial Growth Factor Inhibitors**

Medicines that block vascular endothelial growth factor and reduce abnormal blood-vessel growth or leakage. They are used for some eye diseases and cancers.

**Alternate Names:**

Anti-VEGF Agents; VEGF Neutralizers

**Vascular Forceps**

Surgical forceps designed to handle blood vessels, clots, tissue, or medical materials during vascular procedures. Appropriate use helps avoid crushing or tearing delicate vessels.

**Alternate Names:**

Vascular Surgical Forceps; Atraumatic Clamp

**Vascular Graft**

A tube used to replace or bypass a damaged or blocked blood vessel. It may be made from a person's vein or from a synthetic material; possible problems include infection, bleeding, narrowing, or a clot blocking the graft.

**Vascular Graft Patency**

The continued openness and useful blood flow through a blood-vessel graft or bypass. A clot, narrowing, infection, or scar growth can block it, so examination and sometimes ultrasound are used for monitoring.

**Vascular Hemostat**

A surgical clamp used to temporarily occlude a blood vessel and control bleeding.

**Alternate Names:**

Vascular Clamp; Atraumatic Hemostasis Forceps

**Vascular Imaging**

Tests that show blood vessels and sometimes measure blood flow. Ultrasound, CT, MRI, or catheter-based imaging can find narrowing, blockage, aneurysm, dissection, clot, bleeding, or abnormal vessels; the safest test depends on the body area, urgency, kidney function, and contrast risks.

**Vascularity**

The amount and arrangement of blood vessels supplying a tissue or growth. A highly vascular area may heal or bleed differently and may look different on imaging; blood supply can also affect some treatments.

**Vascular Laser Therapy**

A treatment that uses laser energy to close or change

---

selected blood vessels. The laser may be applied through the skin or through a thin tube.

**Alternate Names:**

Pulsed Dye Laser; PDL; Vascular Laser Therapy

**Vascular Ligation**

Surgical tying, clipping, or sealing of a blood vessel to stop bleeding or redirect blood flow. It may be used during surgery or to treat selected abnormal vessels.

**Alternate Names:**

Vascular Occlusion; Vessel Clipping

**Vascular Malformation**

A blood-vessel or lymph-vessel structure that developed abnormally before birth. It may cause a skin mark, swelling, pain, bleeding, or pressure on nearby organs; effects depend on its type and location.

**Vascular Needle Holder**

A surgical instrument used to hold and guide fine suture needles during repair of blood vessels. It permits precise stitches while minimizing crushing or tearing of delicate vessel walls.

**Alternate Names:**

Vascular Needle Holder; Castroviejo Needle Holder

**Vascular Pain**

Pain caused by impaired blood flow, often appearing in a limb with exertion as claudication or, in more severe disease, at rest.

**Vascular Patch Graft**

A piece of biological or synthetic material sewn onto a blood vessel to widen or repair it.

**Alternate Names:**

Vascular Patch Graft; Angioplasty Patch

**Vascular Retraction**

To pull a vessel or tissue away from an operative field using a surgical retractor.

**Alternate Names:**

Vessel Retraction; Surgical Vessel Exposure

**Vascular Ring**

A birth defect in which blood vessels around the heart form a ring that can press on the windpipe or food pipe. It may cause noisy breathing, cough, repeated chest infections, swallowing difficulty, or feeding problems. Some people need no treatment; symptomatic rings may require surgery.

**Vascular Scissors**

A surgical instrument with two blades used to cut tissue, sutures, dressings, or other materials during vascular procedures.

**Alternate Names:**

Vascular Scissors; Potts Scissors

**Vascular Shunt**

A surgically created or implanted pathway that redirects blood or another fluid from one vessel or body space to another.

**Alternate Names:**

Vascular Shunt; Bypass Shunt

**Vascular Sponge**

A sterile absorbent material used during blood-vessel procedures to absorb blood, apply pressure, or protect delicate tissue.

**Alternate Names:**

Vascular Laparotomy Sponge; Surgical Cottonoid

**Vascular Staple**

A small metal or polymer device used to close tissue or join selected blood-vessel structures. It provides rapid closure but can cause narrowing, bleeding, thrombosis, or tissue injury if unsuitable or misplaced.

**Alternate Names:**

Vascular Surgical Staple; Vessel Closure Device

**Vascular Stent**

A small mesh or tube placed inside a narrowed or blocked vessel to help keep it open.

**Alternate Names:**

Endovascular Stent; Vascular Prosthesis

---

### **Vascular Stent Balloon**

A balloon on a catheter used to expand a vascular stent against the vessel wall and restore an open channel. Inflation can cause vessel injury, bleeding, clotting, or rarely rupture.

#### **Alternate Names:**

Stent-Expansion Balloon; Angioplasty Balloon

### **Vascular Stent-Graft**

A fabric-covered metal frame placed inside a blood vessel to reinforce a weak wall or seal an aneurysm, tear, or leak. Follow-up imaging checks for movement, blockage, or leakage around the graft.

#### **Alternate Names:**

Endovascular Graft; Aortic Stent-Graft

### **Vascular Suction**

A suction device used during a vascular procedure to remove blood or other fluid from the operative field.

#### **Alternate Names:**

Vascular Suction Cannula; Frazier Suction

### **Vascular Surgery**

Operations that repair, widen, replace, or reroute blood vessels to restore circulation or prevent bleeding. Procedures may treat aneurysms, blocked arteries, blood-vessel injuries, or problems with veins and dialysis access.

### **Vascular System**

The network of arteries, veins, and capillaries that carries blood throughout the body. It delivers oxygen, nutrients, and hormones to tissues and carries carbon dioxide and other waste products away. Disease may narrow, weaken, block, or inflame these vessels.

### **Vascular Thrombectomy**

Removal of a blood clot from a blood vessel using surgery or a catheter-based procedure. The aim is to restore blood flow and protect tissue from damage when a clot blocks an important artery or vein.

#### **Alternate Names:**

Vascular Thrombectomy; Clot Extraction

### **Vascular Tissue Adhesive**

A liquid material used to block selected blood vessels during image-guided embolization. Doctors inject it through a catheter to control bleeding, close an abnormal vessel, or reduce blood flow to a tumor.

#### **Alternate Names:**

Cyanoacrylate Embolization Glue; Endovascular Embolic Adhesive

### **Vascular Tone**

The usual degree of tightening in the muscle within blood-vessel walls. It helps control vessel width, blood flow, and blood pressure.

### **Vascular Trauma**

Injury to an artery or vein caused by a blunt blow, penetrating wound, crush injury, or medical procedure. It may cause severe bleeding or loss of blood flow and needs urgent assessment of circulation and nearby nerves.

### **Vascular Ulcer**

A persistent open wound caused or maintained by poor blood flow, high pressure in the veins, loss of protective feeling, or another blood-vessel problem. It usually affects the legs or feet and may heal slowly or become infected.

#### **Alternate Names:**

Vascular Ulceration; Ischemic Ulcer; Arterial Ulcer

### **Vasculitic Neuropathy**

Nerve damage caused by inflammation of the small blood vessels that supply nerves. It often causes painful, uneven numbness or weakness and may affect several limbs as the vessel inflammation continues.

### **Vasculitis**

Inflammation of blood vessels that can narrow them, reduce blood flow, or cause bleeding and tissue damage. It may affect large, medium, or small vessels and can involve the skin, nerves, kidneys, lungs, or other organs.

#### **Alternate Names:**

Angiitis; Arteritis; Blood Vessel Inflammation

---

## **Vas Deferens**

A muscular tube that carries sperm from the epididymis toward the urethra during ejaculation.

## **Vasectomy**

A permanent contraceptive procedure in which the vas deferens are divided or occluded to prevent sperm from entering the ejaculate. It does not immediately provide sterility and does not protect against sexually transmitted infections.

## **Vasectomy Reversal**

Surgery to reconnect the tubes that carry sperm after a vasectomy. It may restore sperm to the semen, but pregnancy is not guaranteed.

### **Alternate Names:**

Vasovasostomy; Vasoepididymostomy

## **Vasoactive Intestinal Peptide Test**

A blood test measuring a hormone that affects the intestine and blood vessels. It is used mainly when a rare hormone-producing tumor is suspected in someone with severe watery diarrhea, dehydration, or abnormal body salts.

## **Vasoconstriction**

Narrowing of blood vessels caused by contraction of vascular smooth muscle, which can raise blood pressure or reduce blood flow to affected tissues.

## **Vasoconstrictor**

A substance that narrows blood vessels by making their muscle walls contract. This can raise blood pressure and reduce local blood flow; vasoconstrictor medicines are used in selected emergencies or with local anesthetics and require careful monitoring.

## **Vasoconstrictors**

Substances or medicines that narrow blood vessels by making their muscular walls contract. They can raise blood pressure or reduce local blood flow and are used only for selected conditions because excessive narrowing can be harmful.

## **Vasodepressor Syncope**

Fainting caused mainly by sudden widening of blood vessels and a fall in blood pressure. It often occurs as part of a vasovagal reaction and may be preceded by nausea, sweating, warmth, or lightheadedness.

## **Vasodilatation**

Widening of blood vessels caused by relaxation of vascular smooth muscle, increasing blood flow and potentially lowering blood pressure.

## **Vasodilation**

Widening of blood vessels when their muscular walls relax. It can increase blood flow and lower blood pressure and occurs with exercise, heat, medicines, and inflammation.

## **Vasodilator**

A substance that relaxes blood-vessel walls and makes the vessels wider. It can lower blood pressure or improve blood flow and is used for selected heart, blood-pressure, or circulation problems; excessive widening may cause headache, flushing, dizziness, or fainting.

## **Vasodilators**

Medicines or substances that relax vascular smooth muscle and widen blood vessels. They are used for conditions such as hypertension, angina, heart failure, or pulmonary hypertension depending on the agent.

## **Vasodilatory Shock**

A form of distributive shock characterized by severe systemic arterial hypotension and tissue hypoperfusion resulting from profound, refractory peripheral vasodilation and loss of vascular tone despite preserved or elevated cardiac output, most frequently seen in severe sepsis, anaphylaxis, and neurogenic injury.

## **Vasomotor**

Relating to control of blood-vessel width by nerves, hormones, and local chemicals. Vasomotor changes narrow or widen vessels and help regulate blood pressure, body temperature, tissue blood flow, and

---

skin color.

### **Vasomotor Center**

A group of nerve cells in the lower brain that helps control the widening and narrowing of blood vessels and helps maintain blood pressure.

### **Vasomotor Centre**

Groups of nerve cells in the brainstem that help control blood-vessel narrowing and widening. By changing vessel tone, they influence blood pressure, heart and tissue blood flow, and the body's response to standing or stress.

### **Vasomotor Nerves**

Autonomic nerves controlling blood-vessel widening and narrowing by changing smooth-muscle activity, helping regulate blood pressure and temperature.

### **Vasomotor Rhinitis**

Common subtype term; see *Nonallergic rhinitis*.

### **Vaso-Occlusive Crisis**

The classic acute, excruciatingly painful hallmark complication of sickle cell disease, caused by the intravascular sludging and adherence of rigid, polymer-filled sickled erythrocytes in the microvasculature, producing focal tissue ischemia, infarction, and systemic inflammation in bones, joints, or viscera.

#### **Alternate Names:**

Sickle Cell Pain Crisis; VOC

### **Vasopressin**

A hormone that increases water reabsorption in the kidney and can constrict blood vessels. It is released from the posterior pituitary.

#### **Alternate Names:**

Arginine Vasopressin; AVP

### **Vasopressin Receptor**

A cell sensor that responds to the hormone vasopressin. Different types help the kidneys retain water and help blood vessels narrow, influencing body-fluid balance and blood pressure.

### **Vasopressin Receptor Antagonists**

A pharmacological class of aquaretic medications (vaptans, such as tolvaptan and conivaptan) that competitively block vasopressin V2 receptors in the renal collecting duct, promoting solute-free water excretion without electrolyte loss in euvolemic and hypervolemic hyponatremia (SIADH, heart failure, cirrhosis).

#### **Alternate Names:**

Vaptans

### **Vasopressor**

A medicine or body signal that raises blood pressure, mainly by narrowing blood vessels and sometimes by strengthening the heartbeat. These medicines are used in selected emergencies such as shock and require close monitoring.

### **Vasopressors In Anesthesia**

Medicines that increase vascular tone or cardiac output to support blood pressure during anesthesia or critical illness. The choice and dose depend on the cause of hypotension, heart rate, cardiac function, and local protocol; monitoring is required.

### **Vasospasm**

Sudden narrowing of a blood vessel caused by contraction of its muscular wall. It can temporarily reduce blood flow and cause chest pain, headache, stroke-like symptoms, or tissue injury, depending on the vessel affected.

### **Vasovagal Attack**

A transient reflex episode caused by reduced sympathetic tone and increased vagal activity, producing hypotension and often bradycardia with fainting or near-fainting. Common triggers include prolonged standing, pain, fear, heat, dehydration, or emotional stress.

### **Vasovagal Reaction**

A reflex that slows the heart and widens blood vessels, briefly lowering blood pressure and sometimes causing fainting; triggers include pain, fear, prolonged standing, heat, or blood exposure.

---

## Vasovagal Syncope

A common faint caused by a reflex that briefly slows the heart or widens blood vessels, lowering blood pressure and brain blood flow. Triggers include standing still, heat, pain, fear, or seeing blood; warning symptoms often occur first.

### Alternate Names:

Neurocardiogenic Syncope

## Vater-Pacini Corpuscles

Large, rapidly adapting, encapsulated mechanoreceptors located deep in the dermis, subcutaneous tissue, periosteum, and joint capsules. Morphologically composed of a central unmyelinated nerve axon surrounded by multiple concentric lamellae of flattened Schwann-like cells separated by fluid; specialized physiologically to detect high-frequency mechanical vibration (200 to 300 Hz) and transient acceleration.

### Alternate Names:

Pacinian Corpuscles; Lamellar Corpuscles

## VATS

Alternate name; see *Video-Assisted Thoracoscopic Surgery*.

## Vaughan Williams Classification

A system that groups medicines used for abnormal heart rhythms according to their main effect on the heart's electrical activity. It helps describe treatment but does not replace patient-specific prescribing.

## VCJD

Alternate name; see *Creutzfeldt-Jakob disease*.

## VDRL Test

A blood or spinal-fluid test that looks for an immune reaction associated with syphilis. A reactive result needs a more specific confirmatory test and may help monitor response to treatment.

## Vector

A living carrier, usually an insect or tick, that spreads an infection between people or animals.

Some vectors allow the germ to develop inside them; others transfer it mechanically without the germ changing.

## Vector-Borne Disease

An infection spread by a living carrier such as a mosquito, tick, flea, or sandfly. Examples include malaria, dengue, Lyme disease, and some forms of viral encephalitis; prevention focuses on avoiding bites and controlling the carrier.

## Vector Control

Measures that reduce insects or other animals that spread infection. Examples include removing standing water, using insecticides, treating breeding areas, improving buildings, wearing protective clothing, and using repellents or bed nets.

## Vegetarian

Vegetarian describes a diet that avoids meat and may include dairy products, eggs, or both. Nutritional needs can usually be met with varied foods, but iron, vitamin B12, protein, and other nutrients may need attention.

## Vegetation

A mass that can form on a heart valve during infective endocarditis. It may contain bacteria, blood-clotting material, and immune cells and can damage the valve or break off and travel to other organs.

## Vein

A blood vessel that carries blood toward the heart. Most veins carry oxygen-poor blood, but veins from the lungs and placenta carry oxygen-rich blood. Veins often contain valves that help prevent backward flow.

## Vein-Graft Aneurysms

An abnormal widening of a vein used to bypass a blocked heart artery. It may contain a clot, block blood flow, press on nearby structures, or rupture and therefore needs specialist assessment.



---

## **Vein of Galen Malformation**

A rare congenital intracranial vascular anomaly resulting from an embryonic arteriovenous fistula that drains directly into the primitive median prosencephalic vein (the precursor of the great cerebral vein of Galen), presenting in neonates with high-output congestive heart failure and hydrocephalus.

### **Alternate Names:**

VGM; Vein of Galen Aneurysmal Malformation

## **VeIP Regimen**

A cancer-treatment combination of vinblastine, ifosfamide, and cisplatin, used mainly for selected germ-cell tumors. It can lower blood counts and damage the kidneys, nerves, or bladder, so specialist monitoring is required.

## **Velamentous Cord Insertion**

An abnormal attachment in which the umbilical blood vessels travel through the membranes before reaching the placenta without the usual protective tissue. The vessels can be compressed or torn, increasing risks of poor fetal growth, bleeding, or vasa previa and requiring specialist pregnancy monitoring.

### **Alternate Names:**

Velamentous Insertion of the Umbilical Cord

## **Velocardiofacial Syndrome**

Alternate name; see *DiGeorge syndrome (22q11.2 deletion syndrome)*.

## **Velopharyngeal Insufficiency**

A structural or functional anatomical failure of the soft palate (velum) and posterior pharyngeal wall to achieve complete closure during speech and deglutition, resulting in hypernasal speech resonance, nasal air emission, and nasal regurgitation of fluids.

### **Alternate Names:**

VPI; Velopharyngeal Incompetence

## **Vemurafenib**

A targeted cancer medicine for tumors with a specific BRAF V600 gene change, including some melanomas. It can cause joint pain, sun sensitiv-

ity, abnormal heart rhythm, new skin cancers, or treatment resistance.

## **Vena Cava**

Either of the two largest veins returning blood to the right side of the heart. The superior vena cava drains the upper body, and the inferior vena cava drains the abdomen, pelvis, and legs.

### **Alternate Names:**

Venae Cavae; Superior and Inferior Vena Cava

## **Vena Cava Filter**

A small device placed in the large vein leading to the heart to trap blood clots traveling from the legs toward the lungs. It may be used when blood-thinning medicines cannot be used or have not worked.

### **Alternate Names:**

IVC Filter; Greenfield Filter

## **Venepuncture**

Inserting a needle into a vein to collect blood or place intravenous access. It may cause brief pain, bruising, bleeding, infection, or fainting, and clean technique reduces complications.

## **Venereal Disease Research Laboratory Test**

A blood test that looks for antibodies associated with syphilis. It is used for screening and monitoring treatment, but a reactive result needs confirmation with a treponemal test.

## **Venesection**

The controlled removal of blood from a vein, used diagnostically or therapeutically, for example in selected cases of hemochromatosis or polycythemia.

## **Venipuncture**

Puncture of a vein with a needle to obtain blood, administer fluids or medicines, or establish intravenous access. Bruising, infection, bleeding, and nerve or artery injury are uncommon complications.

## **Venogram**

Contrast imaging of veins used to detect thrombosis,

---

obstruction, abnormal anatomy, or selected venous flow disorders.

### **Venography**

Imaging of veins, usually after contrast material is administered, to assess venous anatomy, obstruction, thrombosis, or malformation.

### **Venom**

A poisonous substance delivered by an animal through a bite, sting, or similar structure. Its effects vary by species and may include pain, swelling, nerve problems, bleeding, or dangerous changes in breathing or blood pressure.

### **Venomous Bite**

A bite that injects venom from an animal such as a snake, spider, or arthropod. Effects range from local pain and swelling to bleeding, paralysis, or systemic toxicity.

### **Veno-Occlusive Disease**

Blockage and injury of small veins in the liver, especially after some chemotherapy or stem-cell transplantation. It can cause painful liver enlargement, rapid weight gain, fluid retention, and jaundice and can be life-threatening.

### **Veno-occlusive Hepatic Disease (Sinusoidal Obstruction Syndrome)**

A potentially severe liver injury caused by narrowing or blockage of hepatic sinusoids, classically after hematopoietic-cell transplantation or certain chemotherapy. Painful liver enlargement, weight gain, fluid retention, jaundice, and rising bilirubin may occur.

### **Venous**

Relating to veins, which usually carry blood back toward the heart. Vein problems include blood clots, varicose veins, poor return of blood, and ulcers; pulmonary veins are an exception because they carry oxygen-rich blood from the lungs to the heart.

### **Venous Air Embolism**

Obstruction of the venous circulation by intravascular gas, usually after procedures, trauma, or pressure changes. It can impair pulmonary or cardiac function and, if gas reaches the arterial circulation, cause neurologic or systemic ischemia.

### **Venous Aneurysm**

A localized abnormal dilatation of a vein. It may be asymptomatic or present as a soft mass, pain, thrombosis, embolism, or bleeding depending on its location and size.

### **Venous Blood Gas**

The clinical laboratory measurement of pH, partial pressure of carbon dioxide ( $p\text{CO}_2$ ), bicarbonate ( $\text{HCO}_3^-$ ), and base excess in a peripheral or central venous blood sample to assess systemic acid-base balance, ventilatory status, and tissue perfusion.

#### **Alternate Names:**

Venous Blood Gas; VBG; Central Venous Blood Gas

### **Venous Capacitance**

The physical and physiological capacity of the systemic venous circulation to accommodate large changes in blood volume with minimal alterations in venous pressure, functioning as the primary high-capacitance, low-resistance reservoir holding approximately 60 to 70 percent of total blood volume at rest.

### **Venous Diseases**

Conditions affecting veins, including blood clots, varicose veins, poor return of blood, inflammation, ulcers, injuries, and birth-related malformations. They may cause pain, swelling, skin changes, bleeding, or dangerous clots traveling to the lungs.

### **Venous Hum**

A harmless continuous humming sound heard over the neck veins, most often in healthy children. It changes or disappears with body position or gentle pressure on the vein, helping distinguish it from heart murmurs that need further evaluation.

#### **Alternate Names:**

---

Cervical Venous Hum; Innocent Venous Hum

### **Venous Insufficiency**

Impaired return of blood through the veins, usually in the legs, because valves allow backward flow, veins are blocked, or the calf-muscle pump is ineffective. Blood may pool and cause aching, heaviness, swelling, varicose veins, itching, darkened or hardened skin, stasis dermatitis, and difficult-to-heal ulcers near the ankle.

### **Venous Leak**

Difficulty maintaining an erection because blood leaves the penis too readily during erection. It may reflect vascular, nerve, hormonal, or medication-related problems and should be evaluated as erectile dysfunction.

#### **Alternate Names:**

Veno-Occlusive Dysfunction; Cavernous Venous Leakage

### **Venous Pressure**

The pressure of blood inside the veins. It depends on blood volume, vein width, gravity, and how well the right side of the heart pumps; abnormally high pressure can cause swelling, enlarged veins, or fluid buildup.

### **Venous Return**

The flow of blood through the veins back to the heart. It is helped by vein valves, muscle movement, breathing, and pressure changes in the chest.

### **Venous Stasis Dermatitis**

Inflammation of the skin of the lower legs caused by chronic swelling and pooling of blood and fluid, usually from chronic venous insufficiency. It may cause itching, redness, scaling, skin darkening, thickening, or ulcers; treatment addresses the swelling and protects the skin.

### **Venous Stasis Ulcer**

A long-lasting wound near the ankle caused by poor blood return through the leg veins. The leg may also be swollen, painful, or have darkened, thickened skin; compression and treatment of the vein problem

may help.

### **Venous Thromboembolism**

A blood clot that forms in a vein, usually deep in the leg, and may travel to the lungs. A clot in the lungs can cause sudden breathlessness, chest pain, collapse, or death.

### **Venous Thromboembolism In Pregnancy**

A blood clot in a deep vein or lungs during pregnancy or after birth. Pregnancy increases clot risk, and symptoms such as one-sided leg swelling, chest pain, sudden breathlessness, or coughing blood can indicate a serious clot.

### **Venous Thromboembolism Prophylaxis**

Measures used to prevent dangerous vein clots, especially around surgery or hospitalization. They may include walking early, compression devices or stockings, and blood-thinning medicine when bleeding risk is acceptable.

### **Venous Thrombosis**

The formation of a blood clot (thrombus) within a vein, most commonly in the deep veins of the lower extremities (deep vein thrombosis, DVT).

### **Venous Ulcer**

A slow-healing sore caused by poor blood return through the leg veins, usually near the ankle. It may occur with leg swelling, varicose veins, or skin discoloration. Arterial circulation should be assessed because it affects whether compression treatment is safe.

### **Ventilation**

Movement of air into and out of the lungs; effective ventilation brings oxygen to the alveoli and removes carbon dioxide.

### **Ventilation-Perfusion Mismatch**

A mismatch between air reaching parts of the lungs and blood reaching those same parts. It can lower blood oxygen and occur with pneumonia, COPD, asthma, heart failure, or a lung clot.

---

## Ventilation-Perfusion Ratio

The relationship between air reaching the lungs and blood reaching the lung tissues. Good matching allows efficient oxygen exchange; poor matching can cause low blood oxygen.

## Ventilation-Perfusion Scan

A nuclear-medicine scan that compares air movement in the lungs with blood flow through them. Areas with blood flow blocked but preserved air movement may suggest a pulmonary embolism. The test uses a breathed tracer and an injected tracer and is interpreted with symptoms and other findings.

## Ventilator

A machine that supports or replaces spontaneous breathing by delivering air, oxygen, or an air-oxygen mixture under positive pressure. Noninvasive ventilators deliver support through a mask, while invasive ventilators connect through an endotracheal or tracheostomy tube. Ventilators can provide different patterns and levels of pressure or volume according to the device and clinical purpose.

### Alternate Names:

Mechanical Ventilator Breathing Machine

## Ventilator-Associated Pneumonia

A lung infection that develops after a person has used a breathing machine for at least two days. It may cause fever, worsening oxygen levels, cough, or abnormal lung findings and requires prompt evaluation and treatment.

## Ventilator-Induced Lung Injury

Lung damage caused or worsened by a breathing machine when pressure, volume, or oxygen levels are excessive. Clinicians use protective settings and close monitoring to support breathing while limiting further injury.

## Ventouse

A vacuum extractor placed on the baby's scalp to assist a vaginal birth when delivery needs help. It is used only when specific safety conditions are met and can cause temporary scalp swelling or bruising.

## Ventral

Relating to the belly or front side of the body; synonym for anterior in human anatomy.

## Ventral Body Cavity

The body cavity toward the front. It includes the thoracic cavity and the abdominopelvic cavity, which contain many of the body's major organs.

## Ventral Hernia

Protrusion of abdominal tissue through a weakness in the front abdominal wall, including incisional and umbilical hernias. It may cause a bulge or pain and requires urgent assessment if incarceration or strangulation is suspected.

## Ventral Root

The front root of a spinal nerve. It carries movement and other outgoing signals from the spinal cord.

## Ventricle

One of the two lower chambers of the heart that pump blood out of it. The right ventricle sends blood to the lungs to receive oxygen; the left ventricle sends oxygen-rich blood to the rest of the body.

## Ventricular

Relating to a ventricle. In the heart, it concerns one of the lower pumping chambers; in the brain, it concerns the fluid-filled spaces that contain cerebrospinal fluid. The organ gives the term its specific meaning.

## Ventricular Aneurysm

A localized, thin, fibrous, dyskinetic or akinetic bulging outpouching of the scarred left ventricular wall developing as a late mechanical complication of transmural myocardial infarction (typically anterior MI), predisposing to heart failure, refractory ventricular arrhythmias, and apical mural thrombus formation.

## Ventricular Arrhythmia

An abnormal heart rhythm arising from the ventri-

---

cles. It may cause palpitations, dizziness, fainting, chest discomfort, or sudden cardiac arrest; fainting or severe symptoms require emergency care.

### **Ventricular Arrhythmias**

Abnormal heart rhythms arising from the ventricles. They range from premature ventricular contractions to ventricular tachycardia and fibrillation, which can cause sudden cardiac arrest.

### **Ventricular Assist Device**

A mechanical pump that helps a weakened heart move blood. It may support the left side, right side, or both and can be temporary or long-term, but bleeding, infection, stroke, and clots are important risks.

### **Ventricular Bigeminy**

A cardiac rhythm pattern on electrocardiogram in which every normal sinus beat is consistently followed by a premature ventricular complex (PVC) in a regular alternating 1:1 sequence, commonly provoked by electrolyte derangements, digitalis toxicity, or underlying ischemic heart disease.

### **Ventricular Fibrillation**

A chaotic, disorganized ventricular rhythm in which the ventricles quiver rather than contract, producing no effective cardiac output and cardiac arrest. It is a shockable rhythm; immediate CPR and unsynchronized defibrillation are required, with treatment of the underlying cause.

**Alternate Names:**

V-Fib

### **Ventricular Hypertrophy**

Thickening of the wall of one of the heart's lower chambers. It may develop from long-term high blood pressure, valve disease, inherited heart muscle disease, or intense training and can lead to stiffness, abnormal rhythms, or heart failure.

### **Ventricular Myocardium**

The heart muscle forming the walls of the ventricles. Its contraction pumps blood to the lungs and the rest of the body.

### **Ventricular Non-Compaction**

A primary genetic cardiomyopathy characterized by a thickened myocardial wall with a non-compacted endocardial layer displaying prominent, deep intertrabecular recesses and deep trabeculations, typically affecting the left ventricular apex and lateral wall.

**Alternate Names:**

LVNC; Non-Compaction Cardiomyopathy

### **Ventricular Premature Contraction**

Alternate name; see *Premature ventricular contractions (PVCs)*.

### **Ventricular Remodeling**

Progressive alterations in left ventricular geometry, mass, chamber volume, and spherical shape occurring after acute myocardial infarction or chronic pressure/volume overload, driven by neurohormonal activation, cardiomyocyte hypertrophy, and interstitial fibrosis.

**Alternate Names:**

Post-Infarction Remodeling; Cardiac Remodeling

### **Ventricular Rupture**

A tear in a wall of one of the heart's lower chambers, often after a heart attack. It can cause life-threatening bleeding around the heart.

### **Ventricular Septal Defect**

A hole in the wall between the heart's lower chambers. Blood may flow from the stronger left side to the right and increase blood flow to the lungs. Small holes may cause no problems; larger ones can cause heart failure or lung-vessel damage.

### **Ventricular Septal Defect Cross-Reference**

Alternate name; see *Ventricular septal defect*.

### **Ventricular Tachycardia**

A rapid heart rhythm that starts in the lower chambers of the heart. It may cause palpitations, dizziness, fainting, shock, or cardiac arrest and can be life-threatening.



---

## Ventriculoatrial

Relating to a connection between a ventricle and an atrium. In brain surgery, a ventriculoatrial shunt is a tube that drains extra fluid from a brain cavity into the right upper chamber of the heart.

## Ventriculography

Imaging used to measure how well the heart's ventricles pump blood. A radionuclide scan or cardiac MRI can estimate the amount of blood ejected with each heartbeat and show abnormal wall movement; the method is chosen for the clinical question.

## Ventriculoperitoneal Shunt

A surgically implanted system that drains cerebrospinal fluid from a brain ventricle into the abdominal cavity, where the fluid can be absorbed.

## Ventrosuspension

Surgical fixation of a displaced or prolapsed uterus to the anterior abdominal wall or nearby supporting tissues.

## Venturi Mask

A high-flow oxygen delivery device that uses the Venturi principle to supply a precise, fixed fraction of inspired oxygen ( $\text{FiO}_2$ , typically 24% to 50%). Oxygen passing through a calibrated narrow jet nozzle creates a pressure drop that draws in a predictable volume of room air through color-coded entrainment ports. Because the total gas flow meets or exceeds the patient's peak inspiratory flow rate, the delivered oxygen concentration remains constant regardless of the patient's breathing rate or depth, making it vital for managing hypoxemia in patients with chronic obstructive pulmonary disease at risk of hypercapnic respiratory failure.

### Alternate Names:

Air-Entrainment Mask; Venturi Oxygen Mask; High-Flow Oxygen Mask

## Venturi Principle

A physical principle in fluid dynamics stating that as a fluid or gas flows through a constricted section of a pipe or nozzle, its velocity increases and its static lateral pressure decreases. In medicine, this

localized pressure drop creates suction that entrains ambient air or fluids, forming the operational basis for air-entrainment oxygen masks (Venturi masks), jet nebulizers, suction aspirators, and explaining hemodynamic phenomena such as systolic anterior motion of the mitral valve in hypertrophic obstructive cardiomyopathy.

### Alternate Names:

Venturi Effect; Bernoulli-Venturi Principle; Air-Entrainment Principle

## Venule

A small blood vessel that collects blood from capillaries and carries it into larger veins. Venules also allow white blood cells and fluid to leave the bloodstream during inflammation.

## Venules

Small blood vessels that collect blood from capillaries and join to form veins. Their walls also participate in inflammation by allowing white blood cells to leave the bloodstream.

## Verapamil

A non-dihydropyridine calcium-channel blocker used for selected hypertension, angina, and supraventricular tachyarrhythmias; it can cause bradycardia and constipation.

## Verapamil Toxicity

Acute severe poisoning from the non-dihydropyridine calcium channel blocker verapamil, characterized by profound peripheral vasodilation, profound myocardial depression, high-grade atrioventricular block, refractory bradycardia, and cardiogenic shock; managed with intravenous calcium, high-dose insulin euglycemia therapy, and lipid emulsion.

## Vermicides

Medicines or chemicals that kill parasitic worms. The correct treatment depends on the worm, the body site affected, the person's age, and other health conditions.



---

### Vernal Keratoconjunctivitis

A severe, recurrent allergic eye inflammation that primarily affects young boys in warm, sunny climates and often flares up during the spring and summer (spring catarrh). It causes intense eye itching, light sensitivity, watering, and thick, stringy mucus. Characteristic findings include large, cobblestone-like bumps on the underside of the upper eyelid (giant papillae) and small white dots along the border of the cornea (Horner-Trantas dots). In severe cases, inflammation and eye-rubbing can create a non-healing sore on the clear front of the eye (a shield ulcer), which can lead to corneal scarring and vision changes. The condition typically improves on its own after puberty.

#### Alternate Names:

VKC; Spring Catarrh; Vernal Catarrh; Vernal Conjunctivitis

### Verner-Morrison Syndrome

A rare neuroendocrine tumor syndrome (WDHA syndrome) caused by a vasoactive intestinal peptide-secreting pancreatic islet cell tumor (VIPoma), characterized by the tetrad of severe watery diarrhea ('pancreatic cholera'), hypokalemia, achlorhydria (hypochlorhydria), and metabolic acidosis.

#### Alternate Names:

VIPoma Syndrome; WDHA Syndrome

### Vernix

A white, greasy protective coating on the skin of a fetus or newborn, composed mainly of water, lipids, and shed skin cells.

### Vernix Caseosa

A white, greasy protective coating on the skin of a fetus or newborn, composed mainly of water, lipids, and shed skin cells.

### Verruca

Alternate name; see *Wart*.

### Verruca Plana

Small, smooth, relatively flat warts caused by HPV, often occurring in groups on the face, hands, or legs.

#### Alternate Names:

Flat Warts; Juvenile Warts

### Verrucas

Verrucas are rough skin growths on the feet caused by human papillomavirus. They may be painful with pressure and often resolve without treatment, although treatment can reduce discomfort or spread.

### Verruca Vulgaris

A common, benign, hyperkeratotic, exophytic cutaneous papilloma with a rough, verrucous ('warty') surface and thrombosed capillary black dots, caused by localized epidermal infection by human papillomavirus (HPV types 2, 4, 7, and 1).

#### Alternate Names:

Common Wart; Viral Verruca

### Verruciform Xanthoma

A rare, noncancerous growth usually found in the mouth or genital area. It may look like a wart, so examination and sometimes a tissue sample are needed to distinguish it from cancer.

### Verrucose

Having a rough, thickened, wart-like, or finger-like surface. It is a descriptive term for a skin, mucosal, or other lesion and does not by itself identify whether the lesion is benign, infectious, or cancerous.

### Verrucous

Having a rough, thickened, irregular, or wart-like surface. The appearance can occur in benign growths, infections, chronic inflammation, or some cancers.

### Verrucous Carcinoma

A low-grade, well-differentiated variant of squamous cell carcinoma with a warty, exophytic appearance. It is locally invasive but rarely metastasizes, commonly affecting the oral cavity, skin, or genitalia.

### Version

A change in the position or presentation of a body

---

structure. In pregnancy, fetal version is a procedure or movement that changes the baby's position, often to place the head downward before birth.

### **Vertebra**

One of the bones forming the spinal column. Each vertebra helps support the body, allows movement, provides attachment for muscles, and surrounds and protects the spinal cord.

### **Vertebrae**

Vertebrae are the bones of the spinal column that protect the spinal cord, support the body, allow movement, and provide attachment for muscles and ligaments.

### **Vertebral Artery**

One of two arteries that travel through the neck and join to supply the brainstem, cerebellum, and back of the brain. Narrowing or blockage can cause dizziness, imbalance, or stroke.

### **Vertebral Artery Dissection**

A tear in the inner lining of a vertebral artery that lets blood enter the vessel wall, forming an intramural hematoma and possibly narrowing the artery or creating a pseudoaneurysm. It may occur spontaneously or after neck injury and can cause sudden neck or occipital headache, dizziness, imbalance, or a posterior-circulation transient ischemic attack or stroke.

#### **Alternate Names:**

VAD; Spontaneous Vertebral Artery Dissection

### **Vertebral Canal**

The passage formed by the stacked spinal bones that surrounds and protects the spinal cord. Narrowing, bleeding, tumors, fractures, or disc disease inside it can compress nerves and cause weakness or pain.

### **Vertebral Column**

The chain of bones running from the neck to the pelvis. It supports the head and trunk, protects the spinal cord, allows bending and twisting, and provides attachment for muscles and ribs.

### **Vertebral Compression Fracture**

Collapse or cracking of one of the bones in the spine, often from osteoporosis, injury, or cancer. It may cause sudden or gradual back pain, loss of height, or a rounded back and can sometimes press on nerves.

### **Vertebral Hemangioma**

Alternate name; see *Spinal hemangioma*.

### **Vertebral Tumor**

A growth arising in or near a spinal bone. It may be benign or cancerous, start in the spine or spread there, and cause pain, fracture, nerve compression, weakness, or paralysis.

### **Vertebrobasilar Circulatory Disorders**

Conditions that reduce blood flow through the vertebral or basilar arteries supplying the brainstem, cerebellum, and posterior brain; symptoms may include vertigo, double vision, imbalance, weakness, or stroke.

### **Vertebrobasilar Insufficiency**

Reduced blood flow through arteries supplying the back of the brain. It may cause dizziness, double vision, imbalance, weakness, trouble speaking, or stroke-like symptoms and can be serious when sudden.

### **Vertebrobasilar Insufficiency**

Source spelling variant of *vertebrobasilar insufficiency*, reduced blood flow in the vertebral or basilar circulation supplying the brainstem, cerebellum, and posterior brain.

### **Vertebroplasty**

A procedure in which medical cement is injected into a collapsed spinal bone to help stabilize it and sometimes reduce pain. It is used selectively because cement leakage, infection, nerve injury, or other complications can occur.

### **Vertical**

Oriented from superior to inferior or perpendicular to a horizontal plane. In medicine, it may describe

---

position, transmission from parent to child, or a body-oriented imaging or movement axis.

### **Vertical Talus**

A stiff flatfoot present from birth in which one of the ankle bones points downward and the middle of the foot is bent upward. It may occur alone or with other conditions and usually needs early orthopedic treatment.

### **Vertical Transmission**

Passage of an infection or genetic condition from parent to child during pregnancy, delivery, or breastfeeding.

### **Vertigo**

A false feeling that you or your surroundings are moving, spinning, tilting, or shifting when they are not. It is a symptom rather than a disease and often comes from the inner ear, but brain and other conditions can also cause it.

### **Vertigo-Associated Disorders**

Vertigo-associated disorders cause an illusion of movement from vestibular, neurologic, visual, medication-related, or systemic dysfunction and may be accompanied by imbalance, nausea, or hearing symptoms.

#### **Alternate Names:**

Vestibular Disorders; Peripheral Vestibulopathy; Labyrinthine Disorders

### **Very Long-Chain Acyl-CoA Dehydrogenase Deficiency**

A rare inherited disorder in which cells cannot use certain long-chain fats for energy, especially during fasting or illness. Infants may develop low blood sugar, heart disease, or liver problems; older children or adults may have exercise-triggered muscle breakdown and dark urine.

#### **Alternate Names:**

VLCAD Deficiency; VLCADD

### **Very-Low-Density Lipoprotein**

A particle made by the liver that carries mainly

triglycerides through the blood. High levels can contribute to fatty deposits in artery walls and increase cardiovascular risk, although the result is interpreted with the full lipid profile.

#### **Alternate Names:**

VLDL

### **Vesical**

Relating to the urinary bladder. For example, vesical inflammation affects the bladder lining, a vesical tumor is a bladder growth, and the vesical neck is the bladder outlet leading toward the urethra.

### **Vesicant**

A substance that causes blistering and tissue injury on contact. In oncology, vesicant drugs can cause severe extravasation injury if they escape from a vein into surrounding tissue.

### **Vesicants**

Substances that can cause severe blistering and tissue damage when they touch skin or escape from a vein. Some chemotherapy medicines and industrial chemicals are vesicants.

### **Vesicle**

A small, circumscribed, raised skin lesion containing clear, serous, or sometimes bloody fluid, usually less than 1 cm in diameter. A vesicle is a small blister; causes include viral infection, contact irritation, eczema, burns, friction, and immune-mediated skin disease.

### **Vesicles**

Plural of vesicle; see *Vesicle*.

### **Vesicoureteral Reflux**

Backward flow of urine from the bladder toward the kidneys because the valve where a ureter enters the bladder does not work normally. It can cause repeated kidney infections and scarring, especially in children.

#### **Alternate Names:**

VUR; Urinary Reflux

---

### **Vesico-Ureteric Reflux**

Backward flow of urine from the bladder into one or both ureters and sometimes the kidneys, increasing the risk of recurrent urinary infection and renal scarring.

### **Vesicovaginal and Ureterovaginal Fistula**

Abnormal connections between the urinary tract and vagina: a vesicovaginal fistula connects bladder to vagina, while a ureterovaginal fistula connects ureter to vagina. Both commonly cause continuous urine leakage and require imaging and reconstructive urologic or gynecologic care.

### **Vesicovaginal Fistula**

An abnormal connection between the bladder and vagina that causes continuous or frequent leakage of urine through the vagina. It may follow childbirth injury, pelvic surgery, radiation, or advanced infection and usually requires specialist repair.

### **Vesicular**

Describing or containing small, fluid-filled blisters. A vesicular rash may occur with viral infection, allergy, eczema, burns, or other conditions; its appearance alone does not establish the cause.

### **Vesicular Breathing**

The normal soft, gentle breath sound heard over most outer lung areas with a stethoscope. Changes in loudness or quality may suggest fluid, collapse, blockage, infection, or other lung disease.

### **Vesicular Palmoplantar Eczema**

An itchy eczema affecting the palms or soles with small, deep-seated fluid-filled vesicles, followed by peeling, scaling, or fissures. Triggers include sweating, irritants, allergy, stress, and fungal infection elsewhere; persistent disease may need specialist treatment.

### **Vesiculobullous Dermatoses**

A group of skin diseases that cause small or large fluid-filled blisters. Causes include infection, allergy, autoimmune disease, inherited conditions, medicines, and injury; examination and sometimes

a biopsy identify the cause.

### **Vessel**

A vessel is a tube carrying blood, lymph, air, urine, bile, or another fluid through the body, depending on context.

### **Vestibular**

Relating to the vestibular system of the inner ear and brain, which detects head motion and position and helps maintain balance and gaze stability.

### **Vestibular Disorder**

A problem affecting the inner-ear balance organs or the brain pathways that process balance signals. It may cause dizziness, vertigo, unsteadiness, nausea, abnormal eye movements, or difficulty walking. Causes include inner-ear disease, infection, head injury, medicines, migraine, reduced blood flow, or disorders of the brain or nerves.

#### **Alternate Names:**

Vestibular Balance Disorder

### **Vestibular Migraine**

Repeated attacks of dizziness or vertigo related to migraine, with or without headache. Episodes may include sensitivity to movement, light, sound, or visual patterns; other inner-ear and brain causes must be considered.

### **Vestibular Neuritis**

Sudden inflammation or dysfunction of the balance nerve, causing severe spinning, nausea, vomiting, and unsteadiness that may last for days. Hearing is usually unaffected, but urgent assessment may be needed to exclude a stroke.

### **Vestibular Papillomatosis**

A harmless normal variation in which several soft, small, finger-like bumps occur on the inner vulvar folds. It can resemble genital warts but is not caused by an infection and usually needs no treatment.

### **Vestibular Rehabilitation**

Exercise-based therapy that improves balance, gaze

---

stability, and adaptation in people with vestibular disorders or dizziness.

### **Vestibular Schwannoma**

A benign, slow-growing tumor of the vestibular portion of the eighth cranial nerve. See Acoustic Neuroma.

### **Vestibular System**

The inner-ear sensors and connected brain pathways that detect head movement and position and help maintain balance, posture, and steady vision. Problems can cause vertigo, unsteadiness, nausea, or blurred vision during movement.

### **Vestibule**

A balance area of the inner ear between the cochlea and semicircular canals. It contains sensors that detect head position and straight-line movement and send this information to the brain.

### **Vestibulocochlear Nerve**

The eighth cranial nerve, carrying information for hearing and balance from the inner ear to the brain.

### **Vestibulodynia**

Persistent burning, stinging, or tenderness at the entrance to the vagina, often triggered by touch, penetration, or inserting a tampon. Examination rules out infection and other causes; care may include pelvic-floor therapy, pain treatment, and support for sexual and emotional health.

#### **Alternate Names:**

Provoked Vestibulodynia; Vulvar Vestibulitis Syndrome; PVD

### **Vestibulo-Ocular Reflex**

An automatic response that moves the eyes opposite to head movement so vision stays steady while a person turns or walks. Problems with it can cause blurred vision, dizziness, or imbalance.

### **Vestigial**

Describing a structure or feature that has become greatly reduced through evolution and has little or

no apparent original function.

### **VEXAS Syndrome**

A serious inflammatory illness that begins in adulthood because of a gene change acquired in blood-forming cells. It can cause persistent fever, painful joints, skin rashes, inflamed blood vessels or cartilage, anemia, and other blood abnormalities.

### **V-Fib**

Alternate name; see *Ventricular fibrillation*.

### **Viable**

Capable of surviving or developing under suitable conditions. In clinical contexts it may describe a viable pregnancy, tissue, organism, cell, or treatment option.

### **Vibration Problem**

Symptoms or injury caused by repeated exposure to vibrating tools or equipment, which may affect nerves, blood vessels, muscles, and joints in the hands or arms.

### **Vibrator**

A device that produces mechanical vibration and may be used for sexual stimulation, therapeutic massage, or selected rehabilitation purposes.

### **Vibrio**

A genus of curved, Gram-negative bacteria; species include *V. cholerae*, which causes cholera, and *V. vulnificus*, which can cause severe wound or blood-stream infection.

### **Vibrio Cholerae**

A bacterium that causes cholera, a severe diarrheal illness spread mainly through contaminated water or food. It can cause sudden, large-volume watery diarrhea, dehydration, shock, and death without rapid fluid replacement.

### **Vibrio Infection**

Infection caused by *Vibrio* bacteria, acquired through contaminated seafood or exposure of

---

wounds to warm seawater. It may cause watery diarrhea, wound infection, or rapidly worsening bloodstream infection, especially in people with liver disease or weakened immunity. Severe wound pain, blistering, fever, or shock can occur.

**Alternate Names:**

Vibriosis; Vibrio Bacteremia

### **Vibrio Parahaemolyticus Infection**

An acute halophilic bacterial gastroenteritis acquired through ingestion of raw or undercooked marine seafood, particularly shellfish. It typically manifests with explosive watery diarrhea, abdominal cramps, nausea, vomiting, low-grade fever, and headache within 12 to 24 hours of exposure, usually resolving spontaneously within 3 to 5 days without antibiotics.

**Alternate Names:**

Vibrio Parahaemolyticus Gastroenteritis; Halophilic Vibrio Infection

### **Vibrios**

Comma-shaped bacteria, usually Gram-negative, that include species capable of causing cholera and wound or gastrointestinal infections.

### **Vibrio vulnificus Infection**

Infection caused by a saltwater-associated bacterium that can produce rapidly progressive wound infection, sepsis, or severe gastroenteritis. Risk is higher with liver disease, iron overload, or immune suppression; raw oysters and seawater exposure are important sources.

### **Vibrio Vulnificus Infections**

Infections caused by *Vibrio vulnificus*, a gram-negative bacterium found in warm seawater and raw seafood. It can cause wound infection, bloodstream infection, or gastroenteritis, with particularly severe disease in people with chronic liver disease or other risk factors.

### **Video-Assisted Thoracoscopic Surgery**

A chest operation performed through several small openings using a camera and long instruments rather than a large incision between the ribs. It may be

used for lung or pleural biopsy, removing part of a lung, or other chest procedures and often allows faster recovery, though bleeding, air leakage, infection, or conversion to open surgery can occur.

**Alternate Names:**

VATS; Video Thoracoscopy; Thoracoscopic Surgery

### **Videofluoroscopic Swallow Study**

A moving X-ray test that records swallowing after small amounts of food or liquid containing contrast are taken. It shows whether food enters the airway, where swallowing is delayed, and which textures or techniques may be safer.

### **Video Laryngoscope**

An airway device with a small camera that shows the entrance to the windpipe on a screen. It helps clinicians place a breathing tube, especially when a direct view is difficult, but does not remove the risks of the procedure.

### **Video Laryngoscopy**

A technique for viewing the entrance to the windpipe with a camera on a blade connected to a screen. It can help clinicians place a breathing tube when a direct view is difficult.

### **Videostroboscopy**

A voice examination that uses a camera and flashing light to create a slow-motion view of the vocal cords as they vibrate. It helps investigate hoarseness, voice strain, and vocal-cord injury.

### **Vildagliptin**

An oral dipeptidyl-peptidase-4 inhibitor used in some countries for type 2 diabetes, alone or with other medicines. It improves glucose-dependent insulin secretion; liver function and pancreatitis symptoms may require attention.

### **Villi**

Tiny finger-like projections lining the small intestine that greatly increase the area available to absorb nutrients. Similar projections in the placenta help exchange substances between the pregnant person and fetus.



---

## **Villous Adenoma**

A high-risk subtype of neoplastic colonic adenomatous polyp characterized histologically by long, finger-like, frond-like epithelial projections covering more than 75

### **Alternate Names:**

Villous Colonic Polyp; Papillary Adenoma

## **Villous Atrophy**

Flattening or loss of the tiny projections lining the small intestine. This reduces nutrient absorption and may occur in celiac disease, some infections, immune disorders, or other intestinal conditions.

## **Villus**

A small finger-like projection, especially one of the intestinal mucosa that increases absorptive surface area; chorionic villi mediate exchange between placenta and fetus.

## **Villus Atrophy**

The histopathological flattening, blunting, and loss of the microscopic finger-like villi lining the small intestinal mucosa, accompanied by crypt hyperplasia and intraepithelial lymphocytosis, classically observed in celiac disease, tropical sprue, and autoimmune enteropathy.

## **Vinca Alkaloids**

Cancer medicines including vincristine and vinblastine that disrupt microtubules, structures needed for cell division. They can treat several cancers but may cause nerve damage, low blood counts, constipation, or other toxicity.

## **Vincent Angina**

An acute necrotizing infection of the gums and oropharynx associated with fusobacteria and spirochetes. It causes painful ulceration, bleeding, foul breath, and gingival necrosis and requires dental or medical treatment.

## **Vincent Disease**

Acute necrotizing ulcerative gingivitis associated with fusospirochetal bacteria, causing painful bleed-

ing gums, ulceration, and foul breath.

## **Vincent Infection**

Vincent infection is acute necrotizing ulcerative gingivitis caused by mixed oral bacteria, producing painful bleeding gums, ulcers, and foul breath.

## **Vincent's Angina**

Acute necrotizing ulcerative gingivitis or tonsillitis associated with fusiform bacteria and spirochetes, causing painful ulcers, bleeding, and fetid breath.

## **Vincent Stomatitis**

A painful necrotizing infection of the gums caused by mixed anaerobic bacteria, producing foul breath, bleeding, ulceration, and gray necrotic tissue; it is also called acute necrotizing ulcerative gingivitis.

## **Vinorelbine**

A chemotherapy medicine that interferes with cell division. It is used for selected lung and breast cancers and can cause low blood counts, infection, constipation, nerve symptoms, nausea, or irritation if it leaks outside a vein.

## **Viomycin**

An older aminoglycoside antibiotic related to capreomycin that was used against tuberculosis and other mycobacterial infections; kidney and hearing toxicity limit its use.

## **VIPoma**

A rare functioning neuroendocrine tumor arising from the non-beta islet cells of the pancreas (or occasionally extra-pancreatic sympathetic ganglia in children) that hypersecretes vasoactive intestinal peptide (VIP), causing refractory secretory diarrhea and hypokalemia.

### **Alternate Names:**

Vasoactive Intestinal Peptide Tumor; Verner-Morrison Neoplasm

## **VIPoma Syndrome**

A rare neuroendocrine tumor syndrome arising predominantly from the non-beta islet cells of the

---

pancreas that hypersecretes vasoactive intestinal peptide (VIP). Clinically characterized by massive, secretory, watery diarrhea persisting during fasting, severe hypokalemia, and hypochlorhydria or achlorhydria (the WDHA syndrome), accompanied by dehydration and metabolic acidosis.

**Alternate Names:**

WDHA Syndrome; Verner-Morrison Syndrome; Pancreatic Cholera

**Viraemia**

The presence of viruses in the bloodstream. It may be brief or long-lasting and can allow the infection to spread from its original site to other tissues.

**Viral**

Caused by or relating to a virus. Viruses enter living cells to multiply and can cause illnesses ranging from mild colds to serious brain, lung, liver, or bleeding diseases; testing and treatment depend on the specific virus.

**Viral Antibody**

A protective protein made by the immune system after infection or vaccination that recognizes a virus. Blood tests for these antibodies may show past exposure, immune response, or selected current infections, depending on timing and test design.

**Viral Arthritis**

Joint inflammation caused by or associated with a viral infection. It may cause sudden pain, swelling, and stiffness in several joints and often improves as the infection resolves, although some viruses can cause prolonged symptoms.

**Viral Assembly**

The stage of viral reproduction when newly made viral genetic material and proteins are put together to form complete new virus particles.

**Viral Attachment**

The first step of infection, when proteins on a virus bind to matching molecules on a host cell. This binding helps determine which species, tissues, or cell types the virus can infect.

**Viral Carcinogenesis**

Cancer development caused or promoted by a viral infection. Some viruses alter cell growth directly, weaken immune protection, or cause long-lasting inflammation; examples include human papillomavirus, hepatitis B and C viruses, and Epstein-Barr virus.

**Viral Conjunctivitis**

Infection of the thin lining over the eye and inside the eyelids, usually causing redness, watering, irritation, and discharge. It spreads easily through contact with infected secretions or contaminated hands and surfaces.

**Viral Culture**

Growth of a virus in susceptible cell cultures to identify an infectious agent. It can be useful for selected viruses, but molecular and antigen-based tests are often faster or more sensitive; the appropriate method depends on the suspected infection and specimen.

**Viral Encephalitis**

Inflammation of the brain caused by a virus. It may cause fever, headache, confusion, behavior change, seizures, weakness, or reduced consciousness. Severe cases can be life-threatening.

**Viral Entry**

The process by which a virus gets inside a host cell and releases its genetic material so the cell can make more virus.

**Viral Exanthem**

A widespread rash caused by a viral infection. It may occur with fever, tiredness, sore throat, cough, diarrhea, or other symptoms; the rash pattern and overall illness help determine the cause.

**Viral Gastroenteritis**

Alternate name; see *Stomach Flu*.

**Viral Haemorrhagic Fever**

A severe illness caused by certain viruses that can

---

damage blood vessels, disturb clotting, and cause bleeding, shock, or organ failure. The way it spreads and its treatment depend on the virus.

### **Viral Hemorrhagic Fevers**

Severe illnesses caused by certain viruses that can damage blood vessels, disrupt clotting, and lead to bleeding, shock, or organ failure. Some spread easily and may require special infection-control measures.

#### **Alternate Names:**

VHF; Hemorrhagic Fevers; Viral Hemorrhagic Disease

### **Viral Hepatitis**

Liver inflammation caused by hepatitis viruses A, B, C, D, or E. Hepatitis A and E usually spread through contaminated food or water; B, C, and D can spread through blood or body fluids and may become long-term infections that damage the liver.

#### **Alternate Names:**

Infectious Hepatitis

### **Viral Hepatitis Types**

Five viruses that can inflame the liver. They differ in how they spread, whether infection can become long-term, the complications they cause, and whether vaccination or curative treatment is available.

### **Viral Immune Evasion**

Ways viruses avoid or weaken the body's immune defenses, allowing them to survive, multiply, or return after infection. These methods can make infections harder to clear and vaccines or treatments less effective.

### **Viral Infection**

An illness caused by a virus entering body cells and making copies of itself. Viruses can damage cells directly or cause inflammation as the immune system responds. Viral infections range from mild illnesses such as colds and cold sores to serious diseases such as influenza, COVID-19, hepatitis, and HIV. Some viruses remain inactive in the body and can become active again later, such as the chickenpox virus causing shingles.

#### **Alternate Names:**

Viral Infections; Viral Disease; Viral Illness

### **Viral Infections and Pregnancy**

Viral infections during pregnancy may affect maternal illness, fetal development, placenta, or newborn health. Evaluation considers the virus, gestational age, symptoms, immunity, exposure, testing, treatment, vaccination, and measures to reduce perinatal transmission.

### **Viral Infections of the Mouth**

Infections of oral skin or mucosa caused by viruses such as herpes simplex, varicella-zoster, enteroviruses, HPV, or EBV. They may cause blisters, ulcers, warts, sore throat, or systemic symptoms, and treatment depends on the virus and host risk.

### **Viral Latency**

A state in which a virus remains inside cells without actively making new virus particles. Some latent viruses can become active again, especially when the immune system is weakened or the body is under stress.

### **Viral Latency Mechanisms**

The ways some viruses remain inactive inside cells while avoiding immune detection. Reactivation can occur when immune defenses are weakened or after other triggers, depending on the virus.

### **Viral Load**

The amount of virus or viral genetic material in a body fluid or tissue, measured by a laboratory assay; trends can help assess disease activity or response to antiviral treatment.

### **Viral Meningitis**

Infection and inflammation of the coverings of the brain and spinal cord caused by a virus. It may cause fever, headache, neck stiffness, vomiting, light sensitivity, or confusion. Bacterial meningitis can cause similar symptoms and is more dangerous.

### **Viral Myocarditis**

Inflammation of the heart muscle caused by a vi-

---

ral infection or by the immune response after an infection. It may cause tiredness, chest discomfort, palpitations, heart failure, shock, or dangerous abnormal heart rhythms.

**Alternate Names:**

Infectious Viral Myocarditis; Acute Viral Myocarditis

**Viral Pathogenesis**

The ways a virus enters the body, infects and spreads through cells, and causes illness. Damage may come directly from infected cells or from the immune system's response to the infection.

**Viral Pericarditis**

Inflammation of the sac around the heart caused by a viral infection. It often follows a respiratory or stomach illness and may cause sharp chest pain that changes with breathing or position.

**Alternate Names:**

Acute Viral Pericarditis; Viral Pericardial Inflammation

**Viral Pharyngitis**

Pharyngeal inflammation caused by viral pathogens such as adenovirus, rhinovirus, coronavirus, Epstein-Barr virus, or influenza virus. Characterized by cough, rhinorrhea, conjunctivitis, and lack of prominent tonsillar exudates, resolving with supportive care.

**Alternate Names:**

Viral Sore Throat; Acute Viral Pharyngotonsillitis; Viral Upper Respiratory Pharyngitis

**Viral Pneumonia**

Infection and inflammation of lung tissue caused by a virus, producing cough, fever, chills, breathlessness, chest discomfort, fatigue, and variable impairment of oxygenation. Severe disease can cause respiratory failure or secondary bacterial infection.

**Viral Release**

The stage when newly made viruses leave an infected cell and spread to other cells. Some leave by pushing through the cell membrane, while others are released when the infected cell breaks apart.

**Viral Replication**

The process by which a virus uses an infected cell to copy its genetic material and make viral proteins. The exact steps differ among viruses, but the result is production of new virus particles.

**Viral Replication Cycle**

The steps a virus uses to make more copies inside a living cell. Usually the virus attaches to the cell, enters, releases its genetic material, makes new parts, assembles new viruses, and leaves to infect other cells.

**Viral Rhinitis**

Acute inflammation and edema of the nasal mucous membranes caused by respiratory viral pathogens, predominantly rhinoviruses, coronaviruses, respiratory syncytial virus, or parainfluenza viruses. Clinically recognized as acute viral coryza or the common cold, it manifests with clear anterior rhinorrhea, nasal congestion, sneezing, itchy throat, low-grade malaise, and temporary hyposmia, generally resolving without antimicrobial therapy within 7 to 10 days.

**Alternate Names:**

Acute Viral Rhinitis; Acute Coryza; Common Cold Rhinitis

**Viral Structure**

The parts of a virus: genetic material made of DNA or RNA inside a protective protein coat. Some viruses also have an outer fatty covering taken from the infected cell's membrane.

**Viral Uncoating**

The step after a virus enters a cell when its protective outer coat comes apart and releases the genetic material. The cell can then be directed to make new viral components.

**Virchow Node**

A firm, non-tender, pathologically enlarged left supraclavicular lymph node (Troisier sign) located in the left supraclavicular fossa, which receives abdominal lymphatic drainage via the thoracic duct and signals metastatic spread of abdominal malignancies

---

(particularly gastric adenocarcinoma).

**Alternate Names:**

Troisier Node; Left Supraclavicular Sentinel Node

**Virchow-Robin Spaces**

Small fluid-filled spaces around blood vessels as they pass through the brain. They are a normal part of the spaces that surround and support these vessels.

**Alternate Names:**

Perivascular Spaces

**Virchow's Triad**

Three factors that increase the chance of a blood clot forming: slow blood flow, injury to the inside of a blood vessel, and blood that clots too easily. These factors can contribute to clots in deep veins or clots that travel to the lungs.

**Alternate Names:**

Virchow Triad; Triad of Virchow; Thrombogenesis Triad

**Virchow Triad**

Three factors that increase the chance of a blood clot forming: injury to the inside of a blood vessel, slow blood flow, and blood that clots too easily.

**Alternate Names:**

Virchow's Triad; Thrombosis Pathophysiological Triad

**Viremia**

The presence of a virus in the bloodstream. It may be brief or persistent and can allow infection to spread to other organs. Laboratory tests may detect the virus itself or its genetic material.

**Viridans Streptococci**

Viridans streptococci are a group of bacteria commonly found in the mouth that can cause dental infections and, after entering the blood, infect heart valves.

**Virilisation**

The development of male secondary sexual characteristics in females, resulting from excessive androgen production or exposure.

**Virilism**

Development of masculinizing features in a female, usually from excess androgen exposure or production, such as hirsutism, deep voice, acne, and menstrual disturbance.

**Virilism Foetal**

Prenatal or neonatal masculinization of a genetically female fetus or infant due to excess androgen exposure. Causes include congenital adrenal hyperplasia, maternal androgen-producing disease, or placental and fetal steroidogenic disorders.

**Virilization**

Virilization is development of typically male secondary sexual features, such as hirsutism, deep voice, acne, increased muscle mass, or clitoromegaly, from androgen excess or androgen exposure.

**Virilize**

To virilize is to develop or cause typically male physical changes from androgen excess or treatment, including facial hair, voice deepening, acne, muscle change, or genital enlargement.

**Virology**

The study of viruses, including their structure, reproduction, spread, effects on the body, diagnosis, treatment, and prevention.

**Virotherapy**

Treatment that uses a virus, often modified, to attack cancer cells or deliver a useful gene into cells. It is an evolving field; the possible benefits, risks, and availability depend on the specific therapy and disease.

**Virtual Colonoscopy**

Virtual colonoscopy is CT colonography, which uses bowel preparation and gas or carbon-dioxide insufflation to create images of the colon; it can detect polyps and masses but cannot remove lesions or biopsy them.

---

## Virulence

How severely a microorganism can cause disease or tissue damage. It depends on the organism's harmful properties and on the person's immune defenses and other health factors.

## Virulence Factor

A feature of a microorganism that helps it survive in the body, avoid immune defenses, attach to cells, damage tissue, or cause disease. Examples include toxins, protective coatings, and enzymes.

## Virulent

Capable of causing severe disease or extensive harm. Virulence describes the harmfulness of a microorganism and is different from infectivity, which describes how easily it spreads or establishes infection.

## Virus

An infectious agent made of genetic material inside a protective protein coat, sometimes with an outer membrane. It can reproduce only inside living cells and may cause illness by damaging cells or triggering inflammation.

## Virus Activation

Virus activation is the switch from a dormant or inactive viral state to production of new viral components and particles.

## Virus Assembly

Virus assembly is the stage of the viral life cycle when viral genetic material and proteins are packaged into new virus particles.

## Virus Diseases

Illnesses caused by viruses. They range from mild infections of one area, such as a cold, to severe diseases affecting the lungs, liver, brain, blood, or several organs; prevention and treatment depend on the virus.

## Viruses

Viruses are infectious particles containing DNA or RNA within a protein coat, sometimes with a lipid

envelope; they replicate only inside host cells and cause disease through direct injury or immune responses.

## Virus Integration

Virus integration is insertion of viral genetic material into the DNA of a host cell, a process used by some viruses and important in certain cancers.

## Virus Protein

A virus protein is a protein made from viral genetic instructions that helps form the virus, enter cells, replicate, or evade immunity.

## Virus Receptor

A virus receptor is a molecule on a host cell that a virus binds to in order to attach to and enter the cell.

## Virus Replication

Virus replication is the process by which a virus copies its genetic material and makes proteins and new virus particles inside a host cell.

## Virus Uncoating

Virus uncoating is removal of the virus's protective outer layer after entry into a host cell, releasing its genetic material for replication.

## Viscera

The internal organs of the body, particularly those within the thoracic and abdominal cavities. The thoracic viscera include the heart, lungs, esophagus, trachea, and thymus.

## Visceral

Relating to the internal organs, especially those within the thorax or abdomen, rather than the body wall or limbs.

## Visceral Afferents

Sensory nerve fibers that carry information from internal organs to the spinal cord and brain. They convey signals related to stretch, chemical changes, inflammation, and internal pain and help regulate



---

autonomic and reflex responses.

### **Visceral Larva Migrans**

A syndrome caused by migration of animal nematode larvae through human tissues, most often from *Toxocara* species. It may produce fever, eosinophilia, liver lesions, or ocular and neurologic disease.

### **Visceral Leishmaniasis**

A severe, potentially fatal systemic protozoan infection caused by species of the *Leishmania* donovani complex (*Leishmania donovani* and *Leishmania infantum*) transmitted to humans by the bite of female phlebotomine sandflies. Characterized by the intracellular dissemination of amastigotes within the reticuloendothelial system, it presents with prolonged irregular fever, progressive massive splenomegaly, hepatomegaly, profound pancytopenia (anemia, leukopenia, and thrombocytopenia), hypergammaglobulinemia, and severe wasting, fatal in over 95% of untreated cases.

#### **Alternate Names:**

Kala-Azar; Black Fever; Dumdum Fever; Systemic Leishmaniasis

### **Visceromegaly**

Abnormal enlargement of one or more internal organs, such as the liver, spleen, or kidneys. It is a physical finding with causes that include infection, congestion, storage disease, inflammation, and cancer.

### **Viscus**

An internal organ, especially one in the chest or abdomen. The plural is viscera; examples include the heart, lungs, liver, stomach, and intestines.

### **Vision**

The sensory ability to perceive light and interpret visual information processed by the eyes and brain, enabling recognition of shapes, colors, depth, and motion. Normal vision depends on the proper function of the cornea, lens, retina, and optic nerve, as well as intact neural pathways to the visual cortex.

#### **Alternate Names:**

Eyesight; Sight; Visual Perception

### **Vision And Hearing**

Two major senses used to understand and communicate with the world. Problems may develop separately or together because of aging, illness, injury, medicines, or inherited conditions and can affect learning, safety, communication, and daily activities.

### **Vision Impairment**

Reduced visual acuity or visual-field function that affects daily activities. Causes include refractive error, cataract, glaucoma, retinal disease, optic-nerve injury, and neurologic disease. Evaluation and treatment depend on the cause; rehabilitation and visual aids may improve function.

### **Vision Loss**

A decrease in the ability to see that cannot be fully corrected with eyeglasses, contact lenses, or medical treatment. It ranges from partial impairment to complete blindness and can result from conditions such as macular degeneration, glaucoma, diabetic retinopathy, cataracts, and traumatic injury.

### **Vision Problems**

Changes in visual acuity, field, color, or clarity caused by refractive error, eye disease, neurologic disease, or systemic illness. Sudden vision loss is an emergency.

### **Vision Receptors**

Light-sensing cells in the retina, mainly rods and cones, that convert light into nerve signals. Rods support dim-light vision, while cones provide color and sharp detail in brighter conditions.

### **Vision Screening**

Brief checks used to find possible eye or vision problems before symptoms are obvious. Newborn checks may detect cataracts or other serious disorders; screening in young children looks for poor vision, misaligned eyes, or risks for reduced vision and leads to a full eye examination when needed.

### **Vision Test**

An examination that measures visual acuity, fields,

---

color, alignment, or eye structures to detect refractive error, retinal disease, optic-nerve dysfunction, glaucoma, or other causes of visual impairment.

### **Visual Acuity**

How clearly a person can see fine details, usually measured by reading letters or symbols at a set distance. It can be reduced by focusing problems, eye disease, or disorders affecting the visual nerves or brain.

### **Visual Acuity Test**

A test measuring the smallest standardized detail a person can resolve at a specified distance, usually reported as a fraction or decimal.

### **Visual Agnosia**

Inability to recognize familiar objects, faces, or symbols by sight even though the eyes may see adequately. It usually results from damage to brain areas that interpret vision, such as after a stroke or head injury.

### **Visual Evoked Potential**

A test that records the brain's electrical response to flashes or changing visual patterns. It helps assess the nerve pathways that carry visual signals from the eyes to the brain.

#### **Alternate Names:**

VEP; Visual Evoked Response (VER)

### **Visual Field**

The entire area that can be seen while the eyes remain fixed in one position. Visual-field loss may result from retinal, optic-nerve, or brain lesions and is assessed by perimetry.

### **Visual Field Defect**

Loss or change in part of the area a person can see while looking straight ahead. It may result from disease of the retina, optic nerve, or brain, and the pattern can help identify where the problem is.

### **Visual Field Perimetry**

A test that maps how well a person sees in different

parts of the visual field while looking at a fixed point. It can detect and monitor missing areas caused by eye-nerve, retinal, or brain disease.

### **Visual Fields**

The total area that can be seen while the eyes are fixed in one position; defects may indicate retinal, optic-nerve, or brain disease.

### **Visual Field Test**

A test that measures central and side vision and maps areas that are missing or less sensitive.

### **Visual Impairment**

Reduced vision that cannot be fully corrected by ordinary glasses or contact lenses. Causes include refractive error, cataract, glaucoma, retinal disease, optic-nerve injury, and disorders of the brain or nervous system.

### **Visualized Laser-Assisted Prostatectomy**

A procedure that uses a laser, guided by a camera, to remove or vaporize prostate tissue blocking urine flow. It is used for some people with an enlarged prostate and may cause bleeding, infection, scar narrowing, or temporary urinary and sexual problems.

#### **Alternate Names:**

VLAP

### **Visual Neglect**

A brain disorder in which a person fails to notice or respond to things on one side, even though the eyes may work normally. It often follows a stroke affecting the right side of the brain and causes neglect of the left side.

#### **Alternate Names:**

Hemispatial Neglect; Spatial Inattention

### **Vital**

Essential for life or relating to vital physiologic functions. Vital signs commonly include temperature, pulse, respiratory rate, blood pressure, and oxygen saturation.

---

## Vital Capacity

The greatest amount of air a person can breathe out after filling the lungs as fully as possible. It is measured during lung-function testing and may be reduced by lung disease, weak breathing muscles, or restriction of chest movement.

## Vital Signs

Basic measurements of physiologic status, commonly including temperature, pulse, respiratory rate, blood pressure, and oxygen saturation. Values vary with age, activity, illness, and measurement conditions; trends and clinical context are important.

## Vitamin A

A fat-soluble vitamin needed for vision, immune function, skin and mucosal health, and growth; deficiency causes night blindness, while excess can injure the liver and developing fetus.

### Alternate Names:

Retinol; Retinoic Acid

## Vitamin A Blood Test

A blood test measuring circulating retinol, used to assess suspected vitamin A deficiency or toxicity in appropriate clinical settings.

## Vitamin A Deficiency

A nutritional disorder resulting from inadequate intake or absorption of vitamin A (retinol), causing dry eyes (xerophthalmia), night blindness, dry skin, and weakened immunity. Severe deficiency leads to corneal ulceration and is the leading cause of preventable childhood blindness worldwide; clinical eye stages are categorized using the WHO xerophthalmia classification.

### Alternate Names:

Hypovitaminosis A; Retinol Deficiency; Vitamin A Lack

## Vitamin A: Retinal

Vitamin A (retinol) is a fat-soluble vitamin essential for vision, immune function, cell differentiation, and reproduction. In the eye, retinol is oxidized to retinal (retinaldehyde), which binds to opsin in the

photoreceptor outer segments forming rhodopsin (in rods) and photopsins (in cones).

## Vitamin A Toxicity

Harm caused by taking too much vitamin A or related medicines. It may cause nausea, vomiting, headache, dry skin, hair loss, liver injury, high calcium levels, and serious fetal abnormalities during pregnancy.

## Vitamin B

Vitamin B refers to a group of water-soluble vitamins that support energy metabolism, blood-cell formation, nerve function, and other processes. The group includes several distinct vitamins, not one single substance.

## Vitamin B1

Thiamine, a water-soluble vitamin that acts as a co-factor in carbohydrate and amino-acid metabolism and supports nerve function. Severe deficiency can cause beriberi or Wernicke encephalopathy.

### Alternate Names:

Thiamine; Aneurin

## Vitamin B12

A vitamin needed for red-cell formation, DNA production, and normal nerve function. Deficiency can cause anemia, numbness, balance problems, or cognitive changes.

### Alternate Names:

Cobalamin; Cyanocobalamin

## Vitamin B12: Cobalamin

Alternate name; see *Vitamin B12*.

## Vitamin B12 Deficiency

Too little vitamin B12, which can cause large abnormal red cells, tiredness, numbness, balance problems, or nerve damage. Causes include poor intake, poor absorption, autoimmune disease, stomach surgery, and medicines.

## Vitamin B12 Deficiency Anemia

Anemia caused by too little vitamin B12 or poor

---

absorption of it. It can cause tiredness, pale skin, a sore tongue, numbness, balance problems, and permanent nerve damage if untreated for a long time.

**Alternate Names:**

Vitamin B12 Deficiency Anaemia

**Vitamin B12 Deficiency In Older Adults**

Deficiency caused by inadequate intake, malabsorption, autoimmune intrinsic-factor deficiency, gastric surgery, or medications. It may cause megaloblastic anemia, neuropathy, cognitive changes, gait impairment, and glossitis; neurologic injury can occur without anemia.

**Vitamin B12 Level**

The amount of vitamin B12 measured in blood; results are interpreted with symptoms and related tests because serum levels may not fully reflect tissue deficiency.

**Vitamin B1: Thiamine**

Alternate name; see *Vitamin B1*.

**Vitamin B2**

Riboflavin, a water-soluble vitamin that helps cells produce energy and supports healthy skin, eyes, and the lining of the mouth. Deficiency can cause cracks at the mouth corners, a sore tongue, skin changes, or anemia.

**Alternate Names:**

Riboflavin; Lactoflavin

**Vitamin B2 Riboflavin**

Alternate name; see *Riboflavin*.

**Vitamin B3**

A vitamin that helps cells use food for energy and supports the skin and nervous system. Severe deficiency causes pellagra, with diarrhea, skin inflammation, confusion, and other neurologic problems.

**Alternate Names:**

Nicotinic Acid; Nicotinamide

**Vitamin B5**

A water-soluble vitamin needed to help the body use fats, carbohydrates, and proteins and make hormones and other substances. It is found in many foods, so deficiency is rare; very large supplement doses may cause stomach upset.

**Alternate Names:**

Pantothenate; Calcium Pantothenate

**Vitamin B6**

A group of related vitamins needed for amino-acid metabolism, red-cell production, and nerve function. Excess supplements can cause sensory nerve damage.

**Alternate Names:**

Pyridoxine; Pyridoxal Phosphate

**Vitamin B6: Pyridoxine**

Alternate name; see *Vitamin B6*.

**Vitamin B7**

A vitamin that helps the body use fats, carbohydrates, and proteins. Deficiency is uncommon but may cause hair loss, skin rash, or neurologic symptoms; large doses can interfere with some laboratory tests.

**Alternate Names:**

Vitamin H; Coenzyme R

**Vitamin B9**

A vitamin needed for DNA production and healthy red blood cells. Deficiency can cause large-cell anemia and, during pregnancy, increases the risk of neural-tube defects; folic acid is recommended before conception and early in pregnancy.

**Alternate Names:**

Folic Acid; Pteroylglutamic Acid

**Vitamin B Complex**

A group of eight water-soluble vitamins that act mainly as coenzymes or their precursors in cell metabolism. The group includes thiamine (B1), riboflavin (B2), niacin (B3), pantothenic acid (B5), pyridoxine (B6), biotin (B7), folate (B9), and cobalamin (B12), which together support energy

---

metabolism, red-blood-cell formation, and nervous-system function. These vitamins occur together in whole grains, meat, eggs, and legumes; individual deficiencies cause distinct disorders, and combined supplements can prevent or correct deficiency. The sequence has gaps because early candidate substances were later excluded as nonessential: B8 (inositol) is synthesized by the body, B4 (choline, adenine) is not a true vitamin, and the provisional labels B10 and B11 were abandoned once their activity was identified as folate or para-aminobenzoic acid.

**Alternate Names:**

B Vitamins; Vitamin B Group

**Vitamin C**

A water-soluble vitamin needed for collagen formation, wound healing, immune function, antioxidant activity, and absorption of iron from plant foods. Severe deficiency causes scurvy, with bleeding gums, bruising, poor wound healing, and weakness.

**Alternate Names:**

Ascorbic Acid

**Vitamin C Deficiency**

A lack of vitamin C, a water-soluble vitamin needed for collagen formation, wound healing, iron absorption, and antioxidant activity. It follows prolonged low intake and, when severe, causes scurvy, with bleeding gums, easy bruising, poor wound healing, joint pain, and weakness. Replacement and dietary correction reverse the deficiency.

**Vitamin D**

Vitamin D supports calcium and phosphate balance, bone mineralization, muscle function, and immune processes; deficiency causes rickets in children and osteomalacia or bone loss in adults.

**Vitamin D3**

Cholecalciferol, a form of vitamin D made in skin and found in some foods and supplements. It helps the body absorb calcium and maintain healthy bones.

**Vitamin D: Calcitriol**

Alternate name; see *Calcitriol*.

**Vitamin D Deficiency**

Too little vitamin D for the body's needs. It can result from low intake or sunlight, poor absorption, or liver or kidney disease and may weaken bones, cause muscle weakness, or lead to rickets in children.

**Vitamin Deficiency Anemia**

Anemia caused by inadequate vitamin B-12, folate, or other nutrients needed for normal red blood cell production.

**Vitamin E**

Vitamin E is a group of fat-soluble compounds with antioxidant activity. It is found in foods such as nuts, seeds, and vegetable oils; high-dose supplements can increase bleeding risk and interact with medicines.

**Alternate Names:**

Tocopherol; Alpha-Tocopherol

**Vitamin E Deficiency**

Deficiency of a fat-soluble antioxidant vitamin, usually caused by fat-malabsorption disorders or rare inherited conditions. Severe deficiency can cause peripheral neuropathy, ataxia, muscle weakness, retinal dysfunction, or hemolytic anemia.

**Vitamin E: Tocopherol**

Alternate name; see *Vitamin E*.

**Vitamin K**

A fat-soluble vitamin required to activate several blood-clotting proteins and support bone metabolism; the main dietary form is phylloquinone, found especially in leafy green vegetables and some vegetable oils. Deficiency can cause bleeding and is treated or prevented with replacement.

**Vitamin K Deficiency**

Too little vitamin K for normal production of clotting factors, causing an increased bleeding tendency. It can result from poor intake, malabsorption, liver disease, or certain medicines; newborns receive vi-

---

tamin K prophylaxis.

### **Vitamin K Deficiency Bleeding**

Bleeding in a newborn or young infant caused by inadequate vitamin K-dependent clotting activity. It may occur early, classically, or late and can involve the brain or gastrointestinal tract.

#### **Alternate Names:**

VKDB; Neonatal Hypoprothrombinemia

### **Vitamin K: Phylloquinone**

Alternate name; see *Vitamin K*.

### **Vitamin K Prophylaxis**

Giving vitamin K to a newborn soon after birth to prevent bleeding caused by vitamin K deficiency.

### **Vitamins**

Organic nutrients needed in small amounts for normal metabolism, growth, and tissue maintenance; some are water-soluble and others fat-soluble, and both deficiency and excess can cause disease.

### **Vitamin Toxicity**

Harm from excessive vitamin intake, most often involving fat-soluble vitamins that accumulate in the body. Vitamin A can cause headache, liver injury, and bone effects; vitamin D can cause hypercalcemia; and high-dose vitamin E can increase bleeding risk. Stop the source and treat the affected organ or metabolic abnormality.

### **Vitiligo**

Vitiligo is a condition in which areas of skin lose pigment because melanocytes are destroyed or stop functioning. It is not contagious, and treatment may include camouflage, topical medicines, or light therapy.

### **Vitrectomy**

Eye surgery that removes some or all of the vitreous, the clear gel that fills the middle of the eye. Small instruments are inserted through the wall of the eye to remove cloudy gel, blood, scar tissue, or a foreign object and to repair problems of the retina.

It may be used for retinal detachment, diabetic eye disease, macular hole, macular pucker, severe eye injury, or serious eye infection. The removed gel may be replaced with a salt solution, air or gas, or silicone oil.

### **Vitreous**

Relating to the transparent gel filling most of the space between the lens and retina, or describing a glass-like appearance. Vitreous disorders can cause floaters, hemorrhage, detachment, or retinal traction.

### **Vitreous Body**

The transparent gel filling the large space between the lens and retina. It helps maintain the shape of the eye and transmits light; age-related liquefaction can cause floaters or contribute to retinal traction.

#### **Alternate Names:**

Vitreous Humor; Corpus Vitreum

### **Vitreous Detachment**

Separation of the vitreous gel from the retina, usually related to aging and often causing new floaters or flashes. Sudden symptoms can occur with a retinal tear or detachment.

### **Vitreous Hemorrhage**

Bleeding into the vitreous cavity that causes sudden painless floaters, cobwebs, haze, or loss of vision. Common causes include proliferative diabetic retinopathy, retinal tear or detachment, trauma, and posterior vitreous separation.

### **Vitreous Humor**

The clear gel that fills the space between the lens and retina. It helps maintain the shape of the eye and transmits light to the retina.

### **VKORC1 Polymorphism**

An inherited variation in the vitamin K epoxide reductase complex subunit 1 gene that can alter sensitivity to warfarin. Genotype may contribute to dose selection, but clinical factors, interacting medicines, diet, and INR monitoring remain important.



---

## **VLDL Test**

A blood test that measures or estimates very-low-density lipoprotein, a particle carrying triglycerides. The result may help assess cardiovascular risk but is interpreted with the full cholesterol and triglyceride profile.

## **VO2 Max**

The greatest amount of oxygen the body can use during intense exercise. It is a measure of aerobic fitness and is influenced by the heart, lungs, blood, muscles, age, health, and training.

### **Alternate Names:**

Maximal Oxygen Consumption; Maximal Aerobic Capacity

## **Vocal-Cord Dysfunction**

Older term; see *Inducible Laryngeal Obstruction*.

## **Vocal Cord Nodules**

Benign, usually paired thickened areas of the vocal folds caused by repeated voice strain or overuse. They can cause hoarseness, vocal fatigue, and a rough or breathy voice.

## **Vocal Cord Paralysis**

Inability of one or both vocal folds to move normally, often because of nerve damage. It may cause hoarseness, weak cough, choking, or difficulty protecting the airway; sudden breathing trouble is an emergency.

### **Alternate Names:**

Vocal Fold Paralysis

## **Vocal Cord Paresis**

Weakness or partial paralysis of one or both vocal cords, often from injury to the nerves that control them. It can cause a weak or breathy voice, voice fatigue, coughing during eating, or aspiration.

## **Vocal Cord Polyp**

A usually noncancerous growth on a vocal fold, often after heavy or sudden voice use. It can cause persistent hoarseness, a rough voice, or voice fatigue

and is assessed by examining the larynx.

## **Vocal Cords**

Vocal cords, or vocal folds, are bands of tissue in the larynx that vibrate as air passes between them to produce the voice. Swelling, nodules, or injury can cause hoarseness.

## **Vocal-Fold Nodules**

Alternate name for vocal cord nodules, benign thickened areas of the vocal folds that can cause persistent hoarseness.

## **Vocal Fold Paralysis**

Alternate name; see *Vocal cord paralysis*.

## **Vocal-Fold Polyps**

Alternate name for vocal cord polyps, usually non-cancerous growths on the vocal folds that can cause persistent hoarseness or voice fatigue.

## **Vocal Folds**

Two folds inside the voice box that vibrate as air passes between them to make sound. They also close during swallowing to protect the lungs from food and liquid.

## **Vocal Hemorrhage**

Vocal hemorrhage is bleeding into a vocal fold, often after sudden or forceful voice use. It can cause abrupt hoarseness.

## **Vocal Resonance**

The sound of a person's voice heard through the chest with a stethoscope while they speak. Changes in the sound can provide clues to pneumonia, fluid around the lung, a collapsed lung, or other lung disease.

## **Vogt-Koyanagi-Harada Disease**

A rare, multisystem, cell-mediated autoimmune disorder directed against melanocyte-associated antigens (such as tyrosinase), occurring predominantly in pigmented ethnic populations. Characterized by chronic bilateral granulomatous panuveitis with

---

exudative retinal detachments, accompanied by sensorineural hearing loss, tinnitus, vitiligo, poliosis (whitening of eyelashes and hair), and aseptic meningitis.

**Alternate Names:**

VKH Syndrome; Uveomeningoencephalitic Syndrome

**Vogt-Koyanagi-Harada Syndrome**

A multisystem autoimmune inflammatory disorder mediated by T-cell responses against melanocyte-associated antigens, characterized by bilateral chronic granulomatous panuveitis, exudative retinal detachments, meningismus (CSF pleocytosis), sensorineural dysacusis, poliosis, and patchy vitiligo.

**Alternate Names:**

VKH Syndrome; Uveomeningoencephalitic Syndrome

**Vohwinkel Syndrome**

A rare inherited palmoplantar keratoderma causing thick skin on the palms and soles, constricting bands around fingers or toes, and sometimes hearing loss or other features. Severe constriction can threaten circulation and requires specialist care.

**Voice**

Sound produced by vibration of the vocal folds as air passes through the larynx, then shaped by the throat, mouth, and nose for speech.

**Voice And Speech**

The products of complex coordinated activity of the respiratory, laryngeal, and articulatory systems. Communication produced by coordinated breathing, voice-box vibration, and movements of the mouth, tongue, and other speech structures.

**Voice Box**

The larynx, an organ in the neck containing the vocal folds and forming part of the airway. It produces voice and helps protect the lower respiratory tract during swallowing.

**Voice Disorders**

Conditions affecting voice quality, pitch, loudness, or ease of speaking because of problems involving the vocal folds, airway, nerves, or voice use.

**Alternate Names:**

Dysphonia; Vocal Cord Dysfunction

**Void**

To empty the urinary bladder by urinating; in other medical contexts, void may mean an empty space or absence of a substance.

**Voiding Cystourethrogram**

An X-ray test that fills the bladder with contrast through a catheter and records the bladder and urethra while the person urinates. It can show urine flowing backward toward the kidneys, blockage, leakage, or structural problems, especially in children with selected urinary infections.

**Volar**

Relating to the palm of the hand or sole of the foot, especially the surface opposite the dorsum.

**Volatile Anesthetics**

Inhaled medicines used to produce and maintain unconsciousness during some operations. They can lower blood pressure and breathing and rarely cause a dangerous rise in body temperature in susceptible people.

**Alternate Names:**

Inhalational Anesthetics; Inhalation General Anesthetics

**Volkmann Contracture**

A permanent, severe ischemic contracture of the forearm flexor musculature leading to a claw-like hand deformity, resulting from untreated acute compartment syndrome of the forearm following supracondylar humeral fractures, crush injuries, or tight casting.

**Alternate Names:**

Volkmann's Ischemic Contracture; Ischemic Myodystrophy

**Alternate Names:**

---

Volkmann contracture; Volkmann's ischaemic contracture; Volkmann claw deformity.

### **Volume Of Distribution**

A calculated measure of how widely a medicine appears to spread through the body compared with the amount left in the blood. It helps guide dosing but does not represent a real space inside the body.

### **Volume Overload**

Excess total body sodium and water, causing peripheral or pulmonary oedema; common causes include heart failure, kidney failure, liver disease, and excessive intravenous fluid, and treatment depends on the cause and severity.

### **Voluntary**

Under conscious control or performed by choice, such as voluntary movement. Voluntary action is distinguished from reflex, autonomic, or involuntary activity.

### **Voluntary Psychiatric Admission**

Admission to a hospital or psychiatric facility with the person's agreement for assessment, treatment, or safety support. Legal rules about consent, leaving, capacity, and conversion to involuntary status vary by jurisdiction.

### **Volutrauma**

Lung injury caused by overstretching the lungs, usually when a breathing machine delivers breaths that are too large. Careful ventilator settings can reduce this risk.

### **Volvulus**

Twisting of a loop of intestine that blocks passage through the bowel and may cut off its blood supply. It can cause severe pain, vomiting, swelling, and tissue death and can be life-threatening.

### **Vomit**

To eject stomach or upper gastrointestinal contents through the mouth; the expelled material may also be called vomitus. Persistent vomiting can cause

dehydration and electrolyte abnormalities.

### **Vomiting**

Forceful emptying of stomach contents through the mouth. It may result from infection, digestive disease, medicines, pregnancy, migraine, brain disorders, or poisoning. Repeated vomiting can cause dehydration, disturbed body salts, or inhalation of vomit.

### **Vomiting Blood**

The passage of blood or coffee-ground material in vomit, usually indicating bleeding in the upper digestive tract. It can be a serious symptom.

### **Vomitus**

Material expelled from the stomach or upper gastrointestinal tract during vomiting. Its color and contents can provide clues but do not by themselves establish a diagnosis.

### **Von Economo Encephalitis**

An atypical form of epidemic encephalitis (encephalitis lethargica) characterized by pronounced lethargy, hypersomnolence, ophthalmoplegia, and a frequent late sequela of progressive post-encephalitic parkinsonism, pathologically marked by midbrain and basal ganglia inflammation.

#### **Alternate Names:**

Encephalitis Lethargica; Sleeping Sickness (Von Economo Type)

### **Von Gierke Disease**

Glycogen storage disease type I, an inherited deficiency of glucose-6-phosphatase or its transport system that causes fasting hypoglycemia, lactic acidosis, enlarged liver, and excess uric acid or lipids.

### **Von Graefe Sign**

A delay of the upper eyelid as the eyes look downward, leaving a strip of white visible above the iris. It can occur with an overactive thyroid and related eye disease.

#### **Alternate Names:**

Lid Lag Sign; Von Graefe's Lid Lag

---

### Von Hippel-Lindau Disease

An inherited condition, usually caused by a change in the *VHL* gene, that increases the risk of tumors and cysts in the brain, spinal cord, eyes, kidneys, adrenal glands, and pancreas. These growths can affect vision, blood pressure, or kidney function.

### Von Hippel-Lindau Syndrome

Von Hippel-Lindau syndrome is an inherited disorder caused by changes in the *VHL* gene. It increases the risk of tumors and cysts in organs such as the brain, retina, kidneys, pancreas, and adrenal glands.

### Von Meyenburg Complex

A benign congenital biliary ductal plate malformation (biliary microhamartoma) consisting of small, non-progressive clusters of dilated, cystic bile ducts embedded in a dense collagenous stroma, usually discovered incidentally on liver imaging or biopsy.

#### Alternate Names:

Biliary Microhamartoma; Ductal Plate Malformation

### Von Recklinghausen Disease

Von Recklinghausen disease is another name for neurofibromatosis type 1, an inherited condition causing multiple café-au-lait spots, neurofibromas, and increased risk of certain nerve, bone, eye, and other problems.

### Von Recklinghausen'S Disease

Alternate name; see *Neurofibromatosis type 1*.

### Von Recklinghausen'S Neurofibromatosis

Alternate name; see *Neurofibromatosis type 1*.

### Von Recklinghausen'S Phakomatosis

Alternate name; see *Neurofibromatosis type 1*.

### Von Rosen Splint

A malleable, lightweight orthopedic abduction splint applied in neonates with developmental dysplasia of the hip (DDH) to maintain the femoral head securely centered within the acetabulum in flexion and abduction.

#### Alternate Names:

Von Rosen Frame; DDH Abduction Splint

### Von Willebrand Disease

A bleeding disorder caused by too little or poorly working von Willebrand factor, a blood protein that helps clots form. It is often autosomal dominant, although severe type 3 and some other forms are autosomal recessive. It commonly causes nosebleeds, easy bruising, heavy menstrual bleeding, or prolonged bleeding after procedures.

### Voriconazole

An antifungal medicine used for selected serious fungal infections. It stops fungi making an essential part of their outer structure; vision changes, liver problems, medicine interactions, and skin sensitivity require monitoring.

### VOR Reflex

A reflex that moves the eyes opposite to head movement so vision stays steady while the head turns. Problems with this reflex can cause blurred or bouncing vision, dizziness, or imbalance.

### Voyeurism

A recurrent sexual interest in observing an unsuspecting person who is naked, disrobing, or engaging in sexual activity. The interest alone is not a mental disorder; voyeuristic disorder is the diagnostic term when required criteria are met.

### Voyeuristic Disorder

A paraphilic disorder involving recurrent, intense sexual arousal from observing an unsuspecting person who is naked, disrobing, or engaging in sexual activity. Diagnostic criteria include acting on the urges or clinically significant distress or impairment, with the required duration criteria.

### VPCs (Ventricular Premature Contractions)

Alternate name; see *Premature ventricular contractions (PVCs)*.

---

### **VRE (Vancomycin-Resistant Enterococci)**

Enterococcus bacteria that are no longer killed by vancomycin, an antibiotic. They may live harmlessly in the body or cause urinary, wound, bloodstream, or other infections that require infection-control measures and suitable treatment.

### **VSD (Ventricular Septal Defect)**

A hole in the wall between the heart's lower chambers. Small holes may close or cause no symptoms, while larger holes can make the heart and lungs work too hard and may need treatment.

#### **Alternate Names:**

Ventricular Septal Defect

### **Vulnerable**

A substance traditionally believed to help heal wounds. Such products may lack good evidence and can cause allergy or contamination.

### **Vulva**

The external female genital structures, including the mons pubis, labia, clitoris, and vestibule. It protects the openings of the urinary and reproductive tracts and contains specialized sensory and glandular tissue.

#### **Alternate Names:**

Female External Genitalia; Pudendum

### **Vulval Intraepithelial Neoplasia**

Vulval intraepithelial neoplasia is abnormal cell growth limited to the surface layers of the vulvar skin. Some types are linked to HPV, and untreated lesions may progress to invasive cancer.

### **Vulvar**

Relating to the vulva, the external female genital structures including the labia, clitoris, vestibule, and associated glands.

### **Vulvar Cancer**

Cancer arising in the external female genitalia. It may appear as a persistent lump, ulcer, wart-like lesion, itching, pain, or bleeding.

### **Vulvar Carcinoma**

Cancer of the external female genital area, most often arising from surface skin cells. It may appear as a persistent lump, sore, wart-like growth, itching, pain, or bleeding.

#### **Alternate Names:**

Vulvar Cancer; Squamous Cell Carcinoma of Vulva

### **Vulvar Inclusion Cysts**

Small sacs under the skin of the vulva that contain tissue trapped from the surface lining. They may develop after childbirth tears, surgery, or other injury, or occur without an obvious cause. Most cause no symptoms, but they may become swollen, painful, infected, or uncomfortable during sexual activity. Symptomatic cysts may need removal.

### **Vulvar Intraepithelial Neoplasia**

Abnormal cells limited to the surface layer of the vulva. Some forms can progress to invasive cancer, especially without treatment. It may cause itching, burning, sores, warty growths, or a color change, but some people have no symptoms.

### **Vulvar Melanoma**

A rare cancer of the pigment-producing cells in the vulva. It may appear as a dark or skin-colored lump, patch, sore, or bleeding area; diagnosis requires a biopsy, and treatment depends on size and spread.

### **Vulvar Squamous Cell Carcinoma**

The most common invasive vulvar malignancy, arising through HPV-related or HPV-independent pathways. It usually presents as a persistent mass, ulcer, plaque, or wart-like lesion and requires biopsy for diagnosis.

### **Vulvectomy**

Surgical removal of part or all of the vulva, performed for selected cancers, precancerous lesions, or severe persistent disease.

### **Vulvitis**

Inflammation or irritation of the external female genital area. It can cause itching, burning, redness,

---

swelling, soreness, or discharge and may result from infection, allergy, skin disease, hormones, or friction.

### **Vulvodynia**

Persistent or recurring pain in the vulva without an identifiable cause or specific disorder. The pain may be localized or widespread, triggered by touch or pressure, or occur without touch; it is often described as burning, rawness, or soreness.

#### **Alternate Names:**

Chronic Vulvar Pain

### **Vulvovaginal Candidiasis**

A yeast infection of the vulva and vagina, usually caused by *Candida*. It commonly causes intense itching, burning, soreness, redness, and thick discharge; recurrent, severe, unusual, or treatment-resistant symptoms need testing to confirm the cause.

### **Vulvovaginitis**

Simultaneous inflammation or infection of the vulva and vagina, commonly presenting with pruritus, burning, dysuria, dyspareunia, and abnormal vaginal discharge, resulting from infectious etiologies (such as *Candida albicans*, Bacterial vaginosis, *Trichomonas vaginalis*) or non-infectious causes (atrophic vaginitis, contact irritants).

#### **Alternate Names:**

Vaginitis and Vulvitis; Vulvovaginal Inflammation

### **Vocal Cord Disorders**

Disorders affecting the vocal folds or their nerves, including nodules, polyps, cysts, inflammation, paralysis, spasmodic dysphonia, and paradoxical vocal-fold movement. They may cause hoarseness, voice loss, throat discomfort, noisy breathing, or difficulty breathing and swallowing.

### **Vocal Cord Dysfunction**

Inappropriate closing or narrowing of the vocal folds, especially during inhalation, causing sudden throat tightness, noisy breathing, cough, or shortness of breath that can resemble asthma. It is also called inducible laryngeal obstruction and needs assessment when breathing is difficult.

### **Vocal Nodules**

Benign callus-like growths on the vocal folds caused by repeated voice strain. They can produce hoarseness, a rough or breathy voice, vocal fatigue, and loss of range; voice therapy and reducing strain are common treatments.

### **Varicose Veins (Spider Veins)**

Alternate name; see *Varicose Veins* and *Spider Veins*.



# W

## Waardenburg Syndrome

A group of inherited conditions that may cause hearing loss and unusual coloring of the eyes, hair, or skin. Some forms also affect the arms or the nerves controlling the bowel; findings vary by type.

## Waddell Signs

A standardized group of physical examination signs evaluated during back assessment to identify non-organic or behavioral components contributing to lower back pain. Signs include superficial tenderness, simulation testing, distraction maneuvers, regional sensory or motor deficits inconsistent with neuroanatomy, and overreaction during examination.

### Alternate Names:

Waddell Non-Organic Signs; Waddell Score

## WAGR Syndrome

A rare genetic condition that may cause an increased risk of Wilms kidney tumor, missing or poorly formed irises, urinary or reproductive differences, and developmental delay. Children need regular specialist monitoring, especially for kidney tumors.

## Waist Circumference

The distance around the abdomen, measured at a standard level. A larger measurement can indicate more abdominal fat and a higher risk of diabetes and heart disease, but it must be interpreted with other health information.

## Waist-Hip Ratio

A measurement comparing waist size with hip size.

It provides an estimate of where body fat is stored and may add information about diabetes and cardiovascular risk, but it does not diagnose disease by itself.

## Wait List

A list of candidates registered to receive an organ transplant. Placement and allocation priority are determined by the applicable transplant system and may consider medical urgency, expected benefit, blood or tissue compatibility, organ size, waiting time, and other policy-based factors; it is not necessarily a simple first-come, first-served queue.

### Alternate Names:

Waiting List; Transplant Waiting List; Transplant Waitlist

## Wakefulness

The state of being conscious and able to respond to the surroundings. Sleep, medicines, illness, brain injury, and metabolic problems can change the level of wakefulness.

## Waldenström's Macroglobulinemia

A slow-growing blood cancer in which abnormal immune cells make too much of an antibody called IgM. The blood can become unusually thick, causing blurred vision, headaches, bleeding, confusion, or numbness.

## Waldenstrom Macroglobulinemia

A rare blood cancer in which abnormal immune cells build up in the bone marrow and make too much of an antibody called IgM. It may cause anemia, bleeding, vision or nerve problems, enlarged lymph

---

nodes, or no symptoms at first.

**Alternate Names:**

Lymphoplasmacytic Lymphoma

**Waldenstrom's Macroglobulinemia**

A lymphoplasmacytic cancer of the bone marrow that produces excess IgM and may cause anemia, enlarged lymph nodes, or hyperviscosity symptoms.

**Waldeyer Ring**

An annular anatomical arrangement of lymphoid tissue surrounding the opening of the pharynx and nasal cavity, consisting of the pharyngeal tonsils (adenoids), tubal tonsils, palatine tonsils, and lingual tonsils. It functions as the first line of immune defense against ingested or inhaled pathogens.

**Alternate Names:**

Waldeyer Lymphatic Ring; Pharyngeal Lymphoid Ring

**Walker-Warburg Syndrome**

A rare inherited disorder causing severe muscle weakness together with major abnormalities of the brain and eyes from birth. Babies may have low muscle tone, seizures, poor feeding, enlarged fluid spaces in the brain, and abnormal eye development; the condition is usually life-threatening in infancy.

**Alternate Names:**

WWS; Muscle-Eye-Brain Disease (Severe Type); HARD +/- E Syndrome

**Walking Abnormalities**

Walking abnormalities include changes in speed, balance, stride, posture, or coordination caused by pain, weakness, sensory loss, joint disease, or neurologic illness.

**Walking Pneumonia**

An informal clinical term for a mild, atypical community-acquired pneumonia (most frequently caused by *Mycoplasma pneumoniae* or *Chlamydia pneumoniae*), in which the patient presents with persistent dry cough, low-grade fever, and prominent radiographic interstitial infiltrates while remaining ambulatory and non-toxic.

**Alternate Names:**

Atypical Pneumonia

**Walking Speed**

The rate at which a person walks, often measured over a defined distance. It can provide information about mobility and function, but no single speed determines disability in every person.

**Wallenberg Syndrome**

A neurological condition caused by infarction of the lateral medulla oblongata, most commonly resulting from occlusion of the posterior inferior cerebellar artery (PICA) or vertebral artery. Manifestations include ipsilateral facial sensory loss, Horner syndrome, ataxia, dysphagia, hoarseness, and contralateral loss of pain and temperature sensation in the trunk and limbs.

**Alternate Names:**

Lateral Medullary Syndrome; PICA Syndrome; Posterior Inferior Cerebellar Artery Syndrome

**Wallerian Degeneration**

Breakdown of the part of a nerve beyond a severe injury after it loses contact with the nerve cell. The nerve may regrow slowly if the surrounding pathway remains intact, but recovery depends on the injury.

**Walleyed**

Walleyed describes eyes that appear misaligned or divergent; clinically, the condition may be strabismus and requires eye assessment when persistent.

**Wall Stress**

The strain placed on the wall of a heart chamber by the pressure inside it and the chamber's size. High wall stress makes pumping harder and may contribute to heart enlargement or failure.

**Wandering Spleen**

A spleen that can move unusually because its supporting bands are loose or missing. It may cause no symptoms or lead to abdominal pain, enlargement, twisting, loss of blood supply, or sudden severe illness.

---

## Warburg Effect

A hallmark metabolic phenotype of malignant neoplastic cells in which cells preferentially convert glucose to lactate via aerobic glycolysis even in the presence of abundant oxygen, rather than utilizing mitochondrial oxidative phosphorylation, generating glycolytic intermediates for rapid cell proliferation.

## Warfarin

A blood-thinning medicine that reduces production of vitamin-K-dependent clotting proteins. It prevents and treats clots but requires regular blood tests because food, illnesses, and many medicines alter its effect.

## Warfarin and Superwarfarin Toxicity

Excess anticoagulation from warfarin or long-acting rodenticides, causing an elevated INR and potentially delayed bleeding, including gastrointestinal or intracranial hemorrhage. Serial coagulation testing is essential; vitamin K reverses the effect, while life-threatening bleeding may require four-factor prothrombin complex concentrate and prolonged treatment for superwarfarins.

## Warfarin-Induced Skin Necrosis

A rare, catastrophic thrombotic complication occurring 2 to 5 days after starting warfarin without heparin bridging, caused by the rapid initial decline in the short-half-life natural anticoagulant protein C before procoagulant factors fall, triggering microvascular thrombosis and cutaneous infarction in breasts, buttocks, and thighs.

### Alternate Names:

Coumadin-Induced Skin Necrosis

## Warfarin Syndrome

A pattern of fetal abnormalities caused by warfarin exposure during pregnancy, including nasal and skeletal abnormalities, eye or neurologic defects, and fetal or neonatal bleeding.

## Warm and Cold Autoantibodies

Antibodies that make red blood cells clump at warmer or colder temperatures. They can inter-

fere with laboratory testing or cause autoimmune destruction of red cells and anemia.

## Warm Autoimmune Hemolytic Anemia

An illness in which the immune system makes antibodies that cause red blood cells to be destroyed too early, mainly in the spleen. It can cause tiredness, pale or yellow skin, dark urine, shortness of breath, or a fast heartbeat and requires medical evaluation.

### Alternate Names:

wAIHA; Warm Antibody Hemolytic Anemia

## Warm Ischemia

Lack of blood and oxygen reaching tissue while it remains at body temperature. It can quickly cause cell damage and is important during some injuries, operations, and organ transplantation.

## Wart

A usually harmless rough skin growth caused by human papillomavirus. Warts may spread by contact and often disappear over time. Common types include common, plantar, and flat warts; appearance and symptoms vary by site.

### Alternate Names:

Verruca

## Warthin Tumor

A usually harmless tumor of a salivary gland, most often the parotid near the ear. It is strongly associated with tobacco use and may occur on both sides; evaluation distinguishes it from other growths.

## Wart Remover Poisoning

Toxic exposure to a wart-removal product, often involving salicylic acid or another keratolytic. Severity depends on the substance, amount, route, symptoms, and duration of exposure.

## Warty Dyskeratoma

A usually harmless skin growth, often on the head or neck, with a small central dip filled with keratin. It may look like skin cancer, so examination and sometimes a biopsy are needed.

---

## Wasting

Abnormal loss of body weight, muscle, or other tissue, often associated with chronic disease, inadequate nutrition, inflammation, or cancer. Severe wasting may impair strength, immunity, and recovery.

## Watchful Waiting

Planned observation without immediate active treatment, with clinical review and treatment if symptoms develop or the condition progresses. It may be used for selected cancers and other slowly progressive conditions.

## Water Birth

Delivery of a baby while the birthing person remains in a tub or pool of warm water. It may reduce perceived labor pain for some people, but evidence about delivery-related benefits and risks is limited. Eligibility, monitoring, and facility policies determine whether it is offered.

### Alternate Names:

Aquatic Delivery; Water Labor and Birth

## Water Blister

A fluid-filled blister, usually containing clear serum, caused by friction, pressure, burns, or some skin disorders. The blister should be protected from further trauma and assessed if infected or widespread.

## Waterborne Bacterial Disease

An infection acquired from water contaminated with bacteria, such as cholera, typhoid fever, or leptospirosis; prevention depends on safe water, sanitation, and food hygiene.

## Water-Borne Disease

An illness acquired through water contaminated with infectious organisms or harmful chemicals. Examples include cholera, typhoid fever, and hepatitis A; prevention includes safe drinking water, sanitation, hand hygiene, and appropriate treatment of contaminated water.

## Water Brash

A sudden rise of watery or salty-tasting saliva into the mouth, usually because stomach acid has refluxed into the esophagus. It may occur with heartburn, sour taste, throat irritation, or regurgitation.

## Water Channel

A protein pore in a cell membrane that lets water cross quickly. These pores help the kidneys and other tissues move water and maintain the body's fluid balance.

## Water Consumption

Intake of water and other fluids to replace normal losses and maintain hydration. Requirements vary with body size, diet, activity, climate, pregnancy, illness, and kidney or heart function; thirst and urine concentration can help guide intake, while excessive intake can cause dangerous hyponatremia.

## Water-Deprivation Test

A supervised test that briefly limits drinking while urine concentration and blood results are monitored. It helps identify why a person produces excessive amounts of dilute urine and must be done under medical supervision because dehydration and high sodium can occur.

## Water Hammer Pulse

A physical sign of severe chronic aortic regurgitation or high-output states characterized by a bounding, hyperdynamic peripheral pulse with an abrupt, forceful systolic upstroke followed by a rapid, precipitous diastolic collapse.

### Alternate Names:

Corrigan Pulse; Collapsing Pulse

## Water-Hammer Pulse

A pulse that feels forceful at first and then suddenly collapses. It is often associated with severe leakage of the aortic valve or other causes of a wide pulse pressure.

### Alternate Names:

Corrigan's Pulse; Collapsing Pulse; Cannon Pulse

---

### Waterhouse-Friderichsen Syndrome

Waterhouse-Friderichsen syndrome is adrenal hemorrhage and acute adrenal failure associated with overwhelming sepsis, classically meningococcal infection, causing shock and coagulopathy.

### Water Intoxication

A dangerous fall in blood sodium caused by drinking or retaining more water than the body can remove. Severe cases can cause headache, confusion, seizures, or coma and need urgent treatment.

### Water On The Brain

A nontechnical term usually referring to hydrocephalus, an abnormal accumulation or impaired circulation of cerebrospinal fluid that enlarges the ventricles and may increase intracranial pressure.

### Water On The Knee

Alternate name; see *Swollen knee*.

### Water Retention

Excess accumulation of fluid in the body's tissues or cavities, commonly causing swelling and sometimes resulting from heart, kidney, liver, venous, hormonal, or medication-related disorders.

### Watershed Infarction

An ischemic cerebral or intestinal infarct occurring at the border zones between the terminal distal capillary distributions of two non-anastomosing major arterial territories (e.g., between ACA and MCA in the brain, or at the splenic flexure / Griffiths point in the colon), provoked by systemic hypoperfusion and severe hypotension.

#### Alternate Names:

Border-Zone Infarction

### Water-Soluble Vitamin

A vitamin that dissolves in water and is stored only in limited amounts. The group includes vitamin C and the B vitamins, so regular intake is generally needed.

### Waterston Shunt

A historical palliative surgical systemic-to-pulmonary shunt created between the ascending aorta and the right pulmonary artery to increase pulmonary blood flow in infants with cyanotic congenital heart defects. It has largely been replaced by the modified Blalock-Taussig shunt due to lower complication rates.

#### Alternate Names:

Waterston-Cooley Anastomosis; Ascending Aorta to Right Pulmonary Artery Shunt

### Watery Eyes

Excessive tearing caused by irritation, allergy, infection, dry-eye reflex tearing, or blockage of the tear-drainage system. Persistent unilateral tearing or pain requires ocular assessment.

### Wax

A water-resistant substance produced by ceruminous glands in the ear canal; earwax protects the canal and usually clears naturally.

### Wax Poisoning

Toxic exposure to a wax product, usually causing little harm after a small accidental ingestion but potentially causing choking, aspiration, irritation, or toxicity when mixed with other chemicals.

### Waxy Casts

Large, smooth, pale tube-shaped particles seen under a microscope in urine. They form in damaged kidney tubules and can suggest severe or long-standing kidney disease, especially when other urine or blood tests are abnormal.

### WBC Count

A measurement of the number of white blood cells in a volume of blood, used with the differential count to evaluate infection, inflammation, immune disorders, marrow disease, or treatment effects.

### Weakness

Reduced ability to generate force, which may be generalized or limited to one body region. Causes

---

include disorders of muscle, nerve, neuromuscular junction, brain, or spinal cord, as well as metabolic illness, infection, medication effects, and deconditioning; sudden one-sided weakness or weakness with breathing, swallowing, or speech difficulty requires emergency assessment.

### Weaning

Gradual reduction and eventual discontinuation of breastfeeding, bottle-feeding, or a medicine or other treatment, according to the relevant clinical context.

### Wear Debris

Tiny particles produced when an artificial joint or other implant wears down. The particles can trigger inflammation, damage nearby bone, loosen the implant, and eventually cause it to fail.

### Weaver's Bottom

Inflammation of the ischial bursa over the sitting bone, causing pain in the buttock that may worsen with prolonged sitting. It is also called ischial bursitis.

### Weaver Syndrome

A rare genetic condition causing rapid growth before and after birth, a large head, distinctive facial features, low muscle tone, and delayed development or learning. Bone age may be advanced. Most cases involve a change in the *EZH2* gene; care is tailored to the child's needs.

### Webbed Fingers

Partial or complete joining of adjacent fingers by skin or soft tissue, known as syndactyly; it may be congenital or acquired.

### Webbed Neck

Extra folds of skin extending from the sides of the neck toward the shoulders. They may be present from birth and can occur with conditions such as Turner syndrome or Noonan syndrome.

### Webbing Of The Fingers Or Toes

Partial or complete joining of adjacent fingers or

toes by skin or other tissue. It is called syndactyly and may be congenital or acquired.

### Weber

A name used in hearing assessment, especially the Weber test, in which a vibrating tuning fork is placed on the forehead or skull. It helps compare hearing between the two ears but is not a unit of measurement.

### Weber-Christian Disease

Alternate name; see *Weber-Christian Disease*.

### Weber Syndrome

A brainstem stroke syndrome caused by occlusion of paramedian branches of the posterior cerebral artery affecting the anterior midbrain. It is characterized by ipsilateral oculomotor (cranial nerve III) palsy and contralateral spastic hemiparesis of the upper and lower extremities.

#### Alternate Names:

Superior Alternating Hemiplegia; Ventral Midbrain Syndrome

### Weber Test

A quick hearing test using a vibrating tuning fork placed on the middle of the forehead or skull. The examiner asks which ear hears the sound louder; the result helps distinguish blockage-related hearing loss from inner-ear or nerve-related loss.

### Wedge Excision

Surgical removal of a triangular or wedge-shaped portion of tissue; commonly performed for breast biopsies, cervical lesions, and skin tumors.

### Wedge Resection

Surgery that removes a small wedge-shaped piece of tissue, commonly from the lung or breast, to diagnose or treat a localized lesion. It removes less tissue than removal of an entire lobe.

### Weever Fish

A venomous marine fish whose dorsal spines can cause a painful sting, local swelling, and occasionally



---

systemic symptoms.

### Weight-Bearing As Tolerated

An orthopaedic postoperative rehabilitation prescription allowing the patient to place as much weight and mechanical force on an injured or operated lower extremity as is comfortable and tolerated without severe pain, typically using crutches or a walker for support.

**Alternate Names:**  
WBAT

### Weight Gain

Increase in body weight; may be intentional (muscle building, pregnancy) or unintentional (fluid retention, endocrine disorders, medication effects).

### Weight Gain In Pregnancy

Recommended weight gain depends on weight before pregnancy and whether one or more babies are expected. For one baby, usual ranges are about 12.5–18 kg if underweight, 11.5–16 kg at usual weight, 7–11.5 kg if overweight, and 5–9 kg with obesity; prenatal clinicians monitor the pattern and fetal growth.

### Weight Gain – Unintentional

Increase in body weight without a planned change in diet or activity, often caused by fluid retention, medicines, endocrine disease, reduced activity, or changes in eating; rapid swelling or breathlessness needs prompt assessment.

### Weight Loss

A decrease in body weight that may be intentional or unintentional. Intentional loss is usually achieved through sustainable changes in eating, physical activity, and behavior; unintentional or rapid loss can indicate illness, inadequate nutrition, or medication effects and may require medical evaluation.

### Weight Loss Medications

Medicines prescribed for selected people with obesity to help reduce weight, usually together with nutrition and activity changes. Choice depends on health conditions, other medicines, side effects, cost,

and treatment goals.

### Weight-Loss Medicines

Alternate name; see *Weight Loss Medications*.

### Weight Loss Surgery

Weight-loss surgery, or bariatric surgery, changes the stomach or digestive tract to help treat severe obesity and related conditions. It requires long-term nutritional, medical, and lifestyle follow-up.

### Weight-Reducing Diets

Dietary plans intended to create a sustained energy deficit and reduce excess body weight, ideally individualized with attention to nutrition, health conditions, and long-term adherence.

### Weil Disease

Alternate name; see *Leptospirosis*.

### Weil-Felix Reaction

A historical agglutination serological test for rickettsial infections (such as typhus and spotted fevers) that utilizes cross-reacting somatic O-antigens of *Proteus vulgaris* strains (OX-19, OX-2, and OX-K) to detect serum anti-rickettsial antibodies.

### Weil-Felix Test

A classic serological agglutination assay that utilizes antigens from cross-reacting *Proteus vulgaris* strains (OX-19, OX-2, OX-K) to detect serum antibodies against various *Rickettsia* species responsible for typhus and spotted fever infections.

**Alternate Names:**

Weil-Felix Agglutination Reaction; Proteus Agglutination Test

### Weill-Marchesani Syndrome

A rare inherited systemic connective tissue disorder characterized by the constellation of brachydactyly (short fingers and toes), short stature, joint stiffness, microspherophakia (small spherical lenses), and ectopia lentis with high predisposition to secondary pupillary block angle-closure glaucoma. Caused by autosomal recessive mutations in *ADAMTS10* or

---

autosomal dominant mutations in *FBN1*.

**Alternate Names:**

WMS; Spherophakia-Brachymorphia Syndrome

**Weil's Disease**

A zoonotic bacterial infection caused by spirochaetes of the genus *Leptospira*, transmitted through contact with water or soil contaminated with the urine of infected animals (rodents, cattle, dogs). See *Lep-tospirosis*.

**Wellens Syndrome**

A specific pattern on a heart tracing that can signal severe narrowing of a major heart artery, even after chest pain stops. It indicates high heart-attack risk and needs urgent specialist assessment; exercise testing can be dangerous.

**Alternate Names:**

Wellens Pattern; LAD Coronary T-Wave Syndrome; Wellens Sign

**Wells Score**

A validated clinical prediction rule used to estimate the probability that a patient has a dangerous blood clot, specifically deep vein thrombosis (DVT) in the legs or pulmonary embolism (PE) in the lungs. By calculating points for clinical risk factors (including active cancer, recent surgery or bed rest, localized leg swelling or tenderness, fast heart rate, coughing blood, and prior clots), patients are categorized as “unlikely” (low risk) or “likely” (high risk). In low-risk patients, a normal/negative D-dimer blood test safely rules out a clot without scans; high-risk patients proceed directly to imaging (leg compression ultrasound for DVT or CT pulmonary angiography for PE).

**Alternate Names:**

Wells Criteria; Wells Clinical Score; Wells Rule for DVT and PE

**Wells Score For Pulmonary Embolism**

Alternate name; see *Wells Score*.

**Wells Syndrome (Eosinophilic Cellulitis)**

A recurrent inflammatory skin disorder with itchy or burning red plaques, swelling, and tissue eosinophils

that can mimic bacterial cellulitis. It is not usually an infection; biopsy helps diagnosis and systemic corticosteroids may be used for severe flares.

**Welt**

A raised, often itchy area of skin caused by localized swelling, commonly seen in hives or after an insect bite. A welt may fade within hours, but persistent or severe reactions need evaluation.

**Wen**

A common, usually harmless lump beneath the skin filled with a soft protein called keratin. It may become swollen, painful, or infected and can be removed if necessary.

**Alternate Names:**

Infundibular Cyst; Epidermal Inclusion Cyst; Sebaceous Wen

**Wenckebach Phenomenon**

A heart-rhythm pattern in which each heartbeat takes slightly longer to pass from the upper to lower chambers until one beat is missed. It may be harmless or reflect heart disease, depending on the setting.

**Alternate Names:**

Second-Degree AV Block Type 1; Mobitz I Block

**Wens**

Alternate name; see *Epidermoid cyst*.

**Werdnig-Hoffmann Disease**

The most severe early form of inherited spinal muscular atrophy, also called type 1 spinal muscular atrophy. It usually begins before 6 months of age and causes very weak muscles, poor head control, reduced reflexes, a weak cry, feeding difficulty, and breathing problems. Babies generally cannot sit without support. New treatments can improve survival and function, but affected infants need urgent specialist care and breathing and feeding support.

**Alternate Names:**

Spinal Muscular Atrophy Type 1 (SMA 1); Severe Infantile Spinal Muscular Atrophy

---

### Wermer's Syndrome

Alternate name; see *Multiple endocrine neoplasia, type 1 (MEN 1)*.

### Wermer Syndrome

Alternate name; see *Multiple endocrine neoplasia, type 1 (MEN 1)*.

### Werner's Syndrome

A rare inherited condition that causes signs of aging earlier than usual, often after puberty. It may cause early cataracts, short stature, thin skin, gray hair, diabetes, weak bones, and increased cancer risk.

### Werner Syndrome

A rare inherited condition that causes signs of aging earlier than usual, often after puberty. Features may include short stature, early cataracts, thin or tight skin, gray hair, diabetes, weak bones, and increased cancer risk. Regular medical monitoring and treatment of each complication are needed.

### Wernicke Aphasia

A fluent receptive aphasic syndrome resulting from an ischemic stroke or lesion in the posterior portion of the superior temporal gyrus of the dominant hemisphere (Brodmann area 22), characterized by fluent, effortless speech that lacks semantic meaning ("word salad"), accompanied by profound impairment in language comprehension and repetition.

#### Alternate Names:

Receptive Aphasia; Sensory Aphasia

### Wernicke Area

A brain region, usually in the left temporal lobe, that helps a person understand spoken and written language. Damage can cause fluent speech that lacks meaning and poor understanding of language.

### Wernicke Encephalopathy

A sudden brain disorder caused by severe vitamin B1 deficiency. It may cause confusion, difficulty walking, and abnormal eye movements and can lead to permanent damage without prompt treatment.

### Wernicke-Korsakoff Encephalopathy

A spectrum of neurocognitive impairment caused by severe thiamine (vitamin B1) deficiency (most commonly in chronic alcohol use disorder), progressing from the acute reversible triad of Wernicke encephalopathy (confusion, ataxia, ophthalmoplegia) to the chronic irreversible Korsakoff syndrome (anterograde amnesia, confabulation).

### Wernicke-Korsakoff Syndrome

A brain disorder caused by severe vitamin B1 deficiency. It may begin with confusion and poor coordination and leave lasting memory problems, difficulty learning, and inaccurate made-up explanations; alcohol misuse, poor nutrition, and prolonged vomiting are common causes.

### Wernicke's Area

A brain region important for understanding spoken and written language.

#### Alternate Names:

Wernicke's Speech Area; Sensory Speech Area

### Wernicke's Encephalopathy

An acute neurologic disorder caused by thiamine deficiency, classically involving confusion, eye-movement abnormalities, and gait or coordination impairment. It requires prompt thiamine treatment.

### Westergren Method

The reference laboratory procedure used to determine the erythrocyte sedimentation rate (ESR). Anticoagulated whole blood is placed in an upright, calibrated Westergren tube for one hour, measuring the distance in millimeters that red blood cells settle under gravity to assess systemic inflammation.

#### Alternate Names:

Westergren ESR; Westergren Sedimentation Technique

### Westermark Sign

An area of unusually reduced blood-vessel markings on a chest X-ray, sometimes seen beyond a clot in a lung artery. It may suggest pulmonary embolism but is not reliable enough to diagnose it alone.

---

## Western Blot

A laboratory test that separates proteins and uses antibodies to find a specific protein. It can help study infections, immune reactions, or genetic and research problems, depending on the test.

### Alternate Names:

Protein Immunoblot

## Westley Croup Score

A clinical score used to estimate how severely croup is obstructing a child's airway. It considers noisy breathing, chest retractions, air movement, bluish coloring, and alertness; a higher score suggests more severe illness and helps guide urgent treatment and observation.

### Alternate Names:

Westley Clinical Scoring System for Croup; Croup Severity Score

## West Nile Encephalitis

Inflammation of the brain caused by neuroinvasive West Nile virus, transmitted mainly by mosquitoes. It may cause fever, headache, confusion, tremor, weakness, seizures, or coma. There is no specific routine antiviral treatment; severe illness requires hospital-based supportive care.

## West Nile Neuroinvasive Disease

A severe, potentially fatal manifestation of mosquito-borne West Nile virus infection affecting less than 1 percent of infected individuals, presenting as aseptic meningitis, encephalitis (with tremors and parkinsonism), or acute flaccid paralysis resembling poliomyelitis (anterior horn cell loss).

### Alternate Names:

WNND

## West Nile Virus

A mosquito-borne flavivirus that can cause asymptomatic infection, a febrile illness, or neuroinvasive disease such as meningitis, encephalitis, or acute flaccid paralysis. Risk of severe disease is higher in older adults and people with certain medical conditions.

## West Nile Virus Infection

West Nile virus infection is a mosquito-borne illness ranging from fever and body aches to neuroinvasive meningitis, encephalitis, weakness, or paralysis.

## West Syndrome

A serious seizure disorder beginning in infancy, with repeated sudden spasms and an abnormal brain-wave pattern. Development may slow or regress; causes include brain injury, genetic conditions, and metabolic disease, and urgent treatment is needed.

## Wet Age-Related Macular Degeneration

Alternate name; see *Wet macular degeneration*.

## Wet AMD

Alternate name; see *Wet macular degeneration*.

## Wet Beriberi

The predominantly cardiovascular form of severe thiamine deficiency. It can cause a fast heartbeat, enlarged or weakened heart, fluid retention, swelling, and high-output heart failure.

## Wet Dream

Involuntary ejaculation during sleep, usually occurring in adolescence or adulthood and generally a normal physiological event.

## Wet Gangrene

Tissue death complicated by bacterial infection and tissue swelling, producing moist, discolored, rapidly spreading necrosis. It requires urgent medical assessment.

## Wet Macular Degeneration

The neovascular form of age-related macular degeneration in which abnormal blood vessels grow beneath or within the retina and leak fluid or blood, causing rapid central-vision loss.

### Alternate Names:

Neovascular AMD; Exudative Age-Related Macular Degeneration; Wet AMD

---

### Wet-to-Dry Dressing

A mechanical wound debridement technique in which saline-moistened gauze is applied directly to an open wound bed, allowed to dry, and subsequently removed. Dead tissue and exudate adhering to the dried dressing are pulled away during dressing changes.

#### Alternate Names:

Wet-to-Dry Debridement; Mechanical Gauze Dressing

### Wharton Duct

The primary excretory duct of the submandibular salivary gland, approximately 5 cm in length, that courses anteromedially along the floor of the mouth to empty through the sublingual caruncle at the base of the lingual frenulum; the most frequent anatomical site of salivary sialolithiasis.

#### Alternate Names:

Submandibular Duct

### Wharton Jelly

The soft, jelly-like tissue surrounding the blood vessels in the umbilical cord. It cushions the vessels and helps prevent them from being squeezed or bent during pregnancy and labor.

### Wharton's Jelly

Gelatinous connective tissue within the umbilical cord that surrounds and supports the umbilical blood vessels and helps protect them from compression.

### Wheal

A temporary, raised, often itchy area of skin caused by swelling in the skin. It commonly occurs with hives, an allergic reaction, or an insect bite and may move or disappear within hours.

### Wheal And Flare

A temporary raised, swollen area of skin surrounded by redness. It is usually caused by release of histamine during hives, an allergy, or another immediate skin reaction.

### Wheat Allergy

An immune reaction to proteins in wheat that may cause hives, stomach symptoms, vomiting, or breathing problems.

#### Alternate Names:

Wheat Hypersensitivity; Cereal Grain Allergy

### Wheelchairs

Alternate name; see *Wheelchair*.

### Wheeled Stretcher

Mobile patient transport device with wheels; used in hospitals and emergency services for moving patients who cannot walk.

### Wheeze

A high-pitched whistling sound caused by narrowed airways. It often occurs with asthma, infection, allergy, or chronic lung disease; sudden wheezing after choking or with severe breathlessness needs urgent care.

### Wheezing

A musical or whistling sound caused by narrowed lower airways, usually louder when breathing out. It may result from asthma, infection, chronic lung disease, allergy, airway swelling, mucus, or an inhaled object.

### Whiplash

A neck injury caused by the head moving suddenly backward and forward, often in a vehicle crash but also after a fall or sports injury. Pain, stiffness, headache, dizziness, or arm tingling may appear immediately or later. Most cases improve over weeks, but weakness, numbness, trouble walking, or severe pain after major trauma needs urgent assessment.

#### Alternate Names:

Cervical Acceleration-Deceleration Injury; CAD Injury; Neck Sprain

### Whiplash Injury

Alternate name; see *Whiplash*.



---

## Whipple Disease

A rare bacterial infection that can damage the small intestine, joints, brain, heart, or other organs. It may cause years of joint pain followed by weight loss, persistent diarrhea, abdominal pain, and poor nutrient absorption.

## Whipple Procedure

A major operation that removes the head of the pancreas and nearby parts of the small intestine, bile duct, and gallbladder, then reconnects the digestive tract. It is used for selected cancers and other serious diseases in this area.

## Whipple's Disease

A rare bacterial infection caused by *Tropheryma whipplei*, damaging intestinal absorption and potentially affecting joints, brain, or heart.

### Alternate Names:

*Tropheryma Whipplei* Infection; Whipple Disease

## Whipple Triad

Three findings supporting that symptoms are caused by low blood sugar: typical symptoms occur, blood sugar is low at that time, and symptoms improve after blood sugar is raised.

## Whipple Triad for Insulinoma

Three findings that support that symptoms are caused by low blood sugar: symptoms occur, blood sugar is low at the same time, and the symptoms improve when blood sugar rises. Doctors use this pattern with other tests to find the cause, including a possible insulinoma.

## Whipple Triad of Hypoglycemia

The three cardinal clinical criteria required to substantiate a clinical diagnosis of genuine hypoglycemia: symptoms and signs consistent with hypoglycemia (such as diaphoresis, tremor, tachycardia, confusion); a documented low plasma glucose concentration (typically below 55 mg/dL [3.0 mmol/L]) at the exact time of symptoms; and prompt resolution of symptoms following administration of exogenous glucose.

### Alternate Names:

## Whipple Hypoglycemic Triad

## Whipworm

A parasitic roundworm of the genus *Trichuris*, usually *Trichuris trichiura* in humans. Infection occurs by ingesting embryonated eggs and may cause abdominal pain, diarrhea, anemia, or rectal prolapse when severe.

## Whipworm Infection

Intestinal infection caused by the soil-transmitted helminth *Trichuris trichiura* (whipworm), endemic in tropical and subtropical regions with poor sanitation. Humans acquire infection by ingesting embryonated eggs from contaminated soil, food, or water.

### Alternate Names:

*Trichuris Trichiura* Infection; Whipworm Infection

## Whirlpool Bath

Hydrotherapy device providing agitated water for therapeutic bathing; used in wound care, physical therapy, and rehabilitation.

## Whispered Pectoriloquy

An examination finding in which whispered words sound unusually clear through a stethoscope over the lung. It may suggest that air-filled lung has become solid, as in pneumonia, but it is not diagnostic alone.

## White Blood Cell

A cell of the immune system that helps protect the body from infections, foreign substances, and abnormal cells. White blood cells are made mainly in the bone marrow and include neutrophils, lymphocytes, monocytes, eosinophils, and basophils. Their number or function may change with infection, inflammation, immune disorders, or bone-marrow disease.

## White Blood Cell Count

A blood test measuring the number of white blood cells. A high or low result can occur with infection, inflammation, medicines, immune disorders, or bone-marrow disease and must be interpreted with the



---

differential count and clinical findings.

### **White Blood Cell Labeling**

A scan method in which a person's white blood cells are removed, marked with a small amount of radioactive material, and returned to the body. The cells may collect at areas of infection or inflammation, which can then be seen on images.

### **White Cell**

Another name for a white blood cell, an immune cell made mainly in the bone marrow that helps defend the body against infection, foreign substances, and abnormal cells.

### **White Cell Count**

A blood test measuring the number of white blood cells and often the proportion of each type. Results help assess infection, inflammation, immune disorders, and bone-marrow problems.

### **White Clot Syndrome**

A severe arterial or venous thrombotic complication associated with heparin-induced thrombocytopenia (HIT). It is characterized by platelet-dense, fibrin-rich arterial thrombi (white thrombi) that can lead to limb ischemia, stroke, or myocardial infarction.

#### **Alternate Names:**

HIT-Associated Thrombosis; Heparin-Associated White Thrombus Syndrome

### **White Coat Hypertension**

A clinical blood pressure pattern characterized by persistently elevated blood pressure readings recorded in a medical office or clinic setting in an individual whose out-of-office automated ambulatory or home blood pressure monitoring readings are consistently normotensive.

#### **Alternate Names:**

White Coat Effect; Isolated Clinic Hypertension

### **White-Coat Hypertension**

Blood pressure that is high in a medical setting but normal outside it. Home or ambulatory readings help confirm the pattern, which may still predict

higher future hypertension risk.

#### **Alternate Names:**

White-Coat Syndrome; Isolated Clinic Hypertension

### **White Finger**

Episodic pallor or whitening of the fingers caused by vasospasm, commonly associated with Raynaud phenomenon or hand-arm vibration syndrome.

### **Whiteheads**

Whiteheads are closed comedones formed when oil and dead skin block a hair follicle, producing small pale or skin-colored acne bumps.

### **White Infarction**

Tissue death caused by loss of blood flow in a solid organ such as the heart, kidney, or spleen. The damaged area looks pale because little blood can reach it through alternate vessels.

#### **Alternate Names:**

Anemic Infarction; Pale Infarction

### **White Matter**

Tissue in the brain and spinal cord made mainly of nerve fibers covered by a protective substance called myelin. It carries signals between different brain areas and between the brain and spinal cord.

### **White Matter Hyperintensities**

Areas of increased signal intensity on T2-weighted and FLAIR brain magnetic resonance imaging scans located in the deep periventricular and subcortical white matter, representing microvascular lipohyalinosis, chronic ischemia, and demyelination from cerebral small vessel disease.

#### **Alternate Names:**

WMHs; Leukoaraiosis

### **White Matter Of The Brain**

The brain tissue composed mainly of myelinated axons and supporting glial cells, connecting different brain regions and the brain with the spinal cord; disease or injury can impair cognition, sensation, coordination, or movement.

---

## White Of The Eye

The sclera, the tough, opaque outer coat of the eye surrounding the cornea and helping maintain the shape of the globe.

## White Spot Disease

Alternate name; see *Lichen sclerosus*.

## White Tongue

A white coating or patches on the tongue caused by debris, dry mouth, irritation, infection, inflammation, or other oral conditions.

## Whitlow

An infection of the fingertip or tissues around a finger or toenail, often involving the nail fold. It may be bacterial, fungal, or caused by herpes simplex virus; painful grouped blisters suggest herpetic whitlow, whereas pus and a localized abscess suggest bacterial infection.

## Whitlows

Plural of whitlow; see *Whitlow*.

## WHO Classification of Vitamin A Deficiency

A standardized clinical grading system established by the World Health Organization (WHO) to stage eye damage (xerophthalmia) caused by vitamin A deficiency:

- **XN:** Night blindness (inability to see in dim light or darkness).
- **X1A:** Conjunctival xerosis (dryness and lack of shine on the white of the eye).
- **X1B:** Bitot's spots (foamy, silvery-white patches on the white of the eye).
- **X2:** Corneal xerosis (hazy, dry surface on the clear front of the eye).
- **X3A:** Corneal ulceration or keratomalacia involving less than one-third of the cornea.
- **X3B:** Severe corneal melting (keratomalacia) involving one-third or more of the cornea, causing permanent blindness.
- **XS:** Corneal scars (permanent scarring left by healed ulcers).

- **XF:** Xerophthalmic fundus (small pale spots on the back of the eye/retina).

### Alternate Names:

WHO Xerophthalmia Classification; Vitamin A Deficiency Eye Grading; WHO Vitamin A Staging

## Whole-Body Irradiation

Radiation treatment directed at the entire body, most often given before a stem-cell transplant to suppress the immune system and make room for donated cells. It can affect the bone marrow, fertility, and other organs.

## Whole-Body Magnetic Resonance Imaging

An MRI examination that produces images of much or all of the body without using X-rays. It is used in selected cases to look for widespread disease, tumors, or problems affecting several body areas.

## Whole-Body Scan

An imaging examination that surveys much or all of the body, using a technique such as PET/CT, CT, MRI, or a nuclear medicine scan. It may be used for selected cancer staging, infection localization, or evaluation of systemic disease; the coverage and usefulness depend on the test and clinical question.

## Whole Bowel Irrigation

A gastrointestinal decontamination technique in acute clinical toxicology in which large volumes of an isosmotic polyethylene glycol electrolyte solution are administered orally or via nasogastric tube until the rectal effluent is clear, indicated for toxic ingestions of sustained-release medications, enteric-coated tablets, iron, or illicit drug packets ("body packers").

### Alternate Names:

WBI

## Whole-Bowel Irrigation

A hospital treatment that flushes the intestines with a special polyethylene-glycol drink. It is used only for selected poisonings or swallowed objects after clinicians assess the airway and risks.

---

### Whole-Brain Radiation Therapy

Radiation treatment directed at the entire brain, used in selected people with multiple brain metastases or cancer cells around the brain. It may also be used to reduce brain-spread risk in selected cancers; side effects depend on dose and treatment plan.

### Whole Breast Radiation Therapy

Whole breast radiation therapy treats remaining breast tissue after breast-conserving surgery to reduce local recurrence risk, with planned dosing and monitoring for skin and tissue effects.

### Whole Exome Sequencing

A genetic test that reads most protein-coding sections of a person's DNA. It can identify changes causing some inherited disorders, but it may miss changes outside coding regions, structural changes, or causes not yet understood; results require clinical interpretation.

### Whole Genome Sequencing

Laboratory analysis of nearly all of a person's nuclear DNA, including coding and noncoding regions, and sometimes mitochondrial DNA. It can identify many types of genetic variants but may produce uncertain or incidental findings and does not detect every cause of disease; results require appropriate clinical interpretation.

### Whole-Genome Sequencing

Sequencing of nearly the complete nuclear genome, including coding and noncoding regions, together with mitochondrial DNA when covered by the assay. It can detect a broad range of variants but generates substantial interpretation and incidental-finding challenges.

### Whooping Cough

A highly contagious respiratory infection caused by *Bordetella pertussis*. It causes prolonged bouts of rapid coughing that may be followed by a whooping sound or vomiting; infants may have pauses in breathing without the typical whoop. Also called pertussis.

### Whooping-Cough

Alternate name; see *Whooping Cough*.

### Whorled Pattern

A characteristic histological or clinical swirling, spiraling architectural configuration. In pathology, it describes swirling arrangements of spindle cells seen in meningiomas, fibromas, or leiomyomas; in dermatology, it describes patterns along the lines of Blaschko.

#### Alternate Names:

Whorled Architecture; Storiform-Whorled Pattern

### Wickham Striae

Fine, lacy white reticular lines and arborizing patterns seen on the surface of papules and mucosal plaques in lichen planus, classically observed on the buccal mucosa.

### Widal Reaction

An older blood test for antibodies associated with typhoid or paratyphoid fever. It is not very reliable because results may reflect past infection or vaccination, so culture or molecular testing is usually preferred.

### Widal Test

A historical presumptive serological agglutination test for enteric (typhoid and paratyphoid) fever that detects agglutinating serum antibodies against the somatic O and flagellar H antigens of *Salmonella enterica* serovar Typhi.

### Widely Spaced Teeth

Teeth separated by unusually large gaps, also called diastema when the gap is localized. Causes include normal development, missing teeth, tooth-size differences, habits, or jaw and gum conditions.

### Widened Mediastinum

A radiographic finding on an upright posterior-anterior chest X-ray defined as mediastinal width exceeding 8 cm (or greater than twenty-five percent of thoracic width). It warrants urgent evaluation for acute aortic dissection, mediastinal hematoma,

---

mediastinitis, or mass lesions.

**Alternate Names:**

Mediastinal Widening; Mediastinal Enlargement

**Widened Pulse Pressure**

A pulse pressure (difference between systolic and diastolic blood pressure) greater than 60 mmHg or greater than fifty percent of the systolic value. It is commonly observed in aortic regurgitation, severe arteriosclerosis, thyrotoxicosis, fever, sepsis, and arteriovenous malformations.

**Wide Splitting of S2**

An abnormal prolongation of the interval between the aortic (A2) and pulmonic (P2) components of the second heart sound that varies physiologically with respiration but remains distinctly separated in expiration. It occurs due to delayed right ventricular emptying (such as right bundle branch block or pulmonary stenosis) or early aortic valve closure.

**Wilkie Syndrome**

Vascular compression of the third (horizontal) segment of the duodenum between the abdominal aorta and the overlying superior mesenteric artery (SMA). It leads to partial or complete duodenal obstruction, characterized by postprandial abdominal pain, early satiety, nausea, and vomiting.

**Alternate Names:**

Superior Mesenteric Artery Syndrome; SMA Syndrome; Arteriomesenteric Duodenal Compression

**Williams Syndrome**

A genetic condition caused by loss of a small section of chromosome 7, including a gene important for elastic tissues. It may cause characteristic facial features, heart-vessel narrowing, high blood calcium, developmental differences, learning difficulties, and connective-tissue problems. Regular heart, calcium, hearing, and developmental checks are important.

**Wilms Tumor**

A malignant embryonal tumor of the kidney in children. It may present as an abdominal mass, pain, hematuria, or an incidental finding and may be associated with syndromes such as WAGR or Beckwith-

Wiedemann syndrome. Treatment and prognosis depend on stage and tumor biology.

**Alternate Names:**

Nephroblastoma; Pediatric Renal Adenosarcoma

**Wilm's Tumour**

Source spelling variant of *Wilms tumor*, a malignant embryonal kidney tumor occurring mainly in young children.

**Wilson Disease**

An autosomal recessive disorder usually caused by disease-causing variants in the *ATP7B* gene, which encodes a protein needed to transport copper and excrete excess copper into bile. Copper accumulates in the liver and may later affect the brain, eyes, kidneys, blood, and other tissues. Manifestations include hepatitis, cirrhosis, acute or chronic liver failure, tremor, dystonia, poor coordination, speech changes, mood or behavior changes, and Kayser-Fleischer rings around the cornea.

**Wilson's Disease**

Alternate name; see *Wilson Disease*.

**Winchester Syndrome**

A very rare inherited skeletal dysplasia with progressive osteolysis, short stature, joint contractures, and characteristic facial or skin findings. It requires genetic, orthopedic, and multidisciplinary evaluation.

**Windkessel Effect**

The hemodynamic property of large elastic arteries, particularly the aorta, which expand during ventricular systole to store a portion of stroke volume and recoil during diastole. This elastic recoil sustains downstream blood flow and maintains organ perfusion throughout diastole.

**Alternate Names:**

Windkessel Function; Arterial Elastic Reservoir Effect

**Windowing**

Adjusting the brightness and contrast of a CT image

---

so particular tissues, such as bone, lung, or soft tissue, are easier to see.

### Windpipe

The tube carrying air from the voice box to the two main airways leading to the lungs. It is supported by firm rings that help keep it open.

### Winged Scapula

An abnormal musculoskeletal prominence of the medial border and inferior angle of the scapula protruding posteriorly from the thoracic cage, most commonly caused by traction injury, surgical damage, or neuropathy of the long thoracic nerve of Bell paralyzing the serratus anterior muscle.

#### Alternate Names:

Scapula Alata

### Winterbottom Sign

Prominent enlargement of the posterior cervical lymph nodes that serves as a classic clinical physical sign of early-stage African trypanosomiasis (sleeping sickness), caused by *Trypanosoma brucei*.

#### Alternate Names:

Winterbottom's Sign

### Winter Formula

An established physiological mathematical equation used to assess whether respiratory compensation is appropriate in a patient with pure metabolic acidosis. Calculated as  $\text{Expected } PaCO_2 = (1.5 \times [HCO_3^-]) + 8 \pm 2$ ; a measured  $PaCO_2$  higher than the calculated expected range signifies a concomitant primary respiratory acidosis (ventilatory failure), whereas a measured  $PaCO_2$  lower than expected indicates an accompanying primary respiratory alkalosis.

#### Alternate Names:

Winter's Compensation Formula; Acid-Base Compensation Equation

### Winter's Formula

A clinical physiological equation used to calculate expected respiratory compensation (arterial partial pressure of carbon dioxide,  $PCO_2$ ) in response to metabolic acidosis:  $PCO_2 = (1.5 \times [HCO_3^-]) + 8 \pm 2$ .

If measured  $PCO_2$  is higher or lower than calculated, a coexisting respiratory acidosis or alkalosis is present.

#### Alternate Names:

Winter Formula; Metabolic Acidosis Compensation Formula

### Winter Vomiting Disease

An acute gastroenteritis, usually caused by norovirus, characterized by sudden vomiting, diarrhea, and abdominal cramps.

### Wireless Motility Capsule

A swallowed capsule that records pressure, acidity, temperature, and movement as it travels through the digestive tract. The measurements estimate how long food takes to pass through the stomach and intestines and may help investigate unexplained constipation, diarrhea, nausea, or bloating.

### Wirsung Duct

The main excretory duct of the pancreas, extending from the tail through the body and head of the gland, joining the common bile duct at the hepatopancreatic ampulla of Vater to empty digestive enzymes into the second part of the duodenum at the major duodenal papilla.

#### Alternate Names:

Main Pancreatic Duct

### Wisdom Teeth

Wisdom teeth are the third molars that commonly emerge in late adolescence or adulthood. Impaction, infection, decay, cysts, gum disease, or crowding can cause pain and swelling; dental examination and imaging determine whether monitoring or removal is appropriate.

### Wisdom Tooth

The last molar in each dental quadrant, usually erupting in late adolescence or early adulthood. Impaction may cause pain, infection, or damage to adjacent teeth.

#### Alternate Names:

Dens Serotinus; Wisdom Teeth



---

### Wiskott-Aldrich Syndrome

A rare inherited immune disorder that mainly affects boys and causes eczema, repeated infections, and a low number of unusually small platelets. It can also cause bleeding and autoimmune problems and requires specialist care.

### Wiskott-Aldrich Syndrome Protein

A cytoplasmic actin-regulatory protein encoded by the *WAS* gene on the X chromosome that coordinates actin cytoskeleton remodeling in hematopoietic cells and immune synapse formation; loss-of-function mutations cause Wiskott-Aldrich syndrome and X-linked thrombocytopenia.

#### Alternate Names:

WASP

### Witch's Milk

A temporary milky breast discharge in a newborn caused by maternal hormones crossing the placenta; it usually resolves without squeezing or treatment, which can otherwise cause infection.

### Withdrawal

Physical or psychological symptoms that occur when a person who has developed dependence reduces or stops using a medicine or psychoactive substance. Symptoms vary by substance and may include anxiety, agitation, sweating, tremor, nausea, pain, insomnia, cravings, or seizures; some forms require urgent medical care.

#### Alternate Names:

Abstinence Syndrome; Substance Withdrawal Syndrome

### Withdrawal Reflex

An automatic response in which a painful stimulus makes a limb pull away from danger. The opposite limb may straighten to help maintain balance.

### Withdrawal Symptoms

Physical or psychological symptoms that occur when a person stops or reduces a substance or medicine after the body has adapted to it. The symptoms depend on the substance, duration of use, and the person's health.

### Withdrawal Syndrome

A predictable set of signs and symptoms that occur when a drug is discontinued or the dose is reduced after chronic use, reflecting the physiological rebound from drug adaptation.

### Withdrawal Syndromes

Physical and psychological symptoms that occur when regular exposure to an addictive substance is reduced or stopped. Severity and treatment depend on the substance, duration of use, comorbidities, and risk of seizures, delirium, or autonomic instability.

### Wolff-Chaikoff Effect

An autoregulatory mechanism whereby exposure to high levels of inorganic iodide temporarily inhibits organification and thyroid hormone synthesis within the thyroid gland. This effect typically lasts about 10 to 14 days before escape occurs, preventing iodine-induced hyperthyroidism.

#### Alternate Names:

Wolff-Chaikoff Block; Iodide Organification Inhibition

### Wolffian Ducts

Temporary tubes in the developing embryo that form parts of the male reproductive system, including the sperm-transport ducts. They usually shrink during typical female development, although small remnants may remain.

### Wolff-Parkinson-White Accessory Pathway-Related Tachycardia

Alternate name; see *Wolff-Parkinson-White syndrome*.

### Wolf-Hirschhorn Syndrome

A rare genetic disorder caused by loss of part of chromosome 4. It can cause poor growth, a small head, distinctive facial features, developmental and intellectual disability, seizures, and heart or other organ abnormalities; lifelong specialist support is usually needed.

#### Alternate Names:

4p- Syndrome; 4p Deletion Syndrome



---

## Wolfram Syndrome

A rare inherited disorder that commonly causes diabetes requiring insulin and progressive loss of vision from optic-nerve damage. It may also cause excessive thirst and urination, hearing loss, bladder problems, balance difficulty, or other nerve-system problems. Treatment monitors and supports each affected function.

## Wolinella

Genus of anaerobic, Gram-negative bacteria found in the oral cavity and gastrointestinal tract; associated with periodontal disease.

## Wolman Disease

A severe inherited disorder in which a missing enzyme causes fats to build up in organs. It usually begins in infancy with vomiting, diarrhea, poor growth, an enlarged liver and spleen, and adrenal-gland problems. Without treatment it can be fatal early; enzyme replacement and specialist care are required.

## Wolman's Disease

A severe lysosomal acid-lipase deficiency presenting in infancy with hepatosplenomegaly, adrenal calcification, malabsorption, vomiting, diarrhea, and failure to thrive. It is a severe form of lysosomal acid-lipase deficiency and can be fatal early without treatment.

## Womb

Common name for the uterus; the muscular organ where a fetus develops during pregnancy.

### Alternate Names:

Uterus; Metra; Hystera

## Womb Growths

Alternate name; see *Uterine fibroids*.

## Women Health

Alternate name; see *Women's health*.

## Women's Health

Medical specialty focusing on conditions unique to

or more prevalent in females, including reproductive health, pregnancy, menopause, and gender-specific disease patterns.

### Alternate Names:

Gynecological and Obstetric Health; Female Health Care

## Wood Lamp Examination

Examination of the skin in a dark room using special ultraviolet light. The light can make some color changes and infections easier to see, but the finding usually needs confirmation with the examination and other tests.

## Woods Corkscrew Maneuver

An emergency procedure used when a baby's shoulders become stuck during birth. A trained clinician rotates the baby's shoulders to help release them and complete delivery.

## Wood's Light

A special ultraviolet light used in a darkened room to make some skin colors, scales, and infections easier to see. It can support an examination but usually cannot identify the cause by itself.

## Wood Spirit

An old name for methanol, a toxic alcohol whose ingestion can cause metabolic acidosis, blindness, and death.

## Woods Screw Maneuver

A rotational maneuver used during shoulder dystocia to rotate the fetal shoulders and assist delivery.

## Wood-Unit

A clinical unit of vascular resistance calculated by dividing the pressure gradient across a vascular bed (in mmHg) by cardiac output (in L/min). One Wood unit equals approximately 80 dyn·s/cm<sup>5</sup>, commonly used in evaluating pulmonary hypertension.

### Alternate Names:

Wood Resistance Unit; Hybrid Resistance Unit (HRU)

---

### Woody Tongue

A rare long-lasting infection, usually caused by *Actinomyces*, that makes the tongue and nearby tissues hard and swollen. It can interfere with eating, speaking, or breathing.

### Woolly Hair

A rare structural anomaly of scalp hair characterized by tightly coiled, fine, spiraled curls in individuals of non-African descent. It may be inherited in autosomal dominant or recessive patterns and can occur as an isolated trait or part of syndromic disorders such as Naxos disease or Carvajal syndrome.

#### Alternate Names:

Woolly Hair Syndrome; Hereditary Woolly Hair

### Woolsorters' Disease

A form of inhaled anthrax acquired by breathing spores of *Bacillus anthracis*, historically associated with handling contaminated wool or animal products. It can cause severe breathing illness and requires urgent treatment.

#### Alternate Names:

Inhalation Anthrax; Pulmonary Anthrax; Rag-picker's Disease

### Word Blindness

Older term; see *Alexia*.

### Word Deafness

An acquired inability to understand spoken words even though ordinary hearing is intact. It usually results from damage to brain areas that connect hearing with language, such as after a stroke or head injury.

#### Alternate Names:

Pure Word Deafness; Auditory Agnosia for Speech

### Word Recognition Scoring

A hearing test measuring how accurately a person repeats or identifies spoken words at a comfortable volume. It helps assess how well the hearing system and brain understand speech.

### Word Salad

Speech made of words or phrases that have little or no logical connection, making the person difficult to understand. It can occur with severe psychosis, mania, brain disease, or intoxication and needs assessment in context.

### Working Memory

The ability to hold a small amount of information briefly and use it while completing a task, such as remembering instructions while following them. Attention, stress, sleep, illness, and brain conditions can affect it.

### Wound

Damage to body tissue caused by trauma, surgery, pressure, heat, chemicals, or another cause. A wound may be open or closed and can vary in depth, contamination, bleeding, and injury to nearby nerves, vessels, or organs.

### Wound Care

Assessment and treatment of a wound through cleansing, bleeding control, removal of dead or contaminated tissue when indicated, appropriate closure or dressing, pain control, and follow-up. Care also includes tetanus prevention, pressure relief, and treatment of infection or underlying conditions; antibiotics are reserved for selected contaminated wounds, bites, or infections.

### Wound Classification

A description of common injury types: an abrasion is a scrape, a laceration is a torn cut, a contusion is a bruise from blunt force, and an incised wound is a clean cut from a sharp object.

### Wound Complication

A problem that develops during healing, such as infection, separation of wound edges, bleeding, fluid collection, tissue death, or excessive scarring. Increasing pain, redness, swelling, drainage, fever, or wound opening should be assessed promptly.

### Wound Debridement

The surgical, enzymatic, autolytic, or mechanical re-

---

removal of foreign material, necrotic non-viable tissue, cellular debris, and infected slough from a wound bed to eliminate bacterial burden, reduce infection risk, and promote healthy granulation tissue formation.

### **Wound Dehiscence**

Partial or complete reopening of a surgical wound after it has been closed. It may follow infection, poor blood supply, diabetes, increased pressure, or strain on the wound. New separation, drainage, fever, severe pain, or visible deeper tissue needs prompt medical assessment.

### **Wound Evisceration**

Separation of a surgical wound with protrusion of internal organs or tissue through the opening. It is a surgical emergency requiring protection of the exposed tissue and urgent medical evaluation.

### **Wound Healing**

The body's process of stopping bleeding, controlling inflammation, replacing damaged tissue, covering the surface, and strengthening the repair. Blood supply, infection, nutrition, diabetes, medicines, smoking, wound depth, and movement can affect healing.

### **Wound Healing Phases**

Wound healing has overlapping stages: stopping bleeding, inflammation, building new tissue and skin, and strengthening the repair. The timing and result depend on the wound, blood supply, infection, nutrition, and general health.

### **Wound Infection**

Infection of a wound by microorganisms, causing findings such as increasing pain, redness, warmth, swelling, pus, fever, or delayed healing.

### **Wound Management**

Assessment and treatment of acute or chronic wounds through cleansing, debridement when indicated, moisture and exudate control, infection management, pressure relief, nutrition, and correction of underlying disease.

### **Wounds**

Injuries in which tissue is damaged by trauma, surgery, pressure, heat, chemicals, or other causes; wounds may be open or closed and may require cleansing, closure, or debridement.

### **Wound VAC**

A negative-pressure wound treatment in which a sealed dressing is connected to a pump that gently removes fluid and reduces swelling. It can support new tissue growth in selected complex wounds, but requires proper application and monitoring for bleeding, infection, or skin damage.

### **Wound Vacuum**

A therapeutic wound management system that applies localized controlled subatmospheric (negative) pressure to a sealed wound dressing via an open-cell polyurethane foam sponge, facilitating continuous fluid drainage, reducing tissue edema, increasing microvascular blood flow, and accelerating granulation tissue formation.

#### **Alternate Names:**

Negative Pressure Wound Therapy; NPWT; Vacuum-Assisted Closure; VAC

### **Wright Hyperabduction Test**

A clinical physical examination maneuver used to assess neurovascular compression in thoracic outlet syndrome, specifically involving the pectoralis minor space and retropectoral tunnel. With the patient seated, the examiner palpates the radial pulse while passively hyperabducting and externally rotating the arm beyond 90 degrees; a substantial diminution or complete obliteration of the radial pulse accompanied by paresthesias constitutes a positive test.

#### **Alternate Names:**

Wright Test; Hyperabduction Maneuver

### **Wright Stain**

A laboratory dye applied to a blood sample so different blood cells and abnormal cell changes can be seen and identified under a microscope.

---

### Wright Test

A clinical physical examination maneuver used in the assessment of thoracic outlet syndrome. The examiner palpates the radial pulse while the patient's arm is hyperabducted and externally rotated overhead; a marked diminution or obliteration of the radial pulse suggests neurovascular compression under the pectoralis minor tendon.

#### Alternate Names:

Wright Maneuver; Hyperabduction Test

### Wrinkle Test

A simple clinical bedside test to evaluate peripheral sympathetic autonomic nerve function in the digits. Denervated skin fails to develop normal physiological wrinkling when submerged in warm water for 10 to 30 minutes, indicating loss of sympathetic vasoconstrictor tone.

#### Alternate Names:

Digital Wrinkling Test; Water Immersion Wrinkle Test

### Wrisberg Ganglion

A small cluster of nerve cells connected with the facial nerve. It helps relay taste signals from part of the tongue and sensory signals from nearby areas of the ear.

### Wrist

The joint connecting the hand to the forearm. It contains eight small wrist bones, ligaments, tendons, nerves, and blood vessels and allows the hand to bend, turn, and move from side to side.

### Wrist Arthroscopy

A procedure using a small camera and instruments inserted through tiny wrist openings to examine or treat damaged cartilage, ligaments, joint lining, or other wrist problems.

### Wrist Drop

Inability to lift the wrist and fingers, usually caused by injury or compression of the radial nerve. It may cause weakness, numbness, or difficulty using the hand and needs assessment for the cause.

### Wrist Fracture

A break in one or more bones near the wrist, most commonly the distal radius or ulna, usually caused by a fall, direct trauma, or weakened bone.

### Wrist Injury

Damage to the bones, ligaments, tendons, nerves, or vessels of the wrist from a fall, twist, impact, or overuse; fracture, instability, severe swelling, or neurovascular symptoms require assessment.

### Wrist Joint

The joint where the forearm bones meet the small bones of the hand. It allows bending, straightening, side-to-side movement, and limited rotation while ligaments and cartilage provide stability.

### Wrist Pain

Pain in or around the wrist, commonly caused by sprain, fracture, tendinitis, arthritis, or nerve compression such as carpal tunnel syndrome. Persistent pain, deformity, numbness, or weakness requires assessment.

#### Alternate Names:

Carpal Pain; Radiocarpal Joint Discomfort

### Wrist Sprain

Stretching or tearing of one or more ligaments supporting the wrist, causing pain, swelling, bruising, and reduced movement after injury.

### Writer's Cramp

A movement disorder in which involuntary muscle tightening causes abnormal hand or arm positions while writing. It may also affect other closely related fine hand movements.

### Written Documentation

Formal recording of patient information, clinical decisions, and treatment plans; essential for continuity of care and medicolegal purposes.

### Wry-Neck

Torticollis, an abnormal tilt or rotation of the head caused by involuntary contraction, shortening, or

---

dysfunction of neck muscles.

### **Wt**

A medical abbreviation commonly meaning weight. Its meaning should be confirmed from the surrounding documentation because abbreviations can be ambiguous.

### **Wuchereria**

A genus of filarial nematodes that includes *Wuchereria bancrofti*, a major cause of lymphatic filariasis. Infection is transmitted by mosquitoes and may cause lymphangitis, chronic lymphedema, hydrocele, and elephantiasis.

### **Wuchereria Bancrofti**

A parasitic roundworm spread by mosquitoes and the main cause of lymphatic filariasis. Long-term infection can block lymph drainage and cause swelling of the limbs or genitals, sometimes becoming severe and permanent.

### **Wuhan Coronavirus**

Alternate name; see *SARS-CoV-2*.

### **Wandering Rash Tongue**

A common name for geographic tongue, a benign inflammatory condition in which smooth red patches with pale borders move across the tongue. It is not contagious and may cause sensitivity to spicy or acidic foods.

### **Warts and Plantar Warts**

Warts are noncancerous skin growths caused by human papillomavirus. Plantar warts occur on the soles of the feet and may be painful when standing or walking because pressure pushes them inward.

### **WBS**

Abbreviation for Williams-Beuren syndrome, a genetic disorder associated with characteristic facial features, developmental differences, cardiovascular disease, and sometimes high blood calcium.

### **Weight Management**

Long-term care of body weight through nutrition, physical activity, behavior change, medicines, or bariatric procedures when appropriate. Goals should account for health, function, metabolic risk, and sustainable habits rather than weight alone.

### **Weir-Mitchell Disease**

An older name for restless legs syndrome, a neurologic disorder causing an urge to move the legs with uncomfortable sensations that are worse at rest and in the evening and improve temporarily with movement.

### **Werlhof Disease**

An older name for immune thrombocytopenia (immune thrombocytopenic purpura), in which immune destruction of platelets can cause bruising, petechiae, nosebleeds, heavy menstrual bleeding, or other bleeding.

### **West Nile Fever**

A febrile illness caused by West Nile virus, usually transmitted by mosquitoes. Most infections cause no symptoms or a flu-like illness; a small proportion causes neuroinvasive disease such as meningitis, encephalitis, or acute flaccid paralysis.

### **Whooping Cough (Pertussis)**

Alternate name for whooping cough, a highly contagious respiratory infection caused by *Bordetella pertussis*. Prolonged coughing fits may cause vomiting or a whoop; infants may instead have pauses in breathing.

### **WHS**

Abbreviation for Wolf-Hirschhorn syndrome, a rare chromosome 4 deletion disorder that may cause poor growth, distinctive facial features, developmental disability, seizures, and congenital abnormalities.

### **Williams-Beuren Syndrome**

Alternate name for Williams syndrome, a chromosome 7 microdeletion disorder associated with characteristic facial features, supravalvular aortic stenosis and other cardiovascular findings, developmental

---

differences, and hypercalcemia in some people.

### **Wisdom Tooth Problems**

Problems involving third molars, including impaction, infection, gum inflammation, decay, cysts, or damage to nearby teeth. Pain, swelling, fever, difficulty opening the mouth, or trouble swallowing needs dental assessment.

### **Wisdom Tooth Removal**

Dental extraction of one or more third molars, commonly performed when they are impacted, repeatedly infected, decayed, or damaging adjacent teeth. Risks include pain, infection, bleeding, dry socket, and temporary or rarely persistent nerve symptoms.

### **Wolf Hirschhorn Syndrome**

Alternate spelling of Wolf-Hirschhorn syndrome, a rare disorder caused by deletion of part of chromosome 4 and associated with growth restriction, seizures, developmental disability, and distinctive facial features.

### **Wolf Syndrome**

Alternate name for Wolf-Hirschhorn syndrome, a chromosome 4 deletion disorder affecting growth, development, and multiple organ systems.

### **WPW Syndrome**

Abbreviation for Wolff-Parkinson-White syndrome, a pre-excitation disorder in which an extra electrical pathway can cause episodes of supraventricular tachycardia. Palpitations, fainting, chest pain, or severe breathlessness require prompt assessment.

### **Wells Syndrome**

Eosinophilic cellulitis, an uncommon inflammatory skin disorder causing itchy or burning red plaques, swelling, blisters, or a cellulitis-like eruption without typical bacterial infection. Blood eosinophilia and a skin biopsy may support the diagnosis, and recurrences are possible.

### **Wolff-Parkinson-White Syndrome**

A heart-rhythm disorder caused by an extra elec-

trical connection between the atria and ventricles. It can produce episodes of supraventricular tachycardia, palpitations, dizziness, fainting, or, rarely, dangerous ventricular rhythms; catheter ablation can often eliminate the pathway.



# X

## **Xanthelasma**

A harmless yellowish patch or bump caused by fat deposits in the skin of the eyelids, usually near the inner corners of the eyes. It can occur with high blood fats but may also occur when blood-fat levels are normal.

### **Alternate Names:**

Xanthelasma Palpebrarum; Eyelid Xanthoma; Palpebral Xanthoma

## **Xanthine**

A natural breakdown product of substances found in cells and foods. It normally changes into uric acid, but can build up when a rare enzyme disorder blocks this pathway and may then contribute to kidney or urinary stones.

## **Xanthine Oxidase**

An enzyme that helps convert cell and food breakdown products into uric acid. Medicines used for gout can block this enzyme, lowering uric-acid production and helping prevent urate crystal deposits.

## **Xanthine Oxidase Inhibitors**

Medicines such as allopurinol and febuxostat that lower uric acid by blocking an enzyme involved in its production. They are used for selected people with gout or to prevent high uric acid during some cancer treatments. Choice and timing depend on the person's condition.

### **Alternate Names:**

XOIs

## **Xanthinuria**

A rare inherited disorder in which the body cannot

properly change xanthine into uric acid. Xanthine can build up and form kidney or urinary stones; muscle problems may also occur.

## **Xanthochromia**

Yellow or amber color in the fluid around the brain and spinal cord. It can occur when blood has broken down in the fluid, including after bleeding around the brain, but other causes are possible and the meaning depends on timing and other test results.

### **Alternate Names:**

CSF Xanthochromia; Cerebrospinal Fluid Yellowing

## **Xanthogranulomatous**

Describing inflammation in which tissue contains many immune cells filled with fat. This pattern can occur in long-lasting infections and other inflammatory conditions.

## **Xanthogranulomatous Cholecystitis**

A rare, long-lasting inflammation of the gallbladder that can thicken and damage its wall. It may spread to nearby tissues and can look like gallbladder cancer on scans.

### **Alternate Names:**

XGC

## **Xanthogranulomatous Oophoritis**

A rare long-lasting inflammation of an ovary, often linked to infection or endometriosis. It can form a mass that resembles ovarian cancer on scans and may cause pelvic pain, fever, or abnormal bleeding.

## **Xanthogranulomatous Pyelonephritis**

A severe, long-lasting kidney infection that destroys

---

normal kidney tissue and replaces it with inflamed tissue. It is often associated with blockage or stones and may require antibiotics, drainage, or removal of the damaged kidney.

### **Xanthoma**

A yellow-orange spot, bump, or patch formed by fat deposits in the skin, tissue beneath the skin, or tendons. It may occur with high blood fats, diabetes, some liver diseases, or an inherited lipid disorder.

### **Xanthoma Disseminatum**

A rare disorder in which many yellow, red-brown, or orange bumps develop on the skin, especially in body folds. The mouth, throat, or other organs may also be affected, and some people develop problems controlling water balance.

#### **Alternate Names:**

Montgomery Syndrome; Disseminate Xanthoma

### **Xanthoma Eruptivum**

A sudden outbreak of many small yellow-orange bumps, usually on the buttocks, arms, legs, or shoulders. It is often linked to very high blood triglyceride levels.

#### **Alternate Names:**

Eruptive Xanthoma; Eruptive Xanthomatosis

### **Xanthoma Planum**

Flat or slightly raised yellow skin patches caused by the buildup of fat inside immune cells. They may occur in skin folds, the palms, or more widely on the body and can be linked to high blood fats or some blood disorders.

#### **Alternate Names:**

Plane Xanthoma; Xanthoma Striatum Palmare

### **Xanthomas**

Yellowish papules, plaques, or nodules caused by lipid accumulation in skin immune cells. They may reflect high cholesterol or triglycerides, diabetes, cholestatic liver disease, or an inherited lipid disorder and should prompt appropriate evaluation.

### **Xanthoma Striatum Palmare**

Yellow or orange lines or patches in the creases of the palms caused by fat deposits in the skin. They are strongly linked to some inherited blood-fat disorders, especially type III dysbetalipoproteinemia, but blood tests are needed.

#### **Alternate Names:**

Palmar Crease Xanthoma; Planar Palmar Xanthoma

### **Xanthoma Tendinosum**

Firm, usually painless fat deposits in tendons or ligaments, most often in the Achilles tendon or tendons of the hands. They are strongly linked to inherited high cholesterol.

#### **Alternate Names:**

Tendon Xanthoma; Tendinous Xanthoma

### **Xanthomatosis**

A disorder in which fatty deposits called xanthomas develop in the skin, tendons, or organs. It may be linked to inherited or acquired problems with blood fats.

### **Xanthoma Tuberosum**

Slowly enlarging, painless, firm, yellowish-red nodular lipid deposits occurring over extensor pressure areas, particularly the elbows, knees, heels, and buttocks. They are strongly associated with familial hypercholesterolemia and type III hyperlipoproteinemia.

#### **Alternate Names:**

Tuberous Xanthoma; Tuberous Xanthomatosis

### **Xanthopsia**

A visual disturbance in which objects appear yellow or yellowish. It can occur with some eye disorders, medicine toxicity such as digoxin toxicity, or disorders of the nerves or retina.

### **Xanthosis**

Yellow discoloration of skin or other tissues caused by carotene or another yellow substance. Unlike jaundice, it does not usually make the whites of the eyes yellow; the cause is assessed from the person's

---

diet and health.

### **X Chromosome**

One of the two types of human sex chromosomes. It carries genes involved in sex development as well as many other body functions. Most people have one or two X chromosomes.

### **Xenobiotic**

A chemical that does not naturally belong in the body, such as a medicine, pollutant, pesticide, or industrial compound. The body changes and removes it, but some resulting chemicals can be harmful.

### **Xenobiotic Metabolism**

The body's processing of foreign chemicals, including medicines, pollutants, and toxins. Enzymes, especially in the liver, change them so they can be removed; the changed substances may remain active or become harmful.

### **Xenobiotic Resistance**

The ability of cells or organisms to withstand the effects of foreign chemicals, medicines, or toxins. It may result from breaking down the substance faster, pumping it out of cells, or changing the substance's target.

#### **Alternate Names:**

Multidrug Resistance; Xenobiotic Tolerance

### **Xenograft**

Tissue, cells, or a biological material transplanted between different species, such as animal tissue used in a human. Rejection and infection are important risks, although some materials are used temporarily as tissue coverings or substitutes.

### **Xenon Anesthesia**

Use of xenon gas to cause unconsciousness and prevent pain during an operation. Xenon acts on brain signaling and is cleared mainly through breathing.

### **Xenotransplantation**

Transplanting living cells, tissues, or organs between different species, usually from an animal to a human.

It may increase the supply of transplant material but carries major risks of immune rejection, infection, and poor organ function.

### **Xerochilia**

Abnormal, persistent dryness, scaling, and superficial vertical cracking or fissuring of the lips, resulting from environmental dehydration, persistent lip-licking (cheilitis simplex), chronic mouth breathing, or systemic therapy with oral retinoids (such as isotretinoin).

#### **Alternate Names:**

Dry Lips

### **Xeroderma**

Alternate name; see *Dry skin*.

### **Xeroderma Pigmentosum**

A rare usually autosomal-recessive disorder of nucleotide-excision DNA repair, with a variant form affecting translesion synthesis. Minimal ultraviolet exposure can cause severe sunburn or persistent redness, early freckle-like pigmentation, and rapid photoaging; skin and ocular-surface cancers occur at a young age, and some people develop progressive neurologic disease. Strict ultraviolet protection, regular skin and eye examinations, and early treatment of suspicious lesions are essential.

### **Xeroderma Pigmentosum Complementation Groups**

Genetic subtypes of xeroderma pigmentosum. Each subtype results from a change in a different gene involved in repairing damage caused by ultraviolet light. All subtypes can cause extreme sun sensitivity, freckling, and a very high risk of skin cancer.

#### **Alternate Names:**

XP Complementation Groups; NER Defect Groups

### **Xerophthalmia**

Dryness and damage of the eye caused by severe vitamin A deficiency. It may begin with poor night vision and dry eyes and can progress to corneal damage and permanent sight loss.

#### **Alternate Names:**

Vitamin A Deficiency Eye Disease; Ocular Xerosis;

---

Keratomalacia

### Xerophthalmic Ulceration

Severe progressive ulceration of the cornea resulting from severe vitamin A deficiency (classified as WHO stage X3A when involving less than one-third of the corneal area, and stage X3B when involving more). Untreated, it rapidly causes corneal liquefactive necrosis (keratomalacia) and permanent blindness.

#### Alternate Names:

Vitamin A Corneal Ulcer; Keratomalacic Ulceration

### Xerosis

Alternate name; see *Dry skin*.

### Xerostomia

Alternate name; see *Dry Mouth*.

### Xerotic Eczema

A common, intensely pruritic inflammatory dermatosis occurring predominantly on the pretibial lower extremities of elderly individuals during cold, dry winter months, characterized by profound epidermal dehydration, scaling, and fine superficial erythematous fissures resembling "cracked porcelain" or a "dry riverbed."

#### Alternate Names:

Asteatotic Eczema; Eczema Craquele; Winter Itch

### X-Inactivation

A process in early development that switches off most genes on one of two X chromosomes in each cell. It helps balance gene activity, although some genes remain active and the pattern can differ between cells.

### Xiphisternal Joint

The joint between the lower part of the breastbone and the xiphoid process. It often becomes partly or fully hardened with age.

#### Alternate Names:

Xiphisternal Synchondrosis; Xiphisternal Articulation

### Xiphoidalgia

Pain or tenderness over the xiphoid process, the small lower part of the breastbone. Pain may spread to the upper abdomen, chest, back, or shoulders and may worsen with touch, bending, lifting, or coughing.

#### Alternate Names:

Xiphoidodynia; Hypersensitive Xiphoid Syndrome; Xiphoid Syndrome

### Xiphoidectomy

Surgery to remove the xiphoid process, the small lower part of the breastbone. It may be done for persistent pain in this area or to provide access during some chest or abdominal operations.

### Xiphoid Process

The small pointed lower part of the breastbone. Its shape varies between people and it provides attachment for some muscles and ligaments.

### X-Linked

Describing a gene or condition caused by a change on the X chromosome. The pattern differs between people with one X chromosome and those with two; fathers pass their X chromosome to daughters, not sons.

### X-Linked Adrenoleukodystrophy

An inherited disorder, usually affecting boys, in which certain fats build up and damage the adrenal glands and nervous system. It can cause behavior or learning changes, movement problems, seizures, or adrenal failure.

### X-Linked Agammaglobulinemia

An inherited immune disorder, usually affecting boys, in which the body cannot make enough antibodies. Repeated bacterial infections often begin in infancy or early childhood and are treated with antibody replacement and infection care.

### X-Linked Chronic Granulomatous Disease

An inherited immune disorder, usually affecting boys, caused by a change in the *CYBB* gene. Cer-

---

tain white blood cells cannot kill some bacteria and fungi normally, leading to repeated serious infections and areas of ongoing inflammation.

**Alternate Names:**

X-Linked CGD; CYBB Deficiency; Cytochrome b-245 Beta Deficiency

**X-Linked Dominant Inheritance**

A pattern in which one changed gene on the X chromosome can cause a condition. An affected father passes the changed X chromosome to all daughters and no sons.

**X-Linked Emery-Dreifuss Muscular Dystrophy**

An inherited muscle disorder, usually affecting boys, that causes stiff joints, slowly worsening muscle weakness, and problems with the heart's electrical rhythm.

**Alternate Names:**

EDMD1; Emerin Deficiency Muscular Dystrophy

**X-Linked Hyper-IgM Syndrome**

An inherited immune disorder, usually affecting boys, in which the body cannot change one type of protective antibody into other types. It causes repeated lung and sinus infections and may cause infections by organisms that usually do not cause illness.

**Alternate Names:**

HIGM1; CD40L Deficiency; Hyper-IgM Syndrome Type 1

**X-Linked Hypophosphatemia**

An inherited disorder that causes the kidneys to lose too much phosphate. Low phosphate can lead to soft or poorly growing bones, bone pain, bowed legs, dental problems, and reduced height.

**X-Linked Ichthyosis**

An inherited, usually X-linked recessive skin disorder caused by a variant or deletion affecting the STS gene and deficiency of steroid sulfatase. It usually affects males and causes widespread dry, dark, plate-like scales that often begin in infancy or early childhood.

**X-Linked Inheritance**

Inheritance of a gene change located on the X chromosome. The effects depend on the gene, the type of change, and whether a person has one or two X chromosomes; fathers do not pass an X chromosome to sons.

**X-Linked Lymphoproliferative Disease Type 1**

An inherited immune disorder, usually affecting boys, in which the body can react severely to Epstein-Barr virus. It may cause severe glandular fever, very high inflammation, low antibody levels, or lymphoma.

**Alternate Names:**

XLP1; Duncan Disease; SAP Deficiency

**X-Linked Lymphoproliferative Syndrome**

An inherited immune disorder, usually affecting boys, in which the body reacts abnormally to infections. Severe Epstein-Barr virus illness, persistent fever, liver or blood problems, lymphoma, and intestinal inflammation may occur.

**X-Linked Myotubular Myopathy**

A rare inherited muscle disease, usually affecting boys, that causes severe weakness and low muscle tone from birth. Babies may have drooping eyelids, feeding difficulty, and dangerous weakness of the breathing muscles.

**Alternate Names:**

XLMTM; Myotubularin Deficiency

**X-Linked Recessive Inheritance**

A pattern in which a changed gene on the X chromosome usually causes a condition in people with one X chromosome. People with two X chromosomes are often unaffected carriers, although they can sometimes have symptoms. An affected father does not pass the changed X chromosome to sons.

**X-Linked Retinitis Pigmentosa**

An inherited disorder in which the retina slowly loses its ability to sense light. It usually causes night blindness followed by narrowing of the field of vision and worsening sight.

---

**Alternate Names:**

XLRP; X-Linked Pigmentary Retinopathy

**X-Linked Retinoschisis**

An inherited eye disorder, usually affecting boys, in which layers of the retina separate. It can cause blurred central or distance vision, crossed eyes, bleeding, or retinal detachment in severe cases.

**X-Linked Severe Combined Immunodeficiency**

A severe inherited immune disorder, usually affecting boys, in which important immune cells do not develop normally and antibody protection is poor. Babies are highly vulnerable to serious infections.

**X-Linked Sideroblastic Anemia**

An inherited form of anemia, usually affecting boys, in which red blood cells cannot make hemoglobin normally. Iron builds up in developing blood cells, causing small, pale red blood cells and anemia.

**Alternate Names:**

XLSA; ALAS2 Deficiency; Hereditary Sideroblastic Anemia

**X-Linked Spinal and Bulbar Muscular Atrophy**

An inherited muscle and nerve disorder that usually begins in adulthood and slowly causes weakness. It may also cause muscle twitching, trouble speaking or swallowing, breast enlargement, and reduced fertility.

**Alternate Names:**

Kennedy Disease; SBMA; Bulbo-Spinal Muscular Atrophy

**X-Ray**

An imaging test that uses a small amount of radiation to create a picture of structures inside the body. It is especially useful for bones, lungs, teeth, and some other tissues.

**X-Ray Attenuation**

The reduction in strength of an X-ray beam as it passes through the body. Bone, soft tissue, and air

reduce the beam by different amounts, creating the differences seen in the image.

**X-Ray Crystallography**

A laboratory method that uses X-rays to work out the three-dimensional structure of molecules such as proteins. The molecule is made into a crystal, and the pattern produced by the X-rays is analyzed.

**Alternate Names:**

X-Ray Crystal Diffraction Analysis; Macromolecular Crystallography

**X-Ray Shield**

Protective material, often lead or another dense substance, used to reduce exposure to scattered X-rays during imaging.

**X-Ray Therapy**

Treatment that uses carefully aimed X-ray radiation to damage abnormal cells or relieve symptoms. It is used mainly for cancer and may be given from a machine outside the body or near the target.

**XX Male Syndrome**

A difference in sex development in which a person with two X chromosomes develops typically male physical features, usually because a sex-development gene has moved onto an X chromosome. Infertility is common, and care is individualized.

**XXX Syndrome**

Alternate name; see *Triple X syndrome*, a sex-chromosome variation in which a female has an additional X chromosome.

**XXXX Syndrome**

XXXX syndrome is a rare sex-chromosome variation in which a female has four X chromosomes; it may cause developmental or learning difficulties, distinctive physical features, and ovarian dysfunction, with severity varying widely.

**Xylitol**

A sugar alcohol used as a sweetener and in some oral-care products. Large human ingestions may



---

cause gastrointestinal symptoms; it is highly toxic to dogs.

### **Xylose**

A simple sugar found in some plants. A form called D-xylose has been used in tests of how well the small intestine absorbs nutrients and is also used in laboratory studies.

### **Xylose Absorption Test**

A test of how well the small intestine absorbs a simple sugar called D-xylose. Low amounts in the blood or urine may suggest poor absorption from the intestine; normal results with other signs of poor absorption may suggest a problem with the pancreas.

#### **Alternate Names:**

D-Xylose Absorption Test; D-Xylose Tolerance Test

### **XXY Syndrome**

A sex-chromosome variation in which a male has an extra Y chromosome. Many affected individuals have few or no obvious symptoms, though some have tall stature, learning difficulties, or developmental differences.

### **X-linked Juvenile Retinoschisis**

An inherited retinal disorder, usually affecting boys, in which layers of the retina separate during childhood. It can cause reduced central vision, difficulty seeing in dim light, crossed eyes, or retinal complications.

### **X-linked Adult Spinal Muscular Atrophy**

Alternate name for Kennedy disease, or X-linked spinal and bulbar muscular atrophy. It is an inherited androgen-receptor disorder that gradually causes weakness, muscle wasting, twitching, swallowing or speech difficulty, and sometimes breast enlargement.

### **X-linked Intellectual Disability**

Intellectual disability caused by a change in a gene on the X chromosome. Features vary widely and may include developmental delay, learning difficulty, behavior differences, seizures, or associated physical

findings.

### **XO Syndrome**

A form of Turner syndrome caused by complete or partial absence of one X chromosome. It may cause short stature, ovarian insufficiency, infertility, characteristic physical findings, and heart, kidney, hearing, or thyroid problems.

### **XP**

Abbreviation for xeroderma pigmentosum, a rare DNA-repair disorder causing extreme sensitivity to ultraviolet light and a markedly increased risk of early skin and eye cancers.

### **XXY Males**

People with a typically male pattern of development and an extra X chromosome, usually 47,XXY. This is commonly called Klinefelter syndrome and may cause small testes, low testosterone, infertility, tall stature, or learning and language difficulties.

### **XXY Syndrome**

Alternate name for Klinefelter syndrome, a sex-chromosome variation in which a typically male individual has an extra X chromosome. Effects vary and may include low testosterone, infertility, small testes, tall stature, and developmental or language differences.



# Y

## YAG Capsulotomy

A laser procedure that makes a small opening in the cloudy membrane behind an artificial lens after cataract surgery. It can restore vision when this clouding causes blurring or glare and is usually done without another incision.

## YAG Laser Capsulotomy

Alternate name; see *YAG Capsulotomy*.

## YAG Laser Surgery

Treatment using a focused laser to make a precise opening or alter tissue without a conventional surgical incision. It is used mainly for selected eye conditions, and risks depend on the part of the eye treated.

## Yawn

An involuntary behavior consisting of a deep inhalation followed by a slower exhalation, often with stretching of the jaw and facial muscles. Yawning is common with fatigue but excessive or unusual yawning can occur with some medications or neurologic conditions.

## Yawning

Alternate name; see *Yawn*.

## Yaws

Yaws is a chronic infection caused by *Treponema pallidum* subsp. *pertenue* and spread mainly through close skin contact. It can cause an initial papilloma or ulcer and later destructive lesions of skin, bone, and cartilage; antibiotics can cure the infection.

## Yaws Papilloma

The primary clinical cutaneous lesion of early yaws (*Treponema pallidum* subsp. *pertenue* infection), also termed a mother yaw or frambesioma. It presents as an initial painless, raised, yellow-crust, raspberry-like papule or ulcer at the site of cutaneous inoculation.

### Alternate Names:

Mother Yaw; Frambesioma; Primary Yaws Lesion

## Y-Box Binding Protein

A protein that attaches to DNA and RNA and helps control which genes are active and how proteins are made. It also helps cells respond to stress and repair damaged DNA.

### Alternate Names:

YB-1; Y-Box Transcription Factor; Nuclease-Sensitive Element-Binding Protein 1

## Y Chromosome

One of the two human sex chromosomes. It usually carries genes involved in testis development and is passed by a biological father to a child who inherits it; chromosome patterns and body development can vary.

## Y-Chromosome Microdeletion

A small missing section of the Y chromosome that can interfere with sperm production and cause very low or absent sperm counts.

### Alternate Names:

AZF Microdeletion

## Yeast

A type of fungus made of single cells. Some yeasts

---

normally live on or in the body, while others can cause infections of the mouth, skin, vagina, lungs, blood, or other organs.

### **Yeast Infection**

An infection caused by *Candida* yeast that may affect the mouth, skin, vagina, penis, or other tissues. Symptoms depend on the site and may include itching, soreness, redness, white plaques, discharge, or rash. *Candida* may also be present without causing disease, so the significance of a finding depends on the site, symptoms, and evidence of tissue invasion.

### **Yeast Infection In Women And Men**

Genital *Candida* infection may affect the vulva or vagina in women and the glans penis or foreskin in men. Symptoms can include itching, soreness, redness, discharge, or rash.

### **Yeast Present**

A laboratory report showing yeast in a specimen. This may mean harmless colonization, contamination, or infection; the meaning depends on the symptoms, specimen, and type of yeast.

### **Yeast Vulvitis**

Inflammation of the vulva associated with a *Candida* yeast infection. It may cause itching, burning, redness, swelling, and soreness; other conditions can cause similar symptoms.

### **Yellow Cartilage**

A flexible type of cartilage containing many elastic fibers. It is found in the outer ear, the epiglottis, parts of the voice box, and the tube connecting the middle ear to the throat.

#### **Alternate Names:**

Elastic Cartilage; Cartilago Elastica

### **Yellow Fever**

Yellow fever is an acute mosquito-borne infection caused by yellow fever virus. Fever, headache, muscle aches, nausea, and vomiting may be followed by jaundice, bleeding, shock, and organ failure; vaccination is the main preventive measure.

### **Yellow Fever Vaccine**

A vaccine made from a weakened yellow fever virus. It gives long-lasting protection, but rare serious reactions can affect the brain or internal organs.

#### **Alternate Names:**

17D Vaccine

### **Yellow Fever Virus**

A flavivirus transmitted mainly by infected mosquitoes that can cause yellow fever, ranging from a mild febrile illness to jaundice, bleeding, shock, and organ failure.

### **Yellow Ligament**

A series of elastic bands connecting the back parts of neighboring bones in the spine. They help support the spine and surround the spinal canal.

#### **Alternate Names:**

Ligamentum Flavum; Flavum Ligament

### **Yellow Marrow**

Bone marrow containing mostly fat instead of blood-forming cells. It is common in the hollow centers of adult long bones and can change back toward blood-forming marrow when the body needs more blood cells.

#### **Alternate Names:**

Fatty Bone Marrow; Medulla Ossium Flava

### **Yellow Nail Syndrome**

Alternate name; see *Yellow-Nail Syndrome*.

### **Yellow-Nail Syndrome**

A rare disorder causing yellow, thickened, slowly growing nails together with swelling from lymph fluid, usually in the legs, and lung problems such as repeated infections or fluid around the lungs. The cause is not fully understood.

### **Yellow Nail Syndrome Triad**

The three main features of yellow nail syndrome: slow-growing yellow or thickened nails, swelling from lymph-fluid buildup, and long-term breathing or lung problems.

#### **Alternate Names:**

---

YNS Triad; Lymphedema-Pleural Effusion-Yellow Nails

### **Yellow Spot**

The macula, a small yellowish area near the center of the retina at the back of the eye. It provides the sharp central vision used for reading and seeing fine detail.

#### **Alternate Names:**

Macula Lutea; Anatomical Macula

### **Yellow Tongue**

Yellow discoloration or coating of the tongue that may result from oral hygiene, smoking, dry mouth, diet, medicines, or an underlying illness.

### **Yergason Test**

A shoulder examination in which a person turns the palm upward and rotates the arm outward against resistance. Pain or a snapping sensation near the front of the shoulder may suggest a problem involving the biceps tendon, but the test is not diagnostic by itself.

### **Yersinia**

A group of bacteria that can cause intestinal illness, swollen lymph nodes in the abdomen, or plague. Different species spread through contaminated food or water, infected animals, or fleas.

### **Yersinia Enterocolitica**

A bacterium that can cause diarrhea, fever, and abdominal pain, sometimes mimicking appendicitis. People usually acquire it from contaminated food, water, or contact with infected animals.

### **Yersinia Enterocolitica Pseudoappendicitis**

A Yersinia infection that causes pain in the lower right abdomen and swollen lymph nodes near the intestine, making it resemble appendicitis. It can also cause fever and diarrhea, often after contaminated food such as undercooked pork.

#### **Alternate Names:**

Yersinia Mesenteric Adenitis; Pseudoappendiceal

Syndrome

### **Yersinia Infection**

Infection caused by Yersinia bacteria, usually producing diarrhea, abdominal pain, fever, or vomiting. Some cases resemble appendicitis; bloody diarrhea, dehydration, or severe or persistent illness can occur.

### **Yersinia Outer Proteins**

Proteins made by disease-causing Yersinia bacteria that are injected into human cells. They weaken the immune response and help the bacteria avoid being destroyed.

#### **Alternate Names:**

Yops; Yersinia Effector Proteins; Yop Virulence Factors

### **Yersinia Pestis**

The bacterium that causes plague, a severe infection usually spread by infected fleas or animals. It may cause painful swollen lymph nodes, bloodstream infection, or lung disease and can be life-threatening.

### **Yersinia Pestis Pneumonic Plague**

A severe form of plague affecting the lungs. It can spread through breathing in infected droplets or when plague spreads from another part of the body. It may cause high fever, breathing difficulty, chest pain, cough, shock, and rapidly worsening lung failure.

#### **Alternate Names:**

Pneumonic Plague; Pulmonary Yersiniosis

### **Yersinia Pseudotuberculosis**

A bacterium usually acquired from contaminated food or water. It can cause fever, diarrhea, and pain from swollen lymph nodes in the abdomen, sometimes resembling appendicitis.

### **Yersinia Reactive Arthritis**

Joint inflammation that develops after a Yersinia intestinal infection, usually one to four weeks later. It often affects a few joints, especially in the legs, without bacteria being present inside the joints.

---

## Yersiniosis

Infection caused by *Yersinia* bacteria, most often *Y. enterocolitica* or *Y. pseudotuberculosis*, producing diarrhea, abdominal pain, fever, or pseudoappendicitis.

## Yew Poisoning

Poisoning after eating parts of the yew plant, especially its leaves or seeds. The toxins can cause vomiting, dizziness, seizures, dangerous heart rhythms, collapse, and death.

## Y-Linked Gene

A gene located specifically on the non-recombining region of the Y chromosome (such as the *SRY* gene, *ZFY*, or *AZF* azoospermia factor region). These genes are transmitted strictly from biological fathers to sons and are essential for testicular differentiation, gonadal development, and male spermatogenesis.

### Alternate Names:

Holandric Gene; Y-Chromosomal Gene

## Y-Linked Inheritance

Inheritance of a condition caused by a gene on the Y chromosome. A biological father passes a Y-linked gene to sons who inherit his Y chromosome, but not to daughters who inherit his X chromosome.

## Y-Maze Test

A test used mainly in laboratory animals to study movement, exploration, and short-term memory. The animal moves through three arms of a Y-shaped maze, and the pattern of movement is recorded.

### Alternate Names:

Y-Maze Spontaneous Alternation; Y-Maze Behavioral Paradigm

## Yolk Sac

A temporary structure that supports an early embryo before the placenta is fully working. It helps form the first blood cells and part of the early intestine and can be seen on early pregnancy ultrasound.

## Yolk Sac Carcinoma

An aggressive germ-cell cancer that often produces

high levels of alpha-fetoprotein (AFP). AFP can support the diagnosis and help track treatment, but it is interpreted with examination, imaging, and tissue testing.

### Alternate Names:

Endodermal Sinus Tumor; Yolk Sac Germ Cell Tumor

## Yolk Sac Tumor

A cancer of germ cells that often occurs in the ovary or testicle in children or young adults and produces alpha-fetoprotein. It can also occur outside the reproductive organs.

## Yolk-Sac Tumor

Alternate name; see *Yolk Sac Tumor*.

## Yolk Sac Tumor Pathology

A malignant germ-cell tumor with variable reticular, microcystic, papillary, glandular, or solid patterns. Schiller-Duval bodies, hyaline globules, and alpha-fetoprotein expression may support the diagnosis.

## Yolk Sac Tumor Schiller-Duval Bodies

A characteristic microscope finding in yolk sac tumors. It looks somewhat like a tiny kidney filter, with tumor cells arranged around a small blood vessel. The finding helps confirm the diagnosis when considered with other test results.

### Alternate Names:

Schiller-Duval Bodies; Glomeruloid Endodermal Bodies

## Yolk Stalk

A temporary connection between the early yolk sac and the developing intestine. It normally disappears before birth; a remaining portion can form an abnormal intestinal pouch or other problem.

## Young-Burgess Classification

A system used to describe pelvic-ring fractures according to how the injury happened and how stable the pelvis is. Groups include side-to-side compression, front-to-back compression, vertical separation, and mixed injuries.



---

**Alternate Names:**

Young-Burgess Pelvic Fracture Classification;  
Burgess Pelvic Injury Scale

**Young Syndrome**

A rare condition, usually affecting males, that combines blockage of sperm transport with repeated sinus and lung infections. It may also cause bronchiectasis, a long-term widening and damage of the airways.

**Alternate Names:**

Sinopulmonary-Infertility Syndrome; Barry-Perkins-Young Syndrome

**Y-Plate Osteosynthesis**

Surgical repair of a complex broken bone using a metal plate shaped like the letter Y. The plate holds the bone pieces in place while they heal.

**Alternate Names:**

Y-Plate Fixation; Y-Shaped Contoured Bone Plate

**Yttrium-90**

A radioactive substance used in selected treatments to deliver radiation to abnormal tissue, such as some liver tumors or inflamed joints.

**Yttrium-90 Microspheres**

Tiny glass or resin beads containing radioactive yttrium-90. They are delivered through blood vessels to selected liver tumors, where they release radiation.

**Alternate Names:**

Y-90 Radioembolization Microspheres; TheraSphere; SIR-Spheres

**Yttrium-90 Radioembolization**

A treatment in which tiny beads containing radioactive yttrium-90 are delivered through liver blood vessels to release radiation inside liver tumors.

**Alternate Names:**

SIRT; Y-90 SIRT; Y-90 Transarterial Radioembolization

**Yttrium-Aluminum-Garnet Laser**

A laser that uses a crystal to produce a focused

beam of light. It is used in eye care for procedures such as opening a cloudy membrane after cataract surgery and making a small opening for some types of glaucoma.

**Alternate Names:**

Nd:YAG Laser; Neodymium:YAG Laser

**Yuzpe Regimen**

An older form of emergency contraception using two doses of combined birth-control hormones. It has largely been replaced by emergency contraception containing levonorgestrel or ulipristal.

**Y-V Plasty**

A surgical technique in which a Y-shaped cut is closed as a V. This lengthens or widens nearby tissue and may help release a tight scar or other narrowing.

**Alternate Names:**

Y-V Advancement Flap; Y-V Lengthening Procedure

**Yeast Infections**

Infections caused by an overgrowth of yeast, most often *Candida*. They commonly affect the vagina, mouth, skin, or bloodstream; symptoms and treatment depend on the site and the person's risk factors.



## Z

### Zafirlukast

A leukotriene-receptor antagonist used for long-term control and prevention of asthma symptoms. It is taken orally and is not a rescue treatment for an acute asthma attack; liver injury and drug interactions are important safety considerations.

#### Alternate Names:

Accolate

### Zahn Lines

Alternating light and dark layers seen in a blood clot that formed while blood was flowing. The light layers contain platelets and fibrin, while the dark layers contain more red blood cells.

#### Alternate Names:

Lines of Zahn; Striae of Zahn

### Zanamivir

An inhaled neuraminidase inhibitor used to treat or prevent influenza A and B when started appropriately. It may cause bronchospasm and is used cautiously in people with underlying airway disease.

#### Alternate Names:

Relenza

### ZAP-70 Deficiency

A rare inherited immune disorder in which important immune cells do not work properly. Children may have repeated, serious infections.

### Zavanelli Maneuver

An emergency procedure for severe shoulder dystocia that does not respond to other measures. The clinician gently returns the baby's head to the birth canal, followed by immediate cesarean delivery.

#### Alternate Names:

Cephalic Replacement Maneuver

### Z-Disc

A thin structure inside muscle fibers that anchors the protein strands involved in muscle contraction. It marks the boundary of each repeating unit of a muscle fiber.

#### Alternate Names:

Z-Line; Z-Band; Zwischenscheibe

### Zellweger Syndrome

A severe inherited disorder in which cells cannot make peroxisomes, structures that help process fats and other substances. It begins in infancy and may cause weak muscle tone, feeding problems, seizures, liver disease, and impaired brain development.

#### Alternate Names:

ZSD; Cerebrohepatorenal Syndrome

### Zenker Degeneration

Severe damage and death of skeletal muscle fibers, formerly described in serious infections such as typhoid fever or widespread blood infection.

#### Alternate Names:

Zenker's Hyaline Necrosis; Waxy Muscle Degeneration

### Zenker Diverticulum

A pouch that forms in the upper swallowing tube near the throat, usually because the nearby muscle does not relax normally. It may cause difficulty swallowing, return of undigested food, bad breath, a neck lump, cough, or repeated lung infections.

#### Alternate Names:

---

Zenker's Diverticulum; Pharyngoesophageal Diverticulum; Hypopharyngeal Diverticulum; Pulsion Diverticulum

### **Zenker Peroral Endoscopic Myotomy**

A procedure that treats a pouch in the upper swallowing tube called a Zenker diverticulum. A flexible camera is passed through the mouth and the tight muscle between the pouch and swallowing tube is divided so food can pass more easily. Symptoms can return and the procedure has risks such as bleeding or a tear.

#### **Alternate Names:**

Z-POEM; Endoscopic Zenker Myotomy

### **Ziconotide**

A strong pain medicine delivered continuously into the fluid around the spinal cord for selected severe chronic pain. It can cause dizziness, confusion, hallucinations, and other serious nervous-system or mental-health effects.

#### **Alternate Names:**

Prialt

### **Zidovudine**

An HIV medicine that interferes with the virus's ability to copy itself. It may be used in selected treatment plans and to reduce transmission around childbirth; important risks include anemia, low white-cell counts, muscle problems, and other toxicity.

#### **Alternate Names:**

Azidothymidine; AZT; Retrovir

### **Ziehl-Neelsen Stain**

A laboratory stain used to find bacteria with a waxy coating that resists removal by acid, especially the bacteria that cause tuberculosis. These bacteria appear red against a contrasting background under a microscope.

#### **Alternate Names:**

Acid-Fast Stain; AFB Stain

### **Zieve Syndrome**

A temporary complication of heavy alcohol use and sudden liver injury. It causes breakdown of red

blood cells, yellowing of the skin or eyes, and increased blood fats.

#### **Alternate Names:**

Hemolytic Anemia with Hyperlipidemia and Jaundice

### **Zika**

An infection caused by Zika virus. It often causes no symptoms or a mild fever, rash, red eyes, joint pain, and muscle pain. Infection during pregnancy can harm the developing baby, and it can rarely trigger Guillain-Barré syndrome.

#### **Alternate Names:**

Zika Virus Disease; ZVD; Congenital Zika Syndrome

### **Zika Virus**

A virus spread mainly by infected *Aedes* mosquitoes and also through sexual contact and, less commonly, blood or pregnancy-related transmission. Infection is often mild or symptom-free but can injure a developing fetus.

#### **Alternate Names:**

ZIKV

### **Zileuton**

A medicine used for long-term prevention and control of asthma. It does not relieve a sudden asthma attack and can cause liver problems or interact with other medicines.

#### **Alternate Names:**

Zyflo

### **Zinc**

An essential mineral needed for many body reactions, normal growth, immunity, reproduction, and wound healing. Too little can cause deficiency; excessive supplements can cause stomach symptoms and copper deficiency.

#### **Alternate Names:**

Zn

### **Zinc Acetate**

A zinc salt used to treat zinc deficiency and, in some forms, to reduce copper absorption in Wilson

---

disease. It may cause nausea or stomach irritation.

### **Zinc Deficiency**

A state of inadequate available zinc caused by insufficient intake, malabsorption, excessive losses, or increased requirements. Manifestations may include impaired growth, dermatitis, alopecia, diarrhea, altered taste, delayed wound healing, and increased infection susceptibility.

#### **Alternate Names:**

Hypozincemia; Zinc Depletion

### **Zinc Finger Motif**

A small part of a protein held in shape by a zinc atom. It often helps the protein attach to DNA and control gene activity.

### **Zinc Oxide**

An insoluble zinc compound used topically as a skin protectant and barrier, including in preparations for diaper dermatitis and minor irritation. It can reduce contact with moisture and irritants but does not treat every cause of a rash.

#### **Alternate Names:**

ZnO

### **Zinc Oxide Ointment**

A topical preparation containing zinc oxide that forms a protective barrier over irritated or moist skin. It is commonly used for diaper dermatitis and minor skin irritation.

#### **Alternate Names:**

Topical Zinc Barrier Cream; Zinc Oxide Paste

### **Zinc Protoporphyrin**

A substance made in developing red blood cells when zinc is used instead of iron during hemoglobin production. High levels can occur with iron deficiency or long-term lead exposure.

#### **Alternate Names:**

ZPP; Erythrocyte Zinc Protoporphyrin

### **Zinc Sulfate**

An oral or parenteral zinc salt used to prevent or treat zinc deficiency when clinically indicated. Gas-

trointestinal upset is common with oral therapy, and prolonged excessive use can cause copper deficiency.

### **Zinc Toxicity**

Adverse effects caused by excessive zinc ingestion or exposure. Acute toxicity may cause nausea, vomiting, abdominal pain, and diarrhea; chronic excess can impair copper absorption and lead to anemia or neurologic and immune abnormalities.

#### **Alternate Names:**

Hyperzincemia; Zinc Poisoning

### **Zinner Syndrome**

A rare condition present from birth in which one kidney is absent, a nearby seminal-vesicle cyst forms, and the sperm-carrying duct is blocked. It may cause pelvic pain, painful urination, or infertility after puberty.

#### **Alternate Names:**

Mesonephric Duct Malformation Syndrome; Seminal Vesicle Cyst with Renal Agenesis

### **Zirconia**

A zirconium-oxide ceramic used in dentistry for crowns, bridges, implants, and other restorations. Its strength and tissue compatibility make it useful, although restoration design, surface treatment, and clinical indication affect performance.

#### **Alternate Names:**

Zirconium Dioxide; Ceramic Biomaterial

### **Zoledronic Acid**

A potent intravenous bisphosphonate used for osteoporosis and certain cancer-related conditions, including hypercalcemia of malignancy and skeletal complications of bone metastases. Renal injury, hypocalcemia, acute-phase reactions, and rare osteonecrosis of the jaw are important risks.

#### **Alternate Names:**

Zoledronate; Zometa; Reclast

### **Zollinger-Ellison Secretin Stimulation Test**

A test used when a gastrin-producing tumor is suspected but a blood gastrin result is unclear. Secretin

---

is given through a vein and blood gastrin levels are measured more than once.

**Alternate Names:**

Secretin Provocation Test; Secretin Gastrinoma Test

**Zollinger-Ellison Syndrome**

A rare condition in which a hormone-producing tumor makes too much gastrin, causing the stomach to produce excess acid. It can lead to repeated ulcers, severe reflux, abdominal pain, and diarrhea.

**Zolpidem**

A nonbenzodiazepine sedative-hypnotic used for short-term treatment of insomnia. It can cause next-day impairment, complex sleep behaviors, dependence, and respiratory or central-nervous-system depression, especially with other sedatives.

**Alternate Names:**

Ambien

**Zona Fasciculata**

The middle and largest layer of the outer part of the adrenal gland. It makes glucocorticoid hormones, including cortisol.

**Alternate Names:**

Adrenal Zona Fasciculata; Fascicular Zone

**Zona Glomerulosa**

The outer layer of the adrenal gland. It makes mineralocorticoid hormones, mainly aldosterone, which helps control salt, water, and blood pressure.

**Alternate Names:**

Adrenal Zona Glomerulosa; Glomerular Zone

**Zona Pellucida**

A protective layer of proteins surrounding an egg cell. It helps sperm attach to the egg and changes after fertilization to prevent more than one sperm from entering.

**Alternate Names:**

Oocyte Glycoprotein Coat; Pellucid Zone

**Zona Reticularis**

The inner layer of the outer part of the adrenal gland. It makes some hormones related to sex hormones,

including adrenal androgens.

**Alternate Names:**

Adrenal Zona Reticularis; Reticular Zone

**Zonisamide**

An antiseizure medicine used mainly for focal seizures. It can cause sleepiness, thinking problems, kidney stones, reduced sweating with overheating, acid imbalance, severe rash, or other serious reactions.

**Alternate Names:**

Zonegran

**Zonula Adherens**

A belt-like connection between neighboring cells that helps hold them together. It is especially important in sheets of lining cells.

**Alternate Names:**

Belt Desmosome; Intermediate Junction; Adherens Junction

**Zonula Occludens**

A tight connection between neighboring lining cells that helps prevent substances from passing freely between them.

**Alternate Names:**

Tight Junction

**Zonule**

Fine fibers that hold the eye's lens in place. They transfer tension from the ciliary muscle so the lens changes shape and the eye can focus at different distances.

**Alternate Names:**

Zonule of Zinn; Ciliary Zonule; Suspensory Ligament of Lens

**Zoon Balanitis**

A chronic benign inflammation of the glans penis, usually seen in uncircumcised men. It commonly appears as a sharply demarcated shiny orange-red or rust-colored patch; diagnosis is clinical, with biopsy used when the diagnosis is uncertain.

**Alternate Names:**

Plasma Cell Balanitis; Balanitis Circumscripta Plas-



---

macellularis

### Zoonosis

An infection that naturally spreads between animals with backbones and people. Examples include rabies, brucellosis, leptospirosis, toxoplasmosis, and some influenza infections; spread may involve bites, food, water, droplets, or direct contact.

#### Alternate Names:

Zoonotic Infection; Zoonotic Disease

### Zoonotic Spillover

The spread of a germ from an animal or insect carrier to a person. If the germ then spreads easily between people, it can cause an outbreak or epidemic.

#### Alternate Names:

Zoonotic Transmission Event; Cross-Species Spillover

### Zoster

A painful, usually unilateral vesicular eruption caused by reactivation of latent varicella-zoster virus in a sensory nerve ganglion. It may be complicated by postherpetic neuralgia, ocular disease, or neurologic involvement.

#### Alternate Names:

Herpes Zoster; Shingles; Zona

### Zoster Ophthalmicus

Shingles affecting the nerve that supplies the forehead, upper eyelid, and eye. It causes a painful one-sided rash around the eye and can damage vision.

#### Alternate Names:

Herpes Zoster Ophthalmicus; HZO

### Z-Plasty

A reconstructive operation that rearranges nearby skin in a Z-shaped pattern. It can lengthen a tight scar, change the direction of a scar, and improve movement or appearance.

### Z-Score (DEXA)

A statistical bone mineral density (BMD) score obtained from dual-energy X-ray absorptiometry

that compares a patient's measured BMD to the mean BMD of an age-, sex-, and ethnicity-matched reference population. A Z-score of  $-2.0$  or lower is defined as "below the expected range for age" in children, premenopausal women, and men younger than 50 years.

#### Alternate Names:

Age-Matched Bone Density Z-Score; DEXA Z-Score

### Z-Score (Echocardiography)

A number that compares the size of a child's heart or blood-vessel structure with the expected size for children of a similar body size. It can help describe unusually small or large structures.

#### Alternate Names:

Pediatric Echo Z-Score; Cardiovascular Z-Score

### Z-Thumb Deformity

A characteristic deformity of the thumb in which fixed flexion occurs at the metacarpophalangeal (MCP) joint accompanied by hyperextension at the interphalangeal (IP) joint, resembling the letter Z. It is a hallmark joint manifestation of advanced rheumatoid arthritis.

#### Alternate Names:

Boutonniere-Type Thumb Deformity; Hitchhiker's Thumb Deformity

### Z-Tracking Technique

An injection method in which the skin is moved to one side before an injection and allowed to return afterward. This creates a zigzag path that helps keep the medicine in the muscle.

#### Alternate Names:

Z-Track Injection Method; Z-Track Technique

### Zygapophyseal Joint

One of the small joints at the back of the spine that connects neighboring vertebrae. These facet joints guide spinal movement and can become painful with arthritis.

#### Alternate Names:

Facet Joint; Z-Joint; Spinal Articular Facet

---

### Zygapophyseal Joint Arthropathy

Arthritis of the small facet joints at the back of the spine. It can cause local or nearby neck or back pain and stiffness.

#### Alternate Names:

Facet Joint Arthropathy; Spinal Facet Osteoarthritis

### Zygomatic Arch

A prominent horizontal bony bridge on the lateral aspect of the skull formed by the articulation of the temporal process of the zygomatic bone with the zygomatic process of the temporal bone, providing the origin for the masseter muscle and transmitting the temporalis muscle tendon beneath it.

#### Alternate Names:

Cheekbone Arch

### Zygomatic Bone

A paired facial bone that forms the prominence of the cheek and part of the lateral orbital wall and floor. It articulates with the frontal, sphenoid, temporal, and maxillary bones.

#### Alternate Names:

Cheekbone; Malar Bone

### Zygomatic Fracture

A break in the cheekbone or the group of bones joining it to the eye socket, usually caused by facial trauma. It may flatten the cheek, numb the skin below the eye, cause double vision, limit mouth opening, or injure the eye socket.

#### Alternate Names:

ZMC Fracture; Malar Fracture; Tripod Fracture

### Zygomaxillary Complex Fracture

A break involving the cheekbone and its connections to the eye socket, upper jaw, and skull. It may flatten the cheek and cause double vision, numbness below the eye, or difficulty opening the mouth.

#### Alternate Names:

ZMC Fracture; Tetrapod Fracture; Tripod Fracture of the Face

### Zygomycosis

A serious infection caused by molds called Mucorales. It mainly affects people with weakened immunity or poorly controlled diabetes and can rapidly damage the nose, brain, lungs, skin, or other tissues.

#### Alternate Names:

Mucormycosis; Invasive Mucorales Infection

### Zygote

The first cell formed when a sperm joins an egg. It contains genetic material from both parents, divides repeatedly, and develops into an embryo.

#### Alternate Names:

Fertilized Oocyte; Conceptus

### Zymogen

An inactive form of an enzyme that becomes active after a chemical change. Examples include pepsinogen, trypsinogen, and some blood-clotting proteins.

#### Alternate Names:

Proenzyme; Inactive Enzyme Precursor

### Zymogen Granule

A small storage sac inside certain gland cells, such as cells in the pancreas. It stores inactive digestive enzymes until the cell releases them.

#### Alternate Names:

Zymogenic Secretory Vesicle; Proenzyme Granule

### Zika Virus Infection

Infection caused by Zika virus, usually spread by infected *Aedes* mosquitoes and sometimes through sexual or pregnancy-related transmission. It is often mild or asymptomatic but can cause congenital infection and neurologic complications.

### Zika Fever

The usually mild febrile illness caused by Zika virus. It may include rash, red eyes, joint pain, muscle pain, headache, and fever; infection during pregnancy can harm the developing fetus.

### Zinc Poisoning

Illness caused by excessive zinc ingestion or exposure. It may cause nausea, vomiting, abdominal pain, and

---

diarrhea acutely; prolonged excess can interfere with copper absorption and cause anemia or neurologic problems.

### **Zoster (Shingles)**

A painful rash caused by reactivation of varicella-zoster virus in a sensory nerve. It typically affects one side of the body and may lead to postherpetic neuralgia or eye and neurologic complications.



# Drugs

## Generic Medicines and Medication Terms

### 5-Alpha Reductase Inhibitor

A medicine that lowers production of dihydrotestosterone, a hormone that affects the prostate and hair follicles. It is used mainly for an enlarged prostate and selected types of hair loss.

### Abortifacient

A medicine or substance used to end a pregnancy. Some block progesterone, others cause uterine contractions and cervical opening, and certain medicines stop rapidly dividing pregnancy tissue in selected ectopic pregnancies.

### Abortion - Medication

Medication abortion uses prescribed medicines, commonly mifepristone followed by misoprostol or misoprostol alone, to end an early intrauterine pregnancy; confirmation of completion and assessment for ectopic pregnancy are important.

### Abuse Steroid (Anabolic Steroid Abuse)

Alternate index label for Anabolic Steroid Abuse.

### A&E Medicine

The medical specialty concerned with the immediate assessment, resuscitation, diagnosis, and initial treatment of acute illness and injury.

### Alternate Names:

A&E Medicine

### ACE Inhibitor

A medicine that relaxes blood vessels and reduces strain on the heart by lowering production of angiotensin II. It treats high blood pressure, heart failure, and some kidney disease; cough, high potassium, kidney problems, or swelling can occur.

### Acitretin

An oral retinoid used mainly for severe psoriasis and selected keratinization disorders. It is highly teratogenic and requires strict pregnancy precautions and monitoring for adverse effects.

### Adenovirus Vaccine

A vaccine used primarily to prevent adenovirus types 4 and 7 among selected military personnel. It is an oral, live vaccine and is not part of routine vaccination for the general public. Vaccine use and eligibility depend on current public-health recommendations.

### Adverse Drug Event

Harm related to the use of a medication, including harm caused by an adverse drug reaction, medication error, overdose, or inappropriate use.

### Adverse Drug Reaction

An unwanted and harmful response to a medicine taken at a usual dose for prevention, testing, or treatment. It may result from the medicine's expected effects, an allergy, an interaction, or an unusual individual response.

### Agent, Tocolytic

A medicine that reduces uterine contractions, used

---

selectively to delay preterm birth long enough for corticosteroids, transfer, or other preparations to take effect.

### **Agonist-Antagonist**

A medicine that activates some receptors while blocking others, or activates a receptor only partly. Its mixed action can produce a desired effect while limiting another effect; examples include some opioid medicines.

### **AIDS Vaccine**

A vaccine designed to prevent HIV infection or reduce its effects; no generally effective licensed vaccine currently exists.

### **Alendronate**

A bisphosphonate that reduces bone resorption and is used to treat or prevent osteoporosis and selected other disorders of excessive bone turnover.

#### **Alternate Names:**

Alendronic acid; oral bisphosphonate therapy.

### **Anabolic Steroid Abuse**

Anabolic-androgenic steroid misuse can cause infertility, testicular shrinkage, acne, mood changes, high blood pressure, adverse cholesterol changes, liver injury, and cardiovascular events. Stopping may produce withdrawal symptoms and is safest with medical support.

#### **Alternate Names:**

Abuse Steroid (Anabolic Steroid Abuse)

### **Analgesic**

A medicine that relieves pain without necessarily causing loss of consciousness. Analgesics include nonopioid medicines, opioids, and other medicines chosen according to the type and severity of pain.

### **Analgesic Nephropathy**

Chronic tubulointerstitial renal injury associated with prolonged or excessive use of analgesic medicines. It may cause impaired urinary concentration, sterile pyuria, haematuria, papillary necrosis, hypertension, and progressive chronic kidney dis-

ease.

### **Anesthetic**

A medicine or technique that numbs part or all of the body. Depending on the method, a person may remain awake or become unconscious during a procedure; breathing, blood pressure, and other vital functions may need monitoring.

### **Angiotensin Converting-Enzyme Inhibitor**

A medicine that lowers the production of angiotensin II, a hormone that narrows blood vessels and promotes salt retention. It can treat high blood pressure, heart failure, and some kidney disease but may cause cough, high potassium, kidney problems, or swelling.

### **Angiotensin Receptor Blocker**

A medicine that blocks angiotensin II at the AT1 receptor, reducing vasoconstriction and aldosterone effects; it is used for hypertension, heart failure, and selected kidney disease.

### **Antibiotic**

A medicine that kills bacteria or stops them from multiplying. Antibiotics do not treat viral infections and should be selected for the suspected bacteria, infection site, allergies, and resistance risk.

### **Antibiotic-Associated Colitis**

Alternate Mayo Clinic label; see *Pseudomembranous colitis*.

### **Antibiotic-Associated Diarrhea**

Loose or frequent stools that occur during or soon after antibiotic treatment.

#### **Alternate Names:**

Diarrhea, Antibiotic-Associated

### **Antibiotic Cycling**

A strategy in which the preferred antibiotic for a particular clinical use is changed at planned intervals. It is intended to limit selection of resistant organisms, although its benefit depends on local



---

conditions and implementation.

### **Antibiotic De-Escalation**

The process of narrowing antimicrobial spectrum based on microbiological results and clinical response. De-escalation aims to reduce selective pressure for resistance while maintaining therapeutic efficacy.

### **Antibiotic Mixing**

A strategy in which different antibiotics are assigned concurrently to different patients with the same indication. It is intended to reduce uniform selection pressure for resistance in a patient population.

### **Antibiotic Resistance**

The ability of bacteria to survive or grow despite an antibiotic that would normally kill them or stop their growth. It can result from genetic changes and the spread of resistance genes; unnecessary or incorrect antibiotic use can accelerate it.

#### **Alternate Names:**

Resistance Antibiotic (Antibiotic Resistance)

### **Antibiotic Resistance Genes**

Genes that help bacteria survive antibiotics. They may change the antibiotic's target, destroy or modify the medicine, or remove it from the cell. Bacteria can inherit these genes or acquire them from other bacteria.

### **Antibiotic Stewardship**

Coordinated efforts to use antibiotics appropriately by selecting the right medicine, dose, route, and duration while reducing unnecessary treatment and resistance.

### **Anticholinergic Cognitive Burden Scale**

A medication-review tool that estimates the total anticholinergic load of a patient's medicines. It supports risk assessment and deprescribing but does not replace individualized clinical judgment.

### **Anticoagulant**

A medicine that slows the body's blood-clotting

process. It helps prevent or treat clots in veins, lungs, heart, or arteries but can cause bruising or serious bleeding.

#### **Alternate Names:**

Anticoagulants; Blood Thinners

### **Anticoagulant Agent**

See *Anticoagulant*.

### **Anticoagulant Monitoring**

Laboratory assessment used to measure the effect or safety of an anticoagulant. The appropriate test depends on the drug and may include aPTT, anti-Xa activity, INR, or measurements of kidney function and blood counts.

### **Anticoagulant Rodenticides Poisoning**

Poisoning from rat or mouse bait that interferes with blood clotting. Symptoms such as easy bruising, nosebleeds, blood in urine or stool, or weakness may appear later; suspected exposure requires immediate poison-control or emergency advice.

### **Anticonvulsant**

A medicine used to prevent or control seizures. Some are also used for nerve pain, migraine prevention, or mood disorders; choice depends on the condition, other medicines, pregnancy, and side effects.

#### **Alternate Names:**

Antiseizure medicine

### **Anticonvulsant Mood Stabilizers**

Antiepileptic drugs used as mood stabilizers in bipolar disorder. Valproate (divalproex) is effective for acute mania and rapid cycling. Carbamazepine is effective for acute mania but has numerous drug interactions via CYP3A4 induction. Lamotrigine is effective for bipolar depression prevention but not acute mania.

### **Antidepressant**

A medicine used to treat depressive disorders and some other conditions involving mood, anxiety, pain, or sleep.

---

## Antidiabetic Agent

See *Antidiabetic Drugs*.

## Antidiarrheal

A medicine that reduces the frequency or amount of diarrhea. It may be unsafe when stools contain blood, there is high fever or severe infection, or the bowel is blocked, so the cause should be considered first.

### Alternate Names:

Antidiarrheals; Antidiarrheal Medicines

## Antidiarrheal Medicine Overdose

Toxic effects caused by excessive exposure to Antidiarrheal medicine. Severity depends on the dose, route, timing, coexisting conditions, and clinical findings; suspected poisoning requires prompt assessment.

## Antiemetic

A medicine that prevents or reduces nausea and vomiting. The choice depends on the cause, such as infection, migraine, pregnancy, surgery, or cancer treatment, and some medicines can cause sleepiness or movement problems.

### Alternate Names:

Antiemetics; Antiemetic Medicines

## Anti estrogen

A drug that blocks estrogen effects or reduces estrogen production; it may be used in hormone-sensitive breast cancer or to alter reproductive hormone signaling.

## Antifungal

A medicine that kills fungi or stops them from growing. It may be applied to the skin or given by mouth or injection; the choice depends on the fungus, the body site, and the severity of infection.

## Antifungal Agent

A drug or chemical that kills fungi or inhibits their growth, used to treat fungal infections of the skin, mucosa, or internal organs.

## Antifungal Drug Classes

Major classes of antifungal agents include polyenes (amphotericin B), azoles (fluconazole, voriconazole), echinocandins (caspofungin, micafungin), and allylamines (terbinafine). Polyenes bind to ergosterol in fungal cell membranes. Azoles inhibit ergosterol synthesis. Echinocandins inhibit cell wall synthesis.

## Antifungal Medication

A medicine used to treat fungal infection; the choice and route depend on the organism and site, and examples include azoles, terbinafine, polyenes, and echinocandins.

### Alternate Names:

Antifungals; Antifungal Medicines; Antifungal Drugs

## Antihistamine

A medicine that blocks histamine, a chemical involved in allergies. It can relieve sneezing, itching, hives, or watery eyes; older types may cause sleepiness, dry mouth, blurred vision, urinary retention, or confusion.

## Antihypertensive

A medicine used to lower blood pressure. Classes include ACE inhibitors, angiotensin-receptor blockers, calcium-channel blockers, thiazide-type diuretics, beta-blockers, and other agents. Selection depends on blood pressure, kidney function, cardiovascular disease, pregnancy, and other clinical factors.

### Alternate Names:

Antihypertensives; Blood Pressure-Lowering Medicines

## Antihypertensive Agent

See *Antihypertensive*.

### Alternate Names:

Blood-pressure medicine

## Anti-Infective

A medicine or other agent used to prevent or treat an infection. The choice depends on whether the cause is a bacterium, virus, fungus, or parasite, as well as the infection site and the person's health.

---

### Antimalarial

A drug used to prevent or treat malaria caused by Plasmodium parasites; selection depends on species, disease severity, resistance, and geographic exposure.

### Antimicrobial Drug

A medicine used to treat infection by bacteria, viruses, fungi, or parasites; the drug is selected according to the organism, infection site, and patient factors.

### Antimicrobial Drug Interactions

Changes in the effect or level of an infection medicine caused by another medicine, food, or supplement. Some combinations reduce treatment effectiveness or increase toxicity, so the full medicine list should be checked.

### Antiparasitic

A medicine used to prevent or treat infection caused by parasites. The choice depends on the parasite, body site, disease severity, resistance patterns, pregnancy status, and other patient factors.

#### Alternate Names:

Antiparasitics; Antiparasitic Medicines; Antiparasitic Drugs

### Antiparkinsonian

A medicine used to improve movement or other symptoms of Parkinson disease. Medicines may replace dopamine, imitate its effects, or alter other brain signals; benefits and adverse effects vary with the drug and the person.

### Antiplatelet Agent

A medicine that makes blood platelets less likely to clump together. It lowers the risk of clots in arteries but can increase bruising and serious bleeding.

#### Alternate Names:

Antiplatelet medicine

### Antiprotozoal Drug

A medicine used to prevent or treat disease caused by protozoa, including drugs directed against

malaria, intestinal protozoa, or tissue parasites.

### Antipruritic

A medicine or skin treatment that relieves itching. It may work by reducing inflammation, blocking allergy signals, numbing the skin, or changing nerve signals; treatment should also address the cause of the itching.

### Antipsychotic

A medicine used to reduce hallucinations, delusions, severe agitation, or mania and to treat some mood disorders. Possible effects include sleepiness, movement problems, weight or blood-sugar changes, and rarely a dangerous fever-and-stiffness reaction.

#### Alternate Names:

Antipsychotics; Antipsychotic Medicines

### Antipsychotic Agent

See *Antipsychotic*.

### Antipsychotic-Induced Parkinsonism

Drug-induced bradykinesia, rigidity, tremor, or postural instability caused by dopamine-receptor blockade, usually from antipsychotic treatment. Management involves reviewing the medication, dose, alternatives, and symptomatic treatment when appropriate.

### Antispasmodic

A medicine that relaxes involuntary muscle and may reduce cramping in the intestine, bladder, or other organs. Some types can cause dry mouth, blurred vision, constipation, urinary retention, or confusion.

#### Alternate Names:

Antispasmodics

### Antithyroid

See *Antithyroid Drug*.

### Antithyroid Drug

A medicine that reduces thyroid hormone production, used to treat some forms of hyperthyroidism. Examples include methimazole and propylthiouracil. Radioactive iodine is a separate treatment, not an

---

antithyroid drug.

**Alternate Names:**

Antithyroid Medicines; Antithyroid Drugs

**Antiviral**

A medicine that slows or stops viruses from multiplying. It works only against certain viruses and is chosen according to the infection, timing, resistance, and the patient's health.

**Alternate Names:**

Antivirals; Antiviral Medicines; Antiviral Drugs

**Antiviral Drug Classes**

Groups of antiviral medicines classified by the step of viral growth they block. Some stop copying of viral genetic material, some block viral proteins or entry into cells, and others prevent newly formed viruses from being released.

**Anxiolytic**

A medicine used to reduce anxiety. Different medicines work in different ways and have different risks; choice depends on the condition, urgency, other health problems, and patient factors. Psychotherapy is a separate, non-medicine treatment for anxiety.

**Alternate Names:**

Anxiolytics; Anti-Anxiety Medicines

**ARB-Nepriylsin Inhibitor**

Sacubitril/valsartan combines an ARB with a neprilysin inhibitor, increasing natriuretic peptide levels while blocking the renin-angiotensin system. The PARADIGM-HF trial demonstrated superior reduction in cardiovascular death and heart failure hospitalization compared to enalapril.

**Aromatase Inhibitor**

A class of medications that inhibit aromatase, the enzyme that converts androgens (androstenedione, testosterone) to estrogens (estrone, estradiol) in peripheral tissues (adipose, breast, muscle, bone). Used primarily in hormone receptor-positive breast cancer in postmenopausal women.

**Attenuated Vaccine**

A vaccine made with a weakened microorganism that retains the ability to stimulate immunity.

**Alternate Names:**

Live attenuated vaccine

**Bacterial Vaccine**

A vaccine made from killed bacteria, weakened bacteria, purified bacterial parts, or bacterial toxins. It trains the immune system to prevent specific infections; protection and dosing depend on the vaccine and age group.

**BCG Vaccine**

A vaccine made from a weakened relative of the tuberculosis bacterium. It helps protect children from severe tuberculosis, especially meningitis, but protection against lung tuberculosis varies; use depends on local risk and policy.

**Behavioral Medicine**

An interdisciplinary field integrating behavioral science with medical practice to prevent, diagnose, and treat illness. It addresses psychological factors such as stress, coping, adherence, and health behaviors that influence disease outcomes.

**Benzodiazepines**

Medicines that strengthen the brain's main calming signal. They can reduce anxiety, stop some seizures, relax muscles, or cause sleepiness, but can also cause dependence, falls, confusion, and dangerously slow breathing, especially with opioids or alcohol.

**Beta-Adrenergic Blocker**

A drug that antagonizes beta-adrenergic receptors, reducing heart rate, myocardial contractility, and renin release. It is used in selected patients with hypertension, angina, arrhythmias, post-myocardial infarction disease, and heart failure; abrupt withdrawal may worsen ischemia.

**Alternate Names:**

Beta-Adrenergic Antagonist

---

## Beta Agonists

Drugs that activate beta-adrenergic receptors, producing effects such as relaxation of airway smooth muscle or increased cardiac stimulation depending on receptor selectivity.

### Alternate Names:

Beta-adrenergic agonists; beta-agonist drugs.

## Beta Blocker

See *Beta Blockers*.

## Beta-Blocker Overdose

Toxic exposure to a beta-adrenergic blocker that may cause slow heart rate, low blood pressure, low blood glucose, bronchospasm, seizures, or altered consciousness. It requires urgent poison-control and medical management.

## Beta Blockers

Medicines that block some effects of adrenaline at beta receptors. They commonly slow the heart rate and lower blood pressure and are used for several heart conditions and other problems. Some types can worsen asthma or mask warning signs of low blood sugar.

## Beta-Lactamase Inhibitor Combinations

Combinations of beta-lactam antibiotics with beta-lactamase inhibitors that extend activity against beta-lactamase producing organisms. The major combinations include amoxicillin-clavulanate, ampicillin-sulbactam, piperacillin-tazobactam, cefoperazone-sulbactam, and ceftolozane-tazobactam.

## C1 Esterase Inhibitor

A blood protein that helps control inflammation, swelling, and clotting reactions. Too little or abnormal activity can cause hereditary angioedema, with repeated attacks of swelling in the skin, bowel, or airway.

## Calcium Channel Blocker

A medicine that reduces calcium movement into heart and blood-vessel muscle cells. It may relax

arteries, lower blood pressure, reduce chest pain, or slow some abnormal heart rhythms; ankle swelling, constipation, dizziness, or a slow heartbeat may occur.

## Calcium Channel Blocker Overdose

Toxic exposure to a calcium-channel blocker that may cause low blood pressure, slow heart rate, conduction abnormalities, high blood glucose, or shock. It requires urgent medical and poison-control management.

## Calcium Channel Blockers

Medicines that reduce the movement of calcium into heart and blood-vessel muscle cells. Dihydropyridines, such as amlodipine and nifedipine, mainly relax arteries and are used for high blood pressure and angina. Other calcium channel blockers, such as diltiazem and verapamil, also slow the heart rate and electrical conduction and may be used for certain abnormal heart rhythms.

## Camptothecin Analogue

A cancer medicine related to camptothecin that blocks an enzyme cancer cells need to copy DNA. It can damage rapidly dividing cells and commonly causes low blood counts, diarrhea, nausea, or other gastrointestinal effects.

## Cancer Vaccine

A vaccine designed either to prevent infections that can cause cancer or to help the immune system attack an existing cancer. Its use depends on the specific vaccine and cancer; it does not replace other cancer treatment when needed.

## Cardiotonic Agent

A drug or substance that increases the force or efficiency of cardiac contraction, used selectively in heart failure or shock because benefits must be balanced against arrhythmia and other risks.

## Carvedilol Phosphate

The phosphate formulation of carvedilol, a medicine that blocks beta and alpha-1 receptors. It lowers heart rate and blood pressure and is used for heart

---

failure and hypertension; dizziness, slow heartbeat, and worsening fluid retention can occur.

### **Cebocephaly**

A rare birth defect in which the middle part of the face is severely underdeveloped, sometimes with a single nostril or a nose-like structure between closely spaced eyes. It usually occurs with major brain and facial-development abnormalities.

### **Celecoxib**

A selective COX-2 anti-inflammatory medicine used for pain and arthritis. It can cause stomach, kidney, blood-pressure, and heart problems and may increase cardiovascular risk, especially with long-term use or in people with existing risk factors.

### **Cholesterol - Drug Treatment**

Medicines used to lower harmful blood cholesterol or triglycerides and reduce cardiovascular risk. Statins are used most often; other options may be added when needed, according to cholesterol levels, overall risk, other illnesses, and treatment tolerance.

### **Cholinesterase Inhibitor**

A medicine that slows the breakdown of acetylcholine, a chemical used for nerve signalling. It can improve nerve communication in selected conditions but may cause nausea, diarrhoea, slow heartbeat, sweating, or muscle cramps.

### **Cold/Cough Medicines in Children**

Many nonprescription cold and cough medicines are unsafe or ineffective for young children and can cause sedation, agitation, abnormal heart rate, overdose, or breathing problems. Age-appropriate fluids, saline, humidification, and clinician-guided treatment are safer approaches.

### **Conjugate Vaccine**

A vaccine that joins a bacterial sugar to a carrier protein so the immune system makes a stronger, longer-lasting response. This design is especially useful in infants and young children.

#### **Alternate Names:**

Protein-conjugate vaccine; conjugated polysaccha-

ride vaccine.

### **Conscious Sedation**

Medicine given to make a person relaxed and drowsy during a procedure while allowing them to respond to voice or touch and usually breathe independently. Breathing, blood pressure, and alertness must be monitored because deeper sedation can occur.

### **Controlled Substance**

A medicine or chemical whose production, supply, possession, or use is restricted by law because it may cause dependence, misuse, overdose, or other serious harm. Rules differ between countries and substances.

### **Cox-1 Inhibitor**

A medicine that blocks cyclooxygenase-1, an enzyme involved in pain, stomach protection, kidney function, and platelet clotting. Blocking it may relieve pain but can increase stomach irritation, bleeding, or kidney problems.

### **COX Inhibitor**

A class of medicines that reduce production of chemicals involved in pain, fever, and inflammation. They can irritate the stomach, affect kidney function, raise blood pressure, or increase bleeding and cardiovascular risk.

### **Crack (Drug)**

A smokable, rapidly acting form of cocaine; it can cause dependence, severe cardiovascular effects, seizures, stroke, psychiatric symptoms, and overdose.

### **Cutaneous Drug Eruptions**

Skin rashes or other skin reactions caused by medicines. They may appear as a widespread red rash, hives, sun-sensitive areas, or a lichen-planus-like eruption. Blisters, skin peeling, mouth sores, fever, or facial swelling require urgent assessment.

### **Cyclooxygenase Inhibitor**

A medicine that reduces production of



---

prostaglandins and may relieve pain and inflammation.

### Designer Steroid

A chemically modified anabolic-androgenic steroid created to produce performance-enhancing effects or evade detection; products may be unapproved, contaminated, and medically dangerous.

### Diarrhea, Antibiotic-Associated

Alternate index label; see *Antibiotic-associated diarrhea*.

### Digitalis

A group of heart medicines from foxglove plants, including digoxin. They can strengthen heart contraction and slow some heart rhythms but have a narrow safety range and may cause dangerous toxicity.

### Diphtheria Vaccine

A vaccine containing an inactivated form of diphtheria toxin. It trains the immune system to neutralize the toxin and is usually given in combination vaccines with protection against tetanus and pertussis.

### Disease-Modifying Antirheumatic Drug

A medicine used for inflammatory rheumatic diseases to reduce disease activity and help prevent lasting joint or organ damage, rather than only relieving symptoms. Different types are chosen according to the disease, severity, other conditions, and monitoring needs.

#### Alternate Names:

DMARD; DMARDs; Disease-Modifying Antirheumatic Drugs

### Disulfiram

A medicine used as an adjunct for alcohol-use disorder that inhibits aldehyde dehydrogenase and can cause an unpleasant reaction if alcohol is consumed.

### Diuretic

A medicine that helps the kidneys remove extra salt and water in urine. It can treat swelling, high

blood pressure, heart failure, or selected kidney and liver conditions but may disturb salts or cause dehydration.

#### Alternate Names:

Diuretics; Water Pills

### Doctor Of Osteopathic Medicine

A physician who has completed an accredited osteopathic medical degree and residency training and is licensed to diagnose, prevent, and treat disease.

### Drug Administration Route

The pathway by which a medicine enters the body, such as oral, intravenous, intramuscular, inhaled, topical, or subcutaneous administration.

### Drug Allergies

Drug allergies are immune-mediated reactions to a medicine, ranging from rash to anaphylaxis; they must be distinguished from predictable side effects and nonimmune intolerance.

### Drug Allergy

An adverse immune-mediated reaction to a medication, representing a hypersensitivity response that is distinct from side effects, toxicities, and idiosyncratic reactions.

### Drug Antagonism

An interaction where one drug blocks or decreases the effectiveness of another, such as when an antidote blocks a poison.

### Drug Bioavailability

The proportion of a dose that reaches the bloodstream in an active form. It depends on how the medicine is given, how well it is absorbed, and how much is broken down before reaching the blood; intravenous doses enter the bloodstream completely.

### Drug Classification Systems

Systems that group medicines according to properties such as their chemical structure, mechanism of action, therapeutic use, or effects on the body.

---

## Drug Clearance

The body's ability to remove a medicine from the bloodstream, mainly through the liver and kidneys. Clearance affects how long a medicine works and how often it should be taken; liver or kidney disease may require dose adjustment.

## Drug Dependence

A state of physiological or psychological adaptation to a drug, characterized by a withdrawal syndrome upon discontinuation or dose reduction. Physical dependence: the body has adapted to the presence of the drug, and abrupt cessation produces a predictable withdrawal syndrome specific to the drug class.

## Drug-Drug Interactions

Changes in the effect or amount of one medicine caused by another medicine. The combination may make treatment less effective or increase side effects, so all prescription, nonprescription, and supplement use should be reviewed.

## Drug-Eluting Balloon (Vascular)

A vascular balloon catheter that delivers a medicine to the vessel wall during dilation, without leaving a permanent implant.

### Alternate Names:

Vascular

## Drug Emulsion

A medicine made by dispersing tiny droplets of one liquid throughout another liquid that normally would not mix with it. An added stabilizer keeps the mixture uniform so the medicine can be given safely.

## Drug Eruption

A skin eruption caused by a medication, ranging from a mild morbilliform rash to severe reactions such as Stevens–Johnson syndrome or toxic epidermal necrolysis.

## Drug-Facilitated Sexual Assault

Sexual contact occurring when a person's capacity to

consent or resist is impaired by alcohol, medication, or another drug, whether administered covertly or knowingly consumed. Medical care includes urgent safety assessment, treatment of injuries, toxicology sampling when indicated, prophylaxis, and trauma-informed support.

## Drug Hypersensitivity Reactions

Immune-mediated adverse reactions to medicines. They may be immediate or delayed and can range from a mild rash to severe systemic illness.

## Drug Hypersensitivity Syndrome

A serious delayed reaction to a medicine, usually beginning two to eight weeks after it is started. It may cause fever, widespread rash, facial swelling, enlarged lymph nodes, abnormal blood cells, and injury to the liver, kidneys, lungs, or heart.

## Drug-Induced Autoimmunity

An immune reaction caused by a medicine in which the body attacks its own tissues. It may resemble lupus, cause joint pain, fever, rash, or organ inflammation, and often improves after the responsible medicine is stopped under medical guidance.

## Drug-Induced Diarrhea

Diarrhea caused by a medicine or drug interaction. Common causes include antibiotics, laxatives, metformin, and medicines that alter intestinal motility.

## Drug-Induced Immune Hemolytic Anemia

An uncommon immune-mediated destruction of red blood cells triggered by a medication; evaluation includes blood tests and review of exposures, with specialist management.

## Drug Induced Liver Disease

Drug-induced liver injury occurs when a medicine, supplement, or toxin damages liver cells or bile flow. It may cause fatigue, nausea, itching, dark urine, jaundice, or no symptoms; identifying and stopping the responsible exposure under medical guidance is central to care.

---

### **Drug-Induced Low Blood Sugar**

Low blood glucose caused by a medicine, most often insulin or an insulin secretagogue. Symptoms may include sweating, tremor, confusion, seizure, or loss of consciousness.

### **Drug-Induced Pulmonary Disease**

Lung injury caused by a medicine or treatment. It may produce cough, breathlessness, fever, or abnormal lung imaging; the medicine, timing, and other possible causes must be reviewed promptly because some cases can progress to respiratory failure.

### **Drug-Induced Thrombocytopenia**

A low platelet count caused by a medicine through immune destruction, reduced production, or other mechanisms, increasing bleeding risk.

### **Drug-Induced Tremor**

Tremor caused or worsened by a medicine, stimulant, or toxin. Assessment considers timing, dose, other causes, and whether the medicine can be changed.

### **Drug Interaction**

A change in the effect or concentration of one medicine caused by another medicine, food, substance, or disease.

### **Drug Metabolism**

The enzymatic conversion of medicines into metabolites, mainly in the liver but also in the intestine, kidneys, lungs, and other tissues.

### **Drug Metabolism Pharmacogenomics**

The study of how inherited genetic differences change the way a person breaks down medicines. This information can sometimes help clinicians choose a safer or more effective medicine or dose.

### **Drug Overdose**

Harm caused by taking too much medicine or taking it by the wrong route. Effects range from sleepiness, vomiting, and confusion to seizures, organ damage, or stopped breathing. Suspected overdose requires urgent poison-control or emergency advice.

### **Drug Packaging**

The container, labeling, and protective materials used to preserve a medicine and support correct identification, storage, and use.

### **Drug, Prescription**

A medicine legally supplied under the order of an authorized prescriber, with its indication, dose, duration, interactions, and risks specified for the patient.

### **Drug Resistance**

The ability of microorganisms, cancer cells, or parasites to survive and proliferate despite exposure to drugs that were previously effective.

### **Drug, Sulfa**

A sulfonamide drug; some are antibacterial medicines, while others are used as diuretics or for other indications, and sulfonamide reactions vary by drug and patient.

### **Drug Therapeutics**

The use of medicines to prevent, diagnose, or treat disease while balancing expected benefit and potential harm.

### **Drug Tolerance**

A decreased pharmacological response to a drug after repeated administration, requiring dose escalation to achieve the same effect.

### **Drug Toxicity**

Harmful effects caused by a medicine or its metabolites. Toxicity may result from excessive exposure, accumulation, interactions, or an unpredictable individual reaction.

### **Drug Use First Aid**

Immediate care for suspected drug intoxication, including assessment of breathing and responsiveness, emergency activation, overdose-specific support, and naloxone when opioid toxicity is suspected.

---

## Drug Vehicle

An inactive substance or carrier used to deliver an active drug ingredient in a formulation.

## Drug Withdrawal

A clinically significant syndrome or return of symptoms caused by abrupt reduction or cessation of a drug after physiologic adaptation or dependence has developed. Tapering is required for some medications to reduce withdrawal and rebound effects.

## DTaP Vaccine

A vaccine against diphtheria, tetanus, and pertussis (whooping cough), given mainly to infants and young children in a multi-dose series. It reduces severe illness, complications, and spread; schedules vary by country.

## Eflornithine

A medicine that slows cell growth by blocking an enzyme needed to make certain cell components. A skin cream can reduce unwanted facial hair; an intravenous form has been used for selected African trypanosome infections.

## Enzyme Inhibitor

A substance that slows or blocks an enzyme, a protein that helps chemical reactions occur in the body. Medicines use this action to change processes such as blood pressure, inflammation, infection, hormone production, or cancer-cell growth.

## Estrogen Inhibitor

A medicine that reduces estrogen production or blocks estrogen's effects. It may treat selected hormone-sensitive cancers or other conditions; side effects can include hot flashes, bone loss, joint symptoms, or blood clots.

## Etidronate

A bisphosphonate medicine that slows the breakdown and rebuilding of bone. It is used selectively for some disorders of abnormal bone remodeling; treatment depends on the condition and can affect the stomach, kidneys, or minerals.

## Evidence-Based Medicine

The use of reliable research together with clinical expertise and the patient's values and circumstances to make healthcare decisions. It supports, but does not replace, discussion between patient and clinician.

## Expectorant

A medicine intended to loosen or thin mucus so it is easier to cough up. Benefit varies with the cause of cough and the product; persistent breathlessness, fever, chest pain, or blood needs medical assessment.

## Extensively Drug-Resistant TB

Tuberculosis caused by bacteria resistant to rifampicin and usually isoniazid, at least one fluoroquinolone, and at least one of the key drugs bedaquiline or linezolid. Diagnosis requires drug-susceptibility testing and specialist treatment.

## Family Medicine

A medical specialty providing comprehensive continuing care for people of all ages, including prevention, acute illness, chronic disease, and coordination of referrals.

## Fentanyl

A potent synthetic opioid analgesic used for severe pain and anesthesia. It can cause respiratory depression, sedation, constipation, nausea, tolerance, dependence, and fatal overdose, especially when combined with alcohol or other sedatives.

## Fertility Agent

A medicine or biologic intervention intended to improve the chance of conception by inducing ovulation, supporting sperm production, or treating a reproductive mechanism.

## Fidaxomicin

A narrow-spectrum oral macrocyclic antibiotic used to treat *Clostridioides difficile* infection. It has minimal systemic absorption and reduces disruption of normal intestinal flora compared with many alternatives, but recurrence can still occur.

---

## Filgrastim

A recombinant granulocyte colony-stimulating factor that increases production and release of neutrophils from bone marrow. It is used to reduce or treat selected forms of chemotherapy-related neutropenia; bone pain and rare splenic complications can occur.

## Finasteride

A medicine that lowers the conversion of testosterone to dihydrotestosterone. It can shrink an enlarged prostate and help some types of hair loss but may reduce libido, affect sexual function, or alter prostate-test results.

## Fixed Drug Eruption

A medicine-related skin reaction that returns in the same place whenever the person takes the responsible medicine. It may appear as a sharply outlined red or dark patch, sometimes with a blister or sore, and can leave lasting discoloration.

## Fluticasone

A corticosteroid available in inhaled, nasal, topical, and other formulations to reduce inflammation. Adverse effects depend on route and may include local irritation, oral candidiasis, skin thinning, adrenal suppression, or growth effects with prolonged exposure.

## Food-Drug Interaction

A change in how a medicine works or is absorbed because of food, drink, nutrients, or supplements. Examples include grapefruit affecting some medicines, vitamin K changing warfarin's effect, and calcium reducing absorption of some antibiotics.

## Forensic Medicine

Application of medical knowledge to legal questions, including injury interpretation, cause of death, toxicology, identification, sexual assault evaluation, and documentation.

## Forensic Medicine (Legal Medicine)

Branch of medicine that applies medical, biological, and scientific knowledge to legal questions and the

administration of justice.

## Alternate Names:

Legal Medicine

## Foscarnet

An intravenous antiviral agent used for selected cytomegalovirus or herpesvirus infections, especially when resistance or intolerance limits other treatment. Important toxicities include kidney injury and disturbances of calcium, magnesium, potassium, or phosphate.

## Furosemide

A diuretic that makes the kidneys remove more salt and water in urine. It treats swelling and some blood-pressure or heart conditions but may cause dehydration, abnormal body salts, kidney problems, or hearing damage.

## Ganglionic Blocker

A medicine that interrupts nerve signals at switching points controlling automatic body functions. It can affect blood pressure, heart rate, digestion, sweating, vision, and bladder function, so these medicines are rarely used clinically.

## Generic Drug

A medicine equivalent to a brand-name medicine in active ingredient, strength, dosage form, and intended use, usually sold under a nonproprietary name.

## Alternate Names:

Generic medicine

## Generic Medicine

A medicine with the same active ingredient, strength, form, route, and intended effect as an approved reference medicine. It may look different or have different inactive ingredients but must meet regulatory quality and performance requirements.

## Generic Name, Drug

The official nonproprietary name of a medicine's active ingredient, used consistently across manufacturers and countries. It identifies the medicine itself

---

rather than a company's brand name and helps prevent confusion when products have different trade names.

### **Genomic Medicine**

Use of a person's genetic or genomic information to guide disease diagnosis, risk assessment, prevention, or treatment.

### **Geriatric Medicine**

The medical specialty focused on the health, function, diseases, medications, and goals of older adults, often using comprehensive and coordinated care.

### **Growth Inhibitor**

A medicine or biological substance that slows or stops cell growth. In cancer treatment, it may block cell division or a growth signal, but it can also affect healthy cells and cause side effects.

### **Haemophilus Vaccine**

A vaccine that protects against serious disease caused by *Haemophilus influenzae* type b, including meningitis, pneumonia, and bloodstream infection. It is given mainly in infancy according to the recommended immunization schedule.

### **Hepatic Drug Metabolism**

The liver changes medicines into forms that the body can use or remove. This may stop a medicine's effect, activate a medicine that works after conversion, or produce a harmful product; age, illness, genetics, and other medicines can alter the process.

### **Hepatitis B Vaccine**

A recombinant vaccine that prevents hepatitis B infection, which can cause chronic liver disease, cirrhosis, and liver cancer. It is given in a series beginning at birth in many schedules; infants exposed at birth may also need hepatitis B immune globulin.

### **Hepatitis B Vaccine (Birth Dose)**

The first hepatitis B vaccine dose, given soon after birth, helps prevent infection passed from a parent to the newborn. If the birth parent has hepatitis B,

the infant also needs hepatitis B immune globulin promptly; local schedules determine later doses.

### **Alternate Names:**

Birth Dose

### **HER2 Inhibitor**

A targeted cancer treatment that blocks signaling through human epidermal growth factor receptor 2 in cancers with HER2 overexpression or amplification.

### **High Blood Pressure - Medicine-Related**

Hypertension caused or worsened by a medicine or substance, such as NSAIDs, steroids, decongestants, stimulants, oral contraceptives, or immunosuppressants; review and substitution may improve control.

### **Histamine-2 Receptor Antagonist**

A medicine that reduces stomach-acid production by blocking histamine signals in the stomach. It may treat reflux, indigestion, or ulcers; headache, diarrhea, confusion, and medicine interactions can occur.

### **Hormone Inhibitor**

A medicine or biologic that blocks hormone synthesis, release, receptor binding, or downstream signaling, used in selected endocrine disorders and hormone-sensitive cancers.

### **HPV Vaccine**

A vaccine that prevents infection with human papillomavirus types that cause several cancers and genital warts. It is routinely offered in early adolescence, can start from age 9, and may require two or three doses depending on age and immune status; it does not treat existing HPV infection.

### **Hypoglycemic Agent**

A medicine that lowers blood glucose, including insulin, insulin secretagogues, insulin sensitizers, incretin-based drugs, and sodium-glucose cotransporter inhibitors; choice depends on diabetes type and patient risks.



---

### **Hypolipidemic Agent**

A medicine that lowers fats in the blood, such as LDL cholesterol or triglycerides. It helps reduce cardiovascular risk when combined with appropriate diet, activity, and treatment of other risk factors.

### **Immunosuppressant**

A medicine that reduces immune-system activity. It may be used to prevent organ transplant rejection or treat autoimmune and inflammatory diseases; effects and infection risks vary by medicine and dose.

### **Immunosuppressant Drug**

A medicine that reduces immune-system activity, often to prevent transplant rejection or treat autoimmune disease.

### **Immunosuppressive Drug**

A drug that suppresses immune responses, used for organ transplantation, graft-versus-host disease, and autoimmune/inflammatory diseases.

### **Inactivated Polio Vaccine**

An injected vaccine made from killed polioviruses. It trains the immune system to prevent poliomyelitis and is used in routine childhood immunization programs; the schedule and use of other polio vaccines vary by country.

### **Influenza Vaccine**

A seasonal vaccine that reduces the risk of influenza and its complications. It is updated as needed and is offered each year to most people from early infancy; some young children need two doses in their first vaccinated season, according to local recommendations.

### **Inotrope**

A drug that changes myocardial contractility; positive inotropes increase the force of contraction. Dobutamine and milrinone may be used short term under specialist monitoring for cardiogenic shock or selected cases of acute decompensated heart failure with low output. Dopamine, epinephrine, and norepinephrine are primarily vasopressors or emer-

gency agents with additional inotropic effects, while digoxin is a separate cardiac glycoside used in selected chronic heart-failure or rate-control settings. These medicines can cause arrhythmias, ischemia, or other serious effects and are not routine long-term treatment for chronic heart failure.

### **Intensive Care Medicine**

Specialty managing critically ill patients. Components: monitoring, ventilatory support, hemodynamic support, infection control, nutrition, sedation, pain management. Common conditions: sepsis, ARDS, traumatic brain injury, post-cardiac arrest, multiorgan failure.

### **Internal Medicine**

The medical specialty focused on prevention, diagnosis, and nonsurgical treatment of diseases in adults, including complex multisystem illness.

### **Japanese Encephalitis Vaccine**

A vaccine that protects against Japanese encephalitis, a mosquito-borne viral infection of the central nervous system. Vaccine type, schedule, and eligibility depend on age, destination or exposure risk, and local recommendations.

### **Laxative**

See *Laxatives*.

### **Legal Medicine**

The application of medical knowledge to legal questions, including forensic examination, injury assessment, death investigation, and issues of capacity or consent.

### **Liquid Medication Administration**

Giving a medicine in liquid form by mouth, feeding tube, injection, or another prescribed route. The dose must be measured accurately with the supplied device because liquid strengths differ; household spoons can cause dosing errors.

### **Local Anesthetic**

A medicine that temporarily numbs a specific area

---

without usually causing loss of consciousness. It blocks pain signals in nearby nerves and is used for procedures, dental work, wound repair, and some forms of regional pain relief.

**Alternate Names:**

Local Anesthetics

**Luteinizing Hormone-Releasing Hormone Agonist**

A medicine that first stimulates and then, with continued use, suppresses signals from the pituitary gland that control the ovaries or testes. It is used for selected cancers, endometriosis, fibroids, and puberty disorders.

**Alternate Names:**

GnRH agonist; LHRH agonist

**Macrolide Antibiotic**

An antibiotic of the macrolide class that binds the bacterial 50S ribosomal subunit and usually inhibits protein synthesis; activity and resistance vary by organism and drug.

**Malaria Vaccine**

A vaccine intended to reduce malaria caused by *Plasmodium* parasites, especially *P. falciparum*; the RTS,S/AS01 and R21/Matrix-M vaccines are used in recommended childhood immunization programs in some endemic settings, alongside other preventive measures.

**MAO Inhibitor**

A medicine that slows breakdown of certain brain chemicals and is used for selected depression or Parkinson disease. It can interact dangerously with other medicines and some foods, so it requires careful prescribing and instructions.

**Measles-Rubella Vaccine**

A combined live attenuated vaccine containing measles and rubella antigens without the mumps component, widely used in countries where mumps is not a significant public health burden, including India's Universal Immunization Schedule.

**Measles Vaccine**

Measles vaccine is a live attenuated vaccine that induces active immunity against measles virus. It is usually given as part of the measles–mumps–rubella (MMR) or measles–mumps–rubella–varicella (MMRV) vaccine.

**Medication**

A substance or biological product used to prevent, diagnose, treat, or control a disease or symptom. Medicines may be prescribed, bought without a prescription, applied to the body, inhaled, swallowed, or given by injection.

**Medication Adherence**

The extent to which a patient's medication-taking behavior corresponds with the agreed treatment plan. Adherence is influenced by understanding, adverse effects, cost, access, regimen complexity, beliefs, and cognitive or social factors.

**Medication Adverse-Effect Profile**

The clinically relevant pattern of predictable pharmacologic effects and harmful reactions associated with a medication. It includes common dose-related effects, serious rare events, organ-specific toxicity, interactions, and monitoring requirements.

**Medication Cascade**

A prescribing sequence in which an adverse drug effect is misinterpreted as a new medical condition and another drug is prescribed to treat it. Recognizing cascades is an important goal of medication reconciliation and deprescribing.

**Medication Deprescribing**

A planned, supervised process of reducing the dose or stopping a medicine when its current or future harms outweigh its benefits or when treatment goals change. Tapering may be required to prevent withdrawal or rebound disease.

**Medication Error**

A preventable event that may cause or lead to inappropriate medication use or patient harm while the medication is in the control of a healthcare pro-

---

fessional, patient, or consumer. Errors can occur during prescribing, transcribing, dispensing, administration, or monitoring.

### **Medication Management**

The systematic process of optimizing pharmacotherapy to achieve therapeutic goals while minimizing adverse effects, encompassing prescribing, dispensing, administration, monitoring, and deprescribing.

### **Medication Overuse Headaches**

Frequent headaches caused or worsened by regular overuse of headache medicines.

**Alternate Names:**

Rebound Headache

### **Medication Reconciliation**

A structured process of obtaining and comparing the most accurate medication list with current prescriptions at transitions of care. It identifies omissions, duplications, incorrect doses, interactions, and unintended changes.

### **Medication Review**

Systematic examination of all medications for appropriateness. Components: indication, efficacy, safety, adherence, interactions. Tools: Beers Criteria, STOPP/START. Goals: deprescribing, optimizing therapy, reducing adverse effects.

### **Medication Safety**

Practices that reduce medication errors and prevent avoidable harm during prescribing, dispensing, administration, and use.

### **Medication Side Effects**

Unintended harmful responses to medications occurring at standard doses, ranging from mild (nausea, rash, dry mouth) to severe (anaphylaxis, Stevens-Johnson syndrome, QT prolongation, hepatotoxicity).

### **Meningococcal Vaccine**

Conjugate vaccine (MenACWY) and serogroup B vaccine (MenB) given at 11-12 years with booster

at 16 years. Protects against *Neisseria meningitidis*, causing meningitis and septicemia. MenACWY booster at 16 years covers the high-risk period for serogroup C and Y disease.

### **Mineralocorticoid Receptor Antagonist**

A medicine that blocks aldosterone, a hormone that makes the body retain salt and water. It is used for selected heart failure, high blood pressure, and hormone disorders but can raise blood potassium and worsen kidney problems.

### **MMR Vaccine**

A live attenuated vaccine against measles, mumps, and rubella. It is given according to the immunization schedule and is generally avoided during pregnancy and in people with certain severe immune deficiencies.

### **Molecular Medicine**

Molecular medicine applies molecular and cellular biology to understand, diagnose, prevent, and treat disease. It includes genomics, precision oncology, molecular biomarkers, targeted therapy, and other mechanism-based approaches.

### **Monoamine Oxidase Inhibitor**

A medicine that blocks monoamine oxidase, an enzyme that breaks down certain brain chemicals. It can treat selected depression or neurologic disorders but has important interactions with other medicines and some foods.

### **MRNA Vaccine**

A vaccine that gives cells temporary instructions to make a harmless piece of a germ. The immune system learns to recognize that piece and respond later; the messenger RNA does not enter the cell nucleus or change DNA.

### **Multi-Drug Resistance**

Multidrug resistance is resistance of a microorganism, cancer, or other disease process to multiple drugs that would otherwise be active. Mechanisms include target changes, drug inactivation, reduced uptake, efflux, and altered repair or survival path-

---

ways.

### **Multidrug-Resistant Extensively Drug-Resistant TB**

Multidrug-resistant tuberculosis is resistant to key first-line drugs, while extensively drug-resistant TB has additional resistance that greatly limits options. Diagnosis requires molecular or culture testing, and treatment uses prolonged individualized combinations with public-health oversight.

### **Neuropathy Secondary To Drugs**

Drug-induced neuropathy is peripheral nerve damage caused by a medicine or toxin, often producing numbness, burning pain, weakness, or altered reflexes; management may require dose reduction, withdrawal, and treatment of symptoms.

### **Nonsteroidal Anti-Inflammatory Drug**

A medicine that inhibits cyclooxygenase enzymes and reduces prostaglandin production, producing analgesic, anti-inflammatory, and fever-reducing effects. NSAIDs can cause gastrointestinal, kidney, cardiovascular, or other adverse effects.

### **NSAIDs**

See *Nonsteroidal Anti-Inflammatory Drug*.

### **Occupational Medicine**

The medical specialty concerned with the effects of work on health, prevention of work-related illness and injury, and safe return to work.

### **Opioid Analgesics**

Strong pain medicines that act on opioid receptors in the brain, spinal cord, and other tissues. They can relieve severe pain but may cause sleepiness, constipation, dependence, and dangerously slow breathing or overdose.

### **Opioid Antagonist**

A medicine that blocks opioid receptors and can reverse opioid effects such as respiratory depression. Naloxone is a commonly used example.

### **Oral Hypoglycemic**

A medicine taken by mouth to lower blood glucose, used mainly in type 2 diabetes. Different medicines improve insulin sensitivity, increase insulin release, slow glucose absorption, or increase glucose loss in urine; choice depends on the patient's risks and kidney function.

### **Oral Medicine**

Specialty diagnosing oral mucosal diseases, dental causes of facial pain, oral manifestations of systemic disease.

### **Oral Polio Vaccine**

A live attenuated poliovirus vaccine administered by mouth. Its formulation and use depend on the public-health program and circulating poliovirus strains; it has played an important role in polio control.

### **Orphan Drug**

A pharmaceutical agent developed specifically to treat a rare medical condition (affecting fewer than 200,000 people in the United States, as defined by the Orphan Drug Act of 1983).

### **Over-The-Counter Drug**

A medication that can be purchased without a prescription, deemed safe and effective for self-medication when used according to label instructions. OTC drugs have established safety profiles, low toxicity, and clear dosing guidelines suitable for consumer use.

### **Over-The-Counter (OTC) Drug**

A medicine sold without a prescription for uses and doses described on its label. "Over the counter" does not mean risk-free; incorrect use, interactions, or excessive doses can still cause serious harm.

### **P2Y<sub>12</sub> Inhibitor**

An antiplatelet drug that blocks the platelet P2Y<sub>12</sub> ADP receptor, reducing activation and aggregation. Clopidogrel, prasugrel, and ticagrelor are used in selected acute coronary syndrome and percutaneous coronary intervention settings, with bleeding and

---

drug-specific contraindications considered.

### **Palifermin**

A laboratory-made form of a natural growth factor that helps mouth and throat lining cells repair themselves. It is used in selected people receiving intensive treatment before blood-forming stem-cell transplantation to reduce severe mouth sores.

### **Palonosetron**

A long-acting anti-nausea medicine that blocks serotonin signals involved in vomiting. It is used to prevent nausea and vomiting from chemotherapy or surgery; headache, constipation, and rare heart-rhythm changes can occur.

### **Parasympatholytic**

A medicine that blocks some signals of the parasympathetic nervous system. It may widen the airways, relax the bladder, reduce motion sickness, or speed a slow heartbeat, but can cause dry mouth, blurred vision, constipation, urinary retention, or confusion.

### **Parasympathomimetic**

A drug that mimics or enhances parasympathetic (cholinergic) activity. Direct agonists: muscarinic (pilocarpine—glaucoma, dry mouth) and nicotinic agonists.

### **Paroxetine**

Paroxetine is a selective serotonin-reuptake inhibitor used for depression, anxiety disorders, obsessive-compulsive disorder, PTSD, and some other conditions; sexual dysfunction, withdrawal symptoms, serotonin syndrome, and pregnancy-related risks require consideration.

### **Pediatric Nuclear Medicine**

Special considerations for children: lower doses, sedation often needed, distraction techniques. Dose reduction: weight-based activity, optimized protocols.

### **Pefloxacin**

A fluoroquinolone antibiotic that blocks bacterial

enzymes needed to copy and repair DNA. Its use is limited by resistance and potentially serious effects involving tendons, nerves, the heart, and the nervous system.

### **PEG**

An abbreviation that may refer to polyethylene glycol, a laxative that draws water into stool, or to a polyethylene-glycol chemical group used to modify medicines. The exact meaning depends on the clinical context.

### **Pegaspargase**

A long-acting enzyme medicine used in combination treatment for some acute lymphoblastic leukemias. It lowers asparagine needed by leukemia cells; allergic reactions, pancreatitis, liver injury, clotting problems, and bleeding are important risks.

### **Pegfilgrastim**

A long-acting growth factor that stimulates the bone marrow to make neutrophils. It helps prevent infection caused by chemotherapy-related low white-cell counts; bone pain and rare spleen rupture or serious inflammation can occur.

### **Penicillamine**

A medicine that binds certain metals and can also reduce immune activity. It is used selectively for copper overload and some autoimmune or connective-tissue disorders; serious blood, kidney, skin, or allergic reactions can occur.

### **Pentavalent Vaccine**

A WHO-prequalified five-in-one combination vaccine containing Diphtheria toxoid, Tetanus toxoid, whole-cell Pertussis antigen, Hepatitis B (HBsAg), and Haemophilus influenzae type b (Hib) conjugate, administered as a single intramuscular injection.

### **Pentobarbital**

A barbiturate medicine that depresses brain activity and can cause sedation or control seizures. It has a high risk of slowed breathing, coma, dependence, and fatal overdose, especially with alcohol or other sedatives.



---

### Perioperative Antibiotic Prophylaxis

Administration of an appropriate antimicrobial within a defined interval before incision to reduce infection risk for selected procedures. Choice, timing, redosing, and discontinuation depend on procedure, organisms, patient factors, and local guidance.

### Pertussis Vaccine

A vaccine containing selected antigens from *Bordetella pertussis*, given in combination vaccines to prevent whooping cough and reduce severe disease and transmission.

### Pharmaceutical

Relating to medicines, including their development, preparation, manufacture, testing, storage, or use.

### Pharmacodynamic Interaction

A drug interaction in which one drug changes the effect of another at its site of action without necessarily changing its concentration. Effects may be additive, synergistic, or antagonistic, such as combined respiratory depression or bleeding risk.

### Pharmacokinetic Interaction

A drug interaction that changes absorption, distribution, metabolism, or excretion of another drug, thereby altering its concentration over time. Enzyme inhibition, enzyme induction, transporter effects, and altered gastric pH are common mechanisms.

### Phosphodiesterase Inhibitor

A medicine that blocks enzymes which normally break down chemical signals inside cells. Different types relax airways or blood vessels, improve heart pumping, or treat erectile dysfunction; effects and risks depend on the specific type.

### Pneumococcal Conjugate Vaccine

A vaccine that joins parts of the pneumococcus bacterium to a carrier protein so the immune system can build lasting protection. It helps prevent pneumonia, meningitis, bloodstream infection, and other disease; product and schedule depend on age and risk.

### Pneumococcal Vaccine

A vaccine that protects against selected serotypes of *Streptococcus pneumoniae* (pneumococcus), which can cause pneumonia, meningitis, bloodstream infection, sinusitis, and otitis media. Available products include pneumococcal conjugate vaccines and pneumococcal polysaccharide vaccine; the product and schedule depend on age, previous doses, and medical risk.

### Poliovirus Vaccine

A vaccine that trains the immune system to prevent poliovirus infection and paralysis. Inactivated injectable and live oral forms are used in different programs; the schedule and product depend on age, health, and local public-health guidance.

### Prazosin

An alpha-1 blocker that relaxes blood vessels and may relax the bladder outlet. It is used for high blood pressure and, in selected patients, trauma-related nightmares or urinary symptoms; dizziness and fainting are most likely after starting or increasing the dose.

### Pre-Medication

Medicine given before a procedure, treatment, or exposure to prevent a predictable problem such as pain, nausea, allergy, infection, or anxiety.

### Prescription Drug

A medication whose dispensing or use is subject to prescription requirements under the applicable law. It requires professional oversight because of its risks, complexity, or need for monitoring; regulatory approval and prescribing rules vary by country and product.

### Procalcitonin-Guided Antibiotic Therapy

Use of blood procalcitonin measurements, together with examination and other tests, to help decide whether to start or stop antibiotics in selected infections. The result cannot by itself confirm bacterial infection or safely replace clinical judgment.



---

### Prostaglandin Antagonist

A drug that blocks a prostaglandin receptor or pathway, reducing selected prostaglandin-mediated effects such as inflammation, pain, uterine contraction, or gastric secretion.

### Protease Inhibitor

An antiviral medicine that blocks a virus enzyme needed to cut newly made proteins into working parts. In HIV treatment, it prevents formation of mature infectious virus and is used with other medicines.

### Proton Pump Inhibitor

See *Proton Pump Inhibitors*.

### Psychotropic Drug

A medicine that affects brain function and may change mood, thinking, behavior, sleep, or perception. The term includes antidepressants, antipsychotics, mood stabilizers, stimulants, and some medicines for anxiety or sleep.

### Psychotropic Medication

A medicine that changes brain activity and may affect mood, thoughts, behaviour, sleep, or perception. Examples include antidepressants, antipsychotics, mood stabilizers, and some medicines for anxiety or attention disorders.

### Rabies Vaccine

A vaccine containing inactivated rabies virus used for preexposure protection or postexposure prophylaxis after a possible rabies exposure, together with wound care and immune globulin when indicated.

### Rehabilitation Medicine

The medical specialty that helps people restore or adapt movement, communication, thinking, work, and daily activities after illness or injury. It is also called physiatry.

**Alternate Names:**  
physiatry.

### Renal Drug Excretion

Removal of medicines from the body by the kidneys into urine. Kidney disease can slow this process, allowing some medicines to build up and requiring dose changes or closer monitoring.

### Renin Inhibitor

A blood-pressure medicine that blocks renin, an enzyme involved in narrowing blood vessels and retaining salt. It lowers blood pressure but can raise potassium, affect kidney function, and harm a developing fetus.

### Resistance Antibiotic (Antibiotic Resistance)

Alternate index label for Antibiotic Resistance.

### Rock (Drug)

A street term commonly used for crack cocaine, a smokeable form of cocaine. It produces intense, short-lived stimulation but can cause dependence, dangerous heart rhythms, heart attack, stroke, seizures, overheating, or severe agitation.

### Scabicide

A medicine applied to the skin or taken by mouth to kill scabies mites. Treatment usually includes all close household or sexual contacts and washing or isolating recently used clothing and bedding to prevent reinfection.

### Selective Serotonin Reuptake Inhibitor

A medicine that increases serotonin signalling between nerve cells by slowing serotonin reuptake. It treats depression and several anxiety-related disorders; nausea, sexual difficulties, withdrawal symptoms, and rarely serotonin toxicity can occur.

### SGLT2 Inhibitor

A medicine that makes the kidneys pass more glucose and salt into the urine. It lowers blood sugar and can protect the heart and kidneys in selected people, but may cause dehydration, genital infections, or rarely ketoacidosis.

---

## Skeletal Muscle Relaxant

A medicine that reduces excessive contraction of skeletal muscles. It may relieve short-term muscle spasm or longer-term spasticity, but sleepiness, weakness, dizziness, and impaired breathing can occur with some drugs.

## Sleep Drug

A medicine used to help a person fall asleep, stay asleep, or maintain a regular sleep pattern. Benefits and risks vary widely; drowsiness, falls, impaired thinking, dependence, and interactions with alcohol or sedatives are important concerns.

## Steroid

A group of chemicals with a characteristic structure. In medicine, steroids include corticosteroids that reduce inflammation and natural or replacement hormones such as cortisol, estrogen, progesterone, and testosterone.

## Steroid Abuse

Steroid misuse is non-prescribed or excessive use of corticosteroids or anabolic-androgenic steroids, which can cause endocrine suppression, psychiatric effects, infection, metabolic disease, liver injury, cardiovascular harm, or tissue damage.

## Steroid Hormones

Lipid-soluble hormones derived from cholesterol, including cortisol, aldosterone, estrogen, progesterone, and testosterone.

## Steroid Injections - Tendon, Bursa, Joint

Injection of a corticosteroid, often with local anesthetic, into or around a joint, bursa, or tendon sheath to reduce inflammation and pain; injections directly into tendons are generally avoided because of rupture risk.

## Tamsulosin

Tamsulosin is a selective alpha-1A blocker that relaxes prostate and bladder-neck smooth muscle to improve urinary symptoms of benign prostatic hyperplasia; dizziness, postural hypotension, and ejaculatory dysfunction may occur.

## TCAs

Tricyclic antidepressants are medicines that increase the effects of certain brain chemicals. They can treat depression, nerve pain, and some other conditions but may cause sleepiness, dry mouth, constipation, low blood pressure, abnormal heart rhythms, or overdose toxicity.

## Tdap Vaccine

Tetanus, Diphtheria, and acellular Pertussis booster given at 11-12 years and during each pregnancy (27-36 weeks gestation). Replaces one dose of Td. Important for pertussis protection, particularly for protecting newborns through maternal antibody transfer. Also given to adolescents and adults to maintain tetanus and diphtheria immunity.

## Terazosin

An alpha-blocking medicine that relaxes the prostate and blood vessels. It can improve urine flow in prostate enlargement and lower blood pressure, but dizziness or fainting may occur, especially after the first dose or dose increases.

## Therapeutic Drug Levels

The range of medicine concentrations in blood associated with benefit while limiting toxicity for a particular drug. A measured level must be interpreted with the dose, timing, symptoms, kidney or liver function, and the person's response.

## Thrombolytic

A medicine that helps dissolve a blood clot by changing plasminogen into plasmin, which breaks down fibrin. It may be used urgently for selected strokes, heart attacks, or other life-threatening clots when benefits outweigh bleeding risk.

## Thrombolytic Drugs For Heart Attack

Medicines that dissolve a clot blocking a coronary artery during selected heart attacks. They can restore blood flow when urgent artery-opening procedures are unavailable or delayed, but may cause dangerous bleeding and are not suitable for everyone.

---

### Transfusion Medicine

The medical specialty that selects, prepares, tests, and gives blood or blood components such as red cells, platelets, plasma, and cryoprecipitate. It includes matching, preventing transfusion reactions, and monitoring patients.

### Travel Medicine

Medical care concerned with prevention and treatment of health problems related to travel.

### Tricyclic Antidepressant Overdose

Poisoning with a tricyclic antidepressant can cause drowsiness, confusion, dry mouth, seizures, low blood pressure, and dangerous heart rhythms. It is an emergency requiring immediate poison-control or hospital treatment.

### Tumor Necrosis Factor Inhibitor

A biological medicine that blocks tumor-necrosis factor, an immune signal involved in inflammation. It treats selected arthritis, psoriasis, and bowel diseases but can increase serious infection risk, so screening and monitoring are important.

### Tyrosine Kinase Inhibitor

A targeted medicine that blocks enzymes carrying growth signals inside cells. It can slow cancers driven by abnormal signaling, but side effects vary and may include diarrhea, skin changes, liver injury, high blood pressure, or heart problems.

### Uricosuric Agent

A medicine that increases renal excretion of uric acid, lowering serum urate in selected patients with gout. Adequate kidney function and hydration are important, and urinary uric-acid stones may be a concern.

### Urine Drug Screen

A preliminary urine test that looks for selected medicines or recreational drugs and their breakdown products. Results can be falsely positive or negative and may need confirmation with a more specific laboratory test.

### Uterotonic

A medicine that makes the uterus contract. It is used under medical supervision to start or strengthen labor, prevent or treat heavy bleeding after birth, or complete selected pregnancy-related procedures; excessive contraction can harm the mother or fetus.

### Vaccine Adjuvant

A substance added to some vaccines to enhance and prolong the immune response to the antigen. Examples include aluminum salts and newer adjuvant systems.

### Vaccine Therapy

Treatment that uses a vaccine to stimulate an immune response against a disease. It may prevent infection or, in selected cancer treatments, help the immune system recognize abnormal cells; its benefits depend on the specific vaccine and condition.

### Varicella Vaccine

A live attenuated vaccine that prevents or reduces the severity of chickenpox. The number and timing of doses depend on age, vaccination history, immune status, pregnancy, and local recommendations. It is generally avoided during pregnancy and in people with certain immune deficiencies.

### Xanthine Oxidase Inhibitor

A medicine that lowers uric-acid production by blocking an enzyme involved in breaking down substances from cells and food. It is used mainly to prevent gout attacks and, in selected cases, complications from rapid cancer-cell breakdown.

## Brand Medicines



# Medical Prefix

| Prefix          | Meaning                                                                                                                    | Example of use in medical terms                                                            |
|-----------------|----------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|
| Ab-             | A medical prefix meaning away from, off, or separating from,                                                               | abduction (movement away from the midline) and ablation (removal or destruction of tissue) |
| Acou-           | A medical combining form referring to hearing or sound,                                                                    | acoustic nerve, audiology, and acoumeter                                                   |
| Acro-           | Acro- is a medical word element meaning extremity, tip, or highest part,                                                   | acromegaly and acrocyanosis                                                                |
| Acu-            | A combining form meaning sharp, pointed, or related to acuity; in medical terms it may refer to sharp sensation or vision, | acuity, acute, or acupuncture, depending on the word formed                                |
| Ad-             | A medical prefix meaning toward, to, or near,                                                                              | adduction (movement toward the midline) and adrenal (near the kidney)                      |
| Aden-<br>Adeno- | A combining form meaning gland or glandular tissue,<br>A combining form referring to a gland or glandular tissue,          | adenitis and adenoma<br>adenovirus, adenoid, and adenocarcinoma                            |
| Aer-, Aero-     | A combining form meaning air or gas,                                                                                       | aerobic, aerosol, and aerophagia                                                           |
| Alb-<br>Amyl-   | A prefix derived from the Latin albus, meaning white,<br>A combining form referring to starch or a starch-like substance,  | albino and albinism<br>amylase and amyloid                                                 |
| Amylo-          | A combining form referring to starch-like material or amyloid protein deposits                                             | amyloidosis and amyloidoma                                                                 |
| An-             | A prefix meaning without, lacking, or not,                                                                                 | anoxia, anesthesia, and anencephaly                                                        |
| Aniso-          | A combining form meaning unequal or unequal in size, number, or degree                                                     | anisocytosis and anisometropia                                                             |
| Antero-         | A combining form meaning anterior or toward the front                                                                      | anteroposterior and anterolateral                                                          |
| Anti-           | A prefix meaning against, opposed to, or preventing a specified process or condition                                       | antibiotic and anticoagulant                                                               |
| Arthro-<br>Bi-  | A combining form meaning joint<br>A prefix meaning two, twice, or involving two sides or components                        | arthritis and arthroscopy<br>bilateral and bicuspid                                        |

| Prefix                             | Meaning                                                                                                                                    | Example of use in medical terms                                   |
|------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------|
| Bio-                               | A combining form meaning life or living organisms                                                                                          | biochemistry and biopsy                                           |
| Brachy-                            | A combining form meaning short or abnormally short                                                                                         | brachycephaly and brachydactyly                                   |
| Cephal-                            | A combining form meaning head                                                                                                              | cephalic and cephalalgia                                          |
| Cholangi-                          | A combining form meaning bile duct or bile vessel                                                                                          | cholangitis and cholangiography                                   |
| Circum-                            | A combining form meaning around or surrounding                                                                                             | circumcision and circumoral                                       |
| Colpo-                             | A combining form meaning vagina                                                                                                            | colposcopy and colpotomy                                          |
| Cranio-                            | A combining form meaning skull or cranium                                                                                                  | craniotomy and craniostylosis                                     |
| Cyto-                              | A combining form meaning cell                                                                                                              | cytology and cytotoxic                                            |
| Dacryo-                            | A combining form meaning tear or tear duct                                                                                                 | dacryocystitis and dacryostenosis                                 |
| Dactyl-, -Dactyl                   | A combining form or suffix meaning finger or toe                                                                                           | polydactyly and brachydactyly                                     |
| Dextro-                            | A combining form meaning right, toward the right, or sometimes clockwise                                                                   | dextrocardia and dextroposition                                   |
| Di-                                | A combining form meaning two, twice, or double                                                                                             | diplopia and diplegia                                             |
| Dia-                               | A combining form with meanings including through, across, apart, or complete, depending on the word                                        | dialysis and diagnosis                                            |
| Disto-                             | A combining form meaning distal, distant, or away from a point of attachment or origin                                                     | distal and distobuccal                                            |
| Dys-                               | A prefix meaning abnormal, difficult, painful, or impaired,                                                                                | dysphagia (difficulty swallowing) and dysuria (painful urination) |
| Entero-                            | A combining form meaning intestine or bowel,                                                                                               | enteropathy and enteroscopy                                       |
| Epi-                               | A prefix meaning on, upon, over, or above,                                                                                                 | epidermis and epigastric                                          |
| Extra-                             | A clinical condition or finding involving extra-. Its defining features, causes, evaluation, and management depend on the clinical context | extraocular and extracellular                                     |
| Gyne-                              | A combining form meaning woman or female,                                                                                                  | gynecology                                                        |
| Haem- / Haema- / Haemato- / Haemo- | Prefixes derived from the Greek word haima (blood), used in medical terminology to denote blood-related conditions and procedures          | hematology, hematoma, and hemoglobin                              |
| (Prefixes Relating To Blood)       |                                                                                                                                            |                                                                   |
| Hemi-                              | A prefix meaning half or one side,                                                                                                         | hemiplegia and hemithorax                                         |
| Hepat-                             | A combining form meaning liver,                                                                                                            | hepatic and hepatocellular                                        |
| Hepato-                            | A combining form meaning liver,                                                                                                            | hepatobiliary and hepatomegaly                                    |
| Hetero-                            | A prefix meaning different, other, or unequal,                                                                                             | heterozygous and heterogeneous                                    |
| Histo-                             | A combining form meaning tissue,                                                                                                           | histology and histopathology                                      |
| Homo-                              | A prefix meaning same or alike,                                                                                                            | homozygous and homologous                                         |
| Hyper-                             | A prefix meaning excessive, above normal, or increased,                                                                                    | hyperglycemia and hypertension                                    |
| Hypo-                              | A prefix meaning below, deficient, reduced, or less than normal,                                                                           | hypoglycemia and hypothermia                                      |



| Prefix          | Meaning                                                                                                                                                     | Example of use in medical terms                                                             |
|-----------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------|
| Iatr-           | A combining form meaning medical treatment or physician,                                                                                                    | iatrogenic and iatrogeny                                                                    |
| Infra-          | A medical term referring to infra-. Its precise meaning and clinical importance depend on the context in which it is used                                   | infrapatellar and infraorbital                                                              |
| Intra-          | Intra- is an anatomical structure, tissue relationship, physiologic process, or body function; its location and mechanism explain its clinical significance | intraocular and intravenous                                                                 |
| Iso-            | A prefix meaning equal, same, or uniform,                                                                                                                   | isotonic and isometric                                                                      |
| Juxta-          | A prefix meaning next to, beside, or adjoining,                                                                                                             | juxtaglomerular                                                                             |
| Kerato-         | A combining form relating to the cornea or to horny or keratinized tissue,                                                                                  | keratitis and keratosis                                                                     |
| Leuk-           | A combining form meaning white, especially in terms relating to white blood cells,                                                                          | leukocyte                                                                                   |
| Leuko-          | A combining form meaning white,                                                                                                                             | leukoderma or leukopenia                                                                    |
| Levo-           | A prefix meaning left or turning toward the left, often used in chemistry or to describe spatial orientation                                                | levocardia and levodopa                                                                     |
| Lieno-          | A combining form meaning relating to the spleen                                                                                                             | lienorenal and lienectomy                                                                   |
| Lipo-           | A combining form meaning relating to fat or lipids                                                                                                          | lipoma and lipoprotein                                                                      |
| Litho-          | A combining form meaning stone,                                                                                                                             | lithotripsy and nephrolithiasis                                                             |
| Macro- (Prefix) | A prefix meaning large, abnormally large, or on a large scale,                                                                                              | macrocytic, meaning involving abnormally large cells                                        |
| Medical Prefix  | A prefix used in medical terminology to modify a word's meaning                                                                                             | hyper- (above or excessive), hypo- (below or deficient), macro- (large), and micro- (small) |
| Mega-           | A combining form meaning large, great, or abnormally enlarged                                                                                               | megacolon and megakaryocyte                                                                 |
| Mega- (Prefix)  | A prefix derived from Greek meaning large, great, or abnormally enlarged,                                                                                   | megacolon, megacardia, and megaloccephaly                                                   |
| Melan- (Prefix) | A combining form meaning dark or black, derived from the Greek melas; it refers especially to melanin or dark pigmentation,                                 | melanoma and melanocyte                                                                     |
| Mes-            | A combining form meaning middle                                                                                                                             | mesencephalon and mesoderm                                                                  |
| Meso-           | A combining form meaning middle                                                                                                                             | mesothelioma and mesocolon                                                                  |
| Mi-             | A combining form meaning small or relating to muscle, depending on the word in which it appears                                                             | micturition and miosis                                                                      |
| Micro-          | A combining form meaning small or millionth, depending on context                                                                                           | microscopy and microcephaly                                                                 |
| Myc-            | A combining form meaning fungus or pertaining to fungi, derived from the Greek mykes                                                                        | mycosis and mycology                                                                        |
| Myco-           | The term Myco- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                        | mycosis and mycology                                                                        |
| Myel-           | A combining form relating to bone marrow or the spinal cord, depending on the word                                                                          | myelitis and myelopathy                                                                     |

| Prefix           | Meaning                                                                                                                                                 | Example of use in medical terms             |
|------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------|
| Myo- (Prefix)    | A medical term referring to myo- (prefix). Its clinical meaning depends on the context in which it is used                                              | myopathy and myocarditis                    |
| Naso-            | The term Naso- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                    | nasogastric and nasopharyngeal              |
| Neo- (Prefix)    | Neo- is a prefix meaning new or recently formed                                                                                                         | neoplasm and neonatal                       |
| Nephro-          | The term Nephro- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                  | nephritis and nephrectomy                   |
| Oligo-           | A combining form meaning few, scanty, or deficient                                                                                                      | oliguria and oligospermia                   |
| Oligo- (Prefix)  | Oligo- is a prefix meaning few, scanty, or small in number or amount                                                                                    | oliguria and oligospermia                   |
| Onco-            | A combining form relating to tumors or cancer                                                                                                           | oncology and oncogenesis                    |
| Onycho-          | A combining form relating to the nail                                                                                                                   | onychomycosis and onycholysis               |
| Onycho- (Prefix) | Onycho- is a prefix meaning relating to a nail or claw                                                                                                  | onychomycosis and onycholysis               |
| Ortho- (Prefix)  | Ortho- is a prefix meaning straight, correct, upright, or related to orthopedic care depending on the term                                              | orthopedics and orthostatic                 |
| Osteo- (Prefix)  | Osteo- is a prefix meaning relating to bone                                                                                                             | osteomyelitis and osteoporosis              |
| Ot-              | The term Ot- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                      | otitis and otalgia                          |
| Oto-             | The term Oto- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                     | otology and otoscope                        |
| Oxy              | Oxy- is a prefix relating to oxygen, sharpness, or acidity depending on the medical or chemical term                                                    | oxygen and oxyhemoglobin                    |
| Pachy- (Prefix)  | A word-forming prefix meaning thick or thickness,                                                                                                       | pachyonychia, pachyderma, and pachycephaly  |
| Para             | Para- is a prefix meaning beside, beyond, abnormal, or related to childbirth depending on the word; its exact meaning must be read in context           | parathyroid and paranasal                   |
| Para-            | A combining form meaning beside, beyond, abnormal, or closely related, depending on the word                                                            | paracentesis and paraesthesia               |
| Para- (Prefix)   | Para- is a medical prefix whose meaning varies by use, including beside, beyond, abnormal, or the number of births reaching a specified gestational age | parathyroid and parity in obstetric history |
| Patho-           | The term Patho- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                   | pathology and pathogenesis                  |
| Peri-            | The term Peri- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                    | pericardium and periosteum                  |
| Phago-           | The term Phago- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                   | phagocytosis and phagocyte                  |
| Phlebo- (Prefix) | Phlebo- is a medical prefix meaning relating to a vein,                                                                                                 | phlebitis, phlebectomy, or phlebography     |

| Prefix            | Meaning                                                                                                                                                 | Example of use in medical terms                     |
|-------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------|
| Pneumo-           | The term Pneumo- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                  | pneumothorax and pneumonia                          |
| Poly-             | A combining form meaning many or much                                                                                                                   | polyuria and polyneuropathy                         |
| Post              | Post- is a prefix meaning after,                                                                                                                        | post-operative, postpartum, or postprandial         |
| Post-             | A combining form meaning after or behind                                                                                                                | postnatal and postprandial                          |
| Pre-              | A combining form meaning before or in front of                                                                                                          | prenatal and preoperative                           |
| Prefix            | A prefix is a word element placed before a root to modify meaning; medical prefixes commonly indicate location, number, time, direction, or abnormality | hyper-, hypo-, peri-, and brady-                    |
| Prefixes, Medical | Word beginnings that modify medical terms by indicating location, number, time, speed, size, direction, deficiency, excess, or other meaning,           | hyper-, hypo-, peri-, and brady-                    |
| Pro-              | A combining form meaning before, forward, or in favor of                                                                                                | prolactin and prognosis                             |
| Pseudo-           | A combining form meaning false or resembling but not truly                                                                                              | pseudocyst and pseudarthrosis                       |
| Retro-            | A combining form meaning backward, behind, or posterior                                                                                                 | retroperitoneal and retrosternal                    |
| Salpingo-         | A combining form relating to a fallopian tube                                                                                                           | salpingitis and salpingectomy                       |
| Sarco-            | A combining form meaning flesh or soft tissue                                                                                                           | sarcoma and sarcoidosis                             |
| Schizo-           | A combining form relating to splitting or schizophrenia                                                                                                 | schizophrenia and schizoid                          |
| Scler-            | The term Scler- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                   | sclerosis and scleritis                             |
| Sclero-           | The term Sclero- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                  | scleroderma and sclerotherapy                       |
| Sub-              | A combining form meaning under, below, or less than normal                                                                                              | subcutaneous and subdural                           |
| Super-            | A combining form meaning above, excessive, or superior                                                                                                  | superinfection and supernumerary                    |
| Supra-            | A combining form meaning above or beyond                                                                                                                | suprarenal and suprapubic                           |
| Syn-              | A combining form meaning together, with, or joined                                                                                                      | syndactyly and synovitis                            |
| Teno-             | A combining form relating to a tendon                                                                                                                   | tenosynovitis and tenodesis                         |
| Thermo-           | A combining form relating to heat or temperature                                                                                                        | thermoregulation and thermotherapy                  |
| Toc- (Prefix)     | Toc- is a prefix used in some medical terms and often relates to childbirth or labor,                                                                   | tocolysis; the complete word determines its meaning |
| Toco-             | The term Toco- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                    | tocography and tocolysis                            |
| Toko- (Prefix)    | Toko- is a prefix relating to childbirth or labor, derived from Greek terms for birth                                                                   | tocolysis and tokodynamometry                       |
| Tom-              | The term Tom- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used                     | tomography and tomodensitometry                     |

| Prefix           | Meaning                                                                                                                                         | Example of use in medical terms                             |
|------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------|
| Tomo-            | The term Tomo- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used            | tomography and tomographic                                  |
| Trans- (Prefix)  | The term Trans- (prefix) is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used  | transdermal and transcatheter                               |
| Trich- (Prefix)  | The term Trich- (prefix) is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used  | trichiasis and trichology                                   |
| Tricho-          | A combining form relating to hair                                                                                                               | trichoscopy and trichotillomania                            |
| Tricho- (Prefix) | The term Tricho- (prefix) is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used | trichorrhexis and trichomycosis                             |
| Tympano-         | The term Tympano- is a medical word element or prefix whose precise meaning depends on the word or clinical context in which it is used         | tympanoplasty and tympanometry                              |
| Xanth- (Prefix)  | Xanth- is a medical word element meaning yellow or yellowish,                                                                                   | xanthoma, xanthelasma, and xanthopsia                       |
| Xen- (Prefix)    | A medical word element meaning foreign, strange, or originating outside the usual organism or system,                                           | xenograft (a tissue transplanted between different species) |
| Xeno- (Prefix)   | Xeno- is a medical word element meaning foreign, strange, or from another species,                                                              | xenograft and xenotransplantation                           |
| Xero-            | Xero- is a medical word element meaning dry or dryness,                                                                                         | xeroderma, xerophthalmia, and xerostomia                    |

## Complete reference tables

These categorized tables reproduce the prefix reference structure from the Pressbooks resource cited below. Examples are retained as learning aids; the meaning of a word part can vary by term.

| Prefix                 | Meaning                  | Example of use in medical terms |
|------------------------|--------------------------|---------------------------------|
| <b>Common prefixes</b> |                          |                                 |
| A-, An-                | Without; lacking         | Anemia                          |
| Andr/o-                | Male                     | Androgen                        |
| Anti-                  | Against                  | Anticholinergic drugs           |
| Auto-                  | Self                     | Autocrine                       |
| Bio-                   | Life                     | Biology                         |
| Chem/o-                | Chemistry                | Chemotherapy                    |
| Contra-                | Against                  | Contraception                   |
| Cyt/o-                 | Cell                     | Cytokine                        |
| Dis-                   | Separation; taking apart | Dissection                      |
| Dys-                   | Difficult; abnormal      | Dyspnea                         |
| Eu-                    | Good; well               | Eupnea                          |
| Fibr/o-                | Fiber                    | Fibrosis                        |
| Gluco-, Glyco-         | Glucose; sugar           | Glycogen                        |

| Prefix                                     | Meaning                             | Example of use in medical terms |
|--------------------------------------------|-------------------------------------|---------------------------------|
| Gyn/o-, Gynec-                             | Female                              | Gynecology                      |
| Hydr/o-                                    | Water                               | Hydrocephalus                   |
| Idio-                                      | Self; one's own                     | Idiopathic                      |
| Lyso-, Lys-                                | Break down; destruction; dissolving | Lysosome                        |
| Mal-                                       | Bad; abnormal                       | Malignant                       |
| Myc/o-                                     | Fungus                              | Mycetoma                        |
| Necr/o-                                    | Death                               | Necrosis                        |
| Neo-                                       | New                                 | Neonate                         |
| Oxy-                                       | Sharp; acute; oxygen                | Oxytocin                        |
| Pan-, Pant/o-                              | All or everywhere                   | Pancytopenia                    |
| Pharmaco-                                  | Drug; medicine                      | Pharmacist                      |
| Re-                                        | Again; backward                     | Rejuvenation                    |
| Somat/o-, Somatico-                        | Body; bodily                        | Somatic cell                    |
| <b>Body-part prefixes</b>                  |                                     |                                 |
| Acous/o-                                   | Hearing                             | Acoustic meatus                 |
| Aden/o-                                    | Gland                               | Adenoid                         |
| Adip/o-                                    | Fat                                 | Adipocyte                       |
| Adren/o-                                   | Gland                               | Adrenal cortex                  |
| Angi/o-                                    | Blood vessel                        | Angioplasty                     |
| Arteri/o-                                  | Artery                              | Arteriole                       |
| Arthr/o-                                   | Joint                               | Arthroplasty                    |
| Bucc/o-                                    | Cheek                               | Buccal cavity                   |
| Bronch/i-                                  | Bronchus                            | Bronchioles                     |
| Burs/o-                                    | Bursa                               | Bursa                           |
| Carcin/o-                                  | Cancer                              | Basal cell carcinoma            |
| Cardi/o-                                   | Heart                               | Cardiology                      |
| Cephal/o-                                  | Head                                | Cephalic flexure                |
| Chol-                                      | Bile                                | Cholesterol                     |
| Chondri-                                   | Cartilage                           | Chondrosarcoma                  |
| Coron-                                     | Heart                               | Coronary arteries               |
| Cost-                                      | Rib                                 | Costal cartilage                |
| Crani/o-                                   | Brain                               | Cranium                         |
| Cutane-                                    | Skin                                | Cutaneous                       |
| Cyst/o-, Cysti-                            | Bladder or sac                      | Cystoscopy                      |
| Derm-, Dermat/o-                           | Skin                                | Dermatologist                   |
| Duoden/o-                                  | Duodenum                            | Duodenitis                      |
| Gastr-                                     | Stomach                             | Gastrectomy                     |
| Gloss-                                     | Tongue                              | Glossectomy                     |
| Hem-, Hema-,<br>Hemat-, Hemo-,<br>Hemat/o- | Blood                               | Hematopoiesis                   |
| Hepat/o-, Hepatico-                        | Liver                               | Hepatic portal system           |
| Hist/o-, Histio-                           | Tissue                              | Histology                       |
| Hyster/o-                                  | Uterus                              | Hysterectomy                    |
| Ileo-                                      | Ileum                               | Ileostomy                       |
| Ischi/o-                                   | Ischium                             | Ischial tuberosity              |
| Kerat/o-                                   | Cornea (eye or skin)                | Keratin                         |
| Lacrim/o-                                  | Tear                                | Lacrimal fluid                  |
| Lact/o-, Lacti-                            | Milk                                | Lactose                         |
| Laryng/o-                                  | Larynx                              | Laryngitis                      |
| Lingu/o-                                   | Tongue                              | Lingual tonsil                  |

| Prefix                       | Meaning                    | Example of use in medical terms |
|------------------------------|----------------------------|---------------------------------|
| Lip/o-                       | Fat                        | Lipolysis                       |
| Lymph/o-                     | Lymph                      | Lymphocyte                      |
| Mamm-, Mast/o-               | Breast                     | Mammary glands                  |
| Mening/o-                    | Meninges                   | Meningitis                      |
| Muscul/o-                    | Muscle                     | Musculoskeletal                 |
| My/o-                        | Muscle                     | Myocardium                      |
| Myel/o-                      | Spinal cord or bone marrow | Myelin                          |
| Nephr/o-                     | Kidney                     | Nephron                         |
| Neur/i-, Neur/o-             | Nerve                      | Neuron                          |
| Oculo-                       | Eye                        | Oculomotor nerve                |
| Onco-                        | Tumor; bulk; volume        | Oncogene                        |
| Onych/o-                     | Fingernail; toenail        | Onychodystrophy                 |
| Oo-                          | Egg; ovary                 | Oocyte                          |
| Oophor/o-                    | Ovary                      | Oophorectomy                    |
| Op-, Opt-                    | Vision                     | Optic nerve                     |
| Ophthalm/o-                  | Eye                        | Ophthalmic artery               |
| Orchid/o-, Orchio-           | Testis                     | Orchidectomy                    |
| Orth/o-                      | Straight; normal; correct  | Orthostatic                     |
| Osseo-                       | Bony                       | Osseous tissue                  |
| Ossi-                        | Bone                       | Ossicles                        |
| Ost-, Oste/o-                | Bone                       | Osteoporosis                    |
| Ot/o-                        | Ear                        | Otolaryngologist                |
| Ovar/i-, Ovario-, Ovi-, Ovo- | Ovary                      | Ovarian follicle                |
| Phalang-                     | Phalanx                    | Phalanges                       |
| Pharyng/o-                   | Pharynx; throat            | Pharyngeal tonsil               |
| Phleb/o-                     | Vein                       | Phlebotomist                    |
| Phren/i-, Phreno-, Phrenico- | Diaphragm                  | Phrenic nerve                   |
| Pleur-, Pleur/a-, Pleur/o-   | Rib; pleura                | Pleural cavity                  |
| Pneum/a-, Pneumat/o-         | Air; lung                  | Pneumonia                       |
| Proct/o-                     | Anus; rectum               | Proctoscopy                     |
| Prostat-                     | Prostate                   | Prostatectomy                   |
| Pseudo-                      | False                      | Pseudostratified                |
| Psych/o-, Psyche-            | Mind                       | Psychiatrist                    |
| Radio-                       | Radiation; radius          | Radioisotopes                   |
| Ren/o-                       | Kidney                     | Renal cortex                    |
| Retin-                       | Retina                     | Retinitis pigmentosa            |
| Rhin/o-                      | Nose                       | Rhinoscope                      |
| Salping/o-                   | Tube                       | Salpingo-oophorectomy           |
| Sarco-                       | Muscular; flesh-like       | Sarcomere                       |
| Schiz/o-                     | Split; cleft               | Schizophrenia                   |
| Sclera-, Sclero-             | Hardness                   | Sclerosis                       |
| Sigmoid/o-                   | Sigmoid colon              | Sigmoidoscopy                   |
| Sperma-, Sperm-, Spermato-   | Sperm                      | Spermatocyte                    |
| Splen/o-                     | Spleen                     | Splenomegaly                    |
| Sten/o-                      | Narrowed; blocked          | Stenosis                        |
| Stern-                       | Sternum                    | Sternoclavicular joint          |



| Prefix                                      | Meaning              | Example of use in medical terms |
|---------------------------------------------|----------------------|---------------------------------|
| Stom/a-, Stomat/o-                          | Mouth                | Stomatitis                      |
| Thorac/o-, Thoracico-                       | Chest                | Thoracic cavity                 |
| Thromb/o-                                   | Blood clot           | Thrombolytic                    |
| Thyr/o-                                     | Thyroid gland        | Thyroiditis                     |
| Trache/o-                                   | Trachea              | Trachealis                      |
| Tympan/o-                                   | Eardrum              | Tympanic membrane               |
| Ur/o-                                       | Urine                | Urologist                       |
| Vagin-                                      | Vagina               | Vaginal                         |
| Varic/o-                                    | Duct; blood vessel   | Varicose veins                  |
| Vasculo-                                    | Blood vessel         | Vasculitis                      |
| Ven/o-                                      | Vein                 | Venae cavae                     |
| Vertbr-                                     | Vertebra; spine      | Vertebral column                |
| <b>Color prefixes</b>                       |                      |                                 |
| Chlor/o-                                    | Green                | Chlorophyll                     |
| Chrom-, Chromato-                           | Color                | Chromosome                      |
| Cyano-                                      | Blue                 | Cyanosis                        |
| Erythr/o-                                   | Red                  | Erythrocyte                     |
| Leuk/o-                                     | White                | Leukocyte                       |
| Melan/o-                                    | Black                | Melanin                         |
| <b>Physical property and shape prefixes</b> |                      |                                 |
| Cry/o-                                      | Cold                 | Cryotherapy                     |
| Elect-                                      | Electrical activity  | Electrocardiogram               |
| Kin/o-, Kine-, Kinesi/o-                    | Movement             | Kinetic energy                  |
| Kyphy/o-                                    | Humped               | Kyphosis                        |
| Rhabd/o-                                    | Rod-shaped; striated | Rhabdomyosarcoma                |
| Phot/o-                                     | Light                | Photoreceptor                   |
| Reticul/o-                                  | Net                  | Reticulocytes                   |
| Scoli/o-                                    | Twisted              | Scoliosis                       |
| Therm/o-                                    | Heat                 | Thermotherapy                   |
| <b>Direction and position prefixes</b>      |                      |                                 |
| Ab-, Abs-                                   | Away from            | Abductor                        |
| Ad-                                         | Towards              | Adductor                        |
| Ante-                                       | Before; forward      | Antenatal                       |
| Circum-                                     | Around               | Circumcision                    |
| Cycl-                                       | Circle; cycle        | Cyclic neutropenia              |
| De-                                         | Away from; ending    | Dehydration                     |
| Dia-                                        | Across; through      | Diagnosis                       |
| Ect/o-, Exo-                                | Outer; outside       | Exocrine gland                  |
| End/o-, Ent-, Enter/o-                      | Within; inner        | Endocrine gland                 |
| Epi-                                        | Upon; outside of     | Epidermis                       |
| Ex-, Extra-                                 | Beyond               | Expiration                      |
| Infra-                                      | Beneath; below       | Infratemporal fossa             |
| Inter-                                      | Between              | Interstitial fluid              |
| Intra-                                      | Within               | Intracellular fluid             |
| Meso-                                       | Middle               | Mesoderm                        |
| Meta-                                       | Beyond; change       | Metabolism                      |
| Para-                                       | Alongside; abnormal  | Parathyroid glands              |

---

| Prefix                      | Meaning                  | Example of use in medical terms |
|-----------------------------|--------------------------|---------------------------------|
| Path/o-                     | Disease                  | Pathologist                     |
| Peri-                       | Around                   | Pericardium                     |
| Post-                       | Behind; after            | Postpartum                      |
| Pre-                        | Before; in front         | Precancerous                    |
| Retro-                      | Backward; behind         | Retroperitoneum                 |
| Sub-                        | Under                    | Subcutaneous layer              |
| Super-                      | Above                    | Superior                        |
| Supra-                      | Above; upon              | Supraglottis                    |
| Sy-, Syl-, Sym-, Syn-, Sys- | Together                 | Syndrome                        |
| Trans-                      | Across; through          | Transdermal                     |
| <b>Quantity prefixes</b>    |                          |                                 |
| Bi-                         | Two                      | Biceps                          |
| Brady-                      | Slow                     | Bradycardia                     |
| Diplo-                      | Double                   | Diploid                         |
| Hemi-                       | Half                     | Hemihypertrophy                 |
| Hetero-                     | Other; different         | Heterogeneous                   |
| Homo-                       | Same                     | Homozygous genotype             |
| Hyper-                      | Above; beyond; excessive | Hypertension                    |
| Hypo-                       | Under; deficient         | Hypotension                     |
| Iso-                        | Equal; like              | Isointense                      |
| Macro-                      | Large; long; big         | Macrophage                      |
| Mic-, Micro-                | Small                    | Microglia                       |
| Mon-, Mono-                 | One                      | Monocyte                        |
| Olig/o-                     | Few; little              | Oliguria                        |
| Poly-                       | Many; excessive          | Polyuria                        |
| Quadri-                     | Four                     | Quadriceps                      |
| Semi-                       | Half                     | Semilunar valves                |
| Tachy-                      | Fast                     | Tachycardia                     |
| Tetra-                      | Four                     | Tetralogy of Fallot             |
| Tri-                        | Three                    | Triceps                         |
| Uni-                        | One                      | Unicellular                     |

---

---

Source and license: Andrea Nelson and Katherine Greene, *Medical Terminology for Healthcare Professions*, “Prefixes and Suffixes,” University of West Florida Libraries, CC BY 4.0. <https://pressbooks.uwf.edu/medicalterminology/chapter/prefixes-and-suffixes/>

# Medical Suffixes

| Suffix           | Meaning                                                                                                                              | Example of use in medical terms             |
|------------------|--------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------|
| -Stomy           | A suffix meaning surgical creation of an opening,                                                                                    | colostomy or gastrostomy                    |
| -Tomy            | A suffix meaning incision or cutting into                                                                                            | laparotomy or tracheotomy                   |
| Cidal            | A suffix meaning killing or destroying,                                                                                              | bactericidal or fungicidal                  |
| Cide             | A suffix meaning an agent or process that kills,                                                                                     | pesticide, bactericide, or homicide         |
| Cyte             | A suffix meaning cell,                                                                                                               | erythrocyte or leukocyte                    |
| Cytosis          | A suffix or term indicating a condition involving cells or, in some contexts, an increase in the number of a particular type of cell | leukocytosis or erythrocytosis              |
| Dactyl-, -Dactyl | A combining form or suffix meaning finger or toe                                                                                     | polydactyly or brachydactyly                |
| Ectomy           | A suffix meaning surgical removal or excision,                                                                                       | appendectomy or mastectomy                  |
| Emia             | A suffix meaning a condition of the blood or a substance present in the blood,                                                       | anemia or bacteremia                        |
| Iatrics          | A suffix or word element referring to medical treatment or a medical specialty,                                                      | geriatrics and pediatrics                   |
| Iatry            | A suffix meaning medical treatment or a medical specialty,                                                                           | psychiatry and podiatry                     |
| Itis             | A suffix meaning inflammation,                                                                                                       | dermatitis, bronchitis, and arthritis       |
| Kalemia          | A suffix or term referring to the concentration of potassium in the blood,                                                           | hypokalemia or hyperkalemia                 |
| MAB              | Abbreviation for monoclonal antibody; the suffix -mab in a generic drug name indicates a monoclonal antibody,                        | adalimumab or rituximab                     |
| Oma              | -oma is a suffix commonly indicating a tumor or mass, although some words ending in -oma describe non-neoplastic conditions          | lipoma or carcinoma                         |
| Pathy            | A suffix meaning disease, disorder, or abnormality,                                                                                  | neuropathy, cardiomyopathy, and nephropathy |
| Plegia           | -plegia is a suffix meaning paralysis,                                                                                               | paraplegia, hemiplegia, or quadriplegia     |

## Complete reference tables

| Suffix                    | Meaning                                       | Example of use in medical terms |
|---------------------------|-----------------------------------------------|---------------------------------|
| <b>Common suffixes</b>    |                                               |                                 |
| -ac                       | Pertaining to                                 | Cardiac                         |
| -blast, -blasto, -blastic | Bud; germ                                     | Myeloblast                      |
| -cyte, -cytic             | Cell                                          | Thrombocyte                     |
| -dynia                    | Pain; swelling                                | Thoracodynia                    |
| -eal, -ial                | Pertaining to                                 | Esophageal                      |
| -ectasis                  | Expansion; dilation                           | Atelectasis                     |
| -emia                     | Blood condition                               | Anemia                          |
| -ia                       | Condition                                     | Hemophilia                      |
| -iasis                    | Condition; formation of                       | Psoriasis                       |
| -ism                      | Condition                                     | Hypothyroidism                  |
| -ites, -itis              | Inflammation                                  | Arthritis                       |
| -ity                      | Pertaining to                                 | Immunity                        |
| -ium                      | Structure or tissue                           | Epithelium                      |
| -lysis, -lytic            | Breakdown; destruction; dissolving            | Osteolytic                      |
| -malacia                  | Softening                                     | Osteomalacia                    |
| -megaly                   | Enlargement                                   | Acromegaly                      |
| -oid                      | Resembling                                    | Arachnoid trabeculae            |
| -oma                      | Tumor                                         | Angiosarcoma                    |
| -osis                     | Condition; usually abnormal                   | Endometriosis                   |
| -ous                      | Pertaining to                                 | Aqueous                         |
| -pathy                    | Disease                                       | Lymphadenopathy                 |
| -penia                    | Deficiency; lack of                           | Thrombocytopenia                |
| -phagia, -phagy           | Eating; swallowing                            | Dysphagia                       |
| -phasia                   | Speech                                        | Aphasia                         |
| -plasia, -plastic         | Growth                                        | Hyperplasia                     |
| -plegia                   | Paralysis                                     | Hemiplegia                      |
| -pnea                     | Breathing                                     | Sleep apnea                     |
| -poiesis                  | Production                                    | Hemopoiesis                     |
| -ptosis                   | Falling; drooping                             | Apoptosis                       |
| -rrhage, -rrhagic         | Bleeding                                      | Hemorrhage                      |
| -rrhea                    | Flow or discharge                             | Diarrhea                        |
| -sclerosis                | Hardening                                     | Arteriosclerosis                |
| -sis                      | Condition                                     | Agranulocytosis                 |
| -stasis                   | Level; unchanging                             | Homeostasis                     |
| -trophy                   | Growth                                        | Hypertrophy                     |
| -uria                     | In the urine                                  | Anuria                          |
| <b>Procedure suffixes</b> |                                               |                                 |
| -centesis                 | Surgical puncture to remove fluid             | Thoracentesis                   |
| -desis                    | Surgical binding                              | Pleurodesis                     |
| -ectomy                   | Cut out; removal                              | Mastectomy                      |
| -gram                     | Record; picture                               | Electrocardiogram               |
| -graph                    | Instrument used to create a record or picture | Electrocardiograph              |
| -graphy                   | To record or take a picture                   | Echocardiography                |
| -meter                    | Device used for measuring                     | Sphygmomanometer                |
| -opsy                     | Visual examination                            | Biopsy                          |

---

| Suffix  | Meaning                 | Example of use in medical terms |
|---------|-------------------------|---------------------------------|
| -ostomy | Opening                 | Colostomy                       |
| -otomy  | Incision                | Laparotomy                      |
| -pexy   | Surgical fixation       | Oophoropexy                     |
| -plasty | Surgical reconstruction | Vertebroplasty                  |
| -scope  | For examining           | Endoscope                       |
| -scopy  | Examine                 | Endoscopy                       |

---

---

Source and license: Andrea Nelson and Katherine Greene, *Medical Terminology for Healthcare Professions*, “Prefixes and Suffixes,” University of West Florida Libraries, CC BY 4.0. <https://pressbooks.uwf.edu/medicalterminology/chapter/prefixes-and-suffixes/>





# Alternate Names

| Primary term                            | Alternate names                         |
|-----------------------------------------|-----------------------------------------|
| ACE                                     | angiotensin-converting enzyme           |
| ACE Inhibitor                           | angiotensin-converting enzyme inhibitor |
| Acquired Immunodeficiency Syndrome      | AIDS                                    |
| Adenosine Triphosphate                  | ATP                                     |
| AIDS                                    | acquired immunodeficiency syndrome      |
| Alcoholism                              | alcohol dependence: A chronic           |
| Alimentary Canal                        | gastrointestinal                        |
| AMD                                     | age-related macular degeneration        |
| Angiotensin-Converting Enzyme           | ACE                                     |
| Angiotensin-Converting Enzyme Inhibitor | ACE                                     |
| Amyloid Precursor Protein               | APP                                     |
| ARB                                     | angiotensin II receptor blockers        |
| ATP                                     | adenosine triphosphate                  |
| AV Node                                 | atrioventricular node                   |
| Balloon Dilation                        | Balloon dilatation                      |
| BMD                                     | bone mineral density                    |
| BMI                                     | body-mass index                         |
| BPH                                     | benign prostatic hyperplasia            |
| CABG                                    | coronary artery bypass graft            |
| CBC                                     | complete blood count                    |
|                                         | mdash                                   |
| CCU                                     | coronary care unit                      |
| Cerebral Hemorrhage                     | hemorrhagic stroke                      |
| CFS                                     | chronic fatigue syndrome                |
| CK                                      | creatine kinase                         |
| COX-2 Inhibitors                        | cyclooxygenase-2 inhibitors             |
| CPAP                                    | continuous positive airway pressure     |
| CPR                                     | cardiopulmonary resuscitation           |
| CRP                                     | C-reactive protein                      |
| DNA                                     | deoxyribonucleic acid                   |
| DNR                                     | do-not-resuscitate order                |
| DRI                                     | dietary reference intake                |
| EBCT                                    | electron-beam computed tomography       |
| ECG                                     | electrocardiogram                       |
| ED                                      | erectile dysfunction                    |

---

| Primary term                 | Alternate names                                                                     |
|------------------------------|-------------------------------------------------------------------------------------|
| EEG                          | electroencephalogram                                                                |
| EKG                          | electrocardiogram                                                                   |
| EMG                          | electromyography                                                                    |
|                              | mdash                                                                               |
| FSH                          | follicle-stimulating hormone                                                        |
| GERD                         | gastroesophageal reflux disease                                                     |
| GFR                          | glomerular filtration rate                                                          |
| Gingiva                      | gums: a form of oral tissue that covers the roots of teeth and surrounding bone     |
| GnRH                         | gonadotropin-releasing hormone                                                      |
| HbA1C                        | hemoglobin A1c                                                                      |
| HDL                          | high-density lipoprotein                                                            |
| HEPA Filter                  | high-efficiency particulate air                                                     |
| HIV                          | human immunodeficiency virus                                                        |
| HLA                          | human leukocyte antigen                                                             |
| HPA                          | hypothalamic-pituitary-adrenal axis                                                 |
| Human Immunodeficiency Virus | HIV                                                                                 |
| IBD                          | inflammatory bowel disease                                                          |
| ICD                          | implantable cardioverter defibrillator                                              |
| IgE                          | immunoglobulin E                                                                    |
| IU                           | international unit                                                                  |
| LDL                          | low-density lipoprotein                                                             |
| LH                           | luteinizing hormone                                                                 |
| LVAD                         | left-ventricular assist device                                                      |
| MAOIs                        | monoamine oxidase inhibitors                                                        |
| METs                         | metabolic equivalents                                                               |
| MHC Molecule                 | major histocompatibility complex molecule                                           |
| Mitral Valve Prolapse        | Barlow's Syndrome; Barlow Syndrome (Mitral Valve Prolapse)                          |
| MRI                          | magnetic resonance imaging—a scan that creates detailed pictures of internal organs |
| MUSE                         | medicated urethral system for erection                                              |
| NK                           | natural killer cells                                                                |
| NMDA Receptor                | N-methyl-D-aspartame receptor                                                       |
| NO                           | nitric oxide                                                                        |
| NSAID                        | nonsteroidal anti-inflammatory drug                                                 |
| PAD                          | peripheral artery disease                                                           |
| PCOS                         | polycystic ovary syndrome                                                           |
| PDE5 Inhibitors              | phosphodiesterase type 5 inhibitors                                                 |
| PET Scan                     | positron emission tomography                                                        |
| PSA                          | prostate-specific antigen                                                           |
| PTCA                         | percutaneous transluminal coronary angioplasty                                      |
| PTSD                         | post-traumatic stress disorder                                                      |
| RAST                         | radioallergosorbent test                                                            |
| RCT                          | randomized controlled trial                                                         |
| RDA                          | recommended dietary allowance                                                       |
| REM                          | rapid eye movement                                                                  |
| RLS                          | restless legs syndrome                                                              |
| SAD                          | seasonal affective disorder                                                         |
| SERMs                        | selective estrogen receptor modulators                                              |
| SSRIs                        | selective serotonin reuptake inhibitors                                             |

---

---

| Primary term              | Alternate names                                                         |
|---------------------------|-------------------------------------------------------------------------|
| Steroids                  | corticosteroids                                                         |
| T Cell                    | mdash<br>T lymphocyte                                                   |
| TDD                       | telecommunications device for the deaf                                  |
| TENS                      | mdash<br>transcutaneous electrical nerve stimulation                    |
| TIA                       | mdash<br>transient ischemic attack                                      |
| tPA                       | tissue plasminogen activator                                            |
| TSH                       | thyroid-stimulating hormone                                             |
| VLDL                      | very-low-density lipoprotein                                            |
| VSD                       | ventricular septal defect                                               |
| ABO Blood Group System    | mdash<br>ABO Blood Group; ABO Blood Groups; ABO Blood Types; ABO System |
| Absolute Neutrophil Count | ANC                                                                     |
| Aarskog-Scott syndrome    | Aarskog Syndrome                                                        |
| Banti Syndrome            | Banti disease; Banti's syndrome                                         |
| Barber itch               | Barber's itch                                                           |
| Glycated Hemoglobin (A1C) | A1C; A1C test; Hemoglobin A1C; HbA1C                                    |

---



# British and US Medical Terms

This table lists common differences between United States and British medical terminology. Some terms are used in both countries, and local hospitals may use additional variations.

| United States term | British term      |
|--------------------|-------------------|
| Acetaminophen      | Paracetamol       |
| Anemia             | Anaemia           |
| Anesthesia         | Anaesthesia       |
| Anesthetic         | Anaesthetic       |
| Apnea              | Apnoea            |
| Behavior           | Behaviour         |
| Cesarean section   | Caesarean section |
| Celiac disease     | Coeliac disease   |
| Color              | Colour            |
| Diarrhea           | Diarrhoea         |
| Edema              | Oedema            |
| Epinephrine        | Adrenaline        |
| Esophagus          | Oesophagus        |
| Fetal              | Foetal            |
| Fetus              | Foetus            |
| Gynecology         | Gynaecology       |
| Hemoglobin         | Haemoglobin       |
| Hemolysis          | Haemolysis        |
| Hemorrhage         | Haemorrhage       |
| Labor              | Labour            |
| Leukocyte          | Leucocyte         |
| Maneuver           | Manoeuvre         |
| Orthopedics        | Orthopaedics      |
| Pediatric          | Paediatric        |
| Pediatrician       | Paediatrician     |
| Practice (verb)    | Practise (verb)   |
| Sulfur             | Sulphur           |
| Tumor              | Tumour            |

## Common vocabulary differences

| United States term    | British term        |
|-----------------------|---------------------|
| Pap test or Pap smear | Cervical smear test |

---

| United States term           | British term                                                    |
|------------------------------|-----------------------------------------------------------------|
| Emergency department (ED)    | Accident and emergency department (A&E)                         |
| Emergency room (ER)          | Accident and emergency department (A&E)                         |
| Family physician             | General practitioner (GP)                                       |
| Operating room (OR)          | Operating theatre                                               |
| Primary care physician (PCP) | General practitioner (GP)                                       |
| Skilled nursing facility     | Nursing home                                                    |
| Urgent care center           | Walk-in centre or minor injuries unit, depending on the service |

---



# Common Medical Abbreviations

| Abbreviation  | Full form                                                                        |
|---------------|----------------------------------------------------------------------------------|
| ABG           | Arterial blood gas                                                               |
| ACE           | Angiotensin-converting enzyme                                                    |
| ACE inhibitor | Angiotensin-converting enzyme inhibitor                                          |
| ACL           | Anterior cruciate ligament                                                       |
| ACS           | Acute coronary syndrome                                                          |
| ADHD          | Attention-deficit/hyperactivity disorder                                         |
| ADL           | Activity of daily living                                                         |
| AED           | Automated external defibrillator                                                 |
| AFib          | Atrial fibrillation                                                              |
| AIDS          | Acquired immunodeficiency syndrome                                               |
| AKI           | Acute kidney injury                                                              |
| ALP           | Alkaline phosphatase                                                             |
| ALT           | Alanine aminotransferase                                                         |
| AMI           | Acute myocardial infarction                                                      |
| ALS           | Amyotrophic lateral sclerosis; advanced life support, depending on context       |
| ANC           | Absolute neutrophil count                                                        |
| ANA           | Antinuclear antibody                                                             |
| ANP           | Atrial natriuretic peptide                                                       |
| APGAR         | Appearance, pulse, grimace, activity, and respiration                            |
| ARDS          | Acute respiratory distress syndrome                                              |
| AST           | Aspartate aminotransferase                                                       |
| ATN           | Acute tubular necrosis                                                           |
| AV            | Atrioventricular                                                                 |
| AVM           | Arteriovenous malformation                                                       |
| AVN           | Atrioventricular node                                                            |
| BID           | Twice a day; prescription shorthand that should be avoided when possible         |
| BP            | Blood pressure                                                                   |
| BMI           | Body mass index                                                                  |
| BMR           | Basal metabolic rate                                                             |
| BPH           | Benign prostatic hyperplasia                                                     |
| BRCA          | Breast cancer susceptibility gene; commonly referring to the BRCA1 or BRCA2 gene |
| BUN           | Blood urea nitrogen                                                              |
| CABG          | Coronary artery bypass graft                                                     |
| CA            | Cancer antigen or calcium, depending on context                                  |
| CA-125        | Cancer antigen 125                                                               |

---

| Abbreviation | Full form                                                                   |
|--------------|-----------------------------------------------------------------------------|
| CAD          | Coronary artery disease                                                     |
| CAT          | Computed tomography; an older term for computerized axial tomography        |
| CBC          | Complete blood count                                                        |
| CBT          | Cognitive behavioural therapy                                               |
| CDC          | Centers for Disease Control and Prevention                                  |
| CEA          | Carcinoembryonic antigen                                                    |
| CHF          | Congestive heart failure                                                    |
| CHD          | Congenital heart disease                                                    |
| CIN          | Cervical intraepithelial neoplasia                                          |
| CJD          | Creutzfeldt–Jakob disease                                                   |
| CK           | Creatine kinase                                                             |
| CMV          | Cytomegalovirus                                                             |
| CNS          | Central nervous system                                                      |
| COPD         | Chronic obstructive pulmonary disease                                       |
| CPAP         | Continuous positive airway pressure                                         |
| CPK          | Creatine phosphokinase; an older name for creatine kinase                   |
| CPR          | Cardiopulmonary resuscitation                                               |
| CRF          | Chronic renal failure; chronic kidney disease is preferred in current usage |
| CRP          | C-reactive protein                                                          |
| CSF          | Cerebrospinal fluid                                                         |
| CT           | Computed tomography                                                         |
| CVS          | Chorionic villus sampling                                                   |
| CVA          | Cerebrovascular accident                                                    |
| CXR          | Chest radiograph                                                            |
| D&C          | Dilatation and curettage                                                    |
| DKA          | Diabetic ketoacidosis                                                       |
| DJD          | Degenerative joint disease                                                  |
| DM           | Diabetes mellitus                                                           |
| DNA          | Deoxyribonucleic acid                                                       |
| DTP          | Diphtheria, tetanus, and pertussis                                          |
| DVT          | Deep-vein thrombosis                                                        |
| DX           | Diagnosis                                                                   |
| ECG          | Electrocardiogram                                                           |
| EKG          | Electrocardiogram                                                           |
| ECHO         | Echocardiogram                                                              |
| EEG          | Electroencephalogram                                                        |
| eGFR         | Estimated glomerular filtration rate                                        |
| EMG          | Electromyography                                                            |
| ENT          | Ear, nose, and throat                                                       |
| ERCP         | Endoscopic retrograde cholangiopancreatography                              |
| ESR          | Erythrocyte sedimentation rate                                              |
| ESRD         | End-stage renal disease                                                     |
| FBC          | Full blood count                                                            |
| FEV1         | Forced expiratory volume in the first second                                |
| FSH          | Follicle-stimulating hormone                                                |
| GABHS        | Group A beta-haemolytic streptococcus                                       |
| GERD         | Gastroesophageal reflux disease                                             |
| GFR          | Glomerular filtration rate                                                  |
| GI           | Gastrointestinal                                                            |
| GIST         | Gastrointestinal stromal tumour                                             |
| GU           | Genitourinary                                                               |
| HbA1c        | Glycated haemoglobin                                                        |

---

---

| Abbreviation | Full form                                                                                       |
|--------------|-------------------------------------------------------------------------------------------------|
| HDL          | High-density lipoprotein                                                                        |
| HAV          | Hepatitis A virus                                                                               |
| HBV          | Hepatitis B virus                                                                               |
| HCT          | Hematocrit                                                                                      |
| HCV          | Hepatitis C virus                                                                               |
| HGB          | Hemoglobin                                                                                      |
| HIV          | Human immunodeficiency virus                                                                    |
| HLA          | Human leukocyte antigen                                                                         |
| HPV          | Human papillomavirus                                                                            |
| HRT          | Hormone replacement therapy                                                                     |
| HTN          | Hypertension                                                                                    |
| IBD          | Inflammatory bowel disease                                                                      |
| IBS          | Irritable bowel syndrome                                                                        |
| ICD          | Implantable cardioverter-defibrillator                                                          |
| ICU          | Intensive care unit                                                                             |
| IDDM         | Insulin-dependent diabetes mellitus; type 1 diabetes mellitus is preferred in current usage     |
| IM           | Intramuscular                                                                                   |
| IU           | International unit                                                                              |
| IUD          | Intrauterine device                                                                             |
| IV           | Intravenous                                                                                     |
| IVP          | Intravenous pyelography or intravenous pyelogram                                                |
| LDL          | Low-density lipoprotein                                                                         |
| LH           | Luteinizing hormone                                                                             |
| LFT          | Liver function test                                                                             |
| MI           | Myocardial infarction                                                                           |
| MMR          | Measles, mumps, and rubella                                                                     |
| MRI          | Magnetic resonance imaging                                                                      |
| MRSA         | Methicillin-resistant <i>Staphylococcus aureus</i>                                              |
| MS           | Multiple sclerosis; may have other meanings depending on context                                |
| NG           | Nasogastric                                                                                     |
| NIDDM        | Non-insulin-dependent diabetes mellitus; type 2 diabetes mellitus is preferred in current usage |
| NKDA         | No known drug allergies                                                                         |
| NSAID        | Nonsteroidal anti-inflammatory drug                                                             |
| OCD          | Obsessive-compulsive disorder                                                                   |
| PAD          | Peripheral arterial disease                                                                     |
| PAP          | Papanicolaou                                                                                    |
| PCOS         | Polycystic ovary syndrome                                                                       |
| PCR          | Polymerase chain reaction                                                                       |
| PE           | Pulmonary embolism                                                                              |
| PET          | Positron emission tomography                                                                    |
| PFT          | Pulmonary function test                                                                         |
| PID          | Pelvic inflammatory disease                                                                     |
| PMS          | Premenstrual syndrome                                                                           |
| PPD          | Purified protein derivative                                                                     |
| PRN          | As needed (from the Latin <i>pro re nata</i> )                                                  |
| PSA          | Prostate-specific antigen                                                                       |
| PT           | Prothrombin time                                                                                |
| PTCA         | Percutaneous transluminal coronary angioplasty                                                  |
| PTH          | Parathyroid hormone                                                                             |
| PTT          | Partial thromboplastin time                                                                     |

---

---

| Abbreviation     | Full form                                                                                                                  |
|------------------|----------------------------------------------------------------------------------------------------------------------------|
| PUD              | Peptic ulcer disease                                                                                                       |
| PVC              | Premature ventricular contraction                                                                                          |
| QID              | Four times a day; prescription shorthand that should be avoided when possible                                              |
| RA               | Rheumatoid arthritis; may have other meanings depending on context                                                         |
| RAST             | Radioallergosorbent test                                                                                                   |
| RBC              | Red blood cell                                                                                                             |
| RCT              | Randomized controlled trial                                                                                                |
| REM              | Rapid eye movement                                                                                                         |
| RLS              | Restless legs syndrome                                                                                                     |
| RNA              | Ribonucleic acid                                                                                                           |
| RSV              | Respiratory syncytial virus                                                                                                |
| Rx               | Prescription or treatment, depending on context                                                                            |
| SaO <sub>2</sub> | Arterial oxygen saturation                                                                                                 |
| SARS-CoV-2       | Severe acute respiratory syndrome coronavirus 2                                                                            |
| SIDS             | Sudden infant death syndrome                                                                                               |
| SLE              | Systemic lupus erythematosus                                                                                               |
| SOB              | Shortness of breath                                                                                                        |
| SSRI             | Selective serotonin reuptake inhibitor                                                                                     |
| STD              | Sexually transmitted disease; sexually transmitted infection is preferred in current usage                                 |
| STI              | Sexually transmitted infection                                                                                             |
| T <sub>3</sub>   | Triiodothyronine                                                                                                           |
| T <sub>4</sub>   | Thyroxine; commonly referring to tetraiodothyronine                                                                        |
| TAH              | Total abdominal hysterectomy                                                                                               |
| TB               | Tuberculosis                                                                                                               |
| TIA              | Transient ischemic attack                                                                                                  |
| TIBC             | Total iron-binding capacity                                                                                                |
| TID              | Three times a day; prescription shorthand that should be avoided when possible                                             |
| TMJ              | Temporomandibular joint                                                                                                    |
| TORCH            | Toxoplasmosis, other infections, rubella, cytomegalovirus, and herpes simplex; a group of congenital-infection evaluations |
| TSH              | Thyroid-stimulating hormone                                                                                                |
| TURP             | Transurethral resection of the prostate                                                                                    |
| URI              | Upper respiratory infection                                                                                                |
| UTI              | Urinary tract infection                                                                                                    |
| VSD              | Ventricular septal defect                                                                                                  |
| WBC              | White blood cell                                                                                                           |
| WDWN             | Well-developed, well-nourished                                                                                             |
| WNL              | Within normal limits; a finding is within the expected reference range or clinical appearance                              |
| WHO              | World Health Organization                                                                                                  |
| XRT              | Radiotherapy                                                                                                               |

---

**A.C.** A prescription abbreviation meaning “before meals,” from the Latin *ante cibum*.

**ALA** An abbreviation for alpha-linolenic acid, an essential omega-3 fatty acid, or aminolevulinic acid, a heme-pathway compound used topically in some photodynamic treatments; context determines the meaning.

**AP** An abbreviation commonly meaning anteroposterior, or directed from front to back; in some contexts it may have another medical expansion.

**ATM** An abbreviation that may mean the temporomandibular joint or ataxia-telangiectasia mutated, a protein kinase involved in DNA-damage repair; context determines the meaning.

- 
- BF** Physician's shorthand for Black female; demographic abbreviations should be used cautiously and replaced with precise, patient-preferred language when possible.
- BPD** An abbreviation commonly meaning bronchopulmonary dysplasia, chronic lung disease after premature birth; it can also mean borderline personality disorder in psychiatric contexts.
- CA** A medical abbreviation that may mean cancer, carcinoma, or calcium; the intended meaning depends on the surrounding clinical context.
- CKD** Abbreviation for Chronic Kidney Disease: a progressive loss of kidney function over months to years, defined by a glomerular filtration rate (GFR) less than 60 mL/min/1.73m<sup>2</sup> or markers of kidney damage (albuminuria, haematuria, structural abnormalities) persisting for at least 3 months.
- CLA** An abbreviation that commonly means conjugated linoleic acid, a group of fatty-acid isomers; in other medical contexts it may have a different expansion.
- CTG** Abbreviation for cardiotocography; continuous electronic monitoring of fetal heart rate and uterine contractions during labour.
- CVA** An abbreviation for cerebrovascular accident, meaning a stroke caused by interruption of blood flow to the brain or bleeding within the brain. It may cause sudden neurologic deficits.
- CVS** An abbreviation that may mean cardiovascular system or chorionic-villous sampling, a prenatal diagnostic procedure; context determines the meaning.
- DDAVP** The abbreviation for desmopressin, a synthetic vasopressin-like medicine used for some bleeding disorders, central diabetes insipidus, and nighttime urination.
- DES** An abbreviation that may mean diethylstilbestrol, a synthetic estrogen, or drug-eluting stent; diethylstilbestrol exposure during pregnancy is associated with later reproductive and cancer risks.
- DHHS** The abbreviation for the U.S. Department of Health and Human Services.
- DNR** An abbreviation for do-not-resuscitate, indicating that cardiopulmonary resuscitation should not be attempted if breathing or heartbeat stops.
- DNS** An abbreviation that may mean deviated nasal septum or dextrose in normal saline; the intended expansion depends on the clinical context.
- DPT** An older abbreviation for the combined diphtheria, pertussis, and tetanus vaccine; current formulations include DTaP and Tdap.
- DT** An abbreviation that may mean delirium tremens or diphtheria–tetanus vaccine; the intended meaning depends on context.
- DTC** An abbreviation that may mean differentiated thyroid carcinoma or direct-to-consumer, such as direct-to-consumer genetic testing; the intended meaning depends on context.
- DTP** An older abbreviation for diphtheria, tetanus, and pertussis vaccine; current childhood formulations are generally called DTaP.
- DV** An abbreviation that may mean domestic violence, ductus venosus, or another context-specific term; the full expression should be specified.
- DYS** An abbreviation or word root related to dysplasia, dystonia, dyslexia, or another condition beginning with dys-; the intended meaning depends on context.
- EC** An abbreviation that may mean enteric-coated, emergency contraception, or endothelial cell; the intended meaning depends on context.
- EFT** An abbreviation whose medical meaning depends on context, such as endocrine-function testing or emotional freedom technique; the intended expansion should be specified.
- ENT** Abbreviation for ear, nose, and throat; it commonly refers to the otolaryngology specialty or an otolaryngologist.
-

---

**EOS** Commonly an abbreviation for eosinophils or an eosinophil count in laboratory reports; the intended meaning should be confirmed from the report's context.

**EPA** Abbreviation for eicosapentaenoic acid, an omega-3 polyunsaturated fatty acid found in oily fish and fish-oil supplements.

**EPR** Abbreviation for electrophrenic respiration, also called diaphragm pacing, in which electrical stimulation helps activate the diaphragm in selected patients with respiratory muscle paralysis.

**EPT** Abbreviation for estrogen/progestin therapy, treatment using both an estrogen and a progestin, commonly discussed in menopausal hormone therapy when the uterus is present.

**ER** Abbreviation for emergency room or emergency department, the hospital service that evaluates and treats urgent illness and injury.

**FCC** In medicine, an abbreviation for familial colorectal cancer; in other contexts it may have nonmedical meanings.

**FFI** Abbreviation for fatal familial insomnia, a rare inherited prion disease causing progressively severe insomnia, autonomic dysfunction, and neurologic decline.

**FISH** Abbreviation for fluorescence in situ hybridization, a test that uses fluorescent DNA probes to detect specific genes, chromosomes, or chromosomal rearrangements in cells.

**FLAP** Abbreviation for 5-lipoxygenase-activating protein, a protein encoded by *ALOX5AP* that helps produce leukotrienes involved in inflammation.

**GALT** Abbreviation for gut-associated lymphoid tissue, immune tissue in the gastrointestinal tract that samples intestinal contents and helps defend against pathogens.

**GBV-C** Abbreviation for GB virus C, also called human pegivirus, a blood-borne virus that can persist in people but is not established as a cause of chronic hepatitis.

**GD** In the RxList medical-dictionary context, abbreviation for Gaucher disease, an inherited lysosomal storage disorder caused by deficient glucocerebrosidase.

**GI** Abbreviation for gastrointestinal, relating to the digestive tract from the mouth through the anus and its associated organs.

**GPO** Abbreviation for group purchasing organization, an organization that combines buyers' purchasing power to negotiate prices and contracts for medicines, supplies, or services.

**HAART** Abbreviation for highly active antiretroviral therapy, an older term for combination antiretroviral treatment used to suppress HIV replication and preserve immune function.

**HAPE** Abbreviation for high-altitude pulmonary edema, a life-threatening accumulation of fluid in the lungs after rapid ascent; immediate descent and supplemental oxygen are urgent treatments.

**HAV** Abbreviation for hepatitis A virus, an RNA virus spread mainly by the fecal-oral route that causes acute hepatitis but usually does not cause chronic infection.

**HBV** Abbreviation for hepatitis B virus, a DNA virus transmitted through infected blood or body fluids that can cause acute or chronic hepatitis and cirrhosis or liver cancer.

**HEPA** Abbreviation for high-efficiency particulate air, describing a filter designed to remove a very high proportion of airborne particles of specified sizes from an air stream.

**HGD** Commonly an abbreviation for high-grade dysplasia, a marked abnormality of cell development that may be a precursor to invasive cancer but is not itself invasive cancer.

**HNP** Abbreviation for herniated nucleus pulposus, in which the inner material of an intervertebral disk protrudes through the outer ring and may compress a nerve root.

**HPS** Common medical abbreviations include hantavirus pulmonary syndrome, a severe respiratory illness, and Hermansky-Pudlak syndrome, an inherited disorder with oculocutaneous albinism and platelet-storage abnormalities; context determines the meaning.

---



- 
- HS** Commonly an abbreviation for hereditary spherocytosis, an inherited hemolytic anemia; it can also mean half-strength, heart sounds, hamstrings, heparan sulfate, hernia sac, or house surgeon, so context is essential.
- HSA** Abbreviation that may mean a US health savings account or human serum albumin, the main plasma protein that helps maintain oncotic pressure; context determines the meaning.
- ICU** Abbreviation for intensive care unit, a hospital unit providing continuous monitoring and specialized support for critically ill patients.
- ID** A context-dependent abbreviation that may mean infectious disease, infectious diseases specialist, intellectual disability, or identification; the surrounding clinical text determines the meaning.
- IFA** Abbreviation for indirect fluorescent antibody, a laboratory method that detects antibodies or antigens using fluorescent labels, commonly in serologic testing.
- IHS** Abbreviation for the Indian Health Service, a US federal agency providing health services to American Indians and Alaska Native people and communities.
- IPT** Abbreviation for interpersonal therapy, a structured psychotherapy that focuses on relationships, grief, role transitions, and interpersonal difficulties.
- IPV** Abbreviation for inactivated polio vaccine, an injectable vaccine containing killed poliovirus and used to prevent poliomyelitis.
- IQ** Abbreviation for intelligence quotient, a standardized score from cognitive testing; it describes performance on specified tests and is not a complete measure of a person's abilities or worth.
- ITC** Abbreviation for in-the-canal, describing a hearing aid or other device positioned within the ear canal.
- ITN** Abbreviation for insecticide-treated bednet, a net treated with insecticide to reduce mosquito bites and malaria transmission.
- IVC** Commonly abbreviation for inferior vena cava, the large vein returning blood from the lower body to the heart; it can also mean intravenous cholangiogram, an X-ray study of the biliary system.
- JIP** Abbreviation for juvenile intestinal polyposis, an inherited condition in which children develop multiple hamartomatous polyps in the gastrointestinal tract and may have increased cancer risk.
- KB** In medical usage, an abbreviation for keratoderma blennorrhagicum or kilobase of DNA; lower-case *kb* is preferred for kilobase, so context is important.
- KUB** Abbreviation for kidney, ureter, and bladder, commonly referring to a plain abdominal X-ray used to assess urinary tract calculi, bowel gas patterns, or other abdominal findings.
- L** Abbreviation for liter, a metric unit of volume equal to 1,000 milliliters.
- LAVH** Abbreviation for laparoscopically assisted vaginal hysterectomy, removal of the uterus through the vagina with laparoscopic assistance.
- LL** Commonly an abbreviation for lower lid of the eye or lower limit; the intended meaning depends on the clinical context.
- LLL** Abbreviation for the left lower lobe of the lung.
- LLQ** Abbreviation for left lower quadrant, usually referring to the lower-left quarter of the abdomen.
- LMP** Abbreviation for last menstrual period, the first day of the most recent menstrual bleeding, used in estimating gestational age when the date is reliable.
- LPA** LPA abbreviates Little People of America and may also refer to the left pulmonary artery.
- LUL** Abbreviation for the left upper lobe of the lung.
- LUQ** Abbreviation for the left upper quadrant of the abdomen.
- LVF** Abbreviation for left ventricular failure, meaning failure of the left side of the heart; it may also mean left ventricular function, depending on context.
-

- 
- MAOI** Abbreviation for monoamine oxidase inhibitor, a drug that inhibits monoamine oxidase and is used in selected treatment of depression and other disorders.
- MAP** Common abbreviation for *Mycobacterium avium* subspecies *paratuberculosis*, an organism of the *M. avium* complex; the abbreviation can have other meanings in different clinical contexts.
- MCD** An abbreviation with several medical meanings, including minimal change disease, mean corpuscular diameter, and medullary cystic disease; the intended meaning depends on context.
- MCP** Membrane cofactor protein, also called CD46, is a cell-surface protein that regulates complement activation and helps protect host cells from complement-mediated injury; the abbreviation can have other meanings in different contexts.
- MDI** An abbreviation that may mean metered-dose inhaler, multiple daily injections, or mental developmental index; the intended meaning depends on context.
- MEDVAC** Abbreviation for medical evacuation: the transport of ill or injured people to a medical facility or to a location providing a higher level of care.
- MI** Abbreviation for myocardial infarction, or heart attack: death of heart-muscle tissue caused by prolonged inadequate blood flow, usually from acute coronary artery blockage.
- MM** An abbreviation that may mean meningomyelocele, malignant melanoma, or multiple myeloma; in lowercase, *mm* commonly means millimeter. The intended meaning depends on context.
- MP** Abbreviation for mercaptopurine, a purine-antimetabolite drug, or for molecular pathology; the intended meaning depends on context.
- MPS** Abbreviation for mucopolysaccharidosis, a group of inherited lysosomal storage disorders caused by deficient breakdown of glycosaminoglycans.
- MSG** Abbreviation for monosodium glutamate, the sodium salt of glutamic acid used as a flavor enhancer and contributing umami taste.
- MTX** Abbreviation for methotrexate, a folate-antagonist drug used in selected cancers, autoimmune diseases, and other conditions.
- MVP** Abbreviation for mitral valve prolapse, in which one or both mitral-valve leaflets bow into the left atrium during ventricular systole; it is often mild but can sometimes cause mitral regurgitation or arrhythmias.
- MYH** MYH is a gene-family abbreviation involving myosin or DNA-repair proteins; the specific gene and variant determine the clinical meaning.
- ND** ND is a context-dependent abbreviation that may mean naturopathic doctor, neurodevelopmental, or another term; the full meaning should be specified.
- NEP** NEP is a context-dependent abbreviation that may mean neuroendocrine pancreatic tumor, nasoendotracheal procedure, or another clinical term.
- NF** NF is a context-dependent abbreviation, including neurofibromatosis, a group of genetic disorders causing nerve-sheath tumors and other features.
- NG** NG commonly means nasogastric, referring to a tube passed through the nose into the stomach, but the abbreviation has other uses.
- NS** NS is a context-dependent abbreviation, commonly normal saline, a sterile sodium-chloride solution, or not significant; the full term should be stated.
- OB** OB is an abbreviation commonly meaning obstetrics, the care of pregnancy, labor, birth, and the postpartum period, but it may have other meanings.
- OBS** OBS is a context-dependent abbreviation that may mean observation, obstetric service, or organic brain syndrome; the intended term should be written out.

- 
- OCG** OCG is a context-dependent abbreviation that may refer to oral contraceptive guidance, oculocardiac testing, or another clinical term; the full term should be specified.
- OMS** An abbreviation for oral and maxillofacial surgery; the intended meaning depends on the clinical context.
- PAP** PAP is an abbreviation with context-dependent meanings, including Papanicolaou testing, pulmonary alveolar proteinosis, or prostatic acid phosphatase; the full term should be specified.
- PAS** PAS is an abbreviation that may mean periodic-acid–Schiff staining, para-aminosalicylic acid, or pulmonary-artery systolic pressure; context determines the meaning.
- PAT** PAT is a context-dependent abbreviation that may refer to paroxysmal atrial tachycardia, a rapid rhythm beginning in the atria, or another clinical term.
- PCA** PCA is a context-dependent abbreviation, commonly meaning patient-controlled analgesia, in which a patient activates prescribed doses from an infusion device.
- PCT** PCT is a context-dependent abbreviation that may mean prolactin, porphyria cutanea tarda, or patient-care technician; the full term should be specified.
- PDR** PDR commonly means Physicians' Desk Reference, a drug-information reference, but the abbreviation has other uses and should be interpreted in context.
- PEP** PEP is a context-dependent abbreviation, including post-exposure prophylaxis, a preventive medicine course after a possible infection exposure.
- PHS** PHS is an abbreviation whose meaning depends on context, including public health service or primary hyperhidrosis; the full term should be specified.
- PI** PI is a context-dependent abbreviation, including principal investigator, pulmonary insufficiency, or protease inhibitor; the intended meaning must be specified.
- PLP** PLP commonly means pyridoxal 5'-phosphate, the active coenzyme form of vitamin B6 required by many enzymes, but the abbreviation has other uses.
- PMD** PMD is a context-dependent abbreviation, including Pelizaeus–Merzbacher disease, a rare inherited disorder of central myelination, or a professional medical designation.
- PON** PON is a context-dependent abbreviation that may refer to postoperative nausea, paraoxonase, or another clinical term; the full meaning should be stated.
- PTU** PTU is the abbreviation for propylthiouracil, an antithyroid medicine used selectively because of liver toxicity and other safety considerations.
- PV** PV is a context-dependent abbreviation, including polycythemia vera, plasma volume, or peripheral vision; the full term should be specified.
- PVB** PVB is a context-dependent abbreviation that may mean paravertebral block, a regional anesthetic technique, or another medical term.
- PX** PX is a context-dependent abbreviation that may mean prognosis, patient, prescription, or a genetic marker; clinical documentation should spell out the intended term.
- Q.2H** A prescription abbreviation meaning “every 2 hours.” Medication instructions should use clear wording because abbreviation misinterpretation can cause dosing errors.
- Q.3H.** A prescription abbreviation meaning “every 3 hours.” Medication instructions should use clear wording because abbreviation misinterpretation can cause dosing errors.
- Q.S.** A prescription abbreviation from the Latin “quantum sufficit,” meaning “quantity sufficient” to make the required amount or concentration. The exact quantity must be specified by the formulation or prescription context.
- R/O** R/O is an abbreviation for rule out, indicating that a condition is being considered but has not been confirmed; clear documentation should state the actual diagnostic question.
-

---

**RAD** Acronym whose medical meaning depends on context, commonly radiation absorbed dose or right axis deviation on an ECG.

**RBC** RBC is the abbreviation for red blood cell, the cell that transports oxygen with hemoglobin and carries carbon dioxide back toward the lungs.

**RF** RF is a context-dependent abbreviation, including rheumatoid factor, radiofrequency, or risk factor; the complete term should be stated.

**RLL** RLL is the abbreviation for right lower lobe, usually referring to the lower lobe of the right lung.

**RLQ** RLQ is the abbreviation for right lower quadrant of the abdomen, containing structures including the appendix, cecum, and right adnexa.

**RML** RML is the abbreviation for right middle lobe, the middle lobe of the right lung.

**RP** RP is a context-dependent abbreviation, commonly retinitis pigmentosa, an inherited retinal degeneration causing night blindness and progressive visual-field loss.

**RUQ** RUQ is the abbreviation for right upper quadrant of the abdomen, containing the liver, gallbladder, duodenum, and portions of bowel and kidney.

**SGOT** SGOT is the former abbreviation for AST, an enzyme measured in blood that can rise with injury to liver, heart, muscle, or other tissues.

**SGPT** SGPT is the former abbreviation for ALT, a liver-associated enzyme that rises in many forms of hepatocellular injury.

**SHP** SHP is a context-dependent abbreviation that may refer to a healthcare professional, a gene or protein, or another clinical term; the full meaning should be specified.

**SLV** SLV is a context-dependent abbreviation that may mean single-lung ventilation, serum lithium value, or another term; the full meaning should be specified.

**STD(S)** Abbreviation for sexually transmitted disease(s), now often called sexually transmitted infections.

**STS** STS is a context-dependent abbreviation that may mean soft-tissue sarcoma, sodium thiosulfate, or another clinical term; the full term should be stated.

**TMR** TMR is a context-dependent abbreviation, including targeted muscle reinnervation, a surgical technique used to improve control of a prosthesis and reduce amputation pain.

**TPA/TPA** Abbreviation for tissue plasminogen activator.

**TSD** TSD is a context-dependent abbreviation that may mean total sleep deprivation, traumatic stress disorder, or another clinical term; the full term should be specified.

**TSE** TSE is a context-dependent abbreviation, including transmissible spongiform encephalopathy, the prion-disease group that includes Creutzfeldt-Jakob disease.

**UL** UL is a context-dependent abbreviation that may mean upper limit, ulnar or ulna-related, or another clinical term; the full meaning should be specified.

**URG** URG is a context-dependent abbreviation that may refer to urinary urgency or an urgent clinical priority; documentation should state the full term.

**URTI** Abbreviation for upper respiratory tract infection, commonly involving the nose, sinuses, pharynx, or larynx and often caused by viruses.

**VD** Abbreviation for venereal disease, an older term for a sexually transmitted infection.

**WBS** Abbreviation for Williams–Beuren syndrome, a genetic condition associated with characteristic facial features, developmental differences, and cardiovascular disease. In other clinical contexts, WBS may have a different meaning, so the full term should be specified.

# Modern Terms

| Old or obsolete term                | Modern or preferred term                                                                          |
|-------------------------------------|---------------------------------------------------------------------------------------------------|
| Acute renal failure                 | Acute kidney injury                                                                               |
| Adult-onset diabetes mellitus       | Type 2 diabetes mellitus                                                                          |
| Ague                                | Malaria; a historical term for intermittent fever, usually associated with malaria                |
| Apoplexy                            | Stroke or intracerebral haemorrhage, depending on the historical context                          |
| Arteriosclerosis obliterans         | Peripheral arterial disease                                                                       |
| Bright's disease                    | Glomerular kidney disease; the specific modern diagnosis should be stated when known              |
| Chronic renal failure               | Chronic kidney disease                                                                            |
| Congestive heart failure            | Heart failure                                                                                     |
| Consumption                         | Tuberculosis, especially pulmonary tuberculosis                                                   |
| Crippling arthritis                 | Disabling arthritis; describe the functional limitation specifically                              |
| Diabetic retinopathy, background    | Mild or nonproliferative diabetic retinopathy, with severity stated                               |
| Dropsy                              | Oedema or fluid accumulation, with the affected site specified                                    |
| Eclampsia toxemia                   | Preeclampsia or eclampsia, depending on the clinical condition                                    |
| Essential hypertension              | Primary hypertension                                                                              |
| Fallopian tube                      | Uterine tube; fallopian tube remains an accepted anatomical term                                  |
| Female hysteria                     | No single modern diagnosis; reassess using the presenting symptoms and an appropriate diagnosis   |
| Gastric ulcer disease               | Peptic ulcer disease, with the ulcer site specified                                               |
| Heart attack                        | Myocardial infarction in formal medical usage                                                     |
| Herpes zoster                       | Shingles in common language; herpes zoster remains the preferred medical term                     |
| Hyperlipaemia                       | Hyperlipidaemia                                                                                   |
| Idiopathic thrombocytopenic purpura | Immune thrombocytopenia                                                                           |
| Imbecile                            | Intellectual disability; the historical term is offensive and must not be used clinically         |
| Insulin-dependent diabetes mellitus | Type 1 diabetes mellitus                                                                          |
| Junkie's abscess                    | Injection-related abscess or skin and soft-tissue infection                                       |
| Leprosy                             | Hansen disease; leprosy remains a recognized medical term, but person-first language is preferred |

---

| Old or obsolete term                    | Modern or preferred term                                                             |
|-----------------------------------------|--------------------------------------------------------------------------------------|
| Manic depression                        | Bipolar disorder                                                                     |
| Mongolism                               | Down syndrome; the historical term is unacceptable and must not be used              |
| Multiple personality disorder           | Dissociative identity disorder                                                       |
| Non-insulin-dependent diabetes mellitus | Type 2 diabetes mellitus                                                             |
| Old age dementia                        | Dementia or major neurocognitive disorder, with the cause specified when known       |
| Palsy                                   | Paralysis or the specific neurological disorder, depending on the context            |
| Paralytic stroke                        | Stroke with paralysis; specify the neurological deficit and stroke type              |
| Phthisis                                | Pulmonary tuberculosis; historically, progressive wasting disease                    |
| Senile dementia                         | Dementia or major neurocognitive disorder; age alone should not define the diagnosis |
| Senility                                | Cognitive impairment, dementia, or another specific diagnosis, after assessment      |
| Spastic colon                           | Irritable bowel syndrome                                                             |
| Stomach flu                             | Acute gastroenteritis                                                                |
| Sunstroke                               | Heatstroke                                                                           |
| Terminal illness                        | Advanced or life-limiting illness, with the diagnosis and prognosis specified        |
| Toxemia of pregnancy                    | Preeclampsia or eclampsia                                                            |
| Venereal disease                        | Sexually transmitted infection                                                       |
| Wassermann reaction                     | A modern syphilis serologic test, such as a nontreponemal or treponemal test         |
| Water on the brain                      | Hydrocephalus                                                                        |

---



# Medical Specialties

| Medical specialty           | Principal area of practice                                                   |
|-----------------------------|------------------------------------------------------------------------------|
| Allergy and Immunology      | Allergic disease, asthma, immune deficiency, and immune-mediated disorders   |
| Anesthesiology              | Perioperative care, anesthesia, critical care, and pain medicine             |
| Cardiology                  | Diseases of the heart and circulatory system                                 |
| Cardiothoracic Surgery      | Surgical treatment of diseases of the heart, lungs, and chest                |
| Clinical Genetics           | Diagnosis, counseling, and management of inherited and genomic disorders     |
| Clinical Pharmacology       | Effects, uses, interactions, and safe clinical use of medicines              |
| Colon and Rectal Surgery    | Surgical treatment of diseases of the colon, rectum, and anus                |
| Critical Care Medicine      | Assessment and treatment of life-threatening organ dysfunction               |
| Dermatology                 | Diseases of the skin, hair, nails, and mucous membranes                      |
| Emergency Medicine          | Immediate assessment and treatment of acute illness and injury               |
| Endocrinology               | Hormones, metabolism, diabetes, and endocrine glands                         |
| Family Medicine             | Comprehensive continuing care for individuals and families of all ages       |
| Forensic Medicine           | Application of medical knowledge to legal and medicolegal questions          |
| Gastroenterology            | Diseases of the digestive tract, liver, gallbladder, and pancreas            |
| General Practice            | First-contact and continuing medical care across a broad range of conditions |
| General Surgery             | Operative treatment of diseases involving multiple body systems              |
| Geriatric Medicine          | Health care of older adults, including multimorbidity and functional decline |
| Hematology                  | Blood cells, blood-forming organs, clotting, and blood disorders             |
| Hepatology                  | Diseases of the liver and biliary system                                     |
| Hospital Medicine           | Comprehensive medical care of hospitalized adults                            |
| Infectious Disease Medicine | Diagnosis, treatment, and prevention of infections                           |
| Internal Medicine           | Prevention, diagnosis, and treatment of adult medical disorders              |
| Interventional Cardiology   | Catheter-based diagnosis and treatment of cardiovascular disease             |
| Interventional Radiology    | Image-guided minimally invasive diagnosis and treatment                      |
| Medical Oncology            | Systemic treatment and supportive care for cancer                            |
| Neonatology                 | Care of newborn infants, especially premature or critically ill infants      |
| Nephrology                  | Diseases of the kidneys, fluid balance, electrolytes, and hypertension       |
| Neurology                   | Diseases of the brain, spinal cord, peripheral nerves, and muscles           |

---

| Medical specialty                          | Principal area of practice                                                          |
|--------------------------------------------|-------------------------------------------------------------------------------------|
| Neurosurgery                               | Surgical treatment of disorders of the nervous system                               |
| Nuclear Medicine                           | Diagnosis and treatment using radioactive substances and molecular imaging          |
| Obstetrics and Gynecology                  | Pregnancy, childbirth, and diseases of the female reproductive system               |
| Occupational Medicine                      | Health, safety, and disease prevention in the workplace                             |
| Ophthalmology                              | Medical and surgical care of the eyes and visual system                             |
| Oral and Maxillofacial Surgery             | Surgical treatment of diseases and injuries of the mouth, jaws, face, and neck      |
| Orthopedic Surgery                         | Surgical and nonsurgical treatment of the musculoskeletal system                    |
| Otolaryngology–Head and Neck Surgery       | Diseases and surgery of the ear, nose, throat, and head and neck                    |
| Pain Medicine                              | Assessment and treatment of acute, chronic, and cancer-related pain                 |
| Palliative Medicine                        | Relief of symptoms and support for people with serious or life-limiting illness     |
| Pathology                                  | Laboratory examination of tissues, cells, blood, and body fluids                    |
| Pediatric Surgery                          | Surgical care of infants, children, and adolescents                                 |
| Pediatrics                                 | Medical care of infants, children, and adolescents                                  |
| Physical Medicine and Rehabilitation       | Restoration of function and independence after illness or injury                    |
| Plastic and Reconstructive Surgery         | Reconstruction, restoration, and alteration of body tissues                         |
| Preventive Medicine                        | Prevention of disease and promotion of health in individuals and populations        |
| Psychiatry                                 | Diagnosis and treatment of mental, emotional, and behavioral disorders              |
| Public Health Medicine                     | Protection and improvement of population health                                     |
| Pulmonary Medicine                         | Diseases of the lungs and respiratory system                                        |
| Radiation Oncology                         | Use of ionizing radiation to treat cancer                                           |
| Radiology                                  | Diagnostic and image-guided evaluation of disease                                   |
| Reproductive Endocrinology and Infertility | Hormonal and reproductive disorders and fertility care                              |
| Rheumatology                               | Autoimmune, inflammatory, and musculoskeletal disorders                             |
| Sleep Medicine                             | Sleep disorders and their effects on health                                         |
| Sports Medicine                            | Prevention, diagnosis, treatment, and rehabilitation of activity-related conditions |
| Thoracic Surgery                           | Surgery of the chest, lungs, esophagus, and mediastinum                             |
| Transfusion Medicine                       | Collection, testing, processing, and clinical use of blood components               |
| Trauma Surgery                             | Surgical management of serious and complex injuries                                 |
| Urology                                    | Diseases of the urinary tract and male reproductive system                          |
| Vascular Surgery                           | Surgical and endovascular treatment of blood-vessel disease                         |

---

# Human Organs

This reference maps individual organ headwords to related entries in the dictionary. Each headword is followed by a semicolon-separated list of related dictionary terms.

## **Adrenal Gland**

Adrenal Gland; Addison's Disease; Cushing's Syndrome; Pheochromocytoma; Adrenaline.

## **Appendix**

Appendix; Appendicitis; Appendectomy.

## **Bladder**

Bladder; Cystitis; Urinary Retention; Bladder Cancer.

## **Brain**

Brain; Brain Tumour; Stroke; Seizure; Meningitis; Cerebral Palsy; Parkinson's Disease.

## **Breast**

Breast; Breast Cancer; Mastitis; Mammography.

## **Cervix**

Cervix; Cervical Cancer; Cervicitis; Pap Smear.

## **Colon**

Colon; Colitis; Constipation; Diarrhea; Colorectal Cancer; Colonoscopy.

## **Ear**

Ear; Otitis Media; Tinnitus; Hearing Loss; Vertigo.

## **Esophagus**

Esophagus; Dysphagia; Esophagitis; Achalasia; Gastroesophageal Reflux Disease.

## **Eye**

Eye; Cataract; Glaucoma; Conjunctivitis; Retinal Detachment; Macular Degeneration.

## **Gallbladder**

Gallbladder; Cholelithiasis; Cholecystitis; Bile Duct; Cholangiocarcinoma.

## **Heart**

Heart; Heart Failure; Arrhythmia; Myocardial Infarction; Pericarditis; Congenital Heart Defect.

## **Kidney**

Kidney; Acute Kidney Injury; Chronic Kidney Disease; Glomerulonephritis; Nephritis; Dialysis.

## **Liver**

Liver; Hepatitis; Cirrhosis; Fatty Liver Disease; Jaundice; Portal Hypertension.

## **Lung**

Lung; Asthma; COPD; Pneumonia; Bronchitis; Pulmonary Embolism; Pleural Effusion.

---

**Mouth**

Mouth; Dental Caries; Gingivitis; Periodontitis.

**Nose**

Nose; Sinusitis; Rhinitis; Nasal Polyps; Anosmia; Epistaxis.

**Ovary**

Ovary; Ovarian Cyst; Ovarian Cancer; Fallopian Tube; Ectopic Pregnancy.

**Pancreas**

Pancreas; Pancreatitis; Diabetes Mellitus; Pancreatic Cancer; Insulin.

**Pituitary Gland**

Pituitary Gland; Pituitary Tumor; Acromegaly; Diabetes Insipidus.

**Prostate**

Prostate; Benign Prostatic Hyperplasia; Prostatitis; Prostate Cancer.

**Rectum**

Rectum; Rectal Bleeding; Proctitis; Rectal Cancer.

**Skin**

Skin; Dermatitis; Eczema; Psoriasis; Cellulitis; Pressure Ulcer; Melanoma.

**Small Intestine**

Small Intestine; Enteritis; Celiac Disease; Malabsorption; Intestinal Obstruction.

**Spinal Cord**

Spinal Cord; Spinal Cord Injury; Myelitis; Spinal Stenosis; Paraplegia.

**Spleen**

Spleen; Splenomegaly; Splenectomy; Hypersplenism.

**Stomach**

Stomach; Gastritis; Peptic Ulcer Disease; Gastroesophageal Reflux Disease; Gastric Cancer.

**Testis**

Testis; Testicular Torsion; Testicular Cancer; Epididymis.

**Thymus**

Thymus; Thymoma.

**Thyroid Gland**

Thyroid Gland; Hyperthyroidism; Hypothyroidism; Goitre; Thyroid Cancer.

**Ureter**

Ureter; Ureteral Calculi; Ureteral Obstruction; Ureteritis; Ureteral Stricture.

**Urethra**

Urethra; Urethral Stricture; Urethritis; Urethral Obstruction.

**Uterus**

Uterus; Endometriosis; Fibroids; Uterine Cancer; Ectopic Pregnancy.

**Vagina**

Vagina; Vaginitis; Vaginal Discharge; Vaginal Cancer.

# Common Diseases

*Scope:* This reference-oriented list organizes common human diseases by organ system and clinical domain. It is a curated, high-importance list rather than a prevalence ranking, and some diseases appear under more than one heading where they are clinically relevant to several systems. It is not an exhaustive index of every term in the dictionary. Entries named as syndromes are covered in the companion appendix, Common Syndromes; conditions conventionally classified as disorders are covered in Common Disorders; and clinical findings, lesions, congenital anomalies, and injuries are covered in their own companion appendices. Individual symptoms, signs, and laboratory findings are defined as dictionary entries rather than listed here.

**Cardiovascular Diseases** Hypertension; Coronary Artery Disease; Stable Angina; Unstable Angina; Myocardial Infarction; Atherosclerosis; Heart Failure; Right Heart Failure; Cardiogenic Shock; Atrial Fibrillation; Atrial Flutter; Supraventricular Tachycardia; Ventricular Tachycardia; Ventricular Fibrillation; Sinus Bradycardia; Heart Block; Cardiomyopathy; Dilated Cardiomyopathy; Hypertrophic Cardiomyopathy; Restrictive Cardiomyopathy; Myocarditis; Pericarditis; Cardiac Tamponade; Endocarditis; Rheumatic Heart Disease; Aortic Stenosis; Aortic Regurgitation; Mitral Stenosis; Mitral Regurgitation; Mitral Valve Prolapse; Tricuspid Valve Disease; Pulmonary Hypertension; Peripheral Artery Disease; Aortic Aneurysm; Aortic Dissection; Deep Vein Thrombosis; Chronic Venous Insufficiency; Varicose Veins; Thrombophlebitis; Raynaud Phenomenon; Congenital Heart Disease.

**Respiratory Diseases** Asthma; Acute Bronchitis; Chronic Bronchitis; Emphysema; COPD; Pneumonia; Aspiration Pneumonia; Bronchopneumonia; Tuberculosis; COVID-19; Influenza; Bronchiolitis; Bronchiectasis; Cystic Fibrosis; Interstitial Lung Disease; Idiopathic Pulmonary Fibrosis; Pneumoconiosis; Silicosis; Asbestosis; Hypersensitivity Pneumonitis; Sarcoidosis; Pulmonary Edema; Pulmonary Embolism; Pulmonary Hypertension; Respiratory Failure; Pleurisy; Empyema; Lung Abscess; Occupational Lung Disease.

**Gastrointestinal Diseases** Gastroesophageal Reflux Disease; Esophagitis; Barrett Esophagus; Achalasia; Esophageal Spasm; Gastritis; Peptic Ulcer Disease; Gastric Ulcer; Duodenal Ulcer; Gastroenteritis; Gastroparesis; Crohn Disease; Ulcerative Colitis; Inflammatory Bowel Disease; Celiac Disease; Lactose Intolerance; Food Intolerance; Intestinal Obstruction; Ileus; Intussusception; Volvulus; Diverticulosis; Diverticulitis; Appendicitis; Peritonitis; Hemorrhoids; Anal Fissure; Anal Fistula; Rectal Prolapse; Mesenteric Ischemia.

**Liver, Gallbladder & Biliary Diseases** Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD); Alcoholic Liver Disease; Hepatitis A; Hepatitis B; Hepatitis C; Hepatitis D; Hepatitis E; Autoimmune Hepatitis; Acute Liver Failure; Chronic Liver Disease; Cirrhosis; Portal Hypertension; Hepatic Encephalopathy; Hemochromatosis; Wilson Disease; Primary Biliary Cholangitis; Primary Sclerosing Cholangitis; Cholelithiasis; Cholecystitis; Cholangitis.

**Pancreatic Diseases** Acute Pancreatitis; Chronic Pancreatitis; Autoimmune Pancreatitis; Pancreatic Insufficiency; Pancreatic Cancer; Pancreatic Neuroendocrine Tumors.

**Kidney & Urinary Diseases** Acute Kidney Injury; Chronic Kidney Disease; End-Stage Kidney Disease; Glomerulonephritis; Diabetic Nephropathy; Hypertensive Nephropathy; Interstitial Nephritis; Pyelonephritis; Urinary Tract Infection; Cystitis; Urethritis; Nephrolithiasis; Hydronephrosis; Polycystic Kidney Disease; Renal Tubular Acidosis; IgA Nephropathy; Minimal Change Disease; Focal Segmental Glomerulosclerosis; Membranous Nephropathy; Vesicoureteral Reflux.

---

**Endocrine Diseases** Type 1 Diabetes Mellitus; Type 2 Diabetes Mellitus; Gestational Diabetes; Diabetes Insipidus; Diabetic Ketoacidosis; Hyperosmolar Hyperglycemic State; Hypothyroidism; Hyperthyroidism; Hashimoto Thyroiditis; Graves Disease; Thyroiditis; Goiter; Addison Disease; Cushing Disease; Primary Aldosteronism; Hypoparathyroidism; Hyperparathyroidism; Pheochromocytoma; Pituitary Adenoma; Acromegaly; Gigantism; Growth Hormone Deficiency.

**Metabolic Diseases** Obesity; Gout; Dyslipidemia; Hypercholesterolemia; Hypertriglyceridemia; Porphyria; Amyloidosis.

**Nutritional Diseases & Deficiencies** Protein-Energy Malnutrition; Marasmus; Kwashiorkor; Iron Deficiency; Iron-Deficiency Anemia; Vitamin A Deficiency; Thiamine Deficiency; Beriberi; Riboflavin Deficiency; Niacin Deficiency; Pellagra; Folate Deficiency; Vitamin B12 Deficiency; Vitamin C Deficiency; Scurvy; Vitamin D Deficiency; Rickets; Osteomalacia; Vitamin K Deficiency; Iodine Deficiency; Zinc Deficiency; Copper Deficiency; Selenium Deficiency; Magnesium Deficiency.

**Neurological Diseases** Tension-Type Headache; Cluster Headache; Status Epilepticus; Stroke; Ischemic Stroke; Hemorrhagic Stroke; Transient Ischemic Attack; Cerebral Aneurysm; Parkinson Disease; Alzheimer Disease; Vascular Dementia; Frontotemporal Dementia; Lewy Body Dementia; Multiple Sclerosis; Amyotrophic Lateral Sclerosis; Huntington Disease; Diabetic Neuropathy; Myasthenia Gravis; Bell Palsy; Trigeminal Neuralgia; Radiculopathy; Meningitis; Encephalitis; Hydrocephalus; Cerebral Palsy.

**Musculoskeletal Diseases** Osteoarthritis; Osteoporosis; Osteomalacia; Rheumatoid Arthritis; Ankylosing Spondylitis; Psoriatic Arthritis; Reactive Arthritis; Septic Arthritis; Gout; Bursitis; Tendinitis; Tendinopathy; Frozen Shoulder; Tennis Elbow; Golfer's Elbow; Cervical Spondylosis; Lumbar Spondylosis; Spinal Stenosis; Herniated Disc; Scoliosis; Kyphosis; Spondylolisthesis; Fibromyalgia; Muscular Dystrophy; Myositis; Polymyositis; Dermatomyositis.

**Rheumatologic & Autoimmune Diseases** Rheumatoid Arthritis; Systemic Lupus Erythematosus; Polymyalgia Rheumatica; Giant Cell Arteritis; Polyarteritis Nodosa; Granulomatosis with Polyangiitis; Microscopic Polyangiitis; Behçet Disease; Inflammatory Myopathies; Ankylosing Spondylitis; Psoriatic Arthritis; Reactive Arthritis; Relapsing Polychondritis; Adult-Onset Still Disease.

**Skin, Hair & Nail Diseases** Acne Vulgaris; Rosacea; Atopic Dermatitis; Contact Dermatitis; Seborrheic Dermatitis; Psoriasis; Urticaria; Angioedema; Impetigo; Cellulitis; Erysipelas; Folliculitis; Boils; Carbuncle; Ringworm; Athlete's Foot; Jock Itch; Candidiasis; Scabies; Pediculosis; Wart; Molluscum Contagiosum; Herpes Simplex; Herpes Zoster; Hidradenitis Suppurativa; Pressure Ulcers; Onychomycosis.

**Eye Diseases** Amblyopia; Strabismus; Cataract; Open-Angle Glaucoma; Angle-Closure Glaucoma; Conjunctivitis; Keratitis; Corneal Ulcer; Dry Eye; Uveitis; Iritis; Retinal Detachment; Diabetic Retinopathy; Hypertensive Retinopathy; Macular Degeneration; Retinitis Pigmentosa; Central Retinal Artery Occlusion; Central Retinal Vein Occlusion; Optic Neuritis; Keratoconus; Blepharitis.

**Ear & Hearing Diseases** Otitis Externa; Acute Otitis Media; Chronic Otitis Media; Otitis Media with Effusion; Mastoiditis; Presbycusis; Meniere Disease; Labyrinthitis; Vestibular Neuritis; Benign Paroxysmal Positional Vertigo; Otosclerosis; Cholesteatoma; Acoustic Neuroma.

**Nose, Sinus & Throat Diseases** Allergic Rhinitis; Nonallergic Rhinitis; Acute Sinusitis; Chronic Sinusitis; Pharyngitis; Streptococcal Pharyngitis; Tonsillitis; Peritonsillar Abscess; Laryngitis; Epiglottitis; Vocal Cord Paralysis.

**Oral & Dental Diseases** Dental Caries; Pulpitis; Dental Abscess; Gingivitis; Periodontitis; Pericoronitis; Aphthous Ulcer; Oral Candidiasis; Leukoplakia; Oral Lichen Planus.

**Blood Diseases** Iron-Deficiency Anemia; Megaloblastic Anemia; Vitamin B12 Deficiency Anemia; Folate-Deficiency Anemia; Aplastic Anemia; Hemolytic Anemia; Sick Cell Disease; Thalassemia; Hereditary Spherocytosis; Glucose-6-Phosphate Dehydrogenase Deficiency; Polycythemia Vera; Immune Thrombocytopenia; Agranulocytosis; Hemophilia A; Hemophilia B; Von Willebrand Disease; Disseminated Intravascular Coagulation (DIC); Thrombotic Thrombocytopenic Purpura (TTP).

**Immune-System Diseases** Primary Immunodeficiency; Severe Combined Immunodeficiency; Common Variable Immunodeficiency; Selective IgA Deficiency; AIDS (Acquired Immunodeficiency Syndrome); HIV/AIDS; Chronic Granulomatous Disease; Graft-Versus-Host Disease.

---



---

**Infectious & Parasitic Diseases** Common Cold; Influenza; COVID-19; Measles; Mumps; Rubella; Chickenpox; Shingles; Herpes Simplex; Infectious Mononucleosis; Cytomegalovirus Infection; Polio; Rabies; Dengue; Chikungunya; Zika; Hepatitis A; Hepatitis B; Hepatitis C; Hepatitis D; Hepatitis E; HIV; Syphilis; Gonorrhea; Chlamydia; Tuberculosis; Leprosy; Diphtheria; Pertussis; Tetanus; Typhoid Fever; Cholera; Plague; Bacterial Meningitis; Meningococcemia; Scarlet Fever; Salmonellosis; Shigellosis; Campylobacteriosis; Botulism; Brucellosis; Leptospirosis; Lyme Disease; Malaria; Amebiasis; Giardiasis; Toxoplasmosis; Leishmaniasis; Trichomoniasis; Ascariasis; Hookworm; Pinworm Infection; Schistosomiasis; Taeniasis; Echinococcosis; Candidiasis; Aspergillosis; Cryptococcosis; Histoplasmosis.

**Female Reproductive & Gynecological Diseases** Endometriosis; Adenomyosis; Uterine Fibroids; Ovarian Torsion; Pelvic Inflammatory Disease; Vaginitis; Bacterial Vaginosis; Vulvovaginal Candidiasis; Vulvodynia; Pelvic Organ Prolapse; Female Infertility; Premature Ovarian Insufficiency.

**Male Reproductive & Urological Diseases** Male Infertility; Benign Prostatic Hyperplasia; Prostatitis; Epididymitis; Orchitis; Testicular Torsion; Cryptorchidism; Phimosis; Paraphimosis; Peyronie Disease; Balanitis; Hypogonadism.

**Pregnancy & Postpartum Diseases** Hyperemesis Gravidarum; Gestational Diabetes; Gestational Hypertension; Preeclampsia; Eclampsia; Puerperal Infection; Postpartum Endometritis; Postpartum Depression; Mastitis.

**Pediatric & Neonatal Diseases** Respiratory Distress Syndrome; Neonatal Sepsis; Birth Asphyxia; Necrotizing Enterocolitis; Diaper Rash; Croup; Bronchiolitis; Cerebral Palsy; Congenital Hypothyroidism; Kawasaki Disease; Rheumatic Fever; Hand-Foot-and-Mouth Disease.

**Genetic & Chromosomal Diseases** Cystic Fibrosis; Sickle Cell Disease; Thalassemia; Huntington Disease; Duchenne Muscular Dystrophy; Becker Muscular Dystrophy; Neurofibromatosis; Tuberous Sclerosis; Tay-Sachs Disease; Gaucher Disease; Achondroplasia; Hemochromatosis; Wilson Disease.

**Cancers & Other Neoplasms** Breast Cancer; Lung Cancer; Colorectal Cancer; Prostate Cancer; Cervical Cancer; Ovarian Cancer; Endometrial Cancer; Vulvar Cancer; Vaginal Cancer; Testicular Cancer; Penile Cancer; Bladder Cancer; Kidney Cancer; Thyroid Cancer; Oral Cancer; Oropharyngeal Cancer; Laryngeal Cancer; Esophageal Cancer; Stomach Cancer; Liver Cancer; Gallbladder Cancer; Pancreatic Cancer; Small-Intestinal Cancer; Anal Cancer; Melanoma; Basal-Cell Carcinoma; Squamous-Cell Carcinoma; Osteosarcoma; Ewing Sarcoma; Soft-Tissue Sarcoma; Glioma; Glioblastoma; Meningioma; Leukemia; Hodgkin Lymphoma; Non-Hodgkin Lymphoma; Multiple Myeloma; Neuroblastoma; Retinoblastoma.

**Multisystem & Systemic Diseases** Fibromyalgia; Sarcoidosis; Amyloidosis; Sepsis; Septic Shock.



# Common Disorders

*Scope:* This reference-oriented list organizes human conditions conventionally classified as clinical disorders by medical specialty and organ system. In medical nomenclature, disorders represent disruptions to regular bodily structure or function (frequently functional, metabolic, neurobehavioral, or regulatory disturbances), complementing the disease classifications in Common Diseases and clinical signs in Common Syndromes. Every entity listed corresponds to a defined medical concept in the dictionary.

**Mental & Behavioral Disorders** Generalized Anxiety Disorder; Panic Disorder; Social Anxiety Disorder; Agoraphobia; Specific Phobia; Major Depressive Disorder; Persistent Depressive Disorder; Bipolar Disorder; Cyclothymic Disorder; Obsessive-Compulsive Disorder; Post-Traumatic Stress Disorder; Acute Stress Disorder; Attention-Deficit/Hyperactivity Disorder; Autism Spectrum Disorder; Schizophrenia; Schizoaffective Disorder; Delusional Disorder; Brief Psychotic Disorder; Eating Disorders; Alcohol Use Disorder; Opioid Use Disorder; Substance Abuse; Dissociative Identity Disorder; Depersonalization.

**Personality Disorders** Borderline Personality Disorder; Antisocial Personality Disorder; Narcissistic Personality Disorder; Avoidant Personality Disorder; Dependent Personality Disorder; Obsessive-Compulsive Personality Disorder; Paranoid Personality Disorder; Schizoid Personality Disorder; Schizotypal Personality Disorder; Histrionic Personality Disorder.

**Eating & Nutrition Disorders** Anorexia Nervosa; Bulimia Nervosa; Binge-Eating Disorder; Avoidant/Restrictive Food Intake Disorder; Pica; Rumination.

**Impulse-Control & Conduct Disorders** Intermittent Explosive Disorder; Kleptomania; Pyromania; Oppositional Defiant Disorder; Conduct Disorder; Trichotillomania; Excoriation Disorder.

**Somatic & Related Disorders** Somatic Symptom Disorder; Illness Anxiety Disorder; Conversion Disorder; Factitious Disorder.

**Sleep-Wake Disorders** Insomnia; Hypersomnia; Narcolepsy; Obstructive Sleep Apnea; Central Sleep Apnea; Sleepwalking; Night Terrors; Nightmare Disorder; REM Sleep Behavior Disorder; Restless Legs Syndrome; Delayed Sleep Phase.

**Neurological & Movement Disorders** Migraine; Cluster Headache; Tension-Type Headache; Epilepsy; Seizure Disorder; Essential Tremor; Parkinson's Disease; Dystonia; Chorea; Myoclonus; Tic Disorder; Tourette Disorder; Ataxia; Stiff-Person Syndrome.

**Neurodevelopmental & Learning Disorders** Intellectual Disability; Language Disorder; Stuttering; Specific Learning Disorder; Dyslexia; Developmental Coordination Disorder; Fetal Alcohol Spectrum Disorder.

**Autonomic Disorders** Dysautonomia; Postural Orthostatic Tachycardia Syndrome; Orthostatic Hypotension; Neurocardiogenic Syncope; Multiple System Atrophy; Autonomic Neuropathy.

**Cardiovascular & Hemodynamic Disorders** Heart Arrhythmia; Atrioventricular Block; Bundle Branch Block; Microvascular Angina; Cardiac Syndrome X; Orthostatic Intolerance.

---

**Respiratory & Ventilatory Disorders** Chronic Obstructive Pulmonary Disorder; Hypoventilation; Obesity Hypoventilation Syndrome; Pulmonary Fibrosis; Upper Airway Resistance Syndrome; Hyperventilation.

**Digestive & Functional GI Disorders** Irritable Bowel Syndrome; Functional Dyspepsia; Functional Constipation; Functional Diarrhea; Cyclic Vomiting Syndrome; Rumination; Achalasia; Diffuse Esophageal Spasm; Pelvic Floor Dysfunction; Gastroparesis.

**Endocrine & Metabolic Disorders** Hypothyroidism; Hyperthyroidism; Euthyroid Sick Syndrome; Hypopituitarism; Hyperprolactinemia; Adrenal Insufficiency; Cushing Syndrome; Hypoparathyroidism; Hyperparathyroidism; Polycystic Ovary Syndrome; Precocious Puberty; Delayed Puberty; Metabolic Syndrome.

**Lipid & Inborn Metabolic Disorders** Dyslipidemia; Hypercholesterolemia; Hypertriglyceridemia; Familial Hypercholesterolemia; Galactosemia; Phenylketonuria; Maple Syrup Urine Disease; Glycogen Storage Disease; Lysosomal Storage Disorders; Porphyria.

**Electrolyte & Acid–Base Disorders** Hyponatremia; Hypernatremia; Hypokalemia; Hyperkalemia; Hypocalcemia; Hypercalcemia; Hypomagnesemia; Hypermagnesemia; Hypophosphatemia; Hyperphosphatemia; Metabolic Acidosis; Metabolic Alkalosis; Respiratory Acidosis; Respiratory Alkalosis.

**Bone Mineral Density Disorders** Osteopenia; Osteoporosis; Osteomalacia; Renal Osteodystrophy; Fibrous Dysplasia; Paget Disease of Bone.

**Musculoskeletal & Soft-Tissue Disorders** Temporomandibular Joint Disorders; Rotator Cuff Tear; Rotator Cuff Injury; Herniated Disc; Degenerative Disc Disease; Myofascial Pain Syndrome; Tendinitis; Tendinopathy; Carpal Tunnel Syndrome; Cubital Tunnel Syndrome; Hypermobility Syndrome.

**Blood & Coagulation Disorders** Hemophilia A; Hemophilia B; Von Willebrand Disease; Thrombophilia; Glanzmann Thrombasthenia; Immune Thrombocytopenic Purpura; Aplastic Anemia; Myelodysplastic Syndrome.

**Immune, Allergic & Eosinophilic Disorders** Autoimmune Disease; Primary Immunodeficiency; Anaphylaxis; Angioedema; Mastocytosis; Eosinophilic Esophagitis; Eosinophilic Gastroenteritis.

**Sexual Disorders** Male Hypoactive Sexual Desire Disorder; Erectile Dysfunction; Premature Ejaculation; Delayed Ejaculation; Female Sexual Dysfunction; Female Orgasmic Disorder; Vaginismus; Dyspareunia; Gender Dysphoria.

**Menstrual & Reproductive Disorders** Dysmenorrhea; Amenorrhea; Abnormal Uterine Bleeding; Menorrhagia; Premenstrual Syndrome; Premenstrual Dysphoric Disorder; Anovulation; Premature Ovarian Insufficiency; Androgen Insensitivity Syndrome.

**Urinary & Micturition Disorders** Overactive Bladder; Urinary Incontinence; Stress Incontinence; Neurogenic Bladder; Urinary Retention; Bladder Outlet Obstruction.

**Vision & Ocular Motility Disorders** Refractive Disorder; Myopia; Hyperopia; Astigmatism; Presbyopia; Amblyopia; Strabismus; Color Vision Deficiency; Nystagmus; Dry Eye.

**Hearing, Balance & Auditory Disorders** Conductive Hearing Loss; Sensorineural Hearing Loss; Tinnitus; Auditory Processing Disorder; Benign Paroxysmal Positional Vertigo; Meniere Disease.

**Speech, Language & Voice Disorders** Language Disorder; Speech Disorder; Stuttering; Dysarthria; Apraxia; Aphasia; Voice Disorders; Dysphonia; Vocal Cord Dysfunction.

**Hair, Nail & Pigmentation Disorders** Alopecia; Alopecia Areata; Androgenetic Alopecia; Telogen Effluvium; Cicatricial Alopecia; Melasma; Vitiligo; Albinism.

**Connective Tissue & Fascial Disorders** Ehlers-Danlos Syndrome; Marfan Syndrome; Loeys-Dietz Syndrome; Mixed Connective Tissue Disease; Dupuytren Contracture; Plantar Fasciitis.

---

# Mental Health Terms

*Scope:* This appendix groups commonly used mental-health terms for quick reference. It is an index, not a diagnostic or treatment guide. Preferred modern terms are used where an older or informal term may be misleading.

**Foundations and assessment** Acute; Affect; Chronic; Cognition; Diagnosis; Disorder; Distress; Functional Impairment; Mental Disorder; Mental Health; Mental Illness; Mood; Risk Factor; Syndrome.

**Brain and nervous-system structures** Amygdala; Axon; Brain; Brain Stem; Cerebellum; Cerebrum; Dendrite; Frontal Lobe; Gray Matter; Hippocampus; Hypothalamus; Limbic System; Locus Coeruleus; Myelin; Neuron; Neurotransmission; Neurotransmitter; Occipital Lobe; Parietal Lobe; Postsynaptic Membrane; Presynaptic Terminals; Receptor; Serotonin; Synapse; Temporal Lobe; Thalamus.

**Common psychological and behavioral symptoms** Anhedonia; Anxiety; Avolition; Cognitive Symptoms; Compulsions; Delusion; Euphoria; Obsessions; Panic Attack; Psychomotor Agitation; Psychomotor Retardation; Psychosis; Self-Harm; Stress; Trauma.

**Anxiety, trauma, and obsessive-compulsive disorders** Anxiety Disorders; Generalized Anxiety Disorder; Panic Disorder; Specific Phobia; Social Anxiety Disorder (Social Phobia); Separation Anxiety Disorder; Post-Traumatic Stress Disorder; Obsessive-Compulsive Disorder; Trichotillomania.

**Mood disorders** Bipolar Disorder; Bipolar II Disorder; Cyclothymic Disorder; Depression; Major Depressive Disorder; Major Depressive Episode; Persistent Depressive Disorder; Postpartum Depression; Seasonal Affective Disorder.

**Psychotic and personality disorders** Antisocial Personality Disorder; Borderline Personality Disorder; Narcissistic Personality Disorder; Personality Disorders; Brief Psychotic Disorder; Schizoaffective Disorder; Schizophrenia.

**Neurodevelopmental and related conditions** Conduct Disorder; Neurodevelopmental Disorders; Intellectual Disability; Neurocognitive Disorder; Functional Neurologic Disorder.

**Psychiatric and psychological treatments** Antidepressants; Antipsychotic Medicine; Atypical Antipsychotics; Benzodiazepine; Cognitive Behavioral Therapy; Dialectical Behavior Therapy; Electroconvulsive Therapy; Psychotherapy; Systematic Desensitization; Transcranial Magnetic Stimulation.

**Care settings and professionals** Community Mental-Health Treatment; Hospitalization; Psychiatry; Psychiatrist; Clinical Psychologist; Social Worker; Voluntary Psychiatric Admission.

**Investigations and brain imaging** Computed Tomography (CT); Electroencephalography; Functional MRI; Positron Emission Tomography; Single-Photon Emission Computed Tomography.

**Suicide and safety terminology** Suicidality; Suicide; Suicide Attempt; Suicidal Ideation; Suicidal Intent; Suicidal Plan; Suicide Risk Assessment.

**Related scientific concepts** Evidence-Based Practice; Neurosciences; Tolerance; Withdrawal.





# Common Syndromes

*Scope:* This reference-oriented list organizes clinically significant medical syndromes by organ system and medical specialty. In medical terminology, a syndrome represents a characteristic constellation of concurrent signs, symptoms, or anatomical features that consistently occur together and point toward a common pathophysiological mechanism or etiology.

**Cardiovascular Syndromes** Acute Coronary Syndrome; Acute Aortic Syndrome; Sick Sinus Syndrome; Wolff–Parkinson–White Syndrome; Long QT Syndrome; Short QT Syndrome; Brugada Syndrome; Takotsubo Syndrome (Stress Cardiomyopathy); Eisenmenger Syndrome; Postural Orthostatic Tachycardia Syndrome (POTS); Postpericardiotomy Syndrome; Dressler Syndrome; Cardiac Syndrome X; Cardiorenal Syndrome.

**Respiratory Syndromes** Acute Respiratory Distress Syndrome (ARDS); Infant Respiratory Distress Syndrome; Obesity Hypoventilation Syndrome; Upper Airway Resistance Syndrome; Hepatopulmonary Syndrome; Acute Chest Syndrome; Kartagener Syndrome.

**Gastrointestinal Syndromes** Irritable Bowel Syndrome (IBS); Malabsorption Syndrome; Short Bowel Syndrome; Dumping Syndrome; Cyclic Vomiting Syndrome; Rumination Syndrome; Mallory–Weiss Syndrome; Zollinger–Ellison Syndrome; Boerhaave Syndrome; Afferent Loop Syndrome; Cannabinoid Hyperemesis Syndrome; Superior Mesenteric Artery Syndrome; Ogilvie Syndrome.

**Liver & Biliary Syndromes** Hepatorenal Syndrome; Hepatopulmonary Syndrome; Budd–Chiari Syndrome; Gilbert Syndrome; Crigler–Najjar Syndrome; Dubin–Johnson Syndrome; Rotor Syndrome; Reye Syndrome; Mirizzi Syndrome.

**Kidney & Urinary Syndromes** Nephrotic Syndrome; Nephritic Syndrome; Hemolytic Uremic Syndrome (HUS); Alport Syndrome; Fanconi Syndrome; Goodpasture Syndrome; Hepatorenal Syndrome; Nutcracker Syndrome.

**Endocrine Syndromes** Cushing Syndrome; Metabolic Syndrome; Syndrome of Inappropriate Antidiuretic Hormone Secretion (SIADH); Polycystic Ovary Syndrome (PCOS); Carcinoid Syndrome; Empty Sella Syndrome; Androgen Insensitivity Syndrome; Sheehan Syndrome; Nelson Syndrome; Polyglandular Autoimmune Syndromes (PAPS Type 1 and 2); Insulin Resistance Syndrome.

**Neurological Syndromes** Guillain–Barré Syndrome; Miller Fisher Syndrome; Lambert–Eaton Myasthenic Syndrome (LEMS); Restless Legs Syndrome; Locked-In Syndrome; Post-Concussion Syndrome; Cauda Equina Syndrome; Conus Medullaris Syndrome; Anterior Cord Syndrome; Central Cord Syndrome; Central Pain Syndrome; Alien Hand Syndrome; CADASIL Syndrome; Foix–Alajouanine Syndrome; Horner Syndrome; Tourette Syndrome; Lennox–Gastaut Syndrome; West Syndrome; Wernicke–Korsakoff Syndrome; Brown–Séquard Syndrome; Wallenberg Syndrome (Lateral Medullary Syndrome); Isaacs Syndrome.

**Pain & Neuromuscular Syndromes** Carpal Tunnel Syndrome; Cubital Tunnel Syndrome; Complex Regional Pain Syndrome (CRPS); Myofascial Pain Syndrome; Piriformis Syndrome; Compartment Syndrome; Thoracic Outlet Syndrome; Burning Mouth Syndrome; Tarsal Tunnel Syndrome; Greater Trochanteric Pain Syndrome.

**Psychiatric & Behavioral Syndromes** Serotonin Syndrome; Neuroleptic Malignant Syndrome; Capgras Syndrome; Cotard Syndrome; Münchausen Syndrome (Factitious Disorder); Korsakoff Syndrome; Amotivational Syndrome; Ekblom Syndrome (Delusional Parasitosis); Ganser Syndrome.

---

**Rheumatologic & Autoimmune Syndromes** Sjögren Syndrome; Antiphospholipid Syndrome; CREST Syndrome; Felty Syndrome; Behçet Syndrome; SAPHO Syndrome; Reactive Arthritis (formerly Reiter Syndrome); Churg–Strauss Syndrome (EGPA); Caplan Syndrome.

**Hematological Syndromes** Myelodysplastic Syndromes; Antiphospholipid Syndrome; Evans Syndrome; Hyperviscosity Syndrome; Tumor Lysis Syndrome; Hemophagocytic Syndrome (HLH); Acute Chest Syndrome; Thrombotic Thrombocytopenic Purpura (TTP).

**Immunological Syndromes** Severe Combined Immunodeficiency Syndrome; Acquired Immunodeficiency Syndrome (AIDS); Alpha-Gal Syndrome; Antisynthetase Syndrome; Capillary Leak Syndrome; Hyper-IgE Syndrome (Job Syndrome); Autoimmune Lymphoproliferative Syndrome (ALPS); Cytokine Release Syndrome (CRS); Immune Reconstitution Inflammatory Syndrome; Wiskott–Aldrich Syndrome.

**Infectious & Postinfectious Syndromes** Toxic Shock Syndrome (TSS); Post-COVID-19 Condition; Post-Polio Syndrome; Congenital Rubella Syndrome; Congenital Zika Syndrome; Waterhouse–Friderichsen Syndrome; Lemierre Syndrome.

**Genetic & Chromosomal Syndromes** Down Syndrome; Turner Syndrome; Klinefelter Syndrome; Edwards Syndrome; Patau Syndrome; DiGeorge Syndrome (22q11.2 Deletion); Fragile X Syndrome; Williams Syndrome; Prader–Willi Syndrome; Angelman Syndrome; Noonan Syndrome; Rett Syndrome; Cri-du-Chat Syndrome; Beckwith–Wiedemann Syndrome; McCune–Albright Syndrome.

**Connective Tissue Syndromes** Ehlers–Danlos Syndromes; Marfan Syndrome; Loeys–Dietz Syndrome; Stickler Syndrome; Joint Hypermobility Syndrome.

**Pediatric Syndromes** Down Syndrome; Fetal Alcohol Syndrome; Sudden Infant Death Syndrome (SIDS); Reye Syndrome; DiGeorge Syndrome; Beckwith–Wiedemann Syndrome; Nephrotic Syndrome; Hand-Foot-and-Mouth Syndrome.

**Obstetric Syndromes** HELLP Syndrome; Antiphospholipid Syndrome; Twin-to-Twin Transfusion Syndrome (TTTS); Amniotic Band Syndrome; Fetal Alcohol Syndrome.

**Gynecological Syndromes** Polycystic Ovary Syndrome; Premenstrual Syndrome (PMS); Asherman Syndrome; Genitourinary Syndrome of Menopause; Mayer–Rokitansky–Küster–Hauser (MRKH) Syndrome.

**Male Reproductive Syndromes** Klinefelter Syndrome; Androgen Insensitivity Syndrome; Kallmann Syndrome; Post-Vasectomy Pain Syndrome.

**Cancer-Predisposition & Paraneoplastic Syndromes** Lynch Syndrome (HNPCC); Li–Fraumeni Syndrome; Von Hippel–Lindau Syndrome; Multiple Endocrine Neoplasia (MEN 1 and 2); Carcinoid Syndrome; Tumor Lysis Syndrome; Superior Vena Cava Syndrome; Paraneoplastic Neurological Syndromes; Cancer Cachexia Syndrome.

**Metabolic & Nutritional Syndromes** Metabolic Syndrome; Refeeding Syndrome; Wernicke–Korsakoff Syndrome; Milk–Alkali Syndrome; Hungry Bone Syndrome; Fanconi–Bickel Syndrome.

**Toxicological Syndromes (Toxidromes)** Serotonin Syndrome; Neuroleptic Malignant Syndrome; Anticholinergic Syndrome; Cholinergic Syndrome; Sympathomimetic Syndrome; Sedative-Hypnotic Syndrome; Opioid Withdrawal Syndrome; Alcohol Withdrawal Syndrome.

**Critical Care Syndromes** Systemic Inflammatory Response Syndrome (SIRS); Multiple Organ Dysfunction Syndrome (MODS); Acute Respiratory Distress Syndrome (ARDS); Abdominal Compartment Syndrome; Fat Embolism Syndrome.

**Sleep Syndromes** Obstructive Sleep Apnea Syndrome; Central Sleep Apnea Syndromes; Restless Legs Syndrome; Kleine–Levin Syndrome; Delayed Sleep Phase Syndrome.

**Eye & Cranial Nerve Syndromes** Horner Syndrome; Adie Syndrome; Brown Syndrome; Duane Retraction Syndrome; Charles Bonnet Syndrome; Dry Eye Syndrome; Tolosa–Hunt Syndrome.

**ENT & Otolaryngologic Syndromes** Ramsay Hunt Syndrome; Eagle Syndrome; Empty Nose Syndrome; Gradenigo Syndrome; Samter Triad (Aspirin-Exacerbated Respiratory Disease).

---

---

**Dermatological Syndromes** Stevens–Johnson Syndrome; Toxic Epidermal Necrolysis; Staphylococcal Scalded Skin Syndrome; Sweet Syndrome; Gorlin Syndrome; Peutz–Jeghers Syndrome; DRESS Syndrome (Drug Reaction with Eosinophilia and Systemic Symptoms).

**Vascular Syndromes** Raynaud Syndrome; Superior Vena Cava Syndrome; Thoracic Outlet Syndrome; Subclavian Steal Syndrome; Leriche Syndrome; May–Thurner Syndrome; Nutcracker Syndrome.

**Multisystem Syndromes** Chronic Fatigue Syndrome (ME/CFS); Postural Orthostatic Tachycardia Syndrome (POTS); Ehlers–Danlos Syndromes; Metabolic Syndrome; Antiphospholipid Syndrome; Behcet Syndrome.



# Rare Diseases

*Scope:* This reference-oriented list organizes clinically significant rare diseases (orphan diseases) by organ system and clinical domain. A rare disease is typically defined as one affecting fewer than 1 in 2,000 individuals (EU criteria) or fewer than 200,000 people in the United States. It highlights high-importance benchmark conditions characterized by distinct genetic, metabolic, or autoimmune etiologies.

**Genetic & Chromosomal Diseases** Turner Syndrome; Klinefelter Syndrome; Fragile X Syndrome; Williams Syndrome; Prader–Willi Syndrome; Angelman Syndrome; Noonan Syndrome; 22q11.2 Deletion Syndrome; Cri-du-Chat Syndrome; Hutchinson–Gilford Progeria Syndrome; Cornelia de Lange Syndrome.

**Neurological & Neurogenetic Diseases** Huntington Disease; Amyotrophic Lateral Sclerosis (ALS); Spinal Muscular Atrophy (SMA); Friedreich Ataxia; Ataxia–Telangiectasia; Rett Syndrome; Batten Disease (Neuronal Ceroid Lipofuscinosis); Creutzfeldt–Jakob Disease; Fatal Familial Insomnia; Progressive Supranuclear Palsy; Multiple System Atrophy; Stiff-Person Syndrome.

**Neuromuscular & Peripheral Nerve Diseases** Duchenne Muscular Dystrophy; Becker Muscular Dystrophy; Facioscapulohumeral Muscular Dystrophy; Limb-Girdle Muscular Dystrophies; Congenital Muscular Dystrophies; Myasthenia Gravis; Charcot–Marie–Tooth Disease; Amyloid Polyneuropathy.

**Inborn Errors of Metabolism & Storage Diseases** Gaucher Disease; Fabry Disease; Pompe Disease; Tay–Sachs Disease; Niemann–Pick Disease; Krabbe Disease; Metachromatic Leukodystrophy; X-Linked Adrenoleukodystrophy; Alkaptonuria; Homocystinuria; Phenylketonuria; Maple Syrup Urine Disease; Von Gierke Disease (GSD I); Mucopolysaccharidoses (Hurler, Hunter, Sanfilippo Syndromes).

**Blood & Bone Marrow Diseases** Hemophilia A; Hemophilia B; Paroxysmal Nocturnal Hemoglobinuria (PNH); Fanconi Anemia; Diamond–Blackfan Anemia; Pure Red Cell Aplasia; Thrombotic Thrombocytopenic Purpura (TTP); Atypical Hemolytic Uremic Syndrome; Cryoglobulinemia; Castleman Disease.

**Immune, Autoinflammatory & Histiocytic Diseases** Severe Combined Immunodeficiency (SCID); Common Variable Immunodeficiency; Chronic Granulomatous Disease; Wiskott–Aldrich Syndrome; Familial Mediterranean Fever; Cryopyrin-Associated Periodic Syndromes (CAPS); Hemophagocytic Lymphohistiocytosis (HLH); Histiocytosis X (Langerhans Cell Histiocytosis).

**Rheumatologic & Autoimmune Diseases** Systemic Sclerosis (Scleroderma); Granulomatosis with Polyangiitis; Microscopic Polyangiitis; Polyarteritis Nodosa; Behçet Disease; Takayasu Arteritis; Eosinophilic Granulomatosis with Polyangiitis (Churg–Strauss); Relapsing Polychondritis; Adult-Onset Still Disease.

**Connective Tissue & Dermatological Genetic Diseases** Ehlers–Danlos Syndromes (Vascular and Classic Types); Marfan Syndrome; Loeys–Dietz Syndrome; Osteogenesis Imperfecta; Pseudoxanthoma Elasticum; Epidermolysis Bullosa; Ichthyosis; Xeroderma Pigmentosum; Pemphigus Vulgaris; Bullous Pemphigoid.

**Endocrine Diseases** Addison Disease; Congenital Adrenal Hyperplasia; Acromegaly; Cushing Disease (Pituitary-Dependent); Hypoparathyroidism; Multiple Endocrine Neoplasia (MEN 1 and 2); Pheochromocytoma; Autoimmune Polyglandular Syndromes.

---

**Hepatic & Biliary Diseases** Wilson Disease; Alpha-1 Antitrypsin Deficiency; Primary Biliary Cholangitis; Primary Sclerosing Cholangitis; Biliary Atresia; Progressive Familial Intrahepatic Cholestasis; Caroli Disease.

**Renal & Urinary Diseases** Alport Syndrome; Cystinosis; Primary Hyperoxaluria; Bartter Syndrome; Gitelman Syndrome; Polycystic Kidney Disease (Autosomal Recessive Type); Medullary Cystic Kidney Disease.

**Pulmonary Diseases** Idiopathic Pulmonary Fibrosis; Pulmonary Arterial Hypertension; Lymphangioleiomyomatosis (LAM); Pulmonary Alveolar Proteinosis; Primary Ciliary Dyskinesia.

**Cardiovascular Diseases** Brugada Syndrome; Long QT Syndrome; Short QT Syndrome; Arrhythmogenic Right Ventricular Cardiomyopathy; Fibromuscular Dysplasia; Takayasu Arteritis.

**Gastrointestinal Diseases** Hirschsprung Disease; Achalasia; Eosinophilic Esophagitis; Short Bowel Syndrome; Hereditary Pancreatitis; Microvillus Inclusion Disease.

**Eye & Visual Pathway Diseases** Retinitis Pigmentosa; Stargardt Disease; Leber Hereditary Optic Neuropathy; Leber Congenital Amaurosis; Retinoblastoma; Aniridia; Best Disease.

**Bone, Skeletal & Cartilage Diseases** Achondroplasia; Fibrodysplasia Ossificans Progressiva (FOP); Hypophosphatasia; Osteopetrosis (Marble Bone Disease); Cleidocranial Dysplasia; Thanatophoric Dysplasia.

**Rare Infectious & Prion Diseases** Creutzfeldt–Jakob Disease; Variant CJD; Kuru; Whipple Disease; Progressive Multifocal Leukoencephalopathy (PML); Ebola Virus Disease; Marburg Virus Disease; Lassa Fever.

**Rare Cancers & Neoplasms** Malignant Mesothelioma; Gastrointestinal Stromal Tumor (GIST); Neuroendocrine Tumors; Ewing Sarcoma; Osteosarcoma; Retinoblastoma; Uveal Melanoma; Adrenocortical Carcinoma; Thymoma; Merkel Cell Carcinoma; Kaposi Sarcoma; Burkitt Lymphoma.

**Multisystem & Inflammatory Diseases** Erdheim–Chester Disease; POEMS Syndrome; VEXAS Syndrome; IgG4-Related Disease; Systemic Amyloidosis; Castleman Disease.



# Genetic Diseases

*Scope:* This reference-oriented list organizes human genetic diseases by mode of inheritance, cytogenetic classification, and molecular pathophysiological mechanism. Genetic diseases result from alterations in DNA sequences, ranging from single-nucleotide variants to gross chromosomal rearrangements and non-Mendelian mechanisms, providing a focused reference complementing Common Diseases and Rare Diseases.

**Autosomal Dominant Diseases** Huntington Disease; Marfan Syndrome; Neurofibromatosis Type 1 (von Recklinghausen Disease); Neurofibromatosis Type 2; Familial Hypercholesterolemia; Polycystic Kidney Disease (Autosomal Dominant, ADPKD); Achondroplasia; Tuberous Sclerosis Complex; Hereditary Spherocytosis; Von Hippel–Lindau Syndrome; Li–Fraumeni Syndrome; Multiple Endocrine Neoplasia (MEN 1 and 2); Myotonic Dystrophy (Type 1 and 2); Hereditary Non-Polyposis Colorectal Cancer (Lynch Syndrome); Familial Adenomatous Polyposis (FAP); Osteogenesis Imperfecta (Type I); Ehlers–Danlos Syndrome (Hypermobile and Classic Types).

**Autosomal Recessive Diseases** Cystic Fibrosis; Sickle-Cell Disease; Beta-Thalassemia; Alpha-Thalassemia; Phenylketonuria (PKU); Gaucher Disease; Tay–Sachs Disease; Niemann–Pick Disease; Wilson Disease; Hereditary Hemochromatosis; Alpha-1 Antitrypsin Deficiency; Spinal Muscular Atrophy (SMA); Friedreich Ataxia; Alkaptonuria; Homocystinuria; Maple Syrup Urine Disease; Congenital Adrenal Hyperplasia; Galactosemia; Glycogen Storage Diseases (von Gierke, Pompe, Cori); Ataxia–Telangiectasia; Fanconi Anemia; Xeroderma Pigmentosum; Bloom Syndrome; Ellis–van Creveld Syndrome; Kartagener Syndrome.

**X-Linked Recessive Diseases** Duchenne Muscular Dystrophy; Becker Muscular Dystrophy; Hemophilia A (Factor VIII Deficiency); Hemophilia B (Factor IX Deficiency); Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency; Red-Green Color Blindness; Fragile X Syndrome; Lesch–Nyhan Syndrome; Wiskott–Aldrich Syndrome; X-Linked Agammaglobulinemia (Bruton Tyrosine Kinase Deficiency); X-Linked Adrenoleukodystrophy; Hunter Syndrome (MPS II); Fabry Disease; Menkes Disease; Alport Syndrome (X-Linked Type); Chronic Granulomatous Disease (X-Linked Form).

**X-Linked Dominant Diseases** Rett Syndrome (MECP2 Mutation); Incontinentia Pigmenti; X-Linked Hypophosphatemic Rickets; Aicardi Syndrome; Focal Dermal Hypoplasia (Goltz Syndrome); Fragile X-Associated Tremor/Ataxia Syndrome (FXTAS).

**Y-Linked (Holandric) Conditions** Y-Chromosome Infertility (AZF Deletions); SRY Gene Deletions/Mutations; Swyer Syndrome (46,XY Gonadal Dysgenesis); Retinitis Pigmentosa (Y-Linked Form); Hypertrichosis Pinnarum.

**Mitochondrial (Maternal) Genetic Diseases** Leber Hereditary Optic Neuropathy (LHON); Mitochondrial Encephalomyopathy, Lactic Acidosis, and Stroke-Like Episodes (MELAS); Myoclonic Epilepsy with Ragged Red Fibers (MERRF); Neuropathy, Ataxia, and Retinitis Pigmentosa (NARP); Leigh Syndrome (Subacute Necrotizing Encephalomyelopathy); Kearns–Sayre Syndrome; Pearson Syndrome; Chronic Progressive External Ophthalmoplegia (CPEO).

**Chromosomal Aneuploidies (Autosomal)** Trisomy 21 (Down Syndrome); Trisomy 18 (Edwards Syndrome); Trisomy 13 (Patau Syndrome); Trisomy 8 (Warkany Syndrome 2); Trisomy 9 Mosaicism.

---

**Sex Chromosome Aneuploidies** 45,X (Turner Syndrome); 47,XXY (Klinefelter Syndrome); 47,XXX (Triple X Syndrome); 47,XYY (Jacob Syndrome); 48,XXYY Syndrome; 48,XXXY Syndrome; Tetrasomy X.

**Microdeletion & Microduplication Syndromes** 22q11.2 Deletion Syndrome (DiGeorge Syndrome, Velocardio-facial Syndrome); 5p Deletion Syndrome (Cri-du-Chat Syndrome); 4p Deletion Syndrome (Wolf–Hirschhorn Syndrome); 7q11.23 Deletion Syndrome (Williams Syndrome); 17p11.2 Deletion Syndrome (Smith–Magenis Syndrome); 17p11.2 Duplication Syndrome (Potocki–Lupski Syndrome); 1p36 Deletion Syndrome; 15q11–q13 Deletion; Rubinstein–Taybi Syndrome.

**Genomic Imprinting Disorders** Prader–Willi Syndrome (Maternal Imprinting / Paternal 15q11–q13 Loss); Angelman Syndrome (Paternal Imprinting / Maternal UBE3A Loss); Beckwith–Wiedemann Syndrome (11p15.5 Alteration); Silver–Russell Syndrome; Pseudohypoparathyroidism (Albright Hereditary Osteodystrophy); Uniparental Disomy Syndromes.

**Trinucleotide Repeat Expansion Diseases** Huntington Disease (CAG Repeat); Fragile X Syndrome (CGG Repeat); Myotonic Dystrophy Type 1 (CTG Repeat); Friedreich Ataxia (GAA Repeat); Spinocerebellar Ataxias (SCA Types 1, 2, 3, 6, 7); Dentatorubral–Pallidoluysian Atrophy (DRPLA); Spinal and Bulbar Muscular Atrophy (Kennedy Disease).

**Lysosomal Storage Diseases** Gaucher Disease (Glucocerebrosidase Deficiency); Tay–Sachs Disease (Hexosaminidase A Deficiency); Niemann–Pick Disease (Types A, B, C); Fabry Disease (Alpha-Galactosidase A Deficiency); Pompe Disease (Acid Alpha-Glucosidase Deficiency); Krabbe Disease; Metachromatic Leukodystrophy; Mucopolysaccharidoses (Hurler, Scheie, Hunter, Sanfilippo, Morquio, Maroteaux–Lamy, Sly Syndromes); Fucosidosis; Sialidosis.

**Connective Tissue Genetic Disorders** Marfan Syndrome (FBN1 Mutation); Ehlers–Danlos Syndromes; Loeys–Dietz Syndrome; Osteogenesis Imperfecta (Brittle Bone Disease); Cutis Laxa; Stickler Syndrome; Pseudoxanthoma Elasticum; Alport Syndrome.

**Hereditary Cancer Predisposition Syndromes** Hereditary Breast and Ovarian Cancer Syndrome (BRCA1, BRCA2); Lynch Syndrome (MLH1, MSH2, MSH6, PMS2); Familial Adenomatous Polyposis (APC Mutation); Li–Fraumeni Syndrome (TP53 Mutation); Multiple Endocrine Neoplasia Types 1, 2A, 2B; Von Hippel–Lindau Disease; Cowden Syndrome (PTEN Hamartoma Tumor Syndrome); Peutz–Jeghers Syndrome; Nevoid Basal Cell Carcinoma Syndrome (Gorlin Syndrome); Retinoblastoma (RB1 Mutation); Wilms Tumor Predisposition (WT1).

# Inherited Diseases

*Classification:* The first groups show the most common inheritance pattern. The later clinical categories overlap, and several conditions have rare alternate patterns or genetically heterogeneous forms.

**Autosomal Dominant** Achondroplasia; Autosomal Dominant Polycystic Kidney Disease (ADPKD); Familial Adenomatous Polyposis; Familial Hypercholesterolemia; Hereditary Angioedema (most forms); Huntington Disease; Marfan Syndrome; Multiple Endocrine Neoplasia; Neurofibromatosis Type 1; Neurofibromatosis Type 2; Tuberous Sclerosis Complex; Von Hippel–Lindau Disease.

**Autosomal Recessive** Alpha-Thalassemia; Beta-Mannosidosis; Beta-Thalassemia; Congenital Adrenal Hyperplasia; Cystic Fibrosis; Familial Mediterranean Fever; Gaucher Disease; Hemochromatosis; Phenylketonuria (PKU); Sickle-Cell Disease; Spinal Muscular Atrophy; Tay–Sachs Disease; Wilson Disease.

**X-Linked** Becker Muscular Dystrophy; Duchenne Muscular Dystrophy; Fragile X Syndrome; Glucose-6-Phosphate Dehydrogenase (G6PD) Deficiency; Hemophilia A; Hemophilia B; X-Linked Agammaglobulinemia.

**Variable or Genetically Heterogeneous** Alport Syndrome (X-linked, autosomal dominant, or autosomal recessive); Charcot–Marie–Tooth Disease (multiple inheritance patterns); Hereditary Spherocytosis (usually autosomal dominant, sometimes autosomal recessive); Retinitis Pigmentosa (autosomal dominant, autosomal recessive, X-linked, or mitochondrial); Von Willebrand Disease (usually autosomal dominant, with some autosomal recessive forms).

**Mitochondrial** MELAS; MERRF; Leber Hereditary Optic Neuropathy (LHON).

**Inherited Metabolic Disorders** Maple Syrup Urine Disease; Galactosemia; Homocystinuria; Glycogen Storage Diseases; Lysosomal Storage Diseases; Urea-Cycle Disorders.

**Inherited Neurological or Neuromuscular Disorders** Charcot–Marie–Tooth Disease; Spinal Muscular Atrophy; Hereditary Spastic Paraplegia; Friedreich Ataxia.

**Inherited Blood Disorders** Sickle-Cell Disease; Alpha-Thalassemia; Beta-Thalassemia; Hereditary Spherocytosis; Hemophilia A; Hemophilia B; some forms of Von Willebrand Disease.

**Inherited Connective-Tissue or Skeletal Disorders** Osteogenesis Imperfecta; Ehlers–Danlos Syndromes; Marfan Syndrome; Achondroplasia.

**Inherited Renal Disorders** Alport Syndrome; Autosomal Dominant Polycystic Kidney Disease (ADPKD); Autosomal Recessive Polycystic Kidney Disease (ARPKD); Cystinuria.

**Inherited Eye Disorders** Retinitis Pigmentosa; Stargardt Disease; congenital color-vision deficiencies; some forms of congenital cataract.

**Hereditary Cancer Syndromes** Lynch Syndrome; Familial Adenomatous Polyposis; Hereditary Breast and Ovarian Cancer Syndrome; Multiple Endocrine Neoplasia (MEN) Syndromes; Li–Fraumeni Syndrome; Von Hippel–Lindau Disease.



# Common Symptoms

*Scope:* For a medical dictionary or disease-reference project, symptoms are best organized by body system, because a flat alphabetical list quickly becomes difficult to use. Here is a broad list of common medical symptoms.

**General / Systemic** Fatigue; Weakness; Malaise; Fever; Chills; Night Sweats; Excessive Sweating; Hot Flashes; Flushing; Loss of Appetite; Increased Appetite; Unexplained Weight Loss; Unexplained Weight Gain; Excessive Thirst; Dehydration; Exercise Intolerance; Lethargy; Generalized Body Aches.

**Pain** Headache; Chest Pain; Pleuritic Chest Pain; Abdominal Pain; Cramping or Colicky Pain; Back Pain; Neck Pain; Joint Pain; Muscle Pain; Bone Pain; Pelvic Pain; Flank Pain; Facial Pain; Eye Pain; Earache; Sore Throat; Toothache; Painful Urination; Rectal Pain; Burning Pain; Radiating Pain; Throbbing Pain.

**Neurological** Dizziness; Vertigo; Fainting; Light-Headedness; Orthostatic Lightheadedness; Numbness; Tingling; Burning Sensation; Electric Shock Sensations; Tremor; Muscle Weakness; Paralysis; Seizures; Confusion; Memory Problems; Forgetfulness; Difficulty Concentrating; Brain Fog; Loss of Balance; Poor Coordination; Slurred Speech; Altered Consciousness; Aura.

**Eye / Vision** Blurred Vision; Double Vision; Loss of Vision; Reduced Visual Acuity; Night Blindness; Eye Redness; Watery Eyes; Dry Eyes; Gritty or Foreign-Body Sensation; Eye Strain; Itchy Eyes; Photophobia; Floaters; Flashes of Light; Halos Around Lights; Sensitivity to Glare.

**Ear / Hearing** Hearing Loss; Tinnitus; Ear Discharge; Ear Fullness; Ear Pain; Sensitivity to Sound; Popping or Clicking in Ears.

**Nose / Sinus** Runny Nose; Nasal Congestion; Sneezing; Nosebleed; Loss of Smell; Reduced Smell; Sinus Pressure; Postnasal Drip; Frequent Throat Clearing.

**Mouth / Throat** Dry Mouth; Mouth Ulcers; Bad Breath; Altered Taste; Loss of Taste; Bleeding Gums; Hoarseness; Sore Throat; Difficulty Swallowing; Painful Swallowing; Lump in Throat; Burning Mouth Sensation.

**Respiratory** Cough; Dry Cough; Productive Cough; Coughing up Blood; Shortness of Breath; Shortness of Breath on Exertion; Wheezing; Rapid Breathing; Noisy Breathing; Sputum Production; Chest Tightness; Air Hunger; Choking Sensation; Inability to Take a Deep Breath.

**Cardiovascular** Palpitations; Rapid Heartbeat; Slow Heartbeat; Irregular Heartbeat; Heart Fluttering; Chest Pressure; Chest Squeezing; Shortness of Breath on Exertion; Orthopnea; Paroxysmal Nocturnal Dyspnea; Fainting; Ankle or Leg Swelling; Cold Extremities; Intermittent Claudication; Cyanosis.

**Gastrointestinal** Nausea; Vomiting; Vomiting Blood; Heartburn; Acid Regurgitation; Indigestion; Bloating; Abdominal Distension; Abdominal Cramps; Excessive Gas; Belching; Difficulty Passing Gas; Diarrhea; Constipation; Straining at Stool; Tenesmus; Blood in Stool; Black or Tarry Stool; Mucus in Stool; Fecal Urgency; Fecal Incontinence; Early Satiety; Anal Itching.

**Urinary** Frequent Urination; Urinary Urgency; Painful Urination; Excessive Urination; Reduced Urination; Inability to Urinate; Urinary Hesitancy; Post-Void Dribbling; Nighttime Urination; Blood in Urine; Urinary Incontinence; Weak Urinary Stream; Bladder Spasms; Foamy Urine.

---

**Musculoskeletal** Joint Stiffness; Morning Stiffness; Joint Swelling; Reduced Range of Motion; Joint Clicking or Grating; Joint Instability or Giving Way; Muscle Cramps; Muscle Spasms; Muscle Stiffness; Muscle Twitches; Restless Legs Sensation; Limb Pain; Difficulty Walking; Limping.

**Skin** Itching; Rash; Redness; Dry Skin; Scaling or Flaking; Skin Discoloration; Bruising; Easy Bruisability; Swelling; Hives; Blisters; Skin Ulcers; Lumps; Excessive Sweating; Reduced Sweating; Hair Loss; Hair Shedding; Brittle Nails; Crawling Sensation on Skin; Skin Tenderness.

**Endocrine / Metabolic** Excessive Thirst; Excessive Hunger; Frequent Urination; Heat Intolerance; Cold Intolerance; Unexplained Weight Change; Excessive Sweating; Hot Flashes.

**Sleep-Related** Insomnia; Difficulty Falling Asleep; Difficulty Staying Asleep; Frequent Awakening; Early Morning Awakening; Non-Restorative Sleep; Excessive Daytime Sleepiness; Snoring; Gasping in Sleep; Restless Sleep; Nightmares; Sleepwalking.

**Psychological / Behavioral** Anxiety; Depressed Mood; Irritability; Agitation; Mood Changes; Loss of Interest or Pleasure; Poor Concentration; Racing Thoughts; Restlessness; Sleep Disturbance; Appetite Changes; Feeling Detached.

**Reproductive / Gynecological** Pelvic Pain; Painful Menstruation; Heavy Menstrual Bleeding; Irregular Menstruation; Missed Periods; Vaginal Discharge; Vaginal Bleeding; Bleeding Between Periods; Vaginal Itching; Painful Intercourse; Hot Flashes.

**Male Reproductive** Erectile Dysfunction; Reduced Libido; Testicular Pain; Testicular Swelling; Penile Discharge; Painful Ejaculation; Scrotal Heaviness.

**Blood / Lymphatic** Easy Bruising; Prolonged Bleeding; Pale Skin; Swollen Lymph Nodes; Recurrent Infections; Petechiae.

**Allergic** Itching; Sneezing; Watery Eyes; Nasal Congestion; Hives; Facial Swelling; Lip or Tongue Swelling; Wheezing; Shortness of Breath; Anaphylactic Reactions Sensation.

*Clinical Note:* For a comprehensive medical dictionary, it is essential to distinguish symptoms (subjective experiences reported by the patient, such as nausea or pain) from clinical signs (objectively observed physical findings detected on examination, such as jaundice, edema, or tachycardia). Mixing them is common in general health lists, but an authoritative reference maintains clear separation among Symptoms, Clinical Signs, Syndromes, and Diseases/Disorders categories.



# Clinical Signs

*Scope:* A clinical sign is an objective abnormality that can be observed, measured, elicited, or detected by a clinician, unlike a symptom, which is experienced and reported by the patient. For a medical dictionary, the following system-based classification is useful.

**General / Vital Signs** Fever; Hypothermia; Tachycardia; Bradycardia; Tachypnea; Bradypnea; Hypertension; Hypotension; Orthostatic Hypotension; Pulsus Paradoxus; Wide Pulse Pressure; Narrow Pulse Pressure; Hypoxemia; Central Cyanosis; Peripheral Cyanosis; Altered Level of Consciousness; Cachexia; Pallor; Jaundice; Dehydration; Pitting Edema; Non-Pitting Edema; Generalized Lymphadenopathy; Clubbing.

**Skin Signs** Erythema; Pallor; Cyanosis; Jaundice; Petechiae; Purpura; Ecchymosis; Spider Angiomas; Palmar Erythema; Splinter Hemorrhages; Osler Nodes; Janeway Lesions; Rash; Urticaria; Vesicles; Bullae; Pustules; Papules; Nodules; Ulcers; Scaling; Excoriations; Lichenification; Skin Thickening; Hyperpigmentation; Hypopigmentation; Alopecia; Diaphoresis; Reduced Skin Turgor.

**Head & Face Signs** Facial Asymmetry; Facial Edema; Ptosis; Proptosis; Moon Facies; Mask-Like Facies; Temporal Wasting; Kayser-Fleischer Rings; Arcus Senilis; Xanthelasma; Subconjunctival Hemorrhage.

**Eye Signs** Conjunctival Pallor; Conjunctival Injection; Scleral Icterus; Anisocoria; Nystagmus; Strabismus; Ptosis; Proptosis; Papilledema; Absent Pupillary Reflex; Abnormal Red Reflex; Corneal Opacity; Roth Spots; Argyll Robertson Pupil.

**Ear, Nose & Throat Signs** Tympanic Membrane Erythema; Tympanic Membrane Perforation; Otorrhea; Nasal Polyps; Epistaxis; Tonsillar Enlargement; Tonsillar Exudate; Pharyngeal Erythema; Oral Ulcers; Oral Thrush; Strawberry Tongue; Coated Tongue; Koplik Spots; Gingival Hyperplasia; Peritonsillar Bulge; Deviated Uvula.

**Neck Signs** Cervical Lymphadenopathy; Goiter; Thyroid Nodule; Neck Stiffness; Elevated Jugular Venous Pressure; Carotid Bruit; Tracheal Deviation; Pemberton Sign.

**Cardiovascular Signs** Tachycardia; Bradycardia; Irregular Pulse; Weak Pulse; Bounding Pulse; Pulsus Alternans; Corrigan Pulse; Water-Hammer Pulse; Pulse Deficit; Displaced Apex Beat; Heart Murmur; Systolic Murmur; Diastolic Murmur; Gallop Rhythm; Third Heart Sound; Fourth Heart Sound; Pericardial Friction Rub; Cardiac Thrill; Ventricular Heave; Elevated Jugular Venous Pressure; Peripheral Edema; Delayed Capillary Refill.

**Respiratory Signs** Tachypnea; Bradypnea; Apnea; Cyanosis; Accessory-Muscle Use; Nasal Flaring; Chest Retractions; Barrel Chest; Pectus Excavatum; Pectus Carinatum; Asymmetric Chest Expansion; Subcutaneous Emphysema; Reduced Air Entry; Wheezing; Crackles; Rhonchi; Stridor; Bronchial Breath Sounds; Dullness to Percussion; Hyperresonance; Pleural Friction Rub.

**Abdominal & Gastrointestinal Signs** Abdominal Distension; Abdominal Tenderness; Guarding; Rigidity; Rebound Tenderness; Shifting Dullness; Fluid Wave; Caput Medusae; Palpable Abdominal Mass; Hepatomegaly; Splenomegaly; Ascites; Succussion Splash; Abnormal Bowel Sounds; Abdominal Bruit; Visible Peristalsis; Rectal Bleeding; Melena.

---

**Neurological Signs** Altered Consciousness; Disorientation; Abnormal Gait; Festinating Gait; Spastic Hemiplegic Gait; Ataxia; Tremor; Asterixis; Fasciculations; Muscle Wasting; Pronator Drift; Cogwheel Rigidity; Lead-Pipe Rigidity; Abnormal Muscle Tone; Hyperreflexia; Hyporeflexia; Areflexia; Pathological Reflexes; Clonus; Dysmetria; Dysdiadochokinesia; Sensory Deficit; Motor Weakness on Examination; Cranial Nerve Deficit; Nystagmus; Neck Rigidity.

**Musculoskeletal Signs** Joint Swelling; Joint Deformity; Joint Effusion; Restricted Range of Motion; Crepitus; Muscle Wasting; Muscle Spasm; Heberden Nodes; Bouchard Nodes; Swan-Neck Deformity; Boutonniere Deformity; Baker Cyst; Limb Deformity; Abnormal Posture; Scoliosis; Kyphosis; Lordosis; Abnormal Gait.

**Renal & Urinary Signs** Bladder Distension; Costovertebral-Angle Tenderness; Peripheral Edema; Hematuria on Testing; Proteinuria; Oliguria; Anuria.

**Endocrine Signs** Goiter; Exophthalmos; Tremor; Abnormal Hair Distribution; Hirsutism; Striae; Acanthosis Nigricans; Moon Facies; Buffalo Hump; Peripheral Edema.

**Hematological Signs** Pallor; Petechiae; Purpura; Ecchymoses; Generalized Lymphadenopathy; Splenomegaly; Hepatomegaly; Active Bleeding.

**Peripheral Vascular Signs** Peripheral Edema; Pitting Edema; Varicose Veins; Absent or Diminished Peripheral Pulses; Cool Extremities; Delayed Capillary Refill; Venous Ulcers; Arterial Ulcers; Peripheral Cyanosis.

**Gynecological Signs** Abnormal Vaginal Bleeding on Examination; Vaginal or Cervical Discharge; Cervical Lesions; Pelvic Mass; Uterine Enlargement; Adnexal Mass; Cervical-Motion Tenderness; Chandelier Sign.

**Male Genital Signs** Testicular Swelling; Testicular Mass; Scrotal Edema; Penile Discharge; Genital Ulceration; Phimosis; Hydrocele; Varicocele.

**Pediatric & Neonatal Signs** Poor Weight Gain on Growth Charts; Growth Retardation; Bulging Fontanelle; Sunken Fontanelle; Neonatal Jaundice; Central Cyanosis; Chest Retractions; Abnormal Primitive Reflexes; Gowers Sign.

**Named Clinical Signs** Babinski Sign; Murphy Sign; Kernig Sign; Brudzinski Sign; Rovsing Sign; Cullen Sign; Grey Turner Sign; Trousseau Sign; Chvostek Sign; Romberg Sign; Hoffmann Sign; Courvoisier Sign; Kussmaul Sign; Battle Sign; Nikolsky Sign; Auspitz Sign; Ortolani Sign; Barlow Sign; Trendelenburg Sign; McBurney Sign; Tinel Sign; Phalen Sign; Lhermitte Sign; Blumberg Sign; Kehr Sign; Hamman Sign; Homans Sign; Psoas Sign; Obturator Sign; Lachman Sign; Anterior Drawer Sign.

*Clinical Note:* A comprehensive medical reference can contain 500–1,000+ clinical signs, encompassing physical examination findings, named signs, dermatological signs, radiological signs, and laboratory markers.

# Common Congenital Anomalies

*Scope:* A congenital anomaly is a structural abnormality present at birth, whether genetic, chromosomal, or environmental in origin. This list complements the appendices of Common Diseases, Common Disorders, and Common Syndromes.

**Orofacial & Head Anomalies** Cleft Lip; Cleft Palate.

**Neural Tube & Brain Anomalies** Spina Bifida; Anencephaly; Encephalocele; Hydrocephalus.

**Thorax & Abdominal Wall Anomalies** Congenital Diaphragmatic Hernia; Omphalocele; Gastroschisis.

**Digestive System Anomalies** Imperforate Anus; Hirschsprung Disease.

**Urinary & Genital Anomalies** Hypospadias; Epispadias.

**Limb & Skeletal Anomalies** Clubfoot; Developmental Dysplasia of the Hip; Polydactyly; Syndactyly.

**Eye Anomalies** Congenital Cataract.

**Heart & Great-Vessel Anomalies** Congenital Heart Defects; Ventricular Septal Defect; Atrial Septal Defect; Patent Ductus Arteriosus; Tetralogy of Fallot; Coarctation of the Aorta.



# Common Findings, Lesions & Conditions

*Scope:* This companion list collects clinical findings, structural lesions, symptom patterns, and obstetric events that are not diseases in themselves but are frequently encountered in clinical work, imaging, and laboratory testing. It complements the appendices of Common Diseases, Common Disorders, and Common Syndromes.

**Cardiovascular & Respiratory Findings** Pericardial Effusion; Pleural Effusion; Pneumothorax; Hemothorax; Atelectasis.

**Abdominal & Pancreatic Findings** Esophageal Varices; Gallstones; Biliary Obstruction; Cholestasis; Pancreatic Pseudocyst; Pancreatic Cysts; Pancreatic Necrosis.

**Renal & Urinary Findings** Kidney Stones; Hydronephrosis.

**Head, Neck & Airway Findings** Thyroid Nodules; Nasal Polyps; Deviated Nasal Septum; Vocal-Cord Nodules; Earwax Impaction; Tympanic Membrane Perforation.

**Oral & Dental Findings** Tooth Erosion; Impacted Teeth; Malocclusion; Geographic Tongue.

**Eye & Skin Findings** Chalazion; Stye; Keloids; Ingrown Toenail.

**Genital & Reproductive Findings** Ovarian Cysts; Bartholin Cyst; Varicocele; Hydrocele; Spermatocoele.

**Pain & Symptom Conditions** Low-Back Pain; Sciatica.

**Metabolic Risk & Allergic States** Prediabetes; Anaphylaxis.

**Obstetric & Labor Events** Ectopic Pregnancy; Miscarriage; Recurrent Pregnancy Loss; Placenta Previa; Placental Abruption; Placenta Accreta Spectrum; Premature Rupture of Membranes; Preterm Labor; Fetal Growth Restriction; Obstructed Labor; Postpartum Hemorrhage.





# Common Injuries, Poisoning & Environmental Conditions

*Scope:* This companion list covers injuries (trauma), poisoning and envenomation, and adverse effects of environmental exposure and extremes. These are events or conditions rather than diseases; it complements the appendices of Common Diseases, Common Disorders, and Common Syndromes.

**Bone, Joint & Muscle Injuries** Fracture; Dislocation; Sprain; Strain.

**Skin & Soft-Tissue Injuries** Contusion; Laceration; Abrasion; Crush Injury.

**Nervous-System Injuries** Traumatic Brain Injury; Concussion; Spinal-Cord Injury.

**Thermal & Electrical Injuries** Burns; Electrical Injury; Hypothermia; Frostbite; Heat Exhaustion; Heatstroke.

**Environmental Conditions** Drowning; Dehydration; Altitude Sickness; Motion Sickness.

**Poisoning & Envenomation** Carbon Monoxide Poisoning; Lead Poisoning; Mercury Poisoning; Arsenic Poisoning; Pesticide Poisoning; Snakebite; Scorpion Sting.

**Drug & Alcohol Overdose** Drug Overdose; Alcohol Poisoning.



# Conditions That May Be Asymptomatic

*Scope:* This reference list groups diseases, infections, and other conditions that may be present without noticeable symptoms, particularly during early, latent, or compensated stages. Being asymptomatic is a clinical state or mode of presentation rather than a separate category of disease, so every condition here also belongs under its usual organ-system or specialty heading, and “May be asymptomatic” is best treated as an attribute or tag in addition to that classification. Many of these conditions cause symptoms in some people and none in others, and some come to attention only through screening or when a complication appears. The list is a curated, high-importance reference rather than a prevalence ranking, and some conditions appear under more than one heading where they are clinically relevant to several systems. Individual symptoms, signs, and laboratory findings are defined as dictionary entries rather than listed here.

**Important distinction:** “Asymptomatic disease” is not a distinct disease; it describes how a disease presents. Hypertension is a cardiovascular disease that is often asymptomatic, chlamydia is an infection that is frequently asymptomatic, osteoporosis is a skeletal disease that is typically asymptomatic until a fracture, and open-angle glaucoma is an eye disease that is often asymptomatic in its early stages.

**Cardiovascular** Hypertension; Atherosclerosis; Coronary Artery Disease; Peripheral Artery Disease; Some Arrhythmias; Early Valvular Heart Disease

**Metabolic & Endocrine** Prediabetes; Early Type 2 Diabetes; Dyslipidemia; Subclinical Hypothyroidism; Subclinical Hyperthyroidism; Hyperuricemia

**Kidney & Urinary** Early Chronic Kidney Disease; Diabetic Kidney Disease; Mild Proteinuria; Microscopic Hematuria; Some Kidney Stones

**Liver** Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD); Chronic Hepatitis B; Chronic Hepatitis C; Early Cirrhosis; Hemochromatosis

**Infectious** HIV Infection During Clinical Latency; Hepatitis B; Hepatitis C; Chlamydia; Gonorrhea; HPV Infection; Latent Tuberculosis Infection; Some COVID-19 Infections

**Cancer & Precancer** Early Colorectal Cancer; Cervical Precancer; Early Cervical Cancer; Early Prostate Cancer; Early Kidney Cancer; Some Thyroid Cancers; Early Breast Cancer

**Gynecological** HPV Infection; Some Ovarian Cysts; Uterine Fibroids; Early Cervical Abnormalities; Some Sexually Transmitted Infections

**Male Reproductive** Early Prostate Cancer; Some Sexually Transmitted Infections; Varicocele; Some Cases of Benign Prostatic Enlargement

**Blood** Mild Anemia; Monoclonal Gammopathy of Undetermined Significance (MGUS); Early Chronic Leukemia; Mild Thrombocytopenia

**Bone** Osteoporosis; Osteopenia

**Eye** Open-Angle Glaucoma; Early Diabetic Retinopathy; Early Age-Related Macular Degeneration

---

**Neurological & Vascular** Silent Cerebral Infarction; Asymptomatic Carotid Stenosis; Some Unruptured Intracranial Aneurysms

**Gastrointestinal** Diverticulosis; Colon Polyps; Some Gallstones; Some Hiatal Hernias

**Gallbladder** Asymptomatic Gallstones (“Silent Gallstones”); Gallbladder Polyps

**Genetic** Hemochromatosis Before Organ Damage; Some Inherited Thrombophilias; Some Genetic Carrier States

**Parasitic & Latent Infections** Some Malaria Infections; Intestinal Parasitic Infections; Toxoplasmosis and Other Latent Infections

# Lay Language Medical Terms

| Lay term                | Medical term                       | Lay-language explanation                                                                         |
|-------------------------|------------------------------------|--------------------------------------------------------------------------------------------------|
| Belly area              | ABDOMEN/ABDOMINAL                  | The body cavity below the diaphragm containing the stomach, intestines, liver, and other organs. |
| Take in                 | ABSORB                             | To take up or take in a substance, such as fluid.                                                |
| Too much acid in blood  | ACIDOSIS                           | A condition in which the blood is more acidic than normal.                                       |
| Sharpness or clarity    | ACUITY                             | The clearness or keenness of a sense, especially vision.                                         |
| Recent or sudden        | ACUTE                              | New, recent, sudden, or requiring urgent attention.                                              |
| Swollen glands          | ADENOPATHY                         | Enlarged or swollen lymph nodes.                                                                 |
| Helpful added treatment | ADJUVANT                           | Something that supports or assists the main treatment.                                           |
| Added treatment         | ADJUVANT TREATMENT                 | Treatment given in addition to the main or standard treatment.                                   |
| Germ-killing drug       | ANTIBIOTIC                         | A drug used to kill or control bacteria and some other germs.                                    |
| Germ-killing treatment  | ANTIMICROBIAL                      | A substance or treatment that acts against microorganisms such as bacteria, fungi, or parasites. |
| Antiviral drug          | ANTIRETROVIRAL                     | A drug that slows or stops the growth of certain viruses.                                        |
| Side effect             | ADVERSE EFFECT                     | An unwanted or harmful response to a treatment or medicine.                                      |
| Walking                 | AMBULATE/AMBULATION/<br>AMBULATORY | To walk, or to be able to walk.                                                                  |
| Severe allergy          | ANAPHYLAXIS                        | A serious, potentially life-threatening allergic reaction.                                       |
| Low red-cell count      | ANEMIA                             | A reduced number of red blood cells or a low red-cell blood count.                               |
| Numbing medicine        | ANESTHETIC                         | A drug or agent that reduces or removes the feeling of pain, sometimes by causing sleep.         |
| Heart pain              | ANGINA                             | Chest pain caused by inadequate blood flow to the heart muscle.                                  |
| Heart pain              | ANGINA PECTORIS                    | Another medical term for chest pain caused by reduced blood flow to the heart.                   |
| Loss of appetite        | ANOREXIA                           | Lack of appetite or, in some contexts, a disorder involving refusal to eat.                      |

| Lay term                          | Medical term                 | Lay-language explanation                                                             |
|-----------------------------------|------------------------------|--------------------------------------------------------------------------------------|
| Inner elbow area                  | ANTECUBITAL                  | Relating to the inner side of the elbow or forearm.                                  |
| Protective protein                | ANTIBODY                     | A protein made by the body in response to a foreign substance.                       |
| Seizure-prevention drug           | ANTICONVULSANT               | A medicine used to prevent or control seizures.                                      |
| Cholesterol-lowering drug         | ANTILIPEMIC                  | A drug that lowers fat or cholesterol levels in the blood.                           |
| Cough medicine                    | ANTITUSSIVE                  | A medicine used to reduce coughing.                                                  |
| Irregular heartbeat               | ARRHYTHMIA                   | Any abnormal change in the normal heartbeat or rhythm.                               |
| Breathing material into the lungs | ASPIRATION                   | Entry of fluid, food, or other material into the lungs, often after vomiting.        |
| Laboratory test                   | ASSAY                        | A laboratory test used to measure or identify something.                             |
| Check or evaluate                 | ASSESS                       | To examine, measure, evaluate, or look at something carefully.                       |
| Asthma                            | ASTHMA                       | A lung disease in which narrowed or tightened airways make breathing difficult.      |
| Without symptoms                  | ASYMPTOMATIC                 | Having no noticeable symptoms.                                                       |
| Armpit                            | AXILLA                       | The armpit area.                                                                     |
| Not cancerous                     | BENIGN                       | Not malignant and generally not cancerous.                                           |
| Twice daily                       | BID                          | Prescription shorthand meaning twice a day.                                          |
| Carried or stuck together         | BINDING/BOUND                | Carried by something, or made to stick or join together.                             |
| Drug availability                 | BIOAVAILABILITY              | The amount or proportion of a drug that becomes available for the body to use.       |
| Series of blood tests             | BLOOD PROFILE                | A group or panel of blood tests.                                                     |
| Large dose at once                | BOLUS                        | A relatively large amount given all at one time.                                     |
| Amount of mineral in bone         | BONE MASS                    | The amount of calcium and other minerals in a given amount of bone.                  |
| Slow irregular heartbeats         | BRADYARRHYTHMIAS             | Slow heart rhythms that may also be irregular.                                       |
| Slow heartbeat                    | BRADYCARDIA                  | A slower-than-usual heartbeat.                                                       |
| Narrowed airways                  | BRONCHOSPASM                 | Breathing difficulty caused by tightening of the airway muscles.                     |
| Cancer-causing                    | CARCINOGENIC                 | Capable of causing cancer.                                                           |
| Cancer type                       | CARCINOMA                    | A cancer that begins in lining or covering cells of the body.                        |
| Related to the heart              | CARDIAC                      | Relating to the heart.                                                               |
| Resetting the heartbeat           | CARDIOVERSION                | Restoring a normal heart rhythm, often with a controlled electrical shock.           |
| Tube                              | CATHETER                     | A thin tube used to give or remove fluids, or to perform certain medical procedures. |
| Epidural tube                     | CATHETER                     | A catheter placed near the spinal cord to deliver anesthesia during surgery.         |
| Brain and spinal cord             | CENTRAL NERVOUS SYSTEM (CNS) | The brain and spinal cord.                                                           |
| Brain injury                      | CEREBRAL TRAUMA              | Damage to the brain caused by an injury.                                             |
| Stopping                          | CESSATION                    | The act of stopping something.                                                       |
| Coronary heart disease            | CHD                          | Disease of the blood vessels that supply the heart.                                  |
| Cancer drug treatment             | CHEMOTHERAPY                 | Treatment of disease, usually cancer, with chemical medicines.                       |



| Lay term                      | Medical term          | Lay-language explanation                                                                         |
|-------------------------------|-----------------------|--------------------------------------------------------------------------------------------------|
| Long-lasting                  | CHRONIC               | Continuing for a long time or ongoing.                                                           |
| Related to patient care       | CLINICAL              | Pertaining to medical care or the care of patients.                                              |
| Human research study          | CLINICAL TRIAL        | A research study involving human participants.                                                   |
| Unconscious state             | COMA                  | A prolonged state of unconsciousness in which a person cannot be awakened normally.              |
| All disease gone              | COMPLETE RESPONSE     | Total disappearance of the detectable disease being treated.                                     |
| Present from birth            | CONGENITAL            | Present before or at birth.                                                                      |
| Pink, irritated eye           | CONJUNCTIVITIS        | Redness and irritation of the thin membrane covering the eye.                                    |
| Treatment to keep improvement | CONSOLIDATION PHASE   | A treatment phase intended to make a remission last, usually after initial treatment.            |
| Comparison study              | CONTROLLED TRIAL      | A study comparing an experimental treatment with a standard or control treatment.                |
| Research partnership          | COOPERATIVE GROUP     | An association of institutions that work together on clinical research.                          |
| Related to heart arteries     | CORONARY              | Relating to the heart or the blood vessels that supply it.                                       |
| Computerized X-ray scan       | CT SCAN (CAT)         | A series of computer-processed X-ray images that show sections of the body.                      |
| Germ-growth test              | CULTURE               | A test that grows or detects organisms that may cause infection.                                 |
| Added together                | CUMULATIVE            | Added up from the beginning.                                                                     |
| Related to the skin           | CUTANEOUS             | Relating to the skin.                                                                            |
| Stroke                        | CVA                   | A stroke, also called a cerebrovascular accident.                                                |
| Related to the skin           | DERMATOLOGIC          | Pertaining to the skin.                                                                          |
| Lower blood-pressure number   | DIASTOLIC             | Relating to the lower blood-pressure number, when the heart relaxes.                             |
| Farther away                  | DISTAL                | Toward the end or away from the center of the body.                                              |
| Water pill                    | DIURETIC              | A drug that increases urination.                                                                 |
| Sound-wave blood-flow test    | DOPPLER               | A device or test using sound waves to assess blood flow.                                         |
| Masked study                  | DOUBLE BLIND          | A study in which neither the participants nor investigators know which treatment is being given. |
| Improper function             | DYSFUNCTION           | A state in which something does not work normally.                                               |
| Abnormal cells                | DYSPLASIA             | Abnormal changes in the size, shape, or organization of cells.                                   |
| Heart sound-wave test         | ECHOCARDIOGRAM        | An ultrasound test that creates pictures of the heart.                                           |
| Swelling from fluid           | EDEMA                 | Excess fluid collecting in body tissues.                                                         |
| Brain-wave test               | EEG                   | A recording of the brain's electrical activity.                                                  |
| Effectiveness                 | EFFICACY              | How well a treatment works under specified conditions.                                           |
| Heart electrical tracing      | ELECTROCARDIOGRAM     | A tracing of the heart's electrical activity, also called an ECG or EKG.                         |
| Mineral imbalance             | ELECTROLYTE IMBALANCE | An abnormal level or balance of minerals in the blood.                                           |
| Vomiting                      | EMESIS                | The act of vomiting.                                                                             |

| Lay term                   | Medical term          | Lay-language explanation                                                                         |
|----------------------------|-----------------------|--------------------------------------------------------------------------------------------------|
| Based on experience        | EMPIRIC               | Based on experience or likely cause rather than definitive test results.                         |
| Viewing inside the body    | ENDOSCOPIC            | Looking inside the body with a lighted tube or camera.                                           |
| Through the intestines     | ENTERAL               | Given through or involving the digestive tract.                                                  |
| Outside the spinal cord    | EPIDURAL              | Located outside the spinal cord covering, or medicine given into that space.                     |
| Remove completely          | ERADICATE             | To get rid of a disease, infection, or other problem.                                            |
| Examined                   | EVALUATED, ASSESSED   | Examined or checked for a medical condition.                                                     |
| Fast ethics review         | EXPEDITED REVIEW      | A rapid research-protocol review allowed for certain lower-risk studies.                         |
| Outside the body           | EXTERNAL              | Located or occurring outside the body.                                                           |
| Leak out                   | EXTRAVASATE           | To leak outside the intended area, such as from a blood vessel into nearby tissue.               |
| U.S. drug agency           | FDA                   | The U.S. Food and Drug Administration, which regulates medicines and medical products.           |
| Scar-like                  | FIBROUS               | Containing many fibers, often resembling scar tissue.                                            |
| Irregular quivering        | FIBRILLATION          | Very rapid, uncoordinated activity of the heart or another muscle.                               |
| Being put fully to sleep   | GENERAL ANESTHESIA    | Use of medicines to cause unconsciousness and prevent pain during a procedure.                   |
| Related to pregnancy       | GESTATIONAL           | Pertaining to pregnancy.                                                                         |
| Red-cell percentage        | HEMATOCRIT            | The proportion of blood made up of red blood cells.                                              |
| Bruise or blood collection | HEMATOMA              | A collection of blood under the skin or in a tissue, often appearing as a bruise.                |
| Blood-flow measurement     | HEMODYNAMIC MEASURING | Measuring blood flow and circulation.                                                            |
| Red-cell breakdown         | HEMOLYSIS             | Breakdown or destruction of red blood cells.                                                     |
| Heparin tube or port       | HEPARIN LOCK          | An intravenous access device kept open with a prescribed solution, sometimes containing heparin. |
| Liver tumor                | HEPATOMA              | A tumor of the liver, often referring to hepatocellular carcinoma.                               |
| Inherited disease          | HERITABLE DISEASE     | A disease that can be passed through genes to future children.                                   |
| Tissue-disease examination | HISTOPATHOLOGIC       | Relating to examination of tissues or cells for disease.                                         |
| Portable heart monitor     | HOLTER MONITOR        | A portable device that records heartbeats, usually for a day or longer.                          |
| High blood calcium         | HYPERCALCEMIA         | An abnormally high calcium level in the blood.                                                   |
| High blood potassium       | HYPERKALEMIA          | An abnormally high potassium level in the blood.                                                 |
| High blood sodium          | HYPERNATREMIA         | An abnormally high sodium level in the blood.                                                    |
| High blood pressure        | HYPERTENSION          | Blood pressure that is higher than recommended.                                                  |
| Low blood calcium          | HYPOCALCEMIA          | An abnormally low calcium level in the blood.                                                    |
| Low blood potassium        | HYPOKALEMIA           | An abnormally low potassium level in the blood.                                                  |
| Low blood sodium           | HYPONATREMIA          | An abnormally low sodium level in the blood.                                                     |
| Low blood pressure         | HYPOTENSION           | Blood pressure that is lower than expected.                                                      |
| Low blood oxygen           | HYPOXEMIA             | A low level of oxygen in the blood.                                                              |
| Low tissue oxygen          | HYPOXIA               | Too little oxygen reaching body tissues.                                                         |

| Lay term                         | Medical term                   | Lay-language explanation                                                                             |
|----------------------------------|--------------------------------|------------------------------------------------------------------------------------------------------|
| Removal of the uterus            | HYSTERECTOMY                   | Surgical removal of the uterus; the ovaries and other structures may or may not also be removed.     |
| Treatment-caused                 | IATROGENIC                     | Caused by medical treatment or a healthcare procedure.                                               |
| Device research permission       | IDE                            | Permission to study an unapproved medical device in people.                                          |
| Cause unknown                    | IDIOPATHIC                     | Having no known cause.                                                                               |
| Body protection                  | IMMUNITY                       | The body's defense against infection or other foreign substances.                                    |
| Antibody protein                 | IMMUNOGLOBIN                   | A protein involved in making antibodies and immune defense; the standard spelling is immunoglobulin. |
| Immune-suppressing               | IMMUNOSUPPRESSIVE              | A drug or treatment that reduces the body's immune response.                                         |
| Immune-based treatment           | IMMUNOTHERAPY                  | Treatment that uses or changes the immune system, often against cancer or immune disease.            |
| Abnormal function                | IMPAIRED FUNCTION              | Function that is reduced or not working normally.                                                    |
| Placed inside the body           | IMPLANTED                      | Surgically or otherwise placed inside the body.                                                      |
| New-drug research permission     | IND                            | Authorization to test an unapproved new drug in people.                                              |
| Hardening                        | INDURATION                     | An area that has become unusually firm or hard.                                                      |
| Left in place                    | INDWELLING                     | Remaining in a particular location, such as a catheter.                                              |
| Dead tissue from poor blood flow | INFARCT                        | Death of tissue caused by inadequate blood supply.                                                   |
| Disease that can spread          | INFECTIOUS DISEASE             | An illness caused by organisms that may spread between people or through other routes.               |
| Swelling and irritation          | INFLAMMATION                   | The body's response to injury or infection, often causing swelling, warmth, redness, or pain.        |
| Slow delivery into the body      | INFUSION                       | Slow delivery of a substance, usually through a vein or catheter.                                    |
| Taking by mouth                  | INGESTION                      | Eating, drinking, or taking something into the body by mouth.                                        |
| Antiviral immune protein         | INTERFERON                     | A medicine or natural protein that helps the body act against viruses.                               |
| On and off                       | INTERMITTENT                   | Occurring repeatedly with pauses between episodes.                                                   |
| Inside the body                  | INTERNAL                       | Located or occurring within the body.                                                                |
| Inside                           | INTERIOR                       | Located on the inside.                                                                               |
| Into a muscle                    | INTRAMUSCULAR                  | Given into a muscle.                                                                                 |
| Into the abdomen                 | INTRAPERITONEAL                | Given into or located within the abdominal cavity.                                                   |
| Into spinal fluid                | INTRATHECAL                    | Given into the fluid space around the spinal cord.                                                   |
| Into a vein                      | INTRAVENOUS (IV)               | Given through a vein.                                                                                |
| Into the bladder                 | INTRAVESICAL                   | Given into the bladder.                                                                              |
| Place a breathing tube           | INTUBATE                       | To place a tube into the airway.                                                                     |
| Breaking the skin                | INVASIVE PROCEDURE             | A procedure that punctures, opens, or cuts the skin or enters the body.                              |
| Unapproved-drug study            | INVESTIGATIONAL NEW DRUG (IND) | A new drug being studied before it has full approval for general use.                                |

| Lay term                      | Medical term           | Lay-language explanation                                                                     |
|-------------------------------|------------------------|----------------------------------------------------------------------------------------------|
| Unproven treatment method     | INVESTIGATIONAL METHOD | A treatment method still being studied and not established as standard care.                 |
| Reduced blood flow            | ISCHEMIA               | Too little blood and oxygen reaching a tissue, usually because blood flow is reduced.        |
| Open abdominal surgery        | LAPAROTOMY             | Surgery using an incision in the abdominal wall to reach the organs inside.                  |
| Abnormal area                 | LESION                 | A wound, injury, or area of abnormal tissue.                                                 |
| Very tired or sleepy          | LETHARGY               | Sleepiness, tiredness, or reduced energy.                                                    |
| Low white-cell count          | LEUKOPENIA             | A lower-than-normal number of white blood cells.                                             |
| Fat                           | LIPID                  | A fat or fat-like substance.                                                                 |
| Amount of fat                 | LIPID CONTENT          | The amount of fat in a substance, tissue, or blood sample.                                   |
| Cholesterol and fat panel     | LIPID PROFILE (PANEL)  | Blood tests measuring fats and cholesterol.                                                  |
| Numbing a small area          | LOCAL ANESTHESIA       | Numbing medicine used to prevent pain in a limited area.                                     |
| Limited to one area           | LOCALIZED              | Restricted to one area.                                                                      |
| Inner space                   | LUMEN                  | The hollow space inside a tube or organ, such as a blood vessel.                             |
| Lymph-vessel X-ray            | LYMPHANGIOGRAPHY       | An X-ray study of lymph vessels or nodes after contrast dye is injected.                     |
| Immune white blood cell       | LYMPHOCYTE             | A type of white blood cell that helps protect against infection.                             |
| Lymph-node cancer             | LYMPHOMA               | A cancer of lymph nodes or related lymphatic tissue.                                         |
| Feeling unwell                | MALAISE                | A vague feeling of illness or bodily discomfort.                                             |
| Not working correctly         | MALFUNCTION            | A condition in which a device, organ, or system does not work properly.                      |
| Cancerous growth              | MALIGNANCY             | A cancer or other tumor capable of growing and spreading.                                    |
| Brain tumor                   | MEDULLABLASTOMA        | A type of tumor that begins in the brain.                                                    |
| Abnormal red-cell development | MEGALOBLASTOSIS        | A change in blood-forming cells that produces unusually large, abnormal red-cell precursors. |
| Break down in the body        | METABOLIZE             | To chemically process or break down a substance in the body.                                 |
| Cancer spread                 | METASTASIS             | Spread of cancer cells from one body part to another.                                        |
| Antibiotic medicine           | METRONIDAZOLE          | A medicine used for certain bacterial and parasitic infections.                              |
| Heart attack                  | MI                     | Myocardial infarction, or death of heart muscle from blocked blood flow.                     |
| Small or slight               | MINIMAL                | Very small or slight.                                                                        |
| Reduce                        | MINIMIZE               | To reduce something as much as possible.                                                     |
| Check or watch                | MONITOR                | To observe, measure, or keep track of something over time.                                   |
| Ease of movement              | MOBILITY               | The ability to move easily.                                                                  |
| Illness or complication       | MORBIDITY              | Illness, disability, or an unwanted result of disease or treatment.                          |
| Death rate or death           | MORTALITY              | Death, or the rate of death in a population.                                                 |
| Ability to move               | MOTILITY               | The ability of a body part, cell, or organism to move.                                       |
| Magnetic body scan            | MRI                    | Pictures of the inside of the body made with magnets and radio waves rather than X-rays.     |

| Lay term                      | Medical term            | Lay-language explanation                                                                        |
|-------------------------------|-------------------------|-------------------------------------------------------------------------------------------------|
| Moist body lining             | MUCOSA, MUCOUS MEMBRANE | The moist lining of the digestive, respiratory, reproductive, and urinary passages.             |
| Muscle aches                  | MYALGIA                 | Pain or aching in the muscles.                                                                  |
| Related to heart muscle       | MYOCARDIAL              | Pertaining to the muscle of the heart.                                                          |
| Heart attack                  | MYOCARDIAL INFARCTION   | Death of heart muscle caused by inadequate blood flow.                                          |
| Nose-to-stomach tube          | NASOGASTRIC TUBE        | A tube passed through the nose into the stomach.                                                |
| U.S. health institute         | NCI                     | The National Cancer Institute.                                                                  |
| Tissue death                  | NECROSIS                | Death of cells or tissue.                                                                       |
| Tumor                         | NEOPLASIA/NEOPLASM      | An abnormal new growth that may be benign or malignant.                                         |
| Nerve-tissue cancer           | NEUROBLASTOMA           | A cancer that develops from immature nerve tissue.                                              |
| Related to the nervous system | NEUROLOGICAL            | Pertaining to the brain, spinal cord, or nerves.                                                |
| Low infection-fighting cells  | NEUTROPENIA             | A decrease in neutrophils, a major type of infection-fighting white blood cell.                 |
| U.S. health agency            | NIH                     | The National Institutes of Health.                                                              |
| Not entering the body         | NONINVASIVE             | Not breaking, cutting, or entering the skin or body.                                            |
| Hospital-acquired             | NOSOCOMIAL              | Acquired during a stay in a healthcare facility.                                                |
| Blockage                      | OCCLUSION               | Closing, blockage, or obstruction.                                                              |
| Cancer specialty              | ONCOLOGY                | The study and treatment of cancer and tumors.                                                   |
| Related to the eye            | OPHTHALMIC              | Pertaining to the eye.                                                                          |
| Best choice                   | OPTIMAL                 | Best, most favorable, or most suitable.                                                         |
| By mouth                      | ORAL ADMINISTRATION     | Giving a medicine through the mouth.                                                            |
| Related to bones              | ORTHOPEDIC              | Pertaining to bones, joints, muscles, or related structures.                                    |
| Rare dense-bone disorder      | OSTEOPETROSIS           | A rare disorder in which bones become unusually dense but may be fragile.                       |
| Weak bones                    | OSTEOPOROSIS            | A condition in which bones lose strength and become more likely to break.                       |
| Female sex glands             | OVARIES                 | The organs that produce eggs and reproductive hormones.                                         |
| Given by injection            | PARENTERAL              | Given by injection rather than through the digestive tract.                                     |
| Open passage                  | PATENCY                 | The condition of being open and unobstructed.                                                   |
| How a disease develops        | PATHOGENESIS            | The process by which a disease or unhealthy condition develops.                                 |
| Through the skin              | PERCUTANEOUS            | Done through or across the skin.                                                                |
| Away from the center          | PERIPHERAL              | At or toward the outer part rather than the center.                                             |
| By mouth                      | PER OS (PO)             | A prescription direction meaning by mouth.                                                      |
| How the body handles a drug   | PHARMACOKINETICS        | The study of how the body absorbs, distributes, changes, and removes a drug.                    |
| First drug-study phase        | PHASE I                 | The first human study phase, usually focused on safety, effects, and dosing.                    |
| Early effectiveness study     | PHASE II                | A study phase examining safety and early evidence of benefit in people with a condition.        |
| Large confirmation study      | PHASE III               | A larger study phase used to confirm safety and effectiveness and identify common side effects. |

| Lay term                     | Medical term      | Lay-language explanation                                                                         |
|------------------------------|-------------------|--------------------------------------------------------------------------------------------------|
| After-approval study         | PHASE IV          | A study performed after approval to monitor safety or examine additional uses.                   |
| Vein irritation              | PHLEBITIS         | Irritation or inflammation of a vein.                                                            |
| Inactive treatment           | PLACEBO           | An inactive substance or treatment that looks like the real treatment.                           |
| Improvement from expectation | PLACEBO EFFECT    | An improvement that occurs after an inactive treatment, often because of expectation or context. |
| Clotting blood particles     | PLATELETS         | Small blood components that help stop bleeding by forming clots.                                 |
| Possible                     | POTENTIAL         | Possible or capable of happening.                                                                |
| Make stronger                | POTENTIATE        | To increase or multiply the effect of a drug, treatment, or substance.                           |
| Effect booster               | POTENTIATOR       | An agent that helps another agent work more strongly.                                            |
| Before birth                 | PRENATAL          | Occurring before birth.                                                                          |
| Preventive treatment         | PROPHYLAXIS       | A medicine or action used to prevent disease or infection.                                       |
| As needed                    | PRN               | Prescription shorthand meaning when needed.                                                      |
| Expected outlook             | PROGNOSIS         | The expected course and outcome of a disease.                                                    |
| Lying face down              | PRONE             | Lying on the stomach or front.                                                                   |
| Study looking forward        | PROSPECTIVE STUDY | A study that follows participants forward in time.                                               |
| Artificial body part         | PROSTHESIS        | An artificial replacement for a body part.                                                       |
| Study plan                   | PROTOCOL          | The written plan for a study or treatment.                                                       |
| Closer to the center         | PROXIMAL          | Closer to the center of the body or point of attachment.                                         |
| Related to the lungs         | PULMONARY         | Pertaining to the lungs.                                                                         |
| Every day                    | QD                | Prescription shorthand meaning every day; clearer wording is preferred.                          |
| Four times daily             | QID               | Prescription shorthand meaning four times a day.                                                 |
| Radiation treatment          | RADIATION THERAPY | Treatment using high-energy radiation to damage or destroy abnormal cells.                       |
| By chance                    | RANDOM            | Selected or occurring by chance.                                                                 |
| Chance assignment            | RANDOMIZATION     | Assigning participants to groups by chance.                                                      |
| Red blood cell               | RBC               | A red blood cell.                                                                                |
| Genetically combined         | RECOMBINANT       | Made by combining genetic material from different sources.                                       |
| Put back together            | RECONSTITUTION    | Restoring something by putting its parts back together or adding a liquid.                       |
| Happen again                 | RECUR             | To happen again.                                                                                 |
| Not responding               | REFRACTORY        | Not responding adequately to treatment.                                                          |
| Regrowth or repair           | REGENERATION      | Growth again or replacement of lost or damaged tissue.                                           |
| Treatment schedule           | REGIMEN           | The planned pattern, dose, and timing of treatment.                                              |
| Return of disease            | RELAPSE           | Return of a disease after improvement or remission.                                              |
| No detectable disease        | REMISSION         | Disappearance or reduction of evidence of disease; it may be partial or complete.                |
| Related to kidneys           | RENAL             | Pertaining to the kidneys.                                                                       |
| Can be repeated              | REPLICABLE        | Able to be repeated with similar results.                                                        |



| Lay term                    | Medical term        | Lay-language explanation                                                                                    |
|-----------------------------|---------------------|-------------------------------------------------------------------------------------------------------------|
| Remove surgically           | RESECT              | To remove part or all of an organ or tissue by surgery.                                                     |
| Study looking back          | RETROSPECTIVE STUDY | A study that examines past records or past events.                                                          |
| Connective-tissue cancer    | SARCOMA             | A cancer arising from connective, bone, muscle, fat, or related tissues.                                    |
| Calming medicine            | SEDATIVE            | A medicine that calms or makes a person sleepy.                                                             |
| Testicular cancer           | SEMINOMA            | A type of cancer arising in the testicle.                                                                   |
| In order                    | SEQUENTIALLY        | Happening one after another in order.                                                                       |
| Sleepiness                  | SOMNOLENCE          | Unusual sleepiness or a tendency to fall asleep.                                                            |
| Air-measuring instrument    | SPIROMETER          | An instrument that measures the amount of air moved into and out of the lungs.                              |
| Extent of disease           | STAGING             | Assessment of how far a disease has progressed or spread.                                                   |
| Usual accepted treatment    | STANDARD OF CARE    | A treatment plan generally accepted by the medical community as appropriate.                                |
| Narrowing                   | STENOSIS            | Narrowing of a passage, duct, or blood vessel.                                                              |
| Mouth inflammation          | STOMATITIS          | Sores or inflammation inside the mouth.                                                                     |
| Group for comparison        | STRATIFY            | To arrange people or results into groups for analysis.                                                      |
| Barely responsive state     | STUPOR              | A greatly reduced level of alertness in which a person is difficult to awaken or respond to.                |
| Under the collarbone        | SUBCLAVIAN          | Located beneath the collarbone.                                                                             |
| Under the skin              | SUBCUTANEOUS        | Beneath the skin.                                                                                           |
| Lying on the back           | SUPINE              | Lying face up on the back.                                                                                  |
| Care focused on comfort     | SUPPORTIVE CARE     | General care that relieves symptoms and supports the person without directly curing the underlying disease. |
| Having symptoms             | SYMPTOMATIC         | Showing or having symptoms.                                                                                 |
| Group of related findings   | SYNDROME            | A condition identified by a characteristic group of symptoms and signs.                                     |
| Upper blood-pressure number | SYSTOLIC            | Relating to the upper blood-pressure number when the heart contracts.                                       |
| Birth-defect-causing        | TERATOGENIC         | Capable of causing abnormalities in a developing fetus.                                                     |
| Male sex glands             | TESTES/TESTICLES    | The organs that produce sperm and male reproductive hormones.                                               |
| Clot formation              | THROMBOSIS          | Formation of a blood clot inside a blood vessel or other location.                                          |
| Blood clot                  | THROMBUS            | A blood clot formed within a blood vessel or the heart.                                                     |
| Three times daily           | TID                 | Prescription shorthand meaning three times a day.                                                           |
| Gradual dose adjustment     | TITRATION           | Gradually changing a dose or solution to reach a desired effect.                                            |
| Immune white blood cells    | T-LYMPHOCYTES       | White blood cells that help coordinate or carry out immune responses.                                       |
| On the surface              | TOPICAL             | Applied to the surface, usually the skin or a mucous membrane.                                              |
| Surface numbing medicine    | TOPICAL ANESTHETIC  | A numbing medicine applied to a particular surface area.                                                    |

---

| Lay term                 | Medical term          | Lay-language explanation                                                                          |
|--------------------------|-----------------------|---------------------------------------------------------------------------------------------------|
| Harmful effects          | TOXICITY              | Harmful or unwanted effects caused by a drug, chemical, or treatment.                             |
| Through the skin         | TRANSDERMAL           | Delivered through the skin.                                                                       |
| Temporarily              | TRANSIENTLY           | For a short time or temporarily.                                                                  |
| Injury                   | TRAUMA                | Physical injury or wound.                                                                         |
| Walking exercise machine | TREADMILL             | A moving walking surface used for exercise or heart testing.                                      |
| Absorption into tissue   | UPTAKE                | Absorbing or taking in a substance by living tissue.                                              |
| Heart-valve repair       | VALVULOPLASTY         | A procedure that repairs or widens a heart valve.                                                 |
| Enlarged veins           | VARICES               | Abnormally enlarged or twisted veins.                                                             |
| Blood-vessel spasm       | VASOSPASM             | Sudden narrowing of blood vessels.                                                                |
| Disease carrier          | VECTOR                | An organism or carrier that can transmit a disease-causing microorganism.                         |
| Blood draw               | VENIPUNCTURE          | Puncturing a vein with a needle to draw blood or give treatment.                                  |
| Parent-to-child spread   | VERTICAL TRANSMISSION | Spread of an infection from a parent to a child, often during pregnancy, birth, or breastfeeding. |
| White blood cell         | WBC                   | A white blood cell involved in immune defense.                                                    |

---

*Source and adaptation:* Terms and concepts were checked against the University of Iowa Human Subjects Office, “Medical Terms in Lay Language,” available at <https://hso.research.uiowa.edu/get-started/guides-and-standard-operating-procedures-sops/medical-terms-lay-language>. The explanations in this appendix are paraphrased for this dictionary and are not the only acceptable plain-language alternatives.

# Bibliography and Sources



# Bibliography

- [1] CSV610. *Medical Dictionary Book*. GitHub repository, commit 9c065bb (“Expand and standardize urology terms”). Available at: [https://github.com/csv610/med\\_dict\\_book](https://github.com/csv610/med_dict_book) (accessed 16 September 2026).
- [2] American College of Rheumatology. *Diseases and Conditions*. Available at: <https://rheumatology.org/patients/diseases-and-conditions> (accessed 16 September 2026).

# Index

1p36 Deletion Syndrome, [501](#)  
2,3-Bisphosphoglycerate, [501](#)  
2,3-Diphosphoglycerate, [501](#)  
22q11.2 Deletion Syndrome, [501](#)  
5-Alpha Reductase Inhibitor, [2109](#)

A & E Medicine, [1](#)  
A Fib, [1](#)  
A-T, [1](#)  
A1AT Deficiency, [1](#)  
A&E Medicine, [2109](#)  
Aagenaes Syndrome, [1](#)  
Aarskog Syndrome, [1](#)  
Aarskog-Scott Syndrome, [1](#)  
Aase Syndrome, [1](#)  
Aase-Smith Syndrome I, [1](#)  
Aase-Smith Syndrome II, [1](#)  
AAT, [1](#)  
AAT Deficiency, [1](#)  
Abadie Sign, [1](#)  
Abandoned Lead, [2](#)  
Abasia, [2](#)  
Abate, [2](#)  
Abatement, [2](#)  
ABCDE Assessment, [2](#)  
ABCDE Rule For Melanoma, [2](#)  
Abdomen, [2](#)  
Abdominal, [2](#)  
Abdominal Abscess, [2](#)  
Abdominal Adhesion, [2](#)  
Abdominal Adhesions, [2](#)  
Abdominal Angina, [2](#)  
Abdominal Aorta, [3](#)  
Abdominal Aortic Aneurysm, [3](#)  
Abdominal Aortic Aneurysm Repair, [3](#)  
Abdominal Aortic Aneurysm Screening, [3](#)  
Abdominal Aortic Ultrasound, [3](#)  
Abdominal Binder, [3](#)  
Abdominal Bloating, [3](#)  
Abdominal Bruit, [3](#)  
Abdominal Cavity, [3](#)  
Abdominal Circumference, [3](#)  
Abdominal Colic, [3](#)  
Abdominal Compartment Syndrome, [3](#)

Abdominal CT, [3](#)  
Abdominal CT Scan, [4](#)  
Abdominal Distension, [4](#)  
Abdominal Distention, [4](#)  
Abdominal Exploration, [4](#)  
Abdominal Fat, [4](#)  
Abdominal Fibromatosis, [4](#)  
Abdominal Fluid, [4](#)  
Abdominal Girth, [4](#)  
Abdominal Guarding, [4](#)  
Abdominal Hernia, [4](#)  
Abdominal Hysterectomy, [4](#)  
Abdominal Mass, [4](#)  
Abdominal Migraine, [4](#)  
Abdominal MRI Scan, [4](#)  
Abdominal Muscles, [5](#)  
Abdominal Neoplasm, [5](#)  
Abdominal Neoplasms, [5](#)  
Abdominal Pain, [5](#)  
Abdominal Palpation, [5](#)  
Abdominal Point Tenderness, [5](#)  
Abdominal Pregnancy, [5](#)  
Abdominal Quadrant, [5](#)  
Abdominal Radiography, [5](#)  
Abdominal Reflex, [5](#)  
Abdominal Regions, [5](#)  
Abdominal Retractor, [5](#)  
Abdominal Rigidity, [5](#)  
Abdominal Sounds, [6](#)  
Abdominal Swelling, [6](#)  
Abdominal Tap, [6](#)  
Abdominal Thrusts, [6](#)  
Abdominal Ultrasound, [6](#)  
Abdominal Wall, [6](#)  
Abdominal Wall Fat Pad Biopsy, [6](#)  
Abdominal Wall Hernia, [6](#)  
Abdominal Wall Surgery, [6](#)  
Abdominal X-Ray, [6](#)  
Abdominocentesis, [6](#)  
Abdominopelvic Cavity, [6](#)  
Abdominoperineal Resection, [6](#)  
Abdominoplasty, [6](#)  
Abducens Nerve, [6](#)  
Abducens Nerve Diseases, [7](#)



---

Abducens Nerve Palsy, [7](#)  
 Abducent Nerve, [7](#)  
 Abduct, [7](#)  
 Abduction, [7](#)  
 Abductor Muscle, [7](#)  
 Abercrombie Syndrome, [7](#)  
 Aberrant Conduction, [7](#)  
 Aberration, [7](#)  
 Abetalipoproteinemia, [7](#)  
 Abfraction, [7](#)  
 Abiotrophy, [7](#)  
 Ablate, [7](#)  
 Ablation, [8](#)  
 Ablative Surgery, [8](#)  
 Ablepharon, [8](#)  
 Abnormal, [8](#)  
 Abnormal (Dysfunctional) Uterine Bleeding, [8](#)  
 Abnormal Color, [8](#)  
 Abnormal Fissure, [8](#)  
 Abnormal Hemoglobins Testing, [8](#)  
 Abnormal Labor, [8](#)  
 Abnormal Pap Test, [8](#)  
 Abnormal Posturing, [8](#)  
 Abnormal Reflex, [8](#)  
 Abnormal Tooth Shape, [8](#)  
 Abnormal Uterine Bleeding, [8](#)  
 Abnormal Vaginal Bleeding, [9](#)  
 Abnormal Virilization, [9](#)  
 Abnormality, [9](#)  
 Abnormally Dark Or Light Skin, [9](#)  
 Abnormally Excessive Sweating, [9](#)  
 ABO Blood Group System, [9](#)  
 ABO Blood Typing, [9](#)  
 ABO Incompatibility, [9](#)  
 Abortifacient, [2109](#)  
 Abortion, [9](#)  
 Abortion - Medication, [2109](#)  
 Abortion Complications, [9](#)  
 Abortive, [9](#)  
 Abortive Polio, [9](#)  
 Above The Knee Amputation, [9](#)  
 ABR Test, [10](#)  
 Abrade, [10](#)  
 Abrasion, [10](#)  
 Abrin, [10](#)  
 Abruptio Placenta, [10](#)  
 Abruptio Placentae, [10](#)  
 Abruption, Placental, [1538](#)  
 Abs, [10](#)  
 Abscess, [10](#)  
 Abscess, Bartholin's, [177](#)  
 Abscess, Cutaneous, [489](#)  
 Abscessed Tooth, [10](#)  
 Abscission, [10](#)  
 Absence Epilepsy, [10](#)  
 Absence of Menstruation, Primary, [1592](#)  
 Absence Of Sweating, [10](#)  
 Absence Of The Breast, [10](#)  
 Absence Of The Nipple, [11](#)  
 Absence Seizure, [11](#)  
 Absence Seizure Cross-Reference, [11](#)  
 Absent Breath Sounds, [11](#)  
 Absent Eye, [11](#)  
 Absent Menstrual Periods, [11](#)  
 Absent Menstrual Periods, Primary, [1592](#)  
 Absent Menstrual Periods, Secondary, [1759](#)  
 Absent Pulmonary Valve, [11](#)  
 Absent Reflex, [11](#)  
 Absent Thirst, [11](#)  
 Absolute Eosinophil Count, [11](#)  
 Absolute Neutrophil Count, [11](#)  
 Absorb, [11](#)  
 Absorption, [11](#)  
 Abstinence, [11](#)  
 Abulia, [11](#)  
 Abuse Steroid (Anabolic Steroid Abuse), [2109](#)  
 Abuse, Child, [379](#)  
 Abusive Head Trauma, [12](#)  
 Abutment, [12](#)  
 AC Joint, [12](#)  
 Acalculia, [12](#)  
 Acalculous Cholecystitis, [12](#)  
 Acanthamoeba, [12](#)  
 Acanthamoeba Keratitis, [12](#)  
 Acanthocyte, [12](#)  
 Acanthocytosis, [12](#)  
 Acantholysis, [12](#)  
 Acanthoma, [12](#)  
 Acanthosis Nigricans, [13](#)  
 Acapnia, [13](#)  
 Acardia, [13](#)  
 Acariasis, [13](#)  
 Acaricide, [13](#)  
 Acatalasemia, [13](#)  
 Acatalasia, [13](#)  
 ACC – Adenoid Cystic Carcinoma, [13](#)  
 ACC – Agenesis of Corpus Callosum, [13](#)  
 ACC – Aplasia Cutis Congenita, [13](#)  
 Accelerated Growth, [13](#)  
 Accelerated Idioventricular Rhythm, [13](#)  
 Accelerated-Phase Chronic Myeloid Leukemia, [13](#)  
 Accelerometry, [14](#)  
 Acceptable Daily Intake, [14](#)  
 Acceptance-Based Therapies, [14](#)  
 Access Needle, [14](#)  
 Accessory, [14](#)  
 Accessory Digestive Organ, [14](#)  
 Accessory Dwelling Unit, [14](#)

---

---

Accessory Muscles, [14](#)  
Accessory Nerve, [14](#)  
Accessory Pathway, [14](#)  
Accessory Placenta, [14](#)  
Accident And Emergency Medicine, [14](#)  
Accidental Death, [14](#)  
Accidental Fall, [15](#)  
Accommodation, [15](#)  
Accommodation Reflex, [15](#)  
Accommodative Insufficiency, [15](#)  
Accouchement, [15](#)  
Accretion, [15](#)  
ACE Blood Test, [15](#)  
ACE Inhibitor, [2109](#)  
ACE Inhibitors, [15](#)  
Acebutolol, [15](#)  
Acellular, [15](#)  
Acentric Chromosome, [15](#)  
Acephalostomia, [15](#)  
Acetabular Labrum, [15](#)  
Acetabular Notch, [15](#)  
Acetabuloplasty, [15](#)  
Acetabulum, [16](#)  
Acetaldehyde, [16](#)  
Acetaminophen, [16](#)  
Acetaminophen And Codeine Overdose, [16](#)  
Acetaminophen Overdose, [16](#)  
Acetaminophen Poisoning, [16](#)  
Acetaminophen Toxicity, [16](#)  
Acetate, [16](#)  
Acetazolamide, [16](#)  
Acetone, [16](#)  
Acetone Body, [16](#)  
Acetone Breath, [16](#)  
Acetone Poisoning, [17](#)  
Acetonuria, [17](#)  
Acetyl-CoA Carboxylase, [17](#)  
Acetyl-Coenzyme A, [17](#)  
Acetylation, [17](#)  
Acetylcholine, [17](#)  
Acetylcholine Receptor Antibody, [17](#)  
Acetylcholinesterase, [17](#)  
Acetylcholinesterase Inhibitors, [17](#)  
Acetylcysteine, [17](#)  
Acetylsalicylic Acid, [17](#)  
ACH, [17](#)  
ACh (Acetylcholine), [17](#)  
Achalasia, [17](#)  
Achalasia Of The Cardia, [18](#)  
AChE (Acetylcholinesterase), [18](#)  
Achenbach's Syndrome, [18](#)  
Achilles Tendinitis, [18](#)  
Achilles Tendinopathy, [18](#)  
Achilles Tendon, [18](#)  
Achilles Tendon Problems, [18](#)  
Achilles Tendon Repair, [18](#)  
Achilles Tendon Rupture, [18](#)  
Achilles Tendonitis, [18](#)  
Achillobursitis, [18](#)  
Achlorhydria, [18](#)  
Achondrogenesis, [19](#)  
Achondroplasia, [19](#)  
Achoo Syndrome, [19](#)  
Achromatopsia, [19](#)  
Achromia, [19](#)  
Aciclovir, [19](#)  
Acid, [19](#)  
Acid Base Balance, [19](#)  
Acid Indigestion, [19](#)  
Acid Loading Test, [19](#)  
Acid Mucopolysaccharides, [19](#)  
Acid Phosphatase, [19](#)  
Acid Reflux, [19](#)  
Acid Soldering Flux Poisoning, [19](#)  
Acid Sphingomyelinase Deficiency, [20](#)  
Acid-Base Analysis, [20](#)  
Acid-Base Disorder, [20](#)  
Acid-Base Imbalance, [20](#)  
Acid-Base, Fluid, and Electrolyte Disorders, [569](#)  
Acid-Fast Bacilli, [20](#)  
Acid-Fast Bacillus Smear, [20](#)  
Acid-Fast Stain, [20](#)  
Acidemia, [20](#)  
Acidophil, [20](#)  
Acidophilus, [20](#)  
Acidosis, [20](#)  
Aciduria, [20](#)  
Acinar Cell, [20](#)  
Acinar Cell Carcinoma, [21](#)  
Acinetobacter Baumannii Resistance, [21](#)  
Acinus, [21](#)  
Acitretin, [2109](#)  
ACL Injury, [21](#)  
ACL Reconstruction, [21](#)  
Acne, [21](#)  
Acne Conglobata, [21](#)  
Acne Fulminans, [21](#)  
Acne Inversa, [21](#)  
Acne Keloid, [21](#)  
Acne Keloidalis Nuchae (AKN), [21](#)  
Acne Rosacea, [21](#)  
Acne Vulgaris, [21](#)  
Acneiform Eruptions, [21](#)  
Acoustic, [22](#)  
Acoustic Nerve, [22](#)  
Acoustic Neuroma, [22](#)  
Acoustic Reflex, [22](#)  
Acoustic Shock, [22](#)

---

---

Acoustic Trauma, [22](#)  
 Acquired, [22](#)  
 Acquired C1-Inhibitor Deficiency Angioedema, [22](#)  
 Acquired Cystic Kidney Disease, [22](#)  
 Acquired Feminization, [22](#)  
 Acquired Immunity, [22](#)  
 Acquired Immunodeficiency Syndrome, [22](#)  
 Acquired Lipodystrophy, [22](#)  
 Acquired Methemoglobinemia, [23](#)  
 Acquired Nasolacrimal Duct Obstruction, [23](#)  
 Acquired Platelet Function Defect, [23](#)  
 Acquired Tracheomalacia, [23](#)  
 Acquisition, [23](#)  
 Acquisition Duration, [23](#)  
 Acquisition Time, [23](#)  
 Acral Lentiginous Melanoma, [23](#)  
 Acro-Osteolysis, [23](#)  
 Acrocentric Chromosome, [23](#)  
 Acrochordon, [23](#)  
 Acrocyanosis, [24](#)  
 Acrodermatitis, [24](#)  
 Acrodermatitis Chronica Atrophicans, [24](#)  
 Acrodermatitis Enteropathica, [24](#)  
 Acrodynia, [24](#)  
 Acrodysostosis, [24](#)  
 Acrokeratosis Paraneoplastica, [24](#)  
 Acromegaly, [24](#)  
 Acromioclavicular Joint, [24](#)  
 Acromioclavicular Joint Injuries, [24](#)  
 Acromion, [24](#)  
 Acroparaesthesia, [24](#)  
 Acrophobia, [24](#)  
 Acropustulosis of Infancy, [25](#)  
 Acrosin, [25](#)  
 Acrosome, [25](#)  
 Acrosome Reaction, [25](#)  
 ACTH Blood Test, [25](#)  
 ACTH Stimulation Test, [25](#)  
 Actigraphy, [25](#)  
 Actin, [25](#)  
 Actin Myofilament, [25](#)  
 Actin-Binding Protein, [25](#)  
 Acting Out, [25](#)  
 Actinic, [25](#)  
 Actinic Cheilitis, [25](#)  
 Actinic Granuloma, [25](#)  
 Actinic Keratosis, [26](#)  
 Actinic Keratosis Pathology, [26](#)  
 Actinic Purpura, [26](#)  
 Actinomycetoma, [26](#)  
 Actinomycin D, [26](#)  
 Actinomycosis, [26](#)  
 Action Potential, [26](#)  
 Action Tremor, [26](#)  
 Activated Charcoal, [26](#)  
 Activated Partial Thromboplastin Time, [26](#)  
 Activated Protein C Resistance, [26](#)  
 Active Caries, [27](#)  
 Active Management Of Third Stage, [27](#)  
 Active Phase Of Labor, [27](#)  
 Active Site, [27](#)  
 Active Surveillance, [27](#)  
 Active Transport, [27](#)  
 Activities Of Daily Living, [27](#)  
 Acupressure, [27](#)  
 Acupuncture, [27](#)  
 Acute, [27](#)  
 Acute Abdomen, [27](#)  
 Acute Acalculous Cholecystitis, [27](#)  
 Acute Adrenal Crisis, [27](#)  
 Acute Agitation, [28](#)  
 Acute Alcohol Intoxication, [28](#)  
 Acute Angle-Closure Glaucoma, [28](#)  
 Acute Aortic Syndrome, [28](#)  
 Acute Bacterial Prostatitis, [28](#)  
 Acute Bacterial Rhinosinusitis, [28](#)  
 Acute Bilirubin Encephalopathy, [28](#)  
 Acute Blood Loss, [28](#)  
 Acute Bronchitis, [28](#)  
 Acute Bursitis, [28](#)  
 Acute Caries, [29](#)  
 Acute Cerebellar Ataxia, [29](#)  
 Acute Chest Syndrome, [29](#)  
 Acute Cholangitis, [29](#)  
 Acute Cholecystitis, [29](#)  
 Acute Colonic Pseudo-Obstruction, [29](#)  
 Acute Compartment Syndrome, [29](#)  
 Acute Coronary Syndrome, [29](#)  
 Acute Coronary Syndromes, [29](#)  
 Acute Cough, [29](#)  
 Acute Cutaneous Lupus Erythematosus (ACLE), [29](#)  
 Acute Cystitis, [29](#)  
 Acute Dacryocystitis, [30](#)  
 Acute Decompensated Heart Failure, [30](#)  
 Acute Diarrhea, [30](#)  
 Acute Disease, [30](#)  
 Acute Disseminated Encephalomyelitis, [30](#)  
 Acute Ear Infection, [30](#)  
 Acute Endometritis, [30](#)  
 Acute Epididymitis, [30](#)  
 Acute Esophageal Necrosis, [30](#)  
 Acute Exacerbation, [30](#)  
 Acute Exacerbation Of Asthma, [30](#)  
 Acute Exacerbation Of COPD, [30](#)  
 Acute Fatty Liver Of Pregnancy, [30](#)  
 Acute Febrile Illness, [31](#)  
 Acute Febrile Neutrophilic Dermatitis (Sweet Syndrome), [31](#)

---

---

Acute Fever in Adults, [31](#)  
Acute Flaccid Myelitis, [31](#)  
Acute Gastritis, [31](#)  
Acute Generalized Exanthematous Pustulosis, [31](#)  
Acute Glomerulonephritis, [31](#)  
Acute Granulocytic Leukemia, [31](#)  
Acute Hemorrhagic Edema of Infancy, [31](#)  
Acute Hemorrhagic Leukoencephalitis, [31](#)  
Acute Hepatitis, [31](#)  
Acute HIV Syndrome, [32](#)  
Acute Hypercapnic Respiratory Failure, [32](#)  
Acute Hypoxemic Respiratory Failure, [32](#)  
Acute Idiopathic Polyneuritis, [32](#)  
Acute Illness, [32](#)  
Acute Inflammation, [32](#)  
Acute Inflammatory Demyelinating Polyneuropathy, [32](#)  
Acute Inflammatory Demyelinating Polyradiculoneuropathy, [32](#)  
Acute Intermittent Porphyria, [32](#)  
Acute Interstitial Nephritis, [32](#)  
Acute Ischemic Stroke, [32](#)  
Acute Kidney Failure, [32](#)  
Acute Kidney Injury, [32](#)  
Acute Kidney Injury Staging, [32](#)  
Acute Leukemia, [33](#)  
Acute Limb Ischemia, [33](#)  
Acute Limb Ischemia Six Ps, [33](#)  
Acute Liver Failure, [33](#)  
Acute Low Back Pain, [33](#)  
Acute Lymphangitis, [33](#)  
Acute Lymphoblastic Leukemia, [33](#)  
Acute Lymphocytic Leukemia, [33](#)  
Acute Lymphoid Leukemia, [33](#)  
Acute Megakaryoblastic Leukemia, [33](#)  
Acute Mesenteric Ischemia, [33](#)  
Acute Mitral Regurgitation, [33](#)  
Acute Mountain Sickness, [34](#)  
Acute Myeloblastic Leukemia, [34](#)  
Acute Myelogenous Leukemia, [34](#)  
Acute Myeloid Leukemia, [34](#)  
Acute Myocardial Infarction, [34](#)  
Acute Myocardial Infarction Pathology, [34](#)  
Acute Necrotizing Ulcerative Gingivitis, [34](#)  
Acute Nephritic Syndrome, [34](#)  
Acute Nonlymphocytic Leukemia, [34](#)  
Acute Obstructive Uropathy, [34](#)  
Acute Onset Pain, [34](#)  
Acute Otitis Media, [34](#)  
Acute Pain, [35](#)  
Acute Pancreatitis, [35](#)  
Acute Pericarditis, [35](#)  
Acute Poisoning, [35](#)  
Acute Proctitis, [35](#)  
Acute Promyelocytic Leukemia, [35](#)  
Acute Pulmonary Edema, [35](#)  
Acute Pulmonary Histoplasmosis, [35](#)  
Acute Pyelonephritis, [35](#)  
Acute Radiation Sickness, [35](#)  
Acute Radiation Syndrome, [35](#)  
Acute Rehabilitation, [35](#)  
Acute Renal Artery Occlusion, [35](#)  
Acute Renal Failure, [36](#)  
Acute Respiratory Disease, [36](#)  
Acute Respiratory Distress Syndrome, [36](#)  
Acute Respiratory Failure, [36](#)  
Acute Retinal Necrosis, [36](#)  
Acute Retroviral Syndrome, [36](#)  
Acute Rhinitis, [36](#)  
Acute Rhinosinusitis, [36](#)  
Acute Sinusitis, [36](#)  
Acute Stress Disorder, [36](#)  
Acute Symptomatic Hyponatremia, [36](#)  
Acute Transverse Myelitis, [36](#)  
Acute Tubular Necrosis, [36](#)  
Acute Urinary Retention, [37](#)  
Acute Viral Hepatitis, [37](#)  
Acute Viral Illness, [37](#)  
Acute Vision Loss, [37](#)  
Acute Weeping Dermatitis, [37](#)  
Acute-On-Chronic Liver Failure, [37](#)  
Acute-Phase Reactants, [37](#)  
Acyclovir, [37](#)  
Ad Lib, [37](#)  
ADA Deficiency, [37](#)  
Adactyly, [37](#)  
Adam's Apple, [37](#)  
Adamantine, [37](#)  
Adamantinoma, [37](#)  
Adams-Stokes Syndrome, [38](#)  
Adaptation, [38](#)  
Adaptive Design, [38](#)  
Adaptive Equipment, [38](#)  
Adaptive Immunity, [38](#)  
ADD/ADHD, [38](#)  
Addiction, [38](#)  
Addiction, Alcohol, [62](#)  
Addiction, Gambling, [799](#)  
Addiction, Nicotine, [1358](#)  
Addiction, Sexual, [441](#)  
Addison Anemia, [38](#)  
Addison Disease, [38](#)  
Addison's Disease, [38](#)  
Addisonian Anemia, [39](#)  
Addisonian Crisis, [39](#)  
Additive Genetic Effects, [39](#)  
Adduct, [39](#)  
Adduction, [39](#)  
Adductor Muscle, [39](#)

---

---

Adductor Muscles, [39](#)  
Adenine, [39](#)  
Adenitis, [39](#)  
Adenitis, Mesenteric, [1237](#)  
Adenocarcinoma, [39](#)  
Adenocarcinoma In Situ Of Cervix, [39](#)  
Adenocarcinoma Of The Colon, [39](#)  
Adenocarcinoma Of The Lung, [39](#)  
Adenohypophysis, [39](#)  
Adenoid Basal Carcinoma, [40](#)  
Adenoid Cystic Carcinoma, [40](#)  
Adenoid Cystic Carcinoma Of The Vulva, [40](#)  
Adenoid Hypertrophy, [40](#)  
Adenoid Removal, [40](#)  
Adenoidectomy, [40](#)  
Adenoiditis, [40](#)  
Adenoids, [40](#)  
Adenolipoma, [40](#)  
Adenolymphoma, [40](#)  
Adenoma, [40](#)  
Adenoma Malignum, [40](#)  
Adenoma-Carcinoma Sequence, [41](#)  
Adenomas, [41](#)  
Adenomatosis, [41](#)  
Adenomatous Polyp, [41](#)  
Adenomatous Polyposis, [41](#)  
Adenomyoepithelioma Of The Breast, [41](#)  
Adenomyoma, [41](#)  
Adenomyosis, [41](#)  
Adenopathy, [41](#)  
Adenosine, [41](#)  
Adenosine Deaminase Deficiency, [41](#)  
Adenosine Deaminase Severe Combined Immunodeficiency (ADA-SCID), [41](#)  
Adenosine Diphosphate, [41](#)  
Adenosine Monophosphate, [42](#)  
Adenosine Triphosphate, [42](#)  
Adenosis, [42](#)  
Adenovirus, [42](#)  
Adenovirus DNA PCR (Feces), [42](#)  
Adenovirus Infection, [42](#)  
Adenovirus Vaccine, [2109](#)  
Adenoviruses, [42](#)  
Adenylate Cyclase, [42](#)  
Adequate Intake, [42](#)  
ADHD in Children, [42](#)  
Adherence, [43](#)  
Adherens Junction, [43](#)  
Adhesiolysis, [43](#)  
Adhesion, [43](#)  
Adhesion Molecules, [43](#)  
Adhesive Capsulitis, [43](#)  
Adhesive Tape, [43](#)  
Adie Tonic Pupil, [43](#)  
Adipocere, [43](#)  
Adipocyte, [43](#)  
Adipocytes, [43](#)  
Adipogenesis, [43](#)  
Adipokine, [43](#)  
Adiponectin, [44](#)  
Adipose, [44](#)  
Adipose Tissue, [44](#)  
Adiposis Dolorosa, [44](#)  
Adiposity, [44](#)  
Adipsia, [44](#)  
Adjacent, [44](#)  
Adjustment Disorder, [44](#)  
Adjustment Disorder With Anxiety, [44](#)  
Adjustment Disorders, [44](#)  
Adjuvant, [44](#)  
Adjuvant Chemotherapy, [44](#)  
Adjuvant Therapy, [44](#)  
Adlerian Theory, [44](#)  
ADME, [45](#)  
Administered Activity, [45](#)  
Administration Procedure, [45](#)  
Adnexa, [45](#)  
Adnexal Dysplasia, [45](#)  
Adnexal Mass, [45](#)  
Adnexal Torsion, [45](#)  
Adnexal Tumors, [45](#)  
Adnexal Tumors And Masses, [45](#)  
Adnexitis, [45](#)  
Adolescence, [45](#)  
Adolescent Development, [45](#)  
Adolescent Health, [45](#)  
Adolescent Medicine, [46](#)  
Adolescent Obesity, [46](#)  
Adolescent Schizophrenia, [46](#)  
Adolescent Substance Use, [46](#)  
Adoptive Cell Therapy, [46](#)  
Adoptive Immunotherapy, [46](#)  
ADP-Ribosylation, [46](#)  
Adrenal Cancer, [46](#)  
Adrenal Carcinoma, [46](#)  
Adrenal Cortex, [46](#)  
Adrenal Cortex Physiology, [46](#)  
Adrenal Cortical Carcinoma, [46](#)  
Adrenal Crisis, [46](#)  
Adrenal Failure, [46](#)  
Adrenal Gland, [47](#)  
Adrenal Gland Cancer, [47](#)  
Adrenal Gland Removal, [47](#)  
Adrenal Gland Tumor, [47](#)  
Adrenal Glands, [47](#)  
Adrenal Hemorrhage, [47](#)  
Adrenal Hyperplasia, [47](#)  
Adrenal Incidentaloma, [47](#)

---

---

Adrenal Insufficiency, [47](#)  
Adrenal Mass, [47](#)  
Adrenal Medulla, [47](#)  
Adrenal Medulla Physiology, [47](#)  
Adrenal Myelolipoma, [47](#)  
Adrenal Nodule, [48](#)  
Adrenal Tumors, [48](#)  
Adrenal Venous Sampling, [48](#)  
Adrenalectomy, [48](#)  
Adrenaline, [48](#)  
Adrenarche, [48](#)  
Adrenergic Agent, [48](#)  
Adrenergic Agonist, [48](#)  
Adrenergic Antagonists, [48](#)  
Adrenergic Bronchodilator Overdose, [48](#)  
Adrenergic Receptor, [48](#)  
Adrenergic Receptors, [48](#)  
Adrenocortical Carcinoma, [49](#)  
Adrenocorticotrophic Hormone, [49](#)  
Adrenogenital Syndrome, [49](#)  
Adrenoleukodystrophy, [49](#)  
Adrenomedullin, [49](#)  
Adrenomyeloneuropathy, [49](#)  
Adson Test, [49](#)  
Adsorption, [49](#)  
Adult Day Services, [49](#)  
Adult Hypertension, [49](#)  
Adult Immunisation, [50](#)  
Adult Informed Consent, [50](#)  
Adult Nutritional Support in Illness, [50](#)  
Adult Obstructive Sleep Apnea, [50](#)  
Adult Primary Brain Tumor, [50](#)  
Adult Spinal Deformity, [50](#)  
Adult T-Cell Leukemia, [50](#)  
Adult-Onset Asthma, [50](#)  
Adult-Onset Still Disease, [50](#)  
Advance Directive, [51](#)  
Advance Statements, [51](#)  
Advanced Cancer, [51](#)  
Advanced Cardiovascular Life Support, [51](#)  
Advanced Glycation End-Products, [51](#)  
Advanced Prostate Cancer, [51](#)  
Advanced Sleep-Wake Phase Disorder, [51](#)  
Adventitia, [51](#)  
Adventitious, [51](#)  
Adverse Drug Event, [2109](#)  
Adverse Drug Reaction, [2109](#)  
Adverse Effect, [51](#)  
Adverse Event, [51](#)  
Adverse Reaction, [51](#)  
Aedes Aegypti, [52](#)  
Aegophony, [52](#)  
Aerobe, [52](#)  
Aerobic, [52](#)  
Aerobic Bacteria, [52](#)  
Aerobic Bacterium, [52](#)  
Aerobic Exercise, [52](#)  
Aerobiosis, [52](#)  
Aerophagia, [52](#)  
Aerophagy, [52](#)  
Aerophobia, [52](#)  
Aerosol, [52](#)  
Aerospace Medicine, [52](#)  
Aerotitis, [52](#)  
Aesthetic Dentistry, [53](#)  
Aesthetic Medicine, [53](#)  
Aetiology, [53](#)  
Affect, [53](#)  
Affective Disorder, [53](#)  
Affective Flattening, [53](#)  
Afferent, [53](#)  
Afferent Arteriole, [53](#)  
Afferent Loop Syndrome, [53](#)  
Afferent Lymphatic Vessel, [53](#)  
Afferent Neuron, [53](#)  
Afferent Pathway, [53](#)  
Afferent Pupillary Defect, [53](#)  
Affinity, [54](#)  
Affinity Chromatography, [54](#)  
AFib, [54](#)  
Afibrinogenaemia, [54](#)  
Afibrinogenemia, [54](#)  
Aflatoxin, [54](#)  
African Sleeping Sickness, [54](#)  
African Trypanosomiasis, [54](#)  
After-Cataract, [54](#)  
Afterbirth, [54](#)  
Afterdepolarization, [54](#)  
Afterload, [54](#)  
Afterpains, [54](#)  
Aftershave Poisoning, [55](#)  
Agammaglobulinaemia, [55](#)  
Agammaglobulinemia, [55](#)  
Aganglionic Megacolon, [55](#)  
Agar, [55](#)  
Agatston Score, [55](#)  
Age Regression, [55](#)  
Age Spots, [55](#)  
Age-Related Hearing Loss, [55](#)  
Age-Related Macular Degeneration, Dry, [592](#)  
Age-Related Macular Degeneration, Wet, [2072](#)  
Age-Standardized Rate, [55](#)  
Ageing, [55](#)  
Agenesis, [55](#)  
Agenesis of Commissura Magna Cerebri, [56](#)  
Agenesis Of The Corpus Callosum, [56](#)  
Agent, Tocolytic, [2109](#)  
Ageusia, [56](#)

---



---

Agger Nasi, 56  
Agglutination, 56  
Agglutination Test, 56  
Agglutinin, 56  
Agglutininogen, 56  
Aggression, 56  
Aggressive, 56  
Aggressive Angiomyxoma, 56  
Aggressive Cancer, 56  
Agility, 56  
Agitation, 56  
Agnathia, 57  
Agnogenic Myeloid Metaplasia (AMM), 57  
Agnosia, 57  
Agnosia, Primary Visual, 2052  
Agnosis, Primary, 57  
Agonist, 57  
Agonist Muscle, 57  
Agonist-Antagonist, 2110  
Agoraphobia, 57  
Agostini's Reaction, 57  
Agranulocytes, 57  
Agranulocytosis, 57  
Agraphesthesia, 57  
Agraphia, 57  
Aicardi Syndrome, 57  
Aicardi-Goutieres Syndrome, 57  
AIDS, 58  
AIDS Dementia, 58  
AIDS Vaccine, 2110  
AIDS-Associated Nephropathy, 58  
Ainhum, 58  
Air Abrasion, 58  
Air Bag, 58  
Air Bronchogram, 58  
Air Contaminants, 58  
Air Embolism, 58  
Air Passages, 58  
Air Polishing, 58  
Air Pollution, 58  
Air Quality, 58  
Air Sac, 59  
Air Trapping, 59  
Air-Conditioner Lung, 59  
Air-Sickness, 59  
Airborne Disease, 59  
Airborne Precautions, 59  
Airplane Ear, 59  
Airsickness, 59  
Airway, 59  
Airway Clearance Techniques, 59  
Airway Management, 59  
Airway Obstruction, 59  
Airway Resistance, 59  
Akathisia, 60  
Akinesia, 60  
Akinetic, 60  
Akinetic Mutism, 60  
Akinetopsia, 60  
Alagille Syndrome, 60  
Alanine, 60  
Alanine Aminotransferase, 60  
Alanine Transaminase Blood Test, 60  
ALARA Principle, 60  
Alarmin, 60  
Albendazole, 60  
Albers-Schönberg Disease, 61  
Albinism, 61  
Albinismus, 61  
Albino, 61  
Albright Hereditary Osteodystrophy, 61  
Albright's Hereditary Osteodystrophy, 61  
Albumen, 61  
Albumin, 61  
Albumin Clearance, 61  
Albumin-To-Creatinine Ratio, 61  
Albuminocytologic Dissociation, 61  
Albuminuria, 62  
ALCAPA Syndrome, 62  
Alcohol, 62  
Alcohol Dehydrogenase, 62  
Alcohol Intolerance, 62  
Alcohol Intoxication, 62  
Alcohol Poisoning, 62  
Alcohol Swab, 62  
Alcohol Use Disorder, 62  
Alcohol Withdrawal, 62  
Alcohol-Associated Liver Disease, 63  
Alcohol-Related Psychosis, 63  
Alcoholic Cardiomyopathy, 63  
Alcoholic Foetor, 63  
Alcoholic Hepatitis, 63  
Alcoholic Ketoacidosis, 63  
Alcoholic Liver Disease, 63  
Alcoholic Myopathy, 63  
Alcoholic Neuropathy, 63  
Alcuronium, 63  
ALD, 63  
Aldolase Blood Test, 63  
Aldosterone, 63  
Aldosterone Blood Test, 64  
Aldosterone Escape, 64  
Aldosterone-to-Renin Ratio, 64  
Aldosteronism With Normal Blood Pressure, 64  
Alendronate, 2110  
Aleukemic Leukemia, 64  
Alexander Disease, 64  
Alexia, 64

---

---

Alexia Without Agraphia, [64](#)  
Alexithymia, [64](#)  
Alezzandrini Syndrome, [64](#)  
Alfacalcidol, [64](#)  
Algia, [64](#)  
Alglucerase, [65](#)  
Algodystrophy, [65](#)  
Algoneurodystrophy, [65](#)  
Algor Mortis, [65](#)  
Aliasing Artifact, [65](#)  
Alien Hand Syndrome, [65](#)  
Alimentary, [65](#)  
Alimentary Canal, [65](#)  
Alkalemia, [65](#)  
Alkali, [65](#)  
Alkaline Diet, [65](#)  
Alkaline Phosphatase, [65](#)  
Alkaline Phosphatase Test, [65](#)  
Alkaloid, [65](#)  
Alkalosis, [66](#)  
Alkaptonuria, [66](#)  
Alkylating Agents, [66](#)  
Allantoin, [66](#)  
Allantois, [66](#)  
Allele, [66](#)  
Allelic, [66](#)  
Allen Test, [66](#)  
Allergen, [66](#)  
Allergen Immunotherapy, [66](#)  
Allergic and Environmental Asthma, [66](#)  
Allergic Bronchopulmonary Aspergillosis, [66](#)  
Allergic Cascade, [66](#)  
Allergic Conjunctivitis, [67](#)  
Allergic Contact Dermatitis, [67](#)  
Allergic Dermatitis, [67](#)  
Allergic Diseases, [67](#)  
Allergic Proctocolitis, [67](#)  
Allergic Reaction, [67](#)  
Allergic Rhinitis, [67](#)  
Allergic Urticaria, [67](#)  
Allergies to Insect Stings, [67](#)  
Allergy, [67](#)  
Allergy Shots, [68](#)  
Allergy Skin Test, [68](#)  
Allergy Test, [68](#)  
Allergy to Natural Rubber (Latex), [68](#)  
Allis Clamp, [68](#)  
Allocheiria, [68](#)  
Allodynia, [68](#)  
Allogeneic, [68](#)  
Allograft, [68](#)  
Allosteric Regulation, [68](#)  
AloBar Holoprosencephaly, [68](#)  
Aloe, [68](#)  
Aloe Vera, [68](#)  
Alogia, [69](#)  
Alongshan Virus, [69](#)  
Alopecia, [69](#)  
Alopecia Areata, [69](#)  
Alopecia Celsi, [69](#)  
Alopecia Cicatrisata, [69](#)  
Alopecia Circumscripta, [69](#)  
Alopecia Mucinoso (Follicular Mucinoso), [69](#)  
Alopecia Totalis, [69](#)  
Alopecia Universalis, [69](#)  
ALP Blood Test, [69](#)  
ALP Isoenzyme Test, [69](#)  
Alpha Blockers, [69](#)  
Alpha Cells, [70](#)  
Alpha Hydroxy Acids, [70](#)  
Alpha Waves, [70](#)  
Alpha-1, [70](#)  
Alpha-1 Adrenergic Antagonist, [70](#)  
Alpha-1 Antitrypsin Blood Test, [70](#)  
Alpha-1 Antitrypsin Deficiency, [70](#)  
Alpha-Agonists, [70](#)  
Alpha-Fetoprotein, [70](#)  
Alpha-Gal Syndrome, [70](#)  
Alpha-Globulin, [70](#)  
Alpha-Glucosidase Inhibitors, [70](#)  
Alpha-Ketoglutarate, [71](#)  
Alpha-Synuclein, [71](#)  
Alpha-Thalassemia, [71](#)  
Alpha1-Antitrypsin (AAT) Deficiency, [71](#)  
Alport Syndrome, [71](#)  
Alprazolam, [71](#)  
Alstrom Syndrome, [71](#)  
Alternating Hemiplegia Of Childhood, [71](#)  
Altitude Headache, [71](#)  
Altitude Illness, [71](#)  
Aluminum Phosphide, [71](#)  
Aluminum Toxicity, [71](#)  
Alvarado Score, [72](#)  
Alveolar, [72](#)  
Alveolar Abnormalities, [72](#)  
Alveolar Bone, [72](#)  
Alveolar Bone Loss, [72](#)  
Alveolar Dead Space, [72](#)  
Alveolar Echinococcosis, [72](#)  
Alveolar Gas Equation, [72](#)  
Alveolar Hypoventilation, [72](#)  
Alveolar Macrophage, [72](#)  
Alveolar Osteitis, [72](#)  
Alveolar Proteinosis, [72](#)  
Alveolar Rhabdomyosarcoma, [72](#)  
Alveolar Ventilation, [72](#)  
Alveolar Ventilation Equation, [73](#)  
Alveolar-Arterial Oxygen Gradient, [73](#)

---

---

Alveolar-Capillary Barrier, 73  
 Alveolitis, 73  
 Alveolus, 73  
 Alzheimer Disease, 73  
 Alzheimer's, 73  
 Amalgam, 73  
 Amastia, 73  
 Amaurosis, 73  
 Amaurosis Fugax, 73  
 Ambidextrous, 74  
 Ambient, 74  
 Ambiguous Genitalia, 74  
 Amblyopia, 74  
 Ambu Bag, 74  
 Ambulatory Blood Pressure Monitoring, 74  
 Amebiasis, 74  
 Amebic Colitis, 74  
 Amebic Dysentery, 74  
 Amebic Liver Abscess, 74  
 Ameboma, 75  
 Amelanotic Melanoma, 75  
 Amelia, 75  
 Amelioration, 75  
 Ameloblastoma, 75  
 Amelogenesis, 75  
 Amelogenesis Imperfecta, 75  
 Amenorrhea, 75  
 American Trypanosomiasis, 75  
 Ametropia, 75  
 Amino Acid, 75  
 Amino Acid Catabolism, 75  
 Amino Acid Symbols, 76  
 Aminoaciduria, 76  
 Aminoglycoside Nephrotoxicity, 76  
 Aminoglycoside Ototoxicity, 76  
 Aminoglycosides, 76  
 Aminophylline Overdose, 76  
 Aminotransferase, 76  
 Amiodarone Pulmonary Toxicity, 76  
 Amiodarone-Induced Skin Discoloration, 76  
 Amiodarone-Induced Thyroid Dysfunction, 76  
 Amiodarone-Induced Vortex Keratopathy, 76  
 Amitriptyline, 76  
 Amitriptyline And Perphenazine Overdose, 76  
 Amitriptyline Hydrochloride Overdose, 77  
 Ammonia, 77  
 Ammonia Blood Test, 77  
 Ammonia Poisoning, 77  
 Ammonia Toxicity, 77  
 Ammonium Hydroxide Poisoning, 77  
 Amnesia, 77  
 Amnestic Syndrome, 77  
 Amniocentesis, 77  
 Amnioinfusion, 77  
 Amnion, 77  
 Amnion Nodosum, 77  
 Amnioscope, 77  
 Amniotic Band, 77  
 Amniotic Band Sequence, 78  
 Amniotic Band Syndrome, 78  
 Amniotic Fluid, 78  
 Amniotic Fluid Embolism, 78  
 Amniotic Fluid Index, 78  
 Amniotic Sac, 78  
 Amok, 78  
 AMP-Activated Protein Kinase, 78  
 Amphetamine, 78  
 Amphetamine Substance Use, 78  
 Amphetamine Toxicity, 78  
 Amphetamine-Related Psychiatric Disorders, 79  
 Amplified Musculoskeletal Pain Syndrome, 79  
 Ampulla, 79  
 Ampulla of Vater, 79  
 Ampullary Carcinoma, 79  
 Amputation, 79  
 Amputation Stump, 79  
 Amsler Grid, 79  
 Amusia, 79  
 Amygdala, 79  
 Amylase, 79  
 Amylase And Lipase, 79  
 Amylase Test, 79  
 Amylin Analogues, 80  
 Amyloid, 80  
 Amyloid Nephropathy, 80  
 Amyloid Neuropathy, 80  
 Amyloid PET, 80  
 Amyloid Plaque, 80  
 Amyloid Precursor Protein, 80  
 Amyloid Purpura, 80  
 Amyloidosis, 80  
 Amyotrophic Lateral Sclerosis, 80  
 Anabolic Steroid Abuse, 2110  
 Anabolic Steroid Induced Hypogonadism, 80  
 Anabolic Steroid Use and Abuse, 81  
 Anabolic Steroids, 81  
 Anabolism, 81  
 Anaemia, 81  
 Anaerobe, 81  
 Anaerobic, 81  
 Anaerobic Bacteria, 81  
 Anaerobic Glycolysis, 81  
 Anagen, 81  
 Anagen Effluvium, 81  
 Anal Abscess, 81  
 Anal Canal, 81  
 Anal Cancer, 81  
 Anal Carcinoma, 82

---

---

Anal Condyloma, [82](#)  
Anal Fissure, [82](#)  
Anal Fistula, [82](#)  
Anal Intercourse, [82](#)  
Anal Intraepithelial Neoplasia, [82](#)  
Anal Itching, [82](#)  
Anal Mucositis, [82](#)  
Anal Necrosis, [82](#)  
Anal Pain, [82](#)  
Anal Papilla, [82](#)  
Anal Sphincter, [82](#)  
Analgesia, [82](#)  
Analgesic, [2110](#)  
Analgesic Nephropathy, [2110](#)  
Analgesics, [82](#)  
Analog, [83](#)  
Analogous, [83](#)  
Analysis, [83](#)  
Analysis Of Urine, [83](#)  
Analytic Sensitivity, [83](#)  
Anaphia, [83](#)  
Anaphylactic Shock, [83](#)  
Anaphylactoid Reaction, [83](#)  
Anaphylatoxin, [83](#)  
Anaphylaxis, [83](#)  
Anaphylaxis During Anesthesia, [83](#)  
Anaplasia, [84](#)  
Anaplasmosis, [84](#)  
Anaplastic, [84](#)  
Anaplastic Astrocytoma, [84](#)  
Anaplastic Ependymoma, [84](#)  
Anaplastic Large Cell Lymphoma, [84](#)  
Anaplastic Large-cell Lymphoma, [84](#)  
Anaplastic Thyroid Cancer, [84](#)  
Anaplastic Thyroid Carcinoma, [84](#)  
Anarithmetia, [84](#)  
Anasarca, [84](#)  
Anastomosis, [85](#)  
Anastomotic Leak, [85](#)  
Anastrozole, [85](#)  
Anatomic Orientation Terms, [85](#)  
Anatomical Axis, [85](#)  
Anatomical Conduit, [85](#)  
Anatomical Orifice, [85](#)  
Anatomical Planes, [85](#)  
Anatomical Position, [85](#)  
Anatomical Snuffbox, [85](#)  
Anatomist, [85](#)  
Anatomy, [85](#)  
ANCA-Associated Vasculitis, [85](#)  
Ancillary Pathology Studies, [85](#)  
Ancillary Testing, Pathology, [86](#)  
Ancylostomiasis, [86](#)  
Andersen-Tawil Syndrome, [86](#)  
Anderson Disease, [86](#)  
Androgen, [86](#)  
Androgen Deficiency, [86](#)  
Androgen Excess, [86](#)  
Androgen Insensitivity Syndrome, [86](#)  
Androgen Insufficiency, [86](#)  
Androgen Receptor, [86](#)  
Androgen Suppression, [86](#)  
Androgenetic Alopecia, [87](#)  
Androgenic, [87](#)  
Android Pelvis, [87](#)  
Andropause, [87](#)  
Anejaculation, [87](#)  
Anemia, [87](#)  
Anemia and Thrombocytopenia in Pregnancy, [87](#)  
Anemia In Pregnancy, [87](#)  
Anemia Of Chronic Disease, [87](#)  
Anemia Of Inflammation, [87](#)  
Anemia of Malignancy, [87](#)  
Anemic, [87](#)  
Anencephaly, [88](#)  
Anergy, [88](#)  
Anesthesia, [88](#)  
Anesthesia Gas Analyzers, [88](#)  
Anesthesia Informatics, [88](#)  
Anesthesia Machine, [88](#)  
Anesthesia Monitoring, [88](#)  
Anesthesia Pharmacogenomics, [88](#)  
Anesthesiologist, [88](#)  
Anesthesiology, [88](#)  
Anesthetic, [2110](#)  
Anetoderma, [88](#)  
Aneuploidy, [89](#)  
Aneurysm, [89](#)  
Aneurysm Clip, [89](#)  
Aneurysm Clipping, [89](#)  
Aneurysm Coils, [89](#)  
Aneurysm Repair, [89](#)  
Angelman Syndrome, [89](#)  
Anger, [89](#)  
Angiitis, [89](#)  
Angina, [89](#)  
Angina Bullosa Hemorrhagica, [89](#)  
Angina Symptoms, [90](#)  
Angioblastoma, [90](#)  
Angiocardiology, [90](#)  
Angiodysplasia, [90](#)  
Angiodysplasia Of The Colon, [90](#)  
Angioedema, [90](#)  
Angioedema And Hives, [90](#)  
Angioedema, Hereditary, [920](#)  
Angiofibroma, [90](#)  
Angiofollicular Lymph Node Hyperplasia, [90](#)  
Angiogenesis, [90](#)

---

---

Angiogenesis Inhibition, 90  
Angiogenesis Inhibitor, 91  
Angiogram, 91  
Angiography, 91  
Angioid Streak, 91  
Angioimmunoblastic T-Cell Lymphoma, 91  
Angiokeratoma of the Scrotum, 91  
Angiolymphoid Hyperplasia With Eosinophilia, 91  
Angiomyofibroblastoma, 91  
Angiomyolipoma, 91  
Angioneurotic Edema, Hereditary, 920  
Angiopathy, 91  
Angioplasty, 91  
Angioplasty Balloon, 92  
Angiosarcoma, 92  
Angiosarcoma Of The Breast, 92  
Angiosarcoma Of The Heart, 92  
Angiotensin, 92  
Angiotensin Converting Enzyme, 92  
Angiotensin Converting-Enzyme Inhibitor, 2110  
Angiotensin I, 92  
Angiotensin II, 92  
Angiotensin Receptor Blocker, 2110  
Angiotensin Receptor Blockers, 92  
Angiotensin-Converting Enzyme 2, 92  
Angle-Closure Glaucoma, 92  
Angular Cheilitis, 92  
Anhedonia, 92  
Anhidrosis, 93  
Animal-Assisted Therapy, 93  
Anion, 93  
Anion Gap, 93  
Anion Gap Metabolic Acidosis, 93  
Aniridia, 93  
Anisocoria, 93  
Anisocytosis, 93  
Anisometropia, 93  
Ankle, 93  
Ankle Arthroplasty, 93  
Ankle Arthroscopy, 93  
Ankle Bone, 93  
Ankle Dislocation, 94  
Ankle Fracture, 94  
Ankle Fractures, 94  
Ankle Joint, 94  
Ankle Pain, 94  
Ankle Replacement, 94  
Ankle Sprain, 94  
Ankle Swelling, 94  
Ankle-Brachial Index, 94  
Ankle-Foot Orthosis, 94  
Ankyloblepharon, 94  
Ankyloglossia, 94  
Ankylose, 94  
Ankylosing, 94  
Ankylosing Spondylitis, 94  
Ankylosis, 95  
Ankyrin, 95  
Annihilation, 95  
Annular Pancreas, 95  
Annulus, 95  
Annulus Fibrosus, 95  
ANOCA, 95  
Anodontia, 95  
Anogenital Tract, 95  
Anomalous Anatomy, 95  
Anomalous Coronary Artery, 95  
Anomaly, 96  
Anomaly Scan, 96  
Anomia, 96  
Anophthalmia, 96  
Anorchia, 96  
Anorectal Abscess, 96  
Anorectal Fistula, 96  
Anorectal Infection, 96  
Anorectal Malformation, 96  
Anorectal Manometry, 96  
Anorexia, 96  
Anorexia Nervosa, 96  
Anorgasmia, 96  
Anorgasmy, 97  
Anoscopy, 97  
Anoscopy And High-Resolution Anoscopy, 97  
Anosmia, 97  
Anosodiaphoria, 97  
Anosognosia, 97  
Anotia, 97  
Anovulation, 97  
Anoxia, 97  
Anrep Effect, 97  
Anserine Bursa, 97  
Antacids, 97  
Antagonist, 97  
Antagonist Muscle, 98  
Antalgic Gait, 98  
Antebrachium, 98  
Antecubital Fossa, 98  
Antegrade, 98  
Antemortem Injury, 98  
Antenatal, 98  
Antepartum Fetal Surveillance, 98  
Antepartum Hemorrhage, 98  
Anterior, 98  
Anterior Cerebral Artery Syndrome, 98  
Anterior Chamber, 98  
Anterior Chamber Angle, 99  
Anterior Commissure, 99  
Anterior Cord Syndrome, 99

---

---

Anterior Cruciate Ligament, 99  
 Anterior Cruciate Ligament Injury, 99  
 Anterior Cruciate Ligament Tear, 99  
 Anterior Drawer Sign, 99  
 Anterior Drawer Test, 99  
 Anterior Interosseous Nerve, 99  
 Anterior Interosseous Nerve Syndrome, 99  
 Anterior Knee Pain, 99  
 Anterior Mediastinum, 99  
 Anterior Midline, 100  
 Anterior Myocardial Infarction, 100  
 Anterior Pituitary, 100  
 Anterior Talofibular Ligament, 100  
 Anterior Tibial Artery, 100  
 Anterior Uveitis, 100  
 Anterior Vaginal Prolapse, 100  
 Anterior Vaginal Wall Repair, 100  
 Anterior/Posterior Drawer Test, 100  
 Anterograde Amnesia, 100  
 Anterolisthesis, 100  
 Anteroposterior, 101  
 Anthocyanin, 101  
 Anthracosis, 101  
 Anthrax, 101  
 Anthrax Blood Test, 101  
 Anthropometry, 101  
 Anti-Anxiety Agents, 101  
 Anti-CCP, 101  
 Anti-CCP Antibody, 101  
 Anti-Citrullinated Peptide Antibody, 101  
 Anti-Cyclic Citrullinated Peptide Antibody, 101  
 Anti-D, 101  
 Anti-DNase B Blood Test, 101  
 Anti-Double-Stranded DNA Antibody, 102  
 Anti-GBM Disease, 102  
 Anti-Glomerular Basement Membrane Blood Test, 102  
 Anti-Glomerular Basement Membrane Disease, 102  
 Anti-HMGCR Myopathy, 102  
 Anti-IgE Therapy, 102  
 Anti-IL4 Receptor Therapy, 102  
 Anti-IL5 Therapy, 102  
 Anti-Infective, 2112  
 Anti-Inflammatory, 102  
 Anti-Insulin Antibody Test, 102  
 Anti-MDA5 Dermatomyositis, 102  
 Anti-Mullerian Hormone, 103  
 Anti-NMDA Receptor Encephalitis, 103  
 Anti-Reflux Surgery, 103  
 Anti-Reflux Surgery, Pediatric, 1479  
 Anti-Rust Product Poisoning, 103  
 Anti-Smooth Muscle Antibody, 103  
 Anti-SRP Myopathy, 103  
 Anti-Thyroid Peroxidase Antibody, 103  
 Anti-VEGF Injections, 103  
 Antiandrogen, 103  
 Antianginal, 103  
 Antiarrhythmic, 103  
 Antibacterial, 104  
 Antibigram, 104  
 Antibiotic, 2110  
 Antibiotic Cycling, 2110  
 Antibiotic De-Escalation, 2111  
 Antibiotic Misuse in Viral URIs, 104  
 Antibiotic Mixing, 2111  
 Antibiotic Resistance, 2111  
 Antibiotic Resistance Genes, 2111  
 Antibiotic Stewardship, 2111  
 Antibiotic-Associated Colitis, 2110  
 Antibiotic-Associated Diarrhea, 2110  
 Antibiotic-Resistant Bacterial Infection, 104  
 Antibiotics, 104  
 Antibodies, 104  
 Antibody, 104  
 Antibody Engineering, 104  
 Antibody Therapy, 104  
 Antibody Titer Blood Test, 104  
 Antibody-Dependent Cell-Mediated Cytotox., 104  
 Antibody-Drug Conjugate, 104  
 Anticholinergic, 104  
 Anticholinergic Burden, 105  
 Anticholinergic Cognitive Burden Scale, 2111  
 Anticholinergic Toxicity, 105  
 Anticholinergic Toxidrome, 105  
 Anticoagulant, 2111  
 Anticoagulant Agent, 2111  
 Anticoagulant Monitoring, 2111  
 Anticoagulant Rodenticides Poisoning, 2111  
 Anticoagulants and Thrombolytics in Pregnancy, 105  
 Anticoagulation, 105  
 Anticodon, 105  
 Anticonvulsant, 2111  
 Anticonvulsant Mood Stabilizers, 2111  
 Anticonvulsants, 105  
 Antideformity Position, 105  
 Antidepressant, 2111  
 Antidepressants, 105  
 Antidiabetic, 105  
 Antidiabetic Agent, 2112  
 Antidiabetic Drugs, 105  
 Antidiarrheal, 2112  
 Antidiarrheal Medicine Overdose, 2112  
 Antidiuretic Hormone, 105  
 Antidiuretic Hormone Blood Test, 105  
 Antidiuretic Hormone Physiology, 106  
 Antidiuretic Hormone Resistance, 106  
 Antidote, 106  
 Antiemetic, 2112  
 Antiemetics, 106

---



---

Antiepileptic Drugs, 106  
Antiestrogen, 2112  
Antifreeze Poisoning, 106  
Antifungal, 2112  
Antifungal Agent, 2112  
Antifungal Drug Classes, 2112  
Antifungal Medication, 2112  
Antifungal Prophylaxis, 106  
Antifungal Resistance, 106  
Antigen, 106  
Antigen Detection Test, 106  
Antigen Presentation, 106  
Antigen Receptor, 106  
Antigen-Presenting Cell, 106  
Antigenic Drift, 107  
Antigenic Shift, 107  
Antigenic Variation, 107  
Antigout, 107  
Antihistamine, 2112  
Antihistamines, 107  
Antihistamines For Allergies, 107  
Antihypertensive, 2112  
Antihypertensive Agent, 2112  
Antihypertensives In Pregnancy, 107  
Antileukotriene, 107  
Antimalarial, 2113  
Antimetabolite, 107  
Antimicrobial, 107  
Antimicrobial Adverse Effects, 107  
Antimicrobial Agent, 107  
Antimicrobial Desensitization, 107  
Antimicrobial Dosing Optimization, 108  
Antimicrobial Drug, 2113  
Antimicrobial Drug Interactions, 2113  
Antimicrobial Hypersensitivity Reactions, 108  
Antimicrobial Pharmacodynamics, 108  
Antimicrobial Pharmacokinetics, 108  
Antimicrobial Resistance, 108  
Antimicrobial Resistance Mechanisms, 108  
Antimicrobial Resistance Surveillance, 108  
Antimicrobial Sensitivity Test, 108  
Antimicrobial Stewardship, 108  
Antimicrobial Susceptibility Testing, 108  
Antimicrobial Therapy, 108  
Antimitochondrial Antibody, 108  
Antimony, 108  
Antineoplastic, 108  
Antineoplastic Drugs, 109  
Antinuclear Antibody, 109  
Antinuclear Antibody Panel, 109  
Antioxidant, 109  
Antiparasitic, 2113  
Antiparietal Cell Antibody Test, 109  
Antiparkinson, 109  
Antiparkinsonian, 2113  
Antiphospholipid Antibodies, 109  
Antiphospholipid Antibody, 109  
Antiphospholipid Antibody Syndrome, 109  
Antiphospholipid Syndrome, 109  
Antiphospholipid Syndrome and Pregnancy, 109  
Antiplatelet, 110  
Antiplatelet Agent, 2113  
Antiplatelet Agents, 110  
Antiplatelet Agents, P2Y12 Inhibitors, 1445  
Antiplatelet Therapy, 110  
Antiprotozoal, 110  
Antiprotozoal Drug, 2113  
Antipruritic, 2113  
Antipsychotic, 2113  
Antipsychotic Agent, 2113  
Antipsychotic Medicine, 110  
Antipsychotic-Induced Parkinsonism, 2113  
Antipyretic, 110  
Antiretroviral, 110  
Antiretroviral Therapy, 110  
Antisepsis, 110  
Antiseptic, 110  
Antiserum, 110  
Antisocial Personality, 110  
Antisocial Personality Disorder, 110  
Antispasmodic, 2113  
Antispasmodic Drugs, 110  
Antistreptolysin O Titer, 111  
Antisynthetase Syndrome, 111  
Antithrombin, 111  
Antithrombin Deficiency, 111  
Antithrombin III Blood Test, 111  
Antithrombin III Deficiency, 111  
Antithymocyte Globulin, 111  
Antithyroglobulin Antibody Test, 111  
Antithyroid, 2113  
Antithyroid Drug, 2113  
Antitoxin, 111  
Antitubercular Agents, 111  
Antitussive, 111  
Antiviral, 2114  
Antiviral Drug Classes, 2114  
Anton Syndrome, 111  
Antral Follicle Count, 112  
Antral Web, 112  
Antrochoanal Polyp, 112  
Antrum, 112  
Anuria, 112  
Anus, 112  
Anxiety, 112  
Anxiety & Panic, 112  
Anxiety Disorder, 112  
Anxiety Disorders, 112

---

---

Anxiolytic, [2114](#)  
Aorta, [112](#)  
Aortal, [112](#)  
Aortic, [112](#)  
Aortic Aneurysm, [112](#)  
Aortic Aneurysm Rupture, [113](#)  
Aortic Aneurysm Screening, [113](#)  
Aortic Angiography, [113](#)  
Aortic Arch, [113](#)  
Aortic Arch Syndrome, [113](#)  
Aortic Balloon Pump, [113](#)  
Aortic Body, [113](#)  
Aortic Coarctation, [113](#)  
Aortic Disease, [113](#)  
Aortic Dissection, [113](#)  
Aortic Embolism, [113](#)  
Aortic Graft Infection, [113](#)  
Aortic Injury, [114](#)  
Aortic Insufficiency, [114](#)  
Aortic Regurgitation, [114](#)  
Aortic Regurgitation Murmur, [114](#)  
Aortic Rupture, [114](#)  
Aortic Sclerosis, [114](#)  
Aortic Stenosis, [114](#)  
Aortic Stenosis Grading, [114](#)  
Aortic Valve, [114](#)  
Aortic Valve Disease, [114](#)  
Aortic Valve Insufficiency, [114](#)  
Aortic Valve Regurgitation, [114](#)  
Aortic Valve Replacement, [114](#)  
Aortic Valve Stenosis, [115](#)  
Aortic Valve Surgery, Minimally Invasive, [1264](#)  
Aortic Valve Surgery, Open, [1404](#)  
Aortitis, [115](#)  
Aortocaval Compression Syndrome, [115](#)  
Aortoenteric Fistula, [115](#)  
Aortography, [115](#)  
Aortoiliac Occlusive Disease, [115](#)  
Aortopulmonary Window, [115](#)  
AP X-Ray, [115](#)  
APACHE II Score, [115](#)  
Apathy, [115](#)  
APC Tumor Suppressor, [115](#)  
Apert Syndrome, [115](#)  
Apex, [115](#)  
Apex Beat, [116](#)  
Apgar Score, [116](#)  
Aphagia, [116](#)  
Aphakia, [116](#)  
Aphantasia, [116](#)  
Aphasia, [116](#)  
Apheresis, [116](#)  
Apheresis Donation, [116](#)  
Aphonia, [116](#)  
Aphrasia, [116](#)  
Aphthous Stomatitis, [116](#)  
Aphthous Ulcer, [116](#)  
Apical, [116](#)  
Apical Ballooning Syndrome, [116](#)  
Apical Periodontitis, [117](#)  
Apical Pulse, [117](#)  
Apicoectomy, [117](#)  
Aplasia, [117](#)  
Aplasia Cutis Congenita, [117](#)  
Aplastic, [117](#)  
Aplastic Anemia, [117](#)  
Aplastic Crisis, [117](#)  
Apley Compression Test, [117](#)  
Apley Distraction Test, [117](#)  
Apley Grind Test, [117](#)  
Apley Scratch Test, [118](#)  
APLS, [118](#)  
Apnea, [118](#)  
Apnea of Infancy, [118](#)  
Apnea Of Prematurity, [118](#)  
Apnea, Infantile, [118](#)  
Apnea-Hypopnea Index, [118](#)  
Apneic, [118](#)  
Apneustic Breathing, [118](#)  
Apnoea, [118](#)  
Apocrine Hidrocystoma, [118](#)  
Apoenzyme, [118](#)  
Apolipoprotein B100, [118](#)  
Apolipoprotein CII, [119](#)  
Apolipoproteins, [119](#)  
Aponeurosis, [119](#)  
Apophenia, [119](#)  
Apophyses, [119](#)  
Apoptosis, [119](#)  
Apoptosis Extrinsic Pathway, [119](#)  
Apoptosis Intrinsic Pathway, [119](#)  
Apoptotic Body, [119](#)  
Appalachian Type Amyloidosis, [119](#)  
Apparent Mineralocorticoid Excess, [119](#)  
Appendageal, [119](#)  
Appendectomy, [119](#)  
Appendiceal, [119](#)  
Appendiceal Adenocarcinoma, [120](#)  
Appendiceal Cancer, [120](#)  
Appendiceal Rupture, [120](#)  
Appendiceal Tumors, [120](#)  
Appendicitis, [120](#)  
Appendicular Skeleton, [120](#)  
Appendix, [120](#)  
Appendix Cancer, [120](#)  
Appendix Testis, [120](#)  
Appetite, [120](#)  
Appetite Regulation, [120](#)

---

---

Applied Behavior Analysis, [120](#)  
 Apposition, [120](#)  
 Apprehension Test, [120](#)  
 Appropriate For Gestational Age, [121](#)  
 Apraxia, [121](#)  
 Apraxia of Speech, [121](#)  
 Apraxic Gait, [121](#)  
 Apraxia, [121](#)  
 APS, [121](#)  
 Aquagenic Keratoderma, [121](#)  
 Aquaporin, [121](#)  
 Aquatic Therapy, [121](#)  
 Aqueous Flare, [121](#)  
 Aqueous Humor, [122](#)  
 Arachidonic Acid Pathway, [122](#)  
 Arachnitis, [122](#)  
 Arachnodactyly, [122](#)  
 Arachnoid Cyst, [122](#)  
 Arachnoid Cysts, [122](#)  
 Arachnoid Granulations, [122](#)  
 Arachnoid Mater, [122](#)  
 Arachnoidal Fibroblastoma, [122](#)  
 Arachnoiditis, [122](#)  
 ARB-Nephrilysin Inhibitor, [2114](#)  
 Arbovirus, [122](#)  
 Arbovirus A Chikungunya Type, [122](#)  
 ARBs, [123](#)  
 Arcanobacterium haemolyticum Infection, [123](#)  
 Arciform, [123](#)  
 Arcuate Uterus, [123](#)  
 Arcus Senilis, [123](#)  
 ARDS, [123](#)  
 Area Under The Concentration-Time Curve, [123](#)  
 Aregenerative Anemia, [123](#)  
 Arenaviruses, [123](#)  
 Areola, [123](#)  
 ARG Deficiency, [123](#)  
 Argentine Hemorrhagic Fever, [123](#)  
 Argininosuccinate Lyase Deficiency, [123](#)  
 Argininosuccinic Aciduria, [124](#)  
 Argon Plasma Coagulation, [124](#)  
 Argyll Robertson Pupil, [124](#)  
 Arias-Stella Reaction, [124](#)  
 Arm, [124](#)  
 Arm CT Scan, [124](#)  
 Arm Fracture, [124](#)  
 Arm MRI Scan, [124](#)  
 Arm Pain, [124](#)  
 Armpit Lump, [124](#)  
 Arnold-Chiari Malformation, [124](#)  
 Aromatase, [124](#)  
 Aromatase Excess Syndrome, [124](#)  
 Aromatase Inhibitor, [2114](#)  
 Arousal Index, [125](#)  
 Arrector Pili, [125](#)  
 Arrector Pili Muscle, [125](#)  
 Arrested Caries, [125](#)  
 Arrhythmia, [125](#)  
 Arrhythmias, [125](#)  
 Arrhythmogenic Right Ventricular Cardiomyopathy, [125](#)  
 Arsenic, [125](#)  
 Arsenic Poisoning, [125](#)  
 Arsenic Toxicity, [125](#)  
 Artemisinin, [125](#)  
 Arteria Femoralis, [125](#)  
 Arterial Blood Gas, [125](#)  
 Arterial Blood Gas Analyzer, [126](#)  
 Arterial Catheter, [126](#)  
 Arterial Catheterization, [126](#)  
 Arterial Dissection, [126](#)  
 Arterial Duplex, [126](#)  
 Arterial Embolism, [126](#)  
 Arterial Embolization, [126](#)  
 Arterial Injury, [126](#)  
 Arterial Insufficiency, [126](#)  
 Arterial Line, [126](#)  
 Arterial Resistance, [126](#)  
 Arterial Spin Labeling, [126](#)  
 Arterial Stenosis, [126](#)  
 Arterial Stick, [126](#)  
 Arterial Stiffness, [127](#)  
 Arterial Switch Operation, [127](#)  
 Arterial Tension, [127](#)  
 Arterial Thromboembolism, [127](#)  
 Arterial Thrombosis, [127](#)  
 Arterial Ulcer, [127](#)  
 Arteries, [127](#)  
 Arteriogram, [127](#)  
 Arteriole, [127](#)  
 Arterioles, [127](#)  
 Arteriolosclerosis, [127](#)  
 Arteriosclerosis, [127](#)  
 Arteriosclerosis / Atherosclerosis, [127](#)  
 Arteriosclerotic Retinopathy, [128](#)  
 Arteriovenous Fistula, [128](#)  
 Arteriovenous Fistula Creation, [128](#)  
 Arteriovenous Graft, [128](#)  
 Arteriovenous Malformation, [128](#)  
 Arteriovenous Malformation Resection, [128](#)  
 Arteriovenous Nicking, [128](#)  
 Arteritis, [128](#)  
 Arteritis, Giant Cell, [827](#)  
 Arteritis, Takayasu, [1887](#)  
 Arteritis, Takayasu's, [1887](#)  
 Artery, [128](#)  
 Artery Occlusion, [128](#)  
 Arthralgia, [129](#)  
 Arthritis, [129](#)

---

---

Arthritis (Osteoarthritis), [129](#)  
Arthritis of the Wrist, [129](#)  
Arthritis Urethritica, [129](#)  
Arthritis, Ankylosing Spondylitis, [95](#)  
Arthritis, Basal Joint, [1919](#)  
Arthritis, Degenerative, [1426](#)  
Arthritis, Gouty, [853](#)  
Arthritis, Infectious, [1014](#)  
Arthritis, Juvenile Idiopathic, [1079](#)  
Arthritis, Juvenile Rheumatoid, [1079](#)  
Arthritis, Lyme, [1135](#)  
Arthritis, Mixed Connective Tissue Disease, [1273](#)  
Arthritis, Osteoarthritis, [1426](#)  
Arthritis, Plant Thorn, [1541](#)  
Arthritis, Pseudogout, [298](#)  
Arthritis, Psoriatic, [1622](#)  
Arthritis, Reactive, [1672](#)  
Arthritis, Reiter, [1672](#)  
Arthritis, Rheumatoid, [1711](#)  
Arthritis, Sarcoid, [1744](#)  
Arthritis, Scleroderma, [1879](#)  
Arthritis, Septic, [1770](#)  
Arthritis, Sjogren, [1794](#)  
Arthritis, Still, [1843](#)  
Arthritis, Systemic Lupus Erythematosus, [1112](#)  
Arthritis, Thumb, [1919](#)  
Arthrocentesis, [129](#)  
Arthrodesis, [129](#)  
Arthrogram, [129](#)  
Arthrography, [129](#)  
Arthrogryposis, [129](#)  
Arthrogryposis Multiplex Congenita, [129](#)  
Arthrolysis, [129](#)  
Arthropathy, [129](#)  
Arthroplasty, [130](#)  
Arthropod Vector, [130](#)  
Arthroscope, [130](#)  
Arthroscopy, [130](#)  
Arthrosis, [130](#)  
Arthus Reaction, [130](#)  
Articular Cartilage, [130](#)  
Articular Cartilage Injuries, [130](#)  
Articular Ligament, [130](#)  
Articulation, [130](#)  
Artifacts, [130](#)  
Artificial Abortion, [130](#)  
Artificial Heart, [130](#)  
Artificial Insemination, [130](#)  
Artificial Insemination By Donor, [130](#)  
Artificial Insemination By Husband, [131](#)  
Artificial Intelligence, [131](#)  
Artificial Kidney, [131](#)  
Artificial Pacemaker, [131](#)  
Artificial Rupture Of Membranes, [131](#)  
Artificial Saliva, [131](#)  
Artificial Skin, [131](#)  
Artificial Sweeteners, [131](#)  
Artificial Urinary Sphincter, [131](#)  
Artificial Ventilation, [131](#)  
Artificial Ventricle, [131](#)  
Aryepiglottic Fold, [131](#)  
Arytenoid Cartilage, [132](#)  
ASA Physical Status Classification System, [132](#)  
Asbestos-Related Disorders, [132](#)  
Asbestosis, [132](#)  
Ascariasis, [132](#)  
Ascending Aorta, [132](#)  
Ascending Aortic Aneurysm, [132](#)  
Ascending Cholangitis, [132](#)  
Ascending Colon, [132](#)  
Ascites, [132](#)  
Ascitic Fluid, [132](#)  
Ascorbic Acid, [132](#)  
Ascus, [133](#)  
Asepsis, [133](#)  
Aseptic, [133](#)  
Aseptic Loosening, [133](#)  
Aseptic Meningitis, [133](#)  
Aseptic Necrosis, [133](#)  
Aseptic Technique, [133](#)  
Asherman Syndrome, [133](#)  
Ashman Phenomenon, [133](#)  
ASL, [133](#)  
ASOT, [133](#)  
Asp, [133](#)  
Aspartame, [133](#)  
Aspartate Aminotransferase, [134](#)  
Aspartate Aminotransferase Blood Test, [134](#)  
Aspartic Acid, [134](#)  
Asperger Syndrome, [134](#)  
Aspergillosis, [134](#)  
Aspergillosis Precipitin, [134](#)  
Aspergillus, [134](#)  
Aspergillus Infection, [134](#)  
Asphalt Cement Poisoning, [134](#)  
Asphyxia, [134](#)  
Asphyxia Neonatorum, [134](#)  
Asphyxial Death Pathology, [134](#)  
Asphyxiating Thoracic Dysplasia, [135](#)  
Aspirate, [135](#)  
Aspiration, [135](#)  
Aspiration Pneumonia, [135](#)  
Aspiration Pneumonitis, [135](#)  
Aspiration Pneumonitis and Pneumonia, [135](#)  
Aspiration Risk, [135](#)  
Aspirin, [135](#)  
Aspirin Overdose, [135](#)  
Aspirin Poisoning, [135](#)

---

---

Aspirin-Exacerbated Respiratory Disease, [135](#)  
Assay, [135](#)  
Assist-Control Ventilation, [136](#)  
Assisted Breech Delivery, [136](#)  
Assisted Circulation, [136](#)  
Assisted Delivery With Forceps, [136](#)  
Assisted Living, [136](#)  
Assisted Reproductive Technology, [136](#)  
Assistive Device, [136](#)  
Assistive Device Training, [136](#)  
Association, [136](#)  
Association Cortex, [136](#)  
Association Study, [136](#)  
Asteatotic Eczema, [136](#)  
Astereognosis, [136](#)  
Asterixis, [137](#)  
Asteroid Hyalosis, [137](#)  
Asthenia, [137](#)  
Asthenic, [137](#)  
Asthenopia, [137](#)  
Asthenozoospermia, [137](#)  
Asthma, [137](#)  
Asthma Attack, [137](#)  
Asthma Controller Medications, [137](#)  
Asthma Exacerbation, [137](#)  
Asthma in Older Adults, [137](#)  
Asthma in Pregnancy, [137](#)  
Asthma Medications, [138](#)  
Asthma Quick-Relief Medications, [138](#)  
Asthma, Adult-Onset, [50](#)  
Asthma, Exercise-Induced, [693](#)  
Asthma, Occupational, [1388](#)  
Asthma-COPD Overlap, [138](#)  
Astigmatism, [138](#)  
Astringent, [138](#)  
Astrocyte, [138](#)  
Astrocytoma, [138](#)  
Astrovirus Infection, [138](#)  
Asymmetrical, [138](#)  
Asymptomatic, [138](#)  
Asymptomatic Adults, [138](#)  
Asymptomatic Bacteriuria, [138](#)  
Asymptomatic Bacteriuria In Pregnancy, [139](#)  
Asymptomatic Carrier, [139](#)  
Asymptomatic HIV Infection, [139](#)  
Asymptomatic Infections, [139](#)  
Asystole, [139](#)  
At-Home COVID-19 Testing, [139](#)  
Ataxia, [139](#)  
Ataxia Telangiectasia, [139](#)  
Ataxic Breathing, [139](#)  
Ataxic Gait, [139](#)  
Atelectasis, [140](#)  
Athelia, [140](#)  
Atherectomy, [140](#)  
Atheroembolic Renal Disease, [140](#)  
Atheroma, [140](#)  
Atheromatous Plaque, [140](#)  
Atherosclerosis, [140](#)  
Atherosclerosis Pathology, [140](#)  
Atherosclerotic, [140](#)  
Atherosclerotic Heart Disease, [140](#)  
Atherosclerotic Plaque, [140](#)  
Atherothrombotic Stroke, [140](#)  
Athetosis, [141](#)  
Athlete Foot, [141](#)  
Athlete's Foot, [141](#)  
Athlete's Heart, [141](#)  
Atlanto-Occipital Dislocation, [141](#)  
Atlanto-Occipital Joints, [141](#)  
Atlantoaxial Instability, [141](#)  
Atlantoaxial Instability in Down Syndrome, [141](#)  
Atlantoaxial Joint, [141](#)  
Atlas, [141](#)  
Atom, [141](#)  
Atonic, [141](#)  
Atonic Seizure, [142](#)  
Atopic, [142](#)  
Atopic Dermatitis, [142](#)  
Atopic Eruption Of Pregnancy, [142](#)  
Atopic Rhinitis, [142](#)  
Atopy, [142](#)  
Atorvastatin, [142](#)  
Atovaquone–Proguanil, [142](#)  
ATP, [142](#)  
ATP Synthase, [142](#)  
Atransferrinemia, [142](#)  
Atresia, [142](#)  
Atresia Of Bile Ducts, [142](#)  
Atria, [142](#)  
Atrial, [143](#)  
Atrial Ablation, [143](#)  
Atrial Arrhythmia, [143](#)  
Atrial Fib, [143](#)  
Atrial Fibrillation, [143](#)  
Atrial Fibrillation Ablation, [143](#)  
Atrial Fibrillation Medicines, [143](#)  
Atrial Flutter, [143](#)  
Atrial Myxoma, [143](#)  
Atrial Natriuretic Peptide, [143](#)  
Atrial Septal Defect, [143](#)  
Atrial Septum, [143](#)  
Atrial Standstill, [143](#)  
Atrial Switch Operation, [144](#)  
Atrial Tachycardia, [144](#)  
Atrial Thrombosis, [144](#)  
Atrioventricular, [144](#)  
Atrioventricular Block, [144](#)

---

---

Atrioventricular Bundle, [144](#)  
Atrioventricular Canal Defect, [144](#)  
Atrioventricular Dissociation, [144](#)  
Atrioventricular Nodal Reentrant Tachycardia, [144](#)  
Atrioventricular Nodal Reentry Tachycardia, [144](#)  
Atrioventricular Node, [144](#)  
Atrioventricular Septal Defect, [145](#)  
Atrioventricular Valves, [145](#)  
Atrium, [145](#)  
Atrophic Gastritis, [145](#)  
Atrophic Glossitis, [145](#)  
Atrophic Vaginitis, [145](#)  
Atrophoderma Maculatum, [145](#)  
Atrophy, [145](#)  
Atrophy, Vaginal, [2017](#)  
Atropine, [145](#)  
Atropine In Organophosphate Poisoning, [145](#)  
Attachment Disorder, [145](#)  
Attachment, Dental, [526](#)  
Attack Rate, [145](#)  
Attention, [145](#)  
Attention Deficit Hyperactivity Disorder, [145](#)  
Attention-Deficit/Hyperactivity Disorder, [145](#)  
Attenuated, [146](#)  
Attenuated Vaccine, [2114](#)  
Attenuation Correction, [146](#)  
Attributable Risk, [146](#)  
Atypia, [146](#)  
Atypia Morphology, [146](#)  
Atypical, [146](#)  
Atypical (Dysplastic) Nevus, [146](#)  
Atypical Antipsychotics, [146](#)  
Atypical Endometrial Hyperplasia, [146](#)  
Atypical Facial Pain, [146](#)  
Atypical Features, [146](#)  
Atypical Fibroxanthoma, [146](#)  
Atypical Genitalia, [146](#)  
Atypical Hemolytic Uremic Syndrome, [147](#)  
Atypical Hyperplasia Of The Breast, [147](#)  
Atypical Lymphocyte, [147](#)  
Atypical Mole, [147](#)  
Atypical Nevi, [147](#)  
Atypical Pneumonia, [147](#)  
Atypical Spitz Nevus, [147](#)  
Atypical Teratoid Rhabdoid Tumor, [147](#)  
Audiogram, [147](#)  
Audiologist, [147](#)  
Audiology, [147](#)  
Audiometry, [147](#)  
Auditory Acuity, [147](#)  
Auditory Area, [147](#)  
Auditory Fatigue, [148](#)  
Auditory Hallucinations, [148](#)  
Auditory Information Processing Disorder, [148](#)  
Auditory Nerve, [148](#)  
Auditory Neuropathy Spectrum Disorder, [148](#)  
Auditory Ossicle, [148](#)  
Auditory Pathway, [148](#)  
Auditory Perception, [148](#)  
Auditory Processing Disorder, [148](#)  
Auditory Threshold, [148](#)  
Auer Rods, [148](#)  
Auerbach Plexus, [148](#)  
Auger Electrons, [149](#)  
Augmentation Mammoplasty, [149](#)  
Augmentation Mammoplasty, [149](#)  
Augmentation Of Labor, [149](#)  
Aura, [149](#)  
Aural Fullness, [149](#)  
Aural Polyps, [149](#)  
Auricle, [149](#)  
Auricular Hematoma, [149](#)  
Auscultation, [149](#)  
Auscultatory Gap, [149](#)  
Auspitz Sign, [149](#)  
Austin Flint Murmur, [150](#)  
Autism, [150](#)  
Autism Spectrum Disorder, [150](#)  
Autism Spectrum Disorder, Childhood Disintegrative Disorder, [380](#)  
Autistic Savant, [150](#)  
Autoantibodies, [150](#)  
Autoantibody, [150](#)  
Autoantigen, [150](#)  
Autoclave, [150](#)  
Autocrine Signaling, [150](#)  
Autoerythrocyte Sensitivity, [150](#)  
Autogenic Drainage, [150](#)  
Autograft, [150](#)  
Autoimmune, [150](#)  
Autoimmune Associated Epilepsy, [151](#)  
Autoimmune Autonomic Ganglionopathy, [151](#)  
Autoimmune Disease, [151](#)  
Autoimmune Diseases, [151](#)  
Autoimmune Disorder, [151](#)  
Autoimmune Encephalitis, [151](#)  
Autoimmune Encephalopathy, [151](#)  
Autoimmune Endocrine Disease, [151](#)  
Autoimmune Enteropathy, [151](#)  
Autoimmune Epilepsy, [151](#)  
Autoimmune Gastritis, [151](#)  
Autoimmune Hemolytic Anemia, [151](#)  
Autoimmune Hepatitis, [151](#)  
Autoimmune Keratitis, [151](#)  
Autoimmune Liver Disease Panel, [152](#)  
Autoimmune Lymphoproliferative Syndrome, [152](#)  
Autoimmune Pancreatitis, [152](#)  
Autoimmune Polyendocrine Syndromes, [152](#)

---



---

Autoimmune Polyendocrinopathy, [152](#)  
 Autoimmune Polyglandular Syndrome, [152](#)  
 Autoimmune Thyroid Disease and Pregnancy, [152](#)  
 Autoimmune Thyroiditis, [152](#)  
 Autoimmunity, [152](#)  
 Autoimmunity: Mechanisms, [152](#)  
 Autoinflammatory Diseases, [152](#)  
 Autoinoculation, [152](#)  
 Autologous, [153](#)  
 Autologous Blood Donation, [153](#)  
 Autologous Breast Reconstruction, [153](#)  
 Autologous Fat Transplant, [153](#)  
 Autologous Stem Cell Transplantation, [153](#)  
 Autolysis, [153](#)  
 Automated External Defibrillator, [153](#)  
 Automaticity, [153](#)  
 Autonomic Disorder, [153](#)  
 Autonomic Dysreflexia, [153](#)  
 Autonomic Ganglion, [153](#)  
 Autonomic Nervous System, [153](#)  
 Autonomic Neuropathy, [154](#)  
 Autonomic Neuropathy, Diabetic, [543](#)  
 Autonomy, Patient, [1475](#)  
 Autophagy, [154](#)  
 Autopsy, [154](#)  
 Autopsy, Medicolegal, [769](#)  
 Autoradiography, [154](#)  
 Autosomal Dominant, [154](#)  
 Autosomal Dominant Inheritance, [154](#)  
 Autosomal Dominant PKD, [154](#)  
 Autosomal Dominant Polycystic Kidney Disease, [154](#)  
 Autosomal Dominant Tubulointerstitial Kidney Disease, [154](#)  
 Autosomal Inheritance, [154](#)  
 Autosomal Recessive, [154](#)  
 Autosomal Recessive Disorders, [154](#)  
 Autosomal Recessive Inheritance, [155](#)  
 Autosomal Recessive PKD, [155](#)  
 Autosomal Recessive Polycystic Kidney Disease, [155](#)  
 Autosome, [155](#)  
 Avascular Necrosis, [155](#)  
 Avastin, [155](#)  
 Avian Flu, [155](#)  
 Avian Influenza, [155](#)  
 AVM, [155](#)  
 AVN, [155](#)  
 Avoidance Behavior, [155](#)  
 Avoidant Personality Disorder, [155](#)  
 Avoidant/Restrictive Food Intake Disorder, [155](#)  
 Avolition, [156](#)  
 Avulsed Tooth, [156](#)  
 Avulsion, [156](#)  
 Avulsion Fracture, [156](#)  
 Avuncular, [156](#)  
 Awake Craniotomy, [156](#)  
 Awake Fiberoptic Intubation, [156](#)  
 Awareness, [156](#)  
 Awareness Under Anesthesia, [156](#)  
 Axial Skeleton, [156](#)  
 Axial Spondyloarthritis, [157](#)  
 Axial Tomography, Computerized, [441](#)  
 Axilla, [157](#)  
 Axillary, [157](#)  
 Axillary Artery, [157](#)  
 Axillary Dissection, [157](#)  
 Axillary Lymph Node, [157](#)  
 Axillary Lymphadenectomy, [157](#)  
 Axillary Lymphadenopathy, [157](#)  
 Axillary Nerve, [157](#)  
 Axillary Nerve Dysfunction, [157](#)  
 Axillary Tail Of The Breast, [157](#)  
 Axillary Vein, [157](#)  
 Axis, [157](#)  
 Axis Vertebra, [158](#)  
 Axolemma, [158](#)  
 Axon, [158](#)  
 Axon Fasciculation, [158](#)  
 Axon Guidance, [158](#)  
 Axon Hillock, [158](#)  
 Axon Terminal, [158](#)  
 Axonal Neuropathy, Giant, [827](#)  
 Axonal Transport, [158](#)  
 Axoneme, [158](#)  
 Axonotmesis, [158](#)  
 Ayurveda, [158](#)  
 Ayurvedic Medicine, [158](#)  
 Azole Antifungals, [158](#)  
 Azoospermia, [158](#)  
 Azotaemia, [159](#)  
 Azotemia, [159](#)  
 Azygos Vein, [159](#)  
 Azygous Vein, [159](#)  
 B And T Cell Screen, [161](#)  
 B And T Cells, [161](#)  
 B And T Lymphocytes, [161](#)  
 B Cell, [161](#)  
 B Hemophilia, [161](#)  
 B Lymphocyte, [161](#)  
 B Virus, [161](#)  
 B-Cell Activation, [161](#)  
 B-Cell Acute Lymphoblastic Leukemia, [161](#)  
 B-Cell Acute Lymphoblastic Leukemia for Adults, [161](#)  
 B-Cell Leukemia/Lymphoma Panel, [161](#)  
 B-Cell Lymphoma, [162](#)  
 B-Cell Receptor, [162](#)  
 B-Lymphocyte, [162](#)  
 B-Lynch Suture, [162](#)

---

---

B-Mode Ultrasound, [162](#)  
B-Type Natriuretic Peptide, [162](#)  
Ba, [162](#)  
Babcock Clamp, [162](#)  
Babesia, [162](#)  
Babesia Divergens, [162](#)  
Babesia Microti, [162](#)  
Babesiosis, [162](#)  
Babinski Reflex, [162](#)  
Babinski Response, [162](#)  
Babinski Sign, [163](#)  
Baby Acne, [163](#)  
Baby Blues, [163](#)  
Baby Health, [163](#)  
Baby Holding Techniques, [163](#)  
Babyhood, [163](#)  
Bacillaceae, [163](#)  
Bacillary Angiomatosis, [163](#)  
Bacillary Dysentery, [163](#)  
Bacille Calmette-Guérin, [163](#)  
Bacille Calmette-Guérin Vaccine, [163](#)  
Bacilluria, [163](#)  
Bacillus, [163](#)  
Bacillus Anthracis, [163](#)  
Bacillus Cereus, [164](#)  
Bacillus Cereus Food Poisoning, [164](#)  
Bacitracin, [164](#)  
Bacitracin Overdose, [164](#)  
Bacitracin Zinc Overdose, [164](#)  
Back Injury, [164](#)  
Back Labor, [164](#)  
Back Pain, [164](#)  
Back Pressure, [164](#)  
Back Spasms, [164](#)  
Back Strains and Sprains, [164](#)  
Back-Blows, [164](#)  
Backache, [165](#)  
Backwash Ileitis, [165](#)  
Baclofen, [165](#)  
Bacteraemia, [165](#)  
Bacteremia, [165](#)  
Bacteria, [165](#)  
Bacterial Adherence, [165](#)  
Bacterial Arthritis, [165](#)  
Bacterial Biofilm, [165](#)  
Bacterial Capsule, [165](#)  
Bacterial Cell Wall, [165](#)  
Bacterial Classification, [165](#)  
Bacterial Conjugation, [165](#)  
Bacterial Conjunctivitis, [166](#)  
Bacterial Dermatitis, [166](#)  
Bacterial Dormancy, [166](#)  
Bacterial Endocarditis, [166](#)  
Bacterial Endocarditis Vegetations, [166](#)  
Bacterial Endospores, [166](#)  
Bacterial Flagella, [166](#)  
Bacterial Gastroenteritis, [166](#)  
Bacterial Genetics, [166](#)  
Bacterial Growth Phases, [166](#)  
Bacterial Infection, [166](#)  
Bacterial Infections and Pregnancy, [166](#)  
Bacterial Meningitis, [167](#)  
Bacterial Meningococcal Meningitis, [167](#)  
Bacterial Metabolism, [167](#)  
Bacterial Morphology, [167](#)  
Bacterial Overgrowth, [167](#)  
Bacterial Pathogenesis, [167](#)  
Bacterial Peritonitis, [167](#)  
Bacterial Pharyngitis, [167](#)  
Bacterial Pili, [167](#)  
Bacterial Pneumonia, [167](#)  
Bacterial Prostatitis, [167](#)  
Bacterial Protein, [167](#)  
Bacterial Spore, [167](#)  
Bacterial Toxemia, [168](#)  
Bacterial Toxin, [168](#)  
Bacterial Toxins, [168](#)  
Bacterial Tracheitis, [168](#)  
Bacterial Transduction, [168](#)  
Bacterial Transformation, [168](#)  
Bacterial Vaccine, [2114](#)  
Bacterial Vaginosis, [168](#)  
Bacterial Virulence Factors, [168](#)  
Bactericidal, [168](#)  
Bactericide, [168](#)  
Bacteriocin, [168](#)  
Bacteriology, [168](#)  
Bacteriophage, [168](#)  
Bacteriophage Lambda, [168](#)  
Bacteriophage T4, [169](#)  
Bacteriophages, [169](#)  
Bacteriorhodopsin, [169](#)  
Bacteriostasis, [169](#)  
Bacteriostatic, [169](#)  
Bacterium, [169](#)  
Bacteriuria, [169](#)  
Bacteroidaceae, [169](#)  
Bacteroides, [169](#)  
Bacteroides Caccaae, [169](#)  
Bacteroides Dorei, [169](#)  
Bacteroides Fragilis, [169](#)  
Bacteroides Gingivalis, [169](#)  
Bacteroides Ovatus, [169](#)  
Bacteroides Thetaiotaomicron, [170](#)  
Bacteroides Ureolyticus, [170](#)  
Bacteroides Vulgatus, [170](#)  
Bad Breath, [170](#)  
BAER, [170](#)

---

---

Bag-Valve Mask, [170](#)  
Bag-Valve-Mask Ventilation, [170](#)  
Bagassosis, [170](#)  
Bags Under Eyes, [170](#)  
Bainbridge Reflex, [170](#)  
Bak Protein, [170](#)  
Baker Cyst, [170](#)  
Baker's Cyst, [171](#)  
Baking Powder Overdose, [171](#)  
Balamuthia, [171](#)  
Balamuthia Mandrillaris, [171](#)  
Balamuthia Mandrillaris Infection, [171](#)  
Balance, [171](#)  
Balance Assessment, [171](#)  
Balance Disorders, [171](#)  
Balance Disorders And Vertigo, [171](#)  
Balance Impairment, [171](#)  
Balance Problems, [171](#)  
Balance Training, [171](#)  
Balanced Study, [171](#)  
Balanitis, [172](#)  
Balanitis Xerotica Obliterans, [172](#)  
Balanitis, Non-Infectious / Infectious, [172](#)  
Balanoposthitis, [172](#)  
Balantidiasis, [172](#)  
Balantidium, [172](#)  
Balantidium Coli, [172](#)  
Balbiani Body, [172](#)  
Baldness, [172](#)  
Balint's Syndrome, [172](#)  
Ball-And-Socket Joint, [172](#)  
Ballance Sign, [172](#)  
Ballard Score, [172](#)  
Ballism, [172](#)  
Ballismus, [173](#)  
Balloon Aortic Valvuloplasty, [173](#)  
Balloon Brachytherapy, [173](#)  
Balloon Catheter, [173](#)  
Balloon Dilation, [173](#)  
Balloon Occlusion, [173](#)  
Balloon Tamponade, [173](#)  
Balneology, [173](#)  
Balneotherapy, [173](#)  
Balo Disease, [173](#)  
Balsam, [173](#)  
Bamboo Spine, [173](#)  
Band Cell, [174](#)  
Band Heterotopia, [174](#)  
Band Keratopathy, [174](#)  
BANF, [174](#)  
Bankart Lesion, [174](#)  
Bankart Repair, [174](#)  
Bannayan-Riley-Ruvalcaba Syndrome, [174](#)  
Banti Syndrome, [174](#)  
Bar Graph, [174](#)  
Barakat Syndrome, [174](#)  
Barber Itch, [174](#)  
Barber Say Syndrome, [174](#)  
Barbiturate, [174](#)  
Barbiturate Intoxication And Overdose, [174](#)  
Barbiturate Toxicity, [175](#)  
Barbiturates, [175](#)  
Bardet-Biedl Syndrome, [175](#)  
Bare-Metal Stent, [175](#)  
Bariatric Embolization, [175](#)  
Bariatric Surgery, [175](#)  
Bariatrics, [175](#)  
Barium, [175](#)  
Barium Enema, [175](#)  
Barium Meal, [175](#)  
Barium Study, [175](#)  
Barium Sulfate, [175](#)  
Barium Swallow, [176](#)  
Barium-133, [176](#)  
Barlow Maneuver, [176](#)  
Barlow Sign, [176](#)  
Barlow Test, [176](#)  
Barlow's Disease, [176](#)  
Barognosis, [176](#)  
Baroreceptor, [176](#)  
Baroreceptor Reflex, [176](#)  
Baroreceptors, [176](#)  
Baroreflex, [176](#)  
Barotitis Media, [176](#)  
Barotrauma, [176](#)  
Barr Body, [176](#)  
Barraquer-Simons Disease, [177](#)  
Barrel Chest, [177](#)  
Barrett Esophagus, [177](#)  
Barrett Esophagus Surveillance, [177](#)  
Barrett's Esophagus, [177](#)  
Barrett's Oesophagus, [177](#)  
Barrier Creams, [177](#)  
Barrier Nursing, [177](#)  
Bart's Hemoglobinopathy, [177](#)  
Barth Syndrome, [177](#)  
Barthel Index, [177](#)  
Bartholin Abscess, [177](#)  
Bartholin Cyst Or Abscess, [177](#)  
Bartholin Gland, [178](#)  
Bartholin Gland Cyst, [178](#)  
Bartholin Gland Hyperplasia, [178](#)  
Bartholin Gland Mucinous Cyst, [178](#)  
Bartholin's Cyst, [178](#)  
Bartholin's Glands, [178](#)  
Bartholinitis, [178](#)  
Bartonella, [178](#)  
Bartonella Elizabethae, [178](#)

---

---

Bartonella Henselae, 178  
Bartonella Infections, 178  
Bartonella Quintana, 178  
Bartonellaceae, 178  
Bartonellosis, 178  
Bartter Disease, 178  
Bartter Syndrome, 178  
Basal, 178  
Basal Body, 179  
Basal Cell, 179  
Basal Cell Carcinoma, 179  
Basal Cell Skin Cancer, 179  
Basal Ganglia, 179  
Basal Ganglia Dysfunction, 179  
Basal Joint Arthritis, 179  
Basal Lamina, 179  
Basal Metabolic Rate, 179  
Basal Nuclei, 179  
Basal Tear, 179  
Basal Temperature, 179  
Base, 179  
Base Excision Repair, 179  
Base Pair, 179  
Base Pairing, 180  
Base Sequence, 180  
Basedow's Disease, 180  
Baseline, 180  
Baseline EKG, 180  
Basement Membrane, 180  
Basic First Aid, 180  
Basic Life Support, 180  
Basic Metabolic Panel, 180  
Basic Reproduction Number, 180  
Basidiomycota, 180  
Basilar, 180  
Basilar Artery, 180  
Basilar Artery Aneurysm, 180  
Basilar Artery Migraine, 181  
Basilar Invagination, 181  
Basilar Membrane, 181  
Basilar Skull Fracture, 181  
Basilic Vein, 181  
Basiliximab, 181  
Basis Pontis, 181  
Basket Cell, 181  
Basophil, 181  
Basophil Adenoma, 181  
Basophilia, 181  
Basophilic Focus, 181  
Basophilic Stippling, 181  
Basosquamous Carcinoma, 182  
Bassen-Kornzweig Syndrome, 182  
Bateman Purpura, 182  
Bath Salts Abuse And Addiction, 182  
Bather's Itch, 182  
Bathing A Patient In Bed, 182  
Bathing An Infant, 182  
Batrachotoxin, 182  
Batroxobin, 182  
Batson Plexus, 182  
Batson Venous Plexus, 182  
Batten Disease, 182  
Battered Child Syndrome, 183  
Battey Bacillus, 183  
Battle Sign, 183  
Bauhin Valve, 183  
Bauxite-Workers' Lung, 183  
Bax Protein, 183  
Bayes Theorem, 183  
Bazin Disease, 183  
Bazin's Disease, 183  
BCG Vaccine, 2114  
Bcgosis, 183  
Beach Chair Position, 183  
Beacopp Regimen, 183  
Beals Syndrome, 183  
Beam Hardening Artifact, 184  
Beat Number, 184  
Beats, Cardiac, 895  
Beau Lines, 184  
Beau's Lines, 184  
Becaplermin, 184  
Bechterew Disease, 184  
Beck Triad, 184  
Becker Muscular Dystrophy, 184  
Becker Myotonia, 184  
Becker Nevus, 184  
Becker's Muscular Dystrophy, 184  
Becker'S Nevus, 184  
Beckwith-Wiedemann Spectrum, 184  
Beckwith-Wiedemann Syndrome, 185  
Beclomethasone, 185  
Becquerel, 185  
Bed Rest, 185  
Bedbug, 185  
Bedbug Bites, 185  
Bedbugs, 185  
Bednar Aphthae, 185  
Bedpan, 185  
Bedsore, 185  
Bedsores, 185  
Bedwetting, 185  
Bee And Wasp Sting, 185  
Bee Sting, 186  
Bee Stings, 186  
Beef or Pork Tapeworm Infection, 186  
Beer-Drinker's Heart, 186  
Beers Criteria, 186

---

---

Beeswax Poisoning, 186  
 Behavior, 186  
 Behavior Disorder, 186  
 Behavior Test, 186  
 Behavior Therapy, 186  
 Behavioral Abnormality, 186  
 Behavioral Disorder, 186  
 Behavioral Genetics, 186  
 Behavioral Medicine, 2114  
 Behavioral Rehabilitation, 186  
 Behcet Disease, 187  
 Behcet Syndrome, 187  
 Behcet's Syndrome, 187  
 Behr Syndrome, 187  
 Behçet Disease, 187  
 Behçet's Syndrome, 187  
 Beijerinck, Martinus W., 1202  
 Bejel, 187  
 Bekam, 187  
 Belching, 187  
 Belinostat, 187  
 Bell Palsy, 187  
 Bell's Palsy, 187  
 Bell's Phenomenon, 187  
 Bell-Clapper Deformity, 188  
 Belladonna, 188  
 Bells Palsy, 188  
 Belly, 188  
 Below-Knee Amputation, 188  
 Bence Jones Protein, 188  
 Bence-Jones Protein, 188  
 Benchmarking, 188  
 Bendamustine, 188  
 Bendamustine Hydrochloride, 188  
 Bendiocarb, 188  
 Bendopnea, 188  
 Bendroflumethiazide, 188  
 Bends, 189  
 Benedikt Syndrome, 189  
 Benedikt's Syndrome, 189  
 Beneficence And Non-Maleficence, 189  
 Benfotiamine, 189  
 Benfuracarb, 189  
 Benign, 189  
 Benign Adrenal Tumors, 189  
 Benign Ameloblastoma, 189  
 Benign Brain Tumor, 189  
 Benign Chondroblastoma, 189  
 Benign CNS Cyst, 189  
 Benign Ear Cyst Or Tumor, 189  
 Benign Epithelioma, 190  
 Benign Esophageal Stricture, 190  
 Benign Essential Tremor, 679  
 Benign Fasciculation Syndrome, 190  
 Benign Hemangiopericytoma, 190  
 Benign Lesions of the Ovaries, 190  
 Benign Liver Tumors, 190  
 Benign Melanocytic Lesion, 190  
 Benign Melanocytic Nevi Pathology, 190  
 Benign Meningioma, 190  
 Benign Mesenchymoma, 190  
 Benign Migratory Glossitis, 190  
 Benign Mixed Tumor, 190  
 Benign Mullerian Papilloma, 190  
 Benign Myoepithelioma, 190  
 Benign Neoplasm, 190  
 Benign Orgasmic Headache, 191  
 Benign Paraganglioma, 191  
 Benign Paroxysmal Positional Vertigo, 191  
 Benign Peripheral Nerve Tumor, 191  
 Benign Positional Vertigo, 191  
 Benign Prostatic Hyperplasia, 191  
 Benign Prostatic Obstruction, 191  
 Benign Rolandic Epilepsy, 191  
 Benign Skin Lesions, 191  
 Benign Tumor, 191  
 Benign Urethral Lesions, 191  
 Benign Vulvar Lesions, 192  
 Bennett Fracture, 192  
 Benserazide, 192  
 Bent Penis, 192  
 Bent Spine Syndrome, 192  
 Bentall Procedure, 192  
 Benzaldehyde, 192  
 Benzazepines, 192  
 Benzene, 192  
 Benzene Poisoning, 192  
 Benzhexol, 192  
 Benzo[a]pyrene, 192  
 Benzoates, 193  
 Benzocaine, 193  
 Benzodiazepine, 193  
 Benzodiazepine Toxicity, 193  
 Benzodiazepine Use Disorder, 193  
 Benzodiazepine Withdrawal, 193  
 Benzodiazepines, 2114  
 Benzopyran, 193  
 Benzopyrene, 193  
 Benzopyrenes, 193  
 Benzoyl Peroxide, 193  
 Benzphetamine, 193  
 Benztropine, 193  
 Benzydamine, 193  
 Benzyl Benzoate, 194  
 Benzylamines, 194  
 Benzylpenicillin, 194  
 BEP Regimen, 194  
 Berberine, 194

---

---

Bereavement, [194](#)  
 Bereavement Reaction, [194](#)  
 Berger Disease, [194](#)  
 Berger's Disease, [194](#)  
 Beriberi, [194](#)  
 Berloque Dermatitis, [194](#)  
 Bernard-Soulier Disease, [194](#)  
 Bernard-Soulier Syndrome, [194](#)  
 Bernard-Soulier Syndrome, [195](#)  
 Berry Aneurysm, [195](#)  
 Berylliosis, [195](#)  
 Beryllium, [195](#)  
 Best Disease, [195](#)  
 Beta 2-Microglobulin, [195](#)  
 Beta Agonists, [2115](#)  
 Beta Amyloid, [195](#)  
 Beta Blocker, [2115](#)  
 Beta Blockers, [2115](#)  
 Beta Carotene, [195](#)  
 Beta Cell, [195](#)  
 Beta Globulin, [195](#)  
 Beta Glucans, [195](#)  
 Beta Interferons, [195](#)  
 Beta Particle, [196](#)  
 Beta Radiation, [196](#)  
 Beta Thalassemia, [196](#)  
 Beta Tocopherol, [196](#)  
 Beta-Adrenergic Blocker, [2114](#)  
 Beta-Adrenoceptor-Blocking Drugs, [196](#)  
 Beta-Agonist, [196](#)  
 Beta-Alanine, [196](#)  
 Beta-Blocker Overdose, [2115](#)  
 Beta-Blocker Toxicity, [196](#)  
 Beta-Blockers, [196](#)  
 Beta-Blockers Overdose, [196](#)  
 Beta-Carotene, [196](#)  
 Beta-Carotene Blood Test, [196](#)  
 Beta-Chemokines, [196](#)  
 Beta-Endorphin, [197](#)  
 Beta-Glucans, [197](#)  
 Beta-Haemolytic Streptococcus, [197](#)  
 Beta-Human Chorionic Gonadotrophin, [197](#)  
 Beta-Lactam Antibiotics, [197](#)  
 Beta-Lactamase, [197](#)  
 Beta-Lactamase Inhibitor, [197](#)  
 Beta-Lactamase Inhibitor Combinations, [2115](#)  
 Beta-Mannosidosis, [197](#)  
 Beta-Oxidation, [197](#)  
 Beta-Sheet, [197](#)  
 Beta-Thalassemia, [197](#)  
 Beta-Thalassemia Intermedia, [197](#)  
 Beta-Thalassemia Major, [197](#)  
 Beta-Thalassemia Minor, [197](#)  
 Beta-Thalassemia Trait, [198](#)  
 Betacoronavirus, [198](#)  
 Betaherpesvirinae, [198](#)  
 Betaine, [198](#)  
 Betamethasone, [198](#)  
 Betamethasone Valerate, [198](#)  
 Betaxolol, [198](#)  
 Bethanechol, [198](#)  
 Bethesda System For Reporting Thyroid Cytopathology, [198](#)  
 Bethlem Myopathy, [198](#)  
 Bevacizumab, [198](#)  
 Bexarotene, [198](#)  
 Bezafibrate, [199](#)  
 Bezoar, [199](#)  
 Bezold-Jarisch Reflex, [199](#)  
 BH4 Deficiency, [199](#)  
 Biacetyl, [199](#)  
 Bias, [199](#)  
 Bibliotherapy, [199](#)  
 Bicalutamide, [199](#)  
 Bicarbonate, [199](#)  
 Bicarbonate Buffer System, [199](#)  
 Biceps, [199](#)  
 Biceps Brachii Muscle, [199](#)  
 Biceps Reflex, [199](#)  
 Bicipital, [200](#)  
 Bicipital Groove, [200](#)  
 Bicipital Tendinitis, [200](#)  
 Bicipital Tuberosity, [200](#)  
 Bickerstaff Brainstem Encephalitis, [200](#)  
 Bicolor, [200](#)  
 Biconcave, [200](#)  
 Biconvex, [200](#)  
 Bicornuate, [200](#)  
 Bicornuate Uterus, [200](#)  
 Bicuspid, [200](#)  
 Bicuspid Aortic Valve, [200](#)  
 Bicuspid Valve, [200](#)  
 Bid Protein, [200](#)  
 Bielschowsky Head Tilt Test, [200](#)  
 Bier Spots, [201](#)  
 Biermer's Anemia, [201](#)  
 Bietti's Crystalline Dystrophy, [201](#)  
 Bifascicular Block, [201](#)  
 Bifocal, [201](#)  
 Bifurcation, [201](#)  
 Big Toe Sign, [201](#)  
 Bigeminy, [201](#)  
 Biglycan, [201](#)  
 Biguanide, [201](#)  
 Biguanides, [201](#)  
 Bilateral, [201](#)  
 Bilateral Acoustic Neurofibromatosis, [201](#)  
 Bilateral Acoustic Neuroma, [202](#)

---



---

Bilateral Ankle Swelling, [202](#)  
Bilateral Gynecomastia, [202](#)  
Bilateral Hydronephrosis, [202](#)  
Bilateral Microphthalmos, [202](#)  
Bilateral Oophorectomy, [202](#)  
Bilateral Peritonsillar Abscesses, [202](#)  
Bilateral Pneumonia, [202](#)  
Bilateral Polycystic Ovarian Syndrome, [202](#)  
Bilateral Tonic-Clonic Seizure, [202](#)  
Bile, [202](#)  
Bile Acid Sequestrants For Cholesterol, [202](#)  
Bile Acids, [202](#)  
Bile Culture, [202](#)  
Bile Duct, [203](#)  
Bile Duct Cancer, [203](#)  
Bile Duct Carcinoma, [203](#)  
Bile Duct Obstruction, [203](#)  
Bile Duct Stricture, [203](#)  
Bile Ducts, [203](#)  
Bile Pigment, [203](#)  
Bile Reflux, [203](#)  
Bile Reflux Gastritis, [203](#)  
Bile Salt Emulsification, [203](#)  
Bile Salt Malabsorption, [203](#)  
Bile Salts, [203](#)  
Bilevel Positive Airway Pressure, [203](#)  
Bilharzia, [203](#)  
Bili Lights, [204](#)  
Biliary, [204](#)  
Biliary Atresia, [204](#)  
Biliary Atresia, Extrahepatic, [204](#)  
Biliary Calculi, [204](#)  
Biliary Cirrhosis, [204](#)  
Biliary Cirrhosis, Primary, [1592](#)  
Biliary Colic, [204](#)  
Biliary Decompression, [204](#)  
Biliary Dyskinesia, [204](#)  
Biliary Fever, [204](#)  
Biliary Fistula, [204](#)  
Biliary Sand, [204](#)  
Biliary Scintigraphy, [204](#)  
Biliary Stasis, [205](#)  
Biliary Stent, [205](#)  
Biliary System, [205](#)  
Biliary Tract, [205](#)  
Biliary Tract Biopsy, [205](#)  
Biliary Tract Injuries, [205](#)  
Biliary Tree, [205](#)  
Bilious, [205](#)  
Bilious Vomiting, [205](#)  
Bilirubin, [205](#)  
Bilirubin Blood Test, [205](#)  
Bilirubin Encephalopathy, [205](#)  
Bilirubin Metabolism, [205](#)  
Bilirubinuria, [205](#)  
Biliverdin, [206](#)  
Billroth I, [206](#)  
Billroth I Operation, [206](#)  
Billroth II, [206](#)  
Billroth II Operation, [206](#)  
Billroth's Operation, [206](#)  
Bilophila, [206](#)  
Bilophila Wadsworthia, [206](#)  
Bimanual, [206](#)  
Bimatoprost, [206](#)  
Bimedial, [206](#)  
Bimellar, [206](#)  
Binary Fission, [206](#)  
Binder Excipient, [206](#)  
Binding Protein, [207](#)  
Binding Site, [207](#)  
Binet-Simon Scale, [207](#)  
Binge Drinking, [207](#)  
Binge Eating Disorder, [207](#)  
Binge-Eating Disorder, [207](#)  
Binge-Purge Disorder, [207](#)  
Binocular Vision, [207](#)  
Binocular Visual Loss, [207](#)  
Binswanger Disease, [207](#)  
Binswanger's Dementia, [207](#)  
Bio-Informatics, [207](#)  
Bioabsorbable Vascular Stent, [207](#)  
Bioaccumulation, [208](#)  
Bioactive, [208](#)  
Bioassay, [208](#)  
Bioavailability, [208](#)  
Bioburden, [208](#)  
Biochanin A, [208](#)  
Biochemical, [208](#)  
Biochemical Assessment, [208](#)  
Biochemical Reaction, [208](#)  
Biochemical Recurrence, [208](#)  
Biochemistry, [208](#)  
Biocompatibility, [208](#)  
Biodegradable Implant, [208](#)  
Biodistribution, [208](#)  
Bioengineering, [209](#)  
Bioequivalence, [209](#)  
Bioethics, [209](#)  
Biofeedback, [209](#)  
Biofilm, [209](#)  
Biofilm-Associated Infections, [209](#)  
Biofluid, [209](#)  
Biologic Therapy, [209](#)  
Biologic Therapy In Dermatology, [209](#)  
Biologic Variation, [209](#)  
Biological, [209](#)  
Biological Availability, [209](#)

---

---

Biological Clock, [209](#)  
Biological Half-Life, [209](#)  
Biological Model, [210](#)  
Biological Process, [210](#)  
Biological Relative, [210](#)  
Biological Therapy, [210](#)  
Biological Transport, [210](#)  
Biological Valve, [210](#)  
Biological Variability, [210](#)  
Biological Warfare, [210](#)  
Bioluminescence, [210](#)  
Biomarker, [210](#)  
Biomarker Testing, [210](#)  
Biomaterial, [210](#)  
Biomechanical Compliance, [210](#)  
Biomechanics, [210](#)  
Biomedical Cement, [211](#)  
Biomedical Device, [211](#)  
Biomedical Engineering, [211](#)  
Biomedical Tape, [211](#)  
Biomedicine, [211](#)  
Biometry, [211](#)  
Biomicroscopy, [211](#)  
Biomolecule, [211](#)  
Biomonitoring, [211](#)  
Biomphalaria, [211](#)  
Bionics, [211](#)  
Biophotonics, [211](#)  
Biophysical Profile, [211](#)  
Biopsy, [211](#)  
Biopsy Lung, [212](#)  
Biopsy Needle, [212](#)  
Biopsy Site, [212](#)  
Biopsy Specimen, [212](#)  
Biosafety, [212](#)  
Biosafety Level, [212](#)  
Biosecurity, [212](#)  
Biosensor, [212](#)  
Biosimilar, [212](#)  
Biospecimen, [212](#)  
Biostatistics, [212](#)  
Biosynthesis, [212](#)  
Biot Breathing, [212](#)  
Biot's Breathing, [212](#)  
Biotechnology, [213](#)  
Biotelemetry, [213](#)  
Bioterrorism, [213](#)  
Bioterrorism Agents, [213](#)  
Biotherapy, [213](#)  
Biotin, [213](#)  
Biotin Vitamin B7, [213](#)  
Biotinidase Deficiency, [213](#)  
Biotinylation, [213](#)  
Biotransformation, [213](#)  
Biotrauma, [213](#)  
BiPAP, [213](#)  
BiPAP Machine, [213](#)  
Biparental, [214](#)  
Biparietal Diameter, [214](#)  
Bipartite Ossification, [214](#)  
Biphenyl Compounds, [214](#)  
Bipolar, [214](#)  
Bipolar Affective Disorder, [214](#)  
Bipolar Disorder, [214](#)  
Bipolar Disorder in Child and Teens, [214](#)  
Bipolar I Disorder, [214](#)  
Bipolar II Disorder, [214](#)  
Bipolar Neuron, [214](#)  
Bipolaris, [214](#)  
Birbeck Granules, [214](#)  
Bird Fancier's Lung, [214](#)  
Bird Flu, [215](#)  
Birt-Hogg-Dube Syndrome, [215](#)  
Birt-Hogg-Dube Syndrome (BHDS), [215](#)  
Birt-Hogg-Dubé Syndrome, [215](#)  
Birth, [215](#)  
Birth Asphyxia, [215](#)  
Birth Control, [215](#)  
Birth Control And Family Planning, [215](#)  
Birth Control Methods, [215](#)  
Birth Control Pill Overdose, [215](#)  
Birth Control Pills, [215](#)  
Birth Control, Extended-Release, [1157](#)  
Birth Defect, [215](#)  
Birth Defects, [216](#)  
Birth Defects Testing, [216](#)  
Birth Injury, [216](#)  
Birth Order, [216](#)  
Birth Pains, [216](#)  
Birth Rate, [216](#)  
Birth Spacing, [216](#)  
Birth Weight, [216](#)  
Birthmark, [216](#)  
Birthmarks, [216](#)  
Birthmarks And Other Skin Pigmentation Problems, [216](#)  
Bis In Die, [216](#)  
Bisacodyl, [216](#)  
Bisexual, [217](#)  
Bisexuality, [217](#)  
Bishop Score, [217](#)  
Bismuth, [217](#)  
Bisoprolol, [217](#)  
Bispecific Antibody Therapy, [217](#)  
Bisphenol A, [217](#)  
Bisphosphonate, [217](#)  
Bisphosphonate-Related Osteonecrosis of the Jaw, [217](#)  
Bisphosphonates, [217](#)  
Bite, Chigger, [378](#)

---

---

Bite, Flea, [751](#)  
Bite, Snake, [1809](#)  
Bitemporal Hemianopia, [217](#)  
Bitemporal Hemianopsia, [217](#)  
Bitewing, [218](#)  
Bitewing Radiograph, [218](#)  
Bitot's Spots, [218](#)  
Bituminosis, [218](#)  
Biuret Test, [218](#)  
Bivalve, [218](#)  
Bizarre Parosteal Osteochondromatous Proliferation, [218](#)  
BK Virus, [218](#)  
BK Virus Nephropathy, [218](#)  
Black Death, [218](#)  
Black Eye, [218](#)  
Black Fly, [218](#)  
Black Hairy Tongue, [218](#)  
Black Lung, [219](#)  
Black Lung Disease, [219](#)  
Black Nightshade Poisoning, [219](#)  
Black Or Tarry Stools, [219](#)  
Black Stain, [219](#)  
Black Triangle, [219](#)  
Black Widow Spider, [219](#)  
Black Widow Spider Bite, [219](#)  
Blackflesh, [219](#)  
Blackhead, [219](#)  
Blackheads, [219](#)  
Blackout, [219](#)  
Blackwater Fever, [219](#)  
Bladder, [219](#)  
Bladder Augmentation, [220](#)  
Bladder Biopsy, [220](#)  
Bladder Calculi, [220](#)  
Bladder Cancer, [220](#)  
Bladder Carcinoma, [220](#)  
Bladder Control, Loss of, [2002](#)  
Bladder Diary, [220](#)  
Bladder Diseases, [220](#)  
Bladder Disorders, [220](#)  
Bladder Diverticula, [220](#)  
Bladder Diverticulum, [220](#)  
Bladder Exstrophy, [220](#)  
Bladder Exstrophy Repair, [220](#)  
Bladder Fistula, [220](#)  
Bladder Health And Disorders, [221](#)  
Bladder Infection, [221](#)  
Bladder Infections in Children, [221](#)  
Bladder Infections in Teens and Adults, [221](#)  
Bladder Inflammation, [221](#)  
Bladder Neck, [221](#)  
Bladder Neck Contracture, [221](#)  
Bladder Neck Obstruction, [221](#)  
Bladder Neoplasm, [221](#)  
Bladder Outlet Obstruction, [221](#)  
Bladder Pain, [221](#)  
Bladder Pain Syndrome, [221](#)  
Bladder Perforation, [221](#)  
Bladder Problems, [221](#)  
Bladder Prolapse, [222](#)  
Bladder Retraining, [222](#)  
Bladder Spasm, [222](#)  
Bladder Stones, [222](#)  
Bladder Trauma, [222](#)  
Bladder, Urinary, [2001](#)  
Blalock-Taussig Operation, [222](#)  
Blalock-Taussig Shunt, [222](#)  
Bland Diet, [222](#)  
Blaschko Lines, [222](#)  
BLAST, [222](#)  
Blast Cell, [222](#)  
Blast Crisis, [222](#)  
Blast Injuries, [222](#)  
Blast Injury, [222](#)  
Blast Phase, [223](#)  
Blastema, [223](#)  
Blastic Plasmacytoid Dendritic-Cell Neoplasm, [223](#)  
Blastocyst, [223](#)  
Blastocyst Transfer, [223](#)  
Blastocystis, [223](#)  
Blastocystis Hominis, [223](#)  
Blastocystosis, [223](#)  
Blastoma, [223](#)  
Blastomyces Dermatitis, [223](#)  
Blastomycosis, [223](#)  
Blasts, [223](#)  
Blautia Producta, [223](#)  
Bleb, [223](#)  
Bleeder's Disease, [224](#)  
Bleeding, [224](#)  
Bleeding Disorder, [224](#)  
Bleeding Disorders, [224](#)  
Bleeding Esophageal Varices, [224](#)  
Bleeding Gums, [224](#)  
Bleeding Into A Thyroid Nodule, [224](#)  
Bleeding Tendency, [224](#)  
Bleeding Time, [224](#)  
Bleeding Varices, [224](#)  
Blennorrhoea, [224](#)  
Bleomycin, [224](#)  
Bleomycin Sulfate, [224](#)  
Blepharitis, [224](#)  
Blepharoconjunctivitis, [225](#)  
Blepharophimosis, [225](#)  
Blepharoplasty, [225](#)  
Blepharoptosis, [225](#)  
Blepharospasm, [225](#)  
Blind, [225](#)

---

---

Blind Loop Syndrome, [225](#)  
Blind Spot, [225](#)  
Blinded, [225](#)  
Blinding, [225](#)  
Blindness, [225](#)  
Blindness And Vision Loss, [225](#)  
Blinking, [225](#)  
Blister, [226](#)  
Blisters, [226](#)  
Bloating, [226](#)  
Block, [226](#)  
Blockage Of Upper Airway, [226](#)  
Blocked Tear Duct, [226](#)  
Blocked Tear Ducts, [226](#)  
Blocked Ureter, [226](#)  
Blocking Agent, [226](#)  
Blocking Agents, [226](#)  
Blocking Antibody, [226](#)  
Blood, [226](#)  
Blood Alcohol Concentration, [226](#)  
Blood Alcohol Level, [226](#)  
Blood Bank, [227](#)  
Blood Banking, [227](#)  
Blood Biomarker, [227](#)  
Blood Blister, [227](#)  
Blood Brain Barrier, [227](#)  
Blood Cancer, [227](#)  
Blood Cell, [227](#)  
Blood Cholesterol, [227](#)  
Blood Cholinesterase Test, [227](#)  
Blood Chromium Test, [227](#)  
Blood Circulation, [227](#)  
Blood Cleaner, [227](#)  
Blood Clot, [227](#)  
Blood Clot in Lung, [228](#)  
Blood Clots, [228](#)  
Blood Coagulation, [228](#)  
Blood Coagulation Disorder, [228](#)  
Blood Count, [228](#)  
Blood Culture, [228](#)  
Blood Culture Contamination, [228](#)  
Blood Differential Test, [228](#)  
Blood Donation, [228](#)  
Blood Donation Before Surgery, [228](#)  
Blood Donor, [228](#)  
Blood Draw, [228](#)  
Blood Dyscrasia, [228](#)  
Blood Film, [229](#)  
Blood Flow, [229](#)  
Blood Flow Velocity, [229](#)  
Blood Gas Analysis, [229](#)  
Blood Gases, [229](#)  
Blood Glucose, [229](#)  
Blood Glucose Monitoring, [229](#)  
Blood Glucose Monitors, [229](#)  
Blood Group, [229](#)  
Blood Group Incompatibility, [229](#)  
Blood Group, ABO, [9](#)  
Blood Groups, [229](#)  
Blood In Semen, [230](#)  
Blood in Stool, [230](#)  
Blood In The Eye, [230](#)  
Blood in Urine, [230](#)  
Blood Lancet, [230](#)  
Blood Lead Level Test, [230](#)  
Blood Loss, [230](#)  
Blood Low Red Cell Count, [230](#)  
Blood PH, [230](#)  
Blood Plasma, [230](#)  
Blood Preservation, [230](#)  
Blood Pressure, [230](#)  
Blood Pressure Cuff, [230](#)  
Blood Pressure Measurement, [230](#)  
Blood Pressure Monitors For Home, [231](#)  
Blood Pressure Regulation, [231](#)  
Blood Product, [231](#)  
Blood Products, [231](#)  
Blood Protein, [231](#)  
Blood Results, [231](#)  
Blood Sedimentation, [231](#)  
Blood Smear, [231](#)  
Blood Spatter Analysis, [231](#)  
Blood Substitutes, [231](#)  
Blood Sugar, [231](#)  
Blood Sugar Test, [231](#)  
Blood Test, [231](#)  
Blood Thinner, [232](#)  
Blood Transfusion, [232](#)  
Blood Typing, [232](#)  
Blood Urea Nitrogen, [232](#)  
Blood Urea Nitrogen Test, [232](#)  
Blood Vessel, [232](#)  
Blood Vessels, [232](#)  
Blood Viscosity, [232](#)  
Blood Volume, [232](#)  
Blood-Air Barrier, [232](#)  
Blood-Borne Disease, [232](#)  
Blood-Borne Pathogens, [233](#)  
Blood-Brain Barrier, [233](#)  
Blood-Retinal Barrier, [233](#)  
Blood-Testis Barrier, [233](#)  
Bloodborne Pathogens, [233](#)  
Bloom Syndrome, [233](#)  
Blount Disease, [233](#)  
Blue Baby, [233](#)  
Blue Bottle, [233](#)  
Blue Discoloration Of The Skin, [233](#)  
Blue Nevi, [233](#)

---

---

Blue Nevus, [233](#)  
Blue Nightshade Poisoning, [234](#)  
Blue NæVus, [234](#)  
Blue Rubber Bleb Nevus Syndrome, [234](#)  
Blue Sclera, [234](#)  
Blue Sclerae, [234](#)  
Blue Toe Syndrome, [234](#)  
Blue-Green Algae, [234](#)  
Blumberg Sign, [234](#)  
Blunt Abdominal Trauma, [234](#)  
Blunt Force Trauma, [234](#)  
Blunted Affect, [234](#)  
Blurred Vision, [234](#)  
Blush, [234](#)  
Blushing, [235](#)  
Board-Like Rigidity, [235](#)  
Boas' Sign, [235](#)  
Bobbing, Ocular, [1390](#)  
Bobble-Head Doll Syndrome, [235](#)  
Bocavirus, [235](#)  
Bocavirus Infection, [235](#)  
Bochdalek Hernia, [235](#)  
Bock's Disease, [235](#)  
BODE Index, [235](#)  
Bodily Secretion, [235](#)  
Body, [235](#)  
Body Aches, [235](#)  
Body Cavity, [236](#)  
Body Cell Mass, [236](#)  
Body Composition, [236](#)  
Body Dysmorphia, [236](#)  
Body Dysmorphic Disorder, [236](#)  
Body Fat, [236](#)  
Body Fluid, [236](#)  
Body Fluid Exposures, [236](#)  
Body Fluids, [236](#)  
Body Habitus, [236](#)  
Body Height, [236](#)  
Body Identification, [236](#)  
Body Image, [236](#)  
Body Lice, [237](#)  
Body Mass Index, [237](#)  
Body Odor, [237](#)  
Body Odor and Sweating, [237](#)  
Body Odour, [237](#)  
Body Of The Epididymis, [237](#)  
Body Of The Sternum, [237](#)  
Body Piercing, [237](#)  
Body Plethysmography, [237](#)  
Body Position, [237](#)  
Body Region, [237](#)  
Body Ringworm, [237](#)  
Body Space, [237](#)  
Body Substance, [237](#)  
Body Surface, [238](#)  
Body Surface Area, [238](#)  
Body Temperature, [238](#)  
Body Temperature Norms, [238](#)  
Body Wall, [238](#)  
Body Weight, [238](#)  
Boeck's Sarcoid, [238](#)  
Boerhaave Syndrome, [238](#)  
Boerhaave's Syndrome, [238](#)  
Bogros' Space, [238](#)  
Bohn Nodules, [238](#)  
Bohr Effect, [238](#)  
Bohring Syndrome, [238](#)  
Boil, [238](#)  
Boils, [239](#)  
Boils and Carbuncles, [239](#)  
Bolt Device, [239](#)  
Bolus, [239](#)  
Bolus Gastrostomy Feeding, [239](#)  
Bolus Infusion, [239](#)  
Bombay Blood Group, [239](#)  
Bombesin Receptor, [239](#)  
Bone, [239](#)  
Bone And Mineral Disorders, [239](#)  
Bone Callus, [239](#)  
Bone Cancer, [239](#)  
Bone Cement, [239](#)  
Bone Conduction, [240](#)  
Bone Cyst, [240](#)  
Bone Densitometry, [240](#)  
Bone Density, [240](#)  
Bone Density Screening, [240](#)  
Bone Development, [240](#)  
Bone Disease, [240](#)  
Bone Fracture, [240](#)  
Bone Fractures, [240](#)  
Bone Graft, [240](#)  
Bone Health, [240](#)  
Bone Infarction, [240](#)  
Bone Infection, [240](#)  
Bone Lesion Biopsy, [241](#)  
Bone Markers in Osteoporosis, [241](#)  
Bone Marrow, [241](#)  
Bone Marrow Aspiration, [241](#)  
Bone Marrow Aspiration And Biopsy, [241](#)  
Bone Marrow Biopsy, [241](#)  
Bone Marrow Culture, [241](#)  
Bone Marrow Depression, [241](#)  
Bone Marrow Disorder, [241](#)  
Bone Marrow Donation, [241](#)  
Bone Marrow Failure, [241](#)  
Bone Marrow Hyperplasia, [241](#)  
Bone Marrow Necrosis, [241](#)  
Bone Marrow Transplantation, [242](#)

---

---

Bone Mass, [242](#)  
Bone Matrix, [242](#)  
Bone Metastases, [242](#)  
Bone Metastasis, [242](#)  
Bone Mineral Density, [242](#)  
Bone Mineral Density Test, [242](#)  
Bone Morphogenetic Protein, [242](#)  
Bone Necrosis, [242](#)  
Bone Neoplasm, [242](#)  
Bone Pain, [242](#)  
Bone Pain Or Tenderness, [242](#)  
Bone Plate, [242](#)  
Bone Regeneration, [243](#)  
Bone Remodeling, [243](#)  
Bone Resorption, [243](#)  
Bone Sarcoma, [243](#)  
Bone Saw, [243](#)  
Bone Scan, [243](#)  
Bone Screws, [243](#)  
Bone Spurs, [243](#)  
Bone Transplantation, [243](#)  
Bone Tumor, [243](#)  
Bone Tumor-Epidermoid Cyst-Polyposis, [243](#)  
Bone Turnover Markers, [243](#)  
Bone Type, [243](#)  
Bone X-Ray, [244](#)  
Bone-Anchored Hearing Implant, [244](#)  
Bone-Marrow Transplantation, [244](#)  
Bones And Joints, [244](#)  
Bones of the Skeleton, [244](#)  
Bongkrek Poisoning, [244](#)  
Bonnieville-Ullrich Syndrome, [244](#)  
Bony Exostosis, [244](#)  
Bony Labyrinth, [244](#)  
Borborygmi, [244](#)  
Borborygmus, [244](#)  
Border Definition, [244](#)  
Borderline, [244](#)  
Borderline Ovarian Cancer, [244](#)  
Borderline Personality Disorder, [245](#)  
Bordetella, [245](#)  
Bordetella Avium, [245](#)  
Bordetella Bronchiseptica, [245](#)  
Bordetella Hinzei, [245](#)  
Bordetella Parapertussis, [245](#)  
Bordetella Pertussis, [245](#)  
Bordetella Petrii, [245](#)  
Boric Acid, [245](#)  
Boric Acid Poisoning, [245](#)  
Borna Disease, [245](#)  
Bornholm Disease, [245](#)  
Boron, [245](#)  
Borrelia, [246](#)  
Borrelia Burgdorferi, [246](#)  
Borrelia Infection, [246](#)  
Borrelia Miyamotoi, [246](#)  
Borreliosis, [246](#)  
Bortezomib, [246](#)  
Bosentan, [246](#)  
Bosniak Classification, [246](#)  
Boston Criteria, [246](#)  
Bot Fly, [246](#)  
Botfly, [246](#)  
Bothriocephalus, [246](#)  
Botryomycosis, [246](#)  
Bottle Feeding, [247](#)  
Botulinum Toxin, [247](#)  
Botulism, [247](#)  
Bouchard Nodes, [247](#)  
Bouchard's Node, [247](#)  
Bouchard's Nodes, [247](#)  
Bougie, [247](#)  
Bougienage, [247](#)  
Bouin's Solution, [247](#)  
Bounding Pulse, [247](#)  
Bourbon Virus, [247](#)  
Boutonneuse Fever, [247](#)  
Boutonniere Deformity, [248](#)  
Bouveret Syndrome, [248](#)  
Bovine, [248](#)  
Bovine Anaplasmosis, [248](#)  
Bovine Papillomavirus, [248](#)  
Bovine Spongiform Encephalopathy, [248](#)  
Bow Legs, [248](#)  
Bowditch Effect, [248](#)  
Bowel, [248](#)  
Bowel Burn, [248](#)  
Bowel Cancer, [248](#)  
Bowel Death, [248](#)  
Bowel Disorders, [248](#)  
Bowel Habit Changes and Bloating, [249](#)  
Bowel Habits, [249](#)  
Bowel Incontinence, [249](#)  
Bowel Infarction, [249](#)  
Bowel Ischemia, [249](#)  
Bowel Movement, [249](#)  
Bowel Obstruction, [249](#)  
Bowel Preparation, [249](#)  
Bowel Program, [249](#)  
Bowel Retraining, [249](#)  
Bowel Sounds, [249](#)  
Bowel Transit Time, [249](#)  
Bowel Urgency, [249](#)  
Bowen's Disease, [250](#)  
Bowenoid Papulosis, [250](#)  
Bowen's Disease, [250](#)  
Bowlegs, [250](#)  
Bowler's Thumb, [250](#)

---



---

Bowman Capsule, [250](#)  
Bowman Membrane, [250](#)  
Bowman's Capsule, [250](#)  
Bowman's Gland, [250](#)  
Boxed Warning, [250](#)  
Boxer Fracture, [250](#)  
Boxer's Ear, [250](#)  
Boxer's Fracture, [250](#)  
BPH (Benign Prostatic Hyperplasia), [250](#)  
Braces and Retainers, [251](#)  
Brachial, [251](#)  
Brachial Artery, [251](#)  
Brachial Plexopathy, [251](#)  
Brachial Plexus, [251](#)  
Brachial Plexus Cords, [251](#)  
Brachial Plexus Injuries, [251](#)  
Brachial Plexus Injury, [251](#)  
Brachial Plexus Injury In Newborns, [251](#)  
Brachial Plexus Neuropathies, [251](#)  
Brachial Plexus Roots, [251](#)  
Brachial Plexus Trunks, [251](#)  
Brachial Vein, [251](#)  
Brachialgia, [251](#)  
Brachialis Muscle, [252](#)  
Brachiocephalic Artery, [252](#)  
Brachiocephalic Trunk, [252](#)  
Brachiocephalic Veins, [252](#)  
Brachioplasty, [252](#)  
Brachioradial Pruritus, [252](#)  
Brachioradialis Reflex, [252](#)  
Brachium, [252](#)  
Brachycephalic, [252](#)  
Brachycephaly, [252](#)  
Brachydactyly, [252](#)  
Brachytherapy, [252](#)  
Brachytherapy Seed, [252](#)  
Bradycardia, [253](#)  
Bradycardia (Slow Heart Rate), [253](#)  
Bradycardia Of Prematurity, [253](#)  
Bradycardia-Tachycardia Syndrome, [253](#)  
Bradykinesia, [253](#)  
Bradykinesia and Athetosis, [253](#)  
Bradykinetic, [253](#)  
Bradykinin, [253](#)  
Bradyphrenia, [253](#)  
Bradypnea, [253](#)  
Bradypnoea, [253](#)  
BRAF Gene, [253](#)  
Braille System, [254](#)  
Brain, [254](#)  
Brain Abscess, [254](#)  
Brain Aneurysm, [254](#)  
Brain Aneurysm Repair, [254](#)  
Brain Arteriovenous Malformation, [254](#)  
Brain AVM, [254](#)  
Brain Cancer, [254](#)  
Brain Concussion, [254](#)  
Brain Death, [254](#)  
Brain Disorder, [254](#)  
Brain Edema, [254](#)  
Brain Embolism, [254](#)  
Brain Fog, [254](#)  
Brain Fornix, [255](#)  
Brain Haemorrhage, [255](#)  
Brain Health, [255](#)  
Brain Hemorrhage, [255](#)  
Brain Herniation, [255](#)  
Brain Imaging, [255](#)  
Brain Injuries, [255](#)  
Brain Injury, [255](#)  
Brain Lesions, [255](#)  
Brain Malformations, [255](#)  
Brain Mapping, [255](#)  
Brain Metastases, [255](#)  
Brain Metastasis, [256](#)  
Brain Monitoring, [256](#)  
Brain Natriuretic Peptide Test, [256](#)  
Brain Neoplasm, [256](#)  
Brain Neoplasms, [256](#)  
Brain PET Scan, [256](#)  
Brain Sand, [256](#)  
Brain SPECT, [256](#)  
Brain Stem, [256](#)  
Brain Stem Glioma, [256](#)  
Brain Stimulation, [256](#)  
Brain Surgery, [256](#)  
Brain Tumor, [256](#)  
Brain Tumors in Adults, [257](#)  
Brain Tumour, [257](#)  
Brain Ventricles, [257](#)  
Brain Wave, [257](#)  
Brain Waves, [257](#)  
Brainstem, [257](#)  
Brainstem Auditory Evoked Potential, [257](#)  
Brainstem Auditory Evoked Response, [257](#)  
Brainstem Death, [257](#)  
Brainstem Gliomas, [257](#)  
Brainstem Tumor, [257](#)  
Branchial, [257](#)  
Branchial Arch, [258](#)  
Branchial Cleft, [258](#)  
Branchial Cleft Anomalies, [258](#)  
Branchial Cleft Anomaly, [258](#)  
Branchial Cleft Cyst, [258](#)  
Branchial Cyst, [258](#)  
Branchio-Oto-Renal Syndrome, [258](#)  
Branchioma, [258](#)

---

---

Brass-Founder's Ague, [258](#)  
Brassicasterol, [258](#)  
Brassicidic Acid, [258](#)  
Braxton Hicks Contractions, [258](#)  
Brazilian Pemphigus Foliaceus, [258](#)  
Brazilian Trypanosomiasis, [259](#)  
BRCA, [259](#)  
BRCA1 And BRCA2, [259](#)  
BRCA1 And BRCA2 Gene Testing, [259](#)  
BRCA1 Gene, [259](#)  
BRCA2 Gene, [259](#)  
Breakbone Fever, [259](#)  
Breakthrough Bleeding, [259](#)  
Breakthrough Infection, [259](#)  
Breakthrough Pain, [259](#)  
Breast, [259](#)  
Breast Abscess, [259](#)  
Breast Anatomy, [259](#)  
Breast And Thoracic Cancer Surgery, [260](#)  
Breast Augmentation, [260](#)  
Breast Augmentation Surgery, [260](#)  
Breast Bone, [260](#)  
Breast Calcifications, [260](#)  
Breast Cancer, [260](#)  
Breast Cancer And Lymphedema, [260](#)  
Breast Cancer and Pregnancy, [260](#)  
Breast Cancer and Your Genes, [260](#)  
Breast Cancer Genetics, [260](#)  
Breast Cancer Grade, [260](#)  
Breast Cancer in Men, [260](#)  
Breast Cancer Prevention, [260](#)  
Breast Cancer Recurrence, [261](#)  
Breast Cancer Screening, [261](#)  
Breast Cancer Staging, [261](#)  
Breast Cancer, Inflammatory, [1018](#)  
Breast Cancer, Male, [1192](#)  
Breast Cancer, Metastatic or Recurrent, [1677](#)  
Breast Carcinoma, [261](#)  
Breast Cyst, [261](#)  
Breast Cysts, [261](#)  
Breast Cytology, [261](#)  
Breast Density, [261](#)  
Breast Development, [261](#)  
Breast Development in Boys, [261](#)  
Breast Duct, [261](#)  
Breast Engorgement, [261](#)  
Breast Enlargement In Males, [261](#)  
Breast Examination, [261](#)  
Breast Feed, [262](#)  
Breast Finding, [262](#)  
Breast Health, [262](#)  
Breast Imaging-Reporting And Data System, [262](#)  
Breast Infection, [262](#)  
Breast Infection (Mastitis), [262](#)  
Breast Lift, [262](#)  
Breast Lobe, [262](#)  
Breast Lobule, [262](#)  
Breast Lump, [262](#)  
Breast Lump Removal, [262](#)  
Breast Lumps, [262](#)  
Breast Lumps (Benign), [262](#)  
Breast Milk, [262](#)  
Breast Milk Expression and Storage, [263](#)  
Breast Milk Jaundice, [263](#)  
Breast MRI, [263](#)  
Breast MRI Scan, [263](#)  
Breast Neoplasm, [263](#)  
Breast Pain, [263](#)  
Breast Pain (Mastalgia), [263](#)  
Breast Quadrant, [263](#)  
Breast Rash, [263](#)  
Breast Reconstruction, [263](#)  
Breast Reduction, [263](#)  
Breast Screening, [264](#)  
Breast Self-Examination, [264](#)  
Breast Skin And Nipple Changes, [264](#)  
Breast Surgeon, [264](#)  
Breast Surgery, [264](#)  
Breast Ultrasound, [264](#)  
Breast-Conserving Treatment, [264](#)  
Breastfeeding, [264](#)  
Breastfeeding With Rheumatoid Arthritis, [264](#)  
Breastfeeding-Associated Nipple and Skin Changes, [264](#)  
Breath, [264](#)  
Breath Alcohol Test, [265](#)  
Breath Analysis, [265](#)  
Breath Focus, [265](#)  
Breath Odor, [265](#)  
Breath Sounds, [265](#)  
Breath Stacking, [265](#)  
Breath Test, [265](#)  
Breath-Holding Spell, [265](#)  
Breath-Holding Spells, [265](#)  
Breathalyzer, [265](#)  
Breathing, [265](#)  
Breathing Difficulty, [265](#)  
Breathing Difficulty First Aid, [265](#)  
Breathing Difficulty, Recumbent, [1421](#)  
Breathing Exercises, [266](#)  
Breathlessness, [266](#)  
Breech Baby, [266](#)  
Breech Birth, [266](#)  
Breech Delivery, [266](#)  
Breech Presentation, [266](#)  
Bregma, [266](#)  
Bremsstrahlung, [266](#)  
Brenner Tumor, [266](#)  
Brenner Tumour, [266](#)

---

---

Breslau Bacillus, 266  
Breslow Thickness, 266  
Bretylum Tosylate, 266  
Brevibacillus, 267  
Brevibacterium, 267  
Brevibacterium Casei, 267  
Brevundimonas, 267  
Brevundimonas Diminuta, 267  
Brevundimonas Vesicularis, 267  
Bricklayer's Disease, 267  
Bridge To Transplant, 267  
Bridge Work, 267  
Brief Psychotic Disorder, 267  
Brief Resolved Unexplained Event, 267  
Brill's Disease, 267  
Brill-Zinsser Disease, 268  
Brimonidine Tartrate, 268  
Brine Bath, 268  
Brittle Asthma, 268  
Brittle Bone Disease, 268  
Brittle Nail Syndrome, 268  
Brivudine, 268  
Broad Casts, 268  
Broad Ligament, 268  
Broad Nasal Bridge, 268  
Broad-Based Gait, 268  
Broad-Complex Tachycardia, 268  
Broca Aphasia, 268  
Broca's Aphasia, 269  
Broca's Area, 269  
Brodie Abscess, 269  
Brodie's Abscess, 269  
Brodmann Area, 269  
Brody Disease, 269  
Broken Ankle, 269  
Broken Arm, 269  
Broken Blood Vessel in Eye, 269  
Broken Bone, 269  
Broken Clavicle, 269  
Broken Collarbone, 269  
Broken Finger, 269  
Broken Foot, 270  
Broken Hand, 270  
Broken Heart Syndrome, 270  
Broken Hip, 270  
Broken Leg, 270  
Broken Little Finger, 270  
Broken Nose, 270  
Broken Or Dislocated Jaw, 270  
Broken Or Knocked Out Tooth, 270  
Broken Ribs, 270  
Broken Toe, 270  
Broken Wrist, 270  
Bromelain, 270  
Bromhexine, 270  
Bromhidrosis, 271  
Bromides, 271  
Bromine, 271  
Bromism, 271  
Bromobenzenes, 271  
Bromocriptine, 271  
Bromoderma, 271  
Bromodosis, 271  
Bromophos, 271  
Brompheniramine Overdose, 271  
Bromthymol Blue, 271  
Bronchi, 271  
Bronchial, 271  
Bronchial Artery, 272  
Bronchial Artery Embolization, 272  
Bronchial Asthma, 272  
Bronchial Biopsy, 272  
Bronchial Breath Sounds, 272  
Bronchial Carcinoid, 272  
Bronchial Carcinoma, 272  
Bronchial Circulation, 272  
Bronchial Fistula, 272  
Bronchial Hemorrhage, 272  
Bronchial Hyperresponsiveness, 272  
Bronchial Obstruction, 272  
Bronchial Stenosis, 272  
Bronchial Stricture, 272  
Bronchial Tree, 273  
Bronchial Tubes, 273  
Bronchial Vein, 273  
Bronchiectasis, 273  
Bronchiole, 273  
Bronchiolectasis, 273  
Bronchioles, 273  
Bronchiolitis, 273  
Bronchiolitis Obliterans, 273  
Bronchitis, 273  
Bronchitis, Acute, 28  
Bronchitis, Chronic, 397  
Bronchoalveolar Lavage, 273  
Bronchocele, 273  
Bronchocentric Granulomatosis, 273  
Bronchoconstriction, 274  
Bronchoconstrictor Agents, 274  
Bronchodilator, 274  
Bronchodilator Response, 274  
Bronchodilator Reversibility, 274  
Bronchoesophageal Fistula, 274  
Bronchogenic Carcinoma, 274  
Bronchogenic Cyst, 274  
Bronchomalacia, 274  
Bronchophony, 274  
Bronchopleural Fistula, 274

---

---

Bronchopneumonia, [274](#)  
 Bronchoprovocation Test, [274](#)  
 Bronchopulmonary Dysplasia, [275](#)  
 Bronchopulmonary Segments, [275](#)  
 Bronchoscope, [275](#)  
 Bronchoscopic Culture, [275](#)  
 Bronchoscopy, [275](#)  
 Bronchospasm, [275](#)  
 Bronchostenosis, [275](#)  
 Bronchus, [275](#)  
 Bronchus Disease, [275](#)  
 Bronchus Neoplasm, [275](#)  
 Brooke Disease, [275](#)  
 Brooke-Spiegler Syndrome, [275](#)  
 Broselow Tape, [275](#)  
 Brow Presentation, [276](#)  
 Brown Adipose Tissue, [276](#)  
 Brown Fat, [276](#)  
 Brown Lung Disease, [276](#)  
 Brown Recluse Spider, [276](#)  
 Brown Recluse Spider Bite, [276](#)  
 Brown Sequard Syndrome, [276](#)  
 Brown Stain, [276](#)  
 Brown Syndrome, [276](#)  
 Brown Teeth, [276](#)  
 Brown-Sequard Syndrome, [276](#)  
 Brown-SéQuard Syndrome, [276](#)  
 Bruce Protocol, [276](#)  
 Brucella, [277](#)  
 Brucella Abortus, [277](#)  
 Brucella Canis, [277](#)  
 Brucella Melitensis, [277](#)  
 Brucella Suis, [277](#)  
 Brucellaceae, [277](#)  
 Brucellosis, [277](#)  
 Bruch's Membrane, [277](#)  
 Bruck's Syndrome, [277](#)  
 Brudzinski Sign, [277](#)  
 Brugada ECG Pattern Type 1, [277](#)  
 Brugada Phenocopy, [277](#)  
 Brugada Syndrome, [277](#)  
 Brugia, [278](#)  
 Bruise, [278](#)  
 Bruises, [278](#)  
 Bruising, [278](#)  
 Bruit, [278](#)  
 Brunner Gland, [278](#)  
 Brunner Glands, [278](#)  
 Brunner's Gland, [278](#)  
 Brunner's Glands, [278](#)  
 Bruns Syndrome, [278](#)  
 Brush Burns, [278](#)  
 Brushfield's Spots, [278](#)  
 Bruton Agammaglobulinemia, [278](#)  
 Bruxism, [279](#)  
 Bryant's Traction, [279](#)  
 Bubo, [279](#)  
 Bubonic Plague, [279](#)  
 Buccal, [279](#)  
 Buccal Gland, [279](#)  
 Buccal Mucosa, [279](#)  
 Buccal Smear, [279](#)  
 Buccalmucosa, [279](#)  
 Buccinator, [279](#)  
 Buckle Fracture, [279](#)  
 Budd-Chiari Syndrome, [279](#)  
 Budesonide, [279](#)  
 Budvicia, [280](#)  
 Budvicia Aquatica, [280](#)  
 Buerger Disease, [280](#)  
 Buerger Test, [280](#)  
 Buerger's Disease, [280](#)  
 Buerger's Disease, [280](#)  
 Buffalo Hump, [280](#)  
 Buffer, [280](#)  
 Buffer System, [280](#)  
 Buffy Coat, [280](#)  
 Bufonidae, [280](#)  
 Buformin, [280](#)  
 Bug Spray Poisoning, [280](#)  
 Bulbar, [281](#)  
 Bulbar Conjunctiva, [281](#)  
 Bulbar Palsy, [281](#)  
 Bulbar Paralysis, [281](#)  
 Bulbocavernosus, [281](#)  
 Bulbospinal Muscular Atrophy, X-linked, [1821](#)  
 Bulbospongiosus, [281](#)  
 Bulbourethral Gland, [281](#)  
 Bulbourethral Glands, [281](#)  
 Bulging, [281](#)  
 Bulging Eyes, [281](#)  
 Bulging Fontanelle, [281](#)  
 Bulimia, [281](#)  
 Bulimia Nervosa, [281](#)  
 Bulla, [281](#)  
 Bullae, [281](#)  
 Bullectomy, [282](#)  
 Bullet Track, [282](#)  
 Bullet Wounds, [282](#)  
 Bullosa Hereditaria, [282](#)  
 Bullous, [282](#)  
 Bullous Disease of Diabetes (Bullosis Diabeticorum), [282](#)  
 Bullous Emphysema, [282](#)  
 Bullous Impetigo, [282](#)  
 Bullous Pemphigoid, [282](#)  
 Bullous Systemic Lupus Erythematosus (BSLE), [282](#)  
 Bully, [282](#)  
 Bullying, [282](#)

---

---

Bumetanide, [283](#)  
Bumps, [283](#)  
BUN, [283](#)  
BUN-to-Creatinine Ratio, [283](#)  
Bundle Branch Block, [283](#)  
Bundle Branch Reentry Tachycardia, [283](#)  
Bundle Of His, [283](#)  
Bundle of Kent, [283](#)  
Bundle-Branch Block, [283](#)  
Bungarotoxins, [283](#)  
Bunion, [283](#)  
Bunion Removal, [283](#)  
Bunionectomy, [283](#)  
Bunionette, [283](#)  
Bunions, [284](#)  
Bunioplasty, [284](#)  
Bunyaviridae, [284](#)  
Bunyavirus Infection, [284](#)  
Bunyaviruses, [284](#)  
Buphthalmos, [284](#)  
Bupivacaine, [284](#)  
Bupivacaine Hydrochloride, [284](#)  
Buprenorphine, [284](#)  
Buprenorphine/Naloxone Toxicity, [284](#)  
Bupropion, [284](#)  
Buried Penis, [284](#)  
Burkholderia, [284](#)  
Burkholderia Cenocepacia, [285](#)  
Burkholderia Cepacia, [285](#)  
Burkholderia Gladioli, [285](#)  
Burkholderia Mallei, [285](#)  
Burkholderia Pseudomallei, [285](#)  
Burkholderiaceae, [285](#)  
Burkitt Lymphoma, [285](#)  
Burkitt's Lymphoma, [285](#)  
Burkitt's Tumor, [285](#)  
Burn, [285](#)  
Burn Injury, [285](#)  
Burn Shock, [285](#)  
Burn Surgery, [285](#)  
Burn Wound Infections, [285](#)  
Burning Feet, [286](#)  
Burning Mouth Syndrome, [286](#)  
Burning Thigh Pain, [286](#)  
Burnout, [286](#)  
Burns, [286](#)  
Burns, First Aid, [746](#)  
Burnt-Out Joint, [286](#)  
Burping, [286](#)  
Burr Cells, [286](#)  
Burr Hole Evacuation, [286](#)  
Burroughs, [286](#)  
Bursa, [286](#)  
Bursa Inflammation, [287](#)  
Bursectomy, [287](#)  
Bursitis, [287](#)  
Bursitis Of The Heel, [287](#)  
Bursitis of the Knee, [287](#)  
Bursitis, Calcific, [295](#)  
Bursitis, Shoulder, [1782](#)  
Bursopathy, [287](#)  
Burton Line, [287](#)  
Buruli Ulcer, [287](#)  
Burundanga, [287](#)  
Buserelin, [287](#)  
Buspirone, [287](#)  
Busulfan, [287](#)  
Butanol, [288](#)  
Butazolidin Overdose, [288](#)  
Butocarboxim, [288](#)  
Butter, Margarine, And Cooking Oils, [553](#)  
Buttiauxella, [288](#)  
Buttiauxella Agrestis, [288](#)  
Buttiauxella Ferragutiae, [288](#)  
Buttiauxella Gaviniae, [288](#)  
Buttiauxella Noackiae, [288](#)  
Buttock, [288](#)  
Buttocks, [288](#)  
Button Batteries, [288](#)  
Button Magnets In The Nasal Cavity, [288](#)  
Buttonhole Deformity, [288](#)  
Buttressing, [288](#)  
Butylamine, [289](#)  
Butyric Acid, [289](#)  
Bx, [289](#)  
By-Pass, [289](#)  
Bypass, [289](#)  
Bypass Graft, [289](#)  
Bypass Surgery, [289](#)  
Byssinosis, [289](#)  
  
C Fiber, [291](#)  
C-11 Hydroxylase Deficiency, [291](#)  
C-17 Hydroxylase Deficiency, [291](#)  
C-Arm Fluoroscopy, [291](#)  
C-Peptide, [291](#)  
C-Reactive Protein, [291](#)  
C-Reactive Protein Test, [291](#)  
C-Section, [291](#)  
C. Diff, [291](#)  
C. Difficile, [291](#)  
C. Difficile Colitis, [291](#)  
C. Difficile Infection, [291](#)  
C1 Esterase Inhibitor, [2115](#)  
C1-INH, [292](#)  
C1Q Nephropathy, [292](#)  
C282Y Mutation, [292](#)  
CA 19-9, [292](#)

---

---

CA 27-29, 292  
CA-125, 292  
CA-125 Blood Test, 292  
Cabazitaxel, 292  
Cabergoline, 292  
CABG, 292  
Cabot Ring, 292  
Cachectic, 292  
Cachetic, 292  
Cachexia, 292  
Cacosmia, 293  
CAD, 293  
CAD (Coronary Artery Disease), 293  
CADASIL Syndrome, 293  
Cadherin, 293  
Cadmium, 293  
Caduceus, 293  
Cadusafos, 293  
Caecal, 293  
Caecitis, 293  
Caecopexy, 293  
Caecostomy, 293  
Caecotomy, 293  
Caecum, 293  
Caesarean Birth, 294  
Caesarean Section, 294  
Cafe-Au-Lait Macules, 294  
Cafe-Au-Lait Spot, 294  
Caffeine, 294  
Caffeine Intoxication, 294  
Caffeine Overdose, 294  
Caffeine Toxicity, 294  
Caffeine Withdrawal, 294  
Café Au Lait Spot, 294  
Café Coronary, 294  
Café-Au-Lait Macules, 294  
Caisson Disease, 294  
Caladium Plant Poisoning, 294  
Calamine, 294  
Calbindin, 295  
Calcaneal Spur, 295  
Calcaneal Tendon, 295  
Calcaneal Valgus, 295  
Calcaneum, 295  
Calcaneus, 295  
Calcarine Sulcus, 295  
Calcifediol, 295  
Calciferol, 295  
Calcific Bursitis, 295  
Calcific Tendinitis, 295  
Calcific Uremic Arteriopathy, 295  
Calcification, 295  
Calcified Granuloma, 296  
Calcineurin, 296  
Calcinosis, 296  
Calcinosis Cutis, 296  
Calciphylaxis, 296  
Calcitonin, 296  
Calcitonin Blood Test, 296  
Calcitriol, 296  
Calcitriol And Analogues, 296  
Calcium, 296  
Calcium Blood Test, 296  
Calcium Carbonate, 297  
Calcium Carbonate Overdose, 297  
Calcium Carbonate With Magnesium Overdose, 297  
Calcium Channel, 297  
Calcium Channel Blocker, 2115  
Calcium Channel Blocker Overdose, 2115  
Calcium Channel Blocker Toxicity, 297  
Calcium Channel Blockers, 2115  
Calcium Chloride, 297  
Calcium Citrate, 297  
Calcium Deficiency, 297  
Calcium for Osteoporosis, 297  
Calcium Gluconate, 297  
Calcium Hydroxide Poisoning, 297  
Calcium Phosphate, 297  
Calcium Pyrophosphate Arthritis, 297  
Calcium Pyrophosphate Deposition, 298  
Calcium Pyrophosphate Deposition Disease, 298  
Calcium Sensing Receptor, 298  
Calcium Signaling, 298  
Calcium Stones, 298  
Calcium, Elevated, 962  
Calculi, 298  
Calculus, 298  
Calculus Removal, 298  
Caldwell-Luc Operation, 298  
Calf, 298  
Calf Pseudohypertrophy, 298  
Calibration, 298  
Caliciviridae, 298  
Calipers, 299  
Calla Lily Poisoning, 299  
Callosal, 299  
Callosity, 299  
Callosotomy, 299  
Callus, 299  
Calluses and Corns, 299  
Calmodulin, 299  
Calnexin, 299  
Calor, 299  
Calor, Dolor, Rubor, And Tumor, 323  
Caloric Restriction, 299  
Caloric Stimulation, 299  
Calorie, 299

---



---

Calorimetry, [299](#)  
 Calot Triangle, [300](#)  
 Calpain, [300](#)  
 Calpainopathy, [300](#)  
 Calponin, [300](#)  
 Calprotectin, [300](#)  
 Calreticulin, [300](#)  
 Calretinin, [300](#)  
 Calvaria, [300](#)  
 Calvarium, [300](#)  
 Calyx, [300](#)  
 Cambridge Classification, [300](#)  
 Cameron Lesions, [300](#)  
 Campho-Phenique Overdose, [300](#)  
 Camphor, [301](#)  
 Camphor Overdose, [301](#)  
 Campimetry, [301](#)  
 Campomelic Dysplasia, [301](#)  
 Camptocormia, [301](#)  
 Camptodactyly, [301](#)  
 Camptothecin Analogue, [2115](#)  
 Campylobacter, [301](#)  
 Campylobacter Coli, [301](#)  
 Campylobacter Enteritis, [301](#)  
 Campylobacter Fetus, [301](#)  
 Campylobacter Food Poisoning, [301](#)  
 Campylobacter Infection, [301](#)  
 Campylobacter Jejuni, [302](#)  
 Campylobacter Serology Test, [302](#)  
 Campylobacteriosis, [302](#)  
 Canagliflozin, [302](#)  
 Canal Of Schlemm, [302](#)  
 Canaliculus, [302](#)  
 Canalith Repositioning Maneuver, [302](#)  
 Canalith Repositioning Maneuvers, [302](#)  
 Canalization, [302](#)  
 Canavan Disease, [302](#)  
 Cancellous Bone, [302](#)  
 Cancer, [302](#)  
 Cancer And Lymph Nodes, [302](#)  
 Cancer Care Team, [303](#)  
 Cancer Cell, [303](#)  
 Cancer Cluster, [303](#)  
 Cancer Genetics, [303](#)  
 Cancer Grading, [303](#)  
 Cancer Immunotherapy, [303](#)  
 Cancer in Lung, [303](#)  
 Cancer Mutations, [303](#)  
 Cancer Nutrition, [303](#)  
 Cancer of the Ear, [303](#)  
 Cancer of the Tongue, [303](#)  
 Cancer Of Unknown Origin, [303](#)  
 Cancer Of Unknown Primary, [303](#)  
 Cancer Pain, [303](#)  
 Cancer Prevention, [304](#)  
 Cancer Progression, [304](#)  
 Cancer Recurrence, [304](#)  
 Cancer Risk, [304](#)  
 Cancer Screening, [304](#)  
 Cancer Staging, [304](#)  
 Cancer Surveillance, [304](#)  
 Cancer Survivor, [304](#)  
 Cancer Susceptibility Genes, [304](#)  
 Cancer Treatment, [304](#)  
 Cancer Treatment-Related Bleeding, [304](#)  
 Cancer Treatment-Induced Premature Menopause, [304](#)  
 Cancer Treatments, [305](#)  
 Cancer Vaccine, [2115](#)  
 Cancer, Carcinoid Tumors, [316](#)  
 Cancer, Chronic Lymphocytic Leukemia, [400](#)  
 Cancer, Chronic Myelogenous Leukemia, [401](#)  
 Cancer, Colon, [429](#)  
 Cancer, Endometrial, [642](#)  
 Cancer, Esophageal, [675](#)  
 Cancer, Eye Melanoma, [705](#)  
 Cancer, Floor of the Mouth, [753](#)  
 Cancer, Gallbladder, [797](#)  
 Cancer, Gastric, [1844](#)  
 Cancer, Hairy Cell Leukemia, [877](#)  
 Cancer, Hodgkin Lymphoma, [941](#)  
 Cancer, Hurthle Cell, [952](#)  
 Cancer, Inflammatory Breast, [1018](#)  
 Cancer, Kidney, [1089](#)  
 Cancer, Liver, [1151](#)  
 Cancer, Lung, [1168](#)  
 Cancer, Merkel Cell, [1235](#)  
 Cancer, Mouth, [1289](#)  
 Cancer, Multiple Myeloma, [1301](#)  
 Cancer, Nasopharyngeal, [1331](#)  
 Cancer, Neuroblastoma, [1346](#)  
 Cancer, Neuroendocrine, [1347](#)  
 Cancer, Non-Hodgkin Lymphoma, [1367](#)  
 Cancer, Ovarian, [1434](#)  
 Cancer, Pancreatic, [1453](#)  
 Cancer, Paraneoplastic Syndromes, [1461](#)  
 Cancer, Prostate, [1609](#)  
 Cancer, Rectal, [1674](#)  
 Cancer, Retinoblastoma, [1705](#)  
 Cancer, Skin, [1797](#)  
 Cancer, Soft Tissue Sarcoma, [1812](#)  
 Cancer, Squamous Cell, [1833](#)  
 Cancer, Stomach, [1844](#)  
 Cancer, Testicular, [1904](#)  
 Cancer, Throat, [1915](#)  
 Cancer, Thyroid, [1920](#)  
 Cancer, Uterine, [642](#)  
 Cancer, Vagina, [2017](#)  
 Cancer, Vulvar, [2061](#)

---

---

Cancer, Wilms' Tumor, [2078](#)  
Cancer-Associated Thrombosis, [305](#)  
Cancer-Related Fatigue, [305](#)  
Cancers of the Oral Mucosa, [305](#)  
Cancrum Oris, [305](#)  
Candela, [305](#)  
Candida, [305](#)  
Candida Albicans, [305](#)  
Candida Albicans Infection, [305](#)  
Candida Auris, [305](#)  
Candida Auris Infection, [305](#)  
Candida Dubliniensis, [305](#)  
Candida Esophagitis, [305](#)  
Candida Glabrata, [306](#)  
Candida Infection Of The Skin, [306](#)  
Candida Parapsilosis, [306](#)  
Candida Tropicalis, [306](#)  
Candidal Intertrigo, [306](#)  
Candidal Vaginitis, [306](#)  
Candidal Vulvovaginitis, [306](#)  
Candidemia, [306](#)  
Candidiasis, [306](#)  
Candidiasis, Oral, [1412](#)  
Candidiasis, Vaginal, [2017](#)  
Canefield Fever, [306](#)  
Canicola Fever, [306](#)  
Canine Teeth, [306](#)  
Canker, [306](#)  
Canker Sore, [306](#)  
Canker Sores, [307](#)  
Cannabidiol, [307](#)  
Cannabinoid, [307](#)  
Cannabinoid Hyperemesis Syndrome, [307](#)  
Cannabinoid Receptor, [307](#)  
Cannabinol, [307](#)  
Cannabis, [307](#)  
Cannabis Use Disorder, [307](#)  
Cannabis-Related Disorders, [307](#)  
Cannot Intubate, Cannot Oxygenate, [405](#)  
Cannula, [307](#)  
Cannula Hub, [307](#)  
Cannulation, [307](#)  
Canola Oil, [307](#)  
Canrenone, [307](#)  
Cantharides, [308](#)  
Canthotomy, [308](#)  
Canthus, [308](#)  
Cantlie Line, [308](#)  
Cantu Syndrome, [308](#)  
Capacitation, [308](#)  
Capacity, [308](#)  
Capacity And Competence, [308](#)  
Capecitabine, [308](#)  
Capgras Delusion, [308](#)  
Capgras Syndrome, [308](#)  
Capillaria, [308](#)  
Capillaria Philippinensis, [308](#)  
Capillaroscopy, [309](#)  
Capillary, [309](#)  
Capillary Bed, [309](#)  
Capillary Blood Gas, [309](#)  
Capillary Electrophoresis, [309](#)  
Capillary Exchange, [309](#)  
Capillary Hemangioma, [309](#)  
Capillary Leak Syndrome, [309](#)  
Capillary Malformation, [309](#)  
Capillary Microscope, [309](#)  
Capillary Nail Refill Test, [309](#)  
Capillary Permeability, [309](#)  
Capillary Plasma, [310](#)  
Capillary Refill, [310](#)  
Capillary Refill Time, [310](#)  
Capillary Sample, [310](#)  
Capillary Vessel, [310](#)  
Capitate, [310](#)  
Capitation, [310](#)  
Caplan Syndrome, [310](#)  
Caplan's Syndrome, [310](#)  
Capnocytophaga, [310](#)  
Capnocytophaga Canimorsus, [310](#)  
Capnocytophaga Infection, [310](#)  
Capnocytophaga Ochracea, [310](#)  
Capnograph, [311](#)  
Capnography, [311](#)  
Caps, [311](#)  
Capsaicin, [311](#)  
Capsid, [311](#)  
Capsid Proteins, [311](#)  
Capsular Contracture, [311](#)  
Capsular Warning Syndrome, [311](#)  
Capsule, [311](#)  
Capsule Endoscope, [311](#)  
Capsule Endoscopy, [311](#)  
Capsule Of Glisson, [311](#)  
Capsule Of Tenon, [311](#)  
Capsulectomy, [312](#)  
Capsulotomy, [312](#)  
Captopril, [312](#)  
Captopril Renal Scan, [312](#)  
Caput Medusae, [312](#)  
Caput Succedaneum, [312](#)  
CAR T-Cell, [312](#)  
CAR T-Cell Therapy, [312](#)  
Carbachol, [312](#)  
Carbamate, [312](#)  
Carbamazepine, [312](#)  
Carbamazepine Toxicity, [312](#)  
Carbamide, [312](#)

---

---

Carbamoyl Phosphate Synthetase Deficiency, [313](#)  
Carbapenem, [313](#)  
Carbapenem-Resistant Enterobacteriaceae, [313](#)  
Carbapenem-Resistant Enterobacteriaceae Infection, [313](#)  
Carbapenem-Resistant Organisms, [313](#)  
Carbaryl, [313](#)  
Carbidopa, [313](#)  
Carbimazole, [313](#)  
Carbofuran, [313](#)  
Carbohydrate, [313](#)  
Carbohydrate Counting, [313](#)  
Carbohydrate Metabolism, [313](#)  
Carbohydrates, [313](#)  
Carbolic Acid, [313](#)  
Carbolic Acid Poisoning, [314](#)  
Carbon, [314](#)  
Carbon Dioxide, [314](#)  
Carbon Dioxide Absorber, [314](#)  
Carbon Dioxide Transport, [314](#)  
Carbon Ion Therapy, [314](#)  
Carbon Monoxide, [314](#)  
Carbon Monoxide Poisoning, [314](#)  
Carbon Monoxide Toxicity, [314](#)  
Carbon Tetrachloride, [314](#)  
Carbonate, [314](#)  
Carbonic Anhydrase Inhibitors, [314](#)  
Carbophenothion, [314](#)  
Carboplatin, [315](#)  
Carboprost, [315](#)  
Carbosulfan, [315](#)  
Carboxy-Lyases, [315](#)  
Carboxyhaemoglobin, [315](#)  
Carboxyhemoglobin, [315](#)  
Carboxyhemoglobinemia, [315](#)  
Carboxyl Group, [315](#)  
Carboxylesterase, [315](#)  
Carboxylic Acid, [315](#)  
Carboxypeptidase H, [315](#)  
Carbuncle, [315](#)  
Carbuncles and Boils, [315](#)  
Carcinoembryonal Antigen, [315](#)  
Carcinoembryonic Antigen, [316](#)  
Carcinoembryonic Antigen Tumor Marker, [316](#)  
Carcinogen, [316](#)  
Carcinogenesis, [316](#)  
Carcinogenic, [316](#)  
Carcinoid, [316](#)  
Carcinoid Flushing, [316](#)  
Carcinoid Heart Disease, [316](#)  
Carcinoid Syndrome, [316](#)  
Carcinoid Tumor, [316](#)  
Carcinoid Tumors, [316](#)  
Carcinoma, [316](#)  
Carcinoma En Cuirasse, [317](#)  
Carcinoma Erysipeloides, [317](#)  
Carcinoma In Situ, [317](#)  
Carcinoma Of Unknown Primary, [317](#)  
Carcinomatosis, [317](#)  
Carcinosarcoma, [317](#)  
Cardia, [317](#)  
Cardiac, [317](#)  
Cardiac Ablation, [317](#)  
Cardiac Ablation Procedures, [317](#)  
Cardiac Allograft, [317](#)  
Cardiac Allograft Vasculopathy, [317](#)  
Cardiac Amyloidosis, [317](#)  
Cardiac Amyloidosis Diagnosis, [318](#)  
Cardiac Amyloidosis Staging, [318](#)  
Cardiac Arrest, [318](#)  
Cardiac Arrhythmia, [318](#)  
Cardiac Arrhythmias, [318](#)  
Cardiac Axis, [318](#)  
Cardiac Biomarkers, [318](#)  
Cardiac Cachexia, [318](#)  
Cardiac Catheter, [318](#)  
Cardiac Catheterization, [318](#)  
Cardiac Conduction System, [318](#)  
Cardiac Contractility, [318](#)  
Cardiac CT, [318](#)  
Cardiac CT Angiography, [319](#)  
Cardiac Cycle, [319](#)  
Cardiac Defibrillation, [319](#)  
Cardiac Device Infections, [319](#)  
Cardiac Diet, [319](#)  
Cardiac Electrophysiology Study, [319](#)  
Cardiac Enzymes, [319](#)  
Cardiac Event Monitors, [319](#)  
Cardiac Failure, [319](#)  
Cardiac Fibroma, [319](#)  
Cardiac Gated SPECT, [319](#)  
Cardiac Glycoside, [319](#)  
Cardiac Glycoside Overdose, [319](#)  
Cardiac Glycoside Plant Poisoning, [319](#)  
Cardiac Hemangioma, [320](#)  
Cardiac Imaging Tests, [320](#)  
Cardiac Index, [320](#)  
Cardiac Intravascular Ultrasound, [320](#)  
Cardiac Ischemia, [320](#)  
Cardiac Markers, [320](#)  
Cardiac Metastases, [320](#)  
Cardiac MRI, [320](#)  
Cardiac Muscle, [320](#)  
Cardiac Myocyte, [320](#)  
Cardiac Myocytes, [320](#)  
Cardiac Myxoma, [320](#)  
Cardiac Output, [320](#)  
Cardiac Output Measurement, [320](#)  
Cardiac Pacemaker, [321](#)

---

---

Cardiac Pacemaker Types, 321  
Cardiac Perforation, 321  
Cardiac PET, 321  
Cardiac PET Imaging, 321  
Cardiac Power Output, 321  
Cardiac Radiation Injury, 321  
Cardiac Radionuclide Imaging, 321  
Cardiac Rehab Outcomes, 321  
Cardiac Rehabilitation, 321  
Cardiac Rehabilitation Phases, 321  
Cardiac Resynchronization Therapy, 321  
Cardiac Rhabdomyoma, 322  
Cardiac Sarcoidosis, 322  
Cardiac Sarcoma, 322  
Cardiac Septum, 322  
Cardiac Skeleton, 322  
Cardiac SPECT, 322  
Cardiac Stress Test, 322  
Cardiac Stress Testing, 322  
Cardiac Syncope, 322  
Cardiac Syndrome X, 322  
Cardiac Tamponade, 322  
Cardiac Tamponade Physiology, 323  
Cardiac Transplantation, 323  
Cardiac Tumors, 323  
Cardiac Ventriculography, 323  
Cardiac Volume, 323  
Cardiac Wall, 323  
Cardinal Body Part, 323  
Cardinal Cell Part, 323  
Cardinal Ligament, 323  
Cardinal Signs of Inflammation, 323  
Cardio-Oncology, 323  
Cardiobacteriaceae, 323  
Cardiobacterium, 323  
Cardiobacterium Hominis, 323  
Cardiofaciocutaneous Syndrome, 324  
Cardiogenic Shock, 324  
Cardiolipin, 324  
Cardiologist, 324  
Cardiology, 324  
Cardiomegaly, 324  
Cardiometabolic, 324  
Cardiomyopathy, 324  
Cardiomyopathy Classification, 324  
Cardiomyopathy, Dilated, 560  
Cardiomyopathy, Hypertrophic, 971  
Cardiomyopathy, Restrictive, 1700  
Cardiopathy, 324  
Cardioplegia, 324  
Cardiopulmonary, 324  
Cardiopulmonary Bypass, 324  
Cardiopulmonary Bypass Machine, 325  
Cardiopulmonary Exercise Testing, 325  
Cardiopulmonary Resuscitation, 325  
Cardiopulmonary Resuscitation, Child, 1479  
Cardiopulmonary Resuscitation, Infant, 1011  
Cardiorenal Syndrome, 325  
Cardiorespiratory, 325  
Cardiorespiratory Endurance, 325  
Cardioselective Beta-Blockers, 325  
Cardiospasm, 325  
Cardiotocography, 325  
Cardiotonic Agent, 2115  
Cardiotoxicity, 325  
Cardiotoxicity From Chemotherapy, 325  
Cardiotoxin, 326  
Cardiovascular, 326  
Cardiovascular Agents, 326  
Cardiovascular Changes In Pregnancy, 326  
Cardiovascular Disease, 326  
Cardiovascular Disease and Pregnancy, 326  
Cardiovascular Health, 326  
Cardiovascular Infection, 326  
Cardiovascular Injury, 326  
Cardiovascular Neoplasm, 326  
Cardiovascular Pharmacology, 326  
Cardiovascular Prevention in Diabetes, 326  
Cardiovascular Risk Reduction, 326  
Cardiovascular Screening, 327  
Cardiovascular Surgery, 327  
Cardiovascular System, 327  
Cardiovascular-Kidney-Metabolic (CKM) Syndrome, 327  
Cardioversion, 327  
Cardiovirus, 327  
Carditis, 327  
Care Coordination, 327  
Caregiver, 327  
Caregiver Burden, 327  
Caregiver Stress, 327  
Caries, 327  
Caries Risk Assessment, 327  
Carina, 328  
Cariogenic, 328  
Cariogram, 328  
Carisoprodol, 328  
Carminative, 328  
Carmine, 328  
Carmustine, 328  
Carnett Sign, 328  
Carney Complex, 328  
Carney Triad, 328  
Carnitine, 328  
Carnitine Deficiency, 328  
Carnitine Palmitoyltransferase II Deficiency, 329  
Carnosine, 329  
Carnosinemia, 329  
Caroli Disease, 329

---

---

Carotene, [329](#)  
 Carotenemia, [329](#)  
 Carotenoid, [329](#)  
 Carotenoids, [329](#)  
 Carotid, [329](#)  
 Carotid Artery, [329](#)  
 Carotid Artery Disease, [329](#)  
 Carotid Artery Dissection, [329](#)  
 Carotid Artery Stenosis, [329](#)  
 Carotid Artery Stenting, [330](#)  
 Carotid Body, [330](#)  
 Carotid Body Tumor, [330](#)  
 Carotid Bruit, [330](#)  
 Carotid Cavernous Fistula, [330](#)  
 Carotid Duplex Ultrasound, [330](#)  
 Carotid Endarterectomy, [330](#)  
 Carotid Glomus, [330](#)  
 Carotid Occlusion, [330](#)  
 Carotid Sinus Hypersensitivity, [330](#)  
 Carotid Sinus Massage, [330](#)  
 Carotid Stenosis, [331](#)  
 Carpal Bones, [331](#)  
 Carpal Joints, [331](#)  
 Carpal Tunnel, [331](#)  
 Carpal Tunnel Biopsy, [331](#)  
 Carpal Tunnel Release, [331](#)  
 Carpal Tunnel Syndrome, [331](#)  
 Carpal Tunnel Syndrome Tests, [331](#)  
 Carpometacarpal Joint, [331](#)  
 Carpus, [331](#)  
 Carrageenan, [331](#)  
 Carrier, [331](#)  
 Carrier Excipient, [332](#)  
 Carrier Screening, [332](#)  
 Carrier State, [332](#)  
 Carrier-Mediated Transport, [332](#)  
 Carry-Over Effect, [332](#)  
 Carteolol, [332](#)  
 Cartilage, [332](#)  
 Cartilage Disease, [332](#)  
 Cartilaginous Joint, [332](#)  
 Caruncles, [332](#)  
 Carvallo Sign, [332](#)  
 Carvedilol, [332](#)  
 Carvedilol Phosphate, [2115](#)  
 CAS, [332](#)  
 Casanthranol, [332](#)  
 Cascade, [333](#)  
 Cascara, [333](#)  
 Case Definition, [333](#)  
 Case Fatality Rate, [333](#)  
 Case Management, [333](#)  
 Case-Fatality Rate, [333](#)  
 Casein, [333](#)  
 Casein Kinase, [333](#)  
 Caseous, [333](#)  
 Caseous Necrosis, [333](#)  
 Cashew Allergy, [333](#)  
 Caspase, [333](#)  
 Caspase Activation, [333](#)  
 Caspofungin, [333](#)  
 Caspofungin Acetate, [334](#)  
 Cassette, [334](#)  
 Cast, [334](#)  
 Cast Nephropathy, [334](#)  
 Castleman Disease, [334](#)  
 Castor Oil, [334](#)  
 Castor Oil Overdose, [334](#)  
 Castration, [334](#)  
 Cat Allergy, [334](#)  
 CAT Scan, [334](#)  
 Cat Scratch Disease, [334](#)  
 Cat's Eye Reflex, [334](#)  
 Cat-Nap, [334](#)  
 Cat-Scratch Disease, [334](#)  
 Catabolism, [335](#)  
 Catagen, [335](#)  
 Catalase, [335](#)  
 Catalepsy, [335](#)  
 Catalyst, [335](#)  
 Catamenial Epilepsy, [335](#)  
 Catamenial Pneumothorax, [335](#)  
 Cataplexy, [335](#)  
 Cataract, [335](#)  
 Cataract Removal, [335](#)  
 Cataract Surgery, [335](#)  
 Cataracts, [335](#)  
 Catastrophic Aging, [335](#)  
 Catastrophic Reaction, [336](#)  
 Catatonia, [336](#)  
 Catbite, [336](#)  
 Catch-Up Immunization, [336](#)  
 Catechin, [336](#)  
 Catechol, [336](#)  
 Catecholamine, [336](#)  
 Catecholamine Blood Test, [336](#)  
 Catecholamine Crisis, [336](#)  
 Catecholaminergic Polymorphic Ventricular Tachycardia, [336](#)  
 Catecholamines, [336](#)  
 Category Five Hurricane, [336](#)  
 Category I Fetal Heart Tracing, [337](#)  
 Category II Fetal Heart Tracing, [337](#)  
 Category III Fetal Heart Tracing, [337](#)  
 Category Three Hurricane, [337](#)  
 Category Two Hurricane, [337](#)  
 Catenin, [337](#)  
 Caterpillar Dermatitis, [337](#)

---

---

Caterpillars, [337](#)  
Cathartic, [337](#)  
Cathepsin, [337](#)  
Cathepsin D, [337](#)  
Cathepsin H, [337](#)  
Cathepsin K, [337](#)  
Cathepsin L, [337](#)  
Cathepsin Z, [338](#)  
Cathepsins B, [338](#)  
Catheter, [338](#)  
Catheter-Associated Urinary Tract Infection, [338](#)  
Catheter-Associated UTI, [338](#)  
Catheter-Directed Thrombolysis, [338](#)  
Catheter-Related Infections, [338](#)  
Catheter-Related UTI, [338](#)  
Catheterization, [338](#)  
Catheterized Urine Culture, [338](#)  
Cathexis, [339](#)  
Cation, [339](#)  
Cauda Equina, [339](#)  
Cauda Equina Syndrome, [339](#)  
Cauda Equina Syndrome Cross-Reference, [339](#)  
Caudad, [339](#)  
Caudal, [339](#)  
Caudal Dysplasia, [339](#)  
Caudate Nucleus, [339](#)  
Cauliflower Ear, [339](#)  
Caulking Compound Poisoning, [339](#)  
Causalgia, [339](#)  
Cause Of Death, [340](#)  
Causes of Cancer, [340](#)  
Caustic, [340](#)  
Cauterization, [340](#)  
Cautery, [340](#)  
Cautery Device, [340](#)  
Caveola, [340](#)  
Caveolins, [340](#)  
Cavernous Hemangioma, [340](#)  
Cavernous Malformations, [340](#)  
Cavernous Sinus, [340](#)  
Cavernous Sinus Thrombosis, [340](#)  
Cavities, [340](#)  
Cavities And Tooth Decay, [341](#)  
Cavity, [341](#)  
Cavum, [341](#)  
CBC, [341](#)  
CBC Blood Test, [341](#)  
CCP, [341](#)  
CCTA, [341](#)  
CCU, [341](#)  
Cd Antigens, [341](#)  
CD4 Count, [341](#)  
CD4-Positive T Cell, [341](#)  
CD8-Positive T Cell, [341](#)  
Cea, [341](#)  
CEA Assay, [341](#)  
CEA Blood Test, [341](#)  
Cebocephaly, [2116](#)  
Cecal Hemorrhage, [342](#)  
Cecal Infection, [342](#)  
Cecal Neoplasm, [342](#)  
Cecal Volvulus, [342](#)  
Cecostomy, [342](#)  
Cecum, [342](#)  
Cedar Leaf Oil Poisoning, [342](#)  
Cefaclor, [342](#)  
Cefadroxil, [342](#)  
Cefamandole, [342](#)  
Cefatrizine, [342](#)  
Cefazolin, [342](#)  
Cefdinir, [342](#)  
Cefepime, [342](#)  
Cefixime, [343](#)  
Cefmenoxime, [343](#)  
Cefmetazole, [343](#)  
Cefonicid, [343](#)  
Cefoperazone, [343](#)  
Cefotaxime, [343](#)  
Cefotetan, [343](#)  
Cefotiam, [343](#)  
Cefoxitin, [343](#)  
Ceftazidime, [343](#)  
Ceftazidime-Avibactam, [343](#)  
Ceftibuten, [343](#)  
Ceftizoxime, [343](#)  
Ceftriaxone, [344](#)  
Cefuroxime, [344](#)  
Celebrex, [344](#)  
Celecoxib, [2116](#)  
Celiac Artery, [344](#)  
Celiac Crisis, [344](#)  
Celiac Disease, [344](#)  
Celiac Plexus, [344](#)  
Celiac Sprue, [344](#)  
Celiac Trunk, [344](#)  
Celiac Trunk Stenosis, [344](#)  
Cell, [344](#)  
Cell Adhesion, [345](#)  
Cell Aggregation, [345](#)  
Cell Body, [345](#)  
Cell Count, [345](#)  
Cell Culture, [345](#)  
Cell Cycle, [345](#)  
Cell Death, [345](#)  
Cell Dedifferentiation, [345](#)  
Cell Density, [345](#)  
Cell Differentiation, [345](#)  
Cell Division, [345](#)

---



---

Cell Fusion, [345](#)  
Cell Growth, [345](#)  
Cell Maturation, [345](#)  
Cell Migration, [345](#)  
Cell Motility, [346](#)  
Cell Movement, [346](#)  
Cell Nucleus, [346](#)  
Cell Pellet, [346](#)  
Cell Physiology, [346](#)  
Cell Polarity, [346](#)  
Cell Proliferation, [346](#)  
Cell Secretion, [346](#)  
Cell Senescence, [346](#)  
Cell Signaling, [346](#)  
Cell Sorting, [346](#)  
Cell Structure, [346](#)  
Cell Surface, [346](#)  
Cell Survival, [346](#)  
Cell Therapy, [347](#)  
Cell Transplantation, [347](#)  
Cell Wall, [347](#)  
Cell-Cell Junction, [347](#)  
Cell-Cell Signaling, [347](#)  
Cell-Free DNA, [347](#)  
Cell-Free Fetal DNA Screening, [347](#)  
Cell-Matrix Junction, [347](#)  
Cell-Mediated Immune Response, [347](#)  
Cell-Mediated Immunity, [347](#)  
Cellophane, [347](#)  
Cells, Stem, [1839](#)  
Cellular, [347](#)  
Cellular Adaptation, [347](#)  
Cellular Immunity, [347](#)  
Cellular Metabolic Process, [348](#)  
Cellular Quiescence, [348](#)  
Cellular Respiration, [348](#)  
Cellular Swelling, [348](#)  
Cellular Vesicle, [348](#)  
Cellulite, [348](#)  
Cellulitis, [348](#)  
Cellulomonas, [348](#)  
Cellulomonas Hominis, [348](#)  
Cellulose, [348](#)  
Cellulosimicrobium, [348](#)  
Cellulosome, [348](#)  
Celosomy, [348](#)  
Cement, [349](#)  
Cemento-Enamel Junction, [349](#)  
Cementoma, [349](#)  
Cementum, [349](#)  
Cenesthesia, [349](#)  
Centenarian, [349](#)  
Center, [349](#)  
Centigram, [349](#)  
Centimeter, [349](#)  
Centimeter, Unit, [349](#)  
Centipede, [349](#)  
Centipoise, [349](#)  
Central, [349](#)  
Central Apnea, [349](#)  
Central Auditory Processing Disorder, [349](#)  
Central Cord Syndrome, [349](#)  
Central Core Disease, [350](#)  
Central Cyanosis, [350](#)  
Central Deafness, [350](#)  
Central Diabetes Insipidus, [350](#)  
Central Fatigue, [350](#)  
Central Hypothyroidism, [350](#)  
Central Line, [350](#)  
Central Line Placement, [350](#)  
Central Line-Associated Bloodstream Infection, [350](#)  
Central Nervous System, [350](#)  
Central Nervous System Germinoma, [350](#)  
Central Nervous System Lymphoma, [350](#)  
Central Nervous System Vascular Malformations, [351](#)  
Central Neurofibromatosis, [351](#)  
Central Pain Syndrome, [351](#)  
Central Pontine Myelinolysis, [351](#)  
Central Precocious Puberty, [351](#)  
Central Retinal Artery Occlusion, [351](#)  
Central Retinal Vein Occlusion, [351](#)  
Central Serous Chorioretinopathy, [351](#)  
Central Serous Choroidopathy, [351](#)  
Central Sleep Apnea, [351](#)  
Central Sleep Apnea Syndromes, [351](#)  
Central Sulcus, [351](#)  
Central Venous Catheter, [352](#)  
Central Venous Catheter Dressing Change, [352](#)  
Central Venous Catheter Flushing, [352](#)  
Central Venous Line, [352](#)  
Central Venous Pressure, [352](#)  
Central Vertigo, [352](#)  
Central Vision, [352](#)  
Centriacinar Emphysema, [352](#)  
Centric-Fusion Translocation, [352](#)  
Centrifugation, [352](#)  
Centrifuge, [352](#)  
Centrilobular Emphysema, [352](#)  
Centriole, [353](#)  
Centromere, [353](#)  
Centronuclear Myopathy, [353](#)  
Centrosome, [353](#)  
Centrum, [353](#)  
Cephacetrile, [353](#)  
Cephalalgia, [353](#)  
Cephalexin, [353](#)  
Cephalgia, [353](#)  
Cephalhaematoma, [353](#)

---

---

Cephalhematoma, 353  
Cephalic, 353  
Cephalic Vein, 353  
Cephalohematoma, 353  
Cephalometry, 354  
Cephalopelvic Disproportion, 354  
Cephaloridine, 354  
Cephalosporin, 354  
Cephalosporins, 354  
Cephalothin, 354  
Cephapirin, 354  
Cephradine, 354  
Ceramide, 354  
Cercaria, 354  
Cercarial Dermatitis, 354  
Cerclage, 354  
Cerebellar, 354  
Cerebellar Ataxia, 354  
Cerebellar Cortex, 354  
Cerebellar Disease, 354  
Cerebellar Dysmetria, 354  
Cerebellar Hemangioblastoma, 355  
Cerebellar Hypoplasia, 355  
Cerebellar Neoplasm, 355  
Cerebellar Nuclei, 355  
Cerebellar Peduncle, 355  
Cerebellar Tonsillar Herniation, 355  
Cerebellar Tumor, 355  
Cerebellar Vermis, 355  
Cerebellitis, 355  
Cerebellopontine Angle, 355  
Cerebellopontine Angle Tumor, 355  
Cerebellum, 355  
Cerebral, 355  
Cerebral Amyloid Angiopathy, 355  
Cerebral Aneurysm, 356  
Cerebral Angiography, 356  
Cerebral Aqueduct, 356  
Cerebral Arteriovenous Malformation, 356  
Cerebral Artery, 356  
Cerebral Atrophy, 356  
Cerebral Autoregulation, 356  
Cerebral Calcification, 356  
Cerebral Cavernous Malformation, 356  
Cerebral Cavernous Malformations, 356  
Cerebral Cortex, 356  
Cerebral Crus, 356  
Cerebral Diplegia, 356  
Cerebral Edema, 357  
Cerebral Embolism, 357  
Cerebral Haemorrhage, 357  
Cerebral Hemisphere, 357  
Cerebral Hemorrhage, 357  
Cerebral Hypoxia, 357  
Cerebral Infarction, 357  
Cerebral Ischemia, 357  
Cerebral Lobes, 357  
Cerebral Malaria, 357  
Cerebral Palsy, 357  
Cerebral Palsy Entry, 357  
Cerebral Peduncle, 357  
Cerebral Perfusion Pressure, 357  
Cerebral Salt Wasting, 358  
Cerebral Salt-Wasting Syndrome, 358  
Cerebral Thrombosis, 358  
Cerebral Toxoplasmosis, 358  
Cerebral Tumor, 358  
Cerebral Tumours, 358  
Cerebral Vasospasm, 358  
Cerebral Venous Sinus Thrombosis, 358  
Cerebral Ventricle, 358  
Cerebral Ventricles, 358  
Cerebral Ventriculography, 358  
Cerebral Ventriculomegaly, 359  
Cerebral Visual Impairment, 359  
Cerebriform Nevus Sebaceous, 359  
Cerebritis, 359  
Cerebroplacental Ratio, 359  
Cerebrosides, 359  
Cerebrospinal Fluid, 359  
Cerebrospinal Fluid Analysis, 359  
Cerebrospinal Fluid Collection, 359  
Cerebrospinal Fluid Culture, 359  
Cerebrospinal Fluid Cytology, 359  
Cerebrospinal Fluid Leak, 359  
Cerebrotendinous Xanthomatosis (CTX), 359  
Cerebrovascular, 360  
Cerebrovascular Accident, 360  
Cerebrovascular Accident Prevention, 360  
Cerebrovascular Disease, 360  
Cerebrovascular Disorder, 360  
Cerebrovascular Occlusion, 360  
Cerebrum, 360  
Certification, 360  
Certified Nurse-Midwife, 360  
Certolizumab Pegol, 360  
Ceruloplasmin, 360  
Ceruloplasmin Blood Test, 360  
Cerumen, 360  
Cerumen Impaction, 360  
Ceruminous Adenocarcinoma, 361  
Cervical, 361  
Cervical Adenocarcinoma, 361  
Cervical Arthritis, 361  
Cervical Atlas, 361  
Cervical Biopsy, 361  
Cervical Cancer, 361  
Cervical Cancer Screening and Prevention, 361

---

---

Cervical Cancer Screening, Cytology, 361  
 Cervical Cap, 361  
 Cervical Carcinoma, 361  
 Cervical Cerclage, 361  
 Cervical Change, 361  
 Cervical Dilatation, 361  
 Cervical Dilation, 361  
 Cervical Disc Herniation, 362  
 Cervical Dislocation, 362  
 Cervical Dysplasia, 362  
 Cervical Dystocia, 362  
 Cervical Dystonia, 362  
 Cervical Ectropion, 362  
 Cervical Effacement, 362  
 Cervical Endometriosis, 362  
 Cervical Exam In Labor, 362  
 Cervical Ganglion, 362  
 Cervical Incompetence, 362  
 Cervical Insufficiency, 362  
 Cervical Intraepithelial Neoplasia, 362  
 Cervical Length Screening, 362  
 Cervical Lymphadenopathy, 362  
 Cervical MRI Scan, 363  
 Cervical Mucus, 363  
 Cervical Myelopathy, 363  
 Cervical Osteoarthritis, 363  
 Cervical Pain, 363  
 Cervical Plexus, 363  
 Cervical Polyp, 363  
 Cervical Polyps, 363  
 Cervical Pregnancy, 363  
 Cervical Radiculopathy, 363  
 Cervical Rib, 363  
 Cervical Ripening, 363  
 Cervical Ripening Agents, 363  
 Cervical Screening, 363  
 Cervical Spine, 363  
 Cervical Spine CT Scan, 364  
 Cervical Spine Involvement in Rheumatoid Arthritis, 364  
 Cervical Spine Pain, 364  
 Cervical Spine Trauma, 364  
 Cervical Spondyloarthropathy, 364  
 Cervical Spondylosis, 364  
 Cervical Spondylotic Myelopathy, 364  
 Cervical Squamous Cell Carcinoma, 364  
 Cervical Squamous Intraepithelial Lesion, 364  
 Cervical Stenosis, 364  
 Cervical Strain (Whiplash), 364  
 Cervical Vertebrae, 364  
 Cervicitis, 364  
 Cervicogenic Headache, 365  
 Cervix, 365  
 Cervix Carcinoma, 365  
 Cervix Cryosurgery, 365  
 Cervix Disease, 365  
 Cervix Neoplasm, 365  
 Cervix Uteri, 365  
 Cesarean Delivery, 365  
 Cesarean Section, 365  
 Cesium, 365  
 Cessation of Menstruation, 365  
 Cestoda, 365  
 Cestode, 365  
 Cestodiasis, 365  
 Cetirizine, 366  
 Cetrimide, 366  
 Cetorelix, 366  
 Cetuximab, 366  
 Cevimeline, 366  
 Cevimeline Hydrochloride, 366  
 CF, 366  
 CF (Cystic Fibrosis), 366  
 CFS, 366  
 CFS (Chronic Fatigue Syndrome), 366  
 CFTR Modulator Therapy, 366  
 CGD – Growth Delay, Constitutional, 450  
 CHA2DS2-VASc Score, 366  
 Chaddock Reflex, 366  
 Chaddock Sign, 367  
 Chadwick Sign, 367  
 Chafing, 367  
 Chagas Cardiomyopathy, 367  
 Chagas Disease, 367  
 Chagasic Megacolon, 367  
 Chagoma, 367  
 Chain Of Custody, 367  
 Chain-Termination Codon, 367  
 Chalazion, 367  
 Challenge Agent, 368  
 Challenge Testing, 368  
 Chamberlain Procedure, 368  
 Chamomile Extract, 368  
 Chanarin-Dorfman Disease, 368  
 Chance Fracture, 368  
 Chancre, 368  
 Chancroid, 368  
 Chandelier Sign, 368  
 Chandipura Virus Infection, 368  
 Change in Bowel Habits, 369  
 Channelopathy, 369  
 Chaplain, 369  
 Chapped Hands, 369  
 Chapped Lips, 369  
 Char Syndrome, 369  
 Charcoal, 369  
 Charcot Arthropathy, 369  
 Charcot Foot, 369  
 Charcot Joint, 369

---

---

Charcot Triad, [369](#)  
Charcot Triad of Cholangitis, [369](#)  
Charcot's Joints, [369](#)  
Charcot's Triad, [369](#)  
Charcot-Bouchard Aneurysm, [370](#)  
Charcot-Leyden Crystals, [370](#)  
Charcot-Marie-Tooth Disease, [370](#)  
Charcot-Wilbrand Syndrome, [370](#)  
Chargaff Rule, [370](#)  
CHARGE Syndrome, [370](#)  
Charles Bonnet Syndrome, [370](#)  
Charley Horse, [370](#)  
Chasing The Dragon, [370](#)  
Chaulmoogra Oil, [370](#)  
CHD, [370](#)  
Checkpoint Inhibitor, [370](#)  
Checkpoints, [370](#)  
Chediak-Higashi Syndrome, [371](#)  
Cheek, [371](#)  
Cheilectropion, [371](#)  
Cheilitis, [371](#)  
Cheilitis Glandularis, [371](#)  
Cheilitis Granulomatosa, [371](#)  
Cheilodysplasia, [371](#)  
Cheilognathopalatoschisis, [371](#)  
Cheilognathoschisis, [371](#)  
Cheiloplasty, [371](#)  
Cheiloschisis, [371](#)  
Cheilosis, [371](#)  
Cheiromegaly, [371](#)  
Cheilopompholyx, [371](#)  
Chelating Agent, [372](#)  
Chelating Agents, [372](#)  
Chelation, [372](#)  
Chelation Study, [372](#)  
Chelation Therapy, [372](#)  
Chemexfoliation, [372](#)  
Chemical, [372](#)  
Chemical Ablation, [372](#)  
Chemical Burn, [372](#)  
Chemical Burn Or Reaction, [372](#)  
Chemical Burns, [372](#)  
Chemical Carcinogenesis, [372](#)  
Chemical Cleavage, [372](#)  
Chemical Cofactor, [372](#)  
Chemical Eye Injury, [372](#)  
Chemical Fractionation, [373](#)  
Chemical Impurity, [373](#)  
Chemical Injury, [373](#)  
Chemical Peel, [373](#)  
Chemical Peels, [373](#)  
Chemical Pleurodesis, [373](#)  
Chemical Pneumonitis, [373](#)  
Chemistry, [373](#)  
Chemo, [373](#)  
Chemo Brain, [373](#)  
Chemoattractant, [373](#)  
Chemodectoma, [373](#)  
Chemoembolization, [373](#)  
Chemokine, [373](#)  
Chemokine Receptor, [374](#)  
Chemokinesis, [374](#)  
Chemoprevention, [374](#)  
Chemoradiation, [374](#)  
Chemoradiotherapy, [374](#)  
Chemoreceptor, [374](#)  
Chemoreceptor Cell, [374](#)  
Chemoreceptor Reflex, [374](#)  
Chemoreceptor Trigger Zone, [374](#)  
Chemoreceptors, [374](#)  
Chemoresistance, [374](#)  
Chemoresensitivity, [374](#)  
Chemoresensitization, [374](#)  
Chemosis, [374](#)  
Chemosurgery, [375](#)  
Chemotactic Factors, [375](#)  
Chemotaxis, [375](#)  
Chemotherapy, [375](#)  
Chemotherapy Extravasation, [375](#)  
Chemotherapy Regimen, [375](#)  
Chemotherapy Treatment, [375](#)  
Chemotherapy-Induced Nausea and Vomiting, [375](#)  
Chemotherapy-Induced Oral Mucositis, [375](#)  
Chemotherapy-Induced Peripheral Neuropathy, [375](#)  
Cherry Angioma, [375](#)  
Cherry Eye, [375](#)  
Cherry-Red Spot, [375](#)  
Cherubism, [376](#)  
Chest, [376](#)  
Chest Cold, [376](#)  
Chest CT, [376](#)  
Chest Infection, [376](#)  
Chest MRI, [376](#)  
Chest Pain, [376](#)  
Chest Physiotherapy, [376](#)  
Chest Tightness, [376](#)  
Chest Tube, [376](#)  
Chest Tube Insertion, [376](#)  
Chest Tube Thoracostomy, [376](#)  
Chest Wall, [376](#)  
Chest Wall Deformity, [376](#)  
Chest Wall Pain, [377](#)  
Chest X-Ray, [377](#)  
Chewing, [377](#)  
Chewing Tobacco, [377](#)  
Cheyne-Stokes Respiration, [377](#)  
CHF, [377](#)  
CHH, [377](#)

---

---

Chiari I Malformation, [377](#)  
Chiari Malformation, [377](#)  
Chiari-Frommel Syndrome, [377](#)  
Chicken Soup And Sickness, [377](#)  
Chickenpox, [377](#)  
Chickenpox (Varicella), [377](#)  
Chief Cell, [377](#)  
Chigger Bites, [377](#)  
Chiggers, [378](#)  
Chignon, [378](#)  
CHIK, [378](#)  
Chikungunya, [378](#)  
Chikungunya Fever, [378](#)  
Chikungunya Virus, [378](#)  
Chilaiditi Syndrome, [378](#)  
Chilblain, [378](#)  
Chilblains, [378](#)  
Child Abuse, [378](#)  
Child Adoption, [378](#)  
Child Behavior, [378](#)  
Child Development, [379](#)  
Child Health, [379](#)  
Child Health Services, [379](#)  
Child Maltreatment, [379](#)  
CHILD Naevus, [379](#)  
Child Neglect, [379](#)  
Child Neglect And Emotional Abuse, [379](#)  
Child Physical Abuse, [379](#)  
Child Poisoning, [379](#)  
Child Psychiatry, [379](#)  
Child Psychology, [379](#)  
CHILD Syndrome, [379](#)  
Child-Pugh Classification, [379](#)  
Child-Pugh Score, [379](#)  
Childbed Fever, [380](#)  
Childbirth, [380](#)  
Childhood, [380](#)  
Childhood Absence Epilepsy, [380](#)  
Childhood ADHD, [380](#)  
Childhood Angiosarcoma, [380](#)  
Childhood Apraxia Of Speech, [380](#)  
Childhood Asthma, [380](#)  
Childhood Cataracts, [380](#)  
Childhood Cyclic Vomiting, [380](#)  
Childhood Depression, [380](#)  
Childhood Disintegrative Disorder, [380](#)  
Childhood Ependymoma, [380](#)  
Childhood Fibrosarcoma, [381](#)  
Childhood Ganglioglioma, [381](#)  
Childhood Giant Axonal Neuropathy, [381](#)  
Childhood Interstitial Lung Disease, [381](#)  
Childhood Leiomyosarcoma, [381](#)  
Childhood Leukemia, [381](#)  
Childhood Liposarcoma, [381](#)  
Childhood Muscular Dystrophy, [381](#)  
Childhood Obesity, [381](#)  
Childhood Osteosarcoma, [381](#)  
Childhood Rhabdomyosarcoma, [381](#)  
Childhood Schizophrenia, [381](#)  
Children, [381](#)  
Children And Grief, [381](#)  
Children's Cancer Centers, [381](#)  
Chill, [382](#)  
Chills, [382](#)  
Chilomastix Mesnili, [382](#)  
Chimera, [382](#)  
Chimeric Antigen Receptor, [382](#)  
Chimeric Antigen Receptor T-Cell Therapy, [382](#)  
Chimerism, [382](#)  
Chin, [382](#)  
Chin Augmentation, [382](#)  
Chinese Herbal Medicine, [382](#)  
Chinese Restaurant Syndrome, [382](#)  
Chionablepsia, [382](#)  
CHIP, [382](#)  
Chipped, Broken, Or Cracked Tooth, [1937](#)  
Chiropody, [383](#)  
Chiropractic, [383](#)  
Chiropractic Care For Back Pain, [383](#)  
Chiropractor, [383](#)  
Chitin, [383](#)  
Chlamydia, [383](#)  
Chlamydia (Chlamydial Genitourinary Infections), [383](#)  
Chlamydia Pneumoniae Infection, [383](#)  
Chlamydia Psittaci, [383](#)  
Chlamydia Trachomatis, [383](#)  
Chlamydia Trachomatis Infection, [383](#)  
Chlamydiaceae, [383](#)  
Chlamydial Pneumonia, [383](#)  
Chlamydial Pneumonias, [384](#)  
Chlamydiosis, [384](#)  
Chloasma, [384](#)  
Chloracne, [384](#)  
Chloral Hydrate, [384](#)  
Chlorambucil, [384](#)  
Chloramphenicol, [384](#)  
Chloranemia, [384](#)  
Chlordane, [384](#)  
Chlordecone, [384](#)  
Chlordiazepoxide, [384](#)  
Chlordiazepoxide Overdose, [384](#)  
Chlorethoxyfos, [384](#)  
Chlorfenvinphos, [384](#)  
Chlorhexidine, [384](#)  
Chloride, [385](#)  
Chloride Channel, [385](#)  
Chloride In Diet, [385](#)  
Chloride Ion, [385](#)

---

---

Chloride Shift, [385](#)  
Chlorinated Lime, [385](#)  
Chlorinated Lime Poisoning, [385](#)  
Chlorine, [385](#)  
Chlorine Poisoning, [385](#)  
Chlormadinone Acetate, [385](#)  
Chlormethine, [385](#)  
Chlorodeoxyadenosine, [385](#)  
Chloroform, [385](#)  
Chlorogenic Acid, [385](#)  
Chloroleukemia, [385](#)  
Chloroma, [386](#)  
Chlorophyll Poisoning, [386](#)  
Chlorophyllin, [386](#)  
Chloropicrin, [386](#)  
Chloroprene, [386](#)  
Chloropsia, [386](#)  
Chloroquine, [386](#)  
Chlorothiazide, [386](#)  
Chlorpheniramine, [386](#)  
Chlorphentermine, [386](#)  
Chlorpromazine, [386](#)  
Chlorpromazine Overdose, [386](#)  
Chlorpropamide, [386](#)  
Chlorpyrifos, [386](#)  
Chlortalidone, [387](#)  
Chlorthalidone, [387](#)  
Chlorzoxazone, [387](#)  
CHO Cell, [387](#)  
Choanal Atresia, [387](#)  
Chocolate Cyst, [387](#)  
Choice Behavior, [387](#)  
Choke, [387](#)  
Choking, [387](#)  
Choking First Aid for Adults and Children, [387](#)  
Choking First Aid for an Unresponsive Person, [387](#)  
Choking First Aid for Infants, [387](#)  
Choking First Aid, Adult and Child, [387](#)  
Choking First Aid, Infant, [387](#)  
Choking Resuscitation, Unresponsive Patient, [387](#)  
Cholagogue, [387](#)  
Cholangitis, [388](#)  
Cholangiocarcinoma, [388](#)  
Cholangiography, [388](#)  
Cholangiohepatitis, [388](#)  
Cholangitis, [388](#)  
Cholangitis, Primary Sclerosing, [1595](#)  
Cholate, [388](#)  
Cholecalciferol, [388](#)  
Cholecyst, [388](#)  
Cholecystectomy, [388](#)  
Cholecystitis, [388](#)  
Cholecystography, [389](#)  
Cholecystokinin, [389](#)  
Cholecystoses, [389](#)  
Cholecystostomy, [389](#)  
Choledochal Cyst, [389](#)  
Choledochal Cysts, [389](#)  
Choledocholithiasis, [389](#)  
Cholelithiasis, [389](#)  
Cholera, [389](#)  
Choleresis, [389](#)  
Choleretic, [389](#)  
Cholestasis, [389](#)  
Cholestasis Of Pregnancy, [390](#)  
Cholesteatoma, [390](#)  
Cholesterol, [390](#)  
Cholesterol - Drug Treatment, [2116](#)  
Cholesterol And Lifestyle, [390](#)  
Cholesterol And Lipids, [390](#)  
Cholesterol Crystal Embolization, [390](#)  
Cholesterol Crystals, [390](#)  
Cholesterol Emboli, [390](#)  
Cholesterol Embolism, [390](#)  
Cholesterol Embolization, [390](#)  
Cholesterol Lowering, [390](#)  
Cholesterol Synthesis, [390](#)  
Cholesterol Testing And Results, [390](#)  
Cholesterolemia, [390](#)  
Cholesterolosis, [390](#)  
Cholestyramine, [391](#)  
Choline, [391](#)  
Cholinergic, [391](#)  
Cholinergic Crisis, [391](#)  
Cholinergic Neuron, [391](#)  
Cholinergic Toxidrome, [391](#)  
Cholinergic Urticaria, [391](#)  
Cholinesterase, [391](#)  
Cholinesterase Inhibitor, [2116](#)  
Chondritis, [391](#)  
Chondroblastic Osteosarcoma, [391](#)  
Chondroblastoma, [391](#)  
Chondrocalcinosis, [391](#)  
Chondrocyte, [391](#)  
Chondrodermatitis Nodularis Helicis, [391](#)  
Chondrodysplasia, [392](#)  
Chondroitin Sulfate, [392](#)  
Chondroma, [392](#)  
Chondromalacia, [392](#)  
Chondromalacia Patellae, [392](#)  
Chondromyxoid Fibroma, [392](#)  
Chondrosarcoma, [392](#)  
Chondrosarcoma Of The Breast, [392](#)  
Chopart's Joint, [392](#)  
Chorangioma, [392](#)  
Chordae Tendineae, [392](#)  
Chordee, [392](#)  
Chordoma, [392](#)

---



---

Chorea, 393  
 Chorea and Myoclonus, 393  
 Chorea, Huntington's, 952  
 Choreoathetosis, 393  
 Chorioamnionitis, 393  
 Choriocarcinoma, 393  
 Chorion, 393  
 Chorion Villus Sampling, 393  
 Chorionic Gonadotrophin, 393  
 Chorionic Villus Sampling, 393  
 Chorioretinitis, 393  
 Choroid, 393  
 Choroid Plexus, 393  
 Choroid Plexus Carcinoma, 393  
 Choroid Plexus Cyst, 394  
 Choroid Plexus Tumor, 394  
 Choroidal Dystrophies, 394  
 Choroidal Melanoma, 394  
 Choroidal Neovascularization, 394  
 Choroideremia, 394  
 Choroiditis, 394  
 Christmas Disease, 394  
 Chromaffin, 394  
 Chromaffinoma, 394  
 Chromate, 394  
 Chromatic, 394  
 Chromatids, 394  
 Chromatin, 394  
 Chromatography, 394  
 Chromatopsia, 395  
 Chromhidrosis, 395  
 Chromium, 395  
 Chromium In Diet, 395  
 Chromoblastomycosis, 395  
 Chromopertubation, 395  
 Chromosomal Abnormality, 395  
 Chromosomal Breakage Syndromes, 395  
 Chromosomal Microarray, 395  
 Chromosome, 395  
 Chromosome 1, 395  
 Chromosome 13, 395  
 Chromosome 2, 395  
 Chromosome 3, 396  
 Chromosome 4, 396  
 Chromosome 5, 396  
 Chromosome Complement, 396  
 Chromosome Duplication, 396  
 Chromosome X, 396  
 Chromosome Y, 396  
 Chronic, 396  
 Chronic Adrenal Insufficiency, 396  
 Chronic Allograft Nephropathy, 396  
 Chronic Atrophic Gastritis, 396  
 Chronic Autoimmune Hashimoto Thyroiditis, 396  
 Chronic Autoimmune Thyroiditis, 396  
 Chronic Bacterial Prostatitis, 396  
 Chronic Bronchitis, 397  
 Chronic Bronchitis With Cor Pulmonale, 397  
 Chronic Bronchitis With Emphysema, 397  
 Chronic Caries, 397  
 Chronic Cholecystitis, 397  
 Chronic Compartment Syndrome, 397  
 Chronic Coronary Syndrome, 397  
 Chronic Cough, 397  
 Chronic Daily Headaches, 397  
 Chronic Disease, 397  
 Chronic Ear Infection, 397  
 Chronic Endometritis, 397  
 Chronic Eosinophilic Leukaemia, 397  
 Chronic Eosinophilic Pneumonia, 398  
 Chronic Exertional Compartment Syndrome, 398  
 Chronic Fatigue Syndrome, 398  
 Chronic Fever, 398  
 Chronic Gastritis, 398  
 Chronic Glomerulonephritis, 398  
 Chronic Granulomatous Disease, 398  
 Chronic Hypertension, 398  
 Chronic Hypertension In Pregnancy, 398  
 Chronic Idiopathic Constipation, 398  
 Chronic Inflammation, 398  
 Chronic Inflammatory Demyelinating Polyneuropathy, 398  
 Chronic Inflammatory Response Syndrome, 399  
 Chronic Interstitial Nephritis, 399  
 Chronic Intestinal Pseudo-Obstruction, 399  
 Chronic Kidney Disease, 399  
 Chronic Kidney Disease Progression, 399  
 Chronic Kidney Disease Staging, 399  
 Chronic Kidney Disease-Mineral and Bone Disorder, 399  
 Chronic Kidney Failure, 399  
 Chronic Lead Poisoning, 399  
 Chronic Leukemia, 399  
 Chronic Limb-Threatening Ischemia, 399  
 Chronic Liver Disease, 400  
 Chronic Low Back Pain, 400  
 Chronic Lymphocytic Leukemia, 400  
 Chronic Lymphocytic Thyroiditis, 400  
 Chronic Megacolon, 400  
 Chronic Meningitis, 400  
 Chronic Mesenteric Ischemia, 400  
 Chronic Migraine, 400  
 Chronic Mitral Regurgitation, 400  
 Chronic Motor Or Vocal Tic Disorder, 400  
 Chronic Mucocutaneous Candidiasis, 400  
 Chronic Myelogenous Leukemia, 400  
 Chronic Myeloid Leukemia, 401  
 Chronic Myelomonocytic Leukemia, 401  
 Chronic Neutrophilic Leukaemia, 401

---

---

Chronic Obstructive Pulmonary Disease, [401](#)  
Chronic Otitis Media, [401](#)  
Chronic Pain, [401](#)  
Chronic Pain Syndrome, [401](#)  
Chronic Pancreatitis, [401](#)  
Chronic Paroxysmal Hemicrania, [401](#)  
Chronic Pelvic Pain, [401](#)  
Chronic Pelvic Pain in Women, [401](#)  
Chronic Pelvic Pain Syndrome, [402](#)  
Chronic Progressive External Ophthalmoplegia, [402](#)  
Chronic Pyelonephritis, [402](#)  
Chronic Recurrent Multifocal Osteomyelitis, [402](#)  
Chronic Renal Failure, [402](#)  
Chronic Respiratory Failure, [402](#)  
Chronic Rhinitis, [402](#)  
Chronic Rhinosinusitis, [402](#)  
Chronic Rhinosinusitis With Nasal Polyps, [402](#)  
Chronic Sinusitis, [402](#)  
Chronic Spontaneous Urticaria, [402](#)  
Chronic Stiff Neck, [403](#)  
Chronic Subdural Hematoma, [403](#)  
Chronic Suppurative Otitis Media, [403](#)  
Chronic Tension-Type Headache, [403](#)  
Chronic Thromboembolic Pulmonary Hypertension, [403](#)  
Chronic Toxicity, [403](#)  
Chronic Traumatic Encephalopathy, [403](#)  
Chronic Urticaria, [403](#)  
Chronic Venous Disease, [403](#)  
Chronic Venous Insufficiency, [403](#)  
Chronic Venous Stasis, [404](#)  
Chronic Vulvar Pain, [404](#)  
Chronicity, [404](#)  
Chronobiology, [404](#)  
Chronotropic, [404](#)  
Churg-Strauss Disease, [404](#)  
Churg-Strauss Syndrome, [404](#)  
Chvostek Sign, [404](#)  
Chvostek's Sign, [404](#)  
Chylangioma, [404](#)  
Chyle, [404](#)  
Chylomicron, [404](#)  
Chylomicronemia Syndrome, [404](#)  
Chylothorax, [404](#)  
Chylous Ascites, [404](#)  
Chyluria, [405](#)  
Chyme, [405](#)  
Chymotrypsin, [405](#)  
CIC, [405](#)  
Cicatricial Alopecia, [405](#)  
Cicatrizization, [405](#)  
Ciclopia, [405](#)  
CICO Emergency, [405](#)  
Cidofovir, [405](#)  
CIDP, [405](#)  
Cigarette, [405](#)  
Cigarette Smoking, [405](#)  
Ciguatera, [405](#)  
Ciguatera Poisoning, [405](#)  
Ciguatera Toxicity, [406](#)  
Ciguatera Toxin, [406](#)  
Ciguatoxin, [406](#)  
Cilia, [406](#)  
Ciliary Body, [406](#)  
Ciliary Body Melanoma, [406](#)  
Ciliary Dyskinesia, [406](#)  
Ciliary Flush, [406](#)  
Ciliary Muscle, [406](#)  
Ciliary Neuralgia, [406](#)  
Ciliopathy, [406](#)  
Cimetidine, [406](#)  
Cinacalcet, [406](#)  
Cineangiocardiology, [406](#)  
Cineangiography, [406](#)  
Cinefluorography, [407](#)  
Cineplasty, [407](#)  
Cingulotomy, [407](#)  
Cinnarizine, [407](#)  
Ciprofloxacin, [407](#)  
Circadian Rhythm, [407](#)  
Circadian Rhythm Sleep Disorder, [407](#)  
Circadian Rhythms, [407](#)  
Circle of Willis, [407](#)  
Circulating Tumor Cell, [407](#)  
Circulating Tumor DNA, [407](#)  
Circulation, [407](#)  
Circulatory, [407](#)  
Circulatory System, [407](#)  
Circumcision, [407](#)  
Circumduction, [408](#)  
Circumflex, [408](#)  
Circumflex Artery, [408](#)  
Circumvallate Papillae, [408](#)  
Circumvallate Placenta, [408](#)  
Cirrhosis, [408](#)  
Cirrhosis Of The Liver, [408](#)  
Cirrhosis, Primary Biliary, [1592](#)  
Cisplatin, [408](#)  
Cisterna Chyli, [408](#)  
Cisterna Magna, [408](#)  
Citalopram, [408](#)  
Citrate, [408](#)  
Citrate Synthase, [408](#)  
Citric Acid Cycle, [409](#)  
Citric Acid Urine Test, [409](#)  
Citrullinaemia, [409](#)  
Citrullinemia, [409](#)  
CJD, [409](#)  
CK, [409](#)

---

---

CKD-Mineral Bone Disorder, [409](#)  
Cladribine, [409](#)  
Clagett Procedure, [409](#)  
Clammy Skin, [409](#)  
Clamp, [409](#)  
Clang Association, [409](#)  
Clarithromycin, [409](#)  
Clark Level Of Invasion, [409](#)  
Class IA Antiarrhythmics, [409](#)  
Class IB Antiarrhythmics, [410](#)  
Class IC Antiarrhythmics, [410](#)  
Class III Antiarrhythmics, [410](#)  
Classic Galactosemia, [410](#)  
Classic Migraine, [410](#)  
Classic Type Pfeiffer Syndrome, [410](#)  
Clathrin, [410](#)  
Claude Syndrome, [410](#)  
Claudication, [410](#)  
Claudication Distance, [410](#)  
Claustrophobia, [410](#)  
Clavicle, [410](#)  
Clavicle Fracture, [410](#)  
Clavus, [411](#)  
Claw Foot, [411](#)  
Claw Hand, [411](#)  
Claw Toe, [411](#)  
Clay-Colored Stool, [411](#)  
Clean Catch Urine Sample, [411](#)  
Clean-Catch Specimen, [411](#)  
Cleaning Supplies And Equipment, [411](#)  
Clear Cell Carcinoma, [411](#)  
Clear Cell Carcinoma Of The Ovary, [411](#)  
Clear Cell Renal Cell Carcinoma, [411](#)  
Clear Cell Renal Cell Carcinoma Pathology, [411](#)  
Clear Liquid Diet, [411](#)  
Clear Margins, [411](#)  
Clearance, [412](#)  
Cleft Lip, [412](#)  
Cleft Lip And Cleft Palate, [412](#)  
Cleft Lip And Palate Repair, [412](#)  
Cleft Palate, [412](#)  
Cleidocranial Dysostosis, [412](#)  
Cleidocranial Dysplasia, [412](#)  
Clemastine, [412](#)  
Click-Murmur Syndrome, [412](#)  
Climacteric, [412](#)  
Climate Change, [412](#)  
Clindamycin, [412](#)  
Clinical, [412](#)  
Clinical Communication, [412](#)  
Clinical Counseling, [412](#)  
Clinical Depression, [413](#)  
Clinical Deterioration, [413](#)  
Clinical Diagnostics, [413](#)  
Clinical Disease, [413](#)  
Clinical Frailty Scale, [413](#)  
Clinical Handover, [413](#)  
Clinical Imaging, [413](#)  
Clinical Nutrition, [413](#)  
Clinical Pharmacology, [413](#)  
Clinical Presentation, [413](#)  
Clinical Psychologist, [413](#)  
Clinical Reasoning, [413](#)  
Clinical Trial, [413](#)  
Clinical Trial Phases, [413](#)  
Clinicopathologic Correlation, [414](#)  
Clinistix, [414](#)  
Clinitest, [414](#)  
Clinodactyly, [414](#)  
Clinoid Process, [414](#)  
CLIP, [414](#)  
Clitoral, [414](#)  
Clitorectomy, [414](#)  
Clitoridectomy, [414](#)  
Clitoris, [414](#)  
Clitorodynia, [414](#)  
Clitoromegaly, [414](#)  
Clivus, [414](#)  
CLL, [414](#)  
Cloacal Abnormalities, [414](#)  
Cloacal Exstrophy, [414](#)  
Cloacitis, [415](#)  
Clodronate, [415](#)  
Clofazimine, [415](#)  
Clomifene, [415](#)  
Clomipramine, [415](#)  
Clonal Evolution, [415](#)  
Clonal Expansion, [415](#)  
Clonazepam, [415](#)  
Clonic Seizure, [415](#)  
Clonidine, [415](#)  
Clonidine Toxicity, [415](#)  
Clonorchiasis, [415](#)  
Clonus, [415](#)  
Clopidogrel, [415](#)  
Closed Comedo, [416](#)  
Closed Fracture, [416](#)  
Closed Reduction, [416](#)  
Closed Reduction Of A Fractured Bone, [416](#)  
Closed Suction Drain With Bulb, [416](#)  
Closed Surgical Wound Care, [416](#)  
Closed-Angle Glaucoma, [416](#)  
Closed-Loop Insulin Delivery, [416](#)  
Clostridial Infection, [416](#)  
Clostridial Myonecrosis, [416](#)  
Clostridioides Difficile, [416](#)  
Clostridioides Difficile Infection, [416](#)  
Clostridioides Difficile Pathogenesis, [416](#)

---

---

Clostridium, [417](#)  
Clostridium Botulinum, [417](#)  
Clostridium Difficile, [417](#)  
Clostridium Difficile Colitis, [417](#)  
Clostridium Difficile Infection, [417](#)  
Clostridium Perfringens, [417](#)  
Clostridium Perfringens Food Poisoning, [417](#)  
Clostridium Tetani, [417](#)  
Clot, [417](#)  
Clot Buster, [417](#)  
Clot, Blood, [1918](#)  
Cloth Dye Poisoning, [417](#)  
Clotrimazole, [417](#)  
Clotting, [417](#)  
Cloudy Cornea, [418](#)  
Clove Oil, [418](#)  
Clozapine, [418](#)  
Club Drugs List And Side Effects, [418](#)  
Club Foot, [418](#)  
Clubbing, [418](#)  
Clubbing Of The Fingers Or Toes, [418](#)  
Clubbing of the Nails, [418](#)  
Clubfoot, [418](#)  
Clubfoot Repair, [418](#)  
Clubhand, [418](#)  
Clumping, [418](#)  
Cluster, [418](#)  
Cluster Headache, [418](#)  
Cluster Headaches, [418](#)  
Clutton's Joints, [419](#)  
CMTC, [419](#)  
CMV, [419](#)  
CMV Blood Test, [419](#)  
CMV Colitis, [419](#)  
CMV Pneumonia, [419](#)  
CMV Retinitis, [419](#)  
CNS Infection, [419](#)  
CNS Lymphoma, [419](#)  
CO Poisoning, [419](#)  
CO2 Blood Test, [419](#)  
Coagulate, [419](#)  
Coagulation, [419](#)  
Coagulation Cascade, [420](#)  
Coagulation Defects, [420](#)  
Coagulation Factor, [420](#)  
Coagulation Factor Deficiencies, [420](#)  
Coagulation Factor V Leiden, [420](#)  
Coagulation Profile, [420](#)  
Coagulation Time, [420](#)  
Coagulative Necrosis, [420](#)  
Coagulopathy, [420](#)  
Coal Worker's Pneumoconiosis, [420](#)  
Coal Workers' Pneumoconiosis (Black Lung Disease), [420](#)  
Coarctation, [420](#)  
Coarctation Of The Aorta, [420](#)  
Coarse Crackles, [421](#)  
Coats Disease, [421](#)  
Cobalamin, [421](#)  
Cobalamin Deficiency, [421](#)  
Cobalt, [421](#)  
Cobalt Poisoning, [421](#)  
Cobalt-57, [421](#)  
Cobalt-Chromium, [421](#)  
Cocaine, [421](#)  
Cocaine Cardiotoxicity, [421](#)  
Cocaine Intoxication, [421](#)  
Cocaine Substance Use, [421](#)  
Cocaine Toxicity, [421](#)  
Cocaine Use Disorder, [421](#)  
Cocaine Withdrawal, [421](#)  
Cocaine-Associated Cardiomyopathy, [422](#)  
Coccarboxylase, [422](#)  
Coccidioides Complement Fixation, [422](#)  
Coccidioides Immitis, [422](#)  
Coccidioides Precipitin Test, [422](#)  
Coccidioidomycosis, [422](#)  
Coccidioidomycosis and Valley Fever, [422](#)  
Coccidiosis, [422](#)  
Coccus, [422](#)  
Coccydynia, [422](#)  
Coccygectomy, [422](#)  
Coccyx, [422](#)  
Cochlea, [422](#)  
Cochlear Duct, [423](#)  
Cochlear Implant, [423](#)  
Cochlear Implants, [423](#)  
Cockayne Syndrome, [423](#)  
Cockcroft-Gault Formula, [423](#)  
Code Black, [423](#)  
Code Blue, [423](#)  
Codeine Overdose, [423](#)  
Codman Triangle, [423](#)  
Codon, [423](#)  
Coefficient of Inbreeding, [423](#)  
Coenzyme, [423](#)  
Coenzyme Q, [423](#)  
Coenzyme Q10, [424](#)  
Cofactor, [424](#)  
Coffee-Ground Emesis, [424](#)  
Coffin-Lowry Syndrome, [424](#)  
Cogan's Syndrome, [424](#)  
Cognition, [424](#)  
Cognitive, [424](#)  
Cognitive Behavior Therapy, [424](#)  
Cognitive Behavioral Therapy, [424](#)  
Cognitive Behavioral Therapy For Back Pain, [424](#)  
Cognitive Behavioral Therapy For Insomnia, [424](#)

---

---

Cognitive Behavioural Therapy, [424](#)  
Cognitive Decline, [424](#)  
Cognitive Distortions, [424](#)  
Cognitive Fitness, [425](#)  
Cognitive Function, [425](#)  
Cognitive Health And Memory, [425](#)  
Cognitive Impairment, [425](#)  
Cognitive Rehabilitation, [425](#)  
Cognitive Reserve, [425](#)  
Cognitive Screening, [425](#)  
Cognitive Symptoms, [425](#)  
Cognitive Therapy, [425](#)  
Cogwheel Rigidity, [425](#)  
Cohort, [425](#)  
Cohort Study, [425](#)  
Coincidence Detection, [425](#)  
Coitus, [425](#)  
Coitus Interruptus, [426](#)  
Coke, [426](#)  
Cold, [426](#)  
Cold Abscess, [426](#)  
Cold Agglutinin, [426](#)  
Cold Agglutinin Disease, [426](#)  
Cold Agglutinin Syndrome, [426](#)  
Cold Agglutinins, [426](#)  
Cold Allergy, [426](#)  
Cold Exposure, [426](#)  
Cold Hands, [426](#)  
Cold Injury, [426](#)  
Cold Intolerance, [426](#)  
Cold Ischemia Time, [426](#)  
Cold Medicines And Children, [426](#)  
Cold Panniculitis, [426](#)  
Cold Sore, [427](#)  
Cold Sores, [427](#)  
Cold Urticaria, [427](#)  
Cold, Common, [434](#)  
Cold, Flu, Allergy, [499](#)  
Cold/Cough Medicines in Children, [2116](#)  
Colectomy, [427](#)  
Colibacillosis, [427](#)  
Colic, [427](#)  
Colicky Pain, [427](#)  
Coliform, [427](#)  
Colitis, [427](#)  
Colitis, Collagenous, [1256](#)  
Colitis, Ischemic, [1060](#)  
Colitis, Lymphocytic, [1256](#)  
Colitis, Microscopic, [1256](#)  
Colitis, Pseudomembranous, [1620](#)  
Colitis, Ulcerative, [1984](#)  
Collagen, [427](#)  
Collagen Disease, [427](#)  
Collagen Implant, [427](#)  
Collagenase, [427](#)  
Collagenous Colitis, [428](#)  
Collapse Of The Lung, [428](#)  
Collapsed Arch, [428](#)  
Collapsed Lung, [428](#)  
Collapsing Glomerulopathy, [428](#)  
Collapsing Pulse, [428](#)  
Collateral, [428](#)  
Collateral Circulation, [428](#)  
Collateral Ligament Injury, [428](#)  
Collecting Duct, [428](#)  
Colles Fracture, [428](#)  
Collimator, [428](#)  
Colloids, [429](#)  
Coloboma, [429](#)  
Coloboma Of The Iris, [429](#)  
Colocolonic Intussusception, [429](#)  
Cologne Poisoning, [429](#)  
Cologuard, [429](#)  
Colon, [429](#)  
Colon Cancer, [429](#)  
Colon Cancer Prevention, [429](#)  
Colon Cancer Screening, [429](#)  
Colon Polyps, [429](#)  
Colonic Diverticulitis, [429](#)  
Colonic Inertia, [430](#)  
Colonic Ischemia, [430](#)  
Colonic Mucosa, [430](#)  
Colonic Polyps, [430](#)  
Colonic Pseudo-Obstruction, [430](#)  
Colonic Stenosis, [430](#)  
Colonic Tissue Culture, [430](#)  
Colonization, [430](#)  
Colonoscope, [430](#)  
Colonoscopy, [430](#)  
Colony-Forming Unit, [430](#)  
Colony-Stimulating Factors, [430](#)  
Color Blindness, [430](#)  
Color Doppler, [430](#)  
Color Vision Deficiency, [431](#)  
Color Vision Test, [431](#)  
Colorado Tick Fever, [431](#)  
Colorectal Adenocarcinoma, [431](#)  
Colorectal Adenoma, [431](#)  
Colorectal Cancer, [431](#)  
Colorectal Cancer Genetics, [431](#)  
Colorectal Cancer Screening, [431](#)  
Colorectal Cancer Screening Tests, [431](#)  
Colorectal Polyp, [431](#)  
Colorectal Polyps, [431](#)  
Colorectal Surgery, [431](#)  
Colostomy, [432](#)  
Colostrum, [432](#)  
Colpitis, [432](#)

---

---

Colporrhaphy, [432](#)  
Colposcope, [432](#)  
Colposcopy, [432](#)  
Colposcopy-Directed Biopsy, [432](#)  
Columnar Epithelium Lined Lower Esophagus, [432](#)  
Coma, [432](#)  
Coma Scale, [432](#)  
Coma, Myxedema, [1320](#)  
Combination Inhalers, [432](#)  
Combination Therapy, [432](#)  
Combined Electrolyte Imbalances, [432](#)  
Combined Hormone Therapy, [433](#)  
Combined Hyperlipidemia, [433](#)  
Combined Oral Contraceptive Pill, [433](#)  
Combined Spinal-Epidural Analgesia, [433](#)  
Comedocarcinoma, [433](#)  
Comedonal Acne, [433](#)  
Comedones, [433](#)  
Comfort Measures Only, [433](#)  
Commando Procedure, [433](#)  
Commensal, [433](#)  
Comminuted, [433](#)  
Comminuted Fracture, [433](#)  
Commissurotomy, [433](#)  
Commode, [434](#)  
Common Bile Duct, [434](#)  
Common Carotid Artery, [434](#)  
Common Cold, [434](#)  
Common Cold In Babies, [434](#)  
Common Fibular Nerve, [434](#)  
Common Iliac Arteries, [434](#)  
Common Iliac Vein, [434](#)  
Common Medical Abbreviations And Terms, [434](#)  
Common Migraine, [434](#)  
Common Peroneal Nerve Dysfunction, [434](#)  
Common Peroneal Neuropathy, [435](#)  
Common Uterine and Ovarian Growths, [435](#)  
Common Variable Immunodeficiency, [435](#)  
Commotio Cordis, [435](#)  
Communicable Disease, [435](#)  
Communicating Hydrocephalus, [435](#)  
Communicating With Patients, [435](#)  
Communicating With Someone With Aphasia, [435](#)  
Communicating With Someone With Dysarthria, [435](#)  
Community Acquired Infection, [435](#)  
Community Health Worker, [435](#)  
Community Mental-Health Treatment, [435](#)  
Community Rehabilitation, [436](#)  
Community Reintegration, [436](#)  
Community-Acquired, [436](#)  
Community-Acquired Pneumonia, [436](#)  
Community-Acquired Pneumonia In Adults, [436](#)  
Comorbidity, [436](#)  
Compact Bone, [436](#)  
Comparison, [436](#)  
Compartment Pressure, [436](#)  
Compartment Syndrome, [436](#)  
Compartment Syndrome Cross-Reference, [436](#)  
Compensated Heart Failure, [436](#)  
Competence, [436](#)  
Competitive Antagonist, [436](#)  
Competitive Inhibition, [437](#)  
Complement, [437](#)  
Complement C1Q Deficiency, [437](#)  
Complement C2/C4 Deficiency, [437](#)  
Complement C3, [437](#)  
Complement Cascade, [437](#)  
Complement Component 4, [437](#)  
Complement Component C3, [437](#)  
Complement Component Deficiency, [437](#)  
Complement Deficiencies, [437](#)  
Complement Deficiency, [437](#)  
Complement Fixation Test For C Burnetii, [437](#)  
Complement System, [437](#)  
Complementary Sequence, [438](#)  
Complete Blood Count, [438](#)  
Complete Breech, [438](#)  
Complete Cystectomy, [438](#)  
Complete Denture, [438](#)  
Complete Miscarriage, [438](#)  
Complete Response, [438](#)  
Complex Carbohydrates, [438](#)  
Complex Febrile Seizure, [438](#)  
Complex Regional Pain Syndrome, [438](#)  
Compliance, [439](#)  
Complicated Bereavement, [439](#)  
Complicated Grief, [439](#)  
Complicated Migraine, [439](#)  
Complicated Urinary Tract Infection, [439](#)  
Complication, [439](#)  
Composite Resin, [439](#)  
Compound Heterozygote, [439](#)  
Compound Muscle Action Potential, [439](#)  
Compound Presentation, [439](#)  
Compound Presentations, [439](#)  
Compounding, [439](#)  
Compounding Pharmacy, [439](#)  
Comprehensive Geriatric Assessment, [439](#)  
Comprehensive Metabolic Panel, [440](#)  
Compress, [440](#)  
Compressed Nerve, [440](#)  
Compression, [440](#)  
Compression Bandage, [440](#)  
Compression Fracture, [440](#)  
Compression Fractures Of The Back, [440](#)  
Compression Stockings, [440](#)  
Compression Therapy, [440](#)  
Compression Ultrasound, [440](#)

---



---

Compressive Atelectasis, [440](#)  
Compton Scattering, [440](#)  
Compulsions, [440](#)  
Compulsive Gambling, [440](#)  
Compulsive Obsessive Disorder, [441](#)  
Compulsive Overeating, [441](#)  
Compulsive Sexual Behavior, [441](#)  
Compulsive Sexual Behavior Disorder, [441](#)  
Compulsive Stealing, [441](#)  
Computed Tomography, [441](#)  
Computed Tomography Angiography, Abdomen and Pelvis, [484](#)  
Computed Tomography Angiography, Extremities, [484](#)  
Computed Tomography Angiography, Head and Neck, [484](#)  
Computed Tomography Pulmonary Angiography, [441](#)  
Computed Tomography Scan, [441](#)  
Computed Tomography Urography, [441](#)  
Computerized Axial Tomography, [441](#)  
Concato's Disease, [441](#)  
Concealed Conduction, [441](#)  
Concealed Pregnancy, [441](#)  
Concentration, [441](#)  
Concentric Muscle Action, [442](#)  
Conception, [442](#)  
Conchae, [442](#)  
Conclusion, [442](#)  
Concomitant, [442](#)  
Concordance, [442](#)  
Concrescence, [442](#)  
Concretio Cordis, [442](#)  
Concussion, [442](#)  
Conditioning, [442](#)  
Conditions Caused By Deletion Mutations, [442](#)  
Condom, [442](#)  
Conduct Disorder, [442](#)  
Conduction Aphasia, [443](#)  
Conduction Deafness, [443](#)  
Conductive Deafness, [443](#)  
Conductive Hearing Loss, [443](#)  
Condylar Process, [443](#)  
Condyle, [443](#)  
Condylloid Joint, [443](#)  
Condyloma, [443](#)  
Condyloma Acuminata, [443](#)  
Condyloma Acuminatum, [443](#)  
Condyloma Lata, [443](#)  
Condyloma Latum, [443](#)  
Condylomata Acuminata, [443](#)  
Condylomata Lata, [443](#)  
Cone Biopsy, [444](#)  
Cone-Beam Computed Tomography, [444](#)  
Cones, [444](#)  
Confabulation, [444](#)  
Confidentiality, [444](#)  
Confirmatory Test, [444](#)  
Confluent and Reticulated Papillomatosis, [444](#)  
Conformal Radiation Therapy, [444](#)  
Confounding, [444](#)  
Confusion, [444](#)  
Confusion Assessment Method, [444](#)  
Congenital, [444](#)  
Congenital Adrenal Hyperplasia, [444](#)  
Congenital Antithrombin III Deficiency, [445](#)  
Congenital Biliary Atresia, [445](#)  
Congenital Birth Defect, [445](#)  
Congenital Cataract, [445](#)  
Congenital CMV, [445](#)  
Congenital CNS Malformation, [445](#)  
Congenital Contractural Arachnodactyly, [445](#)  
Congenital Cytomegalovirus, [445](#)  
Congenital Cytomegalovirus Infection, [445](#)  
Congenital Diaphragmatic Hernia, [445](#)  
Congenital Diaphragmatic Hernia Repair, [445](#)  
Congenital Disorder, [445](#)  
Congenital Fibrinogen Deficiency, [445](#)  
Congenital Glaucoma, [445](#)  
Congenital Heart Defect, [445](#)  
Congenital Heart Defect Corrective Surgery, [446](#)  
Congenital Heart Defects, [446](#)  
Congenital Heart Defects In Children, [446](#)  
Congenital Heart Disease, [446](#)  
Congenital Heart Disease In Adults, [446](#)  
Congenital Heart Murmur, [446](#)  
Congenital Hemolytic Jaundice, [446](#)  
Congenital Hip Dislocation, [446](#)  
Congenital Hydrocephalus, [446](#)  
Congenital Hyperinsulinism, [446](#)  
Congenital Hyperthyroidism, [446](#)  
Congenital Hypertrichosis Lanuginosa, [446](#)  
Congenital Hypothyroidism, [446](#)  
Congenital Malaria, [446](#)  
Congenital Malformation, [447](#)  
Congenital Megacolon, [447](#)  
Congenital Melanocytic Nevus, [447](#)  
Congenital Mitral Valve Anomalies, [447](#)  
Congenital Muscular Dystrophy, [447](#)  
Congenital Myasthenic Syndromes, [447](#)  
Congenital Myopathies, [447](#)  
Congenital Nasolacrimal Duct Obstruction, [447](#)  
Congenital Nephrotic Syndrome, [447](#)  
Congenital Nevi, [447](#)  
Congenital Nevus, [447](#)  
Congenital Penile Curvature, [447](#)  
Congenital Plagiocephaly, [448](#)  
Congenital Platelet Function Defects, [448](#)  
Congenital Pneumonia, [448](#)  
Congenital Protein C Or S Deficiency, [448](#)

---

---

Congenital Rubella, 448  
Congenital Rubella Syndrome, 448  
Congenital Scoliosis, 448  
Congenital Syphilis, 448  
Congenital Toxoplasmosis, 448  
Congenital Tracheomalacia, 448  
Congenital Vaginal Obstruction, 448  
Congenital Varicella Syndrome, 448  
Congenital Zika Syndrome, 449  
Congestion, 449  
Congestive Heart Failure, 449  
Congestive Hepatopathy, 449  
Conization, 449  
Conjoined Twins, 449  
Conjugate Vaccine, 2116  
Conjugated Equine Estrogens, 449  
Conjugated Hyperbilirubinemia, 449  
Conjugated Linoleic Acid, 449  
Conjunctiva, 449  
Conjunctival, 449  
Conjunctival Melanoma, 449  
Conjunctival Squamous-Cell Carcinoma, 449  
Conjunctivitis, 449  
Conjunctivitis, Acute, 450  
Conjunctivoplasty, 450  
Conn Syndrome, 450  
Conn's Syndrome, 450  
Connective Tissue, 450  
Connective Tissue Disease, 450  
Connexin 26 Deficiency, 450  
Consanguinity, 450  
Conscious Sedation, 2116  
Conscious Sedation For Surgical Procedures, 450  
Consolidation, 450  
Consolidation Therapy, 450  
Constipation, 450  
Constitutional Delay In Puberty, 450  
Constitutional Growth Delay, 450  
Constraint-Induced Movement Therapy, 451  
Constrictive Pericarditis, 451  
Consultation-Liaison Psychiatry, 451  
Contac Overdose, 451  
Contact Allergen Testing, 451  
Contact Dermatitis, 451  
Contact Lenses, 451  
Contact Precautions, 451  
Contact Tracing, 451  
Contact Urticaria, 451  
Contagious Disease, 451  
Contamination, 451  
Continuing Medical Education, 451  
Continuity Of Care, 451  
Continuous Bladder Irrigation, 452  
Continuous Combined Hormone Therapy, 452  
Continuous Electronic Fetal Monitoring, 452  
Continuous Glucose Monitoring, 452  
Continuous Kidney Replacement Therapy, 452  
Continuous Murmurs, 452  
Continuous Passive Motion Machine, 452  
Continuous Positive Airway Pressure, 452  
Continuous Renal Replacement Therapy, 452  
Continuous Subcutaneous Insulin Infusion, 452  
Contraception, 452  
Contraception And Birth Control, 453  
Contraception For Adolescents, 453  
Contraceptive Implant, 453  
Contraceptive, Pill, 1412  
Contraceptives, 453  
Contracted Pelvis, 453  
Contractile Proteins, 453  
Contraction, 453  
Contraction Stress Test, 453  
Contractions, 453  
Contractions, Braxton Hicks, 258  
Contracture, 453  
Contracture Deformity, 453  
Contracture Prevention, 453  
Contraindication, 453  
Contralateral, 454  
Contrast, 454  
Contrast Agent, 454  
Contrast Echocardiography, 454  
Contrast Medium, 454  
Contrast Radiology, 454  
Contrast Reaction, 454  
Contrast-Associated Acute Kidney Injury, 454  
Contrast-Enhanced Computed Tomography, 454  
Contrast-Enhanced Magnetic Resonance Imaging, 454  
Contrast-Enhanced Ultrasound, 454  
Contrast-Induced Nephropathy, 454  
Contrecoup, 454  
Control Group, 454  
Controlled Cord Traction, 455  
Controlled Ovarian Hyperstimulation, 455  
Controlled Substance, 2116  
Controlled-Substance Schedule I, 455  
Controlled-Substance Schedule II, 455  
Controlled-Substance Schedule III, 455  
Controlled-Substance Schedule IV, 455  
Controlled-Substance Schedule V, 455  
Controllers, 455  
Contusion, 455  
Conus Medullaris Syndrome, 455  
Convergence, 455  
Convergence Insufficiency, 455  
Conversion Disorder, 455  
Convulsion, 455  
Cooking Without Salt, 456

---

---

Cooley's Anemia, [456](#)  
Coombs Test, [456](#)  
Cooper's Ligament, [456](#)  
Coordination, [456](#)  
COPD, [456](#)  
COPD And Other Health Problems, [456](#)  
COPD Exacerbation, [456](#)  
COPD Flare-Ups, [456](#)  
COPD Maintenance Medications, [456](#)  
COPD Rehabilitation, [456](#)  
COPD Rescue Medications, [456](#)  
Copper, [456](#)  
Copper Deficiency, [457](#)  
Copper In Diet, [457](#)  
Copper Poisoning, [457](#)  
Coprolalia, [457](#)  
Copy Number Variant, [457](#)  
Copy Number Variation, [457](#)  
CoQ10, [457](#)  
Cor, [457](#)  
Cor Pulmonale, [457](#)  
Cor Triatriatum, [457](#)  
Coracoid Process, [457](#)  
Cord, [457](#)  
Cord Blood Stem Cells, [457](#)  
Cord Blood Testing, [457](#)  
Cord Presentation, [458](#)  
Cordocentesis, [458](#)  
Cordotomy, [458](#)  
Core Biopsy, [458](#)  
Core Needle Biopsy, [458](#)  
Cori Cycle, [458](#)  
Cori Disease, [458](#)  
Corn, [458](#)  
Corn-Cockle Poisoning, [458](#)  
Cornea, [458](#)  
Corneal Abrasion, [458](#)  
Corneal Cross-Linking, [458](#)  
Corneal Disease, [458](#)  
Corneal Dystrophy, [459](#)  
Corneal Edema, [459](#)  
Corneal Endothelium, [459](#)  
Corneal Epithelium, [459](#)  
Corneal Implant, [459](#)  
Corneal Inflammation, [459](#)  
Corneal Injury, [459](#)  
Corneal Laceration, [459](#)  
Corneal Reflex, [459](#)  
Corneal Stroma, [459](#)  
Corneal Transplant, [459](#)  
Corneal Transplantation, [459](#)  
Corneal Ulcer, [459](#)  
Corneal Ulcers And Infections, [459](#)  
Corneitis, [460](#)  
Cornelia De Lange Syndrome, [460](#)  
Corns, [460](#)  
Corns And Calluses, [460](#)  
Cornual Pregnancy, [460](#)  
Coronal, [460](#)  
Coronal Caries, [460](#)  
Coronal Plane, [460](#)  
Coronary, [460](#)  
Coronary Aneurysm, [460](#)  
Coronary Angiography, [460](#)  
Coronary Angioplasty, [460](#)  
Coronary Arteries, [460](#)  
Coronary Artery, [461](#)  
Coronary Artery Aneurysm, [461](#)  
Coronary Artery Anomaly, [461](#)  
Coronary Artery Bypass, [461](#)  
Coronary Artery Bypass Grafting, [461](#)  
Coronary Artery Bypass Surgery, [461](#)  
Coronary Artery Calcium Score, [461](#)  
Coronary Artery Disease, [461](#)  
Coronary Artery Fistula, [461](#)  
Coronary Artery Spasm, [461](#)  
Coronary Calcium Score, [461](#)  
Coronary Calcium Scoring, [461](#)  
Coronary Care Unit, [461](#)  
Coronary Circulation, [462](#)  
Coronary Computed Tomography Angiography, [462](#)  
Coronary Dominance, [462](#)  
Coronary Endarterectomy, [462](#)  
Coronary Insufficiency, [462](#)  
Coronary Microvascular Angina, [462](#)  
Coronary Microvascular Disease, [462](#)  
Coronary Microvascular Dysfunction, [462](#)  
Coronary Occlusion, [462](#)  
Coronary Perfusion Pressure, [462](#)  
Coronary Revascularization, [462](#)  
Coronary Sinus, [462](#)  
Coronary Slow Flow Phenomenon, [462](#)  
Coronary Spasm, [463](#)  
Coronary Steal Phenomenon, [463](#)  
Coronary Steal Syndrome, [463](#)  
Coronary Stent, [463](#)  
Coronary Thrombosis, [463](#)  
Coronavirus, [463](#)  
Coronavirus (COVID-19) and Asthma, [463](#)  
Coronavirus (COVID-19) and High Blood Pressure, [463](#)  
Coronavirus (COVID-19) and Pregnancy, [463](#)  
Coronavirus (COVID-19) in Children, Infants, [469](#)  
Coronavirus (COVID-19), Complications, [469](#)  
Coronavirus (COVID-19), Recovery, [469](#)  
Coronavirus (COVID-19), Symptoms, [469](#)  
Coronavirus (COVID-19), Testing, [469](#)  
Coronavirus (COVID-19), Testing at Home, [139](#)  
Coronavirus Disease 2019, [463](#)

---

---

Coronavirus Disease COVID-19, [463](#)  
 Coronavirus, Types, [463](#)  
 Coronaviruses, [463](#)  
 Corporeal, [463](#)  
 Corpus, [464](#)  
 Corpus Albicans, [464](#)  
 Corpus Callosum, [464](#)  
 Corpus Cavernosum, [464](#)  
 Corpus Luteum, [464](#)  
 Corpus Luteum Cyst, [464](#)  
 Corpus Spongiosum, [464](#)  
 Corrected Age, [464](#)  
 Corrigan Pulse, [464](#)  
 Corrosive, [464](#)  
 Corrosive Ingestion, [464](#)  
 Corrugator Muscle, [464](#)  
 Cortex, [464](#)  
 Corti's Ganglion, [464](#)  
 Cortical, [465](#)  
 Cortical Bone, [465](#)  
 Cortical Cataract, [465](#)  
 Corticobasal Degeneration, [465](#)  
 Corticospinal Tract, [465](#)  
 Corticosteroid, [465](#)  
 Corticosteroids, [465](#)  
 Corticosteroids Mechanism, [465](#)  
 Corticosteroids Overdose, [465](#)  
 Corticotropin-Releasing Factor, [465](#)  
 Cortisol, [465](#)  
 Cortisol Blood Test, [465](#)  
 Cortisol Urine Test, [465](#)  
 Cortisone, [465](#)  
 Coryza, [466](#)  
 Cosmeceutical, [466](#)  
 Cosmetic Allergies, [466](#)  
 Cosmetic Dermatology, [466](#)  
 Cosmetic Ear Surgery, [466](#)  
 Cosmetic Surgery, [466](#)  
 Costal Cartilage, [466](#)  
 Costal Groove, [466](#)  
 Costal Margin, [466](#)  
 Costello Syndrome, [466](#)  
 Costochondritis, [466](#)  
 Costochondritis And Tietze Syndrome, [466](#)  
 Costosternal Chondrodynia, [467](#)  
 Costosternal Syndrome, [467](#)  
 Costotransverse Joint, [467](#)  
 Costovertebral Joint, [467](#)  
 Costovertebral-Angle Tenderness, [467](#)  
 Cot Death, [467](#)  
 Cotard Syndrome, [467](#)  
 Cotton-Wool Spots, [467](#)  
 Cough, [467](#)  
 Cough Augmentation, [467](#)  
 Cough Headaches, [467](#)  
 Cough In Adults, [467](#)  
 Cough Suppressant, [467](#)  
 Cough Syncope, [467](#)  
 Cough, Chronic, [397](#)  
 Coughing Up Blood, [467](#)  
 Councilman Body, [468](#)  
 Count Rate, [468](#)  
 Countercurrent Multiplication, [468](#)  
 Counterfeit Drug, [468](#)  
 Counterpulsation, [468](#)  
 Countertransference, [468](#)  
 Counting Carbohydrates, [468](#)  
 Coup-Contrecoup Injury, [468](#)  
 Courvoisier Sign, [468](#)  
 Cover Test, [468](#)  
 Covered Vascular Stent, [468](#)  
 COVID Toe, [468](#)  
 COVID-19, [468](#)  
 COVID-19 Antibody Test, [468](#)  
 COVID-19 Complications, [469](#)  
 COVID-19 in Infants and Children, [469](#)  
 COVID-19 In Pregnancy, [469](#)  
 COVID-19 Recovery, [469](#)  
 COVID-19 Symptoms, [469](#)  
 COVID-19 Test, [469](#)  
 COVID-19 Testing, [469](#)  
 COVID-19 Vaccines, [469](#)  
 COVID-19 Virus Test, [469](#)  
 COVID-19, Antibody Testing, [468](#)  
 COVID-19-Associated Anosmia and Ageusia, [469](#)  
 Cow's Milk Allergy, [469](#)  
 Cow's Milk And Children, [469](#)  
 Cow's Milk Introduction in Infancy, [469](#)  
 Cowden Disease (Multiple Hamartoma Syndrome), [470](#)  
 Cowden Syndrome, [470](#)  
 Cowper's Gland, [470](#)  
 Cowpox, [470](#)  
 COX Inhibitor, [2116](#)  
 Cox-1 Inhibitor, [2116](#)  
 Cox-2, [470](#)  
 COX-2 Inhibitors, [470](#)  
 Coxa Vara, [470](#)  
 Coxalgia, [470](#)  
 Coxiella Burnetii, [470](#)  
 Cocksackie, [470](#)  
 Cocksackie Virus Type A16, [470](#)  
 Cocksackievirus, [470](#)  
 Cozen Test, [470](#)  
 CP, [470](#)  
 CPAP, [471](#)  
 CPAP Machine, [471](#)  
 CpG, [471](#)  
 CPK Isoenzymes Test, [471](#)

---

---

CPR, [471](#)  
CPVT, [471](#)  
Crabs, [471](#)  
Crabs (Pubic Lice), [471](#)  
Crack (Drug), [2116](#)  
Cracked Heels, [471](#)  
Cracked Tooth Syndrome, [471](#)  
Crackles, [471](#)  
Cradle Cap, [471](#)  
Cramp, [471](#)  
Cramp, Muscle, [1305](#)  
Cramping Leg Pain, [471](#)  
Cramps But No Period, [472](#)  
Cramps, Heat, [895](#)  
Cramps, Menstrual, [1232](#)  
Cranberries, [472](#)  
Cranial, [472](#)  
Cranial Arteritis, [472](#)  
Cranial Cavity, [472](#)  
Cranial Fasciitis, [472](#)  
Cranial Implant, [472](#)  
Cranial Mononeuropathy III, [472](#)  
Cranial Nerve, [472](#)  
Cranial Nerve Palsy, [472](#)  
Cranial Nerve X, [472](#)  
Cranial Nerves, [472](#)  
Cranial Sutures, [472](#)  
Craniectomy, [472](#)  
Craniocerebral Injury, [473](#)  
Craniofacial, [473](#)  
Craniofacial Abnormalities, [473](#)  
Craniofacial Disorder, [473](#)  
Craniofacial Dysostosis, [473](#)  
Craniofacial Microsomia, [473](#)  
Craniopharyngioma, [473](#)  
Cranioplasty, [473](#)  
Craniosynostosis, [473](#)  
Craniosynostosis Repair, [474](#)  
Craniotabes, [474](#)  
Craniotomy, [474](#)  
Cranium, [474](#)  
Crapulence, [474](#)  
Craving, [474](#)  
Crazy Paving Pattern, [474](#)  
CRC, [474](#)  
CRE, [474](#)  
CRE Infection, [474](#)  
Cream, [474](#)  
Creatine, [474](#)  
Creatine Kinase, [474](#)  
Creatine Phosphokinase Test, [475](#)  
Creating A Family Health History, [475](#)  
Creatinine, [475](#)  
Creatinine Blood Test, [475](#)  
Creatinine Clearance Test, [475](#)  
Creatinine Test, [475](#)  
Creatinine Urine Test, [475](#)  
Creeping Eruption, [475](#)  
Cremasteric Reflex, [475](#)  
Crepitation, [475](#)  
Crepitus, [475](#)  
Crescentic Glomerulonephritis, [475](#)  
CREST Syndrome, [475](#)  
Creutzfeld-Jakob Disease, [475](#)  
Creutzfeldt Jakob Disease, [476](#)  
Creutzfeldt-Jakob Disease, [476](#)  
Cri Du Chat Syndrome, [476](#)  
Cri-Du-Chat Syndrome, [476](#)  
CRIA Syndrome, [476](#)  
Crib Death, [476](#)  
Cribiform Carcinoma, [476](#)  
Cribiform Plate, [476](#)  
Cricoid Cartilage, [476](#)  
Cricoid Pressure, [476](#)  
Cricothyroid Membrane, [476](#)  
Cricothyroidotomy, [476](#)  
Cricothyrotomy, [476](#)  
Crigler-Najjar Syndrome, [477](#)  
Crime Scene Investigation, [477](#)  
Crimean-Congo Hemorrhagic Fever, [477](#)  
Criminal Abortion, [477](#)  
CRISPR, [477](#)  
CRISPR Therapy, [477](#)  
CRISPR-Cas9, [477](#)  
Crista Galli, [477](#)  
Critical Care, [477](#)  
Critical Care Nursing, [477](#)  
Critical Findings, [477](#)  
Critical Illness Myopathy, [477](#)  
Critical Illness Polyneuropathy, [478](#)  
Critical Illness-Related Corticosteroid Insufficiency, [478](#)  
Critical Limb Ischemia, [478](#)  
Critical Results, [478](#)  
Critical Value, [478](#)  
Crohn Disease, [478](#)  
Crohn Disease Cobblestoning, [478](#)  
Crohn Disease Pathology, [478](#)  
Crohn Ileitis, [478](#)  
Crohn Ileocolitis, [478](#)  
Crohn's Disease, [478](#)  
Cronkhite-Canada Syndrome, [478](#)  
Cross-Eyes (Strabismus), [479](#)  
Cross-Matching, [479](#)  
Cross-Section, [479](#)  
Cross-Sectional Study, [479](#)  
Cross-Tolerance, [479](#)  
Crossed Extensor Reflex, [479](#)  
Crossmatch, [479](#)

---

---

Crosti-Gianotti Syndrome, [479](#)  
Crotch, [479](#)  
Croup, [479](#)  
Crouzon Syndrome, [479](#)  
Crow-Fukase Syndrome, [479](#)  
Crowe Sign, [479](#)  
Crown, [480](#)  
Crown Lengthening, [480](#)  
Crown-Rump Length, [480](#)  
Crowning, [480](#)  
CRP, [480](#)  
Cruciate Ligament, [480](#)  
Crush Injury, [480](#)  
Crush Syndrome, [480](#)  
Crushing Injury, [480](#)  
Crust, [480](#)  
Crusted Scabies, [480](#)  
Cruveilhier-Baumgarten Sign, [480](#)  
Cry For Help, [480](#)  
Crying In Childhood, [480](#)  
Crying In Infancy, [481](#)  
Cryoablation, [481](#)  
Cryobiopsy, [481](#)  
Cryoglobulinemia, [481](#)  
Cryoglobulinemic Vasculitis, [481](#)  
Cryoglobulins, [481](#)  
Cryopecty, [481](#)  
Cryoprecipitate, [481](#)  
Cryopreservation, [481](#)  
Cryopyrin-Associated Autoinflammatory Syndromes, [481](#)  
Cryopyrin-Associated Periodic Syndromes, [481](#)  
Cryostat, [481](#)  
Cryosurgery, [481](#)  
Cryotherapy, [481](#)  
Cryotherapy For Prostate Cancer, [482](#)  
Cryotherapy For The Skin, [482](#)  
Crypt, [482](#)  
Cryptitis, [482](#)  
Crypto, [482](#)  
Cryptococcal Infection, [482](#)  
Cryptococcal Meningitis, [482](#)  
Cryptococcic Meningitis, [482](#)  
Cryptococcosis, [482](#)  
Cryptococcus Neoformans, [482](#)  
Cryptogenic Organizing Pneumonia, [482](#)  
Cryptorchidism, [482](#)  
Cryptosporidiosis, [482](#)  
Cryptosporidium Enteritis, [483](#)  
Crystal Arthropathy, [483](#)  
Crystal-Induced Synovitis, [483](#)  
Crystalline Lens, [483](#)  
Crystalloids, [483](#)  
Crystalluria, [483](#)  
CSF Analysis, [483](#)  
CSF Cell Count, [483](#)  
CSF Coccidioides Complement Fixation Test, [483](#)  
CSF Glucose Test, [483](#)  
CSF Leak, [483](#)  
CSF Myelin Basic Protein, [483](#)  
CSF Oligoclonal Banding, [483](#)  
CSF Rhinorrhea, [483](#)  
CSF Smear, [484](#)  
CSF Total Protein, [484](#)  
CSF-VDRL Test, [484](#)  
CT, [484](#)  
CT Angiography, [484](#)  
CT Angiography of Abdomen and Pelvis, [484](#)  
CT Angiography of Head and Neck, [484](#)  
CT Angiography of the Extremities, [484](#)  
CT Angiography – Chest, [484](#)  
CT Chest, [484](#)  
CT Colonography, [484](#)  
CT Contrast, [484](#)  
CT Coronary Angiography, [485](#)  
CT Dose-Length Product, [485](#)  
CT Enterography, [485](#)  
CT Perfusion, [485](#)  
CT Pulmonary Angiography, [485](#)  
CT Scan, [485](#)  
CTD-Associated Interstitial Lung Disease, [485](#)  
Ctenocephalides Flea Bite, [485](#)  
CTL, [485](#)  
CTLA-4, [485](#)  
CTPA, [485](#)  
CTS, [485](#)  
Cubital, [485](#)  
Cubital Tunnel Syndrome, [485](#)  
Cubitus, [485](#)  
Cubitus Valgus, [486](#)  
Cubitus Varus, [486](#)  
Cuboid, [486](#)  
Cul-De-Sac, [486](#)  
Culdocentesis, [486](#)  
Cullen Sign, [486](#)  
Cullen's Sign, [486](#)  
Cultural Psychiatry, [486](#)  
Culture, [486](#)  
Culture And Sensitivity, [486](#)  
Culture Media, [486](#)  
Culture Swab, [486](#)  
Culture-Negative Endocarditis, [486](#)  
Cumulative Dose, [486](#)  
Cumulative Trauma Disorder, [487](#)  
Cup-To-Disc Ratio, [487](#)  
Cupping, [487](#)  
Cupping Therapy, [487](#)  
CURB-65 Score, [487](#)  
Curcumin, [487](#)

---



---

Cure, [487](#)  
Curettage, [487](#)  
Curettage Specimen, [487](#)  
Curette, [487](#)  
Curie, [487](#)  
Curling Ulcer, [487](#)  
Curling's Ulcer, [487](#)  
Currant Jelly Sputum, [487](#)  
Curschmann Spirals, [488](#)  
Curvature Of The Penis, [488](#)  
Curvature of the Spine (Scoliosis), [488](#)  
Curvularia Alcornii, [488](#)  
Cushing Disease, [488](#)  
Cushing Syndrome, [488](#)  
Cushing Syndrome Complications, [488](#)  
Cushing Syndrome Due To Adrenal Tumor, [488](#)  
Cushing Triad, [488](#)  
Cushing Ulcer, [488](#)  
Cushing's Disease, [488](#)  
Cushingoid, [489](#)  
Cusp, [489](#)  
Cusp Fracture, [489](#)  
Cut, [489](#)  
Cutaneous, [489](#)  
Cutaneous Abscess, [489](#)  
Cutaneous and Oral Melanocytic Nevi, [489](#)  
Cutaneous Angiosarcoma, [489](#)  
Cutaneous Anthrax, [489](#)  
Cutaneous B-Cell Lymphoma, [489](#)  
Cutaneous Bleeding, [489](#)  
Cutaneous Candidiasis, [489](#)  
Cutaneous Diphtheria, [489](#)  
Cutaneous Drug Eruptions, [2116](#)  
Cutaneous Drug Reaction, [489](#)  
Cutaneous Horn, [489](#)  
Cutaneous Langerhans Cell Histiocytosis, [490](#)  
Cutaneous Larva Migrans, [490](#)  
Cutaneous Leiomyoma, [490](#)  
Cutaneous Leishmaniasis, [490](#)  
Cutaneous Lipomas, [490](#)  
Cutaneous Lupus, [490](#)  
Cutaneous Lupus Erythematosus, [490](#)  
Cutaneous Lymphoma, [490](#)  
Cutaneous Manifestations of HIV, [490](#)  
Cutaneous Melanoma, [490](#)  
Cutaneous Metastases, [490](#)  
Cutaneous Metastasis, [490](#)  
Cutaneous Polyarteritis Nodosa, [491](#)  
Cutaneous Pseudolymphoma, [491](#)  
Cutaneous Skin Tag, [491](#)  
Cutaneous Small-Vessel Vasculitis, [491](#)  
Cutaneous Squamous Cell Carcinoma, [491](#)  
Cutaneous T-Cell Lymphoma, [491](#)  
Cutaneous Vasculitis, [491](#)  
Cuticle, [491](#)  
Cuticle Remover Poisoning, [491](#)  
Cutis Anserina, [491](#)  
Cutis Laxa, [491](#)  
Cutis Marmorata, [491](#)  
Cutis Verticis Gyrata, [491](#)  
Cuts And Puncture Wounds, [492](#)  
Cuts and Wounds, [492](#)  
Cuts, Scrapes And Puncture Wounds, [492](#)  
Cutting Balloon Catheter, [492](#)  
CVC, [492](#)  
Cyanide, [492](#)  
Cyanide Poisoning, [492](#)  
Cyanide Toxicity, [492](#)  
Cyanoacrylates Poisoning, [492](#)  
Cyanosis, [492](#)  
Cyanotic, [492](#)  
Cyanotic Heart Disease, [492](#)  
CyberKnife Radiosurgery, [492](#)  
Cycle Time, [492](#)  
Cyclic Citrullinated Peptide, [492](#)  
Cyclic Edema, [492](#)  
Cyclic Guanosine Monophosphate, [493](#)  
Cyclic Hormone Therapy, [493](#)  
Cyclic Migraines, [493](#)  
Cyclic Neutropenia, [493](#)  
Cyclic Vomiting Syndrome, [493](#)  
Cyclic Vomiting Syndrome Cross-Reference, [493](#)  
Cyclin-Dependent Kinase, [493](#)  
Cyclist's Syndrome, [493](#)  
Cyclooxygenase, [493](#)  
Cyclooxygenase Inhibitor, [2116](#)  
Cyclooxygenase-2, [493](#)  
Cyclops, [493](#)  
Cyclospora Infection, [493](#)  
Cyclosporiasis, [494](#)  
Cyclothymia, [494](#)  
Cyclothymic Disorder, [494](#)  
Cylindroma, [494](#)  
Cylindroma and Spiradenoma, [494](#)  
CYP2C19 Polymorphism, [494](#)  
CYP2C9 Polymorphism, [494](#)  
CYP2D6 Polymorphism, [494](#)  
CYP450 System, [494](#)  
Cyproheptadine Overdose, [494](#)  
Cyprus Fever, [494](#)  
Cyst, [494](#)  
Cyst, Bartholin's, [178](#)  
Cyst, Ganglion, [801](#)  
Cyst, Giardia, [829](#)  
Cyst, Kidney, [1089](#)  
Cyst, Ovarian, [1434](#)  
Cyst, Pancreatic, [1453](#)  
Cyst, Pilonidal, [1532](#)

---

---

Cyst, Spermatic, 1819  
Cystadenoma, 494  
Cystectomy, 495  
Cystic Acne, 495  
Cystic Diseases of the Kidney, 495  
Cystic Echinococcosis, 495  
Cystic Fibrosis, 495  
Cystic Fibrosis Carrier Screening, 495  
Cystic Hygroma, 495  
Cystic Kidney Disease, 495  
Cystic Lung Disease, 495  
Cystic Medial Necrosis, 495  
Cystic Teratoma, 495  
Cysticercosis, 496  
Cystine and Struvite Urinary Crystals, 496  
Cystine Stones, 496  
Cystinosis, 496  
Cystinuria, 496  
Cystitis, 496  
Cystitis Pathology, 496  
Cystitis, Interstitial, 1037  
Cystocele, 496  
Cystography, 496  
Cystoid Macular Edema, 496  
Cystoisosporiasis, 496  
Cystometric Study, 497  
Cystometrogram, 497  
Cystometry, 497  
Cystoscope, 497  
Cystoscopy, 497  
Cystostomy, 497  
Cystotome, 497  
Cystourethrography, 497  
Cysts, 497  
Cysts, Choledochal, 389  
Cysts, Epidermoid, 659  
Cysts, Ovarian, Functional, 786  
Cysts, Ovary, 1434  
Cysts, Pancreatic Inflammatory, 1454  
Cysts, Sebaceous, 659  
Cysts, True Pancreas, 1453  
Cytochrome B5 Reductase, 497  
Cytochrome C, 497  
Cytochrome P450, 497  
Cytogenetic Karyotyping, 497  
Cytogenetics, 497  
Cytokine, 498  
Cytokine Release Syndrome, 498  
Cytokine Storm, 498  
Cytokines, 498  
Cytologic Evaluation, 498  
Cytology, 498  
Cytology Exam Of Pleural Fluid, 498  
Cytology Exam Of Urine, 498  
Cytomegalovirus, 498  
Cytomegalovirus Gastroenteritis and Colitis, 498  
Cytomegalovirus Infection, 498  
Cytomegalovirus Retinitis, 498  
Cytopathology, 499  
Cytopenia, 499  
Cytoplasm, 499  
Cytoreductive Surgery, 499  
Cytoskeleton, 499  
Cytotoxic, 499  
Cytotoxic Alopecia, 499  
Cytotoxic T Cell, 499  
Cytotoxicity, 499  
D & C, 501  
D And C, 501  
D And E, 501  
D Trisomy Syndrome, 501  
D-Dimer, 501  
D-Dimer Test, 501  
D-Lactic Acidosis, 501  
D-Xylose Absorption, 502  
D.O., 502  
D2 Receptor, 502  
D&C Instruments, 502  
Da Costa's Syndrome, 502  
Dacarbazine, 502  
Daclizumab, 502  
Dacomitinib, 502  
Dacrocytes, 502  
Dacryoadenitis, 502  
Dacryocystitis, 502  
Dacryocystocele, 502  
Dacryocystography, 502  
Dacryocystorhinostomy, 503  
Dacryocyte Analysis, 503  
Dacryolithiasis, 503  
Dacryostenosis, 503  
Dactinomycin, 503  
Dactylitis, 503  
Dactylomegaly, 503  
Dactylosymphysis, 503  
Daidzein, 503  
Daily Bowel Care Program, 503  
Dakin Solution, 503  
Dalrymple Sign, 503  
Dalteparin, 503  
Dalteparin Sodium, 503  
Daltonism, 504  
DALY, 504  
Damage To The Amygdala, 504  
Damage-Control Resuscitation, 504  
Damage-Control Surgery, 504  
Damus-Kaye-Stansel Procedure, 504

---

---

Danazol, [504](#)  
 Dance Sign, [504](#)  
 Dandruff, [504](#)  
 Dandruff And Scalp Conditions, [504](#)  
 Dandy Fever, [504](#)  
 Dandy-Walker Malformation, [504](#)  
 Dandy-Walker Syndrome, [504](#)  
 Dane Particle, [505](#)  
 Dangerous Abbreviations, [505](#)  
 Danon Disease, [505](#)  
 Dantrolene, [505](#)  
 Dapsone, [505](#)  
 Daptomycin, [505](#)  
 Daptomycin Resistance, [505](#)  
 Darbepoetin Alfa, [505](#)  
 Darier Sign, [505](#)  
 Darier's Disease, [505](#)  
 Dark Adaptation, [505](#)  
 Dartos, [505](#)  
 Darunavir, [505](#)  
 Darwin's Tubercle, [506](#)  
 Dasatinib, [506](#)  
 DASH Diet, [506](#)  
 Dating Ultrasound, [506](#)  
 Daubenton's Plane, [506](#)  
 Daunorubicin, [506](#)  
 Daunorubicin Hydrochloride, [506](#)  
 Dav Regimen, [506](#)  
 David Procedure, [506](#)  
 Daw, [506](#)  
 Dawn Phenomenon, [506](#)  
 Dawson Fingers, [506](#)  
 Day Hospital, [506](#)  
 Day-Blindness, [507](#)  
 Daytime Sleepiness, [507](#)  
 DBA, [507](#)  
 Dcc Gene, [507](#)  
 DCIS, [507](#)  
 DDH, [507](#)  
 De Lange Syndrome, [507](#)  
 De Musset Sign, [507](#)  
 De Novo, [507](#)  
 De Novo Variant, [507](#)  
 De Quervain Tendinitis, [507](#)  
 De Quervain Tenosynovitis, [507](#)  
 De Quervain Thyroiditis, [507](#)  
 De Quervain's Tendonitis, [507](#)  
 De Quervain's Tenosynovitis, [507](#)  
 De Winter ECG Pattern, [508](#)  
 De Winter T-Wave Pattern, [508](#)  
 De-Escalation, [508](#)  
 Deacetylase Inhibitor, [508](#)  
 Dead Space, [508](#)  
 Dead Space Ventilation, [508](#)  
 Dead Time, [508](#)  
 Deafblindness, [508](#)  
 Deafferentation, [508](#)  
 Deafness, [508](#)  
 Deaner, [508](#)  
 Death, [508](#)  
 Death Rate, [509](#)  
 Death Rattle, [509](#)  
 Death, Sudden Cardiac, [1859](#)  
 Deauville Criteria, [509](#)  
 Deaver Retractor, [509](#)  
 Debility, [509](#)  
 Debride, [509](#)  
 Debridement, [509](#)  
 Debulk, [509](#)  
 Debulking, [509](#)  
 Decalcification, [509](#)  
 Decannulation, [509](#)  
 Decanoic Acid, [509](#)  
 Decapitation, [509](#)  
 Decay Correction, [509](#)  
 Deceased Donor, [510](#)  
 Decenoic Acid, [510](#)  
 Decerebrate Posture, [510](#)  
 Decerebrate Posturing, [510](#)  
 Decerebrate Rigidity, [510](#)  
 Decerebrate State, [510](#)  
 Decerebration, [510](#)  
 Decibel, [510](#)  
 Decidua, [510](#)  
 Decidua Basalis, [510](#)  
 Deciduitis, [510](#)  
 Deciduositis, [510](#)  
 Deciduous Teeth, [510](#)  
 Deciduous Tooth, [510](#)  
 Decision Making, [511](#)  
 Decision-Making Capacity, [511](#)  
 Decitabine, [511](#)  
 Declarative Memory, [511](#)  
 Decoction, [511](#)  
 Decompensated Cirrhosis, [511](#)  
 Decomposition, [511](#)  
 Decompression, [511](#)  
 Decompression Sickness, [511](#)  
 Deconditioning, [511](#)  
 Decongestant, [511](#)  
 Decontamination, [511](#)  
 Decorin, [511](#)  
 Decorticate Posture, [511](#)  
 Decorticate Posturing, [512](#)  
 Decorticate Rigidity, [512](#)  
 Decreased Alertness, [512](#)  
 Decreased Appetite, [512](#)  
 Decreased Breath Sounds, [512](#)

---

---

Decreased Concentration, [512](#)  
Decreased Dentin, [512](#)  
Decreased Libido, [512](#)  
Decreased Tear Production, [512](#)  
Decubitus, [512](#)  
Decubitus Ulcer, [512](#)  
Deep, [512](#)  
Deep Brain Stimulation, [512](#)  
Deep Femoral Artery, [512](#)  
Deep Infiltrating Endometriosis, [512](#)  
Deep Neck Space Infection, [513](#)  
Deep Sleep, [513](#)  
Deep Tendon Reflex, [513](#)  
Deep Tendon Reflexes, [513](#)  
Deep Vein Thrombosis, [513](#)  
Deep Vein Thrombosis Ultrasound, [513](#)  
Deep Venous Reflux, [513](#)  
Deerfly Fever, [513](#)  
Defaecation, [513](#)  
Defecation, [513](#)  
Defecation Syncope, [513](#)  
Defecography, [514](#)  
Defective Virus, [514](#)  
Defence Mechanisms, [514](#)  
Defense Mechanism, [514](#)  
Defense Mechanisms, [514](#)  
Defense Wounds, [514](#)  
Defenses, [514](#)  
Defensin, [514](#)  
Deferasirox, [514](#)  
Deferiprone, [514](#)  
Deferoxamine, [514](#)  
Defibrillation, [514](#)  
Defibrillator, [514](#)  
Defibrillator Implant, [514](#)  
Deficiency Diseases, [514](#)  
Deficiency, FAO, [723](#)  
Deficiency, Primary Immunodeficiency, [1594](#)  
Deficient Color Vision, [515](#)  
Deformation, [515](#)  
Deformity, [515](#)  
Degarelix, [515](#)  
Degeneration, [515](#)  
Degenerative Arthritis, [515](#)  
Degenerative Cervical Myelopathy, [515](#)  
Degenerative Chorea, [515](#)  
Degenerative Disc, [515](#)  
Degenerative Disc Disease, [515](#)  
Degenerative Disease, [515](#)  
Degenerative Disk Disease, [515](#)  
Degenerative Disorder, [515](#)  
Degenerative Joint Disease, [515](#)  
Degenerative Spine Disease, [516](#)  
Degenerative Spondylolisthesis, [516](#)  
Deglutition, [516](#)  
Degos Disease (Malignant Atrophic Papulosis), [516](#)  
Degranulation, [516](#)  
Degrees Fahrenheit, [516](#)  
Dehiscence, [516](#)  
DEHP, [516](#)  
Dehydration, [516](#)  
Dehydroascorbic Acid, [516](#)  
Dehydroepiandrosterone, [516](#)  
Deinstitutionalize, [516](#)  
Deiodinase, [516](#)  
Deja Vu, [517](#)  
Dejerine-Roussy Syndrome, [517](#)  
Delavirdine, [517](#)  
Delayed Cord Clamping, [517](#)  
Delayed Ejaculation, [517](#)  
Delayed Emergence, [517](#)  
Delayed Gastric Emptying, [517](#)  
Delayed Graft Function, [517](#)  
Delayed Growth, [517](#)  
Delayed Hemolytic Transfusion Reaction, [517](#)  
Delayed Hypersensitivity, [517](#)  
Delayed or Absent Tooth Formation, [517](#)  
Delayed Orgasm, [517](#)  
Delayed Puberty, [518](#)  
Delayed Puberty In Boys, [518](#)  
Delayed Puberty In Girls, [518](#)  
Delayed Sleep Phase, [518](#)  
Delayed Sleep Phase Syndrome, [518](#)  
Delayed Sleep-Wake Phase Sleep Disorder, [518](#)  
Deletion, [518](#)  
Deletion Mutation, [518](#)  
Delftia, [518](#)  
Delirium, [518](#)  
Delirium In Anesthesia, [518](#)  
Delirium Superimposed On Dementia, [518](#)  
Delirium Tremens, [518](#)  
Delirium Tremens (DTs), [519](#)  
Delivery, [519](#)  
Delivery Of Posterior Arm, [519](#)  
Delivery Presentations, [519](#)  
Delta Agent, [519](#)  
Delta Waves, [519](#)  
Delta-ALA Urine Test, [519](#)  
Delta-Tocopherol, [519](#)  
Deltacoronavirus, [519](#)  
Deltaretrovirus, [519](#)  
Deltoid, [519](#)  
Deltoid Ligament, [519](#)  
Deltoid Muscle, [519](#)  
Delusion, [519](#)  
Delusional Disorder, [520](#)  
Delusional Infestation, [520](#)  
Demarcation, [520](#)

---

---

Demeclocycline, [520](#)  
Demecolcine, [520](#)  
Dementia, [520](#)  
Dementia And Driving, [520](#)  
Dementia Behavioral and Sleep Changes, [520](#)  
Dementia Due To Metabolic Causes, [520](#)  
Dementia with Lewy Bodies, [520](#)  
Dementia, Frontotemporal, [781](#)  
Dementia, Lewy Body, [1135](#)  
Dementia, Major Neurocognitive, [520](#)  
Dementia, Vascular, [2026](#)  
Demethylation, [520](#)  
Demineralization, [520](#)  
Demulcent, [521](#)  
Demyelinating CNS Disease, [521](#)  
Demyelinating Disease, [521](#)  
Demyelination, [521](#)  
Denaturant, [521](#)  
Dendrite, [521](#)  
Dendrites, [521](#)  
Dendritic Cell, [521](#)  
Dendritic Cells, [521](#)  
Dendritic Spine, [521](#)  
Dendritic Ulcer, [521](#)  
Denervate, [521](#)  
Denervation, [521](#)  
Dengue, [521](#)  
Dengue Fever, [522](#)  
Dengue Hemorrhagic Fever, [522](#)  
Dengue Shock Syndrome, [522](#)  
Dengue Virus, [522](#)  
Denial, [522](#)  
Denileukin Diftitox, [522](#)  
Dennie-Morgan Fold, [522](#)  
Denominator, [522](#)  
Denosumab, [522](#)  
Denosumab For Osteoporosis, [522](#)  
Dens, [522](#)  
Dens Evaginatus, [522](#)  
Dens Invaginatus, [522](#)  
Densitometry, [522](#)  
Density, [523](#)  
Dent Disease, [523](#)  
Dental Abscess, [523](#)  
Dental Age, [523](#)  
Dental Alloy, [523](#)  
Dental Alveolus, [523](#)  
Dental Amalgam, [523](#)  
Dental Anesthesia, [523](#)  
Dental Anxiety, [523](#)  
Dental Arch, [523](#)  
Dental Articulator, [523](#)  
Dental Avulsion, [523](#)  
Dental Bone Graft, [523](#)  
Dental Braces, [523](#)  
Dental Bridge, [524](#)  
Dental Calculus, [524](#)  
Dental Caries, [524](#)  
Dental Cavities, [524](#)  
Dental Cement, [524](#)  
Dental Ceramic, [524](#)  
Dental Clasp, [524](#)  
Dental Crown, [524](#)  
Dental Crowns, [524](#)  
Dental Deposit, [524](#)  
Dental Disease, [524](#)  
Dental Drill, [524](#)  
Dental Dysplasia, [524](#)  
Dental Erosion, [525](#)  
Dental Extraction, [525](#)  
Dental Fillings, [525](#)  
Dental Fissure, [525](#)  
Dental Floss, [525](#)  
Dental Fluorosis, [525](#)  
Dental Hygiene, [525](#)  
Dental Implant, [525](#)  
Dental Impression Material, [525](#)  
Dental Injuries, [525](#)  
Dental Inlay, [525](#)  
Dental Material, [525](#)  
Dental Occlusion, [525](#)  
Dental Overjet, [525](#)  
Dental Papilla, [526](#)  
Dental Pellicle, [526](#)  
Dental Plaque, [526](#)  
Dental Polishing, [526](#)  
Dental Porcelain, [526](#)  
Dental Precision Attachment, [526](#)  
Dental Procedure, [526](#)  
Dental Prosthesis, [526](#)  
Dental Public Health, [526](#)  
Dental Pulp, [526](#)  
Dental Pulp Necrosis, [526](#)  
Dental Pulpitis, [526](#)  
Dental Radiographs, [526](#)  
Dental Radiography, [526](#)  
Dental Restorative Materials, [527](#)  
Dental Root, [527](#)  
Dental Sac, [527](#)  
Dental Screening, [527](#)  
Dental Sealant, [527](#)  
Dental Sealants, [527](#)  
Dental Sleep Apnea Therapy, [527](#)  
Dental Splint, [527](#)  
Dental Trauma, [527](#)  
Dental Traumatology, [527](#)  
Dental X-Rays, [527](#)  
Dentate Gyrus, [527](#)

---

---

Dentate Line, [527](#)  
Dentatorubral-Pallidolusian Atrophy, [527](#)  
Dentifrice, [528](#)  
Dentigerous Cyst, [528](#)  
Dentin, [528](#)  
Dentin Hypersensitivity, [528](#)  
Dentin Sensitivity, [528](#)  
Dentinogenesis, [528](#)  
Dentinogenesis Imperfecta, [528](#)  
Dentist, [528](#)  
Dentistry, [528](#)  
Dentition, [528](#)  
Denture, [528](#)  
Denture Base, [528](#)  
Denture Problems, [528](#)  
Dentures, [528](#)  
Denturist, [528](#)  
Denver Developmental Screening Test, [529](#)  
Denys-Drash Syndrome, [529](#)  
Deoxycoformycin, [529](#)  
Deoxycytidine, [529](#)  
Deoxycytidine Kinase, [529](#)  
Deoxyglucose, [529](#)  
Deoxyguanosine, [529](#)  
Deoxyhemoglobin, [529](#)  
Deoxyribonuclease, [529](#)  
Deoxyribonucleic Acid, [529](#)  
Deoxyribonucleosides, [529](#)  
Deoxyribonucleotides, [529](#)  
Dependence, [529](#)  
Dependence, Nicotine, [1358](#)  
Dependent Personality Disorder, [529](#)  
Dependovirus, [529](#)  
Depersonalization, [530](#)  
Depersonalization-Derealization Disorder, [530](#)  
Dephosphorylation, [530](#)  
Depigmentation, [530](#)  
Depilation, [530](#)  
Depilatory Poisoning, [530](#)  
Depolarization, [530](#)  
Depolarizing Neuromuscular Blocker, [530](#)  
Depot Implant, [530](#)  
Deprescribing, [530](#)  
Depressed Fracture, [530](#)  
Depression, [530](#)  
Depression Screening, [531](#)  
Depression, Postpartum, [1572](#)  
Depression, Teen, [1895](#)  
Depressive Disorder, [531](#)  
Depressive Illness, [531](#)  
Depth Perception, [531](#)  
Depurination, [531](#)  
Derbyshire Neck, [531](#)  
Dercum Disease (Adiposis Dolorosa), [531](#)  
Dercum's Disease, [531](#)  
Derealization, [531](#)  
Dermabrasion, [531](#)  
Dermal, [531](#)  
Dermal Fillers, [531](#)  
Dermal Melanocytosis, [531](#)  
Dermal Papilla, [531](#)  
Dermal Structure, [531](#)  
Dermatan Sulfate, [532](#)  
Dermatitis, [532](#)  
Dermatitis Herpetiformis, [532](#)  
Dermatitis, Atopic, [142](#)  
Dermatitis, Cercarial, [1871](#)  
Dermatitis, Contact, [451](#)  
Dermatitis, Scratch, [1347](#)  
Dermatitis, Seborrheic, [1759](#)  
Dermatofibroma, [532](#)  
Dermatofibrosarcoma, [532](#)  
Dermatofibrosarcoma Protuberans, [532](#)  
Dermatofibrosarcoma Protuberans Of The Breast, [532](#)  
Dermatoglyphics, [532](#)  
Dermatographia, [532](#)  
Dermatographic Urticaria, [532](#)  
Dermatographism, [532](#)  
Dermatologic, [532](#)  
Dermatologic Disorder, [532](#)  
Dermatologic Manifestations of Hematologic Disease, [532](#)  
Dermatologist, [533](#)  
Dermatology, [533](#)  
Dermatoma, [533](#)  
Dermatome, [533](#)  
Dermatomycosis, [533](#)  
Dermatomyositis, [533](#)  
Dermatopathology, [533](#)  
Dermatophyte, [533](#)  
Dermatophytosis, [533](#)  
Dermatoplasty, [533](#)  
Dermatoscope, [533](#)  
Dermatosis, [533](#)  
Dermis, [533](#)  
Dermoid, [533](#)  
Dermoid Cyst, [534](#)  
Dermoid Cyst Of The Ovary, [534](#)  
Dermoscopy, [534](#)  
Descemet Membrane Endothelial Keratoplasty, [534](#)  
Descending Colon, [534](#)  
Descending Thoracic Aorta, [534](#)  
Desensitization, [534](#)  
Desferrioxamine, [534](#)  
Desflurane, [534](#)  
Desiccate, [534](#)  
Desiccated Thyroid, [535](#)  
Designer Steroid, [2117](#)  
Desipramine, [535](#)

---



---

Desipramine Hydrochloride Overdose, [535](#)  
 Desmin, [535](#)  
 Desmoid Tumor, [535](#)  
 Desmoid Tumors, [535](#)  
 Desmoid Tumour, [535](#)  
 Desmoplasia, [535](#)  
 Desmoplastic, [535](#)  
 Desmoplastic Fibroblastoma, [535](#)  
 Desmoplastic Fibroma, [535](#)  
 Desmoplastic Infantile Ganglioglioma, [535](#)  
 Desmoplastic Melanoma, [535](#)  
 Desmoplastic Reaction, [535](#)  
 Desmoplastic Small Round Cell Tumor, [536](#)  
 Desmoplastic Small Round Cell Tumors, [536](#)  
 Desmoplastic Trichoepithelioma, [536](#)  
 Desmopressin, [536](#)  
 Desmosome, [536](#)  
 Desogestrel, [536](#)  
 Desonide, [536](#)  
 Desoximetasone, [536](#)  
 Desoxycorticosterone, [536](#)  
 Desquamation, [536](#)  
 Desquamative Interstitial Pneumonia, [536](#)  
 Desulfovibrio, [536](#)  
 Desvenlafaxine Succinate, [536](#)  
 Desynchronosis, [537](#)  
 Detached, [537](#)  
 Detached Retina, [537](#)  
 Detergent, [537](#)  
 Deterministic Effects, [537](#)  
 Detoxification, [537](#)  
 Detritic Synovitis, [537](#)  
 Detrusor Instability, [537](#)  
 Detrusor Muscle, [537](#)  
 Detrusor Sphincter Dyssynergia, [537](#)  
 Detumescence, [537](#)  
 Deubiquitinating Enzymes, [537](#)  
 Deubiquitination, [537](#)  
 Deuteranopia, [537](#)  
 Deuterium, [537](#)  
 Development, [538](#)  
 Developmental Coordination Disorder, [538](#)  
 Developmental Delay, [538](#)  
 Developmental Disabilities, [538](#)  
 Developmental Disorder, [538](#)  
 Developmental Disorders, [538](#)  
 Developmental Dysplasia Of The Hip, [538](#)  
 Developmental Expressive Language Disorder, [538](#)  
 Developmental Genital Tract Anomalies, [538](#)  
 Developmental History, [538](#)  
 Developmental Milestone, [538](#)  
 Developmental Milestones, [538](#)  
 Developmental Milestones Record, [539](#)  
 Developmental Process, [539](#)  
 Developmental Psychology, [539](#)  
 Developmental Reading Disorder, [539](#)  
 Developmental Regression, [539](#)  
 Developmental Stage, [539](#)  
 Developmental Surveillance, [539](#)  
 Deviated Nasal Septum, [539](#)  
 Deviated Septum, [539](#)  
 Deviation, [539](#)  
 Devic Disease, [539](#)  
 Device Embolisation, [539](#)  
 Devices For Hearing Loss, [539](#)  
 Dexamethasone, [540](#)  
 Dexamethasone Suppression Test, [540](#)  
 Dexamphetamine, [540](#)  
 Dexibuprofen, [540](#)  
 Dexketoprofen, [540](#)  
 Dexlansoprazole, [540](#)  
 Dexmedetomidine, [540](#)  
 Dexmethylphenidate, [540](#)  
 Dexrazoxane, [540](#)  
 Dexrazoxane Hydrochloride, [540](#)  
 Dextrin, [540](#)  
 Dextroamphetamine, [540](#)  
 Dextrocardia, [540](#)  
 Dextromethorphan, [541](#)  
 Dextromethorphan Overdose, [541](#)  
 Dextroposition Of The Heart, [541](#)  
 Dextrose, [541](#)  
 DFSP, [541](#)  
 DH, [541](#)  
 DHA, [541](#)  
 DHEA, [541](#)  
 DHEA-Sulfate Test, [541](#)  
 Diabetes And Alcohol, [541](#)  
 Diabetes And Exercise, [541](#)  
 Diabetes And Eye Disease, [541](#)  
 Diabetes And Eye Problems, [541](#)  
 Diabetes And Kidney Disease, [541](#)  
 Diabetes And Metabolic Health, [542](#)  
 Diabetes And Nerve Damage, [542](#)  
 Diabetes Complications, [542](#)  
 Diabetes Eye Exams, [542](#)  
 Diabetes in Children: Dealing with Low Blood Sugar, [542](#)  
 Diabetes in Children: Preventing Low Blood Sugar, [542](#)  
 Diabetes in Pregnancy, [542](#)  
 Diabetes in Pregnancy (Gestational Diabetes), [542](#)  
 Diabetes Insipidus, [542](#)  
 Diabetes Medications, [542](#)  
 Diabetes Mellitus, [542](#)  
 Diabetes Mellitus and Pregnancy, [542](#)  
 Diabetes Mellitus Type 1, [543](#)  
 Diabetes Mellitus Type 2, [543](#)  
 Diabetes Prevention, [543](#)

---

---

Diabetes Treatment, [543](#)  
Diabetes, Gestational, [826](#)  
Diabetes, Pre, [1581](#)  
Diabetes, Type 1, [1978](#)  
Diabetes, Type 2, [1978](#)  
Diabetic Amyotrophy, [543](#)  
Diabetic Angiopathy, [543](#)  
Diabetic Autonomic Neuropathy, [543](#)  
Diabetic Cardiomyopathies, [543](#)  
Diabetic Cardiomyopathy, [543](#)  
Diabetic Coma, [543](#)  
Diabetic Complications, [543](#)  
Diabetic Diet, [543](#)  
Diabetic Eye Disease, [543](#)  
Diabetic Focal Neuropathy, [544](#)  
Diabetic Foot, [544](#)  
Diabetic Foot Disease, [544](#)  
Diabetic Foot Infections, [544](#)  
Diabetic Foot Ulcer, [544](#)  
Diabetic Gastroparesis, [544](#)  
Diabetic Hyperglycemic Hyperosmolar Syndrome, [544](#)  
Diabetic Hyperosmolar Hyperglycemic State, [544](#)  
Diabetic Hypoglycemia, [544](#)  
Diabetic Ketoacidosis, [544](#)  
Diabetic Ketoacidosis Monitoring, [544](#)  
Diabetic Kidney Disease, [544](#)  
Diabetic Macular Edema, [545](#)  
Diabetic Mastopathy, [545](#)  
Diabetic Microvascular Complications, [545](#)  
Diabetic Nephropathy, [545](#)  
Diabetic Neuropathy, [545](#)  
Diabetic Neuropathy Entry, [545](#)  
Diabetic Peripheral Neuropathy, [545](#)  
Diabetic Proximal Neuropathy, [545](#)  
Diabetic Retinopathy, [545](#)  
Diabetic Third Nerve Palsy, [545](#)  
Diabetogenic, [545](#)  
Diabetologist, [545](#)  
Diabrosis, [545](#)  
Diacetoxyscirpenol, [545](#)  
Diacylglycerol, [546](#)  
Diagnosis, [546](#)  
Diagnostic, [546](#)  
Diagnostic Imaging, [546](#)  
Diagnostic Laparoscopy, [546](#)  
Diagnostic Microbiology, [546](#)  
Diagnostic Peritoneal Lavage, [546](#)  
Diagnostic Procedure, [546](#)  
Diagnostic Radiology, [546](#)  
Diagnostic Service, [546](#)  
Diagnostic Test, [546](#)  
Diagnostic Trial, [546](#)  
Diagnostic Uncertainty, [546](#)  
Diagnostics, [547](#)  
Diakinesis, [547](#)  
Dialectical Behavior Therapy, [547](#)  
Dialister Pneumosintes, [547](#)  
Dialysis, [547](#)  
Dialysis Access, [547](#)  
Dialysis Adequacy, [547](#)  
Dialysis Complications of Chronic Renal Failure, [547](#)  
Dialysis Disequilibrium Syndrome, [547](#)  
Dialysis Machine, [547](#)  
Dialysis Solutions, [547](#)  
Dialysis Therapy, [547](#)  
Dialysis-Related Beta-2m Amyloidosis, [547](#)  
Diamine, [547](#)  
Diamond-Blackfan Anemia, [548](#)  
Diamond-Gardner Syndrome, [548](#)  
Diamorphine, [548](#)  
Dianisidine, [548](#)  
Diapedesis, [548](#)  
Diaper Rash, [548](#)  
Diaphoresis, [548](#)  
Diaphoretic, [548](#)  
Diaphragm, [548](#)  
Diaphragmatic Crus, [548](#)  
Diaphragmatic Eventration, [548](#)  
Diaphragmatic Hernia, [548](#)  
Diaphragmatic Paralysis, [549](#)  
Diaphragmatic Rupture, [549](#)  
Diaphysis, [549](#)  
Diarrhea, [549](#)  
Diarrhea, Antibiotic-Associated, [2117](#)  
Diarrhea, Constipation, Gas, and Bloating, [249](#)  
Diarrhea, Traveler's, [1982](#)  
Diarrhea-Predominant IBS, [549](#)  
Diarrhoea, [549](#)  
Diarthrosis, [549](#)  
Diascopy, [549](#)  
Diastase, [549](#)  
Diastasis, [549](#)  
Diastasis Recti, [549](#)  
Diastematomyelia, [549](#)  
Diastole, [549](#)  
Diastolic, [549](#)  
Diastolic Blood Pressure, [549](#)  
Diastolic Dysfunction, [550](#)  
Diastolic Heart Failure, [550](#)  
Diastolic Murmur, [550](#)  
Diastolic Murmurs, [550](#)  
Diathermy, [550](#)  
Diathesis, [550](#)  
Diatom Test, [550](#)  
Diatomaceous Earth, [550](#)  
Diatrizoate Meglumine, [550](#)  
Diazepam, [550](#)  
Diazepam Overdose, [550](#)

---

---

Diazinon, [550](#)  
Diazinon Poisoning, [550](#)  
Diazonium Compounds, [551](#)  
Diazoxide, [551](#)  
Dibekacin, [551](#)  
Dibenzepin, [551](#)  
Dibucaine, [551](#)  
Dicarboxylic Acid, [551](#)  
Dichlorophen, [551](#)  
Dichlorvos, [551](#)  
Dichorionic Diamniotic Twin Placenta, [551](#)  
Dichorionic Twins, [551](#)  
Diclofenac, [551](#)  
Diclofenac Sodium, [551](#)  
Diclofenac Sodium Overdose, [551](#)  
Dicloxacin, [551](#)  
DICOM, [552](#)  
Dicoumarol, [552](#)  
Dicrotic, [552](#)  
Dicrotophos, [552](#)  
Dicyclomine, [552](#)  
Dieffenbachia Poisoning, [552](#)  
Diencephalon, [552](#)  
Dientamoebiasis, [552](#)  
DIEP Flap, [552](#)  
Diesel Oil, [552](#)  
Diestrus, [552](#)  
Diet Therapy, [552](#)  
Diet, Diabetic, [543](#)  
Diet, Ulcerative Colitis, [1018](#)  
Dietary Assessment, [552](#)  
Dietary Calcium, [553](#)  
Dietary Carbohydrate, [553](#)  
Dietary Fat, [553](#)  
Dietary Fats, [553](#)  
Dietary Fats and Oils, [553](#)  
Dietary Fiber, [553](#)  
Dietary Fibre, [553](#)  
Dietary Inflammation, [553](#)  
Dietary Iron, [553](#)  
Dietary Phospholipid, [553](#)  
Dietary Phosphorus, [553](#)  
Dietary Protein, [553](#)  
Dietary Reference Intakes, [553](#)  
Dietary Sterol, [554](#)  
Dietary Sugars, [554](#)  
Dietary Supplement, [554](#)  
Dietary Supplementation, [554](#)  
Dietary Supplements, [554](#)  
Dietetics, [554](#)  
Diethylcarbamazine, [554](#)  
Diethylhexylphthalate, [554](#)  
Diethylpropion, [554](#)  
Diethylstilbestrol, [554](#)  
Dietitian, [554](#)  
Dietl Crisis, [554](#)  
Dieulafoy Lesion, [554](#)  
Diff Dx, [555](#)  
Differences Of Sex Development, [555](#)  
Differential Blood Count, [555](#)  
Differential Diagnosis, [555](#)  
Differential Diagnosis of Viral Respiratory Infection and Allergic Rhinitis, [499](#)  
Differentiated Cancer, [555](#)  
Differentiation, [555](#)  
Differentiation Antigens, [555](#)  
Difficile, Clostridium, [416](#)  
Difficult Airway, [555](#)  
Difficult Intubation, [555](#)  
Difficult Urination, [555](#)  
Difficult-to-Control Asthma, [555](#)  
Difficulty Speaking, [555](#)  
Difficulty Swallowing, [555](#)  
Difficulty Swallowing (Dysphagia), [555](#)  
Difficulty Urinating, [555](#)  
Difficulty Walking, [556](#)  
Diffuse, [556](#)  
Diffuse Alopecia, [556](#)  
Diffuse Alveolar Hemorrhage, [556](#)  
Diffuse Astrocytoma, [556](#)  
Diffuse Axonal Injury, [556](#)  
Diffuse Esophageal Spasm, [556](#)  
Diffuse Hair Loss, [556](#)  
Diffuse Idiopathic Skeletal Hyperostosis, [556](#)  
Diffuse Large B-Cell Lymphoma, [556](#)  
Diffuse Midline Glioma, [556](#)  
Diffuse Parenchymal Lung Disease, [556](#)  
Diffuse Parenchymal Lung Diseases, [557](#)  
Diffuse Proliferative Glomerulonephritis, [557](#)  
Diffuse Scleroderma, [557](#)  
Diffusing Capacity, [557](#)  
Diffusing Capacity Of The Lung, [557](#)  
Diffusion, [557](#)  
Diffusion Anisotropy, [557](#)  
Diffusion Capacity, [557](#)  
Diffusion Restriction, [557](#)  
Diffusion Tensor Imaging, [557](#)  
Diffusion-Weighted Imaging, [557](#)  
Diflunisal, [557](#)  
Digastric Muscle, [557](#)  
DiGeorge Syndrome, [557](#)  
Digestion, [557](#)  
Digestive Diseases, [558](#)  
Digestive Disorder, [558](#)  
Digestive Health, [558](#)  
Digestive System, [558](#)  
Digestive Tract, [558](#)  
Digit, [558](#)

---

---

Digital Anorectal Examination, [558](#)  
Digital Breast Tomosynthesis, [558](#)  
Digital Clubbing, [558](#)  
Digital Fibrokeratoma, [558](#)  
Digital Ischemia, [558](#)  
Digital Mammography, [558](#)  
Digital Mucous Cyst, [558](#)  
Digital Papillary Adenocarcinoma, [559](#)  
Digital Radiography, [559](#)  
Digital Rectal Exam, [559](#)  
Digital Rectal Examination, [559](#)  
Digital Subtraction Angiography, [559](#)  
Digitalis, [2117](#)  
Digitalis Preparation, [559](#)  
Digitalis Toxicity, [559](#)  
Digitate Warts, [559](#)  
Digitoxin, [559](#)  
Digoxin, [559](#)  
Digoxin Immune Fab, [559](#)  
Digoxin Test, [559](#)  
Digoxin Toxicity, [559](#)  
Dihydrocodeine, [560](#)  
Dihydroergotamine, [560](#)  
Dihydropyrimidine Dehydrogenase Deficiency (DPYD-Related), [560](#)  
Dihydrotestosterone, [560](#)  
Dihydroxyacetone, [560](#)  
Dilantin Overdose, [560](#)  
Dilatation, [560](#)  
Dilate, [560](#)  
Dilated Cardiomyopathy, [560](#)  
Dilated Cardiomyopathy Prognosis, [560](#)  
Dilated Pore of Winer, [560](#)  
Dilation, [560](#)  
Dilation and Curettage, [560](#)  
Dilation And Evacuation, [561](#)  
Dilator, [561](#)  
Diltiazem, [561](#)  
Dilute Russell Viper Venom Time, [561](#)  
Dilution, [561](#)  
Dilution Anemia, [561](#)  
Dimenhydrinate, [561](#)  
Dimenhydrinate Overdose, [561](#)  
Dimercaprol, [561](#)  
Dimesna, [561](#)  
Dimethoate, [561](#)  
Dimethoxon, [561](#)  
Dimethyl Fumarate, [561](#)  
Dimethyl Fumarate Allergy, [562](#)  
Dimethyl Sulfoxide, [562](#)  
Dimethyl Sulphoxide, [562](#)  
Dimethylhydrazines, [562](#)  
Dimetilan, [562](#)  
Diminished Ovarian Reserve, [562](#)  
Dimocillin, [562](#)  
Dimorphic, [562](#)  
Dinophyceae, [562](#)  
Dinoprost, [562](#)  
Dinoprostone, [562](#)  
Dinucleoside Phosphates, [562](#)  
Dinutuximab, [562](#)  
Dioctyl Sodium Sulphosuccinate, [563](#)  
Diopter, [563](#)  
Dioxin, [563](#)  
Dipeptide, [563](#)  
Dipeptidyl Peptidase-4 Inhibitors, [563](#)  
Dipetalonema Infection, [563](#)  
Diphallia, [563](#)  
Diphenhydramine, [563](#)  
Diphenhydramine Overdose, [563](#)  
Diphenoxylate, [563](#)  
Diphosphate, [563](#)  
Diphosphonate, [563](#)  
Diphtheria, [563](#)  
Diphtheria Toxin, [563](#)  
Diphtheria Vaccine, [2117](#)  
Diphtheroid Bacillus, [564](#)  
Diphyllobothriasis, [564](#)  
Diphyllobothrium Latum, [564](#)  
Diplacosis, [564](#)  
Diplegia, [564](#)  
Diplococcus, [564](#)  
Diploid Cell, [564](#)  
Diploidy, [564](#)  
Diplopia, [564](#)  
Dipole, [564](#)  
Dipper, [564](#)  
Diprosopus, [564](#)  
Dipsosis, [564](#)  
Dipstick Device, [564](#)  
Diptera, [564](#)  
Dipylidium Caninum, [565](#)  
Dipyridamole, [565](#)  
Direct Acting Antivirals, [565](#)  
Direct Action, [565](#)  
Direct Antiglobulin Test, [565](#)  
Direct Coombs Test, [565](#)  
Direct Hernia, [565](#)  
Direct Inguinal Hernia, [565](#)  
Direct Oral Anticoagulants, [565](#)  
Direct Transmission, [565](#)  
Direct Veneer, [565](#)  
Directly Observed Therapy, [565](#)  
Dirofilaria Repens, [565](#)  
Dirt-Eating, [566](#)  
Dirty Necrosis, [566](#)  
Disability, [566](#)  
Disability Evaluation, [566](#)

---

---

Disability-Adjusted Life Year, [566](#)  
 Disaccharidase, [566](#)  
 Disaccharide, [566](#)  
 Disarticulation, [566](#)  
 Disc, [566](#)  
 Discectomy, [566](#)  
 Discitis, [566](#)  
 Discography, [566](#)  
 Discoid, [566](#)  
 Discoid Eczema, [566](#)  
 Discoid Lupus, [567](#)  
 Discoid Lupus Erythematosus, [567](#)  
 Discoloration, [567](#)  
 Discomfort, [567](#)  
 Discordance, [567](#)  
 Discordant, [567](#)  
 Discordant Couple, [567](#)  
 Disease, [567](#)  
 Disease Progression, [567](#)  
 Disease Reporting, [567](#)  
 Disease Reservoir, [567](#)  
 Disease Response, [567](#)  
 Disease Surveillance, [567](#)  
 Disease Susceptibility, [567](#)  
 Disease Vector, [568](#)  
 Disease-Free Survival, [568](#)  
 Disease-Modifying Antirheumatic Drug, [2117](#)  
 Disfigurement, [568](#)  
 Disinfectant, [568](#)  
 Disinfection, [568](#)  
 Disk, [568](#)  
 Disk Battery Ingestion, [568](#)  
 Disk Diffusion Susceptibility Test, [568](#)  
 Diskectomy, [568](#)  
 Diskitis, [568](#)  
 Dislocated Ankle, [568](#)  
 Dislocated Elbow, [568](#)  
 Dislocated Hip, [568](#)  
 Dislocated Knee, [568](#)  
 Dislocated Shoulder, [569](#)  
 Dislocation, [569](#)  
 Dislocation Of The Lens, [569](#)  
 Disofenin Scan, [569](#)  
 Disopyramide, [569](#)  
 Disorder, [569](#)  
 Disorder, Antisocial Personality, [110](#)  
 Disorder, Asperger, [134](#)  
 Disordered Thinking, [569](#)  
 Disorders of Fluid, Electrolyte, and Acid-Base Balance, [569](#)  
 Disorders of Oral Pigmentation, [569](#)  
 Disorders, Sleep, [1801](#)  
 Disorganized Symptoms in Schizophrenia, [569](#)  
 Disorientation, [569](#)  
 Dispensary, [569](#)  
 Dispensed Amount, [569](#)  
 Dispensing, [570](#)  
 Displacement, [570](#)  
 Disruption, [570](#)  
 Disruptive, Impulse-Control, And Conduct Disorders, [570](#)  
 Disruptive, Impulse-Control, and Conduct Disorders, [570](#)  
 Dissect, [570](#)  
 Dissecting Aneurysm, [570](#)  
 Dissection, [570](#)  
 Dissection, Aortic, [113](#)  
 Disseminate, [570](#)  
 Disseminated Coccidioidomycosis, [570](#)  
 Disseminated Cutaneous Blastomycosis, [570](#)  
 Disseminated Gonococcal Infection, [570](#)  
 Disseminated Intravascular Coagulation, [570](#)  
 Disseminated Mycobacterium Avium–Intracellular Infection, [570](#)  
 Disseminated Neuroblastoma, [570](#)  
 Disseminated Sclerosis, [571](#)  
 Disseminated Tuberculosis, [571](#)  
 Disseminated Varicella Infection, [571](#)  
 Disseminated Zoster, [571](#)  
 Dissociation, [571](#)  
 Dissociative Amnesia, [571](#)  
 Dissociative Disorder, [571](#)  
 Dissociative Disorders, [571](#)  
 Dissociative Identity Disorder, [571](#)  
 Distal, [571](#)  
 Distal Convoluted Tubule, [571](#)  
 Distal Median Nerve Dysfunction, [571](#)  
 Distal Pancreatectomy, [571](#)  
 Distal Radius Fractures, [572](#)  
 Distal Renal Tubular Acidosis, [572](#)  
 Distal Splenorenal Shunt, [572](#)  
 Distal Tubule, [572](#)  
 Distamycins, [572](#)  
 Distant, [572](#)  
 Distended, [572](#)  
 Distension, [572](#)  
 Distention, [572](#)  
 Distichia, [572](#)  
 Distortion, [572](#)  
 Distraction Osteogenesis, [572](#)  
 Distress, [572](#)  
 Distributive Shock, [572](#)  
 District Nurse, [573](#)  
 Disulfiram, [2117](#)  
 Disulfiram Toxicity, [573](#)  
 Disulfiram-Like Reaction, [573](#)  
 Disulfoton, [573](#)  
 Dithiolethione, [573](#)  
 Ditiocarb, [573](#)

---

---

Diuresis, [573](#)  
Diuretic, [2117](#)  
Diuretic Drug, [573](#)  
Diuretic Resistance, [573](#)  
Diurnal, [573](#)  
Diverticula, [573](#)  
Diverticular Bleeding, [573](#)  
Diverticular Disease, [573](#)  
Diverticulitis, [574](#)  
Diverticulosis, [574](#)  
Diverticulosis And Diverticulitis, [574](#)  
Diverticulum, [574](#)  
Divisum, Pancreas, [1452](#)  
Dix-Hallpike Maneuver, [574](#)  
Dizygotic Twin, [574](#)  
Dizygotic Twins, [574](#)  
Dizziness, [574](#)  
Dizzy, [574](#)  
DKD, [574](#)  
DLCO, [574](#)  
DLI, [574](#)  
DM, [575](#)  
DM2, [575](#)  
DMD, [575](#)  
DMFT Index, [575](#)  
DMG, [575](#)  
DMSA Scan, [575](#)  
DNA, [575](#)  
DNA Adduct, [575](#)  
DNA Alkylation, [575](#)  
DNA Amplification, [575](#)  
DNA Binding, [575](#)  
DNA Damage, [575](#)  
DNA Demethylation, [575](#)  
DNA Double Helix, [575](#)  
DNA Fragmentation, [576](#)  
DNA Gyrase, [576](#)  
DNA Helicase, [576](#)  
DNA Integration, [576](#)  
DNA Ligases, [576](#)  
DNA Ligation, [576](#)  
DNA Methylation, [576](#)  
DNA Packaging, [576](#)  
DNA Polymerase, [576](#)  
DNA Primase, [576](#)  
DNA Primers, [576](#)  
DNA Probe, [576](#)  
DNA Profiling, [576](#)  
DNA Repair, [576](#)  
DNA Repair Mechanisms, [577](#)  
DNA Replication, [577](#)  
DNA Sequencing, [577](#)  
DNA Virus, [577](#)  
Do Not Resuscitate, [577](#)  
Do-Not-Resuscitate Order, [577](#)  
DOACs, [577](#)  
DOB, [577](#)  
Dobutamine, [577](#)  
Dobutamine Stress Echocardiography, [577](#)  
Doc, [577](#)  
Docetaxel, [577](#)  
Doconexent, [577](#)  
Docosadienoic Acid, [577](#)  
Docosahexaenoic Acid, [578](#)  
Docosapentaenoic Acid, [578](#)  
Docosatrienoic Acid, [578](#)  
Doctor Of Osteopathic Medicine, [2117](#)  
Docusate, [578](#)  
Dog Allergy, [578](#)  
Dog Bite, [578](#)  
Dolasetron, [578](#)  
Dolasetron Mesylate, [578](#)  
Dolichocephalic, [578](#)  
Dolichol, [578](#)  
Dolichostenomelia, [578](#)  
Doll Eye Reflex, [578](#)  
Doll's Eye Sign, [578](#)  
Dolor, [579](#)  
Domain, [579](#)  
Domestic Violence, [579](#)  
Dominant, [579](#)  
Domperidone, [579](#)  
Donath-Landsteiner Antibody, [579](#)  
Donath-Landsteiner Test, [579](#)  
Donepezil, [579](#)  
Donnan Equilibrium, [579](#)  
Donohue Syndrome, [579](#)  
Donor, [579](#)  
Donor Insemination, [579](#)  
Donor Suitability Assessment, [579](#)  
Donor Testing, [580](#)  
Donovan Bodies, [580](#)  
Donovanosis, [580](#)  
Dopa, [580](#)  
Dopa-Responsive Dystonia, [580](#)  
Dopamine, [580](#)  
Dopamine Agonist, [580](#)  
Dopamine Agonists, [580](#)  
Dopamine Antagonist, [580](#)  
Dopamine Receptor, [580](#)  
Dopamine Transporter, [580](#)  
Dopaminergic Neuron, [580](#)  
Dopaminergic Pathways, [580](#)  
Doppler Echocardiography, [581](#)  
Doppler Ultrasound, [581](#)  
Doppler Ultrasound of a Limb, [581](#)  
Doppler Velocimetry, [581](#)  
Dor Procedure, [581](#)

---



---

Doravirine, [581](#)  
Dorea, [581](#)  
Doripenem, [581](#)  
Dormant, [581](#)  
Dorsal, [581](#)  
Dorsal Blocking Splint, [581](#)  
Dorsal Body Cavity, [581](#)  
Dorsal Column-Medial Lemniscal Pathway, [581](#)  
Dorsal Funiculus, [582](#)  
Dorsal Kyphosis, [582](#)  
Dorsal Root, [582](#)  
Dorsal Root Ganglion, [582](#)  
Dorsalis Pedis Artery, [582](#)  
Dorsiflexion, [582](#)  
Dorsocervical Fat Pad, [582](#)  
Dorsolateral, [582](#)  
Dorsum, [582](#)  
Dorzolamide, [582](#)  
Dosage, [582](#)  
Dosage Forms, [582](#)  
Dose, [582](#)  
Dose Rate, [582](#)  
Dose Titration, [582](#)  
Dose-Limiting Toxicity, [583](#)  
Dose-Response Curve, [583](#)  
Dose-Response Relationship, [583](#)  
Dosimeter, [583](#)  
Dosimetry, [583](#)  
DOTATATE, [583](#)  
DOTS, [583](#)  
Double Aortic Arch, [583](#)  
Double Blind, [583](#)  
Double Blind Trial, [583](#)  
Double Bubble Sign, [583](#)  
Double Inlet Left Ventricle, [583](#)  
Double Outlet Right Ventricle, [583](#)  
Double Uterus, [584](#)  
Double Vision, [584](#)  
Double-Balloon Enteroscopy, [584](#)  
Double-Blind Method, [584](#)  
Double-Blind Study, [584](#)  
Double-Contrast Barium Enema, [584](#)  
Double-Outlet Right Ventricle, [584](#)  
Douche, [584](#)  
Dowager Hump, [584](#)  
Dowager'S Hump, [584](#)  
Dowling-Meara Syndrome, [584](#)  
Down Syndrome, [584](#)  
Downregulation, [585](#)  
Doxapram, [585](#)  
Doxazosin, [585](#)  
Doxazosin Mesylate, [585](#)  
Doxepin, [585](#)  
Doxepin Overdose, [585](#)  
Doxercalciferol, [585](#)  
Doxorubicin, [585](#)  
Doxorubicin Hydrochloride, [585](#)  
Doxycycline, [585](#)  
Doxycycline Hyclate, [585](#)  
Doxylamine, [585](#)  
DPP-4 Inhibitor, [585](#)  
DPP-4 Inhibitors, [585](#)  
DPT Immunization, [586](#)  
DR, [586](#)  
Dracunculiasis, [586](#)  
Dracunculosis, [586](#)  
Dracunculus Medinensis, [586](#)  
Draf Procedure, [586](#)  
Drain, [586](#)  
Drain Opener Poisoning, [586](#)  
Drainage, [586](#)  
Drainage Procedures, [586](#)  
Drainpipe Cleaners, [586](#)  
Drawer Test, [586](#)  
Dream, [586](#)  
Dreams, [587](#)  
Drepanocytosis, [587](#)  
DRESS Syndrome, [587](#)  
Dressing, [587](#)  
Dressler Syndrome, [587](#)  
Dressler's Syndrome, [587](#)  
DRI, [587](#)  
Dried Specimen, [587](#)  
Drinking Water, [587](#)  
Drip, [587](#)  
Driving And Older Adults, [587](#)  
Driving Evaluation, [587](#)  
Driving Rehabilitation, [587](#)  
Dronabinol, [588](#)  
Drool, [588](#)  
Drooling, [588](#)  
Drooping Eyelids, [588](#)  
Drop Attack, [588](#)  
Drop Foot, [588](#)  
Drop Hand Syndrome, [588](#)  
Droperidol, [588](#)  
Droplet Precautions, [588](#)  
Dropper Bottle, [588](#)  
Drowning, [588](#)  
Drowning Signs, [588](#)  
Drowsiness, [588](#)  
Drug, [588](#)  
Drug Abuse, Teen, [46](#)  
Drug Administration Route, [2117](#)  
Drug Allergies, [2117](#)  
Drug Allergy, [2117](#)  
Drug Antagonism, [2117](#)  
Drug Bioavailability, [2117](#)

---

---

Drug Caution Code, 589  
Drug Classification Systems, 2117  
Drug Clearance, 2118  
Drug Delivery, 589  
Drug Dependence, 2118  
Drug Desensitization, 589  
Drug Discovery, 589  
Drug Emulsion, 2118  
Drug Eruption, 2118  
Drug Eruptions, 589  
Drug Hypersensitivity, 589  
Drug Hypersensitivity Reactions, 2118  
Drug Hypersensitivity Syndrome, 2118  
Drug Implant, 589  
Drug Implants, 589  
Drug Induced Liver Disease, 2118  
Drug Interaction, 2119  
Drug Metabolism, 2119  
Drug Metabolism Pharmacogenomics, 2119  
Drug Overdose, 2119  
Drug Packaging, 2119  
Drug Repurposing, 589  
Drug Resistance, 2119  
Drug Safety, 589  
Drug Screening, 589  
Drug Testing, 589  
Drug Therapeutics, 2119  
Drug Tolerance, 2119  
Drug Toxicity, 2119  
Drug Use First Aid, 2119  
Drug Vehicle, 2120  
Drug Withdrawal, 2120  
Drug, Prescription, 2119  
Drug, Sulfa, 2119  
Drug- and Nephrotoxin-Associated Kidney Disorders, 590  
Drug-Drug Interactions, 2118  
Drug-Eluting Balloon (Vascular), 2118  
Drug-Eluting Stent, 590  
Drug-Facilitated Sexual Assault, 2118  
Drug-Food Interaction, 590  
Drug-Induced Autoimmunity, 2118  
Drug-Induced Bullous Disorders, 590  
Drug-Induced Diarrhea, 2118  
Drug-Induced Gingival Hyperplasia, 590  
Drug-Induced Glaucoma, 590  
Drug-Induced Hepatotoxicity, 590  
Drug-Induced Immune Hemolytic Anemia, 2118  
Drug-Induced Liver Injury, 590  
Drug-Induced Low Blood Sugar, 2119  
Drug-Induced Lupus, 590  
Drug-Induced Lupus Erythematosus, 590  
Drug-Induced Pemphigus, 591  
Drug-Induced Photosensitivity, 591  
Drug-Induced Pigmentation, 591  
Drug-Induced Pulmonary Disease, 2119  
Drug-Induced Pulmonary Toxicity, 591  
Drug-Induced Thrombocytopenia, 2119  
Drug-Induced Tremor, 2119  
Drugs That May Cause Erection Problems, 591  
Drusen, 591  
Dry Age-Related Macular Degeneration, 591  
Dry AMD, 591  
Dry Beriberi, 591  
Dry Cell Battery Poisoning, 591  
Dry Cough, 591  
Dry Cupping, 591  
Dry Drowning, 591  
Dry Eye, 591  
Dry Eye Disease, 592  
Dry Eye Syndrome, 592  
Dry Gangrene, 592  
Dry Hair, 592  
Dry Ice, 592  
Dry Macular Degeneration, 592  
Dry Mouth, 592  
Dry Mouth Xerostomia, 592  
Dry Powder Inhaler, 592  
Dry Skin, 592  
Dry Skin and Itching, 592  
Dry Socket, 593  
Dry-Powder Inhaler, 593  
Dryness, Vaginal, 2017  
DSM, 593  
DSPS, 593  
DSRCT, 593  
DSWPD, 593  
DTaP, 593  
DTaP Vaccine, 2120  
Dual Antiplatelet Therapy, 593  
Dual Energy X-Ray Absorptiometry, 593  
Dual-Energy CT, 593  
Dual-Energy X-Ray Absorptiometry, 593  
Dual-Photon Absorptiometry, 593  
Duane Retraction Syndrome, 593  
Dubin-Johnson Syndrome, 593  
Dubowitz Syndrome, 593  
Duchenne Muscular Dystrophy, 594  
Duck Embryo Vaccine, 594  
Duct, 594  
Duct Ectasia, 594  
Ductal Carcinoma, 594  
Ductal Carcinoma In Situ, 594  
Ductal-Dependent Lesions, 594  
Ductless Glands, 594  
Ductus Arteriosus, 594  
Ductus Venosus, 594  
Duffy-Null Associated Neutrophil Count, 594

---

---

Dugas Sign, [594](#)  
 Duke Criteria, [594](#)  
 Dukes Classification, [595](#)  
 Dull Pain, [595](#)  
 Duloxetine, [595](#)  
 Duloxetine Hydrochloride, [595](#)  
 Dumbbell Ossification, [595](#)  
 Dumdum Fever, [595](#)  
 Dumping Syndrome, [595](#)  
 Dunbar Syndrome, [595](#)  
 Dunphy Sign, [595](#)  
 Duodenal, [595](#)  
 Duodenal Adenocarcinoma, [595](#)  
 Duodenal Atresia, [595](#)  
 Duodenal Bulb, [595](#)  
 Duodenal Disease, [596](#)  
 Duodenal Diverticulum, [596](#)  
 Duodenal Fistula, [596](#)  
 Duodenal Hemorrhage, [596](#)  
 Duodenal Infection, [596](#)  
 Duodenal Neoplasm, [596](#)  
 Duodenal Obstruction, [596](#)  
 Duodenal Papilla, [596](#)  
 Duodenal Perforation, [596](#)  
 Duodenal Stenosis, [596](#)  
 Duodenal Tissue Culture, [596](#)  
 Duodenal Ulcer, [596](#)  
 Duodenitis, [596](#)  
 Duodenoscopy, [597](#)  
 Duodenostomy, [597](#)  
 Duodenum, [597](#)  
 Duplex Doppler Ultrasound Scanning, [597](#)  
 Duplex Surveillance, [597](#)  
 Duplex Ultrasound, [597](#)  
 Duplication, [597](#)  
 Dupuytren Contracture, [597](#)  
 Dupuytren's Disease, [597](#)  
 Dupuytren, Guillaume, [866](#)  
 Dupuytren's Disease, [597](#)  
 Dura, [597](#)  
 Dura Mater, [597](#)  
 Durable Power Of Attorney, [597](#)  
 Dural Arteriovenous Fistula, [598](#)  
 Dural Arteriovenous Fistulas, [598](#)  
 Dural Sac, [598](#)  
 Dural Venous Sinuses, [598](#)  
 Duramorph, [598](#)  
 Durapatite, [598](#)  
 Duroziez Sign, [598](#)  
 Durules, [598](#)  
 Dust, [598](#)  
 Dust Disease, [598](#)  
 Dust Mite Allergy, [598](#)  
 Dutasteride, [598](#)  
 DVT, [599](#)  
 DVT Prevention, [599](#)  
 Dwarfism, [599](#)  
 DWI FLAIR Mismatch, [599](#)  
 Dx, [599](#)  
 DXA Scan, [599](#)  
 Dyad, [599](#)  
 Dye, [599](#)  
 Dye Lasers, [599](#)  
 Dye Remover Poisoning, [599](#)  
 Dying, [599](#)  
 Dynamin Gtpase, [599](#)  
 Dynamometer, [599](#)  
 Dynein Atpase, [599](#)  
 Dynorphin, [600](#)  
 Dyphylline, [600](#)  
 Dysaesthesia, [600](#)  
 Dysania, [600](#)  
 Dysarthria, [600](#)  
 Dysautonomia, [600](#)  
 Dysbarism, [600](#)  
 Dysbiosis, [600](#)  
 Dyscalculia, [600](#)  
 Dyschezia, [600](#)  
 Dyscrasia, [600](#)  
 Dyscrasias, [600](#)  
 Dysdiadochokinesia, [600](#)  
 Dysdiadochokinesis, [601](#)  
 Dysembryoplastic Neuroepithelial Tumor, [601](#)  
 Dysentery, [601](#)  
 Dysesthesia, [601](#)  
 Dysferlinopathy, [601](#)  
 Dysfibrinogenemia, [601](#)  
 Dysfunction, [601](#)  
 Dysfunctional Uterine Bleeding, [601](#)  
 Dysfunctional Voiding, [601](#)  
 Dysgammaglobulinemia, [601](#)  
 Dysgerminoma, [601](#)  
 Dysgeusia, [601](#)  
 Dysgraphia, [601](#)  
 Dyshidrosis, [601](#)  
 Dyshidrotic Eczema, [602](#)  
 Dyskeratosis Congenita, [602](#)  
 Dyskinesia, [602](#)  
 Dyslexia, [602](#)  
 Dyslipidemia, [602](#)  
 Dyslipidemias, [602](#)  
 Dysmenorrhea, [602](#)  
 Dysmenorrhoea, [602](#)  
 Dysmetria, [602](#)  
 Dysmyelogenic Leukodystrophy, [602](#)  
 Dysorexia, [602](#)  
 Dysostoses, [603](#)  
 Dyspareunia, [603](#)

---

---

Dyspepsia, [603](#)  
 Dyspeptic, [603](#)  
 Dysphagia, [603](#)  
 Dysphagia Diet, [603](#)  
 Dysphagia Lusoria, [603](#)  
 Dysphagia Rehabilitation, [603](#)  
 Dysphasia, [603](#)  
 Dysphonia, [603](#)  
 Dysphonia Spastica, [604](#)  
 Dysphoria, [604](#)  
 Dysplasia, [604](#)  
 Dysplasia Epiphysealis Hemimelica (Trevor Disease), [604](#)  
 Dysplasia, Cervical, [362](#)  
 Dysplastic Melanocytic Nevi Pathology, [604](#)  
 Dysplastic Nevi, [604](#)  
 Dysplastic Nevus, [604](#)  
 Dyspnea, [604](#)  
 Dyspnea On Exertion, [604](#)  
 Dyspnoea, [604](#)  
 Dyspraxia, [604](#)  
 Dysraphism, [604](#)  
 Dysrhythmia, [604](#)  
 Dyssomnia, [604](#)  
 Dyssomnias, [605](#)  
 Dysthyroid Eye Disease, [605](#)  
 Dystocia, [605](#)  
 Dystonia, [605](#)  
 Dystonia Disorder, [605](#)  
 Dystonia, Cervical, [362](#)  
 Dystonin, [605](#)  
 Dystrophia Unguium, [605](#)  
 Dystrophic Calcification, [605](#)  
 Dystrophin, [605](#)  
 Dystrophin-Glycoprotein Complex, [606](#)  
 Dystrophy, [606](#)  
 Dysuria, [606](#)  
 Déjà Vu, [606](#)  
  
 E Coli Enteritis, [607](#)  
 E Coli Infection, [607](#)  
 E-Cigarettes And E-Hookahs, [607](#)  
 E-D Syndrome, [607](#)  
 E. Coli, [607](#)  
 E. coli Infection, [607](#)  
 E. Coli O157:H7, [607](#)  
 E. Coli, Shiga Toxin-Producing (STEC), [1780](#)  
 Eagle Syndrome, [607](#)  
 Eales Disease, [607](#)  
 Ear, [607](#)  
 Ear Barotrauma, [607](#)  
 Ear Canal, [608](#)  
 Ear Cartilage, [608](#)  
 Ear Disease, [608](#)  
 Ear Diseases, [608](#)  
 Ear Disorders, [608](#)  
 Ear Drainage Culture, [608](#)  
 Ear Examination, [608](#)  
 Ear Foreign Body, [608](#)  
 Ear Infection, [608](#)  
 Ear Infection, Outer Ear, [1431](#)  
 Ear Microbiome, [608](#)  
 Ear Neoplasm, [608](#)  
 Ear Speculum, [608](#)  
 Ear Tag, [608](#)  
 Ear Thermometer, [608](#)  
 Ear Tube Insertion, [609](#)  
 Ear Wax, [609](#)  
 Ear, Diseases Of, [608](#)  
 Ear, Nose, And Throat, [1432](#)  
 Earache, [609](#)  
 Eardrum, [609](#)  
 Eardrum Repair, [609](#)  
 Eardrum, Ruptured, [1977](#)  
 Earliest Signs Of Increased Intracranial Pressure, [609](#)  
 Earlobes, [609](#)  
 Early Afterdepolarization, [609](#)  
 Early Childhood Tooth Decay, [609](#)  
 Early Decelerations, [609](#)  
 Early Detection, [609](#)  
 Early Diagnosis, [609](#)  
 Early Endosome, [609](#)  
 Early Fetal Ultrasound, Deciding About, [736](#)  
 Early Gastric Cancer, [610](#)  
 Early Infantile Epileptic Encephalopathy, [610](#)  
 Early Intervention, [610](#)  
 Early Pregnancy Loss, [610](#)  
 Early Puberty, [610](#)  
 Early Repolarization Pattern, [610](#)  
 Early Repolarization Syndrome, [610](#)  
 Early Satiety, [610](#)  
 Early-Onset Dementia, [1646](#)  
 Early-Onset Tremor, [1646](#)  
 Early-Stage Prostate Cancer, [610](#)  
 Earwax, [610](#)  
 Earwax Blockage, [610](#)  
 Earwax Removal, [610](#)  
 Eating Disorder, [610](#)  
 Eating Disorder, Pica Type, [1530](#)  
 Eating Disorders, [611](#)  
 Eating Disorders, Anorexia, [96](#)  
 Eating Disorders, Binge Eating, [207](#)  
 Eating Disorders, Bulimia, [281](#)  
 Eaton-Lambert Syndrome, [611](#)  
 EB, [611](#)  
 EBCT, [611](#)  
 Ebola, [611](#)  
 Ebola Hemorrhagic Fever, [611](#)  
 Ebola Virus, [611](#)

---

---

Ebola Virus Disease, [611](#)  
 Ebola Virus Infection, [611](#)  
 Ebolavirus, [611](#)  
 Ebstein Anomaly, [611](#)  
 Ebstein's Anomaly, [611](#)  
 EBUS-TBNA, [611](#)  
 Ecbolics, [612](#)  
 Eccentric, [612](#)  
 Eccentric Action, [612](#)  
 Eccentrocyte Count, [612](#)  
 Ecchymosis, [612](#)  
 Eccrine Carcinoma, [612](#)  
 Eccrine Gland, [612](#)  
 Eccrine Glands, [612](#)  
 Eccrine Porocarcinoma, [612](#)  
 Eccrine Poroma, [612](#)  
 ECG, [612](#)  
 ECG/EKG Machine, [612](#)  
 Echinacea, [613](#)  
 Echinocandins, [613](#)  
 Echinococcal Cysts, [613](#)  
 Echinococcosis, [613](#)  
 Echinococcus, [613](#)  
 Echinococcus Granulosus, [613](#)  
 Echinococcus Multilocularis, [613](#)  
 Echinococcus Vogeli, [613](#)  
 Echinocytes, [613](#)  
 Echo, [613](#)  
 Echocardiogram, [613](#)  
 Echocardiography, [613](#)  
 Echoencephalography, [613](#)  
 Echogenic Bowel, [613](#)  
 Echogenic Intracardiac Focus, [614](#)  
 Echography, [614](#)  
 Echolalia, [614](#)  
 Echophenomena, [614](#)  
 Echopraxia, [614](#)  
 Echovirus, [614](#)  
 Eclampsia, [614](#)  
 Eclampsia in Pregnancy, [614](#)  
 ECMO, [614](#)  
 ECOG Performance Status, [614](#)  
 Econazole, [614](#)  
 Economy Class Syndrome, [614](#)  
 Ecstasy, [614](#)  
 Ectasia, [614](#)  
 Ecthyma, [614](#)  
 Ecthyma Gangrenosum, [615](#)  
 Ectoderm, [615](#)  
 Ectodermal Dysplasia, [615](#)  
 Ectodermal Dysplasia, Rapp-Hodgkin Type, [1669](#)  
 Ectodermal Dysplasias, [615](#)  
 Ectodermosis Erosiva Pluriorificialis, [615](#)  
 Ectomesenchymoma, [615](#)  
 Ectoparasite, [615](#)  
 Ectopia Cordis, [615](#)  
 Ectopia Lentis, [615](#)  
 Ectopic, [615](#)  
 Ectopic ACTH Syndrome, [615](#)  
 Ectopic Beat, [615](#)  
 Ectopic Cushing Syndrome, [615](#)  
 Ectopic Heartbeat, [615](#)  
 Ectopic Heartbeats, [616](#)  
 Ectopic Kidney, [616](#)  
 Ectopic Pacemaker, [616](#)  
 Ectopic Pregnancy, [616](#)  
 Ectopic Thyroid, [616](#)  
 Ectopic Thyroid Tissue, [616](#)  
 Ectopic Tissue, [616](#)  
 Ectopic Ureter, [616](#)  
 Ectopion, [616](#)  
 Ectrodactyly, [616](#)  
 Ectromelia, [616](#)  
 Ectropion, [616](#)  
 Ectropion (Drooping Eyelids), [616](#)  
 Eculizumab, [617](#)  
 Eczema, [617](#)  
 Eczema Herpeticum, [617](#)  
 Eczema, Atopic, [142](#)  
 Eczematous Dermatitis, [617](#)  
 ED, [617](#)  
 Edema, [617](#)  
 Edema Trunk, [617](#)  
 Edema, Idiopathic, [707](#)  
 Edema, Pulmonary, [1630](#)  
 Edentulism, [617](#)  
 Edentulous, [617](#)  
 Edentulousness, [617](#)  
 Edetic Acid, [617](#)  
 EDMD, [617](#)  
 EDTA, [617](#)  
 Edward's Syndrome, [618](#)  
 Edwards Syndrome, [618](#)  
 Edwardsiella, [618](#)  
 Edwardsiella Tarda, [618](#)  
 EEG, [618](#)  
 EFAST Examination, [618](#)  
 Efavirenz, [618](#)  
 Effacement, [618](#)  
 Effective Dose, [618](#)  
 Effective Half-Life, [618](#)  
 Effective Reproduction Number, [618](#)  
 Effectiveness, [618](#)  
 Effector, [618](#)  
 Efferent, [618](#)  
 Efferent Arteriole, [619](#)  
 Efferent Ductules, [619](#)  
 Efferent Lymphatic Vessel, [619](#)

---

---

Efferent Nerve, [619](#)  
Efferent Neuron, [619](#)  
Efferent Pathway, [619](#)  
Effervescent Tablets, [619](#)  
Efficacy, [619](#)  
Effort Angina, [619](#)  
Effort Syncope, [619](#)  
Effusion, [619](#)  
Effusion, Pleural, [1546](#)  
Effusive-Constrictive Pericarditis, [619](#)  
Eflornithine, [2120](#)  
EGD, [619](#)  
EGFR, [620](#)  
EGFR Inhibition, [620](#)  
Egg, [620](#)  
Egg Allergy, [620](#)  
Egg Shell, [620](#)  
Egg-Cell, [620](#)  
Eggerthella, [620](#)  
Eggshell Nail, [620](#)  
Egophony, [620](#)  
EHBA, [620](#)  
Ehlers-Danlos Syndrome, [620](#)  
EHR, [620](#)  
Ehrlichia, [620](#)  
Ehrlichia And Anaplasma, [621](#)  
Ehrlichiosis, [621](#)  
Ehrlichiosis And Anaplasmosis, [621](#)  
Eicosanoid, [621](#)  
Eicosanoid Degradation, [621](#)  
Eicosanoid Metabolism, [621](#)  
Eicosanoid Receptor, [621](#)  
Eicosapentaenoic Acid, [621](#)  
Eicosatetraenoic Acid, [621](#)  
Eighth Cranial Nerve, [621](#)  
Eikenella, [621](#)  
Eikenella Corrodens, [621](#)  
Eisenmenger Complex, [621](#)  
Eisenmenger syndrome, [621](#)  
Eisenmenger's Syndrome, [622](#)  
Ejaculation, [622](#)  
Ejaculatory Duct, [622](#)  
Ejaculatory Ducts, [622](#)  
Ejaculatory Dysfunction, [622](#)  
Ejection Click, [622](#)  
Ejection Fraction, [622](#)  
Ejection Murmur, [622](#)  
EKG, [622](#)  
Ekman-Lobstein Disease, [622](#)  
Elastance, [622](#)  
Elastic Cartilage, [622](#)  
Elastic Fiber, [622](#)  
Elastic Skin, [622](#)  
Elastic Tissue, [623](#)  
Elasticity, [623](#)  
Elastics, [623](#)  
Elastin, [623](#)  
Elastofibroma, [623](#)  
Elastography, [623](#)  
Elastomer, [623](#)  
Elastosis, [623](#)  
Elastosis Perforans Serpiginosa, [623](#)  
Elbow, [623](#)  
Elbow (Radial Head or Neck) Fracture, [623](#)  
Elbow Dislocation, [623](#)  
Elbow Extensor, [623](#)  
Elbow Flexors, [624](#)  
Elbow Joint, [624](#)  
Elbow Pain, [624](#)  
Elbow Replacement, [624](#)  
Elder Abuse, [624](#)  
Elective, [624](#)  
Elective Abortion, [624](#)  
Elective Surgery, [624](#)  
Electric Axis of the Heart, [624](#)  
Electric Impedance, [624](#)  
Electric Shock, [624](#)  
Electrical Alternans, [624](#)  
Electrical Cardioversion, [625](#)  
Electrical Conductance, [625](#)  
Electrical Injury, [625](#)  
Electrical Stimulation, [625](#)  
Electrical Synapse, [625](#)  
Electrocardiogram, [625](#)  
Electrocardiographic Gating, [625](#)  
Electrocardiography, [625](#)  
Electrocauterization, [625](#)  
Electrocautery, [625](#)  
Electrocoagulation, [625](#)  
Electroconvulsive Therapy, [625](#)  
Electrocution, [625](#)  
Electrode, [626](#)  
Electrodessication, [626](#)  
Electroencephalogram, [626](#)  
Electroencephalography, [626](#)  
Electrogastrogram, [626](#)  
Electrolysis, [626](#)  
Electrolyte, [626](#)  
Electrolyte Balance, [626](#)  
Electrolyte Imbalance, [626](#)  
Electrolyte Panel, [626](#)  
Electrolytes, [626](#)  
Electromagnetic Navigation Bronchoscopy, [626](#)  
Electromechanical Dissociation, [627](#)  
Electromyography, [627](#)  
Electron Microscope, [627](#)  
Electron Transport Chain, [627](#)  
Electron-Beam Computed Tomography, [627](#)

---



---

Electronic Cigarette, [627](#)  
Electronic Fetal Monitoring, [627](#)  
Electronystagmography, [627](#)  
Electrophoresis, [627](#)  
Electrophysiologic Testing, [627](#)  
Electrophysiology, [627](#)  
Electrophysiology Study, [627](#)  
Electroretinography, [627](#)  
Electrosurgery, [627](#)  
Electrosurgical Unit, [628](#)  
Elejalde Syndrome, [628](#)  
Element, [628](#)  
Elephant Ear Poisoning, [628](#)  
Elephantiasis, [628](#)  
Elevated Blood Pressure, [628](#)  
Elevated Hemidiaphragm, [628](#)  
Elevated Liver Enzymes, [628](#)  
Elevation, [628](#)  
Elevator, [628](#)  
Eligible, [628](#)  
Elimination, [628](#)  
Elimination Diets, [628](#)  
Elimination Disorders, [629](#)  
Elimination Half-Life, [629](#)  
Elimination Patterns, [629](#)  
ELISA Blood Test, [629](#)  
Elixir, [629](#)  
Elizabethkingia, [629](#)  
Elliptocyte Count, [629](#)  
Elliptocytosis, [629](#)  
Ellis-Van Creveld Syndrome, [629](#)  
Ellsworth-Howard Test, [629](#)  
Elongin, [629](#)  
Elotuzumab, [629](#)  
Eltrombopag, [630](#)  
Eltrombopag Olamine, [630](#)  
Emaciation, [630](#)  
Embolectomy, [630](#)  
Emboli, [630](#)  
Embolic Stroke, [630](#)  
Embolic Stroke of Undetermined Source, [630](#)  
Embolism, [630](#)  
Embolism, Pulmonary, [1630](#)  
Embolization, [630](#)  
Embolization Therapy, [630](#)  
Embolus, [630](#)  
Embryo, [630](#)  
Embryo Neoplasm, [630](#)  
Embryo Transfer, [631](#)  
Embryogenesis, [631](#)  
Embryology, [631](#)  
Embryoma, [631](#)  
Embryonal Carcinoma, [631](#)  
Embryonal CNS Tumor, [631](#)  
Embryonal Rhabdomyosarcoma, [631](#)  
Embryonal Tumors, [631](#)  
Embryonic, [631](#)  
Embryonic Cell, [631](#)  
Embryonic Induction, [631](#)  
Embryonic Mosaic, [631](#)  
Embryonic Stage, [631](#)  
Emergence Agitation, [631](#)  
Emergency Airway Puncture, [631](#)  
Emergency Care, [632](#)  
Emergency Code, [632](#)  
Emergency Contraception, [632](#)  
Emergency Department, [632](#)  
Emergency Medicine, [632](#)  
Emergency Procedures, [632](#)  
Emerging Infectious Disease, [632](#)  
Emery-Dreifuss Muscular Dystrophy, [632](#)  
Emesis, [632](#)  
Emetic, [632](#)  
Emetics, [632](#)  
EMG, [632](#)  
Eminentia, [632](#)  
Emission, [633](#)  
Emmenagogue, [633](#)  
Emmetropia, [633](#)  
Emodin, [633](#)  
Emollient, [633](#)  
Emotion, [633](#)  
Emotional Adjustment, [633](#)  
Emotional Dependency, [633](#)  
Emotional Disorders, [633](#)  
Emotional Dysregulation Disorder, [633](#)  
Emotional Eating, [633](#)  
Empathy, [633](#)  
Emphysema, [633](#)  
Emphysematous Aortitis, [633](#)  
Emphysematous Cholecystitis, [634](#)  
Emphysematous Hepatitis, [634](#)  
Emphysematous Prostatitis, [634](#)  
Emphysematous Pyelonephritis, [634](#)  
Empiric Antibiotic Therapy, [634](#)  
Empiric Risk, [634](#)  
Empirical, [634](#)  
Empirical Treatment, [634](#)  
Employee, [634](#)  
Empty Calories, [634](#)  
Empty Can Test, [634](#)  
Empty Nose Syndrome, [634](#)  
Empty Sella Syndrome, [634](#)  
Empyema, [635](#)  
Empyema Necessitans, [635](#)  
Empyema Necessitatis, [635](#)  
EMR, [635](#)  
Emtricitabine, [635](#)

---

---

Emulsion, [635](#)  
Enalapril, [635](#)  
Enalaprilat, [635](#)  
Enamel, [635](#)  
Enamel Hypoplasia, [635](#)  
Enamel Organ, [635](#)  
Enamel Pearl, [635](#)  
Enanthem, [635](#)  
Enanthema, [636](#)  
Enarthrosis, [636](#)  
Encapsulated, [636](#)  
Encapsulated Peritoneal Sclerosis, [636](#)  
Encapsulating Peritoneal Sclerosis, [636](#)  
Encephalitis, [636](#)  
Encephalitis Lethargica, [636](#)  
Encephalitis Virus, [636](#)  
Encephalitozoon, [636](#)  
Encephalocele, [636](#)  
Encephalofacial Angiomatosis, [636](#)  
Encephalomalacia, [636](#)  
Encephalomyelitis, [636](#)  
Encephalopathy, [636](#)  
Encephalotrigeminal Angiomatosis, [637](#)  
Enchondroma, [637](#)  
Enchondromatosis, [637](#)  
Encoding, [637](#)  
Encopresis, [637](#)  
End Point, [637](#)  
End Stage, [637](#)  
End-Of-Life Care, [637](#)  
End-Stage Kidney Disease, [637](#)  
End-Stage Renal Disease, [637](#)  
Endangiitis, [637](#)  
Endarterectomy, [637](#)  
Endarteritis, [637](#)  
Endemic, [638](#)  
Endemic Diseases, [638](#)  
Endemic Fungi, [638](#)  
Endemic Goiter, [638](#)  
Endergonic, [638](#)  
Ending Pregnancy With Medicines, [638](#)  
Endobronchial Foreign Body, [638](#)  
Endobronchial Tumor, [638](#)  
Endobronchial Ultrasound, [638](#)  
Endobronchial Valve, [638](#)  
Endocannabinoid, [638](#)  
Endocardial Cushion Defect, [638](#)  
Endocardial Fibroelastosis, [638](#)  
Endocarditis, [639](#)  
Endocarditis Prophylaxis, [639](#)  
Endocarditis Surgical Indications, [639](#)  
Endocardium, [639](#)  
Endocervical, [639](#)  
Endocervical Culture, [639](#)  
Endocervical Gram Stain, [639](#)  
Endocrine, [639](#)  
Endocrine Disease, [639](#)  
Endocrine Disorders, [639](#)  
Endocrine Disrupting Chemicals, [639](#)  
Endocrine Disruptor, [639](#)  
Endocrine Finding, [639](#)  
Endocrine Function, [639](#)  
Endocrine Gland, [640](#)  
Endocrine Glands, [640](#)  
Endocrine Hypertension, [640](#)  
Endocrine Myopathies, [640](#)  
Endocrine Neoplasm, [640](#)  
Endocrine Pancreas, [640](#)  
Endocrine System, [640](#)  
Endocrine Testing, [640](#)  
Endocrine Therapy, [640](#)  
Endocrine Tissues, [640](#)  
Endocrinologist, [640](#)  
Endocrinologists, [640](#)  
Endocrinology, [640](#)  
Endocrinopathy, [640](#)  
Endocytosis, [641](#)  
Endoderm, [641](#)  
Endodontics, [641](#)  
Endodontist, [641](#)  
Endogenous, [641](#)  
Endogenous Cushing Syndrome, [641](#)  
Endogenous Opiates, [641](#)  
Endogenous Opioids, [641](#)  
Endogenous Retrovirus, [641](#)  
Endoleak, [641](#)  
Endolimax, [641](#)  
Endolimax Nana, [641](#)  
Endolymph, [641](#)  
Endolymphatic Hydrops, [641](#)  
Endometrial, [642](#)  
Endometrial Ablation, [642](#)  
Endometrial Adenocarcinoma, [642](#)  
Endometrial Biopsy, [642](#)  
Endometrial Cancer, [642](#)  
Endometrial Carcinoma, [642](#)  
Endometrial Disorder, [642](#)  
Endometrial Hyperplasia, [642](#)  
Endometrial Implants, [642](#)  
Endometrial Intraepithelial Neoplasia, [642](#)  
Endometrial Neoplasm, [642](#)  
Endometrial Polyp, [642](#)  
Endometrial Polyps, [643](#)  
Endometrial Receptivity Analysis, [643](#)  
Endometrial Stromal Nodule, [643](#)  
Endometrial Stromal Sarcoma, [643](#)  
Endometrioid Carcinoma Of The Vulva, [643](#)  
Endometrioid Ovarian Carcinoma, [643](#)

---

---

Endometrioma, [643](#)  
 Endometriosis, [643](#)  
 Endometritis, [643](#)  
 Endometrium, [643](#)  
 Endomyocardial Biopsy, [643](#)  
 Endomyocardial Fibrosis, [644](#)  
 Endomysium, [644](#)  
 Endoneurium, [644](#)  
 Endoparasite, [644](#)  
 Endopeptidase, [644](#)  
 Endophthalmitis, [644](#)  
 Endoplasmic Reticulum, [644](#)  
 Endoprosthesis, [644](#)  
 Endoribonuclease, [644](#)  
 Endorphin, [644](#)  
 Endorphins, [644](#)  
 Endoscope, [644](#)  
 Endoscopic Endonasal Surgery, [644](#)  
 Endoscopic Mucosal Resection, [644](#)  
 Endoscopic Retrograde Cholangiopancreatography, [645](#)  
 Endoscopic Submucosal Dissection, [645](#)  
 Endoscopic Third Ventriculostomy, [645](#)  
 Endoscopic Thoracic Sympathectomy, [645](#)  
 Endoscopic Ultrasound, [645](#)  
 Endoscopy, [645](#)  
 Endosome, [645](#)  
 Endosteum, [645](#)  
 Endostosis, [645](#)  
 Endothelial, [645](#)  
 Endothelial Cell, [645](#)  
 Endothelial Nitric Oxide Synthase, [645](#)  
 Endothelin, [646](#)  
 Endothelin Receptor Antagonists, [646](#)  
 Endothelium, [646](#)  
 Endothelium-Derived Relaxing Factor, [646](#)  
 Endotoxin, [646](#)  
 Endotracheal Intubation, [646](#)  
 Endotracheal Tube, [646](#)  
 Endourologist, [646](#)  
 Endovaginal Ultrasound, [646](#)  
 Endovascular, [646](#)  
 Endovascular Aneurysm Repair, [646](#)  
 Endovascular Aortic Aneurysm Repair, [646](#)  
 Endovascular Coiling, [647](#)  
 Endovascular Embolization, [647](#)  
 Endovascular Graft, [647](#)  
 Endovascular Repair, [647](#)  
 Endovenous Laser Ablation, [647](#)  
 Endurance, [647](#)  
 Enema, [647](#)  
 Energy, [647](#)  
 Energy Conservation, [647](#)  
 Energy Expenditure, [647](#)  
 Energy Increased, [647](#)  
 Energy Intake, [648](#)  
 Energy Metabolism, [648](#)  
 Energy Resolution, [648](#)  
 Energy Window, [648](#)  
 Enfuvirtide, [648](#)  
 Engagement, [648](#)  
 Enhanced Elimination, [648](#)  
 Enhanced Pause, [648](#)  
 Enhanced Recovery After Surgery, [648](#)  
 Enhancement Artifact, [648](#)  
 Enhancer, [648](#)  
 Enkephalin, [648](#)  
 Enlarged Adenoids, [648](#)  
 Enlarged Fontanelle, [648](#)  
 Enlarged Heart, [649](#)  
 Enlarged Liver, [649](#)  
 Enlarged Prostate, [649](#)  
 Enlarged Spleen, [649](#)  
 Enophthalmos, [649](#)  
 Enoxaparin, [649](#)  
 Enrichment, [649](#)  
 ENT Physician, [649](#)  
 Entamoeba, [649](#)  
 Entamoeba Coli, [649](#)  
 Entamoeba Hartmanni, [649](#)  
 Entamoeba Histolytica, [649](#)  
 Entamoeba Polecki, [649](#)  
 Enteral, [650](#)  
 Enteral Formula, [650](#)  
 Enteral Nutrition, [650](#)  
 Enteric, [650](#)  
 Enteric Bacteria, [650](#)  
 Enteric Duplication Cyst, [650](#)  
 Enteric Fever, [650](#)  
 Enteric Nervous System, [650](#)  
 Enteric-Coated, [650](#)  
 Enteritis, [650](#)  
 Enterobacter, [650](#)  
 Enterobacter Aerogenes, [650](#)  
 Enterobacter Cloacae, [650](#)  
 Enterobacter Hormaechei, [650](#)  
 Enterobacteriaceae, [651](#)  
 Enterobacteriaceae Carbapenem Resistance, [651](#)  
 Enterobiasis, [651](#)  
 Enterobius, [651](#)  
 Enterobius Vermicularis, [651](#)  
 Enterocoele, [651](#)  
 Enterochromaffin Cells, [651](#)  
 Enterochromaffin-Like Cells, [651](#)  
 Enteroclysis, [651](#)  
 Enterococcaceae, [651](#)  
 Enterococcal Infection, [651](#)  
 Enterococcus, [652](#)  
 Enterococcus Durans, [652](#)

---

---

Enterococcus Faecalis, [652](#)  
 Enterococcus Faecium, [652](#)  
 Enterocolitis, [652](#)  
 Enterocyte, [652](#)  
 Enterocytozoon, [652](#)  
 Enteroendocrine Cell, [652](#)  
 Enterohepatic Circulation, [652](#)  
 Enteromonas Hominis, [652](#)  
 Enteropathic, [652](#)  
 Enteropathic Arthritis, [652](#)  
 Enteropathy, [652](#)  
 Enteropathy-Associated T-Cell Lymphoma, [653](#)  
 Enteroscopy, [653](#)  
 Enterospasm, [653](#)  
 Enterostomy, [653](#)  
 Enterostomy Procedure, [653](#)  
 Enterotoxin, [653](#)  
 Enterovesical Fistula, [653](#)  
 Enterovirulent E. Coli, [653](#)  
 Enterovirus, [653](#)  
 Enterovirus D68, [653](#)  
 Enterovirus Infection, [653](#)  
 Enthesis, [653](#)  
 Enthesitis, [653](#)  
 Enthesitis And Enthesopathy, [653](#)  
 Enthesitis-Related Arthritis, [654](#)  
 Entorhinal Cortex, [654](#)  
 Entrance Wound, [654](#)  
 Entropion, [654](#)  
 Enucleation, [654](#)  
 Enucleation Procedure, [654](#)  
 Enuresis, [654](#)  
 Environment, [654](#)  
 Environmental Exposure, [654](#)  
 Environmental Health, [654](#)  
 Environmental Illness, [654](#)  
 Enzyme, [654](#)  
 Enzyme Activity, [655](#)  
 Enzyme Deficiency, [655](#)  
 Enzyme Induction, [655](#)  
 Enzyme Inhibition, [655](#)  
 Enzyme Inhibitor, [2120](#)  
 Enzyme Kinetics, [655](#)  
 Enzyme Markers, [655](#)  
 Enzyme Precursors, [655](#)  
 Enzyme Therapy, [655](#)  
 Enzyme-Linked Immunosorbent Assay, [655](#)  
 Eosinophil, [655](#)  
 Eosinophilia, [655](#)  
 Eosinophilia-Myalgia Syndrome, [655](#)  
 Eosinophilic Asthma, [655](#)  
 Eosinophilic Bronchitis, [655](#)  
 Eosinophilic Esophagitis, [656](#)  
 Eosinophilic Esophagitis Pathology, [656](#)  
 Eosinophilic Fasciitis, [656](#)  
 Eosinophilic Gastroenteritis, [656](#)  
 Eosinophilic Gastrointestinal Disorders, [656](#)  
 Eosinophilic Granuloma, [656](#)  
 Eosinophilic Granuloma (Pulmonary Histiocytosis X), [656](#)  
 Eosinophilic Granulomatosis With Polyangiitis, [656](#)  
 Eosinophilic Pneumonia Pathology, [656](#)  
 Eosinophils, [656](#)  
 Ependyma, [656](#)  
 Ependymal Cell, [656](#)  
 Ependymal Tumor, [656](#)  
 Ependymoblastoma, [657](#)  
 Ependymoma, [657](#)  
 Ephedrine, [657](#)  
 Ephedrinum, [657](#)  
 Ephelides, [657](#)  
 Ephrins, [657](#)  
 Epiblast, [657](#)  
 Epicanthal Folds, [657](#)  
 Epicardium, [657](#)  
 Epichlorohydrin, [657](#)  
 Epicondyle, [657](#)  
 Epicondylitis, [657](#)  
 Epicondylitis (Tennis Elbow), [657](#)  
 Epicondylitis, Lateral, [1118](#)  
 Epidemic, [658](#)  
 Epidemic Cerebrospinal Meningitis, [658](#)  
 Epidemic Cholera, [658](#)  
 Epidemic Keratoconjunctivitis, [658](#)  
 Epidemiologic Method, [658](#)  
 Epidemiologic Studies, [658](#)  
 Epidemiological Study, [658](#)  
 Epidemiologist, [658](#)  
 Epidemiology, [658](#)  
 Epidermal, [658](#)  
 Epidermal Acanthosis, [658](#)  
 Epidermal Cells, [658](#)  
 Epidermal Growth Factor, [658](#)  
 Epidermal Inclusion Cyst, [658](#)  
 Epidermal Layers, [658](#)  
 Epidermal Nevus, [658](#)  
 Epidermal Nevus Syndrome, [659](#)  
 Epidermis, [659](#)  
 Epidermodysplasia Verruciformis, [659](#)  
 Epidermoid Cyst, [659](#)  
 Epidermoid Cysts, [659](#)  
 Epidermolysis Bullosa, [659](#)  
 Epidermolysis Bullosa Acquisita, [659](#)  
 Epidermolytic Ichthyosis, [659](#)  
 Epidermophyton, [659](#)  
 Epidermophyton Floccosum, [659](#)  
 Epididymal Cyst, [659](#)  
 Epididymal Neoplasm, [659](#)

---

---

Epididymides, [660](#)  
Epididymis, [660](#)  
Epididymitis, [660](#)  
Epididymitis And Orchitis, [660](#)  
Epididymo-Orchitis, [660](#)  
Epidural, [660](#)  
Epidural Abscess, [660](#)  
Epidural Analgesia, [660](#)  
Epidural Analgesia For Labor, [660](#)  
Epidural Anesthesia, [660](#)  
Epidural Blood Patch, [660](#)  
Epidural Hematoma, [660](#)  
Epidural Neoplasm, [661](#)  
Epidural Space, [661](#)  
Epidural Steroid Injection, [661](#)  
Epigastric, [661](#)  
Epigastric Artery, [661](#)  
Epigastric Pain, [661](#)  
Epigastric Region, [661](#)  
Epigastrium, [661](#)  
Epigenetics, [661](#)  
Epigenome, [661](#)  
Epigenomics, [661](#)  
Epiglottic Vallecula, [662](#)  
Epiglottis, [662](#)  
Epiglottitis, [662](#)  
Epilepsia Partialis Continua, [662](#)  
Epilepsy, [662](#)  
Epilepsy and Seizures, [662](#)  
Epilepsy Surgery, [662](#)  
Epilepsy, Frontal Lobe, [781](#)  
Epileptic Spasms, [662](#)  
Epileptiform, [662](#)  
Epimysium, [662](#)  
Epinephrin, [662](#)  
Epinephrine, [662](#)  
Epineurium, [663](#)  
Epiphora, [663](#)  
Epiphyseal Plate, [663](#)  
Epiphyseal Plate Disorders, [663](#)  
Epiphysis, [663](#)  
Epiphysitis, [663](#)  
Epiplonic Foramen, [663](#)  
Epipodophyllotoxin, [663](#)  
Epiretinal Membrane, [663](#)  
Epirubicin, [663](#)  
Epirubicin Hydrochloride, [663](#)  
Episcleritis, [663](#)  
Episiotomy, [664](#)  
Episiotomy Repair, [664](#)  
Episodic Ataxia, [664](#)  
Epispadias, [664](#)  
Epistasis, [664](#)  
Epistaxis, [664](#)  
Epithalamic, [664](#)  
Epithalamus, [664](#)  
Epithelial, [664](#)  
Epithelial Cells, [664](#)  
Epithelial Cyst, [664](#)  
Epithelial Tissue, [664](#)  
Epithelioid Cell, [664](#)  
Epithelioid Histiocyte, [664](#)  
Epithelioid Sarcoma, [665](#)  
Epithelioid Trophoblastic Tumor, [665](#)  
Epithelioma, [665](#)  
Epitheliopathy, [665](#)  
Epithelium, [665](#)  
Epitope, [665](#)  
Epitope Mapping, [665](#)  
Eplerenone, [665](#)  
Epley Maneuver, [665](#)  
EPO Test, [665](#)  
Epoetin Alfa, [665](#)  
Epoetin Beta, [665](#)  
Eponym, [665](#)  
Epoprostenol, [665](#)  
Epoxide Hydrolase, [666](#)  
Epoxy Compounds, [666](#)  
Epsom Salts, [666](#)  
Epstein Pearls, [666](#)  
Epstein-Barr Virus, [666](#)  
Epstein-Barr Virus Antibody Test, [666](#)  
Eptifibatide, [666](#)  
Epulis, [666](#)  
Equilibrium Receptors, [666](#)  
Equilin, [666](#)  
Equine Morbillivirus, [666](#)  
Equinovarus Deformity, [666](#)  
Equipment Safety, [666](#)  
Equivalence Trial, [666](#)  
Eradication, [667](#)  
Erb Palsy, [667](#)  
Erb's Palsy, [667](#)  
Erbb Receptors, [667](#)  
Erbium, [667](#)  
ERCP, [667](#)  
Erdheim-Chester Disease, [667](#)  
Erectile Disorder, [667](#)  
Erectile Dysfunction, [667](#)  
Erection, [667](#)  
Erector Spinae, [667](#)  
Ergocalciferol, [667](#)  
Ergolines, [668](#)  
Ergometrine, [668](#)  
Ergonomics, [668](#)  
Ergosterol, [668](#)  
Ergot, [668](#)  
Ergotamine, [668](#)

---

---

Ergotism, [668](#)  
 Ergots, [668](#)  
 Eribulin Mesylate, [668](#)  
 Erlotinib Hydrochloride, [668](#)  
 Erosion, [668](#)  
 Erosion Lesion, [668](#)  
 Erosive Esophagitis, [668](#)  
 Erosive Gastritis, [668](#)  
 Erosive Lichen Planus, [669](#)  
 Erratic Results, [669](#)  
 Ertapenem, [669](#)  
 Erucic Acid, [669](#)  
 Erucic Acids, [669](#)  
 Eructation, [669](#)  
 Eruption Cyst, [669](#)  
 Eruptive Fever, [669](#)  
 Eruptive Vellus Hair Cysts, [669](#)  
 Eruptive Xanthomas, [669](#)  
 Eruptive Xanthomatosis, [669](#)  
 Eryptosis, [669](#)  
 Erysipelas, [669](#)  
 Erysipeloid, [670](#)  
 Erysipelothrix, [670](#)  
 Erythema, [670](#)  
 Erythema Ab Igne, [670](#)  
 Erythema Annulare Centrifugum, [670](#)  
 Erythema Chronicum Migrans, [670](#)  
 Erythema Dose, [670](#)  
 Erythema Dyschromicum Perstans, [670](#)  
 Erythema Elevatum Diutinum, [670](#)  
 Erythema Gydatum Repens, [670](#)  
 Erythema Induratum, [670](#)  
 Erythema Infectiosum, [670](#)  
 Erythema Infectiosum (Fifth Disease), [671](#)  
 Erythema Marginatum, [671](#)  
 Erythema Migrans, [671](#)  
 Erythema Multiforme, [671](#)  
 Erythema Nodosum, [671](#)  
 Erythema Toxicum, [671](#)  
 Erythema Toxicum Neonatorum, [671](#)  
 Erythrasma, [671](#)  
 Erythroblast, [671](#)  
 Erythroblastosis Fetalis, [671](#)  
 Erythroblastosis Foetalis, [671](#)  
 Erythrocyte, [671](#)  
 Erythrocyte Alloimmunization and Pregnancy, [671](#)  
 Erythrocyte Casts, [672](#)  
 Erythrocyte Enzyme Deficiency, [672](#)  
 Erythrocyte Membrane, [672](#)  
 Erythrocyte Sedimentation Rate, [672](#)  
 Erythrocytosis, [672](#)  
 Erythroderma, [672](#)  
 Erythrodermic Psoriasis, [672](#)  
 Erythrodysplasia, [672](#)  
 Erythroid Hyperplasia, [672](#)  
 Erythrokeratoderma Variabilis et Progressiva, [672](#)  
 Erythroleukemia, [672](#)  
 Erythroleukoplakia, [672](#)  
 Erythromelalgia, [672](#)  
 Erythromycin, [673](#)  
 Erythromycin Estolate, [673](#)  
 Erythromycin Ophthalmic Prophylaxis, [673](#)  
 Erythroplakia, [673](#)  
 Erythroplasia, [673](#)  
 Erythroplasia Of Queyrat, [673](#)  
 Erythropoiesis, [673](#)  
 Erythropoiesis-Stimulating Agents, [673](#)  
 Erythropoietin, [673](#)  
 Erythropoietin Deficiency, [673](#)  
 Erythropoietin Receptor, [673](#)  
 Erythropoietin Test, [673](#)  
 Erythrosine, [673](#)  
 ESB, [673](#)  
 ESB-Producing Organisms, [674](#)  
 Escape Reaction, [674](#)  
 Eschar, [674](#)  
 Escharotomy, [674](#)  
 Escherichia, [674](#)  
 Escherichia Albertii, [674](#)  
 Escherichia Blattae, [674](#)  
 Escherichia Coli, [674](#)  
 Escherichia Coli Infection, [674](#)  
 Escherichia Coli O157:H7 Infection, [674](#)  
 Escherichia Fergusonii, [674](#)  
 Escherichia Hermannii, [674](#)  
 Escherichia Vulneris, [674](#)  
 Escitalopram, [675](#)  
 ESKD, [675](#)  
 Esomeprazole, [675](#)  
 Esomeprazole Magnesium, [675](#)  
 Esophageal, [675](#)  
 Esophageal Achalasia, [675](#)  
 Esophageal Adenocarcinoma, [675](#)  
 Esophageal Atresia, [675](#)  
 Esophageal Bleeding, [675](#)  
 Esophageal Cancer, [675](#)  
 Esophageal Candidiasis, [675](#)  
 Esophageal Carcinoma, [675](#)  
 Esophageal Culture, [675](#)  
 Esophageal Dilation, [676](#)  
 Esophageal Diseases, [676](#)  
 Esophageal Diverticulum, [676](#)  
 Esophageal Duplication, [676](#)  
 Esophageal Fistula, [676](#)  
 Esophageal Foreign Body, [676](#)  
 Esophageal Infection, [676](#)  
 Esophageal Intramural Pseudodiverticulosis, [676](#)  
 Esophageal Leiomyoma, [676](#)

---



---

Esophageal Leukoplakia, [676](#)  
Esophageal Manometry, [676](#)  
Esophageal Motility Disorder, [676](#)  
Esophageal Necrosis, [676](#)  
Esophageal Perforation, [677](#)  
Esophageal pH Monitoring, [677](#)  
Esophageal Spasm, [677](#)  
Esophageal Spasms, [677](#)  
Esophageal Squamous Cell Carcinoma, [677](#)  
Esophageal Stenosis, [677](#)  
Esophageal Stent, [677](#)  
Esophageal Stricture, [677](#)  
Esophageal Surgery, [677](#)  
Esophageal Ulceration, [677](#)  
Esophageal Varices, [677](#)  
Esophageal Varices Cross-Reference, [677](#)  
Esophageal Web, [677](#)  
Esophagectomy, [678](#)  
Esophagitis, [678](#)  
Esophagogastric Bypass, [678](#)  
Esophagogastric Junction, [678](#)  
Esophagogastric Tamponade, [678](#)  
Esophagogastroduodenoscopy, [678](#)  
Esophagoscope, [678](#)  
Esophagoscopy, [678](#)  
Esophagram, [678](#)  
Esophagus, [678](#)  
Esophagus Disease, [678](#)  
Esophagus Neoplasm, [678](#)  
Esotropia, [678](#)  
ESR, [679](#)  
ESRD, [679](#)  
Essential, [679](#)  
Essential Fats, [679](#)  
Essential Fatty Acid, [679](#)  
Essential Hypertension, [679](#)  
Essential Medicines, [679](#)  
Essential Mixed Cryoglobulinemia, [679](#)  
Essential Nutrient, [679](#)  
Essential Thrombocythemia, [679](#)  
Essential Thrombocytosis, [679](#)  
Essential Tremor, [679](#)  
Essiac, [679](#)  
Estazolam, [679](#)  
Ester, [680](#)  
Esterase, [680](#)  
Esterified Estrogens, [680](#)  
Esthesia, [680](#)  
Esthesiometer, [680](#)  
Esthesioneuroblastoma, [680](#)  
Estimated Average Glucose, [680](#)  
Estimated Date Of Confinement, [680](#)  
Estimated Due Date, [680](#)  
Estimated Fetal Weight, [680](#)  
Estimated Glomerular Filtration Rate, [680](#)  
Estimation of Fetal Weight, [680](#)  
Estradiol, [680](#)  
Estradiol Blood Test, [681](#)  
Estramustine, [681](#)  
Estriol, [681](#)  
Estrogen, [681](#)  
Estrogen Inhibitor, [2120](#)  
Estrogen Overdose, [681](#)  
Estrogen Receptor, [681](#)  
Estrogen Replacement Therapy, [681](#)  
Estrogen Test, [681](#)  
Estrogen-Replacement Therapy, [681](#)  
Estrogens, [681](#)  
Estrone, [681](#)  
Estrous Cycle, [681](#)  
Eszopiclone, [681](#)  
Etacrynic Acid, [682](#)  
Etanercept, [682](#)  
Ethacrynic Acid, [682](#)  
Ethambutol, [682](#)  
Ethambutol Toxicity, [682](#)  
Ethane, [682](#)  
Ethanol, [682](#)  
Ethanol Metabolism, [682](#)  
Ethanol Poisoning, [682](#)  
Ethanamine, [682](#)  
Ether, [682](#)  
Ethers, [682](#)  
Ethics, [682](#)  
Ethinyl Estradiol, [683](#)  
Ethiodized Oil, [683](#)  
Ethion, [683](#)  
Ethionamide, [683](#)  
Ethmocephaly, [683](#)  
Ethmoid Bone, [683](#)  
Ethmoid Sinus, [683](#)  
Ethmoid Sinusitis, [683](#)  
Ethmoidal Air Cells, [683](#)  
Ethmoidal Sinus, [683](#)  
Ethmoiditis, [683](#)  
Ethnicity, [683](#)  
Ethoglucid, [683](#)  
Ethosuximide, [683](#)  
Ethoxzalamide, [684](#)  
Ethyl Chloride, [684](#)  
Ethyl Ether, [684](#)  
Ethylene, [684](#)  
Ethylene Chlorohydrin, [684](#)  
Ethylene Dibromide, [684](#)  
Ethylene Glycol, [684](#)  
Ethylene Glycol Blood Test, [684](#)  
Ethylene Glycol Poisoning, [684](#)  
Ethylene Glycol Toxicity, [684](#)

---

---

Ethylene Oxide, [684](#)  
Ethynodiol Diacetate, [684](#)  
Etidronate, [2120](#)  
Etidronic Acid, [685](#)  
Etilefrine, [685](#)  
Etiology, [685](#)  
Etodolac, [685](#)  
Etomidate, [685](#)  
Etonogestrel, [685](#)  
Etoposide, [685](#)  
Etoposide Phosphate, [685](#)  
Etoricoxib, [685](#)  
Eubacterium, [685](#)  
Eubacterium Limosum, [685](#)  
Eubacterium Rectale, [685](#)  
Eucalyptus Oil, [685](#)  
Eucalyptus Oil Overdose, [685](#)  
Euchromatin, [686](#)  
Eugenicist, [686](#)  
Eugenics, [686](#)  
Eugenol Oil Overdose, [686](#)  
Euglycemic Diabetic Ketoacidosis, [686](#)  
Euphoria, [686](#)  
Eustachian Tube, [686](#)  
Eustachian Tube Dysfunction, [686](#)  
Eustachian Tube Patency, [686](#)  
Eustachian Tube Problems, [686](#)  
Eustachian Valve, [686](#)  
Eustachian-Tube Dysfunction, [686](#)  
Euthanasia, [687](#)  
Euthanasia And Assisted Dying, [687](#)  
Euthyroid, [687](#)  
Euthyroid Goiter, [687](#)  
Euthyroid Hyperthyroxinemia, [687](#)  
Euthyroid Sick Syndrome, [687](#)  
Eutocia, [687](#)  
Euvolemic Hyponatremia, [687](#)  
Evacuation, [687](#)  
Evacuation Of Uterus, [687](#)  
Evaluation of Fetal Death, [687](#)  
Evans Syndrome, [687](#)  
Event Monitor, [687](#)  
Event-Free Survival, [688](#)  
Everolimus, [688](#)  
Eversion, [688](#)  
Evidence-Based Medicine, [2120](#)  
Evidence-Based Practice, [688](#)  
Evisceration, [688](#)  
Evoked Potential, [688](#)  
Evoked Potentials, [688](#)  
Evolution, [688](#)  
Evolutionarily Conserved Gene, [688](#)  
Ewart Sign, [688](#)  
Ewing Sarcoma, [688](#)  
Ewing's Family of Tumors, [688](#)  
Ewing's Sarcoma, [688](#)  
Ewing's Tumour, [689](#)  
Ewing's Family of Tumors, [689](#)  
Exacerbate, [689](#)  
Exacerbation, [689](#)  
Exacerbation Of COPD, [689](#)  
Exaggerated Placental Site, [689](#)  
Examination, [689](#)  
Examination Table, [689](#)  
Exanthem, [689](#)  
Exanthem Subitum, [689](#)  
Exanthema, [689](#)  
Exanthema Subitum, [689](#)  
Exanthematous Pustulosis, [689](#)  
Excavation, [690](#)  
Exceptional Release, [690](#)  
Excess Post-Exercise Oxygen Consumption, [690](#)  
Excessive Bruising, [690](#)  
Excessive Daytime Sleepiness, [690](#)  
Excessive Menstrual Bleeding, [690](#)  
Excessive Scarring, [690](#)  
Excessive Sweating, [690](#)  
Excessive Tearing, [690](#)  
Excessive Thirst, [690](#)  
Excessive Yawning, [690](#)  
Exchange Transfusion, [690](#)  
Excise, [690](#)  
Excision, [690](#)  
Excision Biopsy, [690](#)  
Excisional Biopsy, [690](#)  
Excitation, [691](#)  
Excitation-Contraction Coupling, [691](#)  
Excitatory Neurotransmitter, [691](#)  
Excitatory Synapse, [691](#)  
Excitotoxicity, [691](#)  
Excitotoxin, [691](#)  
Exclamation Point Hair, [691](#)  
Exclusion Diet, [691](#)  
Excoriate, [691](#)  
Excoriation, [691](#)  
Excoriation Disorder, [691](#)  
Excreta, [691](#)  
Excretion, [691](#)  
Executive Function, [692](#)  
Executive Functions, [692](#)  
Exemestane, [692](#)  
Exenatide, [692](#)  
Exencephaly, [692](#)  
Exenteration, [692](#)  
Exercise, [692](#)  
Exercise And Age, [692](#)  
Exercise And Immunity, [692](#)  
Exercise Headaches, [692](#)

---

---

Exercise Intolerance, 692  
Exercise Prescription, 692  
Exercise Stress Test, 693  
Exercise Tolerance, 693  
Exercise-Induced Anaphylaxis, 693  
Exercise-Induced Arrhythmias, 693  
Exercise-Induced Asthma, 693  
Exercise-Induced Bronchoconstriction, 693  
Exercises To Help Prevent Falls, 693  
Exfoliation Syndrome, 693  
Exfoliative Cytology, 693  
Exfoliative Dermatitis, 693  
Exhalation, 693  
Exhaled Nitric Oxide, 693  
Exhaustion, 693  
Exhaustion, Heat, 895  
Exhibitionism, 693  
Exhibitionistic Disorder, 694  
Exhumation, 694  
Existential Distress, 694  
Exit Wound, 694  
Exocervix, 694  
Exocrine, 694  
Exocrine Gland, 694  
Exocrine Glands, 694  
Exocrine Pancreas, 694  
Exocrine Pancreatic Insufficiency, 694  
Exocrine Pancreatic Insufficiency EPI, 694  
Exocytosis, 694  
Exogenous, 694  
Exogenous Cushing Syndrome, 694  
Exomphalos, 695  
Exon, 695  
Exopeptidase, 695  
Exophiala, 695  
Exophthalmic Goitre, 695  
Exophthalmos, 695  
Exosomes, 695  
Exostosis, 695  
Exotoxins, 695  
Exotropia, 695  
Expanded Rubella Syndrome, 695  
Expectant Treatment, 695  
Expectorant, 2120  
Expectorants, 695  
Expectoration, 695  
Experience, 696  
Expiration, 696  
Expiratory Grunting, 696  
Expiratory Reserve Volume, 696  
Explant, 696  
Explantation, 696  
Exploratory Laparotomy, 696  
Exploratory Operation, 696  
Exposure, 696  
Exposure Route, 696  
Expression, 696  
Expression Hopping, 696  
Expressivity, 696  
Expulsion, Stage Of, 1762  
Exsanguination, 697  
Exserohilum, 697  
Extend, 697  
Extension, 697  
Extensively Drug-Resistant TB, 2120  
Extensively Drug-Resistant Tuberculosis, 697  
Extensor, 697  
Extensor Digitorum, 697  
Extensor Plantar Response, 697  
Extent, 697  
External Auditory Canal, 697  
External Beam Partial Breast Irradiation, 697  
External Beam Radiation Therapy, 697  
External Capsule, 697  
External Carotid Artery, 698  
External Cephalic Version, 698  
External Ear, 698  
External Fixation, 698  
External Iliac Artery, 698  
External Iliac Vein, 698  
External Incontinence Devices, 698  
External Jugular Vein, 698  
External Nose, 698  
External Oblique Muscle, 698  
External Otitis, 698  
External Resorption, 698  
External Ventricular Drain, 698  
Extinction, 699  
Extirpation, 699  
Extra-Dural, 699  
Extra-Gastrointestinal Stromal Tumor, 699  
Extraamniotic, 699  
Extracapsular Cataract Surgery, 699  
Extracellular, 699  
Extracellular Exosome, 699  
Extracellular Fluid, 699  
Extracellular Matrix, 699  
Extracellular Pathogens, 699  
Extracellular Region, 699  
Extracellular Space, 699  
Extracellular Vesicle, 700  
Extracorporeal, 700  
Extracorporeal Circulation, 700  
Extracorporeal Membrane Oxygenation, 700  
Extracorporeal Shock Wave Lithotripsy, 700  
Extracranial, 700  
Extracranial Hematoma, 700  
Extract, 700

---

---

Extraction, 700  
Extragonadal Germ Cell Tumors, 700  
Extrahepatic Biliary Atresia, 700  
Extrahepatic Cholestasis, 700  
Extramammary Paget Disease, 701  
Extramammary Paget's Disease, 701  
Extramedullary Hematopoiesis, 701  
Extranodal, 701  
Extranodal Marginal Zone Lymphoma, 701  
Extranodal Natural Killer T-Cell Lymphoma, 701  
Extraocular, 701  
Extraocular Muscle, 701  
Extraocular Muscle Function Testing, 701  
Extraocular Muscles, 701  
Extrapulmonary Manifestation Of Tuberculosis, 701  
Extrapyramidal Disorder, 701  
Extrapyramidal Symptoms, 701  
Extrapyramidal System, 702  
Extrapyramidal Tracts, 702  
Extraskkeletal Chondroma, 702  
Extraskkeletal Mesenchymal Chondrosarcoma, 702  
Extraskkeletal Osteosarcoma, 702  
Extrasystole, 702  
Extrauterine Pregnancy, 702  
Extravasation, 702  
Extravascular Hemolysis, 702  
Extraventricular Drain, 702  
Extremities, 702  
Extremity, 702  
Extremity Angiography, 702  
Extremity Hardware Removal, 702  
Extremity X-Ray, 703  
Extrinsic, 703  
Extrinsic Allergic Alveolitis, 703  
Extrinsic Factor, 703  
Extrinsic Ureteral Obstruction, 703  
Extubate, 703  
Extubation, 703  
Extubation Criteria, 703  
Exudate, 703  
Exudative Pleural Effusion, 703  
Eye, 703  
Eye Allergies, 703  
Eye And Orbit Ultrasound, 704  
Eye Bank, 704  
Eye Bleed, 704  
Eye Cancer, 704  
Eye Color, 704  
Eye Development, 704  
Eye Disease, 704  
Eye Disorder, 704  
Eye Disorders, 704  
Eye Drops, 704  
Eye Emergencies, 704  
Eye Floaters, 704  
Eye Health, 704  
Eye Hemorrhage, 704  
Eye Infection, 705  
Eye Injury, 705  
Eye Irritation, 705  
Eye Melanoma, 705  
Eye Movement, 705  
Eye Muscle Repair, 705  
Eye Neoplasm, 705  
Eye Pain, 705  
Eye Prosthesis, 705  
Eye Redness, 705  
Eye Retraction Syndrome, 705  
Eye Trauma, 705  
Eye Twitching, 705  
Eye, Black, 218  
Eye, Diseases Of, 704  
Eyeball, 705  
Eyebrow, 706  
Eyeglass, 706  
Eyeglasses and Sunglasses, 707  
Eyeglasses, Sunglasses, And Magnifiers, 707  
Eyelash, 706  
Eyelashes, 706  
Eyelid, 706  
Eyelid Bump, 706  
Eyelid Cancer, 706  
Eyelid Coloboma, 706  
Eyelid Disorder, 706  
Eyelid Drooping, 706  
Eyelid Inflammation, 706  
Eyelid Lift, 706  
Eyelid Neoplasm, 706  
Eyelid Problems, 706  
Eyelid Swelling, 706  
Eyelid Twitch, 707  
Eyelids, 707  
Eyesight, 707  
Eyestrain, 707  
Ezetimibe, 707  
Fab Immunoglobulins, 709  
Fabry Disease, 709  
Fabry's Disease, 709  
Face, 709  
Face and Brow Presentation, 709  
Face Pain, 709  
Face Powder Poisoning, 709  
Face Presentation, 709  
Facelift, 709  
Facet Joint, 709  
Facet Joints, 710  
Facet Rhizotomy, 710

---

---

Facial Anatomy in Cutaneous Surgery, [710](#)  
Facial Asymmetry, [710](#)  
Facial Cleft, [710](#)  
Facial Colliculus, [710](#)  
Facial Edema, [710](#)  
Facial Expression, [710](#)  
Facial Fractures, [710](#)  
Facial Muscle, [710](#)  
Facial Neoplasm, [710](#)  
Facial Nerve, [710](#)  
Facial Nerve Palsy, [710](#)  
Facial Nerve Palsy From Birth Trauma, [710](#)  
Facial Nerve Problems, [711](#)  
Facial Nucleus, [711](#)  
Facial Pain, [711](#)  
Facial Palsy, [711](#)  
Facial Paralysis, [711](#)  
Facial Paresis, [711](#)  
Facial Soft Tissue Injuries, [711](#)  
Facial Spasm, [711](#)  
Facial Swelling, [711](#)  
Facial Tics, [711](#)  
Facial Trauma, [711](#)  
Facial Vein, [711](#)  
Facies, [711](#)  
Facies Hippocratica, [712](#)  
Facilitated Diffusion, [712](#)  
Facio-Scapulo-Humeral Dystrophy, [712](#)  
Facioscapulohumeral Dystrophy, [712](#)  
Facioscapulohumeral Muscular Dystrophy, [712](#)  
Factitious Disorder, [712](#)  
Factitious Disorder by Proxy, [712](#)  
Factitious Disorder Imposed on Another, [712](#)  
Factitious Disorders, [712](#)  
Factitious Hyperthyroidism, [712](#)  
Factor, [712](#)  
Factor H Deficiency, [712](#)  
Factor I Deficiency, [712](#)  
Factor II Assay, [713](#)  
Factor IX, [713](#)  
Factor IX Assay, [713](#)  
Factor V, [713](#)  
Factor V Assay, [713](#)  
Factor V Deficiency, [713](#)  
Factor V Leiden, [713](#)  
Factor V Leiden Mutation, [713](#)  
Factor VII, [713](#)  
Factor VII Assay, [713](#)  
Factor VII Deficiency, [713](#)  
Factor VIII, [713](#)  
Factor VIII Assay, [713](#)  
Factor X, [714](#)  
Factor X Assay, [714](#)  
Factor X Deficiency, [714](#)  
Factor XI Deficiency, [714](#)  
Factor XII, [714](#)  
Factor XII Assay, [714](#)  
Factor XII Deficiency, [714](#)  
Factor XIII, [714](#)  
Factor XIII Deficiency, [714](#)  
Facultative Anaerobe, [714](#)  
Faecal Fistula, [714](#)  
Faecalibacterium, [714](#)  
Faeces, [714](#)  
Faget Sign, [715](#)  
Fahr Disease, [715](#)  
Fahr Syndrome, [715](#)  
Fahr's Disease, [715](#)  
Fahr's Syndrome, [715](#)  
Failed Induction Of Labor, [715](#)  
Failed Intubation, [715](#)  
Failure To Thrive, [715](#)  
Fainting, [715](#)  
Falciform Ligament, [715](#)  
Falciparum Malaria, [715](#)  
Fall, [715](#)  
Fall Prevention, [715](#)  
Fall Risk, [716](#)  
Fall Screening, [716](#)  
Fallen Arches, [716](#)  
Fallen Bladder, [716](#)  
Fallopian Tube, [716](#)  
Fallopian Tube Cancer, [716](#)  
Fallopian Tube Disorders, [716](#)  
Fallopian Tube Torsion, [716](#)  
Fallopian Tubes, [716](#)  
Fallopian-Tube Endometriosis, [716](#)  
Fallot's Tetralogy, [716](#)  
Falls, [716](#)  
False Labor, [716](#)  
False Negative, [717](#)  
False Passages, [717](#)  
False Positive, [717](#)  
False Ribs, [717](#)  
False-Negative Result, [717](#)  
False-Positive Result, [717](#)  
Faltering Weight, [717](#)  
Falx Cerebelli, [717](#)  
Falx Cerebri, [717](#)  
Famciclovir, [717](#)  
Familial, [717](#)  
Familial Adenomatous Polyposis, [717](#)  
Familial Alobar Holoprosencephaly, [717](#)  
Familial Benign Pemphigus (Hailey-Hailey Disease), [718](#)  
Familial Chordoma, [718](#)  
Familial Combined Hyperlipidemia, [718](#)  
Familial Congestive Cardiomyopathy, [718](#)  
Familial Cutaneous Amyloidosis, [718](#)

---

---

Familial Dilated Cardiomyopathy (FDC), 718  
Familial Dysautonomia, 718  
Familial Dysbetalipoproteinemia, 718  
Familial Emphysema, 718  
Familial Hypercholesterolaemia, 718  
Familial Hypercholesterolemia, 718  
Familial Hypertriglyceridemia, 718  
Familial Hypocalciuric Hypercalcemia, 718  
Familial Jaundice, 718  
Familial Lichen Amyloidosis (Type IX), 719  
Familial Lipoprotein Lipase Deficiency, 719  
Familial Mediterranean Fever, 719  
Familial Multiple Lipomatosis, 719  
Familial Multiple Polyposis, 719  
Familial Multiple Trichoepithelioma, 719  
Familial Paroxysmal Peritonitis, 719  
Familial Primary Pulmonary Hypertension, 719  
Familial Renal Amyloidosis, 719  
Familial Testotoxicosis, 719  
Familial Tremor, 719  
Family Involvement, 719  
Family Medicine, 2120  
Family Planning, 719  
Family Study, 720  
Family Therapy, 720  
Famophos, 720  
Famotidine, 720  
Fanconi Anemia, 720  
Fanconi Syndrome, 720  
Fanconi-Bickel Syndrome, 720  
FAP, 720  
Farad, 720  
Farber Lipogranulomatosis, 720  
Farmer Lung, 720  
Farmer's Lung, 720  
Farsightedness, 721  
Fascia, 721  
Fascia Adherens, 721  
Fascia Lata, 721  
Fascial Loop, 721  
Fasciculation, 721  
Fasciculus Cuneatus, 721  
Fasciculus Gracilis, 721  
Fasciitis, 721  
Fasciola, 721  
Fasciola Hepatica, 721  
Fascioliasis, 721  
Fasciolopsis, 721  
Fasciolopsis Buski, 722  
Fasciotomy, 722  
Fast Heart Rate, 722  
Fast-Twitch Fiber, 722  
Fasting, 722  
Fasting Blood Glucose, 722  
Fasting Lipid Profile, 722  
Fasting Plasma Glucose Test, 722  
Fat, 722  
Fat Embolism, 722  
Fat Embolism Syndrome, 722  
Fat Measurement, 722  
Fat Necrosis, 723  
Fat Pad, 723  
Fat-Soluble Toxins, 723  
Fat-Soluble Vitamin, 723  
Fat-Soluble Vitamins, 723  
Fatal Catatonia, 723  
Fatal Familial Insomnia, 723  
Fatigue, 723  
Fatty Acid, 723  
Fatty Acid Oxidation, 723  
Fatty Acid Oxidation Disorder, 723  
Fatty Acid Synthase, 723  
Fatty Acid Synthesis, 724  
Fatty Acids, 724  
Fatty Degeneration, 724  
Fatty Liver, 724  
Fatty Liver Disease, 724  
Fatty Streak, 724  
Favism, 724  
Favre-Racouchot Syndrome, 724  
Favus, 724  
Fc Receptor, 724  
Fear, 724  
Fears, 724  
Febrile, 724  
Febrile Convulsion, 725  
Febrile Convulsions, 725  
Febrile Neutropenia, 725  
Febrile Seizure, 725  
Febrile Seizure Cross-Reference, 725  
Febrile Seizures, 725  
Febuxostat, 725  
FEC, 725  
Fec Protocol, 725  
Fecal Calprotectin, 725  
Fecal Culture, 725  
Fecal Elastase, 725  
Fecal Fat, 725  
Fecal Immunochemical Test, 725  
Fecal Impaction, 726  
Fecal Incontinence, 726  
Fecal Lactoferrin, 726  
Fecal Microbiota Transplant, 726  
Fecal Microbiota Transplantation, 726  
Fecal Occult Blood Test, 726  
Fecal Smear, 726  
Fecal Transplant, 726  
Fecalith, 726

---



---

Fecaluria, [726](#)  
 FECD, [726](#)  
 Feces, [726](#)  
 Fecolith, [726](#)  
 Fecundity, [727](#)  
 Feedback, [727](#)  
 Feedback Inhibition, [727](#)  
 Feedback Mechanism, [727](#)  
 Feeding Jejunostomy, [727](#)  
 Feeding Methods, [727](#)  
 Feeding Poor, [727](#)  
 Feeding Tube, [727](#)  
 Feingold Diet, [727](#)  
 Felbamate, [727](#)  
 Felodipine, [727](#)  
 Felon, [727](#)  
 Felty Syndrome, [727](#)  
 Felty's Syndrome, [728](#)  
 Female, [728](#)  
 Female Athlete Triad, [728](#)  
 Female Breast, [728](#)  
 Female Circumcision, [728](#)  
 Female Condom, [728](#)  
 Female Condoms, [728](#)  
 Female Genital Mutilation, [728](#)  
 Female Genital Sores, [728](#)  
 Female Genitalia, [728](#)  
 Female Gonad, [728](#)  
 Female Infertility, [728](#)  
 Female Orgasmic Disorder, [728](#)  
 Female Pattern Baldness, [729](#)  
 Female Pelvis, [729](#)  
 Female Phenotype, [729](#)  
 Female Reproductive Organs, [729](#)  
 Female Reproductive System, [729](#)  
 Female Self-Catheterization, [729](#)  
 Female Sexual Dysfunction, [729](#)  
 Female Sexual Interest/Arousal Disorder, [729](#)  
 Female Urethral Meatus, [729](#)  
 Female Urethral Opening, [729](#)  
 Feminization, [729](#)  
 Femoral, [729](#)  
 Femoral Anteversion, [729](#)  
 Femoral Artery, [730](#)  
 Femoral Canal, [730](#)  
 Femoral Hernia, [730](#)  
 Femoral Hernia Repair, [730](#)  
 Femoral Nerve, [730](#)  
 Femoral Nerve Dysfunction, [730](#)  
 Femoral Nerve Stretch Test, [730](#)  
 Femoral Neuropathy, [730](#)  
 Femoral Pulse, [730](#)  
 Femoral Shaft Fractures, [730](#)  
 Femoral Sheath, [730](#)  
 Femoral Triangle, [730](#)  
 Femoral Vein, [730](#)  
 Fentogram, [731](#)  
 Fentoliter, [731](#)  
 Femur, [731](#)  
 Femur Fracture, [731](#)  
 Femur Length, [731](#)  
 Fencer's Pose, [731](#)  
 Fenfluramine, [731](#)  
 Fenitrothion, [731](#)  
 Fenobucarb, [731](#)  
 Fenofibrate, [731](#)  
 Fenoldopam, [731](#)  
 Fenopropfen, [731](#)  
 Fenopropfen Calcium Overdose, [731](#)  
 Fenoterol, [731](#)  
 Fenoxycarb, [732](#)  
 Fentanyl, [2120](#)  
 Fentanyl Citrate, [732](#)  
 Fentanyl Derivatives, [732](#)  
 Fenthion, [732](#)  
 Fermentation, [732](#)  
 Ferriman-Gallwey Score, [732](#)  
 Ferritin, [732](#)  
 Ferritin Blood Test, [732](#)  
 Ferroptosis, [732](#)  
 Ferrous Sulfate, [732](#)  
 Fertile Period, [732](#)  
 Fertilisation, [732](#)  
 Fertility, [732](#)  
 Fertility Agent, [2120](#)  
 Fertility Domain, [733](#)  
 Fertility Drugs, [733](#)  
 Fertility Preservation, [733](#)  
 Fertility Treatment, [733](#)  
 Fertilization, [733](#)  
 Ferumoxytol, [733](#)  
 FESS, [733](#)  
 Festinating Gait, [733](#)  
 Fetal Acid-Base Status, [733](#)  
 Fetal Alcohol Spectrum Disorder, [733](#)  
 Fetal Alcohol Spectrum Disorders, [733](#)  
 Fetal Alcohol Syndrome, [733](#)  
 Fetal Anatomy Survey, [733](#)  
 Fetal Attitude, [734](#)  
 Fetal Bradycardia, [734](#)  
 Fetal Bradycardia, [734](#)  
 Fetal Circulation, [734](#)  
 Fetal Death, [734](#)  
 Fetal Development, [734](#)  
 Fetal Disease, [734](#)  
 Fetal Distress, [734](#)  
 Fetal Echocardiography, [734](#)  
 Fetal Fibronectin, [734](#)

---

---

Fetal Growth Restriction, [734](#)  
 Fetal Heart, [734](#)  
 Fetal Heart Monitoring, [734](#)  
 Fetal Heart Rate, [735](#)  
 Fetal Hemoglobin, [735](#)  
 Fetal Hydantoin Syndrome, [735](#)  
 Fetal Hydrops, [735](#)  
 Fetal Hypoxia, [735](#)  
 Fetal Kick Counts, [735](#)  
 Fetal Lie, [735](#)  
 Fetal Lung Maturity, [735](#)  
 Fetal Macrosomia, [735](#)  
 Fetal Malpresentation, [735](#)  
 Fetal Monitor, [735](#)  
 Fetal Monitoring, [736](#)  
 Fetal Presentation, [736](#)  
 Fetal Pyelectasis, [736](#)  
 Fetal Scalp Electrode, [736](#)  
 Fetal Scalp PH Testing, [736](#)  
 Fetal Scalp Stimulation, [736](#)  
 Fetal Stage, [736](#)  
 Fetal Tachycardia, [736](#)  
 Fetal Tissue, [736](#)  
 Fetal Ultrasound, [736](#)  
 Fetal Viability, [736](#)  
 Fetal Weight, [736](#)  
 Fetal-Maternal Erythrocyte Distribution Blood Test, [736](#)  
 Fetishism, [736](#)  
 Fetishistic Disorder, [737](#)  
 Fetomaternal Hemorrhage, [737](#)  
 Fetomaternal Transfusion, [737](#)  
 Fetor Hepaticus, [737](#)  
 Fetoscope, [737](#)  
 Fetus, [737](#)  
 FEV1/FVC Ratio, [737](#)  
 Fever, [737](#)  
 Fever And Headache, [737](#)  
 Fever Blister, [737](#)  
 Fever Of Unknown Origin, [737](#)  
 Fever-Induced Seizure, [737](#)  
 Fexofenadine, [737](#)  
 Fiber, [738](#)  
 Fiber Optic, [738](#)  
 Fibrates, [738](#)  
 Fibre, [738](#)  
 Fibrillary Glomerulonephritis, [738](#)  
 Fibrillation, [738](#)  
 Fibrin, [738](#)  
 Fibrin Degradation Products Blood Test, [738](#)  
 Fibrin Sealant, [738](#)  
 Fibrin Split Products, [738](#)  
 Fibrinogen, [738](#)  
 Fibrinogen Assay, [738](#)  
 Fibrinogen Blood Test, [738](#)  
 Fibrinogen Decreased, [739](#)  
 Fibrinoid Leukodystrophy, [739](#)  
 Fibrinoid Necrosis, [739](#)  
 Fibrinolysis, [739](#)  
 Fibrinolytic Agent, [739](#)  
 Fibrinolytic Therapy, [739](#)  
 Fibrinopeptide A Blood Test, [739](#)  
 Fibrinous Pericarditis, [739](#)  
 Fibroadenoma, [739](#)  
 Fibroadenoma Of The Breast, [739](#)  
 Fibroblast, [739](#)  
 Fibrocartilage, [739](#)  
 Fibrocystic Breast Changes, [739](#)  
 Fibrocystic Breast Condition, [739](#)  
 Fibrocystic Breasts, [740](#)  
 Fibrodysplasia Ossificans Progressiva, [740](#)  
 Fibroepithelial Polyp Of The Vulva, [740](#)  
 Fibrofolliculoma, [740](#)  
 Fibroid, [740](#)  
 Fibroid Tumor, [740](#)  
 Fibroids, [740](#)  
 Fibroids, Cysts, And Polyps, [435](#)  
 Fibrolamellar Carcinoma, [740](#)  
 Fibrolamellar Hepatocellular Carcinoma, [740](#)  
 Fibrolipoma, [740](#)  
 Fibroma, [740](#)  
 Fibroma Of Tendon Sheath, [741](#)  
 Fibromas, [741](#)  
 Fibromatoses, [741](#)  
 Fibromatosis, [741](#)  
 Fibromuscular Dysplasia, [741](#)  
 Fibromyalgia, [741](#)  
 Fibromyalgia Syndrome, [741](#)  
 Fibromyxosarcoma, [741](#)  
 Fibronectin, [741](#)  
 Fibroproliferation, [741](#)  
 Fibrosarcoma, [741](#)  
 FibroScan, [741](#)  
 Fibrosing Alopecia, [741](#)  
 Fibrosing Cholangitis, [741](#)  
 Fibrosis, [742](#)  
 Fibrous Caverositis, [742](#)  
 Fibrous Dysplasia, [742](#)  
 Fibrous Dysplasia Pathology, [742](#)  
 Fibrous Interstitial Pneumonia, [742](#)  
 Fibrous Joint, [742](#)  
 Fibrous Plaque, [742](#)  
 Fibrous Plaques of the Penis, [742](#)  
 Fibrous Sclerosis of the Penis, [742](#)  
 Fibula, [742](#)  
 Fibular Artery, [742](#)  
 Fick Principle, [742](#)  
 Fidaxomicin, [2120](#)  
 Fiducial Markers, [742](#)

---

---

Field Fever, [742](#)  
 Field Of View, [743](#)  
 Fifth Disease, [743](#)  
 FIGO Leiomyoma Classification, [743](#)  
 Filamentous Fungus, [743](#)  
 Filarial Elephantiasis, [743](#)  
 Filarial Lymphedema, [743](#)  
 Filariasis, [743](#)  
 Filgrastim, [2121](#)  
 Filler, [743](#)  
 Filler Excipient, [743](#)  
 Filopodium, [743](#)  
 Filoviridae, [743](#)  
 Filoviruses, [743](#)  
 Filtered Back Projection, [743](#)  
 Filtering Procedure, [744](#)  
 Filtration, [744](#)  
 Filtration Fraction, [744](#)  
 Filum Terminale, [744](#)  
 Fimbria, [744](#)  
 Fimbriae, [744](#)  
 Fimbriae Proteins, [744](#)  
 Finasteride, [2121](#)  
 Fine Crackles, [744](#)  
 Fine Motor Control, [744](#)  
 Fine Motor Skills, [744](#)  
 Fine Needle Aspiration, [744](#)  
 Fine Needle Aspiration Of The Thyroid, [744](#)  
 Fine-Needle Aspiration, [744](#)  
 Finegoldia, [744](#)  
 Finegoldia Magna, [745](#)  
 Finger, [745](#)  
 Finger Dislocation, [745](#)  
 Finger Flexors, [745](#)  
 Finger Injury, [745](#)  
 Finger Joint, [745](#)  
 Finger Pain, [745](#)  
 Finger Stick, [745](#)  
 Finger Sucking, [745](#)  
 Finger-Nose Test, [745](#)  
 Fingernail, [745](#)  
 Fingers That Change Color, [745](#)  
 Finkelstein Test, [745](#)  
 Finnish Type Amyloidosis (Type V), [745](#)  
 Fire Ants, [746](#)  
 Firesetting Behavior, [746](#)  
 First Adult Seizure, [746](#)  
 First Aid, [746](#)  
 First Aid for Burns, [746](#)  
 First Aid Kit, [746](#)  
 First Menstrual Period, [746](#)  
 First Stage Of Labor, [746](#)  
 First Trimester, [746](#)  
 First-Degree Atrioventricular Block, [746](#)  
 First-Degree AV Block, [746](#)  
 First-Degree Burn, [746](#)  
 First-Degree Heart Block, [746](#)  
 First-Episode Psychosis, [747](#)  
 First-Pass Effect, [747](#)  
 First-Pass Metabolism, [747](#)  
 Fish Odour Syndrome, [747](#)  
 Fish Oil, [747](#)  
 Fish Poisoning, [747](#)  
 Fish Tank Granuloma, [747](#)  
 Fish Tapeworm Infection, [747](#)  
 Fish Venom, [747](#)  
 Fish-Handler's Disease, [747](#)  
 Fisher Syndrome, [747](#)  
 Fisher's Syndrome, [747](#)  
 Fishhook Removal, [748](#)  
 Fissure, [748](#)  
 Fissured Tongue, [748](#)  
 Fistula, [748](#)  
 Fistula, Arteriovenous, [128](#)  
 Fistula-In-Ano, [748](#)  
 Fistulotomy, [748](#)  
 Fit, [748](#)  
 Fitz Hugh Curtis Syndrome, [748](#)  
 Fitz-Hugh-Curtis Syndrome, [748](#)  
 Five-Alpha Reductase, [748](#)  
 Five-Year Survival Rate, [748](#)  
 Fixation, [748](#)  
 Fixative, [748](#)  
 Fixator Muscle, [749](#)  
 Fixed Drug Eruption, [2121](#)  
 Fixed Drug Eruptions, [749](#)  
 Fixed Joint, [749](#)  
 Fixed Macrophage, [749](#)  
 Flaccid, [749](#)  
 Flagellin, [749](#)  
 Flagellum, [749](#)  
 Flail Chest, [749](#)  
 Flakka, [749](#)  
 Flanged Needle, [749](#)  
 Flank, [749](#)  
 Flank Pain, [749](#)  
 Flap Operation, [749](#)  
 Flapping Tremor, [749](#)  
 Flare, [750](#)  
 Flare-Up, [750](#)  
 Flashes, [750](#)  
 Flashing, [750](#)  
 Flat Affect, [750](#)  
 Flat Bones, [750](#)  
 Flat Feet, [750](#)  
 Flatfoot, [750](#)  
 Flatfoot (Pes Planus), [750](#)  
 Flatulence, [750](#)

---

---

Flatus, [750](#)  
Flatus and Bloating, [750](#)  
Flavin Adenine Dinucleotide, [750](#)  
Flaviviridae, [750](#)  
Flavivirus, [751](#)  
Flaviviruses, [751](#)  
Flavobacteriaceae, [751](#)  
Flavobacterium, [751](#)  
Flavoxate, [751](#)  
Flaxseed Extract, [751](#)  
Flea, [751](#)  
Flea Bites, [751](#)  
Flea Bites In Humans, [751](#)  
Fleas, [751](#)  
Flecainide, [751](#)  
Fleischner Sign, [751](#)  
Flesh-Eating Bacteria, [751](#)  
Flesh-Eating Bug, [752](#)  
Flex, [752](#)  
Flexed Breech, [752](#)  
Flexibility, [752](#)  
Flexible Nasolaryngoscopy, [752](#)  
Flexible Sigmoidoscopy, [752](#)  
Flexion, [752](#)  
Flexor, [752](#)  
Flexor Digitorum Profundus, [752](#)  
Flexor Digitorum Superficialis, [752](#)  
Flexor Tendon Anatomy, [752](#)  
Flexor Tenosynovitis, [752](#)  
Flexural, [752](#)  
Flibanserin, [752](#)  
Flight-Or-Fight Response, [753](#)  
Floater, [753](#)  
Floaters, [753](#)  
Floating Kidney, [753](#)  
Floating Rib, [753](#)  
Floating Ribs, [753](#)  
Flocculation, [753](#)  
Floor Of The Mouth Cancer, [753](#)  
Floppy Baby Syndrome, [753](#)  
Floppy Iris Syndrome, [753](#)  
Floppy Valve Syndrome, [753](#)  
Flora, [753](#)  
Flossing, [753](#)  
Flotation, [754](#)  
Flow Cytometry, [754](#)  
Flow-Volume Loop, [754](#)  
Flowmeter, [754](#)  
Floxacin, [754](#)  
Floxuridine, [754](#)  
Flu, [754](#)  
Flu (Influenza), [754](#)  
Flucloxacillin, [754](#)  
Fluconazole, [754](#)  
Flucytosine, [754](#)  
Fludarabine, [754](#)  
Fludarabine Phosphate, [754](#)  
Fludeoxyglucose F-18, [754](#)  
Fludrocortisone, [755](#)  
Fluid Around The Heart, [755](#)  
Fluid Balance, [755](#)  
Fluid Imbalance, [755](#)  
Fluid in Lungs, [755](#)  
Fluid Management, [755](#)  
Fluid Overload, [755](#)  
Fluid Therapy, [755](#)  
Fluid Wave, [755](#)  
Fluid, Electrolyte, and Acid-Base Disorders, [569](#)  
Fluke, [755](#)  
Flumazenil, [755](#)  
Flunarizine, [755](#)  
Flunisolide, [755](#)  
Fluocinolone, [755](#)  
Fluocinolone Acetonide, [756](#)  
Fluocinonide, [756](#)  
Fluocortolone, [756](#)  
Fluorescein, [756](#)  
Fluorescein Angiography, [756](#)  
Fluorescein Eye Stain, [756](#)  
Fluorescence, [756](#)  
Fluorescence In Situ Hybridization, [756](#)  
Fluorescence Microscopy, [756](#)  
Fluorescent Dye, [756](#)  
Fluoridation, [756](#)  
Fluoride, [756](#)  
Fluoride In Diet, [756](#)  
Fluoride Overdose, [756](#)  
Fluoride Poisoning, [757](#)  
Fluoride Toxicity, [757](#)  
Fluoride Varnish, [757](#)  
Fluorimetry, [757](#)  
Fluorine, [757](#)  
Fluorine-18, [757](#)  
Fluorometholone, [757](#)  
Fluorophotometry, [757](#)  
Fluoropyrimidine, [757](#)  
Fluoroquinolone, [757](#)  
Fluoroquinolone-Associated Disability, [757](#)  
Fluoroquinolones, [757](#)  
Fluoroscopy, [758](#)  
Fluorosis, [758](#)  
Fluorouracil, [758](#)  
Fluorouracil Side Effect, [758](#)  
Fluoxetine, [758](#)  
Fluoxymesterone, [758](#)  
Flupentixol, [758](#)  
Fluphenazine, [758](#)  
Fluphenazine Hydrochloride, [758](#)

---

---

Flurandrenolide, [758](#)  
Flurazepam, [758](#)  
Flurbiprofen, [758](#)  
Flush, [758](#)  
Flushable Reagent Stool Blood Test, [758](#)  
Flushing, [759](#)  
Flutamide, [759](#)  
Fluticasone, [2121](#)  
Flutter, [759](#)  
Flutter Valve, [759](#)  
Fluvastatin, [759](#)  
Fluvoxamine, [759](#)  
Flux, [759](#)  
FM, [759](#)  
fMRI, [759](#)  
FMS, [759](#)  
FNA, [759](#)  
FND, [759](#)  
Foam Cell, [759](#)  
Foam Cells, [760](#)  
Focal, [760](#)  
Focal Adhesion, [760](#)  
Focal Aware Seizures, [760](#)  
Focal Cortical Dysplasia, [760](#)  
Focal Dermal Hypoplasia Syndrome, [760](#)  
Focal Glomerulosclerosis, [760](#)  
Focal Impaired Awareness Seizures, [760](#)  
Focal Neurologic Deficits, [760](#)  
Focal Neurological Deficit, [760](#)  
Focal Neuropathy, [760](#)  
Focal Nodular Hyperplasia, [760](#)  
Focal Sclerosis, [760](#)  
Focal Segmental Glomerular Sclerosis, [761](#)  
Focal Segmental Glomerulosclerosis, [761](#)  
Focal Seizure, [761](#)  
Focal-Onset Seizure, [761](#)  
Focus Words, [761](#)  
Focused Assessment with Sonography for Trauma, [761](#)  
FODMAP, [761](#)  
FODMAP Diet, [761](#)  
FODMAPs, [761](#)  
Foetal Acidosis, [761](#)  
Foetal Alcohol Syndrome, [761](#)  
Foetor Hepaticus, [761](#)  
Foetus, [762](#)  
Foix-Alajouanine Syndrome, [762](#)  
Folate, [762](#)  
Folate Antagonist, [762](#)  
Folate Deficiency, [762](#)  
Folate Deficiency Anemia, [762](#)  
Folate Vitamin B9, [762](#)  
Folate-Deficiency Anemia, [762](#)  
Foley Catheter, [762](#)  
Folic Acid, [762](#)  
Folic Acid And Birth Defect Prevention, [762](#)  
Folic Acid Deficiency Anemia, [762](#)  
Folic Acid In Diet, [762](#)  
Folic Acid Supplementation, [762](#)  
Folic Acid Test, [763](#)  
Folie a Deux, [763](#)  
Folinic Acid, [763](#)  
Follicle, [763](#)  
Follicle Stimulating Hormone, [763](#)  
Follicle-Stimulating Hormone, [763](#)  
Follicle-Stimulating Hormone Abnormalities, [763](#)  
Follicle-Stimulating Hormone Blood Test, [763](#)  
Follicular, [763](#)  
Follicular Cyst, [763](#)  
Follicular Fluid, [763](#)  
Follicular Hyperkeratosis, [763](#)  
Follicular Lymphoma, [763](#)  
Follicular Thyroid Carcinoma, [764](#)  
Follicular Unit, [764](#)  
Follicular Unit Transplantation, [764](#)  
Folliculitis, [764](#)  
Folliculitis Decalvans, [764](#)  
Fomentation, [764](#)  
Fomepizole, [764](#)  
Fomite, [764](#)  
Fomites, [764](#)  
Fondaparinux, [764](#)  
Fondaparinux Sodium, [764](#)  
Fong Disease, [764](#)  
Fonofos, [764](#)  
Fonsecaea, [764](#)  
Fontan Procedure, [765](#)  
Fontanel, [765](#)  
Fontanelle, [765](#)  
Food Addiction, [765](#)  
Food Additives, [765](#)  
Food Allergies, [765](#)  
Food Allergy, [765](#)  
Food Contamination, [765](#)  
Food Hygiene, [765](#)  
Food Intolerance, [765](#)  
Food Jags, [765](#)  
Food Poisoning, [765](#)  
Food Protein-Induced Allergic Proctocolitis, [766](#)  
Food Protein-Induced Enterocolitis Syndrome, [766](#)  
Food Safety, [766](#)  
Food Supplement, [766](#)  
Food-Borne Illness, [766](#)  
Food-Drug Interaction, [2121](#)  
Foodborne Disease, [766](#)  
Foot, [766](#)  
Foot Drop, [766](#)  
Foot Fracture, [766](#)  
Foot Pain, [766](#)

---

---

Foot Pain (Plantar Fasciitis), 766  
Foot, Leg, And Ankle Swelling, 1506  
Footling Breech, 766  
FOP, 766  
Foramen, 766  
Foramen Lacerum, 767  
Foramen of Monro, 767  
Foramen Ovale, 767  
Foramen Rotundum, 767  
Foramen Spinosum, 767  
Foraminotomy, 767  
Forbes-Albright Syndrome, 767  
Force Plates, 767  
Forced Expiratory Volume, 767  
Forced Oscillation Technique, 767  
Forced Vital Capacity, 767  
Forceps, 767  
Forceps Delivery, 767  
Forceps-Assisted Delivery, 768  
Forchheimer Spots, 768  
Forcible Intercourse, 768  
Fordyce Granules, 768  
Fordyce's Angiokeratomas, 768  
Forearm, 768  
Forearm Circumference, 768  
Forehead, 768  
Forehead Lift, 768  
Foreign Body, 768  
Foreign Body Airway Obstruction, 768  
Foreign Body Aspiration, 768  
Foreign Body in Ear, 768  
Foreign Body In The Nose, 768  
Foreign Body Reaction, 769  
Foreign-Body Aspiration, 769  
Forensic, 769  
Forensic Anthropology, 769  
Forensic Autopsy, 769  
Forensic Autopsy of Blunt Force Trauma, 769  
Forensic Autopsy of Sharp Force Injuries, 769  
Forensic Defense Wounds, 769  
Forensic Evidence, 769  
Forensic Examination, 769  
Forensic Genetics, 769  
Forensic Laceration, 769  
Forensic Medicine, 2121  
Forensic Medicine (Legal Medicine), 2121  
Forensic Odontology, 769  
Forensic Pathology, 770  
Forensic Pathology of Firearm Wounds, 770  
Forensic Pathology of Thermal Injuries, 770  
Forensic Psychiatry, 770  
Forensic Radiography, 770  
Forensic Toxicology, 770  
Forensic Toxicology Screen, 770  
Forensics, 770  
Foreskin, 770  
Formaldehyde, 770  
Formalin, 770  
Formalin-Fixed Paraffin-Embedded Tissue, 770  
Formate, 770  
Formed Elements Of Blood, 771  
Formetanate, 771  
Formication, 771  
Formoterol Fumarate, 771  
Formothion, 771  
Formparanate, 771  
Formula Feed, 771  
Formula Feeding, 771  
Formulae, 771  
Formulary, 771  
Fornix, 771  
Forrest Classification, 771  
Forrester Classification, 771  
Fortification, 772  
Fortified Wine, 772  
Fosaprepitant Dimeglumine, 772  
Foscarnet, 2121  
Foscarnet Sodium, 772  
Fosfomycin, 772  
Fosinopril, 772  
Fossa, 772  
Fossa Navicularis, 772  
Fossula, 772  
Foster Kennedy Syndrome, 772  
Fotemustine, 772  
Fothergill Sign, 772  
Foul-Smelling Stools, 773  
Found Dead, 773  
Four-Dimensional Ultrasound, 773  
Fourchette, 773  
Fourier Analysis, 773  
Fournier Disease, 773  
Fournier Gangrene, 773  
Fourth Finger, 773  
Fourth Heart Sound, 773  
Fourth Stage Of Labor, 773  
Fourth Venereal Disease, 773  
Fourth-Degree Tear, 773  
Fovea, 773  
Fovea Centralis, 773  
Foville Syndrome, 773  
Fowler Position, 774  
Fowler's Syndrome, 774  
Fox-Fordyce Disease, 774  
Foxglove Poisoning, 774  
Fractional Excretion Of Sodium, 774  
Fractional Excretion of Urea, 774  
Fractional Exhaled Nitric Oxide, 774

---



---

Fractional Flow Reserve, [774](#)  
 Fractional IPV, [774](#)  
 Fractional Shortening, [774](#)  
 Fracture, [774](#)  
 Fracture Displacement, [775](#)  
 Fracture Fixation, [775](#)  
 Fracture Liaison Service, [775](#)  
 Fracture Malunion, [775](#)  
 Fracture Nonunion, [775](#)  
 Fracture Of The Temporal Bone, [775](#)  
 Fracture Prevention, [775](#)  
 Fracture-Fixation Device, [775](#)  
 Fractured Clavicle In The Newborn, [775](#)  
 Fractured Elbow, [775](#)  
 Fractured Ilium, [775](#)  
 Fractured Nose, [775](#)  
 Fractured Rib, [775](#)  
 Fractures, [775](#)  
 Fragile X Syndrome, [776](#)  
 Fragile X-Associated Tremor/Ataxia Syndrome, [776](#)  
 FRAIL Scale, [776](#)  
 Frailty, [776](#)  
 Frambesia, [776](#)  
 Framboesia, [776](#)  
 Frameshift Mutation, [776](#)  
 Framycetin, [776](#)  
 Francisella, [776](#)  
 Francisella Tularensis, [776](#)  
 Francisellaceae, [776](#)  
 Francois Dyscephaly Syndrome, [776](#)  
 Frank Breech, [776](#)  
 Frank Haemoptysis, [777](#)  
 Frank Sign, [777](#)  
 Frank-Starling Law, [777](#)  
 Frank-Starling Mechanism, [777](#)  
 Frasier Syndrome, [777](#)  
 Fraternal Twins, [777](#)  
 FRAX Tool, [777](#)  
 FRAXA, [777](#)  
 Freckle, [777](#)  
 Freckles, [777](#)  
 Free Fragment, [777](#)  
 Free Radical, [777](#)  
 Free Radical Injury, [777](#)  
 Free T4 Test, [778](#)  
 Free Thyroxine Index, [778](#)  
 Free Water Clearance, [778](#)  
 Freeze Drying, [778](#)  
 Freezing, [778](#)  
 Freiberg Disease, [778](#)  
 Fremitus, [778](#)  
 French Disease, [778](#)  
 Frenectomy, [778](#)  
 Frenulum, [778](#)  
 Frenulum Breve, [778](#)  
 Frenum, [778](#)  
 Frequency, [778](#)  
 Frequency Of Urination, [779](#)  
 Frequency Range, [779](#)  
 Frequency Response, [779](#)  
 Frequent Bowel Movements, [779](#)  
 Frequent Or Urgent Urination, [779](#)  
 Frequent Urination, [779](#)  
 Fresh Frozen Plasma, [779](#)  
 Fresh Specimen, [779](#)  
 Frey Syndrome, [779](#)  
 Frey'S Syndrome, [779](#)  
 Friable, [779](#)  
 Friction, [779](#)  
 Friction Blisters, [779](#)  
 Friedlander Pneumonia, [780](#)  
 Friedman Curve, [780](#)  
 Friedreich Ataxia, [780](#)  
 Friedreich's Ataxia, [780](#)  
 Friedreichs Ataxia Syndrome, [780](#)  
 Froin Syndrome, [780](#)  
 Froment Sign, [780](#)  
 Frontal, [780](#)  
 Frontal Bone, [780](#)  
 Frontal Bossing, [780](#)  
 Frontal Lobe, [780](#)  
 Frontal Lobe Disorder, [780](#)  
 Frontal Lobe Epilepsy, [780](#)  
 Frontal Lobe Seizures, [781](#)  
 Frontal Lobe Syndrome, [781](#)  
 Frontal Sinus, [781](#)  
 Frontal Sinusitis, [781](#)  
 Frontalis Suspension, [781](#)  
 Frontotemporal Dementia, [781](#)  
 Frontotemporal Lobar Degeneration, [781](#)  
 Frostbite, [781](#)  
 Frotteuristic Disorder, [781](#)  
 Frozen Section, [781](#)  
 Frozen Shoulder, [781](#)  
 Fructosamine, [781](#)  
 Fructose, [782](#)  
 Fructose 1,6-Diphosphatase Deficiency, [782](#)  
 Fructose Intolerance, [782](#)  
 Fructose Malabsorption, [782](#)  
 Fructose Metabolism, [782](#)  
 Fructose-1,6-Bisphosphatase Deficiency, [782](#)  
 Fruity Breath, [782](#)  
 Frustration, [782](#)  
 Frys Syndrome, [782](#)  
 FSH, [782](#)  
 Ft., [782](#)  
 FTA-ABS Blood Test, [782](#)  
 FTA-ABS Test, [782](#)

---

---

Fuchs Endothelial Corneal Dystrophy, 782  
 Fuchs Endothelial Dystrophy, 783  
 Fuchs' Dystrophy, 783  
 Fucose, 783  
 Fucosidosis, 783  
 Fucosyltransferase, 783  
 Fuel Oil Poisoning, 783  
 Fugue, 783  
 Fukuda Stepping Test, 783  
 Fukuyama Congenital Muscular Dystrophy, 783  
 Fulguration, 783  
 Full Agonist, 783  
 Full Blood Count, 784  
 Full Liquid Diet, 784  
 Full Sibling, 784  
 Full-Thickness Skin Graft, 784  
 Fulminant, 784  
 Fulminant Hepatitis, 784  
 Fulminant Myocarditis, 784  
 Fulminating Hyperpyrexia, 784  
 Fulvestrant, 784  
 Fumigation, 784  
 Functional Asplenia, 784  
 Functional Assessment, 784  
 Functional Capacity, 784  
 Functional Constipation, 784  
 Functional Decline, 784  
 Functional Diarrhea, 785  
 Functional Disorder, 785  
 Functional Dyspepsia, 785  
 Functional Electrical Stimulation, 785  
 Functional Endoscopic Sinus Surgery, 785  
 Functional Gastrointestinal Disorders, 785  
 Functional Hypothalamic Amenorrhea, 785  
 Functional Impairment, 785  
 Functional Incontinence, 785  
 Functional Independence Measure, 785  
 Functional MRI, 786  
 Functional Murmur, 786  
 Functional Neurologic Disorder, 786  
 Functional Neurologic Disorder/Conversion Disorder, 786  
 Functional Neurological Disorder, 786  
 Functional Neurological Symptom Disorder, 786  
 Functional Ovarian Cysts, 786  
 Functional Residual Capacity, 786  
 Functional Screening, 786  
 Functional Status, 786  
 Functional Urinary Incontinence, 786  
 Functioning Carcinoid, 786  
 Fundal Height, 786  
 Fundoplication, 786  
 Fundoscopy, 787  
 Fundus, 787  
 Fundus Oculi, 787  
 Fundus Photography, 787  
 Fundus Uteri, 787  
 Fungal, 787  
 Fungal Arthritis, 787  
 Fungal Corneal Ulcer, 787  
 Fungal Endophthalmitis, 787  
 Fungal Infection, 787  
 Fungal Nail Infection, 788  
 Fungal Nail Infections, 788  
 Fungal Nails, 788  
 Fungal Pathogenesis, 788  
 Fungal Pneumonia, 788  
 Fungal Resistance Mechanisms, 788  
 Fungal Skin Infections, 788  
 Fungating, 788  
 Fungemia, 788  
 Fungi, 788  
 Fungicide, 788  
 Fungoid, 788  
 Fungus, 788  
 Funiculus, 789  
 Funisitis, 789  
 Funnel Chest, 789  
 Funnel-Web Spider Bite, 789  
 Furan, 789  
 Furocoumarin, 789  
 Furosemide, 2121  
 Furrowed Tongue, 789  
 Fursultiamin, 789  
 Furuncle, 789  
 Furuncle And Carbuncle, 789  
 Furuncular Myiasis, 789  
 Furunculosis, 789  
 Fusariosis, 789  
 Fusarium, 789  
 Fusarium Infection, 790  
 Fusarium Solani, 790  
 Fused Placenta, 790  
 Fused Teeth, 790  
 Fusidic Acid, 790  
 Fusiform, 790  
 Fusion, 790  
 Fusion Gene, 790  
 Fusion Of The Ear Bones, 790  
 Fusion Protein, 790  
 Fusobacterium, 790  
 Fusobacterium Infection, 790  
 Fusobacterium Necrophorum, 790  
 Fusobacterium Nucleatum, 790  
 Fusospirochetal Gingivitis, 791  
 G-Protein Coupled Receptors, 793  
 G6PD Deficiency, 793

---

---

G6PD Deficiency in the Newborn, [793](#)  
 Gaba Receptor, [793](#)  
 Gabapentin, [793](#)  
 GAD, [793](#)  
 Gadolinium, [793](#)  
 Gadolinium-Based Contrast Agent, [793](#)  
 Gadolinium-Induced Nephrogenic Systemic Fibrosis, [793](#)  
 Gadopentetate Dimeglumine, [793](#)  
 Gag Gene, [794](#)  
 Gag Protein, [794](#)  
 Gag Reflex, [794](#)  
 Gagging, [794](#)  
 Gain, [794](#)  
 Gain-Of-Function Mutation, [794](#)  
 Gaisbock Syndrome, [794](#)  
 Gait, [794](#)  
 Gait Abnormality, [794](#)  
 Gait Apraxia, [794](#)  
 Gait Assessment, [794](#)  
 Gait Ataxia, [794](#)  
 Gait Cycle, [794](#)  
 Gait Disturbance, [794](#)  
 Gait Speed, [795](#)  
 Gait Training, [795](#)  
 Gaits, [795](#)  
 Galactocele, [795](#)  
 Galactocerebrosidase (GALC) Deficiency, [795](#)  
 Galactocerebrosidase Deficiency, [795](#)  
 Galactocerebroside Beta-Galactosidase Deficiency, [795](#)  
 Galactogogue, [795](#)  
 Galactokinase Deficiency, [795](#)  
 Galactorrhea, [795](#)  
 Galactorrhea-Amenorrhea Syndrome, [795](#)  
 Galactorrhoea, [795](#)  
 Galactosaemia, [795](#)  
 Galactose, [796](#)  
 Galactose Metabolism, [796](#)  
 Galactose-1-Phosphate Uridyl Transferase Deficiency, [796](#)  
 Galactose-1-Phosphate Uridyltransferase Blood Test, [796](#)  
 Galactosemia, [796](#)  
 Galactosialidosis, [796](#)  
 Galactoside, [796](#)  
 Galactosyltransferase, [796](#)  
 Galant Reflex, [796](#)  
 Galeazzi Fracture, [796](#)  
 Galeazzi Fracture-Dislocation, [796](#)  
 Galeazzi Sign, [796](#)  
 Gall Bladder Disease, [796](#)  
 Gall-Bladder, [797](#)  
 Gallavardin Phenomenon, [797](#)  
 Gallbladder Absence, [797](#)  
 Gallbladder Adenomyomatosis, [797](#)  
 Gallbladder Cancer, [797](#)  
 Gallbladder Carcinoma, [797](#)  
 Gallbladder Diseases, [797](#)  
 Gallbladder Duplication, [797](#)  
 Gallbladder Hydrops, [797](#)  
 Gallbladder Infection, [797](#)  
 Gallbladder Inflammation, [797](#)  
 Gallbladder Mucocele, [797](#)  
 Gallbladder Neck, [798](#)  
 Gallbladder Necrosis, [798](#)  
 Gallbladder Pain, [798](#)  
 Gallbladder Perforation, [798](#)  
 Gallbladder Polyp, [798](#)  
 Gallbladder Radionuclide Scan, [798](#)  
 Gallbladder Volvulus, [798](#)  
 Gallium, [798](#)  
 Gallium Nitrate, [798](#)  
 Gallium Scan, [798](#)  
 Gallium-68, [798](#)  
 Gallop Rhythm, [798](#)  
 Gallstone Ileus, [798](#)  
 Gallstone Pancreatitis, [798](#)  
 Gallstones, [799](#)  
 GALT Deficiency, [799](#)  
 Galvanocautery, [799](#)  
 Gambling Disorder, [799](#)  
 Gamete, [799](#)  
 Gamete Intrafallopian Transfer, [799](#)  
 Gametogenesis, [799](#)  
 Gamma Camera, [799](#)  
 Gamma Globulin, [799](#)  
 Gamma Hydroxybutyrate, [799](#)  
 Gamma Knife, [799](#)  
 Gamma Knife Radiosurgery, [799](#)  
 Gamma Rays, [799](#)  
 Gamma-Aminobutyric Acid, [799](#)  
 Gamma-Benzene Hexachloride, [800](#)  
 Gamma-Chemokines, [800](#)  
 Gamma-Globulin, [800](#)  
 Gamma-Glutamyl Transferase Blood Test, [800](#)  
 Gamma-Hydroxybutyrate Toxicity, [800](#)  
 Gamma-Interferon, [800](#)  
 Gamma-Ray Therapy, [800](#)  
 Gammaherpesvirinae, [800](#)  
 Gamna-Gandy Bodies, [800](#)  
 GAN, [800](#)  
 Ganciclovir, [800](#)  
 Ganglia, [800](#)  
 Ganglioglioma, [800](#)  
 Ganglion, [801](#)  
 Ganglion Cell, [801](#)  
 Ganglion Cyst, [801](#)  
 Ganglionectomy, [801](#)  
 Ganglioneuroblastoma, [801](#)  
 Ganglioneuroma, [801](#)

---

---

Ganglioneuromatosis, [801](#)  
Ganglionic Blocker, [2121](#)  
Ganglionic Blockers, [801](#)  
Ganglions, [801](#)  
Ganglioside, [801](#)  
Gangliosidosis, [801](#)  
Gangliosidosis GM2 Type 2, [801](#)  
Gangrene, [801](#)  
Ganser Syndrome, [801](#)  
Gap Junction, [801](#)  
Gardasil, [802](#)  
Gardner Syndrome, [802](#)  
Gardnerella, [802](#)  
Gardnerella Vaginalis, [802](#)  
Gargle, [802](#)  
Garré's Osteomyelitis, [802](#)  
Gartner Duct, [802](#)  
Gartner Duct Cyst, [802](#)  
Gas, [802](#)  
Gas (Flatus), [802](#)  
Gas And Gas Pains, [802](#)  
Gas Chromatography, [802](#)  
Gas Exchange, [802](#)  
Gas Gangrene, [802](#)  
Gas Gangrene (Clostridial Myonecrosis), [802](#)  
Gas Poisoning, [803](#)  
Gas Scavenger, [803](#)  
Gas Transport, [803](#)  
Gas – Flatulence, [803](#)  
Gaskin Maneuver, [803](#)  
Gasoline Poisoning, [803](#)  
Gasserian Ganglion, [803](#)  
Gastralgia, [803](#)  
Gastrectomy, [803](#)  
Gastric Acid, [803](#)  
Gastric Adenocarcinoma, [803](#)  
Gastric Analysis, [803](#)  
Gastric Antral Vascular Ectasia, [804](#)  
Gastric Antrum, [804](#)  
Gastric Balloon, [804](#)  
Gastric Bezoar, [804](#)  
Gastric Biopsy, [804](#)  
Gastric Body, [804](#)  
Gastric Bypass, [804](#)  
Gastric Bypass Surgery, [804](#)  
Gastric Cancer, [804](#)  
Gastric Carcinoid Tumor, [804](#)  
Gastric Cardia, [804](#)  
Gastric Culture, [804](#)  
Gastric Dysplasia, [804](#)  
Gastric Emptying, [804](#)  
Gastric Emptying Study, [805](#)  
Gastric Fistula, [805](#)  
Gastric Fundus, [805](#)  
Gastric Glands, [805](#)  
Gastric Hemorrhage, [805](#)  
Gastric Intestinal Metaplasia, [805](#)  
Gastric Juice, [805](#)  
Gastric Lavage, [805](#)  
Gastric Leiomyosarcoma, [805](#)  
Gastric Lymphoma, [805](#)  
Gastric Malignancy, [805](#)  
Gastric Motility, [805](#)  
Gastric Mucosa, [805](#)  
Gastric Muscularis, [805](#)  
Gastric Necrosis, [805](#)  
Gastric Obstruction, [806](#)  
Gastric Outlet Obstruction, [806](#)  
Gastric Perforation, [806](#)  
Gastric Pneumatosis, [806](#)  
Gastric Polyps, [806](#)  
Gastric Rugae, [806](#)  
Gastric Secretion, [806](#)  
Gastric Serosa, [806](#)  
Gastric Stenosis, [806](#)  
Gastric Submucosa, [806](#)  
Gastric Suction, [806](#)  
Gastric Surgery, [806](#)  
Gastric Tissue Biopsy And Culture, [806](#)  
Gastric Ulcer, [806](#)  
Gastric Varices, [806](#)  
Gastric Volvulus, [807](#)  
Gastric Wall, [807](#)  
Gastrin, [807](#)  
Gastrin Blood Test, [807](#)  
Gastrin Secretion, [807](#)  
Gastrin-Releasing Peptide, [807](#)  
Gastrinoma, [807](#)  
Gastritis, [807](#)  
Gastrocnemius, [807](#)  
Gastrocnemius Muscle, [807](#)  
Gastroduodenal Artery, [807](#)  
Gastroduodenal Crohn Disease, [808](#)  
Gastroduodenostomy, [808](#)  
Gastroenteritis, [808](#)  
Gastroenteritis Bacterial, [808](#)  
Gastroenteritis Viral, [808](#)  
Gastroenteritis, Acute, [808](#)  
Gastroenterologist, [808](#)  
Gastroenterology, [808](#)  
Gastroenteropathy, Protein Losing, [1613](#)  
Gastroenterostomy, [808](#)  
Gastroepiploic Artery, [808](#)  
Gastroesophageal, [808](#)  
Gastroesophageal Junction Adenocarcinoma, [808](#)  
Gastroesophageal Reflux, [808](#)  
Gastroesophageal Reflux Disease, [808](#)  
Gastroesophageal Reflux Disease (GERD), [808](#)

---

---

Gastroesophageal Z-Line, 809  
 Gastrogastic Intussusception, 809  
 Gastrografin Study, 809  
 Gastrointestinal, 809  
 Gastrointestinal Agents, 809  
 Gastrointestinal Amyloidosis, 809  
 Gastrointestinal Bleeding, 809  
 Gastrointestinal Bleeding Scan, 809  
 Gastrointestinal Cancer, 809  
 Gastrointestinal Decontamination, 809  
 Gastrointestinal Disease, 809  
 Gastrointestinal Disease and Pregnancy, 809  
 Gastrointestinal Disorder, 809  
 Gastrointestinal Fistula, 810  
 Gastrointestinal Foreign Bodies, 810  
 Gastrointestinal Gas, 810  
 Gastrointestinal Health, 810  
 Gastrointestinal Hemorrhage, 810  
 Gastrointestinal Hormone, 810  
 Gastrointestinal Motility Disorder, 810  
 Gastrointestinal Neoplasm, 810  
 Gastrointestinal Obstruction, 810  
 Gastrointestinal Pain, 810  
 Gastrointestinal Perforation, 810  
 Gastrointestinal Stromal Tumor, 810  
 Gastrointestinal Stromal Tumors, 810  
 Gastrointestinal Stromal Tumors (GISTs), 810  
 Gastrointestinal System, 811  
 Gastrointestinal Tract, 811  
 Gastrointestinal Villi, 811  
 Gastrojejunostomy, 811  
 Gastromalacia, 811  
 Gastroparesis, 811  
 Gastropathy, 811  
 Gastroplasty, 811  
 Gastropoptosis, 811  
 Gastroschisis, 811  
 Gastroschisis Repair, 811  
 Gastroscope, 811  
 Gastroscopy, 812  
 Gastrosphenic Fistulization, 812  
 Gastrostomy, 812  
 Gastrostomy Tube, 812  
 Gastrostomy Tube Insertion, 812  
 Gastrula, 812  
 Gastrulation, 812  
 Gated Blood Pool Scan, 812  
 Gatifloxacin, 812  
 Gaucher Disease, 812  
 Gaucher Splenomegaly, 812  
 Gaucher's Disease, 812  
 Gaucher-Schlagenhauser, 812  
 Gaultheria Oil, 812  
 Gauss, 813  
 Gauze, 813  
 Gavage, 813  
 Gayet-Wernicke Syndrome, 813  
 Gaze Palsy, 813  
 GBM, 813  
 Gefitinib, 813  
 Gegenhalten, 813  
 Geiger Counter, 813  
 Gel, 813  
 Gel Electrophoresis, 813  
 Gelastic Seizure, 813  
 Gelatin, 813  
 Gelfoam, 814  
 Gelineau Syndrome, 814  
 Gemcitabine, 814  
 Gemcitabine Hydrochloride, 814  
 Gemella, 814  
 Gemfibrozil, 814  
 Gemifloxacin, 814  
 Gemination, 814  
 Gemistocytic Astrocytoma, 814  
 Gemtuzumab Ozogamicin, 814  
 Gender, 814  
 Gender Dysphoria, 814  
 Gender Identity, 814  
 Gender Incongruence, 814  
 Gene, 815  
 Gene Amplification, 815  
 Gene Conversion, 815  
 Gene Deletion, 815  
 Gene Dosage, 815  
 Gene Duplication, 815  
 Gene Expression, 815  
 Gene Frequency, 815  
 Gene Fusion, 815  
 Gene Modification, 815  
 Gene Mutation, 815  
 Gene Panel, 815  
 Gene Rearrangement, 815  
 Gene Silencing, 815  
 Gene Targeting, 815  
 Gene Therapy, 815  
 Gene Transfer, 816  
 Gene-Environment Interaction, 816  
 Gene-Modified, 816  
 General Anesthesia, 816  
 General Anesthetic, 816  
 General Paresis, 816  
 Generalised, 816  
 Generalized, 816  
 Generalized Anxiety Disorder, 816  
 Generalized Anxiety Disorder In Children, 816  
 Generalized Essential Telangiectasia, 816  
 Generalized Granuloma Annulare, 816

---

---

Generalized Lipodystrophy, 816  
 Generalized Lymphadenopathy, 817  
 Generalized Seizure, 817  
 Generalized Tonic-Clonic Seizure, 817  
 Generalized Weakness, 817  
 Generalized-Onset Seizure, 817  
 Generator, 817  
 Generic, 817  
 Generic Drug, 2121  
 Generic Medicine, 2121  
 Generic Name, Drug, 2121  
 Genes, 817  
 Genesis, 817  
 Genetic Anticipation, 817  
 Genetic Carrier, 817  
 Genetic Code, 817  
 Genetic Counseling, 817  
 Genetic Counselling, 817  
 Genetic Counselor, 818  
 Genetic Disease, 818  
 Genetic Disorder, 818  
 Genetic Drift, 818  
 Genetic Engineering, 818  
 Genetic Heterogeneity, 818  
 Genetic Linkage, 818  
 Genetic Marker, 818  
 Genetic Polymorphism, 818  
 Genetic Profile, 818  
 Genetic Recombination, 818  
 Genetic Screening, 818  
 Genetic Sequencing, 818  
 Genetic Sonogram, 818  
 Genetic Susceptibility, 819  
 Genetic Technique, 819  
 Genetic Testing, 819  
 Genetic Transcription, 819  
 Genetic Variant Of Uncertain Significance, 819  
 Genetics, 819  
 Geniculate Body, 819  
 Geniculate Ganglion, 819  
 Genital, 819  
 Genital Edema, 819  
 Genital Herpes, 819  
 Genital Herpes in Pregnancy, 819  
 Genital Herpes in Women, 819  
 Genital Injury, 820  
 Genital Mycoplasmas, 820  
 Genital Prolapse, 820  
 Genital System, 820  
 Genital Ulcer Disease, 820  
 Genital Wart, 820  
 Genital Warts, 820  
 Genitalia, 820  
 Genitals, 820  
 Genito-Pelvic Pain/Penetration Disorder, 820  
 Genito-Urinary Medicine, 820  
 Genitofemoral Nerve, 821  
 Genitourinary, 821  
 Genitourinary Fistula, 821  
 Genitourinary Neoplasm, 821  
 Genitourinary Syndrome Of Menopause, 821  
 Genitourinary Tract, 821  
 Genodermatoses, 821  
 Genome, 821  
 Genomic Imprinting, 821  
 Genomic Instability, 821  
 Genomic Medicine, 2122  
 Genomics, 821  
 Genotoxicity, 821  
 Genotype, 821  
 Genotype Determination, 821  
 Genotype-Phenotype Correlation, 822  
 Gentamicin, 822  
 Gentamicin Sulfate, 822  
 Gentian, 822  
 Gentian Violet, 822  
 Genu, 822  
 Genu Recurvatum, 822  
 Genu Valgum, 822  
 Genu Varum, 822  
 Geographic Atrophy, 822  
 Geographic Tongue, 822  
 Geophagia, 822  
 Geophagy, 822  
 Geotrichum, 822  
 Geriatric Assessment, 823  
 Geriatric Bioethics, 823  
 Geriatric Care Manager, 823  
 Geriatric Dentistry, 823  
 Geriatric Depression, 823  
 Geriatric Depression Scale, 823  
 Geriatric Dream Changes, 823  
 Geriatric Dysphagia, 823  
 Geriatric Hospitalization, 823  
 Geriatric Infection, 823  
 Geriatric Informed Consent, 823  
 Geriatric Medical Ethics, 824  
 Geriatric Medicine, 2122  
 Geriatric Nursing, 824  
 Geriatric Nutrition, 824  
 Geriatric Osteoarthritis, 824  
 Geriatric Rehabilitation, 824  
 Geriatric Screening, 824  
 Geriatric Sleep Changes, 824  
 Geriatric Syndrome, 824  
 Geriatric Trauma, 824  
 Geriatrician, 824  
 Geriatrics, 824

---



---

Germ, [824](#)  
 Germ Cell, [824](#)  
 Germ Cell Tumor, [825](#)  
 Germ Cell Tumor Staging, [825](#)  
 Germ Cell Tumors, [825](#)  
 Germ Layer, [825](#)  
 Germ Theory Of Disease, [825](#)  
 German Measles, [825](#)  
 Germanium, [825](#)  
 Germinoma, [825](#)  
 Germline Mutation, [825](#)  
 Germline Testing, [825](#)  
 Germline Variant, [825](#)  
 Gerontologist, [825](#)  
 Gerstmann Syndrome, [825](#)  
 Gerstmann-Straussler-Scheinker Syndrome, [826](#)  
 Gesell Developmental Schedules, [826](#)  
 Gestation, [826](#)  
 Gestation Period, [826](#)  
 Gestational Age, [826](#)  
 Gestational Diabetes, [826](#)  
 Gestational Diabetes Mellitus, [826](#)  
 Gestational Hypertension, [826](#)  
 Gestational Pyelonephritis, [826](#)  
 Gestational Sac, [826](#)  
 Gestational Trophoblastic Disease, [827](#)  
 Gestational Trophoblastic Neoplasia, [827](#)  
 GFR, [827](#)  
 GHD, [827](#)  
 Ghon Complex, [827](#)  
 Ghon Focus, [827](#)  
 Ghrelin, [827](#)  
 Gianotti Crosti Syndrome, [827](#)  
 Gianotti-Crosti Syndrome, [827](#)  
 Giant Axonal Disease, [827](#)  
 Giant Axonal Neuropathy, [827](#)  
 Giant Cell Arteritis, [827](#)  
 Giant Cell Arteritis Emergency, [828](#)  
 Giant Cell Glioblastoma, [828](#)  
 Giant Cell Glioblastoma, Astrocytoma, [828](#)  
 Giant Cell Glioblastoma, Multiforme, [828](#)  
 Giant Cell Interstitial Pneumonia, [828](#)  
 Giant Cell Myocarditis, [828](#)  
 Giant Cell Tumor, [828](#)  
 Giant Cell Tumor Of Bone, [828](#)  
 Giant Cell Tumor Of Soft Tissue, [828](#)  
 Giant Cell Tumor Of Tendon Sheath, [828](#)  
 Giant Cells, [828](#)  
 Giant Congenital Nevus, [828](#)  
 Giant Coronary Aneurysm, [828](#)  
 Giant Cyst of the Retina, [828](#)  
 Giant Fibroadenoma, [828](#)  
 Giant Hypertrophic Gastritis, [829](#)  
 Giant Lymph Node Hyperplasia, [829](#)  
 Giant Platelet Syndrome, [829](#)  
 Giant Right Atrium, [829](#)  
 Giant Urticaria, [829](#)  
 Giardia, [829](#)  
 Giardia Infection, [829](#)  
 Giardia Lamblia, [829](#)  
 Giardiasis, [829](#)  
 Gibbus Deformity, [829](#)  
 Gibraltar Fever, [829](#)  
 Giemsa Stain, [829](#)  
 Gigabecquerel, [829](#)  
 Gigantism, [830](#)  
 Gigli Saw, [830](#)  
 Gilbert Syndrome, [830](#)  
 Gilbert's Disease, [830](#)  
 Gilbert's Syndrome, [830](#)  
 Gilbert-Lereboullet Syndrome, [830](#)  
 Gilles De La Tourette Syndrome, [830](#)  
 Gingiva, [830](#)  
 Gingiva Disease, [830](#)  
 Gingival Disease, [830](#)  
 Gingival Hemorrhage, [830](#)  
 Gingival Hyperplasia, [830](#)  
 Gingival Infections, [830](#)  
 Gingival Margin, [830](#)  
 Gingival Melanoma, [831](#)  
 Gingival Neoplasm, [831](#)  
 Gingival Overgrowth, [831](#)  
 Gingival Pocket, [831](#)  
 Gingival Recession, [831](#)  
 Gingivectomy, [831](#)  
 Gingivitis, [831](#)  
 Gingivitis Transmission, [831](#)  
 Gingivoplasty, [831](#)  
 Gingivostomatitis, [831](#)  
 Ginkgo Biloba, [831](#)  
 Ginseng, [831](#)  
 Ginseng Preparation, [831](#)  
 Girdle Pain, [831](#)  
 Gitelman Syndrome, [832](#)  
 Glabella, [832](#)  
 Gland, [832](#)  
 Glanders, [832](#)  
 Glandular Fever, [832](#)  
 Glandular Plague, [832](#)  
 Glans Penis, [832](#)  
 Glanzmann Thrombasthenia, [832](#)  
 Glasgow Coma Scale, [832](#)  
 Glass Eye, [832](#)  
 Glass Ionomer, [832](#)  
 Glassy Cell Carcinoma, [832](#)  
 Glaucoma, [832](#)  
 Glaucoma Drainage Implant, [832](#)  
 Glaucoma Surgery, [832](#)

---

---

Glaze Poisoning, [832](#)  
Gleason Grading System, [833](#)  
Gleason Score, [833](#)  
Gleason Tumor Grade, [833](#)  
Glenn Procedure, [833](#)  
Glenohumeral Joint, [833](#)  
Glenoid, [833](#)  
Glenoid Cavity, [833](#)  
Glenoid Labrum, [833](#)  
Gliadin, [833](#)  
Glial Cell, [833](#)  
Glial Cells, [833](#)  
Glial Fibrillary Acidic Protein, [834](#)  
Glibenclamide, [834](#)  
Gliclazide, [834](#)  
GLIM Criteria, [834](#)  
Glimepiride, [834](#)  
Glioblastoma, [834](#)  
Glioblastoma Multiforme, [834](#)  
Glioma, [834](#)  
Glioma-Polyposis Syndrome, [834](#)  
Gliomatosis Cerebri, [834](#)  
Gliosarcoma, [834](#)  
Gliosis, [834](#)  
Glipizide, [834](#)  
Glisson's Capsule, [835](#)  
Global Aphasia, [835](#)  
Global Developmental Delay, [835](#)  
Global Health, [835](#)  
Global Longitudinal Strain, [835](#)  
Globe Rupture, [835](#)  
Globin, [835](#)  
Globoid Cell Leukoencephalopathy, [835](#)  
Globozoospermia, [835](#)  
Globulin, [835](#)  
Globus, [835](#)  
Globus Hystericus, [835](#)  
Globus Pallidus, [835](#)  
Globus Sensation, [835](#)  
Glomangiomyoma, [836](#)  
Glomangiosarcoma, [836](#)  
Glomerular Basement Membrane, [836](#)  
Glomerular Capsule, [836](#)  
Glomerular Filtrate, [836](#)  
Glomerular Filtration, [836](#)  
Glomerular Filtration Rate, [836](#)  
Glomerular Hyperfiltration, [836](#)  
Glomeruli, [836](#)  
Glomerulomegaly, [836](#)  
Glomerulonephritis, [836](#)  
Glomerulonephritis Acute, [836](#)  
Glomerulopathy, [836](#)  
Glomerulosclerosis, [837](#)  
Glomerulus, [837](#)  
Glomus Jugulare Tumor, [837](#)  
Glomus Tumor, [837](#)  
Glomus Tumors, [837](#)  
Glomus Tympanicum, [837](#)  
Glomus Tympanicum Tumor, [837](#)  
Glossal, [837](#)  
Glossalgia, [837](#)  
Glossectomy, [837](#)  
Glossinidae, [837](#)  
Glossitis, [837](#)  
Glossitis Areata Migrans, [837](#)  
Glossodynia, [838](#)  
Glossopharyngeal Nerve, [838](#)  
Glossopharyngeal Neuralgia, [838](#)  
Glossoptosis, [838](#)  
Glossopyrosis, [838](#)  
Glottis, [838](#)  
GLP-1 Agonists, [838](#)  
GLP-1 Receptor Agonists, [838](#)  
GLS, [838](#)  
Glu, [838](#)  
Glucagon, [838](#)  
Glucagon Blood Test, [838](#)  
Glucagon Signaling, [838](#)  
Glucagon-like Peptide-1, [838](#)  
Glucagonoma, [839](#)  
Glucan, [839](#)  
Glucarpidase, [839](#)  
Glucocerebrosidase Deficiency, [839](#)  
Glucocorticoid, [839](#)  
Glucocorticoid Remediabale Aldosteronism, [839](#)  
Glucocorticoid Replacement Therapy, [839](#)  
Glucocorticoid-Induced Osteoporosis, [839](#)  
Glucocorticoid-Remediabale Aldosteronism, [839](#)  
Glucocorticoid-Sparing Therapy, [839](#)  
Glucocorticoids, [839](#)  
Glucometer, [840](#)  
Gluconates, [840](#)  
Gluconeogenesis, [840](#)  
Glucosamine, [840](#)  
Glucosamine and Chondroitin, [840](#)  
Glucose, [840](#)  
Glucose Challenge Test, [840](#)  
Glucose Intolerance, [840](#)  
Glucose Management Indicator, [840](#)  
Glucose Screening Tests During Pregnancy, [840](#)  
Glucose Sensor, [840](#)  
Glucose Test, [840](#)  
Glucose Tolerance Test, [840](#)  
Glucose Transport, [841](#)  
Glucose Transporter, [841](#)  
Glucose Urine Test, [841](#)  
Glucose-6-Phosphatase, [841](#)  
Glucose-6-Phosphate Dehydrogenase Deficiency, [841](#)

---

---

Glucose-6-Phosphate Dehydrogenase Test, [841](#)  
 Glucosuria, [841](#)  
 Glucosyl Ceramide Lipidosis, [841](#)  
 Glucosylceramidase, [841](#)  
 Glucuronate, [841](#)  
 Glucuronic Acid, [841](#)  
 Glucuronide, [841](#)  
 Glucuronyl Transferase, [841](#)  
 Glue Ear, [842](#)  
 GLUT4 Transporter, [842](#)  
 Glutamate, [842](#)  
 Glutamate Decarboxylase, [842](#)  
 Glutamate Receptor, [842](#)  
 Glutamic Acid, [842](#)  
 Glutamic Acid Decarboxylase, [842](#)  
 Glutamine, [842](#)  
 Glutaral, [842](#)  
 Glutaric Acidemia Type 1, [842](#)  
 Glutaric Aciduria, [842](#)  
 Glutathione, [842](#)  
 Glutathione Disulfide, [842](#)  
 Glutathione Peroxidase, [843](#)  
 Glutathione Synthetase Deficiency, [843](#)  
 Glutathione Transferase, [843](#)  
 Gluteal, [843](#)  
 Gluteal Gait, [843](#)  
 Gluteal Injury, [843](#)  
 Gluteal Muscle, [843](#)  
 Gluteal Tendinopathy, [843](#)  
 Gluten, [843](#)  
 Gluten Enteropathy, [843](#)  
 Gluten Intolerance, [843](#)  
 Gluten Sensitivity, [843](#)  
 Gluten-Free Diet, [843](#)  
 Gluten-Sensitive Enteropathy, [844](#)  
 Gluteus, [844](#)  
 Gluteus Maximus Muscle, [844](#)  
 Gluteus Medius Muscle, [844](#)  
 Gluteus Minimius Muscle, [844](#)  
 Glyburide, [844](#)  
 Glycaemia, [844](#)  
 Glycated Hemoglobin, [844](#)  
 Glycation, [844](#)  
 Glycemia, [844](#)  
 Glycemic Control, [844](#)  
 Glycemic Index, [844](#)  
 Glycemic Index And Diabetes, [844](#)  
 Glycemic Load, [845](#)  
 Glyceraldehyde, [845](#)  
 Glyceride, [845](#)  
 Glycerin, [845](#)  
 Glycerol, [845](#)  
 Glycerol Kinase, [845](#)  
 Glycerophosphates, [845](#)  
 Glycerophospholipids, [845](#)  
 Glyceryl Trinitrate, [845](#)  
 Glycerylphosphorylcholine, [845](#)  
 Glycine, [845](#)  
 Glycine Receptor, [845](#)  
 Glycocalyx, [845](#)  
 Glycogen, [845](#)  
 Glycogen Storage Disease, [845](#)  
 Glycogen Storage Disease Type 0 (GSD-0), [846](#)  
 Glycogen Storage Diseases, [846](#)  
 Glycogenesis, [846](#)  
 Glycogenolysis, [846](#)  
 Glycohemoglobin, [846](#)  
 Glycolipid, [846](#)  
 Glycolysis, [846](#)  
 Glycopeptide, [846](#)  
 Glycophorin, [846](#)  
 Glycoprotein, [846](#)  
 Glycopyrrolate, [846](#)  
 Glycosaminoglycan, [846](#)  
 Glycosaminoglycans, [846](#)  
 Glycoside, [846](#)  
 Glycosphingolipid, [846](#)  
 Glycosuria, [847](#)  
 Glycosylated Hemoglobin, [847](#)  
 Glycosylation, [847](#)  
 Glycosylphosphatidylinositol, [847](#)  
 Glycosyltransferase, [847](#)  
 Glycyrrhetic Acid, [847](#)  
 Glycyrrhizic Acid, [847](#)  
 Glyphosate, [847](#)  
 Gm, [847](#)  
 GN, [847](#)  
 Gnathic, [847](#)  
 Gnathion, [847](#)  
 Gnathitis, [847](#)  
 Gnathoplasty, [847](#)  
 Gnathostoma, [847](#)  
 Gnathostoma Spinigerum, [847](#)  
 Gnathostomiasis, [848](#)  
 Gnawing Pain, [848](#)  
 GNE Myopathy, [848](#)  
 GnRH, [848](#)  
 GnRH Agonist, [848](#)  
 GnRH Antagonist, [848](#)  
 GnRH Antagonists, [848](#)  
 Goal Attainment Scaling, [848](#)  
 Goal Setting, [848](#)  
 Goals-Of-Care Discussion, [848](#)  
 Goblet Cell, [848](#)  
 Goblet Cell Adenocarcinoma, [848](#)  
 Goblet Cells, [848](#)  
 Goeckerman Regimen, [849](#)  
 Goiter, [849](#)

---

---

Goitre, 849  
 Goitrogenic Foods, 849  
 Goitrous Autoimmune Thyroiditis, 849  
 Gold, 849  
 Gold Alloy, 849  
 Gold Alloys, 849  
 Gold Therapy, 849  
 Golden Hour, 849  
 Goldenhar Syndrome, 849  
 Goldmann Tonometry, 849  
 Golfer Elbow, 849  
 Golfer's Elbow, 849  
 Golgi Apparatus, 850  
 Golgi Cisternae, 850  
 Golgi Tendon Organ, 850  
 Gomori's Stain, 850  
 Gomphosis, 850  
 Gonad, 850  
 Gonad Disorder, 850  
 Gonadal Agenesis, 850  
 Gonadal Dysgenesis, 850  
 Gonadal Vein, 850  
 Gonadoblastoma, 850  
 Gonadorelin, 850  
 Gonadotrophin, 850  
 Gonadotrophin-Releasing Factors, 850  
 Gonadotrophins, 850  
 Gonadotrophs, 851  
 Gonadotropin, 851  
 Gonadotropin Hormone, 851  
 Gonadotropin-Releasing Hormone, 851  
 Gonadotropins, 851  
 Gonads, 851  
 Gondoic Acid, 851  
 Goniometer, 851  
 Gonioscopy, 851  
 Gonococcal, 851  
 Gonococcal Arthritis, 851  
 Gonococcal Pharyngitis, 851  
 Gonococcal Urethritis, 851  
 Gonococcus, 851  
 Gonorrhea, 851  
 Gonorrhoea, 852  
 Good Samaritan, 852  
 Good Samaritan Law, 852  
 Goodell Sign, 852  
 Goodpasture Syndrome, 852  
 Goodpasture's Syndrome, 852  
 Goose Bump, 852  
 Goose-Skin, 852  
 Gooseflesh, 852  
 Gorlin Syndrome, 852  
 Gorlin's Syndrome, 852  
 Goserelin, 852  
 Gottron Papules, 852  
 Gottron Sign, 852  
 Gougerot-Carteaud Syndrome, 852  
 Gout, 852  
 Gout Flare, 853  
 Gout Nodulosis, 853  
 Gout Suppressants, 853  
 Gouty Tophus, 853  
 Gower Syndrome, 853  
 Gowers Sign, 853  
 Graafian Follicle, 853  
 GRACE Score, 853  
 Gracilis Muscle, 853  
 Grade, 853  
 Grade IV Astrocytoma, 853  
 Graded Potentials, 854  
 Gradenigo Syndrome, 854  
 Graefe's Sign, 854  
 Graft, 854  
 Graft Versus Host, 854  
 Graft Versus Host Disease, 854  
 Graft-Versus-Host Disease, 854  
 Graft-Versus-Host Reaction, 854  
 Grafting, 854  
 Graham-Little-Piccardi-Lasseur Syndrome, 854  
 Gram, 854  
 Gram Mass Unit, 854  
 Gram Metric Unit, 854  
 Gram Stain, 854  
 Gram Stain Of Skin Lesion, 855  
 Gram Stain Of Tissue Biopsy, 855  
 Gram's Stain, 855  
 Gram-Negative, 855  
 Gram-Negative Bacilli, 855  
 Gram-Negative Bacteria, 855  
 Gram-Negative Cocci And Coccobacilli, 855  
 Gram-Negative Folliculitis, 855  
 Gram-Negative Meningitis, 855  
 Gram-Negative Toe Web Infection, 855  
 Gram-Positive, 855  
 Gram-Positive Bacilli, 855  
 Gram-Positive Bacteria, 855  
 Gram-Positive Cocci, 855  
 Gramicidin, 856  
 Grand Multipara, 856  
 Grand Rounds, 856  
 Granisetron, 856  
 Granisetron Hydrochloride, 856  
 Granular Casts, 856  
 Granular Cell Tumor Of The Breast, 856  
 Granular Cell Tumors, 856  
 Granular Leukocyte, 856  
 Granular Parakeratosis, 856  
 Granulation, 856

---

---

Granulation Tissue, 856  
Granulicatella, 856  
Granulicatella Adiacens, 856  
Granulicatella Elegans, 857  
Granulocyte, 857  
Granulocyte Colony-Stimulating Factor, 857  
Granulocyte Count, 857  
Granulocytic Sarcoma, 857  
Granulocytopenia, 857  
Granuloma, 857  
Granuloma Annulare, 857  
Granuloma Fungoides, 857  
Granuloma Gluteale Infantum, 857  
Granuloma Inguinale, 857  
Granulomatosis, 857  
Granulomatosis With Polyangiitis, 857  
Granulomatous Amebic Encephalitis, 858  
Granulomatous Colitis, 858  
Granulomatous Inflammation, 858  
Granulomatous Mastitis, 858  
Granulosa Cell, 858  
Granulosa Cell Tumor, 858  
Granulosa Cell Tumors, 858  
Granulosa Tumor, 858  
Granzyme, 858  
Grasp Reflex, 858  
Grass Allergy, 858  
Grass And Weed Killer Poisoning, 858  
Grave Wax, 858  
Graves Disease, 858  
Graves Ophthalmopathy, 859  
Gravida, 859  
Gravidity, 859  
Gray, 859  
Gray Baby Syndrome, 859  
Gray Matter, 859  
Great Saphenous Vein, 859  
Greater Curvature, 859  
Greater Omentum, 859  
Greater Trochanter, 859  
Greater Trochanteric Pain Syndrome, 859  
Greater Vestibular Glands, 859  
Green Blindness, 859  
Green Nail Syndrome, 860  
Green Stool, 860  
Greenstick Fracture, 860  
Grenz Zone, 860  
Grey Ankle Phenomenon, 860  
Grey Matter, 860  
Grey Turner Sign, 860  
Grey Turner's Sign, 860  
Grid Implant, 860  
Grief, 860  
Grief and Bereavement, 860  
Grimontia Hollisae, 860  
Grip Strength, 860  
Gripping, 860  
Griscelli Syndrome, 861  
Griseofulvin, 861  
Groin, 861  
Groin Lump, 861  
Groin Pain, 861  
Grommet Tube, 861  
Groove, 861  
Grooved Tongue, 861  
Gross Anatomy, 861  
Gross Examination, 861  
Gross Hematuria, 861  
Gross Motor Control, 861  
Gross Motor Skills, 861  
Ground Itch, 862  
Ground Substance, 862  
Ground-Glass Opacity, 862  
Group 1 Pulmonary Hypertension (PAH), 862  
Group 2 Pulmonary Hypertension, 862  
Group 3 Pulmonary Hypertension, 862  
Group 4 Pulmonary Hypertension, 862  
Group A Strep, 862  
Group A Streptococcal Infections, 862  
Group A Streptococcus, 862  
Group B Strep, 862  
Group B Strep Disease, 862  
Group B Streptococcal Disease, 863  
Group B Streptococcal Infection In Pregnancy, 863  
Group B Streptococcal Infections in Newborns, 863  
Group B Streptococcal Neonatal Sepsis, 863  
Group B Streptococcal Septicemia of the Newborn, 863  
Group B Streptococcus, 863  
Group B Streptococcus in Pregnancy, 863  
Group Practice, 863  
Group Therapy, 863  
Grover Disease, 863  
Grover's Disease, 863  
Growing Pains, 863  
Growth, 864  
Growth Chart, 864  
Growth Cone, 864  
Growth Disorder, 864  
Growth Factor, 864  
Growth Failure, 864  
Growth Faltering, 864  
Growth Hormone, 864  
Growth Hormone Deficiency, 864  
Growth Hormone Deficiency in Adults, 864  
Growth Hormone Physiology, 864  
Growth Hormone Stimulation Test, 864  
Growth Hormone Suppression Test, 864  
Growth Hormone Test, 865

---

---

Growth Hormone-Releasing Hormone, 865  
 Growth Inhibitor, 2122  
 Growth Medium, 865  
 Growth Plate, 865  
 Growth Plate Fracture, 865  
 Growth Plate Fractures, 865  
 Growth Retardation, 865  
 Growth Scan, 865  
 Gtp Binding, 865  
 Gtt., 865  
 Guaifenesin, 865  
 Guanfacine, 865  
 Guanine, 865  
 Guanosine Triphosphate, 865  
 Guanosinetriphosphatase, 866  
 Guanylate Cyclase, 866  
 Guarana, 866  
 Guardianship, 866  
 Guarding, 866  
 Guenther Porphyria, 866  
 Guevedoces, 866  
 Guide Device, 866  
 Guided Imagery, 866  
 Guided Tissue Regeneration, 866  
 Guillain-Barre Syndrome, 866  
 Guillain-Barré Syndrome, 866  
 Guillaume Dupuytren, 866  
 Guinea Worm, 866  
 Guinea Worm Infection, 866  
 Gulf War Syndrome, 867  
 Gullet, 867  
 Gum, 867  
 Gum Biopsy, 867  
 Gum Disease, 867  
 Gum Problems, 867  
 Gumma, 867  
 Gummata, 867  
 Gummy Smile, 867  
 Gums, 867  
 Gunn Sign, 867  
 Gunshot Wound, 867  
 Gunther Disease, 868  
 Gustation, 868  
 Gustatory, 868  
 Gustatory Lacrimation, 868  
 Gustatory Sweating, 868  
 Gut, 868  
 Gut Epithelium, 868  
 Gut Microbiome, 868  
 Gut-Brain Axis, 868  
 Guthrie Test, 868  
 Guttae, 868  
 Guttate Psoriasis, 868  
 Guttate Scleroderma, Lichen Sclerosus Type, 1137  
 Guttural, 868  
 Guyon Canal Syndrome, 868  
 GVHD, 869  
 Gymnast Wrist, 869  
 Gynaecologist, 869  
 Gynaecology, 869  
 Gynaecomastia, 869  
 Gynatresia, 869  
 Gynecic, 869  
 Gynecoid Pelvis, 869  
 Gynecologic Pain, 869  
 Gynecologic Tumor Markers, 869  
 Gynecological Examination, 869  
 Gynecologist, 869  
 Gynecology, 869  
 Gynecomastia, 869  
 Gynecomastia, Male, 870  
 Gyrate Atrophy, 870  
 Gyrus, 870  
  
 H And E Staining, 871  
 H And H, 871  
 H And P, 871  
 H Influenzae Meningitis, 871  
 H-Reflex, 871  
 H. Pylori, 871  
 H. Pylori Stool Antigen, 871  
 H1N1 Flu, 871  
 H1N1 Influenza, 871  
 H1N2, 871  
 H2 Blockers, 872  
 H2 Receptor Antagonists Overdose, 872  
 H5N1, 872  
 Habenula, 872  
 Haberland Syndrome (Encephalocraniocutaneous Lipomatosis), 872  
 Habit, 872  
 Habituation, 872  
 Habitus, 872  
 HAE, 872  
 Haem, 872  
 Haemangioma, 872  
 Haemarthrosis, 872  
 Haematemesis, 872  
 Haematinics, 872  
 Haematocoele, 873  
 Haematocolpos, 873  
 Haematocrit, 873  
 Haematogenous, 873  
 Haematologist, 873  
 Haematoma, 873  
 Haematometra, 873  
 Haematomyelia, 873  
 Haematopericardium, 873

---



---

Haematopoiesis, 873  
 Haematoporphyrin, 873  
 Haematosalpinx, 873  
 Haematuria, 873  
 Haemochromatosis, 873  
 Haemodialysis, 873  
 Haemoflagellate, 874  
 Haemoglobin, 874  
 Haemoglobinopathy, 874  
 Haemoglobinuria, 874  
 Haemolysis, 874  
 Haemolytic Anaemia, 874  
 Haemolytic Disease Of The Newborn, 874  
 Haemoperitoneum, 874  
 Haemophilia, 874  
 Haemophilia A, 874  
 Haemophilus, 874  
 Haemophilus Aegyptius, 874  
 Haemophilus Ducreyi, 874  
 Haemophilus Haemolyticus, 874  
 Haemophilus Infections, 874  
 Haemophilus Influenzae, 875  
 Haemophilus Parainfluenzae, 875  
 Haemophilus Vaccine, 2122  
 Haemoptysis, 875  
 Haemorrhage, 875  
 Haemorrhagic Fever, 875  
 Haemorrhagic Fever With Renal Syndrome, 875  
 Haemorrhagic Stroke, 875  
 Haemorrhoids, 875  
 Haemosiderin, 875  
 Haemosiderosis, 875  
 Haemostasis, 875  
 Haemostat, 875  
 Haemothorax, 875  
 Haff Disease, 875  
 Hafnia, 875  
 Hafnia Alvei, 876  
 Hageman Factor, 876  
 Haglund Deformity, 876  
 Hailey-Hailey Disease, 876  
 Hair, 876  
 Hair Bleach Poisoning, 876  
 Hair Bulb, 876  
 Hair Cells, 876  
 Hair Disease, 876  
 Hair Dye Poisoning, 876  
 Hair Follicle, 876  
 Hair Loss, 876  
 Hair Loss Classification, 876  
 Hair Removal, 876  
 Hair Shaft, 877  
 Hair Spray Poisoning, 877  
 Hair Straightener Poisoning, 877  
 Hair Tonic Poisoning, 877  
 Hair Transplant, 877  
 Hair-Ball, 877  
 Hair-Pulling Syndrome, 877  
 Hairy Cell Leukaemia, 877  
 Hairy Cell Leukemia, 877  
 Hairy Leukoplakia, 877  
 Hairy Tongue, 877  
 Hajdu-Cheney Syndrome, 877  
 Halcinonide, 877  
 Haldane Effect, 877  
 Half-Life, 878  
 Halitosis, 878  
 Hallermann's Syndrome, 878  
 Hallpike Test, 878  
 Hallucination, 878  
 Hallucinations, 878  
 Hallucinogen, 878  
 Hallux, 878  
 Hallux Rigidus, 878  
 Hallux Valgus, 878  
 Hallux Varus, 878  
 Halo Nevus, 878  
 Halo Sign, 878  
 Halogen, 879  
 Haloperidol, 879  
 Halothane, 879  
 Halushi-Behcet's Syndrome, 879  
 Hamamelis, 879  
 Hamartoma, 879  
 Hamartomatous Polyposis, 879  
 Hamate, 879  
 Hamate Bone, 879  
 Hamman Sign, 879  
 Hamman's Sign, 879  
 Hamman-Rich Syndrome, 879  
 Hammer, 879  
 Hammer Toe, 879  
 Hammer Toe Repair, 879  
 Hammer, Claw and Mallet Toes, 1112  
 Hammertoe, 879  
 Hammertoes, 880  
 Hampton Hump, 880  
 Hampton Line, 880  
 Hampton's Hump, 880  
 Hamstring, 880  
 Hamstring Injury, 880  
 Hamstring Muscles, 880  
 Hamstring Strain, 880  
 Hand, 880  
 Hand Deformity, 880  
 Hand Foot And Mouth Disease, 880  
 Hand Hygiene, 880  
 Hand Infections, 880

---

---

Hand Microsurgery, 880  
Hand of Benediction, 881  
Hand Or Foot Spasms, 881  
Hand Pain, 881  
Hand Prolapse, 881  
Hand X-Ray, 881  
Hand-Arm Vibration Syndrome, 881  
Hand-Foot Syndrome, 881  
Hand-Foot-And-Mouth Disease, 881  
Hand-Foot-And-Mouth Syndrome, 881  
Hand-Foot-Mouth Disease, 881  
Hand-Held Retractor, 881  
Hand-Schuller-Christian Disease, 882  
Hand-Schüller-Christian Disease, 882  
Handling Sharps And Needles, 882  
Handwashing, 882  
HANE, 882  
Hang-Nail, 882  
Hanging And Strangulation Signs, 882  
Hanging Injuries and Strangulation, 882  
Hangman Fracture, 882  
Hangman's Fracture, 882  
Hangover, 882  
Hansen Disease, 882  
Hantavirus, 882  
Hantavirus Cardiopulmonary Syndrome, 883  
Hantavirus Infection, 883  
Hantavirus Pulmonary Syndrome, 883  
Haploid, 883  
Haploidentical Donor, 883  
Haploidy, 883  
Haploinsufficiency, 883  
Haplotype, 883  
Hapten, 883  
Haptoglobin, 883  
Haptoglobin Analyses, 883  
Haptoglobin Blood Test, 883  
Haptoglobin Decreased, 883  
Hard Palate, 883  
Harlequin Color Change, 884  
Harlequin Foetus, 884  
Harlequin Ichthyosis, 884  
Harm Reduction, 884  
Harm Reduction Therapy, 884  
Harmless Murmur, 884  
Harrison's Sulcus, 884  
Hartmann Pouch, 884  
Hartmann Procedure, 884  
Hartnup Disease, 884  
Hartnup Disorder, 884  
HAS-BLED Score, 884  
Hashimoto Disease, 884  
Hashimoto Encephalopathy, 885  
Hashimoto Thyroiditis, 885  
Hashimoto's Disease, 885  
Hashimoto's Thyroiditis, 885  
Hashish, 885  
Hashitoxicosis, 885  
Hassall Corpuscles, 885  
Hassall's Corpuscles, 885  
Haustrea, 885  
Haverhill Fever, 885  
Haversian Canal, 885  
Hawkins Test, 885  
Hay Fever, 886  
Hazardous Materials, 886  
Hazardous Substance, 886  
Hb, 886  
HbA1c, 886  
HCG, 886  
HCG In Urine, 886  
Hct, 886  
HCV PCR, 886  
HCV-Associated Rheumatic Disease, 886  
HD, 886  
HDL, 886  
HDL Cholesterol, 886  
HDL Test, 886  
HDR Syndrome, 887  
Head, 887  
Head And Face Reconstruction, 887  
Head And Neck Cancer, 887  
Head And Neck Oncologic Surgery, 887  
Head Circumference, 887  
Head CT Scan, 887  
Head Impulse Test, 887  
Head Injury, 887  
Head Injury First Aid, 887  
Head Lice, 887  
Head Movement, 888  
Head MRI, 888  
Head Of The Epididymis, 888  
Head Thrust Test, 888  
Head Trauma, 888  
Head-Lice, 888  
Headache, 888  
Headache and Migraine, 888  
Headache Red Flags, 888  
Headaches, Tension-type, 1900  
Headaches/Migraines, 888  
Healing, 888  
Health Anxiety Disorder, 888  
Health Behavior, 888  
Health Behavior Change, 888  
Health Behavior Motivation, 889  
Health Care, 889  
Health Care Agents, 889  
Health Care Facility, 889

---

---

Health Care Personnel, 889  
 Health Care Proxy, 889  
 Health Checkup, 889  
 Health Disparities, 889  
 Health Disparity, 889  
 Health Education, 889  
 Health Inequity, 889  
 Health Information Technology, 889  
 Health Literacy, 889  
 Health Maintenance Organization, 889  
 Health Promotion, 890  
 Health Risk, 890  
 Health Risks Of Alcohol Use, 890  
 Health Surveillance, 890  
 Health Survey, 890  
 Health-Related QoL, 890  
 Healthcare Associated Infection, 890  
 Healthcare Provider, 890  
 Healthcare Proxy, 890  
 Healthy Aging, 890  
 Hearing, 890  
 Hearing Aid, 890  
 Hearing Aids And Devices, 890  
 Hearing Disorder, 890  
 Hearing Impairment, 891  
 Hearing Loss, 891  
 Hearing Loss And Music, 891  
 Hearing Loss: Sudden Deafness, 891  
 Hearing Screening, 891  
 Hearing Test, 891  
 Heart, 891  
 Heart And Vascular Services, 891  
 Heart Aneurysm, 891  
 Heart Arrhythmia, 891  
 Heart Attack, 891  
 Heart Attack First Aid, 891  
 Heart Attack Prevention, 891  
 Heart Block, 892  
 Heart Bypass Surgery, 892  
 Heart Catheterization, 892  
 Heart Chambers, 892  
 Heart Contraction, 892  
 Heart CT Scan, 892  
 Heart Defect, 892  
 Heart Disease, 892  
 Heart Disease And Depression, 892  
 Heart Disease And Intimacy, 892  
 Heart Disease And Women, 892  
 Heart Failure, 892  
 Heart Failure Due To Valvular Heart Disease, 892  
 Heart Failure In Children, 892  
 Heart Failure Stages, 893  
 Heart Failure With Mildly Reduced Ejection Fraction, 893  
 Heart Failure With Preserved Ejection Fraction, 893  
 Heart Failure With Reduced Ejection Fraction, 893  
 Heart Injury, 893  
 Heart MRI, 893  
 Heart Murmur, 893  
 Heart Murmur Entry, 893  
 Heart Murmurs, 893  
 Heart Muscle, 893  
 Heart Neoplasm, 893  
 Heart Pacemaker, 893  
 Heart Palpitations, 893  
 Heart PET Scan, 893  
 Heart Rate, 893  
 Heart Rate Variability, 894  
 Heart Rhythm, 894  
 Heart Rhythm Disorders, 894  
 Heart Septum, 894  
 Heart Sounds, 894  
 Heart Surgery, 894  
 Heart Tamponade, 894  
 Heart Transplant, 894  
 Heart Transplantation, 894  
 Heart Valve, 894  
 Heart Valve Disease, 894  
 Heart Valve Disorder, 894  
 Heart Valve Prosthesis, 894  
 Heart Valve Surgery, 894  
 Heart Ventricle, 895  
 Heart Ventricles, 895  
 Heart-Lung Machine, 895  
 Heart-Smart Substitutions, 895  
 Heartbeat, 895  
 Heartburn, 895  
 Heartburn/GERD, 895  
 Heartland Virus, 895  
 Heat, 895  
 Heat Cramps, 895  
 Heat Emergencies, 895  
 Heat Exhaustion, 895  
 Heat Intolerance, 895  
 Heat Pack, 896  
 Heat Prostration, 896  
 Heat Rash, 896  
 Heat Shock Protein, 896  
 Heat Stress, 896  
 Heat Stroke, 896  
 Heavy Menstrual Bleeding, 896  
 Heavy Metal Poisoning, 896  
 Heberden Node, 896  
 Heberden Nodes, 896  
 Heberden's Node, 896  
 Heberden's Nodes, 896  
 Hecht's Pneumonia, 896  
 Hedgehog Proteins, 896

---

---

Heel, [897](#)  
Heel Pain, [897](#)  
Heel Spur, [897](#)  
Heel-to-Shin Test, [897](#)  
Heerfordt Syndrome, [897](#)  
Hegar's Dilators, [897](#)  
Hegar's Sign, [897](#)  
Height, [897](#)  
Heimlich Maneuver, [897](#)  
Heimlich Maneuver On Self, [897](#)  
Heimlich Manoeuvre, [897](#)  
Heine-Medin Disease, [897](#)  
Heinz Bodies, [897](#)  
Hejama, [898](#)  
Helcococcus, [898](#)  
Helcococcus Kunzii, [898](#)  
Helcococcus Pyogenes, [898](#)  
Helicase, [898](#)  
Helicobacter, [898](#)  
Helicobacter Cinaedi, [898](#)  
Helicobacter Fennelliae, [898](#)  
Helicobacter Pylori, [898](#)  
Helicobacter Pylori Eradication, [898](#)  
Helicobacter Pylori Infection, [898](#)  
Helicobacter Pylori-Associated Active Gastritis Pathology, [898](#)  
Helicobacteraceae, [898](#)  
Helicotrema, [898](#)  
Heliotherapy, [899](#)  
Heliotrope Rash, [899](#)  
Helium, [899](#)  
Helix, [899](#)  
Heller Myotomy, [899](#)  
Heller's Operation, [899](#)  
HELLP syndrome, [899](#)  
Helminth, [899](#)  
Helminthiasis, [899](#)  
Helminths, [899](#)  
Help Prevent Hospital Errors, [899](#)  
Helper T Cell, [899](#)  
Helper T Cells, [899](#)  
Helper Virus, [899](#)  
HELPERR Mnemonic, [900](#)  
Hemagglutination Test, [900](#)  
Hemagglutinin, [900](#)  
Hemangioblastoma, [900](#)  
Hemangioendothelioma, [900](#)  
Hemangioma, [900](#)  
Hemangiopericytoma, [900](#)  
Hemangiosarcoma, [900](#)  
Hemarthrosis, [900](#)  
Hematemesis, [900](#)  
Hematidrosis, [900](#)  
Hematocele, [900](#)  
Hematochezia, [900](#)  
Hematocolpos, [901](#)  
Hematocrit, [901](#)  
Hematogenous, [901](#)  
Hematogenous Osteomyelitis, [901](#)  
Hematologic Agents, [901](#)  
Hematologic Changes In Pregnancy, [901](#)  
Hematologic Malignancy, [901](#)  
Hematologic Response, [901](#)  
Hematological Disease, [901](#)  
Hematologist, [901](#)  
Hematology, [901](#)  
Hematology-Oncology, [901](#)  
Hematoma, [901](#)  
Hematometra, [901](#)  
Hematometrocolpos, [901](#)  
Hematomyelia, [902](#)  
Hematophobia, [902](#)  
Hematopoiesis, [902](#)  
Hematopoietic Growth Factor, [902](#)  
Hematopoietic Neoplasm, [902](#)  
Hematopoietic Stem Cell Transplantation, [902](#)  
Hematopoietic System, [902](#)  
Hematoporphyrin Derivative, [902](#)  
Hematosalpinx, [902](#)  
Hematosi, [902](#)  
Hematospermia, [902](#)  
Hematoxylin, [902](#)  
Hematuria, [902](#)  
Heme, [902](#)  
Hemeralopia, [903](#)  
Hemianesthesia, [903](#)  
Hemianopia, [903](#)  
Hemianopsia, [903](#)  
Hemiballism, [903](#)  
Hemiballismus, [903](#)  
Hemicolectomy, [903](#)  
Hemidesmosome, [903](#)  
Hemidiaphragm, [903](#)  
Hemifacial Atrophy, [903](#)  
Hemifacial Microsomia, [903](#)  
Hemifacial Spasm, [903](#)  
Hemihypertrophy, [903](#)  
Hemilaryngectomy, [903](#)  
Hemimandibulectomy, [903](#)  
Hemimegalencephaly, [904](#)  
Hemimelia, [904](#)  
Hemineglect, [904](#)  
Hemiparesis, [904](#)  
Hemiparesis with Conjugate Gaze Deviation, [904](#)  
Hemipelvectomy, [904](#)  
Hemiplegia, [904](#)  
Hemispatial Neglect, [904](#)  
Hemispherectomy, [904](#)

---

---

Hemisternebra, [904](#)  
Hemithorax, [904](#)  
Hemivertebrae, [904](#)  
Hemizygous, [904](#)  
Hemobilia, [904](#)  
Hemochromatosis, [905](#)  
Hemochromatosis Cardiomyopathy, [905](#)  
Hemocyanin, [905](#)  
Hemocytometer, [905](#)  
Hemodialysis, [905](#)  
Hemodialysis Access Care, [905](#)  
Hemodialysis Access Procedures, [905](#)  
Hemodialysis Clearance, [905](#)  
Hemodialysis Solution, [905](#)  
Hemodynamics, [905](#)  
Hemofiltration, [905](#)  
Hemoglobin, [905](#)  
Hemoglobin A, [906](#)  
Hemoglobin A1c, [906](#)  
Hemoglobin A1C Variability, [906](#)  
Hemoglobin C, [906](#)  
Hemoglobin C Disease, [906](#)  
Hemoglobin Electrophoresis, [906](#)  
Hemoglobin H, [906](#)  
Hemoglobin Increased, [906](#)  
Hemoglobin M, [906](#)  
Hemoglobin Normal Values, [906](#)  
Hemoglobin S, [906](#)  
Hemoglobin SC Disease, [906](#)  
Hemoglobin Trait, [906](#)  
Hemoglobinopathies, [907](#)  
Hemoglobinopathies: Thalassemias, [907](#)  
Hemoglobinopathy, [907](#)  
Hemoglobinuria, [907](#)  
Hemoglobinuria Test, [907](#)  
Hemolysin, [907](#)  
Hemolysis, [907](#)  
Hemolytic, [907](#)  
Hemolytic Anemia, [907](#)  
Hemolytic Anemia From Toxins, [907](#)  
Hemolytic Anemias: Autoimmune, [907](#)  
Hemolytic Anemias: Hereditary, [907](#)  
Hemolytic Crisis, [907](#)  
Hemolytic Disease Of The Newborn, [907](#)  
Hemolytic Index, [908](#)  
Hemolytic Jaundice, [908](#)  
Hemolytic Transfusion Reaction, [908](#)  
Hemolytic Uremic Syndrome, [908](#)  
Hemolytic-Uremic Syndrome, [908](#)  
Hemoperfusion, [908](#)  
Hemopericardium, [908](#)  
Hemoperitoneum, [908](#)  
Hemopexin, [908](#)  
Hemophagocytic Lymphohistiocytosis, [908](#)  
Hemophagocytosis Of Sickle Cells, [908](#)  
Hemophilia, [908](#)  
Hemophilia A, [909](#)  
Hemophilia B, [909](#)  
Hemophilic Arthritis, [909](#)  
Hemophilus, [909](#)  
Hemophilus Ducreyi, [909](#)  
Hemophobia, [909](#)  
Hemopoietic Stem Cell, [909](#)  
Hemopoietic Tissue, [909](#)  
Hemoprotein, [909](#)  
Hemoptysis, [909](#)  
Hemorrhage, [909](#)  
Hemorrhagic, [909](#)  
Hemorrhagic Anemia, [909](#)  
Hemorrhagic Disorder, [909](#)  
Hemorrhagic Shock, [909](#)  
Hemorrhagic Stroke, [910](#)  
Hemorrhagic Telangiectasia, Hereditary (HHT), [921](#)  
Hemorrhagic Thyroid Nodule, [910](#)  
Hemorrhoid, [910](#)  
Hemorrhoid Surgery, [910](#)  
Hemorrhoidal Hemorrhage, [910](#)  
Hemorrhoidectomy, [910](#)  
Hemorrhoids, [910](#)  
Hemosiderin, [910](#)  
Hemosiderosis, [910](#)  
Hemosiderotic Synovitis, [910](#)  
Hemospermia, [911](#)  
Hemostasis, [911](#)  
Hemostasis And Coagulation, [911](#)  
Hemostasis Valve, [911](#)  
Hemostat, [911](#)  
Hemostatic Agent, [911](#)  
Hemothorax, [911](#)  
Hemovac Drain, [911](#)  
Henderson-Hasselbalch Equation, [911](#)  
Hendra Virus, [911](#)  
Henipavirus, [911](#)  
Henle Loop, [911](#)  
Henna, [911](#)  
Henoch-Schonlein Purpura, [912](#)  
Henoch-Schonlein Purpura (HSP), [912](#)  
Henoch-SchöNlein Purpura, [912](#)  
HEPA Filter, [912](#)  
Hepar, [912](#)  
Heparan Sulfate, [912](#)  
Heparin, [912](#)  
Heparin Binding, [912](#)  
Heparin Resistance, [912](#)  
Heparin-Induced Thrombocytopenia, [912](#)  
Hepatectomy, [912](#)  
Hepatic, [912](#)  
Hepatic Adenoma, [912](#)

---

---

Hepatic Artery, [913](#)  
 Hepatic Artery Embolization, [913](#)  
 Hepatic Artery Occlusion, [913](#)  
 Hepatic Biopsy, [913](#)  
 Hepatic Coma, [913](#)  
 Hepatic Cyst, [913](#)  
 Hepatic Dose Adjustment, [913](#)  
 Hepatic Drug Metabolism, [2122](#)  
 Hepatic Duct, [913](#)  
 Hepatic Encephalopathy, [913](#)  
 Hepatic Fissure, [913](#)  
 Hepatic Flexure, [913](#)  
 Hepatic Granulomas, [913](#)  
 Hepatic Hemangioma, [913](#)  
 Hepatic Hydrothorax, [914](#)  
 Hepatic Ischemia, [914](#)  
 Hepatic Lamina, [914](#)  
 Hepatic Necrosis, [914](#)  
 Hepatic Portal System, [914](#)  
 Hepatic Steatosis, [914](#)  
 Hepatic Surgery, [914](#)  
 Hepatic Synthetic Dysfunction, [914](#)  
 Hepatic Trauma, [914](#)  
 Hepatic Vein, [914](#)  
 Hepatic Vein Obstruction, [914](#)  
 Hepatitis, [914](#)  
 Hepatitis A, [914](#)  
 Hepatitis A Virus, [914](#)  
 Hepatitis B, [915](#)  
 Hepatitis B Vaccine, [2122](#)  
 Hepatitis B Vaccine (Birth Dose), [2122](#)  
 Hepatitis B Virus, [915](#)  
 Hepatitis B Window Period, [915](#)  
 Hepatitis C, [915](#)  
 Hepatitis C Virus, [915](#)  
 Hepatitis D, [915](#)  
 Hepatitis D Virus, [915](#)  
 Hepatitis E, [915](#)  
 Hepatitis E Virus, [916](#)  
 Hepatitis Serology, [916](#)  
 Hepatitis Virus, [916](#)  
 Hepatitis Virus Panel, [916](#)  
 Hepatitis Viruses, [916](#)  
 Hepatobiliary, [916](#)  
 Hepatobiliary Scintigraphy, [916](#)  
 Hepatobiliary System, [917](#)  
 Hepatoblastoma, [917](#)  
 Hepatocellular Adenoma, [917](#)  
 Hepatocellular Carcinoma, [917](#)  
 Hepatocellular Jaundice, [917](#)  
 Hepatocerebral Degeneration, [917](#)  
 Hepatocyte, [917](#)  
 Hepatoduodenal Ligament, [917](#)  
 Hepatojugular Reflux, [918](#)  
 Hepatolenticular Degeneration, [918](#)  
 Hepatologist, [918](#)  
 Hepatology, [918](#)  
 Hepatoma, [918](#)  
 Hepatomegaly, [918](#)  
 Hepatopulmonary Syndrome, [918](#)  
 Hepatorenal Recess, [918](#)  
 Hepatorenal Syndrome, [919](#)  
 Hepatosplenomegaly, [919](#)  
 Hepatotoxicity, [919](#)  
 Hepatovirus, [919](#)  
 Hepcidin, [919](#)  
 Heptachlor, [919](#)  
 HER2, [919](#)  
 HER2 Inhibitor, [2122](#)  
 HER2-Positive, [920](#)  
 HER2-Positive Breast Cancer, [920](#)  
 Herbal Supplement, [920](#)  
 Herbaspirillum, [920](#)  
 Herbicide, [920](#)  
 Herd Immunity, [920](#)  
 Hereditary, [920](#)  
 Hereditary Amyloidosis, [920](#)  
 Hereditary and Acquired Ichthyosis Vulgaris, [920](#)  
 Hereditary Angioedema, [920](#)  
 Hereditary Cancer, [920](#)  
 Hereditary Coproporphyrria, [920](#)  
 Hereditary Diffuse Gastric Cancer, [920](#)  
 Hereditary Diseases, [921](#)  
 Hereditary Elliptocytosis, [921](#)  
 Hereditary Fructose Intolerance, [921](#)  
 Hereditary Hemorrhagic Telangiectasia, [921](#)  
 Hereditary Hyperbilirubinemia, [921](#)  
 Hereditary Leiomyomatosis And Renal Cell Cancer, [921](#)  
 Hereditary Motor and Sensory Neuropathy, [921](#)  
 Hereditary Multiple Exostoses, [921](#)  
 Hereditary Neuropathy with Liability to Pressure Palsies, [921](#)  
 Hereditary Nonpolyposis Colorectal Cancer (Lynch Syndrome), [921](#)  
 Hereditary Nonspherocytic Hemolytic Anemia (HNHA), [922](#)  
 Hereditary Ovalocytosis, [922](#)  
 Hereditary Paraganglioma-Pheochromocytoma Syndrome, [922](#)  
 Hereditary Periodic Fever Syndromes, [922](#)  
 Hereditary Persistence of Fetal Hemoglobin, [922](#)  
 Hereditary Spastic Paraplegia, [922](#)  
 Hereditary Spherocytic Anemia, [922](#)  
 Hereditary Spherocytosis, [922](#)  
 Hereditary Urea Cycle Abnormality, [922](#)  
 Heredity, [922](#)  
 Heredo-Ataxia, [922](#)  
 Heredopolyneuritis, [922](#)

---



---

Hering-Breuer Reflex, [923](#)  
Hermanski-Pudlak Syndrome, [923](#)  
Hermansky-Pudlak Syndrome, [923](#)  
Hernia, [923](#)  
Hernia Mesh, [923](#)  
Hernia Repair, [923](#)  
Herniated Disc, [923](#)  
Herniated Disk, [923](#)  
Herniation, [923](#)  
Herniorrhaphy, [923](#)  
Heroin, [923](#)  
Heroin Toxicity, [923](#)  
Herpangina, [923](#)  
Herpes, [924](#)  
Herpes Encephalitis, [924](#)  
Herpes Gestationis, [924](#)  
Herpes Labialis, [924](#)  
Herpes Simplex, [924](#)  
Herpes Simplex Encephalitis, [924](#)  
Herpes Simplex Infection, [924](#)  
Herpes Simplex Keratitis, [924](#)  
Herpes Simplex Skin Infection, [924](#)  
Herpes Simplex Type 1, [924](#)  
Herpes Simplex Virus, [924](#)  
Herpes Simplex Virus Infection, [924](#)  
Herpes Vegetans, [925](#)  
Herpes Viral Culture Of Lesion, [925](#)  
Herpes Zoster, [925](#)  
Herpes Zoster Ophthalmicus, [925](#)  
Herpesviridae, [925](#)  
Herpesviridae Infection, [925](#)  
Herpetic Gingivostomatitis, [925](#)  
Herpetic Glossitis, [925](#)  
Herpetic Keratitis, [925](#)  
Herpetic Stomatitis, [925](#)  
Herpetic Whitlow, [925](#)  
Herpetiform, [925](#)  
Herring Bodies, [926](#)  
Hertel Exophthalmometer, [926](#)  
Hertz, [926](#)  
Herxheimer Reaction, [926](#)  
Hesitation Wounds, [926](#)  
Hesselbach Triangle, [926](#)  
Heterochromatin, [926](#)  
Heterochromia, [926](#)  
Heterocyclic Amine, [926](#)  
Heterogeneity, [926](#)  
Heterogeneous Echotexture, [926](#)  
Heterograft, [926](#)  
Heterologous, [926](#)  
Heterophil Antibody, [926](#)  
Heterophile Antibody, [927](#)  
Heterophyes Heterophyes, [927](#)  
Heteroplasia, [927](#)  
Heteroplasmy, [927](#)  
Heterosexual, [927](#)  
Heterotaxy Syndrome, [927](#)  
Heterotopic, [927](#)  
Heterotopic Breast Tissue Of The Vulva, [927](#)  
Heterotopic Pregnancy, [927](#)  
Heterotransplant, [927](#)  
Heterozygote, [927](#)  
Heterozygous, [927](#)  
Hexachlorobenzene, [927](#)  
Hexachlorophane, [927](#)  
Hexachlorophene, [927](#)  
Hexadactyly, [928](#)  
Hexadecadienoic Acid, [928](#)  
Hexadecatetraenoic Acid, [928](#)  
Hexamethonium, [928](#)  
Hexamine, [928](#)  
Hexane, [928](#)  
Hexanoic Acid, [928](#)  
Hexokinase, [928](#)  
Hexose, [928](#)  
Hexose Monophosphate Shunt, [928](#)  
Hexylresorcinol, [928](#)  
Heyde Syndrome, [928](#)  
Heyde Syndrome Triad, [928](#)  
HHT, [929](#)  
Hiatal, [929](#)  
Hiatal Hernia, [929](#)  
Hiatus, [929](#)  
Hiatus Hernia, [929](#)  
Hib, [929](#)  
Hibernoma, [929](#)  
Hiccough, [929](#)  
Hiccup, [929](#)  
Hiccups, [929](#)  
Hidradenitis, [929](#)  
Hidradenitis Suppurativa, [929](#)  
Hidradenocarcinoma, [929](#)  
Hidradenoma, [929](#)  
Hidradenoma Papilliferum, [929](#)  
Hidrocystoma, [930](#)  
Hidrosis, [930](#)  
High Altitude, [930](#)  
High Altitude Cerebral Edema, [930](#)  
High Altitude Illness, [930](#)  
High Arch, [930](#)  
High Blood Cholesterol Levels, [930](#)  
High Blood Potassium, [930](#)  
High Blood Pressure, [930](#)  
High Blood Pressure (Hypertension), [930](#)  
High Blood Pressure - Medicine-Related, [2122](#)  
High Blood Pressure And Eye Disease, [930](#)  
High Blood Protein, [930](#)  
High Blood Sugar, [930](#)

---

---

High Cholesterol, [930](#)  
 High HDL Cholesterol (Hyperalphalipoproteinemia), [931](#)  
 High Hemoglobin Count, [931](#)  
 High Output Failure, [931](#)  
 High Potassium, [931](#)  
 High Potassium Level, [931](#)  
 High Protein Diet, [931](#)  
 High Red Blood Cell Count, [931](#)  
 High Triglycerides, [931](#)  
 High Uric Acid Level, [931](#)  
 High White Blood Cell Count, [931](#)  
 High-Altitude Cerebral Edema, [931](#)  
 High-Altitude Pulmonary Edema, [931](#)  
 High-Density Lipoprotein, [931](#)  
 High-Density Lipoprotein Cholesterol, [931](#)  
 High-Dependency Unit, [932](#)  
 High-Fiber Foods, [932](#)  
 High-Flow Nasal Cannula, [932](#)  
 High-Fructose Corn Syrup, [932](#)  
 High-Grade Endometrial Stromal Sarcoma, [932](#)  
 High-Grade Serous Ovarian Carcinoma, [932](#)  
 High-Grade Surface Osteosarcoma, [932](#)  
 High-Intensity Focused Ultrasound, [932](#)  
 High-Level Disinfection, [932](#)  
 High-Output Heart Failure, [932](#)  
 High-Pitched Cry, [932](#)  
 High-Resolution Computed Tomography, [932](#)  
 High-Risk Pregnancy Factors, [933](#)  
 High-Sensitivity Cardiac Troponin, [933](#)  
 Highly Pathogenic Avian Influenza, [933](#)  
 Highly Selective Vagotomy, [933](#)  
 Hilar, [933](#)  
 Hilar Cholangiocarcinoma, [933](#)  
 Hilar Lymphadenopathy, [933](#)  
 Hill's Sign, [933](#)  
 Hill-Sachs Lesion, [933](#)  
 Hilum, [933](#)  
 Hilus, [933](#)  
 Hindbrain, [933](#)  
 Hindgut, [933](#)  
 Hinge Joint, [934](#)  
 Hip, [934](#)  
 Hip Arthroscopy, [934](#)  
 Hip Bursitis, [934](#)  
 Hip Circumference, [934](#)  
 Hip Dislocation, [934](#)  
 Hip Disorders, [934](#)  
 Hip Dysplasia, [934](#)  
 Hip Dysplasia, Congenital, [538](#)  
 Hip Fracture, [934](#)  
 Hip Fracture In Older Adults, [934](#)  
 Hip Fracture Surgery, [934](#)  
 Hip Impingement, [934](#)  
 Hip Injury, [934](#)  
 Hip Joint, [935](#)  
 Hip Joint Injection, [935](#)  
 Hip Joint Replacement, [935](#)  
 Hip Labral Tear, [935](#)  
 Hip Pain, [935](#)  
 Hip Prosthesis, [935](#)  
 Hip Replacement, [935](#)  
 Hip-Fracture Rehabilitation, [935](#)  
 Hippocampal Sclerosis, [935](#)  
 Hippocampus, [935](#)  
 Hippocrates, [935](#)  
 Hippocratic Oath, [935](#)  
 Hippotherapy, [935](#)  
 Hippius, [936](#)  
 Hirata Disease, [936](#)  
 Hirayama Disease, [936](#)  
 Hirschsprung Disease, [936](#)  
 Hirschsprung's Disease, [936](#)  
 Hirsutism, [936](#)  
 Hirudin, [936](#)  
 Hirudins, [936](#)  
 Hirudo Medicinalis, [936](#)  
 His, [936](#)  
 His Bundle Electrophysiology, [936](#)  
 Histamine, [937](#)  
 Histamine Degradation, [937](#)  
 Histamine Receptor, [937](#)  
 Histamine Release, [937](#)  
 Histamine-2 Receptor Antagonist, [2122](#)  
 Histidine, [937](#)  
 Histidine Decarboxylase, [937](#)  
 Histidinemia, [937](#)  
 Histiocyte, [937](#)  
 Histiocytic Sarcoma, [937](#)  
 Histiocytoma, [937](#)  
 Histiocytosis, [937](#)  
 Histiocytosis X, [937](#)  
 Histochemical, [937](#)  
 Histocompatibility, [937](#)  
 Histocompatibility Antigen Test, [937](#)  
 Histocompatibility Testing, [937](#)  
 Histogenesis, [938](#)  
 Histological Techniques, [938](#)  
 Histology, [938](#)  
 Histology Cassette, [938](#)  
 Histone, [938](#)  
 Histone Acetylation, [938](#)  
 Histone Acetyltransferase, [938](#)  
 Histone Deacetylases, [938](#)  
 Histone Deacetylation, [938](#)  
 Histone Modification, [938](#)  
 Histopathologic Grade, [938](#)  
 Histopathology, [938](#)  
 Histoplasma, [938](#)

---

---

Histoplasma Capsulatum, [938](#)  
 Histoplasma Complement Fixation, [938](#)  
 Histoplasmosis, [938](#)  
 Histrionic Personality Disorder, [939](#)  
 Hitchhiker's Thumb, [939](#)  
 HIV, [939](#)  
 Hiv Antibodies, [939](#)  
 HIV in Pregnancy, [939](#)  
 HIV Infection, [939](#)  
 Hiv Seropositivity, [939](#)  
 HIV Test, [939](#)  
 HIV Transmission And Prevention, [939](#)  
 HIV Viral Load, [939](#)  
 HIV-Associated Rheumatic Disease, [939](#)  
 HIV-Related Kidney Disease, [939](#)  
 HIV/AIDS, [940](#)  
 HIV/AIDS Immunopathogenesis, [940](#)  
 HIV/AIDS In Pregnant Women And Infants, [940](#)  
 HIV/AIDS Nutrition, [940](#)  
 Hives, [940](#)  
 Hives And Angioedema, [940](#)  
 HLA, [940](#)  
 Hla Antigen, [940](#)  
 HLA System, [940](#)  
 HLA-B27, [940](#)  
 HLA-B27 Antigen, [940](#)  
 HLHS, [940](#)  
 HMG-CoA Reductase, [940](#)  
 Hmgb Proteins, [941](#)  
 Hmgn Proteins, [941](#)  
 Hoarding, [941](#)  
 Hoarding Disorder, [941](#)  
 Hoarseness, [941](#)  
 Hodgkin Lymphoma, [941](#)  
 Hodgkin Lymphoma In Children, [941](#)  
 Hodgkin's And Non-Hodgkin's Lymphoma Comparison, [941](#)  
 Hodgkin's Disease, [941](#)  
 Hodgkin's Lymphoma, [941](#)  
 Hoffmann Sign, [941](#)  
 Hoffmann Syndrome, [941](#)  
 Hofmann's Syndrome, [941](#)  
 Holiday Heart Syndrome, [941](#)  
 Holistic Medicine, [942](#)  
 Hollenhorst Plaque, [942](#)  
 Holmium, [942](#)  
 Holmium Laser Enucleation Of The Prostate, [942](#)  
 Holoprosencephaly, [942](#)  
 Holorachischisis, [942](#)  
 Holosystolic Murmur Differential, [942](#)  
 Holt-Oram Syndrome, [942](#)  
 Holter Electrocardiography, [942](#)  
 Holter Monitor, [942](#)  
 Holter Monitoring, [942](#)  
 Homeobox Gene, [942](#)  
 Homeodomain Proteins, [942](#)  
 Homeopathy, [943](#)  
 Homeostasis, [943](#)  
 Homeotherm, [943](#)  
 Homicidal Ideation, [943](#)  
 Homicide, [943](#)  
 Homo Sapiens, [943](#)  
 Homocysteine, [943](#)  
 Homocystinuria, [943](#)  
 Homograft, [943](#)  
 Homologous, [943](#)  
 Homologous Recombination, [943](#)  
 Homologous Transplantation, [943](#)  
 Homonymous Hemianopia, [943](#)  
 Homosexual, [943](#)  
 Homosexuality, [944](#)  
 Homovanillic Acid, [944](#)  
 Homozygote, [944](#)  
 Homozygous, [944](#)  
 Homozygous Alpha-1-Antitrypsin Deficiency, [944](#)  
 Homozygous Familial Hypercholesterolemia, [944](#)  
 Honey-Bee Stings, [944](#)  
 Honeycomb Lung, [944](#)  
 Honeycomb Lung Pathology, [944](#)  
 Honolulu Fever, [944](#)  
 Hookworm, [944](#)  
 Hookworm Infection, [944](#)  
 Hookworm Infestation, [944](#)  
 Hordeolum, [944](#)  
 Horizontal Gene Transfer, [945](#)  
 Horizontal Pursuit, [945](#)  
 Hormonal Drugs, [945](#)  
 Hormonal Effects In Newborns, [945](#)  
 Hormone, [945](#)  
 Hormone Inhibitor, [2122](#)  
 Hormone Levels, [945](#)  
 Hormone Receptor, [945](#)  
 Hormone Replacement Therapy, [945](#)  
 Hormone Therapy, [945](#)  
 Hormone Therapy For Breast Cancer, [945](#)  
 Hormone Therapy for Endometriosis, [945](#)  
 Hormone Therapy For Prostate Cancer, [945](#)  
 Hormone-Receptor Positive, [945](#)  
 Hormone-Sensitive Lipase, [945](#)  
 Hormones, [946](#)  
 Horn, [946](#)  
 Horner Syndrome, [946](#)  
 Horner's Syndrome, [946](#)  
 Hornet Stings, [946](#)  
 Horripilation, [946](#)  
 Horseshoe Kidney, [946](#)  
 Hospice, [946](#)  
 Hospice Care, [946](#)

---

---

Hospice Rehabilitation, 946  
Hospital Code, 946  
Hospital Latex Allergy Precautions, 946  
Hospital- and Ventilator-Associated Pneumonia, 946  
Hospital-Acquired Infection, 947  
Hospital-Acquired Infections, 947  
Hospital-Acquired Pneumonia, 947  
Hospital-Associated Disability, 947  
Host, 947  
Host Cell, 947  
Host Response, 947  
Hostility, 947  
Hot Flash, 947  
Hot Flashes, 947  
Hot Flushes, 947  
Hot Tub Folliculitis, 947  
Hounsfield Unit, 947  
Hounsfield Units, 947  
Hour, 947  
Hour-Glass Contraction, 948  
Housemaid's Knee, 948  
Howell-Jolly Bodies, 948  
Howship-Romberg Sign, 948  
HPA (Hypothalamic-Pituitary-Adrenal Axis), 948  
HPLC, 948  
HPV (Human Papillomavirus), 948  
HPV Infection, 948  
HPV Test, 948  
HPV Vaccine, 2122  
HPV/Genital Warts, 948  
HPylori, 948  
HRCT (High-Resolution Computed Tomography), 948  
HRT (Hormone Replacement Therapy), 948  
Hs-cTn, 949  
Ht (Hematocrit), 949  
Huber Needle, 949  
Hughes-Stovin Syndrome, 949  
Human Bites, 949  
Human Calcitonin, 949  
Human Chorionic Gonadotropin, 949  
Human Chromosome Count, 949  
Human Cyclic Neutropenia, 949  
Human Cytomegalovirus Infection, 949  
Human Embryo, 949  
Human Gastrin, 949  
Human Genome, 949  
Human Herpesvirus 6, 949  
Human Herpesvirus 8, 949  
Human Immunodeficiency Virus, 950  
Human Immunodeficiency Virus (HIV), 950  
Human Leukocyte Antigen, 950  
Human Metapneumovirus, 950  
Human Microbiome, 950  
Human Nociceptin, 950  
Human Papillomavirus, 950  
Human Papillomavirus (HPV/Genital Warts), 950  
Human Papillomavirus Infection, 950  
Human Placental Lactogen, 950  
Humectant, 950  
Humeral Epicondyle, 950  
Humeral Fracture, 950  
Humeroradial Joint, 950  
Humeroulnar Joint, 951  
Humerus, 951  
Humerus Fracture, 951  
Humidifier, 951  
Humidifier Fever, 951  
Humidity, 951  
Humor, 951  
Humoral Immunity, 951  
Hunchback, 951  
Hunger, 951  
Hungry Bone Syndrome, 951  
Hunter Syndrome, 951  
Hunter's Syndrome, 951  
Huntington Disease, 951  
Huntington's Chorea (HD), 952  
Huntington's Disease (HD), 952  
Hurler Disease, 952  
Hurler Syndrome, 952  
Hurthle Cell, 952  
Hurthle Cell Cancer, 952  
Hurthle Cell Carcinoma (Oncocytic Carcinoma), 952  
Hutchinson Sign, 952  
Hutchinson Teeth, 952  
Hutchinson Triad, 952  
Hutchinson's Teeth, 952  
Hutchinson-Gilford Progeria Syndrome, 952  
Hutchinson-Gilford Progeria Syndrome (HGPS), 953  
Hutchinson-Gilford Syndrome (HGPS), 953  
Hyaline, 953  
Hyaline Cartilage, 953  
Hyaline Casts, 953  
Hyaline Membrane, 953  
Hyalitis, 953  
Hyaloid Canal, 953  
Hyaluronic Acid, 953  
Hyaluronidase, 953  
Hyaluronoglucosaminidase, 953  
Hybridoma, 953  
Hycanthone, 953  
Hydatid Cyst, 953  
Hydatid Disease, 954  
Hydatidiform Mole, 954  
Hydralazine, 954  
Hydranmios, 954  
Hydranencephaly, 954  
Hydrarthrosis, 954

---

---

Hydration, 954  
 Hydrazine, 954  
 Hydrazine Sulfate, 954  
 Hydrazines, 954  
 Hydrazones, 954  
 Hydroa Vacciniforme, 954  
 Hydrocarbon, 954  
 Hydrocarbon Pneumonia, 954  
 Hydrocarbon Toxicity, 955  
 Hydrocele, 955  
 Hydrocele Repair, 955  
 Hydrocephalus, 955  
 Hydrocephalus Acquired, 955  
 Hydrocephalus Ex-Vacuo, 955  
 Hydrocephalus, Congenital, 446  
 Hydrocephaly, 955  
 Hydrochloric Acid, 955  
 Hydrochloric Acid Poisoning, 955  
 Hydrochlorothiazide, 955  
 Hydrocodone, 955  
 Hydrocodone And Acetaminophen Overdose, 955  
 Hydrocodone Bitartrate, 956  
 Hydrocodone/Oxycodone Overdose, 956  
 Hydrocolpos, 956  
 Hydrocortisone, 956  
 Hydrocyanic Acid Poisoning, 956  
 Hydroflumethiazide, 956  
 Hydrofluoric Acid Exposure, 956  
 Hydrofluoric Acid Poisoning, 956  
 Hydrogel, 956  
 Hydrogen Bond, 956  
 Hydrogen Breath Test, 956  
 Hydrogen Cyanide, 956  
 Hydrogen Peroxide, 956  
 Hydrogen Peroxide Poisoning, 956  
 Hydrogen Sulfide, 957  
 Hydrogen-Ion Concentration, 957  
 Hydrogenation, 957  
 Hydromorphone, 957  
 Hydromorphone Overdose, 957  
 Hydromyelia, 957  
 Hydronephrosis, 957  
 Hydronephrosis In Newborns, 957  
 Hydronephrosis Of One Kidney, 957  
 Hydrophobia, 957  
 Hydropneumothorax, 957  
 Hydrops, 957  
 Hydrops Fetalis, 957  
 Hydrops Of The Gallbladder, 958  
 Hydrosalpinx, 958  
 Hydrotherapy, 958  
 Hydrothorax, 958  
 Hydroureter, 958  
 Hydroxamic Acid, 958  
 Hydroxocobalamin, 958  
 Hydroxyapatite, 958  
 Hydroxychloroquine, 958  
 Hydroxycholesterols, 958  
 Hydroxylase Deficiency (21-Hydroxylase), 958  
 Hydroxylation, 958  
 Hydroxyproline, 958  
 Hydroxyprolinemia, 958  
 Hydroxysteroid, 959  
 Hydroxyurea, 959  
 Hydroxyzine, 959  
 Hydroxyzine Overdose, 959  
 Hydrozoa, 959  
 Hygiene, 959  
 Hygiene Hypothesis, 959  
 Hygroma, 959  
 Hygroma Colli, 959  
 Hymen, 959  
 Hymenectomy, 959  
 Hymenolepiasis, 959  
 Hymenolepis Diminuta, 959  
 Hymenolepis Nana, 959  
 Hymenolepis Tapeworm Infection, 960  
 Hymenoptera, 960  
 Hyoid Bone, 960  
 Hyoscine, 960  
 Hyoscyamine, 960  
 Hyper IgM Syndrome, 960  
 Hyper-IgE Syndrome, 960  
 Hyper-IgM Syndrome, 960  
 Hyper-IgM Syndromes, 960  
 Hyperactivity, 960  
 Hyperactivity, ADHD, 145  
 Hyperacusis, 960  
 Hyperadrenocorticism, 960  
 Hyperaemia, 960  
 Hyperaesthesia, 960  
 Hyperaldosteronism, 961  
 Hyperaldosteronism Complications, 961  
 Hyperalgesia, 961  
 Hyperammonemia, 961  
 Hyperammonemia due to Ornithine Transcarbamylase Deficiency, 961  
 Hyperamylasemia, 961  
 Hyperargininemia, 961  
 Hyperbaric, 961  
 Hyperbaric Chamber, 961  
 Hyperbaric Oxygen Therapy, 961  
 Hyperbilirubinaemia, 961  
 Hyperbilirubinemia, 961  
 Hypercalcaemia, 961  
 Hypercalcemia, 961  
 Hypercalcemia Of Malignancy, 962  
 Hypercalcemia-Supravalvar Aortic Stenosis, 962

---

---

Hypercalcemic Small Cell Ovarian Carcinoma, 962  
Hypercalciuria, 962  
Hypercapnia, 962  
Hypercementosis, 962  
Hyperchloremic Acidosis, 962  
Hypercholesterolaemia, 962  
Hypercholesterolemia, 962  
Hypercoagulable State, 962  
Hypercyanotic Tetralogy, 962  
Hyperelastic Skin, 962  
Hyperemesis, 962  
Hyperemesis Gravidarum, 963  
Hyperemia, 963  
Hyperemia Grade, 963  
Hyperesthesia, 963  
Hyperextension, 963  
Hyperfibrinogenemia, 963  
Hyperflexion, 963  
Hypergammaglobulinemia, 963  
Hyperglycaemia, 963  
Hyperglycemia, 963  
Hyperglycinemia, 963  
Hypergonadotropic Hypogonadism, 963  
Hypergraphia, 963  
Hyperhidrosis, 963  
Hyperhomocysteinemia, 963  
Hypericum Perforatum, 964  
Hyperimmunization, 964  
Hyperimmunoglobulin D Syndrome, 964  
Hyperimmunoglobulin E Syndrome, 964  
Hyperinsulinemia, 964  
Hyperinsulinism, 964  
Hyperkalaemia, 964  
Hyperkalemia, 964  
Hyperkalemic Periodic Paralysis, 964  
Hyperkeratosis, 964  
Hyperkinesia, 964  
Hyperlipasaemia, 964  
Hyperlipidaemia, 964  
Hyperlipidemia, 965  
Hyperlipoproteinemia, 965  
Hyperlordosis, 965  
Hyperlordotic Posture, 965  
Hyperlysinemia, 965  
Hypermagnesemia, 965  
Hypermetropia, 965  
Hypermobile Ehlers-Danlos Syndrome, 965  
Hypermobile Joints, 965  
Hypermobility, 965  
Hypermobility Spectrum Disorder, 965  
Hypermobility Syndrome, 965  
Hypermotility, 965  
Hypernatraemia, 965  
Hypernatremia, 966  
Hypernatremia Result, 966  
Hyperopia, 966  
Hyperornithinemia, 966  
Hyperosmolar, 966  
Hyperosmolar Hyperglycemic State, 966  
Hyperosmolar Non-Ketotic Coma, 966  
Hyperostosis, 966  
Hyperoxaluria, 966  
Hyperoxia, 966  
Hyperparakeratosis, 966  
Hyperparathyroid Bone Disease, 966  
Hyperparathyroidism, 966  
Hyperpathia, 966  
Hyperphagia, 967  
Hyperphosphatemia, 967  
Hyperpigmentation, 967  
Hyperpituitarism, 967  
Hyperplasia, 967  
Hyperplastic Polyp, 967  
Hyperpnea, 967  
Hyperpnoea, 967  
Hyperpolarization, 967  
Hyperprolactinaemia, 967  
Hyperprolactinemia, 967  
Hyperproteinemia, 967  
Hyperpyrexia, 967  
Hyperreactio Luteinalis, 967  
Hyperreflexia, 967  
Hypersensitivity, 968  
Hypersensitivity Pneumonitis, 968  
Hypersensitivity Reactions, 968  
Hypersensitivity Type I, 968  
Hypersensitivity Type II, 968  
Hypersensitivity Type III, 968  
Hypersensitivity Type IV, 968  
Hypersensitivity Vasculitis, 968  
Hypersomnia, 968  
Hypersplenism, 968  
Hypertelorism, 968  
Hypertension, 968  
Hypertension and Pregnancy, 968  
Hypertension and the Kidney, 968  
Hypertension In Anesthesia, 968  
Hypertension With Chronic Kidney Disease, 969  
Hypertensive, 969  
Hypertensive Cerebellar Hemorrhage, 969  
Hypertensive Crisis, 969  
Hypertensive Emergency, 969  
Hypertensive Encephalopathy, 969  
Hypertensive Heart Disease, 969  
Hypertensive Intracerebral Hemorrhage, 969  
Hypertensive Kidney Disease, 969  
Hypertensive Nephropathy, 969  
Hypertensive Nephrosclerosis, 969

---



---

Hypertensive Retinopathy, 969  
 Hypertensive Urgency, 969  
 Hyperthermia, 969  
 Hyperthermia For Treating Cancer, 970  
 Hyperthermia In Anesthesia, 970  
 Hyperthermic Intraperitoneal Chemotherapy, 970  
 Hyperthyroidism, 970  
 Hyperthyroxinemia, 970  
 Hypertonia, 970  
 Hypertonic Saline, 970  
 Hypertonic Saline Inhalation, 970  
 Hypertonic Solution, 970  
 Hypertrichosis, 970  
 Hypertriglyceridemia, 970  
 Hypertrophic, 970  
 Hypertrophic Cardiomyopathy, 970  
 Hypertrophic Cardiomyopathy Genetics, 971  
 Hypertrophic Cardiomyopathy Treatment, 971  
 Hypertrophic Gastritis, 971  
 Hypertrophic Myocardiopathy, 971  
 Hypertrophic Obstructive Cardiomyopathy, 971  
 Hypertrophic Osteoarthropathy, 971  
 Hypertrophic Pyloric Stenosis, 971  
 Hypertrophic Scar, 971  
 Hypertrophy, 971  
 Hypertropia, 971  
 Hyperuricemia, 971  
 Hyperventilating, 971  
 Hyperventilation, 971  
 Hyperventilation Syndrome, 972  
 Hypervigilance, 972  
 Hyperviscosity Syndrome, 972  
 Hypervitaminosis A, 972  
 Hypervitaminosis D, 972  
 Hypervolaemia, 972  
 Hypervolemia, 972  
 Hypha, 972  
 Hyphema, 972  
 Hypnagogic, 972  
 Hypnagogic Hallucinations, 972  
 Hypnic Headache, 972  
 Hypnogram, 972  
 Hypnosis, 973  
 Hypnotherapy, 973  
 Hypnotic, 973  
 Hypoactive Sexual Desire Disorder, 973  
 Hypoadrenalism, 973  
 Hypoalbuminemia, 973  
 Hypoalphalipoproteinemias, 973  
 Hypoblast, 973  
 Hypocalcaemia, 973  
 Hypocalcemia, 973  
 Hypocapnia, 973  
 Hypochloremic Alkalosis, 973  
 Hypochlorite, 973  
 Hypcholesterolemia, 974  
 Hypochondria, 974  
 Hypochondroplasia, 974  
 Hypochromia, 974  
 Hypochromic Anaemia, 974  
 Hypochromic Anemia, 974  
 Hypocitraturia, 974  
 Hypocupremia, 974  
 Hypodermic Needle, 974  
 Hypodermis, 974  
 Hypodontia, 974  
 Hypoesthesia, 974  
 Hypogalactia, 974  
 Hypogammaglobulinemia, 974  
 Hypogastric Region, 975  
 Hypoglossal Canal, 975  
 Hypoglossal Nerve, 975  
 Hypoglossal Nerve Injury, 975  
 Hypoglossal Nerve Palsy, 975  
 Hypoglycaemia, 975  
 Hypoglycemia, 975  
 Hypoglycemia In Diabetes, 975  
 Hypoglycemia Unawareness, 975  
 Hypoglycemic, 975  
 Hypoglycemic Agent, 2122  
 Hypogonadism, 975  
 Hypogonadism In Men, 975  
 Hypogonadism In Women, 976  
 Hypogonadotropic Hypogonadism, 976  
 Hypohidrosis, 976  
 Hypokalaemia, 976  
 Hypokalemia, 976  
 Hypokalemic Periodic Paralysis, 976  
 Hypokinesia, 976  
 Hypolipidemic Agent, 2123  
 Hypolipidemic Agents, 976  
 Hypolipoproteinemia, 976  
 Hypomagnesaemia, 976  
 Hypomagnesemia, 976  
 Hypomania, 976  
 Hypomaniac Episode, 976  
 Hypomelanosis Of Ito, 976  
 Hypomethylation, 977  
 Hypomnesia, 977  
 Hyponatraemia, 977  
 Hyponatremia, 977  
 Hypoparathyroidism, 977  
 Hypopharyngeal Neoplasm, 977  
 Hypopharynx, 977  
 Hypophosphataemia, 977  
 Hypophosphatasia (HPP), 977  
 Hypophosphatemia, 977  
 Hypophosphatemic Rickets, 977

---

---

Hypophysectomy, 977  
 Hypophysis, 977  
 Hypophysitis, 978  
 Hypopigmentation, 978  
 Hypopituitarism, 978  
 Hypoplasia, 978  
 Hypoplastic Left Heart Syndrome, 978  
 Hypoplastic Left Heart Syndrome (HLHS), 978  
 Hypoploidy, 978  
 Hypopnea, 978  
 Hypoproliferative Anemia, 978  
 Hypoproteinaemia, 978  
 Hypoproteinemia, 978  
 Hypopyon, 978  
 Hyporeninemic Hypoaldosteronism, 978  
 Hyposmia, 979  
 Hypospadias, 979  
 Hypospadias Repair, 979  
 Hypotension, 979  
 Hypotension In Anesthesia, 979  
 Hypotensive, 979  
 Hypothalamic Amenorrhea, 979  
 Hypothalamic Dysfunction, 979  
 Hypothalamic Hamartoma, 979  
 Hypothalamic Hormone, 979  
 Hypothalamic Neoplasm, 979  
 Hypothalamic Tumor, 979  
 Hypothalamic-Pituitary-Adrenal Axis, 979  
 Hypothalamic-Pituitary-Adrenal Axis Physiology, 980  
 Hypothalamic-Pituitary-Gonadal Axis Physiology, 980  
 Hypothalamic-Pituitary-Thyroid Axis Physiology, 980  
 Hypothalamus, 980  
 Hypothermia, 980  
 Hypothermia In Anesthesia, 980  
 Hypothyroidism, 980  
 Hypotonia, 980  
 Hypotonia-Hypomnesia-Hypogonadism-Obesity Syndrome (HHHO), 980  
 Hypotonic Solution, 980  
 Hypotony Maculopathy, 980  
 Hypotrichosis, 981  
 Hypotrophy, 981  
 Hypotropia, 981  
 Hypoventilation, 981  
 Hypoventilation Syndromes, 981  
 Hypovolaemia, 981  
 Hypovolemia, 981  
 Hypovolemic Shock, 981  
 Hypoxaemia, 981  
 Hypoxanthine, 981  
 Hypoxemia, 981  
 Hypoxia, 981  
 Hypoxia And Hypoxemia, 981  
 Hypoxia-Inducible Factor, 982  
 Hypoxic Cell Injury, 982  
 Hypoxic Ischemic Encephalopathy, 982  
 Hypoxic Pulmonary Vasoconstriction, 982  
 Hypoxic-Ischemic Brain Injury, 982  
 Hypsarrhythmia, 982  
 Hysterectomy, 982  
 Hysterectomy And Surgical Menopause, 982  
 Hysterosalpingo-Contrast Sonography, 982  
 Hysterosalpingo-Contrast-Sonography, 982  
 Hysterosalpingogram, 982  
 Hysterosalpingography, 983  
 Hysteroscope, 983  
 Hysteroscopic Metroplasty, 983  
 Hysteroscopic Myomectomy, 983  
 Hysteroscopy, 983  
 Hysterotomy, 983  
 Hürthle Cell Tumour, 983  
 I-Cell Disease, 986  
 Iatrogenic, 985  
 Iatrogenic Cushing Syndrome, 985  
 Iatrogenic Disease, 985  
 Iatrogenic Injury, 985  
 Iatrogenic Pneumothorax, 985  
 Ibandronic Acid, 985  
 IBD, 985  
 Ibritumomab Tiuxetan, 985  
 IBS, 985  
 IBS (Irritable Bowel Syndrome), 1065  
 IBS-D, 985  
 Ibuprofen, 986  
 Ibuprofen Overdose, 986  
 Ibuprofen Toxicity, 986  
 IC, 1065  
 ICD, 986  
 ICD Shock Therapy, 986  
 ICD-10, 986  
 ICD-11, 986  
 ICD-O Neoplasm Morphology Classification, 986  
 Ice Pack, 986  
 Ice Pick Headache, 986  
 Iceland Disease, 1065  
 Icelandic Type Amyloidosis (Type VI), 1066  
 ICF Framework, 986  
 Ichthyosis, 986  
 Ichthyosis Vulgaris, 987  
 Ictal State, 987  
 Icteric, 987  
 Icteric Index, 987  
 Icterus, 987  
 Icterus (Chronic Familial), 1066  
 Ictus, 987  
 ICU-Acquired Weakness, 987  
 Id Reaction (Autoeczematization), 988

---

---

Idarubicin, 987  
IDC, 987  
Idea Of Reference, 987  
Ideas Of Reference, 987  
Ideation, 987  
Identical Twins, 987  
Idiopathic, 987  
Idiopathic Amyloidosis, 1066  
Idiopathic Anaphylaxis, 988  
Idiopathic Aplastic Anemia, 1066  
Idiopathic Autoimmune Hemolytic Anemia, 1066  
Idiopathic Basal Ganglia Calcification, 1066  
Idiopathic Chronic Erosive Gastritis, 1066  
Idiopathic Dilated Cardiomyopathy, 1066  
Idiopathic Edema, 707  
Idiopathic Facial Paralysis, 988  
Idiopathic Guttate Hypomelanosis, 988  
Idiopathic Hypercalciuria, 988  
Idiopathic Hypersomnia, 988  
Idiopathic Interstitial Pneumonias, 988  
Idiopathic Intracranial Hypertension, 988  
Idiopathic Myelofibrosis, 988  
Idiopathic Pulmonary Arterial Hypertension, 988  
Idiopathic Pulmonary Fibrosis, 988  
Idiopathic Thrombocytopenic Purpura, 988  
Idiopathic Toe Walking, 988  
Idiosyncrasy, 988  
Idiosyncratic Reaction, 988  
Idioventricular Rhythm, 988  
Ifosfamide, 989  
iFR, 989  
Ig, 989  
IgA, 989  
IgA Nephropathy, 989  
IgA Pemphigus, 989  
IgA Vasculitis, 989  
IgD, 989  
IgE, 989  
IgG, 989  
IgG and Complement Deposition, 989  
IgG4-Related Disease, 989  
IgG4-Related Disease Arteritis, 989  
IgG4-Related Ophthalmic Disease, 989  
IgM, 990  
IGRT, 990  
IIH, 990  
Ileal Atresia, 990  
Ileal Conduit, 990  
Ileal Fistula, 990  
Ileal Hemorrhage, 990  
Ileal Neoplasm, 990  
Ileal Obstruction, 990  
Ileal Pouch-Anal Anastomosis, 990  
Ileal Stenosis, 990  
Ileal Ulcer, 990  
Ileitis, 990  
Ileocaecal, 990  
Ileocecal Junction, 990  
Ileocecal Valve, 991  
Ileostomy, 991  
Ileum, 991  
Ileus, 991  
Iliac, 991  
Iliac Artery, 991  
Iliac Crest, 991  
Iliac Crest Bone Graft, 991  
Iliac Vein, 991  
Iliopsoas Muscles, 991  
Iliopsoas Tendinitis, 991  
Iliotibial Band, 991  
Iliotibial Band Friction Syndrome, 991  
Iliotibial Band Syndrome, 991  
Ilium, 991  
Illinois Type Amyloidosis, 1066  
Illness Anxiety Disorder, 992  
Illusion, 992  
Iloprost, 992  
Image-Guided Ablation, 992  
Image-Guided Biopsy, 992  
Image-Guided Drainage, 992  
Image-Guided Therapy, 992  
Imaging, 992  
Imaging And Radiology, 992  
Imaging Technique, 992  
Imaging-Guided Biopsy, 992  
Imatinib Mesylate, 992  
Imerslund-Gräsbeck Syndrome, 992  
Imidazoles, 992  
Iminoacetic Acid Scan, 993  
Imipenem, 993  
Imipramine, 993  
Imipramine Overdose, 993  
Imiquimod, 993  
Immature Ovarian Teratoma, 993  
Immediate Hypersensitivity, 993  
Immersion Foot, 993  
Immobilization, 993  
Immobilize, 993  
Immotile Cilia Syndrome, 993  
Immune, 993  
Immune And Infectious Diseases, 993  
Immune Checkpoint, 993  
Immune Checkpoint Inhibitor, 994  
Immune Complex, 994  
Immune Disorder, 994  
Immune Evasion, 994  
Immune Hemolytic Anemia, 994  
Immune Privileged Sites, 994

---

---

Immune Reconstitution Inflammatory Syndrome, 994  
Immune Response, 994  
Immune Status, 994  
Immune System, 994  
Immune Thrombocytopenia, 995  
Immune Thrombocytopenia in Pregnancy, 995  
Immune Thrombocytopenic Purpura, 995  
Immune Tolerance, 995  
Immune-Mediated Necrotizing Myopathy, 994  
Immune-Related Adverse Event, 994  
Immunity, 995  
Immunization, 995  
Immunizations, 1066  
Immunizations For People With Diabetes, 995  
Immunoassay, 995  
Immunoassays, 995  
Immunoblast Count, 995  
Immunoblastic Lymphadenopathy, 1066  
Immunoblotting, 995  
Immunochemistry, 995  
Immunocompetence, 996  
Immunocompetent, 996  
Immunocompromised, 996  
Immunocompromised Host, 996  
Immunocompromised Pneumonia, 996  
Immunoconjugate, 996  
Immunocytochemistry, 996  
Immunodeficiency, 996  
Immunodeficiency And Infections, 996  
Immunodeficiency Diseases, 996  
Immunodeficiency Disorders, 996  
Immunodeficiency with Ataxia Telangiectasia, 1066  
Immuno-electrophoresis, 996  
Immuno-electrophoresis – Blood, 996  
Immunofixation Blood Test, 996  
Immunofluorescence, 996  
Immunofluorescence Technique, 997  
Immunogenetics, 997  
Immunogenicity, 997  
Immunoglobulin, 997  
Immunoglobulin A, 997  
Immunoglobulin A Deficiency, 997  
Immunoglobulin D, 997  
Immunoglobulin E, 997  
Immunoglobulin G, 997  
Immunoglobulin G Deficiency, 997  
Immunoglobulin G4-Related Disease, 997  
Immunoglobulin Isotype, 997  
Immunoglobulin M, 997  
Immunoglobulin M Deficiency, 998  
Immunoglobulin Replacement Therapy, 998  
Immuno-hemolysis, 998  
Immunohistochemistry, 998  
Immunologic Adjuvant, 998  
Immunologic Factor, 998  
Immunologic Memory, 998  
Immunologic Skin Test, 998  
Immunologic Surveillance, 998  
Immunologic Technique, 998  
Immunologic Test, 999  
Immunological Memory, 998  
Immunological Synapse, 998  
Immunological Tolerance, 998  
Immunologically Privileged Site, 998  
Immunology, 999  
Immunomodulation, 999  
Immunomodulators, 999  
Immunomodulators In Dermatology, 999  
Immunomodulatory, 999  
Immunophenotyping, 999  
Immunoprotein, 999  
Immunoradiometric Assay, 999  
Immunoregulation, 999  
Immunosenescence, 999  
Immunosuppressant, 2123  
Immunosuppressant Drug, 2123  
Immunosuppression, 999  
Immunosuppression In Transplant, 999  
Immunosuppression Screening, 1000  
Immunosuppressive, 1000  
Immunosuppressive Drug, 2123  
Immunotherapy, 1000  
Immunotherapy And Biologic Treatments, 1000  
Immunotherapy For Cancer, 1000  
Immunotherapy In Dermatology, 1000  
Immunotoxicity, 1000  
Immunotoxin, 1000  
Immunoturbidimetry, 1000  
IMP Dehydrogenase, 1001  
Impacted, 1000  
Impacted Fracture, 1000  
Impacted Tooth, 1000  
Impacted Wisdom Teeth, 1000  
Impacted Wisdom Teeth Cross-Reference, 1000  
Impaction, 1001  
Impaired Fasting Glucose, 1001  
Impaired Glucose Tolerance, 1001  
Impaired Smell, 1001  
Impaired Taste, 1001  
Impedance Hearing Testing, 1001  
Impedance Plethysmography, 1001  
Impella Device, 1001  
Imperforate Anus, 1001  
Imperforate Anus Repair, 1001  
Imperforate Hymen, 1001  
Imperforate Vagina, 1001  
Impetigo, 1001  
Impingement Syndrome, 1002

---

---

Impingement Syndrome of the Shoulder, [1002](#)  
 Implant, [1002](#)  
 Implant Failure, [1003](#)  
 Implant Sterilization, [1003](#)  
 Implant-Associated Infection, [1002](#)  
 Implant-Based Breast Reconstruction, [1003](#)  
 Implant-Supported Prosthesis, [1003](#)  
 Implantable Cardioverter Defibrillator, [1002](#)  
 Implantable Cardioverter-Defibrillator, [1002](#)  
 Implantable Defibrillator, [1002](#)  
 Implantable Glucose Sensor, [1002](#)  
 Implantable Loop Recorder, [1002](#)  
 Implantable Port, [1002](#)  
 Implantable Venous Access Port, [1002](#)  
 Implantation, [1002](#)  
 Implantation Procedure, [1002](#)  
 Implants, [1003](#)  
 Importance Weight, [1003](#)  
 Impotence, [1066](#)  
 Impotence (Erection Problems), [1066](#)  
 Impression, [1003](#)  
 Imprinting, [1003](#)  
 Impulse, [1003](#)  
 Impulse Control Disorder, [1003](#)  
 Impulse Oscillometry, [1003](#)  
 Impulsive Behavior, [1003](#)  
 Impulsivity, [1003](#)  
 In, [1003](#)  
 In Situ, [1025](#)  
 In Situ Carcinoma, [1026](#)  
 In Situ Hybridization, [1026](#)  
 In Utero, [1051](#)  
 In Vitro, [1052](#)  
 In Vitro Fertilisation, [1052](#)  
 In Vitro Fertilization, [1053](#)  
 In Vivo, [1053](#)  
 In-Vivo, [1053](#)  
 Inactivated Polio Vaccine, [2123](#)  
 Inactivated Vaccines, [1003](#)  
 Inactivator, [1004](#)  
 Inamrinone, [1004](#)  
 Inanition, [1004](#)  
 Inappropriate Antidiuretic Hormone Secretion, [1004](#)  
 Inappropriate Sinus Tachycardia, [1004](#)  
 Inborn Error Of Metabolism, [1004](#)  
 Inborn Errors Of Immunity, [1004](#)  
 Inborn Errors of Metabolism, [1004](#)  
 Inbreed, [1004](#)  
 Inbreeding, [1004](#)  
 Incarcerated Hernia, [1004](#)  
 Incentive Spirometry, [1004](#)  
 Incest, [1004](#)  
 Incidence, [1005](#)  
 Incidental Finding, [1005](#)  
 Incidentaloma, [1005](#)  
 Incised Wound, [1005](#)  
 Incision, [1005](#)  
 Incision and Drainage, [1005](#)  
 Incision Care After Surgery, [1066](#)  
 Incisional Biopsy, [1005](#)  
 Incisional Hernia, [1005](#)  
 Incisor, [1005](#)  
 Incisors, [1005](#)  
 Inclusion Body, [1005](#)  
 Inclusion Body Myositis, [1006](#)  
 Inclusion Conjunctivitis, [1006](#)  
 Incompetence, [1006](#)  
 Incompetent Cervix, [1006](#)  
 Incomplete Abortion, [1006](#)  
 Incomplete Heart Block, [1006](#)  
 Incomplete Miscarriage, [1006](#)  
 Incomplete Ossification, [1006](#)  
 Incontinence, [1006](#)  
 Incontinence in Men, [1066](#)  
 Incontinence, Fecal, [726](#)  
 Incontinence, Urinary, [2002](#)  
 Incontinence/OAB, [1066](#)  
 Incontinentia Pigmenti, [1006](#)  
 Incoordination, [1006](#)  
 Incoordinatio, [1006](#)  
 Increased Appetite, [1006](#)  
 Increased Head Circumference, [1006](#)  
 Increased Intracranial Pressure, [1007](#)  
 Increased Libido, [1007](#)  
 Increased Osteoid, [1007](#)  
 Increased Vascular Permeability, [1007](#)  
 Increment, [1007](#)  
 Incretin Effect, [1007](#)  
 Incretins, [1007](#)  
 Incubation, [1007](#)  
 Incubation Period, [1007](#)  
 Incubator, [1007](#)  
 Incus, [1007](#)  
 Indapamide, [1007](#)  
 Indel Mutation, [1007](#)  
 Independence, [1007](#)  
 Indeterminate, [1008](#)  
 India Immunization Schedule, [1008](#)  
 Indian Tick Typhus, [1008](#)  
 Indiana Type Amyloidosis (Type II), [1066](#)  
 Indican, [1008](#)  
 Indicate, [1008](#)  
 Indication, [1008](#)  
 Indigestion, [1008](#)  
 Indigo Carmine, [1008](#)  
 Indirect Antiglobulin Test, [1008](#)  
 Indirect Calorimetry, [1008](#)  
 Indirect Coombs Test, [1008](#)

---

---

Indirect Inguinal Hernia, [1008](#)  
Indirect Veneer, [1008](#)  
Indium, [1009](#)  
Indium-111, [1009](#)  
Indium-111 Scan, [1009](#)  
Indium-Labelled WBC Scan, [1009](#)  
Indocyanine Green, [1009](#)  
Indocyanine Green Clearance, [1009](#)  
Indole, [1009](#)  
Indole-3-Carbinol, [1009](#)  
Indolent, [1009](#)  
Indolent Systemic Mastocytosis, [1067](#)  
Indoles, [1009](#)  
Indometacin, [1009](#)  
Indomethacin, [1009](#)  
Indomethacin Overdose, [1009](#)  
Indoor Allergens, [1009](#)  
Induced Abortion, [1010](#)  
Induced Hyperthermia, [1010](#)  
Induced Labor, [1010](#)  
Induced Sputum Analysis, [1010](#)  
Inducible Laryngeal Obstruction, [1010](#)  
Inducing Labor, [1010](#)  
Induction, [1010](#)  
Induction Of Labor, [1010](#)  
Induction Therapy, [1010](#)  
Indurated, [1010](#)  
Induration, [1010](#)  
Industrial Bronchitis, [1010](#)  
Industrial Disease, [1010](#)  
Industrial Health, [1010](#)  
Indwelling Catheter, [1010](#)  
Indwelling Catheter Care, [1011](#)  
Indwelling Pleural Catheter, [1011](#)  
Inert Gas Narcosis, [1011](#)  
Inertia, [1011](#)  
Infant, [1011](#)  
Infant Botulism, [1011](#)  
Infant Cardiopulmonary Resuscitation, [1011](#)  
Infant Feeding, [1011](#)  
Infant Feeding Patterns and Diet, [1011](#)  
Infant Formula, [1011](#)  
Infant Formulas, [1011](#)  
Infant Gynecomastia, [1012](#)  
Infant Harlequin Syndrome, [1012](#)  
Infant Hearing Loss, [1012](#)  
Infant Incubators, [1013](#)  
Infant Jaundice, [1013](#)  
Infant Leukemia, [1013](#)  
Infant Mortality, [1013](#)  
Infant Nutrition, [1013](#)  
Infant Of A Substance-Using Mother, [1013](#)  
Infant Of Diabetic Mother, [1013](#)  
Infant Reflex, [1013](#)  
Infant Reflexes, [1013](#)  
Infant Reflux, [1013](#)  
Infant Respiratory Distress Syndrome, [1013](#)  
Infant Sleep Position, [1013](#)  
Infant Test Procedure Preparation, [1013](#)  
Infanticide, [1012](#)  
Infantile Bradycardia, [1012](#)  
Infantile Colic, [1012](#)  
Infantile Cortical Hyperostosis (Caffey Disease), [1012](#)  
Infantile Fibrosarcoma, [1012](#)  
Infantile Gaucher Disease, [1067](#)  
Infantile Hemangioma, [1012](#)  
Infantile Hepatic Hemangiomas, [1012](#)  
Infantile Myoclonic Encephalopathy, [1067](#)  
Infantile Paralysis, [1012](#)  
Infantile Sleep Apnea, [1067](#)  
Infantile Spasm, [1012](#)  
Infantile Spasms, [1012](#)  
Infantile Spinal Muscular Atrophy, [1067](#)  
Infantile Spinal Muscular Atrophy Type I, [1067](#)  
Infarct, [1013](#)  
Infarction, [1013](#)  
Infection, [1014](#)  
Infection Control, [1014](#)  
Infection Imaging, [1014](#)  
Infection Prevention, [1014](#)  
Infections in Diabetes, [1014](#)  
Infections In Pregnancy, [1014](#)  
Infectious Arteritis, [1014](#)  
Infectious Arthritis, [1014](#)  
Infectious Chorea, [1067](#)  
Infectious Conjunctivitis, [1014](#)  
Infectious Disease, [1014](#)  
Infectious Diseases, [1014](#)  
Infectious Dose, [1014](#)  
Infectious Encephalitis, [1014](#)  
Infectious Encephalomyelitis, [1014](#)  
Infectious Enterocolitis, [1015](#)  
Infectious Esophagitis, [1015](#)  
Infectious Hepatitis, [1015](#)  
Infectious Mediastinitis, [1015](#)  
Infectious Meningitis, [1015](#)  
Infectious Mononucleosis, [1015](#)  
Infectious Myringitis, [1015](#)  
Infective Cervicitis, [1015](#)  
Infective Cystitis, [1015](#)  
Infective Endocarditis, [1015](#)  
Infective Endocarditis Duke Criteria, [1015](#)  
Infective Endocarditis Pathology, [1015](#)  
Infective Phlebitis, [1015](#)  
Infective Vaginitis, [1016](#)  
Inferior, [1016](#)  
Inferior Alveolar Nerve Block, [1016](#)  
Inferior Colliculus, [1016](#)

---



---

Inferior Hemifield Loss, [1016](#)  
 Inferior Mediastinum, [1016](#)  
 Inferior Mesenteric Artery, [1016](#)  
 Inferior Myocardial Infarction, [1016](#)  
 Inferior Petrosal Sinus Sampling, [1016](#)  
 Inferior Vena Cava, [1016](#)  
 Inferior Vena Cava Filter, [1016](#)  
 Inferior Vena Cava Syndrome, [1016](#)  
 Inferior Wall Myocardial Infarction, [1017](#)  
 Inferolateral, [1017](#)  
 Infertility, [1017](#)  
 Infertility, Female, [728](#)  
 Infertility, Male, [1192](#)  
 Infestation, [1017](#)  
 Infiltrate, [1017](#)  
 Infiltrating Cancer, [1017](#)  
 Infiltrating Ductal Carcinoma, [1017](#)  
 Infiltration, [1017](#)  
 Inflamed Gallbladder, [1017](#)  
 Inflamed Pancreas, [1017](#)  
 Inflamed Pericardium, [1017](#)  
 Inflammaging, [1017](#)  
 Inflammasomes, [1017](#)  
 Inflammation, [1017](#)  
 Inflammation Imaging, [1017](#)  
 Inflammatory Back Pain, [1018](#)  
 Inflammatory Bowel Disease, [1018](#)  
 Inflammatory Bowel Disease Diet, [1018](#)  
 Inflammatory Bowel Disease/Crohn's Disease, [1067](#)  
 Inflammatory Bowel Disease: Intestinal Problems, [1018](#)  
 Inflammatory Breast Cancer, [1018](#)  
 Inflammatory Myopathies, [1018](#)  
 Inflammatory Myopathy, [1018](#)  
 Inflammatory Response, [1018](#)  
 Infliximab, [1018](#)  
 Influenza, [1018](#)  
 Influenza A Virus, [1018](#)  
 Influenza Test, [1018](#)  
 Influenza Vaccination In Pregnancy, [1019](#)  
 Influenza Vaccine, [2123](#)  
 Influenza Virus, [1019](#)  
 Influenzavirus A, [1019](#)  
 Influenzavirus B, [1019](#)  
 Informatics, [1019](#)  
 Informed Consent, [1019](#)  
 Infrared Rays, [1019](#)  
 Infrared Spectrometry, [1019](#)  
 Infrared Therapy, [1019](#)  
 Infrarenal Aorta, [1019](#)  
 Infrapinatus Muscle, [1019](#)  
 Infratentorial Neoplasm, [1019](#)  
 Infrequent, [1019](#)  
 Infundibular Stenosis, [1019](#)  
 Infundibulopelvic Ligament, [1020](#)  
 Infundibulum, [1020](#)  
 Infuse, [1020](#)  
 Infusion, [1020](#)  
 Infusion Pump, [1020](#)  
 Ingested Foreign Body, [1020](#)  
 Ingestion, [1020](#)  
 Ingredient, [1020](#)  
 Ingrown Hair, [1020](#)  
 Ingrown Nail, [1020](#)  
 Ingrown Toenail (Onychocryptosis), [1020](#)  
 Ingrown Toenails, [1020](#)  
 Inguinal, [1021](#)  
 Inguinal Canal, [1021](#)  
 Inguinal Hernia, [1021](#)  
 Inguinal Hernia Repair, [1021](#)  
 Inguinal Orchiectomy, [1021](#)  
 Inguinal Region, [1021](#)  
 Inhalant, [1021](#)  
 Inhalant Substance Use, [1021](#)  
 Inhalation, [1021](#)  
 Inhalation Injury, [1021](#)  
 Inhalational Anesthesia, [1021](#)  
 Inhaled Corticosteroids, [1021](#)  
 Inhaled Foreign Body, [1021](#)  
 Inhaled Nitric Oxide, [1021](#)  
 Inhaler, [1022](#)  
 Inherited Disease, [1022](#)  
 Inherited Lipodystrophy, [1067](#)  
 Inhibin, [1022](#)  
 Inhibition, [1022](#)  
 Inhibitory Neurochemical, [1022](#)  
 Inhibitory Synapse, [1022](#)  
 Iniencephaly, [1022](#)  
 Initial Insomnia, [1022](#)  
 Injectable Fillers, [1022](#)  
 Injection, [1022](#)  
 Injection Drug Use, [1022](#)  
 Injury, [1022](#)  
 Injury Dating, [1022](#)  
 Injury Site, [1022](#)  
 Ink Poisoning, [1023](#)  
 Ink Remover Poisoning, [1023](#)  
 Innate Immune Evasion, [1023](#)  
 Innate Immunity, [1023](#)  
 Innate Lymphoid Cells, [1023](#)  
 Inner Ear, [1023](#)  
 Inner Ear Infection, [1023](#)  
 Innervation, [1023](#)  
 Innominate, [1023](#)  
 INOCA, [1023](#)  
 Inoculation, [1023](#)  
 Inoperable, [1023](#)  
 Inorganic, [1023](#)  
 Inorganic Chemical, [1023](#)

---

---

Inosculation, [1024](#)  
Inosine, [1024](#)  
Inosine Monophosphate, [1024](#)  
Inosine Pranobex, [1024](#)  
Inositol, [1024](#)  
Inositol 1,4,5-Trisphosphate, [1024](#)  
Inositol Phosphates, [1024](#)  
Inositol Trisphosphate, [1024](#)  
Inotrope, [2123](#)  
Inotropes In Anesthesia, [1024](#)  
Inotropic, [1024](#)  
Inotropic Agent, [1024](#)  
Inpatient, [1024](#)  
Inpatient Rehabilitation, [1024](#)  
INR, [1024](#)  
Insect Bite, [1025](#)  
Insect Bites, [1025](#)  
Insect Bites And Stings, [1025](#)  
Insect Sting Allergies, [1025](#)  
Insect Virus, [1025](#)  
Insecta, [1024](#)  
Insecticide, [1025](#)  
Insecticide Poisoning, [1025](#)  
Insemination, [1025](#)  
Insensible Fluid Loss, [1025](#)  
Insertion, [1025](#)  
Insidious, [1025](#)  
Insinuate, [1025](#)  
Insomnia, [1026](#)  
Insomnia Disorder, [1026](#)  
Inspection, [1026](#)  
Inspiratory Capacity, [1026](#)  
Inspiratory Crackles, [1026](#)  
Inspiratory Reserve Volume, [1026](#)  
Inspiratory Time, [1026](#)  
Inspiratory-to-Expiratory Ratio, [1026](#)  
Instantaneous Wave-Free Ratio, [1026](#)  
Instillation, [1026](#)  
Instrument, [1026](#)  
Instrument Sterilization, [1027](#)  
Instrumental Activities Of Daily Living, [1027](#)  
Instrumental ADL, [1027](#)  
Insufficient Cervix, [1027](#)  
Insula, [1027](#)  
Insular Cortex, [1027](#)  
Insulin, [1027](#)  
Insulin Antagonist, [1027](#)  
Insulin Aspart, [1027](#)  
Insulin Autoantibodies, [1027](#)  
Insulin C-Peptide Test, [1027](#)  
Insulin Coma, [1027](#)  
Insulin Detemir, [1028](#)  
Insulin Glargine, [1028](#)  
Insulin Injection Technique, [1028](#)  
Insulin Isophane, [1028](#)  
Insulin Lipohypertrophy, [1028](#)  
Insulin Lispro, [1028](#)  
Insulin Pen Devices, [1028](#)  
Insulin Preparations, [1028](#)  
Insulin Pump Therapy, [1029](#)  
Insulin Pumps, [1029](#)  
Insulin Receptor, [1029](#)  
Insulin Resistance, [1029](#)  
Insulin Resistance Syndrome, [1029](#)  
Insulin Resistance Syndromes, [1029](#)  
Insulin Secretion, [1029](#)  
Insulin Signaling, [1029](#)  
Insulin Storage and Syringe Safety, [1029](#)  
Insulin Therapy in Diabetes, [1029](#)  
Insulin Tolerance Test, [1029](#)  
Insulin-Dependent Diabetes, [1027](#)  
Insulin-Induced Lipohypertrophy, [1028](#)  
Insulin-Like Growth Factor, [1028](#)  
Insulin-Like Growth Factor 1, [1028](#)  
Insulinoma, [1028](#)  
Integrase, [1029](#)  
Integrative Medicine, [1030](#)  
Integrative Rehabilitation, [1030](#)  
Integrin, [1030](#)  
Integrin Binding, [1030](#)  
Integron, [1030](#)  
Integrins, [1030](#)  
Integumentary System, [1030](#)  
Intein, [1030](#)  
Intellectual Disability, [1030](#)  
Intelligence, [1030](#)  
Intelligence Test, [1030](#)  
Intense Pulsed Light, [1030](#)  
Intensity, [1030](#)  
Intensity-Modulated Radiation Therapy, [1030](#)  
Intensive Care, [1030](#)  
Intensive Care Medicine, [2123](#)  
Intensive Care Rehabilitation, [1031](#)  
Intensive Care Unit, [1031](#)  
Intention Tremor, [1031](#)  
Intentional Obstetrical Clavicle Fracture, [1031](#)  
Interatrial Septum, [1031](#)  
Intercalated Disc, [1031](#)  
Intercalating Agent, [1031](#)  
Intercellular, [1031](#)  
Intercostal, [1031](#)  
Intercostal Muscles, [1031](#)  
Intercostal Retraction, [1031](#)  
Intercostal Retractions, [1031](#)  
Intercostal Space, [1031](#)  
Intercurrent Disease, [1031](#)  
Interdental Brush, [1032](#)  
Interdigital Neuroma, [1032](#)

---

---

Interference, [1032](#)  
 Interferometry, [1032](#)  
 Interferon, [1032](#)  
 Interferon Gamma Release Assay, [1032](#)  
 Interferon Inducer, [1032](#)  
 Interferon Signaling, [1032](#)  
 Interferon-Alpha, [1032](#)  
 Interferon-Beta, [1032](#)  
 Interferon-Gamma Release Assay, [1032](#)  
 Interferonopathies, [1032](#)  
 Interferons, [1032](#)  
 Interleukin, [1032](#)  
 Interleukin 1, [1032](#)  
 Interleukin 13, [1033](#)  
 Interleukin 2, [1033](#)  
 Interleukin 7, [1033](#)  
 Interleukin-6, [1033](#)  
 Interleukins, [1033](#)  
 INTERMACS Profile, [1033](#)  
 Intermediate, [1033](#)  
 Intermediate Cuneiform, [1033](#)  
 Intermediate Filament, [1033](#)  
 Intermediate Uveitis, [1033](#)  
 Intermediate-Density Lipoprotein, [1033](#)  
 Intermittent Auscultation, [1033](#)  
 Intermittent Claudication, [1033](#)  
 Intermittent Explosive Disorder, [1033](#)  
 Intermittent Fasting, [1034](#)  
 Intermittent Fever, [1034](#)  
 Intermittent Mandatory Ventilation, [1034](#)  
 Intern, [1034](#)  
 Internal, [1034](#)  
 Internal Bleeding, [1034](#)  
 Internal Capsule, [1034](#)  
 Internal Capsule Stroke, [1034](#)  
 Internal Carotid Artery, [1034](#)  
 Internal Carotid Artery Dissection, [1034](#)  
 Internal Environment, [1034](#)  
 Internal Fixation, [1034](#)  
 Internal Fixator, [1035](#)  
 Internal Hernia, [1035](#)  
 Internal Iliac Artery, [1035](#)  
 Internal Iliac Vein, [1035](#)  
 Internal Jugular Vein, [1035](#)  
 Internal Jugular Vein Cannulation, [1035](#)  
 Internal Jugular Vein Thrombosis, [1035](#)  
 Internal Medicine, [2123](#)  
 Internal Oblique Muscle, [1035](#)  
 Internal Podalic Version, [1035](#)  
 Internal Resorption, [1035](#)  
 International Classification of Diseases, [1035](#)  
 International Normalized Ratio, [1035](#)  
 International Unit, [1035](#)  
 Interneuron, [1036](#)  
 Interneurons, [1036](#)  
 Internist, [1036](#)  
 Internuclear Ophthalmoplegia, [1036](#)  
 Interosseous, [1036](#)  
 Interosseous Nerve Syndrome, [1036](#)  
 Interpersonal Psychotherapy, [1036](#)  
 Interphalangeal Dislocation, [1036](#)  
 Interphalangeal Joint, [1036](#)  
 Interphase, [1036](#)  
 Interphase Cell, [1036](#)  
 Interproximal Caries, [1036](#)  
 Interrupted Aortic Arch, [1036](#)  
 Intersexual, [1036](#)  
 Interstitial, [1036](#)  
 Interstitial (Nonidiopathic) Pulmonary Fibrosis, [1037](#)  
 Interstitial Cystitis, [1036](#)  
 Interstitial Fluid, [1037](#)  
 Interstitial Granulomatous Dermatitis, [1037](#)  
 Interstitial Keratitis, [1037](#)  
 Interstitial Lung Disease, [1037](#)  
 Interstitial Nephritis, [1037](#)  
 Interstitial Pneumonia, [1037](#)  
 Interstitial Pulmonary Fibrosis, [1037](#)  
 Interstitium, [1037](#)  
 Intertrigo, [1037](#)  
 Intertrochanteric Fracture, [1038](#)  
 Interval, [1038](#)  
 Intervening Sequence, [1038](#)  
 Intervention, [1038](#)  
 Interventional, [1038](#)  
 Interventional Cardiology, [1038](#)  
 Interventional Procedure, [1038](#)  
 Interventional Procedures, [1038](#)  
 Interventional Radiologist, [1038](#)  
 Interventional Radiology, [1038](#)  
 Interventricular Septum, [1038](#)  
 Intervertebral Disc, [1038](#)  
 Intervertebral Disk, [1038](#)  
 Intervertebral Foramen, [1039](#)  
 Intestinal Absorption, [1039](#)  
 Intestinal Angina, [1039](#)  
 Intestinal Atresia, [1039](#)  
 Intestinal Cancer, [1039](#)  
 Intestinal Carcinoid Tumor, [1039](#)  
 Intestinal Crypts, [1039](#)  
 Intestinal Diseases, [1039](#)  
 Intestinal Duplication, [1039](#)  
 Intestinal Failure, [1039](#)  
 Intestinal Fistula, [1039](#)  
 Intestinal Flukes, [1039](#)  
 Intestinal Ganglioneuromatosis, [1039](#)  
 Intestinal Gas, [1040](#)  
 Intestinal Ischemia, [1040](#)  
 Intestinal Leiomyoma, [1040](#)

---

---

Intestinal Lipodystrophy, [1040](#)  
Intestinal Lipomatosis, [1040](#)  
Intestinal Lymphangiectasia, [1040](#)  
Intestinal Lymphoma, [1040](#)  
Intestinal Malrotation, [1040](#)  
Intestinal Metaplasia, [1040](#)  
Intestinal Microbiome, [1040](#)  
Intestinal Motility, [1040](#)  
Intestinal Mucosa, [1040](#)  
Intestinal Obstruction, [1040](#)  
Intestinal Obstruction And Ileus, [1040](#)  
Intestinal Obstruction Repair, [1040](#)  
Intestinal Parasite, [1041](#)  
Intestinal Perforation, [1041](#)  
Intestinal Polyp, [1041](#)  
Intestinal Polyposis, [1041](#)  
Intestinal Pseudo-Obstruction, [1041](#)  
Intestinal Resection, [1041](#)  
Intestinal Villi, [1041](#)  
Intestinal Wall, [1041](#)  
Intestine, [1041](#)  
Intestine Disorder, [1041](#)  
Intestine Neoplasm, [1041](#)  
Intestine Obstruction, [1041](#)  
Intimal Dissection, [1041](#)  
Intimal Hyperplasia, [1041](#)  
Intimate Partner Violence In Pregnancy, [1042](#)  
Intolerance, [1042](#)  
Intoxication, [1042](#)  
Intra-Abdominal Abscess, [1042](#)  
Intra-Abdominal Hypertension, [1042](#)  
Intra-Aortic Balloon Pump, [1042](#)  
Intra-Arterial, [1042](#)  
Intra-Articular Injection, [1042](#)  
Intraabdominal Cancer, [1042](#)  
Intraabdominal Hemorrhage, [1042](#)  
Intraabdominal Hernia, [1042](#)  
Intraabdominal Infection, [1042](#)  
Intraarticular, [1042](#)  
Intrabronchial, [1042](#)  
Intracardiac Echocardiography, [1043](#)  
Intracardiac Electrophysiology Study, [1043](#)  
Intracartilaginous, [1043](#)  
Intracavernosal Injection, [1043](#)  
Intracavitary, [1043](#)  
Intracavity Radiotherapy, [1043](#)  
Intracellular, [1043](#)  
Intracellular Fluid, [1043](#)  
Intracellular Membrane, [1043](#)  
Intracellular Pathogens, [1043](#)  
Intracellular Transport, [1043](#)  
Intracerebral, [1043](#)  
Intracerebral Hemorrhage, [1043](#)  
Intracisternal, [1043](#)  
Intracoronary, [1044](#)  
Intracranial, [1044](#)  
Intracranial Aneurysm, [1044](#)  
Intracranial Hemorrhage, [1044](#)  
Intracranial Pressure, [1044](#)  
Intracranial Pressure Monitoring, [1044](#)  
Intracranial Venous Malformations, [1044](#)  
Intractable, [1044](#)  
Intracytoplasmic Sperm Injection, [1044](#)  
Intradermal, [1044](#)  
Intradermal Nevus, [1044](#)  
Intradermal Skin Test, [1044](#)  
Intraductal Carcinoma, [1044](#)  
Intraductal Papillary Mucinous Neoplasm, [1044](#)  
Intraductal Papilloma, [1045](#)  
Intraduodenal, [1045](#)  
Intraepidermal, [1045](#)  
Intraepithelial, [1045](#)  
Intraesophageal, [1045](#)  
Intrahepatic, [1045](#)  
Intrahepatic Cholestasis, [1045](#)  
Intrahepatic Cholestasis Of Pregnancy, [1045](#)  
Intralesional, [1045](#)  
Intralymphatic, [1045](#)  
Intramedullary, [1045](#)  
Intramedullary Nail, [1045](#)  
Intramedullary Spinal Cord Abscesses, [1045](#)  
Intramural Hematoma, [1045](#)  
Intramural Plaque, [1046](#)  
Intramuscular, [1046](#)  
Intramuscular Injection, [1046](#)  
Intramuscular Injection Technique, [1046](#)  
Intramuscular Myxoma, [1046](#)  
Intraocular, [1046](#)  
Intraocular Infection, [1046](#)  
Intraocular Lens, [1046](#)  
Intraocular Lens Dislocation, [1046](#)  
Intraocular Lens Implant, [1046](#)  
Intraocular Lymphoma, [1046](#)  
Intraocular Pressure, [1046](#)  
Intraocular Tumor, [1046](#)  
Intraoperative, [1046](#)  
Intraoperative Blood Transfusion, [1046](#)  
Intraoperative Neuromonitoring, [1047](#)  
Intraoral, [1047](#)  
Intraosseous Access, [1047](#)  
Intraosseous Infusion, [1047](#)  
Intrapartum, [1047](#)  
Intrapartum Fever, [1047](#)  
Intrapartum Stillbirth, [1047](#)  
Intrapericardial, [1047](#)  
Intraperitoneal, [1047](#)  
Intraperitonealroute, [1047](#)

---

---

Intraprostatic, [1047](#)  
Intrapulmonary, [1047](#)  
Intrapulmonary Shunt, [1047](#)  
Intrasynovial, [1047](#)  
Intratendinous, [1047](#)  
Intratesticular, [1048](#)  
Intrathecal, [1048](#)  
Intrathecal Chemotherapy, [1048](#)  
Intrathoracic, [1048](#)  
Intrathoracic Localized Obstruction, [1048](#)  
Intratracheal, [1048](#)  
Intratympanic, [1048](#)  
Intrauterine, [1048](#)  
Intrauterine Contraceptive Device, [1048](#)  
Intrauterine Device, [1048](#)  
Intrauterine Devices, [1048](#)  
Intrauterine Fetal Demise / Stillbirth, [1048](#)  
Intrauterine Growth Restriction, [1048](#)  
Intrauterine Insemination, [1048](#)  
Intrauterine Pressure Catheter, [1049](#)  
Intrauterine System, [1049](#)  
Intrauterine Transfusion, [1049](#)  
Intravascular, [1049](#)  
Intravascular Catheter Infections, [1049](#)  
Intravascular Filter, [1049](#)  
Intravascular Hemolysis, [1049](#)  
Intravascular Lymphoma, [1049](#)  
Intravascular Ultrasound, [1049](#)  
Intravenous, [1049](#)  
Intravenous Anesthesia, [1049](#)  
Intravenous Catheters, [1049](#)  
Intravenous Drip, [1049](#)  
Intravenous Immunoglobulin, [1049](#)  
Intravenous Leiomyomatosis, [1050](#)  
Intravenous Lipid Emulsion Therapy, [1050](#)  
Intravenous Pyelogram, [1050](#)  
Intravenous Pyelography, [1050](#)  
Intravenous Sedation, [1050](#)  
Intravenous Thrombolysis, [1050](#)  
Intraventricular Hemorrhage, [1050](#)  
Intraventricular Hemorrhage Of The Newborn, [1050](#)  
Intravesical Therapy, [1050](#)  
Intravitreal Injection, [1050](#)  
Intrinsic, [1050](#)  
Intrinsic Factor, [1050](#)  
Intrinsic Pathway of Coagulation, [1050](#)  
Intrinsic Renal AKI, [1050](#)  
Intrinsic Sphincter Deficiency, [1051](#)  
Intrinsic Sympathomimetic Activity, [1051](#)  
Introitus, [1051](#)  
Intron, [1051](#)  
Intrusive Thought, [1051](#)  
Intubate, [1051](#)  
Intubation, [1051](#)  
Intussusception, [1051](#)  
Inulin, [1051](#)  
Invasive, [1051](#)  
Invasive Candidiasis, [1051](#)  
Invasive Ductal Carcinoma, [1051](#)  
Invasive Lobular Carcinoma, [1051](#)  
Invasive Mechanical Ventilation, [1052](#)  
Invasive Mole, [1052](#)  
Invasive Procedure, [1052](#)  
Inverse Agonist, [1052](#)  
Inverse Psoriasis, [1052](#)  
Inversion, [1052](#)  
Invert, [1052](#)  
Invert Sugar, [1052](#)  
Inverted Nipple, [1052](#)  
Inverted Papilloma, [1052](#)  
Investigational, [1052](#)  
Inveterate, [1052](#)  
Involucrum, [1053](#)  
Involuntary, [1053](#)  
Involuntary Movement, [1053](#)  
Involuntary Movements, [1053](#)  
Involuntary Muscle, [1053](#)  
Involuntary Psychiatric Treatment, [1053](#)  
Involute, [1053](#)  
Involution, [1053](#)  
Iodamoeba Butschlii, [1053](#)  
Iodide, [1053](#)  
Iodide-Associated Sialadenitis, [1053](#)  
Iodides, [1054](#)  
Iodinated Contrast Media, [1054](#)  
Iodine, [1054](#)  
Iodine Deficiency, [1054](#)  
Iodine Perfusion, [1054](#)  
Iodine Poisoning, [1054](#)  
Iodine-123, [1054](#)  
Iodine-131, [1054](#)  
Iododerma, [1054](#)  
Iodoform, [1054](#)  
Iohexol, [1054](#)  
Ion, [1054](#)  
Ion Channel, [1054](#)  
Ion Transport, [1055](#)  
Ionization, [1055](#)  
Ionized Calcium Test, [1055](#)  
Ionizing Radiation, [1055](#)  
Ionophore, [1055](#)  
Ions, [1055](#)  
Iontophoresis, [1055](#)  
Iopamidol, [1055](#)  
Iothalamate Clearance, [1055](#)  
Ioxaglic Acid, [1055](#)  
IPC, [1055](#)  
Ipecac, [1055](#)

---

---

Ipilimumab, [1055](#)  
IPMN, [1055](#)  
Ipratropium, [1056](#)  
Ipratropium Bromide, [1056](#)  
Iproniazid, [1056](#)  
Ipsilateral, [1056](#)  
IQ Testing, [1056](#)  
Irbesartan, [1056](#)  
Iridium, [1056](#)  
Iridocyclitis, [1056](#)  
Iridodialysis, [1056](#)  
Iridodonesis, [1056](#)  
Iridoschisis, [1056](#)  
Iridotomy, [1056](#)  
Irinotecan, [1056](#)  
Irinotecan Hydrochloride, [1056](#)  
Iris, [1056](#)  
Iris Atrophy, [1057](#)  
Iris Cyst, [1057](#)  
Iris Disease, [1057](#)  
Iris Leiomyoma, [1057](#)  
Iris Melanoma, [1057](#)  
Iris Neoplasm, [1057](#)  
Iris Prolapse, [1057](#)  
Iritis, [1057](#)  
Iron, [1057](#)  
Iron Deficiency, [1057](#)  
Iron Deficiency Anemia, [1057](#)  
Iron Overdose, [1058](#)  
Iron Overload, [1058](#)  
Iron Overload Anemia, [1067](#)  
Iron Toxicity, [1058](#)  
Iron-Deficiency Anaemia, [1057](#)  
Iron-Deficiency Anaemia In Adolescents, [1057](#)  
Iron-Deficiency Anemia, [1058](#)  
Iron-Deficiency Anemia in Children, [1058](#)  
Iron-Deficiency Anemia in Young Children, [1058](#)  
Irradiated Foods, [1058](#)  
Irradiation, [1058](#)  
Irradiation Fibrosis, [1058](#)  
Irregular, [1058](#)  
Irregular Menstrual Periods, [1067](#)  
Irregular Menstruation, [1058](#)  
Irregular Respiration, [1058](#)  
Irregular Sleep-Wake Syndrome, [1058](#)  
Irreversible Cell Injury, [1058](#)  
Irrigate, [1059](#)  
Irrigation, [1059](#)  
Irrigation Solution, [1059](#)  
Irritability, [1059](#)  
Irritable Bowel Syndrome, [1059](#)  
Irritable Bowel Syndrome (IBS), [1067](#)  
Irritable Bowel Syndrome Diet, [1059](#)  
Irritable Mood, [1059](#)  
Irritant Contact Dermatitis, [1059](#)  
Irritants, [1059](#)  
Irritation, [1059](#)  
Isaacs Syndrome, [1059](#)  
Isavuconazole, [1059](#)  
Isazophos, [1059](#)  
Ischaemia, [1060](#)  
Ischaemic Heart Disease, [1060](#)  
Ischemia, [1060](#)  
Ischemia-Reperfusion Injury, [1060](#)  
Ischemic Bone Necrosis, [1060](#)  
Ischemic Cardiomyopathy, [1060](#)  
Ischemic Cascade, [1060](#)  
Ischemic Chest Pain, [1060](#)  
Ischemic Cholangiopathy, [1060](#)  
Ischemic Colitis, [1060](#)  
Ischemic Fasciitis, [1060](#)  
Ischemic Gangrene, [1060](#)  
Ischemic Heart Disease, [1060](#)  
Ischemic Hepatitis, [1060](#)  
Ischemic Necrosis of Bone, [1067](#)  
Ischemic Optic Neuropathy, [1061](#)  
Ischemic Preconditioning, [1061](#)  
Ischemic Priapism, [1061](#)  
Ischemic Stroke, [1061](#)  
Ischiorectal Abscess, [1061](#)  
Ischium, [1061](#)  
Ishak Score, [1061](#)  
Islet Cell, [1061](#)  
Islet Cell Cancer, [1061](#)  
Islets Of Langerhans, [1061](#)  
Iso-Sulfan Blue, [1063](#)  
Isoaminile, [1061](#)  
Isoantibodies, [1061](#)  
Isoantigen, [1061](#)  
Isocarboxazid, [1061](#)  
Isochromosomes, [1061](#)  
Isocitrate Dehydrogenase, [1061](#)  
Isocyanate, [1062](#)  
Isoelectric Point, [1062](#)  
Isoenzyme, [1062](#)  
Isofenphos, [1062](#)  
Isoflurane, [1062](#)  
Isogeneic Graft, [1062](#)  
Isohemagglutinins, [1062](#)  
Isokinetic Exercise, [1062](#)  
Isolate, [1062](#)  
Isolated Systolic Hypertension, [1062](#)  
Isolation, [1062](#)  
Isolation Precautions, [1062](#)  
Isoleucine, [1062](#)  
Isoleucine 33 Amyloidosis, [1067](#)  
Isomerase, [1062](#)  
Isometric, [1062](#)

---



---

Isometropia, [1063](#)  
 Isoniazid, [1063](#)  
 Isoniazid Toxicity, [1063](#)  
 Isoprenaline, [1063](#)  
 Isoprocarb, [1063](#)  
 Isopropanol Alcohol Poisoning, [1063](#)  
 Isopropyl Alcohol, [1063](#)  
 Isoproterenol, [1063](#)  
 Isosorbide, [1063](#)  
 Isosorbide Dinitrate, [1063](#)  
 Isospora, [1063](#)  
 Isospora Belli, [1063](#)  
 Isosporiasis, [1063](#)  
 Isotonic, [1063](#)  
 Isotonic Exercise, [1063](#)  
 Isotonic Solution, [1064](#)  
 Isotope, [1064](#)  
 Isotretinoin, [1064](#)  
 Isovaleric Acidemia, [1064](#)  
 Isovolumetric Contraction, [1064](#)  
 Isozyme, [1064](#)  
 Ispaghula Husk, [1064](#)  
 Isradipine, [1064](#)  
 ISUP Grade Group, [1064](#)  
 Itch, [1064](#)  
 Itching, [1064](#)  
 Itchy Ear, [1064](#)  
 Itchy Skin, [1065](#)  
 Iterative Reconstruction, [1065](#)  
 ITP, [1065](#)  
 Itraconazole, [1065](#)  
 IUCD, [1065](#)  
 IV Fluids, [1065](#)  
 IV Treatment At Home, [1065](#)  
 Ivabradine, [1065](#)  
 IVC Filter, [1065](#)  
 Ivermectin, [1065](#)  
 IVF, [1065](#)  
 Ixabepilone, [1065](#)  
 Ixodes, [1065](#)  
  
 J Receptor, [1076](#)  
 J Wave, [1080](#)  
 Jaccoud Arthropathy, [1069](#)  
 Jaccoud's Arthropathy, [1069](#)  
 Jack-In-The-Pulpit Poisoning, [1069](#)  
 Jackknife Position, [1069](#)  
 Jacksonian Epilepsy, [1069](#)  
 Jacksonian March, [1069](#)  
 Jacksonian Seizure, [1069](#)  
 Jacobsen Syndrome, [1069](#)  
 Jadassohn's Naevus, [1070](#)  
 Jadassohn-Lewandowsky Syndrome, [1069](#)  
 Jaffe-Lichtenstein Syndrome, [1070](#)  
  
 JAK-STAT Pathway, [1070](#)  
 JAK2, [1070](#)  
 Jakob's Disease, [1079](#)  
 Jakob-Creutzfeldt Disease, [1079](#)  
 Jammed Finger, [1070](#)  
 Janeway Lesions, [1070](#)  
 Jansen Disease, [1079](#)  
 Jansen Metaphyseal Chondrodysplasia, [1079](#)  
 Janus Kinase, [1070](#)  
 Janus Kinase Inhibitors, [1070](#)  
 Japanese B Encephalitis, [1080](#)  
 Japanese Encephalitis, [1070](#)  
 Japanese Encephalitis Vaccine, [2123](#)  
 Jarcho-Levin Syndrome, [1070](#)  
 Jargon Aphasia, [1070](#)  
 Jarisch-Herxheimer Reaction, [1070](#)  
 Jatene Procedure, [1070](#)  
 Jaundice, [1071](#)  
 Jaundice And Breastfeeding, [1071](#)  
 Jaundice Causes, [1071](#)  
 Jaundice In Adults, [1071](#)  
 Jaundice In Neonates, [1071](#)  
 Jaundice In Newborns, [1071](#)  
 Jaundice in Newborns (Hyperbilirubinemia), [1080](#)  
 Jaw, [1071](#)  
 Jaw Fracture, [1071](#)  
 Jaw Jerk Reflex, [1071](#)  
 Jaw Neoplasm, [1071](#)  
 Jaw Pain, [1071](#)  
 Jaw Thrust Maneuver, [1071](#)  
 Jaw Tumors And Cysts, [1071](#)  
 Jaw-Winking Syndrome, [1072](#)  
 Jehovah's Witness, [1072](#)  
 Jejunal Atresia, [1072](#)  
 Jejunal Diverticulum, [1072](#)  
 Jejunal Fistula, [1072](#)  
 Jejunal Hemorrhage, [1072](#)  
 Jejunal Neoplasm, [1072](#)  
 Jejunal Stenosis, [1072](#)  
 Jejunal Ulcer, [1072](#)  
 Jejunitis, [1072](#)  
 Jejunoileal Atresia, [1072](#)  
 Jejunojejunostomy, [1072](#)  
 Jejunostomy, [1073](#)  
 Jejunostomy Feeding Tube, [1073](#)  
 Jejunum, [1073](#)  
 Jellyfish Sting, [1073](#)  
 Jellyfish Stings, [1073](#)  
 Jendrassik Maneuver, [1073](#)  
 Jendrassik's Maneuver, [1073](#)  
 Jerk Nystagmus, [1073](#)  
 Jerusalem Cherry Poisoning, [1073](#)  
 Jervell and Lange-Nielsen Syndrome, [1073](#)  
 Jet Lag, [1073](#)

---

---

Jet Lag Prevention, [1073](#)  
Jeune Syndrome, [1073](#)  
Jimsonweed Poisoning, [1073](#)  
Job Syndrome, [1074](#)  
Jock Itch, [1074](#)  
Jod-Basedow Phenomenon, [1074](#)  
Joffroy Sign, [1074](#)  
Johanson-Blizzard Syndrome, [1074](#)  
Joint, [1074](#)  
Joint Aspiration, [1074](#)  
Joint Capsule, [1074](#)  
Joint Contracture, [1074](#)  
Joint Crepitus, [1074](#)  
Joint Diseases, [1074](#)  
Joint Effusion, [1074](#)  
Joint Fluid Culture, [1075](#)  
Joint Fluid Gram Stain, [1075](#)  
Joint Inflammation, [1075](#)  
Joint Injection, [1075](#)  
Joint Mouse, [1075](#)  
Joint Pain, [1075](#)  
Joint Prosthesis, [1075](#)  
Joint Replacement, [1075](#)  
Joint Replacement Rehabilitation, [1075](#)  
Joint Stiffness, [1075](#)  
Joint Swelling, [1075](#)  
Joint X-Ray, [1075](#)  
Jones Criteria, [1075](#)  
Jones Fracture, [1076](#)  
Jones Metatarsal Fracture, [1076](#)  
Jones Tube, [1076](#)  
Jonston's Alopecia, [1080](#)  
Joubert Syndrome, [1076](#)  
Joule-Thomson Effect, [1076](#)  
JRA Juvenile Rheumatoid Arthritis, [1080](#)  
Jugular, [1076](#)  
Jugular Bulb, [1076](#)  
Jugular Foramen, [1076](#)  
Jugular Foramen Syndrome, [1076](#)  
Jugular Vein, [1076](#)  
Jugular Venous Distension, [1077](#)  
Jugular Venous Distention, [1077](#)  
Jugular Venous Pressure, [1077](#)  
Jugular Venous Waveform, [1077](#)  
Jugulodigastric Node, [1077](#)  
Jumper's Knee, [1077](#)  
Jumping Frenchmen of Maine, [1080](#)  
Junctional Bradycardia, [1077](#)  
Junctional Ectopic Tachycardia, [1077](#)  
Junctional Epidermolysis Bullosa, [1077](#)  
Junctional Escape Rhythm, [1077](#)  
Junctional Nevus, [1078](#)  
Junctional Premature Complex, [1078](#)  
Junctional Rhythm, [1078](#)  
Junctional Tachycardia, [1078](#)  
Jungian Analysis, [1078](#)  
Juvenile Absence Epilepsy, [1078](#)  
Juvenile Angiofibroma, [1078](#)  
Juvenile Arthritis, [1078](#)  
Juvenile Astrocytoma, [1078](#)  
Juvenile Dermatomyositis, [1078](#)  
Juvenile Diabetes, [1080](#)  
Juvenile Granulosa Cell Tumor, [1078](#)  
Juvenile Hypermobility Spectrum Disorder, [1079](#)  
Juvenile Hypermobility Syndrome, [1079](#)  
Juvenile Idiopathic Arthritis, [1079](#)  
Juvenile Laryngeal Papillomatosis, [1079](#)  
Juvenile Myoclonic Epilepsy, [1079](#)  
Juvenile Nasopharyngeal Angiofibroma, [1079](#)  
Juvenile Osteoporosis, [1079](#)  
Juvenile Polyposis Syndrome, [1079](#)  
Juvenile Retinoschisis, [1080](#)  
Juvenile Rheumatoid Arthritis, [1080](#)  
Juvenile Schizophrenia, [1079](#)  
Juvenile Xanthogranuloma, [1079](#)  
Juxta-Articular, [1080](#)  
Juxtaglomerular Apparatus, [1080](#)  
Juxtaglomerular Cell Tumor, [1080](#)  
Juxtaglomerular Cells, [1080](#)  
Juxtavesical, [1080](#)  
K-Complex, [1083](#)  
Kabuki Syndrome, [1081](#)  
Kahler Disease, [1081](#)  
Kahler's Disease, [1081](#)  
Kala-Azar, [1081](#)  
Kallikrein-Kinin System, [1081](#)  
Kallmann Syndrome, [1081](#)  
Kallmann Syndrome and IHH, [1081](#)  
Kanamycin, [1081](#)  
Kangaroo Care, [1081](#)  
Kaolin, [1081](#)  
Kaposi Sarcoma, [1082](#)  
Kaposi's Sarcoma, [1082](#)  
Kaposiform Hemangioendothelioma, [1082](#)  
Karnofsky Performance Status, [1082](#)  
Kartagener Syndrome, [1082](#)  
Kartagener Triad, [1082](#)  
Karyomegaly, [1082](#)  
Karyotype, [1082](#)  
Karyotype Result, [1082](#)  
Karyotyping, [1082](#)  
Kasabach-Merritt Phenomenon, [1082](#)  
Kasabach-Merritt Syndrome, [1082](#)  
Kasai Procedure, [1083](#)  
Kashin-Beck Disease, [1083](#)  
Katayama Fever, [1083](#)  
Katz Index, [1083](#)

---

---

Kava, [1102](#)  
Kawasaki Disease, [1083](#)  
Kawasaki Disease Diagnostic Criteria, [1083](#)  
Kayser-Fleischer Rings, [1083](#)  
KC, [1102](#)  
Kearns-Sayre Syndrome, [1083](#)  
Keel Breast, [1083](#)  
Kegel, [1083](#)  
Kegel Exercise, [1083](#)  
Kegel Exercises, [1102](#)  
Kehr Sign, [1083](#)  
Keloid, [1084](#)  
Keloid and Hypertrophic Scar, [1084](#)  
Keloid Scar, [1084](#)  
Keratan Sulfate, [1084](#)  
Keratan Sulphate, [1084](#)  
Keratectomy, [1084](#)  
Keratic Precipitates, [1084](#)  
Keratin, [1084](#)  
Keratin Pearl, [1084](#)  
Keratinocyte, [1084](#)  
Keratinocytes, [1084](#)  
Keratitis, [1084](#)  
Keratoacanthoma, [1084](#)  
Keratocele, [1085](#)  
Keratoconjunctivitis, [1085](#)  
Keratoconjunctivitis Sicca, [1085](#)  
Keratoconus, [1085](#)  
Keratocyte Count, [1085](#)  
Keratoderma, [1085](#)  
Keratoderma Blennorrhagica, [1085](#)  
Keratoderma Blennorrhagicum, [1085](#)  
Keratoderma Blenorrhagicum, [1085](#)  
Keratoderma Blennorrhagicum, [1085](#)  
Keratolysis, [1085](#)  
Keratolytics, [1085](#)  
Keratoma, [1085](#)  
Keratomalacia, [1085](#)  
Keratometer, [1085](#)  
Keratometry, [1086](#)  
Keratoplasty, [1086](#)  
Keratoprosthesis, [1086](#)  
Keratosis, [1086](#)  
Keratosis Follicularis, [1086](#)  
Keratosis Follicularis (Darier Disease), [1086](#)  
Keratosis Obturans, [1086](#)  
Keratosis Pilaris, [1086](#)  
Keratosis Seborrhic, [1102](#)  
Kerion, [1086](#)  
Kerley A Lines, [1086](#)  
Kerley B Lines, [1086](#)  
Kernicterus, [1086](#)  
Kernig Sign, [1086](#)  
Kernig's Sign, [1086](#)  
Kernohan Notch Phenomenon, [1087](#)  
Kerosene Poisoning, [1087](#)  
Keshan Disease, [1087](#)  
Ketamine, [1087](#)  
Ketamine Hydrochloride, [1087](#)  
Ketamine Toxicity, [1087](#)  
Keto Acid, [1087](#)  
Ketoacidosis, [1087](#)  
Ketoconazole, [1087](#)  
Ketogenesis, [1087](#)  
Ketogenic Diet, [1087](#)  
Ketone, [1087](#)  
Ketone Bodies, [1087](#)  
Ketones, [1088](#)  
Ketones Blood Test, [1088](#)  
Ketones Urine Test, [1088](#)  
Ketonuria, [1088](#)  
Ketoprofen, [1088](#)  
Ketoprofen Overdose, [1088](#)  
Ketorolac, [1088](#)  
Ketorolac Tromethamine, [1088](#)  
Ketosis, [1088](#)  
Ketosis-Prone Type 2 Diabetes, [1088](#)  
Ketotic Hypoglycemia, [1088](#)  
Ketotifen, [1088](#)  
Kidney, [1088](#)  
Kidney Abscess, [1089](#)  
Kidney Biopsy, [1089](#)  
Kidney Calculi, [1089](#)  
Kidney Cancer, [1089](#)  
Kidney Cancer In Children, [1089](#)  
Kidney Cysts, [1089](#)  
Kidney Disease, [1089](#)  
Kidney Disease and Pregnancy, [1089](#)  
Kidney Disease, Chronic, [399](#)  
Kidney Disease, Diabetic, [545](#)  
Kidney Disease, Hypertension-Related, [969](#)  
Kidney Disease-Related Anemia, [1089](#)  
Kidney Diseases, [1089](#)  
Kidney Failure, [1089](#)  
Kidney Failure, Acute, [32](#)  
Kidney Function Tests, [1089](#)  
Kidney Infection, [1089](#)  
Kidney Neoplasm, [1090](#)  
Kidney Pain, [1090](#)  
Kidney Position, [1090](#)  
Kidney Removal, [1090](#)  
Kidney Replacement Therapy, [1090](#)  
Kidney Stone, [1090](#)  
Kidney Stones, [1090](#)  
Kidney Stones In Children, [1090](#)  
Kidney Transplant, [1090](#)  
Kidney Transplant Rejection, [1090](#)  
Kidney Transplantation, [1090](#)

---

---

Kidney Trauma, [1090](#)  
Kienbock Disease, [1090](#)  
Kieselguhr, [1090](#)  
Kiesselbach Plexus, [1091](#)  
Kiesselbach Triangle, [1091](#)  
Kikuchi Disease, [1091](#)  
Kikuchi-Fujimoto Disease, [1091](#)  
Killer Cold Virus, [1091](#)  
Killian Triangle, [1091](#)  
Killian's Triangle, [1091](#)  
Killip Classification, [1091](#)  
Kilocalorie, [1091](#)  
Kilojoule, [1091](#)  
Kimmelstiel-Wilson Lesion, [1091](#)  
Kimmelstiel-Wilson Nodules, [1091](#)  
Kimmelstiel-Wilson Syndrome, [1091](#)  
Kimura Disease, [1092](#)  
Kinase, [1092](#)  
Kinase Inhibitor, [1092](#)  
Kinesia Paradoxa, [1092](#)  
Kinesiology, [1092](#)  
Kinesthesia, [1092](#)  
Kinetics, [1092](#)  
Kinetochore, [1092](#)  
King's College Criteria, [1092](#)  
Kingella, [1092](#)  
Kingella Denitrificans, [1092](#)  
Kingella Kingae, [1092](#)  
Kingella Oralis, [1092](#)  
Kingella Potus, [1092](#)  
Kinin, [1093](#)  
Kinin-Kallikrein System, [1093](#)  
Kinins, [1093](#)  
Kinocilium, [1093](#)  
Kissing Disease, [1093](#)  
Kite Angle, [1093](#)  
Klüver-Bucy Syndrome, [1094](#)  
Klatskin Tumor, [1093](#)  
Klebsiella, [1093](#)  
Klebsiella Oxytoca, [1093](#)  
Klebsiella Pneumonia, [1093](#)  
Klebsiella Pneumoniae, [1093](#)  
Klebsiella Rhinoscleromatis, [1093](#)  
Klebsiella Singaporensis, [1093](#)  
Klebsiella Variicola, [1093](#)  
Kleihauer-Betke Test, [1094](#)  
Kleine-Levin Syndrome, [1094](#)  
Kleptomania, [1094](#)  
Klinefelter Syndrome, [1094](#)  
Klinefelter's Syndrome, [1094](#)  
Klippel-Feil Syndrome, [1094](#)  
Klippel-Trenaunay Syndrome, [1094](#)  
Klippel-Trenaunay-Weber Syndrome, [1094](#)  
Klumpke Palsy, [1095](#)  
Klumpke Paralysis, [1095](#)  
Klumpke's Paralysis, [1095](#)  
Kluyvera, [1095](#)  
Kluyvera Ascorbata, [1095](#)  
Kluyvera Cryocrescens, [1095](#)  
Kluyvera Georgiana, [1095](#)  
Kluyvera Intermedia, [1095](#)  
Knee, [1095](#)  
Knee Arthroplasty, [1095](#)  
Knee Arthroscopy, [1095](#)  
Knee Bursitis, [1095](#)  
Knee CT Scan, [1096](#)  
Knee Dislocation, [1096](#)  
Knee Disorders, [1096](#)  
Knee Effusion, [1096](#)  
Knee Hyperextension, [1096](#)  
Knee Injury And Meniscus Tears, [1096](#)  
Knee Jerk Reflex, [1096](#)  
Knee Joint, [1096](#)  
Knee Joint Replacement, [1096](#)  
Knee Meniscus, [1096](#)  
Knee Microfracture Surgery, [1097](#)  
Knee MRI Scan, [1097](#)  
Knee Pain, [1097](#)  
Knee Prosthesis, [1097](#)  
Knee Replacement, [1097](#)  
Knee Replacement Surgery, [1102](#)  
Knee-Chest Position, [1096](#)  
Knee-Jerk, [1096](#)  
Kneecap, [1095](#)  
Kneecap Dislocation, [1096](#)  
Kneecap Instability, [1102](#)  
Kneecap Tracking Disorder, [1102](#)  
Kniest Dysplasia, [1097](#)  
Knife-Cut Ulcers From Crohn'S Disease, [1097](#)  
Knock Knees, [1097](#)  
Knock-Knee, [1097](#)  
Knotless Anchor, [1097](#)  
Knuckle, [1097](#)  
Knuckle Pad, [1097](#)  
Koch's Bacillus, [1098](#)  
Kocher Clamp, [1097](#)  
Kocher Incision, [1098](#)  
Kocher Maneuver, [1098](#)  
Kocher-Debre-Semelaigne Syndrome, [1097](#)  
Kocher-Debré-SéméLaigne Syndrome, [1098](#)  
Koebner Phenomenon, [1098](#)  
Koebnerisation, [1098](#)  
KOH Preparation, [1098](#)  
Kohler Disease, [1098](#)  
Koilocyte, [1098](#)  
Koilonychia, [1098](#)  
Kombucha, [1098](#)  
Konno Procedure, [1099](#)

---

---

Koplik Spots, [1099](#)  
Koplik's Spots, [1099](#)  
Korotkoff Sounds, [1099](#)  
Korsakoff Syndrome, [1099](#)  
Korsakoff's Syndrome, [1099](#)  
Kostmann Syndrome, [1099](#)  
Koumpounophobia, [1099](#)  
Kounis Syndrome, [1099](#)  
KP, [1099](#)  
Krabbe Disease, [1099](#)  
Krabbe's Disease, [1099](#)  
KRAS Gene, [1099](#)  
Kratom, [1102](#)  
Kraurosis, [1100](#)  
Kraurosis Vulvae, [1100](#)  
Krebs Cycle, [1100](#)  
Kringles, [1100](#)  
Krukenberg Spindle, [1100](#)  
Krukenberg Tumor, [1100](#)  
Krukenberg's Tumor, [1100](#)  
Kt/V, [1100](#)  
KTW Syndrome, [1102](#)  
Kugelberg-Welander Disease, [1100](#)  
Kulchitsky Cells, [1100](#)  
Kupffer Cells, [1100](#)  
Kupffer's Cell, [1100](#)  
Kuru, [1101](#)  
Kussmaul Breathing, [1101](#)  
Kussmaul Respiration, [1101](#)  
Kussmaul Sign, [1101](#)  
Kveim-Siltzbach Test, [1101](#)  
Kwashiorkor, [1101](#)  
Kyasanur Forest Disease, [1101](#)  
Kynurenine, [1101](#)  
Kyphoplasty, [1101](#)  
Kyphoscoliosis, [1101](#)  
Kyphosis, [1101](#)  
Kyrle Disease, [1101](#)  
Kytococcus, [1102](#)  
Kytococcus Sedentarius, [1102](#)  
  
L-Dopa, [1120](#)  
L-theanine, [1165](#)  
La Crosse Encephalitis, [1106](#)  
LAAO, [1103](#)  
Lab Results, [1179](#)  
Labbe Vein, [1103](#)  
Labetalol, [1103](#)  
Labhart-Willi Syndrome, [1179](#)  
Labia, [1103](#)  
Labia Majora, [1103](#)  
Labia Minora, [1103](#)  
Labial Sounds, [1103](#)  
Labiaplasty, [1103](#)  
  
Labile, [1103](#)  
Labile Hypertension, [1103](#)  
Lability, [1103](#)  
Labium, [1103](#)  
Labium Majus, [1104](#)  
Labium Minus, [1104](#)  
Labor, [1104](#)  
Labor and Delivery, [1179](#)  
Labor and Delivery Complications, [1179](#)  
Labor Complication, [1104](#)  
Labor Dystocia, [1104](#)  
Labor Pain, [1104](#)  
Labor: Signs And Symptoms, [1104](#)  
Laboratory, [1104](#)  
Laboratory Medicine, [1104](#)  
Laboratory Procedure, [1104](#)  
Laboratory Test, [1104](#)  
Labour, [1104](#)  
Labra, [1104](#)  
Labral Tear, [1104](#)  
Labrum, [1104](#)  
Labyrinth, [1105](#)  
Labyrinth Disorder, [1105](#)  
Labyrinthine Membrane, [1105](#)  
Labyrinthine Positional Nystagmus, [1179](#)  
Labyrinthitis, [1105](#)  
Laceration, [1105](#)  
Laceration Suture and Staple Care, [1105](#)  
Lachman Sign, [1105](#)  
Lachman Test, [1105](#)  
Lachrymal, [1105](#)  
Lacosamide, [1105](#)  
Lacquer Poisoning, [1105](#)  
Lacrimal, [1105](#)  
Lacrimal Apparatus, [1105](#)  
Lacrimal Disease, [1105](#)  
Lacrimal Duct, [1105](#)  
Lacrimal Gland, [1105](#)  
Lacrimal Gland Tumor, [1106](#)  
Lacrimal Sac, [1106](#)  
Lacrimation, [1106](#)  
Lactam, [1106](#)  
Lactase, [1106](#)  
Lactase Deficiency, [1106](#)  
Lactate, [1106](#)  
Lactate Dehydrogenase, [1106](#)  
Lactate Dehydrogenase Test, [1106](#)  
Lactation, [1106](#)  
Lactation Disorder, [1106](#)  
Lactation Mastitis, [1106](#)  
Lactation Nutrition, [1106](#)  
Lactational Amenorrhea, [1106](#)  
Lacteals, [1106](#)  
Lactic Acid, [1107](#)

---

---

Lactic Acid Test, [1107](#)  
Lactic Acidosis, [1107](#)  
Lactiferous Duct, [1107](#)  
Lactobacillus, [1107](#)  
Lactoferrin, [1107](#)  
Lactose, [1107](#)  
Lactose Fermentation, [1107](#)  
Lactose Intolerance, [1107](#)  
Lactose Tolerance Test, [1107](#)  
Lactose Tolerance Tests, [1107](#)  
Lactulose, [1107](#)  
Lacuna, [1107](#)  
Lacunar Infarct, [1107](#)  
Lacunar Stroke, [1108](#)  
Lacunar Syndromes (Small-Vessel Strokes), [1108](#)  
Ladd Bands, [1108](#)  
Ladd Procedure, [1108](#)  
Laetrile, [1108](#)  
Lafora Disease, [1108](#)  
Lagophthalmos, [1108](#)  
LAM, [1179](#)  
Lambda, [1108](#)  
Lambert-Eaton Myasthenic Syndrome, [1108](#)  
Lamellar, [1108](#)  
Lamellar Body, [1108](#)  
Lamellar Bone, [1108](#)  
Lamellar Ichthyosis, [1108](#)  
Lamin, [1109](#)  
Lamina, [1109](#)  
Lamina Cribrosa, [1109](#)  
Lamina Propria, [1109](#)  
Laminectomy, [1109](#)  
Laminin, [1109](#)  
Lamivudine, [1109](#)  
Lamotrigine, [1109](#)  
Lancet, [1109](#)  
Landolt Ring, [1109](#)  
Langer Lines, [1110](#)  
Langerhans Cell Histiocytosis, [1109](#)  
Langerhans Cells, [1109](#)  
Langerhans' Cell, [1109](#)  
Langerhans-Cell Histiocytosis, [1109](#)  
Language Barrier, [1110](#)  
Language Development, [1110](#)  
Language Disorder, [1110](#)  
Language Therapy, [1110](#)  
Lanolin, [1110](#)  
Lanolin Poisoning, [1110](#)  
Lansoprazole, [1110](#)  
Lanthanum, [1110](#)  
Lanugo, [1110](#)  
Lanugo Hair, [1110](#)  
Laparoscope, [1110](#)  
Laparoscopic Cholecystectomy, [1110](#)  
Laparoscopic Colectomy, [1110](#)  
Laparoscopic Gallbladder Removal, [1110](#)  
Laparoscopic Gastric Banding, [1110](#)  
Laparoscopic Hysterectomy, [1111](#)  
Laparoscopic Nephrectomy, [1111](#)  
Laparoscopic Surgery, [1111](#)  
Laparoscopy, [1111](#)  
Laparotomy, [1111](#)  
Lapatinib, [1111](#)  
Lapatinib Ditosylate, [1111](#)  
Laplace Law, [1111](#)  
Large Airway Obstruction, [1111](#)  
Large Bowel Obstruction, [1111](#)  
Large Bowel Resection, [1111](#)  
Large Colon, [1111](#)  
Large For Gestational Age, [1111](#)  
Large Intestine, [1112](#)  
Laron Syndrome, [1112](#)  
Larva Currens From Strongyloides, [1112](#)  
Larva Migrans, [1112](#)  
Laryngeal, [1112](#)  
Laryngeal Botulinum Toxin Injection, [1112](#)  
Laryngeal Cancer, [1112](#)  
Laryngeal Candidiasis, [1112](#)  
Laryngeal Carcinoma, [1112](#)  
Laryngeal Cartilage, [1112](#)  
Laryngeal Cleft, [1112](#)  
Laryngeal Edema, [1113](#)  
Laryngeal Fistula, [1113](#)  
Laryngeal Hemorrhage, [1113](#)  
Laryngeal Masks, [1113](#)  
Laryngeal Mucositis, [1113](#)  
Laryngeal Nerve, [1113](#)  
Laryngeal Nerve Damage, [1113](#)  
Laryngeal Obstruction, [1113](#)  
Laryngeal Papillomatosis, [1113](#)  
Laryngeal Prosthesis, [1113](#)  
Laryngeal Stridor, [1113](#)  
Laryngeal Web, [1113](#)  
Laryngectomy, [1113](#)  
Laryngectomy Tube, [1113](#)  
Laryngitis, [1114](#)  
Laryngocele, [1114](#)  
Laryngomalacia, [1114](#)  
Laryngopharyngeal Reflux, [1114](#)  
Laryngopharynx, [1114](#)  
Laryngoscope, [1114](#)  
Laryngoscopy, [1114](#)  
Laryngoscopy And Nasolaryngoscopy, [1114](#)  
Laryngospasm, [1114](#)  
Laryngostenosis, [1114](#)  
Laryngotracheobronchitis, [1114](#)  
Larynx, [1114](#)

---



---

Larynx Cancer, [1114](#)  
 Larynx Disease, [1114](#)  
 Larynx Muscle, [1115](#)  
 Larynx Neoplasm, [1115](#)  
 Lasague Sign, [1115](#)  
 Laser, [1115](#)  
 Laser Ablation, [1115](#)  
 Laser Assisted Uvula Palatoplasty, [1115](#)  
 Laser Fiber, [1115](#)  
 Laser Photocoagulation, [1115](#)  
 Laser Surgery, [1115](#)  
 Laser Surgery For The Skin, [1115](#)  
 Laser Therapy, [1115](#)  
 Laser Therapy For Cancer, [1115](#)  
 Laser Trabeculoplasty, [1115](#)  
 Laser-Assisted In Situ Keratomileusis, [1115](#)  
 LASIK, [1116](#)  
 LASIK Eye Surgery, [1116](#)  
 Lassa Fever, [1116](#)  
 Lassa Virus, [1116](#)  
 Lassitude, [1116](#)  
 Last Menstrual Period, [1116](#)  
 Latanoprost, [1116](#)  
 Latarjet Procedure, [1116](#)  
 Late Decelerations, [1116](#)  
 Late Effects, [1116](#)  
 Late Gadolinium Enhancement, [1116](#)  
 Late Preterm Birth, [1117](#)  
 Late-Onset Hypogonadism, [1117](#)  
 Latent, [1116](#)  
 Latent Autoimmune Diabetes In Adults, [1116](#)  
 Latent Infection, [1117](#)  
 Latent Phase Of Labor, [1117](#)  
 Latent Syphilis, [1117](#)  
 Latent Tuberculosis, [1117](#)  
 Latent Tuberculosis Infection, [1117](#)  
 Lateral, [1117](#)  
 Lateral Canthotomy And Cantholysis, [1117](#)  
 Lateral Collateral Ligament Injury, [1117](#)  
 Lateral Cuneiform, [1117](#)  
 Lateral Cutaneous Nerve of the Thigh, [1117](#)  
 Lateral Decubitus Position, [1117](#)  
 Lateral Elbow Tendinopathy, [1118](#)  
 Lateral Epicondylitis, [1118](#)  
 Lateral Epicondylolysis, [1118](#)  
 Lateral Geniculate Nucleus, [1118](#)  
 Lateral Medullary Syndrome, [1118](#)  
 Lateral Meniscus, [1118](#)  
 Lateral Pontine Syndrome, [1118](#)  
 Lateral Rotation, [1118](#)  
 Lateral Sclerosis, [1118](#)  
 Lateral Traction, [1118](#)  
 Lateral Ventricle, [1118](#)  
 Lateral X-Ray, [1118](#)  
 Latex Agglutination Test, [1118](#)  
 Latex Allergy, [1118](#)  
 Lathyrism, [1119](#)  
 Latissimus Dorsi, [1119](#)  
 Latissimus Dorsi Muscle, [1119](#)  
 Lattice Degeneration, [1119](#)  
 Lavage, [1119](#)  
 Lavage Fluid, [1119](#)  
 Lawton IADL Scale, [1119](#)  
 Laxative, [1123](#)  
 Laxative Overdose, [1119](#)  
 Laxatives, [1179](#)  
 Laxatives For Constipation, [1119](#)  
 Lazy Eye, [1119](#)  
 Lazy Eye (Amblyopia), [1179](#)  
 LCH, [1179](#)  
 LCIS, [1119](#)  
 LDH Isoenzyme Blood Test, [1119](#)  
 LDL, [1119](#)  
 LDL Cholesterol, [1179](#)  
 LDL Test, [1119](#)  
 Le Fort Fractures, [1121](#)  
 Lead, [1120](#)  
 Lead And Tap Water, [1120](#)  
 Lead Pipe Rigidity, [1120](#)  
 Lead Poisoning, [1120](#)  
 Lead Toxicity and Nutrition, [1120](#)  
 Lead-Pipe Rigidity, [1120](#)  
 Lead-Time Bias, [1120](#)  
 Leadless Pacemaker, [1120](#)  
 Learned Insomnia, [1120](#)  
 Learning Difficulties, [1120](#)  
 Learning Disabilities, [1120](#)  
 Learning Disorder, [1121](#)  
 Leber Congenital Amaurosis, [1121](#)  
 Leber Hereditary Optic Neuropathy, [1121](#)  
 Leber's Hereditary Optic Neuropathy, [1121](#)  
 Lecithin-Sphingomyelin Ratio, [1121](#)  
 Leech, [1121](#)  
 Leech Therapy, [1121](#)  
 Leflunomide, [1121](#)  
 Left Anterior Descending Artery, [1121](#)  
 Left Anterior Descending Coronary Artery, [1121](#)  
 Left Anterior Fascicular Block, [1122](#)  
 Left Atrial Appendage Occlusion, [1122](#)  
 Left Atrial Myxoma, [1122](#)  
 Left Atrium, [1122](#)  
 Left Bundle Branch, [1122](#)  
 Left Bundle Branch Block, [1122](#)  
 Left Circumflex Coronary Artery, [1122](#)  
 Left Heart Catheterization, [1122](#)  
 Left Heart Ventricular Angiography, [1122](#)  
 Left Hypochondriac Region, [1122](#)  
 Left Iliac Region, [1122](#)

---

---

Left Main Coronary Artery, [1122](#)  
Left Testicular Torsion, [1123](#)  
Left Ventricle, [1123](#)  
Left Ventricular Assist Device, [1123](#)  
Left Ventricular Hypertrophy, [1123](#)  
Left Ventricular Non-Compaction, [1123](#)  
Left Ventricular Noncompaction, [1123](#)  
Left-Sided Superior Vena Cava, [1122](#)  
Left-Ventricular Assist Device, [1123](#)  
Leg, [1123](#)  
Leg CT Scan, [1123](#)  
Leg Fracture, [1123](#)  
Leg Lengthening And Shortening, [1124](#)  
Leg MRI Scan, [1124](#)  
Leg Or Foot Amputation, [1124](#)  
Leg Pain, [1124](#)  
Leg Pain With Cramping, [1124](#)  
Leg Swelling, [1124](#)  
Leg Ulcer, [1124](#)  
Legal Medicine, [2123](#)  
Legg-Calve-Perthes Disease, [1123](#)  
Legg-Calvé-Perthes Disease, [1123](#)  
Legionella, [1123](#)  
Legionella Longbeachae, [1124](#)  
Legionella Pneumonia, [1124](#)  
Legionella Pneumophila, [1124](#)  
Legionellosis, [1124](#)  
Legionnaire Disease And Pontiac Fever, [1124](#)  
Legionnaires Disease, [1124](#)  
Legionnaires' Disease, [1124](#)  
Leigh Disease, [1125](#)  
Leigh Syndrome, [1125](#)  
Leigh's Syndrome, [1125](#)  
Leiomyoma, [1125](#)  
Leiomyomatosis, [1125](#)  
Leiomyomatosis Peritonealis Disseminata, [1125](#)  
Leiomyosarcoma, [1125](#)  
Leiomyosarcoma Of The Vulva, [1125](#)  
Leishmania, [1125](#)  
Leishmania Donovanii, [1125](#)  
Leishmania Infection, [1125](#)  
Leishmania Major, [1125](#)  
Leishmania Mexicana, [1125](#)  
Leishmania Tropica, [1126](#)  
Leishmaniasis, [1126](#)  
Lemaitre's Sign, [1126](#)  
Lemierre Syndrome, [1126](#)  
Lenalidomide, [1126](#)  
Lenegre Disease, [1126](#)  
Length-Time Bias, [1126](#)  
Lengthening Of Limb, [1126](#)  
Lennox-Gastaut Syndrome, [1126](#)  
Lenograstim, [1126](#)  
Lens, [1126](#)  
Lens Coloboma, [1126](#)  
Lens Disease, [1126](#)  
Lens Dislocation, [1126](#)  
Lens Prescription, [1127](#)  
Lens Subluxation, [1127](#)  
Lensometer, [1127](#)  
Lentiform Nucleus, [1127](#)  
Lentigines, [1127](#)  
Lentigo, [1127](#)  
Lentigo Maligna, [1127](#)  
Lentigo Maligna Melanoma, [1127](#)  
LEOPARD Syndrome, [1127](#)  
Leopold Maneuvers, [1127](#)  
Lepidoptera, [1127](#)  
Lepirudin, [1127](#)  
Lepromatous Leprosy, [1127](#)  
Lepromin Skin Test, [1127](#)  
Leprosy, [1128](#)  
Leptin, [1128](#)  
Leptin Receptor, [1128](#)  
Leptomeningeal, [1128](#)  
Leptomeningeal Metastases, [1128](#)  
Leptomeningeal Metastasis, [1128](#)  
Leptomeningeal Sarcoma, [1128](#)  
Leptomeninges, [1128](#)  
Leptomeningitis, [1128](#)  
Leptospira, [1128](#)  
Leptospirosis, [1128](#)  
Leri-Weill Dyschondrosteosis, [1129](#)  
Leriche Syndrome, [1128](#)  
Lesch-Nyhan Syndrome, [1129](#)  
Lesion, [1129](#)  
Lesion Identification, [1129](#)  
Lesser Curvature, [1129](#)  
Lesser Omentum, [1129](#)  
Lesser Peritoneal Cavity, [1129](#)  
Lesser Toe Deformities, [1112](#)  
Lesser Trochanter, [1129](#)  
Let-Down Reflex, [1129](#)  
Lethal, [1129](#)  
Lethal Triad, [1129](#)  
Lethal-Means Counseling, [1129](#)  
Lethargy, [1129](#)  
Letrozole, [1130](#)  
Letterer-Siwe Disease, [1130](#)  
Leucocyte, [1130](#)  
Leucocytosis, [1130](#)  
Leucoderma, [1130](#)  
Leucomalacia, [1130](#)  
Leucopenia, [1130](#)  
Leucoplakia, [1130](#)  
Leucopoiesis, [1130](#)  
Leucovorin, [1130](#)  
Leucovorin Calcium, [1130](#)

---

---

Leukaemia, [1130](#)  
Leukapheresis, [1130](#)  
Leukemia, [1130](#)  
Leukemia Cutis, [1131](#)  
Leukemia Genetics, [1131](#)  
Leukemic Infiltration Of The Gingiva, [1131](#)  
Leukemoid Reaction, [1131](#)  
Leukoaraiosis, [1131](#)  
Leukocoria, [1131](#)  
Leukocyte, [1131](#)  
Leukocyte Adhesion Deficiency, [1131](#)  
Leukocyte Alkaline Phosphatase, [1131](#)  
Leukocyte Disorder, [1131](#)  
Leukocyte Esterase Urine Test, [1131](#)  
Leukocyte Transfusion, [1132](#)  
Leukocytes, [1131](#)  
Leukocytoclastic Vasculitis, [1132](#)  
Leukocytosis, [1132](#)  
Leukocytosis in Pregnancy, [1132](#)  
Leukoderma, [1132](#)  
Leukodystrophy, [1132](#)  
Leukoencephalopathy, [1132](#)  
Leukonychia, [1132](#)  
Leukonychia Striata, [1132](#)  
Leukonychia Totalis, [1132](#)  
Leukopenia, [1132](#)  
Leukoplakia, [1132](#)  
Leukopoiesis, [1132](#)  
Leukorrhea, [1133](#)  
Leukostasis, [1133](#)  
Leukotriene, [1133](#)  
Leukotriene Blockers, [1133](#)  
Leukotriene Modifiers, [1133](#)  
Leukotriene Receptor, [1133](#)  
Leukotrienes, [1133](#)  
Leuprolide, [1133](#)  
Leuprolide Acetate, [1133](#)  
Lev Disease, [1134](#)  
Levalbuterol, [1133](#)  
Levamisole, [1133](#)  
Levator Ani, [1133](#)  
Levator Ani Muscle, [1133](#)  
Levator Ani Syndrome, [1133](#)  
Levetiracetam, [1134](#)  
Levine Sign, [1134](#)  
Levobunolol, [1134](#)  
Levocardia, [1134](#)  
Levocarnitine, [1134](#)  
Levodopa, [1134](#)  
Levofloxacin, [1134](#)  
Levoleucovorin, [1134](#)  
Levoleucovorin Calcium, [1134](#)  
Levomepromazine, [1134](#)  
Levonorgestrel, [1134](#)  
Levorphanol, [1134](#)  
Levothyroxine, [1135](#)  
Levothyroxine Sodium, [1135](#)  
Levulose, [1135](#)  
Lewy Bodies, [1135](#)  
Lewy Body, [1135](#)  
Lewy Body Dementia, [1135](#)  
Lewy Body Disease, [1135](#)  
Leydig Cell, [1135](#)  
Leydig Cell Testicular Tumor, [1135](#)  
Leydig Cell Tumor, [1135](#)  
Leydig Cells, [1135](#)  
LGE, [1135](#)  
LGMD, [1179](#)  
LGS, [1179](#)  
LH, [1136](#)  
LH Response To GnRH Blood Test, [1136](#)  
Lhermitte Sign, [1136](#)  
Lhermitte's Sign, [1136](#)  
Lhermitte-Duclos Disease, [1136](#)  
Li-Fraumeni Syndrome, [1138](#)  
LI-RADS, [1148](#)  
Libido, [1136](#)  
Libido And Sexual Performance, [1136](#)  
Libman-Sacks Endocarditis, [1136](#)  
Lice, [1136](#)  
Lice Infestation, [1136](#)  
Lice Infestations, [1136](#)  
Licensed Clinical Social Worker, [1136](#)  
Lichen, [1136](#)  
Lichen Amyloidosis, [1136](#)  
Lichen Annularis, [1179](#)  
Lichen Myxedematosus, [1137](#)  
Lichen Nitidus, [1137](#)  
Lichen Planopilaris, [1137](#)  
Lichen Planus, [1137](#)  
Lichen Planus Sclerosus Atrophicus, [1179](#)  
Lichen Psoriasis, [1179](#)  
Lichen Ruber Planus, [1137](#)  
Lichen Sclerosus, [1137](#)  
Lichen Simplex Chronicus, [1137](#)  
Lichen Spinulosus, [1137](#)  
Lichen Striatus, [1137](#)  
Lichen Urticatus, [1180](#)  
Lichenification, [1137](#)  
Lichenified, [1137](#)  
Lichenoid Drug Eruption, [1137](#)  
Licorice Poisoning, [1138](#)  
Liddle Syndrome, [1138](#)  
Liddle Syndrome Treatment, [1138](#)  
Lidocaine, [1138](#)  
Lieberkühn's Crypts, [1138](#)  
Life Expectancy, [1138](#)  
Life Support, [1180](#)

---

---

Lifetime Risk, [1138](#)  
Ligament, [1138](#)  
Ligamentum Flavum, [1138](#)  
Ligamentum Teres Hepatis, [1138](#)  
Ligand, [1138](#)  
Ligase, [1138](#)  
Ligation, [1139](#)  
Ligature, [1139](#)  
Light, [1139](#)  
Light Chain-Associated Kidney Disease, [1139](#)  
Light Coagulation, [1139](#)  
Light Cycle, [1139](#)  
Light Microscope, [1139](#)  
Light Microscopy, [1139](#)  
Light Reflex, [1140](#)  
Light Scattering, [1140](#)  
Light Therapy, [1140](#)  
Light's Criteria, [1140](#)  
Light-Chain Deposition Disease, [1139](#)  
Light-Near Dissociation, [1139](#)  
Lightening, [1139](#)  
Lighter Fluid Poisoning, [1139](#)  
Lightheadedness, [1139](#)  
Lightning Injuries, [1139](#)  
Lightning Pains, [1139](#)  
Lignocaine, [1140](#)  
Likelihood Ratio, [1140](#)  
Lilliputian Hallucination, [1140](#)  
Lily Of The Valley Poisoning, [1140](#)  
Limb, [1140](#)  
Limb Bud, [1140](#)  
Limb Development, [1140](#)  
Limb Fracture, [1140](#)  
Limb Girdle, [1140](#)  
Limb Injury, [1141](#)  
Limb Pain, [1141](#)  
Limb Plethysmography, [1141](#)  
Limb Prosthesis, [1141](#)  
Limb Reduction Defects, [1141](#)  
Limb-Girdle Muscular Dystrophies, [1140](#)  
Limb-Length Discrepancy, [1141](#)  
Limbic Encephalitis, [1140](#)  
Limbic Lobe, [1140](#)  
Limbic System, [1141](#)  
Limbs, [1141](#)  
Limbus, [1141](#)  
Limbus Corneae, [1141](#)  
Limited Range Of Motion, [1141](#)  
Linagliptin, [1141](#)  
Lincomycin, [1141](#)  
Lindane, [1141](#)  
Lindsay's Nails, [1141](#)  
Linea, [1142](#)  
Linea Alba, [1142](#)  
Linea Aspera, [1142](#)  
Linea Nigra, [1142](#)  
Linear, [1142](#)  
Linear Accelerator, [1142](#)  
Linear Energy Transfer, [1142](#)  
Linear Fracture, [1142](#)  
Linear IgA Bullous Dermatitis, [1142](#)  
Linear Nevus Sebaceous Syndrome, [1142](#)  
Lined Canister, [1142](#)  
Lines of Blaschko, [1142](#)  
Lines of Zahn, [1142](#)  
Linezolid, [1143](#)  
Lingua, [1143](#)  
Lingua Fissurata, [1180](#)  
Lingua Nigra, [1180](#)  
Lingua Villosa Nigra, [1143](#)  
Lingual, [1143](#)  
Lingual Artery, [1143](#)  
Lingual Thyroid, [1143](#)  
Lingual Tonsil, [1143](#)  
Lingual Tonsils, [1143](#)  
Linguofacial Dyskinesia, [1180](#)  
Liniment, [1143](#)  
Linitis Plastica, [1143](#)  
Linoleic Acid, [1143](#)  
Linolenic Acid, [1143](#)  
Liothyronine Sodium, [1143](#)  
Lip, [1143](#)  
Lip Cancer, [1144](#)  
Lip Disease, [1144](#)  
Lip Infection, [1145](#)  
Lip Moisturizer Poisoning, [1145](#)  
Lip Neoplasm, [1145](#)  
Lip Swelling, [1147](#)  
Lipase, [1143](#)  
Lipase Test, [1144](#)  
Lipedema, [1144](#)  
Lipemia Retinalis, [1144](#)  
Lipemic Index, [1144](#)  
Lipid, [1144](#)  
Lipid A, [1144](#)  
Lipid Bilayer, [1144](#)  
Lipid Disorder, [1144](#)  
Lipid Emulsion Therapy, [1144](#)  
Lipid Metabolism, [1144](#)  
Lipid Panel, [1144](#)  
Lipid Profile, [1145](#)  
Lipid Profile Test, [1145](#)  
Lipid Transport, [1145](#)  
Lipidosis, [1144](#)  
Lipids, [1145](#)  
Lipitor, [1145](#)  
Lipoatrophy, [1145](#)  
Lipoblastoma, [1145](#)

---

---

Lipodermatosclerosis, [1145](#)  
Lipodystrophy, [1145](#)  
Lipodystrophy Syndromes, [1145](#)  
Lipofuscin, [1145](#)  
Lipogranulomatosis, [1146](#)  
Lipohypertrophy, [1146](#)  
Lipoid Nephrosis, [1146](#)  
Lipoid Pneumonia, [1146](#)  
Lipoidosis, [1146](#)  
Lipolysis, [1146](#)  
Lipoma, [1146](#)  
Lipomas, [1146](#)  
Lipomastia, [1146](#)  
Lipomatosis, [1146](#)  
Lipomatous Neoplasm, [1146](#)  
Lipophilicity, [1146](#)  
Lipopolysaccharide, [1146](#)  
Lipoprotein, [1146](#)  
Lipoprotein A, [1146](#)  
Lipoprotein Analysis, [1147](#)  
Lipoprotein Lipase, [1147](#)  
Lipoprotein Metabolism, [1147](#)  
Lipoprotein(a), [1147](#)  
Liposarcoma, [1147](#)  
Liposarcoma Of The Breast, [1147](#)  
Liposomal Amphotericin B, [1147](#)  
Liposome, [1147](#)  
Liposuction, [1147](#)  
Lips, [1147](#)  
Lipschütz Bodies, [1147](#)  
Liquefaction, [1147](#)  
Liquefactive Necrosis, [1148](#)  
Liquid Bandage for Lacerations, [1148](#)  
Liquid Biopsy, [1148](#)  
Liquid Chromatography, [1148](#)  
Liquid Diet, [1148](#)  
Liquid Incense, [1148](#)  
Liquid Medication Administration, [2123](#)  
Liquid Nitrogen, [1148](#)  
Liquor, [1148](#)  
Liraglutide, [1148](#)  
Lisch Nodule, [1148](#)  
Lisch Nodules, [1148](#)  
Lisfranc Fracture, [1148](#)  
Lisfranc Injury, [1149](#)  
Lisinopril, [1149](#)  
Lissencephaly, [1149](#)  
Listeria, [1149](#)  
Listeria Infection, [1149](#)  
Listeria Monocytogenes, [1149](#)  
Listeriosis, [1149](#)  
Listeriosis In Pregnancy, [1149](#)  
Lithiasis, [1149](#)  
Lithium, [1149](#)  
Lithium Carbonate, [1149](#)  
Lithium Nephropathy, [1149](#)  
Lithium Toxicity, [1149](#)  
Lithotomy, [1150](#)  
Lithotomy Position, [1150](#)  
Lithotripsy, [1150](#)  
Lithotripsy Probe, [1150](#)  
Lithotripter, [1150](#)  
Lithotripter Probe, [1150](#)  
Litmus, [1150](#)  
Little Area, [1150](#)  
Little's Disease, [1150](#)  
Live Attenuated Vaccines, [1150](#)  
Live Birth, [1150](#)  
Livedo Racemosa, [1151](#)  
Livedo Reticularis, [1151](#)  
Livedoid Vasculopathy, [1150](#)  
Liver, [1151](#)  
Liver Abscess, [1151](#)  
Liver Anatomy and Physiology, [1151](#)  
Liver Biopsy, [1151](#)  
Liver Cancer, [1151](#)  
Liver Cancer Treatment, [1180](#)  
Liver Circulation, [1151](#)  
Liver Cirrhosis, [1151](#)  
Liver Damage, [1151](#)  
Liver Disease, [1151](#)  
Liver Disease and Pregnancy, [1151](#)  
Liver Disease Diet, [1151](#)  
Liver Diseases, [1151](#)  
Liver Disorder, [1152](#)  
Liver Dysfunction, [1152](#)  
Liver Elastography, [1152](#)  
Liver Enzymes, [1152](#)  
Liver Failure, [1152](#)  
Liver Fibrosis, [1152](#)  
Liver Fluke, [1152](#)  
Liver Function, [1152](#)  
Liver Function Tests, [1152](#)  
Liver Health, [1152](#)  
Liver Infection, [1152](#)  
Liver Lobule, [1152](#)  
Liver Mass, [1152](#)  
Liver Metastases, [1152](#)  
Liver Neoplasm, [1153](#)  
Liver Pain, [1153](#)  
Liver Regeneration, [1153](#)  
Liver Scan, [1153](#)  
Liver Shunt, [1153](#)  
Liver Spots, [1153](#)  
Liver Transplant, [1153](#)  
Liver Transplantation, [1153](#)  
Living Donor, [1153](#)  
Living Will, [1153](#)

---

---

Livor Mortis, [1153](#)  
Lizard Bite, [1153](#)  
LLETZ, [1153](#)  
Lloyd-Davies Position, [1153](#)  
Loa Loa, [1154](#)  
Loading Dose, [1153](#)  
Lobaplatin, [1154](#)  
Lobar Hemorrhage, [1154](#)  
Lobar Pneumonia, [1154](#)  
Lobe, [1154](#)  
Lobectomy, [1154](#)  
Lobotomy, [1154](#)  
Lobular Carcinoma, [1154](#)  
Lobular Carcinoma In Situ, [1154](#)  
Lobule, [1154](#)  
Lobules, [1154](#)  
Local, [1154](#)  
Local Anesthesia, [1154](#)  
Local Anesthetic, [2123](#)  
Local Anesthetic Systemic Toxicity, [1154](#)  
Local Anesthetic Toxicity, [1155](#)  
Local Invasion, [1155](#)  
Local Therapy, [1155](#)  
Localised, [1155](#)  
Localization, [1155](#)  
Localized, [1155](#)  
Localized Edema, [1155](#)  
Localized Epidermolysis Bullosa, [1180](#)  
Localized Lipodystrophy, [1155](#)  
Localized Scleroderma, [1155](#)  
Lochia, [1155](#)  
Locked-In Syndrome, [1155](#)  
Lockjaw, [1155](#)  
Locomotion, [1155](#)  
Locomotive System, [1155](#)  
Locomotor Ataxia, [1156](#)  
Locoregional, [1156](#)  
Locus, [1156](#)  
Locus Ceruleus, [1156](#)  
Locus Coeruleus, [1156](#)  
Locus Minoris Resistentiae, [1156](#)  
Lod Score, [1156](#)  
Loeffler Endocarditis, [1156](#)  
Loeffler Syndrome, [1156](#)  
Loeys-Dietz Syndrome, [1156](#)  
Log Roll Test, [1156](#)  
Loiasis, [1157](#)  
Loin, [1157](#)  
Lomotil Overdose, [1157](#)  
Lomustine, [1157](#)  
Lonafarnib, [1157](#)  
Long Arm Of A Chromosome, [1157](#)  
Long Bones, [1157](#)  
Long COVID, [1157](#)  
Long QT Syndrome, [1158](#)  
Long-Acting Beta-2 Agonists, [1157](#)  
Long-Acting Reversible Contraception, [1157](#)  
Long-Sightedness, [1158](#)  
Long-Term Care, [1158](#)  
Long-Term Care Facility, [1158](#)  
Long-Term Complications Of Diabetes, [1158](#)  
Long-Term Effect, [1158](#)  
Long-Term Memory, [1158](#)  
Long-Term Potentiation, [1158](#)  
Long-Term Rehabilitation, [1158](#)  
Longitudinal Axis, [1157](#)  
Longitudinal Section, [1157](#)  
Longitudinal Studies, [1157](#)  
Longitudinal Vaginal Septum, [1157](#)  
Loop Diuretic, [1158](#)  
Loop Diuretics, [1158](#)  
Loop Electrosurgical Excision Procedure, [1158](#)  
Loop Of Henle, [1158](#)  
Loose Bodies, [1158](#)  
Loose Body, [1159](#)  
Loperamide, [1159](#)  
Loperamide Hydrochloride, [1159](#)  
Lopinavir, [1159](#)  
Loratadine, [1159](#)  
Lorazepam, [1159](#)  
Lordosis, [1159](#)  
Losartan, [1159](#)  
Losartan Potassium, [1159](#)  
Loss Of Appetite, [1159](#)  
Loss of Bladder Control, [1159](#)  
Loss Of Consciousness, [1159](#)  
Loss Of Heterozygosity, [1159](#)  
Loss of Smell, [1160](#)  
Loss-Of-Function Mutation, [1159](#)  
Loss-Of-Function Variant, [1159](#)  
Loteprednol Etabonate, [1160](#)  
Lou Gehrig Disease, [1160](#)  
Lou Gehrig's Disease, [1160](#)  
Louse, [1160](#)  
Lovastatin, [1160](#)  
Low Back Pain, [1160](#)  
Low Back Pain Deciding About MRI, [1180](#)  
Low Back Pain Exercises, [1180](#)  
Low Birth Weight, [1160](#)  
Low Blood Oxygen, [1160](#)  
Low Blood Potassium, [1160](#)  
Low Blood Pressure, [1160](#)  
Low Blood Pressure Causes, [1160](#)  
Low Blood Sodium, [1160](#)  
Low Blood Sugar, [1160](#)  
Low Body Temperature, [1160](#)  
Low Eye Tension, [1162](#)  
Low FODMAP Diet, [1162](#)

---



---

Low Grade, [1162](#)  
Low HDL Cholesterol (Hypoalphalipoproteinemia), [1163](#)  
Low Hemoglobin Count, [1163](#)  
Low LDL Cholesterol (Hypobetalipoproteinemia), [1163](#)  
Low Molecular Weight Heparin, [1163](#)  
Low Nasal Bridge, [1163](#)  
Low Neutrophil Count, [1163](#)  
Low Platelet Count, [1163](#)  
Low Potassium, [1163](#)  
Low Potassium Syndrome, [1180](#)  
Low Sperm Count, [1164](#)  
Low T, [1164](#)  
Low Testosterone, [1164](#)  
Low Vision, [1164](#)  
Low White Blood Cell Count, [1164](#)  
Low White Cell Count in Cancer, [1164](#)  
Low-Calorie Diet, [1160](#)  
Low-Density Lipoprotein, [1161](#)  
Low-Dose Aspirin Therapy, [1180](#)  
Low-Dose CT Scan, [1161](#)  
Low-Fiber Diet, [1162](#)  
Low-Flow Low-Gradient Aortic Stenosis, [1162](#)  
Low-Flow Priapism, [1162](#)  
Low-Grade Appendiceal Mucinous Neoplasm, [1162](#)  
Low-Grade Astrocytoma, [1162](#)  
Low-Grade Central Osteosarcoma, [1162](#)  
Low-Grade Endometrial Stromal Sarcoma, [1163](#)  
Low-Molecular-Weight Heparin, [1163](#)  
Low-Phosphorus Diet, [1163](#)  
Low-Potassium Diet, [1163](#)  
Low-Residue Diet, [1164](#)  
Low-Salt Diet, [1164](#)  
Low-Set Ear, [1164](#)  
Low-Set Ears And Pinna Abnormalities, [1164](#)  
Low-Tension Glaucoma, [1164](#)  
Lowe's Syndrome, [1162](#)  
Lower Airway Obstruction, [1161](#)  
Lower Esophageal Ring, [1161](#)  
Lower Esophageal Sphincter, [1161](#)  
Lower Extremity, [1161](#)  
Lower Extremity Amputation Dressing Care, [1161](#)  
Lower Extremity Arterial Bypass, [1161](#)  
Lower Extremity Peripheral Artery Disease, [1161](#)  
Lower Extremity Venogram, [1161](#)  
Lower Extremity Venous Doppler, [1161](#)  
Lower Gastrointestinal Bleeding, [1161](#)  
Lower Leg, [1162](#)  
Lower Motor Neuron Syndrome, [1162](#)  
Lower Segment Cesarean Section, [1162](#)  
Lower Urinary Tract Symptoms, [1162](#)  
Lown-Ganong-Levine Syndrome, [1163](#)  
Lozenges, [1164](#)  
LRINEC Score, [1164](#)  
LSD, [1164](#)  
LSD Substance Use, [1164](#)  
LSD Toxicity, [1165](#)  
Lubiprostone, [1165](#)  
Lubricant, [1165](#)  
Lucilia, [1165](#)  
Ludwig Angina, [1165](#)  
Ludwig's Angina, [1165](#)  
Lugano Classification, [1165](#)  
Lugol's Iodine, [1165](#)  
Lugol's Solution, [1165](#)  
Lumbago, [1165](#)  
Lumbar, [1165](#)  
Lumbar Disc Herniation, [1165](#)  
Lumbar Disk Replacement, [1165](#)  
Lumbar Hernia, [1166](#)  
Lumbar Lordosis, [1166](#)  
Lumbar Lordosis Lower Spine, [1166](#)  
Lumbar MRI Scan, [1166](#)  
Lumbar Pain, [1180](#)  
Lumbar Plexus, [1166](#)  
Lumbar Puncture, [1166](#)  
Lumbar Puncture Opening Pressure, [1166](#)  
Lumbar Radiculopathy, [1166](#)  
Lumbar Region, [1166](#)  
Lumbar Spinal Stenosis, [1166](#)  
Lumbar Spine, [1166](#)  
Lumbar Spine CT Scan, [1167](#)  
Lumbar Spondylosis, [1167](#)  
Lumbar Stenosis, [1167](#)  
Lumbar Vertebrae, [1167](#)  
Lumbosacral Radiculopathy, [1167](#)  
Lumbosacral Region, [1167](#)  
Lumbosacral Spine CT, [1167](#)  
Lumbosacral Spine X-Ray, [1167](#)  
Lumbricals, [1167](#)  
Lumen, [1167](#)  
Luminescence, [1167](#)  
Luminescence Type, [1167](#)  
Lump In The Abdomen, [1167](#)  
Lumpectomy, [1167](#)  
Lumpy Jaw, [1167](#)  
Lunate, [1168](#)  
Lunate Bone, [1168](#)  
Lung, [1168](#)  
Lung Abscess, [1168](#)  
Lung Anatomy, [1168](#)  
Lung Cancer, [1168](#)  
Lung Capacity, [1168](#)  
Lung Carcinoid Tumors, [1180](#)  
Lung Collapse, [1168](#)  
Lung Compliance, [1168](#)  
Lung Cyst, [1168](#)  
Lung Development, [1168](#)  
Lung Diffusion Testing, [1168](#)

---

---

Lung Disease, [1168](#)  
Lung Disease/Respiratory Problems, [1180](#)  
Lung Diseases, [1168](#)  
Lung Fibroma, [1168](#)  
Lung Fluke, [1169](#)  
Lung Function Tests, [1169](#)  
Lung Herniation, [1169](#)  
Lung Injury, [1169](#)  
Lung Lavage, [1169](#)  
Lung Metastases, [1169](#)  
Lung Needle Biopsy, [1169](#)  
Lung Neoplasm, [1169](#)  
Lung Nocardiosis, [1180](#)  
Lung Nodule, [1169](#)  
Lung PET Scan, [1169](#)  
Lung Plethysmography, [1169](#)  
Lung Problems And Volcanic Smog, [1169](#)  
Lung Protective Ventilation, [1169](#)  
Lung Resection, [1169](#)  
Lung Surgery, [1170](#)  
Lung Transplant, [1170](#)  
Lung Transplantation, [1170](#)  
Lung Volume Reduction Surgery, [1170](#)  
Lungs, [1170](#)  
Lungs Design And Purpose, [1170](#)  
Lunula, [1170](#)  
Lupus, [1170](#)  
Lupus Anticoagulant, [1170](#)  
Lupus Arthritis, [1112](#)  
Lupus Band Test, [1170](#)  
Lupus Erythematosus, [1170](#)  
Lupus Miliaris Disseminatus Faciei, [1171](#)  
Lupus Nephritis, [1171](#)  
Lupus Panniculitis, [1171](#)  
Lupus Pernio, [1171](#)  
Lupus Vulgaris, [1171](#)  
Luteal Cell, [1171](#)  
Luteal Phase Defect, [1171](#)  
Lutein, [1171](#)  
Luteinization, [1171](#)  
Luteinized Follicle Cyst, [1112](#)  
Luteinizing Hormone, [1171](#)  
Luteinizing Hormone Blood Test, [1171](#)  
Luteinizing Hormone Deficiency, [1171](#)  
Luteinizing Hormone-Releasing Hormone, [1171](#)  
Luteinizing Hormone-Releasing Hormone Agonist, [2124](#)  
Lutembacher Syndrome, [1171](#)  
Luteolysis, [1172](#)  
Luteoma, [1172](#)  
Luteoma Of Pregnancy, [1172](#)  
Luteotropic Hormone, [1172](#)  
Lutetium-177, [1172](#)  
Luxation, [1172](#)  
LVAD, [1172](#)  
LVNC, [1172](#)  
LVRS, [1172](#)  
Lyase, [1172](#)  
Lyme Arthritis, [1135](#)  
Lyme Disease, [1172](#)  
Lyme Disease Blood Test, [1172](#)  
Lyme Neuroborreliosis, [1172](#)  
Lymph, [1173](#)  
Lymph Leakage, [1175](#)  
Lymph Node, [1175](#)  
Lymph Node / Gland, [1175](#)  
Lymph Node Biopsy, [1175](#)  
Lymph Node Cancer, [1181](#)  
Lymph Node Culture, [1175](#)  
Lymph Node Dissection, [1175](#)  
Lymph Node Structure, [1175](#)  
Lymph Nodes, [1175](#)  
Lymph System, [1178](#)  
Lymphadenectomy, [1173](#)  
Lymphadenitis, [1173](#)  
Lymphadenopathy, [1173](#)  
Lymphangiectasia, [1173](#)  
Lymphangiectasis, [1173](#)  
Lymphangiogenesis, [1173](#)  
Lymphangiogram, [1173](#)  
Lymphangiography, [1173](#)  
Lymphangiomyomatosis, [1181](#)  
Lymphangiomyomatosis, [1173](#)  
Lymphangioma, [1173](#)  
Lymphangiomas, [1173](#)  
Lymphangiomyoma, [1173](#)  
Lymphangiosarcoma, [1174](#)  
Lymphangitis, [1174](#)  
Lymphatic Basin, [1174](#)  
Lymphatic Capillary, [1174](#)  
Lymphatic Circulation, [1174](#)  
Lymphatic Disease, [1174](#)  
Lymphatic Drainage, [1174](#)  
Lymphatic Filariasis, [1174](#)  
Lymphatic Malformation, [1174](#)  
Lymphatic Mapping, [1174](#)  
Lymphatic Obstruction, [1174](#)  
Lymphatic System, [1174](#)  
Lymphatic Tissue, [1174](#)  
Lymphatic Trunk, [1174](#)  
Lymphatic Valve, [1174](#)  
Lymphatic Vessel, [1174](#)  
Lymphedema, [1175](#)  
Lymphoblast, [1175](#)  
Lymphoblast Count, [1175](#)  
Lymphoblastic Lymphoma, [1175](#)  
Lymphocele, [1175](#)  
Lymphocyte, [1175](#)  
Lymphocyte Activation, [1176](#)

---

---

Lymphocyte Count, [1176](#)  
Lymphocyte Proliferation, [1176](#)  
Lymphocytic, [1176](#)  
Lymphocytic Choriomeningitis, [1176](#)  
Lymphocytic Colitis, [1176](#)  
Lymphocytic Interstitial Pneumonia, [1176](#)  
Lymphocytic Leukemia, [1176](#)  
Lymphocytic Mastitis, [1176](#)  
Lymphocytoma Cutis, [1176](#)  
Lymphocytopenia, [1176](#)  
Lymphocytosis, [1176](#)  
Lymphoedema, [1177](#)  
Lymphofollicular Hyperplasia, [1177](#)  
Lymphogranuloma Venereum, [1177](#)  
Lymphoid, [1177](#)  
Lymphoid Hyperplasia, [1177](#)  
Lymphoid Organs, [1177](#)  
Lymphoid Tissue, [1177](#)  
Lymphokine, [1177](#)  
Lymphoma, [1177](#)  
Lymphoma Overview, [1177](#)  
Lymphoma, Non-Hodgkin's, [1367](#)  
Lymphomatoid Granulomatosis, [1177](#)  
Lymphomatoid Papulosis, [1177](#)  
Lymphopenia, [1177](#)  
Lymphoplasmacytic Lymphoma, [1177](#)  
Lymphopoiesis, [1177](#)  
Lymphoproliferative Disorder, [1177](#)  
Lymphoscintigraphy, [1177](#)  
Lynch Syndrome, [1178](#)  
Lysergic Acid Diethylamide, [1178](#)  
Lysine, [1178](#)  
Lysis, [1178](#)  
Lysogenic Conversion, [1178](#)  
Lysogeny, [1178](#)  
Lysophagy, [1178](#)  
Lysophosphatidylcholine, [1178](#)  
Lysophospholipids, [1178](#)  
Lysosomal Storage Disorders, [1178](#)  
Lysosome, [1178](#)  
Lysozyme, [1178](#)  
Lysozyme Test, [1178](#)  
Lyssavirus, [1179](#)  
Lytic, [1179](#)  
Lytic Lesion, [1179](#)  
Löfgren Syndrome, [1135](#)  
  
MacConkey Agar, [1183](#)  
Macerate, [1183](#)  
Maceration, [1183](#)  
Macewen Sign, [1183](#)  
Machupo Virus, [1183](#)  
Macroamylasemia, [1183](#)  
Macroangiopathy, [1183](#)  
Macroautophagy, [1183](#)  
Macrocephalus, [1183](#)  
Macrocephaly, [1183](#)  
Macrocyte, [1183](#)  
Macrocytic, [1183](#)  
Macrocytic Anaemia, [1183](#)  
Macrocytic Anemia, [1184](#)  
Macrocytosis, [1184](#)  
Macroductyly, [1184](#)  
Macrodonia, [1184](#)  
Macrodonitosomia, [1184](#)  
Macroglobulin, [1184](#)  
Macroglossia, [1184](#)  
Macroglossia, [1184](#)  
Macrolide, [1184](#)  
Macrolide Antibiotic, [2124](#)  
Macrolides, [1184](#)  
Macronutrients, [1184](#)  
Macrophage, [1184](#)  
Macrophage Activation, [1184](#)  
Macrophage Activation Syndrome, [1184](#)  
Macrophage Count, [1185](#)  
Macropsia, [1185](#)  
Macroscopic, [1185](#)  
Macroscopic Hematuria, [1185](#)  
Macrosomia, [1185](#)  
Macrostomia, [1185](#)  
Macula, [1185](#)  
Macula Densa, [1185](#)  
Macula Lutea, [1185](#)  
Macular, [1185](#)  
Macular Amyloidosis, [1185](#)  
Macular Cyst, [1185](#)  
Macular Degeneration, [1185](#)  
Macular Drusen, [1186](#)  
Macular Edema, [1186](#)  
Macular Hole, [1186](#)  
Macular Pucker, [1186](#)  
Macular Skin Lesion, [1186](#)  
Macular Sparing, [1186](#)  
Macule, [1186](#)  
Macules, [1186](#)  
Maculopapular Lesion, [1186](#)  
Mad Cow Disease, [1186](#)  
Madarosis, [1186](#)  
Maddrey Discriminant Function, [1186](#)  
Madelung Deformity, [1186](#)  
Madura Foot, [1187](#)  
Maduramycosis, [1187](#)  
Madurella, [1187](#)  
Madurella Mycetomatis, [1187](#)  
Maffucci Syndrome, [1187](#)  
MAG3 Renal Scan, [1187](#)  
Magenblase Syndrome, [1187](#)

---

---

Magnesium, [1187](#)  
Magnesium Blood Test, [1187](#)  
Magnesium Carbonate, [1187](#)  
Magnesium Chloride, [1187](#)  
Magnesium Deficiency, [1187](#)  
Magnesium Hydroxide, [1187](#)  
Magnesium In Diet, [1187](#)  
Magnesium Oxide, [1187](#)  
Magnesium Sulfate, [1188](#)  
Magnesium Sulfate For Neuroprotection, [1188](#)  
Magnesium Sulphate, [1188](#)  
Magnetic Resonance Angiography, [1188](#)  
Magnetic Resonance Cholangiopancreatography, [1188](#)  
Magnetic Resonance Imaging, [1188](#)  
Magnetic Resonance Safety, [1188](#)  
Magnetic Resonance Spectroscopy, [1188](#)  
Magnetic Resonance Urography, [1188](#)  
Magnetic Resonance Venography, [1188](#)  
Magnetic Seed, [1188](#)  
Magnetic Therapy, [1188](#)  
Magnetocardiography, [1188](#)  
Magnetoencephalography, [1189](#)  
Magnusiomyces Capitatus, [1189](#)  
Mahvash Disease, [1189](#)  
Main-En-Griffe, [1189](#)  
Maintenance, [1189](#)  
Maintenance Dose, [1189](#)  
Maintenance Of Wakefulness Test, [1189](#)  
Maintenance Therapy, [1189](#)  
Maisonneuve Fracture, [1189](#)  
Maize, [1189](#)  
Majeed Syndrome, [1189](#)  
Majewski Syndrome, [1189](#)  
Majocchi Granuloma, [1189](#)  
Major, [1189](#)  
Major Depression, [1189](#)  
Major Depression With Psychotic Features, [1190](#)  
Major Depressive Disorder, [1190](#)  
Major Depressive Episode, [1190](#)  
Major Histocompatibility Complex, [1190](#)  
Major Histocompatibility Complex Molecule, [1190](#)  
Major Trauma, [1190](#)  
Malabsorption, [1190](#)  
Malabsorption Syndrome, [1190](#)  
Malacia, [1190](#)  
Malacoplakia, [1190](#)  
Maladaptation, [1190](#)  
Maladaptive Stress Response, [1190](#)  
Malaise, [1190](#)  
Malakoplakia, [1191](#)  
Malar, [1191](#)  
Malar Rash, [1191](#)  
Malaria, [1191](#)  
Malaria Vaccine, [2124](#)  
Malassezia, [1191](#)  
Malassezia (Pityrosporum) Folliculitis, [1191](#)  
Malassezia Furfur, [1191](#)  
Malassimilation, [1191](#)  
Malate Dehydrogenase, [1191](#)  
Malathion, [1191](#)  
Malathion Poisoning, [1191](#)  
MALDI-TOF Mass Spectrometry, [1191](#)  
Male, [1191](#)  
Male Breast Cancer, [1192](#)  
Male Breast Pathology, [1192](#)  
Male Chlamydial Infection, [1192](#)  
Male Condom, [1192](#)  
Male Genital Sores, [1192](#)  
Male Genitalia, [1192](#)  
Male Groin Pain, [1192](#)  
Male Hypoactive Sexual Desire Disorder, [1192](#)  
Male Hypogonadism, [1192](#)  
Male Infertility, [1192](#)  
Male Menopause, [1192](#)  
Male Pattern Baldness, [1192](#)  
Male Pelvis, [1192](#)  
Male Phenotype, [1193](#)  
Male Reproductive System, [1193](#)  
Male Self-Catheterization, [1193](#)  
Male Sexual Problems, [1321](#)  
Male Urethritis, [1193](#)  
Malformation, [1193](#)  
Malignancy, [1193](#)  
Malignancy-Associated Hypercoagulability, [1193](#)  
Malignant, [1193](#)  
Malignant Acanthosis Nigricans, [1193](#)  
Malignant Ameloblastoma, [1193](#)  
Malignant Appendical Tumor, [1321](#)  
Malignant Ascites, [1193](#)  
Malignant Bone Neoplasm, [1193](#)  
Malignant Carcinoid Syndrome, [1321](#)  
Malignant Fibrous Histiocytoma, [1193](#)  
Malignant Giant Cell Tumor Of Tendon Sheath, [1193](#)  
Malignant Glioma, [1194](#)  
Malignant Hypertension, [1194](#)  
Malignant Hyperthermia, [1194](#)  
Malignant Large Bowel Tumor, [1321](#)  
Malignant Mastocytosis, [1194](#)  
Malignant Melanoma, [1194](#)  
Malignant Mesothelioma, [1194](#)  
Malignant Myoepithelioma, [1194](#)  
Malignant Neoplasm, [1194](#)  
Malignant Otitis Externa, [1194](#)  
Malignant Peripheral Nerve Sheath Tumor, [1194](#)  
Malignant Pleural Effusion, [1194](#)  
Malignant Pustule, [1194](#)  
Malignant Syphilis, [1194](#)  
Malignant Teratoma Of The Mediastinum, [1195](#)

---

---

Malignant Thymoma, [1195](#)  
Malignant Tumor, [1195](#)  
Malignant Vulvar Lesions, [1195](#)  
Malingering, [1195](#)  
Mallampati Classification, [1195](#)  
Malleolus, [1195](#)  
Mallet, [1195](#)  
Mallet Finger, [1195](#)  
Mallet Toe, [1195](#)  
Mallet Toe and Hammertoe, [1195](#)  
Mallet Toes, [1321](#)  
Malleus, [1195](#)  
Mallory Bodies, [1195](#)  
Mallory Body, [1195](#)  
Mallory-Weiss Syndrome, [1196](#)  
Mallory-Weiss Tear, [1196](#)  
Malnutrition, [1196](#)  
Malocclusion, [1196](#)  
Malocclusion and Orthodontics, [1321](#)  
Malocclusion Of Teeth, [1196](#)  
Malondialdehyde, [1196](#)  
Malpighian Bodies, [1196](#)  
Malposition, [1196](#)  
Malposition of the Uterus, [1196](#)  
Malpresentation, [1196](#)  
Malrotation, [1196](#)  
MALS, [1196](#)  
MALT Lymphoma, [1196](#)  
Malt-Worker's Lung, [1197](#)  
Malta Fever, [1196](#)  
Maltase, [1196](#)  
Maltodextrin, [1196](#)  
Maltose, [1197](#)  
Maltreatment, Child, [379](#)  
Mammaglobin A, [1197](#)  
Mammaplasty, [1197](#)  
Mammary, [1197](#)  
Mammary Duct Ectasia, [1197](#)  
Mammary Epithelium, [1197](#)  
Mammary Gland, [1197](#)  
Mammary Glands, [1197](#)  
Mammary Implant, [1197](#)  
Mammary Paget Disease, [1197](#)  
Mammary Paget's Disease, [1321](#)  
Mamillary Bodies, [1197](#)  
Mamillary Body, [1198](#)  
Mammogram, [1198](#)  
Mammographic Calcifications, [1198](#)  
Mammography, [1198](#)  
Mammoplasty, [1198](#)  
Management of Third-Stage Labor, [1198](#)  
Mandatory Reporting, [1198](#)  
Mandible, [1198](#)  
Mandibular, [1198](#)  
Mandibular Advancement Device, [1198](#)  
Mandibular Ameloblastoma, [1321](#)  
Mandibular Condyle, [1198](#)  
Mandibular Fracture, [1198](#)  
Mandibular Prosthesis, [1198](#)  
Mandibulofacial Dysostosis, [1198](#)  
Manganese, [1199](#)  
Mania, [1199](#)  
Manic, [1199](#)  
Manic Depression in Children and Teens, [1321](#)  
Manic Disorder, [1199](#)  
Manic Episode, [1199](#)  
Manic Mood, [1199](#)  
Manic-Depression, [1199](#)  
Manic-Depressive Disorder, [1321](#)  
Manic-Depressive Illness, [1199](#)  
Manifest, [1199](#)  
Manifestation, [1199](#)  
Manipulation, [1199](#)  
Manipulation Procedure, [1199](#)  
Mannan, [1199](#)  
Mannheimia, [1199](#)  
Mannitol, [1199](#)  
Mannose, [1200](#)  
Mannose-Binding Lectin Deficiency, [1200](#)  
Mannosidosis, [1200](#)  
Manometer, [1200](#)  
Manometry, [1200](#)  
Mansonella Perstans, [1200](#)  
Mansonella Streptocerca, [1200](#)  
Mansonelliasis, [1200](#)  
Mantle Cell Lymphoma, [1200](#)  
Mantle Field, [1200](#)  
Mantoux Test, [1200](#)  
Manual, [1200](#)  
Manual Removal Of Placenta, [1200](#)  
Manual Therapy or Bodywork, [1323](#)  
Manual Vacuum Aspiration, [1200](#)  
Manubrium, [1201](#)  
Manus, [1201](#)  
MAO Inhibitor, [2124](#)  
MAOIs, [1201](#)  
Map-Dot-Fingerprint Type Corneal Dystrophy, [1201](#)  
Maple Syrup Urine Disease, [1201](#)  
Mapping, [1201](#)  
Marantic, [1201](#)  
Marasmus, [1201](#)  
Maraviroc, [1201](#)  
Marble Bone Disease, [1201](#)  
Marburg Virus, [1201](#)  
Marburgvirus, [1201](#)  
Marcus Gunn Jaw-Winking Phenomenon, [1202](#)  
Marcus Gunn Pupil, [1202](#)  
Marden-Walker Syndrome, [1202](#)

---

---

Marek's Disease, [1202](#)  
Marfan Syndrome, [1202](#)  
Marfan's Syndrome, [1202](#)  
Marfanoid Habitus, [1202](#)  
Marginal Ulcer, [1202](#)  
Marginal Zone Lymphoma, [1202](#)  
Maribavir, [1202](#)  
Marie-Strumpell Disease, [1202](#)  
Marijuana, [1202](#)  
Marijuana Intoxication, [1203](#)  
Marijuana Substance Use, [1203](#)  
Marijuana/Cannabis, [1203](#)  
Marine Animal Stings Or Bites, [1203](#)  
Marine Toxin, [1203](#)  
Marital Disharmony, [1203](#)  
Marital Status, [1203](#)  
Marital Therapy, [1203](#)  
Marjolin Ulcer, [1203](#)  
Marked, [1203](#)  
Marker, [1203](#)  
Maroteaux-Lamy Syndrome, [1203](#)  
Marrow, [1204](#)  
Marrow Hyperplasia, [1204](#)  
Marshall Syndrome, [1204](#)  
Marsupialization, [1204](#)  
Martin-Bell Syndrome, [1204](#)  
Martinus Beijerinck, [1202](#)  
MASA Syndrome, [1204](#)  
Masculinization, [1204](#)  
MASH, [1204](#)  
Mask, [1204](#)  
Masked Data, [1204](#)  
Masklike Face, [1204](#)  
MASLD, [1204](#)  
Masochism, [1204](#)  
Mason Fracture, [1204](#)  
Masoprocol, [1204](#)  
Mass, [1205](#)  
Mass Casualty Incident, [1205](#)  
Mass Screening, [1205](#)  
Mass Spectrometer, [1205](#)  
Mass Spectrometry, [1205](#)  
Massage, [1205](#)  
Massage Therapy, [1205](#)  
Masseter, [1205](#)  
Masseter Muscle, [1205](#)  
Massive Hemoptysis, [1205](#)  
Massive Ovarian Edema, [1205](#)  
Massive Pulmonary Embolism, [1205](#)  
Massive Pulmonary Embolus, [1205](#)  
Massive Transfusion Protocol, [1205](#)  
Mast Cell, [1206](#)  
Mast Cell Activation Syndrome, [1206](#)  
Mast Cell Sarcoma, [1206](#)  
Mast-Cell Sarcoma, [1206](#)  
Mastalgia, [1206](#)  
Mastectomy, [1206](#)  
Masticate, [1206](#)  
Mastication, [1206](#)  
Masticatory Muscle, [1206](#)  
Mastitis, [1206](#)  
Mastocytoma, [1206](#)  
Mastocytosis, [1206](#)  
Mastodynia, [1206](#)  
Mastoid, [1207](#)  
Mastoid Antrum, [1207](#)  
Mastoid Bone, [1207](#)  
Mastoid Process, [1207](#)  
Mastoidectomy, [1207](#)  
Mastoiditis, [1207](#)  
Mastopexy, [1207](#)  
Masturbation, [1207](#)  
Materia Medica, [1207](#)  
Maternal, [1207](#)  
Maternal Age, [1207](#)  
Maternal Expulsive Efforts, [1207](#)  
Maternal Health, [1207](#)  
Maternal Mortality, [1207](#)  
Maternal Mortality Rate, [1208](#)  
Maternal Obesity, [1208](#)  
Maternal Physiologic Changes In Pregnancy, [1208](#)  
Maternal Sepsis, [1208](#)  
Maternal Serum Alpha-Fetoprotein, [1208](#)  
Maternal Vaccine, [1208](#)  
Maternity, [1208](#)  
Maternity Services, [1208](#)  
Materteral, [1208](#)  
Mathematics Disorder, [1208](#)  
Mating, [1208](#)  
Matrix Band, [1208](#)  
Matrix Metalloproteinase, [1208](#)  
Maturation, [1208](#)  
Maturation Disorders Of Erythropoiesis, [1208](#)  
Mature Onset Diabetes, [1209](#)  
Mature Teratoma, [1209](#)  
Maxilla, [1209](#)  
Maxillary, [1209](#)  
Maxillary Sinus, [1209](#)  
Maxillary Sinusitis, [1209](#)  
Maxillofacial, [1209](#)  
Maxillofacial Prosthesis, [1209](#)  
Maximal Breathing Capacity, [1209](#)  
Maximum, [1209](#)  
Maximum Plasma Concentration, [1209](#)  
Maximum Thickness, [1209](#)  
Maximum Tolerated Dose, [1209](#)  
May-Hegglin Anomaly, [1210](#)  
May-Thurner Syndrome, [1210](#)

---



---

Mayer-Rokitansky-Kuster-Hauser Syndrome, [1210](#)  
Mayer-Rokitansky-Küster-Hauser Syndrome, [1210](#)  
Mayo Scissors, [1210](#)  
Maytansine, [1210](#)  
Maze Procedure, [1210](#)  
Mazindol, [1210](#)  
MB ChB, [1210](#)  
MC Lymphoma, [1211](#)  
MCA Peak Systolic Velocity, [1210](#)  
MCAD Deficiency, [1210](#)  
McArdle Disease, [1210](#)  
McBurney Point, [1211](#)  
McBurney Sign, [1211](#)  
McBurney's Point, [1211](#)  
Mcc Gene, [1211](#)  
McCune-Albright Syndrome, [1211](#)  
McDonald Cerclage, [1211](#)  
MCH, [1211](#)  
MCHC, [1211](#)  
MCL Injury, [1211](#)  
McMurray Test, [1211](#)  
McRoberts Maneuver, [1211](#)  
MCRPC, [1211](#)  
MCV, [1211](#)  
MDMA, [1212](#)  
MDMA Toxicity, [1212](#)  
MDS, [1212](#)  
ME/CFS, [1213](#)  
Mead Acid, [1212](#)  
Meadow Saffron, [1212](#)  
Mean, [1212](#)  
Mean Arterial Pressure, [1212](#)  
Mean Cell Volume, [1212](#)  
Mean Corpuscular Hemoglobin Concentration, [1212](#)  
Mean Corpuscular Volume, [1212](#)  
Mean Diameter, [1212](#)  
Mean Thickness, [1212](#)  
Measles, [1212](#)  
Measles Vaccine, [1214](#)  
Measles Virus, [1212](#)  
Measles-Rubella Vaccine, [1214](#)  
Measurable Residual Disease, [1213](#)  
Measurement, [1213](#)  
Meatal, [1213](#)  
Meatal Stenosis, [1213](#)  
Meatus, [1213](#)  
Meatuses, [1213](#)  
Mebendazole, [1213](#)  
Mebrofenin Scan, [1213](#)  
Mecamylamine, [1213](#)  
Mecarazole, [1213](#)  
Mechanical Circulatory Support, [1213](#)  
Mechanical Insufflation-Exsufflation, [1213](#)  
Mechanical Ventilation, [1213](#)  
Mechanical Ventilator, [1214](#)  
Mechanism Of Death, [1214](#)  
Mechanism Of Toxicity, [1214](#)  
Mechanoreceptor, [1214](#)  
Mechlorethamine, [1214](#)  
Meckel Diverticulectomy, [1214](#)  
Meckel Diverticulum, [1214](#)  
Meckel's Diverticulum, [1214](#)  
Meckel-Gruber Syndrome, [1214](#)  
Meclizine, [1214](#)  
Meclofenamate Overdose, [1214](#)  
Meconium, [1214](#)  
Meconium Aspiration, [1215](#)  
Meconium Aspiration Syndrome, [1215](#)  
Meconium Ileus, [1215](#)  
Meconium Peritonitis, [1215](#)  
Meconium-Stained Amniotic Fluid, [1215](#)  
MECP2 Duplication Syndrome, [1215](#)  
Medial, [1215](#)  
Medial Collateral Ligament Injury, [1215](#)  
Medial Cuneiform, [1215](#)  
Medial Epicondylitis, [1215](#)  
Medial Gastrocnemius Strain, [1215](#)  
Medial Geniculate Nucleus, [1215](#)  
Medial Lemniscus, [1215](#)  
Medial Medullary Syndrome, [1215](#)  
Medial Meniscus, [1216](#)  
Medial Rotation, [1216](#)  
Medial Tibial Stress Syndrome, [1216](#)  
Median, [1216](#)  
Median Arcuate Ligament Syndrome, [1216](#)  
Median Nerve, [1216](#)  
Median Nerve Compression, [1216](#)  
Median Neuropathy, [1216](#)  
Median Plane, [1216](#)  
Median Rhomboid Glossitis, [1216](#)  
Median Survival, [1216](#)  
Mediastinal Cyst, [1216](#)  
Mediastinal Disease, [1216](#)  
Mediastinal Emphysema, [1216](#)  
Mediastinal Hemorrhage, [1217](#)  
Mediastinal Lymph Nodes, [1217](#)  
Mediastinal Lymphadenopathy, [1217](#)  
Mediastinal Lymphoma, [1217](#)  
Mediastinal Mass, [1217](#)  
Mediastinal Neoplasm, [1217](#)  
Mediastinal Pleura, [1217](#)  
Mediastinal Shift, [1217](#)  
Mediastinal Tumor, [1217](#)  
Mediastinitis, [1217](#)  
Mediastinoscopes, [1217](#)  
Mediastinoscopy, [1217](#)  
Mediastinoscopy With Biopsy, [1217](#)  
Mediastinotomy, [1217](#)

---

---

Mediastinum, [1217](#)  
Mediators Of Inflammation, [1218](#)  
Medical Abbreviations, [1218](#)  
Medical Cannabis And Cannabidiol, [1218](#)  
Medical Condition, [1218](#)  
Medical Device, [1218](#)  
Medical Entomology, [1218](#)  
Medical Ethics, [1218](#)  
Medical Examiner, [1218](#)  
Medical History, [1218](#)  
Medical Illness, [1218](#)  
Medical Jargon, [1218](#)  
Medical Jurisprudence, [1218](#)  
Medical Malpractice, [1218](#)  
Medical Marijuana, [1218](#)  
Medical Nutrition Therapy, [1219](#)  
Medical Officer, [1219](#)  
Medical Physicist, [1219](#)  
Medical Physics, [1219](#)  
Medical Pin, [1219](#)  
Medical Practitioner, [1219](#)  
Medical Privacy, [1219](#)  
Medical Procedure, [1219](#)  
Medical Records, [1219](#)  
Medical Symbol, [1219](#)  
Medical Tattooing, [1219](#)  
Medical Tests, [1220](#)  
Medication, [1224](#)  
Medication Adherence, [1224](#)  
Medication Adverse-Effect Profile, [1224](#)  
Medication Cascade, [1224](#)  
Medication Deprescribing, [1224](#)  
Medication Error, [1224](#)  
Medication Management, [1225](#)  
Medication Overuse Headaches, [1225](#)  
Medication Reconciliation, [1225](#)  
Medication Review, [1225](#)  
Medication Safety, [1225](#)  
Medication Side Effects, [1225](#)  
Medicinal, [1220](#)  
Medicine, [1220](#)  
Medicines For Back Pain, [1220](#)  
Medicines For Osteoporosis, [1220](#)  
Medicines For Sleep, [1220](#)  
Medicolegal, [1220](#)  
Meditation, [1220](#)  
Mediterranean Anemia, [1220](#)  
Mediterranean Diet, [1220](#)  
Mediterranean Fever, [1220](#)  
Medium, [1220](#)  
MEDLARS, [1220](#)  
Medroxyprogesterone, [1220](#)  
Medroxyprogesterone Acetate, [1220](#)  
Medulla, [1221](#)  
Medulla Oblongata, [1221](#)  
Medullary, [1221](#)  
Medullary Carcinoma, [1221](#)  
Medullary Cavity, [1221](#)  
Medullary Sponge Kidney, [1221](#)  
Medullary Thyroid Carcinoma, [1221](#)  
Medulloblastoma, [1221](#)  
Medulloepithelioma, [1221](#)  
Mees' Lines, [1221](#)  
Mefenamic Acid, [1221](#)  
Mefloquine, [1221](#)  
Megabecquerel, [1221](#)  
Megacalycosis, [1221](#)  
Megacolon, [1222](#)  
Megakaryocyte, [1222](#)  
Megakaryocytopoiesis, [1222](#)  
Megaloblast, [1222](#)  
Megaloblastic Anaemia, [1222](#)  
Megaloblastic Anemia, [1222](#)  
Megalocephaly, [1222](#)  
Megamonas Funiformis, [1222](#)  
Megamonas Hypermegale, [1222](#)  
Megasphaera, [1222](#)  
Megaureter, [1222](#)  
Megestrol, [1222](#)  
Megestrol Acetate, [1222](#)  
Meglitinides, [1222](#)  
Megrim, [1222](#)  
Meibomian Gland Dysfunction, [1223](#)  
Meibomian Glands, [1223](#)  
Meibomianitis, [1223](#)  
Meige Syndrome, [1223](#)  
Meigs Syndrome, [1223](#)  
Meigs' Syndrome, [1223](#)  
Meiosis, [1223](#)  
Meiotic Recombination, [1223](#)  
Meissner Corpuscle, [1223](#)  
Meissner Plexus, [1223](#)  
Meissner's Corpuscles, [1223](#)  
MEK Inhibitor, [1223](#)  
Melancholia, [1223](#)  
Melancholic Features, [1223](#)  
Melanin, [1224](#)  
Melanoblast, [1224](#)  
Melanocyte, [1224](#)  
Melanocytes, [1224](#)  
Melanocytic CNS Tumor, [1224](#)  
Melanocytic Nevi, [1224](#)  
Melanocytic Nevus, [1224](#)  
Melanoderma, [1224](#)  
Melanoma, [1224](#)  
Melanoma Of The Eye, [1224](#)  
Melanoma Skin Cancer, [1224](#)  
Melanoma Surveillance, [1224](#)

---

---

Melanoma Treatment, [1224](#)  
Melanoma/Skin Cancer, [1324](#)  
Melanonychia, [1225](#)  
Melanophore, [1225](#)  
Melanosis, [1225](#)  
Melanosis Coli, [1225](#)  
Melanosome, [1225](#)  
MELAS Syndrome, [1225](#)  
Melasma, [1225](#)  
Melatonin, [1225](#)  
Melatonin Receptor, [1225](#)  
Melatonin Receptor Agonists, [1225](#)  
MELD, [1225](#)  
MELD Score, [1226](#)  
MELD-Na Score, [1225](#)  
Melena, [1226](#)  
Melibiose, [1226](#)  
Melio, [1226](#)  
Meliodosis, [1226](#)  
Melitensis, [1226](#)  
Melitten, [1226](#)  
Melkersson-Rosenthal Syndrome, [1226](#)  
Melorheostosis, [1226](#)  
Meloxicam, [1226](#)  
Melphalan, [1226](#)  
Memantine, [1226](#)  
Memantine Hydrochloride, [1226](#)  
Membrane, [1226](#)  
Membrane Channel, [1227](#)  
Membrane Fusion, [1227](#)  
Membrane Lipids, [1227](#)  
Membrane Microdomain, [1227](#)  
Membrane Part, [1227](#)  
Membrane Potential, [1227](#)  
Membrane Protein, [1227](#)  
Membrane Raft, [1227](#)  
Membranoproliferative Glomerulonephritis, [1227](#)  
Membranous, [1227](#)  
Membranous Labyrinth, [1227](#)  
Membranous Nephropathy, [1227](#)  
Memory, [1227](#)  
Memory Cell, [1228](#)  
Memory Disorder, [1228](#)  
Memory Impairment, [1228](#)  
Memory Loss, [1228](#)  
Memory Span, [1228](#)  
Memory T Cells And B Cells, [1228](#)  
MEN 2, [1228](#)  
Menarche, [1228](#)  
Mendelian Inheritance, [1228](#)  
Mendelism, [1228](#)  
Mendelson Syndrome, [1228](#)  
Menetrier Disease, [1228](#)  
Meniere Disease, [1228](#)  
Meniere's Disease, [1229](#)  
Meningeal Carcinomatosis, [1229](#)  
Meningeal Leukemia, [1229](#)  
Meningeal Lymphoma, [1229](#)  
Meningeal Neoplasm, [1229](#)  
Meninges, [1229](#)  
Meningioma, [1229](#)  
Meningioma Pathology, [1229](#)  
Meningism, [1229](#)  
Meningismus, [1229](#)  
Meningitis, [1229](#)  
Meningocele, [1229](#)  
Meningocele Repair, [1229](#)  
Meningococcal Disease, [1230](#)  
Meningococcal Infection, [1230](#)  
Meningococcal Meningitis, [1230](#)  
Meningococcal Vaccine, [1225](#)  
Meningococcemia, [1230](#)  
Meningococcus, [1230](#)  
Meningoencephalitis, [1230](#)  
Meningomyelocele, [1230](#)  
Meniscal Allograft Transplantation, [1230](#)  
Meniscal Injuries, [1230](#)  
Meniscal Tear, [1230](#)  
Meniscectomy, [1230](#)  
Meniscus, [1230](#)  
Meniscus Tear, [1230](#)  
Menkes Disease, [1230](#)  
Menkes Kinky Hair Syndrome, [1231](#)  
Menogaril, [1231](#)  
Menometrorrhagia, [1231](#)  
Menopausal, [1231](#)  
Menopausal Hormone Therapy, [1231](#)  
Menopausal Hot Flashes, [1231](#)  
Menopause, [1231](#)  
Menopause and Mood Disorders, [1231](#)  
Menopause And Perimenopause, [1231](#)  
Menopause Status, [1231](#)  
Menopause Syndrome, [1231](#)  
Menorrhagia, [1231](#)  
Menotropin, [1231](#)  
Menses Duration, [1231](#)  
Menstrual, [1232](#)  
Menstrual Cramps, [1232](#)  
Menstrual Cycle, [1232](#)  
Menstrual Disorders, [1232](#)  
Menstruation, [1232](#)  
Menstruation Disturbance, [1232](#)  
Mental, [1232](#)  
Mental Age, [1232](#)  
Mental Disorder, [1232](#)  
Mental Disorders, [1232](#)  
Mental Disorders In Children, [1232](#)  
Mental Fatigue, [1232](#)

---

---

Mental Health, [1232](#)  
 Mental Health Crisis, [1233](#)  
 Mental Illness, [1233](#)  
 Mental Illness In Children, [1233](#)  
 Mental Process, [1233](#)  
 Mental Recall, [1233](#)  
 Mental Status Examination, [1233](#)  
 Mental Status Testing, [1233](#)  
 Mentation, [1233](#)  
 Menthol, [1233](#)  
 Menthol Poisoning, [1233](#)  
 Mepacrine, [1233](#)  
 Meperidine, [1233](#)  
 Meperidine Hydrochloride, [1233](#)  
 Meperidine Hydrochloride Overdose, [1233](#)  
 Mephenesin, [1233](#)  
 Mephentermine, [1233](#)  
 Mephenytoin, [1234](#)  
 Mepivacaine, [1234](#)  
 Meprobamate, [1234](#)  
 Meprobamate Overdose, [1234](#)  
 Mepyramine, [1234](#)  
 Meralgia Paresthetica, [1234](#)  
 Merbromin Poisoning, [1234](#)  
 Mercaptopurine, [1234](#)  
 Mercurialism, [1234](#)  
 Mercuric Chloride Poisoning, [1234](#)  
 Mercuric Oxide Poisoning, [1234](#)  
 Mercury, [1234](#)  
 Mercury Poisoning, [1234](#)  
 Meridian, [1235](#)  
 Merkel Cell Carcinoma, [1235](#)  
 Merkel Cells, [1235](#)  
 Mermaid Syndrome, [1235](#)  
 Meropenem, [1235](#)  
 Meropenem-Vaborbactam, [1235](#)  
 MERRF Syndrome, [1235](#)  
 MERS-CoV, [1235](#)  
 Merthiolate Poisoning, [1235](#)  
 Merycism, [1235](#)  
 Mesalamine, [1235](#)  
 Mesangial Cell, [1235](#)  
 Mesangiolytic, [1235](#)  
 Mesangium, [1235](#)  
 Mescaline, [1236](#)  
 Mesencephalon, [1236](#)  
 Mesenchymal Cell, [1236](#)  
 Mesenchymal Chondrosarcoma, [1236](#)  
 Mesenchymal CNS Tumor, [1236](#)  
 Mesenchyme, [1236](#)  
 Mesenchymoma, [1236](#)  
 Mesenteric, [1236](#)  
 Mesenteric Angiography, [1236](#)  
 Mesenteric Artery, [1236](#)  
 Mesenteric Artery Aneurysm, [1236](#)  
 Mesenteric Artery Ischemia, [1236](#)  
 Mesenteric Cyst, [1236](#)  
 Mesenteric Ischemia, [1236](#)  
 Mesenteric Lymph Nodes, [1237](#)  
 Mesenteric Lymphadenitis, [1237](#)  
 Mesenteric Veins, [1237](#)  
 Mesenteric Venous Thrombosis, [1237](#)  
 Mesentery, [1237](#)  
 Mesh, [1237](#)  
 Mesial, [1237](#)  
 Mesial Surface, [1237](#)  
 Mesna, [1237](#)  
 Mesoderm, [1237](#)  
 Mesometrium, [1237](#)  
 Mesonephric, [1237](#)  
 Mesonephric Adenocarcinoma, [1237](#)  
 Mesonephric Tumor, [1238](#)  
 Mesothelial Neoplasm, [1238](#)  
 Mesothelin, [1238](#)  
 Mesothelioma, [1238](#)  
 Mesothelium, [1238](#)  
 Messenger RNA, [1238](#)  
 Met, [1238](#)  
 Metabolic, [1238](#)  
 Metabolic Acidosis, [1238](#)  
 Metabolic Alkalosis, [1238](#)  
 Metabolic Bone Disease, [1238](#)  
 Metabolic Bone Disease of Prematurity, [1238](#)  
 Metabolic Disease, [1238](#)  
 Metabolic Disorders, [1239](#)  
 Metabolic Dysfunction-Associated Steatohepatitis, [1239](#)  
 Metabolic Dysfunction-Associated Steatohepatitis (MASH), [1321](#)  
 Metabolic Dysfunction-Associated Steatotic Liver Disease, [1239](#)  
 Metabolic Equivalent of Task, [1239](#)  
 Metabolic Equivalents, [1239](#)  
 Metabolic Myopathies, [1239](#)  
 Metabolic Neuropathies, [1239](#)  
 Metabolic Rate, [1239](#)  
 Metabolic Response, [1239](#)  
 Metabolic Syndrome, [1239](#)  
 Metabolic Tumor Volume, [1239](#)  
 Metabolism, [1239](#)  
 Metabolite, [1239](#)  
 Metacarpal, [1239](#)  
 Metacarpal Bone, [1240](#)  
 Metacarpals, [1240](#)  
 Metacarpophalangeal Joint, [1240](#)  
 Metacarpophalangeal Joint Dislocation, [1240](#)  
 Metacarpus, [1240](#)  
 Metachromatic Leukodystrophy, [1240](#)  
 Metagonimus Yokogawai, [1240](#)

---

---

Metal Artifact, [1240](#)  
 Metal Fume Fever, [1240](#)  
 Metalloendopeptidase, [1240](#)  
 Metalloporphyrins, [1240](#)  
 Metalloprotease, [1240](#)  
 Metalloprotein, [1240](#)  
 Metallothionein, [1240](#)  
 Metanephrine, [1241](#)  
 Metaphase, [1241](#)  
 Metaphase Plate, [1241](#)  
 Metaphyseal Corner Fractures, [1241](#)  
 Metaphyseal Widening, [1241](#)  
 Metaphysis, [1241](#)  
 Metaplasia, [1241](#)  
 Metaplastic Carcinoma, [1241](#)  
 Metapneumovirus, [1241](#)  
 Metastasectomy, [1241](#)  
 Metastases To The Breast, [1241](#)  
 Metastasis, [1241](#)  
 Metastasis Of Colorectal Cancer, [1241](#)  
 Metastasis-Free Survival, [1241](#)  
 Metastatic Adenocarcinoma, [1242](#)  
 Metastatic Brain Tumor, [1242](#)  
 Metastatic Breast Cancer, [1242](#)  
 Metastatic Calcification, [1242](#)  
 Metastatic Cancer, [1242](#)  
 Metastatic Cancer With Unknown Primary Site, [1242](#)  
 Metastatic Castration-Resistant Prostate Cancer, [1242](#)  
 Metastatic Colon Cancer, [1242](#)  
 Metastatic Colorectal Cancer, [1242](#)  
 Metastatic Inguinal Lymphadenopathy, [1242](#)  
 Metastatic Lung Cancer, [1242](#)  
 Metastatic Melanoma, [1321](#)  
 Metastatic Pleural Tumor, [1242](#)  
 Metastin, [1242](#)  
 Metatarsal, [1243](#)  
 Metatarsal Bone, [1243](#)  
 Metatarsal Valgus, [1243](#)  
 Metatarsalgia, [1243](#)  
 Metatarsals, [1243](#)  
 Metatarsophalangeal Joint, [1243](#)  
 Metatarsus, [1243](#)  
 Metatarsus Adductus, [1243](#)  
 Metatarsus Varus, [1243](#)  
 Metencephalon, [1243](#)  
 Meteorism, [1243](#)  
 Meter, [1243](#)  
 Metered-Dose Inhaler, [1243](#)  
 Metestrus, [1243](#)  
 Metformin, [1243](#)  
 Metformin Hydrochloride, [1243](#)  
 Metformin In Prediabetes, [1244](#)  
 Methacholine Challenge Test, [1244](#)  
 Methacholine Chloride, [1244](#)  
 Methadone, [1244](#)  
 Methadone Hydrochloride, [1244](#)  
 Methadone Overdose, [1244](#)  
 Methaemoglobinaemia, [1244](#)  
 Methamidophos, [1244](#)  
 Methamphetamine, [1244](#)  
 Methamphetamine Overdose, [1244](#)  
 Methamphetamine Toxicity, [1244](#)  
 Methamphetamines, [1244](#)  
 Methane, [1245](#)  
 Methane Sulfonate, [1245](#)  
 Methanobrevibacter Smithii, [1245](#)  
 Methanol, [1245](#)  
 Methanol Poisoning, [1245](#)  
 Methanol Test, [1245](#)  
 Methazolamide, [1245](#)  
 Methemoglobin, [1245](#)  
 Methemoglobinemia, [1245](#)  
 Methenamine, [1245](#)  
 Methicillin-Resistant Staphylococcus Aureus, [1245](#)  
 Methicillin-Resistant Staphylococcus Aureus Infection, [1245](#)  
 Methidathion, [1245](#)  
 Methimazole, [1245](#)  
 Methiocarb, [1246](#)  
 Methionine, [1246](#)  
 Methionine Enkephalin, [1246](#)  
 Methocarbamol, [1246](#)  
 Methohexital, [1246](#)  
 Methomyl, [1246](#)  
 Methotrexate, [1246](#)  
 Methotrexate In Dermatology, [1246](#)  
 Methotrimeprazine, [1246](#)  
 Methoxamine, [1246](#)  
 Methoxsalen, [1246](#)  
 Methoxyflurane, [1246](#)  
 Methyclothiazide, [1246](#)  
 Methyl Alcohol, [1247](#)  
 Methyl Bromide, [1247](#)  
 Methyl Chloride, [1247](#)  
 Methyl Isocyanate, [1247](#)  
 Methyl Parathion, [1248](#)  
 Methyl Salicylate Overdose, [1248](#)  
 Methylation, [1247](#)  
 Methylcellulose, [1247](#)  
 Methyldopa, [1247](#)  
 Methylene Blue, [1247](#)  
 Methylene Blue Test, [1247](#)  
 Methylene Chloride, [1247](#)  
 Methylenedioxymethamphetamine, [1247](#)  
 Methylergometrine, [1247](#)  
 Methylmalonic Acid Blood Test, [1247](#)  
 Methylmalonic Acid Level, [1248](#)  
 Methylmalonic Acidemia, [1247](#)

---

---

Methylmalonic Aciduria, [1248](#)  
Methylmercury Compounds, [1248](#)  
Methylmercury Poisoning, [1248](#)  
Methylmethacrylate, [1248](#)  
Methylmorphine, [1248](#)  
Methylnaltrexone, [1248](#)  
Methylnaltrexone Bromide, [1248](#)  
Methylphenidate, [1248](#)  
Methylprednisolone, [1248](#)  
Methyltestosterone, [1248](#)  
Metoclopramide, [1249](#)  
Metolazone, [1249](#)  
Metolcarb, [1249](#)  
Metopic Ridge, [1249](#)  
Metoprolol, [1249](#)  
Metric System, [1249](#)  
Metritis, [1249](#)  
Metronidazole, [1249](#)  
Metronidazole-Associated Encephalopathy, [1249](#)  
Metronomic Chemotherapy, [1249](#)  
Metrorrhagia, [1249](#)  
METs, [1249](#)  
Metyrapone, [1250](#)  
Metzenbaum Scissors, [1250](#)  
Mevalonate Kinase Deficiency, [1250](#)  
Mevinphos, [1250](#)  
Mexacarbate, [1250](#)  
Mexiletine, [1250](#)  
Meyerozyma Guilliermondii, [1250](#)  
MF, [1250](#)  
MFS, [1250](#)  
MG, [1250](#)  
MGUS, [1250](#)  
MHC Molecule, [1250](#)  
Mianserin, [1250](#)  
MIBG Scan, [1250](#)  
MIBG Scintiscan, [1250](#)  
Mica, [1251](#)  
Micafungin, [1251](#)  
Micelle, [1251](#)  
Michaelis-Menten Kinetics, [1251](#)  
Miconazole, [1251](#)  
Micro-Surgery, [1257](#)  
Microabscess, [1251](#)  
Microalbuminuria, [1251](#)  
Microalbuminuria Test, [1251](#)  
Microaneurysm, [1251](#)  
Microangiopathic Hemolysis, [1251](#)  
Microangiopathic Hemolytic Anemia, [1251](#)  
Microangiopathy, [1251](#)  
Microarray, [1251](#)  
Microarray Analysis, [1252](#)  
Microbacterium, [1252](#)  
Microbe, [1252](#)  
Microbial Biofilms, [1252](#)  
Microbicide, [1252](#)  
Microbiology, [1252](#)  
Microbiome, [1252](#)  
Microcalcification, [1252](#)  
Microcephaly, [1252](#)  
Microcirculation, [1252](#)  
Microcolon, [1252](#)  
Microcornea, [1252](#)  
Microcurie, [1252](#)  
Microcystic Adnexal Carcinoma, [1252](#)  
Microcytic Anemia, [1253](#)  
Microdermabrasion, [1253](#)  
Microdialysis, [1253](#)  
Microdissectomy, [1253](#)  
Microdissection, [1253](#)  
Microdissection Testicular Sperm Extraction, [1253](#)  
Microdontia, [1253](#)  
Microfibril, [1253](#)  
Microfilament, [1253](#)  
Microfilament Proteins, [1253](#)  
Microfilaria, [1253](#)  
Microglandular Adenosis, [1253](#)  
Microglia, [1254](#)  
Microgliosis, [1254](#)  
Micrognathia, [1254](#)  
Micrognathism, [1254](#)  
Micrograms, [1254](#)  
Micrographia, [1254](#)  
Microinjection, [1254](#)  
Micrometastasis, [1254](#)  
Micrometer, [1254](#)  
Micromolar, [1254](#)  
Micromole, [1254](#)  
Micron, [1254](#)  
Microneme, [1254](#)  
Micronization, [1254](#)  
Micronized, [1255](#)  
Micronutrient, [1255](#)  
Micronutrients, [1255](#)  
Microorchidism, [1255](#)  
Microorganism, [1255](#)  
Microorganism Genetics, [1255](#)  
Micropenis, [1255](#)  
Microphallus, [1255](#)  
Microphthalmia, [1255](#)  
Microphthalmos, [1255](#)  
Microplastics, [1255](#)  
Micropolygyria with Muscular Dystrophy, [1321](#)  
Micropsia, [1255](#)  
MicroRNA, [1255](#)  
Microsatellite Instability, [1255](#)  
Microscope, [1256](#)  
Microscopic, [1256](#)

---



---

Microscopic Anatomy, [1256](#)  
Microscopic Colitis, [1256](#)  
Microscopic Examination, [1256](#)  
Microscopic Hematuria, [1256](#)  
Microscopic Polyangiitis, [1256](#)  
Microscopy, [1256](#)  
Microsecond, [1256](#)  
Microspectrophotometry, [1256](#)  
Microsphere, [1256](#)  
Microspherophakia, [1256](#)  
Microsporidia, [1256](#)  
Microsporidiosis, [1256](#)  
Microsporum, [1256](#)  
Microsporum Audouinii, [1257](#)  
Microsporum Distortum, [1257](#)  
Microsporum Ferrugineum, [1257](#)  
Microsporum Gypseum, [1257](#)  
Microstomia, [1257](#)  
Microtechnology, [1257](#)  
Microtia, [1257](#)  
Microtome, [1257](#)  
Microtubule, [1257](#)  
Microtubule Bundle, [1257](#)  
Microvascular, [1257](#)  
Microvascular Disease, [1257](#)  
Microvasculature, [1257](#)  
Microvilli, [1257](#)  
Microvillus, [1258](#)  
Microvolt, [1258](#)  
Microwave Ablation, [1258](#)  
Microwave Therapy, [1258](#)  
Micturition, [1258](#)  
Mid-Dorsal, [1259](#)  
Mid-Forearm, [1259](#)  
Mid-Gut, [1259](#)  
Mid-Leg, [1259](#)  
Mid-Line, [1259](#)  
Mid-Stream Urine Specimen, [1259](#)  
Mid-Systolic Click, [1259](#)  
Mid-Upper-Arm Circumference, [1260](#)  
Midaxillary Line, [1258](#)  
Midazolam, [1258](#)  
Midazolam Hydrochloride, [1258](#)  
Midbrain, [1258](#)  
Midclavicular Line, [1258](#)  
Middle Ear, [1258](#)  
Middle Ear Implant, [1258](#)  
Middle Ear Infection, [1258](#)  
Middle East Respiratory Syndrome, [1258](#)  
Middle Insomnia, [1259](#)  
Middle Lobe Syndrome, [1259](#)  
Middle Mediastinum, [1259](#)  
Midgut Volvulus, [1259](#)  
Midomafetamine, [1259](#)  
Midostaurin, [1259](#)  
Midriff, [1259](#)  
Midtrimester, [1259](#)  
Midwife, [1260](#)  
Midwifery, [1260](#)  
Mifepristone, [1260](#)  
Migraine, [1260](#)  
Migraine And Stroke Symptoms, [1260](#)  
Migraine Headaches, [1322](#)  
Migraine With Aura, [1260](#)  
Migraine Without Aura, [1260](#)  
Migrainous Neuralgia, [1260](#)  
Migrating Motor Complex, [1260](#)  
Migration, [1260](#)  
Migratory Thrombophlebitis, [1260](#)  
Milan System for Salivary Cytopathology, [1260](#)  
Mild, [1261](#)  
Mild Cognitive Impairment, [1261](#)  
Mildew Remover Poisoning, [1261](#)  
Mile, [1261](#)  
Milia, [1261](#)  
Miliaria, [1261](#)  
Miliaria Crystallina, [1261](#)  
Miliaria Profunda, [1261](#)  
Miliaria Rubra, [1261](#)  
Miliary, [1261](#)  
Miliary TB, [1261](#)  
Miliary Tuberculosis, [1261](#)  
Milieu Therapy, [1261](#)  
Miliun, [1261](#)  
Milk Allergy, [1322](#)  
Milk Intolerance, [1322](#)  
Milk Of Magnesia, [1262](#)  
Milk-Alkali Syndrome, [1261](#)  
Milker's Node, [1262](#)  
Milkman Syndrome, [1262](#)  
Milky Urine, [1262](#)  
Millard-Gubler Syndrome, [1262](#)  
Miller Fisher Syndrome, [1262](#)  
Miller's Lung, [1262](#)  
Millicurie, [1262](#)  
Milliequivalent, [1262](#)  
Milligram, [1262](#)  
Milligram-Percent, [1262](#)  
Milligray, [1262](#)  
Milliliter, [1262](#)  
Millimeter, [1262](#)  
Millimolar, [1263](#)  
Millimole, [1263](#)  
Million, [1263](#)  
Milliosmole, [1263](#)  
Millipede Toxin, [1263](#)  
Millisecond, [1263](#)  
Milliunit, [1263](#)

---

---

Millivolt, [1263](#)  
 Milrinone, [1263](#)  
 Milroy Disease, [1263](#)  
 Milroy's Disease, [1263](#)  
 Mimesis, [1263](#)  
 Mimicry, [1263](#)  
 Minamata Disease, [1263](#)  
 Mind, [1263](#)  
 Mindful Eating, [1264](#)  
 Miner's Asthma, [1264](#)  
 Miner's Nystagmus, [1264](#)  
 Mineral, [1264](#)  
 Mineral Oil, [1264](#)  
 Mineral Oil Overdose, [1264](#)  
 Mineral Spirits Poisoning, [1264](#)  
 Mineralocorticoid, [1264](#)  
 Mineralocorticoid Receptor, [1264](#)  
 Mineralocorticoid Receptor Antagonist, [2125](#)  
 Mineralocorticoids, [1264](#)  
 Minerals, [1264](#)  
 Mini-Mental State Examination, [1265](#)  
 Mini-Stroke, [1266](#)  
 Mini-Stroke (TIA), [1322](#)  
 Minimal Change Disease, [1264](#)  
 Minimal Residual Disease, [1265](#)  
 Minimal-Change Disease, [1264](#)  
 Minimally Invasive Aortic Valve Surgery, [1264](#)  
 Minimally Invasive Esophagectomy, [1265](#)  
 Minimally Invasive Gastrointestinal Surgery, [1265](#)  
 Minimally Invasive Heart Bypass Surgery, [1265](#)  
 Minimally Invasive Hip Replacement, [1265](#)  
 Minimally Invasive Mitral Valve Surgery, [1265](#)  
 Minimally Invasive Prostate Resection, [1265](#)  
 Minimally Invasive Spine Surgery, [1265](#)  
 Minimally Invasive Surgery, [1265](#)  
 Minimization, [1265](#)  
 Minimum Alveolar Concentration, [1265](#)  
 Minimum Bactericidal Concentration, [1266](#)  
 Minimum Effect Concentration, [1266](#)  
 Minimum Inhibitory Concentration, [1266](#)  
 Minimum Lethal Concentration, [1266](#)  
 Minimum Lethal Dose, [1266](#)  
 Minisatellite Repeat, [1266](#)  
 MINOCA, [1266](#)  
 Minocycline, [1266](#)  
 Minor, [1266](#)  
 Minor Response, [1266](#)  
 Minority Health, [1266](#)  
 Minoxidil, [1266](#)  
 Minute Ventilation, [1266](#)  
 Miosis, [1267](#)  
 Miosis Disorder, [1267](#)  
 Miotic, [1267](#)  
 Miracidium, [1267](#)  
 Mirage Syndrome, [1267](#)  
 MIRD Schema, [1267](#)  
 Mirizzi Syndrome, [1267](#)  
 Mirror Therapy, [1267](#)  
 Mirror-Writing, [1267](#)  
 Mirtazapine, [1267](#)  
 Misalignment, [1267](#)  
 Miscarriage, [1267](#)  
 Misdiagnosis, [1267](#)  
 Mismatch Repair, [1267](#)  
 Mismatch Repair Deficiency, [1268](#)  
 MISME, [1268](#)  
 Misophonia, [1268](#)  
 Misoprostol, [1268](#)  
 Missed Dose, [1268](#)  
 Missed Miscarriage, [1268](#)  
 Missed Period, [1322](#)  
 Missense Mutation, [1268](#)  
 Missense Variant, [1268](#)  
 Mistletoe Poisoning, [1268](#)  
 Mitchell Disease, [1322](#)  
 Mite, [1268](#)  
 Mites, [1268](#)  
 Mithramycin, [1268](#)  
 Mitobronitol, [1268](#)  
 Mitochondria, [1269](#)  
 Mitochondrial Cristae, [1269](#)  
 Mitochondrial Diabetes, [1269](#)  
 Mitochondrial Disease, [1269](#)  
 Mitochondrial DNA, [1269](#)  
 Mitochondrial DNA Depletion Syndrome MDS MDDS, [1269](#)  
 Mitochondrial Fission, [1269](#)  
 Mitochondrial Fusion, [1269](#)  
 Mitochondrial Inheritance, [1269](#)  
 Mitochondrial Matrix, [1269](#)  
 Mitochondrial Membrane, [1269](#)  
 Mitochondrial Myopathy, [1269](#)  
 Mitochondrial Ribosome, [1269](#)  
 Mitochondrial Swelling, [1270](#)  
 Mitochondrion, [1270](#)  
 Mitogen, [1270](#)  
 Mitolactol, [1270](#)  
 Mitomycin, [1270](#)  
 Mitomycins, [1270](#)  
 Mitophagy, [1270](#)  
 Mitosis, [1270](#)  
 Mitosporic Fungi, [1270](#)  
 Mitotane, [1270](#)  
 Mitotic Anaphase, [1270](#)  
 Mitotic Figure, [1270](#)  
 Mitotic Metaphase, [1270](#)  
 Mitotic Prophase, [1270](#)  
 Mitotic Recombination, [1271](#)

---

---

Mitotic Telophase, [1271](#)  
 Mitoxantrone, [1271](#)  
 MitraClip, [1271](#)  
 Mitral, [1271](#)  
 Mitral Annular Calcification, [1271](#)  
 Mitral Atresia, [1271](#)  
 Mitral Facies, [1271](#)  
 Mitral Insufficiency, [1271](#)  
 Mitral Regurgitation, [1271](#)  
 Mitral Stenosis, [1271](#)  
 Mitral Stenosis Murmur, [1271](#)  
 Mitral Valve, [1271](#)  
 Mitral Valve Insufficiency, [1272](#)  
 Mitral Valve Prolapse, [1272](#)  
 Mitral Valve Regurgitation, [1272](#)  
 Mitral Valve Stenosis, [1272](#)  
 Mitral Valvotomy, [1272](#)  
 Mitral Valvuloplasty, [1272](#)  
 Mitral-Valve Repair, [1272](#)  
 Mitsukella, [1272](#)  
 Mittelschmerz, [1272](#)  
 Mixed Aortic Valve Disease, [1272](#)  
 Mixed Connective Tissue Disease, [1272](#)  
 Mixed Connective Tissue Disease (MCTD), [1322](#)  
 Mixed Connective Tissue Disease-Associated Arthritis, [1272](#)  
 Mixed Hearing Loss, [1273](#)  
 Mixed Hepatic Porphyria, [1322](#)  
 Mixed Mitral Valve Disease, [1273](#)  
 Mixed Oedema, [1273](#)  
 Mixed Urinary Incontinence, [1273](#)  
 Mixed-Lineage Leukemia, [1273](#)  
 Miyoshi Myopathy, [1273](#)  
 MME, [1273](#)  
 MMR Vaccine, [2125](#)  
 Mobile, [1273](#)  
 Mobile Genetic Elements, [1273](#)  
 Mobile Health, [1273](#)  
 Mobile Health Unit, [1274](#)  
 Mobility, [1274](#)  
 Mobilization, [1274](#)  
 Mobiluncus, [1274](#)  
 Mobiluncus Curtisii, [1274](#)  
 Mobiluncus Mulieris, [1274](#)  
 Mobitz Type I, [1274](#)  
 Mobitz Type I Block, [1274](#)  
 Mobitz Type II, [1274](#)  
 Mobitz Type II Block, [1274](#)  
 Mobius Syndrome, [1274](#)  
 Modafinil, [1275](#)  
 Modality, [1275](#)  
 Mode, [1275](#)  
 Model, [1275](#)  
 Moderate To Severe Disabilities, [1275](#)  
 Moderation, [1275](#)  
 Moderator, [1275](#)  
 Modes Of Inheritance, [1275](#)  
 Modified Biophysical Profile, [1275](#)  
 Modulator, [1275](#)  
 Modus Operandi, [1275](#)  
 Modus Vivendi, [1275](#)  
 Moebius Syndrome, [1275](#)  
 Moeller's Glossitis, [1276](#)  
 Moellerella, [1275](#)  
 Moellerella Wisconsensis, [1275](#)  
 MOG Antibody-Associated Disorder, [1276](#)  
 MOGAD, [1276](#)  
 Mogibacterium, [1276](#)  
 Mohs Micrographic Surgery, [1276](#)  
 Mohs Surgery, [1276](#)  
 Moisture, [1276](#)  
 Molar, [1276](#)  
 Molar Pregnancy, [1276](#)  
 Molasses, [1276](#)  
 Mold, [1276](#)  
 Mold Allergy, [1276](#)  
 Mold Exposure, [1276](#)  
 Mole, [1276](#)  
 Molecular, [1276](#)  
 Molecular Chaperone, [1276](#)  
 Molecular Diagnosis, [1277](#)  
 Molecular Dynamics, [1277](#)  
 Molecular Imaging, [1277](#)  
 Molecular Medicine, [2125](#)  
 Molecular Pathology, [1277](#)  
 Molecular Pathology Of Colorectal Cancer, [1277](#)  
 Molecular Probe, [1277](#)  
 Molecular Response, [1277](#)  
 Molecular Testing, [1277](#)  
 Molecular Weight, [1277](#)  
 Molecule, [1277](#)  
 Moles, [1277](#)  
 Molimina, [1277](#)  
 Mollusc, [1277](#)  
 Mollusca, [1277](#)  
 Molluscipoxvirus, [1278](#)  
 Molluscum Contagiosum, [1278](#)  
 Molten Metal Fumes, [1278](#)  
 Molteno Implant, [1278](#)  
 Molybdenum, [1278](#)  
 Molybdenum Cofactor Deficiency, [1278](#)  
 Molybdenum-99, [1278](#)  
 Molybdosis, [1278](#)  
 Mometasone, [1278](#)  
 Mometasone Furoate, [1278](#)  
 Momordica, [1279](#)  
 Monascus, [1279](#)  
 Monckeberg Medial Calcific Sclerosis, [1279](#)

---

---

Mondor Disease, [1279](#)  
Mondor'S Disease, [1279](#)  
Mongolian Spot, [1279](#)  
Monilethrix, [1279](#)  
Monitored Anesthesia Care, [1279](#)  
Monitoring Device, [1279](#)  
Monitoring Test, [1279](#)  
Monkeypox, [1279](#)  
Monkeypox (Mpox), [1279](#)  
Mono, [1279](#)  
Mono (Infectious Mononucleosis), [1324](#)  
Monoamine Oxidase Inhibitor, [2125](#)  
Monoamine Oxidase Inhibitor (MAOI) Toxicity, [1279](#)  
Monoamine Oxidase Inhibitors, [1280](#)  
Monoarticular Joint Pain, [1280](#)  
Monobactams, [1280](#)  
Monoblast, [1280](#)  
Monoblast Count, [1280](#)  
Monochorionic Twin Pregnancy, [1280](#)  
Monoclonal, [1280](#)  
Monoclonal Antibodies, [1280](#)  
Monoclonal Antibody, [1280](#)  
Monoclonal Gammopathy Of Undetermined Significance, [1280](#)  
Monoclonal Gammopathy of Undetermined Significance (MGUS), [1280](#)  
Monocrotophos, [1280](#)  
Monocular, [1280](#)  
Monocyte, [1281](#)  
Monocytes, [1281](#)  
Monocytopenia, [1281](#)  
Monocytosis, [1281](#)  
Monogenic, [1281](#)  
Monogenic Obesity, [1281](#)  
Monograph, [1281](#)  
Monoiodotyrosine, [1281](#)  
Monokines, [1281](#)  
Monomer, [1281](#)  
Monomethylhydrazine, [1281](#)  
Monomodal Visual Amnesia, [1322](#)  
Mononegavirales, [1281](#)  
Mononeuritis Multiplex, [1281](#)  
Mononeuropathy, [1281](#)  
Mononuclear Leukocyte, [1282](#)  
Mononucleosis, [1282](#)  
Mononucleosis Spot Test, [1282](#)  
Monoplegia, [1282](#)  
Monopodium, [1282](#)  
Monorchidism, [1282](#)  
Monosaccharide, [1282](#)  
Monosodium Glutamate, [1282](#)  
Monosodium Urate, [1282](#)  
Monosomy, [1282](#)  
Monospot Test, [1282](#)  
Monotest, [1282](#)  
Monounsaturated Fat, [1282](#)  
Monozygotic Twin, [1282](#)  
Mons Veneris, [1283](#)  
Monteggia Fracture, [1283](#)  
Monteggia Fracture-Dislocation, [1283](#)  
Montelukast, [1283](#)  
Montelukast Sodium, [1283](#)  
Montevideo Units, [1283](#)  
Montgomery Tubercles, [1283](#)  
Montreal Cognitive Assessment, [1283](#)  
Mood, [1283](#)  
Mood Disorder, [1283](#)  
Mood Disorders, [1283](#)  
Mood Stabilizers, [1283](#)  
Mood Swing, [1283](#)  
Morale, [1284](#)  
Moraxella, [1284](#)  
Moraxella Atlantae, [1284](#)  
Moraxella Lincolnii, [1284](#)  
Moraxella Oblonga, [1284](#)  
Moraxella Osloensis, [1284](#)  
Moraxellaceae, [1284](#)  
Morbidity, [1284](#)  
Morbili, [1284](#)  
Morbili Form, [1284](#)  
Morbillivirus, [1284](#)  
Morbus, [1284](#)  
Morcellation, [1284](#)  
Mordant, [1284](#)  
Morgagni Hernia, [1284](#)  
Morgagni-Turner-Albright Syndrome, [1322](#)  
Morganella, [1284](#)  
Morganella Morganii, [1285](#)  
Morgellons Disease, [1285](#)  
Moribund, [1285](#)  
Morning Sickness, [1285](#)  
Moro Reflex, [1285](#)  
Morphea, [1285](#)  
Morpheus, [1285](#)  
Morphinans, [1285](#)  
Morphine, [1285](#)  
Morphine Overdose, [1285](#)  
Morphine Sulfate, [1285](#)  
Morphogenesis, [1285](#)  
Morphologic Response, [1285](#)  
Morphology, [1286](#)  
Morquio Syndrome, [1286](#)  
Mortality, [1286](#)  
Mortality Rate, [1286](#)  
Morton Neuroma, [1286](#)  
Morton's Metatarsalgia, [1286](#)  
Morton's Neuroma, [1286](#)  
Morton's Toe, [1286](#)

---

---

Morula, [1286](#)  
Morvan Disease – Syringomyelia, [1322](#)  
Morvan's Disease, [1286](#)  
Mosaic, [1286](#)  
Mosaicism, [1286](#)  
Mosquito, [1286](#)  
Mosquito Clamp, [1287](#)  
Mosquito-Borne, [1286](#)  
Motilin, [1287](#)  
Motility, [1287](#)  
Motion Analysis, [1287](#)  
Motion Artifact, [1287](#)  
Motion Correction, [1287](#)  
Motion Perception, [1287](#)  
Motion Sickness, [1287](#)  
Motivation, [1287](#)  
Motivational Interviewing, [1287](#)  
Motor, [1287](#)  
Motor Aphasia, [1287](#)  
Motor Cortex, [1287](#)  
Motor Delay, [1288](#)  
Motor End-Plate, [1288](#)  
Motor Endplate, [1288](#)  
Motor Incoordination, [1288](#)  
Motor Learning, [1288](#)  
Motor Neuron, [1288](#)  
Motor Neuron Disease, [1288](#)  
Motor Neuron Disorders Pathology, [1288](#)  
Motor Neurone Disease, [1288](#)  
Motor Neurones, [1288](#)  
Motor Stereotypy, [1288](#)  
Motor Tic, [1288](#)  
Motor Unit, [1288](#)  
Motor Vehicle Accident, [1289](#)  
Mottled Enamel, [1289](#)  
Mottled Teeth, [1289](#)  
Mouflage, [1289](#)  
Mould, [1289](#)  
Mountain Fever, [1322](#)  
Mountain Sickness, [1289](#)  
Mountain Sickness, Acute, [34](#)  
Mouth, [1289](#)  
Mouth (Oral) Cancer, [1322](#)  
Mouth Cancer, [1289](#)  
Mouth Diseases, [1289](#)  
Mouth Neoplasm, [1289](#)  
Mouth Sores, [1289](#)  
Mouth Ulcer, [1289](#)  
Mouth-To-Mouth Resuscitation, [1289](#)  
Mouthwash, [1289](#)  
Mouthwash Overdose, [1289](#)  
Movement, [1289](#)  
Movement Disorder, [1290](#)  
Movement Disorders, [1290](#)  
Movement Incoordination, [1290](#)  
Moxalactam, [1290](#)  
Moxibustion, [1290](#)  
Moxifloxacin, [1290](#)  
Moyamoya Disease, [1290](#)  
Moynihan's Hump, [1290](#)  
MPH, [1290](#)  
MPO, [1290](#)  
Mpox, [1290](#)  
MPS, [1322](#)  
MPS Disorder, [1322](#)  
MPS Disorder I, [1322](#)  
MPS Disorder II, [1323](#)  
MPS I, [1323](#)  
MPS II, [1323](#)  
MPTP, [1291](#)  
MR Angiography, [1291](#)  
MR Enterography, [1291](#)  
MR Neurography, [1292](#)  
MRA, [1291](#)  
MRCP, [1291](#)  
MRD, [1291](#)  
MRI, [1291](#)  
MRI And Low Back Pain, [1291](#)  
MRI Brain, [1291](#)  
MRI Contrast, [1291](#)  
MRI Joint, [1291](#)  
MRI Sequences, [1291](#)  
MRI-Targeted Fusion Prostate Biopsy, [1291](#)  
Mrna Precursor, [1291](#)  
MRNA Vaccine, [2125](#)  
MRSA, [1292](#)  
MRSA Decolonization, [1292](#)  
MRSA Infection, [1292](#)  
MS (Multiple Sclerosis), [1323](#)  
MS Alternative Therapies, [1292](#)  
MSA, [1323](#)  
MSG Symptom Complex, [1292](#)  
MSL, [1323](#)  
MST, [1292](#)  
MTOR, [1292](#)  
Mu-Chain Disease, [1292](#)  
Mucha-Habermann Disease, [1292](#)  
Mucin, [1292](#)  
Mucinoses, [1292](#)  
Mucinous, [1292](#)  
Mucinous Adenocarcinoma, [1293](#)  
Mucinous Borderline Ovarian Tumor, [1293](#)  
Mucinous Cyst Of The Vulva, [1293](#)  
Mucinous Cystadenoma, [1293](#)  
Mucinous Cystadenoma of Pancreas, [1293](#)  
Mucinous Ovarian Carcinoma, [1293](#)  
Muckle-Wells Syndrome, [1293](#)  
Mucoccele, [1293](#)

---

---

Mucocoele and Ranula, [1293](#)  
 Mucociliary Escalator, [1293](#)  
 Mucocutaneous Leishmaniasis, [1293](#)  
 Mucocutaneous Lymph Node Syndrome, [1293](#)  
 Mucoepidermoid Carcinoma, [1293](#)  
 Mucoid, [1294](#)  
 Mucolipidoses, [1294](#)  
 Mucolytic Therapy, [1294](#)  
 Mucolytics, [1294](#)  
 Mucopolysaccharides, [1294](#)  
 Mucopolysaccharidoses, [1294](#)  
 Mucopolysaccharidosis, [1294](#)  
 Mucopolysaccharidosis Type I, [1294](#)  
 Mucopolysaccharidosis Type II, [1294](#)  
 Mucopolysaccharidosis Type III, [1294](#)  
 Mucopolysaccharidosis Type IV, [1294](#)  
 Mucopolysaccharidosis VII, [1294](#)  
 Mucopus, [1294](#)  
 Mucor, [1295](#)  
 Mucor Ramosissimus, [1295](#)  
 Mucorales, [1295](#)  
 Mucormycosis, [1295](#)  
 Mucosa, [1295](#)  
 Mucosa-Associated Lymphoid Tissue, [1295](#)  
 Mucosa-Associated Lymphoid Tissue Lymphoma, [1295](#)  
 Mucosal, [1295](#)  
 Mucosal Candidiasis, [1295](#)  
 Mucosal Immunity, [1295](#)  
 Mucosal Infection, [1295](#)  
 Mucosal Irritation, [1295](#)  
 Mucosal Melanoma, [1295](#)  
 Mucosal Warts, [1295](#)  
 Mucositis, [1296](#)  
 Mucous, [1296](#)  
 Mucous Cysts, [1296](#)  
 Mucous Membrane, [1296](#)  
 Mucous Membrane (Cicatricial) Pemphigoid (MMP), [1296](#)  
 Mucous Membrane Pemphigoid, [1296](#)  
 Mucus, [1296](#)  
 Mucus Plugging, [1296](#)  
 Mud Fever, [1323](#)  
 Muehrcke Lines, [1296](#)  
 Muehrcke Lines of the Fingernails, [1296](#)  
 Muehrcke's Nails, [1296](#)  
 Muenke Syndrome, [1296](#)  
 MUGA Scan, [1296](#)  
 Muir-Torre Syndrome, [1297](#)  
 Mulberry Molars, [1297](#)  
 Mulibrey Nanism, [1297](#)  
 Muller Cells, [1297](#)  
 Mullerian Agenesis, [1297](#)  
 Mullerian Duct Anomalies, [1297](#)  
 Multi-Drug Resistance, [2125](#)  
 Multi-Infarct Dementia, [1298](#)  
 Multi-Minicore Disease, [1299](#)  
 Multicenter Study, [1297](#)  
 Multicentric, [1297](#)  
 Multicentric Reticulohistiocytosis, [1297](#)  
 Multicystic Dysplastic Kidney, [1297](#)  
 Multidetector Computed Tomography, [1297](#)  
 Multidisciplinary Disaster Victim Identification, [1298](#)  
 Multidisciplinary Team, [1298](#)  
 Multidrug-Resistant Extensively Drug-Resistant TB, [2126](#)  
 Multidrug-Resistant TB, [1298](#)  
 Multidrug-Resistant Tuberculosis, [1298](#)  
 Multienzyme Complexes, [1298](#)  
 Multifactorial, [1298](#)  
 Multifocal, [1298](#)  
 Multifocal Atrial Tachycardia, [1298](#)  
 Multifocal Motor Neuropathy, [1298](#)  
 Multigravida, [1298](#)  
 Multilocular, [1298](#)  
 Multimodal Analgesia, [1299](#)  
 Multimodal Imaging, [1299](#)  
 Multimorbidity, [1299](#)  
 Multinodular Goiter, [1299](#)  
 Multinucleated Hepatocyte, [1299](#)  
 Multipara, [1299](#)  
 Multiparametric MRI Of The Prostate, [1299](#)  
 Multiparous, [1299](#)  
 Multiple Bacterial Abscesses, [1299](#)  
 Multiple Biliary Hamartomas, [1299](#)  
 Multiple Endocrine Deficiency Syndrome, [1323](#)  
 Multiple Endocrine Neoplasia, [1299](#)  
 Multiple Endocrine Neoplasia I, [1299](#)  
 Multiple Endocrine Neoplasia II, [1300](#)  
 Multiple Endocrine Neoplasia Type 1, [1300](#)  
 Multiple Endocrine Neoplasia Type 2, [1300](#)  
 Multiple Endocrine Neoplasia Type 2 (MEN2), [1300](#)  
 Multiple Endocrine Neoplasia Type 2A, [1300](#)  
 Multiple Endocrine Neoplasia Type 2B, [1300](#)  
 Multiple Endocrine Neoplasia Type 4, [1300](#)  
 Multiple Exostoses, [1300](#)  
 Multiple Fibrofolliculomas, [1300](#)  
 Multiple Inherited Schwannomas, [1300](#)  
 Multiple Lentigines Syndrome, [1323](#)  
 Multiple Mononeuropathy, [1300](#)  
 Multiple Myeloma, [1300](#)  
 Multiple Myeloma: Deadly To Chronic, [1301](#)  
 Multiple Organ Dysfunction Syndrome, [1301](#)  
 Multiple Organ Failure, [1301](#)  
 Multiple Peripheral Neuritis, [1323](#)  
 Multiple Polyposis of the Colon, [1323](#)  
 Multiple Pregnancy, [1301](#)  
 Multiple Pregnancy: Twins or More, [1323](#)  
 Multiple Sclerosis, [1301](#)

---



---

Multiple Sclerosis (MS), [1323](#)  
Multiple Sclerosis MS And Pregnancy, [1301](#)  
Multiple Sleep Latency Test, [1301](#)  
Multiple Sulfatase Deficiency, [1323](#)  
Multiple Symmetric Lipomatosis, [1323](#)  
Multiple System Atrophy, [1301](#)  
Multiple System Atrophy Cerebellar Type, [1301](#)  
Multiple System Atrophy Parkinsonian Type, [1302](#)  
Multiple System Failure, [1302](#)  
Multiple Vitamin Overdose, [1302](#)  
Multiplex PCR Panel, [1302](#)  
Multiplicity, [1302](#)  
Multipolar Neuron, [1302](#)  
Multisystem Inflammatory Syndrome In Children, [1302](#)  
Multivesicular Body, [1302](#)  
Mumps, [1302](#)  
Mumps Orchitis, [1302](#)  
Mumps Parotitis, [1303](#)  
Mumps Virus, [1303](#)  
Munchausen Syndrome, [1303](#)  
Munchausen Syndrome by Proxy, [1323](#)  
Mupirocin, [1303](#)  
Mural, [1303](#)  
Mural Thrombosis, [1303](#)  
Mural Thrombus, [1303](#)  
Muramidase, [1303](#)  
Murine Typhus, [1303](#)  
Murk Jansen Metaphyseal Chondrodysplasia, [1323](#)  
Murmur, [1303](#)  
Murmur Classification, [1303](#)  
Murphy Sign, [1303](#)  
Murphy's Sign, [1303](#)  
Musca, [1303](#)  
Muscae Volitantes, [1303](#)  
Muscarine, [1303](#)  
Muscarinic Antagonist, [1304](#)  
Muscidae, [1304](#)  
Muscimol, [1304](#)  
Muscle, [1304](#)  
Muscle Aches, [1304](#)  
Muscle Atrophy, [1304](#)  
Muscle Belly, [1304](#)  
Muscle Biopsy, [1304](#)  
Muscle Cell, [1304](#)  
Muscle Contraction, [1304](#)  
Muscle Contracture, [1304](#)  
Muscle Cramp, [1304](#)  
Muscle Cramps, [1305](#)  
Muscle Development, [1305](#)  
Muscle Disorder, [1305](#)  
Muscle Fascicle, [1305](#)  
Muscle Fatigue, [1305](#)  
Muscle Fiber, [1305](#)  
Muscle Fibers, [1305](#)  
Muscle Function Loss, [1305](#)  
Muscle Hypertrophy, [1305](#)  
Muscle Insertion, [1305](#)  
Muscle Neoplasm, [1305](#)  
Muscle Origin, [1305](#)  
Muscle Pain, [1305](#)  
Muscle Protein, [1306](#)  
Muscle Relaxant, [1306](#)  
Muscle Relaxants, [1306](#)  
Muscle Relaxation, [1306](#)  
Muscle Rigidity, [1306](#)  
Muscle Spasm, [1306](#)  
Muscle Spasms, [1306](#)  
Muscle Spindle, [1306](#)  
Muscle Strain Treatment, [1306](#)  
Muscle Strength, [1306](#)  
Muscle Tissue, [1306](#)  
Muscle Tone, [1306](#)  
Muscle Tonus, [1307](#)  
Muscle Twitching, [1307](#)  
Muscle Weakness, [1307](#)  
Muscle-Contraction Headache, [1304](#)  
Muscle-Invasive Bladder Cancer, [1305](#)  
Muscle-Tendon Unit, [1306](#)  
Muscular, [1307](#)  
Muscular Dystrophies, [1323](#)  
Muscular Dystrophy, [1307](#)  
Muscular Endurance, [1307](#)  
Muscular System, [1307](#)  
Muscularis, [1307](#)  
Musculocutaneous Nerve, [1307](#)  
Musculoskeletal, [1307](#)  
Musculoskeletal Deformity, [1307](#)  
Musculoskeletal Disease, [1307](#)  
Musculoskeletal Injury, [1307](#)  
Musculoskeletal Pain, [1308](#)  
Musculoskeletal System, [1308](#)  
Musculoskeletal Tumor, [1308](#)  
Musculus, [1308](#)  
MUSE, [1308](#)  
Mushroom, [1308](#)  
Mushroom Poisoning, [1308](#)  
Mushroom Toxicity, [1308](#)  
Music Therapy, [1308](#)  
MuSK-Positive Myasthenia Gravis, [1308](#)  
Mustard Compounds, [1308](#)  
Mustard Gas, [1308](#)  
Mutagen, [1308](#)  
Mutant, [1309](#)  
Mutant Proteins, [1309](#)  
Mutate, [1309](#)  
Mutation, [1309](#)  
Mutation Abnormality, [1309](#)  
Mutation Rate, [1309](#)

---

---

Mute, [1309](#)  
 Mutism, [1309](#)  
 Myalgia, [1309](#)  
 Myalgic Encephalomyelitis, [1309](#)  
 Myalgic Encephalomyelitis/Chronic Fatigue Syndrome, [1309](#)  
 Myasthenia, [1309](#)  
 Myasthenia Crisis, [1309](#)  
 Myasthenia Gravis, [1310](#)  
 Myasthenia Gravis and Pregnancy, [1310](#)  
 Myasthenic Crisis, [1310](#)  
 Myasthenic Syndrome, [1310](#)  
 Mycelium, [1310](#)  
 Mycetoma, [1310](#)  
 Mycobacteriaceae, [1310](#)  
 Mycobacterial Culture, [1310](#)  
 Mycobacterial Infection, [1310](#)  
 Mycobacterium, [1310](#)  
 Mycobacterium Abscessus, [1310](#)  
 Mycobacterium Avium, [1311](#)  
 Mycobacterium Avium Complex, [1311](#)  
 Mycobacterium Bovis, [1311](#)  
 Mycobacterium Chelonae, [1311](#)  
 Mycobacterium Fortuitum, [1311](#)  
 Mycobacterium Gordona, [1311](#)  
 Mycobacterium Haemophilum, [1311](#)  
 Mycobacterium Infection, [1311](#)  
 Mycobacterium Kansasii, [1311](#)  
 Mycobacterium Lepae, [1311](#)  
 Mycobacterium Marinum, [1311](#)  
 Mycobacterium Scrofulaceum, [1311](#)  
 Mycobacterium Szulgai, [1311](#)  
 Mycobacterium Tuberculosis, [1311](#)  
 Mycobacterium Ulcerans, [1311](#)  
 Mycology, [1311](#)  
 Mycophenolate Mofetil, [1312](#)  
 Mycophenolic Acid, [1312](#)  
 Mycoplasma, [1312](#)  
 Mycoplasma Genitalium, [1312](#)  
 Mycoplasma Hominis, [1312](#)  
 Mycoplasma Pneumonia, [1312](#)  
 Mycoplasma Pneumoniae, [1312](#)  
 Mycoplasma Pneumoniae-Associated Mucositis, [1312](#)  
 Mycoplasmal Pneumonia, [1312](#)  
 Mycoplasmataceae, [1312](#)  
 Mycoses, [1312](#)  
 Mycosis, [1312](#)  
 Mycosis Fungoides, [1312](#)  
 Mycosis Fungoides Lymphoma, [1312](#)  
 Mycotic Aneurysm, [1313](#)  
 Mycotoxin, [1313](#)  
 Mydriasis, [1313](#)  
 Mydriatic Agents, [1313](#)  
 Myelin, [1313](#)  
 Myelin Sheath, [1313](#)  
 Myelination, [1313](#)  
 Myelitis, [1313](#)  
 Myeloablative Agonist, [1313](#)  
 Myeloablative Chemotherapy, [1313](#)  
 Myeloblast, [1313](#)  
 Myelocyte, [1313](#)  
 Myelocytic Leukaemia, [1313](#)  
 Myelodysplastic Neoplasms, [1314](#)  
 Myelodysplastic Or Myeloproliferative Neoplasms, [1314](#)  
 Myelodysplastic Syndrome, [1314](#)  
 Myelodysplastic Syndrome (MDS), [1323](#)  
 Myelodysplastic Syndromes, [1314](#)  
 Myeloencephalitis, [1314](#)  
 Myelofibrosis, [1314](#)  
 Myelogenous, [1314](#)  
 Myelogram, [1314](#)  
 Myelography, [1314](#)  
 Myeloid, [1314](#)  
 Myeloid Cell, [1314](#)  
 Myeloid Leukemia, [1314](#)  
 Myeloid Sarcoma, [1314](#)  
 Myelolipoma, [1314](#)  
 Myeloma, [1314](#)  
 Myeloma Cast Nephropathy, [1314](#)  
 Myelomalacia, [1315](#)  
 Myelomeningocele, [1315](#)  
 Myelomonocyte, [1315](#)  
 Myelopathy, [1315](#)  
 Myeloperoxidase, [1315](#)  
 Myelopoiesis, [1315](#)  
 Myeloproliferative Disorders, [1315](#)  
 Myeloproliferative Neoplasm, [1315](#)  
 Myelosuppression, [1315](#)  
 Myelosuppressive Therapy, [1315](#)  
 Myenteric Plexus, [1315](#)  
 Myhre Syndrome, [1315](#)  
 Myiasis, [1315](#)  
 Myoblast, [1315](#)  
 Myoblastoma, [1316](#)  
 Myocardial, [1316](#)  
 Myocardial Biopsy, [1316](#)  
 Myocardial Bridging, [1316](#)  
 Myocardial Contusion, [1316](#)  
 Myocardial Infarction, [1316](#)  
 Myocardial Ischemia, [1316](#)  
 Myocardial Perfusion Imaging, [1316](#)  
 Myocardial Reperfusion, [1316](#)  
 Myocardial Rupture, [1316](#)  
 Myocardial Viability, [1316](#)  
 Myocarditis, [1316](#)  
 Myocarditis Diagnosis, [1316](#)  
 Myocarditis Pathology, [1316](#)  
 Myocarditis Treatment, [1317](#)

---

---

Myocardium, [1317](#)  
Myoclonic Epilepsy, [1317](#)  
Myoclonic Seizure, [1317](#)  
Myoclonus, [1317](#)  
Myocyte, [1317](#)  
Myodegeneration, [1317](#)  
Myoepithelial Carcinoma Pathology, [1317](#)  
Myoepithelial Cell, [1317](#)  
Myoepithelioma, [1317](#)  
Myoepithelioma Of Soft Tissue, [1317](#)  
Myofascial Pain Syndrome, [1317](#)  
Myofibril, [1317](#)  
Myofibrillar Myopathy, [1317](#)  
Myofibrils, [1318](#)  
Myofibroblastoma, [1318](#)  
Myofibroblastoma Of The Breast, [1318](#)  
Myofibroblasts, [1318](#)  
Myofibromatosis, [1318](#)  
Myofilament, [1318](#)  
Myofilaments, [1318](#)  
Myogenin, [1318](#)  
Myoglobin, [1318](#)  
Myoglobin Blood Test, [1318](#)  
Myoglobin Urine Test, [1318](#)  
Myoglobinuria, [1318](#)  
Myokymia, [1318](#)  
Myoma, [1318](#)  
Myomalacia, [1318](#)  
Myomas, [1319](#)  
Myomectomy, [1319](#)  
Myometrium, [1319](#)  
Myopathic Facies, [1319](#)  
Myopathy, [1319](#)  
Myopericytoma, [1319](#)  
Myopia, [1319](#)  
Myosin, [1319](#)  
Myosin Atpase, [1319](#)  
Myositis, [1319](#)  
Myositis Ossificans, [1319](#)  
Myotonia, [1319](#)  
Myotonia Congenita, [1319](#)  
Myotonia Fluctuans, [1319](#)  
Myotonia Permanens, [1320](#)  
Myotonic Disorder, [1320](#)  
Myotonic Dystrophy, [1320](#)  
Myotonic Dystrophy Type 2, [1320](#)  
Myringitis, [1320](#)  
Myringotomy, [1320](#)  
Myringotomy With Tympanostomy Tube, [1320](#)  
Myristica Oil Poisoning, [1320](#)  
Myxedema, [1320](#)  
Myxedema Coma, [1320](#)  
Myxofibroma, [1320](#)  
Myxoid Cyst, [1321](#)  
Myxoma, [1321](#)  
Myxopapillary Ependymoma, [1321](#)  
Myxosarcoma, [1321](#)  
MZL, [1321](#)  
MéNièRe'S Disease, [1273](#)  
MéNéTrier'S Disease, [1273](#)  
N-Acetylcysteine, [1325](#)  
N-Acetylglutamate Synthetase Deficiency, [1325](#)  
Nabothian Cyst, [1325](#)  
Nabumetone, [1325](#)  
Nadir, [1325](#)  
Naegle Rule, [1325](#)  
Naegeli-Franceschetti-Jadassohn Syndrome, [1325](#)  
Naegleria Fowleri, [1325](#)  
Naegleria Infection, [1325](#)  
Naevus, [1326](#)  
Naevus Flammeus, [1326](#)  
Naffziger Sign, [1326](#)  
Nail, [1326](#)  
Nail Abnormalities, [1326](#)  
Nail Bed Injury, [1326](#)  
Nail Changes, [1326](#)  
Nail Clipping For Fungal Testing, [1326](#)  
Nail Clubbing, [1326](#)  
Nail Fungus, [1326](#)  
Nail Furrows, [1326](#)  
Nail Infection, [1327](#)  
Nail Injuries, [1327](#)  
Nail Patella Syndrome, [1327](#)  
Nail Pits, [1327](#)  
Nail Pitting, [1327](#)  
Nail Psoriasis, [1327](#)  
Nail Surgery, [1327](#)  
Nail-Biting, [1381](#)  
Nail-Fold Capillaroscopy, [1326](#)  
Nail-Patella Syndrome, [1327](#)  
Nails, [1327](#)  
Nalbuphine, [1327](#)  
Naloxone, [1327](#)  
Naltrexone, [1327](#)  
Nanocolloid, [1327](#)  
Nanomedicine, [1327](#)  
Nanoparticle, [1328](#)  
Nanoscience, [1328](#)  
Nanotechnology, [1328](#)  
Napelline, [1328](#)  
Naphazoline, [1328](#)  
Naphthalene Poisoning, [1328](#)  
Naproxen, [1328](#)  
Naproxen Sodium Overdose, [1328](#)  
Narcissism, [1328](#)  
Narcissistic Personality Disorder, [1328](#)  
Narcolepsy, [1328](#)

---

---

Narcoleptic Syndrome, [1381](#)  
NARP Syndrome, [1328](#)  
Narrow Angle Glaucoma, [1329](#)  
Narrow Pulse Pressure, [1329](#)  
Narrow Therapeutic Index, [1329](#)  
Narrow-Complex Tachycardia, [1329](#)  
Narrowband UVB Phototherapy, [1329](#)  
Nasal, [1329](#)  
Nasal Bleeding, [1329](#)  
Nasal Cannula, [1329](#)  
Nasal Cavity, [1329](#)  
Nasal Congestion, [1329](#)  
Nasal Congestion And Rhinorrhea, [1329](#)  
Nasal Corticosteroid Sprays, [1329](#)  
Nasal Discharge, [1329](#)  
Nasal Disorders, [1329](#)  
Nasal Endoscopy, [1329](#)  
Nasal Flaring, [1330](#)  
Nasal Foreign Bodies, [1330](#)  
Nasal Fracture, [1330](#)  
Nasal Implant, [1330](#)  
Nasal Mucosal Biopsy, [1330](#)  
Nasal Obstruction, [1330](#)  
Nasal Passage, [1330](#)  
Nasal Polyps, [1330](#)  
Nasal Septal Hematoma, [1330](#)  
Nasal Septal Perforation Repair, [1381](#)  
Nasal Septum, [1330](#)  
Nasal Septum Perforation, [1330](#)  
Nasal Spray, [1330](#)  
Nasal Stent, [1330](#)  
Nasal Vestibulitis, [1330](#)  
Nasogastric Feeding Tube, [1331](#)  
Nasogastric Tube, [1331](#)  
Nasojejunal Feeding Tube, [1331](#)  
Nasolacrimal Duct, [1331](#)  
Nasopharyngeal Airway, [1331](#)  
Nasopharyngeal Cancer, [1331](#)  
Nasopharyngeal Carcinoma, [1331](#)  
Nasopharyngeal Culture, [1331](#)  
Nasopharynx, [1331](#)  
Natal Cleft, [1331](#)  
Natal Teeth, [1331](#)  
Natriuresis, [1331](#)  
Natriuretic, [1331](#)  
Natural Immunity, [1332](#)  
Natural Killer Cell, [1332](#)  
Natural Killer Cells, [1332](#)  
Naturopathic Medicine, [1381](#)  
Nausea, [1332](#)  
Nausea And Vomiting, [1332](#)  
Nausea And Vomiting During Early Pregnancy, [1332](#)  
Nausea And Vomiting – Adults, [1332](#)  
Navel, [1332](#)  
Navicular, [1332](#)  
Navicular Fracture, [1332](#)  
Navicular Fracture of the Wrist, [1381](#)  
NCS, [1332](#)  
NDI, [1381](#)  
NDM-1, [1332](#)  
Near-Drowning, [1332](#)  
Nearsightedness, [1333](#)  
Nearsightedness (Myopia), [1381](#)  
Nebulizer, [1333](#)  
NEC, [1333](#)  
Necator, [1333](#)  
Necator Americanus, [1333](#)  
Neck, [1333](#)  
Neck And Back Pain, [1333](#)  
Neck Cancer, [1333](#)  
Neck Dissection, [1333](#)  
Neck Lump, [1333](#)  
Neck Mass, [1333](#)  
Neck Pain, [1333](#)  
Neck Pain And Dizziness, [1334](#)  
Neck Sprain, [1334](#)  
Neck Stiffness, [1333](#)  
Neck Strain, [1334](#)  
Neck X-Ray, [1334](#)  
Necrobiosis Lipoidica, [1334](#)  
Necrobiosis Lipoidica Diabeticorum, [1334](#)  
Necrolytic Acral Erythema, [1334](#)  
Necrolytic Migratory Erythema, [1334](#)  
Necrosis, [1334](#)  
Necrotizing Anterior Scleritis, [1334](#)  
Necrotizing Enterocolitis, [1334](#)  
Necrotizing Fasciitis, [1334](#)  
Necrotizing Fasciitis of the Perineum, [1381](#)  
Necrotizing Granuloma, [1334](#)  
Necrotizing Infections, [1334](#)  
Necrotizing Pneumonia, [1334](#)  
Necrotizing Sialometaplasia, [1335](#)  
Necrotizing Soft Tissue Infection, [1335](#)  
Necrotizing Vasculitis, [1335](#)  
Needle, [1335](#)  
Needle Aspiration Biopsy, [1381](#)  
Needle Biopsy, [1335](#)  
Needle Decompression, [1335](#)  
Needle Driver, [1335](#)  
Needle Holder, [1335](#)  
Needlestick Injury, [1335](#)  
Neer Sign, [1335](#)  
Neer Test, [1335](#)  
Negative Feedback, [1335](#)  
Negative Predictive Value, [1336](#)  
Negative Pressure Pulmonary Edema, [1336](#)  
Negative Pressure Ventilation, [1336](#)  
Negative Symptoms, [1336](#)

---

---

Negative-Strand RNA Virus, [1336](#)  
Neglect, [1336](#)  
Neglect Syndrome, [1336](#)  
Neglected Tropical Disease, [1336](#)  
Neisseria Gonorrhoeae, [1336](#)  
Neisseria Infection, [1336](#)  
Neisseria Meningitidis, [1336](#)  
Nelson Syndrome, [1336](#)  
Nemaline Myopathy, [1337](#)  
Neoadjuvant Therapy, [1337](#)  
Neocortex, [1337](#)  
Neoehrlichiosis, [1337](#)  
Neologisms, [1337](#)  
Neonatal, [1337](#)  
Neonatal Abstinence Syndrome, [1337](#)  
Neonatal Cephalohematoma, [1337](#)  
Neonatal Conjunctivitis, [1337](#)  
Neonatal Cystic Fibrosis Screening Test, [1337](#)  
Neonatal Diabetes Mellitus, [1337](#)  
Neonatal Encephalopathy, [1338](#)  
Neonatal Galactorrhea, [1337](#)  
Neonatal Herpes Simplex, [1337](#)  
Neonatal Herpes Simplex Virus, [1338](#)  
Neonatal Hyperbilirubinemia, [1338](#)  
Neonatal Hyperglycemia, [1338](#)  
Neonatal Hypertension, [1338](#)  
Neonatal Hypocalcemia, [1338](#)  
Neonatal Hypoglycemia, [1338](#)  
Neonatal Hypothermia, [1338](#)  
Neonatal Hypothyroidism, [1338](#)  
Neonatal Hypoxic-Ischemic Encephalopathy, [1338](#)  
Neonatal Infections, [1338](#)  
Neonatal Jaundice, [1339](#)  
Neonatal Listeriosis, [1381](#)  
Neonatal Lupus, [1339](#)  
Neonatal Mechanical Ventilation, [1339](#)  
Neonatal Neutropenia, [1339](#)  
Neonatal Opioid Withdrawal Syndrome, [1339](#)  
Neonatal Peripheral Arterial Line, [1339](#)  
Neonatal Peripheral IV Line, [1339](#)  
Neonatal PICC Line, [1339](#)  
Neonatal Pneumothorax, [1339](#)  
Neonatal Polycythemia, [1340](#)  
Neonatal Respiratory Distress Syndrome, [1340](#)  
Neonatal Resuscitation, [1340](#)  
Neonatal Seizures, [1340](#)  
Neonatal Sepsis, [1340](#)  
Neonatal Skin Conditions, [1340](#)  
Neonatal Testicular Torsion, [1340](#)  
Neonatal Thyrotoxicosis, [1340](#)  
Neonatal Transition, [1340](#)  
Neonate, [1340](#)  
Neonatologist, [1340](#)  
Neoplasia, [1340](#)  
Neoplasm, [1340](#)  
Neoplastic, [1341](#)  
Neostigmine, [1341](#)  
Nephoptosis, [1341](#)  
Nephrectomy, [1341](#)  
Nephric, [1341](#)  
Nephritic Syndrome, [1341](#)  
Nephritis, [1341](#)  
Nephroblastoma, [1341](#)  
Nephrocalcinosis, [1341](#)  
Nephrocarcinoma, [1381](#)  
Nephrogenic Diabetes Insipidus, [1341](#)  
Nephrogenic Fibrosing Dermopathy, [1341](#)  
Nephrogenic Systemic Fibrosis, [1341](#)  
Nephrolithiasis, [1341](#)  
Nephrologist, [1341](#)  
Nephrology, [1341](#)  
Nephrolysis, [1342](#)  
Nephron, [1342](#)  
Nephron Function, [1342](#)  
Nephronophthisis, [1342](#)  
Nephropathy, [1342](#)  
Nephropathy Diabetic, [1381](#)  
Nephroptosis, [1342](#)  
Nephrosclerosis, [1342](#)  
Nephrostomy, [1342](#)  
Nephrostomy Tube, [1342](#)  
Nephrotic Range Proteinuria, [1342](#)  
Nephrotic Syndrome, [1342](#)  
Nephrotoxic, [1342](#)  
Nephrotoxicity, [1342](#)  
Nernst Equation, [1343](#)  
Nerve, [1343](#)  
Nerve Biopsy, [1343](#)  
Nerve Block, [1343](#)  
Nerve Compression, [1343](#)  
Nerve Conduction Studies, [1343](#)  
Nerve Conduction Study, [1343](#)  
Nerve Conduction Velocity, [1343](#)  
Nerve Deafness, [1343](#)  
Nerve Endings, [1343](#)  
Nerve Fiber, [1343](#)  
Nerve Fiber Types, [1343](#)  
Nerve Impulse, [1343](#)  
Nerve Injury, [1343](#)  
Nerve Plexus, [1344](#)  
Nerve Root, [1344](#)  
Nerve Sheath Myxoma, [1344](#)  
Nerve Sparing, [1344](#)  
Nerve Tract, [1344](#)  
Nervous Colon Syndrome, [1344](#)  
Nervous System, [1344](#)  
Nervous Tissue, [1344](#)  
Nesidioblastosis, [1344](#)

---

NET, [1344](#)  
Netherton Syndrome, [1344](#)  
Nettle Rash, [1344](#)  
Neural, [1344](#)  
Neural Crest, [1345](#)  
Neural Pathway, [1345](#)  
Neural Tube Defects, [1345](#)  
Neuralgia, [1345](#)  
Neuralgia Amyotrophica, [1345](#)  
Neuraminidase Inhibitors, [1345](#)  
Neurapraxia, [1345](#)  
Neurilemma, [1345](#)  
Neurilemmoma, [1345](#)  
Neurinomatosis, [1345](#)  
Neuritic Plaques, [1345](#)  
Neuritis, [1345](#)  
Neuro-Ophthalmology, [1352](#)  
Neuroacanthocytosis, [1345](#)  
Neuroanatomy, [1345](#)  
Neuroblastoma, [1346](#)  
Neurocardiogenic Syncope, [1346](#)  
Neurocognitive Disorder, [1346](#)  
Neurocognitive Disorders, [1346](#)  
Neurocutaneous Syndrome, [1346](#)  
Neurocutaneous Syndromes Pathology, [1346](#)  
Neurocysticercosis, [1346](#)  
Neurocytoma, [1346](#)  
Neurodegeneration, [1346](#)  
Neurodegeneration With Brain Iron Accumulation, [1346](#)  
Neurodegenerative Disease, [1346](#)  
Neurodegenerative Disorder, [1346](#)  
Neurodermatitis, [1346](#)  
Neurodevelopmental Disorders, [1347](#)  
Neuroendocrine, [1347](#)  
Neuroendocrine Carcinoma, [1347](#)  
Neuroendocrine Carcinoma of the Skin, [1347](#)  
Neuroendocrine Tumor, [1347](#)  
Neuroendocrine Tumors, [1347](#)  
Neuroendocrine Tumors Grading, [1347](#)  
Neuroendocrinology, [1347](#)  
Neurofibrillary Tangles, [1347](#)  
Neurofibroma, [1347](#)  
Neurofibromatosis, [1347](#)  
Neurofibromatosis 2, [1348](#)  
Neurofibromatosis Type 1, [1348](#)  
Neurofibromatosis Type 2, [1348](#)  
Neurofibromatosis Type I, [1348](#)  
Neurofibromatosis Type II, [1348](#)  
Neurofibromatosis-1, [1347](#)  
Neurofibrosarcoma, [1348](#)  
Neurogenesis, [1348](#)  
Neurogenic, [1348](#)  
Neurogenic Bladder, [1348](#)  
Neurogenic Claudication, [1348](#)  
Neurogenic Megacolon, [1348](#)  
Neurogenic Orthostatic Hypotension, [1348](#)  
Neurogenic Pulmonary Edema, [1349](#)  
Neurogenic Shock, [1349](#)  
Neuroglia, [1349](#)  
Neurography, [1349](#)  
Neurohypophysis, [1349](#)  
Neuroimmunology, [1348](#)  
Neurolathyrism, [1349](#)  
Neuroleptic, [1349](#)  
Neuroleptic Agent Toxicity, [1349](#)  
Neuroleptic Agents, [1349](#)  
Neuroleptic Malignant Syndrome, [1349](#)  
Neurologic Deficit, [1350](#)  
Neurologic Disorders, [1350](#)  
Neurologic Localization, [1350](#)  
Neurological, [1349](#)  
Neurological Disease, [1349](#)  
Neurological Examination, [1349](#)  
Neurological Rehabilitation, [1350](#)  
Neurologist, [1350](#)  
Neurology, [1350](#)  
Neurology PET, [1350](#)  
Neurolysis, [1350](#)  
Neuroma, [1350](#)  
Neuromodulation, [1350](#)  
Neuromuscular, [1350](#)  
Neuromuscular Blockade Reversal, [1350](#)  
Neuromuscular Blocking Agents, [1350](#)  
Neuromuscular Disorders, [1351](#)  
Neuromuscular Hamartoma, [1351](#)  
Neuromuscular Junction, [1351](#)  
Neuromyelitis Optica, [1351](#)  
Neuromyelitis Optica Spectrum Disorder, [1351](#)  
Neuromyotonia, [1351](#)  
Neuron, [1351](#)  
Neuron-Specific Enolase, [1351](#)  
Neuron-Specific Enolase Tumor Marker, [1352](#)  
Neuronal Ceroid Lipofuscinoses, [1351](#)  
Neuronal Migration Disorder, [1351](#)  
Neurone, [1351](#)  
Neuronopathy, [1351](#)  
Neurons, [1351](#)  
Neuropathic, [1352](#)  
Neuropathic Pain, [1352](#)  
Neuropathic Ulcer, [1352](#)  
Neuropathy, [1352](#)  
Neuropathy Secondary To Drugs, [2126](#)  
Neuropeptide, [1352](#)  
Neuropeptides, [1352](#)  
Neuropharmacology, [1352](#)  
Neurophysiology, [1352](#)  
Neurophysiotherapy, [1352](#)  
Neuroplasticity, [1352](#)



---

Neuroplegic, [1352](#)  
 Neuropsychiatry, [1353](#)  
 Neuropsychopharmacology, [1353](#)  
 Neuroradiology, [1353](#)  
 Neuroretinitis, [1353](#)  
 Neurosarcoidosis, [1353](#)  
 Neuroscience, [1353](#)  
 Neurosciences, [1353](#)  
 Neurosurgeon, [1353](#)  
 Neurosurgery, [1353](#)  
 Neurosyphilis, [1353](#)  
 Neurotoxicity, [1353](#)  
 Neurotransmission, [1353](#)  
 Neurotransmitter, [1353](#)  
 Neurotransmitter Receptors, [1353](#)  
 Neurotrophic, [1353](#)  
 Neurotrophic Keratopathy, [1354](#)  
 Neurotropic, [1354](#)  
 Neurovascular Examination, [1354](#)  
 Neutral Alignment, [1354](#)  
 Neutral Endopeptidase, [1354](#)  
 Neutralization, [1354](#)  
 Neutralizing Antibodies, [1354](#)  
 Neutropenia, [1354](#)  
 Neutropenic Enterocolitis, [1354](#)  
 Neutropenic Fever, [1354](#)  
 Neutrophil, [1354](#)  
 Neutrophil Extracellular Traps, [1354](#)  
 Neutrophil Recruitment, [1355](#)  
 Neutrophilic Dermatitis, [1355](#)  
 Neutrophilic Eccrine Hidradenitis, [1355](#)  
 Neutrophils, [1355](#)  
 Nevi of Ota and Ito, [1355](#)  
 Nevoid Basal Cell Carcinoma Syndrome, [1355](#)  
 Nevus, [1355](#)  
 Nevus Anemicus, [1355](#)  
 Nevus Araneus (Spider Nevus), [1355](#)  
 Nevus Comedonicus, [1355](#)  
 Nevus Comedonicus Syndrome, [1355](#)  
 Nevus of Ito, [1355](#)  
 Nevus of Ota, [1356](#)  
 Nevus Sebaceus, [1356](#)  
 New Daily Persistent Headache, [1357](#)  
 New York Heart Association Functional Classification, [1357](#)  
 Newborn, [1356](#)  
 Newborn Circumcision, [1356](#)  
 Newborn Head Molding, [1356](#)  
 Newborn Hearing Screen, [1356](#)  
 Newborn Infant Development, [1356](#)  
 Newborn Jaundice, [1356](#)  
 Newborn Metabolic Screen, [1356](#)  
 Newborn Screening, [1356](#)  
 Newborn Screening Panel, [1356](#)  
 Newborn Screening Tests, [1356](#)  
 Nexium, [1357](#)  
 Next-Generation Sequencing, [1357](#)  
 Nexus, [1357](#)  
 NF1, [1357](#)  
 NF2, [1357](#)  
 NF2-Related Schwannomatosis, [1357](#)  
 Niacin, [1357](#)  
 Niacin Deficiency, [1357](#)  
 Niacin For Cholesterol, [1357](#)  
 Niacin Vitamin B3, [1357](#)  
 Niacinamide, [1357](#)  
 Nialamide, [1357](#)  
 Nicardipine, [1358](#)  
 Nickel Allergy, [1358](#)  
 Nickel-Chromium, [1358](#)  
 Niclosamide, [1358](#)  
 Nicolau Syndrome, [1358](#)  
 Nicotinamide, [1358](#)  
 Nicotinamide Adenine Dinucleotide, [1358](#)  
 Nicotine, [1358](#)  
 Nicotine Addiction, [1358](#)  
 Nicotine And Tobacco, [1358](#)  
 Nicotine Dependence, [1358](#)  
 Nicotine Poisoning, [1358](#)  
 Nicotine Replacement Therapy, [1358](#)  
 Nicotine Withdrawal, [1382](#)  
 Nicotinic, [1358](#)  
 Nicotinic Acetylcholine Receptor, [1359](#)  
 Nicotinic Acid, [1359](#)  
 NICU Staff, [1359](#)  
 Nidation, [1359](#)  
 Nidus, [1359](#)  
 Niemann-Pick Disease, [1359](#)  
 Nifedipine, [1359](#)  
 Night Blindness, [1359](#)  
 Night Eating, [1382](#)  
 Night Eating Syndrome, [1359](#)  
 Night Leg Cramps, [1359](#)  
 Night Sweats, [1359](#)  
 Night Sweats in Men, [1360](#)  
 Night Terrors, [1360](#)  
 Night Terrors In Children, [1360](#)  
 Night-Blindness, [1359](#)  
 Night-Sweats, [1360](#)  
 Nightmare Disorder, [1359](#)  
 Nightmares, [1359](#)  
 NIH Stroke Scale, [1360](#)  
 Nijmegen Breakage Syndrome, [1360](#)  
 Nikethamide, [1360](#)  
 Nikolsky Sign, [1360](#)  
 Nikolsky's Sign, [1360](#)  
 Nipah Virus, [1360](#)  
 Nipah Virus Infection, [1360](#)

---

---

Nipple, [1360](#)  
 Nipple Discharge, [1361](#)  
 Nipple Inversion Or Retraction, [1361](#)  
 Nipple Retraction, [1361](#)  
 Nipple-Areola Complex, [1361](#)  
 Niridazole, [1361](#)  
 Nissen Fundoplication, [1361](#)  
 Nit, [1361](#)  
 Nitabuch Layer, [1361](#)  
 Nitrate Tolerance, [1361](#)  
 Nitrates, [1361](#)  
 Nitrazepam, [1361](#)  
 Nitric Acid, [1361](#)  
 Nitric Acid Poisoning, [1362](#)  
 Nitric Oxide, [1362](#)  
 Nitric Oxide Synthase, [1362](#)  
 Nitrofurantoin, [1362](#)  
 Nitrogen, [1362](#)  
 Nitrogen Balance, [1362](#)  
 Nitrogenous Base, [1362](#)  
 Nitroglycerin for Chest Pain, [1382](#)  
 Nitroglycerin Overdose, [1362](#)  
 Nitromors, [1362](#)  
 Nitrosoureas, [1362](#)  
 Nitrous Oxide, [1362](#)  
 Nitrous Oxide For Labor, [1362](#)  
 Nizatidine, [1362](#)  
 NMR, [1362](#)  
 No-Reflow Phenomenon, [1372](#)  
 Nocardia Asteroides Infection, [1362](#)  
 Nocardia Infection, [1363](#)  
 Nocardiosis, [1363](#)  
 Nocebo, [1363](#)  
 Nociception, [1363](#)  
 Nociceptor, [1363](#)  
 Nociceptors, [1363](#)  
 Nocturia, [1363](#)  
 Nocturnal, [1363](#)  
 Nocturnal Emission, [1363](#)  
 Nocturnal Enuresis, [1363](#)  
 Nocturnal Enuresis Entry, [1363](#)  
 Nocturnal Hypoventilation, [1363](#)  
 Nocturnal Incontinence, [1363](#)  
 Nocturnal Lagophthalmos, [1363](#)  
 Nocturnal Polyuria, [1364](#)  
 Nodal Rhythm, [1364](#)  
 Node, [1364](#)  
 Node of Ranvier, [1364](#)  
 Nodular, [1364](#)  
 Nodular Fasciitis, [1364](#)  
 Nodular Localized Cutaneous Amyloidosis, [1364](#)  
 Nodular Lymphoid Hyperplasia, [1364](#)  
 Nodular Melanoma, [1364](#)  
 Nodule, [1364](#)  
 Noise-Induced Hearing Loss, [1364](#)  
 Noise-Related Hearing Loss, [1364](#)  
 Noma, [1364](#)  
 Nomenclature, [1365](#)  
 Nomogram, [1365](#)  
 Non Communicating Hydrocephalus, [1366](#)  
 Non-Accidental Trauma, [1365](#)  
 Non-Adherence, [1365](#)  
 Non-Alcoholic Steatohepatitis, [1365](#)  
 Non-Allergic Rhinitis, [1365](#)  
 Non-Articular Rheumatism, [1365](#)  
 Non-Asthmatic Eosinophilic Bronchitis, [1366](#)  
 Non-Celiac Gluten Intolerance, [1366](#)  
 Non-Celiac Gluten Sensitivity, [1366](#)  
 Non-Coding RNA (ncRNA), [1366](#)  
 Non-Communicable Disease, [1366](#)  
 Non-Compliant Angioplasty Balloon, [1367](#)  
 Non-Contact Tonometry, [1367](#)  
 Non-Convulsive Status Epilepticus, [1367](#)  
 Non-Fatal Drowning, [1367](#)  
 Non-Functioning Pituitary Adenoma, [1367](#)  
 Non-Hodgkin Lymphoma, [1367](#)  
 Non-Hodgkin Lymphoma In Children, [1367](#)  
 Non-Hodgkin's Lymphoma, [1367](#)  
 Non-Hodgkins Lymphomas, [1367](#)  
 Non-Infectious Cystitis, [1367](#)  
 Non-Invasive Positive Pressure Ventilation, [1368](#)  
 Non-Invasive Prenatal Testing, [1368](#)  
 Non-Invasive Procedure, [1368](#)  
 Non-Invasive Ventilation, [1368](#)  
 Non-Ketotic Hyperosmolar State, [1368](#)  
 Non-Muscle-Invasive Bladder Cancer, [1369](#)  
 Non-Obstructive Azoospermia, [1369](#)  
 Non-Pitting Edema, [1369](#)  
 Non-Polio Enterovirus Infection, [1369](#)  
 Non-Proliferative Diabetic Retinopathy, [1369](#)  
 Non-Radiographic Axial Spondyloarthritis, [1369](#)  
 Non-Rapid Eye Movement Sleep, [1369](#)  
 Non-Rebreather Mask, [1369](#)  
 Non-REM Sleep, [1369](#)  
 Non-Responder, [1369](#)  
 Non-Scarring Epidermolysis Bullosa, [1382](#)  
 Non-Seminomatous Germ Cell Tumor, [1370](#)  
 Non-Small Cell Lung Cancer, [1370](#)  
 Non-Small Cell Lung Carcinoma, [1370](#)  
 Non-Specific Urethritis, [1370](#)  
 Non-ST Elevation Myocardial Infarction, [1370](#)  
 Non-ST-Elevation Myocardial Infarction, [1371](#)  
 Non-ST-Segment Elevation Myocardial Infarction, [1371](#)  
 Non-Stress Test, [1371](#)  
 Nonaccidental Injury, [1365](#)  
 Nonalcoholic Steatohepatitis Pathology, [1365](#)  
 Nonallergic Rhinitis, [1365](#)  
 Nonallergic Rhinopathy, [1365](#)

---

---

Nonarticular Rheumatism/Regional Pain Syndrome, [1365](#)  
 Nonbacterial Prostatitis, [1366](#)  
 Nonbacterial Thrombotic Endocarditis, [1366](#)  
 Noncandidal Fungal Infections of the Mouth, [1366](#)  
 Noncompetitive Antagonist, [1366](#)  
 Noncompetitive Inhibition, [1366](#)  
 Nonconvulsive Status Epilepticus, [1367](#)  
 Nondisjunction, [1367](#)  
 Nongenital Warts, [1367](#)  
 Noninfectious Cystitis, [1368](#)  
 Noninfectious Uveitis, [1382](#)  
 Noninvasive, [1368](#)  
 Nonischemic Priapism, [1368](#)  
 Nonketotic Hyperglycemic Hyperosmolar Syndrome, [1368](#)  
 Nonmelanoma Skin Cancer, [1368](#)  
 Nonpathogenic, [1369](#)  
 Nonproliferative Retinopathy, [1369](#)  
 Nonselective Beta-Blockers, [1370](#)  
 Nonseminomatous Germ Cell Tumor, [1370](#)  
 Nonseminomatous Testicular Tumors, [1370](#)  
 Nonsense Variant, [1370](#)  
 Nonspecific Interstitial Pneumonia, [1370](#)  
 Nonspecific Interstitial Pneumonitis, [1370](#)  
 Nonsteroidal Anti-Inflammatory Drug, [2126](#)  
 Nonsteroidal Anti-inflammatory Drug (NSAID) Toxicity, [1371](#)  
 Nonsteroidal Anti-Inflammatory Drugs, [1371](#)  
 Nonsteroidal Anti-Inflammatory Drugs And Ulcers, [1371](#)  
 Nonsyndromic, [1371](#)  
 Nonthrombocytopenic Idiopathic Purpura, [1382](#)  
 Nontraditional Medicine, [1382](#)  
 Nontropical Sprue, [1371](#)  
 Nontuberculous Mycobacteria, [1371](#)  
 Nontuberculous Mycobacterial (NTM) Diseases, [1371](#)  
 Nontuberculous Mycobacterial Infection, [1371](#)  
 Nontuberculous Mycobacterial Lung Disease, [1371](#)  
 Nonulcer Dyspepsia, [1372](#)  
 Nonulcer Stomach Pain, [1372](#)  
 Nonunion And Malunion, [1372](#)  
 Nonunion Fracture, [1372](#)  
 Noonan Syndrome, [1372](#)  
 Noonan Syndrome With Multiple Lentigines, [1372](#)  
 Noradrenaline, [1372](#)  
 Norepinephrine, [1372](#)  
 Norethisterone, [1372](#)  
 Norgestrel, [1372](#)  
 Norketamine, [1372](#)  
 Normal, [1372](#)  
 Normal and Abnormal Puerperium, [1373](#)  
 Normal Flora, [1373](#)  
 Normal Growth And Development, [1373](#)  
 Normal Labor and Delivery, [1373](#)  
 Normal Labour, [1373](#)  
 Normal Menstrual Cycle, [1382](#)  
 Normal Pressure Hydrocephalus, [1373](#)  
 Normal Range, [1373](#)  
 Normal Saline, [1373](#)  
 Normal-Pressure Hydrocephalus, [1373](#)  
 Normal-Tension Glaucoma, [1373](#)  
 Normalization, [1373](#)  
 Normoblast, [1373](#)  
 Normocalcemic Hyperparathyroidism, [1373](#)  
 Normocytic Anaemia, [1373](#)  
 Normocytic Anemia, [1373](#)  
 Normotensive, [1374](#)  
 Norovirus, [1374](#)  
 Norovirus Infection, [1374](#)  
 Noroviruses, [1382](#)  
 Norrbottnian Gaucher Disease, [1382](#)  
 Norrie Disease, [1374](#)  
 Nortriptyline, [1374](#)  
 Norwegian Scabies, [1374](#)  
 Norwood Procedure, [1374](#)  
 Nose, [1374](#)  
 Nose Fracture, [1375](#)  
 Nosebleed, [1375](#)  
 Nosebleeds, [1375](#)  
 Nosocomial, [1375](#)  
 Nosocomial Infection, [1375](#)  
 Nosocomial Norovirus Infection, [1375](#)  
 Nosocomial Vancomycin-Resistant Enterococci, [1375](#)  
 Nostril, [1375](#)  
 Notalgia Paresthetica, [1375](#)  
 Notencephalocele, [1375](#)  
 Notifiable Disease, [1375](#)  
 Notochord, [1375](#)  
 Novel Biomarkers of Kidney Function, [1375](#)  
 NOWS, [1376](#)  
 Noxious, [1376](#)  
 NPS, [1382](#)  
 NREM Sleep, [1376](#)  
 NS, [1382](#)  
 NSAID, [1376](#)  
 NSAID-Induced Ulcers, [1376](#)  
 NSAIDs, [2126](#)  
 NSTEMI, [1376](#)  
 NTD, [1376](#)  
 Nuchal, [1376](#)  
 Nuchal Cord, [1376](#)  
 Nuchal Rigidity, [1376](#)  
 Nuchal Translucency Screening, [1376](#)  
 Nuchal Translucency Test, [1376](#)  
 Nuclear, [1376](#)  
 Nuclear Factor Kappa B, [1376](#)  
 Nuclear Medicine, [1377](#)  
 Nuclear Medicine Imaging, [1377](#)

---

---

Nuclear Pore Complex, [1377](#)  
Nuclear Sclerosis, [1377](#)  
Nuclear Stress Test, [1377](#)  
Nuclear Ventriculography, [1377](#)  
Nucleic Acid, [1377](#)  
Nucleic Acid Hybridization, [1377](#)  
Nucleolus, [1377](#)  
Nucleoside, [1377](#)  
Nucleoside Analogs, [1377](#)  
Nucleotide, [1377](#)  
Nucleus, [1378](#)  
Nucleus Accumbens, [1378](#)  
Nucleus Ambiguus, [1378](#)  
Nucleus Pulposus, [1378](#)  
Null Mutation, [1378](#)  
Nulligravida, [1378](#)  
Nullipara, [1378](#)  
Numb, [1378](#)  
Number Needed To Harm, [1378](#)  
Number Needed To Treat, [1378](#)  
Numbness, [1378](#)  
Numbness And Tingling, [1378](#)  
Numbness In Hands, [1378](#)  
Nummular, [1378](#)  
Nummular Dermatitis, [1378](#)  
Nummular Eczema, [1379](#)  
Nurse Practitioner, [1379](#)  
Nursemaid's Elbow, [1379](#)  
Nursery, [1379](#)  
Nursing, [1379](#)  
Nursing Home, [1379](#)  
Nursing Home Acquired Pneumonia, [1379](#)  
Nut Allergies, [1379](#)  
Nut-Allergy, [1379](#)  
Nutation, [1379](#)  
Nutcracker Esophagus, [1379](#)  
Nutcracker Syndrome, [1379](#)  
Nutrient, [1379](#)  
Nutrigenomics, [1379](#)  
Nutrition, [1380](#)  
Nutrition In Pregnancy, [1380](#)  
Nutrition Risk Screening 2002, [1380](#)  
Nutrition Support, [1380](#)  
Nutrition-Focused Physical Examination, [1380](#)  
Nutritional Anemia, [1380](#)  
Nutritional Assessment, [1380](#)  
Nutritional Deficiency, [1380](#)  
Nutritional Dermatoses, [1380](#)  
Nutritional Disorders, [1380](#)  
Nutritional Rickets, [1382](#)  
Nutritional Screening, [1380](#)  
Nutritive, [1380](#)  
Nux Vomica, [1380](#)  
Nyctalopia, [1380](#)  
NYHA Functional Classification, [1381](#)  
Nystagmus, [1381](#)  
Nystagmus Benign Paroxysmal Positional, [1382](#)  
Nystagmus Retractorius, [1381](#)  
OAB, [1383](#)  
OB/GYN, [1383](#)  
Ober Test, [1383](#)  
Obesity, [1383](#)  
Obesity Hypoventilation Syndrome, [1383](#)  
Obex Region, [1383](#)  
Objective Response Rate, [1383](#)  
Objects Or Insects In Ear, [1383](#)  
Oblique Fracture, [1383](#)  
Oblique Plane, [1383](#)  
Obliteration, [1384](#)  
Observation, [1384](#)  
Obsession, [1384](#)  
Obsessions, [1384](#)  
Obsessive Compulsive Disorder, [1384](#)  
Obsessive-Compulsive And Related Disorders, [1384](#)  
Obsessive-Compulsive Disorder, [1384](#)  
Obsessive-Compulsive Personality Disorder, [1384](#)  
Obstetric Anal Sphincter Injury, [1384](#)  
Obstetric Cholestasis, [1384](#)  
Obstetric Epidural Anesthesia, [1384](#)  
Obstetric Fistula, [1384](#)  
Obstetric Forceps, [1385](#)  
Obstetric Nursing, [1385](#)  
Obstetric Ultrasound, [1385](#)  
Obstetrical Forceps, [1384](#)  
Obstetrician, [1385](#)  
Obstetrics, [1385](#)  
Obstipation, [1385](#)  
Obstructed Infected Urinary System, [1385](#)  
Obstructed Labour, [1385](#)  
Obstruction, [1385](#)  
Obstructive Azoospermia, [1385](#)  
Obstructive Hydrocephalus, [1385](#)  
Obstructive Jaundice, [1385](#)  
Obstructive Nephropathy, [1385](#)  
Obstructive Parotitis, [1386](#)  
Obstructive Shock, [1386](#)  
Obstructive Sleep Apnea, [1386](#)  
Obstructive Sleep Apnea Syndrome, [1386](#)  
Obstructive Uropathy, [1386](#)  
Obtunded, [1386](#)  
Obturator, [1386](#)  
Obturator Canal, [1386](#)  
Obturator Foramen, [1386](#)  
Obturator Hernia, [1386](#)  
Obturator Nerve, [1386](#)  
Obturator Sign, [1386](#)  
Obtuse Marginal Artery, [1387](#)

---

---

Occipital, 1387  
Occipital Bone, 1387  
Occipital Diastasis, 1387  
Occipital Hematoma, 1387  
Occipital Lobe, 1387  
Occipital Neuralgia, 1387  
Occiput, 1387  
Occiput Posterior Position, 1387  
Occiput Transverse Position, 1387  
Occlude, 1387  
Occlusal Adjustment, 1387  
Occlusal Guard, 1387  
Occlusal Splint, 1387  
Occlusion, 1387  
Occlusive Vascular Disease, 1388  
Occult, 1388  
Occult Blood, 1388  
Occult Blood Screen, 1388  
Occult Gastrointestinal Bleeding, 1388  
Occult Primary Cancer, 1388  
Occupational Asthma, 1388  
Occupational COPD, 1388  
Occupational Dermatoses, 1388  
Occupational Disease, 1388  
Occupational Exposure, 1388  
Occupational Exposure Limit, 1388  
Occupational Health, 1388  
Occupational Hearing Loss, 1388  
Occupational Lung Disease, 1389  
Occupational Medicine, 2126  
Occupational Rehabilitation, 1389  
Occupational Safety And Health, 1389  
Occupational Therapist, 1389  
Occupational Therapy, 1389  
Occupational Therapy Assessment, 1389  
OCD, 1389  
Ochre, 1389  
Ochronosis, 1389  
OCP, 1389  
Octreoscan, 1389  
Octreotide, 1389  
Ocular, 1389  
Ocular Albinism, 1390  
Ocular Bobbing, 1390  
Ocular Burns and Chemical Injuries, 1390  
Ocular Cicatricial Pemphigoid, 1390  
Ocular Fixation, 1390  
Ocular Flutter, 1390  
Ocular Foreign Body, 1390  
Ocular Histoplasmosis Syndrome, 1390  
Ocular Hypertension, 1390  
Ocular Hypotony, 1390  
Ocular Ischemic Syndrome, 1390  
Ocular Larva Migrans, 1390  
Ocular Melanoma, 1391  
Ocular Migraine, 1391  
Ocular Motility, 1391  
Ocular Myasthenia Gravis, 1391  
Ocular Myiasis, 1391  
Ocular Orbit, 1391  
Ocular Pain, 1391  
Ocular Rosacea, 1391  
Ocular Surface Disease, 1391  
Ocular Toxoplasmosis, 1391  
Oculocardiac Reflex, 1391  
Oculocephalic Reflex, 1391  
Oculocerebrorenal Syndrome, 1391  
Oculocutaneous Albinism, 1391  
Oculogyric Crisis, 1391  
Oculomotor Nerve, 1392  
Oculomotor Nerve Palsy, 1392  
Oculomotor Nucleus, 1392  
Oculopharyngeal Muscular Dystrophy, 1392  
Oculoplastic Procedures, 1392  
Oculosympathetic Palsy, 1392  
Oculovestibular Reflex, 1392  
ODD, 1392  
Odontalgia, 1392  
Odontoblast, 1392  
Odontoclast, 1392  
Odontodysplasia, 1392  
Odontogenesis, 1392  
Odontogenic Cutaneous Fistula, 1392  
Odontogenic Cyst, 1393  
Odontogenic Fibroma, 1393  
Odontogenic Infection, 1393  
Odontogenic Keratocyst, 1393  
Odontogenic Keratocyst Pathology, 1393  
Odontogenic Neoplasm, 1393  
Odontogenic Tumors and Cysts, 1393  
Odontoid Process, 1393  
Odontoma, 1393  
Odontopathy, 1393  
Odynophagia, 1393  
Oedema, 1393  
Oesophageal, 1393  
Oesophageal Atresia, 1393  
Oesophageal Ulcer, 1393  
Oesophageal Varices, 1394  
Oesophagitis, 1394  
Oesophagoscopy, 1394  
Oesophagus, 1394  
Oestradiol, 1394  
Oestriol, 1394  
Oestrogen, 1394  
Oestrogen Replacement Therapy, 1394  
Oestrogens, 1394  
Oestrone, 1394

---

---

Ofatumumab, [1394](#)  
Off-Label, [1394](#)  
Off-Label Use, [1394](#)  
Offspring, [1394](#)  
Ofloxacin, [1394](#)  
Ogilvie Syndrome, [1395](#)  
Ogilvie'S Syndrome, [1395](#)  
Ohm, [1395](#)  
OHSS, [1395](#)  
Oidium, [1395](#)  
Oil, [1395](#)  
Oily Hair, [1395](#)  
Oily Skin, [1395](#)  
Ointment, [1395](#)  
Ointment Bases, [1395](#)  
Okadaic Acid, [1395](#)  
Okiihiro Syndrome, [1395](#)  
Olanzapine, [1395](#)  
Olaparib, [1395](#)  
Old Age, [1396](#)  
Oleander Poisoning, [1396](#)  
Olecranon, [1396](#)  
Olecranon Bursitis, [1396](#)  
Olecranon Fossa, [1396](#)  
Oleic Acid, [1396](#)  
Oleum, [1396](#)  
Olfaction, [1396](#)  
Olfactory, [1396](#)  
Olfactory Bulb, [1396](#)  
Olfactory Cortex, [1396](#)  
Olfactory Gyrus, [1396](#)  
Olfactory Learning, [1396](#)  
Olfactory Mucosa, [1396](#)  
Olfactory Nerve, [1396](#)  
Olfactory Neuroblastoma, [1396](#)  
Olfactory Tract, [1397](#)  
Olfactory Tubercle, [1397](#)  
Oligoasthenoteratozoospermia, [1397](#)  
Oligoclonal Bands, [1397](#)  
Oligocythaemia, [1397](#)  
Oligodendrocyte, [1397](#)  
Oligodendroglia, [1397](#)  
Oligodendroglial Neoplasm, [1397](#)  
Oligodendroglioma, [1397](#)  
Oligodontia, [1397](#)  
Oligogenic Inheritance, [1397](#)  
Oligohydramnios, [1397](#)  
Oligomenorrhea, [1397](#)  
Oligomenorrhoea, [1397](#)  
Oligometastasis, [1398](#)  
Oligonucleotide, [1398](#)  
Oligopeptide, [1398](#)  
Oligosaccharide, [1398](#)  
Oligospermia, [1398](#)  
Oliguria, [1398](#)  
Olivary, [1398](#)  
Olivary Nucleus, [1398](#)  
Olive Pontocerebellar Atrophy, [1398](#)  
Olivopontocerebellar Atrophy, [1398](#)  
Ollier Disease, [1398](#)  
Ollier's Disease, [1398](#)  
Olmesartan Medoxomil, [1398](#)  
Olopatadine Hydrochloride, [1398](#)  
OLP, [1399](#)  
Olsenella, [1399](#)  
Olsenella Uli, [1399](#)  
Omacetaxine Mepesuccinate, [1399](#)  
Omalizumab, [1399](#)  
Omasum, [1399](#)  
Omega Sign, [1399](#)  
Omega-3, [1399](#)  
Omega-3 Fatty Acid, [1399](#)  
Omega-3 Fatty Acids, [1399](#)  
Omega-6, [1399](#)  
Omega-6 Fatty Acids, [1399](#)  
Omega-N-Methylarginine, [1399](#)  
Omenn Syndrome, [1399](#)  
Omental Caking, [1400](#)  
Omentectomy, [1400](#)  
Omentum, [1400](#)  
Omeprazole, [1400](#)  
Ommaya Reservoir, [1400](#)  
Omphalitis, [1400](#)  
Omphalocele, [1400](#)  
Omphalocele Repair, [1400](#)  
Onchocerca Volvulus, [1400](#)  
Onchocerciasis, [1400](#)  
Oncocytic Neoplasm, [1400](#)  
Oncocytoma, [1400](#)  
Oncogene, [1400](#)  
Oncogenic, [1401](#)  
Oncogenic Osteomalacia, [1401](#)  
Oncogenic Virus, [1401](#)  
Oncogenicity, [1401](#)  
Oncologic Emergency, [1401](#)  
Oncologist, [1401](#)  
Oncology, [1401](#)  
Oncology Nursing, [1401](#)  
Oncology Nutrition, [1401](#)  
Oncology PET, [1401](#)  
Oncology Rehabilitation, [1401](#)  
Oncolytic Virotherapy, [1401](#)  
Oncolytic Virus, [1401](#)  
Oncolytic Viruses, [1402](#)  
Oncoprotein, [1402](#)  
Oncosis, [1402](#)  
Oncotic Pressure, [1402](#)  
Oncotoxic, [1402](#)

---



---

Ondansetron, [1402](#)  
Ondansetron Hydrochloride, [1402](#)  
Ondine Curse, [1402](#)  
Ondine'S Curse, [1402](#)  
One-Hour Glucose Challenge Test, [1402](#)  
Oneirophrenia, [1402](#)  
Onset, [1402](#)  
Onychia, [1402](#)  
Onychocryptosis, [1402](#)  
Onychogryphosis, [1403](#)  
Onycholysis, [1403](#)  
Onychomadesis, [1403](#)  
Onychomycosis, [1403](#)  
Oocyte, [1403](#)  
Oocyte Cryopreservation, [1403](#)  
Oocyte Retrieval, [1403](#)  
Oogenesis, [1403](#)  
Oogonium, [1403](#)  
Oophorectomy, [1403](#)  
Oophoritis, [1403](#)  
Ooplasm, [1403](#)  
Opacity, [1403](#)  
Open Abdominal Aortic Aneurysm Repair, [1404](#)  
Open Aortic Valve Surgery, [1404](#)  
Open Bite, [1404](#)  
Open Esophagectomy, [1404](#)  
Open Fracture, [1404](#)  
Open Fractures, [1404](#)  
Open Gallbladder Removal, [1404](#)  
Open Lung Biopsy, [1404](#)  
Open Mitral Valve Surgery, [1405](#)  
Open Pleural Biopsy, [1405](#)  
Open Prostatectomy, [1405](#)  
Open Reduction And Internal Fixation, [1405](#)  
Open Surgery, [1405](#)  
Open Surgical Wound Care, [1405](#)  
Open-Angle Glaucoma, [1404](#)  
Open-Book Fracture, [1404](#)  
Open-Heart Surgery, [1404](#)  
Opening Snap, [1404](#)  
Operable, [1405](#)  
Operant Conditioning, [1405](#)  
Operating Room, [1405](#)  
Operating Tables, [1405](#)  
Operation, [1405](#)  
Operative Dentistry, [1405](#)  
Operative Vaginal Delivery, [1405](#)  
Operator, [1405](#)  
Operculum, [1406](#)  
Ophthalmia, [1406](#)  
Ophthalmia Neonatorum, [1406](#)  
Ophthalmic, [1406](#)  
Ophthalmic Emergency, [1406](#)  
Ophthalmic Solution, [1406](#)  
Ophthalmitis, [1406](#)  
Ophthalmologist, [1406](#)  
Ophthalmology, [1406](#)  
Ophthalmomyiasis, [1406](#)  
Ophthalmoplegia, [1406](#)  
Ophthalmoscope, [1406](#)  
Ophthalmoscopy, [1406](#)  
Opiate, [1407](#)  
Opiate Alkaloid, [1407](#)  
Opiate And Opioid Withdrawal, [1407](#)  
Opiate Assay, [1407](#)  
Opiate Overdose, [1407](#)  
Opiate Overdose Triad, [1407](#)  
Opioid, [1407](#)  
Opioid Addiction, [1443](#)  
Opioid Agonist Treatment, [1407](#)  
Opioid Analgesic Medications, [1407](#)  
Opioid Analgesics, [2126](#)  
Opioid Antagonist, [2126](#)  
Opioid Dependence, [1442](#)  
Opioid Intoxication, [1407](#)  
Opioid Overdose, [1407](#)  
Opioid Poisoning, [1407](#)  
Opioid Receptor, [1407](#)  
Opioid Reversal, [1407](#)  
Opioid Toxicity, [1408](#)  
Opioid Toxidrome, [1408](#)  
Opioid Use Disorder, [1408](#)  
Opioid Withdrawal, [1408](#)  
Opioids, [1408](#)  
Opisthorchis Felineus, [1408](#)  
Opisthotonos, [1408](#)  
Opisthotonus, [1408](#)  
Opium, [1408](#)  
Oppenheim Sign, [1408](#)  
Opponens, [1409](#)  
Opportunistic Infection, [1409](#)  
Opportunistic Pathogen, [1409](#)  
Opposition, [1409](#)  
Oppositional Defiant Disorder, [1409](#)  
Opsoclonus, [1409](#)  
Opsoclonus-Myoclonus Syndrome, [1409](#)  
Opsoclonus-Myoclonus-Ataxia Syndrome, [1409](#)  
Opsonin, [1409](#)  
Opsonization, [1409](#)  
Optic Atrophy, [1410](#)  
Optic Chiasm, [1410](#)  
Optic Chiasma, [1410](#)  
Optic Disc, [1410](#)  
Optic Disc Drusen, [1410](#)  
Optic Disc Edema, [1410](#)  
Optic Disc Swelling, [1410](#)  
Optic Disk, [1410](#)  
Optic Glioma, [1410](#)

---

---

Optic Nerve, [1410](#)  
Optic Nerve Atrophy, [1411](#)  
Optic Nerve Disorder, [1411](#)  
Optic Nerve Glioma, [1411](#)  
Optic Nerve Swelling, [1443](#)  
Optic Neuritis, [1411](#)  
Optic Pathway Glioma, [1411](#)  
Optic Tract, [1411](#)  
Optical Aids, [707](#)  
Optical Coherence Tomography, [1409](#)  
Optical Coherence Tomography Angiography, [1409](#)  
Optical Illusion, [1410](#)  
Optical Rotation, [1410](#)  
Optician, [1410](#)  
Optokinetic Nystagmus, [1411](#)  
Optometrist, [1411](#)  
Optometry, [1411](#)  
Oral, [1411](#)  
Oral Administration, [1411](#)  
Oral Allergy Syndrome, [1442](#)  
Oral Antidiabetic Agents, [1411](#)  
Oral Appliance Therapy, [1411](#)  
Oral Burn, [1411](#)  
Oral Cancer, [1411](#)  
Oral Candidiasis, [1411](#)  
Oral Cavity, [1412](#)  
Oral Cavity Cancer, [1412](#)  
Oral Cholecystogram, [1412](#)  
Oral Contraceptive, [1412](#)  
Oral Contraceptive Pill, [1412](#)  
Oral Cutaneous Fistulas, [1412](#)  
Oral Fibromas and Fibromatoses, [1412](#)  
Oral Glucose Tolerance Test, [1412](#)  
Oral Hairy Leukoplakia, [1412](#)  
Oral Health, [1412](#)  
Oral Hemangiomas, [1412](#)  
Oral Hemorrhage, [1412](#)  
Oral Herpes, [1413](#)  
Oral Human Papillomavirus Infection, [1413](#)  
Oral Hygiene, [1413](#)  
Oral Hypoglycemic, [2126](#)  
Oral Hypoglycemics Overdose, [1413](#)  
Oral Leukoplakia, [1413](#)  
Oral Leukoplakia and Erythroplakia Pathology, [1413](#)  
Oral Lichen Planus, [1413](#)  
Oral Medications for Type 2 Diabetes, [1413](#)  
Oral Medicine, [2126](#)  
Oral Moniliasis, [1413](#)  
Oral Mucosa, [1413](#)  
Oral Mucous Cyst, [1413](#)  
Oral Nutritional Supplement, [1413](#)  
Oral Pathology, [1413](#)  
Oral Polio Vaccine, [2126](#)  
Oral Potentially Malignant Disorders, [1414](#)  
Oral Pyogenic Granuloma, [1414](#)  
Oral Radiology, [1414](#)  
Oral Rehydration Salts, [1414](#)  
Oral Rehydration Therapy, [1414](#)  
Oral Route, [1414](#)  
Oral Sex Practices, [1414](#)  
Oral Spray, [1414](#)  
Oral Squamous Cell Carcinoma, [1414](#)  
Oral Submucous Fibrosis, [1414](#)  
Oral Surgery, [1414](#)  
Oral Thrush, [1414](#)  
Oral Toxicity, [1415](#)  
Oral Yeast Infection, [1415](#)  
Oral-Facial-Digital Syndrome, [1412](#)  
Orange Peel Skin, [1415](#)  
Orbicularis, [1415](#)  
Orbit, [1415](#)  
Orbit CT Scan, [1416](#)  
Orbital, [1415](#)  
Orbital Apex Syndrome, [1415](#)  
Orbital Blowout Fracture, [1415](#)  
Orbital Cavity, [1415](#)  
Orbital Cellulitis, [1415](#)  
Orbital Contusion, [1415](#)  
Orbital Floor Fracture, [1415](#)  
Orbital Fracture, [1415](#)  
Orbital Fractures, [1416](#)  
Orbital Irradiation, [1416](#)  
Orbital Metastasis, [1416](#)  
Orbital Myositis, [1416](#)  
Orbital Neoplasm, [1416](#)  
Orbital Pseudotumor, [1416](#)  
Orbital Tumors, [1416](#)  
Orbital Varix, [1416](#)  
Orchialgia, [1416](#)  
Orchidectomy, [1416](#)  
Orchiectomy, [1416](#)  
Orchitis, [1416](#)  
Order Coleoptera, [1416](#)  
Order Spirochaetales, [1417](#)  
Orexin, [1417](#)  
Orexin Receptor Antagonists, [1417](#)  
Orf, [1417](#)  
Organ, [1417](#)  
Organ Failure, [1417](#)  
Organ of Corti, [1417](#)  
Organ System, [1418](#)  
Organ Transplant, [1418](#)  
Organ Transplantation, [1418](#)  
Organ Transplants, [1418](#)  
Organelle, [1417](#)  
Organic, [1417](#)  
Organic Chemical, [1417](#)  
Organic Disease, [1417](#)

---

---

Organic Impurity, [1417](#)  
Organization Name, [1417](#)  
Organogenesis, [1417](#)  
Organophosphate, [1418](#)  
Organophosphate Poisoning, [1418](#)  
Organophosphate Toxicity, [1418](#)  
Organosilicon Compounds, [1418](#)  
Orgasm, [1418](#)  
Orgasmic Dysfunction, [1418](#)  
Orgasmic Dysfunction in Women, [1418](#)  
Orientation, [1418](#)  
Orifice, [1418](#)  
Orlistat, [1418](#)  
Ornithine, [1418](#)  
Ornithine Decarboxylase, [1419](#)  
Ornithine Transcarbamylase Deficiency, [1419](#)  
Orogastric Tube, [1419](#)  
Orolabial Herpes Simplex, [1419](#)  
Oropharyngeal, [1419](#)  
Oropharyngeal Airway, [1419](#)  
Oropharyngeal Cancer, [1419](#)  
Oropharyngeal Carcinoma, [1419](#)  
Oropharyngeal Dysphagia, [1419](#)  
Oropharyngeal Neoplasm, [1419](#)  
Oropharynx, [1419](#)  
Oropharynx Lesion Biopsy, [1419](#)  
Oropouche Fever, [1419](#)  
Orotic Aciduria, [1419](#)  
Orphan Disease, [1420](#)  
Orphan Drug, [2126](#)  
Orphenadrine, [1420](#)  
Orthodontic, [1420](#)  
Orthodontic Appliance, [1420](#)  
Orthodontic Bracket, [1420](#)  
Orthodontic Retainer, [1420](#)  
Orthodontic Treatment, [1420](#)  
Orthodontic Wires, [1420](#)  
Orthodontics, [1420](#)  
Orthodontist, [1420](#)  
Orthodromic AVRT, [1420](#)  
Orthognathic Surgery, [1420](#)  
Orthohepadnavirus, [1420](#)  
Orthokeratosis, [1420](#)  
Orthomyxoviridae, [1421](#)  
Orthopaedic Surgery, [1421](#)  
Orthopaedics, [1421](#)  
Orthopedic Biologics, [1421](#)  
Orthopedic Implant, [1421](#)  
Orthopedic Infections, [1421](#)  
Orthopedic Rehabilitation, [1421](#)  
Orthopedic Screws, [1421](#)  
Orthopedic Services, [1421](#)  
Orthopedics, [1421](#)  
Orthopedist, [1421](#)  
Orthopnea, [1421](#)  
Orthopneic Position, [1421](#)  
Orthopnoea, [1421](#)  
Orthopoxvirus, [1422](#)  
Orthoptera, [1422](#)  
Orthoptic, [1422](#)  
Orthoptist, [1422](#)  
Orthoreovirus, [1422](#)  
Orthorexia, [1422](#)  
Orthorexia Nervosa, [1422](#)  
Orthoscopic, [1422](#)  
Orthosis, [1422](#)  
Orthostatic Hypotension, [1422](#)  
Orthostatic Hypotension and Autonomic Failure, [1422](#)  
Orthostatic Intolerance, [1422](#)  
Orthotic, [1422](#)  
Orthotic Devices, [1422](#)  
Orthotic Training, [1423](#)  
Ortolani Maneuver, [1423](#)  
Ortolani Sign, [1423](#)  
Ortolani's Sign, [1423](#)  
Ortolani's Test, [1423](#)  
Os, [1423](#)  
Os Odontoideum, [1425](#)  
Osborn Wave, [1423](#)  
Oscillation, [1423](#)  
Oscillator, [1423](#)  
Oscillograph, [1423](#)  
Oscillometry, [1423](#)  
Oscillopsia, [1423](#)  
Oscitation, [1423](#)  
Oseltamivir, [1423](#)  
Oseltamivir Phosphate, [1423](#)  
OSFED, [1423](#)  
Osgood-Schlatter Disease, [1424](#)  
Osler Nodes, [1424](#)  
Osler Triad, [1424](#)  
Osler-Weber-Rendu Disease, [1424](#)  
Osmidrosis, [1424](#)  
Osmium, [1424](#)  
Osmolal Gap, [1424](#)  
Osmolality, [1424](#)  
Osmolality Blood Test, [1424](#)  
Osmolality Urine Test, [1424](#)  
Osmolar, [1424](#)  
Osmolarity, [1424](#)  
Osmole, [1424](#)  
Osmoreceptor, [1425](#)  
Osmoregulation, [1425](#)  
Osmosis, [1425](#)  
Osmotic Demyelination Syndrome, [1425](#)  
Osmotic Diarrhea, [1425](#)  
Osmotic Diuresis, [1425](#)  
Osmotic Fragility Test, [1425](#)

---

---

Osmostic Pressure, [1425](#)  
Osseointegration, [1425](#)  
Osseous, [1425](#)  
Osseous Metaplasia, [1425](#)  
Ossicle, [1425](#)  
Ossicles, [1425](#)  
Ossicular Discontinuity, [1425](#)  
Ossicular Fusion, [1426](#)  
Ossicular Prosthesis, [1426](#)  
Ossification, [1426](#)  
Ossifying Fibroma, [1426](#)  
Osteitis, [1426](#)  
Osteitis Deformans, [1426](#)  
Osteitis Fibrosa, [1426](#)  
Osteitis Fibrosa Cystica, [1426](#)  
Osteitis Pubis, [1426](#)  
Osteoarthritis, [1426](#)  
Osteoarthropathy, [1426](#)  
Osteoarthrosis, [1426](#)  
Osteoblast, [1426](#)  
Osteoblastoma, [1426](#)  
Osteoblasts, [1427](#)  
Osteocalcin, [1427](#)  
Osteocalcin Assay, [1427](#)  
Osteochondritis, [1427](#)  
Osteochondritis Dissecans, [1427](#)  
Osteochondroma, [1427](#)  
Osteochondromatosis, [1427](#)  
Osteochondrosis, [1427](#)  
Osteoclasia, [1427](#)  
Osteoclast, [1427](#)  
Osteoclastoma, [1427](#)  
Osteoclasts, [1427](#)  
Osteocyte, [1427](#)  
Osteocytes, [1427](#)  
Osteodystrophy, [1428](#)  
Osteogenesis, [1428](#)  
Osteogenesis Imperfecta, [1428](#)  
Osteogenic Sarcoma, [1428](#)  
Osteoid, [1428](#)  
Osteoid Osteoma, [1428](#)  
Osteology, [1428](#)  
Osteolysis, [1428](#)  
Osteolytic, [1428](#)  
Osteoma, [1428](#)  
Osteoma Cutis, [1428](#)  
Osteomalacia, [1428](#)  
Osteomyelitis, [1428](#)  
Osteomyelitis In Children, [1428](#)  
Osteon, [1429](#)  
Osteonecrosis, [1429](#)  
Osteonecrosis Of The Jaw, [1429](#)  
Osteonectin, [1429](#)  
Osteons, [1429](#)  
Osteopath, [1429](#)  
Osteopathy, [1429](#)  
Osteopenia, [1429](#)  
Osteopetrosis, [1429](#)  
Osteophyte, [1429](#)  
Osteoplasty, [1429](#)  
Osteopoikilosis, [1429](#)  
Osteopontin, [1429](#)  
Osteoporosis, [1430](#)  
Osteoporosis Fracture Risk Assessment, [1430](#)  
Osteoporotic Fracture, [1430](#)  
Osteoradionecrosis, [1430](#)  
Osteosarcoma, [1430](#)  
Osteosclerosis, [1430](#)  
Osteosynthesis, [1430](#)  
Osteotome, [1430](#)  
Osteotomy, [1430](#)  
Osteotomy Of The Knee, [1430](#)  
Ostium, [1430](#)  
Ostomy, [1430](#)  
Ostomy Appliance, [1430](#)  
Ostraceous Psoriasis, [1431](#)  
Otagia, [1431](#)  
Othello Syndrome, [1431](#)  
Other Pain Conditions, [1431](#)  
Other Race, [1431](#)  
Otic Barotrauma, [1431](#)  
Otic Capsule, [1431](#)  
Otic Ointment, [1431](#)  
Otitis, [1431](#)  
Otitis Externa, [1431](#)  
Otitis Media, [1431](#)  
Otitis Media With Effusion, [1431](#)  
Otoacoustic Emissions, [1432](#)  
Otocephaly, [1432](#)  
Otolaryngologist, [1432](#)  
Otolaryngology, [1432](#)  
Otolith, [1432](#)  
Otolithiasis, [1432](#)  
Otoliths, [1432](#)  
Otology, [1432](#)  
Otomycosis, [1432](#)  
Otoplasty, [1432](#)  
Otorhinolaryngology, [1432](#)  
Otorrhea, [1432](#)  
Otorrhoea, [1432](#)  
Otosalpingitis, [1433](#)  
Otosclerosis, [1433](#)  
Otoscope, [1433](#)  
Otoscopy, [1433](#)  
Ototoxicity, [1433](#)  
Ottawa Ankle Rules, [1433](#)  
Ounce, [1433](#)  
Outcome, [1433](#)

---

---

Outcome Measurement, 1433  
Outer Ear, 1433  
Outer Ear Infection, 1433  
Outpatient, 1433  
Outpatient Rehabilitation, 1433  
Output, 1433  
Ova, 1433  
Ovalocyte Count, 1434  
Ovarian, 1434  
Ovarian (Adnexal) Torsion, 1434  
Ovarian Cancer, 1434  
Ovarian Carcinoid, 1434  
Ovarian Cyst, 1434  
Ovarian Cyst Rupture, 1434  
Ovarian Cysts, 1434  
Ovarian Dysgerminomas, 1434  
Ovarian Failure, 1434  
Ovarian Fibromata, 1434  
Ovarian Follicle, 1434  
Ovarian Gonadoblastoma, 1434  
Ovarian Hemorrhage, 1435  
Ovarian Hyperstimulation Syndrome, 1435  
Ovarian Infection, 1435  
Ovarian Insufficiency, 1435  
Ovarian Metastasis, 1435  
Ovarian Neuroendocrine Tumor, 1435  
Ovarian Overproduction Of Androgens, 1435  
Ovarian Pregnancy, 1435  
Ovarian Reserve Testing, 1435  
Ovarian Rupture, 1435  
Ovarian Teratoma, 1435  
Ovarian Thecoma, 1435  
Ovarian Torsion, 1435  
Ovaries, 1436  
Ovariectomy, 1436  
Ovary, 1436  
Ovary Disease, 1436  
Ovary Neoplasm, 1436  
Over-Hydration, 1437  
Over-Nutrition, 1437  
Over-The-Counter, 1437  
Over-The-Counter (OTC) Drug, 2126  
Over-The-Counter Drug, 2126  
Overactive Bladder, 1436  
Overactive Bladder Syndrome, 1436  
Overall Survival, 1436  
Overbite, 1436  
Overdenture, 1436  
Overdose, 1436  
Overflow Incontinence, 1436  
Overflow Urinary Incontinence, 1437  
Overgrowth Syndrome, 1437  
Overlap Myositis, 1437  
Overlapping Autoimmune Syndrome, 1437  
Overlay Denture, 1437  
Overriding Aorta, 1437  
Oversedate, 1437  
Overt Proteinuria, 1437  
Overuse Syndrome, 1437  
Overweight, 1437  
Oviduct, 1437  
Ovomucin, 1437  
Ovotesticular Disorder of Sex Development, 1970  
Ovotesticular Disorder of Sexual Development, 1438  
Ovotestis, 1438  
Ovulation, 1438  
Ovulation Home Test, 1438  
Ovulation Induction, 1438  
Ovum, 1438  
Oxacillin, 1438  
Oxalate, 1438  
Oxalate Nephropathy, 1438  
Oxalates, 1438  
Oxalic Acid, 1438  
Oxalic Acid Poisoning, 1438  
Oxaliplatin, 1438  
Oxaloacetate, 1438  
Oxandrolone, 1438  
Oxaprozin, 1439  
Oxazepam, 1439  
Oxazepam Overdose, 1439  
Oxazole, 1439  
Oxcarbazepine, 1439  
Oxidant, 1439  
Oxidants, 1439  
Oxidation, 1439  
Oxidation-Reduction, 1439  
Oxidative Phosphorylation, 1439  
Oxidative Stress, 1439  
Oxidoreductase, 1439  
Oxime, 1439  
Oximeter, 1439  
Oximetry, 1439  
Oxter, 1440  
Oxybuprocaine, 1440  
Oxycephaly, 1440  
Oxycodone, 1440  
Oxycodone Hydrochloride, 1440  
Oxygen, 1440  
Oxygen Consumption, 1440  
Oxygen Content, 1440  
Oxygen Delivery, 1440  
Oxygen Desaturation Index, 1440  
Oxygen Dissociation Curve, 1440  
Oxygen Enhancement Ratio, 1441  
Oxygen Mask, 1441  
Oxygen Pulse, 1441  
Oxygen Safety, 1441

---

---

Oxygen Saturation, [1441](#)  
Oxygen Tent, [1441](#)  
Oxygen Therapy, [1441](#)  
Oxygen Therapy In Infants, [1441](#)  
Oxygen Toxicity, [1441](#)  
Oxygen Transport, [1441](#)  
Oxygen-Hemoglobin Dissociation Curve, [1441](#)  
Oxygenases, [1440](#)  
Oxygenated Blood, [1440](#)  
Oxygenation, [1440](#)  
Oxygenator, [1440](#)  
Oxyhaemoglobin, [1441](#)  
Oxyhemoglobin, [1441](#)  
Oxylipins, [1441](#)  
Oxymetazoline, [1441](#)  
Oxymetholone, [1442](#)  
Oxymorphone, [1442](#)  
Oxyphil Cell, [1442](#)  
Oxyphil Cell Carcinoma, [1442](#)  
Oxyphilic Adenoma, [1442](#)  
Oxytocic, [1442](#)  
Oxytocics, [1442](#)  
Oxytocin, [1442](#)  
Oxytocin Infusion, [1442](#)  
Oxyuriasis, [1442](#)  
Ozaena, [1442](#)  
Ozena, [1442](#)  
Ozone, [1442](#)

P Arm Of A Chromosome, [1466](#)  
P Wave, [1639](#)  
P-Glycoprotein, [1515](#)  
P2Y12 Inhibitor, [2126](#)  
P2Y12 Inhibitors, [1445](#)  
P53 Tumor Suppressor, [1445](#)  
PA X-Ray, [1476](#)  
PA/IVS, [1448](#)  
Pacemaker, [1445](#)  
Pacemaker Complications, [1445](#)  
Pacemaker Implantation, [1445](#)  
Pacemaker Malfunction, [1445](#)  
Pacemaker Syndrome, [1445](#)  
Pacemaker-Mediated Tachycardia, [1445](#)  
Pacemakers and Implantable Cardioverter-Defibrillators, [1445](#)  
Pachyderma, [1446](#)  
Pachydermoperiostosis, [1446](#)  
Pachygyria, [1446](#)  
Pachymeningitis, [1446](#)  
Pachymetry, [1446](#)  
Pachyonychia Congenita, [1446](#)  
Pacing, [1446](#)  
Pacinian Corpuscle, [1446](#)  
Paclitaxel, [1446](#)  
Paclitaxel Liposome, [1446](#)  
PACS, [1446](#)  
PAES, [1446](#)  
Page Kidney, [1446](#)  
Paget Disease Of Bone, [1446](#)  
Paget Disease Of The Anus, [1447](#)  
Paget Disease Of The Nipple, [1447](#)  
Paget Disease Of The Vulva, [1447](#)  
Paget's Disease, [1447](#)  
Paget's Disease of the Nipple, [1447](#)  
Paget's Disease of the Vulva, [1447](#)  
Paget-Schroetter Syndrome, [1447](#)  
Pagetoid Reticulosis, [1447](#)  
Pain, [1447](#)  
Pain Assessment Scales, [1447](#)  
Pain Management, [1448](#)  
Pain Medicine, [1448](#)  
Pain Perception, [1448](#)  
Pain Threshold, [1448](#)  
Pain Unit, [1448](#)  
Painful Bladder Syndrome, [1447](#)  
Painful Intercourse, [1448](#)  
Painful Menstrual Periods, [1447](#)  
Painful Swallowing, [1447](#)  
Painsomnia, [1448](#)  
Palatal Muscle, [1448](#)  
Palatal Myoclonus, [1448](#)  
Palate, [1448](#)  
Palate Neoplasm, [1448](#)  
Palatine, [1448](#)  
Palatine Tonsil, [1448](#)  
Palatorrhaphy, [1449](#)  
Palatoschisis, [1449](#)  
Pale or Clay-Colored Stools, [1449](#)  
Paleness, [1449](#)  
Paleo Diet, [1449](#)  
Palifermin, [2127](#)  
Palindromic Rheumatism, [1449](#)  
Paliperidone, [1449](#)  
Palivizumab, [1449](#)  
Palla Sign, [1449](#)  
Pallesthesia, [1449](#)  
Palliative Care, [1449](#)  
Palliative Radiation, [1449](#)  
Palliative Rehabilitation, [1449](#)  
Palliative Surgery, [1450](#)  
Palliative Treatment, [1450](#)  
Pallidotomy, [1450](#)  
Pallister-Hall Syndrome, [1450](#)  
Pallor, [1450](#)  
Palm, [1450](#)  
PALM-COEIN Classification, [1450](#)  
Palmar, [1450](#)  
Palmar Erythema, [1450](#)

---



---

Palmar Erythema Of Pregnancy, [1450](#)  
Palmar Grasp Reflex, [1450](#)  
Palmar Hyperhidrosis, [1450](#)  
Palmar Reflex, [1450](#)  
Palmitate, [1450](#)  
Palmitic Acid, [1450](#)  
Palmoplantar Hyperkeratosis, [1451](#)  
Palmoplantar Keratoderma (Keratosis Palmaris et Plantaris), [1451](#)  
Palonosetron, [2127](#)  
Palonosetron Hydrochloride, [1451](#)  
Palpable, [1451](#)  
Palpable Purpura, [1451](#)  
Palpate, [1451](#)  
Palpation, [1451](#)  
Palpebra, [1451](#)  
Palpebral Conjunctiva, [1451](#)  
Palpebral Slant, [1451](#)  
Palpitation, [1451](#)  
PALS, [1451](#)  
Palsy, [1451](#)  
Palsy Of The Long Thoracic Nerve, [1451](#)  
Pamidronic Acid, [1452](#)  
Pampiniform Plexus, [1452](#)  
Panacinar Emphysema, [1452](#)  
Pancoast Syndrome, [1452](#)  
Pancoast Tumor, [1452](#)  
Pancolitis, [1452](#)  
Pancreas, [1452](#)  
Pancreas Divisum, [1452](#)  
Pancreas Enzyme, [1452](#)  
Pancreas Infection, [1452](#)  
Pancreas Inflammation, [1452](#)  
Pancreas Neoplasm, [1452](#)  
Pancreas Transplant, [1452](#)  
Pancreas Transplantation, [1452](#)  
Pancreatectomy, [1452](#)  
Pancreatic Abscess, [1453](#)  
Pancreatic Adenocarcinoma, [1453](#)  
Pancreatic Cancer, [1453](#)  
Pancreatic Carcinoma, [1453](#)  
Pancreatic Cysts, [1453](#)  
Pancreatic Diseases, [1453](#)  
Pancreatic Duct, [1453](#)  
Pancreatic Ductal Adenocarcinoma, [1453](#)  
Pancreatic Elastase, [1453](#)  
Pancreatic Enzyme, [1453](#)  
Pancreatic Exocrine Insufficiency, [1453](#)  
Pancreatic Fistula, [1453](#)  
Pancreatic Hormones, [1453](#)  
Pancreatic Inflammatory Cysts, [1454](#)  
Pancreatic Insufficiency, [1454](#)  
Pancreatic IPMN, [1454](#)  
Pancreatic IPMN Entry, [1454](#)  
Pancreatic Islet, [1454](#)  
Pancreatic Juice, [1454](#)  
Pancreatic Mass, [1454](#)  
Pancreatic Necrosis, [1454](#)  
Pancreatic Neuroendocrine Tumor, [1454](#)  
Pancreatic Neuroendocrine Tumors, [1454](#)  
Pancreatic Non-Inflammatory Cysts, [1454](#)  
Pancreatic Pseudocyst, [1454](#)  
Pancreatic Serous Cystadenoma, [1454](#)  
Pancreatic Surgery, [1455](#)  
Pancreatic Trauma, [1455](#)  
Pancreaticobiliary Cytology, [1454](#)  
Pancreaticoduodenectomy, [1454](#)  
Pancreatitis, [1455](#)  
Pancreatoblastoma, [1455](#)  
Pancreatography, [1455](#)  
Pancreozymin, [1455](#)  
Pancrelipase, [1455](#)  
Pancuronium, [1455](#)  
Pancytopenia, [1455](#)  
PANDAS, [1455](#)  
Pandemic, [1455](#)  
Panendoscopy, [1455](#)  
Paneth Cell, [1455](#)  
Panhypopituitarism, [1456](#)  
Panic, [1456](#)  
Panic And Anxiety Disorders, [1645](#)  
Panic Attack, [1456](#)  
Panic Attacks, [1456](#)  
Panic Attacks And Panic Disorder, [1456](#)  
Panic Disorder, [1456](#)  
Panlobular Emphysema, [1456](#)  
Panmyelopathy, [1456](#)  
Panniculectomy, [1456](#)  
Panniculitis, [1456](#)  
Panniculus, [1456](#)  
Pannus, [1456](#)  
Panophthalmitis, [1456](#)  
Panoramic Radiograph, [1456](#)  
Panretinal Photocoagulation, [1457](#)  
Pantothenic Acid, [1457](#)  
Pantothenic Acid And Biotin, [1457](#)  
Panuveitis, [1457](#)  
Pap Smear, [1458](#)  
Pap Test, [1458](#)  
Papain, [1457](#)  
Papillary Fibroelastoma, [1457](#)  
Papillary Muscle, [1457](#)  
Papillary Thyroid Carcinoma, [1457](#)  
Papillectomy, [1457](#)  
Papilledema, [1457](#)  
Papilloedema, [1457](#)  
Papilloma, [1457](#)  
Papilloma With Atypia / DCIS Pathology, [1458](#)

---

---

Papillomatosis, [1457](#)  
Papillomatosis Cutis Lymphostatica, [1457](#)  
Papillomavirus, [1457](#)  
Papular, [1458](#)  
Papular Urticaria, [1458](#)  
Papule, [1458](#)  
Papules, [1458](#)  
PAPVR, [1458](#)  
Paracentesis, [1458](#)  
Paracetamol, [1458](#)  
Paracetamol Overdose, [1458](#)  
Paracetamol Toxicity, [1458](#)  
Paracusia, [1458](#)  
Paradichlorobenzene Poisoning, [1458](#)  
Paraesophageal Hernia, [1459](#)  
Paraesthesia, [1459](#)  
Paraffin, [1459](#)  
Paraffin Poisoning, [1459](#)  
Paraganglioma, [1459](#)  
Paragonimiasis, [1459](#)  
Paragraphia, [1459](#)  
Parainfectious, [1459](#)  
Parainfluenza, [1459](#)  
Paraldehyde, [1459](#)  
Paralysis, [1459](#)  
Paralysis Agitans, [1459](#)  
Paralytic Ileus, [1459](#)  
Paramedical, [1459](#)  
Parametritis, [1459](#)  
Paramnesia, [1460](#)  
Paramyotonia Congenita, [1460](#)  
Paranasal Sinuses, [1460](#)  
Paraneoplastic Autonomic Neuropathy, [1460](#)  
Paraneoplastic Cerebellar Degeneration, [1460](#)  
Paraneoplastic Disorder, [1460](#)  
Paraneoplastic Encephalomyelitis, [1460](#)  
Paraneoplastic Endocrine Syndromes, [1460](#)  
Paraneoplastic Neurologic Syndrome, [1460](#)  
Paraneoplastic Neurologic Syndromes, [1460](#)  
Paraneoplastic Neurological Disorder, [1460](#)  
Paraneoplastic Neurological Syndrome, [1460](#)  
Paraneoplastic Pemphigus, [1460](#)  
Paraneoplastic Syndrome, [1461](#)  
Paraneoplastic Syndromes Of The Nervous System, [1461](#)  
Paranoia, [1461](#)  
Paranoid, [1461](#)  
Paranoid Ideation, [1461](#)  
Paranoid Personality Disorder, [1461](#)  
Paranoid Psychosis, [1461](#)  
Paraparesis, [1461](#)  
Paraphasia, [1461](#)  
Paraphilia, [1461](#)  
Paraphilic Disorder, [1461](#)  
Paraphilic Disorders, [1461](#)  
Paraphimosis, [1461](#)  
Paraplegia, [1462](#)  
Parapneumonic Effusion, [1462](#)  
Parapneumonic Pleural Effusion, [1462](#)  
Parapneumonic Pleural Effusions and Empyema Thoracis, [1462](#)  
Parapsoriasis, [1462](#)  
Paraquat, [1462](#)  
Paraquat Poisoning, [1462](#)  
Paraseptal Emphysema, [1462](#)  
Parasite, [1462](#)  
Parasitemia, [1462](#)  
Parasitic, [1462](#)  
Parasitic Infection, [1462](#)  
Parasitic Worm, [1463](#)  
Parasitology, [1463](#)  
Parasitosis, [1463](#)  
Parasomnia, [1463](#)  
Parasomnias, [1463](#)  
Parasternal Heave, [1463](#)  
Parasuicide, [1463](#)  
Parasympathetic Nervous System, [1463](#)  
Parasympatholytic, [2127](#)  
Parasympathomimetic, [2127](#)  
Parathyroid, [1463](#)  
Parathyroid Adenoma, [1463](#)  
Parathyroid Cancer, [1463](#)  
Parathyroid Carcinoma, [1463](#)  
Parathyroid Crisis, [1464](#)  
Parathyroid Gland, [1464](#)  
Parathyroid Gland Removal, [1464](#)  
Parathyroid Glands, [1464](#)  
Parathyroid Hormone, [1464](#)  
Parathyroid Hormone Blood Test, [1464](#)  
Parathyroid Hormone Physiology, [1464](#)  
Parathyroid Hormone-Related Protein Blood Test, [1464](#)  
Parathyroid Hyperplasia, [1464](#)  
Parathyroid Surgery, [1464](#)  
Parathyroid, Underactivity Of, [977](#)  
Parathyroidectomy, [1464](#)  
Paratubal Cyst, [1464](#)  
Paratyphoid Fever, [1464](#)  
Parechovirus, [1464](#)  
Parenchyma, [1465](#)  
Parenchymal, [1465](#)  
Parenteral, [1465](#)  
Parenteral Nutrition, [1465](#)  
Parenting, [1465](#)  
Paresis, [1465](#)  
Paresthesia, [1465](#)  
Paresthesia (Pins and Needles), [1465](#)  
Paricalcitol, [1465](#)  
Parietal, [1465](#)  
Parietal Bone, [1465](#)

---

---

Parietal Cell Antibodies, [1465](#)  
 Parietal Lobe, [1465](#)  
 Parietal Pericardium, [1465](#)  
 Parietal Pleura, [1465](#)  
 Parinaud Oculoglandular Syndrome, [1466](#)  
 Parinaud Syndrome, [1466](#)  
 Paris System For Reporting Urinary Cytology, [1466](#)  
 Parity, [1466](#)  
 Park-Bench Position, [1466](#)  
 Parkinson Disease, [1466](#)  
 Parkinson's Disease, [1466](#)  
 Parkinson's Disease Center, [1645](#)  
 Parkinsonian Gait, [1466](#)  
 Parkinsonism, [1466](#)  
 Paromomycin, [1466](#)  
 Paronychia, [1466](#)  
 Parosteal Osteosarcoma, [1466](#)  
 Parotid Gland, [1467](#)  
 Parotid Neoplasm, [1467](#)  
 Parotid Tumor, [1467](#)  
 Parotid Tumors, [1467](#)  
 Parotidectomy, [1467](#)  
 Parotitis, [1467](#)  
 Paroxetine, [2127](#)  
 Paroxetine Hydrochloride, [1467](#)  
 Paroxysm, [1467](#)  
 Paroxysmal Atrial Fibrillation, [1467](#)  
 Paroxysmal Atrial Tachycardia, [1467](#)  
 Paroxysmal Cold Hemoglobinuria, [1467](#)  
 Paroxysmal Hemicrania, [1467](#)  
 Paroxysmal Nocturnal Dyspnea, [1467](#)  
 Paroxysmal Nocturnal Haemoglobinuria, [1467](#)  
 Paroxysmal Nocturnal Hemoglobinuria, [1468](#)  
 Paroxysmal Nocturnal Hemoglobinuria (PNH), [1645](#)  
 Paroxysmal Positional Nystagmus, [1646](#)  
 Paroxysmal Supraventricular Tachycardia, [1468](#)  
 Pars, [1468](#)  
 Pars Compacta, [1468](#)  
 Pars Planitis, [1468](#)  
 Parsonage-Turner Syndrome, [1468](#)  
 Partial, [1468](#)  
 Partial Agonist, [1468](#)  
 Partial Anomalous Pulmonary Venous Connection, [1468](#)  
 Partial Breast Brachytherapy, [1468](#)  
 Partial Cystectomy, [1468](#)  
 Partial Denture, [1468](#)  
 Partial Hysterectomy, [1468](#)  
 Partial Knee Replacement, [1469](#)  
 Partial Nephrectomy, [1469](#)  
 Partial Remission, [1469](#)  
 Partial Response, [1469](#)  
 Partial Seizure, [1469](#)  
 Partial Thromboplastin Time, [1469](#)  
 Partial Volume Artifact, [1469](#)  
 Partial-Volume Effect, [1469](#)  
 Participation, [1469](#)  
 Particle Size, [1469](#)  
 Particulate Matter, [1469](#)  
 Partogram, [1469](#)  
 Parturition, [1469](#)  
 Paruresis, [1470](#)  
 Parvimonas Micra, [1470](#)  
 Parvoviridae, [1470](#)  
 Parvovirus, [1470](#)  
 Parvovirus B19, [1470](#)  
 Parvovirus B19 Infection, [1470](#)  
 Parvovirus Infection, [1470](#)  
 Passive Exercise, [1470](#)  
 Passive Heat Therapy, [1470](#)  
 Passive Immunity, [1470](#)  
 Passive Immunization, [1470](#)  
 Passive Smoking, [1470](#)  
 Pasteurella, [1470](#)  
 Pasteurella Aerogenes, [1470](#)  
 Pasteurella Canis, [1470](#)  
 Pasteurella Multocida, [1471](#)  
 Pasteurellaceae, [1471](#)  
 Pasteurellosis, [1471](#)  
 Pasteurization, [1471](#)  
 Pastia Lines, [1471](#)  
 Patau Syndrome, [1471](#)  
 Patau's Syndrome, [1471](#)  
 Patch, [1471](#)  
 Patch Graft, [1471](#)  
 Patch Test, [1471](#)  
 Patch Testing, [1471](#)  
 Patches, [1471](#)  
 Patchy Skin Color, [1471](#)  
 Patella, [1471](#)  
 Patellar Apprehension Test, [1472](#)  
 Patellar Bulge Test, [1472](#)  
 Patellar Dislocation, [1472](#)  
 Patellar Reflex, [1472](#)  
 Patellar Tap Test, [1472](#)  
 Patellar Tendinitis, [1472](#)  
 Patellar Tendinopathy, [1472](#)  
 Patellofemoral Instability, [1472](#)  
 Patellofemoral Osteoarthritis, [1472](#)  
 Patellofemoral Pain Syndrome, [1472](#)  
 Patellofemoral Syndrome, [1472](#)  
 Patent, [1473](#)  
 Patent Ductus Arteriosus, [1473](#)  
 Patent Foramen Ovale, [1473](#)  
 Patent Urachus, [1473](#)  
 Patent Urachus Repair, [1473](#)  
 Paternal Depression, [1473](#)  
 Paternal Postnatal Depression, [1473](#)  
 Paternal Uniparental Disomy 14, [1473](#)

---

---

Paternity Testing, [1473](#)  
Pathogen, [1473](#)  
Pathogenesis, [1473](#)  
Pathogenic, [1473](#)  
Pathogenic Variant, [1473](#)  
Pathogenicity, [1473](#)  
Pathognomonic, [1474](#)  
Pathologic, [1474](#)  
Pathologic Dilatation, [1474](#)  
Pathologic Fracture, [1474](#)  
Pathologic Hypertrophy, [1474](#)  
Pathologic Nipple Discharge, [1474](#)  
Pathologic Ossification, [1474](#)  
Pathologic Process, [1474](#)  
Pathologic Response, [1474](#)  
Pathological Fracture, [1474](#)  
Pathological Laughter And Crying, [1474](#)  
Pathological Stealing, [1474](#)  
Pathologist, [1474](#)  
Pathology, [1474](#)  
Pathology Artifact, [1474](#)  
Pathophysiology, [1474](#)  
Patient, [1475](#)  
Patient Advocacy, [1475](#)  
Patient Autonomy, [1475](#)  
Patient Education, [1475](#)  
Patient Observation, [1475](#)  
Patient Positioning, [1475](#)  
Patient Preparation, [1475](#)  
Patient Rights, [1475](#)  
Patient Safety, [1475](#)  
Patient-Centered Care, [1475](#)  
Patient-Centered Outcome, [1475](#)  
Patient-Controlled Analgesia, [1475](#)  
Patient-Reported Outcome Measure, [1475](#)  
Patlak-Rutland Analysis, [1475](#)  
Patrick Test, [1476](#)  
Pattern Recognition Receptors, [1476](#)  
Pattern-Recognition Receptors, [1476](#)  
Pauci-Immune Glomerulonephritis, [1476](#)  
Paxillin, [1476](#)  
Pazopanib, [1476](#)  
PBA, [1476](#)  
PBG Urine Test, [1476](#)  
PCBs, [1476](#)  
PCL Injuries, [1476](#)  
PCO, [1476](#)  
PCO Disease, [1476](#)  
PCOS, [1477](#)  
PCP, [1477](#)  
PCP and Ketamine Toxicity, [1477](#)  
PDA, [1477](#)  
PDE5 Inhibitors, [1477](#)  
PE Anticoagulation, [1477](#)  
Peak Expiratory Flow Rate, [1477](#)  
Peak Flow Meter, [1477](#)  
Peak Plasma Concentration, [1477](#)  
Peanut Allergy, [1477](#)  
Peanut Oil, [1477](#)  
Pearly Penile Papules, [1477](#)  
Peau D'Orange, [1477](#)  
Pectinate Line, [1477](#)  
Pectoral, [1478](#)  
Pectoral Girdle, [1478](#)  
Pectoralis, [1478](#)  
Pectoralis Major Muscle, [1478](#)  
Pectoralis Minor Muscle, [1478](#)  
Pectoralis Muscles, [1478](#)  
Pectoriloquy, [1478](#)  
Pectus Carinatum, [1478](#)  
Pectus Excavatum, [1478](#)  
Pectus Excavatum Repair, [1478](#)  
Pediatric ADHD, [1645](#)  
Pediatric Advanced Life Support, [1478](#)  
Pediatric AIDS, [1478](#)  
Pediatric Anesthesia, [1478](#)  
Pediatric Antireflux Surgery, [1478](#)  
Pediatric Apnea, [1479](#)  
Pediatric Appendicitis, [1479](#)  
Pediatric Assessment Triangle, [1479](#)  
Pediatric Brain Tumor, [1479](#)  
Pediatric Brain Tumors, [1479](#)  
Pediatric Cancer, [1479](#)  
Pediatric Cardiopulmonary Resuscitation, [1479](#)  
Pediatric Cataract, [1479](#)  
Pediatric Central Venous Line, [1479](#)  
Pediatric Crutch Ambulation, Chair Transfers, [1479](#)  
Pediatric Crutch Ambulation, Gait and Fitting, [1480](#)  
Pediatric Crutch Ambulation, Stair Navigation, [1480](#)  
Pediatric Crutch Gait Training and Fitting, [1480](#)  
Pediatric Crutch Stair Training, [1480](#)  
Pediatric Crutch Transfers, [1479](#)  
Pediatric Dehydration, [1480](#)  
Pediatric Dentistry, [1480](#)  
Pediatric Dermatology, [1480](#)  
Pediatric Dermatology Conditions, [1480](#)  
Pediatric Diabetic Ketoacidosis, [1480](#)  
Pediatric Early Warning Score, [1480](#)  
Pediatric Echocardiogram, [1480](#)  
Pediatric Eczema, [1480](#)  
Pediatric Feeding Tube, [1480](#)  
Pediatric Fever, [1481](#)  
Pediatric Foreign Body Ingestion, [1481](#)  
Pediatric Gastroesophageal Reflux Disease, [1481](#)  
Pediatric Glaucoma, [1481](#)  
Pediatric Growth Hormone Deficiency, [1481](#)  
Pediatric Health, [1481](#)  
Pediatric Heart Surgery, [1481](#)

---

---

Pediatric Hepatitis A, [1481](#)  
Pediatric Hepatitis B, [1481](#)  
Pediatric Hepatitis C, [1482](#)  
Pediatric Hypercholesterolemia, [1482](#)  
Pediatric Hypertension, [1482](#)  
Pediatric Limp, [1482](#)  
Pediatric Meningitis, [1482](#)  
Pediatric Midline Catheters, [1482](#)  
Pediatric Myocarditis, [1482](#)  
Pediatric Nasal Congestion, [1482](#)  
Pediatric Nuclear Medicine, [2127](#)  
Pediatric Nursing, [1482](#)  
Pediatric Obesity, [1482](#)  
Pediatric Obstructive Sleep Apnea, [1483](#)  
Pediatric Ophthalmology, [1483](#)  
Pediatric Orthopedics, [1483](#)  
Pediatric Pancreatitis, [1483](#)  
Pediatric Pneumonia, [1483](#)  
Pediatric Poisoning, [1483](#)  
Pediatric Ptosis, [1483](#)  
Pediatric Pump Gastrostomy Feeding, [1483](#)  
Pediatric Rash Evaluation, [1483](#)  
Pediatric Rehabilitation, [1483](#)  
Pediatric Sepsis, [1484](#)  
Pediatric Shock, [1484](#)  
Pediatric Skin Tumors, [1484](#)  
Pediatric Sleep Apnea, [1484](#)  
Pediatric Total Parenteral Nutrition, [1484](#)  
Pediatric Toxicology, [1484](#)  
Pediatric Volvulus, [1484](#)  
Pediatrician, [1482](#)  
Pediatrics, [1484](#)  
Pedicle, [1484](#)  
Pedicle Skin Graft, [1484](#)  
Pediculosis, [1484](#)  
Pediculosis and Pthiriasis (Lice Infestation), [1484](#)  
Pediculosis Capitis, [1484](#)  
Pediculus Humanus Capitis, [1484](#)  
Pedigree, [1485](#)  
Pedodontics, [1485](#)  
Pedophilia, [1485](#)  
Pedophilic Disorder, [1485](#)  
Pedunculate, [1485](#)  
Pedunculated, [1485](#)  
Peeling Skin, [1485](#)  
PEEP, [1485](#)  
PEF, [1485](#)  
Pefloxacin, [2127](#)  
PEG, [2127](#)  
PEG Tube, [1485](#)  
Pegaspargase, [2127](#)  
Pegfilgrastim, [2127](#)  
Pel-Ebstein Fever, [1485](#)  
Peliosis Hepatis, [1485](#)  
Pelizaeus-Merzbacher Disease, [1485](#)  
Pellagra, [1486](#)  
Pellet, [1486](#)  
Pelvic, [1486](#)  
Pelvic Binder, [1486](#)  
Pelvic Bone, [1486](#)  
Pelvic Cavity, [1486](#)  
Pelvic Congestion, [1486](#)  
Pelvic Congestion Syndrome, [1486](#)  
Pelvic CT Scan, [1486](#)  
Pelvic Diaphragm, [1486](#)  
Pelvic Examination, [1486](#)  
Pelvic Exenteration, [1486](#)  
Pelvic Floor, [1486](#)  
Pelvic Floor Dysfunction, [1486](#)  
Pelvic Floor Health And Disorders, [1487](#)  
Pelvic Floor Muscle Training Exercises, [1487](#)  
Pelvic Floor Muscles, [1487](#)  
Pelvic Fracture, [1487](#)  
Pelvic Girdle, [1487](#)  
Pelvic Girdle Pain In Pregnancy, [1487](#)  
Pelvic Infection, [1487](#)  
Pelvic Inflammatory Disease, [1487](#)  
Pelvic Inflammatory Disease (PID), [1487](#)  
Pelvic Inlet, [1487](#)  
Pelvic Kidney, [1487](#)  
Pelvic Laparoscopy, [1487](#)  
Pelvic Lymphadenectomy, [1487](#)  
Pelvic Mass, [1487](#)  
Pelvic Neoplasm, [1487](#)  
Pelvic Organ Prolapse, [1488](#)  
Pelvic Organ Prolapse Quantification System, [1488](#)  
Pelvic Outlet, [1488](#)  
Pelvic Pain, [1488](#)  
Pelvic Pain During Early Pregnancy, [1488](#)  
Pelvic Ring Fracture, [1488](#)  
Pelvic Spleen, [1488](#)  
Pelvic Viscus, [1488](#)  
Pelvimeter, [1488](#)  
Pelvimetry, [1488](#)  
Pelvis, [1488](#)  
Pelvis MRI Scan, [1488](#)  
Pelvis X-Ray, [1488](#)  
Pemberton Sign, [1488](#)  
Pemberton'S Sign, [1489](#)  
Pemetrexed, [1489](#)  
Pemetrexed Disodium, [1489](#)  
Pemoline, [1489](#)  
Pemphigoid, [1489](#)  
Pemphigoid Gestationis, [1489](#)  
Pemphigus, [1489](#)  
Pemphigus Erythematosus, [1489](#)  
Pemphigus Foliaceus, [1489](#)  
Pemphigus Vulgaris, [1489](#)

---

---

Penbutolol, [1489](#)  
Pencil Eraser Swallowing, [1489](#)  
Pencil Swallowing, [1489](#)  
Pendred Syndrome, [1490](#)  
Pendular, [1490](#)  
Pendular Nystagmus, [1490](#)  
Pendulous, [1490](#)  
Penetrance, [1490](#)  
Penetrating Abdominal Trauma, [1490](#)  
Penetrating Wounds, [1490](#)  
Penicillamine, [2127](#)  
Penicillin, [1490](#)  
Penicillin Allergy, [1490](#)  
Penicillin G, [1490](#)  
Penicillin V, [1490](#)  
Penicillium, [1490](#)  
Penile, [1490](#)  
Penile Adhesions, [1490](#)  
Penile Augmentation, [1491](#)  
Penile Cancer, [1491](#)  
Penile Disease, [1491](#)  
Penile Edema, [1491](#)  
Penile Erection, [1491](#)  
Penile Fracture, [1491](#)  
Penile Hygiene, [1491](#)  
Penile Implant, [1491](#)  
Penile Infection, [1491](#)  
Penile Intraepithelial Neoplasia, [1491](#)  
Penile Intraepithelial Neoplasia (PeIN or PIN), [1491](#)  
Penile Leukoplakia, [1491](#)  
Penile Lichen Sclerosus, [1491](#)  
Penile Prosthesis, [1491](#)  
Penile Torsion, [1492](#)  
Penile Trauma, [1492](#)  
Penis, [1492](#)  
Penis Carcinoma, [1492](#)  
Penis Pain, [1492](#)  
Penoscrotal Transposition, [1492](#)  
Pentad Of TTP, [1492](#)  
Pentalogy of Cantrell, [1492](#)  
Pentamidine, [1492](#)  
Pentamidine Isethionate, [1492](#)  
Pentatrachomonas Hominis, [1492](#)  
Pentavalent Vaccine, [2127](#)  
Pentazocine, [1492](#)  
Pentazocine Overdose, [1492](#)  
Pentetic Acid, [1493](#)  
Pentobarbital, [2127](#)  
Pentobarbital Overdose, [1493](#)  
Pentosan Polysulfate, [1493](#)  
Pentose, [1493](#)  
Pentose Phosphate Pathway, [1493](#)  
Pentoxifylline, [1493](#)  
Penumbra, [1493](#)  
Peplomycin, [1493](#)  
Peppermint, [1493](#)  
Peppermint Oil Overdose, [1493](#)  
Pepsin, [1493](#)  
Pepsin A, [1493](#)  
Pepsinogen, [1493](#)  
Peptic Esophagitis, [1493](#)  
Peptic Stricture, [1493](#)  
Peptic Ulcer, [1493](#)  
Peptic Ulcer Bleeding, [1494](#)  
Peptic Ulcer Cross-Reference, [1494](#)  
Peptic Ulcer Disease, [1494](#)  
Peptidase, [1494](#)  
Peptide, [1494](#)  
Peptide Biosynthesis, [1494](#)  
Peptide Bond, [1494](#)  
Peptide Hormone, [1494](#)  
Peptide Metabolism, [1494](#)  
Peptide Transport, [1494](#)  
Peptide Yy, [1494](#)  
Peptidoglycan, [1494](#)  
Peptococcaceae, [1494](#)  
Peptococcus, [1494](#)  
Peptococcus Niger, [1494](#)  
Peptoniphilus, [1495](#)  
Peptoniphilus Harei, [1495](#)  
Peptoniphilus Indolicus, [1495](#)  
Peptostreptococcus, [1495](#)  
Peracetic Acid, [1495](#)  
Peracute, [1495](#)  
PERC Rule, [1495](#)  
Percentile, [1495](#)  
Perception, [1495](#)  
Perceptual, [1495](#)  
Perceptual Disorder, [1495](#)  
Perceptual Distortion, [1495](#)  
Perceptual Masking, [1495](#)  
Percussion, [1495](#)  
Percussion Myotonia, [1495](#)  
Percussion Note, [1495](#)  
Percutaneous, [1496](#)  
Percutaneous Biopsy, [1496](#)  
Percutaneous Closure, [1496](#)  
Percutaneous Coronary Intervention, [1496](#)  
Percutaneous Discectomy, [1496](#)  
Percutaneous Diskectomy, [1496](#)  
Percutaneous Drainage, [1496](#)  
Percutaneous Electrical Nerve Stimulation, [1496](#)  
Percutaneous Endoscopic Gastrostomy, [1496](#)  
Percutaneous Kidney Procedures, [1496](#)  
Percutaneous Nephrolithotomy, [1496](#)  
Percutaneous Nephrolithotripsy, [1496](#)  
Percutaneous Nephrostomy, [1496](#)  
Percutaneous Transhepatic Cholangiogram, [1497](#)

---



---

Percutaneous Transhepatic Cholangiography, [1497](#)  
 Percutaneous Transluminal Coronary Angioplasty, [1497](#)  
 Percutaneous Umbilical Blood Sampling, [1497](#)  
 Perforant Pathway, [1497](#)  
 Perforated Bowel, [1497](#)  
 Perforated Eardrum, [1497](#)  
 Perforating Folliculitis, [1497](#)  
 Perforation, [1497](#)  
 Perforation Of A Viscus, [1497](#)  
 Performance, [1497](#)  
 Performance Anxiety, [1497](#)  
 Performance Status, [1497](#)  
 Perfusion, [1497](#)  
 Perfusion Defect, [1498](#)  
 Perfusion Imaging, [1498](#)  
 Perfusion PET, [1498](#)  
 Peri-Implantitis, [1501](#)  
 Perianal, [1498](#)  
 Perianal Abscess, [1498](#)  
 Perianal Glands, [1498](#)  
 Perianal Streptococcal Cellulitis, [1498](#)  
 Periapical Abscess, [1498](#)  
 Periapical Granuloma, [1498](#)  
 Periappendiceal, [1498](#)  
 Periarteritis Nodosa, [1498](#)  
 Periarticular, [1498](#)  
 Pericardial, [1498](#)  
 Pericardial Cavity, [1498](#)  
 Pericardial Effusion, [1498](#)  
 Pericardial Effusion Etiology, [1499](#)  
 Pericardial Fluid Analysis, [1499](#)  
 Pericardial Fluid Culture, [1499](#)  
 Pericardial Fluid Gram Stain, [1499](#)  
 Pericardial Friction Rub, [1499](#)  
 Pericardial Inflammation, [1499](#)  
 Pericardial Knock, [1499](#)  
 Pericardial Knock Sound, [1499](#)  
 Pericardial Mesothelioma, [1499](#)  
 Pericardial Metastases, [1499](#)  
 Pericardial Rub, [1499](#)  
 Pericardial Stripping, [1499](#)  
 Pericardial Tamponade, [1500](#)  
 Pericardial Wall, [1500](#)  
 Pericardial Window, [1500](#)  
 Pericardiectomy, [1500](#)  
 Pericardiocentesis, [1500](#)  
 Pericardiostomy, [1500](#)  
 Pericarditis, [1500](#)  
 Pericardium, [1500](#)  
 Perichondritis, [1500](#)  
 Perichondritis Of The Ear, [1500](#)  
 Perichondrium, [1500](#)  
 Pericoronitis, [1500](#)  
 Pericytes, [1501](#)  
 Periductal Mastitis, [1501](#)  
 Perifolliculitis Capitis Abscedens et Suffodiens, [1501](#)  
 Perilymph, [1501](#)  
 Perilymphatic Fistula, [1501](#)  
 Perimenopausal, [1501](#)  
 Perimenopause, [1501](#)  
 Perimetry, [1501](#)  
 Perimysium, [1501](#)  
 Perinatal, [1501](#)  
 Perinatal Asphyxia, [1501](#)  
 Perinatal Death, [1502](#)  
 Perinatal Psychiatry, [1502](#)  
 Perinatal Transmission, [1502](#)  
 Perinatologist, [1502](#)  
 Perindopril, [1502](#)  
 Perineal, [1502](#)  
 Perineal Hemostat, [1502](#)  
 Perineal Laceration, [1502](#)  
 Perineal Laceration Classification, [1502](#)  
 Perineal Needle Holder, [1502](#)  
 Perineal Protection, [1502](#)  
 Perineal Suture, [1502](#)  
 Perineal Tear, [1502](#)  
 Perineum, [1502](#)  
 Perineural, [1503](#)  
 Perineuronal Satellitosis, [1503](#)  
 Perinuclear Space, [1503](#)  
 Periodic Abstinence, [1503](#)  
 Periodic Acid, [1503](#)  
 Periodic Breathing, [1503](#)  
 Periodic Limb Movement Disorder, [1503](#)  
 Periodic Limb Movement Sleep Disorder, [1503](#)  
 Periodic Paralysis Syndrome, [1503](#)  
 Periodontal Abscess, [1503](#)  
 Periodontal Cyst, [1503](#)  
 Periodontal Disease, [1503](#)  
 Periodontal Dressing, [1503](#)  
 Periodontal Flap Surgery, [1503](#)  
 Periodontal Graft, [1504](#)  
 Periodontal Ligament, [1504](#)  
 Periodontal Pocket, [1504](#)  
 Periodontal Probing, [1504](#)  
 Periodontics, [1504](#)  
 Periodontist, [1504](#)  
 Periodontitis, [1504](#)  
 Periodontium, [1504](#)  
 Periodontium Disease, [1504](#)  
 Perionychium, [1504](#)  
 Perioperative, [1504](#)  
 Perioperative Antibiotic Prophylaxis, [2128](#)  
 Perioral Dermatitis, [1504](#)  
 Periorbital, [1504](#)  
 Periorbital Cellulitis, [1505](#)  
 Periorbital Edema, [1505](#)

---

---

Periorbital Infection, [1505](#)  
 Periosteal Osteosarcoma, [1505](#)  
 Periosteal Reaction, [1505](#)  
 Periosteum, [1505](#)  
 Periostitis, [1505](#)  
 Peripancreatic Malignancy, [1505](#)  
 Peripartum, [1505](#)  
 Peripartum (Postpartum) Cardiomyopathy (PPCM), [1505](#)  
 Peripartum Cardiomyopathy, [1505](#)  
 Peripartum Cardiomyopathy Prognosis, [1505](#)  
 Peripheral, [1505](#)  
 Peripheral Angioplasty, [1506](#)  
 Peripheral Artery Disease, [1506](#)  
 Peripheral Blood, [1506](#)  
 Peripheral Blood Smear, [1506](#)  
 Peripheral Blood Stem Cells, [1506](#)  
 Peripheral Cyanosis, [1506](#)  
 Peripheral Edema, [1506](#)  
 Peripheral Ischemia, [1506](#)  
 Peripheral Nerve, [1506](#)  
 Peripheral Nerve Blocks, [1506](#)  
 Peripheral Nerve Injuries, [1506](#)  
 Peripheral Nerve Tumors, [1506](#)  
 Peripheral Nervous System, [1506](#)  
 Peripheral Neuritis, [1507](#)  
 Peripheral Neurofibromatosis, [1507](#)  
 Peripheral Neuropathy, [1507](#)  
 Peripheral Oedema, [1507](#)  
 Peripheral Vascular Disease, [1507](#)  
 Peripheral Vision, [1507](#)  
 Peripherally Inserted Central Catheter, [1506](#)  
 Periplasm, [1507](#)  
 Periprosthetic Joint Infection, [1507](#)  
 Perirenal Abscess, [1507](#)  
 Perirenal Extramedullary Hematopoiesis, [1507](#)  
 Perirenal Hematoma, [1507](#)  
 Peristalsis, [1507](#)  
 Peritoneal, [1507](#)  
 Peritoneal Cancer, [1507](#)  
 Peritoneal Carcinomatosis, [1508](#)  
 Peritoneal Cavity, [1508](#)  
 Peritoneal Dialysis, [1508](#)  
 Peritoneal Dialysis Catheter, [1508](#)  
 Peritoneal Dialysis-Associated Peritonitis, [1508](#)  
 Peritoneal Diseases, [1508](#)  
 Peritoneal Fluid, [1508](#)  
 Peritoneal Fluid Analysis, [1508](#)  
 Peritoneal Fluid Culture, [1508](#)  
 Peritoneal Infection, [1508](#)  
 Peritoneal Lavage, [1508](#)  
 Peritoneal Necrosis, [1508](#)  
 Peritoneal Sac, [1508](#)  
 Peritoneum, [1509](#)  
 Peritoneum Neoplasm, [1509](#)  
 Peritonitis, [1509](#)  
 Peritonitis In PD, [1509](#)  
 Peritonsillar Abscess, [1509](#)  
 Periungual Fibroma, [1509](#)  
 Periurethral, [1509](#)  
 Periventricular Leukomalacia, [1509](#)  
 Perivitelline Space, [1509](#)  
 Perleche, [1509](#)  
 Permanent Teeth, [1509](#)  
 Permease, [1509](#)  
 Permethrin, [1509](#)  
 Permissive Hypercapnia, [1509](#)  
 Pernicious Anaemia, [1510](#)  
 Pernicious Anemia, [1510](#)  
 Pernio, [1510](#)  
 Peroneal, [1510](#)  
 Peroneal Muscular Atrophy, [1510](#)  
 Peroneal Nerve, [1510](#)  
 Peroneal Nerve Palsy, [1510](#)  
 Peroneal Tendon Syndromes, [1510](#)  
 Peroxidase, [1510](#)  
 Peroxisomal Disorder, [1510](#)  
 Peroxisome, [1510](#)  
 Peroxisome Proliferation, [1510](#)  
 Perphenazine, [1510](#)  
 Perseveration, [1510](#)  
 Persistent Asthma, [1510](#)  
 Persistent Cloaca, [1511](#)  
 Persistent Depressive Disorder, [1511](#)  
 Persistent Fetal Vasculature, [1511](#)  
 Persistent Generalized Lymphadenopathy, [1511](#)  
 Persistent Pulmonary Hypertension Of The Newborn, [1511](#)  
 Persistent Pupillary Membranes, [1511](#)  
 Persistent Vegetative State, [1511](#)  
 Person With Disability, [1512](#)  
 Personal Emergency Response System, [1511](#)  
 Personal Protective Equipment, [1512](#)  
 Personality, [1511](#)  
 Personality Change, [1511](#)  
 Personality Disorder, [1511](#)  
 Personality Disorders, [1511](#)  
 Personality Test, [1512](#)  
 Personalized Medicine, [1512](#)  
 Perspiration, [1512](#)  
 Pertussis, [1512](#)  
 Pertussis Toxin, [1512](#)  
 Pertussis Vaccine, [2128](#)  
 Pertuzumab, [1512](#)  
 Pes, [1512](#)  
 Pes Anserinus, [1512](#)  
 Pes Cavus, [1512](#)  
 Pes Planus, [1512](#)

---

---

Pessaries, [1512](#)  
Pessary, [1512](#)  
Pessimism, [1512](#)  
Pesticide, [1513](#)  
Pesticide Poisoning, [1513](#)  
Pesticides, [1513](#)  
Pesticides On Fruits And Vegetables, [1513](#)  
Pestilence, [1513](#)  
Pet Allergy, [1513](#)  
PET Scan, [1514](#)  
PET Scan For Breast Cancer, [1514](#)  
PET Scanner, [1514](#)  
PET Tracers, [1514](#)  
PET/CT, [1513](#)  
PET/MRI, [1513](#)  
Petechia, [1513](#)  
Petechiae, [1513](#)  
Petit Mal, [1513](#)  
Petit Mal Seizure, [1513](#)  
Petrissage, [1513](#)  
Petrolatum, [1513](#)  
Petroleum Jelly Overdose, [1514](#)  
Petrosal Sinus, [1514](#)  
Petrositis, [1514](#)  
Petrous, [1514](#)  
Peutz-Jeghers Syndrome, [1514](#)  
Peutz-Jeghers's Syndrome, [1514](#)  
PEWS, [1514](#)  
Pexophagy, [1514](#)  
Peyer Patches, [1514](#)  
Peyer's Patches, [1514](#)  
Peyronie Disease, [1515](#)  
Peyronie's Disease, [1515](#)  
PFAPA Syndrome, [1515](#)  
Pfeiffer Syndrome, [1515](#)  
Pfeiffer Syndrome Type I, [1646](#)  
pH, [1515](#)  
PH Meter, [1524](#)  
PH Monitoring, [1524](#)  
Phacoemulsification, [1515](#)  
Phacomatosis, [1515](#)  
Phagocyte, [1515](#)  
Phagocytes, [1516](#)  
Phagocytic Vesicle, [1516](#)  
Phagocytosis, [1516](#)  
Phagolysosome, [1516](#)  
Phagophore, [1516](#)  
Phakomatosis TS, [1646](#)  
Phalangeal Fractures, [1516](#)  
Phalanges, [1516](#)  
Phalanx, [1516](#)  
Phalanx Unspecified, [1516](#)  
Phalen Maneuver, [1516](#)  
Phalen Sign, [1516](#)  
Phalen Test, [1516](#)  
Phantom, [1516](#)  
Phantom Limb, [1516](#)  
Phantom Limb Pain, [1517](#)  
Phantom Pain, [1517](#)  
Phantom Pregnancy, [1517](#)  
Phantosmia, [1517](#)  
Pharmaceutical, [2128](#)  
Pharmaceutical Adjuvant, [1517](#)  
Pharmaceutical Excipient, [1517](#)  
Pharmaceutical Solutions, [1517](#)  
Pharmacist, [1517](#)  
Pharmacodynamic Drug Interaction, [1517](#)  
Pharmacodynamic Interaction, [2128](#)  
Pharmacodynamics, [1517](#)  
Pharmacoepidemiology, [1517](#)  
Pharmacogenetic Testing, [1517](#)  
Pharmacogenetics, [1517](#)  
Pharmacogenic Myopathy, [1646](#)  
Pharmacogenomic Testing, [1518](#)  
Pharmacogenomics, [1517](#)  
Pharmacokinetic Drug Interaction, [1518](#)  
Pharmacokinetic Interaction, [2128](#)  
Pharmacokinetics, [1518](#)  
Pharmacologic Substance, [1518](#)  
Pharmacological Stress Test, [1518](#)  
Pharmacologist, [1518](#)  
Pharmacology, [1518](#)  
Pharmacopoeia, [1518](#)  
Pharmacoproteomics, [1518](#)  
Pharmacotherapy, [1518](#)  
Pharmacovigilance, [1518](#)  
Pharmacy, [1518](#)  
Pharmacy Facility, [1518](#)  
Pharyngeal, [1518](#)  
Pharyngeal Carcinoma, [1519](#)  
Pharyngeal Disease, [1519](#)  
Pharyngeal Hemorrhage, [1519](#)  
Pharyngeal Mucositis, [1519](#)  
Pharyngeal Stenosis, [1519](#)  
Pharyngeal Tonsil, [1519](#)  
Pharyngectomy, [1519](#)  
Pharyngitis, [1519](#)  
Pharyngolaryngeal Pain, [1519](#)  
Pharyngomaxillary Space Abscess, [1519](#)  
Pharynx, [1519](#)  
Pharynx Necrosis, [1519](#)  
Pharynx Neoplasm, [1519](#)  
Phase I Metabolism, [1520](#)  
Phase II Metabolism, [1519](#)  
Phase-Contrast Microscopy, [1519](#)  
Phe, [1520](#)  
Phelan-McDermid Syndrome, [1520](#)  
Phenacetin, [1520](#)

---

---

Phenazopyridine, [1520](#)  
Phencyclidine, [1520](#)  
Phencyclidine Overdose, [1520](#)  
Phencyclidine Substance Use, [1520](#)  
Phenelzine, [1520](#)  
Phenethylamines, [1520](#)  
Phenformin, [1520](#)  
Phenindione, [1520](#)  
Pheniramine Overdose, [1520](#)  
Phenobarbital, [1520](#)  
Phenobarbital Overdose, [1521](#)  
Phenobarbitone, [1521](#)  
Phenol, [1521](#)  
Phenols, [1521](#)  
Phenomenon, [1521](#)  
Phenothiazine, [1521](#)  
Phenothiazine Overdose, [1521](#)  
Phenothiazines, [1521](#)  
Phenotype, [1521](#)  
Phenoxybenzamine, [1521](#)  
Phenoxymethylpenicillin, [1521](#)  
Phenprocoumon, [1521](#)  
Phentermine, [1521](#)  
Phenthoate, [1521](#)  
Phentolamine, [1521](#)  
Phenylacetate, [1522](#)  
Phenylalanine, [1522](#)  
Phenylbutazone, [1522](#)  
Phenylbutyrate, [1522](#)  
Phenylephrine, [1522](#)  
Phenylketonuria, [1522](#)  
Phenylketonuria (PKU), [1646](#)  
Phenylpropanolamine, [1522](#)  
Phenytoin, [1522](#)  
Phenytoin Overdose, [1522](#)  
Phenytoin Sodium, [1522](#)  
Phenytoin Toxicity, [1522](#)  
Pheochromocytoma, [1522](#)  
Pheresis, [1523](#)  
Pheromone, [1523](#)  
Phialophora, [1523](#)  
Phialophora Verrucosa, [1523](#)  
Philadelphia Chromosome, [1523](#)  
Philodendron Poisoning, [1523](#)  
Phimosis, [1523](#)  
Phimosis and Paraphimosis, [1523](#)  
Phlebitis, [1523](#)  
Phlebitis And Thrombophlebitis, [1523](#)  
Phlebography, [1523](#)  
Phlebotomy, [1523](#)  
Phlebovirus, [1523](#)  
Phlegm, [1523](#)  
Phlegmasia Alba Dolens, [1523](#)  
Phlegmasia Cerulea Dolens, [1524](#)  
Phobia, [1524](#)  
Phobias, [1524](#)  
Phobic Disorder, [1524](#)  
Phocomelia, [1524](#)  
Phonation, [1524](#)  
Phonocardiogram, [1524](#)  
Phonocardiography, [1524](#)  
Phonological Disorder, [1524](#)  
Phonophobia, [1524](#)  
Phorate, [1524](#)  
Phorbols, [1524](#)  
Phorometer, [1524](#)  
Phosgene, [1524](#)  
Phosgene Oxime, [1525](#)  
Phosmet, [1525](#)  
Phosphamidon, [1525](#)  
Phosphate, [1525](#)  
Phosphate Binders, [1525](#)  
Phosphate Homeostasis, [1525](#)  
Phosphatidic Acid, [1525](#)  
Phosphatidylcholine, [1525](#)  
Phosphatidylethanolamines, [1525](#)  
Phosphatidylinositol, [1525](#)  
Phosphatidylserine, [1525](#)  
Phosphatidylserines, [1525](#)  
Phosphaturic Mesenchymal Tumor, [1525](#)  
Phosphine, [1525](#)  
Phosphodiesterase Inhibitor, [2128](#)  
Phosphodiesterase-5 Inhibitors, [1525](#)  
Phosphofructokinase-1, [1526](#)  
Phospholipase C Pathway, [1526](#)  
Phospholipid, [1526](#)  
Phosphorus, [1526](#)  
Phosphorus Blood Test, [1526](#)  
Phosphorus In Diet, [1526](#)  
Phosphorylation, [1526](#)  
Photoallergy, [1526](#)  
Photocoagulation, [1526](#)  
Photodermatitis, [1526](#)  
Photodynamic Therapy, [1526](#)  
Photodynamic Therapy For Cancer, [1526](#)  
Photoelectric Effect, [1526](#)  
Photographic Fixative Poisoning, [1526](#)  
Photomultiplier Tube, [1526](#)  
Photopeak, [1526](#)  
Photopheresis, [1527](#)  
Photophobia, [1527](#)  
Photoplethysmography, [1527](#)  
Photoreceptors, [1527](#)  
Photorefractive Keratectomy, [1527](#)  
Photosensitivity, [1527](#)  
Photosensitization, [1527](#)  
Photosensitizing Drugs, [1527](#)  
Phototherapy, [1527](#)

---

---

Phototoxic Dermatitis, 1527  
Phototoxicity, 1527  
Phrenic, 1527  
Phrenic Nerve, 1527  
Phrenic Nerve Palsy, 1527  
PHT, 1528  
Phthiriasis, 1528  
Phyllodes Tumor, 1528  
Phyllodes Tumors, 1528  
Physiatrist, 1528  
Physical Abuse, 1647  
Physical Activity, 1528  
Physical Allergy Urticaria, 1645  
Physical Child Abuse, 1528  
Physical Dependence, 1528  
Physical Exam Frequency, 1528  
Physical Examination, 1528  
Physical Half-Life, 1528  
Physical Medicine, 1528  
Physical Therapy, 1528  
Physical Therapy Assessment, 1528  
Physical Urticaria, Cholinergic Type, 391  
Physician Assistant, 1528  
Physicians' Desk Reference, 1529  
Physiologic, 1529  
Physiologic Jaundice Of Newborn, 1529  
Physiologic Jaundice Of The Newborn, 1529  
Physiological Dead Space, 1529  
Physiology, 1529  
Physiotherapy, 1529  
Physostigmine, 1529  
Phytomenadione, 1529  
Phytophotodermatitis, 1529  
Phytotherapy, 1518  
PI, 1647  
PI-RADS, 1535  
Pia Mater, 1529  
Pica, 1530  
PICC Line Dressing Care, 1530  
PICC Line Flushing, 1530  
PICC Line Insertion, 1530  
Pick Disease, 1530  
Pick's Disease, 1530  
Pickwickian Syndrome, 1530  
Pick's Disease, 1647  
PID, 1530  
PID (Pelvic Inflammatory Disease), 1645  
Piebaldism, 1530  
Piedra, 1530  
Pierre Robin Sequence, 1530  
Pierson Syndrome, 1530  
Piezogenic Pedal Papules, 1531  
Pigeon Breast, 1531  
Pigment, 1531  
Pigment Cirrhosis, 1531  
Pigment Dispersion Syndrome, 1531  
Pigmentary Glaucoma, 1531  
Pigmentary Mosaicism, 1531  
Pigmentation, 1531  
Pigmentation Disorder, 1531  
Pigmented Birthmarks, 1531  
Pigmented Purpuric Dermatitis, 1531  
Pilar Cyst, 1531  
Piles, 1531  
Piles (Haemorrhoids), 1532  
Pill, 1532  
Pilocarpine, 1532  
Pilocytic Astrocytoma, 1532  
Piloerection, 1532  
Pilomatrix Carcinoma, 1532  
Pilomatrixoma, 1532  
Pilomatrixoma (Pilomatricoma), 1532  
Pilonidal Cyst, 1532  
Pilonidal Dimple, 1532  
Pilonidal Disease, 1532  
Pilonidal Fistula, 1532  
Pilonidal Sinus Disease, 1532  
Pilus, 1532  
Pimozide, 1532  
Pimple, 1533  
Pimples (Acne), 1645  
Pin Care, 1533  
Pinch Strength, 1533  
Pinched Nerve, 1533  
Pindolol, 1533  
Pine Oil Poisoning, 1533  
Pineal Cyst, 1533  
Pineal Germinoma, 1533  
Pineal Gland, 1533  
Pineal Gland Tumor, 1533  
Pineal Parenchymal Tumor, 1533  
Pinealoma, 1533  
Pineoblastoma, 1533  
Pineocytoma, 1533  
Pinguecula, 1534  
Pinhole Meatus, 1534  
Pink Eye, 1534  
Pink Puffer, 1534  
Pinna, 1534  
Pinocytosis, 1534  
Pins and Needles, 1534  
Pint, 1534  
Pinta, 1534  
Pinworm, 1534  
Pinworm Infection, 1534  
Pinworm Infestation, 1534  
Pinworm Test, 1534  
Pinworms, 1534

---

---

Pioglitazone, [1534](#)  
 Piperacillin, [1535](#)  
 Piperazine, [1535](#)  
 Piperonyl Butoxide, [1535](#)  
 Piperonyl Butoxide With Pyrethrins Poisoning, [1535](#)  
 Pipotiazine, [1535](#)  
 Pipradrol, [1535](#)  
 Piracetam, [1535](#)  
 Pirfenidone, [1535](#)  
 Piriformis Syndrome, [1535](#)  
 Pirimicarb, [1535](#)  
 Piromidic Acid, [1535](#)  
 Piroxicam, [1535](#)  
 Piroxicam Overdose, [1535](#)  
 Pisa Syndrome, [1536](#)  
 Pisiform, [1536](#)  
 Pisiform Bone, [1536](#)  
 Pits, [1536](#)  
 Pitted Keratolysis, [1536](#)  
 Pitting Edema, [1536](#)  
 Pituicytoma, [1536](#)  
 Pituitary, [1536](#)  
 Pituitary Adenoma, [1536](#)  
 Pituitary Apoplexy, [1536](#)  
 Pituitary Carcinoma, [1536](#)  
 Pituitary Disease, [1536](#)  
 Pituitary Dwarfism, [1536](#)  
 Pituitary Gland, [1536](#)  
 Pituitary Hormone, [1536](#)  
 Pituitary Insufficiency, [1537](#)  
 Pituitary Macroadenoma, [1537](#)  
 Pituitary Microadenoma, [1537](#)  
 Pituitary Neoplasm, [1537](#)  
 Pituitary Neuroendocrine Tumor, [1537](#)  
 Pituitary Tumor, [1537](#)  
 Pituitary Tumors Pathology, [1537](#)  
 Pityriasis, [1537](#)  
 Pityriasis Alba, [1537](#)  
 Pityriasis Lichenoides, [1537](#)  
 Pityriasis Rosea, [1537](#)  
 Pityriasis Rubra Pilaris, [1537](#)  
 Pityriasis Versicolor, [1537](#)  
 Pivampicillin, [1537](#)  
 Pivot Joint, [1537](#)  
 Pixantrone, [1537](#)  
 PJI, [1538](#)  
 PJS, [1538](#)  
 PKD, [1538](#)  
 Placebo, [1538](#)  
 Placebo Control, [1538](#)  
 Placebo Effect, [1538](#)  
 Placenta, [1538](#)  
 Placenta Abruptio, [1538](#)  
 Placenta Accreta, [1538](#)  
 Placenta Accreta Spectrum, [1538](#)  
 Placenta Disorder, [1538](#)  
 Placenta Increta, [1538](#)  
 Placenta Percreta, [1539](#)  
 Placenta Praevia, [1539](#)  
 Placenta Previa, [1539](#)  
 Placental Abnormalities, [1538](#)  
 Placental Abruption, [1538](#)  
 Placental Accreta Spectrum, [1538](#)  
 Placental Hormone, [1539](#)  
 Placental Infarction, [1539](#)  
 Placental Insufficiency, [1539](#)  
 Placental Lactogen, [1539](#)  
 Placental Mesenchymal Dysplasia, [1539](#)  
 Placental Separation Signs, [1539](#)  
 Placental Site Nodule, [1539](#)  
 Placentation, [1539](#)  
 Plagiocephaly, [1539](#)  
 Plague, [1539](#)  
 Plain Radiography, [1539](#)  
 Planar, [1540](#)  
 Plane Joint, [1540](#)  
 Plant Fertilizer Poisoning, [1540](#)  
 Plant Poisoning, [1540](#)  
 Plant Thorn Arthritis, [1541](#)  
 Plant Thorn Synovitis, [1541](#)  
 Plantar, [1540](#)  
 Plantar Fasciitis, [1540](#)  
 Plantar Fibromatosis, [1540](#)  
 Plantar Flexion, [1540](#)  
 Plantar Hyperhidrosis, [1540](#)  
 Plantar Nerve, [1540](#)  
 Plantar Neuroma, [1540](#)  
 Plantar Reflex, [1540](#)  
 Plantar Response, [1540](#)  
 Plantar Wart, [1540](#)  
 Plantaris Muscle, [1540](#)  
 Plaque, [1541](#)  
 Plaque And Tartar On Teeth, [1541](#)  
 Plaque Assay, [1541](#)  
 Plaque Psoriasis, [1541](#)  
 Plasma, [1541](#)  
 Plasma / Urine Catecholamine Assay, [1542](#)  
 Plasma Amino Acids, [1541](#)  
 Plasma Cell, [1541](#)  
 Plasma Cell Mastitis, [1541](#)  
 Plasma Cell Myeloma, [1541](#)  
 Plasma Cell Neoplasm, [1541](#)  
 Plasma Cells, [1541](#)  
 Plasma Concentration, [1542](#)  
 Plasma Expanders, [1542](#)  
 Plasma Kallikrein, [1542](#)  
 Plasma Membrane, [1542](#)  
 Plasma Osmolarity, [1542](#)

---



---

Plasma Substitutes, 1542  
 Plasmablastic Lymphoma, 1541  
 Plasmacytoma, 1542  
 Plasmalogen, 1542  
 Plasmapheresis, 1542  
 Plasmid, 1542  
 Plasmids, 1543  
 Plasmin, 1543  
 Plasminogen, 1543  
 Plasminogen Activator, 1543  
 Plasminogen Inactivators, 1543  
 Plasmodium, 1543  
 Plasmodium Vivax, 1543  
 Plasmodium Vivax Malaria, 1543  
 Plaster Of Paris, 1543  
 Plastic and Reconstructive Surgery, 1543  
 Plastic Bronchitis, 1543  
 Plastic Casting Resin Poisoning, 1543  
 Plastic Resin Hardener Poisoning, 1543  
 Plastic Surgery, 1543  
 Platelet, 1544  
 Platelet Aggregation, 1544  
 Platelet Aggregation Test, 1544  
 Platelet Antibodies Blood Test, 1544  
 Platelet Count, 1544  
 Platelet Function Assay, 1544  
 Platelet Function Disorders, 1544  
 Platelet Plug Formation, 1544  
 Platelet-Rich Plasma, 1544  
 Platelets, 1544  
 Platybasia, 1544  
 Platypnea, 1544  
 Platypnea-Orthodeoxia Syndrome, 1544  
 Platysma, 1545  
 Pleiotropic Cytokine, 1545  
 Pleiotropy, 1545  
 Pleomorphic, 1545  
 Pleomorphic Adenoma, 1545  
 Pleomorphic Adenoma Of The Vulva, 1545  
 Pleomorphic Adenoma Pathology, 1545  
 Pleomorphic Dermal Sarcoma, 1545  
 Pleomorphic Xanthoastrocytoma, 1545  
 Pleomorphism, 1545  
 Plethoric, 1545  
 Plethysmography, 1545  
 Pleura, 1545  
 Pleural, 1545  
 Pleural Cavity, 1545  
 Pleural Decortication, 1546  
 Pleural Effusion, 1546  
 Pleural Fluid Analysis, 1546  
 Pleural Fluid Culture, 1546  
 Pleural Fluid Glucose, 1546  
 Pleural Fluid Gram Stain, 1546  
 Pleural Fluid PH, 1546  
 Pleural Fluid Smear, 1546  
 Pleural Friction Rub, 1546  
 Pleural Mesothelioma, 1546  
 Pleural Needle Biopsy, 1546  
 Pleural Neoplasm, 1546  
 Pleural Perforation, 1546  
 Pleural Plaque, 1547  
 Pleural Rub, 1547  
 Pleural Sac, 1547  
 Pleural Space, 1547  
 Pleural Tuberculosis, 1547  
 Pleurectomy, 1547  
 Pleurisy, 1547  
 Pleuritic Pain, 1547  
 Pleuritis, 1547  
 Pleurodesis, 1547  
 Pleurodynia, 1547  
 Pleuropulmonary Blastoma, 1547  
 Plexiform Fibrohistiocytic Tumor, 1547  
 Plexiform Neurofibroma, 1547  
 Plexus, 1548  
 Pliability, 1548  
 Plica, 1548  
 Plitidepsin, 1548  
 Ploidies, 1548  
 PLS, 1548  
 Plummer-Vinson Syndrome, 1548  
 Pluralibacter Gergoviae, 1548  
 Pluripotentia, 1548  
 Plutonium, 1548  
 Pm, 1548  
 PMBCL, 1548  
 PMBL, 1548  
 PMDD, 1548  
 PML, 1548  
 PMS, 1645  
 PN, 1548  
 Pneumatic, 1548  
 Pneumatic Otoscopy, 1549  
 Pneumatic Retinopexy, 1549  
 Pneumatic Tourniquet, 1549  
 Pneumatocele, 1549  
 Pneumatoxis, 1549  
 Pneumatoxis Cystoides Coli, 1549  
 Pneumatoxis Cystoides Intestinalis, 1549  
 Pneumatoxis Intestinalis, 1549  
 Pneumocephalus, 1549  
 Pneumococcal Conjugate Vaccine, 2128  
 Pneumococcal Infection, 1549  
 Pneumococcal Infections (Streptococcus pneumoniae), 1549  
 Pneumococcal Meningitis, 1549  
 Pneumococcal Pneumonia, 1549

---

---

Pneumococcal Sepsis, [1550](#)  
Pneumococcal Vaccine, [2128](#)  
Pneumoconiosis, [1550](#)  
Pneumocystis, [1550](#)  
Pneumocystis Infection, [1550](#)  
Pneumocystis Jirovecii Pneumonia, [1550](#)  
Pneumocystis Pneumonia, [1550](#)  
Pneumocyte, [1550](#)  
Pneumolysis, [1550](#)  
Pneumomediastinum, [1550](#)  
Pneumonectomy, [1550](#)  
Pneumonia, [1550](#)  
Pneumonia in Immunocompromised Patients, [1550](#)  
Pneumonitis, [1550](#)  
Pneumopericardium, [1551](#)  
Pneumoperitoneum, [1551](#)  
Pneumoplasty, [1551](#)  
Pneumotachometer, [1551](#)  
Pneumothorax, [1551](#)  
Pneumovirinae, [1551](#)  
Pneumovirus, [1551](#)  
PNF, [1551](#)  
Pocket Erosion, [1551](#)  
Podagra, [1551](#)  
Podiatrist, [1551](#)  
Podiatry, [1551](#)  
Podocyte, [1551](#)  
Podofilox, [1551](#)  
Podophyllin, [1551](#)  
Podophyllum, [1552](#)  
Podosome, [1552](#)  
POEMS Syndrome, [1552](#)  
Poikilocytosis, [1552](#)  
Poikiloderma, [1552](#)  
Poikilotherm, [1552](#)  
Poinsettia Plant Exposure, [1552](#)  
Point, [1552](#)  
Point Mutation, [1552](#)  
Point-Of-Care Testing, [1552](#)  
Point-Of-Care Ultrasound, [1552](#)  
Poison, [1552](#)  
Poison Control, [1552](#)  
Poison Control Emergency Services, [1553](#)  
Poison Ivy Rash, [1553](#)  
Poison Ivy, Oak, and Sumac, [1944](#)  
Poison-Center Consultation, [1552](#)  
Poisoning, [1553](#)  
Poisoning First Aid, [1553](#)  
Pokeweed Poisoning, [1553](#)  
Pol Gene, [1553](#)  
Poland Syndrome, [1553](#)  
Polar, [1553](#)  
Polarity, [1553](#)  
Polarography, [1553](#)  
Polidocanol, [1553](#)  
Polio, [1553](#)  
Poliomyelitis, [1554](#)  
Poliosis, [1554](#)  
Poliovirus, [1554](#)  
Poliovirus Vaccine, [2128](#)  
Pollakisuria, [1554](#)  
Pollakiuria, [1554](#)  
Pollen, [1554](#)  
Pollen-Food Allergy Syndrome, [1554](#)  
Pollex, [1554](#)  
Pollution, [1554](#)  
Poly A, [1554](#)  
Polyadenylation, [1554](#)  
Polyamine, [1554](#)  
Polyamine Catabolism, [1554](#)  
Polyarteritis, [1554](#)  
Polyarteritis Nodosa, [1554](#)  
Polyarthritis, [1555](#)  
Polyarticular Joint Pain, [1555](#)  
Polybrominated Biphenyl, [1555](#)  
Polychlorinated Biphenyl, [1555](#)  
Polychondritis, [1555](#)  
Polychromasia, [1555](#)  
Polychromatophilia, [1555](#)  
Polycystic Kidney Disease, [1555](#)  
Polycystic Liver Disease, [1555](#)  
Polycystic Ovarian Syndrome, [1555](#)  
Polycystic Ovary, [1555](#)  
Polycystic Ovary Syndrome, [1555](#)  
Polycystic Ovary Syndrome (PCOS), [1647](#)  
Polycythaemia, [1556](#)  
Polycythemia, [1556](#)  
Polycythemia Vera, [1556](#)  
Polydactyly, [1556](#)  
Polydipsia, [1556](#)  
Polyene, [1556](#)  
Polyethylene, [1556](#)  
Polyethylene Glycol, [1556](#)  
Polygenic, [1556](#)  
Polygenic Disease, [1556](#)  
Polygenic Hypercholesterolemia, [1556](#)  
Polygenic Inheritance, [1556](#)  
Polygenic Obesity, [1557](#)  
Polygenic Risk Score, [1557](#)  
Polyglandular Autoimmune Syndrome, [1557](#)  
Polyglandular Autoimmune Syndrome Type 1, [1557](#)  
Polyglandular Autoimmune Syndrome Type 2, [1557](#)  
Polygraphy, [1557](#)  
Polyhydramnios, [1557](#)  
Polyketide Synthase, [1557](#)  
Polymastia, [1557](#)  
Polymenorrhea, [1557](#)  
Polymenorrhoea, [1557](#)

---

---

Polymer, [1557](#)  
Polymerase Chain Reaction, [1557](#)  
Polymerization, [1557](#)  
Polymicrobial Infections, [1558](#)  
Polymicrogyria, [1558](#)  
Polymorphic, [1558](#)  
Polymorphic Eruption of Pregnancy, [1558](#)  
Polymorphic Light Eruption, [1558](#)  
Polymorphism, [1558](#)  
Polymorphonuclear Leukocyte, [1558](#)  
Polymorphous, [1558](#)  
Polymorphous Light Eruption, [1558](#)  
Polymyalgia Rheumatica, [1558](#)  
Polymyositis, [1558](#)  
Polymyxin, [1558](#)  
Polymyxin B, [1558](#)  
Polyneuritis, [1558](#)  
Polyneuropathy, [1559](#)  
Polynucleotide, [1559](#)  
Polyoma Virus, [1559](#)  
Polyomavirus Hominis 2, [1559](#)  
Polyp, [1559](#)  
Polyp Biopsy, [1559](#)  
Polyp Surveillance Colonoscopy, [1559](#)  
Polypeptide, [1559](#)  
Polyphagia, [1559](#)  
Polypharmacology, [1559](#)  
Polypharmacy, [1559](#)  
Polyposis, [1559](#)  
Polypropylene, [1559](#)  
Polyradiculitis, [1559](#)  
Polyradiculopathy, [1559](#)  
Polysaccharide, [1559](#)  
Polysome, [1560](#)  
Polysomnography, [1560](#)  
Polysplenia, [1560](#)  
Polystyrene, [1560](#)  
Polytene Chromosome, [1560](#)  
Polythelia, [1560](#)  
Polythiazide, [1560](#)  
Polytrauma, [1560](#)  
Polyunsaturated Fat, [1560](#)  
Polyurethane, [1560](#)  
Polyuria, [1560](#)  
Polyvalent, [1560](#)  
Polyvinyl, [1560](#)  
Pomalidomide, [1560](#)  
Pompe Disease, [1560](#)  
Pompholyx, [1561](#)  
Pompholyx Eczema, [1561](#)  
Pons, [1561](#)  
Pontiac Fever, [1561](#)  
Pontine, [1561](#)  
Pontine Myelinolysis, [1561](#)  
Poor Circulation, [1561](#)  
Poor Color Vision, [1561](#)  
Popcorn Lung Bronchiolitis Obliterans, [1561](#)  
Popliteal, [1561](#)  
Popliteal Aneurysm, [1561](#)  
Popliteal Artery, [1561](#)  
Popliteal Artery Aneurysm, [1561](#)  
Popliteal Artery Entrapment Syndrome, [1561](#)  
Popliteal Cyst, [1562](#)  
Popliteal Fossa, [1562](#)  
Popliteal Vein, [1562](#)  
Population Attributable Fraction, [1562](#)  
Population Attributable Risk, [1562](#)  
Population Genetics, [1562](#)  
Population Group, [1562](#)  
Porcelain Gallbladder, [1562](#)  
Porcelain Inlay, [1645](#)  
Porcine, [1562](#)  
Pore, [1562](#)  
Porencephaly, [1562](#)  
Porfimer Sodium, [1562](#)  
Porin, [1562](#)  
Pork Tapeworm, [1562](#)  
Porokeratosis, [1563](#)  
Poroma, [1563](#)  
Porphyria, [1563](#)  
Porphyria Cutanea Tarda, [1563](#)  
Porphyrias, [1563](#)  
Porphyrin, [1563](#)  
Porphyrins Blood Test, [1563](#)  
Porphyrins Urine Test, [1563](#)  
Porphyromonas, [1563](#)  
Porphyromonas Bennonis, [1563](#)  
Porphyromonas Catoniae, [1563](#)  
Porphyromonas Endodontalis, [1563](#)  
Porphyromonas Levii, [1563](#)  
Porphyromonas Somerae, [1563](#)  
Port-Wine Stain, [1564](#)  
Portacaval Shunting, [1563](#)  
Portal Circulation, [1564](#)  
Portal Hypertension, [1564](#)  
Portal Hypertensive Colopathy, [1564](#)  
Portal Hypertensive Gastropathy, [1564](#)  
Portal System, [1564](#)  
Portal Vein, [1564](#)  
Portal Vein Obstruction, [1564](#)  
Portal Vein Thrombosis, [1564](#)  
Portion Of Cytosol, [1564](#)  
Portion Size, [1564](#)  
Portopulmonary Syndrome, [1564](#)  
Position, [1564](#)  
Positional Asphyxia, [1565](#)  
Positive, [1565](#)  
Positive Airway Pressure Treatment, [1565](#)

---

---

Positive End-Expiratory Pressure, [1565](#)  
Positive Feedback, [1565](#)  
Positive Finding, [1565](#)  
Positive Predictive Value, [1565](#)  
Positive Predictive Value In Surveillance, [1565](#)  
Positive Pressure Ventilation, [1565](#)  
Positive Psychology, [1565](#)  
Positive Symptoms, [1565](#)  
Positive-Strand RNA Virus, [1565](#)  
Positron, [1565](#)  
Positron Camera, [1565](#)  
Positron Emission Mammography, [1565](#)  
Positron Emission Tomography, [1566](#)  
Post Cibus, [1566](#)  
Post-Acute Rehabilitation, [1566](#)  
Post-Adoptive Depression, [1566](#)  
Post-Cardiac Arrest Management, [1566](#)  
Post-Cardiac Injury Syndrome, [1566](#)  
Post-Chemotherapy Cognitive Impairment, [1566](#)  
Post-Concussion Syndrome, [1566](#)  
Post-Concussional Syndrome, [1566](#)  
Post-Dural Puncture Headache, [1566](#)  
Post-Exposure Prophylaxis, [1568](#)  
Post-Finasteride Syndrome, [1569](#)  
Post-Hoc, [1569](#)  
Post-Infectious Cough, [1569](#)  
Post-Intensive Care Syndrome, [1569](#)  
Post-Kala-Azar Dermal Leishmaniasis, [1569](#)  
Post-Mortem, [1570](#)  
Post-Myocardial Infarction Pericarditis, [1570](#)  
Post-Nasal Drip, [1570](#)  
Post-Natal Depression, [1571](#)  
Post-Obstructive Pneumonia, [1571](#)  
Post-Operative Atelectasis, [1571](#)  
Post-Orgasmic Illness Syndrome, [1572](#)  
Post-Partum Haemorrhage, [1572](#)  
Post-Pneumonectomy Syndrome, [1573](#)  
Post-Polio Syndrome, [1573](#)  
Post-Prandial, [1573](#)  
Post-Splenectomy Complications, [1574](#)  
Post-Streptococcal Glomerulonephritis, [1574](#)  
Post-Surgical Pain, [1574](#)  
Post-Surgical Rehabilitation, [1574](#)  
Post-Term Infant, [1574](#)  
Post-Term Pregnancy, [1574](#)  
Post-Test Probability, [1574](#)  
Post-Thrombotic Syndrome, [1574](#)  
Post-Transfusion Purpura, [1575](#)  
Post-Traumatic Headache, [1575](#)  
Post-Traumatic Stress Disorder, [1575](#)  
Post-Traumatic Stress Disorder (PTSD), [1575](#)  
Post-Vasectomy Pain Syndrome, [1575](#)  
Post-Void Residual, [1575](#)  
Post-Void Residual Volume, [1575](#)  
Postanesthesia Care Unit, [1566](#)  
Postcholecystectomy Syndrome, [1566](#)  
Postcoital Bleeding, [1566](#)  
Postcoital Test, [1645](#)  
Posterior, [1567](#)  
Posterior Chamber, [1567](#)  
Posterior Cord Syndrome, [1567](#)  
Posterior Cruciate, [1567](#)  
Posterior Cruciate Ligament Injury, [1567](#)  
Posterior Cruciate Ligament Tear, [1567](#)  
Posterior Descending Artery, [1567](#)  
Posterior Drawer Test, [1567](#)  
Posterior Epistaxis, [1567](#)  
Posterior Fossa Tumor, [1567](#)  
Posterior Heel Bursitis, [1567](#)  
Posterior Keyhole Foraminotomy, [1567](#)  
Posterior Mediastinum, [1567](#)  
Posterior Myocardial Infarction, [1568](#)  
Posterior Pituitary, [1568](#)  
Posterior Prolapse, [1568](#)  
Posterior Reversible Encephalopathy Syndrome, [1568](#)  
Posterior Rhinorrhea, [1568](#)  
Posterior Subcapsular Cataract, [1568](#)  
Posterior Synechiae, [1568](#)  
Posterior Tibial Artery, [1568](#)  
Posterior Urethral Valves, [1568](#)  
Posterior Uveitis, [1568](#)  
Posterior Vaginal Prolapse, [1568](#)  
Posterior Vitreous Detachment, [1568](#)  
Posteroanterior, [1568](#)  
Postganglionic Fiber, [1569](#)  
Postherpetic Neuralgia, [1569](#)  
Posthitis, [1569](#)  
Posthumous, [1569](#)  
Postinflammatory Hyperpigmentation, [1569](#)  
Postmenopausal, [1569](#)  
Postmenopausal Bleeding, [1569](#)  
Postmenopausal Osteoporosis, [1569](#)  
Postmenopause, [1569](#)  
Postmortem Artifact, [1570](#)  
Postmortem Changes, [1570](#)  
Postmortem Interval, [1570](#)  
Postmortem Lividity, [1570](#)  
Postmortem Toxicology, [1570](#)  
Postmortem Vitreous Analysis, [1570](#)  
Postmyocardial Infarction Syndrome, [1570](#)  
Postnatal, [1570](#)  
Postnatal Anxiety, [1570](#)  
Postnatal Care, [1570](#)  
Postnatal Depression, [1570](#)  
Postop, [1571](#)  
Postoperative, [1571](#)  
Postoperative Care, [1571](#)  
Postoperative Cognitive Dysfunction, [1571](#)

---

---

Postoperative Complications, 1571  
 Postoperative Corneal Edema, 1571  
 Postoperative Delirium, 1571  
 Postoperative Hemorrhage, 1571  
 Postoperative Ileus, 1571  
 Postoperative Infection, 1571  
 Postoperative Nausea And Vomiting, 1571  
 Postoperative Period, 1572  
 Postoperative Pulmonary Complication, 1572  
 Postpartum, 1572  
 Postpartum Anxiety, 1572  
 Postpartum Blues, 1572  
 Postpartum Care, 1572  
 Postpartum Contraception, 1572  
 Postpartum Depression, 1572  
 Postpartum Endometritis, 1572  
 Postpartum Hemorrhage, 1572  
 Postpartum Infection, 1572  
 Postpartum Infections, 1573  
 Postpartum Preeclampsia, 1573  
 Postpartum Psychosis, 1573  
 Postpartum Thyroiditis, 1573  
 Postpartum Urinary Retention, 1573  
 Postpericardiotomy Syndrome, 1573  
 Postprandial Period, 1573  
 Postrenal AKI, 1573  
 Poststreptococcal Glomerulonephritis, 1574  
 Poststreptococcal Pustulosis, 1574  
 Postsynaptic Membrane, 1574  
 Postsynaptic Potential, 1574  
 Postterm Pregnancy, 1574  
 Posttraumatic Stress Disorder (PTSD), 1645  
 Posttraumatic Stress Syndrome, 1645  
 Postural, 1575  
 Postural Drainage, 1575  
 Postural Hypotension, 1575  
 Postural Orthostatic Tachycardia Syndrome, 1575  
 Postural Tremor, 1575  
 Posture, 1575  
 Potassium, 1576  
 Potassium Acetate, 1576  
 Potassium Binders, 1576  
 Potassium Carbonate Poisoning, 1576  
 Potassium Channel, 1576  
 Potassium Chloride, 1576  
 Potassium Citrate, 1576  
 Potassium Cyanide, 1576  
 Potassium Deficiency, 1576  
 Potassium Dichromate, 1576  
 Potassium Hydroxide Poisoning, 1576  
 Potassium In Diet, 1576  
 Potassium Iodide, 1576  
 Potassium Ion, 1576  
 Potassium Permanganate, 1577  
 Potassium Test, 1577  
 Potassium Urine Test, 1577  
 Potassium-Sparing Diuretic, 1577  
 Potassium-Sparing Diuretics, 1577  
 Potbellies And Toddlers, 1577  
 Potency, 1577  
 Potential, 1577  
 Potential Acuity Meter, 1577  
 Potocki-Lupski Syndrome, 1577  
 Pott Disease, 1577  
 Pott Puffy Tumor, 1577  
 Pott's Disease, 1578  
 Potter Sequence, 1577  
 Potter Syndrome, 1577  
 Pouch, 1578  
 Pouch of Douglas, 1578  
 Pound, 1578  
 Povidone, 1578  
 Povidone-Iodine, 1578  
 Powassan Encephalitis, 1578  
 Powassan Virus, 1578  
 Powassan Virus Disease, 1578  
 Powder Diffraction, 1578  
 Power, 1578  
 Power Training, 1578  
 Pox, 1578  
 Poxviridae, 1578  
 Poxviridae Infection, 1578  
 Poxvirus, 1579  
 Ppar Alpha, 1579  
 Ppar Delta, 1579  
 PPD, 1579  
 PPD Skin Test, 1579  
 PPH, 1647  
 PPS (Post-Polio Syndrome), 1646  
 PR Interval, 1596  
 Practitioner, 1579  
 Prader Willi Syndrome, 1646  
 Prader-Labhart-Willi Fancone Syndrome, 1647  
 Prader-Labhart-Willi Syndrome, 1579  
 Prader-Willi Syndrome, 1579  
 Pragmatic, 1579  
 Pragmatic Trial, 1579  
 Pralatrexate, 1579  
 Pralidoxime, 1579  
 Pramipexole, 1579  
 Prandial, 1579  
 Prasugrel, 1579  
 Pravastatin, 1579  
 Pravastatin Sodium, 1580  
 Praziquantel, 1580  
 Prazosin, 2128  
 Pre-Cancerous, 1580  
 Pre-eclampsia, 1582

---

---

Pre-Existing Diabetes And Pregnancy, [1582](#)  
Pre-Medication, [2128](#)  
Pre-MiRNA, [1586](#)  
Pre-Synaptic, [1590](#)  
Prealbumin, [1580](#)  
Prealbumin Tyr-77 Amyloidosis, [1646](#)  
Preauricular Sinuses, [1580](#)  
Prebiotics, [1580](#)  
Prebiotics And Probiotics, [1580](#)  
Precancerous Condition, [1580](#)  
Precancerous Polyp, [1580](#)  
Precapillary Sphincter, [1580](#)  
Precaution, [1580](#)  
Precipitate Labor, [1580](#)  
Precipitating Factor, [1581](#)  
Precision Medicine, [1581](#)  
Precision Oncology, [1581](#)  
Preclinical, [1581](#)  
Precocious Puberty, [1581](#)  
Preconception Care, [1581](#)  
Preconception Counseling, [1581](#)  
Precordial Catch Syndrome, [1581](#)  
Precordium, [1581](#)  
Precursor, [1581](#)  
Prediabetes, [1581](#)  
Prediabetes Syndrome, [1581](#)  
Predictive Test, [1581](#)  
Predispose, [1581](#)  
Prednimustine, [1582](#)  
Prednisolone, [1582](#)  
Prednisone, [1582](#)  
Preeclampsia, [1582](#)  
Preeclampsia and Eclampsia, [1582](#)  
Preeclampsia And High Blood Pressure, [1645](#)  
Preeclampsia With Severe Features, [1582](#)  
Preexcitation Syndrome, [1582](#)  
Prefrontal, [1582](#)  
Prefrontal Cortex, [1582](#)  
Pregabalin, [1582](#)  
Preganglionic Fiber, [1582](#)  
Pregnancy, [1582](#)  
Pregnancy And Herpes, [1582](#)  
Pregnancy And Travel, [1583](#)  
Pregnancy and Urolithiasis, [1583](#)  
Pregnancy And Work, [1583](#)  
Pregnancy at Age 35 or Older, [1583](#)  
Pregnancy Complication, [1583](#)  
Pregnancy Diagnosis, [1583](#)  
Pregnancy Nutrition, [1583](#)  
Pregnancy Of Unknown Location, [1583](#)  
Pregnancy Outcome, [1583](#)  
Pregnancy Status, [1584](#)  
Pregnancy Test, [1584](#)  
Pregnancy Trauma, [1584](#)  
Pregnancy Trimester, [1584](#)  
Pregnancy With Asthma, [1645](#)  
Pregnancy, Cervical, [363](#)  
Pregnancy, Ectopic, [616](#)  
Pregnancy, Extrauterine, [702](#)  
Pregnancy, Tubal, [1971](#)  
Pregnancy-Induced Hypertension, [1583](#)  
Pregnancy-Related Hypertension, [1584](#)  
Pregnancy: Bleeding During The First Trimester, [1583](#)  
Pregnancy: Placenta Previa, [1583](#)  
Pregnancy: Preeclampsia And Eclampsia, [1583](#)  
Pregnane, [1584](#)  
Pregnanolone, [1584](#)  
Pregnant, [1584](#)  
Pregnenolone, [1584](#)  
Prehn Sign, [1584](#)  
Prehypertension, [1584](#)  
Preimplantation Genetic Diagnosis, [1584](#)  
Preimplantation Genetic Testing, [1584](#)  
Preload, [1584](#)  
Preload Reduction, [1585](#)  
Premalignant, [1585](#)  
Premalignant Lesions of the Endometrium, [1585](#)  
Premature, [1585](#)  
Premature Adrenarche, [1585](#)  
Premature Atrial Contraction, [1585](#)  
Premature Baby, [1585](#)  
Premature Birth, [1585](#)  
Premature Closure, [1585](#)  
Premature Ejaculation, [1585](#)  
Premature Infant, [1585](#)  
Premature Labor, [1585](#)  
Premature Menopause, [1585](#)  
Premature Ovarian Failure, [1585](#)  
Premature Ovarian Insufficiency, [1586](#)  
Premature Puberty, [1586](#)  
Premature Rupture Of Membranes, [1586](#)  
Premature Thelarche, [1586](#)  
Premature Ventricular Contraction, [1586](#)  
Premature Ventricular Contractions, [1586](#)  
Prematurity, [1586](#)  
Premenstrual, [1586](#)  
Premenstrual Breast Changes, [1586](#)  
Premenstrual Dysphoric Disorder, [1586](#)  
Premenstrual Syndrome, [1586](#)  
Premenstrual Syndrome (PMS), [1645](#)  
Premolar Tooth, [1586](#)  
Prenatal, [1586](#)  
Prenatal Care, [1587](#)  
Prenatal Depression, [1587](#)  
Prenatal Development, [1587](#)  
Prenatal Diagnosis, [1587](#)  
Prenatal Exposure, [1587](#)  
Prenatal Genetic Counseling, [1587](#)

---



---

Prenatal Testing, [1587](#)  
 Prenatal Vitamins, [1587](#)  
 Prenylation, [1587](#)  
 Preop, [1587](#)  
 Preoperative Assessment, [1587](#)  
 Preoperative Care, [1587](#)  
 Preoperative Evaluation, [1587](#)  
 Preoptic Area, [1588](#)  
 Prepubertal, [1588](#)  
 Prepuce, [1588](#)  
 Prerenal Azotemia, [1588](#)  
 Presbycusis, [1588](#)  
 Presbycusis, [1588](#)  
 Presbyopia, [1588](#)  
 Presbyphonia, [1588](#)  
 Preschooler Development, [1588](#)  
 Preschooler Test Or Procedure Preparation, [1588](#)  
 Prescription, [1588](#)  
 Prescription Drug, [2128](#)  
 Prescription Drug Misuse, [1588](#)  
 Prescription Medications, [1588](#)  
 Presenilin, [1589](#)  
 Presentation, [1589](#)  
 Preseptal Cellulitis, [1589](#)  
 Preservation, [1589](#)  
 Preservative, [1589](#)  
 Pressor, [1589](#)  
 Pressoreceptor, [1589](#)  
 Pressure, [1589](#)  
 Pressure Atrophy, [1589](#)  
 Pressure Injury, [1589](#)  
 Pressure Sore, [1589](#)  
 Pressure Sores, [1646](#)  
 Pressure Support Ventilation, [1589](#)  
 Pressure Ulcer, [1589](#)  
 Pressure Ulcer Prevention, [1590](#)  
 Pressure Urticaria, [1590](#)  
 Pressure-Flow Study, [1589](#)  
 Presynaptic Terminals, [1590](#)  
 Presyncope, [1590](#)  
 Preterm Birth, [1590](#)  
 Preterm Infant, [1590](#)  
 Preterm Labor, [1590](#)  
 Preterm Labor and Birth, [1590](#)  
 Preterm Premature Rupture Of Membranes, [1590](#)  
 Pretest Probability, [1590](#)  
 Pretibial Myxedema, [1590](#)  
 Pretibial Myxoedema, [1591](#)  
 Prevalence, [1591](#)  
 Prevention Of Liver Cancer, [1645](#)  
 Preventive Dentistry, [1591](#)  
 Preventive Health Care, [1591](#)  
 Preventive Medicine, [1591](#)  
 Prevertebral Air, [1591](#)  
 Priapism, [1591](#)  
 Prick Test, [1591](#)  
 Prickly Heat, [1591](#)  
 Primary, [1591](#)  
 Primary Adrenal Insufficiency, [1591](#)  
 Primary Adrenal Lymphoma, [1591](#)  
 Primary Aldosteronism, [1591](#)  
 Primary Alveolar Hypoventilation, [1591](#)  
 Primary Amebic Meningoencephalitis, [1591](#)  
 Primary Amenorrhea, [1592](#)  
 Primary Amyloidosis, [1592](#)  
 Primary and Secondary Fibrinolysis, [1592](#)  
 Primary And Secondary Hyperaldosteronism, [1592](#)  
 Primary Angioplasty, [1592](#)  
 Primary Biliary Cholangitis, [1592](#)  
 Primary Biliary Cirrhosis, [1592](#)  
 Primary Biliary Cirrhosis (PBC), [1646](#)  
 Primary Biliary Cirrhosis Treatment, [1592](#)  
 Primary Care, [1592](#)  
 Primary Ciliary Dyskinesia, [1592](#)  
 Primary Ciliary Dyskinesia (Kartagener Syndrome), [1592](#)  
 Primary CNS Lymphoma, [1593](#)  
 Primary Cutaneous Amyloidosis, [1646](#)  
 Primary Cutaneous B-Cell Lymphoma, [1593](#)  
 Primary Endpoint, [1593](#)  
 Primary Hyperaldosteronism, [1593](#)  
 Primary Hyperoxaluria, [1593](#)  
 Primary Hyperparathyroidism, [1593](#)  
 Primary Hypersomnia, [1593](#)  
 Primary Hypertension, [1593](#)  
 Primary Hypertrophic Osteoarthropathy, [1593](#)  
 Primary Hypoparathyroidism, [1593](#)  
 Primary Hypothyroidism, [1593](#)  
 Primary Immune Response, [1594](#)  
 Primary Immunodeficiency, [1594](#)  
 Primary Immunodeficiency Disease, [1594](#)  
 Primary Immunodeficiency Disease PIDD, [1594](#)  
 Primary Lactase Deficiency, [1594](#)  
 Primary Lateral Sclerosis, [1594](#)  
 Primary Lymphoid Organs, [1594](#)  
 Primary Lymphoma Of The Brain, [1594](#)  
 Primary Metabolism, [1594](#)  
 Primary Microrna, [1594](#)  
 Primary Myelofibrosis, [1594](#)  
 Primary Open-Angle Glaucoma, [1594](#)  
 Primary Osteoporosis, [1594](#)  
 Primary Ovarian Insufficiency, [1594](#)  
 Primary Polycythemia, [1595](#)  
 Primary Prevention, [1595](#)  
 Primary Progressive Aphasia, [1595](#)  
 Primary Progressive Multiple Sclerosis, [1595](#)  
 Primary Sclerosing Cholangitis, [1595](#)  
 Primary Systemic Amyloidosis, [1595](#)

---

---

Primary Tuberculosis, [1595](#)  
Primary Tumor, [1595](#)  
Primary Vulvar Clear Cell Adenocarcinoma, [1592](#)  
Primidone, [1595](#)  
Primigravida, [1595](#)  
Primipara, [1595](#)  
Primitive, [1595](#)  
Primitive Node, [1595](#)  
Primitive Streak, [1595](#)  
Primordial, [1595](#)  
Primordial Prevention, [1596](#)  
Prinzmetal Angina, [1596](#)  
Prion, [1596](#)  
Prion Disease, [1596](#)  
Prism, [1596](#)  
Privacy, [1596](#)  
Privacy Breach, [1596](#)  
Private Mutation, [1597](#)  
PRK, [1597](#)  
Pro Re Nata, [1607](#)  
Probe, [1597](#)  
Probenecid, [1597](#)  
Probiotic, [1597](#)  
Probiotics, [1597](#)  
Probucol, [1597](#)  
PROC, [1597](#)  
Procainamide, [1597](#)  
Procaine, [1597](#)  
Procalcitonin, [1597](#)  
Procalcitonin-Guided Antibiotic Therapy, [2128](#)  
Procarbazine, [1597](#)  
Procarbazine Hydrochloride, [1597](#)  
Procedural Memory, [1597](#)  
Procedural Sedation, [1597](#)  
Procedure, [1598](#)  
Procentriole, [1598](#)  
Procerus Muscle, [1598](#)  
Process, [1598](#)  
Processing Bodies, [1598](#)  
Prochlorperazine, [1598](#)  
Prochlorperazine Overdose, [1598](#)  
Procidentia, [1598](#)  
Procollagen, [1598](#)  
Proctalgia, [1598](#)  
Proctalgia Fugax, [1598](#)  
Proctitis, [1598](#)  
Proctocolitis, [1599](#)  
Proctology, [1599](#)  
Proctoscope, [1599](#)  
Proctoscopy, [1599](#)  
Proctostomy, [1599](#)  
Prodromal, [1599](#)  
Prodrome, [1599](#)  
Prodrug, [1599](#)  
Productive Cough, [1599](#)  
Profunda, [1599](#)  
Progenitor, [1599](#)  
Progeny, [1599](#)  
Progeria, [1599](#)  
Progestational, [1600](#)  
Progesterone, [1600](#)  
Progesterone Antagonist, [1600](#)  
Progesterone Receptor, [1600](#)  
Progesterone Supplementation, [1600](#)  
Progesterone Test, [1600](#)  
Progestin, [1600](#)  
Progestogen, [1600](#)  
Progestogen-Only Pill, [1600](#)  
Proglottid, [1600](#)  
Prognathism, [1600](#)  
Prognosis, [1600](#)  
Prognostic Factor, [1600](#)  
Prognostic Test, [1601](#)  
Programmed Cell Death, [1601](#)  
Progression, [1601](#)  
Progression-Free Survival, [1601](#)  
Progressive, [1601](#)  
Progressive Disease, [1601](#)  
Progressive Lenticular Degeneration, [1601](#)  
Progressive Multifocal Leukoencephalopathy, [1601](#)  
Progressive Muscular Atrophy, [1601](#)  
Progressive Supranuclear Palsy, [1601](#)  
Proinflammatory, [1601](#)  
Proinsulin, [1601](#)  
Projectile Vomiting, [1601](#)  
Projection, [1602](#)  
Prokaryote, [1602](#)  
Prokinetic, [1602](#)  
Prolactin, [1602](#)  
Prolactin Blood Test, [1602](#)  
Prolactin Deficiency, [1602](#)  
Prolactin Physiology, [1602](#)  
Prolactin Receptor, [1602](#)  
Prolactinoma, [1602](#)  
Prolactinoma Treatment, [1602](#)  
Prolapse, [1602](#)  
Prolapse Of Orbital Fat, [1603](#)  
Prolapsed Bladder, [1602](#)  
Prolapsed Cord, [1602](#)  
Prolapsed Disk, [1602](#)  
Prolapsed Uterus, [1602](#)  
Proliferating Pilar Tumor, [1603](#)  
Proliferating Trichilemmal Tumor, [1603](#)  
Proliferation, [1603](#)  
Proliferative, [1603](#)  
Proliferative Diabetic Retinopathy, [1603](#)  
Proliferative Fasciitis, [1603](#)  
Proliferative Myositis, [1603](#)

---

---

Proliferative Retinopathy, 1603  
Proline, 1603  
Prolonged Capillary Refill Time, 1603  
Prolonged Deceleration, 1603  
Prolonged Second Stage, 1603  
Polymorphocyte Count, 1603  
Polymorphocytic Leukemia, 1604  
Promazine, 1604  
Promecarb, 1604  
Prometaphase, 1604  
Promethazine, 1604  
Promethazine Overdose, 1604  
Prominent Iris Collarette, 1604  
Promonocyte Count, 1604  
Promontory, 1604  
Promoter, 1604  
Promoters, 1604  
Promyelocytes, 1604  
Pronation, 1604  
Pronator, 1604  
Pronator Drift, 1605  
Prone, 1605  
Prone Position, 1605  
Prone Positioning, 1605  
Pronephric, 1605  
Pronephros, 1605  
Propafenone, 1605  
Propane, 1605  
Propane Poisoning, 1605  
Propanol, 1605  
Propanolol, 1605  
Propantheline, 1605  
Propellant, 1605  
Properdin, 1605  
Properdin Deficiency, 1605  
Propetamphos, 1606  
Prophage, 1606  
Prophase, 1606  
Prophenofos, 1606  
Prophylactic, 1606  
Prophylactic Mastectomy, 1606  
Prophylaxis, 1606  
Propionates, 1606  
Propionibacterium, 1606  
Propionic Acidemia, 1606  
Propofol, 1606  
Propofol Infusion Syndrome, 1606  
Propoxur, 1606  
Propoxycaine, 1606  
Propoxyphene, 1606  
Propranolol, 1607  
Proprioception, 1607  
Proprioceptor, 1607  
Proptosis, 1607  
Proptosis of Eye, 1607  
Propulsion, 1607  
Propyl Alcohol, 1607  
Propylene Glycol, 1607  
Propylthiouracil, 1607  
Prosencephalon, 1607  
Prosopagnosia, 1607  
Prosoplasia, 1607  
Prosoposchisis, 1607  
Prostacyclin Analogs, 1608  
Prostaglandin, 1608  
Prostaglandin Analogs, 1608  
Prostaglandin Antagonist, 2129  
Prostaglandin Degradation, 1608  
Prostaglandin E2, 1608  
Prostaglandin F2-Alpha, 1608  
Prostaglandin Receptor, 1608  
Prostaglandins, 1608  
Prostaglandins E, 1608  
Prostaglandins F, 1608  
Prostate, 1608  
Prostate Adenocarcinoma, 1608  
Prostate Biopsy, 1608  
Prostate Brachytherapy, 1608  
Prostate Cancer, 1609  
Prostate Cancer Screening, 1609  
Prostate Cancer Staging, 1609  
Prostate Cancer Treatment, 1609  
Prostate Carcinoma, 1609  
Prostate Disease, 1609  
Prostate Enlargement, 1609  
Prostate Enlargement/BPH, 1647  
Prostate Gland, 1609  
Prostate Gland Enlargement, 1609  
Prostate Infection, 1609  
Prostate Neoplasm, 1609  
Prostate-Specific Antigen, 1609  
Prostate-Specific Antigen Blood Test, 1609  
Prostatectomy, 1609  
Prostatic Fluid, 1609  
Prostatic Obstruction, 1610  
Prostatic Urethral Stent, 1610  
Prostatism, 1610  
Prostatitis, 1610  
Prostatodynia, 1610  
Prosthesis, 1610  
Prosthetic, 1610  
Prosthetic Graft, 1610  
Prosthetic Heart Valve, 1610  
Prosthetic Limb, 1610  
Prosthetic Training, 1610  
Prosthetic Valve Endocarditis, 1610  
Prosthetics, 1610  
Prosthodontics, 1611

---

---

Protamine, [1611](#)  
Protanopia, [1611](#)  
Protean, [1611](#)  
Proteaphagy, [1611](#)  
Protease, [1611](#)  
Protease Inhibitor, [2129](#)  
Protease Inhibitors, [1611](#)  
Proteasome, [1611](#)  
Proteasome Binding, [1611](#)  
Proteasome Inhibitor, [1611](#)  
Protective, [1611](#)  
Protective Agent, [1611](#)  
Protegrin, [1611](#)  
Protein, [1611](#)  
Protein Acetylation, [1612](#)  
Protein Binding, [1612](#)  
Protein Biosynthesis, [1612](#)  
Protein C, [1612](#)  
Protein C Blood Test, [1612](#)  
Protein C Deficiency, [1612](#)  
Protein Deficiency, [1612](#)  
Protein Denaturation, [1612](#)  
Protein Dephosphorylation, [1612](#)  
Protein Electrophoresis, [1612](#)  
Protein Engineering, [1612](#)  
Protein Expression, [1612](#)  
Protein Folding, [1612](#)  
Protein Glycosylation, [1613](#)  
Protein Import, [1613](#)  
Protein In Diet, [1613](#)  
Protein Isoform, [1613](#)  
Protein Kinase, [1613](#)  
Protein Malnutrition, [1613](#)  
Protein Metabolism, [1613](#)  
Protein Methylation, [1613](#)  
Protein Overexpression, [1613](#)  
Protein Phosphorylation, [1613](#)  
Protein Prenylation, [1613](#)  
Protein Purification, [1613](#)  
Protein Refolding, [1613](#)  
Protein S, [1613](#)  
Protein S Blood Test, [1614](#)  
Protein S Deficiency, [1614](#)  
Protein Secretion, [1614](#)  
Protein Sequence, [1614](#)  
Protein Splicing, [1614](#)  
Protein Subunit, [1614](#)  
Protein Synthesis, [1614](#)  
Protein Translocation, [1614](#)  
Protein Transport, [1614](#)  
Protein Unfolding, [1614](#)  
Protein-Energy Malnutrition, [1612](#)  
Protein-Losing Enteropathy, [1613](#)  
Proteinaceous Plug, [1612](#)  
Proteinuria, [1614](#)  
Proteobacteria, [1614](#)  
Proteogenomics, [1614](#)  
Proteoglycan, [1614](#)  
Proteolipid Protein, [1614](#)  
Proteolipids, [1615](#)  
Proteolysis, [1615](#)  
Proteolytic, [1615](#)  
Proteome, [1615](#)  
Proteomic Profile, [1615](#)  
Proteomics, [1615](#)  
Proteostasis, [1615](#)  
Proteus, [1615](#)  
Proteus Hauseri, [1615](#)  
Proteus Mirabilis, [1615](#)  
Proteus Penneri, [1615](#)  
Proteus Syndrome, [1615](#)  
Proteus Vulgaris, [1615](#)  
Prothionamide, [1615](#)  
Prothrombin, [1615](#)  
Prothrombin Deficiency, [1615](#)  
Prothrombin Time, [1616](#)  
Proto-Oncogene, [1616](#)  
Proton Pump Inhibitor, [2129](#)  
Proton Pump Inhibitors, [1616](#)  
Proton Therapy, [1616](#)  
Protozoa, [1616](#)  
Protozoal Infection, [1616](#)  
Protractile, [1616](#)  
Protractor, [1616](#)  
Protrusion, [1616](#)  
Protuberance, [1616](#)  
Provisional Diagnosis, [1616](#)  
Provisional Tic Disorder, [1616](#)  
Provocation, [1616](#)  
Proximal, [1616](#)  
Proximal Deep Vein Thrombosis, [1617](#)  
Proximal DVT, [1617](#)  
Proximal Humerus Fractures, [1617](#)  
Proximal Radioulnar Joint, [1617](#)  
Proximal Renal Tubular Acidosis, [1617](#)  
Proximal Tubule, [1617](#)  
Proximal Tubule Reabsorption, [1617](#)  
Prune Belly Syndrome, [1617](#)  
Prurigo, [1617](#)  
Prurigo Nodularis, [1617](#)  
Pruritic, [1617](#)  
Pruritic Eruption, [1617](#)  
Pruritis, [1617](#)  
Pruritis Ani, [1617](#)  
Pruritus, [1618](#)  
Pruritus and Systemic Disease, [1618](#)  
Pruritus Ani, [1618](#)  
Pruritus In CKD, [1618](#)

---

---

Prussian Blue, 1618  
 PSA, 1618  
 PSA Test, 1618  
 PSA Velocity, 1618  
 Psammoma, 1618  
 Pseudarthrosis, 1618  
 Pseudo-Cushing Syndrome, 1619  
 Pseudo-Kaposi Sarcoma (Acroangiodermatitis), 1620  
 Pseudoaneurysm, 1618  
 Pseudoangiomatous Stromal Hyperplasia, 1618  
 Pseudobulbar Affect, 1618  
 Pseudocholinesterase Deficiency, 1619  
 Pseudoclubbing, 1619  
 Pseudocyesis, 1619  
 Pseudodementia, 1619  
 Pseudoephedrine, 1619  
 Pseudofolliculitis of the Beard, 1619  
 Pseudogout, 1619  
 Pseudogout Flare, 1619  
 Pseudogynecomastia, 1619  
 Pseudohypertrophic Muscular Dystrophy, 1619  
 Pseudohypoglycemia, 1619  
 Pseudohypoparathyroidism, 1620  
 Pseudomelanosis Duodeni, 1620  
 Pseudomembranous Colitis, 1620  
 Pseudomonas, 1620  
 Pseudomonas Aeruginosa, 1620  
 Pseudomonas Aeruginosa Infection, 1620  
 Pseudomonas Aeruginosa Resistance, 1620  
 Pseudomonas Folliculitis, 1620  
 Pseudomyxoma Peritonei, 1620  
 Pseudophakia, 1620  
 Pseudoporphyria, 1620  
 Pseudopseudohypoparathyroidism, 1620  
 Pseudorabies, 1621  
 Pseudothrombocytosis Due To Red Cell Fragments, 1621  
 Pseudotumor Cerebri, 1621  
 Pseudoxanthoma Elasticum, 1621  
 Psittacosis, 1621  
 Psittacosis (Parrot Fever), 1621  
 PSMA, 1621  
 Psoas, 1621  
 Psoas Abscess, 1621  
 Psoas Sign, 1621  
 Psoriasis, 1621  
 Psoriatic Arthritis, 1622  
 Psoriatic Plaque, 1622  
 PSP, 1622  
 Psychiatric, 1622  
 Psychiatric Emergency, 1622  
 Psychiatrist, 1622  
 Psychiatry, 1622  
 Psycho-Oncology, 1624  
 Psychoacoustics, 1622  
 Psychoactive, 1622  
 Psychoactive Substance, 1622  
 Psychoanalysis, 1622  
 Psychodermatology, 1622  
 Psychodidae, 1622  
 Psychodrama, 1622  
 Psychodynamic Therapy, 1622  
 Psychogenic, 1623  
 Psychogenic Erectile Dysfunction, 1623  
 Psychogenic Non-Epileptic Seizures, 1623  
 Psycholinguistics, 1623  
 Psychological, 1623  
 Psychological Adjustment, 1623  
 Psychological Dependence, 1623  
 Psychological Health, 1623  
 Psychological Homeostasis, 1623  
 Psychological Support, 1623  
 Psychological Transfer, 1623  
 Psychological Trauma, 1623  
 Psychology, 1623  
 Psychometrics, 1624  
 Psychomotor, 1624  
 Psychomotor Agitation, 1624  
 Psychomotor Disorder, 1624  
 Psychomotor Performance, 1624  
 Psychomotor Retardation, 1624  
 Psychoneuroimmunology, 1624  
 Psychopathic, 1624  
 Psychopathology, 1624  
 Psychopharmacology, 1624  
 Psychophysics, 1624  
 Psychophysiology, 1624  
 Psychosexual, 1624  
 Psychosexual Development, 1624  
 Psychosis, 1625  
 Psychosocial Rehabilitation, 1625  
 Psychosocial Support, 1625  
 Psychosomatic, 1625  
 Psychosomatic Disorder, 1625  
 Psychosomatic Medicine, 1625  
 Psychosurgery, 1625  
 Psychotherapy, 1625  
 Psychotic, 1625  
 Psychotic Disorder, 1625  
 Psychotic Disorders, 1625  
 Psychotropic Drug, 2129  
 Psychotropic Medication, 2129  
 PTCA, 1625  
 PTCL, 1625  
 PTE, 1625  
 Pteridine, 1626  
 Pterygium, 1626  
 PTH-Related Peptide, 1626  
 Pthirus Pubis, 1626

---

---

Ptilosis, 1626  
 PTLN, 1626  
 Ptosis, 1626  
 PTSD, 1626  
 Ptyalin, 1626  
 Ptyalism, 1626  
 Pubertal Failure, 1626  
 Puberty, 1626  
 Puberty In Boys, 1626  
 Puberty In Girls, 1626  
 Pubes, 1626  
 Pubic, 1627  
 Pubic Lice, 1627  
 Pubic Symphysis, 1627  
 Pubis, 1627  
 Public Assistance, 1627  
 Public Exposure, 1627  
 Public Health, 1627  
 Public Health Emergency, 1627  
 Puborectalis Muscle, 1627  
 Pudendal, 1627  
 Pudendal Arteries, 1627  
 Pudendal Block, 1627  
 Pudendal Nerve, 1627  
 Pudendal Nerve Block, 1627  
 Pudendal Neuralgia, 1646  
 Puerperal, 1627  
 Puerperal Disorder, 1627  
 Puerperal Fever, 1628  
 Puerperal Infection, 1628  
 Puerperal Psychosis, 1628  
 Puerperal Sepsis, 1628  
 Puerperium, 1628  
 Puffy Eyes, 1628  
 Pugilistic Stance, 1628  
 Pull Test, 1628  
 Pulling A Patient Up In Bed, 1628  
 Pulmonary, 1628  
 Pulmonary Acinus, 1628  
 Pulmonary Actinomyces, 1628  
 Pulmonary Alveolar Hemorrhage, 1628  
 Pulmonary Alveolar Microlithiasis, 1628  
 Pulmonary Alveolar Proteinosis, 1629  
 Pulmonary Angiography, 1629  
 Pulmonary Arterial Hypertension, 1629  
 Pulmonary Arteries, 1629  
 Pulmonary Arteriovenous Fistula, 1629  
 Pulmonary Arteriovenous Malformation, 1629  
 Pulmonary Artery, 1629  
 Pulmonary Artery Catheter, 1629  
 Pulmonary Artery Pulsatility Index, 1629  
 Pulmonary Artery Wedge Pressure, 1629  
 Pulmonary Aspergilloma, 1630  
 Pulmonary Atresia, 1630  
 Pulmonary Blastoma, 1630  
 Pulmonary Capillary Wedge Pressure, 1630  
 Pulmonary Circuit, 1630  
 Pulmonary Circulation, 1630  
 Pulmonary Compliance, 1630  
 Pulmonary Contusion, 1630  
 Pulmonary Disease and Pregnancy, 1630  
 Pulmonary Edema, 1630  
 Pulmonary Embolism, 1630  
 Pulmonary Embolism Treatment, 1630  
 Pulmonary Embolus, 1630  
 Pulmonary Emphysema, 1631  
 Pulmonary Eosinophilia, 1631  
 Pulmonary Fibrosis, 1631  
 Pulmonary Fistula, 1631  
 Pulmonary Function Tests, 1631  
 Pulmonary Hemorrhage, 1631  
 Pulmonary Hemosiderosis, 1631  
 Pulmonary Hypertension, 1631  
 Pulmonary Hypertension Classification, 1631  
 Pulmonary Hypertension Due to Another Condition, 1646  
 Pulmonary Hypertension, Primary, 988  
 Pulmonary Hypertension, Secondary, 1647  
 Pulmonary Hypoplasia, 1631  
 Pulmonary Infarction, 1631  
 Pulmonary Infection, 1631  
 Pulmonary Langerhans Cell Histiocytosis, 1632  
 Pulmonary Lobes, 1632  
 Pulmonary Lymphangioleiomyomatosis, 1647  
 Pulmonary Lymphangiomyomatosis, 1647  
 Pulmonary Mass, 1632  
 Pulmonary Nocardiosis, 1632  
 Pulmonary Nodule, 1632  
 Pulmonary Oedema, 1632  
 Pulmonary Reactance, 1632  
 Pulmonary Regurgitation (Pulmonic Regurgitation), 1632  
 Pulmonary Rehabilitation, 1632  
 Pulmonary Sequestration, 1632  
 Pulmonary Stenosis, 1632  
 Pulmonary Surfactant, 1632  
 Pulmonary Thromboendarterectomy, 1633  
 Pulmonary Trunk, 1633  
 Pulmonary Tuberculosis, 1633  
 Pulmonary Valve, 1633  
 Pulmonary Valve Insufficiency, 1633  
 Pulmonary Valve Regurgitation, 1633  
 Pulmonary Valve Stenosis, 1633  
 Pulmonary Vascular Resistance, 1633  
 Pulmonary Vasodilator Therapy, 1633  
 Pulmonary Vein, 1633  
 Pulmonary Vein Isolation, 1633  
 Pulmonary Veins, 1633

---



---

Pulmonary Veno-Occlusive Disease, 1633  
 Pulmonary Ventilation, 1634  
 Pulmonary Ventilation/Perfusion Scan, 1634  
 Pulmonary Wedge Pressure, 1634  
 Pulmonary-Artery Dissection, 1629  
 Pulmonary-Renal Syndrome, 1632  
 Pulmonic Valve Stenosis, 1634  
 Pulmonologist, 1634  
 Pulmonology, 1634  
 Pulp, 1634  
 Pulp Canal, 1634  
 Pulp Disease, 1634  
 Pulp Stone, 1635  
 Pulpectomy, 1634  
 Pulpitis, 1634  
 Pulposus, 1634  
 Pulpotomy, 1634  
 Pulsatile Proptosis, 1635  
 Pulsatile Tinnitus, 1635  
 Pulse, 1635  
 Pulse Examination, 1635  
 Pulse Irregular, 1635  
 Pulse Oximeter, 1635  
 Pulse Oximetry, 1635  
 Pulse Pressure, 1635  
 Pulse Rate, 1635  
 Pulseless Disease, 1635  
 Pulseless Electrical Activity, 1635  
 Pulsus, 1635  
 Pulsus Alternans, 1635  
 Pulsus Bisferiens, 1636  
 Pulsus Paradoxus, 1636  
 Pulsus Parvus et Tardus, 1636  
 Pulvinar, 1636  
 Pumice, 1636  
 Pump Failure, 1636  
 Pump Lung, 1647  
 Pump-Oxygenator, 1636  
 Punch Biopsy, 1636  
 Punctal Occlusion, 1636  
 Punctate, 1636  
 Punctate Palmoplantar Keratoderma, 1636  
 Puncture, 1636  
 Puncture Biopsy, 1636  
 Puncture Wound, 1637  
 Pungent, 1637  
 Punicic Acid, 1637  
 Punishment, 1637  
 Pupil, 1637  
 Pupillary, 1637  
 Pupillary Light Reflex, 1637  
 Pure Cutaneous Histiocytosis, 1647  
 Pure Red Cell Aplasia, 1637  
 Pure Tone Audiometry, 1637  
 Pure-Tone Audiometry, 1637  
 Purgative, 1637  
 Purge, 1637  
 Purging, 1637  
 Purging Disorder, 1637  
 Purine, 1638  
 Purine Nucleosides, 1638  
 Purine Nucleotides, 1638  
 Purines, 1638  
 Purinoceptor, 1638  
 Purkinje Cell, 1638  
 Purkinje Fibers, 1638  
 Purkinje Fibres, 1638  
 Purpura, 1638  
 Purpura Fulminans, 1638  
 Purpura Hemorrhagica ITP, 1647  
 Purpura, Henoch-Schonlein, 989  
 Purpura, Idiopathic Thrombocytopenic, 995  
 Purpura, Thrombotic Thrombocytopenic, 1918  
 Purulent, 1638  
 Purulent Pericarditis, 1638  
 Pus, 1638  
 Push Enteroscopy, 1639  
 Pustular Psoriasis, 1639  
 Pustule, 1639  
 Pustules, 1639  
 Pustulosis, 1639  
 Putamen, 1639  
 Putrefaction, 1639  
 Putrescine, 1639  
 PUVA, 1639  
 PVC, 1639  
 PVD, 1639  
 PVD (Peripheral Vascular Disease), 1647  
 PVI, 1639  
 PVPS, 1639  
 Pyarthrosis, 1639  
 Pycnodysostosis, 1639  
 Pyelitis, 1640  
 Pyelo, 1640  
 Pyelogram, 1640  
 Pyelography, 1640  
 Pyelolithotomy, 1640  
 Pyelonephritis, 1640  
 Pyelotomy, 1640  
 Pylephlebitis, 1640  
 Pyloric Antrum, 1640  
 Pyloric Canal, 1640  
 Pyloric Sphincter, 1640  
 Pyloric Stenosis, 1640  
 Pyloric Stenosis In Infants, 1640  
 Pyloroplasty, 1640  
 Pylorus, 1641  
 Pyoderma, 1641

---

---

Pyoderma Gangrenosum, [1641](#)  
 Pyogenesis, [1641](#)  
 Pyogenic, [1641](#)  
 Pyogenic Granuloma, [1641](#)  
 Pyogenic Granuloma (Lobular Capillary Hemangioma), [1641](#)  
 Pyogenic Liver Abscess, [1641](#)  
 Pyometra, [1641](#)  
 Pyonephrosis, [1641](#)  
 Pyosalpinx, [1641](#)  
 Pyostomatitis Vegetans, [1641](#)  
 Pyramid, [1641](#)  
 Pyramidal Cell, [1642](#)  
 Pyramidal Tract, [1642](#)  
 Pyramidal Tracts, [1642](#)  
 Pyrazinamide, [1642](#)  
 Pyrazines, [1642](#)  
 Pyrazole, [1642](#)  
 Pyrazoles, [1642](#)  
 Pyrenochaeta, [1642](#)  
 Pyrethrins, [1642](#)  
 Pyrethrum, [1642](#)  
 Pyrexia, [1644](#)  
 Pyrexia Of Unknown Origin In A Hemodialysis Patient, [1644](#)  
 Pyridine, [1642](#)  
 Pyridostigmine, [1642](#)  
 Pyridostigmine Bromide, [1642](#)  
 Pyridoxal Phosphate, [1642](#)  
 Pyridoxine, [1642](#)  
 Pyridoxine Deficiency, [1643](#)  
 Primethamine, [1643](#)  
 Pyrimidine, [1643](#)  
 Pyrimidine Analog, [1643](#)  
 Pyrimidine Dimer, [1643](#)  
 Pyrimidine Nucleosides, [1643](#)  
 Pyrimidine Nucleotides, [1643](#)  
 Pyrimidines, [1643](#)  
 Pyrogen, [1643](#)  
 Pyroglyphidae, [1643](#)  
 Pyromania, [1643](#)  
 Pyrophosphatase, [1643](#)  
 Pyroptosis, [1644](#)  
 Pyrosequencing, [1644](#)  
 Pyrosis, [1644](#)  
 Pyrroles, [1644](#)  
 Pyrrolidonecarboxylic Acid, [1644](#)  
 Pyrrolnitrin, [1644](#)  
 Pyruvate, [1644](#)  
 Pyruvate Carboxylase Deficiency, [1644](#)  
 Pyruvate Decarboxylase, [1644](#)  
 Pyruvate Dehydrogenase, [1644](#)  
 Pyruvate Dehydrogenase Complex Deficiency (PDCD), [1644](#)  
 Pyruvate Kinase, [1644](#)  
 Pyruvate Kinase Blood Test, [1644](#)  
 Pyruvate Kinase Deficiency, [1645](#)  
 Pyuria, [1645](#)  
 PYY, [1645](#)  
 Q Angle, [1649](#)  
 Q Arm Of A Chromosome, [1649](#)  
 Q Fever, [1649](#)  
 Q Wave, [1656](#)  
 Q Waves, [1656](#)  
 Q-Angle Measurement, [1649](#)  
 Q-Banding, [1649](#)  
 Q-Switch Laser, [1649](#)  
 QRS Axis, [1649](#)  
 QRS Complex, [1649](#)  
 QRS Duration, [1649](#)  
 QT Interval, [1650](#)  
 QT Prolongation, [1650](#)  
 QTc Dispersion, [1650](#)  
 Quad Screen, [1652](#)  
 Quadrant, [1650](#)  
 Quadrantanopia, [1650](#)  
 Quadrantectomy, [1650](#)  
 Quadrangle Lobe, [1650](#)  
 Quadratus Femoris, [1650](#)  
 Quadratus Lumborum, [1650](#)  
 Quadratus Plantae, [1650](#)  
 Quadriceps, [1650](#)  
 Quadriceps Active Test, [1650](#)  
 Quadriceps Contusion, [1651](#)  
 Quadriceps Femoris, [1651](#)  
 Quadriceps Femoris Muscle, [1651](#)  
 Quadriceps Injury, [1651](#)  
 Quadriceps Tendinitis, [1651](#)  
 Quadriceps Tendon Rupture, [1651](#)  
 Quadriceps Tendon Ruptures, [1651](#)  
 Quadrigeminal Cistern, [1651](#)  
 Quadrigeminal Plate, [1651](#)  
 Quadrilateral Space Syndrome, [1651](#)  
 Quadriplegia, [1652](#)  
 Quadruple Screen Test, [1652](#)  
 Quadruplet Pregnancy, [1652](#)  
 Qualitative HCG Blood Test, [1652](#)  
 Quality Of Life, [1652](#)  
 Quality-Adjusted Life Year, [1652](#)  
 Quantitative Bence-Jones Protein Test, [1652](#)  
 Quantitative Computed Tomography, [1652](#)  
 Quantitative HCG Blood Test, [1652](#)  
 Quantitative Immunoglobulins, [1652](#)  
 Quantitative Nephelometry Test, [1652](#)

---

---

Quantitative PET, [1653](#)  
 Quantitative Sensory Testing, [1653](#)  
 Quarantine, [1653](#)  
 Quartan Fever, [1653](#)  
 Queckenstedt Test, [1653](#)  
 Queensland Tick Typhus, [1653](#)  
 Quellung Reaction, [1653](#)  
 Quetiapine Fumarate, [1653](#)  
 Quick Relievers, [1653](#)  
 Quick Sequential Organ Failure Assessment, [1653](#)  
 Quick Test, [1654](#)  
 Quickening, [1653](#)  
 Quinalphos, [1654](#)  
 Quinapril, [1654](#)  
 Quincke Edema, [1654](#)  
 Quincke Needle, [1654](#)  
 Quincke Pulse, [1654](#)  
 Quincke Pulse Sign, [1654](#)  
 Quincke's Disease, [1654](#)  
 Quincke's Edema, [1654](#)  
 Quincke's Pulse, [1654](#)  
 Quincke's Sign, [1654](#)  
 Quinidine, [1655](#)  
 Quinidine Syncope, [1655](#)  
 Quinine, [1655](#)  
 Quinine Amblyopia, [1655](#)  
 Quinolone, [1655](#)  
 Quinolone Antibiotics, [1655](#)  
 Quinolone-Induced Tendinopathy, [1655](#)  
 Quinsy, [1655](#)  
 Quinsy Peritonsillar Abscess, [1655](#)  
 Quinupristin-Dalfopristin, [1655](#)  
 Quit Smoking, [1656](#)  
 Quorum Sensing, [1656](#)  
 Quotidian Fever, [1656](#)  
 Quotidian Malaria, [1656](#)  
  
 R-Factor, [1710](#)  
 Rabeprazole, [1657](#)  
 Rabies, [1657](#)  
 Rabies Vaccination, [1657](#)  
 Rabies Vaccine, [2129](#)  
 Rabies Virus, [1657](#)  
 Rachitic Rosary, [1657](#)  
 Radial, [1657](#)  
 Radial Artery, [1657](#)  
 Radial Artery Thrombosis, [1657](#)  
 Radial Club Hand, [1657](#)  
 Radial Head Subluxation, [1658](#)  
 Radial Nerve, [1658](#)  
 Radial Nerve Dysfunction, [1658](#)  
 Radial Nerve Palsy, [1658](#)  
 Radial Tunnel Syndrome, [1658](#)  
 Radial Vein, [1658](#)  
  
 Radiation, [1658](#)  
 Radiation Biology, [1658](#)  
 Radiation Contamination, [1658](#)  
 Radiation Dermatitis Skin Care, [1658](#)  
 Radiation Dosage, [1658](#)  
 Radiation Dose, [1658](#)  
 Radiation Emergency, [1658](#)  
 Radiation Enteritis, [1659](#)  
 Radiation Exposure, [1659](#)  
 Radiation Fibrosis, [1659](#)  
 Radiation Field, [1659](#)  
 Radiation Leukemia, [1659](#)  
 Radiation Monitoring, [1659](#)  
 Radiation Necrosis, [1659](#)  
 Radiation Oncologist, [1659](#)  
 Radiation Oncology, [1659](#)  
 Radiation Physics, [1659](#)  
 Radiation Pneumonitis, [1659](#)  
 Radiation Poisoning, [1659](#)  
 Radiation Pregnancy Screening, [1659](#)  
 Radiation Proctitis, [1660](#)  
 Radiation Protection, [1660](#)  
 Radiation Recall, [1660](#)  
 Radiation Safety, [1660](#)  
 Radiation Sickness, [1660](#)  
 Radiation Therapy, [1660](#)  
 Radiation Tolerance, [1660](#)  
 Radiation-Induced Angiosarcoma, [1659](#)  
 Radiation-Recall Dermatitis, [1660](#)  
 Radical Cystectomy, [1660](#)  
 Radical Cystoprostatectomy, [1660](#)  
 Radical Hysterectomy, [1660](#)  
 Radical Lymphadenectomy, [1660](#)  
 Radical Mastectomy, [1660](#)  
 Radical Nephrectomy, [1660](#)  
 Radical Prostatectomy, [1661](#)  
 Radicular Cyst, [1661](#)  
 Radiculitis, [1661](#)  
 Radiculopathy, [1661](#)  
 Radio-Femoral Delay, [1662](#)  
 Radio-Opaque, [1664](#)  
 Radio-Radial Delay, [1665](#)  
 Radioactive Iodine, [1661](#)  
 Radioactive Iodine Ablation, [1661](#)  
 Radioactive Iodine Uptake, [1661](#)  
 Radioactive Iodine Uptake Test, [1661](#)  
 Radioactivity, [1661](#)  
 Radioallergosorbent Test, [1661](#)  
 Radiobiology, [1661](#)  
 Radiocarpal Joint, [1661](#)  
 Radiochemical Purity, [1661](#)  
 Radiochemistry, [1662](#)  
 Radiocontrast, [1662](#)  
 Radiodermatitis, [1662](#)

---

---

Radioembolisation, [1662](#)  
Radioembolization, [1662](#)  
Radiofrequency, [1662](#)  
Radiofrequency Ablation, [1662](#)  
Radiofrequency Ablation For Chronic Pain, [1662](#)  
Radiograph, [1662](#)  
Radiographer, [1662](#)  
Radiography, [1662](#)  
Radioimmunoassay, [1662](#)  
Radioimmunoguided Surgery, [1662](#)  
Radioimmunotherapy, [1663](#)  
Radioiodine, [1663](#)  
Radioiodine Therapy, [1663](#)  
Radioisotope, [1663](#)  
Radioligand Therapy, [1663](#)  
Radiologic Health, [1663](#)  
Radiologic Response, [1663](#)  
Radiologic Technologist, [1663](#)  
Radiologic-Pathologic Correlation, [1663](#)  
Radiologist, [1663](#)  
Radiology, [1663](#)  
Radiology Report, [1663](#)  
Radiology-Patient Identification, [1663](#)  
Radiolucent, [1664](#)  
Radiomimetic, [1664](#)  
Radionuclide, [1664](#)  
Radionuclide Angiography, [1664](#)  
Radionuclide Cystogram, [1664](#)  
Radionuclide Generator, [1664](#)  
Radionuclide Heart Scan, [1664](#)  
Radionuclide Imaging, [1664](#)  
Radionuclide Imaging Or Scan, [1664](#)  
Radionuclide Therapy, [1664](#)  
Radionuclides, [1664](#)  
Radionuclidic Purity, [1664](#)  
Radiopaque, [1664](#)  
Radiopharmaceutical, [1665](#)  
Radiopharmaceutical Extravasation, [1665](#)  
Radiopharmaceutical Labeling Kit, [1665](#)  
Radiopharmaceutical Quality Control, [1665](#)  
Radiopharmacy, [1665](#)  
Radioprotective Agent, [1665](#)  
Radiosensitization, [1665](#)  
Radiosurgery, [1665](#)  
Radiotherapy, [1665](#)  
Radiotherapy Bolus, [1665](#)  
Radiotherapy Dosage, [1665](#)  
Radiotracer, [1665](#)  
Radioulnar Joint, [1665](#)  
Radium, [1666](#)  
Radium-223, [1666](#)  
Radius, [1666](#)  
Radius Fracture, [1666](#)  
Radon, [1666](#)  
Ragged Red Fibers, [1666](#)  
Raised Hemidiaphragm, [1666](#)  
Rale, [1666](#)  
Rales, [1666](#)  
Raloxifene, [1666](#)  
Raloxifene Hydrochloride, [1666](#)  
Raltegravir Potassium, [1666](#)  
Raltitrexed, [1666](#)  
Ramipril, [1666](#)  
Rampant Caries, [1667](#)  
Ramsay Hunt Syndrome, [1667](#)  
Ramsay Hunt Syndrome Type I, [1667](#)  
Ramsay Hunt Syndrome Type II, [1667](#)  
Ramucirumab, [1667](#)  
Ramus, [1667](#)  
Random Allocation, [1667](#)  
Randomised Controlled Trial, [1667](#)  
Randomization Validation, [1667](#)  
Randomized Controlled Trial, [1667](#)  
Range Of Motion, [1667](#)  
Ranibizumab, [1667](#)  
Ranke Complex, [1667](#)  
Ranolazine, [1668](#)  
Ranson Criteria, [1668](#)  
RANTES, [1668](#)  
Ranula, [1668](#)  
Raoultella, [1668](#)  
Rape, [1668](#)  
Raphe, [1668](#)  
Raphe Nuclei, [1668](#)  
Rapid Eye Movement Sleep, [1668](#)  
Rapid Eye Movement Sleep Behavior Disorder, [1668](#)  
Rapid Gastric Emptying, [1668](#)  
Rapid Heartbeat, [1668](#)  
Rapid Plasma Reagin Test, [1668](#)  
Rapid Sequence Induction, [1668](#)  
Rapid Sequence Intubation, [1669](#)  
Rapid Shallow Breathing, [1669](#)  
Rapid Strep Test, [1669](#)  
Rapid-Sequence Intubation, [1669](#)  
Rapidly Progressive Glomerulonephritis, [1668](#)  
Rapp-Hodgkin Syndrome, [1669](#)  
Rare Disease, [1669](#)  
Rarefaction, [1669](#)  
Ras Peptide, [1670](#)  
Rasburicase, [1669](#)  
Rash, [1669](#)  
Rasmussen Encephalitis, [1669](#)  
Rasp, [1669](#)  
RAST, [1670](#)  
Rastelli Procedure, [1670](#)  
Rat Bite Fever, [1670](#)  
Rat Lungworm Disease, [1670](#)  
Rat-Bite Fever, [1670](#)

---

---

Rathke Cleft Cyst, [1670](#)  
Rathke Pouch, [1670](#)  
Ratio, [1670](#)  
Rationalization, [1670](#)  
RATS, [1670](#)  
Ravuconazole, [1670](#)  
Raw Egg, [1670](#)  
Raynaud Disease, [1733](#)  
Raynaud Phenomenon, [1670](#)  
Raynaud Syndrome, [1671](#)  
Raynaud's Disease, [1671](#)  
Raynaud's Phenomenon, [1671](#)  
Raynaud's Syndrome, [1671](#)  
Razor Bumps, [1671](#)  
RBC Count, [1671](#)  
RBC Indices, [1671](#)  
RBC Nuclear Scan, [1671](#)  
RBC Urine Test, [1671](#)  
RDA, [1671](#)  
Reabsorption, [1671](#)  
Reaction, [1671](#)  
Reactivation Tuberculosis, [1671](#)  
Reactive Arthritis, [1672](#)  
Reactive Attachment Disorder, [1672](#)  
Reactive Gastropathy Pathology, [1672](#)  
Reactive Hypoglycemia, [1672](#)  
Reactive Oxygen Species, [1672](#)  
Reactive Perforating Collagenosis, [1672](#)  
Reality Testing, [1672](#)  
Rebound Effect, [1672](#)  
Rebound Headache, [1672](#)  
Rebound Insomnia, [1672](#)  
Rebound Tenderness, [1672](#)  
Recalcitrant, [1672](#)  
Receiver Operating Characteristic, [1672](#)  
Recent Memory, [1672](#)  
Receptor, [1673](#)  
Receptor Binding, [1673](#)  
Receptor Desensitization, [1673](#)  
Receptor Downregulation, [1673](#)  
Receptor Tyrosine Kinases, [1673](#)  
Receptor Upregulation, [1673](#)  
Receptors, [1673](#)  
Recession Of Medial Rectus Muscle, [1673](#)  
Recessive, [1673](#)  
RECIST, [1673](#)  
Recognizing Medical Emergencies, [1673](#)  
Recognizing Teen Depression, [1673](#)  
Recombinant, [1673](#)  
Recombinant DNA, [1673](#)  
Recombinant Protein, [1674](#)  
Recombinant Tissue Plasminogen Activator, [1674](#)  
Recombinase, [1674](#)  
Recombinational Repair, [1674](#)  
Recommended Daily Allowance, [1674](#)  
Recommended Dietary Allowance, [1674](#)  
Record, [1674](#)  
Recovery Position, [1674](#)  
Recreation Therapy, [1674](#)  
Recrudescence, [1674](#)  
Rectal, [1674](#)  
Rectal Biopsy, [1674](#)  
Rectal Bleeding, [1674](#)  
Rectal Cancer, [1674](#)  
Rectal Carcinoma, [1675](#)  
Rectal Culture, [1675](#)  
Rectal Disease, [1675](#)  
Rectal Fistula, [1675](#)  
Rectal Hemorrhage, [1675](#)  
Rectal Inflammation, [1675](#)  
Rectal Itching, [1675](#)  
Rectal Mucositis, [1675](#)  
Rectal Necrosis, [1675](#)  
Rectal Neoplasm, [1675](#)  
Rectal Obstruction, [1675](#)  
Rectal Perforation, [1675](#)  
Rectal Polyposis, [1675](#)  
Rectal Polyps, [1675](#)  
Rectal Prolapse, [1675](#)  
Rectal Prolapse Repair, [1675](#)  
Rectal Route, [1675](#)  
Rectal Stenosis, [1675](#)  
Rectal Tenesmus, [1676](#)  
Rectal Ulcer, [1676](#)  
Rectocele, [1676](#)  
Rectosigmoid Junction, [1676](#)  
Rectosigmoid Neoplasm, [1676](#)  
Rectouterine Pouch, [1676](#)  
Rectovaginal Fistula, [1676](#)  
Rectum, [1676](#)  
Rectus Abdominis, [1676](#)  
Rectus Abdominis Muscle, [1676](#)  
Rectus Abdominis Sheath, [1676](#)  
Rectus Sheath Hematoma, [1676](#)  
Recur, [1676](#)  
Recurrence, [1676](#)  
Recurrence Risk, [1677](#)  
Recurrence-Free Survival, [1676](#)  
Recurrent, [1677](#)  
Recurrent Breast Cancer, [1677](#)  
Recurrent Caries, [1677](#)  
Recurrent Disease, [1677](#)  
Recurrent Early Pregnancy Loss, [1677](#)  
Recurrent Fractures, [1677](#)  
Recurrent Laryngeal Nerve, [1677](#)  
Recurrent Laryngeal Nerve Palsy, [1677](#)  
Recurrent or Metastatic Breast Cancer, [1677](#)  
Recurrent Oral Ulcers, [1677](#)

---

---

Recurrent Pregnancy Loss, [1677](#)  
Recurrent Prostate Cancer, [1677](#)  
Recurrent Respiratory Papillomatosis, [1677](#)  
Recurrent Urinary Tract Infection, [1678](#)  
Red Birthmarks, [1678](#)  
Red Blood Cell, [1678](#)  
Red Blood Cell Casts, [1678](#)  
Red Blood Cell Labeling, [1678](#)  
Red Bone Marrow, [1678](#)  
Red Cell Distribution Width, [1678](#)  
Red Cell Indices, [1678](#)  
Red Eye, [1678](#)  
Red Flag, [1678](#)  
Red Hepatization, [1678](#)  
Red Infarction, [1678](#)  
Red Man Syndrome, [1679](#)  
Red Nucleus, [1679](#)  
Red-Man Syndrome, [1679](#)  
Reductase Deficiency, [1679](#)  
Reduction, [1679](#)  
Reduction Division, [1679](#)  
Reduction Mammoplasty, [1679](#)  
Redundant Mitral Valve, [1679](#)  
Reduviidae, [1679](#)  
Reed-Sternberg Cells, [1679](#)  
Reexpansion Pulmonary Edema, [1679](#)  
Refeeding Syndrome, [1679](#)  
Reference Interval, [1679](#)  
Referral, [1680](#)  
Referred Otalgia, [1680](#)  
Referred Pain, [1680](#)  
Reflex, [1680](#)  
Reflex Action, [1680](#)  
Reflex Anoxic Seizure, [1680](#)  
Reflex Arc, [1680](#)  
Reflex Epilepsy, [1680](#)  
Reflexes, [1680](#)  
Reflux, [1680](#)  
Reflux Esophagitis Pathology, [1680](#)  
Reflux Laryngitis, [1680](#)  
Reflux Nephropathy, [1680](#)  
Refraction, [1681](#)  
Refractive Disorder, [1681](#)  
Refractive Error, [1681](#)  
Refractive Errors And Astigmatism, [1681](#)  
Refractive Surgery, [1681](#)  
Refractometer, [1681](#)  
Refractometry, [1681](#)  
Refractory, [1681](#)  
Refractory Anemia, [1681](#)  
Refractory Cancer, [1681](#)  
Refractory Disease, [1681](#)  
Refractory Period, [1681](#)  
Refrigerant Poisoning, [1681](#)  
Refsum Disease, [1681](#)  
Regadenoson Pharmacological Stress, [1681](#)  
Regeneration, [1682](#)  
Regimen, [1682](#)  
Regional, [1682](#)  
Regional Anesthesia, [1682](#)  
Regional Chemotherapy, [1682](#)  
Regional Enteritis, [1682](#)  
Regional Ileitis, [1682](#)  
Regulation Of Health Professions, [1682](#)  
Regulatory Approval, [1682](#)  
Regulatory Authority, [1682](#)  
Regulatory Gene, [1682](#)  
Regulatory T Cells, [1682](#)  
Regurgitation, [1682](#)  
Rehab, [1682](#)  
Rehabilitation, [1682](#)  
Rehabilitation Clinic, [1683](#)  
Rehabilitation Engineering, [1683](#)  
Rehabilitation Medicine, [2129](#)  
Rehabilitation Motivation, [1683](#)  
Rehabilitation Potential, [1683](#)  
Rehabilitation Unit, [1683](#)  
Rehydration, [1683](#)  
Reid Index, [1683](#)  
Reifenstein Syndrome, [1683](#)  
Reiki, [1683](#)  
Reinfection, [1683](#)  
Reinke Edema, [1683](#)  
Reinnervation, [1683](#)  
Rejection, [1683](#)  
Relapse, [1684](#)  
Relapse-Free Survival, [1684](#)  
Relapsing Fever, [1684](#)  
Relapsing Polychondritis, [1684](#)  
Relapsing-Remitting Multiple Sclerosis, [1684](#)  
Relative Afferent Pupillary Defect, [1684](#)  
Relative Biological Effectiveness, [1684](#)  
Relative Bradycardia, [1684](#)  
Relative Risk, [1684](#)  
Relaxation, [1684](#)  
Relaxation Response, [1684](#)  
Relaxation Techniques For Stress, [1684](#)  
Relaxation Therapy, [1685](#)  
Relaxin, [1685](#)  
Religion, [1685](#)  
Relocation Test, [1685](#)  
REM, [1685](#)  
REM Behavior Disorder, [1685](#)  
REM Rebound, [1686](#)  
REM Sleep, [1686](#)  
REM Sleep Behavior Disorder, [1686](#)  
Remedy, [1685](#)  
Remifentanil, [1685](#)

---



---

Remission, [1685](#)  
Remittent Fever, [1685](#)  
Remodeling, [1685](#)  
Remote Surgery, [1685](#)  
Removal Technique, [1685](#)  
Renal, [1686](#)  
Renal Abscess, [1686](#)  
Renal Adenoma, [1686](#)  
Renal Agenesis, [1686](#)  
Renal and Ureteral Injury, [1686](#)  
Renal And Urological Diseases, [1686](#)  
Renal Anemia, [1686](#)  
Renal Arteriography, [1686](#)  
Renal Artery, [1686](#)  
Renal Artery Aneurysm, [1686](#)  
Renal Artery Disease, [1687](#)  
Renal Artery Doppler, [1687](#)  
Renal Artery Embolism, [1687](#)  
Renal Artery Stenosis, [1687](#)  
Renal Biopsy, [1687](#)  
Renal Calculi, [1687](#)  
Renal Calculus, [1687](#)  
Renal Cancer, [1687](#)  
Renal Carcinoma, [1687](#)  
Renal Cell Carcinoma, [1687](#)  
Renal Cell Carcinoma Pathology, [1687](#)  
Renal Changes In Pregnancy, [1687](#)  
Renal Circulation, [1687](#)  
Renal Clearance, [1687](#)  
Renal Colic, [1687](#)  
Renal Corpuscle, [1687](#)  
Renal Cortex, [1688](#)  
Renal Cryptococcosis, [1688](#)  
Renal Cyst, [1688](#)  
Renal Dialysis, [1688](#)  
Renal Diet, [1688](#)  
Renal Disease, [1688](#)  
Renal Diseases, [1688](#)  
Renal Dose Adjustment, [1688](#)  
Renal Drug Excretion, [2129](#)  
Renal Dysplasia, [1688](#)  
Renal Dysplasia And Cystic Disease, [1688](#)  
Renal Failure, [1688](#)  
Renal Function, [1688](#)  
Renal Glomerulus, [1688](#)  
Renal Hamartoma, [1688](#)  
Renal Hypertension, [1688](#)  
Renal Hypertrophy, [1689](#)  
Renal Hypoperfusion, [1689](#)  
Renal Infarction, [1689](#)  
Renal Insufficiency, [1689](#)  
Renal Leukemia Infiltration, [1689](#)  
Renal Lithiasis, [1689](#)  
Renal Manifestations of Sick Cell Disease, [1689](#)  
Renal Mass, [1689](#)  
Renal Medulla, [1689](#)  
Renal Necrosis, [1689](#)  
Renal Oncocytoma, [1689](#)  
Renal Osteodystrophy, [1689](#)  
Renal Papillary Necrosis, [1689](#)  
Renal Parenchyma, [1689](#)  
Renal Pelvis, [1690](#)  
Renal Pelvis Or Ureter Cancer, [1690](#)  
Renal Perfusion Scintiscan, [1690](#)  
Renal Pyramids, [1690](#)  
Renal Reabsorption, [1690](#)  
Renal Replacement Therapy, [1690](#)  
Renal Replacement Therapy Indications, [1690](#)  
Renal Salt-Wasting Syndrome, [1690](#)  
Renal Sarcoma, [1690](#)  
Renal Scan, [1690](#)  
Renal Scintigraphy, [1690](#)  
Renal Stone, [1690](#)  
Renal Transitional Cell Carcinoma, [1690](#)  
Renal Transplantation, [1690](#)  
Renal Trauma, [1690](#)  
Renal Tubular Acidosis, [1691](#)  
Renal Tubular Acidosis Type 1, [1691](#)  
Renal Tubular Acidosis Type 2, [1691](#)  
Renal Tubular Acidosis Type 4, [1691](#)  
Renal Tubular Disorders, [1691](#)  
Renal Tubule, [1691](#)  
Renal Tubule Acidosis, [1691](#)  
Renal Ultrasound, [1691](#)  
Renal Vascular Disease, [1691](#)  
Renal Vein, [1691](#)  
Renal Vein Thrombosis, [1691](#)  
Renal Venogram, [1691](#)  
Renaut Body, [1692](#)  
Renin, [1692](#)  
Renin Blood Test, [1692](#)  
Renin Inhibitor, [2129](#)  
Renin-Angiotensin System, [1692](#)  
Renin-Angiotensin-Aldosterone System, [1692](#)  
Renin-Angiotensin-Aldosterone System Physiology, [1692](#)  
Rennin, [1692](#)  
Renography, [1692](#)  
Renovascular Disease, [1692](#)  
Renovascular Hypertension, [1692](#)  
Renshaw Cell, [1692](#)  
Reoviridae Infection, [1692](#)  
Rep, [1692](#)  
Repair Of Webbed Fingers Or Toes, [1692](#)  
Repeat Surgery, [1693](#)  
Reperfusion, [1693](#)  
Reperfusion Injury, [1693](#)  
Reperfusion Therapy, [1693](#)  
Repetitive Motion Disorders, [1693](#)

---

---

Repetitive Nerve Stimulation, [1693](#)  
Repetitive Strain Injury, [1693](#)  
Replantation Of An Avulsed Tooth, [1693](#)  
Replantation Of Digits, [1693](#)  
Replication Initiation, [1693](#)  
Replicon, [1693](#)  
Repolarization, [1693](#)  
Reportable Diseases, [1693](#)  
Reporting Standards, [1693](#)  
Repressed Memory Therapy, [1694](#)  
Repression, [1694](#)  
Reproduction, [1694](#)  
Reproductive Cells, [1694](#)  
Reproductive History, [1694](#)  
Reproductive Hormones, [1694](#)  
Reproductive Organs, [1694](#)  
Reproductive System, [1694](#)  
Reproductive Technique, [1694](#)  
Reproductive Toxicity, [1694](#)  
Reproductive Years, [1694](#)  
Reprogram, [1694](#)  
Resectable, [1694](#)  
Resection, [1694](#)  
Resectoscope, [1695](#)  
Reserve Volume, [1695](#)  
Reservoir, [1695](#)  
Reservoir Of Infection, [1695](#)  
Resident, [1695](#)  
Residential Mobility, [1695](#)  
Residual, [1695](#)  
Residual Limb, [1695](#)  
Residual Limb Pain, [1695](#)  
Residual Neoplasm, [1695](#)  
Residual Risk, [1695](#)  
Residual Tumor, [1695](#)  
Residual Volume, [1695](#)  
Resistance, [1695](#)  
Resistance Antibiotic (Antibiotic Resistance), [2129](#)  
Resistance Training, [1695](#)  
Resistant Hypertension, [1695](#)  
Resistant Starch, [1696](#)  
Resistin, [1696](#)  
Resolution, [1696](#)  
Resonance, [1696](#)  
Resorcinol, [1696](#)  
Resorption, [1696](#)  
Resorption Atelectasis, [1696](#)  
Respiration, [1696](#)  
Respiratory, [1696](#)  
Respiratory Acidosis, [1696](#)  
Respiratory Alkalosis, [1696](#)  
Respiratory Arrest, [1696](#)  
Respiratory Burst, [1696](#)  
Respiratory Changes In Pregnancy, [1696](#)  
Respiratory Compliance, [1696](#)  
Respiratory Depression, [1697](#)  
Respiratory Disease, [1697](#)  
Respiratory Disorder, [1697](#)  
Respiratory Distress, [1697](#)  
Respiratory Distress Syndrome, [1697](#)  
Respiratory Drive, [1697](#)  
Respiratory Failure, [1697](#)  
Respiratory Failure Types 1 and 2, [1697](#)  
Respiratory Gating, [1697](#)  
Respiratory Health, [1697](#)  
Respiratory Illness, [1697](#)  
Respiratory Infection, [1697](#)  
Respiratory Infections, [1698](#)  
Respiratory Insufficiency, [1698](#)  
Respiratory Muscle, [1698](#)  
Respiratory Neoplasm, [1698](#)  
Respiratory Paralysis, [1698](#)  
Respiratory Physiology, [1698](#)  
Respiratory Protection, [1698](#)  
Respiratory Quotient, [1698](#)  
Respiratory Rate, [1698](#)  
Respiratory Rehabilitation, [1698](#)  
Respiratory Syncytial Virus, [1698](#)  
Respiratory Syncytial Virus (RSV), [1733](#)  
Respiratory Syncytial Virus Bronchiolitis, [1698](#)  
Respiratory Syncytial Virus Infection, [1698](#)  
Respiratory System, [1698](#)  
Respiratory Therapy, [1699](#)  
Respiratory Tract Infections, [1699](#)  
Respiratory Transport, [1699](#)  
Respiratory Viruses, [1699](#)  
Respiratory Viruses And Face Masks, [1699](#)  
Respirologist, [1699](#)  
Respirovirus, [1699](#)  
Respite Care, [1699](#)  
Response Assessment, [1699](#)  
Response Evaluation Criteria, [1699](#)  
Responsible Drinking, [1699](#)  
Rest, [1699](#)  
Restenosis, [1699](#)  
Resting Energy Expenditure, [1699](#)  
Resting Membrane Potential, [1699](#)  
Resting Phase, [1700](#)  
Resting Tremor, [1700](#)  
Restless Anal Syndrome, [1700](#)  
Restless Leg Syndrome, [1700](#)  
Restless Legs, [1700](#)  
Restless Legs Syndrome, [1700](#)  
Restless Legs Syndrome (RLS), [1733](#)  
Restlessness, [1700](#)  
Restorative Dentistry, [1700](#)  
Restraint, [1700](#)  
Restraint-Related Harm, [1700](#)

---

---

Restriction Enzyme, [1700](#)  
Restrictive Cardiomyopathy, [1700](#)  
Restrictive Lung Disease, [1700](#)  
Resuscitation, [1701](#)  
Resuscitative Thoracotomy, [1701](#)  
Retained Placenta, [1701](#)  
Retained Products Of Conception, [1701](#)  
Retainer, [1701](#)  
Retching, [1701](#)  
Rete Testis, [1701](#)  
Retention Of Urine, [1701](#)  
Retention Time, [1701](#)  
Reticular, [1701](#)  
Reticular Activating System, [1701](#)  
Reticular Dermis, [1701](#)  
Reticular Erythematous Mucinosi, [1701](#)  
Reticular Formation, [1701](#)  
Reticulin, [1702](#)  
Reticulo-Endothelial System, [1702](#)  
Reticulocyte, [1702](#)  
Reticulocyte Count, [1702](#)  
Reticulocyte Production Index, [1702](#)  
Reticulocytes, [1702](#)  
Reticulocytopenia, [1702](#)  
Reticulocytosis, [1702](#)  
Reticuloendothelial System, [1702](#)  
Reticulophagy, [1702](#)  
Reticulum, [1702](#)  
Retina, [1702](#)  
Retina Disorder, [1702](#)  
Retina Hemorrhage, [1703](#)  
Retinal Artery Occlusion, [1703](#)  
Retinal Cone, [1703](#)  
Retinal Damage, [1703](#)  
Retinal Degeneration, [1703](#)  
Retinal Detachment, [1703](#)  
Retinal Detachment Repair, [1703](#)  
Retinal Diseases, [1703](#)  
Retinal Disorders, [1703](#)  
Retinal Dysplasia, [1703](#)  
Retinal Dystrophy, [1703](#)  
Retinal Fold, [1703](#)  
Retinal Fundus, [1703](#)  
Retinal Ganglion, [1704](#)  
Retinal Hemorrhages, [1704](#)  
Retinal Implant, [1704](#)  
Retinal Ischemia, [1704](#)  
Retinal Laser Photocoagulation, [1704](#)  
Retinal Layers, [1704](#)  
Retinal Macula, [1704](#)  
Retinal Neoplasm, [1704](#)  
Retinal Perforation, [1704](#)  
Retinal Pigment, [1704](#)  
Retinal Pigment Epithelium, [1704](#)  
Retinal Prosthesis, [1704](#)  
Retinal Tear, [1704](#)  
Retinal Vasculitis, [1704](#)  
Retinal Vein Occlusion, [1705](#)  
Retinaldehyde, [1703](#)  
Retinitis, [1705](#)  
Retinitis Pigmentosa, [1705](#)  
Retinoblastoma, [1705](#)  
Retinoblastoma Protein, [1705](#)  
Retinoic Acid, [1705](#)  
Retinoid, [1705](#)  
Retinoids, [1705](#)  
Retinol, [1705](#)  
Retinopathy, [1705](#)  
Retinopathy Of Prematurity, [1705](#)  
Retinoschisis, [1706](#)  
Retinoscope, [1706](#)  
Retinoscopy, [1706](#)  
Retirement, [1706](#)  
Retracted Nipple, [1706](#)  
Retractile Testicle, [1706](#)  
Retractile Testis, [1706](#)  
Retractor, [1706](#)  
Retrobulbar Hemorrhage, [1706](#)  
Retrobulbar Neuritis, [1706](#)  
Retroesophageal, [1706](#)  
Retrognathia, [1706](#)  
Retrograde, [1706](#)  
Retrograde Amnesia, [1706](#)  
Retrograde Cystography, [1706](#)  
Retrograde Degeneration, [1706](#)  
Retrograde Ejaculation, [1706](#)  
Retrograde Intrarenal Surgery, [1707](#)  
Retrograde Urethrogram, [1707](#)  
Retroperitoneal Fibrosis, [1707](#)  
Retroperitoneal Hematoma, [1707](#)  
Retroperitoneal Hemorrhage, [1707](#)  
Retroperitoneal Inflammation, [1707](#)  
Retroperitoneal Neoplasm, [1707](#)  
Retroperitoneal Space, [1707](#)  
Retroperitoneum, [1707](#)  
Retropharyngeal Abscess, [1707](#)  
Retropharyngeal Space, [1707](#)  
Retrospective Studies, [1707](#)  
Retrospective Study, [1707](#)  
Retrosternal Thyroid Surgery, [1707](#)  
Retrotracheal, [1708](#)  
Retrotransposition, [1708](#)  
Retroversion, [1708](#)  
Retroversion Of The Uterus, [1708](#)  
Retroviridae, [1708](#)  
Retroviridae Infection, [1708](#)  
Retrovirus, [1708](#)  
Rett Syndrome, [1708](#)

---

---

Return Of Spontaneous Circulation, [1708](#)  
Reuptake, [1708](#)  
Revalidation, [1708](#)  
Revascularisation, [1708](#)  
Revascularization, [1708](#)  
Revascularization Surgery, [1708](#)  
Reverberation Artifact, [1709](#)  
Reversal Learning, [1709](#)  
Reverse Total Shoulder Replacement, [1709](#)  
Reverse Transcriptase, [1709](#)  
Reverse Transcriptase Inhibitor, [1709](#)  
Reverse Transcriptase Inhibitors, [1709](#)  
Reverse Transcription, [1709](#)  
Reverse Trendelenburg Position, [1709](#)  
Reversible Cell Injury, [1709](#)  
Revision Arthroplasty, [1709](#)  
Revision Total Hip Arthroplasty, [1709](#)  
Revision Total Knee Arthroplasty, [1709](#)  
Reye Syndrome, [1709](#)  
Reye's Syndrome, [1709](#)  
Reye-Johnson Syndrome, [1709](#)  
Reynolds Pentad, [1710](#)  
Rezafungin, [1710](#)  
Rgs Proteins, [1710](#)  
Rh Factor, [1712](#)  
Rh Incompatibility, [1712](#)  
Rh Isoimmunization, [1714](#)  
Rhabdoid Tumor, [1710](#)  
Rhabdomyolysis, [1710](#)  
Rhabdomyolysis And AKI, [1710](#)  
Rhabdomyoma, [1710](#)  
Rhabdomyosarcoma, [1710](#)  
Rhabdomyosarcoma Pathology, [1710](#)  
Rhegmatogenous Retinal Detachment, [1710](#)  
Rhesus Factor, [1710](#)  
Rhesus Incompatibility, [1711](#)  
Rhesus System, [1711](#)  
Rheumatic Chorea, [1711](#)  
Rheumatic Disease, [1711](#)  
Rheumatic Fever, [1711](#)  
Rheumatic Heart Disease, [1711](#)  
Rheumatism, [1711](#)  
Rheumatoid Arthritis, [1711](#)  
Rheumatoid Arthritis (RA), [1733](#)  
Rheumatoid Arthritis and Pregnancy, [1711](#)  
Rheumatoid Arthritis Cause Gastrointestinal Issues, [1711](#)  
Rheumatoid Arthritis Inflammation Of The Brain, [1711](#)  
Rheumatoid Factor, [1711](#)  
Rheumatoid Lung Disease, [1712](#)  
Rheumatoid Nodule, [1712](#)  
Rheumatoid Nodules, [1712](#)  
Rheumatoid Pneumoconiosis, [1712](#)  
Rheumatoid Synovitis, [1712](#)  
Rheumatoid Vasculitis, [1712](#)  
Rheumatologist, [1712](#)  
Rheumatology, [1712](#)  
Rhinencephalon, [1712](#)  
Rhinitis, [1712](#)  
Rhinitis Medicamentosa, [1712](#)  
Rhinocephaly, [1712](#)  
Rhinocerebral Mucormycosis, [1713](#)  
Rhinophyma, [1713](#)  
Rhinoplasty, [1713](#)  
Rhinorrhea, [1713](#)  
Rhinorrhoea, [1713](#)  
Rhinoscleroma, [1713](#)  
Rhinoscopy, [1713](#)  
Rhinosporidiosis, [1713](#)  
Rhinovirus, [1713](#)  
Rhinovirus A, [1713](#)  
Rhinovirus C, [1713](#)  
Rhinoviruses, [1713](#)  
Rhizomucor, [1714](#)  
Rhizomucor Pusillus, [1714](#)  
Rhizopus, [1714](#)  
Rhizopus Oryzae, [1714](#)  
Rhizotomy, [1714](#)  
Rhodium, [1714](#)  
Rhodococcus, [1714](#)  
Rhodococcus Equi, [1714](#)  
Rhodopsin, [1714](#)  
Rhodospirillaceae, [1714](#)  
Rhodospirillum, [1714](#)  
RhoGAM, [1714](#)  
Rhombencephalon, [1714](#)  
Rhonchi, [1714](#)  
Rhonchus, [1714](#)  
Rhubarb Leaves Poisoning, [1715](#)  
Rhythm Method, [1715](#)  
Rib, [1715](#)  
Rib Angle, [1715](#)  
Rib Cage, [1715](#)  
Rib Fracture, [1715](#)  
Rib Head, [1715](#)  
Rib Neck, [1715](#)  
Rib Tubercle, [1716](#)  
Ribavirin, [1715](#)  
Ribcage Pain, [1715](#)  
Riboflavin, [1715](#)  
Riboflavin Deficiency, [1715](#)  
Ribonuclease, [1715](#)  
Ribonuclease H, [1715](#)  
Ribonuclease III, [1715](#)  
Ribonucleic Acid, [1716](#)  
Ribonucleoprotein, [1716](#)  
Ribonucleoside, [1716](#)  
Ribonucleotide Reductase, [1716](#)

---

---

Ribonucleotides, [1716](#)  
 Ribophagy, [1716](#)  
 Ribose, [1716](#)  
 Ribosomal Protein, [1716](#)  
 Ribosomal RNA, [1716](#)  
 Ribosome, [1716](#)  
 Ribosome Subunit, [1716](#)  
 Richter Transformation, [1716](#)  
 Richter's Syndrome, [1716](#)  
 Ricin, [1716](#)  
 Rickets, [1717](#)  
 Rickettsia, [1717](#)  
 Rickettsia Felis, [1717](#)  
 Rickettsia Infections, [1717](#)  
 Rickettsia Japonica, [1717](#)  
 Rickettsia Prowazekii, [1717](#)  
 Rickettsia Rickettsii, [1717](#)  
 Rickettsia Typhi, [1717](#)  
 Rickettsiaceae, [1717](#)  
 Rickettsialpox, [1717](#)  
 Riddoch Phenomenon, [1717](#)  
 Ridge Preservation, [1718](#)  
 Ridged Cranial Sutures, [1717](#)  
 Ridged Nail, [1717](#)  
 Riedel Thyroiditis, [1718](#)  
 Riedel's Thyroiditis, [1718](#)  
 Rieger Syndrome, [1718](#)  
 Riehl Melanosis (Pigmented Contact Dermatitis), [1718](#)  
 Rifabutin, [1718](#)  
 Rifampicin, [1718](#)  
 Rifampin, [1718](#)  
 Rifampin Toxicity, [1718](#)  
 Rifamycin, [1718](#)  
 Rifaximin, [1718](#)  
 Rift Valley Fever, [1718](#)  
 Right Atrium, [1718](#)  
 Right Bundle Branch, [1719](#)  
 Right Bundle Branch Block, [1719](#)  
 Right Coronary Artery, [1719](#)  
 Right Coronary Artery Occlusion, [1719](#)  
 Right Heart Failure, [1719](#)  
 Right Heart Ventricular Angiography, [1719](#)  
 Right Hypochondriac Region, [1719](#)  
 Right Iliac Region, [1719](#)  
 Right Lower Quadrant Abdominal Pain, [1719](#)  
 Right Lymphatic Duct, [1719](#)  
 Right Sixth Cranial Nerve Palsy, [1719](#)  
 Right Ureterovesical Calculus, [1719](#)  
 Right Ventricle, [1719](#)  
 Right Ventricular Strain, [1720](#)  
 Righting Reflex, [1719](#)  
 Rigid And Flexible Sigmoidoscopy, [1720](#)  
 Rigidity, [1720](#)  
 Rigler Sign, [1720](#)  
 Rigler Triad, [1720](#)  
 Rigor, [1720](#)  
 Rigor Mortis, [1720](#)  
 Rilpivirine, [1720](#)  
 Riluzole, [1720](#)  
 Rimmed Vacuoles, [1720](#)  
 Ring Block, [1720](#)  
 Ring Chromosome, [1720](#)  
 Ring Vaccination, [1721](#)  
 Ringing In The Ear, [1720](#)  
 Ringing In The Ears, [1720](#)  
 Ringworm, [1721](#)  
 Ringworm (Tinea), [1733](#)  
 Ringworm Of The Body, [1721](#)  
 Ringworm Of The Foot, [1721](#)  
 Ringworm Of The Scalp, [1721](#)  
 Rinne and Weber Tests, [1721](#)  
 Rinne Test, [1721](#)  
 Rinnes Test, [1721](#)  
 RIPE Therapy, [1721](#)  
 Rippling Muscle Disease, [1721](#)  
 Risedronate Sodium, [1721](#)  
 Risk, [1721](#)  
 Risk Adjustment, [1721](#)  
 Risk Factor, [1721](#)  
 Risk Management, [1721](#)  
 Risk Register, [1722](#)  
 Risk-Taking, [1722](#)  
 Risks Of Hip And Knee Replacement, [1722](#)  
 Risks Of Tobacco, [1722](#)  
 Risks Of Underage Drinking, [1722](#)  
 Risky Drinking, [1722](#)  
 Risperidone, [1722](#)  
 Ristocetin-Induced Platelet Aggregation, [1722](#)  
 Risus Sardonicus, [1722](#)  
 Ritonavir, [1722](#)  
 Ritual, [1722](#)  
 Rituximab, [1722](#)  
 Rivaroxaban, [1722](#)  
 Rivastigmine, [1722](#)  
 River Blindness, [1723](#)  
 RLS, [1723](#)  
 RMDs, [1723](#)  
 RNA, [1723](#)  
 RNA Binding, [1723](#)  
 RNA Biosynthesis, [1723](#)  
 RNA Degradation, [1723](#)  
 RNA Folding, [1723](#)  
 RNA Helicase, [1723](#)  
 RNA Interference, [1723](#)  
 RNA Splicing, [1723](#)  
 RNA Transport, [1723](#)  
 RNA Virus, [1723](#)  
 RNase P, [1723](#)

---

---

Road Rage, [1723](#)  
Robinow Syndrome, [1723](#)  
Robot-Assisted Rehabilitation, [1723](#)  
Robotic Hysterectomy, [1724](#)  
Robotic Surgery, [1724](#)  
Robotic-Assisted Thoracoscopic Surgery, [1724](#)  
Rock (Drug), [2129](#)  
Rocker-Bottom Foot, [1724](#)  
Rocky Mountain Spotted Fever, [1724](#)  
Rod, [1724](#)  
Rod Photoreceptor, [1724](#)  
Rodent Ulcer, [1724](#)  
Rodenticide Toxicity, [1724](#)  
Rods, [1724](#)  
Rods And Cones, [1724](#)  
Roentgen Rays, [1724](#)  
Rofecoxib, [1724](#)  
Romano-Ward Syndrome, [1725](#)  
Romanus Lesion, [1725](#)  
Romberg Sign, [1725](#)  
Romberg Test, [1725](#)  
Romberg's Sign, [1725](#)  
Rombergism, [1725](#)  
Romidepsin, [1725](#)  
Romiplostim, [1725](#)  
Rongeur, [1725](#)  
Ronnell, [1725](#)  
Root Canal, [1725](#)  
Root Canal Therapy, [1725](#)  
Root Canal Treatment, [1725](#)  
Root Caries, [1725](#)  
Root Filling, [1725](#)  
Root Fracture, [1725](#)  
Rooting Reflex, [1726](#)  
Ropinirole, [1726](#)  
Ropivacaine, [1726](#)  
Ropivacaine Hydrochloride, [1726](#)  
Rorschach Test, [1726](#)  
Rosacea, [1726](#)  
Rosai-Dorfman Disease, [1726](#)  
Rose Bengal, [1726](#)  
Roseola, [1726](#)  
Roseola Infantum, [1726](#)  
Roseolovirus, [1726](#)  
Rosiglitazone, [1726](#)  
Rosiglitazone Maleate, [1727](#)  
Ross Procedure, [1727](#)  
Ross River Virus, [1727](#)  
Rostral, [1727](#)  
Rosuvastatin Calcium, [1727](#)  
Rotation, [1727](#)  
Rotator Cuff, [1727](#)  
Rotator Cuff Arthropathy, [1727](#)  
Rotator Cuff Exercises, [1727](#)  
Rotator Cuff Injury, [1727](#)  
Rotator Cuff Problems, [1727](#)  
Rotator Cuff Repair, [1727](#)  
Rotator Cuff Tear, [1727](#)  
Rotator Cuff Tendinitis, [1727](#)  
Rotavirus, [1727](#)  
Rotavirus Antigen Test, [1728](#)  
Rotavirus Infection, [1728](#)  
Rotavirus Vaccine, [1728](#)  
Rotaviruses, [1728](#)  
Rotenone, [1728](#)  
Roth Spots, [1728](#)  
Roth's Spots, [1728](#)  
Rothmund-Thomson Syndrome, [1728](#)  
Rotor Syndrome, [1728](#)  
Rotter Nodes, [1728](#)  
Rotterdam Criteria, [1728](#)  
Rouleau, [1728](#)  
Rouleaux, [1728](#)  
Rouleaux Formation, [1728](#)  
Round, [1729](#)  
Round Back, [1729](#)  
Round Ligament, [1729](#)  
Round Ligament of the Uterus, [1729](#)  
Round Ligament Pain, [1729](#)  
Round Pneumonia, [1729](#)  
Roundworm, [1729](#)  
Roundworms, [1729](#)  
Roussy-Levy Syndrome, [1729](#)  
Route Of Administration, [1729](#)  
Roux Stasis Syndrome, [1729](#)  
Roux-en-Y Gastric Bypass, [1729](#)  
Rovsing Sign, [1730](#)  
Roxithromycin, [1730](#)  
RPR Test, [1730](#)  
RS3PE Syndrome, [1730](#)  
RSI, [1730](#)  
RSV, [1730](#)  
RSV Antibody Test, [1730](#)  
RSV Infection, [1730](#)  
RT-PCR, [1730](#)  
Rubber, [1730](#)  
Rubber Band Ligation, [1730](#)  
Rubber Cement Poisoning, [1731](#)  
Rubber Dams, [1731](#)  
Rubella, [1731](#)  
Rubella Virus, [1731](#)  
Rubeola, [1731](#)  
Rubeosis Iridis, [1731](#)  
Rubidium, [1731](#)  
Rubin Maneuver, [1731](#)  
Rubinstein-Taybi Syndrome, [1731](#)  
Rubivirus, [1731](#)  
Rubredoxins, [1731](#)

---



---

Rubricyte Count, [1731](#)  
 Rule Of Nines, [1731](#)  
 Rule Out, [1731](#)  
 Rumination, [1732](#)  
 Rumination Disorder, [1732](#)  
 Rumination Syndrome, [1732](#)  
 Ruminococcus Gnavus, [1732](#)  
 Rumpel-Leede Sign, [1732](#)  
 Rumpel-Leede Test, [1732](#)  
 Runner's Knee, [1732](#)  
 Runny Nose, [1732](#)  
 Rupture, [1732](#)  
 Ruptured Abdominal Aortic Aneurysm, [1732](#)  
 Ruptured Disk, [1732](#)  
 Ruptured Eardrum, [1732](#)  
 Ruptured Perforated Eardrum, [1732](#)  
 Russell Viper Venom Time, [1733](#)  
 Russell-Silver Syndrome, [1732](#)  
 Rx, [1733](#)  
 Ryle's Tube, [1733](#)  
 RâLe, [1723](#)  
  
 S Phase, [1819](#)  
 S-Adenosylmethionine, [1737](#)  
 S3 Gallop Pathology, [1735](#)  
 S4 Gallop Pathology, [1735](#)  
 SA Node, [1742](#)  
 SAA, [1735](#)  
 Sabin Vaccine, [1735](#)  
 Sabre Tibia, [1735](#)  
 Saccade, [1735](#)  
 Saccular Aneurysm, [1735](#)  
 Saccule, [1735](#)  
 Sachet Poisoning, [1735](#)  
 Sacral, [1735](#)  
 Sacral Agenesis, [1735](#)  
 Sacral Dimple, [1736](#)  
 Sacral Fistula, [1736](#)  
 Sacral Nerve Stimulation, [1736](#)  
 Sacral Nerve Stimulator, [1736](#)  
 Sacral Nerves, [1736](#)  
 Sacral Plexus, [1736](#)  
 Sacral Region, [1736](#)  
 Sacral Vertebra, [1736](#)  
 Sacral Vertebrae, [1736](#)  
 Sacrococcygeal, [1736](#)  
 Sacrococcygeal Chordoma, [1880](#)  
 Sacrococcygeal Teratoma, [1736](#)  
 Sacrocolpopexy, [1736](#)  
 Sacroiliac Joint, [1736](#)  
 Sacroiliac Joint Dysfunction, [1736](#)  
 Sacroiliac Joint Injury, [1736](#)  
 Sacroiliac Joint Pain, [1737](#)  
 Sacroiliitis, [1737](#)  
  
 Sacrospinous Ligament Fixation, [1737](#)  
 Sacrum, [1737](#)  
 SAD, [1737](#)  
 SAD (Seasonal Affective Disorder), [1880](#)  
 Sad Mood, [1738](#)  
 Saddle Anesthesia, [1737](#)  
 Saddle Joint, [1737](#)  
 Saddle Nose Deformity, [1737](#)  
 Saddle Pulmonary Embolism, [1737](#)  
 Sadism, [1737](#)  
 Safe Sex, [1738](#)  
 Safety Of Drugs, [1738](#)  
 Safety Plan, [1738](#)  
 Safety Study, [1738](#)  
 Saffron, [1738](#)  
 Safrole, [1738](#)  
 Sagittal, [1738](#)  
 Sagittal Axis, [1738](#)  
 Sagittal Plane, [1738](#)  
 Sagittal Sinus Thrombosis, [1738](#)  
 Salaam Attacks, [1738](#)  
 Salbutamol, [1738](#)  
 Salicylate, [1738](#)  
 Salicylate Toxicity, [1739](#)  
 Salicylic Acid, [1739](#)  
 Saline, [1739](#)  
 Saline Infusion Sonohysterography, [1739](#)  
 Saline Nasal Washes, [1739](#)  
 Saliva, [1739](#)  
 Salivary Duct Stones, [1739](#)  
 Salivary Gland, [1739](#)  
 Salivary Gland Biopsy, [1739](#)  
 Salivary Gland Cancer, [1739](#)  
 Salivary Gland Infection, [1739](#)  
 Salivary Gland Infections, [1739](#)  
 Salivary Gland Neoplasm, [1739](#)  
 Salivary Gland Scintigraphy, [1740](#)  
 Salivary Gland Stones, [1740](#)  
 Salivary Gland Tumors, [1740](#)  
 Salivary Glands, [1739](#)  
 Salivation, [1740](#)  
 Salk Vaccine, [1740](#)  
 Salmon Patches, [1741](#)  
 Salmonella, [1740](#)  
 Salmonella Choleraesuis, [1740](#)  
 Salmonella Enterica, [1740](#)  
 Salmonella Enterocolitis, [1740](#)  
 Salmonella Food Poisoning, [1740](#)  
 Salmonella Infection, [1740](#)  
 Salmonella Infections, [1740](#)  
 Salmonella Typhi, [1740](#)  
 Salmonella Typhimurium, [1740](#)  
 Salmonella Typhisuis, [1740](#)  
 Salmonella Vaccines, [1741](#)

---

---

Salmonellosis, [1741](#)  
 Salpingectomy, [1741](#)  
 Salpingitis, [1741](#)  
 Salpingitis Isthmica Nodosa, [1741](#)  
 Salpingo-Oophorectomy, [1741](#)  
 Salpingo-OöPhorectomy, [1741](#)  
 Salpingography, [1741](#)  
 Salpingostomy, [1741](#)  
 Salt, [1741](#)  
 Saltatory Conduction, [1741](#)  
 Salter-Harris Fracture, [1741](#)  
 Salter-Harris Fractures, [1742](#)  
 Salvage Therapy, [1742](#)  
 SAM-e, [1880](#)  
 Sample Fixation, [1742](#)  
 Samter Triad, [1742](#)  
 Sandfly Fever, [1742](#)  
 Sandhoff Disease, [1742](#)  
 Sandifer Syndrome, [1742](#)  
 Sandpaper Tooth, [1742](#)  
 Sanfilippo Syndrome, [1742](#)  
 Sanguineous, [1742](#)  
 Sanitation, [1742](#)  
 Sano Shunt, [1743](#)  
 Santorini Duct, [1743](#)  
 Saphenous, [1743](#)  
 Saphenous Nerve, [1743](#)  
 Saphenous Vein, [1743](#)  
 SAPHO Syndrome, [1743](#)  
 Sapovirus, [1743](#)  
 Sapovirus Infection, [1743](#)  
 Sapropterin, [1743](#)  
 Saquinavir, [1743](#)  
 Saquinavir Mesylate, [1743](#)  
 Sarcocystis Hominis, [1743](#)  
 Sarcocystis Suihominis, [1743](#)  
 Sarcocystosis, [1744](#)  
 Sarcoglycan Complex, [1744](#)  
 Sarcoglycanopathy, [1744](#)  
 Sarcoid Arthropathy, [1744](#)  
 Sarcoid Of Boeck, [1880](#)  
 Sarcoidosis, [1744](#)  
 Sarcoidosis Skin Manifestations, [1744](#)  
 Sarcolemma, [1744](#)  
 Sarcoma, [1744](#)  
 Sarcomere, [1744](#)  
 Sarcopenia, [1744](#)  
 Sarcopenic Obesity, [1744](#)  
 Sarcoplasm, [1744](#)  
 Sarcoplasmic Reticulum, [1744](#)  
 Sarcoptes, [1745](#)  
 Sarcoptes Scabiei, [1745](#)  
 Sarcoptic Mange, [1745](#)  
 Sargramostim, [1745](#)  
 Sarnat Staging, [1745](#)  
 SARS, [1745](#)  
 SARS (Severe Acute Respiratory Syndrome), [1880](#)  
 SARS Virus, [1745](#)  
 SARS-CoV-2, [1745](#)  
 Sartorius Muscle, [1745](#)  
 Sassafras Oil Overdose, [1745](#)  
 Satellite DNA, [1745](#)  
 Satellite Virus, [1745](#)  
 Satiation, [1745](#)  
 Satiety, [1745](#)  
 Saturated Fat, [1746](#)  
 Saturation, [1746](#)  
 Sausage Digit, [1746](#)  
 Saxitoxin, [1746](#)  
 SBBO, [1746](#)  
 Scab, [1746](#)  
 Scabicide, [2129](#)  
 Scabies, [1746](#)  
 SCAD, [1746](#)  
 Scaffold, [1746](#)  
 Scaffold Seeding, [1746](#)  
 Scalded Mouth Syndrome, [1746](#)  
 Scalded Skin Syndrome, [1746](#)  
 Scalded-Skin Syndrome, [1746](#)  
 Scalds, [1746](#)  
 Scalene Muscles, [1746](#)  
 Scales, [1747](#)  
 Scaling, [1747](#)  
 Scaling And Root Planing, [1747](#)  
 Scalp, [1747](#)  
 Scalp and Body Ringworm, [1747](#)  
 Scalp Psoriasis, [1747](#)  
 Scalp Ringworm, [1747](#)  
 Scalp Tenderness, [1747](#)  
 Scalpel, [1747](#)  
 Scalpel Handle, [1747](#)  
 Scan, [1747](#)  
 Scan Gallium, [1747](#)  
 Scanning, [1747](#)  
 Scanning Electron Microscopy, [1748](#)  
 Scanning Speech, [1748](#)  
 Scanning Techniques, [1748](#)  
 Scaphoid, [1748](#)  
 Scaphoid Abdomen, [1748](#)  
 Scaphoid Bone, [1748](#)  
 Scaphoid Fractures, [1748](#)  
 Scapholunate Advanced Collapse, [1748](#)  
 Scaphycephaly, [1748](#)  
 Scapula, [1748](#)  
 Scapular Winging, [1748](#)  
 Scapulothoracic Joint, [1748](#)  
 Scar, [1748](#)  
 Scar Revision, [1749](#)

---

---

Scarlatina, [1748](#)  
Scarlet Fever, [1749](#)  
Scarring Alopecia, [1749](#)  
Scarring Bullosa, [1880](#)  
Scars, [1749](#)  
Scatter Correction, [1749](#)  
Scatter Events, [1749](#)  
Scavenger Receptor, [1749](#)  
Scavenging System, [1749](#)  
Scedosporium, [1749](#)  
Scedosporium Apiospermum, [1749](#)  
Scedosporium Prolifican, [1749](#)  
Schaalia Radingae, [1749](#)  
Schaalia Turicensis, [1749](#)  
Schatzki Ring, [1749](#)  
Schereshevskii-Turner Syndrome, [1881](#)  
Scheuermann Disease, [1750](#)  
Scheuermann's Disease, [1750](#)  
Schiff Base, [1750](#)  
Schiller Test, [1750](#)  
Schilling Test, [1750](#)  
Schirmer Test, [1750](#)  
Schistocyte Count, [1750](#)  
Schistocytes, [1750](#)  
Schistomicide, [1750](#)  
Schistosoma, [1750](#)  
Schistosoma Haematobium, [1750](#)  
Schistosoma Intercalatum, [1750](#)  
Schistosomiasis, [1750](#)  
Schistosomiasis (Bilharzia), [1751](#)  
Schizencephaly, [1751](#)  
Schizoaffective Disorder, [1751](#)  
Schizogony, [1751](#)  
Schizoid Personality Disorder, [1751](#)  
Schizophrenia, [1751](#)  
Schizotypal Personality Disorder, [1751](#)  
Schlemm Canal, [1751](#)  
Schlemm's Canal, [1751](#)  
Schmorl Node, [1751](#)  
Schnitzler Syndrome, [1752](#)  
School Sores, [1752](#)  
School-Age Children Development, [1752](#)  
School-Based Rehabilitation, [1752](#)  
Schwann Cell, [1752](#)  
Schwann Cells, [1752](#)  
Schwannoma, [1752](#)  
Schwannomatosis, [1752](#)  
SCI, [1752](#)  
Sciatic Nerve, [1752](#)  
Sciatic Neuritis, [1752](#)  
Sciatica, [1752](#)  
Scimitar Syndrome, [1752](#)  
Scintigraphy, [1752](#)  
Scintillation, [1752](#)  
Scintillation Crystal, [1753](#)  
Scintillations, [1753](#)  
Scirrhus, [1753](#)  
Scissors, [1753](#)  
Sclera, [1753](#)  
Scleral Buckling, [1753](#)  
Scleral Icterus, [1753](#)  
Scleral Inflammation, [1753](#)  
Scleredema, [1753](#)  
Scleredema Diabeticorum, [1753](#)  
Scleritis, [1753](#)  
Sclerodactyly, [1753](#)  
Scleroderma, [1753](#)  
Scleroderma Renal Crisis, [1754](#)  
Scleroma, [1754](#)  
Scleromyxedema, [1754](#)  
Sclerosing Cholangitis, [1754](#)  
Sclerosing Mesenteritis, [1754](#)  
Sclerosing Peritonitis, [1754](#)  
Sclerosing Sialadenitis, [1754](#)  
Sclerosis, [1754](#)  
Sclerosteosis, [1754](#)  
Sclerotherapy, [1754](#)  
Sclerotomy, [1754](#)  
Scoliosis, [1755](#)  
Scombroid Poisoning, [1755](#)  
Scopolamine, [1755](#)  
Scopolamine Hydrobromide, [1755](#)  
Scopulariopsis, [1755](#)  
Scorbutic, [1755](#)  
Scorpion Fish Sting, [1755](#)  
Scorpion Sting, [1755](#)  
Scorpion Stings, [1755](#)  
Scorpions, [1755](#)  
Scotoma, [1755](#)  
Scotty Dog Sign, [1755](#)  
Scrape, [1756](#)  
Screen Time And Children, [1756](#)  
Screening, [1756](#)  
Screening And Diagnosis For HIV, [1756](#)  
Screening Procedure, [1756](#)  
Screening Program, [1756](#)  
Screening Study, [1756](#)  
Screening Test, [1756](#)  
Screws, [1756](#)  
Script, [1756](#)  
Scrivener's Palsy, [1756](#)  
Scrotal Calcinosiis, [1756](#)  
Scrotal Infection, [1756](#)  
Scrotal Masses, [1756](#)  
Scrotal Pain, [1756](#)  
Scrotal Septum, [1757](#)  
Scrotal Swelling, [1757](#)  
Scrotal Ultrasound, [1757](#)

---

---

Scrotum, [1757](#)  
 Scrub In, [1757](#)  
 Scrub Typhus, [1757](#)  
 Scrub Up, [1757](#)  
 Scrubs, [1757](#)  
 Scurf, [1757](#)  
 Scurvy, [1757](#)  
 Sea-Blue Histiocyte, [1757](#)  
 Sea-Sickness, [1757](#)  
 Seabather's Eruption, [1757](#)  
 Seafood Poisoning, [1757](#)  
 Seasonal Affective Disorder, [1758](#)  
 Seasonal Affective Disorder (SAD), [1758](#)  
 Seasonal Affective Disorder Syndrome, [1758](#)  
 Seasonal Allergies, [1758](#)  
 Sebaceous Adenoma, [1758](#)  
 Sebaceous Carcinoma, [1758](#)  
 Sebaceous Cyst, [1758](#)  
 Sebaceous Epithelioma, [1758](#)  
 Sebaceous Gland, [1758](#)  
 Sebaceous Gland Carcinoma, [1758](#)  
 Sebaceous Glands, [1758](#)  
 Sebaceous Hyperplasia, [1758](#)  
 Seborrhea, [1758](#)  
 Seborrheic Dermatitis, [1759](#)  
 Seborrheic Keratoses, [1759](#)  
 Seborrheic Keratosis, [1759](#)  
 Seborrhoea, [1759](#)  
 Seborrhoeic, [1759](#)  
 Seborrhoeic Keratosis, [1759](#)  
 Sebum, [1759](#)  
 Seckel Syndrome, [1759](#)  
 Secobarbital Overdose, [1759](#)  
 Second Malignancy, [1761](#)  
 Second Messenger Systems, [1762](#)  
 Second Messengers, [1761](#)  
 Second Opinion, [1762](#)  
 Second Stage of Labor, [1762](#)  
 Second Trimester, [1762](#)  
 Second-Degree AV Block, [1761](#)  
 Second-Degree Burn, [1761](#)  
 Second-Hand Smoke, [1761](#)  
 Second-Look Surgery, [1761](#)  
 Secondaries In The Liver, [1152](#)  
 Secondary Adrenal Insufficiency, [1759](#)  
 Secondary Amenorrhea, [1759](#)  
 Secondary Attack Rate, [1759](#)  
 Secondary Brain Cancer, [1759](#)  
 Secondary Cancer, [1759](#)  
 Secondary Hyperparathyroidism, [1760](#)  
 Secondary Hypertension, [1760](#)  
 Secondary Hypothyroidism, [1760](#)  
 Secondary Immune Response, [1760](#)  
 Secondary Immunodeficiency, [1760](#)  
 Secondary Infections, [1760](#)  
 Secondary Lymphoid Organs, [1760](#)  
 Secondary Osteoporosis, [1760](#)  
 Secondary Osteosarcoma, [1760](#)  
 Secondary Parkinsonism, [1760](#)  
 Secondary Peritonitis, [1760](#)  
 Secondary Polycythemia, [1760](#)  
 Secondary Postpartum Hemorrhage, [1761](#)  
 Secondary Prevention, [1761](#)  
 Secondary Progressive Multiple Sclerosis, [1761](#)  
 Secondary Psychosis, [1881](#)  
 Secondary Sexual Characteristics, [1761](#)  
 Secondary Shock, [1761](#)  
 Secondary Syphilis, [1761](#)  
 Secondary Systemic Amyloidosis, [1761](#)  
 Secondary Tumor, [1761](#)  
 Secretin, [1762](#)  
 Secretin Stimulation Test, [1762](#)  
 Secretion, [1762](#)  
 Secretory Diarrhea, [1762](#)  
 Section, [1762](#)  
 Sed Rate, [1763](#)  
 Sedation, [1762](#)  
 Sedation Dentistry, [1762](#)  
 Sedation Levels, [1762](#)  
 Sedative, [1762](#)  
 Sedative, Hypnotic, Anxiolytic Use Disorders, [1763](#)  
 Sedative, Hypnotic, or Anxiolytic Use Disorder, [1763](#)  
 Sedative-Hypnotic, [1762](#)  
 Sedative-Hypnotic Toxidrome, [1763](#)  
 Sedatives, [1763](#)  
 Sedimentation Rate, [1763](#)  
 Segmental Innervation, [1763](#)  
 Segmental Pressures, [1763](#)  
 Segmentation, [1763](#)  
 Segmentectomy, [1763](#)  
 SEID, [1763](#)  
 Seidel Test, [1763](#)  
 Seizure, [1763](#)  
 Seizure Disorder, [1764](#)  
 Seizure Disorders, [1764](#)  
 Seizure Disorders in Pregnancy, [1764](#)  
 Seizures, [1764](#)  
 Selective Antibody Deficiency With Normal Immunoglobulins, [1764](#)  
 Selective Deficiency Of IgA, [1764](#)  
 Selective Estrogen Receptor Modulator, [1764](#)  
 Selective Estrogen Receptor Modulators, [1764](#)  
 Selective IgA Deficiency, [1764](#)  
 Selective Mutism, [1764](#)  
 Selective Serotonin Reuptake Inhibitor, [2129](#)  
 Selective Serotonin Reuptake Inhibitor (SSRI), [1764](#)  
 Selective Serotonin Reuptake Inhibitor Toxicity, [1765](#)  
 Selective Serotonin Reuptake Inhibitors, [1764](#)

---

---

Selective Serotonin-Reuptake Inhibitors, [1764](#)  
Selective Tubal Occlusion Procedure, [1765](#)  
Selegiline, [1765](#)  
Selenium, [1765](#)  
Selenium Deficiency, [1765](#)  
Selenium In Diet, [1765](#)  
Selenium Sulphide, [1765](#)  
Selenium-75, [1765](#)  
Self Harming, [1765](#)  
Self-Diagnosing Body Dysmorphic Disorder, [1765](#)  
Self-Diagnosis, [1765](#)  
Self-Efficacy, [1765](#)  
Self-Harm, [1765](#)  
Self-Help Group, [1765](#)  
Self-Injury/Cutting, [1766](#)  
Self-Management, [1766](#)  
Self-Poisoning, [1766](#)  
Self-Retaining Retraction, [1766](#)  
Self-Testing For COVID-19, [1766](#)  
Sella Turcica, [1766](#)  
Semaglutide, [1766](#)  
Semen, [1766](#)  
Semen Analysis, [1766](#)  
Semen DNA Analysis, [1766](#)  
Semi-Fowler Position, [1766](#)  
Semicircular Canals, [1766](#)  
Semilunar Cartilages, [1766](#)  
Semilunar Valves, [1766](#)  
Seminal Fluid Analysis, [1767](#)  
Seminal Vesicle, [1767](#)  
Seminal Vesicles, [1767](#)  
Seminiferous Tubules, [1767](#)  
Seminoma, [1767](#)  
Semont Maneuver, [1767](#)  
Senescence, [1767](#)  
Senile Lentigines, [1767](#)  
Senile Purpura, [1767](#)  
Senna, [1767](#)  
Sensate Focus Techniques, [1767](#)  
Sensation, [1767](#)  
Sense, [1767](#)  
Sensitisation, [1768](#)  
Sensitivity, [1768](#)  
Sensitivity Analysis, [1768](#)  
Sensitized, [1768](#)  
Sensorimotor Polyneuropathy, [1768](#)  
Sensorineural Deafness, [1768](#)  
Sensorineural Hearing Loss, [1768](#)  
Sensorium, [1768](#)  
Sensory, [1768](#)  
Sensory Cortex, [1768](#)  
Sensory Deprivation, [1768](#)  
Sensory Level, [1768](#)  
Sensory Nerves, [1768](#)  
Sensory Neuron, [1768](#)  
Sensory Processing Disorder, [1769](#)  
Sensory Receptors, [1769](#)  
Sentinel Event, [1769](#)  
Sentinel Injury, [1769](#)  
Sentinel Loop, [1769](#)  
Sentinel Lymph Node, [1769](#)  
Sentinel Lymph Node Biopsy For Melanoma, [1769](#)  
Sentinel Lymph-Node Biopsy, [1769](#)  
Sentinel Node, [1769](#)  
Sentinel Node Biopsy, [1769](#)  
Sentinel Surveillance, [1769](#)  
Separated Cranial Sutures, [1769](#)  
Separation Anxiety, [1769](#)  
Separation Anxiety Disorder, [1770](#)  
Separation Anxiety In Children, [1770](#)  
Separation Of The Iris, [1770](#)  
Sepsis, [1770](#)  
Sepsis Syndrome, [1770](#)  
Septal Defect, [1770](#)  
Septal Hematoma, [1770](#)  
Septate Uterus, [1770](#)  
Septic, [1770](#)  
Septic Abortion, [1770](#)  
Septic Arthritis, [1770](#)  
Septic Embolism, [1770](#)  
Septic Pulmonary Infarction, [1771](#)  
Septic Shock, [1771](#)  
Septic Shock Criteria, [1771](#)  
Septic Thrombophlebitis, [1771](#)  
Septicaemia, [1770](#)  
Septicemia, [1771](#)  
Septo-Optic Dysplasia, [1771](#)  
Septoplasty, [1771](#)  
Septum, [1771](#)  
Sequela, [1771](#)  
Sequelae, [1771](#)  
Sequential Therapy, [1771](#)  
Sequester, [1771](#)  
Sequestrum, [1771](#)  
Ser, [1771](#)  
Serine, [1772](#)  
Serious Adverse Event, [1772](#)  
Serious Health Problems Symptoms, [1772](#)  
SERMs, [1772](#)  
Seroconversion, [1772](#)  
Serodiagnostic, [1772](#)  
Serologic Test, [1772](#)  
Serology, [1772](#)  
Serology For Brucellosis, [1772](#)  
Seroma, [1772](#)  
Seronegative Arthritis, [1772](#)  
Serosanguineous, [1772](#)  
Serositis, [1772](#)

---

---

Serotonin, [1772](#)  
 Serotonin Blood Test, [1772](#)  
 Serotonin Syndrome, [1773](#)  
 Serotonin-Norepinephrine Reuptake Inhibitors, [1773](#)  
 Serotype, [1773](#)  
 Serous, [1773](#)  
 Serous Borderline Ovarian Tumor, [1773](#)  
 Serous Cystadenoma of Pancreas, [1773](#)  
 Serous Fluid Cytology, [1773](#)  
 Serous Membranes, [1773](#)  
 Serous Tubal Intraepithelial Carcinoma, [1773](#)  
 Serpentine Supravenous Hyperpigmentation, [1773](#)  
 Serrated Pathway, [1773](#)  
 Serrated Polyposis Syndrome, [1773](#)  
 Serration, [1773](#)  
 Serratus Anterior Muscle, [1773](#)  
 Sertoli Cell, [1774](#)  
 Sertoli Cells, [1774](#)  
 Sertoli-Cell Tumor, [1774](#)  
 Sertoli-Leydig Cell Tumor, [1774](#)  
 Sertoli-Leydig Tumor, [1774](#)  
 Serum, [1774](#)  
 Serum Albumin Test, [1774](#)  
 Serum Chloride Test, [1774](#)  
 Serum Concentration, [1774](#)  
 Serum Free Hemoglobin Test, [1774](#)  
 Serum Glutamic Oxaloacetic Transaminase, [1774](#)  
 Serum Glutamic Pyruvic Transaminase, [1774](#)  
 Serum Herpes Simplex Antibodies, [1774](#)  
 Serum Iron Test, [1774](#)  
 Serum Osmolal Gap, [1775](#)  
 Serum Phenylalanine Screening, [1775](#)  
 Serum Progesterone, [1775](#)  
 Serum Protein Electrophoresis, [1775](#)  
 Serum Sickness, [1775](#)  
 Serum Therapy, [1775](#)  
 Serum Uric Acid, [1775](#)  
 Serving Size, [1775](#)  
 Sesamoid Bones, [1775](#)  
 Sesamoiditis, [1775](#)  
 Sesquipedalian, [1775](#)  
 Sessile, [1775](#)  
 Set, [1775](#)  
 Sever Disease, [1775](#)  
 Severe Acute Respiratory Syndrome, [1776](#)  
 Severe Brain Injury Coma, [1776](#)  
 Severe Combined Immunodeficiency, [1776](#)  
 Severe Combined Immunodeficiency (SCID), [1776](#)  
 Severe Dengue, [1776](#)  
 Severe Hearing Loss, [1881](#)  
 Severe Obesity, [1776](#)  
 Sevoflurane, [1776](#)  
 Sex Change, [1776](#)  
 Sex Chromosomes, [1776](#)  
 Sex Cord-Stromal Ovarian Tumor, [1776](#)  
 Sex Headaches, [1776](#)  
 Sex Hormones, [1776](#)  
 Sex Therapy, [1777](#)  
 Sex-Linked Dominant, [1777](#)  
 Sex-Linked Inheritance, [1777](#)  
 Sex-Linked Recessive, [1777](#)  
 Sexual Abuse, [1777](#)  
 Sexual Addiction, [1777](#)  
 Sexual Assault, [1777](#)  
 Sexual Assault Prevention, [1777](#)  
 Sexual Dysfunction, [1777](#)  
 Sexual Dysfunction Disorders, [1777](#)  
 Sexual Health, [1777](#)  
 Sexual Intercourse, [1777](#)  
 Sexual Masochism Disorder, [1778](#)  
 Sexual Obsession, [1778](#)  
 Sexual Rehabilitation, [1778](#)  
 Sexual Sadism, [1778](#)  
 Sexual Sadism Disorder, [1778](#)  
 Sexual Violence, [1778](#)  
 Sexually Transmitted Disease, [1777](#)  
 Sexually Transmitted Disease (STD), [1777](#)  
 Sexually Transmitted Disease Treatments, [1778](#)  
 Sexually Transmitted Diseases, [1777](#)  
 Sexually Transmitted Diseases in Pregnancy, [1777](#)  
 Sexually Transmitted Diseases in Women, [1778](#)  
 Sexually Transmitted Infection, [1778](#)  
 Sexually Transmitted Infections, [1778](#)  
 Sezary Disease, [1778](#)  
 Sezary Syndrome, [1778](#)  
 Sgarbossa Criteria, [1778](#)  
 SGLT2 Inhibitor, [2129](#)  
 Shadowing Artifact, [1778](#)  
 Shaken Baby Syndrome, [1778](#)  
 Shaken Impact Syndrome, [1778](#)  
 Shampoo Ingestion, [1779](#)  
 Shared Decision-Making, [1779](#)  
 Shark Attack, [1779](#)  
 Sharp Force Trauma, [1779](#)  
 Sharp-Force Defense Wounds, [1779](#)  
 Sharps Container, [1779](#)  
 Sharps Injury, [1779](#)  
 Shave Biopsy, [1779](#)  
 Sheehan Syndrome, [1779](#)  
 Sheehan's Syndrome, [1779](#)  
 Shellac Poisoning, [1779](#)  
 Shellfish Allergy, [1779](#)  
 Shellfish Poisoning, [1780](#)  
 Shielding, [1780](#)  
 Shift Work Sleep Disorder, [1780](#)  
 Shifting Dullness, [1780](#)  
 Shiga Toxin-Producing Escherichia coli Infection, [1780](#)  
 Shigella, [1780](#)

---



---

Shigellosis, [1780](#)  
Shilajit, [1780](#)  
Shin Bone Fever, [1780](#)  
Shin Splints, [1780](#)  
Shin Spot, [1780](#)  
Shingles, [1780](#)  
Shingles And Pregnancy, [1780](#)  
Shirodkar Cerclage, [1781](#)  
Shisha, [1781](#)  
Shivering, [1781](#)  
Shock, [1781](#)  
Shock and Pregnancy, [1781](#)  
Shock Lung, [1781](#)  
Shock Wave Lithotripsy, [1781](#)  
Shone Complex, [1781](#)  
Short Arm Of A Chromosome, [1781](#)  
Short Bones, [1781](#)  
Short Bowel Syndrome, [1781](#)  
Short Philtrum, [1781](#)  
Short QT Syndrome, [1782](#)  
Short Stature, [1782](#)  
Short-Chain Fatty Acids, [1781](#)  
Short-Sight, [1782](#)  
Short-Term Memory, [1782](#)  
Shortness Of Breath, [1781](#)  
Shotty, [1782](#)  
Shoulder, [1782](#)  
Shoulder Arthroscopy, [1782](#)  
Shoulder Bursitis, [1782](#)  
Shoulder CT Scan, [1782](#)  
Shoulder Dislocation, [1782](#)  
Shoulder Dystocia, [1782](#)  
Shoulder Impingement, [1782](#)  
Shoulder Impingement Syndrome, [1783](#)  
Shoulder Instability, [1783](#)  
Shoulder MRI Scan, [1783](#)  
Shoulder Pain, [1783](#)  
Shoulder Replacement, [1783](#)  
Shulman Syndrome, [1783](#)  
Shunt, [1783](#)  
Shwachman Syndrome, [1783](#)  
Shy-Drager Syndrome, [1783](#)  
SI Units, [1794](#)  
SIADH, [1783](#)  
Sialadenitis, [1783](#)  
Sialogogues, [1783](#)  
Sialidosis (Mucopolipidosis Type I), [1784](#)  
Sialogram, [1783](#)  
Sialography, [1783](#)  
Sialolithiasis, [1784](#)  
Sialorrhea, [1784](#)  
Sialorrhoea, [1784](#)  
Sialuria, [1784](#)  
Sibling, [1784](#)  
SIBO, [1784](#)  
Sicca Syndrome, [1784](#)  
Sick Sinus Syndrome, [1785](#)  
Sickle Cell, [1784](#)  
Sickle Cell Anemia, [1784](#)  
Sickle Cell Crisis, [1784](#)  
Sickle Cell Dactylitis, [1784](#)  
Sickle Cell Disease, [1784](#)  
Sickle Cell Test, [1785](#)  
Sickle Cell Trait, [1785](#)  
Sickle-Cell Anaemia, [1784](#)  
Side Effect, [1785](#)  
Side Effects, [1785](#)  
Side-Effects, [1785](#)  
Sideroblastic Anemia, [1785](#)  
Siderosis, [1785](#)  
SIDS, [1785](#)  
Sievert, [1785](#)  
Sight, [1785](#)  
Sigmoid, [1785](#)  
Sigmoid Colon, [1785](#)  
Sigmoid Volvulus, [1786](#)  
Sigmoidoscopy, [1785](#)  
Sign, [1786](#)  
Sign Language, [1786](#)  
Sign of Leser-Trelat, [1786](#)  
Signal Peptide, [1786](#)  
Signal Transduction, [1786](#)  
Signal-To-Noise Ratio, [1786](#)  
Signature, [1786](#)  
Signature Strengths, [1786](#)  
Signet Ring Cell Carcinoma, [1786](#)  
Signs Of An Asthma Attack, [1786](#)  
Sildenafil Citrate, [1786](#)  
Silent Heart Attack, [1786](#)  
Silent Ischemia, [1786](#)  
Silent Mutation, [1787](#)  
Silent Myocardial Infarction, [1787](#)  
Silent Sinus Syndrome, [1787](#)  
Silent Thyroiditis, [1787](#)  
Silicate, [1787](#)  
Silicon Dioxide, [1787](#)  
Silicone, [1787](#)  
Silicone Elastomer, [1787](#)  
Silicone Implants, [1787](#)  
Silicone Mastitis, [1787](#)  
Silicone Oil Hyperoleon, [1787](#)  
Silicones, [1787](#)  
Silicosis, [1787](#)  
Silk, [1787](#)  
Silk Road Disease, [1788](#)  
Siltuximab, [1788](#)  
Silver, [1788](#)  
Silver Diamine Fluoride, [1788](#)

---

---

Silver Nitrate, [1788](#)  
Silver Poisoning, [1788](#)  
Silver Sulfadiazine, [1788](#)  
Silver-Russell Syndrome, [1788](#)  
Silymarin, [1788](#)  
Simeprevir, [1788](#)  
Simethicone, [1788](#)  
Simian Crease, [1788](#)  
Simmonds Disease, [1788](#)  
Simmonds Test, [1788](#)  
Simmonds-Thompson Test, [1788](#)  
Simple Febrile Seizure, [1789](#)  
Simple Goiter, [1789](#)  
Simple Mastectomy, [1789](#)  
Simple Prostatectomy, [1789](#)  
Simple Pulmonary Eosinophilia, [1789](#)  
Simplexvirus, [1789](#)  
Simpson Test, [1789](#)  
Simuliidae, [1789](#)  
Simultanagnosia, [1789](#)  
Simvastatin, [1789](#)  
Sin Nombre Virus Infection, [1790](#)  
Sincalide, [1789](#)  
Sindbis Virus, [1789](#)  
Sinding-Larsen-Johansson Syndrome, [1789](#)  
Sinecatechins, [1789](#)  
Singer's Nodule, [1790](#)  
Singing, [1790](#)  
Single Nucleotide Polymorphism, [1790](#)  
Single Palmar Crease, [1790](#)  
Single Photon Emission Computed Tomography, [1790](#)  
Single Ventricle, [1790](#)  
Single-Dose Toxicity, [1790](#)  
Single-Fiber Electromyography, [1790](#)  
Single-Photon Absorptiometry, [1790](#)  
Single-Photon Emission Computed Tomography, [1790](#)  
Single-Photon Emission Computed Tomography/Computed Tomography, [1790](#)  
Singlet Oxygen, [1790](#)  
Singultus, [1790](#)  
Sinoatrial Block, [1791](#)  
Sinoatrial Node, [1791](#)  
Sinuatrial Node, [1791](#)  
Sinus, [1791](#)  
Sinus Arrest, [1791](#)  
Sinus Arrhythmia, [1791](#)  
Sinus Bradycardia, [1791](#)  
Sinus CT Scan, [1791](#)  
Sinus Headache, [1791](#)  
Sinus Histiocytosis, [1791](#)  
Sinus Lift, [1792](#)  
Sinus MRI Scan, [1792](#)  
Sinus Node, [1792](#)  
Sinus Pain, [1792](#)  
Sinus Rhomboidalis, [1792](#)  
Sinus Rhythm, [1792](#)  
Sinus Tachycardia, [1792](#)  
Sinus X-Ray, [1792](#)  
Sinuses, [1791](#)  
Sinusitis, [1791](#)  
Sinusitis And Sinus Problems, [1791](#)  
Sinusoid, [1792](#)  
Sinusoidal Fetal Heart Rate Pattern, [1792](#)  
Sinusoidal Obstruction Syndrome, [1792](#)  
Sinusoids, [1792](#)  
Sipuleucel-T, [1792](#)  
Sirenomelia, [1792](#)  
Sirolimus, [1793](#)  
Sirtuins, [1793](#)  
Sisomicin, [1793](#)  
Sister Mary Joseph Nodule, [1793](#)  
Sister Mary Joseph's Nodule, [1793](#)  
Sit-to-Stand Postural Drift, [1793](#)  
Sitagliptin Phosphate, [1793](#)  
Sitosterolemia, [1793](#)  
Sitting Position, [1793](#)  
Situational Syncope, [1793](#)  
Situs Ambiguus, [1793](#)  
Situs Inversus, [1793](#)  
Situs Inversus Totalis, [1793](#)  
Sitz Bath, [1793](#)  
Six Fingers Or Toes, [1794](#)  
Six-Minute Walk Test, [1794](#)  
Sixth Disease, [1794](#)  
Sixth Nerve Palsy, [1794](#)  
Size Perception, [1794](#)  
Sjogren Syndrome, [1794](#)  
Sjogren's Syndrome, [1794](#)  
Sjogren-Larsson Syndrome, [1794](#)  
SJS, [1794](#)  
Sjögren Disease, [1794](#)  
Sjögren Disease-Associated Arthritis, [1794](#)  
SjöGren Syndrome, [1794](#)  
SjöGren's Disease, [1794](#)  
SjöGren's Syndrome, [1794](#)  
Skeletal Development, [1794](#)  
Skeletal Dysplasia, [1795](#)  
Skeletal Limb Abnormalities, [1795](#)  
Skeletal Muscle, [1795](#)  
Skeletal Muscle Relaxant, [2130](#)  
Skeletal Muscle Tissue, [1795](#)  
Skeletal Muscles, [1795](#)  
Skeletal Myocytes, [1795](#)  
Skeletal Survey, [1795](#)  
Skeletal System, [1795](#)  
Skeletal X-Ray, [1795](#)  
Skeleton, [1795](#)  
Skeletonization, [1795](#)

---

---

Skene Gland Cyst, [1795](#)  
Skilled Nursing Facility, [1795](#)  
Skilled Nursing Or Rehabilitation Facilities, [1795](#)  
Skin, [1796](#)  
Skin Abscess, [1796](#)  
Skin Absorption, [1796](#)  
Skin Aging, [1796](#)  
Skin Aging And Appearance, [1796](#)  
Skin Allergies And Contact Dermatitis, [1796](#)  
Skin Allergy Testing, [1796](#)  
Skin And Aging, [1796](#)  
Skin And Environment, [1796](#)  
Skin Appendages, [1796](#)  
Skin Barrier, [1796](#)  
Skin Barrier Function, [1796](#)  
Skin Biopsy, [1796](#)  
Skin Biopsy Methods, [1796](#)  
Skin Biopsy Techniques, [1797](#)  
Skin Blushing/Flushing, [1797](#)  
Skin Cancer, [1797](#)  
Skin Cancer Epidemiology, [1797](#)  
Skin Cancer Screening, [1797](#)  
Skin Cancer Staging, [1797](#)  
Skin Cancer Surgery, [1797](#)  
Skin Cancer, Melanoma, [1224](#)  
Skin Cancer, Squamous Cell Type, [1833](#)  
Skin Carcinoma, [1797](#)  
Skin Care, [1797](#)  
Skin Care And Incontinence, [1797](#)  
Skin Cells, [1797](#)  
Skin Color, [1797](#)  
Skin Culture, [1797](#)  
Skin Cyst, [1798](#)  
Skin Diseases, [1798](#)  
Skin Erosion, [1798](#)  
Skin Examination, [1798](#)  
Skin Findings In Newborns, [1798](#)  
Skin Flaps, [1798](#)  
Skin Graft, [1798](#)  
Skin Immune System, [1798](#)  
Skin Induration, [1798](#)  
Skin Infection, [1798](#)  
Skin Integrity, [1798](#)  
Skin Irritation, [1798](#)  
Skin Lesion, [1798](#)  
Skin Lesion Aspiration, [1798](#)  
Skin Lesion Biopsy, [1799](#)  
Skin Lesion KOH Exam, [1799](#)  
Skin Lesion Of Blastomycosis, [1799](#)  
Skin Lesion Removal, [1799](#)  
Skin Lumps, [1799](#)  
Skin Mastocytoma, [1799](#)  
Skin Microbiome, [1799](#)  
Skin Mycosis, [1799](#)  
Skin Necrosis, [1799](#)  
Skin Neoplasm, [1799](#)  
Skin Nodules, [1799](#)  
Skin Of Color, [1799](#)  
Skin Or Nail Culture, [1799](#)  
Skin Pigmentation, [1799](#)  
Skin Rash, [1881](#)  
Skin Resurfacing, [1799](#)  
Skin Scraping, [1800](#)  
Skin Self-Exam, [1800](#)  
Skin Tag, [1800](#)  
Skin Tags, [1881](#)  
Skin Test, [1800](#)  
Skin Turgor, [1800](#)  
Skin Writing, [1800](#)  
Skin-Grafting, [1798](#)  
Skipped Heartbeats, [1800](#)  
Skull, [1800](#)  
Skull Fracture, [1800](#)  
Skull X-Ray, [1800](#)  
Slanted Ear, [1800](#)  
SLAP Lesion, [1800](#)  
SLAP Tears, [1800](#)  
Slapped Cheek Disease, [1800](#)  
Slapped Cheek Syndrome, [1800](#)  
SLE, [1800](#)  
Sleep, [1800](#)  
Sleep Apnea, [1801](#)  
Sleep Apnoea, [1801](#)  
Sleep Architecture, [1801](#)  
Sleep Bruxism, [1801](#)  
Sleep Debt, [1801](#)  
Sleep Diary, [1801](#)  
Sleep Disorder, [1801](#)  
Sleep Disorders, [1801](#)  
Sleep Disorders in Children, [1801](#)  
Sleep Disorders In Older Adults, [1801](#)  
Sleep Drug, [2130](#)  
Sleep Efficiency, [1801](#)  
Sleep Epilepsy, [1881](#)  
Sleep Fragmentation, [1801](#)  
Sleep Hygiene, [1801](#)  
Sleep Onset Latency, [1802](#)  
Sleep Paralysis, [1802](#)  
Sleep Related Breathing Disorders, [1802](#)  
Sleep Spindles, [1802](#)  
Sleep Studies, [1802](#)  
Sleep-Related Hypoventilation, [1802](#)  
Sleep-Related Hypoxemia, [1802](#)  
Sleeping Sickness, [1801](#)  
Sleeplessness, [1802](#)  
Sleepwalking, [1802](#)  
Sleepwalking And Children, [1802](#)  
Sleepwalking Disorder, [1802](#)

---

---

Sleeve Gastrectomy, [1802](#)  
Sleeve Lobectomy, [1802](#)  
Sliding Filament, [1803](#)  
Sliding Hiatal Hernia, [1803](#)  
Sling, [1803](#)  
Slipped Capital Femoral Epiphysis, [1803](#)  
Slipped Disc, [1881](#)  
Slipped Disk, [1803](#)  
Slipped Vertebra, [1803](#)  
Slipping Rib Syndrome, [1803](#)  
Slit Lamp, [1803](#)  
Slit Lamp Biomicroscopy, [1803](#)  
Slit Lamp Examination, [1803](#)  
Slit-Lamp Biomicroscope, [1803](#)  
Slit-Lamp Exam, [1803](#)  
Slit-Lamp Examination, [1803](#)  
SLL, [1804](#)  
Slough, [1804](#)  
Slow Heart Rate, [1804](#)  
Slow Heartbeat, [1804](#)  
Slow-Twitch Fiber, [1804](#)  
Slow-Wave Sleep, [1804](#)  
Slowed or Arrested Breathing, [1804](#)  
Small Airway Disease, [1804](#)  
Small Bowel Adenocarcinoma, [1804](#)  
Small Bowel Bacterial Overgrowth, [1804](#)  
Small Bowel Bleeding, [1804](#)  
Small Bowel Follow-Through, [1804](#)  
Small Bowel Obstruction, [1805](#)  
Small Bowel Prolapse, [1805](#)  
Small Bowel Resection, [1805](#)  
Small Bowel Series, [1805](#)  
Small Bowel Tissue Smear/Biopsy, [1805](#)  
Small Calorie, [1805](#)  
Small Cell Carcinoma, [1805](#)  
Small Cell Lung Cancer, [1805](#)  
Small Cell Lung Carcinoma, [1805](#)  
Small Eye, [1805](#)  
Small Fiber Neuropathy, [1805](#)  
Small For Gestational Age, [1806](#)  
Small Incision Lenticule Extraction Surgery, [1806](#)  
Small Intestinal Bacterial Overgrowth, [1806](#)  
Small Intestinal Fungal Overgrowth, [1806](#)  
Small Intestinal Ischemia And Infarction, [1806](#)  
Small Intestinal Mucosa, [1806](#)  
Small Intestine, [1806](#)  
Small Intestine Aspirate And Culture, [1806](#)  
Small Intestine Cancer, [1806](#)  
Small Intestine Prolapse, [1806](#)  
Small Lymphocytic Lymphoma, [1881](#)  
Small Round Blue Cell Tumor, [1806](#)  
Small Saphenous Vein, [1806](#)  
Small Vessel Disease, [1807](#)  
Small-Bowel Transplantation, [1805](#)  
Small-Bowel Volvulus, [1805](#)  
Small-Cell Carcinoma, [1805](#)  
Small-Intestinal Cancer, [1806](#)  
Small-Vessel Vasculitis, [1807](#)  
Smallpox, [1806](#)  
SMART, [1807](#)  
Smashed Fingers, [1807](#)  
Smear Of Duodenal Fluid Aspirate, [1807](#)  
Smegma, [1807](#)  
Smell, [1807](#)  
SMILE, [1807](#)  
Smile Design, [1807](#)  
Smith Fracture, [1807](#)  
Smith-Lemli-Opitz Syndrome, [1807](#)  
Smith-Magenis Syndrome, [1807](#)  
Smoke Evacuator, [1807](#)  
Smoke Inhalation, [1808](#)  
Smoke Inhalation Injury, [1808](#)  
Smokeless Tobacco, [1808](#)  
Smoker Keratosis, [1808](#)  
Smoker Lung, [1808](#)  
Smoking, [1808](#)  
Smoking And Asthma, [1808](#)  
Smoking And COPD, [1808](#)  
Smoking and Pregnancy, [1808](#)  
Smoking And Surgery, [1808](#)  
Smoking Cessation, [1808](#)  
Smoking Cessation Medications, [1808](#)  
Smoldering Multiple Myeloma, [1808](#)  
Smooth Muscle, [1808](#)  
Smothering, [1808](#)  
Snake Bite, [1809](#)  
Snake Bites, [1809](#)  
Snake-Bite, [1809](#)  
Snapping Finger, [1809](#)  
Snapping Hip Syndrome, [1809](#)  
Sneddon Syndrome, [1809](#)  
Sneezing, [1809](#)  
Snellen Chart, [1809](#)  
Snoring, [1809](#)  
Snoring And Sleep Apnea, [1809](#)  
Snort, [1809](#)  
Snot, [1809](#)  
Snow Blindness, [1809](#)  
SNRIs, [1809](#)  
Snuff, [1809](#)  
Snuffles, [1810](#)  
Social Anxiety Disorder, [1810](#)  
Social History, [1810](#)  
Social Isolation, [1810](#)  
Social Media Addiction, [1810](#)  
Social Phobia, [1810](#)  
Social Worker, [1810](#)  
Sociophobia, [1810](#)

---

---

Sodium, [1810](#)  
Sodium Bicarbonate Therapy, [1810](#)  
Sodium Bisulfate Poisoning, [1810](#)  
Sodium Blood Test, [1810](#)  
Sodium Carbonate Poisoning, [1810](#)  
Sodium Chloride, [1810](#)  
Sodium Cromoglycate, [1810](#)  
Sodium Diatrizoate, [1810](#)  
Sodium Hydroxide Poisoning, [1811](#)  
Sodium Hypochlorite, [1811](#)  
Sodium Hypochlorite Poisoning, [1811](#)  
Sodium In Diet, [1811](#)  
Sodium Polystyrene Sulfonate, [1811](#)  
Sodium Urine Test, [1811](#)  
Sodium Valproate, [1811](#)  
Sodium-Potassium Pump, [1811](#)  
Soft Markers For Aneuploidy, [1811](#)  
Soft Tissue Augmentation, [1811](#)  
Soft Tissue Injury, [1811](#)  
Soft Tissue Rheumatism, [1811](#)  
Soft Tissue Sarcoma, [1811](#)  
Soft Tissue Tumors, [1812](#)  
Soft-Tissue Calcinosis, [1811](#)  
Soil Contamination, [1812](#)  
Solanine Plant Poisoning, [1812](#)  
Solar Elastosis, [1812](#)  
Solar Keratosis, [1812](#)  
Solar Lentigines, [1812](#)  
Solar Plexus, [1812](#)  
Solar Purpura, [1812](#)  
Solar Urticaria, [1812](#)  
Solder Poisoning, [1812](#)  
Soleus Muscle, [1812](#)  
Solid Pseudopapillary Pancreatic Tumor, [1812](#)  
Solitary Fibrous Tumor, [1812](#)  
Solitary Kidney, [1812](#)  
Solitary Pulmonary Nodule, [1812](#)  
Solution, [1813](#)  
Solvent Abuse, [1813](#)  
Solvent Exposure, [1813](#)  
Somatic, [1813](#)  
Somatic Mutation, [1813](#)  
Somatic Nervous System, [1813](#)  
Somatic Symptom And Related Disorders, [1813](#)  
Somatic Symptom Disorder, [1813](#)  
Somatic Variant, [1813](#)  
Somatization, [1813](#)  
Somatostatin, [1813](#)  
Somatostatin Analog, [1813](#)  
Somatostatinoma, [1813](#)  
Somatotropic Pituitary Adenoma, [1814](#)  
Somatotropin, [1814](#)  
Somatotype, [1814](#)  
Somnambulism, [1814](#)  
Somniloquy, [1814](#)  
Somnolence, [1814](#)  
Somnolent, [1814](#)  
Somogyi Effect, [1814](#)  
Somogyi Phenomenon, [1814](#)  
Sonographer, [1814](#)  
Sonohysterography, [1814](#)  
Soporifics, [1814](#)  
Sorbitol, [1815](#)  
Sore, [1815](#)  
Sore Throat, [1815](#)  
Sotalol, [1815](#)  
Sotos Syndrome, [1815](#)  
Sound, [1815](#)  
Source Control, [1815](#)  
Soy, [1815](#)  
Soy Protein Intolerance, [1815](#)  
Soya, [1815](#)  
Soybeans, [1815](#)  
Space Medicine, [1815](#)  
Spacer, [1815](#)  
Spanish Fly, [1815](#)  
Spasm, [1816](#)  
Spasmodic Dysphonia, [1816](#)  
Spasmodic Torticollis, [1816](#)  
Spasmus Nutans, [1816](#)  
Spastic, [1816](#)  
Spastic Colon, [1816](#)  
Spastic Diplegia, [1816](#)  
Spastic Dysphonia, [1881](#)  
Spastic Gait, [1816](#)  
Spasticity, [1816](#)  
Spatial Resolution, [1816](#)  
Speaking with a Tracheostomy Tube, [1816](#)  
Special Care Baby Unit, [1816](#)  
Special Stain, [1816](#)  
Specific Absorbed Fraction, [1816](#)  
Specific Activity, [1817](#)  
Specific Gravity, [1817](#)  
Specific Learning Disorder, [1817](#)  
Specific Phobia, [1817](#)  
Specific Reading Disability, [1817](#)  
Specificity, [1817](#)  
Specimen Adequacy, [1817](#)  
Specimen Container, [1817](#)  
Specimen Transport Medium, [1817](#)  
Speckled Lentiginous Nevus (Nevus Spilus), [1817](#)  
SPECT Scan, [1817](#)  
Spectral Analysis, [1817](#)  
Speculum, [1817](#)  
Speech, [1818](#)  
Speech And Language Delay, [1818](#)  
Speech Audiometry, [1818](#)  
Speech Delay, [1818](#)

---

---

Speech Discrimination, 1818  
Speech Disorders, 1818  
Speech Impairment In Adults, 1818  
Speech Therapy, 1818  
Speech-Language Pathologist, 1818  
Speech-Language Pathology Assessment, 1818  
Sperm, 1818  
Sperm Cell, 1819  
Sperm Retrieval, 1819  
Spermarche, 1818  
Spermatid, 1818  
Spermatid Cord, 1818  
Spermatocoele, 1819  
Spermatocoele (Epididymal Cyst), 1881  
Spermatogenesis, 1819  
Spermatogonium, 1819  
Spermatorrhea, 1819  
Spermatozoa, 1819  
Spermatozoon, 1819  
Spermicide, 1819  
Sphenoid, 1819  
Sphenoid Bone, 1819  
Sphenoid Sinusitis, 1819  
Sphenoidal Sinus, 1819  
Spherocytes, 1820  
Spherocytic Anemia, 1881  
Sphincter, 1820  
Sphincter Of Oddi, 1820  
Sphingomyelinase Deficiency, 1882  
Sphygmograph, 1820  
Sphygmomanometer, 1820  
Spider Angioma, 1820  
Spider Bite, 1820  
Spider Bites, 1820  
Spider Naevi, 1820  
Spider Naevus, 1820  
Spider Telangiectasia, 1820  
Spider Veins, 1882  
Spielmeyer-Vogt Disease, 1882  
Spielmeyer-Vogt-Batten Syndrome, 1883  
Spigelian Hernia, 1820  
Spina Bifida, 1820  
Spina Bifida Meningocele, 1821  
Spina Bifida Myelomeningocele, 1821  
Spina Bifida Occulta, 1821  
Spinal Analgesia, 1821  
Spinal And Bulbar Muscular Atrophy, 1821  
Spinal And Epidural Anesthesia, 1821  
Spinal Anesthesia, 1821  
Spinal Anesthesia For Cesarean Delivery, 1821  
Spinal Angiography, 1821  
Spinal Arachnoiditis, 1881  
Spinal Artery, 1821  
Spinal Block, 1821  
Spinal Canal, 1821  
Spinal Cavity, 1822  
Spinal Chordoma, 1881  
Spinal Column, 1822  
Spinal Cord, 1822  
Spinal Cord Abscess, 1822  
Spinal Cord and Spine Disorders, 1822  
Spinal Cord Compression, 1822  
Spinal Cord Disease, 1822  
Spinal Cord Infarction, 1822  
Spinal Cord Injuries, 1822  
Spinal Cord Injury, 1822  
Spinal Cord Injury: Treatments And Rehabilitation, 1822  
Spinal Cord Neoplasms, 1822  
Spinal Cord Stimulation, 1822  
Spinal Cord Trauma, 1822  
Spinal Curvature, 1823  
Spinal Disease, 1823  
Spinal Dysraphism, 1823  
Spinal Epidural Abscess, 1823  
Spinal Fluid Leak, 1823  
Spinal Fracture, 1823  
Spinal Fusion, 1823  
Spinal Ganglion, 1823  
Spinal Injury, 1823  
Spinal Meningitis, 1882  
Spinal Metastasis, 1823  
Spinal Muscular Atrophy, 1823  
Spinal Muscular Atrophy (SMA), 1882  
Spinal Neoplasms, 1823  
Spinal Nerve, 1823  
Spinal Precautions, 1823  
Spinal Reflex, 1824  
Spinal Shock, 1824  
Spinal Stenosis, 1824  
Spinal Stenosis Lumbar, 1824  
Spinal Tap, 1824  
Spinal Tumor, 1824  
Spinal Tumors, 1824  
Spindle Pole, 1824  
Spindle Pole Body, 1824  
Spindle-Cell Sarcoma, 1824  
Spine, 1824  
Spine Disorders, 1824  
Spine Injury, 1824  
Spinocerebellar Ataxia, 1824  
Spinocerebellar Ataxias, 1824  
Spinothalamic Tract, 1825  
Spinous Process, 1825  
Spiradenocarcinoma, 1825  
Spiradenoma, 1825  
Spiral Organ, 1825  
Spiramycin, 1825

---



---

Spirillum Minus, [1825](#)  
 Spirit, [1825](#)  
 Spiritual Pain, [1825](#)  
 Spirochaetaceae, [1825](#)  
 Spirochaetales Infection, [1825](#)  
 Spirochaete, [1825](#)  
 Spirochete, [1825](#)  
 Spirogram, [1825](#)  
 Spirometer, [1825](#)  
 Spirometry, [1826](#)  
 Spironolactone, [1826](#)  
 Spirulina, [1826](#)  
 Spit, [1826](#)  
 Spitting, [1826](#)  
 Spitz Nevus, [1826](#)  
 Spitz Nevus Pathology, [1826](#)  
 Splanchnic, [1826](#)  
 Splanchnic Nerve, [1826](#)  
 Spleen, [1826](#)  
 Spleen Development, [1826](#)  
 Spleen Diseases, [1826](#)  
 Spleen Disorder, [1826](#)  
 Spleen Hemangioma, [1827](#)  
 Spleen Hilum, [1827](#)  
 Spleen Removal, [1827](#)  
 Splenectomy, [1827](#)  
 Splenic Abscess, [1827](#)  
 Splenic Artery Aneurysm, [1827](#)  
 Splenic Cyst, [1827](#)  
 Splenic Flexure, [1827](#)  
 Splenic Flexure Syndrome, [1827](#)  
 Splenic Infarction, [1827](#)  
 Splenic Infection, [1827](#)  
 Splenic Neoplasm, [1827](#)  
 Splenic Rupture, [1827](#)  
 Splenomegaly, [1827](#)  
 Splenomegaly With Rheumatoid Arthritis, [1882](#)  
 Splice-Site Variant, [1828](#)  
 Spliceosome, [1828](#)  
 Splint, [1828](#)  
 Splinter, [1828](#)  
 Splinter Haemorrhages, [1828](#)  
 Splinter Hemorrhage, [1828](#)  
 Splinter Hemorrhages, [1828](#)  
 Splinter Removal, [1828](#)  
 Splints, [1828](#)  
 Split Brain, [1828](#)  
 Split-Thickness Skin Graft, [1828](#)  
 Split-Thickness Skin Grafts, [1828](#)  
 Splitting, [1828](#)  
 Spondee, [1828](#)  
 Spondylarthritis, [1829](#)  
 Spondylitis, [1829](#)  
 Spondyloarthritis, [1829](#)  
 Spondylolisthesis, [1829](#)  
 Spondylolysis, [1829](#)  
 Spondylosis, [1829](#)  
 Sponge, [1829](#)  
 Spongiform Encephalopathy, [1829](#)  
 Spongy Bone, [1829](#)  
 Spontaneous Abortion, [1829](#)  
 Spontaneous Bacterial Peritonitis, [1829](#)  
 Spontaneous Bacterial Peritonitis Diagnostic Criteria, [1829](#)  
 Spontaneous Breathing Trial, [1830](#)  
 Spontaneous Coronary Artery Dissection, [1830](#)  
 Spontaneous Miscarriage, [1830](#)  
 Spontaneous Perforation, [1830](#)  
 Spontaneous Pneumothorax, [1830](#)  
 Spontaneous Vaginal Delivery, [1830](#)  
 Sporadic, [1830](#)  
 Sporothrix, [1830](#)  
 Sporothrix Schenckii, [1830](#)  
 Sporotrichosis, [1830](#)  
 Sporozoa, [1830](#)  
 Sporozoites, [1830](#)  
 Sports Cream Overdose, [1830](#)  
 Sports Medicine, [1831](#)  
 Sports Physical, [1831](#)  
 Sports Rehabilitation, [1831](#)  
 Sportsman's Groin, [1830](#)  
 Spots Before The Eyes, [1831](#)  
 Spots In Front Of The Eyes, [1831](#)  
 Spotted Fever, [1831](#)  
 Spousal Abuse, [1831](#)  
 Spouse, [1831](#)  
 Sprain, [1831](#)  
 Sprained Ankle, [1831](#)  
 Sprains, [1831](#)  
 Spray Drying, [1831](#)  
 Sprengel Deformity, [1831](#)  
 Spring Fever Henoch-Schonlein, [1831](#)  
 Sprue, [1832](#)  
 Spur Cell, [1832](#)  
 Spur-Cell Hemolytic Anemia, [1832](#)  
 Spurling Test, [1832](#)  
 Sputum, [1832](#)  
 Sputum Culture, [1832](#)  
 Sputum Direct Fluorescent Antibody Test, [1832](#)  
 Sputum Fungal Smear, [1832](#)  
 Sputum Gram Stain, [1832](#)  
 Sputum Stain For Mycobacteria, [1832](#)  
 Squamous Cell, [1832](#)  
 Squamous Cell Carcinoma, [1832](#)  
 Squamous Cell Carcinoma in Situ, [1832](#)  
 Squamous Cell Carcinoma Of The Lung, [1832](#)  
 Squamous Cell Carcinoma Of The Skin, [1833](#)  
 Squamous Cell Skin Cancer, [1833](#)

---

---

Squamous Cells, [1833](#)  
 Square Foot, [1833](#)  
 Square Inch, [1833](#)  
 Squint, [1833](#)  
 SRT, [1833](#)  
 SSRI, [1833](#)  
 SSRIs, [1833](#)  
 St John's Wort, [1844](#)  
 ST Segment, [1851](#)  
 St Vitus's Dance, [1852](#)  
 ST-Elevation Myocardial Infarction, [1838](#)  
 ST-Segment Elevation, [1851](#)  
 Stab Wound, [1834](#)  
 Stable Angina, [1833](#)  
 Stable Coronary Artery Disease, [1833](#)  
 Stable Disease, [1833](#)  
 Stabs, [1833](#)  
 Stafne Bone Defect, [1834](#)  
 Stage 4 Prostate Cancer, [1834](#)  
 Stage I, [1834](#)  
 Stage II, [1834](#)  
 Stage III, [1834](#)  
 Stage IV, [1834](#)  
 Stages Of Change, [1834](#)  
 Stages of Labor, [1834](#)  
 Staghorn Calculi, [1834](#)  
 Staghorn Calculus, [1834](#)  
 Staging, [1834](#)  
 Staining, [1834](#)  
 Stainless Steel, [1834](#)  
 Stammering, [1835](#)  
 Standard Deviation, [1835](#)  
 Standard Eye Exam, [1835](#)  
 Standard Of Care, [1835](#)  
 Standard Precautions, [1835](#)  
 Standardized Extract, [1835](#)  
 Standardized Uptake Value, [1835](#)  
 Stannosis, [1835](#)  
 Stanazolol, [1835](#)  
 Stapedectomy, [1835](#)  
 Stapedius, [1835](#)  
 Stapes, [1835](#)  
 Staph Infection, [1835](#)  
 Staph Infections In The Hospital, [1835](#)  
 Staphylococcal Food Poisoning, [1835](#)  
 Staphylococcal Infection, [1836](#)  
 Staphylococcal Meningitis, [1836](#)  
 Staphylococcal Pneumonia, [1836](#)  
 Staphylococcal Scalded Skin, [1836](#)  
 Staphylococcal Scalded Skin Syndrome, [1836](#)  
 Staphylococcus, [1836](#)  
 Staphylococcus Aureus, [1836](#)  
 Stapling, [1836](#)  
 Starch Poisoning, [1836](#)  
 Stargardt Disease, [1836](#)  
 Starling Equation, [1836](#)  
 Starling Forces, [1836](#)  
 Starvation, [1836](#)  
 Stasis, [1837](#)  
 Stasis Dermatitis, [1837](#)  
 Stasis Dermatitis And Ulcers, [1837](#)  
 Stasis Ulcer, [1837](#)  
 STAT, [1837](#)  
 Statin, [1837](#)  
 Statin Drugs, [1837](#)  
 Statin-Induced Myopathy, [1837](#)  
 Statins, [1837](#)  
 Statins In Diabetes, [1837](#)  
 Station, [1837](#)  
 Status Asthmaticus, [1837](#)  
 Status Epilepticus, [1837](#)  
 Status Migrainosus, [1838](#)  
 Stay Away From Asthma Triggers, [1838](#)  
 STDs, [1838](#)  
 Steady State, [1838](#)  
 Steal Syndrome, [1838](#)  
 Steatocystoma Multiplex, [1838](#)  
 Steatoma, [1838](#)  
 Steatorrhea, [1838](#)  
 Steatorrhoea, [1838](#)  
 Steatotic Liver Disease (SLD), [1838](#)  
 Steele-Richardson-Olszewski Syndrome, [1838](#)  
 Stein-Leventhal Syndrome, [1838](#)  
 Stem Cell, [1838](#)  
 Stem Cell Donation, [1838](#)  
 Stem Cell Transplantation, [1839](#)  
 Stem Cells, [1839](#)  
 STEMI, [1839](#)  
 Stenosing Tenosynovitis, [1839](#)  
 Stenosis, [1839](#)  
 Stenosis of the LAD, [1839](#)  
 Stenotic, [1839](#)  
 Stenotrophomonas Maltophilia Resistance, [1839](#)  
 Stensen Duct, [1839](#)  
 Stent, [1839](#)  
 Stent Graft, [1839](#)  
 Stent Placement, [1839](#)  
 Stent Restenosis, [1839](#)  
 Stenting, [1839](#)  
 Stepping Reflex, [1839](#)  
 Stereognosis, [1840](#)  
 Stereotactic, [1840](#)  
 Stereotactic Biopsy, [1840](#)  
 Stereotactic Body Radiation Therapy, [1840](#)  
 Stereotactic Brain Biopsy, [1840](#)  
 Stereotactic Breast Biopsy, [1840](#)  
 Stereotactic Radiosurgery, [1840](#)  
 Stereotaxis, [1840](#)

---

---

Stereotypic Movement Disorder, [1840](#)  
Sterile Field, [1840](#)  
Sterile Site, [1840](#)  
Sterile Technique, [1841](#)  
Sterilisation, [1841](#)  
Sterility, [1841](#)  
Sterilization, [1841](#)  
Sternal Angle, [1841](#)  
Sternal Exploration Or Closure, [1841](#)  
Sternal Fracture, [1841](#)  
Sternal Line, [1841](#)  
Sternoclavicular Joint, [1841](#)  
Sternocleidomastoid Muscle, [1841](#)  
Sternocostal Joint, [1841](#)  
Sternotomy, [1841](#)  
Sternum, [1841](#)  
Sternutation, [1841](#)  
Steroid, [2130](#)  
Steroid Abuse, [2130](#)  
Steroid Hormones, [2130](#)  
Steroid Injections - Tendon, Bursa, Joint, [2130](#)  
Steroid Myopathy, [1841](#)  
Steroids, [1842](#)  
Sertoli, [1842](#)  
Stethoscope, [1842](#)  
Stevens Johnson Syndrome, [1882](#)  
Stevens-Johnson Syndrome, [1842](#)  
Stewart-Treves Syndrome, [1842](#)  
STI, [1842](#)  
Stickler Syndrome, [1842](#)  
Stiff Baby Syndrome, [1842](#)  
Stiff-Person Syndrome, [1842](#)  
Stiffness, [1842](#)  
Stigma, [1842](#)  
Stilboestrol, [1842](#)  
Stilet, [1843](#)  
Still Disease, [1843](#)  
Still Disease-Associated Arthritis, [1843](#)  
Still's Disease, [1843](#)  
Still's Murmur, [1843](#)  
Stillbirth, [1843](#)  
Stimulant, [1843](#)  
Stimulant Use Disorder, [1843](#)  
Stimulants, [1843](#)  
Stimulus, [1843](#)  
Stingray, [1843](#)  
Stings, [1843](#)  
Stippling, [1844](#)  
Stitch, [1844](#)  
Stochastic Effects, [1844](#)  
Stoddard Solvent Poisoning, [1844](#)  
Stokes-Adams Syndrome, [1844](#)  
Stoma, [1844](#)  
Stoma Complications, [1845](#)  
Stomach, [1844](#)  
Stomach Cancer, [1844](#)  
Stomach Cramps, [1844](#)  
Stomach Diseases, [1844](#)  
Stomach Flu, [1844](#)  
Stomach Pain, [1844](#)  
Stomach Paralysis, [1844](#)  
Stomach Tube, [1845](#)  
Stomach Ulcer, [1845](#)  
Stomach Washout, [1845](#)  
Stomatitis, [1845](#)  
Stomatodynia, [1845](#)  
Stone, [1845](#)  
Stonefish Sting, [1845](#)  
Stool, [1845](#)  
Stool C Difficile Toxin, [1845](#)  
Stool Examination, [1845](#)  
Stool Gram Stain, [1845](#)  
Stool Guaiac Test, [1845](#)  
Stool Holding, [1845](#)  
Stool Occult Blood Test, [1845](#)  
Stool Ova And Parasites Exam, [1845](#)  
Stools, [1846](#)  
Stools – Floating, [1846](#)  
STOPP/START, [1846](#)  
Storage Diseases, [1846](#)  
Stork Bite, [1846](#)  
Strabismus, [1846](#)  
Straddle Injury, [1846](#)  
Straight Leg Raise Test, [1846](#)  
Strain, [1846](#)  
Strain Imaging, [1846](#)  
Strains, [1846](#)  
Strangulated Hernia, [1846](#)  
Strangulation, [1847](#)  
Strangury, [1846](#)  
Strapping, [1847](#)  
Stratum Corneum, [1847](#)  
Strawberry Hemangioma, [1847](#)  
Strawberry Tongue, [1847](#)  
Strength, [1847](#)  
Strength Testing, [1847](#)  
Strength Training, [1847](#)  
Strep Throat, [1847](#)  
Streptococcal Cellulitis With Lymphangitis, [1847](#)  
Streptococcal Infection, [1847](#)  
Streptococcal Infections, [1847](#)  
Streptococcal Pharyngitis, [1847](#)  
Streptococcal Screen, [1847](#)  
Streptococcal Toxic Shock Syndrome, [1847](#)  
Streptococcus, [1848](#)  
Streptococcus Agalactiae, [1848](#)  
Streptococcus pneumoniae, [1848](#)  
Streptococcus Pyogenes, [1848](#)

---

---

Streptokinase, [1848](#)  
 Streptomycin, [1848](#)  
 Stress, [1848](#)  
 Stress Cardiomyopathy, [1848](#)  
 Stress Echocardiography, [1848](#)  
 Stress Fracture, [1848](#)  
 Stress Fractures, [1848](#)  
 Stress Incontinence, [1848](#)  
 Stress Injuries, [1848](#)  
 Stress Related, [1848](#)  
 Stress Response, [1849](#)  
 Stress Shielding, [1849](#)  
 Stress Tests, [1849](#)  
 Stress Ulcer Prophylaxis, [1849](#)  
 Stress Urinary Incontinence, [1849](#)  
 Stressors, [1848](#)  
 Stretch Mark, [1849](#)  
 Stretch Marks, [1849](#)  
 Stretch Reflex, [1849](#)  
 Stria, [1849](#)  
 Striae, [1849](#)  
 Striae Atrophicae, [1849](#)  
 Striae Distensae (Stretch Marks), [1849](#)  
 Striae Gravidarum, [1849](#)  
 Stricture, [1850](#)  
 Stridor, [1850](#)  
 String Sign, [1850](#)  
 String Test For Intestinal Parasites, [1850](#)  
 Stripping of Membranes, [1850](#)  
 Stroke, [1850](#)  
 Stroke Index, [1850](#)  
 Stroke Rehabilitation, [1850](#)  
 Stroke Risk Factors, [1850](#)  
 Stroke Volume, [1850](#)  
 Stroma, [1850](#)  
 Strongyloides Stercoralis, [1850](#)  
 Strongyloidiasis, [1851](#)  
 Structured Reporting, [1851](#)  
 Struma Ovarii, [1851](#)  
 Struvite Stones, [1851](#)  
 Strychnine, [1851](#)  
 Stucco Keratosis, [1851](#)  
 Student's Elbow, [1851](#)  
 Stuffy Nose, [1851](#)  
 Stuffy Or Runny Nose – Adult, [1851](#)  
 Stump Pain, [1851](#)  
 Stunted Growth, [1851](#)  
 Stupor, [1851](#)  
 Sturge-Weber Syndrome, [1852](#)  
 Stutter, [1852](#)  
 Stuttering, [1852](#)  
 Sty, [1852](#)  
 Stye, [1852](#)  
 Styptics, [1852](#)  
 Subacute, [1852](#)  
 Subacute Bacterial Endocarditis, [1852](#)  
 Subacute Combined Degeneration, [1852](#)  
 Subacute Combined Degeneration Of The Cord, [1852](#)  
 Subacute Cutaneous Lupus Erythematosus (SCLE), [1852](#)  
 Subacute Rehabilitation, [1852](#)  
 Subacute Sclerosing Panencephalitis, [1852](#)  
 Subacute Thyroiditis, [1853](#)  
 Subaortic Stenosis, [1853](#)  
 Subarachnoid, [1853](#)  
 Subarachnoid Cisterns, [1853](#)  
 Subarachnoid Haemorrhage, [1853](#)  
 Subarachnoid Hemorrhage, [1853](#)  
 Subarachnoid Space, [1853](#)  
 Subareolar Abscess, [1853](#)  
 Subchorionic Hemorrhage, [1853](#)  
 Subclavian, [1853](#)  
 Subclavian Artery, [1853](#)  
 Subclavian Steal Syndrome, [1853](#)  
 Subclavian Vein, [1853](#)  
 Subclavian Vein Thrombosis, [1853](#)  
 Subclinical, [1854](#)  
 Subclinical Disease, [1854](#)  
 Subclinical Hypothyroidism, [1854](#)  
 Subclinical Infection, [1854](#)  
 Subconjunctival Hemorrhage, [1854](#)  
 Subconscious, [1854](#)  
 Subcorneal Pustular Dermatitis, [1854](#)  
 Subcortical Dementia, [1882](#)  
 Subcortical Ischemic Vascular Disease, [1882](#)  
 Subcu, [1854](#)  
 Subculture, [1854](#)  
 Subcutaneous, [1854](#)  
 Subcutaneous Emphysema, [1854](#)  
 Subcutaneous Implantable Cardioverter-Defibrillator, [1855](#)  
 Subcutaneous Injections, [1855](#)  
 Subcutaneous Layer, [1855](#)  
 Subcutaneous Tissue, [1855](#)  
 Subdural, [1855](#)  
 Subdural Effusion, [1855](#)  
 Subdural Haematoma, [1855](#)  
 Subdural Hematoma, [1855](#)  
 Subdural Hemorrhage, [1855](#)  
 Subendocardial Myocytes, [1855](#)  
 Subendocardial Sclerosis, [1882](#)  
 Subependymal Giant Cell Astrocytoma, [1855](#)  
 Subependymoma, [1855](#)  
 Subfertility, [1855](#)  
 Subgaleal Hematoma, [1855](#)  
 Subgaleal Hemorrhage, [1856](#)  
 Subinvolution, [1856](#)  
 Subjacent, [1856](#)  
 Subjective, [1856](#)

---

---

Sublimation, [1856](#)  
Sublingual, [1856](#)  
Sublingual Epidermoid Cyst, [1856](#)  
Sublingual Gland, [1856](#)  
Subluxation, [1856](#)  
Submandibular Gland, [1856](#)  
Submucosa, [1856](#)  
Subphrenic Abscess, [1856](#)  
Subq, [1856](#)  
Subscapular Artery, [1856](#)  
Substance, [1857](#)  
Substance Abuse, [1857](#)  
Substance Abuse Disorder, [1857](#)  
Substance Abuse Psychiatry, [1857](#)  
Substance Dependence, [1857](#)  
Substance Intoxication, [1857](#)  
Substance Use, [1857](#)  
Substance Use Disorder, [1857](#)  
Substance Use In Pregnancy, [1857](#)  
Substance Withdrawal, [1857](#)  
Substantia Nigra, [1857](#)  
Substernal Goiter, [1857](#)  
Substrate, [1857](#)  
Subtalar Joint, [1858](#)  
Subungual Exostosis, [1858](#)  
Succenturiate Lobe, [1858](#)  
Succenturiate Placenta, [1858](#)  
Succinate Dehydrogenase, [1858](#)  
Succinylcholine, [1858](#)  
Succumb, [1858](#)  
Succussion Splash, [1858](#)  
Sucking Blister, [1858](#)  
Sucking Reflex, [1858](#)  
Sucrose, [1858](#)  
Sucrose Intolerance, [1882](#)  
Sucrose Lysis Test, [1858](#)  
Suction, [1858](#)  
Suction Device, [1858](#)  
Suction Tip, [1859](#)  
Sudden Cardiac Death, [1859](#)  
Sudden Infant Death Syndrome, [1859](#)  
Sudden Infant Death Syndrome (SIDS), [1859](#)  
Sudden Natural Death Pathology, [1859](#)  
Sudden Sensorineural Hearing Loss, [1859](#)  
Sudden Unexpected Death in Epilepsy, [1859](#)  
Sudden Unexpected Death In Infancy, [1859](#)  
Sudden Unexpected Infant Death, [1859](#)  
Sudden Visual Loss, [1859](#)  
Sudeck Atrophy, [1859](#)  
Sudek's Atrophy, [1859](#)  
Sudorifics, [1859](#)  
Suffocation, [1860](#)  
Sugammadex, [1860](#)  
Sugar Substitutes, [1860](#)  
Sugar-Water Hemolysis Test, [1860](#)  
Suicidal Ideation, [1860](#)  
Suicidal Intent, [1860](#)  
Suicidal Plan, [1860](#)  
Suicidality, [1860](#)  
Suicide, [1860](#)  
Suicide And Suicidal Behavior, [1860](#)  
Suicide Attempt, [1860](#)  
Suicide Risk Assessment, [1860](#)  
Sulcus, [1860](#)  
Sulfa Drugs, [1861](#)  
Sulfamethoxazole, [1861](#)  
Sulfasalazine, [1861](#)  
Sulfatide, [1861](#)  
Sulfite Oxidase Deficiency, [1861](#)  
Sulfite Sensitivity, [1861](#)  
Sulfonamide, [1861](#)  
Sulfonamides, [1861](#)  
Sulfonylurea, [1861](#)  
Sulfonylureas, [1861](#)  
Sulfur Colloid Scan, [1861](#)  
Sulfuric Acid Poisoning, [1861](#)  
Sulindac Overdose, [1861](#)  
Sulphonamides, [1861](#)  
Sulphonylureas, [1862](#)  
Sumatriptan, [1862](#)  
Summer Diarrhoea, [1862](#)  
Sun Protection, [1862](#)  
Sunburn, [1862](#)  
Sundowning, [1862](#)  
Sunflower Syndrome, [1862](#)  
Sunken Chest, [1862](#)  
Sunken Fontanelle, [1862](#)  
Sunscreens, [1862](#)  
Sunstroke, [1862](#)  
Superantigen, [1862](#)  
Superficial, [1862](#)  
Superficial Fascia, [1862](#)  
Superficial Spreading Melanoma, [1863](#)  
Superficial Thrombophlebitis, [1863](#)  
Superimposed Preeclampsia, [1863](#)  
Superinfection, [1863](#)  
Superior, [1863](#)  
Superior Mediastinum, [1863](#)  
Superior Mesenteric Artery, [1863](#)  
Superior Mesenteric Artery Syndrome, [1863](#)  
Superior Semicircular Canal Dehiscence, [1863](#)  
Superior Vena Cava, [1863](#)  
Superior Vena Cava Syndrome, [1863](#)  
Superior Vena Caval Obstruction, [1863](#)  
Supernumerary Digit, [1863](#)  
Supernumerary Nipple, [1864](#)  
Supernumerary Nipples, [1864](#)  
Supernumerary Rib, [1864](#)

---

---

Supernumerary Teeth, [1864](#)  
Supervised Exercise Therapy, [1864](#)  
Supination, [1864](#)  
Supine, [1864](#)  
Supine Hypotensive Syndrome, [1864](#)  
Supine Position, [1864](#)  
Supplemental Oxygen Therapy, [1864](#)  
Supplementation, [1864](#)  
Supplements, [1864](#)  
Supportive Care, [1864](#)  
Suppository, [1864](#)  
Suppression, [1865](#)  
Suppuration, [1865](#)  
Suppurative Arthritis, [1882](#)  
Supracervical Hysterectomy, [1865](#)  
Supracondylar Humerus Fracture, [1865](#)  
Supraglottic Airway Device, [1865](#)  
Supranuclear Ophthalmoplegia, [1865](#)  
Supranuclear Palsy, [1865](#)  
Suprapubic, [1865](#)  
Suprapubic Catheter, [1865](#)  
Suprapubic Catheter Care, [1865](#)  
Suprarenal Glands, [1865](#)  
Suprascapular Nerve, [1865](#)  
Suprascapular Neuropathy, [1865](#)  
Supraspinatus Muscle, [1865](#)  
Suprasternal Notch, [1866](#)  
Supraventricular Tachycardia, [1866](#)  
Sural Nerve, [1866](#)  
Suramin, [1866](#)  
Surfactant, [1866](#)  
Surgeon, [1866](#)  
Surgery, [1866](#)  
Surgery For Pancreatic Cancer, [1866](#)  
Surgery For Pilonidal Cyst, [1866](#)  
Surgical Abortion, [1866](#)  
Surgical Drain, [1866](#)  
Surgical Drill, [1866](#)  
Surgical Excision, [1866](#)  
Surgical Fixation, [1866](#)  
Surgical Forceps, [1867](#)  
Surgical Hemostasis, [1867](#)  
Surgical Hemostat, [1867](#)  
Surgical Instruments, [1867](#)  
Surgical Margin, [1867](#)  
Surgical Repair, [1867](#)  
Surgical Retractor, [1867](#)  
Surgical Risk Stratification, [1867](#)  
Surgical Safety Checklist, [1867](#)  
Surgical Scissors, [1867](#)  
Surgical Site Infection, [1867](#)  
Surgical Stapler, [1867](#)  
Surgical Time-Out, [1867](#)  
Surgical Wound Classification, [1868](#)  
Surgical Wound Infection Treatment, [1868](#)  
Surrogacy, [1868](#)  
Surrogate, [1868](#)  
Surrogate Decision-Maker, [1868](#)  
Surveillance, [1868](#)  
Surveillance Imaging, [1868](#)  
Survival Rate, [1868](#)  
Survivorship, [1868](#)  
Survivorship Care Plan, [1868](#)  
Susac Syndrome, [1868](#)  
Susceptibility, [1868](#)  
Susceptibility Testing, [1868](#)  
Susceptibility-Weighted Imaging, [1868](#)  
Suspected Cause, [1869](#)  
Suspension, [1869](#)  
Sustained Virologic Response, [1869](#)  
Suture, [1869](#)  
Suture Anchor, [1869](#)  
SUV, [1869](#)  
SVC Obstruction, [1869](#)  
SVC Syndrome, [1869](#)  
SVT, [1869](#)  
Swab, [1869](#)  
Swallow Study, [1870](#)  
Swallowing, [1869](#)  
Swallowing Assessment, [1869](#)  
Swallowing Chalk, [1869](#)  
Swallowing Difficulty, [1869](#)  
Swallowing Problems, [1869](#)  
Swallowing Soap, [1870](#)  
Swallowing Sunscreen, [1870](#)  
Swallowing Syncope, [1870](#)  
Swallowing Therapy, [1870](#)  
Swan Neck Deformity, [1870](#)  
Swan-Ganz Catheter, [1870](#)  
Swan-Ganz Catheterization, [1870](#)  
Swan-Neck Deformity, [1870](#)  
Sweat, [1870](#)  
Sweat Chloride Test, [1870](#)  
Sweat Electrolytes Test, [1870](#)  
Sweat Gland, [1870](#)  
Sweat Gland Tumor, [1871](#)  
Sweat Glands, [1871](#)  
Sweating, [1871](#)  
Sweating And Body Odor, [1871](#)  
Sweet Syndrome, [1871](#)  
Sweet's Syndrome, [1871](#)  
Sweetened Beverages, [1871](#)  
Swelling, [1871](#)  
Swimmer's Ear, [1871](#)  
Swimmer's Itch, [1871](#)  
Swimming Pool Granuloma, [1871](#)  
Swine Flu, [1871](#)  
Swollen Ankles And Swollen Feet, [1871](#)

---



---

Swollen Glands, [1872](#)  
 Swollen Gums, [1872](#)  
 Swollen Lymph Glands, [1872](#)  
 Swollen Lymph Nodes, [1872](#)  
 Swollen Lymph Nodes Entry, [1872](#)  
 Swollen Tongue, [1872](#)  
 Swyer Syndrome, [1872](#)  
 Sycosis Barbae, [1872](#)  
 Sydenham Chorea, [1872](#)  
 Sydenham's Chorea, [1872](#)  
 Symblepharon, [1872](#)  
 Symmastia, [1882](#)  
 Sympathectomy, [1873](#)  
 Sympathetic, [1873](#)  
 Sympathetic Ganglion, [1873](#)  
 Sympathetic Nervous System, [1873](#)  
 Sympatholytic, [1873](#)  
 Sympathomimetic Drugs, [1873](#)  
 Sympathomimetic Syndrome, [1873](#)  
 Sympathomimetic Toxicity, [1873](#)  
 Sympathomimetic Toxidrome, [1873](#)  
 Symphysis, [1873](#)  
 Symphysis Pubis, [1873](#)  
 Symphysis-Fundal Height, [1873](#)  
 Symptom, [1873](#)  
 Symptomatic, [1873](#)  
 Symptomatic Porphyria, [1882](#)  
 Symptomatic Treatment, [1873](#)  
 Synaesthesia, [1874](#)  
 Synapse, [1874](#)  
 Synapse Assembly, [1874](#)  
 Synapsins, [1874](#)  
 Synaptic Cleft, [1874](#)  
 Synaptic Membrane, [1874](#)  
 Synaptic Plasticity, [1874](#)  
 Synaptic Transmission, [1874](#)  
 Synaptic Vesicle, [1874](#)  
 Synaptonemal Complex, [1874](#)  
 Synaptosome, [1874](#)  
 Synaptotagmin, [1874](#)  
 Synarthrosis, [1874](#)  
 Sychondroses, [1874](#)  
 Synchronized Cardioversion, [1874](#)  
 Syncope, [1875](#)  
 Syndactyly, [1875](#)  
 Syndecan, [1875](#)  
 Syndesmophyte, [1875](#)  
 Syndesmoses, [1875](#)  
 Syndrome, [1875](#)  
 Syndrome Of Inappropriate ADH Secretion, [1875](#)  
 Syndrome of Inappropriate Antidiuresis (SIAD), [1875](#)  
 Syndrome Of Inappropriate Antidiuretic Hormone, [1875](#)  
 Syndrome Of Inappropriate Antidiuretic Hormone Secretion, [1875](#)  
 Syndrome X, [1875](#)  
 Synechia, [1875](#)  
 Synechiae, [1876](#)  
 Synergist, [1876](#)  
 Synergistic Necrotizing Fasciitis, [1883](#)  
 Synesthesia, [1876](#)  
 Synostoses, [1876](#)  
 Synostosis, [1876](#)  
 Synovectomy, [1876](#)  
 Synovial Biopsy, [1876](#)  
 Synovial Bursa, [1876](#)  
 Synovial Chondromatosis, [1876](#)  
 Synovial Cyst, [1876](#)  
 Synovial Fluid, [1876](#)  
 Synovial Fluid Analysis, [1876](#)  
 Synovial Joint, [1876](#)  
 Synovial Joints, [1876](#)  
 Synovial Membrane, [1876](#)  
 Synovial Reaction To Implanted Prostheses, [1876](#)  
 Synovial Sarcoma, [1877](#)  
 Synoviocytes, [1877](#)  
 Synovitis, [1877](#)  
 Synovium, [1877](#)  
 Synovography, [1877](#)  
 Synthetic Androgen, [1877](#)  
 Synthetic Biology, [1877](#)  
 Syphilis, [1877](#)  
 Syphilis Serology, [1877](#)  
 Syphilis, Congenital, [448](#)  
 Syphilitic Alopecia, [1877](#)  
 Syphilitic Aseptic Meningitis, [1877](#)  
 Syphilitic Gumma, [1877](#)  
 Syringe, [1877](#)  
 Syringe Drivers, [1878](#)  
 Syringocystadenoma Papilliferum, [1878](#)  
 Syringoma, [1878](#)  
 Syringomyelia, [1878](#)  
 Syrinx, [1878](#)  
 Syrup, [1878](#)  
 Systematic Desensitization, [1878](#)  
 Systemic, [1878](#)  
 Systemic Aleukemic Reticuloendotheliosis, [1883](#)  
 Systemic Antifungals, [1878](#)  
 Systemic Chemotherapy, [1878](#)  
 Systemic Circuit, [1878](#)  
 Systemic Dermatoses, [1878](#)  
 Systemic Disease, [1878](#)  
 Systemic Effects, [1878](#)  
 Systemic Exertional Intolerance Disease, [1879](#)  
 Systemic Inflammatory Response Syndrome, [1879](#)  
 Systemic Juvenile Idiopathic Arthritis, [1879](#)  
 Systemic Lupus, [1879](#)  
 Systemic Lupus Erythematosus, [1879](#)

---

---

Systemic Lupus Erythematosus (SLE) and Pregnancy, 1879

Systemic Mastocytosis, 1879

Systemic Scleroderma, 1879

Systemic Sclerosis, 1879

Systemic Sclerosis-Associated Arthritis, 1879

Systemic Therapy, 1879

Systemic Vascular Resistance, 1879

Systemic Vasculitis, 1880

Systemic Venous, 1880

Systole, 1880

Systolic, 1880

Systolic Anterior Motion, 1880

Systolic Murmur, 1880

Systolic Murmurs, 1880

Systolic Pressure, 1880

SéZary Syndrome, 1880

T Cell, 1894

T Wave, 1976

T-Cell, 1894

T-Cell Count, 1894

T-Cell Lymphoma, 1894

T-Cell Lymphomas, 1894

T-Cell Receptor, 1894

T-Cells, 1894

T-Lymphocyte, 1930

T-Wave Alternans, 1976

T3, 1885

T3 Test, 1885

T3RU Test, 1885

T4, 1885

TAB Vaccine, 1885

Tabes Dorsalis, 1885

Tablet, 1885

Tablet Splitting, 1885

TAC Regimen, 1886

Tacalcitol, 1885

Tache Noire, 1885

Tachyarrhythmia, 1885

Tachycardia, 1886

Tachycardia, Supraventricular, 1866

Tachycardia-Bradycardia Syndrome, 1886

Tachycardia-Induced Cardiomyopathy, 1886

Tachykinin, 1886

Tachypnea, 1886

Tachypnoea, 1886

Tachysystole, 1886

TACO, 1886

Tacrolimus, 1886

Tactile, 1887

Tactile Agnosia, 1887

Tactile Fremitus, 1887

Tactile Sense, 1887

Tadalafil, 1887

Taenia, 1887

Taenia Saginata, 1887

Taenia Solium, 1887

Taeniasis, 1887

Tail Of The Epididymis, 1887

Tailbone Trauma, 1887

Takayasu Arteritis, 1887

Takayasu Disease, 1887

Takayasu'S Arteritis, 1887

Takotsubo Cardiomyopathy, 1888

Takotsubo Syndrome, 1888

Talactoferrin Alfa, 1888

Talaporfin Sodium, 1888

Talar Dome Lesion, 1888

Talar Tilt Test, 1888

Talc, 1888

Talcosis, 1888

Talcum Powder Poisoning, 1888

Talipes, 1888

Talipes Calcaneovalgus, 1888

Talipes Equinovarus, 1888

Tall Stature, 1889

Talofibular Ligament Injury, 1889

Talus, 1889

Tamiflu, 1889

Tamoxifen, 1889

Tamoxifen Citrate, 1889

Tamoxifen Toxicity, 1889

Tampon, 1889

Tamponade, 1889

Tamsulosin, 2130

Tamsulosin Hydrochloride, 1889

Tangier Disease, 1889

Tanner Staging, 1889

Tantalum, 1889

Tapentadol, 1889

Tapeworm, 1890

Tapeworm Infection, 1890

Tapotement, 1890

Tapping, 1890

TAPSE, 1890

TAPVR, 1890

Tar Remover Poisoning, 1891

Tar Staining, 1892

TAR Syndrome, 1892

Tarantula Spider Bite, 1890

Tardieu Spots, 1890

Tardive Dyskinesia, 1890

Tardive Dystonia, 1890

Tardive Oral Dyskinesia, 1981

Target Cell, 1890

---

---

Target Cells, [1890](#)  
Target Organ, [1891](#)  
Target Sign, [1891](#)  
Target Tissue, [1891](#)  
Target-Controlled Infusion, [1890](#)  
Targeted Radiation Therapy, [1891](#)  
Targeted Temperature Management, [1891](#)  
Targeted Therapies For Cancer, [1891](#)  
Targeted Therapy, [1891](#)  
Tarlov Cyst, [1891](#)  
Tarlov Cysts, [1891](#)  
Tars, [1891](#)  
Tarsal, [1891](#)  
Tarsal Bones, [1891](#)  
Tarsal Coalition, [1891](#)  
Tarsal Cyst, [1891](#)  
Tarsal Joints, [1892](#)  
Tarsal Tunnel Syndrome, [1892](#)  
Tarsorrhaphy, [1892](#)  
Tarsus, [1892](#)  
Tartar, [1892](#)  
Tarui Disease, [1892](#)  
Tassinari Syndrome, [1892](#)  
Taste, [1892](#)  
Taste Bud, [1892](#)  
Taste Disorder, [1892](#)  
Taste Receptor, [1892](#)  
Tattoos, [1893](#)  
Taurine, [1893](#)  
Taurodontism, [1893](#)  
Taussig-Bing Anomaly, [1893](#)  
TAVR, [1893](#)  
Taxanes, [1893](#)  
Taxotere, [1893](#)  
Tay-Sachs Disease, [1893](#)  
Tazarotene, [1893](#)  
TB, [1893](#)  
TB (Tuberculosis), [1981](#)  
TB Test, [1893](#)  
TBG Blood Test, [1893](#)  
TBI, [1893](#)  
TBM, [1981](#)  
TCA Cycle, [1894](#)  
TCAs, [2130](#)  
TCOF1, [1981](#)  
TCS, [1981](#)  
Td, [1894](#)  
Td Immunization, [1894](#)  
Tdap Vaccination In Pregnancy, [1894](#)  
Tdap Vaccine, [2130](#)  
Tear, [1894](#)  
Tear Film Break-Up Time, [1894](#)  
Teardrop Cells, [1894](#)  
Tearing, [1895](#)  
Tears, [1895](#)  
Technetium-99, [1895](#)  
Technetium-99M, [1895](#)  
Technique, [1895](#)  
Tectum, [1895](#)  
TEE, [1895](#)  
Teen Depression, [1895](#)  
Teenage Pregnancy, [1895](#)  
Teenagers And Drugs, [1895](#)  
Teenagers And Sleep, [1895](#)  
TEER, [1895](#)  
Teeth, [1895](#)  
Teeth Grinding, [1896](#)  
Teeth-Grinding, [1896](#)  
Teething, [1896](#)  
Teichopsia, [1896](#)  
Teicoplanin, [1896](#)  
Telangiectasia, [1896](#)  
Telangiectasis, [1896](#)  
Tele-Rehabilitation, [1896](#)  
Teledermatology, [1896](#)  
Telegraphic Speech, [1896](#)  
Telehealth, [1896](#)  
Telemedicine, [1896](#)  
Teleradiology, [1896](#)  
Telogen and Anagen Effluvium, [1896](#)  
Telogen Effluvium, [1897](#)  
Telomere, [1897](#)  
Temazepam, [1897](#)  
Temperature, [1897](#)  
Temperature Measurement, [1897](#)  
Temple, [1897](#)  
Temple Syndrome, [1897](#)  
Temporal, [1897](#)  
Temporal Arteritis, [1897](#)  
Temporal Arteritis Pathology, [1897](#)  
Temporal Artery, [1897](#)  
Temporal Bones, [1897](#)  
Temporal Lobe, [1897](#)  
Temporal Lobe Epilepsy, [1897](#)  
Temporal Lobe Seizure, [1897](#)  
Temporalis, [1898](#)  
Temporary Blindness, [1982](#)  
Temporary Loss Of Consciousness, [1898](#)  
Temporary Unconsciousness, [1898](#)  
Temporomandibular Disorders, [1898](#)  
Temporomandibular Joint, [1898](#)  
Temporomandibular Joint Disorder, [1898](#)  
Temporomandibular Joint Dysfunction, [1898](#)  
Temporomandibular Joint Replacement, [1898](#)  
Temporomandibular Joint Syndrome, [1898](#)  
TEN, [1898](#)  
Tenascin, [1898](#)  
Tender, [1898](#)

---

---

Tenderness, [1898](#)  
Tendinitis, [1898](#)  
Tendinopathy, [1899](#)  
Tendinosis, [1899](#)  
Tendon, [1899](#)  
Tendon Injury, [1899](#)  
Tendon Repair, [1899](#)  
Tendon Rupture, [1899](#)  
Tendon Sheath, [1899](#)  
Tendon Transfer, [1899](#)  
Tendon Xanthoma, [1899](#)  
Tendonitis, [1899](#)  
Tenecteplase, [1899](#)  
Tenericutes, [1899](#)  
Tenesmus, [1899](#)  
Teniposide, [1900](#)  
Tennis Elbow, [1900](#)  
Tennis Elbow, [1118](#)  
Tennis Elbow Surgery, [1900](#)  
Tenodesis, [1900](#)  
Tenofovir, [1900](#)  
Tenosynovitis, [1900](#)  
Tenotomy, [1900](#)  
Tenovaginitis, [1900](#)  
TENS, [1981](#)  
Tensile Strength, [1900](#)  
Tension, [1900](#)  
Tension Headache, [1900](#)  
Tension Pneumothorax, [1900](#)  
Tension-Type Headache, [1900](#)  
Tentorium, [1900](#)  
Tentorium Cerebelli, [1900](#)  
Teprotumumab, [1901](#)  
Teratocarcinoma, [1901](#)  
Teratogen, [1901](#)  
Teratogenesis, [1901](#)  
Teratogenic, [1901](#)  
Teratogenic Drugs, [1901](#)  
Teratogenicity, [1901](#)  
Teratology and Drug Use During Pregnancy, [1901](#)  
Teratoma, [1901](#)  
Teratozoospermia, [1901](#)  
Terazosin, [2130](#)  
Terazosin Hydrochloride, [1901](#)  
Terbinafine, [1901](#)  
Terbufos, [1901](#)  
Terbutaline, [1902](#)  
Teres Major Muscle, [1902](#)  
Terfenadine, [1902](#)  
Teriparatide For Osteoporosis, [1902](#)  
Terlipressin, [1902](#)  
Term, [1902](#)  
Term Birth, [1902](#)  
Terminal Care, [1902](#)  
Terminal Delirium, [1902](#)  
Terminal Duct Lobular Unit, [1902](#)  
Terminal Ileitis, [1902](#)  
Terminal Illness, [1902](#)  
Terminal Insomnia, [1902](#)  
Terry's Lines, [1902](#)  
Terry's Nails, [1902](#)  
Terson Syndrome, [1903](#)  
Tertian Fever, [1903](#)  
Tertiary Hyperparathyroidism, [1903](#)  
Tertiary Prevention, [1903](#)  
Test Meal, [1905](#)  
Test Strips, [1906](#)  
Testes, [1903](#)  
Testicle, [1903](#)  
Testicle Lump, [1903](#)  
Testicle Pain, [1903](#)  
Testicles, [1903](#)  
Testicular Appendage Torsion, [1903](#)  
Testicular Artery, [1903](#)  
Testicular Atrophy, [1903](#)  
Testicular Biopsy, [1903](#)  
Testicular Cancer, [1904](#)  
Testicular Cancer In Children, [1904](#)  
Testicular Choriocarcinoma, [1904](#)  
Testicular Descent, [1904](#)  
Testicular Diseases, [1904](#)  
Testicular Disorders, [1904](#)  
Testicular Failure, [1904](#)  
Testicular Function, [1904](#)  
Testicular Gonadoblastoma, [1904](#)  
Testicular Hydrocele, [1904](#)  
Testicular Infection, [1904](#)  
Testicular Interstitium, [1904](#)  
Testicular Lobule, [1904](#)  
Testicular Mass, [1904](#)  
Testicular Microlithiasis, [1905](#)  
Testicular Pain, [1905](#)  
Testicular Self-Exam, [1905](#)  
Testicular Seminoma, [1905](#)  
Testicular Torsion, [1905](#)  
Testicular Torsion Emergency, [1905](#)  
Testicular Torsion Overview, [1905](#)  
Testicular Torsion Repair, [1905](#)  
Testicular Trauma, [1905](#)  
Testicular Tumor, [1905](#)  
Testicular Vein, [1905](#)  
Testis, [1905](#)  
Testis Disorder, [1905](#)  
Testis Neoplasm, [1905](#)  
Testosterone, [1906](#)  
Testosterone Deficiency, [1906](#)  
Testosterone Propionate, [1906](#)  
Tests For H Pylori, [1906](#)

---

---

Tet Spell, [1907](#)  
Tetanus, [1906](#)  
Tetanus Toxin, [1906](#)  
Tetany, [1906](#)  
Tethered Cord Syndrome, [1906](#)  
Tetrabenazine, [1906](#)  
Tetracaine, [1906](#)  
Tetrachloroethylene, [1906](#)  
Tetrachlorvinphos, [1906](#)  
Tetracycline, [1907](#)  
Tetracycline Stain, [1907](#)  
Tetracyclines, [1907](#)  
Tetrahydrozoline Poisoning, [1907](#)  
Tetralogy Of Fallot, [1907](#)  
Tetraplegia, [1907](#)  
TGA, [1907](#)  
Th1/Th2/Th17/Tfh/Treg Balance, [1907](#)  
Thalamic Pain Syndrome, [1907](#)  
Thalamus, [1907](#)  
Thalassaemia, [1907](#)  
Thalassemia, [1907](#)  
Thalassemia Major, [1908](#)  
Thalidomide, [1908](#)  
Thallium, [1908](#)  
Thallium Poisoning, [1908](#)  
Thallium Scan, [1908](#)  
Thallium Toxicity, [1908](#)  
Thallium-201, [1908](#)  
Thanatophoric Dysplasia, [1908](#)  
Theca, [1908](#)  
Theca Lutein Cyst, [1908](#)  
Thecoma, [1908](#)  
Thelarche, [1908](#)  
Thenar Eminence, [1908](#)  
Theophylline, [1908](#)  
Theophylline Toxicity, [1909](#)  
Theranostic Pair, [1909](#)  
Theranostics, [1909](#)  
Therapeutic Abortion, [1909](#)  
Therapeutic Alliance, [1909](#)  
Therapeutic Drug Levels, [2130](#)  
Therapeutic Drug Monitoring, [1909](#)  
Therapeutic Equivalence, [1909](#)  
Therapeutic Exercise, [1909](#)  
Therapeutic Hypothermia, [1909](#)  
Therapeutic Index, [1909](#)  
Therapeutic Window, [1909](#)  
Therapeutics, [1909](#)  
Therapy, [1909](#)  
Thermal Injury, [1910](#)  
Thermal Therapy, [1910](#)  
Thermogenesis, [1910](#)  
Thermography, [1910](#)  
Thermoluminescent Dosimeter, [1910](#)  
Thermometer Scales, [1910](#)  
Thermoneutral Zone, [1910](#)  
Thermoplasty, [1910](#)  
Thermoreceptor, [1910](#)  
Thermoregulation, [1910](#)  
Thiabendazole, [1910](#)  
Thiamin, [1910](#)  
Thiamine, [1910](#)  
Thiamine Deficiency, [1911](#)  
Thiazide Diuretics, [1911](#)  
Thiazide Overdose, [1911](#)  
Thiazides, [1911](#)  
Thiazolidinedione Drugs, [1911](#)  
Thiazolidinediones, [1911](#)  
Thigh, [1911](#)  
Thimerosal, [1911](#)  
ThinPrep Pap Test, [1911](#)  
Thiopental Sodium, [1911](#)  
Thiopentone Sodium, [1911](#)  
Thioridazine, [1911](#)  
Thioridazine Overdose, [1911](#)  
Thiotepa, [1911](#)  
Third Heart Sound, [1912](#)  
Third Stage Of Labor, [1912](#)  
Third Trimester, [1912](#)  
Third-Degree Atrioventricular Block, [1912](#)  
Third-Degree AV Block, [1912](#)  
Third-Degree Burn, [1912](#)  
Third-Degree Perineal Tear, [1912](#)  
Thirst, [1912](#)  
Thomas Test, [1912](#)  
Thompson Test, [1912](#)  
Thomsen Disease, [1912](#)  
Thoracentesis, [1913](#)  
Thoracic, [1913](#)  
Thoracic Aortic Aneurysm, [1913](#)  
Thoracic Cage, [1913](#)  
Thoracic Cavity, [1913](#)  
Thoracic Duct, [1913](#)  
Thoracic Endometriosis, [1913](#)  
Thoracic Inlet, [1913](#)  
Thoracic Outlet, [1913](#)  
Thoracic Outlet Syndrome, [1913](#)  
Thoracic Skeleton, [1914](#)  
Thoracic Spine, [1914](#)  
Thoracic Spine CT Scan, [1914](#)  
Thoracic Spine X-Ray, [1914](#)  
Thoracic Surgeon, [1914](#)  
Thoracic Vertebrae, [1914](#)  
Thoracolumbar Spine Trauma, [1914](#)  
Thoracoscopy, [1914](#)  
Thoracotomy, [1914](#)  
Thorax, [1914](#)  
Thought Blocking, [1914](#)

---

---

Thought Disorders, [1914](#)  
Threadworm, [1914](#)  
Threadworm Infection, [1914](#)  
Threadworms, [1914](#)  
Threatened Miscarriage, [1915](#)  
Three Swab Test, [1915](#)  
Three-Day Measles, [1915](#)  
Three-Dimensional Ultrasound, [1915](#)  
Threonine, [1915](#)  
Threshold, [1915](#)  
Thrill, [1915](#)  
Throat, [1915](#)  
Throat Cancer, [1915](#)  
Throat Or Larynx Cancer, [1915](#)  
Throat Swab Culture, [1915](#)  
Throbbing, [1915](#)  
Throckmorton Sign, [1915](#)  
Thrombectomy, [1915](#)  
Thrombin, [1916](#)  
Thrombin Time, [1916](#)  
Thromboangiitis Obliterans, [1916](#)  
Thrombocyte, [1916](#)  
Thrombocytes, [1916](#)  
Thrombocythemia, [1916](#)  
Thrombocytopathy, [1916](#)  
Thrombocytopenia, [1916](#)  
Thrombocytopenia in Pregnancy, [1916](#)  
Thrombocytopenic, [1916](#)  
Thrombocytosis, [1916](#)  
Thromboelastography, [1917](#)  
Thromboembolism, [1917](#)  
Thromboembolism in Pregnancy, [1917](#)  
Thrombolysis, [1917](#)  
Thrombolysis For PE, [1917](#)  
Thrombolysis For Stroke, [1917](#)  
Thrombolytic, [2130](#)  
Thrombolytic Agents, [1917](#)  
Thrombolytic Drugs For Heart Attack, [2130](#)  
Thrombolytic Therapy, [1917](#)  
Thrombophilia, [1917](#)  
Thrombophilia Testing, [1917](#)  
Thrombophilias in Pregnancy, [1917](#)  
Thrombophlebitis, [1917](#)  
Thrombopoiesis, [1917](#)  
Thrombosed Intracranial Aneurysm, [1917](#)  
Thrombosis, [1918](#)  
Thrombosis Of The Inferior Vena Cava, [1918](#)  
Thrombotic Microangiopathy, [1918](#)  
Thrombotic Thrombocytopenic Purpura, [1918](#)  
Thrombotic Thrombocytopenic Purpura Pentad, [1918](#)  
Thrombotic Vasculopathy, [1918](#)  
Thromboxane, [1918](#)  
Thrombus, [1918](#)  
Thrombus Organization, [1918](#)  
Thrush, [1918](#)  
Thrush in Children, [1918](#)  
Thrush In Newborns, [1919](#)  
Thumb Arthritis, [1919](#)  
Thumb Carpometacarpal Arthritis, [1919](#)  
Thumb CMC Arthritis, [1919](#)  
Thumb-Sucking, [1919](#)  
Thymic Carcinoma, [1919](#)  
Thymic Hyperplasia, [1919](#)  
Thymic Neoplasm, [1919](#)  
Thymine, [1919](#)  
Thymocyte, [1919](#)  
Thymoma, [1919](#)  
Thymus, [1919](#)  
Thymus Gland, [1919](#)  
Thyrocalcitonin, [1919](#)  
Thyroglobulin Surveillance, [1919](#)  
Thyroglossal Duct Cyst, [1920](#)  
Thyroid, [1920](#)  
Thyroid Cancer, [1920](#)  
Thyroid Cancer Staging, [1920](#)  
Thyroid Cartilage, [1920](#)  
Thyroid Diseases, [1920](#)  
Thyroid Disorders, [1920](#)  
Thyroid Drugs, [1921](#)  
Thyroid Eye Disease, [1920](#)  
Thyroid Fine-Needle Aspiration, [1920](#)  
Thyroid Follicular Carcinoma, [1920](#)  
Thyroid Function Tests, [1920](#)  
Thyroid Gland, [1920](#)  
Thyroid Gland Enlargement, [1920](#)  
Thyroid Gland Removal, [1920](#)  
Thyroid Hormone, [1921](#)  
Thyroid Hormone Resistance, [1921](#)  
Thyroid Hormone Toxicity, [1921](#)  
Thyroid Hormones, [1921](#)  
Thyroid Lymphoma, [1921](#)  
Thyroid Nodule, [1921](#)  
Thyroid Nodule Evaluation, [1921](#)  
Thyroid Nodules, [1921](#)  
Thyroid Papillary Carcinoma, [1921](#)  
Thyroid Peroxidase Antibody, [1922](#)  
Thyroid Peroxidase Test, [1922](#)  
Thyroid Preparation Overdose, [1922](#)  
Thyroid Scan, [1922](#)  
Thyroid Scintigraphy, [1922](#)  
Thyroid Stimulating Hormone, [1922](#)  
Thyroid Storm, [1922](#)  
Thyroid Storm Diagnostic Criteria, [1922](#)  
Thyroid Storm Following Thyroidectomy, [1922](#)  
Thyroid Surgery, [1922](#)  
Thyroid Ultrasound, [1922](#)  
Thyroid-Stimulating Hormone, [1922](#)  
Thyroidectomy, [1920](#)

---



---

Thyroiditis, [1921](#)  
 Thyroiditis Classification, [1921](#)  
 Thyrotoxic Myopathy, [1923](#)  
 Thyrotoxic Periodic Paralysis, [1923](#)  
 Thyrotoxicosis, [1923](#)  
 Thyrotroph Adenoma, [1923](#)  
 Thyrotrophin-Releasing Hormone, [1923](#)  
 Thyrotrophin-Stimulating Hormone, [1923](#)  
 Thyrotropin, [1923](#)  
 Thyrotropin Alfa, [1923](#)  
 Thyroxine, [1923](#)  
 Thyroxine-Binding Globulin, [1923](#)  
 TI-RADS, [1928](#)  
 TIA, [1923](#)  
 Tibia, [1923](#)  
 Tibia Vara, [1924](#)  
 Tibial Artery, [1923](#)  
 Tibial Fracture, [1924](#)  
 Tibial Nerve, [1924](#)  
 Tibial Shaft Fractures, [1924](#)  
 Tibialcrural Artery, [1924](#)  
 Tibialis Anterior Muscle, [1924](#)  
 Tibialis Posterior Muscle, [1924](#)  
 Tibiofibular Joint, [1924](#)  
 Tic, [1924](#)  
 Tic Disorder, [1924](#)  
 Tic Douloureux, [1924](#)  
 Ticarcillin, [1924](#)  
 Tick, [1924](#)  
 Tick Bite, [1924](#)  
 Tick Bites, [1924](#)  
 Tick Paralysis, [1925](#)  
 Tick Removal, [1925](#)  
 Tick Typhus, [1925](#)  
 Tick-borne Diseases, [1924](#)  
 Tick-Borne Encephalitis, [1925](#)  
 Tickborne Relapsing Fever, [1925](#)  
 Ticks, [1925](#)  
 Ticlopidine, [1925](#)  
 Tics, [1925](#)  
 Tidal Volume, [1925](#)  
 Tidy Wound, [1925](#)  
 Tietze Syndrome, [1925](#)  
 Tietze's Syndrome, [1925](#)  
 Tigecycline, [1925](#)  
 Tight Junction, [1925](#)  
 Tilorone, [1925](#)  
 Tilt Table Test, [1926](#)  
 Tilt Table Testing, [1926](#)  
 Tilted Disc Syndrome, [1926](#)  
 Time In Range, [1926](#)  
 Time Of Death, [1926](#)  
 Time-Of-Flight PET, [1926](#)  
 Timed Up-And-Go Test, [1926](#)  
 TIMI Risk Score, [1926](#)  
 Timing Of Breastfeeding, [1926](#)  
 Timolol, [1926](#)  
 Timolol Maleate, [1926](#)  
 Timothy Syndrome, [1926](#)  
 Tinea, [1926](#)  
 Tinea Barbae, [1927](#)  
 Tinea Capitis, [1927](#)  
 Tinea Corporis, [1927](#)  
 Tinea Cruris, [1927](#)  
 Tinea Faciei, [1927](#)  
 Tinea Favosa, [1927](#)  
 Tinea Imbricata, [1927](#)  
 Tinea Incognito, [1927](#)  
 Tinea Infection, [1927](#)  
 Tinea Manuum, [1927](#)  
 Tinea Nigra, [1927](#)  
 Tinea Pedis, [1927](#)  
 Tinea Unguium, [1927](#)  
 Tinea Versicolor, [1927](#)  
 Tinel Sign, [1927](#)  
 Tinidazole, [1928](#)  
 Tinnitus, [1928](#)  
 Tinzaparin, [1928](#)  
 Tinzaparin Sodium, [1928](#)  
 Tiopronin, [1928](#)  
 Tiotropium Bromide, [1928](#)  
 Tip Lesion Variant FSGS, [1928](#)  
 TIPS, [1928](#)  
 Tirbanibulin, [1928](#)  
 Tired, [1928](#)  
 Tiredness, [1928](#)  
 Tirofiban, [1928](#)  
 Tissue, [1929](#)  
 Tissue Adhesions, [1929](#)  
 Tissue Banks, [1929](#)  
 Tissue Congestion, [1929](#)  
 Tissue Distribution, [1929](#)  
 Tissue Elastance, [1929](#)  
 Tissue Engineering, [1929](#)  
 Tissue Expander, [1929](#)  
 Tissue Expansion, [1929](#)  
 Tissue Kallikrein, [1929](#)  
 Tissue Membrane, [1929](#)  
 Tissue Plasminogen Activator, [1929](#)  
 Tissue Plasminogen Activator-Associated Angioedema, [1929](#)  
 Tissue Transglutaminase, [1929](#)  
 Tissue Typing, [1929](#)  
 Titanium, [1930](#)  
 Titer, [1930](#)  
 Titration, [1930](#)  
 Titre, [1930](#)  
 Titubation, [1930](#)

---

---

Tizanidine, [1930](#)  
Tmax, [1930](#)  
TMD, [1930](#)  
TMJ Disorder, [1981](#)  
TMJ Disorders, [1930](#)  
TNBC, [1930](#)  
TNF Alpha, [1930](#)  
TNM Classification, [1930](#)  
TNM Classification Of Cancer Stages, [1930](#)  
TNM Staging, [1930](#)  
Tobacco, [1931](#)  
Tobacco Use, [1931](#)  
Tobramycin, [1931](#)  
Tocolysis, [1931](#)  
Tocolytic Agents, [1931](#)  
Tocolytic Therapy, [1931](#)  
Tocolytics, [1931](#)  
Tocopherol, [1931](#)  
Todd Paralysis, [1931](#)  
Todd Paresis, [1932](#)  
Toddler Feeding Patterns and Diet, [1931](#)  
Toddler Test Or Procedure Preparation, [1931](#)  
Toddler's Fracture, [1931](#)  
Toe Joint, [1932](#)  
Toe Pressure, [1932](#)  
Toe-Brachial Index, [1932](#)  
Toenail, [1932](#)  
Toenail Fungus, [1932](#)  
Toes, [1932](#)  
Togaviridae, [1932](#)  
Togaviridae Infections, [1932](#)  
Tokophobia, [1932](#)  
Tolazamide, [1932](#)  
Tolcapone, [1932](#)  
Tolerable Upper Intake Level, [1932](#)  
Tolerance, [1932](#)  
Toll-Like Receptor, [1933](#)  
Toll-Like Receptors, [1933](#)  
Tolmetin, [1933](#)  
Tolmetin Overdose, [1933](#)  
Tolnaftate, [1933](#)  
Tolosa-Hunt Syndrome, [1933](#)  
Tolterodine Tartrate, [1933](#)  
Toluene, [1933](#)  
Toluene And Xylene Poisoning, [1933](#)  
Toluene Toxicity, [1933](#)  
Tolvaptan, [1933](#)  
Tomogram, [1933](#)  
Tomography, [1933](#)  
Tomosynthesis, [1934](#)  
Tomotherapy, [1934](#)  
Tongue, [1934](#)  
Tongue Biopsy, [1934](#)  
Tongue Carcinoma, [1934](#)  
Tongue Disease, [1934](#)  
Tongue Neoplasm, [1934](#)  
Tongue Problems, [1934](#)  
Tongue, Fissured, [748](#)  
Tongue, Geographic, [822](#)  
Tongue, Hairy, [877](#)  
Tongue-Tie, [1934](#)  
Tonic Neck Reflex, [1934](#)  
Tonic Pupil, [1934](#)  
Tonic Pupil Syndrome, [1981](#)  
Tonic Seizures, [1935](#)  
Tonic-Clonic Seizure, [1934](#)  
Tonics, [1934](#)  
Tonometer, [1935](#)  
Tonometry, [1935](#)  
Tonsil, [1935](#)  
Tonsil Stones, [1935](#)  
Tonsillar Cancer, [1935](#)  
Tonsillar Neoplasm, [1935](#)  
Tonsillectomy, [1935](#)  
Tonsillitis, [1935](#)  
Tonus, [1935](#)  
Tooth, [1935](#)  
Tooth Abrasion, [1935](#)  
Tooth Abscess, [1935](#)  
Tooth Anatomy, [1936](#)  
Tooth Ankylosis, [1936](#)  
Tooth Attrition, [1936](#)  
Tooth Avulsion, [1936](#)  
Tooth Brushing, [1936](#)  
Tooth Calcification, [1936](#)  
Tooth Component, [1936](#)  
Tooth Crowding, [1936](#)  
Tooth Crown, [1936](#)  
Tooth Decay, [1936](#)  
Tooth Discoloration, [1936](#)  
Tooth Disease, [1936](#)  
Tooth Enamel, [1936](#)  
Tooth Erosion, [1936](#)  
Tooth Eruption, [1937](#)  
Tooth Extraction, [1937](#)  
Tooth Fracture, [1937](#)  
Tooth Infection, [1937](#)  
Tooth Loss, [1937](#)  
Tooth Luxation, [1937](#)  
Tooth Replantation, [1937](#)  
Tooth Resorption, [1937](#)  
Tooth Root, [1937](#)  
Tooth Sensibility Testing, [1937](#)  
Tooth Sensitivity, [1937](#)  
Toothache, [1936](#)  
Toothaches, [1936](#)  
Toothpaste Overdose, [1937](#)  
Tophaceous Gout, [1937](#)

---

---

Tophus, [1937](#)  
 Topical, [1938](#)  
 Topical Antibiotics, [1938](#)  
 Topical Antifungals, [1938](#)  
 Topical Calcineurin Inhibitors, [1938](#)  
 Topical Corticosteroid Potency, [1938](#)  
 Topical Retinoids, [1938](#)  
 Topiramate, [1938](#)  
 Topographic Spinal Cord Anatomy, [1938](#)  
 Topoisomerase, [1938](#)  
 Topoisomerase Ii, [1938](#)  
 Topoisomerase Inhibitors, [1938](#)  
 Topotecan, [1938](#)  
 Topotecan Hydrochloride, [1938](#)  
 TORCH Panel, [1938](#)  
 TORCH Screen, [1939](#)  
 TORCH Syndrome, [1981](#)  
 Toremifene, [1939](#)  
 Torn ACL, [1939](#)  
 Torn Meniscus, [1939](#)  
 Torn Rotator Cuff, [1981](#)  
 Torpor, [1939](#)  
 Torsade De Pointes, [1939](#)  
 Torsades De Pointes, [1939](#)  
 Torsemide, [1939](#)  
 Torsion, [1939](#)  
 Torsion Dystonia, [1939](#)  
 Torsion of Appendix Testis, [1939](#)  
 Torsion of Testicular Appendage, [1939](#)  
 Torsion Of The Appendix Testis, [1939](#)  
 Torticollis, [1940](#)  
 Torulaspora, [1940](#)  
 Torus Fracture, [1940](#)  
 Torus Mandibularis, [1940](#)  
 Torus Palatinus, [1940](#)  
 Total Abdominal Colectomy, [1940](#)  
 Total Anomalous Pulmonary Venous Connection, [1940](#)  
 Total Anomalous Pulmonary Venous Return, [1940](#)  
 Total Artificial Heart, [1940](#)  
 Total Body Surface Area, [1940](#)  
 Total Breech Extraction, [1940](#)  
 Total Cholesterol, [1941](#)  
 Total Daily Energy Expenditure, [1941](#)  
 Total Haemolytic Complement Test, [1941](#)  
 Total Hip Arthroplasty, [1941](#)  
 Total Hysterectomy, [1941](#)  
 Total Intravenous Anesthesia, [1941](#)  
 Total Iron Binding Capacity, [1941](#)  
 Total Iron-Binding Capacity, [1941](#)  
 Total Knee Arthroplasty, [1941](#)  
 Total Knee Replacement, [1941](#)  
 Total Laryngectomy, [1941](#)  
 Total Lesion Glycolysis, [1941](#)  
 Total Lung Capacity, [1942](#)  
 Total Pancreatectomy, [1942](#)  
 Total Parenteral Nutrition, [1942](#)  
 Total Proctocolectomy And Ileal-Anal Pouch, [1942](#)  
 Total Proctocolectomy With Ileostomy, [1942](#)  
 Total Protein, [1942](#)  
 Total Sleep Time, [1942](#)  
 Total Thyroidectomy, [1942](#)  
 Total-Body Skin Examination, [1940](#)  
 Touch, [1942](#)  
 Tourette Disorder, [1942](#)  
 Tourette Syndrome, [1942](#)  
 Tourette's, [1942](#)  
 Tourette's Syndrome, [1942](#)  
 Tourniquet, [1942](#)  
 Tourniquet Test, [1943](#)  
 Towel Clamp, [1943](#)  
 Townes Syndrome, [1943](#)  
 Toxaphene, [1943](#)  
 Toxbase, [1943](#)  
 Toxic, [1943](#)  
 Toxic Adenoma, [1943](#)  
 Toxic Anterior Segment Syndrome, [1943](#)  
 Toxic Dose, [1943](#)  
 Toxic Effect, [1943](#)  
 Toxic Encephalopathy, [1943](#)  
 Toxic Epidermal Necrolysis, [1944](#)  
 Toxic Megacolon, [1944](#)  
 Toxic Metabolic Encephalopathy, [1944](#)  
 Toxic Neuropathy, [1944](#)  
 Toxic Nodular Goiter, [1944](#)  
 Toxic Shock Syndrome, [1945](#)  
 Toxic Synovitis, [1945](#)  
 Toxicity, [1944](#)  
 Toxicity Grade, [1944](#)  
 Toxicodendron Dermatitis, [1944](#)  
 Toxicodendron Poisoning, [1944](#)  
 Toxicodendron Species, [1944](#)  
 Toxicokinetic Description, [1944](#)  
 Toxicology, [1944](#)  
 Toxicology Screen, [1945](#)  
 Toxicology Specimen Preservation, [1945](#)  
 Toxidrome, [1945](#)  
 Toxin, [1945](#)  
 Toxocara, [1945](#)  
 Toxocara Canis, [1945](#)  
 Toxocara Cati, [1945](#)  
 Toxocariasis, [1945](#)  
 Toxoid, [1945](#)  
 Toxoid Vaccines, [1945](#)  
 Toxoplasma, [1945](#)  
 Toxoplasma Blood Test, [1946](#)  
 Toxoplasma Chorioretinitis, [1946](#)  
 Toxoplasma Gondii, [1946](#)  
 Toxoplasma Meningoencephalitis, [1946](#)

---

---

Toxoplasmosis, [1946](#)  
 Toxoplasmosis Pathology, [1946](#)  
 TPA, [1946](#)  
 TPMT Polymorphism, [1946](#)  
 TPN, [1946](#)  
 Tra Gene, [1949](#)  
 Trabectedin, [1946](#)  
 Trabecula, [1946](#)  
 Trabecular Bone, [1946](#)  
 Trabecular Meshwork, [1946](#)  
 Trabeculectomy, [1947](#)  
 Trabeculoplasty, [1947](#)  
 Trace Elements, [1947](#)  
 Trace Mineral, [1947](#)  
 Tracer, [1947](#)  
 Trachea, [1947](#)  
 Tracheal Agenesis With Bronchoesophageal Fistula, [1947](#)  
 Tracheal Foreign Body, [1947](#)  
 Tracheal Rupture, [1947](#)  
 Tracheal Stenosis, [1947](#)  
 Tracheal Tug, [1947](#)  
 Tracheitis, [1947](#)  
 Trachelectomy, [1947](#)  
 Tracheobronchial Injury, [1948](#)  
 Tracheobronchomalacia, [1948](#)  
 Tracheobronchomegaly, [1948](#)  
 Tracheoesophageal Fistula, [1948](#)  
 Tracheoesophageal Fistula And Esophageal Atresia Repair, [1948](#)  
 Tracheomalacia, [1948](#)  
 Tracheostomy, [1948](#)  
 Tracheostomy Care, [1948](#)  
 Tracheostomy Tube, [1948](#)  
 Tracheotomy, [1948](#)  
 Trachoma, [1949](#)  
 Trachyonychia, [1949](#)  
 Traction, [1949](#)  
 Traction Alopecia, [1949](#)  
 Tractional Retinal Detachment, [1949](#)  
 Tractotomy, [1949](#)  
 Traditional Medicine, [1949](#)  
 Train-Of-Four Monitoring, [1949](#)  
 Trained Immunity, [1949](#)  
 Trajectory, [1949](#)  
 TRALI, [1949](#)  
 Tram-Track Appearance, [1949](#)  
 Trance, [1950](#)  
 Tranexamic Acid, [1950](#)  
 Trans Fat, [1951](#)  
 Trans Fatty Acid, [1951](#)  
 Transabdominal Cerclage, [1950](#)  
 Transabdominal Pelvic Ultrasound, [1950](#)  
 Transabdominal Scan, [1950](#)  
 Transaminitis, [1950](#)  
 Transarterial Chemoembolization, [1950](#)  
 Transbronchial Biopsy, [1950](#)  
 Transcatheter Aortic Valve Replacement, [1950](#)  
 Transcatheter Edge-To-Edge Repair, [1950](#)  
 Transcatheter Pulmonary Valve, [1950](#)  
 Transcervical Foley Catheter For Cervical Ripening, [1951](#)  
 Transcervical Resection Of Endometrium, [1951](#)  
 Transcranial Doppler, [1951](#)  
 Transcranial Doppler Scanning, [1951](#)  
 Transcranial Doppler Ultrasound, [1951](#)  
 Transcranial Magnetic Stimulation, [1951](#)  
 Transcription, [1951](#)  
 Transcription Factor, [1951](#)  
 Transcutaneous Electrical Nerve Stimulation, [1951](#)  
 Transdermal, [1951](#)  
 Transesophageal Echocardiography, [1951](#)  
 Transfer Training, [1952](#)  
 Transferase, [1952](#)  
 Transference, [1952](#)  
 Transferrin, [1952](#)  
 Transferrin Saturation, [1952](#)  
 Transforming Growth Factor-Beta, [1952](#)  
 Transfusion, [1952](#)  
 Transfusion Medicine, [2131](#)  
 Transfusion Of Neonate, [1952](#)  
 Transfusion Reaction, [1952](#)  
 Transfusion-Related Acute Lung Injury, [1952](#)  
 Transient Acantholytic Dermatitis, [1981](#)  
 Transient Acantholytic Dermatitis (Grover Disease), [1952](#)  
 Transient Familial Hyperbilirubinemia, [1952](#)  
 Transient Global Amnesia, [1953](#)  
 Transient Hypogammaglobulinemia Of Infancy, [1953](#)  
 Transient Ischaemic Attack, [1953](#)  
 Transient Ischaemic Attacks Or Episodes, [1953](#)  
 Transient Ischemic Attack, [1953](#)  
 Transient Ischemic Attack (TIA), [1982](#)  
 Transient Ischemic Attack Cross-Reference, [1953](#)  
 Transient Neonatal Pustular Melanosis, [1953](#)  
 Transient Pain, [1953](#)  
 Transient Synovitis, [1953](#)  
 Transient Tachypnea Of The Newborn, [1953](#)  
 Transillumination, [1953](#)  
 Transitional Care, [1953](#)  
 Transitional Epithelium, [1954](#)  
 Transjugular Intrahepatic Portosystemic Shunt, [1954](#)  
 Translation, [1954](#)  
 Translation Initiation, [1954](#)  
 Translocation, [1954](#)  
 Transmembrane Transport, [1954](#)  
 Transmission, [1954](#)  
 Transmural Infarction, [1954](#)  
 Transplacental, [1954](#)  
 Transplant, [1954](#)

---

---

Transplant Immunology, 1955  
 Transplant Rejection, 1955  
 Transplant Surgery, 1955  
 Transplantation, 1954  
 Transplantation Conditioning, 1954  
 Transport, 1955  
 Transport Protein, 1955  
 Transport Vesicle, 1955  
 Transposition Of Great Vessels, 1955  
 Transposition Of The Great Arteries, 1955  
 Transposition of the Great Vessels, 1955  
 Transposons, 1955  
 Transrectal Ultrasonography, 1955  
 Transsphenoidal Hypophysectomy, 1955  
 Transthoracic Echocardiogram, 1956  
 Transthoracic Echocardiography, 1956  
 Transthyretin Amyloidosis, 1956  
 Transthyretin Familial Amyloid Polyneuropathy, 1982  
 Transthyretin Methionine-30 Amyloidosis (Type I), 1982  
 Transtracheal, 1956  
 Transtympanic, 1956  
 Transudate, 1956  
 Transudation, 1956  
 Transudative Pleural Effusion, 1956  
 Transurethral, 1956  
 Transurethral Incision Of The Prostate, 1956  
 Transurethral Microwave Thermotherapy, 1956  
 Transurethral Needle Ablation, 1956  
 Transurethral Resection, 1957  
 Transurethral Resection of Bladder Tumor, 1957  
 Transurethral Resection Of The Prostate, 1957  
 Transvaginal, 1957  
 Transvaginal Ultrasound, 1957  
 Transverse, 1957  
 Transverse Colon, 1957  
 Transverse Lie, 1957  
 Transverse Myelitis, 1957  
 Transverse Plane, 1957  
 Transverse Vaginal Septum, 1957  
 Transversus Abdominis Muscle, 1957  
 Transvestic Disorder, 1957  
 Tranylcypromine, 1958  
 TRAP Sequence, 1958  
 Trapezium Bone, 1958  
 Trapezius Muscle, 1958  
 Trapezoid, 1958  
 Trapezoid Bone, 1958  
 Trapped Lung, 1958  
 Trastuzumab, 1958  
 Trauma, 1958  
 Trauma Center, 1958  
 Trauma Surgery, 1958  
 Trauma- And Stressor-Related Disorders, 1958  
 Traumatic Amputation, 1958  
 Traumatic Bladder and Urethral Injury, 1959  
 Traumatic Bleb, 1959  
 Traumatic Brain Injury, 1959  
 Traumatic Brain Injury (Concussion), 1982  
 Traumatic Events And Children, 1959  
 Traumatic Grief, 1959  
 Traumatic Hemorrhagic Shock, 1959  
 Traumatic Injury, 1959  
 Traumatic Shock, 1959  
 Traumatology, 1959  
 Traumeel S, 1959  
 Travel Medicine, 2131  
 Travel Sickness, 1959  
 Travel Vaccinations, 1959  
 Traveler's Diarrhea, 1982  
 Travelers' Diarrhea, 1959  
 Travoprost, 1960  
 Trazodone, 1960  
 Trazodone Overdose, 1960  
 Trb Gene, 1960  
 Trd Gene, 1960  
 Treacher Collins Syndrome, 1960  
 Treacher Collins-Franceschetti Syndrome 1, 1982  
 Treacle, 1960  
 Treat-To-Target Strategy, 1960  
 Treatment-Resistant Depression, 1960  
 Treatment-Resistant Schizophrenia, 1960  
 Trematoda, 1960  
 Tremelimumab, 1960  
 Tremor, 1960  
 Tremor, Familial, 719  
 Tremor, Familial Essential, 679  
 Tremor, Hereditary Benign, 679  
 Trench Fever, 1960  
 Trench Foot, 1961  
 Trench Mouth, 1961  
 Trendelenberg Position, 1961  
 Trendelenburg, 1961  
 Trendelenburg Gait, 1961  
 Trendelenburg Position, 1961  
 Trendelenburg Sign, 1961  
 Trendelenburg Test, 1961  
 Treosulfan, 1961  
 Trephine, 1961  
 Treponema, 1961  
 Treponema Pallidum, 1961  
 Treponema Pallidum Infection, 1962  
 Treponemal Infections, 1961  
 Trepopnea, 1962  
 Tretinoin, 1962  
 TRG Gene, 1962  
 Triage, 1962  
 Trial Of Labor After Cesarean, 1962  
 Triamcinolone, 1962

---

---

Triamcinolone Acetonide, 1962  
 Triamterene, 1962  
 Triangle of Doom, 1962  
 Triangle of Pain, 1962  
 Triazenes, 1962  
 Triaziquone, 1962  
 Triazolam, 1963  
 Triazole, 1963  
 Triazophos, 1963  
 Triceps, 1963  
 Triceps Brachii Muscle, 1963  
 Triceps Reflex, 1963  
 Trichiasis, 1963  
 Trichilemmal Carcinoma, 1963  
 Trichilemmal Cyst (Pilar Cyst), 1963  
 Trichilemmoma, 1963  
 Trichinellosis, 1963  
 Trichiniasis, 1963  
 Trichinosis, 1963  
 Trichobezoar, 1963  
 Trichoblastoma, 1963  
 Trichoepithelioma, 1964  
 Trichomonas, 1964  
 Trichomonas Vaginalis, 1964  
 Trichomoniasis, 1964  
 Trichophyton, 1964  
 Trichorrhaxis Nodosa, 1964  
 Trichostasis Spinulosa, 1964  
 Trichothiodystrophy Syndrome, 1964  
 Trichotillomania, 1964  
 Trichuriasis, 1964  
 Tricuspid Annular Plane Systolic Excursion, 1964  
 Tricuspid Atresia, 1964  
 Tricuspid Incompetence, 1964  
 Tricuspid Regurgitation, 1965  
 Tricuspid Stenosis, 1965  
 Tricuspid Valve, 1965  
 Tricuspid Valve Disease, 1965  
 Tricyclic Antidepressant Drugs, 1965  
 Tricyclic Antidepressant Overdose, 2131  
 Tricyclic Antidepressant Toxicity, 1965  
 Tricyclic Antidepressants, 1965  
 Trifluoperazine, 1965  
 Trigeminal Nerve, 1965  
 Trigeminal Neuralgia, 1965  
 Trigger, 1965  
 Trigger Finger, 1965  
 Trigger Finger Release, 1966  
 Trigger Thumb, 1966  
 Triglyceride, 1966  
 Triglyceride Level, 1966  
 Triglycerides, 1966  
 Trigone, 1966  
 Triiodothyronine, 1966  
 Trimester, 1966  
 Trimethoprim, 1966  
 Trimethoprim-Sulfamethoxazole, 1966  
 Trimethylaminuria, 1966  
 Trimipramine, 1966  
 Trinitrin, 1966  
 Trinitrophenol, 1966  
 Trinucleotide Repeat, 1967  
 Trip, 1967  
 Tripe Palms, 1967  
 Triple X Syndrome, 1967  
 Triple-Negative Breast Cancer, 1967  
 Triplets, 1967  
 Triploidy Syndrome, 1967  
 Triptan, 1967  
 Triptans, 1967  
 Triquetrum, 1967  
 Trismus, 1967  
 Trisodium Phosphate Poisoning, 1967  
 Trisomy, 1967  
 Trisomy 13, 1968  
 Trisomy 13 Syndrome, 1982  
 Trisomy 18, 1968  
 Trisomy 18 Syndrome, 1982  
 Trisomy 21, 1968  
 Trisomy 21 Syndrome, 1968  
 Trisomy X, 1968  
 Trocar, 1968  
 Trochanter, 1968  
 Trochanteric Bursitis, 1968  
 Troche, 1968  
 Trochlear Nerve, 1968  
 Trochlear Nerve Palsy, 1968  
 Trophic, 1968  
 Trophoblast, 1968  
 Trophoderm, 1968  
 Trophozoite, 1968  
 Tropical Disease, 1969  
 Tropical Diseases, 1969  
 Tropical Medicine, 1969  
 Tropical Sprue, 1969  
 Tropical Ulcer, 1969  
 Troponin, 1969  
 Troponin I, 1969  
 Troponin Test, 1969  
 Trouble Swallowing, 1969  
 Trousseau Sign, 1969  
 Trousseau Sign Of Latent Tetany, 1969  
 Trousseau Sign of Malignancy, 1969  
 Trousseau Syndrome, 1969  
 Trp, 1970  
 True Ribs, 1970  
 Truncation, 1970  
 Truncus Arteriosus, 1970

---



---

Trunk, [1970](#)  
 Trypanosoma, [1970](#)  
 Trypanosoma Brucei, [1970](#)  
 Trypanosomiasis, [1970](#)  
 Trypsin, [1970](#)  
 Trypsin And Chymotrypsin In Stool, [1970](#)  
 Trypsinogen Test, [1970](#)  
 Tryptophan, [1970](#)  
 Tsetse Fly, [1970](#)  
 TSH, [1971](#)  
 TSH Test, [1971](#)  
 TSH-Receptor Antibody, [1971](#)  
 TSI Test, [1971](#)  
 Tsutsugamushi Fever, [1971](#)  
 Tubal Ligation, [1971](#)  
 Tubal Ligation Reversal, [1971](#)  
 Tubal Pregnancy, [1971](#)  
 Tubercle, [1971](#)  
 Tuberculide, [1971](#)  
 Tuberculin, [1971](#)  
 Tuberculin Skin Test, [1971](#)  
 Tuberculin Test, [1971](#)  
 Tuberculoid Leprosy, [1971](#)  
 Tuberculosis, [1972](#)  
 Tuberculosis (TB), [1972](#)  
 Tuberculosis In Children, [1972](#)  
 Tuberculous Meningitis, [1972](#)  
 Tuberculous Pleural Effusion, [1972](#)  
 Tuberosc Sclerosis, [1982](#)  
 Tuberous Sclerosis, [1972](#)  
 Tuberous Sclerosis Complex, [1972](#)  
 Tuberous Sclerosis-1, [1982](#)  
 Tubes, [1972](#)  
 Tubo-Ovarian Abscess, [1972](#)  
 Tubography, [1972](#)  
 Tubular Adenoma, [1972](#)  
 Tubular Maximum, [1972](#)  
 Tubular Reabsorption, [1972](#)  
 Tubular Secretion, [1973](#)  
 Tubule, [1973](#)  
 Tubuloglomerular Feedback, [1973](#)  
 Tubulointerstitial Diseases of the Kidney, [1973](#)  
 Tubulointerstitial Nephritis, [1973](#)  
 Tubulointerstitial Nephropathy, [1973](#)  
 Tularaemia, [1973](#)  
 Tularemia, [1973](#)  
 Tularemia Blood Test, [1973](#)  
 Tulle Gras, [1973](#)  
 Tullio Phenomenon, [1973](#)  
 Tumescence, [1973](#)  
 Tumor, [1973](#)  
 Tumor Board, [1974](#)  
 Tumor Burden, [1974](#)  
 Tumor Differentiation, [1974](#)  
 Tumor Embolism, [1974](#)  
 Tumor Grade, [1974](#)  
 Tumor Grading, [1974](#)  
 Tumor Heterogeneity, [1974](#)  
 Tumor Lysis Syndrome, [1974](#)  
 Tumor Marker, [1974](#)  
 Tumor Mutational Burden, [1974](#)  
 Tumor Necrosis Factor, [1974](#)  
 Tumor Necrosis Factor Inhibitor, [2131](#)  
 Tumor Necrosis Factor Receptor-Associated Periodic Syndrome, [1974](#)  
 Tumor Staging, [1975](#)  
 Tumor Suppressor, [1975](#)  
 Tumor Suppressor Gene, [1975](#)  
 Tumor-Infiltrating Lymphocyte Therapy, [1974](#)  
 Tumor-Suppressor Gene, [1975](#)  
 Tumour, [1975](#)  
 Tungiasis, [1975](#)  
 Tunica Adventitia, [1975](#)  
 Tunica Albuginea, [1975](#)  
 Tunica Media, [1975](#)  
 Tunica Vaginalis, [1975](#)  
 Tunnel Vision, [1975](#)  
 Turbinate Surgery, [1975](#)  
 Turbinates, [1975](#)  
 Turbinates Hypertrophy, [1975](#)  
 Turcot Syndrome, [1975](#)  
 Turgor, [1976](#)  
 Turner Hypoplasia, [1976](#)  
 Turner Syndrome, [1976](#)  
 Turner's Syndrome, [1976](#)  
 Turner-Varny Syndrome, [1982](#)  
 Turning Patients Over In Bed, [1976](#)  
 Turpentine Oil Poisoning, [1976](#)  
 Tussis, [1976](#)  
 TVS, [1976](#)  
 Twin, [1976](#)  
 Twin Placenta, [1976](#)  
 Twin Pregnancy, [1976](#)  
 Twin Twin Transfusion Syndrome, [1982](#)  
 Twin-To-Twin Transfusion Syndrome, [1976](#)  
 Twin-Twin Transfusion Syndrome, [1977](#)  
 Twins, [1976](#)  
 Twitch, [1977](#)  
 Twitching, [1977](#)  
 Tympanic, [1977](#)  
 Tympanic Bulla, [1977](#)  
 Tympanic Cavity, [1977](#)  
 Tympanic Hyperectasis, [1977](#)  
 Tympanic Membrane, [1977](#)  
 Tympanic Membrane Perforation, [1977](#)  
 Tympanites, [1977](#)  
 Tympanometer, [1977](#)  
 Tympanometry, [1977](#)

---

---

Tympanoplasty, [1977](#)  
Tympanosclerosis, [1977](#)  
Tympanostomy Tube Insertion, [1978](#)  
Tympanostomy Tubes, [1978](#)  
Tympanum, [1978](#)  
Tympany, [1978](#)  
Tyndall Effect, [1978](#)  
Type 1 Diabetes, [1978](#)  
Type 1 Diabetes In Children, [1978](#)  
Type 1 Diabetes Mellitus, [1978](#)  
Type 2 Diabetes, [1978](#)  
Type 2 Diabetes In Children, [1978](#)  
Type 2 Diabetes Meal Planning, [1978](#)  
Type 2 Diabetes Medications, [1979](#)  
Type 2 Diabetes Mellitus, [1979](#)  
Type 3B Diabetes, [1979](#)  
Type I Hypersensitivity, [1979](#)  
Type I Muscle Fiber, [1979](#)  
Type Ia Glycogen Storage Disease, [1979](#)  
Type Ib Glycogen Storage Disease, [1979](#)  
Type II Hypersensitivity, [1979](#)  
Type II Muscle Fiber, [1979](#)  
Type III GSD (Forbes-Cori), [1979](#)  
Type III Hypersensitivity, [1979](#)  
Type III Hypersensitivity Reaction, [1979](#)  
Type IV GSD (Andersen), [1979](#)  
Type IV Hypersensitivity, [1979](#)  
Type V Glycogen Storage Disease, [1980](#)  
Type VI Glycogen Storage Disease, [1980](#)  
Type VII Glycogen Storage Disease, [1980](#)  
Types Of Ileostomy, [1979](#)  
Typhlitis, [1980](#)  
Typhoid, [1980](#)  
Typhoid Fever, [1980](#)  
Typhus, [1980](#)  
Typhus Fever, [1980](#)  
Typical Antipsychotics, [1980](#)  
Tyramine, [1980](#)  
Tyrosine, [1980](#)  
Tyrosine Kinase Inhibitor, [2131](#)  
Tyrosine Kinase Inhibitors, [1981](#)  
Tyrosinemia, [1981](#)  
Tzanck Smear, [1981](#)

U Wave, [2013](#)  
UA (Urinalysis), [1983](#)  
Ubiquitin-Proteasome System, [1983](#)  
UBT (Urea Breath Test), [1983](#)  
UDP-Glucuronosyltransferase, [1983](#)  
UDP-Glucuronosyltransferase Deficiency, [1983](#)  
Uhl Anomaly, [1983](#)  
Uhthoff Phenomenon, [1983](#)  
Ulcer, [1983](#)  
Ulcer Healing Drugs, [1984](#)  
Ulceration, [1983](#)  
Ulcerative Colitis, [1983](#)  
Ulcerative Colitis (UC), [2013](#)  
Ulcerative Colitis Pathology, [1984](#)  
Ulcerative Keratitis, [1984](#)  
Ulcerative Proctitis, [1984](#)  
Ulcers, [1984](#)  
Ulerythema, [1984](#)  
Ullrich Congenital Muscular Dystrophy, [1984](#)  
Ulna, [1984](#)  
Ulnar, [1984](#)  
Ulnar Artery, [1984](#)  
Ulnar Claw Hand, [1984](#)  
Ulnar Collateral Ligament, [1984](#)  
Ulnar Deviation, [1985](#)  
Ulnar Drift, [1985](#)  
Ulnar Hypoplasia, [1985](#)  
Ulnar Nerve, [1985](#)  
Ulnar Nerve Dysfunction, [1985](#)  
Ulnar Nerve Entrapment, [1985](#)  
Ulnar Neuropathy, [1985](#)  
Ulnar Neuropathy at Guyon Canal, [1985](#)  
Ulnar Paradox, [1985](#)  
Ulnar Tunnel Syndrome, [1985](#)  
Ulnar Vein, [1986](#)  
Ulnar Wrist Pain, [1986](#)  
Ulnar-Sided Wrist Pain, [1985](#)  
Ultra-processed food, [1986](#)  
Ultrafiltration, [1986](#)  
Ultrasonic Therapy, [1986](#)  
Ultrasonography, [1986](#)  
Ultrasound, [1986](#)  
Ultrasound Biomicroscopy, [1986](#)  
Ultrasound Contrast, [1986](#)  
Ultrasound Scan, [2013](#)  
Ultrasound Therapy, [1986](#)  
Ultrasound Transducer, [1987](#)  
Ultrasound-Guided Biopsy, [1986](#)  
Ultrasound-Guided Breast Biopsy, [1986](#)  
Ultraviolet Rays, [1987](#)  
Ultraviolet Therapy, [1987](#)  
Umbilical, [1987](#)  
Umbilical Artery Catheter, [1987](#)  
Umbilical Artery Doppler, [1987](#)  
Umbilical Catheters, [1987](#)  
Umbilical Cord, [1987](#)  
Umbilical Cord Blood Gas, [1987](#)  
Umbilical Cord Complications, [1987](#)  
Umbilical Cord Prolapse, [1987](#)  
Umbilical Granuloma, [1987](#)  
Umbilical Hernia, [1987](#)  
Umbilical Region, [1988](#)  
Umbilical Vein, [1988](#)  
Umbilication, [1988](#)

---

---

Umbilicus, [1988](#)  
 Uncal Herniation, [1988](#)  
 Uncal Transtentorial Herniation, [1988](#)  
 Unconjugated Hyperbilirubinemia, [1988](#)  
 Unconscious, [1988](#)  
 Unconsciousness, [1988](#)  
 Uncoupling Protein, [1988](#)  
 Uncover Test, [1988](#)  
 Underactive Thyroid, [2013](#)  
 Underventilation, [1989](#)  
 Underweight, [1989](#)  
 Undescended Testicle, [1989](#)  
 Undescended Testicle Repair, [1989](#)  
 Undescended Testis, [1989](#)  
 Undifferentiated, [1989](#)  
 Undifferentiated Arthritis, [1989](#)  
 Undifferentiated Cancer, [1989](#)  
 Undifferentiated Carcinoma, [1989](#)  
 Undifferentiated Connective Tissue Disease, [1989](#)  
 Undifferentiated Connective-Tissue Disease, [1989](#)  
 Undifferentiated Pleomorphic Sarcoma, [1989](#)  
 Undifferentiated Pleomorphic Sarcoma Of The Breast, [1989](#)  
 Undifferentiated Spondyloarthritis, [1989](#)  
 Undulant Fever, [1989](#)  
 Unerupted Tooth, [1990](#)  
 Ungual, [1990](#)  
 Unicameral Bone Cyst, [1990](#)  
 Unicornuate Uterus, [1990](#)  
 Unilateral, [1990](#)  
 Unilateral Gynecomastia, [1990](#)  
 Unilateral Microphthalmos, [1990](#)  
 Unilateral Spatial Neglect, [1990](#)  
 Uninodular Goiter, [1990](#)  
 Unintentional Weight Loss, [1990](#)  
 Uniparental Disomy, [1990](#)  
 Universal Donor, [1990](#)  
 Universal Precautions, [2013](#)  
 Universal Recipient, [1991](#)  
 Unloader Knee Braces, [1991](#)  
 Unopposed Estrogen, [1991](#)  
 Unresectable, [1991](#)  
 Unresponsive Coma, [1991](#)  
 Unsaturated Fat, [1991](#)  
 Unstable Angina, [1991](#)  
 Unsteadiness, [1991](#)  
 Untidy Wound, [1991](#)  
 Upper Airway Biopsy, [1991](#)  
 Upper Airway Cough Syndrome, [1991](#)  
 Upper Airway Obstruction, [1991](#)  
 Upper Airway Resistance Syndrome, [1992](#)  
 Upper Arm, [1992](#)  
 Upper Endoscopy, [1992](#)  
 Upper Esophageal Sphincter, [1992](#)  
 Upper Extremity, [1992](#)  
 Upper Extremity Deep Vein Thrombosis, [1992](#)  
 Upper Extremity Venous Doppler, [1992](#)  
 Upper Gastrointestinal Bleeding, [1992](#)  
 Upper GI And Small Bowel Series, [1992](#)  
 Upper GI Series, [1992](#)  
 Upper Leg, [1992](#)  
 Upper Limb Disorders, [1992](#)  
 Upper Motor Neuron, [1992](#)  
 Upper Motor Neuron Lesion, [1993](#)  
 Upper Motor Neuron Syndrome, [1993](#)  
 Upper Respiratory Infection, [1993](#)  
 Upper Respiratory Tract Infection, [1993](#)  
 Upper Tract Urothelial Carcinoma, [2008](#)  
 Upper-Tract Urothelial Carcinoma, [1993](#)  
 Upright Labor Positions, [1993](#)  
 Upward Gaze, [1993](#)  
 Urachal Abnormalities, [1993](#)  
 Urachal Carcinoma, [1993](#)  
 Urachal Cyst, [1993](#)  
 Urachal Fistula, [1993](#)  
 Urachal Mucinous Cystic Tumor, [1993](#)  
 Urachus, [1993](#)  
 Uracil-DNA Glycosylase, [1994](#)  
 Uraemia, [1994](#)  
 Uraemic Foetor, [1994](#)  
 Uraemic Syndrome, [1994](#)  
 Uranium, [1994](#)  
 Urate, [1994](#)  
 Urate Nephropathy, [1994](#)  
 Urate-Lowering Therapy, [1994](#)  
 Urates, [1994](#)  
 Urea, [1994](#)  
 Urea Breath Test, [1994](#)  
 Urea Cycle, [1994](#)  
 Urea Cycle Disorder, [1995](#)  
 Urea Nitrogen Urine Test, [1995](#)  
 Ureaplasma, [1995](#)  
 Ureaplasma Urealyticum, [1995](#)  
 Uremia, [1995](#)  
 Uremic Bleeding, [1995](#)  
 Uremic Encephalopathy, [1995](#)  
 Uremic Fetor, [1995](#)  
 Uremic Frost, [1995](#)  
 Uremic Pericarditis, [1995](#)  
 Uremic Platelet Dysfunction, [1995](#)  
 Uremic Pruritus, [1995](#)  
 Uremic Stomatitis, [1996](#)  
 Ureter, [1996](#)  
 Ureter Duplex, [1997](#)  
 Ureter Fissus, [1997](#)  
 Ureter Obstruction, [1997](#)  
 Ureteral Calculi, [1996](#)  
 Ureteral Catheters, [1996](#)

---

---

Ureteral Injury During Gynecologic Surgery, [1996](#)  
Ureteral Obstruction, [1996](#)  
Ureteral Orifice, [1996](#)  
Ureteral Perforation, [1996](#)  
Ureteral Polyp, [1996](#)  
Ureteral Reimplantation, [1996](#)  
Ureteral Retrograde Brush Biopsy, [1996](#)  
Ureteral Stenosis, [1996](#)  
Ureteral Stent, [1996](#)  
Ureteral Stones, [2013](#)  
Ureteral Stricture, [1996](#)  
Ureteric Bud, [1997](#)  
Ureteric Calculi, [1997](#)  
Ureteric Colic, [1997](#)  
Ureteric Perforation, [1997](#)  
Ureteritis, [1997](#)  
Ureterocele, [1997](#)  
Ureterolithiasis, [1997](#)  
Ureteropelvic Junction Obstruction, [1997](#)  
Ureteroscope, [1997](#)  
Ureteroscopes, [1997](#)  
Ureteroscopy, [1998](#)  
Ureterostomy Procedure, [1998](#)  
Urethra, [1998](#)  
Urethra Disorder, [1998](#)  
Urethral Cancer, [1998](#)  
Urethral Carcinoma, [1998](#)  
Urethral Caruncle, [1998](#)  
Urethral Catheter, [1998](#)  
Urethral Discharge, [1999](#)  
Urethral Diverticulum, [1998](#)  
Urethral Fistula, [1998](#)  
Urethral Hypermobility, [1998](#)  
Urethral Infection, [1998](#)  
Urethral Injury, [1998](#)  
Urethral Meatus, [1998](#)  
Urethral Obstruction, [1999](#)  
Urethral Pressure Profile, [1999](#)  
Urethral Prolapse, [1999](#)  
Urethral Sling, [1999](#)  
Urethral Sphincter, [1999](#)  
Urethral Stenosis, [1999](#)  
Urethral Stent, [1999](#)  
Urethral Stricture, [1999](#)  
Urethral Syndrome, [1999](#)  
Urethral Trauma, [1999](#)  
Urethritis, [1999](#)  
Urethrocele, [1999](#)  
Urethrography, [2000](#)  
Urethroplasty, [2000](#)  
Urethroscopy, [2000](#)  
Urge Incontinence, [2000](#)  
Urge Urinary Incontinence, [2000](#)  
Urgency, [2000](#)  
URI (Upper Respiratory Infection), [2006](#)  
Uric Acid, [2000](#)  
Uric Acid Nephrolithiasis, [2000](#)  
Uric Acid Nephropathy, [2000](#)  
Uric Acid Stones, [2000](#)  
Uric Acid Urine Test, [2000](#)  
Uricosuric Agent, [2131](#)  
Uricosuric Drug, [2000](#)  
Uridine Triacetate, [2000](#)  
Urinal, [2000](#)  
Urinalysis, [2001](#)  
Urinary, [2001](#)  
Urinary Anion Gap, [2001](#)  
Urinary Bladder, [2001](#)  
Urinary Bladder Disorder, [2001](#)  
Urinary Bladder Squamous Cell Carcinoma Pathology, [2001](#)  
Urinary Calculi, [2001](#)  
Urinary Casts, [2001](#)  
Urinary Catheter, [2001](#)  
Urinary Catheterization, [2001](#)  
Urinary Catheters, [2001](#)  
Urinary Diversion, [2001](#)  
Urinary Fistula, [2001](#)  
Urinary Frequency, [2001](#)  
Urinary Hesitation, [2002](#)  
Urinary Incontinence, [2002](#)  
Urinary Lithiasis, [2002](#)  
Urinary Retention, [2002](#)  
Urinary Retraining, [2002](#)  
Urinary Stasis, [2002](#)  
Urinary System, [2002](#)  
Urinary Tract, [2002](#)  
Urinary Tract Disorder, [2002](#)  
Urinary Tract Imaging, [2002](#)  
Urinary Tract Infection, [2002](#)  
Urinary Tract Infection (UTI), [2013](#)  
Urinary Tract Infection In Adults, [2002](#)  
Urinary Tract Infection In Children, [2002](#)  
Urinary Tract Infection In Pregnancy, [2003](#)  
Urinary Tract Infections in Pregnancy, [2003](#)  
Urinary Tract Neoplasm, [2003](#)  
Urinary Tract Obstruction, [2003](#)  
Urinary Urgency, [2003](#)  
Urination, [2003](#)  
Urination Disorder, [2003](#)  
Urine, [2003](#)  
Urine 24-Hour Volume, [2003](#)  
Urine Albuminuria, [2003](#)  
Urine Alkalinization, [2003](#)  
Urine Amylase Test, [2003](#)  
Urine Analysis, [2003](#)  
Urine Bilirubin Test, [2003](#)  
Urine Calcium Test, [2004](#)

---

[Urine Catecholamine Test, 2004](#)  
[Urine Chemistry, 2004](#)  
[Urine Chloride Test, 2004](#)  
[Urine Concentration Test, 2004](#)  
[Urine Conductivity, 2004](#)  
[Urine Culture, 2004](#)  
[Urine Cytology, 2004](#)  
[Urine Dipstick, 2004](#)  
[Urine Drainage Bags, 2004](#)  
[Urine Drug Screen, 2131](#)  
[Urine Electrolytes Test, 2004](#)  
[Urine Formation, 2005](#)  
[Urine HCG Test, 2005](#)  
[Urine Hemoglobin, 2005](#)  
[Urine Immunoelectrophoresis, 2005](#)  
[Urine Immunofixation, 2005](#)  
[Urine Osmolality, 2005](#)  
[Urine PH, 2005](#)  
[Urine PH Test, 2005](#)  
[Urine Protein Dipstick Test, 2005](#)  
[Urine Protein Electrophoresis Test, 2005](#)  
[Urine Retention, 2005](#)  
[Urine Sediment, 2005](#)  
[Urine Sodium, 2005](#)  
[Urine Specific Gravity Test, 2005](#)  
[Urine Toxicology, 2006](#)  
[Urinoma, 2006](#)  
[Urinometer, 2006](#)  
[Urobilinogen, 2006](#)  
[Urodynamic Testing, 2006](#)  
[Urodynamics, 2006](#)  
[Uroflow, 2006](#)  
[Uroflowmetry, 2006](#)  
[Urogenital, 2006](#)  
[Urogenital Fistula, 2006](#)  
[Urogenital Sinus, 2006](#)  
[Urogenital System, 2006](#)  
[Urogenital Tract, 2006](#)  
[Urography, 2007](#)  
[Urokinase, 2007](#)  
[Urolithiasis, 2007](#)  
[Urologic Diagnostic Yield, 2007](#)  
[Urologist, 2007](#)  
[Urology, 2007](#)  
[Uropathy, 2007](#)  
[Urosepsis, 2007](#)  
[Urostomy, 2013](#)  
[Urostomy Leak, 2007](#)  
[Urostomy Procedure, 2007](#)  
[Urostomy Prolapse, 2007](#)  
[Urostomy Stenosis, 2007](#)  
[Urothelial Carcinoma, 2007](#)  
[Urothelium, 2008](#)  
[Urotrauma, 2008](#)

---

UV Light, [2012](#)  
Uvea, [2012](#)  
Uveal Disease, [2012](#)  
Uveal Melanoma, [2012](#)  
Uveal Neoplasm, [2012](#)  
Uveitic Glaucoma, [2012](#)  
Uveitis, [2012](#)  
Uveoparotid Fever, [2012](#)  
Uvula, [2012](#)  
Uvulitis, [2012](#)  
Uvulopalatopharyngoplasty, [2013](#)

V-Fib, [2043](#)  
Vaccination, [2015](#)  
Vaccination Schedule, [2015](#)  
Vaccinations/Immunizations During Pregnancy, [2015](#)  
Vaccine, [2015](#)  
Vaccine Adjuvant, [2131](#)  
Vaccine Allergy, [2015](#)  
Vaccine Therapy, [2131](#)  
Vaccine-Associated Paralytic Poliomyelitis, [2015](#)  
Vaccines, [2015](#)  
Vaccines and Immunizations, [2015](#)  
Vaccinia, [2015](#)  
Vaccinia Virus, [2015](#)  
VACTERL Association, [2016](#)  
Vacuum Curettage, [2016](#)  
Vacuum Delivery, [2016](#)  
Vacuum Erection Devices, [2016](#)  
Vacuum Extractor, [2016](#)  
Vacuum-Assisted Delivery, [2016](#)  
Vagal Maneuver, [2016](#)  
Vagal Nerve Stimulation, [2016](#)  
Vagal Paraganglioma, [2016](#)  
Vagina, [2016](#)  
Vagina Carcinoma, [2016](#)  
Vagina Neoplasm, [2020](#)  
Vaginal Agenesis, [2016](#)  
Vaginal Atresia, [2016](#)  
Vaginal Atrophy, [2017](#)  
Vaginal Bleeding, [2017](#)  
Vaginal Bleeding Between Periods, [2017](#)  
Vaginal Bleeding During Early Pregnancy, [2017](#)  
Vaginal Bleeding During Late Pregnancy, [2017](#)  
Vaginal Bleeding In Early Pregnancy, [2017](#)  
Vaginal Bleeding In Late Pregnancy, [2017](#)  
Vaginal Bleeding In Pregnancy, [2017](#)  
Vaginal Cancer, [2017](#)  
Vaginal Candidiasis, [2017](#)  
Vaginal Cuff, [2017](#)  
Vaginal Cuff Dehiscence, [2017](#)  
Vaginal Cysts, [2017](#)  
Vaginal Dilator, [2018](#)  
Vaginal Discharge, [2018](#)  
Vaginal Dryness, [2018](#)  
Vaginal Fistula, [2018](#)  
Vaginal Foreign Body, [2018](#)  
Vaginal Fornix, [2018](#)  
Vaginal Fusion And Duplication, [2018](#)  
Vaginal Health, [2018](#)  
Vaginal Hematoma, [2018](#)  
Vaginal Hemorrhage, [2018](#)  
Vaginal Hypoplasia, [2018](#)  
Vaginal Hysterectomy, [2018](#)  
Vaginal Inflammation, [2018](#)  
Vaginal Intercourse, [2018](#)  
Vaginal Intraepithelial Neoplasia, [2018](#)  
Vaginal Introitus, [2019](#)  
Vaginal Itching, [2019](#)  
Vaginal Laceration, [2019](#)  
Vaginal Lubrication, [2019](#)  
Vaginal Membrane, [2019](#)  
Vaginal Microbiome, [2019](#)  
Vaginal Mucosa, [2019](#)  
Vaginal Odor, [2019](#)  
Vaginal Opening, [2019](#)  
Vaginal Pessary, [2019](#)  
Vaginal pH, [2019](#)  
Vaginal Prolapse, [2019](#)  
Vaginal Rugae, [2019](#)  
Vaginal Seeding, [2019](#)  
Vaginal Speculum, [2019](#)  
Vaginal Stenosis, [2020](#)  
Vaginal Swab, [2020](#)  
Vaginal Trauma, [2020](#)  
Vaginal Ultrasound Probe, [2020](#)  
Vaginal Vault Prolapse, [2020](#)  
Vaginal Vestibule, [2020](#)  
Vaginal Wall, [2020](#)  
Vaginal Yeast Infection, [2020](#)  
Vaginectomy, [2020](#)  
Vaginismus, [2020](#)  
Vaginitis, [2020](#)  
Vaginoplasty, [2020](#)  
Vaginospasm, [2020](#)  
Vagotomy, [2021](#)  
Vagus Nerve, [2021](#)  
Vagus Nerve Stimulation, [2021](#)  
Valacyclovir, [2021](#)  
Valganciclovir, [2021](#)  
Valgus, [2021](#)  
Valgus Stress Test, [2021](#)  
Valinemia, [2021](#)  
Valley Fever, [2021](#)  
Valproate, [2021](#)  
Valproate Toxicity, [2021](#)  
Valproic Acid, [2021](#)  
Valproic Acid Toxicity, [2021](#)

---



---

Valrubicin, [2022](#)  
Valsalva, [2022](#)  
Valsalva Maneuver, [2022](#)  
Valsalva Retinopathy, [2022](#)  
Valsalva's Manoeuvre, [2022](#)  
Valsartan, [2022](#)  
Valves, [2022](#)  
Valvotomy, [2022](#)  
Valvulae Conniventes, [2022](#)  
Valvular Disease, [2022](#)  
Valvular Endocarditis, [2022](#)  
Valvular Heart Disease, [2023](#)  
Valvular Insufficiency, [2023](#)  
Valvuloplasty, [2023](#)  
Valvulotomy, [2023](#)  
Van der Woude Syndrome, [2023](#)  
Vancomycin, [2023](#)  
Vancomycin Flushing Reaction, [2023](#)  
Vancomycin-Resistant Enterococcus, [2023](#)  
Vandetanib, [2023](#)  
Vaping, [2023](#)  
Vaporiser, [2023](#)  
Variable Decelerations, [2023](#)  
Variant Angina Pectoris, [2023](#)  
Variant CJD, [2024](#)  
Variant Classification, [2024](#)  
Varibaculum Cambriense, [2024](#)  
Variceal Band Ligation, [2024](#)  
Variceal Hemorrhage, [2024](#)  
Varicella, [2024](#)  
Varicella Vaccine, [2131](#)  
Varicella-Zoster Virus, [2024](#)  
Varicellovirus, [2024](#)  
Varices, [2024](#)  
Varicocele, [2024](#)  
Varicocele Embolization, [2024](#)  
Varicose Ulcer, [2024](#)  
Varicose Veins, [2024](#)  
Varicose Veins (Spider Veins), [2062](#)  
Varicose Veins and Venous Insufficiency, [2024](#)  
Variegate Porphyria, [2025](#)  
Variola, [2025](#)  
Varix, [2025](#)  
Varus, [2025](#)  
Varus Stress Test, [2025](#)  
Vas, [2025](#)  
Vas Deferens, [2030](#)  
Vasa Deferentia, [2025](#)  
Vasa Efferentia, [2025](#)  
Vasa Previa, [2025](#)  
Vasa Recta, [2025](#)  
Vasa Vasorum, [2025](#)  
Vascular, [2025](#)  
Vascular Access, [2025](#)  
Vascular Access For Hemodialysis, [2025](#)  
Vascular Balloon Catheter, [2026](#)  
Vascular Bed, [2026](#)  
Vascular Bypass, [2026](#)  
Vascular Claudication, [2026](#)  
Vascular Clip, [2026](#)  
Vascular Compliance, [2026](#)  
Vascular Complications, [2026](#)  
Vascular Dementia, [2026](#)  
Vascular Disease, [2026](#)  
Vascular Ehlers-Danlos Syndrome, [2026](#)  
Vascular Embolic Particles, [2026](#)  
Vascular Embolization, [2026](#)  
Vascular Embolization Coil, [2027](#)  
Vascular Embolization Plug, [2027](#)  
Vascular Endothelial Growth Factor, [2027](#)  
Vascular Endothelial Growth Factor Inhibitor, [2027](#)  
Vascular Endothelial Growth Factor Inhibitors, [2027](#)  
Vascular Forceps, [2027](#)  
Vascular Graft, [2027](#)  
Vascular Graft Patency, [2027](#)  
Vascular Hemostat, [2027](#)  
Vascular Imaging, [2027](#)  
Vascular Laser Therapy, [2027](#)  
Vascular Ligation, [2028](#)  
Vascular Malformation, [2028](#)  
Vascular Needle Holder, [2028](#)  
Vascular Pain, [2028](#)  
Vascular Patch Graft, [2028](#)  
Vascular Retraction, [2028](#)  
Vascular Ring, [2028](#)  
Vascular Scissors, [2028](#)  
Vascular Shunt, [2028](#)  
Vascular Sponge, [2028](#)  
Vascular Staple, [2028](#)  
Vascular Stent, [2028](#)  
Vascular Stent Balloon, [2029](#)  
Vascular Stent-Graft, [2029](#)  
Vascular Suction, [2029](#)  
Vascular Surgery, [2029](#)  
Vascular System, [2029](#)  
Vascular Thrombectomy, [2029](#)  
Vascular Tissue Adhesive, [2029](#)  
Vascular Tone, [2029](#)  
Vascular Trauma, [2029](#)  
Vascular Ulcer, [2029](#)  
Vascularity, [2027](#)  
Vasculitic Neuropathy, [2029](#)  
Vasculitis, [2029](#)  
Vasectomy, [2030](#)  
Vasectomy Reversal, [2030](#)  
Vaso-Occlusive Crisis, [2031](#)  
Vasoactive Intestinal Peptide Test, [2030](#)  
Vasoconstriction, [2030](#)

---

---

Vasoconstrictor, [2030](#)  
 Vasoconstrictors, [2030](#)  
 Vasodepressor Syncope, [2030](#)  
 Vasodilatation, [2030](#)  
 Vasodilation, [2030](#)  
 Vasodilator, [2030](#)  
 Vasodilators, [2030](#)  
 Vasodilatory Shock, [2030](#)  
 Vasomotor, [2030](#)  
 Vasomotor Center, [2031](#)  
 Vasomotor Centre, [2031](#)  
 Vasomotor Nerves, [2031](#)  
 Vasomotor Rhinitis, [2031](#)  
 Vasopressin, [2031](#)  
 Vasopressin Receptor, [2031](#)  
 Vasopressin Receptor Antagonists, [2031](#)  
 Vasopressor, [2031](#)  
 Vasopressors In Anesthesia, [2031](#)  
 Vasospasm, [2031](#)  
 Vasovagal Attack, [2031](#)  
 Vasovagal Reaction, [2031](#)  
 Vasovagal Syncope, [2032](#)  
 Vater-Pacini Corpuscles, [2032](#)  
 VATS, [2032](#)  
 Vaughan Williams Classification, [2032](#)  
 VCJD, [2032](#)  
 VDRL Test, [2032](#)  
 Vector, [2032](#)  
 Vector Control, [2032](#)  
 Vector-Borne Disease, [2032](#)  
 Vegetarian, [2032](#)  
 Vegetation, [2032](#)  
 Vein, [2032](#)  
 Vein of Galen Malformation, [2033](#)  
 Vein-Graft Aneurysms, [2032](#)  
 VeIP Regimen, [2033](#)  
 Velamentous Cord Insertion, [2033](#)  
 Velocardiofacial Syndrome, [2033](#)  
 Velopharyngeal Insufficiency, [2033](#)  
 Vemurafenib, [2033](#)  
 Vena Cava, [2033](#)  
 Vena Cava Filter, [2033](#)  
 Venepuncture, [2033](#)  
 Venereal Disease Research Laboratory Test, [2033](#)  
 Venesection, [2033](#)  
 Venipuncture, [2033](#)  
 Veno-Occlusive Disease, [2034](#)  
 Veno-occlusive Hepatic Disease (Sinusoidal Obstruction Syndrome), [2034](#)  
 Venogram, [2033](#)  
 Venography, [2034](#)  
 Venom, [2034](#)  
 Venomous Bite, [2034](#)  
 Venous, [2034](#)  
 Venous Air Embolism, [2034](#)  
 Venous Aneurysm, [2034](#)  
 Venous Blood Gas, [2034](#)  
 Venous Capacitance, [2034](#)  
 Venous Diseases, [2034](#)  
 Venous Hum, [2034](#)  
 Venous Insufficiency, [2035](#)  
 Venous Leak, [2035](#)  
 Venous Pressure, [2035](#)  
 Venous Return, [2035](#)  
 Venous Stasis Dermatitis, [2035](#)  
 Venous Stasis Ulcer, [2035](#)  
 Venous Thromboembolism, [2035](#)  
 Venous Thromboembolism In Pregnancy, [2035](#)  
 Venous Thromboembolism Prophylaxis, [2035](#)  
 Venous Thrombosis, [2035](#)  
 Venous Ulcer, [2035](#)  
 Ventilation, [2035](#)  
 Ventilation-Perfusion Mismatch, [2035](#)  
 Ventilation-Perfusion Ratio, [2036](#)  
 Ventilation-Perfusion Scan, [2036](#)  
 Ventilator, [2036](#)  
 Ventilator-Associated Pneumonia, [2036](#)  
 Ventilator-Induced Lung Injury, [2036](#)  
 Ventouse, [2036](#)  
 Ventral, [2036](#)  
 Ventral Body Cavity, [2036](#)  
 Ventral Hernia, [2036](#)  
 Ventral Root, [2036](#)  
 Ventricle, [2036](#)  
 Ventricular, [2036](#)  
 Ventricular Aneurysm, [2036](#)  
 Ventricular Arrhythmia, [2036](#)  
 Ventricular Arrhythmias, [2037](#)  
 Ventricular Assist Device, [2037](#)  
 Ventricular Bigeminy, [2037](#)  
 Ventricular Fibrillation, [2037](#)  
 Ventricular Hypertrophy, [2037](#)  
 Ventricular Myocardium, [2037](#)  
 Ventricular Non-Compaction, [2037](#)  
 Ventricular Premature Contraction, [2037](#)  
 Ventricular Remodeling, [2037](#)  
 Ventricular Rupture, [2037](#)  
 Ventricular Septal Defect, [2037](#)  
 Ventricular Septal Defect Cross-Reference, [2037](#)  
 Ventricular Tachycardia, [2037](#)  
 Ventriculoatrial, [2038](#)  
 Ventriculography, [2038](#)  
 Ventriculoperitoneal Shunt, [2038](#)  
 Ventrosuspension, [2038](#)  
 Venturi Mask, [2038](#)  
 Venturi Principle, [2038](#)  
 Venule, [2038](#)  
 Venules, [2038](#)

---

---

Verapamil, [2038](#)  
 Verapamil Toxicity, [2038](#)  
 Vermicides, [2038](#)  
 Vernal Keratoconjunctivitis, [2039](#)  
 Verner-Morrison Syndrome, [2039](#)  
 Vernix, [2039](#)  
 Vernix Caseosa, [2039](#)  
 Verruca, [2039](#)  
 Verruca Plana, [2039](#)  
 Verruca Vulgaris, [2039](#)  
 Verrucas, [2039](#)  
 Verruciform Xanthoma, [2039](#)  
 Verrucose, [2039](#)  
 Verrucous, [2039](#)  
 Verrucous Carcinoma, [2039](#)  
 Version, [2039](#)  
 Vertebra, [2040](#)  
 Vertebrae, [2040](#)  
 Vertebral Artery, [2040](#)  
 Vertebral Artery Dissection, [2040](#)  
 Vertebral Canal, [2040](#)  
 Vertebral Column, [2040](#)  
 Vertebral Compression Fracture, [2040](#)  
 Vertebral Hemangioma, [2040](#)  
 Vertebral Tumor, [2040](#)  
 Vertebrobasilar Circulatory Disorders, [2040](#)  
 Vertebrobasilar Insufficiency, [2040](#)  
 Vertebrobasilar Insufficiency, [2040](#)  
 Vertebroplasty, [2040](#)  
 Vertical, [2040](#)  
 Vertical Talus, [2041](#)  
 Vertical Transmission, [2041](#)  
 Vertigo, [2041](#)  
 Vertigo-Associated Disorders, [2041](#)  
 Very Long-Chain Acyl-CoA Dehydrogenase Deficiency, [2041](#)  
 Very-Low-Density Lipoprotein, [2041](#)  
 Vesical, [2041](#)  
 Vesicant, [2041](#)  
 Vesicants, [2041](#)  
 Vesicle, [2041](#)  
 Vesicles, [2041](#)  
 Vesico-Ureteric Reflux, [2042](#)  
 Vesicoureteral Reflux, [2041](#)  
 Vesicovaginal and Ureterovaginal Fistula, [2042](#)  
 Vesicovaginal Fistula, [2042](#)  
 Vesicular, [2042](#)  
 Vesicular Breathing, [2042](#)  
 Vesicular Palmoplantar Eczema, [2042](#)  
 Vesiculobullous Dermatoses, [2042](#)  
 Vessel, [2042](#)  
 Vestibular, [2042](#)  
 Vestibular Disorder, [2042](#)  
 Vestibular Migraine, [2042](#)  
 Vestibular Neuritis, [2042](#)  
 Vestibular Papillomatosis, [2042](#)  
 Vestibular Rehabilitation, [2042](#)  
 Vestibular Schwannoma, [2043](#)  
 Vestibular System, [2043](#)  
 Vestibule, [2043](#)  
 Vestibulo-Ocular Reflex, [2043](#)  
 Vestibulocochlear Nerve, [2043](#)  
 Vestibulodynia, [2043](#)  
 Vestigial, [2043](#)  
 VEXAS Syndrome, [2043](#)  
 Viable, [2043](#)  
 Vibration Problem, [2043](#)  
 Vibrator, [2043](#)  
 Vibrio, [2043](#)  
 Vibrio Cholerae, [2043](#)  
 Vibrio Infection, [2043](#)  
 Vibrio Parahaemolyticus Infection, [2044](#)  
 Vibrio vulnificus Infection, [2044](#)  
 Vibrio Vulnificus Infections, [2044](#)  
 Vibrios, [2044](#)  
 Video Laryngoscope, [2044](#)  
 Video Laryngoscopy, [2044](#)  
 Video-Assisted Thoracoscopic Surgery, [2044](#)  
 Videofluoroscopic Swallow Study, [2044](#)  
 Videostroboscopy, [2044](#)  
 Vildagliptin, [2044](#)  
 Villi, [2044](#)  
 Villous Adenoma, [2045](#)  
 Villous Atrophy, [2045](#)  
 Villus, [2045](#)  
 Villus Atrophy, [2045](#)  
 Vinca Alkaloids, [2045](#)  
 Vincent Angina, [2045](#)  
 Vincent Disease, [2045](#)  
 Vincent Infection, [2045](#)  
 Vincent Stomatitis, [2045](#)  
 Vincent's Angina, [2045](#)  
 Vinorelbine, [2045](#)  
 Viomycin, [2045](#)  
 VIPoma, [2045](#)  
 VIPoma Syndrome, [2045](#)  
 Viraemia, [2046](#)  
 Viral, [2046](#)  
 Viral Antibody, [2046](#)  
 Viral Arthritis, [2046](#)  
 Viral Assembly, [2046](#)  
 Viral Attachment, [2046](#)  
 Viral Carcinogenesis, [2046](#)  
 Viral Conjunctivitis, [2046](#)  
 Viral Culture, [2046](#)  
 Viral Encephalitis, [2046](#)  
 Viral Entry, [2046](#)  
 Viral Exanthem, [2046](#)

---

---

Viral Gastroenteritis, [2046](#)  
Viral Haemorrhagic Fever, [2046](#)  
Viral Hemorrhagic Fevers, [2047](#)  
Viral Hepatitis, [2047](#)  
Viral Hepatitis Types, [2047](#)  
Viral Immune Evasion, [2047](#)  
Viral Infection, [2047](#)  
Viral Infections and Pregnancy, [2047](#)  
Viral Infections of the Mouth, [2047](#)  
Viral Latency, [2047](#)  
Viral Latency Mechanisms, [2047](#)  
Viral Load, [2047](#)  
Viral Meningitis, [2047](#)  
Viral Myocarditis, [2047](#)  
Viral Pathogenesis, [2048](#)  
Viral Pericarditis, [2048](#)  
Viral Pharyngitis, [2048](#)  
Viral Pneumonia, [2048](#)  
Viral Release, [2048](#)  
Viral Replication, [2048](#)  
Viral Replication Cycle, [2048](#)  
Viral Rhinitis, [2048](#)  
Viral Structure, [2048](#)  
Viral Uncoating, [2048](#)  
Virchow Node, [2048](#)  
Virchow Triad, [2049](#)  
Virchow's Triad, [2049](#)  
Virchow-Robin Spaces, [2049](#)  
Viremia, [2049](#)  
Viridans Streptococci, [2049](#)  
Virilisation, [2049](#)  
Virilism, [2049](#)  
Virilism Foetal, [2049](#)  
Virilization, [2049](#)  
Virilize, [2049](#)  
Virology, [2049](#)  
Virotherapy, [2049](#)  
Virtual Colonoscopy, [2049](#)  
Virulence, [2050](#)  
Virulence Factor, [2050](#)  
Virulent, [2050](#)  
Virus, [2050](#)  
Virus Activation, [2050](#)  
Virus Assembly, [2050](#)  
Virus Diseases, [2050](#)  
Virus Integration, [2050](#)  
Virus Protein, [2050](#)  
Virus Receptor, [2050](#)  
Virus Replication, [2050](#)  
Virus Uncoating, [2050](#)  
Viruses, [2050](#)  
Viscera, [2050](#)  
Visceral, [2050](#)  
Visceral Afferents, [2050](#)  
Visceral Larva Migrans, [2051](#)  
Visceral Leishmaniasis, [2051](#)  
Visceral Pain, [2022](#)  
Visceromegaly, [2051](#)  
Viscus, [2051](#)  
Vision, [2051](#)  
Vision And Hearing, [2051](#)  
Vision Impairment, [2051](#)  
Vision Loss, [2051](#)  
Vision Problems, [2051](#)  
Vision Receptors, [2051](#)  
Vision Screening, [2051](#)  
Vision Test, [2051](#)  
Visual Acuity, [2052](#)  
Visual Acuity Test, [2052](#)  
Visual Agnosia, [2052](#)  
Visual Evoked Potential, [2052](#)  
Visual Field, [2052](#)  
Visual Field Defect, [2052](#)  
Visual Field Perimetry, [2052](#)  
Visual Field Test, [2052](#)  
Visual Fields, [2052](#)  
Visual Impairment, [2052](#)  
Visual Neglect, [2052](#)  
Visualized Laser-Assisted Prostatectomy, [2052](#)  
Vital, [2052](#)  
Vital Capacity, [2053](#)  
Vital Signs, [2053](#)  
Vitamin A, [2053](#)  
Vitamin A Blood Test, [2053](#)  
Vitamin A Deficiency, [2053](#)  
Vitamin A Toxicity, [2053](#)  
Vitamin A: Retinal, [2053](#)  
Vitamin B, [2053](#)  
Vitamin B Complex, [2054](#)  
Vitamin B1, [2053](#)  
Vitamin B12, [2053](#)  
Vitamin B12 Deficiency, [2053](#)  
Vitamin B12 Deficiency Anemia, [2053](#)  
Vitamin B12 Deficiency In Older Adults, [2054](#)  
Vitamin B12 Level, [2054](#)  
Vitamin B12: Cobalamin, [2053](#)  
Vitamin B1: Thiamine, [2054](#)  
Vitamin B2, [2054](#)  
Vitamin B2 Riboflavin, [2054](#)  
Vitamin B3, [2054](#)  
Vitamin B5, [2054](#)  
Vitamin B6, [2054](#)  
Vitamin B6: Pyridoxine, [2054](#)  
Vitamin B7, [2054](#)  
Vitamin B9, [2054](#)  
Vitamin C, [2055](#)  
Vitamin C Deficiency, [2055](#)  
Vitamin D, [2055](#)

---

---

Vitamin D Deficiency, [2055](#)  
Vitamin D3, [2055](#)  
Vitamin D: Calcitriol, [2055](#)  
Vitamin Deficiency Anemia, [2055](#)  
Vitamin E, [2055](#)  
Vitamin E Deficiency, [2055](#)  
Vitamin E: Tocopherol, [2055](#)  
Vitamin K, [2055](#)  
Vitamin K Deficiency, [2055](#)  
Vitamin K Deficiency Bleeding, [2056](#)  
Vitamin K Prophylaxis, [2056](#)  
Vitamin K: Phylloquinone, [2056](#)  
Vitamin Toxicity, [2056](#)  
Vitamins, [2056](#)  
Vitiligo, [2056](#)  
Vitrectomy, [2056](#)  
Vitreous, [2056](#)  
Vitreous Body, [2056](#)  
Vitreous Detachment, [2056](#)  
Vitreous Hemorrhage, [2056](#)  
Vitreous Humor, [2056](#)  
VKORC1 Polymorphism, [2056](#)  
VLDL Test, [2057](#)  
VO2 Max, [2057](#)  
Vocal Cord Disorders, [2062](#)  
Vocal Cord Dysfunction, [2062](#)  
Vocal Cord Nodules, [2057](#)  
Vocal Cord Paralysis, [2057](#)  
Vocal Cord Paresis, [2057](#)  
Vocal Cord Polyp, [2057](#)  
Vocal Cords, [2057](#)  
Vocal Fold Paralysis, [2057](#)  
Vocal Folds, [2057](#)  
Vocal Hemorrhage, [2057](#)  
Vocal Nodules, [2062](#)  
Vocal Resonance, [2057](#)  
Vocal-Cord Dysfunction, [2057](#)  
Vocal-Fold Nodules, [2057](#)  
Vocal-Fold Polyps, [2057](#)  
Vogt-Koyanagi-Harada Disease, [2057](#)  
Vogt-Koyanagi-Harada Syndrome, [2058](#)  
Vohwinkel Syndrome, [2058](#)  
Voice, [2058](#)  
Voice And Speech, [2058](#)  
Voice Box, [2058](#)  
Voice Disorders, [2058](#)  
Void, [2058](#)  
Voiding Cystourethrogram, [2058](#)  
Volar, [2058](#)  
Volatile Anesthetics, [2058](#)  
Volkmann Contracture, [2058](#)  
Volume Of Distribution, [2059](#)  
Volume Overload, [2059](#)  
Voluntary, [2059](#)  
Voluntary Psychiatric Admission, [2059](#)  
Volutrauma, [2059](#)  
Volvulus, [2059](#)  
Vomit, [2059](#)  
Vomiting, [2059](#)  
Vomiting Blood, [2059](#)  
Vomitus, [2059](#)  
Von Economo Encephalitis, [2059](#)  
Von Gierke Disease, [2059](#)  
Von Graefe Sign, [2059](#)  
Von Hippel-Lindau Disease, [2060](#)  
Von Hippel-Lindau Syndrome, [2060](#)  
Von Meyenburg Complex, [2060](#)  
Von Recklinghausen Disease, [2060](#)  
Von Recklinghausen'S Disease, [2060](#)  
Von Recklinghausen'S Neurofibromatosis, [2060](#)  
Von Recklinghausen'S Phakomatosis, [2060](#)  
Von Rosen Splint, [2060](#)  
Von Willebrand Disease, [2060](#)  
VOR Reflex, [2060](#)  
Voriconazole, [2060](#)  
Voyeurism, [2060](#)  
Voyeuristic Disorder, [2060](#)  
VPCs (Ventricular Premature Contractions), [2060](#)  
VRE (Vancomycin-Resistant Enterococci), [2061](#)  
VRE Infection, [2022](#)  
VSD (Ventricular Septal Defect), [2061](#)  
Vulnerable, [2061](#)  
Vulva, [2061](#)  
Vulval Intraepithelial Neoplasia, [2061](#)  
Vulvar, [2061](#)  
Vulvar Cancer, [2061](#)  
Vulvar Carcinoma, [2061](#)  
Vulvar Inclusion Cysts, [2061](#)  
Vulvar Intraepithelial Neoplasia, [2061](#)  
Vulvar Melanoma, [2061](#)  
Vulvar Squamous Cell Carcinoma, [2061](#)  
Vulvectomy, [2061](#)  
Vulvitis, [2061](#)  
Vulvodynia, [2062](#)  
Vulvovaginal Candidiasis, [2062](#)  
Vulvovaginitis, [2062](#)  
Waardenburg Syndrome, [2063](#)  
Waddell Signs, [2063](#)  
WAGR Syndrome, [2063](#)  
Waist Circumference, [2063](#)  
Waist-Hip Ratio, [2063](#)  
Wait List, [2063](#)  
Wakefulness, [2063](#)  
Waldenstrom Macroglobulinemia, [2063](#)  
Waldenstrom's Macroglobulinemia, [2064](#)  
Waldenström's Macroglobulinemia, [2063](#)  
Waldeyer Ring, [2064](#)

---

---

Walker-Warburg Syndrome, [2064](#)  
Walking Abnormalities, [2064](#)  
Walking Pneumonia, [2064](#)  
Walking Speed, [2064](#)  
Wall Stress, [2064](#)  
Wallenberg Syndrome, [2064](#)  
Wallerian Degeneration, [2064](#)  
Walleyed, [2064](#)  
Wandering Rash Tongue, [2085](#)  
Wandering Spleen, [2064](#)  
Warburg Effect, [2065](#)  
Warfarin, [2065](#)  
Warfarin and Superwarfarin Toxicity, [2065](#)  
Warfarin Syndrome, [2065](#)  
Warfarin-Induced Skin Necrosis, [2065](#)  
Warm and Cold Autoantibodies, [2065](#)  
Warm Autoimmune Hemolytic Anemia, [2065](#)  
Warm Ischemia, [2065](#)  
Wart, [2065](#)  
Wart Remover Poisoning, [2065](#)  
Warthin Tumor, [2065](#)  
Warts and Plantar Warts, [2085](#)  
Warts, Genital (Human Papillomavirus), [820](#)  
Warty Dyskeratoma, [2065](#)  
Wasting, [2066](#)  
Watchful Waiting, [2066](#)  
Water Birth, [2066](#)  
Water Blister, [2066](#)  
Water Brash, [2066](#)  
Water Channel, [2066](#)  
Water Consumption, [2066](#)  
Water Hammer Pulse, [2066](#)  
Water Intoxication, [2067](#)  
Water On The Brain, [2067](#)  
Water On The Knee, [2067](#)  
Water Retention, [2067](#)  
Water-Borne Disease, [2066](#)  
Water-Deprivation Test, [2066](#)  
Water-Hammer Pulse, [2066](#)  
Water-Soluble Vitamin, [2067](#)  
Waterborne Bacterial Disease, [2066](#)  
Waterhouse-Friderichsen Syndrome, [2067](#)  
Watershed Infarction, [2067](#)  
Waterston Shunt, [2067](#)  
Watery Eyes, [2067](#)  
Wax, [2067](#)  
Wax Poisoning, [2067](#)  
Waxy Casts, [2067](#)  
WBC Count, [2067](#)  
WBS, [2085](#)  
Weakness, [2067](#)  
Weaning, [2068](#)  
Wear Debris, [2068](#)  
Weaver Syndrome, [2068](#)  
Weaver's Bottom, [2068](#)  
Webbed Fingers, [2068](#)  
Webbed Neck, [2068](#)  
Webbing Of The Fingers Or Toes, [2068](#)  
Weber, [2068](#)  
Weber Syndrome, [2068](#)  
Weber Test, [2068](#)  
Weber-Christian Disease, [2068](#)  
Wedge Excision, [2068](#)  
Wedge Resection, [2068](#)  
Weever Fish, [2068](#)  
Weight Gain, [2069](#)  
Weight Gain In Pregnancy, [2069](#)  
Weight Gain – Unintentional, [2069](#)  
Weight Loss, [2069](#)  
Weight Loss Medications, [2069](#)  
Weight Loss Surgery, [2069](#)  
Weight Management, [2085](#)  
Weight-Bearing As Tolerated, [2069](#)  
Weight-Loss Medicines, [2069](#)  
Weight-Reducing Diets, [2069](#)  
Weil Disease, [2069](#)  
Weil's Disease, [2070](#)  
Weil-Felix Reaction, [2069](#)  
Weil-Felix Test, [2069](#)  
Weill-Marchesani Syndrome, [2069](#)  
Weir-Mitchell Disease, [2085](#)  
Wellens Syndrome, [2070](#)  
Wells Score, [2070](#)  
Wells Score For Pulmonary Embolism, [2070](#)  
Wells Syndrome, [2086](#)  
Wells Syndrome (Eosinophilic Cellulitis), [2070](#)  
Welt, [2070](#)  
Wen, [2070](#)  
Wenckebach Phenomenon, [2070](#)  
Wens, [2070](#)  
Werdnig-Hoffmann Disease, [2070](#)  
Werlhof Disease, [2085](#)  
Wermer Syndrome, [2071](#)  
Wermer's Syndrome, [2071](#)  
Werner Syndrome, [2071](#)  
Werner's Syndrome, [2071](#)  
Wernicke Aphasia, [2071](#)  
Wernicke Area, [2071](#)  
Wernicke Encephalopathy, [2071](#)  
Wernicke's Encephalopathy, [2071](#)  
Wernicke-Korsakoff Encephalopathy, [2071](#)  
Wernicke-Korsakoff Syndrome, [2071](#)  
Wernicke'S Area, [2071](#)  
West Nile Encephalitis, [2072](#)  
West Nile Fever, [2085](#)  
West Nile Neuroinvasive Disease, [2072](#)  
West Nile Virus, [2072](#)  
West Nile Virus Infection, [2072](#)

---



---

West Syndrome, [2072](#)  
Westergren Method, [2071](#)  
Westermarck Sign, [2071](#)  
Western Blot, [2072](#)  
Westley Croup Score, [2072](#)  
Wet Age-Related Macular Degeneration, [2072](#)  
Wet AMD, [2072](#)  
Wet Beriberi, [2072](#)  
Wet Dream, [2072](#)  
Wet Gangrene, [2072](#)  
Wet Macular Degeneration, [2072](#)  
Wet-to-Dry Dressing, [2073](#)  
Wharton Duct, [2073](#)  
Wharton Jelly, [2073](#)  
Wharton's Jelly, [2073](#)  
Wheal, [2073](#)  
Wheal And Flare, [2073](#)  
Wheat Allergy, [2073](#)  
Wheelchairs, [2073](#)  
Wheeled Stretcher, [2073](#)  
Wheeze, [2073](#)  
Wheezing, [2073](#)  
Whiplash, [2073](#)  
Whiplash Injury, [2073](#)  
Whipple Disease, [2074](#)  
Whipple Procedure, [2074](#)  
Whipple Triad, [2074](#)  
Whipple Triad for Insulinoma, [2074](#)  
Whipple Triad of Hypoglycemia, [2074](#)  
Whipple's Disease, [2074](#)  
Whipworm, [2074](#)  
Whipworm Infection, [2074](#)  
Whirlpool Bath, [2074](#)  
Whispered Pectoriloquy, [2074](#)  
White Blood Cell, [2074](#)  
White Blood Cell Count, [2074](#)  
White Blood Cell Labeling, [2075](#)  
White Cell, [2075](#)  
White Cell Count, [2075](#)  
White Clot Syndrome, [2075](#)  
White Coat Hypertension, [2075](#)  
White Finger, [2075](#)  
White Infarction, [2075](#)  
White Matter, [2075](#)  
White Matter Hyperintensities, [2075](#)  
White Matter Of The Brain, [2075](#)  
White Of The Eye, [2076](#)  
White Spot Disease, [2076](#)  
White Tongue, [2076](#)  
White-Coat Hypertension, [2075](#)  
Whiteheads, [2075](#)  
Whitlow, [2076](#)  
Whitlows, [2076](#)  
WHO Classification of Vitamin A Deficiency, [2076](#)  
Whole Bowel Irrigation, [2076](#)  
Whole Breast Radiation Therapy, [2077](#)  
Whole Exome Sequencing, [2077](#)  
Whole Genome Sequencing, [2077](#)  
Whole-Body Irradiation, [2076](#)  
Whole-Body Magnetic Resonance Imaging, [2076](#)  
Whole-Body Scan, [2076](#)  
Whole-Bowel Irrigation, [2076](#)  
Whole-Brain Radiation Therapy, [2077](#)  
Whole-Genome Sequencing, [2077](#)  
Whooping Cough, [2077](#)  
Whooping Cough (Pertussis), [2085](#)  
Whooping-Cough, [2077](#)  
Whorled Pattern, [2077](#)  
WHS, [2085](#)  
Wickham Striae, [2077](#)  
Widal Reaction, [2077](#)  
Widal Test, [2077](#)  
Wide Splitting of S2, [2078](#)  
Widely Spaced Teeth, [2077](#)  
Widened Mediastinum, [2077](#)  
Widened Pulse Pressure, [2078](#)  
Wilkie Syndrome, [2078](#)  
Williams Syndrome, [2078](#)  
Williams-Beuren Syndrome, [2085](#)  
Willis-Ekbom Disease, [1733](#)  
Wilm's Tumour, [2078](#)  
Wilms Tumor, [2078](#)  
Wilson Disease, [2078](#)  
Wilson's Disease, [2078](#)  
Winchester Syndrome, [2078](#)  
Windkessel Effect, [2078](#)  
Windowing, [2078](#)  
Windpipe, [2079](#)  
Winged Scapula, [2079](#)  
Winter Formula, [2079](#)  
Winter Vomiting Disease, [2079](#)  
Winter's Formula, [2079](#)  
Winterbottom Sign, [2079](#)  
Wireless Motility Capsule, [2079](#)  
Wirsung Duct, [2079](#)  
Wisdom Teeth, [2079](#)  
Wisdom Tooth, [2079](#)  
Wisdom Tooth Problems, [2086](#)  
Wisdom Tooth Removal, [2086](#)  
Wiskott-Aldrich Syndrome, [2080](#)  
Wiskott-Aldrich Syndrome Protein, [2080](#)  
Witch's Milk, [2080](#)  
Withdrawal, [2080](#)  
Withdrawal Reflex, [2080](#)  
Withdrawal Symptoms, [2080](#)  
Withdrawal Syndrome, [2080](#)  
Withdrawal Syndromes, [2080](#)  
Wolf Hirschhorn Syndrome, [2086](#)

---

---

Wolf Syndrome, [2086](#)  
Wolf-Hirschhorn Syndrome, [2080](#)  
Wolff-Chaikoff Effect, [2080](#)  
Wolff-Parkinson-White Accessory Pathway-Related Tachycardia, [2080](#)  
Wolff-Parkinson-White Syndrome, [2086](#)  
Wolffian Ducts, [2080](#)  
Wolfram Syndrome, [2081](#)  
Wolinella, [2081](#)  
Wolman Disease, [2081](#)  
Wolman's Disease, [2081](#)  
Womb, [2081](#)  
Womb Growths, [2081](#)  
Women Health, [2081](#)  
Women's Health, [2081](#)  
Wood Lamp Examination, [2081](#)  
Wood Spirit, [2081](#)  
Wood's Light, [2081](#)  
Wood-Unit, [2081](#)  
Woods Corkscrew Maneuver, [2081](#)  
Woods Screw Maneuver, [2081](#)  
Woody Tongue, [2082](#)  
Woolly Hair, [2082](#)  
Woolsorters' Disease, [2082](#)  
Word Blindness, [2082](#)  
Word Deafness, [2082](#)  
Word Recognition Scoring, [2082](#)  
Word Salad, [2082](#)  
Working Memory, [2082](#)  
Wound, [2082](#)  
Wound Care, [2082](#)  
Wound Classification, [2082](#)  
Wound Complication, [2082](#)  
Wound Debridement, [2082](#)  
Wound Dehiscence, [2083](#)  
Wound Evisceration, [2083](#)  
Wound Healing, [2083](#)  
Wound Healing Phases, [2083](#)  
Wound Infection, [2083](#)  
Wound Management, [2083](#)  
Wound VAC, [2083](#)  
Wound Vacuum, [2083](#)  
Wounds, [2083](#)  
WPW Syndrome, [2086](#)  
Wright Hyperabduction Test, [2083](#)  
Wright Stain, [2083](#)  
Wright Test, [2084](#)  
Wrinkle Test, [2084](#)  
Wrisberg Ganglion, [2084](#)  
Wrist, [2084](#)  
Wrist Arthroscopy, [2084](#)  
Wrist Drop, [2084](#)  
Wrist Fracture, [2084](#)  
Wrist Injury, [2084](#)  
Wrist Joint, [2084](#)  
Wrist Pain, [2084](#)  
Wrist Sprain, [2084](#)  
Writer's Cramp, [2084](#)  
Written Documentation, [2084](#)  
Wry-Neck, [2084](#)  
Wt, [2085](#)  
Wuchereria, [2085](#)  
Wuchereria Bancrofti, [2085](#)  
Wuhan Coronavirus, [2085](#)  
  
X Chromosome, [2089](#)  
X-Inactivation, [2090](#)  
X-Linked, [2090](#)  
X-Linked Adrenoleukodystrophy, [2090](#)  
X-linked Adult Spinal Muscular Atrophy, [2093](#)  
X-Linked Agammaglobulinemia, [2090](#)  
X-Linked Chronic Granulomatous Disease, [2090](#)  
X-Linked Dominant Inheritance, [2091](#)  
X-Linked Emery-Dreifuss Muscular Dystrophy, [2091](#)  
X-Linked Hyper-IgM Syndrome, [2091](#)  
X-Linked Hypophosphatemia, [2091](#)  
X-Linked Ichthyosis, [2091](#)  
X-Linked Inheritance, [2091](#)  
X-linked Intellectual Disability, [2093](#)  
X-linked Juvenile Retinoschisis, [2093](#)  
X-Linked Lymphoproliferative Disease Type 1, [2091](#)  
X-Linked Lymphoproliferative Syndrome, [2091](#)  
X-Linked Myotubular Myopathy, [2091](#)  
X-Linked Recessive Inheritance, [2091](#)  
X-Linked Retinitis Pigmentosa, [2091](#)  
X-Linked Retinoschisis, [2092](#)  
X-Linked Severe Combined Immunodeficiency, [2092](#)  
X-Linked Sideroblastic Anemia, [2092](#)  
X-Linked Spinal and Bulbar Muscular Atrophy, [2092](#)  
X-Ray, [2092](#)  
X-Ray Attenuation, [2092](#)  
X-Ray Crystallography, [2092](#)  
X-Ray Shield, [2092](#)  
X-Ray Therapy, [2092](#)  
Xanthelasma, [2087](#)  
Xanthine, [2087](#)  
Xanthine Oxidase, [2087](#)  
Xanthine Oxidase Inhibitor, [2131](#)  
Xanthine Oxidase Inhibitors, [2087](#)  
Xanthinuria, [2087](#)  
Xanthochromia, [2087](#)  
Xanthogranulomatous, [2087](#)  
Xanthogranulomatous Cholecystitis, [2087](#)  
Xanthogranulomatous Oophoritis, [2087](#)  
Xanthogranulomatous Pyelonephritis, [2087](#)  
Xanthoma, [2088](#)  
Xanthoma Disseminatum, [2088](#)  
Xanthoma Eruptivum, [2088](#)

---

---

Xanthoma Planum, 2088  
Xanthoma Striatum Palmare, 2088  
Xanthoma Tendinosum, 2088  
Xanthoma Tuberosum, 2088  
Xanthomas, 2088  
Xanthomatosis, 2088  
Xanthopsia, 2088  
Xanthosis, 2088  
Xenobiotic, 2089  
Xenobiotic Metabolism, 2089  
Xenobiotic Resistance, 2089  
Xenograft, 2089  
Xenon Anesthesia, 2089  
Xenotransplantation, 2089  
Xerochilia, 2089  
Xeroderma, 2089  
Xeroderma Pigmentosum, 2089  
Xeroderma Pigmentosum Complementation Groups, 2089  
Xerophthalmia, 2089  
Xerophthalmic Ulceration, 2090  
Xerosis, 2090  
Xerostomia, 2090  
Xerotic Eczema, 2090  
Xiphisternal Joint, 2090  
Xiphoid Process, 2090  
Xiphoidalgia, 2090  
Xiphoidectomy, 2090  
XO Syndrome, 2093  
XP, 2093  
XX Male Syndrome, 2092  
XXX Syndrome, 2092  
XXXX Syndrome, 2092  
XXY Males, 2093  
XXY Syndrome, 2093  
Xylitol, 2092  
Xylose, 2093  
Xylose Absorption Test, 2093  
XYY Syndrome, 2093  
  
Y Chromosome, 2095  
Y-Box Binding Protein, 2095  
Y-Chromosome Microdeletion, 2095  
Y-Linked Gene, 2098  
Y-Linked Inheritance, 2098  
Y-Maze Test, 2098  
Y-Plate Osteosynthesis, 2099  
Y-V Plasty, 2099  
YAG Capsulotomy, 2095  
YAG Laser Capsulotomy, 2095  
YAG Laser Surgery, 2095  
Yawn, 2095  
Yawning, 2095  
Yaws, 2095  
  
Yaws Papilloma, 2095  
Yeast, 2095  
Yeast Infection, 2096  
Yeast Infection In Women And Men, 2096  
Yeast Infections, 2099  
Yeast Present, 2096  
Yeast Vulvitis, 2096  
Yellow Cartilage, 2096  
Yellow Fever, 2096  
Yellow Fever Vaccine, 2096  
Yellow Fever Virus, 2096  
Yellow Ligament, 2096  
Yellow Marrow, 2096  
Yellow Nail Syndrome, 2096  
Yellow Nail Syndrome Triad, 2096  
Yellow Spot, 2097  
Yellow Tongue, 2097  
Yellow-Nail Syndrome, 2096  
Yergason Test, 2097  
Yersinia, 2097  
Yersinia Enterocolitica, 2097  
Yersinia Enterocolitica Pseudoappendicitis, 2097  
Yersinia Infection, 2097  
Yersinia Outer Proteins, 2097  
Yersinia Pestis, 2097  
Yersinia Pestis Pneumonic Plague, 2097  
Yersinia Pseudotuberculosis, 2097  
Yersinia Reactive Arthritis, 2097  
Yersiniosis, 2098  
Yew Poisoning, 2098  
Yolk Sac, 2098  
Yolk Sac Carcinoma, 2098  
Yolk Sac Tumor, 2098  
Yolk Sac Tumor Pathology, 2098  
Yolk Sac Tumor Schiller-Duval Bodies, 2098  
Yolk Stalk, 2098  
Yolk-Sac Tumor, 2098  
Young Syndrome, 2099  
Young-Burgess Classification, 2098  
Young-Onset Dementia, 1588  
Yttrium-90, 2099  
Yttrium-90 Microspheres, 2099  
Yttrium-90 Radioembolization, 2099  
Yttrium-Aluminum-Garnet Laser, 2099  
Yuzpe Regimen, 2099  
  
Z-Disc, 2101  
Z-Plasty, 2105  
Z-Score (DEXA), 2105  
Z-Score (Echocardiography), 2105  
Z-Thumb Deformity, 2105  
Z-Tracking Technique, 2105  
Zafirlukast, 2101  
Zahn Lines, 2101

---

---

Zanamivir, [2101](#)  
ZAP-70 Deficiency, [2101](#)  
Zavanelli Maneuver, [2101](#)  
Zellweger Syndrome, [2101](#)  
Zenker Degeneration, [2101](#)  
Zenker Diverticulum, [2101](#)  
Zenker Peroral Endoscopic Myotomy, [2102](#)  
Ziconotide, [2102](#)  
Zidovudine, [2102](#)  
Ziehl-Neelsen Stain, [2102](#)  
Zieve Syndrome, [2102](#)  
Zika, [2102](#)  
Zika Fever, [2106](#)  
Zika Virus, [2102](#)  
Zika Virus Infection, [2106](#)  
Zileuton, [2102](#)  
Zinc, [2102](#)  
Zinc Acetate, [2102](#)  
Zinc Deficiency, [2103](#)  
Zinc Finger Motif, [2103](#)  
Zinc Oxide, [2103](#)  
Zinc Oxide Ointment, [2103](#)  
Zinc Poisoning, [2106](#)  
Zinc Protoporphyrin, [2103](#)  
Zinc Sulfate, [2103](#)  
Zinc Toxicity, [2103](#)  
Zinner Syndrome, [2103](#)  
Zirconia, [2103](#)  
Zoledronic Acid, [2103](#)  
Zollinger-Ellison Secretin Stimulation Test, [2103](#)  
Zollinger-Ellison Syndrome, [2104](#)  
Zolpidem, [2104](#)  
Zona Fasciculata, [2104](#)  
Zona Glomerulosa, [2104](#)  
Zona Pellucida, [2104](#)  
Zona Reticularis, [2104](#)  
Zonisamide, [2104](#)  
Zonula Adherens, [2104](#)  
Zonula Occludens, [2104](#)  
Zonule, [2104](#)  
Zoon Balanitis, [2104](#)  
Zoonosis, [2105](#)  
Zoonotic Spillover, [2105](#)  
Zoster, [2105](#)  
Zoster (Shingles), [2107](#)  
Zoster Ophthalmicus, [2105](#)  
Zygapophyseal Joint, [2105](#)  
Zygapophyseal Joint Arthropathy, [2106](#)  
Zygomatic Arch, [2106](#)  
Zygomatic Bone, [2106](#)  
Zygomatic Fracture, [2106](#)  
Zygomaticomaxillary Complex Fracture, [2106](#)  
Zygomycosis, [2106](#)  
Zygote, [2106](#)  
Zymogen, [2106](#)  
Zymogen Granule, [2106](#)

---