

Sri Chaitanya IIT Academy.,India.

A.P T.S KARNATAKA TAMILNADU MAHARASTRA DELHI RANCHI

A right Choice for the Real Aspirant

ICON Central Office - Madhapur - Hyderabad

Sec: Sr. XL-500, Sr.S60(N&S), Elite, IAS & LIIT **Paper -2(Adv-2019-P2-Model)** **Date: 29-03-2026**

Time: 02.30Pm to 05.30Pm **GTA-21** **Max. Marks: 186**

29-03-2026_Sr. XL-500, Sr.S60(N&S), Elite, IAS & LIIT_Jee-Adv(2019-P2)_GTA-21_Syllabus,

PHYSICS : TOTAL SYLLABUS

CHEMISTRY : TOTAL SYLLABUS

MATHEMATICS : TOTAL SYLLABUS

Rankers Academy JEE

Name of the Student: _____ **H.T. NO:** | | | | | | | |

Sri Chaitanya Educational Institutions
Infinity Learn
41



For More Join: @IITJEE_Advanced

JEE-ADVANCE-2019-P2-Model

Time: 3:00Hr's **IMPORTANT INSTRUCTIONS** **Max Marks: 186**

PHYSICS:

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 1 – 8)	Questions with Multiple Correct Choice **+1 partial marks**	4	-1	8	32
Sec – II(Q.N : 9 – 14)	Questions with Numerical Value Answer Type	3	0	6	18
Sec – III(Q.N : 15 – 18)	Matching Type	3	-1	4	12
Total				**18**	**62**

CHEMISTRY:

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 19 – 26)	Questions with Multiple Correct Choice **+1 partial marks**	4	-1	8	32
Sec – II(Q.N : 27 – 32)	Questions with Numerical Value Answer Type	3	0	6	18
Sec – III(Q.N : 33 – 36)	Matching Type	3	-1	4	12
Total				**18**	**62**

MATHEMATICS:

Section	Question Type	+Ve Marks	- Ve Marks	No.of Qs	Total marks
Sec – I(Q.N : 37 – 44)	Questions with Multiple Correct Choice **+1 partial marks**	4	-1	8	32
Sec – II(Q.N : 45 – 50)	Questions with Numerical Value Answer Type	3	0	6	18
Sec – III(Q.N : 51 – 54)	Matching Type	3	-1	4	12
Total				**18**	**62**

Sri Chaitanya Educational Institutions
Infinity Learn
41


For More Join: @IITJEE_Advanced

PHYSICS **Max Marks: 62**


SECTION – I
(ONE OR MORE CORRECT ANSWER TYPE)

This section contains 8 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which ONE OR MORE than ONE option can be correct. **Marking scheme: +4 for all correct options & +1 partial marks, 0 if not attempted and -1 in all wrong cases**

Section 1 (Max Marks: 32)

- Section 1 contains eight questions.
- Each Question has Four Options and Only One of these four will be the correct answer.
- For each question, choose the option corresponding to the correct answer
- The Marking scheme to evaluate the Answer to each question:
  Full Marks: +4 (If the answer is correct)
  Partial Marks: +3 (If all the four options are correct but only 3 are chosen)
  Partial Marks: +2 (If THREE or More options are correct but only 2 correct answers are chosen)
  Partial Marks: +1 (If TWO or More options are correct but only 1 correct answer is chosen)
  Zero Marks: 0 (If the question is unanswered)
  Negative Marks: -1 (In all other cases)

1. Consider a satellite 'S' which is just above the Brazil and rotating on equatorial plane of earth of orbital radius r_0. The sense of rotation of earth and satellite are similar means both are rotating from west to east. The time period of satellite is 6 hours. At $t = 0$, satellite is just above the Brazil. Now choose the correct option(s).



A) The time after which the satellite will be above the Brazil again is 8 hours.


B) The above satellite may be a geostationary satellite.

C) The height of geostationary satellite above the sea level is 35,800 km approximately.

D) If the radius of orbit of the satellite 'S' is changed to $4r_0$, then the time after which the satellite will be above the Brazil again is 24 hours.

2. A concave mirror of focal length 3 meter is moving in the x-y plane with a velocity $\left(4\hat{i} + 4\hat{j}\right) m/s$. A point source '$S$' is also moving in the x-y plane with a velocity 10 m/s as shown in the figure. At a given instant, the object is at 5 m along the negative x-axis from pole and 20 cm along positive y-axis from the pole. The pole is assumed to be origin at this instant. Assuming paraxial approximation, choose the correct option(s).

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced



A) Velocity of image with respect to mirror at the same instant is $\left(-9\hat{i}-\frac{18}{5}\hat{j}\right)m/s$

B) Velocity of image with respect to mirror at the same instant is $\left(-18\hat{i}-\frac{51}{5}\hat{j}\right)m/s$

C) Velocity of image with respect to ground at the same instant is $\left(-5\hat{i}+\frac{2}{5}\hat{j}\right)m/s$

D) Velocity of image with respect to source at the same instant is $\left(-13\hat{i}-\frac{28}{5}\hat{j}\right)m/s$

3. The position vector of a particle of mass m is given by the following equation $\vec{r}(t)=\alpha t^3\hat{i}+\beta t^2\hat{j}$, where $\alpha=\frac{10}{3}m/s^3, \beta=5m/s^2$ and $m=0.1kg$. At $t=1s$, which of the following statement(s) is(are) true about the particle ?

A) The velocity $\vec{v}$ is given by $\vec{v}=\left(10\hat{i}+10\hat{j}\right)ms^{-1}$.


B) The angular momentum $\vec{L}$ with respect to the origin is given by $\vec{L}=-\left(\frac{5}{3}\right)\hat{k}Nms$.

C) The force $\vec{F}$ is given by $\vec{F}=\left(\hat{i}+2\hat{j}\right)N$.


D) The torque $\vec{\tau}$ with respect to the origin given by $\vec{\tau}=-\left(\frac{20}{3}\right)\hat{k}Nm$.

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1 IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced


4. Figure shows a LCR circuit, in which capacitor is charged, and at $t = 0$ switch is closed. Initially there is no energy stored in inductor. We draw graph $V_C(t)$ *vs* time for different value of resistance, which graph is/are nearly correct ?




A)




B)



C)



D)



5. A man standing on the edge of a swimming pool looks at a stone lying at the bottom. The depth of the swimming pool is equal to h. If the line of vision makes an angle θ with the normal to the surface, then which of the following is/are correct? (μ is refractive index of water).

A) Distance from the surface of water to the image of the stone formed is $\frac{h}{\mu}$, if angle θ is very small


B) Distance from the surface of water to the image of the stone formed is $h\mu$, if angle θ is very small

C) Distance from the surface of water to the image of the stone formed is $\frac{h\mu^3\cos^3\theta}{\left\{\mu^2-\sin^2\theta\right\}^{3/2}}$, if angle θ is large

D) Distance from the surface of water to the image of the stone formed is $\frac{h\mu^2\cos^3\theta}{\left\{\mu^2-\sin^2\theta\right\}^{3/2}}$, if angle θ is large.

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced

6. Four balls of mass m, then another four balls of M and then one more ball of mass m are strung on a horizontal, frictionless, fixed rod, as shown in the figure. The balls are close to each other, they are made of perfectly elastic material and $m < M$. From the left another ball of mass m is coming at a speed of v and collides with the first one of the ball string.



A) The final velocity of ball 1 is zero.

B) The final velocity of ball 5 is zero.

C) The final velocity of ball 9 is $\frac{m(M-m)v}{(M+m)^2}$.

D) The final velocity of ball 10 is $\frac{4Mmv}{(M+m)^2}$.

7. The powder charge of a cartridge supplies the bullet with a definite amount of energy. The barrel of the pistol can be imagined as a tube with an inner radius R. We mount the pistol so that the horizontal axis of the barrel passes through the center of the target. If the barrel is smooth inside, the bullet will hit the target at a distance h_1 below its center. For better flight properties of the bullet the barrel is rifled inside so that the bullet spins before leaving the barrel. Before leaving the barrel, the bullet makes N revolutions around its horizontal axis. If we carry out the same experiment with a rifled barrel, as we did the smooth one, the bullet will strike the target at a distance h_2 below the center of the target. The moment of inertia of bullet about its axis is kmR^2.

A) The length of the pistol barrel is $\sqrt{k}\,2\pi NR\sqrt{\frac{h_1}{h_2-h_1}}$.


B) The length of the pistol barrel is $\sqrt{k}\,2\pi NR\frac{h_1}{(h_2-h_1)}$.

C) The fraction of energy gained from combustion, consumed in spinning the bullet is $\frac{h_1}{h_2}$.

D) The fraction of energy gained from combustion, consumed in spinning the bullet is $1-\frac{h_1}{h_2}$.

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced

8. Moment of inertia of an elliptical disc of semi-axes a & b having mass m about



A) The major axis is $\frac{mb^2}{4}$.

B) The major axis is $\frac{ma^2}{4}$.

C) The minor axis is $\frac{mb^2}{4}$.

D) The minor axis is $\frac{ma^2}{4}$.

SECTION – II
(NUMERICAL VALUE TYPE)

This section contains 6 questions. Each question is numerical value type. For each question, enter the correct numerical value **(e.g. 003.2343, 6.250023, 7.000000, -0.33, -.30, 30.27000, -127.0030).**

Marking scheme: +3 for correct answer, 0 if not attempted and 0 in all other cases. Section 2 (Max Marks: 18)


- Section two contains six (6) questions. The answer to each question is a NUMERICAL VALUE
- For each question enter the correct numerical value of the answer using the mouse and the on-screen virtual numerical keypad in the place designated to enter the answer. If the numerical values has more than two decimal places, truncate/round off the value to TWO decimal places.

Answer to each question will be evaluated according to the following marking scheme

Full Marks: +3 (If only the correct numerical value is entered)

Zero Marks: 0 (In all other cases)


9. Two identical cars, each of mass m, connected by an undeformed spring of stiffness k, move without friction along a straight horizontal rail track with speed v_0 and collide with a stationary car of mass $\frac{m}{2}$. The collision is perfectly elastic.



The maximum compression of spring is $\sqrt{\frac{\beta m v_0^2}{k}}$. The value of β is

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1 IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced


10. Two identical shafts are constructed in the form of three concentric cylinders with radii equal to R, 2R and 3R and the distance between their centeres is $L = 3\pi R$. Both shafts rotate with the same rotation period T_0, and three belts are fastened to the shafts as shown in the figure. On each belt, there is a mark that, at the instant $t = 0$, is aligned with reference point O. Assuming that the belts rotate without slipping on the shafts, what is the smallest time (in multiple of T_0) after which the three marks are again aligned with the reference point O ?



11. A nucleus X, initially at rest decays into a nucleus Y with the emission of an α particle and energy Q is released. If m is mass of α particle and M is that of nucleus Y. The energy of the emitted α particle is given by $E_\alpha = \dfrac{QM}{M + xm}$. Find value of x.

12. An electron moving along x-axis is confined between two walls at x = 0 and x = L. there is no loss of energy during collision between electron and the wall. Initially the electron is in quantum state n = 2, and it jumps to quantum state n = 1. The emitted photon is incident on a metal having work function equal to energy of electron in quantum state n = 1. If maximum possible kinetic energy of emitted photoelectron is x times the energy of electron in quantum state n = 1, then find x.


13. Two current carrying wires of infinite length are placed perpendicular to x-y plane at $x = 0$ and $x = R\sqrt{3}$ as shown in figure. The circle has radius R. Current in both the wires is same and inwards to the x-y plane. Find ratio of $\left|\int \vec{B}.\vec{dl}\right|$ in segment CDA to segment ABC.



14. Consider a uniform charged sphere of charge density ρ and radius R. A cavity of radius R/2 is made inside the sphere and the removed part is again placed near the charge distribution as shown in the figure. If the electric field at the centre of the cavity is $\dfrac{\rho R}{n\varepsilon_0}$, find n.

Sri Chaitanya Educational Institutions
Infinity Learn by Sri Chaitanya
41


For More Join: @IITJEE_Advanced



SECTION-III
(MATCHING LIST TYPE)

This section contains 4 questions, each having two matching lists (List-1 & List-II). The options for the correct match are provided as (A), (B),(C) and (D) out of which ONLY ONE is correct.

Marking scheme: +3 for correct answer, 0 if not attempted and -1 in all other cases.

Section 3 (Max Marks: 12)

- This Section contains two (02) List-Match sets.
- Each List-Match set has TWO (02) Multiple-choice Questions
- Each List-Match set has two lists: List 1 and List 2
- List 1 has four entries (I), (II), (III), (IV) and List 2 has six entries (P), (Q), (R), (S), (T) and (U)
- Four options are given in each multiple choice question based on List 1 and List 2 and Only ONE of these four options satisfies the condition asked in the Multiple Choice Question.

Answer to each question will be evaluated according to the following marking scheme

Full Marks: +3 (If only the option corresponding to the correct combination is chosen))

Zero Marks: 0 (If none of the options is chosen (i.e. the question is unanswered)

Negative Marks: -1 (In all other cases)

Answer Q.15 and Q.16 by appropriately matching the list based on the information given in the paragraph

A musical instrument is made using four different metal strings, 1, 2, 3 and 4 with mass per unit length $\mu, 2\mu, 3\mu$ and 4μ respectively. The instrument is played by vibrating the strings by varying the free length in between the range L_0 and $2L_0$. It is found that in string – 1 (μ) at free length L_0 and tension T_0 the fundamental mode frequency is f_0. List – I gives the above four strings while List – II lists the magnitude of some quantity.

	List-I		List-II
(I)	String – 1 (μ)	(P)	1
(II)	String – 2(2μ)	(Q)	1/2
(III)	String – 3(3μ)	(R)	$1/4$
(IV)	String – 4 (4μ)	(S)	$1/2\sqrt{3}$
		(T)	3/16
		(U)	1/16
		(V)	$\frac{1}{2\sqrt{2}}$

15. If the tension in each strings is T_0, the correct match for the lowest fundamental frequency in f_0 units will be,

A) I – P, II – R, III – S, IV – Q **B)** I – Q, II – V, III – S, IV – R

C) I – P, II – Q, III – T, IV – S **D)** I – Q, II – P, III – R, IV – V

Sri Chaitanya Educational Institutions Infinity Learn 41

THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1 IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023

JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1

JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1

NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1

For More Join: @IITJEE_Advanced

16. The length of the string 1, 2, 3 and 4 are kept fixed at $L_0, \frac{3L_0}{2}, \frac{5L_0}{4}$ and $\frac{7L_0}{4}$, respectively. Strings 1, 2, 3 and 4 are vibrated at their $1^{st}, 3^{rd}, 5^{th}$ and 14th harmonics, respectively such that all the strings have same frequency. The correct match for the tension in the four strings in the units of T_0 will be.

A) I – P, II – Q, III – R, IV – T **B)** I – T, II – Q, III – R, IV – U

C) I – P, II – R, III – T, IV – U **D)** I – P, II – Q, III – T, IV – U

Answer Q.17 and Q.18 by appropriately matching the list based on the information given in the paragraph


In a thermodynamic process on an ideal monatomic gas, the infinitesimal heat absorbed by the gas is given by $T\Delta X$, where T is temperature of the system and ΔX is the infinitesimal change in a thermodynamic quantity X of the system. For a mole of monatomic ideal gas $X = \frac{3}{2}R\ln\left(\frac{T}{T_A}\right) + R\ln\left(\frac{V}{V_A}\right)$. Here, R is gas constant, V is volume of gas. T_A and V_A are constants.

The List – I below gives some quantities involved in a process and List – II gives some possible values of these quantities.

List-I		List-II	
(I)	Work done by the system in process $1 \to 2 \to 3$	(P)	$\frac{1}{3}RT_0\ell n2$
(II)	Change in internal energy in process $1 \to 2 \to 3$	(Q)	RT_0
(III)	Heat absorbed by the system in process $1 \to 2 \to 3$	(R)	$3RT_0$
(IV)	Heat absorbed by the system in process $1 \to 2$	(S)	$4RT_0$
		(T)	$\frac{1}{3}RT_0(3+\ell n2)$
		(U)	$\frac{5}{2}RT_0$

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced

17. If the process carried out on one mole of monoatomic ideal gas is as shown in figure in the PV-diagram with $P_0V_0 = RT_0$,




the correct match is

A) I-Q; II-S; III-R;IV-U **B)** I-Q; II-R; III-S; IV-U

C) I-Q; II-R; III-P; IV-U **D)** I-S; II-R;III-Q; IV-T

18. If the process is one mole of monoatomic ideal gas is as shown in figure in the TV-diagram with $P_0V_0 = \frac{1}{3}RT_0$,



the correct match is

A) I-P; II-T; III-Q; IV-T **B)** I-S; II-T; III-Q;IV-U

C) I-P; II-Q;III-T; IV-P **D)** I-P; II-R; III-T; IV-S

Rankers Academy JEE

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced

CHEMISTRY **Max Marks: 62**

SECTION – I
(ONE OR MORE CORRECT ANSWER TYPE)

This section contains 8 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which ONE OR MORE than ONE option can be correct. **Marking scheme: +4 for all correct options & +1 partial marks, 0 if not attempted and -1 in all wrong cases**

Section 1 (Max Marks: 32)

- Section 1 contains eight questions.
- Each Question has Four Options and Only One of these four will be the correct answer.
- For each question, choose the option corresponding to the correct answer
- The Marking scheme to evaluate the Answer to each question:
  Full Marks: +4 (If the answer is correct)
  Partial Marks: +3 (If all the four options are correct but only 3 are chosen)
  Partial Marks: +2 (If THREE or More options are correct but only 2 correct answers are chosen)
  Partial Marks: +1 (If TWO or More options are correct but only 1 correct answer is chosen)
  Zero Marks: 0 (If the question is unanswered)
  Negative Marks: -1 (In all other cases)

19. Among the following correct statement is/are


A) Metal hydroxides like $Mg(OH)_2$ and $Al(OH)_3$ are better antacids than $NaHCO_3$


B) Cimetidine can cure the reason for acidity but $Al(OH)_3$ cannot cure the reason for acidity.


C) The interaction of histidine with receptor protein in the stomach walls is the reason for production acid in the stomach.


D) Cimetidine is agonist drug.

20. Which of the following on reaction with HBr gives $2R,3S-2,3-$Dibromobutane as the major product.

A)


B)


C)


D)


21. 2.5 g of salt of weak acid (NaA) (Molecular weight = 50) is dissolved in 100 ml water and titrated with 0.5 M HCl at 25°C. The pH was found to be 9 after addition of 20ml of HCl solution (log 2 = 0.3). Which of the following statements are correct?

A) K_a of acid is 10^{-9}.

B) pH at equivalence point is 4.5

C) Volume of HCl used at equivalence point is 100 ml.

D) pH after addition of 101 ml of HCl solution is nearly 2.6.

Sri Chaitanya Educational Institutions
Infinity Learn
41


For More Join: @IITJEE_Advanced

22. The correct information regarding a solid element 'A' (atomic radius = r) having ABCABC…. Type packing (FCC unit cell edge length = a) is


A) The distance between nearest octahedral and tetrahedral void is $\frac{\sqrt{3}a}{4}$.

B) The distance between two nearest octahedral void is $\frac{a}{\sqrt{2}}$.

C) The distance between two nearest tetrahedral void is $\frac{\sqrt{3}a}{2}$.

D) If anions X^- are placed in all tetrahedral voids and cations P^{2+} in all octahedral voids and all the atoms of A are removed the resulting PX_2 lattice has the same structure as CaF_2 lattice.

23.



A) A, B and C are $Mg; MgH_2; Mg(OH)_2$

B) D, E and F are $MgCO_3; Mg(HCO_3)_2; MgO$

C) G and H are $CaC_2, CaCN_2$

D) Equal volume of mixture of 0.1 M solution of D and E form buffer solution

24. Which of the following is/are correct?

A) Leaching is the method to remove quantitatively Fe_2O_3 from red bauxite

B) In the metallurgy of Aluminium, purified Al_2O_3 is mixed with Na_3AlF_6 or CaF_2 which lowers the melting point of the mix and also increases conductivity

C) Metallurgy of gold involves hydrometallurgical process involving soluble complex formation and its convenient reduction

D) Self reduction is not possible during metallurgy of Mg

Sri Chaitanya Educational Institutions
Infinity Learn
41


For More Join: @IITJEE_Advanced

25. $2CH_3Cl + Si \xrightarrow[570\,K]{Cu\ powder} A(major)$

$3CH_3MgCl + Si \rightarrow B + 3MgCl_2$

$CH_3MgCl + SiCl_4 \rightarrow D + MgCl_2$

Which of the following are correct ?

A) The compound 'A' on hydrolysis followed by polymerization give linear or cyclic silicones.

B) The chain length of linear silicone can be controlled by adding the compound 'B'.

C) The compound 'D' on hydrolysis followed by polymerization give cross-linked silicone.

D) The compound 'D' can be used to control the chain length of linear silicone.

26. Which of the following is/are correct

A) $H_2 > B_2 > F_2 > Li_2$ order of bond energy

B) $O-O$ bond length in KO_2 will be smaller than Na_2O_2

C) $O-O$ bond length in $O_2[PtF_6]$ will be more than KO_2

D) $N-O$ bond length in NO gaseous molecule will be smaller than in NO^+ gaseous ion

SECTION – II
(NUMERICAL VALUE TYPE)

This section contains 6 questions. Each question is numerical value type. For each question, enter the correct numerical value **(e.g. 003.2343, 6.250023, 7.000000, -0.33, -.30, 30.27000, -127.0030).**

Marking scheme: +3 for correct answer, 0 if not attempted and 0 in all other cases. Section 2 (Max Marks: 18)


- Section two contains six (6) questions. The answer to each question is a NUMERICAL VALUE
- For each question enter the correct numerical value of the answer using the mouse and the on-screen virtual numerical keypad in the place designated to enter the answer. If the numerical values has more than two decimal places, truncate/round off the value to TWO decimal places.

Answer to each question will be evaluated according to the following marking scheme

Full Marks: +3 (If only the correct numerical value is entered)


Zero Marks: 0 (In all other cases)

27. For the following compound, number of enantiomers possible are is/are X, number of diasteromers possible is/are Y. $(X+Y)$ value is:





Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced

28.



In this reaction among the following how many are more reactive than benzenediazonium chloride?





Rankers Academy JEE

29. The standard molar enthalpies of formations of $H_2O(l)$ and $H_2O_2(l)$ are – 286 and – 188 kJ/mol, respectively. Molar enthalpies of vaporization of H_2O and H_2O_2 are 44 and 53 kJ, respectively. The bond dissociation enthalpy of $O_2(g)$ is 498 kJ/mol. Calculate the bond dissociation enthalpy (in kJ/mol) of $O-O$ bond in H_2O_2, assuming that the bond dissociation enthalpy of $O-H$ bond is same in both H_2O and H_2O_2.

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced

30. Equal masses of two aqueous solutions, one of 3.6 g glucose per 100 g of water and the other of 3.6 g urea per 100 g of water are mixed. If the freezing point of resulting solution is $-x°C$, then the value of '$1000x$' is (K_f of water = 1.86 kgKmol^{-1})

31. Consider the following oxides of nitrogen

$N_2O, NO, N_2O_3, NO_2, N_2O_4, N_2O_5$

Find the value of $\frac{(x+y+z+w)}{3}$

Where x = Total no. of colorless compound (s)

y = Total no. of acidic compound(s)

z = Total no.of gaseous compound(s) at room temperature

w = Total no. of compounds(s) which is/are Diamagnetic

32. Among the following the number of sulphides which are not soluble in yellow ammonium sulphide is $As_2S_3, Sb_2S_3, CuS, SnS_2, CdS, PbS$

SECTION-III
(MATCHING LIST TYPE)

This section contains 4 questions, each having two matching lists (List-1 & List-II). The options for the correct match are provided as (A), (B),(C) and (D) out of which ONLY ONE is correct.
Marking scheme: +3 for correct answer, 0 if not attempted and -1 in all other cases.
Section 3 (Max Marks: 12)












- This Section contains two (02) List-Match sets.
- Each List-Match set has TWO (02) Multiple-choice Questions
- Each List-Match set has two lists: List 1 and List 2
- List 1 has four entries (I), (II), (III), (IV) and List 2 has six entries (P), (Q), I, (S), (T) and (U)
- Four options are given in each multiple choice question based on List 1 and List 2 and Only ONE of these four options satisfies the condition asked in the Multiple Choice Question.

Answer to each question will be evaluated according to the following marking scheme
Full Marks: +3 (If only the option corresponding to the correct combination is chosen))
Zero Marks: 0 (If none of the options is chosen (i.e. the question is unanswered)
Negative Marks: -1 (In all other cases)

Answer Q.33 and Q.34 by appropriately matching the list based on the information given in the paragraph

In the List – I some sequential reactions given. In the List – II intermediate products and final products formed in those sequential reactions given.

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced

LIST – I		LIST – II	
I)		P)	
II)		Q)	
III)		R)	
IV)		S)	
		T)	
		U)	
		V)	

Rankers Academy JEE
Sri Chaitanya Education INDIA

Sri Chaitanya Educational Institutions
Infinity Learn
41


For More Join: @IITJEE_Advanced

33. Which one of the following options has correct combination considering **LIST – I** and **LIST – II** ?

A) I – RQ **B)** II – PR **C)** I – P **D)** II – QR

34. Which one of the following options has correct combination considering **LIST – I** and **LIST – II** ?

A) IV – TU **B)** III – TV **C)** IV – ST **D)** III – S

Answer Q.35 and Q.36 by appropriately matching the list based on the information given in the paragraph

Consider the reactions A $\rightarrow$ products.

Column I (Graph)		Column II (Magnitude of slope)		Column III (Intercept)	
I)	A^2 is plotted against time 't' in zero – order reaction	**P)**	$\frac{k}{2.303}$	**U)**	${A_0}^2$
II)	$\log A$ is plotted against 't' in first-order reaction	**Q)**	$\frac{2k}{A_0}$	**V)**	$\frac{1}{A_0}$
III)	$\frac{1}{A}$ is plotted against 't' in second order reaction	**R)**	$2kA_0$	**W)**	$\frac{1}{{A_0}^2}$
IV)	$\frac{1}{A^2}$ is plotted against 't' in second order reaction	**S)**	k	**X)**	$\log(A_0)$

Here A_0 = initial concentration of reactant, A = concentration at time 't', k = rate constant for the reaction.

35. Which of the following options is correct, if slope is measured at $t = 0$

A) I – SU **B)** II – RX **C)** III – PW **D)** IV – QW

36. How many of the following are incorrect if slope is measured at $t = \frac{2A_0}{k}$

i)) I – RU ii) II – PX iii) III – SV iv) IV – QW

A) 1 **B)** 2 **C)** 3 **D)** 4

Sri Chaitanya Educational Institutions
Infinity Learn
41


For More Join: @IITJEE_Advanced

MATHEMATICS

Max Marks: 62

SECTION – I
(ONE OR MORE CORRECT ANSWER TYPE)

This section contains 8 multiple choice questions. Each question has 4 options (A), (B), (C) and (D) for its answer, out of which ONE OR MORE than ONE option can be correct. **Marking scheme: +4 for all correct options & +1 partial marks, 0 if not attempted and -1 in all wrong cases**

Section 1 (Max Marks: 32)

- Section 1 contains eight questions.
- Each Question has Four Options and Only One of these four will be the correct answer.
- For each question, choose the option corresponding to the correct answer
- The Marking scheme to evaluate the Answer to each question:
  Full Marks: +4 (If the answer is correct)
  Partial Marks: +3 (If all the four options are correct but only 3 are chosen)
  Partial Marks: +2 (If THREE or More options are correct but only 2 correct answers are chosen)
  Partial Marks: +1 (If TWO or More options are correct but only 1 correct answer is chosen)
  Zero Marks: 0 (If the question is unanswered)
  Negative Marks: -1 (In all other cases)

---

37. **For** $0<\alpha<\frac{3}{2}$, define $x_n=(n+1)^{\alpha}-n^{\alpha}$. For different values of α, $\lim\limits_{n\to\infty} x_n$ can take one or more than one finite value or the limit may be infinite. Let $a=$ Number of possible finite limit value(s) and (b,c) be the exhaustive set of values of α such that the limit tends to infinity. Then

A) $a=1$ **B)** $a=2$ **C)** $b+2c=4$ **D)** $b+2c=\frac{9}{2}$

38. Consider the system of three cubic equations

$$x^3+(\alpha+1)x^2+(\beta+3)x+(\gamma+2)=0$$

$$x^3+(\alpha+2)x^2+(\beta+1)x+(\gamma+3)=0$$

$$x^3+(\alpha+3)x^2+(\beta+2)x+(\gamma+1)=0$$

Each of the cubic equations has three distinct real roots, but the total number of distinct real roots of the system is 5. Using the above information, we can uniquely find out the real numbers α,β,γ. Then

A) $2\alpha+3=0$ **B)** $2\alpha-4\beta=11$ **C)** $2\beta-5\gamma=4$ **D)** $\gamma-2\alpha=1$

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
Rankers Academy JEE
For More Join: @IITJEE_Advanced

39. Consider a discrete frequency distribution

Variable	X_1	X_2	X_3	....	.....	....	X_n
Frequency	f_1	f_2	f_3	.....	.....	.....	f_n

Where $X_i, f_i \in \mathbb{R} \forall i = 1,2,3,....,n.$

If deviations of X_i from its mean M say 'x_i' are small compared to mean M such that $\left(\frac{x_i}{M}\right)^3$ and higher powers of $\frac{x_i}{M}$ can be neglected for $\forall i = 1,2,3,...,n.$

If $G =$ Geometric mean and b is standard deviation of the distribution, then

A) $G = M\left(1 - \frac{b^2}{M^2}\right)$ **B)** $G = M\left(1 - \frac{b^2}{2M^2}\right)$

C) $M^2 - b^2 = G^2$ **D)** $b^2 - M^2 = G^2$

40. A circle passing through the centre of a rectangular hyperbola cuts the curve at points A, B, C and D. Circumcircle of the triangle formed by tangents at A, B and C to the hyperbola is drawn.

A) The circumcircle passes through the point D.

B) The circumcircle passes through centre of the rectangular hyperbola.

C) Centre of the circumcircle is at D.

D) Centre of the circumcircle is at the other end of the chord drawn from D passing through centre of hyperbola.

41. If n_{f_i} = number of non-differentiabilities of function f_i in it's domain &

$$f_1(x) = \lim_{n\to\infty} \frac{n^x - n^{-x}}{n^x + n^{-x}}; x \in \mathbb{R}$$

$$f_2(x) = \lim_{n\to\infty} \sqrt[n]{4^n + x^{2n} + \frac{1}{x^{2n}}}; x \in \mathbb{R} - \{0\}$$

$$f_3(x) = \lim_{n\to\infty} \frac{\ell n\left(e^n + x^n\right)}{n}; x \geq 0$$

$$f_4(x) = \lim_{n\to\infty} \sqrt[2n]{\sin^{2n} x + \cos^{2n} x}; x \in (0, 2\pi), \text{ Then}$$

A) $n_{f_1} = 1$ **B)** $n_{f_2} = 5$ **C)** $n_{f_3} = 1$ **D)** $n_{f_4} = 2$

Sri Chaitanya Educational Institutions Infinity Learn 41



For More Join: @IITJEE_Advanced

42. Let L_1 be the line of intersection of the planes given by the equations $2x+3y+z=4$ and $x+2y+z=5$. Let L_2 be the line passing through the point $P(2,-1,3)$ and parallel to L_1. Let M denote the plane given by the equation $2x+y-2z=6$. Suppose that the line L_2 meets the plane M at the point Q. Let R be the foot of the perpendicular drawn from P to the plane M. Then which of the following statements is(are) TRUE?

A) The length of the line segment PQ is $9\sqrt{3}$.

B) The length of the line segment QR is 15.

C) The area of ΔPQR is $\frac{3}{2}\sqrt{234}$

D) The acute angle between the line segments PQ and PR is $\cos^{-1}\left(\frac{1}{2\sqrt{3}}\right)$

43. Let $\mathbb{N}$ denote the set of all natural numbers, and $\mathbb{Z}$ denote the set of all integers. Consider the functions $f:\mathbb{N}\to\mathbb{Z}$ and $g:\mathbb{Z}\to\mathbb{N}$ defined by

$$f(n)=\begin{cases}(n+1)/2 & \text{if } n \text{ is odd,}\\ (4-n)/2 & \text{if } n \text{ is even,}\end{cases} \text{ and } g(n)=\begin{cases}3+2n & \text{if } n\geq 0,\\ -2n & \text{if } n<0.\end{cases}$$

Define $(g\circ f)(n)=g(f(n))$ for all $n\in\mathbb{N}$, and $(f\circ g)(n)=f(g(n))$ for all $n\in\mathbb{Z}$.

Then which of the following statements is(are) TRUE?

A) $g\circ f$ is NOT one-one and $g\circ f$ is NOT onto

B) $f\circ g$ is NOT one-one but $f\circ g$ is onto

C) g is one-one and g is onto

D) f is NOT one-one but f is onto

44. Let $I=\begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix}$ and $P=\begin{pmatrix}2 & 0\\ 0 & 3\end{pmatrix}$. Let $Q=\begin{pmatrix}x & y\\ z & 4\end{pmatrix}$ for some non-zero real numbers x, y, and z, for which there is a 2×2 matrix R with all entries being non-zero real numbers, such that $QR=RP$. Then which of the following statements is (are) TRUE?

A) The determinant of $Q-2I$ is zero **B)** the determinant of $Q-6I$ is 12.

C) The determinant of $Q-3I$ is 15 **D)** $yz=2$.

Sri Chaitanya Educational Institutions
Infinity Learn
41


For More Join: @IITJEE_Advanced

SECTION – II
(NUMERICAL VALUE TYPE)

This section contains 6 questions. Each question is numerical value type. For each question, enter the correct numerical value **(e.g. 003.2343, 6.250023, 7.000000, -0.33, -.30, 30.27000, -127.0030).**

Marking scheme: +3 for correct answer, 0 if not attempted and 0 in all other cases. Section 2 (Max Marks: 18)

- Section two contains six (6) questions. The answer to each question is a NUMERICAL VALUE
- For each question enter the correct numerical value of the answer using the mouse and the on-screen virtual numerical keypad in the place designated to enter the answer. If the numerical values has more than two decimal places, truncate/round off the value to TWO decimal places.

Answer to each question will be evaluated according to the following marking scheme

Full Marks: +3 (If only the correct numerical value is entered)

Zero Marks: 0 (In all other cases)

45. Let $f:\mathbb{R}^+ \to \mathbb{R}$ be a differentiable function satisfying $f(y)-f(x)=\frac{x^x}{y^y}f\left(\frac{y^y}{x^x}\right)$ for all $x,y>0, f'(1)=1$. Find the value of $\frac{f\left(2022^2\right)}{1011 f(2022)}$.

46. If the locus of the foci of all conics which touch the four lines $x=\pm 5$ and $y=\pm 3$ is the conic $S=0$, then find the length of latus rectum of $S=0$.

47. Three distinct normals are drawn to the parabola $y^2=20x\cos\theta$ from any point lying on the straight line $y=4\sin\theta$. Find the eccentricity of the locus of the orthocenter of the triangle formed by the corresponding tangents $\left(\theta\in\left(0,\frac{\pi}{2}\right)\right)$

48. Consider the parabola $x^2-2x-4y+9=0$. Two chords of the parabola LOL' and MOM' are drawn passing through a point $O(1,5)$ on its axis. Then radical axis of the circles described on LL' and MM' as diameter passes through a fixed point $P(\alpha,\beta)$, where $2\alpha+3\beta=$____.

49. Let the set of all relations R on the set $\{a,b,c,d,e,f\}$, such that R is reflexive and symmetric, and R contains exactly 10 elements, be denoted by S. Then the number of elements in S is ____.

50. A circle contains the points $(0,11)$ and $(0,-11)$ on its circumference and all points (x,y) satisfying $x^2+y^2<1$ lies in the interior of the circle. Find the largest possible radius of the circle.

SECTION-III

Sri Chaitanya Educational Institutions
Infinity Learn
41


For More Join: @IITJEE_Advanced

(MATCHING LIST TYPE)

This section contains 4 questions, each having two matching lists (List-1 & List-II). The options for the correct match are provided as (A), (B),(C) and (D) out of which ONLY ONE is correct.

Marking scheme: +3 for correct answer, 0 if not attempted and -1 in all other cases.

Section 3 (Max Marks: 12)

- This Section contains two (02) List-Match sets.
- Each List-Match set has TWO (02) Multiple-choice Questions
- Each List-Match set has two lists: List 1 and List 2
- List 1 has four entries (I), (II), (III), (IV) and List 2 has six entries (P), (Q), I, (S), (T) and (U)
- Four options are given in each multiple choice question based on List 1 and List 2 and Only ONE of these four options satisfies the condition asked in the Multiple Choice Question.

Answer to each question will be evaluated according to the following marking scheme

Full Marks: +3 (If only the option corresponding to the correct combination is chosen))

Zero Marks: 0 (If none of the options is chosen (i.e. the question is unanswered)

Negative Marks: -1 (In all other cases)

Answer Q.51 and Q.52 by appropriately matching the list based on the information given in the paragraph

Let α,β and γ be real numbers. Consider the following system of linear equations

$$x+2y+z=7$$

$$x+\alpha z=11$$

$$2x-3y+\beta z=\gamma$$

Match each entry in **List-I** to the correct entries in **List-II**

List-I		List-II	
(P)	If $\beta=\frac{1}{2}(7\alpha-3)$ and $\gamma=28$, then the system has	(1)	A unique solution
(Q)	If $\beta=\frac{1}{2}(7\alpha-3)$ and $\gamma\neq 28$, then the system has	(2)	No solution
(R)	If $\beta\neq\frac{1}{2}(7\alpha-3)$ where $\alpha=1$ and $\gamma\neq 28$, then the system has	(3)	Infinitely many solutions.
(S)	If $\beta\neq\frac{1}{2}(7\alpha-3)$ where $\alpha=1$ and $\gamma=28$, then the system has	(4)	$x=11, y=-2$ and $z=0$ as a solution.
		(5)	$x=-15, y=4$ and $z=0$ as a solution.
		(6)	Trivial Solution

51. Which of the following is the only CORRECT combination?

Sri Chaitanya Educational Institutions
Infinity Learn
41
THE PERFECT HAT-TRICK WITH ALL- INDIA RANK 1
IN JEE MAIN 2023 JEE ADVANCED 2023 AND NEET 2023
JEE MAIN 2023 SINGARAJU VENKAT KOUNDINYA RANK 1
JEE Advanced 2023 VAVILALA CHIDVILAS REDDY RANK 1
NEET 2023 BORA VARUN CHAKRAVARTHI RANK 1
For More Join: @IITJEE_Advanced

A) $P \to 4$ **B)** $Q \to 3$ **C)** $R \to 1$ **D)** $S \to 5$

52. Which of the following is the only CORRECT combination?

A) $P \to 2$ **B)** $Q \to 2$ **C)** $R \to 4$ **D)** $S \to 6$

Answer Q.53 and Q.54 by appropriately matching the list based on the information given in the paragraph

Let $\gamma \in R$ be such that the lines $L_1 : \frac{x+11}{1} = \frac{y+21}{2} = \frac{z+29}{3}$ and $L_2 : \frac{x+16}{3} = \frac{y+11}{2} = \frac{z+4}{\gamma}$ intersect. Let R_1 be the point of intersection of L_1 and L_2.

Let $O = (0,0,0)$, and $\hat{n}$ denote a unit normal vector to the plane containing both the lines L_1 and L_2.

Match each entry in **List-I** to the correct entries in **List-II**

List-I		List-II	
(P)	γ **equals**	(1)	$-\hat{i} - \hat{j} + \hat{k}$
(Q)	A possible choice for $\hat{n}$ **is**	(2)	$\sqrt{\frac{3}{2}}$
(R)	$\overrightarrow{OR_1}$ **equals**	(3)	1
(S)	A possible value of $\overrightarrow{OR_1} \cdot \hat{n}$ **is**	(4)	$\frac{1}{\sqrt{6}}\hat{i} - \frac{2}{\sqrt{6}}\hat{j} + \frac{1}{\sqrt{6}}\hat{k}$
		(5)	$\sqrt{\frac{2}{3}}$
		(6)	$-\frac{2}{\sqrt{14}}\hat{i} + \frac{3}{\sqrt{14}}\hat{j} - \frac{1}{\sqrt{14}}\hat{k}$

53. Which of the following is the only CORRECT combination?

A) $P \to 2$ **B)** $Q \to 6$ **C)** $R \to 4$ **D)** $S \to 5$

54. Which of the following is the only CORRECT combination?

A) $P \to 5$ **B)** $Q \to 4$ **C)** $R \to 6$ **D)** $S \to 2$

Sri Chaitanya Educational Institutions
Infinity Learn
41


For More Join: @IITJEE_Advanced

Sri Chaitanya Educational Institutions | Infinity Learn

41 Years of Excellence 1986-2026

LEADING BY MILES SRI CHAITANYA DOMINATES

JEE ADVANCED 2025

29 Ranks in Top 100 in All-India Open Category



4 Students in Top 11 in JEE-Advanced 2025, All India Open Category

Rankers Academy JEE

16 RANK DEVDUTTA MAJHI HT. No. 255053116*	18 RANK DHARMANA GNANA RUTVIK SAI HT. No. 256055278	19 RANK VANGALA AJAY REDDY HT. No. 256131009	23 RANK AKSH GOGI HT. No. 252071075*	26 RANK P HEMA SAI SURYA KARTHIK HT. No. 256033006	27 RANK SARKARSAMUDRA HT. No. 252071105*
30 RANK OM PRAKASH BEHERA HT. No. 252021018*	32 RANK SUNKARA SAI RISHANTH REDDY HT. No. 256165327	34 RANK DHRUBA JYOTHI PANJA HT. No. 252048248*	35 RANK BHAVESH JAYANTHI HT. No. 251043080	36 RANK ADVAY MAYANK HT. No. 252104113*	37 RANK KARMANYA GUPTA HT. No. 252081477*
42 RANK MD ANAS HT. No. 252046210*	45 RANK RAMIT GOYAL HT. No. 257001113*	52 RANK MAULIK JAIN HT. No. 252079407*	54 RANK GARV HT. No. 252056188*	59 RANK LARISSA HT. No. 252079071*	60 RANK ARYAN BALABADRULA HT. No. 256132077
63 RANK SAMYAJYOTI BISWAS HT. No. 255058456*	64 RANK AARUSH ANAND HT. No. 251006176*	72 RANK RUSHMITH BANDARI HT. No. 256168048	78 RANK KORIKANA RASAGNYA HT. No. 256057046	87 RANK LAKSHYA SHARMA HT. No. 252070079*	91 RANK AVANEESH BANSAL HT. No. 251113130*	95 RANK KAVYA AGGARWAL HT. No. 252079121*

BELOW 100 ALL INDIA OPEN CATEGORY RANKS: 29	BELOW 500 ALL INDIA OPEN CATEGORY RANKS: 113	BELOW 1000 ALL INDIA OPEN CATEGORY RANKS: 205	BELOW 1000 ALL INDIA CATEGORY RANKS COUNT: 745	NUMBER OF QUALIFIED RANKS: 4,212

040 66 15 15 15 www.srichaitanya.net

For More Join: @IITJEE_Advanced