SOLUTION

SECTION-I

Single Correct Answer Type **15 Q. [3 M (–1)]**

1. Ans. (C)

2. Ans. (B)

3. Ans. (D)

4. Ans. (B)

5. Ans. (A)

6. Ans. (B)

7. Ans. (C)

8. Ans. (B)

9. Ans. (B)

Sol. Equation of demped oscillation

$$x=\left[Ae^{-b/2mt}\right]\sin\omega t$$

$$I=\left[I_0e^{-\frac{R}{2L}t}\right]\sin\omega t$$

$I_0e^{-(R/2L)t}=\frac{I_0}{e}$ For maximum

$\frac{R}{2L}t=1$ $\phi=\omega t$

$t=\frac{2L}{R}$ $\phi=\frac{2L\omega}{R}\left(X_L=\omega L\right)$

$(2\pi)n=\frac{2\times200}{20}=n=\frac{10}{\pi}$

10. Ans. (D)

Sol. $\omega'=\sqrt{\frac{K}{m}-\frac{b^2}{4m^2}}$ and $\omega'=\frac{2\pi}{T}$

11. Ans. (D)

Sol. This is the case of Damped oscillation.

12. Ans. (C)

Sol. Considering gravity, buoyant force and viscouse force oscillation will be damped.

$\Omega_1^2=\omega_2^2-C$, ω_2 is natural frequency.

$\Rightarrow T_1>T_2$ C is a constant.

Rankers Academy JEE

For More Material Join: @JEEAdvanced_2024

13. Ans. (B)

14. Ans. (C)

15. Ans. (B)

Sol. $T_0 = 2\pi\sqrt{\frac{m}{k}}$

$= \frac{2\pi}{\sqrt{10}}$

$A = A_0 e^{-t/\gamma}$

$\therefore$ for $A = \frac{A_0}{e}$, $t = \gamma$

$t = \gamma = \frac{2m}{b} = \frac{2m}{\frac{B^2\ell^2}{R}} = 10^4 s$

$\therefore$ No of oscillation $\frac{t}{T_0} = \frac{10^4}{2\pi/\sqrt{10}} \approx 5000$.

Multiple Correct Answer Type **2 Q. [4 M (–1)]**

16. Ans. (A,B,D)

17. Ans. (A,D)

Linked Comprehension Type **(1 Para × 2Q.) [3 M (-1)]**

(Single Correct Answer Type)

18. Ans. (A)

19. Ans. (D)

SECTION-II

Linked Comprehension Type **(1 Para × 2Q.) [3 M (-1)]**

(upto second decimal place)

20. Ans. 50.00

21. Ans. 30.00

Numerical Answer Type Question **1Q.[3(0)]**

(upto second decimal place)

22. Ans. 0.4

Rankers Academy JEE

For More Material Join: @JEEAdvanced_2024