
SOLUTION
SECTION-I

Single Correct Answer Type **5 Q. [3 M (–1)]**

1. Ans. (B)

Sol. $F\cos 37 = \mu\left(1000 - \frac{3F}{5}\right)$



$\frac{4}{5}F = \frac{1000}{3} - \frac{F}{5}$

$F = \frac{1000}{3}$

$\frac{1000}{3} = 25(g + a)$

$a = \frac{10}{3}$

2. Ans. (B)

Sol. Tension = T – mg = ma

T = m(g + a)

T = 10(10 + 2)

T = 10 × 12 = 120

Upper support 120 + 80 = 200


(80) weight of latter monkey

3. Ans. (A)

Sol. Acceleration of C in horizontal direction = acceleration of A=a (right direction) = $a\hat{i}$

Accelration of C in vertical downward direction = $2(a+b)\hat{j}$

Hence, net acceleration of $C = a\hat{i} - 2(a+b)\hat{j}$ (see figure)



From length constraint $\ell_1 + \ell_2 + \ell_3 + \ell_4 = C$

$\ddot{\ell}_1 + \ddot{\ell}_2 + \ddot{\ell}_3 + \ddot{\ell}_4 = 0$ –a –b + 0 –a –b + c = 0


c = 2a + 2b

From wedge constraints, acceleration of C towards right side is a. Acceleration of C w.r.t. ground $= a\hat{i} - 2(a+b)\hat{j}$

For More Material Join: @JEEAdvanced_2025

4. Ans. (A)

Sol.



$N = F\cos\theta$

$F\sin\theta = \mu N$

$$\frac{\sin\theta}{\cos\theta} = \mu$$


$\tan\theta = 0.9$

$\theta = \tan^{-1}(0.9)$


5. Ans. (A)

Sol. Initially $m_1 g = kx$

When support is removed, spring force does not change.



New FBD



For m_1 : $m_1 g - kx = m_1 a_1$ $\Rightarrow a_1 = 0$

For m_2 : $m_2 g + kx = m_2 a_2$ $\Rightarrow a_2 = \dfrac{(m_1 + m_2)g}{m_2}\downarrow$

Multiple Correct Answer Type 5 Q. [4 M (–1)]

6. Ans. (A,B)

Sol. Man $N + T = mg$

$N + T = 500$ .....(i)

Weghing machine $T = mg + N$

$T - N = mg$

$T - N = 200$ ....(ii)



$N + T = 500$

$T - N = 200$

$2T = 700$

$T = 350$

$N + 350 = 500$


$N = 150$

Reading 15 kg

$T = 350$

7. Ans. (A,B,C)

Sol.



For More Material Join: @JEEAdvanced_2025

Net force by surface is resultant (R) of normal and friction

$\tan\theta = \frac{f_r}{N} = \frac{f_r}{mg}$

for $F \geq \mu mg$ $\qquad f_r = \mu mg = \frac{1}{\sqrt{3}} mg$

$\tan\theta = \frac{\frac{1}{\sqrt{3}} mg}{mg} = \frac{1}{\sqrt{3}} \Rightarrow \theta = 30°$

For $F < \mu mg$ $\qquad f_r = F$ and $\tan\theta = \frac{F}{mg} < \frac{\mu mg}{mg}$

$\tan\theta < \frac{1}{\sqrt{3}} \Rightarrow \theta < 30°$

(D) For $F < \mu mg$, θ depends on m but for $F \geq \mu mg$ value of θ is fixed ($= 30°$) and does not depend on m

8. Ans. (A,B,C)

Sol. $F - f = ma$

$30 - 10 = m \times 10$ $\qquad \mu mg = 10$

$20 = 10m$ $\qquad \mu 20 = 10$

$m = 2kg$ $\qquad \mu = \frac{1}{2}$

block start at force = 10. Limiting friction is 10.

9. Ans. (B,C,D)

Sol. Spring force does not change instantaneously.


Thus for m_1 ; $a_1 = a_0$

for m_2 $F_{S_p} = m_2 a_2$ .....(i)

instantaneously aftr F_2 is withdrawn

Initially $F_{S_p} - F_2 = m_2 a_0$


$F_{S_p} = F_2 + m_2 a_0$ .....(ii)



from (i) and (ii) $a_2 = \frac{F_2}{m_2} + a_0$

10. Ans. (A,B,C)

Sol.



Mass 2 kg is at rest, so $T_2 = 20N$

So for (2)

$T_2 - mg = ma$

$20 - 10 = 1 \times a$

$a = 10\ m/s^2$


For More Material Join: @JEEAdvanced_2025


$T_1 = 2\ T_2 = 40$ N
So for (1)
$T_2 - mg = ma$
$40 - 10 = 1 \times a$
$a = 30$ m/s^2
End of rope is coming down with 5 m/s^2
So acceleration of (1) w.r.t. his rope is
$30 + 5 = 35$ m/s^2 upwards
Acceleration of monkey (2) with respect to his rope is zero because he is just holding the string.

Linked Comprehension Type (2 Para × 3Q.) [3 M (-1)]
(Single Correct Answer Type)

11. Ans. (D)

Sol. No slipping between ribbon and blocks implies ribbon also moving with blocks and blocks and ribbon have acceleration of same magnitude





As ribbon is massless, net force on it is zero
$\Rightarrow \quad f_1 = f_2 = f$
by F = ma $\qquad Mg\sin\theta - f = Ma$
$f - mg\sin\theta = ma$

$$\Rightarrow a = \frac{g\sin\theta(M-m)}{M+m}$$

12. Ans. (C)

Sol. Tension = f

$$Mg\sin\theta - f = Ma = \frac{Mg\sin\theta(M-m)}{M+m}$$

$$f = \frac{2Mmg\sin\theta}{(M+m)}$$

13. Ans. (A)


Sol. As θ increases $\Rightarrow$ f increases
as value $f_1)_{max} > f_2)_{max}$
So first m block slip on ribbon

14. Ans. (A)

Sol. At F = 10N, motion between blocks starts

15. Ans. (C)

Sol.



So the block will never move

16. Ans. (A)

Sol. $30 - 10 = 10\ a$
$20 = 10\ a$
$a = 2$


For More Material Join: @JEEAdvanced_2025

Matching List Type (4 × 4) 1 Q. [3 M (–1)]

17. Ans. (D)

Sol. The system slip down if

$(m_1 + m_2)\, g \sin\theta > \mu m_2 g\cos\theta$



$\tan\theta > \dfrac{\mu m_2}{m_1 + m_2} > \dfrac{0.3\times 2}{3}$

$\tan\theta > 0.2$

$\Rightarrow \theta > 11.5°$

For P and Q system will remain stationary hence friction = $(m_1 + m_2)\, g \sin\theta$


For R and S system will move hence limiting friction acts friction = $\mu m_2 g \cos\theta$

SECTION-II

Numerical Answer Type Question (upto second decimal place) 2 Q. [3(0)]

18. Ans. 0.5 s

Sol. $a_A = \dfrac{20-16}{1} = 4 m/s^2$



$a_B = \dfrac{8}{1} = 8 m/s^2$


$a_{B/A} = 4 m/s^2$

$0.5 = 1/2 \times 4t^2$

$t = \dfrac{1}{2}\text{sec} = 0.5\,\text{sec}$

19. Ans. 10 m/s²

Sol.



$ma\cos 37 - \mu\, ma \sin 37 - \mu mg = ma_0$

$25\times\dfrac{4}{5} - \dfrac{2}{5}\times 25\times\dfrac{3}{5} - \dfrac{2}{5}\times 10 = a_0$

$20 - 6 - 4 = 90$

$a_0 = 10\ m/s^2$

For More Material Join: @JEEAdvanced_2025

SECTION-III

Numerical Grid Type (Ranging from 0 to 9) **4 Q. [4 M (0)]**

20. Ans. 8


Sol. a = –3

$$u=\sqrt{\left(25\times\frac{4}{5}-8\right)^2+\left(25\times\frac{3}{5}\right)^2}=\sqrt{(12)^2+(15)^2}=3\sqrt{41}$$

at t = $\frac{3\sqrt{41}}{3}$ so it will stop relative motion before 10 sec. So it will have same velocity as plank.

21. Ans. 5

Sol.



$$\frac{2F_S}{\sqrt{2}}=\frac{mg}{\sqrt{2}};\ F_S=\frac{mg}{2}\quad\therefore\ \text{acceleration}=\frac{F_S}{m}=\frac{g}{2}=5\ \text{m/s}^2$$


22. Ans. 5

Sol. The girl will catch the ball when their horizontal displacement becomes equal

$$\therefore 12t=\frac{1}{2}\left(g\sin 37^\circ\cos 37^\circ\right)t^2\Rightarrow t=5\,\text{sec}$$

23. Ans. 0

Sol. f_k = 6 N which will act till velocity is not zero.




Let velocity becomes zero at t, then $0-4.5=\int_0^t(3-3t)dt\Rightarrow t=3$ so at t = 3, v = 0

SECTION-IV

Matrix Match Type (4 × 5) **1 Q. [8 M (for each entry +2(0))]**

24. Ans. (A) - (S) ; (B) - (S) ; (C) - (Q) ; (D) - (T)

Sol.



$$\text{force}=100\times\frac{2}{100}+100\times\frac{2}{100}$$

4 =2a ; a = 2

$$\text{(C) } 100\times\frac{4\times1}{100}+100\times\frac{4}{100}$$

8 = 2a; a = 4

$$\text{(D) } 100\times\frac{1}{100}+100\times\frac{1}{100}$$

2 = 2a; a = 1

Subjective Type **6 Q. [4 M (0)]**

25. Ans. 3/4

For More Material Join: @JEEAdvanced_2025

Sol. If m notsliding


$a = g \sin\theta$

$mg - N = mg\sin^2\theta$

$N = mg(1 - \sin^2\theta) = mg\cos 2\theta$


$\mu N = mg \sin\theta \cos\theta$

$$\mu = \frac{\sin\theta}{\cos 2\theta} = \tan\theta = \frac{3}{4}$$



26. **Ans.** $M_B = 15$ kg, $T = 15$, $a_A = 5\ m/s^2$, $a_B = 9\ m/s^2$

Sol.



By string constrain

$2Ta_1 + Ta_2 - Ta_3 = 0$

$2a_1 + a_2 = 1$ .... (i)

by $F = ma$

$18 - T = 3 \times 1 \Rightarrow T = 15$

$$2T = 6a_1 \Rightarrow a_1 = \frac{2\times 15}{6} = 5 m/s^2$$

by equation (i) $2(5) + a_2 = 1 \Rightarrow a_2 = -9$

$T - mg = ma_2 \Rightarrow 15 - 10\ m = m(-9)$

$\Rightarrow m = 15$ kg

27. **Ans.** (i) $a_A = g\downarrow$, $a_B = \frac{2g}{3}\uparrow$, $a_c = \frac{2g}{3}\downarrow$, $a_D = \frac{2g}{3}\downarrow$, $a_E = 0$

(ii) $a_A = 0$, $a_B = \frac{g}{3}\downarrow$, $a_C = \frac{g}{3}\uparrow$, $a_D = \frac{g}{3}\uparrow$, $a_E = g\downarrow$

(iii) $a_A = 0, a_B = g\downarrow, a_C = g\uparrow, a_D = 2g\downarrow, a_E = 0$

(iv) $a_A = 0, a_B = 3g\downarrow, a_C = \frac{3g}{2}\downarrow, a_D = \frac{3g}{2}\downarrow, a_E = 0$

Sol. (i) spring 1 is cut

$\vec{a}_A = g\downarrow$

acceleration of B,C,D are a.


$mg - T_1 = ma$

$T_1 - mg - T_2 = ma$

$T_2 - 2mg = ma$

$-2mg = 3ma$

$$a = \frac{2g}{3}$$



For More Material Join: @JEEAdvanced_2025

$a_B = \frac{2g}{3}\uparrow$ $a_C = \frac{2g}{3}\downarrow$ $a_D = \frac{2g}{3}$

$a_E = 0$

(ii) spring (2) is cut $a_A = 0$

all of B.C.D. are let a

$3mg - T_1 = ma$ $\quad \vec{a}_B = \frac{g}{3}\downarrow$ $\quad \vec{a}_C = \frac{g}{3}\uparrow$

$T_1 - T_2 - mg = ma$ $\quad \vec{a}_D = g/3\uparrow$

$T_2 - mg = ma$ $\quad \vec{a}_E = g\downarrow$

$a = \frac{g}{3}$

(iii) String between C & D is cut

$a_A = 0$ $\quad a_E = 0$ $\quad a_D = 2g\downarrow$

let acceleration of B & C

$3mg - T_1 = mg$ $\quad a_B = g\downarrow$ $\quad a_C = g\uparrow$

$T_1 - mg = ma$

(iv) Between B & C is cut

$a_A = 0$ $\quad a_E = 0$

For a_B

$3mg = ma_B$

$a_B = 3g\downarrow$

acceleration of C & D are (a) (let)

$2mg - T = ma$

$mg + T = ma$

$a = \frac{3g}{2}$ $\quad a_c = \frac{3g}{2}\downarrow$ $\quad a_D = \frac{3g}{2}\downarrow$

28. **Ans.** 5 m/s² downwards, 0 m/s², 10 N

Sol. $20 - T = 2a$

for sleeve

$10 - f = 1a_1$

$T = f$

$a_1 + a = 5$


$a_1 = 5 - a$

$20 - f = 2a$ $\quad$ acceleration objective


$10 - f = 5 - a$ $\quad a_1 = 5 - 5 = 0$

$10 = 3a - 5$ $\quad 20 - T = 2 \times 5$


$a = 5$ m/s² downwards $\quad T = 10N$



For More Material Join: @JEEAdvanced_2025

29. **Ans.** (i) 

(ii) a = 3/5 m/s² , T = 18 N, F = 60N

Sol. 

$f_1 = 2f_2$

for bigger block

$f_1 - T = 20a$

$F - f_1 = 50\ a$ $\quad$ $T - f_2 = 5a$

$F - 2f_2 = 50\ a$ $\quad$ ....(i)

$2f_2 - T = 20a$ $\quad$ ...(ii)

$T - f_2 = 5a$ $\quad$ ....(iii)

$f_2 = 25\ a \Rightarrow 15 = 25\ a \Rightarrow a = a = \frac{3}{5} m/s$

from equation (i) $\quad$ $F_2 - 50a = 50\ a$

$F_2 = 100\ a = 60\ N$


$f_2 = \mu\ m_2 g = 0.3 \times 50$ $\quad$ $T - 15 = 3$

$= 15\ N$ $\quad$ $T = 18\ N$

30. Ans. $F = 2\mu_s mg\left(\frac{m+M}{2m+M}\right)$

Sol. $\mu_s mg = (M+2m)a \Rightarrow a = \frac{\mu_s mg}{M+2m}$

Also $F - \mu_s mg = Ma \Rightarrow 2\mu_s mg \frac{(m+M)}{(M+2m)}$



For More Material Join: @JEEAdvanced_2025