EXERCISE (O-1)

SINGLE CORRECT TYPE QUESTIONS

1. If R is a relation from a finite set A having m elements to a finite set B having n elements, then the number of relations from A to B is-

(A) 2^{mn-1} (B) $2^{mn}-1$ (C) 2^{mn} (D) m^n

MRL001

2. If R be a relation '<' from $A = \{1, 2, 3, 4\}$ to $B = \{1, 3, 5\}$ i.e. $(a, b) \in R$ iff $a < b$, then RoR^{-1} is-

(A) {(1, 3), (1, 5), (2, 3), (2, 5), (3, 5), (4, 5)} (B) {(3, 1), (5, 1), (3, 2), (5, 2), (5, 3), (5, 4)}

(C) {(3, 3), (3, 5), (5, 3), (5, 5)} (D) {(3, 3), (3, 4), (4, 5)}

MRL002

3. Let R be a relation from the set {1, 2, 3.....,30} to itself such that $R = \{(a, b) : b = pq$, where $p, q \geq 3$ are prime numbers}. Then, the number of elements in R is :

(A) 110 (B) 120 (C) 130 (D) 140

MRL003

4. Let R_1 and R_2 be relations on the set {1, 2, ..., 7} such that $R_1 = \{(p, p^n) : p$ is a prime and $n \geq 0$ is an integer} and $R_2 = \{(p, p^n) : p$ is a prime and $n = 0$ or 1}. Then, the number of elements in $R_1 - R_2$ is ____.

(A) 0 (B) 1 (C) 2 (D) 4

MRL004

Rankers Academy JEE

5. The relation R defined on the set $A = \{1, 2, 3, 4, 5\}$ by $R = \{(x, y) : |x^2 - y^2| < 16\}$ is given by

(A) {(1, 1), (2, 1), (3, 1), (4, 1), (2, 3)} (B) {(2, 2), (3, 2), (4, 2), (2, 4)}

(C) {(3, 3), (3, 4), (5, 4), (4, 3), (3, 1)} (D) None of these

MRL042

6. Let $R = \{(1, 3), (4, 2), (2, 4), (2, 3), (3, 1)\}$ be a relation on the set $A = \{1, 2, 3, 4\}$. The relation R is-

(A) transitive (B) not symmetric (C) reflexive (D) a function

MRL005

7. Let $R = \{(3, 3), (6, 6), (9, 9), (12, 12), (6, 12), (3, 9), (3, 12), (3, 6)\}$ be relation on the set $A = \{3, 6, 9, 12\}$. The relation is-

(A) reflexive and transitive only (B) reflexive only

(C) an equivalence relation (D) reflexive and symmetric only

MRL006

8. Let L denote the set of all straight lines in a plane. Let a relation R be defined by $\alpha R \beta \Leftrightarrow \alpha \perp \beta$, $\alpha, \beta \in L$. Then R is-

(A) Reflexive (B) Symmetric (C) Transitive (D) none of these

MRL007

9. If R is an equivalence relation in a set A, then R^{-1} is-

(A) reflexive but not symmetric (B) symmetric but not transitive

(C) an equivalence relation (D) none of these

MRL008

For More Join: @IITJEE_Advanced

10. Let $R = \{(x, y) : x, y \in A, x + y = 5\}$, where $A = \{1, 2, 3, 4, 5\}$, then
(A) R is not reflexive, symmetric and not transitive
(B) R is an equivalence relation
(C) R is reflexive, symmetric but not transitive
(D) R is not reflexive, not symmetric but transitive

MRL010

11. In the set $A = \{1, 2, 3, 4, 5\}$ a relation R is defined by $R = \{(x, y) | x, y \in A \text{ and } x < y\}$. Then R is
(A) Reflexive (B) Symmetric (C) Transitive (D) Equivalence relation

MRL011

12. The relation R defined in N as $aRb \Leftrightarrow b$ is divisible by a is
(A) Reflexive but not symmetric (B) Symmetric but not transitive
(C) Symmetric and transitive (D) Equivalence relation

MRL012

13. Which one of the following relations on R is equivalence relation-
(A) $xR_1y \Leftrightarrow x^2 = y^2$ (B) $xR_2y \Leftrightarrow x \geq y$
(C) $xR_3y \Leftrightarrow x | y (x \text{ divides } y)$ (D) $xR_4y \Leftrightarrow x < y$

MRL013

14. Two points P and Q in a plane are related if $OP = OQ$, where O is a fixed point. This relation is-
(A) Reflexive but symmetric (B) Symmetric but not transitive
(C) An equivalence relation (D) none of these

MRL014

15. Let S be the set of all real numbers. Then the relation $R = \{(a, b) : 1 + ab > 0\}$ on S is
(A) Reflexive and symmetric but not transitive
(B) Reflexive, transitive but not symmetric
(C) Symmetric, transitive but not reflexive
(D) Reflexive, transitive and symmetric

MRL015

16. **Statement–1 :** In the set $N \times N$ consider relation R defined as $(a, b)\ R\ (c, d) \Rightarrow ad = bc$; then R is equivalence relation.
Statement–2 : Relation R is reflexive, symmetric & transitive.
(A) Statement–1 is true, Statement–2 is true; Statement–2 is not the correct explanation of Statement–1
(B) Statement–1 is false, Statement–2 is true
(C) Statement–1 is true, Statement–2 is false
(D) Statement–1 is true, Statement–2 is true; Statement–2 is the correct explanation of Statement–1.

MRL016

17. Let R and S be two equivalence relations in a set A. Then-
(A) $R \cup S$ is an equivalence relation in A (B) $R \cap S$ is an equivalence relation in A
(C) $R - S$ is an equivalence relation in A (D) none of these

MRL017

18. Let $A = \{p, q, r\}$. Which of the following is an equivalence relation in A ?
(A) $R_1 = \{(p, q), (q, r), (p, r), (p, p)\}$ (B) $R_2 = \{(r, q)\ (r, p), (r, r), (q, q)\}$
(C) $R_3 = \{(p, p), (q, q), (r, r), (p, q)\}$ (D) none of these

MRL018

Bankers Academy JEE

For More Join: @IITJEE_Advanced

19. Let $R = \{(x, y) : x, y \in N \text{ and } x^2 - 4xy + 3y^2 = 0\}$,
where N is the set of all natural numbers. Then the relation R is :
(A) reflexive and transitive.
(B) symmetric and transitive
(C) reflexive but neither symmetric nor transitive
(D) reflexive and symmetric.

MRL019

20. Let W denote the words in the English dictionary. Define the relation R by : $R = \{(x, y) \in W \times W :$ the words x and y have at least one letter in common$\}$. Then R is-
(A) reflexive, symmetric and not transitive
(B) reflexive, symmetric and transitive
(C) reflexive, not symmetric and transitive
(D) not reflexive, symmetric and transitive

MRL020

Rankers Academy JEE

For More Join: @IITJEE_Advanced

EXERCISE (O-2)

MULTIPLE CORRECT TYPE QUESTIONS

1. Let r be a relation from R(Set of real number) to R defined by $r = \{(x, y) \mid x, y \in R$ and xy is an irrational number$\}$, then which of the following is/are incorrect -
(A) relation is reflexive and symmetric only
(B) relation is symmetric only
(C) relation is symmetric and transitive only
(D) relation is an equivalence relation

MRL049

2. Let S be the set of all subsets of first 10 natural numbers. If a relation R is defined on set S such that $(A,B) \in R$ if $A \cap B \neq \phi$; $A, B \in S$, then which of the following is/are incorrect -
(A) relation is symmetric and transitive
(B) relation is symmetric and reflexive only
(C) relation is symmetric only
(D) relation is reflexive only

MRL050

3. Let $A = N \times N$ be the Cartesian product of N and N. Let
$S = \{((m, n), (p, q)) \in A \times A : m + q = n + p\}$
Consider the following statements:
I. If $((m,n), (p, q)) \in S$, and $((p, q), (r, s)) \in S$ then $((r, s), (m, n)) \in S$
II. There exists at least one element $((m, n), (p, q)) \in S$ such that $((p, q), (m, n)) \notin S$
Then select false options?
(A) **I** is correct
(B) **II** is correct
(C) Both **I** and **II** are correct
(D) Neither **I** nor **II** is correct

MRL051

4. Let $A = Z$, the set of integers. Let $R_1 = \{(m, n) \in Z \times Z: (m + 4n)$ is divisible by 5 in $Z\}$.
Let $R_2 = \{(m, n) \in Z \times Z: (m + 9n)$ is divisible by 5 in $Z\}$.
Which of the following is/are incorrect ?
(A) R_1 is a proper subset of R_2
(B) R_2 is a proper subset of R_1
(C) $R_1 = R_2$
(D) R_1 is not a symmetric relation on Z

MRL052

5. Let X be the set of all persons living in a state. Elements x, y in X are said to be related if '$x < y$', whenever y is 5 years older than x. Which of the following is/are incorrect?
(A) The relation is an equivalence relation
(B) The relation is transitive only
(C) The relation is transitive and symmetric, but not reflexive
(D) The relation is neither reflexive, nor symmetric, nor transitive

MRL053

Rankers Academy JEE
For More Join: @IITJEE_Advanced

6. Which of the following is/are correct for relation R on the set of real numbers ?

(A) $(x, y) \in R \Leftrightarrow 0 < |x| - |y| \leq 1$ is neither transitive nor symmetric.

(B) $(x, y) \in R \Leftrightarrow 0 < |x - y| \leq 1$ is symmetric and transitive.

(C) $(x, y) \in R \Leftrightarrow |x| - |y| \leq 1$ is reflexive but not symmetric.

(D) $(x, y) \in R \Leftrightarrow |x - y| \leq 1$ is reflexive and symmetric.

MRL054

COMPREHENSION TYPE QUESTIONS

Paragraph for Question No. 7 to 9

A relation R is defined on set $A = \{1,2,3...,n\}$.

7. Total number of reflexive relations that can be defined from $A \to A$ is-

(A) 2^{n^2-n} (B) 2^{n^2} (C) 2^{n^2-n+1} (D) 2^{n^2+n}

MRL055

8. Find the maximum number of antisymmetric relation in R is

(A) $3^{\frac{n^2-n}{2}}.2^n$ (B) $3^{\frac{n^2-n}{2}}$ (C) $3^{\frac{n^2+n}{2}}.2^n$ (D) 2^{n^2}

MRL056

9. Total number of symmetric relations that can be defined from $A \to A$ is-

(A) $2^{\frac{n(n+2)}{2}}$ (B) $2^{\frac{n(n-1)}{2}}$ (C) $2^{\frac{n(n+1)}{2}}$ (D) $2^{n(n+2)}$

MRL057

Rankers Academy JEE

MATCHING LIST TYPE QUESTION

10. Match the relation defined on set $A = \{a, b, c\}$ in **List-I** with the corresponding type in **List-II**

	List-I		**List-II**
(I)	$\{(a, b), (b, a)\}$	(P)	symmetric but not reflexive and transitive
(II)	$\{(a, b), (b, a), (a, a), (b, b)\}$	(Q)	equivalence
(III)	$\{(a, b), (b, c), (a, c)\}$	(R)	symmetric and transitive but not reflexive
(IV)	$\{(a, a), (b, b), (c, c)\}$	(S)	transitive but not reflexive and symmetric

(A) I $\to$ P; II $\to$ Q; III $\to$ R; IV $\to$ S (B) I $\to$ P; II $\to$ R; III $\to$ S; IV $\to$ Q

(C) I $\to$ R; II $\to$ S; III $\to$ P; IV $\to$ Q (D) I $\to$ Q; II $\to$ R; III $\to$ S; IV $\to$ P

MRL058

For More Join: @IITJEE_Advanced

EXERCISE (S)

1. If $A = \{2, 3\}$ and $B = \{1, 2\}$, then $n(A \times B)$ is

MRL021

2. A and B are two sets having 3 and 4 elements respectively and having 2 elements in common. If the number of relations which can be defined from A to B is 4^n then value of n -

MRL022

3. The relation R defined in $A = \{1, 2, 3\}$ by a R b if $|a^2 - b^2| \leq 5$. Number of elements in range of R

MRL023

4. Let $A = \{2, 3, 4, 5, \ldots, 16\}$ and '$\simeq$' be an equivalence relation on $A \times A$, defined by $(a, b) \simeq (c, d)$, if and only if $ad = bc$. Then the number of ordered pairs which satisfy this equivalence relation with ordered pair (4, 3) is equal to :

MRL024

5. Given the relation $R = \{(1, 2), (2, 3)\}$ on the set $A = \{1, 2, 3\}$, the minimum number of ordered pairs which when added to R make it an equivalence relation is -

MRL025

6. Let $A=\{0, 3,4,6,7,8,9,10\}$ and R be the relation defined on A such that $R=\{(x,y)\in A\times A: x-y$ is odd positive integer or $x - y = 2\}$. The minimum number of elements that must be added to the relation R, so that it is a symmetric relation, is equal to ____.

MRL043

7. Let $A = \{2, 3, 4\}$ and $B = \{8, 9, 12\}$. Then the number of elements in the relation $R = \{((a_1, b_1), (a_2, b_2)) \in (A \times B, A \times B): a_1$ divides b_2 and a_2 divides $b_1\}$ is :

MRL044

8. Let $A= \{-4, -3, -2, 0, 1, 3, 4\}$ and $R = \{(a, b) \in A \times A: b = |a|$ or $b^2 = a + 1\}$ be a relation on A. Then the minimum number of elements, that must be added to the relation R so that it becomes reflexive and symmetric, is____.

MRL045

9. The minimum number of elements that must be added to the relation $R = \{(a, b), (b, c), (b, d)\}$ on the set $\{a, b, c, d\}$ so that it is an equivalence relation, is _____.

MRL046

10. Let $A=\sum_{i=1}^{10}\sum_{j=1}^{10}\min\{i, j\}$ and $B=\sum_{i=1}^{10}\sum_{j=1}^{10}\max\{i, j\}$. Then $A + B$ is equal to _____.

MRL047

11. Let $A = \{1, 2, 3, 4\}$ and R be a relation on the set $A \times A$ defined by $R=\{((a,b),(c,d)): 2a+3b=4c+5d\}$. Then the number of elements in R is:

MRL048

Rankers Academy JEE

For More Join: @IITJEE_Advanced

EXERCISE - JEE (Main) PYQ

1. Let Z be the set of integers. If $A = \{x \in Z : 2^{(x+2)(x^2-5x+6)} = 1\}$ and $B = \{x \in Z : -3 < 2x - 1 < 9\}$, then the number of subsets of the set $A \times B$, is: **[JEE (Main) 2019]**

(1) 2^{18} (2) 2^{10} (3) 2^{15} (4) 2^{12}

MRL026

2. If $R = \{(x, y) : x, y \in Z, x^2 + 3y^2 \le 8\}$ is a relation on the set of integers Z, then the domain of R^{-1} is: **[JEE (Main) 2020]**

(1) {–2, –1, 1,2} (2) {–1, 0, 1} (3) {–2, –1, 0, 1, 2} (4) {0, 1}

MRL027

3. Let R_1 and R_2 be two relations defined as follows :
$R_1 = \{(a, b) \in R^2 : a^2 + b^2 \in Q\}$ and $R_2 = \{(a, b) \in R^2 : a^2 + b^2 \notin Q\}$,
where Q is the set of all rational numbers. Then: **[JEE (Main) 2021]**

(1) R_2 is transitive but R_1 is not transitive (2) R_1 is transitive but R_2 is not transitive
(3) R_1 and R_2 are both transitive (4) Neither R_1 nor R_2 is transitive

MRL028

4. Let $A = \{2, 3, 4, 5, \ldots, 30\}$ and '$\simeq$' be an equivalence relation on $A \times A$, defined by $(a, b) \simeq (c, d)$, if and only if $ad = bc$. Then the number of ordered pairs which satisfy this equivalence relation with ordered pair (4, 3) is equal to : **[JEE (Main) 2021]**

(1) 5 (2) 6 (3) 8 (4) 7

Rankers Academy JEE

MRL029

5. Let R_1 and R_2 be two relations defined on $\mathbb{R}$ by $a\ R_1\ b \Leftrightarrow ab \ge 0$ and $a\ R_2 b \Leftrightarrow a \ge b$, then **[JEE (Main) 2022]**

(1) R_1 is an equivalence relation but not R_2
(2) R_2 is an equivalence relation but not R_1
(3) both R_1 and R_2 are equivalence relations
(4) neither R_1 nor R_2 is an equivalence relation

MRL030

6. For $\alpha \in N$, consider a relation R on N given by $R = \{(x, y) : 3x + \alpha y$ is a multiple of 7$\}$. The relation R is an equivalence relation if and only if : **[JEE (Main) 2022]**

(1) $\alpha = 14$
(2) α is a multiple of 4
(3) 4 is the remainder when α is divided by 10
(4) 4 is the remainder when α is divided by 7

MRL031

7. Let R be a relation from the set {1, 2, 3.....,60} to itself such that $R = \{(a, b) : b = pq$, where $p, q \ge 3$ are prime numbers$\}$. Then, the number of elements in R is : **[JEE (Main) 2022]**

(1) 600 (2) 660 (3) 540 (4) 720

MRL032

For More Join: @IITJEE_Advanced

8. Let $R_1 = \{(a,b) \in N \times N : |a-b| \leq 13\}$ and $R_2 = \{(a,b) \in N \times N : |a-b| \neq 13\}$. Then on N:

[JEE (Main) 2022]

(1) Both R_1 and R_2 are equivalence relations
(2) Neither R_1 nor R_2 is an equivalence relation
(3) R_1 is an equivalence relation but R_2 is not
(4) R_2 is an equivalence relation but R_1 is not

MRL033

9. Let R_1 and R_2 be relations on the set {1, 2, ..., 50} such that $R_1 = \{(p, p^n) : p$ is a prime and $n \geq 0$ is an integer} and $R_2 = \{(p, p^n) : p$ is a prime and $n = 0$ or 1. Then, the number of elements in $R_1 - R_2$ is.

[JEE (Main) 2022]

MRL034

10. Let a set $A = A_1 \cup A_2 \cup ... \cup A_k$, where $A_i \cap A_j = \phi$ for $i \neq j$ $1 \leq i, j \leq k$. Define the relation R from A to A by $R = \{(x, y) : y \in A_i$ if and only if $x \in A_i, 1 \leq i \leq k\}$. Then, R is : **[JEE (Main) 2022]**

(1) reflexive, symmetric but not transitive
(2) reflexive, transitive but not symmetric
(3) reflexive but not symmetric and transitive
(4) an equivalence relation

MRL035

11. Let R be a relation on $\mathbb{R}$, given by $R = \{(a, b) : 3a - 3b + \sqrt{7}$ is an irrational number}. Then R is

[JEE (Main) 2023]

(1) Reflexive but neither symmetric nor transitive
(2) Reflexive and transitive but not symmetric
(3) Reflexive and symmetric but not transitive
(4) An equivalence relation

Bankers Academy JEE

MRL036

12. Let $P(S)$ denote the power set of $S = \{1,2,3,\ldots,10\}$. Define the relations R_1 and R_2 on $P(S)$ as AR_1B if $(A \cap B^c) \cup (B \cap A^c) = \varnothing$ and AR_2B if $A \cup B^c = B \cup A^c$, $\forall$ $A, B \in P(S)$. Then :

[JEE (Main) 2023]

(1) both R_1 and R_2 are equivalence relations
(2) only R_1 is an equivalence relation
(3) only R_2 is an equivalence relation
(4) both R_1 and R_2 are not equivalence relations

MRL037

13. Among the relations $S = \left\{(a,b) : a,b \in \mathbb{R} - \{0\}, 2 + \frac{a}{b} > 0\right\}$ And $T = \{(a,b) : a,b \in \mathbb{R}, a^2 - b^2 \in Z\}$,

[JEE (Main) 2023]

(1) S is transitive but T is not
(2) T is symmetric but S is not
(3) Neither S nor T is transitive
(4) Both S and T are symmetric

MRL038

14. Let $A = \{1, 3, 4, 6, 9\}$ and $B = \{2, 4, 5, 8, 10\}$. Let R be a relation defined on $A \times B$ such that $R = \{((a_1, b_1), (a_2, b_2)) : a_1 \leq b_2$ and $b_1 \leq a_2\}$. Then the number of elements in the set R is

[JEE (Main) 2023]

(1) 26 (2) 160 (3) 180 (4) 52

MRL039

For More Join: @IITJEE_Advanced

15. The relation $R = \{(a,b) : \gcd(a,b) = 1, 2a \neq b, a,b \in \mathbb{Z}\}$ is: ___

[JEE (Main) 2023]

(1) transitive but not reflexive
(2) symmetric but not transitive
(3) reflexive but not symmetric
(4) neither symmetric nor transitive

MRL040

16. Let $A = \{1, 2, 3, 4, 5, 6, 7\}$. Then the relation $R = \{(x, y) \in A \times A : x + y = 7\}$is

[JEE (Main) 2023]

(1) transitive but neither symmetric nor reflexive
(2) reflexive but neither symmetric nor transitive
(3) an equivalence relation
(4) symmetric but neither reflexive nor transitive

MRL041

17. Let a relation R on $\mathbb{N} \times \mathbb{N}$ be defined as :
$(x_1,y_1)\ R(x_2,y_2)$ if and only if $x_1 \leq x_2$ or $y_1 \leq y_2$
Consider the two statements :
(I) R is reflexive but not symmetric.
(II) R is transitive
Then which one of the following is true ? **[JEE (Main) 2024]**

(1) Only (II) is correct.
(2) Only (I) is correct.
(3) Both (I) and (II) are correct.
(4) Neither (I) nor (II) is correct.

MRL059

Rankers Academy JEE

18. Let $A = \{1, 2, 3, 4, 5\}$. Let R be a relation on A defined by xRy if and only if $4x \leq 5y$. Let m be the number of elements in R and n be the minimum number of elements from $A \times A$ that are required to be added to R to make it a symmetric relation. Then $m + n$ is equal to: **[JEE (Main) 2024]**

(1) 24 (2) 23 (3) 25 (4) 26

MRL060

19. Let the relations R_1 and R_2 on the set
$X = \{1, 2, 3, ..., 20\}$ be given by
$R_1 = \{(x, y) : 2x - 3y = 2\}$ and
$R_2 = \{(x, y) : -5x + 4y = 0\}$. If M and N be the minimum number of elements required to be added in R_1 and R_2, respectively, in order to make the relations symmetric, then $M + N$ equals

[JEE (Main) 2024]

(1) 8 (2) 16 (3) 12 (4) 10

MRL061

20. Let $A = \{2, 3, 6, 8, 9, 11\}$ and $B = \{1, 4, 5, 10, 15\}$ Let R be a relation on $A \times B$ define by $(a, b)R(c, d)$ if and only if $3ad - 7bc$ is an even integer. Then the relation R is **[JEE (Main) 2024]**

(1) reflexive but not symmetric.
(2) transitive but not symmetric.
(3) reflexive and symmetric but not transitive.
(4) an equivalence relation.

MRL062

For More Join: @IITJEE_Advanced

EXERCISE - JEE (Advanced) PYQ

Paragraph for Q.1 & Q.2

Let $S = \{1, 2, 3, 4, 5, 6\}$ and X be the set of all relations R from S to S that satisfy both the following properties.

i. R has exactly 6 elements.

ii. For each $(a, b) \in R$, we have $|a - b| \geq 2$.

Let $Y = \{R \in X : \text{The range of } R \text{ has exactly one element}\}$ and

$Z = \{R \in X : R \text{ is a function from } S \text{ to } S.\}$

Let $n(A)$ denote the number of elements in a set A.

(There are two questions based on PARAGRAPH, the question given below is one of them)

1. If $n(X) = {}^{m}C_6$, then the value of m is _____. **[JEE (Advanced) 2024]**

MRL063

2. If the value of $n(Y) + n(Z)$ is k^2, then $|k|$ is _____. **[JEE (Advanced) 2024]**

MRL064

Rankers Academy JEE

For More Join: @IITJEE_Advanced

ANSWER KEY

EXERCISE (O-1)

Que.	1	2	3	4	5	6	7	8	9	10
Ans.	C	C	B	B	D	B	A	B	C	A
Que.	11	12	13	14	15	16	17	18	19	20
Ans.	C	A	A	C	A	D	B	D	C	A

EXERCISE (O-2)

Que.	1	2	3	4	5	6	7	8	9	10
Ans.	A,C,D	A,B,D	B,C,D	A,B,D	A,B,C	A,C,D	A	A	C	B

EXERCISE (S)

1. 4 **2.** 6 **3.** 3 **4.** 4 **5.** 7
6. 19 **7.** 36 **8.** 7 **9.** 13 **10.** 1100
11. 6

EXERCISE - JEE (Main) PYQ

Que.	1	2	3	4	5	6	7	8	9	10
Ans.	3	2	4	4	4	4	2	2	8	4
Que.	11	12	13	14	15	16	17	18	19	20
Ans.	1	1	2	2	4	4	2	3	4	3

Rankers Academy JEE

EXERCISE - JEE (Advanced) PYQ

Que.	1	2
Ans.	20.00	36.00

For More Join: @IITJEE_Advanced