



Red Ring Disease Management Strategy for Trinidad and Tobago

Assim Dilbar

Technical Officer, Horticulture
Agricultural Services Division
Ministry of Food Production

Coconut Industry Development for the Caribbean:

Towards a Shared Vision & Road Map

Georgetown, Guyana

7-8 October, 2013



Red ring disease

- **Causal agent**(nematode):

Bursaphelenchus(= *Rhadinaphelenchus*)
cocophilus **Host range:**

- Major Host: Coconuts (*Cocos nucifera*); African Oil Palm (*Elaeis guineensis*) ; Date palm (*Phoenix dactylifera*).
- Under green house conditions *West Indian Royal palm* (*Roystonea oleracea*); Gru Gru palm (*Acrocomia aculeata*); Moriche Palms (*Mauritia flexuosa*) and Cucurite palm (*Maximiliana maripa*) were infected artificially
- Minor Host: affects 17 species in Palmae Family



Red ring disease vector



Adult



Larva



Distribution of red ring disease vector:
palm weevil (*Rhynchophorus palmarum*)

- **Central America & Caribbean**

- Barbados, Belize, Costa Rica, Cuba, Dominica, Dominican Republic, El Salvador, Grenada, Guadeloupe, Guatemala, Honduras, Martinique, Nicaragua, Panama, Puerto Rico, Saint Lucia, Saint Vincent and the Grenadines, Trinidad and Tobago

- **North America**

- Mexico



Distribution of red ring disease vector: palm weevil (*Rhynchophorus palmarum*)

■ **South America**

- Argentina, Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Paraguay, Peru, Suriname, Uruguay and Venezuela





Symptoms

- Common in trees 2.5-10 years old
- Chlorosis of oldest leaves tips and spread towards their bases
- Browning of lower leaves
- Breaking of the petiole
- Nut shedding
- Crown toppling
- In cross section of stem, brick-red coloured ring
- In roots, white soft cortex becomes orange to faint red in colour

Chlorosis and necrosis of older leaves



Advanced field symptoms-RRD



Advanced symptoms - RRD



Discolouration of stem - RRD





Spread

- Vector, the palm weevil (*Rynchophorus palmarum*) carries the nematode, *Bursaphelenchus cocophilus* (Cobb) Baujard 1989 in the haemocoel of the weevil larvae via the gut tract
- In adult weevils, the nematode can be found in the gut, body cavity and the region of the ovipositor
- Nematode is transferred through burrowing and feeding by adult and larvae of the palm weevil



Management

- Aggressive phytosanitation
- Mass trapping with baited traps (e.g. chopped sugarcane and synthetic aggregation pheromone, racemic 6-methyl-2-hepten-4-ol (Rhyncholure®))
- Combination of perimeter and internal traps is most effective for mass trapping

Baited traps with insecticide





Trinidad and Tobago Strategy

- Coconut Industry Transformation Committee – Appointed by Food Production Minister
 - Chair-Permanent Secretary – Ministry of Food Production and members from Planning, Research and Extension Divisions, Regional Administration North and South
 - St. Patrick Coconut Growers Association
 - Coconut Growers Association
 - CARDI



Term of Reference - Purpose

- To drive the transformation of the coconut industry in TT to facilitate the development of a competitive and sustainable sub-sector



Term of Reference - Mandate

- To recommend, develop and oversee the implementation of plans and initiatives for the redevelopment of the industry



Term of Reference

- Identify critical areas e.g. RRD
- Develop project proposal(s)
- Seek funding
- Identify areas of responsibility for implementation, operations and delivery of work programmes



Project

**THE MINISTRY OF FOOD PRODUCTION COCONUT
INDUSTRY TRANSFORMATION COMMITTEE**

**PUBLIC SECTION INVESTMENT PROGRAMME
DRAFT ESTIMATES
2013-2014**

**ERADICATION
OF RED RING DISEASE AND POTENTIAL PESTS OF
QUARANTINE SIGNIFICANCE IN COCONUT**

Ministry of Food Production

St. Clair Circle

St. Clair, POS

Telephone numbers: 622-1221/4;622-5596

Email: ps@fplma.gov.tt



Contents

- SUMMARY OF FUNDS
- TERMS OF REFERENCE
- INTRODUCTION
- THE PROJECT
- DEMAND ANALYSIS
- TECHNICAL ASPECTS
- INPUT REQUIREMENTS
- IMPLEMENTATION



Contents

LIST OF TABLES

- Table 1
 - LOGICAL FRAMEWORK NETWORK FOR THE PERIOD 2013-2014 & 2014-2015
- Table 2
 - FUNDING REQUIREMENTS FOR PHASE 1
OCTOBER 2013- SEPTEMBER 2014
- Table 3
 - FUNDING REQUIREMENTS FOR PHASE 2
OCTOBER 2014- SEPTEMBER 2015



Contents

LIST OF APPENDICES:

- APPENDIX 1 – GEOGRAPHICAL DISTRIBUTION OF EXTENSION FIELD **OFFICERS (EFO's)**
- APPENDIX 2 – ORGANOGRAM
- APPENDIX 3- LIST OF OFFICE AND FIELD EQUIPMENT
- APPENDIX 4 - CONTRACTED HUMAN RESOURCE REQUIREMENTS
- APPENDIX 5- JOB DESCRIPTIONS
- APPENDIX 6 - SALARY CLASSIFICATIONS
- APPENDIX 7 – **CONTRACTOR'S EQUIPMENT AND MATERIALS PER CREW**



Contents

LIST OF APPENDICES: (continued)

- APPENDIX 8- ETIS COSTING – TRAINING AND MASS COMMUNICATION
- APPENDIX 9- RRD TRAP SETTING /ANNUAL & COSTS
- APPENDIX 10- **CONTRACTOR'S EQUIPMENT AND LABOUR COSTS**
- APPENDIX 11- COST OF PHEROMONE PER ANNUM
- APPENDIX 12- Terms of Reference for Integrated Strategic Plan & Integrated Feasibility Study
- APPENDIX 13-Gantt Charts (Phase 1 October 2013-September 2014)
- APPENDIX 14- Gantt Charts (Phase 2 October 2014-September 2015)



OTHER ACTIVITIES

- Replanting
 - Identify local seed sources
 - Develop seed gardens
 - Transplant (GAP)
 - ID new/improved germplasm
 - Conduct PRA
 - Introduce
 - Evaluate
 - Expand



OTHER ACTIVITIES

- Continuous monitoring for RRD symptoms
- Removal and destruction
- Baiting and trapping
- Surveillance for **new pests**

Most estates with old trees



Small acreages replanted



Seed garden - Manzanilla



Need to achieve – productive trees




Visit by team from Embrapa



Visit by team from India





Thank You