Smashing single cells into k-mer sketches

This manuscript (<u>permalink</u>) was automatically generated from <u>czbiohub/primate-brain-organoid-paper@4864d14</u> on November 1, 2019.

Authors

• Olga Borisovna Botvinnik

© 0000-0003-4412-7970 · ♥ olgabot · ♥ olgabot

Data Sciences Platform, Chan Zuckerberg Biohub

Abstract

Single-cell RNA-sequencing is a powerful technology for identifying novel and known cell types, however its power is limited to organisms with well-annotated genomes. We demonstrate the utility of using annotation-agnostic methods which quantify cell-cell similarity using k-mer profiles. We benchmark a few methods and demonstrate the utility of converting cell types from mouse to human and back, and compare to using purely 1:1 mapped orthologous genes.

Introduction

There are a predicted 8.7 million Eukaryotic species on earth [1], yet only 14% (1,233,500) have been catalogued and 0.001% (9,449) have genomes deposited in the National Center for Biotechnology Information Genome Assembly [2]. And yet, the genome sequence is not enough. To truly understand the diversity of life on this planet, we need to determine not just the DNA blueprints of life, but understand the instantiation of the DNA, the cell types of the species. While sequencing DNA gives a quantitative measure of the nucleotide differences, it does not inform the functional strategies that change with DNA sequence. As new species can be defined by a new cell type. For example, the existence of a single cell type, the Cnidocyte [3], a stinging cell of a single-celled biological weapon, defines the phylum Cnidaria. Thus, entire clades, not only species, can be defined by the introduction of an additional cell type or state.

Novel organizations of existing cell states can also define cell types. For example, the development of genitalia in amniotes, while using similar cell types, ultimately uses a different physical organization of cell types to generate genitalia in mammals compared to reptiles [4]

Determining common gene ancestry ("orthology") is a difficult problem. Many approaches exist [5].

Determining common ancestry of cell types ("orthologous cell types") [6] is an additional difficult problem. Comparative transcriptomics begins with finding a common feature set for embedding molecular profiles across divergent species into a common space. Many researchers take the approach of using one-to-one orthologous genes [Cite: brawand2011, CCA, LIGER, Scanorama, basically all the single cell "alignment" packages], others use clusters of orthologous groups [8], others map reads onto a common genome derived from whole-genome alignment [cite: recent primate brain paper from Barbara Treutlein], or map onto native genomes [9] and re-annotate using a tool such as Comparative Annotation Toolkit [10].

k-mers have been proposed for comparing single cells [11] as they are a fast, simple way to create cell-cell similarities. However, the work so far has focused on using annotated organisms and not cross-species analyses.

Methods

Methods go here.

We used ENSEMBL version 97.

Results

To determine whether short segments of sequences could detect gene orthologues, we k-merized orthologous genes derived from the ENSEMBL version 97 [12] COMPARA database [13] (Fig.~). We compared human protein sequences to orthologous chimpanzee, mouse, (orangutan, bonobo, gorilla, macaque, opossum, platypus, chicken) protein sequences, as these are species used in [14]. As a background, we randomly chose 10 non-orthologous genes relative to the human gene. In addition to k-merizing the protein-coding sequence, we also re-encoded the protein-coding sequence into Dayhoff [15] and hydrophobic-polar encodings [16], show in Table~.

Table 1: Dayhoff and hydrophobic-polar encodings are a reduced amino acid alphabet allowing for permissive cross-species sequence comparisons. For example, the amino acid sequence SASHAFIERCE would be Dayhoff-encoded to bbbdbfecdac, and HP-encoded to phpphhhpppp.

| Amino acid | Property | Dayhoff | Hydrophobic-polar (HP) |
|---------------|-----------------------|---------|------------------------|
| С | Sulfur polymerization | a | р |
| A, G, P, S, T | Small | b | A, G, P: h |
| | | | S,T: p |
| D, E, N, Q | Acid and amide | С | р |
| H, K, R | Basic | d | р |
| I, L, M, V | Hydrophobic | е | h |
| F, W, Y | Aromatic | f | h |




Figure 1: Figure 1.

We found that, consistent with previous knowledge, that 1:1 orthologues had higher k-mer similarities as determined by the Jaccard Index.

Additionally, more recently diverged genes had higher k-mer similarity as well.



Extract coding sequence using human proteins, k-merize coding sequence and minhash, com pare samples/cells across minhashes




Figure 2: Figure 2.

Discussion

References

1. How Many Species Are There on Earth and in the Ocean?

Camilo Mora, Derek P. Tittensor, Sina Adl, Alastair G. B. Simpson, Boris Worm *PLoS Biology* (2011-08-23) https://doi.org/fpr4z8

DOI: <u>10.1371/journal.pbio.1001127</u> · PMID: <u>21886479</u> · PMCID: <u>PMC3160336</u>

2. Genome List - Genome - NCBI https://www.ncbi.nlm.nih.gov/genome/browse/

3. Cnidocyte

Wikipedia

(2019-10-20) https://en.wikipedia.org/w/index.php?title=Cnidocyte&oldid=922123291

4. A relative shift in cloacal location repositions external genitalia in amniote evolution

Patrick Tschopp, Emma Sherratt, Thomas J. Sanger, Anna C. Groner, Ariel C. Aspiras, Jimmy K. Hu, Olivier Pourquié, Jérôme Gros, Clifford J. Tabin

Nature (2014-11-05) https://doi.org/f6sg74

DOI: <u>10.1038/nature13819</u> · PMID: <u>25383527</u> · PMCID: <u>PMC4294627</u>

5. Standardized benchmarking in the quest for orthologs

Adrian M AltenhoffBrigitte Boeckmann, Salvador Capella-Gutierrez, Daniel A Dalquen, Todd DeLuca, Kristoffer Forslund, Jaime Huerta-Cepas, Benjamin Linard, Cécile Pereira, ... Christophe Dessimoz *Nature Methods* (2016-04-04) https://doi.org/f3rpzx

DOI: 10.1038/nmeth.3830 · PMID: 27043882 · PMCID: PMC4827703

6. The origin and evolution of cell types

Detlev Arendt, Jacob M. Musser, Clare V. H. Baker, Aviv Bergman, Connie Cepko, Douglas H. Erwin, Mihaela Pavlicev, Gerhard Schlosser, Stefanie Widder, Manfred D. Laubichler, Günter P. Wagner *Nature Reviews Genetics* (2016-11-07) https://doi.org/f9b62x

DOI: <u>10.1038/nrg.2016.127</u> · PMID: <u>27818507</u>

7. How Single-Cell Genomics Is Changing Evolutionary and Developmental Biology

John C. Marioni, Detlev Arendt

Annual Review of Cell and Developmental Biology (2017-10-06) https://doi.org/ggb632

DOI: <u>10.1146/annurev-cellbio-100616-060818</u> · PMID: <u>28813177</u>

8. The COG database: a tool for genome-scale analysis of protein functions and evolution

R. L. Tatusov

Nucleic Acids Research (2000-01-01) https://doi.org/fr3gqz

DOI: 10.1093/nar/28.1.33 · PMID: 10592175 · PMCID: PMC102395

9. Establishing Cerebral Organoids as Models of Human-Specific Brain Evolution

Alex A. Pollen, Aparna Bhaduri, Madeline G. Andrews, Tomasz J. Nowakowski, Olivia S. Meyerson, Mohammed A. Mostajo-Radji, Elizabeth Di Lullo, Beatriz Alvarado, Melanie Bedolli, Max L. Dougherty, ... Arnold R. Kriegstein

Cell (2019-02) https://doi.org/gfvgzr

DOI: <u>10.1016/j.cell.2019.01.017</u> · PMID: <u>30735633</u> · PMCID: <u>PMC6544371</u>

10. Comparative Annotation Toolkit (CAT)—simultaneous clade and personal genome annotation

Ian T. Fiddes, Joel Armstrong, Mark Diekhans, Stefanie Nachtweide, Zev N. Kronenberg, Jason G. Underwood, David Gordon, Dent Earl, Thomas Keane, Evan E. Eichler, ... Benedict Paten

Genome Research (2018-06-08) https://doi.org/gdpg7n

DOI: 10.1101/gr.233460.117 · PMID: 29884752 · PMCID: PMC6028123

11. K-mer counting with low memory consumption enables fast clustering of single-cell sequencing data without read alignment

Christina Huan Shi, Kevin Y. Yip

bioRxiv (2019-08-02) https://www.biorxiv.org/content/10.1101/723833v1

DOI: <u>10.1101/723833</u>

12. Ensembl 2018

Daniel R Zerbino, Premanand Achuthan, Wasiu Akanni, M Ridwan Amode, Daniel Barrell, Jyothish Bhai, Konstantinos Billis, Carla Cummins, Astrid Gall, Carlos García Girón, ... Paul Flicek

Nucleic Acids Research (2017-11-16) https://doi.org/gcwg6r

DOI: 10.1093/nar/gkx1098 · PMID: 29155950 · PMCID: PMC5753206

13. Ensembl comparative genomics resources

Javier Herrero, Matthieu Muffato, Kathryn Beal, Stephen Fitzgerald, Leo Gordon, Miguel Pignatelli, Albert J. Vilella, Stephen M. J. Searle, Ridwan Amode, Simon Brent, ... Paul Flicek *Database* (2016) https://doi.org/ggb9tv

DOI: 10.1093/database/bav096 · PMID: 26896847 · PMCID: PMC4761110

14. The evolution of gene expression levels in mammalian organs

David Brawand, Magali Soumillon, Anamaria Necsulea, Philippe Julien, Gábor Csárdi, Patrick Harrigan, Manuela Weier, Angélica Liechti, Ayinuer Aximu-Petri, Martin Kircher, ... Henrik Kaessmann *Nature* (2011-10) https://doi.org/fcvk54

DOI: 10.1038/nature10532 · PMID: 22012392

15. Atlas of protein sequence and structure

Margaret O Dayhoff
National Biomedical Research Foundation. (1969)

16. Physical biology of the cell

Rob Phillips, Julie Theriot, Jane Kondev, Hernan Garcia *Garland Science* (2012)