

Title: Spinocerebellar Ataxia Type 14 *GeneReview* Table 2  
Authors: Chen D-H, Bird TD, Raskind WH  
Updated: April 2013  
Note: The following information is provided by the authors listed above and has not been reviewed by *GeneReviews* staff.

Table 2. SCA14 Genotypes and Phenotypes

Familial (F)/ Sporadic (S)	Predicted Amino Acid Changes	Nucleotide Changes	Exon	Mean Age of Onset (Range in yrs)	Gait Ataxia	Dysarthria	Abnormal Eye Movement	Myoclonus	Tremor / Chorea	Decreased Sense	Depression	Cognitive Deficits	Additional Signs	Tendon Reflexes	MRI Cerebellar Atrophy	Author
F	p.R26G	c.76a>g	1	37(15-68)	+	+	9/10	–	1/10	–	–	–	ptosis	↓/↑	+	Sailer et al 2012
F	p.R41P	c.122g>c	1	46(29-58)	+	+	–	–	–	–	–	–	–	nl	+	Chen et al 2005
F	p.G63V	c.188g>t	2	Childhood-50	+	+	+	–	–	–	–	–	Dystonia	↓	+	Nolte et al 2007b
F	p.C77S	c.229t>a	3	40 (26, 56)	+	+	+	–	–	+	+	–	–	N/A	+	Wieczorek et al 2007
S	p.K100-H101del	c.296-301del	4	38	+	+	+	–	–	–	+	–	–	↑	+	Chen et al 2005
F	p.H101Y	c.301c>t	4	31(10-50)	+	+	+	–	–	–	–	–	–	↓/↑	+	Chen et al 2003
F	p.H101Y	c.301c>t	4	adolescence	+	N/A	N/A	–	+	–	+	–	Psychosis	N/A	+	Nolte et al 2007a
F	p.H101Q	c.303c>g	4	32 (40-20)	+	+	+	–	–	–	–	–	–	N/A	+	Alonso et al 2004
F	p.C114Y	c.341g>a	4	28.8(17-37)	+	+	2/5	–	2/5	–	–	1/5	Plantar+	nl/↑	+	Klebe et al 2005
F	p.G118D	c.353g>a	4	38.3(21-59)	+	+	+	2/23	1/23	8/22	–	–	Dystonia	nl/↑/↓	+	van de Warrenburg et al 2003, Verbeek et al 2005, Visser et al 2007
F	p..S119P	c.355t>c	4	42(35-51)	+	+	–	–	–	+	+	–		↑	+	Chen et al 2003
F	p.S119F	c.356c>t	4	52.7(14-70)	+	–	–	–	–	–	–	1/5	Seizures	nl/↑	+	Hiramoto et al 2006
F	p.G123R	c.367g>a	4	48.3(40-60)	+	+/–	2/3	–	–	2/3	–	1/3	Deafness	N/A	+	Klebe et al 2005
F	p.G123E	c.368g>a	4	36(24-48)	+	+/–	–	1/2	1/2	1/2	–	2/2		N/A	+	Klebe et al 2005
F	p.Q127R	c.380a>g	4	28(12-42)	+	+	+	5/11	–	–	–	–		↓	+	Yabe et al 2003
S	p.G128D	c.383g>a	4	22, 44	+	+	+	–	–	–	–	–		nl	+	Chen et al 2003, Morita et al 2006
F	p.G128D	c.383g>a	4	32 (20-49)	+	+	+	–	+	–	+	2/3	Dystonia Vertigo ↓vision	↑	+	Miura et al 2009
F	p.C131R	c.391t>c	4	childhood-50	+	+	–	–	+	–	–	–	Seizures?	N/A	+	Dalski et al 2006
F	p.C131Y	c.392g>a	4	adult (18, 50)	+	+	–	–	–	–	–	–	–	↑	+	Klebe et al 2005
F	p.V138E	c.413t>a	4	22(3-45)	+	+/–	+	1/7	2/7	2/7	–	–	Plantar+	nl/↑	+	Vlak et al 2006
F	p.H139G	c.47c>t	5	28.1(10-45)	+	+/–	+	–	1/13	6/13	–	3/13		nl/↑	+	Koht e al 2012
F	p.C150F	c.449,450gc>tt	5	31.3(19-44)	+	+/–	+	–	1/6	4/6	–	–		↑	+	Fahey et al 2005
F	p.G360S	c.1078g>a	10	36.5(20-53)	+	+/–	–	–	2/2	–	–	–	Rippling	N/A	+	Klebe et al 2005
F	p.S361G	c.1081a>g	10	28(5-60)	+	+	–	–	–	–	+	–		nl	+	Chen et al 2005

Familial (F)/ Sporadic (S)	Predicted Amino Acid Changes	Nucleotide Changes	Exon	Mean Age of Onset (Range in yrs)	Gait Ataxia	Dysarthria	Abnormal Eye Movement	Myoclonus	Tremor / Chorea	Decreased Sense	Depression	Cognitive Deficits	Additional Signs	Tendon Reflexes	MRI Cerebellar Atrophy	Author
F	p.F643L	c.1928t>c	18	34(child-60)	+	+/–	10/18	Facial myokymia	4/18	4/18	–	13/18		↑	+	Stevanin et al 2004
F	p.F643L	c.1928t>c	18	20	+	+	–	–	–	+	–	–		N/A	+	Klebe et al 2005
F	p.V692G	c.2075t>g	18	15	+	+	–	–	2/2	–	–	N/A	Plantar+	↑	+	Klebe et al 2005
F	p.M697I-ex13	c.2089-2192del	18	Childhood-60	+	N/A	N/A	1/3	+	N/A		–		↓	+	Asai et al 2009

Literature Cited

Alonso I, Costa C, Coutinho P, Silveira I, Sequeiros J (2004) Molecular analysis of the PRKCG gene in Portuguese patients with spinocerebellar ataxia. Am J Hum Genet 75 Suppl:2425

Asai H, Hirano M, Shimada K, Kiriyaama T, Furiya Y, Ikeda M, Iwamoto T, Mori T, Nishinaka K, Konishi N, Udaka F, Ueno S (2009) Protein kinase C gamma, a protein causative for dominant ataxia, negatively regulates nuclear import of recessive-ataxia-related aprataxin. Hum Mol Genet. 18:3533-43.

Chen DH, Brkanac Z, Verlinde CL, Tan XJ, Bylenok L, Nochlin D, Matsushita M, Lipe H, Wolff J, Fernandez M, Cimino PJ, Bird TD, Raskind WH (2003) Missense mutations in the regulatory domain of PKC gamma: a new mechanism for dominant nonepisodic cerebellar ataxia. Am J Hum Genet 72:839-49

Chen DH, Cimino PJ, Ranum LP, Zoghbi HY, Yabe I, Schut L, Margolis RL, Lipe HP, Feleke A, Matsushita M, Wolff J, Morgan C, Lau D, Fernandez M, Sasaki H, Raskind WH, Bird TD (2005) The clinical and genetic spectrum of spinocerebellar ataxia 14. Neurology 64:1258-60

Dalski A, Mitulla B, Burk K, Schattenfroh C, Schwinger E, Zuhlke C (2006) Mutation of the highly conserved cysteine residue 131 of the SCA14 associated PRKCG gene in a family with slow progressive cerebellar ataxia. J Neurol 253:1111-2

Fahey MC, Knight MA, Shaw JH, Gardner RJ, du Sart D, Lockhart PJ, Delatycki MB, Gates PC, Storey E (2005) Spinocerebellar ataxia type 14: study of a family with an exon 5 mutation in the PRKCG gene. J Neurol Neurosurg Psychiatry 76:1720-2

Hiramoto K, Kawakami H, Inoue K, Seki T, Maruyama H, Morino H, Matsumoto M, Kurisu K, Sakai N (2006) Identification of a new family of spinocerebellar ataxia type 14 in the Japanese spinocerebellar ataxia population by the screening of PRKCG exon 4. Mov Disord 21:1355-60

Klebe S, Durr A, Rentschler A, Hahn-Barma V, Abele M, Bouslam N, Schols L, Jedynak P, Forlani S, Denis E, Dussert C, Agid Y, Bauer P, Globas C, Wullner U, Brice A, Riess O, Stevanin G (2005) New mutations in protein kinase Cgamma associated with spinocerebellar ataxia type 14. Ann Neurol 58:720-9

Koht J, Stevanin G, Durr A, Mundwiller E, Brice A, Tallaksen CME (2012) SCA14 in Norway, two families with autosomal dominant cerebellar ataxia and a novel mutation in the PRKCG gene. Acta Neurologica Scandinavica 125:116-122.

Miura S, Nakagawara H, Kaida H, Sugita M, Noda K, Motomura K, Ohyagi Y, Ayabe M, Aizawa H, Ishibashi M, Taniwaki T (2009) Expansion of the phenotypic spectrum of SCA14 caused by the Gly128Asp mutation in PRKCG. Clin Neurol Neurosurg. 111:211-5.

Morita H, Yoshida K, Suzuki K, Ikeda S (2006) A Japanese case of SCA14 with the Gly128Asp mutation. J Hum Genet 51:1118-21

Nolte D, Klebe S, Baron R, Deuschl G, Müller U (2007a) Codon 101 of PRKCG, a preferential mutation site in SCA14. Mov Disord. 22:1831-2

Nolte D, Landendinger M, Schmitt E, Müller U (2007b) Spinocerebellar ataxia 14: novel mutation in exon 2 of PRKCG in a German family. Mov Disord. 22:265-7

Sailer A, Scholz SW, Gibbs JR, Tucci A, Johnson JO, Wood NW, Plagnol V, Hummerich H, Ding J, Hernandez D, Hardy J, Federoff HJ, Traynor BJ, Singleton AB, Houlden H (2012) Exome sequencing in an SCA14 family demonstrates its utility in diagnosing heterogeneous diseases. Neurology. 79:127-31

Stevanin G, Hahn V, Lohmann E, Bouslam N, Gouttard M, Soumphonphakdy C, Welter ML, Ollagnon-Roman E, Lemainque A, Ruberg M, Brice A, Durr A (2004) Mutation in the catalytic domain of protein kinase C gamma and extension of the phenotype associated with spinocerebellar ataxia type 14. Arch Neurol 61:1242-8

van de Warrenburg BP, Verbeek DS, Piersma SJ, Hennekam FA, Pearson PL, Knoers NV, Kremer HP, Sinke RJ (2003) Identification of a novel SCA14 mutation in a Dutch autosomal dominant cerebellar ataxia family. Neurology 61:1760-5

Verbeek DS, Warrenburg BP, Hennekam FA, Dooijes D, Ippel PF, Verschuuren-Bemelmans CC, Kremer HP, Sinke RJ (2005) Gly118Asp is a SCA14 founder mutation in the Dutch ataxia population. Hum Genet 117:88-91

Visser JE, Bloem BR, van de Warrenburg BP (2007) PRKCG mutation (SCA-14) causing a Ramsay Hunt phenotype. Mov Disord. 22:1024-6

Vlak MH, Sinke RJ, Rabelink GM, Kremer BP, van de Warrenburg BP (2006) Novel PRKCG/SCA14 mutation in a Dutch spinocerebellar ataxia family: expanding the phenotype. Mov Disord 21:1025-8