

Title: Fanconi Anemia *GeneReview* – Molecular Genetics of Less Commonly Involved Genes

Authors: Mehta PA, Tolar J

Updated: September 2016

Note: The following information is provided by the authors and has not been reviewed by *GeneReviews* staff.

ERCC4

Gene structure. *ERCC4* has 11 exons ([NM_005236.2](#))

Pathogenic allelic variants. See [Table A](#).

Normal gene product. *ERCC4* ([NP_005227.1](#); 916 amino acids) is an excision and cross-link repair protein that plays a role in DNA stability and repair of DNA damage. Aside from Fanconi anemia, aberrant *ERCC4* has been associated with Conskayne syndrome, xeroderma pigmentosum, and breast/ovarian cancer [Bogliolo et al 2013, Kashiwama et al 2013].

Abnormal gene product. See [Molecular Genetic Pathogenesis](#).

FANCL

Gene structure. *FANCL* has 14 exons ([NM_018062.2](#)).

Pathogenic allelic variants. See [Table A](#).

Normal gene product. The E3 ubiquitin-protein ligase FANCL has 375 amino acids. It is a component of the FA core complex with three WD40 (Tryptophan-Aspartate -40) repeats and a PHD finger motif (a variant RING finger motif) [Meetei et al 2003] and is presumed to be the catalytic subunit of the FA core complex as an ubiquitin ligase for FANCD2 and FANCI [Hodson et al 2011, Miles et al 2015]. FANCL directly interact with UBE2T (E2 ubiquitin conjugating enzyme) [Machida et al 2006, Hodson et al 2014]. A baculoviral generated protein has been shown to have in vitro monoubiquitylation activity [Alpi et al 2008]. FANCL is relevant to normal development as is evidenced by its dysfunction linked to VACTERL association [Vetro et al 2015].

Abnormal gene product. See [Molecular Genetic Pathogenesis](#).

FANCM

Gene structure. *FANCM* has 23 exons ([NM_020937.1](#)).

Pathogenic allelic variants. See [Table A](#).

Normal gene product. The Fanconi anemia group M protein (FANCM) has 2048 amino acids. It is a component of the FA core complex, contains the seven helicase-specific motifs, one degenerate endonuclease domain, and ssDNA and dsDNA-stimulated ATPase activity and DNA translocase activity [Meetei et al 2005, Singh et al 2013]. FANCM is phosphorylated in response to DNA damage and complexes with the

histone-fold complex MHF [Tao et al 2012, Fox et al 2014, Girard et al 2014, Xue et al 2015]. In concert with FAAP24, a FANCM binding protein [Coulthard et al 2013], FANCM participates in a checkpoint reaction to DNA damage, nucleotide excision repair, and maintenance of genome stability [Deans & West 2009, Huang et al 2010, Kelsall et al 2012, Wang et al 2013].

Abnormal gene product. See [Molecular Genetic Pathogenesis](#).

***PALB2* (previously FANCN)**

Gene structure. *PALB2* has 13 exons ([NM_024675.3](#)).

Pathogenic allelic variants. See [Table A](#).

Normal gene product. PALB2, the partner and localizer of BRCA2 protein, has 1186 amino acids. It regulates localization and stability of BRCA2 protein. Short sections of the PALB2 N-terminus share homologies with a segment of prefoldin and the light chain 3 (LC3) of microtubule-associated protein MAP1. PALB2 also has two WD40 repeat-like segments at the C terminus [Xia et al 2006]. *PALB2* is one of the FA-related genes that is also a breast cancer susceptibility gene. It is a known protein partner of FANCD1/BRCA2 and plays a role in the homologous recombination repair pathway [Xia et al 2007, Guo et al 2015].

Abnormal gene product. See [Molecular Genetic Pathogenesis](#).

RAD51C

Gene structure. *RAD51C* has nine exons ([NM_058216](#)).

Pathogenic allelic variants. See [Table A](#).

Normal gene product. RAD51C is a protein of 376 amino acids shown to participate in several distinct protein complexes involved in homologous recombination [Somyajit et al 2012]. RAD51 variants have been demonstrated to bind to single strand overhangs that occur after processing of DNA lesions in concert with BRCA2 [Somyajit et al 2010].

Abnormal gene product. See [Molecular Genetic Pathogenesis](#).

SLX4

Gene structure. *SLX4* has 15 exons (reference sequence [NM_032444](#)).

Pathogenic allelic variants. See [Table A](#).

Normal gene product. SLX4 is a protein of 1834 amino acids involved in resolution of homologous recombination intermediates, such as Holliday junctions [Kim et al 2011, Stoepker et al 2011]. SLX4 interacts with other endonuclease complexes, including MUS81-EME1 and XPF-ERCC1 [Kim et al 2011, Stoepker et al 2011], interacts with XPF (ERCC4, FANCD1) [Bogliolo et al 2013, Kashiya et al 2013, Hashimoto et al 2015], and is operational in multiple aspects of genome maintenance [Cybulski & Howlett 2011, Kim et al 2011, Salewski et al 2012, Schuster et al 2013, Kim 2014].

Abnormal gene product. See [Molecular Genetic Pathogenesis](#).

UBE2T

Gene structure. *UBE2T* has six exons ([NM_001310326](#)).

Pathogenic allelic variants. See [Table A](#).

Normal gene product. UBE2T is a protein of 197 amino acids. It functions as E2 conjugase that interacts with FANCL and mediates monoubiquitination of FANCD2 and FANCI. Critically, it is not a member of the FA core complex of proteins, but rather is recruited to damaged chromatin independently [Machida et al 2006, Alpi et al 2007, Longerich et al 2014, Hira et al 2015, Miles et al 2015, Rickman et al 2015, Virts et al 2015].

Abnormal gene product. See [Molecular Genetic Pathogenesis](#).

References

- Alpi A, Langevin F, Mosedale G, Machida YJ, Dutta A, Patel KJ. UBE2T, the Fanconi anemia core complex, and FANCD2 are recruited independently to chromatin: a basis for the regulation of FANCD2 monoubiquitination. *Mol Cell Biol*. 2007;27:8421-30.
- Alpi AF, Pace PE, Babu MM, Patel KJ. Mechanistic insight into site-restricted monoubiquitination of FANCD2 by Ube2t, FANCL, and FANCI. *Mol Cell*. 2008;32:767-77.
- Bogliolo M, Schuster B, Stoepker C, Derkunt B, Su Y, Raams A, Trujillo JP, Minguillón J, Ramírez MJ, Pujol R, Casado JA, Baños R, Rio P, Knies K, Zúñiga S, Benítez J, Bueren JA, Jaspers NG, Schärer OD, de Winter JP, Schindler D, Surrallés J. Mutations in ERCC4, encoding the DNA-repair endonuclease XPF, cause Fanconi anemia. *Am J Hum Genet*. 2013;92:800-6.
- Coulthard R, Deans AJ, Swuec P, Bowles M, Costa A, West SC, McDonald NQ. Architecture and DNA recognition elements of the Fanconi anemia FANCM-FAAP24 complex. *Structure*. 2013;21:1648-58.
- Cybulski KE, Howlett NG. FANCP/SLX4: a Swiss army knife of DNA interstrand crosslink repair. *Cell Cycle*. 2011;10:1757-63.
- Deans AJ, West SC. FANCM connects the genome instability disorders Bloom's Syndrome and Fanconi Anemia. *Mol Cell*. 2009;36:943-53.
- Fox D 3rd, Yan Z, Ling C, Zhao Y, Lee DY, Fukagawa T, Yang W, Wang W. The histone-fold complex MHF is remodeled by FANCM to recognize branched DNA and protect genome stability. 2014;24:560-75.
- Girard C, Crismani W, Froger N, Mazel J, Lemhemdi A, Horlow C, Mercier R. FANCM-associated proteins MHF1 and MHF2, but not the other Fanconi anemia factors, limit meiotic crossovers. *Nucleic Acids Res*. 2014;42:9087-95.
- Guo Y, Feng W, Sy SM, Huen MS. ATM-dependent phosphorylation of the Fanconi anemia protein PALB2 promotes the DNA damage response. *J Biol Chem*. 2015;290:27545-56.
- Hashimoto K, Wada K, Matsumoto K, Moriya M. Physical interaction between SLX4 (FANCP) and XPF (FANCP) proteins and biological consequences of interaction-defective missense mutations. *DNA Repair (Amst)*. 2015;35:48-54.
- Hira A, Yoshida K, Sato K, Okuno Y, Shiraishi Y, Chiba K, Tanaka H, Miyano S, Shimamoto A, Tahara H, Ito E, Kojima S, Kurumizaka H, Ogawa S, Takata M, Yabe H, Yabe M. Mutations in the gene encoding the E2 conjugating enzyme UBE2T cause Fanconi anemia. *Am J Hum Genet*. 2015;96:1001-7.
- Hodson C, Cole AR, Lewis LP, Miles JA, Purkiss A, Walden H. Structural analysis of human FANCL, the E3 ligase in the Fanconi anemia pathway. *J Biol Chem*. 2011;286:32628-37.

Hodson C, Purkiss A, Miles JA, Walden H. Structure of the human FANCL RING-Ube2T complex reveals determinants of cognate E3-E2 selection. *Structure*. 2014;22:337-44.

Huang M, Kim JM, Shiotani B, Yang K, Zou L, D'Andrea AD. The FANCM/FAAP24 complex is required for the DNA interstrand crosslink-induced checkpoint response. *Mol Cell*. 2010;39:259-68.

Kashiyama K, Nakazawa Y, Pilz DT, Guo C, Shimada M, Sasaki K, Fawcett H, Wing JF, Lewin SO, Carr L, Li TS, Yoshiura K, Utani A, Hirano A, Yamashita S, Greenblatt D, Nardo T, Stefanini M, McGibbon D, Sarkany R, Fassihi H, Takahashi Y, Nagayama Y, Mitsutake N, Lehmann AR, Ogi T. Malfunction of nuclease ERCC1-XPF results in diverse clinical manifestations and causes Cockayne syndrome, xeroderma pigmentosum, and Fanconi anemia. *Am J Hum Genet*. 2013;92:807-19.

Kelsall IR, Langenick J, MacKay C, Patel KJ, Alpi AF. The Fanconi anaemia components UBE2T and FANCM are functionally linked to nucleotide excision repair. *PLoS One*. 2012;7:e36970.

Kim Y. Nuclease delivery: versatile functions of SLX4/FANCP in genome maintenance. *Mol Cells*. 2014;37:569-74.

Kim Y, Lach FP, Desetty R, Hanenberg H, Auerbach AD, Smogorzewska A. Mutations of the SLX4 gene in Fanconi anemia. *Nat Genet*. 2011;43:142-6.

Longerich S, Kwon Y, Tsai MS, Hlaing AS, Kupfer GM, Sung P. Regulation of FANCD2 and FANCI monoubiquitination by their interaction and by DNA. *Nucleic Acids Res*. 2014;42:5657-70.

Machida YJ, Machida Y, Chen Y, Gurtan AM, Kupfer GM, D'Andrea AD, Dutta A. UBE2T is the E2 in the Fanconi anemia pathway and undergoes negative autoregulation. *Mol Cell*. 2006;23:589-96.

Meetei AR, Medhurst AL, Ling C, Xue Y, Singh TR, Bier P, Steltenpool J, Stone S, Dokal I, Mathew CG, Hoatlin M, Joenje H, de Winter JP, Wang W. A human ortholog of archaeal DNA repair protein Hef is defective in Fanconi anemia complementation group M. *Nat Genet*. 2005;37:958-63.

Meetei AR, de Winter JP, Medhurst AL, Wallisch M, Waisfisz Q, van de Vrugt HJ, Oostra AB, Yan Z, Ling C, Bishop CE, Hoatlin ME, Joenje H, Wang W. A novel ubiquitin ligase is deficient in Fanconi anemia. *Nat Genet* 2003;35:165-70

Miles JA, Frost MG, Carroll E, Rowe ML, Howard MJ, Sidhu A, Chaugule VK, Alpi AF, Walden H. The Fanconi anemia DNA repair pathway is regulated by an interaction between ubiquitin and the E2-like fold domain of FANCL. *J Biol Chem*. 2015;290:20995-1006.

Rickman KA, Lach FP, Abhyankar A, Donovan FX, Sanborn EM, Kennedy JA, Sougnez C, Gabriel SB, Elemento O, Chandrasekharappa SC, Schindler D, Auerbach AD, Smogorzewska A. Deficiency of UBE2T, the E2 ubiquitin ligase necessary for FANCD2 and FANCI ubiquitination, causes FA-T subtype of Fanconi anemia. *Cell Rep*. 2015;12:35-41.

Salewsky B, Schmiester M, Schindler D, Digweed M, Demuth I. The nuclease hSNM1B/Apollo is linked to the Fanconi anemia pathway via its interaction with FANCP/SLX4. *Hum Mol Genet*. 2012;21:4948-56.

Schuster B, Knies K, Stoecker C, Velleuer E, Friedl R, Gottwald-Mühlhauser B, de Winter JP, Schindler D. Whole exome sequencing reveals uncommon mutations in the recently identified Fanconi anemia gene SLX4/FANCP. *Hum Mutat*. 2013;34:93-6.

Singh TR, Ali AM, Paramasivam M, Pradhan A, Wahengbam K, Seidman MM, Meetei AR. ATR-dependent phosphorylation of FANCM at serine 1045 is essential for FANCM functions. *Cancer Res*. 2013;73:4300-10.

Somyajit K, Subramanya S, Nagaraju G. Distinct roles of FANCO/RAD51C protein in DNA damage signaling and repair: implications for Fanconi anemia and breast cancer susceptibility. *J Biol Chem*. 2012;287:3366-80.

Somyajit K, Subramanya S, Nagaraju G. RAD51C: a novel cancer susceptibility gene is linked to Fanconi anemia and breast cancer. *Carcinogenesis*. 2010;31:2031-8.

Stoecker C, Hain K, Schuster B, Hilhorst-Hofstee Y, Rooimans MA, Steltenpool J, Oostra AB, Eirich K, Korthof ET, Nieuwint AW, Jaspers NG, Bettecken T, Joenje H, Schindler D, Rouse J, de Winter JP. SLX4,

a coordinator of structure-specific endonucleases, is mutated in a new Fanconi anemia subtype. *Nat Genet.* 2011;43:138-41.

Tao Y, Jin C, Li X, Qi S, Chu L, Niu L, Yao X, Teng M. The structure of the FANCM-MHF complex reveals physical features for functional assembly. *Nat Commun.* 2012;3:782.

Vetro A, Iascone M, Limongelli I, Ameziane N, Gana S, Della Mina E, Giussani U, Ciccone R, Forlino A, Pezzoli L, Rooimans MA, van Essen AJ, Messa J, Rizzuti T, Bianchi P, Dorsman J, de Winter JP, Lalatta F, Zuffardi O. Loss-of-function FANCL mutations associate with severe Fanconi anemia overlapping the VACTERL association. *Hum Mutat.* 2015;36:562-8.

Virts EL, Jankowska A, Mackay C, Glaas MF, Wiek C, Kelich SL, Lottmann N, Kennedy FM, Marchal C, Lehnert E, Scharf RE, Dufour C, Lanciotti M, Farruggia P, Santoro A, Savasan S, Scheckenbach K, Schipper J, Wagenmann M, Lewis T, Leffak M, Farlow JL, Foroud TM, Honisch E, Niederacher D, Chakraborty SC, Vance GH, Pruss D, Timms KM, Lanchbury JS, Alpi AF, Hanenberg H. AluY-mediated germline deletion, duplication and somatic stem cell reversion in UBE2T defines a new subtype of Fanconi anemia. *Hum Mol Genet.* 2015;24:5093-108.

Wang Y, Leung JW, Jiang Y, Lowery MG, Do H, Vasquez KM, Chen J, Wang W, Li L. FANCM and FAAP24 maintain genome stability via cooperative as well as unique functions. *Mol Cell.* 2013;49:997-1009.

Xia B, Dorsman JC, Ameziane N, de Vries Y, Rooimans MA, Sheng Q, Pals G, Errami A, Gluckman E, Llera J, Wang W, Livingston DM, Joenje H, de Winter JP. Fanconi anemia is associated with a defect in the BRCA2 partner PALB2. *Nat Genet.* 2007;39:159-61.

Xia B, Sheng Q, Nakanishi K, Ohashi A, Wu J, Christ N, Liu X, Jasin M, Couch FJ, Livingston DM. Control of BRCA2 cellular and clinical functions by a nuclear partner, PALB2. *Mol Cell.* 2006;22:719-29.

Xue X, Sung P, Zhao X. Functions and regulation of the multitasking FANCM family of DNA motor proteins. *Genes Dev.* 2015;29:1777-88.