

Title: Tuberous Sclerosis Complex – Somatic Mosaicism

Authors: Northrup H, Koenig MK, Pearson DA, Au KS

Date: September 2015

Note: The following information is provided by the authors listed above and has not been reviewed by *GeneReviews* staff.

Somatic mosaicism has been described in many individuals with TSC and parents of children with TSC. However, an estimate of the frequency of somatic mosaicism is complicated by:

- The sensitivity of the pathogenic variant screening methods and the difficulty of detecting low-level mosaicism;
- The relative ease of detecting mosaicism for exonic / whole-gene deletions or duplications and the difficulty in detecting mosaicism for single nucleotide variants (SNVs) and small intra-exonic deletions or duplications;
- The likely overestimate of the percent of mosaicism in some studies because only affected persons who did not have a pathogenic variant identified by other molecular genetic test methods (sequencing and/or gene-targeted deletion/duplication analysis) were evaluated for mosaicism; and
- The fact that the phenotypes caused by mutation of *TSC1* and *TSC2* cannot be differentiated clinically; therefore, the proportion of individuals with each phenotype is unknown at the outset of studies screening for somatic mosaicism.

Note: (1) It is difficult to identify with confidence low-level mosaicism for pathogenic missense and other variants involving a few nucleotides because these sequence variants may be artifacts possibly introduced by the PCR processes. (2) It was estimated that levels of mosaicism greater than 20% in lymphocyte DNA can be detected with current exon sequencing and next-generation sequencing methods [Qin et al 2010, Authors, personal observation]. Using targeted deep next-generation sequencing of *TSC1* and *TSC2*, somatic mosaic pathogenic variants at 9.5% in an affected individual who did not have a pathogenic variant identified by exon sequence analysis or gene-targeted deletion/duplication analysis was reported [Nellist et al 2015]. However, it has not yet been determined whether levels of somatic mosaicism below 9.5% can be detected by deep next-generation sequencing.

The frequency of somatic mosaicism for large deletions and duplications of *TSC1* and *TSC2* in affected individuals (who did not have a pathogenic variant identified by sequence analysis or other similar methods) has been reported as about 5% (N=165) [Kozlowski et al 2007; Author, personal observation]. Seven of the eight individuals with mosaic pathogenic variants had deletions in *TSC2*; one had a duplication in *TSC2* [Kozlowski et al 2007].

Because somatic mosaicism for a pathogenic variant in *TSC2* has been reported in seven of 26 families with a combined TSC/PKD ([autosomal dominant polycystic kidney disease](#)) phenotype and six of 62 probands in another series, DNA testing of other

tissues (e.g., tumors, saliva, skin, and/or hair follicles) is warranted when somatic mosaicism is suspected and routine molecular genetic testing has not revealed a pathogenic variant.

No assay kit or service laboratory for testing somatic mosaicism exists at this time. Somatic mosaicism testing is currently only available from research groups on research basis.

References

- Kozlowski P, Roberts P, Dabora S, Franz D, Bissler J, Northrup H, Au KS, Lazarus R, Domanska-Pakiela D, Kotulska K, Jozwiak S, Kwiatkowski DJ. Identification of 54 large deletions/duplications in TSC1 and TSC2 using MLPA, and genotype-phenotype correlations. *Hum Genet.* 2007 May;121(3-4):389-400.
- Nellist M, Brouwer RW, Kockx CE, van Veghel-Plandsoen M, Withagen-Hermans C, Prins-Bakker L, Hoogeveen-Westerveld M, Mrcic A, van den Berg MM, Koopmans AE, de Wit MC, Jansen FE, Maat-Kievit AJ, van den Ouweland A, Halley D, de Klein A, van IJcken WF. Targeted Next Generation Sequencing reveals previously unidentified TSC1 and TSC2 mutations. *BMC Med Genet.* 2015 Feb 25;16:10.
- Qin W, Kozlowski P, Taillon BE, Bouffard P, Holmes AJ, Janne P, Camposano S, Thiele E, Franz D, Kwiatkowski DJ. Ultra deep sequencing detects a low rate of mosaic mutations in tuberous sclerosis complex. *Hum Genet.* 2010 Mar;127(5):573-82.