

Title: *PROP1*-Related Combined Pituitary Hormone Deficiency *GeneReview* – *LHX3* Variants

Author: de Graaff L

Date: August 2014

Note: The following information is provided by the authors listed above and has not been reviewed by *GeneReviews* staff.

Specific variants in *LHX3*

- Netchine et al [2000] identified homozygosity for two different *LHX3* variants in affected members of two unrelated consanguineous families with rigidity of the cervical spine and CPHD that involved all the anterior pituitary hormones except ACTH: one was a missense variant (p.Tyr116Cys) and one a 23-bp intragenic deletion.
- Bhangoo et al [2006] identified a homozygous 1-bp *LHX3* deletion (g.159delT) in a boy with CPHD and rigid cervical spine.
- Pfaeffle et al [2007] identified the *LHX3* variants p.Glu173Ter, p.Ala210Val, p.Trp224Ter, and a deletion of the entire gene in seven affected individuals from four families in a cohort of 366 persons from 342 families with CPHD. They concluded that *LHX3* variants are a rare cause of CPHD and that they are associated with mild or severe deficiencies of GH, PRL, TSH, and LH/FSH in all cases. They noted that limited neck rotation was not present in three sibs with CPHD who had an *LHX3* nonsense variant.
- Rajab et al [2008] identified a homozygous large intragenic deletion (3088-bp del) and a homozygous nonsense variant (p.Lys50Ter) in four persons from two unrelated consanguineous families with CPHD and neonatal hypoglycemia, short neck with limited rotation, and mild sensorineural hearing loss. Based on the observations that sensorineural hearing loss was found when three of the individuals with pathogenic variants studied by Netchine et al [2000] were reexamined and that ACTH deficiency was present in one of the persons reported by Rajab et al [2008], they proposed that the phenotypic spectrum of *LHX3* variants includes CPHD (with or without ACTH deficiency) with limited neck rotation and mild sensorineural hearing loss.
- Kriström et al [2009] found a novel, recessive, A-G splice-acceptor site variation in exon 3, resulting in deletion of the homeodomain and the C-terminus. Like the individuals reported by Rajab et al [2008], the individuals reported by Kriström et al [2009] had CPHD, restricted neck rotation, and a severe hearing defect.
- In 2011, Bonfig et al reported a novel *LHX3* mutation (R77X), which is associated with combined pituitary hormone deficiency including ACTH deficiency, short neck, and sensorineural hearing loss.
- In 2012, Bechtold-Dalla Pozza et al found a novel homozygous *LHX3* mutation (T194R) was detected in two female siblings from related parents. The index

patient had CPHD featuring deficiencies of GH, LH, FSH, PRL, and TSH, with later onset of ACTH deficiency. She also had a hypoplastic anterior pituitary, respiratory distress, hearing impairment, and limited neck rotation. Both patients had neonatal complications. The affected amino acid is conserved in the DNA-binding homeodomain. Computer modeling predicted that the T194R change would alter the homeodomain structure. The T194R protein did not bind tested LHX3 DNA recognition sites and did not activate the α -glycoprotein and PRL target genes.

- In 2012, Sobrier et al described two new LHX3 defects in a patient with syndromic CPHD, severe scoliosis, and normal intelligence. The paternally inherited c.252-3C>G mutation, which disrupts an acceptor splice site, would lead to severely truncated proteins containing a single LIM domain, resembling LIM-only proteins. Coexpression studies revealed the dominant-negative effect of this LIM-only protein over the wild-type LHX3. The maternally inherited p.Cys118Tyr mutation results in partial loss of transcriptional activity and synergy with POU1F1.

References

- Bechtold-Dalla Pozza S, Hiedl S, Roeb J, Lohse P, Malik RE, Park S, Durán-Prado M, Rhodes SJ. A recessive mutation resulting in a disabling amino acid substitution (T194R) in the LHX3 homeodomain causes combined pituitary hormone deficiency. *Horm Res Paediatr.* 2012;77:41-51
- Bhangoo AP, Hunter CS, Savage JJ, Anhalt H, Pavlakis S, Walvoord EC, Ten S, Rhodes SJ. Clinical case seminar: a novel LHX3 mutation presenting as combined pituitary hormonal deficiency. *J Clin Endocrinol Metab.* 2006;91:747-53.
- Bonfig W, Krude H, Schmidt H A novel mutation of LHX3 is associated with combined pituitary hormone deficiency including ACTH deficiency, sensorineural hearing loss, and short neck-a case report and review of the literature. *Eur J Pediatr.* 2011;170:1017-21.
- Kriström B, Zdunek AM, Rydh A, Jonsson H, Sehlin P, Escher SA. A novel mutation in the LIM homeobox 3 gene is responsible for combined pituitary hormone deficiency, hearing impairment, and vertebral malformations. *J Clin Endocrinol Metab.* 2009;94:1154-61.
- Netchine I, Sobrier ML, Krude H, Schnabel D, Maghnie M, Marcos E, Duriez B, Cacheux V, Moers A, Goossens M, Grüters A, Amselem S (2000) Mutations in LHX3 result in a new syndrome revealed by combined pituitary hormone deficiency. *Nat Genet.* 25:182-6.
- Pfaeffle RW, Savage JJ, Hunter CS, Palme C, Ahlmann M, Kumar P, Bellone J, Schoenau E, Korsch E, Brämwig JH, Stobbe HM, Blum WF, Rhodes SJ. Four novel mutations of the LHX3 gene cause combined pituitary hormone deficiencies with or without limited neck rotation. *J Clin Endocrinol Metab.* 2007;92:1909-19.
- Rajab A, Kelberman D, de Castro SC, Biebermann H, Shaikh H, Pearce K, Hall CM, Shaikh G, Gerrelli D, Grueters A, Krude H, Dattani MT. Novel mutations in LHX3 are associated with hypopituitarism and sensorineural hearing loss. *Hum Mol Genet.* 2008;17:2150-9.
- Sobrier ML, Brachet C, Vié-Luton MP, Perez C, Copin B, Legendre M, Heinrichs C, Amselem S. Symptomatic heterozygotes and prenatal diagnoses in a nonconsanguineous family with syndromic combined pituitary hormone deficiency resulting from two novel LHX3 mutations. *J Clin Endocrinol Metab.* 2012;97:E503-9.