



References

Gaujoux S, Salenave S, Ronot M, Rangheard AS, Cros J, Belghiti J, Sauvanet A, Ruszniewski P, Chanson P. Hepatobiliary and pancreatic neoplasms in patients with McCune-Albright syndrome. *J Clin Endocrinol Metab.* 2014.

Parvanescu A, Cros J, Ronot M, Hentic O, Grybek V, Couvelard A, Levy P, Chanson P, Ruszniewski P, Sauvanet A, Gaujoux S. Lessons from McCune-Albright syndrome-associated intraductal papillary mucinous neoplasms: GNAS-activating mutations in pancreatic carcinogenesis. *JAMA Surg.* 2014.

Legend

GERD = gastroesophageal reflux disease; H&P = history & physical exam; LFTs = liver function tests; MAS = McCune-Albright syndrome; MRCP = magnetic resonance cholangiopancreatography

¹Performed at initial presentation, regardless of clinical symptoms. ² Age is not based upon clinical evidence, but on age at which patients may undergo MRI/MRCP without requiring sedation. ³Age of onset of pancreatic cyst development is not established, therefore clinical monitoring for signs of involvement in these patients is indicated. ⁴Patients should continue to be monitored clinically for new signs of gastrointestinal/pancreatic involvement including pancreatitis and diabetes.